

## RESTRICTED USE PESTICIDE

**Due to toxicity to fish and aquatic organisms.**

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.



EPA REG. NO. 279-3426

## **2(ee) Recommendation**

**FOR DISTRIBUTION AND USE IN  
ALL REGISTERED STATES (Except Texas)**

**THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL  
DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY  
FMC CORPORATION UNDER FIFRA SECTION 2(ee) AND HAS  
NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.**

**This recommendation for MUSTANG® MAXX Insecticide is valid until 12/31/2030, or until withdrawn, canceled or suspended.**

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND  
PRECAUTIONS ON THE EPA REGISTERED MUSTANG® MAXX INSECTICIDE (EPA REG. NO. 279-3426) LABEL MUST BE  
FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.**

**For the suppression or control of spotted lanternfly in listed crops.**

CROP	RATE OF APPLICATION	RESTRICTIONS
Grape	4.0 fl. oz/A (0.025 lb ai/A)	<ul style="list-style-type: none"><li>• <b>Do not</b> apply more than 4.0 fl. oz/A (0.025 lb ai/A) of product per application.</li><li>• <b>Do not</b> apply more than 24 fl. oz/A (0.15 lb ai/A) of product per year.</li><li>• <b>Do not</b> make more than 6 applications per year.</li><li>• <b>Do not</b> make applications less than 7 days apart.</li><li>• <b>Do not</b> apply within 1 day of harvest.</li></ul>

### **DIRECTIONS FOR USE:**

Apply by ground or air equipment using sufficient water to obtain full coverage of foliage.

**Application by ground:** Apply in a minimum of 10 gal/A of finished spray by ground.

**Application by air:** Apply in a minimum of 2 gal/A of finished spray by air.



FMC Corporation  
2929 Walnut Street  
Philadelphia, PA, 19103

CROP	RATE OF APPLICATION	RESTRICTIONS
<b>Pome Fruit Group, including: Apple, Crabapple, Loquat, Mayhaw, Pear, Asian Pear, and Quince</b>	<b>4.0 fl. oz/A (0.025 lb ai/A)</b>	<ul style="list-style-type: none"> <li>• <b>Do not</b> apply more than 4.0 fl. oz/A (0.025 lb ai/A) of product per application.</li> <li>• <b>Do not</b> apply more than 24 fl. oz/A (0.15 lb ai/A) of product per year.</li> <li>• <b>Do not</b> make more than 6 applications per year.</li> <li>• <b>Do not</b> apply as a ULV spray.</li> <li>• <b>Do not</b> feed or allow livestock to graze on cover crops from treated orchards.</li> <li>• <b>Do not</b> make applications less than 7 days apart.</li> <li>• <b>Do not</b> apply within 14 days of harvest.</li> </ul>

**DIRECTIONS FOR USE:**

Avoid applications when honey bees are actively foraging by applying during the early morning or evening hours.

Apply by ground or air equipment using sufficient water to obtain full coverage of foliage.

**Application by ground:** Apply in a minimum of 20 gal/A of finished spray for concentrate spray, or in a minimum of 100 gal/A of finished spray for dilute spray.

**Application by air:** Apply in a minimum of 10 gal/A of finished spray.

CROP	RATE OF APPLICATION	RESTRICTIONS
<b>Stone Fruit Group, including: Apricot, Cherry (Sweet and Tart), Nectarine, Peach, Plum (including Chickasaw Plum, Damson Plum and Japanese Plum), Plumcot, and Prune (Fresh)</b>	<b>4.0 fl. oz/A (0.025 lb ai/A)</b>	<ul style="list-style-type: none"> <li>• <b>Do not</b> apply more than 4.0 fl. oz/A (0.025 lb ai/A) of product per application.</li> <li>• <b>Do not</b> apply more than 24 fl. oz/A (0.15 lb ai/A) of product per year.</li> <li>• <b>Do not</b> make more than 6 applications per year.</li> <li>• <b>Do not</b> apply as a ULV spray.</li> <li>• <b>Do not</b> feed or allow livestock to graze on cover crops from treated orchards.</li> <li>• <b>Do not</b> make applications less than 7 days apart.</li> <li>• <b>Do not</b> apply within 3 days of harvest for cherries and 14 days of harvest for all other listed stone fruits.</li> </ul>

**DIRECTIONS FOR USE:**

Apply by ground or air equipment using sufficient water to obtain full coverage of foliage.

**Application by ground:** Apply in a minimum of 20 gal/A of finished spray for concentrate spray, or in a minimum of 100 gal/A of finished spray for dilute spray.

**Application by air:** Apply in a minimum of 10 gal/A of finished spray.

**REMARKS:**

Thorough coverage is essential to achieve control.

Multiple applications may be required to maintain control and prevent reinfestation from migrating adults into the crop. Spotted lanternflies are mobile and may reinfest crops quickly. If another treatment is needed before the minimum retreatment interval, use another insecticide to maintain control.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED MUSTANG® MAXX INSECTICIDE LABEL (EPA REG. NO. 279-3426) MUST BE FOLLOWED. THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.**

Mustang and FMC are trademarks of FMC Corporation or an affiliate. ©2025 FMC Corporation.



FMC Corporation  
2929 Walnut Street  
Philadelphia, PA, 19103