

EPA Reg. No. 2935-560

# For Distribution and Use Only in the States of Arizona and California

# REDUCED DILUENT FOR LABELED DOSAGE ON LEAFY VEGETABLES – EXCEPT BRASSICA VEGETABLES (Crop Group 4)

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Before using SPÄRRA in accordance with this recommendation, read and follow all applicable directions for use, restrictions, and precautions on the EPA registered label, as found on the package.
- This recommendation should be in the possession of the user at the time of application.
- This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

### **DIRECTIONS FOR USE**

## LEAFY VEGETABLES - EXCEPT BRASSICA VEGETABLES (Crop Group 4)

Including, but not limited to, Amaranth, Leafy, Chinese Spinach; Arugula; Cardoon; Celery; Celery, Chinese; Celtuce; Chervil; Chrysanthemum, Edible-Leaved, Garland; Corn Salad; Cress, Garland, Upland; Dandelion; Dock (Sorrel); Endive; Fennel, Florence; Lettuce, Head and Lettuce; Orach; Parsley; Purslane, Garden, Winter; Radicchio (Red Chicory); Rhubarb; Spinach; Spinach, New Zealand and Vine; and Swiss Chard.

NOTE: Refer to APPLICATION PROGRAM for disease prevention and control programs.

DISEASE	APPLICATION METHOD	RATE/APPLICATION
Downy Mildew Fusarium Phytophthora Powdery Mildew Pythium Rhizoctonia	Foliar – Ground	1-3 quarts per acre in a minimum of 20 gallons of water.

#### **APPLICATION PROGRAM**

**Disease Prevention Program:** Apply the lower specified rate at 7- to 28-day intervals after plants become established. Do not apply SPÄRRA at intervals less than 3 days.

**Disease Control Program:** Apply the higher specified rate at 7- to 14-day intervals until control is reached. Under severe circumstances, applications can be made at intervals of up to every three days. Do not apply SPÄRRA at intervals less than 3 days. Consult with Farm Advisor, Licensed Agricultural Pest Control Advisor or Certified Crop Advisor to determine disease severity.

WILBUR-ELLIS® logo and FUNGICIDE® logo are registered trademarks and SPÄRRA is a trademark of Wilbur-Ellis Company LLC.

2023-0830

