

Sumilarv® 0.5G

Sachets (WSP)

Sumilarv® WSP is an easy to apply sachet containing a powerful granule formulation larvicide for achieving longer lasting vector control. Delivering proven efficacy at low dose rates with a quickly dissolving WSP sachet, Sumilarv WSP can dramatically reduce re-application costs because of its unique formulation and non-flushing properties.

- Provides up to 150 days residual control of mosquitoes with one dose (3 WSP) in storm water catch basins
- Prevents mosquitoes from becoming breeding, biting adults
- Works well even in areas that dry out and rewet
- Non-flushing formulation
- Remains active after large rain events
- Distinct Mode Of Action (MOA)
- Contains Pyriproxyfen IGR
- Effective in polluted and non-polluted water
- Remains active through seasonal temperature changes

Product Specifications

Signal Word	Caution
Packaging	25 gram WSP (sachet) 40 WSP per bag (1 kg) 10 bags per case (10 kg)
EPA Registration Number	1021-2818
Product Handling	For use in outdoor mosquito breeding sites
Stability of Undiluted Product	Not applicable – sand granule
Stability of Diluted Product	Not applicable – sand granule
Appearance	Fine sand granule
Odor	No odor
Active Ingredients	Pyriproxyfen 0.5%
Flammability	Not applicable
Mode of Action	Insect growth regulator
Class of Chemistry	IGR
Respirator Required	Not required
Mix or Dilute in	Not applicable – ready-to-use
Activity	IGR and residual activity
Shelf Life	3 years or more if stored at room temperature



Use Areas

Outdoor Mosquito Breeding Areas, such as:

- Manmade temporary or permanent water holding sites, including
 - Catch basins, water retention or detention ponds, ornamental ponds, fountains, gutters, sewers, waste water impoundments, waste settling ponds, livestock ponds, animal waste lagoons, abandoned swimming pools, and French drains
- Natural & Artificial Water Holding Containers:
 - Tire piles, hollow trees & tree holes, potted plants, bird baths, rain barrels, flooded roof tops, flower pots, buckets, abandoned vehicles and salvage yards

Method of Application

Outdoor:

- Simply toss WSP sachet
- Hand applied for cryptic or frequently treated breeding sites

Best Practices When Using Sumilarv WSP

Identify area as a breeding site for mosquitoes and verify larval activity.

Application Tips

Estimate the volume of water for the treatment site, prior to application using the following calculation method: Total cu. Ft. of target treatment site (ft.³) = surface area (ft.²) x average depth (ft.)

Note: 7.5 gallons per cu. ft.

Total water volume = Total cu. ft. of treatment site x 7.5 gallons

Application Rates to Use for the Control of Mosquito Larvae

Treatment site (Volume of Water in Gallons)	Water Depth Less than or Equal to (ft.)	Amount of Sumilarv 0.5G Sachets (WSP)	lb. AI Sumilarv
0 – 500	1	1	0.00028
500 – 1500	2	2	0.00055
1500 – 3000	4	3	0.00083
3000 – 4500	6	4	0.00110
4500 – 6000	8	5	0.00138

Apply at the rate specified in the application rate table above:

Use Sumilarv WSP as part of an IPM program


For best results, use Sumilarv WSP as part of an integrated pest management program to control the targeted pest and the diseases they vector.

Effective Mosquito Application

- Identify mosquito breeding site.
- Calculate the size & water volume of treatment area.
- Simply toss the ready-to-use WSP sachet into the treatment site.
- Continue to monitor or surveil treatment areas to verify adult emergence inhibition has ceased.
- Please note: Sumilarv WSP will not affect mosquito breeding activities, mosquito larval development, but will prevent adult mosquitoes from emerging from the pupae stage.

Key Insects Controlled

Mosquitoes



Source: Field Study, 2020, Dr. Tianyun (Steven) Su.

Always read and follow label and SDS directions.

To learn more, visit www.mgk.com, call 1-800-645-6466 or send an email to brands@mgk.com.

Sumilarv® 0.5G
Sachets (WSP)