

SUPPLEMENTAL LABELING

Buckle® Microactiv Herbicide

EPA Reg. No. 10163-289

GRANULAR SELECTIVE HERBICIDE FOR FALL PREPLANT APPLICATION TO CONTROL WILD OATS AND PIGEONGRASS IN SPRING WHEAT.

READ AND FOLLOW THE ENTIRE LABEL FOR BUCKLE® MICROACTIV HERBICIDE BEFORE PROCEEDING WITH THE USE AND DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Buckle Microactiv Herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

FOR USE ONLY WITHIN THE FOLLOWING COUNTIES IN NORTH AND SOUTH DAKOTA:

North Dakota:

Adams, Billings, Bottineau, Bowman, Burleigh, Burke, Divide, Dunn, Emmons, Grant, Golden Valley, Hettinger, Kidder, McHenry, McKenzie, McLean, Mercer, Morton, Mountrail, Oliver, Renville, Sheridan, Sioux, Slope, Stark, Ward, and Williams.

South Dakota:

Corson, Dewey, Harding, Perkins, and Ziebach.

KEEP OUT OF REACH OF CHILDREN CAUTION

In case of a medical emergency involving this product, call toll free 1-888-487-0798.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

This label must be in possession of the user at the time of herbicide application.

Apply Buckle Microactiv Herbicide at a rate of 10 pounds per acre in the fall prior to spring planting of Spring Wheat. Average soil temperature at the 2-inch depth must be 40° F or less. Application is limited to one per growing season and must not exceed 10 lbs. of this product per acre.

Incorporate fall Buckle Microactiv Herbicide applications within 24 hours with a field cultivator. For the best results a second incorporation should be done in the spring at a right angle with a field cultivator with harrow attachment. If no spring tillage is anticipated, both incorporations should be done in the fall with a minimum of 5 days between first and second incorporations. For best results, care must be taken to avoid working the soil deeper than the first tillage operation.

Do not fall-apply Buckle Microactiv Herbicide for spring wheat on soil that has been treated with 0.5 lbs ai/acre or greater of trifluralin, or other dinitroaniline herbicide, during that calendar year.

NOTE: Spring wheat varieties identified as susceptible to early injury from preplant applications of Buckle Microactiv Herbicide include: Alex, Amidon, Butte 86, Ellar, Era, Erik, Lew and Newana.

To minimize stress and allow for uniform germination, a seeding depth of no more than 1.5 to 2 inches is recommended. Delaying spring planting until soil temperatures have reached 45°F at the two inch depth will reduce the potential for crop injury.

DO NOT APPLY TO HILLTOPS.

Seed treatments are recommended to reduce potential stress to germinating seedlings from disease or insects.

Do not use on fields if the field will be irrigated during the spring wheat growing season.

Note: Slight stand reductions do not normally affect yield.

DO NOT APPLY BUCKLE MICROACTIV HERBICIDE TO A STRIP OR AREA IN THE FIELD THAT WILL SERVE AS AN UNTREATED COMPARISON FOR WEED CONTROL AND CROP TOLERANCE.

Read the "NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS" in the label for Buckle Microactiv Herbicide before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Buckle® is a registered trademark of Gowan Company LLC.