SUPPLEMENTAL LABELING

RESTRICTED USE PESTICIDE DUE TO ACUTE TOXICITY

TO BE USED BY CERTIFIED APPLICATORS ONLY – NOT TO BE USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A CERTIFIED APPLICATOR, EXCEPT THAT UNCERTIFIED PERSONS MAY TRANSPORT CONTAINERS AS PROVIDED UNDER DIRECTIONS FOR USE.

Syngenta Crop Protection, LLC

P. O. Box 18300 Greensboro, North Carolina, 27419-8300 SCP 1652A-S13A 0822

PARAQUAT DICHLORIDE	GROUP	22	HERBICIDE	
---------------------	-------	----	-----------	--

Gramoxone® SL 3.0 HERBICIDE

EPA Reg. No. 100-1652

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

Active Ingredients:

Paraquat dichloride*	
Other Ingredients:	56.1%
Total:	100.0%

*CAS No. 4685-14-7

Gramoxone SL 3.0 is formulated as a liquid containing 3.0 pounds paraquat cation per gallon as 4.134 pounds paraquat dichloride per gallon.

Gramoxone SL 3.0 contains alerting agent (odor), emetic, and dye.

KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.) See additional precautionary statements and directions for use inside booklet.

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Gramoxone SL 3.0 as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



DIRECTIONS FOR USE RESTRICTED USE PESTICIDE

Flowers, WA Only

Crops (Including all cultivars, varieties, and/or hybrids of these)			
Iris	Narcissus	Tulip	

Target Weed	Rate pt/A	Application Timing	Use Directions
Refer to Section 7.0.	1.3 – 2.5	Prior to bulb emergence	For use in Washington Only for control of annual and broadleaf weeds and grasses in tulips, narcissus and iris.
			Apply by ground as a broadcast spray in a minimum of 30 gal/A.
			When using this product in Eastern Washington, refer to the Washington State Department of Agriculture's restrictions on the use of desiccant herbicides.
			Apply by ground as a broadcast spray in a minimum of 30 gal/A.
			Add a nonionic surfactant at least 50% surface active ingredient at 1-2 pints per 100 gallons of dilute spray.
			It is essential to obtain complete coverage of target weeds to get adequate control.
			See Section 4.4.4 of the Gramoxone SL 3.0 label for adjuvant recommendations.

Tank Mix Options:

Gramoxone SL 3.0 can be tank mixed with other herbicides labeled for use on flowers applied by the same methods and at the same timings. Refer to Section 4.4.2 for tank-mix precautions in the Gramoxone SL 3.0 label.

Resistance Management:

Refer to Section 3.1.

USE RESTRICTIONS

- Refer to Section 6.1 for additional product use
 Maximum Single Application Rate: 2.5 pt/A Refer to **Section 6.1** for additional product use restrictions.
- 3. **REI:** 48 hours
- 4. Minimum Application Interval: 7 days
- 5. Maximum Annual Rate: 5.0 pt/A
 - **DO NOT** exceed 1.8 lb cation/A/year of paraguat-containing products.
- 6. **DO NOT** make more than 2 application per year.
- 7. **DO NOT** apply this product through any type of irrigation system.
- 8. Pre-Harvest Interval (PHI): Harvest crop at normal harvest maturity.

Gramoxone® and the SYNGENTA Logo are Trademarks of a Syngenta Group Company.

©2022 Syngenta

For non-emergency (e.g., current product information), call Syngenta Crop Protection at 1-800-334-9481.

Manufactured for: Syngenta Crop Protection, LLC P.O. Box 18300 Greensboro, North Carolina, 27419-8300