



RESTRICTED USE PESTICIDE

TOXIC TO FISH AND AQUATIC ORGANISMS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS, OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

PRODUCT BULLETIN

FIFRA SECTION 2(ee) RECOMMENDATION

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. Follow all applicable directions, restrictions, and precautions on the EPA-registered label. This product bulletin must be in the possession of the user at the time of application.

Date: January 27, 2026

Product: Bifender® SM EPA Reg. No. 89118-8

Use: Bifender SM for In-Season Control of Corn Rootworm Larvae

State: CO, CT, DE, MI, MN, IA, ID, IN, KS, MO, ND, NE, NV, OH, SD, WA, WI

DIRECTIONS FOR USE

CORN

FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED, SWEET CORN, SWEET CORN GROWN FOR SEED

SOIL APPLICATION	
TARGET PEST	USE RATES
Corn Rootworm Larvae (Northern, Southern, and Western)	11.6 fl oz/A (0.1 lb ai/A)
Instructions: Apply 11.6 fl oz/a Bifender SM in a minimum of 0.5 acre-inch water via overhead irrigation (sprinklers including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, drip irrigation, or hand move irrigation systems). For optimum control of Corn Rootworm Larvae in a single application, target application of Bifender SM when soils are moist and when accumulated soil growing degree days (base 52°F) are between 550 and 684 for the calendar year. Refer to the Bifender SM EPA-registered label for all sprinkler chemigation directions and restrictions	
RESTRICTIONS: DO NOT make more than 3 foliar applications to field corn and 2 foliar applications to sweet corn per year.	



- **DO NOT** apply more than 0.20 pound active ingredient per acre per year as an at-plant application.
- **For field corn: DO NOT** apply more than 0.30 pound active ingredient per acre per year including at-plant, PPI & PRE, and foliar applications of Bifender SM and other bifenthrin containing products.
- **For sweet corn: DO NOT** apply more than 0.20 pound active ingredient per acre total per year including at-plant, PPI & PRE, and foliar applications of other bifenthrin products.
- **DO NOT** apply within 30 days of harvest for field corn (grain and silage), popcorn, or field corn grown for seed.
- **DO NOT** apply within 1 day of harvest for sweet corn, or sweet corn grown for seed.
- **DO NOT** graze livestock in treated areas or cut treated crops for feed within 1 day of harvest for sweet corn, or sweet corn grown for seed.
- Use of ultra low volume (ULV) application on corn is prohibited.
- **DO NOT** make aerial or ground applications to corn if heavy rainfall is imminent.