

**RESTRICTED USE PESTICIDE**

DUE TO TOXICITY TO NON-TARGET INVERTEBRATES, MAMMALS, AND AQUATIC ORGANISMS
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION,
AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

PRODUCT BULLETIN**FIFRA SECTION 2(ee) RECOMMENDATION**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. Follow all applicable directions, restrictions, and precautions on the EPA-registered label. This product bulletin must be in the possession of the user at the time of application.

Date: February 17, 2026

Product: Averland® SM EPA Reg. No. 89118-7

Use: Averland SM for suppression of nematodes in Potatoes

States: CO, DE, ID, FL, MO, NC, ND, NV, RI, WA, WI, WY

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Potatoes**Soil Application**

PESTS	USE RATES
	fl oz product/A (lb a.i./A)
Suppression of nematodes including: Columbia root-knot nematode, northern root-knot nematode and root-lesion nematode	2.34-2.4 (0.0186 – 0.019)
Sprinkler Chemigation Instructions: • Apply 2.34 to 2.4 fl oz/A Averland SM in a minimum of 0.5 acre-inch water via overhead irrigation (sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; furrow; border or drip (trickle) irrigation systems) • For optimum suppression on nematodes, apply Averland SM through chemigation when soils are moist • Make up to 3 sequential overhead sprinkler applications 7-14 days apart for the suppression of nematodes • Refer to the Averland SM EPA-registered label for all sprinkler chemigation directions and restrictions • Make applications as part of an overall nematode management program.	
Use Restrictions:	



- **Maximum Single Application Rate:** 2.4 fl oz Averland SM/A (0.019 lb a.i./A) per application.
- **Minimum Application Interval:** 7 days.
- **Maximum Annual Rate:** 7.0 fl oz Averland SM/A/calendar year.
- **DO NOT** exceed 0.056 lb a.i./A/calendar year from all abamectin containing products including all application types (seed treatments, soil, foliar).
- **DO NOT** make more than 2 sequential applications of Averland SM or any other foliarly applied abamectin-containing product.
- **DO NOT** feed or allow livestock to graze treated foliage.
- **Pre-Harvest Interval (PHI):** 14 days.

Follow all applicable directions, restrictions, and precautions on the EPA-registered label.