

Drexel Simazine 4L is a state restricted use pesticide and is to be distributed only by licensed pesticide dealers. Only certified applicators or persons under the direct supervision of a certified applicator may apply Drexel Simazine 4L.

**FIFRA SECTION 24(c) SPECIAL LOCAL NEED (SLN) LABEL
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON**

SIMAZINE	GROUP	5	HERBICIDE
----------	-------	---	-----------



Drexel

Simazine 4L

Herbicide

EPA Reg. No. 19713-60
EPA SLN No. WA-260006

This label for Drexel Simazine 4L expires and must not be distributed in or used in accordance with this SLN registration after December 31, 2030.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the EPA registered label.
- Read and follow all applicable directions and restrictions in the Environmental Hazards section of the EPA registered label to protect groundwater, surface water and non-target organisms.

DIRECTIONS FOR USE

Brussels Sprout Grown for Seed Only and Cabbage Grown for Seed Only / Annual Weeds (such as Annual bluegrass, Chickweed, Groundsel, Pineappleweed, Prostrate knotweed, Smartweed, Wild buckwheat)

Make two applications of 1.6 pints (0.8 lb. active ingredient) of Simazine 4L per acre. Apply as a broadcast spray in a minimum 20 gallons of water per acre by ground application only.

Make the first application in the Fall after the start of the Winter rainfall period (to control Fall germinating weeds), but before the ground is frozen. Make the second application in the early Spring just after the last cultivation. Because these applications are broadcast over-the-top sprays and not directed or shielded applications, they should be made before there is any significant emergence of seedling annual weeds.

Restrictions / Precautions

- Do not enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of 12 hours.
- Do not apply Simazine 4L to *Sand, Loamy sand* or gravelly areas of the field. Do not apply to areas of the field where water will accumulate.
- Make only two applications of Simazine 4L per acre.
- For use under this SLN label, do not apply Simazine 4L through any type of irrigation system.
- Aerial application is prohibited.
- Land treated with Simazine 4L should not be planted to any crop except Corn until the following year as injury may occur. After the last harvest of the treated crop, plow and thoroughly till the soil in the Fall or Spring to minimize possible injury to Spring-seeded rotational crops. Avoid rotating to crops susceptible to injury from Simazine.
- All Brussels sprout / Cabbage seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the WSDA immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter, or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the growers(s), and the date of disposal.
- No portion of the Brussels sprout / Cabbage plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
- Brussels sprout / Cabbage seed shall bear a tag or container label which forbids the use of the seed for human consumption or animal feed.
- Brussels sprout / Cabbage seed may not be distributed for human consumption or animal feed.

60SLNWA-0326*P(BRUSSELS SPROUT)
Simazine 4L SLN Label

WSDA Container Disposal Guidance

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling a cleaned container is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA website under the Waste Pesticide Program. Cleaned containers may also be disposed of in a sanitary landfill if permitted by the county. Burning is not a legal method of container disposal in Washington.



24(c) Registrant:
Drexel Chemical Company
P.O. Box 13327, Memphis, TN 38113-0327
SINCE 1972

The DREXEL logo is a registered trademark of Drexel Chemical Company.