

Aggressor[®] AX

HERBICIDE



EPA Reg. No. 45002-51

Product Bulletin

UPDATED WINTER WHEAT USE RATE (according to EPA approved label dated 2/26/2026)

For use in the states of: Alabama, Colorado, Idaho, Indiana, Kansas, Minnesota, Montana, North Carolina, North Dakota, Nebraska, New Mexico, Ohio, Oklahoma, Oregon, South Carolina, South Dakota, Texas, Utah, Virginia, Washington, Wyoming

KEEP OUT OF REACH OF CHILDREN CAUTION/ PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

**For 24-hour chemical spill, leak, fire, exposure or accident response information,
call CHEMTREC toll free at 1-800-424-9300.**

DIRECTIONS FOR USE

It is a violation of Federal law to use Aggressor[®] AX in a manner inconsistent with its labeling. Read the label affixed to the container of Aggressor[®] AX before applying. Use of Aggressor[®] AX according to this labeling is subject to the use precautions imposed by the label affixed to the container of Aggressor[®] AX.

FOR DISTRIBUTION AND USE ONLY IN THE STATES LISTED ABOVE

This PRODUCT BULLETIN, which accompanies the sale and packaging of Aggressor[®] AX, must be in the possession of the user at the time of pesticide application.

Refer to Albaugh Technical Bulletins or contact your regional Albaugh representative for approved herbicide tank-mix options with Aggressor[®] AX

CoAXium WINTER WHEAT with the AXigen Trait

(based on US EPA approved label dated 2/26/2026)

USE RATE

Apply 8 - 16 fl. oz. Aggressor[®] AX (0.055 – 0.11 lb AI) in a single application per acre. **See Weeds Controlled section (below) for detailed use rate specifications.**

Fall applications to winter wheat can be made with Aggressor[®] AX at rates from 8 – 16 fl. oz. (0.055 – 0.11 lb AI) / Acre.

Spring applications made on winter wheat can be made with Aggressor[®] AX at rates from 8 – 16 fl. oz. (0.055 – 0.11 lb AI) / Acre.

Fall and spring split applications can be made with Aggressor[®] AX at a single rate of 8 fl. oz. (0.055 lb AI) / Acre in the fall followed by 8 fl. oz. (0.055 lb AI) / Acre in the spring.

See WEEDS CONTROLLED AND RATE SELECTION Table (below) for detailed use rate specifications.

RESTRICTIONS:

- **DO NOT** apply more than 16 fl. oz. Aggressor[®] AX/acre (0.11 lb AI) / Acre per year.
- **DO NOT** apply more than 16 fl. oz. Aggressor[®] AX /acre (0.11 lb AI) / Acre in a single application.
- **DO NOT** make more than 1 application per year of Aggressor[®] AX at the 16 fl. oz. (0.11 lb AI) or 2 applications per year at 8 fl. oz. (0.055 lb AI). **DO NOT** exceed 16 fl. oz. (0.11 lbs. AI) per acre per year.
- **DO NOT** make a second application within 14 days of first application of Aggressor[®] AX.
- **DO NOT** harvest treated wheat for forage or hay within 60 days of the last application of Aggressor[®] AX. Treated wheat can be harvested for grain or straw at maturity.

WEEDS CONTROLLED AND RATE SELECTION (based on US EPA approved label dated 2/26/2026)
(Refer to the "Winter Wheat Restrictions" (below) for application limitations for single and split applications)

Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to quizalofop-P-ethyl or other Group 1 herbicides have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of Aggressor® AX specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed.

	Size at Application (height inches)	Aggressor® AX Applied Alone fl. oz. product / A (lbs AI / A)	Aggressor® AX Tank Mixed with Broadleaf Herbicide* fl.oz. product / A (lbs AI / A)
ANNUAL GRASSES**			
Volunteer Corn (<i>Zea mays</i>)***	6-30	8 – 16 (0.055 – 0.11 lbs AI)	10 – 16 (0.068 – 0.11 lbs AI)
Foxtail, Giant (<i>Setaria faberi</i>)	2-4 (pre-tiller)		
Johnsongrass, Seedling (<i>Sorghum halepense</i>)	2-8		
Shattercane (<i>Sorghum bicolor</i>)	6-12		
Wild Proso Millet (<i>Panicum miliaceum</i>)	2-6		
Crowfootgrass (<i>Dactyloctenium aegyptium</i>)	2-6	8 – 16 (0.055 – 0.11 lbs AI)	16 (0.11 lbs AI)
Fall Panicum (<i>Panicum dichotomifolium</i>)	2-6		8-16 (0.055 – 0.11 lbs AI)
Field Sandbur (<i>Cenchrus incertus</i>)	2-6		
Foxtail, Bristly (<i>Setaria verticillata</i>)	2-4		16 (0.11 lbs AI)
Foxtail, Giant (<i>Setaria faberi</i>)	2-8		†
Foxtail, Green (<i>Setaria viridis</i>)	2-4		10 – 16 (0.068 – 0.11 lbs AI)
Foxtail, Yellow (<i>Setaria lutescens</i>)	2-4		
Goosegrass (<i>Eleusine indica</i>)	2-6 ‡		
Itchgrass (<i>Rottboellia exaltata</i>)	2-8		
Sprangletop (<i>Leptochloa filiformis</i>)	2-6		
Volunteer Barley (<i>Hordeum vulgare</i>)	2-6		
Volunteer Oats (<i>Avena sativa</i>)	2-6		
Volunteer Rye (<i>Secale cereale</i>)	2-6		
Volunteer Wheat (<i>Triticum aestivum</i>) (non-QPE tolerant)	2-6		
Wild Oat (<i>Avena fatua</i>)	2-6		
Witchgrass (<i>Panicum capillare</i>)	2-6		
Rye, Feral or Cereal (<i>Secale cereale</i>)	1-4	10 – 16 (0.068 – 0.11 lbs AI)	12-16 (0.0825 - 0.11 lbs AI)
Barnyardgrass (<i>Echinochloa crus-galli</i>)	2-6	8 – 16 (0.055 – 0.11 lbs AI)	†
Crabgrass, Large (<i>Digitaria sanguinalis</i>)	2-6 ‡		10 – 16 (0.068 – 0.11 lbs AI)
Crabgrass, Smooth (<i>Digitaria ischaemum</i>)	2-6 ‡		
Junglerice (<i>Echinochloa colonum</i>)	2-6		†
Texas Panicum (<i>Panicum texanum</i>) ††	2-4	8 – 10 (0.055 – 0.068 lbs AI)	†
Red Rice (<i>Oryza sativa</i>)	1-4		†
Woolly Cupgrass (<i>Eriochloa villosa</i>)	2-4 §		†
Broadleaf Signalgrass (<i>Brachiaria platyphylla</i>)	2-6	10 (0.068 lbs AI)	†
Jointed goatgrass (<i>Aegilops cylindrica</i>) †††	2-6	10 – 16 (0.068 – 0.11 lbs AI)	12 - 16 (0.0825 -0.11 lbs AI)

WEEDS CONTROLLED AND RATE SELECTION (based on US EPA approved label dated 2/26/2026)
(Refer to the "Winter Wheat Restrictions" (below) for application limitations for single and split applications)

Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to quizalofop-P-ethyl or other Group 1 herbicides have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of Aggressor® AX specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed.

	Size at Application (height inches)	Aggressor® AX Applied Alone fl. oz. product / A (lbs AI / A)	Aggressor® AX Tank Mixed with Broadleaf Herbicide* fl.oz. product / A (lbs AI / A)
Italian ryegrass (<i>Lolium multiflorum</i>) Rescue grass (<i>Bromus catharticus</i>) Downy brome (<i>Bromus tectorum</i>) Windgrass (<i>Bromus mollis</i>) Japanese brome (<i>Bromus arvensis</i>)	2-6	8 – 16 (0.055 – 0.11 lbs AI)	
PERENNIAL GRASSES**			
Wirestem Muhly (<i>Muhlenbergia frondosa</i>)	4-8	8 – 10 (0.055 – 0.068 lbs AI)	†
Bermudagrass (<i>Cynodon dactylon</i>)	3" tall (or 6" runners)	10 – 16 (0.068 – 0.11 lbs AI)	†
Johnsongrass, Rhizome (<i>Sorghum halepense</i>)	10-24		10 (0.068 lbs AI)
Quackgrass (<i>Agropyron repens</i>)	6-10		†

* See "Applications With Broadleaf Herbicides"

** For annual and perennial grasses, up to 16 fl. oz. Aggressor® AX (0.11 lbs AI) / Acre may be applied, based on local recommendations. Under arid conditions apply at the higher rate.

*** Control includes Roundup Ready (glyphosate resistant), Liberty Link, and IMI-Corn varieties. Apply 8 fl. oz. Aggressor® AX (0.055 lb AI) for control of volunteer corn.

† May not be controlled adequately using a tank mix with broadleaf herbicides. For best results, alternate application of Aggressor® AX with a broadleaf herbicide, ensuring that Aggressor® AX is applied either 24 hours before or 7 days after the broadleaf herbicide.

‡ Length of lateral growth

§ Size in height or diameter, whichever is more restrictive. Applications to plants with more than three tillers may result in unsatisfactory control.

†† In Texas and other areas of the arid west, apply at 10 fluid ounces per acre for control of Texas panicum, use of lower rates may result in unsatisfactory control.

††† Use COC's at 1% v/v (1 gallon per 100 gallons spray solution) when targeting control of Jointed goatgrass.

AD022626