

FIFRA Sec. 2(ee) Recommendation

Priaxor[®]

Xemium[®] Brand Fungicide

Reduced use rate for infurrow applications to aid in the control of soilborne Rhizoctonia in potatoes in Colorado, Delaware, Florida, Idaho, Maine, Maryland, Massachusetts, Michigan, Minnesota, Montana, New Jersey, North Carolina, North Dakota, Oregon, Pennsylvania, Virginia, Washington, and Wisconsin

This Section 2(ee) Recommendation expires on December 31, 2028.

EPA Reg. No. 7969-311

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

All applicable directions, restrictions, precautions, and Conditions of Sale and Warranty on the EPA registered label are to be followed.

Information contained in this Section 2(ee) Recommendation is not intended to replace or amend any product labeling. Always read and follow all label directions when using any pesticide alone or in tank mix combinations.

The user must have this recommendation in their possession at the time of use.

Directions For Use

Use 0.46 to 0.55 fl oz of **Priaxor[®] Xemium[®] brand fungicide** per 1000 row feet to aid in control of soilborne Rhizoctonia in potatoes. Refer to the chart below to determine the product rate per acre.

Product Rate per 1000 row feet	Product Rate per Acre (fl ozs product)			
(fl oz product)	34-inch rows or less	36-inch rows	38-inch rows	40-inch rows
0.46	7.1	6.7	6.3	6.0
0.55	See footnote ¹	8.0	7.6	7.2
¹ For 34-inch rows or less, use a maximum of 0.46 fl oz product per 1000 row feet.				

Priaxor and Xemium are registered trademarks of BASF.

© 2025 BASF Agricultural Solutions US LLC
All rights reserved.

007969-00311.20251010.**NVA 2025-04-0372-0261**
Supersedes: NVA 2022-04-0372-0247


We create chemistry