

Product Bulletin

SULFOXAFLOR GROUP 4C INSECTICIDE



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Transform[®] WG

EPA Reg. No. 62719-625

Special 2(ee) Recommendation[†]
(FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY)

**For Management of Cotton Jassid (Two-Spot Cotton Leafhopper)
in Cotton and Okra**

This 2(ee) recommendation expires on December 31, 2029.

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Transform WG insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Transform WG according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Transform WG.

Directions for Use

Refer to product label for Transform WG for General Use Precautions, Mixing Directions, and Application Directions.

Pests and Application Rates

Pests	Transform WG (oz/acre)
Cotton Jassid (Two-Spot Cotton Leafhopper)	1.50 – 2.25 (0.047 – 0.071 lb. ai/acre)

Advisory Pollinator Statement: Notifying known beekeepers within 1 mile of the treatment area 48 hours before the product is applied will allow them to take additional steps to protect their bees. Also, limiting application to times when managed bees and native pollinators are least active, e.g. 2 hours prior to sunset or when the temperature is below 50° F at the site of application will minimize risk to bees. The RT₂₅ for this product is less than or equal to 3 hours.

Application Timing: Treat in accordance with local economic thresholds. Consult your company representative, cooperative extension service, certified crop advisor or state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: 1.50 oz/A. – 2.25 oz/A. (0.047 lb. ai/acre – 0.071 lb. ai/acre)

Restrictions:

- **Preharvest Interval:** Do not apply within 14 days of harvest.
- **Minimum Treatment Interval:** Do not make applications less than 5 days apart.

- Do not make more than four applications per acre per year.
- Do not make more than two consecutive applications per crop.
- Do not apply more than a total of 8.5 oz of Transform WG (0.266 lb ai of sulfoxaflor) per acre per year.

TM® Trademarks of Corteva Agriscience and its affiliated companies

R396-287

Approved: 03/12/2026

Initial printing.