

Supplemental Label



ETHABOXAM	GROUP	22	FUNGICIDE
-----------	-------	----	-----------



Elumin Fungicide

EPA Reg. No. 59639-211

**V-10208 4 SC FUNGICIDE FOR USE ON LEAF PETIOLE VEGETABLE
(CROP SUBGROUP 22B)
(Greenhouse use only)**

This supplemental label expires on **August 9, 2027** and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR V-10208 4 SC FUNGICIDE BEFORE APPLYING. USE OF V-10208 4 SC FUNGICIDE ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR V-10208 4 SC FUNGICIDE.

SPECIFIC REQUIREMENTS – All Crops

When to Apply: See crop-specific requirements. When applying as a foliar spray, apply in sufficient water to obtain thorough coverage. Chemigation is not advised when the primary disease is downy mildew or other foliar diseases.

Resistance Management: V-10208 4 SC Fungicide must be used as part of an Integrated Pest Management (IPM) program.

Restrictions and Limitations

- The maximum V-10208 4 SC Fungicide application rate is 8 fl oz per acre (0.25 lb ai/A) per application and 16 fl oz per acre (0.5 lb ai/A) per year.
- If noted in the CROP SPECIFIC REQUIREMENTS section, do not apply sequentially before rotating to a non-Group 22 fungicide that is registered for the same use and that is effective against the pathogens of concern.
- Do not make more than 2 applications of V-10208 4 SC Fungicide per acre per year.
- Do not apply this product when weather conditions favor spray drift from treated areas.
- When applying by air, observe drift management restrictions and precautions listed in the AERIAL APPLICATION section. Best control is achieved through ground application.

CROP SPECIFIC REQUIREMENTS

Leaf petiole vegetable (Crop subgroup 22B) Cardoon; celery; celery, Chinese; fuki; rhubarb; udo; zuiki; cultivars, varieties, and hybrids of these commodities. (Greenhouse use only)				
Diseases	Product Application Rate	Application Volume	Specific Use Instructions	Use Restrictions
	fl oz/A	gal/A		
Pythium Root Rot (<i>Pythium spp.</i>)	8 (0.25 lb ai/A)	Ground: 20 to 100	Pythium Root Rot For Pythium root rot control in the greenhouse, make 2 soil drench applications to seedlings (that have at least two true leaves) at an interval of 14 days when conditions are favorable for disease development and prior to disease onset. Following the second application, after V-10208 4 SC Fungicide has dried on the plant, transplant celery into the field.	• Do not apply V-10208 4 SC Fungicide at intervals of less than 14 days. • Do not make more than 2 applications per year. • Do not apply more than 8 fl oz per acre (0.25 lb ai/A) per application. • Do not apply more than 16 fl oz (0.5 lb ai) of V-10208 4 SC Fungicide per acre per year. • Preharvest Interval (PHI): N/A

PLEASE CONTACT VALENT U.S.A. LLC AT 800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

©2025 Valent U.S.A. LLC

Manufactured for:
Valent U.S.A. LLC
 P.O. Box 5075
 San Ramon CA 94583

Made in U.S.A.

2025-ELUMIN-0011-LeafPetiole