

SPINOSAD	GROUP	5	INSECTICIDE
----------	-------	---	-------------

FIFRA Section 24(c) Special Local Need Label



Corteva Agriscience LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA 1-800-258-3033

Gf-120[®] NF Naturalyte[®] Fruit Fly Bait

EPA Reg. No. 62719-498

EPA SLN No. NY-180004

For Distribution and Use Only in the State of New York

For Use Only in Regulated Areas Under Quarantine

For use only by or under the direct supervision of the state/federal cooperative program officials involved in the control of exotic fruit fly members of the family *Tephritidae*, such as the European Cherry Fruit Fly

For eradication/suppression efforts in the 200-meter zone around positive detections.

Expiration date: This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2030.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read this SLN and the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this FIFRA 24(c) Special Local Need labeling is subject to all use precautions and limitations imposed by the label affixed to the container for this product.
- For ground application only. Aerial application is prohibited.
- This product is toxic to bees exposed to direct treatment. Avoid applying this product or allowing it to drift to blooming crops or weeds if bees are foraging within the treatment area.
- The state/federal cooperative program shall ensure that all beekeepers that operate hives in the treatment area are notified about the application at least 48 hours in advance of a planned treatment.

Refer to product label for GF-120 NF for General Information, Mixing and Application directions.

Directions for Use

GF-120[®] NF Naturalyte[®] Fruit Fly Bait is a bait concentrate that should be diluted with water prior to application.

- Once diluted, GF-120 NF Naturalyte Fruit Fly Bait should be used within 24 hours.
- A large droplet spray of 4000 to 6000 um (4 to 6 mm) is recommended to optimize length of bait attract ability.
- Direct spray application to bottoms of leaves and leaves inside the foliage canopy to reduce direct exposure to sun and rain. This product resists wash off, but will lose effectiveness if exposed to rain and overhead irrigation.
- Begin applications as soon as monitoring traps indicate flies are present. Repeat applications every 7-14 days are recommended, however spot spray applications may be made as

frequently as every two days when program monitoring determines a need for more applications. Spot spray applications may be shortened to daily after unanticipated rainfall events.

Amount of Undiluted GF-120 NF Naturalyte Fruit Fly Bait			
Broadcast Application		Spot Spray of Individual Plants	
fl oz/acre	liters/hectare	fl oz/acre	ml/tree
10 – 20	0.12 – 0.24	1 – 3	30 - 90

- Choose rate based on pest pressure and amount of foliage needed to cover.
- For larger trees where the 3 fl oz rate is insufficient for adequate coverage, up to 21 fl oz/tree may be applied. The higher rate may only be used for limited spot treatment in regulated quarantine areas: **Do Not** exceed 2,300 fl oz/acre per application.

Rates of Final Diluted Spray Solution per Acre					
Dilution Ratio GF-120 NF Naturalyte : Water	10 fl oz/acre	12 fl oz/acre	15 fl oz/acre	18 fl oz/acre	20 fl oz/acre
Dilution Ratio of 1 : 1.5	10 + 15	12 + 18	15 + 22.5	18 + 27	20 + 30
Amount of Final Solution	25	30	37.5	45	50

™®Trademarks of Corteva Agriscience and its affiliated companies

R188-007

Approved: 02/04/2026

Initial Printing/Replaces R188-006