



SEED SAFENER

Viludo™



SEED TREATMENT FOR SAFENING RICE AGAINST HERBICIDE INJURY

For use by commercial seed treaters only

SAFENING INGREDIENT:

FENCLORIM (CAS No. 3740-92-9) 42.0%*

OTHER INGREDIENTS: 58.0%

TOTAL: 100.0%

*Contains 4.2 lbs. active per gallon or 500 grams per liter.

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See front panel for First Aid Instructions and Booklet for complete Precautionary Statements and Directions for Use.

FIRST AID	
IF ON SKIN:	• Remove contaminated clothing. Wash with soap and plenty of water. Call a Poison Control Center or doctor.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move to fresh air. Call a Poison Control Center or doctor.
IF SWALLOWED:	• Call a POISON CONTROL CENTER or doctor immediately. Do not induce vomiting without medical advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE CALL ROCKY MOUNTAIN POISON AND DRUG SAFETY: 1-866-673-6671. FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire, exposure or accident) CALL CHEMTREC: 1-800-424-9300 or +1-703-527-3887.	
For Product Use Information Call 1-800-438-6071	

	Warning Can cause moderate eye irritation. Wash face and hands thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection. If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present, and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.
--	---

Read SDS before use.

Produced For: UPL NA Inc.
PO Box 12219, Research Triangle Park
NC 27709 U.S.A. • 1-800-438-6071



PRECAUTIONARY STATEMENTS

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Socks and shoes,
- Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, or Viton,
- Protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. Do not contaminate water when disposing of equipment washwater or rinsate.

USE INFORMATION

VILUDO™ ST safens rice seed germination and emergence by specifically reducing seed and seedling injury caused by herbicide products containing the active ingredient clomazone. Without treatment with **VILUDO**, seedlings may suffer injury from the referenced herbicides. Using labeled herbicides that have expired, been improperly stored, or demonstrate degraded formulation quality may result in insufficient crop safety, even when used in combination with **VILUDO**. **VILUDO** may cause a slight delay of seedling emergence. The use of a seed treatment containing gibberellic acid in addition to **VILUDO** has shown the ability to mitigate this delay in emergence. Plants rapidly recover with no impact to final yield.

DIRECTIONS FOR USE – RICE (Dry-Seeded Rice Only)

Apply **VILUDO** uniformly to high quality rice seed at the following application rate:

Crop	Quantity of Seed (lbs)	Amount of VILUDO	Grams of ai/ 100 kg of seed
Rice	100	7.67 fl oz	250

- **DO NOT** plant rice treated with **VILUDO** in fields where crustacea (including crawfish) production or fish farming occurs.
- **DO NOT** plant rice treated with **VILUDO** in fields where water from a treated field will be directed to fields or ponds where crustacea (including crawfish) production or fish farming will occur.

MIXING INSTRUCTIONS

CONSTANT AGITATION IS REQUIRED whenever a custom blend of **VILUDO** is in the mix-tank. Application equipment must be cleaned out immediately after treatment.

COMPATIBILITY WITH FUNGICIDES AND INSECTICIDES

VILUDO is compatible with most StartUP™ and RANCONA® branded fungicide and insecticide seed treatment products labeled for use on rice. However, conducting appropriate compatibility testing with all seed treatment products is warranted when using **VILUDO** unless compatibility has been verified previously. Always conduct a standard compatibility test before mixing commercial quantities of this product with another product, to assure that there is acceptable compatibility. If incompatibility is found, a compatibility agent must be used, and the drying time should be extended before bagging the treated seed. Read and carefully follow the label directions of each combination product. When using combinations of products, the most restrictive of label and precautions must be followed.

GERMINATION, VARIETAL DIFFERENCES, AND QUALITY CONTROL

To assure quality standards, treat a sample of each seed lot with **VILUDO** and check for germination and safening protection. Germination of rice treated with **VILUDO** may be affected by genetic differences among varieties (endosperm lines), seed vigor, adverse growing conditions, or seed lots with seed coat damage. For best results, treat a sample of each commercial seed lot with **VILUDO** and check for germination. Herbicide protection can be checked by running a standard bioassay procedure in either a laboratory or greenhouse. Only those seed lots meeting acceptable germination standards from these tests should be treated in commercial quantities.

GERMINATION AND ENVIRONMENTAL DIFFERENCES

Optimum germination and emergence of rice occurs when the soil temperature at 4-inch depth is at least 60°F for rice planted at 0.25 to 1.5-inch depth. Planting prior to a heavy rainfall event can create adverse germination and stand establishment conditions, which can lead to potential stand reduction. Planting rice too early or planting into cold/wet soils will delay emergence and subject the seed to additional stress. When cool soils persist prior to planting, we recommend that you use only those varieties that exhibit acceptable cold tolerance, to minimize possible germination reduction.

CARRY OVER SEED

Rice seed treated with **VILUDO** at the labeled rate may be carried over and planted the year following initial seed treatment provided that it's germination is acceptable.

TREATED SEED LABELING

Seed treated with **VILUDO** may be tagged and sold only for planting, provided the following statement appears on the tag or container:

"WARNING: TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES.

- **DO NOT** plant rice treated with **VILUDO** in fields where crustacea (including crawfish) production or fish farming occurs.

- **DO NOT** plant rice treated with **VILUDO** in fields where water from a treated field will be directed to fields or ponds where crustacea (including crawfish) production or fish farming will occur."

The labels of registered seed treatment fungicides and insecticides may contain specific wording for the labeling of treated seed, based on the requirements of federal seed laws and regulations. **VILUDO** is not in the same regulatory category as fungicides and insecticides, and there are no federal requirements for labeling safened seed. However, because some state seed laws may be more restrictive than the Federal law, the above WARNING statement should be used with **VILUDO** treated seeds.

UPL NA Inc. recommends that the seed laws be reviewed for each state in which safened seed is to be sold, and that clarification be sought from a qualified attorney or representative of the State with respect to seed safeners and labeling requirements, because some states may have reduced seed labeling requirements for safened seed under their seed laws.

IMPORTANT

Substances listed below are those identified as hazardous chemicals under the criteria of the OSHA Hazardous Communication Standard (29 CFR 1910.1200):

FENCLORIM: Pyrimidine, 4,6-dichloro-2-phenyl-; 4,6-dichloro-2-phenylpyrimidine (CAS Reg. No. 3740-92-9)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PRODUCT STORAGE:

Store in the original container only. After partial use, close the container tightly. Store in a secure place that is temperate, dry and out of direct sunlight. DO NOT STORE ABOVE 85°F.

PRODUCT DISPOSAL:

Wastes resulting from the use of this product that cannot be used according to label instructions or chemically reprocessed must be disposed of in an approved waste disposal facility, in accordance with applicable federal, state and local procedures.

CONTAINER HANDLING:

Nonrefillable container:

DO NOT reuse or refill this container. Triple rinse (or equivalent), adding rinsate to spray tank. Offer rinsed containers for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT INFORMATION READ BEFORE USING PRODUCT

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of UPL NA Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of UPL NA Inc. and Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold UPL NA Inc. and Seller harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.

To the extent consistent with applicable law, UPL NA Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UPL NA INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UPL NA INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

UPL NA Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of UPL NA Inc.

UPL, the UPL logo, VILUDO, STARTUP and RANCONA are trademarks of a UPL Corporation Limited Group Company.

©2025 UPL Corporation Limited Group Company

2917298-A (09/25)

REV080725-13205-091025

NOTES



SEED TREATMENT FOR SAFENING RICE AGAINST HERBICIDE INJURY

For use by commercial seed treaters only

SAFENING INGREDIENT:

FENCLOIRIM (CAS No. 3740-92-9) 42.0%*

OTHER INGREDIENTS: 58.0%

TOTAL: 100.0%

*Contains 4.2 lbs. active per gallon or 500 grams per liter.

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See front panel for First Aid Instructions and Booklet for complete Precautionary Statements and Directions for Use.

FIRST AID

IF ON SKIN:	• Remove contaminated clothing. Wash with soap and plenty of water. Call a Poison Control Center or doctor.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move to fresh air. Call a Poison Control Center or doctor.
IF SWALLOWED:	• Call a POISON CONTROL CENTER or doctor immediately. Do not induce vomiting without medical advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE CALL ROCKY MOUNTAIN POISON AND DRUG SAFETY: 1-866-673-6671.

FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire, exposure or accident) CALL CHEMTREC: 1-800-424-9300 or +1-703-527-3887.

For Product Use Information Call 1-800-438-6071



Warning

Can cause moderate eye irritation.

Wash face and hands thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection.

If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present, and easy to do. Continue rinsing.

If eye irritation persists: Get medical advice/attention.

Read SDS before use.

PRECAUTIONARY STATEMENTS

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or incorporate spilled treated seed. Do not contaminate water when disposing of equipment washwater or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PRODUCT STORAGE:

Store in the original container only. After partial use, close the container tightly. Store in a secure place that is temperate, dry and out of direct sunlight. DO NOT STORE ABOVE 85°F.

PRODUCT DISPOSAL:

Wastes resulting from the use of this product that cannot be used according to label instructions or chemically reprocessed must be disposed of in an approved waste disposal facility, in accordance with applicable federal, state and local procedures.

CONTAINER HANDLING:

Nonrefillable container:

DO NOT reuse or refill this container. Triple rinse (or equivalent), adding rinsate to spray tank. Offer rinsed containers for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Produced For: UPL NA Inc. • PO Box 12219, Research Triangle Park, NC 27709 U.S.A. • 1-800-438-6071



Net Contents: 2.5 Gallons* □ 15 Gallons (FG# 6022177)

*unless alternate checked