

SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

MANDIPROPAMID	GROUP	40	FUNGICIDE
---------------	-------	----	-----------

Revus®

Fungicide

This supplemental label expires on 03/30/2027 and must not be used or distributed after this date.

Active Ingredient:

Mandipropamid*	23.3%
----------------	-------

<i>Other Ingredients:</i>	76.7%
---------------------------	-------

Total:	100.0%
--------	--------

* CAS No. 374726-62-2

Revus is formulated as a suspension concentrate and contains 2.08 lb of active ingredient per gallon or 250 grams per liter of active ingredient.

Contains 1,2-benzisothiazolin-3-one at 0.017% as a preservative.

KEEP OUT OF REACH OF CHILDREN.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 100-1254

All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Revus as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

In-Furrow Application

- Apply Revus as an in-furrow spray in 5-15 gallons water per acre at planting.
- Mount the spray nozzle so the spray is directed into the furrow just before the seed are covered.

Table 1: Soil application rates for Revus /1,000 feet of row, based on plant row spacing

Revus Conversion Chart for Drip (Trickle) Chemigation, Continuous Transplant Water, and Direct/Banded/In-Furrow Application							
Corresponding field rate (fl oz/A)	Rate in fl oz product/1,000 row ft based on planted row spacing of:						
	30"	34"	36"	48"	60"	72"	84"
8.0	0.46	0.52	0.55	0.73	0.92	1.1	1.28
16.0	0.92	1.04	1.1	1.47	1.83	2.2	2.57

Transplant Water Application

- Transplants should be adequately watered before transplanting. Ensure transplant water volume is sufficient to thoroughly wet the root zone.
- See **Table 1** for continuous-stream transplanters. Ensure 4-8 fl oz transplant water/transplant depending on sandy (4 fl oz) vs silty soil (6-8 fl oz).
- For water-wheel transplanters, use the plant population to determine the rate of product per plant.

Example:

$$\frac{16 \text{ fl oz product}}{\text{acre}} \times \frac{\text{acre}}{4,356 \text{ plants}} = \frac{0.0036 \text{ fl oz product}}{\text{plant}}$$

Surface Band or Directed Application

- Apply in a 6- to 12-inch band. See **Table 1** for rates.
- For ground broadcast application, apply in a minimum of 20 gallons of water per acre, unless specified otherwise. If rain is not expected within 24 hours after application, mechanically incorporate (before planting) or sprinkler irrigate (after planting) with 0.25-0.5 inch of water to move Revus into target disease zone.

SPECIFIC DIRECTIONS FOR USE

Tree Nuts Crop Group 14-12, Non-bearing

Crops (including cultivars, varieties, and/or hybrids of these)			
African nut-tree	Coconut	Monkey puzzle nut	
Almond	Coquito nut	Okari nut	
Beechnut	Dika nut	Pachira nut	
Brazil nut	Ginkgo	Peach palm nut	
Brazilian Pine	Guiana Chestnut	Pecan	
Bunya	Hazelnut (Filbert)	Pequi Nut	
Bur Oak	Heartnut	Pili Nut	
Butternut	Hickory nut	Pine nut	
Cajou nut	Japanese horse-chestnut	Pistachio	
Candlenut	Macadamia Nut	Sapucaia nut	
Cashew	Mongongo nut	Tropical almond	
Chestnut	Monkey pot	Walnut, black	
Chinquapin		Walnut, English	
		Yellowhorn	
Target Diseases	Rate fl oz/100 gal (lb ai/A)	Application Timing	Use Directions
Phytophthora root rot (<i>Phytophthora</i> spp.)	16.0 fl oz/A (0.26)	Apply 2 times a year, with a minimum interval of 3 months, during root flushes to protect the new root growth and the crown of the plant.	Apply as a drench, soil directed spray, or through irrigation water (micro-sprinkler, or drip). For effective disease control, ensure that the product solution thoroughly wets the target root zone.
	0.36 fl oz per 1,000 sq ft (0.26)		
	Drench in potted container 16.0 fl oz/100 gal (0.26)	Apply to the container 2 times a year, with a minimum interval of 3 months, during root flushes to protect the new root growth and the crown of the plant.	Apply as a drench (see Table below for volume) to the container. Use a sufficient volume of water to wet the soil in the container. In order to maintain protection in the field, drench pot prior to transplanting.

Volume of Revus Drench by Container Size

Pot Diameter (inches)	Drench Solution (fl oz) per Container	Maximum Number of Potted Trees that can be Treated per Acre
4	4	3,200
5	5	2,560
6	6	2,133
8	10	1,280
10	20	640
12	30	426

Amount of Revus to Achieve Desired Volume of Drench Solution

Rate/100 gal	5 gal	10 gal	25 gal	50 gal	75 gal	100 gal
16 fl oz	0.8 fl oz	1.6 fl oz	4 fl oz	8 fl oz	12 fl oz	16 fl oz

- Drench applications to potted containers cannot exceed the maximum number of treated trees per acre listed in the table.

Resistance Management:

- Refer to **Section 3.1**.

Integrated Pest Management

- Refer to **Section 3.2**.

USE RESTRICTIONS

- Refer to **Section 6.1** for additional product use restrictions.
- Maximum Single Application Rate: DO NOT** exceed maximum rate listed on table for this application type.
- Maximum Number of Applications per Year: DO NOT** make more than 2 applications at the maximum rate per year.
- Minimum Application Interval: 90 days**
- Maximum Annual Rate:** 32 fl oz/A/year (0.52 lb ai/A/year)
 - DO NOT** exceed 0.52 lb ai/A/year of mandipropamid-containing products.
- DO NOT** apply by air.
- Preharvest Interval (PHI): 365 days**

Tuberous and Corm Vegetables, Crop Subgroup 1C

Crops (including cultivars, varieties, and/or hybrids of these)			
Arracacha		Cassava, sweet	Potato
Arrowroot		Chayote, root	Sweet potato
Artichoke, Chinese		Chufa	Tanier
Artichoke, Jerusalem		Dasheen (Taro)	Turmeric
Canna, edible		Ginger	Yam bean
Cassava, bitter		Leren	Yam, true
Target Diseases	Rate fl oz/A (lb ai/A)	Application Timing	Use Directions
Suppression Only: Pink rot <i>(Phytophthora erythroseptica)</i>	8.0 (0.13)	At-Planting Soil Application: Apply in-furrow at planting.	Soil Application: <i>Provided mandipropamid was not used as a potato seed treatment,</i> apply as a 6- to 8-inch band directly over the seed pieces in the furrow and then close the furrows. Make application in 3-15 gal/A. Refer to Section 4.1.1 for additional instructions on in-furrow application. You may need to follow this in-furrow application of Revus with a foliar application of a different product such as a mefenoxam-containing (e.g., Ridomil Gold® Bravo® SC (EPA Reg. No. 100-1221)) Refer to label for specific use instructions at tuber initiation: • When conditions are conducive for disease development. • When the variety is susceptible or moderately susceptible to pink rot or • In areas with a long growing season.
Late blight <i>(Phytophthora infestans)</i>	8.0 (0.13)	Foliar Application: Begin applications prior to disease development and continue throughout the season on a 7-10 day interval. <i>If mandipropamid was used as a potato seed treatment or at planting,</i> make up to three post-planting foliar applications, otherwise follow use restrictions below.	Apply by ground, air, or chemigation. The addition of a spreading/penetrating type adjuvant such as a non-ionic surfactant or crop oil concentrate or blend is advised when applying by ground or air. Use the shorter interval and/or higher rates under high pressure or when conditions are conducive to disease.
Resistance Management: • Refer to Section 3.1. • Make no more than 2 consecutive applications before switching to another effective non-Group 40 fungicide.			

Integrated Pest Management Refer to Section 3.2.
USE RESTRICTIONS
1) Refer to Section 6.1 for additional product use restrictions. 2) Maximum Single Application Rate: DO NOT exceed maximum rate listed on table for this application type. 3) Maximum Number of Applications per Year: DO NOT make more than 4 applications at the maximum rate per year regardless of application method. 4) Minimum Application Interval: 7 days 5) Maximum Annual Rate: Soil application: 8 fl oz/A/year (0.13 lb ai/A/year) Foliar: 32 fl oz/A/year (0.52 lb ai/A/year) a. DO NOT exceed 0.13 lb ai/A/year soil applied and DO NOT exceed a total of 0.52 lb ai/A/year of mandipropamid-containing products. 6) Mandipropamid may be used as a potato seed treatment or applied in soil, but NOT both within a crop cycle. 7) Preharvest Interval (PHI): 14 days

Revus®, Ridomil Gold®, and the Syngenta logo are trademarks of a Syngenta Group Company

Bravo®, is a trademark of Makhteshim Agan of North America, Inc. d/b/a ADAMA.

©2026 Syngenta