

RIMON[®] 0.83EC

INSECTICIDE

Insecticide for use on: Avocado*; Berries (Low-Growing); Brassica, head and stem (Group 5-16); Brassica, Leafy Greens (Subgroup 4-16B); Bushberries; Carrot*; Individual Legume Vegetable Commodities; Pears*; Pome Fruits; Potatoes/Sweet Potatoes; Stonefruits; Strawberry; Sweet Corn; Swiss Chard*; Turnip Greens*; Vegetable, cucurbit (Group 9); Vegetable, fruiting (Group 8-10); Kohlrabi; Tropical Fruits (Subgroup 24A)* including Lychee; Vegetable, foliage of legume, subgroup 7A, forage; and Vegetable, foliage of legume, subgroup 7A.

* Not registered for use in California.

ACTIVE INGREDIENT:

Novaluron: 1-[3-chloro-4-(1,1,2-trifluoro-2-trifluoromethoxyethoxy)phenyl]-3-(2,6-difluorobenzoyl)urea**	% BY WT.
	9.3%

OTHER INGREDIENTS:

	90.7%
--	-------

TOTAL:	100.0%
---------------	---------------

** Contains 0.83 lb. novaluron per gallon.

EPA Reg. No. 66222-35-70506

EPA Est. No. 037429-GA-002

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID - IF IN EYES: • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. **IF ON SKIN:** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. • **DO NOT** give anything by mouth to an unconscious person. **IF INHALED:** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

Emergency Assistance: Have the product container or label with you when calling a poison control center or doctor or going for treatment. For non-emergency general information on this pesticide product (including health concerns or pesticide incidents), you may call 1-877-250-9291, 24 hours per day, 7 days per week. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

In case of spills, fire, leaks or accidents call 1-800-535-5053.

For additional precautionary, handling and use statements, see inside of this booklet.

Net Contents: 2.5 Gallons

INSECTICIDE

Distributed by: UPL NA Inc. • 630 Freedom Business Center, Suite 402
King of Prussia, PA 19406 U.S.A. • 1-800-438-6071



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING: Causes substantial but temporary eye injury. **DO NOT** get in eyes or on clothing. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Chemical-resistant gloves such as barrier laminate, butyl rubber $\geq$ 14 mils, nitrile rubber $\geq$ 14 mils, neoprene rubber $\geq$ 14 mils, or Viton $\geq$ 14 mils;
- Shoes plus socks; and
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater and estuarine/marine invertebrates. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment wash waters or rinsate. This product may impact surface water quality due to runoff or rain water. This is especially true for poorly draining soils and soils with shallow ground water.

Pollinator Advisory: Because of its mode of action as an insect growth regulator, and since it is not systemic, RIMON® 0.83EC Insecticide has the potential to impact larval bees (i.e., brood). In order to minimize the possibility of effects to honeybee brood, **DO NOT** use RIMON 0.83EC Insecticide on blooming crops when bees are actively foraging.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff for several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of novaluron from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Alternatively, for products with water-in requirements, avoid watering to the point of runoff.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS:

- Carefully read this product label for crop specific instructions and precautions, as failure to do so may result in crop injury.
- RIMON 0.83EC Insecticide has demonstrated some phytotoxic effects to new, expanding leaves, when mixed with products that are formulated as emulsifiable concentrates, systemic in nature, and/or intended to improve plant uptake, e.g., foliar nutrients/amendments, and/or petroleum/plant oil-based products.
- Use low rates of non-ionic, silicone, and other non-oil and non-penetrating adjuvants and/or surfactants known to be safe on listed crops.
- Carefully read the adjuvant and/or surfactant label to determine the presence of oil and/or penetrant activity before use; or consult the adjuvant and/or surfactant manufacturer.
- When an adjuvant is to be used with this product, the manufacturer recommends the use of a Council of Producers & Distributors of Agrotechnology certified adjuvant.
- Apply the spray solution with adjuvant and/or surfactant to a small area of the crop and wait 7 to 10 days and observe for signs of phytotoxicity before treating the entire field.

USE RESTRICTIONS:

- Apply this product outdoors only as specified the EPA approved label.
- **DO NOT** apply this product in a way that it will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- The use of novaluron on crops grown for food in greenhouses, except tomatoes and cucumbers, is prohibited.
- **DO NOT** allow RIMON 0.83EC to drift on grapes as leaf spotting may occur.
- **DO NOT** mix RIMON 0.83EC Insecticide with oil-based adjuvants or surfactants intended for plant absorption. Crop injury is typically exhibited as, but may not be limited to, chlorosis or mottling of new, expanding leaves.
- **DO NOT** allow to enter indoor or outdoor drains. (No permita la entrada a desagües internos o externos.)
- Follow proper disposal procedures on this label (*Siga las indicaciones del etiquetado para el desecho apropiado del producto.*)
- For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Water Protection Statements:

- **DO NOT** spray the product into fish pools, ponds, streams, or lakes.
- **DO NOT** apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- **DO NOT** allow the product to enter any drain during or after application.
- **DO NOT** apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
- **DO NOT** apply or irrigate to the point of runoff.

Rain Related Statements:

- **DO NOT** make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).
- Rainfall within 24 hours after application may cause unintended runoff of pesticide application.



BUFFER ZONES:

Vegetative Buffer Zones. Construct and maintain a minimum 25-foot vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; and estuarine/marine habitats). Only apply products containing nivalonuron onto fields where a well-maintained vegetative buffer strip of at least 25 feet exists between the field and down gradient aquatic habitat. For guidance, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21pp.

Buffer Zone for Ground Application (All Crops). **DO NOT** apply within 75 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds; and estuarine/marine habitats). All applications must include a 25-foot vegetative buffer strip within the buffer zone to decrease runoff.

Buffer Zone for Aerial Application. **DO NOT** apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds; and estuarine/marine habitats). All applications must include a 25-foot vegetative buffer strip within the buffer zone to decrease runoff.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over long-sleeved shirt and long pants;
- Chemical-resistant gloves such as barrier laminate, butyl rubber $\geq$ 14 mils, nitrile rubber $\geq$ 14 mils, neoprene rubber $\geq$ 14 mils, or Viton $\geq$ 14 mils;
- Shoes plus socks; and
- Protective eyewear

PRODUCT INFORMATION

RIMON 0.83EC Insecticide must be ingested and/or contacted by insects to be effective. Proper application techniques help ensure thorough spray coverage and correct dosage necessary to obtain optimum control. Apply at the required rates when insect populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area. Apply follow-up treatments of RIMON 0.83EC Insecticide per **DIRECTIONS FOR USE**, to keep pest populations within threshold limits. Scout fields regularly to determine optimum application timing based on pest levels and stages of growth.

The primary mode of action is by disrupting cuticle formation and deposition occurring when insects molt, resulting in their death. Due to this mode of action, RIMON 0.83EC Insecticide has no direct effect on adults.

NOTE: The compatibility of RIMON 0.83EC Insecticide with concurrent releases of insects for biocontrol of plant pests has not been established. When used as directed, RIMON 0.83EC Insecticide affects developing immature stages of insects by disrupting the molting process. Consequently, fully developed adult stages of pest and beneficial species are not affected.

Rotational Crops: Only registered crops may be rotated in a treated field within 30 days of the final application.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance-management, RIMON 0.83EC Insecticide contains a Group 15 insecticide. Any insect population may contain individuals naturally resistant to RIMON 0.83EC Insecticide and any other Group 15 insecticide. The resistant individuals may dominate the insect population if these groups of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of RIMON 0.83EC Insecticide or other Group 15 insecticide within a growing season, or among growing seasons, with different groups that control the same pests. Avoid application of more than the maximum seasonal use rate or the total number of consecutive sprays of RIMON 0.83EC Insecticide or other insecticides in the same group in a season.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. **DO NOT** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact UPL NA Inc. representatives at 1-800-438-6071.

APPLICATION PROCEDURES

MIXING PROCEDURES

Prepare solution concentrations in a clean, empty spray tank. Use clean spray filters. Add water to 1/2 level of tank. Add the appropriate amount of RIMON 0.83EC Insecticide to the tank and agitate to ensure proper mixture. Continue filling tank with water until desired dilution is achieved. Shake or re-agitate material in the sprayer before use if application is interrupted. Make up only the amount of application volume as required. Dispose of any unused spray material at the end of each day according to the instructions found in the **STORAGE AND DISPOSAL** section of this label.

For those crops where an adjuvant can be used, UPL NA Inc. suggests the use of a Chemical Producers and Distributors Association certified adjuvant.

SPRAY COVERAGE

All parts of the crop must receive uniform spray coverage or else desired result may not occur. Higher water volumes and increased spray pressure generally provide better coverage. Consult your local agricultural specialist for specific information on the best rates, timings, and spray volumes for your region.

ORCHARD APPLICATION

Make applications of RIMON 0.83EC insecticide by conventional orchard sprayers that are calibrated to deliver 50 to 400 gallons of carrier to the trees. Apply at a carrier volume that ensures complete coverage to trees. Operate spray equipment at proper ground speeds, adequate spray pressures and spray volumes that assure that the air volume within the tree canopy is completely replaced by the output from the air-blast sprayer resulting in proper coverage of the target crop. **DO NOT** use RIMON 0.83EC insecticide in alternate row middle application patterns since this method will result in off-timing application and poor performance.

GROUND APPLICATION

Apply required dosage by conventional ground sprayer equipment capable of delivering sufficient water to obtain thorough, uniform coverage of the target crop. Orient spray equipment boom and nozzles in a manner to minimize boom height to optimize coverage uniformity, maximize deposition and reduce spray drift. Drop nozzles may be required to obtain uniform coverage against certain pests that develop down in the canopy. Use a minimum of 10 gallons per acre in potatoes and vegetables. Higher gallonages will provide better coverage and performance. Use hollow cone, disc-cone hollow cone or twin jet fan nozzles suitable for insecticide spraying.

AERIAL APPLICATION

For aerial application apply in a total of 2 to 10 gallons per acre using a nozzle configuration that will provide a median droplet size of 200 - 300 microns. Use a minimum of 5 gallons of water per acre for potatoes. Higher gallonages will provide better coverage and performance. Adhere to the minimum safe application height - not greater than 12 feet above crop canopy. Boom length must be less than 75% of wing span and swath markers. Use flagging or GPS system during application. Make applications when wind speed is between 2 and 10 mph. **DO NOT** make applications when wind speed exceeds 10 mph. Under low humidity and high temperatures, adjust spray volume upward to compensate for evaporation of spray droplets.

APPLICATION THROUGH

IRRIGATION SYSTEMS - CHEMIGATION

RIMON 0.83EC insecticide may be applied through properly equipped chemigation systems for insect control in cranberries, potatoes and sweet corn. Apply this product only through sprinkler (including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move) irrigation systems. **DO NOT** apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

In order to calibrate the irrigation system and injector to apply the mixture, determine the following: 1) Calculate the number of acres irrigated by the system; 2) Set the irrigation rate and determine the number of minutes for the system to cover the intended treatment area; 3) Calculate the total gallons of the mixture needed to cover the desired acreage. Divide the total gallons of mixture needed by the number of minutes to cover the treated area. This value equals the gallons per minute that the injector must deliver. Convert the gallons per minute to ounces per minute. Calibrate the injector pump with the system in operation at the desired irrigation rate. Calibrate the injector pump at least twice before operation, and the system be monitored during operation.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers, or other experts.

DO NOT connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

If the chemigation system is connected to a public water supply, the following conditions must also be met:

- Public water systems mean a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from a point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Upon completion of insecticide application, remove scale, pesticide residues, and other foreign matter from the supply tank and entire injector system. Flush thoroughly with clean water.
- **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

SPRINKLER CHEMIGATION

For continuously moving systems, the mixture containing RIMON 0.83EC insecticide must be injected continuously and uniformly into the irrigation water line as the sprinkler is moving. If continuously moving irrigation equipment is used, apply in no more than 0.25 inch of water. For sprinkler systems that **DO NOT** move during operation, apply in no more than 0.25 inch of irrigation immediately before the end of the irrigation cycle.

Maintain continuous agitation of the pesticide supply tank for the duration of the application period.

To apply a pesticide using sprinkler chemigation, the chemigation system must meet the following specifications:

- The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

SPRAY DRIFT

DO NOT allow RIMON 0.83EC Insecticide to drift on grapes as leaf spotting may occur.

Mandatory Spray Drift Management

Airblast Applications:

- Spray must be directed into the canopy.
- **DO NOT** apply when wind speeds exceed 15 miles per hr. at the application site.
- User must turn off outward pointing nozzles at row ends and when spraying outer row.
- **DO NOT** apply during temperature inversions.

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for the fixed-wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- If the windspeed is 10 miles per hour or less, applicators must use 1/2 swath displacement upwind at the downwind edge of the field. When the windspeed is between 11 - 15 miles per hour, applicators must use 3/4 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications:

- **DO NOT** release spray at a height greater than 4 feet above the ground or crop canopy.
- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Boomless Ground Applications:

- Applicators must select nozzle and pressure that deliver medium or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT - Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications:

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

Take precautions to minimize spray drift.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Ultra Low Volume (ULV) application is not permitted.

CROPS

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
AVOCADO*	DO NOT apply within 1 day of harvest.	Lepidoptera Larva Including: Western Avocado Leafroller Avocado Looper Omnivorous Looper Orange Tortrix	19.3	Use a minimum spray volume of 100 GPA. Repeat applications as needed, but not less than 14 days apart.
		Avocado Use Restrictions: • DO NOT apply more than 57.9 fl. ozs. (0.38 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. * Not registered for use in California.		

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
BERRIES (CROP SUBGROUP 13-7H), INCLUDING: CRANBERRY, LINGONBERRY, MUNTRIES, PARTRIDGEBERRY, BEARBERRY, BILBERRY, LOWBUSH BLUEBERRY, CLOUDBERRY, EXCEPT STRAWBERRY (see separate direction for STRAWBERRY)	DO NOT apply within 1 day of harvest.	Blackheaded Fireworm Spotted Fireworm	12	1st generation larvae (May - June): Apply when the majority of overwintering eggs have hatched in early spring. 2nd generation larvae (late June - July): Apply at the first sign of oviposition through early egg hatch. Spray with a sufficient volume of water to ensure thorough coverage of fruit and leaf surfaces. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		Cranberry Blossomworm Cranberry Fruitworm Cranberry Spanworm Gypsy Moth Sparganothis Fruitworm	12	Apply when the majority of the target pest population is at egg hatch to early instars. Spray with a sufficient volume of water to ensure thorough coverage of fruit and leaf surfaces. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		Cranberry Fleabeetle Cranberry Tipworm Sap Beetle	12	Apply when adults appear and prior to egg hatch. For adult control, tank mix with an adulticide. Spray with a sufficient volume of water to ensure thorough coverage of fruit and leaf surfaces. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		Drosophila spp.* Including Spotted Wing Drosophila	12	Apply when adults appear. For adult control, tank mix with an adulticide. Spray with a sufficient volume of water to ensure thorough coverage of fruit and leaf surfaces. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		Berries (Crop Subgroup 13-7H) (Except Strawberries) Use Restrictions: • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. NOTE: For application to cranberries through irrigation systems, refer to the section entitled APPLICATION THROUGH IRRIGATION SYSTEMS - CHEMIGATION. * Not registered for use in California.		

Crops	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
BRASSICA, HEAD AND STEM (GROUP 5-16) INCLUDING: BROCCOLI, CHINESE BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAVALO BROCCOLO, CAULIFLOWER, CHINESE BROCCOLI (GAI LON), CHINESE CABBAGE (NAPA), CHINESE MUSTARD (GAI CHOY), AND KOHLRABI	DO NOT apply within 7 days of harvest.	Alfalfa Looper Armyworms Cabbage Loopers Cabbage Webworm Corn Earworm Cucumber Beetles Diamondback Moth Imported Cabbageworm Leafminers (Lepidopteran)	6 to 12	Apply when the majority of the population is at egg hatch to the second instar. Use higher rates and higher spray volumes when larvae are large, when target pests populations are 2X or more above state threshold level or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		Bagrada Bugs Leafminers (Dipteran) Lygus Bugs Stink Bugs Thrips Vegetable Weevil Whiteflies	12	
		Brassica, head and stem (Group 5-16) Use Restrictions: • DO NOT apply more than two applications against whiteflies or thrips per year. • DO NOT apply more than 24 fl. ozs. (0.16 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year.		

Crops	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
BRASSICA, LEAFY GREENS (SUBGROUP 4-16B) INCLUDING: ARUGULA; BROCCOLI, CHINESE; BROCCOLI RAAB; CABBAGE, ABYSSINIAN; CABBAGE, CHINESE, BOK CHOY; CABBAGE, SEAKALE; COLLARDS; CRESS, GARDEN; CRESS, UPLAND; HANOVER SALAD; KALE; MACA, LEAVES; MIZUNA; MUSTARD GREENS; RADISH, LEAVES; RAPE GREENS; ROCKET, WILD; SHEPHERD'S PURSE; CULTIVARS, VARIETIES, AND HYBRIDS OF THESE COMMODITIES.	DO NOT apply within 7 days of harvest.	Alfalfa Looper Armyworms Cabbage Loopers Cabbage Webworm Corn Earworm Cucumber Beetles Diamondback Moth Imported Cabbageworm Leafminers (Dipteran and Lepidopteran) Southern Cabbageworm	6 to 12	Apply when the majority of the population is at egg hatch to the second instar. Use higher rates and higher spray volumes when larvae are large, when target pests populations is 2X or more above state threshold level or foliage canopy is tall or dense. Repeat applications as needed to protect new growth, but not less than 7 days apart.
		Bagrada Bugs Leafminers (Dipteran) Lygus Bugs Stink Bugs Thrips Vegetable Weevil Whiteflies	12	
		Brassica, Leafy Greens (Subgroup 4-16B) Use Restrictions: • DO NOT apply more than 2 applications against whiteflies per year. • DO NOT apply more than 36 fl. ozs. per acre per calendar year (0.23 lb. a.i. per acre per calendar year). • DO NOT apply more than three applications per calendar year. • DO NOT apply to turnips harvested for the root. • DO NOT feed turnip tops to livestock.		

Crops	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
BUSHBERRIES (CROP SUBGROUP 13-07B), INCLUDING: BLUEBERRY (HIGHBUSH AND LOWBUSH), CURRANT, ELDERBERRY, GOOSEBERRY, AND HUCKLEBERRY	DO NOT apply within 8 days of harvest.	Blueberry Flea Beetle (Larvae) Blueberry Spanworm Cranberry Fruitworm Oblique-banded Leafroller Sparganothis Fruitworm	20 to 30	Make application when the majority of the population is at egg hatch to the second instar. Use higher rates and higher spray volumes when larvae are large, or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 10 days apart.
		Blueberry Maggot Fly Sap Beetle	20 to 30	Make application when adults are observed and prior to egg laying. Use higher rates and higher spray volumes when larvae are large, or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 10 days apart.
		Plum Curculio (Larvae)	20 to 30	Apply at pre-bloom to the newly expanded foliage and unopened blooms/buds. Adult females will deposit non-viable eggs after contact with, and feeding on, treated plants, providing control of eggs and larvae on early season harvested varieties. RIMON 0.83EC Insecticide will not control adult stages. A subsequent post-bloom spray using an adjuvant is recommended to achieve optimum control of all life stages. Use higher rates and higher spray volumes when larvae are large, or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 10 days apart.
		Drosophila spp.* Including Spotted Wing Drosophila	20 to 30	Apply when adults appear. For adult control, tank mix with an adjuvant. Use higher rates and higher spray volumes when larvae are large, or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 10 days apart.
		Bushberries Use Restrictions: - Some phytotoxic symptoms to foliage in the form of mottled chlorosis may be observed when RIMON 0.83EC Insecticide is applied to blueberries under conditions of high temperatures and/or drought stress, particularly during periods of new, tender shoot growth. Such phytotoxic symptoms will not occur on future growth, and will not affect fruiting or yields. Higher spray volumes and lower spray concentration will minimize the risk of transient phytotoxic symptoms on newly expanded foliage. DO NOT apply more than 90 fl. ozs. (0.58 lb. a.i.) of formulated product per acre per calendar year. DO NOT apply more than three applications per calendar year. * Not registered for use in California.		

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
CARROT*	DO NOT apply within 3 day of harvest.	Carrot Weevil Root Weevil White Grub Wireworm	12.3	Use a minimum spray volume of 20 GPA. Repeat applications, but not less than 7 days apart.
		Carrot Use Restrictions: • DO NOT apply more than 36.9 fl. ozs. (0.24 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. * Not registered for use in California.		

Crops	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
CUCURBIT VEGETABLES (CROP SUBGROUP 9B), INCLUDING BALSAM APPLE, BALSAM PEAR, CHAYOTE (FRUIT) CANTALOUPE, CUCUMBER, CHINESE CUCUMBER, GHERKIN (WEST INDIAN), EDIBLE GOURD, MELON, CITRON MELON, MUSKMELON, BITTERMELON, PUMPKIN, SQUASH, SUMMER SQUASH, WINTER SQUASH, WATERMELON AND CHINESE WAXGOURD	DO NOT apply within 1 day of harvest.	Armyworms Cucumber Beetles Leafminers (Lepidopteran) Loopers	9 to 12	Apply when the majority of the population is at egg hatch to the second instar. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 14 days apart. Apply sufficient spray volume to ensure full coverage of foliage, and flower buds. Use higher rates and higher spray volumes when larvae are large, or foliage canopy is tall or dense.
		Leafminer (Dipteran) Melonworm Pickleworm Sap Beetles Squash Bugs Thrips Whiteflies	12	Apply at the first sign of egg lay or egg hatch. For adult control, tank mix with an adulticide. DO NOT apply more than two applications against whiteflies or thrips per year. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 14 days apart. Apply sufficient spray volume to ensure full coverage of foliage, and flower buds. Use higher rates and higher spray volumes when larvae are large, or foliage canopy is tall or dense.
		Cucurbit Vegetables Use Restrictions: • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • The use of novaluron on crops grown for food in greenhouses, except tomatoes and cucumbers, is prohibited. • DO NOT apply more than three applications per calendar year.		

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
DRIED SHELLED BEAN* ADZUKI BEAN, DRY SEED AFRICAN YAM-BEAN, DRY SEED AMERICAN POTATO BEAN, DRY SEED ANDEAN LUPIN BEAN, DRY SEED ASPARAGUS BEAN, DRY SEED BLACK BEAN, DRY SEED BLACKEYED PEA, DRY SEED BLUE LUPIN BEAN, DRY SEED BROAD BEAN, DRY SEED CATJANG BEAN, DRY SEED CHINESE LONGBEAN, DRY SEED COWPEA, DRY SEED CRANBERRY BEAN, DRY SEED CROWDER PEA, DRY SEED DRY BEAN, DRY SEED FIELD BEAN, DRY SEED FRENCH BEAN, DRY SEED GARDEN BEAN, DRY SEED GOA BEAN, DRY SEED GRAIN LUPIN BEAN, DRY SEED GREAT NORTHERN BEAN, DRY SEED GREEN BEAN, DRY SEED GUAR BEAN, DRY SEED HORSE GRAM, DRY SEED JACKBEAN, DRY SEED KIDNEY BEAN, DRY SEED LABLAB BEAN, DRY SEED LIMA BEAN, DRY SEED MORAMA BEAN, DRY SEED MOTH BEAN, DRY SEED MUNG BEAN, DRY SEED NAVY BEAN, DRY SEED PINK BEAN, DRY SEED PINTO BEAN, DRY SEED RED BEAN, DRY SEED RICE BEAN, DRY SEED SCARLET RUNNER BEAN, DRY SEED SOUTHERN PEA, DRY SEED SWEET LUPIN BEAN, DRY SEED SWORD BEAN, DRY SEED TEPARY BEAN, DRY SEED URD BEAN, DRY SEED VEGETABLE SOYBEAN, DRY SEED VELVET BEAN, SEED, DRY SEED WHITE LUPIN BEAN, DRY SEED WHITE SWEET LUPIN BEAN, DRY SEED WINGED PEA, DRY SEED YARDLONG BEAN, DRY SEED YELLOW BEAN, DRY SEED YELLOW LUPIN BEAN, DRY SEED	DO NOT apply within 1 day of harvest.	Armyworms Loopers Webworms Bean Leaf Beetle Bean Plataspid Cucumber Beetle Mexican Bean Beetle Lygus Thrips Whiteflies	6 to 12 9 to 12 12 12	Apply when the majority of the target pest population is at egg hatch to early instars. Use higher rates and higher spray volumes when larvae are large or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart. Apply when the majority of the target pest population is at egg hatch to early instars. Use higher rates and higher spray volumes when larvae are large or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart. Apply when plant bugs appear and oviposition is initiated. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart. Apply when the majority of the target pest population is at egg hatch to early instars. DO NOT apply more than two applications against whiteflies or thrips per year. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
Dried Shelled Bean Use Restrictions: • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. * Not registered for use in California.				

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
DRIED SHELLED PEA* CHICKPEA, DRY SEED DRY PEA, DRY SEED FIELD PEA, DRY SEED GARDEN PEA, DRY SEED GRASS-PEA, DRY SEED GREEN PEA, DRY SEED LENTIL, DRY SEED PIGEON PEA, DRY SEED	DO NOT apply within 1 day of harvest.	Armyworms Loopers Webworms Bean Leaf Beetle Bean Plataspid Cucumber Beetle Mexican Bean Beetle Lygus Thrips Whiteflies Plant Bugs*, including Western Tarnished Plant Bug	12 in 10 gallons of water	Apply when the majority of the target pest population is at egg hatch to early instars. Use higher rates and higher spray volumes when larvae are large or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart. DO NOT apply more than two applications against whiteflies or thrips per year.
Dried Shelled Pea Use Restrictions: • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. * Not registered for use in California.				

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
EDIBLE PODDED BEAN* ASPARAGUS BEAN, EDIBLE PODDED CATJANG BEAN, EDIBLE PODDED CHINESE LONGBEAN, EDIBLE PODDED COWPEA, EDIBLE PODDED FRENCH BEAN, EDIBLE PODDED GARDEN BEAN, EDIBLE PODDED GOA BEAN, EDIBLE PODDED GREEN BEAN, EDIBLE PODDED GUAR BEAN, EDIBLE PODDED JACKBEAN, EDIBLE PODDED KIDNEY BEAN, EDIBLE PODDED LABLAB BEAN, EDIBLE PODDED MOTH BEAN, EDIBLE PODDED MUNG BEAN, EDIBLE PODDED NAVY BEAN, EDIBLE PODDED RICE BEAN, EDIBLE PODDED SCARLET RUNNER BEAN, EDIBLE PODDED SNAP BEAN, EDIBLE PODDED SWORD BEAN, EDIBLE PODDED URD BEAN, EDIBLE PODDED VEGETABLE SOYBEAN, EDIBLE PODDED VELVET BEAN, EDIBLE PODDED WAX BEAN, EDIBLE PODDED WINGED PEA, EDIBLE PODDED YARDLONG BEAN, EDIBLE PODDED	DO NOT apply within 1 day of harvest.	Armyworms Loopers Webworms Bean Leaf Beetle Bean Plataspid Cucumber Beetle Mexican Bean Beetle Lygus Thrips Whiteflies	12 in 30 gallons of water	Apply when the majority of the target pest population is at egg hatch to early instars. Use higher rates and higher spray volumes when larvae are large or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart. DO NOT apply more than two applications against whiteflies or thrips per year.
Edible Poddled Bean Use Restrictions: • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. * Not registered for use in California.				

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
EDIBLE PODDED PEA* CHICKPEA, EDIBLE PODDED DWARF PEA, EDIBLE PODDED EDIBLE PODDED PEA GRASS-PEA, EDIBLE PODDED GREEN PEA, EDIBLE PODDED LENTIL, EDIBLE PODDED PIGEON PEA, EDIBLE PODDED SNAP PEA, EDIBLE PODDED SNOW PEA, EDIBLE PODDED SUGAR SNAP PEA, EDIBLE PODDED	DO NOT apply within 1 day of harvest.	Armyworms Loopers Webworms Bean Leaf Beetle Bean Plataspid Cucumber Beetle Mexican Bean Beetle Lygus Thrips Whiteflies Plant Bugs*, including Western Tarnished Plant Bug, and Colorado Potato Beetle*	12 in 30 gallons of water	Apply when the majority of the target pest population is at egg hatch to early instars. Use higher rates and higher spray volumes when larvae are large or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
Edible Poddled Pea Use Restrictions: • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. * Not registered for use in California.				

Crops	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
FRUITING VEGETABLES (FIELD GROWN), INCLUDING TOMATOES (including BUSH, CURRANT and TREE TOMATOES), PEPPERS, EGGPLANTS (including AFRICAN, PEA and SCARLET EGGPLANTS), TOMATILLO, GROUNDCHERRY, PEPINO, OKRA, COCONA, GOJI BERRY, GARDEN HUCKLEBERRY, MARTYNIA, NARANJILLA, ROSELLE, and SUNBERRY	DO NOT apply within 1 day of harvest.	Armyworms Colorado Potato Beetle European Corn Borer Foliage Feeding Caterpillars Leafminers (Lepidopterous) Loopers Tomato Fruitworm Tomato Hornworm Tomato Pinworm	9 to 12	Apply when the majority of the population is at egg hatch to the second instar. For Colorado potato beetle, DO NOT apply more than twice to a single generation and DO NOT apply to successive generations. Use higher rates and higher spray volumes when populations are heavy, larvae are large, or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		Pepper Weevil	9 to 12	Apply at initial flowering stage.
		Leafminers (Dipteran) Stink Bugs Thrips Whiteflies	12	Apply when the majority of the target pest population is at egg hatch to early instars. DO NOT apply more than two applications against whiteflies or thrips per year. Use higher rates and higher spray volumes when populations are heavy, larvae are large, or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		Fruiting Vegetables (Field Grown) Use Restrictions: • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. • The use of novaluron on crops grown for food in greenhouses, except tomatoes and cucumbers, is prohibited.		

Crops	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
TROPICAL FRUITS (SUBGROUP 24A)* INCLUDING LYCHEE; AISEN; BAEL FRUIT; BURMESE GRAPE; CAT'S-EYES; INGÁ; LONGAN; MADRAS-THORN; MANDURO; MATISIA; MESQUITE; MONGONGO, FRUIT; PAWPAW, SMALLFLOWER; SATINLEAF; SIERRA LEONE-TAMARIND; SPANISH LIME; VELVET TAMARIND; WAMPI; WHITE STAR APPLE; CULTIVARS, VARIETIES, AND HYBRIDS OF THESE COMMODITIES.	DO NOT apply within 1 day of harvest.	Sri Lanka Weevils	1.25 quarts (40 fl. ozs.) in 100 gallons of water	Apply when the majority of the population is at egg hatch to the second instar. Repeat applications maximum 4 times if needed to protect new growth, but not less than 10 days apart.
		Tropical Fruits (Subgroup 24A) Use Restrictions: • DO NOT apply more than 3 applications against Sri Lanka weevils per year. • DO NOT apply more than 3.75 quarts (120 fl. ozs. (0.78 lb. a.i.)) per acre per calendar year. * Not registered for use in California.		

Crops	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
PEARS (GROUP 11-10 Pear; Asian Pear)* (for use only in Colorado, Michigan, New York, Pennsylvania, Washington and Oregon)	DO NOT apply within 14 days of harvest.	Codling Moth	20 to 32	Begin applications prior to egg deposition or shortly thereafter to prevent codling moth damage to fruit. However, best protection is achieved when application is initiated at the beginning of oviposition.
		Leafrollers (Oblique-Banded, Pandemis)	20 to 32	Initiate applications at cluster bud timing up to "Pear turn down" stage of development.
		Pear Psylla	20 to 32	Set the timing to occur during dormant through pear turn-down stage with the initiation of pear psylla oviposition.
		Pears (Group 11-10; Asian Pear) Use Restrictions: • DO NOT apply more than 96 fl. ozs. (0.62 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply this product when crop is in bloom. • DO NOT apply more than three applications per calendar year. NOTE: • If your growing region uses a Degree Day (DD) or Biofix model, or no model is available, consult the local cooperative extension, professional consultants, or qualified advisories to ensure the proper timing for the intended target pest. • One repeat application can be made to protect new foliage growth, but not less than 10 days after the first application. Phytotoxicity: Given the right set of environmental conditions phytotoxicity may occur when applied after pear turn-down. Factors increasing the probability of crop injury are: 1) varietal sensitivity; 2) excessive rainfall, high temperatures and/or drought, and; 3) incompatibility with other products (e.g., oils or strobilurin fungicides). * Not registered for use in California.		

Crops	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
POME FRUIT, GROUP 11-10 EXCEPT PEARS (see separate directions for PEARS) (Apple; azarole; crabapple; loquat; mayhaw; medlar; quince; Chinese quince; Japanese quince; tejocote; cultivars, varieties, and/or hybrids of these)	DO NOT apply with- in 14 days of harvest.	Budmoths (Eyespotted, Tufted Apple)	20 to 40	For each generation, make an application at the beginning of egg hatch.
		Codling Moth	20 to 40 (Eastern USA) 20 to 50 (Western USA)	For all generations, best protection is achieved when applications are initiated at the beginning of oviposition. RIMON 0.83EC Insecticide must be applied prior to egg deposition or shortly thereafter to prevent codling moth damage to fruit. Apply RIMON 0.83EC Insecticide at the following timings: 1st Generation: Begin applications at 50 - 100 DD from Biofix, or 225 - 275 DD from January 1. NOTE: Biofix is defined as the date of first sustained adult catch in pheromone traps - typically five moths in three traps in a seven-day period. 2nd Generation: Begin applications at 1000 DD from Biofix, or 1175 DD from January 1. Follow with subsequent applications at approximately 14 to 17-day intervals, if sustained moth pressure is high.
		Lacanobia Fruitworm	20 to 50	Begin applications when the majority of eggs have hatched and larvae are in the first to third instar stages.
		Leafminers (Spotted Tentiform, Western Tentiform)	15 to 40	Application timing for leafminers varies between species and geographic locations. Monitor the moth flights and treat at egg hatch for each generation.
		Leafrollers (European, Fruittree, Redbanded, Variegated)	20 to 40	For control of the surface or foliar feeding leafroller larval complex, application can be made at any time larvae are feeding. However, most effective crop protection results from application made at the initiation of egg hatch.
		Leafrollers (Oblique-banded, Pandemis)	20 to 50 (Eastern USA) 30 to 50 (Western USA)	Apply RIMON 0.83EC Insecticide treatments at the following timings: 1st Generation: Begin applications during pink to petal fall period. 2nd Generation: Begin application targeting 20% egg hatch.
		Oriental Fruit Moth	20 to 40	Begin applications before egg hatch of each generation to prevent larval penetration of the fruit.
		Plant Bug, White Apple Leafhopper	20 to 50	Populations of immature stages of plant bugs and/or white apple leafhopper may be suppressed with applications of RIMON 0.83EC Insecticide. RIMON 0.83EC Insecticide will not control adults of these pests due to its mode of action.
		Stink Bug spp.* Including Brown Marmorated Stink Bug	20 to 30	Apply when adults are first detected. For adult control, tank mix with an adulticide.
Pome Fruit (Group 11-10 Except Pears) Use Restrictions: • DO NOT apply more than 150 fl. ozs. (0.97 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply this product when crop is in bloom. • DO NOT apply more than three applications per calendar year. NOTE: • The Degree Days (DD) listed in the above Application Instructions are based on Biofix dates for specific target pests. If your growing region uses a different DD or Biofix model, or no model is available, consult the local cooperative extension, professional consultants, or qualified advisories to ensure the proper timing for the intended target pest. • Best protection is achieved when applications are initiated at the beginning of egg oviposition. • RIMON 0.83EC Insecticide will provide up to 14 days of protection depending on the application rate and rate of foliage growth and fruit expansion. • Repeat applications as needed to protect new foliage growth, and fruit, but not less than 10 days apart. • Use the higher rates and shorter application intervals for heavy infestations or under continuous pest pressure. • For situations of heavy infestations and continuous moth flight and egg oviposition, and where it is difficult to obtain thorough coverage, use the highest labeled rate and maintain coverage with timely reapplications at 10 to 14 day intervals. • RIMON 0.83EC Insecticide may be alternated or tank mixed with other insecticides targeted against the same pest as long as the application interval does not exceed the period of effectiveness of the alternate product. *Not registered for use in California.				

Crops	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
POTATOES/ SWEET POTATOES	DO NOT apply within 14 days of harvest.	Armyworms	6 to 12	Apply when the majority of the population is at egg hatch to the second instar. Use higher rates and higher spray volumes when larvae are large, or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, but not less than 7 days apart.
		Colorado Potato Beetle		
		European Corn Borer		
		Foliage Feeding Caterpillars		
		Loopers		
Potato Tuberworm				
Sweet Potato Leafminer				
Whiteflies		12		
Potato Psyllid*		12	Apply on a preventative basis or when first evidence of zebra chip disease and/or live psyllids are detected in the growing area. Repeat application at 7 - 14-day interval or alternate with an adulticide product for optimum control.	
Potatoes/Sweet Potatoes Use Restrictions: • DO NOT apply to successive generations of Colorado Potato Beetle. • DO NOT apply more than two applications against whiteflies per year. • DO NOT apply more than 24 fl. ozs. (0.16 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than two applications per calendar year. NOTE: • For application to potatoes through irrigation systems, refer to the section entitled APPLICATION THROUGH IRRIGATION SYSTEMS - CHEMIGATION. * Not registered for use in California.				

Crops	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
STONE FRUITS (capulin; black cherry; Nanking cherry; sweet cherry; tart cherry; cultivars, varieties, and/or hybrids of these; nectarine; peach; cultivars, varieties, and/or hybrids of these; apricot; Japanese apricot; Chinese jujube; plum; American plum; beach plum; Canada plum; cherry plum; Chickasaw plum; Damson plum; Japanese plum; Klamath plum; prune plum; plumcot; sloe; cultivars, varieties, and/or hybrids of these)	DO NOT apply within 8 days of harvest.	Fruit Flies (Cherry, W. Cherry, <i>Drosophila</i> spp. Including <i>Spotted Wing Drosophila</i> *)	20 to 40	Begin applications when adults are detected in the orchard, or after 950 degree days (DD) from March 1st. Adult females will deposit non-viable eggs after contact with, and feeding on, treated foliage and fruit, providing control of eggs and larvae. For adult control, tank mix with an adulticide. Thorough coverage is needed to achieve optimum effect. DO NOT use spray volumes below 100 GPA. DO NOT make alternate row treatments.
		Leafrollers (Oblique-banded, Pandemis)	20 to 50 (Eastern USA) 30 to 50 (Western USA)	Control of leafrollers is best when applications are timed against early (first to fourth) instar larvae. Apply RIMON 0.83EC Insecticide at the following timings: 1st Generation: Begin applications during the pink to petal fall period. 2nd Generation: Begin application targeting 20% egg hatch.
		Leafrollers (European, Fruittree, Redbanded, Variegated)	20 to 40	For control of the surface or foliar feeding leafroller larval complex, application can be made at any time larvae are feeding. However, most effective crop protection results from application made at the initiation of egg hatch.
		Lesser Peachtree Borer	20	In southeast, apply in a tank mix with either a pyrethroid or phosmet after April 1st, and again in 2 to 4 weeks.
		Oriental Fruit Moth	20 to 40	Begin applications before egg hatch of each generation to prevent larval penetration of the fruit.
		Peachtree Borer	20	In southeast, apply in pre-harvest applications to cultivars ripening after July 1st.

(continued)

Crops	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
STONE FRUITS (continued) (capulin; black cherry; Nanking cherry; sweet cherry; tart cherry; cultivars, varieties, and/or hybrids of these; nectarine; peach; cultivars, varieties, and/or hybrids of these; apricot; Japanese apricot; Chinese jujube; plum; American plum; beach plum; Canada plum; cherry plum; Chickasaw plum; Damson plum; Japanese plum; Klamath plum; prune plum; plumcot; sloe; cultivars, varieties, and/or hybrids of these)	DO NOT apply within 8 days of harvest.	Peach Twig Borer	20 to 40	Dormant/Delayed Dormant: Apply RIMON 0.83EC Insecticide with 4 to 6 gallons per acre of narrow range oil. Always use the higher rates if the orchard has a history of heavy populations. In-season: Monitor orchard from bloom onward for shoot strikes at the end of each generation. Shoot strikes first appear when the degree-day accumulation from moths in traps approaches 400 DD ₅₀ but more will be evident around 700 - 800 DD ₅₀ . If larvae or their damage are observed at this time, make application in sufficient spray volume for thorough coverage.
		Sap Beetle	20	Apply in a tank mix with adulticides to help effect egg hatch.
		Stink Bug spp.* Including Brown Marmorated (Immature)	20 to 40	Apply when thresholds are reached. For adult control, tank mix with an adulticide.
		Stone Fruits Use Restrictions: - Repeat applications as needed to protect new foliage growth and fruit, but not less than 7 days apart. DO NOT apply more than 150 fl. ozs. (0.97 lb. a.i.) of formulated product per acre per calendar year. DO NOT apply this product when crop is in bloom. DO NOT apply more than three applications per calendar year. NOTE: - The Degree Days (DD) listed in the above Application Instructions are based on timing for specific target pests. If your growing region uses a different DD or Biofix model, or no model is available, consult the local cooperative extension, professional consultants, or qualified advisories to ensure the proper timing for the intended target pest. - Best protection is achieved when applications are initiated at the beginning of egg oviposition. - RIMON 0.83EC Insecticide will provide up to 14 days of protection depending on the application rate and rate of foliage growth and fruit expansion. - Use the higher rates and shorter application intervals for heavy infestations or under continuous pest pressure. - For situations of heavy infestations and continuous moth flight and egg oviposition, and where it is difficult to obtain thorough coverage, use the highest labeled rate and maintain coverage with timely reapplications at 10 to 14-day intervals. - RIMON 0.83EC Insecticide may be alternated or tank mixed with other insecticides targeted against the same pest as long as the application interval does not exceed the period of effectiveness of the alternate product. * Not registered for use in California.		

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
STRAWBERRY	DO NOT apply within 1 day of harvest.	Armyworms Corn Earworm Loopers <i>Lygus</i> <i>Thrips</i> Webworms	9 to 12	Apply when the majority of the population is at egg hatch to the second instar. For <i>Lygus</i> , apply when adults are observed in the field and just prior to egg hatch. Optimum control will be achieved with the 12 fl. ozs./A rate. Spray with a sufficient volume of water to ensure thorough coverage of fruit and leaf surfaces. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		<i>Thrips</i> (Western Flower, Chili, etc.) spp.*	6 to 12	Apply when <i>Thrips</i> populations begin to build. For adult control, tank mix with an adulticide. Spray with a sufficient volume of water to ensure thorough coverage of fruit and leaf surfaces. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		Asian Cockroach* Sap Beetles*	6 to 12	Apply when adults appear and prior to egg hatch. For adult control of all life stages, tank mix with an adulticide. Spray with a sufficient volume of water to ensure thorough coverage of fruit and leaf surfaces. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		Strawberry Use Restrictions: • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. * Not registered for use in California.		

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
SUCCULENT SHELLED BEAN* ANDEAN LUPIN, SUCCULENT SHELLED BLACKKEYED PEA, SUCCULENT SHELLED BLUE LUPIN, SUCCULENT SHELLED BROAD BEAN, SUCCULENT SHELLED CATJANG BEAN, SUCCULENT SHELLED COWPEA, SUCCULENT SHELLED CROWDER PEA, SUCCULENT SHELLED GOA BEAN, SUCCULENT SHELLED GRAIN LUPIN, SUCCULENT SHELLED JACKBEAN, SUCCULENT SHELLED LABLAB BEAN, SUCCULENT SHELLED LIMA BEAN, SUCCULENT SHELLED MOTH BEAN, SUCCULENT SHELLED SCARLET RUNNER BEAN, SUCCULENT SHELLED SOUTHERN PEA, SUCCULENT SHELLED SWEET LUPIN, SUCCULENT SHELLED VEGETABLE SOYBEAN, SUCCULENT SHELLED VELVET BEAN, SUCCULENT SHELLED WAX BEAN, SUCCULENT SHELLED WHITE LUPIN, SUCCULENT SHELLED WHITE SWEET LUPIN, SUCCULENT SHELLED YELLOW LUPIN, SUCCULENT SHELLED	DO NOT apply within 1 day of harvest.	Armyworms Loopers Webworms Bean Leaf Beetle Bean Plataspid Cucumber Beetle Mexican Bean Beetle <i>Lygus</i> <i>Thrips</i> Whiteflies	12 in 30 gallons of water	Apply when the majority of the target pest population is at egg hatch to early instars. Use higher rates and higher spray volumes when larvae are large or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		Succulent Shelled Bean Use Restrictions: • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. * Not registered for use in California.		

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
SUCCULENT SHELLED PEA* CHICKPEA, SUCCULENT SHELLED ENGLISH PEA, SUCCULENT SHELLED GARDEN PEA, SUCCULENT SHELLED GREEN PEA, SUCCULENT SHELLED LENTIL, SUCCULENT SHELLED PIGEON PEA, SUCCULENT SHELLED	DO NOT apply within 1 day of harvest.	Armyworms Loopers Webworms Bean Leaf Beetle Bean Plataspid Cucumber Beetle Mexican Bean Beetle Lygus Thrips Whiteflies	12 in 30 gallons of water	Apply when the majority of the target pest population is at egg hatch to early instars. Use higher rates and higher spray volumes when larvae are large or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, and fruit, but not less than 7 days apart.
		Succulent Shelled Pea Use Restrictions: • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. * Not registered for use in California.		

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
SWEET CORN	DO NOT apply within 1 day of harvest.	Armyworms Corn Earworms Eur. Corn Borers Foliage Feeding Caterpillars Grasshoppers* (Nymphs Only)	6 to 12	Pre-tassel Timing: Apply when adult activity is first observed or when the majority of the immature population is at egg hatch to second instar. For optimum corn earworm and corn borer control, tank mix with a knockdown and/or adulticide. Silking/Post-tassel Timing: Apply when adult activity is first observed or when eggs begin to hatch. Apply only in a tank mix with knockdown or adulticide products. Apply in sufficient volume to ensure full coverage of foliage and developing ears. Use higher rates and higher spray volumes when larvae are large or foliage canopy is tall or dense.
		Sap Beetle* Cucumber Beetle*	6 to 12	Apply when adults first appear and prior to egg hatch. Apply in sufficient volume to ensure full coverage of foliage and developing ears. Use higher rates and higher spray volumes when larvae are large or foliage canopy is tall or dense.
		Sweet Corn Use Restrictions: • Repeat applications as needed to protect new growth, but not less than 7 days apart. • DO NOT apply more than 60 fl. ozs. (0.39 lb. a.i.) of formulated product per acre per calendar year. • The retreatment of sweet corn with novaluron is prohibited (i.e., only 1 application at 0.078 lb. a.i./A) in arid areas which receive less than 20 inches of precipitation (including irrigation) per year. • DO NOT apply more than five applications per calendar year. NOTE: • For application to sweet corn through irrigation systems, refer to the section entitled APPLICATION THROUGH IRRIGATION SYSTEMS - CHEMIGATION. * Not registered for use in California.		

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
SWISS CHARD*	DO NOT apply within 1 day of harvest.	Armyworms Cucumber Beetle Loopers	9 to 12	Apply when the majority of the population is at egg hatch to the early instars.
		Beet Webworm	12	Apply during oviposition through early instar stages. Use higher spray volumes and increased pressure to ensure complete coverage and penetration to immature leaves at the base of the plant.
		Swiss Chard Use Restrictions: • Repeat applications as needed to protect new foliage growth, but not less than 7 days apart. • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply more than three applications per calendar year. * Not registered for use in California.		

Crop	PHI	Target Pests	Rates (fl. ozs./A)	Application Instructions
TURNIP GREENS*	DO NOT apply within 7 days of harvest.	Alfalfa Looper Armyworms Cabbage Loopers Cabbage Webworm Corn Earworm Cucumber Beetles Diamondback Moth Imported Cabbageworm Leafminers (Dipteran and Lepidopteran) Southern Cabbageworm	6 to 12	Apply when the majority of the population is at egg hatch to the second instar. Use higher rates and higher spray volumes when larvae are large, when target pests populations is 2X or more above state threshold level or foliage canopy is tall or dense. Repeat applications as needed to protect new growth, but not less than 7 days apart.
		Lygus Bugs Stink Bugs <i>Thrips</i> Vegetable Weevil Whiteflies	12	
		Turnip Greens Use Restrictions: • DO NOT apply more than 2 applications against whiteflies per year. • DO NOT apply more than 36 fl. ozs. (0.23 lb. a.i.) of formulated product per acre per calendar year. • DO NOT apply to turnips harvested for the root. • DO NOT feed turnip tops to livestock. • DO NOT apply more than three applications per calendar year. * Not registered for use in California.		

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals, in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Keep above freezing.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility. **DO NOT** pour or dispose of down the drain or sewer. Call your local solid waste agency for local disposal options.

CONTAINER HANDLING:

Nonrefillable Container (five gallons or less): Nonrefillable container. **DO NOT** re-use or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available or reconditioning, or puncture and dispose of in a sanitary landfill, or other procedures allowed by State and local authorities.

Nonrefillable Container (greater than five gallons): Nonrefillable container. **DO NOT** reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling, if available or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures allowed by State and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES** and **LIMITATIONS OF LIABILITY**.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of UPL NA Inc. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, UPL NA Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of UPL NA Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, UPL NA Inc. disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at UPL NA Inc.'s election, the replacement of product.

RIMON is a registered trademark of an ADAMA Group Company.

UPL and the UPL logo are trademarks of a UPL Corporation Limited Group Company.

©2024 UPL Corporation Limited Group Company

2022957-1 (0424)

ESL101222-11331-040324

RIMON® 0.83EC

INSECTICIDE

Insecticide for use on: Avocado*; Berries (Low-Growing); Brassica, head and stem (Group 5-16); Brassica, Leafy Greens (Subgroup 4-16B); Bushberries; Carrot*; Individual Legume Vegetable Commodities; Pears*; Pome Fruits; Potatoes/Sweet Potatoes; Stonefruits; Strawberry; Sweet Corn; Swiss Chard*; Turnip Greens*; Vegetable, cucurbit (Group 9); Vegetable, fruiting (Group 8-10); Kohlrabi; Tropical Fruits (Subgroup 24A)* including Lychee; Vegetable, foliage of legume, subgroup 7A, forage; and Vegetable, foliage of legume, subgroup 7A.

* Not registered for use in California.

ACTIVE INGREDIENT:	% BY WT.
Novaluron: 1-[3-chloro-4-(1,1,2-trifluoro-2-trifluoromethoxyethoxy)phenyl]-3-(2,6-difluorobenzoyl)urea**	9.3%
OTHER INGREDIENTS:	90.7%
TOTAL:	100.0%

** Contains 0.83 lb. novaluron per gallon.

EPA Reg. No. 66222-35-70506

EPA Est. No. 037429-GA-002

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

IF ON SKIN: • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.

IF SWALLOWED: • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • **DO NOT** induce vomiting unless told to do so by a poison control center or doctor. • **DO NOT** give anything by mouth to an unconscious person.

IF INHALED: • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

Emergency Assistance: Have the product container or label with you when calling a poison control center or doctor or going for treatment. For non-emergency general information on this pesticide product (including health concerns or pesticide incidents), you may call 1-877-250-9291, 24 hours per day, 7 days per week. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

In case of spills, fire, leaks or accidents call 1-800-535-5053.

For additional precautionary, handling and use statements, see inside of this booklet.

Net Contents: 2.5 Gallons

Distributed by: UPL NA Inc. • 630 Freedom Business Center, Suite 402, King of Prussia, PA 19406 U.S.A. • 1-800-438-6071

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING: Causes substantial but temporary eye injury. **DO NOT** get in eyes or on clothing. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater and estuarine/marine invertebrates. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** contaminate water when disposing of equipment wash waters or rinsate. This product may impact surface water quality due to runoff or rain water. This is especially true for poorly draining soils and soils with shallow ground water.

Pollinator Advisory: Because of its mode of action as an insect growth regulator, and since it is not systemic, RIMON® 0.83EC Insecticide has the potential to impact larval bees (i.e., brood). In order to minimize the possibility of effects to honeybee brood, **DO NOT** use RIMON 0.83EC Insecticide on blooming crops when bees are actively foraging.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff for several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of novaluron from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Alternatively, for products with water-in requirements, avoid watering to the point of runoff.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE: Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals, in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Keep above freezing.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility. **DO NOT** pour or dispose of down the drain or sewer. Call your local solid waste agency for local disposal options.

CONTAINER HANDLING:

Nonrefillable Container (five gallons or less): Nonrefillable container. **DO NOT** re-use or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available or reconditioning, or puncture and dispose of in a sanitary landfill, or other procedures allowed by State and local authorities.

