



A biochemical pesticide that

- Is especially formulated for use on a wide range of agricultural crops to provide protection against crop yield loss from damage caused by plant-parasitic nematodes,
- Enhances resistance of treated crops to nematode damage through activating plant's natural defense against nematode feeding and egg hatching,
- Regulates genes involved in drought tolerance pathways, stimulates plant secondary metabolites that directly inhibit nematode egg hatching,
- Enhances crop growth, quality, and yield.

Wettable Dry Granule

ACTIVE INGREDIENT:

PDHP 68949 1.0%

OTHER INGREDIENTS: 99.0%

Total: 100.0%

EPA Reg. No. 71771-16-2935

EPA Est. No. 88746-GA-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION / PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

Read the entire label before using this product. Read Conditions of Sale and Limitation of Warranty and Liability before buying or using. If terms are not acceptable, return at once unopened.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if inhaled. Avoid breathing dust or spray mist. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID	
If inhaled	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment information, call the National Pesticide Information Center at 1-800-858-7378, 6:30 AM to 4:30 PM Pacific Time, seven days a week. During other times, call the poison control center at 1-800-222-1222.	

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes and socks
- Waterproof gloves
- A NIOSH approved particulate respirator with any N, R, or P filter with NIOSH approval number prefix TC-84A; or a NIOSH approved powered air-purifying respirator with HE filter with NIOSH approval number prefix TC-21C. Repeated exposures to high concentrations of microbial proteins can cause allergic sensitization.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval and notification to workers. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water) is:

- Coveralls
- Waterproof gloves, and
- Shoes and socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

PRODUCT INFORMATION

ATALIOL is effective against plant-parasitic nematodes. This product is a wettable dry granule to be diluted with water. Shake well before use. When a rate range is given for a specific crop, use the higher listed rate when conditions favor higher pest pressure. ATALIOL is a plant response-elicitor peptide based on a naturally occurring harpin protein. It reduces the incidence and severity of plant nematode damage by eliciting the plant's own natural defense system, referred to as systemic acquired resistance (SAR). Early applications prior to nematode infestation either alone or in combination with another registered nematicide can delay or reduce nematode damage and severity. This product is most effective when combined with nematicide programs having alternative modes of action. The plant's enhanced defense response system due to the application of ATALIOL adds to nematicide programs, leading to improved overall nematode control.

ATALIOL is a tool for use in Integrated Crop Management (ICM). It has no direct killing effect on nematodes. Product efficacy and protection against damage from plant parasitic nematodes is based on its ability to activate the plant's own internal defense and growth mechanisms. This product works best when used in conjunction with good crop management practices.

Growth Enhancement and Marketable Yield: Use ATALIOL to enhance plant growth and quality and increase plant health and vigor to assist in the management of certain nematode species and to enhance crop quality and marketable yield.

Sites: Use ATALIOL for greenhouse, shadehouse, nursery, and field production of all plants listed on this label.

Coverage: Use spray volume adequate to obtain coverage without runoff. Uniform or full leaf coverage is helpful but is not required.

Days to Harvest: This product can be applied up to the day of harvest.

RESISTANCE MANAGEMENT RECOMMENDATIONS

Nematode Pressure and Management

A variety of nematode population pressures can exist in field conditions. The visible efficacy of a single product will depend upon the effectiveness of the product and the level of nematode population. Under extremely high nematode pressure no single product will provide high-level nematode control, and in these circumstances a range of nematode management measures should be undertaken by the grower to reduce the nematode pressure, such as use of rotations, fallow periods, resistant or tolerant varieties, application of additional chemical and biological nematode control agents.

Integrated Nematode Management: Consider the following recommended management practices to sustainably control or suppress plant-parasitic nematode populations:

- Take soil samples regularly to determine the plant-parasitic nematode species present and population density from the previous crop.
- Consider using nematode resistant or tolerant crop varieties.
- Consider cultural methods to reduce nematode populations, e.g., fallow periods, rotations, or soil amendments.
- Consider a combination of effective products in a nematode management program – pre-planting and in the crop, e.g., approved fumigants, nematicides, and other products that protect crop roots. This may be critical under high pest pressure situations where no single product may provide sufficient control.
- To minimize the possibility of nematode resistance from occurring, avoid long-term repeated applications of the same product or group of compounds in the same field. If any reduction in product performance is noted, it is important to contact the local company representative.
- Monitor treated nematode populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and / or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Wilbur-Ellis Company LLC at (720) 306-6340. You can also contact your pesticide distributor or university extension specialist to report nematode resistance.

USE PRECAUTIONS

Use Promptly: Use product on the same day it is mixed with water and use opened packages within 3 weeks. Carefully reseal opened packages to minimize exposure to air and moisture.

TANK MIXING

Use in Mixtures: ATALIOL is believed to be compatible with most other labeled pesticides, such as post-emergence herbicides, insecticides, acaricides, and fungicides, as well as most foliar nutritional products. To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. Add dry flowables first, then flowables, and then emulsifiable concentrates last. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Once compatibility has been proven, use the same procedure for adding products to the spray tank. If ATALIOL cannot be put in the tank first, before adding it to the tank, slurry the product in a small amount of clean water and add the slurry to the tank. Do not mix this product with any other product containing a label prohibition against such mixing. When tank mixing this product with any other approved pesticide, always read and follow all use directions, restrictions, and precautions of both ATALIOL and the tank mix partner(s). The resulting tank mix must be used in accordance with the most restrictive label limitations and precautions. Do not exceed label dosage rates.

Glyphosate Herbicides: When tank mixing ATALIOL with glyphosate for application on crops designated Roundup Ready®, only use formulations of glyphosate herbicide that are fully labeled for use on Roundup Ready® crops. Never spray this mixture on crops that are not designated Roundup Ready® as severe injury or death of the crop can occur. Some glyphosate formulations allow for the addition of surfactants. Please refer to the surfactant recommendations from the manufacturer.

Precautions: Do not use this product in tank mixes or water below pH 5 or above pH 10. Do not mix this product with pyrophosphates, phosphoric acid, or other strong oxidizers.

Surfactants: Use only non-ionic adjuvants approved for use on growing crops.

DILUTION WATER QUALITY: If dilution water is high in total mineral content, salinity, suspended solids and/or exhibits any other factors that reduce the solubility of this product, then first dissolve ATALIOL in an appropriate volume of “clean water” such as municipal tap water. Pour the dissolved ATALIOL solution into the dilution water for the spray (see step 1 of “Mixing Instructions”).

DILUTION WATER QUANTITY: If 1 ounce of ATALIOL is diluted in more than 35 gallons of chlorinated water (e.g., municipal water), add a labeled water treatment product such as sodium sulfite, sodium bisulfite, or sodium metabisulfite to remove excess chlorine before adding this product. Consult your Wilbur-Ellis Company representative if you need further directions on water treatment.

RAIN: Do not apply during rain. Reapplication is not necessary if the spray has dried before rain begins.

STRESSED PLANTS: Plants must be actively growing at the time of foliar applications. Applications made to plants that are stressed by extreme heat, cold, moisture, or nutrient deficiency can be less effective.

SPRAY DRIFT: Avoiding spray drift at the application site is the responsibility of the applicator. The interactions of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering these factors when making decisions.

AERIAL APPLICATIONS:

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must not exceed 65% of the wingspan for fixed wing aircraft or 75% of the rotor diameter for helicopters. Otherwise, the boom length must not exceed 75% of the wingspan for fixed wing aircraft or 90% of the rotor diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

AIRBLAST APPLICATIONS:

- Sprays must be directed into the canopy.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at the row end and when spraying the outer row.
- **DO NOT** apply during temperature inversions.

GROUND BOOM APPLICATIONS:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

BOOMLESS GROUND APPLICATIONS:

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

MIXING INSTRUCTIONS

- Step 1. Fill at least one-half of the mix tank with clean water. Provide gentle agitation.
- Step 2. If 1 ounce of ATALIOL is diluted in more than 35 gallons of chlorinated water, add a labeled dechlorination water treatment product (such as sodium sulfite, sodium bisulfite, or sodium metabisulfite) and continue gentle agitation (see Use Precautions on "Dilution Water Quantity").
- Step 3. If dilution water quality is suspect, dissolve ATALIOL in clean water before adding to the mix tank (see Use Precautions on "Dilution Water Quality"). Add the required amount of this product. Agitate until dissolved and avoid excessive foaming.
- Step 4. If tank mixing, add other materials to the mix tank. Add remaining water to mix tank.
- Step 5. Continue gentle agitation and apply promptly.

APPLICATION INSTRUCTIONS

Apply ATALIOL as a pre-transplant root dip or as a foliar application.

Pre-Plant Dip: Apply this product as a pre-plant foliar dip or root dip to vegetable crops (e.g., cucurbit and cole vegetables), ornamentals, tobacco, and small fruit crops at the rate of 0.5-3 oz per 100 gallons of water immediately prior to transplanting. Wash transplants to remove excess soil prior to dipping. Completely immerse planting stock in solution. Dip, soak, or expose plants for a minimum of 2 to 5 minutes. **DO NOT** reuse dip solution. Dispose of dip solution according to local restrictions. Plant treated plants as quickly as possible. For continued crop management, follow with foliar applications of this product as specified in the tables that follow.

Foliar Application: Apply this product as a greenhouse or field application using conventional ground or aerial equipment. Use this product as part of an integrated pest management (IPM) program to assist nematode management.

Application via Sprinkler or Drip (Trickle) Chemigation Systems: Apply this product only through sprinkler (including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move) and drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury and lack of effectiveness can result from non-uniform distribution of treated water. If you have questions about your system's calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Sprinkler Chemigation System Requirements: The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment. Apply product continuously for the duration of the water application.

Drip (Trickle) Chemigation System Requirements: The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back-flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Apply product continuously for the duration of the water application.

CROPS AND APPLICATION RATES

Use the following tables to decide upon application rates and timing. Follow applicable specifications for your crop.

TRANSPLANT APPLICATIONS

USE SITE & CROP	RATE	TIMING
Transplant Any Crop Listed Elsewhere on This Label	0.5-3 oz per 100 gal water	Apply 5-7 days before transplanting. Can also be applied as a plant drench at transplanting.

FOOD CROP RECOMMENDATIONS

CEREAL GRAINS (Crop Group 15) and FORAGE, FODDER AND STRAW OF CEREAL GRAINS (Crop Group 16) Including, but not limited to, Barley; Buckwheat; Corn; Millet, Pearl; Millet, Proso; Oats; Popcorn; Rice; Rye; Sorghum (milo); Teosinte; Triticale; Wheat; and Wild rice.			
NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
CORN (Grain, Silage, Seed, or Pop)			
Awl (<i>Dolichodorus</i> spp.) Dagger (<i>Xiphinema</i> spp.) Lance (<i>Hoplolaimus</i> spp.) Lesion (<i>Pratylenchus</i> spp.) Needle (<i>Longidorus</i> spp.) Ring (<i>Mesocriconema</i> spp.) Sheath (<i>Hemicyclophora</i> spp.) Spiral (<i>Helicotylenchus</i> spp.) Sting (<i>Belonolaimus</i> spp.) Stubby-root (<i>Paratrichodorus</i> spp.) Stunt (<i>Tylenchorynchus</i> spp.)	0.5-3	5-200 by ground 2-50 by air	Apply beginning early in the season at V-2 to V-6 stage.
ALL OTHER CEREAL GRAINS			
Cereal Cyst (<i>Heterodera</i> spp.) Root-knot (<i>Meloidogyne</i> spp.) Root Lesion (<i>Pratylenchus</i> spp.) Stunt (<i>Tylenchorhynchus</i> spp.)	0.5-3	5-200 by ground 2-50 by air	Apply early at 3-5 leaf and every 14-35 days or in combination with nematicide programs.

LEGUME VEGETABLES – SUCCULENT OR DRIED (Crop Group 6) Including, but not limited to, Bean (Includes Grain Lupin, Sweet Lupin, White Lupin, and White Sweet Lupin); Bean (Includes Field, Kidney, Lima, Navy, Pinto, Runner, Snap, Tepary, and Wax); Bean (Includes Adzuki Bean, Asparagus Bean, Blackeyed Pea, Catjang, Chinese Longbean, Cowpea, Crowder Pea, Moth Bean, Mung Bean, Rice Bean, Southern Pea, Urd Bean, and Yardlong Bean); Broad Bean; Chickpea (Garbanzo Bean); Guar; Jackbean; Lablab Bean; Lentil; Pea (Includes Dwarf Pea, Edible-Pod Pea, English Pea, Field Pea, Garden-Pea, Green Pea, Snow Pea, and Sugar Snap Pea); Pigeon Pea; Soybean; Soybean (immature seed); and Sword Bean.			
SOYBEAN			
NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Awl (<i>Dolichodorus</i> spp.) Lance (<i>Hoplolaimus</i> spp.) Root-knot (<i>Meloidogyne</i> spp.) Root Lesion (<i>Pratylenchus</i> spp.) Soybean Cyst (<i>Heterodera</i> spp.) Stunt (<i>Tylenchorhynchus</i> spp.)	0.5-3	10-200 by ground 2-50 by air	Begin applications at VC to V6 or prior to nematode infestation, whichever is earlier.
ALL OTHER LEGUMES			
NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Root-knot (<i>Meloidogyne</i> spp.) Root Lesion (<i>Pratylenchus</i> spp.) Reniform (<i>Rotylenchulus</i> spp.) Stem / Bulb (<i>Ditylenchus</i> spp.)	1-3	10-200 by ground 2-50 by air	Begin applications at early emergence or first new growth and repeat at 14-28 day intervals in combination with nematicide programs.

NON-GRASS ANIMAL FEEDS – FORAGE, FODDER, STRAW AND HAY (Crop Group 18)
Including, but not limited to, Alfalfa; Bean, Velvet; Clover; Kudzu; Lespedeza; Lupin; Sainfoin; Trefoil; Vetch; Vetch, Crown; Vetch, Milk.

NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Clover Cyst (<i>Heterodera</i> spp.) Root-knot (<i>Meloidogyne</i> spp.) Root Lesion (<i>Pratylenchus</i> spp.)	0.5-3	10-200 by ground 2-50 by air	Begin applications at early emergence or first new growth and repeat at 14-35 day intervals in combination with nematicide programs.

OILSEED (Crop Group 20)
Including, but not limited to, Cottonseed; Flax Seed; Jojoba; Milkweed; Mustard Seed; Niger Seed; Poppy Seed; Rapeseed (canola varieties only); Safflower; Sesame; Sunflower.

COTTON

NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Awl (<i>Dolichodorus</i> spp.) Lance (<i>Hoplolaimus</i> spp.) Reniform (<i>Rotylenchulus</i> spp.) Root-knot (<i>Meloidogyne</i> spp.),	0.5-3	10-200 by ground 2-50 by air	For suppression of nematodes, begin applications prior to infestation and repeat at 14-35 day intervals in combination with nematicide programs.

SUNFLOWER

NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Potato Cyst (<i>Globodera</i> spp.) Reniform (<i>Rotylenchulus</i> spp.) Root-knot (<i>Meloidogyne</i> spp.)	0.5-3	10-200 by ground 2-50 by air	Apply the first application between stage V-6 to V-8 and repeat when the terminal bud forms (R1 stage). All applications can be in combination with nematicide programs.

ALL OTHER OILSEEDS

NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Root-knot (<i>Meloidogyne</i> spp.)	0.5-3	10-200 by ground 2-50 by air	Begin applications at early emergence or first new growth and repeat at 14-28 day intervals in combination with nematicide programs.

PEANUT

NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Peanut Root-knot (<i>Meloidogyne</i> spp.) Root Lesion (<i>Pratylenchus</i> spp.) Sting (<i>Belonolaimus</i> spp.)	1-3	10-200 by ground 2-50 by air	Begin applications at early emergence or first new growth and repeat at 14-35 day intervals in combination with nematicide programs.

POTATO (Including seed production)			
NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Awl (<i>Dolichodorus</i> spp.) Potato Cyst (<i>Globodera</i> spp.) Root-knot (<i>Meloidogyne</i> spp.) Root Lesion (<i>Pratylenchus</i> spp.) Sheath (<i>Hemicycliophora</i> spp.) Spiral (<i>Helicotylenchus</i> spp.) Sting (<i>Belonolaimus</i> spp.) Stubby-root (<i>Paratrichodorus</i> spp. and <i>Trichodorus</i> spp.) Stunt (<i>Tylenchorhynchus</i> spp.)	1-3	10-200 by ground 2-50 by air	Apply prior to nematode infestation. Repeat applications as needed every 14-28 days in combination with nematicide programs.

ROOT AND TUBER VEGETABLES (Crop Group 1)			
Including, but not limited to, Arracacha; Arrowroot; Artichoke, Chinese and Jerusalem; Beet, Garden and Sugar; Burdock, Edible; Canna, Edible; Carrot; Cassava, Bitter and Sweet; Celeriac (Celery Root); Chayote (Root); Chervil, Turnip-Rooted; Chicory; Chufa; Dasheen (Taro); Ginger; Ginseng; Horseradish; Leren; Parsley, Turnip-Rooted; Parsnip; Radish; Radish, Oriental; Rutabaga; Salsify; Salsify, Black and Spanish; Skirret; Sweet Potato; Tanier; Turmeric; Turnip; Yam (all types).			
NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Awl (<i>Dolichodorus</i> spp.) Needle (<i>Longidorus</i> spp.) Pin (<i>Paratylenchus</i> spp.) Reniform (<i>Rotylenchulus</i> spp.) Root-knot (<i>Meloidogyne</i> spp.) Root Lesion (<i>Pratylenchus</i> spp.) Sting (<i>Belonolaimus</i> spp.) Stubby-root (<i>Paratrichodorus</i> spp. and <i>Trichodorus</i> spp.) Sugar Beet Cyst (<i>Heterodera</i> spp.)	1-3	10-200 by ground 2-50 by air	Apply at early emergence prior to nematode infestation, then repeat applications every 14-28 days in combination with the normal nematicide program.

SUGARCANE			
NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Lance (<i>Hoplolaimus</i> spp.) Reniform (<i>Rotylenchulus</i> spp.) Ring (<i>Mesocriconema</i> spp., <i>Criconemoides</i> spp.) Root-knot (<i>Meloidogyne</i> spp.) Root Lesion (<i>Pratylenchus</i> spp.) Spiral (<i>Helicotylenchus</i> spp. and <i>Rotylenchus</i> spp.) Sting (<i>Belonolaimus</i> spp.) Stubby-root (<i>Paratrichodorus</i> and <i>Trichodorus</i> spp.) Stunt (<i>Tylenchorhynchus</i> spp.)	1-3	10-200 by ground 2-50 by air	Apply after new growth commences. Repeat at 14-35 day intervals as needed in combination with nematicide programs. Apply at in-furrow over seed cane stalks or pieces and ensure coverage of all buds.

TOBACCO (all types)			
NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Root-knot (<i>Meloidogyne</i> spp.) Stunt (<i>Tylenchorhynchus</i> spp.) Tobacco Cyst (<i>Globodera</i> spp.)	0.5-3	10-200 by ground 2-50 by air	Begin applications when plants are established. Repeat at 14-35 day intervals as needed in combination with nematicide programs.

NON FOOD CROP RECOMMENDATIONS

GRASSES GROWN FOR SEED			
NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Ring (<i>Mesocriconema</i> spp., <i>Criconemoides</i> spp., <i>Criconemella</i> spp.) Root-knot (<i>Meloidogyne</i> spp.) Spiral (<i>Helicotylenchus</i> spp. and <i>Rotylenchus</i> spp.) Sting (<i>Belonolaimus</i> spp.) Stubby-root (<i>Paratrichodorus</i> spp. and <i>Trichodorus</i> spp.)	0.5-3	10-200 by ground 2-50 by air	Apply at 14-28 day intervals, beginning at early emergence in combination with nematicide programs.

GRASS FORAGE, FODDER, AND HAY (Crop Group 17)			
Including, but not limited to bermuda grass, bluegrass and brome grass or fescue. Any grass, Gramineae family (either green or cured) except sugarcane and those included in the cereal grains group, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage.			
NEMATODE	RATE (Oz/Acre)	DILUTION (Gal/Acre)	TIMING
Ring (<i>Mesocriconema</i> spp., <i>Criconemoides</i> spp., <i>Criconemella</i> spp.) Root-knot (<i>Meloidogyne</i> spp.) Spiral (<i>Helicotylenchus</i> spp. and <i>Rotylenchus</i> spp.) Sting (<i>Belonolaimus</i> spp.) Stubby-root (<i>Paratrichodorus</i> spp. and <i>Trichodorus</i> spp.)	0.5-3	10-200 by ground 2-50 by air	Begin applications at early emergence. Repeat at 14-28 day intervals in combination with nematicide programs.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep product away from exposure to air, sunlight, moisture, or heat. Do not store in temperatures over 110°F for more than 7 days. Use product within 3 weeks of opening and the same day as mixing.

Pesticide Disposal: Wastes resulting from use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using the product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

ALL STATEMENTS MADE HEREIN ARE SUBJECT TO APPLICABLE LAW, AND TO THE EXTENT THERE IS ANY INCONSISTENCY OR CONTENTION, APPLICABLE LAW SHALL GOVERN.

The Directions for Use of the product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of many different factors including, without limitation, manner of use or application, weather, combination with other products, or crop conditions. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Manufacturer and Seller harmless from any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label. EXCEPT FOR THIS WARRANTY, THE PRODUCT IS FURNISHED "AS-IS", AND NEITHER SELLER NOR MANUFACTURER MAKES ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, WITH RESPECT TO THE SELECTION, PURCHASE OR USE OF THIS PRODUCT; SELLER AND MANUFACTURER SPECIFICALLY DISCLAIM ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE BEYOND WHAT IS STATED ON THE LABEL. Buyer and User accept all risks arising from any use of this product, including without limitation, uses contrary to label instructions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Manufacturer.

Neither Manufacturer nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE BUYER OR USER, AND THE EXCLUSIVE LIABILITY OF MANUFACTURER AND SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT, OR, AT THE ELECTION OF MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

These Conditions of Sale and Limitation of Warranty and Liability shall be interpreted, unless otherwise required by the law of the state of purchase, in accordance with the laws of the State of California, excluding its conflicts of laws rules, and may not be amended by any oral or written agreement.

All trademarks, service marks, trade names, trade dress, product names and logos appearing on this label are the property of their respective owners.

Always read and follow label instructions before buying or using this product. Roundup Ready® is a registered trademark of Monsanto Technology LLC. WILBUR-ELLIS logo is a registered trademark and NEMATOCIDE logo and ATALIOL are trademarks of Wilbur-Ellis Company LLC.

NET WEIGHT: 5 lb (2.26 kg)

BATCH CODE: _____

PIA:2026-0219
2026-0224

Manufactured for:

WILBUR-ELLIS COMPANY LLC

4582 S. Ulster Street, Suite 400, Denver, CO 80237
(720) 306-6340



WILBUR-ELLIS.