

FIFRA Section 24(c) Special Local Need (SLN) Label



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

PYROXSULAM	GROUP	2	HERBICIDE
FLUROXYPYR	GROUP	4	HERBICIDE

OpenSky®

EPA Reg. No. 62719-721

EPA 24(c) Special Local Need Registration SLN WA-250001

For Distribution and Use Only Within the State of Washington

Postemergence Grass and Broadleaf Weed Control or Suppression in Teff

This label for OpenSky expires and must not be distributed or used in accordance with this SLN registration after December 31, 2027.

Environmental Hazards

Aquatic Organism Advisory: This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Toxic to aquatic organisms and non-target terrestrial plants.

Non-Target Organism Advisory: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Ground Water Advisory: This chemical has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow water. This product is classified as having high potential for reaching surface water via runoff for several weeks after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of pyroxsulam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is expected to occur within 48 hours.

Directions for Use

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the EPA registered label.

Apply 1 pint per acre (0.12 lb/acre fluroxypyr, 0.013 lb/acre pyroxsulam) of OpenSky postemergence to the main flush of actively growing weeds. Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. For best results and to minimize spray drift, apply in a spray volume of 10 gallons per acre or more. Apply to Teff from the tillering to jointing stage (Zadoks scale 21 to 31) of growth according to the weed application timings shown on the federal label in the section entitled Weeds Controlled (C) or Suppressed (S). Best results are obtained when application is made to weeds that are actively growing.

Extreme growing conditions such as drought or temperatures near or below freezing prior to, at, or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Refer to the spray equipment manufacturer's directions for detailed information on nozzle types, arrangement, spacing, and operating height and pressure. Operate equipment at spray pressures no greater than is necessary to produce a uniform spray pattern. Operate the spray boom no higher than is necessary to produce a uniformly overlapping pattern between spray nozzles.

Ground Applications:

Volume: Apply this product in a total spray volume of 10 or more gallons per acre using spray equipment designed to produce large-droplet, low pressure sprays. Spot treatments should be applied only with a calibrated boom to prevent over application.

Restriction:

- Do not apply with hollow cone-type insecticide nozzles or other nozzles that produce a fine-droplet spray.
- For use in accordance with this SLN label **do not** apply this product through aerial application.

Mandatory Spray Drift Management

Ground Boom Applications:

- User must apply with the release height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- Applicators are required to use a Medium or coarser droplet size (ASABE S572.3).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Use Restrictions

- **Preharvest Interval:** Do not apply within 60 days of harvest.
- Do not apply more than 1 pint/acre of OpenSky (0.12 lb/acre fluroxypyr, 0.013 lb/acre pyroxsulam) per application.
- Do not make more than one application of OpenSky per year.
- Do not apply more than 1 pint/acre of OpenSky (0.12 lb/acre fluroxypyr, 0.013 lb/acre pyroxsulam) per year.
- Do not allow livestock to graze the treated crop within 7 days following application.
- Do not cut the treated crop for hay within 28 days following application.
- For crop rotation restrictions follow the directions for wheat on the federal label for OpenSky.
- **Chemigation:** For use in accordance with this SLN label **do not** apply this product through any type of irrigation system.
- For use in accordance with this SLN label **do not** apply this product through aerial application.
- This product is persistent and may be present in plant materials for over 30 days after application. Do not use treated plant material or manure from animals that have grazed or consumed forage from treated areas for compost, mulch, or mushroom spawn until 30 days after application.
- Animals that have been fed Fluroxypyr treated forage must be fed forage free of Fluroxypyr for at least 3 days before they are moved off the treated property.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA website under the Waste Pesticide Program. Cleaned containers may also be disposed of in a sanitary landfill if permitted by the county. Burning is not a legal method of container disposal in Washington.

Special Conditions and Risks of Use

USE OF OpenSky® herbicide (THE “PRODUCT”) ON TEFF (THE “CROP”) MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.

Corteva Agriscience makes the Product available for use in the manner described in this SLN Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this SLN Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this SLN Labeling, you acknowledge and accept that, to the extent consistent with applicable law:

- (1) You assume all risk of Crop injury, Crop yield reduction, and/or Crop loss;
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this SLN Labeling;
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to inability to sell the Crop, Crop injury, Crop yield reduction and/or Crop loss;
- (5) These Special Conditions and Risks of Use supersede any contrary representations or recommendations by Corteva Agriscience, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Corteva Agriscience and not specified or enforced by the U.S. EPA or the State of Washington.

**24(c) Registrant: Corteva Agriscience LLC
9330 Zionsville Road
Indianapolis, IN 46268**

™® Trademarks of Corteva Agriscience and its affiliated companies

R430-007
Approved: 11/12/2025
Initial Printing