

InflameTM

280 SL



Inflame 280 SL is a nonselective herbicide that provides control of a broad spectrum of broadleaf and grassy weeds.

Inflame 280 SL is registered for use as a:

- Burndown treatment prior to planting or prior to emergence of canola, corn, sweet corn*, cotton, soybean, and sugar beet
- Postemergence weed control herbicide to be applied on LibertyLink® or glufosinate-resistant crops including LibertyLink canola, LibertyLink corn, LibertyLink sweet corn*, LibertyLink cotton*, LibertyLink soybeans, and LibertyLink sugar beets
- Postemergence weed control herbicide to be applied in cotton with a hooded sprayer only
- Postemergence weed control herbicide to be applied in listed pome fruit, citrus, grapes, stone fruit, tree nuts, and bushberry crops
- Postemergence weed control herbicide to be applied in olives, avocado*, fig*, and hops*
- Vine desiccant in potatoes

*Not Registered for use by California

ACTIVE INGREDIENT:

Glufosinate-ammonium (CAS No. 77182-82-2) 24.5%

OTHER INGREDIENTS:

TOTAL: 100.0%

Contains 2.34 lbs. of Glufosinate-ammonium per gallon.

EPA Reg. No.: 91234-82

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See inside label booklet for Precautionary Statements and Directions for Use.

FIRST AID

If on skin:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at **1-844-685-9173** for emergency medical treatment information.

For Chemical Emergency Spill, Leak, Fire, Exposure, or Accident Call CHEMTREC Day or Night
Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using toilet. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long sleeved shirt and long pants, socks, shoes;
- Chemical-resistant gloves such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils; chemical resistant footwear plus socks;
- Protective eyewear (goggles, face shield or safety glasses);
- Wear a chemical resistant apron when mixing/loading and cleaning equipment.

Mixers/loaders supporting aerial applications to canola, corn, cotton, and soybean must use closed mixing/loading systems.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water or to areas where surface water is present. **DO NOT** apply to intertidal areas below the mean high water mark. **DO NOT** contaminate water by cleaning of equipment or disposal of equipment wash waters or rinsate.

This product is moderately toxic to bees on a chronic basis, and may cause chronic risk to pollinators or other terrestrial invertebrates. **DO NOT** apply this product to blooming vegetation or if bees or other pollinating insects are visiting the treatment area.

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift and runoff.

This pesticide is toxic to vascular plants and must be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

Under some conditions, this product may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no till, limited till and contour plowing; these methods also reduce pesticide run-off. Use of vegetation filter strips along rivers, creeks, streams, wetlands, etc. or on the downhill side of fields where run-off could occur to minimize water runoff is advised.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT use this product until you have read the entire label. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

In the State of New York Only: Not for Use in Nassau and Suffolk Counties.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses; and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls worn over short-sleeved shirt and short pants;
- chemical resistant gloves including barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils;
- chemical resistant footwear plus socks,
- protective eyewear (goggles, face shield or safety glasses).

IMPORTANT CROP SAFETY INFORMATION READ BEFORE USING THIS PRODUCT

Burndown treatments

Inflame 280 SL may be applied as a burndown treatment prior to planting of canola, corn, sweet corn, cotton, soybean, sugar beets, fruiting vegetables, cucurbits, olives, trees, vines, berries tropical and subtropical fruits listed on this label, **or after planting but prior to emergence** of canola, corn, sweet corn, cotton, soybean, and sugar beet.

Post emergent treatments

Post emergence row crop applications of **Inflame 280 SL** may be made only to crops containing the LibertyLink trait. The basis of selectivity of **Inflame 280 SL** in LibertyLink crops is the presence of a gene that makes crops not sensitive to glufosinate. **Crops not containing the LibertyLink trait will be sensitive to Inflame 280 SL and severe crop injury and/or death may occur. DO NOT allow spray to contact foliage or green tissue of desirable vegetation other than crops not sensitive to the active ingredient in this product.**

Post emergent applications of **Inflame 280 SL** may be applied to conventional or other transgenic cotton sensitive to the active ingredient in **Inflame 280 SL** using a hooded sprayer.

Tree, Nut, Vine and Berry treatments

When applying **Inflame 280 SL** to apples, berries, tree nuts and vines, and tropical and subtropical fruits, avoid contact of solution, spray, drift or mist with green bark, stems or foliage, as injury may occur. Only trunks with calloused, mature brown bark may be sprayed unless protected from spray contact by nonporous wraps, grow tubes or waxed containers. Contact of **Inflame 280 SL** with parts of trees, berries, vines, or tropical and subtropical fruits other than mature brown bark can result in serious damage.

PRODUCT INFORMATION

Inflame 280 SL is a water-soluble non-selective, broad-spectrum herbicide used for control of annual and perennial grass and broadleaf weeds in a variety of crops. Uses include applications as foliar sprays in trees, vines, hops, tropical and subtropical fruits, and berry crops for control of emerged weeds; postemergence weed control herbicide to be applied between rows of fruiting vegetables and cucurbits; broadcast burndown applications prior to planting in labeled conventional row crops; burndown treatment after planting but prior to emergence of canola, corn, sweet corn, cotton, soybean, and sugar beet; and as over-the-top applications in canola, corn, cotton, soybeans and sugar beets designated as LibertyLink® or glufosinate-tolerant. **Inflame 280 SL** may be used for weed control in cotton when applied with a hooded sprayer in-crop.

Inflame 280 SL may also be applied for potato vine desiccation.

It is important to always follow a responsible integrated weed management program. Contact your local agronomic advisor for more specific information on integrated weed management in your area.

ROTATIONAL CROP RESTRICTIONS*

Rotational crop planting intervals following application of **Inflame 280 SL** are listed below. Failure to comply with these restrictions may result in illegal residues in rotated crops.

Rotational Crop	Plant-back Interval (Minimum Rotational Crop Planting Interval from Last Application)
Canola, Sweet Corn, Corn, Cotton, Rice, Soybeans, Sugar Beets, Fruiting Vegetables**, and Cucurbits**	May be planted at any time
Cover Crops***	7 Days
Transplanted Perennial Crops on label (bushberries group 13-07B, citrus group 10-10, olives, pome fruit group 11-10, stone fruit group 12-12, tree nuts group 14-12, fruit, grape (table, wine and raisins), hops, avocado, fig, and tropical and subtropical fruits 23B/24A/24B)	14 Days
Root and Tuber Vegetables, Leafy Vegetables, Brassica Leafy Vegetables, Small Grains (barley, buckwheat, oats, rye, teosinte, triticale, and wheat)	70 Days
All Other Crops	180 Days

*See **Potato** under **CROP USE DIRECTIONS** for rotational crop restrictions specifically after **Inflame 280 SL** applications to potatoes for vine desiccation. See **Sugar Beets** under **CROP USE DIRECTIONS** for rotational crop restrictions specifically for sugar beet.

For in-crop applications for these crops, follow the respective **CROP USE DIRECTIONS section of the label.

***Planting of cover crops for conservation purposes may be planted in fields previously treated with **Inflame 280 SL** as long as these cover crops are not grazed by livestock nor harvested for food. For best results, **DO NOT** plant cover crops less than 7 days after an application of **Inflame 280 SL** nor before 1/2 inch of rainfall or irrigation has occurred. Planting sooner than this may result in stand reduction. Planting of crops listed in the **ROTATIONAL CROP RESTRICTIONS** that follow the listed planting intervals and other restrictions are considered a rotational crop and therefore may be harvested.

WEED RESISTANCE MANAGEMENT

Inflame 280 SL is a Group 10 Herbicide, i.e., a glutamine synthetase inhibitor. Any weed population may contain plants naturally resistant to glufosinate and other Group 10 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies must be followed. Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed. If levels of control provided by applications of this product is reduced and cannot be accounted for by factors such as misapplication, abnormal levels of target species or extremes of weather, it may be the case that target species have developed a strain resistant to applications of **Inflame 280 SL**.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of **Inflame 280 SL** or other Group 10 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Atticus, LLC at 984-465-4800.

WEEDS CONTROLLED

The following weeds controlled charts are outlined by crop or crop group.

Volunteer LibertyLink or glufosinate-tolerant crop plants (corn, cotton, soybeans, sugar beets, canola) from the previous season will not be controlled by applications of **Inflame 280 SL**.

For best results, apply to emerged, small and actively growing weeds less than 3 inches in height. Warm temperatures, high humidity, and bright sunlight improve the performance of **Inflame 280 SL**. Uniform, thorough spray coverage of weeds is necessary to achieve consistent weed control.

Weed control may be reduced when applications are made to weeds under stress including drought or cool temperatures and in dense populations. Stressed conditions may also include prior treatments of other contact or systemic herbicides. Regrowth of weeds may occur due to the weed stage of growth at application, use rate, or environmental conditions at the time of application.

When any of these conditions exist, select a higher rate within the label rate range to improve weed control.

Table 1. Weeds Controlled
(Including glyphosate-, triazine-, PPO-, ALS-, HPPD-, and auxin- resistant biotypes)

Weeds Controlled at 22 to 28 fl. oz./A	
Broadleaf Weeds	
Common Name	Scientific Name
Anoda, spurred	<i>Anoda cristata</i>
Beggarweed, Florida	<i>Desmodium tortuosum</i>
Black medic	<i>Medicago lupulina</i> L.
Blueweed, Texas	<i>Helianthus ciliaris</i> DC.
Buckwheat, wild	<i>Polygonum convolvulus</i>
Buffalobur	<i>Solanum cornutum</i>
Burcucumber	<i>Sicyos angulatus</i>
Canola, volunteer ¹	<i>Brassica</i> spp.
Carpetweed	<i>Mollugo verticillata</i>
Catchweed bedstraw (cleavers)	<i>Galium aparine</i> L.
Chickweed, common	<i>Stellaria media</i>
Cocklebur, common	<i>Xanthium strumarium</i>
Copperleaf, hophornbeam	<i>Acalypha ostryaefolia</i>
Cotton, volunteer ¹	<i>Gossypium</i> spp.
Croton, tropic	<i>Croton glandulosus</i>
Croton, woolly	<i>Croton capitatus</i>
Devil's claw	<i>Proboscidea louisiana</i>
Eclipta	<i>Eclipta alba</i>
Fleabane, annual	<i>Erigeron annuus</i>
Galinsoga, hairy	<i>Galinsoga ciliata</i>
Galinsoga, smallflower	<i>Galinsoga parviflora</i>
Geranium, cutleaf	<i>Geranium dissectum</i> L.
Groundcherry, cutleaf	<i>Physalis angulata</i>
Hempnettle	<i>Galeopsis</i> spp.
Horsenettle, Carolina ²	<i>Solanum carolinense</i>
Jimsonweed	<i>Datura stramonium</i>
Knotweed	<i>Polygonum</i> spp.
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, common	<i>Chenopodium album</i>
Mallow, common	<i>Malva</i> spp.
Mallow, Venice	<i>Hibiscus trionum</i>
Marsh elder, annual	<i>Iva annua</i>
Morningglory, entireleaf	<i>Ipomoea hederacea</i> var. <i>integriuscula</i>

(continued)

Table 1. Weeds Controlled (continued)
(Including glyphosate-, triazine-, PPO-, ALS-, HPPD-, and auxin-resistant biotypes)

Weeds Controlled at 22 to 28 fl. oz. / A (continued)	
Broadleaf Weeds (continued)	
Common Name	Scientific Name
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>
Morningglory, pitted	<i>Ipomoea lacunosa</i>
Morningglory, sharppod	<i>Ipomoea cordatotriloba</i>
Morningglory, smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall	<i>Ipomoea purpurea</i>
Mustard, wild	<i>Sinapis arvensis</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, eastern black	<i>Solanum ptycanthum</i>
Nightshade, hairy	<i>Solanum sarrachoides</i>
Pennycress	<i>Thlaspi arvense</i>
Pigweed, prostrate	<i>Amaranthus blitoides</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Pigweed, spiny	<i>Amaranthus spinosus</i>
Pigweed, tumble	<i>Amaranthus albus</i>
Puncturevine	<i>Tribulus terrestris</i>
Purslane, common	<i>Portulaca oleracea</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Ragweed, giant	<i>Ambrosia trifida</i>
Senna, coffee	<i>Cassia occidentalis</i>
Sesbania, hemp	<i>Sesbania herbacea</i>
Shepherd's purse	<i>Capsella bursa-pastoris</i>
Sicklepod (java bean)	<i>Senna obtusifolia</i>
Sida, prickly	<i>Sida spinosa</i> L.
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Smell melon	<i>Cucumis melo</i> L. var. <i>dudaim</i>
Sowthistle, annual	<i>Sonchus oleraceus</i> L.
Soybeans, volunteer ¹	<i>Glycine max</i>
Spurge, prostrate	<i>Euphorbia humifusa</i>
Spurge, spotted	<i>Euphorbia maculata</i> L.
Starbur, bristly	<i>Acanthospermum hispidum</i>
Sunflower, common	<i>Helianthus annuus</i>
Sunflower, prairie	<i>Corythucha pura</i>

(continued)

Table 1. Weeds Controlled (continued)
(Including glyphosate-, triazine-, PPO-, ALS-, HPPD-, and auxin-resistant biotypes)

Weeds Controlled at 22 to 28 fl. oz. / A (continued)	
Broadleaf Weeds (continued)	
Common Name	Scientific Name
Sunflower, volunteer	<i>Helianthus annuus</i>
Velvetleaf	<i>Abutilon theophrasti</i>

Weeds Controlled at 22 to 28 fl. oz. / A (continued)	
Grass Weeds	
Common Name	Scientific Name
Barley, volunteer ²	<i>Hordeum vulgare</i>
Barnyardgrass	<i>Echinochloa</i> spp.
Bluegrass, annual	<i>Poa annua</i> L.
Corn, volunteer ¹	<i>Zea mays</i> L.
Crabgrass, large ⁴	<i>Digitaria sanguinalis</i>
Crabgrass, smooth ⁴	<i>Digitaria ischaemum</i>
Cupgrass, woolly	<i>Eriochloa villosa</i>
Foxtail, bristly	<i>Setaria verticillata</i>
Foxtail, giant	<i>Setaria faberi</i>
Foxtail, green	<i>Setaria viridis</i>
Foxtail, robust purple	<i>Setaria viridis</i>
Foxtail, yellow ⁴	<i>Setaria pumila</i>
Goosegrass ²	<i>Eleusine indica</i>
Johnsongrass, seedling	<i>Sorghum halepense</i>
Junglerice	<i>Echinochloa colonum</i>
Millet, proso volunteer	<i>Milium vernale</i>
Millet, wild proso	<i>Panicum miliaceum</i> L.
Oat, wild ⁴	<i>Avena fatua</i>
Panicum, fall	<i>Panicum dichotomiflorum</i>
Panicum, Texas	<i>Panicum texanum</i>
Rice, red	<i>Oryza sativa</i> L.
Rice, volunteer ¹	<i>Oryza sativa</i>
Sandbur, field ^{*4}	<i>Cenchrus pauciflorus</i>
Shattercane	<i>Sorghum vulgare</i> Pers.
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
Sorghum, volunteer	<i>Sorghum</i> spp.
Sprangletop	<i>Leptochloa</i> spp.

(continued)

Table 1. Weeds Controlled (continued)
(Including glyphosate-, triazine-, PPO-, ALS-, HPPD-, and auxin- resistant biotypes)

Weeds Controlled at 22 to 28 fl. oz. / A (continued)	
Grass Weeds	
Common Name	Scientific Name
Stinkgrass	<i>Eragrostis ciliaris</i>
Wheat, volunteer ⁴	<i>Triticum</i> spp.
Witchgrass	<i>Panicum virgatum</i> L.

Additional Weeds Controlled at 29 to 43 fl. oz. / A	
Broadleaf Weeds	
Common Name	Scientific Name
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Kochia	<i>Kochia scoparia</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatus</i>
Marestail ³	<i>Conyza canadensis</i>
Pusley, Florida	<i>Richardia scabra</i>
Thistle, Russian ²	<i>Salsola kali</i>

Additional Weeds Controlled at 29 to 43 fl. oz. / A (continued)	
Grass Weeds	
Common Name	Scientific Name
Sandbur, field ⁴	<i>Cenchrus pauciflorus</i>

Additional Weeds Controlled at 29 to 43 fl. oz. / A (continued)	
Biennial and Perennial Weeds	
Common Name	Scientific Name
Alfalfa	<i>Medicago sativa</i> L.
Bermudagrass	<i>Cynodon dactylon</i>
Bindweed, field	<i>Convolvulus arvensis</i> L.
Bindweed, hedge	<i>Calystegia sepium</i>
Bluegrass, Kentucky	<i>Poa pratensis</i> L.
Blueweed, Texas	<i>Helianthus ciliaris</i> DC.
Bromegrass, smooth	<i>Bromus inermis</i>
Burdock	<i>Arctium</i> spp.
Bursage, woollyleaf	<i>Ambrosia grayi</i>
Chickweed, mouse-ear	<i>Cerastium vulgatum</i> L.
Clover, red	<i>Trifolium pratense</i> L.
Dandelion	<i>Taraxacum officinale</i>

(continued)

Table 1. Weeds Controlled (continued)
(Including glyphosate-, triazine-, PPO-, ALS-, HPPD-, and auxin- resistant biotypes)

Additional Weeds Controlled at 29 to 43 fl. oz. / A (continued)	
Biennial and Perennial Weeds (continued)	
Common Name	Scientific Name
Dock, smooth*	<i>Rumex</i> spp.
Dogbane, hemp*	<i>Apocynum cannabinum</i>
Johnsongrass, rhizome	<i>Sorghum halepense</i>
Milkweed, common*	<i>Asclepias syriaca</i>
Milkweed, honeyvine*	<i>Ampelamus albidus</i>
Muhly, wirestem*	<i>Muhlenbergia frondosa</i>
Nightshade, silverleaf	<i>Solanum elaeagnifolium</i>
Nutsedge, purple*	<i>Cyperus rotundus</i>
Nutsedge, yellow*	<i>Cyperus ferax</i>
Orchardgrass	<i>Dactylis glomerata</i> L.
Poinsettia, wild*	<i>Euphorbia heterophylla</i> L.
Pokeweed	<i>Phytolacca</i> L.
Quackgrass	<i>Agropyron repens</i>
Sowthistle, perennial	<i>Sonchus arvensis</i> L.
Thistle, bull*	<i>Cirsium vulgare</i>
Thistle, Canada	<i>Cirsium arvense</i>
Timothy*	<i>Phleum pratense</i> L.
Wormwood, biennial	<i>Artemisia biennis</i>

Additional Weeds Controlled at 48 to 82 fl. oz. / A	
Broadleaf Weeds	
Common Name	Scientific Name
Alkali sida	<i>Sida hederacea</i>
Ammannia, purple	<i>Ammannia robusta</i>
Arrowhead, California	<i>Sagittaria montevidensis</i>
Burclover, California	<i>Medicago polymorpha</i>
Chinese thornapple	<i>Datura quercifolia</i>
Copperleaf, Virginia	<i>Acalypha virginica</i>
Cudweed	<i>Gnaphalium</i> sp.
Cutleaf evening primrose	<i>Oenothera laciniata</i>
Dodder	<i>Cuscuta</i> sp.
Fiddleneck	<i>Amsinckia intermedia</i>
Filaree	<i>Erodium</i> sp.
Filaree, redstem	<i>Erodium cicutarium</i>

(continued)

Table 1. Weeds Controlled (continued)
(Including glyphosate-, triazine-, PPO-, ALS-, HPPD-, and auxin- resistant biotypes)

Additional Weeds Controlled at 48 to 82 fl. oz. / A (continued)	
Broadleaf Weeds (continued)	
Common Name	Scientific Name
Goosefoot	<i>Chenopodium</i> sp.
Gromwell, field	<i>Lithospermum arvense</i>
Groundsel, common	<i>Senecio vulgaris</i>
Henbit	<i>Lamium amplexicaule</i>
Lettuce, miner's	<i>Claytonia perfoliata</i>
Lettuce, prickly	<i>Lactuca serriola</i>
London rocket	<i>Sisymbrium irio</i>
Malva (little mallow)	<i>Malva parviflora</i>
Mayweed	<i>Anthemis cotula</i>
Mullein, turkey	<i>Croton setigerus</i>
Nettle	<i>Urtica</i> sp.
Pineapple-weed	<i>Matricaria discoidea</i>
Radish, wild	<i>Raphanus raphanistrum</i>
Redmaids	<i>Calandrinia ciliata</i>
Starthistle, yellow	<i>Centaurea solstitialis</i>
Swinecress	<i>Lepidium</i> sp.
Turnip, wild	<i>Rapistrum rugosum</i>
Vervain	<i>Verbena</i> sp.
Vetch	<i>Vicia sativa</i>
Willowherb, panicle	<i>Epilobium brachycarpum</i>

Additional Weeds Controlled at 48 to 82 fl. oz. / A (continued)	
Grass Weeds	
Common Name	Scientific Name
Brome, ripgut	<i>Bromus diandrus</i>
Bromegrass, downy	<i>Bromus tectorum</i>
Canarygrass	<i>Phalaris canariensis</i>
Chess, soft	<i>Bromus hordeaceus</i>
Rush, toad*	<i>Juncus bufonius</i>
Ryegrass, annual*	<i>Lolium multiflorum</i> subsp. <i>gaudini</i>
Windgrass	<i>Apera spica-venti</i>

Table 1. Weeds Controlled (continued)
(Including glyphosate-, triazine-, PPO-, ALS-, HPPD-, and auxin- resistant biotypes)

Additional Weeds Controlled at 48 to 82 fl. oz. / A (continued)	
Biennial and Perennial Weeds (continued)	
Common Name	Scientific Name
Aster, white heath	<i>Symphotrichum pilosum</i>
Bluegrass, Kentucky	<i>Poa pratensis</i>
Bulrush*	<i>Scirpus</i> sp.
Clover, Alsike	<i>Trifolium hybridum</i>
Clover, white	<i>Trifolium repens</i>
Dallisgrass	<i>Paspalum dilatatum</i>
Dock, curly	<i>Rumex crispus</i>
Fescue	<i>Festuca</i> sp.
Guineagrass	<i>Megathyrsus maximus</i>
Horsetail	<i>Equisetum</i> sp.
Lovegrass	<i>Eragrostis</i> sp.
Mugwort	<i>Artemisia vulgaris</i>
Mullein, common	<i>Verbascum Thapsus</i>
Mustard, tansy	<i>Descurainia pinnata</i>
Onion, wild	<i>Allium canadense</i>
Orchardgrass	<i>Dactylis glomerata</i>
Paragrass	<i>Urochloa mutica</i>
Plantain	<i>Plantago</i> sp.
Poison ivy	<i>Toxicodendron</i> sp.
Poison oak	<i>Toxicodendron</i> sp.
Rocket, yellow	<i>Barbarea vulgaris</i>
Rose, wild	<i>Rosa multiflora</i>
Rubus spp.	<i>Rubus</i> sp.
Spurge, leafy	<i>Euphorbia esula</i>
Thistle, musk	<i>Carduus nutans</i>
Torpedograss	<i>Panicum repens</i>
Vaseygrass	<i>Paspalum urvillei</i>
Woodsorrel	<i>Oxalis</i> sp.

(continued)

Table 1. Weeds Controlled (continued)
(Including glyphosate-, triazine-, PPO-, ALS-, HPPD-, and auxin-resistant biotypes)

Additional Weeds Controlled at 48 to 82 fl. oz./A (continued)	
Biennial and Perennial Weeds (continued)	
Common Name	Scientific Name
Yarrow, common	<i>Achillea millefolium</i>

* Suppression only.

¹Volunteer LibertyLink® or glufosinate-tolerant crops from the previous season will not be controlled. A timely cultivation 7 to 10 days after an application and/or retreatment 10 to 21 days after the first application can be made for controlling dense clumps of volunteer corn or volunteer rice.

²May require sequential applications for control.

³For optimum control apply **Inflame 280 SL** on 6-inch marestalk.

⁴For best control of yellow foxtail, field sandbur, crabgrass, wild oats, and volunteer wheat, treat prior to tiller initiation.

⁵For control of biennial and perennial weeds, use tank mixes or sequential applications of **Inflame 280 SL**.

Table 2. Weeds Controlled
(Specified Non-Crop Uses)

Broadleaf Weeds	
Chickweed	London Rocket
Clover	Malva (little mallow)
Filaree	Marestail
Jimsonweed	Shepherdspurse
Kochia	Smartweed

Grasses and Sedges	
Annual bluegrass	Kentucky bluegrass
Bahiagrass	Lovegrass
Barley	Nutsedge
Bermudagrass	Paragrass
Barnyardgrass	Quackgrass
Carpetgrass	Ryegrass
Crabgrass	Sandbur
Cupgrass	Shattercane
Dallisgrass	Smallflower Alexandergrass (Signalgrass)
Downy brome	Smooth brome
Fall panicum	Stinkgrass
Fescue	Torpedograss
Giant foxtail	Vaseygrass
Goosegrass	Wheat
Green foxtail	Wild oat

(continued)

Table 2. Weeds Controlled
(Specified Non-Crop Uses)

Grasses and Sedges (continued)	
Guineagrass	Windgrass
Johnsongrass (rhizome)	Yellow foxtail

Broadleaf Weeds	
Annual sowthistle	Musk thistle
Bindweed	Nettle
Buffalobur	Nightshade
Burdock	Pennycress
Canada thistle	Pigweed, redroot
Curly dock	Plantain
Dandelion	Prickly Lettuce
Dogbane (hemp)	Ragweed
Field gromwell	Russian Thistle
Fleabane	Tansy mustard
Goldenrod	Velvetleaf
Horsetail	Vervain
Lambsquarters	Virginia copperleaf
Leafy spurge	White heath aster
Mugwort	Wild buckwheat
Wild mustard	Wild turnip
Wild Onion	Woodsorrel
Wild Rose	Yellow Rocket

Brush* Controlled or Suppressed	
Blackberry	Pine
Deer brush	Roundleaf Vine
Douglas fir	Greenbrier
Gallberry	Salmonberry
Hazel	Sweetgum
Honeysuckle	Sumac
Huckleberry	Thimbleberry
Maple	Trumpet creeper
Multiflora rose	Maple
Oak	Western Red Cedar
Poison ivy/oak	

*Not registered for use by California

APPLICATION INSTRUCTIONS

DO NOT use flood jet nozzles, controlled droplet application equipment, or air-assisted spray equipment. Uniform, thorough spray coverage is important to achieve consistent weed control.

GROUND APPLICATION

Refer to the **WEEDS CONTROLLED** section (Table 1) or **CROP USE DIRECTIONS** for application rates. **DO NOT** apply when winds are gusty, or when conditions favor movement of spray particles off the desired spray target.

Apply early when weeds are small with directed rates as identified in the **WEEDS CONTROLLED** section. Apply **Inflame 280 SL** broadcast in a minimum of 15 gallons of water per acre using a minimum spray pressure of 40 psi and a maximum ground speed of 10 mph.

NOZZLE SELECTION

Apply with nozzles and pressure that deliver medium to coarse spray droplets as indicated in nozzle manufacturer's catalogues and in accordance with ASABE Standard 572.1 unless otherwise mandated by tank mix product. Addition of some drift retardants can significantly increase the droplet size and reduce spray coverage and efficacy. If a drift retardant is used, ensure that it is compatible for use with **Inflame 280 SL** and spray equipment being used. Under dense weed/crop canopies, when covering large weeds, or using larger spray droplets, use a broadcast rate of 20 gallons of water per acre so that thorough spray coverage will be obtained. **DO NOT** use raindrop nozzles. See the **Spray Drift Management** section of this label for additional information on proper application of **Inflame 280 SL**.

AERIAL APPLICATION

Thorough coverage is necessary for best weed control. For optimal weed control, apply **Inflame 280 SL** in a minimum of 10 gallons per acre. See the **Spray Drift Management** section of this label for additional information on proper application of **Inflame 280 SL**.

ADJUVANT INSTRUCTIONS

Ammonium sulfate (AMS) can be used at 1.5 lbs./A to 3 lbs./A. Rates are dependent on tank mix partners, environmental conditions, temperatures and potential for leaf burn. AMS has shown to improve weed control of difficult-to-control weeds, like velvetleaf and lambsquarters, under difficult environmental conditions (low relative humidity) or hard water. Anti-foam agent is advised. No additional surfactant is needed with any tank mix partner. The use of additional surfactants or crop oils may increase the risk of crop response. Please refer to the surfactant label for more detailed information.

Use the **USE RATE EQUIVALENCY FOR INFLAME 280 SL** (Table 3) to determine the corresponding amounts of active ingredient (glufosinate) from **Inflame 280 SL** product use rates.

Table 3. USE RATE EQUIVALENCY FOR INFLAME 280 SL

Amount of Inflame 280 SL (fl. oz./A)	Amount of Glufosinate (lbs. ai/A)
22	0.40
29	0.53
30	0.55
32	0.59
36	0.66
43	0.79
48	0.88

(continued)

Table 3. USE RATE EQUIVALENCY FOR INFLAME 280 SL (continued)

Amount of Inflame 280 SL (fl. oz./A)	Amount of Glufosinate (lbs. ai/A)
49	0.90
55	1.00
56	1.02
58	1.06
60	1.10
62	1.13
64	1.17
72	1.32
82	1.50
87	1.59
165	3.02
246	4.50

MIXING INSTRUCTIONS

COMPATIBILITY TESTING

If **Inflame 280 SL** will be mixed with pesticide products not listed on this label, test the compatibility of the intended tank mixture before mixing the products in the spray tank. The following procedure assumes a spray volume of 25 gallons per acre. For other spray volumes, adjust the amount of the water used accordingly. Check compatibility using this process:

1. In a clear 1-quart jar, place 1.0 pint of water from the source that will be used to prepare the spray solution.
2. For each pound of a dry tank mix partner to be applied per acre, add 1.5 teaspoons to the jar.
3. For each 16 fl. oz. of a liquid tank mix partner to be applied per acre, add 0.5 teaspoon to the jar.
4. For each 16 fl. oz. of **Inflame 280 SL** to be applied per acre, add 0.5 teaspoon to the jar.
5. After adding all the ingredients, place a lid on the jar and tighten, then invert 10 times to mix.
6. Allow the mixture to stand for 15 minutes, then evaluate the solution for uniformity and stability. Look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. If the tank mix partners are not compatible, **DO NOT** use the mixture in a spray tank.
7. Once compatibility testing is complete, dispose of any pesticide wastes in accordance with the **STORAGE AND DISPOSAL** section of this label.

TANK MIX INSTRUCTIONS

Inflame 280 SL may be applied in tank mix combinations with labeled rates of other products provided these other products are labeled for the timing and method of application for the crop to be treated. Use the tank mix partner in accordance with label limitations and restrictions. **DO NOT** exceed label dosage rates.

Inflame 280 SL may not be mixed with any product containing a label prohibition against such mixing. Refer to the specific **CROP USE DIRECTIONS** section for rates and other restrictions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Inflame 280 SL must be applied with properly calibrated and clean equipment. **Inflame 280 SL** is formulated to mix readily in water. Prior to adding **Inflame 280 SL** to the spray tank, ensure that the spray tank is thoroughly clean, particularly if a herbicide with the potential to injure crops was previously used (see **CLEANING INSTRUCTIONS**).

Mix **Inflame 280 SL** with water to make a finished spray solution as follows:

1. Fill the spray tank half full with water.
2. Begin agitation.
3. If mixing with a flowable/wettable powder tank mix partner, prepare a slurry of the proper amount of the product in a small amount of water. Add the slurry to the spray tank.
4. Add the appropriate amount of ammonium sulfate (AMS) to the spray tank.
5. If mixing with a liquid tank mix partner, add the liquid mix partner next.
6. Complete filling the spray tank with water.
7. Add the proper amount of **Inflame 280 SL** and continue agitation.
8. If foaming occurs, use a silicone-based antifoam agent.

Ensure that all spray system lines including pipes, booms, etc. have the correct concentration of spray solution by flushing out the spray system lines before starting the crop application.

If tank mix partners listed on this label are added, maintain good agitation at all times until contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to re-suspend the mixture before spraying is resumed. Keep bypass line on or near bottom of tank to minimize foaming. Screen size in nozzles or line strainers must be 50 mesh or larger.

CLEANING INSTRUCTIONS

Before using **Inflame 280 SL**, thoroughly clean bulk storage tank, refillable tank, nurse tanks, spray tank, lines, and filter, particularly if a herbicide with the potential to injure crops was previously used. Ensure that equipment is thoroughly rinsed using a commercial tank cleaner.

After using **Inflame 280 SL**, triple rinse the spray equipment and clean with a commercial tank cleaner before using for crops not labeled as LibertyLink or glufosinate-tolerant. Make sure any rinsate or foam is thoroughly removed from spray tank and boom. Rinsate may be disposed following the pesticide disposal directions on this label.

TANK MIX PARTNERS

For all crops, certain herbicide tank mixes may aid in the performance of **Inflame 280 SL** or be added to provide residual herbicide activity. When tank mixing with a residual herbicide, no additional surfactant is needed. **Inflame 280 SL** may be applied in tank mix combinations with labeled rates of other products labeled for the timing and method of application for the crop to be treated. The tank mix partner must be used in accordance with the label limitations and restrictions. **DO NOT** exceed label dosage rates. **Inflame 280 SL** may not be mixed with any product containing a label prohibition against such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels

involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

flumioxazin	glyphosate
napropamide	indaziflam
oxyfluorfen	quizalofop-P-ethyl
diuron	terbacil
penoxsulam	norflurazon
saflufenacil	oryzalin
simazine	pendimethalin
clethodim	rimsulfuron

MANDATORY SPRAY DRIFT MITIGATION

- When applying to crops via aerial application equipment, the spray boom must be mounted on the aircraft so as to minimize drift caused by wing tip or rotor blade vortices. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
- When applying to crops via aerial application equipment, applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.
- For aerial applications, **DO NOT** release spray at a height greater than 10 feet above the crop canopy, unless a greater application height is required for pilot safety.
- For aerial applications:
 - o Applicators must select nozzle and pressure that deliver medium to coarse droplets in accordance with American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- For ground applications:
 - o Applicators must select nozzle and pressure that deliver medium to coarse droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Spray at the appropriate boom height based on nozzle selection and nozzle spacing, but **DO NOT** exceed a boom height of 24 inches above target pest or crop canopy. Set boom to lowest effective height over the target pest or crop canopy based on equipment manufacturer's directions. Automated boom height controllers are advised with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for spray drift.
- For non-crop vegetation management ground applications, apply with the nozzle height no more than 4 feet above the ground or target vegetation, unless necessitated by the application equipment. Examples would include roadside, railroad, utility rights of way, forestry and other industrial vegetation management applications where safety or natural barriers obstruct application.

ADVISORY SPRAY DRIFT

POLLINATOR ADVISORY STATEMENT: This product contains an herbicide. Follow all label directions and precautions to minimize potential off-target exposure in order to prevent effects to non-target plants adjacent to the treated site which may serve as habitat or forage for pollinators.

Spray Drift Management

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. **APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!** See **Wind**, **Temperature and Humidity**, and **Temperature Inversions** sections of this label.

Techniques for Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Use the lower spray pressures advised for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. **WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.**
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

Controlling Droplet Size -Aircraft

- **Number of Nozzles** - Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations. **AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.**
- **Nozzle Type** - Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- **Boom Length** - Longer booms increase drift potential. Therefore, a shorter boom length is advised.
- **Application Height** - Application more than 10ft. above the canopy increases the potential for spray drift.

Boom Height

Setting the boom at the lowest referenced height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom must remain level with the crop and have minimal bounce.

Drift Reduction Technology (DRT)

The EPA Drift Reduction Technology (DRT) Program was developed to encourage the manufacture, marketing, and use of spray technologies scientifically verified to significantly reduce pesticide drift. The use of DRTs should result in significantly less pesticide from spray applications drifting and being deposited in areas not targeted by those applications, compared to spray technologies that do not meet the minimum DRT standard. EPA-verified drift reduction technologies (DRTs) and their ratings will be added to the following webpage as they become available: <https://www.epa.gov/reducing-pesticide-drift/epa-verified-and-rated-drift-reduction-technologies>

Wind

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. **AVOID APPLICATIONS DURING GUSTY OR WINDLESS CONDITIONS.** Note: Local terrain can influence wind patterns. Every applicator needs to be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

CROP USE DIRECTIONS

The following tables indicate use patterns, rates, minimum spray volumes, preharvest intervals and other precautions, restrictions and comments specific to each crop. Read and follow directions carefully.

Inflame 280 SL is a foliar active herbicide with little to no soil residual activity. For best results, apply to emerged, young, actively growing weeds. Warm temperatures, high humidity and bright sunlight improves the performance of **Inflame 280 SL**. Necrosis of leaves and young shoots occurs within 2 to 4 days after application under growing conditions.

Weeds that emerge after application will not be controlled. **Inflame 280 SL** will have an effect on weeds that are larger than the specified leaf stage, however, speed of activity and control may be reduced.

Weed control may be reduced if application is made when heavy dew, fog, mist or rain are present or when weeds are under stress due to drought, cool temperatures, or extended periods of cloudiness.

When applying for control of lambsquarters and velvetleaf, make applications between dawn and 2 hours before sunset to avoid the possibility of reduced control.

The addition of ammonium sulfate may improve weed control if weeds are under stress. For optimal yield, early season weed removal is important.

To maximize weed control, **DO NOT** cultivate from 5 days before an application to 7 days after an application.

Refer to **WEEDS CONTROLLED Table 1** for proper application rate based upon the weeds present. Refer to **APPLICATION INSTRUCTIONS** section for additional information on tank mixes, adjuvants and applications.

Inflame 280 SL is rainfast 4 hours after application; therefore rainfall within 4 hours may necessitate retreatment. Refer to specific use **CROP USE DIRECTIONS** sections of this label for minimum intervals required before reapplication of this product and use rates.

Consult your local Cooperative Extension Service for guidelines on optimum application timing for **Inflame 280 SL** in your region.

COTTON

Application Methods For Cotton

Application of **Inflame 280 SL** to cotton varieties not labeled as LibertyLink or glufosinate-tolerant requires the use of hooded spray equipment designed to minimize exposure of the spray to the cotton stand. A hooded sprayer directs the spray onto weeds, while shielding the cotton stand from contact. Use nozzles that provide uniform coverage within the treated area. Keep hoods on these sprayers adjusted to protect desirable vegetation. Extreme care must be exercised to avoid exposure of the desirable vegetation to the spray.

With a hooded sprayer, the spray pattern is completely enclosed on the top and all 4 sides by a hood. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground as this may cause spray particles to escape and come into contact with the cotton, causing damage or destruction of the crop.

Herbicide rates and spray volume instructions are presented as broadcast equivalents and must be reduced in proportion to the area actually treated. Use the following formulas to calculate the correct rate and volume per planted (field) acre:

<u>Band Width in Inches</u>	X	Broadcast Rate	=	Amount of Banded Product
<u>Row Width in Inches</u>		per Acre		Needed per Acre
<u>Band Width in Inches</u>	X	Broadcast Spray	=	Banded Spray Volume
<u>Row Width in Inches</u>		Volume per Acre		needed Per Acre

Use Pattern	Rate (Fl. Oz./Acre)	Directions
Burndown (Prior to Planting or Prior to Crop Emergence)	29.0 - 43.0 (0.53 - 0.79 lb. ai)	Apply to emerged, young, actively growing weeds up to early bloom. Uniform, thorough spray coverage is necessary to achieve consistent weed control.
In-Season (Post Emergent to the Crop)		When applying In-Season to non-LibertyLink® or non glufosinate-resistant cotton, a hooded sprayer designed to minimize exposure of the spray to the cotton stand must be used. Refer to Application Methods for Cotton section.
Post harvest Burndown (After Cotton Harvest)	Up to 43.0 (0.79 lb. ai)	Post Emergent Application: apply from crop emergence to early bloom stage. Severe injury or death may result if the Inflame 280 SL contacts the foliage or stems of cotton NOT labeled as LibertyLink or glufosinate-tolerant.
Seed Propagation	43.0 (0.79 lb. ai)	Apply as a foliar spray to selectively eliminate cotton plants that do not carry the LibertyLink trait or not glufosinate-tolerant. Can be applied to remove susceptible segregates during cotton seed propagation. Breeding material not possessing the LibertyLink gene or that are not glufosinate-tolerant will be severely injured or killed if treated with this herbicide.

Restrictions:

- **DO NOT** apply more than 43.0 fl. oz./A (0.79 lb. ai/A) in one application.
- **DO NOT** apply more than three applications per year when using reduced rates.
- **DO NOT** apply more than 87.0 fl. oz./A (1.59 lbs. ai/A) through any combination of use patterns per year.
- **DO NOT** apply to Cotton labeled as LibertyLink or glufosinate-tolerant in Florida, South of Tampa (Florida Route 60), or in Hawaii, except for test plots or breeding nurseries.
- **DO NOT** apply within 70 days of harvest.
- **DO NOT** apply through any type of irrigation system.
- **Burndown:**
 - o **DO NOT** make more than 1 application per year.
- **In-crop and Seed Propagation:**
 - o Applications must be a minimum of 5 days apart.
 - o For in-crop applications to cotton labeled as LibertyLink* or glufosinate-tolerant, allow a minimum of 10 days between applications if additional herbicides or acephate is included in the application.
 - o **DO NOT** make more than 2 applications per year for Seed Propagation.

CORN

Crop	Use Pattern	Rate (Fl. Oz./Acre)	Directions ¹
Corn Field Silage	Burndown (Prior to Planting or Prior to Crop Emergence)	29.0 - 43.0 (0.53 - 0.79 lb. ai)	Apply to emerged, young, actively growing weeds. Uniform, thorough spray coverage is necessary to achieve consistent weed control. Post Emergent Application: apply broadcast or with drop nozzles from emergence up to 24" tall or in the V6 stage of growth whichever comes first.
	In-Season to LibertyLink or glufosinate-tolerant Corn Only (Post Emergent to the Crop)	29.0 - 43.0 (0.53 - 0.79 lb. ai)	For corn 24" to 36" tall, apply using drop nozzles or ground application and nozzles and avoid spraying into the whorl or leaf axils of the corn stalks. Must be applied with ammonium sulfate (AMS). A second In-Season application may be needed to control weeds that have not yet emerged at time of application.
	Seed Propagation	22.0 (0.40 lb. ai)	Apply Inflame 280 SL plus AMS at 3 lbs./A (17 lbs./100 gals) when corn is in the V-3 to V-4 stage of growth, i.e., 3 - 4 developed collars. Make a second treatment of Inflame 280 SL plus AMS at 3 lbs./A when the corn is in the V-6 to V-7 stage of growth or up to 24" tall. The rate of AMS can be reduced to 1.5 lbs./A (8.5 lbs./100 gals) when temperatures exceed 85°F to reduce potential leaf burn. Inbred lines (plants not possessing the LibertyLink trait or not glufosinate-tolerant) will be severely injured or killed if treated with this herbicide. Hooded sprayer may be used to protect plants from coming into contact with the herbicide application.

Restrictions:

- **DO NOT** apply more than 43.0 fl. oz./A (0.79 lb. ai/A) per application.
- If used as a burndown application up to two In-Season applications may be applied.
- **DO NOT** apply more than 3 applications when using reduced rates including burndown use per year.
- **DO NOT** apply more than one burndown application per year.
- **DO NOT** apply more than 87.0 fl. oz./A (1.59 lbs. ai/A) through any combination of use patterns per year.
- Retreatment interval is 7 days.
- **DO NOT** apply with 60 days of harvesting corn forage, and within 70 days of harvesting corn grain or corn fodder.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** use nitrogen solutions as spray carriers. A silicone based anti foam agent may be added if needed.
- **DO NOT** apply if corn shows injury from environmental stress or prior herbicide applications.
- **Seed Propagation:**
 - o **DO NOT** apply more than 22.0 fl. oz./A (0.40 lb. ai/A) per application.
 - o **DO NOT** apply more than 2 applications per year.
 - o **DO NOT** apply more than 44 fl. oz./A (0.80 lb. ai/A) per year.
 - o Retreatment interval is 10 days.

Crop	Use Pattern	Rate (Fl. Oz./Acre)	Directions ¹
Corn, Sweet*	Burndown (Prior to Planting or Prior to Crop Emergence)	29.0 - 43.0 (0.53 - 0.79 lb. ai)	Apply to emerged, young, actively growing weeds. Uniform, thorough spray coverage is necessary to achieve consistent weed control. Post Emergent Application: apply from emergence up to 24" tall or in the V6 stage of growth whichever comes first.
	In-Season to LibertyLink or glufosinate-tolerant Sweet Corn Only (Post Emergent to the Crop)	22.0 (0.40 lb. ai)	A second In-Season application may be needed to control weeds that have not yet emerged at time of application.
	Seed Propagation	22.0 (0.40 lb. ai)	Apply Inflame 280 SL plus AMS at 3 lbs./A (17 lbs./100 gals) when corn is in the V-3 to V-4 stage of growth, i.e., 3 - 4 developed collars. Make a second treatment of Inflame 280 SL plus AMS at 3 lbs./A when the corn is in the V-6 to V-7 stage of growth or up to 24" tall. The rate of AMS can be reduced to 1.5 lbs./A (8.5 lbs./100 gals) when temperatures exceed 85°F to reduce potential leaf burn. Inbred lines (plants not possessing the LibertyLink trait or not glufosinate-tolerant) will be severely injured or killed if treated with this herbicide. Hooded sprayer may be used to protect plants from coming into contact with the herbicide application.

Restrictions:

- **DO NOT** apply more than 44.0 fl. oz./A (0.80 lb. ai/A) through any combination of use patterns per year.
- **DO NOT** apply within 50 days of harvesting sweet corn ears and within 55 days of harvesting stover.
- **DO NOT** apply through any type of irrigation system.
- If used as a burndown application no In-Season applications may be applied.
- **DO NOT** apply more than 2 applications per year.
- **DO NOT** use nitrogen solutions as spray carriers. A silicone based anti-foam agent may be added if needed.
- **DO NOT** apply if corn shows injury from environmental stress or prior herbicide applications.
- Must be applied with ammonium sulfate (AMS).
- **Burndown:**
 - o **DO NOT** apply more than 43.0 fl. oz./A (0.79 lb. ai/A) per application.
 - o **DO NOT** more than one burndown application per year.
- **In-Season:**
 - o **DO NOT** apply more than 22.0 fl. oz./A (0.40 lb. ai/A) per application.
 - o Applications must be at least 7 days apart.
- **Seed Propagation:**
 - o **DO NOT** apply more than 22.0 fl. oz./A (0.40 lb. ai/A) per application.
 - o Retreatment interval is 10 days.

*Not Registered for Use by California

¹Refer to **APPLICATION INSTRUCTIONS** and **Tank Mix Partners for Inflame 280 SL on LibertyLink or Glufosinate-Tolerant Corn** sections for additional information on tank mixes.

Tank Mix Partners for Inflamm 280 SL on LibertyLink or Glufosinate-Tolerant Corn

2,4-D	tembotrione + thiencarbazone-methyl	tembotrione	Pendimethalin ¹	dicamba, sodium salt + halosulfuron-methyl
acetochlor	dicamba, sodium salt + diflufenzopyr-sodium	atrazine + mesotrione + s-metolachlor ²	halosulfuron-methyl	mesotrione + s-metolachlor ²
carfentrazone-ethyl	atrazine + dimethenamid-P	atrazine + mesotrione + s-metolachlor ²	flumetsulam	
atrazine	glyphosate + mesotrione + s-metolachlor	Metolachlor ²	s-metolachlor ²	
mesotrione	flumetsulam + clopyralid potassium	nicosulfuron	primisulfuron-methyl + prosulfuron	
mesotrione + s-metolachlor ²	topramezone	dicamba, sodium salt + primisulfuron-methyl	dicamba, sodium salt + diflufenzopyr-sodium	

¹Tank mixing with pendimethalin may result in reduced control of barnyardgrass, fall panicum, field sandbur, yellow foxtail, and volunteer corn.

²For best results refer to the product label when tank mixing these products.

SPECIMEN

CANOLA, GOLD OF PLEASURE (CAMELINA)

Use Pattern	Rate (Fl. Oz./Acre)	Directions ¹
Burndown (Prior to Planting or Prior to Crop Emergence)	29.0 - 43.0 (0.53 - 0.79 lb. ai)	Apply to emerged, young, actively growing weeds. Uniform, thorough spray coverage is necessary to achieve consistent weed control. Up to 2 additional in-crop applications to LibertyLink or glufosinate-tolerant canola at 22.0 to 29.0 fl. oz./A may be made if a burndown application is made.
In-Season to LibertyLink or glufosinate-tolerant Canola or Gold of Pleasure (Camelina) Only (Post Emergent to the Crop)	22.0 - 29.0 (0.40 - 0.53 lb. ai)	Post Emergent Application: apply from cotyledon stage up to early bolting stage. Slight discoloration of the canola may be visible after application. This effect is temporary and will not influence crop growth, maturity, or yield. May be applied with feed grade or spray grade ammonium sulfate (AMS) at 1.5 - 3 lbs./A. Additional surfactants or crop oils may increase risk of crop response. Up to 58 fl. oz./A (1.06 lbs. ai/A) may be used per year if no burndown application was used. If a burndown application was used, the maximum per year is 87 fl. oz./A. A second In-Season application may be needed to control weeds that have not yet emerged at time of application.
Seed Propagation	29.0 (0.53 lbs ai)	Foliar Spray: Make up to 3 applications from the cotyledon stage up to the early bolting stage (e.g., BBCH 18-30, between just prior to stem elongation/bolting, eight or more leaves and beginning of stem elongation, no internodes). May be used to selectively eliminate canola plants that do not carry the LibertyLink gene or are not glufosinate-tolerant. Can be applied to remove susceptible segregate during canola seed propagation. Breeding material not possessing the LibertyLink gene or that are not glufosinate-tolerant will be severely injured or killed if treated with this herbicide.

Restrictions:

- **DO NOT** apply more than 3 applications including burndown per year.
- **DO NOT** apply more than 87 fl. oz./A (1.59 lbs. ai/A) through any combination of use patterns per year.
- **DO NOT** use on LibertyLink or glufosinate-resistant canola in states of AL, DE, GA, KY, MD, NJ, NC, SC, TN, VA, WV.
- **DO NOT** apply within 65 days of harvest.
- **DO NOT** graze the treated crop or cut for hay.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** apply if canola shows injury from environmental stress or prior herbicide applications.
- **Burndown:**
 - **DO NOT** apply more than 43.0 fl. oz./A (0.79 lb. ai/A) per application.
 - **DO NOT** apply more than 1 application per year.
- **In-Season and Seed Propagation:**
 - **DO NOT** apply more than 29.0 fl. oz./A (0.53 lb. ai/A) per application.
 - Applications must be at least 7 days apart.
 - For Seed Propagation, **DO NOT** make more than 3 applications per year.
 - **DO NOT** apply beyond the early bolting stage for Seed Propagation.
 - **DO NOT** use Seed Propagation treated canola seeds for food, feed or oil purposes.

*Not Registered for Use by California

¹Refer to **APPLICATION INSTRUCTIONS** and **Tank Mix Partners for Inflamm 280 SL on LibertyLink or Glufosinate-Tolerant Canola** sections for additional information on tank mixes, adjuvants and applications.

Tank Mix Partners for Inflamm 280 SL on LibertyLink or Glufosinate-Tolerant Canola

Tank Mix Partner	Rate (fl. oz./A)
quizalofop-p-ethyl	Refer to tank mix partner product label
sethoxydim	
Clethodim (26.4%)	
Clethodim (12.6%)	



SOYBEAN

Use Pattern	Rate (Fl. Oz./Acre)	Directions
Burndown (Prior to Planting or Prior to Crop Emergence)	29.0 - 43.0 (0.53 - 0.79 lb. ai)	Apply to emerged, young, actively growing weeds. Time application at emergence up to bloom or R1 growth stage of soybeans. Uniform, thorough spray coverage is necessary to achieve consistent weed control. A silicone-based antifoam agent may be added if needed. Post Emergent Application: apply from crop emergence up to but not including bloom stage.
In-Season to LibertyLink or glufosinate-tolerant Soybeans Only (Post Emergent to the Crop)		
Seed Propagation	43.0 (0.79 lb. ai)	Apply as a foliar spray to selectively eliminate soybean plants that do not carry a gene that imparts resistance to glufosinate-ammonium. Can be applied to remove susceptible segregates during soybean seed propagation. Soybeans not possessing the glufosinate-ammonium resistance gene will be severely injured or killed if treated with this herbicide.

Restrictions:

- **DO NOT** apply more than 43.0 fl. oz./A (0.79 lb. ai/A) per application.
- **DO NOT** make more than three* applications per year when using reduced rates.
- **DO NOT** make more than 1 burndown application per year.
- **DO NOT** apply more than 2 applications per year for Seed Propagation.
- **DO NOT** apply more than 87.0 fl. oz./A (1.59 lbs. ai/A) through any combination of use patterns per year.
- Make sequential applications at least 5 days apart.
- **DO NOT** apply within 70 days of harvesting soybean seed.
- **DO NOT** graze the treated crop or cut for hay.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** use nitrogen solutions as spray carriers.
- **DO NOT** apply if soybeans show injury from environmental stress or prior herbicide applications.

*Not Registered for Use by California

SPECIMEN

SUGARBEETS

Use Pattern	Rate (Fl. Oz./Acre)	Directions
Burndown (Prior to Planting or Prior to Crop Emergence)	29.0 - 36.0 (0.53 - 0.66lb. ai)	Apply to emerged, young, actively growing weeds. Time application at cotyledon up to 10 leaf stage of LibertyLink or glufosinate-resistant sugar beets.
In-Season to LibertyLink or glufosinate-tolerant Sugar Beets Only (Post Emergent to the Crop)	29.0 (0.53lb. ai)	For best control application must begin when weeds are up to 1 inch in height or diameter. Repeat applications when newly germinated weeds again reach 1 inch in height or diameter. Uniform, thorough spray coverage is necessary to achieve consistent weed control. Post Emergent Application: apply from cotyledon stage up to 10 leaf stage of sugar beet. Anti foams or drift control agents may be added if needed. A second In-Season application may be needed to control weeds that have not yet emerged at time of application.

Restrictions:

- **DO NOT** apply more than 60.0 fl. oz./A (1.10 lbs. ai/A) through any combination of use patterns per year.
- **DO NOT** make more than 2 applications per year.
- **DO NOT** make more than 1 in-season application if burndown application is made.
- **DO NOT** apply within 60 days of harvesting sugar beets.
- **DO NOT** plant rotation crops in a field treated with **Inflame 280 SL** within 120 days after the last application of this product with the exception of wheat, barley, buckwheat, millet, oats, rye, sorghum, and triticale, which may be planted 70 days after the last application of this product. Canola, corn, soybeans, and sugar beets resistant to glufosinate may be planted at any time.
- **DO NOT** graze the treated crop or cut for hay.
- **DO NOT** apply product through any type of irrigation system.
- **DO NOT** add surfactants.
- **DO NOT** apply if sugar beets show injury from environmental stress or prior herbicide applications.
- Applications must be a minimum of 10 days apart.
- **Burndown:**
 - o **DO NOT** apply more than 36.0 fl. oz./A (0.66 lb. ai/A) per application.
 - o **DO NOT** make more than one burndown application per year.
- **In-Season:**
 - o **DO NOT** apply more than 29.0 fl. oz./A (0.53 lb. ai/A) per application.

TREE, NUT, VINE AND BERRIES

Application Methods for Tree, Nut, Vine, and Berries

Refer to **APPLICATION INSTRUCTIONS** section for additional information on tank mixes, adjuvants and applications.

Banded Spray Applications

Banded applications may be used using the following formula to calculate the amount of herbicide needed for orchard or vineyard strip sprays:

$$\frac{\text{Band Width in Inches}}{\text{Row Width in Inches}} \times \frac{\text{Rate per Acre}}{\text{Broadcast}} = \text{Amount of Herbicide Needed for Treatment}$$

Spot or Directed Spray Applications

For spot or directed spray applications mix **Inflame 280 SL** at 1.7 fl. oz. (0.03 lb. ai) of product per gallon of water. Apply to undesirable vegetation foliage until wet but prior to runoff. Ensure uniform and complete coverage. Thoroughly clean the sprayer following use. **DO NOT** make spot or directed spray applications to tree or vine trunk as injury may occur.

Crop	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Pome Fruit (Crop Group 11-10) Apples Crabapple Loquat Mayhaw Quince Pear Oriental Pear Azarole Medlar Tejocote Cultivars, varieties and/or hybrids of these	Broadcast Banded Directed Spray Spot Treatments	Weeds < 3" in Height: 48.0 (0.88 lb. ai) Weeds < 6" in Height: 49.0 - 56.0 (0.90 - 1.02 lbs. ai) Weeds > 6" in Height and/or Grasses that have Tillered: 56.0 - 82.0 (1.02 - 1.50 lbs. ai)	Apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Inflame 280 SL . Uniform, thorough spray coverage is necessary to achieve consistent weed control. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. DO NOT retreat these weeds with Inflame 280 SL until sufficient regrowth has occurred. Avoid direct spray, drift or mist to desirable vegetation, green bark, stems or foliage, as injury may occur.
	Sucker Control Broadcast	56.0 (1.02 lbs. ai)	When applied to suckers that are young, green, and uncalled, Inflame 280 SL will reduce or eliminate sucker growth. Make a split application approximately 4 weeks apart. Thorough coverage of all sucker foliage is necessary for optimum control.

Restrictions:

- **DO NOT** apply more than 82 fl. oz./A (1.50 lbs. ai/A) per application.
- **DO NOT** make more than 5 applications per year at reduced application rates.
- **DO NOT** apply more than 246 fl. oz./A (4.5 lbs. ai/A) through any combination of use patterns per year.
- Applications must be a minimum of 14 days apart.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** graze, harvest and/or feed treated orchard cover crops to livestock.
- **DO NOT** aerially apply.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** allow spray to contact trunks other than those that have callused, mature brown bark or are protected from spray contact by nonporous wraps, grow tubes, or waxed containers.
- **Spot Treatment:**
 - o **DO NOT** exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application.
 - o **DO NOT** exceed the equivalent of 5.65 fl. oz. (0.10 lb. ai) per 1,000 sq. ft. per year.
- **Sucker Control:**
 - o Suckers must not exceed 12 inches in length.
 - o **DO NOT** make spot or directed-spray applications to suckers as tree injury may occur.

¹See **Application Methods for Tree, Nut, Vine, and Berries** section for additional information on banded, directed spray and spot treatment applications.

Crop	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Citrus (Crop Group 10-10) Calamondin Citrus citron Citrus hybrids (chironja, tangelo, tangor) Grapefruit Kumquat Lemon Lime Mandarin (tangerine) Orange (sour, sweet) Pummelo Satsuma mandarin Cultivars, varieties and/or hybrids of these)	Broadcast Banded Directed Spray Spot Treatments	Weeds < 3" in Height: 48.0 (0.88 lb. ai) Weeds < 6" in Height: 49.0 - 56.0 (0.90 - 1.02 lbs. ai) Weeds > 6" in Height and/or Grasses that have Tillered: 56.0 - 82.0 (1.02 - 1.50 lbs. ai)	Apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Inflame 280 SL . Uniform, thorough spray coverage is necessary to achieve consistent weed control. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. DO NOT retreat these weeds with Inflame 280 SL until sufficient regrowth has occurred. Avoid direct spray, drift or mist to desirable vegetation, green bark, stems or foliage, as injury may occur.
	Sucker Control Broadcast	56.0 (1.02 lbs. ai)	When applied to suckers that are young, green, and uncalled, Inflame 280 SL will reduce or eliminate sucker growth. Make a split application approximately 4 weeks apart. Thorough coverage of all sucker foliage is necessary for optimum control.

Restrictions:

- **DO NOT** apply more than 82 fl. oz./A (1.50 lbs. ai/A) per application.
- **DO NOT** make more than 5 applications per year at reduced application rates.
- **DO NOT** apply more than 246 fl. oz./A (4.5 lbs. ai/A) through any combination of use patterns per year.
- Applications must be a minimum of 14 days apart.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** graze, harvest and/or feed treated orchard cover crops to livestock.
- **DO NOT** aerially apply.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** allow spray to contact trunks other than those that have callused, mature brown bark or are protected from spray contact by nonporous wraps, grow tubes, or waxed containers.
- **Spot Treatment:**
 - o **DO NOT** exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application.
 - o **DO NOT** exceed the equivalent of 5.65 fl. oz. (0.10 lb. ai) per 1,000 sq. ft. per year.
- **Sucker Control:**
 - o Suckers must not exceed 12 inches in length.
 - o **DO NOT** make spot or directed-spray applications to suckers as tree injury may occur.

¹See **Application Methods for Tree, Nut, Vine, and Berries** section for additional information on banded, directed spray and spot treatment applications.

Crop	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
GRAPES Table, Wine, Raisin	Broadcast Banded Directed Spray Spot Treatments	Weeds < 3" in Height: 48.0 (0.88 lb. ai) Weeds < 6" in Height: 49.0 - 56.0 (0.90 - 1.02 lbs. ai) Weeds > 6" in Height and/or Grasses that have Tillered: 56.0 - 82.0 (1.02 - 1.50 lbs. ai)	Apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Inflame 280 SL . Uniform, thorough spray coverage is necessary to achieve consistent weed control. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. DO NOT retreat these weeds with Inflame 280 SL until sufficient regrowth has occurred. Avoid direct spray, drift or mist to desirable vegetation, green bark, stems, or foliage as injury may occur. Only trunks with callused, mature brown bark may be sprayed unless protected from spray contact by nonporous wraps, grow tubes, or waxed containers.
	Sucker Control Broadcast	56.0 (1.02 lbs. ai)	When applied to suckers that are young, green, and uncalled, Inflame 280 SL will reduce or eliminate sucker growth. Make a split application approximately 4 weeks apart. Thorough coverage of all sucker foliage is necessary for optimum control.
Restrictions: • DO NOT apply more than 82 fl. oz./A (1.50 lbs. ai/A) per application. • DO NOT make more than 5 applications per year at reduced application rates. • DO NOT apply more than 246 fl. oz./A (4.5 lbs. ai/A) through any combination of use patterns per year. • Applications must be a minimum of 28 days apart. • DO NOT apply within 14 days of harvest. • DO NOT aerially apply. • DO NOT apply through any type of irrigation system. • Spot Treatment: ◦ DO NOT exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application. ◦ DO NOT exceed the equivalent of 5.65 fl. oz. (0.10 lb. ai) per 1,000 sq. ft. per year. • Sucker Control: ◦ Suckers must not exceed 12 inches in length. ◦ DO NOT make spot or directed-spray applications to suckers or vine trunk as injury may occur.			

¹See **Application Methods for Tree, Nut, Vine, and Berries** section for additional information on banded, directed spray and spot treatment applications.

Crop	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Stone Fruit (Crop Group 12-12) Apricot Cherry (sweet, tart) Nectarine Peach Plum (chickasaw, damson, Japanese) Capulin* Jujube* Sloe* Plumcot Prune (fresh)	Broadcast Banded Directed Spray Spot Treatments	Weeds < 3" in Height: 48.0 (0.88 lb. ai) Weeds < 6" in Height: 49.0 - 56.0 (0.90 - 1.02 lbs. ai) Weeds > 6" in Height and/or Grasses that have Tillered: 56.0 - 82.0 (1.02 - 1.50 lbs. ai)	Apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Inflame 280 SL . Uniform, thorough spray coverage is necessary to achieve consistent weed control. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. DO NOT retreat these weeds with Inflame 280 SL until sufficient regrowth has occurred. Avoid direct spray, drift or mist to desirable vegetation, green bark, stems, or foliage as injury may occur.
	Sucker Control Broadcast	56.0 (1.02 lbs. ai)	When applied to suckers that are young, green, and uncalled, Inflame 280 SL will reduce or eliminate sucker growth. Make a split application approximately 4 weeks apart. Thorough coverage of all sucker foliage is necessary for optimum control.

Restrictions:

- **DO NOT** apply more than 82 fl. oz./A (1.50 lbs. ai/A) per application.
- **DO NOT** make more than 2 applications per year.
- **DO NOT** apply more than 164 fl. oz./A (3.0 lb. ai/A) through any combination of use patterns per year.
- Applications must be a minimum of 28 days apart.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** graze, harvest and/or feed treated orchard cover crops to livestock.
- **DO NOT** aerially apply.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** allow spray to contact trunks other than those that have callused, mature brown bark or are protected from spray contact by nonporous wraps, grow tubes, or waxed containers.
- **Spot Treatment:**
 - o **DO NOT** exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application.
 - o **DO NOT** exceed the equivalent of 5.65 fl. oz. (0.10 lb. ai) per 1,000 sq. ft. per year.
- **Sucker Control:**
 - o Suckers must not exceed 12 inches in length.
 - o **DO NOT** make spot or directed-spray applications to suckers as tree injury may occur.

*Not Registered for use by California

¹See **Application Methods for Tree, Nut, Vine, and Berries** section for additional information on banded, directed spray and spot treatment applications.

Crop	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Tree Nuts (Crop Group 14-12) (including Pistachio) Almond Beech nut Brazil nut Butternut Cashew Chestnut Chinquapin Filbert (hazelnut) Hickory nut Macadamia (bush nut) Pecan Pistachio Walnut (black and English (Persian))	Broadcast Banded Directed Spray Spot Treatments	Weeds < 3" in Height: 48.0 (0.88 lb. ai) Weeds < 6" in Height: 49.0 - 56.0 (0.90 - 1.02 lbs. ai) Weeds > 6" in Height and/or Grasses that have Tillered: 56.0 - 82.0 (1.02 - 1.50 lbs. ai)	Apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Inflame 280 SL . Uniform, thorough spray coverage is necessary to achieve consistent weed control. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. DO NOT retreat these weeds with Inflame 280 SL until sufficient regrowth has occurred. Avoid direct spray, drift or mist to desirable vegetation, green bark, stems, or foliage, as injury may occur.
	Sucker Control Broadcast	56.0 (1.02 lbs. ai)	When applied to suckers that are young, green, and uncalled, Inflame 280 SL will reduce or eliminate sucker growth. Make a split application approximately 4 weeks apart. Thorough coverage of all sucker foliage is necessary for optimum control.

Restrictions:

- **DO NOT** apply more than 82 fl. oz./A (1.50 lbs. ai/A) per application.
- **DO NOT** make more than 5 applications per year at reduced application rates.
- **DO NOT** apply more than 246 fl. oz./A (4.5 lbs. ai/A) through any combination of use patterns per year.
- Applications must be at least 28 days apart.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** graze, harvest and/or feed treated orchard cover crops to livestock.
- **DO NOT** aerially apply.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** allow spray to contact trunks other than those that have callused, mature brown bark or are protected from spray contact by nonporous wraps, grow tubes, or waxed containers.
- **Spot Treatment:**
 - o **DO NOT** exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application.
 - o **DO NOT** exceed the equivalent of 5.65 fl. oz. (0.10 lb. ai) per 1,000 sq. ft. per year.
- **Sucker Control:**
 - o Suckers must not exceed 12 inches in length.
 - o **DO NOT** make spot or directed-spray applications to suckers as tree injury may occur.

¹See **Application Methods for Tree, Nut, Vine, and Berries** section for additional information on banded, directed spray and spot treatment applications.

Crop	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Bushberry Subgroup 13-07B Aronia berry* Blueberry, highbush and lowbush Buffalo currant* Chilean guava* Cranberry, highbush* Currant (black and red) Elderberry European barberry* Gooseberry Honeysuckle, edible* Huckleberry Jostaberry* Lingonberry Juneberry Native currant* Salal Seabuckthorn* Cultivars, varieties, and/or hybrids of these.*	Broadcast Banded Directed Spray Spot Treatments	Weeds < 3" in Height: 48.0 (0.88 lb. ai) Weeds < 6" in Height: 49.0 - 56.0 (0.90 - 1.02 lbs. ai) Weeds > 6" in Height and/or Grasses that have Tillered: 56.0 - 82.0 (1.02 - 1.50 lbs. ai)	Apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Inflame 280 SL . Uniform, thorough spray coverage is necessary to achieve consistent weed control. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. DO NOT retreat these weeds with Inflame 280 SL until sufficient regrowth has occurred. Avoid direct spray, drift or mist to desirable vegetation, green bark, stems, or foliage, as injury may occur.
	Sucker Control Broadcast	56.0 (1.02 lbs. ai)	When applied to suckers that are young, green, and uncalled, Inflame 280 SL will reduce or eliminate sucker growth. Make a split application approximately 4 weeks apart. Thorough coverage of all sucker foliage is necessary for optimum control.

Restrictions:

- **DO NOT** apply more than 82 fl. oz./A (1.50 lbs. ai/A) per application.
- **DO NOT** make more than 3 applications per year at reduced application rates.
- **DO NOT** apply more than 164 fl. oz./A (3.0 lbs. ai/A) through any combination of use patterns per year.
- Applications must be a minimum of 28 days apart.
- **DO NOT** apply within 14 days of harvest.
- **DO NOT** aerially apply.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** allow spray to contact trunks other than those that have callused, mature brown bark or are protected from spray contact by nonporous wraps, grow tubes, or waxed containers.
- **Spot Treatment:**
 - o **DO NOT** exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application.
 - o **DO NOT** exceed the equivalent of 5.65 fl. oz. (0.10 lb. ai) per 1,000 sq. ft. per year.
 - o **DO NOT** make spot applications to trunk as injury may occur.
- **Sucker Control:**
 - o Suckers must not exceed 12 inches in length.
 - o **DO NOT** make spot or directed-spray applications to suckers as injury may occur.

*Not Registered for Use by California

¹See **Application Methods for Tree, Nut, Vine, and Berries** section for additional information on banded, directed spray and spot treatment applications.

Crop	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
OLIVES	Broadcast Banded Directed Spray Spot Treatments	Weeds < 3" in Height: 48.0 (0.88 lb. ai) Weeds < 6" in Height: 49.0 - 56.0 (0.90 - 1.02 lbs. ai) Weeds > 6" in Height and/or Grasses that have Tillered: 56.0 - 82.0 (1.02 - 1.50 lbs. ai)	Apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Inflame 280 SL . Uniform, thorough spray coverage is necessary to achieve consistent weed control. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures. Weeds under stress or in dense populations will require application at the highest specified label use rate. Stressed conditions also include prior treatments of other contact or systemic herbicides. DO NOT retreat these weeds with Inflame 280 SL until sufficient regrowth has occurred. Avoid direct spray, drift or mist to desirable vegetation, green bark, stems, or foliage, as injury may occur.
	Sucker Control Broadcast	56.0 (1.02 lbs. ai)	When applied to suckers that are young, green, and uncalled, Inflame 280 SL will reduce or eliminate sucker growth. Make a split application approximately 4 weeks apart. Thorough coverage of all sucker foliage is necessary for optimum control.

Restrictions:

- **DO NOT** apply more than 82 fl. oz./A (1.50 lbs. ai/A) per application.
- **DO NOT** make more than 5 applications per year at reduced application rates.
- **DO NOT** apply more than 246 fl. oz./A (4.50 lbs. ai/A) through any combination of use patterns per year.
- Applications must be a minimum of 14 days apart.
- **DO NOT** graze, harvest and/or feed treated orchard cover crops to livestock.
- **DO NOT** aerially apply.
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** allow spray to contact trunks other than those that have callused, mature brown bark or are protected from spray contact by nonporous wraps, grow tubes, or waxed containers.
- **Spot Treatment:**
 - o **DO NOT** exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application.
 - o **DO NOT** exceed the equivalent of 5.65 fl. oz. (0.10 lb. ai) per 1,000 sq. ft. per year.
- **Sucker Control:**
 - o Suckers must not exceed 12 inches in length.
 - o **DO NOT** make spot or directed-spray applications to suckers as tree injury may occur.

¹See **Application Methods for Tree, Nut, Vine, and Berries** section for additional information on banded, directed spray and spot treatment applications.

TROPICAL AND SUBTROPICAL FRUITS*

Crop	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Avocado* (including Guatemalan, Mexican, and West Indian)	Broadcast Banded Directed Spray Spot Treatments	Weeds < 3" in Height: 48.0 - 82.0 (0.88 - 1.50 lbs. ai) Weeds < 6" in Height Pre-tiller Grasses: 56.0 - 82.0 (1.02 - 1.50 lbs. ai) Weeds > 6" in Height and/or Grasses that have Tillered: 64.0 - 82.0 (1.17 - 1.50 lbs. ai)	Inflame 280 SL may be applied in a single application or in sequential applications. Avoid contact of Inflame 280 SL solution, spray, drift, or mist with green bark, stems, foliage, or fruit as injury may occur to trees. Only trunks with callused, mature brown bark may be sprayed unless protected from spray contact by nonporous wraps, grow tubes, or waxed containers. Contact of Inflame 280 SL with parts of trees other than mature brown bark can result in serious damage. For postemergence control of weeds present, apply Inflame 280 SL as a broadcast directed spray anytime during the season up to the day of harvest. Inflame 280 SL may also be applied as a banded or spot treatment to target emerged weeds. Sequential Applications: Apply Inflame 280 SL at a minimum of 30 days apart. Regrowth may occur due to the weed stage of growth at application, low use rate, or environmental conditions. Repeat applications of Inflame 280 SL may be necessary to control plants generating from underground parts or seed.

Restrictions:

- **DO NOT** apply more than 82 fl. oz./A (1.50 lbs. ai/A) per application.
- **DO NOT** apply more than a maximum cumulative amount of 246 fl. oz./A of **Inflame 280 SL** (4.50 lbs. ai/A) from sequential applications per year.
- **DO NOT** make more than 3 applications per year.
- Separate sequential applications by at least 30 days.
- **DO NOT** apply this product aerially.
- **Pre-Harvest Interval (PHI):** 1 day.
- **Spot Treatment:**
 - o **DO NOT** exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application.
 - o **DO NOT** exceed the equivalent of 5.65 fl. oz. (0.10 lb. ai) per 1,000 sq. ft. per year.

*Not Registered for use by California

¹See **Application Methods for Tree, Nut, Vine, and Berries** section for additional information on banded, directed spray and spot treatment applications.

Crop	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Fig*	Broadcast Banded Directed Spray Spot Treatments	Weeds < 3" in Height: 48.0 - 82.0 (0.88 - 1.50 lbs. ai) Weeds < 6" in Height Pre-tiller Grasses: 56.0 - 82.0 (1.02 - 1.50 lbs. ai) Weeds > 6" in Height and/or Grasses that have Tillered: 64.0 - 82.0 (1.17 - 1.50 lbs. ai)	Apply in a single application or in sequential applications. For postemergence control of weeds present, apply Inflame 280 SL as a broadcast directed spray anytime during the season up to the day of harvest. Inflame 280 SL may also be applied as a banded or spot treatment to target emerged weeds. Avoid contact of Inflame 280 SL solution, spray, drift, or mist with green bark, stems, foliage, or fruit as injury may occur to trees. Only trunks with callused, mature brown bark may be sprayed unless protected from spray contact by nonporous wraps, grow tubes, or waxed containers. Contact of Inflame 280 SL with parts of trees other than mature brown bark can result in serious damage. Sequential Applications: Apply Inflame 280 SL at a minimum of 30 days apart. Regrowth may occur due to the weed stage of growth at application, low use rate, or environmental conditions. Repeat applications of Inflame 280 SL may be necessary to control plants generating from underground parts or seed.

Restrictions:

- **DO NOT** apply more than 82 fl. oz./A (1.50 lbs. ai/A) per application.
- **DO NOT** apply more than a maximum cumulative amount of 246 fl. oz./A of **Inflame 280 SL** (4.50 lbs. ai/A) from sequential applications per year.
- **DO NOT** make more than 3 applications per year.
- Separate sequential applications by at least 30 days.
- **DO NOT** apply this product aerially.
- **Pre-Harvest Interval (PHI):** 1 day.
- **Spot Treatment:**
 - o **DO NOT** exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application.
 - o **DO NOT** exceed the equivalent of 5.65 fl. oz. (0.10 lb. ai) per 1,000 sq. ft. per year.

*Not Registered for use by California

¹See **Application Methods for Tree, Nut, Vine, and Berries** section for additional information on banded, directed spray and spot treatment applications.

HOPS*

Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Broadcast Directed Spray	Weeds < 3" in Height and Hop Sucker Control: 32.0 - 55.0 (0.59 - 1.00 lb. ai) Weeds < 6" in Height Pre-tiller Grasses: 55.0 (1.0 lb. ai)	Apply in a single application or in sequential applications. For postemergence control of weeds present between hops rows and/or for control of hop sucker growth, apply Inflame 280 SL as a broadcast directed spray to the lower portion of the hop plant. Inflame 280 SL may be applied with a hooded sprayer to prevent spray drift to susceptible vegetation. Avoid contact of Inflame 280 SL solution, spray, drift, or mist with green bark, stems, foliage, or fruit as injury may occur to trees. Only trunks with callused, mature brown bark may be sprayed unless protected from spray contact by nonporous wraps, grow tubes, or waxed containers. Contact of Inflame 280 SL with parts of trees other than mature brown bark can result in serious damage.
Burndown (14 days prior to planting)	Weeds < 3" in Height: 32.0 (0.59 lb. ai) Weeds < 6" in Height Pre-tiller Grasses: 55.0 (1.0 lb. ai)	Sequential Applications: Apply Inflame 280 SL at a minimum of 25 days apart. Regrowth may occur due to the weed stage of growth at application, low use rate, or environmental conditions. Repeat applications of Inflame 280 SL may be necessary to control plants generating from underground parts or seed.
Restrictions: • DO NOT apply more than 55 fl. oz./A of Inflame 280 SL (1.00 lb. ai/A) per application. • DO NOT apply more than a maximum cumulative amount of 165 fl. oz./A of Inflame 280 SL (3.02 lbs. ai/A) from sequential applications in hops per year. • DO NOT make more than 3 applications per year. • Separate sequential applications by at least 25 days. • DO NOT apply aerially. • Pre-Harvest Interval (PHI): 10 days. • DO NOT apply to hops that are less than 6 feet tall, and then only apply to the lower 18 inches of hops plants that are over 6 feet tall. • DO NOT apply to hop suckers prior to training hops on the string/wire and before hop height is 6 feet tall on string/wire. • DO NOT use Inflame 280 SL to burn back existing vines to obtain even emergence of subsequent vines. • Burndown: ◦ DO NOT make more than 1 burndown application per year. *Not Registered for use by California		

¹See **Application Methods for Tree, Nut, Vine, and Berries** section for additional information on banded, directed spray and spot treatment applications.

POTATOES

Use Pattern	Rate (Fl. Oz./Acre)	Directions
Vine Desiccation	21.0 (0.38 lb. ai)	Apply at the beginning of natural senescence of potato vines and when petiole nitrate levels are below 15,000 ppm. Potato varieties with heavy or dense vines may require an application of another desiccation product to complete vine desiccation. Thorough coverage of the potato vines to be desiccated is essential. Use sufficient volume of water (20 to 100 gpa). Vary the gallons of water per acre and spray pressure as indicated by the density of the potato vines. Increase spray volume to at least 30 gallons of water per acre when potato canopy is dense or under cool and dry conditions. Apply with the spray boom as low as possible to achieve thorough coverage of the potato vines for best control and to minimize drift potential. The use of additives or adjuvants may improve the performance of Inflame 280 SL in desiccating potatoes. However, the combination of Inflame 280 SL with adjuvants, other than ammonium sulfate (AMS), have been known to cause injury in potatoes under specific conditions and in certain geographies. To the extent consistent with applicable law, the user assumes all risks associated with adding adjuvants, other than AMS, to Inflame 280 SL. Atticus, LLC cannot be held responsible for crop injury on potatoes when using these adjuvants.
Restrictions: • DO NOT apply to potatoes grown for seed. • DO NOT split application or make more than 1 application per year. • DO NOT apply more than 21.0 fl. oz./A (0.38 lb. ai/A) per year. • DO NOT harvest potatoes until 9 days or more after application. • Replanting Interval after application of Inflame 280 SL as a potato vine desiccant: - o Potatoes, canola, corn, cotton, rice, soybean and sugar beets: any time - o Wheat, barley, buckwheat, millet, oats, rye sorghum or triticale: 30 days or more - o DO NOT plant treated areas to root and tuber vegetables, leafy vegetables, and Brassica vegetables until 70 days after an application of Inflame 280 SL potato vine desiccant. - o All other crops: 120 or more days		

FALLOW FIELDS AND POST HARVEST

Use Site	Use Pattern	Rate (Fl. Oz./Acre)	Directions ¹
Fallow Fields Any crop listed on this label: Cotton Corn (field, silage, Sweet) Canola Soybean Sugar Beets Pome Fruit Citrus Small Vine Climbing Fruit (except Fuzy Kiwifruit) Stone Fruit Tree Nuts (including Pistachio) Bushberries Tropical and Subtropical Fruits Olives Cucurbits Fruiting Vegetables Hops Potatoes	Post-harvest Before Planting or emergence	22.0 - 29.0 (0.40 - 0.53 lb. ai)	Inflame 280 SL may be used as a substitute for tillage in fallow fields to control or suppress weeds listed in the WEEDS CONTROLLED Table 1 section of this label. Applications may be made in fallow fields, post-harvest, before planting or emergence of any crop listed on this label. Inflame 280 SL must be applied with ammonium sulfate or an adjuvant containing ammonium sulfate at a rate that provides 1.5 to 3 lbs. ammonium sulfate/acre. Tank mixes with 2,4-D, glyphosate or atrazine and Inflame 280 SL will enhance total weed control. See the APPLICATION INSTRUCTIONS section for additional information on tank mixes and how to apply this product.
Restrictions: • DO NOT apply more than 29 fl. oz./A (0.53 lb. ai/A) in one application. • DO NOT apply more than 145 fl. oz./A (2.65 lbs. ai/A) per year. • Applications must be a minimum of 10 days apart. • DO NOT apply more than 5 applications per year at 29 fl. oz./A (0.53 lb. ai/A).			

¹See the **PRODUCT INFORMATION** section of this label for rotational crop restrictions.

NON-CROP USE DIRECTIONS

When applied as listed, **Inflame 280 SL** controls undesirable plant vegetation in non-crop areas including the following: access roads, golf courses (excluding greens, tees, aprons, fairways, and roughs)*, railroad rights of way, airfields, gravel yards[†], ramps, airports, refineries, alleys, habitat restoration and management areas, resorts, along fences fence rows, and fence lines[†], highways and roadsides (including aprons, medians, guardrails, and rights of way), sewage disposal areas, bare ground[†], industrial areas and industrial plant sites, around commercial or industrial structures or outbuildings, greenhouses, shade houses, barrier strips, landscapes[†], shelter belts, beaches[*], lanes, sidewalks[†], campgrounds, mulched areas[†], canals, natural areas and brush control[†], sports areas, commercial plants, nurseries, storage and lumber yards, ornamental gardens[†], substations, construction sites, parking lots and parking areas, tennis courts, ditch banks, parks and recreation areas, trails, drive-in theaters, paths[†], uncropped farmstead areas, driveways[†], paved areas[†], vacant lots, dry ditches, petroleum and other tank farms, walkways[†], educational facilities and schools, pipeline, power, telephone and utility rights of way, wastelands, around farm and ranch structures and outbuildings, farmsteads and farmstead areas (barnyards, buildings, building foundations, driveways, facilities, farmyards, machinery or implement yards, windbreaks, shelter belts), power stations, wildlife food plots*, firebreaks, preplant to turf and ornamental plants[†], wildlife habitat areas, fuel storage areas, pumping installations, wildlife openings, ornamentals and Christmas trees, conservation reserve program (CRP)*

*Not Registered for use by California

[†]Residential uses are limited to spot treatment only.

Application Methods for Non-Crop Uses

Inflame 280 SL is a foliar-active material and works best when weeds are actively growing. Weed control may be reduced when applications are made to weeds under stress due to drought or cool temperatures.

Weeds under stress or in dense populations will require application of the highest rate indicated. Always apply at the labeled rate. Repeat applications or tank mixes of **Inflame 280 SL** plus one or more appropriate residual herbicides will be needed to control weeds emerging from underground parts or seeds. When tank mixing with other herbicides, follow the label with the most restrictive directions for use and precautions. No label dosage rates may be exceeded.

Regrowth may occur due to the weed stage of growth at application, low use rate, or environmental conditions. Repeat treatments may be necessary to control weeds generating from underground parts or seeds.

Trimming and Edging

Inflame 280 SL may be used for trimming and edging listed non-crop areas. For control of weeds emerging from seed, the use of **Inflame 280 SL** in a tank mix with pre-emergence herbicides is advised. If spraying in areas adjacent to desirable plants, use a shield made of cardboard, plywood, or sheet metal while spraying to help prevent spray from contacting foliage of desirable plants.

Directed Spray and Spot Treatment

Use rate depends on weed species being controlled. Spray undesirable vegetation foliage on a spray-to-wet basis. Ensure uniform and complete coverage. Use a coarse spray. Backpack, pump-up, and hydraulic sprayers may be used. Thoroughly clean the sprayer following use.

Broadcast

Use rate depends on weed species being controlled. Use 30-psi spray pressure minimum. For smaller weeds 3 inches or less, use the lower rate. For weeds 6 inches or less use the upper end of the rate range.

Drift control additives may be used. If a drift control additive is used, observe and follow all directions and precautions as specified on the additive label.

Use Notes

Use higher rates within the specified rate range for weed sized listen when vegetation cover is sense or when weeds are growing under stressed conditions including drought or when average temperatures are below 50°F.

The addition of 8.5 to 17 pounds of ammonium sulfate (spray grade) per 100 gallons of water (1 to 2% by weight) or 2 to 4 pounds of ammonium sulfate per acre may improve the level of weed control.

Use Site	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions ²
Farmsteads, Recreational, and Public Areas	Broadcast Banded Directed Spray Spot Treatment	Weeds <3" in Height: 48.0 (0.88 lb. ai) Weeds <6" in Height: 49.0 - 56.0 (0.90 - 1.02 lbs. ai) Weeds >6" in Height and/or Grasses that have Tillered: 56.0 - 82.0 (1.02 - 1.50 lbs. ai)	For best results, apply to emerged, young, actively growing weeds. Warm temperatures, high humidity, and bright sunlight improve the performance of Inflame 280 SL . Apply Inflame 280 SL as a directed spray to control undesirable vegetation in farmsteads, recreational, and public areas listed on this label. Apply as a broadcast, banded, or spot treatment application depending on the situation to control weeds listed under the heading Table 1. WEEDS CONTROLLED . Avoid direct spray or drift to desirable vegetation. For spot or directed-spray applications, mix Inflame 280 SL at 1.7 fl. oz. of product per gallon of water.

Restrictions:

- **DO NOT** apply more than 82 fl. oz./A (1.5 lbs. ai/A) in one application.
- Applications must be a minimum of 14 days apart.
- **DO NOT** apply more than 246 fl. oz./A (4.5 lbs. ai/A) per year.
- **DO NOT** apply more than 3 applications per year.
- **DO NOT** apply this product through any type of irrigation system.
- **Directed Spray and Spot Treatment:**
 - **DO NOT** exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application.
 - **DO NOT** exceed the equivalent of 5.65 fl. oz. (0.10 lb. ai) per 1,000 sq. ft. per year.

¹See **Application Methods for Non-Crop Uses** section for additional information on broadcast, directed spray and spot treatment applications.

²See the **PRODUCT INFORMATION** section of this label for rotational crop restrictions.

Use Site	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Military and Private Areas	Broadcast Banded	48.0 - 72.0 fl. oz./A (0.88 - 1.32 lbs. ai)	For ground broadcast applications, apply in a minimum of 40 gallons of water per acre. For aerial broadcast applications, apply in a minimum of 5 gallons of water per acre. Refer to WEEDS CONTROLLED Table 2 section for list of weeds controlled.
	Directed Spray Spot Treatment	1.0 - 2.0 fl. oz. (0.02 - 0.04 lb. ai) Per gallon of water	

Restrictions:

- **DO NOT** apply more than 72 fl. oz./A (1.5 lbs. ai/A) in one application.
- **DO NOT** apply more than 72 fl. oz./A (1.5 lbs. ai/A) per year.
- **DO NOT** apply more than 1 application per year.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation.
- **DO NOT** allow grazing of vegetation treated with **Inflame 280 SL**.
- **Directed Spray and Spot Treatment:**
 - o **DO NOT** apply more than 2 fl. oz. (0.04 lb. ai) per gallon of water per application.
 - o **DO NOT** apply more than 1.7 fl. oz./1,000 sq. ft. per year.
 - o **DO NOT** apply beyond runoff.
 - o **DO NOT** spray during windy conditions.

¹See **Application Methods for Non-Crop Uses** section for additional information on broadcast, directed spray and spot treatment applications.

Use Site	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Conservation Reserve Program (CRP)*	Broadcast	48.0 - 56.0 (0.88 - 1.0 lb. ai)	Inflame 280 SL may be used to suppress competitive growth and seed production of undesirable vegetation when rotating out of CRP acres. Apply at labeled rates in early spring, before CRP grasses break dormancy, for selective applications with broadcast spray equipment. After desirable perennial grasses have reached dormancy, late fall applications may be made. Some stunting of CRP perennial grasses may occur if applications are made when plants are not dormant. For ground broadcast applications, apply in a minimum of 40 gallons of water per acre. For aerial broadcast applications, apply in a minimum of 5 gallons of water per acre. Refer to WEEDS CONTROLLED Table 2 section for list of weeds controlled.

Restrictions:

- **DO NOT** apply more than 72 fl. oz./A (1.32 lbs. ai) in a single application.
- **DO NOT** apply more than 72 fl. oz./A (1.5 lbs. ai/A) per year.
- **DO NOT** apply more than 1 application per year
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation.
- **DO NOT** allow grazing of vegetation treated with **Inflame 280 SL**.

*Not Registered for use by California

¹See **Application Methods for Non-Crop Uses** section for additional information on broadcast applications.

Use Site	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Dormant Bermudagrass	Broadcast Banded Directed Spray Spot Treatment	56.0 - 82.0 (1.02 - 1.50 lbs. ai)	Inflame 280 SL may be used to control winter annual weeds in well-established ornamental dormant hybrid or common Bermudagrass. Apply only when the turf is fully dormant and weather is cool, and prior to spring green-up or severe turfgrass injury or delayed green-up may occur. For best results, apply Inflame 280 SL at labeled rates after most weeds have germinated and are in an early growth stage. Applications of Inflame 280 SL may also be used to suppress or control target biennial or perennial weeds. Avoid high volume and spot applications where spray volume exceeds 80 gallons per acre or injury or delayed green-up may occur. Refer to WEEDS CONTROLLED Table 2 section for list of weeds controlled.

Restrictions:

- **DO NOT** apply more than 82 fl. oz./A (1.5 lbs. ai/A) in a single application.
- **DO NOT** apply more than 82 fl. oz./A (1.5 lbs. ai/A) per year.
- **DO NOT** make more than one application per year.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation.
- **DO NOT** allow grazing of vegetation treated with **Inflame 280 SL**.
- **Directed Spray and Spot Treatment:**
 - o **DO NOT** exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application.
 - o **DO NOT** exceed the equivalent of 1.88 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per year.

¹See **Application Methods for Non-Crop Uses** section for additional information on Broadcast, Directed Spray and Spot Treatments.

Use Site	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Ornamentals and Christmas Trees	Directed Spray (Pre-Plant Site Preparation) Directed Spray (Control in-row weeds in field-grown wood plants)	1.0 - 2.0 fl. oz. (0.02 - 0.04 lb. ai) Per gallon of water	When applied as advised by this label, this product may be used for the control of undesired vegetation in site preparation prior to planting, around and within shade and greenhouses, and as a directed spray around containers and field-grown established ornamentals and Christmas trees. For pre-plant site preparation applications for control of annual and perennial weeds listed on this label, ornamental and Christmas trees may be planted into the treated area after the restricted-entry interval (REI) of 12 hours has elapsed. Inflame 280 SL may be used between and around containers and in site preparation for new plantings, and to control in-row weeds in field-grown wood plants. Apply Inflame 280 SL as a directed spray. Refer to WEEDS CONTROLLED Table 2 section for list of weeds controlled.

Restrictions:

- **DO NOT** apply more than 2 fl. oz. (0.04 lb. ai) per gallon of water in any single application.
- **DO NOT** apply more than 1.7 fl. oz./1,000 sq. ft. per year.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation or injury may result.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** allow grazing of vegetation treated with **Inflame 280 SL**.
- **DO NOT** apply **Inflame 280 SL** as an over-the-top broadcast spray in ornamentals and shade or Christmas trees.
- **DO NOT** apply beyond runoff.
- **DO NOT** spray during windy conditions.

¹See **Application Methods for Non-Crop Uses** section for additional information on directed spray applications.

Use Site	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Greenhouse and Shadehouse	Directed Spray	1.0 - 2.0 fl. oz. (0.02 - 0.04 lb. ai) Per gallon of water	For greenhouse and shade house applications where Inflame 280 SL is used to control weeds, air circulation fans must be turned off during application. Apply Inflame 280 SL as a directed spray, using large droplet and low-pressure type nozzles. Avoid drift and direct contact with desirable vegetation. Refer to WEEDS CONTROLLED Table 2 section for list of weeds controlled.

Restrictions:

- **DO NOT** apply more than 2 fl. oz. (0.04 lb. ai) per gallon of water in any single application.
- **DO NOT** apply more than 1.7 fl. oz./1,000 sq. ft. in a single year.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation.
- **DO NOT** allow grazing of vegetation treated with **Inflame 280 SL**.
- **DO NOT** use in greenhouses or shade houses containing edible crops.
- **DO NOT** apply beyond runoff.
- **DO NOT** spray during windy conditions.

¹See **Application Methods for Non-Crop Uses** section for additional information on directed spray applications.

Use Site	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Site Preparation for Conifer and Hardwood Production Areas	Broadcast Banded	48.0 - 72.0 fl. oz. (0.88 - 1.32 lbs. ai)	Prior to planting conifer and hardwood species, Inflame 280 SL can be used as a site preparation treatment.
	Directed Spray Spot Treatment	1.0 - 2.0 fl. oz. (0.02 - 0.04 lb. ai) Per gallon of water	For ground broadcast applications, apply in a minimum of 40 gallons of water per acre. For aerial broadcast applications, apply in a minimum of 5 gallons of water per acre. Refer to WEEDS CONTROLLED Table 2 section for list of weeds controlled.

Restrictions:

- **DO NOT** apply more than 72 fl. oz./A (1.32 lbs. ai) in any single application.
- **DO NOT** apply more than 72 fl. oz./A (1.32 lbs. ai) in a single year.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation.
- **DO NOT** allow grazing of vegetation treated with **Inflame 280 SL**.
- **DO NOT** apply **Inflame 280 SL** as an over-the-top broadcast spray to desirable conifer or hardwood plantings.
- Restricted-Entry Interval (REI) for seedling conifer and hardwood treats to be planted into the treated area: 12 hours.
- **Directed Spray and Spot Treatment:**
 - o **DO NOT** apply more than 2 fl. oz. (0.04 lb. ai) per gallon of water per application.
 - o **DO NOT** apply more than 1.7 fl. oz./1,000 sq. ft. per year.
 - o **DO NOT** apply beyond runoff.
 - o **DO NOT** spray during windy conditions.

¹See **Application Methods for Non-Crop Uses** section for additional information on Broadcast, Directed Spray and Spot Treatments.

Use Site	Use Pattern ¹	Rate (Fl. Oz./Acre)	Directions
Brush Control or Suppression*	Broadcast Banded Directed Spray Spot Treatment	32.0 - 96.0 (0.59 - 1.76 lbs. ai)	Inflame 280 SL will provide control or suppression of listed perennial wood weed species when applied as directed. When conditions are not optimum for good spray penetration, use the higher directed use rate. Refer to WEEDS CONTROLLED Table 2 section for list of weeds controlled.

Restrictions:

- **DO NOT** apply more than 246 fl. oz./A (4.5 lbs. ai/A) per year.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply directly to or allow drift to contact desirable green tissue or green, thin, or uncalloused bark of desirable vegetation.
- **DO NOT** allow grazing of vegetation treated with **Inflame 280 SL**.
- **Directed Spray and Spot Treatment:**
 - o **DO NOT** exceed the equivalent of 2.2 fl. oz. (0.03 lb. ai) per 1,000 sq. ft. per application.
 - o **DO NOT** exceed the equivalent of 5.65 fl. oz. (0.10 lb. ai) per 1,000 sq. ft. per year.

*Not Registered for Use by California

¹See **Application Methods for Non-Crop Uses** section for additional information on Broadcast, Directed Spray and Spot Treatments.

SPECIMEN

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Do not use or store near heat or open flame. Keep the container tightly closed and dry in a cool, well-ventilated place. Storage temperature should not exceed 125°F. If storage temperature for bulk **Inflame 280 SL** is below 32°F, the material must not be pumped until its temperature exceeds 32° F. Protect against direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER HANDLING:

Rigid, Non-refillable plastic containers small enough to shake (i.e., with capacities equal to or less than 5 gallons) Non-refillable container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Once container is rinsed, then offer for recycling if available or reconditioning if appropriate; or puncture and dispose of in a sanitary landfill, or by incineration; or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Rigid, Non-refillable plastic containers (i.e., with capacities greater than 5 gallons) triple rinse or pressure rinse as follows: Triple rinse: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container back on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers. Pressure rinse: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

SEED DISPOSAL: To dispose of out-of-date or otherwise unmarketable seed from plants, which have been treated with **Inflame 280 SL**, broadcast and lightly incorporate seed into field soils using disc or other suitable implement. Any resulting crop may be destroyed by chemical or mechanical means. Alternatively, seed may be destroyed by deep burial, incineration or landfill disposal.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of Atticus, LLC. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Atticus, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither Atticus, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Inflame is a trademark of Atticus, LLC.

Liberty, LibertyLink, Laudis, and the LibertyLink design are registered trademarks of Bayer CropScience.

Impact is a registered trademark of Amvac Chemical Company.

20250902bp1