

Product Bulletin



Corteva Agriscience, LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quelex

EPA Reg. No. 62719-661

Special 2(ee) Recommendation

(For Distribution and Use Only in the States of ID, OR, and WA)

For Improved Control of Annual Mustard Species, Prickly Lettuce, and Common Lambsquarters

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of this product according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for this product.

Directions for Use

Refer to Quelex product label for General Use Precautions, Mixing Directions and Application Directions.

Quelex may be applied at 0.55 oz/acre in tank mix combination with 16.4 fl oz/acre Rezuvant to wheat or barley, or in tank mix combination with 6.0 fl oz/acre Pixxaro EC to wheat, barley, or triticale. Apply to actively growing labeled crops from the 2-leaf crop growth stage up to flag leaf emergence (Zadoks scale 39). Apply to susceptible broadleaf weed seedlings which are actively growing and less than 8 inches tall. For best results apply when susceptible broadleaf weed seedlings are actively growing and less than 4 inches tall. Only weeds emerged at the time of treatment will be controlled.

Warm, moist growing conditions promote active weed growth and enhance the activity of Quelex by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds.

R411-014

Issued: 01/27/26

Replaces R411-013