

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call poison control center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call poison control center or doctor for treatment advice. **IF SWALLOWED:** Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Call poison control center or doctor for treatment advice. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. CORROSIVE. Causes irreversible eye damage and skin burns. May be fatal if inhaled or absorbed through the skin. Harmful if swallowed. Do not breathe vapors or spray mist. Do not get in eyes on skin or on clothing. Wear Coveralls worn over long-sleeved shirt and long pants, socks, chemical-resistant footwear, waterproof or chemical-resistant gloves, a NIOSH approved respirator with any N,R,P filter with NIOSH approval number prefix TC-84A; or a NIOSH-approved powered air purifying respirator with an HE filter, with NIOSH approval number prefix TC-21c. Wear goggles and/or face shield and rubber gloves when handling. Do not enter an enclosed area without proper respiratory protection. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

PHYSICAL AND CHEMICAL HAZARDS. STRONG OXIDIZING AGENT. CORROSIVE. Mix only with water. Product must be diluted in accordance with label directions prior to use. **Jet-Oxide® 15** is not combustible; however, at temperatures exceeding 156°F, decomposition occurs releasing oxygen. The oxygen released could initiate combustion.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds, fish, and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the US Environmental Protection Agency. Sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the US Environmental Protection Agency.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Wear Coveralls worn over long-sleeved shirt and long pants, socks, chemical-resistant footwear, waterproof or chemical resistant gloves, a NIOSH approved respirator with any N, R, P filter with NIOSH approval number prefix TC-84A; or a NIOSH-approved powered air purifying respirator with an HE filter, with NIOSH approval number prefix TC-21c.

Bactericidal*, VirucidalΔ, Fungicidal, Tuberculocidal For use on hard, non-porous surfaces.

SANITIZATION

NOTE: FOR MECHANICAL OPERATIONS prepared use solution may not be used for subsequent sanitizing but may be reused for other purposes such as cleaning.

FOR MANUAL OPERATIONS fresh sanitizing solutions should be prepared at least daily or more often if the solution becomes diluted or soiled.

Jet-Oxide® 15 peroxyacetic acid sanitizer is recommended for use on pre-cleaned surfaces such as equipment, pipelines, tanks, vats, fillers, evaporators, pasteurizers and aseptic equipment in dairies, breweries, wineries, beverage and food processing/packing plants, egg processing/packing equipment surfaces, and eating establishments. This product is effective as a sanitizer when solution is prepared in water of up to 400 ppm hardness as CaCO3. This product has demonstrated greater than a 99.99% reduction of survivors after a 30 second exposure period in the AOAC Germicidal and Detergent Sanitizing Action of Disinfectants study.

Sanitizing Food Contact Surfaces: Effective against Staphylococcus aureus [ATCC 6538] and Escherichia coli [ATCC11229]. Prior to sanitizing, remove gross food particles, then wash with a detergent solution, followed by a potable water rinse. Sanitize with a concentration of 0.33 - 1.87 fluid ounce **Jet-Oxide® 15** dissolved in 5 gallons of water. This will provide 88 - 500 ppm of peroxyacetic acid. Use immersion, coarse spray or circulation techniques as appropriate to the equipment. All surfaces should be exposed to the sanitizing solution for a period of at least 60 seconds or more if specified by governing sanitary code. Surfaces should remain visibly wet for the duration of the contact time. Drain thoroughly. Do not rinse.

Sanitizing, Eating, Drinking, And Food Prep Utensils: Remove gross food particles by a prescrape, a preflush and, when necessary, a presoak treatment. Wash with a recommended detergent. Rinse with clean water. Sanitize in a solution of 0.33 - 1.87 fluid ounce **Jet-Oxide® 15** dissolved in 5 gallons of water. Immerse all utensils for at least 60 seconds or contact time specified by governing sanitary code. Surfaces should remain visibly wet for the duration of the contact time. Drain and air dry.

Sanitizing Tableware: For sanitizing tableware in low temperature warewashing machines, inject **Jet-Oxide® 15** into the final rinse water at a concentration of 0.33 - 1.87 fluid ounce **Jet-Oxide® 15** dissolved in 5 gallons of water. Do not exceed 500 ppm. This will provide 88 - 500 ppm of peroxyacetic acid. Surfaces should remain visibly wet for the duration of the contact time. Air dry.

To insure that the **Jet-Oxide® 15** sanitizer concentration does not fall below 48 ppm peroxyacetic acid, periodically test the rinse solution with a suitable test kit and adjust the dispensing rate accordingly. Consult your technical service representative for assistance and further information on sanitizing table-ware in warewashing machines

BATCH SANITIZATION (NON-FOOD CONTACT SURFACES) OF ULTRA FILTRATION AND REVERSE OSMOSIS (RO) MEMBRANES:** **Jet-Oxide® 15** can be used for the sanitization of ultra filtration, medical and non-medical institutional/industrial reverse osmosis (RO) membranes and their associated distribution systems.

This product has been shown to be an effective disinfectant when tested according to AOAC. This product may not eliminate all vegetative microorganisms in reverse osmosis membranes and their associated piping systems due to their construction and/or assembly, but can be relied upon to reduce the number of microorganisms to acceptable levels when used as directed.

Check with equipment manufacturer for membrane compatibility with **Jet-Oxide® 15**. Remove biological or organic fouling from the membrane or other parts of the system with an appropriate cleaner. Flush the system with RO permeate or similar quality water.

Remove mineral deposits with suitable acidic cleaner prior to sanitizing the membranes with **Jet-Oxide® 15**. Flush the system again with the RO permeate or similar quality water. Prepare an appropriate volume of 1% solution of the product 42.52 fl. oz of **Jet-Oxide® 15** to 100 gallons of water. This will provide 568 ppm of peroxyacetic acid and 833.1 ppm hydrogen peroxide. Fill the entire water circuit to be sanitized with the dilute solution and allow the solution to reach a minimum of 20 °C (68°F). Recirculate the dilute solution of **Jet-Oxide® 15** for a minimum of 10 minutes. Allow membrane elements to soak in the solution for a minimum of 20 minutes. Rinse the RO system and test for residuals to ensure that there is less than 3 ppm per oxygen. Diverting product water to drain can reduce residuals.

Continuous/Intermittent Addition to Minimize the Accumulation of Biological Matter Between Intermittent Sanitizing Episodes in Piping Systems Associated with RO Membranes (Non-food Contact Surfaces):** **Jet-Oxide® 15**, as received or diluted, may be added continuously to the feed water system, between system sanitizing episodes, to aid in minimizing the re-growth/ accumulation of biological matter. The peroxigen residual in the system which will be effective will vary with the design and usage characteristics of the system. Adjust the addition rate of **Jet-Oxide® 15** or the solution and periodically monitor residual peroxigen so that the desired effect is obtained. For continuous addition, do not exceed 7 ppm peroxyacetic acid (0.33 fluid ounce of product per 440 gallons of water) **Jet-Oxide® 15**. This will give 1 ppm peroxyacetic acid and 1.4 ppm hydrogen peroxide. For intermittent feed, do not exceed 750 ppm (8.5 fluid ounces of product per 100 gallons of water) **Jet-Oxide® 15**. This will give 110 ppm peroxyacetic acid and 160 ppm hydrogen peroxide.



(PEROXYACETIC ACID SOLUTION)

**Jet-Oxide® 15 IS A PEROXYACETIC ACID-BASED
SANITIZER/DISINFECTANT DEVELOPED FOR THE FOLLOWING USES:**

Institutional/Industrial Sanitizing of Previously Cleaned Non-Porous Food Contact Surfaces in Dairies, Wineries, Breweries, Food and Beverage Plants, Disinfecting Poultry Premises, Poultry Hatchery surfaces, Animal Housing Facilities, and Reverse Osmosis Membranes and Ultra Filtration. Hard Surface Disinfection in Hospitals, Health Care Facilities, Schools, Colleges, Veterinary Clinics, Animal Life Science Laboratories, Industrial Facilities, Office Buildings, Recreational Facilities, and Retail and Wholesale Establishments. Non-public Health Bacteria, Fungi & Slime Control in Pulp and Paper Mill Systems.

Active Ingredients:	
Hydrogen Peroxide	22.0%
Peroxyacetic Acid.....	15.0%
Inert Ingredients:	63.0%
Total:	100.0%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

Before Using This Product, Please Read This Entire Label Carefully.

**not approved in the State of California

PN105862
MBI-LBL-138 r2

EPA Registration
No. 54289-4-84059

EPA Est. No.
☐ 70253-CA-001
☒ 70547-IL-001

Manufactured for:



Pro Farm Group, Inc.
1530 Drew Ave.
Davis, CA 95618
1-877-664-4476; info@profarmgroup.com

- Net Contents:**
- ☐ 2.5 gallons
 - ☐ 5 gallons
 - ☐ 30 gallons
 - ☐ 55 gallons
 - ☐ 265 gallons
 - ☐ 330 gallons
 - ☒ **Volume stated on container**

Batch:

Hard Surface Disinfection: Jet-Oxide® 15 disinfects as it cleans in one operation. **Jet-Oxide® 15** can be used to disinfect floors, walls and other hard nonporous surfaces such as tables, chairs, countertops, bathroom fixtures, sinks, bed frames, shelves, racks, carts, refrigerators, coolers, glazed tile, linoleum, vinyl, non-porous glazed porcelain, plastic (such as polypropylene and polyethylene), stainless steel, or glass. Refrigerators and coolers should be at room temperature for treatment.

For Health Care, Institutional, and Industrial Use: Jet-Oxide® 15 may be used for surgical and obstetrical suites, house-keeping services, physical therapy departments, nursing services, and autopsy facilities. Also, use **Jet-Oxide® 15** in nursing homes, other healthcare facilities, schools, colleges, veterinary clinics, animal life science laboratories, industrial facilities, dietary areas, office buildings, recreational facilities, retail and wholesale establishments.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

COMBINATION DISINFECTION AND CLEANING

Jet-Oxide® 15 is effective against Staphylococcus aureus, Salmonella choleraesuis, Pseudomonas aeruginosa, Trichophyton interdigitale, and Escherichia coli 0157:H7 at 0.08% (0.5 fl. oz./5 gal.) in hard water (400 ppm as CaCO3) and 5% fetal bovine serum on hard nonporous surfaces which will provide 137 ppm peroxyacetic acid. For visibly soiled areas a pre-cleaning step is required. Apply solution with mop, cloth, sponge, brush, scrubber, or coarse spray device, or by soaking so as to wet all surfaces thoroughly. Allow to remain visibly wet for 10 minutes, then remove solution and entrapped soil with a clean wet mop, cloth, or wet vacuum pickup. Surfaces should remain visibly wet for the duration of the contact time. Prepare a fresh solution daily or when it becomes soiled or diluted.

Jet-Oxide® 15 is designed for use in animal hospitals, animal laboratories, kennels, pet shops, zoos, pet animal quarters, poultry premises, poultry hatcheries (not eggs), and livestock quarters. When used as directed, **Jet-Oxide® 15** is specifically designed to disinfect, deodorize and clean inanimate, hard, surfaces such as walls, floors, sink tops, furniture, operating tables, kennel runs, cages, and feeding and watering equipment. In addition **Jet-Oxide® 15** will deodorize those areas which are generally hard to keep smelling fresh such as garbage storage areas, empty garbage bins and cans, and any other areas which are prone to odors caused by microorganisms.

All treated equipment that will contact food, feed, or drinking water must be rinsed with potable water before reuse. For visibly soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use. Surfaces should remain visibly wet for the duration of the contact time.

DISINFECTION OF POULTRY PREMISES, TRUCKS, COOPS AND CRATES

Poultry Hatchery Disinfection: Not for hatching eggs. Remove all poultry and feeds from premises, trucks, coops and crates. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with a detergent and rinse with water. Saturate surfaces with a 0.08% (0.5 fl. oz./5 gal.) solution of **Jet-Oxide® 15** for a period of 10 minutes. Surfaces should remain visibly wet for the duration of the contact time. Ventilate buildings, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with a detergent and rinse with potable water before reuse. See your technical representative for specific recommendations for all cleaning and rinsing requirements.

Disinfection and Deodorizing of Animal Housing Facilities (Barns, Kennels, Hutches, Etc.): Do not use in milking stalls, milking parlors, or milk houses (for phenolics, cresylic acid, and pine oils). Remove animals and feed from premises, vehicles, and enclosures. Remove litter, waste matter, and gross soils from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering equipment. Thoroughly clean all surfaces with a detergent and rinse with water. Saturate surfaces with a 0.08% 0.5 fl. oz./5 gal. solution of **Jet-Oxide® 15** for a period of 10 minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

BIFOULING CONTROL IN PULP AND PAPERMILL SYSTEMS:** For use in the manufacture of paper and paperboard intended for food-contact and non-food contact. **Jet-Oxide® 15** can be used to control bacteria, fungi, and fresh water organisms in paper, paperboard, or nonwoven process water and influent water systems. Suitable dosing points include but are not limited to: stock chests, pulpers, the white water loop and white water storage systems and influent water streams.

- CONTAMINATION WITH OTHER CHEMICALS COULD RESULT IN PRODUCT DECOMPOSITION.
- Add **Jet-Oxide® 15** at a point in the system where uniform mixing and even distribution will occur.
 - Use 0.1 to 0.5 lb. (1.4 to 6.85 fl. oz.) of **Jet-Oxide® 15** per 1000 gallons of solution as a continuous or intermittent slug treatment. This will provide 1.8 to 9 ppm peroxyacetic acid (12 to 60 ppm **Jet-Oxide® 15**).

Repeat treatment as required to maintain control.

Antimicrobial Rinse of Pre-Cleaned, New Returnable or Non-Returnable Containers: To reduce the number of non-pathogenic beverage spoilage organisms. Effective against Aspergillus versicolor (ATCC 9577), Byssoschlamys fulva (ATCC 10099), Pediococcus damnosus (ATCC 29358), Lactobacillus buchneri (ATCC 4005), and Saccharomyces cerevisiae.

1. Prepare **Jet-Oxide® 15** solution by adding 9.85 fl. oz. to 5 gallons potable water. This provides 2,632 ppm peroxyacetic acid.
2. Apply antimicrobial rinse at a temperature of 40 °C to 60 °C (104 °F to 140 °F) and allow a minimum seven-second contact period.(ATCC 47058)
3. Allow containers to drain thoroughly, and then rinse with sterile or potable water.

TREATMENT OF FRUIT AND VEGETABLE PROCESS WATER SYSTEMS

Jet-Oxide® 15 can be used in water or ice that contacts raw or fresh, post-harvest or further processed fruits and vegetables for the control of spoilage bacteria and fungi in commercial operations and packinghouses.

Batch, Continuous or Spray System Processes: Fill vessel containing fruits and vegetables with known amount of water. Ensure that water is circulating in vessel if using the submersion method. Add this product at a rate no more than 80 ppm peroxyacetic acid to the use solution. This can be accomplished by initially adding 1.0 fl. oz. per 16.4 gallons of water. The fruits and vegetables can be continuously sprayed (using coarse spray) or submerged (dipped) in the resulting solution. Periodic or continuous addition of this product to maintain the required concentration may be added as necessary. Contact time of 60 seconds is recommended to insure efficacy. A potable water rinse is not required. This product is not intended for use in primary flumes prior to the point of the first dewatering stage.

AGRICULTURAL OR HORTICULTURAL USES

There is a Restricted-Entry-Interval of zero (0) hours after the use of this product. This product should never be mixed or combined with any other pesticide or fertilizer. Upon soil contact this diluted product decomposes rapidly to oxygen, carbon dioxide and water. This product may be harmful to fish if exposed on a continuous basis at concentrations of 1 ppm or more of active peroxyacetic acid. Meter this product into pressurized pipes using a plastic or stainless steel injection/backflow device installed far enough upstream from the equipment to insure thorough mixing. For open flowing bodies of water, apply this product as far upstream as possible to allow adequate mixing prior to the flow entering any larger body of water. If open pouring of this product is required, pour product as close to the surface of the water as possible to reduce odor exposure.

Treatment of Agricultural or Irrigation Water Systems (Sand Filters, Humidification Systems, Storage Tanks, Ponds, Reservoirs, Canals): For the control of sulfides, odor, slime and algae in water systems, apply this product at 2-10 ppm active peroxyacetic acid. This feed rate equals 15-75 fl. oz. per 10,000 gallons of water. Repeat dose as necessary to maintain control, which will vary with seasonal conditions. For prevention of algae, some systems may require continuous low level dosing during warm sunny periods (2-5 ppm peroxyacetic acid).

Drip Irrigation Systems: To clean slime and algae from drip system filters, tapes and emitters, meter this product at the rate of 7.5-15 fl. oz. per 1000 gallons of water (10-20 ppm peroxyacetic acid). When required during normal irrigation cycles, use this product at the recommended dose for a minimum of 30 minutes. Thereafter, the irrigation cycle should be discontinued and the line should not be flushed.

For Treatment of Raw, Unprocessed Fruit and Vegetable Surfaces: Jet-Oxide® 15 can be applied as a dip or spray to control the growth of non-public health microorganisms that may cause decay and/or spoilage on raw, post-harvest fruits and vegetables during the washing process. This product can be applied during physical cleaning processes, including at the roller spreader, washer manifold, dip tank, on the brushes or elsewhere in the washing process prior to, simultaneously with or after detergent wash.

1. Prepare treating solution by diluting 1 fluid ounces per 16 gallons of potable water. This will provide 85 ppm peroxyacetic acid and 125 ppm hydrogen peroxide.
2. Apply the treating solution using a coarse spray directed at the fruits or vegetables, or by soaking the fruits and vegetables in the solution. Allow a contact time of at least 45 seconds.
3. The treated produce can be drain dried without a potable water rinse.
4. Do not reuse solution after treatment.

For Treatment of Processed Fruits and Vegetables and Process Waters to Control Growth of Non-Public Health Micro-organisms that Can Cause Spoilage:

1. Prepare treating solution by diluting 1.5 fluid ounces per 25 gallons of potable water. This will provide 80 ppm peroxyacetic acid and 117 ppm hydrogen peroxide.
2. Apply the treating solution as a spray or dip. Allow a contact time of at least 45 seconds. No rinse following application is required. This use complies with the requirements of 21 CFR173.315 (a) 5.
3. The treated produce can be drain dried without a potable water rinse.
4. Do not reuse solution after treatment.

Bactericidal, Virucidal^Δ, Fungicidal, Tuberculocidal FOR USE ON HARD, NON-POROUS SURFACES

Application: Jet-Oxide® 15 disinfects as it cleans when used according to disinfection directions for use. **Jet-Oxide® 15** can be used to disinfect floors, walls and other hard nonporous surfaces such as tables, chairs, countertops, bathroom fixtures, sinks, bed frames, doors, shelves, racks, carts, refrigerators and coolers that are at room temp for treatment) glazed tile, linoleum, vinyl, non-porous glazed porcelain, glazed ceramic, plastic (such as polypropylene and polyethylene), stainless steel, glass, aluminum, non-porous baked enamel, chrome, laminated or painted surfaces or sealed stone. **Jet-Oxide® 15** should not be used on marble or brass surfaces.

FOR HEALTH CARE, INSTITUTIONAL, AND INDUSTRIAL USE

Areas of Use in Hospitals: Jet-Oxide® 15 may be used for surgical and obstetrical suites; housekeeping surfaces; physical therapy departments; nursing services; dental facilities; autopsy facilities; intensive care units; pharmacies; and clinical laboratories. Also, use **Jet-Oxide® 15** in nursing homes, other health-care facilities, schools, colleges, veterinary clinics, animal life science laboratories, industrial facilities, office buildings, recreational facilities, industrial facilities; hotels; retail facilities; office buildings; retail and wholesale establishments.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection. Dilute Jet Oxide 15 with the appropriate amount of water to an effective concentration of 1135 ppm Peracetic acid and 1665 ppm hydrogen peroxide 0.85 fl. oz per 1 gallon of water. Apply solution with a cloth, mop, sponge, auto-scrubber, or hand pumped trigger sprayer such that all surfaces remain visibly wet for 1 minute to kill bacteria, viruses, and fungi as cited on the label. Use a 3-minute contact time for Streptococcus pneumoniae, and Vancomycin-Resistant Enterococcus faecalis. Use a 10-minute contact time for Tuberculocidal activity. Jet Oxide 15 is effective against Mycobacterium bovis (TB surrogate) at ambient temperature (22°C). For Poliovirus Type 1, dilute Jet Oxide 15 with the appropriate amount of water to an effective concentration of 3000 ppm Peracetic acid and 4400 ppm hydrogen peroxide 2.75 fl. oz per 1 gallon of water for a 3 minute contact time. Surfaces should remain visibly wet for the duration of the contact time. Allow surface to air dry. For visibly soiled areas, a preliminary cleaning is required. Prepare a fresh solution daily or more often if the use solution becomes visibly soiled clouded or diluted.

Jet-Oxide® 15 is effective against the following organisms in 1 minute with 5% organic soil load and 400 ppm hard water: For use on hard, non-porous surfaces.

Ω Contact time is increased to 10 minutes to be effective against Mycobacterium bovis (Tuberculosis surrogate).

§ Contact time is increased to 3 minutes to be effective against Streptococcus pneumoniae, Vancomycin resistant Enterococcus faecalis (VRE) and Poliovirus Type 1

BACTERIA*	HUMAN VIRUSES ^Δ
Acinetobacter baumanii ATCC 19606	Herpes simplex virus Type 1 ATCC VR-260
	Herpes simplex virus Type 2 ATCC VR-734
Community-associated MRSA, USA300 strain (CA-MRSA) CI 08001	Human immunodeficiency virus type 1 (HIV-1) (Zeptomatrix Corporation)
Escherichia coli 0157:H7 ATCC 35150	Human Influenza A virus (A/Hong Kong/8/68-H3N2) SPAFAS
Klebsiella pneumoniae ATCC 4352	Human Rotavirus ATCC VR-2018
Listeria monocytogenes ATCC 19111	Poliovirus Type 1§ ATCC 1562
Methicillin-resistant Staphylococcus aureus (MRSA) ATCC 33592	Respiratory Syncytial Virus ATCC VR-26
Methicillin-resistant Staphylococcus epidermidis (MRSE) ATCC 51625	Rhinovirus Type 37 ATCC 1147
Mycobacterium bovis (TB surrogate)Ω (Organon Teknika Corp)	
	Vaccinia virus ATCC VR-156
Pseudomonas aeruginosa ATCC 15442	Human Coronavirus strain 229e
ATCC VR-740 229e	
Salmonella enterica serovar Typhimurium ATCC 13311	Duck hepatitis B virus (Human Hepatitis B surrogate) (Hepanda Virus Testing)
Serratia marcescens ATCC 13880	Feline calicivirus (Norovirus surrogate) (University of Ottawa)
Shigella dysenteriae serotype 1 ATCC 29026	
Staphylococcus aureus ATCC 6538	Animal viruses ^Δ
Streptococcus pneumoniae§ ATCC 6304	Avian influenza A (Turkey/Wis/66-H9N2) SPAFAS
Streptococcus pyogenes ATCC 19615	
Vancomycin-intermediate Staphylococcus aureus (VISA) ATCC 700787	Fungi ∞
Vancomycin-resistant Enterococcus faecalis (VRE) ATCC 51575	Candida albicans ATCC 10231
Vibrio cholerae ATCC 14035	Trichophyton interdigitale ATCC 9533
Yersinia enterocolitica ATCC 35669	

KILLS HIV-1 (AIDS VIRUS), HBV (Hepatitis B), and HCV(Hepatitis C):

On pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings or other settings in which there is an expected likelihood of soiling of surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus, or Hepatitis C Virus.

Special Instructions for Cleaning and Decontamination Against HIV-1 (AIDS Virus), HBV (Hepatitis B), and HCV(Hepatitis C) of Surfaces/Objects Soiled with Blood/Body Fluids: Personal Protection: Disposable protective gloves, gowns, face masks, or eye coverings as appropriate must be worn during all cleaning of body fluids, blood and decontamination procedures. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application. **Contact Time:** HIV-1, HBV, and HCV are inactivated after 1 minute of contact. **Infectious Material:** Blood and other bodily fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in original containers in a cool, well-vented area, away from direct sunlight. Do not allow product to become overheated in storage. This may cause increased degradation of the product, which will decrease product effectiveness. In case of spill, flood area with large quantities of water. Do not store in a manner where cross-contamination with other pesticides or fertilizers could occur.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited. If wastes cannot be disposed of according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Disposal: Nonrefillable container. Do not reuse or refill this container unless the directions for use allow a different (concentrated) product to be diluted in the container.

Containers Less Than 5 Gallons: Triple Rinse As Follows – Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Containers Greater Than 5 Gallons: Triple Rinse (or equivalent) Promptly After Emptying. Triple Rinse As Follows: Empty the remaining contents into application equipment or mix tank. Fill the container ¾ full with water. Replace and tighten closures. Tip the container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand container on its end and tip back and for the several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use of disposal. Repeat the procedure two more times. Then offer for recycling or dispose in a sanitary landfill, or by incineration, if allowed by state and local authorities by burning.