

Verdato™

For control of certain foliar, stem, and root disease in turfgrass including golf courses, institutional, commercial, and residential lawns, sod farms, sports fields, parks, municipal grounds and cemeteries.

Not Registered for Use by the State of New York For Use in Nassau and Suffolk Counties.

ACTIVE INGREDIENT:

Fludioxonil.....	By Wt.
	40.3%

OTHER INGREDIENTS:	59.7%
---------------------------------	-------

TOTAL:	100.0%
---------------------	--------

Contains 4 pounds active ingredient per gallon (480 grams per liter).

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS

For Medical Emergencies, Call (877) 325-1840

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

EPA REG. NO. 228-723



Net Contents
64 Fl Oz (1.89 L)
Nonrefillable Container

Manufactured for
Nufarm Americas Inc.
11901 S. Austin Avenue
Alsip, IL 60803

 **nufarm**

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything to an unconscious person.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call poison control center or doctor for treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, viton ≥ 14 mils
- Shoes plus socks

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and oysters and shrimp. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. **Groundwater Advisory:** This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. **Surface Water Advisory:** This chemical may contaminate water through drift or spray in wind. This chemical has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this chemical. A level, well maintained vegetative buffer strip between areas to which this chemical is applied and surface water features such as ponds, streams, and springs, will reduce the potential for contamination of water from rainfall runoff. Runoff of such as when soils are saturated and/or significant rainfall is forecast in the next 48 hours. Sound erosion control practices will reduce this chemical's contribution to surface water contamination.

Physical or Chemical Hazards

Do not use or store near heat or open flame. Do not use with or store near any oxidizing agents, hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, viton ≥ 14 mils
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR DISEASE CONTROL OR PLANT INJURY**

USE RESTRICTIONS

- **Do not apply Verdato with any type of aircraft.**
- Not registered for use by the State of Hawaii.
- Not registered for use by the State of New York for use in Nassau and Suffolk counties.

**OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF
AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS,
MARSHES OR NATURAL PONDS, ESTUARIES, AND COMMERCIAL FISH FARM PONDS.**

- Do not apply within 75 ft of bodies of water such as lakes reservoirs rivers, permanent streams natural ponds, marshes or estuaries.
- For all plantings within 150 ft of bodies of water as described above, spray crops from outside the planting away from the bodies of water.
- Shut off the sprayer when at row ends.
- Spray the last three rows windward of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas.
- Do not cultivate within 10 ft of aquatic areas as to allow a vegetative filter strip.
- Do not apply when weather conditions favor drift to aquatic areas. Do not apply when gusts or sustained winds exceed 10 mph.
- Do not apply during a temperature inversion. Mist or fog may indicate the presence of an inversion in humid areas.

RESISTANCE MANAGEMENT

For resistance management, this product contains a Group 12 fungicide. Any fungal population may contain individuals naturally resistant to this product and other Group 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 12 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as, cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact Nufarm at 855-280-6609 to report suspected resistance.
- Contact your local extension specialist or certified pest control advisor for any additional pesticide resistance-management and/or IPM recommendations for specific sites and pathogens.

MIXING PROCEDURES

1. Prepare no more spray mixture than is needed for the immediate operation.
2. Thoroughly clean spray equipment before using this product.
3. Vigorous agitation is necessary to maintain uniformity of the spray mixture.
4. Maintain maximum agitation throughout the spraying operation.
5. Do not let the spray mixture stand overnight in the spray tank.
6. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

To determine the physical compatibility of Verdato with other products, use a jar test as described below.

Jar Compatibility Test

1. Using a quart Jar, add the proportionate amounts of the products to 1 qt of water.
2. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last.
3. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily it is physically compatible.

4. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.
5. If using Verdato in a tank mixture, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations which appear on the tank-mix product label.
6. Do not exceed the dosage rates on any label.
7. Follow the most restrictive label restrictions.
8. Do not mix this product with any other product whose label prohibits such mixing.
9. Tank mixtures are permitted only in those states where the tank mix partner is registered.

Verdato Alone:

1. Add 2/3 of the required amount of water to the spray or mixing tank.
2. With the agitator running, add the labeled amount of Verdato into the spray tank all at once.
3. Continue agitation while adding the remainder of the water.
4. Begin application of the spray solution after the material has completely dispersed into the mix water.
5. Maintain agitation until all of the mixture has been applied.

Verdato + Tank Mixtures:

Verdato is usually compatible with Strider®, Torque®, Tourney, Adorn, Spectro 90™, 3336®, Alude™, and other commonly used fungicides, insecticides and foliar nutrient products. However, the physical and biological compatibility of Verdato with tank-mix partners should be tested before use. If in doubt, run a jar compatibility test or consult with Nufarm company representatives, other users, university or extension personnel before proceeding.

1. Add 2/3 of the required amount of water to the spray or mixing tank.
2. With the agitator running add the labeled amount of Verdato into the tank all at once.
3. Continue agitation while adding the remainder of the water.
4. Allow Verdato to dissolve and the product to completely disperse into the mix water. Then add the desired amount of other products recommended for tank mixture and allow them to become completely dispersed.
5. Continue agitation to maintain a uniform suspension until all of the spray solution has been applied.

6. Other wettable powders or water dispersible granules should be added to the water in the tank next, followed by flowable products, and emulsifiable concentrates added last.
7. Provide sufficient mechanical or bypass agitation during mixing and application.

SPRAY DRIFT ADVISORIES:

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE:

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for spray drift will be greater if applications are made improperly or under unfavorable environmental conditions.

SHIELDED SPRAYERS:

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY:

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS:

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND:

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

TURFGRASS

Verdato is a protectant fungicide for control of certain diseases of turfgrass. Verdato provides control of brown patch, leaf spot, yellow patch, summer patch, anthracnose, gray leaf spot, bentgrass dead spot, and pink and gray snow mold when used in an integrated, preventive disease management program.

DO NOT USE ON TURFGRASS IN HAWAII.

DO NOT USE ON TURFGRASS IN NASSAU AND SUFFOLK COUNTIES IN NEW YORK.

Use Verdato as a foliar spray in an integrated, preventive disease management program alone or in combination with other fungicides, such as Affirm, Tourney EZ , Torque, or Traction.

- Apply in sufficient water to ensure thorough coverage.
- Apply after mowing.
- For control of foliar diseases, allow sprayed area to completely dry before irrigation.
- Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.
- For optimum turfgrass quality and disease control, use Verdato in conjunction with turfgrass management practices that promote good plant health and optimum disease control.
- Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease is essential to determine the best control measures.

Notes: (1) To avoid possible illegal residues, do not graze animals on treated areas, and (2) Do not feed clippings from treated areas to livestock or poultry. (3) **Note:** Do not apply more than 1.5 fl oz/1000 sq ft/year (4 lbs product/A/year) on turfgrass.

Turfgrass Disease Control with Verdato

Disease	Rate product/1,000 sq ft	Application Interval	Instructions
Brown Patch (<i>Rhizoctonia solani</i>)	0.20 – 0.25 fl oz (0.27-0.34 lbs ai/A)	7 days	Begin applications prior to disease development. For extended and more broad spectrum disease control, including dollar spot, tank mix 0.20-0.50 fl oz of Verdato with 0.18-0.37 fl oz Tourney EZ or 1.3 fl oz Traction, or 0.6 fl oz Torque per 1,000 sq ft.
	0.50 fl oz (0.68 lbs ai/A)	14 days	
Brown Ring Patch (<i>Waitea circinata</i> var. <i>circinata</i>) Rhizoctonia Leaf and Sheath Spot (<i>Waitea circinata</i> var. <i>zeae</i> and <i>Rhizoctonia zeae</i>)	0.25-0.50 fl oz (0.34-0.68 lbs ai/A)	7-14 days	Apply when conditions are favorable for disease development. For best control, use as a component of a preventive disease control program including Torque, Tourney EZ, or Affirm.
Bentgrass Dead Spot (<i>Ophiosphaerella agrostis</i>)	0.30-0.50 fl oz (0.41-0.68 lbs ai/A)	14 days	Begin applications prior to disease development.
Anthrachnose (<i>Colletotrichum graminicola</i>)	0.25-0.50 fl oz (0.34-0.68 lbs ai/A)	14 days	Begin applications prior to disease development. For more broad spectrum disease control, use as a component of a preventive disease control program.
Leaf Spot (<i>Bipolaris</i> spp., <i>Drechslera</i> spp.)	0.25-0.50 fl oz (0.34-0.68 lbs ai/A)	14-21 days	Apply when conditions are favorable for disease development.

(continued)

Turfgrass Disease Control with Verdato *(continued)*

Disease	Rate product/1,000 sq ft	Application Interval	Instructions
Summer Patch (<i>Magnaporthe poae</i>)	0.50 fl oz (0.68 lbs ai/A)	14 days	Apply when conditions are favorable for disease development. For best disease control, tank mix 0.30-0.50 fl oz of Verdato with 0.37 fl oz Tourney EZ or 0.6 fl oz Torque per 1,000 sq ft.
Yellow Patch (<i>Rhizoctonia cerealis</i>)	0.50 fl oz (0.68 lbs ai/A)	14 days	Apply when yellow patch rings are first observed in the spring or fall. A second application can be made 14 days later. For best disease control, tank mix Verdato with Tourney EZ or Torque.
Snow Mold: Gray (<i>Typhula incarnata</i> , <i>T. ishikariensis</i>) Pink (<i>Microdochium nivale</i>)	0.50 fl oz (0.68 lbs ai/A)	Late Fall	Apply one to two applications in late fall before snow cover. Do not apply on top of snow. For best disease control, tank mix with one to two registered snow mold products. For example, mix with a combination of Tourney EZ, Torque or Traction. Verdato can be used when tank mixing with Tourney EZ, Torque, and/or Traction when disease conditions are expected to be light to moderate.

(continued)

Turfgrass Disease Control with Verdato *(continued)*

Disease	Rate product/1,000 sq ft	Application Interval	Instructions
Microdochium Patch (<i>Microdochium nivale</i>)	0.25 – 0.50 fl oz (0.34-0.68 lbs ai/A)	7-14 days	For the control of Microdochium Patch without snow cover. Under heavy infestations combine with Tourney EZ or Traction or Torque. Repeat applications can be made at 7-14 day intervals when conditions are favorable for disease development.
Gray Leafspot (<i>Pyricularia grisea</i>)	0.25 – 0.50 fl oz (0.34-0.68 lbs ai/A)	14 days	Apply when conditions are favorable for disease development. For best control, use as a component of a preventive disease control program.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool place. **DO NOT** store in direct sunlight. Protect from freezing temperatures.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Nonrefillable Containers 5 Gallons or Less

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

RV101725[2]

Verdato is a trademark of Nufarm Americas, Inc. All other trademarks are the property of their respective owners.

Verdato™

Active Ingredient:

Fludioxonil..... 40.3%

Other Ingredients:..... 59.7%**Total:**..... 100.0%

Contains 4 pounds active ingredient per gallon (480 grams per liter).

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

**SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY
STATEMENTS**

For Medical Emergencies, Call (877) 325-1840

**For Chemical Spill, Leak, Fire, or Exposure,
Call CHEMTREC (800) 424-9300**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

EPA REG. NO. 228-723

(continued)

**Net Contents 64 Fl Oz (1.89 L)
Nonrefillable Container**

FLUDIOXONIL GROUP 12 FUNGICIDE

FIRST AID *(continued)*

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. **HOT LINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool place. **DO NOT** store in direct sunlight. Protect from freezing temperatures.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Nonrefillable Containers 5 Gallons or Less

Nonrefillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

15853000

RV101725[2]

Manufactured for
Nufarm Americas Inc.
11901 S. Austin Avenue
Alsip, IL 60803