

FIFRA SECTION 24(C) SPECIAL LOCAL NEED (SLN) LABEL

FLUTOLANIL GROUP 7 FUNGICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

MONCUT® SC FUNGICIDE

TO CONTROL *RHIZOCTONIA SPP.* IN FIELD-GROWN FLOWER BULBS

EPA Reg. No. 71711-28

EPA SLN No. WA-210003

This label for Moncut SC Fungicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2030.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of application.
- FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, AND PRECAUTIONS ON THIS SLN LABEL AND THE EPA REGISTERED LABEL.

DIRECTIONS FOR USE

CROP AND DISEASE CONTROLLED

For in-furrow use of Moncut SC Fungicide on field grown flower bulbs (including but not limited to iris, lilies, tulips) to control *Rhizoctonia spp.* (including gray bulb rot).

USE RATE AND APPLICATION METHOD

Apply 38.5 – 154 fl. oz. (1.14 - 4.57 lbs. ai) of Moncut SC Fungicide per acre (see Application Rate Chart below) as an in-furrow spray by directing spray uniformly over the tops of planted bulbs in an 8-to-18-inch band prior to covering with soil (hilling). Apply in sufficient amount of water which may range from 100 to 150 gallons of spray volume per acre. Use highest rate when disease conditions are expected to be severe or if field has a history of *Rhizoctonia* infestation.

In-Furrow Application Rate Chart					
Row Spacing	Rate of Moncut SC Fungicide per acre (fl. oz. product)				
	38.5	64	96	128	154
	Rate of Moncut SC Fungicide per 1000 Row Feet (fl. oz. product)				
42" Row	3.1	5.1	7.7	10.3	12.4
40" Row	2.9	4.9	7.3	9.8	11.8
38" Row	2.8	4.7	7.0	9.3	11.2
36" Row	2.7	4.4	6.6	8.8	10.6
34" Row	2.5	4.2	6.2	8.3	10.0

SPRAY DRIFT MANAGEMENT

Spray equipment and weather affect spray drift. Consider all factors when making application decisions. Where states have more stringent regulations, they must be observed. Avoiding spray drift is the responsibility of the applicator or grower. To reduce the potential for drift, the application equipment must be set to apply medium or larger droplets (i.e., ASABE Standard 572) with corresponding spray pressure. Use high flow rate nozzles to apply the highest practical spray volume, using the appropriate droplet size to ensure adequate canopy distribution, coverage, and penetration. With most nozzle types, narrow spray angles produce larger droplets. Follow the nozzle manufacturer's directions on pressure, orientation, spray volume, etc., in order to minimize drift and optimize coverage and control.

USE PRECAUTIONS AND RESTRICTIONS

- Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
 - For use rates over 131 fl. oz. / acre (3.88 lb. ai / acre) do not apply within 100 feet of any estuary or marine habitat or within 100 feet of any ditches, drainage tiles, or other waterways that drain directly into (within ½ mile) estuaries and marine habitats.
 - Make no more than 4 applications of Moncut SC Fungicide per year to ornamental plantings.
 - Apply no more than a combined total of 261 fl. oz. / acre (7.76 lb. ai / acre) during a single growing season.
 - Do not use oil as carrier or add other additives to the finished spray solution.
 - For use in accordance with this SLN label **do not** apply this product through any type of irrigation system.
-
- Do not plant rotational crops other than brassica (cole) leafy vegetables, peanuts, potatoes, soybeans, or turnip greens for 12 months following last application of Moncut SC Fungicide, with the following exceptions:
 - Wheat may be planted 30 days after the last application.
 - Leafy vegetables (such as lettuce, spinach, or celery) or small grain crops other than wheat (such as barley, rye, or oats) may be planted 150 days after the last application.
 - Corn (such as field, sweet, or popcorn) or sorghum may be planted 240 days after the last application.

WSDA Aquatic Advisory: This pesticide is toxic to fish and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater.

This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of flutolanil from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling a cleaned container is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA website under the Waste Pesticide Program. Cleaned containers may also be disposed of in a sanitary landfill if permitted by the county. Burning is not a legal method of container disposal in Washington.

©2025 Nichino America, Inc. Moncut is a registered trademark of Nichino America, Inc.



NICHINO

Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700