



US012390586B2

(12) **United States Patent**
Jacobson et al.

(10) **Patent No.:** **US 12,390,586 B2**

(45) **Date of Patent:** ***Aug. 19, 2025**

(54) **CONCURRENT INFUSION WITH COMMON LINE AUTO FLUSH**

(71) Applicant: **ICU Medical, Inc.**, San Clemente, CA (US)

(72) Inventors: **James Duane Jacobson**, Lindenhurst, IL (US); **Gerald William Brann**, Antioch, IL (US)

(73) Assignee: **ICU Medical, Inc.**, San Clemente, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 230 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **17/493,692**

(22) Filed: **Oct. 4, 2021**

(65) **Prior Publication Data**

US 2022/0176037 A1 Jun. 9, 2022

Related U.S. Application Data

(63) Continuation of application No. 17/114,359, filed on Dec. 7, 2020, now Pat. No. 11,135,360.

(51) **Int. Cl.**
A61M 5/168 (2006.01)
A61M 5/14 (2006.01)

(Continued)

(52) **U.S. Cl.**
CPC **A61M 5/16827** (2013.01); **A61M 5/1407** (2013.01); **G16H 20/17** (2018.01);
(Continued)

(58) **Field of Classification Search**
CPC A61M 5/14; A61M 5/1407; A61M 5/1409; A61M 5/142; A61M 5/145; A61M 5/168;
(Continued)

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,401,337 A 9/1968 Beusman et al.
3,484,681 A 12/1969 Grady, Jr. et al.
(Continued)

FOREIGN PATENT DOCUMENTS

AU 2013216679 9/2013
BR PI0704229-9 11/2009
(Continued)

OTHER PUBLICATIONS

Alaadeen et al., "Total Parenteral Nutrition-Associated Hyperglycemia Correlates with Prolonged Mechanical Ventilation and Hospital Stay in Septic Infants", *Journal of Pediatric Surgery*, Jan. 2006, vol. 41, No. 1, pp. 239-244.

(Continued)


Primary Examiner — Scott J Medway

(74) *Attorney, Agent, or Firm* — Knobbe, Martens, Olson & Bear, LLP

(57) **ABSTRACT**

An infusion pump system and method provide concurrent infusion with common line auto flush. The infusion pump system has a first reservoir, a second reservoir, a junction, a mixing chamber, a common line having one end in fluid connection with the mixing chamber and having a terminal fluid delivery end, and an infusion pump. The method includes infusing the first fluid at a first rate along a first flow path; determining a common line flush volume value for the common line; switching to a concurrent infusion mode to drive a combination of the first fluid and the second fluid at the first rate along a second flow path including the common line; monitoring a volume of the combination of the first and second fluids driven at the first rate; and driving the combination of the first and second fluids at a combined rate along the second flow path when the monitored volume is equal to or greater than the common line flush volume value.

23 Claims, 11 Drawing Sheets



- (51) **Int. Cl.**
G06K 7/10 (2006.01)
G06K 19/07 (2006.01)
G16H 20/17 (2018.01)
G16H 40/40 (2018.01)
G16H 40/67 (2018.01)
- (52) **U.S. Cl.**
CPC **G16H 40/40** (2018.01); **G16H 40/67** (2018.01); **A61M 2005/1403** (2013.01); **A61M 2205/3334** (2013.01); **A61M 2205/502** (2013.01); **A61M 2205/52** (2013.01); **G06K 7/10297** (2013.01); **G06K 2007/10504** (2013.01); **G06K 19/0723** (2013.01)
- (58) **Field of Classification Search**
CPC A61M 5/16804; A61M 5/16827; A61M 5/16877; A61M 5/172; G06F 19/3468
See application file for complete search history.
- (56) **References Cited**
U.S. PATENT DOCUMENTS
- | | | | | | | | |
|-----------|---|---------|------------------------|-----------|---|---------|---------------------|
| 3,699,320 | A | 10/1972 | Zimmerman et al. | 4,217,993 | A | 8/1980 | Jess et al. |
| 3,727,074 | A | 4/1973 | Keller et al. | 4,240,294 | A | 12/1980 | Grande |
| 3,731,679 | A | 5/1973 | Wilhelmson et al. | 4,240,438 | A | 12/1980 | Updike et al. |
| 3,768,084 | A | 10/1973 | Haynes | 4,244,365 | A | 1/1981 | McGill |
| 3,770,354 | A | 11/1973 | Tsuruta et al. | 4,256,437 | A | 3/1981 | Brown |
| 3,778,702 | A | 12/1973 | Finger | 4,261,356 | A | 4/1981 | Turner et al. |
| 3,806,821 | A | 4/1974 | Niemeyer et al. | 4,264,861 | A | 4/1981 | Radu et al. |
| 3,838,565 | A | 10/1974 | Carlyle | 4,265,240 | A | 5/1981 | Jenkins |
| 3,854,038 | A | 12/1974 | McKinley | 4,270,532 | A | 6/1981 | Franetzki et al. |
| 3,886,459 | A | 5/1975 | Hufford et al. | 4,277,226 | A | 7/1981 | Archibald et al. |
| 3,890,554 | A | 6/1975 | Yoshitake et al. | 4,278,085 | A | 7/1981 | Shim |
| 3,894,431 | A | 7/1975 | Muston et al. | 4,280,495 | A | 7/1981 | Lampert |
| 3,898,637 | A | 8/1975 | Wolstenholme | 4,282,872 | A | 8/1981 | Franetzki et al. |
| 3,901,231 | A | 8/1975 | Olson | 4,286,202 | A | 8/1981 | Clancy et al. |
| 3,909,693 | A | 9/1975 | Yoshitake et al. | 4,290,346 | A | 9/1981 | Bujan |
| 3,910,701 | A | 10/1975 | Henderson | 4,291,692 | A | 9/1981 | Bowman et al. |
| 3,911,343 | A | 10/1975 | Oster | 4,292,405 | A | 9/1981 | Mascoli |
| 3,919,608 | A | 11/1975 | Usami et al. | 4,298,357 | A | 11/1981 | Permic |
| 3,921,622 | A | 11/1975 | Cole | 4,308,866 | A | 1/1982 | Jeliffe |
| 3,930,404 | A | 1/1976 | Ryden, Jr. | 4,312,341 | A | 1/1982 | Zissimopoulos |
| 3,933,431 | A | 1/1976 | Trujillo et al. | 4,319,568 | A | 3/1982 | Tregoning |
| 3,935,876 | A | 2/1976 | Massie et al. | 4,322,201 | A | 3/1982 | Archibald |
| 3,944,963 | A | 3/1976 | Hively | 4,323,849 | A | 4/1982 | Smith |
| 3,966,358 | A | 6/1976 | Heimes et al. | 4,324,662 | A | 4/1982 | Schnell |
| 3,971,980 | A | 7/1976 | Jungfer et al. | 4,328,800 | A | 5/1982 | Marx |
| 3,974,681 | A | 8/1976 | Namery | 4,328,801 | A | 5/1982 | Marx |
| 3,974,683 | A | 8/1976 | Martin | 4,333,045 | A | 6/1982 | Oltendorf |
| 3,985,467 | A | 10/1976 | Lefferson | 4,343,316 | A | 8/1982 | Jespersen |
| 3,990,444 | A | 11/1976 | Vial | 4,344,429 | A | 8/1982 | Gupton et al. |
| 3,997,888 | A | 12/1976 | Kremer | 4,346,707 | A | 8/1982 | Whitney et al. |
| 4,005,724 | A | 2/1977 | Courtot | 4,360,019 | A | 11/1982 | Portner et al. |
| 4,014,206 | A | 3/1977 | Taylor | 4,366,384 | A | 12/1982 | Jensen |
| 4,038,982 | A | 8/1977 | Burke | 4,367,736 | A | 1/1983 | Gupton |
| 4,039,269 | A | 8/1977 | Pickering | 4,370,983 | A | 2/1983 | Lichtenstein et al. |
| 4,048,474 | A | 9/1977 | Olesen | 4,373,527 | A | 2/1983 | Fischell |
| 4,049,954 | A | 9/1977 | Da Costa Vieira et al. | 4,379,452 | A | 4/1983 | DeVries |
| 4,055,175 | A | 10/1977 | Clemens et al. | 4,381,005 | A | 4/1983 | Bujan |
| 4,057,228 | A | 11/1977 | Völker et al. | 4,384,578 | A | 5/1983 | Winkler |
| 4,068,521 | A | 1/1978 | Cosentino et al. | 4,385,247 | A | 5/1983 | Satomi |
| 4,078,562 | A | 3/1978 | Friedman | 4,391,598 | A | 7/1983 | Thompson |
| 4,089,227 | A | 5/1978 | Falgari et al. | 4,392,849 | A | 7/1983 | Petre et al. |
| 4,094,318 | A | 6/1978 | Burke | 4,394,862 | A | 7/1983 | Shim |
| 4,105,028 | A | 8/1978 | Sadlier et al. | 4,395,259 | A | 7/1983 | Prestele et al. |
| 4,114,144 | A | 9/1978 | Hyman | 4,397,194 | A | 8/1983 | Soltz |
| 4,151,845 | A | 5/1979 | Clemens | 4,399,362 | A | 8/1983 | Cormier et al. |
| 4,155,362 | A | 5/1979 | Jess | 4,407,659 | A | 10/1983 | Adam |
| 4,164,986 | A | 8/1979 | Eloy | 4,411,651 | A | 10/1983 | Schulman |
| 4,173,224 | A | 11/1979 | Marx | 4,418,565 | A | 12/1983 | St. John |
| 4,181,610 | A | 1/1980 | Shintani et al. | 4,432,699 | A | 2/1984 | Beckman et al. |
| 4,183,244 | A | 1/1980 | Kohn et al. | 4,432,761 | A | 2/1984 | Dawe |
| 4,195,515 | A | 4/1980 | Smoll | 4,432,762 | A | 2/1984 | Dawe |
| 4,210,138 | A | 7/1980 | Jess et al. | 4,443,218 | A | 4/1984 | Decant, Jr. et al. |
| 4,213,454 | A | 7/1980 | Shim | 4,444,546 | A | 4/1984 | Pazemenas |
| | | | | 4,447,191 | A | 5/1984 | Bilstad et al. |
| | | | | 4,447,224 | A | 5/1984 | Decant, Jr. et al. |
| | | | | 4,453,931 | A | 6/1984 | Pastrone |
| | | | | 4,457,751 | A | 7/1984 | Rodler |
| | | | | 4,463,301 | A | 7/1984 | Moriguchi et al. |
| | | | | 4,464,170 | A | 8/1984 | Clemens |
| | | | | 4,467,654 | A | 8/1984 | Murakami et al. |
| | | | | 4,468,222 | A | 8/1984 | Lundquist |
| | | | | 4,468,601 | A | 8/1984 | Chamran et al. |
| | | | | 4,469,481 | A | 9/1984 | Kobayashi |
| | | | | 4,475,666 | A | 10/1984 | Bilbrey et al. |
| | | | | 4,475,901 | A | 10/1984 | Kraegen et al. |
| | | | | 4,477,756 | A | 10/1984 | Moriguchi |
| | | | | 4,479,760 | A | 10/1984 | Bilstad et al. |
| | | | | 4,480,218 | A | 10/1984 | Hair |
| | | | | 4,480,483 | A | 11/1984 | McShane |
| | | | | 4,483,202 | A | 11/1984 | Ogua et al. |
| | | | | 4,487,601 | A | 12/1984 | Lindemann |
| | | | | 4,492,909 | A | 1/1985 | Hartwig |
| | | | | 4,496,346 | A | 1/1985 | Mosteller |
| | | | | 4,498,843 | A | 2/1985 | Schneider et al. |
| | | | | 4,501,531 | A | 2/1985 | Bilstad et al. |
| | | | | 4,504,263 | A | 3/1985 | Steuer |
| | | | | 4,507,112 | A | 3/1985 | Hillel |
| | | | | 4,510,266 | A | 4/1985 | Eertink |
| | | | | 4,513,796 | A | 4/1985 | Miller et al. |

(56)

References Cited

U.S. PATENT DOCUMENTS

4,515,584 A	5/1985	Abe et al.	4,785,969 A	11/1988	McLaughlin
4,519,792 A	5/1985	Dawe	4,786,800 A	11/1988	Kamen
4,521,212 A	6/1985	Ruschke	4,789,014 A	12/1988	DiGianfilippo
4,525,163 A	6/1985	Slavik et al.	4,797,655 A	1/1989	Orndal et al.
4,526,568 A	7/1985	Clemens et al.	4,803,389 A	2/1989	Ogawa et al.
4,526,574 A	7/1985	Pekkarinen	4,803,625 A	2/1989	Fu et al.
4,529,401 A	7/1985	Leslie et al.	4,818,186 A	4/1989	Pastrone et al.
4,533,350 A	8/1985	Danby et al.	4,820,281 A	4/1989	Lawler
4,543,955 A	10/1985	Schroeppel	4,821,558 A	4/1989	Pastrone et al.
4,551,134 A	11/1985	Slavik et al.	4,828,545 A	5/1989	Epstein et al.
4,553,958 A	11/1985	LeCocq	4,828,693 A	5/1989	Lindsay
4,559,036 A	12/1985	Wunsch	4,829,448 A	5/1989	Balding et al.
4,559,037 A	12/1985	Franetzki et al.	4,838,856 A	6/1989	Mulreany et al.
4,559,044 A	12/1985	Robinson	4,838,857 A	6/1989	Strowe et al.
4,559,454 A	12/1985	Kramer	4,840,542 A	6/1989	Abbott
4,565,500 A	1/1986	Jeensalute et al.	4,842,584 A	6/1989	Pastrone et al.
4,583,981 A	4/1986	Urquhart et al.	4,845,487 A	7/1989	Frantz et al.
4,587,473 A	5/1986	Turvey	4,846,792 A	7/1989	Bobo et al.
4,607,520 A	8/1986	Dam	4,850,805 A	7/1989	Madsen et al.
4,617,014 A	10/1986	Cannon et al.	4,851,755 A	7/1989	Fincher
4,624,661 A	11/1986	Arimond	4,854,324 A	8/1989	Hirschman et al.
4,627,835 A	12/1986	Fenton, Jr.	4,856,339 A	8/1989	Williams
4,633,878 A	1/1987	Bombardieri	4,857,048 A	8/1989	Simons et al.
4,634,426 A	1/1987	kamen	4,857,050 A	8/1989	Lentz et al.
4,634,427 A	1/1987	Hannula et al.	4,858,154 A	8/1989	Anderson et al.
4,636,144 A	1/1987	Abe et al.	4,863,425 A	9/1989	Slate et al.
4,637,813 A	1/1987	DeVries	4,865,584 A	9/1989	Epstein et al.
4,645,489 A	2/1987	Krumme	4,869,722 A	9/1989	Heyman
4,648,869 A	3/1987	Bobo, Jr.	4,874,359 A	10/1989	White et al.
4,652,260 A	3/1987	Fenton, Jr. et al.	4,881,413 A	11/1989	Georgi et al.
4,658,244 A	4/1987	Meijer	4,882,575 A	11/1989	Kawahara
4,668,216 A	5/1987	Martin	4,884,013 A	11/1989	Jackson et al.
4,668,945 A	5/1987	Aldrovandi et al.	4,884,065 A	11/1989	Crouse et al.
4,673,334 A	6/1987	Allington et al.	4,886,422 A	12/1989	Takeuchi et al.
4,673,389 A	6/1987	Archibald et al.	4,898,576 A	2/1990	Philip
4,676,776 A	6/1987	Howson et al.	4,898,578 A	2/1990	Rubalcaba, Jr.
4,677,359 A	6/1987	Enami et al.	4,906,103 A	3/1990	Kao
4,678,979 A	7/1987	Hori	4,908,017 A	3/1990	Howson et al.
4,678,998 A	7/1987	Muramatsu	4,908,019 A	3/1990	Urquhart et al.
4,679,562 A	7/1987	Luksha	4,910,475 A	3/1990	Lin
4,683,428 A	7/1987	Gete	4,919,595 A	4/1990	Likuski et al.
4,685,903 A	8/1987	Cable et al.	4,919,596 A	4/1990	Slate et al.
4,690,673 A	9/1987	Blomquist	4,925,444 A	5/1990	Orkin et al.
4,691,153 A	9/1987	Nishimura	4,927,411 A	5/1990	Pastrone et al.
4,692,145 A	9/1987	Weyant	4,930,358 A	6/1990	Motegi et al.
4,696,671 A	9/1987	Epstein et al.	4,936,820 A	6/1990	Dennehey
4,697,129 A	9/1987	Enami et al.	4,936,828 A	6/1990	Chiang
4,702,675 A	10/1987	Aldrovandi et al.	4,938,079 A	7/1990	Goldberg
4,705,506 A	11/1987	Archibald et al.	4,943,279 A	7/1990	Samiotes et al.
4,710,106 A	12/1987	Iwata et al.	4,946,439 A	8/1990	Eggers
4,714,462 A	12/1987	DiDomenico	4,947,856 A	8/1990	Beard
4,714,463 A	12/1987	Archibald et al.	4,950,235 A	8/1990	Slate et al.
4,718,576 A	1/1988	Tamura et al.	4,950,244 A	8/1990	Fellingham
4,720,636 A	1/1988	Benner	4,959,050 A	9/1990	Bobo, Jr.
4,722,224 A	2/1988	Scheller et al.	4,966,579 A	10/1990	Polaschegg
4,722,734 A	2/1988	Kolin	4,968,941 A	11/1990	Rogers
4,731,051 A	3/1988	Fischell	4,972,842 A	11/1990	Korten et al.
4,731,057 A	3/1988	Tanaka et al.	4,976,687 A	12/1990	Martin
4,737,711 A	4/1988	O'Hare	4,978,335 A	12/1990	Arthur, III
4,739,346 A	4/1988	Buckley	4,979,940 A	12/1990	Lapp et al.
4,741,732 A	5/1988	Crankshaw et al.	4,981,467 A	1/1991	Bobo et al.
4,741,736 A	5/1988	Brown	5,000,663 A	3/1991	Gorton
4,748,857 A	6/1988	Nakagawa	5,000,739 A	3/1991	Kulisz et al.
4,751,445 A	6/1988	Sakai	5,006,050 A	4/1991	Cooke et al.
4,756,706 A	7/1988	Kerns et al.	5,010,473 A	4/1991	Jacobs
4,758,228 A	7/1988	Williams	5,014,714 A	5/1991	Millay et al.
4,763,525 A	8/1988	Cobb	5,014,798 A	5/1991	Glynn
4,764,166 A	8/1988	Spani et al.	5,018,945 A	5/1991	D'Silva
4,764,697 A	8/1988	Christiaens	5,026,348 A	6/1991	Venegas
4,769,001 A	9/1988	Prince	5,028,857 A	7/1991	Taghezout
4,776,842 A	10/1988	Franetzki et al.	5,032,112 A	7/1991	Fairchild et al.
4,781,687 A	11/1988	Wall	5,034,004 A	7/1991	Crankshaw
4,784,576 A	11/1988	Bloom et al.	5,035,143 A	7/1991	Latimer et al.
4,785,184 A	11/1988	Bien et al.	5,040,699 A	8/1991	Gangemi
4,785,799 A	11/1988	Schoon et al.	5,041,086 A	8/1991	Koenig et al.
			5,043,706 A	8/1991	Oliver
			5,045,069 A	9/1991	Imparato
			5,049,047 A	9/1991	Polaschegg et al.
			5,052,230 A	10/1991	Lang

(56)

References Cited

U.S. PATENT DOCUMENTS

5,053,747	A	10/1991	Slate et al.	5,292,306	A	3/1994	Wynkoop et al.
5,055,761	A	10/1991	Mills	5,295,967	A	3/1994	Rondelet et al.
5,056,992	A	10/1991	Simons	5,298,021	A	3/1994	Sherer
5,058,161	A	10/1991	Weiss	5,303,585	A	4/1994	Lichte
5,059,171	A	10/1991	Bridge	5,304,126	A	4/1994	Epstein et al.
5,063,603	A	11/1991	Burt	5,304,216	A	4/1994	Wallace
5,064,412	A	11/1991	Henke et al.	5,308,333	A	5/1994	Skakoon
5,078,683	A	1/1992	Sancoff et al.	5,317,506	A	5/1994	Coutre et al.
5,084,663	A	1/1992	Olsson	5,319,363	A	6/1994	Welch et al.
5,084,828	A	1/1992	Kaufman et al.	5,319,979	A	6/1994	Abrahamson
5,088,981	A	2/1992	Howson et al.	5,321,392	A	6/1994	Skakoon et al.
5,096,385	A	3/1992	Georgi et al.	5,325,170	A	6/1994	Bornhop
5,097,505	A	3/1992	Weiss	5,325,728	A	7/1994	Zimmerman et al.
5,100,380	A	3/1992	Epstein et al.	5,328,460	A	7/1994	Lord et al.
5,102,392	A	4/1992	Sakai et al.	5,330,634	A	7/1994	Wong et al.
5,103,211	A	4/1992	Daoud et al.	5,333,497	A	8/1994	Braend et al.
5,104,374	A	4/1992	Bishko et al.	5,336,051	A	8/1994	Tamari
5,108,367	A	4/1992	Epstein et al.	5,338,157	A	8/1994	Blomquist
5,109,850	A	5/1992	Blanco et al.	5,342,298	A	8/1994	Michaels
5,116,203	A	5/1992	Nartwick et al.	5,343,734	A	9/1994	Maeda et al.
5,116,312	A	5/1992	Blakenship et al.	5,343,885	A	9/1994	Grant
5,116,316	A	5/1992	Sertic	5,346,466	A	9/1994	Yerlikaya et al.
5,123,275	A	6/1992	Daoud et al.	5,356,378	A	10/1994	Doan et al.
5,124,627	A	6/1992	Okada	5,359,271	A	10/1994	Husher
5,125,499	A	6/1992	Saathoff et al.	D352,778	S	11/1994	Irvin et al.
5,131,816	A	7/1992	Brown	5,364,346	A	11/1994	Schrezenmeir
5,132,603	A	7/1992	Yoshimoto	5,366,346	A	11/1994	Danby
5,153,827	A	10/1992	Coutre et al.	5,368,562	A	11/1994	Blomquist et al.
5,158,441	A	10/1992	Aid	5,374,865	A	12/1994	Yoshimura et al.
5,161,222	A	11/1992	Montejo et al.	5,376,070	A	12/1994	Purvis et al.
5,174,472	A	12/1992	Raque et al.	5,378,231	A	1/1995	Johnson et al.
5,176,631	A	1/1993	Koenig	5,382,232	A	1/1995	Hague et al.
5,176,646	A	1/1993	Kuroda	5,383,369	A	1/1995	Khuri-Yakub et al.
5,179,340	A	1/1993	Rogers	5,389,071	A	2/1995	Kawahara et al.
5,180,287	A	1/1993	Natwick et al.	5,389,078	A	2/1995	Zalesky et al.
5,181,910	A	1/1993	Scanlon	5,392,638	A	2/1995	Kawahara
5,186,057	A	2/1993	Everhart	5,394,732	A	3/1995	Johnson et al.
5,188,603	A	2/1993	Vaillancourt	5,395,320	A	3/1995	Padda et al.
5,190,522	A	3/1993	Wocicki et al.	5,399,171	A	3/1995	Bowman et al.
5,191,795	A	3/1993	Fellingham et al.	5,406,954	A	4/1995	Tomita
5,192,340	A	3/1993	Grant et al.	5,408,326	A	4/1995	Priestley
5,194,796	A	3/1993	Domeki et al.	5,415,528	A	5/1995	Ogden et al.
5,198,776	A	3/1993	Carr	5,417,119	A	5/1995	Smoll
5,200,090	A	4/1993	Ford	5,417,222	A	5/1995	Dempsey et al.
5,205,819	A	4/1993	Ross et al.	5,417,395	A	5/1995	Fowler et al.
5,206,522	A	4/1993	Danby et al.	5,418,443	A	5/1995	Kikuchi
5,207,642	A	5/1993	Orkin et al.	5,421,208	A	6/1995	Packard et al.
5,211,626	A	5/1993	Frank et al.	5,423,748	A	6/1995	Uhala
5,213,573	A	5/1993	Sorich et al.	5,423,749	A	6/1995	Merte et al.
5,215,450	A	6/1993	Tamari	5,423,759	A	6/1995	Campbell
5,216,597	A	6/1993	Beckers	5,428,284	A	6/1995	Kaneda et al.
5,219,099	A	6/1993	Spence et al.	5,429,485	A	7/1995	Dodge
5,219,327	A	6/1993	Okada	5,429,601	A	7/1995	Conley
5,221,268	A	6/1993	Barton et al.	5,429,602	A	7/1995	Hauser
5,229,713	A	7/1993	Bullock et al.	5,431,627	A	7/1995	Pastrone et al.
5,232,476	A	8/1993	Grant	5,434,508	A	7/1995	Ishida
5,233,571	A	8/1993	Wirtschafter	5,437,624	A	8/1995	Langley et al.
5,237,309	A	8/1993	Frantz et al.	5,444,316	A	8/1995	Ohya et al.
5,242,406	A	9/1993	Gross et al.	5,444,378	A	8/1995	Rogers
5,242,408	A	9/1993	Jhuboo et al.	5,445,621	A	8/1995	Poli et al.
5,243,982	A	9/1993	Möstl et al.	5,450,758	A	9/1995	Smoll
5,244,463	A	9/1993	Cordner, Jr. et al.	5,451,881	A	9/1995	Finger
5,244,568	A	9/1993	Lindsay et al.	5,455,423	A	10/1995	Mount et al.
5,254,096	A	10/1993	Rondelet et al.	5,455,851	A	10/1995	Chaco et al.
5,256,155	A	10/1993	Yerlikaya et al.	5,463,906	A	11/1995	Spani et al.
5,256,156	A	10/1993	Kern et al.	5,464,392	A	11/1995	Epstein et al.
5,256,157	A	10/1993	Samiotes et al.	5,465,082	A	11/1995	Chaco
5,257,206	A	10/1993	Hanson	5,469,851	A	11/1995	Lipschutz
5,260,665	A	11/1993	Goldberg	5,473,948	A	12/1995	Moss et al.
5,267,980	A	12/1993	Dirr et al.	5,480,294	A	1/1996	Di Perna et al.
5,274,316	A	12/1993	Evans et al.	5,482,438	A	1/1996	Anderson et al.
5,276,610	A	1/1994	Maeda et al.	5,485,408	A	1/1996	Blomquist
5,280,728	A	1/1994	Sato et al.	5,486,286	A	1/1996	Peterson et al.
5,283,510	A	2/1994	Tamaki et al.	5,489,265	A	2/1996	Montalvo et al.
5,287,851	A	2/1994	Beran et al.	5,495,566	A	2/1996	Kwatinetz
				5,496,273	A	3/1996	Pastrone et al.
				5,505,696	A	4/1996	Miki
				5,505,828	A	4/1996	Wong et al.
				5,507,288	A	4/1996	Bocker et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

5,507,412 A	4/1996	Ebert et al.	5,722,417 A	3/1998	Rudolph
5,520,637 A	5/1996	Pager et al.	5,728,074 A	3/1998	Castellano et al.
5,522,798 A	6/1996	Johnson et al.	5,728,948 A	3/1998	Bignell et al.
5,522,799 A	6/1996	Furukawa	5,733,257 A	3/1998	Stemby
5,527,630 A	6/1996	Nagata	5,733,259 A	3/1998	Valcke et al.
5,533,389 A	7/1996	Kamen et al.	5,734,464 A	3/1998	Gibbs
5,537,853 A	7/1996	Finburgh et al.	5,738,659 A	4/1998	Neer et al.
5,542,040 A	7/1996	Chang et al.	5,743,856 A	4/1998	Oka et al.
5,545,140 A	8/1996	Conero et al.	5,744,027 A	4/1998	Connell et al.
5,547,470 A	8/1996	Johnson et al.	5,744,929 A	4/1998	Miyazaki
5,551,850 A	9/1996	Williamson et al.	5,745,378 A	4/1998	Barker et al.
5,554,013 A	9/1996	Owens et al.	5,752,813 A	5/1998	Tyner et al.
5,554,115 A	9/1996	Thomas et al.	5,752,918 A	5/1998	Fowler et al.
5,558,638 A	9/1996	Evers et al.	5,752,919 A	5/1998	Schrimpf
5,562,615 A	10/1996	Nassif	5,755,691 A	5/1998	Hilborne
5,563,486 A	10/1996	Yamamoto et al.	5,758,643 A	6/1998	Wong et al.
5,572,105 A	11/1996	Nojima et al.	5,761,072 A	6/1998	Bardsley, Jr. et al.
5,573,502 A	11/1996	LeCocq et al.	5,764,034 A	6/1998	Bowman et al.
5,583,280 A	12/1996	Mo et al.	5,766,155 A	6/1998	Hyman et al.
5,584,667 A	12/1996	Davis	5,772,635 A	6/1998	Dastur et al.
5,584,806 A	12/1996	Amano	5,778,256 A	7/1998	Darbee
5,586,868 A	12/1996	Lawless et al.	5,781,442 A	7/1998	Engleson et al.
5,590,653 A	1/1997	Aida et al.	5,782,805 A	7/1998	Meinzer et al.
5,594,786 A	1/1997	Chaco et al.	5,788,669 A	8/1998	Peterson
5,600,073 A	2/1997	Hill	5,788,674 A	8/1998	McWilliams
5,601,420 A	2/1997	Warner et al.	5,789,923 A	8/1998	Shimoyama et al.
5,609,575 A	3/1997	Larson et al.	5,792,069 A	8/1998	Greenwald et al.
5,609,576 A	3/1997	Voss	5,793,211 A	8/1998	Shimoyama et al.
5,611,784 A	3/1997	Barresi et al.	5,795,327 A	8/1998	Wilson et al.
5,616,124 A	4/1997	Hague et al.	5,798,934 A	8/1998	Saigo et al.
5,620,312 A	4/1997	Hyman et al.	5,800,387 A	9/1998	Duffy et al.
5,620,608 A	4/1997	Rosa et al.	5,803,712 A	9/1998	Davis et al.
5,626,140 A	5/1997	Feldman et al.	5,803,917 A	9/1998	Butterfield
5,626,151 A	5/1997	Linden	5,805,455 A	9/1998	Lipps
5,626,563 A	5/1997	Dodge et al.	5,807,322 A	9/1998	Lindsey et al.
5,627,443 A	5/1997	Kimura et al.	5,810,770 A	9/1998	Chin et al.
5,628,309 A	5/1997	Brown	5,813,972 A	9/1998	Nazarian et al.
5,628,731 A	5/1997	Dodge et al.	5,814,004 A	9/1998	Tamari
5,630,710 A	5/1997	Tune et al.	5,814,015 A	9/1998	Gargano et al.
5,634,896 A	6/1997	Bryant et al.	5,816,779 A	10/1998	Lawless et al.
5,637,095 A	6/1997	Nason et al.	5,822,715 A	10/1998	Worthington et al.
5,640,075 A	6/1997	Brasseur et al.	5,827,179 A	10/1998	Lichter et al.
5,640,150 A	6/1997	Atwater	5,827,223 A	10/1998	Butterfield
5,643,212 A	7/1997	Coutre et al.	5,832,448 A	11/1998	Brown
5,648,710 A	7/1997	Ikeda	5,836,910 A	11/1998	Duffy et al.
5,649,536 A	7/1997	Ogura et al.	5,841,261 A	11/1998	Nojima et al.
5,651,775 A	7/1997	Walker et al.	5,841,284 A	11/1998	Takahashi
5,657,000 A	8/1997	Ellingboe	5,843,035 A	12/1998	Bowman
5,658,133 A	8/1997	Anderson et al.	5,848,971 A	12/1998	Fowler et al.
5,658,250 A	8/1997	Blomquist et al.	5,850,344 A	12/1998	Conkright
5,659,234 A	8/1997	Cresens	5,857,843 A	1/1999	Leason et al.
5,661,245 A	8/1997	Svoboda et al.	5,864,330 A	1/1999	Haynes
D384,052 S	9/1997	Kodosky	5,865,805 A	2/1999	Ziemba
5,662,612 A	9/1997	Niehoff	5,867,821 A	2/1999	Ballantyne et al.
5,665,065 A	9/1997	Colman et al.	5,871,465 A	2/1999	Vasko
5,669,877 A	9/1997	Blomquist	5,872,453 A	2/1999	Shimoyama et al.
5,672,154 A	9/1997	Sillén et al.	5,875,195 A	2/1999	Dixon
5,672,832 A	9/1997	Cucci et al.	5,882,300 A	3/1999	Malinouskas et al.
5,681,285 A	10/1997	Ford et al.	5,882,339 A	3/1999	Beiser et al.
5,681,286 A	10/1997	Niehoff	5,885,245 A	3/1999	Lynch et al.
5,685,844 A	11/1997	Marttila	5,889,379 A	3/1999	Yanagi et al.
5,685,866 A	11/1997	Lopez	5,891,051 A	4/1999	Han et al.
5,687,717 A	11/1997	Halpern et al.	5,894,209 A	4/1999	Takagi et al.
5,689,229 A	11/1997	Chaco et al.	5,897,493 A	4/1999	Brown
5,691,613 A	11/1997	Gutwillinger	5,897,498 A	4/1999	Canfield, II et al.
5,695,464 A	12/1997	Viallet	5,898,292 A	4/1999	Takemoto et al.
5,695,473 A	12/1997	Olsen	5,899,665 A	5/1999	Makino et al.
5,697,899 A	12/1997	Hillman et al.	5,901,150 A	5/1999	Jhuboo et al.
5,697,916 A	12/1997	Schrage	5,904,666 A	5/1999	DeDecker et al.
5,712,795 A	1/1998	Layman et al.	5,904,668 A	5/1999	Hyman et al.
5,713,856 A	2/1998	Eggers et al.	5,905,207 A	5/1999	Schalk
5,714,691 A	2/1998	Hill	5,906,598 A	5/1999	Giesier
5,718,562 A	2/1998	Lawless et al.	5,910,252 A	6/1999	Truitt et al.
5,718,569 A	2/1998	Holst	5,915,240 A	6/1999	Karpf
5,720,721 A	2/1998	Dumas et al.	5,920,263 A	7/1999	Huttenhoff et al.
			5,923,159 A	7/1999	Ezell
			5,924,074 A	7/1999	Evans
			5,927,349 A	7/1999	Martucci
			5,932,119 A	8/1999	Kaplan et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

5,932,987 A	8/1999	McLoughlin	6,248,067 B1	6/2001	Causey, III et al.
5,935,066 A	8/1999	Harris	6,250,132 B1	6/2001	Drzewiecki
5,935,099 A	8/1999	Peterson et al.	6,259,355 B1	7/2001	Chaco et al.
5,935,106 A	8/1999	Olsen	6,259,587 B1	7/2001	Sheldon et al.
5,938,634 A	8/1999	Packard	6,261,065 B1	7/2001	Nayak
5,938,636 A	8/1999	Kramer et al.	6,262,946 B1	7/2001	Khuri-Yakub et al.
5,941,846 A	8/1999	Duffy et al.	6,267,559 B1	7/2001	Mossman et al.
5,944,660 A	8/1999	Kimball et al.	6,267,725 B1	7/2001	Dubberstein et al.
5,947,911 A	9/1999	Wong et al.	6,269,340 B1	7/2001	Ford et al.
5,954,527 A	9/1999	Jhuboo et al.	6,270,455 B1	8/2001	Brown
5,954,696 A	9/1999	Ryan et al.	6,271,813 B1	8/2001	Palalau
5,956,023 A	9/1999	Lyle et al.	6,277,072 B1	8/2001	Bardy
5,956,501 A	9/1999	Brown	6,277,099 B1	8/2001	Strowe et al.
5,957,885 A	9/1999	Bollish et al.	6,280,380 B1	8/2001	Bardy
5,957,890 A	9/1999	Mann et al.	6,280,391 B1	8/2001	Olson et al.
5,971,594 A	10/1999	Sahai et al.	6,280,408 B1	8/2001	Sipin
5,973,497 A	10/1999	Bergk et al.	6,283,761 B1	9/2001	Joao
5,975,081 A	11/1999	Hood et al.	6,285,155 B1	9/2001	Maske et al.
5,989,222 A	11/1999	Cole et al.	6,312,378 B1	11/2001	Bardy
5,990,838 A	11/1999	Burns et al.	6,322,516 B1	11/2001	Masuda et al.
5,991,525 A	11/1999	Shah et al.	6,330,351 B1	12/2001	Yasunaga
5,993,393 A	11/1999	Ryan et al.	6,336,053 B1	1/2002	Beatty
5,994,876 A	11/1999	Canny et al.	6,337,675 B1	1/2002	Toffolo et al.
5,997,476 A	12/1999	Brown	6,345,539 B1	2/2002	Rawes et al.
6,000,828 A	12/1999	Leet	6,347,553 B1	2/2002	Morris et al.
6,003,006 A	12/1999	Colella et al.	6,349,740 B1	2/2002	Cho et al.
6,003,388 A	12/1999	Oeftering	6,358,225 B1	3/2002	Butterfield
6,012,034 A	1/2000	Hamparian et al.	6,358,387 B1	3/2002	Kopf-Sill et al.
6,017,318 A	1/2000	Gauthier et al.	6,362,591 B1	3/2002	Moberg
6,017,493 A	1/2000	Cambron	6,385,505 B1	5/2002	Lipps
6,021,392 A	2/2000	Lester et al.	6,386,050 B1	5/2002	Yin et al.
6,023,977 A	2/2000	Langdon et al.	6,394,958 B1	5/2002	Bratteli et al.
6,024,539 A	2/2000	Blomquist	6,396,583 B1	5/2002	Clare
6,027,441 A	2/2000	Cantu	D459,362 S	6/2002	Platz
6,028,412 A	2/2000	Shine et al.	6,398,760 B1	6/2002	Danby
6,032,676 A	3/2000	Moore	6,405,076 B1	6/2002	Taylor et al.
6,033,561 A	3/2000	Schoendorfer	6,408,679 B1	6/2002	Kline-Schoder et al.
6,036,017 A	3/2000	Bayliss, IV	6,409,699 B1	6/2002	Ash
6,068,612 A	5/2000	Bowman	6,413,238 B1	7/2002	Maget
6,068,615 A	5/2000	Brown et al.	6,416,291 B1	7/2002	Butterfield et al.
6,073,106 A	6/2000	Rozen et al.	6,418,334 B1	7/2002	Unger et al.
6,077,246 A	6/2000	Kullas et al.	6,418,535 B1	7/2002	Kulakowski et al.
6,083,206 A	7/2000	Molko	6,445,053 B1	9/2002	Cho
6,089,104 A	7/2000	Chang	6,456,245 B1	9/2002	Crawford
6,104,295 A	8/2000	Gaisser et al.	6,457,346 B1	10/2002	Kline-Schoder et al.
6,110,152 A	8/2000	Kovelman	6,463,785 B1	10/2002	Kline-Schoder et al.
6,110,153 A	8/2000	Davis	6,467,331 B1	10/2002	Kline-Schoder et al.
RE36,871 E	9/2000	Epstein et al.	6,468,242 B1	10/2002	Wilson et al.
6,120,459 A	9/2000	Nitzan et al.	6,475,178 B1	11/2002	Krajewski
6,122,536 A	9/2000	Sun et al.	6,481,980 B1	11/2002	Vandlik
6,142,008 A	11/2000	Cole et al.	6,482,158 B2	11/2002	Mault
6,150,942 A	11/2000	O'Brien	6,482,185 B1	11/2002	Hartmann
6,157,914 A	12/2000	Seto et al.	6,485,263 B1	11/2002	Bryant et al.
6,158,288 A	12/2000	Smith	6,485,418 B2	11/2002	Yasushi et al.
6,158,965 A	12/2000	Butterfield et al.	6,485,465 B2	11/2002	Moberg et al.
6,159,147 A	12/2000	Lichter et al.	6,487,916 B1	12/2002	Gomm et al.
6,159,186 A	12/2000	Wickham et al.	6,489,896 B1	12/2002	Platt
6,164,921 A	12/2000	Moubayed et al.	6,494,694 B2	12/2002	Lawless et al.
6,168,561 B1	1/2001	Cantu	6,494,831 B1	12/2002	Koritzinsky
6,178,827 B1	1/2001	Feller	6,497,680 B1	12/2002	Holst et al.
6,182,667 B1	2/2001	Hanks et al.	6,503,221 B1	1/2003	Briggs
6,186,141 B1	2/2001	Pike et al.	6,512,944 B1	1/2003	Kovtun et al.
6,189,105 B1	2/2001	Lopes	6,516,667 B1	2/2003	Broad et al.
6,192,752 B1	2/2001	Blaine	6,517,482 B1	2/2003	Eiden et al.
6,195,589 B1	2/2001	Ketcham	6,519,569 B1	2/2003	White et al.
6,202,711 B1	3/2001	Martucci	6,529,751 B1	3/2003	Van Driel et al.
6,203,528 B1	3/2001	Deckert	6,531,708 B1	3/2003	Malmstrom
6,208,107 B1	3/2001	Maske et al.	6,539,315 B1	3/2003	Adams et al.
6,212,936 B1	4/2001	Meisberger	D473,238 S	4/2003	Cockerill
6,213,972 B1	4/2001	Butterfield	6,540,672 B1	4/2003	Simonsen et al.
6,231,320 B1	5/2001	Lawless et al.	6,544,212 B2	4/2003	Galley et al.
6,234,176 B1	5/2001	Domae et al.	6,544,228 B1	4/2003	Heitmeier
6,236,326 B1	5/2001	Murphy et al.	6,558,125 B1	5/2003	Futterknecht
6,237,398 B1	5/2001	Porat et al.	6,558,351 B1	5/2003	Steil et al.
6,241,704 B1	6/2001	Peterson et al.	6,562,012 B1	5/2003	Brown et al.
			6,564,825 B2	5/2003	Lowery et al.
			6,565,509 B1	5/2003	Say et al.
			6,568,416 B2	5/2003	Tucker et al.
			6,572,542 B1	6/2003	Houben et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

6,572,545 B2	6/2003	Knobbe et al.	6,852,104 B2	2/2005	Blomquist
6,572,576 B2	6/2003	Brugger et al.	6,854,338 B2	2/2005	Khuri-Yakub et al.
6,578,422 B2	6/2003	Lam et al.	6,857,318 B1	2/2005	Silber et al.
6,578,435 B2	6/2003	Gould et al.	6,869,425 B2	3/2005	Briggs et al.
6,581,117 B1	6/2003	Klein et al.	6,873,268 B2	3/2005	Lebel et al.
RE38,189 E	7/2003	Walker et al.	6,883,376 B2	4/2005	He
6,585,675 B1	7/2003	O'Mahony et al.	6,885,881 B2	4/2005	Leonhardt
6,589,229 B1	7/2003	Connelly et al.	6,887,216 B2	5/2005	Hochman et al.
6,589,792 B1	7/2003	Malachowski	6,898,301 B2	5/2005	Iwanaga
6,599,281 B1	7/2003	Struys et al.	6,907,361 B2	6/2005	Molenaar
6,599,282 B2	7/2003	Burko	6,907,792 B2	6/2005	Ohnishi
6,602,191 B2	8/2003	Quy	6,915,170 B2	7/2005	Engleson et al.
6,605,072 B2	8/2003	Struys et al.	6,920,795 B2	7/2005	Bischoff et al.
6,606,047 B1	8/2003	Börjesson et al.	6,923,763 B1	8/2005	Kovatchev et al.
6,609,047 B1	8/2003	Lipps	6,928,338 B1	8/2005	Buchser et al.
6,615,674 B2	9/2003	Ohnishi	6,929,619 B2	8/2005	Fago et al.
6,616,633 B1	9/2003	Butterfield et al.	6,929,751 B2	8/2005	Bowman
6,617,564 B2	9/2003	Ockerse et al.	6,932,114 B2	8/2005	Sparks
6,618,916 B1	9/2003	Eberle et al.	6,932,796 B2	8/2005	Sage et al.
6,622,542 B2	9/2003	Derek	6,935,192 B2	8/2005	Sobek et al.
6,622,561 B2	9/2003	Lam et al.	6,936,029 B2	8/2005	Mann et al.
D481,121 S	10/2003	Evans	6,941,005 B2	9/2005	Lary et al.
6,629,449 B1	10/2003	Kline-Schoder et al.	6,942,636 B2	9/2005	Holst et al.
6,634,233 B2	10/2003	He	6,945,954 B2	9/2005	Hochman et al.
6,640,246 B1	10/2003	Gardy, Jr. et al.	6,958,705 B2	10/2005	Lebel et al.
6,641,533 B2	11/2003	Causey, III et al.	6,964,204 B2	11/2005	Clark et al.
6,641,541 B1	11/2003	Lovett et al.	6,973,374 B2	12/2005	Ader
6,648,861 B2	11/2003	Platt et al.	6,974,437 B2	12/2005	Lebel et al.
6,652,455 B1	11/2003	Kocher	6,975,922 B2	12/2005	Duncan et al.
6,653,937 B2	11/2003	Nelson et al.	6,978,779 B2	12/2005	Haveri et al.
6,659,980 B2	12/2003	Moberg et al.	6,979,326 B2	12/2005	Mann et al.
D485,356 S	1/2004	Evans	6,981,960 B2	1/2006	Cho et al.
6,685,668 B1	2/2004	Cho et al.	6,984,218 B2	1/2006	Nayak et al.
6,685,678 B2	2/2004	Evans et al.	6,985,768 B2	1/2006	Hemming et al.
6,689,069 B2	2/2004	Bratteli et al.	6,985,870 B2	1/2006	Martucci et al.
6,689,091 B2	2/2004	Bui et al.	6,986,347 B2	1/2006	Hickle
6,692,241 B2	2/2004	Watanabe et al.	6,986,753 B2	1/2006	Bui
D487,574 S	3/2004	Glaser	6,997,905 B2	2/2006	Gillespie, Jr. et al.
6,716,004 B2	4/2004	Vandlik	6,997,920 B2	2/2006	Mann et al.
6,719,535 B2	4/2004	Rakestraw et al.	7,006,005 B2	2/2006	Nazarian et al.
6,721,582 B2	4/2004	Trepagnier et al.	7,017,623 B2	3/2006	Tribble et al.
6,722,211 B1	4/2004	Ciobanu et al.	7,021,148 B2	4/2006	Kuhn
6,725,200 B1	4/2004	Rost	7,025,743 B2	4/2006	Mann et al.
6,725,721 B2	4/2004	Venczel	7,029,455 B2	4/2006	Flaherty
6,731,989 B2	5/2004	Engleson et al.	7,029,456 B2	4/2006	Ware et al.
6,732,595 B2	5/2004	Lynnworth	7,059,184 B2	6/2006	Kanouda et al.
6,738,052 B1	5/2004	Manke et al.	7,060,059 B2	6/2006	Keith et al.
6,740,072 B2	5/2004	Starkweather et al.	7,069,793 B2	7/2006	Ishikawa et al.
6,741,212 B2	5/2004	Kralovec et al.	7,072,725 B2	7/2006	Bristol et al.
6,748,808 B2	6/2004	Lam et al.	7,074,209 B2	7/2006	Evans et al.
6,749,403 B2	6/2004	Bryant et al.	7,080,557 B2	7/2006	Adnan
6,752,787 B1	6/2004	Causey, III et al.	7,082,843 B2	8/2006	Clark et al.
6,753,842 B1	6/2004	Williams et al.	7,087,444 B2	8/2006	Wong et al.
6,759,007 B1	7/2004	Westberg	7,092,796 B2	8/2006	Vanderveen
6,760,643 B2	7/2004	Lipps	7,092,797 B2	8/2006	Gaines et al.
6,768,920 B2	7/2004	Lange	7,093,502 B2	8/2006	Kupnik et al.
6,773,412 B2	8/2004	O'Mahony	7,096,729 B2	8/2006	Repko et al.
6,780,156 B2	8/2004	Haueter et al.	7,103,419 B2	9/2006	Engleson et al.
6,783,328 B2	8/2004	Lucke et al.	7,104,763 B2	9/2006	Bouton et al.
6,785,573 B2	8/2004	Kovtun et al.	7,104,769 B2	9/2006	Davis
6,786,885 B2	9/2004	Hochman et al.	7,108,680 B2	9/2006	Rohr et al.
6,789,426 B2	9/2004	Yaralioglu et al.	7,109,878 B2	9/2006	Mann et al.
6,790,198 B1	9/2004	White et al.	7,115,113 B2	10/2006	Evans et al.
6,793,625 B2	9/2004	Cavallaro et al.	7,117,041 B2	10/2006	Engleson et al.
6,801,227 B2	10/2004	Bocionek et al.	7,137,964 B2	11/2006	Flaherty
6,805,671 B2	10/2004	Stergiopoulos et al.	7,141,037 B2	11/2006	Butterfield et al.
6,807,965 B1	10/2004	Hickle	7,152,490 B1	12/2006	Freund, Jr. et al.
6,809,653 B1	10/2004	Mann et al.	7,154,397 B2	12/2006	Zerhusen et al.
6,813,964 B1	11/2004	Clark et al.	7,161,488 B2	1/2007	Frasch
6,814,547 B2	11/2004	Childers	7,162,290 B1	1/2007	Levin
6,824,528 B1	11/2004	Faries	7,162,927 B1	1/2007	Selvan et al.
6,830,558 B2	12/2004	Flaherty et al.	7,171,277 B2	1/2007	Engleson et al.
6,840,113 B2	1/2005	Fukumura et al.	7,171,992 B2	2/2007	DiGianfilippo et al.
6,846,161 B2	1/2005	Kline	7,174,789 B2	2/2007	Orr et al.
6,852,094 B2	2/2005	Beck	7,185,288 B2	2/2007	McKeever
			7,197,943 B2	4/2007	Lee et al.
			7,201,734 B2	4/2007	Hickle
			7,204,823 B2	4/2007	Estes et al.
			7,206,715 B2	4/2007	Vanderveen et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

7,213,009 B2	5/2007	Pestotnik	7,775,126 B2	8/2010	Eckhardt
7,220,240 B2	5/2007	Struys et al.	7,775,127 B2	8/2010	Wade
7,229,430 B2	6/2007	Hickle et al.	7,785,284 B2	8/2010	Baralsi et al.
7,230,529 B2	6/2007	Ketcherside	7,785,313 B2	8/2010	Mastrototaro
7,232,430 B2	6/2007	Carlisle	7,786,909 B2	8/2010	Udupa et al.
7,238,164 B2	7/2007	Childers et al.	7,806,886 B2	10/2010	Kanderian, Jr. et al.
7,247,154 B2	7/2007	Hickle	7,826,981 B2	11/2010	Goode, Jr. et al.
7,253,779 B2	8/2007	Greer et al.	7,847,276 B2	12/2010	Carlisle
7,254,425 B2	8/2007	Lowery et al.	7,860,583 B2	12/2010	Concurso et al.
7,258,534 B2	8/2007	Fathallah et al.	7,871,394 B2	1/2011	Halbert et al.
7,267,664 B2	9/2007	Rizzo	7,876,443 B2	1/2011	Bernacki
7,267,665 B2	9/2007	Steil et al.	7,895,053 B2	2/2011	Holland et al.
7,272,529 B2	9/2007	Hogan et al.	7,895,882 B2	3/2011	Carlisle
7,278,983 B2	10/2007	Ireland et al.	7,896,834 B2	3/2011	Smisson, III
7,291,123 B2	11/2007	Baraldi et al.	7,896,842 B2	3/2011	Palmroos et al.
7,293,461 B1	11/2007	Gimdt	7,905,710 B2	3/2011	Wang et al.
7,294,109 B2	11/2007	Lovett et al.	7,933,780 B2	4/2011	de la Hueraga
7,296,482 B2	11/2007	Schaffer et al.	7,938,817 B2	5/2011	Gelfand et al.
7,300,418 B2	11/2007	Zaleski	7,945,452 B2	5/2011	Fathallah et al.
7,305,883 B2	12/2007	Khuri-Yakub et al.	D642,195 S	7/2011	Marks
7,327,273 B2	2/2008	Hung et al.	7,976,508 B2	7/2011	Hoag
D563,986 S	3/2008	Lettau	7,981,073 B2	7/2011	Mollstam
7,338,470 B2	3/2008	Katz	7,981,082 B2	7/2011	Wang et al.
7,343,224 B2	3/2008	DiGianfilippo et al.	7,998,134 B2	8/2011	Fangrow
7,347,836 B2	3/2008	Peterson et al.	8,002,736 B2	8/2011	Patrick et al.
7,347,854 B2	3/2008	Shelton et al.	8,034,020 B2	10/2011	Dewey
7,354,420 B2	4/2008	Steil et al.	8,038,593 B2	10/2011	Friedman et al.
7,356,382 B2	4/2008	Vanderveen	8,065,161 B2	11/2011	Howard et al.
7,360,999 B2	4/2008	Nelson et al.	8,067,760 B2	11/2011	Carlisle
7,364,562 B2	4/2008	Braig et al.	8,075,514 B2	12/2011	Butterfield et al.
7,367,942 B2	5/2008	Grage et al.	8,075,546 B2	12/2011	Carlisle et al.
7,369,948 B1	5/2008	Ferenczi et al.	8,078,983 B2	12/2011	Davis et al.
7,384,410 B2	6/2008	Eggers et al.	8,121,857 B2	2/2012	Galasso et al.
7,397,166 B1	7/2008	Morgan et al.	8,149,131 B2	4/2012	Blomquist
7,398,183 B2	7/2008	Holland et al.	D659,709 S	5/2012	Eby
7,399,277 B2	7/2008	Saidara et al.	8,175,668 B1	5/2012	Nabutovsky et al.
7,402,153 B2	7/2008	Steil et al.	8,177,739 B2	5/2012	Cartledge et al.
7,402,154 B2	7/2008	Mendez	8,180,440 B2	5/2012	McCombie et al.
7,407,489 B2	8/2008	Mendez	8,185,322 B2	5/2012	Schroeder et al.
7,414,534 B1	8/2008	Kroll et al.	8,197,444 B1	6/2012	Bazargan et al.
7,415,895 B2	8/2008	Kurisaki et al.	8,219,413 B2	7/2012	Martinez et al.
7,426,443 B2	9/2008	Simon	8,221,395 B2	7/2012	Shelton et al.
7,430,675 B2	9/2008	Lee et al.	8,226,597 B2	7/2012	Jacobson et al.
7,447,566 B2	11/2008	Knauper et al.	8,231,578 B2	7/2012	Fathallah et al.
7,447,643 B1	11/2008	Olson	8,234,128 B2	7/2012	Martucci et al.
7,452,190 B2	11/2008	Bouton et al.	D667,452 S	9/2012	Wujcik
7,454,314 B2	11/2008	Holland et al.	D667,840 S	9/2012	Anzures
7,471,994 B2	12/2008	Ford et al.	8,271,106 B2	9/2012	Wehba et al.
7,477,997 B2	1/2009	Kaplit	8,287,514 B2	10/2012	Miller et al.
7,482,818 B2	1/2009	Greenwald et al.	8,291,337 B2	10/2012	Gannin et al.
7,483,756 B2	1/2009	Engleson et al.	8,313,308 B2	11/2012	Lawless et al.
7,490,021 B2	2/2009	Holland et al.	8,317,698 B2	11/2012	Lowery
7,491,187 B2	2/2009	Van Den Berghe et al.	8,317,750 B2	11/2012	Ware et al.
7,503,903 B2	3/2009	Carlisle et al.	8,317,752 B2	11/2012	Cozmi et al.
7,517,332 B2	4/2009	Tonelli et al.	8,318,094 B1	11/2012	Bayandorian et al.
7,523,401 B1	4/2009	Aldridge	8,340,792 B2	12/2012	Concurso et al.
D593,125 S	5/2009	Danton	8,347,731 B2	1/2013	Genosar
7,545,075 B2	6/2009	Huang et al.	8,359,338 B2	1/2013	Butterfield et al.
D596,195 S	7/2009	Wall	8,361,021 B2	1/2013	Wang et al.
7,556,616 B2	7/2009	Fathallah et al.	8,378,837 B2	2/2013	Wang et al.
7,561,986 B2	7/2009	Vanderveen et al.	8,388,598 B2	3/2013	Steinkogler
7,571,024 B2	8/2009	Duncan et al.	8,398,616 B2	3/2013	Budiman
7,605,730 B2	10/2009	Tomioka et al.	8,403,908 B2	3/2013	Jacobson et al.
7,614,310 B2	11/2009	Konzelmann	D679,727 S	4/2013	Abratowski
7,645,258 B2	1/2010	White et al.	8,409,164 B2	4/2013	Fangrow
7,654,127 B2	2/2010	Krulevitch et al.	8,449,524 B2	5/2013	Braig et al.
7,657,443 B2	2/2010	Crass	8,469,942 B2	6/2013	Kow et al.
7,668,731 B2	2/2010	Martucci et al.	8,477,307 B1	7/2013	Yufa et al.
7,678,048 B1	3/2010	Urbano et al.	8,494,879 B2	7/2013	Davis et al.
7,693,697 B2	4/2010	Westenskow et al.	8,504,179 B2	8/2013	Blomquist
7,699,806 B2	4/2010	Ware et al.	8,506,552 B2	8/2013	Rebours
7,705,727 B2	4/2010	Pestotnik	8,517,990 B2	8/2013	Teel et al.
D617,807 S	6/2010	Christie	8,518,021 B2	8/2013	Stewart et al.
D621,845 S	8/2010	Anzures	8,522,832 B2	9/2013	Lopez et al.
7,766,873 B2	8/2010	Moberg et al.	8,523,797 B2	9/2013	Lowery et al.
			8,539,812 B2	9/2013	Stringham et al.
			8,543,416 B2	9/2013	Palmroos et al.
			8,577,692 B2	11/2013	Silkaitis et al.
			8,622,990 B2	1/2014	Estes et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

8,630,722 B2	1/2014	Concurso et al.	9,943,269 B2	4/2018	Muhsin et al.
8,665,214 B2	3/2014	Forutanpour et al.	9,995,611 B2	6/2018	Ruchti et al.
8,666,769 B2	3/2014	Butler et al.	10,002,496 B2	6/2018	Humphrey
8,700,421 B2	4/2014	Feng et al.	10,022,498 B2	7/2018	Ruchti et al.
8,706,233 B2	4/2014	Su et al.	10,046,112 B2	8/2018	Oruklu et al.
D705,260 S	5/2014	Gerssen	D827,665 S	9/2018	Segars
8,721,584 B2	5/2014	Braithwaite et al.	10,089,055 B1	10/2018	Fryman
8,728,020 B2	5/2014	Caleffi et al.	10,099,009 B1	10/2018	Anderson et al.
D706,294 S	6/2014	Jewitt	10,166,328 B2	1/2019	Oruklu et al.
8,758,306 B2	6/2014	Lopez et al.	10,241,626 B2	3/2019	Miyazawa
8,761,906 B2	6/2014	Concurso et al.	10,297,350 B2	5/2019	Duke et al.
D709,091 S	7/2014	Kwon	10,342,917 B2	7/2019	Shubinsky et al.
8,768,719 B2	7/2014	Wehba et al.	10,430,761 B2	10/2019	Hume et al.
8,771,251 B2	7/2014	Ruchti et al.	D865,777 S	11/2019	Kovács
8,792,981 B2	7/2014	Yudovsky et al.	10,463,788 B2	11/2019	Day
D711,916 S	8/2014	Matas	10,549,248 B2	2/2020	Brown et al.
D712,926 S	9/2014	Meegan	10,578,474 B2	3/2020	Ruchti et al.
D713,417 S	9/2014	Daniel	10,596,316 B2	3/2020	Dumas, III et al.
D713,418 S	9/2014	Yang	10,635,784 B2	4/2020	Rubalcaba, Jr. et al.
D713,420 S	9/2014	Dallmeyer	10,656,894 B2	5/2020	Fryman
8,821,432 B2	9/2014	Unverdorben	10,682,102 B2	6/2020	Declerck
8,823,382 B2	9/2014	Rondoni et al.	10,709,885 B2	7/2020	Janders et al.
8,857,269 B2	10/2014	Johnson et al.	D898,055 S	10/2020	Connolly
8,858,185 B2	10/2014	Johnson et al.	10,850,024 B2	12/2020	Day et al.
8,905,965 B2	12/2014	Mandro et al.	10,874,793 B2	12/2020	Oruklu et al.
D721,385 S	1/2015	Barling	11,004,035 B2	5/2021	Hume et al.
8,948,734 B2	2/2015	Vaglio	11,007,119 B2	5/2021	Lopez et al.
8,964,185 B1	2/2015	Luo et al.	D922,432 S	6/2021	Kataoka et al.
9,005,150 B2	4/2015	Ware et al.	D923,050 S	6/2021	Kataoka et al.
9,026,370 B2	5/2015	Rubalcaba et al.	11,029,911 B2	6/2021	Fryman
9,084,855 B2	7/2015	Ware et al.	D926,201 S	7/2021	Bryant et al.
9,114,217 B2	8/2015	Sur et al.	D926,224 S	7/2021	Hummel
9,134,735 B2	9/2015	Lowery et al.	D928,813 S	8/2021	Nurutdinov et al.
9,134,736 B2	9/2015	Lowery et al.	D928,840 S	8/2021	Amit et al.
9,138,526 B2	9/2015	Ware et al.	11,090,431 B2	8/2021	Dumas, III et al.
D742,413 S	11/2015	Torres	D931,884 S	9/2021	Bryant et al.
D742,414 S	11/2015	Brunner	D931,892 S	9/2021	Nurutdinov
D742,415 S	11/2015	Cahill	D934,282 S	10/2021	Clymer
D743,414 S	11/2015	Uno	11,135,360 B1	10/2021	Jacobson et al.
9,190,010 B2	11/2015	Vik et al.	11,219,715 B2	1/2022	Gray et al.
D747,339 S	1/2016	Cohen	11,246,985 B2	2/2022	Gylland et al.
9,240,002 B2	1/2016	Hume et al.	D946,608 S	3/2022	Higuchi
D750,099 S	2/2016	Kadosh	11,278,671 B2	3/2022	Cavendish, Jr. et al.
9,272,089 B2	3/2016	Jacobson et al.	11,298,456 B2	4/2022	Shubinsky et al.
9,316,216 B1	4/2016	Cook et al.	11,324,888 B2	5/2022	Shubinsky et al.
D757,099 S	5/2016	Seo	11,344,668 B2	5/2022	Sileika et al.
9,333,291 B2	5/2016	Jacobson et al.	11,344,673 B2	5/2022	Lindo et al.
D758,379 S	6/2016	Kadosh	11,376,361 B2	7/2022	Ruchti et al.
D759,036 S	6/2016	Evanes	11,378,430 B2	7/2022	Ruchti et al.
D760,238 S	6/2016	Suarez	11,395,875 B2	7/2022	Rubalcaba, Jr. et al.
D760,248 S	6/2016	Smith	11,433,177 B2	9/2022	Oruklu et al.
D760,295 S	6/2016	Smith	11,439,570 B2	9/2022	Lopez et al.
D760,788 S	7/2016	Cho	11,596,737 B2	3/2023	Dumas, III et al.
D761,820 S	7/2016	Lee	11,599,854 B2	3/2023	Hume et al.
D762,238 S	7/2016	Day	11,623,042 B2	4/2023	Day
9,381,296 B2	7/2016	Arrizza et al.	11,868,161 B2	1/2024	Fryman
9,393,362 B2	7/2016	Cozmi et al.	11,883,361 B2	1/2024	Janssen
D764,538 S	8/2016	Lee	D1,017,633 S	3/2024	Chung
9,468,718 B2	10/2016	Hung et al.	D1,018,593 S	3/2024	Chiah
9,498,583 B2	11/2016	Sur et al.	11,933,650 B2	3/2024	Ruchti et al.
D773,519 S	12/2016	Hurley	D1,021,917 S	4/2024	Ceniceroz
D777,205 S	1/2017	Orr	D1,023,027 S	4/2024	Slettines
9,545,475 B2	1/2017	Borges et al.	D1,024,096 S	4/2024	Zhong
9,545,476 B2	1/2017	Qi et al.	11,972,395 B2	4/2024	Hume et al.
D779,537 S	2/2017	Wingate-Whyte	D1,027,974 S	5/2024	Correy
D781,874 S	3/2017	Dunn	12,048,831 B2	7/2024	Oruklu et al.
D782,535 S	3/2017	Menz	12,059,551 B2	8/2024	Dumas, III et al.
9,707,341 B2	7/2017	Dumas, III et al.	12,076,531 B2	9/2024	Shubinsky et al.
9,764,087 B2	9/2017	Peterfreund et al.	12,083,310 B2	9/2024	Shubinsky et al.
9,773,330 B1	9/2017	Douglas	2001/0007636 A1	7/2001	Butterfield
D803,881 S	11/2017	Hurley	2001/0014769 A1	8/2001	Bufe et al.
D806,109 S	12/2017	Day	2001/0015099 A1	8/2001	Blaine
9,852,265 B1	12/2017	Treacy et al.	2001/0016056 A1	8/2001	Westphal et al.
D809,006 S	1/2018	Mehta	2001/0032099 A1	10/2001	Joao
9,883,987 B2	2/2018	Lopez et al.	2001/0037060 A1	11/2001	Thompson et al.
			2001/0041869 A1	11/2001	Causey et al.
			2001/0044731 A1	11/2001	Coffman et al.
			2002/0003892 A1	1/2002	Iwanaga
			2002/0007116 A1	1/2002	Zatezalo et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2002/0013545	A1	1/2002	Soltanpour et al.	2004/0077996	A1	4/2004	Jasperson et al.
2002/0013551	A1	1/2002	Zaitso et al.	2004/0082908	A1	4/2004	Whitehurst
2002/0015018	A1	2/2002	Shimazu et al.	2004/0082918	A1	4/2004	Evans et al.
2002/0018720	A1	2/2002	Carlisle et al.	2004/0104271	A1	6/2004	Martucci et al.
2002/0029776	A1	3/2002	Blomquist	2004/0119753	A1	6/2004	Zencke
2002/0031838	A1	3/2002	Meinhart et al.	2004/0120825	A1	6/2004	Bouton et al.
2002/0032583	A1	3/2002	Joao	2004/0128162	A1	7/2004	Schlotterbeck et al.
2002/0040208	A1	4/2002	Flaherty et al.	2004/0128163	A1	7/2004	Goodman et al.
2002/0044059	A1	4/2002	Reeder et al.	2004/0130573	A1	7/2004	Konuma
2002/0045806	A1	4/2002	Baker, Jr. et al.	2004/0133166	A1	7/2004	Moberg et al.
2002/0082728	A1	6/2002	Mueller et al.	2004/0145114	A1	7/2004	Ippolito et al.
2002/0083771	A1	7/2002	Khuri-Yakub et al.	2004/0147034	A1	7/2004	Gore et al.
2002/0085952	A1	7/2002	Ellingboe et al.	2004/0149823	A1	8/2004	Aptekar
2002/0087115	A1	7/2002	Hartlaub	2004/0152970	A1	8/2004	Hunter et al.
2002/0093641	A1	7/2002	Ortyn et al.	2004/0158193	A1	8/2004	Bui et al.
2002/0095486	A1	7/2002	Bahl	2004/0167464	A1	8/2004	Ireland et al.
2002/0099282	A1	7/2002	Knobbe et al.	2004/0167465	A1	8/2004	Kohler
2002/0099334	A1	7/2002	Hanson et al.	2004/0167804	A1	8/2004	Simpson
2002/0143580	A1	10/2002	Bristol et al.	2004/0172222	A1	9/2004	Simpson et al.
2002/0147389	A1	10/2002	Cavallaro et al.	2004/0172283	A1	9/2004	Vanderveen
2002/0152239	A1	10/2002	Bautista-Lloyd et al.	2004/0172289	A1	9/2004	Kozic et al.
2002/0158919	A1	10/2002	Nacey	2004/0172301	A1	9/2004	Mihai et al.
2002/0168278	A1	11/2002	Jeon et al.	2004/0172302	A1	9/2004	Martucci et al.
2002/0173703	A1	11/2002	Lebel et al.	2004/0176984	A1	9/2004	White et al.
2002/0183693	A1	12/2002	Peterson et al.	2004/0181314	A1	9/2004	Zaleski
2003/0009244	A1	1/2003	Engleson	2004/0193025	A1	9/2004	Steil et al.
2003/0013959	A1	1/2003	Grunwald et al.	2004/0193325	A1	9/2004	Bonderud
2003/0018289	A1	1/2003	Ng et al.	2004/0193328	A1	9/2004	Butterfield et al.
2003/0018308	A1	1/2003	Tsai	2004/0204638	A1	10/2004	Diab et al.
2003/0025602	A1	2/2003	Medema et al.	2004/0204673	A1	10/2004	Flaherty et al.
2003/0028082	A1	2/2003	Thompson	2004/0220517	A1	11/2004	Starkweather et al.
2003/0030001	A1	2/2003	Cooper et al.	2004/0225252	A1	11/2004	Gillespie et al.
2003/0045840	A1	3/2003	Burko	2004/0225409	A1	11/2004	Duncan et al.
2003/0050621	A1	3/2003	Lebel et al.	2004/0232219	A1	11/2004	Fowler
2003/0060688	A1	3/2003	Ciarniello et al.	2004/0253123	A1	12/2004	Xie et al.
2003/0060765	A1	3/2003	Campbell et al.	2004/0254434	A1	12/2004	Goodnow et al.
2003/0065537	A1	4/2003	Evans	2004/0254513	A1	12/2004	Shang et al.
2003/0065589	A1	4/2003	Giacchetti	2005/0021006	A1	1/2005	Tonnies
2003/0073954	A1	4/2003	Moberg et al.	2005/0021297	A1	1/2005	Hartlaub
2003/0079746	A1	5/2003	Hickle	2005/0022274	A1	1/2005	Campbell et al.
2003/0083583	A1	5/2003	Kovtun et al.	2005/0038680	A1	2/2005	McMahon
2003/0091442	A1	5/2003	Bush et al.	2005/0055242	A1	3/2005	Bello et al.
2003/0104982	A1	6/2003	Wittmann et al.	2005/0055244	A1	3/2005	Mullan et al.
2003/0106553	A1	6/2003	Vanderveen	2005/0065465	A1	3/2005	Lebel et al.
2003/0125662	A1	7/2003	Bui	2005/0075544	A1	4/2005	Shapiro et al.
2003/0130616	A1	7/2003	Steil	2005/0096593	A1	5/2005	Pope et al.
2003/0135087	A1	7/2003	Hickle et al.	2005/0099624	A1	5/2005	Staehr
2003/0136193	A1	7/2003	Fujimoto	2005/0107923	A1	5/2005	Vanderveen
2003/0139701	A1	7/2003	White et al.	2005/0108057	A1	5/2005	Cohen et al.
2003/0140928	A1	7/2003	Bui et al.	2005/0119597	A1	6/2005	O'Mahony et al.
2003/0140929	A1	7/2003	Wilkes et al.	2005/0119914	A1	6/2005	Batch
2003/0141981	A1	7/2003	Bui et al.	2005/0131739	A1	6/2005	Rabinowitz et al.
2003/0143746	A1	7/2003	Sage, Jr.	2005/0137522	A1	6/2005	Aoki
2003/0144878	A1	7/2003	Wilkes et al.	2005/0143864	A1	6/2005	Blomquist
2003/0158508	A1	8/2003	DiGianfilippo	2005/0145010	A1	7/2005	Vanderveen et al.
2003/0159741	A1	8/2003	Sparks	2005/0171503	A1	8/2005	Van Den Bergh et al.
2003/0160683	A1	8/2003	Blomquist	2005/0171815	A1	8/2005	Vanderveen
2003/0163789	A1	8/2003	Blomquist	2005/0177045	A1	8/2005	Degertekin et al.
2003/0173408	A1	9/2003	Mosher, Jr. et al.	2005/0177096	A1	8/2005	Bollish et al.
2003/0186833	A1	10/2003	Huff et al.	2005/0182306	A1	8/2005	Sloan
2003/0187338	A1	10/2003	Say et al.	2005/0182355	A1	8/2005	Bui
2003/0200116	A1	10/2003	Forrester	2005/0182366	A1	8/2005	Vogt et al.
2003/0204274	A1	10/2003	Ullestad et al.	2005/0187515	A1	8/2005	Varrichio et al.
2003/0204416	A1	10/2003	Acharya	2005/0192529	A1	9/2005	Butterfield et al.
2003/0212364	A1	11/2003	Mann et al.	2005/0192557	A1	9/2005	Brauker et al.
2003/0212379	A1	11/2003	Bylund et al.	2005/0197554	A1	9/2005	Polcha
2003/0216682	A1	11/2003	Junker	2005/0197621	A1	9/2005	Poulsen et al.
2003/0217962	A1	11/2003	Childers et al.	2005/0197649	A1	9/2005	Shelton et al.
2003/0233071	A1	12/2003	Gillespie, Jr. et al.	2005/0209563	A1	9/2005	Hopping et al.
2004/0030277	A1	2/2004	O'Mahony et al.	2005/0209793	A1	9/2005	Yamada
2004/0047736	A1	3/2004	Nose et al.	2005/0224083	A1	10/2005	Crass
2004/0057226	A1	3/2004	Berthou et al.	2005/0235732	A1	10/2005	Rush
2004/0064342	A1	4/2004	Browne et al.	2005/0238506	A1	10/2005	Mescher et al.
2004/0073125	A1	4/2004	Lovett et al.	2005/0240305	A1	10/2005	Bogash et al.
2004/0073161	A1	4/2004	Tachibana	2005/0273059	A1	12/2005	Mernoe et al.
				2005/0277890	A1	12/2005	Stewart et al.
				2005/0279419	A1	12/2005	Tribble et al.
				2006/0002799	A1	1/2006	Schann et al.
				2006/0009727	A1	1/2006	O'Mahony et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2006/0009734	A1	1/2006	Martin	2007/0180916	A1	8/2007	Tian et al.
2006/0042633	A1	3/2006	Bishop et al.	2007/0191770	A1	8/2007	Moberg et al.
2006/0047270	A1	3/2006	Shelton	2007/0191817	A1	8/2007	Martin
2006/0053036	A1	3/2006	Coffman et al.	2007/0197963	A1	8/2007	Griffiths et al.
2006/0064020	A1	3/2006	Burnes et al.	2007/0214003	A1	9/2007	Holland et al.
2006/0064053	A1	3/2006	Bollish et al.	2007/0215545	A1	9/2007	Bissler et al.
2006/0079768	A1	4/2006	Small et al.	2007/0233035	A1	10/2007	Wehba et al.
2006/0079831	A1	4/2006	Gilbert	2007/0233049	A1	10/2007	Wehba et al.
2006/0100746	A1	5/2006	Leibner-Druska	2007/0240497	A1	10/2007	Robinson et al.
2006/0100907	A1	5/2006	Holland et al.	2007/0250339	A1	10/2007	Mallett et al.
2006/0106649	A1	5/2006	Eggers et al.	2007/0255250	A1	11/2007	Moberg et al.
2006/0116639	A1	6/2006	Russell	2007/0257788	A1	11/2007	Carlson
2006/0117856	A1	6/2006	Orr et al.	2007/0267945	A1	11/2007	Sudol
2006/0117867	A1	6/2006	Froehlich et al.	2007/0270747	A1	11/2007	Remde
2006/0122867	A1	6/2006	Eggers et al.	2007/0274843	A1	11/2007	Vanderveen et al.
2006/0135939	A1	6/2006	Brown	2007/0289384	A1	12/2007	Sakai et al.
2006/0135940	A1	6/2006	Joshi	2008/0009684	A1	1/2008	Corsetti et al.
2006/0136095	A1	6/2006	Rob et al.	2008/0028868	A1	2/2008	Konzelmann et al.
2006/0136271	A1	6/2006	Eggers et al.	2008/0033361	A1	2/2008	Evans et al.
2006/0140798	A1	6/2006	Kutsuzawa	2008/0039777	A1	2/2008	Katz et al.
2006/0143051	A1	6/2006	Eggers et al.	2008/0048211	A1	2/2008	Khuri-Yakub et al.
2006/0173260	A1	8/2006	Gaoni et al.	2008/0058773	A1	3/2008	John
2006/0173406	A1	8/2006	Hayes et al.	2008/0060448	A1	3/2008	Wiest et al.
2006/0180916	A1	8/2006	Wyland et al.	2008/0065420	A1	3/2008	Tirinato et al.
2006/0181695	A1	8/2006	Sage, Jr.	2008/0071210	A1	3/2008	Moubayed et al.
2006/0187069	A1	8/2006	Duan	2008/0071496	A1	3/2008	Glascocock
2006/0190302	A1	8/2006	Eggers et al.	2008/0071580	A1	3/2008	Marcus et al.
2006/0195022	A1	8/2006	Trepagnier et al.	2008/0077116	A1	3/2008	Dailey et al.
2006/0200007	A1	9/2006	Brockway et al.	2008/0086087	A1	4/2008	Spohn et al.
2006/0200369	A1	9/2006	Batch et al.	2008/0091466	A1	4/2008	Butler et al.
2006/0211404	A1	9/2006	Cromp et al.	2008/0097288	A1	4/2008	Levin et al.
2006/0224140	A1	10/2006	Junker	2008/0097289	A1	4/2008	Steil et al.
2006/0224141	A1	10/2006	Rush et al.	2008/0097317	A1	4/2008	Alholm et al.
2006/0224181	A1	10/2006	McEwen et al.	2008/0098798	A1	5/2008	Riley et al.
2006/0226088	A1	10/2006	Robinson et al.	2008/0119822	A1	5/2008	Knauper
2006/0226089	A1	10/2006	Robinson et al.	2008/0125701	A1	5/2008	Moberg et al.
2006/0226090	A1	10/2006	Robinson et al.	2008/0139907	A1	6/2008	Rao et al.
2006/0229918	A1	10/2006	Fotsch et al.	2008/0145249	A1	6/2008	Smisson
2006/0235353	A1	10/2006	Gelfand et al.	2008/0169044	A1	7/2008	Osborne et al.
2006/0255149	A1	11/2006	Retter et al.	2008/0172030	A1	7/2008	Blomquist et al.
2006/0258985	A1	11/2006	Russell	2008/0177254	A1	7/2008	Shelton et al.
2006/0260416	A1	11/2006	Sage et al.	2008/0184784	A1	8/2008	Dam
2006/0264895	A1	11/2006	Flanders	2008/0188789	A1	8/2008	Galavotti et al.
2006/0266128	A1	11/2006	Clark et al.	2008/0188796	A1	8/2008	Steil et al.
2006/0270971	A1	11/2006	Gelfand et al.	2008/0208484	A1	8/2008	Butterfield et al.
2006/0271286	A1	11/2006	Rosenberg	2008/0214919	A1	9/2008	Harmon et al.
2006/0272421	A1	12/2006	Frinak et al.	2008/0221521	A1	9/2008	Getz et al.
2006/0275142	A1	12/2006	Bouton et al.	2008/0221522	A1	9/2008	Moberg et al.
2007/0015972	A1	1/2007	Wang et al.	2008/0262469	A1	10/2008	Bristol et al.
2007/0036511	A1	2/2007	Lundquist et al.	2008/0269663	A1	10/2008	Arnold et al.
2007/0060796	A1	3/2007	Kim	2008/0269714	A1	10/2008	Mastrototaro et al.
2007/0060869	A1	3/2007	Tolle et al.	2008/0269723	A1	10/2008	Mastrototaro et al.
2007/0060871	A1	3/2007	Istoc	2008/0275384	A1	11/2008	Mastrototaro et al.
2007/0060872	A1	3/2007	Hall et al.	2008/0300572	A1	12/2008	Rankers et al.
2007/0060874	A1	3/2007	Nesbitt et al.	2009/0001908	A1	1/2009	Shubinsky et al.
2007/0062250	A1	3/2007	Krulevitch et al.	2009/0005703	A1	1/2009	Fasciano
2007/0065363	A1	3/2007	Dalal et al.	2009/0006061	A1	1/2009	Thukral et al.
2007/0078314	A1	4/2007	Grounsell	2009/0006129	A1	1/2009	Thukral
2007/0083152	A1	4/2007	Williams, Jr. et al.	2009/0006133	A1	1/2009	Weinert
2007/0084286	A1	4/2007	Ajay et al.	2009/0015824	A1	1/2009	Shubinsky et al.
2007/0084288	A1	4/2007	Thomas et al.	2009/0043171	A1	2/2009	Rule
2007/0088271	A1	4/2007	Richards	2009/0054743	A1	2/2009	Stewart
2007/0088333	A1	4/2007	Levin et al.	2009/0054754	A1	2/2009	McMahon et al.
2007/0093753	A1	4/2007	Krulevitch et al.	2009/0062727	A1	3/2009	Woo
2007/0094045	A1	4/2007	Cobbs et al.	2009/0069743	A1	3/2009	Krishnamoorthy et al.
2007/0094046	A1	4/2007	Cobbs et al.	2009/0077248	A1	3/2009	Castellucci et al.
2007/0100222	A1	5/2007	Mastrototaro et al.	2009/0082676	A1	3/2009	Bennison
2007/0100665	A1	5/2007	Brown	2009/0088731	A1	4/2009	Campbell et al.
2007/0112298	A1	5/2007	Mueller et al.	2009/0097029	A1	4/2009	Tokhtuev et al.
2007/0118405	A1	5/2007	Campbell et al.	2009/0099866	A1	4/2009	Newman
2007/0129618	A1	6/2007	Goldberger et al.	2009/0105636	A1	4/2009	Hayter et al.
2007/0142822	A1	6/2007	Remde	2009/0112155	A1	4/2009	Zhao
2007/0156452	A1	7/2007	Batch	2009/0114037	A1	5/2009	Smith
2007/0156456	A1	7/2007	McGillin et al.	2009/0119330	A1	5/2009	Sampath et al.
2007/0179436	A1	8/2007	Braig et al.	2009/0124963	A1	5/2009	Hogard et al.
				2009/0124964	A1	5/2009	Leach et al.
				2009/0126825	A1	5/2009	Eliuk et al.
				2009/0131861	A1	5/2009	Braig et al.
				2009/0135196	A1	5/2009	Holland et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2009/0143726	A1	6/2009	Bouton et al.	2011/0071844	A1	3/2011	Cannon et al.
2009/0144025	A1	6/2009	Bouton et al.	2011/0072379	A1	3/2011	Gannon
2009/0144026	A1	6/2009	Bouton et al.	2011/0077480	A1	3/2011	Bloom et al.
2009/0149743	A1	6/2009	Barron et al.	2011/0078608	A1	3/2011	Gannon et al.
2009/0156922	A1	6/2009	Goldberger et al.	2011/0099313	A1	4/2011	Bolanowski
2009/0156975	A1	6/2009	Robinson et al.	2011/0105983	A1	5/2011	Kelly et al.
2009/0177146	A1	7/2009	Nesbitt et al.	2011/0106561	A1	5/2011	Eaton, Jr. et al.
2009/0177188	A1	7/2009	Steinkogler	2011/0107251	A1	5/2011	Guaitoli et al.
2009/0177248	A1	7/2009	Roberts	2011/0137241	A1	6/2011	DelCastillo et al.
2009/0177769	A1	7/2009	Roberts	2011/0144595	A1	6/2011	Cheng
2009/0178485	A1	7/2009	Thomas et al.	2011/0152770	A1	6/2011	Diperna et al.
2009/0183147	A1	7/2009	Davis et al.	2011/0160649	A1	6/2011	Pan
2009/0192367	A1	7/2009	Braig et al.	2011/0162647	A1	7/2011	Huby et al.
2009/0198347	A1	8/2009	Kirzinger	2011/0172918	A1	7/2011	Tome
2009/0205426	A1	8/2009	Balschat et al.	2011/0175728	A1	7/2011	Baker, Jr.
2009/0209938	A1	8/2009	Aalto-Setälä	2011/0190598	A1	8/2011	Shusterman
2009/0209945	A1	8/2009	Lobl et al.	2011/0190694	A1	8/2011	Lanier et al.
2009/0212966	A1	8/2009	Panduro	2011/0218514	A1	9/2011	Rebours
2009/0221890	A1	9/2009	Saffer et al.	2011/0238032	A1	9/2011	McTaggart et al.
2009/0223294	A1	9/2009	Thomas et al.	2011/0264006	A1	10/2011	Ali et al.
2009/0227939	A1	9/2009	Memoe et al.	2011/0264043	A1	10/2011	Kotnick et al.
2009/0264720	A1	10/2009	Torjman et al.	2011/0282321	A1	11/2011	Steil et al.
2009/0270810	A1	10/2009	DeBelser	2011/0313390	A1	12/2011	Roy et al.
2009/0270833	A1	10/2009	DeBelser	2011/0319728	A1	12/2011	Petisce et al.
2010/0022988	A1	1/2010	Wochner	2011/0320049	A1	12/2011	Chossat et al.
2010/0036310	A1	2/2010	Hillman	2012/0016215	A1	1/2012	Condurso et al.
2010/0056992	A1	3/2010	Hayter	2012/0023431	A1	1/2012	Roth
2010/0057042	A1	3/2010	Hayter	2012/0025995	A1	2/2012	Moberg et al.
2010/0069892	A1	3/2010	Steinbach et al.	2012/0059234	A1	3/2012	Barrett et al.
2010/0077866	A1	4/2010	Graboi et al.	2012/0068001	A1	3/2012	Pushkarsky et al.
2010/0079760	A1	4/2010	Bernacki	2012/0083760	A1	4/2012	Ledford et al.
2010/0094251	A1	4/2010	Estes et al.	2012/0085277	A1	4/2012	Abdel-Rahman
2010/0106082	A1	4/2010	Zhou	2012/0089411	A1	4/2012	Srnka et al.
2010/0114027	A1	5/2010	Jacobson et al.	2012/0095433	A1	4/2012	Hungerford et al.
2010/0121170	A1	5/2010	Rule	2012/0123322	A1	5/2012	Scarpaci et al.
2010/0121415	A1	5/2010	Skelton et al.	2012/0143116	A1	6/2012	Ware et al.
2010/0130933	A1	5/2010	Holland et al.	2012/0180790	A1	7/2012	Montgomery
2010/0131434	A1	5/2010	Magent et al.	2012/0185267	A1	7/2012	Kamen et al.
2010/0141460	A1	6/2010	Tokhtuev et al.	2012/0191059	A1	7/2012	Cummings et al.
2010/0147081	A1	6/2010	Thomas et al.	2012/0194341	A1	8/2012	Peichel et al.
2010/0152554	A1	6/2010	Steine et al.	2012/0203177	A1	8/2012	Lanier
2010/0160854	A1	6/2010	Gauthier	2012/0222774	A1	9/2012	Husnu et al.
2010/0168535	A1	7/2010	Robinson et al.	2012/0226350	A1	9/2012	Rudser et al.
2010/0177375	A1	7/2010	Seyfried	2012/0245525	A1	9/2012	Pope et al.
2010/0185142	A1	7/2010	Kamen et al.	2012/0259278	A1	10/2012	Hayes et al.
2010/0185182	A1	7/2010	Alme et al.	2012/0310204	A1	12/2012	Krogh et al.
2010/0198034	A1	8/2010	Thomas et al.	2012/0323212	A1	12/2012	Murphy
2010/0198182	A1	8/2010	Lanigan et al.	2013/0006666	A1	1/2013	Schneider
2010/0198183	A1	8/2010	Lanigan et al.	2013/0009551	A1	1/2013	Knapp
2010/0211002	A1	8/2010	Davis	2013/0012880	A1	1/2013	Blomquist
2010/0212407	A1	8/2010	Stringham et al.	2013/0012917	A1	1/2013	Miller et al.
2010/0212675	A1	8/2010	Walling et al.	2013/0032634	A1	2/2013	McKirdy
2010/0217154	A1	8/2010	Deshmukh et al.	2013/0041342	A1	2/2013	Bernini et al.
2010/0217621	A1	8/2010	Schoenberg	2013/0044111	A1	2/2013	VanGilder et al.
2010/0271218	A1	10/2010	Hoag et al.	2013/0110538	A1	5/2013	Butterfield et al.
2010/0271479	A1	10/2010	Heydlauf	2013/0116649	A1	5/2013	Breton et al.
2010/0273738	A1	10/2010	Valcke et al.	2013/0150766	A1	6/2013	Olde et al.
2010/0280430	A1	11/2010	Caleffi et al.	2013/0150821	A1	6/2013	Bollish et al.
2010/0292634	A1	11/2010	Kircher	2013/0173291	A1	7/2013	Kelly
2010/0295686	A1	11/2010	Sloan et al.	2013/0177455	A1	7/2013	Kamen et al.
2010/0298765	A1	11/2010	Budiman et al.	2013/0184676	A1	7/2013	Kamen et al.
2010/0312039	A1	12/2010	Quirico et al.	2013/0197930	A1	8/2013	Garibaldi et al.
2010/0317093	A1	12/2010	Turewicz et al.	2013/0201482	A1	8/2013	Munro
2010/0317952	A1	12/2010	Budiman et al.	2013/0218080	A1	8/2013	Peterfreund et al.
2010/0318025	A1	12/2010	John	2013/0253430	A1	9/2013	Kouyoumjian et al.
2011/0000560	A1	1/2011	Miller et al.	2013/0253946	A1	9/2013	Broselow
2011/0001605	A1	1/2011	Kiani et al.	2013/0274576	A1	10/2013	Amirouche et al.
2011/0004186	A1	1/2011	Butterfield	2013/0281965	A1	10/2013	Kamen et al.
2011/0009797	A1	1/2011	Kelly et al.	2013/0291116	A1	10/2013	Homer
2011/0028885	A1	2/2011	Eggers et al.	2013/0296823	A1	11/2013	Melker et al.
2011/0046558	A1	2/2011	Gravesen et al.	2013/0296984	A1	11/2013	Burnett et al.
2011/0054311	A1	3/2011	Williams et al.	2013/0318158	A1	11/2013	Teng et al.
2011/0062703	A1	3/2011	Lopez et al.	2013/0322201	A1	12/2013	Hitchcock et al.
2011/0064612	A1	3/2011	Franzoni et al.	2013/0345658	A1	12/2013	Browne et al.
2011/0071464	A1	3/2011	Palerm	2013/0345666	A1	12/2013	Panduro et al.
				2014/0067425	A1	3/2014	Dudar et al.
				2014/0132524	A1	5/2014	Lee
				2014/0145915	A1	5/2014	Ribble et al.
				2014/0180711	A1	6/2014	Kamen et al.

(56)

References Cited**U.S. PATENT DOCUMENTS**

2014/0224829 A1 8/2014 Capone et al.
 2014/0267563 A1 9/2014 Baca et al.
 2014/0303591 A1 10/2014 Peterfreund et al.
 2014/0303754 A1 10/2014 Nixon et al.
 2015/0025453 A1 1/2015 Ledford et al.
 2015/0033073 A1 1/2015 Yang et al.
 2015/0051458 A1 2/2015 Chen
 2015/0057108 A1 2/2015 Regimbal
 2015/0065988 A1 3/2015 Holderle et al.
 2015/0089439 A1 3/2015 Wada
 2015/0114515 A1 4/2015 Phallen
 2015/0141921 A1 5/2015 Stewart et al.
 2015/0168958 A1 6/2015 Downie et al.
 2015/0265765 A1 9/2015 Yavorsky et al.
 2015/0278474 A1 10/2015 Stueckemann
 2015/0289823 A1 10/2015 Rack-Gomer et al.
 2015/0338340 A1 11/2015 Jiang et al.
 2015/0343141 A1 12/2015 Lindo et al.
 2015/0363086 A1 12/2015 Lim
 2015/0371004 A1 12/2015 Jones
 2016/0019352 A1 1/2016 Seo
 2016/0042264 A1 2/2016 Borges et al.
 2016/0051750 A1 2/2016 Tsoukalis
 2016/0110088 A1 4/2016 Vik et al.
 2016/0144101 A1 5/2016 Pananen
 2016/0151560 A1 6/2016 Toro et al.
 2016/0151562 A1 6/2016 Magers et al.
 2016/0151601 A1 6/2016 Cardelius et al.
 2016/0158437 A1 6/2016 Biasi et al.
 2016/0175517 A1 6/2016 Sileika et al.
 2016/0193604 A1 7/2016 McFarland et al.
 2016/0253460 A1 9/2016 Kanada
 2016/0339167 A1 11/2016 Ledford et al.
 2017/0010677 A1 1/2017 Roh
 2017/0043089 A1 2/2017 Handler
 2017/0056604 A1 3/2017 Cowan
 2017/0068498 A1 3/2017 Hashem
 2017/0132867 A1 5/2017 Berg et al.
 2017/0354941 A1 12/2017 Brown et al.
 2018/0018440 A1 1/2018 Sugawara
 2018/0021514 A1 1/2018 Rosinko et al.
 2018/0206798 A1 7/2018 Murai
 2018/0296751 A1 10/2018 Lefort et al.
 2018/0300994 A1 10/2018 Nelson et al.
 2018/0326146 A1 11/2018 Gupta et al.
 2019/0072405 A1 3/2019 Luchner
 2019/0091401 A1 3/2019 Ruchti et al.
 2019/0117890 A1 4/2019 Oruklu et al.
 2019/0160254 A1 5/2019 Anand
 2019/0201607 A1 7/2019 Öberg
 2019/0262535 A1 8/2019 Shubinsky et al.
 2019/0282757 A1 9/2019 Gylland et al.
 2019/0351131 A1 11/2019 Butterfield et al.
 2020/0054825 A1 2/2020 Kamen et al.
 2020/0069864 A1 3/2020 Shubinsky et al.
 2020/0090122 A1 3/2020 Hume
 2020/0113784 A1 4/2020 Lopez et al.
 2020/0238007 A1 7/2020 Day
 2020/0271499 A1 8/2020 Ruchti et al.
 2020/0282137 A1 9/2020 Dumas, III et al.
 2020/0319837 A1 10/2020 Fryman
 2020/0324044 A1 10/2020 Gylland et al.
 2020/0330689 A1 10/2020 Nemoto et al.
 2020/0357500 A1 11/2020 Rubalcaba, Jr. et al.
 2020/0384191 A1 12/2020 Rosinko et al.
 2021/0158481 A1 5/2021 Wang
 2021/0158946 A1 5/2021 Starobinets et al.
 2021/0162115 A1 6/2021 Surine
 2021/0170101 A1 6/2021 Cavendish, Jr. et al.
 2021/0260283 A1 8/2021 Oruklu et al.
 2021/0295263 A1 9/2021 Hume et al.
 2021/0304864 A1 9/2021 Kamen et al.
 2021/0397396 A1 12/2021 Fryman
 2022/0031943 A1 2/2022 Dumas, III
 2022/0184302 A1 6/2022 Cavendish, Jr. et al.

2022/0184304 A1 6/2022 Rinehart
 2022/0296806 A1 9/2022 Shubinsky et al.
 2022/0305200 A1 9/2022 Gylland et al.
 2022/0331518 A1 10/2022 Shubinsky et al.
 2022/0362463 A1 11/2022 Lindo et al.
 2023/0010290 A1 1/2023 Oruklu et al.
 2023/0010638 A1 1/2023 Rubalcaba, Jr. et al.
 2023/0017117 A1 1/2023 Sileika et al.
 2023/0058662 A1 2/2023 Ruchti et al.
 2023/0112979 A1 4/2023 Xavier
 2023/0115595 A1 4/2023 Cousineau et al.
 2023/0181419 A1 6/2023 Fister
 2023/0245741 A1 8/2023 Shigyo
 2023/0270938 A1 8/2023 Dumas, III et al.
 2023/0285669 A1 9/2023 Day
 2023/0310735 A1 10/2023 Cousineau
 2024/0201922 A1 6/2024 Fryman
 2024/0263981 A1 8/2024 Ruchti et al.
 2024/0325246 A1 10/2024 Janssen
 2024/0366858 A1 11/2024 Cousineau et al.

FOREIGN PATENT DOCUMENTS

CA 2 113 473 3/1993
 CA 2 551 817 7/2005
 CA 2 554 407 8/2005
 CN 105682703 6/2016
 CN 107106042 8/2017
 CN 110573195 12/2019
 CN 105848694 1/2020
 CN 111954966 11/2020
 CN 306893275 10/2021
 CN 307412164 6/2022
 CN 307979072 4/2023
 CN 112105405 8/2023
 CN 308499458 3/2024
 CO 0020220004676-0001 11/2022
 CO 0020220008155-0001 11/2022
 DE 31 12 762 1/1983
 DE 34 35 647 7/1985
 DE 35 30 747 3/1987
 DE 37 20 664 1/1989
 DE 38 27 444 2/1990
 DE 197 34 002 9/1998
 DE 199 01 078 2/2000
 DE 198 40 965 3/2000
 DE 198 44 252 3/2000
 DE 199 32 147 1/2001
 DE 102 49 238 5/2004
 DE 103 52 456 7/2005
 EM 008932172-0003 4/2022
 EM 008932172-0004 4/2022
 EP 0 282 323 9/1988
 EP 0 291 727 11/1988
 EP 0 319 272 6/1989
 EP 0 319 275 6/1989
 EP 0 335 385 10/1989
 EP 0 337 092 10/1989
 EP 0 341 582 11/1989
 EP 0 370 162 5/1990
 EP 0 387 724 9/1990
 EP 0 429 866 6/1991
 EP 0 441 323 8/1991
 EP 0 453 211 10/1991
 EP 0 462 405 12/1991
 EP 0 501 234 9/1992
 EP 0 516 130 12/1992
 EP 0 519 765 12/1992
 EP 0 643 301 3/1995
 EP 0 683 465 11/1995
 EP 0 431 310 1/1996
 EP 0 589 439 8/1998
 EP 0 880 936 12/1998
 EP 0 954 090 11/1999
 EP 0 960 627 12/1999
 EP 1 174 817 1/2002
 EP 1 177 802 2/2002
 EP 1 197 178 4/2002
 EP 1 500 025 4/2003

(56)	References Cited		WO	WO 02/087664	11/2002
	FOREIGN PATENT DOCUMENTS		WO	WO 03/006091	1/2003
			WO	WO 03/053498	7/2003
			WO	WO 03/093780	11/2003
EP	1 813 188	8/2007	WO	WO 2004/035115	4/2004
EP	1 490 131	12/2007	WO	WO 2004/060455	7/2004
EP	2 062 527	5/2009	WO	WO 2004/061745	7/2004
EP	2 228 004	9/2010	WO	WO 2004/070556	8/2004
EP	2 243 506	10/2010	WO	WO 2004/070994	8/2004
EP	2 381 260	10/2011	WO	WO 2004/112579	12/2004
ES	254513	10/1981	WO	WO 2005/018716	3/2005
FR	2 717 919	9/1995	WO	WO 2005/030489	4/2005
GB	2 121 971	1/1984	WO	WO 2005/036447	4/2005
GB	2 303 706	2/1997	WO	WO 2005/050526	6/2005
GB	2 312 022	10/1997	WO	WO 2005/057175	6/2005
GB	2 312 046	10/1997	WO	WO 2005/065146	7/2005
GB	6 201 192	4/2022	WO	WO 2005/065749	7/2005
GB	6 201 193	4/2022	WO	WO 2005/082450	9/2005
JP	01-301118	12/1989	WO	WO 2005/118015	12/2005
JP	01-308568	12/1989	WO	WO 2006/016122	2/2006
JP	04-231966	8/1992	WO	WO 2006/022906	3/2006
JP	07-502678	3/1995	WO	WO 2006/026270	3/2006
JP	07-289638	11/1995	WO	WO 2007/000426	1/2007
JP	11-128344	5/1999	WO	WO 2007/033025	3/2007
JP	2000-111374	4/2000	WO	WO 2007/035567	3/2007
JP	2000-510575	8/2000	WO	WO 2007/087443	8/2007
JP	2000-515716	11/2000	WO	WO 2008/004560	1/2008
JP	2001-356034	12/2001	WO	WO 2008/019016	2/2008
JP	2002-506514	2/2002	WO	WO 2008/053193	5/2008
JP	2002-131105	5/2002	WO	WO 2008/059492	5/2008
JP	2003-038642	2/2003	WO	WO 2008/063429	5/2008
JP	2003-050144	2/2003	WO	WO 2008/067245	6/2008
JP	2005-021463	1/2005	WO	WO 2008/088490	7/2008
JP	2005-524081	3/2005	WO	WO 2008/134146	11/2008
JP	2006-517423	7/2006	WO	WO 2009/016504	2/2009
JP	2007-071695	3/2007	WO	WO 2009/023406	2/2009
JP	2007-518471	7/2007	WO	WO 2009/023407	2/2009
JP	2007-520270	7/2007	WO	WO 2009/023634	2/2009
JP	2007-275106	10/2007	WO	WO 2009/039203	3/2009
JP	2008-249400	10/2008	WO	WO 2009/039214	3/2009
JP	4322661	6/2009	WO	WO 2009/049252	4/2009
JP	2009-148592	7/2009	WO	WO 2009/127683	10/2009
JP	2010-063767	3/2010	WO	WO 2009/141504	11/2009
JP	5716879	3/2015	WO	WO 2010/017279	2/2010
TW	201841165	11/2018	WO	WO 2010/075371	7/2010
WO	WO 84/000690	3/1984	WO	WO 2010/099313	9/2010
WO	WO 84/000894	3/1984	WO	WO 2010/114929	10/2010
WO	WO 90/007942	7/1990	WO	WO 2010/119409	10/2010
WO	WO 91/000113	1/1991	WO	WO 2010/124127	10/2010
WO	WO 91/016087	10/1991	WO	WO 2010/135646	11/2010
WO	WO 91/016416	10/1991	WO	WO 2010/135654	11/2010
WO	WO 93/004284	3/1993	WO	WO 2010/135670	11/2010
WO	WO 95/016200	6/1995	WO	WO 2010/135686	11/2010
WO	WO 95/031233	11/1995	WO	WO 2010/148205	12/2010
WO	WO 96/008755	3/1996	WO	WO 2011/017778	2/2011
WO	WO 96/025186	8/1996	WO	WO 2011/080188	7/2011
WO	WO 96/028209	9/1996	WO	WO 2011/109774	9/2011
WO	WO 96/041156	12/1996	WO	WO 2012/042763	4/2012
WO	WO 97/010013	3/1997	WO	WO 2012/082599	6/2012
WO	WO 97/030333	8/1997	WO	WO 2012/108910	8/2012
WO	WO 98/004304	2/1998	WO	WO 2012/167090	12/2012
WO	WO 98/012670	3/1998	WO	WO 2013/036854	3/2013
WO	WO 98/014234	4/1998	WO	WO 2013/096769	6/2013
WO	WO 98/019263	5/1998	WO	WO 2015/134478	9/2015
WO	WO 98/044320	10/1998	WO	WO 2017/051271	3/2017
WO	WO 98/056441	12/1998	WO	WO 2017/087157	5/2017
WO	WO 99/010029	3/1999	WO	WO 2017/144366	8/2017
WO	WO 99/015216	4/1999	WO	WO 2017/197024	11/2017
WO	WO 99/051003	10/1999	WO	WO 2019/063462	4/2019
WO	WO 99/052575	10/1999	WO	WO 2019/092680	5/2019
WO	WO 00/013580	3/2000	WO	WO 2020/214717	10/2020
WO	WO 00/013726	3/2000	WO	WO 2022/020184	1/2022
WO	WO 00/041621	7/2000	WO	WO 2022/072159	4/2022
WO	WO 01/014974	3/2001	WO	WO 2022/125471	6/2022
WO	WO 01/033484	5/2001	WO	WO 2023/064662	4/2023
WO	WO 02/005702	1/2002	WO	WO 2023/108030	6/2023
WO	WO 02/009795	2/2002			
WO	WO 02/027276	4/2002			
WO	WO 02/066101	8/2002			

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO WO 2023/192791 10/2023
 WO WO 2023/244922 12/2023

OTHER PUBLICATIONS

ALARIS® Medical Systems, "Signature Edition® GOLD—Single & Dual Channel Infusion System", San Diego, CA, USA, date unknown, but believed to be at least as early as Nov. 29, 2008, pp. 2-88 & 2-91.

Allegro, "3955—Full-Bridge PWM Microstepping Motor Drive", Datasheet, 1997, pp. 16.

Aragon, Daleen RN, Ph.D., CCRN, "Evaluation of Nursing Work Effort and Perceptions About Blood Glucose Testing in Tight Glycemic Control", *American Journal of Critical Care*, Jul. 2006, vol. 15, No. 4, pp. 370-377.

Baxter, "Baxter Receives 510(k) Clearance for Next-Generation SIGMA Spectrum Infusion Pump with Master Drug Library" Press Release, May 8, 2014, pp. 2. http://web.archive.org/web/20160403140025/http://www.baxter.com/news-media/newsroom/press-releases/2014/05_08_14_sigma.page.

Bequette, Ph.D., "A Critical Assessment of Algorithms and Challenges in the Development of a Closed-Loop Artificial Pancreas", *Diabetes Technology & Therapeutics*, Feb. 28, 2005, vol. 7, No. 1, pp. 28-47.

Bequette, B. Wayne, Ph.D., "Analysis of Algorithms for Intensive Care Unit Blood Glucose Control", *Journal of Diabetes Science and Technology*, Nov. 2007, vol. 1, No. 6, pp. 813-824.

Binder et al., "Insulin Infusion with Parenteral Nutrition in Extremely Low Birth Weight Infants with Hyperglycemia", *Journal of Pediatrics*, Feb. 1989, vol. 114, No. 2, pp. 273-280.

Bode et al., "Intravenous Insulin Infusion Therapy: Indications, Methods, and Transition to Subcutaneous Insulin Therapy", *Endocrine Practice*, Mar./Apr. 2004, vol. 10, Supplement 2, pp. 71-80.

Buhrdorf et al., "Capacitive Micromachined Ultrasonic Transducers and their Application", *Proceedings of the IEEE Ultrasonics Symposium*, Feb. 2001, vol. 2, pp. 933-940.

Cannon, MD et al., "Automated Heparin-Delivery System to Control Activated Partial Thromboplastin Time", *Circulation*, Feb. 16, 1999, vol. 99, pp. 751-756.

"CareAware® Infusion Management", Cerner Store, as printed May 12, 2011, pp. 3, <https://store.cerner.com/items/7>.

Chen et al., "Enabling Location-Based Services on Wireless LANs", *The 11th IEEE International Conference on Networks, ICON 2003*, Sep. 28-Oct. 1, 2003, pp. 567-572.

Cheung et al., "Hyperglycemia is Associated with Adverse Outcomes in Patients Receiving Total Parenteral Nutrition", *Diabetes Care*, Oct. 2005, vol. 28, No. 10, pp. 2367-2371.

Coley et al., "Performance of Three Portable Infusion-Pump Devices Set to Deliver 2 mL/hr", *American Journal of Health-System Pharmacy*, Jun. 1, 1997, vol. 54, No. 11, pp. 1277-1280.

"Continually vs Continuously", https://web.archive.org/web/20090813092423/http://www.diffen.com/difference/Continually_vs_Continuously, as accessed Aug. 13, 2009 in 4 pages.

"Criticore® Monitor: Critical Fluid Output and Core Bladder Temperature Monitor", *BARD Urological Catheter Systems*, Advertisement, 2005, pp. 2.

Daimiwal et al., "Wireless Transfusion Supervision and Analysis Using Embedded System", *IEEE, 2010 International Conference ICBT, China*, Apr. 2010, pp. 56-60.

Davidson et al., "A Computer-Directed Intravenous Insulin System Shown to be Safe, Simple, and Effective in 120,618 h of Operation", *Diabetes Care*, Oct. 2005, vol. 28, No. 10, pp. 2418-2423.

"Decision of the Administrative Council of Oct. 16, 2013 Amending Rule 135 and 164 of the Implementing Regulations to the European Patent Convention (CA/D 17/13)", *Official Journal EPO* Nov. 2013, Nov. 2013, pp. 503-506. http://archive.epo.org/epo/pubs/oj013/11_13/11_5033.pdf.

"Decision of the Administrative Council of Oct. 27, 2009 Amending the Implementing Regulations to the European Patent Convention (CA/D 20/09)", *Official Journal EPO* Dec. 2009, Dec. 2009, pp. 582-584. http://archive.epo.org/epo/pubs/oj009/12_09/12_5829.pdf.

Diabetes Close Up, *Close Concerns AACE Inpatient Management Conference Report, Consensus Development Conference on Inpatient Diabetes and Metabolic Control*, Washington, D.C., Dec. 14-16, 2003, pp. 1-32.

"Differential Pressure Transmitter, Series PD-39 X", *SensorsOne Ltd.*, Advertisement, Dec. 2005, pp. 2.

Dunster et al., "Flow Continuity of Infusion Systems at Low Flow Rates", *Anaesthesia and Intensive Care*, Oct. 1995, vol. 23, No. 5, pp. 5.

Fogt et al., *Development and Evaluation of a Glucose Analyzer for a Glucose-Controlled Insulin Infusion System (Biostator®)*, *Clinical Chemistry*, 1978, vol. 24, No. 8, pp. 1366-1372.

"Froth", <http://www.merriam-webster.com/dictionary/froth>, as accessed May 13, 2015 in 1 page.

Goldberg et al., "Clinical Results of an Updated Insulin Infusion Protocol in Critically Ill Patients", *Diabetes Spectrum*, 2005, vol. 18, No. 3, pp. 188-191.

Halpern et al., "Changes in Critical Care Beds and Occupancy in the United States 1985-2000: Differences Attributable to Hospital Size", *Critical Care Medical*, Aug. 2006, vol. 34, No. 8, pp. 2105-2112.

Hospira, "Plum A+™ Infusion System" as archived Dec. 1, 2012, pp. 2. www.hospira.com/products_and_services/infusion_pumps/plum/index.

Hospira, "Plum XL™ Series Infusion System" Technical Service Manual, Feb. 2005, Lake Forest, Illinois, USA, pp. i-vii, 5-14, 8-3.

Ilfeld et al., "Delivery Rate Accuracy of Portable, Bolus-Capable Infusion Pumps Used for Patient-Controlled Continuous Regional Analgesia", *Regional Anesthesia and Pain Medicine*, Jan.-Feb. 2003, vol. 28, No. 1, pp. 17-23.

Ilfeld et al., "Portable Infusion Pumps Used for Continuous Regional Analgesia: Delivery Rate Accuracy and Consistency", *Regional Anesthesia and Pain Medicine*, Sep.-Oct. 2003, vol. 28, No. 5, pp. 424-432.

JMS Co., Ltd., "Infusion Pump: OT-701", Tokyo, Japan, 2002, pp. 4.

Kim, M.D., et al., "Hyperglycemia Control of the Nil Per Os Patient in the Intensive Care Unit: Introduction of a Simple Subcutaneous Insulin Algorithm", Nov. 2012, *Journal of Diabetes Science and Technology*, vol. 6, No. 6, pp. 1413-1419.

Kutcher et al., "The Effect of Lighting Conditions on Caries Interpretation with a Laptop Computer in a Clinical Setting", *Elsevier*, Oct. 2006, vol. 102, No. 4, pp. 537-543.

Lamsdale et al., "A Usability Evaluation of an Infusion Pump by Nurses Using a Patient Simulator", *Proceedings of the Human Factors and Ergonomics Society 49th Annual Meeting*, Sep. 2005, pp. 1024-1028.

Logan et al., "Fabricating Capacitive Micromachined Ultrasonic Transducers with a Novel Silicon-Nitride-Based Wafer Bonding Process", *IEEE Transactions on Ultrasonics, Ferroelectrics, and Frequency Control*, May 2009, vol. 56, No. 5, pp. 1074-1084.

Magaji et al., "Inpatient Management of Hyperglycemia and Diabetes", *Clinical Diabetes*, 2011, vol. 29, No. 1, pp. 3-9.

Mauseth et al., "Proposed Clinical Application for Tuning Fuzzy Logic Controller of Artificial Pancreas Utilizing a Personalization Factor", *Journal of Diabetes Science and Technology*, Jul. 2010, vol. 4, No. 4, pp. 913-922.

Maynard et al., "Subcutaneous Insulin Order Sets and Protocols: Effective Design and Implementation Strategies", *Journal of Hospital Medicine*, Sep./Oct. 2008, vol. 3, Issue 5, Supplement 5, pp. S29-S41.

Merry et al., "A New, Safety-Oriented, Integrated Drug Administration and Automated Anesthesia Record System", *Anesthesia & Analgesia*, Aug. 2001, vol. 93, No. 2 pp. 385-390.

Microchip Technology Inc., "MTA11200B; TrueGauge™ Intelligent Battery Management I.C.", <https://www.elektronik.ropla.eu/pdf/stock/mcp/mta11200b.pdf>, 1995, pp. 44.

Moghissi, Etie, MD, FACP, FACE, "Hyperglycemia in Hospitalized Patients", *A Supplement to ACP Hospitalist*, Jun. 15, 2008, pp. 32.

(56)

References Cited**OTHER PUBLICATIONS**

Nuckols et al., "Programmable Infusion Pumps in ICUs: An Analysis of Corresponding Adverse Drug Events", *Journal of General Internal Medicine*, 2007, vol. 23, Supp. 1, pp. 41-45.

Pretty et al., "Hypoglycemia Detection in Critical Care Using Continuous Glucose Monitors: An in Silico Proof of Concept Analysis", *Journal of Diabetes Science and Technology*, Jan. 2010, vol. 4, No. 1, pp. 15-24.

Saager et al., "Computer-Guided Versus Standard Protocol for Insulin Administration in Diabetic Patients Undergoing Cardiac Surgery", *Annual Meeting of the American Society of Critical Care Anesthesiologists*, Oct. 13, 2006.

Sebald et al., "Numerical Analysis of a Comprehensive in Silico Subcutaneous Insulin Absorption Compartmental Model", *31st Annual International Conference of the IEEE Engineering in Medicine and Biology Society*, Sep. 2-6, 2009, pp. 3901-3904.

SGS-Thomson Microelectronics, "L6219—Stepper Motor Drive", Datasheet, Dec. 1996, pp. 10.

SGS-Thomson Microelectronics, "PBL3717A—Stepper Motor Drive", Datasheet, Apr. 1993, pp. 11.

Simonsen, Michael Ph.D., POC Testing, New Monitoring Strategies on Fast Growth Paths in European Healthcare Arenas, *Biomedical Business & Technology*, Jan. 2007, vol. 30, No. 1, pp. 1-36.

Smith, Joe, "Infusion Pump Informatics", *CatalyzeCare: Transforming Healthcare*, as printed May 12, 2011, pp. 2.

Tang et al., "Linear Dimensionality Reduction Using Relevance Weighted LDA", *Pattern Recognition*, 2005, vol. 38, pp. 485-493, http://staff.ustc.edu.cn/~ketang/papers/TangSuganYaoQin_PR04.pdf.

Thomas et al., "Implementation of a Tight Glycaemic Control Protocol Using a Web-Based Insulin Dose Calculator", *Anaesthesia*, 2005, vol. 60, pp. 1093-1100.

Van Den Berghe, M.D., Ph.D., et al., "Intensive Insulin Therapy in Critically Ill Patients", *The New England Journal of Medicine*, Nov. 8, 2001, vol. 345, No. 19, pp. 1359-1367.

Van Den Berghe, M.D., Ph.D., et al., "Intensive Insulin Therapy in the Medical ICU", *The New England Journal of Medicine*, Feb. 2, 2006, vol. 354, No. 5, pp. 449-461.

Westbrook et al., "Errors in the Administration of Intravenous Medications in Hospital and the Role of Correct Procedures and Nurse Experience", *BMJ Quality & Safety*, 2011, vol. 20, pp. 1027-1034.

Zakariah et al., "Combination of Biphasic Transmittance Waveform with Blood Procalcitonin Levels for Diagnosis of Sepsis in Acutely Ill Patients", *Critical Care Medicine*, 2008, vol. 36, No. 5, pp. 1507-1512.

Abbott Laboratories, "LifeCare® 5000, Plum®: Concurrent Flow Infusion System with DataPort™", *System Operating Manual*, Version 1.6, Jul. 1998, pp. 76.

International Search Report and Written Opinion received in PCT Application No. PCT/US2021/062072, dated Mar. 18, 2022 in 14 pages.

International Preliminary Report on Patentability and Written Opinion received in PCT Application No. PCT/US2021/062072, dated Jun. 22, 2023 in 11 pages.

Junda, Lin, "Global development trends of green bonds", *Jul. 10, 2018*, pp. 9.

Notice of Opposition in European Patent Application No. 16759381.3 (Patent No. EP3285827), dated May 31, 2023 in 48 pages.

Response to Notice of Opposition in European Patent Application No. 16759381.3 (Patent No. EP3285827), dated Oct. 23, 2023 in 63 pages.

Amazon, Post-it Message "Sign Here" Flags, 30/Dispenser, 4 Dispensers/Pack, Published Sep. 29, 2016, <https://www.amazon.com/Assorted-Color-Colors-Dispenser-MMM684SH/dp/B00006JNMN/?th=1>, * pages.

File:Soldado, Raso Fuerza Aerea Boliviana.jpg, Feb. 9, 2021, https://commons.m.wikimedia.org/wiki/File:Soldado_Raso_Fuerza_A%C3%A9rea_Boliviana.jpg, 1 page.

Fresenius, "Infusion Workstation: Orchestra® Base Intensive", *Operator's Guide*, Jun. 20, 2006, pp. 24. <<https://manualmachine.com/fresenius/orchestrabaseunit/7455278-user-manual/>>.

"ICU Medical Receives FDA Clearance for New Infusion Pump", *Medical Design & Development Staff*, Aug. 29, 2023, <https://www.medicaldesigndevelopment.com/topics/devices/news/22871498/icu-medical-receives-fda-clearance-for-new-infusion-pump>. 2 pages.




FIG. 1A




FIG. 1B




FIG. 2




FIG. 3




FIG. 4A




FIG. 4B




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 6




FIG. 7A

FIG. 7B

FIG. 7C




FIG. 7D




FIG. 7E

1

**CONCURRENT INFUSION WITH COMMON
LINE AUTO FLUSH****CROSS-REFERENCE TO RELATED
APPLICATIONS**

The present application is a continuation of U.S. Ser. No. 17/114,359, filed Dec. 7, 2020, now U.S. Pat. No. 11,135,360, which is incorporated by reference herein. The present application is also related to U.S. application Ser. No. 16/301,379, filed Nov. 13, 2018, which is the national stage of International Application No. PCT/US2017/032017, filed May 10, 2017, which claims the benefit of priority from U.S. Provisional No. 62/336,191, filed May 13, 2016, all of which are incorporated by reference in their entireties.

BACKGROUND**Field**

The present invention relates to medical devices and infusion pump systems.

Infusion pumps are medical devices that deliver fluids, including nutrients and medications such as antibiotics, chemotherapy drugs, and pain relievers, in controlled amounts. Many types of pumps, including large volume, patient-controlled analgesia (PCA), elastomeric, syringe, enteral, and insulin pumps, are used worldwide in healthcare facilities, such as hospitals, and in the home. Clinicians and patients rely on pumps for safe and accurate administration of fluids and medications.

It is often desirable to provide more than one therapeutic fluid to the patient from the same infusion pump. Two fluid reservoirs with different therapeutic fluids are connected to the infusion pump and then delivered through a common line having a terminal fluid delivery end. The terminal fluid delivery end is attached to the patient. The first therapeutic fluid and second therapeutic fluid may be administered concurrently or one at a time by controlling the fluid flow path to draw fluid from both reservoirs or from only one reservoir.

When switching from single to concurrent fluid delivery, the therapeutic fluid remaining in the common line may lead to complexity in controlling delivery volumes or flow rates when switching between fluid sources. For example, the remaining therapeutic fluid must be cleared from the common line before the next therapeutic fluid begins administration (entering the patient's body), which delays the next therapeutic fluid from reaching the patient. In addition, when the therapeutic fluids are administered concurrently, the first therapeutic fluid remaining in the common line will be administered at the combined rate of the first therapeutic fluid infusion rate plus the second therapeutic fluid infusion rate, e.g., the remaining first therapeutic fluid will be administered at the combined rate determined from the rates specified for the first and second therapeutic fluids. This can result in the patient receiving more or less than the optimum therapy with respect to the first therapeutic fluid. Furthermore, the remaining therapeutic fluid may not be correctly accounted for, potentially creating delays in the values indicated at the infusion pump, versus therapeutic fluid received by the patient. Finally, a single medication delivered at a combined rate may actually result in the single medication being infused at a rate that can exceed an upper soft or hard limit specified for such medication until the medication in the common line is displaced by an intended second fluid (in the case of a piggyback infusion) or a

2

mixture of first and second fluids (in the case of a concurrent delivery). While the pump data will be correct in terms of infusion rates over given times, the actual fluid delivery to the terminal fluid delivery end at the patient may not be correctly captured in pump and system data.

In addition, while some infusion therapies specify a particular volume of fluid to infuse to a patient, in some therapies it is preferred to deliver 100% of the volume of fluid contained within a particular fluid reservoir, such that the fluid is delivered until the reservoir is emptied. However, with many infusion pump systems, due to variable fluid volume contained in the reservoir and typical pump delivery accuracy tolerances and system dependencies, it is only possible to achieve 100% fluid delivery by over-programming the pump, or by entering pump programming parameters that do not accurately reflect the volume and duration of fluid actually administered to the patient.

It would be desirable to have infusion pump systems and methods with common line auto flush that would overcome the above disadvantages.

SUMMARY

In one embodiment, a control system is provided to control operation of an infusion pump of an infusion pump system. The infusion pump system includes a first reservoir configured to hold a first fluid, a second reservoir configured to hold a second fluid, a junction in fluid communication with the first reservoir and the second reservoir, a common line in fluid communication with the junction and having a terminal fluid delivery end, and the infusion pump, wherein the infusion pump is operable to drive fluid through the common line toward the terminal fluid delivery end. The control system includes: one or more hardware processors; and a memory storing executable instructions that when executed by the one or more hardware processors, configure the infusion pump to: receive instructions to deliver the first fluid at a first rate, subsequently concurrently deliver a mixture of the first fluid and the second fluid, and concurrently deliver the first fluid at the first rate and the second fluid at a second rate; infuse the first fluid at the first rate along a first flow path, the first flow path including the common line; determine a common line volume corresponding to a volume of the common line; draw the first fluid from the first reservoir the second fluid from the second reservoir to deliver the mixture of the first fluid and the second fluid; infuse the mixture of the first fluid and the second fluid at a flushing rate along a second flow path, the second flow path including the common line; determine that an infused volume of the mixture of the first fluid and the second fluid equals or exceeds the common line volume; and change the infusion rate of the mixture of the first fluid and the second fluid from the flushing rate to a combined rate, wherein the combined rate is the sum of the first rate and the second rate, and continue to infuse the mixture of the first fluid and the second fluid along the second flow path at the combined rate.

The control system may also include a mixing chamber in fluid communication with the first reservoir, the second reservoir, and the common line. The executable instructions may further configure the infusion pump to determine the flushing rate based upon whether the first fluid is a medicinal fluid, determine the flushing rate as the first rate when the first fluid is a medicinal fluid, or determine the flushing rate as the first rate increased by a flushing rate factor when the first fluid is not a medicinal fluid.

The instructions may further configure the infusion pump to receive the common line volume from a user input,

retrieve the common line volume from the memory, or retrieve the common line volume over a network. The common line volume may be predetermined. The instructions may further configure the infusion pump to determine the common volume based on the first fluid. The first rate may be different than the second rate.

The instructions may further configure the infusion pump to receive the instructions for the delivery from an input via a user interface. The executable instructions further configure the infusion pump to: determine that an infusion of the second fluid has completed; draw the first fluid from the first reservoir without drawing the second fluid from the second reservoir; infuse the first fluid at the combined rate; determine that a volume of the first fluid infused at the combined rate equals or exceeds the common line volume; and change the infusion rate of the first fluid from the combined rate to the first rate.

The executable instructions may configure the infusion pump to determine that an infusion of the second fluid has completed by comparing a volume of fluid infused to a programmed volume to infuse, determine that an infusion of the second fluid has completed by receiving an instruction to stop infusing the second fluid, or determine that an infusion of the second fluid has completed by determining that the second reservoir has been depleted of second fluid.

The executable instructions may further configure the infusion pump to: determine that an infusion of the first fluid has completed; draw the second fluid from the second reservoir without drawing the first fluid from the first reservoir; infuse the second fluid at the combined rate; determine that a volume of the second fluid infused at the combined rate equals or exceeds the common line volume; and change the infusion rate of the second fluid from the combined rate to the second rate.

The executable instructions may configure the infusion pump to determine that an infusion of the first fluid has completed by comparing a volume of fluid infused to a programmed volume to infuse, determine that an infusion of the first fluid has completed by receiving an instruction to stop infusing the first fluid, or determine that an infusion of the first fluid has completed by determining that the first reservoir has been depleted of first fluid.

In another embodiment, a method for controlling operation of an infusion pump of an infusion pump system is provided. The infusion pump system includes a first reservoir configured to hold a first fluid, a second reservoir configured to hold a second fluid, a junction in fluid communication with the first reservoir and the second reservoir, a common line in fluid communication with the junction and having a terminal fluid delivery end, and the infusion pump, wherein the infusion pump is operable to drive fluid through the common line toward the terminal fluid delivery end. The method includes: drawing the first fluid from the first reservoir and the second fluid from the second reservoir to form a mixture of the first fluid and the second fluid; infusing the mixture of the first fluid and the second fluid at a combined rate, wherein the combined rate is a sum of a first infusion rate associated with the first fluid and a second infusion rate associated with the second fluid; determining a common line volume corresponding to a volume of the common line; determining that the second reservoir is depleted; drawing the first fluid from the first reservoir without drawing the second fluid from the second reservoir; driving the first fluid at the combined rate along a flow path including the common line; determining that a driven volume of the first fluid equals or exceeds the common line volume; and changing the infusion rate of the first fluid from

the combined rate to the first rate, and continuing to infuse the first fluid along the flow path at the first rate.

The infusion pump may also include a mixing chamber in fluid communication with the first reservoir, the second reservoir, and the common line. Determining the common line volume may include receiving the common line volume from a user input, retrieving the common line volume from a memory, or retrieving the common line volume over a network. The common line volume may be predetermined.

Determining the common line volume may include determining the common line volume based on the first fluid. The first rate may be different than the second rate. Driving the first fluid at the combined rate may include driving the first fluid at a rate that exceeds a drug library rate limit associated with the first fluid. Determining that the second reservoir is depleted may include receiving a sensor signal that air is present in the junction or in a line coupling the junction to the second reservoir.

The method may further include pumping the first fluid from the first reservoir towards the second reservoir in response to receiving the sensor signal that air is present in the junction or in the line coupling the junction to the second reservoir.

In yet another embodiment, a control system for controlling operation of an infusion pump of an infusion pump system is provided. The infusion pump system includes a first reservoir configured to hold a first fluid, a second reservoir configured to hold a second fluid, a junction in fluid communication with the first reservoir and the second reservoir, a common line in fluid communication with the junction and having a terminal fluid delivery end, and the infusion pump, wherein the infusion pump is operable to drive fluid through the common line toward the terminal fluid delivery end. The control system includes: one or more hardware processors; and a memory storing executable instructions that when executed by the one or more hardware processors, configure the infusion pump to: draw the first fluid from the first reservoir and the second fluid from the second reservoir to form a mixture of the first fluid and the second fluid; infuse the mixture of the first fluid and the second fluid at a combined rate, wherein the combined rate is a sum of a first infusion rate associated with the first fluid and a second infusion rate associated with the second fluid; determine a common line volume corresponding to a volume of the common line; draw the first fluid from the first reservoir without drawing the second fluid from the second reservoir; drive the first fluid at the combined rate along a flow path including the common line; determine that a driven volume of the first fluid equals or exceeds the common line volume; and change the infusion rate of the first fluid from the combined rate to the first rate, and continue to infuse the first fluid along the flow path at the first rate.

The infusion pump may also include a mixing chamber in fluid communication with the first reservoir, the second reservoir, and the common line.

The executable instructions may also configure the infusion pump to determine the common line volume by receiving the common line volume from a user input, determine the common line volume by retrieving the common line volume from a memory, or determine the common line volume by retrieving the common line volume over a network. The common line volume may be predetermined.

The executable instructions may also configure the infusion pump to determine the common line volume by determining the common line volume based on the first fluid. The first rate may be different than the second rate.

5

The executable instructions may configure the infusion pump to drive the first fluid at the combined rate by driving the first fluid at a rate that exceeds a drug library rate limit associated with the first fluid. The executable instructions may configure the infusion pump to determine that the second reservoir is depleted by receiving a sensor signal that air is present in the junction or in a line coupling the junction to the second reservoir. The executable instructions may further configure the infusion pump to pump the first fluid from the first reservoir towards the second reservoir in response to receiving the sensor signal that air is present in the junction or in the line coupling the junction to the second reservoir.

In yet another embodiment, a method for controlling operation of an infusion pump of an infusion pump system is provided. The infusion pump system includes a first reservoir configured to hold a first fluid, a second reservoir configured to hold a second fluid, a junction in fluid communication with the first reservoir and the second reservoir, a common line in fluid communication with the junction and having a terminal fluid delivery end, and the infusion pump, wherein the infusion pump is operable to drive fluid through the common line toward the terminal fluid delivery end. The method includes: receiving instructions to deliver the first fluid at a first rate, subsequently concurrently deliver a mixture of the first fluid and the second fluid, and concurrently deliver the first fluid at the first rate and the second fluid at a second rate; infusing the first fluid at the first rate along a first flow path, the first flow path including the common line; determining a common line volume corresponding to a volume of the common line; drawing the first fluid from the first reservoir the second fluid from the second reservoir to deliver the mixture of the first fluid and the second fluid; infusing the mixture of the first fluid and the second fluid at a flushing rate along a second flow path, the second flow path including the common line; determining that an infused volume of the mixture of the first fluid and the second fluid equals or exceeds the common line volume; and changing the infusion rate of the mixture of the first fluid and the second fluid from the flushing rate to a combined rate, wherein the combined rate is the sum of the first rate and the second rate, and continue to infuse the mixture of the first fluid and the second fluid along the second flow path at the combined rate.

The infusion pump system may also include a mixing chamber in fluid communication with the first reservoir, the second reservoir, and the common line. The method may also include determining the flushing rate based upon whether the first fluid is a medicinal fluid, determining the flushing rate as the first rate when the first fluid is a medicinal fluid, or determining the flushing rate as the first rate increased by a flushing rate factor when the first fluid is not a medicinal fluid.

The method may also include receiving the common line volume from a user input, retrieving the common line volume from the memory, or retrieving the common line volume over a network. The common line volume may be predetermined.

The method may also include determining the common volume based on the first fluid. The first rate may be different than the second rate. Infusing the mixture of the first fluid and the second fluid at the flushing rate may include one or more of delivering the first fluid at a first fluid flush rate that exceeds a drug library rate limit associated with the first fluid or delivering the second fluid at a second fluid flush rate that exceeds a drug library rate limit associated with the second fluid.

6

The method may also include: determining that an infusion of the second fluid has completed; drawing the first fluid from the first reservoir without drawing the second fluid from the second reservoir; infusing the first fluid at the combined rate; determining that a volume of the first fluid infused at the combined rate equals or exceeds the common line volume; and changing the infusion rate of the first fluid from the combined rate to the first rate.

The method may also include determining that an infusion of the second fluid has completed by comparing a volume of fluid infused to a programmed volume to infuse, determining that an infusion of the second fluid has completed by receiving an instruction to stop infusing the second fluid, or determining that an infusion of the second fluid has completed by determining that the second reservoir has been depleted of second fluid. Infusing the first fluid at the combined rate may include infusing the first fluid at a rate that exceeds a drug library rate limit associated with the first fluid.

The method may also include: determining that an infusion of the first fluid has completed; drawing the second fluid from the second reservoir without drawing the first fluid from the first reservoir; infusing the second fluid at the combined rate; determining that a volume of the second fluid infused at the combined rate equals or exceeds the common line volume; and changing the infusion rate of the second fluid from the combined rate to the second rate.

The method may also include determining that an infusion of the first fluid has completed by comparing a volume of fluid infused to a programmed volume to infuse, determining that an infusion of the first fluid has completed by receiving an instruction to stop infusing the first fluid, or determining that an infusion of the first fluid has completed by determining that the first reservoir has been depleted of first fluid. Infusing the second fluid at the combined rate may include infusing the second fluid at a rate that exceeds a drug library rate limit associated with the second fluid.

The foregoing and other features and advantages of the invention will become further apparent from the following detailed description of the presently preferred embodiments, read in conjunction with the accompanying drawings. The detailed description and drawings are merely illustrative of the invention rather than limiting. The scope of the invention is defined by the appended claims and equivalents thereof.

In certain embodiments, a control system can control operation of an infusion pump system. The infusion pump system can include a first reservoir that can hold a first fluid, a second reservoir configured to hold a second fluid, a junction in fluid communication with the first reservoir and the second reservoir, a common line in fluid communication with the junction and having a terminal fluid delivery end, and an infusion pump operable to drive fluid through the common line toward the terminal fluid delivery end. The control system can control whether fluids from the reservoirs are drawn individually or concurrently (e.g., simultaneously or in an alternating manner). For example, the control system can include a flow control mechanism to manipulate a flow path at the junction to draw fluid from the first reservoir alone, the second reservoir alone, or both first and second reservoirs in an alternating manner.

The first reservoir may be referred to as the primary source and the second reservoir may be referred to as the secondary source. During a primary infusion, fluid is infused from the first, or primary reservoir, into the junction, and through the common line to the terminal end (and into the patient) at a first infusion rate. During a secondary infusion, fluid is infused from the second, or secondary reservoir, into

the junction, and through the common line to the terminal end (and into the patient) at a second infusion rate. During a concurrent infusion (sometimes referred to as concurrent delivery), first and second fluids are infused simultaneously to a patient at respective first and second infusion rates. A first volume of the first fluid is drawn from the first reservoir, and a second volume of the second fluid is drawn from the second reservoir. The first and second volumes are proportionate to the first and second infusion rates. Once the first and second fluids have been drawn, the pump drives (e.g., pumps or pushes out) the fluid combination through the common line to the terminal and (and into the patient) at a combined rate.

The combined rate can be equal to one of the first or second infusion rates, or it can be determined from the first and second infusion rates. For example, the combined rate can be determined as the sum as the first and second infusion rates. In some cases, a maximum rate may be established, and if the sum of the programmed first and second rates exceeds the maximum rate, then the combined rate may be set to the maximum rate. Other methods of determining a combined rate using the first and second rates are possible, as well. In addition, if the combined rate equals or exceeds a predetermined maximum combined rate, the first and second rates may be reduced proportionally such that their sum is less than or equals the maximum combined rate. In other embodiments, if the combined rate equals or exceeds a predetermined maximum combined rate, only the first rate is reduced until the sum of the first and second rates is less than or equals the maximum combined rate. For example, only the first rate may be reduced based upon a determination of fluid types of the first and second fluids. If the first fluid is a non-medication and the second fluid is a medication, then in some embodiments, the only the first rate is reduced (or the first rate is reduced by an amount or proportion that is greater than an amount or proportion that the second rate is reduced), such that the sum of the first and second rates is less than or equal to the maximum combined rate. In such embodiments, the user would be presented with a suggested first and second rate for approval or confirmation via a user interface before changing and/or initiating an infusion according to such adjusted first and/or second rates.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A and 1B are block diagrams of infusion pump systems with concurrent fluid delivery and common line auto flush in accordance with the present invention.

FIG. 2 is a block diagram of an infusion pump with concurrent fluid delivery and common line auto flush in accordance with the present invention.

FIG. 3 is a schematic diagram of an infusion pump with concurrent fluid delivery and common line auto flush in accordance with the present invention.

FIGS. 4A and 4B are graphs of fluid volume delivered at the terminal fluid delivery end of the common line versus time for a method of use for an infusion pump with concurrent fluid delivery and common line auto flush in accordance with the present invention.

FIG. 5A is a flowchart of a method of concurrent fluid delivery and common line auto flush in accordance with the present invention that may be performed by the infusion pumps of FIGS. 1-3.

FIG. 5B is a flowchart of a method of providing a secondary infusion until the secondary reservoir is depleted in accordance with the present invention that may be performed by the infusion pumps of FIGS. 1-3.

FIG. 5C is a flowchart of a method of providing sequential infusions, where a primary infusion is provided until the primary reservoir is depleted in accordance with the present invention that may be performed by the infusion pumps of FIGS. 1-3.

FIG. 6 is a flowchart of a method of determining fluid drive start times to cause infusions to reach a patient at desired infusion start times in accordance with the present invention that may be performed by the infusion pumps of FIGS. 1-3.

FIGS. 7A-7E are schematic diagrams of use for an infusion pump system with concurrent fluid delivery and common line auto flush in accordance with the present invention.

Like elements share like reference numbers throughout the various figures.

DETAILED DESCRIPTION

Systems and methods that improve an infusion pump system with concurrent delivery and common line auto flush are described herein. An infusion pump can operate in a primary delivery mode and deliver a first fluid from a first reservoir at a first rate, and then switch to a concurrent delivery mode, such as by delivering a combination of the first fluid from the first reservoir and a second fluid from a second reservoir at a combined delivery rate. The pump may switch from a concurrent delivery mode to a primary delivery mode (or to a secondary delivery mode where a second fluid is delivered from a second reservoir at a second rate), as well.

As discussed above, first fluid will remain in the common line at the time the delivery mode is switched from primary delivery mode to concurrent delivery mode. Therefore, if the first and second fluids are delivered at the combined delivery rate as soon as the concurrent delivery mode begins, the first fluid remaining in the common line will be delivered into the patient at the incorrect (i.e., the combined) rate. Furthermore, delivering fluids at rates other than the desired rates may result in inaccurate therapy, which can be dangerous to the patient. The systems and methods described herein improve delivery and accurately account for the fluid remaining in the volume of the common line. Fluid as used herein can be any fluid suitable to be administered to a patient by infusion, including saline fluid, fluid including a drug or other therapeutic agent, or the like.

FIGS. 1A & 1B are block diagrams for embodiments of infusion pump systems with concurrent delivery and a common line. The infusion pump system illustrated in FIG. 1A includes a junction in fluid communication with the first reservoir and the second reservoir. An optional mixing chamber is located at the junction, or between the junction and the common line. The junction and/or mixing chamber is located internal to the infusion pump. In the embodiment of the infusion pump system illustrated in FIG. 1B a junction in fluid communication with the first reservoir and the second reservoir is located external to the infusion pump. An optional mixing chamber is located at the junction, or between the junction and the common line. The location of the junction, in part, determines the length and internal volume of the common line between the junction and the terminal fluid delivery end. The internal cross-sectional shape, which is usually substantially circular, and the diameter and length of the common line determine its internal volume. Other shapes can be used without detracting from the scope of the disclosure.

The infusion pump system 100 of FIG. 1A includes a junction 180 internal to the infusion pump 130 and an optional mixing chamber (not shown) at the junction or between the junction 180 and a common line 140. The infusion pump system 100 includes a first reservoir 110 that contains a first fluid 112; a second reservoir 120 that contains a second fluid 122; a junction 180 in fluid communication with the first reservoir 110 and the second reservoir 120; an optional mixing chamber (not shown); a common line 140 in fluid communication with the mixing chamber and/or the junction 180 at one end 140A and having a terminal fluid delivery end 140B for connection to the patient 102, and an infusion pump 130 operable to drive fluid through the common line 140.

Primary Infusion Mode

The infusion pump 130 is operable to operate in a primary infusion mode during which the infusion pump infuses the first fluid 112 at a first rate along a first flow path 150 that includes the first reservoir 110, the junction 180, the optional mixing chamber, and the common line 140. The infusion pump 130 is further operable to determine a common line flush volume value corresponding to the internal volume of the common line 140. The infusion pump 130 may determine the common line flush volume by receiving the value from an operator, receiving it over a network (e.g., from a drug library or other database), retrieving it from a memory of the infusion pump, or any other method described herein. Primary to Concurrent Infusion Mode with Auto Flush

The infusion pump 130 is further configured to change to a concurrent infusion mode by drawing a second fluid 122 from a second reservoir 120 along a second flow path 160 into the junction 180 and/or mixing chamber and mixing it with the first fluid 112, drawn from the first reservoir 110 via the first flow path 150. The second flow path 160 includes the second reservoir 120, the junction 180, and the optional mixing chamber. The infusion pump is configured to initially infuse the mixture at the first rate until the volume of the first fluid is flushed out of the common line 140. The infusion pump 130 is configured to monitor volume of the mixture of first and second fluids 112, 122 driven at the first rate and subsequently pump the mixture of first and second fluids 112, 122 at the programmed combined rate when the monitored volume of the mixture is equal to or greater than the common line flush volume value. In this case, the delivery rates of fluid 1 and fluid 2 would be reduced (scaled down) from programmed rates during displacement of the common line volume, and the pump system may allow an override of one or both lower rate limits, or other associated limits, defined respectively for each of fluid 1 and fluid 2, during this phase of delivery.

Alternatively, for example in a scenario where the first fluid is not a medication (e.g., saline) and it is desired to initiate delivery of the second fluid (that is a medication) rapidly, the infusion pump could be configured to initially infuse the mixture at a more rapid rate to quickly displace the relatively inert common line volume. In this case, the initial combined rate could be increased (scaled up) to the programmed first rate plus the programmed second rate until the monitored volume of the mixture is equal to the common line flush volume. In this case, the delivery rates of fluid 1 and/or fluid 2 may be increased above upper rate limits defined for those respective fluids and the pump system may allow override of those limits during this phase of delivery. Further, the resulting scaled combined rate will be applied to the common line fluid 1, whose upper rate limit may limit or define allowable increased combined rates during the common line displacement phase. In this case, drug library

defined limits would be considered and applied by the pump system at the point of infusion to the patient as well as per pump programming activity.

Concurrent to Primary Infusion Mode with Auto Flush

The infusion pump 130 is further configured to change from concurrent delivery to a primary infusion mode by refraining from drawing the second fluid 122 from the second reservoir 120, and by infusing only the first fluid 112 from the first reservoir 110 along the first flow path 150. When the infusion pump switches to primary infusion mode, the infusion pump is configured to initially drive the first fluid 112 at the combined rate until the volume of the mixture of first and second fluids 112, 122 is flushed out of the common line 140. The infusion pump 130 is configured to monitor volume of the first fluid 112 driven at the combined rate and subsequently pump the first fluid 112 at the first rate when the monitored first fluid volume is equal to or greater than the common line flush volume value. In one example, the infusion pump 130 can be a fluid displacement pump employing a cassette, such as the Plum 360™ infusion pump available from ICU Medical, Inc. of San Clemente, CA. Those skilled in the art will appreciate that the infusion pump 130 can be any type of pump operable to drive fluid from two reservoirs through a common line 140. In this case, driving of the first fluid at the combined rate during common line displacement may require that the pump system allows override of drug library-defined upper rate limits for the first fluid.

Concurrent to Secondary Infusion Mode with Auto Flush

In another embodiment, the infusion pump 130 is further configured to change from a concurrent delivery to a secondary infusion mode by refraining from drawing the first fluid 112 from the first reservoir 110, and by infusing only the second fluid 122 from the second reservoir 120 along the second flow path 160. For example, if the first fluid 112 infusion is completed or is stopped, the infusion pump 130 may automatically switch to a secondary infusion mode. In such case, the infusion pump 130 will stop drawing first fluid 112 from the first reservoir 110, and it will continue to pump at the combined rate until the common line is cleared of the fluid mixture. In this case, the pump system may need to allow an override of upper rate limits for the second fluid while it is pumped at the combined rate during common line displacement. The infusion pump 130 is configured to monitor volume of the second fluid 122 driven at the combined rate and subsequently pump the second fluid 122 at the second rate when the monitored second fluid volume is equal to or greater than the common line flush volume value.

In one embodiment, the infusion pump 130 can be operably connected to a medication management unit (MMU) 170 or a server over a hospital network and/or the Internet, to receive a drug library (or other database), which may specify an appropriate common line flush volume value. For example, the drug library (or other database) may include information regarding the volume of various tubing assemblies, each tubing assembly including a common line. The infusion pump (or server) may be configured to determine a tubing assembly identifier associated with a tubing assembly that is attached to the infusion pump and the patient 102. The infusion pump may determine the tubing assembly identifier by receiving it from a server over the hospital network and/or the Internet, by receiving it via manual data entry by an operator, and/or by reading the tubing assembly identifier from the tubing assembly (or by other methods). For example, a tag, such as an RFID tag, an NFC tag or other wireless tag, may include the tubing assembly identifier. A

11

tag reader incorporated into or in communication (directly or indirectly) with the infusion pump, may read the tag to determine the tubing assembly identifier. The common line flush volume value may be determined using the tubing assembly identifier and the drug library (or database).

In one embodiment, the infusion pump **130** can be further operable to increment a displayed value of first fluid volume by the monitored volume when the mixture of first and second fluids **112**, **122** are driven at the first rate. The infusion pump **130** can be further operable to increment a displayed value of first and second fluid volumes when the first fluid **112** is driven at the combined rate. In one embodiment, the infusion pump **130** is operable to monitor the volume of infused first fluid **112** and switch to a concurrent infusion mode when the volume of the infused first fluid is equal to a Volume To Be Infused (VTBI) for the first fluid or when the volume of the infused first fluid is equal to the VTBI for the first fluid minus the volume of the common line. In one embodiment, the infusion pump **130** is operable to monitor the volume of infused second fluid **112** during a concurrent infusion mode and switch to a primary infusion mode when the volume of the infused second fluid is equal to a Volume To Be Infused (VTBI) for the second fluid or when the volume of the infused second fluid is equal to the VTBI for the second fluid minus the volume of the common line.

The infusion pump **130** can be operable to receive the common line flush volume value for the common line **140** automatically from the drug library stored in a memory locally in the infusion pump system **100** or remotely on a server. In one example, the drug library associates the common flush volume value with a particular therapeutic agent. In some cases, the drug library may include an indication (e.g., a flag, value, etc.) that a particular fluid is a rate dependent medicinal fluid whose action is rate dependent. The infusion pump **130** may be configured to infuse such fluids (whether alone or concurrently with a second fluid) at the infusion rate specified for such fluids. In another example, the drug library associates the common flush volume value with a particular clinical care area (CCA), such as general care, an intensive care unit (ICU), a neonatal ICU, or the like. In yet another example, the drug library associates the common flush volume value with a particular consumable infusion set, which provides the common line volume. The drug library can include upper and lower dosing limits with hard and soft limits for a number of therapeutic agents. In another embodiment, the infusion pump **130** can be operable to receive the common line flush volume value for the common line **140** from a caregiver via an input on a user interface of the infusion pump.

The common line **140** as illustrated includes the line between the junction **180** and the terminal fluid delivery end **140B** that is generally connectable to the patient **102** and includes any fluid path common to the first flow path **150** and the second flow path **160**. Thus, the common line **140** can include flow paths within the infusion pump **130** (including the associated consumable infusion set, when applicable) common to the first flow path **150** and the second flow path **160**, and is not limited to tubing external to the infusion pump **130**. The common line **140** is any portion of the infusion pump system **100** through which the first fluid **112** or a combination of the first fluid **112** and the second fluid **122** can alternately flow when switched. In one embodiment, the common line flush volume value is an internal volume of the common line **140**. The common line flush volume value can include an associated consumable infusion set volume, extension sets, filters, stopcocks, manifolds, patient access

12

devices, catheters, and the like. In another embodiment, the common line flush volume value is an internal volume of the common line **140** plus an adjustment volume. The adjustment volume can be any volume desired as a safety factor to assure that the common line **140** is free of the first fluid **112** before the second fluid **122** is infused at the second rate.

The infusion pump system **100'** of FIG. 1B has a junction **180'** external to the infusion pump **130'** and an optional mixing chamber (not shown) at the junction **180'** or between the junction **180'** and a common line **140'**. The infusion pump system **100'** includes a first reservoir **110'** containing a first fluid **112'**; a second reservoir **120'** containing a second fluid **122'**; a junction **180'** in fluid communication with the first reservoir **110'** and the second reservoir **120'**; an optional mixing chamber (not shown); a common line **140'** in fluid communication with the mixing chamber and/or the junction **180'** at one end **140a'** of the common line **140'** and a terminal fluid delivery end **140B'** that is generally connectable to the patient **102'**, and an infusion pump **130'** operable to drive fluid through the common line **140'**. The infusion pump **130'** is operable to: infuse the first fluid **112'** at a first rate along a first flow path **150'** including the first reservoir **110'**, the junction **180'**, the optional mixing chamber, and the common line **140'**; determine a common line flush volume value for the common line **140'**. The infusion pump **130'** may determine the common line flush volume by receiving the value from an operator, receiving it over a network (e.g., from a drug library or other database), retrieving it from a memory of the infusion pump, or any other method described herein.

The infusion pump **130'** is further configured to change to a concurrent infusion mode by drawing a second fluid from a second reservoir **120'** along a second flow path **160'** into the optional mixing chamber and mixing it with the first fluid, drawn from the first reservoir **110'** via the first flow path **150'**. The second flow path **160'** includes the second reservoir **120'**, the junction **180'**, and the optional mixing chamber. The infusion pump is configured to initially infuse the mixture at the first rate until the volume of the first fluid is flushed out of the common line **140'**. The infusion pump **130'** is configured to monitor volume of the mixture of first and second fluids **112'**, **122'** driven at the first rate and subsequently pump the mixture of first and second fluids **112'**, **122'** at a combined rate when the monitored volume is equal to or greater than the common line flush volume value.

The infusion pump **130'** is further configured to change to a primary infusion mode by refraining from drawing the second fluid **122'** from the second reservoir **120'**, and by infusing only the first fluid **112'** from the first reservoir **110'** along the first flow path **150'**. When the infusion pump **130'** switches to primary infusion mode, the infusion pump **130'** is configured to initially infuse the first fluid **112'** at the combined rate until the volume of the mixture of first and second fluids **112'**, **122'** is flushed out of the common line **140'**. The infusion pump **130'** is configured to monitor volume of the first fluid **112'** driven at the combined rate and subsequently pump the first fluid **112'** at the first rate when the monitored volume is equal to or greater than a common line flush volume value. The infusion pump **130'** is further configured to determine the common line flush volume value according to any of the methods described herein.

In one embodiment, the junction **180'** can include a two-way valve to manually or automatically switch the infusion pump system **100'** between the first flow path **150'** and the second flow path **160'**. In one example, the infusion pump **130'** can be a peristaltic pump. Those skilled in the art

13

will appreciate that the infusion pump **130'** can be any type of pump operable to drive fluid through the common line **140'**.

FIG. 2 is a block diagram of an embodiment of an infusion pump with concurrent fluid delivery and common line auto flush. The infusion pump **230** is operably connected to a common line **240** in fluid communication with a junction **280** and/or mixing chamber at one end **240A** and having a terminal fluid delivery end **240B** (not shown), the junction **280** being in fluid communication with a first reservoir (not shown) containing a first fluid and a second reservoir (not shown) containing a second fluid. In this example, a first reservoir line **211** provides fluid communication between the first reservoir and the junction **280** and a second reservoir line **221** provides fluid communication between the second reservoir and the junction **280**.

The infusion pump **230** includes a memory **233** operable to store programming code; a flow controller **235** operably connected to the memory **233**; and a fluid driver **232** operably connected to receive a control signal **231** from the flow controller **235**, the fluid driver **232** being operable to drive fluid through the common line **240**. The flow controller **235** is operable to execute the programming code and provide the control signal **231** to the fluid driver **232** in response to the programming code. The fluid driver **232** is responsive to the control signal **231** to infuse the first fluid at a first rate along a first flow path **211** including the first reservoir, the junction **280**, and the common line **240**; receive a common line flush volume value associated with the common line **240**; switch from infusing only the first fluid via the first flow path **250** to infusing a combination of the first fluid from the first reservoir and a second fluid from the second reservoir; drive the fluid combination at the first rate; monitor volume of the fluid combination driven at the first rate; and drive the fluid combination at a combined rate when the monitored volume is equal to or greater than the common line flush volume value. The fluid driver **232** is also responsive to the control signal **231** to infuse the fluid combination at the combined rate; switch to infusing only the first fluid via the first flow path **250**; drive the first fluid at the combined rate; monitor the volume of the first fluid driven at the combined rate; and drive the first fluid at the first rate when the monitored volume is equal to or greater than the common line flush volume value. The combined rate may be retrieved from the memory **233** or determined from a first infusion rate associated with the first fluid and a second infusion rate associated with the second fluid. For example, the combined rate may be determined as the sum of the first and second infusion rates.

In an embodiment, the flow controller **235** monitors the volume based on a time elapsed and a rate of delivery. The flow controller **235** can also monitor volume based on measurements, such as number of turns of a motor or signals from a sensor.

The flow controller **235** can include a hardware processor, microprocessor, or the like responsive to the programming code to generate the control signal **231**. The fluid driver **232** can include a metered pump, such as a cartridge pump, peristaltic pump, or the like, operable to drive fluid at a desired rate in response to the control signal **231**. In one embodiment, the fluid driver **232** can be further responsive to the control signal **231** to increment a displayed first fluid volume by the monitored volume when the fluid combination is driven at the first rate or when the monitored volume is equal to or greater than an internal volume of the common line **240**. The fluid driver **232** can be further responsive to the control signal **231** to increment displayed first and

14

second fluid volumes as the first fluid is driven at the combination rate or when the monitored volume is equal to or greater than the internal volume of the common line **240**. The first fluid displayed volume and/or the second fluid displayed volume can be displayed on a user interface **236**.

The memory **233** can also be operable to store data and other information, such as a drug library **234** (or other database) including the common flush volume value, which can optionally be associated with a particular therapeutic agent, a particular clinical care area, and/or a particular consumable infusion set. Different therapeutic agents may have different fluid properties and thus it may be advantageous in some embodiments to associate particular common flush volume value with particular therapeutic agents. In one embodiment, the infusion pump **230** can receive the common line flush volume value for the common line **240** automatically from the drug library **234**. In another embodiment, the infusion pump **230** can receive the common line flush volume value manually via direct entry of the value on a user interface **236**. The manual entry can be accomplished using a manufacturer provided volume value based upon the length and internal diameter of the common line **240** or a list number or other identifier that is used to access an associated volume value from a lookup table in the pump memory **233**, drug library, stored in a network location, at a server, or MMU. The possibility for manual typographical errors can be reduced by use of a barcode, radio frequency (RFID), optical, touch memory reader, near field communicator, or the like to input or scan a machine readable identifier on the infusion set, common line, or its package to obtain the volume value, the list number or other identifier associated with the volume value.

The infusion pump **230** can include human and/or machine interfaces as desired for a particular application. A user interface **236** operably connected to the flow controller **235** can provide input from and/or output to a caregiver or other user to the infusion pump **230**. Exemplary user interfaces can include display screens, soft keys or fixed keys, touchscreen displays, and the like. An I/O interface **237** operably connected to the flow controller **235** can provide input from and/or output to hardware associated with the infusion pump **230**. Exemplary I/O interfaces can include a wired and/or wireless interface to an electronic network, medication management unit (MMU), medication management system (MMS), or the like.

The common line flush volume value can be selected as desired for a particular application. The common line **240** includes the line between the junction **280** and the terminal fluid delivery end **240B**, and includes any fluid path common to the first flow path **250** and the second flow path **260** and so can include any portion of the infusion pump **230** (including the associated consumable infusion set) through which the first fluid or the second fluid can alternately flow or flow in a combined manner. In one embodiment, the common line flush volume value is equal to the internal volume of the common line **240**, so that the second fluid is infused at the second rate along the second flow path as soon as the first fluid has been cleared from the common line **240**. In another embodiment, the common line flush volume value is equal to the internal volume of the common line **240** plus an adjustment volume (to take into account the added/subtracted volume of other connectors or components), so that the second fluid is infused at the second rate along the second flow path after the first fluid has been cleared from the common line **240** plus the adjustment volume of the second fluid has been delivered at the first rate. In another embodiment, the common line flush volume value is equal

15

to the internal volume of the common line modified by a percentage, which could provide a desired overage or underage. The adjustment volume can be used as a safety factor to assure that the common line 240 is free of the first fluid before the second fluid is infused at the second rate.

FIG. 3 is a schematic diagram of an infusion pump with common line auto flush in accordance with the present invention. In this example, the infusion pump 330 includes a display 340, soft keys 350, and fixed keys 360 as a user interface. The display 340 provides operational and/or programming information to the user. The soft keys 350 perform different functions depending on the command displayed on an adjacent command portion 342 of the display 340. The fixed keys 360 are labeled with an input or function which functions the same, regardless of whatever is displayed on the display 340. In this example, the infusion pump 330 also includes a pump mechanism 370 operable to communicate with the first reservoir line and the second reservoir line and to move the first fluid or the second fluid to the terminal fluid delivery end of the common line.

FIGS. 4A & 4B are graphs of fluid volume delivered at the terminal end of the common line or patient versus time for a method of use for an infusion pump with common line auto flush in accordance with the present invention.

Referring to FIG. 4A, graph 510 is the fluid volume delivered at the terminal fluid delivery end of the common line for a first fluid versus time and graph 520 is the fluid volume delivered at the terminal fluid delivery end of the common line for a mixture of the first fluid and a second fluid versus time. From T1 to T2, the first fluid is infused at a first rate along a first flow path including the first reservoir and the second fluid is not infused. From T2 to T3, the first fluid is infused at a flushing rate greater than the first rate as a mixture of first and second fluids are drawn from first and second reservoirs, respectively, into the junction and/or mixing chamber and driven out at the flushing rate. For example, if the first fluid is a non-medicinal fluid (e.g., a saline solution, etc.), it may be desirable to flush the first fluid from the common line at an increased rate in order to infuse the second fluid into the patient as soon as possible. The flushing rate can be equal to the combined first rate plus second rate (as shown) or it can be determined by increasing the combined rate (e.g., the first rate plus the second rate by a flushing factor (e.g., 10%, 20%, 50%, 100%, etc.). The second fluid cannot be infused (e.g., it will not enter the patient) until the internal volume of the common line is cleared of the first fluid. From T3 to T4, the internal volume of the common line has been cleared of the first fluid and beginning at T3 the mixture of the first and second fluids are infused into the patient at a combined rate. From T4 to T5, auto flush is performed: the mixture of the first fluid and the second fluid is infused into the patient at the combined rate as only the first fluid is drawn into the junction and/or mixing chamber and driven out at the combined rate until the internal volume of the common line is cleared of the first and second fluid mixture. The first fluid cannot be infused by itself (e.g., it cannot enter the patient without the second fluid) until the internal volume is cleared of the first and second fluid mixture. After T5, the first fluid is infused at the first rate along the first flow path including the first reservoir after the internal volume of the common line has been cleared of the first and second fluid mixture. In this example, no additional second fluid is infused after T5, although in other embodiments, additional concurrent infusions (of first and second fluid mixtures) and/or secondary infusions (of just the second fluid) may be programmed to occur, as well.

16

Those skilled in the art will appreciate that the transition between the two infusion modes can be selected as desired for a particular application. In the example of FIG. 4A, a common line auto flush is performed from T4 to T5, but not from T2 to T3. As long as the common line flush volume value is known, the common line auto flush maintaining the first rate between T2 and T3 can be performed as desired.

Referring to FIG. 4B, graph 530 is the fluid volume delivered at the terminal fluid delivery end of the common line for a first fluid versus time and graph 540 is the fluid volume delivered at the terminal fluid delivery end of the common line for a mixture of the first fluid and a second fluid versus time. From T1 to T2, the first fluid is infused at the first rate along a first flow path including the first reservoir and the second fluid is not infused. From T2 to T3, auto flush occurs and the first fluid is infused at the first rate as a mixture of first and second fluids are drawn from first and second reservoirs, respectively, into a junction and/or mixing chamber, and driven out at a combined rate (as discussed above). The first fluid is infused, driven or displaced until the internal volume of the common line has been cleared of the first fluid. After T3, the mixture of the first and second fluids is infused, driven or displaced at the combined rate (as discussed above) after the internal volume of the common line has been cleared of the first fluid. In one embodiment, the common line is cleared of the first fluid when the monitored volume of the mixture of the first and second fluids driven at the first rate between T2 and T3 is equal to or greater than the common line flush volume value. In this example, no additional second fluid is infused after T3, although in other embodiments, additional concurrent infusions (of first and second fluid mixtures) and/or secondary infusions (of just the second fluid) may be programmed to occur, as well.

Concurrent Delivery with Common Line Auto Flush

FIG. 5A is a flowchart of an embodiment of a method for concurrent infusion with common line auto flush. The method 550 can be performed with any infusion pump system described herein. In one embodiment, the infusion pump system includes a first reservoir containing a first fluid, a second reservoir containing a second fluid, a junction in fluid communication with the first reservoir and the second reservoir, an optional mixing chamber at or in fluid communication with the junction, and a common line in fluid communication with the junction and/or mixing chamber at one end and having a terminal fluid delivery end, and an infusion pump operable to drive fluid through the common line. The method 550 can be performed by any of the systems discussed herein. In an embodiment, some or all aspects of the method 550 are stored as programmed instructions to be executed by an infusion pump flow controller (e.g., flow controller 235). The method 550 can be used with an infusion pump system and infusion pump as described in FIGS. 1A, 1B, & 2 above. A drug library may include an indication (e.g., flag, value, etc.) to enable or disable concurrent infusion with auto flush, as described with respect to FIG. 5A. In this example, the infusion pump infuses a first fluid on a first flow path at a first rate and switches to a concurrent infusion mode during which it infuses a mixture of the first fluid and a second fluid, maintaining the first rate long enough to clear the remaining first fluid from the common line before changing to a combined rate for infusing the mixture of the first and second fluids.

Referring to FIG. 5A, at block 552, the flow controller 235 determines a common line flush volume value. As discussed above, the common line flush value can be received based on a user input via any of the user interfaces

discussed above. In an embodiment, the flow controller **235** can automatically retrieve the common line flush volume value from the memory **233** or over a network (e.g., from a drug library or other database), or by wirelessly reading information from a tag associated with the common line and using the information to retrieve the common line flush volume from the memory or over the network. The common line flush volume may be predetermined for particular fluids. The common line flush volume may also depend on the VTBI or rate of the infusion.

At block **554**, a first infusion mode to infuse the first fluid at a first infusion rate begins. The first fluid is infused or driven at a first infusion rate along a first flow path that includes the first reservoir, the junction, the optional mixing chamber, and the common line. The infusion of the first fluid can be controlled by the flow controller **235** based on a control signal to activate the pump or other mechanical system. In some embodiments, the infusion of the first fluid can also be based on a user input or user control of the pump or the mechanical system. During the first infusion mode, the infusion pump drives the first fluid from the first reservoir at the first infusion rate. At block **556**, the flow controller **235** can determine to switch from the first infusion mode to a concurrent infusion mode. During an auto flush period, at block **556**, the infusion pump drives a mixture or combination of the first fluid and the second fluid toward the common line at the first rate. By driving the combination of the first and second fluids at the first rate, the first fluid remaining in the common line is flushed and delivered to the patient at the same rate as therapeutically required. In some embodiments, during the auto flush period, the infusion pump drives the combination of the first fluid and the second fluid at a combined rate, instead of the first rate. For example, it may be advantageous to use a combined rate to more quickly flush the common line, particularly when the fluid being flushed from the common line is a non-medicinal fluid, such as saline, or other non-medicinal fluid. The combined rate can be determined using any of the methods described herein. For example, the combined rate may be determined as the sum as the first and second rates. The flow controller **235** can use control signals to control the driving of the mixture of the first fluid and the second fluid and to control the rate of delivery. It also may be desirable to flush the common line of a non-medicinal first fluid such as saline, at a rate even higher than the combined rate to expedite delivery of the second medication. In scenarios where drug library-defined limits are assigned for one or both of the two fluid delivery rates, the pump system may allow overrides of the upper rate limit for one or both of the fluids during the common line flush. For example, the pump system could effectively apply these delivery limits upon delivery to the patient, versus upon delivery from the pump. In another embodiment, the method **550** of FIG. **5A** may be modified at block **556** such that the infusion pump drives a fluid combination at a rate that is a ratio of a first programmed first fluid rate and a programmed second fluid rate.

At block **558**, the flow controller **235** can monitor volume of the mixture of first and second fluids driven at the first rate. The flow controller **235** can determine when the monitored volume is equal to the common line flush volume value. When it is determined that the monitored volume equals or exceeds the common line flush volume, the method **550** proceeds to block **560**, where the flow controller **235** continues driving the mixture of the first and second fluids, but at the combined rate. In some embodiments, the flow controller **235** can measure an amount of time before changing the rate of the mixture fluid delivery to the

combined rate. In one embodiment, the flow controller **235** can further include incrementing a first fluid displayed volume and a second fluid displayed volume by a proportion of the monitored volume when the monitored volume is equal to or greater than an internal volume of the common line. The proportion of monitored volume to be incremented for each of the first and second fluids can be equal to the proportion of first and second flow rates associated with the first and second fluids, respectively. For example, if the first flow rate is 10 ml/hr and the second flow rate is 5 ml/hr, the proportions of the monitored value incremented on the first and second volume displays will have a 2:1 ratio. If the monitored volume is 3 ml, then the display of the first fluid value will be increased by 2 ml and the display of the second fluid value will be increased by 1 ml. The flow controller **235** can thus accurately track the rate, time, and an amount of each fluid delivered to the patient. In some embodiments, the flow controller **235** executes only some of the steps described above with respect to FIG. **5A**. Furthermore, the flow controller **235** can change the order of the steps, include additional steps, or modify some of the steps discussed above.

The common line flush volume value can be selected as desired for a particular application. In one embodiment, the common line flush volume value is an internal volume of the common line. In another embodiment, the common line flush volume value is an internal volume of the common line plus or minus an adjustment volume. The adjustment volume can be any volume desired as a safety factor to assure that the common line is free of the first fluid before the second fluid is infused at the second rate.

In one embodiment, the method **550** further includes incrementing a first fluid displayed volume by the monitored volume when driving a mixture of the first and second fluids at the first rate. The first fluid displayed volume is incremented by the monitored volume when the monitored volume is equal to or greater than an internal volume of the common line.

In some embodiments, the method **550** ends after the concurrent infusion at the combined rate ends. However, in other embodiments, the method **550** continues concurrent delivery of the first and second fluids until one of the fluids is depleted or until the desired volume of one of the fluids has been delivered. In such case, for example, when the second fluid reservoir is depleted, the infusion continues according to the method **580** discussed below with respect to FIG. **5B**. If instead, the concurrent infusion continues until the desired volume of one of the fluids has been delivered, then the infusion may continue according to a slightly modified method **580**, as discussed below with respect to FIG. **5B**.

Concurrent Delivery to Infusion Completion with Common Line Auto Flush

FIG. **5B** illustrates a method **580** of safely performing a concurrent infusion of first and second fluids until the volume of the second fluid reservoir is depleted (e.g., totally depleted or emptied of the second fluid), such that no second fluid or substantially no second fluid remains in the second reservoir. The method **580** can be performed by a flow controller (e.g., flow controller **235**) alone and/or in conjunction with the method **550** of FIG. **5A**. For example, method **580** may be performed beginning at block **586** and following block **560** of method **550** of FIG. **5A**. A drug library may include an indication (e.g., flag, value, etc.) to enable or disable infusion until depletion functionality, as described with respect to FIG. **5B**.

At block **582**, the method **580** determines a common line flush volume of a common line. Any of the methods described herein may be used to determine the common line flush volume. At block **584**, a concurrent infusion occurs, where a fluid combination is driven by an infusion pump at a combined rate. The fluid combination includes a mixture of a first fluid drawn into a junction and/or mixing chamber from a first reservoir and a second fluid drawn into the junction and/or mixing chamber from a second reservoir. As discussed herein, a first infusion rate may be associated with the infusion of the first fluid and a second infusion rate may be associated with the infusion of the second fluid. The ratio of the volumes of first and second fluids drawn into the mixing chamber is equal to the ratio of the ratio of first and second infusion rates. The fluid combination is driven from the junction and/or mixing chamber to the common line at a combined rate, which may be determined according to any of the methods described herein. For example, the combined rate may be determined as the sum of the first and second infusion rates.

At block **586**, the method **580** determines whether the second reservoir has been depleted. For example, a sensor can detect whether there is air or air bubbles in the line between the junction and the second reservoir. If the method **580** does not determine that the second reservoir is depleted, the method **580** returns to block **584**. If the method **580** determines that the second reservoir has been depleted, the method **580** proceeds to block **588**. The method **580** may also optionally cause the infusion pump to at least partially back-prime the line between the junction and the second reservoir. For example, the infusion pump may pump some fluid from the first reservoir to force fluid into the line between the junction and the second reservoir in order to remove air from the line (or at least the portion of the line near the junction).

In a modified version of method **580**, at block **586** the method **580** instead determines whether a desired or programmed volume of the second fluid has been delivered. For example, if the infusion pump was programmed to delivery only 100 ml of the second fluid during concurrent delivery mode, the method **580** would determine whether 100 ml of the second fluid had been delivered. In another embodiment, the method **580**, determines whether a desired volume of second fluid has been delivered by receiving a command to stop an infusion of the second fluid. When a user provides an input to stop the infusion, the method **580** determines that the desired volume of second fluid has been delivered. If so, the method **580** continues to block **588**. If not, the method **580** returns to block **584**.

At block **588**, the method **580** stops drawing fluid from the second reservoir, and instead only draws fluid from the first reservoir. The method **580** drives the first fluid to the common line at the combined rate in order to auto flush or clear the volume of the common line of the fluid combination remaining in the common line. In the case when there is a drug-library defined limit on the first fluid, the pump system may need to allow an override of this limit in order to support pumping of the first fluid at the combined rate. In other words, drug library-defined delivery limits for the first fluid would apply at the patient, versus at the pump.

At block **590**, the method **580** monitors the volume of first fluid driven at the combined rate and determine when the monitored volume equals or exceeds the common line flush volume. If the monitored volume is not equal to the common line flush volume, the method **580** returns to block **588**. If the monitored volume is equal to or exceeds the common line flush volume, the method **580** proceeds to block **592**.

At block **592**, the method **580** continues to draw the first fluid from the first reservoir, but at the first rate. In some embodiments, the method **580** can measure an amount of time before changing the rate of the first fluid delivery to the first rate. In one embodiment, the method **580** can further include incrementing a first fluid displayed volume and a second fluid displayed volume by a proportion of the monitored volume when the monitored volume is equal to or greater than an internal volume of the common line. The proportion of monitored volume to be incremented for each of the first and second fluids can be equal to the proportion of first and second flow rates associated with the first and second fluids, respectively. For example, if the first flow rate is 10 ml/hr and the second flow rate is 5 ml/hr, the proportions of the monitored value incremented on the first and second volume displays will have a 2:1 ratio. If the monitored volume is 3 ml, then the display of the first fluid value will be increased by 2 ml and the display of the second fluid value will be increased by 1 ml. The method **580** can thus accurately track the rate, time, and an amount of each fluid delivered to the patient. In some embodiments, the method **580** executes only some of the steps described above with respect to FIG. **5B**. Furthermore, the method **580** can change the order of the steps, include additional steps, or modify some of the steps discussed above.

Sequential Delivery to Reservoir Depletion with Common Line Auto Flush

FIG. **5C** illustrates a method **581** of safely performing a sequential infusion (sometimes referred to as a piggyback infusion) of a first fluid at a first infusion rate until the infusion of the first fluid at the first rate is stopped and the infusion switches to an infusion of a second fluid at a second infusion rate. The method **581** can be performed by a flow controller (e.g., flow controller **235**) alone and/or in conjunction with the method **550** of FIG. **5A** or the method **580** of FIG. **5B**. A drug library may include an indication (e.g., flag, value, etc.) to enable or disable infusion until reservoir depletion functionality, as described with respect to FIG. **5B**.

At block **583**, the method **581** determines a common line flush volume of a common line. Any of the methods described herein may be used to determine the common line flush volume. At block **585**, a primary infusion occurs, where a first fluid is driven by an infusion pump at a first infusion rate. The first fluid is drawn into a junction and/or mixing chamber from a first reservoir. The first infusion rate may be associated with the infusion of the first fluid and a second infusion rate may be associated with an infusion of the second fluid. The first fluid is driven from the junction and/or mixing chamber to the common line at the first infusion rate.

At block **587**, the method **581** determines whether to pause the first infusion and initiate a "piggyback" infusion, or infusion of a second fluid at a second rate. If the method **581** determines that the second fluid program should be initiated, the method **581** proceeds to block **589**. If not, the method **581** returns to block **585**.

At block **589**, the method **581** stops drawing fluid from the first reservoir (pauses the primary infusion), and instead only draws fluid from the second reservoir. The method **581** drives the second fluid to the common line at the first infusion rate in order to auto flush or clear the volume of the common line of the first fluid remaining in the common line. If the drug library includes limits on the delivery of fluid **2**, these limits may need to be allowed to be overridden during the common line flush period defined by block **589**. For example, if fluid **1** was programmed at a rate below the lower limit allowed for fluid **2**, or if fluid **1** was programmed

21

at a rate above the upper limit allowed for fluid 2, an override of such a limit would be allowed during the common line flush.

At block 591, the method 581 monitors the volume of second fluid driven at the first infusion rate and determines when the monitored volume equals or exceeds the common line flush volume. If the monitored volume is not equal to the common line flush volume, the method 581 returns to block 589. If the monitored volume is equal to or exceeds the common line flush volume, the method 581 proceeds to block 593.

At block 593, the method 581 continues to draw the second fluid from the second reservoir, but at the second infusion rate. In some embodiments, the method 581 can measure an amount of time before changing the rate of the first fluid delivery to the second infusion rate. In one embodiment, the method 581 can further include incrementing a first fluid displayed volume by the monitored volume when the monitored volume is equal to or greater than an internal volume of the common line. The method 581 can thus accurately track the rate, time, and an amount of each fluid delivered to the patient. In some embodiments, the method 581 executes only some of the steps described above with respect to FIG. 5B. Furthermore, the method 581 can change the order of the steps, include additional steps, or modify some of the steps discussed above. In some embodiments, it may be preferable to infuse the second fluid at a rate that exceeds the first infusion rate until the common line (filled with non-medicinal fluid) is cleared, in order to more quickly introduce the second (medicinal) fluid to the patient. If a drug library defined limit for fluid 2 is present, the pump system may permit an override of this limit to allow pumping of fluid 2 at this increased rate. Similarly, there may be limits on fluid 1 delivery rates that should be considered by the pump system, imposing a limit on fluid 2 pumping rates intended to displace common line volume. At block 595, the method 581 determines whether the piggyback infusion is complete. For example, the method 581 may determine that the second reservoir is depleted of fluid, that a desired volume of fluid has been infused, that a desired infusion duration period has been reached, etc. In one embodiment, a sensor determines that air is detected within the fluid line. If the piggyback infusion is not complete, the method 581 returns to block 593. If the piggyback infusion is complete, the method 581 proceeds to block 597. At block 591, the primary infusion, e.g., the infusion of the first fluid, is resumed, though at the second infusion rate until the driven first fluid volume is equal to or greater than the common line volume. In the case where the first fluid has a drug library defined limit(s), the pump system may need to support an override of such a lower or upper limit to support pumping at the rate programmed for the second fluid. Method 581 then continues to drive the first fluid, but now at the first infusion rate.

The method 581 may also optionally cause the infusion pump to at least partially back-prime the line between the junction and the second reservoir after air is recognized at the depletion of the second fluid reservoir. For example, the infusion pump may pump some fluid from the first reservoir to force fluid into the line between the junction and the second reservoir in order to remove air from the line (or at least the portion of the line near the junction). Intermittent Concurrent Delivery

FIG. 6 illustrates a method 600 of scheduling intermittent concurrent deliveries. Method 600 can be performed by an infusion pump, a flow controller (e.g., flow controller 235), and/or alone or in conjunction with the method 550 of FIG.

22

5A and/or method 580 of FIG. 5B. Method 600 may be performed when it is desired to deliver a secondary infusion (e.g., deliver a second fluid via a concurrent infusion with a first fluid) multiple times per day at specific start times.

Method 600 enables an infusion pump to determine a time to start an auto flush procedure to assure that the second fluid is infused into the patient (e.g., enters the patient) at the desired, specific start times.

For example, if a common line volume will take 10 minutes to flush at the primary (first) infusion rate, then the concurrent infusion will initiate an auto flush process (infusing a mixture of first and second fluids at the first infusion rate to flush the first fluid out of the common line tubing) 10 minutes before the desired secondary infusion start time (e.g., 10 minutes before the second fluid is to enter the patient).

At block 602, the method 600 determines a common line flush volume. However, the method may skip block 602 if the common line flush volume has already been determined.

At block 604, the method 600 determines one or more second fluid infusion start times. For example, the method 600 may receive or download schedule information corresponding to desired start times to infuse a second fluid into a patient. The schedule information may define specific times during the day (e.g., 8 am, noon, 4 pm, 8 pm, etc.), it may define a number of infusions per day (e.g., 2, 3, 4, 6 infusions per day, etc.), or it may define an interval between second fluid infusions (e.g., one bag of second fluid every 4 hours, etc.). The schedule information may be used to determine one or more second fluid infusion start times.

At block 606, the method 600 determines a flush time period based on the first fluid infusion rate (or first fluid flush rate if a faster flush rate is desired for the particular, e.g., non-medicinal, first fluid) and the common line flush volume. For example, the flush time period may be determined by dividing the common line flush volume by the first fluid infusion rate (or first fluid flush rate). The flush time period represents the amount of time it will take to flush remaining fluid from the common line between the junction (or mixing chamber) and the common line distal end when fluid is driven at the first (or first fluid flush) rate.

At block 608, the method 600 determines second fluid drive start times, which correspond to the actual times that the infusion pump will begin to draw first fluid from a first reservoir and second fluid from a second reservoir, and drive the mixture of first and second fluids to the common line at the first (or first fluid flush) rate. In one embodiment, the method 600 may determine the second fluid drive start times by subtracting the flush time period from each of the second fluid infusion start times. For example, if the flush time period is determined to be 20 minutes and the second fluid infusion start times are 8:00 am, 2:00 pm, and 8:00 pm, then the second fluid drive start times may be determined as 7:40 am, 1:40 pm, and 7:40 pm. By initiating an auto flush concurrent infusion at the second fluid drive start times, a mixture of the second fluid and the first fluid will reach the patient and will be infused into the patient (e.g., enter the patient's body) at the second fluid infusion start times. At block 610, the method causes the infusion pump to initiate such auto flush concurrent infusions at the second fluid drive start times.

FIGS. 7A-7E are schematic diagrams of use for an infusion pump system with concurrent infusion and common line auto flush in accordance with the present invention. FIGS. 7A-7E illustrate switching from infusing a first fluid to infusing a mixture or combination of first and second fluids, then switching back to infusing the first fluid, while

accounting for the previously infused fluid in the common line. In this example, the infusion pump is infusing a first fluid on a first flow path at a first rate and switches to infusing a mixture or combination of first and second fluids on a second flow path, maintaining the first rate long enough to clear the remaining first fluid from the common line before changing to a combined rate for infusing the mixture or combination of first and second fluids. The infusion pump then switches to infusing a first fluid on the first flow path, maintaining the combined rate long enough to clear the remaining mixture or combination of first and second fluids from the common line before changing to a first rate for infusing the first fluid.

Referring to FIG. 7A, the first fluid 712 is delivered to the terminal end 740B of a common line 740 at a first rate along a first flow path 750 including the first reservoir 710, the junction 780, an optional mixing chamber (not shown), and the common line 740. The first fluid 712 is indicated by upward from left to right diagonal lines. Referring to FIG. 7B, the infusion has changed to a concurrent mode. During the concurrent mode, first fluid 712 is drawn from the first reservoir 710 and second fluid 722 is drawn from the second reservoir 720 along a second flow path 760. The second fluid 722 is indicated by downward from left to right diagonal lines. The mixture of first and second fluids 712, 722 is driven by the infusion pump into the common line 740. During this auto flush mode of concurrent delivery, the common line 740 contains first common line fluid 741 remaining from the initial infusion of the first fluid 712 and indicated by the upward diagonal lines, and second common line fluid 742 (the mixture of the first and second fluids 712, 722) indicated by the hashed lines. The flow rate remains at the first rate because the remaining first common line fluid 741 is being delivered to the terminal fluid delivery end 740B or to the patient when connected. Referring to FIG. 7C, none of the first fluid remains in the common line 740, so the second common line fluid 743 (the mixture of the first and second fluids 712, 722) is driven at the combined rate.

The infusion pump system can subsequently switch back to infusing only the first fluid (for example, after a predetermined time period, after a predetermined volume of combined first and second fluids are infused, after a predetermined volume of the second fluid is infused, or after the infusion pump determines that the second reservoir has been depleted of the second fluid, etc.). Referring to FIG. 7D, the infusion mode has changed from concurrent delivery to primary delivery (infusing only first fluid 712 from the first reservoir 710). Initially, the common line 740 still contains a mixture of the first and second fluids 712, 722 (represented by the hashed lines) as second common line fluid 744 remaining from the previous infusion, and first common line fluid 745 (the first fluid 712 alone) indicated by the upward diagonal lines. The flow rate remains at the combined rate because the remaining second common line fluid 744 is being delivered. Referring to FIG. 7E, none of the mixture of first and second fluids remains in the common line 740, so the first common line fluid 746 is driven at the first rate along the first flow path 750.

While the embodiments of the invention disclosed herein are presently considered to be preferred, various changes, rearrangement of steps, and modifications can be made without departing from the scope of the invention. The scope of the invention is indicated in the appended claims, and all changes that come within the meaning and range of equivalents are intended to be embraced therein.

What is claimed is:

1. A method for controlling operation of an infusion pump of an infusion pump system, the infusion pump system comprising a first reservoir configured to hold a first fluid, a second reservoir configured to hold a second fluid, a junction in fluid communication with the first reservoir and the second reservoir, a common line in fluid communication with the junction and having a terminal fluid delivery end, and the infusion pump, wherein the infusion pump is operable to drive fluid through the common line toward the terminal fluid delivery end, the method comprising:

receiving instructions to deliver the first fluid at a first rate, subsequently concurrently deliver a mixture of the first fluid and the second fluid, and concurrently deliver the first fluid at the first rate and the second fluid at a second rate;

infusing the first fluid at the first rate along a first flow path, the first flow path including the common line;

determining a common line volume corresponding to a volume of the common line;

drawing the first fluid from the first reservoir and the second fluid from the second reservoir to deliver the mixture of the first fluid and the second fluid into a first end of the common line;

infusing the mixture of the first fluid and the second fluid at a flushing rate into the first end of the common line, wherein infusing the mixture of the first fluid and the second fluid at the flushing rate into the first end of the common line causes displacement of a volume of the first fluid remaining in the common line and infusion of the first fluid out of a terminal end of the common line at the flushing rate;

determining that an infused volume of the mixture of the first fluid and the second fluid equals or exceeds the common line volume; and

changing infusion of the mixture of the first fluid and the second fluid from the flushing rate to a combined rate in response to determining that the infused volume of the mixture of the first fluid and the second fluid equal or exceeds the common line volume, wherein the combined rate is a sum of the first rate and the second rate, and continue to infuse the mixture of the first fluid and the second fluid at the combined rate.

2. The method of claim 1, wherein the infusion pump system further comprises a mixing chamber in fluid communication with the first reservoir, the second reservoir, and the common line.

3. The method of claim 1, further comprising determining the flushing rate based upon whether the first fluid is a medicinal fluid.

4. The method of claim 3, further comprising determining the flushing rate as the first rate when the first fluid is a medicinal fluid.

5. The method of claim 3, further comprising determining the flushing rate as the first rate increased by a flushing rate factor when the first fluid is not a medicinal fluid.

6. The method of claim 1, further comprising receiving the common line volume from a user input.

7. The method of claim 1, further comprising retrieving the common line volume from a memory.

8. The method of claim 1, further comprising retrieving the common line volume over a network.

9. The method of claim 1, wherein the common line volume is predetermined.

10. The method of claim 1, further comprising determining the common line volume based on the first fluid.

11. The method of claim 1, wherein the first rate is different than the second rate.

25

12. The method of claim 1, wherein receiving the instructions further comprises receiving the instructions from an input via a user interface.

13. The method of claim 1, wherein infusing the mixture of the first fluid and the second fluid at the flushing rate comprises one or more of delivering the first fluid at a flush rate outside of drug library defined rate limits associated with the first fluid or delivering the second fluid at a second fluid flush rate outside of drug library defined rate limits associated with the second fluid.

14. The method of claim 1, further comprising:

determining that an infusion of the second fluid has completed;

drawing the first fluid from the first reservoir without drawing the second fluid from the second reservoir;

infusing the first fluid at the combined rate;

determining that a volume of the first fluid infused at the combined rate equals or exceeds the common line volume; and

changing infusion of the first fluid from the combined rate to the first rate.

15. The method of claim 14, further comprising determining that an infusion of the second fluid has completed by comparing a volume of fluid infused to a programmed volume to infuse.

16. The method of claim 14, further comprising determining that an infusion of the second fluid has completed by receiving an instruction to stop infusing the second fluid.

17. The method of claim 14, further comprising determining that an infusion of the second fluid has completed by determining that the second reservoir has been depleted of second fluid.

26

18. The method of claim 14, wherein infusing the first fluid at the combined rate comprises infusing the first fluid at a rate that exceeds a drug library rate limit associated with the first fluid.

19. The method of claim 1, further comprising:

determining that an infusion of the first fluid has completed;

drawing the second fluid from the second reservoir without drawing the first fluid from the first reservoir;

infusing the second fluid at the combined rate;

determining that a volume of the second fluid infused at the combined rate equals or exceeds the common line volume; and

changing infusion of the second fluid from the combined rate to the second rate.

20. The method of claim 19, further comprising determining that an infusion of the first fluid has completed by comparing a volume of fluid infused to a programmed volume to infuse.

21. The method of claim 19, further comprising determining that an infusion of the first fluid has completed by receiving an instruction to stop infusing the first fluid.

22. The method of claim 19, further comprising determining that an infusion of the first fluid has completed by determining that the first reservoir has been depleted of first fluid.

23. The method of claim 19, wherein infusing the second fluid at the combined rate comprises infusing the second fluid at a rate that exceeds a drug library rate limit associated with the second fluid.

* * * * *