



US 20250262245A1

(19) **United States**

(12) **Patent Application Publication**  
**TOM et al.**

(10) **Pub. No.: US 2025/0262245 A1**

(43) **Pub. Date: Aug. 21, 2025**

(54) **METHODS OF MANUFACTURE OF  
THERAPEUTIC PRODUCTS COMPRISING  
VITALIZED PLACENTAL DISPERSIONS**

(71) Applicant: **OSIRIS THERAPEUTICS, INC.**,  
Columbia, MD (US)

(72) Inventors: **Samson TOM**, Baltimore, MD (US);  
**Alla DANILKOVITCH**, Columbia,  
MD (US); **Dana YOO**, Falls Church,  
VA (US); **Timothy JANSEN**,  
Baltimore, MD (US); **Jin-Qiang**  
**KUANG**, Woodstock, MD (US);  
**Jennifer Michelle MARCONI**, Glen  
Burnie, MD (US)

(73) Assignee: **OSIRIS THERAPEUTICS, INC.**,  
Columbia, MD (US)

(21) Appl. No.: **19/197,705**

(22) Filed: **May 2, 2025**

**Related U.S. Application Data**

(63) Continuation of application No. 18/101,727, filed on Jan. 26, 2023, now Pat. No. 12,318,408, which is a continuation of application No. 16/841,359, filed on Apr. 6, 2020, now Pat. No. 11,590,173, which is a continuation of application No. 15/949,808, filed on Apr. 10, 2018, now Pat. No. 10,646,519, which is a continuation of application No. 14/070,040, filed on Nov. 1, 2013, now Pat. No. 9,956,248, which is a continuation of application No. 13/030,595, filed on Feb. 18, 2011, now abandoned.

(60) Provisional application No. 61/369,562, filed on Jul. 30, 2010, provisional application No. 61/338,464, filed on Feb. 18, 2010, provisional application No. 61/338,489, filed on Feb. 18, 2010.

**Publication Classification**

(51) **Int. Cl.**

<i>A61K 35/28</i>	(2015.01)
<i>A01N 1/125</i>	(2025.01)
<i>A61K 35/50</i>	(2015.01)
<i>A61K 38/18</i>	(2006.01)
<i>A61K 38/39</i>	(2006.01)
<i>A61K 38/57</i>	(2006.01)
<i>C12N 5/073</i>	(2010.01)

(52) **U.S. Cl.**

CPC ..... *A61K 35/28* (2013.01); *A01N 1/125*  
(2025.01); *A61K 35/50* (2013.01); *A61K*  
*38/1825* (2013.01); *A61K 38/1841* (2013.01);  
*A61K 38/39* (2013.01); *A61K 38/57* (2013.01);  
*C12N 5/0605* (2013.01); *C12N 2500/02*  
(2013.01); *C12N 2501/115* (2013.01); *C12N*  
*2502/025* (2013.01)

(57)

**ABSTRACT**

This invention provides a fluid therapeutic placental product comprising placental cells and a placental dispersion comprising placental factors. The placental cells and the placental dispersion are derived from placental tissue. A placental tissue can optionally be an amnion, chorion, or a trophoblast-depleted chorion. The placental product of the present invention is useful in treating a patient with a tissue injury (e.g. wound or burn) by applying the placental product to the injury. Similar application is useful with ligament and tendon repair and for engraftment procedures such as bone engraftment.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7






FIG. 8




FIG. 9




FIG. 10





FIG. 11

A



B





FIG. 12A and FIG. 12B

A



B




FIG. 13A and FIG. 13B




FIG. 14A and FIG. 14B






FIG. 15A, FIG. 15B, FIG. 15C






FIG. 16




FIG. 17

## METHODS OF MANUFACTURE OF THERAPEUTIC PRODUCTS COMPRISING VITALIZED PLACENTAL DISPERSIONS

### RELATED APPLICATIONS

**[0001]** This application is a continuation of U.S. patent application Ser. No. 18/101,727, filed Jan. 26, 2023, which is a continuation of U.S. patent application Ser. No. 16/841,359 filed Apr. 6, 2020 (now U.S. Pat. No. 11,590,173), which is a continuation of U.S. patent application Ser. No. 15/949,808 filed Apr. 10, 2018 (now U.S. Pat. No. 10,646,519), which is a continuation of U.S. patent application Ser. No. 14/070,040 filed Nov. 1, 2013 (now issued U.S. Pat. No. 9,956,248), which is a continuation of U.S. patent application Ser. No. 13/030,595 filed on Feb. 18, 2011 (now abandoned), which claims priority to: U.S. Provisional Application No. 61/338,464 filed Feb. 18, 2010; U.S. Provisional Application No. 61/338,489 filed Feb. 18, 2010; and Provisional Application No. 61/369,562 filed Jul. 30, 2010. Each application is hereby incorporated herein by reference.

**[0002]** In addition, this application is related to U.S. patent application Ser. No. 13/030,507 filed Feb. 18, 2011 (now abandoned); U.S. patent application Ser. No. 13/030,539 filed Feb. 18, 2011 (now abandoned); U.S. patent application Ser. No. 13/030,551 filed Feb. 18, 2011 (now abandoned); U.S. patent application Ser. No. 13/030,562 filed Feb. 18, 2011 (now abandoned); and U.S. patent application Ser. No. 13/030,580 filed Feb. 18, 2011 (now abandoned). Each application is incorporated herein by reference.

### TECHNICAL FIELD

**[0003]** The present invention relates to placental products, methods of medical treatment using placental products, and methods of making placental products.

### BACKGROUND

**[0004]** The structural integrity of tissue is achieved, in part, by a dynamic interaction of the tissue with bioactive molecules, extracellular matrix, and a host of circulating cell types. Such interactions are also pivotal during tissue aging, injury, restorative and regenerative treatments. For example, burns produce local tissue damage as well as systemic consequences. Currently, treatment of burn wounds is focused on promoting healing and decreasing the risk of infection. Burn wounds continue to be a frustrating and serious problem in the clinic, and these wounds are often accompanied by high morbidity and mortality rates. The standard of care for burns includes the use of antiseptics and gauze wound dressings. However, for severe and large surface area burns, this treatment is not satisfactory. The gold standard for severe burn treatment continues to be autologous living skin grafts. However, the amount of skin available for grafting is often extremely limited, and this procedure always results in donor site wounds.

**[0005]** Attempts to improve burn wound care have included the use of a single growth factor or cocktail of growth factors as well as biological skin substitutes. Growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF), basic fibroblast growth factor (FGF), vascular endothelial growth factor (VEGF), and other singular factors have been tested in burn wound healing; however, with varying results.

**[0006]** The use of placental membranes for burns and other types of wounds originated more than 100 years ago (reviewed by Kesting et al., 2008). Placental membranes contain components that are present in skin and required for wound healing such as extracellular matrix, growth factors, and cells, including MSCs that are responsible for orchestrating the process of wound healing. The effectiveness of placental membranes such as amniotic membranes for burns was recorded in a number of published reports; however, the use of placental membranes for large surface area burns is limited due to challenges in providing sufficient placental membranes to cover large areas.

**[0007]** What is needed in the art is a therapeutic product that provides the benefits of placental membranes yet can be applied in fluid form. Moreover, needed is a product that provides dynamic therapy throughout more than one, optimally all, of the phases of wound repair: inflammatory, proliferative, and remodeling.

### SUMMARY OF THE INVENTION

**[0008]** The present invention provides methods of manufacturing placental products comprising placental cells and a placental dispersion comprising placental factors. The placental cells and the placental dispersion are derived from placental tissue, e.g. a whole placenta or portion thereof. Placental tissue can be obtained by mechanical manipulation (e.g. dissection) or enzymatic digestion or combinations thereof. A placental tissue can optionally be an amnion, chorion, a mixture of amnion and chorion, or other tissue described here.

**[0009]** The present invention also provides a method of treating a tissue injury (e.g. wound or burn) comprising administering to a patient in need thereof a placental product of the present invention.

**[0010]** Optionally, the placental dispersion is a homogenate.

**[0011]** Optionally, placental factors present include extracellular matrix components.

**[0012]** Optionally, the placental dispersion comprises one or more placental factors set forth in Table 1, Table 2, Table 3, or Table 5.

**[0013]** Optionally, the placental cells comprise stromal cells such as MSCs (mesenchymal stem cells) and PSCs (placental stem cells).

**[0014]** In one embodiment, the method of making a placental product is a parallel processing method that comprises: (i) obtaining a first placental (e.g. amniotic or chorionic) tissue; (ii) obtaining placental cells from the first placental tissue; (iii) obtaining a second placental (e.g. amniotic or chorionic) tissue; (iv) disrupting the second placental tissue to form a dispersion comprising placental factors; (v) combining the placental cells and the dispersion to form the placental product.

**[0015]** Optionally, the first placental tissue and the second placental tissue are autologous to each other, for example, derived from the same donor.

**[0016]** In one embodiment, the method of making a placental product is a serial processing method wherein the second placental tissue is derived from the first placental tissue after said step of isolating the placental cells from a first placental tissue. For example, a first chorionic tissue may be retained after isolating a population of cells thereof, and then disrupted to form a dispersion. The dispersion may then be combined with the placental cells.

**[0017]** Optionally, the step of isolating the placental cells comprises contacting the first placental tissue (e.g. amnion or a chorion or a chorion lacking trophoblasts) with a digestive enzyme, such as a collagenase II. Optionally, the first placental tissue is exposed to a limited digestion with an enzyme such as collagenase II; e.g. exposure for less than about 1 hour (e.g. about 10 minutes or about 20 minutes).

**[0018]** Optionally, the placental tissue (from which the placental dispersion is produced) is chorionic tissue depleted of trophoblasts by treatment with a digestive enzyme such as dispase II followed by physical removal.

**[0019]** In another embodiment, the method of making a placental product comprises: (i) obtaining a placental (e.g. amniotic or chorionic) tissue; (ii) exposing the placental tissue to collagenase; (iii) dividing the placental tissue into a first portion and a second portion; (iv) isolating placental cells from the first placental portion; (v) disrupting the second placental portion to form a dispersion comprising placental factors; and (vi) combining the placental cells and the placental dispersion to form the placental product; (vii) in another embodiment, the method of making a placental product comprises: (viii) obtaining a placental (e.g. amniotic or chorionic) tissue; (ix) exposing the placental tissue to a collagenase for a time sufficient to release placental cells; (x) isolating the released placental cells from the collagenase exposed placental tissue; (xi) disrupting the collagenase exposed placental tissue to form a dispersion comprising placental factors; and (xii) combining the placental cells and the placental dispersion to form the placental product.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**[0020]** FIG. 1 depicts cell viability, before and after a freeze-thaw cycle of a placental product comprising isolated cells and a placental dispersion.

**[0021]** FIG. 2 depicts recovery of viable cells isolated by digestion.

**[0022]** FIG. 3 depicts cell viability, before and after a freeze-thaw cycle of a placental product comprising isolated cells and a placental dispersion.

**[0023]** FIG. 4 depicts recovery of viable cells isolated by digestion.

**[0024]** FIG. 5 depicts the level of viable cells in a placental product made with or without a step of cell isolation before tissue disruption.

**[0025]** FIG. 6 depicts cell phenotype of cells in a placental product.

**[0026]** FIG. 7 depicts cell viability using various cryoprotectants

**[0027]** FIG. 8 depicts placental tissue weight and live cells recovered following collagenase treatment of various incubation times.

**[0028]** FIG. 9 depicts the number of collagenase-released cells from multiple donors.

**[0029]** FIG. 10 depicts viable cell level in a placental product when a placenta is subjected to hypoxic or normoxic conditions.

**[0030]** FIG. 11 depicts cell viability when a placenta is subjected to hypoxic or normoxic conditions.

**[0031]** FIGS. 12A and 12B depict expression of bFGF (12A) and VEGF (12B) in placental products for 14 days in culture.

**[0032]** FIGS. 13A and 13B depict expression of IFN-2 $\alpha$  (13A) and TGF- $\beta$ 3 (13B) in placental products,

**[0033]** FIGS. 14A and 14B depict BMP-2, BMP-4, BMP-7, PLAB, and PLGF (14A), and IGF-1 (14B) expression in placental products derived from the chorionic membrane.

**[0034]** FIGS. 15A, 15B, and 15C show representative images of passage 2 cells isolated and expanded from a placental product derived from the chorionic membrane.

**[0035]** FIG. 16 depicts recovery of viable cells isolated by digestion using various collagenase II enzymes.

**[0036]** FIG. 17 depicts cell viability, before and after a freeze-thaw cycle of a placental product comprising isolated cells and a placental dispersion.

#### DETAILED DESCRIPTION OF THE INVENTION

**[0037]** As used here, the following definitions and abbreviations apply.

**[0038]** “Chorionic tissue” or “Chorionic membrane” means the chorion or a portion thereof, e.g. the trophoblast, the somatic mesoderm, or combinations thereof.

**[0039]** “Exemplary” (or “e.g.” or “by example”) means a non-limiting example.

**[0040]** “Placental dispersion” means a product formed by physical/mechanical disruption of placental tissue. For example, a dispersion may be in the form of a homogenate, a blend, a suspension, a colloid, or a solution.

**[0041]** “Placental tissue” means tissue derived from the placenta in the broadest sense of the word. Placental tissue can be a whole placenta or any portion thereof. “Portions of the placenta” is meant to include chorion, amnion, a chorion and amniotic membrane (e.g. amnio-chorion), Wharton’s jelly, umbilical cord, placental cotyledons or combinations thereof. The placental tissue may be dissected or digested (or combinations thereof) to remove portions, membrane, or structures.

**[0042]** “Placental cells” means any cell that can be obtained from a placenta, without regard to genetic origin (e.g. maternal vs. fetal), developmental origin (e.g. endodermal, ectodermal, or mesodermal), or differentiation. Placental cells may comprise any placental cells known in the art, for example, mesenchymal stem cells (MSCs), endometrial stromal cells (ESCs), placenta-derived mesenchymal progenitor cells, placental mesenchymal stem cells, fibroblasts, epithelial cells, placental mesenchymal cells, macrophages, and the like.

**[0043]** “Placental cells” are further meant to require some feature of live cells such as one or more of metabolic activity, structural integrity (e.g. exclusion of a viability stain such as methylene blue), mitotic activity, signal transduction, and the like.

**[0044]** “Placental factor” means any product that is obtainable from a placental tissue (or placental cells). The product can be an angiogenic factor, chemokine, cytokine, growth factor, protease, protease inhibitor, or matrix component. Exemplary placental factors are listed in Table 1, Table 2, Table 3 and Table 5.

**[0045]** “Tissue injury” means an injury of any tissue such as skin or the outer layer of any organ. By injury, it is meant a pathology that involves or results from an mechanical, metabolic or other insult. Examples of such tissue injuries are burns, wounds, ulcerations, and lacerations, ablations (including laser, freezing, cryo-surgery, heat and electrical ablations), and surgical incisions.

## I. PLACENTAL PRODUCT

### A. Overview

**[0046]** It has been surprisingly discovered that a placental product can now be produced by combining placental cells and a placental dispersion to produce a medicinal product of substantial and superior therapeutic value when administered to a tissue injury. The placental product has several advantageous properties.

**[0047]** Fluidity. The placental product shares certain properties of a fluid such as an ability to deform under an applied stress and can be quantified measurements of viscosity. Thus, the present placental product can be spread over the surface of the surface to which it is applied. For example, one ml of placental product can be spread topically to cover more than about any of about 1 cm<sup>2</sup>, about 10 cm<sup>2</sup>, about 25 cm<sup>2</sup>, about 50 cm<sup>2</sup>, or about 100 cm<sup>2</sup> of skin. This fluid property solves the problem of limited applicability of products that retain the non-elastic properties of tissue (e.g. skin grafts). Moreover, the fluidity of the present placental product now makes it practical for new uses such as application to articulating joints and curved surfaces. It also provides a means of rapid application.

**[0048]** Extended release. Extended release formulations, especially for topical pharmaceutical products, are especially problematic. Moreover, due to natural instabilities as well as metabolic degradation, topical formulations often exhibit substantial loss of activity with time after administration. Without being bound by theory, the inventors believe that the placental cells of the present placental products produce placental components after administration. Thus, the present placental products can contain placental components derived from the placental dispersion and derived from the placental cells and depletion of placental components can be reduced. Additionally, placental cells in the present placental product can produce placental factors (e.g. protease inhibitors) that reduce the metabolic degradation of placental factors.

**[0049]** Dynamic responsivity. Without being bound by theory, the inventors believe that presence of live placental cells provide to the placental product the capacity to respond to physiologic stimuli in a manner somewhat analogous to endogenous cells in situ. Evidence of dynamic responsivity includes stimulated release of placental factors or changes in the placental factor profile with time after administration.

### B. Placental Cells

**[0050]** Placental cells may be obtained from any placental tissue (e.g. chorion). Placental cells may be obtained by processing placental tissue in any manner which retains cell viability of at least one cell type (e.g. MSCs). For example, placental cells may be isolated or purified from placental tissue (e.g. by collagenase digestion of the chorion) or may be obtained without isolation from one or more placental factors (e.g. extracellular matrix) or from other placental cells.

**[0051]** Placental cells may be obtained by any method known in the art. Useful methods of obtaining placental cells (e.g. chorionic cells) are described, for example, by Portmann-Lanz et al. ("Placental mesenchymal stem cells as potential autologous graft for pre- and perinatal neuroregeneration," *American Journal of Obstetrics and Gynecology* (2006) 194, 664-73), ("Isolation and characterization of

mesenchymal cells from human fetal membranes," *Journal Of Tissue Engineering And Regenerative Medicine* 2007; 1:296-305.), and (Concise Review: Isolation and Characterization of Cells from Human Term Placenta: Outcome of the First International Workshop on Placenta Derived Stem Cells 77").

**[0052]** In one embodiment, placental cells are obtained by contacting placental tissue with one or more digestive enzymes, for example, by immersing placental tissue (e.g. a chorion, or placental tissue lacking trophoblasts) in a solution containing the digestive enzyme. The digestive enzyme may be any digestive enzyme known in the art. The digestive enzyme may also be combination of enzymes. Exemplary digestive enzymes include one or more: collagenases (e.g., collagenase I, II, III and IV), matrix metalloprotease, neutral proteases, papains, deoxyribonucleases, serine protease (e.g. trypsin, chymotrypsin, elastase), or any combination thereof.

**[0053]** In one embodiment, placental cells are obtained from a chorion by contacting a chorion (e.g. a chorion lacking trophoblasts) with a collagenase (e.g. collagenase II). The collagenase may present in any suitable concentration, for example, about 100 U/mL to about 1000 mL, and in any suitable collagenase solvent, such as DMEM, and at any suitable temperature, for example 37° C. The chorion may be contacted with the digestive enzyme for any suitable period of time. Optionally, the chorion is contacted with a collagenase (e.g. collagenase II) for less than about any of: about 3 hrs, about 2 hr, or about 1 hr. Optionally, the chorion is contacted with the collagenase (e.g. collagenase II) for less than about 1 hour, for example, less than about any of: about 60 min, about 50 min, about 40 min, about 30 min, about 20 min, about 15 min, about 10 min, or about 5 min. Optionally, the chorion is contacted with a collagenase for a limited period of time such that a substantial portion of the placental tissue is retained on a about 100 micron filter. Optionally, the chorion is contacted with collagenase II for a limited period of time such that a substantial portion of the placental tissue is retained on a 100 micron filter. Optionally, after the placental cells are obtained, the chorion is disrupted to form a dispersion and the population is combined with (e.g. added to) the dispersion.

**[0054]** Surprisingly, a step of obtaining placental cells before subjecting the placental tissue to tissue disruption results in substantially a greater number of cells generally and also results in a population of cells that more resemble the population in the placental tissue than population of cells that are obtained from disrupted placental tissue.

**[0055]** A placental product that comprises placental cells from placental tissue that has not been disrupted surprisingly provides a therapeutically effective amount of viable cells without the need for ex vivo expansion of the placental cells. Although ex vivo expansion is a known method of increasing the number of viable cells in a population, such a step often leads to changes in the population make-up or distribution of cell phenotype. For example, various cells in a population may expand at different rates and expansion may also induce differentiation. Accordingly, one embodiment of the present invention provides a placental product comprising placental cells derived from a placental tissue wherein the placental cells exhibits a phenotypic distribution of cells which is substantially similar to the cells of the placental tissue of origin.

### C. Placental Dispersion

**[0056]** A placental dispersion may be provided by disrupting a placenta (e.g. a chorion). The disruption of placental tissue may be accomplished by any physical/mechanical method of disrupting tissue (i.e. use of a “tissue disruptor” or “means for disruption”). For example, disruption may comprise homogenization, maceration, use of a blender, crushing, or mincing. Disruption may additionally or alternatively comprise shearing, mincing, dicing, or chopping. Disruption may additionally or alternatively comprise sonication.

**[0057]** The placental tissue may be disrupted for any suitable duration which produces a dispersion from the placenta. For example, the placenta may be disrupted (e.g. homogenized) for less than about 20 sec, about 15 sec, about 10 sec, or about 5 seconds.

**[0058]** The placental tissue can be disrupted sufficient to form a placental product with fluid characteristic and yet retain viable cells. Accordingly, live cells in the placental products of the present invention can additionally comprise placental cells that are derived from the placental dispersion.

**[0059]** The extent of tissue disruption may be reduced by a prior enzymatic digestion step with a matrix degrading enzyme such as collagenase(s), a protease(s), or combinations thereof. Indeed, it has surprisingly been discovered that such prior digestion preserves viable cells in the placental dispersion. For example, the length of treatment by a tissue disruptor can be reduced by prior enzymatic digestion.

### D. Placental Factors

**[0060]** A placental product of the present invention may comprise one or more placental factors where the placental factors are components of the placental dispersion or components released into the placental product by the placental cells or a combination thereof.

**[0061]** It has surprisingly been discovered that the content of placental factors in placental products made according to the present invention have an unexpected therapeutic value. Such content of placental factors as taught herein is accordingly referred to here as a “therapeutic profile”.

**[0062]** In one embodiment of the present invention, a therapeutic profile is one that provides two or more, or three or more, or four or more placental factors listed in Table 1, Table 2, or Table 5. Optionally, the placental factors are present in an amount of about 20% to about 500% of the mean concentration set forth in Table 1, Table 2, or Table 5. Optionally, the placental factors are present in an amount of about 20% to about 500% of the minimum and the maximum (respectively) of the values set forth in Table 1, Table 2, or Table 5.

**[0063]** Placental factors, according to the present invention, can be placental-derived factors such as angiogenic factors, chemokines, cytokines, growth factors, matrix metalloproteases, extracellular matrix proteins (or “matrix proteins”), and combinations thereof. The present placental products can comprise any of these placental factors.

**[0064]** The present placental products can optionally comprise a therapeutic profile of one or more of a PDGF (e.g. PDGF-bb), EGF, FGF, TGF- $\beta$ 1, TGF- $\beta$ 3, and VEGF and/or one or more of IL-8, IL-6, and MCP-1.

**[0065]** Useful placental products of the present invention can have a therapeutic profile as set forth in Table 1, Table 2, Table 3, or Table 5.

**[0066]** Useful placental products of the present invention can have a therapeutic profile comprising at least 25% of the minimum concentration of one or more placental factors set forth in Table 1 and optionally no more than 400% of the maximum concentration of one or more placental factors set forth in Table 1. In one embodiment, the one or more placental factors comprise fibronectin, TIMP, TGF $\beta$ 1, bFGF, and MMPs (e.g. MMP1,2,4,7,8,9, and 10).

**[0067]** Useful placental products of the present invention can have a therapeutic profile comprising four or more placental factors where at least two placental factors are extracellular matrix components (or fragment thereof).

**[0068]** Placental products of the present invention can comprise a therapeutic profile of one or more placental factors which promote the migration of epithelial cells into a wound area (e.g. HGF and/or KGF), optionally in combination with a growth factor such as TGF- $\beta$ 1. Optionally the concentration of such placental factors is about 25% of the minimum values set forth in Table 1 and optionally no more than 400% of the maximum concentration set forth in Table 1.

**[0069]** Placental products can comprise a therapeutic profile of placental factors that are mitotic or growth promoting. Placental products can contain HGF and KGF. For example, HGF at a concentration of about 5,000 to about 200,000  $\mu$ g/mL and KGF at a concentration of about 5,000 to about 400,000  $\mu$ g/mL are present in an exemplary placental product as detailed in Example 10. Optionally, such placental products are useful in preventing scarring or a useful therapy aid during re-epithelialization.

**[0070]** Placental products of the present invention can comprise a therapeutic profile of placental factors comprising one or more angiogenic factors (e.g. VEGF and/or bFGF) and can optionally additionally comprise one or more growth factors (e.g. TGF- $\beta$ 1 and/or TGF- $\beta$ 2),

**[0071]** Exemplary placental products of the present invention contain a therapeutic profile of VEGF levels greater than about 10  $\mu$ g/ml or greater than about 50  $\mu$ g/ml or greater than about 100  $\mu$ g/ml. For example, an exemplary placental product can comprise greater than about 200  $\mu$ g/ml as detailed in Example 10.

**[0072]** Exemplary placental products of the present invention contain a therapeutic profile of bFGF levels greater than any of about 10 or 100 or 1,000 or 10,000  $\mu$ g/ml. An exemplary placental product can comprise greater than about 11,000  $\mu$ g/mL, as detailed in Example 10. Optionally such FGF-comprising placental products are useful for burn wound healing.

**[0073]** Placental products of the present invention can comprise a therapeutic profile of TGF- $\beta$ 1 and TGF- $\beta$ 2. An exemplary placental product, as detailed in Example 10, comprises bFGF, TGF- $\beta$ 1 and TGF- $\beta$ 2. Optionally, such placental products are useful when the skin pathology being treated involves an inflammatory or a scarring pathology.

**[0074]** Placental products of the present invention may comprise a therapeutic profile of one or more protease inhibitors, such as tissue inhibitors of matrix metalloproteinases (TIMPs), alpha-2 macroglobulin, and/or thrombospondins.

**[0075]** In one embodiment, a placental product (e.g. derived from chorion) comprises one or more protease inhibitors.

**[0076]** In one embodiment, a placental product (e.g. derived from chorion) comprises one or more protease inhibitors and extracellular matrix proteins

**[0077]** In one embodiment, a placental product (e.g. derived from chorion) comprises one or more protease inhibitors and viable cells.

**[0078]** In one embodiment, a placental product (e.g. derived from chorion) comprises one or more protease inhibitors, extracellular matrix proteins, and viable cells.

**[0079]** Without being bound by theory, the present inventors believe that the surprising efficacy that characterizes placental products of the present invention result in an interaction between the placental cells and the placental factors comprising (1) growth factor(s), (2) protease inhibitor(s), and (3) extracellular matrix components. Growth factors can bind to extracellular matrix thereby protecting the growth factors from degradation and effectively extending the half life of the growth factors. Bioavailability can be further regulated by subsequent release or matrix degradation. Similarly, protease inhibitors in exemplary placental products provide additional protection against protease degradation. The placental cells further can protect growth factors and other placental factors in the placental products from degradation by providing additional protease inhibitors and growth factors. Accordingly, such placental products can optionally maintain surprising product integrity for extended periods of time resulting in placental products that require less frequent applications and superior treatment of tissue injuries such as burns and wounds. Surprisingly, the growth factors in such placental products can demonstrate a longer half-life in comparison to other growth factor therapies such as ACCS.

## II. FORMULATION

**[0080]** The placental products of the present invention are administered as a dermatologically acceptable pharmaceutical product. Optionally, active pharmaceutical ingredients or excipients or combinations thereof can be added.

**[0081]** Viscosity. Viscosity values that are useful and desirable according to the present invention vary as a function of the indication being treated. For example, where broad coverage (i.e. large areas of skin) or lower concentrations of placental products are desired, a less viscous formulation is advantageous. Examples of less viscous formulations are those of about 1,000 cps to about 50,000 cps, or about 2,000 cps to about 25,000 cps, or about 2,000 cps to about 10,000 cps, or about 5,000 cps to about 15,000 cps. Such less viscous compositions facilitate spreading of applied composition.

**[0082]** Where more restricted coverage or higher levels of placental products are desired, a more viscous formulation is advantageous. Examples of more viscous formulations are about 20,000 cps to about 200,000 cps or about 50,000 cps to about 100,000 cps.

**[0083]** The skilled artisan will now readily recognize that the desired viscosity can be attained according to the present invention by adjustments of the dispersion method (discussed elsewhere herein) or by selection of a dermatologically acceptable thickening agent and empirically determining the concentration necessary to achieve the desired thickening agent.

**[0084]** The placental products of the present invention can optionally include one or more antibiotic, emollient, keratolytics agent, humectants, anti-oxidants, preservative, or combinations thereof.

**[0085]** In one embodiment, a placental product comprises albumin, such as HSA or BSA. Optionally, the placental product comprises an electrolyte solution, for example, to provide physiological osmolality and pH (e.g. Plasma-LyteA). Optionally, the placental product comprises a cryopreservative, such as DMSO, glycerol, sericin, sugars, or a mixture thereof.

**[0086]** In one embodiment, a placental product comprises albumin, an electrolyte solution, and a cryopreservative. Optionally, the therapeutic product comprises 1% to about 15% albumin by weight and about 5% to about 20% cryopreservative by volume (e.g. about 10%). Optionally, the albumin is HSA, the electrolyte solution is Plasma-Lyte A, and the cryopreservative is DMSO.

## III. MANUFACTURE

### A. Overview

**[0087]** A placental product of the present invention may be manufactured from a placenta in any suitable manner that provides the technical features taught herein. Any placental tissue is useful according to the present invention. Each of the embodiments of the present invention set forth here are meant to specifically embrace placental products where the placental dispersion is a dispersion of chorion that is depleted of or lacking trophoblastic components.

**[0088]** According to the present invention, the placenta is processed to produce the placental dispersion and the placental cells.

**[0089]** In one embodiment, the placental dispersion and the placental cells are derived from a different placenta or different placental portion (e.g. parallel processing).

**[0090]** In one embodiment, the placental dispersion and the placental cells are derived from the same placenta or the same placental portion (e.g. sequential processing).

**[0091]** Manufacturing Method 1. In one embodiment, a placental product is manufactured by: obtaining a placental (e.g. chorionic) tissue; digesting the placental tissue with one or more matrix degrading enzymes (e.g. a collagenase, optionally collagenase II); obtaining placental cells from the digested placental tissue; disrupting the digested placental tissue with a tissue disruptor to form a placental dispersion comprising placental factors; and combining the placental cells and the placental dispersion to form the placental product.

**[0092]** Optional Manufacturing Method 2. In one embodiment, a placental product is manufactured by: obtaining a first placental (e.g. chorionic) tissue; digesting the first placental tissue with one or more matrix degrading enzymes (e.g. a collagenase, optionally collagenase II); obtaining placental cells from the digested first placental tissue; obtaining a second placental tissue; disrupting the second placental tissue with a tissue disruptor to form a placental dispersion comprising placental factors; and combining the placental cells and the placental dispersion to form the placental product.

**[0093]** For either Manufacture Method, the placental tissue can be a chorion tissue such as a chorion tissue that has been processed to reduce the number of trophoblastic cells.

[0094] Exemplary placental products of the present invention can be manufactured or provided with a bandage or wound dressing.

#### B. Trophoblast Removal

[0095] In one embodiment, trophoblasts are depleted or removed to produce the placental tissue from which the placental cells or the placental dispersion or both are derived. Surprisingly, such a placental product has one or more of the following superior features: (a) is substantially non-immunogenic; (b) provides remarkable healing time; and (c) provides enhanced therapeutic efficacy.

[0096] Trophoblasts may be removed in any suitable manner which substantially diminishes the trophoblast content of the placental product. Optionally, the trophoblasts are selectively removed or otherwise removed without eliminating a substantial portion of one or more therapeutic components from the placenta (e.g. MSCs, placental factors, etc.). Optionally, the trophoblasts are removed before isolating a population of cells and/or disrupting the placental tissue.

[0097] One method of removing trophoblasts comprises treating the placenta (e.g. chorion or amnio-chorion) with a digestive enzyme such as dispase (e.g. dispase II) and separating the trophoblasts from the placenta. Optionally, the step of separating comprises mechanical separation such as scraping. Optionally, scraping comprises scraping with a soft instrument such as a finger.

[0098] Useful methods of removing trophoblasts from a placenta (e.g. chorion) are described by Portmann-Lanz et al. ("Placental mesenchymal stem cells as potential autologous graft for pre- and perinatal neuroregeneration"; American Journal of Obstetrics and Gynecology (2006) 194, 664-73), ("Isolation and characterization of mesenchymal cells from human fetal membranes"; Journal Of Tissue Engineering And Regenerative Medicine 2007; 1:296-305.), and (Concise Review: Isolation and Characterization of Cells from Human Term Placenta: Outcome of the First International Workshop on Placenta Derived Stem Cells").

#### C. Preservation

[0099] A placental product of the present invention may be used fresh or may be preserved for a period of time.

[0100] Also as depicted in FIG. 1, a placental product of the present invention, cell viability is retained surprisingly well after a freeze-thaw cycle

[0101] In one embodiment, a placental product is cryopreserved. A placental product may be cryopreserved by freezing (e.g. a  $-80^{\circ}\text{C}.$ ). Freezing may comprise storage in a cryopreservation medium such as DMSO, glycerol, sericin, sugars, or mixtures thereof. Freezing may comprise, for example, incubating the placental product at  $4^{\circ}\text{C}.$  for 30-60 min, and then incubating at  $-80^{\circ}\text{C}.$  until use. The placental product may then be thawed for use.

[0102] A placental product may be formulated in a cryopreservative before cryopreservation. Exemplary cryopreservatives include DMSO, glycerol, and the like. The cryopreservative may further be formulated with additional components such as albumin (e.g. HSA or BSA), an electrolyte solution (e.g. Plasma-Lyte), or a combination thereof. Optionally, the placental product comprises 1% to about 15% albumin by weight and about 5% to about 20% cryopreservative by volume (e.g. about 10%).

[0103] Optionally, a placental product can be formed by the addition of cryopreserved placental cells of the present invention to a fresh (never frozen) placental dispersion or to a frozen placental dispersion or to a lyophilized placental dispersion.

[0104] Optionally, a placental product can be formed by the addition of fresh placental cells of the present invention to a frozen placental dispersion or to a lyophilized placental dispersion.

#### IV. METHODS OF USE

[0105] The placental products of the present invention may be used to treat any tissue injury. A method of treatment may be provided, for example, by administering to a subject in need thereof, a placental product of the present invention.

[0106] A typical administration method of the present invention is topical administration. Administering the present invention can optionally involve administration to an internal tissue where access is gained by a surgical procedure.

[0107] Placental products can be administered autologously, allogeneically or xenogeneically.

[0108] In one embodiment, a present placental product is administered to a subject to treat a wound. Optionally, the wound is a laceration, scrape, thermal or chemical burn, incision, puncture, or wound caused by a projectile. Optionally, the wound is an epidermal wound, skin wound, chronic wound, acute wound, external wound, internal wounds, congenital wound, ulcer, or pressure ulcer. Such wounds may be accidental or deliberate, e.g., wounds caused during or as an adjunct to a surgical procedure. Optionally, the wound is closed surgically prior to administration.

[0109] In one embodiment, the injury is a burn. Optionally, the burn is a first-degree burn, second-degree burn (partial thickness burns), third degree burn (full thickness burns), infection of burn wound, infection of excised and unexcised burn wound, loss of epithelium from a previously grafted or healed burn, or burn wound impetigo.

[0110] In one embodiment, the injury is an ulcer, for example, a diabetic ulcer (e.g. foot ulcer).

[0111] In one embodiment, a placental product is administered by placing the placental product directly over the skin of the subject, e.g., on the stratum corneum, on the site of the wound, so that the wound is covered, for example, using an adhesive tape. Additionally or alternatively, the placental product may be administered as an implant, e.g., as a subcutaneous implant.

[0112] In one embodiment, a placental product is administered to the epidermis to reduce rhytids or other features of aging skin. Such treatment is also usefully combined with so-called cosmetic surgery (e.g. rhinoplasty, rhytidectomy, etc.).

[0113] In one embodiment, a placental product is administered to the epidermis to accelerate healing associated with a dermal ablation procedure or a dermal abrasion procedure (e.g. including laser ablation, thermal ablation, electric ablation, deep dermal ablation, sub-dermal ablation, fractional ablation, and microdermal abrasion).

[0114] Other pathologies that may be treated with placental products of the present invention include traumatic wounds (e.g. civilian and military wounds), surgical scars and wounds, spinal fusions, spinal cord injury, avascular necrosis, reconstructive surgeries, ablations, and ischemia.



**[0115]** In one embodiment, a placental product of the present invention is used in a tissue graft procedure. Optionally, the placental product is applied to a portion of the graft which is then attached to a biological substrate (e.g. to promote healing and/or attachment to the substrate). By way of non-limiting example, tissues such as skin, cartilage, ligament, tendon, periosteum, perichondrium, synovium, fascia, mesenter and sinew can be used as tissue graft.

**[0116]** In one embodiment, a placental product is used in a tendon or ligament surgery to promote healing of a tendon or ligament. Optionally, the placental product is applied to portion of a tendon or ligament which is attached to a bone. The surgery can be any tendon or ligament surgery, including, e.g. knee surgery, shoulder, leg surgery, arm surgery, elbow surgery, finger surgery, hand surgery, wrist surgery, toe surgery, foot surgery, ankle surgery, and the like. For example, the placental product can be applied to a tendon or ligament in a grafting or reconstruction procedure to promote fixation of the tendon or ligament to a bone.

**[0117]** Through the insight of the inventors, it has surprisingly been discovered that placental products of the present invention provide superior treatment (e.g. healing time and/or healing strength) for tendon and ligament surgeries. Tendon and ligament surgeries can involve the fixation of the tendon or ligament to bone. Without being bound by theory, the present inventors believe that osteogenic and/or chondrogenic potential of MSCs in the present placental products promotes healing process and healing strength of tendons or ligaments. The present inventors believe that the present placental products provide an alternative or adjunctive treatment to periosteum-based therapies. For example, useful periosteum based treatments are described in Chen et al. ("Enveloping the tendon graft with periosteum to enhance tendon-bone healing in a bone tunnel: A biomechanical and histologic study in rabbits"; *Arthroscopy*. 2003 March; 19 (3): 290-6), Chen et al. ("Enveloping of periosteum on the hamstring tendon graft in anterior cruciate ligament reconstruction"; *Arthroscopy*. 2002 May-Jun; 18 (5): 27E), Chang et al. ("Rotator cuff repair with periosteum for enhancing tendon-bone healing: a biomechanical and histological study in rabbits"; *Knee Surgery, Sports Traumatology, Arthroscopy* Volume 17, Number 12, 1447-1453), each of which are incorporated by reference.

**[0118]** As non-limiting example of a method of tendon or ligament surgery, a tendon is sutured to and/or wrapped or enveloped in a placental membrane and the tendon is attached to a bone. Optionally, the tendon is placed into a bone tunnel before attached to the bone.

**[0119]** In one embodiment, the tendon or ligament surgery is a graft procedure, wherein the placental product is applied to the graft. Optionally, the graft is an allograft, xenograft, or an autologous graft.

**[0120]** In one embodiment, the tendon or ligament surgery is repair of a torn ligament or tendon, wherein the placental product is applied to the torn ligament or tendon.

**[0121]** Non-limiting examples of tendons to which a placental product can be applied include a digitorum extensor tendon, a hamstring tendon, a bicep tendon, an Achilles Tendon, an extensor tendon, and a rotator cuff tendon.

**[0122]** In one embodiment, a placental product of the present invention is used to reduce fibrosis by applying the placental product to a wound site.

**[0123]** In one embodiment, a placental product of the present invention is used as an anti-adhesion wound barrier,

wherein the placental product is applied to a wound site, for example, to reduce fibrosis (e.g. postoperative fibrosis).

**[0124]** Non-limiting examples of wound sites to which the placental product can be applied include those that are surgically induced or associated with surgery involving the spine, laminectomy, knee, shoulder, or child birth, trauma related wounds or injuries, cardiovascular procedures, angiogenesis stimulation, brain/neurological procedures, burn and wound care, and ophthalmic procedures. For example, optionally, the wound site is associated with surgery of the spine and the stromal side of the placental product is applied to the dura (e.g. the stromal side facing the dura). Direction for such procedures, including the selection of wound sites and/or methodologies, can be found, for example, in WO 2009/132186 and US 2010/0098743, which are hereby incorporated by reference.

**[0125]** A placental product of the present invention can optionally be used to reduce adhesion or fibrosis of a wound. Postoperative fibrosis is a natural consequence of all surgical wound healing. By example, postoperative peridural adhesion results in tethering, traction, and compression of the thecal sac and nerve roots, which cause a recurrence of hyperesthesia that typically manifests a few months after laminectomy surgery. Repeated surgery for removal of scar tissue is associated with poor outcome and increased risk of injury because of the difficulty of identifying neural structures that are surrounded by scar tissue. Therefore, experimental and clinical studies have primarily focused on preventing the adhesion of scar tissue to the dura matter and nerve roots. Spinal adhesions have been implicated as a major contributing factor in failure of spine surgery. Fibrotic scar tissue can cause compression and tethering of nerve roots, which can be associated with recurrent pain and physical impairment.

**[0126]** The placental products disclosed herein are useful in treating a number of wounds including: tendon repair, cartilage repair (e.g. femoral condyle, tibial plateau), ACL replacement at the tunnel/bone interface, dental tissue augmentation, fistulas (e.g. Crohn's disease, G-tube, tracheoesophageal), missing tissue at adhesion barriers (e.g. nasal septum repair, vaginal wall repair, abdominal wall repair, tumor resection), dermal wounds (e.g. partial thickness burns, toxic epidermal necrolysis, epidermolysis bullosa, pyoderma gangrenosum, ulcers e.g. diabetic ulcers (e.g. foot), venous leg ulcers), surgical wounds, hernia repair, tendon repair, bladder repair, periosteum replacement, keloids, organ lacerations, epithelial defects, and repair or replacement of a tympanic membrane.

**[0127]** The presently described technology and its advantages will be better understood by reference to the following examples. These examples are provided to describe specific embodiments of the present technology. By providing these specific examples, it is not intended limit the scope and spirit of the present technology. It will be understood by those skilled in the art that the full scope of the presently described technology encompasses the subject matter defined by the claims appending this specification, and any alterations, modifications, or equivalents of those claims.

**[0128]** The citations provided herein are hereby incorporated by reference for the cited subject matter.

**[0129]** In the present specification, use of the singular includes the plural except where specifically indicated.

## V. EXAMPLES

## A. Example 1. Obtaining Placental Tissue

**[0130]** A whole placenta was obtained from a registered tissue bank after informed consent. The placenta and placed, with the maternal surface (rough surface) face down, on a sterile tray. The amniotic-chorionic membrane was cut and removed from the placenta. The chorionic membrane was then separated from the amnion and washed twice in PBS.

**[0131]** The chorionic membrane was then soaked in an anticoagulant (ACD-A) solution to remove blood clots and then washed again in PBS.

**[0132]** The chorionic membrane was then digested by incubation with dispase II for 30 min. at 37° C. The trophoblast layer was mechanically removed by scraping with fingers and the chorion was washed again in PBS.

**[0133]** The chorionic membrane was then incubated for 24 hours in an antibiotic cocktail containing gentamicin, vancomycin, and amphotericin B, and washed again in PBS.

## B. Example 2. Digesting Placental Tissue

**[0134]** A chorion membrane (obtained from Example 1) was digested by incubation in 200 mL of a collagenase II solution (300 U/mL in DMEM) for 10 min at 37° C. The chorionic membrane was then removed, leaving a digestion suspension containing collagenase and placental cells.

**[0135]** The volume and container for digestion was determined based on the need to provide a suitable digestion environment for the tissue once placed on a shaker. The digestion was carried out on a standard plate shaker set at moderate speed in a 37° C. cell culture incubator.

## C. Example 3. Obtaining Placental Cells

**[0136]** A digestion suspension comprising placental cells (obtained from Example 2) was centrifuged at 2000 rcf for 5 min to separate the digestive enzyme (collagenase II) from the placental cells. This step centrifugation step may enhance cell viability by preventing over-digestion and ensure that the enzyme is washed away before homogenizing the tissue. This centrifugation step pellets the cells without damaging them, allowing the collagenase II to be removed as supernatant.

**[0137]** The cells were then centrifuged again, the supernatant poured off, and the placental cells were resuspended in a small volume (2 mL) of cryoprotectant (5% DMSO in saline). Two mL provides an adequate volume to resuspend the cells while not over-diluting the chorion membrane dispersion once the cells have been added.

## D. Example 4. Obtaining a Placental Dispersion

**[0138]** A chorionic membrane (obtained from Example 2) was washed twice in PBS to remove residual digestion enzyme and placed in a homogenization container with 1 ml cryoprotectant per gram of chorionic membrane. This volume was determined to be appropriate for diluting the chorion membrane enough to produce a dispersion of ideal consistency while maintaining protein concentration at clinically significant levels.

**[0139]** The temperature of the chorionic membrane was reduced by placing the container on ice for greater than 10 min. The chorionic membrane was then homogenized twice at high speed for 5 sec. using a tissue homogenizer to obtain a chorionic dispersion (homogenate).

**[0140]** Once the chorion membrane is subjected to digestion, it becomes easy to homogenize. Surprisingly, only a small amount of homogenization is needed to create a homogenous solution ideal for clinical use and increases the amount of live cells present in the final dispersion.

## E. Example 5. Providing a Placental Product

**[0141]** A placental dispersion (obtained from Example 4) was combined with viable isolated placental cells (obtained from Example 3) and mixed thoroughly to provide a placental product. The placental product may be used (e.g. for therapy) fresh or may first be preserved (e.g. cryogenically) for a period of time.

## F. Example 6. Cryopreservation

**[0142]** A placental product (obtained from Example 5) was aliquoted into vials and incubated at 4° C. for 30-60 min. The vials were then frozen at -80° C. until use.

## G. Example 7. Isolation of Cells without Complete Digestion of Placenta

**[0143]** The inventors tested whether a limited collagenase II digestion might be performed to obtain a suspension containing live cells and yet preserve the integrity of the placental tissue (e.g. preserve placental factors and remaining live cells). A brief 10 minute digestion with collagenase II left the tissue intact and made further handling possible. In addition, a 10 min. collagenase digestion was able to produce high numbers of viable cells

**[0144]** Two placentas were obtained, each from a different donor, and processed according to the procedure detailed in Example 1 through Example 2, except a collagenase II concentration of 244 U/mL, as described above. A cell count was performed immediately following digestion to determine the number of viable cells per gram of tissue that each enzyme was able to digest away off the tissue. The data are presented in FIG. 2.

**[0145]** The placentas were further processed as described in Example 3 through Example 6. Before freezing and after thawing, cells were counted using a hemocytometer and trypan blue staining was used to distinguish live cells. The data are presented in FIG. 3.

**[0146]** Surprisingly, a substantial population of cells was isolated by digestion of less than 1 hr (e.g. 10 min). Digesting the tissue for only 10 min allowed the loosening and removal cells from the tissue without completely breaking up the tissue. In this manner, it was possible to separate the collagenase II/cell mixture from the chorionic membrane. The inventors discovered that 10 min was an adequate amount of digestion time and allowed for variances introduced as a result of donor variability. The digestion process allows isolation of as many live cells as possible while not disrupting the tissue integrity of the chorion membrane to a degree that makes it impossible to manipulate further. The chorion membrane could then be disrupted to produce a placental dispersion that was rich in placental factors while the cells could be isolated from the enzyme solution and then reintroduced to the dispersion to form the placental product.

## H. Example 8. Isolation of Cells without Complete Digestion of Placental Tissue

**[0147]** Multiple placental products were prepared and cell counts were taken immediately following digestion (FIG. 4)

and before freezing and after thawing (FIG. 1), using the procedure described in Example 7. Cells were counted using a hemocytometer and trypan blue staining was used to distinguish live cells. All cell count data was pooled and a mean was calculated.

[0148] As depicted in FIG. 4, digestion of an intact membrane as taught herein produces a surprising number of cells, and does so without mechanical disruption of the membrane. Also depicted in FIG. 4, digestion of a membrane as taught herein produces a surprisingly high ratio of viable to non-viable cells.

[0149] As depicted in FIG. 1, a fresh placental product of the present invention comprises surprisingly high cell viability. Also as depicted in FIG. 1, a placental product of the present invention subjected to a freeze-thaw cycle comprises surprisingly high cell viability. Also as depicted in FIG. 1, a placental product of the present invention, cell viability is retained surprisingly well after a freeze-thaw cycle.

I. Example 9. Isolation of Placental Cells

[0150] Manufacturing methods were explored to obtain superior recovery of live cells in the placental dispersion. Specifically, an experiment was performed to determine the level of viable cells in a placental product manufactured with or without a step of cell isolation before homogenization. Briefly, a placenta prepared according to the procedure detailed in Example 1. The resulting chorion membrane was then divided into equal halves. Half the tissue was processed as described in Example 2 through Example 5 and the other half was processed in the same manner but without cell isolation (collagenase II digestion) prior to homogenization followed by recombining the isolated cells with the dispersion. Cells were counted using a hemocytometer and trypan blue staining was used to distinguish live cells. The data are presented in FIG. 5.

[0151] Results indicate that without prior digestion, homogenization eliminates virtually all viable cells from the

end dispersion. Surprisingly, a placental product contains a substantially greater number of viable cells and is provides enhanced therapeutic efficacy when manufactured with a step of cell isolation before homogenization.

J. Example 10. Profile of a Placental Product

[0152] Multiple placental products were prepared, each from a different donor, according to the procedure detailed in Example 1 through Example 6 and placental factors were analyzed. Briefly, 1 mL of homogenate from each placental product was centrifuged at 14,000 rpm in a microcentrifuge for 10 min.

[0153] The resulting supernatant from each sample was collected as a test sample. Negative control samples consisted of 5% DMSO in saline (cryopreservation solution) and positive control samples consisted of cryopreservation solution with a known concentration of spiked recombinant proteins (bFGF, EGF, and VEGF). Protein profiles comprising placental factors listed in Table 1 were obtained using the SearchLight protein array assay (Aushon Biosystems). Results are indicated in Table 1 as a minimum and maximum expression levels (pg/mL) in a pool of four donors. Since the supernatant is analyzed rather than the complete homogenate, it is likely that protein level estimates are below actual concentrations in each chorion membrane homogenate containing live cells. The levels of VEGF and bFGF in each sample were confirmed by ELISA.

[0154] Surprisingly, many placental factors were detectable at levels that are known to be influential for burn wound healing as well as in the treatment of other indications.

[0155] As seen from the data in Table 1, placental products of the present invention comprise a therapeutic profile of placental factors.

[0156] Table 2 sets forth a therapeutic profile of placental products. Only now, with the teaching herein, the skilled artisan can examine the placental factors, consider the functional role as set forth in Table 3, and assess the value of a placental factor in wound repair.

TABLE 1

Therapeutic Profile of Factors in the Placental Products				
Protein	Min. (pg/mL)	Max. (pg/mL)	Mean (pg/mL)	Function
MMP1	2210.07	3468.94	2808.12	Matrix and growth factor degradation, facilitate cell migration
MMP2	8207.46	70964.65	25648.74	
MMP3	241.76	615.23	454.49	
MMP7	79.78	4429.02	1190.31	
MMP8	778.03	4661.35	2821.20	
MMP9	32879.10	149579.10	71487.03	
MMP10	6728.94	22686.00	14688.40	
MMP13	TLTD	TLTD	TLTD	
TIMP1	18739.41	315870.30	116341.69	Inhibit activity of MMPs, angiogenic
TIMP2	7160.87	60711.15	21335.46	
TSP1	TLTD	TLTD	TLTD	Regulate TGFβ activity, anti-angiogenic
TSP2	1123.02	18784.67	6190.03	
TGFα	TLTD	TLTD	TLTD	Stimulate growth and migration
TGFβ1	1041.50	6572.83	2661.65	
TGFβ2	91.81	1809.81	558.53	Promote angiogenesis, also proliferative and migration stimulatory effects
TGFβ3	77.02	146.31	104.35	Promote angiogenesis, also proliferative and migration stimulatory effects
bFGF (FGF-2)	3554.58	11856.91	7479.40	Inhibit scar formation
				Promote angiogenesis, also proliferative and migration stimulatory effects

TABLE 1-continued

Therapeutic Profile of Factors in the Placental Products				
Protein	Min. (pg/mL)	Max. (pg/mL)	Mean (pg/mL)	Function
KGF (FGF-7)	14.15	111.58	45.72	Stimulate cell growth and migration
EGF	0.42	3.72	1.57	Stimulate cell growth and migration
HB-EGF	TLTD	TLTD	TLTD	
PDGFAA	39.20	173.52	77.46	Promote angiogenesis, also proliferative and migration stimulatory effects
PDGFAB	495.90	495.90	495.90	
PDGFBB	7.73	235.85	70.56	
VEGF	13.95	211.17	76.73	Promote angiogenesis, also proliferative and migration stimulatory effects
VEGFC	64.77	178.51	118.71	
VEGFD	64.73	85.55	77.34	
HGF	9180.77	71280.10	27480.10	Inhibit scar formation, stimulate cell growth and migration
PEDF	805.18	805.18	805.18	Stimulate growth and migration
ANG2	TLTD	TLTD	TLTD	Stimulate growth and migration
IGFBP1	5022.96	1227128.50	322596.69	Regulate IGF and its proliferative effects
IGFBP2	564.62	564.62	564.62	
IGFBP3	226.20	809.16	603.93	
ACRP30	6403.34	33898.70	16229.15	Regulate growth and activity of keratinocytes
Fibronectin	2950999.50	90198200.00	24973399.00	ECM, cellular adhesion, stimulates growth and migration
Alpha2mac	280783.30	4653881.00	1554151.49	Inhibit protease activity, coordinate growth factor bioavailability
IL1ra	961.93	10035.52	3568.27	Anti-inflammatory
NGAL	420.82	2908.38	1592.17	Anti-bacterial
SDF1b	TLTD	TLTD	TLTD	Recruit cells from circulation to site of tissue damage

TLTD = too low to detect

TABLE 2

Therapeutic Profile of Factors in the Chorionic Membrane			
Protein	Min. (pg/mL)	Max. (pg/mL)	Mean (pg/mL)
MMP1	2882.87	6582.26	4732.56
MMP2	748.82	949.52	849.17
MMP3	TLTD	TLTD	TLTD
MMP7	4.46	9.07	6.76
MMP8	TLTD	TLTD	TLTD
MMP9	1259.30	2676.23	1967.77
MMP10	79.31	87.51	83.41
MMP13	TLTD	TLTD	TLTD
TIMP1	17419.86	50712.30	34066.08
TIMP2	640.73	779.98	710.36
TGF $\alpha$	TLTD	TLTD	TLTD
bFGF (FGF-2)	351.28	375.05	363.17
KGF (FGF-7)	1.53	3.07	2.30
EGF	0.75	0.75	0.75
HB-EGF	15.40	84.49	49.94
PDGFAA	35.25	39.79	37.52
PDGFAB	14.03	14.43	14.23
PDGFBB	1.29	3.99	2.64
VEGF	8.39	125.16	66.78

TABLE 2-continued

Therapeutic Profile of Factors in the Chorionic Membrane			
Protein	Min. (pg/mL)	Max. (pg/mL)	Mean (pg/mL)
VEGFC	51.74	123.45	87.60
VEGFD	14.99	20.42	17.70
HGF	29979.57	50392.75	40186.16
PEDF	TLTD	TLTD	TLTD
ANG2	TLTD	TLTD	TLTD
IGFBP1	934.03	1443.63	1188.83
IGFBP2	134.61	135.86	135.24
IGFBP3	4571.51	11970.15	8270.83
LIF	TLTD	TLTD	TLTD
GCSF	0.74	1.22	0.98
TPO	TLTD	TLTD	TLTD
PIGF	TLTD	TLTD	TLTD
ACRP30	225.35	1213.70	719.52
Alpha2mac	8174.44	9968.59	9071.52
IL1ra	525.53	5168.21	2846.87
NGAL	229.72	938.51	584.11
SDF1b	TLTD	TLTD	TLTD

TLTD = too low to detect

TABLE 3

Functions of Placental Factors	
Specific Proteins	Selected Functions
Matrix Metalloproteinase 1 (MMP1), MMP2, 3, 7, 8, 9, 10, 13	Matrix and growth factor degradation, facilitate cell migration
Tissue Inhibitors of MMPs (TIMP1 and TIMP2)	Inhibit activity of MMPs, angiogenic
Angiotensin-2 (Ang-2), Heparin-Bound Epidermal Growth	Stimulate growth and migration

TABLE 3-continued

Functions of Placental Factors	
Specific Proteins	Selected Functions
Factor (HB-EGF), EGF, FGF-7 (also known as Keratinocyte Growth Factor—KGF), Placenta Growth Factor (PIGF), Pigment Epithelium Derived Factor (PEDF), Thrombopoietin (TPO), Transforming Growth Factor- $\alpha$ (TGF- $\alpha$ )	
Basic Fibroblast Growth Factor basic (bFGF), Platelet Derived Growth Factor (PDGF) AA, AB and BB, Vascular Endothelial Growth Factor (VEGF), VEGF-C and VEGF-D	Promote angiogenesis, also proliferative and migration stimulatory effects
TGF- $\beta$ 3, Hepatocyte Growth Factor (HGF)	Inhibit scar formation
$\alpha$ 2-macroglobulin	Inhibit protease activity, coordinate growth factor bioavailability
Adiponectin (Acrp-30)	Regulate growth and activity of keratinocytes
Granulocyte Colony-Stimulating Factor (G-CSF)	Stimulate stem cell migration and proliferation
Interleukin 1 Receptor Antagonist (IL-1RA)	Anti-inflammatory
Neutrophil Gelatinase-Associated Lipocalin (N-GAL)	Anti-bacterial
Leukemia Inhibitory Factor (LIF)	Support of angiogenic growth factors
SDF-1 $\beta$	Recruit cells from circulation to site of tissue damage
Insulin-like Growth Factor Binding Protein (IGFBP1, 2, 3)	Regulate IGF and its proliferative effects

## K. Example 11. Cell Phenotype

**[0157]** FACS was performed to determine cell phenotype in a placental product of the present invention. Placental products were prepared according to the procedure detailed in Example 1 through Example 6. The products were thawed and subsequently filtered through a 100  $\mu$ m filter to remove tissue debris. Single cell suspensions were then centrifuged using a Beckman TJ-6 at 2000 rpm for 10 min and washed twice with DPBS. Supernatant was discarded after each wash, and cells were resuspended in 2 mL of FACS staining buffer (DPBS+0.09% NaN<sub>3</sub>+1% FBS).

**[0158]** Once the single cell suspensions were prepared, a minimum of  $1 \times 10^5$  cells in 100  $\mu$ L of FACS staining buffer was treated with antibodies labeled with fluorescent dye. Table 4 provides descriptions of the antibodies and the amounts used. For cell surface markers, cells were incubated for 30 min at room temperature in the dark with antibodies followed by washing twice with FACS staining buffer by centrifugation at 1300 rpm for 5 min using a Beckman TJ-6 centrifuge. Cells were then resuspended in 400  $\mu$ L of FACS staining buffer and analyzed using a BD FACSCalibur flow cytometer. Results indicate that a placental product derived from chorion contains live cells which stain positive for MSC markers (FIG. 6), implicating the presence of MSC-like cells.

TABLE 4

FACS Antibodies				
Cell marker antibody and label type	Cat No.	Volume of antibody solution used	Cell marker type	Cell marker specificity
IgG1 isotype-PE	BD 559320	5 $\mu$ L	Cell surface	Isotype control
CD105-PE	Caltag MHCD10504	20 $\mu$ L	Cell surface	MSC marker
CD166-PE	BD 559263	80 $\mu$ L	Cell surface	MSC marker

TABLE 4-continued

FACS Antibodies				
Cell marker antibody and label type	Cat No.	Volume of antibody solution used	Cell marker type	Cell marker specificity
CD45-PE	BD 555483	10 $\mu$ L	Cell surface	Hematopoietic cell marker

## L. Example 12. Optimization of Cryoprotectants

**[0159]** A placenta was processed according to the procedure detailed in Example 1 through Example 2. The resulting digestion suspension comprising cells was divided into several aliquots, and each processed according to the procedure detailed in Example 3 through Example 5 using a different cryoprotectant. Three different cryoprotectants were analyzed for their ability to enhance the number of viable cells recovered after freezing and to preserve protein recovery after freezing. The following cryoprotectant solutions were tested: 1. 10% DMSO and 5% HSA in Plasma-Lyte A (CTR solution); 2. 5% DMSO and 5% HSA in Plasma-Lyte A; 3. 10% DMSO in Saline; 4. 5% DMSO in Saline; and 5. 10% Glycerol in Saline

**[0160]** Before freezing and after thawing, cells were counted using a hemocytometer and trypan blue staining was used to distinguish live cells. The following formula was used to calculate the number of cells per mL of homogenate: Cells per mL=(#Cells counted per four 0.0001 mL squares) $\times 10,000 \times$  dilution factor. The results are depicted in FIG. 7.

## M. Example 13. Time Course Optimization of Collagenase Digestion of Chorionic Tissue

**[0161]** To determine the optimal time to digest a placental tissue such as chorionic tissues in collagenase II, chorionic tissues from three different donors were analyzed. The tissues were incubated overnight in an antibiotic cocktail.

Each chorionic membrane tissue was then washed twice to remove antibiotic solution and split into three pieces. Each piece of tissue was weighed to obtain an initial weight (0 min.) before being digested for 10, 20, or 30 minutes in collagenase II solution (300 U/mL).

**[0162]** At the end of each digestion period, the remaining tissue was separated from the collagenase II solution containing the isolated cells by filtering through a 100  $\mu$ m pore cell filter. The separated tissue was then weighed while the collagenase II solution containing digested cells was centrifuged. The resulting cell pellet was resuspended in PBS and counted using a hemocytometer with trypan blue exclusion.

**[0163]** The weight of each remaining tissue piece, including the weight of tissue remaining on the cell filter, was used to calculate the percent of weight lost by digestion with collagenase II.

**[0164]** As shown in FIG. 8, after 10 min. of digestion, about 10% of the original tissue weight was reduced. Further incubation resulted in a more dramatic loss of weight. By 30 minutes, nearly half of the original weight was lost. It was further noted that tissue digested for longer than 10 min. became extremely difficult to separate from the collagenase II solution.

**[0165]** FIG. 8 also shows the number of cells released by collagenase digestion. After 10 minutes of incubation, a substantial number of cells were released. However, by 20 minutes, the number of cells released increased by about 4-fold.

**[0166]** These results surprisingly demonstrate that by performing only a limited collagenase digestion (e.g. about 10 minutes), a substantial number of placental cells can be released and the integrity of the placental tissue is maintained. Accordingly, when the limited collagenase digested placental tissue is subsequently disrupted, the dispersion retains a substantial amount of its native character. For example, the inventors generally observe that after prolonged collagenase digestion (e.g. 30 minutes), the placental tissue can be passed through a 100 micron filter. This is in contrast to the limited digestion where substantially less (e.g. one half or one quarter or less) of the tissue can be passed through a 100 micron filter.

**[0167]** When this dispersion is combined with the released placental cells, a superior therapeutic product is produced.

**[0168]** In data not shown, no significant change in the viability of the collagenase-released cells was observed through 30 min. of digestion.

#### N. Example 14. Time Course Optimization of Collagenase Digestion of Amniotic Tissue

**[0169]** The limited digestion method of Example 13 was tested for applicability when the placental tissue is amniotic tissue. The following procedure was performed: (1) Process placenta; (a) Remove amniotic tissue and wash twice in PBS; (b) Soak amniotic tissue to loosen red blood cells; (i) If needed, clear red blood cells from tissue using fingers; (c) Incubate amniotic tissue for 24 hrs. in antibiotic cocktail; (2) Remove amniotic tissue from antibiotic cocktail and wash twice in PBS; (3) Incubate amniotic tissue for 30 min at 37° C. in 200 mL trypsin solution (0.25%); (4) Remove amniotic tissue from trypsin solution and wash twice in PBS; (5) Incubate amniotic tissue for 10 min at 37° C. in 200 mL collagenase II solution (300 U/mL in DMEM); (6) Remove amniotic tissue from collagenase II solution and wash twice in PBS; (7) Processing of collagenase II and trypsin live cell

suspensions; (a) Centrifuge each suspension at 2000 rcf for 5 min; (b) Pour off each supernatant and replace with 10 mL PBS; (i) Resuspend cells in PBS to wash; (c) Centrifuge cell suspension at 2000 rcf for 5 min; (d) Pour off supernatants and resuspend cells in 2 mL cryoprotectant (5% DMSO in saline); (e) Combine pellets; (8) Processing of amniotic tissue; (a) Place amniotic tissue in homogenization container with a volume of cryoprotectant (mL) equal to the weight of the amniotic membrane (g). For example, if the amniotic membrane weight 25 g place it in the homogenization container with 25 mL of cryoprotectant; (b) Allow the amniotic tissue and cryoprotectant to sit on ice for at least 10 min; (c) Homogenize at high speed twice for 5 sec. using a tissue homogenizer; (9) Combine isolated live cells with homogenate and mix thoroughly (the “placental product”); (10) Aliquot into vials and place at 4° C. for 30-60 min; (11) Freeze at -80° C. until use.

**[0170]** To determine the mean number of live cells in the amnion homogenate, multiple placentas were prepared. Each amnion was processed in one piece, and cell counts were obtained post thaw after cryopreservation (incubation at 4° C. and subsequent freezing at -80° C.). All cell count data were pooled, and a mean was calculated.

**[0171]** Samples from each donor were also prepared for protein array analysis. Briefly, 1 mL of homogenate from each donor was centrifuged at 14,000 rpm in a microcentrifuge for 10 min. The resulting supernatant from each sample was collected. Supernatants along with positive and negative controls were sent to Aushon Biosystems for analysis using their SearchLight protein array assay. This assay measures the levels of 37 proteins of interest in each sample. For this experiment, negative control samples consisted of 5% DMSO in saline (cryopreservation solution), and positive control samples consisted of cryopreservation solution with known concentrations of spiked recombinant proteins (bFGF, EGF, and VEGF).

**[0172]** FACS analysis of single cell suspensions from the placental product was performed for the markers CD45, CD 105, and CD 166.

#### a. Results

**[0173]** As shown in FIG. 9, limited collagenase digestion of amniotic membrane tissue resulted in release of a substantial number of live placental cells.

**[0174]** As shown in Table 5, limited collagenase digestion of amniotic membrane tissue preserved placental factors in the placental dispersion made therefrom.

**[0175]** When Example 13 and Example 14 are considered together, it is now concluded that limited collagenase digestion of placental tissue, whether it be chorion tissue, amniotic tissue, or other tissue of placental origin, results unexpectedly in: Substantial numbers of release live placental cells; Preserved endogenous placental factors; Preserved endogenous placental protein (e.g. matrix proteins); and A therapeutically effective product.

TABLE 5

Therapeutic Profiles of Amnion-Derived Placental Products			
Protein	Min. (pg/mL)	Max. (pg/mL)	Mean (pg/mL)
MMP1	6697.73	10010.27	8354
MMP2	5456.52	53432.45	29444.49
MMP3	570.97	579.1	575.04
MMP7	74.11	207.31	140.71
MMP8	3829.63	3978.42	3904.03
MMP9	11735.19	43661.63	27698.41
MMP10	38916.81	51526.9	45221.86
MMP13	TLTD	TLTD	TLTD
TIMP1	31427.94	78147	54787.47
TIMP2	6149.25	23167.29	14658.27
TSP1	TLTD	TLTD	TLTD
TSP2	7741.98	13312.64	10527.31
TGF $\alpha$	TLTD	TLTD	TLTD
TGF $\beta$ 1	85.17	350.51	217.84
TGF $\beta$ 2	47.98	58.6	53.29
bFGF (FGF-2)	19305.72	23427.48	21366.6
KGF (FGF-7)	70.39	94.29	82.34
EGF	13.71	69.55	41.63
HB-EGF	TLTD	TLTD	TLTD
PDGFAA	14.47	27.93	21.2
PDGFAB	TLTD	TLTD	TLTD
PDGFBB	7.49	12.34	9.91
VEGF	346.3	1058.85	702.57
VEGFC	114.35	220.27	167.31
VEGFD	49.54	75.29	62.42
HGF	12068.53	17408.53	14738.53
PEDF	TLTD	TLTD	TLTD
ANG2	TLTD	TLTD	TLTD
IGFBP1	128.6	159.84	144.22
IGFBP2	TLTD	TLTD	TLTD
IGFBP3	699.01	1349.06	1024.04
ACRP30	6677.35	11232.13	8954.74
Fibronectin	141595.2	254184.05	197889.63
Alpha2mac	421402.95	790851	606126.98
IL1ra	7542.74	10535.55	9039.14
NGAL	1521.63	3283.59	2402.61
SDF1b	TLTD	TLTD	TLTD

TLTD = too low to detect

#### O. Example 15. Live Cells from the Placental Dispersions and the Placental Cell Components of the Placental Product

[0176] The manufacturing steps taught here (e.g. limited collagenase digestion, removal of placental cells before placental tissue disruption, and limited disruption methods) result in a highly effective therapeutic product. The relative role of the placental dispersion and the placental cells components were evaluated for respective role in providing live cells.

[0177] Chorionic tissue was obtained from placental tissue of 9 subjects and the placental cells (e.g. collagenase-released) and placental dispersion was assessed for the number of live cells.

TABLE 6

Placental Cells from Placental Cell and Placental Dispersion Fractions			
Donor	Cells in the Placental cell fraction (collagenase-released)	Cells in the Placental dispersion fraction	Theoretical cells in the placental product
D144	3.84E+05	7.95E+06	8.33E+06
D145	8.40E+05	1.25E+07	1.33E+07

TABLE 6-continued

Placental Cells from Placental Cell and Placental Dispersion Fractions			
Donor	Cells in the Placental cell fraction (collagenase-released)	Cells in the Placental dispersion fraction	Theoretical cells in the placental product
D146	1.60E+05	7.84E+06	8.00E+06
D147	2.17E+07	5.70E+06	2.74E+07
D153	3.26E+06	1.64E+07	1.97E+07
D154	3.70E+05	1.07E+07	1.11E+07
D155	2.08E+06	7.10E+06	9.18E+06
D156	4.90E+05	1.26E+07	1.31E+07
Mean	3.66E+06	1.01E+07	1.38E+07

[0178] As shown in Table 6, 21% to 98% of the cells in the placental products were derived from the placental dispersion component. Thus, the methods of the present invention unexpectedly preserve important placental factors and live cells in the placental dispersion and also provide substantial numbers of live cells from the placental cell (collagenase-released) component.

#### O. Example 16. Hypoxia Treatment

[0179] Results from private studies indicate that hypoxia induces many proteins having beneficial functions in the process of burn wound healing. However, the extent to which hypoxia effects cell growth and protein expression depends on the specific conditions of its application. Therefore, several experiments were performed to determine if hypoxia could enhance the effectiveness of chorion-derived placental products.

[0180] A placenta was processed according to the procedure detailed in Example 1, except the chorionic membrane was divided into two halves before treatment with the antibiotic cocktail. One half of the chorionic membrane tissue was incubated under hypoxic conditions (1% O<sub>2</sub>) while the other was incubated under normal cell culture conditions (~20% O<sub>2</sub>). Each half was then process as described in Example 2 through Example 5. Before freezing and after thawing, cells were counted using a hemocytometer and trypan blue staining was used to distinguish live cells. The results are depicted in FIG. 10.

#### P. Example 16. Hypoxia Treatment and Cryoprotectants

[0181] A placenta was processed according to the procedure detailed in Example 15 except that the digests from each half of the chorionic membrane were further split and formulated with different cryoprotectants, as in Example 12. Before freezing and after thawing, cells were counted using a hemocytometer and trypan blue staining was used to distinguish live cells. The data are presented in FIG. 11. As depicted in FIG. 11, processing under normoxic conditions provides superior cell viability. Also as depicted in FIG. 11, subjecting the chorion to hypoxic conditions may be detrimental to cell viability.

#### Q. Example 18. Growth Factors are Expressed for a Minimum of 14 days

[0182] Placental products of the present invention demonstrate a durable effect, desirable for wound healing treatments. The extracellular matrix and presence of viable cells

within the placental product derived from the chorionic membrane described in this invention allow for a cocktail of proteins that are known to be important for wound healing to be present for at least 14 days.

**[0183]** Placental product derived from the chorionic membrane were thawed and plated onto tissue culture wells and incubated at 37° C. ±2° C. for 3, 7, and 14 days. At each time point, a sample of the product was collected and centrifuged at 16,000 g for 10 min to collect the supernatant. The supernatants were then run on ELISAs for bFGF and VEGF. FIG. 12 illustrates the duration of two key wound healing proteins, bFGF and VEGF, at 3, 7 and 14 days. Although the expression of bFGF goes down with time, it should be noted that significant levels of bFGF was present even out to 14 days. Interestingly, the expression of VEGF increased with time, which could be due to continued active expression of VEGF from the viable cells within the placental product derived from the chorionic membrane.

#### R. Example 19. Interferon 2α (IFN-2α) and Transforming Growth Factor-β3 (TGF-β3)

**[0184]** Interferon-2α and TGF-β3 have been described in the literature as playing critical roles in the prevention of scar and contracture formation (Kwan et al., *Hand Clin*, 2009, 25:511; Tredget et al., *Surg Clin North Am* 1997, 77:701). IFN-2α is known to decrease collagen and fibronectin synthesis and fibroblast-mediated wound contracture. Clinically, IFN-2α has been administered subcutaneously and shown to improve scar quality (Nedelec et al., *Lab Clin Med* 1995, 126:474). TGF-β3 regulates the deposition of extracellular matrix and has been shown to decrease scar formation when injected in rodent cutaneous wound models. Clinically, TGF-β3 has been shown to improve scar appearance when injected at the wound site (Occleston et al., *J Biomater Sci Polym Ed* 2008, 19:1047). Placental product derived from the chorionic membrane described in this invention has been analyzed for the presence of IFN-2α and TGF-β3. Briefly, placental product derived from the chorionic membrane was thawed and centrifuged at 16,000 g to collect supernatants. Supernatants were analyzed on a commercially available ELISA kit from MabTech (IFN-2α) and R&D Systems (TGF-β3). FIG. 13 shows significant expression of IFN-2α and TGF-β3 in placenta products derived from the chorionic membrane.

#### S. Example 20. Tissue Reporative Proteins in Chorionic Membrane Homogenates

**[0185]** Placental product derived from the chorionic membrane was analyzed for the presence of proteins that are important in tissue repair.

**[0186]** Placental products derived from chorionic membranes described in this invention were analyzed for the presence of these tissue reparative proteins. Briefly, placental product derived from the chorionic membrane was incubated at 37° C. ±2° C. for 72 hrs. The product was centrifuged, and the supernatant was analyzed on commercially available ELISA kits from R&D Systems. FIG. 14 shows significant expression of BMP-2, BMP-4, BMP-7, PLAB, PIGF, and IGF-1 in several donors of placental products derived from chorionic membranes.

**[0187]** Without being bound by theory, the inventors believe that efficacy of the present placental products for wound repair are due, in part, to the role of BMPs, IGF-1,

and PIGF in the development and homeostasis of various tissues by regulating key cellular processes. BMP-2 and BMP-4 may stimulate differentiation of MSCs to osteoblasts in addition to promote cell growth; placental BMP or PLAB is a novel member of the BMP family that is suggested to mediate embryonic development. Insulin-like growth factor 1 (IGF-1) may promotes proliferation and differentiation of osteoprogenitor cells. Placental derived growth factor (PIGF) may acts as a mitogen for osteoblasts.

**[0188]** Without being bound by theory, the inventors believe that efficacy of the present placental products for wound repair are due, in part, to the role of BMPs, IGF-1, and PIGF in the development and homeostasis of various tissues by regulating key cellular processes. BMP-2 and BMP-4 may stimulate differentiation of MSCs to osteoblasts in addition to promote cell growth; placental BMP or PLAB is a novel member of the BMP family that is suggested to mediate embryonic development. Insulin-like growth factor 1 (IGF-1) may promotes proliferation and differentiation of osteoprogenitor cells. Placental derived growth factor (PIGF) may acts as a mitogen for osteoblasts.

#### T. Example 1 Differentiation Capacity of Cells Derived from the Chorionic Membrane

**[0189]** Placental cells, in optional embodiments of the present invention, are adherent, express specific cellular markers such as CD105 and lack expression of other markers such as CD45, and demonstrate the ability to differentiate into adipocytes, osteoblasts, and chondroblasts.

**[0190]** The expression of specific cellular markers has already been described in Example 20. To determine if the cells within the placental product derived from the chorionic membrane can adhere to plastic and differentiate into one of the lineages, cells were isolated from the placental product derived from the chorion as described in this invention and cultured at 37° C. ±2° C. and expanded.

**[0191]** FIG. 15-A shows a representative image of passage 2 cells, demonstrating the ability of the cells to adhere to tissue culture plastic. As a comparison, a representative image of MSCs isolated and expanded from human bone marrow aspirate is shown in FIG. 15-B.

**[0192]** Osteogenic differentiation capacity was demonstrated by staining the cultured cells with alkaline phosphatase labeling following the manufacturer's recommendations (BCIP/NBT Alkaline Phosphatase Substrate Kit IV, Vector Laboratories Cat. No. SK-5400). Alkaline phosphatase is an enzyme involved in bone mineralization (Allori et al., *Tissue Engineering: Part B*, 2008, 8:275), and its expression within cells is indicative of osteo-precursor cells (Majors et al., *J Orthopaedic Res*, 1997, 15:546). Staining for alkaline phosphatase is carried out through an enzymatic reaction with Bromo-4-Chloro-3'-Indolylphosphate p-Toluidine Salt (BCIP) and Nitro-Blue Tetrazolium Chloride (NTP). BCIP is hydrolyzed by alkaline phosphatase to form an intermediate that undergoes dimerization to produce an indigo dye. The NBT is reduced to the NBT-formazan by the two reducing equivalents generated by the dimerization. Together these reactions produce an intense, insoluble black-purple precipitate when reacted with alkaline phosphatase.

**[0193]** FIG. 15-C shows a representative image of passage 2 cells isolated and expanded from placental product derived from the chorionic membrane staining positively for alkaline phosphatase.



1. A method of treating a tissue injury in a patient, the method comprising:

obtaining a flowable placental dispersion comprising a placental tissue homogenate; and  
applying the placental dispersion to at least a portion of the tissue injury.

2. The method of claim 1, wherein the placental tissue does not comprise viable cells.

3. The method of claim 1, wherein the placental dispersion comprises one or more placental-derived factors comprising extracellular matrix proteins, angiogenic factors, chemokines, cytokines, growth factors, matrix metalloproteinases, or combinations thereof.

4. The method of claim 1, wherein the viscosity of the placental dispersion is about 1,000 cps to about 50,000 cps or about 20,000 cps to about 200,000 cps.

5. The method of claim 1, wherein the placental dispersion is applied to about 1 cm<sup>2</sup> to about 100 cm<sup>2</sup> of the tissue injury.

6. The method of claim 1, wherein the placental tissue further comprises amnion, chorion, umbilical cord, or combination thereof.

7. The method of claim 1, wherein the placental dispersion is dehydrated.

8. The method of claim 1, wherein the placental dispersion further comprises a thickening agent, an antibiotic, an emollient, a keratolytic agent, a humectant, an anti-oxidant, a preservative, an electrolyte solution, albumin, or combinations thereof.

9. The method of claim 1, wherein the tissue injury is an ablation, laceration, scrape, burn, incision, puncture, epidermal wound, skin wound, chronic wound, acute wound, external wound, internal wound, congenital wound, ulceration, or surgical wound.

10. The method of claim 1, wherein the placental dispersion is comprised in a product that is wrapped around or attached to the tissue injury.

11. The method of claim 1, wherein the placental tissue homogenate is derived from whole placenta.

12. A method of treating a tissue injury in a patient, the method comprising:

obtaining a placental dispersion comprising a placental tissue homogenate, wherein the placental dispersion lacks viable cells and is substantially free of one or more extracellular matrix proteins, growth factors, or cytokines native to the placental tissue; and  
applying the placental dispersion to at least a portion of the tissue injury.

13. The method of claim 12, wherein:

the extracellular matrix proteins comprise fibronectin.  
the growth factors comprise one or more of PDGF, EGF, FGF, TGF- $\beta$ 1, VEGF, HGF, or IGF; and/or

the cytokines comprise one or more of IL-6, IL-1RA, G-CSF, or IL-8.

14. The method of claim 12, wherein the placental dispersion is applied to about 1 cm<sup>2</sup> to about 100 cm<sup>2</sup> of the tissue injury.

15. The method of claim 12, wherein the placental dispersion is dehydrated.

16. The method of claim 12, wherein the placental dispersion further comprises a thickening agent, an antibiotic, an emollient, a keratolytic agent, a humectant, an anti-oxidant, a preservative, an electrolyte solution, albumin, or combinations thereof.

17. The method of claim 12, wherein the placental dispersion further comprises an electrolyte solution.

18. The method of claim 12, wherein the tissue injury is an ablation, laceration, scrape, burn, incision, puncture, epidermal wound, skin wound, chronic wound, acute wound, external wound, internal wound, congenital wound, ulceration, or surgical wound.

19. The method of claim 12, wherein the placental dispersion is comprised in a product that is wrapped around or attached to the tissue injury.

20. The method of claim 12, wherein the placental tissue homogenate is derived from whole placenta.

21. A method of treating a tissue injury in a patient, the method comprising:

obtaining a placental dispersion comprising enzymatically digested and homogenized placental tissue; and  
applying the placental dispersion to at least a portion of the tissue injury.

22. The method of claim 21, wherein the placental dispersion is at a physiological pH.

23. The method of claim 21, wherein the placental dispersion comprises fibronectin.

24. The method of claim 21, wherein the placental dispersion is dehydrated.

25. The method of claim 21, wherein the placental dispersion further comprises a thickening agent, an antibiotic, an emollient, a keratolytic agent, a humectant, an anti-oxidant, a preservative, an electrolyte solution, albumin, or combinations thereof.

26. The method of claim 21, the tissue injury is an ablation, laceration, scrape, burn, incision, puncture, epidermal wound, skin wound, chronic wound, acute wound, external wound, internal wound, congenital wound, ulceration, or surgical wound.

27. The method of claim 21, wherein the placental dispersion is comprised in a product that is wrapped around or attached to the tissue injury.

28. The method of claim 21, wherein the enzymatically digested and homogenized placental tissue is derived from whole placenta.

\* \* \* \* \*