



US012390428B2

(12) **United States Patent**  
**Tabuteau**

(10) **Patent No.:** **US 12,390,428 B2**

(45) **Date of Patent:** **\*Aug. 19, 2025**

(54) **BUPROPION AND DEXTROMETHORPHAN  
FOR REDUCTION OF SUICIDE RISK IN  
DEPRESSION PATIENTS**

(71) Applicant: **ANTECIP BIOVENTURES II LLC,**  
New York, NY (US)

(72) Inventor: **Herriot Tabuteau,** New York, NY (US)

(73) Assignee: **ANTECIP BIOVENTURES II LLC,**  
New York, NY (US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.  
  
This patent is subject to a terminal dis-  
claimer.

(21) Appl. No.: **18/397,220**

(22) Filed: **Dec. 27, 2023**

(65) **Prior Publication Data**

US 2024/0390297 A1 Nov. 28, 2024

**Related U.S. Application Data**

(63) Continuation of application No. 18/323,714, filed on  
May 25, 2023, now Pat. No. 11,896,563, which is a  
continuation of application No.  
PCT/US2021/061492, filed on Dec. 1, 2021.

(60) Provisional application No. 63/120,160, filed on Dec.  
1, 2020, provisional application No. 63/120,672, filed  
on Dec. 2, 2020, provisional application No.  
63/122,902, filed on Dec. 8, 2020.

(51) **Int. Cl.**

**A61K 31/137** (2006.01)

**A61K 31/485** (2006.01)

**A61P 25/24** (2006.01)

(52) **U.S. Cl.**

CPC ..... **A61K 31/137** (2013.01); **A61K 31/485**  
(2013.01); **A61P 25/24** (2018.01)

(58) **Field of Classification Search**

CPC ..... **A61K 31/137**; **A61K 31/485**; **A61P 25/24**  
See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

5,358,970 A 10/1994 Ruff et al.  
5,731,000 A 3/1998 Ruff et al.  
5,763,493 A 6/1998 Ruff et al.  
6,306,436 B1 10/2001 Chungi et al.  
6,780,871 B2 8/2004 Glick et al.  
8,088,786 B2 1/2012 McKinney et al.  
8,569,328 B1 10/2013 Tabuteau  
9,168,234 B2 10/2015 Tabuteau  
9,198,905 B2 12/2015 Tabuteau  
9,205,083 B2 12/2015 Tabuteau  
9,238,032 B2 1/2016 Tabuteau  
9,278,095 B2 3/2016 Tabuteau  
9,314,462 B2 4/2016 Tabuteau  
9,370,513 B2 6/2016 Tabuteau  
9,375,429 B2 6/2016 Tabuteau

9,402,843 B2 8/2016 Tabuteau  
9,402,844 B2 8/2016 Tabuteau  
9,408,815 B2 8/2016 Tabuteau  
9,421,176 B1 8/2016 Tabuteau  
9,457,023 B1 10/2016 Tabuteau  
9,457,025 B2 10/2016 Tabuteau  
9,474,731 B1 10/2016 Tabuteau  
9,486,450 B2 11/2016 Tabuteau  
9,700,528 B2 7/2017 Tabuteau  
9,700,553 B2 7/2017 Tabuteau  
9,707,191 B2 7/2017 Tabuteau  
9,763,932 B2 9/2017 Tabuteau  
9,861,595 B2 1/2018 Tabuteau  
9,867,819 B2 1/2018 Tabuteau  
9,968,568 B2 5/2018 Tabuteau  
10,058,518 B2 8/2018 Tabuteau  
10,064,857 B2 9/2018 Tabuteau  
10,080,727 B2 9/2018 Tabuteau  
10,092,560 B2 10/2018 Tabuteau  
10,092,561 B2 10/2018 Tabuteau  
10,105,327 B2 10/2018 Tabuteau

(Continued)

**FOREIGN PATENT DOCUMENTS**

BR 102016010170 A2 11/2017  
EP 4183391 A1 5/2023

(Continued)

**OTHER PUBLICATIONS**

Spravato (esketamine), Highlights of Prescribing Information, revised  
Jul. 2020.

Nuedexta (dextromethorphan hydrobromide and quinidine sulfate),  
Highlights of Prescribing Information, revised Dec. 2022.

Aplenzin (bupropion hydrobromide), Highlights of Prescribing Infor-  
mation, revised Mar. 2022.

Tod et al., Quantitative Prediction of Cytochrome P450 (CYP)  
2D6-Mediated Drug Interactions, Clinical Pharmacokinetics, 50(8),  
519-530, Aug. 2011.

Kotlyar et al., Inhibition of CYP2D6 Activity by Bupropion, Journal  
of Clinical Psychopharmacology, 25(2), 226-229, Jun. 2005.

Pope et al., Pharmacokinetics of Dextromethorphan after Single or  
Multiple Dosing in Combination with Quinidine in Extensive and  
Poor Metabolizers, The Journal of Clinical Pharmacology, 44(10),  
1132-1142, Oct. 2004.

(Continued)

*Primary Examiner* — Kevin E Weddington

(74) *Attorney, Agent, or Firm* — PROCOPIO, CORY,  
HARGREAVES & SAVITCH LLP; Brent Johnson;  
Yuefen Zhou

(57)

**ABSTRACT**

This disclosure relates to a method of treating depression  
and/or reducing risk of suicide, comprising administering a  
combination of about 90 mg to about 120 mg of bupropion  
hydrochloride, or a molar equivalent amount of another  
form of bupropion, and about 40 mg to about 50 mg of  
dextromethorphan hydrobromide, or a molar equivalent  
amount of another form of dextromethorphan. The combi-  
nation may be administered twice a day to a human being  
suffering from major depressive disorder and having a score  
of 3 or greater on the Suicidality Item of the Montgomery-  
Åsberg Depression Rating Scale (MADRS-SI).

**17 Claims, No Drawings**

(56)

**References Cited**

## U.S. PATENT DOCUMENTS

10,105,361	B2	10/2018	Tabuteau	11,517,542	B2	12/2022	Tabuteau
10,251,879	B2	4/2019	Tabuteau	11,517,543	B2	12/2022	Tabuteau
10,463,634	B2	11/2019	Tabuteau	11,517,544	B2	12/2022	Tabuteau
10,512,643	B2	12/2019	Tabuteau	11,524,007	B2	12/2022	Tabuteau
10,548,857	B2	2/2020	Tabuteau	11,524,008	B2	12/2022	Tabuteau
10,596,167	B2	3/2020	Tabuteau	11,534,414	B2	12/2022	Tabuteau
10,688,066	B2	6/2020	Tabuteau	11,541,021	B2	1/2023	Tabuteau
10,695,304	B2	6/2020	Tabuteau	11,541,048	B2	1/2023	Tabuteau
10,772,850	B2	9/2020	Tabuteau	11,571,399	B2	2/2023	Tabuteau
10,780,064	B2	9/2020	Tabuteau	11,571,417	B2	2/2023	Tabuteau
10,780,066	B2	9/2020	Tabuteau	11,576,877	B2	2/2023	Tabuteau
10,786,469	B2	9/2020	Tabuteau	11,576,909	B2	2/2023	Tabuteau
10,786,496	B2	9/2020	Tabuteau	11,590,124	B2	2/2023	Tabuteau
10,799,497	B2	10/2020	Tabuteau	11,596,627	B2	3/2023	Tabuteau
10,806,710	B2	10/2020	Tabuteau	11,617,728	B2	4/2023	Tabuteau
10,813,924	B2	10/2020	Tabuteau	11,617,747	B2	4/2023	Tabuteau
10,864,209	B2	12/2020	Tabuteau	11,628,149	B2	4/2023	Tabuteau
10,874,663	B2	12/2020	Tabuteau	11,660,273	B2	5/2023	Tabuteau
10,874,664	B2	12/2020	Tabuteau	11,660,274	B2	5/2023	Tabuteau
10,874,665	B2	12/2020	Tabuteau	11,717,518	B1	8/2023	Tabuteau
10,881,624	B2	1/2021	Tabuteau	11,730,706	B1	8/2023	Tabuteau
10,881,657	B2	1/2021	Tabuteau	11,752,144	B1	9/2023	Tabuteau
10,894,046	B2	1/2021	Tabuteau	11,779,579	B2	10/2023	Tabuteau
10,894,047	B2	1/2021	Tabuteau	11,839,612	B1	12/2023	Tabuteau
10,898,453	B2	1/2021	Tabuteau	11,844,797	B1	12/2023	Tabuteau
10,925,842	B2	2/2021	Tabuteau	11,883,373	B1	1/2024	Tabuteau
10,933,034	B2	3/2021	Tabuteau	11,896,563	B2 *	2/2024	Tabuteau ..... A61P 25/24
10,940,124	B2	3/2021	Tabuteau	11,925,636	B2	3/2024	Tabuteau
10,945,973	B2	3/2021	Tabuteau	11,969,421	B2	4/2024	Tabuteau
10,966,941	B2	4/2021	Tabuteau	11,986,444	B2	5/2024	Tabuteau
10,966,942	B2	4/2021	Tabuteau	12,036,191	B1	7/2024	Tabuteau
10,966,974	B2	4/2021	Tabuteau	12,042,473	B2	7/2024	Tabuteau
10,980,800	B2	4/2021	Tabuteau	12,109,178	B2	10/2024	Tabuteau
11,007,189	B2	5/2021	Tabuteau	12,138,260	B2	11/2024	Tabuteau
11,020,389	B2	6/2021	Tabuteau	12,146,889	B1	11/2024	Tabuteau
11,058,648	B2	7/2021	Tabuteau	12,194,005	B2	1/2025	Tabuteau
11,065,248	B2	7/2021	Tabuteau	12,194,006	B2	1/2025	Tabuteau
11,090,300	B2	8/2021	Tabuteau	12,194,036	B2	1/2025	Tabuteau
11,096,937	B2	8/2021	Tabuteau	2002/0035105	A1	3/2002	Caruso
11,123,343	B2	9/2021	Tabuteau	2003/0044462	A1	3/2003	Subramanian et al.
11,123,344	B2	9/2021	Tabuteau	2008/0044462	A1	2/2008	Trumbore et al.
11,129,826	B2	9/2021	Tabuteau	2010/0040679	A1	2/2010	Chang
11,141,388	B2	10/2021	Tabuteau	2010/0291225	A1	11/2010	Fanda et al.
11,141,416	B2	10/2021	Tabuteau	2015/0126541	A1	5/2015	Tabuteau
11,147,808	B2	10/2021	Tabuteau	2015/0126542	A1	5/2015	Tabuteau
11,185,515	B2	11/2021	Tabuteau	2015/0126543	A1	5/2015	Tabuteau
11,191,739	B2	12/2021	Tabuteau	2015/0126544	A1	5/2015	Tabuteau
11,197,839	B2	12/2021	Tabuteau	2015/0133485	A1	5/2015	Tabuteau
11,207,281	B2	12/2021	Tabuteau	2015/0133486	A1	5/2015	Tabuteau
11,213,521	B2	1/2022	Tabuteau	2015/0150830	A1	6/2015	Tabuteau
11,229,640	B2	1/2022	Tabuteau	2015/0157582	A1	6/2015	Tabuteau
11,234,946	B2	2/2022	Tabuteau	2015/0342947	A1	12/2015	Pollard et al.
11,253,491	B2	2/2022	Tabuteau	2016/0008352	A1	1/2016	Tabuteau
11,253,492	B2	2/2022	Tabuteau	2016/0030420	A1	2/2016	Tabuteau
11,273,133	B2	3/2022	Tabuteau	2016/0030421	A1	2/2016	Tabuteau
11,273,134	B2	3/2022	Tabuteau	2016/0128944	A1	5/2016	Chawrai et al.
11,285,118	B2	3/2022	Tabuteau	2016/0128998	A1	5/2016	Tabuteau
11,285,146	B2	3/2022	Tabuteau	2016/0136155	A1	5/2016	Tabuteau
11,291,638	B2	4/2022	Tabuteau	2016/0199321	A1	7/2016	Tabuteau
11,291,665	B2	4/2022	Tabuteau	2016/0228390	A1	8/2016	Tabuteau
11,298,351	B2	4/2022	Tabuteau	2016/0263099	A1	9/2016	Tabuteau
11,298,352	B2	4/2022	Tabuteau	2016/0263100	A1	9/2016	Tabuteau
11,311,534	B2	4/2022	Tabuteau	2016/0317475	A1	11/2016	Tabuteau
11,344,544	B2	5/2022	Tabuteau	2016/0317476	A1	11/2016	Tabuteau
11,357,744	B2	6/2022	Tabuteau	2016/0324807	A1	11/2016	Tabuteau
11,364,233	B2	6/2022	Tabuteau	2016/0339017	A1	11/2016	Tabuteau
11,382,874	B2	7/2022	Tabuteau	2016/0346276	A1	12/2016	Tabuteau
11,419,867	B2	8/2022	Tabuteau	2016/0361305	A1	12/2016	Tabuteau
11,426,370	B2	8/2022	Tabuteau	2016/0375008	A1	12/2016	Tabuteau
11,426,401	B2	8/2022	Tabuteau	2016/0375012	A1	12/2016	Tabuteau
11,433,067	B2	9/2022	Tabuteau	2017/0007558	A1	1/2017	Tabuteau
11,439,636	B1	9/2022	Tabuteau	2017/0014357	A1	1/2017	Tabuteau
11,478,468	B2	10/2022	Tabuteau	2017/0252309	A1	9/2017	Tabuteau
11,497,721	B2	11/2022	Tabuteau	2017/0281617	A1	10/2017	Tabuteau
11,510,918	B2	11/2022	Tabuteau	2017/0304229	A1	10/2017	Tabuteau
				2017/0304230	A1	10/2017	Tabuteau
				2017/0304298	A1	10/2017	Tabuteau
				2017/0354619	A1	12/2017	Tabuteau
				2017/0360773	A1	12/2017	Tabuteau

## (56) References Cited

## U.S. PATENT DOCUMENTS

2017/0360774	A1	12/2017	Tabuteau	2021/0186956	A1	6/2021	Tabuteau
2017/0360776	A1	12/2017	Tabuteau	2021/0196704	A1	7/2021	Tabuteau
2018/0092906	A1	4/2018	Tabuteau	2021/0196705	A1	7/2021	Tabuteau
2018/0116980	A1	5/2018	Tabuteau	2021/0205239	A1	7/2021	Tabuteau
2018/0133195	A1	5/2018	Tabuteau	2021/0205240	A1	7/2021	Tabuteau
2018/0207151	A1	7/2018	Tabuteau	2021/0205297	A1	7/2021	Tabuteau
2018/0256518	A1	9/2018	Tabuteau	2021/0220293	A1	7/2021	Tabuteau
2018/0360823	A1	12/2018	Tabuteau	2021/0220294	A1	7/2021	Tabuteau
2019/0000835	A1	1/2019	Tabuteau	2021/0220348	A1	7/2021	Tabuteau
2019/0008800	A1	1/2019	Tabuteau	2021/0260054	A1	8/2021	Tabuteau
2019/0008801	A1	1/2019	Tabuteau	2021/0267967	A1	9/2021	Tabuteau
2019/0008805	A1	1/2019	Tabuteau	2021/0338605	A1	11/2021	Tabuteau
2019/0015407	A1	1/2019	Tabuteau	2021/0346370	A1	11/2021	Tabuteau
2019/0083426	A1	3/2019	Tabuteau	2021/0361645	A1	11/2021	Tabuteau
2019/0142768	A1	5/2019	Tabuteau	2021/0401828	A1	12/2021	Tabuteau
2019/0192450	A1	6/2019	Tabuteau	2021/0401829	A1	12/2021	Tabuteau
2019/0192507	A1	6/2019	Tabuteau	2021/0401830	A1	12/2021	Tabuteau
2019/0216798	A1	7/2019	Tabuteau	2021/0401831	A1	12/2021	Tabuteau
2019/0216800	A1	7/2019	Tabuteau	2022/0008363	A1	1/2022	Tabuteau
2019/0216801	A1	7/2019	Tabuteau	2022/0071930	A1	3/2022	Tabuteau
2019/0290601	A1	9/2019	Tabuteau	2022/0071931	A1	3/2022	Tabuteau
2020/0022929	A1	1/2020	Tabuteau	2022/0079892	A1	3/2022	Tabuteau
2020/0093762	A1	3/2020	Tabuteau	2022/0096462	A1	3/2022	Tabuteau
2020/0147008	A1	5/2020	Tabuteau	2022/0105086	A1	4/2022	Tabuteau
2020/0147075	A1	5/2020	Tabuteau	2022/0133655	A1	5/2022	Tabuteau
2020/0206217	A1	7/2020	Tabuteau	2022/0142950	A1	5/2022	Tabuteau
2020/0215055	A1	7/2020	Tabuteau	2022/0193012	A1	6/2022	Tabuteau
2020/0215056	A1	7/2020	Tabuteau	2022/0218631	A1	7/2022	Tabuteau
2020/0215057	A1	7/2020	Tabuteau	2022/0218698	A1	7/2022	Tabuteau
2020/0215058	A1	7/2020	Tabuteau	2022/0233470	A1	7/2022	Tabuteau
2020/0215059	A1	7/2020	Tabuteau	2022/0233474	A1	7/2022	Tabuteau
2020/0222389	A1	7/2020	Tabuteau	2022/0233518	A1	7/2022	Tabuteau
2020/0230078	A1	7/2020	Tabuteau	2022/0233519	A1	7/2022	Tabuteau
2020/0230129	A1	7/2020	Tabuteau	2022/0241220	A1	8/2022	Tabuteau
2020/0230130	A1	7/2020	Tabuteau	2022/0241221	A1	8/2022	Tabuteau
2020/0230131	A1	7/2020	Tabuteau	2022/0241269	A1	8/2022	Tabuteau
2020/0237751	A1	7/2020	Tabuteau	2022/0241270	A1	8/2022	Tabuteau
2020/0237752	A1	7/2020	Tabuteau	2022/0265639	A1	8/2022	Tabuteau
2020/0246280	A1	8/2020	Tabuteau	2022/0280504	A1	9/2022	Tabuteau
2020/0261431	A1	8/2020	Tabuteau	2022/0313689	A1	10/2022	Tabuteau
2020/0297666	A1	9/2020	Tabuteau	2022/0323381	A1	10/2022	Tabuteau
2020/0338022	A1	10/2020	Tabuteau	2022/0378779	A1	12/2022	Tabuteau
2020/0360310	A1	11/2020	Tabuteau	2023/0045675	A1	2/2023	Tabuteau
2020/0397723	A1	12/2020	Tabuteau	2023/0096437	A1	3/2023	Tabuteau
2020/0397724	A1	12/2020	Tabuteau	2023/0099206	A1	3/2023	Tabuteau
2020/0405664	A1	12/2020	Tabuteau	2023/0100008	A1	3/2023	Tabuteau
2021/0000763	A1	1/2021	Tabuteau	2023/0100913	A1	3/2023	Tabuteau
2021/0000764	A1	1/2021	Tabuteau	2023/0114111	A1	4/2023	Tabuteau
2021/0000765	A1	1/2021	Tabuteau	2023/0131854	A1	4/2023	Tabuteau
2021/0000768	A1	1/2021	Tabuteau	2023/0142244	A1	5/2023	Tabuteau
2021/0000820	A1	1/2021	Tabuteau	2023/0210843	A1	7/2023	Tabuteau
2021/0015768	A1	1/2021	Tabuteau	2023/0218550	A1	7/2023	Tabuteau
2021/0015814	A1	1/2021	Tabuteau	2023/0225995	A1	7/2023	Tabuteau
2021/0015815	A1	1/2021	Tabuteau	2023/0233491	A1	7/2023	Tabuteau
2021/0023075	A1	1/2021	Tabuteau	2023/0241010	A1	8/2023	Tabuteau
2021/0023076	A1	1/2021	Tabuteau	2023/0248668	A1	8/2023	Tabuteau
2021/0023077	A1	1/2021	Tabuteau	2023/0248669	A1	8/2023	Tabuteau
2021/0030747	A1	2/2021	Tabuteau	2023/0255905	A1	8/2023	Tabuteau
2021/0030749	A1	2/2021	Tabuteau	2023/0263750	A1	8/2023	Tabuteau
2021/0030750	A1	2/2021	Tabuteau	2023/0270740	A1	8/2023	Tabuteau
2021/0030751	A1	2/2021	Tabuteau	2023/0277478	A1	9/2023	Tabuteau
2021/0046067	A1	2/2021	Tabuteau	2023/0277479	A1	9/2023	Tabuteau
2021/0052521	A1	2/2021	Tabuteau	2023/0277480	A1	9/2023	Tabuteau
2021/0060004	A1	3/2021	Tabuteau	2023/0277481	A1	9/2023	Tabuteau
2021/0060005	A1	3/2021	Tabuteau	2023/0293456	A1	9/2023	Tabuteau
2021/0069125	A1	3/2021	Tabuteau	2024/0000770	A1	1/2024	Tabuteau
2021/0069128	A1	3/2021	Tabuteau	2024/0016797	A1	1/2024	Tabuteau
2021/0077428	A1	3/2021	Tabuteau	2024/0024309	A1	1/2024	Tabuteau
2021/0077429	A1	3/2021	Tabuteau	2024/0041862	A1	2/2024	Tabuteau
2021/0077483	A1	3/2021	Tabuteau	2024/0041863	A1	2/2024	Tabuteau
2021/0106546	A1	4/2021	Tabuteau	2024/0050383	A1	2/2024	Tabuteau
2021/0177834	A1	6/2021	Tabuteau	2024/0066025	A1	2/2024	Tabuteau
2021/0186899	A1	6/2021	Tabuteau	2024/0115524	A1	4/2024	Tabuteau
2021/0186900	A1	6/2021	Tabuteau	2024/0148719	A1	5/2024	Tabuteau
2021/0186901	A1	6/2021	Tabuteau	2024/0156751	A1	5/2024	Tabuteau
2021/0186955	A1	6/2021	Tabuteau	2024/0165104	A1	5/2024	Tabuteau
				2024/0189302	A1	6/2024	Tabuteau
				2024/0197656	A1	6/2024	Tabuteau
				2024/0197720	A1	6/2024	Tabuteau

(56)

**References Cited****U.S. PATENT DOCUMENTS**

2024/0238276	A1	7/2024	Tabuteau
2024/0252451	A1	8/2024	Tabuteau
2024/0269130	A1	8/2024	Tabuteau
2024/0299319	A1	9/2024	Tabuteau
2024/0299320	A1	9/2024	Tabuteau
2024/0307408	A1	9/2024	Tabuteau
2024/0390297	A1	11/2024	Tabuteau
2024/0408040	A1	12/2024	Tabuteau
2025/0009754	A1	1/2025	Tabuteau
2025/0012821	A1	1/2025	Tabuteau

**FOREIGN PATENT DOCUMENTS**

KR	101612197	B1	4/2016
WO	1998050044		11/1998
WO	2003086362	A2	10/2003
WO	2004089873	A1	10/2004
WO	2009006194		1/2009
WO	2009050726	A2	4/2009
WO	2015069809	A1	5/2015
WO	2016125108	A1	8/2016
WO	2020146412	A1	7/2020
WO	2021202329	A1	10/2021
WO	2021202419	A1	10/2021
WO	2023004064	A1	1/2023

**OTHER PUBLICATIONS**

Auvelity (dextromethorphan hydrobromide and bupropion hydrochloride), Highlights of Prescribing Information and Medication Guide, issued Dec. 2022.

International Preliminary Report on Patentability, PCT/US2021/061492, mailed on Jun. 15, 2023.

International Search Report and Written Opinion, PCT/US2021/061492 received on Jun. 15, 2023.

International Search Report and Written Opinion, PCT/US2022/012768 received on Jul. 5, 2023.

International Search Report and Written Opinion, PCT/US2023/067062 mailed on Jul. 12, 2023.

Axsome Therapeutics Announces Topline Results of the Stride-1 Phase 3 Trial in Treatment Resistant Depression and Expert Call to Discuss Clinical Implications, Mar. 2020 (retrieved from internet on Jul. 19, 2023). <axsometherapeuticsinc.gcs-web.com/node/9176/pdf>.

Anderson, A.; et al. "Efficacy and Safety of AXS-05, an Oral NMDA Receptor Antagonist with Multimodal Activity, in Major Depressive Disorder: Results of a Phase 2, Double-Blind, Active-Controlled Trial" ASCP Annual Meeting 2019 (retrieved from internet on Jul. 19, 2023). <d3dybxyjb4kyh.cloudfront.net/pdfs/SOBP+2021+AXS-05+MDD+Poster+FINAL.pdf> (May 2019).

O'Gorman, C.; et al. "Rapid Effects of AXS 05, an Oral NMDA Receptor Antagonist, in Major Depressive Disorder: Results from Two Randomized, Double Blind, Controlled Trials" ASCP Annual Meeting 2021 (retrieved from internet on Jul. 19, 2023). <d3dybxyjb4kyh.cloudfront.net/pdfs/SOBP+2021+AXS-05+MDD+Poster+FINAL.pdf> (Jun. 2021).

O'Gorman, C.; et al. "PMH40 Effects of AXS-05 on Patient Reported Depressive Symptoms in Major Depressive Disorder: Results from the Gemini Trial" <doi.org/10.1016/j.jval.2021.04.662> (retrieved from internet on Jul. 19, 2023). Value in Health, Jun. 2021, vol. 24, Supplement 1, pp. S135.

O'Gorman, C.; et al. "P246. Rapid Antidepressant Effects and MADRS Core Symptom Improvements With AXS-05, an Oral NMDA Receptor Antagonist, in Major Depressive Disorder: Results From Two Randomized, Double-Blind, Controlled Trials" ACNP 60th Annual Meeting: Poster Abstracts P246 <nature.com/articles/s41386-021-01236-7> (retrieved from internet on Jul. 19, 2023). Neuropsychopharmacol. 46 (Suppl 1), 72-217, Dec. 2021.

International Preliminary Report on Patentability, PCT/US2022/012768, mailed on Jul. 27, 2023.

Nofziger et al., Evaluation of dextromethorphan with select anti-depressant therapy for the treatment of depression in the acute care psychiatric setting, Mental Health Clinician, 9(2), 76-81, Mar. 2019. Update: Bupropion Hydrochloride Extended-Release 300 mg Bioequivalence Studies, FDA, retrieved Mar. 2021.

FDA Draft Guidance on Bupropion Hydrochloride, revised Mar. 2013.

Forfivo XL (bupropion hydrochloride) extended-release tablets, for oral use, Highlights of Prescribing Information, revised Dec. 2019. Forfivo XL (Bupropion HCl) extended-release tablet, NDA 22497, Jan. 25, 2010.

Wellbutrin XL (bupropion hydrochloride extended-release), Highlights of Prescribing Information, revised Mar. 2022.

Baker T. E. et al., Human Milk and Plasma Pharmacokinetics of Single-Dose Rimegepant 75mg in Healthy Lactating Women, Breastfeeding Medicine, 17(3), 277-282, 2022.

Berle J. O. et al., Antidepressant Use During Breastfeeding, Current Women's Health Reviews, 7(1), 28-34, Feb. 2011.

Briggs G. G. et al., Excretion of bupropion in breast milk, Annals of Pharmacotherapy, 27(4):431-433, Apr. 1993.

Chad L. et al., Update on antidepressant use during breastfeeding, Canadian Family Physician, 59(6), 633-634, Jun. 2013.

Chaudron L. H. et al., Bupropion and Breastfeeding: A case of a possible Infant Seizure, The Journal of clinical psychiatry, 65(6), 881-882, Jun. 2004.

Davis M. F. et al., Bupropion Levels in Breast Milk for 4 Mother-Infant Pairs: More Answers to Lingering Questions, J. Clin. Psychiatry, 70(2), 297-298, Feb. 2009.

Di Scalea T. L. et al., Antidepressant Medication Use during Breastfeeding, Clinical obstetrics and gynecology, 52 (3): 483-497, Sep. 2009.

Dwoskin L. P. et al., Review of the Pharmacology and Clinical Profile of Bupropion, and Antidepressant and Tobacco Use Cessation Agent, CNS Drug Reviews, 12(3-4), 178-207, Sep. 2006.

Gentile S., The safety of newer antidepressants in pregnancy and breastfeeding, Drug Safety, 28(2), 137-152, Feb. 2005. [doi: 10.2165/00002018-200528020-00005. PMID: 15691224].

Haas J. S. et al., Bupropion in breast milk: an exposure assessment for potential treatment to prevent post-partum tobacco use, Tobacco Control, 13(1), 52-56, Mar. 2004.

Ram D. et al., Antidepressants, anxiolytics, and hypnotics in pregnancy and lactation, Indian J Psychiatry, 57(Suppl 2): S354-S371, Jul. 2015. [doi:10.4103/0019-5545.161504].

Weissman A. M. et al., Pooled Analysis of Antidepressant Levels in Lactating Mothers, Breast Milk, and Nursing Infants, Am J Psychiatry, 161(6), 1066-1078, Jun. 2004.

Horn J. R. et al., Get to Know an Enzyme: CYP2D6, Pharmacy Times, Jul. 2008, retrieved on Aug. 28, 2023.

International Search Report and Written Opinion, PCT/US2023/069286 mailed on Aug. 22, 2023.

International Search Report and Written Opinion, PCT/US2023/069239 mailed on Aug. 28, 2023.

International Search Report and Written Opinion, PCT/US2023/069367 mailed on Aug. 28, 2023.

International Search Report and Written Opinion, PCT/US2023/069655 mailed on Sep. 15, 2023.

International Search Report and Written Opinion, PCT/US2023/069371 mailed on Sep. 26, 2023.

International Search Report and Written Opinion, PCT/US2022/037913 mailed on Sep. 21, 2022.

International Search Report and Written Opinion, PCT/US2022/074713 mailed on Sep. 21, 2022.

International Preliminary Report on Patentability, PCT/US2022/037913, issued on Jan. 18, 2024.

International Preliminary Report on Patentability, PCT/US2022/074713, issued on Feb. 22, 2024.

Chinese Pat. No. 202080004041.1 Invalidation Notice and Request issued on Jan. 15, 2024. (English translation included).

Ward K. and Citrome L.: "AXS-05: an investigational treatment for Alzheimer's disease-associated agitation", Expert Opinion on Investigational Drugs, Jul. 6, 2022, vol. 31, issue 8, pp. 773-780, DOI: 10.1080/13543784.2022.2096006.

(56)

**References Cited**

## OTHER PUBLICATIONS

Defendant Teva Pharmaceuticals, Inc.'s Invalidity Contentions for U.S. Pat. Nos. 11,752,144, 11,717,518, 11,730,706 and Exhibits A-C dated Apr. 11, 2024.

Goodnick, Psychotropic drugs and the ECG: focus on the QTc interval, Expert Opinion on Pharmacotherapy, vol. 3, No. 5, p. 479-498, 2002.

International Search Report and Written Opinion, PCT/US2024/046359 mailed on Nov. 28, 2024.

Tabuteau H, et al. "Effect of AXS-05 (Dextromethorphan-Bupropion) in Major Depressive Disorder: A Randomized Double-Blind Controlled Trial" Am J Psychiatry (2022) vol. 179 pp. 490-499. doi: 10.1176/appi.ajp.21080800.

International Preliminary Report on Patentability, PCT/US2023/067062, mailed on Nov. 28, 2024.

International Search Report and Written Opinion, PCT/US2024/043903 mailed on Nov. 28, 2024.

International Preliminary Report on Patentability, PCT/US2023/069286, mailed on Jan. 9, 2025.

International Preliminary Report on Patentability, PCT/US2023/069371, mailed on Jan. 9, 2025.

International Preliminary Report on Patentability, PCT/US2023/069655, mailed on Jan. 16, 2025.

\* cited by examiner

# **BUPROPION AND DEXTROMETHORPHAN FOR REDUCTION OF SUICIDE RISK IN DEPRESSION PATIENTS**

## **CROSS-REFERENCE TO RELATED APPLICATIONS**

This application is a continuation of U.S. patent application Ser. No. 18/323,714, filed May 25, 2023; which is a continuation of International Pat. App. No. PCT/US2021/061492, filed Dec. 1, 2021; which claims the benefit of U.S. Provisional App. Nos. 63/120,160, filed Dec. 1, 2020, 63/120,672, filed Dec. 2, 2020, and 63/122,902, filed Dec. 8, 2020; all of which are incorporated by reference herein in their entireties.

## **SUMMARY**

This disclosure relates to a method of treating depression and/or reducing risk of suicide, comprising administering combination of about 90 mg to about 120 mg of bupropion hydrochloride, or a molar equivalent amount of another form of bupropion, and about 40 mg to about 50 mg of dextromethorphan hydrobromide, or a molar equivalent amount of another form of dextromethorphan. The combination may be administered twice a day to a human being suffering from major depressive disorder and having a score of 3 or greater on the Suicidality Item of the Montgomery-Åsberg Depression Rating Scale (MADRS-SI).

## **DETAILED DESCRIPTION**

The combination of dextromethorphan and bupropion may be used to treat depression, such as major depressive disorder and/or reduce the risk of suicide. Patients being treated by this combination may suffer from depression and may have a score of 2 or greater, or 3 or greater on the Suicidality Item of the Montgomery-Åsberg Depression Rating Scale (MADRS-SI).

Dextromethorphan is rapidly metabolized in the human liver. This rapid hepatic metabolism may limit systemic drug exposure in individuals who are extensive metabolizers. Human beings can be: 1) extensive metabolizers of dextromethorphan—those who rapidly metabolize dextromethorphan; 2) poor metabolizers of dextromethorphan—those who only poorly metabolize dextromethorphan; or 3) intermediate metabolizers of dextromethorphan—those whose metabolism of dextromethorphan is somewhere between that of an extensive metabolizer and a poor metabolizer. Extensive metabolizers can also be ultra-rapid metabolizers. Non-poor metabolizers of dextromethorphan include extensive metabolizers of dextromethorphan and intermediate metabolizers of dextromethorphan. Extensive metabolizers of dextromethorphan are a significant portion of the human population. Dextromethorphan can, for example, be metabolized to dextrophan.

When given the same oral dose of dextromethorphan, plasma levels of dextromethorphan are significantly higher in poor metabolizers or intermediate metabolizers as compared to extensive metabolizers of dextromethorphan. The low plasma concentrations of dextromethorphan can limit its clinical utility as a single agent for extensive metabolizers, and possibly intermediate metabolizers, of dextromethorphan. Bupropion inhibits the metabolism of dextromethorphan, and raises the plasma concentration of dextromethorphan, and can thus improve its therapeutic efficacy.

Similarly, bupropion may allow dextromethorphan to be given less often or in a lower amount, such as once a day instead of twice a day, once a day instead of three times a day, once a day instead of four times a day, twice a day instead of three times a day, or twice a day instead of four times a day, without loss of therapeutic efficacy.

The MADRS is a clinician-rated scale. The MADRS is used to assess depressive symptomatology during the previous week. Subjects are rated on 10 items to assess symptoms: 1) apparent sadness, 2) reported sadness, 3) inner tension, 4) reduced sleep, 5) reduced appetite, 6) concentration difficulties, 7) lassitude, 8) inability to feel, 9) pessimistic thoughts, 10) suicidal thoughts. Each item yields a score of 0 to 6.

The overall score ranges from 0 to 60. A score of 0 indicates the absence of symptoms, and a score of 60 indicates symptoms of maximum severity. A total score ranging from 0 to 6 indicates that the patient is in the normal range (no depression), a score ranging from 7 to 19 indicates “mild depression,” 20 to 34 indicates “moderate depression,” a score of 35 and greater indicates “severe depression.”

In some embodiments, the human being that is treated with a combination of dextromethorphan and bupropion, e.g. for a type of depression, has, or is selected for having, a MADRS score that is at least about 20, at least about 25, at least about 30, at least about 35, at least about 40, at least about 45, at least about 50, at least about 55, about 20-25, about 25-30, about 30-35, about 35-40, about 40-45, about 45-50, about 50-55, about 55-60, about 25-35, about 35-45, about 45-60, about 25-40, or about 40-60.

In some embodiments, treatment with the combination of dextromethorphan and bupropion results in the human being having a MADRS score that is reduced by at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, about 10-20%, about 20-30%, about 30-40%, about 40-50%, about 50-60%, about 60-80%, about 80-90%, or about 90-100% as compared to baseline or placebo. In some embodiments, the reduction is compared to baseline. In some embodiments, the reduction is compared to the baseline before treatment.

In some embodiments, treatment with the combination of dextromethorphan and bupropion results in the human being having a MADRS score that is less than 34, about 20-34, about 7-19, about 0-6, about 30 or less, about 26 or less, about 25 or less, about 20 or less, about 19 or less, about 17 or less, about 14 or less, about 12 or less, about 10 or less, about 8 or less, about 6 or less, about 5 or less, about 4 or less, about 3 or less, about 2 or less, about 1 or less, about 0, about 7 or less, about 0-6, about 1-2, about 2-3, about 3-4, about 4-5, about 5-6, about 6-7, about 7-8, about 8-9, about 9-10, about 10-11, about 11-12, about 12-13, about 13-14, about 14-15, about 15-16, about 16-17, about 17-18, about 18-19, about 19-20, about 18-20, about 1-3, about 3-6, about 6-9, about 9-12, about 12-14, about 12-15, about 15-19, or about 15-20.

Depression may be manifested by depressive symptoms. These symptoms may include psychological changes such as changes in mood, feelings of intense sadness, despair, mental slowing, loss of concentration, pessimistic worry, agitation, anxiety, irritability, guilt, anger, feelings of worthlessness, reckless behavior, suicidal thoughts or attempts, and/or self-deprecation. Physical symptoms of depression may include insomnia, anorexia, appetite loss, weight loss, weight gain, decreased energy and libido, fatigue, restless-

ness, aches, pains, headaches, cramps, digestive issues, and/or abnormal hormonal circadian rhythms.

Some patients, even after treatment with medications such as antidepressants, may have an inadequate or no response to the treatment. Treatment resistant depression (TRD), or treatment-refractory depression, is a condition generally associated with patients who have failed treatment with at least two antidepressants. Part of the diagnosis for TRD is for the patient to have had an inadequate response to treatment with the antidepressants after an adequate dose and adequate course, e.g. in the current depressive episode. TRD may be more difficult to treat due to the comorbidity of other medical or psychological illnesses, such as drug/alcohol abuse or eating disorders, or TRD being misdiagnosed. Some TRD patients have had an inadequate response to 1, 2, 3, or more adequate antidepressant treatment trials or have failed or had an inadequate response to 1, 2, 3, or more prior antidepressant treatments. In some embodiments, a patient being treated for treatment resistant depression has failed treatment with at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more antidepressant therapies. In addition to other depressed patients, the combination of bupropion and dextromethorphan may be effective in treating patients suffering from treatment resistant depression with suicidal ideation.

Patients who may benefit from the treatments described herein include pediatric patients, such as patients under about 18 years of age, about 0-5 years of age, about 5-10 years of age, about 10-12 years of age, or about 12-18 years of age; adult patients, such as patients having an age of about 18-70 years, about 18-65 years, about 18-30 years, about 18-20 years, about 20-30 years, about 30-40 years, about 40-50 years, about 50-60 years, about 60-70 years, about 70-80 years, about 80-90 years, about 30-50 years, about 50-65 years; elderly patients, such as patients over 65 years of age, about 65-75 years of age, about 75-90 years of age, or over 90 years of age; and about 41 years of age or older.

In some embodiments, the human being that is treated with a combination of dextromethorphan and bupropion, e.g. for a type of depression, is, or is selected for being, of Asian descent. In some embodiments, the human being that is treated with a combination of dextromethorphan and bupropion, e.g. for a type of depression, is, or is selected for being, of Japanese descent. In some embodiments, the human being that is treated with a combination of dextromethorphan and bupropion, e.g. for a type of depression, is, or is selected for being, of Korean descent. In some embodiments, the human being that is treated with a combination of dextromethorphan and bupropion, e.g. for a type of depression, is, or is selected for being, of Chinese descent. The assignment of an individual as having Asian, Chinese, Japanese, or Korean descent may be based upon self-reporting by the individual. In these Asian individuals, the combination of dextromethorphan and bupropion may be effective for treating depression where bupropion alone is not effective for treating depression. This may be of particular importance because patients of Asian descent may suffer from more severe depression than those of other ethnic or cultural groups.

In some embodiments, the human being that is treated with a combination of dextromethorphan and bupropion, e.g. for a type of depression, is suffering from, or is selected for suffering from, a major depressive episode that has lasted between about 8 weeks and about 24 months, about 1-6 months, about 6-12 months, about 1-2 years, at least about 1 week, at least about 2 weeks, at least about 3 weeks, at least about 4 weeks, at least about 6 weeks, at least 7 weeks, at least about 2 months, at least about 3 months, at least

about 4 months, at least about 5 months, at least about 6 months, at least about 9 months, at least about 1 year, at least about 18 months, at least about 2 years, about 1-12 weeks, about 3-6 months, about 6-9 months, about 9-12 months, about 12-18 months, about 18-24 months, about 2-4 years, about 4-6 years, about 6-10 years, about 10-20 years or longer.

In some embodiments, the human being that is treated with a combination of dextromethorphan and bupropion, e.g. for a type of depression, has, or is selected having, about 1-100, or more, lifetime depressive episodes, such as a major depressive episodes, including at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 10, at least 15, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90, at least 100, 1-5, 5-10, 10-20, 20-30, 30-40, 40-50, 50-60, 60-70, 70-80, 80-90, 90-100, or 4-7 lifetime depressive episodes.

In some embodiments, the human being that is treated with a combination of dextromethorphan and bupropion, e.g. for a type of depression, has, or is selected for having, an inadequate response to one or more prior antidepressant therapies, e.g. 1, 2, 3, 4, 5 or more prior antidepressant therapies, including prior antidepressant therapies in the current depressive episode (e.g. the current major depressive episode).

In some embodiments, the human being that is treated with a combination of dextromethorphan and bupropion, e.g. for a type of depression, is, or is selected for being male. In some embodiments, the human being that is treated with a combination of dextromethorphan and bupropion, e.g. for a type of depression, is, or is selected for being female.

The combination of bupropion and dextromethorphan is administered once a day or twice a day for at least one week, at least two weeks, at least three weeks, at least four weeks, at least six weeks, at least eight weeks, at least three months, at least four months, at least five months, or at least six months; and/or may be administered for up to four months, up to six months, up to one year, up to two years, or longer. In some embodiments, the combination of bupropion and dextromethorphan is administered twice a day for at least one week, at least two weeks, at least three weeks, at least four weeks, at least six weeks, at least eight weeks, at least three months, at least four months, at least five months, or at least six months; and/or may be administered for up to four months, up to six months, up to one year, up to two years, or longer. In some embodiments, the combination of bupropion and dextromethorphan is administered once a day for 1, 2, 3, 4, 5, 6, or 7 days, and then administered twice a day thereafter.

In some embodiments, at least about 50 mg, at least about 70 mg, at least about 90 mg, at least about 100 mg, at least about 110 mg, or at least about 120 mg of bupropion hydrochloride, or a molar equivalent amount of another form of bupropion (such as another salt form or the free base form) is administered once a day or twice a day. In some embodiments, the bupropion is administered once a day for 1, 2, 3, 4, 5, 6, or 7 days, and then administered twice a day thereafter.

In some embodiments, up to about 70 mg, up to about 90 mg, up to about 100 mg, up to about 110 mg, up to about 120 mg up to about 130 mg, or up to about 150 mg, of bupropion hydrochloride, or a molar equivalent amount of another form of bupropion (such as another salt form or the free base form) is administered once a day or twice a day. In some embodiments, the bupropion is administered once a day for 1, 2, 3, 4, 5, 6, or 7 days, and then administered twice a day thereafter.

In some embodiments, about  $0.8 \times$  mg to about  $1.2 \times$  mg of bupropion hydrochloride, or a molar equivalent amount of another form of bupropion (such as another salt form or the free base form) is administered once or twice a day, wherein x is about 50 mg (e.g. 40-60 mg), about 60 mg, about 70 mg, about 80 mg, about 90 mg, about 100 mg, about 120 mg, or about 140 mg. In some embodiments, the bupropion is administered once a day for 1, 2, 3, 4, 5, 6, or 7 days, and then administered twice a day thereafter.

In some embodiments, about 50-150 mg, about 90-120 mg, about 100-110 mg, or about 105 mg of bupropion hydrochloride, or a molar equivalent amount of another form of bupropion (such as another salt form or the free base form) is administered once a day or twice a day. In some embodiments, the bupropion is administered once a day for 1, 2, 3, 4, 5, 6, or 7 days, and then administered twice a day thereafter.

In some embodiments, at least about 30 mg, at least about 35 mg, at least about 40 mg, or about 45 mg of dextromethorphan hydrobromide, or a molar equivalent amount of another form of dextromethorphan (such as another salt form or the free base form) is administered once a day or twice a day. In some embodiments, the dextromethorphan is administered once a day for 1, 2, 3, 4, 5, 6, or 7 days, and then administered twice a day thereafter.

In some embodiments, up to about 50 mg, up to about 55 mg, up to about 60 mg, or up to about 45 mg of dextromethorphan hydrobromide, or a molar equivalent amount of another form of dextromethorphan (such as another salt form or the free base form) is administered once a day or twice a day. In some embodiments, the dextromethorphan is administered once a day for 1, 2, 3, 4, 5, 6, or 7 days, and then administered twice a day thereafter.

In some embodiments, about  $0.8 \times$  mg to about  $1.2 \times$  mg of dextromethorphan hydrobromide, or a molar equivalent amount of another form of dextromethorphan (such as another salt form or the free base form) is administered once a day or twice a day, wherein x is about 30 mg (e.g. 24-36 mg), about 40 mg, about 50 mg, or about 60 mg. In some embodiments, the dextromethorphan is administered once a day for 1, 2, 3, 4, 5, 6, or 7 days, and then administered twice a day thereafter.

In some embodiments, about 30-60 mg, about 40-50 mg, about 44-46 mg, or about 45 mg of dextromethorphan hydrobromide, or a molar equivalent amount of another form of dextromethorphan (such as another salt form or the free base form) is administered once a day or twice a day. In some embodiments, the dextromethorphan is administered once a day for 1, 2, 3, 4, 5, 6, or 7 days, and then administered twice a day thereafter.

Administration of the combination of bupropion and dextromethorphan may improve depression symptoms, such as the Montgomery-Åsberg Depression Rating Scale (MADRS) score. It may also reduce the Suicidality Item of the MADRS (MADRS-SI), such as by at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 80%, or up to about 100%, e.g. after the combination is administered daily, such as twice a day, for one week, two weeks, four weeks, six weeks, eight weeks, or twelve weeks, etc.

In some embodiments, after eight days of receiving the combination of bupropion and dextromethorphan twice a day, the  $C_{max}$  of dextromethorphan in the human being is increased about 15- to about 25-fold as compared to administration of 60 mg of dextromethorphan hydrobromide with-

out bupropion or as compared to administration of a single dose of the combination of bupropion and dextromethorphan.

In some embodiments, after eight days of receiving the combination of bupropion and dextromethorphan twice a day, the  $C_{min}$  of dextromethorphan in the human being is increased about 35- to about 45-fold as compared to administration of 60 mg of dextromethorphan hydrobromide without bupropion or as compared to administration of a single dose of the combination of bupropion and dextromethorphan.

In some embodiments, after eight days of receiving the combination of bupropion and dextromethorphan twice a day, the  $AUC_{0-12}$  of dextromethorphan in the human being is increased about 25- to about 45-fold as compared to administration of 60 mg of dextromethorphan hydrobromide without bupropion or as compared to administration of a single dose of the combination of bupropion and dextromethorphan.

In some embodiments, after eight days of receiving the combination of bupropion and dextromethorphan twice a day, the  $C_{max}$  of dextromethorphan in the human being is about 75-80 ng/mL.

In some embodiments, after eight days of receiving the combination of bupropion and dextromethorphan twice a day, the  $C_{min}$  of dextromethorphan is about 42-50 ng/mL.

In some embodiments, after eight days of receiving the combination of bupropion and dextromethorphan twice a day, the  $AUC_{0-12}$  of dextromethorphan in the human being is about 755 ng·hr/mL.

In some embodiments, after eight days of receiving the combination of bupropion and dextromethorphan twice a day, the  $C_{max}$  of bupropion in the human being is about 85-90 ng/mL.

In some embodiments, after eight days of receiving the combination of bupropion and dextromethorphan twice a day, the  $C_{min}$  of bupropion is about 25-35 ng/mL.

In some embodiments, after eight days of receiving the combination of bupropion and dextromethorphan twice a day, the  $AUC_{0-12}$  of bupropion in the human being is about 660-670 ng·hr/mL.

#### Example 1

An open label Phase III clinical trial was conducted using a tablet containing 45 mg dextromethorphan hydrobromide and 105 mg of bupropion hydrochloride with modulated delivery. This tablet was administered twice a day to patients suffering from major depressive disorder. A total of 611 patients participated in the trial, and a total of 597 patients were treated for at least 6 months when the trial was concluded. The mean MADRS score of the total patient population was 32.7 prior to treatment. The results for the total patient population are summarized in Table 1.

TABLE 1

MADRS reduction for all patients after treatment	
Timepoint	MADRS Reduction-all patients
Baseline	0
1 week	-10.4
2 weeks	-14.7
6 weeks	-20.6

A total of 37 of these patients suffered from major depressive disorder with suicidal ideation, defined as a score



of  $\geq 3$  on the Suicidality Item of the Montgomery-Åsberg Depression Rating Scale (MADRS-SI). For the patients with suicidal ideation, the mean MADRS score was 36.8 and the mean MADRS-SI score was 3.4 prior to treatment. The results for these patients are summarized in Table 2.

TABLE 2

MADRS reduction for suicidal ideation patients after treatment	
Timepoint	MADRS Reduction-suicidal ideation patients
Baseline	0
1 week	-12.9
2 weeks	-17.8
6 weeks	-22.8

Additionally, in the patients who suffered from suicidal ideation, the MADRS-SI score was reduced by 67.1% at the end of 1 week of treatment, 73.5% at the end of 2 weeks of treatment, and 82.4% at the end of 4 weeks of treatment as compared with baseline. This is potentially lifesaving because it shows a quick reduction in the risk of suicide.

The invention claimed is:

1. A method of treating depression and reducing risk of suicide, comprising selecting a human being suffering from major depressive disorder and having a score of 3 or greater on the Suicidality Item of the Montgomery-Åsberg Depression Rating Scale (MADRS-SI) and a score of at least 35 on the Montgomery-Åsberg Depression Rating Scale (MADRS), and administering a combination of about 90 mg to about 120 mg of bupropion hydrochloride, or a molar equivalent amount of another form of bupropion, and about 40 mg to about 50 mg of dextromethorphan hydrobromide, or a molar equivalent amount of another form of dextromethorphan, to a human being; wherein the combination is administered twice a day to the human being.

2. The method of claim 1, wherein the combination comprises about 105 mg of bupropion hydrochloride, or a molar equivalent amount of another form of bupropion, and about 45 mg of dextromethorphan hydrobromide, or a molar equivalent amount of another form of dextromethorphan.

3. The method of claim 1, wherein the combination is administered twice a day for at least one week.

4. The method of claim 1, wherein the combination is administered twice a day for at least two weeks.

5. The method of claim 1, wherein the combination is administered twice a day for at least six weeks.

6. The method of claim 1, wherein the MADRS-SI of the human being is reduced by at least 30% after the combination is administered to the human being twice a day for one week.

7. The method of claim 1, wherein the MADRS-SI of the human being is reduced by at least 50% after the combination is administered to the human being twice a day for two weeks.

8. The method of claim 1, wherein the MADRS-SI of the human being is reduced by at least 60% after the combination is administered to the human being twice a day for four weeks.

9. The method of claim 1, wherein after eight days of administration of the combination twice a day, the  $C_{max}$  of dextromethorphan in the human being is increased about 15-to about 25-fold as compared to administration of a single dose of the combination.

10. The method of claim 1, wherein after eight days of administering the combination twice a day, the  $C_{min}$  of dextromethorphan in the human being is increased about 35-to about 45-fold as compared to administering a single dose of the combination.

11. The method of claim 1, wherein after eight days of administering the combination twice a day, the  $AUC_{0-12}$  of dextromethorphan in the human being is increased about 25-to about 45-fold as compared to administering a single dose of the combination.

12. The method of claim 2, wherein after eight days of administering the combination twice a day, the  $C_{max}$  of dextromethorphan in the human being is about 75 ng/ml to about 80 ng/ml.

13. The method of claim 2, wherein after eight days of administering the combination twice a day, the  $C_{min}$  of dextromethorphan is about 42 ng/ml to about 50 ng/mL.

14. The method of claim 2, wherein after eight days of administering the combination twice a day, the  $AUC_{0-12}$  of dextromethorphan in the human being is about 755 ng·hr/mL.

15. The method of claim 2, wherein after eight days of administering the combination twice a day, the  $C_{max}$  of bupropion in the human being is about 85 ng/ml to about 90 ng/ml.

16. The method of claim 2, wherein after eight days of administering the combination twice a day, the  $C_{min}$  of bupropion is about 25 ng/ml to about 35 ng/ml.

17. The method of claim 2, wherein after eight days of administering the combination twice a day, the  $AUC_{0-12}$  of bupropion in the human being is about 660 ng·hr/mL to about 670 ng·hr/mL.

\* \* \* \* \*