



US012391999B2

(12) **United States Patent**
McKernan

(10) **Patent No.:** **US 12,391,999 B2**
(45) **Date of Patent:** ***Aug. 19, 2025**

(54) **METHODS AND KITS FOR CLASSIFYING CANNABINOID PRODUCTION IN CANNABIS PLANTS**

(71) Applicant: **MEDICINAL GENOMICS CORPORATION**, Beverly, MA (US)

(72) Inventor: **Kevin McKernan**, Marblehead, MA (US)

(73) Assignee: **MEDICINAL GENOMICS CORPORATION**, Beverly, MA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1102 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **17/346,709**

(22) Filed: **Jun. 14, 2021**

(65) **Prior Publication Data**

US 2022/0017974 A1 Jan. 20, 2022

Related U.S. Application Data

(62) Division of application No. 16/221,171, filed on Dec. 14, 2018, now Pat. No. 11,035,010.

(60) Provisional application No. 62/598,967, filed on Dec. 14, 2017.

(51) **Int. Cl.**
C12Q 1/6895 (2018.01)
C12N 9/02 (2006.01)

(52) **U.S. Cl.**
CPC **C12Q 1/6895** (2013.01); **C12N 9/0004** (2013.01); **C12Y 121/03007** (2015.07); **C12Y 121/03008** (2015.07); **C12Q 2600/13** (2013.01); **C12Q 2600/156** (2013.01); **C12Q 2600/16** (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,102,784 A	4/1992	George, Jr.
6,410,278 B1	6/2002	Notomi et al.
2012/0157326 A1	6/2012	Tisi et al.
2014/0057251 A1	2/2014	McKernan
2016/0177404 A1	6/2016	McKernan

FOREIGN PATENT DOCUMENTS

EP	0497272 A1	8/1992
WO	2016/189384 A1	12/2016
WO	2016/197258 A1	12/2016
WO	2019/118912 A1	6/2019

OTHER PUBLICATIONS

Gandelman et al. (International Journal of Mol. Sciences, vol. 12, pp. 9108-9124, 2011). (Year: 2011).*

McKernan et al. (bioRxiv, Plant Biology, 28654/1-28654/23, Oct. 2015) (Year: 2015).*

Focke et al. (J. of Agricultural and Food Chemistry, vol. 61, pp. 2943-2949, 2013). (Year: 2013).*

Tighe et al. (J. Biomol Tech. vol. 32, No. 3, pp. 199-205, Sep. 2021) (Year: 2021).*

Jul. 15, 2021 Examination Report issued in European Patent Application No. 18839647.7.

Allen et al. (J. Forensic Investigation, vol. 4, No. 1, pp. 1-7, Feb. 19, 2016.) (Year: 2016).

Taura et al. (FEBS Letters, vol. 581, pp. 2929-2934, 2007). (Year: 2007).

Kitamura et al. (J. Nat. Med. vol. 71, pp. 86-95, Aug. 17, 2016). (Year: 2016).

Stiedl et al. (J. of Vet. Diagnostic Investigation, vol. 29, No. 2, pp. 176-180, 2017). (Year: 2017).

Barany, F., Genetic disease detection and DNA amplification using cloned thermostable ligase, Proc Natl Acad Sci USA, 88(1):189-93 (1991).

Barringer, K.J. et al., Blunt-end and single-strand ligations by *Escherichia coli* ligase: influence on an in vitro amplification scheme, Gene, 89(1):117-22 (1990).

De Meijer, E.P. et al., The inheritance of chemical phenotype in *Cannabis sativa* L, Genetics, 163(1):335-46 (2003).

Gandelman, O. et al., Loop-mediated amplification accelerated by stem primers, Int J Mol Sci, 12(12):9108-24 (2011).

Gill, P. and Ghaemi, A., Nucleic acid isothermal amplification technologies: a review, Nucleosides Nucleotides Nucleic Acids, 27(3):224-43 (2008).

Guatelli, J.C. et al., Isothermal, in vitro amplification of nucleic acids by a multienzyme reaction modeled after retroviral replication, Proc Natl Acad Sci USA, 87(5):1874-8 (1990).

Higuchi, R. et al., Kinetic PCR analysis: real-time monitoring of DNA amplification reactions, Biotechnology (NY), 11(9):1026-30 (1993).

Kato, Y. and Braunstein, G.D., Beta-core fragment is a major form of immunoreactive urinary chorionic gonadotropin in human pregnancy, J Clin Endocrinol Metab, 66(6):1197-201(1988).

Kitamura, M. et al., Development of Loop-Mediated Isothermal Amplification (LAMP) Assay for Rapid Detection of Cannabis sativa, Biol Pharm Bull. 39(7):1144-9 (2016).

Kitamura, M. et al., Rapid identification of drug-type strains in Cannabis sativa using loop-mediated isothermal amplification assay, J Nat Med, 71(1):86-95 (2017).

Kojima, M. et al., DNA polymorphisms in the tetrahydrocannabinolic acid (THCA) synthase gene in "drug-type" and "fiber-type" *Cannabis sativa* L, Forensic Sci Int, 159(2-3):132-40 (2006).

(Continued)

Primary Examiner — Jeanine A Goldberg

(74) *Attorney, Agent, or Firm* — MH2 Technology Law Group, LLP

(57) **ABSTRACT**

The present disclosure provides methods and kits for characterizing *Cannabis* plants. Methods and kits of the present disclosure include detection/amplification of one or more enzymes involved in the production of cannabinoids, such as, for example, tetrahydrocannabinolic acid (THCA) synthase and/or cannabidiolic acid (CBDA) synthase.

20 Claims, 12 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited**OTHER PUBLICATIONS**

Kwoh, D.Y. et al., Transcription-based amplification system and detection of amplified human immunodeficiency virus type 1 with a bead-based sandwich hybridization format, *Proc Natl Acad Sci USA*, 86(4):1173-7 (1989).

Marks, M.D. et al., Identification of candidate genes affecting 9-tetrahydrocannabinol biosynthesis in *Cannabis sativa*, *J Exp Bot*, 60(13):3715-26 (2009).

Nagamine, K. et al., Accelerated reaction by loop-mediated isothermal amplification using loop primers, *Mol Cell Probes*, 16(3):223-9 (2002).

Onofri, C. et al., Sequence heterogeneity of cannabidiolic and tetrahydrocannabinolic acid-synthase in *Cannabis sativa* L. and its relationship with chemical phenotype, *Phytochemistry*, 116:57-68 (2015).

Staginnus, C. et al., A PCR marker linked to a THCA synthase polymorphism is a reliable tool to discriminate potentially THC-rich plants of *Cannabis sativa* L., *J Forensic Sci*, 59(4):919-26 (2014).

Walker, G.T. et al., Strand displacement amplification—an isothermal, in vitro DNA amplification technique, *Nucleic Acids Res*, 20(7):1691-6 (1992).

Walker, G.T., et al., Isothermal in vitro amplification of DNA by a restriction enzyme/DNA polymerase system, *Proc Natl Acad Sci USA*, 89(1):392-6 (1992).

Weiblen, G.D. et al., Gene duplication and divergence affecting drug content in *Cannabis sativa*, *New Phytol*, 208 (4):1241-50 (2015).

Wittwer, C.T. et al., Real-time multiplex PCR assays, *Methods*, 25(4):430-42 (2001).

Wu, D.Y. and Wallace, R.B., The ligation amplification reaction (LAR)-amplification of specific DNA sequences using sequential rounds of template-dependent ligation, *Genomics*, 4(4):560-9 (1989).
Cascini, F. et al., Analysis of THCA synthase gene expression in cannabis: a preliminary study by real-time quantitative PCR, *Forensic Sci Int*, 231(1-3):208-12 (2013).

International Search Report for PCT/US2018/065821 (Methods and Kits for Classifying Cannabinoid Production in Cannabis Plants, filed Dec. 14, 2018), issued by ISA/EP, 7 pages (Mar. 8, 2019).

Jeangkhwa, P. et al., Identification of *Cannabis sativa* L. using the 1-kbTHCA synthase-fluorescence in situ hybridization probe, *Sci Justice*, 57(2):101-106 (2017).

Allen, L.N. et al., Complex Variability within the THCA and CBDA Synthase Genes in Cannabis Species, *J Forensic Investigation*, p. 1, retrieved from Internet: <http://www.avensonline.org/wp-content/uploads/JFI-2330-0396-04--0029.pdf> the whole document (2016).

Written Opinion for PCT/US2018/065821 (Methods and Kits for Classifying Cannabinoid Production in Cannabis Plants, filed Dec. 14, 2018), issued by ISA/EP, 8 pages (Mar. 8, 2019).

Aug. 23, 2023 Office Action issued in Canadian Patent Application No. 3,123,182.

Zhang, Y. et al., "Isothermal Amplification of Long, Discrete DNA Fragments Facilitated by Single-Stranded Binding Protein", *Scientific Reports*, 7(8497), pp. 1-9 (2017).


Jun. 30, 2022 Office Action issued in Canadian Patent Application No. 3,123,182.

Clinton Torres et al., "LAVA: An Open-Source Approach to Designing LAMP (Loop-Mediated Isothermal Amplification) DNA Signatures," *BMC Bioinformatics*, 2011 12:240, pp. 1-7.

Aug. 30, 2022 Office Action issued in European Patent Application No. 18 839 647.7.

* cited by examiner

FIGURE 1



B




FIGURE 2




























CBDA-/CBDA-			THCA+/THCA+		Type Ia THCA plant
CBDA-/CBDA-			THCA+/THCA-		Type Ib THCA plant
CBDA+/CBDA-			THCA+/THCA+		Type IIa or 1:2 CBDA:THCA plant
CBDA+/CBDA-			THCA+/THCA-		Type IIb or 1:1 CBDA/THCA plant
CBDA+/CBDA+			THCA+/THCA+		Type IIb or 1:1 CBDA/THCA plant
CBDA+/CBDA+			THCA+/THCA-		Type IIc or 2:1 CBDA/THCA plant
CBDA+/CBDA-			THCA-/THCA-		Type IIIa CBDA plant (<10%)
CBDA+/CBDA+			THCA-/THCA-		Type IIIb CBDA plant (20%)
CBDA-/CBDA-			THCA-/THCA-		Type IV CBGA plant

FIGURE 3




FIGURE 3 (CONT.)




FIGURE 3 (CONT.)




FIGURE 4




FIGURE 5




FIGURE 6

Samples Mass Dispensaries	BarCode	THC	CBD	Well Location
White Fire OG_21NOV2016	FR13698731	17.3	0.3	A2
MediHaze CBD_21NOV2016	FR13697288	8.7	16.8	B2
Northern Lights #5 X Haze_21NOV2016	FR13695854	23.2	0.3	C2
OG Kush CBD_21NOV2016	FR13695841	4.5	9.8	D2
Phantom Cookies_21NOV2016	FR13694935	19.4	0.1	E2
Critical Mass_21NOV2016	FR13694452	9.84	13.41	F2
Super Silver Haze_21NOV2016	FR13697319	27.4	0.3	G2
Canna-Tsu_21NOV2016	FR13694429	6.5	13.5	H2
Grand Daddy Purple_21NOV2016	FR13695383	17.5	0.2	A3
Herijuana_21NOV2016	FR13698237	16.2	0.1	B3
California Dreaming_21NOV2016	FR13697370	19.2	0.2	C3
Citrix_29NOV2016	FR13695850	13.2	0.1	D3
Acacia Trainwreck_29NOV2016	FR13698721	12.6	0.0	E3
Master Kush_29NOV2016	FR13695413	16.3	0.1	F3
Harfox_29NOV2016	FR13695415	13.8	0.1	G3
Hurricane_29NOV2016	FR13697825	17.2	0.1	H3
Grape Stomper_29NOV2016	FR13697344	15.4	0.1	A4
Erk Train X Deadhead OG_29NOV2016	FR13697371	16.4	0.2	B4
Emerald Cup THC/CBD Samples	BarCode	THC	CBD	Well Location
NOTHERNLIGHTS CBD	FR09352163	6.72%	9.69%	A6
F-CANCER	FR09352247	6.10%	13.39%	B6
UNITY	FR09352367	10.02%	11.13%	C6
CANNATONIC	FR09352545	3.86%	12.47%	D6
GUERILLA MEDS	FR09352527	6.37%	10.07%	E6
TRIDENT	FR09352371	6.34%	17.79%	F6
SIERRA GOLD	FR09352486	8.18%	12.60%	G6
VITAMIN CBD	FR09352537	5.70%	11.54%	H6
HARLEQUIN CBD	FR09352134	3.51%	9.14%	A7
PINEAPPLETSU	FR09352298	6.67%	16.70%	B7






FIGURE 7




C Hemp Samples Run on both CBDA assay and Long Range THCA assay

Hemp Sample Map												
	1	2	3	4	5	6	7	8	9	10	11	12
A	M2	M1	M3	M6	CC8							
B	M5	M4	M7	M8	CC7							
C	M9	M10	M11	M12	CC5							
D	M13	M14	CC1	CC4	CC5							
E	CC3	CC2	CC7	CC6							M = Mother Mary	
F	CW5	CW4	CW3	CW2							CC = Charlotte's Cherries	
G	CW1	CC7	CC6	CC5							CC = Cobler	
H	CC4	CC3	CC2	CC1							CW = Cherry Wine	

Hemp Sample Map												
	1	2	3	4	5	6	7	8	9	10	11	12
A	M2	M1	M3	M6	CC8							
B	M5	M4	M7	M8	CC7							
C	M9	M10	M11	M12	CC5							
D	M13	M14	CC1	CC4	CC5							
E	CC3	CC2	CC7	CC6							M = Mother Mary	
F	CW5	CW4	CW3	CW2							CC = Charlotte's Cherries	
G	CW1	CC7	CC6	CC5							CC = Cobler	
H	CC4	CC3	CC2	CC1							CW = Cherry Wine	

FIGURE 8



THC_BIP	CACACAAGCACGTATTTGGGCTTTAGTTTTCACCTTAACTAACCT	SEQ ID NO: 8	Desalt	16uM
THC_FIP	GACTCGCATGATTAGTTTTTCTATCCTTATGTGTCCCAAATCC	SEQ ID NO: 9	Desalt	16uM
THC_Loop1	TCCCTATAATTGAGATACGCCAAT	SEQ ID NO: 10	Desalt	4uM
THC_F3	ATGGGAACCATATAAAAGTCAAT	SEQ ID NO: 11	Desalt	2uM
THC_B3	TATGATTGTCTACACAGTTCTAATGTAGATTTTC	SEQ ID NO: 12	Desalt	2uM

CBD_BIP	CAATCTTAGATTCACCTCTGACACAAGACATGTGAAGGAGTGAC	SEQ ID NO: 13	Desalt	16uM
CBD_FIP	TGTAATTGTCGAATTTAGGACAGACAATGCAACAAATCTAAACTCGTA	SEQ ID NO: 14	Desalt	16uM
CBD_Loop1	CAATGGGTTGTTTTGAGTGT	SEQ ID NO: 15	Desalt	4uM
CBD_Loop2	ACCCCAAAACCACCTTGT	SEQ ID NO: 16	Desalt	4uM
CBD_F3	CTTCTCGCAATATATTCCCAAT	SEQ ID NO: 17	Desalt	2uM
CBD_B3	GCATAGAATAGTGCCTTGGAT	SEQ ID NO: 18	Desalt	2uM

CBD_BIP	CAATCTTAGATTCACCTCTGACACAAGACATGTGAAGGAGTGAC	SEQ ID NO: 13	Desalt	16uM
CBD-NEG_FIP	TGTAATTGTCGAATTTAGGACAGACAATGCAACAAATCTAAACTTACA	SEQ ID NO: 19	Desalt	16uM
CBD_Loop1	CAATGGGTTGTTTTGAGTGT	SEQ ID NO: 15	Desalt	4uM
CBD_Loop2	ACCCCAAAACCACCTTGT	SEQ ID NO: 16	Desalt	4uM
CBD_F3	CTTCTCGCAATATATTCCCAAT	SEQ ID NO: 17	Desalt	2uM
CBD_B3	GCATAGAATAGTGCCTTGGAT	SEQ ID NO: 18	Desalt	2uM

FIGURE 9

FIGURE 10

Accession code	87	187	366	373	399	706	749	794	1179	1229	1395	1560	Seq. ID	THC(V)/A proportion
E33090	T	A	A	G	A	G	C	A	A	G	T	G	1/1	unknown
55.22.7.2.2	T	A	A	G	A	G	C	A	A	G	T	G	1/1	93.1%
	C	A	A	G	A	G	C	A	A	G	T	G	1/2	
55.24.4.34.7	T	A	A	G	A	G	C	A	A	G	T	G	1/1	89.1%
2009.27.44.2.19	T	A	A	G	A	C	C	A	A	G	T	G	1/3	11.1%
55.28.1.4.12	T	A	A	G	A	G	C	A	A	G	T	G	1/1	95.0%
2010.24a.35/T	T	C	T	G	G	G	C	A	T	G	T	G	3	96.2%
M110	T	A	A	G	A	G	C	A	A	G	T	G	1/1	96.4%
	T	A	T	C	A	G	C	A	A	A	A	A	2/1	
	T	A	T	C	A	G	C	A	A	G	T	G	2/2	
	T	A	T	C	A	G	C	A	A	A	T	G	2/3	
	T	A	A	G	A	G	C	G	A	A	A	A	4	
M124	T	C	T	G	G	G	C	A	T	G	T	G	3	98.6%
	T	A	A	G	A	G	A	A	A	G	T	G	1/4	
M128	T	A	A	G	A	G	C	A	A	G	T	G	1/1	96.5%
	T	C	T	G	G	G	C	A	T	G	T	G	3	
		63 Ileu ↓ Leu		125 Val ↓ Leu		236 Glu ↓ Gln	250 Ala ↓ Asp	265 Glu ↓ Gly		410 Gly ↓ Glu				

Onofri et al., (2015) *Phytochemistry*, 116:57-68, PMID: 25865737

METHODS AND KITS FOR CLASSIFYING CANNABINOID PRODUCTION IN CANNABIS PLANTS

CROSS-REFERENCE TO RELATED APPLICATION

This application is a divisional of U.S. application Ser. No. 16/221,171, filed Dec. 14, 2018, which claims the benefit of priority of U.S. Provisional Application No. 62/598,967, filed Dec. 14, 2017, all of which is incorporated herein by reference in its entirety. The present application contains a Sequence Listing that has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. The ASCII copy, created on Jun. 14, 2021, is named Sequence Listing ST25.txt and is 30,081 bytes in size.

BACKGROUND

C. sativa is an annual herbaceous flowering plant that has been cultivated throughout recorded history for use as fiber, seed oil, food, medicine and recreation. Cannabinoids are secondary metabolites found in *C. sativa* plants that can act on cannabinoid receptors in cells that alter neurotransmitter release in the brain. More than 100 cannabinoids have been isolated from *C. sativa*, which have varied effects on mammals to which they have been administered.

SUMMARY

The present disclosure encompasses the recognition that certain genotypes are associated with altered cannabinoid synthesis or an altered cannabinoid profile. For example, the present disclosure recognizes that certain variants (e.g., wild-type and/or mutated) of tetrahydrocannabinolic acid (THCA) synthase gene sequences and/or cannabidiolic acid (CBDA) synthase gene sequences can provide insight regarding synthesis of tetrahydrocannabinol (THC) and/or cannabidiol (CBD) in a *Cannabis* plant. The present disclosure provides a classification system for *Cannabis*, which designates a *Cannabis* plant as being of a particular type based on its genotype, which includes the classifications illustrated in FIG. 2 and outlined in Table 1 below:

TABLE 1

CBDA Genotype	THCA Genotype	Classification
CBDA ⁻ /CBDA ⁻	THCA ⁺ /THCA ⁺	Type Ia Plant
CBDA ⁻ /CBDA ⁻	THCA ⁺ /THCA ⁻	Type Ib Plant
CBDA ⁺ /CBDA ⁻	THCA ⁺ /THCA ⁺	Type IIa Plant
CBDA ⁺ /CBDA ⁻	THCA ⁺ /THCA ⁻	Type IIb Plant
CBDA ⁺ /CBDA ⁺	THCA ⁺ /THCA ⁺	Type IIc Plant
CBDA ⁺ /CBDA ⁺	THCA ⁺ /THCA ⁻	Type IIc Plant
CBDA ⁺ /CBDA ⁻	THCA ⁻ /THCA ⁻	Type IIIa Plant
CBDA ⁺ /CBDA ⁺	THCA ⁻ /THCA ⁻	Type IIIb Plant
CBDA ⁻ /CBDA ⁻	THCA ⁻ /THCA ⁻	Type IV Plant

Such a classification system can be utilized, e.g., in the selection of a *Cannabis* plant for various purposes. For example, a Type IIIb *Cannabis* plant (CBDA⁺/CBDA⁺ and THCA⁻/THCA⁻) may be useful as a therapeutic, particularly in applications in which psychoactive effects caused by THC are undesirable. The present disclosure also recognizes Type V *Cannabis* plants. In some embodiments, a Type V *Cannabis* plant is characterized by a mutation in one or both of a olivetol synthase gene sequence and a divarinic acid synthase gene sequence. Olivetolic acid and divarinic acid

are produced by enzymes olivetol synthase and divarinic acid synthase, respectively, in the cannabinoid synthesis pathway (see, e.g., FIG. 1, panel (A)).

The present disclosure further recognizes certain limits associated with previous assays to characterize THCA synthase, e.g., an inability to detect a THCA synthase gene sequence (e.g., because assays use primers or probes are complementary to highly variant sequences), and/or a false-positive detection (e.g., by amplification of a THCA synthase pseudogene). The present disclosure provides a method that reliably and accurately detects a THCA synthase gene sequence in a *Cannabis* plant. The present disclosure provides the insight that detection of a THCA synthase gene sequence (e.g., a wild-type or mutated THCA synthase gene sequence) present in a *Cannabis* plant can be achieved by amplification of a THCA synthase gene sequence using one or more primers with sequences complementary to sequences (e.g., promoter) upstream of the coding sequence and (e.g., terminator region, 5' UTR, etc.) downstream of the coding sequence may be particularly useful for in vitro detection of a functional THCA synthase gene sequence. Moreover, the present disclosure provides methods that include detection of a wild-type THCA synthase gene sequence and a mutant version of a THCA synthase gene sequence, thereby indicating the THCA synthase genotype of the plant.

The present disclosure also provides a method that reliably and accurately detects a CBDA synthase gene sequence in a *Cannabis* plant. The present disclosure provides methods that include detection of a wild-type CBDA synthase gene sequence and a mutant version of a CBDA synthase gene sequence, thereby indicating the CBDA synthase genotype of the plant. In some embodiments, detection of a wild-type CBDA synthase gene sequence in a *Cannabis* plant can be achieved by amplification of a CBDA synthase gene sequence using a primer that is complementary (at least in part) to a sequence of a first CBDA synthase gene sequence (e.g., CGTA at residues 330-333 in FIG. 4). In some embodiments, detection of a mutant CBDA synthase gene sequence in a *Cannabis* plant can be achieved by amplification of a CBDA synthase gene sequence using a primer that is complementary (at least in part) to a CBDA synthase gene sequence that includes a deletion of the first sequence (e.g., ACTTAC at residues 327-329 and 334-336 in FIG. 4).

In some aspects, the present disclosure relates to methods and kits for classifying cannabinoid production in a *Cannabis* plant, such as a *C. sativa* plant, a *C. indica* plant, or a *C. ruderalis* plant.

In some aspects, the present disclosure provides methods that include detection of one or more nucleic acid sequences in a sample of a *Cannabis* plant. In some embodiments, a method includes detecting the presence of a tetrahydrocannabinolic acid (THCA) synthase gene sequence, e.g., a THCA synthase gene sequence comprising a THCA synthase coding region (e.g., an entire THCA synthase coding region) and/or a THCA synthase gene sequence promoter. In some embodiments, a method includes detecting the presence of a mutated THCA synthase gene sequence. In some embodiments, a method includes detecting the presence of a wild-type THCA synthase gene sequence. In some embodiments, a method includes detecting the presence of a wild-type cannabidiolic acid (CBDA^{wt}) synthase gene sequence. In some embodiments, a method includes detecting the

3

presence of a mutated cannabidiolic acid (CBDA^{mut}) synthase gene sequence. In some embodiments, a mutated CBDA synthase gene sequence comprises a deletion mutation.

In some embodiments, a method includes detecting a wild-type THCA synthase gene sequence, a mutant THCA synthase gene sequence, a wild-type CBDA synthase gene sequence, a mutant CBDA synthase gene sequence, or any combination thereof from a sample that comprises *Cannabis* nucleic acid (i.e., nucleic acid of a *Cannabis* plant). In some embodiments, a method further includes detecting one, two, three, four, or more of a wild-type olivetol synthase gene sequence, a variant olivetol synthase gene sequence, a wild-type divarinic acid synthase gene sequence, a variant divarinic acid synthase gene sequence, a wild-type limonene synthase gene sequence, and a variant limonene synthase gene sequence from a sample that comprises *Cannabis* nucleic acid.

In some embodiments, detection of one or more nucleic acid sequences in a sample of *Cannabis* plant includes isothermal nucleic acid amplification. In some embodiments, isothermal nucleic acid amplification is a Loop-Mediated Isothermal Amplification (LAMP) assay. In some embodiments, a LAMP assay is a colorimetric LAMP assay.

In some embodiments, detection of a THCA synthase gene sequence includes amplification of a THCA synthase gene sequence that is at least 2000 nucleotides long. In some certain embodiments, an amplified THCA synthase gene sequences is 1500-2500, 1750-2250, 2000-2200, or 2100-2200 nucleotides long. In some embodiments, an amplified THCA synthase gene sequence includes at least part of the THCA synthase promoter and 5' UTR.

In some embodiments, a *Cannabis* plant is a *C. sativa* plant. In some embodiments, a *Cannabis* plant is a *C. indica* plant. In some embodiments, a *Cannabis* plant is a *C. ruderalis* plant.

In some embodiments, detecting the presence of a THCA synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least two THCA synthase primers under conditions sufficient for amplification of a THCA synthase gene sequence. In some certain embodiments, detecting the presence of a THCA synthase gene sequence includes contacting nucleic acid from a *Cannabis* plant with at least two primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to any of SEQ ID NOs: 8-12.

In some certain embodiments, detecting the presence of a THCA synthase gene sequence is by LAMP amplification and includes contacting nucleic acid from a *Cannabis* plant with four or more primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to any of SEQ ID NOs: 8-12.

In some embodiments, detecting the presence of a mutated THCA synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least two THCA synthase primers under conditions sufficient for amplification of a mutated THCA synthase gene sequence.

In some embodiments, detecting the presence of a wild-type cannabidiolic acid (CBDA^{WT}) synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least two CBDA synthase primers under conditions sufficient for amplification of a wild-type CBDA (CBDA^{WT}) synthase gene sequence. In some certain embodiments, detecting the presence of a

4

CBDA^{WT} synthase gene sequence includes contacting nucleic acid from a *Cannabis* plant with at least two primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to any of SEQ ID NOs: 13-18 and 27.

In some embodiments, detecting the presence of a mutated cannabidiolic acid (CBDA^{mut}) synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least two CBDA synthase primers under conditions sufficient for amplification of a CBDA^{mut} synthase gene sequence. In some embodiments a CBDA^{mut} synthase is a deletion (CBDA^{del}). In some certain embodiments, detecting the presence of a CBDA^{del} synthase gene sequence includes contacting nucleic acid from a *Cannabis* plant with at least two primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to any of SEQ ID NOs: 13, 15-19, and 27.

In some embodiments, detecting the presence of a THCA synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least five THCA synthase primers under conditions sufficient for LAMP of a THCA synthase gene sequence. In some certain embodiments, the at least five primer sequences are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to each of SEQ ID NOs: 8-12.

In some embodiments, detecting the presence of a mutated THCA synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least five THCA synthase primers under conditions sufficient for LAMP of a mutated THCA synthase gene sequence.

In some embodiments, detecting the presence of a wild-type cannabidiolic acid (CBDA^{WT}) synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least five CBDA synthase primers under conditions sufficient for LAMP of a CBDA^{WT} synthase gene sequence. In some certain embodiments, detecting the presence of a CBDA^{WT} synthase gene sequence is by LAMP amplification and includes contacting nucleic acid from a *Cannabis* plant with six primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to each of SEQ ID NOs: 13-18 and 27.

In some embodiments, detecting the presence of a mutated cannabidiolic acid (CBDA^{mut}) synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least five CBDA synthase primers under conditions sufficient for LAMP of a CBDA^{mut} synthase gene sequence. In some embodiments a CBDA^{mut} synthase is a deletion (CBDA^{del}). In some certain embodiments, detecting the presence of a CBDA^{del} synthase gene sequence is by LAMP amplification and includes contacting nucleic acid from a *Cannabis* plant with six primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to each of SEQ ID NOs: 13, 15-19, and 27.

In some embodiments, a method of the present disclosure includes a step of analyzing results from one or more detecting steps, thereby characterizing a *Cannabis* plant as a Type Ia, Type Ib, Type IIa, Type IIb, Type IIc, Type IIIa, Type IIIb, Type IV or Type V plant.

In some aspects, the present disclosure provides methods that include contacting a sample that comprises nucleic acid from a *Cannabis* plant with at least two primers sufficient for amplification of a *Cannabis* enzyme gene sequence. In some embodiments, a method includes contacting a sample that comprises nucleic acid from a *Cannabis* plant with at least two primers sufficient for amplification of a THCA synthase gene sequence under conditions sufficient for amplification of a THCA synthase gene sequence, where at least two THCA synthase primers include a forward THCA synthase primer and a reverse THCA synthase primer. In some embodiments, a forward THCA synthase primer is complementary to a sequence that is 200-1000 nucleotides upstream of a THCA synthase open reading frame in a *Cannabis* genome. In some embodiments, a reverse THCA synthase primer is complementary to a sequence that is 50-1000 nucleotides downstream of the THCA synthase open reading frame.

In some embodiments, a method includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least two cannabidiolic acid (CBDA) synthase primers under conditions sufficient for amplification of a Bd variant of CBDA synthase (CBDA^{Bd}) gene sequence. In some embodiments, at least one CBDA^{Bd} synthase primer is complementary to a sequence that bridges a 4 nucleotide deletion found in the CBDA^{Bd} open reading frame.

In some embodiments, a method includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least two cannabidiolic acid (CBDA) synthase primers under conditions sufficient for amplification of a wild-type CBDA (CBDA^{WT}) synthase gene sequence. In some embodiments, at least one CBDA^{WT} synthase primer is complementary to a sequence that includes the 4 nucleotide deleted in the CBDA^{Bd} synthase open reading frame.

In some embodiments, a method includes a combination of steps that include contacting a sample that comprises nucleic acid from the *Cannabis* plant with primers under conditions sufficient for amplification of two or more of a THCA synthase gene sequence, a variant THCA synthase gene sequence, a CBDA^{WT} synthase gene sequence, and/or a CBDA^{mut} synthase gene sequence.

In some embodiments, a method includes amplification of THCA synthase gene sequence that is at least 2000 nucleotide long. In some certain embodiments, an amplified THCA synthase gene sequences is 1500-2500, 1750-2250, 2000-2200, or 2100-2200 nucleotides long. In some embodiments, an amplified THCA synthase gene sequence includes at least part of the THCA synthase promoter and 5' UTR.

In some embodiments, amplification of one or more nucleic acid sequences in a *Cannabis* genome is by isothermal nucleic acid amplification. In some embodiments, amplification includes performing a Loop-Mediated Isothermal Amplification (LAMP) assay. In some embodiments, a LAMP assay is a colorimetric assay.

In some embodiments, a *Cannabis* plant is a *C. sativa* plant. In some embodiments, a *Cannabis* plant is a *C. indica* plant. In some embodiments, a *Cannabis* plant is a *C. ruderalis* plant.

In some embodiments, amplification of a THCA synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least five THCA synthase primers under conditions sufficient for LAMP of a THCA synthase gene sequence. In some certain embodiments, the at least five primer sequences are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to each of SEQ ID NOs: 8-12.

In some embodiments, amplification of a mutated THCA synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least five THCA synthase primers under conditions sufficient for LAMP of a mutated THCA synthase gene sequence.

In some embodiments, amplification of a wild-type cannabidiolic acid (CBDA^{WT}) synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least five CBDA synthase primers under conditions sufficient for LAMP of a wild-type CBDA (CBDA^{WT}) synthase gene sequence. In some certain embodiments, detecting the presence of a CBDA^{WT} synthase gene sequence is by LAMP amplification and includes contacting nucleic acid from a *Cannabis* plant with six primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to each of SEQ ID NOs: 13-18 and 27.

In some embodiments, amplification of a mutated cannabidiolic acid (CBDA^{mut}) synthase gene sequence includes contacting a sample that comprises nucleic acid from the *Cannabis* plant with at least five CBDA synthase primers under conditions sufficient for LAMP of a CBDA^{mut} synthase gene sequence. In some embodiments a CBDA^{mut} synthase is a deletion (CBDA^{del}). In some certain embodiments, detecting the presence of a CBDA^{del} synthase gene sequence is by LAMP amplification and includes contacting nucleic acid from a *Cannabis* plant with six primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to each of SEQ ID NOs: 13, 15-19, and 27.

In some aspects, the present disclosure provides kits that include primers that are complementary to sequences encoding *Cannabis* enzymes that are involved in cannabinoid synthesis. In some embodiments, a kit may be used to amplify a genomic sequence from a *C. sativa*, *C. indica*, and/or a *C. ruderalis* plant.

In some embodiments, a kit includes at least two THCA synthase primers including a forward THCA synthase primer and a reverse THCA synthase primer, wherein the forward THCA synthase primer is complementary to a sequence that is 200-1000 nucleotides upstream of a THCA synthase open reading frame in a *Cannabis* genome and the reverse THCA synthase primer is complementary to a sequence that is 50-1000 nucleotides downstream of the THCA synthase open reading frame.

In some embodiments, a forward THCA synthase primer is or comprises a sequence complementary to 20-60 nucleotides of SEQ ID NO: 6 and wherein the reverse THCA synthase primer is or comprises a sequence complementary to 20-60 nucleotides of SEQ ID NO: 7. In some embodiments, a kit includes at least five THCA synthase primers.

In some embodiments, a kit of the present disclosure may additionally or alternatively include at least two cannabidiolic acid (CBDA) synthase primers for amplification of a Bd variant of CBDA synthase (CBDA^{Bd}) gene sequence and/or at least two cannabidiolic acid (CBDA) synthase primers for amplification of a wild-type CBDA (CBDA^{WT}) synthase gene sequence. In some embodiments, at least one CBDA^{Bd} synthase primer is complementary to a sequence that bridges a 4 nucleotide deletion found in the CBDA^{Bd} open reading frame. In some embodiments, at least one CBDA^{WT} synthase primer is complementary to a sequence that includes the 4 nucleotides deleted in the CBDA^{Bd} synthase open reading frame.

In some certain embodiments, at least one CBDA^{WT} synthase primer is or comprises a sequence that is at least

80% identical to SEQ ID NO: 14. In some certain embodiments, at least one CBDA^{Bd} synthase primer is or comprises a sequence that is at least 80% identical to SEQ ID NO: 19. In some certain embodiments, at least one CBDA^{WT} synthase primer is or comprises a sequence of SEQ ID NO: 14. In some certain embodiments, at least one CBDA^{Bd} synthase primer is or comprises a sequence of SEQ ID NO: 19.

In some embodiments, a kit of the present disclosure may additionally or alternatively include primers for amplification of one or more of a wild-type olivetol synthase gene sequence, a variant olivetol synthase gene sequence, a wild-type divarinic acid synthase gene sequence, a variant divarinic acid synthase gene sequence, a wild-type limonene synthase gene sequence, and a variant limonene synthase gene sequence from a sample that comprises *Cannabis* nucleic acid.

Any of the kits of the present disclosure may additionally include reagents for a Loop-Mediated Isothermal Amplification (LAMP) assay. In some embodiments, a LAMP assay is a colorimetric LAMP assay.

These, and other aspects encompassed by the present disclosure, are described in more detail below and in the claims.

BRIEF DESCRIPTION OF THE DRAWING

The Figures described below, that together make up the Drawing, are for illustration purposes only, not for limitation.

FIG. 1: Panels (A) and (B) provide schematic representations of portion of the cannabinoid synthesis pathway.

FIG. 2: depicts a subtype classification system for *Cannabis* plants.

FIG. 3: depicts an alignment of various primers sequences with the sequence of a *Cannabis* THCA synthase gene sequence (SEQ ID NO: 20). Exemplary mutations in a THCA synthase gene sequence are provided along THCA-FIBER dotted underneath the ORF/CDS sequence.

FIG. 4: depicts a position-specific weight matrix for alignment of CBDA synthase alleles (SEQ ID NOS: 23, 24, and 25) in a *Cannabis* plant.

FIG. 5: depicts isothermic (e.g., LAMP) amplification of a THCA synthase gene sequence. Panel (A) depicts an isothermic (e.g., LAMP) amplification at 65° C. of THCA synthase gene sequence with Kitamura primers (“K primers”) (Kitamura et al., (2017) *Journal of natural medicines*, 71(1):86-95 and Kitamura et al., (2016) *Biological & pharmaceutical bulletin*, 39(7):1144-9, both of which are incorporated herein by reference) and Long Range primers (“LR primers”) on the *Cannabis* strain Granddaddy Purple (“GDP”). Additionally, an isothermic amplification with K primers of various strains of *Cannabis*: Otto, two hemp samples, Grape Stomper (RSP10516), and Erk Train X Deadhead OG (RSP10517) were also determined. Panel (B) depicts an isothermic (LAMP) amplification with LR primers at various temperatures for multiple strains of *Cannabis*.

FIG. 6: depicts an isothermic (LAMP) amplification of THCA gene sequence at 65° C. with Long Range primers on various *Cannabis* strains.

FIG. 7: depicts isothermic (LAMP) amplification of (Panel (A)) a CBDA synthase gene sequence and (Panel (B)) isothermic (LAMP) amplification of THCA gene sequence on various hemp varieties of *Cannabis*. Panel (C) includes a map of the samples run in both Panels (A) and (B).

FIG. 8: depicts isothermic (LAMP) amplification assays for CBDA^{WT} (Panel (A)) and CBDA^{del} (Panel (B)) sequences on various hemp varieties of *Cannabis*.

FIG. 9: depicts exemplary primer sequences (SEQ ID NOS: 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 13, 19, 15, 16, 17, and 18 in that order) useful in the context of the present disclosure.

FIG. 10: depicts a table of exemplary THCA synthase gene sequence.

CERTAIN DEFINITIONS

Associated With: The term “associated with” is used herein to describe an observed correlation between two items or events. For example, a mutation in THCA synthase may be considered to be “associated with” a particular cannabinoid synthesis profile and/or cannabinoid composition.

Cannabis: As used herein, “*Cannabis*” refers to any plant in the genus *Cannabis*. In some embodiments, “*Cannabis*” refers to a part of, a specific compound from, and/or any product from a *Cannabis* plant (e.g., *C. sativa*, *C. indica*, *C. ruderalis*). For example, a “*Cannabis* enzyme” refers to an enzyme from a *Cannabis* plant. Similarly, a “*Cannabis* genome” refers to a genome from a *Cannabis* plant. *Cannabis* includes both “marijuana” and “hemp,” two forms of *Cannabis* that are distinguished on the basis of the relative abundances of different cannabinoids. *Cannabis* includes any variety of *Cannabis* species, cultivar of *Cannabis* species, or hybrid between any *Cannabis* species.

Chemotype: As used herein, the term “chemotype” refers to chemically distinct entity (e.g., plant) with a particular profile of metabolites. In some embodiments, a chemotype is a particular type of *Cannabis* plant with a particular profile of one or more cannabinoids. In some embodiments, plants having different chemotypes may have same or different morphological characteristics. In some embodiments, a chemotype is characterized by a highly abundant chemical produced by that entity (e.g., plant). In some certain embodiments, a chemotype may refer to a *Cannabis* plant with an abundance of one or more cannabinoids (e.g., THC or CBD). In some embodiments, abundance of one or more cannabinoids may be a relative amount (e.g., a ratio of cannabinoids, such as a THC:CBD ratio).

Coding sequence: As used herein, the term “coding sequence” refers to a sequence of a nucleic acid or its complement, or a part thereof, that (i) can be transcribed to an mRNA sequence that can be translated to produce a polypeptide or a fragment thereof, or (ii) an mRNA sequence that can be translated to produce a polypeptide or a fragment thereof. Coding sequences include exons in genomic DNA or immature primary RNA transcripts, which are joined together by the cell’s biochemical machinery to provide a mature mRNA.

Gene and Gene sequence: The term “gene,” as used herein, refers to a part of the genome that codes for a product (e.g., an RNA product and/or a polypeptide product). A “gene sequence” is a sequence that includes at least a portion of a gene (e.g., all or part of a gene) and/or regulatory elements associated with a gene. In some embodiments, a gene includes coding sequence; in some embodiments, a gene includes non-coding sequence. In some particular embodiments, a gene may include both coding (e.g., exonic) and non-coding (e.g., intronic) sequences. In some embodiments, a gene may include one or more regulatory elements (e.g., a promoter) that, for example, may control or impact one or more aspects of gene expression (e.g., cell-type-specific expression, inducible expression, etc.).

Mutation: As used herein, the term “mutation” refers to a change introduced into a parental sequence, including, but

not limited to, substitutions, insertions, deletions (including truncations). The consequences of a mutation include, but are not limited to, the creation of a new character, property, function, phenotype or trait not found in the protein encoded by the parental sequence, or the increase or reduction/elimination of an existing character, property, function, phenotype or trait not found in the protein encoded by the parental sequence.

Nucleic Acid: As used herein, the terms “nucleic acid,” “nucleic acid molecule,” “oligonucleotide,” and “polynucleotide” are each used herein to refer to a polymer of at least three nucleotides. In some embodiments, a nucleic acid comprises deoxyribonucleic acid (DNA). In some embodiments comprises ribonucleic acid (RNA). In some embodiments, a nucleic acid is single stranded. In some embodiments, a nucleic acid is double stranded. In some embodiments, a nucleic acid comprises both single and double stranded portions. Unless otherwise stated, the terms encompass nucleic acid-like structures with synthetic backbones, as well as amplification products. In some embodiments, nucleic acids of the present disclosure are linear nucleic acids.

Plant part: As used herein, the term “plant part” refers to any part of a plant including but not limited to the embryo, shoot, root, stem, seed, stipule, leaf, petal, flower bud, flower, ovule, bract, trichome, branch, petiole, internode, bark, pubescence, tiller, rhizome, frond, blade, ovule, pollen, stamen, and the like. The two main parts of plants grown in some sort of media, such as soil or vermiculite, are often referred to as the “above-ground” part, also often referred to as the “shoots”, and the “below-ground” part, also often referred to as the “roots.” Plant part may also include certain extracts such as kief or hash which includes *cannabis* trichomes or glands.

Primer: The terms “primer,” as used herein, typically refers to oligonucleotides that hybridize in a sequence specific manner to a complementary nucleic acid molecule (e.g., a nucleic acid molecule comprising a target sequence). In some embodiments, a primer will comprise a region of nucleotide sequence that hybridizes to at least 8, e.g., at least 10, at least 15, at least 20, at least 25, or 20 to 60 nucleotides of a target nucleic acid (i.e., will hybridize to a sequence of the target nucleic acid). In general, a primer sequence is identified as being either “complementary” (i.e., complementary to the coding or sense strand (+)), or “reverse complementary” (i.e., complementary to the anti-sense strand (-)). In some embodiments, the term “primer” may refer to an oligonucleotide that acts as a point of initiation of a template-directed synthesis using methods such as PCR (polymerase chain reaction) under appropriate conditions (e.g., in the presence of four different nucleotide triphosphates and a polymerization agent, such as DNA polymerase in an appropriate buffer solution containing any necessary reagents and at suitable temperature(s)). Such a template directed synthesis is also called “primer extension.” For example, a primer pair may be designed to amplify a region of DNA using PCR. Such a pair will include a “forward primer” and a “reverse primer” that hybridize to complementary strands of a DNA molecule and that delimit a region to be synthesized and/or amplified.

Reference: As will be understood from context, a reference sequence, sample, population, agent or individual is one that is sufficiently similar to a particular sequence, sample, population, agent or individual of interest to permit a relevant comparison (i.e., to be comparable). In some embodiments, information about a reference sample is obtained simultaneously with information about a particular

sample. In some embodiments, information about a reference sample is historical. In some embodiments, information about a reference sample is stored for example in a computer-readable medium. In some embodiments, comparison of a particular sample of interest with a reference sample establishes identity with, similarity to, or difference of a particular sample of interest relative to a reference.

Regulatory Sequence: The term “regulatory sequence” is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals).

Wild type: As used herein, the term “wild-type” refers to a typical or common form existing in nature; in some embodiments it is the most common form.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

The disclosure provides useful methods for characterizing certain genetic sequences of *Cannabis* plants. The present disclosure encompasses a recognition that amplification of a region of THCA synthase gene sequence that includes a THCA synthase gene sequence promoter and/or termination region can robustly predict the presence of a functional THCA synthase gene sequence. The present disclosure also provides a method that reliably and accurately detects a CBDA synthase gene sequence in a *Cannabis* plant. Methods of the present disclosure may include characterization (e.g., detection, amplification) of CBDA gene sequences, including wildtype (CDBA^{wt}) and deletion (CDBA^{del}) sequences.

Cannabis Plants and Cannabinoids

Two sub-species of *Cannabis* plants are *C. indica* and *C. sativa*, which are commonly distinguished based on morphology of a *Cannabis* plant. Generally, *C. sativa* plants are taller, loosely branched and have long, narrow leaves, while *C. indica* plants are shorter, more densely branched and have wider leaves. However, there is some doubt about the accuracy of these generalizations. Originally, classification of *C. indica* was made in 1785 by a French biologist named Jean-Baptiste Lamarck who observed that certain marijuana plants from India were intoxicating and could be made into hashish. In contrast, traditional hemp crops in Europe had little or no mind-altering effect. Lamarck came up with the name *C. indica* to distinguish Indian *Cannabis* from European hemp, which was known at the time as *C. sativa*. Therefore, additionally or alternatively, *Cannabis* plants can be characterized by production of one or more chemical metabolites (e.g., cannabinoids). In some embodiments, a *Cannabis* plant is characterized as having a specified level (e.g., high/low) of one or more cannabinoids, flavonoids and/or terpenes.

Cannabinoids are terpenophenolic secondary metabolites, produced by *Cannabis* plants in the sessile and stalked trichomes. Trichomes are generally abundant on the inflorescences of a *Cannabis* plant, present in lower number on leaves, petioles and stems, and generally absent on roots and seeds. As a consequence, roots and seeds generally do not contain cannabinoids. In some embodiments, a *Cannabis* plant for use in a method of the present disclosure may include any plant part, such as, for example, a bloom, leaf, petiole, stem, root and/or seed.

As an annual, *Cannabis* plants follow a solar cycle consisting of two basic stages often referred to as vegetative, and bloom (flowering). Cannabinoid synthesis occurs predominantly in bloom (flowering) phase. In some embodi-

11

ments, a *Cannabis* plant for use in a method of the present disclosure is in a vegetative state. In some embodiments, a *Cannabis* plant for use in a method of the present disclosure is in a flowering state.

Cannabinoids such as, for example, delta-9-tetrahydrocannabinol (Δ^9 -THC or THC), cannabidiol (CBD), cannabitol (CBN), cannabigerol (CBG), and cannabichromene (CBC) have been identified from *Cannabis* plants. In some embodiments, a cannabinoid is an aryl-substituted monoterpene. Generally, cannabinoids are lipid soluble and neutral. Cannabinoids can be divided into at least ten classes: cannabigerol, cannabichromene, cannabidiol, delta-9-tetrahydrocannabinol, delta-8-tetrahydrocannabinol, cannabicyclol, cannabielsoin, cannabinol and cannabinodiol, cannabitriol and miscellaneous cannabinoids.

In some embodiments, *C. indica* plants are characterized as having high THC:CBD ratios and *C. sativa* plants are characterized as having high CBD:THC ratios. However, many strains produce varying amounts of cannabinoids, which may be due to hybridization, or cross breeding. Accordingly, in some embodiments, a *C. sativa* plant may be rich in THC and a *C. indica* plant may have low THC.

Cannabis plants for use in accordance with the methods of the present disclosure may be, for example, a *C. sativa* plant and/or a *C. indica* plant. In some embodiments, a *Cannabis* plant may be characterized as and/or determined to be rich in one or more cannabinoids such as a cannabigerol, cannabichromene, cannabidiol, delta-9-tetrahydrocannabinol, delta-8-tetrahydrocannabinol, cannabicyclol, cannabielsoin, cannabinol and cannabinodiol, and/or cannabitriol. In some embodiments, a *Cannabis* plant is determined to be or characterized as rich in THC. In some embodiments, a *Cannabis* plant is determined to be or characterized as rich in CBD.

In some embodiments, a *Cannabis* plant may be characterized as and/or determined to be expressing a low amount of one or more cannabinoids such as a cannabigerol, cannabichromene, cannabidiol, delta-9-tetrahydrocannabinol, delta-8-tetrahydrocannabinol, cannabicyclol, cannabielsoin, cannabinol and cannabinodiol, and/or cannabitriol. In some embodiments, a *Cannabis* plant is determined to be or characterized as low in THC. In some embodiments, a *Cannabis* plant is determined to be or characterized as low in CBD.

The present disclosure also encompasses the recognition that levels of two or more cannabinoids may be related. FIG. 1 depict portions of the cannabinoid synthesis pathway. As shown in FIG. 1, Panels A and B, cannabigerolic acid is a precursor of both THCA and CBDA. THCA and CBDA each undergo non-enzymatic conversion to THC and CBD, respectively. Since both THC and CBD are synthesized from the same precursor, increased synthesis of one of these compounds may reduce synthesis of the other (e.g., by depletion of precursor). In some embodiments, the rate/amount of THC synthesis is inversely proportional to the rate/amount of CBD synthesis. Moreover, as synthesis of both THC and CBD use cannabigerolic acid, increased synthesis of either or both of these cannabinoids may reduce the amount of cannabigerolic acid. In some embodiments, the rate/amount of THC synthesis is inversely proportional to the rate/amount of cannabigerolic acid. In some embodiments, the rate/amount of CBD synthesis is inversely proportional to the rate/amount of cannabigerolic acid.

Earlier studies proposed a model that the genes coding for functional THCA- and CBDA-synthase (referred to as Bt and Bd, respectively) are allelic and codominant. (de Meijer et al. (2003) *Genetics*, 163(1):335-46, PMID: 12586720,

12

which is incorporated herein by reference). Variations in CBDA/THCA ratios could be directly related to a differential efficiency in transforming cannabigerolic acid by THCA- and CBDA-synthases. Id.

In some embodiments, a *Cannabis* plant genome includes a THCA synthase gene sequence. In some embodiments, a *Cannabis* plant genome includes a wild-type THCA synthase gene sequence. In some embodiments, a *Cannabis* plant genome is homozygous for a wild-type THCA synthase gene sequence. In some embodiments, a *Cannabis* plant genome is heterozygous for a wild-type THCA synthase gene sequence. In some embodiments, a *Cannabis* plant genome is homozygous for a variant THCA synthase gene sequence. In some embodiments, a *Cannabis* plant genome is heterozygous for a variant THCA synthase gene sequence. In some embodiments, a *Cannabis* plant genome includes a THCA synthase gene sequence that is or comprises a sequence that is 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to SEQ ID NO: 1, or a portion thereof.

Exemplary *C. sativa* THCA synthase gene sequence
SEQ ID NO: 1
ATGGGAACCATAATAAACTATAAAAGTCATTATGTGTAAGTCTACCAT
AGGCACCTATATCCACAACTAGCTACCATAGCCAATTCTTTTGTG
TTCCAATATCCAATTTTATTGATGCCAACTATTCAATGTACAATGTA
CATTATTTTCAATAAGGGCTTCACCTAACAAAGGTGCCTAATTTTAGT
TGATTTATTTTATCAGATGTGACTATTATGACTATCAAAATATAA
AATATTTAAGTCAATTTATTTGCCCACTCCAATATATAATATTATAA
ATAGGATAGTTCTCAATTCCTAATAATTCAAAAAATCATTAGGACTGAA
GAAAAATGAATTGCTCAGCATTTTCTTTTGGTTGTGTCAAAATAAT
ATTTTCTTTCTCTCATTCATATCCAAATTTCAATAGCTAATCCTCGA
GAAACTTCCTTAAATGCTTCTCAAAACATATCCCAACATGTAGCAA
ATCCAAACTCGTATACACTCAACACGACCAATGTATATGTCTATCCT
GAATTCGACAATACAAATCTTAGATTATCTCTGATACACCCCAAAA
CCACTCGTTATTGTCACTCCTTCAATAAATCCCATATCCAAGCAACTA
TTTATGCTCTAAGAAAGTTGGCTTGAGATTGCAACTCGAAGCGGTGG
CCATGATGCTGAGGTATGTCCTACATATCTCAAGTCCCATTTGTGTA
GTAGACTTGAGAAACATGCATTCGATCAAAATAGATGTTATAGCCAAA
CTGCGTGGTTGAAGCCGAGCTACCCTTGAGAGAGTTTATTATTGGAT
CAATGAGAAGAAATGAGAATCTTAGTTTCTGGTGGGTATTGCCTACT
GTTGGCGTAGGTGGACACTTTAGTGGAGGAGGCTATGGAGCATTGATGC
GAAATTATGGCCTTGCGGCTGATAATATTATTGATGCACACTTAGTCAA
TGTTGATGGAAAAGTTCTAGATCGAAAATCCATGGGAGAAGATCTGTTT
TGGGCTATACGTGGTGGTGGAGGAGAAAACCTTGAATCATTGCAGCAT
GGAAAATCAAACCTGGTTGATGTCCTCAAAAGTCTACTATATTAGTGT
TAAAAGAACATGGAGATACATGGGCTTGTCAAGTTATTTAACAAATGG
CAAAATATTGCTTACAAGTATGACAAAGATTTAGTACTCATGACTCACT
TCATAACAAAGAATATTACAGATAATCATGGGAAGAATAAGACTACAGT

13

-continued

ACATGGTTACTTCTCTCAATTTTTCATGGTGGAGTGGATAGTCTAGTC
 GACTTGATGAACAAGAGCTTTCCTGAGTTGGGTATTAAAAAACTGATT
 GCAAAGAATTTAGCTGGATTGATACAACCATCTTCTACAGTGGTGTGT
 AAATTTTAACACTGCTAATTTTAAAAAGGAAATTTTGCTTGATAGATCA
 GCTGGGAAGAAGACGGCTTTCCTCAATTAAGTTAGACTATGTTAAGAAAC
 CAATTCAGAACTGCAATGGTCAAAATTTTGGAATAATATATGAAGA
 AGATGTAGGAGCTGGGGTGTGTACCCCTACGGTGGTATAATGGAGGAG
 ATTTGAGAATCAGCAATTCATTCCCTCATCGAGCTGGAATAATGTATG
 AACTTTGGTCACTGCTTCTGGGAGAGCAAGAAGATAATGAAAGCA
 TATAAAGTGGTTCGAAGTGTATATAATTTACGACTCCTTATGTGTCC
 CAAAATCCAAGATTGGCGTATCTCAATTATAGGGACCTTGATTTAGGAA
 AACTAATCATGCGAGTCTTAATAATTACACACAAGCAGTATTTGGGG
 TGAAAAGTATTTTGGTAAAAATTTTAACAGGTAGTTAAGGTGAAAACT
 AAAGTTGATCCCAATAATTTTGTAGAAACGAACAAAGTATCCCACTC
 TTCCACCGCATCATCATTAAATATCTTTAATAGATATATTTCCCTTAT
 CAATTAGTTAATCATTATACCATACATACATTATTGTATATAGTTTAT
 CTACTCATATTATGTATGCTCCCAAGTATGAAAATCTACATTAGAACTG
 GTAGACAATCATA

In some embodiments, a *Cannabis* plant genome includes a CBDA synthase gene sequence. In some embodiments, a *Cannabis* plant genome includes a wild-type CBDA synthase gene sequence. In some embodiments, a *Cannabis* plant genome is homozygous for a wild-type CBDA synthase gene sequence. In some embodiments, a *Cannabis* plant genome is heterozygous for a wild-type CBDA synthase gene sequence. In some embodiments, a *Cannabis* plant genome is homozygous for a variant CBDA synthase gene sequence. In some embodiments, a *Cannabis* plant genome is heterozygous for a variant CBDA synthase gene sequence. In some embodiments, a variant CBDA synthase gene sequence comprises a deletion. In some embodiments, a variant CBDA synthase gene sequence comprises a deletion of a sequence comprising CGTA (SEQ ID NO:3). In some embodiments, a *Cannabis* plant genome includes a CBDA synthase gene sequence that is or comprises a sequence that is 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to SEQ ID NO: 2, or a portion thereof.

Exemplary *C. sativa* CBDA synthase gene sequence
 SEQ ID NO: 2
 GATATATATCTCACACGGATGCACCTAACATGATGCCTAATTTTGTG
 AATTTTTTTTACCACATGACTTAATGATATCAAATTATGAATATTTAG
 TTAATTTATTTGCCCTGCTCCAATATATAAGCTATAAATAGGATAGT
 TCTTAATCCATAGTAATT

CAAAATTCATTAGAACTAAAGAAAAATGAAGTGTCAACATTCTCCTTT
 TGGTTTGTTCGAAGATAATATTTTCTTTTCTCATTCAATATCCAA

14

-continued

CTTCATTGCTAATCCTCGAGAAAACCTTCCTAAATGCTTCTCGCAATA
 TATTCCCAATAATGCAACAAATCTAAAACTCGTATACACTCAAAACAAC
 CCATTGTATATGTCTGTCTTAAATTCGACAATACACAATCTTAGATTCA
 CCTCTGACACAACCCCAAAACCACTTGTATCGTCACTCCTTCACATGT
 CTCTCATATCCAAGGCACTATTCTATGCTCCAAGAAAGTTGGCTTGCAG
 ATTCGAACTCGAAGTGGTGGTCATGATTCTGAGGGCATGTCTACATAT
 CTCAGTCCCATTGTGTATAGTAGACTTGAGAAACATGCGTTCAATCAA
 AATAGATGTTTATAGCCAAATGCATGGGTTGAAGCCGAGCTACCCCTT
 GGAGAAGTTTATTATTTGGGTTAATGAGAAAAATGAGAATCTTAGTTTGG
 CGGCTGGGTATTGCCCTACTGTTTTCGCGAGGTGGCACTTTGGTGGAGG
 AGGCTATGGACCATTGATGAGAACTATGGCCTCGCGGCTGATAATATC
 ATTGATGCACACTTAGTCAACGTTTCATGGAAGTGTAGATCGAAAT
 CTATGGGGGAAGATCTCTTTTGGGCTTACGTGGTGGTGGAGCAGAAAG
 CTCGGAATCATTGTAGCATGGAATAATAGACTGGTTGCTGTCCCAAAG
 TCTACTATGTTTAGTGTAAAAGATCATGGAGATACATGAGCTTGTCA
 AGTTAGTTAACAAATGGCAAAATATTGCTTACAAGTATGACAAAGATT
 ATTACTCATGACTCACTTCATAACTAGGAACATTACAGATAATCAAGGG
 AAGAATAAGACAGCAATACACACTTACTTCTCTTCAGTTTTCTTGGTG
 GAGTGGATAGTCTAGTCTGACTTGATGAACAAGAGTTTCTCGAGTTGGG
 TATTAACAAAAACGGATTGCAGACAATTGAGCTGGATTGATACTATCATC
 TTCTATAGTGGTGTGTAAATACGACACTGATAATTTTAAACAGGAAA
 TTTTCTTGATAGATCCGCTGGGCAGAACGGTGTCTTCAAGATTAAGTT
 AGACTACGTTAAGAAACCAATCCAGAATCTGTATTTGTCCAAATTTTG
 GAAAAATTATGAAGAAGATATAGGAGCTGGGATGTATGCGTTGTACC
 CTTACGGTGGTATAATGGATGAGATTTTCAAGTATGAGCAATTCATTCCC
 TCATCGAGCTGGAATCTTGTATGAGTTATGGTACATATGTAGTTGGGAG
 AAGCAAGAAGATAACGAAAGCATCTAACTGGATTAGAAATATTTATA
 ACTTTCATGACTCCTTATGTGTCCAAAAATCCAAGATTGGCATATCTCAA
 TTATAGAGACCTTGATATAGGAATAAATGATCCCAAGAAATCCAAATAAT
 TACACACAAGCAGTATTTGGGGTGAAGATATTTGGTAAAAATTTTG
 ACAGGCTAGTAAAAGTGAAAACCTGGTTGATCCCAATAACTTTTTTAG
 AAACGAACAAAGCATCCACCTCTTCCACGGCATCGTCATTAATGATCT
 TAAATAGATCTTTTTCTCTTATTAATAGTCTTATAATATACATATAT
 TGATTATATATATAAAAAATAGTTTGTCCGGGTGACTGTGTATGCGATA
 TATATCTCACAC

In some embodiments, a *Cannabis* plant genome includes an olivetol synthase gene sequence. In some embodiments, a *Cannabis* plant genome includes a wild-type olivetol synthase gene sequence. In some embodiments, a *Cannabis* plant genome is homozygous for a wild-type olivetol synthase gene sequence. In some embodiments, a *Cannabis* plant genome is heterozygous for a wild-type olivetol synthase gene sequence. In some embodiments, a *Cannabis* plant genome is homozygous for a variant olivetol synthase gene sequence.

15

gene sequence. In some embodiments, a *Cannabis* plant genome is heterozygous for a variant olivetol synthase gene sequence. In some embodiments, a *Cannabis* plant genome includes an olivetol synthase gene sequence that is or comprises a sequence that is 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to SEQ ID NO: 4, or a portion thereof.

Exemplary *C. sativa* olivetol synthase gene sequence

SEQ ID NO: 4

ATGAATCATCTTCGTGCTGAGGGTCCGGCTCCGTTCTCGCCATTGGCA
CCGCCAATCCGAGAACATTTATTACAAGATGAGTTTCTTGACTACTA
TTTTCGCGTCACCAAAAGTGAACACATGACTCAACTCAAAGAAAAGTTT
CGAAAAATATGTGACAAAAGTATGATAAGGAAACGTAACGTGTTCTTAA
ATGAAGAACACCTAAAGCAAAACCAAGATTGGTGGAGCAGAGATGCA
AACTCTGGATGCACGTCAAGACATGTTGGTAGTTGAGGTTCCAAAACCT
GGGAAGGATGCTTGTGCAAAAGGCCATCAAAGAATGGGGTCAACCAAGT
CTAAATCACTCATTTAATCTTCACTAGCGCATCAACCACTGACATGCC
CGGTGCAGACTACCATTGCGCTAAGCTTCTCGACTGAGTCCCTCAGTG
AAGCGTGTGATGATGATCAACTAGGCTGTTATGGTGGTGAACCGTTC
TACGCATTGCCAAGGACATAGCAGAGAATAACAAAGGCGCACGAGTTCT
CGCCGTGTGTTGTGACATAATGGCTTGCTTGTTCGTGGGCCCTCAGAG
TCTGACCTCGAATTACTAGTGGGACAAGCTATCTTGGTGTGAGGGCTG
CTGCGGTGATTGTTGGAGCTGAACCCGATGAGTCAGTTGGGGAAGGCC
GATATTTGAGTTGGTGTCAACTGGGCAACAATCTTACCAAACTCGGAA
GGAACTATTGGGGGACATATAAGGGAAGCAGGACTGATATTGATTTAC
ATAAGGATGTGCTATGTTGATCTCTAATAATATTGAGAAATGTTTGAT
TGAGGCATTTACTCTATTGGGATTAGTGATTGGAACCTCATATTTTGG
ATTACACACCCAGGTGGGAAAGCTATTTTGGACAAAGTGGAGGAGAAGT
TGCATCTAAAGAGTGATAAGTTTGTGGATTACGTCATGTGCTGAGTGA
GCATGGGAATATGTCTAGCTCAACTGTCTTGTGTTTATGGATGAGTTG
AGGAAGAGGTGCTTGGAGGAAGGGAAGTCTACCACTGGAGATGGATTTG
AGTGGGGTGTCTTTTGGGTTTGGACCAGGTTTGACTGTGCAAGAGAT
GGTCTGTCGTAGTGTTCATCAATATTAA

In some embodiments, a *Cannabis* plant genome includes a divarinic acid synthase gene sequence. In some embodiments, a *Cannabis* plant genome includes a wild-type divarinic acid synthase gene sequence. In some embodiments, a *Cannabis* plant genome is homozygous for a wild-type divarinic acid synthase gene sequence. In some embodiments, a *Cannabis* plant genome is heterozygous for a wild-type divarinic acid synthase gene sequence. In some embodiments, a *Cannabis* plant genome is homozygous for a variant divarinic acid synthase gene sequence. In some embodiments, a *Cannabis* plant genome is heterozygous for a variant divarinic acid synthase gene sequence.

In some embodiments, a *Cannabis* plant genome includes a limonene synthase gene sequence. In some embodiments, a *Cannabis* plant genome includes a wild-type limonene synthase gene sequence. In some embodiments, a *Cannabis*

16

plant genome is homozygous for a wild-type limonene synthase gene sequence. In some embodiments, a *Cannabis* plant genome is heterozygous for a wild-type limonene synthase gene sequence. In some embodiments, a *Cannabis* plant genome is homozygous for a variant limonene synthase gene sequence. In some embodiments, a *Cannabis* plant genome is heterozygous for a variant limonene synthase gene sequence. In some embodiments, a *Cannabis* plant genome includes a limonene synthase gene sequence that is or comprises a sequence that is 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to SEQ ID NO: 5, or a portion thereof.

Exemplary *C. sativa* limonene synthase gene sequence

SEQ ID NO: 5

ATGCAGTGCATAGCTTTTACCAATTTGCTTCATCATCATCCCTCCCTA
TTTGGAGTAGTATTGATAATCGTTTTACACCAAAACTTCTATTACTTC
TATTTCAAACCAAAACCAAACTAAATCAAATCAAACCTGAAATCG
AGATCGAGATCAAGTACTTGTACTCCATACATGTACTGTGGTCGATA
ACCCTAGTTCTACGATTACTAATAATAGTGATCGAAGATCAGCCAACTA
TGGACCTCCCATTTGGTCTTTTGATTTTGTTCAATCTCTTCCAATCCAA
TATAAGGGTGAATCTTATACAAGTCGATTAAATAAGTTGGAGAAAGATG
TGAAAAGGATGCTAATTGGAGTGGAAAACCTTTAGCCCAACTTGAAC
AATTGATACAATACAAGACTTGAATATCTTATCGTTTTGAAAATGAA
ATCATTTCTATTTTGAAGAAAAATTACCAATAATAATGACAACCCCTA
ATCCTAATTATGATTTATATGCTACTGCTCTCCAATTTAGGCTTCTACG
CCAATATGGATTGAAGTACCTCAAGAAATTTCAATAATTTTAAAAAT
CACAAGACAGGAGAGTTCAAGGCAATATAAGTAATGATATTATGGGAG
CATTGGGCTTATATGAAGCTTCATTCATGGGAAAAAGGGTGAAAGTAT
TTTGAAGAAGCAAGAATTTTCAACAATAATGTCTCAAAAAATACAAA
TTAATGTCAAGTAGTAATAATAATATGACATTAATATCATTATTAG
TGAATCATGCTTTGGAGATGCCACTTCAATGGAGAATCACAAGATCAGA
AGCTAAATGGTTTATGAAGAAATATATGAAGAAAAACAAGACATGAAT
CCAACCTTACTTTAGTTTGGCAAAATGGGATTTCAATATGCTGCAATCAA
CATATCAAGAGGAGCTCAAAGTACTCTCTAGTGGTGGGAAGATTCTAA
ACTTGGAGAGAAATGCCTTTGTTAGAGATAGATTGGTGGAGTGTTC
TTATGGCAAGTTGGAGTAAGATTTGAGCCACAATTCAGTTACTTTAGAA
TAATGGATACAAAACCTATGTTCTATTAACAATAATTGATGATATGCA
TGACATTTATGGAACATTGGAGGAACACAACCTTTCACTAATGCTCTT
CAAAGATGGGATTTGAAGAATTAGATAAATTACCAGATTATATGAAGA
CAGCTTTCTACTTTACATACAATTTCAAAATGAATTGGCATTGATGT
ATTACAAGAACATGGTTTGTTCATTTGAATACTTCAAGAAACGATG
GTAGAGTTGTGTAACATCATTTGCAAGAGGCAAAATGGTTTATAGTG
GATACAAACCAACATTGCAAGAATATGTTGAGAATGGATGGTTGTCTGT
GGGAGGACAAGTTATTTCTTATGCATGCATATTTGCTTTTACAAATCCT

-continued

GTTACCAAGAGGCATTGGAATGTCATAAAGACGGTCATCTAACATAG
 TTCGCCATGCATCGATAATATTACGACTTGCGATGATCTAGGAACATT
 GTCGGATGAAGTGAAGAGAGCGATGTTCTTAAATCAATTCAATGTTAT
 ATGCACGATACTGGTGCTTCTGAAGATGAAGCTCGTGAGCACATAAAAT
 ATTTAATAAGTGAATCATGGAAGGAGATGAATAATGAAGATGGAAATAT
 TAACTCTTTTTCTCAAATGAATTTGTTCAAGTTTGCCAAAATCTTGGT
 AGAGCGTCACAATTATATACAGTATGGCGATGGACATGCTTCTCAGA
 ATAATCTATCGAAAGAGCGCGTTTATAGGGTTGATTATTACTCTTATCCC
 CATGTAA

Classification of *Cannabis*

Provided herein is a recognition that certain genotypes are associated with altered cannabinoid synthesis or an altered cannabinoid profile. Genetic information can positively benefit consumers, growers, and regulators of *Cannabis*. There are at least several hundred compounds found in *Cannabis* and currently only a fraction of these compounds are analyzed. As used herein, chemotype of a *Cannabis* plant refers to an abundance and/or deficiency of one or more compounds found in a *Cannabis* plant (e.g., cannabinoids, flavonoids and/or terpenes). In some embodiments, a chemotype is an abundance of one or more cannabinoids, flavonoids and/or terpenes. In some embodiments, a chemotype is a relative amount of two or more cannabinoids, flavonoids and/or terpenes.

Chemotype determination has been limited as quantitative analytical techniques and standards do not currently exist for all *Cannabis* compounds. As a result, previously described chemotypes fail to capture the chemical and potential medicinal complexity in a *Cannabis* plant. The present disclosure encompasses a recognition that genetic analysis, for example, of one, two, three, four, five or more synthetic enzymes, can serve as a proxy for unmeasured chemical complexity.

The present disclosure encompasses the recognition that genetic analysis can be used to assess the potential of a *Cannabis* plant to produce certain compounds, such as cannabinoids, flavonoids and/or terpenes. While a *Cannabis* plant can be raised under conditions that result in varied expression and/or concentrations of certain compounds, the relative ratios of key chemotypic gene sequences, such as, for example, THCA synthase and CBDA synthase, are usually genetically determined. For example, there are currently no known agricultural methods to make a CBD-dominant *Cannabis* strain become a THC-dominant *Cannabis* strain via environmental conditions. These critical chemotypes are governed by genetic alterations in their respective enzymatic syntheses (e.g., CBDA synthase and THCA synthase).

In some embodiments, methods of the present disclosure include genetic analysis of one, two, three, four, five or more enzymes involved in the production of cannabinoids, flavonoids and/or terpenes. In some embodiments, methods of the present disclosure include amplification of a genomic region that encodes an enzyme involved in the production of a cannabinoid, flavonoid and/or terpene. In some embodiments, methods of the present disclosure include

amplification of a portion of a gene sequence that encodes an enzyme involved in the production of a cannabinoid, flavonoid and/or terpene.

Genetic prediction of cannabinoid production has been an active area of study. See, for example, Weiblen et al., (2015), *The New Phytologist*, PMID: 26189495; Marks et al., (2009) *Journal of Experimental Botany*, 60(13):3715-26, PMID: 19581347; Onofri et al., (2015) *Phytochemistry*, 116:57-68, PMID: 25865737; de Meijer et al. (2003) *Genetics*, 163(1): 335-46, PMID: 12586720; and Kojoma et al. (2006), *Forensic Sci. Int.*, 159(2-3):132-40, PMID: 16143478, each of which is incorporated herein by reference. However, there are significant caveats and shortcomings with prior methods of genetic prediction of cannabinoid production. For example, while alleles of Bt (functional THCA synthase) and Bd (functional CBDA synthase) have been proposed as a model for generating Type I, II, III, and IV plants, the exact DNA sequences that govern these alleles remain unknown. As described in the examples herein, single molecule sequencing was performed to identify mutations in gene sequences in cannabinoid synthesis pathway enzyme(s) (e.g., THCA synthase and CBDA synthase). This sequencing revealed a more refined structure of inheritance and numerous additional subtypes that can be identified by genotyping to predict chemical inheritance. See, FIG. 2.

Kitamura describes amplification of a small region of THCA synthase to resolve hemp from drug-type (i.e., THC expressing) strains. Kitamura et al., (2017) *Journal of natural medicines*, 71 (1):86-95, PMID: 27535292 and Kitamura et al., (2016) *Biological & pharmaceutical bulletin*, 39 (7):1144-9, PMID: 27118244, each of which is incorporated herein by reference. However, the present disclosure recognizes that these studies only tested 3 cultivars and also failed to consider CBDA status. The present disclosure encompasses a recognition that an understanding CBDA status is critical for determining allelic balance of CBDA and THCA synthases, which may be important for *Cannabis* breeding efforts.

Cannabis genomes are about 10 fold more variable than human genomes. Moreover, a given *Cannabis* strain is often capable of crossing with highly divergent *Cannabis* strains (e.g., *Cannabis* strains of a different chemotype, genotype, etc.). As a result, even sibling strains cannot be not be assumed to have the same chemotype. The present disclosure provides methods and kits suitable for genetic characterization of individual *Cannabis* plants.

In some embodiments, a method includes detection of one or a plurality of sequences that include a gene sequence or portion thereof in a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, a gene sequence encodes an enzyme involved in the synthesis of one or more cannabinoids.

In some embodiments, a method of the present disclosure includes detecting the presence of a tetrahydrocannabinolic acid (THCA) synthase gene sequence or a portion thereof. The present disclosure encompasses the recognition that THCA synthase is under selective breeding pressure and has approximately a 4 fold higher polymorphism rate than a *Cannabis* genome in general. Moreover, a THCA synthase gene sequence can include a SNP approximately every 25 nucleotides, which complicates design of primers that anneal within the open reading frame of THCA synthase. In some embodiments, a THCA synthase gene sequence encodes a wildtype THCA synthase enzyme (i.e., functional). In some embodiments, a THCA synthase gene sequence includes a THCA synthase gene sequence pro-

motor sequence. In some embodiments, a THCA synthase gene sequence includes a THCA synthase gene sequence 5' UTR sequence.

In some embodiments, a method of the present disclosure includes amplifying a THCA synthase gene sequence or portion thereof from a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, amplification of a THCA synthase gene sequence includes contacting nucleic acid from a *Cannabis* plant with a forward THCA synthase primer that is complementary to a sequence that is 200-1000 nucleotides upstream of a THCA open reading frame. In some embodiments, a forward THCA synthase primer is complementary to a sequence that is 200 to 800 nucleotides, 200 to 600 nucleotides, 200 to 400 nucleotides upstream of a THCA open reading frame. In some certain embodiments, a forward THCA synthase primer is complementary to a sequence that is at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a portion of SEQ ID NO: 6. In some certain embodiments, a forward THCA synthase primer is at 20 to 60 nucleotides long.

Exemplary *C. sativa* sequence upstream of THCA synthase open reading frame

SEQ ID NO: 6

CATAGCGACTATCGTGATGGGAACCATAATAAACTATAAAAGTCATTAT
GTGTACTTGTCTACCATAGGCACCTATATCCCAACAACTAGCTACCATAG
CCAATTTCTTTTTTGTTCCTAATATCAATTTTTTATGATGCCAAACTA
TTCAATGTACAATGTACATTTATTTTCAATAAGGCTTCACCTAACAAA
GGTGCTAATTTTAGTTGATTTATTTTATCAGATGTGACTATTTAAT
GACTATCAAAATATAAAATATTTAAGTCAATTTATTTGCCCAACTCCA
ATATATAATATTATAAATAGGATAGTTCTCAATTCCTAATAATTCAAAA
AATCATTA

In some embodiments, amplification of a THCA synthase gene sequence includes contacting nucleic acid from a *Cannabis* plant with a reverse THCA synthase primer that is complementary to a sequence that is 200-1000 nucleotides downstream of a THCA open reading frame. In some embodiments, a reverse THCA synthase primer is complementary to a sequence that is 200 to 800 nucleotides, 200 to 600 nucleotides, 200 to 400 nucleotides downstream of a THCA open reading frame. In some certain embodiments, a reverse THCA synthase primer is complementary to a sequence that is at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a portion of SEQ ID NO: 7. In some certain embodiments, a reverse THCA synthase primer is about 20 to 60 nucleotides long.

Exemplary *C. sativa* sequence downstream of THCA synthase open reading frame

SEQ ID NO: 7

TTATCAATTAGTTAATCATTATACCATACATCATTTTGTATATAGT
TTATCTACTCATATTATGTATGCTCCCAAGTATGAAAATCTACATTAGA
ACTGTGTAGACAATCATACATAGCGACTATCGTG

In some embodiments, amplification of a THCA synthase gene sequence includes contacting nucleic acid from a *Cannabis* plant with two or more primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to any of SEQ ID NOs: 8-12.

In some embodiments, a method of the present disclosure includes detecting the presence of a variant THCA synthase gene sequence in a sample that comprises nucleic acid from the *Cannabis* plant. In some embodiments, a method of the present disclosure includes amplifying a variant THCA synthase gene sequence or portion thereof from a sample that comprises nucleic acid from a *Cannabis* plant.

In some embodiments, a method of the present disclosure includes detecting the presence of a mutated THCA synthase gene sequence in a sample that comprises nucleic acid from the *Cannabis* plant. In some embodiments, a mutated THCA synthase gene sequence includes at least one mutation that alters the activity of a THCA synthase encoded by the mutated THCA synthase gene sequence relative to a THCA synthase encoded by a wild-type THCA synthase gene sequence.

In some embodiments, a method of the present disclosure includes detecting the presence of a polymorphism within a THCA synthase gene sequence in a sample that comprises nucleic acid from the *Cannabis* plant. In some embodiments, a polymorphism within a THCA synthase gene sequence is or comprises one or more variants as described in FIG. 3 and/or FIG. 10. FIG. 10 prevents a chart describing certain Single Nucleotide Polymorphism (SNP) variants of THCA synthase in different *Cannabis* plant strains (indicated by accession code). SNPs that result in an amino acid change are indicated in bold, and the type of change is with the corresponding position in the amino acid sequence is indicated in the bottom row. In the last column is provided the proportion of THC(V)A product of the total cannabinoid fraction that is accumulated in each *Cannabis* plant variant. In some embodiments, a THCA synthase gene sequence is or comprises a I63L variant, a V125L variant, a E236Q variant, a A250D variant, a E265G variant, and/or a G410E variant. In some certain embodiments, a THCA synthase gene sequence is or comprises a A411V variant.

In some embodiments, a method of the present disclosure includes detecting the presence of a wild-type cannabidiolic acid (CBDA^{WT}) synthase gene sequence in a sample that comprises nucleic acid from the *Cannabis* plant. In some embodiments, a method of the present disclosure includes amplifying a CBDA^{WT} synthase gene sequence or portion thereof from a sample that comprises nucleic acid from a *Cannabis* plant.

In some embodiments, a method of the present disclosure includes detecting the presence of a mutated cannabidiolic acid (CBDA^{mut}) synthase gene sequence in a sample that comprises nucleic acid from the *Cannabis* plant, wherein the mutated CBDA synthase gene sequence comprises a deletion mutation. In some embodiments, a CBDA^{del} sequence includes or comprises a deletion of CGTA (SEQ ID NO:3). In some embodiments, a method of the present disclosure includes amplifying a CBDA^{mut} synthase gene sequence or portion thereof from a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, a method of the present disclosure includes amplifying a CBDA^{del} synthase gene sequence or portion thereof from a sample that comprises nucleic acid from a *Cannabis* plant.

In some embodiments, a method of the present disclosure includes detecting a combination of two or more genomic sequences selected from: a THCA synthase gene sequence, a variant or mutant THCA synthase gene sequence, a CBDA^{WT} synthase, and CBDA^{mut} synthase gene sequence. In some embodiments, a method of the present disclosure includes amplifying two or more genomic sequences selected from: a THCA synthase gene sequence, a variant or mutant THCA synthase gene sequence, a CBDA^{WT} syn-

thase, and CBDA^{mut} synthase gene sequence, from a sample that comprises nucleic acid from a *Cannabis* plant.

In some embodiments, a method of the present disclosure includes detecting the presence of an olivetol synthase gene sequence in a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, a method of the present disclosure includes amplifying an olivetol synthase gene sequence or portion thereof from a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, a method of the present disclosure includes detecting the presence of a variant olivetol synthase gene sequence in a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, a method of the present disclosure includes amplifying a variant olivetol synthase gene sequence or portion thereof from a sample that comprises nucleic acid from a *Cannabis* plant.

In some embodiments, a method of the present disclosure includes detecting the presence of a divaric acid synthase gene sequence in a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, a method of the present disclosure includes amplifying a divaric acid synthase gene sequence or portion thereof from a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, a method of the present disclosure includes detecting the presence of a variant divaric acid synthase gene sequence in a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, a method of the present disclosure includes amplifying a variant divaric acid synthase gene sequence or portion thereof from a sample that comprises nucleic acid from a *Cannabis* plant.

In some embodiments, a method of the present disclosure includes detecting the presence of a limonene synthase gene sequence in a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, a method of the present disclosure includes amplifying a limonene synthase gene sequence or portion thereof from a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, a method of the present disclosure includes detecting the presence of a variant limonene synthase gene sequence in a sample that comprises nucleic acid from a *Cannabis* plant. In some embodiments, a method of the present disclosure includes amplifying a variant limonene synthase gene sequence or portion thereof from a sample that comprises nucleic acid from a *Cannabis* plant.

In some embodiments, a method of the present disclosure includes detecting a combination of two, three, four, five, six, seven, eight, or more genomic sequences selected from: a THCA synthase gene sequence, a variant or mutant THCA synthase gene sequence, a CBDA^{WT} synthase, and CBDA^{mut} synthase gene sequence. In some embodiments, a method of the present disclosure includes amplifying two or more genomic sequences selected from: a THCA synthase gene sequence, a variant or mutant THCA synthase gene sequence, a CBDA^{WT} synthase, a CBDA^{mut} synthase gene sequence, a olivetol synthase gene sequence, a variant olivetol synthase gene sequence, a divaric acid synthase gene sequence, a variant divaric acid synthase gene sequence, a limonene synthase gene sequence, and a variant limonene synthase gene sequence from a sample that comprises nucleic acid from the *Cannabis* plant.

Methods for Genotyping *Cannabis*

In vitro nucleic acid amplification technique can be used for genotyping *Cannabis* plants. In vitro nucleic acid amplification techniques can be grouped according to the temperature requirements of the procedure. For example, polymerase chain reaction (PCR) is the most popular method as a technique of amplifying nucleic acid in vitro. PCR has

high sensitivity based on the effect of exponential amplification. Further, since a PCR amplification product can be recovered as DNA, this method is applied widely for genetic engineering techniques such as gene sequence cloning and structural determination. In PCR, however, temperature cycling or a special temperature controller is necessary for practice; the exponential progress of the amplification reaction causes a problem in quantification. Other PCR-based amplification techniques include, for example, transcription-based amplification (D. Y. Kwok, et al. 1989. *Proc. Natl. Acad. Sci. USA* 86, 1173-1177, which is incorporated herein by reference), ligase chain reaction (LCR; D. Y. Wu, et al. 1989. *Genomics* 4, 560-569; K. Barringer, et al. 1990. *Gene* 89, 117-122; F. Barany. 1991. *Proc. Natl. Acad. Sci. USA* 88, 189-193, each of which are incorporated herein by reference), and restriction amplification (U.S. Pat. No. 5,102,784, which is incorporated herein by reference).

More recently, a number of isothermal nucleic acid amplification techniques have been developed. That is, these techniques do not rely on thermocycling to drive nucleic acid amplification. Isothermal amplification techniques typically utilize DNA polymerases with strand-displacement activity, thus eliminating the high temperature melt cycle that is required for PCR. This allows isothermal techniques to be faster and more energy efficient than PCR, and also allows for more simple and thus lower cost instrumentation since rapid temperature cycling is not required. For example, methods such as Strand Displacement Amplification (SDA; Walker, et al., (1992) *Proc. Natl. Acad. Sci. USA* 89: 392-396; Walker, et al., (1992) *Nuc. Acids. Res.* 20:1691-1696; U.S. Pat. No. 5,648,211 and EP 0 497 272, each of which is incorporated herein by reference); self-sustained sequence replication (3SR; J. C. Guatelli, et al., (1990) *Proc. Natl. Acad. Sci. USA*, 87: 1874-1878, which is incorporated herein by reference); and Q β replicase system (Lizardi, et al., (1988) *BioTechnology*, 6: 1197-1202, which is incorporated herein by reference) are isothermal reactions. See also, Nucleic Acid Isothermal Amplification Technologies—A Review. Nucleosides, Nucleotides and Nucleic Acids (2008) v27(3):224-243, which is incorporated herein by reference.

In some embodiments, an in vitro nucleic acid amplification assay for use in the methods of the present disclosure is an isothermal amplification method. Isothermal amplification methods include, for example, transcription mediated amplification (TMA) or self-sustained sequence replication (3SR), nucleic acid sequence-based amplification (NASBA), signal mediated amplification of RNA technology (SMART), strand displacement amplification (SDA), rolling circle amplification (RCA), loop-mediated isothermal amplification of DNA (LAMP), isothermal multiple displacement amplification (IMDA), helicase-dependent amplification (HDA), single primer isothermal amplification (SPIA), and circular helicase-dependent amplification (cHDA))

LAMP Assay

It is envisioned that LAMP amplification assays as described herein will provides a rapid and scalable means to resolve chemotypes of *Cannabis* plants. In some embodiments, an in vitro nucleic acid amplification assay for use in the methods of the present disclosure is Loop-Mediated Isothermal Amplification (LAMP). Typically, LAMP reactions use a strand-displacing DNA polymerase with four to six primers, which can result in exponential amplification of a target sequence.

In LAMP, a target sequence is amplified at a constant temperature. In some embodiments, a LAMP reaction temperature is between 37° C. and 75° C. In some embodiments,

a LAMP reaction temperature is between 50° C. and 70° C. In some embodiments, a LAMP reaction temperature is between 55° C. and 65° C. In some embodiments, a LAMP reaction temperature is between 60° C. and 65° C. In some embodiments, a LAMP reaction uses two or three primer sets, and a polymerase with high strand displacement activity in addition to a replication activity. (See Nagamine et al., (2002) Mol. Cell. Probes 16 (3): 223-9; and U.S. Pat. No. 6,410,278, each of which is incorporated herein by reference).

LAMP was originally invented and formulated as an isothermal amplification with a requirement for four primers: two loop-generating primers (FIP and BIP comprising F1, F2 and B1, B2 priming sites, correspondingly) and two “Displacement primers” (F3 and B3). In order to increase the speed of LAMP-based assays additional “Loop primers” were added in conjunction with the other primers used in LAMP, which resulted in significantly faster assays.

In some embodiments, LAMP amplification of a THCA synthase gene sequence includes contacting nucleic acid from a *Cannabis* plant with four or more primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to any of SEQ ID NOs: 8-12.

In some embodiments, LAMP amplification of a CBDA synthase gene sequence includes contacting nucleic acid from a *Cannabis* plant with four, five, or six, or more primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to any of SEQ ID NOs: 13-18 and 27.

In some embodiments, LAMP amplification of a CBDA^{mut} (CBDA^{del}) synthase gene sequence includes contacting nucleic acid from a *Cannabis* plant with four, five, or six, or more primers that are at least 70%, 75%, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to any of SEQ ID NOs: 13, 15-19, and 27.

In some embodiments, a LAMP amplification assay as described herein is a colorimetric LAMP assay.

Due to the specific nature of the action of these primers, the amount of DNA produced in LAMP is considerably higher than PCR based amplification. The reaction can be followed in real-time either by measuring the turbidity or by fluorescence using intercalating dyes. Dye molecules intercalate or directly label the DNA, and in turn can be correlated to the number of copies initially present. Hence, LAMP can also be quantitative. Thus, LAMP provides major advantages due to its simplicity, ruggedness, and low cost, and has the potential to be used as a simple screening assay in the field or at the point of care by clinicians.

Primer design for LAMP assays generally involves selection of eight separate regions of a target nucleic acid sequence (the FIP and BIP primers encompass two primer binding sites each), with BIP/FIP and Loop primers having significant restrictions on their positioning relative to each other. “Loop primers” are positioned between B2 and B1 sites and F2 and F1 sites, respectively, and must be oriented in a particular direction. Further, significant care must be taken in primer design to avoid primer-dimers between the six primers (which can be especially difficult as the FIP and BIP primers are generally greater than 40 nucleotides long). As a consequence, LAMP primer design is extremely challenging, especially when targeting highly polymorphic markers and sequences containing complex secondary structure. At least in part because primer design for LAMP is

subject to numerous constraints, software is generally used to assist with LAMP primer design.

LAMP has been observed to be less sensitive than PCR to inhibitors in complex samples such as blood, likely due to use of a different DNA polymerase (typically Bst DNA polymerase rather than Taq polymerase as in PCR). LAMP is useful primarily as a diagnostic or detection technique, but is generally not useful for some molecular biology applications enabled by PCR, such as, for example, cloning.

Also, multiplexing approaches for LAMP are relatively undeveloped. The larger number of primers per target in LAMP increases the likelihood of primer-primer interactions for multiplexed target sets. The product of LAMP is a series of concatemers of the target region, giving rise to a characteristic “ladder” or banding pattern on a gel, rather than a single band as with PCR. Although this is not a problem when detecting single targets with LAMP, “traditional” (endpoint) multiplex PCR applications wherein identity of a target is confirmed by size of a band on a gel are not feasible with LAMP. Multiplexing in LAMP has been achieved by choosing a target region with a restriction site, and digesting prior to running on a gel, such that each product gives rise to a distinct size of fragment, although this approach adds complexity to the experimental design and protocol. The use of a strand-displacing DNA polymerase in LAMP also precludes the use of hydrolysis probes, e.g. TaqMan probes, which rely upon the 5'-3' exonuclease activity of Taq polymerase.

In some embodiments, a modified LAMP technique called LAMP-STEM is used in a method of the present disclosure. LAMP-STEM system utilizes “Stem primers,” which are directed to the stem portion of the LAMP amplicon (or “dumbbell”). Stem primers can be used as an alternative to LAMP “Loop primers.” When used in addition to loop-generating and displacement primers, Stem primers offer similar benefits in speed and sensitivity to the Loop primers. (See Gandelman et al., Loop-Mediated Amplification Accelerated by Stem Primers. Int. J. Mol. Sci. 2011, v12:9108-9124, and US 2012/0157326, each of which is incorporated herein by reference). This beneficial effect of Stem primers is surprising as they do not bind to the single-stranded DNA loops, which define the very nature of the LAMP technology. Stem primers can be employed in either orientation, do not require either the B2/B1 or F2/F1 sites to be a specific distance apart, can be multiplexed, and allow the F1 and B1 sites to be positioned further from each other than in LAMP.

Stem primers significantly accelerate LAMP comprised of loop-generating and displacement primers only. They can be used on their own or synergistically with other Stem primers or even Loop primers. Addition of Stem primers into LAMP has a positive effect on both speed and sensitivity. In some cases they improve reproducibility at low copy number. The action of Stem primers can be rationalized via the proposed mechanism of LAMP. They anneal to transiently single-stranded regions of the amplicon and recopy the entire binding sites for the BIP/FIP primers. An additional unique feature is the extra strong intra-molecular self-priming when Stem primers delimit amplicon.

In general, positioning of Stem primers is less constrained than that of Loop primers. A rather challenging primer design involving selection of at least eight binding sites is thus simplified. Furthermore, Stem primers impose fewer limitations on the primer design in terms of stem length, orientation and distances between B1-B2 and F1-F2 sites. In contradiction to the postulated LAMP mechanism that relies on the involvement of displacement primers Stem primers can occasionally allow displacement primers not to be used

at all, though it is not clear why this is so. This has a major implication for primer design, as it allows the ability to omit one displacement primer or even both, if necessary.

qPCR

In some embodiments, an in vitro nucleic acid amplification assay for use in the methods of the present disclosure is qPCR. Real-time quantitative polymerase chain reaction (qPCR) determines, the fractional cycle number (C_t) at which the well's rising fluorescence (proportional to product formation) crosses a set threshold that is several standard deviations above the baseline fluorescence (Higuchi, et al., (1993) Kinetic PGR analysis: real-time monitoring of DNA amplification reactions, *Biotechnology* (NY), 11: 1026-1030, which is incorporated by reference in its entirety). The C_t versus log (amount of input target DNA) plot is linear, allowing relative quantitation of unknowns by comparison to a standard curve derived from amplifying, in the same plate, serial dilutions of a reference DNA sample.

In some embodiments, a qPCR assay is normalized, for example, a signal from a target sequence can be normalized to a signal from a reference sequence. In some embodiments, a target sequence and reference sequence are measured in separate (monoplex) reactions. In some embodiments, a reaction includes a dye that fluoresces upon intercalation into any double-stranded DNA, e.g., ethidium bromide or SYBR Green I, etc. In some embodiments, a target sequence and reference sequence are measured in the same reaction vessel via a multicolor multiplex qPCR. In some embodiments, a multiplex qPCR uses separate fluorescent dyes with distinct excitation/emission spectra for each of the DNA sequences being quantified (Wittwer, et al., (2001) Real-time multiplex PCR assays. *Methods*, 25, 430-442, which is incorporated by reference in its entirety).

Kits

In some embodiments, the present disclosure provides kits comprising materials useful for amplification and detection and/or sequencing of *Cannabis* plant nucleic acid (e.g., DNA). In some embodiments, *Cannabis* plant nucleic acid sample includes detection of all or part of a THCA synthase gene sequence and/or a CBDA synthase gene sequence as described herein. In some embodiments, a kit in accordance of the present disclosure is portable.

Suitable amplification reaction reagents that can be included in an inventive kit include, for example, one or more of: buffers; enzymes having polymerase activity; enzyme cofactors such as magnesium or manganese; salts; nicotinamide adenine dinucleotide (NAD); and deoxynucleoside triphosphates (dNTPs) such as, for example, deoxyadenosine triphosphate; deoxyguanosine triphosphate, deoxycytidine triphosphate and deoxythymidine triphosphate, biotinylated dNTPs, suitable for carrying out the amplification reactions.

In some embodiments, a kit comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more primer sequences for in vitro nucleic acid amplification. Primer sequences may be suitable for in vitro nucleic acid amplification with any of the methods described herein (e.g., QT-PCR, LAMP, etc.). In some embodiments, a kit of the present disclosure includes reagents suitable to perform a colorimetric LAMP assay for amplification of one or more *Cannabis* gene sequences as described herein.

Depending on the procedure, a kit may further include one or more of: wash buffers and/or reagents, hybridization buffers and/or reagents, labeling buffers and/or reagents, and detection means. The buffers and/or reagents included in a kit are preferably optimized for the particular amplification/detection technique for which a kit is intended. Protocols for

using these buffers and reagents for performing different steps of the procedure may also be included in a kit.

In some embodiments, a kit may further include one or more reagents for preparation of nucleic acid from a plant sample. For example, a kit may further include one or more of a lysis buffer, a DNA preparation solution (e.g., a solution for extraction and/or purification of DNA). Kits may also contain reagents for the isolation of nucleic acids from biological specimen prior to amplification. Protocols for using these reagents for performing different steps of the procedure may also be included in a kit.

Furthermore, kits may be provided with an internal control as a check on the amplification procedure and to prevent occurrence of false negative test results due to failures in the amplification procedure. An optimal control sequence is selected in such a way that it will not compete with the target nucleic acid sequence in the amplification reaction (as described above).

In some embodiments, a kit may further include reagents for an amplification assay to characterize the gender of a *Cannabis* plant.

Reagents may be supplied in a solid (e.g., lyophilized) or liquid form. Kits of the present disclosure may optionally comprise different containers (e.g., vial, ampoule, test tube, flask or bottle) for each individual buffer and/or reagent. In some embodiments, each component will generally be suitable as aliquoted in its respective container or provided in a concentrated form. Other containers suitable for conducting certain steps of inventive amplification/detection assay(s) may also be provided. Individual containers of a kit are preferably maintained in close confinement for commercial sale.

A kit may also comprise instructions for using the amplification reaction reagents, primer sets, primer/probe sets according to the present disclosure. Instructions for using a kit according to one or more methods of the present disclosure may comprise instructions for processing the biological sample, extracting nucleic acid molecules, and/or performing one or more amplification reactions; and/or instructions for interpreting results. In some embodiments, a kit may comprise instruction for determining, assessing and/or classifying a *Cannabis* plant used in the described methods as a Type Ia, Type Ib, Type IIa, Type IIb, Type IIc, Type IIIa, Type IIIb, Type IV or Type V plant.

Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

EXEMPLIFICATION

Example 1: Genotyping *Cannabis* Plants

Bt and Bd alleles have been proposed as a model for generating Type I, II, III, and IV plants. While this classification system can be predictive, the exact DNA sequences that govern them remain unknown. Single molecule sequencing was performed to identify mutations in gene sequences in cannabinoid synthesis pathway(s) (e.g., THCA synthase and CBDA synthase). This sequencing revealed a more refined structure of the inheritance and revealed numerous additional subtypes that can be identified by genotyping to predict chemical inheritance. See, FIG. 2. As illustrated in FIG. 2, a *Cannabis* plant may be characterized

as a Type Ia, Type Ib, Type IIa, Type IIb, Type IIc, Type IIIa, Type IIIb, Type IV or Type V plant.

Example 2: Assay for Genotyping THCA Synthase

This example describes development of superior isothermal nucleic acid amplification assays for detection of THCA synthase. Specifically, a colorimetric LAMP assay for detection of THCA synthase is described. It is envisioned that such a colorimetric LAMP assay may be portable, so that it can be performed at a point of grow to resolve *Cannabis* chemotypes in a rapid and scalable manner.

Challenges with genotyping a THCA synthase include, for example, (1) that there are many pseudogene sequence copies of THCA synthase and (2) that THCA synthase gene sequences are highly polymorphic. For at least this reason, designing primers can be difficult and testing such primers on a few samples does not provide adequate population diversity to guarantee assay performance in the broader markets. The present disclosure addresses these deficiencies with a novel Long Range (2130 bp) LAMP assay that includes a primer that anneals to a promoter region of THCA synthase and internal primers that are less prone to effects resulting from polymorphisms.

Weiblen, Stagginus and Onofri suggested primers for THCA synthase detection but failed to test enough samples to find fatal polymorphisms in their detection approach (Weiblen et al., (2015), *The New Phytologist*, PMID:

genetic studies of THCA synthase gene sequences. THCA synthase gene sequences are under selective breeding pressure and has a 4-fold higher polymorphism rate than the rest of the genome. There is a SNP approximately every 25 nucleotides in THCA synthase gene sequences, making primer design internal to a THCA synthase gene sequence complicated. To address this, primers (e.g., F3 and B3 primers for LAMP) were designed that were external to a THCA synthase gene sequence, where there is more sequence conservation. This results in a THCA synthase gene sequence amplicon of greater than 2000 nucleotides. There was no expectation that such a reaction would work, as there were no reports of Long Range LAMP assays in the literature and the common design tools found at Primer Explorer did not allow primer design of targets this large.

Materials and Methods

LAMP assay

10 ul 2xLAMP mix (NEB Catalog #M1800S)
2 ul of primer cocktail
6 ul ddH₂O (pH 6.5)
2 ul DNA (4 mm leaf biopsy boil in 100 ul ddH₂O, 5% Chelex)
Exemplary THCA synthase Long Range LAMP primers are provided in FIG. 9 and Table 2:

TABLE 2

THC_BIP	CACACAAGCACGTATTGGGCTTTAGTTTTCACCTTAACCT	SEQ ID NO: 8
THC_FIP	GACTCGCATGATTAGTTTTTCTATCCTTATGTGTCCCAAAATCC	SEQ ID NO: 9
THC_Loop1	TCCCTATAATTGAGATACGCCAAT	SEQ ID NO: 10
THC-F3	ATGGGAACCATAATAAACTATAAAAGTCATT	SEQ ID NO: 11
THC_B3	TATGATTGTCTACACAGTTCTAATGTAGATTTTC	SEQ ID NO: 12

26189495; Stagginus et al., (2014) *Journal of Forensic Sciences*, 59(4):919-26, PMID: 24579739; and Onofri et al., (2015), supra, each of which is incorporated by reference in its entirety). While Kitamura et al. described a portable LAMP assay for THCA synthase, the assay was performed using only two THCA+ cultivars and one fiber type. Consequently, Kitamura et al. did not consider the full complexity of *cannabis* genotypes in circulation. Moreover, the primers used in Kitamura's assay annealed to different regions in a THCA synthase gene sequence. As a result, and shown herein, the present disclosure recognizes that the Kitamura primers were insufficient for detecting/amplifying a THCA synthase gene sequence in numerous common *Cannabis* cultivars. In fact, sequence analysis of the Kitamura primers revealed that the 5' end of the Kitamura BIP primer sits on a common I63E and I63F polymorphisms described by Onofri et al., (2015), supra, which is incorporated by reference in its entirety. As this 5' end becomes the 3' end in the second strain synthesis of a LAMP reaction, presence of this polymorphism could impair or stall the LAMP reaction. A similar problem existed with other Kitamura primers, which traversed the P333R variant and the Glu265Gln variants described by Onofri, et al., (2015), supra, incorporated by reference in its entirety.

Amplification issues resulting from polymorphisms in an underlying gene sequence is a common problem with

To compare efficacy of Kitamura primers ("K primers") to exemplary Long Range primers ("LR primers"), LAMP assays were performed with these primers on various strains of *Cannabis* plants. Results are shown in FIG. 5, panel (A). In FIG. 5, panel (A), the text on the left side of the image indicates the strain of *Cannabis* plant for each row of samples that was tested by LAMP amplification. The top row included samples from Granddaddy Purple (GDP), the second row from the top included samples from Otto, the third row from the top included samples from a first hemp strain, the fourth row from the top included samples from a second hemp strain, the fifth row from the top included samples from Grape Stomper (RSP10516), and the bottom row included samples from Erk Train X Deadhead OG (RSP10517). A sample obtained from each strain was placed in the first and third columns. For each strain, a LAMP assay using the K primers was performed in the first column, and a LAMP assay using the LR primers was performed in the third column.

As shown in FIG. 5, panel (A), the two hemp strains displayed a negative result for THCA synthase using either the K primers or the LR primers. As a hemp strain would be expected to be THCA⁺/THCA⁻, this result showed that LAMP amplification using either the K primers or the LR primers was not giving a false positive result. The Otto

strain, which expresses THCA synthase, displayed a positive result using the K primers, which showed that LAMP amplification using the K primers detected the presence of a THCA synthase gene. However, the GDP strain of *Cannabis* plant, which also expresses THCA synthase, had a negative result when LAMP amplification was performed with K primers, while giving a positive result when LAMP amplification was performed with LR primers. This results showed that LR primers are able to successfully detect a THCA gene sequence by LAMP amplification, and do so in strains for which K primers are unable to detect a THCA gene sequence by LAMP amplification.

To determine how a Long Range LAMP assay of THCA synthase would perform at various temperatures, LAMP assays with LR primers were performed on various samples over a range of temperatures (temperatures indicated on right side of FIG. 5, panel (B)). As indicated in FIG. 5, panel (B), the first two columns of samples were obtained from THCA synthase negative hemp strains. The third column included samples from a THCA synthase positive *Cannabis* strain. The fourth column included samples from a Granddaddy purple strain. The eighth column included a negative control, and the ninth column included a positive control. Amplification of the THCA synthase gene sequence was successful for the THCA synthase positive *Cannabis* strain, the Granddaddy Purple strain, and the positive control at each of the temperatures tested. These data showed that LAMP assays using LR primers were able to properly detect THCA synthase positive strains across a broad temperature profile.

The exemplary Long Range LAMP assay of THCA synthase was repeated on various different *Cannabis* strains that are known to be THCA synthase positive. See, FIG. 6. FIG. 6, depicts LAMP amplifications with LR primers on 28 different known THCA synthase positive strains of *Cannabis* at 65° C., for 50 minutes. Wells H6-H9 were control samples. As can be seen, all 28 THCA synthase positive samples had successful amplification with the exemplary LR primers.

Accordingly, this example demonstrates that LAMP assays using long range primers (such as the exemplified LR primers) of THCA synthase gene sequences was robust and reliable. This example also demonstrates the shortcomings of LAMP amplification using the Kitamura primers for accurately detecting THCA synthase gene sequences in *Cannabis* strains.

Example 3: Combined THCA Synthase/CBDA Synthase Assay

This example describes characterization of the CBDA synthase Bd allele as a four nucleotide deletion within the coding sequence of this gene. This example also describes development of amplification assays for detection of wild-type (CBDA^{wt}) and mutant (CBDA^{Bd}) CBDA synthase gene sequences.

Kitamura simply assessed presence of a single gene sequence (THCA synthase) and its deleted pseudogene

sequence copies in certain hemp fiber varieties. Thus, the LAMP assay described in Kitamura (at best) only differentiated Type I or II *Cannabis* from other types (Type II-V), but failed to refine certain chemotype categories. To more rigorously stratify the numerous classes of cannabinoid production other gene sequence(s) that compete for the same cannabinoid precursor (e.g., cannabigerolic acid or CBGA) should be considered. Since both THCA synthase and CBDA synthase compete for cannabigerolic acid, the mutational spectrum of both of these gene sequences with respect to one another was examined.

To characterize CBDA synthase, single molecule sequencing of CBDA synthase in dozens of *Cannabis* cultivars was performed. This sequencing revealed a 4 nucleotide frame shifting deletion that governs the Bd allele. The Bd allele was then surveyed in more than 50 hemp and CBD lines to confirm the Bd allele and its phenotype. A consensus sequence for the region of the CBDA synthase gene sequence that includes this deletion is shown in FIG. 4.

Assays that targeted a 4 nucleotide deletion in the CBDA synthase gene sequence (4bpDel) (See, FIG. 4, residues 330-333) were designed. Various different hemp lines that were known to be THCA synthase negative were tested by both an exemplary CBDA synthase assay and an exemplary LAMP assay using LR primers for a THCA synthase gene sequence. FIG. 7 shows LAMP assays for amplification of wild-type CBDA synthase gene sequence (panel (A)) and THCA synthase gene sequence (panel (B)) on 38 different hemp strains (i.e., THCA synthase negative strains). A plate map describing the hemp strain in each well is provided at the bottom (panel (C)). As shown in FIG. 7, panel (A), all samples but one (sample M2 in well A1) resulted in amplification of a wild-type CBDA synthase gene sequence. Further, as shown in FIG. 7, panel (B), LAMP assay using LR primers for a THCA synthase gene sequence consistently showed no amplification on hemp samples. These results showed that the chemotype of *Cannabis* strains were accurately detected by performance of a CBDA synthase assay and an LAMP amplification using LR primers for a THCA synthase gene sequence.

For the one hemp sample (A1) that failed to amplify in the exemplary CBDA synthase gene sequence assay, when this sample was repeated using a different DNA purification method, amplification of CBDA synthase gene sequence was observed. Thus, it was concluded that the result for this particular sample was a false negative that resulted from other experimental conditions.

With two assays, discernment of THCA synthase positive plants from CBDA synthase positive plants was achieved. The copy number of THCA synthase or CBDA synthase in the genome of *Cannabis* plants could not be detected, however. Copy number can be useful information for breeders looking to breed plants that produce high levels of THC or CBD. In order to assess copy number, a deletion in the CBDA synthase gene sequence was targeted.

Exemplary CBDA^{wt} synthase primers (which hybridize to a wild-type CBDA synthase gene sequence) are provided in FIG. 9 and Table 3:

TABLE 3

CBD_BIP	CAATCTTAGATTACCTCTGACACAAGACATGTGAAGGAG	SEQ ID NO: 13
	TGAC	
CBD_FIP	TGTATTGTGCAATTTAGGACAGACAATGCAACAAATCTAA	SEQ ID NO: 14
	AACTCGTA	
CBD_Loop1	CAATGGGTTGTTTGTGAGTGT	SEQ ID NO: 15

TABLE 3-continued

CBD_Loop2	ACCCCAAACCACTTGT	SEQ ID NO: 16
CBD_F3	CTTCTCGCAATATATCCCAAT	SEQ ID NO: 17
CBD_F3_v2	ATGCTTCTCGCAATATATCCCA	SEQ ID NO: 27
CBD_B3	GCATAGAATAGTGCCTTGGAT	SEQ ID NO: 18

Exemplary CBDA^{del} synthase primers (which hybridize to a mutant CBDA synthase gene sequence including a 4 nucleotide deletion) are provided in FIG. 9 and Table 4:

TABLE 4

CBD_BIP	CAATCTTAGATTACCTCTGACACAAGACATGTGAAGGAG TGAC	SEQ ID NO: 13
CBD- NEG_FIP	TGTATTGTCGAATTTAGGACAGACAATGCAACAAATCTAA AACTTACA	SEQ ID NO: 19
CBD_Loop1	CAATGGGTGTTTTGAGTGT	SEQ ID NO: 15
CBD_Loop2	ACCCCAAACCACTTGT	SEQ ID NO: 16
CBD_F3	CTTCTCGCAATATATCCCAAT	SEQ ID NO: 17
CBD_F3_v2	ATGCTTCTCGCAATATATCCCA	SEQ ID NO: 27
CBD_B3	GCATAGAATAGTGCCTTGGAT	SEQ ID NO: 18

30

The present disclosure encompasses a recognition that detection of both wild-type and mutant CBDA gene sequences will enable genetic differentiation of strains, for example, to differentiate a CBDA⁻/CBDA⁺ genotype from a CBDA⁺/CBDA⁺ genotype.

FIG. 8 depicts results obtained from exemplary LAMP assays for amplification of CBDA^{wt} and CBDA^{del} (e.g., CBDA^{B^d}) gene sequences. FIG. 8, panel (A), depicts results obtained from LAMP assays with CBDA^{wt} primers on various hemp samples. Most of the samples resulted in amplification of a CBDA^{wt} gene sequence. FIG. 8, panel (B), depicts results from LAMP assays with CBDA^{del} primers on various hemp samples. As can be seen, most of these samples did not amplify a CBDA^{del} gene sequence.

The present disclosure encompasses a recognition that detection of both wild-type and mutant THCA gene sequences will enable genetic differentiation of strains. In order to assess THCA synthase copy number, one or more

sequence variants can be targeted. For example, primers can be used that target the *Cannabis* THCA synthase A411V variant.

Thus, combined THCA synthase/CBDA synthase assays provided herein can differentiate additional types of *Cannabis* plants. In some embodiments, methods in the context of the present disclosure can differentiate Type I, Type II, Type III, Type IV, and Type V plants. As described herein, the present disclosure further provides new sub-classification of plants we term Type Ia, Type Ib, Type IIa, Type IIb, Type IIc, Type IIIa, Type IIb, Type IV and Type V *Cannabis* plants, see, e.g., FIG. 2 and Table 1 (above).

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of an invention described herein. The scope of an invention described is not intended to be limited to the above Description, but rather is as set forth in the following claims:

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 27

<210> SEQ ID NO 1

<211> LENGTH: 2121

<212> TYPE: DNA

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 1

atgggaacca taataaacta taaaagtcac tatgtgtact tgctaccata ggcacctata 60

tccccaaaac tagctaccat agccaatttc tttttgtgtt ccaatatcca atttttattg 120

atgccaaaact attcaatgta caatgtacat ttattttcaa taagggtctc acctaacaaa 180

-continued

ggtgcctaatt tttagttgat ttatttttta tcacatgtga ctatttaattg actatcaaatt	240
tataaaatat ttaagtcaat ttatttgccc caactccaat atataaatatt ataaatagga	300
tagttctcaa ttctaataa ttcaaaaaat cattaggact gaagaaaaat gaattgctca	360
gcattttcct tttggtttgt ttgcaaaata atatttttct ttctctcatt ccatatccaa	420
atttcaatag ctaatcctcg agaaaacttc cttaaatgct tctcaaaaca tattcccaac	480
aatgtagcaa atccaaaact cgtatacact caacacgacc aattgtatat gtctatcctg	540
aattcgacaa tacaaaatct tagattcatc tctgatacaa ccccaaaacc actcggtatt	600
gtcactcctt caaataactc ccatatccaa gcaactattt tatgctctaa gaaagtggc	660
ttgcagattc gaactcgaag cggtgcccat gatgctgagg gtatgtccta catatctcaa	720
gtcccatttg ttgtagtaga cttgagaaac atgcattcga tcaaaataga tgttcatagc	780
caaactgcgt ggggtgaagc cggagctacc cttggagaag tttattattg gatcaatgag	840
aagaatgaga atcttagttt tcttggtggg tattgccta ctgttgccgt aggtggacac	900
tttagtggag gaggetatgg agcattgatg cgaaattatg gccttgccgc tgataatatt	960
attgatgcac acttagtcaa tgttgatgga aaagtcttag atcgaaaac catggggagaa	1020
gatctgtttt gggctatacg tgggtgtgga ggagaaaact ttggaatcat tgcagcatgg	1080
aaaatcaaac tgggtgatgt cccatcaaag tctactatat tcagtgttaa aaagaacatg	1140
gagatacatg ggcttgctaa gttatttaac aaatggcaaa atattgctta caagtatgac	1200
aaagatttag tactcatgac tcacttcata acaaagaata ttacagataa tcatgggaag	1260
aataagacta cagtacatgg ttacttctct tcaatttttc atggtggagt ggatagtcta	1320
gtcgacttga tgaacaagag ctttctctgag ttgggtatta aaaaaactga ttgcaagaa	1380
tttagctgga ttgatacaac catctctac agtgggtgtg taaattttaa cactgctaatt	1440
tttaaaaagg aaattttgct tgatagatca gctgggaaga agacggcttt ctcaattaag	1500
ttagactatg ttaagaaacc aattccagaa actgcaatgg tcaaaatttt ggaaaaatta	1560
tatgaagaag atgtaggagc tgggggtgtg tacccttacg gtggtataat ggaggagatt	1620
tcagaatcag caattccatt cctcatcga gctggaataa tgtatgaact ttggtacact	1680
gcttcctggg agaagcaaga agataatgaa aagcatataa actgggttcg aagtgtttat	1740
aattttaca ctccttatgt gtcccaaat ccaagattgg cgtatctcaa ttataggac	1800
cttgatttag gaaaaactaa tcatgcgagt cctaataatt acacacaagc acgtatttgg	1860
ggtgaaaagt attttggtaa aaattttaac aggttagtta aggtgaaaac taaagttag	1920
cccaataatt tttttagaaa cgaacaaagt atcccacctc ttccaccgca tcatcattaa	1980
ttatctttaa atagatatat ttcccttacc aattagttaa tcattatacc atacatacat	2040
ttattgtata tagtttatct actcatatta tgtatgctcc caagtatgaa aatctacatt	2100
agaactgtgt agacaatcat a	2121

<210> SEQ ID NO 2

<211> LENGTH: 1941

<212> TYPE: DNA

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 2

gatatatatc tcacacggat gcacctaaca atgatgccta atttttgtga atttttttta	60
ccacatgact taatgatatc aaattatgaa atatttagtt aatttatttg cccctgctcc	120
aatatataaa gctataaata ggatagttct taatccatag taattcaaaa ttcattagaa	180

-continued

```

ctaaagaaaa atgaagtgtc caacattctc cttttgggtt gtttgcaaga taatattttt 240
ctttttctca ttcaatatcc aaacttccat tgctaatacct cgagaaaact tccttaaatg 300
cttctcgcaa tatattccca ataatgcaac aaatctaaaa ctcttataca ctcaaaacaa 360
cccattgtat atgtctgtcc taaattcgac aatacacaaat cttagattca cctctgacac 420
aaccceaaaa ccacttggtt tctgctactcc ttcacatgtc tctcatatcc aaggcactat 480
tctatgctcc aagaaagtgt gcttgccgat tcgaactcga agtggtgggtc atgattctga 540
gggcatgtcc tacatatctc aagtcctatt tgttatagta gacttgagaa acatgcgttc 600
aatcaaaata gatgttcata gccaaactgc atgggttgaa gccggagcta cccttgagaa 660
agtttattat tgggttaaat agaaaaatga gaactctagt ttggcggctg ggtattgccc 720
tactgtttgc gcagggtggc actttgggtg aggaggctat ggaccattga tgagaaacta 780
tggcctcgcg gctgataata tcattgatgc acacttagtc aacgttcatg gaaaagtgtc 840
agatcgaaaa tctatggggg aagatctctt ttgggcttta cgtgggtgtg gagcagaaa 900
cttcggaatc attgtagcat ggaaaattag actggttgcg gtcccaaagt ctactatgtt 960
tagtgtaaaa aagatcatgg agatacatga gcttgtcaag ttagttaaca aatggcaaaa 1020
tattgcttac aagtatgaca aagatttatt actcatgact cacttcataa ctaggaaat 1080
tacagataat caagggaaga ataagacagc aatacacact tacttctctt cagttttcct 1140
tggtggagtg gatagcttag tgcactgat gaacaagagt tttcctgagt tgggtattaa 1200
aaaaacggat tgcagacaat tgagctggat tgatactatc atcttctata gtggtgttgt 1260
aaattacgac actgataatt ttaacaagga aattttgctt gatagatccg ctgggcagaa 1320
cgggtgcttc aagattaagt tagactacgt taagaaacca attccagaat ctgtatttgt 1380
ccaaattttg gaaaaattat atgaagaaga tataggagct gggatgtatg cgttgtacct 1440
ttacgggtgt ataattgatg agatttcaga atcagcaatt ccattccctc atcgagctgg 1500
aatcttgatg gagttatggt acatatgtag ttgggagaag caagaagata acgaaaagca 1560
tctaaactgg attagaaata tttataactt catgactcct tatgtgtcca aaaatccaag 1620
attggcatat ctcaattata gagaccttga tataggaata aatgatccca agaatccaaa 1680
taattacaca caagcacgta tttgggggtg gaagtatttt ggtaaaaatt ttgacaggct 1740
agtaaaaagt aaaaccctgg ttgatcccaa taactttttt agaaacgaac aaagcatccc 1800
acctcttcca cggcatcgtc attaatgatc ttaaatagat ctttttctct tattaattag 1860
tccttataat atacatatat tgattatata tataaaaaata gtttgtccgg gtgtactgtg 1920
tatgcgatat atatctcaca c 1941

```

```

<210> SEQ ID NO 3
<211> LENGTH: 4
<212> TYPE: DNA
<213> ORGANISM: Cannabis sativa

```

```

<400> SEQUENCE: 3

```

```

cgta

```

4

```

<210> SEQ ID NO 4
<211> LENGTH: 1158
<212> TYPE: DNA
<213> ORGANISM: Cannabis sativa

```

```

<400> SEQUENCE: 4

```


-continued

atgaatcacc	ttcgtgctga	gggtccggcc	tccgttctcg	ccattggcac	cgccaatccg	60
gagaacattt	tattacaaga	tgagtttcct	gactactatt	ttcgcgtcac	caaaagtga	120
cacatgactc	aactcaaaga	aaagtttcga	aaaatatgtg	acaaaagtat	gataaggaaa	180
cgtaactgtt	tcttaaatga	agaacaccta	aagcaaaacc	caagattggg	ggagcacgag	240
atgcaaaact	tgatgcacg	tcaagacatg	ttggtagttg	aggttccaaa	acttgggaa	300
gatgcttggt	caaaggccat	caaagaatgg	ggtcaaccga	agtctaaaat	cactcattta	360
atcttcacta	gcgcatcaac	cactgacatg	cccgggtgcg	actaccattg	cgctaagctt	420
ctcggactga	gtccctcagt	gaagcgtgtg	atgatgtatc	aactaggctg	ttatggtggt	480
ggaaccgttc	tacgcattgc	caaggacata	gcagagaata	acaaaggcgc	acgagttctc	540
gcccgtgtgt	gtgacataat	ggcttgcttg	tttcgtgggc	cttcagagtc	tgacctcgaa	600
ttactagtgg	gacaagctat	ctttggtgat	ggggctgctg	cggtgattgt	tgagactgaa	660
cccgatgagt	cagttgggga	aaggccgata	tttgagttgg	tgtaactggg	gcaacaatc	720
ttaccaaact	cggaaggaa	tattggggga	catataagg	aagcaggact	gatatttgat	780
ttacataagg	atgtgcctat	gttgatctct	aataatattg	agaaatgttt	gattgaggca	840
tttactccta	ttgggattag	tgattggaac	tccatatttt	ggattacaca	cccaggtggg	900
aaagctat	ttggacaaagt	ggaggagaag	ttgcatctaa	agagtataa	gtttgtggat	960
tcacgtcatg	tgctgagtga	gcattgggaat	atgtctagct	caactgtctt	gtttgttatg	1020
gatgagttga	ggaagaggtc	gttgaggagaa	gggaagtcta	ccactggaga	tggaattgag	1080
tggtgtgttc	tttttgggtt	tgaccagggt	ttgactgtcg	aaagagtggg	cgtgcgtagt	1140
gttcccatca	aatattaa					1158

<210> SEQ ID NO 5

<211> LENGTH: 1869

<212> TYPE: DNA

<213> ORGANISM: Cannabis sativa

<400> SEQUENCE: 5

atgcagtgc	tagcttttca	ccaatttgct	tcatcatcat	ccctccctat	ttggagtagt	60
attgataatc	gttttacacc	aaaaacttct	attacttcta	tttcaaaacc	aaaaccaaaa	120
ctaaaatcaa	aatcaaacct	gaaatcgaga	tcgagatcaa	gtacttgcta	ctccatacaa	180
tgtactgtgg	tcgataaacc	tagttctacg	attactaata	atagtgtatg	aagatcagcc	240
aactatggac	ctcccatctt	gtcttttgat	ttgtttcaat	ctcttccaat	ccaatataag	300
ggtgaatctt	atacaagtgc	attaaataag	ttggagaaa	atgtgaaaag	gatgctaatt	360
ggagtggaaa	actcttttagc	ccaacttgaa	ctaattgata	caatacaaa	acttggaata	420
tcttatcggt	ttgaaaatga	aatcatttct	atgttgaaag	aaaaattcac	caataataat	480
gacaacccta	atcctaatta	tgatttatat	gctactgtct	tccaatttag	gcttctacgc	540
caatatggat	ttgaagtacc	tcaagaaatt	ttcaataatt	ttaaaaatca	caagacagga	600
gagttcaagg	caaatataag	taatgatatt	atgggagcat	tggtgttata	tgaagcttca	660
ttccatggga	aaaagggtga	aagtattttg	gaagaagcaa	gaattttcac	aacaaaatgt	720
ctcaaaaaat	acaaattaat	gtcaagtagt	aataataata	atatgacatt	aatatcatta	780
ttagtgaatc	atgcttttga	gatgccactt	caatggagaa	tcacaagatc	agaagctaaa	840
tggtttattg	aagaaatata	tgaagaaaa	caagacatga	atccaacttt	acttgagttt	900
gccaaattgg	atttcaatat	gctgcaatca	acatatcaag	aggagctcaa	agtactctct	960

-continued

```

aggtggtgga aggattctaa acttgagag aaattgcctt tcgttagaga tagattggtg 1020
gagtgtttct tatggcaagt tggagtaaga tttgagccac aattcagtta ctttagaata 1080
atggatacaa aactctatgt tctattaaca ataattgatg atatgcatga ctttatgga 1140
acattggagg aactacaact ttccactaat gctcttcaaa gatgggattt gaaagaatta 1200
gataaattac cagattatat gaagacagct ttctacttta catacaattt cacaaatgaa 1260
ttggcatttg atgtattaca agaacatggt tttgttcaca ttgaatactt caagaaactg 1320
atggtagagt tgtgtaaaca tcatttgcaa gaggcaaaat ggttttatag tggatacaaa 1380
ccaacattgc aagaatatgt tgagaatgga tggttgtctg tgggaggaca agttattctt 1440
atgcatgcat atttcgcttt taaaaatcct gttaccaaag aggcattgga atgtctaaaa 1500
gacggtcatc ctaacatagt tcgccatgca tcgataatat tacgacttgc agatgatcta 1560
ggaacattgt cggatgaact gaaaagaggc gatgttccta aatcaattca atgttatatg 1620
cacgatactg gtgcttctga agatgaagct cgtgagcaca taaaatattt aataagtgaa 1680
tcatggaagg agatgaataa tgaagatgga aatattaact cttttttctc aaatgaattt 1740
gttcaagttt gccaaaatct tggtagagcg tcacaattca tataccagta tggcgatgga 1800
catgcttctc agaataatct atcgaaagag cgcgttttag gggtgattat tactcctatc 1860
cccatgtaa 1869

```

```

<210> SEQ ID NO 6
<211> LENGTH: 351
<212> TYPE: DNA
<213> ORGANISM: Cannabis sativa

```

```

<400> SEQUENCE: 6
catagcgact atcgatgagg gaaccataat aaactataaa agtcattatg tgtacttgct 60
accataggca cctatatccc acaaaactagc taccatagcc aatttctttt ttgtttccaa 120
tatccaattt ttattgatgc caaactattc aatgtacaat gtacatttat ttccaataag 180
ggcttcacct aacaaagggtg cctaatttta gttgatttat tttttatcac atgtgactat 240
ttaatgacta tcaaattata aaatatttaa gtcaatttat ttgcccacac tccaatatat 300
aatattataa ataggatagt tctcaattcc taataattca aaaaatcatt a 351

```

```

<210> SEQ ID NO 7
<211> LENGTH: 132
<212> TYPE: DNA
<213> ORGANISM: Cannabis sativa

```

```

<400> SEQUENCE: 7
ttatcaatta gttaatcatt ataccatata tacatttatt gtatatagtt tatctactca 60
tattatgtat gctcccaagt atgaaaatct acattagaac tgtgtagaca atcacacata 120
gcgactatcg tg 132

```

```

<210> SEQ ID NO 8
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

```

```

<400> SEQUENCE: 8
cacacaagca cgtatttggg ctttagtttt caccttaact aacct 45

```

-continued

<210> SEQ ID NO 9
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 9

gactcgcatg attagttttt cctatcotta tgtgtcccaa aatcc 45

<210> SEQ ID NO 10
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 10

tcctcataat tgagatacgc caat 24

<210> SEQ ID NO 11
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 11

atgggaacca taataaacta taaaagtc t 31

<210> SEQ ID NO 12
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 12

tatgattgtc tacacagttc taatgtagat tttc 34

<210> SEQ ID NO 13
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 13

caatcttaga ttcacctctg acacaagaca tgtgaaggag tgac 44

<210> SEQ ID NO 14
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 14

tgtattgtcg aatttaggac agacaatgca acaaatctaa aactogta 48

<210> SEQ ID NO 15
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

-continued

```

<400> SEQUENCE: 15
caatgggttg ttttgagtgt                20

<210> SEQ ID NO 16
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 16
accccaaaac cacttgt                    17

<210> SEQ ID NO 17
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 17
cttctcgcaa tatattccca at             22

<210> SEQ ID NO 18
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 18
gcatagaata gtgccttgga t              21

<210> SEQ ID NO 19
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 19
tgtattgtcg aatttaggac agacaatgca acaaatctaa aacttaca        48

<210> SEQ ID NO 20
<211> LENGTH: 2153
<212> TYPE: DNA
<213> ORGANISM: Cannabis sp.
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (365)..(1993)

<400> SEQUENCE: 20
catagcgact atcgtgatgg gaaccataat aaactataaa agtcattatg tgtacttgct    60
accataggca cctatatccc acaaaactagc taccatagcc aatttctttt ttgtttccaa    120
tatccaattt ttattgatgc caaactattc aatgtacaat gtacatttat tttcaataag    180
ggcttcacct aacaaagggtg cctaatttta gttgatttat tttttatcac atgtgactat    240
ttaatgacta tcaaatataa aaatatttaa gtcaatttat ttgcccacac tccaatatat    300
aatattataa ataggtagtg tctcaattcc taataattca aaaaatcatt aggactgaag    360
aaaa atg aat tgc tca gca ttt tcc ttt tgg ttt gtt tgc aaa ata ata    409
      Met Asn Cys Ser Ala Phe Ser Phe Trp Phe Val Cys Lys Ile Ile
      1           5           10          15

```

-continued

ttt ttc ttt ctc tca ttc cat atc caa att tca ata gct aat cct cga	457
Phe Phe Phe Leu Ser Phe His Ile Gln Ile Ser Ile Ala Asn Pro Arg	
20 25 30	
gaa aac ttc ctt aaa tgc ttc tca aaa cat att ccc aac aat gta gca	505
Glu Asn Phe Leu Lys Cys Phe Ser Lys His Ile Pro Asn Asn Val Ala	
35 40 45	
aat cca aaa ctc gta tac act caa cac gac caa ttg tat atg tct atc	553
Asn Pro Lys Leu Val Tyr Thr Gln His Asp Gln Leu Tyr Met Ser Ile	
50 55 60	
ctg aat tcg aca ata caa aat ctt aga ttc atc tct gat aca acc cca	601
Leu Asn Ser Thr Ile Gln Asn Leu Arg Phe Ile Ser Asp Thr Thr Pro	
65 70 75	
aaa cca ctc gtt att gtc act cct tca aat aac tcc cat atc caa gca	649
Lys Pro Leu Val Ile Val Thr Pro Ser Asn Asn Ser His Ile Gln Ala	
80 85 90 95	
act att tta tgc tct aag aaa gtt ggc ttg cag att cga act cga agc	697
Thr Ile Leu Cys Ser Lys Lys Val Gly Leu Gln Ile Arg Thr Arg Ser	
100 105 110	
ggt ggc cat gat gct gag ggt atg tcc tac ata tct caa gtc cca ttt	745
Gly Gly His Asp Ala Glu Gly Met Ser Tyr Ile Ser Gln Val Pro Phe	
115 120 125	
gtt gta gta gac ttg aga aac atg cat tcg atc aaa ata gat gtt cat	793
Val Val Val Asp Leu Arg Asn Met His Ser Ile Lys Ile Asp Val His	
130 135 140	
agc caa act gcg tgg gtt gaa gcc gga gct acc ctt gga gaa gtt tat	841
Ser Gln Thr Ala Trp Val Glu Ala Gly Ala Thr Leu Gly Glu Val Tyr	
145 150 155	
tat tgg atc aat gag aag aat gag aat ctt agt ttt cct ggt ggg tat	889
Tyr Trp Ile Asn Glu Lys Asn Glu Asn Leu Ser Phe Pro Gly Gly Tyr	
160 165 170 175	
tgc cct act gtt ggc gta ggt gga cac ttt agt gga gga ggc tat gga	937
Cys Pro Thr Val Gly Val Gly Gly His Phe Ser Gly Gly Gly Tyr Gly	
180 185 190	
gca ttg atg cga aat tat ggc ctt gcg gct gat aat att att gat gca	985
Ala Leu Met Arg Asn Tyr Gly Leu Ala Ala Asp Asn Ile Ile Asp Ala	
195 200 205	
cac tta gtc aat gtt gat gga aaa gtt cta gat cga aaa tcc atg gga	1033
His Leu Val Asn Val Asp Gly Lys Val Leu Asp Arg Lys Ser Met Gly	
210 215 220	
gaa gat ctg ttt tgg gct ata cgt ggt ggt gga gga gaa aac ttt gga	1081
Glu Asp Leu Phe Trp Ala Ile Arg Gly Gly Gly Gly Glu Asn Phe Gly	
225 230 235	
atc att gca gca tgg aaa atc aaa ctg gtt gat gtc cca tca aag tct	1129
Ile Ile Ala Ala Trp Lys Ile Lys Leu Val Asp Val Pro Ser Lys Ser	
240 245 250 255	
act ata ttc agt gtt aaa aag aac atg gag ata cat ggg ctt gtc aag	1177
Thr Ile Phe Ser Val Lys Lys Asn Met Glu Ile His Gly Leu Val Lys	
260 265 270	
tta ttt aac aaa tgg caa aat att gct tac aag tat gac aaa gat tta	1225
Leu Phe Asn Lys Trp Gln Asn Ile Ala Tyr Lys Tyr Asp Lys Asp Leu	
275 280 285	
gta ctc atg act cac ttc ata aca aag aat att aca gat aat cat ggg	1273
Val Leu Met Thr His Phe Ile Thr Lys Asn Ile Thr Asp Asn His Gly	
290 295 300	
aag aat aag act aca gta cat ggt tac ttc tct tca att ttt cat ggt	1321
Lys Asn Lys Thr Thr Val His Gly Tyr Phe Ser Ser Ile Phe His Gly	
305 310 315	
gga gtg gat agt cta gtc gac ttg atg aac aag agc ttt cct gag ttg	1369
Gly Val Asp Ser Leu Val Asp Leu Met Asn Lys Ser Phe Pro Glu Leu	

-continued

320	325	330	335	
ggt att aaa aaa act gat tgc aaa gaa ttt agc tgg att gat aca acc				1417
Gly Ile Lys Lys Thr Asp Cys Lys Glu Phe Ser Trp Ile Asp Thr Thr	340	345	350	
atc ttc tac agt ggt gtt gta aat ttt aac act gct aat ttt aaa aag				1465
Ile Phe Tyr Ser Gly Val Val Asn Phe Asn Thr Ala Asn Phe Lys Lys	355	360	365	
gaa att ttg ctt gat aga tca gct ggg aag aag acg gct ttc tca att				1513
Glu Ile Leu Leu Asp Arg Ser Ala Gly Lys Lys Thr Ala Phe Ser Ile	370	375	380	
aag tta gac tat gtt aag aaa cca att cca gaa act gca atg gtc aaa				1561
Lys Leu Asp Tyr Val Lys Lys Pro Ile Pro Glu Thr Ala Met Val Lys	385	390	395	
att ttg gaa aaa tta tat gaa gaa gat gta gga gct ggg gtg ttg tac				1609
Ile Leu Glu Lys Leu Tyr Glu Glu Asp Val Gly Ala Gly Val Leu Tyr	400	405	410	415
cct tac ggt ggt ata atg gag gag att tca gaa tca gca att cca ttc				1657
Pro Tyr Gly Gly Ile Met Glu Glu Ile Ser Glu Ser Ala Ile Pro Phe	420	425	430	
cct cat cga gct gga ata atg tat gaa ctt tgg tac act gct tcc tgg				1705
Pro His Arg Ala Gly Ile Met Tyr Glu Leu Trp Tyr Thr Ala Ser Trp	435	440	445	
gag aag caa gaa gat aat gaa aag cat ata aac tgg gtt cga agt gtt				1753
Glu Lys Gln Glu Asp Asn Glu Lys His Ile Asn Trp Val Arg Ser Val	450	455	460	
tat aat ttt acg act cct tat gtg tcc caa aat cca aga ttg gcg tat				1801
Tyr Asn Phe Thr Thr Pro Tyr Val Ser Gln Asn Pro Arg Leu Ala Tyr	465	470	475	
ctc aat tat agg gac ctt gat tta gga aaa act aat cat gcg agt cct				1849
Leu Asn Tyr Arg Asp Leu Asp Leu Gly Lys Thr Asn His Ala Ser Pro	480	485	490	495
aat aat tac aca caa gca cgt att tgg ggt gaa aag tat ttt ggt aaa				1897
Asn Asn Tyr Thr Gln Ala Arg Ile Trp Gly Glu Lys Tyr Phe Gly Lys	500	505	510	
aat ttt aac agg tta gtt aag gtg aaa act aaa gtt gat ccc aat aat				1945
Asn Phe Asn Arg Leu Val Lys Val Lys Thr Lys Val Asp Pro Asn Asn	515	520	525	
ttt ttt aga aac gaa caa agt atc cca cct ctt cca ccg cat cat cat				1993
Phe Phe Arg Asn Glu Gln Ser Ile Pro Pro Leu Pro Pro His His His	530	535	540	
taattatctt taaatagata tatttccctt atcaattagt taatcattat accatacata				2053
catttattgt atatagttta tctactcata ttatgtatgc tcccaagtat gaaaactac				2113
attagaactg tgtagacaat catacatagc gactatcg				2153

<210> SEQ ID NO 21

<211> LENGTH: 543

<212> TYPE: PRT

<213> ORGANISM: Cannabis sp.

<400> SEQUENCE: 21

Met Asn Cys Ser Ala Phe Ser Phe Trp Phe Val Cys Lys Ile Ile Phe
1 5 10 15

Phe Phe Leu Ser Phe His Ile Gln Ile Ser Ile Ala Asn Pro Arg Glu
20 25 30

Asn Phe Leu Lys Cys Phe Ser Lys His Ile Pro Asn Asn Val Ala Asn
35 40 45

Pro Lys Leu Val Tyr Thr Gln His Asp Gln Leu Tyr Met Ser Ile Leu
50 55 60

-continued

Asn	Ser	Thr	Ile	Gln	Asn	Leu	Arg	Phe	Ile	Ser	Asp	Thr	Thr	Pro	Lys	65	70	75	80
Pro	Leu	Val	Ile	Val	Thr	Pro	Ser	Asn	Asn	Ser	His	Ile	Gln	Ala	Thr	85	90	95	
Ile	Leu	Cys	Ser	Lys	Lys	Val	Gly	Leu	Gln	Ile	Arg	Thr	Arg	Ser	Gly	100	105	110	
Gly	His	Asp	Ala	Glu	Gly	Met	Ser	Tyr	Ile	Ser	Gln	Val	Pro	Phe	Val	115	120	125	
Val	Val	Asp	Leu	Arg	Asn	Met	His	Ser	Ile	Lys	Ile	Asp	Val	His	Ser	130	135	140	
Gln	Thr	Ala	Trp	Val	Glu	Ala	Gly	Ala	Thr	Leu	Gly	Glu	Val	Tyr	Tyr	145	150	155	160
Trp	Ile	Asn	Glu	Lys	Asn	Glu	Asn	Leu	Ser	Phe	Pro	Gly	Gly	Tyr	Cys	165	170	175	
Pro	Thr	Val	Gly	Val	Gly	Gly	His	Phe	Ser	Gly	Gly	Gly	Tyr	Gly	Ala	180	185	190	
Leu	Met	Arg	Asn	Tyr	Gly	Leu	Ala	Ala	Asp	Asn	Ile	Ile	Asp	Ala	His	195	200	205	
Leu	Val	Asn	Val	Asp	Gly	Lys	Val	Leu	Asp	Arg	Lys	Ser	Met	Gly	Glu	210	215	220	
Asp	Leu	Phe	Trp	Ala	Ile	Arg	Gly	Gly	Gly	Gly	Glu	Asn	Phe	Gly	Ile	225	230	235	240
Ile	Ala	Ala	Trp	Lys	Ile	Lys	Leu	Val	Asp	Val	Pro	Ser	Lys	Ser	Thr	245	250	255	
Ile	Phe	Ser	Val	Lys	Lys	Asn	Met	Glu	Ile	His	Gly	Leu	Val	Lys	Leu	260	265	270	
Phe	Asn	Lys	Trp	Gln	Asn	Ile	Ala	Tyr	Lys	Tyr	Asp	Lys	Asp	Leu	Val	275	280	285	
Leu	Met	Thr	His	Phe	Ile	Thr	Lys	Asn	Ile	Thr	Asp	Asn	His	Gly	Lys	290	295	300	
Asn	Lys	Thr	Thr	Val	His	Gly	Tyr	Phe	Ser	Ser	Ile	Phe	His	Gly	Gly	305	310	315	320
Val	Asp	Ser	Leu	Val	Asp	Leu	Met	Asn	Lys	Ser	Phe	Pro	Glu	Leu	Gly	325	330	335	
Ile	Lys	Lys	Thr	Asp	Cys	Lys	Glu	Phe	Ser	Trp	Ile	Asp	Thr	Thr	Ile	340	345	350	
Phe	Tyr	Ser	Gly	Val	Val	Asn	Phe	Asn	Thr	Ala	Asn	Phe	Lys	Lys	Glu	355	360	365	
Ile	Leu	Leu	Asp	Arg	Ser	Ala	Gly	Lys	Lys	Thr	Ala	Phe	Ser	Ile	Lys	370	375	380	
Leu	Asp	Tyr	Val	Lys	Lys	Pro	Ile	Pro	Glu	Thr	Ala	Met	Val	Lys	Ile	385	390	395	400
Leu	Glu	Lys	Leu	Tyr	Glu	Glu	Asp	Val	Gly	Ala	Gly	Val	Leu	Tyr	Pro	405	410	415	
Tyr	Gly	Gly	Ile	Met	Glu	Glu	Ile	Ser	Glu	Ser	Ala	Ile	Pro	Phe	Pro	420	425	430	
His	Arg	Ala	Gly	Ile	Met	Tyr	Glu	Leu	Trp	Tyr	Thr	Ala	Ser	Trp	Glu	435	440	445	
Lys	Gln	Glu	Asp	Asn	Glu	Lys	His	Ile	Asn	Trp	Val	Arg	Ser	Val	Tyr	450	455	460	
Asn	Phe	Thr	Thr	Pro	Tyr	Val	Ser	Gln	Asn	Pro	Arg	Leu	Ala	Tyr	Leu	465	470	475	480

-continued

Asn Tyr Arg Asp Leu Asp Leu Gly Lys Thr Asn His Ala Ser Pro Asn
485 490 495

Asn Tyr Thr Gln Ala Arg Ile Trp Gly Glu Lys Tyr Phe Gly Lys Asn
500 505 510

Phe Asn Arg Leu Val Lys Val Lys Thr Lys Val Asp Pro Asn Asn Phe
515 520 525

Phe Arg Asn Glu Gln Ser Ile Pro Pro Leu Pro Pro His His His
530 535 540

<210> SEQ ID NO 22

<211> LENGTH: 1635

<212> TYPE: DNA

<213> ORGANISM: Cannabis sp.

<400> SEQUENCE: 22

```

atgaattgct cagcattttc cttttggttt gtttgcaaaa taatattttt ctttctctca    60
ttcaatatcc aaatttcaat agctaactct caagaaaact tccttaaattg cttctcggaa    120
tatattccta acaatccagc aaatccaaaa ttcataatac ctcaacacga ccaattgtat    180
atgtctgtcc tgaattcgac aatacaaaat cttagattca cctctgatac aacccccaaa    240
ccactcggtt ttgtactccc ttcaaatgtc tcccatatcc aggccagtat tctctgctcc    300
aagaaagttg gtttgcagat tcgaactcga agcgggtggc atgatgctga gggtttgtcc    360
tacatatctc aagtoecatt tgctatagta gacttgagaa acatgcatac ggtcaaagta    420
gatattcata gccaaactgc gtgggttgaa gccggagcta cccttgaga agtttattat    480
tggatcaatg agatgaatga gaattttagt ttctctggtg ggtattgccc tactgttggc    540
gtaggtggac acttttagtg aggaggtcat ggagcattga tgcgaaatta tggccttgcg    600
gctgataata tcattgatgc acacttagtc aatgttgatg gaaaagttct agatcgaaaa    660
tccatgggag aagatctatt ttgggctata cgtggtggag gaggagaaaa ctttggaatc    720
attgcagcat ggaaaatcaa acttggtgtt gtcccatcaa aggctactat attcagtgtt    780
aaaaagaaca tggagataca tgggcttgtc aagttattta acaaatggca aaatattgct    840
tacaagtatg acaagattt aatgctcacg actcaactca gaactaggaa tattacagat    900
aatcatggga agaataagac tacagtacat ggttacttct cttccatttt tcttggtgga    960
gtggatagtc tagttgactt gatgaacaag agctttctct agttgggtat taaaaaaact   1020
gattgcgaag aattgagctg gattgatata accatcttct acagtgggtg tgtaaatatc   1080
aacactgcta attttaaaaa ggaaattttg cttgatagat cagctgggaa gaagacggct   1140
ttctcaatta agttagacta tgtaagaaa ctaataacctg aaactgcaat ggtcaaaatt   1200
ttggaaaaat tatatgaaga agaggtagga gttgggatgt atgtgttgta cccttacggg   1260
ggtataatgg atgagatttc agaatcagca attccattcc ctcatcgagc tggaaataatg   1320
tatgaacttt ggtacactgc tacctgggag aagcaagaag ataacgaaaa gcatataaac   1380
tggtttcgaa gtgtttataa tttcacaact cttatgtgtg cccaaaatcc aagattggcg   1440
tatctcaatt atagggacct tgatttagga aaaactaatc ctgagagtcc taataattac   1500
acacaagcac gtatttgggg tgaaaagtat tttggtaaaa attttaacag gttagttaag   1560
gtgaaaacca aagctgatcc caataatttt tttgaaaacg aacaaagtat cccacctctt   1620
ccaccgcac atcat                                     1635

```

<210> SEQ ID NO 23

<211> LENGTH: 96

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Cannabis sp.

<400> SEQUENCE: 23

tatttttctt tctctcattc aatatccaaa cttcaattgc taatcctcga atgtaacaaa      60
tctaaaaactt acacccaaaa caaccaattg tatatg                                96

<210> SEQ ID NO 24
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Cannabis sp.

<400> SEQUENCE: 24

tatttttctt tttctcattc aatatccaaa cttccattgc taatcctcga atgcaacaaa      60
tctaaaaactc gtatacactc aaaacaaccc attgtatatg                            100

<210> SEQ ID NO 25
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (70)..(73)
<223> OTHER INFORMATION: Deletion region

<400> SEQUENCE: 25

tatttttctt tytctcattc aatatccaaa cttcmattgc taatcctcga atgyaacaaa      60
tctaaaaactc gtatacacyc aaaacaaccm attgtatatg                            100

<210> SEQ ID NO 26
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 26

gttttttgtt                                                                10

<210> SEQ ID NO 27
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic primer

<400> SEQUENCE: 27

atgcttctcg caatatattc cca                                              23

```

What is claimed is:

1. A method of detecting whether a tetrahydrocannabinolic acid (THCA) synthase gene sequence is present in a *Cannabis* plant, comprising:

obtaining a sample collected from the *Cannabis* plant that contains nucleic acids;

contacting the nucleic acids with at least five primers specific for the THCA synthase gene sequence;

amplifying the THCA synthase gene sequence when present among the nucleic acids; and

detecting amplicons upon amplification of the THCA synthase gene sequence, wherein:

55 the primers specific for the THCA synthase gene sequence include a primer specific for a promoter sequence of the THCA synthase gene sequence;

60 amplification of the THCA synthase gene sequence indicates that the THCA synthase gene sequence is at least 2000 nucleotides long; and
a colorimetric Loop-Mediated Isothermal Amplification (LAMP) assay is performed.

65 2. The method of claim 1, wherein the primer specific for the promoter sequence of the THCA synthase gene sequence binds to the promoter sequence at a location that is 200-1000 nucleotides upstream of a THCA synthase open reading frame.

55

3. The method of claim 1, wherein the THCA synthase gene sequence is between 2100-2200 nucleotides long.

4. The method of claim 1, wherein the *Cannabis* plant is a *C. sativa*, *C. indica*, or *C. ruderalis* plant.

5. The method of claim 1, wherein the *Cannabis* plant is characterized as a Type Ia, Type Ib, Type IIa, Type IIb, Type IIc, Type IIIa, Type IIIb, Type IV, or Type V plant.

6. The method of claim 1, wherein the primers specific for the THCA synthase gene sequence include two primers comprising sequences that are at least 90% identical to SEQ ID NO: 11 and SEQ ID NO: 12, respectively.

7. The method of claim 1, wherein the primers specific for the THCA synthase gene sequence include six primers comprising sequences that are at least 80% identical to each of SEQ ID NOS: 8-12.

8. The method of claim 1, wherein the primers specific for the THCA synthase gene sequence include six primers comprising sequences that are at least 90% identical to each of SEQ ID NOS: 8-12.

9. The method of claim 1, wherein the primers specific for the THCA synthase gene sequence include a primer comprising a sequence complementary to 20-60 nucleotides of SEQ ID NO: 6 and/or a primer comprising a sequence complementary to 20-60 nucleotides of SEQ ID NO: 7.

10. The method of claim 1, wherein the primers specific for the THCA synthase gene sequence comprise at least one primer comprising a sequence that is at least 70% identical to any of SEQ ID NOS: 8-12.

11. The method of claim 1, wherein the primers specific for the THCA synthase gene sequence comprise at least one primer comprising a sequence that is at least 80% identical to any of SEQ ID NOS: 8-12.

12. The method of claim 1, wherein the primers specific for the THCA synthase gene sequence comprise at least one primer comprising a sequence that is at least 80% identical to SEQ ID NO: 11 or SEQ ID NO: 12.

13. The method of claim 1, wherein the primers specific for the THCA synthase gene sequence comprise at least one primer comprising a sequence that is at least 90% identical to any of SEQ ID NOS: 8-12.

14. A method of detecting whether a tetrahydrocannabinolic acid (THCA) synthase gene sequence is present in a *Cannabis* plant, comprising:

obtaining a sample collected from the *Cannabis* plant that contains nucleic acids;
contacting the nucleic acids with at least five primers specific for the THCA synthase gene sequence;

56

amplifying the THCA synthase gene sequence when present among the nucleic acids; and
detecting amplicons upon amplification of the THCA synthase gene sequence, wherein:

the primers specific for the THCA synthase gene sequence include a primer specific for a promoter sequence of the THCA synthase gene sequence;

the primer specific for the promoter sequence of the THCA synthase gene sequence binds to the promoter sequence at a location that is 200-1000 nucleotides upstream of a THCA synthase open reading frame; and
a colorimetric Loop-Mediated Isothermal Amplification (LAMP) assay is performed.

15. The method of claim 14, wherein the primers specific for the THCA synthase gene sequence further include a primer that binds to a sequence that is 50-1000 nucleotides downstream of the THCA synthase open reading frame.

16. The method of claim 15, wherein the THCA synthase gene sequence is at least 2000 nucleotides long.

17. The method of claim 15, wherein the THCA synthase gene sequence is between 2100-2200 nucleotides long.

18. A method of detecting whether a tetrahydrocannabinolic acid (THCA) synthase gene sequence is present in a *Cannabis* plant, comprising:

obtaining a sample collected from the *Cannabis* plant that contains nucleic acids;

contacting the nucleic acids with at least five primers specific for the THCA synthase gene sequence;

amplifying the THCA synthase gene sequence when present among the nucleic acids; and

detecting amplicons upon amplification of the THCA synthase gene sequence, wherein:

the primers specific for the THCA synthase gene sequence include a primer specific for a promoter sequence of the THCA synthase gene sequence;

the amplicons of the THCA synthase gene sequence include at least part of the THCA synthase promoter and 5' UTR; and

a colorimetric Loop-Mediated Isothermal Amplification (LAMP) assay is performed.

19. The method of claim 18, wherein the THCA synthase gene sequence is at least 2000 nucleotides long.

20. The method of claim 18, wherein the THCA synthase gene sequence is between 2100-2200 nucleotides long.

* * * * *