



US 20250262310A1

(19) United States

(12) Patent Application Publication (10) Pub. No.: US 2025/0262310 A1

Levin

(43) Pub. Date: Aug. 21, 2025

(54) COMPOSITIONS AND METHODS FOR INCREASING EFFICIENCY OF CARDIAC METABOLISM

(71) Applicant: **Imbria Pharmaceuticals, Inc.**, Boston, MA (US)(72) Inventor: **Andrew D. Levin**, Newton, MA (US)(21) Appl. No.: **19/196,944**(22) Filed: **May 2, 2025**

Related U.S. Application Data

(60) Continuation of application No. 18/386,101, filed on Nov. 1, 2023, now Pat. No. 12,318,453, which is a continuation of application No. 17/828,640, filed on May 31, 2022, now Pat. No. 11,844,840, which is a division of application No. 17/074,877, filed on Oct. 20, 2020, now Pat. No. 11,376,330, which is a con-

tinuation of application No. 16/722,754, filed on Dec. 20, 2019, now Pat. No. 10,918,728.

Publication Classification

(51) Int. Cl.	<i>A61K 47/55</i>	(2017.01)
	<i>A61K 31/495</i>	(2006.01)
	<i>A61K 47/60</i>	(2017.01)
	<i>A61P 3/00</i>	(2006.01)
	<i>C07D 401/12</i>	(2006.01)

(52) U.S. Cl.	CPC	<i>A61K 47/551</i> (2017.08); <i>A61K 31/495</i> (2013.01); <i>A61K 47/55</i> (2017.08); <i>A61K 47/60</i> (2017.08); <i>C07D 401/12</i> (2013.01); <i>A61P 3/00</i> (2018.01)
---------------	-----------	--

(57) ABSTRACT

Compositions and methods for increasing efficiency of cardiac metabolism are provided.

Compounds

Compound Id	Alert	OCR AC ₅₀ (µM)	Reserve Capacity AC ₅₀ (µM)	ECAR AC ₅₀ (µM)	Mechanism
Nicotinamide	No (-)	NR	NR	NR	No effect
Trimetazidine + Nicotinamide	No (-)	NR	NR	NR	No effect
Succinate	Yes (+)	>100 †	97.1	>100 †	Other
CV-8816	No (-)	NR	NR	>100 †	No effect
CV-8814	No (-)	NR	NR	>100 †	No effect
Trimetazidine	No (-)	NR	NR	>100 †	No effect
CV-8815	No (-)	NR	NR	>100 †	No effect
Succinate + Nicotinamide + Trimetazidine	Yes (+)	NR	>100 †	>100 †	Other
Trimetazidine analog 2 + Nicotinamide	No (-)	NR	NR	>100 †	No effect
Trimetazidine analog 1 + Nicotinamide	No (-)	NR	NR	>100 †	No effect
Trimetazidine analog 3 + Nicotinamide	No (-)	NR	NR	>100 †	No effect
Succinate + Nicotinamide	No (-)	NR	NR	>100 †	No effect

AC₅₀ The concentration at which 50% maximum effect is observed for each cell health parameter.† An AC₅₀ was calculated, but is greater than the maximum surviving concentration.

↑↓ Direction of response.

NR No response observed.

Compounds

Compound ID	Alert	OGR	Rescue Capacity	EC ₅₀	Mechanism
		AC ₅₀ (nM)	AC ₅₀ (nM)	AC ₅₀ (nM)	
Nicotinamide	No (-)	NR	NR	NR	No effect
Trimetazidine + Nicotinamide	No (-)	NR	NR	NR	No effect
Succinate	Yes (+)	>100†	97.1	>100†	Other
CV-8816	No (-)	NR	NR	>100†	No effect
CV-8814	No (-)	NR	NR	>100†	No effect
Trimetazidine	No (-)	NR	NR	>100†	No effect
CV-8815	No (-)	NR	NR	>100†	No effect
Succinate + Nicotinamide + Trimetazidine	Yes (+)	NR	NR	>100†	Other
Trimetazidine analog 2 + Nicotinamide	No (-)	NR	NR	>100†	No effect
Trimetazidine analog 1 + Nicotinamide	No (-)	NR	NR	>100†	No effect
Trimetazidine analog 3 + Nicotinamide	No (-)	NR	NR	>100†	No effect
Succinate + Nicotinamide	No (-)	NR	NR	>100†	No effect

AC₅₀ The concentration at which 50% maximum effect is observed for each cell health parameter.

† An AC₅₀ was calculated, but is greater than the maximum surviving concentration.

↑ Direction of response.

NR No response observed.

FIG. 1

Nicotinamide (CY0000017658)

Assay Summary		Cell Health Parameters		Potential Mechanism	
	Concentration (μM)	OCR	Reserve Capacity	ECAR	
Incubation time:	0h	No effect	No effect	No effect	No effect
Concentrators (μM):	0.1, 0.316, 1, 3.16, 10, 31.6, 100	No effect	No effect	No effect	No effect
Replicates per concentration:	2	No effect	No effect	No effect	No effect
Cell model:	HepG2	No effect	No effect	No effect	No effect
Certified on:	2017-05-26	No effect	No effect	No effect	No effect
Data Summary		Mechanism		Mechanism	
Concentration (μM)	0.1	0.1	0.1	0.1	0.1
	0.316	No effect	No effect	No effect	No effect
	1	No effect	No effect	No effect	No effect
	3.16	No effect	No effect	No effect	No effect
	10	No effect	No effect	No effect	No effect
	31.6	No effect	No effect	No effect	No effect
	100	No effect	No effect	No effect	No effect
AICN	No (-)				

FIG. 2

MEC Minimum effective concentration that significantly crosses vehicle control threshold.
 AC_{50} The concentration at which 50% maximum effect is observed for each cell health parameter.

† An AC_{50} was calculated, but is greater than the maximum surviving concentration.

↑↓ Direction of response.

NR No response observed.

MR (%) Maximum % response.

Alert: Yes (+) if a feature deviates outside of the vehicle control this is flagged as a potential mitochondrial toxicant*.

*If the AC_{50} values of all responding features are greater than 100x plasma total C_{max} this is considered to have lower potential to exhibit mitochondrial toxicity *in vivo*. In the absence of C_{max} data then this cut-off is 50 μ M (as described by Eakins *et al.*, (2016), *TIV*, 34, 161-170).

FIG. 2 (cont.)

Nicotinamide (CY0000176586)




FIG. 3




FIG. 3 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 3 (cont.)




FIG. 3 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 3 (cont.)

Nicotinamide (CY0000176586)




FIG. 4




FIG. 4 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 4 (cont.)

Trametazidine + Nicotinamide (CY0000176587)

Assay Summary		Cell Health Parameter				ECAR				Potential Mechanism			
		IC50	MIC (µM)	AC50 (µM)	MR (%)	NR	NR	NR	NR	NR	NR	NR	NR
OCR						NR	NR	NR	NR	NR	NR	NR	NR
Reserve Capacity						NR	NR	NR	NR	NR	NR	NR	NR
ECAR						NR	NR	NR	NR	NR	NR	NR	NR
Concentration (µM)	IC50	Reserve Capacity	ECAR	Potential Mechanism									
0.1	No effect	No effect	No effect	No effect									
0.316	No effect	No effect	No effect	No effect									
1	No effect	No effect	No effect	No effect									
3.16	No effect	No effect	No effect	No effect									
10	No effect	No effect	No effect	No effect									
31.6	No effect	No effect	No effect	No effect									
100	No effect	No effect	No effect	No effect									

Alert: No (-)

Mechanism:

FIG. 5

MEC Minimum effective concentration that significantly crosses vehicle control threshold.
 AC_{50} The concentration at which 50% maximum effect is observed for each cell health parameter.

† An AC_{50} was calculated, but is greater than the maximum surviving concentration.

↑↓ Direction of response.

NR No response observed.

MR (%) Maximum % response.

Alert: Yes (+) if a feature deviates outside of the vehicle control this is flagged as a potential mitochondrial toxicant*.

*If the AC_{50} values of all responding features are greater than 100x plasma total C_{max} this is considered to have lower potential to exhibit mitochondrial toxicity *in vivo*. In the absence of C_{max} data then this cut-off is 50 μ M (as described by Eakins et al., (2016), *TIV*, 34, 161-170).

FIG. 5 (cont.)

Trimetazidine + Nicotinamide (CY0000176587)




FIG. 6




FIG. 6 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 6 (cont.)




FIG. 6 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 6 (cont.)

Trimetazidine + Nicotinamide (CY0000176587)




FIG. 7





FIG. 7 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 7 (cont.)

Succinate (CyU000155203)

**FIG. 8**

MEC Minimum effective concentration that significantly crosses vehicle control threshold.
 AC_{50} The concentration at which 50% maximum effect is observed for each cell health parameter.

† An AC_{50} was calculated, but is greater than the maximum surviving concentration.

↑↓ Direction of response.

NR No response observed.

MR (%) Maximum % response.

Alert: Yes (+) if a feature deviates outside of the vehicle control this is flagged as a potential mitochondrial toxicant*.

*If the AC_{50} values of all responding features are greater than 100x plasma total C_{max} this is considered to have lower potential to exhibit mitochondrial toxicity *in vivo*. In the absence of C_{max} data then this cut-off is 50 μ M (as described by Eakins et al., (2016), *TIV*, 34, 161-170).

FIG. 8 (cont.)

Succinate (CY0000155293)




FIG. 9




FIG. 9 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 9 (cont.)




FIG. 9 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 9 (cont.)

Succinate (CY0000155293)




FIG. 10




FIG. 10 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 10 (cont.)

CV-88 | 4 (CY00000174001)

Assay Summary	
Incubation time:	0h
Concentrations (µM):	0.1, 0.316, 1, 3.16, 10, 31.6, 100
Replicates per concentration:	2
Cell model:	HepG2
Certified on:	2017-04-12
Data Summary	
Cell Death Estimated	
OCR	NR
Reserve Capacity	NR
ECAR	↓
Passive Capacity	
Concentration (µM)	ECAR
0.1	No effect
0.316	No effect
1	No effect
3.16	No effect
10	No effect
31.6	No effect
100	No effect
Potential Mechanism	
Alexi	No (-)
Mechanism	↓

FIG. 11

MEC Minimum effective concentration that significantly crosses vehicle control threshold.
 AC_{50} The concentration at which 50% maximum effect is observed for each cell health parameter.

† An AC_{50} was calculated, but is greater than the maximum surviving concentration.

↑↓ Direction of response.

NR No response observed.

MR (%) Maximum % response.

Alert: Yes (+) if a feature deviates outside of the vehicle control this is flagged as a potential mitochondrial toxicant*.

*If the AC_{50} values of all responding features are greater than 100x plasma total C_{max} this is considered to have lower potential to exhibit mitochondrial toxicity *in vivo*. In the absence of C_{max} data then this cut-off is 50 μ M (as described by Eakins et al., (2016), *TIV*, 34, 161-170).

FIG. 11 (cont.)

CV-8814 (CY0000174001)




FIG. 12




FIG. 12 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 12 (cont.)




FIG. 12 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 12 (cont.)

CV-8814 (CY00000174001)




FIG. 13




FIG. 13 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 13 (cont.)

Triflunetazidine (CY0000174002)

Assay Summary				
	0h			
Incubation time:	0h			
Concentrations (µM):	0.1, 0.316, 1, 3.16, 10, 31.6, 100			
Replicates per concentration:	2			
Cell model:	HepG2			
Certified on:	2017-04-12			
Data Summary				
	Cell Health Parameter	MIC (µM)	EC50 (µM)	IC50 (µM)
OCR	NR	NR	NR	NR
Reserve Capacity	NR	NR	NR	NR
ECAR	↓	10.7	>100 ↓	19
Concentration (µM)	OCR	Reserve Capacity	ECAR	Potential Mechanism
0.1	No effect	No effect	No effect	No effect
0.316	No effect	No effect	No effect	No effect
1	No effect	No effect	No effect	No effect
3.16	No effect	No effect	No effect	No effect
10	No effect	No effect	No effect	No effect
31.6	No effect	No effect	No effect	No effect
100	No effect	No effect	No effect	No effect
			↓	↓
All	No (-)			Mechanism

FIG. 14

MEC Minimum effective concentration that significantly crosses vehicle control threshold.

AC₅₀ The concentration at which 50% maximum effect is observed for each cell health parameter.

† An AC₅₀ was calculated, but is greater than the maximum surviving concentration.

↑↓ Direction of response.

NR No response observed.

MR (%) Maximum % response.

Alert: Yes (+) if a feature deviates outside of the vehicle control this is flagged as a potential mitochondrial toxicant*.

*If the AC₅₀ values of all responding features are greater than 100x plasma total C_{max} this is considered to have lower potential to exhibit mitochondrial toxicity *in vivo*. In the absence of C_{max} data then this cut-off is 50µM (as described by Eakins *et al.*, (2016), *TIV*, 34, 161-170).

FIG. 14 (cont.)

Trimetazidine (CY00000174002)




FIG. 15




FIG. 15 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 15 (cont.)




FIG. 15 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 15 (cont.)

Trimetazidine (CY00000174002)




FIG. 16




FIG. 16 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 16 (cont.)

Succinate + Nicotinamide + Trifluoperazine (CY0000176589)

Assay Summary	
Incubation time:	0h
Concentrations (µM):	0.1, 0.316, 1, 3.16, 10, 31.6, 100
Replicates per concentration:	2
Cell model:	HepG2
Certified on:	2017-05-26
Data Summary	
Cell Function Parameter	
OCR	NR
Reserve Capacity	↑
ECAR	↓
Concentration (µM)	
0.1	No effect
0.316	No effect
1	No effect
3.16	No effect
10	No effect
31.6	No effect
100	No effect
	↓
	↑
Mechanism	
Yes (+)	Mechanism
	Other
All	Other

FIG. 17

MEC Minimum effective concentration that significantly crosses vehicle control threshold.

AC₅₀ The concentration at which 50% maximum effect is observed for each cell health parameter.

† An AC₅₀ was calculated, but is greater than the maximum surviving concentration.

↑↓ Direction of response.

NR No response observed.

MR (%) Maximum % response.

Alert: Yes (+) if a feature deviates outside of the vehicle control this is flagged as a potential mitochondrial toxicant*.

*If the AC₅₀ values of all responding features are greater than 100x plasma total C_{max} this is considered to have lower potential to exhibit mitochondrial toxicity *in vivo*. In the absence of C_{max} data then this cut-off is 50µM (as described by Eakins *et al.*, (2016), *TIV*, 34, 161-170).

FIG. 17 (cont.)

Succinate + Nicotinamide + Trimetazidine (CY0000176589)




FIG. 18




FIG. 18 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 18 (cont.)




FIG. 18 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 18 (cont.)

Succinate + Nicotinamide + Trimetazidine (CY0000176589)




FIG. 19




FIG. 19 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 19 (cont.)

Trimectazidine analog 2 + Nicotinamide (CY0000176591)

Assay Summary				
Parameter	Value			
Incubation time:	0h			
Concentrations (μM):	0.1, 0.316, 1, 3.16, 10, 31.6, 100			
Replicates per concentration:	2			
Cell model:	HepG2			
Certified on:	2017-05-26			
Data Summary				
Cell Health Parameter	ECAR	MEC (μM)	AC ₅₀ (μM)	MR (%)
OCR	NR	NR	NR	NR
Reserve Capacity	NR	NR	NR	NR
ECAR	↓	55.5	>100 ↓	13
Concentration (μM)	OCR	Reserve Capacity	ECAR	Potential Mechanism
0.1	No effect	No effect	No effect	No effect
0.316	No effect	No effect	No effect	No effect
1	No effect	No effect	No effect	No effect
3.16	No effect	No effect	No effect	No effect
10	No effect	No effect	No effect	No effect
31.6	No effect	No effect	No effect	No effect
100	No effect	No effect	No effect	No effect
All	No (-)	No (-)	No (-)	Mechanism

FIG. 20

MEC Minimum effective concentration that significantly crosses vehicle control threshold.

AC₅₀ The concentration at which 50% maximum effect is observed for each cell health parameter.

† An AC₅₀ was calculated, but is greater than the maximum surviving concentration.

↑↓ Direction of response.

NR No response observed.

MR (%) Maximum % response.

Alert: Yes (+) if a feature deviates outside of the vehicle control this is flagged as a potential mitochondrial toxicant*.

*If the AC₅₀ values of all responding features are greater than 100x plasma total C_{max} this is considered to have lower potential to exhibit mitochondrial toxicity *in vivo*. In the absence of C_{max} data then this cut-off is 50µM (as described by Eakins *et al.*, (2016), *TIV*, 34, 161-170).

FIG. 20 (cont.)

Trimetazidine analog 2 + Nicotinamide (CY00000176591)




FIG. 21




FIG. 21 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 21 (cont.)




FIG. 21 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 21 (cont.)

Trimetazidine analog 2 + Nicotinamide (CY0000176591)




FIG. 22




FIG. 22 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 22 (cont.)

Trimetazidine analog 1 + Nicotinamide (CV0000176590)

Assay Summary		Cell Health Parameters						ECAR		Potential Mechanism	
		Concentration (μM)	OCR	Reserve Capacity	ECAR	MIC ₅₀ (μM)	AIC ₅₀ (μM)	HR (%)	NR	NR	NR
Incubation time:	0h	0.1	No effect	No effect	No effect	NR	NR	NR	No effect	No effect	No effect
Concentrations (μM):	0.1, 0.316, 1, 3.16, 10, 31.6, 100	0.316	No effect	No effect	No effect	NR	NR	NR	No effect	No effect	No effect
Replicates per concentration:	2	1	No effect	No effect	No effect	NR	NR	NR	No effect	No effect	No effect
Cell model:	HepG2	3.16	No effect	No effect	No effect	NR	NR	NR	No effect	No effect	No effect
Certified on:	2017-05-26	10	No effect	No effect	No effect	>100	1	15	No effect	No effect	No effect
Data Summary		31.6	No effect	No effect	No effect	↓	67.3	↓	No effect	No effect	No effect
		100	No effect	No effect	No effect	↓	↓	↓	No effect	No effect	No effect
Alert		No (-)	Mechanism						No (-)		

FIG. 23

MEC Minimum effective concentration that significantly crosses vehicle control threshold.

AC₅₀ The concentration at which 50% maximum effect is observed for each cell health parameter.

† An AC₅₀ was calculated, but is greater than the maximum surviving concentration.

↑↓ Direction of response.

NR No response observed.

MR (%) Maximum % response.

Alert: Yes (+) if a feature deviates outside of the vehicle control this is flagged as a potential mitochondrial toxicant*.

*If the AC₅₀ values of all responding features are greater than 100x plasma total C_{max} this is considered to have lower potential to exhibit mitochondrial toxicity *in vivo*. In the absence of C_{max} data then this cut-off is 50µM (as described by Eakins *et al.*, (2016), *TIV*, 34, 161-170).

FIG. 23 (cont.)

Trimetazidine analog 1 + Nicotinamide (CY00000176590)




FIG. 24




FIG. 24 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 24 (cont.)




FIG. 24 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 24 (cont.)

Trimetazidine analog 1 + Nicotinamide (CY0000176590)




FIG. 25




FIG. 25 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 25 (cont.)

Trimetazidine analog 3 + Nicotinamide (CY0000176592)

Assay Summary									
Parameter	Value								
Incubation time:	0h								
Concentrations (μM):	0.1, 0.316, 1, 3.16, 10, 31.6, 100								
Replicates per concentration:	2								
Cell model:	HepG2								
Certified on:	2017-05-26								
Data Summary									
Parameter	0.1	0.316	1	3.16	10	31.6	100	ECAR	Potential Mechanism
OCR	NR	NR	NR	NR	NR	NR	NR	↓	No effect
Reserve Capacity	NR	NR	NR	NR	NR	NR	NR	↓	No effect
ECAR	73.1	73.1	73.1	73.1	73.1	73.1	>100	↓	No effect
Concentration (μM)	CCR	CCR	CCR	CCR	CCR	CCR	CCR	ECAR	Potential Mechanism
0.1	No effect	No effect	No effect	No effect	No effect	No effect	No effect	↓	No effect
0.316	No effect	No effect	No effect	No effect	No effect	No effect	No effect	↓	No effect
1	No effect	No effect	No effect	No effect	No effect	No effect	No effect	↓	No effect
3.16	No effect	No effect	No effect	No effect	No effect	No effect	No effect	↓	No effect
10	No effect	No effect	No effect	No effect	No effect	No effect	No effect	↓	No effect
31.6	No effect	No effect	No effect	No effect	No effect	No effect	No effect	↓	No effect
100	No effect	No effect	No effect	No effect	No effect	No effect	No effect	↓	No effect
Alert	No (-)	No (-)	No (-)	No (-)	No (-)	No (-)	No (-)	No (-)	Mechanism

FIG. 26

MEC Minimum effective concentration that significantly crosses vehicle control threshold.

AC₅₀ The concentration at which 50% maximum effect is observed for each cell health parameter.

† An AC₅₀ was calculated, but is greater than the maximum surviving concentration.

↑↓ Direction of response.

NR No response observed.

MR (%) Maximum % response.

Alert: Yes (+) if a feature deviates outside of the vehicle control this is flagged as a potential mitochondrial toxicant*.

*If the AC₅₀ values of all responding features are greater than 100x plasma total C_{max} this is considered to have lower potential to exhibit mitochondrial toxicity *in vivo*. In the absence of C_{max} data then this cut-off is 50µM (as described by Eakins *et al.*, (2016), *TIV*, 34, 161-170).

FIG. 26 (cont.)

Trimetazidine analog 3 + Nicotinamide (CY00000176592)




FIG. 27




FIG. 27 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 27 (cont.)




FIG. 27 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 27 (cont.)

Trimetazidine analog 3 + Nicotinamide (CY0000176592)




FIG. 28




FIG. 28 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 28 (cont.)

Succinate + Nicotinamide (succinonicotinamide)

Assay Summary		Cell Health Parameters		EC50		Potential Mechanism	
Concentration (µM)	EC50	Reserve Capacity	EC50	IC50	MR	No. of Mechanisms	
0.1	No effect	No effect	No effect	NR	NR	No effect	
0.316	No effect	No effect	No effect	NR	NR	No effect	
1	No effect	No effect	No effect	NR	NR	No effect	
3.16	No effect	No effect	No effect	NR	NR	No effect	
10	No effect	No effect	No effect	NR	NR	No effect	
31.6	No effect	No effect	No effect	NR	NR	No effect	
100	No effect	No effect	No effect	NR	NR	No effect	

FIG. 28

MEC Minimum effective concentration that significantly crosses vehicle control threshold.

AC₅₀ The concentration at which 50% maximum effect is observed for each cell health parameter.

† An AC₅₀ was calculated, but is greater than the maximum surviving concentration.

↑↓ Direction of response.

NR No response observed.

MR (%) Maximum % response.

Alert: Yes (+) if a feature deviates outside of the vehicle control this is flagged as a potential mitochondrial toxicant*.

*If the AC₅₀ values of all responding features are greater than 100x plasma total C_{max} this is considered to have lower potential to exhibit mitochondrial toxicity *in vivo*. In the absence of C_{max} data then this cut-off is 50µM (as described by Eakins *et al.*, (2016), *TIV*, 34, 161-170).

FIG. 29 (cont.)

Succinate + Nicotinamide (CY0000176588)




FIG. 30




FIG. 30 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 30 (cont.)




FIG. 30 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 30 (cont.)

Succinate + Nicotinamide (CY00000176588)




FIG. 31




FIG. 31 (cont.)

- Green dashed lines Significant cut-off from vehicle control (used to calculate the MEC).
 - Filled blue diamonds Mean data points for each concentration (plus or minus standard deviation).
 - Blue X Data points excluded from plot due to precipitate in well.
 - Open blue circles Data points excluded from plot due to data plateau, or other reasons. Points lying outside y-axis limits are annotated with small arrows.
 - Red solid lines Historical maximum and minimum responses, used to calculate AC₅₀.
- NS Fit not statistically significant.
- MR Maximum response (ratio of control).

FIG. 31 (cont.)




FIG. 32

*: p<0.01, TNS vs. CON




FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 37




FIG. 38




FIG. 39




FIG. 40






FIG. 41



LVEDD: Left Ventricular End-Systolic Diameter

FIG. 42



IVSd: Interventricular Septal Dimension

FIG. 43





FIG. 44

*: p<0.05, TAC vs. TAC+CV4/TAC+TV8
#: p<0.01, TAC vs. TAC+CV4/TAC+TV8



IVRT: Isovolumic relaxation time

FIG. 45

*: p<0.05, TAC vs. TAC+CV4/TAC+TV8
#: p<0.01, TAC vs. TAC+CV4/TAC+TV8





FIG. 46



LVDP: Left ventricular developed pressure

FIG. 47




FIG. 48




FIG. 49




FIG. 50




FIG. 51




FIG. 52




FIG. 53




FIG. 54




FIG. 55




FIG. 56




FIG. 57




FIG. 58




FIG. 59




FIG. 60




FIG. 61




FIG. 62




FIG. 63




FIG. 64





FIG. 65



66




FIG. 67




FIG. 68




FIG. 69




FIG. 70




FIG. 71




FIG. 72




FIG. 73

COMPOSITIONS AND METHODS FOR INCREASING EFFICIENCY OF CARDIAC METABOLISM

[0001] This application is a continuation of U.S. patent application Ser. No. 16/011,196, which claims the benefit of, and priority to, U.S. Provisional Patent Application No. 62/647,926, filed Mar. 26, 2018, U.S. Provisional Patent Application No. 62/637,434, filed Mar. 2, 2018, U.S. Provisional Patent Application No. 62/710,316, filed Feb. 16, 2018, U.S. Provisional Patent Application No. 62/524,237, filed Jun. 23, 2017, and U.S. Provisional Patent Application No. 62/522,214, filed Jun. 20, 2017, the contents of each of which are incorporated by reference.

FIELD OF THE INVENTION

[0002] This application is related to compositions and methods for increasing the efficiency of cardiac metabolism.

BACKGROUND

[0003] Heart disease is the leading cause of death worldwide, accounting for 15 million deaths across the globe in 2015. In many forms of heart disease, decreased cardiac efficiency stems from changes in mitochondrial energy metabolism. Mitochondria are sub-cellular compartments in which metabolites derived from glucose and fatty acids are oxidized to produce high-energy molecules. Increasing fatty acid oxidation in the heart decreases glucose oxidation, and vice versa. Glucose oxidation is a more efficient source of energy, but in certain types of heart disease, such as heart failure, ischemic heart disease, and diabetic cardiomyopathies, fatty acid oxidation predominates in cardiac mitochondria. As a result, the pumping capacity of the heart is reduced.

[0004] Existing drugs that redress the balance between glucose oxidation and fatty acid oxidation in cardiac mitochondria have serious shortcomings. Foremost among them is that such drugs address only part of the problem: the reliance on fatty acid oxidation in lieu of glucose oxidation causes a 10% reduction in efficiency in energy production, but patients with heart disease often show a decrease in cardiac efficiency of up to 30%. Consequently, existing approaches to improve cardiac function by altering mitochondrial metabolism are unsatisfactory, and millions of people continue to die from heart disease each year.

SUMMARY

[0005] The invention provides compositions that stimulate cardiac glucose oxidation and mitochondrial respiration. The compositions include a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, such as trimetazidine, and a compound that promotes mitochondrial respiration, such as succinate. The compositions may also include a molecule, such as nicotinic acid, that serves as a precursor for synthesis of nicotinamide adenine dinucleotide (NAD^+), which also facilitates mitochondrial respiration. Preferably, the compositions include compounds in which a trimetazidine derivative, succinate, and, optionally, a NAD^+ precursor are covalently linked in a single molecule. Such compounds can be metabolized in the body to allow the individual components to exert distinct biochemical effects to increase glucose oxidation relative to fatty acid oxidation and improve overall mitochondrial

respiration in the heart. The invention also provides methods of altering cardiac metabolism by providing compounds of the invention.

[0006] Because the compositions concomitantly shift cardiac metabolism toward glucose oxidation and increase mitochondrial respiration, they are useful as therapeutic agents for treating heart diseases characterized by elevated fatty acid oxidation, such as heart failure, ischemic heart disease, and diabetic cardiomyopathies. By shifting cardiac metabolism from fatty acid oxidation to glucose oxidation, the compositions allow the use of a more efficient source of energy. In addition, the compositions stimulate metabolic pathways that are common to oxidation of both glucose and fatty acids and that may also be impaired in patients with heart disease. Some compositions of the invention include a compound that comprises trimetazidine covalently coupled to one or more activators of mitochondrial respiration.

[0007] Furthermore, trimetazidine can cause Parkinsonian symptoms for a portion of the population. Without being limited by any particular theory or mechanism of action, it is also believed that delivery of trimetazidine as a component of a larger molecule may improve its efficacy and mitigate its side effects.

[0008] In an aspect, the invention includes compounds represented by formula (I):



in which A is a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, L is a linker, and B is a compound that promotes mitochondrial respiration.


[0009] The compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation may be trimetazidine, etomoxir, perhexiline, a PPAR agonist, a malonyl CoA decarboxylase inhibitor, or dichloroacetate.

[0010] The compound that promotes mitochondrial respiration may be an intermediate of the citric acid cycle or a molecule that can be metabolized to enter the citric acid cycle. For example, the compound may be succinate, fumarate, malate, oxaloacetate, citrate, isocitrate, α -ketoglutarate, pyruvate, acetone, acetoacetic acid, β -hydroxybutyric acid, β -ketopentanoate, or β -hydroxypentanoate.

[0011] The linker may be any suitable linker that can be cleaved in vivo. The linker may be an alkoxy group. The linker may be polyethylene glycol of any length. Preferably, the linker is represented by $(\text{CH}_2\text{CH}_2\text{O})_x$, in which $x=1-15$.


[0012] The compound may include a NAD^+ precursor molecule covalently linked to another component of the compound. The NAD^+ precursor molecule may be nicotinic acid, nicotinamide, or nicotinamide riboside. The NAD^+ precursor molecule may be attached to the compound that shifts cardiac metabolism, the compound that promotes mitochondrial respiration, or the linker. The NAD^+ precursor molecule may be attached to another component via an additional linker. Preferably, the NAD^+ precursor molecule is attached to the compound that promotes mitochondrial respiration via a 1,3-propanediol linkage.

[0013] The compound of formula (I) may be represented by formula (II):




in which $y=1-3$.

[0014] The compound of formula (I) may be represented by formula (III):



in which $y=1-3$.


[0015] In another aspect, the invention includes a compound represented by formula (IV):



in which R^1 , R^2 , and R^3 are independently H or a (C_1-C_4) alkyl group; R^4 and R^5 together are $=O$, $-O(CH_2)_mO-$, or $-(CH_2)_m-$, in which $m=2-4$, or R^4 is H and R^5 is OR^{14} , SR^{14} , or $(CH_2CH_2O)_nH$, in which R^{14} is H or a (C_1-C_4) alkyl group and $n=1-15$; and R^6 is a single or multi-ring structure optionally substituted at one or more ring positions by a heteroatom, in which each ring position optionally comprises one or more substituents.


[0016] One or more ring position of R^6 may include a substituent that includes a compound that promotes mitochondrial respiration, such as succinate, fumarate, malate, oxaloacetate, citrate, isocitrate, α -ketoglutarate, pyruvate, acetone, acetoacetic acid, β -hydroxybutyric acid, β -ketopen-tanoate, or β -hydroxypentanoate. The substituent may include a linker, such as $(CH_2CH_2O)_x$, in which $x=1-15$. The substituent may include a NAD^+ precursor molecule, such as nicotinic acid, nicotinamide, and nicotinamide riboside.

[0017] The substituent on a ring position of R^6 may be




in which $y=1-3$.

[0018] The substituent on a ring position of R^6 may be





in which $y=1-3$.


[0019] R^6 may be



[0020] The compound of formula (IV) may have a structure represented by formula (IX) or formula (X):



[0021] In another aspect, the invention includes compounds represented by formula (V):




in which R¹, R², and R³ are independently H or a (C₁-C₄) alkyl group; R⁷ and R⁸ together are —O—, —O(CH₂)_mO—, or —(CH₂)_m—, in which m=2-4, or R⁴ is H and R⁸ is H, OR¹⁴, SR¹⁴, or (CH₂CH₂O)_nH, in which R¹⁴ is H or a (C₁-C₄)alkyl group and n=1-15; R⁹, R¹⁰, R¹², and R¹³ are independently H or (CH₂CH₂O)_zH, in which z=1-6; and R¹¹ comprises a compound that promotes mitochondrial respiration.

[0022] The compound that promotes mitochondrial respiration may be an intermediate of the citric acid cycle or a molecule that can be metabolized to enter the citric acid cycle. For example, the compound may be succinate, fumarate, malate, oxaloacetate, citrate, isocitrate, α-ketoglutarate, pyruvate, acetone, acetoacetic acid, β-hydroxybutyric acid, β-ketopentanoate, or β-hydroxypentanoate.

[0023] R¹¹ may include a linker, such as polyethylene glycol. For example, R¹¹ may include (CH₂CH₂O)_x, in which x=1-15.


[0024] R¹¹ may be



in which y=1-3.

[0025] R¹¹ may include a NAD⁺ precursor molecule. For example, R¹¹ may include nicotinic acid, nicotinamide, or nicotinamide riboside.

[0026] R¹¹ may be



in which y=1-3.

[0027] In an aspect, the invention includes compounds represented by formula (VII):

A-C

(VII),

in which A is a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, and C is a NAD⁺ precursor molecule. A and C may be covalently linked.

[0028] The compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation may be trimetazidine, etomoxir, perhexiline, a PPAR agonist, a malonyl CoA decarboxylase inhibitor, or dichloroacetate.

[0029] The compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation may be PEGylated with an ethylene glycol moiety. The compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation may have multiple ethylene glycol moieties, such as one, two, three, four, five, or more ethylene glycol moieties. The ethylene glycol moiety may be represented by (CH₂CH₂O)_x, in which x=1-15. The ethylene glycol moiety may form a covalent linkage between the compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation and the NAD⁺ precursor molecule. The ethylene glycol moiety may be separate from a covalent linkage between the compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation and the NAD⁺ precursor molecule. The compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation may be a PEGylated form of trimetazidine.

[0030] The NAD⁺ precursor molecule may be nicotinic acid, nicotinamide, or nicotinamide riboside.

[0031] The compound of formula (VII) may include nicotinic acid that is covalently linked to a PEGylated form of trimetazidine. The nicotinic acid may be covalently linked via the PEGylated moiety, i.e., via an ethylene glycol linkage. The nicotinic acid may be covalently linked via the trimetazidine moiety.

[0032] The compound of formula (VII) may have a structure represented by formula (X), as shown above.

[0033] In an aspect, the invention includes compounds represented by formula (VIII):

A-L-C

(VIII),

in which A is a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, L is a linker, and C is a NAD⁺ precursor molecule. A may be covalently linked to L, and L may be covalently linked to C.

[0034] The compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, the linker, and the NAD⁺ precursor molecule may be as described above in relation to compounds of other formulas.

[0035] The compound of formula (VIII) may have a structure represented by formula (X), as shown above.

[0036] Any of the compounds described above may include one or more atoms that are enriched for an isotope. For example, the compounds may have one or more hydrogen atoms replaced with deuterium or tritium. The isotopically enriched atom or atoms may be located at any position within the compound.

[0037] In an aspect, the invention includes compositions that include at least two of A, B, and C, in which A is a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation as described above, B is a compound that promotes mitochondrial respiration as described above, and C is a NAD⁺ precursor molecule as described above. The compositions may include A, B, and C. Each of components A, B, and C may be provided as a separate molecule, or two or more of the components may be covalently linked in a single molecule. For example, components A and B may be covalently linked in a single molecule, and C may be provided as a separate molecule.

[0038] The compositions may include co-crystals of two or more separate molecules that include two or more of components A, B, and C. For example, a co-crystal may include (1) a compound of formula (I), (III), (IV), or (V) and (2) nicotinic acid, nicotinamide, or nicotinamide riboside. Preferably the co-crystal includes nicotinamide.

[0039] In an aspect, the invention includes methods of increasing efficiency of cardiac metabolism in a subject. The methods include providing a compound represented by formula (I), as described above. In the methods, the compound of formula (I) may include any of the features described above in relation to compounds of the invention.

[0040] In an aspect, the invention includes methods of increasing efficiency of cardiac metabolism in a subject. The methods include providing a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, a compound that promotes mitochondrial respiration, and, optionally, a compound that is a NAD⁺ precursor molecule.

[0041] The compound that shifts cardiac metabolism from fatty acid oxidation to glucose may be trimetazidine, etoximoxir, perhexiline, a PPAR agonist, a malonyl CoA decarboxylase inhibitor, or dichloroacetate.

[0042] The compound that promotes mitochondrial respiration may be an intermediate of the citric acid cycle or a molecule that can be metabolized to enter the citric acid cycle, such as succinate, fumarate, malate, oxaloacetate, citrate, isocitrate, α-ketoglutarate, pyruvate, acetone, acetoacetic acid, β-hydroxybutyric acid, β-ketopentanoate, or β-hydroxypentanoate.

[0043] The NAD⁺ precursor molecule may be nicotinic acid, nicotinamide, or nicotinamide riboside.

[0044] The compounds may be provided in any suitable manner. The compounds may be provided in a single composition. Alternatively, the compounds may not be provided in a single composition. For example, one or two of the compounds may be provided in a single composition, and another compound may be provided in a separate composition. Alternatively, each compound may be provided in a separate composition. The compounds may be provided simultaneously or sequentially. The compounds may be provided at different intervals, with different frequency, or in different quantities.

[0045] It is believed that any disease that may be treated using trimetazidine would benefit from compounds of the invention as described herein with more efficacious results and fewer side effects. Exemplary diseases are those that involve impaired mitochondrial function or altered fatty acid oxidation, such as heart failure diseases, cardiac dysfunction diseases, or muscle myopathy diseases. Exemplary methods involve providing a composition as described herein or any combination of a compound that shifts cardiac metabolism from fatty acid oxidation to glucose metabolism, a compound that promotes mitochondrial respiration, and/or optionally an NAD⁺ precursor molecule.

BRIEF DESCRIPTION OF THE DRAWINGS

[0046] FIG. 1 is a table summarizing the effects of various compounds on mitochondrial function.

[0047] FIG. 2 is a table summarizing the effects of nicotinamide on various mitochondrial functional parameters.

[0048] FIG. 3 is a series of graphs showing the effects of nicotinamide on oxygen consumption rate and reserve capacity.

[0049] FIG. 4 is a series of graphs showing the effects of nicotinamide on extracellular acidification rate.

[0050] FIG. 5 is a table summarizing the effects of a combination of trimetazidine and nicotinamide on various mitochondrial functional parameters.

[0051] FIG. 6 is a series of graphs showing the effects of a combination of trimetazidine and nicotinamide on oxygen consumption rate and reserve capacity.

[0052] FIG. 7 is a series of graphs showing the effects of a combination of trimetazidine and nicotinamide on extracellular acidification rate.

[0053] FIG. 8 is a table summarizing the effects of succinate on various mitochondrial functional parameters.

[0054] FIG. 9 is a series of graphs showing the effects of succinate on oxygen consumption rate and reserve capacity.

[0055] FIG. 10 is a series of graphs showing the effects of succinate on extracellular acidification rate.

[0056] FIG. 11 is a table summarizing the effects of compound CV-8814 on various mitochondrial functional parameters.

[0057] FIG. 12 is a series of graphs showing the effects of compound CV-8814 on oxygen consumption rate and reserve capacity.

[0058] FIG. 13 is a series of graphs showing the effects of compound CV-8814 on extracellular acidification rate.

[0059] FIG. 14 is a table summarizing the effects of trimetazidine on various mitochondrial functional parameters.

[0060] FIG. 15 is a series of graphs showing the effects of trimetazidine on oxygen consumption rate and reserve capacity.

[0061] FIG. 16 is a series of graphs showing the effects of trimetazidine on extracellular acidification rate.

[0062] FIG. 17 is a table summarizing the effects of a combination of succinate, nicotinamide, and trimetazidine on various mitochondrial functional parameters.

[0063] FIG. 18 is a series of graphs showing the effects of a combination of succinate, nicotinamide, and trimetazidine on oxygen consumption rate and reserve capacity.

[0064] FIG. 19 is a series of graphs showing the effects of a combination of succinate, nicotinamide, and trimetazidine on extracellular acidification rate.

[0065] FIG. 20 is a table summarizing the effects of a combination of trimetazidine analog 2 and nicotinamide on various mitochondrial functional parameters.

[0066] FIG. 21 is a series of graphs showing the effects of a combination of trimetazidine analog 2 and nicotinamide on oxygen consumption rate and reserve capacity.

[0067] FIG. 22 is a series of graphs showing the effects of a combination of trimetazidine analog 2 and nicotinamide on extracellular acidification rate.

[0068] FIG. 23 is a table summarizing the effects of a combination of trimetazidine analog 1 and nicotinamide on various mitochondrial functional parameters.

[0069] FIG. 24 is a series of graphs showing the effects of a combination of trimetazidine analog 1 and nicotinamide on oxygen consumption rate and reserve capacity.

[0070] FIG. 25 is a series of graphs showing the effects of a combination of trimetazidine analog 1 and nicotinamide on extracellular acidification rate.

[0071] FIG. 26 is a table summarizing the effects of a combination of trimetazidine analog 3 and nicotinamide on various mitochondrial functional parameters.

- [0072] FIG. 27 is a series of graphs showing the effects of a combination of trimetazidine analog 3 and nicotinamide on oxygen consumption rate and reserve capacity.
- [0073] FIG. 28 is a series of graphs showing the effects of a combination of trimetazidine analog 3 and nicotinamide on extracellular acidification rate.
- [0074] FIG. 29 is a table summarizing the effects of a combination of succinate and nicotinamide on various mitochondrial functional parameters.
- [0075] FIG. 30 is a series of graphs showing the effects of a combination of succinate and nicotinamide on oxygen consumption rate and reserve capacity.
- [0076] FIG. 31 is a series of graphs showing the effects of a combination of succinate and nicotinamide on extracellular acidification rate.
- [0077] FIG. 32 is a schematic of the ischemia-reperfusion (IR) method used to analyze the effects of compositions of the invention on coronary flow.
- [0078] FIG. 33 is a graph of coronary flow of after IR.
- [0079] FIG. 34 is graph of left ventricular developed pressure (LVDP) after IR.
- [0080] FIG. 35 shows images of TTC-stained heart slices after IR.
- [0081] FIG. 36 is graph of infarct size after IR.
- [0082] FIG. 37 is a schematic of the method used to analyze the effects of compositions of the invention on cardiac function.
- [0083] FIG. 38 shows hearts from mice six weeks after transverse aortic constriction.
- [0084] FIG. 39 is of graph of heart weight relative to body weight six weeks after transverse aortic constriction.
- [0085] FIG. 40 is graph of heart weight six weeks after transverse aortic constriction.
- [0086] FIG. 41 shows graphs of fractional shortening (FS) and ejection fraction (EF) at indicated time points after transverse aortic constriction.
- [0087] FIG. 42 is a graph of left ventricular end-systolic diameter at indicated time points after transverse aortic constriction.
- [0088] FIG. 43 is a graph of intraventricular septal dimension at indicated time points after transverse aortic constriction.
- [0089] FIG. 44 is a graph of left ventricular mass at indicated time points after transverse aortic constriction.
- [0090] FIG. 45 is a graph of isovolumic relaxation time at indicated time points after transverse aortic constriction.
- [0091] FIG. 46 is a graph of the ratio peak velocity flow in early diastole vs. late diastole at indicated time points after transverse aortic constriction.
- [0092] FIG. 47 is a graph of left ventricular developed pressure at six weeks after transverse aortic constriction.
- [0093] FIG. 48 is a graph of the rate of left ventricle pressure rise at six weeks after transverse aortic constriction.
- [0094] FIG. 49 is a graph showing levels of CV-8814 and trimetazidine after intravenous administration of CV-8834.
- [0095] FIG. 50 is a graph showing levels of CV-8814 and trimetazidine after oral administration of CV-8834.
- [0096] FIG. 51 is a graph showing levels of CV-8814 and trimetazidine after oral administration of CV-8834.
- [0097] FIG. 52 is a graph showing levels of CV-8814 and trimetazidine after oral administration of CV-8834.
- [0098] FIG. 53 is a graph showing levels of CV-8814 and trimetazidine after oral administration of CV-8834.

- [0099] FIG. 54 is a graph showing levels of trimetazidine after oral administration of CV-8972 or intravenous administration of trimetazidine.
- [0100] FIG. 55 is a graph showing levels of CV-8814 after oral administration of CV-8972 or intravenous administration of CV-8814.
- [0101] FIG. 56 is a graph showing levels of CV-8814 after intravenous administration of CV-8834 or oral administration of CV-8834.
- [0102] FIG. 57 is a graph showing levels of CV-8814 after intravenous administration of CV-8814 or oral administration of CV-8814.
- [0103] FIG. 58 is a graph showing the HPLC elution profile of a batch of CV-8972.
- [0104] FIG. 59 is a graph showing analysis of molecular species present in a batch of CV-8972.
- [0105] FIG. 60 is a pair of graphs showing HPLC elution profiles of molecular species present in a batch of CV-8972.
- [0106] FIG. 61 is a pair of graphs showing HPLC elution profiles of molecular species present in a batch of CV-8972.
- [0107] FIG. 62 is a graph showing X-ray powder diffraction analysis of a batch of CV-8972.
- [0108] FIG. 63 is a graph showing X-ray powder diffraction analysis of batches of CV-8972.
- [0109] FIG. 64 is a graph showing differential scanning calorimetry and thermal gravimetric analysis of a batch of CV-8972.
- [0110] FIG. 65 is a graph showing dynamic vapor sorption (DVS) of a batch of CV-8972.
- [0111] FIG. 66 is a graph showing differential scanning calorimetry and thermal gravimetric analysis of a batch of CV-8972.
- [0112] FIG. 67 is a graph showing dynamic vapor sorption (DVS) of a batch of CV-8972.
- [0113] FIG. 68 is a graph showing X-ray powder diffraction analysis of samples of CV-8972.
- [0114] FIG. 69 is a graph showing differential scanning calorimetry and thermal gravimetric analysis of a batch of CV-8972.
- [0115] FIG. 70 is a graph showing X-ray powder diffraction analysis of samples of CV-8972.
- [0116] FIG. 71 is a graph showing X-ray powder diffraction analysis of samples of CV-8972.
- [0117] FIG. 72 is a graph showing differential scanning calorimetry and thermal gravimetric analysis of samples containing form A of CV-8972.
- [0118] FIG. 73 is a graph showing differential scanning calorimetry and thermal gravimetric analysis of a sample containing form A of CV-8972.

DETAILED DESCRIPTION

- [0119] The invention provides compositions that increase the efficiency of cardiac metabolism by concomitantly shifting cardiac metabolism from fatty acid oxidation to glucose oxidation and increasing mitochondrial respiration. Glucose oxidation and fatty acid oxidation are energy-producing metabolic pathways that compete with each other for substrates. In glucose oxidation, glucose is broken down to pyruvate via glycolysis in the cytosol of the cell. Pyruvate then enters the mitochondria, where it is converted to acetyl coenzyme A (acetyl-CoA). In beta-oxidation of fatty acids, which occurs in the mitochondria, two-carbon units from long-chain fatty acids are sequentially converted to acetyl-CoA.

[0120] The remaining steps in energy production from oxidation of glucose or fatty acids are common to the two pathways. Acetyl-CoA is oxidized to carbon dioxide (CO_2) via the citric acid cycle, which results in the conversion of nicotinamide adenine dinucleotide (NAD^+) to its reduced form, NADH. NADH, in turn, drives the mitochondrial electron transport chain. The electron transport chain comprises a series of four mitochondrial membrane-bound complexes that transfer electrons via redox reactions and pump protons across the membrane to create a proton gradient. The redox reactions of the electron transport chain require molecular oxygen (O_2). Finally, the proton gradient enables another membrane-bound enzymatic complex to form high-energy ATP molecules, the source of energy for most cellular reactions.

[0121] In many types of heart disease, the overall efficiency of energy production by cardiac mitochondria is diminished. In part, this is due to an increased reliance on fatty acid oxidation over glucose oxidation in many types of heart disease. Glucose oxidation is a more efficient pathway for energy production, as measured by the number of ATP molecules produced per O_2 molecule consumed, than is fatty acid oxidation. However, other metabolic changes contribute to decreased cardiac efficiency in patients with heart disease. For example, overall mitochondrial oxidative metabolism can be impaired in heart failure, and energy production is decreased in ischemic heart disease due to a limited supply of oxygen. As indicated above, the final steps in ATP synthesis, which include several redox reactions and oxygen-driven proton transport, are common to both the glucose oxidation and fatty acid oxidation pathways. Thus, shifting the balance from fatty acid oxidation to glucose oxidation by itself is not enough in many circumstances to restore full cardiac efficiency because downstream processes are affected as well.

[0122] The invention provides compositions that improve cardiac efficiency by using multiple mechanisms to alter mitochondrial metabolism. By including a component that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation and one or more other components that promote mitochondrial respiration, the compositions trigger a change in the pathway used to produce energy and concomitantly improve overall mitochondrial oxidative function. Consequently, the compositions of the invention are more effective at restoring cardiac capacity in patients with heart disease, such as heart failure, ischemic heart disease, and diabetic cardiomyopathies, than are compounds that only effect a shift to glucose oxidation.

[0123] In some embodiments, the compositions are compounds represented by formula (I):



in which A is a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, L is a linker, and B is a compound that promotes mitochondrial respiration.

[0124] Component A may be any suitable compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation. Such compounds can be classified based on their mechanism of action. See Fillmore, N., et al., Mitochondrial fatty acid oxidation alterations in heart failure, ischemic heart disease and diabetic cardiomyopathy, *Brit. J. Pharmacol.* 171:2080-2090 (2014), incorporated herein by reference.

[0125] One class of glucose-shifting compounds includes compounds that inhibit fatty acid oxidation directly. Compounds in this class include inhibitors of malonyl CoA decarboxylase (MCD), carnitine palmitoyl transferase 1 (CPT-1), or mitochondrial fatty acid oxidation. Mitochondrial fatty acid oxidation inhibitors include trimetazidine and other compounds described in WO 2002/064576, which is incorporated herein by reference. Trimetazidine binds to distinct sites on the inner and outer mitochondrial membranes and affects both ion permeability and metabolic function of mitochondria. Morin, D., et al., Evidence for the existence of [^3H]-trimetazidine binding sites involved in the regulation of the mitochondrial permeability transition pore, *Brit. J. Pharmacol.* 123:1385-1394 (1998), incorporated herein by reference. MCD inhibitors include CBM-301106, CBM-300864, CBM-301940, 5-(1,1,1,3,3-hexafluoro-2-hydroxypropan-2-yl)-4,5-dihydroisoxazole-3-carboxamides, methyl 5-(N-(4-(1,1,1,3,3-hexafluoro-2-hydroxypropan-2-yl)phenyl)morpholine-4-carboxamido)pentanoate, and other compounds described in Chung, J. F., et al., Discovery of Potent and Orally Available Malonyl-CoA Decarboxylase Inhibitors as Cardioprotective Agents, *J. Med. Chem.* 49:4055-4058 (2006); Cheng J. F. et al., Synthesis and structure-activity relationship of small-molecule malonyl coenzyme A decarboxylase inhibitors, *J. Med. Chem.* 49:1517-1525 (2006); US Publication No. 2004/0082564; and WO 2002/058698, which are incorporated herein by reference. CPT-1 inhibitors include oxfenicine, perhexiline, etomoxir, and other compounds described in WO 2015/018660, WO 2008/109991; WO 2009/015485; US Publication No. 2011/0212072; and WO 2009/156479, which are incorporated herein by reference.

[0126] Another class of glucose-shifting compounds includes compounds that stimulate glucose oxidation directly. Examples of such compounds are described in US Publication No. 2003/0191182; WO 2006/117686; U.S. Pat. No. 8,202,901, which are incorporated herein by reference.

[0127] Another class of glucose-shifting compounds includes compounds that decrease the level of circulating fatty acids that supply the heart. Examples of such compounds include agonists of PPAR α and PPAR γ , including fibrate drugs, such as clofibrate, gemfibrozil, ciprofibrate, bezafibrate, and fenofibrate, and thiazolidinediones, GW-9662, and other compounds described in U.S. Pat. No. 9,096,538, which is incorporated herein by reference.

[0128] Component L may be any suitable linker. Preferably, the linker can be cleaved in vivo to release components A and B. The linker may be an alkoxy group. The linker may be polyethylene glycol of any length. The linker may be represented by $(\text{CH}_2\text{CH}_2\text{O})_x$, in which $x=1-15$ or $(\text{CH}_2\text{CH}_2\text{O})_x$, in which $x=1-3$. Other suitable linkers include 1,3-propanediol, diazo linkers, phosphoramidite linkers, disulfide linkers, cleavable peptides, iminodiacetic acid linkers, thioether linkers, and other linkers described in Leriche, G., et al., Cleavable linkers in chemical biology, *Bioorg. Med. Chem.* 20:571-582 (2012); WO 1995000165; and U.S. Pat. No. 8,461,117, which are incorporated herein by reference.

[0129] Component B may be any compound that promotes mitochondrial respiration. For example, component B may be an intermediate of the citric acid cycle or a molecule that can be metabolized to enter the citric acid cycle, such as succinate, fumarate, malate, oxaloacetate, citrate, isocitrate, α -ketoglutarate, pyruvate, acetone, acetoacetic acid, β -hydroxybutyric acid, β -ketopentanoate, or β -hydroxypentano-


ate. Intermediates of the citric acid cycle may become depleted if these molecules are used for biosynthetic purposes, resulting in inefficient generation of ATP from the citric acid cycle. However, due to the anaplerotic effect, providing one intermediate of the citric acid cycle leads to restoration of all intermediates as the cycle turns. Thus, intermediates of the citric acid cycle can promote mitochondrial respiration.

[0130] The compound may include a NAD⁺ precursor molecule. NAD⁺ is an important oxidizing agent that acts as a coenzyme in multiple reactions of the citric acid cycle. In these reactions, NAD⁺ is reduced to NADH. Conversely, NADH is oxidized back to NAD⁺ when it donates electrons to mitochondrial electron transport chain. In humans, NAD⁺ can be synthesized de novo from tryptophan, but not in quantities sufficient to meet metabolic demands. Consequently, NAD⁺ is also synthesized via a salvage pathway, which uses precursors that must be supplied from the diet. Among the precursors used by the salvage pathway for NAD⁺ synthesis are nicotinic acid, nicotinamide, and nicotinamide riboside. By providing a NAD⁺ precursor, such as nicotinic acid, nicotinamide, or nicotinamide riboside, the compound facilitates NAD⁺ synthesis.

[0131] The inclusion of a NAD⁺ precursor in compounds of the invention allows the compounds to stimulate energy production in cardiac mitochondria in multiple ways. First, component A shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, which is inherently more efficient. Next, component B ensures that the intermediates of the citric acid cycle are present at adequate levels and do not become depleted or limiting. As a result, glucose-derived acetyl CoA is efficiently oxidized. Finally, the NAD⁺ precursor provides an essential coenzyme that cycles between oxidized and reduced forms to promote respiration. In the oxidized form, NAD⁺ drives reactions of the citric acid cycle. In the reduced form, NADH promotes electron trans-


and allows them to remain in circulation longer. The compound may contain a PEG polymer of any size. For example, the PEG polymer may have from 1-500 ($\text{CH}_2\text{CH}_2\text{O}$) units. The PEG polymer may have any suitable geometry, such as a straight chain, branched chain, star configuration, or comb configuration. The compound may be PEGylated at any site. For example, the compound may be PEGylated on component A, component B, component L, or, if present, the NAD⁺ precursor. The compound may be PEGylated at multiple sites. For a compound PEGylated at multiple sites, the various PEG polymers may be of the same or different size and of the same or different configuration.

[0134] The compound may be a PEGylated form of trimetazidine. For example, the compound may be represented by formula (VI):




in which one or more of the carbon atoms at positions A, B, C, D, and E and/or the nitrogen atom at position F are substituted with $-(CH_2CH_2O)_nH$ and n=1-15. The carbon atoms at positions A, B, C, D, and E may have two PEG substituents. In molecules that have multiple PEG chains, the different PEG chains may have the same or different length.

[0135] The compounds of formula (I) may be represented by formula (II):




in which $y=1-3$.

[0136] The compounds of formula (I) may be represented by formula (III):



in which $y=1-3$.

[0137] The invention also provides compounds represented by formula (IV):




in which R¹, R², and R³ are independently H or a (C₁-C₄) alkyl group; R⁴ and R⁵ together are =O, —O(CH₂)_mO—, or —(CH₂)_m—, in which m=2-4, or R⁴ is H and R⁵ is OR¹⁴, SR¹⁴, or (CH₂CH₂O)_nH, in which R¹⁴ is H or a (C₁-C₄) alkyl group and n=1-15; and R⁶ is a single or multi-ring structure optionally substituted at one or more ring positions by a heteroatom, in which each ring position optionally comprises one or more substituents.

[0138] R⁶ may be a single or multi-ring structure of any size. For example, the structure may contain 3-22 atoms, not including hydrogen atoms bonded to atoms in ring positions. The structure may include one or more alkyl, alkenyl, or aromatic rings. The structure may include one or more heteroatoms, i.e., atoms other than carbon. For example, the heteroatom may be oxygen, nitrogen, or sulfur, or phosphorus.


[0139] One or more ring position of R⁶ may include a substituent that includes a compound that promotes mitochondrial respiration, as described above in relation to component B of formula (I). The substituent may include a linker, as described above in relation to component L of formula (I). The substituent may include a NAD⁺ precursor molecule, as described above in relation to compounds of formula (I).

[0140] The substituent on a ring position of R⁶ may be




in which y=1-3.

[0141] The substituent on a ring position of R⁶ may be





in which y=1-3.

[0142] R⁶ may be




[0143] For some compounds of the invention that include trimetazidine prodrugs, analogs, derivatives, it is advantageous to have the trimetazidine moiety substituted with a single ethylene glycol moiety. Thus, preferred compositions of the invention include compounds of formulas (I) and (VII) that contain linkers in which x=1, compounds of formulas (II) and (III) in which y=1, compounds of formula (V) in which z=1, compounds of formula (VI) in which n=1, and compounds of formula (VII) in which A is linked to C via a single ethylene glycol moiety. Without wishing to be bound by theory, the attachment of a single ethylene glycol moiety to the trimetazidine moiety may improve the bioavailability of trimetazidine.

[0144] The compound of formula (IV) may have structure represented by formula (IX) or formula (X):




[0145] The invention also provides compounds represented by formula (V):



in which R¹, R², and R³ are independently H or a (C₁-C₄) alkyl group; R⁴ and R⁸ together are =O, —O(CH₂)_mO—, or —(CH₂)_m—, in which m=2-4, or R⁴ is H and R⁸ is H, OR¹⁴, SR¹⁴, or (CH₂CH₂O)_nH, in which R¹⁴ is H or a (C₁-C₄) alkyl group and n=1-15; R⁹, R¹⁰, R¹², and R¹³ are independently H or (CH₂CH₂O)_zH, in which z=1-15; and R¹ comprises a compound that promotes mitochondrial respiration, as

described above in relation to component B of formula (I). R¹¹ may include a linker, as described above in relation to component L of formula (I).


[0146] R¹ may be



in which y=1-3.

[0147] R¹¹ may include a NAD⁺ precursor molecule, as described above in relation to compounds of formula (I).

[0148] R¹ may be



in which y=1-3.

[0149] In some embodiments described above, compounds of the invention include multiple active agents joined by linkers in a single molecule. It may be advantageous to deliver multiple active agents as components of a single molecule. Without wishing to be bound by a particular theory, there are several reasons why co-delivery of active agents in a single molecule may be advantageous. One possibility is that a single large molecule may have reduced side effects compared to the component agents. Free trimetazidine causes symptoms similar to those in Parkinson's disease in a fraction of patients. However, when trimetazidine is derivatized to include other components, such as succinate, the molecule is bulkier and may not be able to access sites where free trimetazidine can cause unintended effects. Trimetazidine derivatized as described above is also more hydrophilic and thus may be less likely to cross the blood-brain barrier to cause neurological effects. Another possibility is that modification of trimetazidine may alter its pharmacokinetic properties. Because the derivatized molecule is metabolized to produce the active agent, the active agent is released gradually. Consequently, levels of the active agent in the body may not reach peaks as high as when a comparable amount is administered in a single bolus. Another possibility is that less of each active agent, such as trimetazidine, is required because the compounds of the invention include multiple active agents. For example, trimetazidine shifts metabolism from fatty acid oxidation to glucose oxidation, and succinate improves mitochondrial respiration generally. Thus, a compound that provides both agents stimulates a larger increase in glucose-driven ATP production for a given amount of trimetazidine than does a compound that delivers trimetazidine alone.

[0150] The invention also provides compounds represented by formula (VII):

A-C

(VII),

in which A is a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, and C is a NAD⁺ precursor molecule. A and C may be covalently linked.

[0151] The compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation may be PEGylated with an ethylene glycol moiety. The compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation may have multiple ethylene glycol moieties, such as one, two, three, four, five, or more ethylene glycol moieties. The ethylene glycol moiety may be represented by (CH₂CH₂O)_x, in which x=1-15. The ethylene glycol moiety may form a covalent linkage between the compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation and the NAD⁺ precursor molecule. The ethylene glycol moiety may be separate from a covalent linkage between the compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation and the NAD⁺ precursor molecule.

[0152] The compound of formula (VII) may include nicotinic acid that is covalently linked to a PEGylated form of trimetazidine. The nicotinic acid may be covalently linked via a PEGylated moiety, i.e., via an ethylene glycol linkage. The nicotinic acid may be covalently linked via the trimetazidine moiety.

[0153] The invention also provides compounds represented by formula (VIII):

A-L-C

(VIII),

in which A is a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, L is a linker, and C is a NAD⁺ precursor molecule. A may be covalently linked to L, and L may be covalently linked to C.

[0154] The compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, the linker, and the NAD⁺ precursor molecule may be as described above in relation to compounds of other formulas.

[0155] The invention also provides compositions that include at least two of (1) a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, (2) a compound that promotes mitochondrial respiration, and (3) a NAD⁺ precursor molecule. The aforementioned components of the composition may be provided as separate molecules.

[0156] The compositions may include each of a (1) a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, (2) a compound that promotes mitochondrial respiration, and (3) a NAD⁺ precursor molecule. In such compositions, each of the three components may be provided as a separate molecule. Alternatively, in such compositions, two of the components may be covalently linked as part of single molecule, and the third component may be provided as a separate molecule. For example, the compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation may be linked to the compound that promotes mitochondrial respiration, and the NAD⁺ precursor may be provided as a separate molecule.

[0157] The compounds of the invention may be provided as co-crystals with other compounds. Co-crystals are crystalline materials composed of two or more different mol-

ecules in the same crystal lattice. The different molecules may be neutral and interact non-ionically within the lattice. Co-crystals of the invention may include one or more compounds of the invention with one or more other molecules that stimulate mitochondrial respiration or serve as NAD⁺ precursors. For example, a co-crystal may include any of the following combinations: (1) a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation and (2) a NAD⁺ precursor molecule; (1) a compound that promotes mitochondrial respiration and (2) a NAD⁺ precursor molecule; (1) a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation and (2) a compound that promotes mitochondrial respiration; (1) a molecule comprising a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation covalently linked to a compound that promotes mitochondrial respiration and (2) a NAD⁺ precursor molecule. In specific embodiments, a co-crystal may include (1) a compound of formula (I), (III), (IV), or (V) and (2) nicotinic acid, nicotinamide, or nicotinamide riboside.

[0158] The compounds may include one or more atoms that are enriched for an isotope. For example, the compounds may have one or more hydrogen atoms replaced with deuterium or tritium. Isotopic substitution or enrichment may occur at carbon, sulfur, or phosphorus, or other atoms. The compounds may be isotopically substituted or enriched for a given atom at one or more positions within the compound, or the compounds may be isotopically substituted or enriched at all instances of a given atom within the compound.

[0159] The invention provides pharmaceutical compositions containing one or more of the compounds described above. A pharmaceutical composition containing the compounds may be in a form suitable for oral use, for example, as tablets, troches, lozenges, fast-melts, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, syrups or elixirs. Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from sweetening agents, flavoring agents, coloring agents and preserving agents, in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the compounds in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration in the stomach and absorption lower down in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the techniques described in U.S. Pat. Nos. 4,256,108, 4,166,452 and 4,265,874, to form osmotic therapeutic tablets for control release. Preparation and administration of compounds is discussed in U.S. Pat. No. 6,214,841 and U.S. Pub. 2003/0232877, incorporated by reference herein in their entirety.

[0160] Formulations for oral use may also be presented as hard gelatin capsules in which the compounds are mixed with an inert solid diluent, for example calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the compounds are mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

[0161] An alternative oral formulation, where control of gastrointestinal tract hydrolysis of the compound is sought, can be achieved using a controlled-release formulation, where a compound of the invention is encapsulated in an enteric coating.

[0162] Aqueous suspensions may contain the compounds in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as a naturally occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example, polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such a polyoxyethylene with partial esters derived from fatty acids and hexitol anhydrides, for example polyoxyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

[0163] Oily suspensions may be formulated by suspending the compounds in a vegetable oil, for example, arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

[0164] Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the compounds in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified, for example sweetening, flavoring and coloring agents, may also be present.

[0165] The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally occurring phosphatides, for example soya bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

[0166] Syrups and elixirs may be formulated with sweetening agents, such as glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a

preservative, and agents for flavoring and/or coloring. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be in a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or di-glycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

[0167] The compounds of the invention are useful for improving cardiac efficiency. A variety of definitions of cardiac efficiency exist in the medical literature. See, e.g. Schipke, J. D. Cardiac efficiency, Basic Res. Cardiol. 89:207-40 (1994); and Gibbs, C. L. and Barclay, C. J. Cardiac efficiency, Cardiovasc. Res. 30:627-634 (1995), incorporated herein by reference. One definition of cardiac mechanical efficiency is the ratio of external cardiac power to cardiac energy expenditure by the left ventricle. See Lopaschuk G. D., et al., Myocardial Fatty Acid Metabolism in Health and Disease, Phys. Rev. 90:207-258 (2010), incorporated herein by reference. Another definition is the ratio between stroke work and oxygen consumption, which ranges from 20-25% in the normal human heart. Visser, F., Measuring cardiac efficiency: is it useful? Hear Metab. 39:3-4 (2008), incorporated herein by reference. Another definition is the ratio of the stroke volume to mean arterial blood pressure. Any suitable definition of cardiac efficiency may be used to measure the effects of compounds of the invention.

[0168] The invention also provides methods of altering cardiac metabolism in a subject to increase glucose oxidation relative to fatty acid oxidation. The methods may include providing a composition of the invention, such as any the compounds described above, including the compounds represented by formulas (I), (II), (III), (IV), or (V) or formulations thereof.

[0169] The methods may include providing a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, as described above, and a compound that promotes mitochondrial respiration, as described above. The compounds may be provided as components of a single molecule, as separate molecules in a single composition, or as separate compositions.

[0170] The methods may also include providing a NAD⁺ precursor molecule, as described above. In methods that involve providing a compound that shifts cardiac metabolism from fatty acid oxidation to glucose oxidation, a compound that promotes mitochondrial respiration, and a NAD⁺ precursor molecule, compounds may be provided as components of a single molecule, two different molecules, or three different molecules. The compounds may be provided in one, two, three, or any number of different compositions. The compounds may be provided together, separately, or in any combination. The compounds may be provided simultaneously or sequentially. The compounds may be provided at different intervals, with different frequency, in different quantities, or at different dosages.

[0171] The invention also provides methods of treating conditions by providing compositions of the invention. The condition may be heart disease, such as heart failure, ischemic heart disease, diabetic cardiomyopathy, rheumatic heart disease, valvular heart disease, aneurysm, atherosclerosis, high blood pressure (hypertension), peripheral arterial disease, angina, atherosclerosis, coronary artery disease, coronary heart disease, heart attack, atherosclerosis, cerebral vascular disease, stroke, transient ischemic attacks, atherosclerosis, cardiomyopathy, pericardial disease, valvular heart disease, or congenital heart disease.

Examples

Protocol

[0172] The effects of compounds of the invention on mitochondrial function were analyzed. HepG2 cells were dosed with test compound and in real time the extracellular oxygen levels and pH were measured using the XFe96 flux analyzer (Seahorse Biosciences). XFe Technology uses solid-state sensors to simultaneously measure both oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) to determine effects on oxidative phosphorylation (OXPHOS) and glycolysis simultaneously. The cells were then subjected to sequential exposure to various inhibitors of mitochondrial function to assess cellular metabolism.

Data Interpretation.

[0173] A compound was identified as positive mitochondrial-active compound when it caused a change in oxygen consumption rate (OCR) or extracellular acidification rate (ECAR) in the absence of cytotoxicity. Cytotoxicity was determined when both OXPHOS (OCR) and glycolysis (ECAR) were inhibited.

Definition of Mitochondrial Parameters.

[0174] Oxygen consumption rate (OCR) is a measurement of oxygen content in extracellular media. Changes in OCR indicate effects on mitochondrial function and can be bi-directional. A decrease is due to an inhibition of mitochondrial respiration, while an increase may indicate an uncoupler, in which respiration is not linked to energy production.

$$OCR = \frac{\text{compound } OCR - \text{non mitochondrial } OCR}{\text{basal } OCR - \text{non mitochondrial } OCR}$$

[0175] Extracellular acidification rate (ECAR) is the measurement of extracellular proton concentration (pH). An increase in signal means an increase in rate in number of pH ions (thus decreasing pH value) and seen as an increase in glycolysis. ECAR is expressed as a fraction of basal control (rate prior to addition of compound).

$$ECAR = \frac{\text{compound } ECAR}{\text{basal } ECAR}$$

[0176] Reserve capacity is the measured ability of cells to respond to an increase in energy demand. A reduction indicates mitochondrial dysfunction. This measurement demonstrates how close to the bioenergetic limit the cell is.

$$\text{reserve capacity} = \frac{FCCP \text{ } OCR - \text{non mitochondrial } OCR}{\text{basal } OCR - \text{non mitochondrial } OCR}$$

Mitochondrial Stress Test.

[0177] A series of compounds were added sequentially to the cells to assess a bioenergetics profile, effects of test compounds on parameters such as proton leak, and reserve capacity. This can be used to assist in understanding potential mechanisms of mitochondrial toxicity. The following compounds were added in order: (1) oligomycin, (2) FCCP, and (3) rotenone and antimycin A.

[0178] Oligomycin is a known inhibitor of ATP synthase and prevents the formation of ATP. Oligomycin treatment provides a measurement of the amount of oxygen consumption related to ATP production and ATP turnover. The addition of oligomycin results in a decrease in OCR under normal conditions, and residual OCR is related to the natural proton leak.

[0179] FCCP is a protonophore and is a known uncoupler of oxygen consumption from ATP production. FCCP treatment allows the maximum achievable transfer of electrons and oxygen consumption rate and provides a measurement of reserve capacity.

[0180] Rotenone and antimycin A are known inhibitors of complex I and III of the electron transport chain, respectively. Treatment with these compounds inhibits electron transport completely, and any residual oxygen consumption is due to non-mitochondrial activity via oxygen requiring enzymes.

Definition of Mechanisms.

[0181] An electron transport chain inhibitor is an inhibitor of mitochondrial respiration that causes an increase in glycolysis as an adaptive response (e.g. decrease OCR and increase in ECAR).

[0182] The inhibition of oxygen consumption may also be due to reduced substrate availability (e.g. glucose, fatty acids, glutamine, pyruvate), for example, via transporter inhibition. Compounds that reduce the availability of substrates are substrate inhibitors. A substrate inhibitor does not result in an increase in glycolysis (e.g. OCR decrease, no response in ECAR).

[0183] Compounds that inhibit the coupling of the oxidation process from ATP production are known as uncouplers. These result in an increase in mitochondrial respiration (OCR) but inhibition of ATP production.

[0184] FIG. 1 is a table summarizing the effects of various compounds on mitochondrial function.

[0185] FIG. 2 is a table summarizing the effects of nicotinamide on various mitochondrial functional parameters.

[0186] FIG. 3 is a series of graphs showing the effects of nicotinamide on oxygen consumption rate and reserve capacity.

[0187] FIG. 4 is a series of graphs showing the effects of nicotinamide on extracellular acidification rate.

[0188] FIG. 5 is a table summarizing the effects of a combination of trimetazidine and nicotinamide on various mitochondrial functional parameters.

[0189] FIG. 6 is a series of graphs showing the effects of a combination of trimetazidine and nicotinamide on oxygen consumption rate and reserve capacity.

[0190] FIG. 7 is a series of graphs showing the effects of a combination of trimetazidine and nicotinamide on extracellular acidification rate.

[0191] FIG. 8 is a table summarizing the effects of succinate on various mitochondrial functional parameters.

[0192] FIG. 9 is a series of graphs showing the effects of succinate on oxygen consumption rate and reserve capacity.

[0193] FIG. 10 is a series of graphs showing the effects of succinate on extracellular acidification rate.

[0194] FIG. 11 is a table summarizing the effects of compound CV-8814 on various mitochondrial functional parameters.

[0195] FIG. 12 is a series of graphs showing the effects of compound CV-8814 on oxygen consumption rate and reserve capacity.

[0196] FIG. 13 is a series of graphs showing the effects of compound CV-8814 on extracellular acidification rate.

[0197] FIG. 14 is a table summarizing the effects of trimetazidine on various mitochondrial functional parameters.

[0198] FIG. 15 is a series of graphs showing the effects of trimetazidine on oxygen consumption rate and reserve capacity.

[0199] FIG. 16 is a series of graphs showing the effects of trimetazidine on extracellular acidification rate.

[0200] FIG. 17 is a table summarizing the effects of a combination of succinate, nicotinamide, and trimetazidine on various mitochondrial functional parameters.

[0201] FIG. 18 is a series of graphs showing the effects of a combination of succinate, nicotinamide, and trimetazidine on oxygen consumption rate and reserve capacity.

[0202] FIG. 19 is a series of graphs showing the effects of a combination of succinate, nicotinamide, and trimetazidine on extracellular acidification rate.

[0203] FIG. 20 is a table summarizing the effects of a combination of trimetazidine analog 2 and nicotinamide on various mitochondrial functional parameters.

[0204] FIG. 21 is a series of graphs showing the effects of a combination of trimetazidine analog 2 and nicotinamide on oxygen consumption rate and reserve capacity.

[0205] FIG. 22 is a series of graphs showing the effects of a combination of trimetazidine analog 2 and nicotinamide on extracellular acidification rate.

[0206] FIG. 23 is a table summarizing the effects of a combination of trimetazidine analog 1 and nicotinamide on various mitochondrial functional parameters.

[0207] FIG. 24 is a series of graphs showing the effects of a combination of trimetazidine analog 1 and nicotinamide on oxygen consumption rate and reserve capacity.

[0208] FIG. 25 is a series of graphs showing the effects of a combination of trimetazidine analog 1 and nicotinamide on extracellular acidification rate.

[0209] FIG. 26 is a table summarizing the effects of a combination of trimetazidine analog 3 and nicotinamide on various mitochondrial functional parameters.

[0210] FIG. 27 is a series of graphs showing the effects of a combination of trimetazidine analog 3 and nicotinamide on oxygen consumption rate and reserve capacity.

[0211] FIG. 28 is a series of graphs showing the effects of a combination of trimetazidine analog 3 and nicotinamide on extracellular acidification rate.

[0212] FIG. 29 is a table summarizing the effects of a combination of succinate and nicotinamide on various mitochondrial functional parameters.

[0213] FIG. 30 is a series of graphs showing the effects of a combination of succinate and nicotinamide on oxygen consumption rate and reserve capacity.

[0214] FIG. 31 is a series of graphs showing the effects of a combination of succinate and nicotinamide on extracellular acidification rate.

Effect of Compositions on Coronary Flow, Cardiac Function, and Infarct Size.

[0215] The effect of compositions on the coronary flow, cardiac function, and infarct size was analyzed.

[0216] FIG. 32 is a schematic of the ischemia-reperfusion (IR) method used to analyze the effects of compositions of the invention on coronary flow, cardiac function, and infarct size. At time 0, mice were given (1) 20 μ M trimetazidine (TMZ), (2) 2 μ M each of trimetazidine, nicotinamide, and succinate (TNF), (3) 20 μ M each of trimetazidine, nicotinamide, and succinate (TNS), or (4) the delivery vehicle (CON). At 20 minutes, ischemia was induced, and coronary flow was analyzed. At 50 minutes, reperfusion was initiated to restore blood flow. At 170 minutes, coronary flow and cardiac function was analyzed, and then the hearts were preserved, sectioned, and infarct size was measured by triphenyltetrazolium chloride (TTC) staining.

[0217] FIG. 33 is a graph of coronary flow of after IR. Data is expressed as ratio cardiac flow at 170 minutes to cardiac flow at 20 minutes. TNS treatment preserved coronary flow after IR. Raw data is provided in Tables 1-2.

TABLE 1

	CF20 (ml/min)	CF170 (ml/min)	CF170/CF20 (ul/ml)
CON11	2.31E+00	1.11E-01	4.81E+01
CON13	1.07E+00	4.80E-02	4.48E+01
CON14	8.28E-01	4.50E-02	5.43E+01
CON9	2.11E+00	6.96E-02	3.30E+01
CON10	1.85E+00	4.92E-02	2.66E+01
CON7	1.57E+00	5.40E-02	3.44E+01
CON8	3.22E+00	6.78E-02	2.11E+01
CON5	2.18E+00	6.60E-02	3.03E+01
CON3	2.24E+00	7.92E-02	3.53E+01
CON4	2.22E+00	7.84E-02	3.53E+01
CON2	1.68E+00	5.12E-02	3.05E+01
MEAN	1.93E+00	6.54E-02	3.58E+01
SD	6.50E-01	1.94E-02	9.72E+00
SE	1.96E-01	5.86E-03	2.93E+00
TTEST			
TMZ4	2.13E+00	5.16E-02	2.42E+01
TMZ3	1.70E+00	1.00E-01	5.87E+01
TMZ1	2.18E+00	7.78E-02	3.57E+01

TABLE 1-continued

	CF20 (ml/min)	CF170 (ml/min)	CF170/CF20 (ul/ml)
TMZ2	3.83E+00	1.29E-01	3.37E+01
TMZ7	1.72E+00	8.98E-02	5.21E+01
TMZ8	2.40E+00	6.56E-02	2.73E+01
TMZ5	2.14E+00	5.56E-02	2.60E+01
TMZ9	2.03E+00	1.30E-01	6.39E+01
MEAN	2.27E+00	8.74E-02	4.02E+01
SD	6.75E-01	3.06E-02	1.57E+01
SE	2.39E-01	1.08E-02	5.56E+00
TTEST			
TNF1	2.24E+00	4.80E-02	2.14E+01
TNF2	2.24E+00	3.80E-02	1.69E+01
TNF3	7.32E-01	4.80E-02	6.56E+01
TNF4	8.20E-01	4.90E-02	5.98E+01
TNF5	1.09E+00	2.70E-02	2.48E+01
TNF6	9.48E-01	1.50E-01	1.58E+02
TNF7	8.08E-01	3.70E-02	4.58E+01
TNF8	1.20E+00	4.60E-02	3.83E+01
TNF9	1.45E+00	1.21E-01	8.33E+01
TNF10	1.20E+00	1.52E-02	1.27E+01
MEAN	1.27E+00	5.79E-02	5.27E+01
SD	5.56E-01	4.28E-02	4.37E+01
SE	1.76E-01	1.35E-02	1.38E+01
TTEST			
TNS1	1.52E+00	4.70E-02	3.08E+01
TNS2	9.30E-01	2.90E-02	3.12E+01
TNS3	2.24E+00	1.67E-01	7.46E+01
TNS5	5.64E-01	5.00E-02	8.87E+01
TNS6	6.28E-01	4.40E-02	7.01E+01
TNS7	1.08E+00	6.40E-02	5.95E+01
TNS8	8.72E-01	2.30E-02	2.64E+01
TNS9	1.18E+00	8.50E-02	7.23E+01
TNS10	1.70E+00	1.84E-01	1.08E+02
MEAN	1.19E+00	7.70E-02	6.24E+01
SD	5.43E-01	5.89E-02	2.82E+01
SE	1.81E-01	1.96E-02	9.42E+00
TTEST			
vs TMZ			6.82E-02

TABLE 2

	CON	TMZ	TNF	TNS
MEAN	36	40	53	62
SD	10	16	44	28
SE	3	6	14	9

[0218] FIG. 34 is graph of left ventricular developed pressure (LVDP) after IR. Blue bars indicate LVDP at 20 minutes, and orange bars indicate LVDP at 170 minutes. TMZ, TNS, and

[0219] TNF treatment prevented a decline in cardiac function after IR. Raw data is provided in Tables 3-6.

TABLE 3

pre-ischemia	LVESP	LVEDP	HR	LVDP	LVDP \times HR
5-18-CN	CON11	6.61E+01	6.20E+00	3.28E+02	5.99E+01
	CON12	8.15E+01	3.73E+00	3.56E+02	7.78E+01
6-10-CN	CON13	8.00E+01	-3.74E+00	1.37E+02	8.37E+01
	CON14	7.28E+01	6.12E+00	4.54E+02	6.67E+01
5-15-CN	CON9	8.07E+01	5.00E+00	1.42E+02	7.57E+01
	CON10	4.91E+01	1.15E+00	3.21E+02	4.80E+01
5-12-CN	CON7	8.55E+01	6.35E+00	3.05E+02	7.91E+01
	CON8	5.06E+01	1.68E+00	3.04E+02	4.90E+01
5-9-CN	CON5	5.45E+01	5.63E+00	2.75E+02	4.89E+01
	CON6	6.37E+01	4.31E+00	3.08E+02	5.94E+01

TABLE 3-continued

	pre-ischemia	LVESP	LVEDP	HR	LVDP	LVDP × HR
5-7-CN	CON3	7.32E+01	2.70E+00	2.40E+02	7.05E+01	1.69E+04
	CON4	4.91E+01	1.65E-01	3.14E+02	4.89E+01	1.54E+04
5-5-CN	CON1	9.48E+01	7.96E+00	3.04E+02	8.68E+01	2.64E+04
	CON2	4.69E+01	1.64E-01	4.02E+02	4.67E+01	1.88E+04
5-11-CN	MEAN	6.77E+01	3.39E+00	2.99E+02	6.44E+01	1.88E+04
	SD	1.58E+01	3.21E+00	8.52E+01	1.46E+01	6.12E+03
5-14-TMZ	SE	4.21E+00	8.57E-01	2.28E+01	3.91E+00	1.63E+03
	TTEST				2.42E-04	
5-11-TMZ	TMZ3	7.58E+01	6.53E+00	2.63E+02	6.93E+01	1.83E+04
	TMZ4	8.44E+01	5.43E+00	2.93E+02	7.90E+01	2.31E+04
5-8-TMZ	TMZ1	7.15E+01	6.76E+00	1.66E+02	6.48E+01	1.08E+04
	TMZ2	5.47E+01	1.74E+00	3.35E+02	5.30E+01	1.77E+04
5-6-TMZ	TMZ7	6.87E+01	3.58E+00	3.58E+02	6.51E+01	2.33E+04
	TMZ8	4.27E+01	4.71E+00	3.33E+02	3.80E+01	1.26E+04
5-4-TMZ	TMZ5	3.30E+01	4.77E+00	3.48E+02	2.82E+01	9.82E+03
	TMZ6	3.30E+01	1.46E+00	3.21E+02	3.15E+01	1.01E+04
5-19-TNF	TMZ9	6.60E+01	7.25E+00	2.67E+02	5.87E+01	1.57E+04
	TMZ10	7.38E+01	2.70E+00	3.32E+02	7.11E+01	2.36E+04
6-8-TNF	MEAN	6.03E+01	4.49E+00	3.02E+02	5.59E+01	1.68E+04
	SD	1.85E+01	2.07E+00	5.75E+01	1.77E+01	5.56E+03
6-12-TNF	SE	5.84E+00	6.56E-01	1.82E+01	5.58E+00	1.76E+03
					3.85E-01	
6-14-TNF	TNF1	5.02E+01	3.04E+00	4.09E+02	4.72E+01	1.93E+04
	TNF2	4.65E+01	1.76E-01	2.76E+02	4.63E+01	1.28E+04
6-15-TNF	TNF3	7.13E+01	1.53E+00	6.48E+01	6.97E+01	4.52E+03
	TNF4	9.97E+01	4.15E+00	1.54E+02	9.55E+01	1.47E+04
6-18-TNF	TNF5	7.14E+01	-3.42E+00	2.77E+02	7.49E+01	2.07E+04
	TNF6	8.98E+01	8.85E+00	3.10E+02	8.09E+01	2.51E+04
6-19-TNF	TNF7	6.58E+01	7.01E+00	3.98E+02	5.88E+01	2.34E+04
	TNF8	5.99E+01	1.02E+00	2.28E+02	5.89E+01	1.34E+04
6-20-TNS	TNF9	7.89E+01	2.37E-01	2.71E+02	7.87E+01	2.13E+04
	TNF10	4.01E+01	1.88E+00	3.14E+02	3.82E+01	1.20E+04
6-21-TNS	MEAN	6.74E+01	2.45E+00	2.70E+02	6.49E+01	1.67E+04
	SD	1.90E+01	3.54E+00	1.04E+02	1.81E+01	6.32E+03
6-22-TNS	SE	6.00E+00	1.12E+00	3.28E+01	5.73E+00	2.00E+03
					1.38E-01	
6-23-TNS	TNS1	5.59E+01	5.23E+00	3.33E+02	5.07E+01	1.69E+04
	TNS2	5.54E+01	-1.83E+00	1.24E+02	5.72E+01	7.09E+03
6-24-TNS	TNS3	8.78E+01	1.53E+00	1.64E+02	8.63E+01	1.42E+04
	TNS4	1.07E+02	9.86E+00	2.41E+02	9.74E+01	2.35E+04
6-25-TNS	TNS5	8.97E+01	2.34E+00	8.35E+01	8.74E+01	7.29E+03
	TNS6	6.17E+01	6.21E+00	1.85E+02	5.55E+01	1.03E+04
6-26-TNS	TNS7	6.62E+01	4.14E+00	3.36E+02	6.21E+01	2.09E+04
	TNS8	6.54E+01	1.22E+01	1.22E+02	5.32E+01	6.47E+03
6-27-TNS	TNS9	6.16E+01	3.64E+00	3.45E+02	5.80E+01	2.00E+04
	TNS10	5.44E+01	2.47E+00	4.12E+02	5.20E+01	2.14E+04
6-28-TNS	MEAN	7.05E+01	4.58E+00	2.35E+02	6.60E+01	1.48E+04
	SD	1.80E+01	4.09E+00	1.15E+02	1.74E+01	6.61E+03
6-29-TNS	SE	5.69E+00	1.29E+00	3.63E+01	5.49E+00	2.09E+03
					7.89E-02	

TABLE 4

	after 2 h reperfusion	LVESP	LVEDP	HR	LVDP	LVDP × HR
5-18-CN	CON11	7.78E+01	3.68E+01	1.18E+02	4.10E+01	4.82E+03
	CON12	7.07E+01	2.23E+01	9.23E+01	4.84E+01	4.47E+03
6-10-CN	CON13	6.48E+01	5.54E+01	5.72E+02	9.39E+00	5.38E+03
	CON14	9.54E+01	5.64E+01	2.08E+02	3.90E+01	8.12E+03
5-15-CN	CON9	5.18E+01	2.71E+01	1.75E+02	2.47E+01	4.33E+03
	CON10	1.10E+02	3.13E+01	5.76E+01	7.84E+01	4.51E+03
5-12-CN	CON7	3.93E+01	1.42E+01	9.11E+01	2.51E+01	2.29E+03
	CON8	5.29E+01	9.48E+00	6.07E+01	4.34E+01	2.64E+03
5-9-CN	CON5	6.56E+01	4.89E+01	6.50E+01	1.67E+01	1.09E+03
	CON6	7.44E+01	6.56E+01	3.78E+01	8.81E+00	3.33E+02

TABLE 4-continued

		after 2 h reperfusion	LVESP	LVEDP	HR	LVDP	LVDP × HR
5-7-CN	CON3	6.35E+01	9.99E+00	1.15E+02	5.35E+01	6.18E+03	
	CON4	8.76E+01	5.34E+01	1.06E+02	3.43E+01	3.65E+03	
	CON1	9.29E+01	4.38E+01	2.61E+02	4.91E+01	1.28E+04	
	CON2	5.18E+01	4.43E+00	2.57E+02	4.74E+01	1.22E+04	
	MEAN	7.13E+01	3.42E+01	1.58E+02	3.71E+01	5.20E+03	
	SD	1.98E+01	2.02E+01	1.39E+02	1.90E+01	3.68E+03	
	SE	5.29E+00	5.40E+00	3.72E+01	5.08E+00	9.83E+02	
	TTEST						
	5-14-TMZ	TMZ3	5.07E+01	2.93E+01	1.18E+02	2.14E+01	2.52E+03
		TMZ4	7.66E+01	3.31E+01	1.19E+02	4.34E+01	5.15E+03
5-11-TMZ	TMZ1	9.19E+01	3.96E+01	1.01E+02	5.22E+01	5.28E+03	
	TMZ2	4.77E+01	1.80E+01	1.51E+02	2.97E+01	4.49E+03	
5-8-TMZ	TMZ7	5.18E+01	3.36E+00	6.70E+01	4.84E+01	3.24E+03	
	TMZ8	4.86E+01	1.87E+00	9.22E+01	4.67E+01	4.31E+03	
5-6-TMZ	TMZ5	6.09E+01	1.99E+01	2.22E+02	4.10E+01	9.11E+03	
	TMZ6	1.09E+02	3.21E+01	1.70E+02	7.65E+01	1.30E+04	
5-4-TMZ	TMZ9	7.38E+01	1.84E+01	1.16E+02	5.53E+01	6.44E+03	
	TMZ10	7.61E+01	1.77E+00	2.38E+02	7.43E+01	1.77E+04	
	MEAN	6.86E+01	1.97E+01	1.39E+02	4.89E+01	6.82E+03	
	SD	2.05E+01	1.39E+01	5.58E+01	1.73E+01	4.82E+03	
	SE	6.49E+00	4.39E+00	1.77E+01	5.46E+00	1.52E+03	
5-19-TNF	TNF1	8.37E+01	6.66E+01	1.53E+02	1.71E+01	2.62E+03	
	TNF2	6.19E+00	5.54E+00	2.13E+03	6.48E-01	1.38E+03	
6-8-TNF	TNF3	8.99E+01	1.88E+01	1.05E+01	7.11E+01	7.49E+02	
	TNF4	6.06E+01	1.34E+01	8.10E+01	4.72E+01	3.82E+03	
6-12-TNF	TNF5	1.54E+02	4.15E+01	2.20E+01	1.13E+02	2.48E+03	
	TNF6	1.30E+02	4.25E+01	3.33E+01	8.77E+01	2.92E+03	
6-14-TNF	TNF7	5.70E+01	4.00E+01	4.00E+01	1.70E+01	6.80E+02	
	TNF8	3.76E+01	1.87E+01	5.36E+01	1.88E+01	1.01E+03	
6-15-TNF	TNF9	6.23E+01	3.38E+01	1.97E+02	2.85E+01	5.59E+03	
	TNF10	7.85E+01	2.75E+01	7.85E+01	5.10E+01	4.00E+03	
	MEAN	7.60E+01	3.09E+01	2.80E+02	4.52E+01	2.53E+03	
	SD	4.28E+01	1.79E+01	6.54E+02	3.59E+01	1.62E+03	
	SE	1.35E+01	5.65E+00	2.07E+02	1.14E+01	5.12E+02	
5-20-TNS	TNS1	6.47E+01	1.78E+01	1.04E+02	4.69E+01	4.88E+03	
	TNS2	8.95E+01	3.03E+01	5.55E+01	5.92E+01	3.29E+03	
6-7-TNS	TNS3	7.79E+01	6.34E+01	1.28E+02	1.45E+01	1.85E+03	
	TNS4	7.74E+01	2.73E+01	1.02E+02	5.01E+01	5.09E+03	
6-9-TNS	TNS5	1.37E+02	5.63E+01	1.63E+01	8.08E+01	1.32E+03	
	TNS6	8.59E+01	1.23E+01	1.06E+02	7.36E+01	7.79E+03	
6-13-TNS	TNS7	5.76E+01	5.16E+01	1.35E+02	6.00E+00	8.07E+02	
	TNS8	4.96E+01	1.53E+01	1.22E+02	3.43E+01	4.20E+03	
6-15-TNS	TNS9	9.97E+01	3.00E+01	7.46E+01	6.98E+01	5.21E+03	
	TNS10	4.32E+01	-4.32E+00	7.20E+01	4.75E+01	3.42E+03	
	MEAN	7.83E+01	3.00E+01	9.15E+01	4.83E+01	3.79E+03	
	SD	2.74E+01	2.14E+01	3.69E+01	2.45E+01	2.11E+03	
	SE	8.67E+00	6.78E+00	1.17E+01	7.75E+00	6.69E+02	

TABLE 5

	pre-ischemia	+dp/dtm	-dp/dtm	after 2 h reperfusion	+dp/dtm	-dp/dtm
5-18-CN	CON11	2.60E+03	-1.82E+03	CON11	1.44E+03	-8.67E+02
	CON12	2.95E+03	-2.58E+03	CON12	1.63E+03	-1.07E+03
6-10-CN	CON13	3.10E+03	-2.42E+03	CON13	2.25E+02	-2.22E+02
	CON14	3.08E+03	-2.10E+03	CON14	3.44E+02	-2.87E+02
5-15-CN	CON9	2.28E+03	-1.38E+03	CON9	9.45E+02	-5.54E+02
	CON10	2.06E+03	-1.50E+03	CON10	2.29E+03	-1.75E+03
5-12-CN	CON7	2.71E+03	-2.10E+03	CON7	2.51E+02	-2.55E+02
	CON8	1.58E+03	-1.10E+03	CON8	3.63E+02	-3.05E+02
5-9-CN	CON5	2.17E+03	-1.50E+03	CON5	2.39E+02	-2.41E+02
	CON6	2.25E+03	-1.62E+03	CON6	1.47E+02	-1.49E+02

TABLE 5-continued

	pre-ischemia			after 2 h		
		+dp/dtm	-dp/dtm	reperfusion	+dp/dtm	-dp/dtm
5-7-CN	CON3	2.63E+03	-2.06E+03	CON3	1.63E+03	-1.06E+03
	CON4	2.05E+03	-1.38E+03	CON4	1.10E+03	-7.03E+02
5-5-CN	CON1	3.17E+03	-2.37E+03	CON1	1.03E+03	-1.12E+03
	CON2	2.10E+03	-1.50E+03	CON2	1.75E+03	-1.27E+03
	MEAN	2.48E+03	-1.82E+03	MEAN	9.56E+02	-7.04E+02
	SD	4.84E+02	4.56E+02	SD	7.08E+02	4.95E+02
	SE	1.29E+02	1.22E+02	SE	1.89E+02	1.32E+02
	TTEST			TTEST		
5-14-TMZ	TMZ3	2.41E+03	-1.69E+03	TMZ3	4.14E+02	-3.57E+02
	TMZ4	2.77E+03	-2.26E+03	TMZ4	1.48E+03	-1.15E+03
5-11-TMZ	TMZ1	1.80E+03	-1.59E+03	TMZ1	1.38E+03	-7.45E+02
	TMZ2	2.15E+03	-1.80E+03	TMZ2	1.06E+03	-6.85E+02
5-8-TMZ	TMZ7	3.40E+03	-2.59E+03	TMZ7	3.44E+02	-3.39E+02
	TMZ8	1.75E+03	-1.20E+03	TMZ8	7.36E+02	-4.28E+02
5-6-TMZ	TMZ5	1.27E+03	-8.82E+02	TMZ5	1.28E+03	-8.38E+02
	TMZ6	1.24E+03	-6.59E+02	TMZ6	1.85E+03	-1.06E+03
5-4-TMZ	TMZ9	1.98E+03	-1.41E+03	TMZ9	1.13E+03	-6.38E+02
	TMZ10	2.02E+03	-1.56E+03	TMZ10	1.62E+03	-9.83E+02
	MEAN	2.08E+03	-1.56E+03	MEAN	1.13E+03	-7.22E+02
	SD	6.58E+02	5.81E+02	SD	5.01E+02	2.90E+02
	SE	2.08E+02	1.84E+02	SE	1.58E+02	9.16E+01
					5.16E-01	9.18E-01
5-19-TNF	TNF1	2.67E+03	-1.49E+03	TNF1	3.86E+02	-3.75E+02
	TNF2	2.85E+03	-1.44E+03	TNF2	1.46E+02	-1.43E+02
6-8-TNF	TNF3	1.53E+03	-7.24E+02	TNF3	2.28E+02	-2.34E+02
	TNF4	3.86E+03	-2.59E+03	TNF4	2.84E+02	-2.40E+02
6-12-TNF	TNF5	3.29E+03	-2.34E+03	TNF5	2.92E+03	-2.08E+03
	TNF6	3.03E+03	-1.90E+03	TNF6	2.48E+03	-1.84E+03
6-14-TNF	TNF7	3.22E+03	-1.62E+03	TNF7	2.53E+02	-2.48E+02
	TNF8	1.74E+03	-1.12E+03	TNF8	1.53E+02	-1.52E+02
6-15-TNF	TNF9	2.14E+03	-2.33E+03	TNF9	1.04E+03	-6.31E+02
	TNF10	1.86E+03	-9.97E+02	TNF10	2.04E+03	-1.34E+03
	MEAN	2.62E+03	-1.65E+03	MEAN	9.93E+02	-7.29E+02
	SD	7.71E+02	6.26E+02	SD	1.08E+03	7.43E+02
	SE	2.44E+02	1.98E+02	SE	3.41E+02	2.35E+02
		1.09E-03	7.48E-03			
5-20-TNS	TNS1	2.37E+03	-1.60E+03	TNS1	1.79E+03	-1.12E+03
	TNS2	2.87E+03	-2.53E+03	TNS2	1.84E+03	-1.30E+03
6-7-TNS	TNS3	4.00E+03	-2.67E+03	TNS3	2.91E+02	-3.02E+02
	TNS4	3.32E+03	-2.63E+03	TNS4	1.62E+03	-1.30E+03
6-9-TNS	TNS5	3.36E+03	-2.21E+03	TNS5	2.43E+02	-2.46E+02
	TNS6	2.53E+03	-1.89E+03	TNS6	2.36E+03	-1.74E+03
6-13-TNS	TNS7	2.92E+03	-1.75E+03	TNS7	2.49E+02	-2.47E+02
	TNS8	1.12E+03	-7.42E+02	TNS8	1.29E+03	-8.50E+02
6-15-TNS	TNS9	2.29E+03	-1.75E+03	TNS9	2.06E+03	-1.59E+03
	TNS10	2.11E+03	-1.58E+03	TNS10	1.26E+03	-1.10E+03
	MEAN	2.69E+03	-1.94E+03	MEAN	1.30E+03	-9.80E+02
	SD	7.99E+02	5.95E+02	SD	7.86E+02	5.52E+02
	SE	2.53E+02	1.88E+02	SE	2.49E+02	1.75E+02
		9.96E-04	1.55E-03			

TABLE 6

		CON	TMZ	TNF	TNS
T20	Mean	64.36	55.86	64.90	65.96
T20	SE	3.91	5.58	5.73	5.49
T170	Mean	37.09	48.91	45.16	48.27
T170	SE	5.08	5.46	11.36	7.75

[0220] FIG. 35 shows images of TTC-stained heart slices after IR. TMZ and TNS treatment decreased infarct size after IR.

[0221] FIG. 36 is graph of infarct size after IR. TMZ and TNS treatment decreased infarct size after IR. Raw data is provided in Tables 7-55.

TABLE 7

CN11 raw values			
1	Slide11.jpg	1649	
2	Slide11.jpg	10	0.06
3	Slide11.jpg	1385	8.40
4	Slide11.jpg	2808	
5	Slide11.jpg	104	0.81
6	Slide11.jpg	2525	19.78
7	Slide11.jpg	3807	
8	Slide11.jpg	1014	7.99
9	Slide11.jpg	2207	17.39
10	Slide11.jpg	3952	
11	Slide11.jpg	15	0.08
12	Slide11.jpg	3300	17.54
13	Slide11.jpg	3376	
14	Slide11.jpg	103	0.92
15	Slide11.jpg	2816	25.02
16	Slide11.jpg	1616	

TABLE 7-continued

CN11 raw values			
17	Slide11.jpg	975	6.03
18	Slide11.jpg	409	2.53
19	Slide11.jpg	2805	
20	Slide11.jpg	819	6.42
21	Slide11.jpg	1496	11.73
22	Slide11.jpg	3973	
23	Slide11.jpg	1047	7.91
24	Slide11.jpg	2465	18.61
25	Slide11.jpg	3971	
26	Slide11.jpg	1102	5.83
27	Slide11.jpg	2430	12.85
28	Slide11.jpg	3516	
29	Slide11.jpg	1919	16.37
30	Slide11.jpg	920	7.85

TABLE 8

CN11 summary	
non-IS	26.21
IS	70.86
LV	97.07
IS/LV	73%

TABLE 9

CN12 raw values			
1	Slide12.jpg	1562	
2	Slide12.jpg	1059	8.81
3	Slide12.jpg	485	4.04
4	Slide12.jpg	2925	
5	Slide12.jpg	260	1.78
6	Slide12.jpg	2159	14.76
7	Slide12.jpg	3492	
8	Slide12.jpg	263	1.88
9	Slide12.jpg	2886	20.66
10	Slide12.jpg	4855	
11	Slide12.jpg	1992	16.00
12	Slide12.jpg	2292	18.41
13	Slide12.jpg	2934	
14	Slide12.jpg	1405	6.70
15	Slide12.jpg	914	4.36
16	Slide12.jpg	2061	
17	Slide12.jpg	81	0.51
18	Slide12.jpg	1704	10.75
19	Slide12.jpg	2966	
20	Slide12.jpg	105	0.71
21	Slide12.jpg	2810	18.95
22	Slide12.jpg	4099	
23	Slide12.jpg	823	5.02
24	Slide12.jpg	2350	14.33
25	Slide12.jpg	3979	
26	Slide12.jpg	357	3.50
27	Slide12.jpg	2787	27.32
28	Slide12.jpg	2974	
29	Slide12.jpg	490	2.31
30	Slide12.jpg	2112	9.94

TABLE 10

CN12 summary	
non-IS	23.61
IS	71.76
LV	95.37
IS/LV	75%

TABLE 11

TNS1 raw values			
1	Slide15.jpg	1857	
2	Slide15.jpg	58	0.28
3	Slide15.jpg	1672	8.10
4	Slide15.jpg	3383	
5	Slide15.jpg	901	4.53
6	Slide15.jpg	1873	9.41
7	Slide15.jpg	3460	
8	Slide15.jpg	1452	13.43
9	Slide15.jpg	2272	21.01
10	Slide15.jpg	3712	
11	Slide15.jpg	772	8.32
12	Slide15.jpg	2422	26.10
13	Slide15.jpg	3088	
14	Slide15.jpg	498	3.87
15	Slide15.jpg	1733	13.47
16	Slide15.jpg	1762	
17	Slide15.jpg	65	0.33
18	Slide15.jpg	1626	8.31
19	Slide15.jpg	3532	
20	Slide15.jpg	2034	9.79
21	Slide15.jpg	1206	5.80
22	Slide15.jpg	3411	
23	Slide15.jpg	1752	16.44
24	Slide15.jpg	1006	9.44
25	Slide15.jpg	4241	
26	Slide15.jpg	2148	20.26
27	Slide15.jpg	1101	10.38
28	Slide15.jpg	3440	
29	Slide15.jpg	2307	16.10
30	Slide15.jpg	165	1.15

TABLE 12

TNS1 summary	
non-IS	46.67
IS	56.59
LV	103.26
IS/LV	55%

TABLE 13

TNS2 raw values			
1	Slide16.jpg	1565	
2	Slide16.jpg	1058	7.44
3	Slide16.jpg	145	1.02
4	Slide16.jpg	2654	
5	Slide16.jpg	431	3.90
6	Slide16.jpg	2043	18.47
7	Slide16.jpg	3247	
8	Slide16.jpg	1053	8.43
9	Slide16.jpg	1584	12.68
10	Slide16.jpg	3892	
11	Slide16.jpg	2391	22.73
12	Slide16.jpg	863	8.20
13	Slide16.jpg	2505	
14	Slide16.jpg	1488	14.85
15	Slide16.jpg	363	3.62
16	Slide16.jpg	1526	
17	Slide16.jpg	9	0.06
18	Slide16.jpg	1357	9.78
19	Slide16.jpg	2337	
20	Slide16.jpg	16	0.16
21	Slide16.jpg	1899	19.50
22	Slide16.jpg	3558	
23	Slide16.jpg	1453	10.62
24	Slide16.jpg	1504	10.99
25	Slide16.jpg	4041	
26	Slide16.jpg	517	4.73

TABLE 13-continued

TNS2 raw values			
27	Slide16.jpg	2763	25.30
28	Slide16.jpg	2946	
29	Slide16.jpg	631	5.35
30	Slide16.jpg	1326	11.25

TABLE 14

TNS2 summary			
non-IS		39.14	
IS		60.41	
LV		99.56	
IS/LV		61%	

TABLE 15

TNF1 raw values			
1	Slide17.jpg	1326	
2	Slide17.jpg	63	0.24
3	Slide17.jpg	1183	4.46
4	Slide17.jpg	3158	
5	Slide17.jpg	825	5.49
6	Slide17.jpg	2014	13.39
7	Slide17.jpg	4805	
8	Slide17.jpg	1774	12.92
9	Slide17.jpg	1722	12.54
10	Slide17.jpg	4675	
11	Slide17.jpg	1984	15.28
12	Slide17.jpg	2470	19.02
13	Slide17.jpg	2754	
14	Slide17.jpg	269	2.05
15	Slide17.jpg	1377	10.50
16	Slide17.jpg	1373	
17	Slide17.jpg	1067	3.89
18	Slide17.jpg	43	0.16
19	Slide17.jpg	3113	
20	Slide17.jpg	803	5.42
21	Slide17.jpg	2008	13.55
22	Slide17.jpg	4657	
23	Slide17.jpg	1189	8.94
24	Slide17.jpg	2398	18.02
25	Slide17.jpg	4607	
26	Slide17.jpg	1256	9.81
27	Slide17.jpg	1978	15.46
28	Slide17.jpg	2769	
29	Slide17.jpg	2115	16.04
30	Slide17.jpg	72	0.55

TABLE 16

TNF1 summary			
non-IS		40.03	
IS		53.82	
LV		93.86	
IS/LV		57%	

TABLE 17

TNF2 raw values			
1	Slide18.jpg	2133	
2	Slide18.jpg	1861	12.21
3	Slide18.jpg	239	1.57
4	Slide18.jpg	4037	

TABLE 17-continued

TNF2 raw values			
5	Slide18.jpg	753	5.60
6	Slide18.jpg	2304	17.12
7	Slide18.jpg	4663	
8	Slide18.jpg	1548	10.62
9	Slide18.jpg	2917	20.02
10	Slide18.jpg	5017	
11	Slide18.jpg	2648	20.06
12	Slide18.jpg	2480	18.78
13	Slide18.jpg	3629	
14	Slide18.jpg	1698	13.10
15	Slide18.jpg	348	2.69
16	Slide18.jpg	2130	
17	Slide18.jpg	4	0.03
18	Slide18.jpg	1988	13.07
19	Slide18.jpg	4108	
20	Slide18.jpg	253	1.85
21	Slide18.jpg	3796	27.72
22	Slide18.jpg	4612	
23	Slide18.jpg	815	5.65
24	Slide18.jpg	2427	16.84
25	Slide18.jpg	4880	
26	Slide18.jpg	562	4.38
27	Slide18.jpg	3535	27.53
28	Slide18.jpg	3507	
29	Slide18.jpg	497	3.97
30	Slide18.jpg	1837	14.67

TABLE 18

TNF2 summary			
non-IS		38.73	
IS		80.00	
LV		118.73	
IS/LV		73%	

TABLE 19

TNS3 raw values			
1	Slide19.jpg	1484	
2	Slide19.jpg	923	4.98
3	Slide19.jpg	714	3.85
4	Slide19.jpg	3124	
5	Slide19.jpg	990	6.65
6	Slide19.jpg	1845	12.40
7	Slide19.jpg	3414	
8	Slide19.jpg	1282	13.89
9	Slide19.jpg	1833	19.87
10	Slide19.jpg	3380	
11	Slide19.jpg	2123	16.33
12	Slide19.jpg	1042	8.02
13	Slide19.jpg	2105	
14	Slide19.jpg	957	7.73
15	Slide19.jpg	308	2.49
16	Slide19.jpg	1524	
17	Slide19.jpg	10	0.05
18	Slide19.jpg	1530	8.03
19	Slide19.jpg	2860	
20	Slide19.jpg	13	0.10
21	Slide19.jpg	2293	16.84
22	Slide19.jpg	3358	
23	Slide19.jpg	960	10.58
24	Slide19.jpg	2639	29.08
25	Slide19.jpg	2538	
26	Slide19.jpg	296	3.03

TABLE 19-continued

TNS3 raw values			
27	Slide19.jpg	1797	18.41
28	Slide19.jpg	1992	
29	Slide19.jpg	1105	9.43
30	Slide19.jpg	401	3.42

TABLE 20

TNS3 summary			
non-IS		36.39	
IS		61.20	
LV		97.58	
IS/LV		63%	

TABLE 21

TNS4 raw values			
1	Slide20.jpg	1524	
2	Slide20.jpg	47	0.28
3	Slide20.jpg	1417	8.37
4	Slide20.jpg	2478	
5	Slide20.jpg	582	5.17
6	Slide20.jpg	1617	14.36
7	Slide20.jpg	3284	
8	Slide20.jpg	1226	11.20
9	Slide20.jpg	2072	18.93
10	Slide20.jpg	3639	
11	Slide20.jpg	771	7.20
12	Slide20.jpg	2177	20.34
13	Slide20.jpg	3114	
14	Slide20.jpg	491	5.36
15	Slide20.jpg	2189	23.90
16	Slide20.jpg	1648	
17	Slide20.jpg	1244	6.79
18	Slide20.jpg	94	0.51
19	Slide20.jpg	2912	
20	Slide20.jpg	1446	10.92
21	Slide20.jpg	1262	9.53
22	Slide20.jpg	4073	
23	Slide20.jpg	2350	17.31
24	Slide20.jpg	1049	7.73
25	Slide20.jpg	3470	
26	Slide20.jpg	2445	23.96
27	Slide20.jpg	1052	10.31
28	Slide20.jpg	3219	
29	Slide20.jpg	2120	22.39
30	Slide20.jpg	32	0.34

TABLE 22

TNS4 summary			
non-IS		55.29	
IS		57.16	
LV		112.45	
IS/LV		51%	

TABLE 23

TNF3 raw values			
1	Slide21.jpg	1551	
2	Slide21.jpg	3	0.02
3	Slide21.jpg	1502	10.65
4	Slide21.jpg	3054	

TABLE 23-continued

TNF3 raw values			
5	Slide21.jpg	922	6.34
6	Slide21.jpg	2049	14.09
7	Slide21.jpg	3374	
8	Slide21.jpg	1280	12.52
9	Slide21.jpg	1566	15.32
10	Slide21.jpg	2799	
11	Slide21.jpg	1476	14.77
12	Slide21.jpg	1061	10.61
13	Slide21.jpg	2330	
14	Slide21.jpg	398	3.25
15	Slide21.jpg	1012	8.25
16	Slide21.jpg	1689	
17	Slide21.jpg	7	0.05
18	Slide21.jpg	1544	10.06
19	Slide21.jpg	2894	
20	Slide21.jpg	361	2.62
21	Slide21.jpg	1925	13.97
22	Slide21.jpg	3254	
23	Slide21.jpg	1137	11.53
24	Slide21.jpg	1267	12.85
25	Slide21.jpg	2814	
26	Slide21.jpg	1272	12.66
27	Slide21.jpg	1113	11.07
28	Slide21.jpg	2821	
29	Slide21.jpg	1438	9.69
30	Slide21.jpg	174	1.17

TABLE 24

TNF3 summary			
non-IS		36.71	
IS		54.02	
LV		90.74	
IS/LV		60%	

TABLE 25

TNF4 raw values			
1	Slide22.jpg	1354	
2	Slide22.jpg	72	0.37
3	Slide22.jpg	1335	6.90
4	Slide22.jpg	2892	
5	Slide22.jpg	672	3.95
6	Slide22.jpg	2093	12.30
7	Slide22.jpg	3414	
8	Slide22.jpg	1342	9.83
9	Slide22.jpg	2213	16.21
10	Slide22.jpg	3698	
11	Slide22.jpg	1168	10.11
12	Slide22.jpg	2317	20.05
13	Slide22.jpg	2565	
14	Slide22.jpg	243	2.94
15	Slide22.jpg	1398	16.90
16	Slide22.jpg	1486	
17	Slide22.jpg	638	3.01
18	Slide22.jpg	583	2.75
19	Slide22.jpg	2719	
20	Slide22.jpg	26	0.16
21	Slide22.jpg	2164	13.53
22	Slide22.jpg	3514	
23	Slide22.jpg	568	4.04
24	Slide22.jpg	2361	16.80
25	Slide22.jpg	3908	

TABLE 25-continued

TNF4 raw values			
26	Slide22.jpg	1498	12.27
27	Slide22.jpg	1805	14.78
28	Slide22.jpg	2946	
29	Slide22.jpg	16	0.17
30	Slide22.jpg	1969	20.72

TABLE 26

TNF4 summary			
non-IS		23.42	
IS		70.46	
LV		93.88	
IS/LV		75%	

TABLE 27

TNS5 raw values			
1	Slide23.jpg	1615	
2	Slide23.jpg	8	0.04
3	Slide23.jpg	1571	8.75
4	Slide23.jpg	2789	
5	Slide23.jpg	1477	11.65
6	Slide23.jpg	1042	8.22
7	Slide23.jpg	3558	
8	Slide23.jpg	2026	22.21
9	Slide23.jpg	1327	14.55
10	Slide23.jpg	3822	
11	Slide23.jpg	1044	8.74
12	Slide23.jpg	1590	13.31
13	Slide23.jpg	3246	
14	Slide23.jpg	1224	8.67
15	Slide23.jpg	705	5.00
16	Slide23.jpg	1445	
17	Slide23.jpg	1228	7.65
18	Slide23.jpg	200	1.25
19	Slide23.jpg	2732	
20	Slide23.jpg	1951	15.71
21	Slide23.jpg	782	6.30
22	Slide23.jpg	3858	
23	Slide23.jpg	3039	30.72
24	Slide23.jpg	400	4.04
25	Slide23.jpg	3697	
26	Slide23.jpg	2609	22.58
27	Slide23.jpg	943	8.16
28	Slide23.jpg	3358	
29	Slide23.jpg	1492	10.22
30	Slide23.jpg	583	3.99

TABLE 28

TNS5 summary			
non-IS		69.10	
IS		36.78	
LV		105.88	
IS/LV		35%	

TABLE 29

TNS6 raw values			
1	Slide24.jpg	1216	
2	Slide24.jpg	258	1.49
3	Slide24.jpg	770	4.43

TABLE 29-continued

TNS6 raw values			
4	Slide24.jpg	3079	
5	Slide24.jpg	1436	10.26
6	Slide24.jpg	1417	10.12
7	Slide24.jpg	3677	
8	Slide24.jpg	2085	11.34
9	Slide24.jpg	1122	6.10
10	Slide24.jpg	3908	
11	Slide24.jpg	2151	15.96
12	Slide24.jpg	1415	10.50
13	Slide24.jpg	2371	
14	Slide24.jpg	1651	14.62
15	Slide24.jpg	495	4.38
16	Slide24.jpg	1123	
17	Slide24.jpg	879	5.48
18	Slide24.jpg	262	1.63
19	Slide24.jpg	3090	
20	Slide24.jpg	1775	12.64
21	Slide24.jpg	1121	7.98
22	Slide24.jpg	3470	
23	Slide24.jpg	2215	12.77
24	Slide24.jpg	1219	7.03
25	Slide24.jpg	3666	
26	Slide24.jpg	2524	19.97
27	Slide24.jpg	1411	11.16
28	Slide24.jpg	2470	
29	Slide24.jpg	1397	11.88
30	Slide24.jpg	140	1.19

TABLE 30

TNS6 summary			
non-IS		58.20	
IS		32.27	
LV		90.47	
IS/LV		36%	

TABLE 31

CN13 raw values			
1	Slide25.jpg	1010	
2	Slide25.jpg	4	0.04
3	Slide25.jpg	1006	8.96
4	Slide25.jpg	2216	
5	Slide25.jpg	756	5.80
6	Slide25.jpg	1708	13.10
7	Slide25.jpg	3122	
8	Slide25.jpg	744	5.72
9	Slide25.jpg	1674	12.87
10	Slide25.jpg	3214	
11	Slide25.jpg	177	1.87
12	Slide25.jpg	1678	17.75
13	Slide25.jpg	2504	
14	Slide25.jpg	371	3.41
15	Slide25.jpg	770	7.07
16	Slide25.jpg	940	
17	Slide25.jpg	3	0.03
18	Slide25.jpg	902	8.64
19	Slide25.jpg	1907	
20	Slide25.jpg	266	2.37
21	Slide25.jpg	1439	12.83
22	Slide25.jpg	2763	
23	Slide25.jpg	1036	9.00
24	Slide25.jpg	1855	16.11
25	Slide25.jpg	2930	

TABLE 31-continued

CN13 raw values			
26	Slide25.jpg	988	11.46
27	Slide25.jpg	1618	18.78
28	Slide25.jpg	2498	
29	Slide25.jpg	280	2.58
30	Slide25.jpg	1839	16.93

TABLE 32

CN13 summary			
non-IS		21.14	
IS		66.52	
LV		87.66	
IS/LV		76%	

TABLE 33

CN14 raw values			
1	Slide26.jpg	1387	
2	Slide26.jpg	40	0.23
3	Slide26.jpg	1356	7.82
4	Slide26.jpg	2994	
5	Slide26.jpg	699	4.67
6	Slide26.jpg	1620	10.82
7	Slide26.jpg	3017	
8	Slide26.jpg	1087	11.89
9	Slide26.jpg	1443	15.78
10	Slide26.jpg	2871	
11	Slide26.jpg	2644	29.47
12	Slide26.jpg	188	2.10
13	Slide26.jpg	2504	
14	Slide26.jpg	7	0.05
15	Slide26.jpg	1996	13.55
16	Slide26.jpg	1424	
17	Slide26.jpg	490	2.75
18	Slide26.jpg	931	5.23
19	Slide26.jpg	2926	
20	Slide26.jpg	40	0.27
21	Slide26.jpg	2231	15.25
22	Slide26.jpg	3248	
23	Slide26.jpg	782	7.95
24	Slide26.jpg	2137	21.71
25	Slide26.jpg	3401	
26	Slide26.jpg	348	3.27
27	Slide26.jpg	2624	24.69
28	Slide26.jpg	2079	
29	Slide26.jpg	573	4.69
30	Slide26.jpg	1042	8.52

TABLE 34

CN14 summary			
non-IS		32.62	
IS		62.74	
LV		95.36	
IS/LV		66%	

TABLE 35

TNF5 raw values			
1	Slide27.jpg	1504	
2	Slide27.jpg	22	0.13
3	Slide27.jpg	1336	7.99

TABLE 35-continued

TNF5 raw values			
4	Slide27.jpg	2786	
5	Slide27.jpg	390	3.22
6	Slide27.jpg	1956	16.15
7	Slide27.jpg	3792	
8	Slide27.jpg	1444	10.66
9	Slide27.jpg	2232	16.48
10	Slide27.jpg	3470	
11	Slide27.jpg	587	5.41
12	Slide27.jpg	2824	26.04
13	Slide27.jpg	3002	
14	Slide27.jpg	2361	16.52
15	Slide27.jpg	1329	9.30
16	Slide27.jpg	1666	
17	Slide27.jpg	274	1.48
18	Slide27.jpg	1024	5.53
19	Slide27.jpg	2735	
20	Slide27.jpg	9	0.08
21	Slide27.jpg	2897	24.36
22	Slide27.jpg	3575	
23	Slide27.jpg	1217	9.53
24	Slide27.jpg	2163	16.94
25	Slide27.jpg	3350	
26	Slide27.jpg	997	9.52
27	Slide27.jpg	1812	17.31
28	Slide27.jpg	3022	
29	Slide27.jpg	12	0.08
30	Slide27.jpg	1778	12.36

TABLE 36

TNF5 summary			
non-IS		28.32	
IS		76.23	
LV		104.55	
IS/LV		73%	

TABLE 37

TNF6 raw values			
1	Slide28.jpg	1114	
2	Slide28.jpg	62	0.45
3	Slide28.jpg	879	6.31
4	Slide28.jpg	2858	
5	Slide28.jpg	459	3.85
6	Slide28.jpg	1713	14.38
7	Slide28.jpg	3625	
8	Slide28.jpg	369	3.56
9	Slide28.jpg	2924	28.23
10	Slide28.jpg	3948	
11	Slide28.jpg	511	4.27
12	Slide28.jpg	2866	23.96
13	Slide28.jpg	3135	
14	Slide28.jpg	386	3.08
15	Slide28.jpg	1447	11.54
16	Slide28.jpg	1126	
17	Slide28.jpg	10	0.07
18	Slide28.jpg	1043	7.41
19	Slide28.jpg	3156	
20	Slide28.jpg	160	1.22
21	Slide28.jpg	3062	23.29
22	Slide28.jpg	3790	
23	Slide28.jpg	827	7.64
24	Slide28.jpg	2644	24.42
25	Slide28.jpg	3618	

TABLE 37-continued

TNF6 raw values			
26	Slide28.jpg	1607	14.66
27	Slide28.jpg	2452	22.36
28	Slide28.jpg	3440	
29	Slide28.jpg	1023	7.43
30	Slide28.jpg	1770	12.86

TABLE 38

TNF6 summary			
non-IS		23.11	
IS		87.38	
LV		110.50	
IS/LV		79%	

TABLE 39

TNS7 raw values			
1	Slide29.jpg	1713	
2	Slide29.jpg	607	4.61
3	Slide29.jpg	782	5.93
4	Slide29.jpg	2484	
5	Slide29.jpg	195	1.88
6	Slide29.jpg	1842	17.80
7	Slide29.jpg	2807	
8	Slide29.jpg	1568	12.29
9	Slide29.jpg	380	2.98
10	Slide29.jpg	3271	
11	Slide29.jpg	2187	20.06
12	Slide29.jpg	350	3.21
13	Slide29.jpg	2309	
14	Slide29.jpg	610	5.55
15	Slide29.jpg	1008	9.17
16	Slide29.jpg	1923	
17	Slide29.jpg	865	5.85
18	Slide29.jpg	631	4.27
19	Slide29.jpg	3033	
20	Slide29.jpg	1501	11.88
21	Slide29.jpg	780	6.17
22	Slide29.jpg	3287	
23	Slide29.jpg	2214	14.82
24	Slide29.jpg	456	3.05
25	Slide29.jpg	3395	
26	Slide29.jpg	2398	21.19
27	Slide29.jpg	287	2.54
28	Slide29.jpg	2969	
29	Slide29.jpg	1647	11.65
30	Slide29.jpg	67	0.47

TABLE 40

TNS7 summary			
non-IS		54.88	
IS		27.79	
LV		82.68	
IS/LV		34%	

TABLE 41

TNS8 raw values			
1	Slide30.jpg	1123	
2	Slide30.jpg	11	0.05
3	Slide30.jpg	988	4.40

TABLE 41-continued

TNS8 raw values			
4	Slide30.jpg	2352	
5	Slide30.jpg	279	2.25
6	Slide30.jpg	2001	16.16
7	Slide30.jpg	3274	
8	Slide30.jpg	1085	7.29
9	Slide30.jpg	1821	12.24
10	Slide30.jpg	3333	
11	Slide30.jpg	2048	17.20
12	Slide30.jpg	838	7.04
13	Slide30.jpg	2240	
14	Slide30.jpg	793	7.08
15	Slide30.jpg	840	7.50
16	Slide30.jpg	914	
17	Slide30.jpg	866	4.74
18	Slide30.jpg	64	0.35
19	Slide30.jpg	2811	
20	Slide30.jpg	397	2.68
21	Slide30.jpg	2135	14.43
22	Slide30.jpg	3378	
23	Slide30.jpg	588	3.83
24	Slide30.jpg	2250	14.65
25	Slide30.jpg	3241	
26	Slide30.jpg	2671	23.08
27	Slide30.jpg	287	2.48
28	Slide30.jpg	2697	
29	Slide30.jpg	1247	9.25
30	Slide30.jpg	23	0.17

TABLE 42

TNS8 summary			
non-IS		38.73	
IS		39.71	
LV		78.44	
IS/LV		51%	

TABLE 43

TNF7 raw values			
1	Slide31.jpg	1733	
2	Slide31.jpg	15	0.06
3	Slide31.jpg	1704	6.88
4	Slide31.jpg	3401	
5	Slide31.jpg	719	3.38
6	Slide31.jpg	2216	10.43
7	Slide31.jpg	3789	
8	Slide31.jpg	917	7.02
9	Slide31.jpg	2163	16.56
10	Slide31.jpg	4149	
11	Slide31.jpg	719	5.03
12	Slide31.jpg	3423	23.93
13	Slide31.jpg	3309	
14	Slide31.jpg	1479	8.49
15	Slide31.jpg	1771	10.17
16	Slide31.jpg	1777	
17	Slide31.jpg	1049	4.13
18	Slide31.jpg	678	2.67
19	Slide31.jpg	3117	
20	Slide31.jpg	221	1.13
21	Slide31.jpg	2281	11.71
22	Slide31.jpg	3970	
23	Slide31.jpg	2416	17.65
24	Slide31.jpg	796	5.81
25	Slide31.jpg	4354	

TABLE 43-continued

TNF7 raw values			
26	Slide31.jpg	3291	21.92
27	Slide31.jpg	697	4.64
28	Slide31.jpg	3316	
29	Slide31.jpg	2414	13.83
30	Slide31.jpg	62	0.36

TABLE 44

TNF7 summary	
non-IS	41.32
IS	46.57
LV	87.90
IS/LV	53%

TABLE 45

TNF8 raw values			
1	Slide32.jpg	1553	
2	Slide32.jpg	572	2.58
3	Slide32.jpg	873	3.93
4	Slide32.jpg	3334	
5	Slide32.jpg	1084	5.53
6	Slide32.jpg	1525	7.78
7	Slide32.jpg	4166	
8	Slide32.jpg	2437	12.87
9	Slide32.jpg	1557	8.22
10	Slide32.jpg	4558	
11	Slide32.jpg	2698	20.13
12	Slide32.jpg	1306	9.74
13	Slide32.jpg	3405	
14	Slide32.jpg	2991	25.47
15	Slide32.jpg	51	0.43
16	Slide32.jpg	1543	
17	Slide32.jpg	3	0.01
18	Slide32.jpg	1407	6.38
19	Slide32.jpg	3359	
20	Slide32.jpg	581	2.94
21	Slide32.jpg	2011	10.18
22	Slide32.jpg	3986	
23	Slide32.jpg	202	1.11
24	Slide32.jpg	3788	20.91
25	Slide32.jpg	4684	
26	Slide32.jpg	425	3.08
27	Slide32.jpg	3308	24.01
28	Slide32.jpg	3498	
29	Slide32.jpg	920	7.63
30	Slide32.jpg	1731	14.35

TABLE 46

TNF8 summary	
non-IS	40.68
IS	52.97
LV	93.65
IS/LV	57%

TABLE 47

TNS9 raw values			
1	Slide33.jpg	2637	
2	Slide33.jpg	14	0.06
3	Slide33.jpg	2081	9.47

TABLE 47-continued

TNS9 raw values			
4	Slide33.jpg	4101	
5	Slide33.jpg	1571	7.28
6	Slide33.jpg	1516	7.02
7	Slide33.jpg	4527	
8	Slide33.jpg	2519	18.36
9	Slide33.jpg	1555	11.34
10	Slide33.jpg	3326	
11	Slide33.jpg	3188	19.17
12	Slide33.jpg	27	0.16
13	Slide33.jpg	2336	
14	Slide33.jpg	1885	9.68
15	Slide33.jpg	240	1.23
16	Slide33.jpg	2343	
17	Slide33.jpg	2027	10.38
18	Slide33.jpg	21	0.11
19	Slide33.jpg	3393	
20	Slide33.jpg	1928	10.80
21	Slide33.jpg	945	5.29
22	Slide33.jpg	4425	
23	Slide33.jpg	2984	22.25
24	Slide33.jpg	637	4.75
25	Slide33.jpg	3063	
26	Slide33.jpg	773	5.05
27	Slide33.jpg	1885	12.31
28	Slide33.jpg	2324	
29	Slide33.jpg	1390	7.18
30	Slide33.jpg	9	0.05

TABLE 48

TNS9 summary	
non-IS	55.11
IS	25.86
LV	80.97
IS/LV	32%

TABLE 49

TNS10 raw values			
1	Slide34.jpg	1775	
2	Slide34.jpg	1082	4.88
3	Slide34.jpg	348	1.57
4	Slide34.jpg	3607	
5	Slide34.jpg	1823	11.12
6	Slide34.jpg	1483	9.05
7	Slide34.jpg	4313	
8	Slide34.jpg	1087	6.80
9	Slide34.jpg	2173	13.60
10	Slide34.jpg	4275	
11	Slide34.jpg	2471	15.03
12	Slide34.jpg	1734	10.55
13	Slide34.jpg	2864	
14	Slide34.jpg	2424	18.62
15	Slide34.jpg	43	0.33
16	Slide34.jpg	1601	
17	Slide34.jpg	1600	8.00
18	Slide34.jpg	16	0.08
19	Slide34.jpg	3486	
20	Slide34.jpg	933	5.89
21	Slide34.jpg	935	5.90
22	Slide34.jpg	4312	
23	Slide34.jpg	3250	20.35
24	Slide34.jpg	722	4.52
25	Slide34.jpg	4178	

TABLE 49-continued

TNS10 raw values			
26	Slide34.jpg	3996	24.87
27	Slide34.jpg	231	1.44
28	Slide34.jpg	3046	
29	Slide34.jpg	2854	20.61
30	Slide34.jpg	39	0.28

TABLE 50

TNS10 summary	
non-IS	68.08
IS	23.66
LV	91.74
IS/LV	26%

TABLE 51

TNF9 raw values			
1	Slide35.jpg	1737	
2	Slide35.jpg	841	2.91
3	Slide35.jpg	788	2.72
4	Slide35.jpg	3368	
5	Slide35.jpg	1416	7.99
6	Slide35.jpg	1230	6.94
7	Slide35.jpg	4474	
8	Slide35.jpg	1046	8.18
9	Slide35.jpg	3356	26.25
10	Slide35.jpg	4877	
11	Slide35.jpg	1303	6.68
12	Slide35.jpg	3142	16.11
13	Slide35.jpg	3803	
14	Slide35.jpg	2906	16.81
15	Slide35.jpg	15	0.09
16	Slide35.jpg	1719	
17	Slide35.jpg	8	0.03
18	Slide35.jpg	1545	5.39
19	Slide35.jpg	3500	
20	Slide35.jpg	9	0.05
21	Slide35.jpg	3382	18.36
22	Slide35.jpg	4790	
23	Slide35.jpg	9	0.07
24	Slide35.jpg	4476	32.71
25	Slide35.jpg	4213	
26	Slide35.jpg	1798	10.67
27	Slide35.jpg	2840	16.85
28	Slide35.jpg	3714	
29	Slide35.jpg	2917	17.28
30	Slide35.jpg	342	2.03

TABLE 52

TNF9 summary	
non-IS	35.33
IS	63.72

TABLE 52-continued

TNF9 summary	
LV	99.05
IS/LV	64%

TABLE 53

TNF10 raw values			
1	Slide36.jpg	2294	
2	Slide36.jpg	14	0.08
3	Slide36.jpg	2183	12.37
4	Slide36.jpg	4093	
5	Slide36.jpg	189	1.34
6	Slide36.jpg	3572	25.31
7	Slide36.jpg	4330	
8	Slide36.jpg	829	9.38
9	Slide36.jpg	2710	30.67
10	Slide36.jpg	2189	
11	Slide36.jpg	185	1.18
12	Slide36.jpg	1581	10.11
13	Slide36.jpg	1961	
14	Slide36.jpg	344	1.40
15	Slide36.jpg	1293	5.27
16	Slide36.jpg	2188	
17	Slide36.jpg	1766	10.49
18	Slide36.jpg	382	2.27
19	Slide36.jpg	4243	
20	Slide36.jpg	2206	15.08
21	Slide36.jpg	1246	8.52
22	Slide36.jpg	4883	
23	Slide36.jpg	3763	37.76
24	Slide36.jpg	583	5.85
25	Slide36.jpg	2162	
26	Slide36.jpg	2025	13.11
27	Slide36.jpg	18	0.12
28	Slide36.jpg	2558	
29	Slide36.jpg	1179	3.69
30	Slide36.jpg	615	1.92

TABLE 54

TNF10 summary	
non-IS	46.76
IS	51.20
LV	97.96
IS/LV	52%

TABLE 55

Composite image data						
	IS/LV		IS/LV		IS/LV	
CON7	70%	TMZ3	64%	TNF1	57%	TNS1
CON5	65%	TMZ1	68%	TNF2	67%	TNS2
CON6	75%	TMZ2	60%	TNF3	60%	TNS3
CON4	65%	TMZ7	43%	TNF4	75%	TNS4
CON3	64%	TMZ8	51%	TNF5	73%	TNS5
CON1	77%	TMZ5	58%	TNF6	79%	TNS6
CON2	55%	TMZ6	49%	TNF7	53%	TNS7
CON8	68%	TMZ9	44%	TNF8	57%	TNS8
CON9	67%	TMZ10	49%	TNF9	64%	TNS9
CON10	62%	TMZ4	71%	TNF10	52%	TNS10
CON11	73%					
CON12	75%					
CON13	76%					
CON14	66%					
Mean	68%	Mean	56%	Mean	64%	Mean
SD	6%	SD	10%	SD	10%	SD
SE	2%	SE	3%	SE	3%	SE
TTEST			8.77E-04		1.61E-01	
					4.79E-06	TMZ/TNS
						4.00E-02

[0222] The results show that a combination of trimetazidine, nicotinamide, and succinate at 20 μ M preserved coronary flow and cardiac functional recovery and decreased infarct size in isolated hearts after ischemia-reperfusion. This combination was more effective in decreasing infarct size than TMZ alone. A combination of trimetazidine, nicotinamide, and succinate at 2 μ M did not appear to decrease myocardial ischemia-reperfusion injury.

[0223] This study suggested that the combination of trimetazidine, nicotinamide, and succinate at 20 μ M generated better protection against ischemia-reperfusion injury in Langendorff system.

[0224] FIG. 37 is a schematic of the method used to analyze the effects of compositions of the invention on cardiac function. Following transverse aortic constriction (TAC) or a sham procedure, mice were given one of the following via an osmotic mini-pump: CV8814 at 5.85 mg/kg/day (CV4); CV8814 at 5.85 mg/kg/day, nicotinic acid at 1.85 mg/kg/day, and succinate at 2.43 mg/kg/day (TV8); or saline (SA). Echocardiograms were measured immediately following TAC, three weeks after TAC, and 6 weeks after TAC. Mice were sacrificed at 6 weeks, and tissues were analyzed.

[0225] FIG. 38 shows hearts from mice six weeks after a sham procedure (SHAM), TAC followed by saline administration (TAC), TAC followed by CV4 administration (CV4), or TAC followed by TV8 administration.

[0226] FIG. 39 is of graph of heart weight relative to body weight six weeks after transverse aortic constriction. Treatments are as indicated in relation to FIG. 38.

[0227] FIG. 40 is graph of heart weight six weeks after transverse aortic constriction. Treatments are as indicated in relation to FIG. 38.

[0228] FIG. 41 shows graphs of fractional shortening (FS) and ejection fraction (EF) at indicated time points after transverse aortic constriction. Treatments are as indicated in relation to FIG. 38.

[0229] FIG. 42 is a graph of left ventricular end-systolic diameter at indicated time points after transverse aortic constriction. Treatments are as indicated in relation to FIG. 38.

[0230] FIG. 43 is a graph of intraventricular septal dimension at indicated time points after transverse aortic constriction. Treatments are as indicated in relation to FIG. 38.

[0231] FIG. 44 is a graph of left ventricular mass at indicated time points after transverse aortic constriction. Treatments are as indicated in relation to FIG. 38.

[0232] FIG. 45 is a graph of isovolumic relaxation time at indicated time points after transverse aortic constriction. Treatments are as indicated in relation to FIG. 38.


[0233] FIG. 46 is a graph of the ratio peak velocity flow in early diastole vs. late diastole at indicated time points after transverse aortic constriction. Treatments are as indicated in relation to FIG. 38.


[0234] FIG. 47 is a graph of left ventricular developed pressure at six weeks after transverse aortic constriction. Treatments are as indicated in relation to FIG. 38.

[0235] FIG. 48 is a graph of the rate of left ventricle pressure rise at six weeks after transverse aortic constriction. Treatments are as indicated in relation to FIG. 38.

Chemical Synthesis Schemes.

[0236] Compounds of the invention include 2-(4-(2,3,4-trimethoxybenzyl)piperazin-1-yl)ethan-1-ol (referred to herein as CV8814) and 2-(4-(2,3,4-trimethoxybenzyl)piperazin-1-yl)ethyl nicotinate (referred to herein as CV-8972). These compounds may be synthesized according to the following scheme:





[0238] In batch MBA 25, 5% water w/r/t total volume of solvent (23 volumes) containing 30% methanol: 70% MEK was used for precipitation. The yield was 67% of monohydrate of CV-8972. Water content was determined by KF to be 3.46%.

[0239] In batch MBA 26, 1.33% water w/r/t total volume of solvent (30 volumes) containing 20% methanol: 80% MEK was used for precipitation. The yield was 86.5% of monohydrate of CV-8972. Water content was determined by

[0243] In batch MBA 30, 3% water was added. Material was isolated and dried under vacuum at 40° C. for 24 hours. Water content was determined by KF to be 3.51%, showing that monohydrate is forming with addition of excess water.

[0244] In batch MBA 31, 5% water was added. Material was isolated and dried under vacuum at 40° C. for 24 hours. Water content was determined by KF to be 3.30%, showing that monohydrate is forming with addition of excess water.

[0245] Results are summarized in Table 56.

TABLE 56

Sample	Water percentage theoretical (for monohydrate preparation)	KF result (% of water)	KF result (Sample after drying at 40° C. for 24 hours)	Amount of Water used for reaction (based on total volume)	Ratio of MeOH:MEK	Total Volume	Yield obtained (%)	Drying Time (hr)	Drying temperature (° C.)
289-MBA-25	3.32%	3.46	—	5%	30-70	23 vol	67.6	24	22
289-MBA-26	3.32%	4.00	3.75	1.33%	20-80	30 vol	86.5	19	23
289-MBA-27	3.32%	3.93	3.54	3%	22-78	32 vol	87.22	18	23
289-MBA-29	3.32%	—	0.89	1.0 eq based on input weight	20-80	30 vol	84	24	40
289-MBA-30	3.32%	—	3.51	3%	20-80	30 vol	90	24	40
289-MBA-31	3.32%	—	3.30	5%	20-80	30 vol	81	24	40

KF to be 4.0%. The product was dried under vacuum at 40° C. for 24 hours to decrease water content to 3.75%.

[0240] In batch MBA 27, 3% water w/r/t total volume of solvent (32 volumes) containing 22% methanol: 78% MEK was used for precipitation. The yield was 87.22% of monohydrate of CV-8972. Water content was determined by KF to be 3.93% after 18 hours of drying at room temperature under vacuum. The product was further dried under vacuum at 40° C. for 24 hours to decrease water content to 3.54%.

[0241] In other batches, the ratio and total volume of solvent were held constant at 20% methanol: 80% MEK and 30 volumes in batches using 2.5 g of product, and only the percentage of water was varied.

[0242] In batch MBA 29, 1.0 equivalent of water was added. Material was isolated and dried under vacuum at 40° C. for 24 hours. Water content was determined by KF to be 0.89%, showing that the monohydrate form was not forming stoichiometrically.

Metabolism of Compounds in Dogs

[0246] The metabolism of various compounds was analyzed in dogs.

[0247] FIG. 49 is a graph showing levels of CV-8814 (solid triangles, solid lines) and trimetazidine (open triangles, dashed lines) after intravenous administration of CV-8834 at 2.34 mg/kg. CV-8834 is a compound of formula (II) in which y=1.

[0248] FIG. 50 is a graph showing levels of CV-8814 (solid triangles, solid lines) and trimetazidine (open triangles, dashed lines) after oral administration of CV-8834 at 77.4 mg/kg.

[0249] FIG. 51 is a graph showing levels of CV-8814 (solid triangles, solid lines) and trimetazidine (open triangles, dashed lines) after oral administration of CV-8834 at 0.54 mg/kg.

[0250] FIG. 52 is a graph showing levels of CV-8814 (solid triangles, solid lines) and trimetazidine (open triangles, dashed lines) after oral administration of CV-8834 at 1.08 mg/kg.

[0251] FIG. 53 is a graph showing levels of CV-8814 (solid triangles, solid lines) and trimetazidine (open triangles, dashed lines) after oral administration of CV-8834 at 2.15 mg/kg.

[0252] Data from FIGS. 48-53 is summarized in Table 57.

TABLE 57

Compound	Route of admin.	Dose (mg/kg)	Analyte	T _{max} (hours)	C _{max} (ng/mL)	AUC ₀₋₈ (ng × hr/mL)	% F
CV-8834	PO	77.4	8814	0.75	12100	38050	69
CV-8834	PO	77.4	TMZ	1.67	288	1600	72
CV-8834	IV	2.34	8814	0.083	974	1668	—
CV-8834	IV	2.34	TMZ	2.67	13.4	66.7	—
CV-8834	PO	0.54	8814	0.5	74.0	175	45
CV-8834	PO	0.54	TMZ	1.17	3.63	17.6	>100
CV-8834	PO	1.08	8814	0.5	136	335	44
CV-8834	PO	1.08	TMZ	0.866	6.19	30.4	99
CV-8834	PO	2.15	8814	0.583	199	536	35
CV-8834	PO	2.15	TMZ	1.17	9.80	51.6	84

[0253] FIG. 54 is a graph showing levels of trimetazidine after oral administration of CV-8972 at 1.5 mg/kg (triangles) or intravenous administration of trimetazidine at 2 mg/kg (squares).

[0254] FIG. 55 is a graph showing levels of CV-8814 after oral administration of CV-8972 at 1.5 mg/kg (triangles) or intravenous administration of CV-8814 at 2.34 mg/kg (squares).

[0255] FIG. 56 is a graph showing levels of CV-8814 after intravenous administration of CV-8834 at 4.3 mg/kg (squares) or oral administration of CV-8834 at 2.15 mg/kg (triangles). FIG. 57 is a graph showing levels of CV-8814 after intravenous administration of CV-8814 at 2.34 mg/kg (squares) or oral administration of CV-8814 at 2.34 mg/kg (triangles).

[0256] Data from FIGS. 54-57 is summarized in Table 58.

TABLE 58

Compound	Route of admin.	Dose (mg/kg)	Vehicle	Fasted	Analyte	T _{max} (hours)	C _{max} (ng/mL)	AUC ₀₋₂₄ (ng × hr/mL)	% F
CV-8972	PO	1.5	—	—	TMZ	2.0	17.0	117	4.3%
TMZ	IV	2	0.9% NaCl	8 hrs	TMZ	0.083	1002	3612	—
CV-8972	PO	1.5	—	—	8814	1.125	108	534	27%
CV-8814	IV	2.34	0.9% NaCl	8 hrs	8814	0.083	1200	3059	—
CV-8834	PO	4.3	0.9% NaCl	8 hrs	8814	1.0	692	2871	69%
CV-8834	IV	4.3	0.9% NaCl	8 hrs	8814	0.083	1333	4154	—
CV-8834	PO	4.3	0.9% NaCl	8 hrs	8814	1.0	692	2871	51%
CV-8814	IV	2.34	0.9% NaCl	8 hrs	8814	0.083	1200	3059	—
CV-8814	PO	2.34	0.9% NaCl	8 hrs	8814	0.333	672	1919	63%
CV-8814	IV	2.34	0.9% NaCl	8 hrs	8814	0.083	1200	3059	—

Effect of CV-8814 on Enzyme Activity

[0257] The effect of CV-8814 on the activity of various enzymes was analyzed in in vitro assays. Enzyme activity was assayed in the presence of 10 μ M CV-8814 using conditions of time, temperature, substrate, and buffer that were optimized for each enzyme based on published literature. Inhibition of 50% or greater was not observed for any of the following enzymes: ATPase, Na⁺/K⁺, pig heart; Cholinesterase, Acetyl, ACES, human; Cyclooxygenase COX-1, human; Cyclooxygenase COX-2, human; Monoamine Oxidase MAO-A, human; Monoamine Oxidase MAO-B, human; Peptidase, Angiotensin Converting Enzyme, rabbit; Peptidase, CTSG (Cathepsin G), human; Phosphodiesterase

PDE3, human; Phosphodiesterase PDE4, human; Protein Serine/Threonine Kinase, PKC, Non-selective, rat; Protein Tyrosine Kinase, Insulin Receptor, human; Protein Tyrosine Kinase, LCK, human; Adenosine A1, human; Adenosine A_{2A}, human; Adrenergic α_{1A} , rat; Adrenergic α_{1B} , rat;

Adrenergic α_{1D} , human; Adrenergic α_{2A} , human; Adrenergic α_{2B} , human; Adrenergic β_1 , human; Adrenergic β_2 , human; Androgen (Testosterone), human; Angiotensin AT₁, human; Bradykinin B₂, human; Calcium Channel L-Type, Benzothiazepine, rat; Calcium Channel L-Type, Dihydro-pyridine, rat; Calcium Channel L-Type, Phenylalkylamine, rat; Calcium Channel N-Type, rat; Cannabinoid CB₁, human; Cannabinoid CB₂, human; Chemokine CCR1, human; Chemokine CXCR2 (IL-8R_B), human; Cholecystokinin CCK₁ (CCK_A), human; Cholecystokinin CCK₂ (CCK_B), human; Dopamine D₁, human; Dopamine D_{2L}, human; Dopamine D_{2S}, human; Endothelin ET_A, human; Estrogen ER α , human; GABA_A, Chloride Channel, TBOB, rat; GABA_A, Flunitrazepam, Central, rat; GABA_A, Ro-15-1788, Hippocampus, rat; GABA_{B1A}, human; Glucocorticoid, human; Glutamate, AMPA, rat; Glutamate, Kainate,

rat; Glutamate, Metabotropic, mGlu5, human; Glutamate, NMDA, Agonism, rat; Glutamate, NMDA, Glycine, rat; Glutamate, NMDA, Phencyclidine, rat; Glutamate, NMDA, Polyamine, rat; Glycine, Strychnine-Sensitive, rat; Histamine H₁, human; Histamine H₂, human; Melanocortin MC₁, human; Melanocortin MC₄, human; Muscarinic M₁, human; Muscarinic M₂, human; Muscarinic M₃, human; Muscarinic M₄, human; Neuropeptide Y Y₁, human; Nicotinic Acetylcholine, human; Nicotinic Acetylcholine al, Bungarotoxin, human; Opiate δ_1 (OP1, DOP), human; Opiate κ (OP2, KOP), human; Opiate μ (OP3, MOP), human; Platelet Activating Factor (PAF), human; Potassium Channel [KATP], hamster; Potassium Channel hERG, human;

PPAR γ , human; Progesterone PR-B, human; Serotonin (5-Hydroxytryptamine) 5-HT_{1A}, human; Serotonin (5-Hydroxytryptamine) 5-HT_{1B}, human; Serotonin (5-Hydroxytryptamine) 5-HT_{2A}, human; Serotonin (5-Hydroxytryptamine) 5-HT_{2B}, human; Serotonin (5-Hydroxytryptamine) 5-HT_{2C}, human; Serotonin (5-Hydroxytryptamine) 5-HT₃, human; Sodium Channel, Site 2, rat; Tachykinin NK₁, human; Transporter, Adenosine, guinea pig; Transporter, Dopamine (DAT), human; Transporter, GABA, rat; Transporter, Norepinephrine (NET), human; Transporter, Serotonin (5-Hydroxytryptamine) (SERT), human; and Vasopressin V_{1A}, human.

Analysis of CV-8972 Batch Properties

[0258] CV-8972 (2-(4-(2,3,4-trimethoxybenzyl) piperazin-1-yl)ethyl nicotinate, HCl salt, monohydrate) was prepared and analyzed. The batch was determined to be 99.62% pure by HPLC.

[0259] FIG. 58 is a graph showing the HPLC elution profile of a batch of CV-8972.

[0260] FIG. 59 is a graph showing analysis of molecular species present in a batch of CV-8972. FIG. 60 is a pair of graphs showing HPLC elution profiles of molecular species present in a batch of CV-8972.

[0261] FIG. 61 is a pair of graphs showing HPLC elution profiles of molecular species present in a batch of CV-8972.

[0262] FIG. 62 is a graph showing X-ray powder diffraction analysis of a batch of CV-8972.

[0263] FIG. 63 is a graph showing X-ray powder diffraction analysis of batches of CV-8972. Batch 289-MBA-15-A, shown in blue, contains form B of CV-8972, batch 276-MBA-172, shown in black contains form A of CV-8972, and batch 289-MBA-16, shown in red, contains a mixture of forms A and B.

[0264] FIG. 64 is a graph showing differential scanning calorimetry and thermal gravimetric analysis of batch 276-MBA-172 of CV-8972.

[0265] FIG. 65 is a graph showing dynamic vapor sorption (DVS) of batch 276-MBA-172 of CV-8972.

[0266] FIG. 66 is a graph showing differential scanning calorimetry and thermal gravimetric analysis of batch 289-MBA-15-A of CV-8972.

[0267] FIG. 67 is a graph showing dynamic vapor sorption (DVS) of batch 289-MBA-15-A of CV-8972.

[0268] FIG. 68 is a graph showing X-ray powder diffraction analysis of samples of CV-8972. A pre-DVS sample from batch 276-MBA-172 is shown in blue, a pre-DVS sample from batch 289-MBA-15-A is shown in red, and a post-DVS sample from batch 289-MBA-15-A is shown in black.

[0269] FIG. 69 is a graph showing differential scanning calorimetry and thermal gravimetric analysis of batch 289-MBA-16 of CV-8972.

[0270] FIG. 70 is a graph showing X-ray powder diffraction analysis of samples of CV-8972. Form B is shown in green, form A is shown in blue, a sample from an ethanol slurry of batch 289-MBA-15-A is shown in red, and a sample from an ethanol slurry of batch 289-MBA-16 is shown in black.

[0271] The stability of CV-8972 was analyzed.

[0272] Samples from batch 289-MBA-15-A (containing form B) were added to various solvents, incubated under various conditions, and analyzed by X-ray powder diffraction. Results are summarized in Table 59.

TABLE 59

Solvent	Conditions	XRPD results
EtOH	Slurry, RT, 3 d	Form A + Form B
MeOH/H ₂ O (95:5) $A_w = 0.16$	Slurry, RT, 5 d	Form A
IPA/H ₂ O (98:2) $A_w = 0.26$	Slurry, RT, 5 d	Form A
MeOH/H ₂ O (80:20) $A_w = 0.48$	Slurry, RT, 5 d	Form A
EtOH/H ₂ O (90:10) $A_w = 0.52$	Slurry, RT, 5 d	Form A
IPA/H ₂ O (90:10) $A_w = 0.67$	Slurry, RT, 5 d	Form A
Acetone/H ₂ O (90:10) $A_w = 0.72$	Slurry, RT, 5 d	Form A
ACN/H ₂ O (90:10) $A_w = 0.83$	Slurry, RT, 5 d	Form A
EtOAc/H ₂ O (97:3) $A_w = 0.94$	Slurry, RT, 5 d	Form A
MeOH	Slurry, RT, 5 d	Form A + Form B
EtOAc	Slurry, RT, 5 d	Form A + Form B
MEK	Slurry, RT, 5 d	Form A
—	100° C., 20 minutes	Form B, shifted with minor Form A
EtOH	CC from 60° C.	Form C + minor Form A

[0273] Samples from batch 289-MBA-16 (containing forms A and B) were added to various solvents, incubated under various conditions, and analyzed by X-ray powder diffraction. Results are summarized in Table 60.

TABLE 60

Solvent	Conditions	XRPD results
EtOH	Slurry, RT, 3 d	Form A + Form B
MeOH	Vapor diffusion w/ MTBE	Form A
EtOAc	Attempted to dissolve at ~60° C., solids remained, cooled slowly to RT, let stir at RT from 60° C.	Form A + Form B

[0274] FIG. 71 is a graph showing X-ray powder diffraction analysis of samples of CV-8972. A sample containing form B is shown in blue, a sample containing form A is shown in red, and a sample containing a mixture of forms A and C is shown in black.

[0275] The stability of CV-8972 was analyzed. Aqueous samples containing CV-8972 at different concentrations and pH were incubated for various periods and analyzed. Results are shown in Table 61.

TABLE 61

Sample	(hrs)	pH	Retention Time					Decrease in purity of CV-8972 between time points
			2.2	2.6	4.2	4.7	5.6	
276-MBA-172	0	6.6	3.39	0.6	0.23	0.54	95.24	
10 mg/mL pH 6	1	6.8	4.81	0.81	0.23	0.73	93.43	1.81
(Form A)	4	6.8	5.72	0.9	0.21	0.83	91.82	1.61
	6	6.7	6.45	0.81	ND	0.93	91.8	0.02
	22	6.7	7.38	1.54	0.13	1.11	89.66	2.14
276-MBA-172	0	6.1	ND	ND	1.29	ND	98.01	
2 mg/mL pH 6	1	6.1	1.5	ND	1.28	ND	97.22	0.79
(Form A)	4	6.1	2.03	ND	0.95	ND	97.01	0.21
	6	6.1	2.47	ND	1.02	ND	96.51	0.5
	22	6.1						
289-MBA-15-A 10	0	6	3.3	0.6	0.26	0.48	95.36	
mg/mL pH 6 (Form	1	6.1	3.76	0.65	0.25	0.53	94.81	0.55
B)	4	6	3.97	0.59	0.19	0.56	94.69	0.12
	6	5.9	4.3	0.54	0.17	0.6	94.39	0.3
	22	5.9	4.53	0.69	0.19	0.65	93.93	0.46
289-MBA-15-A 2	0	6.9	1.33	ND	1.19	ND	97.48	
mg/mL pH 6 (Form	1	6.9	3.73	ND	1.17	ND	95.1	2.38
B)	4	6.8	5.25	0.67	0.84	0.79	92.45	2.65
	6	6.8	6.63	0.9	0.83	0.99	90.65	1.8
	22	6.7	7.72	1.13	0.86	1.14	89.15	1.5
276-MBA-172 10	0	7.1	5.9	0.94	0.22	0.78	92.85	
mg/mL pH 7 (Form	1	7.2	8.12	1.45	0.21	1.17	89.05	3.8
A)	4	7.1	10.14	1.48	0.13	1.46	86.8	2.25
	6	7.1	11.63	1.78	0.13	1.67	84.79	2.01
	22	7						
276-MBA-172 2	0	6.7	1.42	ND	1.05	ND	97.53	
mg/mL pH 7 (Form	1	6.8	3.31	ND	1.06	0.57	95.06	2.47
A)	4	6.7	4.21	0.58	0.82	0.69	93.7	1.36
	6	6.7	5.63	0.67	0.74	0.85	92.12	1.58
	22	6.8	6.26	0.85	0.85	0.98	91.07	1.05
289-MBA-15-A 10	0	7.4	6.2	1.16	0.27	0.87	91.5	
mg/mL pH 7 (Form	1	7.4	10.47	1.65	0.25	1.44	86.18	5.32
B)	4	7.4	13.64	1.93	0.19	1.89	82.36	3.82
	6	7.3	15.66	2.57	0.2	0.2	79.37	2.99
	22	7.1						
289-MBA-15-A 2	0	6.5	1.62	ND	0.9	ND	97.48	
mg/mL pH 7 (Form	1	6.6	3.16	ND	0.89	0.49	95.46	2.02
B)	4	6.5	4.27	0.53	0.66	0.62	93.92	1.54
	6	6.5						
	22	6.5						

[0276] Samples from batch S-18-0030513 (containing form A) were added to various solvents, incubated under various conditions, and analyzed by X-ray powder diffraction. Results are summarized in Table 62.

TABLE 62

Solvent	Conditions	XRPD results
CHC13	Slurry, RT	Form A
EtOAc	Slurry, RT	Form A
THF	Slurry, RT	Form A
—	VO, RT	Form A
—	80° C., 20 minutes	Form A with slight peak shifting
—	100° C., 20 minutes	Form B + Form A, shifted
—	97% RH Stress of	Form A
	Form A dried at	
	80° C. for 20 min	
EtOH	Crash cool from	Form A + Form C
	70° C.	
MEK/H2O 99:1	Slow cool from	Form A
	70° C.	

[0277] Samples from batch 289-MBA-16 (containing forms A and B) were added to various solvents, incubated under various conditions, and analyzed by X-ray powder diffraction. Results are summarized in Table 63.

TABLE 63

Solvent	Conditions	XRPD results
EtOH	Slurry, RT, 3 d	Form A + Form B
MeOH	VD w/ MTBE	Form A
EtOAc	SC from 60° C.	Form A + Form B
THF	SC from 60° C.	Form B
EtOH	SC from 60° C.	Form A + Form C
MeOH/H2O (95:5)	Slurry, overnight, 1 g scale	Form A

[0278] FIG. 72 is a graph showing differential scanning calorimetry and thermal gravimetric analysis of samples containing form A of CV-8972. A sample from an ethanol acetate-water slurry is shown with solid lines, a sample from a methanol-water slurry is shown with regularly-dashed lines, and a sample from an ethanol-water slurry is shown with dashed-dotted lines.

[0279] FIG. 73 is a graph showing differential scanning calorimetry and thermal gravimetric analysis of a sample containing form A of CV-8972. Prior to analysis, the sample was dried at 100° C. for 20 minutes.

[0280] Samples containing form A of CV-8972 were analyzed for stability in response to humidity. Samples were incubated at 40° C., 75% relative humidity for various periods and analyzed. Results are shown in Table 64.

TABLE 64

Time (days)	Retention Time			
	1.9	3.9	4.5	5.4
0	ND	1.16	ND	98.84
1	ND	0.68	ND	99.32
7	0.63	0.14	0.12	99.12

[0281] Form A of CV-8972 were analyzed for stability in aqueous solution. Aqueous samples containing CV-8972 at different concentrations and pH were incubated for various periods and analyzed. Results are shown in Table 65.

TABLE 65

Concentration of CV-8972	Time (hrs)	Retention Time					% change from t0 of RT 5.4
		1.9	2.2	3.9	4.5	5.4	
21 mg/mL, Initial pH = 2.0	0	ND	ND	1.12	ND	98.88	—
	1	1.03	ND	0.94	ND	98.03	-0.86
	2	1.9	ND	1	ND	97.11	-1.79
	6	5.25	0.83	0.96	0.78	92.18	-6.78
12.5 mg/mL, Initial pH = 2.1	0	ND	ND	1.79	ND	98.21	—
	1	1.38	ND	1.41	ND	97.21	-1.02
	2	2.43	ND	1.67	ND	95.9	-2.35
	6	6.59	1.04	1.74	1.04	89.58	-8.79
4.2 mg/mL, Initial pH = 2.3	0	ND	ND	5.35	ND	94.65	—
	1	ND	ND	4.02	ND	95.98	1.41
	2	3.72	ND	5.09	ND	91.19	-3.66
	6	9.71	ND	5.3	ND	84.99	-10.21

[0282] The amount of CV-8972 present in various dosing compositions was analyzed. Results are shown in Table 66.

TABLE 66

Target Dose (mg/mL)	Vol. API soln. (mL)	Mass CV8972 (mg)	Initial		Total vol.		pH after base soln. addn.	Vol. addl. 1N NaOH added (mL)	Final Dose (mg/mL)
			pH API soln.	Vol. 1N NaOH (mL)	base soln. (mL)	pH base soln. addn.			
10	30	779.06	2.0	2.07	30	3.6	0.7	9.92	
2	30	157.38	2.4	0.19	30	2.8	0.35	2.02	
10	50	777.05	2.1	2.77	10	6.2	—	10.01	
2	50	142.08	2.5	0.99	10	3.0	0.3	1.82	

Brain-to-Plasma Ratio of Compounds in Vivo

[0283] The brain-to-plasma ratio of trimetazidine and CV-8814 was analyzed after intravenous administration of the compounds to rats. Dosing solutions were analyzed by liquid chromatography tandem mass spectrometry (LC-MS/MS). Results are shown in Table 67.

TABLE 67

Test Article	Route of Administration	Vehicle	Nominal	Measured	
			Dosing Conc. (mg/mL)	Solution Conc. (mg/mL)	% of Nominal
TMZ	IV	Normal Saline*	1.0	1.14	114
CV-8814	IV	Normal Saline*	0.585	0.668	114

[0284] The concentrations of compounds in the brain and plasma were analyzed 2 hours after administering compounds at 1 mg/kg to rats. Results from trimetazidine-treated rats are shown in Table 68. Results from CV-8814-treated rats are shown in Table 69.

TABLE 68

Rat#	11	12	13
Brain Weight (g)	1.781	1.775	1.883
Brain Homogenate Volume (mL)	8.91	8.88	9.42
Brain Homogenate Conc. (ng/mL)	7.08	7.35	7.90
Brain Tissue Conc. (ng/g)	35.4	36.8	39.5
Plasma Conc. (ng/g) ¹	22.7	14.0	14.1
B:P Ratio	1.56	2.63	2.80

TABLE 69

Rat#	14	15	16
Brain Weight (g)	1.857	1.902	2.026
Brain Homogenate Volume (mL)	9.29	9.51	10.1
Brain Homogenate Conc. (ng/mL)	4.01	4.21	4.74
Brain Tissue Conc. (ng/g)	20.1	21.1	24
Plasma Conc. (ng/g) ¹	19.3	17.0	14.0
B:P Ratio	1.04	1.24	1.693

[0285] The average B: P ratio for trimetazidine-treated rats was 2.33 ± 0.672 . The average B: P ratio for trimetazidine-treated rats was 1.32 ± 0.335 .

INCORPORATION BY REFERENCE


[0286] References and citations to other documents, such as patents, patent applications, patent publications, journals, books, papers, web contents, have been made throughout this disclosure. All such documents are hereby incorporated herein by reference in their entirety for all purposes.

EQUIVALENTS

[0287] Various modifications of the invention and many further embodiments thereof, in addition to those shown and described herein, will become apparent to those skilled in the art from the full contents of this document, including references to the scientific and patent literature cited herein. The subject matter herein contains important information, exemplification and guidance that can be adapted to the practice of this invention in its various embodiments and equivalents thereof.

1. A HCl salt of a compound represented by formula (X):

(X)



2. The HCl salt of claim **1**, wherein the salt is a monohydrate.

3. A pharmaceutical composition comprising the HCl salt of claim **1**.

4. The pharmaceutical composition of claim **3**, wherein the composition comprises the compound in a therapeutically effective amount to increase cardiac efficiency in a subject.

5. The pharmaceutical composition of claim **3**, wherein the composition is formulated for oral administration.

6. The pharmaceutical composition of claim **5**, wherein the composition comprises a format selected from the group consisting of a tablet, troche, lozenge, hard capsule, and soft capsule.

* * * * *