

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12383543
Kind Code	B2
Date of Patent	August 12, 2025
Inventor(s)	Gottardis; Marco et al.

Methods of treating prostate cancer

Abstract

Disclosed are methods of treating prostate cancer by administering niraparib to a human in need thereof.

Inventors: Gottardis; Marco (Princeton, NJ), Hawkins; Rebecca (Harleysville, PA), Snyder; Linda A. (Pottstown, PA), Yamada; Douglas H. (Philadelphia, PA)

Applicant: Janssen Pharmaceutica NV (Beerse, BE)

Family ID: 59649991

Assignee: Janssen Pharmaceutica NV (Beerse, BE)

Appl. No.: 17/528017

Filed: November 16, 2021

Prior Publication Data

Document Identifier	Publication Date
US 20220071979 A1	Mar. 10, 2022

Related U.S. Application Data

continuation parent-doc US 16131772 20180914 US 11207311 child-doc US 17528017
continuation parent-doc US 15663082 20170728 ABANDONED child-doc US 16131772
us-provisional-application US 62369239 20160801
us-provisional-application US 62368466 20160729

Publication Classification

Int. Cl.: A61K31/454 (20060101); A61K9/00 (20060101); A61P13/08 (20060101); A61P35/00 (20060101); A61P35/04 (20060101)

U.S. Cl.:

CPC **A61K31/454** (20130101); **A61K9/0053** (20130101); **A61P13/08** (20180101);
A61P35/00 (20180101); **A61P35/04** (20180101);

Field of Classification Search

CPC: A61K (31/454); A61K (9/0053); A61K (2300/00); A61K (31/573); A61K (31/58); A61K
(45/06); A61P (35/00); A61P (13/08); A61P (35/04)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
4717721	12/1987	Deluca et al.	N/A	N/A
4851401	12/1988	Deluca et al.	N/A	N/A
4857518	12/1988	Deluca et al.	N/A	N/A
4866048	12/1988	Calverley et al.	N/A	N/A
5120722	12/1991	Baggiolini et al.	N/A	N/A
5145846	12/1991	Baggiolini et al.	N/A	N/A
5190935	12/1992	Binderup et al.	N/A	N/A
5237110	12/1992	Deluca et al.	N/A	N/A
5411949	12/1994	Neef et al.	N/A	N/A
5446035	12/1994	Neef et al.	N/A	N/A
5547947	12/1995	Dore et al.	N/A	N/A
5604213	12/1996	Barrie et al.	N/A	N/A
5618807	12/1996	Barrie et al.	N/A	N/A
5688977	12/1996	Sisti et al.	N/A	N/A
6087350	12/1999	Johnson et al.	N/A	N/A
6310226	12/2000	Calverley et al.	N/A	N/A
6521608	12/2002	Henner et al.	N/A	N/A
6559139	12/2002	Johnson et al.	N/A	N/A
6872568	12/2004	Ni et al.	N/A	N/A
7071333	12/2005	Combs et al.	N/A	N/A
7256193	12/2006	Kyle et al.	N/A	N/A
7482334	12/2008	Frincke et al.	N/A	N/A
7547687	12/2008	Reading et al.	N/A	N/A
7709517	12/2009	Sawyers et al.	N/A	N/A
8071623	12/2010	Jones et al.	N/A	N/A
8183274	12/2011	Sawyers et al.	N/A	N/A
8436185	12/2012	Foley et al.	N/A	N/A
8658128	12/2013	Altschul et al.	N/A	N/A
8822438	12/2013	Auerbach et al.	N/A	N/A
9114147	12/2014	Altschul et al.	N/A	N/A
9126941	12/2014	Sawyers et al.	N/A	N/A
9295680	12/2015	Altschul et al.	N/A	N/A
9314473	12/2015	Altschul et al.	N/A	N/A
9320747	12/2015	Altschul et al.	N/A	N/A
9598459	12/2016	Altschul et al.	N/A	N/A

9636351	12/2016	Altschul et al.	N/A	N/A
9642866	12/2016	Altschul et al.	N/A	N/A
9855284	12/2017	Altschul et al.	N/A	N/A
9861643	12/2017	Altschul et al.	N/A	N/A
10058563	12/2017	Altschul et al.	N/A	N/A
10076528	12/2017	Altschul et al.	N/A	N/A
10231982	12/2018	Altschul et al.	N/A	N/A
10238666	12/2018	Altschul et al.	N/A	N/A
10517881	12/2018	Altschul et al.	N/A	N/A
10537586	12/2019	Altschul et al.	N/A	N/A
11040037	12/2020	Altschul et al.	N/A	N/A
11207311	12/2020	Gottardis et al.	N/A	N/A
11224599	12/2021	Altschul et al.	N/A	N/A
11364252	12/2021	Altschul et al.	N/A	N/A
11576921	12/2022	Altschul et al.	N/A	N/A
2002/0128240	12/2001	Mazess	N/A	N/A
2003/0083231	12/2002	Ahlem et al.	N/A	N/A
2003/0119795	12/2002	Henner et al.	N/A	N/A
2004/0138187	12/2003	Reading et al.	N/A	N/A
2005/0020546	12/2004	Laidlaw et al.	N/A	N/A
2005/0054620	12/2004	Koeffler et al.	N/A	N/A
2005/0101581	12/2004	Reading et al.	N/A	N/A
2005/0233958	12/2004	Ni et al.	N/A	N/A
2006/0003021	12/2005	Strugnell et al.	N/A	N/A
2006/0003950	12/2005	Strugnell et al.	N/A	N/A
2006/0018933	12/2005	Vaya et al.	N/A	N/A
2006/0018934	12/2005	Vaya et al.	N/A	N/A
2006/0024365	12/2005	Vaya et al.	N/A	N/A
2006/0030608	12/2005	Nelson et al.	N/A	N/A
2007/0203107	12/2006	Frincke et al.	N/A	N/A
2007/0213309	12/2006	Reading et al.	N/A	N/A
2007/0265236	12/2006	Reading et al.	N/A	N/A
2007/0275937	12/2006	Reading et al.	N/A	N/A
2007/0275938	12/2006	Reading et al.	N/A	N/A
2008/0004286	12/2007	Wang et al.	N/A	N/A
2008/0004287	12/2007	Ma et al.	N/A	N/A
2008/0051375	12/2007	Auerbach et al.	N/A	N/A
2008/0085873	12/2007	Reading et al.	N/A	N/A
2008/0138426	12/2007	Hara et al.	N/A	N/A
2008/0167345	12/2007	Jones et al.	N/A	N/A
2009/0124587	12/2008	Auerbach et al.	N/A	N/A
2012/0201747	12/2011	Altschul et al.	N/A	N/A
2014/0107086	12/2013	Theise et al.	N/A	N/A
2014/0315920	12/2013	Virca et al.	N/A	N/A
2014/0336154	12/2013	Do et al.	N/A	N/A
2014/0336157	12/2013	Auerbach et al.	N/A	N/A
2015/0344968	12/2014	Johnson	N/A	N/A
2015/0366881	12/2014	Altschul et al.	N/A	N/A
2015/0366882	12/2014	Altschul et al.	N/A	N/A
2016/0113948	12/2015	Altschul et al.	N/A	N/A

2016/0279148	12/2015	Altschul et al.	N/A	N/A
2016/0279149	12/2015	Altschul et al.	N/A	N/A
2017/0051007	12/2016	Altschul et al.	N/A	N/A
2017/0128465	12/2016	Altschul et al.	N/A	N/A
2017/0202859	12/2016	Altschul et al.	N/A	N/A
2017/0232005	12/2016	Altschul et al.	N/A	N/A
2018/0017438	12/2017	Crohn et al.	N/A	N/A
2018/0028521	12/2017	Gottardis et al.	N/A	N/A
2018/0117066	12/2017	Altschul et al.	N/A	N/A
2018/0185392	12/2017	Altschul et al.	N/A	N/A
2018/0296574	12/2017	Snyder et al.	N/A	N/A
2018/0325920	12/2017	Altschul et al.	N/A	N/A
2018/0360852	12/2017	Altschul et al.	N/A	N/A
2019/0022079	12/2018	Gottardis et al.	N/A	N/A
2019/0134062	12/2018	Altschul et al.	N/A	N/A
2019/0151335	12/2018	Altschul et al.	N/A	N/A
2019/0381038	12/2018	Altschul et al.	N/A	N/A
2020/0101087	12/2019	Altschul et al.	N/A	N/A
2021/0353623	12/2020	Altschul et al.	N/A	N/A
2021/0361675	12/2020	Altschul et al.	N/A	N/A
2022/0298203	12/2021	Altschul et al.	N/A	N/A
2023/0142627	12/2022	Altshul et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
101528308	12/2008	CN	N/A
104306977	12/2014	CN	N/A
10061137	12/2001	DE	N/A
0413270	12/1990	EP	N/A
0912535	12/1998	EP	N/A
0914116	12/1998	EP	N/A
1140192	12/2000	EP	N/A
1336602	12/2002	EP	N/A
1379246	12/2003	EP	N/A
1385514	12/2003	EP	N/A
1385515	12/2003	EP	N/A
1385518	12/2003	EP	N/A
1423381	12/2003	EP	N/A
1463733	12/2003	EP	N/A
1466628	12/2003	EP	N/A
1487829	12/2003	EP	N/A
1515949	12/2004	EP	N/A
1556354	12/2004	EP	N/A
1562932	12/2004	EP	N/A
1562936	12/2004	EP	N/A
1572299	12/2004	EP	N/A
1583524	12/2004	EP	N/A
1583763	12/2004	EP	N/A
1598338	12/2004	EP	N/A
1598339	12/2004	EP	N/A

1598340	12/2004	EP	N/A
1631285	12/2005	EP	N/A
1307197	12/2005	EP	N/A
1648879	12/2005	EP	N/A
1664016	12/2005	EP	N/A
1664041	12/2005	EP	N/A
1674479	12/2005	EP	N/A
1676577	12/2005	EP	N/A
1803718	12/2006	EP	N/A
1810970	12/2006	EP	N/A
1412368	12/2006	EP	N/A
1673092	12/2006	EP	N/A
1862458	12/2006	EP	N/A
1867644	12/2006	EP	N/A
1918279	12/2007	EP	N/A
1927858	12/2007	EP	N/A
1930322	12/2007	EP	N/A
1942106	12/2007	EP	N/A
1975164	12/2007	EP	N/A
2336120	12/2010	EP	N/A
2478907	12/2011	EP	N/A
2061561	12/2012	EP	N/A
2805945	12/2013	EP	N/A
2007733	12/2015	EP	N/A
2003-525254	12/2002	JP	N/A
2006-515623	12/2005	JP	N/A
2010-501575	12/2009	JP	N/A
2010-515715	12/2009	JP	N/A
92/00992	12/1991	WO	N/A
93/20097	12/1992	WO	N/A
95/09178	12/1994	WO	N/A
01/64251	12/2000	WO	N/A
01/81364	12/2000	WO	N/A
01/93836	12/2000	WO	N/A
02/03286	12/2001	WO	N/A
02/32861	12/2001	WO	N/A
02/53138	12/2001	WO	N/A
02/85355	12/2001	WO	N/A
02/85361	12/2001	WO	N/A
02/91993	12/2001	WO	N/A
2002/102783	12/2001	WO	N/A
03/20699	12/2002	WO	N/A
03/22835	12/2002	WO	N/A
03/37252	12/2002	WO	N/A
03/39460	12/2002	WO	N/A
03/86388	12/2002	WO	N/A
03/86404	12/2002	WO	N/A
03/92595	12/2002	WO	N/A
2004/012699	12/2003	WO	N/A
2004/012700	12/2003	WO	N/A

2004/016753	12/2003	WO	N/A
2004/037269	12/2003	WO	N/A
2004/041164	12/2003	WO	N/A
2004/062620	12/2003	WO	N/A
2005/016236	12/2004	WO	N/A
2005/021487	12/2004	WO	N/A
2005/107801	12/2004	WO	N/A
2006/004917	12/2005	WO	N/A
2006/021776	12/2005	WO	N/A
2006/027266	12/2005	WO	N/A
2006/050402	12/2005	WO	N/A
2006/081152	12/2005	WO	N/A
2006/116217	12/2005	WO	N/A
2006/116716	12/2005	WO	N/A
2007/014327	12/2006	WO	N/A
2008/024484	12/2007	WO	N/A
2008/039254	12/2007	WO	N/A
2008/062466	12/2007	WO	N/A
2008/084261	12/2007	WO	N/A
2008/100985	12/2007	WO	N/A
2008/109740	12/2007	WO	N/A
2008/127290	12/2007	WO	N/A
2009/087381	12/2008	WO	N/A
2012/009475	12/2011	WO	N/A
2012/106514	12/2011	WO	N/A
2014/089324	12/2013	WO	N/A
2015/164586	12/2014	WO	N/A
2016/094391	12/2015	WO	N/A
2017/023694	12/2016	WO	N/A
2018/067520	12/2017	WO	N/A
2019/074536	12/2018	WO	N/A

OTHER PUBLICATIONS

Erica L. T. Van Den Akker et al., “Differential Inhibition of 17alpha-hydroxylase and 17,20-lyase activities by three novel missense CYP17 mutations identified in patients with P450c17 deficiency”, *The Journal of Clinical Endocrinology & Metabolism* Dec. 2002, vol. 87 No. 12 pp. 5714-5721. cited by applicant

Erie J. Small et al., “Antiandrogen Withdraw-al Alone or in Combination With Ketoconazole in Androgen-Independent Prostate Cancer Patients: A Phase III Trial (CALGB9583)”, *Journal of Clinical Oncology*, *Wockhardt vs. Janssen*, Filed for Case IPR2016-01582, Janssen Exhibit 2172, on Apr. 3, 2017, 18 pages. cited by applicant

Ernel Arinc et al., “Molecular Aspects of Monooxygenases and Bioactivation of Toxic Compounds”. Series A: Life Sciences vol. 202 Mylan Pharms. Inc., 28 pages. cited by applicant
Excerpts from Seifter, *Concepts in Medical Physiology*, Chapter 34, pp. 540-553, Chapter 37, pp. 606-620 (2005), Filed for Case IPR2016-00286, Janssen Exhibit 2058. cited by applicant

F. Labrie et al., “Anti-hormone Treatment for Prostate Cancer Relapsing after Treatment with Flutamide and Castration”, *British Journal of Urology*, (1989) 63 ,pp. 634-638. cited by applicant

Fable Zustovich and Davide Pastorelli, *Therapeutic management of bone metastasis in prostate cancer: an update*, *Expert Review of Anticancer Therapy*,

<http://dx.doi.org/10.1080/14737140.2016.1241148>, Sep. 27, 2016. cited by applicant

Fakih, et al., *Urology* (2002) 60, p. 553-561. cited by applicant

Fallowfield et al., “Patients' preference for administration of endocrine treatments by injection or tablets”, results from a study of women with breast cancer, *Annals of Oncology* vol. 17 No. 2 Feb. 2000, pp. 205-210. cited by applicant

Farwell, et al., “Total Suppression of Cortisol Excretion by Ketoconazole in the Therapy of the Ectopic Adrenocorticotrophic Hormone Syndrome”, *American Journal of Medicine*, filed for case IPR2016-00286, Janssen Exhibit 2065, on Jun. 1988, vol. 84, pp. 1063-1006. cited by applicant

Final Written Decision, United States Patent and Trademark Office Before the Patent Trial and Appeal Board, Case No. IPR2016-01582 U.S. Pat. No. 8,822,438 82, Jan. 17, 2018, 51 pages. cited by applicant

Fizazi et al., “Abiraterone plus Prednisone in Metastatic, Castration-Sensitive Prostate Cancer”, *The new england journal of medicine*, Jul. 2017, vol. 377, No. 4, pp. 352-360. cited by applicant

Fizazi et al., “Low Incidence of Corticosteroid-associated Adverse Events on long-term Exposure to Low-dose Prednisone Given with Abiraterone Acetate to Patients with Metastatic Castration-resistant Prostate Cancer”, *Eur Urol*. Sep. 2016, vol. 70, No. 3, pp. 438-444. cited by applicant

Fossa et al., “Flutamide versus prednisone in patients with prostate cancer symptomatically progressing after androgen-ablative therapy: a phase III study of the European organization for research and treatment of cancer genitourinary group,” *Journal of Clinical Oncology*, vol. 19(1) 62-71 (2001). cited by applicant

Fossa, et al., *Weekly Docetaxel and Prednisone Versus Prednisolone Alone in Androgen-Independent Prostate Cancer: A Randomized Phase II Study*, *European Urology*, 2007, pp. 1691-1699, vol. 52. cited by applicant

Fourteenth Edition Physicians' Desk Reference to Pharmaceutical Specialties and Biologicals, 1960, PDR, 2 pages. cited by applicant

Friel, Patrick N., et al., *Suppression of adrenal function by low-dose prednisone: assessment with 24-hour urinary steroid hormone profiles—A review of five cases* , *Alternative Medicine Review* (2006), 11(1). cited by applicant

G. Sonpavde et al., *Impact of single-agent daily prednisone on outcomes in men with metastatic castration-resistant prostate cancer*, *Prostate Cancer and Prostatic Diseases* (2016) 00, pp. 1-5. cited by applicant

Garnick Deposition Blackhard, Letter to Editor, *The Journal of Urology*. “Official Journal of The American Urological Association, Inc.”, file for Case IPR2016-01332, Janssen Exhibit 2011 on Dec. 1991, vol. 146, No. 6, pp. 1621-1622. cited by applicant

Garnick et al., “Androgen deprivation therapy: the future”, *Prostate Cancer Principles and Practice*, 19 pages. cited by applicant

Genentech Provides Update on Phase III Study of Avastin in Men With late Stage Prostate Cancer Janssen Exhibit 2081, *Amerigen vs. Janssen* filed for Case# IPR2016-00286, Mar. 12, 2010, 3 Pages. cited by applicant

Geoff Cumming, “Inference by eye: reading the overlap of independent confidence intervals”, *Statistics in Medicine*, vol. 28, 2009, pp. 205-220. cited by applicant

Gerber, et al., *Prostate Specific Antigen for Assessing Response to Ketoconazole and Prednisone in Patients with Hormone Refractory Metastatic Prostate Cancer*, *The Journal of Urology*, 1990, pp. 1177-1179, vol. 1444, No. 5. cited by applicant

Ghatana et al... “Effect of Single-agent Daily Prednisone on Outcomes and Toxicities in Metastatic Castration-resistant Prostate Cancer: Pooled Analysis of Prospective Studies”, *Clinical Genitourinary Cancer*, vol. 16, No. 2, April 2018, pp. e277-e287. cited by applicant

Gignac et al., “Castration Resistant, Taxane Naive Metastatic Prostate Cancer: Current Clinical Approaches and Future Directions,” *J. Urology*, vol. 178:S30-S35 (2007). cited by applicant

Gill et al., “Efficacy of Eplerenone in the Management of Mineralocorticoid Excess In Men With Metastatic Castration-resistant Prostate Cancer Treated With Abiraterone Without Prednisone”,

Clinical Genitourinary Cancer; vol. 15, No. 4, Aug. 2017, pp. e599-e602. cited by applicant

Gleave et al., "Use of antisense oligonucleotides targeting the cytoprotective gene, clusterin, to enhance androgen- and chemo-sensitivity in prostate cancer", *World J Urol.* Feb. 2005, vol. 23, No. 1, pp. 38-46. cited by applicant

Gordon Williams et al., "Objective Responses to Ketoconazole Therapy in Patients with Relapsed Progressive Prostatic Cancer" *British Journal of Urology* vol. 58, pp. 45-51 (1986). cited by applicant

Gras Jordi: "Niraparib hydrochloride. Poly [ADP-ribose] polymerase (PARP) inhibitor, Oncolytic", *Drugs of the Future*, Prous Science, ES, vol. 38, No. 10, Oct. 1, 2013 (Oct. 1, 2013), pp. 679-685. cited by applicant

Grinspoon and Biller, "Clinical Review 62 Laboratory Assessment of Adrenal Insufficiency", *Journal of Clinical Endocrinology and Metabolism*, Filed for Case IPR2016-00286, Janssen Exhibit 2052, on 1994, vol. 79, No. 4, pp. 923-931. cited by applicant

Grove, M. et al., "Bioavailability of Seocalcitol i: Relating Solubility in Biorelevant Media with Oral Bioavailability in Rats—Effect of Medium and Long Chain Triglycerides", 94(8) *J. Pharm. Sci.* No. 1830-1838 (2005). cited by applicant

H.I. Scher et al., Design and End Points of Clinical Trials for Patients with Progressive Prostate Cancer and Castrate Levels of Testosterone: Recommendations of the Prostate cancer Clinical Trials Working Group, *Journal of Clinical Oncology*, vol. 26, No. 7, at 1148-1159 (Mar. 1, 2008). cited by applicant

Hadaschik et al., "Novel targets and approaches in advanced prostate cancer", *Current Opinion in Urology* 2007, vol. 17: pp. 182-187. cited by applicant

Haidar et al., "Effects of novel 17alpha-hydroxylase/C17.20-lyase (P450 17 CYP 17) inhibitors on androgen biosynthesis in vitro and in vivo," *J. Steroid Bioch. & Mol. Biol.* vol 84:555-62 (2003). cited by applicant

Haidar et al., "Novel steroidal pyrimidyl inhibitors of P450 17 (17alpha-hydroxylase/C17-20-lyase)," *Arch. Pharm. Pharm. Med. Chern.*, vol. 334:373-374 (2001). cited by applicant

Hakki et al., CYP17- and CYP11B-dependent steroid hydroxylases as drug development targets, Elsevier. 2006, pp. 27-52, vol. 11. cited by applicant

Hansen. C. M. et al., "Seocalcitol (EB 1089): A Vitamin D. Analogue of Anti-Cancer Potential. Background, Design, Synthesis,. Pre-Clinical and Clinical Evaluation", 6 *Current Pharmaceutical Design*, 803-828 (2000). cited by applicant

Harris, et al., Low Dose Ketoconazole with Replacement Doses of Hydrocortisone in Patients with Progressive Androgen Independent Prostate Cancer, *The Journal of Urology*, 2002, pp. 542-545, vol. 168. cited by applicant

Harrison's 15th Edition Principles of Internal Medicine, Braunwald Fauci Kasper Hauser Longo Jameson, *N Engl J. Med.* 341: 156, 1999. cited by applicant

Hartmann et al., "Synthesis and evaluation of novel steroidal oxime inhibitors of P450 17 (17alpha-hydroxylase/C17-20-lyase) and 5alpha-reductase types 1 and 2," *J. Med. Chem.*, vol. 43:4266-4277 (2000). cited by applicant

Harzstark et al., "Therapies in Development for Castrate-Resistant Prostate Cancer," *Expert Rev. Anticancer Ther.*; vol. 8(2):259-268 (2008). cited by applicant

Harzstark et al., "Novel Therapeutic Strategies in Development for Prostate Cancer," *Expert Opin. Invest. Drugs*, vol. 17(1):13-22 (2008). cited by applicant

Haynes et al., "Pharmacology of CB759S, a Highly Potent Inhibitor of Cytochrome P450c17." *Proceedings American Association for Cancer Research*, vol. 35, Eighty-fifth Annual Meeting, abstract No. 2507 (1994). cited by applicant

Hellersledt et al., "The current state of hormonal therapy for prostate cancer", *CA A Cancer Journal for Clinicians*, May-Jun. 2002, vol. 52, No. 3, pp. 154-179. cited by applicant

Heremans GF, Moolenaar AJ. van Gelderen HH. Female phenotype In a male child due to 17-

alpha-hydroxylase deficiency. Arch Dis Child, file for Case IPR2016-00286, Amerige11 Exhibit 1167 on 1976, vol. 51. No. 9 pp. 721-723. cited by applicant

Herr and Pfizenmaier. "Gkicocorticoid use in prostate cancer and other solid tumours: implications for effectiveness o cytotoxic treatment and metastases," The Lancet Oncol, Filed for Case IPR2016-00286, ..Janssen Exhibit 2023, 01: May 2006, vol. 7 pp. 425-430. cited by applicant

Highlights of Alimta prescribing information. cited by applicant

Highlights of Zytiga Prescribing Information, 2015. cited by applicant

Highlights of Zytiga Prescribing Information, 2018. cited by applicant

Hildesheim, "Prostate cancer pill extends life", Lancastria.net. May 26, 2011, 5 pages. cited by applicant

Horsham, Pa, Dec. 10, 2012—US. FDA Approves Expanded Zytiga (Registered) Indication For Treatment Of Metastatic Castration-Resistant Prostate Cancer Johnson & Johnson, Services, Inc. 1997-2017. 5 pages. cited by applicant

Robert Twycross, "Corticosteroids in Advanced Cancer".filed for Case# IPR2016-00286, Vot 305, pp. 969-970, Oct. 24, 1992. cited by applicant

Ross et al., "Hormone Refractory Prostate Cancer: Choosing the Appropriate Treatment Option," Oncology, vol. 21(2):185-193 (2007). cited by applicant

Rowlands et al., Esters of 3-Pyridylacetic Acid that Combine Potent Inhibition of 17alpha-Hydroxylase/C17.20-lyase (Cytochrome P45017alpha) with Resistance to Esterase Hydrolysis, J. Med. Chem., vol. 38:4191-4197 (1995). cited by applicant

Ruben H Munoz, Akin Gump Strauss H,t.\UER & Feld LLP, "Zytiga IPR", Mailed for Cast>.# IPR2016-00286, Jun. 29, 2016, 1 page. cited by applicant

Rumohr et al., "Current Chemotherapeutic approaches for androgen-independent prostate cancer", Current Opinion in Investigational Drug, Filed for Case IPR2016-01332. Janssen Exhibit 2027, on 2006, vol. 7. No. 6, pp. 529-533. cited by applicant

Runge, Marschall S., et al., Principles of Molecular Medicine; Second edition; (2006) Humana Press Inc. ISBN: -1-58829-202-9. pp. 365-376 and 482-484. cited by applicant

Ryan Charles J et al: "Abiraterone acetate plus prednisone versus placebo plus prednisone in chemotherapy-naïve men with metastatic castration-resistant prostate cancer (COU-AA-302): final overall survival analysis of a randomised, double-blind, placebo-controlled phase 3 study." , The Lancet. Oncology, vol. 16, No. 2, Feb. 2015 (Feb. 2015) pp. 152-160. cited by applicant

Ryan et al. "Impact of Prior Ketoconazole Therapy on Response Proportion to Abiraterone Acetate, a 17 alpha hydroxylase C17,20-Lyase Inhibitor in Castration Resistant Prostate Cancer (CRPC)," presentation (2008). cited by applicant

Ryan et al. , "Impad of prior ketoconazole therapy on response proportion to abiraterone acetate, a 17-alpha hydroxy lase C17.20-lyase inhibitor in castration resistant prostate cancer (CRPC)," J. Clin. Oncol. (Meeting abstracts), vol. 26 (May 20 Supplement), abstract No. 5018 (2008). cited by applicant

Ryan et al., "Abiraterone Acetare Plus Prednisone in Chemotherapy (Chemo)-Naive castration Resistant Prostate Cancer (CPRC) Patients Not Exposed to Ketoconazole: Results of a Multi-Center Phase II Study," 2009 Genitourinary cancers Symposium (Feb. 26-28, 2009). abstract Submission (2008). cited by applicant

Ryan et al., "Effect of Concornitant Food Intake on Pharmacokinetics of Abiraterone Acetate, a 17 alpha Hydroxylase C17,20-Lyase Inhibitor in Castration-Resistant Prostate Cancer (CRPC)," Molecular Cancer Therapeutics (Dec. 2007), vol. 6(12):3527s, 2007 AACR-NCI-EORTC International Conference, Poster Session C, abstract No. C2 (2007). cited by applicant

Ryan et al., "Phase I Evaluation of Abiraterone Acetate (CB7630), a 17 alpha hydroxylase C17,20-Lyase Inhibitor as Secondary Hormonal Therapy in Castration Resistant Prostate Cancer (CRPC)," poster (undated). cited by applicant

Ryan et al., "Phase II Study of Abiraterone In Chemotherapy-Naive Metastatic Castration-

Resistant Prostate Cancer Displaying Bone Flare Disocordant with Serologic Response”, Clinical Cancer Research, 2011, Filed for Case IPR2016-00286, Janssen Exhibit 2017, vol. 17, No. 14, pp. 4854-4861. cited by applicant

Ryan et al., “Phase I Evaluation of Abiraterone Acetate (CB7630), a 17 alpha hydroxy lase C17,20-Lyase Inhibitor in Androgen-Independent Prostate Cancer (AIPG).” ASCO 2007 Prostate Cancer Symposium, abstract No. 278 (2007). cited by applicant

Ryan et al., “Phase I Evaluation of Abiraterone Acetate (CB7630), a 17 alpha hydroxylase C17,20-Lyase Inhibitor in Androgen-Independent Prostate Cancer (AIPC).” J. Clin. Oncol.. vol. 25 (18S):250s, ASCO Annual Meeting Proceedings Part I, abstract No. 5064 (2007). cited by applicant

Ryan et al., “Prostate Specific Antigen Only Androgen Independent Prostate Cancer: Natural History, Challenges in Management and Clinical Trial Design.” J, Urology, vol. 178:S25-S29 (2007). cited by applicant

Ryan, “Secondary Hormonal Manipulations in Prostate Cancer,” Hematology/Oncology Clinics of North America, vol. 20(4):925-934 (2006). cited by applicant

S, Halabi et al., Prostate-Specific Antigen Changes As Surrogate for Overall Survival in Men With Metastatic Castration-Resistant Prostate Cancer Treated With Second-Line Chemotherapy, Journal of Clinical Oncology, vol. 31, No. 31, pp. 3944-3950 (Nov. 1, 2013). cited by applicant

S, Wilkinson, S. and G, Chodak., “An Evaluation of Intermediate-Dose Ketoconazole in Hormone Refractory Prostate Cancer,” European Urology, vol. 45, pp. 581-585, Nov. 26, 2004. cited by applicant

S. M. Withelm et al., BAY 43-9006 Exhibits Broad Spectrum Oral Antitumor Activity and Targets the RAF/MEK/ERK Pathway and Receptor Tyrosine Kinases Involved in Tumor Progression and Angiogenesis, Cancer Research, filed for Case# IPR2016-00286, vol. 64, pp. 7099-7109, (Oct. 1, 2004). cited by applicant

S. Oudard et al., Prostate-Specific Antigen Doubling Time Before Onset of Chemotherapy as a Predictor of Survival for Hormone-Refractory Prostate Cancer Patients, Annals of Oncology vol. 18, pp. 1828-1833, (Nov. 2007). cited by applicant

S. Udhane et al., Specificity of Anti-Prostate Cancer CYP17A1 Inhibitors on Androgen Biosynthesis, Biochemical and Biophysical Research Communications, filed for Case# IPR2016-00286, vol. 477, pp. 1005-1010, (Elsevier 2016). cited by applicant

Sabroe, T.P. et al., “An Efficient Synthesis of a Key Intermediate for the Biologically Active Vitamin D Analogue. Seocalcitol”, Organic Process Research & Dev. vol. 8(1): 133-35 (2004). cited by applicant

Sahu, B., et al., “FoxA1 Specifies Unique Androgen and Glucocorticoid Receptor Binding Events in Prostate Cancer Cells” Cancer Research (2013), vol. 73, pp. 1570-1580. cited by applicant

Sandhu et al. “Phase I Study of Poly(ADP)-Ribose Polymerase (PARP) Inhibitor MK-4827(MK) with Antitumor Activity in Sporadic Castration Resistant Prostate Cancer (CRPC)”, Asia-Pac J. Clin. Oncol., vol. 8(Suppl. 1), pp. 29-34, 2012 (Abstract 112). cited by applicant

Sandhu et al., “The poly(ADP-ribose) polymerase inhibitor niraparib (MK4827) in BRCA mutation carriers and patients with sporadic cancer: a phase 1 dose-escalation trial”, Lancet Oncol, vol. 14(9), 2013 , pp. 882-892. cited by applicant

Sartor et al., “Combination therapy: Abiraterone prolongs survival in metastatic prostate cancer”. Nature Reviews Clinical Oncology, Aug. 2, 2011, vol. 8, No. 9, pp. 515-516. cited by applicant

Sartor et al., “Novel Therapeutic Strategies for Metastatic Prostate Cancer in the Post-Docetaxel Setting”, Academia-Pharma Intersect Genitourinary Cancer: Prostate, The Oncologist, Nov. 2, 2011. vol. 16, pp. 1487-1497. cited by applicant

Sartor, et al., Abiraterone Prolongs Survival in Metastatic Prostate Cancer, Nature Reviews Clinical Oncology, 2011, pp. 515-516, vol. 8. cited by applicant

Sartor, Urology, 1998, 252-256. cited by applicant

Saunders et al., “Inhibition of breast and ovarian carcinoma cell growth by 1,25-dihydroxyvitamin

D3 combined with retinoic acid or dexamethasone,” *Anti-Cancer Drugs*, Rapid Communications of Oxford, vol. 6 (4) 562-569 (1995). cited by applicant

Scher et al., “Bicalutamide for Advanced Prostate Cancer: The Natural Versus Treated History of Disease”, *Journal of Clinical Oncology*, file for Case IPR2016-00286, Janssen Exhibit 2055, vol. 15, No. 8 on Aug. 1997, pp. 2928-2938. cited by applicant

Scher et al., “Biology of Progressive, Castration-Resistant Prostate Cancer: Directed Therapies Targeting the Androgen-Receptor Signaling , Axis,” *J. Clin. Oncol.* vol. 23(32):8253-61 (2005). cited by applicant

Scher H.I. et al., “Increased survival with enzalutamide in prostate cancer after chemotherapy” *New Eng. J. Med.*, vol. 367 No. 13, Sep. 27, 2012, pp. 1187-1197. cited by applicant

Schiewer et al, *parp1 promotes prostate tumor growth and progression, with veliparib as the parp inhibitor tested*; *Cancer Discovery* 2:1134, 2012. cited by applicant

Scholz, et al, *Long-Term Outcome for Men with Androgen Independent Prostate Cancer Treated with Ketoconazole and Hydrocortisone*, *The Journal of Urology*, 2005, pp. 1947-1952. vol. 173. cited by applicant

Schroder, “Progress in Understanding Androgen-Independent Prostate Cancer (APC): A Review of Potential Endocrine-Mediated Mechanisms,” *European Urology*, vol. 53: 1129-1137 (2008). cited by applicant

Schulte et al., *The corticotropinreleasing hormone stimulation test a possible aid in the evaluation of patients with adrenal insufficiency* *Journal of Clinical Endocrinology Metabolism* 1984, vol. 58 No. 6, pp. 1064-1067. cited by applicant

Seale and Compton, “Side-effects of corticosteroid agents”, *The Medical Journal of Australia* vol. 144, No. 3 Feb. 3, 1986 pp. 139-142. cited by applicant

Seocalcitrol Versus Placebo in Advanced Hepatocellular Carcinoma LEO Pharma Study [http://www.clinicaltrials.gov/c1/show,No. 100051532](http://www.clinicaltrials.gov/c1/show,No.100051532) (2004). cited by applicant

Sephton, et al., “Diurnal Cortisol Rhythm as a Predictor of Breast Cancer Survival”, *Journal of the National Cancer Institute*, vol. 92, No. 12, Jun. 21, 2000, pp. 994-1000. cited by applicant

Seventy One Edition Physicians' Desk Reference, the trusted drug reference for over 70 years, 2017. PDR, 5 pages. cited by applicant

Sharifi et al., “Secondary Hormonal Therapy for Prostate Cancer: What Lies on the Horizon,” *BJU International*, vol. 101:271-74 (2007). cited by applicant

Sills, Irene N., et al., “17 α -Hydroxylase deficiency in a genetic male and female sibling pair”, *Int J. Gynaecol Obstet.*, (1981), 19, 473:479. cited by applicant

Small et al, “Antiandrogen Withdrawal Alone or in Combination with Ketoconazole in Androgen-Independent Prostate Cancer Patients: A Phase III Trial (CALGB 9583)” *Journal of Clinical Oncology*, file for Case IPR2016-01332, Janssen Exhibit 2063, on Mar. 15, 2004, vol. 22, No. 6, pp. 1025-1033. cited by applicant

Small et al., *The Effect of Secondary Hormonal Therapies in the Chemotherapy Age*, *The Journal of Urology*, 2006; pp. S66-S71, vol. 176. cited by applicant

Small, E.J., et al., “Simultaneous Antiandrogen Withdrawal and Treatment with Ketoconazole and Hydrocortisone in Patients with Advanced Prostate Carcinoma”, *American Cancer Soc.*, 80(9), (1997) pp. 1755-1759. cited by applicant

Sonino et al., “The Use of Ketoconazole as an Inhibitor of Steroid Production”, *New England Journal of Medicine*, file for Case IPR2016 01332, Janssen Exhibit 2163, on Sep. 24, 1987, vol. 317, No. 13, pp. 812-818. cited by applicant

Sonpavde et al., *Impact of single-agent daily prednisone on outcomes in men with metastatic castration-resistant prostate cancer*, *Prostate Cancer and Prostatic Diseases*, Mar. 2017, vol. 20, No. 1, pp. 67-71. cited by applicant

St. Louis, *A Division of J.B. Lippincott, Drug Facts and Comparisons*. 1985 Edition, 13 pages. cited by applicant

Jones et al., "Discovery of 2-{4-[(3S)-Piperidin-3-yl]phenyl}-2H-indazole-7-carboxamide (MK-4827): A Novel Oral Poly(ADP-ribose)polymerase (PARP) Inhibitor Efficacious in BRCA-1 and -2 Mutant Tumors", *J. Med. Chem.* 2009, vol. 52, pp. 7170-7185. cited by applicant

Cougar Biotechnology, Cougar Biotechnology announces presentation of positive phase I and phase II data at ASCO Prostate Cancer Symposium, Cougar Biotechnology, Feb. 23, 2007. cited by applicant

Cougar Biotechnology, Cougar Technology Announces Presentation of Positive CB7630 Clinical Data at ASCO Annual Meeting, The Free library, Jun. 4, 2007. cited by applicant

Cougar Biotechnology, Inc., Clinical Study Report: COU-AA-001 and COU-AA 001 EXT, Nov. 17, 2010, 7 pages. cited by applicant

Cowen & Company, "Biotechnology Quarterly." Jul. 2, 2012. cited by applicant

Cowen & Company, "Quick Take: Zytiga Gets FDA OK For Use In Pre-Chemo Setting On rPFS Data—Johnson & Johnson." Accessed (Dec. 11, 2012). cited by applicant

Crawford et al., "Treating Patients with Metastatic Castration Resistant Prostate Cancer", A Comprehensive Review of Available Therapies, *The Journal of Urology*, vol. 194, Dec. 2015, pp. 1537-1547. cited by applicant

Credit Suisse, "Prostate Cancer—Implications of Zytiga's Pre-Chemo Approval" Dec. 11, 2012. cited by applicant

Czock, et al., "Pharmacokinetics and Pharmacodynamics of Systemically Administered Glucocorticoids", *Pharmacokinetics* (2005), 44(1), p. 61-98. cited by applicant

D. Feldman, "Ketoconazole and Other Imidazole derivatives as Inhibitors of steroidogenesis", filed for Case# IPR2016-00286, vol. 7, No. 4, 12 pages. cited by applicant

D. Lorente et al., Tumour responses following a steroid switch from prednisone to dexamethasone in castration-resistant prostate cancer patients progressing on abiraterone, *British Journal of Cancer*, (2014) 111, pp. 2248-2253. cited by applicant

D.E. Rathkopf et al., "Updated Interim Efficacy Analysis and Long-Term Safety of Abiraterone Acetate in Metastatic Castration-Resistant Prostate Cancer Patients Without Prior Chemotherapy(COU-AA-302)". *European Urology*, filed for Case# IPR20.Math.16-00286, No. 66, pp. 815-825, 2014. cited by applicant

Dalhoff, KA, "Phase II Study of the Vitamin D Analogue Seocalcitol in Patients with Inoperable Hepatocellular Carcinoma". 89(2) *British Journal of Cancer*, 252-257 (2003). cited by applicant

Dalia Buffery, "The 2015 Oncology Drug Pipeline: Innovation Drives the Race to Cure Cancer", *Am Health Drug Benefits*, vol. 8(4), 2015, pp. 216-222. cited by applicant

Daniel C Danila et al., "Prednisone Therapy in Patients With Docetaxel-Treated Castration-Resistant Prostate Cancer", *Journal of Clinical Oncology*, vol. 28, No. 9, Mar. 20, 2010. pp. 1496-1501. cited by applicant

Danielenko et al., "Enhancement by other compounds of the anti-cancer activity of vitamin D3 and its analogs," *Experimental Cell Research*, vol. 298 (2):339-358 (2004). cited by applicant

Danila et al., "Abiraterone acetate and prednisone in patients (Pts) with progressive metastatic castration resistant prostate cancer (CRPC) after failure of docetaxel-based chemotherapy," *J. Clin. Oncol. (Meeting abstracts)*, vol. 26 (May 20 Supplement), abstract No. 5019 (2008). cited by applicant

Danila et al., "Preliminary Phase II Results of Abiraterone Acetate in Patients With Castration Resistant Metastatic Prostate Cancer After Failure of Docetaxel-Based Chemotherapy: COU-AA-004," in *Innovative Cancer Therapy for Tomorrow: Foundation Symposium XXV*; presentation (2007). cited by applicant

Danila et al., "Preliminary Phase II Results of Abiraterone Acetate in Patients with Castration-Resistant Metastatic Prostate Cancer After Failure of Docetaxel-Based Chemotherapy," *ASCO 2008 Genitourinary Cancers Symposium*, abstract No. 3 (2008). cited by applicant

Danila et al., Phase II Multicenter Study of Abiraterone Acetate Plus Prednisone Therapy in

Patients With Docetaxel-Treated Castration-Resistant Prostate Cancer. *Journal of Clinical Oncology*, Filed for Case IPR2016-01332, Janssen Exhibit 2016, on Mar. 20, 2010, vol. 28, No. 9, pp. 1496-1501. cited by applicant

Datta et al., *The Journal of Urology*, 1997, 158, 175-177. cited by applicant

De Bono et al. Antitumor Activity of Abiraterone Acetate, a CYP17 Inhibitor That Blocks Androgen Synthesis, in Castration-Resistant Prostate Cancer: ASCO 2008 Annual Meeting, Presentation. cited by applicant

De Bono et al., "Anti-tumor activity of abiraterone acetate (AA), a CYP17 inhibitor of androgen synthesis, in chemotherapy naive and docetaxel pre-treated castration resistant prostate cancer (CRPC)," *J. Clin. Oncol. (Meeting abstracts)*, vol. 26 (May 20 Supplement), abstract No. 5005 (2008). cited by applicant

De Bono et al., Inhibition of CYP17 by abiraterone administered once daily to castrate patients with prostate cancer resistant to LHRH analogues, anti-androgens and steroid therapy is well tolerated, *The Institute of Cancer Research*, 2007. cited by applicant

De Bono et al., Abiraterone and Increased Survival in Metastatic Prostate Cancer, *New England Journal of Medicine*, filed for case IPR2016-01582, Janssen Exhibit 2159, on May 26, 2011, Established in 1812, vol. 364, No. 21, pp. 1995-2005. cited by applicant

De Coster et al., "P450-Dependent Enzymes as Targets for Prostate Cancer Therapy," *J. Steroid Biochem. Mol. Biol.*, vol. 56(1-6):133-143 (1996). cited by applicant

De Coster, et al., Effects of High-Dose Ketoconazole and Dexamethasone on ACTH-Stimulated Adrenal Steroidogenesis in Orchiectomized Prostatic Cancer Patients, *ACTA Endocrinologica (Copenh)*, 1987, pp. 265-271, vol. 115. cited by applicant

DeBono et al., 'Clinical and endocrine evaluation of abiraterone acetate (AA), a rationally designed small molecule inhibitor of androgen synthesis targeting 17 α hydroxylase (17 α H)/17,20 lyase in patients with hormone refractory prostate cancer, 2005 Prostate Cancer Symposium, Abstract No. 290. cited by applicant

Deltasone—prednisone tablet Pharmacia and Upjohn and Company, Deltasone (Registered) prednisone tablets, USP. 13 pages. cited by applicant

DeMario et al., "Oral Chemotherapy", Rationale and Future Directions, *Journal of Clinical Oncology*, vol. 16. No. 7, Jul. 1998, pp. 2557-2567. cited by applicant

Dennis L Kasper, et al (Eds.), *Harrison's Principles of Internal Medicine*, 16th Edition (2005), 12 pages. cited by applicant

Di Cerbo et al., "Combined 17 α -Hydroxylase/17, 20-lyase deficiency caused by Phe93 Cys mutation in the CYP17 gene", *The Journal of Clinical Endocrinology & Metabolism* 2002, vol. 87, No. 2 pp. 898-905. cited by applicant

Dickstein G et al., "One microgram is the lowest ACTH dose to cause a maximal cortisol response. There is No. diurnal variation of cortisol response to submaximal ACTH stimulation". *European Journal of Endocrinology* 1997, vol. 137 No. 2 pp. 172-175. cited by applicant

Dickstein G., et al., "Low' dose ACTH test—A word of caution to the word of caution" when and how to use it *Journal of Clinical Endocrinology and Metabolism* 1997, vol. 82 No. 1, 322 pages. cited by applicant

Division of Endocrinology et al., "Dose-response aspects in the clinical assessment of the hypothalamo pituitary-adrenal axis and the low-dose adrenocorticotropin test", *European Journal of Endocrinology*, 1996, 135: pp. 27-33. cited by applicant

Dizdar 2015, Is Dexamethasone a Better Partner for Abiraterone Than Prednisolone, *The Oncologist* May 2015 vol. 20, No. 5 e 13. cited by applicant

Donald L. Trump et al., High-Dose Ketoconazole in Advanced Hormone-Refractory Prostate Cancer: Endocrinologic and Clinical Effects, *Journal of Clinical Oncology*, vol. 7, No. Aug. 8, 1989, pp. 1093-1098. cited by applicant

Dorff, TB, Crawford, Ed. Management and challenges of corticosteroid therapy in men with

metastatic castrate-resistant prostate cancer, *Annals of Oncology*, 2013. pp. 31-38, vol. 24(1). cited by applicant

Dorin et al., "Diagnosis of Adrenal Insufficiency", *Academia and Clinic, Annals of Internal Medicine*, Filed for Case IPR2016-00286, Janssen Exhibit 2051, on Aug. 5, 2003, vol. 139, No. 3, pp. 194-204. cited by applicant

Duc et al., In vitro and in vivo models for the evaluation of potent inhibitors of male rat 17 (Alpha)-hydroxylase/C17.20-lyase: *Journal of Steroid Biochemistry & Molecular Biology*, 2003. Filed for Case IPR2016-01332, Janssen Exhibit 2012, vol. 84 pp. 537-542. cited by applicant
ECOG Performance Status, "Developed by the Eastern Cooperative Oncology Group, Robert L Combs, MD, Group Chair", *Wockhardt v. Janssen*, Filed for Case# IPR2016-01582, Janssen Exhibit 2158, <http://ecog-acrin.org/resources/ecog-performance-status>, Mar. 14, 2017, 2 pages. cited by applicant

Edward G. Bigueri, et al., "17-Hydroxylation Deficiency in Man", *Journal of Clinical Investigation*, vol. 45, No. 12. 1966,. 10 pages. cited by applicant

Efstathiou et al., "Candidate Predictors of Response to Abiraterone Acetate (AA) in Castrate Resistant Prostate Cancer (CRPC)," 2009 Gemtounnary Cancers Symposium (Feb. 26-28, 2009), abstract Submission (2008). cited by applicant

Efstathiou et al., Identification of an Androgen Withdrawal Responsive Phenotype in Castrate Resistant Prostate Cancer (CRPC) Patients Treated with Abiraterone Acetate; presentation (undated). cited by applicant

Efstathiou, Eleni, et al. Effects of Abiraterone Acetate on Androgen Signaling in Castrate-Resistant Prostate Cancer in Bone, *American Society of Clinical Oncology, Journal of Clinical Oncology*, 2011, pp. 1-8. cited by applicant

Eighteenth Edition Physicians' Desk Reference to Pharmaceutical Specialties and Biologicals, 1964, PDR Published by Medical Economics, Inc. 3 pages. cited by applicant

EMA-Zytiga Product Information, http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-product_Information/human/002321/WC500112858pdf 37 pages. cited by applicant

Endert E. et al., "Establishment of reference values for endocrine tests. Part IV: Adrenal insufficiency", *Netherlands, Journal of Medicine* 2005, vol. 63, No. 11, pp. 435-443. cited by applicant

Ergun-Longmire. Berrin et al., "Two Novel Mutations Found in a Patient with 17 α -Hydroxylase Enzyme Deficiency", *The Journal of Clinical Endocrinology & Metabolism* (2006), 91(10), p. 4179-4182. cited by applicant

Eric J. Small et al., "Ketoconazole Retains Activity in Advanced Prostate Cancer Patients With Progression Despite Flutamide Withdrawal", *The Journal of Urology*, vol. 157 Apr. 1997, pp. 1204-1207. cited by applicant

Eric J. Small et al., "Second-Line Hormonal therapy for Advanced Prostate Cancer: A Shifting Paradigm", *Journal of Clinical Oncology*, vol. 15, No. 1 Jan. 1997, pp. 382-388. cited by applicant

Burke et al. , "Active-Site Conformation of 17-(3-Pyridyl) Androsta-5, 16-Dien-3 β -ol, a Potent Inhibitor of the P450 Enzyme C17 α -Hydroxylase/C17-20 Lyase," *Bioorganic & Medicinal Chemistry letters*, vol. 5(11):1125-1130 (1995). cited by applicant

C.J. Ryan et al., "Abiraterone acetate plus prednisone versus placebo plus prednisone in chemotherapy-naïve men with metastatic castration-resistant prostate cancer (COU-AA-302): final overall survival analysis of a randomised, double-blind, placebo-controlled phase 3 study", *The Lancet*, filed for Case# IPR2016-00286, vol. 16, pp. 152-160, Feb. 2015. cited by applicant

C.J. Ryan et al., Abiraterone in Metastatic Prostate Cancer without Previous Chemotherapy, *The New England Journal of Medicine*, 368; 2, pp. 138-148 (Jan. 10, 2013). cited by applicant

C.N. Sternberg et al., Phase III Trial of Satraplatin, an Oral Platinum plus Prednisone vs. Prednisone alone in Patients with Hormone-Refractory Prostate Cancer, *Oncology*, 68, pp. 2-9

(2005). cited by applicant

C.W. Ryan et al., "Dose-Ranging Study of the Safety and Pharmacokinetics of Atrasentan in Patients with Refractory Malignancies", *Clinical Cancer Research*, filed for IPR2016-00286, vol. 10, pp. 440&-4411, Jul. 1, 2004. cited by applicant

Campbell-Walsh Urology, Ninth Edition, Saunders, vol. 3, Chapters 104 and 105, 2007. cited by applicant

Cancer.0111(ACS), "What are the key statistics about prostate cancer?", <http://www.cancer.org/cancer/prostatecancer/featured-guides/prostate-cancer-key-statistics> (accessed Jun. 28, 2016), Mylan Pharms. Inc., Exhibit 1041, 2 pages. cited by applicant

Cancer.gov (NIH NCI), "Metastatic cancer," <https://www.cancer.gov/types/metastatic-cancer> (accessed Oct. 3, 2016), pp. 3 For Case# IPR2016-00286. cited by applicant

Cancer.gov (NIH NCI), Metastatic cancer <https://www.cancer.gov/about-cancer/treatment/drugs/docetaxel> (accessed Oct. 3, 2016), pp. 3 For Case# IPR2016-01332. cited by applicant

Cancer.Net "Treatment of Metastatic Castration-Resistant Prostate Cancer" filed for Case# IPR2016-00286 on Sep. 8, 2014 3 pages. cited by applicant

Cancer.net (ASCO Patient Website), "Treatment of Metastatic Castration-Resistant Prostate Cancer Sep. 8, 2014", <http://www.cancer.net/research-and-advocacy/asco-care-and-treatment-recommendations-patients/treatment-metastatic-castration-resistant-prostate-cancer> (accessed Jun. 28, 2016), 4 pages. cited by applicant

Cancer.org (ACS), "Hormone therapy for prostate cancer," <https://www.cancer.org/content/cancer/en/cancer/prostatecancer/treating/hormone-therapy-.html> (accessed Apr. 10, 2017), 11 pages. cited by applicant

Cancer.gov (NIH NCI), "Prostate-specific antigen (PSA) test," <http://www.cancer.gov/types/prostate/psa-fact-sheet> (accessed Apr. 11, 2017), 8 pages. cited by applicant

Cannell, 100th Annual Meeting of the American Association for Cancer Research, Los Angeles, CA, USA; <http://oncology.thelancet.com>, 2007, pp. 471, vol. 8. cited by applicant

Carden et al., "Crossover Pharmacokinetics (PK) Study to Assess Oral Administration of Abiraterone Acetate Capsule and Tablet Formulations in Fasted and Fed States in Patients with Prostate Cancer," *J. Clin. Oncol. (Meeting abstracts)*, vol. 26 (May 20 Supplement), abstract No. 5168 (2008). cited by applicant

Carducci et al., "A Phase 3 Randomized Controlled Trial of the Efficacy and Safety of Atrasentan in Men With Metastatic Hormone . . . refractory Prostate Cancer," vol. 110, No. 9, Nov. 1, 2007, pp. 1959-1966. cited by applicant

Carducci, MA, "What is more exciting? The Activity of Docetaxel in Early Prostate Cancer or the Successful Collaboration between Urologists and Medical Oncologists to complete a study in early Prostate Cancer?", *Journal of Clinical Oncology* (2005), vol. 23, No. 15, pp. 3304-3307. cited by applicant

CB1089 <http://www.cougartbiotechnology.com/eb1089.html> (Jul. 2006). cited by applicant

Cecil Textbook of Medicine, Wyngaarden & Smith 18th edition; Chapter on "Glucocorticosteroid Therapy", Wyngaarden & Smith 18th edition, (1988) p. 128-131. cited by applicant

Centers for Disease Control and Prevention, Prostate Cancer Janssen Exhibit 2100, *Amerigen vs. Janssen* filed for Case# IPR2016-00286, 2 Pages. cited by applicant

Chang, Ching-Yi, et al Glucocorticoids Manifest Androgenic Activity in a Cell Derived from a Metastatic Prostate Cancer, *Cancer Research*, 2001, pp. 8712-8717, vol. 61. cited by applicant

Charles et al., "Abiraterone acetate plus prednisone versus placebo plus prednisone in chemotherapy-naïve men with metastatic castration-resistant prostate cancer (COU-AA-302): final overall survival analysis of a randomised, double-blind, placebo-controlled phase 3 study.", *The Lancet. Oncology*, vol. 16, No. 2, Feb. 2015 (Feb. 2015), pp. 152-160. cited by applicant

Charles J. Ryan et al., “Phase I clinical trial of the CYP17 inhibitor abiraterone acetate demonstrating clinical activity in patients with castration-resistant prostate cancer who received prior ketoconazole therapy”, *Journal of Clinical Oncology* Mar. 20, 2010, vol. 28, No. 9, pp. 1481-1488. cited by applicant

Charles L. James et al., “Antimicrobial Therapy, in *Trauma Critical Care*”, vol. 2, (William C. Wilson et al. eds., 2007), pp. 927-960. cited by applicant

CHARTED—ChemoHormonal Therapy versus Androgen Ablation Randomized Trial for Extensive Disease in Prostate Cancer, Declaration of Christopher J. Sweeney, *Clinical Advances in Hematology and Oncology*, filed for Case IPR2016-01582, Janssen Exhibit 2157, on Aug. 2006, vol. 4, No. 8, pp. 588-590. cited by applicant

Chemicals: Seocalcitol
[http://ctd.mdibl.org/voc.go:isessionid"126741EAF326D9CE517F360251236091?voc=chem&acc=C078903&queryferms=seocaldicol&qureyType=oontains&browser=r](http://ctd.mdibl.org/voc.go:isessionid) (Jul. 17, 2006). cited by applicant

Chen, C.D, et al., Molecular determinants of resistance to antiandrogen therapy, *Nat Med*, 10(1), (2004) pp. 33-39. cited by applicant

Cheol Kwak, et al., “Abiraterone acetate and prednisolone for metastatic castration-resistant prostate cancer failing androgen deprivation and docetaxel-based chemotherapy”, *International Journal of Urology* (2014), vol. 21, pp. 1239-1244. cited by applicant

Chi et al., “A Phase I Pharmacokinetic and Pharmacodynamic Study of OGX-011, a 2'-Methoxyethyl Antisense Oligonucleotide to Clusterin, in Patients With Localized Prostate Cancer”, *Journal of the National Cancer Institute*, vol. 97, Issue 17, Sep. 7, 2005, pp. 1287-1296. cited by applicant

Chou et al, “Desoxyepothilone B: An efficacious microtubule-targeted antitumor agent with a promising in vivo profile relative to epothilone B”, *Proc. Natl. Acad. Sci. USA* vol. 95, No. 16. Aug. 1998 *Pharmacology*, pp. 9642-9647. cited by applicant

Christensen G.L., “Sequential Versus Combined Treatment of Human Breast Cancer Cells with Antiestrogens and the Vitamin D Analogue EB 1089 and Evaluation of Predictive Markers for Vitamin D Treatment”, *Breast Cancer Research and Treatment* 85(1) 53-63 (2004). cited by applicant

Clement et al., “Three dimensional pharmacophore modeling of human CYP17 inhibitors. Potential agents for prostate cancer therapy”, *Journal of Medicinal Chemistry*, vol. 46, No. 12, 2003, pp. 2345-2351. cited by applicant

Clinical Cancer Research, “A Journal of the American Association for Cancer Research”, Letter to Ian Judson, Re: “Manuscript# 030579, Hormonal impact of the 17 α -hydroxylase/C17, 20-lyase inhibitor abiraterone acetate (CB7630) in patients with prostate cancer”, Filed for case IPR2016-01332, Janssen Exhibit 2030, on May 12, 2003, 4 pages. cited by applicant

Clinical trial patient data extract, for Patient 1, Patient 2, 2 pages. cited by applicant

ClinicalTrials.gov Archive, NCT00485303 on Jun. 11, 2007.
<http://clinicaltrials.gov/archive/NCT00485303/>, 3 pages. cited by applicant

ClinicalTrials.gov, “Phase II Clinical Trial of Abiraterone Acetate Without Exogenous Glucocorticoids in Men With Castration-resistant Prostate Cancer With Correlative Assessment of Hormone Intermediates.”, NCT02025010, *Amerigen v. Janssen*, Case# IPR2016-00286. 5 pages. cited by applicant

Cole, “Cancer Expert Doubts Claims About Prostate Cancer Trial,” *BMJ*, vol. 337: a979 (2008). cited by applicant

Coligar Biotechnology, Cougar Biotechnology presents CB7630 Phase I clinical data at the 2005 Prostate Cancer Symposium, All Business, 2005. cited by applicant

Collins et al., “A systematic review of the effectiveness of docetaxel and mitoxantrone for the treatment of metastatic hormone-refractory prostate cancer,” *British J. of Cancer*. 95, pp. 457-462

(2006). cited by applicant

Colston et al, "Mechanisms implicated In the growth regulatory effects of vitamin D In breast cancer," *Endocrine-Related Cancer*, vol. 9(1):45-59 (2001). cited by applicant

Colston, K. W., "Effects of Seocalcitol (EB1089) on Nitrosomethyl Urea-Induced Rat Mammary Tumors", 80(3) *Breast Cancer*, 303-311 (2003). cited by applicant

Conde and Aronson, "Risk factors for male osteoporosis." *Urologic Oncology, Seminars and Original Investigations*, Filed for Case IPR2016-00286, Janssen Exhibit 2025. on 2003, vol. 21, pp. 380-383. cited by applicant

Consider the impact of Zvtiga (Registered) 500 mg film-coated tablets over the course of a month or a year for your patients with mCRPC, Zvtiga (Registered) (abiraterone acetate) Dosing & Monitoring HCP 7 pages. cited by applicant

Continuing Challenge of Hormone-Refractory Prostate Cancer, Declaration of Oliver Sartor, "Clinical Genitourinary Cancer", filed for case Janssen IPR2016-01582, Janssen Exhibit 2156 on Mar. 2006, pp. 238-239. cited by applicant

Cooper et al., "Mechanisms of Disease: Biomarkers and Molecular Targets from Microarray Gene Expression Studies in Prostate Cancer," *Nature Clin Practice Oncol*, vol. 4(12):677-687 (2007). cited by applicant

Costa-Santos, M. et al. "Two Prevalent CYP17 Mutations and Genotype-Phenotype Correlations in 24 Brazilian Patients with 17-Hydroxylase Deficiency," *J. Clin. Endocrin. & Metabol* vol. (89)1, pp. 49-60, 2004. cited by applicant

Cougar Biotechnology Inc. with the U.S. Securities and Exchange Commission, Form 10-QSB, 2013. cited by applicant

Cougar Biotechnology, A Phase III Open Label Study of the 17-Hydroxylase/17 α -C17-20 lyase Inhibitor, Abiraterone Acetate in Patients with Prostate Cancer Who have Failed Hormone Therapy, Letter to Johann De-Bono, Re: Revised abiraterone protocol, Dec. 6, 2004, 86 pages. cited by applicant

Cougar Biotechnology, Cougar Biotechnology Announces Initiation of Phase I/II Trial for C87630 (Abiraterone Acetate), Cougar Biotechnology, Dec. 14, 2004. cited by applicant

Cougar Biotechnology, Cougar Biotechnology Announces Presentation of Positive C87630 Clinical Data at ESMO Conference, Drugs.com, Jul. 2007. cited by applicant

Attard et al., Activity, toxicity, and effect on steroid precursor levels of abiraterone (A), an oral irreversible inhibitor of CYP17 (17 α -hydroxylase/17 α -lyase), in castrate men with castration refractory prostate cancer (CRPC): 2007 Prostate Cancer Symposium, Abstract No. 264. cited by applicant

Attard et al., "Dissecting Prostate Carcinogenesis Through ETS Gene Rearrangement Studies: Implications for Anticancer Drug Development," *J. Clin. Pathol*, Vol. 61:891-896 (2008). cited by applicant

Attard et al., "Improving the Outcome of Patients with Castration-Resistant Prostate Cancer Through Rational Drug Development," *British Journal of Cancer*, vol. 95:767-774 (2006). cited by applicant

Attard et al., "Management Strategies for Hormone-Refractory Prostate Cancer", *Therapy in Practice*, Ann. J. 2006, vol. 5, No. 3, pp. 163-169. cited by applicant

Attard et al., "Phase I Clinical Trial of a Selective Inhibitor of CYP17, Abiraterone Acetate, Confirms That Castration-Resistant Prostate Cancer Commonly Remains Hormone Driven." *J. Clin. Oncol.*, vol. 26(26):4563-4571 (2008). cited by applicant

Attard et al., "Phase I study of continuous oral dosing of an irreversible CYP17 inhibitor, abiraterone (A), in castration refractory prostate cancer (CRPC) patients (p) incorporating the evaluation of androgens and steroid metabolites in plasma and tumor," *J. Clin. Oncology*, 2007 ASCO Annual Meeting Proceedings Part I, vol. 25(188), Abstract No. 5063. cited by applicant

Attard et al., "Predictors of Response and Pharmacodynamic (PD) Endpoints in a Phase I and

Pharmacokinetic Study of Abiraterone Acetate (AA) in Castration-Resistant Prostate Cancer (CRPC),” ASCO 2008 Genitourinary Cancers Symposium, abstract No. 214 (2008). cited by applicant

Attard et al., Selective Inhibition of CYP17 with Abiraterone Acetate Is Well Tolerated and Results in a High Response Rate in Castration-Resistant Prostate Cancer (CRPC), *N Molecular Cancer Therapeutics* (December 2008 Val. 6(12) 3455s, 2007 AACR-NCI-EORTC International Conference, Poster Session B, abstract No. B73 (2007). cited by applicant

Attard et al., “Selective Inhibition of CYP17 With Abiraterone Acetate Is Highly Active in the Treatment of Castration-Resistant Prostate Cancer,” *Journal of Clinical Oncology*, Filed for Case IPR2016-002B6, Janssen Exhibit 2015, vol. 27, No. 23, on Aug. 10, 2009, pp. 3742-3748. cited by applicant

Attard et al., Steroid Hormone Receptors in Prostate Cancer: A Hard Habit to Break?, *Cancer Cell*, vol. 16, Dec. 8, 2009, pgs 458-462. cited by applicant

Attard et al., “Selective blockade of androgenic steroid synthesis by novel lyase inhibitors as a therapeutic strategy for treating metastatic prostate cancer,” *BJU International*, vol. 96, pp. 1241-1246 (2005). cited by applicant

Attard et al... Abiraterone, an oral, irreversible CYP450C17 enzyme inhibitor appears to have activity in post-docetaxel castration refractory prostate cancer (CRPC) patients (pts). *Annals of Oncology*, vol. 18(Supplement 9): ix173-ix174. Abstract No. 51PD (2007). cited by applicant

Attard et al... “Abiraterone Acetate Is Well”, *Journal of Clinical Oncology*, vol. 28, No. 29, Oct. 10, 2010. pp. e560-e561. cited by applicant

Attard G et al., “Clinical and biochemical consequences of CYP17A1 inhibition with abiraterone given with and without exogenous glucocorticoids in castrate men with advanced prostate cancer”. *Journal Of Clinical Endocrinol Metab* Feb. 2012, vol. 97 No. 2 pp. 507-516. cited by applicant

Attard, Poster: A randomized trial of abiraterone acetate (AA) administered with 1 of 4 glucocorticoid (GC) regimens in metastatic castration-resistant prostate cancer (mCRPC) patients (pts). *J. Clin Oncol* 34, 2016 (Suppl 2S: abstr 261). pp. 1-4, Nov. 15, 2016. cited by applicant

Auches et al., “Use of Prednisone With Abiraterone Acetate in Metastatic Castration-Resistant Prostate Cancer”, *The Oncologist*, vol. 19, No. 12, Oct. 31, 2014 (Oct. 31, 2014), pp. 1231-1240. cited by applicant

Auchus et al., “Human steroid biosynthesis for the oncologist”, NIH Public Access, *J Investing Med*. Author manuscript: available in PMC May 14, 2013, 22 pages. cited by applicant

Auchus et al., “Use of Prednisone With Abiraterone Acetate in Metastatic Castration-Resistant Prostate Cancer”, *The Oncologist* published online Oct. 31, 2014, 10 pages. cited by applicant

Auchus, R.J. “The genetics, pathophysiology, and management of human deficiencies of P450c17,” *Endocrinology and Metabolism. Clinics of North America*. vol. 30, No. 1, pp. 101-119, Mar. 2001. cited by applicant

Auchus-3, R. J, “The genetics, pathophysiology, and the management of human deficiencies of P450c17”, *Endocrinol Metab Clin North Am* (2001), 30, p. 101-119. cited by applicant

Auerbach et al., United States Patent and Trademark Office, Before the Patent Trial and Appeal Board, *Amerigen Pharmaceuticals Limited and Argentum Pharmaceuticals LLC* (Petitioners), *Janssen Oncology, Inc.* (Patent Owner) for U.S. Pat. No. 8,822,438, “Methods and Compositions for Treating Cancer”, filed for Case IPR2016-01582, Amerigen Exhibit 2019 Issued on Sep. 2, 2014, 13 pages. cited by applicant

Austin and Hux. “A Brief note on overlapping confidence intervals”, *Mylan vs. Janssen*, Filed for Case IPR2016-01332, Janssen Exhibit 2184, on 2002, *Journal of Vascular Surgery*, vol. 36, No. 1, 194-195 pages. cited by applicant

Ayub, M., Inhibition of testicular 17 α -hydroxylase and 17,20-lyase but not 3 β -hydroxysteroid dehydrogenase-isomerase or 17 β -hydroxysteroid oxidoreductase by ketoconazole and other imidazole drugs, *Journal of Steroid Biochemistry* (1987) 28(5), p. 521-531. cited by applicant

Azad et al., "Outcomes with Abiraterone Acetate in Metastatic Castration-resistant Prostate Cancer Patients Who Have Poor Performance Status", *European Urology*, vol. 67, Issue 3, Mar. 2015, pp. 441-447. cited by applicant

B. A. J. Ponder et al., "Response to aminoglutethimide and cortisone acetate in advanced prostatic cancer", *Br. J. Cancer*, (1984) 50 ,pp. 757-763. cited by applicant

Balaji, *Managing Metastatic Prostate Cancer in Your Urological Oncology Practice* (2016). cited by applicant

Barrie et al., "17-(3-Pyridyl) Substituted Irreversible Inhibitors of Cytochrome P45017alpha" *British J. Cancer*, vol. 78(Suppl. 1):34. abstract No. 33 (1998). cited by applicant

Barrie et al., "Biochemistry and Pharmacokinetics of Potent Non-Steroidal Cytochrome P45017Alpha Inhibitors," *J Steroid Biochem. Malec. Biol.*, vol. 60(5-6):347-351 (1997). cited by applicant

Barrie et al., "Biochemistry of Potent Cytochrome P45017alpha Inhibitors," *British J. Cancer*, vol. 75 (Suppl. 1):6, abstract No. 1.7 (1997). cited by applicant

Barrie et al., "CB7598: A Potent Inhibitor of Steroidal 17alpha-Hydroxylase/C17,20 Lyase. A Potential New Drug for the Treatment of Prostate Cancer," *J. Pharmacy and Pharmacology*, vol. 47(128):1076 (1995). cited by applicant

Barrie et al., "Highly Potent Inhibitors of Human Cytochrome P-450(17alpha): Activity In Vitro and In Vivo," *British J. Cancer* (1993), p. 75, BACR/ACP/BOA Annual Meeting, abstract No. 177 (1993). cited by applicant

Barrie et al., "Inhibitors of Cytochrome P450 17alpha (17alpha-Hydroxylase/C17,20 Lyase)," *Endocrine-Related Cancer*, vol. 3:25-39 (1996). cited by applicant

Barrie et al., *Pharmacology of Novel Steroidal Inhibitors of Cytochrome P450 17alpha (17alpha-Hydroxylase/C17-20 Lyase; J, Steroid Biochem Malec, Biol*, vol. 50(5/6)267-273 (1994). cited by applicant

Beardsley et al., "Systemic Therapy After First-Line Docetaxel in Metastatic castration-Resistant Prostate Cancer," *Current Opinion in Supportive and Palliative Care*, vol. 2:161-166 (2008). cited by applicant

Beer et al., "Weekly high-dose calcitriol and docetaxel in metastatic androgen-independent prostate cancer," *J Clin. Oncology*, vol. 21(1) pp. 123-128 (2003). cited by applicant

Bernard P. Schimmer and Keith L. Parker, *Andrenocorticotrophic Hormone: Adrenocortical Steroids and Their Synthetic Analogs: Inhibitors of the Synthesis and Actions of Adrenocortical Hormones*, in *Goodman & Gilman's the Pharmacological Basis of Therapeutics*, (10th ed. 2001), pp. 1649-1677. cited by applicant

Berry, W. et al.. Phase III Study of Mitoxantrone Plus Low Dose Prednisone Versus low Dose Prednisone Alone in Patients with Asymptomatic Hormone Refractory Prostate Cancer, *The Journal of Urology*, 2002, pp. 2439-2443, vol. 168. cited by applicant

Biff F. Palmer, "Managing J Hyperkalemia Caused by Inhibitors of the Renin-Angiotensin-Aldosterone System", *New England Journal of Medicine*, file for Case IPR2016-01332, Janssen Exhibit 2067 on Aug. 5, 2004, vol. 351, No. 6, pp. 585-592. cited by applicant

Biglieri EG et al., "Herron MA, Brust N. 17-hydroxylation deficiency in man", *Journal of Clinical Investigation* 1966, vol. 45 No. 12, pp. 1946-1954. cited by applicant

Biophosphonates Fact Sheet. The Paget Foundation
<http://www.paget.org/information.FactSheet/bisfact/html> (Jul. 2006). cited by applicant

Blutt et al., "A calcitriol analogue, EB 1089, inhibits the growth of LNCaP tumors in nude mice," *Cancer Res.*, vol. 60(4), pp. 779-782 (2000). cited by applicant

Bonnie Gills, "ASCO GU Highlights New Treatment Options for Prostate Cancer", 2016 Highlights from ASH: CLL News, 2015, 5 Pages. cited by applicant

Booth et al., *Oncology's trials*, 'Nature Reviews', vol. 2: Aug. 2003, pp. 609-610. cited by applicant

Borner et al., "Answering Patients Needs", *Oral Alternatives to Intravenous Therapy*, *The Oncol-*

gisil 2001 vol. 6, suppl 4, pp. 12-16. cited by applicant

Boumpas et al., “Glucocorticoid Therapy for Immune-mediated Diseases: Basic and Clinical Correlates,” *Annals of Internal Medicine*, Filed for Case IPR2016-00286, Janssen Exhibit 2021 on Dec. 15, 1993, vol. 119, No. 12 1198-1208. cited by applicant

Brenner et al. ETS fusion positive prostate tumor models repond to olaparib; *Cancer Cell* 19 664; 2011. cited by applicant

Brooke et al., “A novel point mutationin P450c17 (CYP17) causing combined 17alpha-hydroxylase/17,20.lyase deficiency”,. *The Journal of Clinical Endocrinology & Metabolism*, 2006, vol. 91 No. 6 pp. 2428-2431. cited by applicant

Bruno et al., *Targeting cytochrome P450 enzymes: A new approach in anti-cancer drug development Elsevier*, 2007, pp. 5047-5060, vol. 15. cited by applicant

Bubley et al., “Eligibility and Response Guidelines for Phase II Clinical Trials in Androgen-Independent Prostate Cancer: Recommendations From the Prostate-Specific Aniigen Working Gmup”, *Journal of Clinical Oncology*, file for Case IPR2016-00286, Janssen Exhibit 2057, on Nov. 1999, vol. 17, No. 11, pp. 3461-3467. cited by applicant

Burgess and Roth et al., “Changing Perspectives of the Role of Chemotherapy in Advanced Prostate Cancer,” *urologic Clinics of North America* 2006, filed for case IPR2016-00286, Janssen Exhibit 2007, Vol 33, pp. 227-236. cited by applicant

“Amended marketing authorization for Zytiga”, European Commission, Nov. 9, 2016, pp. 74. cited by applicant

“Clinical Trials: What you need to know”, American cancer society, Aug. 18, 2020, pp. 1-29. cited by applicant

“Full Prescribing Information and Patient Information for Zejula”, Mar. 2017, pp. 1-19. cited by applicant

“Marketing authorization for Zytiga”, European Commission, Sep. 5, 2011, pp. 31. cited by applicant

“Monetizing Focus: Tesaro Adds Runway By Out-licensing Niraparib In Prostate Cancer”, *Pink sheet*, Apr. 6, 2016, pp. 1-3. cited by applicant

“Tesaro Announces Global Prostate Cancer Collaboration and Licensing Agreement With Janssen”, Apr. 6, 2016, pp. 1-2. cited by applicant

Boudadi et al., “Resistance to Novel Antiandrogen Therapies in Metastatic Castration-Resistant Prostate Cancer”, *Clin Med Insights Oncol*, Feb. 9, 2016, vol. 10, (Suppl 1), pp. 1-9. cited by applicant

Palma et al., “ABT-888 Confers Broad In vivo Activity in Combination with Temozolomide in Diverse Tumors”, *Cancer Therapy: Preclinical*, Dec. 1, 2009, pp. 7277-7290. cited by applicant

Raritan, NJ, “Janssen Enters Worldwide Collaboration and License Agreement with Tesaro, Inc., for Niraparib in Prostate Cancer”, Apr. 6, 2016, pp. 2. cited by applicant

Hussain et al., “Co-targeting androgen receptor (AR) and DNA repair: A randomized ETS gene fusion-stratified trial of abiraterone + prednisone (Abi) +/- the PARP1 inhibitor veliparib for metastatic castration-resistant prostate cancer (mCRPC) patients (pts) (NCI9012)—A University of Chicago phase II consortium trial”, *Journal of Clinical Oncology*, vol. 34, No. 15, May 20, 2016. cited by applicant

Study NCT02924766—Submitted Date: Oct. 4, 2016 (v1). cited by applicant

Stamey et al., “Prostate-Specific Antigen As A Serum Marker For Adenocarcinoma Of The Prostate,” *Tile New England journal of Medicine*, vol. 317, No. 15, Oct. 8, 1987 pp. 909-916. cited by applicant

Stein, et al., *Randomized Phase 2 Therapeutic Equivalence Study of Abiralerone Acetate Fine Particle Formulation vs. Originator Abiraterone Acetate In Patients With Metastatic Castration Resistant Prostate Cancer: The STAAR Study*, *Urologic Oncology: Seminars and Original Investigations*. 36, 2018, 81.e9-81.e16. cited by applicant

Sternberg, "Systematic Chemotherapy and New Experimental Approaches in the Treatment of Metastatic Prostate Cancer," *Annals of Oncol.*, vol. 19 (Supp. 7):vii91-vii95 (2008). cited by applicant

Sternberg. "Hormone refractory metastatic prostate cancer", *Annals of Oncology*, file for Case IPR2016-00286, Janssen Exhibit 2056, on 1992, vol. 3, pp. 331-335. cited by applicant

Storlie. J.A., et al., "Prostate Specific Antigen Levels and Clinical Response to Low Dose Dexamethasone for Hormone-Refractory Metastatic Prostate Carcinoma", *Cancer* (1995) vol. 76, No. 1, p. 96-100. cited by applicant

Strother et al., "Novel cytotoxic and biological agents for prostate cancer" Where will the money be in 2005?, *European Journal of Cancer* 2005, filed for Case IPR2.016-00286, Janssen Exhibit 2008. vol. 41, pp. 954-964. cited by applicant

Summary of Product Characteristics for Zytiga 250mg tablets (Jan. 16, 2014). cited by applicant

Swartz and Dluhy, "Corticosteroids: Clinical Pharmacology and Therapeutic Use", *Clinical Pharmacology and Therapeutic, Drugs*, file for Case IPR2016-00286, Janssen Exhibit 2068, on 1978, vol. 16, pp. 238-255. cited by applicant

Sweeney et al., "Chemohomonal Therapy in Metastatic Hormone-Sensitive Prostate Cancer", *New England Journal of Medicine*, filed for Case *Wockhardt vs. Janssen* IPR2016-01582, Janssen Exhibit 2163, on Aug. 20, 2015, vol. 373, No. 8, pp. 737-746. cited by applicant

Szmulewitz et al., "Antiandrogen Therapy in Prostate Cancer," *Update on Cancer Therapeutics*, vol. 2:119-131 (2007). cited by applicant

T. Saika et al., Treatment of androgen-independent prostate cancer with dexamethasone: A prospective study in stage D2 patients, *International Journal of Urology*, 8, pp. 290-294 (2001). cited by applicant

Taichman et al., "The evolving biology and treatment of prostate cancer", *The Journal of Clinical Investigation*, Sep. 2007, vol. 117, No. 9, 2351-2361. cited by applicant

Takeda et al., "Inhibitors of the Key Enzymes of Androgen Synthesis: Potential Agents as Targets for Prostate Cancer," *Japanese Journal of Clinical Medicine*, vol. 58(Suppl.):312-316 (2000); Partial Translation Included (Total pages 14). cited by applicant

Tamic, R, et al., "Hormonal Effects of High Dose Medroxyprogesterone Acetate Treatment in Males with Renal or Prostatic Adenocarcinoma", (1988), vol. 22 (1), Abstract. cited by applicant

Tanagho, EA, et al., "The Leading Single-Volume Resource in Urology" *Smith's General Urology*, 16th Edition, (2004), Chapter 19, pp. 321-323; Chapter 22, pp. 380-385. cited by applicant

Tannock et al., Chemotherapy with mitoxantrone plus prednisone or prednisone alone for symptomatic hormone-resistant prostate cancer: a Canadian randomized trial with palliative end points, *Journal of Clinical Oncology*,. vol. 14,. (6):1756-1764 (1996). cited by applicant

Tannock et al., "Treatment of metastatic prostatic cancer with low-dose prednisone: evaluation of pain and quality of life as pragmatic indices of response," *Journal of Clinical Oncology*, vol. 7 590-597 (1989). cited by applicant

Tannock, IF, et al.. Docetaxel plus prednisone or mitoxantrone plus prednisone for advanced prostate cancer, *The New England Journal of Medicine*, 2004, pp. 1502-1512, vol. 351(15). cited by applicant

Taplin, "Drug Insight Role of the Androgen Receptor in the Development and Progression of Prostate Cancer," *Nature Clin. Practice Oncol*, vol. 4(4):236-244 (2007). cited by applicant

Taxotere (docetaxel) Injection Concentrate, Aventis Pharmaceuticals Inc. (2005). prescribing information. cited by applicant

Taxotere label, May 2004, Mylan Pharms. Inc. Exhibit 1140. cited by applicant

Taxotere prescribing information (Dec. 2015), Taxotere (docetaxel) Injection Concentrate, Intravenous Infusion (IV). Initial U.S. Approval: 1996, 63 pages. cited by applicant

Taxotere(Registered),"RX Taxotere (Docetaxel) Injecton Concentrate", InformatIn as of May 2004, 35 pages. cited by applicant

Tewari et al., “Long-Term Survival Probability in Men with Clinically Localized Prostate Cancer Treated Either Conservatively or with Definitive Treatment (Radiotherapy or Radical Prostatectomy)”, *Urology*, Dec. 2006, vol. 68. Issue 6, 1268-1274. cited by applicant

The Institute of Cancer Research, Abiraterone: a story of scientific innovation and commercial partnership, Making the discoveries that defeat cancer filed for IPR2016-00286 Oct. 3, 2016. cited by applicant

Theodore, “Cancers Genito-Urinaires,” *Oncologie*, vol. 10:497-500 (2008) (partial English translation enclosed). cited by applicant

Therasse et al., “New Guidelines to Evaluate the Response to Treatment in Solid Tumors *Journal of the National Cancer Institute*”, vol. 92, No. 3, Feb. 2, 2000 pp. 205-216. cited by applicant

Thirty Second Edition Physicians' Desk Reference, 1978, PDR, 6 pages. cited by applicant

Toshihiko Yanase et al., “Deletion of a phenylalanine in the N-terminal region of human cytochrome P-450(17 alpha) results in partial combined 17 alpha-hydroxylase/17, 20-lyase deficiency” *Journal of Biological Chemistry* May 24, 1989, vol. 264 No. 30 pages 18076-18082. cited by applicant

Trelstar (Trademark) Depot 3.75 mg (triptorelin pamoate for injectable suspension), Pharmacia & Upjohn Company (2001), package insert. cited by applicant

Trelstar(Trademark) LA 11.25 mg (triptorelin pamoate for injectable suspension), Pharmacia & Upjohn Company (2001), package insert. cited by applicant

Trepanier, “Glucocorticoids”, *Cliniciansbrief.com*, Dec. 2005 5 pages. cited by applicant

Trial of Abiraterone Without Exogenous Glucocorticoids in Men With CRPC With Correlative Assessment of Hormone Intermediates, 10 pages. cited by applicant

Trump et al., “Phase II trial of high-dose, intermittent calcitriol (1,25 dihydroxyvitamin D3) and dexamethasone in androgen-independent prostate cancer,” *Cancer*, vol. 106(10), pp. 2136-2142 (2006). cited by applicant

Truven Commercial and Medicare Data, *Wockhardt vs. Janssen*, Filed for case IPR2016-01332, Janssen Exhibit 2135, 12 pages. cited by applicant

Tucker et al., “Reversible Adrenal Insufficiency Induced by Ketoconazole” Janssen Exhibit 2090, *Wockhardt vs. Janssen*, Case# IPR2016-01582, *JAMA*, vol. 253, No. 16 Apr. 26, 1985, 2 Pages. cited by applicant

Twenty Ninth Edition Physicians' Desk Reference, 1975, PDR, 4 pages. cited by applicant

U.S. Patent. Alan H, Auerbach, et al., U.S. Pat. No. 8,822,438 B2, “Methods and Compositions for Treating Cancer”, filed for Case# 15-cv-05909-KM-JBC, Document# 288-1, filed on Feb. 13, 2017, p. 2 of 13. cited by applicant

U.S. Food and Drug Administration (“FDA”), “FDA Approves New Indication for Taxotere-Prostate Cancer”, FDA News Release dated May 19, 2004, 2 pages. cited by applicant

U.S. Food and Drug Administration FDA News Release, “FDA expands Zytiga's use for late-stage prostate cancer”, Dec. 10, 2012
<http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm331492.htm> (access Jun. 30, 2016), 2 pages. cited by applicant

U.S. Food and Drug Administration, “FDA News Release”, Press Announcements> FDA approves Zytiga for late-stage prostate cancer, Janssen Exhibit 2070, *Amerigen v. Janssen*, for Case# IPR2016-00286, Apr. 28, 2011, 2 Pages. cited by applicant

U.S. Food and Drug Administration, “Orange Book: Approved Drug Products with Therapeutic Equivalence Evaluations” *Wockhardt v. Janssen* IPR2016-00286 Janssen Exhibit 2107 Oct. 3, 2016. cited by applicant

U.S. Food and Drug Administration, FDA Website Drugs@FDA-Zytiga,
<http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.DrugDetails> (accessed Jun. 28, 2016), Mylan Pharms, Inc., Exhibit 1046, 2 pages. cited by applicant

UBS Investment Research, “Johnson & Johnson—Zytiga Label Extended,” Dec. 10, 2012. cited by

applicant

UBS Research, “Medivation—A look at the Growth and Share in Prostate Cancer,” Feb. 3, 2014. cited by applicant

Understanding the role of prednisone In combination with Zytiga(Registered), (abiraterone acetate), Putting Prednisone in Perspective, 13 pages. cited by applicant

Understanding Zytiga Users “Urologist Success” Qualitative Research, Janssen Exhibit 2092, *Amerigen* vs. *Janssen* filed for Case# IPR2016-00286, Jan. 2014. cited by applicant

Veliparib (ABT-888) is a PARP inhibitor being investigated to treat non-small cell hmg cancer, BRCA breast cancer and ovarian cancer, 2019, <https://www.abbvie.com/onr-science/pipeline.html>. cited by applicant

Venkitaraman, R., et al., “Efficacy of low-Dose Dexamethasone in Castration-Refractory Prostate Cancer”, *BJU Int* (2008), 101, pp. 1756-1764. cited by applicant

Viadur(Registered) (leuprolide acetate implant). Bayer Pharmaceuticals Corporation (2004), package insert. cited by applicant

Janssen Announces Preliminary Results from Phase 2 GALAHAD Study in Adults with Metastatic Castration-Resistant Prostate Cancer and DNA-Repair Pathway Defects (DRD), Feb. 14, 2019, 8 pages. cited by applicant

Janssen Announces U.S. FDA Breakthrough Therapy Designation Granted for Niraparib for the Treatment of Metastatic Castration-Resistant Prostate Cancer, Oct. 3, 2019, 4 pages. cited by applicant

Smith et al., “Niraparib in patients with metastatic castration-resistant prostate cancer and DNA repair gene defects (GALAHAD): a multicentre, open-label, phase 2 trial”, *Lancet Oncol.*, Mar. 2022, 23(3): 362-373. cited by applicant

Vidal et al., “Reversing Resisiance to Targeted Therapy”, vol. 16—Supplement No. 4, 2004, pp. 7-12. cited by applicant

Vijayakumar, S. et al., “Clinical Trials Involving Vitamin D Analogs in Prostate Cancer”, *Cancer J.* vol. 11(5): 362-73 (2005). cited by applicant

Vink-Van Wijngaarden et al., “Inhibition of breast cancer cell growth by cornbined treatment with vitamin D3 analogues and tamoxifen,” *Cancer Research*, American Association for Cancer Research, Baltimore, MD:5711-5717, abstract (1994). cited by applicant

Vitamin D Aids Chemotherapy for Advanced Prostate Cancer, <http://www.supplementquality.com/efficacy/VitD.prostatechemo.html> (2002). cited by applicant

Vitamin D Boosts Cancer Treatment, <http://new-s.bbc.co.uk/1/hi/health/2961806.stm> (2003). cited by applicant

Vivek K. Arora et al., Glucocorticoid Receptor Confers Resistance to Anti-Androgens by Bypassing Androgen Receptor Blockade, *Cell*, Dec. 5, 2013: 155(6): 1309-1322. cited by applicant

Vogelzang, N.J., Curriculum Vitae, 15 pages. cited by applicant

W. H.J. Kruit et al.. Effect of combination therapy with aminoglutethimide and hydrocortisnne on prostate-specific antigen response in metastatic prostate cancer refractory to standard endocrine therapy, *Anti-Cancer Drugs*, 2004, vol. 15, No. 9, pp. 843-847. cited by applicant

Walsh et al., “Docetaxel Plus Prednisone or Mitoxantrone Plus Prednisone for Advanced Prostate Cancer,” *J Urology*, vol. 173 :2 p. 456 (2005). cited by applicant

Wang, C., et al., “Hypertension due to 17a-Hydroxylase deficiency”, *Australian and New Zealand Journal of Medicine* (1978), 8(3), p. 295-299. cited by applicant

Wedbush Quick Note, “Medivation: Zytiga Market Share Decline Accelerates From Last Quarter,” Jul. 14, 2015. cited by applicant

Wells Fargo Sewrities, LLC., Equity Research, “Johnson & Johnson,” Jun. 29, 2015. cited by applicant

What You Need to Know About Prostate Cancer, NIH Publication No. 12-1576 (2012) Janssen Exhibit 2091 *Amerigen* vs. *Jans-sen* filed for Case# IPR2016-00286. cited by applicant

White, P.C., "Synthesis and Metabolism of Corticosteroids," Principles and Practice of Endocrinology and Metabolism, Ed. Kenneth L Becker, Philadelphia: Lippincott Williams & Wilkins, Janssen Exhibit 2086 *Amerigen* vs. *Janssen* for Case# IPR2016-00286, 2001, Chapter 72, 704-714. cited by applicant

Whitworth, "Mechanism of glucocorticoid-induced hypertension," *Kidney International*. vol. 31:1213-1224 (1987). cited by applicant

Wikipedia, Corticosteroid, undated. website, 2013. cited by applicant

William Blair, "Biotechnology—Zytiga Fourth-Quarter Sales Imply Xtandi Strength," Jan. 22, 2013. cited by applicant

William Blair, Medivation, Inc. "Looking into Recent Weaknesses: Second-Quarter Preview and Breast Cancer Prospect; Lowering Price Target to \$150 on Adjusting Share Count," Jul. 14, 2015. cited by applicant

William D. Figg et al.. "A Randomized Phase II Trial of Ketoconazole Plus Alendronate Versus Ketoconazole Alone in Patients With Androgen Independent Prostate Cancer and Bone Metastases", the *Journal of Urology*, vol. 173. pp. 790-796, Mar. 2005. cited by applicant

Williams, "Discontinued Drugs in 2007: oncology drugs," *Expert Opinion on Investigational Drugs*, vol. 17 No. 12: pp. 1791-1816 (2008). cited by applicant

Xiao-Yan Zhao et al., Glucocorticoids can promote androgen-independent growth of prostate cancer cells through a mutated androgen receptor, *Nat Med* vol. 6 No. 6, Jun. 2000, pp. 703-706. cited by applicant

Xu et al., Correlation between Prostate-Specific Antigen Kinetics and Overall Survival in Abiraterone Acetate-Treated Castration-Resistant Prostate Cancer Patients, *Clinical Cancer Research*, Jul. 15, 2015, vol. 21, No. 14 pp. 3170-3177. cited by applicant

Yanase T et al., 17 alpha hydroxylase/17-20-lyase deficiency: from clinical investigation to molecular definition, *Endocrine Reviews* 1991; vol. 12 No. 1 pp. 91-108. cited by applicant

Yano et al., "Glucocorticoids Suppress Tumor Angiogenesis and In vivo Growth of Prostate Cancer Cells", *Clinical Cancer Research*, May 15, 2006, vol. 12. No. 10, pp. 3003-3009. cited by applicant

Yano, A., et al., "Glucocorticoids Suppress Tumor Lymphangiogenesis of Prostate Cancer Cells", *Clin Cancer Res* (2006), vol. 12, pp. 6012-6017. cited by applicant

Yap et al., "Abiraterone acetate, an oral irreversible inhibitor of CYP450C17, administered to castration refractory prostate cancer patients is safe, suppresses androgen and steroid precursor levels, and has a high degree of durable antitumor activity," *J Urology*, vol. 177(4) p. 199 (2007). cited by applicant

Yap et al., "Targeting CYP17; Established and Novel Approaches in Prostate Cancer" *Current Opinion in Pharmacology* (Jul. 28, 2008), vol. 8:449-457 (2008). cited by applicant

Z. Zaferiou et al . . . Managing Metastatic Castration-Resistant Prostate Cancer in the Pre-chemotherapy Setting: A Changing Approach in the Era of New Targeted Agents, *Drugs*, 76:421-430, Feb. 12, 2016. cited by applicant

Zafeiris Zaferiou et al.. in Balaji, Chapter 9, Abiraterone for the Treatment of mCRPC, *Amerigen* v. *Janssen*, Case# IPR2016-00286, pp. 125-155. cited by applicant

Zhang et al, parp inhibitor treatment of neuroendocrine prostate cancer, *Clin Cancer Res* 24: 696, 2017. cited by applicant

Zhang, "Poly(ADP-ribos) polymerase inhibitor: an evolving paradigm in the treatment of prostate cancer", *Asian J. of Andrology*, vol. 16, pp. 401-406, 2014. cited by applicant

Zhou et al , *Asian Pacific Journal of Cancer Prevention*, 2014; 15:1313-1320. cited by applicant

Zoladex (Registered) 10.8 mg 3-Month (goserelin acetate implant), AstraZeneca (2004) package insert. cited by applicant

Zoladex (Registered) 3.6 mg (goserelin acetate implant), AstraZeneca (2004), package Insert. cited by applicant

Zytiga Brochure, Putting Prednisone in Perspective, “Prednisone reduces the incidence and severity of mineralocorticoid-related adverse reactions associated with Zytiga(Registered)”, Janssen Biotech, Inc. Mar. 15, 2015. cited by applicant

Zytiga Label May 20, 2015, Zytiga(Registered) (abiraterone acetate) Tablets For Oral Administration Initial U.S. Approval: 2011, May 2015, Mylan Pharms. Inc., Exhibit 1065, 30 pages. cited by applicant

Zytiga Presentation—Key Clinical Findings for Patients with mCRPC who have progressed on Androgen Deprivation Therapy. cited by applicant

Zytiga Promotional Brochure—A comparison of The Mechanisms of Action of Select Prostate Cancer Treatments. cited by applicant

Zytiga Promotional Brochure—for patients with mCRPC who received prior Chemotherapy containing Docetaxel. cited by applicant

Zytiga Promotional Brochure—in Men with mCRPC—Mechanism of Action. cited by applicant

Zytiga Promotional Brochure—Prednisone reduces the incidence and severity of mineralocorticoid-related adverse reactions with Zytiga. cited by applicant

Zytiga Promotional Brochure—Zytiga (abiraterone acetate)—Introducing a New Option for Patients with mCRPC before Chemotherapy. cited by applicant

Zytiga Promotional Brochure—Ztiga abiraterone acetate—an Oral Androgen Biosynthesis Inhibitor. cited by applicant

Zytiga Usage—prednisone information Janssen Exhibit 2095, *Amerigen vs. Janssen* filed for Case# IPR2016-00286. cited by applicant

Zytiga Usage—total promotional spend Janssen Exhibit 2096 *Amerigen vs. Janssen* filed for Case# IPR2016-00286, 3 Pages. cited by applicant

Zytiga Website, About Zytiga(Registered) (abiraterone acetate), “Prescribed Oral Medication for Metastatic Castration-Resistant Prostate Cancer”, <https://www.zytiga.com/choosing-zytiga/results-of-zytiga> (accessed Apr. 3, 2017), 10 pages. cited by applicant

Zytiga Website How Zytiga(Registered) (abiraterone acetate) Works, “Zytiga(Registered) Inhibits Androgen Production at 3 Sources—Including the Tumor Itself” <https://www.zytiga.com/print/about-zytiga/how-zytiga-works> (accessed Jun. 28, 2016). cited by applicant

Zytiga, “Orange Book: Approved Drug Products with Therapeutic Equivalence Evaluations”, Amerigen 1035, Sep. 24, 2015, 2 pages. cited by applicant

Zytiga (Trademark) (abiraterone acetate) Tablets For Oral Administration Initial U.S. Approval Issued Apr. 2011. *Wockhardt vs. Janssen*, Filed for Case IPR2016-01582, Janssen Exhibit 2168, 22 pages. cited by applicant

Hotte and Saad, “Current management of castrate-resistant prostate cancer,” *Current Oncology*—Volume 17, Supplement 2, S72-S79 2010 filed for Case# IPR2016-00286. cited by applicant

How Zytiga(Registered) (abiraterone acetate) Zytiga (Registered) Inhibits Androgen Production at 3 Sources—Including the Tumor Itself Works, <https://www.zytiga.com/print/about-zytiga/how-zytiga-works> (accessed 7/23/2015). cited by applicant

Hsieh et al., “Novel Concepts in Androgen Receptor Blockade,” *The Cancer Journal* (Jan./Feb. 2008), vol. 14(1):11-14. cited by applicant

Huggins, Charles, et al. *Studies on Prostatic Cancer. I. The Effect of Castration. of Estrogen and of Androgen Injection on Serum Phosphatases in Metastatic Carcinoma of the Prostate*, Cancer Research, 1941, pp. 293-297, vol. 1. cited by applicant

Huggins, C., et al. “Studies on Prostatic Cancer. I. The Effect of Castration, of Estrogen and Androgen Injection on Serum Phosphatases in Metastatic Carcinoma of the Prostate”. *Cancer Res.* 22(4), (1972) pp. 232-240. cited by applicant

In the United States District Court, for the District of New Jersey, BTG International Limited, Janssen Biotech, Inc., Janssen Oncology, Inc., Janssen Research & Development, LLC, for Case#

15-cv 05909-KM-JBC, "Plaintiffs' Opening Claim Construction Brief" Document# 209 Filed 06/13/16 p. 1 of 15. cited by applicant

In the United States District Court, for the District of New Jersey, *BTG International United, Janssen Biotech, Inc., Janssen Oncology, Inc., Janssen Research & Development, LLC*, (Plaintiffs) vs. *Actavis Laboratories FL Inc*, Case# 15-cv-05909-KM-JBC, "Plaintiffs' Responsive Claim Construction Brief" Document 220 Filed Aug. 31, 2016 p. 1 of 25. cited by applicant

Information concerning Zytiga (abiraterone acetate) from

<http://www.kampendium.ch/prod/pnr/1183238lde?Platfonn=Desktop> as of Mar. 25, 2014. cited by applicant

Inspira (eplerenone) tablets, Initial U.S. Approval: 2002, *Wockhardt v. Janssen* filed for Case# IPR2016-01332, Janssen Exhibit 2129, 8 Pages. cited by applicant

International Search Report dated Jul. 13, 2018; International Application No.

PCT/US2018/026661 (PRD3446WOPCT). cited by applicant

International Search Report dated Sep. 29, 2017; International Application No.

PCT/US2017/044413 (PRD3423WOPCT). cited by applicant

Isaacs et al., Identification of ABR-215050 as lead second generation quinoline-3-carboxamide anti-angiogenic agent for the treatment of prostate cancer, *The Prostate*. Dec. 2006, vol. 66, No. 16, pp. 1768-1778. cited by applicant

J J. Body, "Low-dose prednisone and increased risk of development of bone metastases", *Annals of Oncology*, file for Case IPR2016-00286, Janssen Exhibit 2061, on 1996, vol. 7, pp. 643-645. cited by applicant

J. Gonzaalbez et al., "Establishment of reference values for standard dose short synacthen test (250 microgram). low dose short synacthen test (1 microgram) and insulin tolerance test for assessment of the hypothalamo-pituitary-adrenal axis in normal subjects", *Clinical Endocrinology* 2000, vol. 53, 199-204. cited by applicant

J. Trachtenberg and A. Pont, *The Lancet*, Ketoconazole Therapy for Advanced Prostate Cancer, Aug. 25, 1984, pp. 433-435. cited by applicant

Jakobsen et al., "Medroxyprogesterone Acetate and Prednisone in Advanced Breast Cancer. A Randomized Trial," *Eur. J Cancer Clio. Oncol.* vol. 22:9. pp. 1067-1072 (1986). cited by applicant
Janssen Exhibit 2006, *Amerigen v. Janssen* IPR2016-00286, *Clinical Cancer Research*, A Journal of the American Association for Cancer Research, Letter to Ian Judson. Re: "Manuscript# 030579, Hormonal impact of the 17 α -hydroxylase/C17, 20-lyase inhibitor abiraterone acetate (CB7630) in patients with prostate cancer", Dated May 12, 2003, 4 pages. cited by applicant

Janssen research & Development, LLC "A Safety and Pharmacokinetics Study of Niraparib Plus an Androgen Receptor—Targeted Therapy in Men With Metastatic Castration-Resistant Prostate Cancer (Bedivere)", *ClinicalTrials.gov*, Oct. 5, 2016 (Oct. 5, 2016). cited by applicant

Janssen Research & Development. Clinical Study Report Synopsis Protocol JNJ-212082-JPN-201; Phase 2, 9 pages. cited by applicant

Jarman et al., "The Mechanism of Irreversible Inhibition of Human Cytochrome P45017 α by Abiraterone, a Potential New Drug for the Treatment of Prostate Cancer:" *Annals of Oncol.* vol. 9(Suppl 2):135, 10th NCL-EORTC Symposium on New Drugs in Cancer Therapy, abstract No. 516 (1998). cited by applicant

Jarman et al., "The 16.17-Double Bond is Needed for Irreversible Inhibition of Human Cytochrome P45017 α by Abiraterone (17-(3-Methoxy-4-oxo-5 α -androstano-16-en-3 β -ol) and Related Steroidal Inhibitors," *J. Med. Chem.* vol. 41:5375-5381 (1998). cited by applicant

Jarman et al., "Enzyme Inhibitors In Endocrinology," *J. Endocrinology*, vol. 148 (Suppl.), abstract No. S23 (1996). cited by applicant

Jemal, A., et al., "Cancer Statistics, 2007", *CA Cancer J Clin*, 57, (2007) pp. 43-66. cited by applicant

Jennifer Craft, "Eplerenone (Inspira), a new aldosterone antagonist for the treatment of systemic

hypertension and heart failure”, Baylor University Medical Center Proceedings, file for Case IPR2016-00286, Janssen Exhibit 2062, on Apr. 2004, vol. 17, No. 2, pp. 217-220. cited by applicant

Jevtana (Registered) (cabazitaxel) injection, for intravenous use Initial U.S. Approval: 2010, Jevtana Label, Sep. 2016, *Amerigen vs. Janssen* filed for case IPR2016-00286, Amerigeo Exhibit 1149, 25 pages. cited by applicant

Jevtana prescribing information (Sep. 2016), Jevtana(Registered) (cabazitaxel) injection, for intravenous use Initial U.S. Approval 2010, 25 pages. cited by applicant

Jevtana Website. “Dosing and Administration”, <http://www.jevtana.com/hcp/dosing/default.aspx> (accessed Jun. 28, 2016), Mylan Pharms, Inc., Exhibit 1049, 4 pages. cited by applicant

Jhun et al., “Gene expression signature of Gleason score is associated with prostate cancer outcomes in a radical prostatectomy cohort”, *Oncotarget*, Jun. 2017, vol. 8, No. 26, pp. 43035-43047. cited by applicant

John S, et al., “Secondary Hormonal Therapy for Advanced Prostate Cancer”, *The Journal of Urology*, vol. 175, Jan. 2006, pp. 27.34. cited by applicant

Johnson & Johnson Announces Definitive Agreement to Acquire Cougar Biotechnology, Inc., Access to Late-Stage, First-in-Class Prostate Cancer Treatment Strengthens Presence in Oncology May 21, 2009, pp 4. cited by applicant

Johnson & Johnson, “Zytiga Approved in the EU for Use in the Treatment of Metastatic Castration-Resistant Prostate Cancer Before Chemotherapy,” Jan. 11, 2013, 4 pages. cited by applicant

Jones P, Wilcoxon K, Rowley M, Toniatti C. Niraparib: A Poly(ADP-ribose) Polymerase (PARP) Inhibitor for the Treatment of Tumors with Defective Homologous Recombination. *J Med Chem.* Apr. 23, 2015;58(8):3302-3314. cited by applicant

Jubelirer and Hogan, “High Dose Ketoconazole For The Treatment of Hormone Refractory Metastatic Prostate Carcinoma; 16 Cases And Review Of The literature,” *The Journal of Urology*, Filed for Case IPR2016-00286, Janssen Exhibit 2018, on Jul. 1989, vol. 142, No. 1. pp. 89-91. cited by applicant

Julian Seifter et al., “Concepts in Medical Physiology”, Lippincott Williams and Wilkins, *Wockhardt vs. Janssen*, Files for Case IPR2016-01582, Janssen Exhibit 2171, 540-620 pages. cited by applicant

Juliet Richards et al., Interactions of Abiraterone, Eplerenone, and Prednisolone with Wild-type and Mutant Androgen Receptor: A Rationale for Increasing Abiraterone Exposure or Combining with MDV3100, *Cancer Res*; 72(9) pp. 2176-2182, May 1, 2012. cited by applicant

K, Akakura et al., Possible Mechanism of Dexamethasone Therapy for Prostate Cancer: Suppression of Circulating Level of Inledeukin-6, *The Prostate*, 56:106.Math.109 (2003). cited by applicant

K. Kobayashi et al., “Mineralocorticoid Insufficiency Due to Suramin Therapy, Cancer”, vol. 78, No. 11 (Dec. 1, 1996). cited by applicant

K. Nishimura et al., Low Doses of Oral Dexamethasone for Hormone-Refractory Prostate Carcinoma, *Cancer* vol. 89, No. 12, at 2570-2576 (Dec. 15, 2000). cited by applicant

Kaye et al., “New Drug Treatment for Cancer in 2007 Real Progress at Last?” *EJC Supplements*, vol. 5(4):35, 14th European Cancer Conference, abstract No. 126 (2007). cited by applicant

Kirby M. et al., “Characterising the castration-resistant prostate cancer population: A systematic review”, *Int'l J. Clinical Practice*, vol. 65 No. (11): pp. 1180-1192, (Nov. 2011). cited by applicant

Kissmeyer, A.M. et al., “The Tissue-specific Distribution of 3H-Seocalcitol (EB 1089) and 3H-calcitriol in Rats”, *J. of Steroid Bioch. & Mol. Biol.* 89-90: 43-47 (2004). cited by applicant

Knol et al., “The Mis-use of Overlap of Confidence Intervals to Assess Effect Modification”, Filed for Case *Wockhardt V. Janssen* IPR2016-01332, Janssen Exhibit 2183, on 2011, *Eur J. Epidemiol.*, vol. 26, 253-254 pages. cited by applicant

Koshizuka et al., "Combined effect of vitamin D3 analogs and paclitaxel on the growth of MCF-7 breast cancer cells in vivo," Breast Cancer Research and Treatment, Springer, New York, NY, vol. 53(2):.Math.113-120 (1999). cited by applicant

Krishnan et al., "A Glucocorticoid-Responsive Mutant Androgen Receptor Exhibits Unique Ligand Specificity Therapeutic Implications for Androgen-Independent Prostate Cancer," Endocrinology, Filed for Case IPR2016-00286, Janssen Exhibit 2024, on May 2002, vol. 143, No. 5, pp. 1889-1900. cited by applicant

Kuzel et al., "A Phase II Study of Continuous Infusion 5-Fluorouracil in Advanced Hormone Refractory Prostate Cancer", Cancer, file for Case IPR2016-00286, Janssen Exhibit 2054, vol. 72, No. 6, on Sep. 15, 1993, 1965-1968 pages. cited by applicant

I. A. Luthy et al., "Androgenic Activity of Synthetic Progestins and Spironoladone In Androgen-Sensitive Mouse Mammary Carcinoma (Shionogi) Cells in Culture", J. Steroid Biochem, vol. 31, No. 5 .Math.1988. pp. 845-852. cited by applicant

L. Collette et al., Is Prostate-Specific Antigen a Valid Surrogate End Point for Survival in Hormonally Treated Patients With Metastatic Prostate Cancer? Joint Research of the European Organisation for Research and Treatment of Cancer, the Limburgs Universitair Centrum, and AstraZeneca Pharmaceuticals, Journal of Clinical Oncology, vol. 23, No. 25, pp. 6139-6148 (Sep. 1, 2005). cited by applicant

L. Collette et al., Prostate-specific antigen (PSA) alone is not an appropriate surrogate marker of long-term therapeutic benefit in prostate cancer trials, European Journal of Cancer, 42, pp. 1344-1350 (2006). cited by applicant

Lara and Meyers, "Treatment Options in Androgen-Independent Prostate Cancer" Cancer Investigation, file for Case IPR2016-00286, Janssen Exhibit 2053, vol. 17, No. 2, on 1999, pp. 137-144. cited by applicant

Li et al. AR-induced bcraness and part inhibition in preclinical studies; Sci Signal 10: eaam7479, 2017. cited by applicant

Teyssonneau et al., "Prostate Cancer and PARP Inhibitors: Progress and Challenges", J Hematol Oncol, Mar. 29, 2021, vol. 14, No. 51, 19 Pages. cited by applicant

Bundgaard, "Design of Prodrugs", Elsevier, 1985, pp. 1-96. cited by applicant

Declaration of Dr. Gerhardt Attard, 2015, 29 pages. cited by applicant

Kluetz P. G. et al., Abiraterone Acetate in Combination with Prednisone for the Treatment of Patients with Metastatic Castration-Resistant Prostate Cancer: U.S. Food and Drug Administration Drug Approval Summary. Clinical Cancer Research, Dec. 15, 2013, vol. 19, No. 24, pp. 6650-6656. cited by applicant

Lupo et al., "Inhibition of poly(ADP-ribosylation) in cancer: old and new paradigms revisited", Biochim Biophys Acta, Aug. 2014, vol. 1846, No. 1, pp. 201-215. cited by applicant

NCT01715285 Clinicaltrials.gov, History of Changes for Study: NCT01715285 A Study of Abiraterone Acetate Plus Low-Dose Prednisone Plus Androgen Deprivation Therapy (ADT) Versus ADT Alone in Newly Diagnosed Participants With High-Risk, Metastatic Hormone-Naive Prostate Cancer (mHNPC), Sep. 19, 2019, 13 pages. cited by applicant

Papeo et al., "Poly(ADP-ribose) Polymerase Inhibition in Cancer Therapy: Are We Close to Maturity?", Expert Opinion on Therapeutic Patents, Oct. 2009, vol. 19, No. 10, pp. 1377-1400. cited by applicant

Perlmutter et al: "New Directions in Prostate Cancer Management Androgen Deprivation Therapy in the Treatment of Advanced Prostate Cancer", Reviews in Urology, vol. 9, Jan. 2007, pp. S3-S8. cited by applicant

Rydzewska et al., "Adding Abiraterone to Androgen Deprivation Therapy in Men with Metastatic Hormone-sensitive Prostate Cancer: A Systematic Review and Meta-analysis," European Journal of Cancer, vol. 84, 2017, pp. 88-101. cited by applicant

Sandhu et al., "Poly (ADP-ribose) polymerase (PARP) inhibitors for the treatment of advanced

germline BRCA2 mutant prostate cancer”, *Annals of Oncology*, 2013, vol. 24, Issue 5, p. 1416-1418. cited by applicant

Study NCT03012321 “Abiraterone/Prednisone, Olaparib, or Abiraterone/Prednisone + Olaparib in Patients With Metastatic Castration-Resistant Prostate Cancer With DNA Repair Defects”, *Clinical Trials*, NCT03012321, 2021, pp. 1-13, first posted Jan. 6, 2017. cited by applicant

Xu Yuxia et al., “Clinical Observation of Abiraterone Acetate and Prednisone on Patients with Castration resistant Prostate Cancer”, *Cancer Research on Prevention and Treatment*, 2015, vol. 42, No. 4, pp. 382-384. cited by applicant

Zhang et al., “Help-seeking behavior for erectile dysfunction: a clinic-based survey in China,” *Asian J Androl*, 2014, vol. 16, Issue 1, pp. 131-135. cited by applicant

“A Phase 3, Randomized, Double-Blind, Placebo-Controlled Study of Abiraterone Acetate (CB7630) Plus Prednisone in Patients with Metastatic Castration-Resistant Prostate Cancer Who Have Failed Docetaxel-Based Chemotherapy,” *Clinical Genitourinary Cancer*, vol. 6(2):140 (2008). cited by applicant

“CB-7630 Shows Activity in Prostate Cancer Trials,” *Daily Essentials* (2007). cited by applicant

“Cougar Bioedmolgy Announces Positive CB7630 Phase I Data at the AACR Innovations in Prostate Cancer Research Conference,” *News Release*, available at <http://www.cougarbiotechnology.com> (2006). cited by applicant

“Cougar Bioedmolgy Announces Presentation of Positive Phase I and Phase II Data at ASCO Prostate Cancer Symposium,” *News Release*, available at <http://www.cougarbiotechnology.com> (2007). cited by applicant

“Cougar Biotechnology Presents Positive CB7630 Phase I and Phase II Data at ASCO 2008 Genitourinary Cancers Symposium,” *News Release*, available at <http://www.cougarbiotechnology.com> (2008). cited by applicant

“Cougar Biotechnology Announces Presentation of Positive CB7630 Phase I Clinical Data at ASCO 2008 Genitourinary Cancers Symposium,” *News Release*, available at <http://www.cougarbiotechnology.com> (2008). cited by applicant

“Cougar Biotechnology Announces Acceptance of CTA for Abiraterone Acetate,” *News Release*, available at: <http://www.cougarbiotechnology.com> (2005). cited by applicant

“Cougar Biotechnology Announces Initiation of Phase III Trial of CB7630 (Abiraterone Acetate),” *News Release*, available at <http://www.cougarbiotechnology.com> (2008). cited by applicant

“Cougar Biotechnology Announces Initiation of Phase I/II Trial for CB7630 (Abiraterone Acetate).” *News Release*, available at: <http://www.cougarbiotechnology.com> (2005). cited by applicant

“Cougar Biotechnology Announces Initiation of Phase I Trial for CB7630 (Abiraterone Acetate),” *News Release*, available at <http://www.cougarbiotechnology.com> (2006). cited by applicant

“Cougar Biotechnology Announces Initiation of Phase I/II Trial of CB7630 (Abiraterone Acetate) in Advanced Breast Cancer Patients,” *News Release*, available at <http://www.cougarbiotechnology.com> (2008). cited by applicant

“Cougar Biotechnology Announces Positive CB7630 Phase I Data Presented at the National Cancer Research Institute Conference,” *News Release* available at: <http://www.cougarbiotechnology.com> (2006). cited by applicant

“Cougar Biotechnology Announces Presentation of Positive CB7630 (Abiraterone Acetate) Phase II Data at ASCO 2009 Genitourinary Cancers Symposium,” *News Release*, available at <http://www.cougarbiotechnology.com> (2009). cited by applicant

“Cougar Biotechnology Announces Presentation of Positive CB7630 Clinical Data at AACR-NCI-EORTC International Conference on Molecular Targets and Cancer Therapeutics.” *News Release*, available at <http://www.Cougarbiotechnology.com> (2007). cited by applicant

“Cougar Biotechnology Announces Presentation of Positive CB7630 Clinical Data at ASCO Annual Meeting.” *News Release*, available at: <http://www.cougarbiotechnology.com> (2007). cited

by applicant

“Cougar Biotechnology Announces Presentation of Positive CB7630 Clinical Data at AUA Annual Meeting,” News Release, available at <http://www.cougarbiotechnology.com> (2007). cited by applicant

“Cougar Biotechnology Announces Presentation of Positive CB7630 Clinical Data at ESMO Conference,” News Release, available at: <http://www.cougarbiotechnology.com> (2007). cited by applicant

“Cougar Biotechnology Announces CB7630 Phase I Data to be Presented at National Cancer Research Institute Conference,” News Release (Sep. 27, 2006), available at <http://www.cougarbiotechnology.com> (2006). cited by applicant

Cougar Biotechnology Initiates Phase II Trial of CB7630 (Abiraterone Acetate), News Release, available at: <http://www.cougarbiotechnology.com> (2006). cited by applicant

“Cougar Biotechnology Presents CB7630 Phase I Clinical Data at the 2005 Prostate Cancer Symposium,” News Release, available at <http://www.cougarbiotechnology.com> (2005). cited by applicant

“Cougar Biotechnology Presents Positive CB7630 (Abiraterone Acetate) Phase I and Phase II Data at ASCO 2008 Annual Meeting,” News Release, available at <http://www.cougarbiotechnology.com> (2008). cited by applicant

“Cougar Biotechnology Presents Positive CB7630 Clinical Data at AACR Annual Meeting Late-Breaking Clinical Trials Session,” News Release, available at: <http://www.cougarbiotechnology.com> (2007). cited by applicant

“Cougar Biotechnology Presents Positive CB7630 Phase II Data at Chemotherapy Foundation Symposium,” News Release, available at: <http://www.cougarbiotechnology.com> (2007). cited by applicant

“Cougar Biotechnology Presents Positive CB7630 Phase II Data at Prostate Cancer Foundation Scientific Retreat,” News Release, available at <http://www.cougarbiotechnology.com> (2007). cited by applicant

“Cougar Biotechnology Presents Positive CB7630 Phase II Data at EORTC-NCI-AACR Symposium,” News Release, available at: <http://www.cougarbiotechnology.com> (2008). cited by applicant

“Cougar Biotechnology to Present Clinical Data on CB7630 (Abiraterone Acetate) at American Society of Clinical Oncology 2008 Annual Meeting,” News Release, available at <http://www.cougarbiotechnology.com> (2008). cited by applicant

“Cougar Biotechnology Announces Agreement with FDA on Special Protocol Assessment for Phase III Trial of CB7630 (Abiraterone Acetate) in Chemotherapy Naive Castration Resistant Prostate Cancer Patients”. News Release, available at <http://www.cougarbiotechnology.com> (2008). cited by applicant

“Cougar Biotechnology Presents CB7630 Phase I Data at Prostate Cancer Foundation Scientific Retreat.” News Release, available at: <http://www.cougarbiotechnology.com> (2004). cited by applicant

“Final Written Decision”, United States Patent and Trademark Office Before the Patent Trial and Appeal Board, Case No. IPR2016,00286 U.S. Pat. No. 8,822,438 B2, Jan. 17, 2018, 48 pages. cited by applicant

“Final Written Decision”, United States Patent and Trademark Office Before the Patent Trial and Appeal Board, Case No. IPR2016-01332 U.S. Pat. No. 8,822,438 B2, Jan. 17, 2018, 50 page. cited by applicant

“OncoGenex Announces Top-Line Survival Results of Phase 3 Synergy Trial Evaluating Enzalutamide for Metastatic Castrate-Resistant Prostate Cancer”, Janssen Exhibit 2077, *Amerigen vs. Janssen*, filed for Case# IPR2016-00286, Apr. 28, 2014, 2 Page. cited by applicant

“Positive Phase II Data on Cougar Biotechnology's CB7630 Presented at Prostate Cancer

Foundation Scientific Retreat,” News Release, available at <http://www.cougarbiotechnology.com> (2008). cited by applicant

2011 Zytiga(Registered) Approval Prescribing Information, Zytiga(Trademarks), (abiraterone acetate) Tablets For Oral Administration Initial U.S. Approval—2011, Apr. 2011, 22 pages. cited by applicant

Abad L et al., “Male pseudohermaphroditism with 17 alpha-hydroxylase deficiency”, A case report. *Br J Obstet Gynaecol* Dec. 1980, vol. 87 No. 12 pp. 1162-1165. cited by applicant

Abiraterone Acetate: Abbreviated Clinical Study Report Synopsis COU-AA-BE (Doc. EDMS-ERI-13494974:2.0) (BE Synopsis) 5 pages. cited by applicant

About Zytiga (Registered) Abiraterone acetate, retrieved from <https://www.zytiga.com/choosing-zytiga#how-zytiga-works>, on Jul. 25, 2017, 7 pages. cited by applicant

Academy of Managed Care Pharmacy (AMCP) Nexus 2016, National Harbor, MD, USA: Oct. 3-6, 2016 filed for Janssen IPR2016 01332. cited by applicant

Aggarwal et al., “Development of Abiraterone Acetate, a 17-alpha Hydroxylase C17,20-Lyase Inhibitor as a Secondary Hormonal Therapy in Prostate Cancer,” *Update on Cancer Therapeutics*. vol 2:171-175 (2007). cited by applicant

Altman et al., “The Revised Consort Statement for Reporting Randomized Trials: Explanation and Elaboration”, *Annals of Internal Medicine* vol. 134, No. 8, Apr. 17, 2001, pp. 663-694. cited by applicant

Annane D et al., “Effect of treatment with low doses of hydrocortisone and fludrocortisone on mortality in patients with septic shock”, *JAMA*, Aug. 21, 2002, vol. 288, No. 7, pp. 862-871. cited by applicant

Antifungal Treatment Should Be Taken Off the Market, Public Citizen Tells FDA filed for Case# IPR2016-01582 on Feb. 24, 2015, 1 page. cited by applicant

Antonarakis and Eisenberger, Phase III Trials With Docetaxel-Based Combinations for Metastatic Castration-Resistant Prostate Cancer: Time to Learn From Past Experiences *Journal of Clinical Oncology*, vol. 31, No. 14. May 10, 2013: pp. 1709-1712. cited by applicant

Arlt, W. et al., Adrenal insufficiency: *Lancet*, vol. 361 May 31, 2003, pp. 1881-1893. cited by applicant

Armstrong and Carducci. “New drugs in prostate cancer,” *Current Opinions Urology*, 2006, Filed for Case IPR2016-00286, Janssen Exhibit 2011, vol. 16, pp. 138-145. cited by applicant

Armstrong et al., “New Drug Development in Metastatic Prostate Cancer,” *Urologic Oncology: Seminars and Original Investigations*, vol. 26 430-437 (2008). cited by applicant

ASCO Cancer Foundation, Poster Session F: Hormone Refractory, ASCO, 2005. cited by applicant

Asim et al, AR signaling and parp inhibition synergize; olaparib inhibited growth of PC3 prostate tumors; *Nature comm* 8:374; 2017. cited by applicant

Assessment Report for Zytiga (abiraterone) published 2011 by the CHMP of the EMA. cited by applicant

Attard et al., “Antitumor Activity with CYP17 Blockade Indicates That Castration-Resistant Prostate Cancer Frequently Remains Hormone Driven”, *Cancer Res* 2009, vol. 69, No. 12, Jun. 15, 2009, pp. 4937-4940. cited by applicant

Attard et al. “Prostate Cancer's Day in the Sun.” *BMJ*, vol. 337: a1256 (2008). cited by applicant

“Phase II Study to Evaluate Olaparib With Abiraterone in Treating Metastatic Castration Resistant Prostate Cancer”, NCT01972217, First Posted Oct. 30, 2013. cited by applicant

Li et al. targeting parp/cmyb in prostate cancer, with olaparib tested in vivo in Vcap model; *Sci Signal* 7: ra47; 2014. cited by applicant

Ling et al., “17-imidazolyl, pyrazolyl, and isoxazolyl androstene derivatives, Novel steroidal inhibitors of human cytochrome C17 α -lyase (P45017 α):” *J. Med. Chem.*, vol. 40:3297-3304 (1997). cited by applicant

Logothetis et al., “Effect of abiraterone acetate and prednisone compared with placebo and

prednisone on pain control and skeletal-related events in patients with metastatic castration-resistant prostate cancer: exploratory analysis of data from the COU-AA-301 randomised trial”, *Lancet* Oneal, 13, pp. 1201-1217, 2012. cited by applicant

Logothetis et al., “Identification of an androgen withdrawal responsive phenotype in castrate resistant prostate cancer (CRPC) patients (pts) treated with abiraterone acetate (AA),” *J. Clin. Oncol.* (Meeting abstracts), vol. 26 (May 20 Supplement), abstract No. 5017 (2008). cited by applicant

M. Morioka et al., Prostate-Specific Antigen Levels and Prognosis in Patients with Hormone-Refractory Prostate Cancer Treated with Low-Dose Dexamethasone, *Urologia Internationalis* vol. 68, at 10-15 (2002). cited by applicant

Madan Ravi A et al. Abiraterone Cougar Biotechnology Idrugs, Current Drugs Ltd, GB, vol. 9, No. 1, 2006, pp. 49-55. cited by applicant

Mantero et al., “Long-term treatment of mineralocorticoid excess syndromes”, Divisions of Endocrinology, Universities of Padua and Ancona, Italy, *Steroids*, file for Case IPR2016-00286, Janssen Exhibit 2066, on Jan. 1995, vol. 60 pp. 81-86. cited by applicant

Marc B-Garnick, “Management of Metastatic Carcinoma of the Prostate—Treatment Options and Controversies”, in *Prostatic Disorders*, (David F. Paulson et al. eds., 1989), pp. 354-367. cited by applicant

Maria I., “Male pseudohermaphroditism due to 17 alpha-hydroxylase deficiency”, *Journal of Clinical Investigation* 1970, vol. 49 No. 10 pp. 1930-1941. cited by applicant

Marik PE et al., Recommendations for the diagnosis and management of corticosteroid insufficiency in critically ill adult patients: consensus statements from an international task force by the American College of Critical Care Medicine. *Critical Care Medicine* 2008, vol. 36, No. 6, pp. 1937-1949. cited by applicant

Marin Fakhri et al., “Glucocorticoids and Treatment of Prostate Cancer: A Preliminary and Clinical Review”. *Urology* 60. 2002, pp. 553-561. cited by applicant

Marini et al., “The effect of adjuvant prednisone combined with GMF on patterns of relapse and occurrence of second malignancies in patients with breast cancer”, *Annals of Oncology*, file for Case IPR2016-00286, Janssen Exhibit 2060, on 1996, vol. 7, pp. 245-250. cited by applicant

Mark. J. Ratain et al., Statistical and Ethical Issues in the Design and Conduct of Phase I and II Clinical Trials of New Anticancer Agents, *Journal of the National Cancer Institute*, vol. 85, No. 20, pp. 1637-1649, Oct. 20, 1993. cited by applicant

Martins et al., “A Validated Liquid Chromatographic-Tandem Mass Spectroscopy Method for the Quantification of Abiraterone Acetate and Abiraterone in Human Plasma,” *J. Chromatography B*, vol. 843:262-257 (2006). cited by applicant

Masuda et al., MPromise of vitamin D analogues in the treatment of hyperproliferative conditions, *Mol. Cancer Ther.*, Vol 5(4) pp. 797-808 (2006). cited by applicant

Mateo, J., et al., “DNA-Repair Defects and Olaparib in Metastatic Prostate Cancer”, *The New England Journal of Medicine*, 2015, vol. 373, No. 18, pp. 1697-1708. cited by applicant

Mayo Clinic “Prednisone and other corticosteroids” for Case# IPR2016-00286 Janssen Exhibit 2102 Oct. 3, 2016 4 Pages. cited by applicant

Mayo Clinic Website, “Prostate cancer”, <http://www.mayoclinic.org/diseasesconditions/prostatecancer/basics/definition/con-20029597?p=1> (accessed Jun. 28, 2016), Mylan Pharms. Inc., Exhibit 1051, 11 pages. cited by applicant

MB. Sawyer et al., “Phase I Study of an Oral Formulation of ZD9331 Administered Daily for 28 Days”, *Journal of Clinical Oncology*, filed for IPR2016-00286, vol. 21, No. 9, pp. 1859-1865, May 1, 2003. cited by applicant

McKay et al., “A phase II trial of abiraterone acetate (AA) without prednisone in castration resistant prostate cancer (CRPC)”, *Genitourinary (Prostate) Cancer, An American Society of Clinical Oncology Journal*, 5 pages. cited by applicant

McPhaul. Michael J., “Mechanisms of Prostate Cancer Progression to Androgen Independence,” Best Practice & Research Clinical Endocrinology & Metabolism. vol. 22(2):373-388 (2008). cited by applicant

Medical Dictionary, “Refractory cancer definition of refractory cancer by Medical dictionary” Janssen Exhibit 2103 *Amerigen* vs. *Janssen* filed for Case# IPR2016-00286. Oct. 3, 2016, 1 page. cited by applicant

MedlinePlus, MACTH stimulation test, Available at <https://medlineplus.gov/ency/article/003696.htm>, last visited Sep. 30, 2016, file for Case IPR2016-00286, Janssen Exhibit 2050, 4 pages. cited by applicant

Melby JC et al., “Comparative studies on adrenal cortical function and cortisol metabolism in healthy adults and in patients with shock due to infection”, Journal Clin Invest 1958, vol. 37 No. 12 pp. 1791-1798. cited by applicant

Mendel et al., “In vivo antitumor activity of SU11248, a novel tyrosine kinase inhibitor targeting vascular endothelial growth factor and platelet-derived growth factor receptors determination of a pharmacokinetic/pharmacodynamic relationship”, Clinical Cancer Research, Jan. 2003. vol. 9, No. 1. pp. 327-337. cited by applicant

Michaelson et al., “Randomized, Placebo-Controlled, Phase III Trial of Sunitinib Plus Prednisone Versus Prednisone Alone in Progressive, Metastatic, Castration-Resistant Prostate Cancer.” Journal of Clinical Oncology, vol. 31. pp. 1-8, (2013). cited by applicant

Millikan, et al., “Randomized phase 2 trial of ketoconazole and ketoconazole/doxorubicin in androgen independent prostate cancer” Urologic Oncology, file for Case IPR2016-00286, Janssen Exhibit 2064, on 2001. vol. 6, pp. 111-115. cited by applicant

Milliken et al., EB1089, a vitamin D receptor agonist, reduces proliferation and decreases tumor growth rate in a mouse model of hormone-induced mammary cancer, *M Cancer letters*, New York, NY, vol. 229 (2):205-215 (2005). cited by applicant

Moreira et al., “CYP17 Inhibitors for Prostate Cancer Treatment—An Update,” *Current Medicinal Chemistry*, vol. 15:888-899 (2008). cited by applicant

Moreira et al., Synthesis and evaluation of novel 17-indazole androstene derivatives designed as CYP17 inhibitors, Elsevier, 2007, pp. 939-948, vol. 72. cited by applicant

Morgan et al., “Impact of prednisone on toxicities and survival in metastatic castration-resistant prostate cancer: A systematic review and meta-analysis of randomized clinical trials”, *Critical Reviews in Oncology/Hematology*, Jun. 2014, vol. 90, No. 3, pp. 253-261. cited by applicant

Mostaghel et al, “Intracrine Androgen Metabolism in Prostate Cancer Progression: Mechanisms of Castration Resistance and Therapeutic Implications,” *Best Practice & Research Clinical Endocrinology & Metabolism*, vol. 22(2) 243-258 (2008). cited by applicant

Mostaghel, E.A., “Abiraterone in the treatment of metastatic castration-resistant prostate cancer”, *Cancer Management Res.* (2014) 6, p. 39-51. cited by applicant

Mostaghel, EA et al. Molecular Pathways Targeting resistance in the androgen receptor for therapeutic benefit, *Clin Cancer Res*, Dec. 4, 2013. cited by applicant

Mottet et al., “Highlights on Prostate Cancer from Urological and Oncological Congresses in 2007,” *European Urology Supplements*, vol. 7:460-476 (2008). cited by applicant

Mulcahy, Phase 3 Trial of Immunotherapy for Metastatic Prostate Cancer Terminated Janssen Exhibit 2082 *Amerigen* vs. *Janssen* for Case#. IPR2016v0028-6, Oct. 17, 2008, 2 Pages. cited by applicant

Murai et al, parp inhibitor resistance mediated by SLFN11 inactivation; *Oncotarget* 7: 76534, 2016. cited by applicant

Murphy W.J. JL Orcutt & P.C. Remus (2012) *Patent Valuation: Improving Decision Making through Analysis* Hoboken NJ: Wiley. cited by applicant

N. Rornero-Laorden et al., Prospective Evaluation of the Response to Prednisone-Dexamethasone Switch in Castration-Resistant Prostate Cancer Patients Treated with abiraterone pre-and post-

docetaxel, *Journal of Clinical Oncology* vol. 34, Nov. 14, 2016. cited by applicant

Nakabayashi, M., et al.. “Response to Low-Dose Ketoconazole and Subsequent Dose Escalation to High-Dose Ketoconazole in Patients with Androgen-Independent Prostate Cancer”, *Amer Cane Soc.*, 107(5), (2006) pp. 975-981. cited by applicant

National Cancer Institute-seer Stat Fact Sheets: Prostate Cancer Janssen Exhibit 2089, *Amerigen vs. Janssen* filed for case # PR2016-00286, Oct. 3, 2016, pp. 1-11. cited by applicant

NCCN Practice Guidelines in Oncology. V. 1.2005, Prostate Cancer, 41 pages. cited by applicant

NCT01576172, “Abiraterone Acetate and Prednisone With or Without Beliparib in Treating Patients With Metastatic Castration-Resistant Prostate Cancer”, *ClinicalTrials.gov*, Apr. 12, 2012. cited by applicant

NCT01867710 *Clinicaltrials.gov*, “A Randomized Phase 2 Study Evaluating Abiraterone Acetate With Different Steroid Regimens for Preventing Symptoms Associated With Mineralocorticoid Excess in Asymptomatic, Chemotherapy-naïve and Metastatic Castration-resistant Prostate Cancer (mCRPC) Patients” file for Case IPR2016-00286, *Amerigen* Exhibit 1187, on Jan. 14, 2017, 4 pages. cited by applicant

NCT02500901, “Enzalutamide and Niraparib in the Treatment of Metastatic Castrate-Resistant Prostate Cancer (CRPC)”, *ClinicalTrials.gov.*, Jul. 17, 2015, downloaded on Sep. 29, 2017 from “clinicaltrials.gov/ct2/show/NCT0200901”, 7 pages. cited by applicant

New Treatment for Prostate Cancer Under Development May 22, 1996 Wed for Case#. IPR2016-01582. cited by applicant

Newell et al. The Cancer Research UK experience of pre-clinical toxicology studies to support early clinical trials with novel cancer therapies, *Elsevier*, 2004, pp. 899-906, vol. 40. cited by applicant

Nimalasena et al., “Paraneoplastic Cushing's Syndrome in Prostate Cancer: A Difficult Management Problem,” *BJU International*, vol. 101:424-427 (2007). cited by applicant

Nishimura, Kazuo, et al. Potential Mechanism for the Effects of Dexamethasone on Growth of Androgen-Independent Prostate Cancer, *Journal of the National Cancer Institute*, 2001, pp. 1739-1746, vol. 93. cited by applicant

Nishiyama, et al., Hormone/Antihormone Withdrawal and Oexamethasone, for Hormone-Refractory Prostate Cancer, *international Journal of Urology*, 10, 1997, vol. 5, 44-47. cited by applicant

“A Safety and Pharmacokinetics Study of Niraparib Plus an Androgen Receptor-Targeted Therapy in Men With Metastatic Castration Resistant Prostate Cancer (Bedivere)”, NCT02924766, Jul. 26, 2019, Version 17, Jul. 26, 2019, 4 pages. cited by applicant

“A Safety and Pharmacokinetics Study of Niraparib Plus an Androgen Receptor-Targeted Therapy in Men With Metastatic Castration-Resistant Prostate Cancer (Bedivere)”, NCT02924766 Version 4, May 12, 2017, 4 pages. cited by applicant

“A Safety and Pharmacokinetics Study of Niraparib Plus Apalutamide in Men With Metastatic Castration-Resistant Prostate Cancer (Bedivere)”, NCT02924766, Version 2 Clinical Trial Nov. 3, 2016, 8 pages. cited by applicant

“A Safety and Pharmacokinetics Study of Niraparib Plus Apalutamide in Men With Metastatic Castration-Resistant Prostate Cancer (Bedivere)”, NCT02924766 Version 1, Oct. 4, 2016, 8 pages. cited by applicant

“Abiraterone/Prednisone, Olaparib, or Abiraterone/Prednisone + Olaparib in Patients With Metastatic Castration-Resistant Prostate Cancer With DNA Repair Defects”, NCT03012321, Version 5, Feb. 8, 2017, 13 pages. cited by applicant

“Ph II Study to Evaluate Olaparib With Abiraterone in Treating Metastatic Castration Resistant Prostate Cancer”, NCT01972217, Version 27, Feb. 2, 2017, 19 pages. cited by applicant

Brown et al., “Targeting DNA Repair in Cancer: Beyond PARP Inhibitors”, *Cancer Discovery*, Jan. 2017, vol. 7, No. 1, pp. 20-37. cited by applicant

Chi et al., "Niraparib and Abiraterone Acetate for Metastatic Castration-Resistant Prostate Cancer", J Clin Oncol., Mar. 23, 2023, vol. 41, No. 18, pp. 3339-3351. cited by applicant

Drean et al., "PARP inhibitor combination therapy", Critical Reviews in Oncology/Hematology, vol. 108, 2016, pp. 73-85. cited by applicant

Geethakumari et al., "PARP Inhibitors in Prostate Cancer", Curr Treat Options Oncol., 2017, vol. 18, 37, pp. 1-16. cited by applicant

History of Changes for Study: NCT01576172, "Abiraterone Acetate and Prednisone With or Without Veliparib in Treating Patients With Metastatic Hormone-Resistant Prostate Cancer", NCT01576172 Version 52, Dec. 22, 2016, 23 pages. cited by applicant

Schiewer et al., "Dual roles of PARP-1 promote cancer growth and progression", Cancer Discovery, Dec. 2012, vol. 2, No. 12, pp. 1134-1149. cited by applicant

Study NCT03012321, "Abiraterone/Prednisone, Olaparib, or Abiraterone/Prednisone + Olaparib in Patients With Metastatic Castration Resistant Prostate Cancer With DNA Repair Defects", NCT03012321 Version 5 Feb. 8, 2017, 13 pages. cited by applicant

Tozer et al., "Introduction to Pharmacokinetics and Pharmacodynamics", The Quantitative Basis of Drug Therapy, 2006, 40 pages. cited by applicant

Michie, C.O., et al., "Final results of the phase I trial of niraparib (MK4827), a poly(ADP)ribose polymerase (PARP) inhibitor incorporating proof of concept biomarker studies and expansion cohorts involving BRCA1/2 mutation carriers, sporadic ovarian, and castration resistant prostate cancer (CRPC)," Journal of Clinical Oncology, vol. 31, No. 15, suppl, May 20, 2013, pp. 2513-2513. cited by applicant

Phillip et al., Targeting PARP in prostate cancer: novelty, pitfalls, and promise. Oncology, vol. 30, No. 5, May 15, 2016 p. 377-377. cited by applicant

Nnane et al., Inhibition of Androgen Synthesis in Human Testicular and Prostatic Microsomes and in Male Rata by Novel Steroidal Compounds, Endocrinology, 1999, pp. 2891-2897. vol. 140 No. 6. cited by applicant

O'Dormen et al., British Journal of Cancer, 2004, 90, :2317-2325. cited by applicant

Oelkers W. et al., Dose-response relationships between plasma adrenocorticotropin (ACTH), cortisol, aldosterone, and 18-hydroxycorticosterone after injection of ACTH-(1-39) of human corticotropin-releasing hormone in man, Journal of Clinical Endocrinology Metabolism, 1988, vol. 66, No. 1, pp. 181-186. cited by applicant

Official Journal of the American Urological Association, "Prostate Cancer", Journal Urology, vol. 177, No. 4, Apr. 2007, 2 pages. cited by applicant

Official Journal of the American Urological Association, Inc., "The Journal of Urology", vol. 135, Number 4, part 2, filed for Case, IPR2016-00286, Janssen Exhibit 2022 on Apr. 1986, 203A pages, Abstract 397. cited by applicant

Oh W. K., "Secondary hormonal therapies in the treatment of prostate cancer", Urology 60(Supp. 3A): pp. 87-93, (Sep. 2002). cited by applicant

Oliver Sartor et al., Effect of Prednisone on Prostate-Specific Antigen in Patients With Hormone-Refractory Prostate Cancer, Urology, 52: 1998, pp. 252-256. cited by applicant

Olmos et al., "Reply: Clinical Outcome and Prognostic Factors for Patients Treated Within a Phase I Study: The Royal Marsden Hospital Experience," British Journal of Cancer, vol. 99:1365 (2008). cited by applicant

Opar, Alisa, ASCO Presentations Highlight Value of Cancer Biomarkers, Nature Reviews, vol. 7:547-548 (2008). cited by applicant

Osaba, D., et al., "Health-Related Quality of Life in Men with Metastatic Prostate Cancer Treated with Prednisone alone or Mitoxantrone and Prednisone" J Clin. Oncol. (1999), 17(6), p. 165-1663. cited by applicant

Ospina et al., "ACTH Stimulation Tests for the Diagnosis of Adrenal Insufficiency: Systematic Review and Meta-Analysis", The Journal of Clinical Endocrinology & Metabolism Feb. 2016, vol.

101 No. 2 pp. 427-434. cited by applicant

Oudar, Stephane, et al. Actualite dans le cancer de la prostate, Synthese, Bull Cancer 2005; 92 (10), pp. 865-873 (relevance in English abstract). cited by applicant

Oudard et al., "Prostate Cancer: update," Bull. Cancer, 92(10), pp. 865-873 (2005). cited by applicant

P. M. Clark et al., "Defining the normal cortisol response to the short Synacthen test", Implications for the investigation of hypothalamic pituitary disorders, clinical Endocrinology 1998 vol. 49 pp. 287-292. cited by applicant

Paltiel et al., Management of severe hypokalemia in hospitalized patients: a study of quality of care based on computerized database, Arch Intern Med. Apr. 23, 2001, vol. 161, No. 8, pp. 1089, 1095. cited by applicant

Papatsoris et al., "Novel Biological Agents for the Treatment of Hormone-Refractory Prostate Cancer (HRPC)", Current Medicinal Chemistry 2007, filed for Case IPR2016-00286, Janssen Exhibit 2010, vol. 12, pp. 277-296. cited by applicant

Patent Owner Janssen Oncology, Inc., for U.S. Pat. No. 8,822,438, *Mylan vs. Janssen*, filed for case ,PR2016-01332, Janssen Exhibit 2189, 86 pages. cited by applicant

PCT International Search Report for PCT/US2007/018770, published Feb. 28, 2008. cited by applicant

PCT Written Opinion of the International Searching Authority for PCT/US2007/018770, published Feb. 28, 2008. cited by applicant

Peehl et al., "Preclinical activity of Ketoconazole in combination with calcitriol or the vitamin D analogue EB 1089 in prostate cancer cells," J. Urology, vol. 168, pp. 1583-1588 (2002). cited by applicant

Petrylak et al., "Docetaxel and Estramustine Compared with Mitoxantrone and Prednisone for Advanced Refractory Prostate Cancer", New England Journal of Medicine, file for Case IPR2016-01582; Janssen Exhibit 2059, on Oct. 7, 2004, vol. 351, No. 15, pp. 1513-1520. cited by applicant

Petrylak, "Future Directions In The Treatment Of Androgen-Independent Prostate Cancer," Urology 65 (Supplement 6A), pp. 8-13 (2005). cited by applicant

Petrylak, D.P., "New Paradigms for Advanced Prostate Cancer", Rev. Urol. (2007), 9, Suppl. 2, S3-S12. cited by applicant

PMLive Website, "Top 50 Pharmaceutical Products by Global Sales", <http://pmlive.com/top-pharma-list/Top-50-pharmaceutical-products-by-global-sales> (accessed Jun. 30, 2016), Mylan Pharms. Inc., Exhibit 1055, 3 pages. cited by applicant

Polkinghorn et al, AR signaling regulates DNA repair in prostate cancer; Cancer discovery 3: 1245, 2013. cited by applicant

Posner. G.H. et al., A non-Calcemic Sulfone Version of the Vitamin D3 Analogue Seocalcitol (E81089): Chemical Synthesis. Biological Evaluation and Potency Enhancement of the Anticancer Drug Adriamycin, Bioorganic & Medicinal Chem. 9: 2365-71 (2001). cited by applicant

Postma, "Treatment of Prostate Cancer", Annals of Oncology, Sep. 2006, vol. 17, Supplement 10, x207-x210. cited by applicant

Potter et al. "A Convenient Large-Scale Synthesis of Abiraterone Acetate [3Beta-acetoxy-17-(3-pyridyl)androsta-5,16-diene], a Potential New Drug for the Treatment of Prostate Cancer," Organic Preparations and Procedures Intl, vol. 29(1):123-134 (1997). cited by applicant

Potter et al. "Discovery of highly potent and selective enzyme inhibitors with potential for the treatment of prostate cancer; the important dual role of transition metal chemistry in both drug design and synthesis," Poster presented at the SmithKline Beecham Research Symposium, Robinson College, Cambridge. England (1993). cited by applicant

Potter et al., "Highly potent inhibitors of human cytochrome P-450 (17alpha)," Poster presented at the third drug discovery and development symposium, San Diego, California (1993). cited by applicant

Potter et al., "Novel Steroidal Inhibitors of Human Cytochrome P4501711 (17alpha-Hydroxylase-C17.20-lyase): Potential Agents for the Treatment of Prostatic Cancer," J. Med. Chem, vol. 38:2463-2471 (1995). cited by applicant

Prednisone Side Effects from <http://www.drugs.com/sfx/prednisone-side-effects.html?printable=1>, (Jul. 8, 2010). cited by applicant

PredniSONE Tablets USP, 1 mg, 2.5 mg, 5 mg, 10 mg, 20 mg and 50 mg, PredniSONE Oral Solubon USP, 5 mg per 5 mL and PredniSONE Intensol (Trademark) Oral Solution (Concentrate), 5 mg per mL, Revised Nov. 2012, 18 pages. cited by applicant

Prelone, Physician's Desk Reference, 54th edition 1959-1960 (2000). cited by applicant

Prostate Cancer End Points Workshop Jun. 21-22, 2004, Bethesda Marriott—Bethesda MD, 66 pages. cited by applicant

Prostate Cancer Principles and Practice, Taylor & Francis (2006) Chapter 93. cited by applicant

Public Citizen Press Room Release—"Antifungal Treatment Should Be Taken Off the Market, Public Citizen Tells FOAD,," Filed for Case IPR2016-00286, Janssen Exhibit 2019. on Feb. 24, 2015, 1 page. cited by applicant

R. J. Auchus et al: "Use of Predni sone With Abiraterone Acetate in Metastatic Castration-Resistant Prostate Cancer", The Oncologist, vol. 19, No. 12, Oct. 31, 2014 (Oct. 31, 2014), pp. 1231-1240. cited by applicant

R. Venkitararnan et al., A Randomised Phase 2 Trial of Dexamethasone Versus Prednisolone in Castration-resistant Prostate Cancer, *European Urology*, 67, pp. 673-679 (2015). cited by applicant

Raghavan et al., "Prostate Cancer: Moving Forward by Reinventing the Wheel . . . But This Time it is Round." J. Clin. Oncol, Vol. 26(28):4535-4536 (2008). cited by applicant

Ramiah et al.. Clinical Endpoints for Drug Development in Prostate Cancer: Current Opinion in Urology, vol. 18:303-308 (2008). cited by applicant

Randy Osborne, "Historic' and 'practice-changing' Zytiga delivers in prostate cancer at ASCO", 2 pages. cited by applicant

Reid et al., "CYP17 Inhibition as a Hormonal Strategy for Prostate Cancer," *Nature Clin. Prac. Urology*, vol. 5(11):610-620 (2008). cited by applicant

Reid et al., "Inhibition of Androgen Synthesis Results in a High Response Rate in Castration Refractory Prostate Cancer (CRPC)," *Annals of Oncol.* vol. 18 (Supp. 9): abstract No. 50PD: 173-74 (2007). cited by applicant

Reid et al., "Selective CYP17 Inhibition with Abiraterone Acetate (AA) Results in a High Response Rate (RR) in Castration-Resistant Prostate Cancer (CRPC) Confirming the Continued Importance of Targeting Androgen Receptor (AR) Signaling," ASCO 2008 Genitourinary Cancers Symposium, abstract No. 50 (2008). cited by applicant

Reid et al., "Significant and Sustained Antitumor Activity in Post-Docetaxel, Castration-Resistant Prostate Cancer With the CYP17 Inhibitor Abiraterone Acetate," J. Clin. Oncol., vol. 28:(9), pp. 1489-1495 (2010). cited by applicant

Reid, A., et al., "Annals of Oncology", Educational and Abstract Book of the ESMO Conference Lugano (ECLU), (2007), 18(Supplement 9), ix173-ix174. Abstract 50PD. cited by applicant

Remington—The Science and Practice of Pharmacy, 20th Edition, filed for Case IPR2016-01582, Janssen Exhibit 2026. on 2000, pp. 1363-1370. cited by applicant

Report to the Nation on Prostate Cancer 2004. Prostate Cancer Foundation. cited by applicant

Richard d. Auchus, "Steroid 17-hydroxylase and 17,20-lyase deficiencies, genetic and pharmacologic", *Journal of Steroid Biochemistry & Molecular Biology*, 2017, vol. 165, pp. 71-78. cited by applicant

U.S. Appl. No. 16/131,772, filed Sep. 14, 2018. cited by applicant

U.S. Appl. No. 15/663,082, filed Jul. 28, 2017. cited by applicant

Antolin et al, "Linking off-target kinase pharmacology to the differential cellular effects observed among PARP inhibitors", *Oncotarget*, vol. 5, No. 10, May 2014, pp. 3023-3028. cited by applicant

Carlson et al, "PARP Inhibitors Show Promise Against Metastatic Triple-Negative Breast Cancer in Early Studies," Oncology Times, vol. 31, Issue 15, Aug. 10, 2009, pp. 10-11. cited by applicant
Carlson, "PARP Inhibitors Show Promise Against Metastatic Triple-Negative Breast Cancer in Early Studies", Oncology Times, Special Edition, American Society of Clinical Oncology 45th Annual Meeting, May 29-Jun. 2, 2009, 3pp. cited by applicant
Carroll, "Updated: J&J grabs prostate cancer rights for Tesaro's niraparib in \$500M deal", Apr. 6, 2016, p. 1. cited by applicant
Hopkins et al, "Mechanistic Dissection of PARP1 Trapping and the Impact on In Vivo Tolerability and Efficacy of PARP Inhibitors", Mol Cancer Res, vol. 13, Issue 11, 2015, pp. 1465-1477. cited by applicant
Johnson & Johnson Q3 2016 Results—Earnings Call Transcript Oct. 18, 2016, pp. 1-32. cited by applicant
Murai, et al., "Differential trapping of PARP1 and PARP2 by clinical PARP inhibitors", National Institute of Health, Cancer Res. Nov. 1, 2012; 72(21): 5588-5599. cited by applicant
Wahlberg et al., "Family-wide chemical profiling and structural analysis of PARP and tankyrase inhibitors", Nature Biotechnology, vol. 30, No. 3, 2012, pp. 283-289. cited by applicant
Fong et al., Inhibition of Poly(ADP-Ribose) Polymerase in Tumors from BRCA Mutation Carriers. New England Journal of Medicine 2009, 361 (2), 123-134. cited by applicant

Primary Examiner: Rao; Savitha M

Attorney, Agent or Firm: BakerHostetler

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS (1) The present application is a continuation of U.S. patent application Ser. No. 16/131,772, filed Sep. 14, 2018, which is a continuation of U.S. patent application Ser. No. 15/663,082, filed Jul. 28, 2017, which claims the benefit of U.S. Provisional Application No. 62/368,466, filed on Jul. 29, 2016, and U.S. Provisional Application No. 62/369,239, filed on Aug. 1, 2016, each of which are incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

(1) The present invention relates to the treatment of metastatic hormone-naïve prostate cancer in a human by administering a safe and/or an effective amount of niraparib to such human.

BACKGROUND OF THE INVENTION

(2) Prostate cancer is the most common non-cutaneous malignancy in men and the second leading cause of death in men from cancer in the western world. Prostate cancer results from the uncontrolled growth of abnormal cells in the prostate gland. Once a prostate cancer tumor develops, androgens, such as testosterone, promote prostate cancer tumor growth. At its early stages, localized prostate cancer is often treated with local therapy including, for example, surgical removal of the prostate gland and radiotherapy. However, when local therapy fails to cure prostate cancer, as it does in up to a third of men, the disease progresses into incurable metastatic disease (i.e., disease in which the cancer has spread from one part of the body to other parts).

(3) Treatment of metastatic prostate cancer Androgen deprivation therapy ("ADT") or androgen suppression therapy is performed to reduce the testicular production of testosterone. ADT includes surgical castration (orchiectomy) or the use of luteinizing hormone-releasing hormone ("LHRH") antagonists or agonists. Examples of LHRH antagonists include degarelix. Examples of LHRH agonists include goserelin acetate, histrelin acetate, leuprolide acetate, and triptorelin palmoate.

(4) Abiraterone acetate is a prodrug of abiraterone, inhibits 17 α hydroxylase/C17, 20-lyase (cytochrome P450c17 [CYP17]), a key enzyme in androgen biosynthesis. Abiraterone acetate in combination with prednisone has been approved for the treatment of men with metastatic castration-resistant prostate cancer (“mCRPC”) who have received prior chemotherapy containing docetaxel. The efficacy and safety of abiraterone acetate (1,000 mg daily tablet dose) and prednisone (5 mg twice daily) therapy in patients with mCRPC is established by the results of COU-AA-301 and COU-AA-302, both Phase 3, multinational, randomized, double-blind, placebo-controlled studies. Study COU-AA-301 was the first Phase 3 study to demonstrate that further lowering testosterone concentrations below that achieved with androgen deprivation therapy (“ADT”) using CYP17 inhibition with abiraterone acetate improves survival in patients with mCRPC. COU-AA-302 demonstrated significantly improved overall survival (“OS”) and radiographic progression-free survival (“rPFS”) in chemotherapy-naïve patients with mCRPC treated with abiraterone acetate plus prednisone compared with placebo plus prednisone. What is needed are data to determine whether abiraterone acetate in combination with low-dose prednisone and ADT is superior to ADT alone in improving rPFS and OS in subjects with mHNPc with high-risk prognostic factors.

(5) Thus, the treatment of prostate cancer, including castrate resistant prostate cancer and metastatic castrate resistant prostate cancer, by way of PARP inhibition with niraparib in mCRPC patients, including those with DNA-repair anomalies. This treatment may follow chemotherapy or may be a chemo-naïve subject. This treatment may follow AR-targeted agents, e.g., enzalutamide, apalutamide, and bicalutamide. Therefore, niraparib may present another treatment option.

SUMMARY OF THE INVENTION

(6) The present invention is directed to a method for treating prostate cancer in a human in need of such treatment comprising, consisting of, and/or consisting essentially of administering to the human a therapeutically effective amount of niraparib.

(7) In an embodiment, the present invention is directed to a method of treating prostate cancer in a human in need of such treatment comprising, consisting of, and/or consisting essentially of administering to the human a therapeutically effective amount of niraparib, wherein the prostate cancer is castration-resistant prostate cancer (“CRPC”), metastatic castration-resistant prostate cancer, and/or antiandrogen-resistant prostate cancer.

(8) In another embodiment, the present invention is directed a method for treating prostate cancer in a human in need of such treatment comprising, consisting of and/or consisting essentially of administering niraparib to a human, wherein the human is carrying at least one DNA repair anomaly selected from the group consisting of BRCA-1, BRCA-2, FANCA, PALB2, CHEK2, BRIP1, HDAC2, and ATM.

(9) In another embodiment, the present invention is directed a method for treating prostate cancer in a human in need of such treatment comprising, consisting of and/or consisting essentially of administering niraparib to a human, wherein the human is carrying at least one DNA repair anomaly that is either BRCA-1 or BRCA-2.

(10) In another embodiment, the present invention is directed to a method of treating prostate cancer in a human in need of such treatment comprising, consisting of, and/or consisting essentially of administering to the human niraparib in an amount of, preferably, from about 30 mg/day to about 400 mg/day, more preferably 300 mg/day, and most preferably once daily oral administration in three 100 mg oral dosage forms.

(11) In another embodiment, the present invention is directed to a composition comprising niraparib for the treatment of prostate cancer, antiandrogen resistant prostate cancer, castration-resistant prostate cancer, and metastatic castration-resistant prostate cancer.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

- (1) FIG. 1. Illustrates that niraparib inhibits the growth of human prostate tumor cell lines in vitro.
- (2) FIG. 2. Illustrates that niraparib suppresses PAR formation in two human prostate tumor cell lines in vitro.
- (3) FIG. 3. Illustrates that niraparib treatment induces increased γ -H2AX in 22RV1 cells in a dose-dependent manner, as measured by flow cytometry.
- (4) FIG. 4. Illustrates that niraparib induces γ -H2AX in 22RV1, LNCaP AR-TB, and C4-2B cells in vitro.
- (5) FIG. 5. Illustrates that niraparib treatment inhibits growth of C4-2B-luc prostate tumors in NSG male mice.

DETAILED DESCRIPTION OF THE INVENTION

- (6) The term “subject” refers to a mammal, most preferably a human, who has been or is the object of treatment, observation or experiment.
- (7) The term “treatment” refers to the treatment of a subject afflicted with a pathological condition and refers to an effect that alleviates the condition by killing the cancerous cells, but also to an effect that results in the inhibition of the progress of the condition, and includes a reduction in the rate of progress, a halt in the rate of progress, amelioration of the condition, and cure of the condition. Treatment as a prophylactic measure (i.e., prophylaxis) is also included.
- (8) The term “therapeutically effective amount” refers to an amount of niraparib that elicits the biological or medicinal response in a tissue system that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation or partial alleviation of the symptoms of the disease, syndrome, condition, or disorder being treated.
- (9) The term “safe and effective amount” refers to an amount of niraparib that elicits the prevention or amelioration of disease progression and acceptable unacceptable toxicity in the human.
- (10) The term “composition” refers to a pharmaceutical product that includes the specified ingredients sometimes in therapeutically effective amounts, as well as any product that results, directly, or indirectly, from combinations of the specified ingredients in the specified amounts.
- (11) The term “pharmaceutically acceptable” as used herein pertains to compound, materials, compositions and/or dosage forms that are, within the scope of sound medical judgement, suitable for use in contact with the tissues of a human without excessive toxicity, irritations, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. Each carrier, excipient, etc. must all be “acceptable” in the sense of being compatible with the other ingredients of the formulation.
- (12) The term “androgen receptor” as used herein is intended to include the wild-type androgen receptor as well as androgen-resistant ARs and/or AR mutants associated with castration-resistant prostate cancer.
- (13) As used herein, the term “antiandrogen” refers to a group of hormone receptor antagonist compounds that is capable of preventing or inhibiting the biologic effects of androgens on normally responsive tissues in the body. In some embodiments, an anti-androgen is a small molecule. Antiandrogens include enzalutamide, apalutamide, and abiraterone acetate.
- (14) As used herein, the term “first-generation anti-androgen” refers to an agent that exhibits antagonist activity against a wild-type AR polypeptide. However, first-generation anti-androgens differ from second-generation anti-androgens in that first-generation anti-androgens can potentially act as agonists in CRPC.
- (15) Exemplary first-generation anti-androgens include, but are not limited to, flutamide, nilutamide and bicalutamide.
- (16) As used herein, the term “second-generation anti-androgen” refers to an agent that exhibits full antagonist activity against a wild-type AR polypeptide. Second-generation anti-androgens differ from first-generation anti-androgens in that second-generation anti-androgens act as full

antagonists in cells expressing elevated levels of AR, such as for example, in CRPC. Exemplary second-generation anti-androgens include 4-[7-(6-cyano-5-trifluoromethylpyridin-3-yl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-5-yl]-2-fluoro-N-methylbenzamide (also known as ARN-509; CAS No. 956104-40-8); 4-(3-(4-cyano-3-(trifluoromethyl)phenyl)-5,5-dimethyl-4-oxo-2-thioxoimidazolidin-1-yl)-2-fluoro-N-methylbenzamide (also known as MDV3100 or enzalutamide; CAS No: 915087-33-1) and RD162 (CAS No. 915087-27-3). In some embodiments, a second-generation anti-androgen binds to an AR polypeptide at or near the ligand binding site of the AR polypeptide.

(17) As used herein, the term “third-generation anti-androgen” refers to an agent that exhibits full antagonist activity against a wild-type AR polypeptide and against mutant forms of the AR polypeptide, with mutations arising in the ligand binding domain (LBD) of the AR polypeptide as set forth below. Third-generation anti-androgens retain the differentiation from first-generation anti-androgens in that third-generation anti-androgens act as full antagonists in cells expressing elevated levels of AR, such as for example, in CRPC.

(18) As used herein, the term “mutant” refers to an altered (as compared with a reference) nucleic acid or polypeptide, or to a cell or organism containing or expressing such altered nucleic acid or polypeptide.

(19) As used herein, unless otherwise noted, the term “affect” or “affected” (when referring to a disease, syndrome, condition or disorder that is affected by antagonism of AR) includes a reduction in the frequency and/or severity of one or more symptoms or manifestations of said disease, syndrome, condition or disorder; and/or include the prevention of the development of one or more symptoms or manifestations of said disease, syndrome, condition or disorder or the development of the disease, condition, syndrome or disorder.

(20) Embodiments of the present invention include prodrugs of niraparib. In general, such prodrugs will be functional derivatives of the compounds that are readily convertible in vivo into the required compound. Thus, in the methods of treating or preventing embodiments of the present invention, the term “administering” encompasses the treatment or prevention of the various diseases, conditions, syndromes and disorders described with the compound specifically disclosed or with a compound that may not be specifically disclosed, but which converts to the specified compound in vivo after administration to a patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in “Design of Prodrugs”, ed. H. Bundgaard, Elsevier, 1985.

(21) Androgen Receptor (AR)

(22) Androgens bind to a specific receptor, the androgen receptor (AR), inside the cells of target tissues. The AR is expressed in numerous tissues of the body and is the receptor through which the physiological as well as the pathophysiological effects of endogenous androgen ligands, such as testosterone (T) and dihydrotestosterone (DHT), are expressed. Structurally, the AR is composed of three main functional domains: the ligand binding domain (LBD), the DNA-binding domain, and amino-terminal domain. A compound that binds to the AR and mimics the effects of an endogenous AR ligand is referred to as an AR agonist, whereas a compound that inhibits the effects of an endogenous AR ligand is termed an AR antagonist. Binding of androgen to the receptor activates it and causes it to bind to DNA binding sites adjacent to target genes. From there it interacts with coactivator proteins and basic transcription factors to regulate the expression of the gene. Thus, via its receptor, androgens cause changes in gene expression in cells. These changes ultimately have consequences on the metabolic output, differentiation or proliferation of the cell that are visible in the physiology of the target tissue. In the prostate, androgens stimulate the growth of prostate tissue and prostate cancer cells by binding to the AR that is present within the cytoplasm of androgen sensitive tissue.

(23) Compounds that selectively modulate AR are of clinical importance in the treatment of or prevention of a variety of diseases, conditions, and cancers, including, but not limited to, prostate

cancer, benign prostatic hyperplasia, hirsutism in women, alopecia, anorexia nervosa, breast cancer, acne, musculoskeletal conditions, such as bone disease, hematopoietic conditions, neuromuscular disease, rheumatological disease, cancer, AIDS, cachexia, for hormone replacement therapy (HRT), employed in male contraception, for male performance enhancement, for male reproductive conditions, and primary or secondary male hypogonadism.

(24) Castration Resistant Prostate Cancer

(25) Agents that block the action (antiandrogens) of endogenous hormones (e.g., testosterone) are highly effective and routinely used for the treatment of prostate cancer (androgen ablation therapy). While initially effective at suppressing tumor growth, these androgen ablation therapies eventually fail in almost all cases, leading CRPC. Most, but not all, prostate cancer cells initially respond to androgen withdrawal therapy. However, with time, surviving populations of prostate cancer cells emerge because they have responded to the selective pressure created by androgen ablation therapy and are now refractory to it. Not only is the primary cancer refractory to available therapies, but cancer cells may also break away from the primary tumor and travel in the bloodstream, spreading the disease to distant sites (especially bone). This is known as metastatic castration resistant prostate cancer ("mCRPC"). Among other effects, this causes significant pain and further bone fragility in the subject.

(26) In some embodiments, the subject's prostate cancer is resistant to or non-responsive to antiandrogen treatment, including, but not limited to, enzalutamide, apalutamide and abiraterone acetate ("antiandrogen resistance").

(27) The preparation of niraparib, 2-[4-[(3S)-piperidin-3-yl]phenyl]indazole-7-carboxamide, may be found in U.S. Pat. No. 8,071,623, issued on Dec. 6, 2011 and entitled Amide Substituted Indazoles as Poly(ADP-Ribose) Polymerase (PARP) Inhibitors, which claims the benefit of U.S. provisional patent application No. 60/921,310, filed on Feb. 16, 2010, as well as U.S. Pat. No. 8,436,185, issued on May 7, 2013 and entitled Pharmaceutically Acceptable Salts of 2-[4-[(3S)-piperidin-3-yl]phenyl]-2H-indazole-7-carboxamide, which claims the benefit of U.S. provisional patent application No. 61/010,333 filed on Jan. 8, 2008, each of which is incorporated herein by reference.

(28) The invention also provides pharmaceutical compositions comprising niraparib and a pharmaceutically acceptable carrier. The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs.

(29) Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, microcrystalline cellulose, sodium croscarmellose, corn starch, or alginic acid; binding agents, for example starch, gelatin, polyvinyl-pyrrolidone or acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to mask the unpleasant taste of the drug or delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a water-soluble taste masking material such as hydroxypropyl-methylcellulose or hydroxypropylcellulose, or a time delay material such as ethyl cellulose, cellulose acetate butyrate may be employed.

(30) Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate

or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water-soluble carrier such as polyethyleneglycol or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

(31) Aqueous suspensions contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanal, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

(32) Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as butylated hydroxyanisole or alpha-tocopherol.

(33) Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

(34) The pharmaceutical compositions of the invention may also be in the form of an oil-in-water emulsion. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally occurring phosphatides, for example soy bean lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavoring agents, preservatives and antioxidants.

(35) Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, flavoring and coloring agents and antioxidant. The pharmaceutical compositions may be in the form of a sterile injectable aqueous solution. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.

(36) The sterile injectable preparation may also be a sterile injectable oil-in-water microemulsion where the active ingredient is dissolved in the oily phase. For example, the active ingredient may be first dissolved in a mixture of soybean oil and lecithin. The oil solution then introduced into a water and glycerol mixture and processed to form a microemulsion.

(37) The injectable solutions or microemulsions may be introduced into a patient's blood stream by local bolus injection. Alternatively, it may be advantageous to administer the solution or microemulsion in such a way as to maintain a constant circulating concentration of the instant compound. In order to maintain such a constant concentration, a continuous intravenous delivery device may be utilized. An example of such a device is the Deltec CADD-PLUS™ model 5400 intravenous pump.

(38) The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension for intramuscular and subcutaneous administration. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

(39) Niraparib may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

(40) For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the instant compounds are employed. (For purposes of this application, topical application shall include mouth washes and gargles.)

(41) Niraparib can be administered in intranasal form via topical use of suitable intranasal vehicles and delivery devices, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen. Niraparib may also be delivered as a suppository employing bases such as cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

(42) When niraparib is administered to a subject, the selected dosage level will depend on a variety of factors including, but not limited to, the activity of the particular compound, the severity of the individual's symptoms, the route of administration, the time of administration, the rate of excretion of the compound, the duration of the treatment, other drugs, compounds, and/or materials used in combination, and the age, sex, weight, condition, general health, and prior medical history of the patient. The amount of niraparib and route of administration will ultimately be at the discretion of the physician, although generally the dosage will be to achieve local concentrations at the site of action which achieve the desired effect without causing substantial harmful or deleterious side-effects.

(43) Administration in vivo can be effected in one dose, continuously or intermittently (e.g. in divided doses at appropriate intervals) throughout the course of treatment. Methods of determining the most effective means and dosage of administration are well known to those of skill in the art and will vary with the formulation used for therapy, the purpose of the therapy, the target cell being treated, and the subject being treated. Single or multiple administrations can be carried out with the dose level and pattern being selected by the treating physician.

(44) In general, a suitable dose of niraparib is in the range of about 100 μg to about 250 mg per kilogram body weight of the subject per day. Where the active compound is a salt, an ester, prodrug, or the like, the amount administered is calculated on the basis of the parent compound and so the actual weight to be used is increased proportionately.

(45) A therapeutically effective amount of niraparib or a pharmaceutical composition thereof for the treatment of prostate cancer includes a dose range from about 30 mg/day to about 400 mg/day of niraparib, or any particular amount or range therein, in particular about 300 mg/day, and once daily oral administration in three 100 mg oral dosage forms. Optimal dosages of niraparib to be administered may be readily determined and will vary with the particular compound used, the mode of administration, the strength of the preparation and the advancement of the disease,

syndrome, condition or disorder. In addition, factors associated with the particular subject being treated, including subject gender, age, weight, diet and time of administration, will result in the need to adjust the dose to achieve an appropriate therapeutic level and desired therapeutic effect. The above dosages are thus exemplary of the average case. There can be, of course, individual instances wherein higher or lower dosage ranges are merited, and such are within the scope of this invention.

(46) Niraparib may be administered in any of the foregoing compositions and dosage regimens or by means of those compositions and dosage regimens established in the art whenever use of niraparib is required for a subject in need thereof.

EXAMPLES

(47) The following Examples are set forth to aid in the understanding of the invention, and are not intended and should not be construed to limit in any way the invention set forth in the claims which follow thereafter.

Example 1

(48) In Vitro Cytotoxicity of Niraparib in Human Prostate Tumor Lines

(49) The cytotoxicity of niraparib was tested in several human prostate tumor lines in vitro. None of the tumor lines is known to be BRCA-1 or BRCA-2 deficient.

(50) Methods:

(51) In vitro cytotoxicity of niraparib was assessed in 5 human prostate cancer cell lines: C4-2B, LNCaP, LNCaP AR.TB, VCaP, and 22Rv1. C4-2B, LNCaP, LNCaP AR.TB, and 22Rv1 cell lines were grown in RPMI1640+GlutaMAX™-I medium (Life Technologies #61870-036) supplemented with 10% heat-inactivated fetal bovine serum (FBS) (Life Technologies #16140-071) and non-essential amino acids (NEAA) (Life Technologies #11140-050); and VCaP cells were grown in DMEM+GlutaMAX™-I medium (Life Technologies #10569-010) with 10% FBS and NEAA. VCaP cells were subcultured every 7 days; other lines were split every 3-4 days.

(52) Cell growth kinetics of each cell line were determined by seeding cells at several densities and monitoring growth at intervals up to 7 days. Growth was determined using the Promega Cell TiterGlo reagent (#G7571) to measure cellular ATP by means of a chemiluminescent luciferin-luciferase reaction. Plates were read on a Perkin-Elmer Envision plate reader, and luminescence values were plotted in order to identify seeding densities that resulted in log-phase growth and a cell density within the linear range of the Cell TiterGlo assay at the desired time point.

(53) For niraparib cytotoxicity experiments, cells were harvested by brief trypsinization and each line was seeded to the inner 60 wells of 96-well plates in 100 μ L of medium at an appropriate density for a 7-day treatment. The outer wells of each plate were filled with Dulbecco's phosphate buffered saline (DPBS; Life Technologies #14190-144) to reduce evaporation from test wells. Cells were rested overnight in the plates at 37° C. in a humidified 5% CO₂ incubator. Treatment was initiated by addition of 50 μ L of 3 \times niraparib (final concentrations 500, 125, 31.3, 7.8, 1.95, 0.49, 0.12, 0.03 μ M) in the appropriate medium to triplicate wells. The final vehicle concentration was 0.5% DMSO.

(54) Cells were cultured for 7 days. Relative cell viability after treatment was determined using Cell TiterGlo reagent, as above. All luminescent output values were normalized to percent inhibition based on mean luminescence of the untreated control wells, and the mean percent inhibition of the vehicle control wells was subtracted from each treatment value. Percent inhibition was plotted vs log μ M concentration in GraphPad Prism 7.00. Nonlinear regression and calculation of EC₅₀ values were performed using the log (agonist) vs. response—Variable slope (four parameters) fit.

(55) Results:

(56) Results of the cytotoxicity assay are shown in FIG. 1 and Table 1. Growth of each cell line was reduced in a dose-dependent fashion by increasing concentrations of niraparib. C4-2B cells appeared to be the most sensitive, with an EC₅₀ value of ~1.2 μ M. VCaP cells appeared to be

the least sensitive with an EC.sub.50 value of 4.1 μ M.

(57) TABLE-US-00001 TABLE 1 EC.sub.50 values for 7-day niraparib treatment of human prostate tumor cell lines. Cell Line EC.sub.50, μ M C4-2B 1.222 LNCaP 3.502 LNCaP AR.TB 2.140 VCaP 4.099 22Rv1 3.517

Example 2

(58) Inhibition of PAR Formation by Niraparib

(59) The ability of niraparib to inhibit the formation of poly (ADP) ribose (PAR) was tested in two human prostate tumor lines in vitro. Neither of the tumor lines is known to be BRCA-1 or BRCA-2 deficient.

(60) Methods:

(61) PAR inhibition using niraparib was assessed in 2 human prostate cancer cell lines, C4-2B and VCaP. The C4-2B cell line was grown in RPMI1640+GlutaMAX™-I medium supplemented with 10% FBS and NEAA and split every 3-4 days. VCaP cells were grown in DMEM+GlutaMAX™-I medium with FBS and NEAA and subcultured every 7 days.

(62) Cells were harvested by brief trypsinization and each line was seeded into 6-well plates in 1 mL of medium at an appropriate density. An extra 500 μ L of complete medium was added, for a total volume per well of 1.5 mL. Cells were rested overnight in the plates at 37° C. in a humidified 5% CO.sub.2 incubator. The next day, medium was removed from plates and cells were washed using 1 mL serum free medium (RPMI or DMEM respectively). Treatment was initiated by addition of 1 mL of niraparib (final concentrations 100, 10, 1, 0.1, 0.01 and 0 μ M in 0.1% DMSO) in the appropriate medium to triplicate wells. Plates were returned to the incubator for two hours.

(63) Following treatment, extracts were prepared using reagents and procedures provided in the HT PARP in vivo Pharmacodynamic Assay II (Trevigen #4520-096-K). Medium was removed from each well and placed into separate labeled microfuge tubes, and the plates were placed on ice. The tubes were spun at 1500 rpm for 4 minutes to pellet any cells that detached from the plate during drug incubation. Lysis buffer was prepared using 24.5 mL of cell lysis reagent with 250 μ L of 100 mM PMSF (in ethanol; Sigma #93482) and 250 μ L 100 \times Protease Inhibitor Cocktail (Thermo Scientific #78429). Lysis buffer (300 μ L) was added to each well of the plates, on ice. Adherent cells were scraped into lysis buffer and kept on ice for at least 15 minutes. Supernatant was removed from the microfuge tubes, and the cell lysates from the 6-well plates were added to each tube from the corresponding tubes. SDS (20% w/v) was added to bring the final SDS concentration to 1%. The cell extracts were heated to 95-100° C. for 5 minutes. After cooling to room temperature, 0.01 volume of 100 \times magnesium cation and 3 μ L DNase were added to each tube. Tubes were briefly vortexed and returned to a 37° C. incubator for 90 minutes. After the incubation, tubes were centrifuged at 10,000 \times g for 10 minutes at room temperature. If a pellet was present, it was removed using a pipette tip and extracts were transferred to a 96 well dilution plate. Cell extracts were frozen at -80° C. until used for protein quantitation and the PAR ELISA assay. The ELISA assay protocol was performed according to the manufacturer's instructions.

(64) Protein quantitation was performed using the detergent-compatible Biorad DC protein assay kit II (#500-0002) with the Biorad Quick Start Bovine Serum Albumin Standard Set (#5000207) according to the manufacturer's 96-well plate protocol. ELISA lysis buffer was spiked into the standards, and an equal volume of PBS was added to all sample wells to correct for any effect of the lysis buffer on protein readings. Samples were assayed in duplicate. Buffer A' (25 μ L) was added to all wells of the plate, and 200 μ L of Buffer B was immediately added to each well. Plates were incubated for 15 minutes at room temperature on a shaker. Absorbance was read at 750 nm on a Molecular Devices M5 plate reader, using the DC Protein Assay protocol in SoftMax Pro version 6.3 software. Linear regression of the standard curve, interpolation of sample protein values, and replicate averaging were performed in the software. Data was exported to Excel, where any corrections for sample dilution were performed.

(65) Luminescence values of PAR ELISA standards and samples were analyzed in GraphPad Prism

version 7, where linear regression of the standard curve and interpolation of sample values were calculated. Interpolated PAR values (pg PAR/mL) were corrected for sample dilution and divided by the corresponding protein concentration to yield pg PAR/mg of protein. These values were graphed in GraphPad Prism v7.

(66) Results:

(67) The results from the PAR assay are shown in FIG. 2. PAR was reduced in a dose-dependent fashion by increasing concentrations of niraparib in each cell line.

Example 3

(68) Niraparib Induces γ -H2AX in Human Prostate Tumor Lines In Vitro

(69) The ability of niraparib to induce double stranded breaks in DNA was measured in 3 human prostate cancer cell lines 22RV1, LNCaP AR.TB, and C4-2B. Double stranded breaks in DNA are followed by phosphorylation of adjacent histone γ -H2AX, and this phosphorylation can be measured by antibody staining and flow cytometry.

(70) Methods:

(71) 22RV1, LNCaP AR.TB, and C4-2B cell lines were grown as outlined above. Cell lines were passaged every 3-4 days.

(72) For each cell line, 2×10^5 cells were seeded in each well of a 12-well plate (Falcon #353043) in a volume of 1 mL of media. Cells were rested overnight in a 37° C. humidified 5% CO₂ incubator, then 1 mL of media containing $2 \times$ concentrated serially diluted niraparib was added to achieve final concentrations of 200, 100, 50, 25, 12.5, 6.25, 3.13, 1.57, 0.78, 0.39, 0.2, and 0.1 μ M in triplicate wells. The final vehicle concentration was 0.2% DMSO and triplicate wells of vehicle and media controls were also obtained for each cell line. Plates were incubated for another 18 hours.

(73) Following the 18 hour incubation with drug, each well of cells was harvested by first transferring the 2 mL of media into a 15 mL conical tube (Corning #430798). 500 μ L of cell dissociation buffer (Gibco #13151-014) was then added to the well and allowed to sit for 5 minutes. Using a 1 mL pipette, 1 mL of media was added to the well, cells were dislodged by pipetting, and the cell-containing media was transferred to the corresponding 15 mL conical tube. Tubes were centrifuged at 1200 rpm for 5 minutes, the supernatant was discarded, and the pelleted cells were resuspended and transferred to a 96-well v-bottom plate (Costar #3896). Plates were centrifuged at 1800 rpm for 3 minutes, supernatant discarded, then wells were washed with 200 μ L of DPBS. This process was repeated for a total of 3 washes. Cells were then stained with 100 μ L of DPBS containing a 1:800 dilution of Invitrogen Live/Dead fixable aqua (Invitrogen #L34957) for 20 minutes at 4° C. Cells were then washed with 150 μ L BD Pharmingen Stain Buffer (stain buffer; BD #554657) and centrifuged at 1800 rpm for 3 minutes. Cells were washed again 2 times with 200 μ L stain buffer and then once with DPBS.

(74) Cells were fixed with 100 μ L -20° C. 70% ethanol/H₂O and plates were stored at -20° C. for 2 hours. Cells were washed 3 times with stain buffer centrifuging at 2200 rpm for 3 minutes between each wash. Cells were then incubated with 100 μ L of a 1:1 dilution stain buffer and AXELL biotin-free Fc receptor blocker (Accurate Chemical & Scientific Corp #NB309) for 20 minutes at 4° C. Cells were washed with 150 μ L of stain buffer then centrifuged at 2200 rpm for 3 minutes, supernatant discarded, then cells were incubated with 50 μ L stain buffer containing 0.2% v/v Triton X-100 (Acros Organics #21568-2500) for 2 hours at room temperature in the dark along with 1:100 dilutions of γ -H2AX antibody (Biolegend #613408).

(75) Cells were washed once with 200 μ L stain buffer containing 0.2% v/v Triton X-100, and then washed once with 200 μ L stain buffer only. Cells were resuspended in 80 μ L of stain buffer and 50 μ L was analyzed on a BD Fortessa flow cytometer. Data were analyzed using TreeStar FlowJo v9.8.5. Data was gated on live cells, then following doublet discrimination, the entire population was assessed for γ -H2AX antibody signal. Results were graphed In GraphPad Prism v7.

(76) Results:

(77) Representative histograms for the 22RV1 cell line are shown in FIG. 3, depicting the effect of different concentrations of niraparib. Drug-treated samples were compared to vehicle and media controls and are graphed in FIG. 4. The lowest concentrations where γ -H2AX signal rises significantly above vehicle control are indicated in Table 2. The results show that, in each prostate tumor line, niraparib induces γ -H2AX in a dose-dependent manner.

(78) TABLE-US-00002 TABLE 2 Minimum concentration of niraparib that induces significant change in γ -H2AX 1.sup.st Significant Cell line Concentration (μ M) 22RV1 1.57 LnCaP.AR.TB 3.13 C4-2B 1.57

Example 4

(79) Niraparib Inhibits the Growth of C4-2B Human Prostate Tumors in Mice

(80) The activity of niraparib was tested in the pre-established human prostate subcutaneous C4-2B model in non-obese diabetic (NOD) severe combined immunodeficient (scid) gamma (NOD.Cg-Prkdc Il2rg/SzJ) (NSG) mice. This tumor model is not believed to be BRCA-1 or BRCA-2 deficient.

(81) Methods:

(82) Vehicle was 0.5% Methyl cellulose (Methocel™ F4M) prepared and kept at 4° C. in the dark. All formulations were made to be dosed at a volume of 10 ml/kg body weight. NSG male mice (Jackson Laboratories) were used. Animals were habituated for one week prior to any experimental procedures being performed. Mice were group housed (5 per cage) in disposable IVC-cages (Innovive, San Diego, CA, USA) under a 12-h light: dark cycle at a temperature of 19 to 22° C. and 35 to 40% humidity. Mice were fed an autoclaved high fat (6%) diet laboratory chow and water ad libitum.

(83) Mice were injected with LNCaP C4-2B-luc tagged cells (1×10^6 tumors cells in a 200 μ l volume of Cultrex®: RPMI 1640 medium (1:1 ratio) on the right flank. Mice were randomized per tumor volumes (tumor volume = 241 ± 14 mm³), with 10 mice per treatment group. Mice were dosed daily by gavage (p.o.) with either vehicle, or vehicle containing niraparib as indicated below at 10 ml/kg dosing volume. Start of treatment = Day 1. Mice were treated through study day 24. Group 1 0 mg/kg Vehicle (0.5% Methocel F4M) dosed QD p.o. Group 2 25 mg/kg niraparib in 0.5% Methocel F4M dosed QD p.o. Group 3 50 mg/kg niraparib in 0.5% Methocel F4M dosed QD p.o.

(84) For each individual animal, body weight and tumor volume [using the formula: Tumor Volume (mm³) = ($a \times b^2$)/2; where 'a' represents the length, and 'b' the width of the tumor as determined by caliper measurements], were monitored twice weekly throughout the study. For the pre-established tumors, a time-course of tumor growth is expressed as mean \pm standard error of the mean (SEM).

(85) Results:

(86) The vehicle treated mice started to reach ethical limits for tumor volume around study day 22 onwards (see FIG. 5 for individual tumor volumes). Tumor volume data was presented up to study day 24 (when 9 of 10 vehicle-treated mice remained on study). After 18, 22, and 24 days of treatment, group 3 dosed daily with 50 mg/kg niraparib p.o. showed significant inhibition/delay in tumor growth, with tumor growth inhibition (TGI) values of ~40% on these days. Significant differences in tumor growth were observed on days 18, 22 and 24 (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$). Mice dosed at 25 mg/kg of niraparib did not show significant tumor growth inhibition, though there was modest TGI of ~12% on days 22 and 24.

Example 5

(87) A multicenter, open-label study is carried out to assess the efficacy and safety of once daily dosing of 300 mg niraparib in male subjects over the age of 18 years with mCRPC and DNA-repair anomalies who have had at least one line of taxane-based chemotherapy and at least one line of antiandrogen therapy (e.g., abiraterone acetate, enzalutamide, apalutamide). The study will enroll approximately 100 subjects. Subjects will be monitored for safety during the study period, and up

to 30 days after the last dose of study drug. Treatment will continue until disease progression, unacceptable toxicity, death, or the sponsor terminates the study.

(88) The study will consist of 4 phases; a Prescreening Phase for biomarker evaluation only, a Screening Phase, a Treatment Phase, and a Follow-up Phase. The efficacy evaluations include the following: Tumor measurements: chest, abdomen and pelvis CT or MRI scans and whole body bone scans (.sup.99mTc), serum PSA, survival status, CTC, and symptomatic skeletal event (SSE).

(89) Niraparib, 300 mg, will be provided as capsules (3×100 mg) for once daily oral administration. The capsules must be swallowed whole. Subjects should take their dose in the morning (with or without food). Although not considered study medication, subjects who have not undergone surgical castration must continue to receive regularly prescribed GnRHa. All GnRHa therapies should be recorded in the concomitant medication section of the eCRF.

(90) A treatment cycle is defined as 28 days. Subjects will begin taking niraparib on Day 1 of Cycle 1. Sufficient quantities of niraparib for each treatment cycle will be distributed on the first day of each cycle. If subjects miss a dose, then that dose should be replaced if the subject remembers within an approximate 12-hour window. Otherwise, subjects should take the next dose the following day, without compensating for the missed dose. Missed doses should be recorded in the eCRF.

(91) Prescreening Phase for Biomarker Evaluation

(92) The Prescreening Phase will evaluate if a potential subject is biomarker-positive for DNA-repair anomalies. All subjects will be required to sign a specific ICF for the Prescreening Phase and provide baseline demographic characteristics and disease-specific medical history. The Prescreening Phase may occur any time prior to the Screening Phase.

(93) The process for determining biomarker-positivity will be different for subjects who enter the Prescreening Phase before a blood-based assay is available, compared with those subjects who enter after a blood-based assay is available. The 2 processes are described below

(94) Process for Determining Biomarker-Positivity Before a Blood-Based Assay is Available

(95) The Subject signs the prescreening ICF. If the subject has had tumor tissue previously analyzed by the FoundationOne® gene panel, then after the subject grants a release, the FoundationOne® data can be reviewed to determine eligibility based on the criteria defined in Table 1. If the subject is biomarker-positive, they are eligible to enter the Screening Phase. If the subject has not had tumor tissue previously analyzed by the FoundationOne® gene panel, then they must have either archival or recently collected (recommended) tumor tissue analyzed for biomarker-positivity by a sponsor-approved test. If the subject is biomarker-positive, they are eligible to enter the Screening Phase.

(96) Blood samples will also be collected from all subjects during the Prescreening Phase and stored for when a blood-based assay becomes available. At the time a blood-based assay becomes available, the stored blood sample will be analyzed for concordance with the tumor tissue sample results. This analysis may occur at any time after the blood-based assay becomes available.

(97) Process for Determining Biomarker-Positivity after a Blood-Based Assay is Available

(98) Subject signs the prescreening ICF. Subject has blood collected and sent for analysis of biomarker-positivity. If the subject has had tumor tissue previously analyzed by the FoundationOne® gene panel, then after the subject grants a release, the FoundationOne® data can be reviewed to determine eligibility based on the criteria defined in Table 1. If the subject is biomarker-positive, they are eligible to enter the Screening Phase and do not need to wait for results of the blood-based analysis. If the FoundationOne® gene panel is negative, the subject may still be considered eligible if they are determined to be biomarker-positive by the blood-based assay. If the subject has not had tumor tissue previously analyzed by the FoundationOne® gene panel and archival tissue is available, then a request for retrieval and analysis of the archived tumor tissue is initiated. If the blood-based assay results are biomarker-positive, then the subject is eligible to enter the Screening Phase and does not need to wait for results of the archival tumor

tissue-based analysis. The results from the archival tumor tissue-based analysis, when they are available, may be used in conjunction with the blood-based results for concordance and bridging studies.

(99) At the discretion of the study sponsor, if the blood-based assay results are negative, then the archival tumor tissue-based results may be used to determine eligibility.

(100) If no archival tumor tissue is available, then the subject must agree to have tumor tissue collected.

(101) If the blood-based assay results are biomarker-positive, then the recent tumor tissue must be collected prior to Cycle 1 Day 1 for later use in concordance and bridging studies. Analysis of the recently collected tumor tissue may occur any time during the study and the results may not be required prior to the subject entering the Screening Phase.

(102) At the discretion of the study sponsor, if the blood-based assay results are negative, then the recently collected tumor tissue may be used to determine eligibility.

(103) Once subjects are identified as biomarker-positive during the Prescreening Phase, the Screening Phase should start within 30 days.

(104) Screening Phase

(105) All biomarker-positive subjects must sign the main study ICF prior to the conduct of any study-related procedures in the Screening Phase. During this phase, eligibility criteria will be reviewed and a complete clinical evaluation will be performed as specified in the Time and Events Schedule. Screening procedures will be performed up to 35 days before Cycle 1 Day 1, unless otherwise specified. Imaging will be accepted up to 8 weeks prior to Cycle 1 Day 1. Screening clinical safety laboratory evaluations can be used for Cycle 1 Day 1 assessments if performed within 14 days of Cycle 1.

(106) Subjects who do not meet all inclusion criteria, or who meet an exclusion criterion, may be rescreened once. Rescreening is at the discretion of the investigator and requires sponsor approval and agreement. Subjects who are to be rescreened must sign a new ICF before rescreening. Subjects rescreened within 35 days of planned enrollment may use the initial screening laboratory results, computed tomography (CT)/magnetic resonance imaging (MRI) and bone scans (if still within 8 weeks of Cycle 1 Day 1) to determine eligibility if not the reason for the rescreening.

(107) Treatment Phase

(108) The Treatment Phase will begin at Cycle 1 Day 1 and will continue until the study drug is discontinued. The last measurements taken on Day 1 of Cycle 1 before administration of the study drug or at screening (whichever value was last) will be defined as the baseline values. Visits for each cycle will have a ± 3 -day window, unless otherwise specified. Study visits will be calculated from the Cycle 1 Day 1 date. Subjects may have imaging performed within ± 7 days of visits requiring images. Refer to the Time and Events Schedule for treatment visits and assessments during the Treatment Phase.

(109) For PK and pharmacodynamics sampling days, the subject must not take the study drug at home on the morning of study visits. Study drug should be taken at the site. Details of PK and pharmacodynamics sampling days and times are provided in the Time and Events Schedule. Details of blood sample handling and storage procedures for PK and pharmacodynamics are provided in the laboratory manual.

(110) Clinical evaluations and laboratory studies may be repeated more frequently, if clinically indicated. Study drug treatment will continue until disease progression, unacceptable toxicity, death, or the sponsor terminates the study. Once the subject discontinues study drug, the subject must complete the End-of-Treatment (EoT) visit within 30 days after the last dose of study drug, and enter the Follow-up Phase.

(111) End-of-Treatment Visit An End of Treatment visit must be scheduled within 30 days after the last dose of study drug or prior to administration of a new anti-prostate cancer therapy, whichever occurs first. If a subject is unable to return to the site for the EoT visit, the subject should be

contacted to collect AEs that occurred within 30 days after the last dose of study drug.

Follow-up Phase

(112) Once a subject has completed the Treatment Phase, survival follow-up and SSEs will be performed every 3 months either via clinic visits, telephone interview, chart review, or other convenient methods. Deaths regardless of causality and SAEs thought to be related to study drugs will be collected and reported within 24 hours of discovery or notification of the event. If the follow-up information is obtained via telephone contact, then written documentation of the communication must be available for review in the source documents.

(113) Biomarker-Positive Sample for DNA-Repair Anomalies

(114) To evaluate if subjects are biomarker-positive, a blood-based assay may become available during the study that will provide a more rapid method than tissue-based analysis for determining biomarker-positive status, while being more convenient for the subjects. Prior to the blood-based assay becoming available, tumor tissue (either archival or recently collected) will require analysis. To ensure that all subjects, regardless of when they enter the study, have the same biomarker data available for analysis (i.e., for concordance and bridging studies), both tumor tissue and blood samples will be collected from all subjects who sign the prescreening informed consent form (ICF). The process for determining biomarker-positivity will be different for subjects who enter the Prescreening Phase before the blood-based assay is available, compared with those subjects who enter after the blood-based assay is available. However, the status of biomarker-positivity in both tumor tissue and blood will be assessed for all subjects.

(115) To be eligible for the study, subjects must be confirmed biomarker-positive by 5 tumor tissue (either archival or recently collected), or blood testing when available. The biomarkers of interest for this study and the biomarker-positivity criteria are listed in Table 3. Analyses will be performed to define a proxy for bi-allelic loss (e.g., mutation co-expression frequency with copy number loss) and these proxies may be used to determine biomarker-positivity as that information becomes available.

(116) TABLE-US-00003 TABLE 3 Biomarker Panel and Criteria for Positivity Genomic Lesion

Genes Definition Required for Positivity BRCA-1 Breast Cancer gene 1 Homozygous deletion

BRCA-2 Breast Cancer gene 2 Heterozygous deletion + FANCA Fanconi Anemia

Complementation deleterious mutation Group A gene Copy neutral loss of PALB2 Partner and

Localizer of BRCA2 heterozygosity + gene deleterious mutation CHEK2 Checkpoint Kinase 2

gene BRIP1 BRCA1 Interacting Protein C-terminal Helicase 1 gene HDAC2 Histone Deacetylase

2 gene ATM Ataxia Telangiectasia Mutated gene ATM Ataxia Telangiectasia Mutated gene Mono-

allelic deleterious mutation in the kinase catalytic domain Control Genes AR Androgen Receptor

gene TP53 Tumor Protein 53 gene Mono-allelic loss in all genes will be acceptable for entry into

the study until the existing algorithm for bi-allelic loss is validated.

Circulating Tumor Cells

(117) Blood samples will be collected in a Cellsave tube at timepoints specified in the Time and Events Schedule. CTC enumeration will be evaluated at the central laboratory, to assess response to study drug.

(118) Whole Blood for RNA

(119) Whole blood samples will be collected in a Paxgene tube. Multiple ribonucleic acid (RNA) transcripts found in prostate tumors are detectable in the RNA and analysis of these samples will allow evaluation of potential mechanisms of resistance that may emerge with niraparib.

(120) Circulating Tumor DNA

(121) Plasma samples collected during the course of treatment will be used to screen for changes in the levels or types of DNA-repair anomalies observed over time by circulating tumor DNA (ctDNA), and to monitor for potential markers of resistance to niraparib.

(122) While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the

invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.

Claims

1. A method of treating prostate cancer in a human in need of such treatment comprising administering to the human a therapeutically effective amount of niraparib or a salt thereof, wherein the prostate cancer is antiandrogen resistant and wherein the human is carrying at least one DNA repair anomaly in a gene selected from the group consisting of BRCA-1 and BRCA-2.
 2. The method of claim 1, wherein the prostate cancer is castration-resistant prostate cancer.
 3. The method of claim 1, wherein the prostate cancer is metastatic castration-resistant prostate cancer.
 4. The method of claim 1, wherein niraparib or a salt thereof is administered in an amount of from about 30 mg niraparib/day to about 400 mg niraparib/day.
 5. The method of claim 1, wherein niraparib or a salt thereof is administered in an amount of about 300 mg niraparib/day.
 6. The method of claim 5, wherein niraparib or a salt thereof is administered orally, once daily, in 100 mg niraparib oral dosage forms.
 7. The method of claim 1, wherein the human has had at least one line of taxane-based chemotherapy.
 8. The method of claim 1, wherein the prostate cancer has been exposed to at least one line of enzalutamide-, apalutamide-, or abiraterone acetate-based chemotherapy.
 9. A method of treating castration- and antiandrogen-resistant prostate cancer in a human comprising administering niraparib or a salt thereof in 100 mg niraparib oral dosage forms once daily to the human, wherein the human is carrying at least one DNA repair anomaly in a gene selected from the group consisting of BRCA-1 and BRCA-2.
 10. The method of claim 9, wherein niraparib or a salt thereof is administered in an amount of from about 30 mg niraparib/day to about 400 mg niraparib/day.
 11. The method of claim 9, wherein niraparib or a salt thereof is administered in an amount of about 300 mg niraparib/day.
 12. The method of claim 9, wherein the human has had at least one line of taxane-based chemotherapy.
 13. The method of claim 9, wherein the prostate cancer has been exposed to at least one line of enzalutamide-, apalutamide-, or abiraterone acetate-based chemotherapy.
 14. The method of claim 1, wherein niraparib or a salt thereof is administered in an amount of about 200 mg niraparib/day.
 15. The method of claim 1, wherein niraparib or a salt thereof is administered in an amount of about 50 mg niraparib/day.
 16. The method of claim 1, wherein the DNA repair anomaly is a genomic lesion.
 17. The method of claim 14, wherein the genomic lesion is a homozygous deletion, heterozygous deletion plus deleterious mutation, or copy neutral loss of heterozygosity plus deleterious mutation.
 18. The method of claim 1, further comprising determining whether the human is carrying at least one DNA repair anomaly in a gene selected from the group consisting of BRCA-1 and BRCA-2.
 19. A method of treating prostate cancer in a human in need of such treatment comprising: determining whether the human is carrying at least one DNA repair anomaly in a gene selected from the group consisting of BRCA-1 and BRCA-2, and administering to the human a therapeutically effective amount of niraparib or a salt thereof to a human who is carrying the at least one DNA repair anomaly, wherein the prostate cancer is antiandrogen resistant.
-