

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent
Kind Code
Date of Patent
Inventor(s)

12391957
B2
August 19, 2025
Durbin; Joan E. et al.

Recombinant Newcastle disease viruses and uses thereof for the prevention of RSV disease or human metapneumovirus disease

Abstract

Described herein are recombinant Newcastle disease viruses (“NDVs”) comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a respiratory syncytial virus (“RSV”) F protein or human metapneumovirus (“hMPV”) F protein. Also described herein are recombinant NDVs comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises (i) an RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains; or (ii) an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. The recombinant NDVs and compositions thereof are useful for the immunizing against RSV or hMPV as well as the prevention of RSV disease or hMPV disease.

| | |
|------------------------------|--|
| Inventors: | Durbin; Joan E. (Sergeantsville, NJ), Durbin; Russell K. (Sergeantsville, NJ), Garcia-Sastre; Adolfo (New York, NY), Mena; Ignacio (New York, NY), Palese; Peter (New York, NY) |
| Applicant: | ICAHN SCHOOL OF MEDICINE AT MOUNT SINAI (New York, NY); RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY (New Brunswick, NJ) |
| Family ID: | 1000008765609 |
| Assignee: | ICAHN SCHOOL OF MEDICINE AT MOUNT SINAI (New York, NY); RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY (New Brunswick, NJ) |
| Appl. No.: | 17/268403 |
| Filed (or PCT Filed): | August 16, 2019 |
| PCT No.: | PCT/US2019/046837 |
| PCT Pub. No.: | WO2020/037215 |

PCT Pub. Date: February 20, 2020

Prior Publication Data

| Document Identifier | Publication Date |
|---------------------|------------------|
| US 20210198323 A1 | Jul. 01, 2021 |

Related U.S. Application Data

us-provisional-application US 62765242 20180817

Publication Classification

Int. Cl.: **C07K14/08** (20060101); **A61K39/12** (20060101); **C07K16/10** (20060101); **C12N15/113** (20100101); **C12N15/86** (20060101); A61K39/00 (20060101)

U.S. Cl.:

CPC **C12N15/86** (20130101); **A61K39/12** (20130101); **C07K14/08** (20130101); **C12N15/113** (20130101); A61K2039/505 (20130101); A61K2039/543 (20130101)

Field of Classification Search

USPC: None

References Cited

U.S. PATENT DOCUMENTS

| Patent No. | Issued Date | Patentee Name | U.S. Cl. | CPC |
|------------|-------------|---------------------|----------|-----|
| 4444887 | 12/1983 | Hoffman | N/A | N/A |
| 4474893 | 12/1983 | Reading | N/A | N/A |
| 4676980 | 12/1986 | Segal et al. | N/A | N/A |
| 4714681 | 12/1986 | Reading | N/A | N/A |
| 4716111 | 12/1986 | Osband et al. | N/A | N/A |
| 4816397 | 12/1988 | Boss et al. | N/A | N/A |
| 4816567 | 12/1988 | Cabilly et al. | N/A | N/A |
| 4925648 | 12/1989 | Hansen et al. | N/A | N/A |
| 5122464 | 12/1991 | Wilson et al. | N/A | N/A |
| 5166057 | 12/1991 | Palese et al. | N/A | N/A |
| 5223409 | 12/1992 | Ladner et al. | N/A | N/A |
| 5225539 | 12/1992 | Winter | N/A | N/A |
| 5273745 | 12/1992 | Schirmacher | N/A | N/A |
| 5403484 | 12/1994 | Ladner et al. | N/A | N/A |
| 5413923 | 12/1994 | Kucherlapati et al. | N/A | N/A |
| 5427908 | 12/1994 | Dower et al. | N/A | N/A |
| 5516637 | 12/1995 | Huang et al. | N/A | N/A |
| 5530101 | 12/1995 | Queen et al. | N/A | N/A |

| | | | | |
|---------|---------|----------------------|-----|-----|
| 5545806 | 12/1995 | Lonberg et al. | N/A | N/A |
| 5565332 | 12/1995 | Hoogenboom et al. | N/A | N/A |
| 5569825 | 12/1995 | Lonberg et al. | N/A | N/A |
| 5571698 | 12/1995 | Ladner et al. | N/A | N/A |
| 5573920 | 12/1995 | Randle | N/A | N/A |
| 5580717 | 12/1995 | Dower et al. | N/A | N/A |
| 5585089 | 12/1995 | Queen et al. | N/A | N/A |
| 5601819 | 12/1996 | Wong et al. | N/A | N/A |
| 5625126 | 12/1996 | Lonberg et al. | N/A | N/A |
| 5633425 | 12/1996 | Lonberg et al. | N/A | N/A |
| 5658727 | 12/1996 | Barbas et al. | N/A | N/A |
| 5661016 | 12/1996 | Lonberg et al. | N/A | N/A |
| 5698426 | 12/1996 | Huse | N/A | N/A |
| 5731168 | 12/1997 | Carter et al. | N/A | N/A |
| 5733743 | 12/1997 | Stuart et al. | N/A | N/A |
| 5750753 | 12/1997 | Kimae et al. | N/A | N/A |
| 5766886 | 12/1997 | Studnicka et al. | N/A | N/A |
| 5780225 | 12/1997 | Wigler et al. | N/A | N/A |
| 5786199 | 12/1997 | Palese | N/A | N/A |
| 5807715 | 12/1997 | Morrison et al. | N/A | N/A |
| 5814318 | 12/1997 | Lonberg et al. | N/A | N/A |
| 5820871 | 12/1997 | Palese et al. | N/A | N/A |
| 5821047 | 12/1997 | Garrard et al. | N/A | N/A |
| 5854037 | 12/1997 | Palese et al. | N/A | N/A |
| 5891680 | 12/1998 | Lieschke et al. | N/A | N/A |
| 5939598 | 12/1998 | Kucherlapati et al. | N/A | N/A |
| 5969108 | 12/1998 | McCafferty et al. | N/A | N/A |
| 6005079 | 12/1998 | Casterman et al. | N/A | N/A |
| 6146642 | 12/1999 | Garcia-Sastre et al. | N/A | N/A |
| 6190901 | 12/2000 | Sundick et al. | N/A | N/A |
| 6287554 | 12/2000 | Sundick et al. | N/A | N/A |
| 6329511 | 12/2000 | Vasquez et al. | N/A | N/A |
| 6331415 | 12/2000 | Cabilly et al. | N/A | N/A |
| 6407213 | 12/2001 | Carter et al. | N/A | N/A |
| 6451323 | 12/2001 | Garcia-Sastre et al. | N/A | N/A |
| 6544785 | 12/2002 | Palese et al. | N/A | N/A |
| 6635416 | 12/2002 | Palese et al. | N/A | N/A |
| 6649372 | 12/2002 | Palese et al. | N/A | N/A |
| 6669943 | 12/2002 | Palese et al. | N/A | N/A |
| 6719979 | 12/2003 | Peeters et al. | N/A | N/A |
| 6737522 | 12/2003 | Sundick et al. | N/A | N/A |
| 6811784 | 12/2003 | Haller et al. | N/A | N/A |
| 6852522 | 12/2004 | Palese et al. | N/A | N/A |
| 6896881 | 12/2004 | Russell et al. | N/A | N/A |
| 7052685 | 12/2005 | Rook | N/A | N/A |
| 7056689 | 12/2005 | Lorence et al. | N/A | N/A |

| | | | | |
|----------|---------|----------------------|-----|-----|
| 7060430 | 12/2005 | Palese et al. | N/A | N/A |
| 7141550 | 12/2005 | Molling et al. | N/A | N/A |
| 7223389 | 12/2006 | Zakay-Rones et al. | N/A | N/A |
| 7244558 | 12/2006 | Samal et al. | N/A | N/A |
| 7332169 | 12/2007 | Peeters et al. | N/A | N/A |
| 7384774 | 12/2007 | Palese et al. | N/A | N/A |
| 7442379 | 12/2007 | Garcia-sastre et al. | N/A | N/A |
| 7442527 | 12/2007 | Palese et al. | N/A | N/A |
| 7470426 | 12/2007 | Roberts et al. | N/A | N/A |
| 7488802 | 12/2008 | Collins et al. | N/A | N/A |
| 7494808 | 12/2008 | Palese et al. | N/A | N/A |
| 7547442 | 12/2008 | Peeters et al. | N/A | N/A |
| 7709226 | 12/2009 | Foote | N/A | N/A |
| 7736640 | 12/2009 | Lorence et al. | N/A | N/A |
| 7780962 | 12/2009 | Roberts et al. | N/A | N/A |
| 7833774 | 12/2009 | Palese et al. | N/A | N/A |
| 7858081 | 12/2009 | Bernard et al. | N/A | N/A |
| 8043612 | 12/2010 | Roberts et al. | N/A | N/A |
| 8105578 | 12/2011 | Roberts et al. | N/A | N/A |
| 8124084 | 12/2011 | Lefrancois et al. | N/A | N/A |
| 8163879 | 12/2011 | Wong et al. | N/A | N/A |
| 8168757 | 12/2011 | Finnefrock et al. | N/A | N/A |
| 8326547 | 12/2011 | Liu et al. | N/A | N/A |
| 8354509 | 12/2012 | Craven et al. | N/A | N/A |
| 8475790 | 12/2012 | Jure-Kunkel | N/A | N/A |
| 8486418 | 12/2012 | Bublot et al. | N/A | N/A |
| 8490289 | 12/2012 | Nystrom et al. | N/A | N/A |
| 8492118 | 12/2012 | Wong et al. | N/A | N/A |
| 8507222 | 12/2012 | Wong et al. | N/A | N/A |
| 8591881 | 12/2012 | Palese et al. | N/A | N/A |
| 8709417 | 12/2013 | Allison et al. | N/A | N/A |
| 8765462 | 12/2013 | Medin et al. | N/A | N/A |
| 8871191 | 12/2013 | Pavlakakis et al. | N/A | N/A |
| 8900587 | 12/2013 | Craven et al. | N/A | N/A |
| 8940288 | 12/2014 | Lefrancois et al. | N/A | N/A |
| 8952136 | 12/2014 | Carven et al. | N/A | N/A |
| 9217136 | 12/2014 | Palese et al. | N/A | N/A |
| 9370563 | 12/2015 | Morales et al. | N/A | N/A |
| 9375475 | 12/2015 | Allison et al. | N/A | N/A |
| 9387242 | 12/2015 | Palese et al. | N/A | N/A |
| 9476033 | 12/2015 | Samal et al. | N/A | N/A |
| 9616118 | 12/2016 | Bublot et al. | N/A | N/A |
| 9642298 | 12/2016 | Martin | N/A | N/A |
| 9821016 | 12/2016 | Lauer et al. | N/A | N/A |
| 9937196 | 12/2017 | Samal et al. | N/A | N/A |
| 10023637 | 12/2017 | Allison et al. | N/A | N/A |
| 10251922 | 12/2018 | Palese et al. | N/A | N/A |
| 10308913 | 12/2018 | Palese et al. | N/A | N/A |

| | | | | |
|--------------|---------|----------------------|-----|-----|
| 10383936 | 12/2018 | Samal et al. | N/A | N/A |
| 10519426 | 12/2018 | Cheng et al. | N/A | N/A |
| 2002/0052030 | 12/2001 | Wonderling et al. | N/A | N/A |
| 2002/0150554 | 12/2001 | Sundick et al. | N/A | N/A |
| 2003/0044384 | 12/2002 | Roberts et al. | N/A | N/A |
| 2003/0078410 | 12/2002 | Garcia-sastre et al. | N/A | N/A |
| 2003/0224017 | 12/2002 | Samal et al. | N/A | N/A |
| 2004/0005545 | 12/2003 | Fouchier et al. | N/A | N/A |
| 2004/0234552 | 12/2003 | Peeters et al. | N/A | N/A |
| 2005/0042664 | 12/2004 | Wu et al. | N/A | N/A |
| 2005/0048549 | 12/2004 | Cao et al. | N/A | N/A |
| 2005/0191617 | 12/2004 | Inoue et al. | N/A | N/A |
| 2005/0235134 | 12/2004 | O'Sullivan | N/A | N/A |
| 2005/0238622 | 12/2004 | Axelrod et al. | N/A | N/A |
| 2006/0159695 | 12/2005 | Delvecchio et al. | N/A | N/A |
| 2006/0216310 | 12/2005 | Lorence et al. | N/A | N/A |
| 2008/0057037 | 12/2007 | Roberts et al. | N/A | N/A |
| 2008/0206201 | 12/2007 | Beier et al. | N/A | N/A |
| 2009/0061521 | 12/2008 | Palese et al. | N/A | N/A |
| 2009/0081161 | 12/2008 | Roberts et al. | N/A | N/A |
| 2009/0082299 | 12/2008 | Felber et al. | N/A | N/A |
| 2009/0175826 | 12/2008 | Subbiah et al. | N/A | N/A |
| 2009/0214590 | 12/2008 | Sundick et al. | N/A | N/A |
| 2009/0238791 | 12/2008 | Jacques et al. | N/A | N/A |
| 2009/0280144 | 12/2008 | Garcia-sastre et al. | N/A | N/A |
| 2010/0092430 | 12/2009 | Beier et al. | N/A | N/A |
| 2010/0297072 | 12/2009 | DePinho et al. | N/A | N/A |
| 2011/0020282 | 12/2010 | Beier et al. | N/A | N/A |
| 2011/0044937 | 12/2010 | Bell et al. | N/A | N/A |
| 2011/0081311 | 12/2010 | Pavlakis et al. | N/A | N/A |
| 2011/0081374 | 12/2010 | Bublott et al. | N/A | N/A |
| 2011/0158938 | 12/2010 | Bernard et al. | N/A | N/A |
| 2011/0189189 | 12/2010 | Jure-Kunkel | N/A | N/A |
| 2012/0034242 | 12/2011 | Jooss et al. | N/A | N/A |
| 2012/0058141 | 12/2011 | Palese et al. | N/A | N/A |
| 2012/0058538 | 12/2011 | Palese et al. | N/A | N/A |
| 2012/0064112 | 12/2011 | Samal et al. | N/A | N/A |
| 2012/0071859 | 12/2011 | Morgan et al. | N/A | N/A |
| 2012/0114648 | 12/2011 | Langermann et al. | N/A | N/A |
| 2012/0122185 | 12/2011 | Palese et al. | N/A | N/A |
| 2013/0084264 | 12/2012 | Schrier et al. | N/A | N/A |
| 2013/0108665 | 12/2012 | Liang | N/A | N/A |
| 2014/0044678 | 12/2013 | Palese et al. | N/A | N/A |
| 2014/0134128 | 12/2013 | Wong et al. | N/A | N/A |
| 2014/0159960 | 12/2013 | Mueller | N/A | N/A |
| 2014/0186303 | 12/2013 | Subbiah et al. | N/A | N/A |
| 2014/0205560 | 12/2013 | Wong et al. | N/A | N/A |
| 2014/0219955 | 12/2013 | Wong et al. | N/A | N/A |

| | | | | |
|--------------|---------|-------------------|------------|----------------|
| 2014/0242025 | 12/2013 | Wong et al. | N/A | N/A |
| 2014/0271677 | 12/2013 | Palese et al. | N/A | N/A |
| 2014/0271699 | 12/2013 | Kwong | 435/254.11 | C07K 14/005 |
| 2014/0377221 | 12/2013 | Tufaro et al. | N/A | N/A |
| 2015/0017121 | 12/2014 | Becher et al. | N/A | N/A |
| 2015/0093357 | 12/2014 | Lefrancois et al. | N/A | N/A |
| 2015/0132257 | 12/2014 | Wong et al. | N/A | N/A |
| 2015/0133531 | 12/2014 | Wiegand | N/A | N/A |
| 2015/0139945 | 12/2014 | Lefrancois et al. | N/A | N/A |
| 2015/0152188 | 12/2014 | Morisseau et al. | N/A | N/A |
| 2015/0250837 | 12/2014 | Nolin et al. | N/A | N/A |
| 2016/0015760 | 12/2015 | Palese et al. | N/A | N/A |
| 2016/0068823 | 12/2015 | Palese et al. | N/A | N/A |
| 2016/0137721 | 12/2015 | Palese et al. | N/A | N/A |
| 2016/0208222 | 12/2015 | Cheng et al. | N/A | N/A |
| 2017/0037379 | 12/2016 | Palese et al. | N/A | N/A |
| 2017/0247425 | 12/2016 | Ungerechts et al. | N/A | N/A |
| 2017/0285037 | 12/2016 | Kulangara et al. | N/A | N/A |
| 2018/0078592 | 12/2017 | Palese et al. | N/A | N/A |
| 2018/0251555 | 12/2017 | Allison et al. | N/A | N/A |
| 2018/0256655 | 12/2017 | Palese et al. | N/A | N/A |
| 2018/0280455 | 12/2017 | Palese et al. | N/A | N/A |
| 2020/0061184 | 12/2019 | Palese et al. | N/A | N/A |
| 2021/0198323 | 12/2020 | Durbin | N/A | A61P 31/14 |

FOREIGN PATENT DOCUMENTS

| Patent No. | Application Date | Country | CPC |
|------------|---------------------|---------|-----|
| 2002307971 | 12/2007 | AU | N/A |
| 101787373 | 12/2009 | CN | N/A |
| 102740887 | 12/2011 | CN | N/A |
| 105734023 | 12/2015 | CN | N/A |
| 106166294 | 12/2015 | CN | N/A |
| 3922444 | 12/1990 | DE | N/A |
| 0239400 | 12/1993 | EP | N/A |
| 0702085 | 12/1995 | EP | N/A |
| 0780475 | 12/1996 | EP | N/A |
| 0974660 | 12/1999 | EP | N/A |
| 0592106 | 12/2003 | EP | N/A |
| 0519596 | 12/2004 | EP | N/A |
| 1248654 | 12/2004 | EP | N/A |
| 1032269 | 12/2006 | EP | N/A |
| 1486211 | 12/2007 | EP | N/A |
| 2085092 | 12/2008 | EP | N/A |
| 0702085 | 12/2009 | EP | N/A |
| 2669381 | 12/2012 | EP | N/A |
| 2579884 | 12/2013 | EP | N/A |
| 2393921 | 12/2014 | EP | N/A |

| | | | |
|---------------|---------|----|-----|
| 2987856 | 12/2015 | EP | N/A |
| 2766035 | 12/2017 | EP | N/A |
| 2012-527465 | 12/2011 | JP | N/A |
| WO 1986005807 | 12/1985 | WO | N/A |
| WO 1989001036 | 12/1988 | WO | N/A |
| WO 1990002809 | 12/1989 | WO | N/A |
| WO 1991000360 | 12/1990 | WO | N/A |
| WO 1991009967 | 12/1990 | WO | N/A |
| WO 1991010737 | 12/1990 | WO | N/A |
| WO 1991010741 | 12/1990 | WO | N/A |
| WO 1992000373 | 12/1991 | WO | N/A |
| WO 1992001047 | 12/1991 | WO | N/A |
| WO 1992008802 | 12/1991 | WO | N/A |
| WO 1992018619 | 12/1991 | WO | N/A |
| WO 1992022324 | 12/1991 | WO | N/A |
| WO 1993011236 | 12/1992 | WO | N/A |
| WO 1993017105 | 12/1992 | WO | N/A |
| WO 1993017715 | 12/1992 | WO | N/A |
| WO 1994004678 | 12/1993 | WO | N/A |
| WO 1994004690 | 12/1993 | WO | N/A |
| WO 1994025591 | 12/1993 | WO | N/A |
| WO 1994025627 | 12/1993 | WO | N/A |
| WO 1995015982 | 12/1994 | WO | N/A |
| WO 1995015982 | 12/1994 | WO | N/A |
| WO 1995020401 | 12/1994 | WO | N/A |
| WO 1996033735 | 12/1995 | WO | N/A |
| WO 1996034096 | 12/1995 | WO | N/A |
| WO 1996034625 | 12/1995 | WO | N/A |
| WO 1997006270 | 12/1996 | WO | N/A |
| WO 1997012032 | 12/1996 | WO | N/A |
| WO 1997013844 | 12/1996 | WO | N/A |
| WO 1997014433 | 12/1996 | WO | N/A |
| WO 1998002530 | 12/1997 | WO | N/A |
| WO 1998013501 | 12/1997 | WO | N/A |
| WO 1998013501 | 12/1997 | WO | N/A |
| WO 1998016654 | 12/1997 | WO | N/A |
| WO 1998024893 | 12/1997 | WO | N/A |
| WO 1998024893 | 12/1997 | WO | N/A |
| WO 1998046645 | 12/1997 | WO | N/A |
| WO 1998046645 | 12/1997 | WO | N/A |
| WO 1998050433 | 12/1997 | WO | N/A |
| WO 1998050433 | 12/1997 | WO | N/A |
| WO 1998053078 | 12/1997 | WO | N/A |
| WO 1999002657 | 12/1998 | WO | N/A |
| WO 1999015672 | 12/1998 | WO | N/A |
| WO 1999018799 | 12/1998 | WO | N/A |
| WO 1999066045 | 12/1998 | WO | N/A |
| WO 2000062735 | 12/1999 | WO | N/A |
| WO 2000062735 | 12/1999 | WO | N/A |
| WO 2000067786 | 12/1999 | WO | N/A |

| | | | |
|----------------|---------|----|-----|
| WO 2001004333 | 12/2000 | WO | N/A |
| WO 2001020989 | 12/2000 | WO | N/A |
| WO 2001044301 | 12/2000 | WO | N/A |
| WO 2002081621 | 12/2001 | WO | N/A |
| WO 2002081621 | 12/2001 | WO | N/A |
| WO 2002102404 | 12/2001 | WO | N/A |
| WO 2003092579 | 12/2002 | WO | N/A |
| WO 2003092579 | 12/2002 | WO | N/A |
| WO 2006/042156 | 12/2005 | WO | N/A |
| WO 2006050984 | 12/2005 | WO | N/A |
| WO 2006050984 | 12/2005 | WO | N/A |
| WO 2007001677 | 12/2006 | WO | N/A |
| WO 2007001677 | 12/2006 | WO | N/A |
| WO 2007008918 | 12/2006 | WO | N/A |
| WO 2007008918 | 12/2006 | WO | N/A |
| WO 2007064802 | 12/2006 | WO | N/A |
| WO 2007084342 | 12/2006 | WO | N/A |
| WO 2007084342 | 12/2006 | WO | N/A |
| WO 2007113648 | 12/2006 | WO | N/A |
| WO 2007113648 | 12/2006 | WO | N/A |
| WO 2008011726 | 12/2007 | WO | N/A |
| WO 2008156712 | 12/2007 | WO | N/A |
| WO 2009002562 | 12/2007 | WO | N/A |
| WO 2009002562 | 12/2007 | WO | N/A |
| WO 2009095167 | 12/2008 | WO | N/A |
| WO 2009101149 | 12/2008 | WO | N/A |
| WO 2009101149 | 12/2008 | WO | N/A |
| WO 2010091262 | 12/2009 | WO | N/A |
| WO 2010135242 | 12/2009 | WO | N/A |
| WO 2011022656 | 12/2010 | WO | N/A |
| WO 2011022656 | 12/2010 | WO | N/A |
| WO 2011041613 | 12/2010 | WO | N/A |
| WO 2011041613 | 12/2010 | WO | N/A |
| WO 2011119628 | 12/2010 | WO | N/A |
| WO 2011119628 | 12/2010 | WO | N/A |
| WO 2011154476 | 12/2010 | WO | N/A |
| WO 2012000188 | 12/2011 | WO | N/A |
| WO 2012000443 | 12/2011 | WO | N/A |
| WO 2012142529 | 12/2011 | WO | N/A |
| WO 2012142529 | 12/2011 | WO | N/A |
| WO 2013053775 | 12/2012 | WO | N/A |
| WO 2013112942 | 12/2012 | WO | N/A |
| WO 2013178344 | 12/2012 | WO | N/A |
| WO 2014047350 | 12/2013 | WO | N/A |
| WO 2014066527 | 12/2013 | WO | N/A |
| WO 2014066527 | 12/2013 | WO | N/A |
| WO 2014158811 | 12/2013 | WO | N/A |
| WO 2014170032 | 12/2013 | WO | N/A |
| WO 2015018528 | 12/2014 | WO | N/A |
| WO 2015018529 | 12/2014 | WO | N/A |

| | | | |
|----------------|---------|----|-----|
| WO 2015032755 | 12/2014 | WO | N/A |
| WO 2015127501 | 12/2014 | WO | N/A |
| WO 2015131994 | 12/2014 | WO | N/A |
| WO 2016018920 | 12/2015 | WO | N/A |
| WO 2016048903 | 12/2015 | WO | N/A |
| WO 2016094377 | 12/2015 | WO | N/A |
| WO 2017019894 | 12/2016 | WO | N/A |
| WO 2017019896 | 12/2016 | WO | N/A |
| WO 2017019897 | 12/2016 | WO | N/A |
| WO 2017062953 | 12/2016 | WO | N/A |
| WO 2017083291 | 12/2016 | WO | N/A |
| WO 2017118867 | 12/2016 | WO | N/A |
| WO 2017123981 | 12/2016 | WO | N/A |
| WO 2017190112 | 12/2016 | WO | N/A |
| WO 2018027316 | 12/2017 | WO | N/A |
| WO 2018209194 | 12/2017 | WO | N/A |
| WO 2018209194 | 12/2017 | WO | N/A |
| WO 2018218151 | 12/2017 | WO | N/A |
| WO 2019197275 | 12/2018 | WO | N/A |
| WO 2019209859 | 12/2018 | WO | N/A |
| WO 2020/012037 | 12/2019 | WO | N/A |
| WO 2020014591 | 12/2019 | WO | N/A |
| WO 2020037215 | 12/2019 | WO | N/A |
| WO 2020043835 | 12/2019 | WO | N/A |
| WO 2020079427 | 12/2019 | WO | N/A |

OTHER PUBLICATIONS

Martinez-Sobrido et al. (Journal of Virology. 2006; 80 (3): 1130-1139). cited by examiner
 SEQ ID No. 6 alignment with Geneseq db access No. AEH30839 Apr. 2006. cited by examiner
 SEQ ID No. 3 alignment with Geneseq db access No. AAZ44617 Apr. 2000. cited by examiner
 SEQ ID No. 10 alignment with Geneseq db access BBN47024 Nov. 2014. cited by examiner
 SEQ ID No. 9 alignment with GenEmbl db access AF295543 2001. cited by examiner
 SEQ ID No. 1 alignment with Geneseq db access AEH30838 2006. cited by examiner
 SEQ ID No. 17 UniProt db acc No. Q8B9N8_9MONO 2003. cited by examiner
 SEQ ID No. 16 alignment with GenEmbl db access AY145301 2003. cited by examiner
 SEQ ID No. 20 alignment with GenEmbl db access AY845400 2005. cited by examiner
 McGinnes et al. (Journal of Virology. 2011; 366-377). cited by examiner
 Kim et al. (Viruses. 2016; 8 (7): 183). cited by examiner
 Instant SEQ 21 alignment with Geneseq db access ARL59598 by Bu et al. in CN1869234-2008.
 cited by examiner
 Abolnik et al., 2012, "Full genomic sequence of an African avian paramyxovirus type 4 strain
 isolated from a wild duck," Virus Genes, 45(3):537-541. cited by applicant
 Afonso et al., 2016, "Taxonomy of the order Mononegavirales: update 2016," Arch Virol.,
 161(8):2351-2360. cited by applicant
 Aigner et al., "An effective tumor vaccine optimized for costimulation via bispecific and trispecific
 fusion proteins," Int. J. Oncol., 32(4): 777-789 (2008). cited by applicant
 Akhras et al., 2010, "Human metapneumovirus and respiratory syncytial virus: subtle differences
 but comparable severity," Infect. Dis. Rep., 2(2):e12 (5 pages). cited by applicant
 Alexander et al., 1978, "Relationship of parakeet/Netherlands/449/75 virus to other avian
 paramyxoviruses," Res. Vet. Sci., 25(1):105-106. cited by applicant

Alexander et al., 1981, "Characterization of viruses from doves representing a new serotype of avian paramyxoviruses," *Arch. Virol.*, 68(3-4):265-269. cited by applicant

Alexander, "Newcastle disease, Newcastle disease virus—an avian paramyxovirus," Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 1-22 (1988). cited by applicant

Al-Lazikani et al., "Standard conformations for the canonical structures of immunoglobulins," *J. Mol. Biol.*, 273(4):927-948 (1997). cited by applicant

Alper et al., 2009, "Rate of concurrent otitis media in upper respiratory tract infections with specific viruses," *Arch. Otolaryngol. Head Neck Surg.*, 135(1):17-21. cited by applicant

Altomonte et al., "Engineered newcastle disease virus as an improved oncolytic agent against hepatocellular carcinoma," *Mol. Ther.*, 18: 275-284 (2010). cited by applicant

Andral et al., 1984, "Isolation of avian paramyxovirus 2 and 3 from turkeys in Brittany," *Vet. Rec.*, 114(23):570-571. cited by applicant

Annels et al., "Oncolytic Immunotherapy for Bladder Cancer Using Coxsackie A21 Virus," *Mol. Ther. Oncolytics*, 9:1-12 (2018). cited by applicant

Assudani et al., "Immunotherapeutic potential of DISC-HSV and OX40L in cancer," *Cancer Immunol. Immunother.*, 55:104-111 (2006). cited by applicant

Ayers et al., "IFN- γ -related mRNA profile predicts clinical response to PD-1 blockade," *J. Clin. Invest.*, 127(8):2930-2940 (2017). cited by applicant

Ayllon et al., 2013, "Rescue of Recombinant Newcastle Disease Virus from cDNA," *J. Vis. Exp.*, (80):e50830. cited by applicant

Baca et al., "Antibody humanization using monovalent phage display," *J. Biol. Chem.*, 272(16):10678-10684 (1997). cited by applicant

Bai et al., 2002, "Dendritic cells pulsed with viral oncolysates potently stimulate autologous T cells from cancer patients," *Int. J. Oncol.*, 21(4):685-694. cited by applicant

Bankowski et al., 1981, "Effect of paramyxovirus yucaipa on fertility, hatchability, and poult yield of turkeys," *Avian. Dis.*, 25(2):517-520. cited by applicant

Barber et al. "Restoring function in exhausted CD8 T cells during chronic viral infection," *Nature* 439:682-687, with Supplemental Material attached (2006). cited by applicant

Bart et al., "Role of interferon in the anti-melanoma effects of poly (I).poly (C) and Newcastle disease virus," *Nat. New. Biol.*, 245:229-230 (1973). cited by applicant

Bauzon et al., 2014, "Armed therapeutic viruses—a disruptive therapy on the horizon of cancer immunotherapy," *Front. Immunol.*, 5(74):1-10. cited by applicant

Blackburn et al., "Tissue-specific differences in PD-1 and PD-L1 expression during chronic viral infection: implications for CD8 T-cell exhaustion," *J. Virol.*, 84(4):2078-2089 (2010) (Epub Dec. 2, 2009). cited by applicant

Blake et al., "Automated kinetic exclusion assays to quantify protein binding interactions in homogeneous solution," *Anal. Biochem.*, 272(2):123-134 (1999). cited by applicant

Bohle et al., 1990, "Postoperative active specific immunization in colorectal cancer patients with virus-modified autologous tumor-cell vaccine. First clinical results with tumor-cell vaccines modified with live but avirulent Newcastle disease virus," *Cancer*, 66(7):1517-1523. cited by applicant

Bohnsack et al., "Adaptation of the Immune-Related Response Criteria: irRECIST," *ESMO, Abstract 4958*, pp. 1-8 (2014). cited by applicant

Boisseau, 1993, "Basis for the evaluation of the microbiological risks due to veterinary drug residues in food," *Vet. Microbiol.*, 35(3-4):187-192. cited by applicant

Brahmer et al., "Phase I study of single-agent anti-programmed death-1 (MDX-1106) in refractory solid tumors: safety, clinical activity, pharmacodynamics, and immunologic correlates," *J. Clin. Oncol.*, 28(19):3167-3175 (2010). cited by applicant

Brahmer et al., "Safety and activity of anti-PD-L1 antibody in patients with advanced cancer," *N. Engl. J. Med.*, 366(26):2455-2465 (2012). cited by applicant

Briand et al., 2012, "Complete genome sequence of a novel avian paramyxovirus," *J Virol.*, 86(14):7710. cited by applicant

Brinkmann et al., "Phage display of disulfide-stabilized Fv fragments," *J. Immunol. Methods*, 182(1):41-50 (1995). cited by applicant

Brown et al., "Role of PD-1 in regulating acute infections," *Curr. Opin. Immunol.*, 22(3):397-401 (2010). cited by applicant

Bryant et al., "Development of intermediate-grade (mantle cell) and low-grade (small lymphocytic and marginal zone) human non-Hodgkin's lymphomas xenotransplanted in severe combined immunodeficiency mouse models," *Lab. Invest.*, 80:557-573 (2000). cited by applicant

Buchman et al., 2002, "Nasal and otologic effects of experimental respiratory syncytial virus infection in adults," *Am. J. Otolaryngol.*, 23(2):70-75. cited by applicant

Buijs et al., "Recombinant Immunomodulating Lentogenic or Mesogenic Oncolytic Newcastle Disease Virus for Treatment of Pancreatic Adenocarcinoma," *Viruses*, 7:2980-2998 (2015). cited by applicant

Burton et al., "Human antibodies from combinatorial libraries," *Adv. Immunol.*, 57:191-280 (1994). cited by applicant

Calain et al., "The Rule of Six, a Basic Feature for Efficient Replication of Sendai Virus Defective Interfering RNA," *J. Virol.*, 67(8):4822-4830 (1993). cited by applicant

Caldas et al., "Design and synthesis of germline-based hemi-humanized single-chain Fv against the CD18 surface antigen," *Protein Eng.*, 13(5):353-360 (2000). cited by applicant

Capua et al., 2004, "Isolation of an avian paramyxovirus type 9 from migratory waterfowl in Italy," *Vet. Rec.*, 155(5):156. cited by applicant

Car et al., "The Toxicology of Interleukin-12: A Review," *Toxicol. Pathol.*, 27(1):58-63 (1999). cited by applicant

Carnero et al., 2009, "Optimization of human immunodeficiency virus gag expression by newcastle disease virus vectors for the induction of potent immune responses," *J Virol.*, 83(2):584-597 (Epub Nov. 12, 2008). cited by applicant

Carpenter et al., "Non-Fc receptor-binding humanized anti-CD3 antibodies induce apoptosis of activated human T cells," *J. Immunol.*, 165(11):6205-6213 (2000). cited by applicant

Carthon et al., "Preoperative CTLA-4 blockade: tolerability and immune monitoring in the setting of a presurgical clinical trial," *Clin. Canc. Res.*, 16(10):2861-2871 (2010). cited by applicant

Caruso et al., "Adenovirus-mediated interleukin-12 gene therapy for metastatic colon carcinoma," *Proc. Natl. Acad. Sci. USA*, 93:11302-11306 (1996). cited by applicant

Cassel et al., 1965, "Newcastle Disease Virus as an Antineoplastic Agent," *Cancer*, 18:863-868. cited by applicant

Chaiwatpongsakorn et al., 2011, "Soluble respiratory syncytial virus fusion protein in the fully cleaved, pretriggered state is triggered by exposure to low-molarity buffer," *J. Virol.*, 85(8):3968-3977. cited by applicant

Chen et al., "CD4 T Cells Require ICOS-Mediated P13K Signaling to Increase T-Bet Expression in the Setting of Anti-CTLA-4 Therapy," *Cancer Immunol. Res.*, 2(2):167-176 (2013). cited by applicant

Cheng et al., "Genetic Modification of Oncolytic Newcastle Disease Virus for Cancer Therapy." *J. Virol.*, 90(1):5343-5352 (2016). cited by applicant

Cho et al., 2018, "Co-expression of the Hemagglutinin and Neuraminidase by Heterologous Newcastle Disease Virus Vectors Protected Chickens against H5 Clade 2.3.4.4 HPAI Viruses," *Sci. Rep.*, 8(1):16854 (9 pages). cited by applicant

Chothia et al., "Canonical structures for the hypervariable regions of immunoglobulins," *J. Mol. Biol.*, 196(4):901-917 (1987). cited by applicant

Chothia et al., "Conformations of immunoglobulin hypervariable regions," *Nature*, 342:877-883 (1989). cited by applicant

Chothia et al., "Structural repertoire of the human VH segments," *J. Mol. Biol.*, 227(3):799-817 (1992). cited by applicant

Chumakov et al., "Oncolytic Enteroviruses," *Mol. Biol. (Mosk)*, 46(5):639-650 (2012). cited by applicant

Clinical Trial NCT01295827, "Study of Pembrolizumab (MK-3475) in Participants With Progressive Locally Advanced or Metastatic Carcinoma, Melanoma, or Non-small Cell Lung Carcinoma (P07990/MK-3475-001/Keynote-001) (Keynote-001)," Merck Sharp & Dohme Corp., updated Sep. 13, 2018 (13 pages). cited by applicant

Cloud et al., 1980, "Characterization of Nine Avian Paramyxoviruses," *Avian Dis.*, 24(1):139-152. cited by applicant

Colbère-Garapin et al., "A new dominant hybrid selective marker for higher eukaryotic cells," *J. Mol. Biol.*, 150(1):1-14 (1981). cited by applicant

Copeland et al., 2005, "Functional chimeras of human immunodeficiency virus type 1 Gp120 and influenza A virus (H3) hemagglutinin," *J Virol.*, 79(10):6459-6471. cited by applicant

Crouse et al., "Expression and amplification of engineered mouse dihydrofolate reductase minigenes," *Mol. Cell Biol.*, 3(2):257-266 (1983). cited by applicant

Crowe, 2001, "Respiratory syncytial virus vaccine development," *Vaccine*, 20 Suppl 1:S32-S37. cited by applicant

Csatary et al., "MTH-68/H oncolytic viral treatment in human high-grade gliomas," *J. Neurooncol.*, 67:83-93 (2004). cited by applicant

Cuadrado-Castano et al., 2015, "Enhancement of the proapoptotic properties of newcastle disease virus promotes tumor remission in syngeneic murine cancer models," *Mol Cancer Ther.*, 14(5):1247-1258. cited by applicant

Cuadrado-Castano et al., 2015, "The therapeutic effect of death: Newcastle disease virus and its antitumor potential," *Virus Res.*, 209:56-66. cited by applicant

Cull et al., "Screening for receptor ligands using large libraries of peptides linked to the C terminus of the lac repressor," *Proc. Natl. Acad. Sci. USA*, 89(5):1865-1869 (1992). cited by applicant

Cuoto et al., "Anti-BA46 monoclonal antibody Mc3: humanization using a novel positional consensus and in vivo and in vitro characterization," *Cancer Res.*, 55(8):1717-1722 (1995). cited by applicant

Cuoto et al., "Designing human consensus antibodies with minimal positional templates," *Cancer Res.*, 55(23 Suppl):5973s-5977s (1995). cited by applicant

Curran et al., "Combination CTLA-4 blockade and 4-1BB activation enhances tumor rejection by increasing T-cell infiltration, proliferation, and cytokine production," *PLoS One*, 6(4):e19499 (2011). cited by applicant

Curran et al., "PD-1 and CTLA-4 combination blockade expands infiltrating T cells and reduces regulatory T and myeloid cells within B16 melanoma tumors," *Proc. Natl. Acad. Sci. USA*, 107(9):4275-4280 (2010). cited by applicant

Cwirla et al., "Peptides on phage: a vast library of peptides for identifying ligands," *Proc. Natl. Acad. Sci. USA*, 87(16):6378-6382 (1990). cited by applicant

De Leeuw et al., "Virulence of Newcastle disease virus is determined by the cleavage site of the fusion protein and by both the stem region and globular head of the haemagglutinin-neuraminidase protein," *J. Gen. Virol.*, 86(5): 1759-1769 (2005). cited by applicant

De Sousa Linares et al., "Not All Immune Checkpoints Are Created Equal," *Front. Immunol.*, 9:1909 (2018). cited by applicant

Dezfouli et al., "Enhancing CTL responses to melanoma cell vaccines in vivo: synergistic increases obtained using IFN γ primed and IFN β treated B7-1+ B16-F10 melanoma cells," *Immunol. Cell. Biol.*, 81(6):459-471 (2003). cited by applicant

Diamond et al., "Type I interferon is selectively required by dendritic cells for immune rejection of tumors," *J. Exp. Med.*, 208(10):1989-2003 (2011). cited by applicant

Dias et al., "Targeted cancer immunotherapy with oncolytic adenovirus coding for a fully human monoclonal antibody specific for CTLA-4," *Gen Ther.*, 19(10):988-998 (2012). cited by applicant

Dortmans et al., "Virulence of Newcastle disease virus: what is known so far?" *Vet. Res.*, 42:122 (2011). cited by applicant

Dortmans et al., 2012, "Newcastle disease virus outbreaks: vaccine mismatch or inadequate application?," *Vet. Microbiol.*, 160(1-2):17-22. cited by applicant

Douin-Echinard et al., "Enhancement of anti-tumor immunity by injection of fibroblasts genetically engineered to produce IL-12 and to express CD70," *Gene Therapy of Cancer*, edited by Walden et al., Plenum Press, New York, 353-357 (1998). cited by applicant

Doyle et al., 1927, "A Hitherto Unrecorded Disease of Fowls Due to a Filter-passing Virus," *J. Comp. Pathol. Ther.*, 40:144-169. cited by applicant

Dulos et al., 2012, "PD-1 blockade augments Th1 and Th17 and suppresses Th2 responses in peripheral blood from patients with prostate and advanced melanoma cancer," *J Immunother*, 35(2):169-178. cited by applicant

Dupraz et al., "Dominant negative MyD88 proteins inhibit interleukin-1 β /interferon- γ -mediated induction of nuclear factor κ B-dependent nitrite production and apoptosis in β cells," *J. Biol. Chem.*, 275:37672-37678 (2000). cited by applicant

Durbin et al., 1996, "Targeted disruption of the mouse Stat1 gene results in compromised innate immunity to viral disease," *Cell*, 84(3):443-450. cited by applicant

Eisenhauer et al., "New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1)," *Eur. J. Cancer*, 45:228-247 (2009). cited by applicant

Elankumaran et al., "Type I interferon-sensitive recombinant newcastle disease virus for oncolytic virotherapy," *J. Virol.*, 84:3835-3844 (2010). cited by applicant

Elmberg et al., 2017, "Potential disease transmission from wild geese and swans to livestock, poultry and humans: a review of the scientific literature from a One Health perspective," *Infect Ecol Epidemiol.*, 7(1):1300450 (21 pages). cited by applicant

Esper et al., 2003, "Human metapneumovirus infection in the United States: clinical manifestations associated with a newly emerging respiratory infection in children," *Pediatrics*, 111(6 Pt 1):1407-1410. cited by applicant

Fan et al., "Engagement of the ICOS pathway markedly enhances efficacy of CTLA-4 blockade in cancer immunotherapy," *J. Exp. Med.*, 211(4):715-725 (2014). cited by applicant

Fecci et al., "Systemic CTLA-4 blockade ameliorates glioma-induced changes to the CD4 $^{+}$ T cell compartment without affecting regulatory T-cell function," *Clin. Cancer Res.*, 13: 2158-2167 (2007). cited by applicant

Felici et al., "Selection of antibody ligands from a large library of oligopeptides expressed on a multivalent exposition vector," *J. Mol. Biol.*, 222(2):301-310 (1991). cited by applicant

Fields et al., "Adenoviridae and Their Replication," *Fundamental Virology*, 2nd Edition, Raven Press, Chapter 31, pp. 771-813 (1991). cited by applicant

Fiola et al., "Tumor selective replication of Newcastle disease virus: association with defects of tumor cells in antiviral defence," *Int. J. Cancer*, 119(2): 328-338 (2006). cited by applicant

Fisher et al., "IL-6 trans-signaling licenses mouse and human tumor microvascular gateways for trafficking of cytotoxic T cells," *J. Clin. Invest.*, 121(10):3846-3859 (2011). cited by applicant

Fodde et al., "Disease model: familial adenomatous polyposis," *Trends Mol. Med.*, 7:369-373 (2001). cited by applicant

Fodor et al., "Multiplexed biochemical assays with biological chips," *Nature*, 364(6437):555-556 (1993). cited by applicant

Foecking et al., "Powerful and versatile enhancer-promoter unit for mammalian expression vectors," *Gene*, 45(1):101-105 (1986). cited by applicant

Foote et al., "Antibody framework residues affecting the conformation of the hypervariable loops," *J. Mol. Biol.*, 224(2):487-499 (1992). cited by applicant

Fournier et al., "Oncolytic Newcastle Disease Virus as Cutting Edge between Tumor and Host," *Biology*, 2:936-975 (2013). cited by applicant

Fournier et al., 2012, "Analysis of three properties of Newcastle disease virus for fighting cancer: tumor-selective replication, antitumor cytotoxicity, and immunostimulation," *Methods Mol Biol.*, 797:177-204. cited by applicant

Foy et al., "Regulation of interferon regulatory factor-3 by the hepatitis C virus serine protease," *Science*, 300(5622):1145-1148 (2003). cited by applicant

Franciszkiewicz et al., "Role of chemokines and chemokine receptors in shaping the effector phase of the antitumor immune response," *Cancer Res.*, 72(24):6325-6332 (2012). cited by applicant

Fransen et al., "Controlled Local Delivery of CTLA-4 Blocking Antibody Induces CD8+ T-Cell-Dependent Tumor Eradication and Decreases Risk of Toxic Side Effects," *Clin. Cancer Res.*, 19(19):5381-5389 (2013). cited by applicant

Freeman et al., "Phase I/II trial of intravenous NDV-HUJ oncolytic virus in recurrent glioblastoma multiforme," *Mol. Ther.*, 13:221-228 (2006). cited by applicant

Fu et al., "The ICOS/ICOSL Pathway is Required for Optimal Antitumor Responses Mediated by Anti-CTLA-4 Therapy," *Cancer Res.*, 71:5445-5454 (2011). cited by applicant

Fuertes et al., "Host type I IFN signals are required for antitumor CD8+ T cell responses through CD8{alpha} + dendritic cells," *J. Exp. Med.*, 208(10):2005-2016 (2011). cited by applicant

Fujimoto et al., 2012, "Induction and maintenance of anti-influenza antigen-specific nasal secretory IgA levels and serum IgG levels after influenza infection in adults," *Influenza Other Respir. Viruses*, 6(6):396-403. cited by applicant

Galivo et al., "Interference of CD40L-Mediated Tumor Immunotherapy by Oncolytic Vesicular Stomatitis Virus," *Human Gene Therapy*, 21:439-450 (2010). cited by applicant

Gambotto et al., "Induction of antitumor immunity by direct intratumoral injection of a recombinant adenovirus vector expressing interleukin-12," *Cancer Gene Ther.*, 6(1):45-53 (1999). cited by applicant

Ganar et al., 2014, "Newcastle disease virus: current status and our understanding," *Virus Res.*, 184:71-81. cited by applicant

Gao et al., "Expression of transgenes from newcastle disease virus with a segmented genome," *J. Virol.*, 82(6): 2692-2698 (2008). cited by applicant

Garcia-Sastre et al., "Introduction of foreign sequences into the genome of influenza A virus," *Dev. Biol. Stand.*, 82:237-246 (1994). cited by applicant

Garcia-Sastre et al., "Use of a mammalian internal ribosomal entry site element for expression of a foreign protein by a transfectant influenza virus," *J. Virol.*, 68:6254-6261 (1994). cited by applicant

Garcia-Sastre et al., 1995, "Influenza virus vectors," *Biologicals*, 23(2):171-178. cited by applicant

Gardiner et al., "A Randomized, Double-Blind, Placebo-Controlled Assessment of BMS-936558, a Fully Human Monoclonal Antibody to Programmed Death-1 (PD-1), in Patients with Chronic Hepatitis C Virus Infection," *PLoS One*, 8(5):e63818 (2013). cited by applicant

Genbank Accession No. AAA52578.1, "GM-CSF [*Homo sapiens*]," Nov. 8, 1994. cited by applicant

Genbank Accession No. AAB19466.1, "granulocyte-macrophage colony-stimulating factor [*Canis lupus familiaris*]," May 7, 1993. cited by applicant

Genbank Accession No. AAB41697.1, "interleukin-15 [*Rattus norvegicus*]," Feb. 1, 1997. cited by applicant

Genbank Accession No. AAB60398.1, "interleukin-15 [*Macaca mulatta*]," Feb. 3, 1996. cited by applicant

Genbank Accession No. AAB94536.1, "interleukin-15 [*Rattus norvegicus*]," Jan. 2, 1998. cited by applicant

Genbank Accession No. AAC06041.1, "granulocyte-macrophage colony-stimulating factor [*Felis catus*]," Mar. 19, 1998. cited by applicant

Genbank Accession No. AAD00274.1, "CD70 [*Mus musculus*]," Jan. 5, 1999. cited by applicant

Genbank Accession No. AAG16626.1, "granulocyte-macrophage colony-stimulating factor [*Macaca mulatta*]," Dec. 23, 2002. cited by applicant

Genbank Accession No. AAH18149.1, "Interleukin 15 [*Homo sapiens*]," Jul. 15, 2006. cited by applicant

Genbank Accession No. AAH23698.1, "Interleukin 15 [*Mus musculus*]," Jul. 15, 2006. cited by applicant

Genbank Accession No. AAI00964.1, "Interleukin 15 [*Homo sapiens*]," Oct. 4, 2006. cited by applicant

Genbank Accession No. AAS67141.1, "fusion protein [Avian avulavirus 1]," Mar. 22, 2004. cited by applicant

Genbank Accession No. AAS67147.1, "fusion protein [Avian avulavirus 1]," Mar. 22, 2004. cited by applicant

Genbank Accession No. AAS67153.1, "fusion protein [Avian avulavirus 1]," Mar. 22, 2004. cited by applicant

Genbank Accession No. AAS67159.1, "fusion protein [Avian avulavirus 1]," Mar. 22, 2004. cited by applicant

Genbank Accession No. AAS67165.1, "fusion protein [Avian avulavirus 1]," Mar. 22, 2004. cited by applicant

Genbank Accession No. ACJ53752.1, "fusion protein [Avian avulavirus 1]," Nov. 25, 2008. cited by applicant

Genbank Accession No. ACJ53758.1, "fusion protein [Avian avulavirus 1]," Nov. 25, 2008. cited by applicant

Genbank Accession No. ACK57498.1, "fusion protein [Avian avulavirus 1]," Apr. 19, 2011. cited by applicant

Genbank Accession No. ADF59234.1, "fusion protein [Avian avulavirus 1]," Aug. 16, 2011. cited by applicant

Genbank Accession No. AF124443.1, "Newcastle disease virus isolate Roakin matrix protein mRNA, complete cds," Mar. 6, 2000. cited by applicant

Genbank Accession No. AF163440.1, "Newcastle disease virus fusion glycoprotein precursor, gene, complete cds," Jul. 6, 1999. cited by applicant

Genbank Accession No. AF309418.1, "Newcastle disease virus B1, complete genome," Dec. 2, 2000. cited by applicant

Genbank Accession No. AIA66858.1, "NBS-LRR resistance protein, partial [*Solanum viarum*]," Jun. 4, 2014. cited by applicant

Genbank Accession No. AIA66951.1, "fusion protein [Avian avulavirus 1]," Jun. 4, 2014. cited by applicant

Genbank Accession No. AIO82815.1, "PPIC-type PPIASE domain protein [*Burkholderia pseudomallei*]," Dec. 1, 2014. cited by applicant

Genbank Accession No. AIQ82815.1, "E7 [Human papillomavirus type 16]," Sep. 29, 2014. cited by applicant

Genbank Accession No. AKN79013.1, "E6 [Human papillomavirus type 16]," Jul. 4, 2015. cited by applicant

Genbank Accession No. AY143159.1, "Newcastle disease virus strain MET95 hemagglutinin-neuraminidase protein HN gene, complete cds," Jul. 25, 2003. cited by applicant

Genbank Accession No. AY351959.1, "Newcastle disease virus hemagglutinin-neuraminidase gene, complete cds," Aug. 25, 2003. cited by applicant

Genbank Accession No. AY390310.1, "Newcastle disease virus strain YG97 from goose fusion protein gene, partial cds," Jul. 26, 2016. cited by applicant

Genbank Accession No. AY845400.2, "Newcastle disease virus strain LaSota, complete genome,"

Mar. 17, 2005., cited by applicant

Genbank Accession No. CAA26820.1, "GM-CSF [*Mus musculus*]," Sep. 24, 2008. cited by applicant

Genbank Accession No. CAA62616.1, "interleukin-15 [*Homo sapiens*]," Nov. 14, 2006. cited by applicant

Genbank Accession No. CAG46642.1, "CD86, partial [*Homo sapiens*]," Jul. 26, 2016. cited by applicant

Genbank Accession No. CAI41082.1, "interleukin 15 receptor, alpha [*Homo sapiens*]," Jan. 13, 2009. cited by applicant

Genbank Accession No. EF033114.1, "Macaca mulatta interleukin 15 receptor alpha (IL15Ra) mRNA, complete cds, alternatively spliced," Nov. 12, 2006. cited by applicant

Genbank Accession No. EU293914.1, "Newcastle disease virus strain Italien, complete genome," Jun. 24, 2008. cited by applicant

Genbank Accession No. EU338414.1, "Avian paramyxovirus 2 strain APMV-2/Chicken/California/Yucaipa/56, complete genome," Sept. 3, 2008. cited by applicant

Genbank Accession No. EU622637.2, "Avian paramyxovirus 6 strain APMV-6/duck/HongKong/18/199/77, complete genome," Feb. 25, 2011. cited by applicant

Genbank Accession No. EU782025.1, "Avian paramyxovirus 3 strain turkey/Wisconsin/68, complete genome," Mar. 26, 2010. cited by applicant

Genbank Accession No. FJ177514.1, "Avian paramyxovirus 4 strain APMV-4/duck/Hongkong/D3/75, complete genome," Nov. 4, 2008. cited by applicant

Genbank Accession No. FJ231524.1, "Avian paramyxovirus 7 strain APMV-7/dove/Tennessee/4/75, complete genome," Sep. 10, 2009. cited by applicant

Genbank Accession No. FJ619036.1, "Avian paramyxovirus 8 isolate APMV-8/Goose/Delaware/1053/76, complete genome," Jun. 1, 2009. cited by applicant

Genbank Accession No. JF950510.1, Newcastle disease virus strain LaSota, complete genome, Aug. 10, 2011. cited by applicant

Genbank Accession No. JN571485.1, "Avian paramyxovirus 4 strain APMV4/mallard/Belgium/15129/07, complete genome," Dec. 8, 2011. cited by applicant

Genbank Accession No. JX133079.1, "Avian paramyxovirus 4 isolate APMV-4/Egyptian goose/South Africa/N1468/2010, complete genome," Apr. 4, 2013. cited by applicant

Genbank Accession No. JX987283.1, "Avian paramyxovirus 4 strain APMV-4/duck/Delaware/549227/2010, complete genome," Mar. 12, 2013. cited by applicant

Genabnk Accession No. KC439346.1, "Avian paramyxovirus 4 strain APMV4/duck/China/G302/2012, complete genome," Apr. 19, 2013. cited by applicant

Genbank Accession No. KM058635.1, "Human papillomavirus type 16 isolate IR-32 E7 (E7) gene, complete cds," Sep. 29, 2014. cited by applicant

Genbank Accession No. KP677555.1, "Human papillomavirus type 16 isolate Af2-ZOAASSOUMOU3 E6 (E6) gene, complete cds," Jul. 4, 2015. cited by applicant

Genbank Accession No. KU601399.1, "Avian paramyxovirus 4 isolate Uria_aalge/Russia/Tyuleny Island/115/2015, complete genome," Jul. 30, 2016. cited by applicant

Genbank Accession No. M11220.1, "Human granulocyte-macrophage colony stimulating factor (GM-CSF) mRNA," Nov. 8, 1994. cited by applicant

Genbank Accession No. NC_002617.1, "Newcastle disease virus B1, complete genome," Nov. 30, 2009. cited by applicant

Genbank Accession No. NC_025390.1, "Avian paramyxovirus 9 strain duck/New York/22/1978, complete genome," Aug. 13, 2018. cited by applicant

Genbank Accession No. NG_016779.1, "*Homo sapiens* interleukin 2 (IL2), RefSeqGene on chromosome 4," Dec. 4, 2019. cited by applicant

Genbank Accession No. NM_000402.2, "*Homo sapiens* glucose-6-phosphate dehydrogenase

(G6PD), nuclear gene encoding mitochondrial protein, mRNA,” May 7, 2006. cited by applicant
Genbank Accession No. NM_000585.4, “*Homo sapiens* interleukin 15 (IL15), transcript variant 3, mRNA,” May 6, 2017. cited by applicant
Genbank Accession No. NM_000586.3, “*Homo sapiens* interleukin 2 (IL2), mRNA,” Apr. 30, 2017. cited by applicant
Genbank Accession No. NM_000590.1, “*Homo sapiens* interleukin 9 (IL9), mRNA,” May 6, 2017. cited by applicant
Genbank Accession No. NM_000594.3, “*Homo sapiens* tumor necrosis factor (TNF), mRNA,” May 7, 2017. cited by applicant
Genbank Accession No. NM_000619.2, “*Homo sapiens* interferon gamma (IFNG), mRNA,” May 13, 2017. cited by applicant
Genbank Accession No. NM_000758.3, “*Homo sapiens* colony stimulating factor 2 (CSF2), mRNA,” Oct. 13, 2018. cited by applicant
Genbank Accession No. NM_000758.4, “*Homo sapiens* colony stimulating factor 2 (CSF2), mRNA,” Feb. 16, 2020. cited by applicant
Genbank Accession No. NM_000880.3, “*Homo sapiens* interleukin 7 (IL7), transcript variant 1, mRNA,” May 10, 2017. cited by applicant
Genbank Accession No. NM_000882.3, “*Homo sapiens* interleukin 12A (IL12A), mRNA,” Apr. 30, 2017. cited by applicant
Genbank Accession No. NM_000937.2, “*Homo sapiens* polymerase (RNA) II (DNA directed) polypeptide A, 220kDa (POLR2A), mRNA,” Feb. 11, 2008. cited by applicant
Genbank Accession No. NM_001024736.1, “*Homo sapiens* CD276 molecule (CD276), transcript variant 1, mRNA,” May 7, 2017. cited by applicant
Genbank Accession No. NM_001090.2, “*Homo sapiens* ATP binding cassette subfamily F member 1 (ABCF1), transcript variant 2, mRNA,” Apr. 27, 2017. cited by applicant
Genbank Accession No. NM_001099270.1, “*Homo sapiens* zinc finger and BTB domain containing 34 (ZBTB34), mRNA,” Apr. 30, 2017. cited by applicant
Genbank Accession No. NM_001172085.1, “*Homo sapiens* TATA-box binding protein (TBP), transcript variant 2, mRNA,” Apr. 17, 2017. cited by applicant
Genbank Accession No. NM_001199886.1, “*Homo sapiens* interleukin 7 (IL7), transcript variant 2, mRNA,” May 10, 2017. cited by applicant
Genbank Accession No. NM_001199887.1, “*Homo sapiens* interleukin 7 (IL7), transcript variant 3, mRNA,” May 10, 2017. cited by applicant
Genbank Accession No. NM_001199888.1, “*Homo sapiens* interleukin 7 (IL7), transcript variant 4, mRNA,” May 10, 2017. cited by applicant
Genbank Accession No. NM_001207006.2, “*Homo sapiens* interleukin 21 (IL21), transcript variant 2, mRNA,” May 7, 2017. cited by applicant
Genbank Accession No. NM_001242.4, “*Homo sapiens* CD27 molecule (CD27), mRNA,” Apr. 28, 2017. cited by applicant
Genbank Accession No. NM_001252.3, “*Homo sapiens* CD70 molecule (CD70), mRNA,” Jan. 11, 2014. cited by applicant
Genbank Accession No. NM_001267706.1, “*Homo sapiens* CD274 molecule (CD274), transcript variant 2, mRNA,” Apr. 23, 2017. cited by applicant
Genbank Accession No. NM_001314029.1, “*Homo sapiens* CD274 molecule (CD274), transcript variant 4, mRNA,” Apr. 23, 2017. cited by applicant
Genbank Accession No. NM_001768.5, “*Homo sapiens* CD8a molecule (CD8A), transcript variant 1, mRNA,” Sep. 28, 2008. cited by applicant
Genbank Accession No. NM_002122.3, “*Homo sapiens* major histocompatibility complex, class II, DQ alpha 1 (HLA-DQA1), mRNA,” May 7, 2017. cited by applicant
Genbank Accession No. NM_002124.1, “*Homo sapiens* major histocompatibility complex, class II,

DR beta 1 (HLA-DRB1), mRNA," Jul. 31, 2007. cited by applicant

Genbank Accession No. NM_002164.3, "*Homo sapiens* indoleamine-pyrrole 2,3 dioxygenase (INDO), mRNA," Aug. 6, 2007. cited by applicant

Genbank Accession No. NM_002187.2, "*Homo sapiens* interleukin 12B (IL12B), mRNA," May 6, 2017. cited by applicant

Genbank Accession No. NM_002189.4, "*Homo sapiens* interleukin 15 receptor subunit alpha (IL15RA), transcript variant 1, mRNA," Jan. 2, 2020. cited by applicant

Genbank Accession No. NM_002190.2, "*Homo sapiens* interleukin 17A (IL17A), mRNA," May 7, 2017. cited by applicant

Genbank Accession No. NM_002286.5, "*Homo sapiens* lymphocyte activating 3 (LAG3), mRNA," May 7, 2017. cited by applicant

Genbank Accession No. NM_002416.1, "*Homo sapiens* chemokine (C-X-C motif) ligand 9 (CXCL9), mRNA," Sep. 25, 2014. cited by applicant

Genbank Accession No. NM_002801.2, "*Homo sapiens* proteasome (prosome, macropain) subunit, beta type, 10 (PSMB10), mRNA," Feb. 10, 2011. cited by applicant

Genbank Accession No. NM_002985.2, "*Homo sapiens* C-C motif chemokine ligand 5 (CCL5), transcript variant 1, mRNA," May 6, 2017. cited by applicant

Genbank Accession No. NM_003326.3, "*Homo sapiens* tumor necrosis factor (ligand) superfamily, member 4 (TNFSF4), mRNA," May 4, 2014. cited by applicant

Genbank Accession No. NM_003811.3, "*Homo sapiens* TNF superfamily member 9 (TNFSF9), mRNA," Mar. 4, 2017. cited by applicant

Genbank Accession No. NM_004072.1, "*Homo sapiens* chemokine-like receptor 1 (CMKLR1), mRNA," Jul. 13, 2008. cited by applicant

Genbank Accession No. NM_004152.2, "*Homo sapiens* ornithine decarboxylase antizyme 1 (OAZ1), mRNA," Jul. 4, 2014. cited by applicant

Genbank Accession No. NM_004168.1, "*Homo sapiens* succinate dehydrogenase complex, subunit A, flavoprotein (Fp) (SDHA), nuclear gene encoding mitochondrial protein, mRNA," Jun. 3, 2007. cited by applicant

Genbank Accession No. NM_005018.2, "*Homo sapiens* programmed cell death 1 (PDCD1), mRNA," May 6, 2017. cited by applicant

Genbank Accession No. NM_005191.3, "*Homo sapiens* CD80 molecule (CD80), mRNA," May 6, 2017. cited by applicant

Genbank Accession No. NM_005516.4, "*Homo sapiens* major histocompatibility complex, class I, E (HLA-E), mRNA," Jul. 5, 2010. cited by applicant

Genbank Accession No. NM_005601.3, "*Homo sapiens* natural killer cell granule protein 7 (NKG7), mRNA," Apr. 26, 2017. cited by applicant

Genbank Accession No. NM_006564.1, "*Homo sapiens* C-X-C motif chemokine receptor 6 (CXCR6), mRNA," Apr. 25, 2017. cited by applicant

Genbank Accession No. NM_007315.2, "*Homo sapiens* signal transducer and activator of transcription 1, 91kDa (STAT1), transcript variant alpha, mRNA," Apr. 29, 2008. cited by applicant

Genbank Accession No. NM_008357.2, "*Mus musculus* interleukin 15 (I115), transcript variant 1, mRNA," Dec. 28, 2019. cited by applicant

Genbank Accession No. NM_008358.2, "*Mus musculus* interleukin 15 receptor, alpha chain (I115ra), transcript variant 1, mRNA," Jan. 12, 2020. cited by applicant

Genbank Accession No. NM_009404.3, "*Mus musculus* tumor necrosis factor (ligand) superfamily, member 9 (Tnfsf9), mRNA," Apr. 29, 2017. cited by applicant

Genbank Accession No. NM_009452.2, "*Mus musculus* tumor necrosis factor (ligand) superfamily, member 4 (Tnfsf4), mRNA," May 12, 2017. cited by applicant

Genbank Accession No. NM_009855.2, "*Mus musculus* CD80 antigen (Cd80), mRNA," Apr. 28, 2017. cited by applicant

Genbank Accession No. NM_009969.4, "Mus musculus colony stimulating factor 2 (granulocyte-macrophage) (Csf2), mRNA," Jan. 27, 2020. cited by applicant

Genbank Accession No. NM_011617.2, "Mus musculus CD70 antigen (Cd70), mRNA," Sept. 1, 2016. cited by applicant

Genbank Accession No. NM_014143.3, "*Homo sapiens* CD274 molecule (CD274), transcript variant 1, mRNA," Apr. 30, 2017. cited by applicant

Genbank Accession No. NM_015259.4, "*Homo sapiens* inducible T-cell co-stimulator ligand (ICOSLG), mRNA," Sept. 2, 2013. cited by applicant

Genbank Accession No. NM_015527.3, "*Homo sapiens* TBC1 domain family member 10B (TBC1D10B), mRNA," May 2, 2017. cited by applicant

Genbank Accession No. NM_015790.3, "Mus musculus icos ligand (ICOSL), mRNA," Feb. 15, 2015. cited by applicant

Genbank Accession No. NM_017970.3, "*Homo sapiens* NRDE-2, necessary for RNA interference, domain containing (NRDE2), mRNA," May 10, 2017. cited by applicant

Genbank Accession No. NM_018955.2, "*Homo sapiens* ubiquitin B (Ubb), mRNA," Apr. 17, 2013. cited by applicant

Genbank Accession No. NM_019388.3, "Mus musculus CD86 antigen (Cd86), mRNA," May 2, 2017. cited by applicant

Genbank Accession No. NM_020525.4, "*Homo sapiens* interleukin 22 (IL22), mRNA," Apr. 26, 2017. cited by applicant

Genbank Accession No. NM_021803.3, "*Homo sapiens* interleukin 21 (IL21), transcript variant 1, mRNA," May 7, 2017. cited by applicant

Genbank Accession No. NM_025239.3, "*Homo sapiens* programmed cell death 1 ligand 2 (PDCDILG2), mRNA," May 3, 2017. cited by applicant

Genbank Accession No. NM_052902.2, "*Homo sapiens* serine/threonine kinase 11 interacting protein (STK11IP), mRNA," Mar. 22, 2015. cited by applicant

Genbank Accession No. NM_053852.1, "Rattus norvegicus colony stimulating factor 2 (Csf2), mRNA," Dec. 28, 2019. cited by applicant

Genbank Accession No. NM_172175.2, "*Homo sapiens* interleukin 15 (IL15), transcript variant 2, mRNA," May 6, 2017. cited by applicant

Genbank Accession No. NM_173799.2, "*Homo sapiens* T cell immunoreceptor with Ig and ITIM domains (TIGIT), mRNA," Jun. 18, 2009. cited by applicant

Genbank Accession No. NP_000576.1, "interleukin-15 isoform 1 preproprotein [*Homo sapiens*]," Feb. 9, 2020. cited by applicant

Genbank Accession No. NP_000577.2, "interleukin-2 precursor [*Homo sapiens*]," Feb. 3, 2020. cited by applicant

Genbank Accession No. NP_000749.2, "granulocyte-macrophage colony-stimulating factor precursor [*Homo sapiens*]," Feb. 16, 2020. cited by applicant

Genbank Accession No. NP_001009207.1, "interleukin-15 precursor [*Felis catus*]," Dec. 25, 2019. cited by applicant

Genbank Accession No. NP_001243.1, "CD70 antigen isoform 1 [*Homo sapiens*]," Oct. 8, 2016. cited by applicant

Genbank Accession No. NP_001254635.1, "programmed cell death 1 ligand 1 isoform b precursor [*Homo sapiens*]," Apr. 23, 2017. cited by applicant

Genbank Accession No. NP_001300958.1, "programmed cell death 1 ligand 1 isoform c precursor [*Homo sapiens*]," Apr. 23, 2017. cited by applicant

Genbank Accession No. NP_002178.2, "interleukin-12 subunit beta precursor [*Homo sapiens*]," May 6, 2017. cited by applicant

Genbank Accession No. NP_002180.1, "interleukin-15 receptor subunit alpha isoform 1 precursor [*Homo sapiens*]," Jan. 2, 2020. cited by applicant

Genbank Accession No. NP_003317.1, "tumor necrosis factor ligand superfamily member 4 isoform 1 [*Homo sapiens*]," Jul. 4, 2014. cited by applicant

Genbank Accession No. NP_003802.1, "tumor necrosis factor ligand superfamily member 9 [*Homo sapiens*]," Mar. 4, 2017. cited by applicant

Genbank Accession No. NP_005009.2, "programmed cell death protein 1 precursor [*Homo sapiens*]," May 6, 2017. cited by applicant

Genbank Accession No. NP_005182.1, "T-lymphocyte activation antigen CD80 precursor [*Homo sapiens*]," Apr. 30, 2017. cited by applicant

Genbank Accession No. NP_032383.1, "interleukin-15 preproprotein [*Mus musculus*]," Dec. 28, 2019. cited by applicant

Genbank Accession No. NP_032384.1, "interleukin-15 receptor subunit alpha isoform 1 precursor [*Mus musculus*]," Jan. 12, 2020. cited by applicant

Genbank Accession No. NP_033430.1, "tumor necrosis factor ligand superfamily member 9 [*Mus musculus*]," Apr. 29, 2017. cited by applicant

Genbank Accession No. NP_033478.1, "tumor necrosis factor ligand superfamily member 4 [*Mus musculus*]," May 12, 2017. cited by applicant

Genbank Accession No. NP_033985.3, "T-lymphocyte activation antigen CD80 precursor [*Mus musculus*]," Apr. 28, 2017. cited by applicant

Genbank Accession No. NP_034099.2, "granulocyte-macrophage colony-stimulating factor precursor [*Mus musculus*]," Jan. 27, 2020. cited by applicant

Genbank Accession No. NP_054862.1, "programmed cell death 1 ligand 1 isoform a precursor [*Homo sapiens*]," Apr. 30, 2017. cited by applicant

Genbank Accession No. NP_056074.1, "ICOS ligand isoform a precursor [*Homo sapiens*]," Apr. 23, 2017. cited by applicant

Genbank Accession No. NP_056605.1, "ICOS ligand precursor [*Mus musculus*]," Feb. 15, 2015. cited by applicant

Genbank Accession No. NP_056606.1, "F-box only protein 8 [*Mus musculus*]," Apr. 8, 2003. cited by applicant

Genbank Accession No. NP_062261.3, "T-lymphocyte activation antigen CD86 precursor [*Mus musculus*]," May 2, 2017. cited by applicant

Genbank Accession No. NP_079515.2, "programmed cell death 1 ligand 2 precursor [*Homo sapiens*]," May 3, 2017. cited by applicant

Genbank Accession No. NP_446304.1, "granulocyte-macrophage colony-stimulating factor precursor [*Rattus norvegicus*]," Dec. 28, 2019. cited by applicant

Genbank Accession No. U25837.1, "Newcastle disease virus isolate Ulster matrix protein mRNA, complete cds," Jul. 12, 1996. cited by applicant

Genbank Accession No. X03021.1, "Human gene for granulocyte-macrophage colony stimulating factor (GM-CSF)," Feb. 10, 1999. cited by applicant

Genbank Accession No., NP_000873.2, "interleukin-12 subunit alpha precursor [*Homo sapiens*]," Apr. 30, 2017. cited by applicant

Ghaneh et al., "Adenovirus-mediated transfer of p53 and p16(INK4a) results in pancreatic cancer regression in vitro and in vivo," *Gene Ther.*, 8:199-208 (2001). cited by applicant

Gillies et al., "High-level expression of chimeric antibodies using adapted cDNA variable region cassettes," *J. Immunol. Methods*, 125(1-2):191-202 (1989). cited by applicant

Ginting et al., 2017, "Proinflammatory response induced by Newcastle disease virus in tumor and normal cells," *Oncolytic Virother*, 6:21-30. cited by applicant

Gitiban et al., 2005, "Chinchilla and murine models of upper respiratory tract infections with respiratory syncytial virus," *J. Virol.*, 79(10):6035-6042. cited by applicant

Goff et al., "A Majority of Infectious Newcastle Disease Virus Particles Contain a Single Genome, while a Minority Contain Multiple Genomes," *J. Virol.*, 86(19):10852-10856 (2012). cited by

applicant

Gogoi et al., 2017, "Avian Paramyxovirus: A Brief Review," *Transbound. Emerg. Dis.*, 64(1):53-67 (Epub 2015). cited by applicant

Gomez et al., 2009, "Phase-I study MEDI-534, of a live, attenuated intranasal vaccine against respiratory syncytial virus and parainfluenza-3 virus in seropositive children," *Pediatr. Infect. Dis. J.*, 28(7):655-658. cited by applicant

Grieves et al., 2015, "Acute and Chronic Airway Disease After Human Respiratory Syncytial Virus Infection in Cotton Rats (*Sigmodon hispidus*)," *Comp. Med.*, 65(4):315-326. cited by applicant

Gruber et al., "Efficient tumor cell lysis mediated by a bispecific single chain antibody expressed in *Escherichia coli*," *J. Immunol.*, 152(11):5368-5374 (1994). cited by applicant

Grund et al., 2014, "Avian paramyxovirus-8 immunization reduces viral shedding after homologous APMV-8 challenge but fails to protect against Newcastle disease," *Virol. J.*, 11:179 (6 pages). cited by applicant

Guo et al., "Oncolytic immunotherapy: dying the right way is a key to eliciting potent antitumor immunity," *Front. Oncol.*, 4(74):1-11 (2014). cited by applicant

Haas et al., "Bispecific antibodies increase T-cell stimulatory capacity in vitro of human autologous virus-modified tumor vaccine," *Clin. Cancer Res.*, 4(3):721-730 (1998). cited by applicant

Haas et al., "A tumor vaccine containing ant-CD# and anti-CD28 bispecific antibodies triggers strong and durable antitumor activity in human lymphocytes," *Int. J. Cancer*, 188(3): 658-667 (2006). cited by applicant

Haas et al., "An effective strategy of human tumor vaccine modification by coupling bispecific costimulatory molecules," *Cancer Gene Therapy*, 6(3):254-262 (1999). cited by applicant

Habibi et al., 2015, "Impaired Antibody-mediated Protection and Defective IgA B-Cell Memory in Experimental Infection of Adults with Respiratory Syncytial Virus," *Am. J. Respir. Crit. Care Med.*, 191(9):1040-1049. cited by applicant

Hall et al., 1991, "Immunity to and frequency of reinfection with respiratory syncytial virus," *J. Infect. Dis.*, 163(4):693-698. cited by applicant

Hall et al., 2001, "Respiratory syncytial virus infections in previously healthy working adults," *Clin. Infect. Dis.*, 33(6):792-796. cited by applicant

Hallak et al., 2000, "Iduronic acid-containing glycosaminoglycans on target cells are required for efficient respiratory syncytial virus infection," *Virology*, 271(2):264-275. cited by applicant

Hamid et al., "Safety and Tumor Responses with Lambrolizumab (Anti-PD-1) in Melanoma," *N. Engl. J. Med.*, 369(2):134-144 (2013). cited by applicant

Hamid et al., 2016, "1050PD—Combination of MEDI0680, an anti-PD-1 antibody, with durvalumab, an anti-PD-L1 antibody: A phase 1, open-label study in advanced malignancies," *Ann. Oncol.*, 27(suppl. 6):vi359-vi378 (1 page). cited by applicant

He et al., "Humanization and pharmacokinetics of a monoclonal antibody with specificity for both E- and P-selectin," *J. Immunol.*, 160(2):1029-1035 (1998). cited by applicant

Heery et al., "Phase I open-label, multiple ascending dose trial of MSB0010718C, an anti-PD-L1 monoclonal antibody, in advanced solid malignancies," *J. Clin. Oncol.*, 32(suppl. 5S):abstract 3064 (2014). cited by applicant

Heicappell et al., 1986, "Prevention of metastatic spread by postoperative immunotherapy with virally modified autologous tumor cells. I. Parameters for optimal therapeutic effects," *Int. J. Cancer*, 37(4):569-577. cited by applicant

Hemminki et al., "Oncolytic Immunotherapy: Where Are We Clinically?," *Scientifica*, 2014, Article ID 862925 (2014). cited by applicant

Herber et al., "Squamous epithelial hyperplasia and carcinoma in mice transgenic for the human papillomavirus type 16 E7 oncogene," *J. Virol.*, 70:1873-1881 (1996). cited by applicant

Herbst et al., "Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients," *Nature*, 515(7528):563-567 (2014). cited by applicant

Herbst et al., "A study of MPDL3280A, an engineered PD-L1 antibody in patients with locally advanced or metastatic tumors," J. Clin. Oncol., 31(suppl):abstract 3000 (2013). cited by applicant

Hines et al., 2012, "Avian paramyxovirus serotype-1: a review of disease distribution, clinical symptoms, and laboratory diagnostics," Vet Med Int., 2012:708216 (17 pages). cited by applicant

Hirano et al., "Blockade of B7-H1 and PD-1 by Monoclonal Antibodies Potentiates Cancer Therapeutic Immunity," Cancer Res., 65(3):1089-1096 (2005). cited by applicant

Hirschhorn-Cymerman et al., "Induction of tumoricidal function in CD4+ T cells is associated with concomitant memory and terminally differentiated phenotype," J. Exp. Med., 209(11):2113-2126 (2012). cited by applicant

Hofmeyer et al., "The PD-1/PD-L1 (B7-H1) Pathway in Chronic Infection-Induced Cytotoxic T Lymphocyte Exhaustion," J. Biomed. Biotechnol., 2011:451694 (2011). cited by applicant

Holliger et al., "Diabodies": small bivalent and bispecific antibody fragments," Proc. Natl. Acad. Sci. USA, 90(14):6444-6448 (1993). cited by applicant

Hollinger et al., "Engineered antibody fragments and the rise of single domains," Nature Biotech., 23(9):1126-1136 (2005). cited by applicant

Hosokawa et al., "In vivo analysis of mammary and non-mammary tumorigenesis in MMTV-cyclin D1 transgenic mice deficient in p53," Transgenic Res., 10:471-478 (2001). cited by applicant

Hotte et al., "An optimized clinical regimen for the oncolytic virus PV701," Clin. Cancer Res., 13:977-985 (2007). cited by applicant

Hou et al., 2009, "Study on the effect of Newcastle disease virus vaccine and interleukin-12 to the transplatable nude mice model of human ovarian cancer," Chin. J. Cancer Prev. Treat., 16(18):1375-1378 (in Chinese with English abstract). cited by applicant

Houdebine, "Production of Pharmaceutical Proteins by Transgenic Animals." Comp. Immunol. Microbiol. Infect. Dis., 32(2):107-121 (2009). cited by applicant

Hough et al., "A model for spontaneous B-lineage lymphomas in IgHmu-HOX11 transgenic mice," Proc. Natl. Acad. Sci. USA, 95:13853-13858 (1998). cited by applicant

Houghten et al. "The use of synthetic peptide combinatorial libraries for the identification of bioactive peptides," Biotechniques, 13(3):412-421 (1992). cited by applicant

Huang et al., "Newcastle disease virus V protein is associated with viral pathogenesis and functions as an alpha interferon antagonist," J. Virol., 77:8676-8685 (2003). cited by applicant

Huang et al., "Preclinical validation: LV/IL-12 transduction of patient leukemia cells for immunotherapy of AML," Mol. Ther., 3:16074 (2016). cited by applicant

Huard et al., "CD4/major histocompatibility complex class II interaction analyzed with CD4- and lymphocyte activation gene-3 (LAG-3)-Ig fusion proteins," Eur. J. Immunol., 25:2718-2721 (1995). cited by applicant

Infante et al., "Clinical and pharmacodynamic (PD) results of a phase I trial with AMP-224 (B7-DC Fc) that binds to the PD-1 receptor," J. Clin. Oncol., 31(suppl):abstract 3044 (2013). cited by applicant

Inouye et al., "Up-promoter mutations in the lpp gene of *Escherichia coli*," Nucleic Acids Res., 13(9):3101-3110 (1985). cited by applicant

International Nonproprietary Names for Pharmaceutical Substances (INN), 2013, WHO Drug Information, vol. 27, No. 2, List 109, pp. 135-209. cited by applicant

International Search Report and Written Opinion for International Patent Application No. PCT/US2019/041568 (Pub No. WO 202001459) mailed Nov. 20, 2019 (12 pages). cited by applicant

International Search Report and Written Opinion for International Patent Application No. PCT/US2019/046837 (Pub. No. WO 2020037215) mailed Jan. 2, 2020 (13 pages). cited by applicant

Ishida et al., "Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death," Embo. J., 11:3887-3895 (1992). cited by applicant

Iwai et al., "PD-1 Inhibits Antiviral Immunity at the Effector Phase in the Liver," *J. Exp. Med.*, 198(1):39-50 (2003). cited by applicant

Iwai et al., "Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade," *Proc. Natl. Acad. Sci USA*, 99:12293-12297 (2002). cited by applicant

Jarahian et al., 2009, "Activation of natural killer cells by newcastle disease virus hemagglutinin-neuraminidase," *J Virol.*, 83(16):8108-8121. cited by applicant

Jones et al., "Molecular interactions within the IL-6/IL-12 cytokine/receptor superfamily," *Immunol. Res.*, 51(1):5-14 (2011). cited by applicant

Kabat et al., "Attempts to locate complementarity-determining residues in the variable positions of light and heavy chains," *Ann. NY Acad. Sci.*, 190:382-393 (1971). cited by applicant

Kado et al., "Intestinal Microflora Are Necessary for Development of Spontaneous Adenocarcinoma of the Large Intestine in T-Cell Receptor β Chain and p53 Double-Knockout Mice," *Cancer Res.*, 61:2395-2398 (2001). cited by applicant

Kamphorst et al., "Rescue of exhausted CD8 T cells by PD-1-targeted therapies is CD28-dependent," *Science*, 355(6332):1423-1427 (2017). cited by applicant

Kapczynski et al., 2013, "Immune responses of poultry to Newcastle disease virus," *Dev Comp Immunol.*, 41(3):447-453. cited by applicant

Karamendin et al., 2016, "Complete Genome Sequence of a Novel Avian Paramyxovirus (APMV-13) Isolated from a Wild Bird in Kazakhstan," *Genome Announc.*, 4(3). pii:e00167-16 (2 pages). cited by applicant

Karcher et al., 2004, "Antitumor vaccination in patients with head and neck squamous cell carcinomas with autologous virus-modified tumor cells," *Cancer Res.*, 64(21):8057-8061. cited by applicant

Kato et al., "Cell type-specific involvement of RIG-I in antiviral response," *Immunity*, 23(1):19-28 (2005). cited by applicant

Kettleborough et al., "Isolation of tumor cell-specific single-chain Fv from immunized mice using phage-antibody libraries and the re-construction of whole antibodies from these antibody fragments," *Eur. J. Immunol.*, 24:952-958 (1994). cited by applicant

KEYTRUDA® (pembrolizumab) Prescribing Information, revised Aug. 2018 (52 pages). cited by applicant

KEYTRUDA® (pembrolizumab) Prescribing Information, revised Dec. 2015 (25 pages). cited by applicant

KEYTRUDA® (pembrolizumab) Prescribing Information, revised Jan. 2020 (85 pages). cited by applicant

KEYTRUDA® (pembrolizumab) Prescribing Information, revised Jun. 2018 (53 pages). cited by applicant

KEYTRUDA® (pembrolizumab) Prescribing Information, revised Oct. 2016 (26 pages). cited by applicant

Khattar et al., "A Y526Q mutation in the Newcastle disease virus HN protein reduces its functional activities and attenuates virus replication and pathogenicity," *J. Virol.*, 83:7779-7782 (2009). cited by applicant

Kilpatrick et al., "Rapid development of affinity matured monoclonal antibodies using RIMMS," *Hybridoma*, 16(4):381-389 (1997). cited by applicant

Kim et al., "Evaluation of fusion protein cleavage site sequences of Newcastle disease virus in genotype matched vaccines," *PLOS One*, 12(3):e0173965 (2017). cited by applicant

Kim et al., "Expression and characterization of a recombinant Fab fragment derived from an anti-human alpha-fetoprotein monoclonal antibody," *Mol. Cells.*, 11:158-163 (2001). cited by applicant

Kim et al., "LaSota fusion (F) cleavage motif-mediated fusion activity is affected by other regions of the F protein from different genotype Newcastle disease virus in a chimeric virus: implication

for virulence attenuation,” *J. Gen. Virol.*, 97(6):1297-1304 (2016). cited by applicant

Kim et al., 1969, “Respiratory syncytial virus disease in infants despite prior administration of antigenic inactivated vaccine,” *Am. J. Epidemiol.*, 89(4):422-434. cited by applicant

Kim et al., 2012, “Replication, neurotropism, and pathogenicity of avian paramyxovirus serotypes 1-9 in chickens and ducks,” *PLoS One*. 2012;7(4):e34927 (13 pages). cited by applicant

Kim et al., 2013, “Mutations in the fusion protein cleavage site of avian paramyxovirus serotype 4 confer increased replication and syncytium formation in vitro but not increased replication and pathogenicity in chickens and ducks,” *PLoS One*, 8(1):e50598 (15 pages). cited by applicant

Kim et al., 2017, “A novel chimeric Newcastle disease virus vectored vaccine against highly pathogenic avian influenza virus,” *Virology*, 503:31-36. cited by applicant

Kohler, “Immunoglobulin chain loss in hybridoma lines,” *Proc. Natl. Acad. Sci. USA*, 77(4):2197-2199 (1980). cited by applicant

Korppi et al., 2004, “Respiratory morbidity 20 years after RSV infection in infancy,” *Pediatr. Pulmonol.*, 38(2):155-160. cited by applicant

Kostelny et al., “Formation of a bispecific antibody by the use of leucine zippers,” *J. Immunol.*, 148(5):1547-1553 (1992). cited by applicant

Kozbor et al., “A human hybrid myeloma for production of human monoclonal antibodies,” *J. Immunol.*, 133(6):3001-3005 (1984). cited by applicant

Krishnamurthy et al., “Differentially regulated interferon response determines the outcome of Newcastle disease virus infection in normal and tumor cell lines,” *J. Virology*, 80:5145-5155 (2006). cited by applicant

Kulangara et al., “Development of the combined positive score (CPS) for the evaluation of PD-L1 in solid tumors with the immunohistochemistry assay PD-L1 IHC 22C3 pharmDx,” *J. Clin. Oncol.*, 35(15_suppl):e 14589 (2017). cited by applicant

Kumar et al., 2010, “Experimental avian paramyxovirus serotype-3 infection in chickens and turkeys,” *Vet Res.*, 41(5):72 (14 pages). cited by applicant

Kumar et al., 2011, “Evaluation of the Newcastle disease virus F and HN proteins in protective immunity by using a recombinant avian paramyxovirus type 3 vector in chickens,” *J. Virol.*, 85(13):6521-6534 (retracted). cited by applicant

Kumar et al., 2020, “Retraction for Kumar et al., “Evaluation of the Newcastle Disease Virus F and HN Proteins in Protective Immunity by Using a Recombinant Avian Paramyxovirus Type 3 Vector in Chickens,”,” *J. Virol.*, 94(6):e01867-19 (1 page). cited by applicant

Kuraguchi et al., “Tumor-associated Apc mutations in Mlh1-/- Apc1638N mice reveal a mutational signature of Mlh1 deficiency,” *Oncogene*, 19(50):5755-5763 (2001). cited by applicant

Lam et al., “A new type of synthetic peptide library for identifying ligand-binding activity,” *Nature*, 354(6348):82-84 (1991). cited by applicant

Lam et al., 2011, “Safety and clinical usage of newcastle disease virus in cancer therapy,” *J Biomed Biotechnol*, 2011:718710 (13 pages). cited by applicant

Lamb et al., *Paramyxoviridae: The Viruses and Their Replication*. In B.N. Fields, D.M. Knipe, & P.M. Howley (Eds.), *Fundamental Virology*, pp. 577-604, Philadelphia, PA (1996). cited by applicant

Leach et al., “Enhancement of antitumor immunity by CTLA-4 blockade,” *Science*, 271:1734-1736 (1996). cited by applicant

Lee et al., “Isolation of cDNA for a human granulocyte-macrophage colony-stimulating factor by functional expression in mammalian cells,” *Proc. Natl. Acad. Sci. USA*, 82(13):4360-4364 (1985). cited by applicant

Lefranc et al., “IMGT, the international ImMunoGeneTics database,” *Nucl. Acids Res.*, 27(1):209-212 (1999). cited by applicant

Lefranc, “The IMGT unique numbering for immunoglobulins, T-cell receptors, and Ig-like domains,” *Immunologist*, 7:132-136 (1999). cited by applicant

Lei et al., "An oncolytic adenovirus expressing granulocyte macrophage colony-stimulating factor shows improved specificity and efficacy for treating human solid tumors," *Cancer Gene Therapy*, 16:33-43 (2009). cited by applicant

Leonard et al., "Effects of single-dose interleukin-12 exposure on interleukin-12-associated toxicity and interferon-gamma production," *Blood*, 90(7):2541-2548 (1997). cited by applicant

Li et al., "Therapeutic effects of a fusogenic Newcastle disease virus in treating head and neck cancer," *Head Neck*, 33(10):1394-1399 (2011). cited by applicant

Li et al., "Decreased dependence on receptor recognition for the fusion promotion activity of L289A-mutated newcastle disease virus fusion protein correlates with a monoclonal antibody-detected conformational change," *J. Virol.*, 79(2):1180-1190 (2005). cited by applicant

Li et al., 1999, "Recombinant influenza A virus vaccines for the pathogenic human A/Hong Kong/97 (H5N1) viruses," *J. Infect. Dis.*, 179(5):1132-1138. cited by applicant

Li et al., 2005, "Chimeric influenza virus hemagglutinin proteins containing large domains of the *Bacillus anthracis* protective antigen: protein characterization, incorporation into infectious influenza viruses, and antigenicity," *J. Virol.*, 79(15):10003-10012. cited by applicant

Liang et al., 2003, "Application of autologous tumor cell vaccine and NDV vaccine in treatment of tumors of digestive tract," *World J Gastroenterol.*, 9(3):495-498. cited by applicant

Lieschke et al., "Bioactive murine and human interleukin-12 fusion proteins which retain antitumor activity in vivo," *Nature*, 15: 35-40 (1997). cited by applicant

Lipson et al., "Durable Cancer Regression Off-treatment and Effective Re-induction Therapy with an Anti-PD-1 Antibody," *Clin. Cancer Res.*, 19(2):462-468 (2013). cited by applicant

Liu et al., "Adenovirus-mediated intratumoral expression of immunostimulatory proteins in combination with systemic Treg inactivation induces tumor-destructive immune responses in mouse models," *Cancer Gene Ther.*, 18(6):407-418 (2011). cited by applicant

Liu et al., "ICP34.5 deleted herpes simplex virus with enhanced oncolytic, immune stimulating, and anti-tumour properties," *Gene Therapy*, 10:292-303 (2003). cited by applicant

Liu et al., 2018, "Chimeric Newcastle disease virus-vectored vaccine protects chickens against H9N2 avian influenza virus in the presence of pre-existing NDV immunity," *Arch. Virol.*, 163(12):3365-3371. cited by applicant

Logan et al., "Adenovirus tripartite leader sequence enhances translation of mRNAs late after infection," *Proc. Natl. Acad. Sci. USA*, 81:3655-3659 (1984). cited by applicant

Lorence et al., "Phase 1 clinical experience using intravenous administration of PV701, an oncolytic Newcastle disease virus," *Curr. Cancer Drug Targets*, 7:157-167 (2007). cited by applicant

Lorence et al., 1988, "Newcastle disease virus as an antineoplastic agent: induction of tumor necrosis factor-alpha and augmentation of its cytotoxicity," *J. Natl. Cancer Inst.*, 80(16):1305-1312. cited by applicant

Lotze et al., "Cytokine Gene Therapy of Cancer Using Interleukin-12: Murine and Clinical Trials," *Ann. N.Y. Acad. Sci.*, 795:440-454 (1995). cited by applicant

Lowy et al., "Isolation of Transforming DNA: Cloning the Hamster *aprt* Gene," *Cell* 22:817-823 (1980). cited by applicant

Lutzky, "A phase 1 study of MEDI4736, an anti-PD-L1 antibody, in patients with advanced solid tumors," *J. Clin. Oncol.*, 32(suppl 5S):abstract 3001 (2014). cited by applicant

Maamary et al., 2011, "Newcastle disease virus expressing a dendritic cell-targeted HIV gag protein induces a potent gag-specific immune response in mice," *J Virol.*, 85(5):2235-2246. cited by applicant

MacCallum et al., "Antibody-antigen interactions: contact analysis and binding site topography," *J. Mol. Biol.*, 262(5):732-745 (1996). cited by applicant

Maeda et al., "Live bivalent vaccine for parainfluenza and influenza virus infections," *J. Viral.*, 79(11):6674-6679 (2005). cited by applicant

Mansour et al., "Oncolytic specificity of newcastle disease virus is mediated by selectivity for apoptosis-resistant cells," J. Virol., 85(12):6015-6023 (2011). cited by applicant

Martin, "Protein Sequence and Structure Analysis of Antibody Variable Domains," Antibody Engineering, Kontermann et al. eds., Springer-Verlag, Berlin, Chapter 31, pp. 33-51 (2001). cited by applicant

Martinez-Sobrido et al., 2006, "Protection against respiratory syncytial virus by a recombinant Newcastle disease virus vector," J Virol., 80(3):1130-1139. cited by applicant

Mazur et al., 2015, "Lower respiratory tract infection caused by respiratory syncytial virus: current management and new therapeutics," Lancet Respir. Med., 3(11):888-900. cited by applicant

Mazzolini et al., "Adenoviral Gene Transfer of Interleukin 12 into Tumors Synergizes with Adoptive T Cell Therapy Both at the Induction and Effector Level," Human Gene Ther., 11:113-125 (2000). cited by applicant

Mazzolini et al., "Regression of colon cancer and induction of antitumor immunity by intratumoral injection of adenovirus expressing interleukin-12," Cancer Gene Ther., 6(6):514-522 (1999). cited by applicant

Mcdermott et al., "Atezolizumab, an Anti-Programmed Death-Ligand 1 Antibody, in Metastatic Renal Cell Carcinoma: Long-Term Safety, Clinical Activity, and Immune Correlates From a Phase Ia Study," J. Clin. Oncol., 34(8):833-842 (2016). cited by applicant

Menne et al., "A comparison of signal sequence prediction methods using a test set of signal peptides," Bioinformatics, 16(8):741-742 (2000). cited by applicant

Meseck et al., "A Functional Recombinant Human 4-1BB Ligand for Immune Costimulatory Therapy of Cancer," J. Immunother., 34(2):175-182 (2011). cited by applicant

Miller et al., 2010, "Evidence for a new avian paramyxovirus serotype 10 detected in rockhopper penguins from the Falkland Islands," J Virol., 84(21):11496-11504. cited by applicant

Mills et al., 1971, "Experimental respiratory syncytial virus infection of adults. Possible mechanisms of resistance to infection and illness" J. Immunol., 107(1):123-130. cited by applicant

Milstein et al., "Hybrid hybridomas and their use in immunohistochemistry," Nature, 305:537-540 (1983). cited by applicant

Morea et al., "Antibody modeling: implications for engineering and design," Methods, 20(3):267-279 (2000). cited by applicant

Morgan et al., "Human gene therapy," Annu. Rev. Biochem., 62:191-217 (1993). cited by applicant

Morris et al., "Lung-specific expression in mice of a dominant negative mutant form of the p53 tumor suppressor protein," J. La. State Med. Soc., 150(4):179-185 (1998). cited by applicant

Morrison, "Transfectomas provide novel chimeric antibodies," Science, 229(4719):1202-1207 (1985). cited by applicant

Mulligan et al., "Selection for animal cells that express the *Escherichia coli* gene coding for xanthine-guanine phosphoribosyltransferase," Proc. Natl. Acad. Sci. USA, 78:2072-2076 (1981). cited by applicant

Mulligan, "The basic science of gene therapy," Science, 260:926-932 (1993). cited by applicant

Muranski et al., "Tumor-specific Th17-polarized cells eradicate large established melanoma," Blood, 112(2):362-373 (2008). cited by applicant

Murawski et al., "Newcastle disease virus-like particles containing respiratory syncytial virus G protein induced protection in BALB/c mice, with No. evidence of immunopathology", J. Virol., 84(2):1110-1123 (2010). cited by applicant

Mustaffa-Babjee et al., 1974, "A pathogenic paramyxovirus from a budgerigar (*Melopsittacus undulatus*), " Avian Dis., 18(2):226-230. cited by applicant

Muyldermans, "Single domain camel antibodies: current status," J. Biotechnol., 74(4):277-302 (2001). cited by applicant

Naganawa et al., "Generation of mouse-human hybridomas secreting human monoclonal antibodies to Japanese cedar pollen allergen Cry j1," Hum. Antibodies, 14:27-31 (2005). cited by applicant

Nakaya et al., "Recombinant Newcastle disease virus as a vaccine vector," *J. Virol.*, 75:11868-11873 (2001). cited by applicant

Narvaiza et al., "Intratumoral Coinjection of Two Adenoviruses, One Encoding the Chemokine IFN- γ -Inducible Protein-10 and Another Encoding IL-12, Results in Marked Antitumoral Synergy," *J. Immunol.*, 164(6):3112-3122 (2000). cited by applicant

Nguyen et al., 2020, "Oncolytic Virus Encoding a Master Pro-Inflammatory Cytokine Interleukin 12 in Cancer Immunotherapy," *Cells*, 9(2), 400. cited by applicant

Niu et al., "Recombinant Newcastle Disease virus Expressing IL15 Demonstrates Promising Antitumor Efficiency in Melanoma Model," *Technol. Cancer Res. Treatment*, 14(5):607-615 (2015). cited by applicant

Nolden et al., 2016, "Reverse genetics in high throughput: rapid generation of complete negative strand RNA virus cDNA clones and recombinant viruses thereof," *Sci Rep.*, 6:23887 (15 pages). cited by applicant

Nuttall et al., "Immunoglobulin VH domains and beyond: design and selection of single-domain binding and targeting reagents," *Curr. Pharm. Biotechnol.*, 1(3):253-263 (2000). cited by applicant

O'Hare et al., "Transformation of mouse fibroblasts to methotrexate resistance by a recombinant plasmid expressing a prokaryotic dihydrofolate reductase," *Proc. Natl. Acad. Sci. USA* 78:1527-1531 (1981). cited by applicant

OPDIVO® (nivolumab) Prescribing Information, revised Apr. 2018 (83 pages). cited by applicant

OPDIVO® (nivolumab) Prescribing Information, revised Dec. 2014 (20 pages). cited by applicant

OPDIVO® (nivolumab) Prescribing Information, revised Sep. 2019 (31 pages). cited by applicant

Oseledchyk et al., "Lysis-independent potentiation of immune checkpoint blockade by oncolytic virus," *Oncotarget* 9(47):28702-28716 with Supplementary Materials (2 pages) (2018). cited by applicant

Overwijk et al., "Tumor regression and autoimmunity after reversal of a functionally tolerant state of self-reactive CD8⁺ T cells," *J. Exp. Med.*, 198(4):568-580 (2003). cited by applicant

Padlan, "A possible procedure for reducing the immunogenicity of antibody variable domains while preserving their ligand-binding properties," *Mol. Immunol.*, 28:489-498 (1991). cited by applicant

Paramore et al., 2004, "Economic impact of respiratory syncytial virus-related illness in the US: an analysis of national databases," *Pharmacoeconomics*, 22(5):275-284. cited by applicant

Pardoll, "The blockade of immune checkpoints in cancer immunotherapy," *Nat. Rev. Cancer*, 12(4):252-264 (2012). cited by applicant

Park et al., "Engineered viral vaccine constructs with dual specificity: avian influenza and Newcastle disease," *Proc. Natl. Acad. Sci. (USA)*, 103:8203-8208 (2006). cited by applicant

Park et al., "Newcastle disease virus V protein is a determinant of host range restriction," *J. Virol.*, 77(17):9522-9532 (2003). cited by applicant

Park et al., 2003, "Newcastle disease virus (NDV)-based assay demonstrates interferon-antagonist activity for the NDV V protein and the Nipah virus V, W, and C proteins," *J Virol.*, 77(2):1501-1511. cited by applicant

Patent Cooperation Treaty, International Search Report for PCT/US2006/045859, dated Mar. 28, 2007. cited by applicant

Patent Cooperation Treaty, International Search Report for PCT/US2010/023335, dated Jun. 7, 2010. cited by applicant

Patent Cooperation Treaty, International Search Report for PCT/US2014/020299, dated Aug. 15, 2014. cited by applicant

Patent Cooperation Treaty, International Search Report for PCT/US2018/032255, dated Nov. 19, 2018. cited by applicant

Patent Cooperation Treaty, Written Opinion for PCT/US2006/045859, dated Mar. 28, 2007. cited by applicant

Patent Cooperation Treaty, Written Opinion for PCT/US2010/023335, dated Jun. 7, 2010. cited by applicant

Patent Cooperation Treaty, Written Opinion for PCT/US2014/020299, dated Jul. 19, 2014. cited by applicant

Patent Cooperation Treaty, Written Opinion for PCT/US2018/032255, dated Nov. 19, 2018. cited by applicant

Pecora et al., "Phase I trial of intravenous administration of PV701, an oncolytic virus, in patients with advanced solid cancers," *J. Clon. Oncol.*, 20:2251-2266 (2002). cited by applicant

Pedersen et al., "Comparison of surface accessible residues in human and murine immunoglobulin Fv domains. Implication for humanization of murine antibodies," *J. Mol. Biol.*, 235(3):959-973 (1994). cited by applicant

Peeters et al., "Generation of a recombinant chimeric Newcastle disease virus vaccine that allows serological differentiation between vaccinated and infected animals," *Vaccine*, 19:1616-1627 (2001). cited by applicant

Peeters et al., "Rescue of Newcastle disease virus from cloned cDNA: evidence that cleavability of the fusion protein is a major determinant for virulence," *J. Virol.*, 73(6):5001-5009 (1999). cited by applicant

Peggs et al., "Cancer immunotherapy: co-stimulatory agonists and co-inhibitory antagonists," *Clin. Immunol.*, 157(1):9-19 (2009). cited by applicant

Pelaia et al., 2006, "Respiratory infections and asthma," *Respir. Med.*, 100(5):775-784 (Epub 2005). cited by applicant

Persic et al., "An integrated vector system for the eukaryotic expression of antibodies or their fragments after selection from phage display libraries," *Gene*, 187:9-18 (1997). cited by applicant

Phuangsab et al., "Newcastle disease virus therapy of human tumor xenografts: antitumor effects of local or systemic administration," *Cancer Lett.*, 172:27-36 (2001). cited by applicant

Plitt et al., "Cancer therapy with Newcastle disease virus: rationale for new immunotherapeutic combinations," *Clinical Investigation*, 5(1):75-87 (2015). cited by applicant

Polack et al., 2002, "A role for immune complexes in enhanced respiratory syncytial virus disease," *J. Exp. Med.*, 196(6):859-865. cited by applicant

Pomer et al., 1995, "Tumor response and 4 year survival-data of patients with advanced renal-cell carcinoma treated with autologous tumor vaccine and subcutaneous R-IL-2 and IFN-alpha(2b)," *Int. J. Oncol.*, 6(5):947-954. cited by applicant

Proudfoot, "Transcriptional interference and termination between duplicated α -globin gene constructs suggests a novel mechanism for gene regulation," *Nature*, 322(7):562-565 (1986). cited by applicant

Puhler et al., "Generation of a recombinant oncolytic Newcastle disease virus and expression of a full IgG antibody from two transgenes," *Gene Ther.*, 15:371-383 (2008). cited by applicant

Puzanov et al., "Phase 1 results of a phase 1b/2, multicenter, open-label trial to evaluate safety and efficacy of talimogene laherparepvec (T-VEC) and ipilimumab (ipi) vs ipi alone in previously untreated, unresected stage IIIB-IV melanoma," *J. Immunother. Cancer*, 1(Suppl 1):P84 (2013). cited by applicant

Quetglas et al., "Virotherapy with a Semliki Forest Virus-Based Vector Encoding IL12 Synergizes with PD-1/PD-L1 Blockade," *Cancer Immunol. Res.*, 3(5):449-454 (2015). cited by applicant

Quezada et al., "CTLA4 blockade and GM-CSF combination immunotherapy alters the intratumor balance of effector and regulatory T cells," *J. Clin. Invest.*, 116(7): 1935-1945 (2006). cited by applicant

Quinn et al., "Rapid quantitation of recombinant retrovirus produced by packaging cell clones," *Biotechniques*, 23(6):1038-1044 (1997). cited by applicant

Ravindra et al., "Newcastle disease virus as an oncolytic agent," *Indian J. Med. Res.*, 130(5):507-513 (2009). cited by applicant

Ren et al., 2016, "Recombinant Newcastle Disease Virus Encoding IL-12 and/or IL-2 as Potential Candidate for Hepatoma Carcinoma Therapy," *Technol. Cancer Res. Treat.*, 15(5):NP83-NP94. cited by applicant

Ribas et al., "Oncolytic Virotherapy Promotes Intratumoral T Cell Infiltration and Improves Anti-PD-1 Immunotherapy," *Cell*, 170(6):1109-1119 (2017). cited by applicant

Riechmann et al., "Single domain antibodies: comparison of camel VH and camelised human VH domains," *J. Immunol. Methods*, 231:25-38 (1999). cited by applicant

Robbins et al., "Human tumor antigens recognized by T cells," *Curr. Opin. Immunol.*, 8(5):628-636 (1996). cited by applicant

Robert et al., "Ipilimumab plus dacarbazine for previously untreated metastatic melanoma," *N. Engl. J. Med.*, 364(26):2517-2526 (2011). cited by applicant

Roguska et al., "A comparison of two murine monoclonal antibodies humanized by CDR-grafting and variable domain resurfacing," *Protein Eng.*, 9(10):895-904 (1996). cited by applicant

Roguska et al., "Humanization of murine monoclonal antibodies through variable domain resurfacing," *Proc. Natl. Acad. Sci USA*, 91(3):969-973 (1994). cited by applicant

Romer-Oberdorfer et al., 1999, "Generation of recombinant lentogenic Newcastle disease virus from cDNA" *J Gen Virol.*, 80(Pt 11):2987-2995. cited by applicant

Ruther et al., "Easy identification of cDNA clones," *EMBO J.*, 2(10):1791-1794 (1983). cited by applicant

Saif et al., 1997, "Natural and experimental infection of turkeys with avian paramyxovirus-7," *Avian Dis.*, 41(2):326-329. cited by applicant

Sandhu et al., 2003, "Influenza A Virus Infection of Domestic Ducks," *JSTOR*, 47:93-99. cited by applicant

Santerre et al., "Expression of prokaryotic genes for hygromycin B and G418 resistance as dominant-selection markers in mouse L cells," *Gene*, 30:147-156 (1984). cited by applicant

Sapoznik et al. "Novel anti-melanoma immunotherapies: disarming tumor escape mechanisms," *Clin Dev Immunol.*, 2012: 818214. doi: 10.1155/2012/818214. Epub Apr. 23, 2012 (2012). cited by applicant

Sawai et al., "Direct production of the Fab fragment derived from the sperm immobilizing antibody using polymerase chain reaction and cDNA expression vectors," *A. J. Reprod. Immunol.*, 34(1):26-34 (1995). cited by applicant

Scapin et al., "Structure of full-length human anti-PD1 therapeutic IgG4 antibody pembrolizumab," *Nat. Struct. Mol. Biol.*, 22(12):953-958 (2015). cited by applicant

Schickli et al., "Plasmid-only rescue of influenza A virus vaccine candidates," *Phil. Trans. R. Soc. Lond.*, 356:1965-1973 (2001). cited by applicant

Schirmacher et al., "Antitumor effects of Newcastle Disease Virus in vivo: Local versus systemic effects," *Int. J. Oncol.*, 18:945-952 (2001). cited by applicant

Schirmacher et al., "Newcastle disease virus: a promising vector for viral therapy, immune therapy, and gene therapy of cancer," *Methods Mol. Biol.*, 542: 565-605 (2009). cited by applicant

Schirmacher et al., 1998, "Immunization with virus-modified tumor cells" *Semin. Oncol.*, 25(6):677-696. cited by applicant

Schirmacher et al., 1999, "Human tumor cell modification by virus infection: an efficient and safe way to produce cancer vaccine with pleiotropic immune stimulatory properties when using Newcastle disease virus," *Gene Ther.*, 6(1):63-73. cited by applicant

Schirmacher et al., 2014, "Long-term remission of prostate cancer with extensive bone metastases upon immuno- and virotherapy: A case report," *Oncol Lett.*, 8(6):2403-2406. cited by applicant

Schirmacher et al., 2014, "Multimodal cancer therapy involving oncolytic newcastle disease virus, autologous immune cells, and bi-specific antibodies," *Front Oncol.*, 4:224 (5 pages). cited by applicant

Schirmacher, 2016, "Fifty Years of Clinical Application of Newcastle Disease Virus: Time to

Celebrate!" *Biomedicines*, 4(3), 14 pages. cited by applicant

Scott et al., "Searching for peptide ligands with an epitope library," *Science*, 249(4967):386-390 (1990). cited by applicant

Seliger et al., "Characterization of the major histocompatibility complex class I deficiencies in B16 melanoma cells," *Cancer Res.*, 61(3):1095-1099 (2001). cited by applicant

Senne et al., 2004, "Control of Newcastle disease by vaccination," *Dev. Biol. (Basel)*, 119:165-170. cited by applicant

Seppi et al., "Direct determination of oxygen by HPLC. 2. Chamber and sample application system for determination of O_2 at trace levels," *Anal. Chem.*, 69(21):4476-4481 (1997). cited by applicant

Sergel et al., "A Single Amino Acid Change in the Newcastle Disease Virus Fusion Protein Alters the Requirement for HN Protein in Fusion," *J. Virol.*, 74(11):5101-5107 (2000). cited by applicant

Sharma et al., "Triggering the interferon antiviral response through an ikk-related pathway," *Science*, 300(5622):1148-1151 (2003). cited by applicant

Shenk, "Adenoviridae: The Viruses and Their Replication," *Fundamental Virology*, Fields et al. eds., Lippincott-Raven, Philadelphia, PA, pp. 979-1016 (1996). cited by applicant

Shim et al., "Inhibitory Receptors Induced by VSV Viroimmunotherapy Are Not Necessarily Targets for Improving Treatment Efficacy," *Mol. Ther.*, 25(4):962-975 (2017). cited by applicant

Shinmoto et al., "Generation of mouse-human hybridomas secreting antibodies against peanut allergen Ara h1," *Cytotechnology*, 46(1):19-23 (2004). cited by applicant

Shinoff et al., 2008, "Young infants can develop protective levels of neutralizing antibody after infection with respiratory syncytial virus," *J. Infect. Dis.*, 198(7):1007-1015. cited by applicant

Shinohara et al., "Structure and chromosomal localization of the human PD-1 gene (PDCD1)," *Genomics*, 23:704-706 (1994). cited by applicant

Shnyrova et al., 2007, "Vesicle formation by self-assembly of membrane-bound matrix proteins into a fluidlike budding domain," *J Cell Biol.*, 179(4):627-633. cited by applicant

Shortridge et al., 1980, "Isolation and properties of viruses from poultry in Hong Kong which represent a new (sixth) distinct group of avian paramyxoviruses," *J Gen Virol.*, 49(2):255-262. cited by applicant

Sigurs et al., 2000, "Respiratory syncytial virus bronchiolitis in infancy is an important risk factor for asthma and allergy at age 7," *Am. J. Respir. Crit. Care Med.*, 161(5):1501-1507. cited by applicant

Sigurs et al., 2005, "Severe respiratory syncytial virus bronchiolitis in infancy and asthma and allergy at age 13," *Am. J. Respir. Crit. Care Med.*, 171(2):137-141 (Epub 2004). cited by applicant

Silberhumer et al., "Genetically engineered oncolytic Newcastle disease virus effectively induces sustained remission of malignant pleural mesothelioma," *Mol. Cancer Ther.*, 9(10):2761-2769 (2010). cited by applicant

Simoes et al., 2003, "Impact of severe disease caused by respiratory syncytial virus in children living in developed countries," *Pediatr. Infect. Dis. J.*, 22(2 Suppl):S13-S20. cited by applicant

Simpson et al., "Regulation of CD4 T cell activation and effector function by inducible costimulator (ICOS)," *Curr. Opin. Immunol.*, 22(3):326-332 (2010). cited by applicant

Sinkovics et al., "Newcastle disease virus (NDV): brief history of its oncolytic strains," *J. Clin. Virol.*, 16:1-15 (2000). cited by applicant

Song et al., "Antitumor efficacy of viral therapy using genetically engineered Newcastle disease virus [NDV(F3aa)-GFP] for peritoneally disseminated gastric cancer," *J. Mol. Med. (Berl.)*, 88(6):589-596 (2010). cited by applicant

Spranger et al., "Up-regulation of PD-L1, IDO, and T(regs) in the melanoma tumor microenvironment is driven by CD8(+) T cells," *Sci. Transl. Med.*, 5(200):200ra116 (2013). cited by applicant

Stanislawek et al., 2002, "Avian paramyxoviruses and influenza viruses isolated from mallard

ducks (*Anas platyrhynchos*) in New Zealand,” *Arch. Virol.*, 147(7):1287-1302. cited by applicant

Steglich et al., 2013, “Chimeric newcastle disease virus protects chickens against avian influenza in the presence of maternally derived NDV immunity,” *PLoS One*, 8(9):e72530 (14 pages). cited by applicant

Steiner et al., 2004, “Antitumor vaccination of patients with glioblastoma multiforme: a pilot study to assess feasibility, safety, and clinical benefit,” *J Clin Oncol.*, 22(21):4272-4281. cited by applicant

Studnicka et al., “Human-engineered monoclonal antibodies retain full specific binding activity by preserving non-CDR complementarity-modulating residues,” *Protein Eng.*, 7(6):805-814 (1994). cited by applicant

Su et al., “Immunoadjuvant activities of a recombinant chicken IL-12 in chickens vaccinated with Newcastle disease virus recombinant HN protein,” *Veterinary Microbiology*, 151:220-228 (2011). cited by applicant

Subbiah et al., 2010, “Pathogenesis of two strains of avian paramyxovirus serotype 2, Yucaipa and Bangor, in chickens and turkeys,” *Avian Dis.*, 54(3):1050-1057. cited by applicant

Suresh et al., “Bispecific monoclonal antibodies from hybrid hybridomas,” *Methods Enzymol.*, 121:210-228 (1986). cited by applicant

Swann et al., “Type I IFN contributes to NK cell homeostasis, activation, and antitumor function,” *J. Immunol.*, 178(12):7540-7549 (2007). cited by applicant

Swayne et al., “Recombinant paramyxovirus type 1-avian influenza-H7 virus as a vaccine for protection of chickens against influenza and Newcastle disease,” *Avian Dis.*, 47:1047-1050 (2003). cited by applicant

Swayne, 2003, “Vaccines for List A poultry diseases: emphasis on avian influenza,” *Dev. Biol. (Basel)*, 114:201-212. cited by applicant

Szybalska et al., “Genetics of human cell line. IV. DNA-mediated heritable transformation of a biochemical trait,” *Proc. Natl. Acad. Sci. USA* 48:2026-2034 (1992). cited by applicant

Tan et al., ““Superhumanized” antibodies: reduction of immunogenic potential by complementarity-determining region grafting with human germline sequences: application to an anti-CD28,” *J. Immunol.*, 169(2):1119-1125 (2002). cited by applicant

Tang et al., “Use of a peptide mimotope to guide the humanization of MRK-16, an anti-P-glycoprotein monoclonal antibody,” *J. Biol. Chem.*, 274(39): 27371-27378 (1999). cited by applicant

TECENTRIQ® (atezolizumab), Prescribing Information, revised Oct. 2016 (23 pages). cited by applicant

TECENTRIQ™ (atezolizumab), Prescribing Information, May 2016 (17 pages). cited by applicant

Terregino et al., 2013, Antigenic and genetic analyses of isolate APMV/wigeon/Italy/3920-1/2005 indicate that it represents a new avian paramyxovirus (APMV-12), *Arch. Virol.*, 158(11):2233-2243. cited by applicant

Therasse et al., “New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada,” *J. Natl. Cancer Inst.*, 92(3):205-216 (2000). cited by applicant

Tolstoshev, “Gene therapy, concepts, current trials and future directions,” *Ann. Rev. Pharmacol. Toxicol.*, 32:573-596 (1993). cited by applicant

Topalian et al., “Safety, activity, and immune correlates of anti-PD-1 antibody in cancer,” *New Eng. J. Med.*, 366:2443-2454 (2012). cited by applicant

Tramontano et al., “Framework residue 71 is a major determinant of the position and conformation of the second hypervariable region in the VH domains of immunoglobulins,” *J. Mol. Biol.*, 215(1):175-182 (1990). cited by applicant

Traunecker et al., “Bispecific single chain molecules (Janusins) target cytotoxic lymphocytes on

HIV infected cells,” *EMBO J.*, 10(12):3655-3659 (1991). cited by applicant

Tumeh et al., “PD-1 blockade induces responses by inhibiting adaptive immune resistance,” *Nature*, 515(7528):568-571 (2014). (Extended Data Figures 1-6 and Extended Data Tables 1-4 attached). cited by applicant

Tumova et al., 1979, “A hitherto unreported paramyxovirus of turkeys,” *Res. Vet. Sci.*, 27(2):135-140. cited by applicant

Turk et al., “Concomitant tumor immunity to a poorly immunogenic melanoma is prevented by regulatory T cells,” *J. Exp. Med.*, 200(6):771-782 (2004). cited by applicant

Tuve et al., “In situ adenovirus vaccination engages T effector cells against cancer,” *Vaccine*, 27:4225-4239 (2009). cited by applicant

UniProtKB/Swiss-Prot No. Q60819.1, “RecName: Full=Interleukin-15 receptor subunit alpha; Short=IL-15 receptor subunit alpha; Short=IL-15R-alpha; Short=IL-15RA; AltName: CD_antigen=CD215; Contains: RecName: Full=Soluble interleukin-15 receptor subunit alpha; Short=sIL-15 receptor subunit alpha; Short=sIL-15R-alpha; Short=sIL-15RA; Flags: Precursor,” Oct. 16, 2019. cited by applicant

U.S. Appl. No. 14/205,776, Final Office Action dated Sep. 23, 2016. cited by applicant

U.S. Appl. No. 14/205,776, Non-Final Office Action dated Mar. 4, 2016. cited by applicant

U.S. Appl. No. 14/205,776, Amendment and Response dated Jan. 8, 2016. cited by applicant

U.S. Appl. No. 14/205,776, Amendment under 37 C.F.R. 1.111 dated Jun. 9, 2015. cited by applicant

U.S. Appl. No. 14/205,776, Final Office Action dated Aug. 10, 2015. cited by applicant

U.S. Appl. No. 14/205,776, Non-Final Office Action dated Jan. 28, 2015. cited by applicant

U.S. Appl. No. 14/774,962, Non-Final Office Action mailed Nov. 9, 2016. cited by applicant

U.S. Appl. No. 15/588,251, Non-Final Office Action dated Jul. 16, 2018. cited by applicant

U.S. Appl. No. 15/789,340, Amendment and Response dated May 17, 2019. cited by applicant

U.S. Appl. No. 15/789,340, Final Office Action dated Jun. 14, 2019. cited by applicant

U.S. Appl. No. 15/789,340, Non-Final Office Action date Feb. 19, 2019. cited by applicant

U.S. Appl. No. 15/789,539, Amendment and Response dated Oct. 30, 2018. cited by applicant

U.S. Appl. No. 15/789,539, Non-Final Office Action mailed Jul. 30, 2018. cited by applicant

U.S. Appl. No. 15/789,539, Notice of Allowance mailed Dec. 11, 2018. cited by applicant

Vail et al., “Spontaneously occurring tumors of companion animals as models for human cancer,” *Cancer Invest.*, 18:781-792 (2000). cited by applicant

Van Den Hoogen et al., 2001, “A newly discovered human pneumovirus isolated from young children with respiratory tract disease,” *Nat. Med.*, 7(6):719-724. cited by applicant

Van Heeke et al., “Expression of human asparagine synthetase in *Escherichia coli*,” *J. Biol. Chem.*, 264(10):5503-5509 (1989). cited by applicant

Velu et al., “Role of PD-1 co-inhibitory pathway in HIV infection and potential therapeutic options,” *Retrovirology*, 12:14 (2015). cited by applicant

Verma et al., “Gene therapy—promises, problems and prospects,” *Nature*, 389:239-242 (1997). cited by applicant

Vigil et al., “Recombinant Newcastle disease virus as a vaccine vector for cancer therapy,” *Molecular Therapy*, 16(11):1883-1890 (2008). cited by applicant

Vigil et al., “Use of reverse genetics to enhance the oncolytic properties of Newcastle disease virus,” *Cancer Research*, 67(17):8285-8292 (2007). cited by applicant

Vlasak et al., “Use of flow cytometry for characterization of human cytomegalovirus vaccine particles,” *Vaccine*, 34:2321-2328 (2016). cited by applicant

Von Heijne, “A new method for predicting signal sequence cleavage sites,” *Nucleic Acids Res.*, 14(11):4683-4690 (1986). cited by applicant

Von Heijne, “Patterns of amino acids near signal-sequence cleavage sites,” *Eur. J. Biochem.*, 133(1):17-21 (1983). cited by applicant

Wagner et al., 1989, "Serum immunoglobulin G antibody subclass response to respiratory syncytial virus F and G glycoproteins after first, second, and third infections," J. Clin. Microbiol., 27(3):589-592. cited by applicant

Waitz et al., "Potent induction of tumor immunity by combining tumor cryoablation with anti-CTLA-4 therapy," Cancer Res., 72(2):430-439 (2012). cited by applicant

Wakamatsu et al., "Convergent and divergent effects of costimulatory molecules in conventional and regulatory CD4+ T cells," Proc. Natl. Acad. Sci. USA, 110(3):1023-1028 (2013). cited by applicant

Wakamatsu et al., "The effect on pathogenesis of Newcastle disease virus LaSota strain from a mutation of the fusion cleavage site to a virulent sequence," Avian Dis., 50(4):483-488 (2006). cited by applicant

Walsh et al., 2004, "Humoral and mucosal immunity in protection from natural respiratory syncytial virus infection in adults," J. Infect. Dis., 190(2):373-378. cited by applicant

Walter et al., "Targeted inhibition of interferon-dependent intercellular adhesion molecule-1 (ICAM-1) expression using dominant-negative stat1," J. Biol. Chem., 272(45):28582-28589 (1997). cited by applicant

Walter et al., "Two avirulent, lentogenic strains of Newcastle disease virus are cytotoxic for some human pancreatic tumor lines in vitro," J. Pancreas, 13(5):502-513 (2012). cited by applicant

Wang et al., "A Novel, Clinically Relevant Animal Model of Metastatic Pancreatic Adenocarcinoma Biology and Therapy," Int. J. Pancreatol., 29(1):37-46 (2001). cited by applicant

Wang et al., "Impact of modification of cleavage site of fusion protein and foreign gene insertion on the virulence of Newcastle Disease Virus LaSota vaccine strain," Wei Sheng Wu Xue Bao, 48(3):362-368 (2008). (English abstract attached). cited by applicant

Washburn et al., 2002, "Human tumor cell infection by Newcastle Disease Virus leads to upregulation of HLA and cell adhesion molecules and to induction of interferons, chemokines and finally apoptosis," Int. J. Oncol., 21(1):85-93. cited by applicant

Weber et al., "Viral suppression of the interferon system," Biochimie, 89(6-7):836-842 (2007). cited by applicant

Webster et al., 1976, "Ortho- and paramyxoviruses from migrating feral ducks: characterization of a new group of influenza A viruses," J Gen Virol., 32(2):217-225. cited by applicant

Welliver et al., 1980, "The antibody response to primary and secondary infection with respiratory syncytial virus: kinetics of class-specific responses," J. Pediatr., 96(5):808-813. cited by applicant

Wheelock et al., 1964, "Observations on the repeated administration of viruses to a patient with acute leukemia. A preliminary report," N. Engl. J. Med., 271:645-651. cited by applicant

Wigler et al., "Transformation of mammalian cells with an amplifiable dominant-acting gene," Proc Natl. Acad. Sci. USA, 77:3567-3570 (1980). cited by applicant

Wigler et al., "Transfer of purified herpes virus thymidine kinase gene to cultured mouse cells," Cell, 11(1):223-232 (1977). cited by applicant

Wilden et al., "Expression of RIG-I, IRF3, IFN-beta and IRF7 determines resistance or susceptibility of cells to infection by Newcastle Disease Virus," Int. J. Oncol., 34(4):971-982 (2009). cited by applicant

Wold et al., "Adenovirus vectors for gene therapy, vaccination and cancer gene therapy," Curr. Gene Ther., 13(6):421-433 (2013). cited by applicant

Woller et al., "Oncolytic viruses as anticancer vaccines," Front. Oncol., 4(188):1-13 (2014). cited by applicant

Woolcock et al., 1996, "Isolation of paramyxovirus serotype 7 from ostriches (*Struthio camelus*)," Avian Dis., 40(4):945-949. cited by applicant

Wren et al., "SIGNAL-Sequence Information and GeNomic Analysis," Comput. Methods Programs Biomed., 68(2):177-181 (2002). cited by applicant

Wright et al., 1976, "Evaluation of a live, attenuated respiratory syncytial virus vaccine in infants,"

J. Pediatr., 88(6):931-936. cited by applicant

Wright et al., 1982, "Administration of a highly attenuated, live respiratory syncytial virus vaccine to adults and children," *Infect. Immun.*, 37(1):397-400. cited by applicant

Wu et al., "Delivery systems for gene therapy," *Biotherapy*, 3(1):87-95 (1991). cited by applicant

Yamaki et al., "The potential of recombinant vesicular stomatitis virus-mediated virotherapy against metastatic colon cancer," *Int. J. Mol. Med.*, 31:299-306 (2013). cited by applicant

Yamamoto et al., 2015, "Characterization of novel avian paramyxovirus strain APMV/Shimane67 isolated from migratory wild geese in Japan," *J Vet Med Sci.*, 77(9):1079-1085. cited by applicant

Yamane et al., 1982, "Characterization of avian paramyxoviruses isolated from feral ducks in northern Japan: the presence of three distinct viruses in nature," *Microbiol Immunol.*, 26(7):557-568. cited by applicant

Yao et al., "Reviving exhausted T lymphocytes during chronic virus infection by B7-H1 blockade," *Trends Mol. Med.*, 12(6):244-246 (2006). cited by applicant

Ying et al., "Adenovirus-mediated intratumoral expression of immunostimulatory proteins in combination with systemic Treg inactivation induces tumor-destructive immune responses in mouse models," *Cancer Gene Ther.*, 18(6):407-418 (2011). cited by applicant

Yoneyama et al., "The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses," *Nature Immunol.*, 5:730-737 (2004). cited by applicant

Yoshida et al., 2017, "Avian Paramyxovirus Type-3 as a Vaccine Vector: Identification of a Genome Location for High Level Expression of a Foreign Gene," *Front Microbiol.*, 8:693 (8 pages). cited by applicant

Yoshida et al., 2019, "Novel avian paramyxovirus-based vaccine vectors expressing the Ebola virus glycoprotein elicit mucosal and humoral immune responses in guinea pigs," *Sci. Rep.*, 9(1):5520 (10 pages). cited by applicant

Zamarin et al., "Potentiation of immunomodulatory antibody therapy with oncolytic viruses for treatment of cancer," *Mol. Therapy-Oncolytics*, 1:14004; doi: 10.1038/mto.2014.4. (2014). cited by applicant

Zamarin et al., "Upregulation of PD-L1 in tumor microenvironment is a resistance mechanism for oncolytic virus immunotherapy," *J. Immunother. Cancer*, 5(Suppl 2):87 (2017). cited by applicant

Zamarin et al., "Enhancement of Oncolytic Properties of Genetically-Engineered Fusogenic Newcastle Disease Virus through Antagonism of Cellular Innate Immune Responses," *Mol. Ther.*, 16(Suppl. 1), Abstract #43 (2008). cited by applicant

Zamarin et al., "Enhancement of oncolytic properties of recombinant Newcastle disease virus through antagonism of cellular innate immune responses," *Mol. Ther.*, 17(4):697-706 (2009). cited by applicant

Zamarin et al., "Genetically engineered Newcastle disease virus for malignant melanoma therapy," *Gene Ther.*, 16(6):796-804 (2009). cited by applicant

Zamarin et al., "Intratumoral modulation of the inducible co-stimulator ICOS by recombinant oncolytic virus promotes systemic anti-tumour immunity," *Nat. Commun.*, 8:14340 (2017). cited by applicant

Zamarin et al., "Localized oncolytic virotherapy inflames distant tumors and synergizes with immune checkpoint blockade leading to systemic tumor rejection," *J. Immunother. Cancer*, 1(Suppl 1):09 (2013). cited by applicant

Zamarin et al., "Oncolytic Newcastle disease virus for cancer therapy: old challenges and new directions," *Future Microbiol.*, 7:347-367 (2012). cited by applicant

Zamarin et al., "PD-L1 in tumor microenvironment mediates resistance to oncolytic immunotherapy," *J. Clin. Invest.*, 128(4):1413-1428, with Supplemental Materials (2018). cited by applicant

Zamarin et al., 2014, "Localized oncolytic virotherapy overcomes systemic tumor resistance to immune checkpoint blockade immunotherapy," *Sci. Transl. Med.*, 6(226):226ra32, with

Supplemental Materials (109 pages). cited by applicant
Zamarin et al., 2017, "P415: Upregulation of PD-L1 in tumor microenvironment is a resistance mechanism for oncolytic virus immunotherapy," J. Immunother. Cancer, 5(Suppl 2):87, p. 202. cited by applicant
Zhang et al., "Anti-Oncogene and Tumor Suppressor Gene Therapy—Examples from a Lung Cancer Animal Model," In-Vivo, 8:755-769 (1994). cited by applicant
Zhao et al., "P and M gene junction is the optimal insertion site in Newcastle disease virus vaccine vector for foreign gene expression," J. Gen. Virol., 96(Pt 1):40-45 (2015) (Epub Oct. 1, 2014). cited by applicant
Zhao et al., 2008, "Recombinant Newcastle disease virus expressing human interleukin-2 serves as a potential candidate for tumor therapy," Virus Res., 136(1-2):75-80. cited by applicant
Zimmer et al., "A chimeric respiratory syncytial virus fusion protein functionally replaces the F and HN glycoproteins in recombinant sendai virus," J Virol., 79(16):10467-10477 (2005). cited by applicant
Zitvogel et al., "Type I interferons in anticancer immunity," Nature Rev. Immunol., 15:405-414 (2015). cited by applicant

Primary Examiner: Foley; Shanon A.

Attorney, Agent or Firm: CANTOR COLBURN LLP

Background/Summary

(1) This application is a U.S. National Stage Application under 35 U.S.C. § 371 of International Patent Application No. PCT/US2019/046837, filed Aug. 16, 2019, which claims the benefit of U.S. Provisional Patent Application No. 62/765,242, filed Aug. 17, 2018, the disclosure of each of which is incorporated by reference herein in its entirety. (2) The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Jul. 31, 2019, is named 6923-280-228_SL.txt and is 375,809 bytes in size.

1. INTRODUCTION

(1) In one aspect, described herein are recombinant Newcastle disease virus ("NDV") comprising a packaged genome, wherein the packaged genome comprises a transgene encoding respiratory syncytial virus ("RSV") F protein. In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene comprising a codon optimized nucleic acid sequence encoding RSV F protein. In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In some embodiments, the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence. Also described herein are compositions comprising such recombinant NDV and the use of such recombinant NDV to induce an immune response to RSV F protein. In another aspect, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding human metapneumovirus (hMPV) F protein. In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene comprising a codon optimized nucleic acid sequence encoding hMPV F protein. In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a

transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In some embodiments, the ectodomain of the hMPV F protein is encoded by a codon optimized nucleic acid sequence. Also described herein are compositions comprising such recombinant NDV and the use of such recombinant NDV to induce an immune response to hMPV F protein.

2. BACKGROUND

(2) 2.1 RSV

(3) Human respiratory syncytial virus (RSV), a negative sense RNA virus in the Pneumoviridae family (Afonso et al., 2016, Arch. Virol. 161: 2351-2360), is the major cause of bronchiolitis and pneumonia in infants (Hall C B, Long C E, Schnabel K C. Respiratory syncytial virus infections in previously healthy working adults. Clinical infectious diseases: an official publication of the Infectious Diseases Society of America. 2001; 33(6):792-6. Epub 2001/08/21. doi: 10.1086/322657. PubMed PMID: 11512084; Paramore L C, Ciuryla V, Ciesla G, Liu L. Economic impact of respiratory syncytial virus-related illness in the US: an analysis of national databases. Pharmacoeconomics. 2004; 22(5):275-84. Epub 2004/04/06. PubMed PMID: 15061677). RSV outbreaks occur on an annual basis and essentially all persons are infected within the first two years of life. While RSV infection is limited to the upper respiratory tract in most healthy adults and children, severe, even fatal, RSV pneumonia occurs in young infants 2 to 4 months of age, transplant recipients and the elderly (Hall et al.). Secondary RSV infections, which are generally limited to the upper respiratory tract, present with mild, cold-like symptoms in healthy adults, but are commonly associated with otitis media in young children (Alper C M, Winther B, Mandel E M, Hendley J O, Doyle W J. Rate of concurrent otitis media in upper respiratory tract infections with specific viruses. Archives of otolaryngology—head & neck surgery. 2009; 135(1):17-21. Epub 2009/01/21. doi: 10.1001/archotol.135.1.17. PubMed PMID: 19153302; Hall C B, Walsh E E, Long C E, Schnabel K C. Immunity to and frequency of reinfection with respiratory syncytial virus. The Journal of infectious diseases. 1991; 163(4):693-8. Epub 1991/04/01. PubMed PMID: 2010624; Simoes E A, Carbonell-Estrany X. Impact of severe disease caused by respiratory syncytial virus in children living in developed countries. The Pediatric infectious disease journal. 2003; 22(2 Suppl):S13-8; discussion S8-20. Epub 2003/04/03. doi: 10.1097/01.inf.0000053881.47279.d9. PubMed PMID: 12671448). In addition, RSV has been associated with the development of asthma, and exacerbation of wheezing in asthmatic patients (Korppi M, Piippo-Savolainen E, Korhonen K, Remes S. Respiratory morbidity 20 years after RSV infection in infancy. Pediatric pulmonology. 2004; 38(2):155-60. Epub 2004/06/24. doi: 10.1002/ppul.20058. PubMed PMID: 15211700; Pelaia G, Vatrella A, Gallelli L, Renda T, Cazzola M, Maselli R, et al. Respiratory infections and asthma. Respiratory medicine. 2006; 100(5):775-84. Epub 2005/11/18. doi: 10.1016/j.rmed.2005.08.025. PubMed PMID: 16289785; Sigurs N, Bjarnason R, Sigurbergsson F, Kjellman B. Respiratory syncytial virus bronchiolitis in infancy is an important risk factor for asthma and allergy at age 7. American journal of respiratory and critical care medicine. 2000; 161(5):1501-7. Epub 2000/05/12. PubMed PMID: 10806145; Sigurs N, Gustafsson P M, Bjarnason R, Lundberg F, Schmidt S, Sigurbergsson F, et al. Severe respiratory syncytial virus bronchiolitis in infancy and asthma and allergy at age 13. American journal of respiratory and critical care medicine. 2005; 171(2):137-41. Epub 2004/11/02. doi: 10.1164/rccm.200406-7300C. PubMed PMID: 15516534).

(4) Immunity to RSV is remarkably ineffective, allowing for repeated infection of immunocompetent children and adults (Buchman C A, Doyle W J, Pilcher O, Gentile D A, Skoner D P. Nasal and otologic effects of experimental respiratory syncytial virus infection in adults. American journal of otolaryngology. 2002; 23(2):70-5. Epub 2002/03/15. PubMed PMID: 1189397; Mills Jt, Van Kirk J E, Wright P F, Chanock R M. Experimental respiratory syncytial virus infection of adults. Possible mechanisms of resistance to infection and illness. Journal of immunology. 1971; 107(1):123-30. Epub 1971/07/01. PubMed PMID: 5091954; Habibi M S,

Jozwik A, Makris S, Dunning J, Paras A, DeVincenzo J P, et al. Impaired Antibody-mediated Protection and Defective IgA B-Cell Memory in Experimental Infection of Adults with Respiratory Syncytial Virus. *American journal of respiratory and critical care medicine*. 2015; 191(9):1040-9. doi: 10.1164/rccm.201412-22560C. PubMed PMID: 25730467; PubMed Central PMCID: PMC4435460). Unlike other viral pathogens, serum antibody levels are very slow to rise following RSV infection, with a gradual accumulation of protective antibodies only after multiple re-infections (Wagner D K, Muelenaer P, Henderson F W, Snyder M H, Reimer C B, Walsh E E, et al. Serum immunoglobulin G antibody subclass response to respiratory syncytial virus F and G glycoproteins after first, second, and third infections. *Journal of clinical microbiology*. 1989; 27(3):589-92. Epub 1989/03/01. PubMed PMID: 2715331; PubMed Central PMCID: PMC267370; Welliver R C, Kaul T N, Putnam T I, Sun M, Riddlesberger K, Ogra P L. The antibody response to primary and secondary infection with respiratory syncytial virus: kinetics of class-specific responses. *The Journal of pediatrics*. 1980; 96(5):808-13. Epub 1980/05/01. PubMed PMID: 7365579). The inability of RSV to induce robust immunity following repeated natural infections likely underlies the difficulties encountered in attempts to design effective, attenuated vaccine strains (Crowe J E, Jr. Respiratory syncytial virus vaccine development. *Vaccine*. 2001; 20 Suppl 1:S32-7. Epub 2001/10/06. PubMed PMID: 11587807).

(5) A recent study suggests that the ability of RSV to re-infect immunocompetent adults is correlated with a defect in B cell memory. Neutralizing serum and nasal antibody levels in experimentally infected healthy adult volunteers increased post-infection, but returned to baseline levels 6 months later (Habibi M S, Jozwik A, Makris S, Dunning J, Paras A, DeVincenzo J P, et al. Impaired Antibody-mediated Protection and Defective IgA B-Cell Memory in Experimental Infection of Adults with Respiratory Syncytial Virus. *American journal of respiratory and critical care medicine*. 2015; 191(9):1040-9. doi: 10.1164/rccm.201412-22560C. PubMed PMID: 25730467; PubMed Central PMCID: PMC4435460). This differs markedly from the lifelong persistence of influenza specific IgG that follows this infection (Fujimoto C, Takeda N, Matsunaga A, Sawada A, Tanaka T, Kimoto T, et al. Induction and maintenance of anti-influenza antigen-specific nasal secretory IgA levels and serum IgG levels after influenza infection in adults. *Influenza Other Respir Viruses*. 2012; 6(6):396-403. doi: 10.1111/j.1750-2659.2011.00330.x. PubMed PMID: 22226319; PubMed Central PMCID: PMC4941696). Thus, it is important that any RSV vaccine candidate induce both systemic and mucosal RSV-specific antibodies, but do so in a manner that induces a B cell memory response superior to that induced by RSV infection.

(6) While an effective RSV vaccine has been eagerly sought, no licensed vaccine yet exists. RSV vaccine development has focused primarily on live-attenuation, a strategy with demonstrated safety in infants, the target population for most RSV vaccines. Unfortunately, live-attenuated RSV vaccines evaluated clinically to date have been poorly immunogenic, an outcome that could be predicted from the poor immune response to natural RSV infection (Gomez M, Mufson M A, Dubovsky F, Knightly C, Zeng W, Losonsky G. Phase-I study MEDI-534, of a live, attenuated intranasal vaccine against respiratory syncytial virus and parainfluenza-3 virus in seropositive children. *The Pediatric infectious disease journal*. 2009; 28(7):655-8. Epub 2009/06/02. doi: 10.1097/INF.0b013e318199c3b1. PubMed PMID: 19483659; Wright P F, Belshe R B, Kim H W, Van Voris L P, Chanock R M. Administration of a highly attenuated, live respiratory syncytial virus vaccine to adults and children. *Infection and immunity*. 1982; 37(1):397-400. Epub 1982/07/01. PubMed PMID: 7107009; PubMed Central PMCID: PMC347542; Wright P F, Shinozaki T, Fleet W, Sell S H, Thompson J, Karzon D T. Evaluation of a live, attenuated respiratory syncytial virus vaccine in infants. *The Journal of pediatrics*. 1976; 88(6):931-6. Epub 1976/06/01. PubMed PMID: 178852). Disappointing immunogenicity of live-attenuated RSV vaccines has led investigators to diversify RSV vaccine strategies (reviewed in Mazur N I, Martinon-Torres F, Baraldi E, Fauroux B, Greenough A, Heikkinen T, et al. Lower respiratory tract infection caused by respiratory syncytial virus: current management and new therapeutics. *Lancet Respir Med*. 2015; 3(11):888-900. doi:

10.1016/52213-2600(15)00255-6. PubMed PMID: 26411809).

(7) An important goal of pre-clinical vaccine evaluation is to establish the safety profile of a vaccine candidate. This is particularly true for RSV vaccine candidates given the history of the 1960's vaccine trial in which enhanced pathology following natural RSV infection of infants that had previously received a formalin-inactivated RSV vaccine (FI-RSV) (Kim H W, Canchola J G, Brandt C D, Pyles G, Chanock R M, Jensen K, et al. Respiratory syncytial virus disease in infants despite prior administration of antigenic inactivated vaccine. *American journal of epidemiology*. 1969; 89(4):422-34. Epub 1969/04/01. PubMed PMID: 4305198). It has been suggested that this vaccine-enhanced disease may have been related to the presence of non-neutralizing antibody, allowing virus replication and simultaneous immune complex deposition in the lung (Polack F P, Teng M N, Collins P L, Prince G A, Exner M, Regele H, et al. A role for immune complexes in enhanced respiratory syncytial virus disease. *The Journal of experimental medicine*. 2002; 196(6):859-65. PubMed PMID: 12235218; PubMed Central PMCID: PMCPMC2194058). FI-RSV has been shown to generate primarily non-neutralizing antibody in mice, and RSV challenge of mice receiving the FI-RSV vaccine resulted in immune complex mediated pathology (Id.). It is therefore important to demonstrate the induction of long-lasting, virus-specific adaptive immunity by any vaccine candidate, but also to verify the absence of disease-promoting immune responses.

(8) 2.2 HMPV

(9) Human metapneumovirus (hMPV) is a nonsegmented negative strand RNA virus in the Pneumovirinae family that also includes human respiratory syncytial virus (RSV) (Afonso et al., 2016, *Arch. Virol.* 161: 2351-2360). hMPV was first isolated in the Netherlands in 2001 (van den Hoogen B G, de Jong J C, Groen J, Kuiken T, de Groot R, Fouchier R A, Osterhaus A D. 2001. "A newly discovered human pneumovirus isolated from young children with respiratory tract disease." *Nat Med* 7:719-724) and has subsequently been identified worldwide as second to RSV in causing respiratory disease and hospitalization in infants, with also a significant impact in young kids together with RSV and influenza viruses (Esper F, Boucher D, Weibel C, Martinello R A, Kahn J S. 2003. Human metapneumovirus infection in the United States: clinical manifestations associated with a newly emerging respiratory infection in children. *Pediatrics* 111:1407-1410). In fact, hMPV was found to be a major pathogen associated with hospitalization of children and with the same severity of illness as RSV but in a slightly older population (Akhras N, Weinberg J B, Newton D. 2010. Human metapneumovirus and respiratory syncytial virus: subtle differences but comparable severity. *Infect Dis Rep* 2:e12). Both RSV and hMPV are an important cause of severe respiratory disease in immunocompromised patients. No vaccines are available at this moment for these two important human respiratory pathogens. Thus, a need exists for a safe and effective vaccine for RSV and hMPV.

(10) 2.3 NDV

(11) Newcastle disease virus (NDV) is a member of the Avulavirus genus in the Paramyxoviridae family, which has been shown to infect a number of avian species (Alexander, DJ (1988). *Newcastle disease, Newcastle disease virus—an avian paramyxovirus*. Kluwer Academic Publishers: Dordrecht, The Netherlands. pp 1-22). NDV possesses a single-stranded RNA genome in negative sense and does not undergo recombination with the host genome or with other viruses (Alexander, DJ (1988). *Newcastle disease, Newcastle disease virus—an avian paramyxovirus*. Kluwer Academic Publishers: Dordrecht, The Netherlands. pp 1-22). The genomic RNA contains genes in the order of 3'-NP-P-M-F-HN-L-5', described in further detail below. Two additional proteins, V and W, are produced by NDV from the P gene by alternative mRNAs that are generated by RNA editing. The genomic RNA also contains a leader sequence at the 3' end.

(12) The structural elements of the virion include the virus envelope which is a lipid bilayer derived from the cell plasma membrane. The glycoprotein, hemagglutinin-neuraminidase (HN) protrudes from the envelope allowing the virus to contain both hemagglutinin (e.g., receptor binding/fusogenic) and neuraminidase activities. The fusion glycoprotein (F), which also interacts

with the viral membrane, is first produced as an inactive precursor, then cleaved post-translationally to produce two disulfide linked polypeptides. The active F protein is involved in penetration of NDV into host cells by facilitating fusion of the viral envelope with the host cell plasma membrane. The matrix protein (M), is involved with viral assembly, and interacts with both the viral membrane as well as the nucleocapsid proteins.

(13) The main protein subunit of the nucleocapsid is the nucleocapsid protein (NP) which confers helical symmetry on the capsid. In association with the nucleocapsid are the P and L proteins. The phosphoprotein (P), which is subject to phosphorylation, is thought to play a regulatory role in transcription, and may also be involved in methylation, phosphorylation and polyadenylation. The L gene, which encodes an RNA-dependent RNA polymerase, is required for viral RNA synthesis together with the P protein. The L protein, which takes up nearly half of the coding capacity of the viral genome is the largest of the viral proteins, and plays an important role in both transcription and replication. The V protein has been shown to inhibit interferon-alpha and to contribute to the virulence of NDV (Huang et al. (2003). Newcastle disease virus V protein is associated with viral pathogenesis and functions as an Alpha Interferon Antagonist. *Journal of Virology* 77: 8676-8685).

3. SUMMARY

(14) In one aspect, presented herein are recombinant Newcastle disease virus ("NDV") comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a respiratory syncytial virus ("RSV") F protein. The RSV F protein may be an RSV F protein of a human or bovine strain of RSV. In a specific embodiment, the transgene encodes the human RSV F protein comprising the amino acid sequence set forth in SEQ ID NO: 6 or the bovine RSV F protein comprising the amino acid sequence set forth in SEQ ID NO: 10. Due to the degeneracy of the nucleic acid code, multiple different nucleic acid sequences may encode for the same human RSV F protein or the same bovine RSV F protein. In one embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a human RSV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:1. In another embodiment described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a human RSV F protein comprising the amino acid sequence set forth in SEQ ID NO:49, 50 or 58. In another embodiment described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a human RSV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 25, 27 or 29. In another embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a bovine RSV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:9. In another embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a bovine RSV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:40 or 42. In a preferred embodiment, a transgene comprises a codon optimized version of a nucleic acid sequence encoding an RSV F protein (e.g., a human or bovine RSV F protein). In a specific embodiment, the RSV F protein is expressed by cells infected with the recombinant NDV.

(15) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene comprising a codon optimized nucleic acid sequence encoding a human RSV F protein. Specific examples of codon optimized nucleic acid sequences encoding a human RSV F protein include SEQ ID NO: 2, 26, 28 or 30. In a preferred embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a human RSV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:2. In a specific embodiment, the human RSV F protein is expressed by cells infected with the recombinant NDV.

(16) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene comprising a codon optimized nucleic acid sequence encoding a bovine RSV F protein. Specific examples of codon optimized nucleic acid sequences encoding a bovine RSV F protein include SEQ ID NO: 11, 41, or 43. In a preferred embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a bovine RSV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:11. In a specific embodiment, the bovine RSV F protein is expressed by cells infected with the recombinant NDV.

(17) In another embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In other words, the NDV F protein transmembrane and cytoplasmic domains replace the RSV F protein transmembrane and cytoplasmic domains so that the chimeric F protein does not include the RSV F protein transmembrane and cytoplasmic domains. The RSV F protein may be an RSV F protein of a human or bovine strain of RSV. In a specific embodiment, the transgene encodes a chimeric F protein comprising the amino acid sequence set forth in SEQ ID NO: 7 or 33. In another embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:51. In another embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 31. In a preferred embodiment, a transgene comprises a codon optimized version of a nucleic acid sequence encoding the RSV F protein ectodomain (e.g., a human or bovine RSV F protein ectodomain). In a specific embodiment, the RSV F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV and the chimeric F protein is incorporated into the NDV virion.

(18) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence. Specific examples of nucleic acid sequences encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence, include SEQ ID NO:4, 44, 45 or 46. In a preferred embodiment, described herein is a recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:4. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV and the chimeric F protein is incorporated into the NDV virion.

(19) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein,

wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence. Specific examples of nucleic acid sequences encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence, include SEQ ID NO:14, 38 or 39. In a preferred embodiment, described herein is a recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:14. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV and the chimeric F protein is incorporated into the NDV virion.

(20) The recombinant NDV may have the backbone of any NDV type or strain, including, but not limited to, naturally-occurring strains, variants or mutants, mutagenized viruses, reassortants or genetically engineered viruses, or any combination thereof. The NDV that serves as the backbone for genetic engineering of the recombinant NDV may be a lentogenic strain, a mesogenic strain, or a velogenic strain. In a specific embodiment, the recombinant NDV comprises an NDV backbone which is lentogenic. In another specific embodiment, the recombinant NDV comprises an NDV backbone of the NDV LaSota strain. See, e.g., SEQ ID NO: 47 for a cDNA sequence of the genomic sequence of NDV LaSota strain. In another specific embodiment, the recombinant NDV comprises an NDV backbone of the NDV Hitchner B1 strain. See, e.g., SEQ ID NO: 48 for a cDNA sequence of the genomic sequence of NDV Hitchner strain. In another specific embodiment, the recombinant NDV comprises an NDV backbone of a lentogenic strain other than the NDV Hitchner B1 strain. In another specific embodiment, the recombinant NDV comprises an NDV backbone of the NDV Fuller strain. In another specific embodiment, the recombinant NDV comprises an NDV backbone of the NDV Ulster strain.

(21) The transgene encoding a RSV F protein or a chimeric F protein may be incorporated into the genome of any NDV type or strain. In a specific embodiment, the transgene is incorporated into the genome of a lentogenic NDV. In another specific embodiment, the transgene is incorporated in the genome of NDV strain LaSota. See, e.g., SEQ ID NO: 47 for a cDNA sequence of the genomic sequence of NDV LaSota strain. Other examples of NDV strains into which the transgene may be incorporated are the NDV Fuller, the NDV Ulster strain or the NDV Hitchner B1 strain. In a specific embodiment, the transgene may be incorporated into the genomic sequence of NDV Hitchner B1 strain. See, e.g., SEQ ID NO:48 for a cDNA sequence of the genomic sequence of NDV Hitchner B1 strain. In a specific embodiment, the transgene may be incorporated into the genome of a lentogenic strain other than the NDV Hitchner B1 strain. The transgene may be incorporated into the NDV genome between two transcription units (e.g., between NDV P and M genes).

(22) In a specific embodiment, a transgene encoding a human RSV F protein is incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). For example, a transgene encoding the human RSV F protein set forth in SEQ ID NO:6 may be incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). In a specific embodiment, such a transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO:1. However, given the degeneracy of the nucleic acid code, there are a number of different nucleic acid sequences that may encode the same human RSV F protein. In certain embodiments, a transgene encoding the human RSV F protein set forth in SEQ ID NO:49, 50 or 58 may be incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). In some embodiments, a transgene encoding the human RSV F protein comprises the amino acid sequence

encoded by the nucleic acid sequence comprising the sequence of SEQ ID NO:25, 27 or 29. In a specific embodiment, transgene encoding a human RSV F protein is codon optimized. For example, the human RSV F protein is encoded by the codon optimized nucleic acid sequence set forth in SEQ ID NO: 2, 26, 28 or 30. The transgene encoding a human RSV F protein may be incorporated between any two NDV transcription units (e.g., between NDV P and M genes). In a specific embodiment, a recombinant NDV described herein comprises a packaged genome, wherein the packaged genome corresponds to a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO: 3. In certain embodiment, the genome of the recombinant NDV does not comprise a heterologous sequence encoding a heterologous protein other than the human RSV F protein. In some embodiments, the genome of the recombinant NDV does not comprise a transgene other than a transgene encoding a human RSV F protein. In certain embodiments, the genome of the recombinant NDV comprises a transgene encoding a human RSV F protein and a transgene encoding an hMPV F protein or a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In such embodiments, the genome of the recombinant NDV may not comprise any additional transgenes.

(23) In a specific embodiment, a transgene encoding a bovine RSV F protein is incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). For example, a transgene encoding the bovine RSV F protein comprising the amino acid sequence set forth in SEQ ID NO:10 may be incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). In a specific embodiment, such a transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO:9. However, given the degeneracy of the nucleic acid code, there are a number of different nucleic acid sequences that may encode the same bovine RSV F protein. In some embodiments, a transgene encoding the bovine RSV F protein comprises the amino acid sequence encoded by the nucleic acid sequence comprising the sequence of SEQ ID NO:40 or 42. In a specific embodiment, transgene encoding a bovine RSV F protein is codon optimized. For example, the bovine RSV F protein is encoded by the codon optimized nucleic acid sequence set forth in SEQ ID NO: 11, 41 or 43. The transgene encoding a bovine RSV F protein may be incorporated between any two NDV transcription units (e.g., between NDV P and M genes). In a specific embodiment, a recombinant NDV described herein comprises a packaged genome, wherein the packaged genome is an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO: 12 or 13. In certain embodiment, the genome of the recombinant NDV does not comprise a heterologous sequence encoding a heterologous protein other than the bovine RSV F protein. In some embodiments, the genome of the recombinant NDV does not comprise a transgene other than a transgene encoding a bovine RSV F protein.

(24) In a specific embodiment, a transgene encoding a chimeric F protein is incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain), wherein the chimeric F protein comprises the ectodomain of an RSV F protein (e.g., a human or bovine RSV F protein) and NDV F protein transmembrane and cytoplasmic domains. For example, a transgene encoding the chimeric F protein comprising the amino acid sequence set forth in SEQ ID NO:7 may be incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). In a specific embodiment, such a transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO: 51. However, given the degeneracy of the nucleic acid code, there are a number of different nucleic acid sequences that may encode the same chimeric F protein. In a specific embodiment, the nucleic acid sequence encoding a human RSV F protein ectodomain is codon optimized. For example, the nucleic acid sequence of one or more of the domains of the chimeric F protein may be codon optimized (e.g. the ectodomain of the chimeric F protein may be encoded by a codon optimized nucleic acid sequence, such as set forth in SEQ ID NO: 4, 44, 45, or 46. The transgene encoding a chimeric F protein may be incorporated between any two NDV transcription units (e.g., between NDV P and M genes). In a specific embodiment, a

recombinant NDV described herein comprises a packaged genome, wherein the packaged genome corresponds to a negative sense RNA sequence transcribed from a cDNA sequence comprising the sequence set forth in SEQ ID NO: 5. In certain embodiments, the genome of the recombinant NDV does not comprise a heterologous sequence encoding a heterologous protein other than the chimeric F protein. In some embodiments, the genome of the recombinant NDV does not comprise a transgene other than a transgene encoding a chimeric F protein. In certain embodiments, the genome of the recombinant NDV comprises a transgene encoding a first chimeric F protein and a transgene encoding an hMPV F protein or a second chimeric F protein, wherein the first chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains and the second chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In such embodiments, the genome of the recombinant NDV may not comprise any additional transgenes.

(25) In a specific embodiment, a transgene encoding a chimeric F protein is incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain), wherein the chimeric F protein comprises the ectodomain of an RSV F protein (e.g., a human or bovine RSV F protein) and NDV F protein transmembrane and cytoplasmic domains. For example, a transgene encoding the chimeric F protein comprising the amino acid sequence set forth in SEQ ID NO:33 may be incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). In a specific embodiment, such a transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO: 31. However, given the degeneracy of the nucleic acid code, there are a number of different nucleic acid sequences that may encode the same chimeric F protein. In a specific embodiment, the nucleic acid sequence encoding a bovine RSV F protein ectodomain is codon optimized. For example, the nucleic acid sequence of one or more of the domains of the chimeric F protein may be codon optimized (e.g. the ectodomain of the chimeric F protein may be encoded by a codon optimized nucleic acid sequence, such as set forth in SEQ ID NO: 14, 38, or 39. The transgene encoding a chimeric F protein may be incorporated between any two NDV transcription units (e.g., between NDV P and M genes). In certain embodiment, the genome of the recombinant NDV does not comprise a heterologous sequence encoding a heterologous protein other than the chimeric F protein. In some embodiments, the genome of the recombinant NDV does not comprise a transgene other than a transgene encoding a chimeric F protein. In certain embodiments, the genome of the recombinant NDV comprises a transgene encoding a first chimeric F protein and a transgene encoding an hMPV F protein or a second chimeric F protein, wherein the first chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains and the second chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In such embodiments, the genome of the recombinant NDV may not comprise any additional transgenes.

(26) The recombinant NDV and uses thereof described herein are based, in part, on the robust, long-lived antigen-specific humoral immunity induced by the recombinant NDV and the prevention of RSV infection of the lower airway and the largely eliminated RSV replication in the upper airway in animals. In addition, the recombinant NDV and uses thereof described herein are based, in part, on the demonstration of vaccine efficacy in previously-infected, seropositive animals. Further, the recombinant NDV and uses thereof described herein are based, in part, on the decreased inflammation induced by the recombinant NDV with the NDV LaSota strain backbone and the enhanced RSV F protein expression resulting from the use of a codon optimized nucleic acid sequence encoding RSV F protein or a nucleic acid sequence encoding a chimeric F protein.

(27) In another aspect, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a human RSV F protein. In one embodiment, the human RSV F protein comprises the amino acid sequence

set forth in SEQ ID NO: 6, 49, 50 or 58. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same human RSV F protein. For example, the human RSV F protein may be encoded a sequence comprising the amino acid sequence set forth SEQ ID NO:1, 25, 27 or 29. Alternatively, the nucleic acid sequence encoding the human RSV F protein may be codon optimized, such as set forth in SEQ ID NO: 2, 26, 28 or 30. In some embodiments, the human RSV F protein is encoded by a sequence comprising the nucleic acid sequence set forth in SEQ ID NO:1, 2, 25, 26, 27, 28, 29, or 30. In a specific embodiment, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a human RSV F protein. In a specific embodiment, the human RSV F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human infant six months old or older. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(28) In another aspect, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a bovine RSV F protein. In one embodiment, the bovine RSV F protein comprises the amino acid sequence set forth in SEQ ID NO: 10. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same bovine RSV F protein. For example, the bovine RSV F protein may be encoded by SEQ ID NO:9. Alternatively, the nucleic acid sequence encoding the bovine RSV F protein may be codon optimized, such as set forth in SEQ ID NO: 11. In some embodiments, the bovine RSV F protein is encoded by a sequence comprising the nucleic acid sequence set forth in SEQ ID NO:9, 11, 40, 41, 42, or 43. In a specific embodiment, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a bovine RSV F protein. In a specific embodiment, the bovine RSV F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(29) In another aspect, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO: 7. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:51. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO:4, 44, 45, or 46). In a specific embodiment, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises

a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human infant six months old or older. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(30) In another aspect, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:33. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:31. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO:14, 38, or 39). In a specific embodiment, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(31) In another aspect, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a human RSV F protein. In one embodiment, the human RSV F protein comprises the amino acid sequence set forth in SEQ ID NO: 6, 49, 50 or 58. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same human RSV F protein. For example, the human RSV F protein may be encoded by SEQ ID NO:1, 25, 27 or 29. Alternatively, the nucleic acid sequence encoding the human RSV F protein may be codon optimized, such as set forth in SEQ ID NO:2, 26, 28, or 30. In some embodiments, the human RSV F protein is encoded by a nucleic acid sequence comprising the sequence of SEQ ID NO: 1, 2, 25, 26, 27, 28, 29, or 30. In a specific embodiment, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a human RSV F protein. In a specific embodiment, the human RSV F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human

infant six months old or older. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(32) In another aspect, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a bovine RSV F protein. In one embodiment, the bovine RSV F protein comprises the amino acid sequence set forth in SEQ ID NO: 10. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same bovine RSV F protein. For example, the bovine RSV F protein may be encoded by SEQ ID NO:9. Alternatively, the nucleic acid sequence encoding the bovine RSV F protein may be codon optimized, such as set forth in SEQ ID NO: 11. In some embodiments, the bovine RSV F protein is encoded by a sequence comprising the nucleic acid sequence set forth in SEQ ID NO:9, 11, 40, 41, 42 or 43. In a specific embodiment, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a bovine RSV F protein. In a specific embodiment, the bovine RSV F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(33) In another aspect, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:7. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:51. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO:4, 44, 45, or 46). In a specific embodiment, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human infant six months old or older. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(34) In another aspect, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant

NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:33. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:31. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO:14, 38, or 39). In a specific embodiment, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(35) In another aspect, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a human RSV F protein. In one embodiment, the human RSV F protein comprises the amino acid sequence set forth in SEQ ID NO: 6, 49, 50 or 58. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same human RSV F protein. For example, the human RSV F protein may be encoded by SEQ ID NO:1, 25, 27 or 29. Alternatively, the nucleic acid sequence encoding the human RSV F protein may be codon optimized, such as set forth in SEQ ID NO: 2, 26, 28 or 30. In some embodiments, the human RSV F protein is encoded by a nucleic acid sequence comprising the sequence of SEQ ID NO: 1, 2, 25, 26, 27, 28, 29, or 30. In a specific embodiment, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a human RSV F protein. In a specific embodiment, the human RSV F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human infant six months old or older. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(36) In another aspect, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a bovine RSV F protein. In one embodiment, the bovine RSV F protein comprises the amino acid sequence set forth in SEQ ID NO: 10. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same bovine RSV F protein. For example, the bovine RSV F protein may be encoded by SEQ ID NO:9. Alternatively, the nucleic acid sequence encoding the bovine RSV F protein may be codon optimized, such as set forth in SEQ ID NO: 11. In some embodiments, the bovine RSV F protein is encoded by a nucleic acid sequence comprising the sequence of SEQ ID NO: 9, 11, 40, 41, 42, or 43. In a specific embodiment, presented herein are

methods for the prevention of RSV disease comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a bovine RSV F protein. In a specific embodiment, the bovine RSV F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(37) In another aspect, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO: 7. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:51. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO: 4, 44, 45, or 46). In a specific embodiment, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human infant six months old or older. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(38) In another aspect, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:33. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:31. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO:14, 38, or 39). In a specific embodiment, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected

with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is seropositive for antibodies to RSV (e.g., the subject is seropositive for antibodies to RSV F protein or RSV G protein).

(39) The recombinant NDV described herein may be administered to a subject in combination with one or more other therapies. The recombinant NDV and one or more other therapies may be administered by the same or different routes of administration to the subject. In a specific embodiment, the recombinant NDV is administered to a subject intranasally. See, e.g., Sections 5.15.1, and 6, *infra* for information regarding recombinant NDV, Section 5.5.4 for information regarding other therapies, Section 5.4, *infra*, for information regarding compositions and routes of administration, and Sections 5.5.1 and 66, *infra*, for information regarding methods of immunizing against RSV.

(40) In another embodiment, presented herein is a recombinant NDV or composition thereof for inducing an immune response to an RSV F protein. In another embodiment, presented herein is a recombinant NDV or composition thereof for immunizing a subject (e.g., a human or bovine subject) against RSV. In another embodiment, presented herein is a recombinant NDV or composition thereof for the prevention of an RSV disease. See, e.g., Sections 5.1, and 6, *infra* for information regarding recombinant NDV, Section 5.5.4, for information regarding other therapies, Section 5.4, *infra*, for information regarding compositions and routes of administration, and Sections 5.5.1 and 6, *infra*, for information regarding methods of immunizing against RSV.

(41) In another aspect, presented herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a human metapneumovirus (“hMPV”) F protein. The hMPV F protein may be an hMPV F protein of any strain of hMPV. In a specific embodiment, the transgene encodes the hMPV F protein set forth in SEQ ID NO:17. Due to the degeneracy of the nucleic acid code, multiple different nucleic acid sequences may encode for the same hMPV F protein. In one embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a hMPV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:16. In some embodiments, the hMPV F protein is encoded by a nucleic acid sequence comprising the sequence of SEQ ID NO:16, 52, 54, or 56. In a preferred embodiment, a transgene comprises a codon optimized version of a nucleic acid sequence encoding an hMPV F protein. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV.

(42) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene comprising a codon optimized nucleic acid sequence encoding a hMPV F protein. Specific examples of codon optimized nucleic acid sequences encoding a hMPV F protein include those set forth in SEQ ID NO:18, 53, 55, or 57. In a specific embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a human RSV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:18. In a specific embodiment, the human RSV F protein is expressed by cells infected with the recombinant NDV.

(43) In another embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In other words, the NDV F protein transmembrane and cytoplasmic domains replace the hMPV F protein transmembrane and cytoplasmic domains so that the chimeric F protein does not include the hMPV F protein transmembrane and cytoplasmic domains. The hMPV F protein may be an hMPV F protein of any of hMPV. In another embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F

protein comprises a hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:32. In a preferred embodiment, a transgene comprises a codon optimized version of a nucleic acid sequence encoding the hMPV F protein ectodomain. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV.

(44) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the hMPV F protein is encoded by a codon optimized nucleic acid sequence. Specific examples of nucleic acid sequences encoding a chimeric F protein, wherein the chimeric F protein comprises a hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the hMPV F protein is encoded by a codon optimized nucleic acid sequence, include SEQ ID NO:19, 34, 35, or 36. In a specific embodiment, described herein is a recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:19. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV and the chimeric F protein is incorporated into the NDV virion.

(45) The recombinant NDV may have the backbone of any NDV type or strain, including, but not limited to, naturally-occurring strains, variants or mutants, mutagenized viruses, reassortants, genetically engineered viruses, or any combination thereof. The NDV that serves as the backbone for genetic engineering of the recombinant NDV may be a lentogenic strain, a mesogenic strain, or a velogenic strain. In a specific embodiment, the recombinant NDV comprises an NDV backbone which is lentogenic. In another specific embodiment, the recombinant NDV comprises an NDV backbone of the NDV LaSota strain. In another specific embodiment, the recombinant NDV comprises an NDV backbone of the NDV Hitchner B1 strain. In another specific embodiment, the recombinant NDV comprises an NDV backbone of a lentogenic strain other than the NDV Hitchner B1 strain. In another specific embodiment, the recombinant NDV comprises an NDV backbone of the NDV Fuller strain. In another specific embodiment, the recombinant NDV comprises an NDV backbone of the NDV Ulster strain.

(46) The transgene encoding a hMPV F protein or a chimeric F protein may be incorporated into the genome of any NDV type or strain. In a specific embodiment, the transgene is incorporated into the genome of a lentogenic NDV. In another specific embodiment, the transgene is incorporated in the genome of NDV strain LaSota. Other examples of NDV strains into which the transgene may be incorporated are the NDV Fuller, the NDV Ulster strain or the NDV Hitchner B1 strain. The transgene may be incorporated into the NDV genome between two transcription units (e.g., between NDV P and M genes).

(47) In a specific embodiment, a transgene encoding a hMPV F protein is incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). For example, a transgene encoding the hMPV F protein comprising the amino acid sequence set forth in SEQ ID NO:17 may be incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). In a specific embodiment, such a transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO:16. However, given the degeneracy of the nucleic acid code, there are a number of different nucleic acid sequences that may encode the same hMPV F protein. In a specific embodiment, transgene encoding a hMPV F protein is codon optimized. For example, the hMPV F protein is encoded by the codon optimized nucleic acid sequence set forth in

SEQ ID NO: 18. In some embodiments, the hMPV F protein is encoded by a nucleic acid sequence comprising the sequence of SEQ ID NO: 16, 18, 52, 53, 54, 55, 56, or 57. The transgene encoding a hMPV F protein may be incorporated between any two NDV transcription units (e.g., between NDV P and M genes). In a specific embodiment, a recombinant NDV described herein comprises a packaged genome, wherein the packaged genome corresponds to a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO: 20 or 21. In certain embodiment, the genome of the recombinant NDV does not comprise a heterologous sequence encoding a heterologous protein other than the hMPV F protein. In some embodiments, the genome of the recombinant NDV does not comprise a transgene other than a transgene encoding a hMPV F protein.

(48) In a specific embodiment, a transgene encoding a chimeric F protein is incorporated into the genome of any NDV type or strain, wherein the chimeric F protein comprises the ectodomain of an hMPV F protein and NDV F protein transmembrane and cytoplasmic domains. For example, a transgene encoding the chimeric F protein comprising the amino acid sequence set forth in SEQ ID NO:15 may be incorporated into the genome of any NDV type or strain. In a specific embodiment, such a transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO:32. However, given the degeneracy of the nucleic acid code, there are a number of different nucleic acid sequences that may encode the same chimeric F protein. In a specific embodiment, the nucleic acid sequence encoding the ectodomain of the hMPV F protein is codon optimized. For example, the chimeric F protein may be encoded by the codon optimized nucleic acid sequence, such as set forth in SEQ ID NO: 19, 34, 35, or 36. The nucleotide sequence comprising the nucleic acid sequence encoding a chimeric F protein may be incorporated between any two NDV transcription units (e.g., between NDV P and M genes). In a specific embodiment, a recombinant NDV described herein comprises a packaged genome, wherein the packaged genome corresponds to a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO:22. In certain embodiment, the genome of the recombinant NDV does not comprise a heterologous sequence encoding a heterologous protein other than the chimeric F protein. In some embodiments, the genome of the recombinant NDV does not comprise a transgene other than a transgene encoding a chimeric F protein.

(49) In another aspect, presented herein are methods for inducing an immune response to an hMPV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a hMPV F protein. In one embodiment, the hMPV F protein comprises the amino acid sequence set forth in SEQ ID NO: 17. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same hMPV F protein. For example, the hMPV F protein may be encoded by SEQ ID NO:16. Alternatively, the nucleic acid sequence encoding the hMPV F protein may be codon optimized, such as set forth in SEQ ID NO: 18. In some embodiments, the hMPV F protein is encoded by a nucleic acid sequence comprising the sequence of SEQ ID NO: 16, 18, 52, 53, 54, 55, 56 or 57. In a specific embodiment, presented herein are methods for inducing an immune response to an hMPV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding an hMPV F protein. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to hMPV.

(50) In another aspect, presented herein are methods for inducing an immune response to an hMPV

F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:15. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:32. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the hMPV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO: 19, 34, 35 or 36). In a specific embodiment, presented herein are methods for inducing an immune response to an hMPV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise an hMPV RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the hMPV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to hMPV.

(51) In another aspect, presented herein are methods for immunizing against hMPV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a hMPV F protein. In one embodiment, the hMPV F protein comprises the amino acid sequence set forth in SEQ ID NO: 17. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same hMPV F protein. For example, the hMPV F protein may be encoded by SEQ ID NO:16. Alternatively, the nucleic acid sequence encoding the hMPV F protein may be codon optimized, such as set forth in SEQ ID NO: 18. In some embodiments, the hMPV F protein is encoded by a nucleic acid sequence comprising the sequence of SEQ ID NO: 16, 18, 52, 53, 54, 55, 56 or 57. In a specific embodiment, presented herein are methods for immunizing against hMPV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding an hMPV F protein. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to hMPV.

(52) In another aspect, presented herein are methods for immunizing against hMPV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:15. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein.

For example, the chimeric F protein may be encoded by SEQ ID NO:32. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the hMPV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO: 19, 34, 35, or 36). In a specific embodiment, presented herein are methods for immunizing against hMPV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise an hMPV RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the hMPV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV and the chimeric F protein is incorporated into the NDV virion. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to hMPV.

(53) In another aspect, presented herein are methods for the prevention of hMPV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a hMPV F protein. In one embodiment, the hMPV F protein comprises the amino acid sequence set forth in SEQ ID NO: 17. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same hMPV F protein. For example, the hMPV F protein may be encoded by SEQ ID NO:16. Alternatively, the nucleic acid sequence encoding the hMPV F protein may be codon optimized, such as set forth in SEQ ID NO: 18. In some embodiments, the hMPV F protein is encoded by a nucleic acid sequence comprising the sequence of SEQ ID NO: 16, 18, 52, 53, 54, 55, 56 or 57. In a specific embodiment, presented herein are methods for the prevention of hMPV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding an hMPV F protein. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to hMPV.

(54) In another aspect, presented herein are methods for the prevention of hMPV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:15. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:32. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the hMPV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO: 19, 34, 35, or 36). In a specific embodiment, presented herein are methods for the prevention of hMPV disease comprising administering to a subject (e.g.,

a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise an hMPV RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the hMPV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV and the chimeric F protein is incorporated into the NDV virion. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In another specific embodiment, the subject is a human infant. In another specific embodiment, the subject is a human toddler. In another specific embodiment, the subject is a human child. In another specific embodiment, the subject is a human adult. In another specific embodiment, the subject is an elderly human. In another specific embodiment, the subject is seropositive for antibodies to hMPV.

(55) The recombinant NDV described herein may be administered to a subject in combination with one or more other therapies. The recombinant NDV and one or more other therapies may be administered by the same or different routes of administration to the subject. In a specific embodiment, the recombinant NDV is administered to a subject intranasally. See, e.g., Sections 5.15.1, and 6, *infra* for information regarding recombinant NDV, Section 5.5.4 for information regarding other therapies, Section 5.4, *infra*, for information regarding compositions and routes of administration, and Sections 5.5.2 5.5.1 and 6, *infra*, for information regarding methods of immunizing against hMPV.

(56) In another embodiment, presented herein is a recombinant NDV or composition thereof for inducing an immune response to an hMPV F protein. In another embodiment, presented herein is a recombinant NDV or composition thereof for immunizing a subject (e.g., a human subject) against hMPV. In another embodiment, presented herein is a recombinant NDV or composition thereof for the prevention of an hMPV disease. See, e.g., Sections 5.1, and 6, *infra* for information regarding recombinant NDV, Section 5.5.4 for information regarding other therapies, Section 5.4, *infra*, for information regarding compositions and routes of administration, and Sections 5.5.2 and 6, *infra*, for information regarding methods of immunizing against hMPV.

(57) In another aspect, presented herein are pharmaceutical compositions (e.g., immunogenic compositions) comprising a recombinant NDV described herein, in admixture with a pharmaceutically acceptable carrier. In a specific embodiment, a pharmaceutical composition (e.g., an immunogenic composition) comprises a recombinant NDV described herein as the sole active ingredient, in admixture with a pharmaceutically acceptable carrier. In another specific embodiment, a pharmaceutical composition (e.g., an immunogenic composition) comprises two different recombinant NDVs described herein, in admixture with a pharmaceutically acceptable carrier. See, e.g., Section 5.4, *infra*, for examples of pharmaceutical compositions. The pharmaceutical compositions may be used to induce an immune response to an RSV F protein or hMPV F protein, immunize against RSV or hMPV, or prevent RSV disease or hMPV disease.

(58) 3.1 Terminology

(59) As used herein, the term “about” or “approximately” when used in conjunction with a number refers to any number within 1, 5 or 10% of the referenced number.

(60) As used herein, the terms “antibody” and “antibodies” refer to molecules that contain an antigen binding site, e.g., immunoglobulins. Antibodies include, but are not limited to, monoclonal antibodies, bispecific antibodies, multispecific antibodies, human antibodies, humanized antibodies, synthetic antibodies, chimeric antibodies, polyclonal antibodies, single domain antibodies, camelized antibodies, single-chain Fvs (scFv), single chain antibodies, Fab fragments, F(ab') fragments, disulfide-linked bispecific Fvs (sdFv), intrabodies, and anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id and anti-anti-Id antibodies to antibodies), and epitope-binding fragments of any of the above. In particular, antibodies include immunoglobulin molecules and

immunologically active fragments of immunoglobulin molecules. Immunoglobulin molecules can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass.

(61) As used herein, the term “heterologous” in the context of NDV refers to an entity not found in nature to be associated with (e.g., encoded by, expressed by the genome of, or both) a naturally occurring NDV. In a specific embodiment, a heterologous sequence encodes a protein that is not found associated with naturally occurring NDV.

(62) As used herein, the term “elderly human” refers to a human 65 years or older.

(63) As used herein, the term “human adult” refers to a human that is 18 years or older.

(64) As used herein, the term “human child” refers to a human that is 1 year to 18 years old.

(65) As used herein, the term “human toddler” refers to a human that is 1 year to 3 years old.

(66) As used herein, the term “human infant” refers to a newborn to 1 year old year human.

(67) “Human Metapneumovirus F protein” and “hMPV F protein” refer to any Human Metapneumovirus F protein known to those of skill in the art. The hMPV F protein is synthesized as a F0 inactive precursor. The F0 inactive precursor requires cleavage during intracellular maturation. The hMPV F is cleaved to form F1 and F2. The hMPV F protein exists in two conformations, prefusion and post-fusion. GenBank™ accession number AY145301.1 and KJ627437.1, provide exemplary nucleic acid sequences encoding hMPV F protein. In a preferred embodiment, an hMPV F protein is encoded by a nucleic acid sequence provided in Section 5.1.4, 5.1.5 or 6, e.g., SEQ ID NOs: 16, 18, 52, 53, 54, 55, 56, or 57. GenBank™ accession numbers AAN52915.1, AHV79975.1, AGJ74035.1, and AGZ48845.1 provide exemplary hMPV F protein amino acid sequences. In a preferred embodiment, an hMPV F protein comprises an amino acid sequence provided in Section 5.1.4 or 6, e.g., SEQ ID NOs: 17. As used herein, the terms “hMPV F protein” and “human metapneumovirus F protein” encompass hMPV F polypeptides that are modified by post-translational processing such as signal peptide cleavage, disulfide bond formation, glycosylation (e.g., N-linked glycosylation), protease cleavage and lipid modification (e.g. S-palmitoylation). In some embodiments, the hMPV F protein includes a signal sequence. In other embodiments, hMPV F protein does not include a signal sequence. The signal sequence can be the naturally occurring signal peptide sequence or a variant thereof. The hMPV F protein signal sequence is typically 18 amino acids in length. See, e.g., SEQ ID NO:24 for an exemplary hMPV protein signal sequence. In some embodiments, the signal peptide is an hMPV F protein signal peptide. In some embodiments, the signal peptide is heterologous to an hMPV F protein signal peptide.

(68) As used herein, the phrases “IFN deficient systems” or “IFN-deficient substrates” refer to systems, e.g., cells, cell lines and animals, such as mice, chickens, turkeys, rabbits, rats, horses etc., which do not produce one, two or more types of IFN, or do not produce any type of IFN, or produce low levels of one, two or more types of IFN, or produce low levels of any IFN (i.e., a reduction in any IFN expression of 5-10%, 10-20%, 20-30%, 30-40%, 40-50%, 50-60%, 60-70%, 70-80%, 80-90% or more when compared to IFN-competent systems under the same conditions), do not respond or respond less efficiently to one, two or more types of IFN, or do not respond to any type of IFN, have a delayed response to one, two or more types of IFN, are deficient in the activity of antiviral genes induced by one, two or more types of IFN, or induced by any type of IFN, or any combination thereof.

(69) As used herein, the phrase “multiplicity of infection” or “MOI” is the average number of virus per infected cell. The MOI is determined by dividing the number of virus added (ml added x Pfu/ml) by the number of cells added (Pfu added/cells).

(70) “RSV F protein” and “respiratory syncytial virus F protein” refer to any respiratory syncytial F protein known to those of skill in the art. The RSV F protein typically exists as a homotrimer. The RSV F protein is synthesized as a F0 inactive precursor which is heavily N-glycosylated. The F0 inactive precursor requires cleavage during intracellular maturation by a furin-like proteases. The

RSV F contains two furin sites, and cleavage by furin-like proteases leads to three polypeptides: F2, p27 and F1, with the latter containing a hydrophobic fusion peptide at its N terminus. The RSV F protein exists in two conformations, prefusion and post-fusion. The RSV F protein may be human RSV F protein or bovine F protein. GenBank™ accession numbers KJ155694.1, KU950686.1, KJ672481.1, KP119747, and AF035006.1 provide exemplary nucleic acid sequences encoding human RSV F protein. In a preferred embodiment, a human RSV F protein may be encoded by the nucleic acid sequence provided in Section 5.1.2, 5.1.3. or 6, e.g., SEQ ID NOs: 1, 2, 25, 26, 27, 28, 29 or 30. GenBank™ accession numbers AHL84194.1, AMT79817.1, AHX57603.1, AIY70220.1 and AAC14902.1 provide exemplary human RSV F protein amino acid sequences. In a preferred embodiment, a human RSV F protein comprises the amino sequence provided in Section 5.1.2 or 6, e.g., SEQ ID NO: 6, 49, 50 or 58. GenBank™ accession numbers AF295543.1, AF092942.1, and Y17970.1 provide exemplary nucleic acid sequences encoding bovine RSV F protein. In a specific embodiment, a bovine RSV F protein is encoded by a nucleic acid sequence comprising the nucleic acid sequence of SEQ ID NO:9, 11, 39, 40, 41, 42 or 43. GenBank™ accession numbers AAL49399.1, NP_048055.1, AAC96308.1, and CAA76980.1 provide exemplary bovine RSV F protein amino acid sequences. In another embodiment, a bovine RSV F protein comprises the amino acid sequence provided in Section 5.1.2 or 6, e.g., SEQ ID NO: 10. As used herein, the terms “RSV F protein” and “respiratory syncytial virus F protein” encompass RSV F polypeptides that are modified by post-translational processing such as signal peptide cleavage, disulfide bond formation, glycosylation (e.g., N-linked glycosylation), protease cleavage and lipid modification (e.g. S-palmitoylation). In some embodiments, the RSV F protein includes a signal sequence. In other embodiments, RSV F protein does not include a signal sequence. The signal sequence can be the naturally occurring signal peptide sequence or a variant thereof. The RSV F protein signal sequence is typically 25 amino acids in length. See, e.g., SEQ ID NO:23 or 60 for an exemplary human RSV F protein signal sequence and bovine RSV F protein signal sequence, respectively. In some embodiments, the signal peptide is an RSV F protein signal peptide. In some embodiments, the signal peptide is heterologous to an RSV F protein signal peptide.

(71) As used herein, the terms “subject” or “patient” are used interchangeably. As used herein, the terms “subject” and “subjects” refers to an animal. In some embodiments, the subject is a mammal including a non-primate (e.g., a camel, donkey, zebra, bovine, horse, horse, cat, dog, rat, and mouse) and a primate (e.g., a monkey, chimpanzee, and a human). In some embodiments, the subject is a non-human mammal. In certain embodiments, the subject is a pet (e.g., dog or cat) or farm animal (e.g., a horse, pig or cow). In specific embodiments, the subject is a human. In other specific embodiments, the subject is a bovine. In certain embodiments, the mammal (e.g., human) is 4 to 6 months old, 6 to 12 months old, 1 to 5 years old, 5 to 10 years old, 10 to 15 years old, 15 to 20 years old, 20 to 25 years old, 25 to 30 years old, 30 to 35 years old, 35 to 40 years old, 40 to 45 years old, 45 to 50 years old, 50 to 55 years old, 55 to 60 years old, 60 to 65 years old, 65 to 70 years old, 70 to 75 years old, 75 to 80 years old, 80 to 85 years old, 85 to 90 years old, 90 to 95 years old or 95 to 100 years old. In specific embodiments, the subject is an animal that is not avian.

(72) As used herein, the term “in combination” in the context of the administration of (a) therapy(ies) to a subject, refers to the use of more than one therapy. The use of the term “in combination” does not restrict the order in which therapies are administered to a subject. A first therapy can be administered prior to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second therapy to a subject. For example, a recombinant NDV described herein may be administered prior to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours,

72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of another therapy.

(73) As used herein, the terms “therapies” and “therapy” can refer to any protocol(s), method(s), agent(s) or a combination thereof that can be used in the treatment or prevention of RSV disease or hMPV disease, or vaccination. In certain embodiments, the term “therapy” refers to a recombinant NDV described herein. In other embodiments, the term “therapy” refers to an agent that is not a recombinant NDV described herein.

(74) As used herein, the term “wild-type” in the context of nucleotide and amino acid sequences of viruses refers to the nucleotide and amino acid sequences of viral strains found in nature. In particular, the sequences described as wild-type herein are sequences that have been reported in public databases as sequences from natural viral isolates.

Description

4. BRIEF DESCRIPTION OF THE FIGURES

(1) FIGS. 1A-1N. Inflammation following NDV-F vaccination and boost. Cohorts of cotton rats were sacrificed 28 days after the priming intranasal immunization, or 1 month after a second, boosting dose given on day 28. Mock-immunized animals were given allantoic fluid alone, or the NDV vaccine vector. Representative sections from the nasal cavity and lung after a single priming dose (FIGS. 1A-1D, and FIGS. 1I-1K), or after both a priming and a boosting dose (FIGS. 1E-1H and FIGS. 1L-1N), are pictured. A lymphohistiocytic submucosal infiltrate is present in the nasal cavity (FIGS. 1C and 1D), and surrounding bronchioles (FIG. 1K), after priming with NDV-F, with increased cellularity 1 month following the boosting dose (FIGS. 1G, 3H, and 3N). A minimal peribronchiolar infiltrate is detected only after priming and boosting with the NDV vector. Rare eosinophils are indicated (*). Images were captured at 100× or 400× (insets) magnification.

(2) FIGS. 2A-2F. Histology of the nasal cavity and lung following primary RSV challenge. Nasal cavity (FIGS. 2A-2C) and lung (FIGS. 2D-2F) were collected 4 days after primary RSV challenge from animals that had been previously mock-vaccinated, or given a priming and boosting dose of NDV-F. Respiratory mucosa in the noses of infected animals was disrupted by lymphoplasmacytic infiltrates (FIGS. 2A and 2B), whereas, the nasal cavities of NDV-F vaccinated animals (FIG. 2C) were similar to those observed following vaccination alone (FIG. 1). RSV infection of the lung induced mild peribronchiolar infiltrates of lymphocytes, plasma cells and rare eosinophils and mucus production, with plugging of small airways (FIGS. 2D and 2E). RSV challenge of NDV-F vaccinated animals resulted in lymphoplasmacytic peribronchial infiltrates (FIG. 2F), but without obstruction of small airways by mucus and necrotic cells. Images were taken at 100× (FIGS. 2D-2E), 200× (FIGS. 2A-2C) or 400× (insets) magnification.

(3) FIGS. 3A-3G. Viral burden following primary RSV challenge. Nasal cavity and lung were collected 4 days after primary RSV challenge from controls and animals that had been previously vaccinated. RSV immunohistochemistry revealed marked to complete reduction in antigen-positive cells in nasal cavity (FIGS. 3A-3C) and lung (FIGS. 3D-3F) collected from NDV-F immunized animals compared to controls. Virus was detected by plaque assay (FIG. 3G) only in samples collected from animals mock-immunized with allantoic fluid or vector alone. Images were taken at 200× (FIGS. 3A-3C), 100× (FIGS. 3D-3F) or 400× (insets) magnification.

(4) FIGS. 4A-4E. Humoral immune responses to vaccination and RSV challenge. Serum, BAL fluid and NW fluid were collected at multiple time points after vaccination and RSV challenge and F-specific IgG levels were determined by ELISA (FIGS. 4A-4C). Virus neutralization assay (FIG.

4D) revealed that F-specific antibodies induced by NDV-F vaccination were neutralizing. Cervical lymph nodes (CLNs) were collected 4 days and 5 months after RSV challenge of vaccinated animals and F-specific IgG secreting cells (ASCs) were enumerated by ELISpot (FIG. 4E).

(5) FIGS. 5A-5B. BAL fluid cellularity following vaccination and RSV challenge. BAL fluid was collected at multiple time points after vaccination and RSV challenge. Cellularity (FIG. 5A) of BAL fluid collected from animals in all cohorts increased slightly, but not significantly, in response to primary RSV challenge then decreased gradually over time. Cellularity increased in BAL fluid collected from mock and NDV, but not NDV-F, vaccinated animals in response to secondary RSV challenge. The percentage of eosinophils (FIG. 5B) in BAL fluid collected from mock and NDV, but not NDV-F vaccinated animals, increased in response to primary and secondary RSV challenge. BAL fluid cellularity and eosinophil percentage were not affected by vaccination alone. * $p < 0.05$

(6) FIG. 6. Immunization of RSV-immune animals. Cotton rats were infected with RSV on day 0. Two months post infection, 20/40 animals were primed with NDV-F, and boosted with a second dose 28 days later. At 4 months and 9 months post infection serum samples were taken from cohorts of immunized and control animals (N=5). At the 9 month time point, all animals were given a second challenge of RSV, and sacrificed day 2 and day 4 post re-infection.

(7) FIGS. 7A-7B. Serum antibody titers following post-infection immunization. Serum was harvested from cotton rats 4 or 9 months following primary RSV infection. (FIG. 7A) Levels of neutralizing antibody were determined by incubation of a 1/800 dilution of each serum with rgRSV, which was then used to inoculate A549 cells. Percentage of GFP-expressing cells was used to calculate % neutralization. * $p = 0.0027$; ** $p < 0.0001$. (FIG. 7B) Levels of RSV F specific antibody in each sample were determined by ELISA. * $p = 0.0114$; ** $p = 0.0004$. Levels of neutralizing and RSV F specific antibody were significantly higher in the NDV-F vaccinated animals at 4 and 9 months post primary infection as determined by the unpaired t test.

(8) FIG. 8. Cloning of the chimeric F protein with the ectodomain of the human RSV F and the transmembrane and cytoplasmic domains of the NDV F (not to scale). After rescue all the recombinant NDVs were amplified by inoculation in 8-10 old embryonated chicken eggs and characterized by RT-PCR and immunofluorescence (FIG. 13).

(9) FIG. 9. A schematic representation of the different rNDV-F, rNDV-F.sub.opt, rNDV-F.sub.chim are shown.

(10) FIG. 10. Vero cells were infected with each construct, or the LaSota NDV virus vector for 24 hours, then stained with the Synagis® antibody to detect RSV F expression by flow cytometry. Mean Fluorescence Intensity (MFI) of the positive gated cells is shown.

(11) FIG. 11. The number of plaque forming units of RSV detected in lung homogenates obtained from cotton rats mock treated or given priming and boosting doses of NDV, rNDV-F, rNDV-F.sub.opt, or rNDV-F.sub.chim following RSV challenge.

(12) FIGS. 12A-12B. FIG. 12A shows the relationship of serum sampling to immunization and challenge. FIG. 12B shows antibody titers from individual cotton rats measured with an RSV F protein specific ELISA.

(13) FIG. 13. Detection of the RSV F protein and NDV HN protein expressed by the different recombinant NDVs by double immunofluorescent microscopy.

(14) FIG. 14. Detection of virus in cells treated with different concentrations of Synagis® followed by infection with GFP expressing viruses (RSV-GFP and NDV-GFP). Representative microscopic fields were photographed to prepare the figure.

(15) FIG. 15. Detection of NDV in cells treated with different concentrations of Synagis® followed by infection with NDV, rNDV-F, rNDV-Fopt, or rNDV-Fchim. Representative microscopic fields were photographed to prepare the figure.

(16) FIGS. 16A-16C. An alignment comparing the wild-type nucleic acid sequence encoding human RSV F protein (SEQ ID NO:1) to the codon optimized nucleic acid sequence encoding human RSV F protein (SEQ ID NO:2).

(17) FIGS. 17A-17C. An alignment comparing the wild-type nucleic acid sequence encoding bovine RSV F protein (SEQ ID NO:9) to the codon optimized nucleic acid sequence encoding bovine RSV F protein (SEQ ID NO:11).

(18) FIG. 18. Detection of NDV in cells treated with different concentrations of Synagis® followed by infection with rNDV-bovine RSV F.sub.opt or rNDV-bovine RSVF.sub.chim. Representative microscopic fields were photographed to prepare the figure.

(19) FIGS. 19A-19C. An alignment comparing the wild-type nucleic acid sequence encoding human metapneumovirus F protein (SEQ ID NO:16) to the codon optimized nucleic acid sequence encoding human metapneumovirus F protein (SEQ ID NO:18).

5. DETAILED DESCRIPTION

(20) In one aspect, described herein are recombinant Newcastle disease viruses (“NDVs”) comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a respiratory syncytial virus (“RSV”) F protein or human metapneumovirus (“hMPV”) F protein. In another aspect, described herein are recombinant NDVs comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises (i) an RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains; or (ii) an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In another aspect, described herein are compositions (e.g., immunogenic compositions) comprising a recombinant NDV described herein. In another aspect, the recombinant NDVs and compositions thereof are useful for inducing an immune response to RSV F protein or hMPV F protein, immunizing against RSV or hMPV, or the prevention of RSV disease or hMPV disease.

(21) 5.1 Recombinant Newcastle Disease Virus

(22) 5.1.1 NDV

(23) Any NDV type or strain may be serve as the “backbone” that is engineered to encode a transgene described herein, including, but not limited to, naturally-occurring strains, variants or mutants, mutagenized viruses, reassortants and/or genetically engineered viruses. In a specific embodiment, the nucleotide sequence is incorporated into the genome of a lentogenic NDV. In another specific embodiment, the nucleotide sequence is incorporated in the genome of NDV strain LaSota. Other examples of NDV strains into which the nucleotide sequence may be incorporated are the NDV Fuller, the NDV Ulster strain or the NDV Hitchner B1 strain. In some embodiments, a lentogenic strain other than NDV Hitchner B1 strain is used as the backbone into which a nucleotide sequence may be incorporated. The nucleotide sequence may be incorporated into the NDV genome between two transcription units (e.g., between the M and P transcription units or between the HN and L transcription units).

(24) In a specific embodiment, the NDV that is engineered to encode a transgene described herein is a naturally-occurring strain. In certain embodiments, the NDV that is engineered to encode a transgene described herein is a lytic strain. In other embodiments, the NDV that is engineered to encode a transgene described herein is a non-lytic strain. In certain embodiments, the NDV that is engineered to encode a transgene described herein is lentogenic strain. In some embodiments, the NDV that is engineered to encode a transgene described herein is a mesogenic strain. In other embodiments, the NDV that is engineered to encode a transgene described herein is a velogenic strain. Specific examples of NDV strains include, but are not limited to, the 73-T strain, NDV HUIJ strain, Ulster strain (see, e.g., GenBank No. U25837), Fuller strain, MTH-68 strain, Italien strain (see, e.g., GenBank No. EU293914), Hickman strain (see, e.g., Genbank No. AF309418), PV701 strain, Hitchner B1 strain (see, e.g., GenBank No. AF309418 or NC_002617), La Sota strain (see, e.g., GenBank Nos. AY845400, AF07761.1 and JF950510.1 and GI No. 56799463), YG97 strain (see, e.g., GenBank Nos. AY351959 or AY390310), MET95 strain (see, e.g., GenBank No. AY143159), Roakin strain (see, e.g., GenBank No. AF124443), and F48E9 strain (see, e.g., GenBank Nos. AF163440 and U25837). In a specific embodiment, the NDV that is engineered to

encode a transgene described herein is the Hitchner B1 strain. In another embodiment, the NDV that is engineered to encode a transgene described herein is a B1 strain as identified by GenBank No. AF309418 or NC_002617. In a specific embodiment, the nucleotide sequence of the Hitchner B1 genome comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO:48. In another specific embodiment, the NDV that is engineered to encode a transgene described herein is the La Sota strain. In another embodiment, the NDV that is engineered to encode a transgene described herein is a LaSota strain as identified by GenBank Nos. AY845400, AF07761.1 or JF950510.1. In a specific embodiment, the nucleotide sequence of the La Sota genome comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO:47. One skilled in the art will understand that the NDV genomic RNA sequence is an RNA sequence corresponding to the negative sense of a cDNA sequence encoding the NDV genome. Thus, any program that converts a nucleotide sequence to its reverse complement sequence may be utilized to convert a cDNA sequence encoding an NDV genome into the genomic RNA sequence (see, e.g., Bioinformatics.org and DNASTar). Accordingly, the nucleotide sequences provided in Tables 1-6, infra, may be readily converted to the negative-sense RNA sequence of the NDV genome by one of skill in the art.

(25) In specific embodiments, the NDV that is engineered to encode a transgene described herein is not pathogenic in birds as assessed by a technique known to one of skill. In certain specific embodiments, the NDV that is engineered to encode a transgene described herein is not pathogenic as assessed by intracranial injection of 1-day-old chicks with the virus, and disease development and death as scored for 8 days. In some embodiments, the NDV that is engineered to encode a transgene described herein has an intracranial pathogenicity index of less than 0.7, less than 0.6, less than 0.5, less than 0.4, less than 0.3, less than 0.2 or less than 0.1. In certain embodiments, the NDV that is engineered to encode a transgene described herein has an intracranial pathogenicity index of zero. See, e.g., OIE Terrestrial Manual 2012, Chapter 2.3.14, entitled "Newcastle Disease (Infection With Newcastle Disease Virus) for a description of this assay, which is incorporated herein by reference in its entirety.

(26) In certain embodiments, the NDV that is engineered to encode a transgene described herein is a mesogenic strain that has been genetically engineered so as not be a considered pathogenic in birds as assessed by techniques known to one skilled in the art. In certain embodiments, the NDV that is engineered to encode a transgene described herein is a velogenic strain that has been genetically engineered so as not be a considered pathogenic in birds as assessed by techniques known to one skilled in the art.

(27) In preferred embodiments, the NDV that is engineered to encode a transgene described herein is non-pathogenic in humans or bovine. In preferred embodiments, the NDV that is engineered to encode a transgene described herein is non-pathogenic in humans, bovines and avians. In certain embodiments, the NDV that is engineered to encode a transgene described herein is attenuated such that the NDV remains, at least partially, infectious and can replicate in vivo, but only generate low titers resulting in subclinical levels of infection that are non-pathogenic (see, e.g., Khattar et al., 2009, J. Virol. 83:7779-7782). Such attenuated NDVs may be especially suited for embodiments wherein the virus is administered to a subject in order to act as an immunogen, e.g., a live vaccine. The viruses may be attenuated by any method known in the art. In a specific embodiment, the NDV genome comprises sequences necessary for infection and replication of the attenuated virus such that progeny is produced and the infection level is subclinical.

(28) 5.1.2 RSV F Protein/Chimeric F Protein with the RSV F Protein Ectodomain

(29) In a specific embodiment, a transgene encoding a human RSV F protein is incorporated into the genome of any NDV type or strain. (e.g., NDV LaSota strain) See, e.g., Section 5.1.1., supra, for types and strains of NDV that may be used. The transgene encoding any RSV F protein may be inserted into any NDV type or strain (e.g., NDV LaSota strain). The RSV F protein may be an RSV F protein of a human or bovine strain of RSV. In a specific embodiment, a transgene encoding a

human RSV F protein is incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). See, e.g., Section 5.8 3.1 for exemplary sequences for human RSV F protein and exemplary nucleic acid sequences encoding human RSV F protein. One of skill in the art would be able to use such sequence information to produce a transgene for incorporation into the genome of any NDV type or strain. For example, a transgene encoding the human RSV F protein comprising the amino acid sequence set forth in SEQ ID NO:6, 49, 50 or 58 may be incorporated into the genome of any NDV type or strain. In a specific embodiment, such a transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO:1, 25, 27 or 29. However, given the degeneracy of the nucleic acid code, there are a number of different nucleic acid sequences that may encode the same human RSV F protein. In a specific embodiment, a transgene encoding a human RSV F protein is codon optimized. See, e.g., Section 5.1.6, *infra*, for a discussion regarding codon optimization. For example, the human RSV F protein may be encoded by a codon optimized nucleic acid sequence, such as set forth in SEQ ID NO: 2, 26, 28 or 30. In some embodiments, the transgene encoding a human RSV F protein comprises the amino acid sequence encoded by the nucleic acid sequence comprising the sequence set forth in SEQ ID NO:1, 2, 25, 26, 27, 28, 29, or 30. The transgene encoding a human RSV F protein may be incorporated between any two NDV transcription units (e.g., between the NDV P and M transcription units, or between the HN and L transcription units).

(30) In a specific embodiment, a transgene encoding a bovine RSV F protein is incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). See, e.g., Section 5.8 3.1 for exemplary sequences for bovine RSV F protein and exemplary nucleic acid sequences encoding bovine RSV F protein. One of skill in the art would be able to use such sequence information to produce a transgene for incorporation into the genome of any NDV type or strain. For example, a transgene encoding the bovine RSV F protein comprising the amino acid sequence set forth in SEQ ID NO:10 may be incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). In a specific embodiment, a transgene encoding the bovine RSV F protein comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO:9, 40 or 42. However, given the degeneracy of the nucleic acid code, there are a number of different nucleic acid sequences that may encode the same bovine RSV F protein. In a specific embodiment, a transgene encoding a bovine RSV F protein is codon optimized. For example, the bovine RSV F protein may be encoded by the codon optimized nucleic acid sequence set forth in SEQ ID NO: 11, 41 or 43. In some embodiments, the transgene encoding a bovine RSV F protein comprises the amino acid sequence encoded by the nucleic acid sequence comprising the sequence set forth in SEQ ID NO:9, 11, 40, 41, 42, or 43. The transgene encoding a bovine RSV F protein may be incorporated between any two NDV transcription units (e.g., between NDV P and M genes).

(31) In another embodiment, described herein are transgenes encoding a chimeric F protein, wherein the chimeric F protein comprises an RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In other words, the NDV F protein transmembrane and cytoplasmic domains replace the RSV F protein transmembrane and cytoplasmic domains so that the chimeric F protein does not include the RSV F protein transmembrane and cytoplasmic domains. The RSV F protein may be an RSV F protein of a human or bovine strain of RSV. The ectodomain, transmembrane and cytoplasmic domains of the RSV F protein and NDV F protein may be determined using techniques known to one of skill in the art. For example, published information, GenBank or websites such as VIPR virus pathogen website, DTU Bioinformatics domain website or programs available to determine the transmembrane domain may be used to determine the ectodomain, transmembrane and cytoplasmic domains of the RSV F protein and NDV F protein. See, e.g., the tables *infra* with the transmembrane and cytoplasmic domains indicated. In specific embodiment, described herein is a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein

transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 51. In another embodiment, described herein is a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises the an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 31. In a preferred embodiment, a transgene comprises a codon optimized version of a nucleic acid sequence encoding the RSV F protein ectodomain (e.g., a human or bovine RSV F protein ectodomain). In a specific embodiment, a transgene encoding a chimeric F protein is incorporated into the genome of any NDV type or strain (e.g., NDV LaSota strain). See, e.g., Section 5.1.1, supra, for types and strains of NDV that may be used. The transgene encoding a chimeric F protein may be incorporated between any two NDV transcription units (e.g., between the NDV P and M transcription units, or between the HN and L transcription units).

(32) In a specific embodiment, described herein is a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence. Specific examples of nucleic acid sequences encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence, include SEQ ID NO:4, 44, 45, or 46. In a preferred embodiment, described herein is a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:4. In a specific embodiment, a transgene encoding a chimeric F protein is incorporated into the genome of any NDV type or strain. See, e.g., Section 5.1.1, supra, for types and strains of NDV that may be used. The transgene encoding a chimeric F protein may be incorporated between any two NDV transcription units (e.g., between the NDV P and M transcription units, or between the HN and L transcription units).

(33) In a specific embodiment, described herein is a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence. Specific examples of nucleic acid sequences encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence, include SEQ ID NO:14, 38, or 39. In a preferred embodiment, described herein is a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:14 or 31. In a specific embodiment, a transgene encoding a chimeric F protein is incorporated into the genome of any NDV type or strain. See, e.g., Section 5.1.1, supra, for types and strains of NDV that may be used. The transgene encoding a chimeric F protein may be incorporated between any two NDV transcription units (e.g., between the NDV P and M transcription units, or between the HN and L transcription units).

(34) In a specific embodiments, a transgene encoding a human RSV F protein or a chimeric F protein is encoded by a nucleic acid sequence comprising a nucleic acid sequence set forth in SEQ ID NO:1, 2, 4, 25, 26, 27, 28, 29, 30, 44, 45, 46, or 51, or an RNA sequence corresponding to the negative sense thereof. In specific embodiments, a transgene encoding a bovine RSV F protein or chimeric F protein is encoded by a nucleic acid sequence comprising a nucleic acid sequence set

forth in SEQ ID NO: 9, 11, 14, 31, 38, 39, 40, 41, 42, or 43, or an RNA sequence corresponding to the negative sense thereof. See, e.g., Tables 1 and 3 for nucleic acid and amino acid sequences of RSV F. Also, see, e.g., Section 5.8 for exemplary human or bovine RSV F proteins, or chimeric F proteins.

(35) In a specific embodiment, a transgene encoding an RSV F protein or a chimeric F protein is as described in Section 6, *infra*.

(36) In certain embodiments, a transgene encoding an RSV F protein or a chimeric F protein comprises NDV regulatory signals (e.g., gene end, intergenic, and gene start sequences) and Kozak sequences. In some embodiments, a transgene encoding an RSV F protein or a chimeric F protein comprises NDV regulatory signals (e.g., gene end, intergenic, and gene start sequences), Kozak sequences and restriction sites to facilitate cloning. See, e.g., FIG. 8. In certain embodiments, a transgene encoding an RSV F protein or a chimeric F protein comprises NDV regulatory signals (gene end, intergenic and gene start sequences), Kozak sequences, restriction sites to facilitate cloning, and additional nucleotides in the non-coding region to ensure compliance with the rule of six. In a preferred embodiment, the transgene complies with the rule of six.

(37) 5.1.3 Recombinant NDV Encoding an RSV F Protein or a Chimeric F Protein with an RSV Ectodomain

(38) In one aspect, presented herein are recombinant Newcastle disease virus (“NDV”) comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a respiratory syncytial virus (“RSV”) F protein. The RSV F protein may be an RSV F protein of a human or bovine strain of RSV. See, e.g., Section 5.1.2 and 6 for transgenes encoding a human or bovine RSV F protein which the packaged genome may comprise. In a specific embodiment, the transgene encodes the human RSV F protein comprising the amino sequence set forth in SEQ ID NO: 6, 49, 50 or 58, or the bovine RSV F protein set forth in SEQ ID NO: 10. Due to the degeneracy of the nucleic acid code, multiple different nucleic acid sequences may encode for the same human RSV F protein or the same bovine RSV F protein. In one embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a human RSV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:1, 25, 27 or 29. In another embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a bovine RSV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:9, 40 or 42. In a preferred embodiment, a transgene comprises a codon optimized version of a nucleic acid sequence encoding an RSV F protein (e.g., a human or bovine RSV F protein). In a specific embodiment, the RSV F protein is expressed by cells infected with the recombinant NDV. In some embodiments, RSV F protein is detected in the virion of recombinant NDV.

(39) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene comprising a codon optimized nucleic acid sequence encoding a human RSV F protein. Specific examples of codon optimized nucleic acid sequences encoding a human RSV F protein include those in SEQ ID Nos:2, 26, 28, or 30. In a preferred embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a human RSV F protein, wherein the transgene comprises the an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:2. In a specific embodiment, the human RSV F protein is expressed by cells infected with the recombinant NDV. In some embodiments, RSV F protein is detected in the virion of recombinant NDV.

(40) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene comprising a codon optimized nucleic acid sequence encoding a bovine RSV F protein. Specific examples of codon optimized nucleic acid sequences encoding a bovine RSV F protein include those in SEQ ID Nos: 11, 41, or

43. In a preferred embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a bovine RSV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:11. In a specific embodiment, the bovine RSV F protein is expressed by cells infected with the recombinant NDV. In some embodiments, RSV F protein is detected in the virion of recombinant NDV.

(41) In another embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In other words, the NDV F protein transmembrane and cytoplasmic domains replace the RSV F protein transmembrane and cytoplasmic domains so that the chimeric F protein does not include the RSV F protein transmembrane and cytoplasmic domains. In a specific embodiment, the NDV F protein transmembrane and cytoplasmic domains are from the same strain of NDV as the NDV backbone. For example, if the NDV backbone is NDV LaSota, then the transmembrane and cytoplasmic domains of the chimeric F protein are NDV LaSota transmembrane and cytoplasmic domains. The RSV F protein may be an RSV F protein of a human or bovine strain of RSV. See, e.g., Section 5.1.2 and 6 for transgenes encoding a chimeric F protein which the packaged genome may comprise. In another embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:51. In another embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:31. In a preferred embodiment, a transgene comprises a codon optimized version of a nucleic acid sequence encoding the RSV F protein ectodomain (e.g., a human or bovine RSV F protein ectodomain). In a specific embodiment, the RSV F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the RSV F protein is expressed by cells infected with the recombinant NDV and the chimeric F protein is incorporated into the NDV virion.

(42) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence. Specific examples of nucleic acid sequences encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence, include SEQ ID NO:4, 44, 45, and 46. In a preferred embodiment, described herein is a recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:4. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV and the chimeric F protein is incorporated into the virion of the NDV. In another specific embodiment, the RSV F protein is expressed by cells infected with the recombinant NDV and the chimeric F protein is incorporated into the NDV virion.

(43) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence. Specific examples of nucleic acid sequences encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the RSV F protein is encoded by a codon optimized nucleic acid sequence, include SEQ ID NO:14, 38, or 39. In a preferred embodiment, described herein is a recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:14. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. In another specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV and the chimeric F protein is incorporated into the virion of the NDV.

(44) In a specific embodiment, a recombinant NDV comprises a packaged genome, wherein the packaged genome comprises a transgene encoding a RSV F protein or a chimeric F protein, wherein the RSV F protein or chimeric F protein is encoded by a nucleic acid sequence comprising a nucleic acid sequence set forth in SEQ ID NO:1, 2, 4, 25, 26, 27, 28, 29, 30, 44, 45, 46, or 51, or an RNA sequence corresponding to the negative sense thereof. In specific embodiments, recombinant NDV comprises a packaged genome, wherein the packaged genome comprises a transgene encoding a RSV F protein or a chimeric F protein, wherein the RSV F protein or chimeric F protein is encoded by a nucleic acid sequence comprising a nucleic acid sequence set forth in SEQ ID NO: 9, 11, 14, 31, 38, 39, 40, 41, 42, or 43, or an RNA sequence corresponding to the negative sense thereof.

(45) In a specific embodiment, a recombinant NDV is as described in Section 6, *infra*.

(46) In a specific embodiment, a recombinant NDV described herein comprises a packaged genome, wherein the packaged genome corresponds to a negative sense RNA transcribed from a cDNA sequence comprising (or consisting of) the sequence set forth in SEQ ID NO: 3. In certain embodiment, the genome of the recombinant NDV does not comprise a heterologous sequence encoding a heterologous protein other than the human RSV F protein. In some embodiments, the genome of the recombinant NDV does not comprise a transgene other than a transgene encoding a human RSV F protein described herein. In certain embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV and a transgene encoding a human RSV F protein. In other words, the recombinant NDV encodes for both NDV F protein and the human RSV F protein. In certain embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV, a transgene encoding a human RSV F protein, and a transgene encoding an hMPV F protein or a chimeric F protein described herein, e.g., in Section 5.1.4, but does not include other transgenes, wherein the chimeric F protein comprises the ectodomain of hMPV F protein and the transmembrane and cytoplasmic domains of NDV F protein. In some embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV and a transgene encoding a human RSV F protein but does not include any other transgenes.

(47) In a specific embodiment, a recombinant NDV described herein comprises a packaged genome, wherein the packaged genome corresponds to a negative sense RNA of a cDNA sequence comprising (or consisting of) the sequence set forth in SEQ ID NO: 12 or 13. In certain embodiment, the genome of the recombinant NDV does not comprise a heterologous sequence encoding a heterologous protein other than the bovine RSV F protein. In some embodiments, the

genome of the recombinant NDV does not comprise a transgene other than a transgene encoding a bovine RSV F protein described herein. In certain embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV and a transgene encoding a bovine RSV F protein. In other words, the recombinant NDV encodes for both NDV F protein and the bovine RSV F protein. In certain embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV, a transgene encoding a bovine RSV F protein, and a transgene encoding an hMPV F protein or a chimeric F protein described herein, e.g., in Section 5.1.4, but does not include other transgenes, wherein the chimeric F protein comprises the ectodomain of hMPV F protein and the transmembrane and cytoplasmic domains of NDV F protein. In some embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV and a transgene encoding a bovine RSV F protein but does not include any other transgenes.

(48) In a specific embodiment, a recombinant NDV described herein comprises a packaged genome, wherein the packaged genome corresponds to a negative sense RNA sequence of a cDNA sequence comprising (or consisting of) the sequence set forth in SEQ ID NO: 5, 37 or 59. In some embodiments, the packaged genome of NDV encodes a chimeric F protein, wherein the chimeric F protein comprises the human or bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In certain embodiment, the genome of the recombinant NDV does not comprise a heterologous sequence encoding a heterologous protein other than the chimeric F protein. In some embodiments, the genome of the recombinant NDV does not comprise a transgene other than a transgene encoding a chimeric F protein described herein. In certain embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV and a transgene encoding a chimeric F protein. In other words, the recombinant NDV encodes for both NDV F protein and the chimeric F protein. In some embodiment, the genome of a recombinant NDV described herein comprises a transgene encoding a chimeric F protein and the genes found in NDV except for the gene encoding NDV F protein. In other words, the NDV encodes the chimeric F protein but not the NDV F protein. In certain embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV, a transgene encoding first chimeric F protein, and a transgene encoding an hMPV F protein or a second chimeric F protein described herein, e.g., in Section 5.1.4, but does not include other transgenes, wherein the first chimeric F protein comprises the ectodomain of human RSV F protein or bovine RSV F protein and the transmembrane and cytoplasmic domains of NDV F protein, and wherein the second chimeric F protein comprises the ectodomain of hMPV F protein and the transmembrane and cytoplasmic domains of NDV F protein. In some embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV and a transgene encoding a chimeric F protein but does not include other transgenes.

(49) 5.1.4 Human Metapneumovirus (hMPV) F Protein

(50) In a specific embodiment, a transgene encoding a hMPV F protein is incorporated into the genome of any NDV type or strain. See, e.g., Section 5.1.1, supra, for types and strains of NDV that may be used. The transgene encoding any hMPV F protein may inserted into any NDV type or strain (e.g., NDV LaSota strain). The hMPV F protein may be an hMPV F protein of any strain of hMPV. See, e.g., Section 5.8 for exemplary sequences for hMPV F protein and exemplary nucleic acid sequences encoding hMPV F protein. One of skill in the art would be able to use such sequence information to produce a transgene for incorporation into the genome of any NDV type or strain. For example, a transgene encoding the hMPV F protein comprising the amino acid sequence set forth in SEQ ID NO:17 may be incorporated into the genome of any NDV type or strain. In a specific embodiment, such a transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO:16, 52, 54, or 56. However, given the

degeneracy of the nucleic acid code, there are a number of different nucleic acid sequences that may encode the same hMPV F protein. In a specific embodiment, a transgene encoding an hMPV F protein is codon optimized. See, e.g., Section 5.1.6, *infra*, for a discussion regarding codon optimization. For example, the hMPV F protein is encoded by the codon optimized nucleic acid sequence, such as set forth in SEQ ID NO: 18, 53, 55, or 57. In some embodiments, the transgene encoding an hMPV F protein comprises the amino acid sequence encoded by the nucleic acid sequence comprising the sequence set forth in SEQ ID NO:16, 18, 52, 53, 54, 55, 56 or 57. In certain embodiments, the hMPV F protein is encoded by a transgene comprising the an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO:16, 18, 52, 53, 54, 55, 56 or 57. The transgene encoding a hMPV F protein may be incorporated between any two NDV transcription units (e.g., between the NDV P and M transcription units, or between the HN and L transcription units).

(51) In another embodiment, described herein are transgenes encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In other words, the NDV F protein transmembrane and cytoplasmic domains replace the hMPV F protein transmembrane and cytoplasmic domains so that the chimeric F protein does not include the hMPV F protein transmembrane and cytoplasmic domains. The hMPV F protein may be an hMPV F protein of any strain of hMPV. The ectodomain, transmembrane and cytoplasmic domains of the hMPV F protein and NDV F protein may be determined using techniques known to one of skill in the art. For example, published information, GenBank or websites such as VIPR virus pathogen website, DTU Bioinformatics domain website, or programs available to determine the transmembrane domain may be used to determine the ectodomain, transmembrane and cytoplasmic domains of the hMPV F protein and NDV F protein. See, e.g., the tables below indicating the transmembrane and cytoplasmic domains. In specific embodiment, described herein is a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:32. In a preferred embodiment, a transgene comprises a codon optimized version of a nucleic acid sequence encoding the hMPV F protein ectodomain. In specific embodiment, described herein is a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:19, 34, 35, or 36. In a specific embodiment, a transgene encoding a chimeric F protein is incorporated into the genome of any NDV type or strain. See, e.g., Section 5.1.1, *supra*, for types and strains of NDV that may be used. The transgene encoding a chimeric F protein may be incorporated between any two NDV transcription units (e.g., between the NDV P and M transcription units, or between the HN and L transcription units).

(52) In a specific embodiment, described herein is a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the hMPV F protein is encoded by a codon optimized nucleic acid sequence. Specific examples of nucleic acid sequences encoding a chimeric F protein, wherein the chimeric F protein comprises a human hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the ectodomain of the hMPV F protein is encoded by a codon optimized nucleic acid sequence, include SEQ ID NO:19, 34, 35, and 36. In a preferred embodiment, described herein is a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:19. In a specific embodiment, a transgene encoding a chimeric F protein is

incorporated into the genome of any NDV type or strain. See, e.g., Section 5.1.1, supra, for types and strains of NDV that may be used. The transgene encoding a chimeric F protein may be incorporated between any two NDV transcription units (e.g., between the NDV P and M transcription units, or between the HN and L transcription units).

(53) In specific embodiments, a transgene encoding an hMPV F protein or a chimeric F protein is encoded by a nucleic acid sequence comprising a nucleic acid sequence set forth in SEQ ID NO:16, 18, 19, 32, 34, 35, 36, 52, 53, 54, 55, 56, or 57, or an RNA sequence corresponding to the negative sense thereof. See, e.g., Section 5.8 for exemplary hMPV F protein or chimeric F proteins.

(54) In a specific embodiment, a transgene encoding an hMPV F protein or a chimeric F protein is as described in Section 6, infra.

(55) In certain embodiments, a transgene encoding an hMPV F protein or a chimeric F protein comprises NDV regulatory signals (e.g., gene end, intergenic, and gene start sequences) and Kozak sequences. In some embodiments, a transgene encoding an hMPV F protein or a chimeric F protein comprises NDV regulatory signals (e.g., gene end, intergenic, and gene start sequences), Kozak sequences and restriction sites to facilitate cloning. In certain embodiments, a transgene encoding an hMPV F protein or a chimeric F protein comprises NDV regulatory signals (gene end, intergenic and gene start sequences), Kozak sequences, restriction sites to facilitate cloning, and additional nucleotides in the non-coding region to ensure compliance with the rule of six. In a preferred embodiment, the transgene complies with the rule of six.

(56) 5.1.5 Recombinant NDV Encoding a hMPV F Protein or a Chimeric F Protein with an hMPV Ectodomain

(57) In one aspect, presented herein are recombinant Newcastle disease virus (“NDV”) comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a human metapneumovirus (“hMPV”) F protein. See, e.g., Section 5.1.4 and 6 for transgenes encoding an hMPV F protein which the packaged genome may comprise. The hMPV F protein may be an hMPV F protein of any strain of hMPV. In a specific embodiment, the transgene encodes the hMPV F protein comprising the sequence set forth in SEQ ID NO: 17. Due to the degeneracy of the nucleic acid code, multiple different nucleic acid sequences may encode for the same hMPV F protein. Specific examples of nucleic acid sequences encoding a hMPV F protein include those set forth in SEQ ID NOs:16, 52, 54 and 56. In one embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a hMPV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:16. In a preferred embodiment, a transgene comprises a codon optimized version of a nucleic acid sequence encoding an hMPV F protein. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV.

(58) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene comprising a codon optimized nucleic acid sequence encoding a hMPV F protein. Specific examples of codon optimized nucleic acid sequences encoding a hMPV F protein include those set forth in SEQ ID Nos:18, 53, 55 and 57. In a specific embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a hMPV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:18. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV.

(59) In another embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In other words, the NDV F protein transmembrane and cytoplasmic domains replace the hMPV F protein transmembrane and cytoplasmic domains so that the chimeric F protein does not include the hMPV F protein transmembrane and cytoplasmic

domains. The chimeric hMPV F protein may comprise the amino acid sequence of SEQ ID NO:15. In a specific embodiment, the NDV F protein transmembrane and cytoplasmic domains are from the strain of NDV as the NDV backbone. The hMPV F protein may be an hMPV F protein of any strain of hMPV. See, e.g., Section 5.1.4 and 6 for transgenes encoding a chimeric F protein which the packaged genome may comprise. In another embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:32. In a preferred embodiment, a transgene comprises a codon optimized version of a nucleic acid sequence encoding the hMPV F protein ectodomain. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV. In some embodiments, hMPV F protein is detected in the NDV virion.

(60) In a specific embodiment, described herein are recombinant NDV comprising a packaged genome, wherein the packaged genome comprises a transgene comprising a codon optimized nucleic acid sequence encoding a hMPV F protein. Specific examples of codon optimized nucleic acid sequences encoding a hMPV F protein include those set forth in SEQ ID NOs: 19, 34, 35 and 36. In a specific embodiment, described herein is a recombinant NDV comprising a packaged genome comprising a transgene encoding a hMPV F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:19. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV. In some embodiments, hMPV F protein is detected in the NDV virion.

(61) In a specific embodiment, a recombinant NDV described herein comprises a packaged genome, wherein the packaged genome corresponds to a negative sense RNA of the cDNA sequence comprising (or consisting of) the sequence set forth in SEQ ID NO: 20 or 21. In certain embodiment, the genome of the recombinant NDV does not comprise a heterologous sequence encoding a heterologous protein other than the hMPV F protein. In some embodiments, the genome of the recombinant NDV does not comprise a transgene other than a transgene encoding a hMPV F protein described herein. In certain embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV and a transgene encoding an hMPV F protein. In other words, the recombinant NDV encodes for both NDV F protein and the hMPV F protein. In certain embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV, a transgene encoding an hMPV F protein, and a transgene encoding a human RSV F protein or a second chimeric F protein described herein, e.g., in Section 5.1.2, but does not include other transgenes, wherein the first chimeric F protein comprises the ectodomain of human RSV F protein and the transmembrane and cytoplasmic domains of NDV F protein. In some embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV and a transgene encoding an hMPV F protein but does not include other transgenes.

(62) In a specific embodiment, a recombinant NDV described herein comprises a packaged genome, wherein the packaged genome corresponds to a negative sense RNA of the cDNA sequence comprising (or consisting of) the sequence set forth in SEQ ID NO: 22. In a specific embodiment, a recombinant NDV is as described in Section 6, *infra*. In certain embodiment, the genome of the recombinant NDV does not comprise a heterologous sequence encoding a heterologous protein other than the chimeric F protein. In some embodiments, the genome of the recombinant NDV does not comprise a transgene other than a transgene encoding a chimeric F protein described herein. In certain embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV and a transgene encoding a chimeric F protein. In other words, the recombinant NDV encodes for both NDV F

protein and the chimeric F protein. In some embodiments, the genome of a recombinant NDV described herein comprises a transgene encoding the chimeric F protein and the genes of NDV except the gene encoding NDV F protein. In other words, the recombinant NDV encodes the chimeric F protein but not the NDV F protein. In certain embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV, a transgene encoding a first chimeric F protein, and a transgene encoding a human RSV F protein or a second chimeric F protein described herein, e.g., in Section 5.1.2, but does not include other transgenes, wherein the first chimeric F protein comprises the ectodomain of hMPV F protein and the transmembrane and cytoplasmic domains of NDV F protein, and wherein the second chimeric F protein comprises the ectodomain of human RSV F protein and the transmembrane and cytoplasmic domains of NDV F protein. In some embodiments, a recombinant NDV described herein comprises a packaged genome, wherein the genome comprises the genes found in NDV and a transgene encoding a chimeric F protein but does not include other transgenes.

(63) 5.1.6 Codon Optimization

(64) Any codon optimization technique known to one of skill in the art may be used to codon optimize a nucleic acid sequence encoding an RSV F protein or a domain thereof (e.g., the ectodomain thereof). Similarly, any codon optimization technique may be used to codon optimize a nucleic acid sequence encoding an hMPV F protein or a domain thereof (e.g., the ectodomain thereof). Methods of codon optimization are known in the art, e.g., the OptimumGene™ (GenScript®) protocol and Genewiz® protocol, which are incorporated by reference herein in its entirety. See also U.S. Pat. No. 8,326,547 for methods for codon optimization, which is incorporated herein by reference in its entirety.

(65) As an exemplary method for codon optimization, each codon in the open frame of the nucleic acid sequence encoding an RSV F protein or a domain thereof (e.g., the ectodomain thereof), or an hMPV F protein or a domain thereof (e.g., the ectodomain thereof) is replaced by the codon most frequently used in mammalian proteins. This may be done using a web-based program (Encorbio) that uses the Codon Usage Database, maintained by the Department of Plant Gene Research in Kazusa, Japan. This nucleic acid sequence optimized for mammalian expression may be inspected for: (1) the presence of stretches of 5xA or more that may act as transcription terminators; (2) the presence of restriction sites that may interfere with subcloning; (3) compliance with the rule of six. Following inspection, (1) stretches of 5xA or more that may act as transcription terminators may be replaced by synonymous mutations; (2) restriction sites that may interfere with subcloning may be replaced by synonymous mutations; (3) NDV regulatory signals (gene end, intergenic and gene start sequences), and Kozak sequences for optimal protein expression may be added; and (4) nucleotides may be added in the non-coding region to ensure compliance with the rule of six. Synonymous mutations are typically nucleotide changes that do not change the amino acid encoded. For example, in the case of a stretch of 6 As (AAAAAA), which sequence encodes Lys-Lys, a synonymous sequence would be AAGAAG, which sequence also encodes Lys-Lys.

(66) In a specific embodiment, codon optimization reduces the AU content of the nucleic acid sequences encoding an RSV F protein (e.g., a human or bovine RSV-F protein) or a domain thereof (e.g., the ectodomain thereof), or a human MPV-F or a domain thereof (e.g., the ectodomain thereof). FIGS. 16A-16C is an alignment comparing the wild-type nucleic acid sequence encoding a wild-type human RSV F protein to the codon optimized nucleic acid sequence encoding human RSV F protein. FIGS. 17A-17C is an alignment comparing the wild-type nucleic acid sequence encoding a bovine RSV F protein to the codon optimized nucleic acid sequence encoding bovine RSV F protein. FIGS. 19A-19C is an alignment comparing the wild-type nucleic acid sequence encoding a wild-type hMPV F protein to the codon optimized nucleic acid sequence encoding hMPV F protein.

(67) In a specific embodiment, the codon optimized nucleic acid sequence encoding human RSV F protein has a 10%, 15%, 20%, 25% or more drop in the AU content relative to the wild-type

nucleic acid sequence encoding human RSV F protein. In another specific embodiment, the codon optimized nucleic acid sequence encoding human RSV F protein has a 10% to 25%, 10% to 30%, 15% to 25% 15% to 30%, or 20% to 30% drop in the AU content relative to the wild-type nucleic acid sequence encoding human RSV F protein. The open reading frame of the wild-type nucleic acid sequence encoding human RSV F protein set forth in SEQ ID NO:1 has a 65.16% A+U, whereas the codon optimized nucleic acid sequence set forth in SEQ ID NO:2 has a 39.43% A+U. In a specific embodiment, the codon optimized nucleic acid sequence encoding bovine RSV F protein has a 10%, 15%, 20%, 25% or more drop in the AU content relative to the wild-type nucleic acid sequence encoding bovine RSV F protein. In another specific embodiment, the codon optimized nucleic acid sequence encoding bovine RSV F protein has a 10% to 25%, 10% to 30%, 15% to 25% 15% to 30%, or 20% to 30% drop in the AU content relative to the wild-type nucleic acid sequence encoding bovine RSV F protein. The open reading frame of the wild-type nucleic acid sequence encoding bovine RSV F protein set forth in SEQ ID NO:9 has a 64.4% A+U, whereas the codon optimized nucleic acid sequence set forth in SEQ ID NO:11 has a 39.66% A+U. In a specific embodiment, the codon optimized nucleic acid sequence encoding hMPV F protein has a 10%, 15%, 20%, 25% or more drop in the AU content relative to the wild-type nucleic acid sequence encoding hMPV F protein. In another specific embodiment, the codon optimized nucleic acid sequence encoding hMPV F protein has a 10% to 25%, 10% to 30%, 15% to 25% 15% to 30%, or 20% to 30% drop in the AU content relative to the wild-type nucleic acid sequence encoding hMPV F protein. The open reading frame of the wild-type nucleic acid sequence encoding hMPV F protein set forth in SEQ ID NO:16 has a 56.86% A+U, whereas the codon optimized nucleic acid sequence set forth in SEQ ID NO:18 has a 37.9% A+U.

(68) 5.2 Construction of NDVs

(69) The recombinant NDVs described herein (see, e.g., Sections 5.1 and 6) can be generated using the reverse genetics technique. The reverse genetics technique involves the preparation of synthetic recombinant viral RNAs that contain the non-coding regions of the negative-strand, viral RNA which are essential for the recognition by viral polymerases and for packaging signals necessary to generate a mature virion. The recombinant RNAs are synthesized from a recombinant DNA template and reconstituted in vitro with purified viral polymerase complex to form recombinant ribonucleoproteins (RNPs) which can be used to transfect cells. A more efficient transfection is achieved if the viral polymerase proteins are present during transcription of the synthetic RNAs either in vitro or in vivo. The synthetic recombinant RNPs can be rescued into infectious virus particles. The foregoing techniques are described in U.S. Pat. No. 5,166,057 issued Nov. 24, 1992; in U.S. Pat. No. 5,854,037 issued Dec. 29, 1998; in U.S. Pat. No. 6,146,642 issued Nov. 14, 2000; in European Patent Publication EP 0702085A1, published Feb. 20, 1996; in U.S. patent application Ser. No. 09/152,845; in International Patent Publications PCT WO97/12032 published Apr. 3, 1997; WO96/34625 published Nov. 7, 1996; in European Patent Publication EP A780475; WO 99/02657 published Jan. 21, 1999; WO 98/53078 published Nov. 26, 1998; WO 98/02530 published Jan. 22, 1998; WO 99/15672 published Apr. 1, 1999; WO 98/13501 published Apr. 2, 1998; WO 97/06270 published Feb. 20, 1997; and EPO 780 475A1 published Jun. 25, 1997, each of which is incorporated by reference herein in its entirety.

(70) The helper-free plasmid technology can also be utilized to engineer a NDV described herein. Briefly, a complete cDNA of a NDV (e.g., the Hitchner B1 strain or LaSota strain) is constructed, inserted into a plasmid vector and engineered to contain a unique restriction site between two transcription units (e.g., the NDV P and M genes; or the NDV HN and L genes). A nucleotide sequence encoding a heterologous amino acid sequence (e.g., a transgene or other sequence described herein such as, e.g., a nucleotide sequence encoding an RSV F protein, a chimeric F protein, hMPV F protein) may be inserted into the viral genome at the unique restriction site. Alternatively, a nucleotide sequence encoding a heterologous amino acid sequence (e.g., a transgene or other sequence described herein such as, e.g., a nucleotide sequence encoding an RSV

F protein, a chimeric F protein, hMPV F protein) may be engineered into a NDV transcription unit so long as the insertion does not affect the ability of the virus to infect and replicate. The single segment is positioned between a T7 promoter and the hepatitis delta virus ribozyme to produce an exact negative or positive transcript from the T7 polymerase. The plasmid vector and expression vectors comprising the necessary viral proteins are transfected into cells leading to production of recombinant viral particles (see, e.g., International Publication No. WO 01/04333; U.S. Pat. Nos. 7,442,379, 6,146,642, 6,649,372, 6,544,785 and 7,384,774; Swayne et al. (2003). *Avian Dis.* 47:1047-1050; and Swayne et al. (2001). *J. Virol.* 11868-11873, each of which is incorporated by reference in its entirety).

(71) Bicistronic techniques to produce multiple proteins from a single mRNA are known to one of skill in the art. Bicistronic techniques allow the engineering of coding sequences of multiple proteins into a single mRNA through the use of IRES sequences. IRES sequences direct the internal recruitment of ribosomes to the RNA molecule and allow downstream translation in a cap independent manner. Briefly, a coding region of one protein is inserted downstream of the ORF of a second protein. The insertion is flanked by an IRES and any untranslated signal sequences necessary for proper expression and/or function. The insertion must not disrupt the open reading frame, polyadenylation or transcriptional promoters of the second protein (see, e.g., Garcia-Sastre et al., 1994, *J. Virol.* 68:6254-6261 and Garcia-Sastre et al., 1994 *Dev. Biol. Stand.* 82:237-246, each of which are incorporated by reference herein in their entirety).

(72) Methods for cloning recombinant NDV to encode a transgene and express a heterologous protein encoded by the transgene (e.g., a transgene encoding an RSV F protein, a chimeric F protein, hMPV F protein) are known to one skilled in the art, such as, e.g., insertion of the transgene into a restriction site that has been engineered into the NDV genome, inclusion an appropriate signals in the transgene for recognition by the NDV RNA-dependent-RNA polymerase (e.g., sequences upstream of the open reading frame of the transgene that allow for the NDV polymerase to recognize the end of the previous gene and the beginning of the transgene, which may be, e.g., spaced by a single nucleotide intergenic sequence), inclusion of a valid Kozak sequence (e.g., to improve eukaryotic ribosomal translation); incorporation of a transgene that satisfies the “rule of six” for NDV cloning; and inclusion of silent mutations to remove extraneous gene end and/or gene start sequences within the transgene. Regarding the rule of six, one skilled in the art will understand that efficient replication of NDV (and more generally, most members of the paramyxoviridae family) is dependent on the genome length being a multiple of six, known as the “rule of six” (see, e.g., Calain, P. & Roux, L. The rule of six, a basic feature of efficient replication of Sendai virus defective interfering RNA. *J. Virol.* 67, 4822-4830 (1993)). Thus, when constructing a recombinant NDV described herein, care should be taken to satisfy the “Rule of Six” for NDV cloning. Methods known to one skilled in the art to satisfy the Rule of Six for NDV cloning may be used, such as, e.g., addition of nucleotides downstream of the transgene. See, e.g., Ayllon et al., Rescue of Recombinant Newcastle Disease Virus from cDNA. *J. Vis. Exp.* (80), e50830, doi:10.3791/50830 (2013) for a discussion of methods for cloning and rescuing of NDV (e.g., recombinant NDV), which is incorporated by reference herein in its entirety.

(73) In a specific embodiment, an NDV described herein (see, e.g., Sections 5.1 and 6) may be generated according to a method described in Section 6, *infra*.

(74) In a specific embodiment, a recombinant NDV comprising a packaged genome comprising a transgene encoding RSV F protein described herein comprises a LaSota strain backbone. In a specific embodiment, the genomic sequence of the La Sota strain backbone (i.e., without the transgene) is as set forth in SEQ ID NO:47.

(75) In a specific embodiment, a recombinant NDV comprising a packaged genome comprising a transgene encoding hMPV F protein described herein comprises a LaSota strain backbone. In a specific embodiment, the genomic sequence of the La Sota strain backbone (i.e., without the transgene) is as set forth in SEQ ID NO:47.

(76) In a specific embodiment, a recombinant NDV comprising a packaged genome comprising a transgene encoding a chimeric F protein described herein comprises a LaSota strain backbone. In a specific embodiment, the genomic sequence of the La Sota strain backbone (i.e., without the transgene) is as set forth in SEQ ID NO:47.

(77) 5.3 Propagation of NDVs

(78) The recombinant NDVs described herein (e.g., Sections 5.1 and 6) can be propagated in any substrate that allows the virus to grow to titers that permit the uses of the viruses described herein. In one embodiment, the substrate allows the recombinant NDVs described herein to grow to titers comparable to those determined for the corresponding wild-type viruses.

(79) The recombinant NDVs described herein (e.g., Sections 5.1 and 6) may be grown in cells (e.g., avian cells, chicken cells, etc.) that are susceptible to infection by the viruses, embryonated eggs (e.g., chicken eggs or quail eggs) or animals (e.g., birds). Such methods are well-known to those skilled in the art. In a specific embodiment, the recombinant NDVs described herein may be propagated in cancer cells, e.g., carcinoma cells (e.g., breast cancer cells and prostate cancer cells), sarcoma cells, leukemia cells, lymphoma cells, and germ cell tumor cells (e.g., testicular cancer cells and ovarian cancer cells). In another specific embodiment, the recombinant NDVs described herein may be propagated in cell lines, e.g., cancer cell lines such as HeLa cells, MCF7 cells, THP-1 cells, U87 cells, DU145 cells, Lncap cells, and T47D cells. In certain embodiments, the cells or cell lines (e.g., cancer cells or cancer cell lines) are obtained, derived, or obtained and derived from a human(s). In another embodiment, the recombinant NDVs described herein are propagated in interferon deficient systems or interferon (IFN) deficient substrates, such as, e.g., IFN deficient cells (e.g., IFN deficient cell lines) or IFN deficient embryonated eggs. In another embodiment, the recombinant NDVs described herein are propagated in chicken cells or embryonated chicken eggs. Representative chicken cells include, but are not limited to, chicken embryo fibroblasts and chicken embryo kidney cells. In a specific embodiment, the recombinant NDVs described herein are propagated in Vero cells. In another specific embodiment, the recombinant NDVs described herein are propagated in chicken eggs or quail eggs. In certain embodiments, a recombinant NDV virus described herein is first propagated in embryonated eggs and then propagated in cells (e.g., a cell line).

(80) The recombinant NDVs described herein may be propagated in embryonated eggs, e.g., from 6 to 14 days old, 6 to 12 days old, 6 to 10 days old, 6 to 9 days old, 6 to 8 days old, 8 to 10 day old, or 10 to 12 days old. In a specific embodiment, 10 day old embryonated chicken eggs are used to propagate the recombinant NDVs described herein. Young or immature embryonated eggs can be used to propagate the recombinant NDVs described herein. Immature embryonated eggs encompass eggs which are less than ten day old eggs, e.g., eggs 6 to 9 days old or 6 to 8 days old that are IFN-deficient. Immature embryonated eggs also encompass eggs which artificially mimic immature eggs up to, but less than ten day old, as a result of alterations to the growth conditions, e.g., changes in incubation temperatures; treating with drugs; or any other alteration which results in an egg with a retarded development, such that the IFN system is not fully developed as compared with ten to twelve day old eggs. The recombinant NDVs described herein can be propagated in different locations of the embryonated egg, e.g., the allantoic cavity. For a detailed discussion on the growth and propagation viruses, see, e.g., U.S. Pat. Nos. 6,852,522 and 7,494,808, both of which are hereby incorporated by reference in their entireties.

(81) For virus isolation, the recombinant NDVs described herein can be removed from embryonated eggs or cell culture and separated from cellular components, typically by well known clarification procedures, e.g., such as centrifugation, depth filtration, and microfiltration, and may be further purified as desired using procedures well known to those skilled in the art, e.g., tangential flow filtration (TFF), density gradient centrifugation, differential extraction, or chromatography.

(82) In a specific embodiment, virus isolation from allantoic fluid of an infected egg (e.g., a

chicken egg) begins with harvesting allantoic fluid, which is clarified using a filtration system to remove cells and other large debris, specifically, comprising a membrane having a net positive charge such that there is a measurable reduction in host cell DNA. The clarified bulk is subsequently processed by tangential flow filtration. The concentrated clarified bulk is then diafiltered against four diavolumes of high salt buffer, followed by four diavolumes of low salt formulation buffer and subsequently concentrated approximately 10-fold. Accordingly, residual egg proteins, e.g., primarily ovalbumin, and residual DNA are reduced to acceptable levels, and the buffer is exchanged to a buffer compatible with formulation of the recombinant NDV for a composition to be administered to a subject. The resulting product is then sterile filtered through a filter, e.g., a 0.2 μm filter, dispensed into appropriate sterile storage containers, frozen, and stored at -70 degrees Celsius.

(83) In a specific embodiment, a recombinant NDV described herein (see, e.g., Sections 5.1 and 6) is propagated, isolated, and/or purified according to a method described in Section 6. In a specific embodiment, a recombinant NDV described herein (see, e.g., Sections 5.1 and 6) is either propagated, isolated, or purified, or any two or all of the foregoing, using a method described in Section 6.

(84) In a specific embodiment, provided herein is a cell (e.g., a cell line) or embryonated egg (e.g., a chicken embryonated egg) comprising a recombinant NDV described herein. In another specific embodiment, provided herein is a method for propagating a recombinant NDV described herein, the method comprising culturing a cell (e.g., a cell line) or embryonated egg (e.g., a chicken embryonated egg) infected with the recombinant NDV. In some embodiments, the method may further comprise isolating or purifying the recombinant NDV from the cell or embryonated egg. In a specific embodiment, provided herein is a method for propagating a recombinant NDV described herein, the method comprising (a) culturing a cell (e.g., a cell line) or embryonated egg infected with a recombinant NDV described herein; and (b) isolating the recombinant NDV from the cell or embryonated egg. The cell or embryonated egg may be one described herein or known to one of skill in the art. In some embodiments, the cell or embryonated egg is IFN deficient.

(85) In a specific embodiment, provided herein is a method for producing a pharmaceutical composition (e.g., an immunogenic composition) comprising a recombinant NDV described herein, the method comprising (a) propagating a recombinant NDV described herein a cell (e.g., a cell line) or embryonated egg; and (b) isolating the recombinant NDV from the cell or embryonated egg. The method may further comprise adding the recombinant NDV to a container along with a pharmaceutically acceptable carrier.

(86) 5.4 Compositions and Routes of Administration

(87) Provided herein are compositions comprising a recombinant NDV described herein (e.g., Section 5.1 or 6). In a specific embodiment, the compositions are pharmaceutical compositions, such as immunogenic compositions (e.g., vaccine compositions). In a specific embodiment, provided herein are immunogenic compositions comprising a recombinant NDV described herein (e.g., Section 5.1 or 6). The compositions may be used in methods of inducing an immune response to RSV F protein or hMPV F protein. The compositions may be used in methods for immunizing against RSV or hMPV. The compositions may be used in methods for preventing an RSV disease or hMPV disease.

(88) In one embodiments, a pharmaceutical composition comprises a recombinant NDV described herein (e.g., Section 5.1 or 6), in an admixture with a pharmaceutically acceptable carrier. In some embodiments, the pharmaceutical composition further comprises one or more additional prophylactic or therapeutic agents, such as described in Section 5.5.4, *infra*. In a specific embodiment, a pharmaceutical composition comprises an effective amount of a recombinant NDV described herein (e.g., Section 5.1 or 6), and optionally one or more additional prophylactic or therapeutic agents, in a pharmaceutically acceptable carrier. In some embodiments, the recombinant NDV (e.g., Sections 5.1 or 6) is the only active ingredient included in the pharmaceutical

composition. In specific embodiments, two or more recombinant NDV are included in the pharmaceutical composition. In a particular embodiment, the pharmaceutical composition is an immunogenic composition.

(89) In a specific embodiment, a pharmaceutical composition comprises a first recombinant NDV and a second recombinant NDV, in an admixture with a pharmaceutically acceptable carrier, wherein the first recombinant NDV comprises a packaged genome comprising a first transgene, wherein the first transgene encodes a human RSV F protein, and wherein the second recombinant NDV comprises a packaged genome comprising a second transgene, wherein the second transgene encodes an hMPV F protein. In another specific embodiment, a pharmaceutical composition comprises a first recombinant NDV and a second recombinant NDV, in an admixture with a pharmaceutically acceptable carrier, wherein the first recombinant NDV comprises a packaged genome comprising a first transgene, wherein the first transgene encodes a human RSV F protein, wherein the second recombinant NDV comprises a packaged genome comprising a second transgene, and wherein the second transgene encodes a chimeric F protein comprising an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In another specific embodiment, a pharmaceutical composition comprises a first recombinant NDV and a second recombinant NDV, in an admixture with a pharmaceutically acceptable carrier, wherein the first recombinant NDV comprises a packaged genome comprising a first transgene, wherein the first transgene encodes an hMPV F protein, wherein the second recombinant NDV comprises a packaged genome comprising a second transgene, and wherein the second transgene encodes a chimeric F protein comprising a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In another specific embodiment, a pharmaceutical composition comprises a first recombinant NDV and a second recombinant NDV, in an admixture with a pharmaceutically acceptable carrier, wherein the first recombinant NDV comprises a packaged genome comprising a first transgene, wherein the first transgene encodes a first chimeric F protein comprising an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, wherein the second recombinant NDV comprises a packaged genome comprising a second transgene, and wherein the second transgene encodes a second chimeric F protein comprising a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. See, e.g., Section 5.1 or 6 for nucleic acid sequences encoding such transgenes. In a particular embodiment, the pharmaceutical composition is an immunogenic composition.

(90) In a specific embodiment, the recombinant NDV included in a pharmaceutical composition described herein is a live virus. In particular, embodiment, the recombinant NDV included in a pharmaceutical composition described herein is an attenuated live virus. In some embodiments, the recombinant NDV included in a pharmaceutical composition described herein is inactivated. In particular embodiments, the RSV F protein or hMPV F protein of the inactivated recombinant NDV is in the pre-fusion conformation.

(91) The pharmaceutical compositions provided herein can be in any form that allows for the composition to be administered to a subject. In a specific embodiment, the pharmaceutical compositions are suitable for veterinary administration, human administration, or both. As used herein, the term “pharmaceutically acceptable” means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeiae for use in animals, and more particularly in humans. The term “carrier” refers to a diluent, adjuvant, excipient, or vehicle with which the pharmaceutical composition is administered. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. Examples of suitable pharmaceutical carriers are described in “Remington's Pharmaceutical

Sciences” by E. W. Martin. The formulation should suit the mode of administration.

(92) In a specific embodiment, the pharmaceutical compositions are formulated to be suitable for the intended route of administration to a subject. For example, the pharmaceutical composition may be formulated to be suitable for parenteral, intravenous, intraarterial, intrapleural, inhalation, intranasal, intraperitoneal, oral, intradermal, colorectal, intraperitoneal, intracranial, and intratumoral administration. In one embodiment, the pharmaceutical composition may be formulated for intravenous, intraarterial, oral, intraperitoneal, intranasal, intratracheal, intrapleural, intracranial, subcutaneous, intramuscular, topical, pulmonary, or intratumoral administration. In a specific embodiment, the pharmaceutical composition may be formulated for intranasal administration.

(93) In a specific embodiment, the pharmaceutical composition comprising a recombinant NDV described herein (see, e.g., Sections 5.1 or 6) is formulated to be suitable for intranasal administration to the subject (e.g., human subject or bovine subject). In a specific embodiment, the pharmaceutical composition comprising a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding an RSV F protein, is formulated to be suitable for intranasal administration to the subject (e.g., human subject). In another specific embodiment, the pharmaceutical composition comprising a recombinant NDV is formulated to be suitable for intranasal administration to the subject (e.g., human subject), wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, and wherein the chimeric F protein comprises an RSV F protein ectodomain and transmembrane and cytoplasmic domains of an NDV F protein. In a particular embodiment, the pharmaceutical composition is an immunogenic composition.

(94) In a specific embodiment, the pharmaceutical composition comprising a recombinant NDV comprising a packaged genome comprising a transgene encoding an hMPV F protein is formulated to be suitable for intranasal administration to the subject (e.g., human subject). In another specific embodiment, the pharmaceutical composition comprising a recombinant NDV is formulated to be suitable for intranasal administration to the subject (e.g., human subject), wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, and wherein the chimeric F protein comprises an hMPV F protein ectodomain and transmembrane and cytoplasmic domains of an NDV F protein. In a particular embodiment, the pharmaceutical composition is an immunogenic composition.

(95) 5.5 Prophylactic Uses of a Recombinant NDV

(96) 5.5.1 Prevention of RSV Disease

(97) In another aspect, presented herein are methods for inducing an immune response in a subject (e.g., a human subject) comprising administering the subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a human RSV F protein. See, e.g., Section 5.1.2 and 6 for transgenes encoding a human RSV F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In a specific embodiment, the transgene comprises a codon optimized nucleic acid sequence encoding the human RSV F protein. In another aspect, presented herein are methods for inducing an immune response in a subject (e.g., a human subject) comprising administering the subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises the ectodomain of human RSV F protein and the transmembrane and cytoplasmic domains of NDV F protein. See, e.g., Section 5.1.2 and 6 for transgenes encoding a chimeric RSV F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In a specific embodiment, the ectodomain of the human RSV F protein is encoded by a codon optimized nucleic acid sequence.

(98) In another aspect, presented herein are methods for inducing an immune response to a RSV F

protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a human RSV F protein. See, e.g., Section 5.1.2 and 6 for transgenes encoding a chimeric RSV F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the human RSV F protein comprises the amino acid sequence set forth in SEQ ID NO: 6, 49, 50 or 58. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same human RSV F protein. For example, the human RSV F protein may be encoded by SEQ ID NO:1, 25, 27 or 29. Alternatively, the nucleic acid sequence encoding the human RSV F protein may be codon optimized, such as set forth in SEQ ID NO: 2, 26, 28, or 30. Specific examples of transgenes encoding a human RSV F protein include transgenes comprising an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID Nos: 1, 2, 25, 26, 27, 28, 29, and 30. In a specific embodiment, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a human RSV F protein. In a specific embodiment, the human RSV F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(99) In another aspect, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. See, e.g., Section 5.1.2 and 6 for transgenes encoding a human RSV F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:7. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:51. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO: 4, 44, 45, and 46). In a specific embodiment, the codon optimized sequence comprises the nucleic acid sequence of SEQ ID NO:4. In a specific embodiment, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(100) In another aspect, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a human RSV F protein. See, e.g., Section 5.1.2 and 6 for transgenes encoding a human RSV F protein which the packaged

genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the human RSV F protein comprises the amino acid sequence set forth in SEQ ID NO: 6, 49, 50 or 58. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same human RSV F protein. For example, the human RSV F protein may be encoded by SEQ ID NO:1, 25, 27 or 29. Alternatively, the nucleic acid sequence encoding the human RSV F protein may be codon optimized, such as set forth in SEQ ID NO: 2, 26, 28, or 30. In a specific embodiment, the codon optimized nucleic acid sequence comprises the nucleic acid sequence of SEQ ID NO:2. Specific examples of transgenes encoding a human RSV F protein include transgenes comprising an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID Nos: 1, 2, 25, 26, 27, 28, 29, and 30. In a specific embodiment, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a human RSV F protein. In a specific embodiment, the human RSV F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(101) In another aspect, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. See, e.g., Section 5.1.2 and 6 for transgenes encoding a chimeric F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO: 7. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:51. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO: 4, 44, 45, or 46). In a specific embodiment, the codon optimized nucleic acid sequence comprises the nucleic acid sequence of SEQ ID NO:4. In a specific embodiment, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(102) In another aspect, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a human RSV F protein. See, e.g., Section 5.1.2 and 6 for transgenes encoding a human RSV F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the human RSV F protein comprises the amino acid sequence set forth in SEQ ID NO: 6, 49, 50 or 58. Due to the degeneracy of the nucleic

acid code, a number of different nucleic acid sequences may encode for the same human RSV F protein. For example, the human RSV F protein may be encoded by SEQ ID NO:1, 25, 27 or 29. Alternatively, the nucleic acid sequence encoding the human RSV F protein may be codon optimized, such as set forth in SEQ ID NO: 2, 26, 28, or 30. Specific examples of transgenes encoding a human RSV F protein include transgenes comprising an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NOs: 1, 2, 25, 26, 27, 28, 29, and 30. In a specific embodiment, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a human RSV F protein. In a specific embodiment, the human RSV F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(103) In another aspect, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. See, e.g., Section 5.1.2 and 6 for transgenes encoding a chimeric F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO: 7. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:51. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO: 4, 44, 45, or 46). In a specific embodiment, the codon optimized nucleic acid sequence comprises the nucleic acid sequence of SEQ ID NO:4. In a specific embodiment, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(104) In another aspect, presented herein are methods for immunizing against RSV in a subject (e.g., a bovine subject) comprising administering the subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a bovine RSV F protein. See, e.g., Section 5.1.2 and 6 for transgenes encoding a bovine RSV F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In a specific embodiment, the transgene comprises a codon optimized nucleic acid sequence encoding the bovine RSV protein. In another aspect, presented herein are methods for inducing an immune response in a subject (e.g., a bovine subject) comprising administering the subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a

chimeric F protein, wherein the chimeric F protein comprises the ectodomain of bovine RSV F protein and the transmembrane and cytoplasmic domains of NDV F protein. See, e.g., Section 5.1.2 and 6 for transgenes encoding a chimeric F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In a specific embodiment, the ectodomain of the bovine RSV F protein is encoded by a codon optimized nucleic acid sequence.

(105) In another aspect, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a bovine RSV F protein. See, e.g., Section 5.1.2 and 6 for transgenes encoding a bovine RSV F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the bovine RSV F protein comprises the amino acid sequence set forth in SEQ ID NO: 10. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same bovine RSV F protein. For example, the bovine RSV F protein may be encoded by SEQ ID NO:9. Alternatively, the nucleic acid sequence encoding the bovine RSV F protein may be codon optimized, such as set forth in SEQ ID NO: 11. Specific examples of transgenes encoding a bovine RSV F protein include transgenes comprising an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NOs: 9, 11, 40, 41, 42, and 43. In a specific embodiment, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a bovine RSV F protein. In a specific embodiment, the bovine RSV F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(106) In another aspect, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. See, e.g., Section 5.1.2 and 6 for transgenes encoding a chimeric F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:33. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:33. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO:14, 38 or 39). In a specific embodiment, the codon optimized nucleic acid sequence comprises the nucleic acid sequence of SEQ ID NO:14. In a specific embodiment, presented herein are methods for inducing an immune response to a RSV F protein comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of

administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(107) In another aspect, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a bovine RSV F protein. See, e.g., Section 5.1.2 and 6 for transgenes encoding a bovine RSV F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the bovine RSV F protein comprises the amino acid sequence set forth in SEQ ID NO: 10. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same bovine RSV F protein. For example, the bovine RSV F protein may be encoded by SEQ ID NO:9. Alternatively, the nucleic acid sequence encoding the bovine RSV F protein may be codon optimized, such as set forth in SEQ ID NO: 11. Specific examples of transgenes encoding a bovine RSV F protein include transgenes comprising an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NOs: 9, 11, 40, 41, 42, and 43. In a specific embodiment, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a bovine RSV F protein. In a specific embodiment, the bovine RSV F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(108) In another aspect, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. See, e.g., Section 5.1.2 and 6 for transgenes encoding a chimeric F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:33. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:31. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO:14, 38, or 39). In a specific embodiment, the codon optimized nucleic acid sequence comprises the nucleic acid sequence of SEQ ID NO:14. In a specific embodiment, presented herein are methods for immunizing against RSV comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(109) In another aspect, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the

recombinant NDV comprises a packaged genome comprising a transgene encoding a bovine RSV F protein. See, e.g., Section 5.1.2 and 6 for transgenes encoding a bovine RSV F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the bovine RSV F protein comprises the amino acid sequence set forth in SEQ ID NO: 10. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same bovine RSV F protein. For example, the bovine RSV F protein may be encoded by SEQ ID NO:9. Alternatively, the nucleic acid sequence encoding the bovine RSV F protein may be codon optimized, such as set forth in SEQ ID NO: 11. Specific examples of transgenes encoding a bovine RSV F protein include transgenes comprising an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NOs: 9, 11, 40, 41, 42, and 43. In a specific embodiment, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding a bovine RSV F protein. In a specific embodiment, the bovine RSV F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(110) In another aspect, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. See, e.g., Section 5.1.2 and 6 for transgenes encoding a chimeric F protein which the packaged genome may comprise. See also Sections 5.1.3 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:33. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:33. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the RSV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO:14, 38 or 39). In a specific embodiment, the codon optimized nucleic acid sequence comprises the nucleic acid sequence of SEQ ID NO:14. In a specific embodiment, presented herein are methods for the prevention of RSV disease comprising administering to a subject (e.g., a bovine subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise a bovine RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(111) In another embodiment, presented herein is a recombinant NDV or composition thereof for inducing an immune response to an RSV F protein. In another embodiment, presented herein is a recombinant NDV or composition thereof for immunizing a subject (e.g., a human or bovine subject) against RSV. In another embodiment, presented herein is a recombinant NDV or composition thereof for the prevention of an RSV disease. See, e.g., Sections 5.1, and 6, *infra* for information regarding recombinant NDV, Section 5.5.4 for information regarding other therapies, and Section 5.4, *infra*, for information regarding compositions and routes of administration.

(112) The recombinant NDV described herein may be administered to a subject in combination with one or more other therapies. The recombinant NDV and one or more other therapies may be administered by the same or different routes of administration to the subject. In a specific embodiment, the recombinant NDV is administered to a subject intranasally. See, e.g., Sections 5.1, and 6, *infra* for information regarding recombinant NDV, Section 5.5.4 for information regarding other therapies, and Section 5.4, *infra*, for information regarding compositions and routes of administration.

(113) The recombinant NDV and one or more additional therapies may be administered concurrently or sequentially to the subject. In certain embodiments, the recombinant NDV and one or more additional therapies are administered in the same composition. In other embodiments, the recombinant NDV and one or more additional therapies are administered in different compositions. The recombinant NDV and one or more other therapies may be administered by the same or different routes of administration to the subject. Any route known to one of skill in the art or described herein may be used to administer the recombinant NDV and one or more other therapies. In a specific embodiment, the recombinant NDV is administered intranasally and the one or more other therapies is administered intravenously.

(114) In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a patient to prevent the onset of one, two or more symptoms of an RSV disease (such a patient may be at risk of developing an RSV infection). In a specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents the onset or development of one, two or more symptoms of RSV disease, reduces the severity of one, two or more symptoms of RSV disease, or prevents the onset or development of one, two or more symptoms of RSV disease and reduces the severity of one, two or more symptoms of RSV disease. Symptoms of RSV disease include congested or runny nose, cough, fever, sore throat, headache, wheezing, rapid or shallow breathing or difficulty breathing, bluish color the skin due to lack of oxygen, lack of appetite, lethargy and irritability.

(115) In a specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents the spread of RSV infection. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents hospitalization. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents pneumonia caused by RSV infection. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents otitis media caused by RSV infection. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents bronchiolitis caused by RSV infection. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents recurring RSV infections. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject reduces the likelihood of asthma linked to RSV infection.

(116) In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein induces antibodies to RSV F protein. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein induces both mucosal and systemic antibodies to RSV F protein (e.g., neutralizing antibodies). In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject induces neutralizing antibody to RSV F

protein. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject induces robust, long-lived (e.g., 6 months, 1 year, 2 years, 3 years or more), antigen-specific humoral immunity. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein does not induce a disease-promoting immune response. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof induces no or a low level of inflammation. In a particular embodiment, the administration of a recombinant NDV described herein induces no inflammation in the lung or nasal cavity as assessed by the lack of evidence of inflammatory infiltrates. In another embodiment, the administration of a recombinant NDV described herein induces a small amount of inflammation in the lung or nasal cavity that is not clinically significant. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject induces an IFN response. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents RSV infection of the lower airway and inhibits RSV replication in the upper airway.

(117) In a specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject results one, two, three or more, or all of the effects reported in Section 6 (e.g., Section 6.1, 6.2, or 6.4). In a specific embodiment, a method of preventing RSV disease or immunizing against RSV involves a protocol similar to or the same as that described in Section 6, *infra*.

(118) In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a subject suffering from an RSV disease. In other embodiments, an NDV (e.g., a recombinant NDV) described herein or a composition thereof, or a combination therapy described herein is administered to a subject predisposed or susceptible to an RSV disease. In some embodiments, an NDV (e.g., a recombinant NDV) or a composition thereof, or a combination therapy described herein is administered to a subject seronegative for RSV antibodies (e.g., antibodies to RSV F protein, RSV G protein, or both). In some embodiments, an NDV (e.g., a recombinant NDV) or a composition thereof, or a combination therapy described herein is administered to a subject seropositive for RSV antibodies (e.g., antibodies to RSV F protein, RSV G protein, or both). In certain embodiments, the subject is assessed for anti-RSV antibodies prior to administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein. In other embodiments, the subject is not assessed for anti-RSV antibodies prior to administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein.

(119) In certain embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a human that is 0 to 6 months old, 2 to 4 months old, 4 to 6 months old, 6 to 12 months old, 6 to 18 months old, 18 to 36 months old, 1 to 5 years old, 5 to 10 years old, 10 to 15 years old, 15 to 20 years old, 20 to 25 years old, 25 to 30 years old, 30 to 35 years old, 35 to 40 years old, 40 to 45 years old, 45 to 50 years old, 50 to 55 years old, 55 to 60 years old, 60 to 65 years old, 65 to 70 years old, 70 to 75 years old, 75 to 80 years old, 80 to 85 years old, 85 to 90 years old, 90 to 95 years old or 95 to 100 years old. In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a human infant. In another specific embodiment, the subject is a human infant six months old or older. In other embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a human toddler. In other embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a human child. In other embodiments, a recombinant NDV described herein or a composition thereof, or a

combination therapy described herein is administered to a human adult. In yet other embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to an elderly human.

(120) In a specific embodiment, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) in close contact with an individual with increased risk of an RSV or disease resulting from RSV infection (e.g., immunocompromised or immunosuppressed individuals). In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) who is pregnant. In other embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) who may or will be pregnant during the RSV season (e.g., generally, October to April (with peak RSV in December through February) in the Northern hemisphere). In specific embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) who is a woman who has given birth 1, 2, 3, 4, 5, 6, 7, or 8 weeks earlier. In specific embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) who is a parent (e.g., a mother or father) of children under the age of 18 years old.

(121) In particular embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) with a condition that increases susceptibility to RSV complications or for which RSV increases complications associated with the condition. Examples of conditions that increase susceptibility to RSV complications or for which RSV increases complications associated with the condition include conditions that affect the lung, such as cystic fibrosis, chronic obstructive pulmonary disease (COPD), emphysema, asthma, or bacterial infections (e.g., infections caused by *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Legionella pneumophila*, and *Chlamydia trachomatis*); cardiovascular disease (e.g., congenital heart disease, congestive heart failure, and coronary artery disease); endocrine disorders (e.g., diabetes); and neurological and neuron-developmental conditions (e.g., disorders of the brain, the spinal cord, the peripheral nerve, and muscle (such as cerebral palsy, epilepsy (seizure disorders), stroke, intellectual disability (e.g., mental retardation), muscular dystrophy, and spinal cord injury)).

(122) In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) that resides in a group home, such as a nursing home. In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) that works in, or spends a significant amount of time in, a group home, e.g., a nursing home. In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) that is a health care worker (e.g., a doctor or nurse). In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) that is a smoker.

(123) In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to (1) a subject (e.g., a human subject) who can transmit RSV to those at high risk for complications, such as, e.g., members of households with high-risk subjects, including households that include or will include human infants (e.g., infants younger than 6 months), (2) a subject coming into contact with human infants (e.g., infants less than 6 months of age), (3) a subject who is or will come into contact with subjects who live in nursing homes or other long-term care facilities, or (4) a subject who is or will come into contact with subjects with long-term disorders of the lungs, heart, or circulation; (5) subjects with metabolic diseases (e.g., diabetes) or subjects with weakened immune systems (including

immunosuppression caused by medications, malignancies such as cancer, organ transplant, or HIV infection); or (6) any combination of 1-5

(124) In certain embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a calf. In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a cow. In certain embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a cow undergoing or about to undergo transportation with other cattle.

(125) In certain embodiments a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a subject in an immunocompromised state or immunosuppressed state or at risk for becoming immunocompromised or immunosuppressed. In certain embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a subject that has or is at risk of getting RSV disease. For example, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a subject with COPD, cystic fibrosis or asthma. In some embodiments, a recombinant NDV described herein or a composition thereof is administered to a patient that is a transplant recipient.

(126) 5.5.2 Prevention of Human Metapneumovirus Disease

(127) In another aspect, presented herein are methods for inducing an immune response in a subject (e.g., a human subject) comprising administering the subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a hMPV F protein. See, e.g., Section 5.1.4 and 6 for transgenes encoding an hMPV F protein which the packaged genome may comprise. See also Sections 5.1.5 and 6 for examples of recombinant NDV that may be used in the methods. In a specific embodiment, the transgene comprises a codon optimized nucleic acid sequence encoding the hMPV F protein. In another aspect, presented herein are methods for inducing an immune response in a subject (e.g., a human subject) comprising administering the subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises the ectodomain of hMPV F protein and the transmembrane and cytoplasmic domains of NDV F protein. In a specific embodiment, the ectodomain of the hMPV F protein is encoded by a codon optimized nucleic acid sequence.

(128) In another aspect, presented herein are methods for inducing an immune response to an hMPV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a hMPV F protein. See, e.g., Section 5.1.4 and 6 for transgenes encoding an hMPV F protein which the packaged genome may comprise. See also Sections 5.1.5 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the hMPV F protein comprises the amino acid sequence set forth in SEQ ID NO: 17. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same hMPV F protein. In some embodiments, the hMPV F protein may be encoded by SEQ ID NO:16, 52, 54 or 56. Alternatively, the nucleic acid sequence encoding the hMPV F protein may be codon optimized, such as set forth in SEQ ID NO: 18, 53, 55 or 57. Specific examples of transgenes encoding an hMPV F protein include transgenes comprising RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NOs:16, 18, 52, 53, 54, 55, and 57. In a specific embodiment, presented herein are methods for inducing an immune response to an hMPV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding an hMPV F protein. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific

embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(129) In another aspect, presented herein are methods for inducing an immune response to an hMPV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. See, e.g., Section 5.1.4 and 6 for transgenes encoding a chimeric F protein which the packaged genome may comprise. See also Sections 5.1.5 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:15. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:32. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the hMPV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO: 19, 34, 35, and 36). In a specific embodiment, presented herein are methods for inducing an immune response to an hMPV F protein comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the hMPV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(130) In another aspect, presented herein are methods for immunizing against hMPV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a hMPV F protein. See, e.g., Section 5.1.4 and 6 for transgenes encoding an hMPV F protein which the packaged genome may comprise. See also Sections 5.1.5 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the hMPV F protein comprises the amino acid sequence set forth in SEQ ID NO: 17. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same hMPV F protein. In some embodiments, the hMPV F protein may be encoded by SEQ ID NO:16, 52, 54 or 56. Alternatively, the nucleic acid sequence encoding the hMPV F protein may be codon optimized, such as set forth in SEQ ID NO: 18, 53, 55, or 57. Specific examples of transgenes encoding an hMPV F protein include transgenes comprising an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NOs:16, 18, 52, 53, 54, 55, and 57. In a specific embodiment, presented herein are methods for immunizing against hMPV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding an hMPV F protein. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(131) In another aspect, presented herein are methods for immunizing against hMPV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein,

wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. See, e.g., Section 5.1.4 and 6 for transgenes encoding a chimeric F protein which the packaged genome may comprise. See also Sections 5.1.5 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:15. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:32. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the hMPV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO: 19, 34, 35, and 36). In a specific embodiment, presented herein are methods for immunizing against hMPV comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the hMPV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(132) In another aspect, presented herein are methods for the prevention of hMPV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a hMPV F protein. See, e.g., Section 5.1.4 and 6 for transgenes encoding an hMPV F protein which the packaged genome may comprise. See also Sections 5.1.5 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the hMPV F protein comprises the amino acid sequence set forth in SEQ ID NO: 17. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for the same hMPV F protein. In some embodiments, the hMPV F protein may be encoded by SEQ ID NO:16, 52, 54 or 56. Alternatively, the nucleic acid sequence encoding the hMPV F protein may be codon optimized, such as set forth in SEQ ID NO: 18, 53, 55 or 57. Specific examples of transgenes encoding an hMPV F protein include transgenes comprising an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NOs:16, 18, 52, 53, 54, 55, and 57. In a specific embodiment, presented herein are methods for the prevention of hMPV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding an hMPV F protein. In a specific embodiment, the hMPV F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly.

(133) In another aspect, presented herein are methods for the prevention of hMPV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. See, e.g., Section 5.1.4 and 6 for transgenes encoding a chimeric F protein which the packaged genome may comprise. See also Sections 5.1.5 and 6 for examples of recombinant NDV that may be used in the methods. In one embodiment, the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:15. Due to the degeneracy of the nucleic acid code, a number of different nucleic acid sequences may encode for

the same chimeric F protein. For example, the chimeric F protein may be encoded by SEQ ID NO:32. Alternatively, the nucleic acid sequence encoding one or more domains of the chimeric F protein may be codon optimized (e.g., the nucleic acid sequence encoding the hMPV F protein ectodomain may be codon optimized, such as set forth in SEQ ID NO: 19, 34, 35, and 36). In a specific embodiment, presented herein are methods for the prevention of hMPV disease comprising administering to a subject (e.g., a human subject) a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprise an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the transgene comprises a codon optimized nucleic acid sequence encoding the hMPV F protein ectodomain. In a specific embodiment, the chimeric F protein is expressed by cells infected with the recombinant NDV. The recombinant NDV may be administered to a subject by any route of administration. In another specific embodiment, the recombinant NDV is administered to a subject intranasally. In some embodiments, the recombinant NDV is administered to a subject intramuscularly. In another embodiment, presented herein is a recombinant NDV or composition thereof for inducing an immune response to an hMPV F protein. In another embodiment, presented herein is a recombinant NDV or composition thereof for immunizing a subject (e.g., a human subject) against hMPV. In another embodiment, presented herein is a recombinant NDV or composition thereof for the prevention of an hMPV disease. See, e.g., Sections 5.1, and 6, *infra* for information regarding recombinant NDV, Section 5.5.4 for information regarding other therapies, and Section 5.4, *infra*, for information regarding compositions and routes of administration.

(134) The recombinant NDV described herein may be administered to a subject in combination with one or more other therapies. The recombinant NDV and one or more other therapies may be administered by the same or different routes of administration to the subject. In a specific embodiment, the recombinant NDV is administered to a subject intranasally. See, e.g., Sections 5.1, and 6, *infra* for information regarding recombinant NDV, Section 5.5.4 for information regarding other therapies, and Section 5.4, *infra*, for information regarding compositions and routes of administration.

(135) In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a patient to prevent the onset of one, two or more symptoms of an hMPV disease (e.g., such a patient is at risk of developing an hMPV infection). In a specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents the onset or development of one, two or more symptoms of hMPV disease, reduces the severity of one, two or more symptoms of hMPV disease, or prevents the onset or development of one, two or more symptoms of hMPV disease and reduces the severity of one, two or more symptoms of hMPV disease. Symptoms of hMPV disease include nasal congestion, runny nose, fever, cough, sore throat, wheezing, difficulty breathing, lack of appetite, lethargy and irritability.

(136) In a specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents the spread of hMPV infection. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents hospitalization. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents pneumonia caused by hMPV infection. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject reduces the likelihood of asthma linked to hMPV infection.

(137) In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein induces antibodies to hMPV F protein. In another specific embodiment, the administration of a recombinant NDV described

herein or a composition thereof, or a combination therapy described herein induces both mucosal and systemic antibodies to hMPV F protein (e.g., neutralizing antibodies). In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject induces neutralizing antibody to hMPV F protein. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein does not induce a disease-promoting immune response. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof induces no or a low level of inflammation. In a particular embodiment, the administration of a recombinant NDV described herein induces no inflammation in the lung or nasal cavity as assessed by the lack of evidence of inflammatory infiltrates. In another embodiment, the administration of a recombinant NDV described herein induces a small amount of inflammation in the lung or nasal cavity that is not clinically significant. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject induces robust, long-lived (e.g., 6 months, 1 year, 2 years, 3 years or more), antigen-specific humoral immunity. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject induces an IFN response. In another specific embodiment, the administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein to a subject prevents hMPV infection of the lower airway and inhibits hMPV replication in the upper airway.

(138) In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a subject suffering from an hMPV disease. In other embodiments, an NDV (e.g., a recombinant NDV) described herein or a composition thereof, or a combination therapy described herein is administered to a subject predisposed or susceptible to an hMPV disease. In some embodiments, an NDV (e.g., a recombinant NDV) or a composition thereof, or a combination therapy described herein is administered to a subject seronegative for hMPV antibodies (e.g., antibodies to hMPV F protein, hMPV G protein or both). In some embodiments, an NDV (e.g., a recombinant NDV) or a composition thereof, or a combination therapy described herein is administered to a subject seropositive for hMPV antibodies (e.g., antibodies to hMPV F protein, hMPV G protein, both). In certain embodiments, the subject is assessed for anti-hMPV antibodies prior to administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein. In other embodiments, the subject is not assessed for anti-hMPV antibodies prior to administration of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein.

(139) In certain embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a human that is 0 to 6 months old, 2 to 4 months old, 4 to 6 months old, 6 to 12 months old, 6 to 18 months old, 18 to 36 months old, 1 to 5 years old, 5 to 10 years old, 10 to 15 years old, 15 to 20 years old, 20 to 25 years old, 25 to 30 years old, 30 to 35 years old, 35 to 40 years old, 40 to 45 years old, 45 to 50 years old, 50 to 55 years old, 55 to 60 years old, 60 to 65 years old, 65 to 70 years old, 70 to 75 years old, 75 to 80 years old, 80 to 85 years old, 85 to 90 years old, 90 to 95 years old or 95 to 100 years old. In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a human infant. In another specific embodiment, the subject is a human infant six months old or older. In other embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a human toddler. In other embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a human child. In other embodiments, a recombinant NDV described herein or a composition thereof, or a

combination therapy described herein is administered to a human adult. In yet other embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to an elderly human.

(140) In a specific embodiment, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) in close contact with an individual with increased risk of an hMPV infection or disease resulting from hMPV infection (e.g., immunocompromised or immunosuppressed individuals). In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) who is pregnant. In other embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) who may or will be pregnant during the hMPV season (e.g., generally, November to July in the Northern hemisphere). In specific embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) who is a woman who has given birth 1, 2, 3, 4, 5, 6, 7, or 8 weeks earlier. In specific embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) who is a parent (e.g., a mother or father) of children under the age of 18 years old.

(141) In particular embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) with a condition that increases susceptibility to hMPV complications or for which hMPV increases complications associated with the condition. Examples of conditions that increase susceptibility to hMPV complications or for which hMPV increases complications associated with the condition include conditions that affect the lung, such as cystic fibrosis, chronic obstructive pulmonary disease (COPD), emphysema, asthma, or bacterial infections (e.g., infections caused by *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Legionella pneumophila*, and *Chlamydia trachomatis*); cardiovascular disease (e.g., congenital heart disease, congestive heart failure, and coronary artery disease); endocrine disorders (e.g., diabetes); and neurological and neuron-developmental conditions (e.g., disorders of the brain, the spinal cord, the peripheral nerve, and muscle (such as cerebral palsy, epilepsy (seizure disorders), stroke, intellectual disability (e.g., mental retardation), muscular dystrophy, and spinal cord injury)).

(142) In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) that resides in a group home, such as a nursing home. In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) that works in, or spends a significant amount of time in, a group home, e.g., a nursing home. In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) that is a health care worker (e.g., a doctor or nurse). In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered a subject (e.g., a human subject) that is a smoker.

(143) In some embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to (1) a subject (e.g., a human subject) who can transmit hMPV to those at high risk for complications, such as, e.g., members of households with high-risk subjects, including households that include or will include human infants (e.g., infants younger than 6 months), (2) a subject coming into contact with human infants (e.g., infants less than 6 months of age), (3) a subject who is or will come into contact with subjects who live in nursing homes or other long-term care facilities, (4) a subject who is or will come into contact with subjects with long-term disorders of the lungs, heart, or circulation; (5) subjects with metabolic diseases (e.g., diabetes) or subjects with weakened immune systems (including immunosuppression

caused by medications, malignancies such as cancer, organ transplant, or HIV infection); or (6) any combination of 1-5.

(144) In certain embodiments a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a subject in an immunocompromised state or immunosuppressed state or at risk for becoming immunocompromised or immunosuppressed. In certain embodiments, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a subject that has or is at risk of getting hMPV disease. For example, a recombinant NDV described herein or a composition thereof, or a combination therapy described herein is administered to a subject with COPD, cystic fibrosis or asthma. In some embodiments, a recombinant NDV described herein or a composition thereof is administered to a patient that is a transplant recipient.

(145) 5.5.3 Dosage and Frequency

(146) The amount of a recombinant NDV or a composition thereof, which will be effective in the prevention of RSV disease or hMPV disease, or immunization against RSV or hMPV will depend on the route of administration, the general health of the subject, etc. Standard clinical techniques, such as in vitro assays, may optionally be employed to help identify dosage ranges. However, suitable dosage ranges of a recombinant NDV for administration are generally about 10^{sup.4} to about 10^{sup.12}, and can be administered to a subject once, twice, three, four or more times with intervals as often as needed. In certain embodiments, dosages similar to those currently being used in clinical trials for NDV are administered to a subject.

(147) In certain embodiments, a recombinant NDV or a composition thereof is administered to a subject as a single dose followed by a second dose 1 to 6 weeks, 1 to 5 weeks, 1 to 4 weeks, 1 to 3 weeks, 1 to 2 weeks, 6 to 12 weeks, 3 to 6 months, 6 to 9 months, 6 to 12 months, or 6 to 9 months later. In accordance with these embodiments, booster inoculations may be administered to the subject at 3 to 6 month or 6 to 12 month intervals following the second inoculation.

(148) In certain embodiments, administration of the same recombinant NDV or a composition thereof may be repeated and the administrations may be separated by at least 1 day, 2 days, 3 days, 5 days, 6 days, 7 days, 10 days, 14 days, 15 days, 21 days, 28 days, 30 days, 45 days, 2 months, 75 days, 3 months, or at least 6 months. In other embodiments, administration of the same recombinant NDV or a composition thereof may be repeated and the administrations may be separated by 1 to 14 days, 1 to 7 days, 7 to 14 days, 1 to 30 days, 15 to 30 days, 15 to 45 days, 15 to 75 days, 15 to 90 days, 1 to 3 months, 3 to 6 months, 3 to 12 months, or 6 to 12 months. In some embodiments, a first recombinant NDV or a composition thereof is administered to a subject followed by the administration of a second recombinant NDV or a composition thereof. In some embodiments, the first and second recombinant NDV are different from each other. For example, the first recombinant NDV may comprise a packaged genome comprising a transgene encoding an RSV F protein, and the second recombinant NDV may comprise a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In another example, the first recombinant NDV may comprise a packaged genome comprising a transgene encoding an hMPV F protein, and the second recombinant NDV may comprise a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In certain embodiments, the first and second recombinant NDVs or compositions thereof may be separated by at least 1 day, 2 days, 3 days, 5 days, 6 days, 7 days, 10 days, 14 days, 15 days, 21 days, 28 days, 30 days, 45 days, 2 months, 75 days, 3 months, or at least 6 months. In other embodiments, the first and second recombinant NDVs or compositions thereof may be separated by 1 to 14 days, 1 to 7 days, 7 to 14 days, 1 to 30 days, 15 to 30 days, 15 to 45 days, 15 to 75 days, 15 to 90 days, 1 to 3 months, 3 to 6 months, 3 to 12 months, or 6 to 12 months.

(149) In certain embodiments, a recombinant NDV or composition thereof is administered to a

subject in combination with one or more additional therapies, such as a therapy described in Section 5.5.4, *infra*. The dosage of the other one or more additional therapies will depend upon various factors including, e.g., the therapy, the route of administration, the general health of the subject, etc. and should be decided according to the judgment of a medical practitioner. In specific embodiments, the dose of the other therapy is the dose and/or frequency of administration of the therapy recommended for the therapy for use as a single agent is used in accordance with the methods disclosed herein. Recommended doses for approved therapies can be found in the Physician's Desk Reference.

(150) In certain embodiments, a recombinant NDV or composition thereof is administered to a subject concurrently with the administration of one or more additional therapies. In a specific embodiment, a first pharmaceutical composition comprising a first recombinant NDV is administered to a subject in combination with a second pharmaceutical composition comprising a second recombinant NDV, wherein the first recombinant NDV comprises a packaged genome comprising a first transgene, wherein the first transgene encodes a human RSV F protein, and wherein the second recombinant NDV comprises a packaged genome comprising a second transgene, wherein the second transgene encodes an hMPV F protein. In another specific embodiment, a first pharmaceutical composition comprising a first recombinant NDV is administered to a subject in combination with a second pharmaceutical composition comprising a second recombinant NDV, wherein the first recombinant NDV comprises a packaged genome comprising a first transgene, wherein the first transgene encodes a human RSV F protein, wherein the second recombinant NDV comprises a packaged genome comprising a second transgene, and wherein the second transgene encodes a chimeric F protein comprising an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In another specific embodiment, a first pharmaceutical composition comprising a first recombinant NDV is administered to a subject in combination with a second pharmaceutical composition comprising a second recombinant NDV, wherein the first recombinant NDV comprises a packaged genome comprising a first transgene, wherein the first transgene encodes an hMPV F protein, wherein the second recombinant NDV comprises a packaged genome comprising a second transgene, and wherein the second transgene encodes a chimeric F protein comprising a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In another specific embodiment, a first pharmaceutical composition comprising a first recombinant NDV is administered to a subject in combination with a second pharmaceutical composition comprising a second recombinant NDV, wherein the first recombinant NDV comprises a packaged genome comprising a first transgene, wherein the first transgene encodes a first chimeric F protein comprising an hMPV RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, wherein the second recombinant NDV comprises a packaged genome comprising a second transgene, and wherein the second transgene encodes a second chimeric F protein comprising a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. See, e.g., Section 5.1 or 6 for nucleic acid sequences encoding such transgenes. In certain embodiments, the first and second pharmaceutical compositions are administered concurrently to the subject, or within 1 minute, 2 minutes, 5 minutes, 10 minutes, 15 minutes, 20 minutes, 30 minutes, 45 minutes, 60 minutes, 1.5 hours, 2 hours, 3 hours, 4 hours, 5 hours, or 6 hours of each other. In certain embodiments, the first and second pharmaceutical compositions are administered to the subject within 12 hours, 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks or 12 weeks of each other. In certain embodiments, the first and second pharmaceutical compositions are administered to the subject within 3-6 months, 6-9 months, 6-12 months, or 3 months, 4 months, 6 months, 9 months, or 12 months of each other.

(151) In certain embodiments, a first pharmaceutical composition is administered to a subject as a priming dose and after a certain period (e.g., 1 month, 2 months, 3 months, 4 months, 5 months, 6

months, or 1-6 months) a booster dose of a second pharmaceutical composition is administered. In some embodiments, the first pharmaceutical composition comprises a first recombinant NDV and a second recombinant NDV, wherein the first recombinant NDV comprises a packaged genome comprising a transgene encoding a human RSV F protein, and the second recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In certain embodiments, the first pharmaceutical composition comprises a first recombinant NDV and a second recombinant NDV, wherein the first recombinant NDV comprises a packaged genome comprising a transgene encoding an hMPV F protein, and the second recombinant NDV comprises a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In some embodiments, the first pharmaceutical composition comprises a first recombinant NDV and a second recombinant NDV, wherein the first recombinant NDV comprises a packaged genome comprising a transgene encoding a human RSV F protein, and the second recombinant NDV comprises a packaged genome comprising a transgene encoding an hMPV F protein. In certain embodiments, the first pharmaceutical composition comprises a first recombinant NDV and a second recombinant NDV, wherein the first recombinant NDV comprises a packaged genome comprising a transgene encoding a first chimeric F protein, and the second recombinant NDV comprises a packaged genome comprising a transgene encoding a second chimeric F protein, wherein the first chimeric F protein comprises a human RSV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains, and wherein the second chimeric F protein comprises an hMPV F protein ectodomain and NDV F protein transmembrane and cytoplasmic domains. In some embodiments, the second pharmaceutical composition comprises the same recombinant NDV as the first pharmaceutical composition. In another embodiment, the second pharmaceutical composition comprises the same recombinant NDV as the first pharmaceutical composition with the exception that the strain of NDV used as the backbone of the virus is different.

(152) 5.5.4 Additional Therapies

(153) Additional therapies that can be used in a combination with a recombinant NDV described herein or a composition thereof include, but are not limited to, acetaminophen, ibuprofen, throat lozenges, cough suppressants, inhalers, antibiotics and oxygen. In a specific embodiment, the additional therapy is a second recombinant NDV described herein.

(154) 5.6 Biological Assays

(155) In a specific embodiment, one, two or more of the assays described in Section 6 may be used to characterize a recombinant NDV described herein, or an F protein or a chimeric F protein.

(156) 5.6.1 In Vitro Viral Assays

(157) Viral assays include those that indirectly measure viral replication (as determined, e.g., by plaque formation) or the production of viral proteins (as determined, e.g., by western blot analysis) or viral RNAs (as determined, e.g., by RT-PCR or northern blot analysis) in cultured cells in vitro using methods which are well known in the art.

(158) Growth of the recombinant NDVs described herein can be assessed by any method known in the art or described herein (e.g., in cell culture (e.g., cultures of chicken embryonic kidney cells or cultures of chicken embryonic fibroblasts (CEF)) (see, e.g., Section 6). Viral titer may be determined by inoculating serial dilutions of a recombinant NDV described herein into cell cultures (e.g., CEF, MDCK, EFK-2 cells, Vero cells, primary human umbilical vein endothelial cells (HUVEC), H292 human epithelial cell line or HeLa cells), chick embryos, or live animals (e.g., avians). After incubation of the virus for a specified time, the virus is isolated using standard methods. Physical quantitation of the virus titer can be performed using PCR applied to viral supernatants (Quinn & Trevor, 1997; Morgan et al., 1990), hemagglutination assays, tissue culture infectious doses (TCID₅₀) or egg infectious doses (EID₅₀). An exemplary method of assessing

viral titer is described in Section 6, below.

(159) Incorporation of nucleotide sequences encoding a heterologous peptide or protein (e.g., a transgene into the genome of a recombinant NDV described herein can be assessed by any method known in the art or described herein (e.g., in cell culture, an animal model or viral culture in embryonated eggs)). For example, viral particles from cell culture of the allantoic fluid of embryonated eggs can be purified by centrifugation through a sucrose cushion and subsequently analyzed for protein expression by Western blotting using methods well known in the art. In a specific embodiment, a method described in Section 6, *infra*, is used to assess the incorporation of a transgene into the genome of a recombinant NDV.

(160) Immunofluorescence-based approaches may also be used to detect virus and assess viral growth. Such approaches are well known to those of skill in the art, e.g., fluorescence microscopy and flow cytometry (see, e.g., Section 6, *infra*). Methods for flow cytometry, including fluorescence activated cell sorting (FACS), are available (see, e.g., Owens, et al. (1994) *Flow Cytometry Principles for Clinical Laboratory Practice*, John Wiley and Sons, Hoboken, NJ; Givan (2001) *Flow Cytometry*, 2nd ed.; Wiley-Liss, Hoboken, NJ; Shapiro (2003) *Practical Flow Cytometry*, John Wiley and Sons, Hoboken, NJ). Fluorescent reagents suitable for modifying nucleic acids, including nucleic acid primers and probes, polypeptides, and antibodies, for use, e.g., as diagnostic reagents, are available (Molecular Probes (2003) *Catalogue*, Molecular Probes, Inc., Eugene, OR; Sigma-Aldrich (2003) *Catalogue*, St. Louis, MO). See, e.g., the assays described in Section 6, *infra*.

(161) Standard methods of histology of the immune system are described (see, e.g., Muller-Harmelink (ed.) (1986) *Human Thymus: Histopathology and Pathology*, Springer Verlag, New York, NY; Hiatt, et al. (2000) *Color Atlas of Histology*, Lippincott, Williams, and Wilkins, Philadelphia, PA; Louis, et al. (2002) *Basic Histology: Text and Atlas*, McGraw-Hill, New York, NY). See also Section 6, *infra*, for histology and immunohistochemistry assays that may be used.

(162) 5.6.2 Interferon Assays

(163) IFN induction and release by a recombinant NDV described herein may be determined using techniques known to one of skill in the art. For example, the amount of IFN induced in cells following infection with a recombinant NDV described herein may be determined using an immunoassay (e.g., an ELISA or Western blot assay) to measure IFN expression or to measure the expression of a protein whose expression is induced by IFN. Alternatively, the amount of IFN induced may be measured at the RNA level by assays, such as Northern blots and quantitative RT-PCR, known to one of skill in the art. In specific embodiments, the amount of IFN released may be measured using an ELISPOT assay. Further, the induction and release of cytokines and/or interferon-stimulated genes may be determined by, e.g., an immunoassay or ELISPOT assay at the protein level and/or quantitative RT-PCR or northern blots at the RNA level.

(164) 5.6.3 Activation Marker Assays and Immune Cell Infiltration Assay

(165) Techniques for assessing the expression of T cell marker, B cell marker, activation marker, co-stimulatory molecule, ligand, or inhibitory molecule by immune cells are known to one of skill in the art. For example, the expression of T cell marker, B cell marker, an activation marker, co-stimulatory molecule, ligand, or inhibitory molecule by an immune cell can be assessed by flow cytometry. In a specific embodiment, an assay described in Section 6, *infra*, is used to assess infiltration of a type(s) of immune cell.

(166) 5.6.4 Toxicity Studies

(167) In some embodiments, the recombinant NDVs described herein or compositions thereof, or combination therapies described herein are tested for cytotoxicity in mammalian, preferably human, cell lines. In certain embodiments, cytotoxicity is assessed in one or more of the following non-limiting examples of cell lines: U937, a human monocyte cell line; primary peripheral blood mononuclear cells (PBMC); Huh7, a human hepatoblastoma cell line; HL60 cells, HT1080, HEK 293T and 293H, MLPC cells, human embryonic kidney cell lines; human melanoma cell lines,

such as SkMel2, SkMel-119 and SkMel-197; THP-1, monocytic cells; a HeLa cell line; and neuroblastoma cells lines, such as MC-IXC, SK-N-MC, SK-N-MC, SK-N-DZ, SH-SY5Y, and BE(2)-C. In some embodiments, the ToxLite assay is used to assess cytotoxicity.

(168) Many assays well-known in the art can be used to assess viability of cells or cell lines following infection with a recombinant NDV described herein or composition thereof, and, thus, determine the cytotoxicity of the recombinant NDV or composition thereof. For example, cell proliferation can be assayed by measuring Bromodeoxyuridine (BrdU) incorporation, (.sup.3H) thymidine incorporation, by direct cell count, or by detecting changes in transcription, translation or activity of known genes such as proto-oncogenes (e.g., fos, myc) or cell cycle markers (Rb, cdc2, cyclin A, D1, D2, D3, E, etc.). The levels of such protein and mRNA and activity can be determined by any method well known in the art. For example, protein can be quantitated by known immunodiagnostic methods such as ELISA, Western blotting or immunoprecipitation using antibodies, including commercially available antibodies. mRNA can be quantitated using methods that are well known and routine in the art, for example, using northern analysis, RNase protection, or polymerase chain reaction in connection with reverse transcription. Cell viability can be assessed by using trypan-blue staining or other cell death or viability markers known in the art. In a specific embodiment, the level of cellular ATP is measured to determined cell viability. In preferred embodiments, a recombinant NDV described herein or composition thereof does not kill healthy (i.e., non-cancerous) cells.

(169) In specific embodiments, cell viability may be measured in three-day and seven-day periods using an assay standard in the art, such as the CellTiter-Glo Assay Kit (Promega) which measures levels of intracellular ATP. A reduction in cellular ATP is indicative of a cytotoxic effect. In another specific embodiment, cell viability can be measured in the neutral red uptake assay. In other embodiments, visual observation for morphological changes may include enlargement, granularity, cells with ragged edges, a filmy appearance, rounding, detachment from the surface of the well, or other changes.

(170) The recombinant NDVs described herein or compositions thereof, or combination therapies can be tested for in vivo toxicity in animal models. For example, animal models, known in the art to test the effects of compounds on RSV infection or hMPV infection can also be used to determine the in vivo toxicity of the recombinant NDVs described herein or compositions thereof, or combination therapies. For example, animals are administered a range of pfu of a recombinant NDV described herein, and subsequently, the animals are monitored over time for various parameters, such as one, two or more of the following: lethality, weight loss or failure to gain weight, and levels of serum markers that may be indicative of tissue damage (e.g., creatine phosphokinase level as an indicator of general tissue damage, level of glutamic oxalic acid transaminase or pyruvic acid transaminase as indicators for possible liver damage). These in vivo assays may also be adapted to test the toxicity of various administration mode and regimen in addition to dosages.

(171) The toxicity, efficacy or both of a recombinant NDV described herein or a composition thereof, or a combination therapy described herein can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Therapies that exhibit large therapeutic indices are preferred.

(172) The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage of the therapies for use in subjects. The dosage of such agents lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any therapy described herein, the therapeutically effective dose can be estimated initially from cell culture assays.

(173) 5.6.5 Biological Activity Assays

(174) The recombinant NDVs described herein or compositions thereof, or combination therapies described herein can be tested for biological activity using animal models for inhibiting RSV disease, hMPV disease, antibody response to the recombinant NDVs, etc. (see, e.g., Section 6). Such animal model systems include, but are not limited to, rats, mice, hamsters, cotton rats, chicken, cows, monkeys (e.g., African green monkey), pigs, dogs, rabbits, etc.

(175) In a specific embodiment, the recombinant NDVs described herein or compositions thereof, or combination therapies described herein may be tested using animal models for the ability to induce a certain geometric mean titer of antibody(ies) that binds to the RSV F protein (e.g., human RSV F protein or bovine F protein) or hMPV F protein. In another specific embodiment, the recombinant NDVs described herein or compositions thereof, or combination therapies described herein may be tested using animal models for the ability to induce antibodies that have neutralizing activity against RSV F protein (e.g., human RSV F protein or bovine F protein) or hMPV F protein in a microneutralization assay. For example, the microneutralization assay described in Section 6, *infra*, may be used to assess neutralizing activity. In some embodiments, the recombinant NDVs described herein or compositions thereof, or combination therapies described herein may be tested using animal models for the ability to induce a certain geometric mean titer of antibody(ies) that binds to the RSV F protein (e.g., human RSV F protein or bovine F protein) or hMPV F protein and neutralizes RSV F protein (e.g., human RSV F protein or bovine F protein) or hMPV F protein in a microneutralization assay. In a specific embodiment, the recombinant NDVs described herein or compositions thereof, or combination therapies described herein may be tested using animal models for the ability to induce a certain fold increase in levels of antibody(ies) that binds to RSV F protein (e.g., human RSV F protein or bovine F protein) or hMPV F protein post-immunization with a recombinant NDV described herein or a composition thereof relative to the levels of such antibody pre-immunization. For example, a 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold or greater increase in levels of antibody(ies) that binds to RSV F protein (e.g., human RSV F protein or bovine F protein) or hMPV F protein post-immunization with a recombinant NDV described herein or a composition thereof relative to the levels of such antibody(ies) pre-immunization.

(176) 5.6.6 Expression of Transgene

(177) Assays for testing the expression of RSV F protein, chimeric F protein, or hMPV F protein in cells infected with a recombinant NDV comprising a packaged genome comprising a transgene encoding RSV F protein, chimeric F protein, or hMPV F protein, respectively may be conducted using any assay known in the art, such as, e.g., western blot, immunofluorescence, and ELISA, or any assay described herein (see, e.g., Section 6).

(178) In a specific aspect, ELISA is utilized to detect expression of RSV F protein, chimeric F protein, or hMPV F protein in cells infected with a recombinant NDV comprising a packaged genome comprising a transgene encoding of RSV F protein, chimeric F protein, or hMPV F protein. In a specific embodiment, RSV F protein is quantified using an ELISA such as described in Section 6, *infra*.

(179) In one embodiment, RSV F protein or chimeric F protein encoded by a packaged genome of a recombinant NDV described herein is assayed for proper folding by testing its ability to bind specifically to an anti-RSV F protein antibody (e.g., Synagis®) using any assay for antibody-antigen interaction known in the art. In another embodiment, hMPV F protein or chimeric F protein encoded by a packaged genome of a recombinant NDV described herein is assayed for proper folding by testing its ability to bind specifically to an anti-hMPV F protein antibody using any assay for antibody-antigen interaction known in the art. In another embodiment, an RSV F protein, a chimeric F protein, or an hMPV F protein encoded by a packaged genome of a recombinant NDV described herein is assayed for proper folding by determination of the structure or conformation of the RSV F protein, chimeric F protein, or hMPV F protein, respectively using any method known in the art such as, e.g., NMR, X-ray crystallographic methods, or secondary structure prediction

aagcaaagctgcagcatatcaaatatagcaactgtgatagagttccaacaaa
agaacaacagactactagagattaccaggggaatttagtgtaatgcagggtg
aactacacctgtaagcacttacatgtaactaatagtgaattattgtcatta
atcaatgatatgcctataacaaatgatcagaaaaagttaatgtccaacaatg
ttcaaataggttagacagcaaagttactctatcatgtccataataaaagagga
agtcttagcatatgtagtacaattaccactatatggtgttatagatacaccc
tggttgaaactacacacatccccctctatgtacaaccaacacaaaagaagggt
ccaacatctgtttaacaagaactgacagaggatggtactgtgacaatgcagg
atcagtatctttctcccacaagctgaaacatgtaaagttcaatcaaatcga
gtattttgtgacacaatgaacagtttaacattaccaagtgaagtaaattctt
gcaatgttgacatatccaaccccaaatatgattgtaaaattatgacttcaa
aacagatgtaagcagctccgttatcacatctctaggagccattgtgtcatgc
tatggcaaaactaaatgtacagcatccaataaaaaatcgtggaatcataaaga
cattttctaacgggtgcgattatgtatcaaataaaggggtggacactgtgtc
tgtaggtaacacattatattatgtaaataagcaagaaggtaaaagtctctat
gtaaaagggtgaaccaataataaatttctatgaccattagtattcccctctg
atgaatttgatgcatcaatatctcaagtcaacgagaagattaaccagagcct
agcatttattcgtaaatccgatgaattattacataatgtaaatgctggtaaa
tccaccataaatatcatgataactactataattatagtgattatagtaatat
tggtatcattaattgctgttgactgctcttatactgtaaggccagaagcac
accagtcacactaagcaaagatcaactgagtggtataaataatattgcattt agtaactaa Codon-
ATGGAGCTGCTGATCCTGAAGGCCAACGCCATCACCCAC 2 optimized
CATCCTGACCGCCGTGACCTTCTGCTTCGCCAGCGGCCA nucleic acid
GAACATCACCGAGGAGTTCTACCAGAGCACCTGCAGCG sequence
CCGTGAGCAAGGGCTACCTGAGCGCCCTGCGCACCGGC encoding human
TGGTACACCAGCGTGATCACCATCGAGCTGAGCAACAT RSV F of A2
CAAGGAGAACAAGTGCAACGGCACCGACGCCAAGGTG strain (codon
AAGCTGATCAAGCAGGAGCTGGACAAGTACAAGAACG optimized
CCGTGACCGAGCTGCAGCTGCTGATGCAGAGCACCCCC version of SEQ
GCCACCAACAACCGCGCCCGCCGCGAGCTGCCCCGCTT ID NO:1)
CATGAACTACACCCTGAACAACGCCAAGAAGACCAACG
TGACCCTGAGCAAGAAGCGCAAGCGCCGCTTCCTGGGC
TTCCTGCTGGGCGTGGGCAGCGCCATCGCCAGCGGCGT
GGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTGA
ACAAGATCAAGAGCGCCCTGCTGAGCACCAACAAGGCC
GTGGTGAGCCTGAGCAACGGCGTGAGCGTGCTGACCAG
CAAGGTGCTGGACCTGAAGAACTACATCGACAAGCAGC
TGCTGCCCATCGTGAACAAGCAGAGCTGCAGCATCAGC
AACATCGCCACCGTGATCGAGTTCCAGCAGAAGAACAA
CCGCCTGCTGGAGATCACCCGCGAGTTCAGCGTGAACG
CCGGCGTGACCACCCCGTGAGCACCTACATGCTGACC
AACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCAT
CACCAACGACCAGAAGAAGCTGATGAGCAACAACGTG
CAGATCGTGCGCCAGCAGAGCTACAGCATCATGAGCAT
CATCAAGGAGGAGGTGCTGGCCTACGTGGTGCAGCTGC
CCCTGTACGGCGTGATCGACACCCCTGCTGGAAGCTG
CACACCAGCCCCCTGTGCACCACCAACACCAAGGAGGG
CAGCAACATCTGCCTGACCCGACCGACAGGGGCTGGT
ACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCCCAG
GCCGAGACCTGCAAGGTGCAGAGCAACCGCGTGTTCTG

CGACACCATGCAACGTGGACATCTTCAACCCCAAGTACGAC
ACCTGTGCAACGTGGACATCTTCAACCCCAAGTACGAC
TGCAAGATCATGACCAGCAAGACCGACGTGAGCAGCAG
CGTGATCACCAGCCTGGGCGCCATCGTGAGCTGCTACG
GCAAGACCAAGTGCACCGCCAGCAACAAGAACAGGGG
CATCATCAAGACCTTCAGCAACGGCTGCGACTACGTGA
GCAACAAGGGCGTGACACCGTGAGCGTGGGCAACAC
CCTGTACTACGTGAACAAGCAGGAGGGCAAGAGCCTGT
ACGTGAAGGGCGAGCCCATCATCAACTTCTACGACCCC
CTGGTGTTCCCCAGCGACGAGTTTCGACGCCAGCATCAG
CCAGGTGAACGAGAAGATCAACCAGAGCCTGGCCTTCA
TCCGCAAGAGCGACGAGCTGCTGCACAACGTGAACGCC
GGCAAGAGCACCATCAACATCATGATCACCACCATCAT
CATCGTGATCATCGTGATCCTGCTGAGCCTGATCGCCGT
GGGCCTGCTGCTGTACTGCAAGGCCCGCAGCACCCCCG
TGACCCTGAGCAAGGACCAGCTGAGCGGCATCAACAAC ATCGCCTTCAGCAACTAA
cDNA of the full accaaacagagaatccgtgagttacgataaaaggcgaaggagcaattg 3 genome of NDV
aagtcgcacgggtagaagggtgaatctcgagtgcgagcccgaagcac strain LaSota
aaactcgagaaaagccttctgccaacatgtctccgtatttgatgagta with the
cgaacagctcctcgcggctcagactcgcccaatggagctcatggagg transgene
gggagaaaaagggagtagcttaaaagtagacgtcccggtattcactct comprising the
taacagtgatgaccagaagatagatggagctttgtggtattctgcct codon-
ccggattgctgttagcgaagatgccaacaaaccactcaggcaaggtgc optimized
tctcatatcttttatgctccactcacaggaatgaggaacatgt nucleic acid
tgccCttgcagggaaacagaatgaagccacattggccgtgcttgagat sequence (SEQ
tgatggctttgccaacggcacgccccagttcaacaataggagtggagt ID NO: 2)
gtctgaagagagagcacagagatttgcatgatagcaggatctctccc encoding human
tcgggcatgcagcaacggaaccccgttcgtcacagccggggcCgaaga RSV F protein
tgatgcaccagaagacatcaccgataccctggagaggatcctctctat of A2 strain
ccaggctcaagtatgggtcacagtagcaaaagccatgactgcgtatga inserted between
gactgcagatgagtcggaaacaaggcgaatcaataagtatatgcagca the NDV P and
aggcaggttcaaaaagaaatacatcctctaccccgtagcaggagcac M genes (The
aatccaactcacgatcagacagctctcttgagtcgcatcttttgggt Sac II
tagcgagctcaagagaggccgcaacacggcaggtggtaccttactta (CCGCGG)
ttataacctggtaggggacgtagactcatacatcaggaataccgggct restriction sites
tactgcattcttctgacactcaagtacggaatcaacaccaagacatc used to insert the
agcccttgacttagtagcctctcaggcgacatccagaagatgaagca codon optimized
gctcatgcgtttgtatcggtgaaaggagataatgcgccgtacatgac human RSV F
attacttggtgatagtaccagatgagctttgcgcctgccgagtatgc protein open
acaactttactcctttgccatgggtatggcatcagtcctagataaagg reading frame
tactgggaaataccaatttgccagggactttatgagcacatcattctg into the genomic
gagacttgagtagagtacgctcaggctcagggaagtagcattaacga sequence are
ggatatggctgccgagctaaagctaaccccgagcagcaaGgaGgggcct double
ggcagctgctgccaacgggtctccgaGgaGaccagcagcataGacat underlined. The
gcctactcaacaagtcggagtcctcactgggcttagcgaggggggggtc initiation and
ccaagctctacaaggcggatcgaatagatcgcaagggaaccagaagc stop codons of
cggggatggggagaccaattcctggatctgatgagagcggtagcaaa the codon
tagcatgagggagggcgccaaactctgcacagggcactcccaatcggg optimized human
gcctcccccaactcctgggccatccaagataacgacaccgactgggg RSV F open
gtattgatggacaaaaccagcctgcttcacaaaaacatcccaatgc reading frame

cctcacccgtagtcgacccctcgatttgcggctctatatgaccacacc are underlined.
ctcaaacaacatccccctcttctcctccccctgctgtacaactA The open reading
cgTacgccctagataccacaggcacaatgcgggctcactaacaatcaaa from of the
acagagccgaggggaattagaaaaaagtacgggtagaagagggatattc codon optimized
agagatcagggcaagtctcccgagtctctgctctctctctacctgat human RSV F is
agaccaggacaaacatggccacctttacagatgcagagatcgacgagc in **bold**.

tatttgagacaagtggaaactgtcattgacaacataattacagcccagg
gtaaaccagcagagactgttgggaaggagtgcaatcccacaaggcaaga
ccaaggtgctgagcgcagcatgggagaagcatgggagcatccagccac
cggccagtcaagacaaccccgatcgacaggacagatctgacaaacaac
catccacacccgagcaaacgaccccgcatgacagcccgcggccacat
ccgccgaccagccccccacccaggccacagacgaagccgtcgacacac
agCtcaggaccggagcaagcaactctctgctgttgatgcttgacaagc
tcagcaataaatcgccaatgctaaaaagggcccatggtcgagcccc
aagaggggaatcaccaacgtccgactcaacagcaggggagtcaacca
gtcgcggaaacagtacaggaaagaccgcagaaccaagtcaaggccgccc
ctggaaaccagggcacagacgtgaacacagcatatcatggacaatggg
aggagtcacaactatcagctgggtgcaacccctcatgctctccgatcaa
ggcagagccaagacaatacccttgtatctgcggatcatgtccagccac
ctgtagactttgtgcaagcgatgatgtctatgatggaggcgatatcac agagagtaagtaagggtgactatcagctagatcttgtcttgaaacaga
catcctccatccctatgatgcgggtccgaaatccaacagctgaaaacat
ctgttcgagtcattggaagccaacttgggaatgatgaagattctggatc ccggttgccaacatttcattctctgagtgatctacgggcagttgccc
gatctcacccgggttttagtttcaggccctggagacccctctccctatg
tgacacaaggaggcgaaatggcacttaataaaactttcgcaaccagtg
cacatccatctgaattgattaaacccgccactgcatgcgggcctgata
taggagtggaaggacactgtccgtgcattgatcatgtcacgccc
tgaccccgagttcttcagccaagctcctaagcaagttagatgcagccg
ggtcgatcgaggaaatcaggaaaatcaagcgcttgctctaaatggct
aattactactgccacacgtagcgggtccctgtccactcggcacacac
ggaatctgcaccgagttccccccgcGgTTAGAAAAAATACGGGTAGA

ACCGCCACC**ATGGAGCTGCTGATCCTGAAGGCCAACGCCATCACCACC**
ATCCTGACCGCCGTGACCTTCTGCTTCGCCAGCGGCCAGAACATCACC
GAGGAGTTCTACCAGAGCACCTGCAGCGCCGTGAGCAAGGGCTACCTG
AGCGCCCTGCGCACCGGCTGGTACACCAGCGTGATCACCATCGAGCTG
AGCAACATCAAGGAGAACAAGTGCAACGGCACCGACGCCAAGGTGAAG
CTGATCAAGCAGGAGCTGGACAAGTACAAGAACGCCGTGACCGAGCTG
CAGCTGCTGATGCAGAGCACCCCCGCCACCAACAACCGCGCCCCGCCGC
GAGCTGCCCCGCTTCATGAACTACACCCTGAACAACGCCAAGAAGACC
AACGTGACCCTGAGCAAGAAGCGCAAGCGCCGCTTCCTGGGCTTCCTG
CTGGGCGTGGGCAGCGCCATCGCCAGCGGCGTGGCCGTGAGCAAGGTG
CTGCACCTGGAGGGCGAGGTGAACAAGATCAAGAGCGCCCTGCTGAGC
ACCAACAAGGCCGTGGTGAGCCTGAGCAACGGCGTGAGCGTGCTGACC
AGCAAGGTGCTGGACCTGAAGAACTACATCGACAAGCAGCTGCTGCCC
ATCGTGAACAAGCAGAGCTGCAGCATCAGCAACATCGCCACCGTGATC
GAGTTCCAGCAGAAGAACAACCGCCTGCTGGAGATCACCCGCGAGTTC
AGCGTGAACGCCGGCGTGACCACCCCCGTGAGCACCTACATGCTGACC
AACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCATCACCAACGAC
CAGAAGAAGCTGATGAGCAACAACGTGCAGATCGTGCGCCAGCAGAGC
TACAGCATCATGAGCATCATCAAGGAGGAGGTGCTGGCCTACGTGGTG
CAGCTGCCCCCTGTACGGCGTGATCGACACCCCTGCTGGAAGCTGCAC

ACCAGCCCCCTGTGACACCAAGGAGGGCAGCAACATCTGC
CTGACCCCGCACCGACaGgGGCTGGTACTGCGACAACGCCGGCAGCGTG
AGCTTCTTCCCCCAGGCCGAGACCTGCAAGGTGCAGAGCAACCGCGTG
TTCTGCGACACCATGAACAGCCTGACCCTGCCCAGCGAGGTGAACCTG
TGCAACGTGGACATCTTCAACCCCAAGTACGACTGCAAGATCATGACC
AGCAAGACCGACGTGAGCAGCAGCGTGATCACCAGCCTGGGCGCCATC
GTGAGCTGCTACGGCAAGACCAAGTGCACCGCCAGCAACAAGAACaGg
GGCATCATCAAGACCTTCAGCAACGGCTGCGACTACGTGAGCAACAAG
GGCGTGGACACCGTGAGCGTGGGCAACACCCTGTACTACGTGAACAAG
CAGGAGGGCAAGAGCCTGTACGTGAAGGGCGAGCCCATCATCAACTTC
TACGACCCCCTGGTGTTCCCCAGCGACGAGTTCGACGCCAGCATCAGC
CAGGTGAACGAGAAGATCAACCAGAGCCTGGCCTTCATCCGCAAGAGC
GACGAGCTGCTGCACAACGTGAACGCCGGCAAGAGCACCATCAACATC
ATGATCACCACCATCATCATCGTGATCATCGTGATCCTGCTGAGCCTG
ATCGCCGTGGGCCTGCTGCTGTACTGCAAGGCCCGCAGCACCCCCGTG
ACCCTGAGCAAGGACCAGCTGAGCGGCATCAACAACATCGCCTTCAGC
AACTAAccccccgcggaaccaaggtccaactctccaagcggcaatcct
ctctcgcttctcagccccactgaatgAtcgcgtaaccgtaattaatc
tagctacatttaagattaagaaaaatacgggtagaattggagtgccc caattgtgccaagatggactcatctaggacaattgggctgtactttga
ttctgccattcttctagcaacctgttagcatttccgatcgtcctaca
agAcacaggagatgggaagaagcaaatcgccccgcaatataggatcca
gcgcttgacttgtggactgataagtaggaggactcagtattcatcac
cacctatggattcatctttcaagttgggaatgaagaagccacCgtcgg
catgatcatgataaacccaagcgcgagttactttccgctgcgatgct
ctgcctaggaagcgtcccaaataccggagaccttattgagctggcaag
ggcctgtctcactatgatagtcacatgcaagaagagtgaactaatac
tgagagaatggttttctcagtagtgaggcaccccaagtgctgcaaag
ctgtaggggttgaggcaaaataactcatcagtgaatgcagtcaagca
cgtgaaagcgccagagaagattcccgggagtggaaaccctagaatacaa
ggatgaactttgtctccttgactgtggtaggaccgaagaGggatgtctacaa gatcccagctgcagttgaagggttctggctcgagtctgtacaatct
tgcgctcaatgtcactattaatgtggaggtagacccgaggagtccctt ggtaaactctCtgtctaagtctgacagcggaatactatgctaacctct
cttgcatattggacttatgaccacTgtagataggaaggggaagaaagt
gacatttgacaagctggaaaagaaaataaggagccttgatctatctgt
cgggctcagtgatgtgctcgggccttccgtgttggtgtaaaagcaagagg
tgcacggactaagcttttggcaccttctctctagcagtgaggacagc ctgctatcccatagcaaagtcttctcctcaggtggccaagatactctg
gagtcaaaccgcgtgcctgcggagcgttaaaatcattatccaagcagg
taccacacgcgctgtcgcagtgaccgccgaccacgaggttacctctac
taagctggagaaggggcacacccttgccaaatacaatccttttaagaa
ataagctgcgtctctgagattgcgctccgcccactcaccagatcatc atgacacaaaaaactaatctgtcttgattattacagttagttacct
gtctatcaagttagaaaaaacacgggtagaagattctggatcccgggt
ggcgccctccaggtgcaagatgggctccagaccttctaccaagaaccc
agcacctatgatgtgactatccgggttgcgctggtagtgcattgcat
ctgtccggcaaacctcattgatggcaggcctcttgagctgcaggaat
tgtggttacaggagacaaagccgtcaacatatacactcatcccagac
aggatcaatcatagttaagctcctcccgaatctgcccaggataagga
ggcatgtgcgaaagcccccttgatgcatacaacaggacattgaccac
tttgctaccccccttggtgactctatccgtaggatacaagagtctgt
gactacatctggaggggggagacagggggcgcttataggcgccattat
tggcgggtgtggctcttgggggtgcaactgccgcacaaataacagcggc
cgcagctctgatacaagccaaacaaatgctgccaacatcctccgact

taaagagagcattgccgcaaccaatgaggctgtgcatgaggtcactga
cggattatcgcaactagcagtgaggcagttgggaagatgcagcagtttgt
taatgaccaatttaataaaaacagctcaggaattagactgcatcaaaat
tgcacagcaagttgggtgtagagctcaacctgtacctaaccgaattgac
tacagtattcggaccacaaatcacttcacctgctttaaacaagctgac tattcaggcactttacaatctagctgggtggaaatatggattacttatt
gactaagttagggtgtaggggaacaatcaactcagctcattaatcggtag cggcttaatcacccggaaccctattctatacgaactcacagactcaact
cttgggtatacaggttaactctaccttcagtcgggaacctaaataatat
gcgtgccacctacttggaaccttatccgtaagcacaaccaggggatt
tgcctcggcacttgtcccAaaagtgggtgacacaggctcgggttctgtgat agaagaacttgacacctatactgtatagaaactgacttagatttata
ttgtacaagaatagtaacgtttccctatgtcccctggattttattctg cttgagcggcaatacgtcggcctgtatgtactcaaagaccgaaggcgc
acttactacaccatacatgactatcaaaggttcagtcacgcgaactg
caagatgacaacatgtagatgtgtaaacccccgggtatcatatcgca
aaactatggagaagccgtgtctctaatagataaacaatcatgcaatgt ttatccttaggcgggataactttaaggctcagtggggaattcgatgt
aacttatcagaagaatatctcaatacaagatttcaagtaataataac aggcaatcttgatatctcaactgagcttgggaatgtcaacaactcgat
cagtaatgctttgaataagtttagaggaaaagcaacagaaaactagacaa agtcaatgtcaaactgactagcacatctgctctcattacctatatcgt
tttgactatcatatctctgtttttgggtatacttagcctgattctagc atgctacctaatgtacaagcaaaaaggcgcacaacaaaagaccttattatg
gcttgggaataataactctagatcagatgagagccactacaaaaatgtg
aacacagatgaggaacgaagggttccctaataagtaatttgtgtgaaag ttctggtagtctgtcagttcagagagttaagaaaaaactaccgggtgt
agatgaccaaaggacgatatacgggtagaacggtaagagaggccgccc
ctcaattgcgagccaggcttcacaacctccgttctaccgcttcaccga
caacagtcctcaatcatggaccgcgccgttagccaagttgcgttagag
aatgatgaaagagaggcaaaaaatacatggcgcttgatattccggatt gcaatcttattcttaacagtagtgaccttggctatatctgtagcctcc
cttttatatagcatgggggctagcacacctagcgatctttaggcata
ccgactaggatttccagggcagaagaaaagattacatctacacttgg
tccaatcaagatgtagtagataggatatataagcaagtggcccttgag tctccgttggcattgttaaatactgagaccacaattatgaacgcaata
acatctctctcttatcagattaatggagctgcaaacaacagtggggtgg
ggggcacctatccatgaccagattatatagggggggataggcaaagaa
ctcattgtagatgatgctagtgtacatcatttctatccctctgca ttcaagaacatctgaattttatcccggcgctactacaggatcaggt
tgcactcgaataccctcatttgacatgagtgtacccattactgtac acccataatgtaattgtctggatgcagagatcactcacattcatat
cagtatttagcacttgggtgtgctccggacatctgcaacaggggagggtta ttcttttactctgcgttccatcaacctggacgacacccaaaatcgg
aagtcttgcagtgtagtgcaactcccctgggttgatgtgtgtgc
tcgaaagtcacggagacagaggaagaagattataactcagctgtccct
acgcggatggtacatgggaggttaggggtcgacggccagtagccacgaa
aaggacctagatgtcacaacattattcggggactgggtggccaactac
ccaggagtaggggggtggatctttattgacagccgcgtatggttctca
gtctacggaggggttaaaacccaattcacccagtgacactgtacaggaa
gggaaatatgtgatatacaagcgatacaatgacacatgccagatgag
caagactaccagattcgaatggccaagtcttcgtataagcctggacgg
tttgggtgggaaacgcatacagcaggctatcttatctatcaaggtgtca
acatccttaggcgaagaccgggtactgactgtaccgccaacacagtc
acactcatgggggcccgaaggcagaattctcacagtagggacatctcat ttcttgatcaacgagggtcatcatacttctctcccgcgttattatat
cctatgacagtcagcaaaaaacagccactcttcatagtccttataca ttcaatgccttcactcggccaggtagatcccttgccaggcttcagca
agatgccccaaactcgtgtgttactggagtctatacagatccatatccc ctaatcttctatagaaaccacaccttgcgaggggtattcgggacaatg
cttgatgggtgtacaagcaagacttaacctgcgtctgcagtattcgat
agcacatcccgcagtcgcattactcgagtgagttcaagcagtagcaaaa
gcagcatacacaacatcaactgttttaaagtgggtcaagactaataag acctattgtctcagcattgctgaaatatctaatactctcttcggagaa
ttcagaatcgccccgttactagttgagatcctcaagatgacgggggt
agagaagccaggctcggtagttgagtcaattataaaggagttggaaa
gatggcattgtatcacctatcttctgcgacatcaagaatcaaaccgaa
tgccggcgcggtgctcgaattccatggtgccaagttgaccacaatcagcc

agtgctcatgcgatcagattaagccttgatcaAtaGtctcttgattaag
aaaaaatgtaagtggcaatgagatacaaggcaaacagctcatgggTa
aCaatacgggtaggacatggcgagctccggctcctgaaagggcagagca
tcagattatcctaccagagTcacacctgtcttcaccattgggtcaagca caaactactctattactggaaattaactgggctaccgcttctgatga
atgtgacttcgaccacctcattctcagccgacaatggaaaaaatact
tgaatcggcctctcctgatactgagagaatgataaaactcggaagggc
agtacaccaaactcttaaccacaattccagaataaccggagtgctcca
ccccaggtgttagaaGaactggctaattagggtcccagattcaac
caacaaatttcggaagattgagaagaagatccaaattcacaacacgag
atatggagaactgttcacaaggctgtgtacgcatatagagaagaaact
gctggggctcatcttggtctaacaatgtcccccggtcagaggagttcag
cagcattcgtacggatccggcattctggtttcactcaaaatgggtccac
agccaagtttgcatggctccatataaaacagatccagaggcatctgat
gggtggcagctaGgacaaggctctgcgccaacaaattgggtgatgctaac
ccataaggtaggccaagtctttgtcactcctgaacttgcgttgtagc gcatacgaatgagaacaagttcacatgtcttaccaggaacttgatt
gatgtatgcagatatgatggagggcagagatatgggtcaacataatc
aaccacggcggtgcatctcagaagcttatcagagaaaattgatgacat
tttgcggttaatagacgctctggcaaaagacttgggtaataagctta cgatgttgatcactaatggagggatttgcatacggagctgtccagct
actcgagccgtcaggtacatttgcaggagatttcttcgcattcaacct gcaggagcttaaagacattctaattggcctcctcccaatgatatagc
agaatccgtgactcatgcaatcgctactgtattctctggttagaaca gaatcaagcagctgagatgttggtgtgtgtgctgtgggggtcaccc
actgcttgagctccgtattgcagcaaaaggcagtcaggagccaaatgtg cgcaccgaaaatggtagactttgatatgatccttcagggtactgtcttt
cttcaagggaacaatcatcaacgggtacagaaagaagaatgcaggtgt
gtggccgcgagtcaaagtggatacaatatatgggaagggtcattgggca
actacatgcagattcagcagagatttcacacgatatcatgttgagaga gtataagagtttatctgcacttgaatttgagccatgtatagaatatga
ccctgtcaccaacctgagcatgttcctaaaagacaaggcaatcgaca
ccccaacgataattggcttgctcgtttaggcggaaccttctctccga
agaccagaagaaacatgtaaaagaagcaacttcgactaatcgctctt gatagagtttttagagtcaaataatttgatccatataaagagatgga
atatctgacgaccttgagtaccttagagatgacaatgtggcagatc
atactcgctcaaggagaagggaagtgaaggttaatggacggatcttcgc
taagctgacaaagaagttaagggaactgtcaggtgatggcggaagggat
cctagccgatcagattgcacctttcttcagggaatggagtcattca ggatagcatatccttgaccaagagtatgctagcgatgagtcactgtc
tttaacagcaataaagaacgtatcactgactgtaaagaaagagtatc
ttcaaacgcgaatcatgatccgaaaagcaagaaccgtcggagagttgc
aaccttcataacaactgacctgcaaaagtactgtcttaattggagata tcagacaatcaaattgttcgctcatgccatcaatcagttgatgggcct
acctcacttctcgaatggattcacctaagactgatggacactacgat gttcgtaggagaccctttcaatcctccaagtgacctactgactgtga
cctctcaagagtcctaatgatgacatatatattgtcagtgccagagg
gggtatcgaaggattatgccagaagctatggacaatgatctcaattgc tgcaatccaacttgctgcagctagatcgcatgtcgtgttgctgtat
ggtacaggggtgataatcaagtaatagcagtaacgagagaggtgaatc
agacgactctccggagatgggtgttgacacagttgcatcaagccagtgta taatttctcaagggaattaattcatgtcaatcatttgattggccataa
tttgaaggatcgtgaaaccatcaggtcagacacattcttcatatacag
caaacgaatcttcaaatgaggcaatcctcagtcagtcctcaaaaaa ttcactctaaattagtgtcaggtgatctcagtgaaaacaccgt
aatgtcctgtgccaacattgcctctactgtagcacggctatgcgagaa cgggcttccaaagacttctgttactatttaaactatataatgagttg
tgtgcagacatactttgactctgagtttccatcaccaacaattcgca ccccgatcttaatcagtcgtggattgaggacatctctttgtgcactc
atatgttctgactcctgccaattagggggactgagtaaccttcaata ctcaaggctctacactagaaatacggtgacccggggactactgcttt
tgacagagatcaagcgactagaagcagtgaggattactgagtcctaact
tatgactaatatcttaactaggccgcctgggaatggagattgggccag tctgtgcaacgaccatactctttcaatttgagactgttgcaagccc
aaatattgttcttaagaaacatacgcaaagagtcctatttgaaacttg ttcaaatccctattgtctggagtgacacagaggataatgaggcaga
agagaaggcattggctgaattcttgcttaatcaagaggtgattcatcc
ccgcgttgcgcatgccatcatggaggcaagctctgtaggtaggagaaa
gcaaatcaagggtgttgacacaacaaacaccgtaattaagattgc

gcttactaggaggccattaggcatcaagaggctgatgcggatagtc aa ttattctagcatgcatgcaatgctgtttagagacgatgtttttcctc
cagtagatccaaccaccccttagtctcttctaataatgtgttctctgac
actggcagactatgcacggaatagaagctggtcacctttgacgggagg
caggaaaatactgggtgtatctaactctgatac gatagaactcgtaga
gggtgagattcttagtgtaagcggagggtgtacaagatgtgacagcgg
agatgaacaatttacttggtccatcttccaagaatataagaattgac
cgatgacaccagcaagaatcctccgatgagggtaccatatctcgggtc
aaagacacaggagaggagagctgcctcacttgcaaaaatagctcatat
gtcgccacatgtaaaggctgcctaagggcacatccgtgttgatctg ggcttatggggataatgaagtaaattggactgctgctcttacgattgc
aaaatctcgggtgaatgtaaacttagagtatcttcgggttactgtcccc ttaccacacggctgggaatcttcaacatagactagatgatggataac
tcagatgacattcaccctgcattctcttacaggGgtgcacttacat tcacatatccaatgattctcaaaggctgttactgaagaaggagtcaa
agagggggaatgtggtttaccaacagatcatgctcttgggtttatctct aatcgaatcgatctttccaatgacaacaaccaggacatatgatgagat
cacactgcacctacatagtaaatttagttgctgtatcagagaagcacc
tgttgccggttccttcgagctacttgggggtggtagccgaactgaggac agtgacctcaaataagtttatgtatgatcctagccctgtatcggagggg
agactttgcgagacttgacttagctatcttcaagagttatgagcttaa tctggagtcatacccacgatagagctaataaacattctttcaatc
cagcggggaagtgtgattggccagtctgtggtttcttatgatgaagatac ctccataaagaatgacgccataatagtgtatgacaatacccgaattg
gatcagtgaaagctcagaattcagatgtgggtccgcctatttgaatatgc agcacttgaagtgtcctcgactgttcttaccactctattacctgag
agtaagaggcctGgacaatatgtcttatatatgggtgattatacaa gaatatgccaggaattctactttcaacattgcagctacaatatctca
tcccgctattcattcaaggttacatgcagtgggcctgggtcaacctga cggatcacaccaacttgagatacggattttatcgaaatgtctgcaa
actattagtatcttgcacccgacgtgtgatctccggcttatattcagg aaataagtatgatctgctgttcccatctgtcttagatgataacctgaa
tgagaagatgcttcagctgatatcccgggttatgtgtctgtacacggg actctttgtacaacaagagaaatcccgaataaagaggcttaactgc
agaagagaaatgttcaactcactgagtatttactgtcggatgtctgt gaaaccattacttagccccgatcaagtgagctctatcatgtctcctaa
cataattacattcccagctaattctgtactacatgtctcggagagcct caatttgatcaggggaaaggaggacagggatactatctggcggtgtt
gttccccaagagccattattagagttcccttctgtgcaagatattgg tgctcgagtgaagatccattcacccgacaacctgcggcattttgca
agagttagatttgagtgtccagcaagggtatgacgcattcacacttag tcagattcatcctgaactcacatctccaaatccggaggaagactactt
agtacgatacttggtcagagggatagggactgcattcttcttggta taaggcatctcatctcttctgtaccttgaggtaagatgtgcaagaca
cgggaactccttatacttagctgaaggggagcggagccatcatgagtct tctcgaactgcatgtaccacatgaaactatctattacaatacgctctt
ttcaaatgagatgaaccccccgcaacgacatttcggggccgaccccaac
tcagttttgaattcggtgtttataggaatctacaggcggaggtaac atgcaaagatggattgtccaagagttccgtccattatggagagaaaa
tacagaggaaagCgacctgacctcagataaagTagtgggggtatattac
atctgcagtgccttacagatctgtatcattgtcgtattgtacattga aattcctccagggtccaatcaaagcttactagatcaactagctatcaa
tttatctctgattgccatgcattctgtaaggaggggcggggtagtaat catcaaagtgttgatgcaatgggatactactttcatctactcatgaa
cttgtttgctccgtgttcacaaaaggatatattctcttaatggta tgcattgtcaggagatatggagtgttacctgggtatttgcattgggtta
cctggggcgggcctacattgtacatgaggtgggtgaggatggcGaaaac
tctggtgcagcggcacggtacgtTttgtctaaatcagatgagatcac
actgaccaggtattcacctcacagcggcagcgtgtgacagacatcct
atccagtcctttaccaagattaataaagtacttgaggaagaatattga
cactgcgctgattgaagccgggggacagcccgtccgtccattctgtgc
ggagagtctggtagcagcgtacgcaacataactcagataaccagat
Catcgctagtcacattgacacagttatccggctgtgtatatatatgga agctgagggtgatctcgctgacacagtatttctatttacccttaca
tctcttactgacgggaaaaagaggacatcacttaAacagtgcacgag
acagatcctagagggttacaataactaggtcttagagtcgaaaatctcaa taaaataggcgatataatcagcctagtgttaaaggcatgatctccat
ggaggaccttatcccactaaggacatacttgaagcatagtacctgccc taaatatttgaaggctgtcctaggtattaccaaactcaaagaaatgtt
tacagacacttctgtaCtgtacttgactcgtgtcaacaaaaattcta catgaaaactataggcaatgcagtcaaaaggatattacagtaactgtga
ctcttaacgaaaatcacatattaataggctcctttttggccaattgt attcttggtgatttaacatattatgttagaaaaaagtgaacctga
ctccttaggactcgaattcgaactcaataaatgtcttaaaaaaagggt tgcgcacaattattcttgagtgtagtctcgtcattcaccaaactcttg
ttgggt Nucleic acid ATGGAGCTGCTGATCCTGAAGGCCAACGCCATCACAC 4 sequence
of CATCCTGACCGCCGTGACCTTCTGCTTCGCCAGCGGCCA chimeric F
GAACATCACCGAGGAGTTCTACCAGAGCACCTGCAGCG sequence,
CCGTGAGCAAGGGCTACCTGAGCGCCCTGCGCACCGGC wherein the

TGGTACACCAAGTGCACCGCACCGACGCCAAGGTG protein CAAGGAGAACAAGTGCAACGGCACCGACGCCAAGGTG protein AAGCTGATCAAGCAGGAGCTGGACAAGTACAAGAACG comprises the CCGTGACCGAGCTGCAGCTGCTGATGCAGAGCACCCCC human RSV F GCCACCAACAACCGCGCCCCGCCGCGAGCTGCCCCGCTT protein CATGAATAACACCTGAACAACGCCAAGAAGACCAACG ectodomain of TGACCCTGAGCAAGAAGCGCAAGCGCCGCTTCCTGGGC the RSV A2 TTCCTGCTGGGCGTGGGCAGCGCCATCGCCAGCGGCGT strain fused to GGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTGA the ACAAGATCAAGAGCGCCCTGCTGAGCACCAACAAGGCC transmembrane GTGGTGAGCCTGAGCAACGGCGTGAGCGTGCTGACCAG and cytoplasmic CAAGGTGCTGGACCTGAAGAATAACATCGACAAGCAGC domains of the TGCTGCCCATCGTGAACAAGCAGAGCTGCAGCATCAGC NDV LaSota AACATCGCCACCGTGATCGAGTTCCAGCAGAAGAACA strain. The CCGCCTGCTGGAGATCACCCGCGAGTTCAGCGTGAACG nucleic acid CCGGCGTGACCACCCCCGTGAGCACCTACATGCTGACC sequence AACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCAT encoding the CACCAACGACCAGAAGAAGCTGATGAGCAACAACGTG ectodomain of CAGATCGTGCGCCAGCAGAGCTACAGCATCATGAGCAT the human RSV CATCAAGGAGGAGGTGCTGGCCTACGTGGTGCAGCTGC F protein of the CCCTGTACGGCGTGATCGACACCCCCTGCTGGAAGCTG RSV A2 strain is CACACCAGCCCCCTGTGCACCACCAACACCAAGGAGGG codon-optimized. CAGCAACATCTGCCTGACCCGACCGACAGGGGCTGGT (The ACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCCCAG transmembrane GCCGAGACCTGCAAGGTGCAGAGCAACCGCGTGTTCTG and cytoplasmic CGACACCATGAACAGCCTGACCCTGCCAGCGAGGTGA domains of NDV ACCTGTGCAACGTGGACATCTTCAACCCCAAGTACGAC F are TGCAAGATCATGACCAGCAAGACCGACGTGAGCAGCAG underlined.) CGTGATCACCAGCCTGGGCGCCATCGTGAGCTGCTACG GCAAGACCAAGTGCACCGCCAGCAACAAGAAGAGGGG CATCATCAAGACCTTCAGCAACGGCTGCGACTACGTGA GCAACAAGGGCGTGACACCGTGAGCGTGGGCAACAC CCTGTACTACGTGAACAAGCAGGAGGGCAAGAGCCTGT ACGTGAAGGGCGAGCCCATCATCAACTTCTACGACCCC CTGGTGTTCCCAGCGACGAGTTCGACGCCAGCATCAG CCAGGTGAACGAGAAGATCAACCAGAGCCTGGCCTTCA TCCGCAAGAGCGACGAGCTGCTGCACAACGTGAACGCC GGCAAGAGCACCATCAACGTTAACCTCATTACCTATAT CGTTTTGACTATCATATCTCTTGTGTTTTTGGTATACTTAGC CTGATTCTAGCATGCTACCTAATGTACAAGCAAAAGGC GCAACAAAAGACCTTATTATGGCTTGGGAATAATACCC TAGATCAGATGAGAGCCACTACAAAAATGTGA cDNA of accaaacagagaatccgtgagttacgataaaaggcgaaggagca 5 genomic attgaagtcgcacgggtagaaggtgtgaatctcgagtgcgagcc sequence of the cgaagcaciaaactcgagaaagccttctgccaacatgtcttccgt NDV strain atttgatgagtacgaacagctcctcgcggtcagactcgcccca LaSota genome atggagctcatggagggggagaaaaaggagtagcttaaaagta with the gacgtcccgggtattcactttaacagtgatgaccagaagatag chimeric F atggagctttgtggtattctgcctccggattgctgttagcgaag sequence (SEQ atgccaacaaaccactcaggcaaggtgctctcatatctttta ID NO:4)

tgctccactcacaggtaatgaggaacatgttgccCttgcagg inserted (The Sac
gaaacagaatgaagccacattggccgtgcttgagattgatggct II (CCGCGG)
ttgccaacggcacgccccagttcaacaataggagtggagtgtct restriction sites
gaagagagagcacagagatttgcgatgatagcaggatctctccc used to insert the
tcgggcatgcagcaacggaaccccgttcgtcacagccggggcCg chimeric F open
aagatgatgcaccagaagacatcaccgataccctggagaggatc reading frame
ctctctatccaggctcaagtatgggtcacagtagcaaaaagccat into the genomic
gactgcgtatgagactgcagatgagtcggaaacaaggcgaatca sequence are
ataagtatatgcagcaaggcaggggtccaaaagaaatacatcctc shown by double
taccccgatgcaggagcacaatccaactcacgatcagacagtc underlining. The
tcttgagtcgcgcatcttttgggttagcgagctcaagagaggcc initiation and
gcaacacggcaggtgggtacctctactattataacctggtaggg stop codons of
gacgtagactcatacatcaggaataccgggcttactgcattctt the chimeric F
cttgacactcaagtacggaatcaacaccaagacatcagcccttg open reading
cacttagtagcctctcaggcgacatccagaagatgaagcagctc frame are
atgcgtttgtatcggatgaaaggagataatgcgccgtacatgac underlined. The
attacttgggtgatagtaccagatgagctttgcgcctgccgagt sequence coding
atgcacaactttactcctttgccatgggtatggcatcagtccta for the RSV-F
gataaaggtactgggaaataccaatttgccagggactttatgag ectodomain is in
cacatcattctggagacttgagtagagtacgctcaggctcagg bold. The
gaagtagcattaacagggatatgggtgccgagctaaagctaacc sequence coding
ccagcagcaaGgaGgggcctggcagctgctgccaacgggtctc the
cgaGgaGaccagcagcataGacatgcctactcaacaagtcggag transmembrane
tcctactgggcttagcgaggggggggtcccaagctctacaaggc and cytoplasmic
ggatcgaatagatcgcaagggaaccagaagccggggatggggga domains of the
gacccaattcctggatctgatgagagcggtagcaaatagcatga NDV F protein is
gggaggcgccaaactctgcacagggcactcccaatcggggcct in italics.)
ccccaaactcctggggccatccaagataacgacaccgactgggg gtattgatggacaaaaccagcctgcttcacaaaaacatccca
atgccctcacccgtagtcgacccctcgatttgcggctctatatg accacaccctcaaaacaacatcccccttttctccctccccct
gctgtacaactAcgTacgccctagataccacaggcacaatgcgg ctactaacaatcaaaacagagccgagggaattagaaaaaagta
cgggtagaagagggatattcagagatcagggaagtcctccgag tctctgctctctctctacctgatagaccaggacaaacatggcc
acctttacagatgcagagatcgacgagctatttgagacaagtgg aactgtcattgacaacataattacagcccagggtaaaccagcag
agactgttgaaggagtgcaatcccacaaggcaagaccaagggtg ctgagcgcagcatgggagaagcatgggagcatccagccaccggc
cagtcaagacaacccgatcgacaggacagatctgacaaacaac catccacaccgagcaaacgaccccgcatgacagcccgcggcc
acatccgcccagaccagccccccaccaggccacagacgaagccgt cgacacacagCtcaggaccggagcaagcaactctctgtgtga
tgcttgacaagctcagcaataaatctccaatgctaaaaagggc ccatggctgagcccccaagaggggaatcaccaacgtccgactca
acagcaggggagtgcaaccagtcgcggaacagtcaggaaagac
cgcagaaccaagtcaaggccgcccctggaaaccaggggcacagac gtgaacacagcatatcatggacaatgggaggagtcacaactatc
agctgggtgaacccctcatgctctccgatcaaggcagagccaag acaataaccctgtatctgcggatcatgtccagccacctgtagac
tttgtgcaagcgatgatgtctatgatggaggcgatatcacagag agtaagtaagggtgactatcagctagatcttgtcttgaaacaga
catcctccatccctatgatgcggtcggaatccaacagctgaaa acatctgttgagtcagtggaagccaacttgggaatgatgaagat
tctggatcccggttgtgccaacatttcattctctgagtgatctac gggcagttgccgatctcaccgggttttagtttcaggccctgga
gacccctctccctatgtgacacaaggaggcgaaatggcacttaa taaactttcgcaaccagtgccacatccatctgaattgattaaac
ccgccactgcatgcgggcctgatataggagtggaaaaggacact gtccgtgcattgatcatgtcacgccaatgcacccgagttcttc
agccaagctcctaagcaagtttagatgcagccgggtcgatcgagg aatcaggaaaatcaagcgccttgctctaaatggctaattacta
ctgccacacgtagcgggtccctgtccactcggcatcacacggaa tctgcaccgagttccccccgcGgacccaaggtccaactctcca
agcggcaatcctctctcgcttctcagccccactgaatgAtcgc gtaaccgtaattaatctagctacatttaagattaagaaaaata
cgggtagaattggagtgcccaCtAgtgccCA**ATGGAGCTGCT**
GATCCTGAAGGCCAACGCCATCACCACCATCCTGACCGCCGTGA
CCTTCTGCTTCGCCAGCGGCCAGAACATCACCGAGGAGTTCTAC

CAGAGACACCTGACGACGCGCGCTGAGCAAGGCGCTACCTGAGCGCCCT
 GCGCACCGGCTGGTACACCAGCGTGATCACCATCGAGCTGAGCA
 ACATCAAGGAGAACAAGTGCAACGGCACCGACGCCAAGGTGAAG
 CTGATCAAGCAGGAGCTGGACAAGTACAAGAACGCCGTGACCGA
 GCTGCAGCTGCTGATGCAGAGCACCCCCGCCACCAACAACCGCG
 CCCGCCGCGAGCTGCCCCGCTTCATGAACTACACCCTGAACAAC
 GCCAAGAAGACCAACGTGACCCTGAGCAAGAAGCGCAAGCGCCG
 CTTCTTGGGCTTCCTGCTGGGCGTGGGCAGCGCCATCGCCAGCG
 GCGTGGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTGAAC
 AAGATCAAGAGCGCCCTGCTGAGCACCAACAAGGCCGTGGTGAG
 CCTGAGCAACGGCGTGAGCGTGCTGACCAGCAAGGTGCTGGACC
 TGAAGAACTACATCGACAAGCAGCTGCTGCCCCATCGTGAACAAG
 CAGAGCTGCAGCATCAGCAACATCGCCACCGTGATCGAGTTCCA
 GCAGAAGAACAACCGCCTGCTGGAGATCACCCGCGAGTTCAGCG
 TGAACGCCGGCGTGACCACCCCCGTGAGCACCTACATGCTGACC
 AACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCATCACCA
 CGACCAGAAGAAGCTGATGAGCAACAACGTGCAGATCGTGCGCC
 AGCAGAGCTACAGCATCATGAGCATCATCAAGGAGGAGGTGCTG
 GCCTACGTGGTGCAGCTGCCCCCTGTACGGCGTGATCGACACCCC
 CTGCTGGAAGCTGCACACCAGCCCCCTGTGCACCACCAACACCA
 AGGAGGGCAGCAACATCTGCCTGACCCGACCCGACaGgGGCTGG
 TACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCCCAGGCCGA
 GACCTGCAAGGTGCAGAGCAACCGCGTGTTCTGCGACACCATGA
 ACAGCCTGACCCTGCCCAGCGAGGTGAACCTGTGCAACGTGGAC
 ATCTTCAACCCCAAGTACGACTGCAAGATCATGACCAGCAAGAC
 CGACGTGAGCAGCAGCGTGATCACCAGCCTGGGCGCCATCGTGA
 GCTGCTACGGCAAGACCAAGTGACCCGCCAGCAACAAGAAaGg
 GGCATCATCAAGACCTTCAGCAACGGCTGCGACTACGTGAGCAA
 CAAGGGCGTGAGACACCGTGAGCGTGGGCAACACCCTGTACTACG
 TGAACAAGCAGGAGGGCAAGAGCCTGTACGTGAAGGGCGAGCCC
 ATCATCAACTTCTACGACCCCCCTGGTGTTCCCCAGCGACGAGTT
 CGACGCCAGCATCAGCCAGGTGAACGAGAAGATCAACCAGAGCC
 TGGCCTTCATCCGCAAGAGCGACGAGCTGCTGCACAACGTGAAC
 GCCGGCAAGAGCACCATCAACGTTAACCTCATTACCTATATCGT
 TTTGACTATCATATCTCTTGTGTTTTGGTATACTTAGCCTGATTC
 TAGCATGCTACCTAATGTACAAGCAAAAGGCGCAACAAAAGACC
 TTATTATGGCTTGGGAATAATACCCTAGATCAGATGAGAGCCAC
 TACAAAATGTGAaccgcgacccaaggtccaactctccaagcgg
 caatcctctctcgcttcctcagccccactgaatgAtcgcgtaac cgtaatatctagctacatttaagattaagaaaaatacgggt
 agaattggagtgcccaattgtgccaagatggactcatctagga caattgggctgtactttgattctgccattcttctagcaacctg
 ttagcatttccgatcgtcctacaagAcacaggagatgggaagaa gcaaatcgccccgcaatataggatccagcgccttgacttgtga
 ctgatagtaaggaggactcagtattcatcaccacctatggattc atctttcaagttgggaatgaagaagccacCgtcggcatgatcga
 tgataaacccaagcgcgagttactttccgctgcgatgctctgcc taggaagcgtcccaaataccggagaccttattgagctggcaagg
 gcctgtctcactatgatagtcacatgcaagaagagtgcaactaa tactgagagaatggttttctcagtagtgagggcaccccaagtg
 tgcaaaagctgtaggggtgtggcaaaacaatactcatcagtgaat gcagtcaagcacgtgaaagcgccagagaagattcccgggagtg
 aaccctagaatacaaggtgaactttgtctccttgactgtggtac cgaagaGggatgtctacaagatcccagctgcagtattgaaggt
 tctggctcgagtgctgtacaatcttgcgctcaatgtcactattaa tgtggaggtagaccgaggagtcctttggttaaatctCtgtcta
 agtctgacagcggatactatgctaacctcttcttgcatttga cttatgaccacTgtagataggaaggggaagaaagtacattga
 caagctggaaaagaaaataaggagccttgatctatctgtcgggc tcagtgatgtgctcgggcctccgtgttggtaaaagcaagaggt
 gcacggactaagcttttggcacctttctctctagcagtgggac agcctgctatcccatagcaaatgcttctcctcaggtggccaaga

tactctggagtcacaaaccgcgtgcctgcggagcgttaaaaatcatt atccaagcagggtacccaacgcgctgtcgcagtgaccgccgacca
cgagggtaccttactaagctggagaaggggacaccccttgcca aatacaatcctttaagaaataagctgcgtctctgagattgcgc
tccgcccactcaccagatcatcatgacacaaaaaactaatctg tcttgattatttacagtttagtttacctgtctatcaagttagaaa
aaacacgggtagaagattctggatcccgggtggcgccctccagg tgcaagatgggctccagaccttctaccaagaacccagcacctat
gatgctgactatccgggtgcgctggtagtgcattctgtc cggcaaacctcattgatggcaggcctcttgacgctgcaggaatt
gtgggttacaggagacaaagccgtcaacatatacacctcatcca gacaggatcaatcatagttaagctcctcccgaatctgcccaagg
ataaggaggcatgtgcgaaagcccccttgatgcatacaacagg acattgaccactttgtcaccccccttggtgactctatccgtag
gatacaagagtctgtgactacatctggaggggggagacaggggc gccttataggcgccattattggcggtgtggctctgggggtgca
actgccgcacaaataacagcggccgcagctctgatacaagccaa acaaaatgctgccaacatcctccgacttaaagagagcattgccg
caaccaatgaggctgtgcatgaggctcactgacggattatcgcaa ctagcagtggcagttgggaagatgcagcagtttgtaatgacca
atttaataaaacagctcaggaattagactgcatcaaaattgcac agcaagttgggtgtagagctcaacctgtacctaaccgaattgact
acagtattcggaccacaaatcattcacctgctttaaacaagct gactattcaggcactttacaatctagctgggtggaaatatggatt
acttattgactaagtttaggtgtaggggaacaatcaactcagctca ttaatcggtagcggcttaatcaccggtaacctattctatacga
ctcacagactcaactcttgggtatacaggtaactctaccttcag tcgggaacctaaataatatgcgtgccacactacttgaaacctta
tccgtaagcacaccaggggatttgcctcggcacttgtcccAaa agtgggtgacacaggctcggttctgtgatagaagaacttgacacct
catactgtatagaaactgacttagatttatattgtacaagaata gtaacgttcctatgtccctgggtatttattctgcttgagcgg
caatacgtcggcctgtatgtactcaaagaccgaaggcgcaactta ctacaccatacatgactatcaaagggtcagtcacgccaaactgc
aagatgacaacatgtagatgtgtaaaccctccgggtatcatatc gcaaaactatggagaagccgtgtctctaatagataaacaatcat
gcaatgtttatccttaggcgggataactttaaggctcagtggg gaattcgatgtaacttatcagaagaatatctcaatacaagattc
tcaagtaataataacaggcaatcttgatatctcaactgagcttg ggaatgtcaacaactcgatcagtaatgcttgtaataagttagag
gaaagcaacagaaaactagacaaagtcaatgtcaaactgactag cacatctgctctcattacctatatcgtttgactatcatatctc
ttgttttgggtatacttagcctgattctagcatgctacctaata tacaagcaaaaggcgcaacaaaagaccttattatggcttgggaa
taatactctagatcagatgagagccactacaaaatgtgaacac agatgaggaacgaaggtttcctaataagtaattgtgtgaaagt
tctggtagtctgtcagttcagagagttaagaaaaaactaccggt ttagatgaccaaaaggacgatatacgggtagaacggtaagagag
gccgcccctcaattgcgagccaggcttcacaacctccgttctac cgcttcaccgacaacagtcctcaatcatggaccgcgccgttagc
caagttgcgttagagaatgatgaaagagaggcaaaaaatacatg gcgcttgatattccggattgcaatcttattcttaacagtagtga
ccttggctatatctgtagcctcccttttatatagcatgggggct agcacacctagcgatctttaggcataccgactaggatttcag
ggcagaagaaaagattacatctacacttgggtccaatcaagatg tagtagataggatatataagcaagtgcccttgagtctccgttg
gcattgttaaatactgagaccacaattatgaacgcaataacatc tctctttatcagattaatggagctgcaaaacagtggggtggg
gggcacctatccatgaccagattatatagggggggataggcaaa gaactcattgtagatgatgctagtgtatgtcacatcattctatcc
ctctgcatttcaagaacatctgaatttatcccggcgcctacta caggatcagggtgcactcgaataccctcatttgacatgagtgt
accattactgtacaccataatgtaattgtctggatgcag agatcactcacattcatatcagttatttagcacttgggtgtgtcc
ggacatctgcaacaggagggtattcttttctactctgcgttcc atcaacctggacgacacccaaaatcggaagtcttgacgtgtgag
tgcaactcccctgggttgatgtgtgtcgtcgaaagtcacgg agacagagggaagaagattataactcagctgtccctacgcggatg
gtacatgggaggttaggggtcgacggccagtagcacgaaaagga cctagatgcacaacattattcggggactgggtggccaactacc
caggagtaggggggtggatcttttattgacagccgcgtatgggtc tcagtctacggagggttaaaacccaattcacccagtgacactgt
acaggaagggaatatgtgatatacaagcgatacaatgacacat gcccagatgagcaagactaccagattcgaatggccaagtcttcg
tataagcctggacggttgggtgggaaacgcatacagcaggctat ctatctatcaaggtgtcaacatccttaggcgaagaccgggtac
tgactgtaccgccaacacagtcacactcatggggggccgaaggc agaattctcacagtagggacatctcatttctgtatcaacgagg
gtcatcatacttctctcccgcgttattatctctatgacagtca gcaacaaaacagccactcttcatagtccttatacattcaatgcc
ttcactcggccaggtagtatcccttgccaggcttcagcaagatg ccccaactcgtgtgttactggagtctatacagatccatatcccc
taatcttctatagaaaccacaccttgcgaggggtattcggggaca atgcttgatgggtgacaagcaagacttaaccctgcgtctgcagt
attcgatagcacatcccgcagtcgcattactcgagtgaagtcaa gcagtaccaaagcagcatacacaacatcaacttgttttaaagtg
gtcaagactaataagacctattgtctcagcattgctgaaatc taatactctcttcggagaattcagaatcgtcccgttactagttg
agatcctcaaagatgacggggtagagaagccagggtctggctag ttgagtcaattataaaggagttggaaagatggcattgtatcacc
tatcttctgcgacatcaagaatcaaaccgaatgccgggcgctgc tcgaattccatgttgccagttgaccacaatcagccagtgctcat
gcgatcagattaagccttgtcaAtaGtctcttgattaagaaaaa atgtaagtggcaatgagatacaaggcaaaacagctcatgggTaa
Caatacgggtaggacatggcgagctccggctctgaaagggcaga gcatcagattatcctaccagagTcacacctgtcttcaccattgg
tcaagcacaactactctattactggaaattaactgggctaccg cttctgtatgaatgtgacttcgaccacctcattctcagccgaca
atggaaaaaaatacttgaatcggcctctctgatactgagagaa tgataaaactcggaagggcagtagacacaaactttaaccacaat

tcagaataaccggagtgctccacccaggtgttttagaaGaact ggctaattagggtccagattcaaccaaaatttcggaaga
ttgagaagaagatccaaattcacacacgagatatggagaactg ttcacaaggctgtgtacgcatatagagaagaaactgctggggtc
atcttggtctaacaatgtcccccggtcagaggagttcagcagca ttcgtacggatccggcattctggtttcactcaaaatgggccaca
gccaaagttgcatggctccatataaaacagatccagaggcatct gatggtggcagctaGgacaaggctctgcggccaacaaattggtga
tgctaaccataaggtaggccaagtctttgtcactcctgaactt gtcgttgtgacgcatacgaatgagaacaagttcacatgtcttac
ccaggaacttgattgatgtatgcagatatgatggagggcgagag atatggtcaacataatatcaaccacggcggtgcatctcagaagc
ttatcagagaaaaattgatgacattttgcggttaatagacgctct ggcaaaagacttgggtaataagttacgatgtttgatcactaa
tggaggggattgcatacggagctgtccagctactcgagccgtca ggtacatttgaggagatttcttcgattcaacctgcaggagct
taaagacattctaattggcctcctccccaatgatatagcagaat ccgtgactcatgcaatcgctactgtattctctggtttagaacag
aatcaagcagctgagatgttgtgtctgttgcgtctgtggggtca cccactgcttgagctccgtattgcagcaaaggcagtcaggagcc
aaatgtgcgcaccgaaaaatggtagactttgatatgatccttcag gtactgtcttttcaagggaacaatcatcaacgggtacagaaa
gaagaatgcagggtgtgtggccgcgagtcaaagtggatacaatat atgggaaggctattgggcaactacatgcagattcagcagagatt
tcacacgatatcatgttgagagagtataagagtttatctgcact tgaatttgagccatgtatagaatatgacctgtcaccaacctga
gcatgttcctaaaagacaaggcaatcgcacaccccaacgataat tggcttgcctcgtttaggcggaaccttctctccgaagaccagaa
gaaacatgtaaaagaagcaacttcgactaatcgctcttgatag agtttttagagtcaaataatttgatccatataaagagatggaa
tatctgacgaccttgagtaccttagagatgacaatgtggcagt atcatactcgctcaaggagaaggaaagtgaaggttaattggacgga
tcttcgctaagctgacaaagaagttaaggaactgtcaggtgatg gcggaagggtacctagccgatcagattgcacctttcttcaggg
aaatggagtcattcagगतatcatatccttgaccaagagtatgc tagcgatgagtcactgtctttaacagcaataagaaacgtatc
actgactgtaaaagaaagagtatcttcaaaccgcaatcatgatcc gaaaagcaagaaccgtcggagagttgcaaccttcataacaactg
acctgcaaaagtactgtcttaattggagatatcagacaatcaaa ttgttcgctcatgccatcaatcagttgatgggcctacctcactt
cttcgaatggattcacctaagactgatggacactacgatgttcg tagggagacctttcaatcctccaagtacacctactgactgtgac
cttcaagagtcctaattgatgacatatatattgtcagtgccag aggggggtatcgaaggattatgccagaagctatggacaatgatct
caattgctgcaatccaacttgctgcagctagatcgattgtcgt gttgcctgtatggtacaggggtgataatcaagtaatagcagtaac
gagagaggtgaagatcagacgactctccggagatggtgttgacac agttgcatcaagccagtataatttctcaaggaattaattcat
gtcaatcatttgattggccataatttgaaggatcgtgaaacat caggtcagacacatttctcatatacagcaaacgaatcttcaaag
atggagcaatcctcagtcgaagtcctcaaaaattcatctaaatta gtgtagtgtcaggtgatctcagtgaaaacaccgtaattgtcctg
tgccaacattgcctctactgtagcacggctatgcgagaacgggc ttcccaaagacttctgttactatttaaactatataatgagttgt
gtgcagacatactttgactctgagtttccatcaccaacaattc gcaccccgatcttaatcagtcgtggattgaggacatctcttttg
tgcactcatatgttctgactcctgccaattaggggggactgagt aaccttcaataactcaaggctctacactagaaatatcggtgacct
ggggactactgcttttgagagatcaagcgactagaagcagtggt gattactgagtcctaacattatgactaataatcttaactaggccg
cctgggaatggagattgggccaagtctgtgcaacgacctatactc tttcaattttgagactgttgcaagcccaaatattgttctaaga
aacatacgcaaagagtcctatttgaaacttggtcaaatccctta ttgtctggagtgcacacagaggataatgaggcagaagagaaggc
attggctgaattctgtctaatacaagaggtgattcatccccgcg ttgcgatgccatcatggaggcaagctctgtaggtaggagaaag
caaattcaagggtgttgacacaacaaacaccgtaattaagat tgcgcttactaggaggccattaggcatcaagaggctgatcgga
tagtcaattattctagcatgcatgcaatgctgtttagagacgat gtttttctccagtagatccaaccaccccttagtctcttctaa
tatgtgttctctgacactggcagactatgcacggaatagaagct ggtcacctttgacgggaggcaggaaaatactgggtgtatcta
cctgatacगतagaactcगतaggggtgagattcttagtgtaag cggaggggtgtacaagatgtgacagcggagatgaacaatttactt
ggttccatcttcaaagcaatatagaattgaccgatgacaccagc aagaatcctccgatgaggggtaccatatctcggtcaaagacaca
ggagaggagagctgcctcacttgcaaaaatagctcatatgtcgc cacatgtaaaggctgccctaagggtcatcatccgtgttgatctgg
gcttatggggataatgaagtaaattggactgctgctttacgat tgcaaaatctcggtgtaattgtaaacttagagtatcttcggttac
tgtcccccttaccacgggtgggaatttcaacatagactagat gatggtataactcagatgacattcacccctgcatctctctacag
gGtgtcaccttacattcacatatccaatgattctcaaaggctgt tcaactgaagaaggagtcaaagaggggaatgtggtttaccaacag
atcatgctcttgggtttatcttaatcgaatcgatctttccaat gacaacaaccaggacatatgatgagatcacactgcacctacata
gtaaatttagttgctgtatcagagaagcacctgttgcggttctc ttcgagctacttgggggtggtaccggaactgaggacagtgcctc
aaataagtttatgatgatccttagccctgatcgaggggagact ttgcgagacttgacttagctatcttcaagagttagagctta
ctggagtcatatcccacगतatagactaatgaacattcttcaat atccagcgggaagttgattggccagtctgtggtttcttatgatg
aagatacctccataaagaatgacgccataatagtgtatgacaat acccgaaattggatcagtgaagctcagaattcagatgtggtccg
cctatttgaatatgcagcacttgaagtgtcctcgactgttctt accaactctattacctgagagtaagaggcctGgacaatattgtc
ttatatatgggtgatttatacaagaatatgccaggaattctact ttccaacattgcagctacaatatctcatcccgtcattcattcaa
gggtacatgcagtgggcctgggtcaacatgacggatcacacca cttgcagatacggattttatcgaaatgtctgcaaaactattagt

atcttgacccgacgtgtgatctccggcttatattcaggaaata agtatgatctgctgttcccatctgtcttagatgataacctgaat
gagaagatgcttcagctgatatcccggttatgtgtctgtacac ggtactctttgctacaacaagagaaatccccgaaaataagaggct
taactgcagaagagaaatgttcaatactcactgagatttactg tcggatgctgtgaaaccattacttagccccgatcaagtgagctc
tatcatgtctcctaacataattacattcccagctaattctgtact acatgtctcggaagagcctcaatttgatcagggaaagggaggac
agggatactatcctggcgtgtgttcccccaagagccattatt agagttcccttctgtgcaagatattgggtgctcagagtgaagatc
cattcacccgacaacctgcggcatttttgcaagagttagatttg agtgctccagcaaggtatgacgcattcacacttagtcagattca
tcctgaactcacatctccaaatccggaggaagactacttagtac gatacttggtcagagggatagggactgcatcttctcttggtat
aaggcatctcatctcttctgtaccgaggtgaagatgtgcaag acacggggaactcctatacttagctgaagggagcggagccatca
tgagtcttctcgaactgcatgtaccacatgaaactatctattac aatacgctcttttcaaagatgaaccccccgcaacgacattt
cgggccgaccccaactcagttttgaattcgggtgtttatagga atctacaggcggaggtaacatgcaaagatggatttgccaagag
ttccgtccattatggagagaaaatacagaggaaagCgacctgac ctcagataaagTagtgggggtatattacatctgcagtgccctaca
gatctgtatcattgctgcattgtgacattgaaattcctccaggg tccaatcaaagcttactagatcaactagctatcaatttatctt
gattgccatgcattctgtaagggagggcggggtagtaatcatca aagtgttgatgcaatgggatactactttcatctactcatgaac
ttgtttgctccgtgttcacaaaaggatatattctctctaattgg ttatgcatgtcgaggagatatggagtgttacctggattttgtca
tgggttacctgggcgggcctacatttgatcatgaggtggtgagg atggcGaaaactctggtgcagcggcacgggtacgctTttgtctaa
atcagatgagatcacactgaccaggttattcacctcacagcggc agcgtgtgacagacatcctatccagtcctttaccaagattaata
aagtacttgaggaagaatattgacactgcgctgattgaagccgg gggacagcccgtccgtccattctgtgcggagagctggtgagca
cgctagcgaacataactcagataaccagatCatcgctagtcac attgacacagttatccggctgtgatatatatggaagctgaggg
tgatctcgctgacacagattttctatttacccttacaatctct ctactgacgggaaaaagaggacatcacttaAacagtgcacgaga
cagatcctagaggttacaatactaggtcttagagtcgaaaatct caataaaataggcgatataatcagcctagtgtctaaaggcatga
tctccatggaggaccttatcccactaaggacatacttgaagcat agtacctgccctaaatatttgaaggctgtcctaggtattaccaa
actcaaagaaatgtttacagacactctgtaCtgtacttgactc gtgtcaacaaaaattctacatgaaaactataggcaatgcagtc
aaaggatattacagtaactgtgactcttaacgaaaatcacatat taataggctcctttttggccaattgtattctgttgattta
catattatgtagaaaaaagttgaaccctgactccttaggactc gaattcgaactcaaataaatgtcttaaaaaaagggtgcgcacaa
ttattcttgagtgtagtctcgtcattcaccaaatctttttgg t Amino acid

MELLILKANAITTILTAVTFCFASGQNITEEFYQSTCSAVSK 6 sequence of
GYLSALRTGWYTSVITIELSNIKENKCNGTDAKVKLIKQEL human RSV F
DKYKNAVTELQLLMQSTPATNNRARRELPRFMNYTLNNA protein of A2
KKTNVTLSSKKRKRRLGFLGVSIAISGVAVSKVLHLEG strain (The
EVNKIKSALLSTNKAVVSLSNGVSVLTSKVLDLKNYIDKQ transmembrane
LLPIV NKQSCSISNIATVIEFQQKNNRLLEITREFSVNAGVT and cytoplasmic
TPVSTYMLTNSELLSLINDMPITNDQKKLMSNNVQIVRQQ domains of the
SYSIMSIIKEEVLAYVVQLPLYGVIDTPCWKLHTSPLCTTN human RSV F
TKEGSNICLTRTDRGWYCDNAGSVSFFPQAETCKVQSNRV protein are
FCDTMNSLTLPSEVNL CNVDIFNP KYDCKIMTSKTDVSSSV underlined. The
ITSLGAIVSCYGKTKCTASNKNRGIKTF SNGCDYVS NKG V amino acid
DTVSVGNTLYYV NKQEGKSLYVKGEPIINFYDPLVFP SDEF sequence of the
DASISQVNEKINQSLAFIRKSDELLHNVNAGKSTINIMITTII RSV F protein
IVIIIVILLSLIAVGLLLYCKARSTPVTLSKDQLSGINNIAFSN encoded by the wild-type
nucleic acid sequence for the human RSV F protein (SEQ ID NO: 1) is identical
to the amino acid sequence of the human RSV F protein encoded by the codon-
optimized nucleic acid sequence for human RSV F protein. (SEQ ID NO: 2))
Amino acid MELLILKANAITTILTAVTFCFASGQNITEEFYQSTCSAVSK 7 sequence of
GYLSALRTGWYTSVITIELSNIKENKCNGTDAKVKLIKQEL chimeric F
DKYKNAVTELQLLMQSTPATNNRARRELPRFMNYTLNNA protein encoded
KKTNVTLSSKKRKRRLGFLGVSIAISGVAVSKVLHLEG by SEQ ID NO:
EVNKIKSALLSTNKAVVSLSNGVSVLTSKVLDLKNYIDKQ 4 (The
LLPIV NKQSCSISNIATVIEFQQKNNRLLEITREFSVNAGVT transmembrane
TPVSTYMLTNSELLSLINDMPITNDQKKLMSNNVQIVRQQ and cytoplasmic
SYSIMSIIKEEVLAYVVQLPLYGVIDTPCWKLHTSPLCTTN domains are

TKEGSNICLRTCDNAGSVSFPPQAE^{TC}TKVQSNRV underlined.)
FCDTMNSLTLPSEVNLCNVDIFNPKYDCKIMTSKTDVSSSV
ITSLGAIVSCYGKTKCTASNKNRGIKTF^SNGCDYVSNKGV
DTVSVGNTLYYV^NKQEGKSLYVKGEPIINFYDPLVFP^SDEF
DASISQVNEKINQSLAFIRKSDELLHNVNAGKSTINVNLITY
IVLTIISLVFGILSLILACYLMYKQKAQ^QKTLLWLGNNTLD QMRATTKM Wild-type
ATGGAGTTGCCAATCCTCAAAACAAATGCTATTACCAC 25 nucleic acid
AATCCTTGCTGCAGTCACACTCTGTTTTGCTTCCAGTCA sequence
AAACATCACTGAAGAATTTTATCAATCAACATGCAGTG encoding human
CAGTTAGCAAAGGCTATCTTAGTGCTCTAAGAACTGGTT RSV F protein
GGTATACTAGTGTTATAACTATAGAATTAAGTAATATCA from strain
AGGAAAATAAGTGTAATGGTACAGACGCTAAGGTAAA RSVA/Homo
ATTAATAAAACAAGAATTAGATAAATATAAAAATGCTG sapiens/USA/T
TAACAGAATTGCAGTTGCTCATGCAAAGCACACCAGCA H_10656/2014
GCCAACAGTCGAGCCAGAAGAGAACTACCAAGATTTAT (transmembrane
GAATTATACACTCAACAATACCAAAAACACCAATGTAA and cytoplasmic
CATTAAGTAAGAAAAGGAAAAGAAGATTTCTTGGATTT domains are
TTGTTAGGTGTTGGATCTGCAATCGCCAGTGGCATTGCC underlined)
GTATCCAAGGTCCTGCACCTAGAAGGGGAAGTGAACAA
AATCAAAAGTGCTCTACTATCCACAAACAAGGCTGTAG
TCAGCTTATCTAATGGAGTCAGTGTCTTAACCAGCAAG
GTGTTAGACCTCAAAAAC^TATATAGATAAACAGTTGTT
ACCTATTGTTAACAAGCAAAGCTGCAGCATATCAAACA
TTGAAACTGTGATAGAGTTCCAACAAAAGAACAACAGA
CTACTAGAGATTACCAGAGAATTTAGTGTTAATGCAGG
TGTA^ACTACACCTGTAAGCACTTATATGTTAACTAATAG
TGAGTTATTATCATTAATCAATGATATGCCTATAACAAA
TGATCAGAAAAAGTTAATGTCCAGCAATGTTCAAATAG
TTAGACAGCAAAGTTACTCTATCATGTCAATAATAAAA
GAGGAAGTCTTAGCATATGTAGTACAATTACCACTATA
TGGTGTAATAGATACTCCTTGTTGGAACTACACACATC
CCCTCTATGTACAACCAACACAAAGGAAGGATCCAACA
TCTGCTTAACAAGAACCGACAGAGGATGGTACTGTGAC
AATGCAGGATCAGTATCCTTTTTCCCACAAGCTGAAAC
ATGTAAAGTTCAATCGAATCGGGTGTTTTGTGACACAAT
GAACAGTTTAAACATTACCAAGTGAGGTAAATCTCTGCA
ACATTGACATATTCAACCCCAAATATGATTGCAAAATT
ATGACTTCAAAAACAGATGTAAGCAGCTCCGTTATCAC
ATCTCTAGGAGCCATTGTGTCATGCTATGGCAAAACCA
AATGTACAGCATCCAATAAAAATCGTGGGATCATAAAG
ACATTCTCTAACGGGTGTGATTATGTATCAAATAAGGG
GGTGGATACTGTGTCTGTAGGTAATACATTATATTATGT
AAATAAGCAAGAAGGCAAAAGTCTCTATGTAAAAGGTG
AACCAATAATAAATTTCTATGATCCATTAGTGTTCCCCT
CTGATGAATTTGATGCATCAATATCTCAAGTCAATGAG
AAAATTAATCAGAGTCTAGCATTTATCCGTAAATCAGA
TGAATTATTACATAATGTAAATGCTGGTAAATCCACCAC
AAATATCATGATAACTACCATAATTATAGTAATTATAGT
AATATTGTTAGCATTAATTGCAGTTGGACTGCTTCTATA
CTGCAAGGCCAGAAGCACACCAGTCACATTAAGTAAGG

ATCAACTGAGTGGTATAAATAACATTGCATTTAGTAACT GA Codon
ATGGAGCTGCCCCATCCTGAAGACCAACGCCATCACCCAC 26 optimized
CATCCTGGCCGCCGTGACCCTGTGCTTCGCCAGCAGCCA nucleic acid
GAACATCACCGAGGAGTTCTACCAGAGCACCTGCAGCG sequence
CCGTGAGCAAGGGCTACCTGAGCGCCCTGAGAACCGGC encoding human
TGGTACACCAGCGTGATCACCATCGAGCTGAGCAACAT RSV F protein
CAAGGAGAACAAGTGCAACGGCACCGACGCCAAGGTG from strain
AAGCTGATCAAGCAGGAGCTGGACAAGTACAAGAACG RSVA/Homo
CCGTGACCGAGCTGCAGCTGCTGATGCAGAGCACCCCC sapiens/USA/T
GCCGCCAACAGCAGAGCCAGAAGAGAGCTGCCCAGATT H_10656/2014
CATGAACTACACCCTGAACAACACCAAGAACACCAACG
TGACCCTGAGCAAGAAGAGAAAGAGAAGATTCCTGGG
CTTCCTGCTGGGCGTGGGCAGCGCCATCGCCAGCGGCA
TCGCCGTGAGCAAGGTGCTGCACCTGGAGGGGCGAGGTG
AACAAGATCAAGAGCGCCCTGCTGAGCACCAACAAGGC
CGTGGTGAGCCTGAGCAACGGCGTGAGCGTGCTGACCA
GCAAGGTGCTGGACCTGAAGAACTACATCGACAAGCAG
CTGCTGCCCCATCGTGAACAAGCAGAGCTGCAGCATCAG
CAACATCGAGACCGTGATCGAGTTCCAGCAGAAGAACA
ACAGACTGCTGGAGATCACCCAGAGAGTTTCAGCGTGAAC
GCCGGCGTGACCACCCCCGTGAGCACCTACATGCTGAC
CAACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCA
TCACCAACGACCAGAAGAAGCTGATGAGCAGCAACGTG
CAGATCGTGAGACAGCAGAGCTACAGCATCATGAGCAT
CATCAAGGAGGAGGTGCTGGCCTACGTGGTGCAGCTGC
CCCTGTACGGCGTGATCGACACCCCCTGCTGGAAGCTG
CACACCAGCCCCCTGTGCACCACCAACACCAAGGAGGG
CAGCAACATCTGCCTGACCAGAACCGACAGAGGCTGGT
ACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCCCAG
GCCGAGACCTGCAAGGTGCAGAGCAACAGAGTGTTCTG
CGACACCATGAACAGCCTGACCCTGCCCAGCGAGGTGA
ACCTGTGCAACATCGACATCTTCAACCCCAAGTACGAC
TGCAAGATCATGACCAGCAAGACCGACGTGAGCAGCAG
CGTGATCACCGCCTGGGCGCCATCGTGAGCTGCTACG
GCAAGACCAAGTGCACCGCCAGCAACAAGAACAGAGG
CATCATCAAGACCTTCAGCAACGGCTGCGACTACGTGA
GCAACAAGGGCGTGACACCGTGAGCGTGGGCAACAC
CCTGTACTACGTGAACAAGCAGGAGGGCAAGAGCCTGT
ACGTGAAGGGCGAGCCCATCATCAACTTCTACGACCCC
CTGGTGTTCCCCAGCGACGAGTTCGACGCCAGCATCAG
CCAGGTGAACGAGAAGATCAACCAGAGCCTGGCCTTCA
TCAGAAAGAGCGACGAGCTGCTGCACAACGTGAACGCC
GGCAAGAGCACCACCAACATCATGATCACCAACCATCAT
CATCGTGATCATCGTGATCCTGCTGGCCCTGATCGCCGT
GGGCCTGCTGCTGTACTGCAAGGCCAGAAGCACCCCCG
TGACCCTGAGCAAGGACCAGCTGAGCGGCATCAACAAC ATCGCCTTCAGCAAC
Amino acid MELPILKTNAITTLAAVTLCFASSQNITEEFYQSTCSAVSK 50 sequence of
GYLSALRTGWYTSVITIELSNIKENKCNGTDAKVKLIKQEL human RSV F
DKYKNAVTELQLLMQSTPAANSRRARRELPRFMNYTLNNT protein from the
KNTNVTLSSKKRKRRLGFLLGVSIAISGIAVSKVLHLEG strain

EVNKIKSLNSKNGVSLNSGVSLVLSKVLDSLKNYIDKQ RSVV/Homo
LLPIV NKQSCSISNIETVIEFQQKNNRLLEITREFSVNAGVTT sapiens/USA/T
PVSTYMLTNSELLSLINDMPITNDQKKLMSSNVQIVRQQS H_10656/2014
YSIMSIIEEVLAYVVQLPLYGVIDTPCWKLHTSPLCTTNT
KEGSNICLTRTRDRGWYCDNAGSVSFFPQAETCKVQSNRVF
CDTMNSLTLPSEVNL CNIDIFNPKYDCKIMTSKTDVSSSVIT
SLGAIVSCYGKTKCTASNKNRGIKTF SNGCDYVSNKGVD
TVSVGNTLYYV NKQEGKSLYVKGEPIINFYDPLVFPSDEFD
ASISQVNEKINQSLAFIRKSDELLHNVNAGKSTTNIMITTH
VIIVILLALIAVGLLLYCKARSTPVTLSKDQLSGINNIAFSN Wild-type
ATGGAGTTGCTGATCCATAGATCAAGTGTAATCTTCCTA 27 nucleic acid
ACTCTTGCTATTAACGCATTGTACCTCACCTCAAGTCAG sequence
AACATAACTGAGGAGTTTTACCAATCGACATGTAGTGC encoding human
AGTTAGCAGAGGTTACTTTAGTGCTTTAAGAACAGGTT RSV F protein
GGTATACTAGTGTCATAACAATAGAATTAAGTAATATA from strain
AAAGAAACCAAATGCAATGGAAGTACACTAAAGTAA RSVB/Homo
AACTTATAAAACAAGAATTAGATAAGTATAAGAATGCA sapiens/USA/L
GTAACAGAATTACAGTTACTTATGCAAAACATACCAGC A2_82/2013
TGCCAACAACCGAGCCAGAAGAGAAGCACACAGTAT (transmembrane
ATGAACTACACAATCAATACCACTAAAAACCTAAATGT and cytoplasmic
ATCAATAAGCAAGAAGAGGAAACGAAGATTTCTGGGCT domains are
TCTTGTTAGGTGTAGGATCTGCAATAGCAAGTGGCATA underlined)
GCTGTATCCAAAGTTCTACACCTTGAAGGAGAAGTGAA
CAAGATCAAAAATGCTTTGCTGTCTACAAACAAAGCTG
TAGTCAGTCTATCAAATGGGGTCAGTGTTTTAACCAGCA
AAGTGTTAGATCTCAAGAATTATATAAACAACCAATTA
TTACCTATAGTAAATCAACAGAGTTGTGCGCATCTCCAAC
ATTGAAACAGTTATAGAATTCCAGCAGAGAAGAACAGCAG
ATTGTTGGAAATCACCAGAGAATTTAGTGTCATGCAG
GTGTAACGACACCTTTAAGCACTTACATGTTAACAAAC
AGTGAGTTGCTATCATTAAATCAATGATATGCCTATAACA
AATGATCAGAAGAAATTAATGTCAAGCAATGTTTCAGAT
AGTAAGGCAACAAAGTTATTCTATCATGTCTATAATAA
AGGAAGAAGTCCTCGCATATGTTGTACAGCTACCTATCT
ATGGTGTAATAGATACACCTTGCTGGAAATTACACACA
TCACCTCTGTGCACCACCAACATCAAGGAAGGTTCAAA
TATTTGTTTAACAAGGACTGATAGAGGATGGTACTGTG
ATAATGCAGGATCAGTATCCTTCTTTCCACAGGCTGACA
CTTGTAAGTACAGTCCAATCGAGTATTTTGTGACACTA
TGAACAGTTTGACATTACCAAGTGAAGTCAGCCTTTGTA
ACACTGACATATTCAATTCCAAGTATGATTGCAAAATTA
TGACATCAAAAACAGACATAAGCAGCTCAGTAATTACT
TCTCTAGGAGCTATAGTGTCATGCTATGGTAAACTAA
ATGCACTGCATCCAACAAAATCGTGGAATTATAAAGA
CATTTTCTAATGGTTGTGATTATGTGTCAAACAAAGGAG
TAGATACTGTATCAGTGGGCAACACTTTATACTATGTCA
ACAAGCTGGAAGGCAAAAACCTTTATGTAAAAGGGGA
ACCTATAATAAATTACTATGACCCTCTAGTGTTTCCTTC
TGATGAGTTTGATGCATCAATATCTCAAGTCAATGAAA
AAATTAATCAAAGTTTAGCTTTTATTTCGTAGATCCGATG

AATTATTATGATAACTGTAATTATTATAGTAATCATTGTA
GTATTGTTATCATTAATAGCTATTGGTTTACTGTTGTATT
GCAAAGCCAAAAACACACCAGTTACACTAAGCAAAGA
CCAATAAGTGGAATCAATAATATTGCATTCAGCAAAT AG Codon
ATGGAGCTGCTGATCCACAGAAGCAGCGTGATCTTCCT 28 optimized
GACCCTGGCCATCAACGCCCTGTACCTGACCAGCAGCC nucleic acid
AGAACATCACCGAGGAGTTCTACCAGAGCACCTGCAGC sequence
GCCGTGAGCAGAGGCTACTTCAGCGCCCTGAGAACCGG encoding human
CTGGTACACCAGCGTGATCACCATCGAGCTGAGCAACA RSV F protein
TCAAGGAGACCAAGTGCAACGGCACCGACACCAAGGT from strain
GAAGCTGATCAAGCAGGAGCTGGACAAGTACAAGAAC RSVB/Homo
GCCGTGACCGAGCTGCAGCTGCTGATGCAGAACATCCC sapiens/USA/L
CGCCGCCAACAACAGAGCCAGAAGAGAGGCCCCCCCAG A2_82/2013
TACATGAACTACACCATCAACACCACCAAGAACCTGAA
CGTGAGCATCAGCAAGAAGAGAAAGAGAAGATTCTTG
GGCTTCCTGCTGGGCGTGGGCAGCGCCATCGCCAGCGG
CATCGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGG
TGAACAAGATCAAGAACGCCCTGCTGAGCACCAACAAG
GCCGTGGTGAGCCTGAGCAACGGCGTGAGCGTGCTGAC
CAGCAAGGTGCTGGACCTGAAGAACTACATCAACAACC
AGCTGCTGCCCATCGTGAACCAGCAGAGCTGCAGAATC
AGCAACATCGAGACCGTGATCGAGTTCCAGCAGAAGAA
CAGCAGACTGCTGGAGATCACCAGAGAGTTCAGCGTGA
ACGCCGGCGTGACCACCCCCCTGAGCACCTACATGCTG
ACCAACAGCGAGCTGCTGAGCCTGATCAACGACATGCC
CATCACCACGACCAGAAGAAGCTGATGAGCAGCAAC
GTGCAGATCGTGAGACAGCAGAGCTACAGCATCATGAG
CATCATCAAGGAGGAGGTGCTGGCCTACGTGGTGCAGC
TGCCCATCTACGGCGTGATCGACACCCCCCTGCTGGAAG
CTGCACACCAGCCCCCTGTGCACCACCAACATCAAGGA
GGGCAGCAACATCTGCCTGACCAGAACCGACAGAGGCT
GGTACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCC
CAGGCCGACACCTGCAAGGTGCAGAGCAACAGAGTGTT
CTGCGACACCATGAACAGCCTGACCCTGCCCAGCGAGG
TGAGCCTGTGCAACACCGACATCTTCAACAGCAAGTAC
GACTGCAAGATCATGACCAGCAAGACCGACATCAGCAG
CAGCGTGATCACCAGCCTGGGCGCCATCGTGAGCTGCT
ACGGCAAGACCAAGTGCACCGCCAGCAACAAGAACAG
AGGCATCATCAAGACCTTCAGCAACGGCTGCGACTACG
TGAGCAACAAGGGCGTGACACCGTGAGCGTGGGCAA
CACCTGTACTACGTGAACAAGCTGGAGGGCAAGAACC
TGTACGTGAAGGGCGAGCCCATCATCAACTACTACGAC
CCCCTGGTGTTCCCCAGCGACGAGTTCGACGCCAGCAT
CAGCCAGGTGAACGAGAAGATCAACCAGAGCCTGGCCT
TCATCAGAAGAAGCGACGAGCTGCTGCACAACGTGAAC
ACCGGCAAGAGCACCAACACATCATGATCACCGTGAT
CATCATCGTGATCATCGTGGTGCTGCTGAGCCTGATCGC
CATCGGCCTGCTGCTGTACTGCAAGGCCAAGAACCCC
CCGTGACCCTGAGCAAGGACCAGCTGAGCGGCATCAAC AACATCGCCTTCAGCAAG

Amino acid MELLISSVIFLTLAINALYLTSSQNTIEFYQSTCSAVSR 58 sequence of
RSV GYFSALRTGWYTSVITIELSNIKETKCNGTDTKVKLIKQEL F protein from
DKYKNAVTELQLLMQNIPAANNRARRREAPQYMNYTINTT strain
KNLNVSISKKRKRRLGFLGVGSAIASGIAVSKVLHLEGE RSVB/Homo
VNKIKNALLSTNKAVVSLNNGVSVLTSKVLDLKNYINNQL sapiens/USA/L
LPIVNQQSCRISNIETVIEFQQKNSRLLEITREFSVNAGVTTP A2_82/2013
LSTYMLTNSELLSLINDMPITNDQKKLMSSNVQIVRQQSYS
IMSIIKEEVLAYVVQLPIYGVIDTPCWKLHTSPLCTTNIKEG
SNICLTRTRDRGWYCDNAGSVSFFPQADTCKVQSNRVFCDT
MNSLTLPSEVSLCNTDIFNSKYDCKIMTSKTDISSSVITSLG
AIVSCYGKTKCTASNKNRGIKTFNNGCDYVSNKGVDTVS
VGNTLYYVKNLEGKNLYVKGEPIINYYDPLVFPSDEFDASI
SQVNEKINQSLAFIRRSDELLHNVNTGKSTTNIMITVIIIIVII
VVLLSLIAIGLLLYCKAKNTPVTL SKDQLSGINNIAFSK Wild-type
ATGGAGTTGCCAATCCTCAAAGCAAATGCTATTACCAC 29 nucleic acid
AATCCTTGCTGCAGTCACACTCTGTTTTGCTTCCAGTCA sequence
AAACATCACTGAAGAATTTTATCAATCAACATGCAGTG encoding human
CAGTTAGCAAAGGCTATCTTAGTGCTCTAAGAACTGGTT RSV F protein
GGTATACTAGTGTTATAACTATAGAATTAAGTAATATCA from strain
AGGAAAATAAGTGTAATGGAACAGATGCTAAGGTAAA HRSV-A-GZ08-
ATTGATAAAACAAGAATTAGATAAATACAAAAATGCTG 18
TAACAGAATTGCAGTTGCTCATGCAAAGCACACCAGCA (transmembrane
GCCAACAATCGAGCCAGAAGGGAATTACCAAGATTTAT and cytoplasmic
GAATTATACACTCAACAATACTGAAAACACCAATGTAA domains are
CATTAAGCAAGAAAAGGAAAAGAAGATTTCTTGGCTTT underlined)
TTGTTAGGTGTTGGATCTGCAATCGCCAGTGGCATTGCC
GTATCCAAGGTCCTGCACCTAGAAGGGGAAGTGAACAA
AATCAAAAAGTGCTCTACTATCCACAAACAAGGCTGTAG
TCAGCTTATCAAATGGAGTCAGTGTCTTAACCAGCAAA
GTGTTAGATCTCAAAAACCTATATAGATAAACAGTTGTT
ACCTATTGTGAACAAGCAAAGCTGCAGCATATCAAACA
TTGAAACTGTAATAGAGTTCCAACAAAAGAACAACAGA
CTACTAGAGATTACCAGAGAATTTAGTGTTAATGCAGG
TGTA ACTACACCTGTAAGCACTTATATGTTAACTAATAG
TGAATTATTATCATTAAATCAATGATATGCCTATAACAAA
TGATCAGAAAAAGTTAATGTCCAACAATGTTCAAATAG
TTAGACAGCAAAGTTACTCTATCATGTCCATAATAAAA
GAGGAAGTCTTAGCATATGTAGTACAATTACCACTATA
TGGTGTAATAGATACACCTTGTTGGAAACTGCACACAT
CCCCTCTATGTACAACCAACACAAAGGAAGGGTCCAAC
ATCTGCTTAACAAGAACCGACAGAGGATGGTACTGTGA
CAATGCAGGATCAGTATCTTTCTTCCCACAAGCCGAAA
CATGTAAAGTTCAATCGAATCGGGTATTTTGTGACACA
ATGAACAGTTTAACATTACCAAGTGAGGTAAAGTCTCTG
CAACGTTGACATATTCAACCCCAAATATGATTGCAAAA
TTATGACTTCAAAAACAGATGTAAGCAGCTCCGTTATC
ACATCTCTGGGAGCCATTGTGTCATGCTATGGCAAAAC
CAAATGTACAGCATCCAATAAAAATCGTGGGATCATAA
AGACATTTTCTAACGGGTGTGATTATGTATCAAATAAG
GGGGTGGATACTGTGTCTGTAGGTAATACATTATATTAT

GTAAATAAGCAAGCAAAAATCTCTATGTAAAAGG
TGAACCAATAATAAATTTCTATGACCCATTAGTGTTCCC
CTCTGATGAATTTGATGCATCAATATCTCAAGTCAATGA
GAAGATTAACCAGAGTCTAGCATTATTCGTAAATCAG
ATGAATTATTACATAATGTAAATGCTGTTAAATCCACTA
CAAATATCATGATAACTACTATAATTATAGTGATTATAG
TAATATTGTTATCATTAATTGTAGTTGGACTGCTTCTAT
ACTGCAAGGCCAGAAGCACACCAGTCACACTAAGTAAG
GATCAACTGAGTGGTATAAATAATATTGCATTTAGTAG CTGA Codon
ATGGAGCTGCCCATCCTGAAGGCCAACGCCATCACCCAC 30 optimized
CATCCTGGCCGCCGTGACCCTGTGCTTCGCCAGCAGCCA nucleic acid
GAACATCACCGAGGAGTTCTACCAGAGCACCTGCAGCG sequence
CCGTGAGCAAGGGCTACCTGAGCGCCCTGAGAACCGGC encoding human
TGGTACACCAGCGTGATCACCATCGAGCTGAGCAACAT RSV F protein
CAAGGAGAACAAGTGCAACGGCACCGACGCCAAGGTG from RSV strain
AAGCTGATCAAGCAGGAGCTGGACAAGTACAAGAACG HRSV-A-GZ08-
CCGTGACCGAGCTGCAGCTGCTGATGCAGAGCACCCCC 18
GCCGCCAACAACAGAGCCAGAAGAGAGCTGCCCAGATT
CATGAACTACACCCTGAACAACACCGAGAACACCAACG
TGACCCTGAGCAAGAAGAGAAAGAGAAGATTCCTGGG
CTTCCTGCTGGGCGTGGGCAGCGCCATCGCCAGCGGCA
TCGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTG
AACAAGATCAAGAGCGCCCTGCTGAGCACCAACAAGGC
CGTGGTGAGCCTGAGCAACGGCGTGAGCGTGCTGACCA
GCAAGGTGCTGGACCTGAAGAACTACATCGACAAGCAG
CTGCTGCCCATCGTGAACAAGCAGAGCTGCAGCATCAG
CAACATCGAGACCGTGATCGAGTTCCAGCAGAAGAACA
ACAGACTGCTGGAGATCACCGAGAGAGTTCAGCGTGAAC
GCCGGCGTGACCACCCCCGTGAGCACCTACATGCTGAC
CAACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCA
TCACCAACGACCAGAAGAAGCTGATGAGCAACAACGTG
CAGATCGTGAGACAGCAGAGCTACAGCATCATGAGCAT
CATCAAGGAGGAGGTGCTGGCCTACGTGGTGCAGCTGC
CCCTGTACGGCGTGATCGACACCCCCTGCTGGAAGCTG
CACACCAGCCCCCTGTGCACCACCAACACCAAGGAGGG
CAGCAACATCTGCCTGACCAGAACCGACAGAGGCTGGT
ACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCCCAG
GCCGAGACCTGCAAGGTGCAGAGCAACAGAGTGTTCTG
CGACACCATGAACAGCCTGACCCTGCCAGCGAGGTGA
GCCTGTGCAACGTGGACATCTTCAACCCCAAGTACGAC
TGCAAGATCATGACCAGCAAGACCGACGTGAGCAGCAG
CGTGATCACCGCCTGGGCGCCATCGTGAGCTGCTACG
GCAAGACCAAGTGCACCGCCAGCAACAAGAACAGAGG
CATCATCAAGACCTTCAGCAACGGCTGCGACTACGTGA
GCAACAAGGGCGTGACACCGTGAGCGTGGGCAACAC
CCTGTACTACGTGAACAAGCAGGAGGGCAAGAACCTGT
ACGTGAAGGGCGAGCCCATCATCAACTTCTACGACCCC
CTGGTGTTCCCCAGCGACGAGTTTCGACGCCAGCATCAG
CCAGGTGAACGAGAAGATCAACCAGAGCCTGGCCTTCA
TCAGAAAGAGCGACGAGCTGCTGCACAACGTGAACGCC

GTGAAGACACCAACATCATGATCACCACCATCAT
CATCGTGATCATCGTGATCCTGCTGAGCCTGATCGTGGT
GGGCCTGCTGCTGTACTGCAAGGCCAGAAGCACCCCCG
TGACCCTGAGCAAGGACCAGCTGAGCGGCATCAACAAC ATCGCCTTCAGCAGC

Amino acid

MELPILKANAITTILAAVTLCFASSQNITEEFYQSTCSAVSKGYLSALRTGWYTSVI 49

sequence of

TIELSNIKENKCNGTDAKVKLIKQELDKYKNAVTELQLLMQSTPAANNRAREL human
RSV F PRFMNYTLNNTENTNVTLSSKKRKRRLGFLLGVGSIAIASGIAVSKVLHLEGEVN
protein from

KIKSALLSTNKAVVSLSNGVSVLTSKVLDLKNYIDKQLLPIVNBKQSCSISNIETVIEF strain
HRSV-A- QQKNNRLLEITREFSVNAGVTTTPVSTYMLTNSELLSLINDMPITNDQKKLMSN
GZ08-18 NVQIVRQQSYSIMSIIKEEVLAYVVQLPLYGVIDTPCWKLHTSPLCTTNTKEGSNI
CLTRTDRGWYCDNAGSVSFFPQAETCKVQSNRVFCDTMNSLTLPSVSLCNV
DIFNPKYDCKIMTSKTDVSSSVITSLGAIVSCYGKTKCTASNKNRGIKTFSSNGCD
YVSNKGVDTVSVGNTLYYVNBKQEGKNLYVKGEPIINFYDPLVFPSEDFDASISQV
NEKINQSLAFIRKSDELLHNVNAVKSTTNIMITTHIIVIVILLSLIVVGLLLYCKARST

PVTLSKDQLSGINNIAFSS Nucleic acid

ATGGAGCTGCCCATCCTGAAGACCAACGCCATCACCACCATCCTGGCCGCC 44 sequence
GTGACCCTGTGCTTCGCCAGCAGCCAGAACATCACCGAGGAGTTCTACCAG encoding
AGCACCTGCAGCGCCGTGAGCAAGGGCTACCTGAGCGCCCTGAGAACCGG chimeric F
CTGGTACACCAGCGTGATCACCATCGAGCTGAGCAACATCAAGGAGAACAA protein,
wherein GTGCAACGGCACCGACGCCAAGGTGAAGCTGATCAAGCAGGAGCTGGACA
the nucleic acid

AGTACAAGAACGCCGTGACCGAGCTGCAGCTGCTGATGCAGAGCACCCCC sequence
GCCGCCAACAGCAGAGCCAGAAGAGAGCTGCCCAGATTCATGAACTACAC comprises
a CCTGAACAACACCAAGAACACCAACGTGACCCTGAGCAAGAAGAGAAAGA codon
optimized GAAGATTCCTGGGCTTCCTGCTGGGCGTGGGCAGCGCCATCGCCAGCGGC
nucleic acid

ATCGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTGAACAAGATCAA sequence
GAGCGCCCTGCTGAGCACCAACAAGGCCGTGGTGAGCCTGAGCAACGGCG encoding
the TGAGCGTGCTGACCAGCAAGGTGCTGGACCTGAAGAACTACATCGACAAG human
RSV F CAGCTGCTGCCCATCGTGAACAAGCAGAGCTGCAGCATCAGCAACATCGAG
protein ACCGTGATCGAGTTCCAGCAGAAGAACAACAGACTGCTGGAGATCACCAG
ectodomain of

AGAGTTCAGCGTGAAACGCCGGCGTGACCACCCCGTGAGCACCTACATGCT strain
GACCAACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCATCACCAACGA
RSVA/Homo

CCAGAAGAAGCTGATGAGCAGCAACGTGCAGATCGTGAGACAGCAGAGCT
sapiens/USA/TH_

ACAGCATCATGAGCATCATCAAGGAGGAGGTGCTGGCCTACGTGGTGAG
10656/2014 and

CTGCCCTGTACGGCGTGATCGACACCCCCTGCTGGAAGCTGCACACCAGC a
nucleic acid

CCCCTGTGCACCACCAACACCAAGGAGGGCAGCAACATCTGCCTGACCAGA sequence
ACCGACAGAGGCTGGTACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCC encoding
the CAGGCCGAGACCTGCAAGGTGCAGAGCAACAGAGTGTTCTGCGACACCAT
transmembrane

GAACAGCCTGACCCTGCCCAGCGAGGTGAACCTGTGCAACATCGACATCTT and
cytoplasmic CAACCCCAAGTACGACTGCAAGATCATGACCAGCAAGACCGACGTGAGCA

domains of F
GCAGCGTGATCACCAGCCTGGGCGCCATCGTGAGCTGCTACGGCAAGACC protein
from AAGTGCACCGCCAGCAACAAGAACAGAGGCATCATCAAGACCTTCAGCAAC
NDV strain GGCTGCGACTACGTGAGCAACAAGGGCGTGGACACCGTGAGCGTGGGCA
LaSota ACACCCTGTACTACGTGAACAAGCAGGAGGGCAAGAGCCTGTACGTGAAG
(transmembrane
GGCGAGCCCATCATCAACTTCTACGACCCCCTGGTGTTCCTCCAGCGACGAG and
cytoplasmic TTCGACGCCAGCATCAGCCAGGTGAACGAGAAGATCAACCAGAGCCTGGC
domains of NDV
CTTCATCAGAAAGAGCGACGAGCTGCTGCACAACGTGAACGCCGGCAAGA F are
underlined) GCACCACCAACATCATGg^{tt}aacCTCATTACCTATATCGTTTTGACTATCATAT
CTCTTGTTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCTAATGTACAAG
CAAAAGGCGCAACAAAAGACCTTATTATGGCTTGGGAATAATACCCTAGAT
CAGATGAGAGCCACTACAAAAATGTGAaccg^{cg} Nucleic acid
ATGGAGCTGCTGATCCACAGAAGCAGCGTGATCTTCCTGACCCTGGCCATC 45 sequence
AACGCCCTGTACCTGACCAGCAGCCAGAACATCACCGAGGAGTTCTACCAG encoding
AGCACCTGCAGCGCCGTGAGCAGAGGCTACTTCAGCGCCCTGAGAACCGG chimeric F
CTGGTACACCAGCGTGATCACCATCGAGCTGAGCAACATCAAGGAGACCAA protein,
wherein GTGCAACGGCACCGACACCAAGGTGAAGCTGATCAAGCAGGAGCTGGACA
the nucleic acid
AGTACAAGAACGCCGTGACCGAGCTGCAGCTGCTGATGCAGAACATCCCCG sequence
CCGCCAACAACAGAGCCAGAAGAGAGGGCCCCCAGTACATGAACTACACC comprises
a ATCAACACCACCAAGAACCTGAACGTGAGCATCAGCAAGAAGAGAAAGAG codon
optimized AAGATTCCTGGGCTTCCTGCTGGGCGTGGGCAGCGCCATCGCCAGCGGCAT
nucleic acid
CGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTGAACAAGATCAAGA sequence
ACGCCCTGCTGAGCACCAACAAGGCCGTGGTGAGCCTGAGCAACGGCGTG encoding
the AGCGTGCTGACCAGCAAGGTGCTGGACCTGAAGAACTACATCAACAACCA human
RSV F GCTGCTGCCCATCGTGAACCAGCAGAGCTGCAGAATCAGCAACATCGAGAC
protein CGTGATCGAGTTCCAGCAGAAGAACAGCAGACTGCTGGAGATCACCAGAG
ectodomain of
AGTTCAGCGTGAACGCCGGCGTGACCACCCCCCTGAGCACCTACATGCTGA strain
CCAACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCATCACCAACGACC
RSVB/Homo
AGAAGAAGCTGATGAGCAGCAACGTGCAGATCGTGAGACAGCAGAGCTAC
sapiens/USA/LA2_
AGCATCATGAGCATCATCAAGGAGGAGGTGCTGGCCTACGTGGTGCAGCT 82/2013
and a GCCCATCTACGGCGTGATCGACACCCCCTGCTGGAAGCTGCACACCAGCCC
nucleic acid
CCTGTGCACCACCAACATCAAGGAGGGCAGCAACATCTGCCTGACCAGAAC sequence
CGACAGAGGCTGGTACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCCCA encoding
the GGCCGACACCTGCAAGGTGCAGAGCAACAGAGTGTTCTGCGACACCATGA
transmembrane
ACAGCCTGACCCTGCCCAGCGAGGTGAGCCTGTGCAACACCGACATCTTCA and
cytoplasmic ACAGCAAGTACGACTGCAAGATCATGACCAGCAAGACCGACATCAGCAGC
domains of F
AGCGTGATCACCAAGCCTGGGCGCCATCGTGAGCTGCTACGGCAAGACCAA protein
from GTGCACCGCCAGCAACAAGAACAGAGGCATCATCAAGACCTTCAGCAACG
NDV strain
GCTGCGACTACGTGAGCAACAAGGGCGTGGACACCGTGAGCGTGGGCAAC LaSota

ACCCTGTACTACGTGAACAAGCTGGAGGGCAAGAACCTGTACGTGAAGGG
 (transmembrane
 CGAGCCCATCATCAACTACTACGACCCCCTGGTGTTCCTCCAGCGACGAGTTC and
 cytoplasmic GACGCCAGCATCAGCCAGGTGAACGAGAAGATCAACCAGAGCCTGGCCTT
 domains of NDV
 CATCAGAAGAAGCGACGAGCTGCTGCACAACGTGAACACCGGCAAGAGCA F are
 underlined) CCACCAACATCATGg^{tt}aaacCTCATTACCTATATCGTTTTGACTATCATATCTCT
TGTTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCTAATGTACAAGCAA
AAGGCGCAACAAAAGACCTTATTATGGCTTGGGAATAATACCCTAGATCAG
ATGAGAGCCACTACAAAAATGTGAccg^{cg}g Nucleic acid
 ATGGAGCTGCCCATCCTGAAGGCCAACGCCATCACCAACATCCTGGCCGCC 46 sequence
 GTGACCCTGTGCTTCGCCAGCAGCCAGAACATCACCGAGGAGTTCTACCAG encoding
 AGCACCTGCAGCGCCGTGAGCAAGGGCTACCTGAGCGCCCTGAGAACCGG chimeric F
 CTGGTACACCAGCGTGATCACCATCGAGCTGAGCAACATCAAGGAGAACAA protein,
 wherein GTGCAACGGCACCGACGCCAAGGTGAAGCTGATCAAGCAGGAGCTGGACA
 the nucleic acid
 AGTACAAGAACGCCGTGACCGAGCTGCAGCTGCTGATGCAGAGCACCCCC sequence
 GCCGCCAACAAACAGAGCCAGAAGAGAGCTGCCCAGATTCATGAACTACACC
 comprises a
 CTGAACAACACCGAGAACACCAACGTGACCCTGAGCAAGAAGAGAAAGAG codon
 optimized AAGATTCCTGGGCTTCCTGCTGGGCGTGGGCAGCGCCATCGCCAGCGGCAT
 nucleic acid
 CGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTGAACAAGATCAAGA sequence
 GCGCCCTGCTGAGCACCAACAAGGCCGTGGTGAGCCTGAGCAACGGCGTG encoding
 the AGCGTGCTGACCAGCAAGGTGCTGGACCTGAAGAACTACATCGACAAGCA human
 RSV F GCTGCTGCCCATCGTGAACAAGCAGAGCTGCAGCATCAGCAACATCGAGAC
 protein CGTGATCGAGTTCCAGCAGAAGAACAACAGACTGCTGGAGATCACCAGAG
 ectodomain of
 AGTTCAGCGTGAACGCCGGCGTGACCACCCCCGTGAGCACCTACATGCTGA strain
 HRSV-A- CCAACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCATCACCAACGACC
 GZ08-18 and a
 AGAAGAAGCTGATGAGCAACAACGTGCAGATCGTGAGACAGCAGAGCTAC nucleic
 acid AGCATCATGAGCATCATCAAGGAGGAGGTGCTGGCCTACGTGGTGAGCT sequence
 GCCCCTGTACGGCGTGATCGACACCCCCTGCTGGAAGCTGCACACCAGCCC encoding
 the CCTGTGCACCACCAACACCAAGGAGGGCAGCAACATCTGCCTGACCAGAAC
 transmembrane
 CGACAGAGGCTGGTACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCCCA and
 cytoplasmic GGCCGAGACCTGCAAGGTGCAGAGCAACAGAGTGTCTGCGACACCATGA
 domains of F
 ACAGCCTGACCCTGCCCAGCGAGGTGAGCCTGTGCAACGTGGACATCTTCA protein
 from ACCCCAAGTACGACTGCAAGATCATGACCAGCAAGACCGACGTGAGCAGC NDV
 strain AGCGTGATCACAGCCTGGGCGCCATCGTGAGCTGCTACGGCAAGACCAA LaSota
 GTGCACCGCCAGCAACAAGAACAGAGGCATCATCAAGACCTTCAGCAACG
 (transmembrane
 GCTGCGACTACGTGAGCAACAAGGGCGTGAGACACCGTGAGCGTGGGCAAC and
 cytoplasmic ACCCTGTACTACGTGAACAAGCAGGAGGGCAAGAACCTGTACGTGAAGGG
 domains of NDV
 CGAGCCCATCATCAACTTCTACGACCCCCTGGTGTTCCTCCAGCGACGAGTTC F are
 underlined) GACGCCAGCATCAGCCAGGTGAACGAGAAGATCAACCAGAGCCTGGCCTT
 CATCAGAAAGAGCGACGAGCTGCTGCACAACGTGAACGCCGTGAAGAGCA

CCACCAATCATGgttaacCTCATATACCTATATCGTTTTGACTATCATATCTCT
TGTTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCTAATGTACAAGCAA
AAGGCGCAACAAAAGACCTTATTATGGCTTGGGAATAATACCCTAGATCAG
ATGAGAGCCACTACAAAAATGTGAAccgcgg Nucleic acid

atggagttgctaactcctcaaagcaaatgcaattaccacaatcctcactgcagtcacattttgtttgct 51 sequence
tctggtcaaaacatcactgaagaattttatcaatcaacatgcagtgtagcaaaggctatctta encoding a
gtgctctgagaactggttggtataccagtggtataactatagaattaagtaatatcaaggaaaataa chimeric F
gtgtaatggaacagatgctaaggtaaaattgataaaacaagaattagataaatataaaaatgctgt protein, wherein
aacagaattgcagttgctcatgcaaagcacaccagcaacaaacaatcgagccagaagagaacta the chimeric F
ccaaggtttatgaattatacactcaacaatgccaaaaaaccaatgtaacattaagcaagaaaag protein
gaaaagaagatttcttggtttttgttaggtgttgatctgcaatcgccagtggtgctgtatctaa comprises the
ggctctgcacctagaaggggaagtgaacaagatcaaaagtgtcttactatccacaacaaggctg human RSV F
tagtcagcttatcaaatggagttagtgcttaaccagcaaagtgttagacctcaaaaactatataga protein
taaacaattgttacctattgtgaacaagcaaagctgcagcatatcaaatatagcaactgtgataga ectodomain of
gttccaacaaaagaacaacagactactagagattaccagggaatttagtgtaatgcaggtgtaac RSV A2 strain
tacacctgtaagcacttacatgtaactaatgtgaattattgtcattaatcaatgatatgcctataac and the
aaatgatcagaaaaagttaatgtccaacaatgttcaaatagttagacagcaaagttactctatcatg transmembrane
tcataataaaaagaggaagtcttagcatatgtagtacaattaccactatatggtgttatagatacac and cytoplasmic
cctgttggaactacacacatcccctctatgtacaaccaacacaaaagaagggtccaacatctgttt domains of
aacaagaactgacagaggatggtactgtgacaatgcaggatcagtatctttctcccacaagctga NDV F protein
aacatgtaaagttaaatcaaatcgagtattttgtgacacaatgaacagtttaacattaccaagtga of NDV LaSota
gtaaatctctgcaatgttgacatattcaaccccaaatatgattgtaaaattatgacttcaaaaacag strain
atgtaagcagctccgttatcacatctctaggagccattgtgtcatgctatggcaaaactaaatgtac
agcatccaataaaaatcgtggaatcataaagacattttctaacgggtgcgattatgtatcaaataaa
ggggtggacactgtgtctgtaggtaacacattatattgtaaataagcaagaaggtaaaagtctct
atgtaaaagggtgaaccaataataaatttctatgaccattagattcccctctgatgaatttgatgca
tcaatatctcaagtcaacgagaagattaaccagagcctagcatttattcgtaaatccgatgaattatt
acataatgtaaatgctggtaaatccaccataaatgttaacCTCATTACCTATATCGTTTTGA
CTATCATATCTCTTGTTTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCTA
ATGTACAAGCAAAAAGGCGCAACAAAAGACCTTATTATGGCTTGGGAATAAT
ACCCTAGATCAGATGAGAGCCACTACAAAAATGTGA

(186) TABLE-US-00002 TABLE 2 NDV LaSota F protein Amino acid
MGSRPSTKNPAPMTLTIRVALVLSCICPANSIDGRPLAAAG 8 sequence of F
IVVTGDKAVNIYTSSQTGSIIVKLLPNLPKDKEACAKAPLD protein of NDV
AYNRTLTLTLLTPLGDSIRRIQESVTTSGGGRQGRLIGAIIGG strain LaSota
VALGVATAAQITAAAALIQAKQNAANILRLKESIAATNEA (transmembrane
VHEVTDGLSQLAVAVGKMQQFVNDQFNKTAQELDCIKIA domain is
QQVGVELNLYLTELTTFVFGPQITSPALNKLTIQALYNLAG underlined and
GNMDYLLTKLGVGNNQLSSLIGSGLITGNPILYDSQTQLLG cytoplasmic
IQVTLPVGNLNNMRATYLETLVSTTRGFASALVPKVVT domain is in
QVGSVIEELDTSYCIETDLDLYCTRIVTFPMSPGIYSCLSGN bold)
TSACMYSKTEGALTTPYMTIKGSVIANCKMTTCRCVNPPG
IISQNYGEAVSLIDKQSCNVLSLGGITLRLSGEFDVITYQKNI
SIQDSQVIITGNLDISTELGNVNNSISNALNKLEESNRKLDK
VNVKLTSTSAILITYIVLTIISLVFGILSLILACYL**MYKQKAQ**
QKTLLWLGNNTLDQMRATTKM

(187) TABLE-US-00003 TABLE 3 Bovine RSV F Sequences Wild-type
ATGGCGACAACAGCCATGAGGATGATCATCAGCATTATCTTCATCTCT 9 Nucleic acid
ACCTATGTGACACATATCACTTTATGCCAAAACATAACAGAAGAATTT sequence
TATCAATCAACATGCAGTGCAGTTAGTAGAGGTTACCTTAGTGCATTA encoding bovine

AGAAACTGATGGATGGATATACAAAGTGTGGTGAACATAGATTGAGTGGACGAGTGGATATA RSV F protein
CAAAAAAATGTGTGTAATAGTACTGATTCAAAAGTGAAATTAATAAAG of bovine RSV
CAAGAACTAGAAAGATACAACAATGCAGTAGTGGAATTGCAGTCACTT strain
ATGCAAAATGAACCGGCCTCCTTCAGTAGAGCAAAAAGAGGGATACCA ATCC51908
GAGTTGATACATTATACAAGAACTCTACAAAAAAGTTTTATGGGCTA (transmembrane
ATGGGCAAGAAGAGAAAAAGGAGATTTTTAGGATTCTTGCTAGGTATT and cytoplasmic
GGATCTGCTATTGCAAGTGGTGTAGCAGTGTCCAAAGTACTACACCTG domains
GAGGGAGAGGTGAATAAAATTA AAAATGCACTGCTATCCACAAATAAA underlined)
GCAGTAGTTAGTCTATCCAATGGAGTTAGTGTCTTACTAGCAAAGTA
CTTGATCTAAAGAACTATATAGACAAAGAGCTTCTACCTAAAGTTAAC
AATCATGATTGTAGGATATCCAAAATAGAACTGTGATAGAATTCCAA
CAAAAAACAATAGATTGTTAGAAATTGCTAGGGAATTTAGTGTAAT
GCTGGTATTACCACACCTCTCAGTACATACATGTTGACCAATAGTGAA
TTACTATCACTAATTAATGATATGCCTATAACGAATGACCAAAAAAAG
CTAATGTCAAGTAATGTTCAAATAGTCAGGCAACAGAGTTATTCCATT
ATGTCAGTGGTCAAAGAAGAAGTCATAGCTTATGTTGTACAATTGCCT
ATTTATGGAGTTATAGACACCCCCTGTTGGAACTACACACCTCTCCG
TTATGCACCACTGATAATAAAGAAGGGTCAAACATCTGCTTAACTAGG
ACAGATCGTGGGTGGTATTGTGACAATGCAGGCTCTGTGTCTTTTTTC
CCACAGACAGAGACATGTAAGGTACAATCAAATAGAGTGTTCTGTGAC
ACAATGAACAGTTTAACTCTGCCTACTGACGTTAACTTATGCAACACT
GACATATTCAATACAAAGTATGACTGTAAAATAATGACATCTAAAAT
GACATAAGTAGCTCTGTGATAACTTCAATTGGAGCTATTGTATCATGC
TATGGGAAGACAAAATGTACAGCTTCTAATAAAAATCGTGGAATCATA
AAGACTTTTTCCAATGGGTGTGATTATGTATCAAACAAAGGAGTAGAT
ACTGTATCTGTTGGTAACACACTATATTATGTAAATAAGCTAGAGGGG
AAAGCACTCTATATAAAGGGTGAACCAATTATTAATTACTATGATCCA
CTAGTGTTTCCTTCTGATGAGTTTGATGCATCAATTGCCCAAGTAAAC
GCAAAAATAAACC AAAGCCTGGCCTTCATACGTCGATCTGATGAGTTA
CTTCACAGTG TAGATGTAGGAAAATCCACCACAAATGTAGTAATTACT
ACTATTATCATAGTGATAGTTGTAGTGATATTAATGTTAATAGCTGTA
GGATTACTGTTTTACTGTAAGACCAAGAGTACTCCTATCATGTTAGGG
AAGGATCAGCTCAGTGGTATCAACAATCTTTCCTTTAGTAAATGA Amino acid
MATTAMRMIISIIIFISTYVTHITLCQNITEEFYQSTCSAVSRGY 10 sequence of
LSALRTGWYTSVVTIELSKIQKNVCNSTDSKVKLIKQELERYNN bovine RSV F
AVVELQSLMQNEPASFSRAKRGIPELIHYTRNSTKKFYGLMGKK protein of
RKRRFLGFLLGIGSAIASGVAVSKVLHLEGEV NKIKNALLSTNK bovine RSV
AVVSLSNGVSVLT SKVLDLKNYIDKELLPKVNNHDCRISKIETV strain
IEFQQKNNRLL EIAREFSVNAGITPLSTYMLTNS ELLSLINDM ATCC51908
PITNDQKKLMSSNVQIVRQQSYSIMSVVKEEVIAYVVQLPIYGV (transmembrane
IDTPCWKLHTSPLCTTDNKEGSNICLTRTDRGWYCDNAGSVSFF and cytoplasmic
PQTETCKVQSNRVFCDTMNSLTLPD VNLCNTDIFNTKYDCKIM domains
TSKTDISSSVITSIGAIVSCYGKTKCTASNKNRGIKTESNGCD underlined. The
YVSNKGVDTVSVGNTLYYV NKLE GKALYIKGEPIINYYDPLVFP amino acid
SDEFDASIAQVNAKINQSLAFIRRSDELLHSVDVGKSTTNVVIT sequence
TIIIVIVVVILMLIAVGLLFYCKTKSTPIMLGKDQLSGINNLSE encoded by SEQ SK ID
NO: 9) Codon ATGGCCACCACCGCCATGCGCATGATCATCAGCATCATCTTCAT 11
optimized CAGCACCTACGTGACCCACATCACCTGTGCCAGAACATCACCG nucleic acid
AGGAGTTCTACCAGAGCACCTGCAGCGCCGTGAGTCGCGGCTAC sequence
CTGAGCGCCCTGCGCACCGGCTGGTACACCAGCGTGGTGACCAT encoding bovine

CGAGCTGAGCTAGGACCAAGATCCAGACGAGACGTGTGCAACAGCAGCA RSV F protein
GCAAGGTGAAGCTGATCAAGCAGGAGCTGGAGCGCTACAACAAC of bovine RSV
GCCGTGGTGGAGCTGCAGAGCCTGATGCAGAACGAGCCCCGCCAG strain
CTTCAGCCGCGCCAAGCGCGGCATCCCCGAGCTGATCCACTACA ATCC51908
CCCGCAACAGCACCAAGAAGTTCTACGGCCTGATGGGCAAGAAG
CGCAAGCGCCGCTTCCTGGGCTTCCTGCTGGGCATCGGCAGCGC
CATCGCCAGCGGCGTGCCCGTGAGCAAGGTGCTGCACCTGGAGG
GCGAGGTGAACAAGATCAAGAACGCCCTGCTGAGCACCAACAAG
GCCGTGGTGGAGCTGAGCAACGGCGTGAGCGTGCTGACCAGCAA
GGTGCTGGACCTGAAGAACTACATCGACAAGGAGCTGCTGCCCA
AGGTGAACAACCACGACTGCCGCATCAGCAAGATCGAGACCGTG
ATCGAGTTCCAGCAGAAGAACAACCGCCTGCTGGAGATCGCCCCG
CGAGTTCAGCGTGAACGCCGCGCATCACCACCCCCCTGAGCACCT
ACATGCTGACCAACAGCGAGCTGCTGAGCCTGATCAACGACATG
CCCATCACCAACGACCAGAAGAAGCTGATGAGCAGCAACGTGCA
GATCGTGCGCCAGCAGAGCTACAGCATCATGAGCGTGGTGAAGG
AGGAGGTGATCGCCTACGTGGTGCAGCTGCCCATCTACGGCGTG
ATCGACACCCCCCTGCTGGAAGCTGCACACCAGCCCCCTGTGCAC
CACCGACAACAAGGAGGGCAGCAACATCTGCCTGACCCGACCCG
ATCGCGGCTGGTACTGCGACAACGCCGGCAGCGTGAGCTTCTTC 12
CCCCAGACCGAGACCTGCAAGGTGCAGAGCAACCGCGTGTTCTG
CGACACCATGAACAGCCTGACCCTGCCACCGACGTGAACCTGT
GCAACACCGACATCTTCAACACCAAGTACGACTGCAAGATCATG
ACCAGCAAGACCGACATCAGCAGCAGCGTGATCACCAGCATCGG
CGCCATCGTGAGCTGCTACGGCAAGACCAAGTGCACCGCCAGCA
ACAAGAATCGCGGCATCATCAAGACCTTCAGCAACGGCTGCGAC
TACGTGAGCAACAAGGGCGTGACACCGTGAGCGTGGGCAACAC
CCTGTACTACGTGAACAAGCTGGAGGGCAAGGCCCTGTACATCA
AGGGCGAGCCCATCATCACTACTACGACCCCCCTGGTGTTCCTC
AGCGACGAGTTCGACGCCAGCATCGCCCAGGTGAACGCCAAGAT
CAACCAGAGCCTGGCCTTCATCCGCCGCAGCGACGAGCTGCTGC
ACAGCGTGGACGTGGGCAAGAGCACCACCAACGTGGTGATCACC
ACCATCATCATCGTGATCGTGGTGGTGATCCTGATGCTGATCGC
CGTGGGCCTGCTGTTCTACTGCAAGACCAAGAGCACCCCCATCA
TGCTGGGCAAGGACCAGCTGAGCGGCATCAACAACCTGAGCTTC AGCAAGTAA
cDNA of accaaacagagaatccgtgagttacgataaaaaggcgaaggagcaattg genomic
aagtcgcacgggtagaaggtgtgaatctcgagtgcgagcccgaagcac sequence of
aaactcgagaaagccttctgccaacatgtcttccgtatttgatgagta NDV of LaSota
cgaacagctcctcgcggtcagactcgcccaatggagctcatggagg strain with the
gggagaaaaagggtaccttaaaagtagacgtcccggtattcactct nucleic acid
taacagtgatgaccagaagatagatggagctttgtggtattctgcct encoding bovine
ccggattgctgtagcgaagatgccaacaaaccactcaggcaaggtgc RSV F protein
tctcatatctctttatgctccactcacaggaatgaggaaccatgt of strain
tgccCttgcagggaacagaatgaagccacattggccgtgcttgagat ATCC51908
tgatggctttgccaacggcagcggcagttcaacaataggagtggagt (nucleic acid
gtctgaagagagagcacagagatttgcatgatagcaggatctctccc sequence
tcgggcatgcagcaacggaacccgttcgtcacagccggggcCgaaga encoding bovine
tgatgcaccagaagacatcaccgataccctggagaggatcctctctat RSV F protein is
ccaggctcaagtatgggtcacagtagcaaaagccatgactgcgtatga underlined)
gactgcagatgagtcggaaacaaggcgaatcaataagtatatgcagca

aggcagggtccaaaagaatacatcctctacccccgtatgcaggagcac
aatccaactcacgatcagacagtctcttgcatccgcatcttttgggt
tagcgagctcaagagaggccgcaacacggcagggtgggtaccttactta
ttataacctggtaggggacgtagactcatacatcaggaataccgggct
tactgcattctcttgacactcaagtacggaatcaacaccaagacatc
agcccttgacacttagtagcctctcaggcgacatccagaagatgaagca
gctcatgcgtttgtatcggtatgaaaggagataatgcgccgtacatgac
attacttggtgatagtaccagatgagctttgcgcctgccgagtatgc acaactttactcctttgccatgggtatggcatcagtcctagataaagg
tactgggaaataccaatttgccagggactttatgagcacatcattctg
gagacttggagtagagtacgctcaggctcagggaagtagcattaacga
ggatatggctgccgagctaaagctaaccacagcagcaaGgaGgggcct
ggcagctgctgcccaacgggtctccgaGgaGaccagcagcataGacat
gcctactcaacaagtcggagtcctcactgggcttagcgaggggggggtc
ccaagctctacaaggcggatcgaatagatcgcaagggaaccagaagc
cgggggatggggagaccaattcctggatctgatgagagcggtagcaaa
tagcatgaggggaggcgccaaactctgcacagggcactcccaatcggg
gcctcccccaactcctgggccatcccaagataacgacaccgactgggg
gtattgatggacaaaaccagcctgcttcacaaaaacatcccaatgc
cctcacccgtagtcgacccctcgatttgcggctctatatgaccacacc ctcaaacaacatccccctctttcctccctccccctgctgtacaactA
cgTacgccctagataccacaggcacatgcggctcactaacaatcaaa
acagagccgaggggaattagaaaaagtagcgggtagaagagggatattc
agagatcagggcaagtctcccgagtctctgctctctcctctacctgat
agaccaggacaaacatggccacctttacagatgcagagatcgacgagc
tatttgagacaagtggaaactgtcattgacaacataattacagcccagg
gtaaaccagcagagactgttgggaaggagtgaatcccacaaggcaaga
ccaaggtgctgagcgcagcatggggagaagcatgggagcatccagccac
cggccagtcagacaaccccgatcgacaggacagatctgacaaacaac
catccacacccgagcaaacgaccccgcatgacagcccgcggccacat
ccgccgaccagccccccacccaggccacagacgaagccgtcgacacac
agCtcaggaccggagcaagcaactctctgctgttgatgcttgacaagc
tcagcaataaatcgccaatgctaaaaagggcccatggctcgagcccc
aagaggggaatcaccaacgtccgactcaacagcaggggagtcaccca
gtcgcggaaacagtcaggaaagaccgcagaaccaagtcaaggccgccc
ctggaaaccagggcacagacgtgaacacagcatatcatggacaatggg
aggagtcacaactatcagctgggtgaacccctcatgctctccgatcaa
ggcagagccaagacaatacccttgtatctgcggatcatgtccagccac
ctgtagactttgtgaagcgatgatgtctatgatggaggcgatatcac agagagtaagtaagggtgactatcagctagatcttgtcttgaaacaga
catcctccatccctatgatgcgggtccgaaatccaacagctgaaaaacat
ctgttgcatggaagccaacttgggaatgatgaagattctggatc ccggttgtgccaacatttcattctctgagtgatctacgggcagttgccc
gatctcacccgggttttagtttcaggccctggagacccctctccctatg
tgacacaaggaggcgaaatggcacttaataaaactttcgcaaccagtg
cacatccatctgaattgattaaacccgccactgcatgcgggcctgata
taggagtggaaaaggacactgtccgtgcattgatcatgtcacgccc
tgaccccgagttcttcagccaagctcctaagcaagttagatgcagccg
ggtcgatcgaggaaatcaggaaaaatcaagcgcttgcctctaaatggct
aattactactgccacacgtagcggggtccctgtccactcggcatcacac
ggaatctgcaccgagttccccccgcGgTTAGAAAAAATACGGGTAGA
ACCGCCACCATGGCGACAACAGCCATGAGGATGATCATCAGCATTATC
TTCATCTCTACCTATGTGACACATATCACTTTATGCCAAAACATAACA
GAAGAATTTTATCAATCAACATGCAGTGCAGTTAGTAGAGGTTACCTT

AGTGCATTAAGAACTGGATGGTATGATCAAGTGTGGTAACAATAGAGTTG
AGCAAAATACAAAAAATGTGTGTAATAGTACTGATTCAAAAGTGAAA
TTAATAAAGCAAGAACTAGAAAGATACAACAATGCAGTAGTGGAATTG
CAGTCACTTATGCAAAATGAACCGGCCTCCTTCAGTAGAGCAAAAAGA
GGGATACCAGAGTTGATACATTATACAAGAACTCTACAAAAAAGTTT
TATGGGCTAATGGGCAAGAAGAGAAAAAGGAGATTTTTAGGATTCTTG
CTAGGTATTGGATCTGCTATTGCAAGTGGTGTAGCAGTGTCCAAAGTA
CTACACCTGGAGGGAGAGGTGAATAAAATTA AAAATGCACTGCTATCC
ACAAATAAAGCAGTAGTTAGTCTATCCAATGGAGTTAGTGTCTTACT
AGCAAAGTACTTGATCTAAAGA ACTATATAGACAAAGAGCTTCTACCT
AAAGTTAACAATCATGATTGTAGGATATCCAAAATAGAACTGTGATA
GAATTCCAACAAAAAACAATAGATTGTTAGAAATTGCTAGGGAATTT
AGTGTAATGCTGGTATTACCACACCTCTCAGTACATACATGTTGACC
AATAGTGAATTACTATCACTAATTAATGATATGCCTATAACGAATGAC
CAAAAAAAGCTAATGTCAAGTAATGTTCAAATAGTCAGGCAACAGAGT
TATTCCATTATGTCAGTGGTCAAAGAAGAAGTCATAGCTTATGTTGTA
CAATTGCCTATTTATGGAGTTATAGACACCCCCTGTTGGAACTACAC
ACCTCTCCGTTATGCACCACTGATAATAAAGAAGGGTCAAACATCTGC
TTAACTAGGACAGATCGTGGGTGGTATTGTGACAATGCAGGCTCTGTG
TCTTTTTTCCCACAGACAGAGACATGTAAGGTACAATCAAATAGAGTG
TTCTGTGACACAATGAACAGTTTAACTCTGCCTACTGACGTAACTTA
TGCAACACTGACATATTCAATACAAAGTATGACTGTAAAATAATGACA
TCTAAAACTGACATAAGTAGCTCTGTGATAACTTCAATTGGAGCTATT
GTATCATGCTATGGGAAGACAAAATGTACAGCTTCTAATAAAAATCGT
GGAATCATAAAGACTTTTTCCAATGGGTGTGATTATGTATCAAACAAA
GGAGTAGATACTGTATCTGTTGGTAACACACTATATTATGTAAATAAG
CTAGAGGGGAAAGCACTCTATATAAAGGGTGAACCAATTATTAATTAC
TATGATCCACTAGTGTTCCTTCTGATGAGTTTGATGCATCAATTGCC
CAAGTAAACGCAAAAATAAACC AAAGCCTGGCCTTCATACGTCGATCT
GATGAGTTACTTCACAGTGTAGATGTAGGAAAATCCACCACAAATGTA
GTAATTACTACTATTATCATAGTGATAGTTGTAGTGATATTAATGTTA
ATAGCTGTAGGATTACTGTTTTACTGTAAGACCAAGAGTACTCCTATC
ATGTTAGGGAAGGATCAGCTCAGTGGTATCAACAATCTTTCCTTTAGT
AAATGAccccccgcggacccaaggtccaactctccaagcggcaatcct
ctctcgcttcctcagccccactgaatgAtcgcgtaaccgtaattaatc
tagctacatttaagattaagaaaaatacgggtagaattggagtgccc caattgtgccaagatggactcatctaggacaattgggctgtactttga
ttctgccattcttctagcaacctgttagcattccgatcgtctaca
agAcacaggagatgggaagaagcaaatcgccccgcaatataggatcca
gcgccttgacttgtggactgatagtaaggaggactcagtattcatcac
cacctatggattcatctttcaagttgggaatgaagaagccacCgtcgg
catgatcgatgataaacccaagcgcgagttactttccgctgcgatgct
ctgcctaggaagcgtcccaataccggagaccttattgagctggcaag
ggcctgtctcactatgatagtcacatgcaagaagagtgcactaatac
tgagagaatggttttctcagtagtcaggcaccccaagtgtctgcaaa
ctgtaggggtgtggcaaaataactcatcagtgaatgcagtcaagca
cgtgaaagcgccagagaagattcccgggagtggaaccctagaatacaa
ggatgaactttgtctccttgactgtggtaccgaagaGggatgtctacaa gatcccagctgcagtattgaaggtttctggctcgagtctgtacaatct
tgcgctcaatgtcactattaatgtggaggtagacccgaggagtccttt ggtaaactctCtgtctaagctgacagcggatactatgctaactctt
cttgcatattggacttatgaccacTgtagataggaagggggaagaaagt
gacatttgacaagctggaaaagaaaataaggagccttgatctatctgt

cgggctcagtgatgtgctcgggccttcctggttggtaaaaagcaagagg
tgcacggactaagcttttggcacctttctctctagcagtgaggacagc ctgctatcccatagcaaagtcttctcctcaggtggccaagatactctg
gagtcaaaccgcgtgcctgcggagcgttaaaatcattatccaagcagg
taccacgcgctgtcgcagtgaccgccgaccacgaggttacctctac
taagctggagaaggggcacacccttgccaaatacaatccttttaagaa
ataagctgctctctgagattgcgctccgcccactcaccagatcatc atgacacaaaaaactaatctgtcttgattatttacagttagtttacct
gtctatcaagttagaaaaaacacgggtagaagattctggatcccgggt
ggcgccctccaggtgcaagatgggctccagaccttctaccaagaaccc
agcacctatgatgctgactatccgggttgcgctgggtactgagttgcat
ctgtccggcaaactccattgatggcaggcctcttgagctgcaggaat
tgtggttacaggagacaaaagccgtcaacatatacacctcatcccagac
aggatcaatcatagttaagctcctcccgaatctgcccaggataagga
ggcatgtgcgaaagcccccttgatgcatacaacaggacattgaccac
tttgctcaccccccttggtgactctatccgtaggatacaagagtctgt
gactacatctggaggggggagacagggggcgcttataggcgccattat
tggcgggtgtggctcttggggttgcaactgccgcacaaataacagcggc
cgagctctgatacaagccaaacaaaatgctccaacatcctccgact
taaagagagcattgccgcaaccaatgaggctgtgcatgaggtcactga
cggattatcgcaactagcagtgggcagttgggaagatgcagcagtttgt
taatgaccaatttaataaaacagctcaggaattagactgcatcaaaat
tgcacagcaagttggtgtagagctcaacctgtacctaaccgaattgac
tacagtattcggaccacaaatcacttcacctgtttaaacaagctgac tattcaggcactttacaatctagctggtggaaatatggattacttatt
gactaagttaggtgtagggaacaatcaactcagctcattaatcggtag cggcttaatcacgggtaaccctattctatacgactcacagactcaact
cttgggtatacaggttaactctaccttcagtcgggaacctaataatat
gcgtgccacctacttggaaccttatccgtaagcacaaccaggggatt
tgcctcggcacttgtcccAaaagtggtagacacaggtcggttctgtgat agaagaacttgacacctatactgtatagaaactgacttagattata
ttgtacaagaatagtaacgttcctatgtcccctgggtattattcctg cttgagcggcaatacgtcggcctgtatgtactcaaagaccgaaggcgc
acttactacaccatacatgactatcaaaggttcagtcacgcgcaactg
caagatgacaacatgtagatgtgtaaacccccgggtatcatatcgca
aaactatggagaagccgtgtctctaatagataaacaatcatgcaatgt ttatccttaggcgggataactttaaggctcagtggggaattcgatgt
aacttatcagaagaatatctcaatacaagattctcaagtaataataac aggcaatcttgatatctcaactgagcttggaatgtcaacaactcgat
cagtaatgctttgaataagttagaggaaaagcaacagaaaactagacaa agtcaatgtcaaactgactagcacatctgctctcattacctatatcgt
tttgactatcatatctctgtttttgggtatacttagcctgattctagc atgctacctaattgtacaagcaaaaggcgcacaacaaagaccttattatg
gcttggaataatactctagatcagatgagagccactacaaaaatgtg
aacacagatgaggaacgaaggtttccctaataagtaatttgtgtgaaag ttctggtagtctgtcagttcagagagttaagaaaaaactaccggttgt
agatgacaaaggacgatatacgggtagaacggtaagagaggccgccc
ctcaattgcgagccaggcttcacaacctccgttctaccgcttcaccga
caacagtcctcaatcatggaccgcgccgttagccaagttgcgttagag
aatgatgaaagagaggcaaaaaatacatggcgcttgatattccggatt gcaatcttattcttaacagtagtgaccttggtctatatctgtagcctcc
cttttatatagcatgggggctagcacacctagcgatctttaggcata
ccgactaggatttcaggggcagaagaaaagattacatctacacttggt
tccaatcaagatgtagtagataggatatataagcaagtggcccttgag tctccgttggcattgttaaatactgagaccacaattatgaacgcaata
acatctctctctatcagattaatggagctgcaacaacagtggggtgg
ggggcacctatccatgaccagattataggggggataggcaaagaa
ctcattgtagatgatgtagtgatgtcacatcattctatccctctgca ttcaagaacatctgaattttatcccggcgctactacaggatcaggt
tgcactcgaataccctcatttgacatgagtgctaccattactgctac acccataatgtaatatgtctggatgcagagatcactcacattcatat
cagtatttagcacttggtgtgctccggacatctgcaacagggagggtta ttcttttactctgcgttccatcaacctggacgacacccaaaaatcgg
aagtcttgagtgtagtgcaactcccctgggttgatgtgctgtgc
tcgaaagtcacggagacagaggaagaagattataactcagctgtccct
acgcggatggtacatgggagggttaggggtcgacggccagtagccacgaa

aaggacctagatgtcacacattattcggggactgggggccaactac
ccaggagtaggggggtggatcttttattgacagccgcgtatggttctca
gtctacggagggttaaaacccaattcacccagtacactgtacaggaa
gggaaatatgtgatatacaagcgatacaatgacacatgccagatgag
caagactaccagattcgaatggccaagtcttcgtataagcctggacgg
tttgggtgggaaacgcatacagcaggctatcttatctatcaaggtgtca
acatccttaggcgaagacccgggtactgactgtaccgccaacacagtc
acactcatggggggccgaaggcagaattctcacagtagggacatctcat ttcttgatcaacgagggtcatcatacttctctcccgcgttattatat
cctatgacagtcagcaaaaaacagccactcttcatagtccttataca ttcaatgccttcactcggccaggtagtatcccttgccaggcttcagca
agatgccccaaactcgtgtgttactggagtgctatacagatccatatccc ctaatcttctatagaaaccacaccttgcgaggggtattcgggacaatg
cttgatgggtgtacaagcaagacttaaccctgcgtctgcagtattcgat
agcacatcccgagtcgcattactcgagtgaagcagtagtcaagcagtagtcaaaa
gcagcatacacaacatcaactgttttaaagtgggtcaagactaataag acctattgtctcagcattgctgaaatatctaatactctcttcggagaa
ttcagaatcgctcccgttactagttgagatcctcaaagatgacgggggt
agagaagccaggctctggctagttgagtcaattataaaggagttggaaa
gatggcattgtatcacctatcttctgcgacatcaagaatcaaaccgaa
tgccggcgcggtgctcgaattccatgttgccagttgaccacaatcagcc
agtgtcatgcgatcagattaagccttgtcaAtaGtctcttgattaag
aaaaaatgtaagtggcaatgagatacaaggcaaaacagctcatgggTa
aCaatacgggtaggacatggcgagctccgggtcctgaaagggcagagca
tcagattatctaccagagTcacacctgtcttcaccattgggtcaagca caaactactcttattactggaaattaactgggctaccgcttctctgatga
atgtgacttcgaccacctcattctcagccgacaatggaaaaaataact
tgaatcggcctctcctgatactgagagaatgataaaactcggaagggc
agtacaccaaactcttaaccacaattccagaataaccggagtgctcca
ccccagggtgttagaaGaactggctaataattgaggtcccagattcaac
caacaaatttcggaagattgagaagaagatccaaattcacaacacgag
atatggagaactgttcacaaggctgtgtacgcataatagagaagaaact
gctggggtcatttgggttaacaatgtcccccggtcagaggagttcag
cagcattcgtacggatccggcattctgggttctactcaaaatgggtccac
agccaagtttgcattgggtccatataaaacagatccagaggcatctgat
gggtggcagctaGgacaaggctctgcggccaacaaattgggtgatgctaac
ccataaggtaggccaagtctttgtcactcctgaactgtcgttgtgac gcatacgaatgagaacaagttcacatgtcttaccaggaacttgatt
gatgtatgcagatatgatggagggcagagatatgggtcaacataatc
aaccacggcggtgcattcagaagcttatcagagaaaattgatgacat
tttgcgggttaatagacgctctggcaaaagacttgggtaatacaagtcta cgatgttgatcactaatggaggggatttgcatacggagctgtccagct
actcgagccgtcaggtacatttgcaggagatttcttcgcattcaacct gcaggagcttaaagacatttctaattggcctcctcccaatgatatagc
agaatccgtgactcatgcaatcgtactgtattctctggtttagaaca gaatcaagcagctgagatgttgtgtctgttgcgtctgtggggtcacc
actgcttgagtcccgtattgcagcaaaggcagtcaggagccaaatgtg cgcaccgaaaatggtagactttgatatgatccttcagggtactgtcttt
cttcaagggaacaatcatcaacgggtacagaaagaagaatgcaggtgt
gtggccgcgagtcaaagtggatacaatatatgggaagggtcattgggca
actacatgcagattcagcagagatttcacacgatatcatgttgagaga gtataagagtttatctgcacttgaatttgagccatgtatagaatatga
ccctgtcaccaacctgagcatgttcctaaaagacaaggcaatcgaca
ccccaacgataattggcttgctcgtttaggcggaaccttctctccga
agaccagaagaaacatgtaaaagaagcaacttcgactaatcgctctt gatagagtttttagagtcaaataatttgatccatataaagagatgga
atatctgacgaccttgagtaccttagagatgacaatgtggcagtatc
atactcgtcgaaggagaagggaagtgaaggttaattggacggatcttcgc
taagctgacaaagaaggttaagggaactgtcaggtgatggcggaaggat
cctagccgatcagattgcacctttcttcagggaatggagtcattca ggatagcatatccttgaccaagagtatgctagcgatgagtcaactgtc
tttaacagcaataagaaacgtatcactgactgtaaagaaagagtatc
ttcaaaccgcaatcatgatccgaaaagcaagaaccgtcggagagttgc

aacctcataacaactgacctgcaaaagtactgtcttaattggagata tcagacaatcaaattgttcgctcatgccatcaatcagttgatgggcct
acctcacttcttcgaatggattcacctaagactgatggacactacgat gttcgtaggagacccttcaatcctccaagtgaccctactgactgtga
cctctcaagagtcctaatgatgacatatattgtcagtgccagagg
gggtatcgaaggattatgccagaagctatggacaatgatctcaattgc tgcaatccaacttgctgcagctagatcgattgtcgtgttgccctgat
ggtacaggggtgataatcaagtaatagcagtaacgagagaggtaagatc
agacgactctccggagatgggtgttgacacagttgcatcaagccagtga taatttctcaaggaattaattcatgtcaatcatttgattggccataa
tttgaaggatcgtgaaaccatcaggtcagacacattcttcatatacag
caaacgaatcttcaaagatggagcaatcctcagtcagtcaagtcctcaaaaa ttcacttaaattagtgtcagtgatctcagtgaaaacaccgt
aatgtcctgtgccaacattgcctctactgtagcacggctatgcgagaa cgggcttcccaaagacttctgttactatttaactatataatgagttg
tgtgcagacatactttgactctgagttctccatcaccaacaattcgca ccccgatcttaatcagtcgtggattgaggacatctctttgtgcactc
atatgttctgactcctgcccattagggggactgagtaaccttcaata ctcaaggctctacactagaaatcgggtgacccggggactactgcttt
tgcagagatcaagcgactagaagcagtgaggattactgagtcctaacat
tatgactaatatcttaactaggccgctgggaatggagattgggccag tctgtgcaacgaccatactctttcaatttgagactgttgcaagccc
aaatattgttcttaagaaacatacgcaaagagtcctatttgaaacttg ttcaaatccctattgtctggagtgcacacagaggataatgaggcaga
agagaaggcattggctgaattcttgcttaatcaagaggtgattcatcc
ccgcgttgcgcatgccatcatggaggcaagctctgtaggtaggagaaa
gcaaattcaagggtgttgacacaacaaacaccgtaattaagattgc
gcttactaggaggccattaggcatcaagaggctgatgcgatagtgcaa ttattctagcatgcatgcaatgctgtttagagacgatgttttctc
cagtagatccaaccaccccttagtctcttctaatatgtgttctctgac
actggcagactatgcacggaatagaagctggtcaccttgacgggagg
caggaaaatactgggtgtatctaactctgatacgatagaactcgtaga
gggtgagattcttagtgtaagcggagggtgtacaagatgtgacagcgg
agatgaacaatttacttggttccatcttccaagcaatatagaattgac
cgatgacaccagcaagaatcctccgatgagggtaccatatctcgggtc
aaagacacaggagaggagagctgcctcacttgcaaaaatagctcatat
gtcgccacatgtaaaggctgcctaagggtcatcatccgtgttgatctg ggcttatggggataatgaagtaaattggactgctgctcttacgattgc
aaaatctcgggtgaatgtaaacttagagtatcttcgggtactgtcccc ttaccacggctgggaatcttcaacatagactagatgatggtataac
tcagatgacattcacccctgcatctcttacaggGgtcaccttacat tcacatatccaatgattctcaaaggctgttactgaagaaggagtcaa
agaggggaatgtggtttaccaacagatcatgctcttgggtttatctct aatcgaatcgatctttccaatgacaacaaccaggacatatgatgagat
cacactgcacctacatagtaaatttagttgctgtatcagagaagcacc
tgttgcggttccttcgagctacttgggggtggtaccggaactgaggac agtgacctcaaataagtttatgtatgatcctagccctgtatcggaggg
agactttgcgagacttgacttagctatctcaagagttatgagcttaa tctggagtcatatccacgatagagctaataaacattcttcaatc
cagcgggaagtgattggccagctctgtggttcttatgatgaagatac ctccataaagaatgacgccataatagtgtatgacaatacccgaattg
gatcagtgaaagctcagaattcagatgtgggtccgcctattgaaatgc agcactgaaagtgcctcgcactgttcttaccactctattacctgag
agtaagaggcctGgacaatattgtcttatatatgggtgattatacaa gaatatgccaggaattctactttcaacattgcagctacaatatctca
tcccgatcattcattcaaggttacatgcagtgggcctggtcaaccatga cggatcacaccaactgcagatacggattttatcgaaatgtctgcaaa
actattagtatcttgacccgacgtgtgatctccggcttatattcagg aaataagtatgatctgctgttcccatctgtcttagatgataacctgaa
tgagaagatgcttcagctgatatcccgggtatgtgtctgtacacgggt actcttctgacacaagagaaaatcccgaataagaggcttaactgc
agaagagaaatgttcaatactactgagtatttactgtcggatgctgt gaaaccattacttagccccgatcaagtgagctctatcatgtctcctaa
cataattacattcccagctaattctgtactacatgtctcggaagagcct caattgatcagggaaaggaggagcagggatactatcctggcgtgtt
gttcccccaagagccattattagagttcccttctgtgcaagatattgg tgctcgagtgaagatccattcacccgacaacctgcggcatttttgca
agagttagatttgagtgtccagcaaggtatgacgcattcacacttag tcagattcatctgaactcacatctccaaatccggaggaagactactt
agtacgatactgttcagagggatagggactgcatcttcttcttggtg taaggcatctcatctccttctgtacctcgaggtaagatgtgcaagaca
cgggaactccttatacttagctgaagggagcggagccatcatgagtct tctcgaactgcatgtaccacatgaaactatctattacaatacgtctt
ttcaaatagagatgaacccccgcaacgacatttcgggcccgaacccaaac
tcagttttgaattcggtgtttataggaatctacaggcggaggtaac atgcaaagatggatttgcgaagagttccgtccattatggagagaaaa
tacagaggaaagCgacctgacctcagataaagTagtggggtatattac
atctgcagtgccctacagatctgtatcattgtcgttgatgtgacattga aattcctccagggtccaatcaaagcttactagatcaactagctatcaa
tttatctctgattgccatgcattctgtaaggaggggcggggtagtaac catcaaagtgttgatgcaatgggatactactttcatctactcatgaa
cttgtttgctccgtgttccacaaaaggatatattctcttaattgggtta tgcattgtcaggagatatggagtgttacctgggtatttgcattgggtta

cctgggcgggcctacattgtacatgaggtggtgaggatggcGaaaac
tctggtgcagcggcacggtacgctTttgtctaaatcagatgagatcac
actgaccaggttattcacctcacagcggcagcgtgtgacagacatcct
atccagtcctttaccaagattaataaagtacttgaggaagaatattga
cactgcgctgattgaagccgggggacagcccgtccgtccattctgtgc
ggagagtctggtgagcacgctagcgaacataactcagataaccagat
Catcgctagtcacattgacacagttatccgggtctgtgatatatatgga agctgaggggtgatctcgctgacacagttattctattacccttataa
tctcttactgacgggaaaaagaggacatcacttaAacagtgcacgag
acagatcctagagggttacaatactaggtcttagagtcgaaaaatctcaa taaaataggcgatataatcagcctagtgcctaaaggcatgatctccat
ggaggaccttatcccactaaggacatacttgaagcatagtacctgccc taaatatttgaaggctgtcctaggtattaccaaactcaaagaaatgtt
tacagacacttctgtaCtgtacttgactcgtgctcaacaaaaattcta catgaaaactataggcaatgcagtcaaaggatattacagtaactgtga
ctcttaacgaaaaatcacatattaataggctccttttttgccaattgt attcttgttgatttaatcatattatgttagaaaaaagttgaaccctga
ctccttaggactcgaattcgaactcaaataaatgtcttaaaaaaagggt tgcgcacaattattcttgagtgtagtctcgctcattcaccaaattcttg
tttgggt CDNA of accaaacagagaatccgtgagttacgataaaaggcgaaggagcaattg 13 genomic
aagtcgcacgggtagaagggtgtaatctcgagtcgagccccgaagcac sequence of
aaactcgagaaaagccttctgccaacatgtcttccgtatttgatgagta NDV LaSota
cgaacagctcctcgcggtcagactcgccccaatggagctcatggagg strain with the
gggagaaaaagggagtaccttaaaagtagacgtcccggtattcactct codon optimized
taacagtgatgaccagaagatagatggagctttgtggtattctgcct nucleic acid
ccggattgctgttagcgaagatgccaacaaaccactcaggcaaggtgc sequence
tctcatatcttttatgtctccactcacaggaatgaggaaccatgt encoding bovine
tgccCttgcagggaacagaatgaagccacattggccgtgcttgagat RSV F protein
tgatggctttgccaacggcacgccccagttcaacaataggagtggagt of strain
gtctgaagagagagcacagagatttgcgatgatagcaggatctctccc ATCC51908
tcgggcatgcagcaacggaaccccgttcgtcacagccggggcCgaaga inserted (codon
tgatgcaccagaagacatcaccgataccctggagaggatcctctctat optimized
ccaggctcaagtatgggtcacagtagcaaaagccatgactgcgtatga nucleic acid
gactgcagatgagtcggaaacaaggcgaatcaataagtatatgcagca sequence of
aggcagggtccaaaagaaatacatcctctaccccgtatgcaggagcac bovine RSV F
aatccaactcacgatcagacagtctcttgagtcgcatcttttgggt protein is
tagcgagctcaagagaggccgcaacacggcaggtggtaccttactta underlined)
ttataacctggtaggggacgtagactcatacatcaggaataccgggct
tactgcattcttctgacactcaagtacggaatcaacaccaagacatc
agcccttgacttagtagcctctcaggcgacatccagaagatgaagca
gctcatgcgtttgtatcggtgaaaggagataatgcgccgtacatgac
attacttgggtgatagtaccagatgagctttgcgcctgccgagtatgc acaactttactcctttgccatgggtatggcatcagtcctagataaagg
tactgggaaataccaatttgccagggactttatgagcacatcattctg
gagacttggagtagagtacgctcaggctcagggaagtagcattaacga
ggatatggctgccgagctaaagctaacccagcagcaaGgaGgggcct
ggcagctgctgccaacgggtctccgaGgaGaccagcagcataGacat
gcctactcaacaagtcggagtcctcactgggcttagcgagggggggtc
ccaagctctacaaggcggatcgaatagatcgcaagggcaaccagaagc
cggggatggggagacccaattcctggatctgatgagagcggtagcaaa
tagcatgagggaggcgccaaactctgcacagggcactccccaatcggg
gcctccccaaactcctgggccatcccaagataacgacaccgactgggg
gtattgatggacaaaaccagcctgcttcacaaaaacatcccaatgc
cctcaccgtagtcgacccctcgatttgcggctctatatgaccacacc ctcaaacaacatccccctctttcctccctccccctgctgtacaactA
cgTacgccctagataccacaggcacaatgcgggtcactaacaatcaaa
acagagccgagggaattagaaaaaagtacgggtagaagagggatattc
agagatcagggcaagtctcccaggtctctgctctcctctacctgat

agaccaggacaaacatggccacctttacagatgcagagatcgacgagc
tatttgagacaagtggaaactgtcattgacaacataattacagcccagg
gtaaaccagcagagactgttggaaggagtgaatcccacaaggcaaga
ccaaggtgctgagcgcagcatgggagaagcatgggagcatccagccac
cggccagtcaagacaaccccgatcgacaggacagatctgacaaacaac
catccacacccgagcaaacgaccccgcatgacagcccgcggccacat
ccgccgaccagccccccacccaggccacagacgaagccgtcgacacac
agCtcaggaccggagcaagcaactctctgctgttgatgcttgacaagc
tcagcaataaatcgccaatgctaaaaagggccccatggctgagcccc
aagaggggaatcaccaacgtccgactcaacagcaggggagtcaccca
gtcgcggaaacagtcaggaaagaccgcagaaccaagtcaaggccgccc
ctggaaaccagggcacagacgtgaacacagcatatcatggacaatggg
aggagtcacaactatcagctgggtgaacccctcatgctctccgatcaa
ggcagagccaagacaatacccttgatatctgcggatcatgtccagccac
ctgtagactttgtgaagcgatgatgtctatgatggaggcgatatcac agagagtaagtaagggtgactatcagctagatcttgtcttgaaacaga
catcctccatccctatgatgcgggtccgaaatccaacagctgaaaacat
ctgttgcatgcatggaagccaacttgggaatgatgaagattctggatc ccggttggtgccaacatttcatctctgagtgatctacgggcagttgccc
gatctcaccgggttttagtttcaggccctggagacccctctccctatg
tgacacaaggaggcgaaatggcacttaataaaactttcgcaaccagtgc
cacatccatctgaattgattaaacccgccactgcatgcgggcctgata
taggagtggaaaaggacactgtccgtgcattgatcatgtcacgccc
tgcacccgagttcttcagccaagctcctaagcaagttagatgcagccg
ggctgatcgaggaaatcaggaaaatcaagcgcttgctctaaatggct
aattactactgccacacgtagcgggtccctgtccactcggcacacac
ggaatctgcaccgagttccccccgcGgTTAGAAAAAATACGGGTAGA

ACCGCCACCATGGCCACCACCGCCATGCGCATGATCATCAGCATCATC
TTCATCAGCACCTACGTGACCCACATCACCTGTGCCAGAACATCAC
GAGGAGTTCTACCAGAGCACCTGCAGCGCCGTGAGTCGCGGCTACCTG
AGCGCCCTGCGCACCGGCTGGTACACCAGCGTGGTGACCATCGAGCTG
AGCAAGATCCAGAAGAACGTGTGCAACAGCACCGACAGCAAGGTGAAG
CTGATCAAGCAGGAGCTGGAGCGCTACAACAACGCCGTGGTGAGCTG
CAGAGCCTGATGCAGAACGAGCCCGCCAGCTTCAGCCGCGCCAAGCGC
GGCATCCCCGAGCTGATCCACTACACCCGCAACAGCACCAAGAAGTTC
TACGGCCTGATGGGCAAGAAGCGCAAGCGCCGCTTCCTGGGCTTCCTG
CTGGGCATCGGCAGCGCCATCGCCAGCGGCGTGGCCGTGAGCAAGGTG
CTGCACCTGGAGGGCGAGGTGAACAAGATCAAGAACGCCCTGCTGAGC
ACCAACAAGGCCGTGGTGAGCCTGAGCAACGGCGTGAGCGTGCTGACC
AGCAAGGTGCTGGACCTGAAGAACTACATCGACAAGGAGCTGCTGCCC
AAGGTGAACAACCACGACTGCCGCATCAGCAAGATCGAGACCGTGATC
GAGTTCCAGCAGAAGAACAACCGCCTGCTGGAGATCGCCCGCGAGTTC
AGCGTGAACGCCGGCATCACCAACCCCTGAGCACCTACATGCTGACC
AACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCATCACCAACGAC
CAGAAGAAGCTGATGAGCAGCAACGTGCAGATCGTGCGCCAGCAGAGC
TACAGCATCATGAGCGTGGTGAAGGAGGAGGTGATCGCCTACGTGGTG
CAGCTGCCCATCTACGGCGTGATCGACACCCCTGCTGGAAGCTGCAC
ACCAGCCCCCTGTGCACCACCGACAACAAGGAGGGCAGCAACATCTGC
CTGACCCGCAACCGATCGCGGCTGGTACTGCGACAACGCCGGCAGCGTG
AGCTTCTTCCCCCAGACCGAGACCTGCAAGGTGCAGAGCAACCGCGTG
TTCTGCGACACCATGAACAGCCTGACCCTGCCACCGACGTGAACCTG
TGCAACACCGACATCTTCAACACCAAGTACGACTGCAAGATCATGACC

AGCAAGACCGACATCAGCAGCGTGATCACCAGCATCGGCGCCATC
GTGAGCTGCTACGGCAAGACCAAGTGCACCGCCAGCAACAAGAATCGC
GGCATCATCAAGACCTTCAGCAACGGCTGCGACTACGTGAGCAACAAG
GGCGTGGACACCGTGAGCGTGGGCAACACCCTGTACTACGTGAACAAG
CTGGAGGGCAAGGCCCTGTACATCAAGGGCGAGCCCATCATCAACTAC
TACGACCCCCTGGTGTTCCTCCAGCGACGAGTTCGACGCCAGCATCGCC
CAGGTGAACGCCAAGATCAACCAGAGCCTGGCCTTCATCCGCCGCAGC
GACGAGCTGCTGCACAGCGTGGACGTGGGCAAGAGCACCACCAACGTG
GTGATCACCACCATCATCATCGTGATCGTGGTGGTGATCCTGATGCTG
ATCGCCGTGGGCCTGCTGTTCTACTGCAAGACCAAGAGCACCCCCATC
ATGCTGGGCAAGGACCAGCTGAGCGGCATCAACAACCTGAGCTTCAGC
AAGTAAccccccgcggaaccaaggtccaactctccaagcggcaatcct
ctctcgcttctcagccccactgaatgAtcgcgtaaccgtaattaatc
tagctacatttaagattaagaaaaatacgggtagaattggagtgccc caattgtgccaagatggactcatctaggacaattgggctgtactttga
ttctgcccattctttagcaacctgttagcatttccgatcgtcctaca
agAcacaggagatgggaagaagcaaatcgccccgcaatataggatcca
gcgcccttgactgtggactgataagtaaggaggactcagtattcatcac
cacctatggattcatctttcaagttgggaatgaagaagccacCgtcgg
catgatcgatgataaacccaagcgcgagttactttccgctgcgatgct
ctgcctaggaagcgtcccaataccggagaccttattgagctggcaag
ggcctgtctcactatgatagtcacatgcaagaagagtgcaactaatac
tgagagaatggttttctcagtagtgagggcaccccaagtgctgcaag
ctgtaggggttgggcaaaataactcatcagtgaatgcagtcaagca
cgtgaaagcgccagagaagattcccgggagtggaaccctagaatacaa
gggtgaactttgtctccttgactgtggtagcgaagaGggatgtctacaa gatcccagctgcagattgaaggtttctggctcgagtctgtacaatct
tgcgctcaatgtcactattaatgtggaggtagacccgaggagtccttt ggtaaactctCtgtctaagctgacagcggatactatgctaaccttt
cttgcatattggacttatgaccacTgtagataggaaggggaagaaagt
gacatttgacaagctggaaaagaaaataaggagccttgatctatctgt
cgggctcagtgatgtgctcgggccttccgtgttggtgtaaaagcaagagg
tgcacggactaagcttttggcacctttctctctagcagtgggacagc ctgctatcccatagcaaagtcttctcctcaggtggccaagatactctg
gagtcaaaccgcgtgcctgcggagcgttaaaatcattatccaagcagg
taccacacgcgctgtcgcagtgaccgccgaccacgaggttacctctac
taagctggagaaggggacaccccttgccaaatacaatccttttaagaa
ataagctgcgtctctgagattgcgctccgcccactcaccagatcatc atgacacaaaaaactaatctgtcttgattatttacagttagttacct
gtctatcaagttagaaaaaacacgggtagaagattctggatcccggtt
ggcgccctccaggtgcaagatgggctccagaccttctaccaagaaccc
agcacctatgatgtgactatccgggttgcgctggtagtgcattgcat
ctgtccggcaaacctcattgatggcaggcctcttgacgtgcaggaat
tgtggttacaggagacaaagccgtcaacatataacctcatcccagac
aggatcaatcatagttaagctcctcccgaatctgcccaggaagga
ggcatgtgcgaaagcccccttggtatgcatacaacaggacattgaccac
tttgctcaccccccttggtgactctatccgtaggatacaagagctgtg
gactacatctggaggggggagacaggggcgcttataggcgccattat
tggcggtgtggctcttgggggtgcaactgccgcacaaataacagcggc
cgcagctctgatacaagccaaacaaatgtgccaacatcctccgact
taaagagagcattgccgcaaccaatgaggctgtgcatgaggtcactga
cggattatcgcaactagcagtggcagttgggaagatgcagcagtttgt
taatgaccaatttaataaaacagctcaggaattagactgcatcaaaat
tgcacagcaagttggtgtagagctcaacctgtacctaaccgaattgac
tacagtattcggaccacaaatcacttcacctgtttaaacaagctgac tattcaggcactttacaatctagctgggtggaaatatggattacttatt

gactaagtttaggtgttaggggaacaaactcaactcagctcattaatcggtag cggttaatcacccggtaacctattctatacgaactcacagactcaact
cttgggtatacaggaactctaccttcagtcgggaacctaataatat
gcgtgccacctacttggaaccttatccgtaagcacaaccaggggatt
tgcctcggcacttgtcccAaaagtgggtgacacaggtcgggttctgtgat agaagaacttgacacctcatactgtatagaaactgacttagattata
ttgtacaagaatagtaacgttccctatgtcccctgggtatttattcctg cttgagcggcaatacgtcggcctgtatgtactcaaagaccgaaggcgc
acttactacaccatacatgactatcaaagggttcagtcatcgccaactg
caagatgacaacatgtagatgtgtaaacccccgggtatcatatcgca
aaactatggagaagccgtgtctctaatagataaacaatcatgcaatgt tttatccttaggcgggataactttaaggctcagtggggaattcgatgt
aacttatcagaagaatatctcaatacaagatttcaagtaataataac aggcaatcttgatatctcaactgagcttgggaatgtcaacaactcgat
cagtaatgctttgaataagttagaggaaagcaacagaaaactagacaa agtcaatgtcaaactgactagcacatctgctctcattacctatatcgt
tttgactatcatatctcttgttttgggtatacttagcctgattctagc atgctacctaattgtacaagcaaaaggcgcaacaaaagaccttattatg
gcttgggaataatactctagatcagatgagagccactacaaaaatgtg
aacacagatgaggaacgaagggttccctaatagtaatttgtgtgaaag ttctggtagtctgtcagttcagagagttaagaaaaaactaccggttgt
agatgaccaaaggacgatatacgggtagaacggtaagagaggccgccc
ctcaattgcgagccagggttcacaacctccgttctaccgcttcaccga
caacagtcctcaatcatggaccgcgccgttagccaagttgcgttagag
aatgatgaaagagaggcaaaaaatacatggcgcttgatattccggatt gcaatcttattcttaacagtagtgaccttggctatatctgtagcctcc
cttttatatagcatgggggctagcacacctagcgaatctttagggcata
ccgactaggatttccagggcagaagaaaagattacatctacacttgg
tccaatcaagatgtagtagataggatatataagcaagtggcccttgag tctccgttggcattgttaaatactgagaccacaattatgaacgcaata
acatctctcttctatcagattaatggagctgcaacaacagtggggtgg
ggggcacctatccatgaccagattatatagggggggataggcaaagaa
ctcattgtagatgatgtagtgatgtcacatcattctatccctctgca tttaagaacatctgaattttatccccggcgctactacaggatcaggt
tgcactcgaataccctcatttgacatgagtgctacccattactgtac acccataatgtaatatgtctggatgcagagatcactcacattcatat
cagtatttagcacttgggtgtgctccggacatctgcaacaggggagggtta ttcttttactctgcgttccatcaacctggacgacacccaaaaatcgg
aagtcttgcagtgtgagtgcaactcccctgggttgatgatgtgtgc
tcgaaagtcacggagacagaggaagaagattataactcagctgtccct
acgcggatggtacatgggagggttaggggttcgacggccagtaccacgaa
aaggacctagatgtcacaacattattcggggactgggtggccaactac
ccaggagtaggggggtggatctttattgacagccgcgtatggttctca
gtctacgggagggttaaaaccaattcaccagtgacactgtacaggaa
gggaaatatgtgatatacaagcgatacaatgacacatgccagatgag
caagactaccagattcgaatggccaagtcttcgtataagcctggacgg
tttgggtgggaaacgcatacagcaggctatcttatctatcaaggtgtca
acatccttaggcgaagacccgggtactgactgtaccgccaacacagtc
acactcatggggggccgaaggcagaattctcacagtagggacatctcat ttcttgatcaacgagggtcatcatacttctctcccgcttattatat
cctatgacagtcagcaaaaaacagccactcttcatagtccttataca ttcaatgccttcactcggccaggtagtatcccttgccaggcttcagca
agatgccccaaactcgtgtgttactggagtcatacagatccatatccc ctaatctctatagaaaccacaccttgcgaggggtattcgggacaatg
cttgatgggtgtacaagcaagacttaaccctgcgtctgcagtattcgat
agcacatcccgcagtcgcattactcgagtgagttcaagcagtagcaaaa
gcagcatacacaacatcaactgttttaaagtgggtcaagactaataag acctattgtctcagcattgtgaaatatctaatactctcttcggagaa
ttcagaatcgctccggttactagttgagatcctcaaagatgacgggggt
agagaagccagggtctggctagttgagtcattataaaggagttggaaa
gatggcattgtatcacctatcttctgcgacatcaagaatcaaaaccgaa
tgccggcgcggtgctcgaattccatgttgccagttgaccacaatcagcc
agtgtcatgcgatcagattaagccttgtcaAtaGtctcttgattaag
aaaaaatgtaagtggcaatgagatacaaggcaaaacagctcatgggTa
aCaatacgggtaggacatggcgagctccggctctgaaagggcagagca
tcagattatcctaccagagTcacacctgtcttcaccattgggtcaagca caaactactctattactggaaattaactgggctaccgcttctgatga
atgtgacttcgaccacctcattctcagccgacaatggaaaaaataact

tgaatcggcctctctgatactgagagaatgataaaactcggaagggc
agtacaccaaactcttaaccacaattccagaataaccggagtgctcca
ccccagggtgttagaaGaactggctaattattgagggtcccagattcaac
caacaaatttcggaagattgagaagaagatccaaattcacacacgag
atatggagaactgttcacaaggctgtgtacgcatatagagaagaaact
gctgggggtcatcttgggtctaacaatgtcccccggtcagaggagttcag
cagcattcgtacggatccggcattctgggttccactcaaaatgggccac
agccaagtttgcattggctccatataaaacagatccagaggcatctgat
gggtggcagctaGgacaaggctcgcggccaacaaattgggtgatgctaac
ccataaggtaggccaagtctttgtcactcctgaacttgtcgttgtag gcatacgaatgagaacaagttcacatgtcttaccaggaacttgtatt
gatgtatgcagatatgatggagggcagagatatggccaacataatc
aaccacggcgggtgcatctcagaagcttatcagagaaaattgatgacat
tttgcggttaatagacgctctggcaaaagacttgggtaatacgaagtcta cgatgtgtatcactaatggagggttgcatacggagctgtccagct
actcgagccgtcaggtacatttgcaggagatttcttcgcattcaacct gcaggagcttaaagacatttctaattggcctcctcccaatgatatagc
agaatccgtgactcatgcaatcgctactgtattctctggttagaaca gaatcaagcagctgagatgttgtgtctgttgcgtctgtgggggtcaccc
actgcttgagtcccgtattgcagcaaaaggcagtcaggagccaaatgtg cgcaccgaaaatggtagactttgatatgatccttcagggtactgtcttt
cttcaagggaacaatcatcaacgggtacagaaagaagaatgcagggtgt
gtggccgcgagtcaaagtggatacaatatatgggaagggtcattgggca
actacatgcagattcagcagagatttcacacgatatcatgttgagaga gtataagagtttatctgcacttgaatttgagccatgtatagaatatga
ccctgtcaccaacctgagcatgttcctaaaagacaaggcaatcgcaca
ccccaacgataattggcctgcctcgtttaggcggaaccttctctccga
agaccagaagaaacatgtaaaagaagcaacttcgactaatcgctctt gatagagtttttagagtcaaataatttgatccatataaagagatgga
atatctgacgaccttgagtaccttagagatgacaatgtggcagtatc
atactcgtcaaggagaagggaagtgaaggttaattggacggatcttcgc
taagctgacaaagaagttaaggaaactgtcaggtgatggcgggaagggat
cctagccgatcagattgcaccttcttcagggaatggagtcattca ggatagcatatccttgaccaagagtatgctagcgatgagtcactgtc
ttttaacagcaataagaaacgtatcactgactgtaaagaaagagtatc
ttcaaaccgcaatcatgatccgaaaagcaagaaccgtcggagagttgc
aaccttcataacaactgacctgcaaaagtactgtcttaattggagata tcagacaatcaaattgttcgctcatgccatcaatcagttgatgggcct
acctcacttcttgaatggattcacctaagactgatggacactacgat gttcgtaggagaccctttcaatcctcaaagtgaccctactgactgtga
cctctcaagagtcctaatgatgacatatatattgtcagtgccagagg
gggtatcgaaggattatgcagaagctatggacaatgatctcaattgc tgcaatccaacttgctgcagctagatcgcattgtcgtgttgctgtat
ggtacagggtgataatcaagtaatagcagtaacgagagaggtaagatc
agacgactctccggagatgggtgtgacacagttgcatcaagccagtga taatttctcaagggaattaattcatgtcaatcatttgattggccataa
tttgaaggatcgtgaaaccatcagggtcagacacattctcatatacag
caaacgaatcttcaaagatggagcaatcctcagtcgaagtcctcaaaaa ttcattctaaattagtgtcaggtgatctcagtgaaaacaccgt
aatgtcctgtgccaacattgcctctactgtagcacggctatgcgagaa cgggcttcccaaagacttctgttactatttaaaactatataatgagttg
tgtgcagacatactttgactctgagttctccatcaccaacaattcgca ccccgatcttaatcagtcgtggattgaggacatctctttgtgcactc
atatgttctgactcctgcccaattagggggactgagtaaccttcaata ctcaaggctctacactagaaatatcggtgacccggggactactgcttt
tgacagagatcaagcgactagaagcagtgaggattactgagtcctaacat
tatgactaataatcttaactaggccgcctgggaatggagattgggccag tctgtgcaacgaccatactctttcaattttgagactgttgcaagccc
aaatattgttcttaagaaacatacgcaaagagtcctatttgaaacttg ttcaaatcccttattgtctggagtgcacacagaggataatgaggcaga
agagaaggcattgggtgaattcttgcttaatcaagagggtgattcatcc
ccgcgttgcgcatgccatcatggaggcaagctctgtaggtaggagaaa
gcaaattcaagggttgttgacacaacaaacaccgtaattaagattgc
gcttactaggaggccattaggcatcaagagggtgatgcggatagtcaa ttattctagcatgcatgcaatgctgtttagagacgatgtttttctc
cagtagatccaaccaccccttagtctcttctaatatgtgttctctgac
actggcagactatgcacggaatagaagctgggtcaccttgacggggagg
caggaaaatactgggtgtatctaactcctgatacgaataaactcgtaga
gggtgagattcttagtgaagcggagggtgtacaagatgtgacagcgg

agatgaacaatttacttgggtccatcttccaagcaatatagaattgac
cgatgacaccagcaagaatcctccgatgaggggtaccatatctcgggtc
aaagacacaggagaggagagctgcctcacttgcaaaaatagctcatat
gtcgccacatgtaaaggctgcctaaggggcatcatccgtgttgatctg ggcttatggggataatgaagtaaattggactgctgctcttacgattgc
aaaatctcgggtgaatgtaaacttagagtatcttcgggtactgtcccc ttaccacaggctgggaatcttcaacatagactagatgatggtataac
tcagatgacattcacccctgcatctctctacaggGgtgcacctacat tcacatatccaatgattctcaaaggctgttactgaagaaggagtaa
agaggggaatgtggtttaccaacagatcatgctcttgggtttatctct aatcgaatcgatctttccaatgacaacaaccaggacatatgatgagat
cacactgcacctacatagtaaattagttgctgtatcagagaagcacc
tgttgcggttcctttcgagctacttgggggtggtaccggaactgaggac agtgacctcaaataagtttatgtatgatcctagccctgtatcggaggg
agactttgagacttgacttagctatcttcaagagttagagcttaa tctggagtcatatccacgatagagctaataaacattctttcaatatc
cagcgggaagttgattggccagtctgtggtttcttatgatgaagatac ctccataaagaatgacgccataatagtgtatgacaatacccgaattg
gatcagtgaagctcagaattcagatgtggtccgcctatttgaatatgc agcacttgaagtgtcctcgactgttcttaccactctattacctgag
agtaagaggcctGgacaatatgtcttatatatgggtgattatacaa gaatatgccaggaattctactttccaacattgcagctacaatatctca
tcccgctattcattcaagggtacatgcagtgggcctgggtcaaccatga cggatcacaccaacttgagatacggattttatcgaaatgtctgcaa
actattagtatcttgaccccgacgtgtgatctccggcttatattcagg aaataagtatgatctgtgttccatctgtcttagatgataacctgaa
tgagaagatgcttcagctgatatcccggttatgtgtctgtacacgggt actctttgtacaacaagagaaatcccgaataagaggcttaactgc
agaagagaaatgttcaatactcactgagtatttactgtcggatgtgt gaaaccattacttagccccgatcaagtgagctctatcatgtctctaa
cataattacattcccagctaattctgtactacatgtctcggaagagcct caatttgatcagggaaaaggaggagcagggatactatcctggcggtgtt
gttcccccaagagccattattagagttcccttctgtgcaagatatgg tgctcgagtgaagatccattcacccgacaacctgcggcatttttgca
agagttagatttgagtgtccagcaagggtatgacgcattcacacttag tcagattcatcctgaactcacatctccaaatccggaggaagactactt
agtacgatacttggtcagagggatagggactgcatcttctcttggtta taaggcatctcatctcttctgtacctcgaggtaagatgtgcaagaca
cgggaactccttatacttagctgaaggggagcggagccatcatgagtct tctcgaactgcatgtaccacatgaaactatctattacaatacgtctct
ttcaaatgagatgaacccccgcaacgacatttcggggccgaccccaac
tcagttttgaattcgggtgtttataggaatctacaggcggaggtaac atgcaaagatggattgtccaagagttccgtccattatggagagaaaa
tacagaggaaagCgacctgacctcagataaagTagtgggggtatattac
atctgcagtgccctacagatctgtatcattgtctgattgtgacattga aattcctccaggggtccaatcaaagcttactagatcaactagctatcaa
ttatctctgattgcatgacattctgtaaggaggggcgggggtagtaat catcaaagtgttgatgcaatgggatactactttcatctactcatgaa
cttgtttgctccgtgttccacaaaaggatatattctcttaattggtta tgcattgtcgaggagatatggagtggtacctggtatttgcattgggtta
cctggggcgggcctacattgtacatgaggtggtgaggatggcGaaaac
tctggtgcagcggcacgggtacgtTttgtctaaatcagatgagatcac
actgaccagggtattcacctcacagcggcagcgtgtgacagacatcct
atccagtcctttaccaagattaataaagtacttgaggaagaatattga
cactgcgctgattgaagccgggggacagcccgtccgtccattctgtgc
ggagagtctggtgagcacgctagcgaacataactcagataaccagat
Catcgctagtcacattgacacagttatccgggtctgtgatatatatgga agctgaggggtgatctcgctgacacagtatttctatttacccttaca
tctcttactgacgggaaaaagaggacatcacttaAacagtgcacgag
acagatcctagagggttacaatactaggtcttagagtcgaaaatctcaa taaaataggcgatataatcagcctagtgttaaaggcatgatctccat
ggaggaccttatcccactaaggacatacttgaagcatagtacctgccc taaatatttgaaggctgtcctaggtattaccaaaactcaaagaaatgtt
tacagacacttctgtaCtgtacttgactcgtgtcaacaaaaattcta catgaaaactataggcaatgcagtcgaaaggatattacagtaactgtga
ctcttaacgaaaatcacatattaataggctcctttttggccaattgt attctgttgatttaacatattatgtagaaaaaagttgaaccctga
ctccttaggactcgaattcgaactcaataaattgtcttaaaaaaagggt tgcgcacaattattcttgagtgtagtctcgtcattcaccaaatctttg
tttgggt Nucleic acid ATGGCCACCACCGCCATGCGCATGATCATCAGCATCAT 14 sequence
CTTCATCAGCACCTACGTGACCCACATCACCCCTGTGCCA encoding
GAACATCACCGAGGAGTTCTACCAGAGCACCTGCAGCG chimeric F
CCGTGAGTCGCGGCTACCTGAGCGCCCTGCGCACCGGC protein
TGGTACACCAGCGTGGTGACCATCGAGCTGAGCAAGAT comprising the
CCAGAAGAACGTGTGCAACAGCACCGACAGCAAGGTG ectodomain of
AAGCTGATCAAGCAGGAGCTGGAGCGCTACAACAACGC bovine RSV F
CGTGGTGGAGCTGCAGAGCCTGATGCAGAACGAGCCCG protein of strain
CCAGCTTCAGCCGCGCCAAGCGCGGCATCCCCGAGCTG ATCC51908 and

ATCCACTACCGCAACAGCACCAAGATTCTACGG the
CCTGATGGGCAAGAAGCGCAAGCGCCGCTTCCTGGGCT transmembrane
TCCTGCTGGGCATCGGCAGCGCCATCGCCAGCGGCGTG and cytoplasmic
GCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTGAA domains of
CAAGATCAAGAACGCCCTGCTGAGCACCAACAAGGCCG NDV F protein
TGGTGAGCCTGAGCAACGGCGTGAGCGTGCTGACCAGC of NDV LaSota
AAGGTGCTGGACCTGAAGAACTACATCGACAAGGAGCT strain, wherein
GCTGCCCAAGGTGAACAACCACGACTGCCGCATCAGCA the nucleotide
AGATCGAGACCGTGATCGAGTTCCAGCAGAAGAACAAC sequence
CGCCTGCTGGAGATCGCCCGCGAGTTCAGCGTGAACGC encoding the
CGGCATCACCAACCCCCCTGAGCACCTACATGCTGACCA ectodomain of
ACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCATC the RSV F
ACCAACGACCAGAAGAAGCTGATGAGCAGCAACGTGC protein is codon
AGATCGTGCGCCAGCAGAGCTACAGCATCATGAGCGTG optimized
GTGAAGGAGGAGGTGATCGCCTACGTGGTGACGCTGCC (transmembrane
CATCTACGGCGTGATCGACACCCCCTGCTGGAAGCTGC and cytoplasmic
ACACCAGCCCCCTGTGCACCACCGACAACAAGGAGGGC domains are
AGCAACATCTGCCTGACCCGCACCGATCGCGGCTGGTA underlined)
CTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCCCAGA
CCGAGACCTGCAAGGTGCAGAGCAACCGCGTGTTCTGC
GACACCATGAACAGCCTGACCCTGCCCACCGACGTGAA
CCTGTGCAACACCGACATCTTCAACACCAAGTACGACT
GCAAGATCATGACCAGCAAGACCGACATCAGCAGCAGC
GTGATCACCAGCATCGGCGCCATCGTGAGCTGCTACGG
CAAGACCAAGTGCACCGCCAGCAACAAGAATCGCGGC
ATCATCAAGACCTTCAGCAACGGCTGCGACTACGTGAG
CAACAAGGGCGTGACACCGTGAGCGTGGGCAACACCC
TGTACTACGTGAACAAGCTGGAGGGCAAGGCCCTGTAC
ATCAAGGGCGAGCCCATCATCAACTACTACGACCCCCT
GGTGTTCCCCAGCGACGAGTTCGACGCCAGCATCGCCC
AGGTGAACGCCAAGATCAACCAGAGCCTGGCCTTCATC
CGCCGCAGCGACGAGCTGCTGCACAGCGTGGACGTGGG
CAAGAGCACCAACCGTTAACCTCATTACCTATATCGT
TTTGACTATCATATCTCTTGTTTTTGGTATACTTAGCCTG
ATTCTAGCATGCTACCTAATGTACAAGCAAAAGGCGCA
ACAAAAGACCTTATTATGGCTTGGGAATAATACCCTAG
ATCAGATGAGAGCCACTACAAAAATGTGA Nucleic acid
ATGGCGACAACAGCCATGAGGATGATCATCAGCATTATCTTCATCTCTACCT 31 sequence
ATGTGACACATATCACTTTATGCCAAAACATAACAGAAGAATTTTATCAATC encoding a
AACATGCAGTGCAGTTAGTAGAGGTTACCTTAGTGCATTAAGAACTGGATG chimeric F
GTATACAAGTGTGGTAACAATAGAGTTGAGCAAAATACAAAAAATGTGTG protein
TAATAGTACTGATTCAAAAGTGAAATTAATAAAGCAAGAAGCTAGAAAGATA comprising
CAACAATGCAGTAGTGGAATTGCAGTCACTTATGCAAAATGAACCGGCCTC bovine
RSV F CTTCAGTAGAGCAAAAAGAGGGATACCAGAGTTGATACATTATACAAGAAA
protein CTCTACAAAAAAGTITTATGGGCTAATGGGCAAGAAGAGAAAAAGGAGAT
ectodomain of
TTTTAGGATTCTTGCTAGGTATTGGATCTGCTATTGCAAGTGGTGTAGCAGT bovine RSV
GTCCAAAGTACTACACCTGGAGGGAGAGGTGAATAAAATTAATAAATGCACT strain
GCTATCCACAAATAAAGCAGTAGTTAGTCTATCCAATGGAGTTAGTGTCCTT
ATCC51908 and

ACTAGCAAACCTTGATCTAAAGAACTATATAGACAAAGAGCTTCTACCTANDV F
protein AAGTTAACAATCATGATTGTAGGATATCCAAAATAGAACTGTGATAGAAT
transmembrane
TCCAACAAAAACAATAGATTGTTAGAAATTGCTAGGGAATTTAGTGTA and
cytoplasmic ATGCTGGTATTACCACACCTCTCAGTACATACATGTTGACCAATAGTGAATT
domains of the
ACTATCACTAATTAATGATATGCCTATAACGAATGACCAAAAAAAGCTAATG NDV
LaSota TCAAGTAATGTTCAAATAGTCAGGCAACAGAGTTATTCCATTATGTCAGTGG
strain TCAAAGAAGAAGTCATAGCTTATGTTGTACAATTGCCTATTTATGGAGTTAT
(transmembrane
AGACACCCCTGTTGGAACTACACACCTCTCCGTTATGCACCACTGATAAT and
cytoplasmic AAAGAAGGGTCAAACATCTGCTTAACTAGGACAGATCGTGGGTGGTATTGT
domains of NDV
GACAATGCAGGCTCTGTGTCTTTTTTCCCACAGACAGAGACATGTAAGGTA F protein
are CAATCAAATAGAGTGTTCTGTGACACAATGAACAGTTTAACTCTGCCTACTG
underlined) ACGTTAACTTATGCAACACTGACATATTCAATACAAAGTATGACTGTAAAAT
AATGACATCTAAAAGTACATAAGTAGCTCTGTGATAACTTCAATTGGAGCT
ATTGTATCATGCTATGGGAAGACAAAATGTACAGCTTCTAATAAAAAATCGT
GGAATCATAAAGACTTTTTCCAATGGGTGTGATTATGTATCAAACAAAGGA
GTAGATACTGTATCTGTTGGTAACACACTATATTATGTAAATAAGCTAGAGG
GGAAAGCACTCTATATAAAGGGTGAACCAATTATTAATTACTATGATCCACT
AGTGTTTCCTTCTGATGAGTTTGATGCATCAATTGCCCAAGTAAACGCAAAA
ATAAACCAAAGCCTGGCCTTCATACGTCGATCTGATGAGTTACTTCACAGTG
TAGATGTAGGAAAATCCACCACAAATgtaacCTCATTACCTATATCGTTTTGA
CTATCATATCTCTTGTTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCTA
ATGTACAAGCAAAAGGCGCAACAAAAGACCTTATTATGGCTTGGGAATAAT
ACCCTAGATCAGATGAGAGCCACTACAAAAATGTGA

(188) TABLE-US-00004 Nucleic acid
ATGGCCACCACCGCCATGAGAATGATCATCAGCATCATCTTCATCAGCACCT 38 sequence
ACGTGACCCACATCACCTGTGCCAGAACATCACCGAGGAGTTCTACCAGA encoding
GCACCTGCAGCGCCGTGAGCAGAGGCTACCTGAGCGCCCTGAGAACCGGC chimeric F
TGGTACACCAGCGTGGTGACCATCGAGCTGAGCAAGATCCAGAAGAACGT protein,
wherein GTGCAAGAGCACCGACAGCAAGGTGAAGCTGATCAAGCAGGAGCTGGAG
the nucleic acid
AGATACAACAACGCCGTGGTGGAGCTGCAGAGCCTGATGCAGAACGAGCC sequence
CGCCAGCTTCAGCAGAGCCAAGAGAGGCATCCCCGAGCTGATCCACTACAC
comprises a
CAGAAACAGCACCAAGAAGTTCTACGGCCTGATGGGCAAGAAGAGAAAGA codon
optimized GAAGATTCCTGGGCTTCTGCTGGGCATCGGCAGCGCCGTGGCCAGCGGC
nucleic acid GTGGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTGAACAAGATCA
sequence AGAACGCCCTGCTGAGCACCAACAAGGCCGTGGTGAGCCTGAGCAACGGC
encoding the
GTGAGCGTGCTGACCAGCAAGGTGCTGGACCTGAAGAACTACATCGACAA bovine
RSV F GGAGCTGCTGCCCCAGGTGAACAACCACGACTGCAGAATCAGCAACATCG
protein AGACCGTGATCGAGTTCCAGCAGAAGAACAACAGACTGCTGGAGATCGCC
ectodomain of
AGAGAGTTCAGCGTGAACGCCGGCATCACCACCCCCTGAGCACCTACATG strain
CTGACCAACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCATCACCAAC
ATue51908 and
GACCAGAAGAAGCTGATGAGCAGCAACGTGCAGATCGTGAGACAGCAGA a nucleic

AGCTACGACATCATGAGCGTGAGGAGGAGGTGATCGCCTACGTGGTG sequence
CAGCTGCCCATCTACGGCGTGATCGACACCCCCTGCTGGAAGCTGCACACC encoding
the AGCCCCCTGTGCACCACCGACAACAAGGAGGGCAGCAACATCTGCCTGACC
transmembrane
AGAACCGACAGAGGCTGGTACTGCGACAACGCCGGCAGCGTGAGCTTCTT and
cytoplasmic CCCCCAGACCGAGACCTGCAAGGTGCAGAGCAACAGAGTGTTCTGCGACA
domains of F
CCATGAACAGCCTGACCCTGCCCACCGACGTGAACCTGTGCAACACCGACA protein
from TCTTCAACACCAAGTACGACTGCAAGATCATGACCAGCAAGACCGACATCA
NDV strain GCAGCAGCGTGATCACCAGCATCGGCGCCATCGTGAGCTGCTACGGCAAG
LaSota ACCAAGTGCACCGCCAGCAACAAGAACAGAGGCATCATCAAGACCTTCAGC
(transmembrane
AACGGCTGCGACTACGTGAGCAACAAGGGCGTGAGACACCGTGAGCGTGG and
cytoplasmic GCAACACCCTGTACTACGTGAACAAGCTGGAGGGCAAGGCCCTGTACATCA
domains of NDV
AGGGCGAGCCCATCATCAACTACTACGACCCCCCTGGTGTTCCCCAGCGACG F are
underlined) AGTTCGACGCCAGCATCGCCCAGGTGAACGCCAAGATCAACCAGAGCCTG
GCCTTCATCAGAAGAAGCGACGAGCTGCTGCACAGCGTGGACGTGGGCAA
GAGCACCACCAACGTGgtaacCTCATTACCTATATCGTTTTGACTATCATATC
TCTTGTTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCTAATGTACAAGC
AAAAGGCGCAACAAAAGACCTTATTATGGCTTGGAATAATACCCTAGATC
AGATGAGAGCCACTACAAAAATGTGAaccgagg Nucleic acid
ATGGCCACCACCGCCATGACCATGATCATCAGCATCATCTTCATCAGCACCT 39 sequence
ACGTGACCCACATCACCTGTGCCAGAACATCACCGAGGAGTTCTACCAGA encoding
GCACCTGCAGCGCCGTGAGCAGAGGCTACCTGAGCGCCCTGAGAACCGGC chimeric F
TGGTACACCAGCGTGGTGACCATCGAGCTGAGCAAGATCCAGAAGAAGCT protein,
wherein GTGCAAGAGCACCGACAGCAAGGTGAAGCTGATCAAGCAGGAGCTGGAG
the nucleic acid
AGATACAACAACGCCGTGGTGGAGCTGCAGAGCCTGATGCAGAACGAGCC sequence
CGCCAGCTTCAGCAGAGCCAAGAGAAGCATCCCCGAGCTGATCCACTACAC
comprises a
CAGAAACAGCACCAAGAAGTTCTACGGCCTGATGGGCAAGAAGAGAAAGA codon
optimized GAAGATTCCTGGGCTTCCTGCTGGGCATCGGCAGCGCCATCGCCAGCGGCG
nucleic acid TGGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTGAACAAGATCAA
sequence GAACGCCCTGCTGAGCACCAACAAGGCCGTGGTGAGCCTGAGCAACGGCG
encoding the
TGAGCGTGCTGACCAGCAAGGTGCTGGACCTGAAGAACTACATCGACAAG bovine
RSV F GAGCTGCTGCCCAAGGTGAACAACACGACTGCAGAATCAGCAACATCGCC
protein ACCGTGATCGAGTTCCAGCAGAAGAACAACAGACTGCTGGAGATCGCCAG
ectodomain of
AGAGTTCAGCGTGAAACGCCGGCATCACACCCCCCTGAGCACCTACATGCT strain
snook and GACCAACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCATCACCAACGA
a nucleic acid
CCAGAAGAAGCTGATGAGCAGCAACGTGCAGATCGTGAGACAGCAGAGCT sequence
ACAGCATCATGAGCGTGGTGAAGGAGGAGGTGATCGCCTACGTGGTGCAG encoding
the CTGCCCATCTACGGCGTGATCGACACCCCCTGCTGGAAGCTGCACACCAGC
transmembrane
CCCCTGTGCACCACCGACAACAAGGAGGGCAGCAACATCTGCCTGACCAGA and
cytoplasmic ACCGACAGAGGCTGGTACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCC
domains of F

CAGGCGGACCTGCAAGGTGCAGAGCAAGAGTGTCTGCGACACCAT protein
from GAACAGCCTGACCCTGCCCACCGACGTGAACCTGTGCAACACCGACATCTT
NDV strain
CAACACCAAGTACGACTGCAAGATCATGACCAGCAAGACCGACATCAGCAG LaSota
CAGCGTGATCACCAGCATCGGCGCCATCGTGAGCTGCTACGGCAAGACCAA
(transmembrane
GTGCACCGCCAGCAACAAGAAGAGAGGCATCATCAAGACCTTCAGCAACG and
cytoplasmic GCTGCGACTACGTGAGCAACAAGGGCGTGGACACCGTGAGCGTGGGCAAC
domains of NDV
ACCCTGTACTACGTGAACAAGCTGGAGGGCAAGGCCCTGTACATCAAGGG F are
underlined) CGAGCCCATCATCAACTACTACGACCCCCTGGTGTTCCTCCAGCGACGAGTTC
GACGCCAGCATCGCCCAGGTGAACGCCAAGATCAACCAGAGCCTGGCCTTC
ATCAGAAGAAGCGACGAGCTGCTGCACAGCGTGGACGTGGGCAAGAGCAC
CACCAACGTGg^{tt}aacCTCATTACCTATATCGTTTTGACTATCATATCTCTTGTT
TTTGGTATACTTAGCCTGATTCTAGCATGCTACCTAATGTACAAGCAAAAAGG
CGCAACAAAAGACCTTATTATGGCTTGGAATAATACCCTAGATCAGATGA
GAGCCACTACAAAATGTGAaccg^{cg} Wild-type
ATGGCGACAACAGCCATGAGGATGATCATCAGCATTAT 40 Nucleic acid
CTTCATCTCTACCTATGTGACACATATCACTTTATGCCA encoding bovine
AAACATAACAGAAGAATTTTATCAATCAACATGCAGTG RSV F protein
CAGTTAGTAGAGGTTACCTTAGTGCATTAAGAACTGGA of strain
TGGTATACAAGTGTGGTAACAATAGAGTTGAGCAAAAT ATue51908 (the
ACAAAAAAATGTGTGTAAAAGTACTGATTCAAAGTGA transmembrane
AATTAATAAAGCAAGAACTAGAAAGATACAACAATGC and cytoplasmic
AGTAGTGGAATTGCAGTCACTTATGCAAAATGAACCGG domains are
CCTCCTTCAGTAGAGCAAAAAGAGGGATACCAGAGTTG underlined)
ATACATTATACAAGAACTCTACAAAAAAGTTTTATGG
GCTAATGGGCAAGAAGAGAAAAAGGAGATTTTTAGGAT
TCTTGCTAGGTATTGGATCTGCTGTTGCAAGTGGTGTAG
CAGTGTCCAAAGTACTACACCTGGAGGGAGAGGTGAAT
AAAATTAAAAATGCACTGCTATCCACAAATAAAGCAGT
AGTTAGTCTATCCAATGGAGTTAGTGTCTTACTAGCAA
AGTACTTGATCTAAAGAAGTATATAGACAAAGAGCTTC
TACCTCAAGTTAACAATCATGATTGTAGGATATCCAAC
ATAGAACTGTGATAGAATTCCAACAAAAAACAATAG
ATTGTTAGAAATTGCTAGGGAATTTAGTGTAATGCTG
GTATTACCACACCTCTCAGTACATACATGTTGACCAATA
GTGAATTACTATCACTAATTAATGATATGCCTATAACGA
ATGACCAAAAAAAGCTAATGTCAAGTAATGTTCAAATA
GTCAGGCAACAGAGTTATTCCATTATGTCAGTGGTCAA
AGAAGAAGTCATAGCTTATGTTGTACAATTGCCTATTTA
TGGAGTTATAGACACCCCCTGTTGGAAACTACACACCT
CTCCGTTATGCACCACTGATAATAAAGAAGGGTCAAAC
ATCTGCTTAACTAGGACAGATCGTGGGTGGTATTGTGA
CAATGCAGGCTCTGTGTCTTTTTTCCCACAGACAGAGAC
ATGTAAGGTACAATCAAATAGAGTGTTCTGTGACACAA
TGAACAGTTTAACTCTGCCTACTGACGTTAACTTATGCA
ACACTGACATATTCAATACAAAGTATGACTGTAAAATA
ATGACATCTAAAAGTACATAAGTAGCTCTGTGATAAC
TTCAATTGGAGCTATTGTATCATGCTATGGGAAGACAA

AATGTAATCAATAAATCGTGGAATCATAAAG
ACTTTTTCCAATGGGTGTGATTATGTATCAAACAAAGGA
GTAGATACTGTATCTGTTGGTAACACACTATATTATGTA
AATAAGCTAGAGGGGAAAGCACTCTATATAAAGGGTGA
ACCAATTATTAATTACTATGATCCACTAGTGTTTCCTTC
TGATGAGTTTGATGCATCAATTGCCCAAGTAAACGCAA
AAATAAACCAAAGCCTGGCCTTCATACGTCGATCTGAT
GAGTTACTTCACAGTGTAGATGTAGGAAAATCCACCAC
AAATGTAGTAATTACTACTATTATCATAGTGATAGTTGT
AGTGATATTAATGTTAATAGCTGTAGGATTACTGTTTTA
CTGTAAGACCAAGAGTACTCCTATCATGTTAGGGAAGG
ATCAGCTCAGTGGTATCAACAATCTTTCCTTTAGTAAAT GA Codon
ATGGCCACCACCGCCATGAGAATGATCATCAGCATCAT 41 optimized
CTTCATCAGCACCTACGTGACCCACATCACCTGTGCCA nucleic acid
GAACATCACCGAGGAGTTCTACCAGAGCACCTGCAGCG sequence
CCGTGAGCAGAGGCTACCTGAGCGCCCTGAGAACCGGC encoding bovine
TGGTACACCAGCGTGGTGACCATCGAGCTGAGCAAGAT RSV F protein
CCAGAAGAACGTGTGCAAGAGCACCGACAGCAAGGTG of strain
AAGCTGATCAAGCAGGAGCTGGAGAGATACAACAACG ATue51908
CCGTGGTGGAGCTGCAGAGCCTGATGCAGAACGAGCCC
GCCAGCTTCAGCAGAGCCAAGAGAGGCATCCCCGAGCT
GATCCACTACACCAGAAACAGCACCAAGAAGTTCTACG
GCCTGATGGGCAAGAAGAGAAAGAGAAGATTCCTGGG
CTTCCTGCTGGGCATCGGCAGCGCCGTGGCCAGCGGCG
TGGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTG
AACAAGATCAAGAACGCCCTGCTGAGCACCAACAAGGC
CGTGGTGAGCCTGAGCAACGGCGTGAGCGTGCTGACCA
GCAAGGTGCTGGACCTGAAGAACTACATCGACAAGGAG
CTGCTGCCCCAGGTGAACAACCACGACTGCAGAATCAG
CAACATCGAGACCGTGATCGAGTTCCAGCAGAAGAACA
ACAGACTGCTGGAGATCGCCAGAGAGTTCAGCGTGAAC
GCCGGCATCACCAACCCCCCTGAGCACCTACATGCTGAC
CAACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCCA
TCACCAACGACCAGAAGAAGCTGATGAGCAGCAACGTG
CAGATCGTGAGACAGCAGAGCTACAGCATCATGAGCGT
GGTGAAGGAGGAGGTGATCGCCTACGTGGTGCAGCTGC
CCATCTACGGCGTGATCGACACCCCCTGCTGGAAGCTG
CACACCAGCCCCCTGTGCACCACCGACAACAAGGAGGG
CAGCAACATCTGCCTGACCAGAACCGACAGAGGCTGGT
ACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCCCAG
ACCGAGACCTGCAAGGTGCAGAGCAACAGAGTGTTCTG
CGACACCATGAACAGCCTGACCCTGCCCACCGACGTGA
ACCTGTGCAACACCGACATCTTCAACACCAAGTACGAC
TGCAAGATCATGACCAGCAAGACCGACATCAGCAGCAG
CGTGATCACAGCATCGGCGCCATCGTGAGCTGCTACG
GCAAGACCAAGTGCACCGCCAGCAACAAGAACAGAGG
CATCATCAAGACCTTCAGCAACGGCTGCGACTACGTGA
GCAACAAGGGCGTGACACCGTGAGCGTGGGCAACAC
CCTGTACTACGTGAACAAGCTGGAGGGCAAGGCCCTGT
ACATCAAGGGCGAGCCCATCATCAACTACTACGACCCC

CTGGTGTTCGAGCCAGAGTTCGACGCCAGCATCGC
CCAGGTGAACGCCAAGATCAACCAGAGCCTGGCCTTCA
TCAGAAGAAGCGACGAGCTGCTGCACAGCGTGGACGTG
GGCAAGAGCACCACCAACGTGGTGATCACCACCATCAT
CATCGTGATCGTGGTGGTGCCTGATGCTGATCGCCGT
GGGCCTGCTGTTCTACTGCAAGACCAAGAGCACCCCCA
TCATGCTGGGCAAGGACCAGCTGAGCGGCATCAACAAC CTGAGCTTCAGCAAG Wild-
type ATGGCGACAACAGCCATGACGATGATCATCAGCATTAT 42 Nucleic acid
CTTCATCTCTACCTATGTGACACATATCACTTTATGCCA sequence
AAACATAACAGAAGAATTTTATCAATCAACATGCAGTG encoding bovine
CAGTTAGTAGAGGTTACCTTAGTGCATTAAGAAGTGG A RSV F protein
TGGTATACAAGTGTGGTAACAATAGAGTTGAGCAAAAT of strain snook
ACAAAAAATGTGTGTAAAAGTACTGATTCGAAAGTGA (transmembrane
AATTAATAAAGCAAGAAGTACTAGAAAGATACAACAATGC and cytoplasmic
AGTAGTGGAATTGCAGTCACTTATGCAAAATGAACCGG domains are
CCTCCTTCAGTAGAGCAAAAAGAAGTATACCAGAGTTG underlined)
ATACATTATACAAGAACTCTACAAAAAAGTTTTATGG
GCTAATGGGCAAGAAGAGAAAAAGGAGATTTTTAGGAT
TCTTACTAGGTATTGGATCTGCTATTGCAAGTGGTGTAG
CAGTGTCCAAAGTACTACACCTGGAGGGAGAGGTGAAT
AAAATTAAAAATGCACTGCTATCCACAAACAAAGCAGT
AGTTAGTCTATCCAATGGAGTTAGTGTCTTACTAGCAA
AGTACTTGATCTAAAGAAGTATATAGACAAAGAGCTTC
TACCTAAAGTTAACAATCATGATTGTAGGATATCCAAC
ATAGCAACTGTGATAGAATTCCAACAAAAAACAATAG
ATTGTTAGAAATTGCTAGGGAATTTAGTGTAATGCTG
GTATTACCACACCCCTCAGTACATACATGTTGACCAATA
GTGAATTACTATCACTAATTAATGATATGCCTATAACGA
ATGACCAAAAAAAGCTAATGTCAAGTAATGTTCAAATA
GTCAGGCAACAGAGTTATTCCATTATGTCAGTGGTCAA
AGAAGAGGTCATAGCTTATGTTGTACAATTGCCTATTTA
TGGAGTTATAGACACCCCCTGTTGGAACTACACACTTC
TCCATTATGCACCACTGATAATAAAGAAGGGTCAAACA
TCTGCTTAACTAGGACAGATCGTGGGTGGTATTGTGAC
AATGCAGGCTCTGTATCTTTTTTCCCACAGGCAGAGACG
TGTAAGGTACAATCAAATAGAGTGTTCTGTGACACAAT
GAACAGTTTAACTCTGCCTACTGATGTTAACTTATGCAA
CACTGACATATTCAATACAAAGTATGACTGTAAAATAA
TGACATCTAAAAGTACATAAGTAGCTCTGTAATAACTT
CAATTGGAGCTATTGTATCATGCTATGGGAAGACAAAA
TGTACAGCTTCTAATAAAAAATCGTGGAATCATAAAGAC
TTTTTCCAATGGGTGTGATTATGTATCAAACAAAGGAGT
TGATACTGTATCTGTTGGTAACACACTATATTATGTAAA
TAAGCTAGAGGGGAAAGCACTCTATATAAAGGGTGAAC
CAATTATTAATTACTATGATCCACTAGTGTTTCCTTCTG
ATGAGTTTGTATGCATCAATTGCCCAAGTAAACGCAAAA
ATAAACCAAAGCCTGGCTTTCATACGTCGATCTGATGA
GTTACTTCACAGTGTAGATGTAGGAAAATCCACCACAA
ATGTAGTAATTACTACTATTATCATAGTGATAGTTGTAG
TGATATTAATGTTAATAGCTGTAGGATTACTGTTTTACT

GTAAGACCAAGTACTCCTATCATGTTAGGGAAGGAT
CAGCTTAGTGGTATCAACAATCTTTCCTTTAGTAAATGA Codon
ATGGCCACCACCGCCATGACCATGATCATCAGCATCAT 43 optimized
CTTCATCAGCACCTACGTGACCCACATCACCTGTGCCA nucleic acid
GAACATCACCGAGGAGTTCTACCAGAGCACCTGCAGCG sequence
CCGTGAGCAGAGGCTACCTGAGCGCCCTGAGAACCGGC encoding bovine
TGGTACACCAGCGTGGTGACCATCGAGCTGAGCAAGAT RSV F protein
CCAGAAGAACGTGTGCAAGAGCACCGACAGCAAGGTG of strain snook
AAGCTGATCAAGCAGGAGCTGGAGAGATACAACAACG
CCGTGGTGGAGCTGCAGAGCCTGATGCAGAACGAGCCC
GCCAGCTTCAGCAGAGCCAAGAGAAGCATCCCCGAGCT
GATCCACTACACCAGAAACAGCACCAAGAAGTTCTACG
GCCTGATGGGCAAGAAGAGAAAGAGAAGATTCTCTGGG
CTTCCTGCTGGGCATCGGCAGCGCCATCGCCAGCGGCG
TGGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGTG
AACAAGATCAAGAACGCCCTGCTGAGCACCAACAAGGC
CGTGGTGAGCCTGAGCAACGGCGTGAGCGTGCTGACCA
GCAAGGTGCTGGACCTGAAGAACTACATCGACAAGGAG
CTGCTGCCCAAGGTGAACAACCACGACTGCAGAATCAG
CAACATCGCCACCGTGATCGAGTTCCAGCAGAAGAACA
ACAGACTGCTGGAGATCGCCAGAGAGTTCAGCGTGAAC
GCCGGCATCACCAACCCCCCTGAGCACCTACATGCTGAC
CAACAGCGAGCTGCTGAGCCTGATCAACGACATGCCCA
TCACCAACGACCAGAAGAAGCTGATGAGCAGCAACGTG
CAGATCGTGAGACAGCAGAGCTACAGCATCATGAGCGT
GGTGAAGGAGGAGGTGATCGCCTACGTGGTGCAGCTGC
CCATCTACGGCGTGATCGACACCCCCCTGCTGGAAGCTG
CACACCAGCCCCCTGTGCACCACCGACAACAAGGAGGG
CAGCAACATCTGCCTGACCAGAACCGACAGAGGCTGGT
ACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCCCAG
GCCGAGACCTGCAAGGTGCAGAGCAACAGAGTGTTCTG
CGACACCATGAACAGCCTGACCCTGCCCACCGACGTGA
ACCTGTGCAACACCGACATCTTCAACACCAAGTACGAC
TGCAAGATCATGACCAGCAAGACCGACATCAGCAGCAG
CGTGATCACCAAGCATCGGCGCCATCGTGAGCTGCTACG
GCAAGACCAAGTGACCGCCAGCAACAAGAACAGAGG
CATCATCAAGACCTTCAGCAACGGCTGCGACTACGTGA
GCAACAAGGGCGTGACACCGTGAGCGTGGGCAACAC
CCTGTACTACGTGAACAAGCTGGAGGGCAAGGCCCTGT
ACATCAAGGGCGAGCCCATCATCACTACTACGACCCC
CTGGTGTTCCCCAGCGACGAGTTCGACGCCAGCATCGC
CCAGGTGAACGCCAAGATCAACCAGAGCCTGGCCTTCA
TCAGAAGAAGCGACGAGCTGCTGCACAGCGTGGACGTG
GGCAAGAGCACCAACCGTGGTGATCACCAACCATCAT
CATCGTGATCGTGGTGGTGATCCTGATGCTGATCGCCGT
GGGCCTGCTGTTCTACTGCAAGACCAGAAGCACCCCCA
TCATGCTGGGCAAGGACCAGCTGAGCGGCATCAACAAC CTGAGCTTCAGCAAG
Amino acid MATTAMRMIISIIIFISTYVTHITLCQNITEEFYQSTCSAVSRG 33 sequence
YLSALRTGWYTSVVTIELSKIQKNVCNSTDSKVKLIKQELE encoding
RYNNAVVELQSLMQNEPASFSRAKRGIPELIHYTRNSTKK chimeric F

FYGLMGKRRFLGFLGSAIASGVAVSKVLHLEGEV protein
NKIKNALLSTNKAVVSLNNGVSVLTskVLDLKNYIDKELL comprising the
PKVNNHDCRISKIETVIEFQQKNNRLLLEIAREFSVNAGITTP ectodomain of
LSTYMLTNSELLSLINDMPITNDQKKLMSSNVQIVRQQSYS bovine RSV F
IMSVVKEEVIAYVVQLPIYGVIDTPCWKLHTSPLCTTDNKE protein of bovine
GSNICLTRDRGWYCDNAGSVSFFPQTETCKVQSNRVFCD RSV strain
TMNSLTLPDVLNLCNTDIFNTKYDCKIMTSKTDISSSVITSI ATCC51908 and
GAIVSCYGKTKCTASNKNRGIKTFsNGCDYVSNKGVDTV the
SVGNTLYYVnkLEGKALYIKGEPIINYDPLVFPSDEFDAS transmembrane
IAQVNAKINQSLAFIRRSDELLHSVDVGKSTTNVNLITYIVL and cytoplasmic
TIISLVFGILSLILACYLMYKQKAQQKTLLWLGNNTLDQM domains of NDV RATTKM
F protein of NDV LaSota strain (transmembrane and cytoplasmic domains of NDV F
protein are underlined) Genome of
accaaacagagaatccgtgagttacgataaaaggcgaaggagcaattgaagtcgcacggg 37 NDV LaSota
tagaaggtgtgaatctcgagtgcgagcccgaagcacaactcgagaaagccttctgccaac with a chimeric
atgtcttccgtatttgatgagtacgaacagctcctcgcggtcagactcgcccaatggagct F protein
catggagggggagaaaaaggagtagctttaaagtagacgtcccgggtattcactcttaaca (underlined).
gtgatgaccagaagatagatggagctttgtggtattctgctcgggattgctgtagcgaag Ectodomain is
atgccaacaaaccactcaggcaaggtgctctcatatctctttatgctccactcacaggtaat codon optimized
gaggaacctgttgccCttgcagggaacagaatgaagccacattggccgtgcttgagatt from bovine
gatggctttgccaacggcagccccagttcaacaataggagtggagtgtctgaagagagag RSV strain
cacagagatttgcgatgatagcaggatctctccctcgggcatgcagcaacggaaccccgctc ATCC51908.
gtcacagccggggcGgaagatgatgcaccagaagacatcaccgataccctggagaggat Transmembrane
cctctctatccaggctcaagtatgggtcacagtagcaaaagccatgactgcgtatgagactg and cytoplasmic
cagatgagtcggaacaaggcgaatcaataagtatatgcagcaaggcagggtccaaaaga domains are
aatacatcctctaccccgatgcaggagcacaatccaactcacgatcagacagtctcttgag from the F
tccgcattcttttggttagcgagctcaagagaggccgcaacacggcagggtggtacctctactt protein of NDV
attataacctggtaggggacgtagactcatacatcaggaataaccgggcttactgcattcttctt strain LaSota
gacactcaagtacggaatcaacaccaagacatcagcccttgacttagtagcctctcaggcg
acatccagaagatgaagcagctcatgcgtttgtatcggtatgaaaggagataatgcgccgtac
atgacattacttggtagatgaccagatgagctttgcgcctgccgagtagcacaactttact
cctttgccatgggtatggcatcagctctagataaaaggtagtgggaaataccaatttgccaggg
actttatgagcacatcattctggagacttggagtagagtacgctcagggtcagggaagtagc
attaacgaggatattggctgccgagctaaagctaaccagcagcaaGgaGgggcctggc
agctgctgccaacgggtctccgaGgaGaccagcagcataGacatgcctactcaacaag
tcggagtcctcactgggcttagcgaggggggggtcccaagctctacaaggcggatcgaata
gatcgcaagggaaccagaagccggggatggggagaccaattcctggatctgatgaga
gcggtagcaaatagcatgagggaggcgccaaactctgcacagggcactcccaatcggg
gcctccccaaactcctgggccatccaagataacgacaccgactgggggtattgatggaca
aaaccagcctgcttcacaaaaacatcccaatgcctcaccgtagtcgacctcgatttg
cggctctatatgaccacacctcaaaaaacatccccctcttctccctccccctgctgtaca
actAcgTacgccttagataccacaggcacaatcggtctactaacaatcaaaacagagc
cgagggaattagaaaaaagtacgggtagaagagggatattcagagatcagggaaggtctc
ccgagtctctgctctctctctacctgatagaccaggacaaacatggccacctttacagatgc
agagatcgacgagctatttgagacaagtgggaactgtcattgacaacataattacagcccagg
gtaaaccagcagagactgttggaaggagtgaatcccacaaggcaagaccaaggtgctga
gcgcagcatgggagaagcatgggagcatccagccaccggccagtcaagacaaccccgat
cgacaggacagatctgacaaacaacctccacacccgagcaaacgaccccgcatgacag
cccgcggccacatccgaccagccccccaccaggccacagacgaagccgctgcac
acacagCtcaggaccggagcaagcaactctctgctgttgatgcttgacaagctcagcaata

aatcgtccaatgctaataaggcccatggtcgagcccccaagaggggaatcaccaacgtc
cgactcaacagcaggggagtcacccagtcgcggaacagtcaggaaagaccgcagaa
ccaagtcaaggccgcccctggaaaccagggcacagacgtgaacacagcatatcatggac
aatgggaggagtcacaactatcagctggtgcaaccctcatgctctccgatcaaggcagag
ccaagacaataacccttgatctgcggatcatgtccagccacctgtagactttgtgcaagcgat
gatgtctatgatggaggcgatatcacagagagtaagtaagggtgactatcagctagatctgtc
ttgaaacagacatcctccatccctatgatgcggtccgaaatccaacagctgaaaacatctgtt
gcagtcatggaagccaacttggaatgatgaagattctggatcccgggttgccaacatttca
tctctgagtgatctacgggcagttgcccgatctcacccggttttagtttcaggccctggagacc
cctctccctatgtgacacaaggaggcgaaatggcacttaataaactttcgcaaccagtgcca
catccatctgaattgattaaacccgccactgcatgcgggcctgatataggagtggaaaagga
cactgtccgtgcattgatcatgtcacgccaatgcacccgagttcttcagccaagctcctaag
caagttagatgcagccgggtcgatcgaggaaatcaggaaaatcaagcgccttgctctaaat
ggctaattactactgccacacgtagcgggtccctgtccactcggcatcacacggaatctgca
ccgagttccccccgcGgTTAGAAAAAATACGGGTAGAACCGCC
ACCATGGCCACCACCGCCATGCGCATGATCATCAGCAT
CATCTTCATCAGCACCTACGTGACCCACATCACCCCTGTG
CCAGAACATCACCGAGGAGTTCTACCAGAGCACCTGCA
GCGCCGTGAGTCGCGGCTACCTGAGCGCCCTGCGCACC
GGCTGGTACACCAGCGTGGTGACCATCGAGCTGAGCAA
GATCCAGAAGAACGTGTGCAACAGCACCGACAGCAAG
GTGAAGCTGATCAAGCAGGAGCTGGAGCGCTACAACAA
CGCCGTGGTGGAGCTGCAGAGCCTGATGCAGAACGAGC
CCGCCAGCTTCAGCCGCGCCAAGCGCGGCATCCCCGAG
CTGATCCACTACACCCGCAACAGCACCAAGAAGTTCTA
CGGCCTGATGGGCAAGAAGCGCAAGCGCCGCTTCCTGG
GCTTCCTGCTGGGCATCGGCAGCGCCATCGCCAGCGGC
GTGGCCGTGAGCAAGGTGCTGCACCTGGAGGGCGAGGT
GAACAAGATCAAGAACGCCCTGCTGAGCACCAACAAG
GCCGTGGTGAGCCTGAGCAACGGCGTGAGCGTGCTGAC
CAGCAAGGTGCTGGACCTGAAGAACTACATCGACAAGG
AGCTGCTGCCCAAGGTGAACAACACGACTGCCGCATC
AGCAAGATCGAGACCGTGATCGAGTTCCAGCAGAAGAA
CAACCGCCTGCTGGAGATCGCCCGCGAGTTCAGCGTGA
ACGCCGGCATCACCACCCCCTGAGCACCTACATGCTG
ACCAACAGCGAGCTGCTGAGCCTGATCAACGACATGCC
CATCACCAACGACCAGAAGAAGCTGATGAGCAGCAAC
GTGCAGATCGTGCGCCAGCAGAGCTACAGCATCATGAG
CGTGGTGAAGGAGGAGGTGATCGCCTACGTGGTGCAGC
TGCCCATCTACGGCGTGATCGACACCCCCTGCTGGAAG
CTGCACACCAGCCCCCTGTGCACCACCGACAACAAGGA
GGGCAGCAACATCTGCCTGACCCGCAACCGATCGCGGCT
GGTACTGCGACAACGCCGGCAGCGTGAGCTTCTTCCCC
CAGACCGAGACCTGCAAGGTGCAGAGCAACCGCGTGTT
CTGCGACACCATGAACAGCCTGACCCTGCCCACCGACG
TGAACCTGTGCAACACCGACATCTTCAACACCAAGTAC
GACTGCAAGATCATGACCAGCAAGACCGACATCAGCAG
CAGCGTGATCACAGCATCGGCGCCATCGTGAGCTGCT
ACGGCAAGACCAAGTGCACCGCCAGCAACAAGAATCG
CGGCATCATCAAGACCTTCAGCAACGGCTGCGACTACG

TGAGCAACAAGGGCGTGACACCGTGAGCGTGGGCAA
CACCCTGTACTACGTGAACAAGCTGGAGGGCAAGGCC
TGTACATCAAGGGCGAGCCCATCATCACTACTACGAC
CCCCTGGTGTTCGCCAGCGACGAGTTCGACGCCAGCAT
CGCCCAGGTGAACGCCAAGATCAACCAGAGCCTGGCCT
TCATCCGCCGCAGCGACGAGCTGCTGCACAGCGTGGAC
GTGGGCAAGAGCACCAACGTTAACCTCATTACCTA
TATCGTTTTGACTATCATATCTCTTGTTTTTGGTATACTT
AGCCTGATTCTAGCATGCTACCTAATGTACAAGCAAAA
GGCGCAACAAAAGACCTTATTATGGCTTGGGAATAATA
CCCTAGATCAGATGAGAGCCACTACAAAAATGTGAGCT
TCAGCAAGTAAccccccgcggaaccaaggtccaactctccaagcggcaatcctc
tctcgcttcctcagccccactgaatgAtcgcgtaaccgtaattaatctagctacatttaagatta
agaaaaaatacgggtagaattggagtgcccaattgtgccaagatggactcatctaggacaa
ttgggctgtactttgattctgcccattctttagcaacctgttagcatttccgatcgctctacaag
Acacaggagatgggaagaagcaaactgccccgcaatataggatccagcgccttgacttgt
ggactgatagtaaggaggactcagtattcatcaccacctatggattcatctttcaagttgggaa
tgaagaagccacCgtcggcatgatcgatgataaaccgaagcgcgagttactttccgctgcg
atgctctgcctaggaagcgtcccaataaccggagaccttattgagctggcaagggcctgtct
cactatgatagtcacatgcaagaagagtgaactaatactgagagaatggttttctcagtagtg
caggcaccccaagtgtgcaaaagctgtaggggtgtggcaaacaatactcatcagtgaatgc
agtcaagcacgtgaaagcgccagagaagattcccgggagtggaaccctagaatacaaggt
gaactttgtctccttgactgtggtaccgaagaGggatgtctacaagatcccagctgcagtatt
gaaggtttctggctcgagtctgtacaactcttgcgctcaatgtcactattaatgtggaggtagac
ccgaggagtcctttggttaaactctCtgtctaagtctgacagcggatactatgctaaccttctt
gcatattggacttatgaccacTgtagataggaaggggaagaaagtacatttgacaagctg
gaaaagaaaataaggagccttgatctatctgtcgggctcagtgatgtgctcgggccttccgtg
ttggtaaaagcaagaggtgcacggactaagcttttggcaccttcttcttagcagtgggaca
gcctgctatcccatagcaaagtcttctcctcaggtggccaagatactctggagtcaaaccgcg
tgcctgcgagcgttaaaatcattatccaagcaggtaccaacgcgctgtcgcagtgaccgc
cgaccacgaggttaccttactaagctggagaaggggcacacccttgccaaatacaatcctt
ttaagaaataagctgcgtctctgagattgcgctccgcccactcaccagatcatcatgacaca
aaaaactaatctgtcttgattattacagttagttacctgtctatcaagttagaaaaaacacggg
tagaagattctggatcccgggtggcgccctccaggtgcaagatgggctccagaccttctacc
aagaaccagcacctatgatgtgactatccgggttgcgctggtagtgactgagttgcatctgtccg
gcaaactccattgatggcaggcctcttgagctgcaggaattgtggttacaggagacaaagc
cgtcaacatatacctcatcccagacaggatcaatcatagttaagctcctcccgaatctgcc
caaggataaggaggcatgtgcgaaagcccccttgatgcatacaacaggacattgaccact
ttgtcaccccccttggtgactctatccgtaggatacaagagtctgtgactacatctggaggg
gggagacagggggcgcttataggcgccattattggcggtgtggctcttgggggtgcaactgc
cgcacaaataacagcggccgcagctctgatacaagccaaacaaaatgctgccaacatcctc
cgacttaaagagagcattgccgcaaccaatgaggctgtgcatgaggtcactgacggattatc
gcaactagcagtggtcagttgggaagatgcagcagtttgtaatgaccaatttaataaaacagc
tcaggaattagactgcatcaaaattgcacagcaagttggtgtagagctcaacctgtacctaac
cgaattgactacagtattcggaaccacaaatcacttcacctgcttaacaagctgactattcag
gcactttacaatctagctgggtggaaatatggattacttattgactaagttaggtgtagggaaca
tcaactcagctcattaatcggtagcggcttaatcaccggtaaccctattctatacgaactcacag
actcaactcttgggtatacaggttaactctaccttcagtcgggaacctaaataatatgcgtgcc
cctacttggaaaccttatccgtaagcacaaccaggggatttgcctcggcacttgtcccAaaa
gtgggtgacacaggtcgggtctgtgatagaagaacttgacacctcatactgtatagaaactgac

ttagatttatattgtacaagaatagtaacgttccctatgtcccctggattatttcttgccttgagcg
gcaatacgtcggcctgtatgtactcaaagaccgaaggcgcacttactacaccatacatgact
atcaaaggttcagtcacgcgaactgcaagatgacaacatgtagatgtgtaaacccccggg
tatcatatcgaaaactatggagaagccgtgtctctaatagataaacaatcatgcaatgtttat
ccttaggcgggataactttaaggctcagtggggaattcgatgtaacttatcagaagaatatctc
aatacaagattctcaagtaataataacaggcaattctgatatctcaactgagcttgggaatgtc
aacaactcgatcagtaatgcttgaataagtttagaggaaagcaacagaaaactagacaaagt
caatgtcaaaactgactagcacatctgctctcattacctatatcgtttgactatcatatcttgtttt
tggtatacttagcctgattctagcatgctacctaattgtacaagcaaaaggcgcaacaaaagac
cttattatggcttgggaataatactctagatcagatgagagccactacaaaaatgtgaacaca
gatgaggaacgaagggttccctaataagtaattgtgtgaaagttctggtagtctgtcagttcaga
gagttaagaaaaaactaccgggttagatgaccaaaggacgatatacgggtagaacggtaa
gagaggccgcccctcaattgcgagccaggcttcacaacctccgttctaccgcttcaccgaca
acagtcctcaatcatggaccgcgccgttagccaagttgcgtagagaatgatgaaagagag
gcaaaaaatacatggcgcttgatattccggattgcaatcttattcttaacagtagtgaccttggc
tatatctgtagcctcccttttatatagcatgggggctagcacacctagcgatctttaggcatac
cgactaggatttccagggcagaagaaaagattacatctacacttgggtccaatcaagatgtag
tagataggatatataagcaagtggcccttgagtctccgttggcattgttaaatactgagaccac
aattatgaacgcaataacatctctctcttatcagattaatggagctgcaacaacagtggtgg
ggggcacctatccatgaccagattatataggggggataggcaaagaactcattgtagatga
tgctagtgatgtcacatcattctatccctctgcatttcaagaacatctgaatttatcccggcgcc
tactacaggatcaggttgcaactcgaataccctcattgacatgagtgtaccattactgtctac
accataatgtaataattgtctggatgcagagatcactcacattcatatcagtatttagcacttgg
gtgctccggacatctgcaacaggggaggggtattcttttctactctgcgttccatcaacctggacg
acacccaaaatcggaagtcttgagtgtagtgcaactcccctgggttgtagatgctgtgct
cgaaagtcacggagacagaggaagaagattataactcagctgtccctacgcggatggtaca
tgggaggttaggggtcgacggccagtagccacgaaaaggacctagatgtcacaacattattcg
gggactgggtggccaactaccagggagtaggggggtggatctttattgacagccgcgtatg
gttctcagttctacggaggggttaaaacccaattcacccagtgacactgtacaggaagggaat
atgtgatatacaagcgatacaatgacacatgccagatgagcaagactaccagattcgaatg
gccaagtcttcgtataagcctggacgggttggtgggaaacgcatacagcaggctatcttatct
atcaaggtgtcaacatccttaggcgaagacccggtactgactgtaccgccaacacagtcac
actcatgggggcccgaaggcagaattctcacagtagggacatctcatttctgtatcaacgagg
gtcatcatacttctctccgcgttattatatcctatgacagtcagcaaaaaacagccactcttc
atagtccttatacattcaatgccttcactcggccaggtagtagtcccttgccaggcttcagcaag
atgccccaaactcgtgtgttactggagttatatacagatccatatcccctaattcttatagaaacc
acaccttgcgaggggtattcgggacaatgcttgatgggtgtacaagcaagacttaacctgcg
tctgcagtattcgatagcacatcccgcagtcgcattactcgagtgagttcaagcagtagcaaaa
gcagcatacacaacatcaactgttttaaagtgggtcaagactaataagacctattgtctcagcat
tgctgaaatatctaactctcttcggagaattcagaatcgccccgttactagttgagatcctcaa
agatgacgggggttagagaagccaggtctggctagttgagtcaattataaaggagttggaag
atggcattgtatcacctatcttctgcgacatcaagaatcaaaccgaatgccggcgcggtgctcg
aattccatgttgccagttgaccacaatcagccagtgctcatgcgatcagattaagccttgtca
AtaGtctcttgattaagaaaaaatgtaagtggcaatgagatacaaggcaaaacagctcatg
gtTaaCaatacgggtaggacatggcgagctccggtcctgaaagggcagagcatcagatta
tcctaccagagTcacacctgtcttcaccattgggtcaagcacaactactctattactggaaatt
aactgggctaccgcttctgatgaatgtgacttcgaccacctattctcagccgacaatggaa
aaaaatacttgaatcggcctctcctgatactgagagaatgataaaactcggaagggcagtag
accaaactcttaaccacaattccagaataaccggagtgctccaccccaggtgtttagaaGaa
ctggctaataattgaggtcccagattcaaccaacaaatttcggaagattgagaagaagatccaa

attcacaacacgagatatggagaactgttcacaaggctgtgtacgcatatagagaagaaact
gctgggggtcatcttgggtctaacaatgtcccccggtcagaggagttcagcagcattcgtacgg
atccggcattctggtttcactcaaaatggtccacagccaagtttgcattggctccatataaaaca
gatccagaggcatctgatgggtggcagctaGgacaaggctgtcggccaacaaattgggtgat
gctaaccataaggtaggccaagtctttgtcactcctgaacttgcgttgtagcgcatacgaat
gagaacaagttcatatgtcttaccaggaactgtattgatgtatgcagatatgatggagggc
agagatatgggtcaacataatatcaaccacggcggtgcatctcagaagcttatcagagaaaatt
gatgacattttgcggttaatagacgctctggcaaaagacttgggtaataagcttacgatgtg
tatcactaatggagggatttgcatacggagctgtccagctactcgagccgtcaggtacatttg
caggagattttcttcgattcaacctgcaggagcttaaagacattctaattggcctcctcccaa
tgatatagcagaatccgtgactcatgcaatcgctactgtattctctggttagaacagaatcaag
cagctgagatgttgtgtctgttgcgtctgtggggtcacccactgcttgagtcccgtattgcagc
aaaggcagtcaggagccaaatgtgcgcaccgaaaatggtagactttgatatgaccttcagg
tactgtctttctcaagggaacaatcatcaacgggtacagaaagaagaatgcaggtgtgtggc
cgcgagtcaaagtggatacaatatatgggaagggtcattggggcaactacatgcagattcagca
gagatttcacacgatatcatgttgagagagtataagagtttatctgcactgaatttgagccatg
tatagaatatgaccctgtcaccaacctgagcatgttctaaaagacaagggaatcgcacacc
ccaacgataattggcttgctcgtttaggcggaaccttctctccgaagaccagaagaaacatg
taaaagaagcaacttcgactaatcgctcttgatagagtttttagagtcaaataatttgatccat
ataaagagatggaatatctgacgacccttgagtaccttagagatgacaatgtggcagtatcat
actcgtcaaggagaaggaaagtgaagttaatggacggatcttcgctaagctgacaaagaa
gttaaggaaactgtcaggtgatggcggaagggtaccttagccgatcagattgcacctttcttca
gggaaatggagtcattcaggatagcatatccttgaccaagagtatgctagcgatgagtcaact
gtcttttaacagcaataagaaacgtatcactgactgtaaagaaagagtatctcaaaccgcaat
catgatccgaaaagcaagaaccgtcgggagagttgcaaccttcataacaactgacctgcaaa
agtactgtcttaattggagatatcagacaatcaaattgttcgctcatgccatcaatcagttgatg
ggcctacctcacttctcgaatggattcacctaagactgatggacactacgatgttcgtaggag
accctttcaatcctccaagtgaccttactgactgtgaccttcaagagtcctaataatgatgacat
atatattgtcagtgccagaggggggtatcgaaggattatgccagaagctatggacaatgatctc
aattgctgcaatccaacttgctgcagctagatcgcttgcgtgttgctgtatggtacaggggt
gataatcaagtaatagcagtaacgagagaggttaagatcagacgactctccggagatgggtgt
gacacagttgcatcaagccagtataatttctcaagggaattaattcatgtcaatcatttgattgg
ccataatttgaaggatcgtgaaaccatcaggtcagacacattcttcatatacagcaaacgaat
cttcaaagatggagcaatcctcagtcaagtcctcaaaaattcatctaaattagtgtagtgtca
gggtgatctcagtgaacacacgtaattgtcctgtgccaacattgccttactgtagcacggctat
gcgagaacgggcttcccaaagacttctgttactattttaaactatataatgagttgtgtgcagac
atactttgactctgagttctccatcaccaacaattcgcacccgatcttaatacagtcgtggattg
aggacatctcttttgcactcatatgttctgactcctgcccaattagggggactgagtaacctt
caatactcaaggctctacactagaaatatcggtagcccggggactactgctttgcagagatc
aagcgactagaagcagtgaggattactgagtcctaacattatgactaatatcttaactaggccg
cctgggaatggagattgggccaagtctgtgcaacgaccatactctttcaattttgagactgttg
caagcccaaataattgttctaagaaacatacgcgaagagtcctatttgaaactgttcaaatccc
ttattgtctggagtgacacagaggataatgaggcagaagagaaggcattggctgaattcttg
cttaatacaagaggtgattcatccccgcgttgcgcatgccatcatggaggcaagctctgtaggt
aggagaaagcaaattcaagggtgttgacacaacaaacaccgtaattaagattgcgcttact
aggaggccattaggcatcaagaggctgatgcggatagtcaattattctagcatgcatgcaat
gctgtttagagacgatgtttttcctccagtagatccaaccaccccttagtctcttctaatatgtgt
tctctgacactggcagactatgcacggaatagaagctgggtcacctttgacgggaggcagga
aaatactgggtgtatctaactcctgatacgaataaactcgtagagggtgagattcttagtgaag
cggagggtgtacaagatgtgacagcggagatgaacaatttacttgggtccatcttccaagcaa

tatagaattgaccgatgacaccagcaagaatcctccgatgagggtaccatatctcgggtcaa
agacacaggagaggagagctgcctcacttgcaaaaatagctcatatgtcgccacatgtaaa
ggctgccctaagggcacatccgtgttgatctgggcttatggggataatgaagtaaattggac
tgctgctcttacgattgcaaaatctcgggtgtaactgtaaacttagagtatcttcggttactgtcccc
tttaccacggctgggaatcttcaacatagactagatgatggtataactcagatgacattcacc
cctgcatctcttacaggGtgtcaccttacattcacatatccaatgattctcaaaggctgttcac
tgaagaaggagtgcaaagaggggaatgtggtttaccacagatcatgctcttgggtttatctcta
atcgaatcgatctttccaatgacaacaaccaggacatatgatgagatcacactgcacctacat
agtaaatttagttgctgtatcagagaagcacctgttgcggttccttcgagctacttgggggtgt
accggaactgaggacagtgcctcaaataagtttatgtatgaccttagccctgtatcgaggagg
agactttgcgagacttgacttagctatcttcaagagttatgagcttaatctggagtcatatccca
cgatagagctaataaacattctttcaatatccagcgggaagtgttgccagtcgtgtggtttct
tatgatgaagatacctccataaagaatgacgccataatagtgtatgacaatacccgaaattgg
atcagtgaagctcagaattcagatgtggtccgcctatttgaatatgcagcacttgaagtgtcc
tcgactgttcttaccactctattacctgagagtaagaggcctGgacaatattgtcttatatg
gggtgatttatacaagaatatgccaggaattctactttccaacattgcagctacaatatctcatcc
cgtcattcattcaagggtacatgcagtgggcctgggtcaaccatgacggatcacaccaacttgc
agatacggattttatcgaaatgtctgcaaaactattagtatcttgcacccgacgtgtgatctccg
gcttatattcaggaaataagtatgatctgctgttcccatctgtcttagatgataacctgaatgaga
agatgcttcagctgatatcccggttatgtgtctgtacacgggtactctttgctacaacaagaga
aatcccgaaaataagaggcttaactgcagaagagaaatgttcaatactcactgagtatttactg
tcggatgctgtgaaaccattacttagccccgatcaagtgagctctatcatgtctcctaacataat
tacattcccagctaattctgtactacatgtctcggaagagcctcaatttgatcagggaagggga
ggacaggggatactatcctggcggtgtgttcccccaagagccattattagagttcccttctgtgc
aagatattggtgctcgagtgaagatccattcacccgacaacctgcggcattttgcaagagtt
agatttgagtgtccagcaagggtatgacgcattcacacttagtcagattcatcctgaactcaca
tctccaaatccggagggaagactacttagtacgatacttgttcagagggatagggactgcatct
tcctcttggtataaggcatctcatctcctttctgtacccgaggtgaagatgtgcaagacacggga
actccttatacttagctgaagggagcggagccatcatgagtcttctcgaactgcatgtaccac
atgaaactatctattacaatacgtcttttcaaatgagatgaacccccgcaacgacatttcgg
gccgaccccaactcagttttgaattcgggtgtttataggaatctacaggcggaggtaacatgc
aaagatggatttgccaagagttccgtccattatggagagaaaaatacagaggaaagCgacc
tgacctcagataaagTagtgggggtatattacatctgcagtgccctacagatctgtatcattgct
gcattgtgacattgaaattcctccaggggtccaatcaaagcttactagatcaactagctatcaatt
tatctctgattgccatgcattctgtaagggagggcggggtagtaatcatcaaagtgttgatgc
aatgggatactactttcatctactcatgaactgtttgtccgtgttcacaaaaggatatattctc
tctaattgggtatgcatgtcgaggagatatggagtgttacctgggtattgtcatgggttacctggg
cgggcctacattgtacatgaggtggtgaggatggcGaaaactctgggtgcagcggcacgggt
acgctTttgtctaaatcagatgagatcacactgaccaggttattcacctcacagcggcagcgt
gtgacagacatcctatccagtcctttaccaagattaataaagtacttgaggaagaatattgaca
ctgcgctgattgaagccgggggacagccccgtccgtccattctgtgcggagagctggtgag
cacgctagcgaacataactcagataaccagatCatcgctagtcacattgacacagttatcc
gggtctgtgatatatatggaagctgaggggtgatctcgctgacacagtatttctattacccttac
aatctcttactgacgggaaaaagaggacatcacttaAacagtgcacgagacagatcctag
aggttacaatactaggtcttagagtcgaaaaatctcaataaaaataggcgatataatcagcctagt
gcttaaaggcatgatctccatggaggaccttatcccactaaggacatacttgaagcatagtac
ctgccctaaatattgaaggctgtcctaggtattacaaactcaaagaaatgtttacagacactt
ctgtaCtgtacttgactcgtgtcaacaaaaattctacatgaaaactataggcaatgcagtcaa
aggatattacagtaactgtgactcttaacgaaaatcacatattaataggctccttttttgccaat
tgtattctgttgatttaacatattatgttagaaaaaagttgaaccctgactccttaggactcgaa

ttcgaactcaataaatgtcttataaaaagggttgcgcacaaattattcttagtgtagtctcgtcat tcaccaaattcttgtttggt Genome of
accaaacagagaatccgtgagttacgataaaaggcgaaggagcaattgaagtcgcacggg 59 NDV LaSota
tagaaggtgtgaatctcgagtgcgagcccgaagcacaactcgagaaaagccttctgccaac with a chimeric
atgtcttccgtatttgatgagtagaacagctcctcgcggctcagactcgcccaatggagct F protein
catggaggggggagaaaaaggagtagccttaaaagtagacgtcccggtattcactcttaaca (underlined).
gtgatgacccagaagatagatggagctttgtggtattctgcctccggattgctgttagcgaag Ectodomain is
atgccaaaccactcaggcaaggtgctctcatatctcttttatgctccactcacaggtaat NOT optimized
gaggaacatgttgccCttgcagggaacagaatgaagccacattggccgtgcttgagatt from bovine
gatggctttgccaacggcacgccccagttcaacaataggagtggagtgtctgaagagagag RSV strain
cacagagatttgcgatgatagcaggatctctccctcgggcatgcagcaacggaaccccgctt ATCC51908.
gtcacagccggggcCgaagatgatgcaccagaagacatcaccgataccctggagaggat Transmembrane
cctctctatccagggtcaagtatgggtcacagtagcaaaagccatgactgcgtatgagactg and cytoplasmic
cagatgagtcggaaacaaggcgaatcaataagtatatgcagcaaggcagggtccaaaaga domains are
aatacatcctctaccccgatgcaggagcacaatccaactcacgatcagacagtctcttgag from the F
tccgcattcttttggttagcgagctcaagagaggccgcaacacggcagggtggtacctctactt protein of NDV
attataacctggtaggggacgtagactcatacatcaggaataccggggttactgcattcttct strain LaSota
gacactcaagtacggaatcaacaccaagacatcagcccttgacttagtagcctctcaggcg
acatccagaagatgaagcagctcatgcgtttgtatcggtatgaaggagataatgcgccgtac
atgacattacttggtgatagtaccagatgagctttgcgcctgccgagtatgcacaactttact
cctttgccatgggtatggcatcagtcctagataaaaggtagtgggaaataccaatttgccaggg
actttatgagcacatcattctggagacttggagtagagtacgctcagggtcagggaagtagc
attaacgaggatattggctgccgagctaaagctaaccagcagcaaGgaGgggcctggc
agctgctgccaacgggtctccgaGgaGaccagcagcataGacatgcctactcaacaag
tcggagtcctcactgggcttagcgaggggggggtcccaagctctacaaggcggatcgaata
gatcgcaagggaaccagaagccgggggatggggagacccaattcctggatctgatgaga
gcggtagcaaatagcatgagggagggcgccaaactctgcacagggcactccccaatcggg
gcctccccaaactcctgggcatcccaagataacgacaccgactgggggtattgatggaca
aaaccagcctgcttcacaaaaacatcccaatgccctcaccgtagtcgacccctcgatttg
cggctctatatgaccacacctcaacaaacatccccctcttctcctccccctgctgtaca
actAcgTaccgcttagataccacaggcacaatgcgggtcactaacaatcaaacagagc
cgagggaattagaaaaaagtacgggtagaagagggatattcagagatcagggaagcttc
ccgagtctctgctctcctctacctgatagaccaggacaaacatggccacctttacagatgc
agagatcgacgagctatttgagacaagtggaaactgtcattgacaacataattacagcccagg
gtaaacagcagagactgttggaaggagtgaatcccaaggcaagaccaaggtgctga
gcgcagcatgggagaagcatgggagcatccagccaccggccagtcaagacaaccccgat
cgacaggacagatctgacaacaacatccacaccgagcaaacgaccccgcatgacag
cccgcgggccacatccgccgaccagccccccaccaggccacagacgaagccgtcgac
acacagCtcaggaccggagcaagcaactctctgctgttgatgcttgacaagctcagcaata
aatcgccaatgctaaaaagggcccatggctgagcccccaagaggggaatcaccaacgtc
cgactcaacagcaggggagtcacccagtcgcggaaacagtcaggaaagaccgcagaa
ccaagtcaaggccgcccctggaaaccaggggcacagacgtgaacacagcatatcatggac
aatgggaggagtcacaactatcagctggtgcaacccctcatgctctccgatcaaggcagag
ccaagacaatacccttgatctgcggatcatgtccagccacctgtagactttgtgcaagcgat
gatgtctatgatggaggcgatatcacagagagtaagtaagggtgactatcagctagatctgtc
ttgaaacagacatcctccatccctatgatgcggtccgaaatccaacagctgaaaacatctgtt
gcagtcatggaagccaacttgggaatgatgaagattctggatccccggttggtgccaacattca
tctctgagtgatctacgggcagttgccccgatctcaccggttttagtttcaggccctggagacc
cctctccctatgtgacacaaggaggcgaaatggcacttaataaaacttcgcaaccagtgcc
catccatctgaattgattaaacccgccactgcatgcgggcctgatataggagtggaaaagga
cactgtccgtgcattgatcatgtcacgccaatgcacccgagttcttcagccaagctcctaag

caagttagatgcagccgggtcgatcgaggaatcaggaaaatcaagcgcttgctctaaat
ggctaattactactgccacacgtagcgggtccctgtccactcggcatcacacggaatctgca
ccgagttcccccccgGgTTAGAAAAAATACGGGTTAGAACCGCC
ACCATGGCGACAACAGCCATGAGGATGATCATCAGCAT
TATCTTCATCTCTACCTATGTGACACATATCACTTTATG
CCAAAACATAACAGAAGAATTTTATCAATCAACATGCA
GTGCAGTTAGTAGAGGTTACCTTAGTGCAATTAAGAACT
GGATGGTATACAAGTGTGGTAACAATAGAGTTGAGCAA
AATACAAAAAAATGTGTGTAATAGTACTGATTCAAAAG
TGAAATTAATAAAGCAAGAACTAGAAAGATACAACAAT
GCAGTAGTGGAATTGCAGTCACTTATGCAAAATGAACC
GGCCTCCTTCAGTAGAGCAAAAAGAGGGATACCAGAGT
TGATACATTATACAAGAACTCTACAAAAAAGTTTTAT
GGGCTAATGGGCAAGAAGAGAAAAAGGAGATTTTTAG
GATTCTTGCTAGGTATTGGATCTGCTATTGCAAGTGGTG
TAGCAGTGTCCAAAGTACTACACCTGGAGGGAGAGGTG
AATAAAATTAATAAATGCCTGCTATCCACAAATAAAGC
AGTAGTTAGTCTATCCAATGGAGTTAGTGTCCTTACTAG
CAAAGTACTTGATCTAAAGAACTATATAGACAAAGAGC
TTCTACCTAAAGTTAACAATCATGATTGTAGGATATCCA
AAATAGAACTGTGATAGAATTCCAACAAAAAAACAAT
AGATTGTTAGAAATTGCTAGGGAATTTAGTGTAATGC
TGGTATTACCACACCTCTCAGTACATACATGTTGACCAA
TAGTGAATTACTATCACTAATTAATGATATGCCTATAAC
GAATGACCAAAAAAAGCTAATGTCAAGTAATGTTCAA
TAGTCAGGCAACAGAGTTATTCCATTATGTCAGTGGTCA
AAGAAGAAGTCATAGCTTATGTTGTACAATTGCCTATTT
ATGGAGTTATAGACACCCCCTGTTGGAACTACACACC
TCTCCGTTATGCACCACTGATAATAAAGAAGGGTCAA
CATCTGCTTAACTAGGACAGATCGTGGGTGGTATTGTG
ACAATGCAGGCTCTGTGTCTTTTTTCCCACAGACAGAGA
CATGTAAGGTACAATCAAATAGAGTGTTCTGTGACACA
ATGAACAGTTTAACTCTGCCTACTGACGTAACTTATGC
AACACTGACATATTCAATACAAAGTATGACTGTAAAAT
AATGACATCTAAAAGTACATAAGTAGCTCTGTGATAA
CTTCAATTGGAGCTATTGTATCATGCTATGGGAAGACA
AAATGTACAGCTTCTAATAAAAATCGTGGAATCATAAA
GACTTTTTTCCAATGGGTGTGATTATGTATCAAACAAAGG
AGTAGATACTGTATCTGTTGGTAACACACTATATTATGT
AAATAAGCTAGAGGGGAAAGCACTCTATATAAAGGGTG
AACCAATTATTAATTACTATGATCCACTAGTGTTTCCTT
CTGATGAGTTTGATGCATCAATTGCCCAAGTAAACGCA
AAAATAAACCAGCCTGGCCTTCATACGTCGATCTGA
TGAGTTACTTCACAGTGTAGATGTAGGAAAATCCACCA
CAAATgttaacCTCATTACCTATATCGTTTTTACTATCATAT
CTCTTGTTTTTGGTATACTTAGCCTGATTCTAGCATGCTA
CCTAATGTACAAGCAAAAAGGCGCAACAAAAGACCTTAT
TATGGCTTGGAATAATACCCTAGATCAGATGAGAGCC
ACTACAAAAATGTGAGCTTCAGCAAGTAAccccccgcggaccc
aagggtccaactctccaagcggcaatcctctctcgcttctcagccccactgaatgAtcgcg

aaccgtaattaatctagctacatfttaagattaagaaaaaacgggtagaattggagtgcccca
attgtgccaatggactcatctaggacaattgggctgtactttgattctgccattcttctagca
acctgttagcatttccgatcgctcctacaagAcacaggagatgggaagaagcaaatcgcccc
gcaatataggatccagcgccttgacttgtggactgataagaggactcagtattcatcac
cacctatggattcatctttcaagttgggaatgaagaagccacCgtcggcatgatcgatgataa
acccaagcgcgagttactttccgctgcgatgctctgcctaggaagcgtcccaaataccggag
accttattgagctggcaaggcctgtctcactatgatagtcacatgcaagaagagtgcacta
atactgagagaatggttttctcagtagtgcaggcaccccaagtgtgcaaagctgtagggttg
tggcaaaaaatactcatcagtgaaatgcagtcaagcacgtgaaagcgccagagaagattcc
cgggagtggaaccctagaatacaaggtgaactttgtctccttgactgtggtaccgaagaGg
gatgtctacaagatcccagctgcagtattgaagggttctggctcgagtctgtacaatcttgcgct
caatgtcactattaatgtggaggttagacccgaggagtcctttggttaaactCtgtctaagtct
gacagcggatactatgctaacctcttctgcatattggacttatgaccacTgtagataggaag
gggaagaaagtgcatttgacaagctggaaaaagaaaataaggagccttgatctatctgtcgg
gctcagtgatgtgctcgggcttccgtgttggtaaaagcaagaggtgcacggactaagctttt
ggcacctttcttcttagcagtgggacagcctgctatcccatagcaaatgcttctcctcaggtg
gccaagatactctggagtcaaaccgctgcctgcggagcgttaaaatcattatccaagcagg
taccacgcgctgtcgcagtgcacggcaccacgaggttaccttactaagctggagaag
gggcacacccttgccaaatacaatccttttaagaaataagctgcgtctctgagattgcgctcc
gcccactcaccagatcatcatgacacaaaaaactaatctgtcttgattatttacagttagtttac
ctgtctatcaagttagaaaaaacacgggtagaagattctggatcccgggtggcgccctccag
gtgcaagatgggctccagaccttctaccaagaaccagcacctatgatgtgactatccggg
ttgcgctgggtactgagttgcatctgtccggcaaaactccattgatggcaggcctcttgcagctgc
aggaattgtggttacaggagacaaagccgtcaacatatacacctcatcccagacaggatcaa
tcatagttaagctcctcccgaatctgcccaggataaggaggcatgtgcgaaagcccccttg
gatgcatacaacaggacattgaccactttgctcaccccccttggtgactctatccgtaggatac
aagagtctgtgactacatctggaggggggagacaggggcgccttataggcgccattattgg
cgggtgtggctcttgggggttgcaactgccgcacaaataacagcggccgcagctctgatacaa
gcaaaacaaaatgctgccaacatcctccgacttaaagagagcattgccgcaaccaatgagg
ctgtgcatgaggtcactgacggattatcgcaactagcagtggcagttgggaagatgcagca
gtttgttaatgaccaatthaataaaacagctcaggaattgactgcatcaaaattgcacagcaa
gttggtgtagagctcaacctgtacctaaccgaattgactacagtattcggaaccacaaatcactt
cacctgctthaacaagctgactattcaggcactttacaatctagctgggtggaaatatggattac
ttattgactaagttaggtgtagggaacaatcaactcagctcattaatcggtagcggcctaatca
ccggtaacctattctatacgaactcacagactcaactcttgggtatacaggttaactctacctca
gtcgggaacctaaataatatgcgtgccacctacttggaaccttatccgtaagcacaccag
gggatttgctcggcacttgtcccAaaaagggtgacacaggtcgggtctgtgataagaagac
ttgacacctcatactgtatagaactgacttagatttatattgtacaagaatagtaacgttcctat
gtcccctgggtattttattctgcttgagcggcaatacgtcggcctgtatgtactcaaagaccgaa
ggcgcacttactacaccatacatgactatcaaagggtcagtcacgccaaactgcaagatgaca
acatgtagatgtgtaaacccccgggtatcatatcgcaaaactatggagaagccgtgtctcta
atagataaacaatcatgcaatgtttatccttaggcgggataactthaaggctcagtggggaatt
cgatgtaacttatcagaagaatatctcaatacaagattctcaagtaataataacaggcaatcttg
atatctcaactgagcttgggaatgtcaacaactcgatcagtaatgctttgaataagttagagga
aagcaacagaaaactagacaaagtcaatgtcaaactgactagcacatctgctctcattaccta
tatcgttttgactatcatactcttgtttttgggtatacttagcctgattctagcatgctacctaata
caagcaaaaaggcgcaacaaaagaccttattatggcttgggaataatactctagatcagatga
gagccactacaaaaatgtgaacacagatgaggaacgaagggttcctaataagtaatttgtgtg
aaagttctggtagtctgtcagttcagagagtttaagaaaaaactaccggtttagatgacaaa
ggacgatatacgggtagaacggtaagagaggccgccccctcaattgcgagccaggctcac

aacctccgttctaccgcttcaccgacacagtcctcaatcatggaccgcgccgttagccaagt
tgcgttagagaatgatgaaagagaggcaaaaaatacatggcgcttgatattccggattgcaat
cttattcttaacagtagtgaccttggtctatatctgtagcctcccttttatatagcatgggggctag
cacacctagcgatctgttaggcataccgactaggatttcagggcagaagaaaagattacat
ctacacttgggtccaatcaagatgtagtagataggatatataagcaagtggcccttgagtctcc
gttggcattgttaaactgagaccacaattatgaacgcaataacatctctctcttatcagattaa
tggagctgcaaacaacagtggggtggggggcacctatccatgaccagattatataggggg
gataggcaaagaactcattgtagatgatgctagtgatgtcacatcattctatccctctgcatttc
aagaacatctgaattttatcccggcgctactacaggatcagggtgcactcgaataccctcatt
tgacatgagtgtaccattactgtacaccataatgtaattgtctggatgcagagatcact
cacattcatatcagtttttagcacttgggtgtgctccggacatctgcaacagggagggtattcttt
tctactctgcgttccatcaacctggacgacacccaaaaatcggaagtcttgcaagtgtgagtga
actcccctgggttgtgatatgctgtgctcgaaagtcacggagacagaggaagaagattataa
ctcagctgtccctacgcggatggtacatgggagggttaggggtcgacggccagtaccacgaa
aaggacctagatgtcacaacattattcggggactgggtggccaactaccagagtagggg
gtggatctttattgacagccgcgtatggttctcagtcacggagggttaaaaccaattcacc
cagtgacactgtacaggaagggaatatgtgatatacaagcgatacaatgacacatgcca
gatgagcaagactaccagattcgaatggccaagtcttcgtataagcctggacggttgggtgg
gaaacgcatacagcaggctatcttatctatcaagggtgcaacatccttaggcgaagaccggt
actgactgtaccgccaacacagtcacactcatgggggcccgaaggcagaattctcacagta
gggacatctcatttctgtatcaacgagggtcatcatacttctctccgcgttattatctctatga
cagtcagcaaaaaacagccactcttcatagtccttatacattcaatgccttcactcggccagg
tagtatcccttgccaggcttcagcaagatgccccaaactcgtgtgttactggagtctatacagat
ccatatcccctaattcttctatagaaaccacaccttgcgaggggtattcgggacaatgcttgatg
gtgtacaagcaagacttaacctgctgtcagatttcgatagcacatcccgcagtcgcatta
ctcgagtgttcaagcagtaaccaagcagcatacacaacatcaactgttttaaagtggta
agactaataagacctattgtctcagcattgtgaaatatctaactctcttcggagaattcaga
atcgtcccgttactagttgagatcctcaaagatgacgggggttagagaagccagggtctggcta
gttgagtcaattataaaggagttggaaagatggcattgtatcacctatcttctgcgacatcaag
aatcaaaccgaatgccggcgcggtgctcgaattccatgttgccagttgaccacaatcagccag
tgctcatgcatcagattaagccttggtcaAtaGtctcttgattaagaaaaaatgtaagtggca
atgagatacaaggcaaaacagctcatgggTaaCaatacgggtaggacatggcgagctcc
ggctctgaaagggcagagcatcagattatcctaccagagTcacacctgtcttcaccattggg
caagcacaaactactctattactggaaattaactgggctaccgcttctctgatgaatgtgacttc
gaccacctcattctcagccgacaatggaaaaaaatacttgaatcggcctctctgatactgag
agaatgataaaactcggaagggcagtacaccaaactcttaaccacaattccagaataaccg
gagtgtccaccccagggtgttagaaGaactggctaattagggtcccagattcaaccaac
aaatttcggaagattgagaagaagatccaaattcacaacacgagatatggagaactgttcac
aaggctgtgtacgcatatagagaagaaactgctgggggtcatcttgggtctaacaatgtcccccg
gtcagaggagttcagcagcattcgtacggatccggcattctgggttactcaaaatgggtccac
agccaagtttgcatgggtccatataaaacagatccagaggcatctgatgggtggcagctaGg
acaaggctctgcggccaacaaattgggtgatgtaaccataaggtaggccaagtctttgtcact
cctgaacttgtcgttgtgacgcatacgaatgagaacaagttcacatgtcttaccaggaacttg
tattgatgtatgcagatatgatggagggcagagatatgggtcaacataatatcaaccacggcg
gtgcatctcagaagcttatcagagaaaattgatgacattttgcgggttaatagacgctctggcaa
aagacttgggtaatacaagtctacgatgttgatcactaatggagggtattgcatacggagctgt
ccagctactcgagccgtcaggtacatttgcaggagatttcttcgattcaacctgcaggagctt
aaagacattctaattggcctcctcccaatgatatagcagaatccgtgactcatgcaatcgcta
ctgtattctctggtttagaacagaatcaagcagctgagatgttgtgtctgttgctgtgtgggt
caccactgcttgagtcccgatttgagcaaaaggcagtcaggagccaaatgtgcgcaccga

aaatggtagactttgatatgatccttcaggtagtcttttcaagggaacaatcatcaacggg
tacagaaagaagaatgcagggtgtgtggccgcgagtgcaagtgagatacaatatatgggaagg
tcattggggcaactacatgcagattcagcagagattcacacgatatcatgttgagagagtataa
gagtttatctgcacttgaatttgagccatgtatagaatatgacctgtcaccaacctgagcatgt
tcctaaaagacaaggcaatcgcacaccccaacgataattggcttgcctcgttaggcggaac
cttctctccgaagaccagaagaaacatgtaaaagaagcaacttcgactaatcgctcttgata
gagtttttagagtcaaagtatttgatccatataaagagatggaatatctgacgaccttgagta
ccttagagatgacaatgtggcagtatcatactcgctcaaggagaagggaagtgaagttaatg
gacggatcttcgctaagctgacaaaagaagtaagggaactgtcaggtgatggcggaagggat
cctagccgatcagattgcacctttcttcagggaaatggagtcattcaggatagcatatccttg
accaagagtatgctagcgatgagtcaactgtctttaacagcaataagaaacgtatcactgact
gtaaagaaagagtatcttcaaaccgcaatcatgatccgaaaagcaagaaccgtcggagagt
tgcaaccttcataacaactgacctgcaaaagtactgtcttaattggagatatcagacaatcaaa
ttgttcgctcatgccatcaatcagttgatgggcctacctcactcttcgaatggattcacctaag
actgatggacactacgatgttcgtaggagaccctttcaatcctccaagtgacctactgactgt
gacctctcaagagtcctaataatgatgacatatatattgtcagtgccagaggggggtatcgaagga
ttatgccagaagctatggacaatgatctcaattgctgcaatccaacttgctgcagctagatcgc
attgtcgtgttgctgtatggtacaggggtataatcaagtaatagcagtaacgagagaggtaa
gatcagacgactctccggagatgggtgttgacacagttgcatcaagccagtataatcttcaa
ggaattaattcatgtcaatcatttgattggccataattgaaggatcgtgaaaccatcaggtcag
acacattcttcatatacagcaaacgaatcttcaaagatggagcaatcctcagtcaagtcctcaa
aaattcatctaaattagtgtagtgctaggtgatctcagtgaacacaccgtaatgtcctgtgcca
acattgccttactgtagcacggctatgcgagaacgggcttcccaaagacttctgttactattta
aactatataatgagttgtgtgcagacatactttgactctgagttctccatcaccaacaattcgca
ccccgatcttaatcagtcgtggattgaggacatctctttgtgcactcatatgttctgactcctgc
ccaattaggggggactgagtaaccttcaatactcaaggctctacactagaaatatcggtgacct
ggggactactgcttttgagagatcaagcgactagaagcagtgaggattactgagtcctaaca
ttatgactaataatcttaactaggccgcctgggaatggagattgggccagtctgtgcaacgacc
catactctttcaatttgagactgttgcaagcccaaattgttcttaagaaacatacgcгааага
gtcctatttgaaactgttcaaatcccttattgtctggagtgcacacagaggataatgaggcag
aagagaaggcattggctgaattctgtctaataagaggtgattcatccccgcgttgcgcatg
ccatcatggaggcaagctctgtaggtaggagaaagcaaattcaagggtgttgacacaac
aaacaccgtaattaagattgcgcttactaggaggccattagggcatcaagaggctgatgcgga
tagtcaattattctagcatgcatgcaatgctgtttagagacgatgttttctccagtagatccaa
ccaccccttagtctcttctaatatgtgttctctgacactggcagactatgcacggaatagaagct
ggtcaccttgacgggagggcaggaaaatactgggtgtatctaactctgatacगतagaactc
gtagaggggtgagattcttagtgtaagcggaggggtgtacaagatgtgacagcggagatgaac
aatttacttggtccatcttccaagcaatatagaattgaccgatgacaccagcaagaatcctcc
gatgaggggtaccatatctcgggtcaaagacacaggagaggagagctgcctcacttgcaaaa
atagctcatatgtcgccacatgtaaaggctgcctaagggcacatccgtgttgatctgggctt
atgggggataatgaagtaaattggactgctgctcttacgattgcaaaatctcggtgtaatgtaaa
cttagagtatcttcggttactgtcccctttaccacggctgggaatcttcaacatagactagatg
atggtataactcagatgacattcacccctgcatctctctacaggGtgtcaccttacattcacata
tccaatgattctcaaaggctgttactgaagaaggagtcaaagaggggaatgtgggtttacca
cagatcatgctcttggtttatctctaatacgaatcgatctttccaatgacaacaaccaggacata
tgatgagatcacactgcacctacatagtaaatttagttgctgtatcagagaagcacctgttgcg
gttcctttcgagctacttgggggtgtaccggaactgaggacagtgcactcaaataagtttatgt
atgatcctagccctgtatcgaggaggagactttgcgagacttgacttagctatcttcaagagttat
gagcttaatctggagtcatatcccacgatagagctaataaacattcttcaatatccagcggga
agttgattggccagtctgtggtttcttatgatgaagatacctccataaagaatgacgccataata

gtgtatgacaatacccgaattggatcagtgaaagctcagaattcagatgtgggtccgcctatttg
aatatgcagcacttgaagtgtcctcactgttcttaccactctattacctgagagtaagagg
cctGgacaatattgtcttatatatgggtgatttatacaagaatatgccaggaattctactttcaa
cattgcagctacaatatctcatcccgctcattcattcaagggtacatgcagtgggcctgggtcaac
catgacgggatcacaccaacttgcatgacggattttatcgaaatgtctgcaaaactattagtat
cttgcacccgacgtgtgatctccggcttatattcaggaaataagtatgatctgctgttcccatct
gtcttagatgataacctgaatgagaagatgcttcagctgatatcccggttatgctgtctgtacac
gggtactctttgtacaacaagagaaatcccgaataaagaggcttaactgcagaagagaaat
gttcaatactcactgagtatcttactgtcggatgctgtgaaaccattacttagccccgatcaagt
agctctatcatgtctcctaacataattacattcccagctaattctgtactacatgtctcggaagag
cctcaatttgatcaggggaaagggaggacagggatactatcctggcggttgtgttcccccaag
agccattattagagttcccttctgtgcaagatattgggtgctcgagtgaagatccattcacccg
acaacctgcggcatttttgaagagttagatttgagtgtccagcaagggtatgacgcattcaca
cttagtcagattcatcctgaactcacatctccaaatccggaggaagactacttagtacgatactt
gttcagaggggatagggactgcatcttctcttgggtataaggcatctcatctctttctgtacccg
aggtaagatgtgcaagacacgggaactccttatacttagctgaaggggagcggagccatcat
gagtcttctcgaactgcatgtaccacatgaaactatctattacaatacgcctctttcaaagt
gaaccccccgcaacgacatttcggggccgaccccaactcagttttgaattcgggtgttatagg
aatctacaggcggaggtacatgcaaagatggattgtccaagagttccgtccattatggaga
gaaaatacagaggaaagCgacctgacctcagataaagTagtgggggtatattacatctgca
gtgccctacagatctgtatcattgctgcattgtgacattgaaattcctccaggggtccaatcaaag
cttactagatcaactagctatcaattatctctgattgccatgcattctgtaaggggagggggg
tagtaatcatcaaagtgtgtatgcaatgggatactactttcatctactcatgaactgtttgctcc
gtgttccacaaaaggatattctctctaatgggtatgcatgtcgaggagatatggagtgttacc
tggtatttgcagtggttacctggggcgccctacattgtacatgaggtggtgaggatggcGa
aaactctggtgcagcggcacgggtacgctTttgtctaaatcagatgagatcacactgaccagg
ttattcacctcacagcgggcagcgtgtgacagacatcctatccagtcctttaccaagattaataa
agtacttgaggaagaatattgacactgcgctgattgaagccggggggacagcccgtccgtcc
attctgtgcggagagctggtgagcacgctagcgaacataactcagataaccagatCatc
gctagtacattgacacagttatccgggtctgtgatatatatggaagctgaggggtgatctcgctg
acacagtatttctatttacccttacaatctcttactgacgggaaaaagaggacatcacttaA
acagtgcacgagacagatcctagaggttacaatactaggtcttagagtcgaaaatctcaataa
aataggcgcataatcagcctagtgtctaaaggcatgatctccatggaggaccttatccctact
aaggacatacttgaagcatagtagctgcctaaatatttgaaggctgtcctaggtattaccaaa
ctcaaagaaatgtttacagacacttctgtgaCtgtacttgactcgtgctcaacaaaaattctacat
gaaaactataggcaatgcagtcaaaggatattacagtaactgtgactcttaacgaaaatcaca
tattaataggctcctttttggccaattgtattcttgtgattaatcatattatgttagaaaaaagttg
aacctgactccttaggactcgaattcgaactcaaataaatgtcttaaaaaaagggtgcgcac
aattattcttgagtgtagtctcgtcattcaccaaatctttgtttgggt

(189) TABLE-US-00005 TABLE 4 hMPV F Sequences Wild-type

ATGTCTTTGGAAAGTGGTGATCATTTTTTTCATTGCTAATAACACCTCAA 16 Nucleic acid
CACGGTCTTAAAGAGAGCTACCTAGAAGAATCATGTAGCACTATAACT encoding
GAGGGATATCTTAGTGTTCTGAGGACAGGTTGGTATACCAACGTTTTT hMPV-F
ACATTAGAGGTGGGTGATGTAGAAAACCTTACATGTTCTGATGGACCT protein of
AGCCTAATAAAAACAGAATTAGATCTGACCAAAAGTGCCTAAGAGAG hMPV strain
CTCAAAACAGTCTCTGCTGACCAATTGGCAAGAGAGGAACAAATTGAG CAN00-16
AATCCCAGACAATCTAGGTTTGTCTAGGAGCAATAGCACTCGGTGTT (transmembrane
GCAACAGCAGCTGCAGTCACAGCAGGTGTTGCAATTGCCAAAACCATC and
cytoplasmic CGGCTTGAGAGTGAAGTCACAGCAATTAAGAATGCCCTCAAACGACC
domains are AATGAAGCAGTATCTACATTGGGGAATGGAGTTCGAGTGTTGGCAACT

indicated) by CCACTGAGAGAGCTGAGTGAAGAACTTGTGTGCAAGAACTTCGTGCA
underlining) ATCAACAAAAACAAGTGCGACATTGATGACCTAAAAATGGCCGTTAGC
TTCAGTCAATTCACAGAAGGTTTCTAAATGTTGTGCGGCAATTTTCA
GACAATGCTGGAATAACACCAGCAATATCTTTGGACTTAATGACAGAT
GCTGAAGTAGCCAGGGCCGTTTCTAACATGCCGACATCTGCAGGACAA
ATAAAATTGATGTTGGAGAACCGCGCGATGGTGCGAAGAAAGGGGTTC
GGAATCCTGATAGGGGTCTACGGGAGCTCTGTAATTTACATGGTGCAG
CTGCCAATCTTTGGCGTTATAGACACGCCTTGCTGGATAGTAAAAGCA
GCCCCTTCTTGTTCCGAAAAAAAGGGAAACTATGCTTGCCTCTTAAGA
GAAGACCAAGGGTGGTATTGTCAGAATGCAGGGTCAACTGTTTACTAC
CCAAATGAGAAAGACTGTGAAACAAGAGGAGACCATGTCTTTTTCGAC
ACAGCAGCAGGAATTAATGTTGCTGAGCAATCAAAGGAGTGCAACATC
AACATATCCACTACAAATTACCCATGCAAAGTCAGCACAGGAAGACAT
CCTATCAGTATGGTTGCACTGTCTCCTCTTGGGGCTCTGGTTGCTTGC
TACAAAGGAGTAAGCTGTTCCATTGGCAGCAACAGAGTAGGGATCATC
AAGCAGCTGAACAAAGGTTGCTCCTATATAACCAACCAAGATGCAGAC
ACAGTGACAATAGACAACACTGTATATCAGCTAAGCAAAGTTGAGGGT
GAACAGCATGTTATAAAAGGCAGACCAGTGTCAAGCAGCTTTGATCCA
ATCAAGTTTCCTGAAGATCAATTCATGTTGCACTTGACCAAGTTTTT
GAGAGCATTGAAAACAGCCAGGCCTTGGTAGATCAATCAAACAGAATC
CTAAGCAGTGCAGAGAAAGGAATACTGGCTTCATCATTGTAATAATT
CTAATTGCTGTCCTTGGCTCTAGCATGATCCTAGTGAGCATCTTCATT
ATAATCAAGAAAACAAAGAAACCAACGGGAGCACCTCCAGAGCTGAGT
GGTGTACAAACAATGGCTTCATACCACACAGTTAG Amino acid

MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVE 17 sequence of
TLEVGDVENLTCSDGPSLIKTELDTLSALRELKTVSADQLAREEQIE huMPV-F
NPRQSRFVLGAIALGVATAAAVTAGVAIAKTIRLESEVTAIKNALKTT protein of
NEAVSTLGNVVRVLATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVS hMPV strain
FSQFNRRELVVRQFSDNAGITPAISLDLMTDAELARAVSNMPTSAGQ CAN00-16
IKLMLENRAMVRRKGFGILIGVYGSSVIYMVQLPIFGVIDTPCWIVKA (transmembrane
APSCSEKKGNYACLLREDQGWYCQNAGSTVYYPNEKDCETRGDHVFCD and
cytoplasmic TAAGINVAEQSKECNINISTTNYPCKVSTGRHPISMVALSPLGALVAC domains
are YKGVSCSIGSNRVGIIKQLNKGCSYITNQDADTVTIDNTVYQLSKVEG underlined;
SEQ EQHVIKGRPVSSSFDPIKFPEDQFNVALDQVFESIENSQALVDQSNRI ID No: 16
or 18 LSSAEKGNTGFIIVIIIAVLGSSMILVSIFIIKKTKKPTGAPPELS encode this
GVTNNGFIPHS amino acid sequence) Codon

ATGAGCTGGAAGGTGGTGATCATCTTCAGCCTGCTGATCACCCCCCAG 18 optimized
CACGGCCTGAAGGAGAGCTACCTGGAGGAGAGCTGCAGCACCATCACC nucleic acid
GAGGGCTACCTGAGCGTGCTGAGAACCGGCTGGTACACCAACGTGTTC encoding
ACCCTGGAGGTGGGCGACGTGGAGAACCTGACCTGCAGCGACGGCCCC hMPV-F
AGCCTGATCAAGACCGAGCTGGACCTGACCAAGAGCGCCCTGAGAGAG protein of
CTGAAGACCGTGAGCGCCGACCAGCTGGCCAGAGAGGAGCAGATCGAG hMPV strain
AACCCCAGACAGAGCAGATTTCGTGCTGGGCGCCATCGCCCTGGGCGTG CAN00-16
GCCACCGCCGCCGCCGTGACCGCCGGCGTGGCCATCGCCAAGACCATC (transmembrane
AGACTGGAGAGCGAGGTGACCGCCATCAAGAACGCCCTGAAGACCAACC and
cytoplasmic AACGAGGCCGTGAGCACCTGGGCAACGGCGTGAGAGTGCTGGCCACC
domains are GCCGTGAGAGAGCTGAAGGACTTCGTGAGCAAGAACCTGACCAGAGCC
underlined) ATCAACAAGAACAAGTGCGACATCGACGACCTGAAGATGGCCGTGAGC
TTCAGCCAGTTCAACAGAAGATTCCTGAACGTGGTGAGACAGTTCAGC
GACAACGCCGGCATCACCCCCGCCATCAGCCTGGACCTGATGACCGAC

CCGAGAGCTGGCCAGAGACCGGCTGAGCAACATGCCACCCAGCGCCGCCAG
ATCAAGCTGATGCTGGAGAACAGAGCCATGGTGAGAAGAAAGGGCTTC
GGCATCCTGATCGGCGTGACGGCAGCAGCGTGATCTACATGGTGACG
CTGCCCATCTTCGGCGTGATCGACACCCCCTGCTGGATCGTGAAGGCC
GCCCCCAGCTGCAGCGAGAAGAAGGGCAACTACGCCTGCCTGCTGAGA
GAGGACCAGGGCTGGTACTGCCAGAACGCCGGCAGCACCGTGTACTAC
CCCAACGAGAAGGACTGCGAGACCAGAGGGCGACCACGTGTTCTGCGAC
ACCGCCGCCGGCATCAACGTGGCCGAGCAGAGCAAGGAGTGCAACATC
AACATCAGCACCACTACCCCTGCAAGGTGAGCACCGGCAGACAC
CCCATCAGCATGGTGGCCCTGAGCCCCCTGGGCGCCCTGGTGGCCTGC
TACAAGGGCGTGAGCTGCAGCATCGGCAGCAACAGAGTGGGCATCATC
AAGCAGCTGAACAAGGGCTGCAGCTACATCACCAACCAGGACGCCGAC
ACCGTGACCATCGACAACACCGTGTAACAGCTGAGCAAGGTGGAGGGC
GAGCAGCACGTGATCAAGGGCAGACCCGTGAGCAGCAGCTTCGACCCC
ATCAAGTTCCCCGAGGACCAGTTCAACGTGGCCCTGGACCAGGTGTTT
GAGAGCATCGAGAACAGCCAGGCCCTGGTGGACCAGAGCAACAGAATC
CTGAGCAGCGCCGAGAAGGGCAACACCGGCTTCATCATCGTGATCATC
CTGATCGCCGTGCTGGGCAGCAGCATGATCCTGGTGAGCATCTTCATC
ATCATCAAGAAGACCAAGAAGCCCCACCGGCGCCCCCCCCCGAGCTGAGC
GGCGTGACCAACAACGGCTTCATCCCCACAGCTGA Nucleic acid
ATGAGCTGGAAGGTGGTGATCATCTTCAGCCTGCTGAT 19 encoding
CACCCCCCAGCACGGCCTGAAGGAGAGCTACCTGGAGG chimeric hMPV
AGAGCTGCAGCACCATCACCGAGGGCTACCTGAGCGTG F protein
CTGAGAACCGGCTGGTACACCAACGTGTTACCCCTGGA comprising the
GGTGGGCGACGTGGAGAACCTGACCTGCAGCGACGGCC ectodomain of
CCAGCCTGATCAAGACCGAGCTGGACCTGACCAAGAGC the F protein of
GCCCTGAGAGAGCTGAAGACCGTGAGCGCCGACCAGCT hMPV strain
GGCCAGAGAGGAGCAGATCGAGAACCCCAGACAGAGC CAN00-16 and
AGATTCGTGCTGGGCGCCATCGCCCTGGGCGTGGCCAC the
CGCCGCCGCGCTGACCGCCGGCGTGGCCATCGCCAAGA transmembrane
CCATCAGACTGGAGAGCGAGGTGACCGCCATCAAGAAC and cytoplasmic
GCCCTGAAGACCACCAACGAGGCCGTGAGCACCTGGG domains of the
CAACGGCGTGAGAGTGCTGGCCACCGCCGTGAGAGAGC F protein of
TGAAGGACTTCGTGAGCAAGAACCTGACCAGAGCCATC NDV LaSota
AACAAGAACAAGTGCGACATCGACGACCTGAAGATGG strain, wherein
CCGTGAGCTTCAGCCAGTTCAACAGAAGATTCCTGAAC the nucleotide
GTGGTGAGACAGTTCAGCGACAACGCCGGCATCACCCC sequence
CGCCATCAGCCTGGACCTGATGACCGACGCCGAGCTGG encoding the
CCAGAGCCGTGAGCAACATGCCACCCAGCGCCGGCCAG ectodomain of
ATCAAGCTGATGCTGGAGAACAGAGCCATGGTGAGAAG the F protein of
AAAGGGCTTCGGCATCCTGATCGGCGTGACGGCAGCA the hMPV is
GCGTGATCTACATGGTGCAGCTGCCATCTTCGGCGTGA codon optimized
TCGACACCCCCTGCTGGATCGTGAAGGCCGCCCCCAGC
TGCAGCGAGAAGAAGGGCAACTACGCCTGCCTGCTGAG
AGAGGACCAGGGCTGGTACTGCCAGAACGCCGGCAGC
ACCGTGTAACCCCCAACGAGAAGGACTGCGAGACCAG
AGGCGACCACGTGTTCTGCGACACCGCCGCCGGCATCA
ACGTGGCCGAGCAGAGCAAGGAGTGCAACATCAACATC
AGCACCACTACCCCTGCAAGGTGAGCACCGGCAG
ACACCCCATCAGCATGGTGGCCCTGAGCCCCCTGGGCG

CCCTGGTGGCTAGTACCAAGGCGTGAAGCTGCAGCATC
GGCAGCAACAGAGTGGGCATCATCAAGCAGCTGAACA
AGGGCTGCAGCTACATCACCAACCAGGACGCCGACACC
GTGACCATCGACAACACCGTGTACCAGCTGAGCAAGGT
GGAGGGCGAGCAGCACGTGATCAAGGGCAGACCCGTG
AGCAGCAGCTTCGACCCCATCAAGTTCCCGAGGACCA
GTTCAACGTGGCCCTGGACCAGGTGTTTCGAGAGCATCG
AGAACAGCCAGGCCCTGGTGGACCAGAGCAACAGAAT
CCTGAGCAGCGCCGAGAAGGGCAACACCGGCGTTAACC
TCATTACCTATATCGTTTTGACTATCATATCTCTTGTTTT
TGGTATACTTAGCCTGATTCTAGCATGCTACCTAATGTA
CAAGCAAAAGGCGCAACAAAAGACCTTATTATGGCTTG
GGAATAATACCCTAGATCAGATGAGAGCCACTACAAAA ATGTGA Amino acid
MSWKVVIIFSLITPQHGLKESYLEESCSTITEGYLSVLRTGWYTNVF 15 sequence of the
TLEVGDVENLTCSDGPSLIKTELDLTKSALRELKTVSADQLAREEQIE chimeric F
NPRQSRFVLGAIALGVATAAAVTAGVAIAKTIRLESEVTAIKNALKTT protein
NEAVSTLGNVVRVLATAVRELKDFVSKNLTRAINKNKCDIDDLKMAVS comprising the
FSQFNRRFLNVVRQFSDNAGITPAISLDLMTDAELARAVSNMPTSAGQ ectodomain of
IKLMLENRAMVRRKGFGILIGVYGSSVIYMVQLPIFGVIDTPCWIVKA hMPV F
APSCSEKKGNVYACLLREDQGWYCQNAGSTVYYPNEKDCETRGDHVFCD protein of
TAAGINVAEQSKECNINISTTNYPCKVSTGRHPISMVALSPLGALVAC CAN00-16
YKGVSCSIGSNRVGIIKQLNKGCSYITNQDADTVTIDNTVYQLSKVEG strain and
EQHVIKGRPVSSSFDPIKFPEDQFNVALDQVFESIENSQALVDQSNRI LaSota strain
LSSAEKGNTGVNLITYIVLTIISLVFGILSLILACYLMYKQKAQQKTL NDV F protein
LWLGNNTLDQMRATTKM transmembrane and cytoplasmic domains cDNA of
accaaacagagaatccgtgagttacgataaaaaggcgaaggagcaattg 20 genomic
aagtcgcacgggtagaagggtgaatctcgagtgcgagcccgaagcac sequence of
aaactcgagaaagccttctgccaatgtctccgtatttgatgagta NDV LaSota
cgaacagctcctcgcggtcagactcgcccaatggagctcatggagg strain with the
gggagaaaaagggagtaccttaaaagtagacgtcccggtattcactct nucleic acid
taacagtgatgaccagaagatagatggagcttgtggtattctgcct encoding hMPV
ccggattgctgttagcgaagatccaacaaaccactcaggcaaggtgc F protein of
tctcatatctctttatgctcccactcacaggaatgaggaacatgt hMPV strain
tgccCttgcagggaaacagaatgaagccacattggccgtgcttgagat CAN00-16
tgatggctttgccaacggcacgccccagttcaacaataggagtggagt inserted (nucleic
gtctgaagagagagcacagagatttgcgatgatagcaggatctctccc acid sequence
tcgggcatgcagcaacggaacccgttcgtcacagccggggcCgaaga encoding hMPV
tgatgcaccagaagacatcaccgataccctggagaggatcctctctat F protein is
ccaggctcaagtatgggtcacagtagcaaaagccatgactgcgtatga indicated by
gactgcagatgagtcggaaacaaggcgaatcaataagtatatgcagca underlining; Sac
aggcagggtccaaaagaaatacatcctctaccccgatgcaggagcac II restriction site
aatccaactcacgatcagacagtctcttgagtcgcatcttttggt is indicated by
tagcgagctcaagagaggccgcaacacggcaggtggtaccttactta double
ttataacctggtaggggacgtagactcatacatcaggaataaccgggct underlining)
tactgcattctcttgacactcaagtacggaatcaacaccaagacatc
agcccttgacttagtagcctctcaggcgacatccagaagatgaagca
gctcatgcgtttgatcggtgaaaggagataatgcgccgtacatgac
attacttggtgatagtaccagatgagctttgcgcctgccgagtatgc acaactttactccttgccatgggtatggcatcagtcctagataaagg
tactgggaaataccaatttgccagggactttatgagcacatcattctg
gagacttgagtagagtacgctcaggctcagggaagtagcattaacga

ggatatggctgccgagctaagctaaccgccagcagcaaGgaGgggcct
ggcagctgctgcccaacgggtctccgaGgaGaccagcagcataGacat
gcctactcaacaagtcggagtcctcactgggcttagcgaggggggggtc
ccaagctctacaaggcggatcgaatagatcgcaagggcaaccagaagc
cggggatggggagaccaattcctggatctgatgagagcggtagcaaa
tagcatgagggagggcgccaaactctgcacagggcactccccaatcggg
gcctccccaaactcctgggccatcccaagataacgacaccgactgggg
gtattgatggacaaaaccagcctgcttcacaaaaacatcccaatgc
cctcacccgtagtcgacccctcgatttgcggctctatatgaccacacc ctaaacaacatccccctctttcctccctccccctgctgtacaactA
cgTacgccctagataccacaggcacaatgcgggtcactaacaatcaaa
acagagccgaggggaattagaaaaaagtacgggtagaagagggatattc
agagatcagggcaagtctcccgagtctctgctctctcctctacctgat
agaccaggacaaacatggccaccttacagatgcagagatcgacgagc
tatttgagacaagtggaaactgtcattgacaacataattacagcccagg
gtaaaccagcagagactgttggaaggagtgaatcccacaaggcaaga
ccaaggtgctgagcgcagcatgggagaagcatgggagcatccagccac
cggccagtcaagacaaccccgatcgacaggacagatctgacaaacaac
catccacaccgagcaaacgaccccgcatgacagcccgcggccacat
ccgccgaccagccccccacccaggccacagacgaagccgtcgacacac
agCtcaggaccggagcaagcaactctctgctgttgatgcttgacaagc
tcagcaataaatcgccaatgctaaaaagggcccatggtcgagcccc
aagaggggaatcaccaacgtccgactcaacagcaggggagtcaacca
gtcgcggaaacagtcaggaaagaccgcagaaccaagtcaaggccgccc
ctggaaaccagggcacagacgtgaacacagcatatcatggacaatggg
aggagtcacaactatcagctgggtgcaaccctcatgctctccgatcaa
ggcagagccaagacaatacccttgatctgcggatcatgtccagccac
ctgtagactttgtgcaagc gatgtctatgatggaggcgatatcac agagagtaagtaagggtgactatcagctagatcttgtcttgaaacaga
catcctccatccctatgatgcgggtccgaaatccaacagctgaaaaacat
ctgttcgagtcatggaagccaacttgggaatgatgaagattctggatc ccggttgccaacatttcattctctgagtgatctacgggcagttgccc
gatctcacccggttttagtttcaggccctggagaccctctccctatg
tgacacaaggaggcgaaatggcacttaataaaactttcgcaaccagtgc
cacatccatctgaattgattaaacccgccactgcatgcgggcctgata
taggagtggaaaaggacactgtccgtgcattgatcatgtcacgccc
tgaccccgagttcttcagccaagctcctaagcaagttagatgcagccg
ggtcgatcgaggaaatcaggaaaatcaagcgcttgctctaaatggct
aattactactgccacacgtagcgggtccctgtccactcggcacacac
ggaatctgcaccgagttcccccccgGgTTAGAAAAAATACGGGTAGA
ACCGCCACCATGTCTTGGAAAGTGGTGATCATTTTTTTCATTGCTAATA
ACACCTCAACACGGTCTTAAAGAGAGCTACCTAGAAGAATCATGTAGC
ACTATAACTGAGGGATATCTTAGTGTTCTGAGGACAGGTTGGTATACC
AACGTTTTTACATTAGAGGTGGGTGATGTAGAAAACCTTACATGTTCT
GATGGACCTAGCCTAATAAAAAACAGAATTAGATCTGACCAAAAAGTGCA
CTAAGAGAGCTCAAAACAGTCTCTGCTGACCAATTGGCAAGAGAGGAA
CAAATTGAGAATCCCAGACAATCTAGGTTTGTCTAGGAGCAATAGCA
CTCGGTGTTGCAACAGCAGCTGCAGTCACAGCAGGTGTTGCAATTGCC
AAAACCATCCGGCTTGAGAGTGAAGTCACAGCAATTAAGAATGCCCTC
AAAACGACCAATGAAGCAGTATCTACATTGGGGAATGGAGTTCGAGTG
TTGGCAACTGCAGTGAGAGAGCTGAAAGACTTTGTGAGCAAGAATTA
ACTCGTGCAATCAACAAAAACAAGTGCGACATTGATGACCTAAAAATG
GCCGTTAGCTTCAGTCAATTCAACAGAAGGTTTCTAAATGTTGTGCGG

CAATTTTCAGACAATGCTGGAATAACACAGCAATATCTTTGGACTTA
ATGACAGATGCTGAACTAGCCAGGGCCGTTTCTAACATGCCGACATCT
GCAGGACAAATAAAATTGATGTTGGAGAACCGCGCGATGGTGCGAAGA
AAGGGGTTCGGAATCCTGATAGGGGTCTACGGGAGCTCTGTAATTTAC
ATGGTGCAGCTGCCAATCTTTGGCGTTATAGACACGCCTTGCTGGATA
GTAAAAGCAGCCCCCTTCTTGTTCCGAAAAAAAGGGAAACTATGCTTGC
CTCTTAAGAGAAGACCAAGGGTGGTATTGTCAGAATGCAGGGTCAACT
GTTTACTACCCAAATGAGAAAGACTGTGAAACAAGAGGAGACCATGTC
TTTTGCGACACAGCAGCAGGAATTAATGTTGCTGAGCAATCAAAGGAG
TGCAACATCAACATATCCACTACAAATTACCCATGCAAAGTCAGCACA
GGAAGACATCCTATCAGTATGGTTGCACTGTCTCCTCTTGGGGCTCTG
GTTGCTTGCTACAAAGGAGTAAGCTGTTCCATTGGCAGCAACAGAGTA
GGGATCATCAAGCAGCTGAACAAAGGTTGCTCCTATATAACCAACCAA
GATGCAGACACAGTGACAATAGACAACACTGTATATCAGCTAAGCAAA
GTTGAGGGTGAACAGCATGTTATAAAAGGCAGACCAGTGTCAAGCAGC
TTTGATCCAATCAAGTTTCCTGAAGATCAATTCAATGTTGCACTTGAC
CAAGTTTTTGAGAGCATTGAAAACAGCCAGGCCTTGGTAGATCAATCA
AACAGAATCCTAAGCAGTGCAGAGAAAGGGAATACTGGCTTCATCATT
GTAATAATTCTAATTGCTGTCCTTGGCTCTAGCATGATCCTAGTGAGC
ATCTTCATTATAATCAAGAAAACAAAGAAACCAACGGGAGCACCTCCA
GAGCTGAGTGGTGTCACAAACAATGGCTTCATACCACACAGTTAGGcc
gcggaccaaggtccaactctccaagcggcaatcctctctcgcttct
cagccccactgaatgAtcgcgtaaccgtaattaatctagctacattta
agattaagaaaaatacgggtagaattggagtgcccccaattgtgcaa gatggactcatctaggacaattgggctgtactttgattctgcccattc
ttctagcaacctgtagcatttccgatcgtcctacaagAcacaggaga
tgggaagaagcaaatcgccccgcaatataggatccagcgccttgactt
gtggactgatagtaaggaggactcagtattcatcaccacctatggatt
catctttcaagttgggaatgaagaagccacCgtcggcatgatcgatga
taaacccaagcgcgagttactttccgctgcgatgctctgcctaggaag
cgtcccaaataccggagaccttattgagctggcaagggcctgtctcac
tatgatagtcacatgcaagaagagtgaactaatactgagagaatggt
tttctcagtagtgaggcaccccaagtgtgcaaagctgtagggtgt
ggcaaacaataactcatcagtgatgcagtcaagcacgtgaaagcgcc
agagaagattcccgggagtggaaaccctagaatacaaggtgaactttgt
ctccttgactgtggtaccgaagaGggatgtctacaagatcccagctgc agtattgaaggtttctggctcgagtctgtacaatcttgcgctcaatgt
cactattaatgtggaggtagaccgaggagtcctttgggttaaattctCt gtctaagtctgacagcggatactatgctaacctcttcttgcatattgg
acttatgaccacTgtagataggaaggggaagaaagtacatttgacaa
gctggaaaagaaaataaggagccttgatctatctgtcgggctcagtg
tgtgctcgggccttccgtgttggtgtaaaagcaagaggtgcacggactaa gcttttggcacctttcttcttagcagtgggacagcctgctatcccat
agcaaagcttctcctcaggtggccaagatactctggagtcaaaccgc
gtgcctgcggagcgttaaaatcattatccaagcaggtacccaacgcgc
tgtcgcagtgaccgccgaccacgaggttaccttactaagctggagaa
ggggcacacccttgccaaatacaatccttttaagaaataagctgcgtc
tctgagattgcgctccgcccactcaccagatcatcatgacacaaaaa actaatctgtcttgattatttacagttagtttacctgtctatcaagtt
agaaaaaacacgggtagaagattctggatcccgggtggcgccctccag
gtgcaagatgggctccagaccttctaccaagaaccagcacctatgat
gctgactatccgggttgcgctgggtactgagttgcatctgtccggcaaa
ctccattgatggcaggccttctgcagctgcaggaattgtggttacagg
agacaaaagccgtcaacatatacacctcatcccagacaggatcaatcat
agttaagctcctcccgaatctgccaaggataaggaggcatgtgcgaa

agcccccttggatgcatacaacaggacattgaccactttgctcacccc
ccttggtgactctatccgtaggatacaagagtctgtgactacatctgg
aggggggagacagggggcgcccttataggcgccattattggcgggtgtggc
tcttgggggtgcaactgccgcacaaataacagcggccgcagctctgat
acaagccaaacaaaatgctgccaacatcctccgacttaaagagagcat
tgccgcaaccaatgaggctgtgcatgaggctactgacggattatcgca
actagcagtggtgagttgggaagatgcagcagtttgtaatgaccaatt
taataaacagctcaggaattagactgcatcaaaattgcacagcaagt
tggtgtagagctcaacctgtacctaaccgaattgactacagtattcgg accacaaatcacttcacctgcttaaacaaagctgactattcaggcact
ttacaatctagctgggtggaaatatggattacttattgactaagttagg tgtagggaacaatcaactcagctcattaatcggtagcggcctaatacac
cggtaaccctattctatacgactcacagactcaactcttgggtataca ggtaactctaccttcagtcgggaacctaataatatgcgtgccaccta
cttggaacctaataccgtaagcacaaccaggggatttgcctcggcact
tgtcccAaaagtgggtgacacaggtcggttctgtgatagaagaacttga cacctcatactgtatagaaactgacttagatttatattgtacaagaat
agtaacgttccctatgtcccctgggtatttattcctgcttgagcggcaa tacgtcggcctgtatgtactcaaagaccgaaggcgcacttactacacc
atacatgactatcaaaggttcagtcacgccaactgcaagatgacaac
atgtagatgtgtaaacccccgggtatcatatcgcaaaactatggaga agccgtgtctctaatagataaacaatcatgcaatgtttatccttagg
cgggataactttaaggctcagtggggaattcgatgtaacttatcagaa gaatatctcaatacaagattctcaagtaataataacaggcaatcttga
tatctcaactgagcttgggaatgtcaacaactcgatcagtaatgcttt
gaataagttagaggaaagcaacagaaaaactagacaaaagtcaatgtcaa actgactagcacatctgctctcattacctatatcgtttgactatcat
atctctgtttttgggtatacttagcctgattctagcatgctacctaata gtacaagcaaaaggcgaacaaaagaccttattatggcttggaataa
tactctagatcagatgagagccactacaaaaatgtgaacacagatgag gaacgaaggttccctaataagtaatttgtgtgaaagtctggtagtct
gtcagttcagagagttaagaaaaaactaccgggtgtagatgaccaaag
gacgatatacgggtagaacggtaagagaggccgcccctcaattgcgag
ccaggcttcacaacctccgttctaccgcttcaccgacaacagtcctca
atcatggaccgcgccgttagccaagtgcgttagagaatgatgaaaga
gaggcaaaaaatacatggcgcttgatattccggattgcaatcttattc ttaacagtagtgaccttggctatatctgtagcctcccttttatatagc
atggggggctagcacacctagcgatctttaggcataccgactaggatt
tccagggcagaagaaaagattacatctacacttgggtccaatcaagat
gtagtagataggatatataagcaagtggcccttgagtctccgttggca ttgttaaatactgagaccacaattatgaacgcaataacatctctct
tatcagattaatggagctgcaacaacagtggggtggggggcacctatc
catgaccagattatataggggggataggcaaagaactcattgtagat gatgctagtgtatgcacatcattctatccctctgcatttcaagaacat
ctgaattttatcccggcgctactacaggatcagggttgactcgaata cctcatttgacatgagtgtacccattactgtacaccataatgta
atattgtctggatgcagagatcactcacattcatatcagtattttagca cttggtgtgctccggacatctgcaacaggggagggtattcttttctact
ctgcgttccatcaacctggacgacacccaaaatcggaagtcttgag
gtgagtgaactcccctgggttgatgctgtgctcgaaagtcacg
gagacagaggaagaagattataactcagctgtccctacgcggatggta
catgggaggttaggggttcgacggccagtaccacgaaaaggacctagat
gtcacaacattattcggggactgggtggccaactaccagagtaggg
gggtggatctttattgacagccgcgtatgggttctcagcttacggaggg
ttaaaaccaattcacccagtgcactgtacaggaagggaatatgtg
atatacaagcgatacaatgacacatgccagatgagcaagactaccag
attcgaatggccaagtcttcgtataagcctggacgggttgggtgggaaa cgcatacagcaggctatcttatcaaggtgtcaacatccttaggc
gaagacccgggtactgactgtaccgccaacacagtcacactcatgggg
gccgaaggcagaattctcacagtagggacatctcatttctgtatcaa cgagggtcatcatacttctctcccgcgttatttatatcctatgacagtc
agcaacaaaacagccactcttcagtccttatacattcaatgccttc
actcggccaggtagtatcccttgccaggcttcagcaagatgccccaac tcgtgtgttactggaggtctatacagatccatatcccctaattcttctat
agaaaccacaccttgcgaggggtattcgggacaatgcttgatgggtga
caagcaagacttaacctgcgtctgcagtattcgatgcacatcccg
agtcgcattactcgagtgagttcaagcagtacaaagcagcatacaca acatcaacttgtttaaaagtggtcaagactaataagacctattgtctc
agcattgctgaaatatctaatactctcttcggagaattcagaatcgtc

ccgttactagttgagatcctcaaaagatgacggggttagagaagccagg
tctggctagttgagtcaattataaaggagttggaaagatggcattgta
tcacctatcttctgcgacatcaagaatcaaaccgaatgccggcgcgtg
ctcgaattccatgttgccagttgaccacaatcagccagtgctcatgcg atcagattaagccttgtcaAtaGtctcttgattaagaaaaaatgtaag
tggcaatgagatacaaggcaaaacagctcatgggTaaCaatacgggta
ggacatggcgagctccggctctgaaagggcagagcatcagattatcct
accagagTcacacctgtcttcaccattgggtcaagcaciaactactcta ttactggaaattaactgggctaccgcttctgatgaatgtgacttcca
ccacctattctcagccgacaatggaaaaaataacttgaatcggcctc
tcctgatactgagagaatgataaaactcggaagggcagtacaccaaac
tcttaaccacaattccagaataaccggagtgctccaccccagggtgtt
agaaGaactggctaattagggtcccagattcaaccaacaaatttcg
gaagattgagaagaagatccaaattcacaacacgagatatggagaact
gttcacaaggctgtgtacgcatatagagaagaaactgctgggggtcatc
ttgggtctaacaatgtccccgggtcagaggagttcagcagcattcgtac
ggatccggcattctggtttcactcaaaatgggtccacagccaagtttgc
atgggtccatataaaacagatccagaggcatctgatgggtggcagctaG
gacaaggctgcggccaacaaattgggtgatgctaaccataaggtagg
ccaagtctttgtcactcctgaacttgcgttgtgacgcatacgaatga gaacaagttcacatgtcttaccagggaactgtattgatgtatgcaga
tatgatggagggcagagatatgggtcaacataatatcaaccacggcgggt
gcatctcagaagcttatcagagaaaattgatgacatttgcgggttaat agacgctctggcaaaaagacttgggtaatcaagtctacgatgttgatc
actaatggagggatttgcatacggagctgtccagctactcgagccgtc
aggtacatttgcaggagatttcttcgattcaacctgcaggagcttaa agacattctaattggcctcctcccaatgatatagcagaatccgtgac
tcatgcaatcgctactgtattctctggtttagaacagaatcaagcagc tgagatgttgtgtctgttgcgtctgtgggggtcaccactgcttgagtc
ccgtattgcagcaaaggcagtcaggagccaaatgtgcgcaccgaaaat
ggtagactttgatatgatccttcagggtactgtctttctcaagggaac
aatcatcaacgggtacagaaagaagaatgcagggtgtgtggccgcgagt
caaagtggatacaatatatgggaagggtcattgggcaactacatgcaga
ttcagcagagatttcacacgatatcatgttgagagagtataagagttt atctgcacttgaatttgagccatgtatagaatatgacctgtcaccaa
cctgagcatgttctaaaagacaaggcaatcgcacaccccaacgataa
ttggcttgccctcgttaggcggaaccttctctccgaagaccagaagaa acatgtaaaagaagcaacttcgactaatcgccctcttgatagagtttt
agagtcaaattgatttgatccatataaagagatggaatatctgacgac ccttgagtaccttagagatgacaatgtggcagtatcatactcgctcaa
ggagaaggaagtgaaggttaattggacggatcttcgctaagctgacaaa
gaagttaagggaactgtcagggtgatggcggaagggtacctagccgatca
gattgcacctttcttcagggaaatggagtcattcagगतatgcatatc cttgaccaagagtatgctagcgatgagtcaactgtcttttaacagcaa
taagaaacgtatcactgactgtaaagaaagagtatcttcaaaccgcaa
tcatgatccgaaaagcaagaaccgtcggagagttgcaaccttcataac
aactgacctgcaaaagtactgtcttaattggagatatcagacaatcaa attgttcgctcatgccatcaatcagttgatgggcctacctcacttctt
cgaatggattcacctaagactgatggacactacgatgttcgtagggaga
ccctttcaatcctcaaagtgacctactgactgtgacctctcaagagt ccctaattgatgacatatatattgtcagtgccagaggggggtatcgaagg
attatgccagaagctatggacaatgatctcaattgtgcaatccaact tgctgcagctagatcgattgtcgtgttgccctgtatggtacaggggtga
taatcaagtaatagcagtaacgagagaggttaagatcagacgactctcc
ggagatgggtgttgacacagttgcatcaagccagtgataatttctcaa ggaattaattcatgtcaatcatttgattggccataatttgaaggatcg
tgaaaccatcaggtcagacacattcttcatatacagcaaacgaatctt caaagatggagcaatcctcagtcgaagtcctcaaaaattcatctaaatt
agtgctagtgtcagggtgatctcagtgaaaacaccgtaatgtcctgtgc
caacattgcctctactgtagcacggctatgcgagaacgggcttcccaa agacttctgttactattttaaactatataatgagttgtgtgcagacata
ctttgactctgagttctccatcaccaacaattcgcaccccgatcttaa tcagtcgtggattgaggacatctcttttgcactcatatgttctgac
tcctgccaattagggggactgagtaaccttcaataactcaaggctcta
cactagaaatatcggtgacccggggactactgcttttgagagatcaa
gcgactagaagcagtgaggattactgagtcctaacattatgactaatat
cttaactaggccgctgggaatggagattgggccaagtctgtgcaacga ccatactctttcaattttgagactgttgcaagcccaaatattgttct

taagaaacatacgcaaagagtcctatttgaaacttggtcaaaccctt
attgtctggagtgcacacagaggataatgaggcagaagagaaggcatt
ggctgaattcttgcttaatcaagagggtattcatccccgcgttgcgca
tgccatcatggaggcaagctctgtaggtaggagaaagcaaattcaagg
gcttggtgacacaacaaacaccgtaattaagattgcgcttactaggag
gccattaggcatcaagaggctgatgcggatagtcaattattctagcat gcatgcaatgctgttttagagacgatgttttccctccagtagatccaa
cccccccttagtctcttctaatatgtgttctctgacactggcagacta
tgcacggaatagaagctgggtcacctttgacgggaggcaggaaaatact
gggtgtatctaactctgatacgatagaactcgtagagggtgagattct
tagtgtaagcggagggtgtacaagatgtgacagcggagatgaacaatt
tacttggttccatcttccaagaatatagaattgaccgatgacaccag
caagaatcctccgatgagggtaccatatctcgggtcaaagacacagga
gaggagagctgcctcacttgcaaaaatagctcatatgtcgccacatgt
aaaggctgccctaaggggcatcatccgtgttgatctgggcttatgggga taatgaagtaaattggactgctgctcttacgattgcaaaatctcgggtg
taatgtaaacttagagtatcttcggttactgtcccccttaccacggc tgggaatctcaacatagactagatgatggtataactcagatgacatt
caccctgcatctctctacaggGtgtcaccttacattcacatatccaa
tgatttctcaaaggctgttactgaagaaggagtgaaagaggggaatgt ggttaccaacagatcatgctcttgggtttatctctaactgaatcgat
ctttccaatgacaacaaccaggacatatgatgagatcacactgcacct acatagtaaatttagttgctgtatcagagaagcacctgttgcggttcc
tttcgagctacttgggggtgggtaccggaactgaggacagtgacctcaaa
taagtttatgtatgatccttagccctgtatcggaggggagactttgcgag acttgacttagctatcttcaagagttatgagcttaactctggagtcata
tcccacgatagagctaataaacattcttcaatatccagcgggaagt gattggccagctctgtggtttcttatgatgaagatacctccataaagaa
tgacgccataatagtgtatgacaatacccgaattggatcagtgaagc tcagaattcagatgtggtcgcctatttgaatatgcagcacttgaagt
gctcctcgactgttcttaccactctattacctgagagtaagaggcct Ggacaatattgtcttatatatgggtgattatacaagaatatgccagg
aattctactttccaacattgcagctacaatatctcatcccgctcattca ttcaaggttacatgcagtgggcctgggtcaaccatgacggatcacacca
acttgacgatacggattttatcgaaaatgtctgcaaaactattagtatc ttgcacccgacgtgtgatctccggcttatattcaggaaataagtatga
tctgctgttcccatctgtcttagatgataacctgaatgagaagatgct tcagctgatatcccgggttatgctgtctgtacacgggtactctttgctac
aacaagagaaaatcccgaataaagagggttaactgcagaagagaaaatg ttcaatactcactgagtatttactgtcggatgtgtgaaaccattact
tagccccgatcaagtgagctctatcatgtctcctaacataattacatt cccagctaactctgtactacatgtctcgggaagagcctcaatttgatcag
ggaaagggaggacagggatactatcctggcggtgtgttcccccaaga
gccattattagagttcccttctgtgcaagatattgggtgctcgagtga agatccattcacccgacaacctgcggcatttttgaagagtttagatt
gagtgtccagcaagggtatgacgcattcacacttagtcagattcatcc tgaactcacatctccaaatccggaggaagactacttagtacgatactt
gttcagaggggatagggtgcatcttcctcttggtataaggcatctca tctcctttctgtacccgaggtgaagatgtgcaagacacgggaactcctt
atacttagctgaaggggagcggagccatcatgagtcttctcgaactgca tgtaccacatgaaactatctattacaatacgtcttttcaaatgagat
gaacccccgcgaacgacatttcgggcccaccccaactcagttttgaa
ttcgggtgtttataggaatctacaggcggaggtaacatgcaaagatgg
attgtccaagagttccgtccattatggagagaaaatacagaggaaaag
CgacctgacctcagataaagTagtgggggtatattacatctgcagtgcc ctacagatctgtatcattgctgcattgtgacattgaaattcctccagg
gtccaatcaaagcttactagatcaactagctatcaatttatctctgat tgccatgcattctgtaaggaggggcggggttagtaatcatcaaagtgtt
gtatgcaatgggatactactttcatctactcatgaactgtttgctcc gtgttcacaaaaggatatattctctctaattggttatgcatgtcgagg
agatatggagtgttacctgggtatttgcaggttacctgggcccggcc
tacattgtacatgaggtggtgaggatggcGaaaactctggtgcagcg
gcacgggtacgctTttgtctaaatcagatgagatcacactgaccaggtt
attcacctcacagcggcagcgtgtgacagacatcctatccagtcctt
accaagattaataaagtacttgaggaagaatattgacactgcgctgat
tgaagccggggggacagccccgtccgtccattctgtgaggagagtggt
gagcacgctagcgaacataactcagataaccagatCatcgctagtca
cattgacacagttatccggtctgtgatataatggaagctgagggtga tctcgctgacacagttatttctatttacccttacaatctctctactga
cgggaaaaagaggacatcacttaAacagtgcacgagacagatcctaga
gggtacaatactagggtcttagagtcgaaaatctcaataaaaataggcga tataatcagcctagtgtctaaaggcatgatctccatggaggaccttat
cccactaaggacatacttgaagcatagtagctgcctaaatatttgaa ggctgtcctagggtattaccaaaactcaaagaaatgtttacagacacttc

tgtaCtgacttgactcgtgctcaaaaaatttcatatgaaaactat aggcattgcagtcacaaaggatattacagtaactgtgactcttaacgaaa
atcacatattaataggctccttttttggccaattgtattcttgttgat ttaatcatattatgtagaaaaaagtgaaccctgactccttaggact
cgaattcgaactcaataaatgtcttaaaaaaagggttgcgcacaatta ttcttgagtgtagtctcgtcattcacaaatctttgtttggt cDNA
of accaaacagagaatccgtgagttacgataaaaggcgaaggagcaattg 21 genomic
aagtcgcacgggtagaagggtgtaatctcgagtgcgagcccgaagcac sequence of
aaactcgagaaaagccttctgccaacatgtcttccgtatttgatgagta NDV LaSota
cgaacagctcctcgcggctcagactcgcccaatggagctcatggagg strain with the
gggagaaaaagggagtagcttaaaagtagacgtcccggtattcactct codon optimized
taacagtgatgaccagaagatagatggagctttgtggtattctgcct nucleic acid
ccggattgctgttagcgaagatgccaacaaaccactcaggcaaggtgc encoding hMPV
tctcatatctctttatgctcccactcacaggaatgaggaacatgt F protein of
tgccCttgcagggaacagaatgaagccacattggccgtgcttgagat CAN00-16
tgatggctttgccaacggcacgccccagttcaacaataggagtggagt inserted (Sac II
gtctgaagagagagcacagagatttgcgatgatagcaggatctctccc restriction sites
tcgggcatgcagcaacggaacccgttcgtcacagccggggcCgaaga used to insert the
tgatgcaccagaagacatcaccgataccctggagaggatcctctctat codon optimized
ccaggctcaagtatgggtcacagtagcaaaagccatgactgcgtatga hMPV F protein
gactgcagatgagtgcgaaacaaggcgaatcaataagtatatgcagca are shown by
aggcagggtccaaaagaaatacatcctctaccccgatgcaggagcac double
aatccaactcacgatcagacagctcttgcagtcgcgatcttttgggt underlining. The
tagcgagctcaagagaggccgcaacacggcaggtggtaccttactta initiation and
ttataacctggtaggggacgtagactcatacatcaggaataccgggct stop codons of
tactgcattcttctgacactcaagtacggaatcaacaccaagacatc the codon
agcccttgacttagtagcctctcaggcgacatccagaagatgaagca optimized hMPV
gctcatgcgtttgtatcggtgaaaaggagataatgcgccgtacatgac F are underlined.
attacttggtgatagtaccagatgagctttgcgcctgccgagtatgc The open reading
acaactttactcctttgccatgggtatggcatcagtcctagataaagg frame of the
tactgggaaataccaatttgccagggactttatgagcacatcattctg codon optimized
gagacttgagtagagtacgctcaggctcagggaagtagcattaacga hMPV F is in
ggatatggctgccgagctaaagctaaccacagcagcaaGgaGgggcct bold.)
ggcagctgctgcccaacgggtctccgaGgaGaccagcagcataGacat
gcctactcaacaagtcggagtcctcactgggcttagcgaggggggggtc
ccaagctctacaaggcggatcgaatagatcgcaagggaaccagaagc
cggggatggggagacccaattcctggatctgatgagagcggtagcaaa
tagcatgaggggagcgccaaactctgcacagggcactcccaatcggg
gcctccccaaactcctgggccatcccaagataacgacaccgactgggg
gtattgatggacaaaaccagcctgcttcacaaaaacatcccaatgc
cctcacccgtagtcgacccctcgatttgcggctctatatgaccacacc ctcaaacaaacatccccctcttctcctccccctgctgtacaactA
cgTacgccctagataccacaggcacaatgcggctcactaacaatcaaa
acagagccgagggaattagaaaaaagtacgggtagaagagggatattc
agagatcagggaagctccccgagtctctgctctctctctacctgat
agaccaggacaaacatggccacctttacagatgcagagatcgacgagc
tatttgagacaagtggaactgtcattgacaacataattacagcccagg
gtaaaccagcagagactgttggaaggagtgaatcccacaaggcaaga
ccaaggtgctgagcgcagcatgggagaagcatgggagcatccagccac
cggccagtcgaagacaaccccgatcgacaggacagatctgacaaacaac
catccacaccgagcaaacgaccccgcatgacagcccggccacat
ccgccgaccagccccccacccaggccacagacgaagccgtcgacacac
agCtcaggaccggagcaagcaactctctgctgttgatgcttgacaagc
tcagcaataaatcgccaatgctaaaaagggcccatggctcgagcccc

aagaggggaatcaccaacgtccgactcaacagcaggggagtcaccca
gtcgcggaaacagtcaggaaagaccgcagaaccaagtcaggccgccc
ctggaaaccagggcacagacgtgaacacagcatatcatggacaatggg
aggagtcacaactatcagctgggtgcaaccctcatgctctccgatcaa
ggcagagccaagacaatacccttgtatctgcggatcatgtccagccac
ctgtagactttgtgcaagcgatgatgtctatgatggaggcgatatcac agagagtaagtaagggtgactatcagctagatcttgtcttgaaacaga
catcctccatccctatgatgcgggtccgaaatccaacagctgaaaacat
ctgttgcatgcatggaagccaacttgggaatgatgaagattctggatc ccggtgtgccaacatttcatctctgagtgatctacgggcagttgcc
gatctcaccgggttttagtttcaggccctggagaccctctccctatg
tgacacaaggaggcgaaatggcacttaataaaactttcgcaaccagtg
cacatccatctgaattgattaaacccgccactgcatgcgggcctgata
taggagtggaagaggacactgtccgtgcattgatcatgtcacgccc
tgcacccgagttcttcagccaagctcctaagcaagttagatgcagccg
ggtcgatcgaggaaatcaggaaaatcaagcgcccttgctctaaatggct
aattactactgccacacgtagcgggtccctgtccactcggcacacac
ggaatctgcaccgagttccccccgcGgTTAGAAAAAATACGGGTAGA
ACCGCCACCATGAGCTGGAAGGTGGTGATCATCTTCAGCCTGCTGATC
ACCCCCAGCACGGCCTGAAGGAGAGCTACCTGGAGGAGAGCTGCAGC
ACCATCACCGAGGGCTACCTGAGCGTGCTGAGAACCGGCTGGTACACC
AACGTGTTCACCCTGGAGGTGGGCGACGTGGAGAACCTGACCTGCAGC
GACGGCCCCAGCCTGATCAAGACCGAGCTGGACCTGACCAAGAGCGCC
CTGAGAGAGCTGAAGACCGTGAGCGCCGACCAGCTGGCCAGAGAGGAG
CAGATCGAGAACCCAGACAGAGCAGATTCTGTGCTGGGCGCCATCGCC
CTGGGCGTGGCCACCGCCGCGCGCGTGACCGCCGGCGTGGCCATCGCC
AAGACCATCAGACTGGAGAGCGAGGTGACCGCCATCAAGAACGCCCTG
AAGACCACCAACGAGGGCCGTGAGCACCCCTGGGCAACGGCGTGAGAGTG
CTGGCCACCGCCGTGAGAGAGCTGAAGGACTTCGTGAGCAAGAACCTG
ACCAGAGCCATCAACAAGAACAAGTGCGACATCGACGACCTGAAGATG
GCCGTGAGCTTCAGCCAGTTCAACAGAAGATTCCTGAACGTGGTGAGA
CAGTTCAGCGACAACGCCGGCATCACCCCGCCATCAGCCTGGACCTG
ATGACCGACGCCGAGCTGGCCAGAGCCGTGAGCAACATGCCCACCAGC
GCCGGCCAGATCAAGCTGATGCTGGAGAACAGAGCCATGGTGAGAAGA
AAGGGCTTCGGCATCCTGATCGGCGTGACGGCAGCAGCGTGATCTAC
ATGGTGACAGCTGCCCATCTTCGGCGTGATCGACACCCCTGCTGGATC
GTGAAGGCCGCCCCAGCTGCAGCGAGAAGAAGGGCAACTACGCCTGC
CTGCTGAGAGAGGACCAGGGCTGGTACTGCCAGAACGCCGGCAGCACC
GTGTACTACCCCAACGAGAAGGACTGCGAGACCAGAGGCGACACGTG
TTCTGCGACACCGCCGCCGGCATCAACGTGGCCGAGCAGAGCAAGGAG
TGCAACATCAACATCAGCACCACTAACCCTGCAAGGTGAGCACC
GGCAGACACCCCATCAGCATGGTGGCCCTGAGCCCCCTGGGCGCCCTG
GTGGCCTGCTACAAGGGCGTGAGCTGCAGCATCGGCAGCAACAGAGTG
GGCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCACCAACCAG
GACGCCGACACCGTGACCATCGACAACACCGTGTACCAGCTGAGCAAG
GTGGAGGGCGAGCAGCACGTGATCAAGGGCAGACCCGTGAGCAGCAGC
TTCGACCCCATCAAGTTCCCCGAGGACCAGTTCAACGTGGCCCTGGAC
CAGGTGTTTCGAGAGCATCGAGAACAGCCAGGCCCTGGTGGACCAGAGC
AACAGAATCCTGAGCAGCGCCGAGAAGGGCAACACCGGCTTCATCATC
GTGATCATCCTGATCGCCGTGCTGGGCAGCAGCATGATCCTGGTGAGC
ATCTTCATCATCATCAAGAAGACCAAGAAGCCCACCGGCGCCCCCCCC
GAGCTGAGCGGCGTGACCAACAACGGCTTCATCCCCACAGCTGAGcc

gcggaccgaaggtccaactctccaagcggaatcctctcgtcttcc
cagccccactgaatgAtcgcgtaaccgtaattaatctagctacattta
agattaagaaaaaacacgggtagaattggagtgcccaattgtgccaa gatggactcatctaggacaattgggctgtactttgattctgccattc
ttctagcaacctgtagcatttccgatcgtcctacaagAcacaggaga
tggaagaagcaaactgccccgcaatataggatccagcgcttgactt
gtggactgatagtaaggaggactcagtattcatcaccacctatggatt
catctttcaagttgggaatgaagaagccacCgtcggcatgatgatga
taaaccgaagcgcgagttactttccgctgcgatgctctgcctaggaag
cgtcccaataaccggagaccttattgagctggcaagggcctgtctcac
tatgatagtcacatgaagaagagtgcaactaatactgagagaatggt
tttctcagtagtgaggcaccccaagtgtgcaaagctgtagggtgt
ggcaaaacaaatactcatcagtgaatgcagtcaagcacgtgaaagcgcc
agagaagattcccgggagtggaaccctagaatacaaggtgaactttgt
ctccttgactgtggtaccgaagaGggatgtctacaagatcccagctgc agtattgaaggtttctggctcgagtctgtacaatcttgcgctcaatgt
cactattaatgtggaggtagaccgaggagtcctttgggttaaatctCt gtctaaagtctgacagcggatactatgctaacctcttcttgcatttgg
acttatgaccacTgtagataggaaggggaagaaagtacatttgaaa
gctggaaaagaaaataaggagccttgatctatctgtcgggctcagtga
tgtgctcgggcttccgtgttggttaaaagcaagaggtgcacggactaa gcttttggcacctttcttcttagcagtgggacagcctgctatcccat
agcaaagtcttctcctcaggtggccaagatactctggagtcacacccgc
gtgcctgcggagcgttaaaatcattatccaagcaggtacccaacgcgc
tgtcgagtgaccgccgaccacgaggttaccttactaagctggagaa
ggggcacacccttgccaaatacaatccttttaagaaataagctgcgtc
tctgagattgcgctccgcccactcaccagatcatcatgacacaaaaa actaatctgtcttgattatttacagtttagttacctgtctatcaagtt
agaaaaaacacgggtagaagattctggatcccgggtggcgccctccag
gtgcaagatgggctccagaccttctaccaagaacccagcacctatgat
gctgactatccgggttgctgtgtactgagttgcatctgtccggcaaa
ctccattgatggcaggcctcttgagctgcaggaattgtggttacagg
agacaaagccgtcaacatatacacctcatcccagacaggatcaatcat
agttaagctcctcccgaatctgccaagataaggagggcatgtgcgaa
agcccccttggtatgcatacaacaggacattgaccactttgctcacc
ccttggtgactctatccgtaggatacaagagtctgtgactacatctgg
agggggggagacagggggcgctttagggcgccattattggcggtgtggc
tcttggggttgcaactgccgcacaaataacagcggccgcagctctgat
acaagccaaacaaaatgctgccaacatcctccgacttaaagagagcat
tgccgcaaccaatgaggctgtgcatgaggtcactgacggattatcgca
actagcagtggcagttgggaagatgcagcagttgttaatgaccaatt
taataaaacagctcaggaattagactgcatcaaaattgcacagcaagt
tggtgtagagctcaacctgtacctaaccgaattgactacagtattcgg accacaaatcacttcacctgctttaacaagctgactattcaggcact
ttacaatctagctggtggaaatattgattacttattgactaagttagg tgtagggaacaatcaactcagctcattaatcggtagcggcttaacac
cggtaaccctattctatacgaactcacagactcaactcttgggtatata ggtaactctaccttcagtcgggaacctaaataatgctgcccaccta
cttggaaacctatccgtaagcacaccaggggatttgcctcggcact
tgtcccAaaagtgggtgacacaggtcgggtctgtgataagaacttga cacctcatactgtatagaaactgacttagatttatattgtacaagaat
agtaacgttccctatgtcccctgggtatttattcctgcttgagcggcaa tacgtcggcctgtatgtactcaaagaccgaaggcgcacttactacacc
atacatgactatcaaagggtcagtcacgccaactgcaagatgacaac
atgtagatgtgtaaacccccgggtatcatatcgcaaaactatggaga agccgtgtctctaatagataaacaatcatgcaatgtttatccttagg
cgggataactttaaggctcagtggggaattcgatgtaacttatcagaa gaatatctcaatacaagattctcaagtaataataacaggcaatcttga
tatctcaactgagcttgggaatgtacaactcgatcagtaattgcttt
gaataagttagaggaaagcaacagaaaaactagacaaagtcaatgtcaa actgactagcacatctgctctcattacctatcgttttgactatcat
atctctgtttttgggtatacttagcctgattctagcatgctacctaata gtacaagcaaaaggcgcaacaaaagaccttattatggccttggaataa
tactctagatcagatgagagccactacaaaaatgtgaacacagatgag gaacgaaggtttccctaataagtaatttgtgtgaaagttctggtagtct

gtcagttcagagagttaagaaaaaactaccggttgtagatgaccaaag
gacgatatacgggtagaacggtaagagaggccgcccctcaattgcgag
ccaggcttcacaacctccgttctaccgcttcaccgacaacagtcctca
atcatggaccgcgccgttagccaagttgcgttagagaatgatgaaaga
gaggcaaaaaatacatggcgcttgatattccggattgcaatcttattc ttaacagtagtgaccttggctatatctgtagcctcccttttatatagc
atgggggctagcacacctagcgatctttagggcataccgactaggatt
tccagggcagaagaaaagattacatctacacttgggttccaatcaagat
gtagtagataggatatataagcaagtggcccttgagtctccgttggca ttgttaaatactgagaccacaattatgaacgcaataacatctctctct
tatcagattaatggagctgcaaacaacagtggggtggggggcacctatc
catgaccagattatataggggggataggcaaagaactcattgtagat gatgctagtgatgtcacatcattctatccctctgcatttcaagaacat
ctgaattttatcccggcgccctactacaggatcagggttgactcgaata ccctcatttgacatgagtgtacccattactgctacaccataatgta
atattgtctggatgcagagatcactcacattcatatcagtattttagca ctgggtgtgctccggacatctgcaacagggaggggtattcttttctact
ctgcgttccatcaacctggacgacacccaaaaatcggaagtcttgagat
gtgagtgcaactcccctgggttgtagatgctgtgctcgaaagtcacg
gagacagaggaagaagattataactcagctgtccctacgcggatggta
catgggaggttaggggtcgacggccagtaccacgaaaaggacntagat
gtcacaacattattcggggactgggtggccaactaccagagtaggg
gggtggatctttattgacagccgcgtatgggttctcagttctacggaggg
ttaaaccctaacctcaccagtgacactgtacaggaagggaatatgtg
atatacaagcgatacaatgacacatgccagatgagcaagactaccag
attcgaatggccaagtcttcgtataagcctggacgggttgggtgggaaa cgcatacagcaggctatcttatctatcaaggtgtcaacatccttaggc
gaagaccgggtactgactgtaccgccaacacagtcacactcatgggg
gccgaaggcagaattctcacagtagggacatctcatttctgtatcaa cgagggtcatcatacttctctcccgcgttattatatcctatgacagtc
agcaaaaaacagccactcttcatagtccttatacattcaatgccttc
actcgccaggtagtatcccttgccaggcttcagcaagatgccccaac tcgtgtgttactggagtgctatacagatccatatcccctaattctctat
agaaaccacaccttgcgaggggtattcgggacaatgcttgatgggtga
caagcaagacttaaccctgcgtctgcagttatcgatgacatcccgc
agtcgcattactcgagtgagttcaagcagtaacaaagcagcatacaca acatcaactgttttaaagtgggtcaagactaataagacctattgtctc
agcattgctgaaatatctaatactctcttcggagaattcagaatcgtc
ccgttactagttgagatcctcaaagatgacgggggttagagaagccagg
tctggctagttgagtcattataaaggagttggaaagatggcattgta
tcacctatcttctgcgacatcaagaatcaaaccgaatgccggcgcgctg
ctcgaattccatgttgccagttgaccacaatcagccagtgctcatgcg atcagattaagccttgtcaAtaGtctcttgattaagaaaaaatgtaag
tggcaatgagatacaaggcaaaacagctcatgggTaaCaatacgggta
ggacatggcgagctccgggtcctgaaagggcagagcatcagattatcct
accagagTcacacctgtcttcaccattgggtcaagcacaactactcta ttactggaaattaactgggctaccgcttctgatgaatgtgacttcca
ccacctattctcagccgacaatggaaaaaataacttgaatcggcctc
tcctgatactgagagaatgataaaactcggaagggcagtacaccaaac
tcttaaccacaattccagaataaccggagtgctccaccccaggtgtt
agaaGaactggctaattagaggtcccagattcaaccaacaaatttcg
gaagattgagaagaagatccaaattcacaacacgagatatggagaact
gttcacaaggctgtgtacgcatatagagaagaaactgctggggctatc
ttggctacaacatgtccccgggtcagaggagttcagcagcattcgtac
ggatccggcattctggtttcactcaaaatgggtccacagccaagtgtgc
atggctccatataaaacagatccagaggcatctgatgggtggcagctaG
gacaaggctgcggccaacaaattgggtgatgctaaccataaggtagg
ccaagtcttctcactcctgaacttgcgttgtgacgcatacgaatga gaacaagttcacatgtcttaccaggaacttgattgatgtatgcaga
tatgatggagggcagagatatggtaacataatatcaaccacggcggt
gcattctcagaagcttatcagagaaaattgatgacatttgcgggttaat agacgctctggcaaaaagacttgggtaatacagctacgatgttgatc
actaatggagggatttcatacggagctgtccagctactcagccgtc

aggtagcatttgcaggagatttctcgcattcaacctgcaggagcttaa agacatttcaattggcctcctcccaatgatatagcagaatccgtgac
tcatgcaatcgcactgtattctctggtttagaacagaatcaagcagc tgagatgttgtgtctgttgcgtctgtgggggtcaccactgcttgagtc
ccgtattgcagcaaaggcagtcaggagccaaatgtgcgcaccgaaaat
ggtagactttgatatgatccttcaggtactgtctttctcaagggaac
aatcatcaacgggtacagaaagaagaatgcagggtgtgtggccgcgagt
caaagtggatacaatatatgggaagggtcattgggcaactacatgcaga
ttcagcagagatttcacacgatatcatgttgagagagtataagagttt atctgcacttgaatttgagccatgtatagaatatgacctgtcaccaa
cctgagcatgttcttaaaagacaaggcaatcgcacaccccaacgataa
ttggcttgcctcgttttaggcggaaccttctctccgaagaccagaagaa acatgtaaaagaagcaacttcgactaatcgctcttgatagagtttt
agagtcaaatgattttgatccatataaagagatggaatatctgacgac cttgagtagccttagagatgacaatgtggcagtatcatactcgctcaa
ggagaaggaagtgaagttaatggacggatcttcgctaagctgacaaa
gaagttaaggaactgtcaggtgatggcggaagggtacctagccgatca
gattgcacctttcttcagggaaatggagtcattcaggatagcatatc cttgaccaagagtatgctagcgtatgagtcactgtctttaacagcaa
taagaaacgtatcactgactgtaaagaaagagtatcttcaaaccgcaa
tcatgatccgaaaagcaagaaccgtcggagagttgcaaccttcataac
aactgacctgcaaaagtactgtcttaattggagatatcagacaatcaa attgttcgctcatgccatcaatcagttgatgggcctacctcacttctt
cgaatggattcacctaagactgatggacactacgatgttcgtagggaga
ccctttcaatcctccaagtgcacctactgactgtgacctctcaagagt ccctaattgatgacatatattgtcagtgccagaggggggtatcgaagg
attatgccagaagctatggacaatgatctcaattgctgcaatccaact tgctgcagctagatcgcatgtcgtgttgctgtatggtacaggggtga
taatcaagtaatagcagtaacgagagaggttaagatcagacgactctcc
ggagatgggtgttgacacagttgcatcaagccagtgataatttctcaa ggaattaattcatgtcaatcatttgattggccataattgaaggatcg
tgaaaccatcaggtcagacacattcttcatatacagcaaacgaatctt caaagatggagcaatcctcagtcgaagtcctcaaaaattcatctaaatt
agtgtcagtgatcaggtgatctcagtgaaaacaccgtaattgtcctgtgc
caacattgcctctactgtagcacggctatgcgagaacgggcttccaa agacttctgttactatttaaactatataatgagttgtgtgcagacata
ctttgactctgagttctccatcaccaacaattcgcaccccgatcttaa tcagtcgtggattgaggacatctcttttgcactcatatgttctgac
tcctgccaattagggggactgagtaaccttcaatactcaaggctcta
cactagaaatatcggtagcccggggactactgctttgcagagatcaa
gcgactagaagcagtgaggattactgagtcttaacattatgactaatat
cttaactaggccgctgggaatggagattggggcagctgtgcaacga ccatactctttcaattttgagactgttgcaagcccaaatattgttct
taagaaacatacgcaaagagtcctatttgaaactgttcaaatccctt
attgtctggagtgacacagaggataatgaggcagaagagaaggcatt
ggctgaattcttgcttaataagaggtgattcatccccgcgttgcgca
tgccatcatggaggcaagctctgtaggtaggagaaaagcaaattcaagg
gcttgtgacacaacaaacaccgtaattaagattgcgcttactaggag
gccattaggcatcaagaggctgatgcggatagtcattattctagcat gcatgcaatgctgttttagagacgatgttttctcctcagtagatccaa
ccacccttagtctcttctaatatgtgttctctgacactggcagacta
tgcacggaatagaagctggtcacctttgacgggaggcaggaaaatact
gggtgtatctaactctgatacgaataactcgtagaggggtgagattct
tagtgaagcggagggtgtacaagatgtgacagcggagatgaacaatt
tacttggttccatcttccaagcaatatagaattgaccgatgacaccag
caagaatcctccgatgagggtaccatatctcgggtcaaagacacagga
gaggagagctgcctcacttgcaaaaatagctcatatgtcgccacatgt
aaaggctgcctaagggcacatccgtgttgatctgggcttatgggga taatgaagtaaattggactgctgctcttacgattgcaaaatctcggtg
taatgtaaacttagagtatcttcggttactgtccctttaccacggc tgggaaatcttcaacatagactagatgatggtataactcagatgacatt
caccctgcatctctacaggGtgtcaccttacattcacatatccaa
tgattctcaaaggctgttactgaagaaggagtcaaagaggggaatgt ggttaccaacagatcatgctcttgggtttatcttaatcgaatcgat
ctttccaatgacaacaaccaggacatatgatgagatcacactgcacct acatagtaaatttagttgctgtatcagagaagcacctgttgcggttcc
tttcgagctacttgggggggtaccggaactgaggacagtgacctcaaa taagtttatgtatgatcctagccctgtatcgaggaggagactttgcgag
acttgacttagctatcttcaagagttatgagcttaatctggagtcata tcccacgatagagctaataacattctttcaatatccagcgggaagtt
gattggccagtcgttggtttcttatgatgaagatacctccataaagaa tgacgccataatagtgtatgacaatacccgaattggatcagtgaaagc

tcagaattcagatgtgggtccgcctatttgaatatgcagcacttgaagt gctcctcgactgttcttaccacactctattacctgagagtaagaggcct
Ggacaatattgtcttatatatgggtgatttatacaagaatatgccagg aattctactttccaacattgcagctacaatatctcatcccgctcattca
ttcaaggttacatgcagtggtggcctgggtcaaccatgacggatcacacca acttgcagatacggattttatcgaaatgtctgcaaaactattagtatc
ttgcacccgacgtgtgatctccggccttatattcaggaaataagtatga tctgctgttcccatctgtcttagatgataacctgaatgagaagatgct
tcagctgatatcccggttatgctgtctgtacacgggtactctttgctac
aacaagagaaaatcccgaaaataagaggcttaactgcagaagagaaatg
ttcaatactcactgagtatttactgtcggatgctgtgaaaccattact tagccccgatcaagtgagctctatcatgtctcctaacataattacatt
cccagctaattctgtactacatgtctcgggaagagcctcaatttgatcag
ggaaaagggaggacagggatactatcctggcggtgtgttcccccaaga
gccattattagagttcccttctgtgcaagatattgggtgctcgagtga agatccattcacccgacaacctgcggcattttgcaagagttagattt
gagtgtccagcaaggtatgacgcattcacacttagtcagattcatcc tgaactcacatctccaaatccggaggaagactacttagtacgatactt
gttcagaggggatagggactgcatcttctcttgggtataaggcatctca tctccttctgtacccgaggtgaagatgtgcaagacacgggaactcctt
atacttagctgaagggagcggagccatcatgagtcttctcgaactgca tgtaccacatgaaactatctattacaatacgtcttttcaaatgagat
gaaccccccgcaacgacatttcggggccgacccaactcagttttgaa
ttcggtgtttataggaatctacaggcggaggtaacatgcaaagatgg
atttgtccaagagttccgtccattatggagagaaaatacagaggaaaag
CgacctgacctcagataaagTagtgggggtatattacatctgcagtgcc ctacagatctgtatcattgtctgcattgtgacattgaaattcctccagg
gtccaatcaaagcttactagatcaactagctatcaatttatctctgat tgccatgcattctgtaagggagggcggggtagtaatcatcaaagtgtt
gtatgcaatgggatactactttcatctactcatgaactgtttgctcc gtgtccacaaaaggatataattctctctaattgggtatgcatgtcgagg
agatatggagtggttacctgggtatttgcagtggttacctggggggcc
tacattgtacatgaggtgggtgaggatggcGaaaactctgggtgcagcg
gcacgggtacgctTttgtctaaatcagatgagatcacactgaccaggtt
attcacctcacagcggcagcgtgtgacagacatcctatccagtccttt
accaagattaataaagtacttgaggaagaatattgacactgcgctgat
tgaagccgggggacagcccgtccgtccattctgtgcggagagtggtt
gagcacgctagcgaacataactcagataaccagatCatcgctagtca
cattgacacagttatccgggtctgtgatatatatggaagctgaggggtga tctcgtgacacagttattctatttacccttacaatctctctactga
cgggaaaaagaggacatcacttaAacagtgcacgagacagatcctaga
ggttacaatactaggtcttagagtcgaaaatctcaataaaataggcga tataatcagcctagtgcttaaaggcatgatctccatggaggaccttat
cccactaaggacatacttgaagcatagtagctgcctaaatatttgaa ggctgtcctaggtattaccaaactcaaagaaatgtttacagacacttc
tgtaCtgtacttgactcgtgtcaacaaaaattctacatgaaaactat aggcaatgcagtcaaaggatattacagtaactgtgactcttaacgaaa
atcacatattaataggctcctttttggccaattgtattcttgttgat ttaatcatattatgtagaaaaaagttgaaccctgactccttaggact
cgaattcgaactcaaataaatgtcttaaaaaaagggtgcgcacaaatta ttcttgagtgtagtctcgtcattcaccaaacttttgtttggt
(190) TABLE-US-00006 Nucleic acid

ATGTCTTGGAAAGTGGTGATCATTTTTTTCATTGCTAATAACACCTCAACACG 32 sequence
GTCTTAAAGAGAGCTACCTAGAAGAATCATGTAGCACTATAACTGAGGGAT encoding a
ATCTTAGTGTTCTGAGGACAGGTTGGTATAACCAACGTTTTTACATTAGAGGT chimeric F
GGGTGATGTAGAAAACCTTACATGTTCTGATGGACCTAGCCTAATAAAAAC protein
AGAATTAGATCTGACCAAAAGTGCCTAAGAGAGCTCAAAACAGTCTCTGC comprising
TGACCAATTGGCAAGAGAGGAACAAATTGAGAATCCCAGACAATCTAGGTT hMPV F
TGTTCTAGGAGCAATAGCACTCGGTGTTGCAACAGCAGCTGCAGTCACAGC protein
AGGTGTTGCAATTGCCAAAACCATCCGGCTTGAGAGTGAAGTCACAGCAAT
ectodomain of
TAAGAATGCCCTCAAAACGACCAATGAAGCAGTATCTACATTGGGGAATGG hMPV
strain AGTTCGAGTGTTGGCAACTGCAGTGAGAGAGCTGAAAGACTTTGTGAGCA
CAN00-16 and
AGAATTTAACTCGTGCAATCAACAAAAACAAGTGCGACATTGATGACCTAA NDV F
protein AAATGGCCGTTAGCTTCAGTCAATTCAACAGAAGGTTTCTAAATGTTGTGCG
transmembrane
GCAATTTTCAGACAATGCTGGAATAACACCAGCAATATCTTTGGACTTAATG and

cytoplasmic ACAGATGCTGAACCTAGCCAGGGCGTTCCTAACATGCCGACATCTGCAGGA domains of the CAAATAAAATTGATGTTGGAGAACC GCGCGATGGTGCGAAGAAAGGGGTT LaSota strain CGGAATCCTGATAGGGGTCTACGGGAGCTCTGTAATTTACATGGTGCAGCT (transmembrane GCCAATCTTTGGCGTTATAGACACGCCTTGCTGGATAGTAAAAGCAGCCCC and cytoplasmic TTCTTGTTCCGAAAAAAAGGGAAACTATGCTTGCCTCTTAAGAGAAGACCA domains of NDV AGGGTGGTATTGTCAGAATGCAGGGTCAACTGTTTACTACCCAAATGAGAA F protein are AGACTGTGAAACAAGAGGAGACCATGTCTTTTGGCAGACAGCAGCAGGAA underlined) TTAATGTTGCTGAGCAATCAAAGGAGTGCAACATCAACATATCCACTACAA ATTACCCATGCAAAGTCAGCACAGGAAGACATCCTATCAGTATGGTTGCAC TGTCTCCTCTTGGGGCTCTGGTTGCTTGCTACAAAGGAGTAAGCTGTTCCAT TGGCAGCAACAGAGTAGGGATCATCAAGCAGCTGAACAAAGGTTGCTCCT ATATAACCAACCAAGATGCAGACACAGTGACAATAGACAACACTGTATATC AGCTAAGCAAAGTTGAGGGTGAACAGCATGTTATAAAAGGCAGACCAGTG TCAAGCAGCTTTGATCCAATCAAGTTTCCTGAAGATCAATTCAATGTTGCAC TTGACCAAGTTTTTTGAGAGCATTGAAAACAGCCAGGCCTTGGTAGATCAAT CAAACAGAATCCTAAGCAGTGCAAGAGAAAGGGAATACTGGCggttaacCTCAT TACCTATATCGTTTTTGACTATCATATCTCTTGTTTTTTGGTATACTTAGCCTGAT TCTAGCATGCTACCTAATGTACAAGCAAAAGGCGCAACAAAAGACCTTATT ATGGCTTGGGAATAATACCCTAGATCAGATGAGAGCCACTACAAAATGTG A Nucleic acid ATGAGCTGGAAGGTGGTGATCATCTTCAGCCTGCTGATCACCCCCCAGCAC 34 sequence GGCCTGAAGGAGAGCTACCTGGAGGAGAGCTGCAGCACCATCACCGAGGG encoding CTACCTGAGCGTGCTGAGAACCGGCTGGTACACCAACGTGTTACCCCTGGA chimeric F GGTGGGCGACGTGGAGAACCTGACCTGCGCCGACGGCCCCAGCCTGATCA protein, wherein AGACCGAGCTGGACCTGACCAAGAGCGCCCTGAGAGAGCTGAAGACCGTG the nucleic acid AGCGCCGACCAGCTGGCCAGAGAGGAGCAGATCGAGAACCCCAGACAGA sequence GCAGATTTCGTGCTGGGCGCCATCGCCCTGGGCGTGGCCACCGCCGCCGCC comprises a GTGACCGCCGGCGTGGCCATCGCCAAGACCATCAGACTGGAGAGCGAGGT codon optimized GACCGCCATCAAGAACGCCCTGAAGAAGACCAACGAGGCCGTGAGCACCC nucleic acid TGGGCAACGGCGTGAGAGTGCTGGCCACCGCCGTGAGAGAGCTGAAGGA sequence CTTCGTGAGCAAGAACCTGACCAGAGCCATCAACAAGAACAAGTGCGACAT encoding the CGACGACCTGAAGATGGCCGTGAGCTTCAGCCAGTTCAACAGAAGATTCCT hMPV F GAACGTGGTGAGACAGTTCAGCGACAACGCCGGCATCACCCCCGCCATCAG protein CCTGGACCTGATGACCGACGCCGAGCTGGCCAGAGCCGTGAGCAACATGC ectodomain of CCACCAGCGCCGGCCAGATCAAGCTGATGCTGGAGAACAGAGCCATGGTG strain AGAAGAAAGGGCTTCGGCATCCTGATCGGCGTGACGGCAGCAGCGTGAT HMPV/Homo CTACATGGTGCAGCTGCCCATCTTCGGCGTGATCGACACCCCCTGCTGGATC sapiens/PER/FPP GTGAAGGCCGCCCCCAGCTGCAGCGAGAAGAAGGGCAACTACGCCTGCCT 00726/2011/A GCTGAGAGAGGACCAGGGCTGGTACTGCCAGAACGCCGGCAGCACCGTGT and a nucleic ACTACCCCAACGAGAAGGACTGCGAGACCAGAGGCGACCACGTGTTCTGC acid sequence

GACACCAGCCCGGCCGATCAACCTGAGCAGGAGTGGCAACAT encoding
the CAACATCAGCACCAACTACCCCCTGCAAGGTGAGCACCGGCAGACACCC
transmembrane
CATCAGCATGGTGGCCCTGAGCCCCCTGGGCGCCCTGGTGGCCTGCTACAA and
cytoplasmic GGGCGTGAGCTGCAGCATCGGCAGCAACAGAGTGGGCATCATCAAGCAGC
domains of F
TGAACAAGGGCTGCAGCTACATCACCAACCAGGACGCCGACACCGTGACCA protein
from TCGACAACACCGTGTACCAGCTGAGCAAGGTGGAGGGCGAGCAGCACGTG
NDV strain ATCAAGGGCAGACCCGTGAGCAGCAGCTTCGACCCCGTGAAGTTCCCCGA
LaSota GGACCAGTTCAACGTGGCCCTGGACCAGGTGTTTCGAGAACATCGAGAACA
(transmembrane
GCCAGGCCCTGGTGGACCAGAGCAACAGAATCCTGAGCAGCGCCGAGAAG and
cytoplasmic GGCAACACCGGCg^{ttaac}CTCATTACCTATATCGTTTTGACTATCATATCTCTT
domains of NDV
GTTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCTAATGTACAAGCAA F are
underlined) AGGCGCAACAAAAGACCTTATTATGGCTTGGGAATAATACCCTAGATCAGA
TGAGAGCCACTACAAAAATGTGAccgcgg Nucleic acid
ATGAGCTGGAAGGTGATGATCATCATCAGCCTGCTGAT 35 sequence
CACCCCCCAGCACGGCCTGAAGGAGAGCTACCTGGAGG encoding
AGAGCTGCAGCACCATCACCGAGGGCTACCTGAGCGTG chimeric F
CTGAGAACCGGCTGGTACACCAACGTGTTCACCCTGGA protein, wherein
GGTGGGCGACGTGGAGAACCTGACCTGCACCGACGGCC the nucleic acid
CCAGCCTGATCAAGACCGAGCTGGACCTGACCAAGAGC sequence
GCCCTGAGAGAGCTGAAGACCGTGAGCGCCGACCAGCT comprises a
GGCCAGAGAGGAGCAGATCGAGAACCCCAGACAGAGC codon optimized
AGATTCGTGCTGGGCGCCATCGCCCTGGGCGTGGCCAC nucleic acid
CGCCGCCGCCGTGACCGCCGGCATCGCCATCGCCAAGA sequence
CCATCAGACTGGAGAGCGAGGTGAACGCCATCAAGGGC encoding the
GCCCTGAAGCAGACCAACGAGGCCGTGAGCACCCCTGGG hMPV F
CAACGGCGTGAGAGTGCTGGCCACCGCCGTGAGAGAGC protein
TGAAGGAGTTCGTGAGCAAGAACCTGACCAGCGCCATC ectodomain of
AACAGAAACAAGTGCGACATCGCCGACCTGAAGATGGC strain
CGTGAGCTTCAGCCAGTTCAACAGAAGATTCCTGAACG HMPV/USA/C2-
TGGTGAGACAGTTCAGCGACAACGCCGGCATCACCCCC 175/2005/B and
GCCATCAGCCTGGACCTGATGACCGACGCCGAGCTGGC a nucleic acid
CAGAGCCGTGAGCTACATGCCCACCAGCGCCGGCCAGA sequence
TCAAGCTGATGCTGGAGAACAGAGCCATGGTGAGAAGA encoding the
AAGGGCTTCGGCATCCTGATCGGCGTGTACGGCAGCAG transmembrane
CGTGATCTACATGGTGCAGCTGCCCATCTTCGGCGTGAT and cytoplasmic
CGACACCCCCTGCTGGATCATCAAGGCCGCCCCCAGCT domains of F
GCAGCGAGAAGAACGGCAACTACGCCTGCCTGCTGAGA protein from
GAGGACCAGGGCTGGTACTGCAAGAACGCCGGCAGCA NDV strain
CCGTGTACTACCCCAACGAGAAGGACTGCGAGACCAGA LaSota
GGCGACCACGTGTTCTGCGACACCGCCGCGGCATCAA (transmembrane
CGTGGCCGAGCAGAGCAGAGAGTGCAACATCAACATCA and cytoplasmic
GCACCACCAACTACCCCTGCAAGGTGAGCACCGGCAGA domains of NDV
CACCCCATCAGCATGGTGGCCCTGAGCCCCCTGGGCGC F are underlined)
CCTGGTGGCCTGCTACAAGGGCGTGAGCTGCAGCATCG
GCAGCAACAGAGTGGGCATCATCAAGCAGCTGCCCAAG
GGCTGCAGCTACATCACCAACCAGGACGCCGACACCGT

GACCATCGACAACACCGTGTTACCAAGCTGAGCAAGGTGG
AGGGCGAGCAGCACGTGATCAAGGGCAGACCCGTGAG
CAGCAGCTTCGACCCCATCAGATTCCCCGAGGACCAGT
TCAACGTGGCCCTGGACCAGGTGTTTCGAGAGCATCGAG
AACAGCCAGGCCCTGGTGGAGCAGAGCAACAAGATCCT
GAACAGCGCCGAGAAGGGCAACACCGGGCgttaacCTCATT
ACCTATATCGTTTTGACTATCATATCTCTTGTTTTTGGTA
TACTTAGCCTGATTCTAGCATGCTACCTAATGTACAAGC
AAAAGGCGCAACAAAAGACCTTATTATGGCTTGGGAAT
AATACCCTAGATCAGATGAGAGCCACTACAAAAATGTG Accgagg Nucleic acid
ATGAGCTGGAAGGTGGTGATCATCTTCAGCCTGCTGATCACCCCCCAGCAC 36 sequence
GGCCTGAAGGAGAGCTACCTGGAGGAGAGCTGCAGCACCATCACCGAGGG encoding
CTACCTGAGCGTGCTGAGAACCGGCTGGTACACCAACGTGTTACCCCTGGA chimeric F
GGTGGGCGACGTGGAGAACCTGACCTGCAGCGACGGCCCCAGCCTGATCA protein,
wherein AGACCGAGCTGGACCTGACCAAGAGCGCCCTGAGAGAGCTGAAGACCGTG
the nucleic acid
AGCGCCGACCAGCTGGCCAGAGAGGAGCAGATCGAGAACCCCAGACAGA sequence
GCAGATTCGTGCTGGGCGCCATCGCCCTGGGCGTGGCCACCGCCGCCGCC comprises a
GTGACCGCCGGCGTGGCCATCGCCAAGACCATCAGACTGGAGAGCGAGGT codon
optimized GACCGCCATCAAGAACGCCCTGAAGACCACCAACGAGGCCGTGAGCACCCCT
nucleic acid
GGGCAACGGCGTGAGAGTGCTGGCCACCGCCGTGAGAGAGCTGAAGGAC sequence
TTCGTGAGCAAGAACCTGACCAGAGCCATCAACAAGAACAAGTGCGACATC
encoding the
GACGACCTGAAGATGGCCGTGAGCTTCAGCCAGTTCAACAGAAGATTCCTG hMPV F
AACGTGGTGAGACAGTTTCAGCGACAACGCCGGCATCACCCCCGCCATCAGC protein
CTGGACCTGATGACCGACGCCGAGCTGGCCAGAGCCGTGAGCAACATGCC
ectodomain of
CACCAGCGCCGGCCAGATCAAGCTGATGCTGGAGAACAGAGCCATGGTGA strain
GAAGAAAGGGCTTCGGCATCCTGATCGGCGTGACGGCAGCAGCGTGATC
HMPV/ARG/10
TACATGGTGCAGCTGCCCATCTTCGGCGTGATCGACACCCCCTGCTGGATC 7/2002/A
and a GTGAAGGCCGCCCCCAGCTGCAGCGAGAAGAAGGGCAACTACGCCTGCCT
nucleic acid
GCTGAGAGAGGACCAGGGCTGGTACTGCCAGAACGCCGGCAGCACCGTGT sequence
ACTACCCCAACGAGAAGGACTGCGAGACCAGAGGGCGACCACGTGTTCTGC encoding
the GACACCGCCGCGGCCATCAACGTGGCCGAGCAGAGCAAGGAGTGCAACAT
transmembrane
CAACATCAGCACCACTACCCCTGCAAGGTGAGCACCGGCAGACACCC and
cytoplasmic CATCAGCATGGTGGCCCTGAGCCCCCTGGGCGCCCTGGTGGCCTGCTACAA
domains of F
GGGCGTGAGCTGCAGCATCGGCAGCAACAGAGTGGGCATCATCAAGCAGC protein
from TGAACAAGGGCTGCAGCTACATCACCAACCAGGACGCCGACACCGTGACCA
NDV strain
TCGACAACACCGTGTACCAGCTGAGCAAGGTGGAGGGCGAGCAGCACGTG LaSota
ATCAAGGGCAGACCCGTGAGCAGCAGCTTCGACCCCATCAAGTTCCCCGAG
(transmembrane
GACCAGTTCAACGTGGCCCTGGACCAGGTGTTTCGAGAACATCGAGAACAG and
cytoplasmic CCAGGCCCTGGTGGACCAGAGCAACAGAATCCTGAGCAGCGCCGAGAAGG
domains of NDV

GCAACCGGCGgttaacCTCATTACCTATCGTTTACTATCTCTTTG F are
underlined) TTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCTAATGTACAAGCAAAA
GGCGCAACAAAAGACCTTATTATGGCTTGGGAATAATACCCTAGATCAGAT
GAGAGCCACTACAAAAATGTGAccgcgg cDNA of
accaaacagagaatccgtgagttacgataaaaggcgaaggagcaattg 22 genomic
aagtcgcacgggtagaagggtgaatctcgagtgcgagcccgaagcac sequence of
aaactcgagaaagccttctgccaacatgtcttccgtatttgatgagta NDV LaSota
cgaacagctcctcgcggctcagactcgcaccaatggagctcatggagg strain with the
gggagaaaaaggagtagccttaaaaagtagacgtcccggtattcactct nucleic acid
taacagtgatgaccagaagatagatggagctttgtggtattctgcct encoding a
ccggattgctgttagcgaagatgccaacaaccactcaggcaaggtgc chimeric F
tctcatatctctttatgctcccactcacaggaatgaggaacatgt protein,
tgccCttgcagggaaacagaatgaagccacattggccgtgcttgagat wherein the
tgatggctttgccaacggcacgccccagttcaacaataggagtggagt chimeric F
gtctgaagagagagcacagagatttgcgatgatagcaggatctctccc protein
tcgggcatgcagcaacggaacccccgttcgtcacagccggggcCgaaga comprises
tgatgcaccagaagacatcaccgataccctggagaggatcctctctat ectodomain of
ccaggctcaagtatgggtcacagtagcaaaagccatgactgcgtatga hMPV F
gactgcagatgagtcggaaacaaggcgaatcaataagtatatgcagca protein of
aggcaggggtccaaaagaaatacatcctctacccccgtatgcaggagcac hMPV CAN00-
aatccaactcacgatcagacagtctcttgagtcgcgtatcttttgggt 16 strain and the
tagcgagctcaagagaggccgcaacacggcaggtggtagccttactta transmembrane
ttataacctggtaggggacgtagactcatacatcaggaataaccgggct and cytoplasmic
tactgcattctcttgacactcaagtacggaatcaaccaagacatc domains of the
agcccttgacttagtagcctctcaggcgacatccagaagatgaagca NDV F protein
gctcatgcgtttgtatcggtgaaaggagataatgcgccgtacatgac of LaSota strain
attacttgggtgatgtgaccagatgagctttgcgcctgccgagtatgc inserted (Sac II
acaactttactcctttgccatgggtatggcatcagtcctagataaagg (CCGCGG)
tactgggaaataccaatttgccagggactttatgagcacatcattctg restriction sites
gagacttggagtagagtacgctcaggctcagggaagtagcattaacga used to insert the
ggatatgggtgccgagctaaagctaaccacagcagcaaGgaGgggcct chimeric RSV-F
ggcagctgctgccaacgggtctccgaGgaGaccagcagcataGacat sequence are
gcctactcaacaagtcggagtcctcactgggcttagcgaggggggggtc double
ccaagctctacaaggcggatcgaatagatcgcaagggcaaccagaagc underlined and
cggggatggggagacccaattctggatctgatgagagcggtagcaaa the open reading
tagcatgagggaggcgccaaactctgcacagggcactccccaatcggg frame for
gcctccccaaactcctgggcatcccaagataacgacaccgactgggg hMPVF protein
gtattgatggacaaaaccagcctgcttcacaaaaacatcccaatgc is underlined)
cctcaccgtagtcgacccctcgatttgcggctctatatgaccacacc ctcaaacaacatccccctcttctcctccccctgctgtacaactA
cgTacgccctagataccacaggcacaatgcgggtcactaacaatcaaa
acagagccgaggggaattagaaaaaagtacgggtagaagagggatattc
agagatcagggaagctccccgagtctctgctctctctctacctgat
agaccaggacaaacatggccaccttacagatgcagagatcgacgagc
tatttgagacaagtggaaactgtcattgacaacataattacagcccagg
gtaaacagcagagactgttggaaggagtgcaatcccacaaggcaaga
ccaaggtgctgagcgcagcatgggagaagcatgggagcatccagccac
cggccagtcaagacaacccccgatcgacaggacagatctgacaaacaac
catccacacccgagcaaacgacccccgatgacagcccgcggccacat
ccgccgaccagccccccacccaggccacagacgaagccgtcgacacac
agCtcaggaccggagcaagcaactctctgctgttgatgcttgacaagc

tcagcaataatcgccaatgctaataaggcccatggctcgagcccc
aagaggggaatcaccaacgtccgactcaacagcaggggagtcaccca
gtcgcggaaacagtcaggaaagaccgcagaaccaagtcaaggccgcc
ctggaaaccagggcacagacgtgaacacagcatatcatggacaatggg
aggagtcacaactatcagctgggtgaacccctcatgctctccgatcaa
ggcagagccaagacaatacccttgatctgcggatcatgtccagccac
ctgtagactttgtgaagcgatgatgtctatgatggaggcgatatcac agagagtaagtaagggtgactatcagctagatcttgtcttgaaacaga
catcctccatccctatgatgcgggtccgaaatccaacagctgaaaacat
ctgttgtagtcatggaagccaacttggaatgatgaagattctggatc ccggttggtgccaacatttcatctctgagtgatctacgggcagttgcc
gatctcaccgggttttagtttcaggccctggagacccctctccctatg
tgacacaaggaggcgaaatggcacttaataaaactttcgcaaccagtgc
cacatccatctgaattgattaaacccgccactgcatgcgggcctgata
taggagtggaaggaactgtccgtgcattgatcatgtcacgcccac
tgcacccgagttcttcagccaagctcctaagcaagttagatgcagccg
ggtcgatcgaggaaatcaggaaaatcaagcgcttgctctaaatggct
aattactactgccacacgtagcgggtccctgtccactcggcacacac
ggaatctgcaccgagttccccccgcGgaccaaggtccaactctcca
agcggcaatcctctctcgttctcagccccactgaatgAtcgcgtaa
ccgtaattaatctagctacatttaagattaagaaaaatacgggtaga
attggagtgccccaCtAgtTTTgcccaCCATGTCTTGGAAAGTGGTGAT
CATTTTTTCATTGCTAATAACACCTCAACACGGTCTTAAAGAGAGCTA
CCTAGAAGAATCATGTAGCACTATAACTGAGGGATATCTTAGTGTTCT
GAGGACAGGTTGGTATACCAACGTTTTTACATTAGAGGTGGGTGATGT
AGAAAACCTTACATGTTCTGATGGACCTAGCCTAATAAAAACAGAATT
AGATCTGACCAAAAAGTGCCTAAGAGAGCTCAAAAACAGTCTCTGCTGA
CCAATTGGCAAGAGAGGAACAAATTGAGAATCCCAGACAATCTAGGTT
TGTTCTAGGAGCAATAGCACTCGGTGTTGCAACAGCAGCTGCAGTCAC
AGCAGGTGTTGCAATTGCCAAAACCATCCGGCTTGAGAGTGAAGTCAC
AGCAATTAAGAATGCCCTCAAAAACGACCAATGAAGCAGTATCTACATT
GGGGAATGGAGTTCGAGTGTTGGCAACTGCAGTGAGAGAGCTGAAAGA
CTTTGTGAGCAAGAATTTAACTCGTGCAATCAACAAAAACAAGTGCGA
CATTGATGACCTAAAAATGGCCGTTAGCTTCAGTCAATTCAACAGAAG
GTTTCTAAATGTTGTGCGGCAATTTTCAGACAATGCTGGAATAACACC
AGCAATATCTTTGGACTTAATGACAGATGCTGAACTAGCCAGGGCCGT
TTCTAACATGCCGACATCTGCAGGACAAATAAAATTGATGTTGGAGAA
CCGCGCGATGGTGCGAAGAAAGGGGTTTCGGAATCCTGATAGGGGTCTA
CGGGAGCTCTGTAATTTACATGGTGCAGCTGCCAATCTTTGGCGTTAT
AGACACGCCTTGCTGGATAGTAAAAGCAGCCCCTTCTTGTTCCGAAAA
AAAGGGAACTATGCTTGCCTCTTAAGAGAAGACCAAGGGTGGTATTG
TCAGAATGCAGGGTCAACTGTTTACTACCCAAATGAGAAAGACTGTGA
AACAAGAGGAGACCATGTCTTTTGCGACACAGCAGCAGGAATTAATGT
TGCTGAGCAATCAAAGGAGTGCAACATCAACATATCCACTACAAATTA
CCCATGCAAAGTCAGCACAGGAAGACATCCTATCAGTATGGTTGCACT
GTCTCCTCTTGGGGCTCTGGTTGCTTGCTACAAAGGAGTAAGCTGTTT
CATTGGCAGCAACAGAGTAGGGATCATCAAGCAGCTGAACAAAGGTTG
CTCCTATATAACCAACCAAGATGCAGACACAGTGACAATAGACAACAC
TGTATATCAGCTAAGCAAAGTTGAGGGTGAACAGCATGTTATAAAAGG
CAGACCAGTGTCAAGCAGCTTTGATCCAATCAAGTTTCCTGAAGATCA
ATTCATGTTGCACTTGACCAAGTTTTTGGAGAGCATTGAAAACAGCCA
GGCCTTGGTAGATCAATCAAACAGAATCCTAAGCAGTGCAGAGAAAGG

GAATACTGGCGTTAATTACTTATATCGTTTTGACTATCATATC
TCTTGTTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCTAATGTA
CAAGCAAAAGGCGCAACAAAAGACCTTATTATGGCTTGGAATAATAC
CCTAGATCAGATGAGAGCCACTACAAAAATGTGAccgcggaaccaagg

tcctaactctcaagcggcaatcctctctcgttcctcagccccactga atgAtcgcgtaaccgtaattaatctagctacatttaagattaagaaaa
aatacgggtagaattggagtgcccaattgtgccaagatggactcatc taggacaattgggctgtactttgattctgcccattctctagcaacct
gtagcatttccgatcgtcctacaagAcacaggagatgggaagaagca
aatcgccccgcaatataggatccagcgccttgacttgggactgatag taaggaggactcagtattcatcaccacctatggattcatctttcaagt
tgggaatgaagaagccacCgtcggcatgatcgatgataaacccaagcg
cgagttactttccgctgcgatgctctgcctaggaagcgtcccaatac
cggagaccttattgagctggcaagggcctgtctcactatgatagtcac
atgcaagaagagtgaactaatactgagagaatggttttctcagtagt
gcaggcaccacaagtgtgctgcaaagctgtaggggtgtggcaaacaaata
ctcatcagtgaatgcagtcaagcacgtgaaagcgccagagaagattcc
cgggagtggaaccctagaatacaaggtgaactttgtctccttgactgt
ggtagcgaagaGggatgtctacaagatccagctgcagtattgaaggt ttctggctcgagtctgtacaatcttgcgctcaatgtcactattaatgt
ggaggttagaccgaggagtcctttgggttaaactctCtgtctaagtctga cagcggatactatgctaacctcttctgcatattggacttatgaccac
Tgtagataggaaaggggaagaaagtacatttgacaagctggaaaagaa
aataaggagccttgatctatctgtcgggctcagtgatgtgctcgggcc
ttccgtgttggtaaaagcaagaggtgcacggactaagcttttggcacc ttcttctctagcagtgggacagcctgctatcccatagcaaatgcttc
tcctcaggtggccaagatactctggagtcacacccgcgtgcctgcggag
cgtaaatacattatccaagcaggtacccaacgcgtgtcgcagtgac
cgccgaccacgaggttacctctactaagctggagaaggggcacaccct
tgccaaatacaatcctttaagaaataagctgcgtctctgagattgcg ctccgcccactcaccagatcatcatgacacaaaaaactaatctgtct
tgattatttacagttagtttacctgtctatcaagttagaaaaaacacg ggtagaagattctggatcccgggtggcgccctccaggtgcaagatggg
ctccagaccttctaccaagaaccagcacctatgatgctgactatccg
ggttgcgctggtagtgagttgcatctgtccggcaaacctcattgatgg
caggcctcttgagctgcaggaattgtggttacaggagacaaagccgt
caacatatacacctcatcccagacaggatcaatcatagttaagctcct
cccgaatctgccaaggataaggaggcatgtgcgaaagcccccttgga
tgcatacaacaggacattgaccactttgctcacccttggtgactc
tatccgtaggatacaagagtctgtgactacatctggagggggggagaca
ggggcgcccttataggcgccattattggcggtgtggctcttgggggtgc
aactgccgcacaaataacagcggccgcagctctgatacaagccaaaca
aaatgctgccaacatcctccgacttaaagagagcattgccgcaaccaa
tgaggctgtgcatgaggtcactgacggattatcgcaactagcagtggc
agttgggaagatgcagcagtttgtaatgaccaatttaataaaacagc
tcaggaattagactgcatcaaaattgcacagcaagttggtgtagagct
caacctgtacctaaccgaattgactacagtattcggaaccacaaatcac ttacctgctttaaacaagctgactattcaggcactttacaatctagc
tggtggaaatatggattactattgactaagttaggtgtagggaaaca tcaactcagctcattaatcggtagcggcttaatcaccggtaaccctat
tctatacgactcacagactcaactcttgggtatacaggttaactctacc ttcagtcgggaacctaaataatatgcgtgccacctacttggaacctt
atccgtaagcacaaccaggggatttgctcggcacttgctcccAaaagt
ggtagacaggtcgggtctgtgataagaacttgacacctcactg tatagaaactgacttagatttatattgtacaagaatagtaacgttccc
tatgtcccctgggtatttattctgcttgagcggcaatacgtcggcctg tatgtactcaaagaccgaaggcgcacttactacaccatacatgactat
caaaggttcagtcatcgccaactgcaagatgacaacatgtagatgtg
aaaccccccggtatcatatcgcaaaactatggagaagccgtgtctct aatagataaacaatcatgcaatgtttatccttaggcgggataacttt
aaggctcagtggggaattcgatgtaacttatcagaagaatatctcaat acaagattctcaagtaataataacaggcaatcttgatatctcaactga
gcttggaatgtcaacaactcgatcagtaattgcttgaataagttaga
ggaaagcaacagaaaactagacaaagtcaatgtcaaaactgactagcac atctgctctcattacctatatcgttttgactatcatatctctgtttt
tggtatacttagcctgattctagcatgtacctaattgtacaagcaaaa ggcgcaacaaaagaccttattatggcttggaataatactctagatca

gatgagagccactacaaaaatgtgaacacagatgaggaacgaagggtt
ccctaatagtaatgtgtgaaagttctggtagctgtcagttcagag agttaagaaaaaactaccggttagatgaccaaaggacgatatacgg
gtagaacggtaagagaggccgccccctcaattgcgagccaggcttcaca
acctccgttctaccgcttcaccgacaacagtcctcaatcatggaccgc
gccgtagccaagttgcgtagagaatgatgaaagagaggcaaaaaat acatggcgcttgatattccggattgcaatcttattcttaacagtagtg
accttggctatatctgtagcctcccttttatatagcatgggggctagc acacctagcgatcttgtaggcataccgactaggattccagggcagaa
gaaaagattacatctacacttggttccaatcaagatgtagtagatagg atatataagcaagtgggcccttgagtctccgttggcattgttaaatact
gagaccacaattatgaacgcaataacatctctctcttatcagattaat
ggagctgcaaacaacagtggggtgggggggcacctatccatgaccagat
tatataggggggtagaggcaaagaactcattgtagatgatgctagtgat gtcacatcattctatccctctgcatttcaagaacatctgaattttatc
ccggcgctactacaggatcaggttgcaactcgaataccctcatttgac atgagtgtaccctactgctacaccataatgtaatatgtctgga
tgacagagatcactcacattcatatcagtttagcacttgggtgtgctc cggacatctgcaacagggagggtattcttttctactctgcgttccatc
aacctggacgacacccaaaaatcggaagcttgcaagtgtgagtgcact
cccctgggttgtagatgtgtgctcgaaagtcacggagacagaggaa
gaagattataactcagctgtccctacgcggatggtacatgggagggtta
gggttcgacggccagtagccacgaaaaggacctagatgtcacaacatta
ttcggggactgggtggccaactaccaggagtaggggggggatcttt
attgacagccgcgtatggttctcagcttacggagggttaaaaccaat
tcacccagtgcactgtacaggaagggaatatgtgatatacaagcga
tacaatgacacatgccagatgagcaagactaccagattcgaatggcc
aagtcttcgtataagcctggacggttgggggaaacgcatacagcag
gctatcttatctatcaaggtgtcaacatccttaggcgaagaccggtta
ctgactgtaccgccaacacagtcacactcatgggggccaaggcaga
attctcacagtagggacatctcatttctgtatcaacgagggtcatca tacttctctccgcgttattatctctatgacagtcagcaaaaaaca
gccactctcatagtccttatacattcaatgccttcactcgccaggt agtatcccttgccaggcttcagcaagatgccccaaactcgtgtgttact
ggagctctatacagatccatatcccctaattctctatagaaaccacacc ttgcgaggggtattcgggacaatgcttgatggtgtacaagcaagactt
aacctgcgtctgcagtattcgatagcacatcccgcagtcgcattact
cgagtgcgttcaagcagtagcacaacacatcaacttgt ttaaagtggtaagactaataagacctattgtctcagcattgtctgaa
atatctaatactctcttcggagaattcagaatcgctccggttactagtt gagatcctcaaagatgacgggggttagagaagccaggctctggctagttg
agtcaattataaaggaggttgaaagatggcattgtatcacctatcttc
tgcgacatcaagaatcaaaccgaatgccggcgctgctcgaattccat
gttgccagttgaccacaatcagccagtgctcatgcgatcagattaagc
cttgtaAtaGtctcttgattaagaaaaaatgtaagtggcaatgagat
acaaggcaaaacagctcatggtTaaCaatacgggtaggacatggcgag
ctccggtcctgaaagggcagagcatcagattatcctaccagagTcaca
cctgtcttcaccattgggtcaagcacaaactactctattactggaaatt aactgggctaccgcttctgatgaatgtgacttcgaccacctcattct
cagccgacaatggaaaaaatacttgaatcggcctctctgatactga
gagaatgataaaactcggaagggcagtagaccaaactcttaaccacaa
ttccagaataaccggagtgctccaccccagggtgttagaaGaactggc
taatattgaggtcccagattcaaccaacaaatttcggaagattgagaa
gaagatccaaattcacaacacgagatatggagaactgttcacaaggct
gtgtacgcatatagagaagaaactgctggggctcatcttggtctaaca
tgtccccggctcagaggagttcagcagcattcgtacggatccggcatt
ctggtttactcaaaatgggtccacagccaagtttgcatgggtccatat
aaaacagatccagaggcatctgatggtggcagctaGgacaaggctctgc
ggccaacaaattggtgatgctaaccataaggtaggccaagtctttgt
cactctgaacttgtctgtgacgcatacgaatgagaacaagttcac atgtcttaccaggaacttgattgatgtatgcagatatgatggaggg
cagagatatggtcaacataatatcaaccacggcggtgcatctcagaag
cttatcagagaaaattgatgacattttgcgggttaatagacgtctggc aaaagacttgggtaataagctacgatgttgatcactaatggaggg
attgcatacggagctgtccagctactcgagccgtcaggtacatttgc aggagatttcttcgattcaacctgcaggagcttaagacattcta

tggcctcctccccaatgatatagcagaatccgtgactcatgcaatcgc tactgtattctctggtttagaacagaatcaagcagctgagatggttg
tctgttgcgctctgtgggggtcacccactgcttgagtcccgtattgcagc
aaaggcagtcaggagccaaatgtgcgaccgaaaatggtagactttga
tatgatccttcagggtactgtctttcttcaaggggaacaatcatcaacgg
gtacagaaagaagaatgcaggtgtgtggccgagtc aaagtggatac
aatatatgggaaggtcattggggcaactacatgcagattcagcagagat ttacacgatatcatgttgagagagtataagagtttatctgcacttga
atttgagccatgtatagaatatgacctgtcaccaacctgagcatgtt
cctaaaagacaaggcaatcgcacaccccaacgataattggcttgctc
gtttaggcggaaccttctctccgaagaccagaagaacatgtaaaaga
agcaacttcgactaatcgctcttgatagagtttttagagtcaaatga tttgatccatataaagagatggaatatctgacgaccttgagtacct
tagagatgacaatgtggcagtatcatactcgctcaaggagaaggaagt
gaaagttaatggacggatcttcgctaagctgacaaagaagttaaggaa
ctgtcaggtgatggcggaagggtacctagccgatcagattgcacctt
ctttcagggaaatggagtcattcaggatagcatatccttgaccaagag tatgctagcgatgagtc aactgtcttttaacagcaataagaaacgtat
cactgactgtaaagaaagagtatcttcaaaccgcaatcatgatccgaa
aagcaagaaccgtcggagagttgcaaccttcataacaactgacctgca
aaagtactgtcttaattggagatatcagacaatcaaattgttcgctca tgccatcaatcagttgatgggcctacctcacttcttgaatggattca
cctaagactgatggacactacgatgttcgtaggagaccctttcaatcc
tccaagtgacctactgactgtgacctctcaagagtccttaatatga
catatatattgtcagtgccagaggggggtatcgaaggattatgccagaa
gctatggacaatgatctcaattgctgcaatccaacttgctgcagctag atcgcatgtcgtgttgccctgtatggtacaggggtgataatcaagtaat
agcagtaacgagagaggtaagatcagacgactctccggagatgggtgtt
gacacagttgcatcaagccagtgataatttcttcaaggaattaattca tgtcaatcatttgattggccataatttgaaggatcgtgaaaccatcag
gtcagacacattcttcatatacagcaaacgaatcttcaaagatggagc aatcctcagtc aagtcctcaaaaattcatctaaattagtgtctagtgtc
aggtgatctcagtgaaaacaccgtaattgtcctgtgccaacattgcctc
tactgtagcacggctatgcgagaacgggcttcccaaagacttctgtta ctatttaaactatataatgagttgtgtgcagacatactttgactctga
gttctccatcaccaacaattcgcaccccgatcttaatcagtcgtggat tgaggacatctctttgtgcactcatatgttctgactcctgcccatt
aggggggactgagtaaccttcaatactcaaggctctacactagaaatat
cgggtgacccggggactactgctttgcagagatcaagcgactagaagc
agtgggattactgagtcctaacattatgactaatatcttaactaggcc gcctgggaatggagattgggccagtcgtgcaacgaccatactctt
caattttgagactgttgcaagcccaaatattgttcttaagaaacatac gcaaagagtcctatttgaaactgttcaaatcccttattgtctggagt
gcacacagaggataatgaggcagaagagaaggcattggctgaattctt
gcttaatcaagaggtgattcatccccgcgttgcgcatgccatcatgga
ggcaagctctgtaggtaggagaaagcaaattcaagggcttgttgacac
aacaacaccgtaattaagattgcgcttactaggaggccattaggcat
caagaggctgatgcggatagtc aattattctagcatgcatgcaatgct gtttagagacgatgtttttcctccagtagatccaaccacccttagt
ctcttctaatatgtgttctctgacactggcagactatgcacggaatag
aagctggtcacctttgacgggaggcaggaaaatactgggtgtatctaa
tcctgatac gatagaactcgtagaggggtgagattcttagtgtaagcgg
aggggtgtacaagatgtgacagcggagatgaacaatttacttggtcca
tcttccaagcaatatagaattgaccgatgacaccagcaagaatcctcc
gatgaggggtaccatatctcgggtcaaagacacaggagaggagagctgc
ctcacttgcaaaaatagctcatatgtcgccacatgtaaaggctgcct
aagggcatcatccgtgttgatctgggcttatggggataatgaagtaaa ttggactgctgctcttacgattgcaaaaatctcgggtgtaatgtaaactt
agagtatcttcgggttactgtcccccttaccacggctgggaattctca acatagactagatgatggtataactcagatgacattcacccctgcatc
tctctacaggGtgtcaccttacattcacatatccaatgatttctcaaag
gctgttactgaagaaggagtc aaagaggggaatgtgggttaccaca gatcatgctcttgggtttatctctaatacgaatc gatctttccaatgac
aacaaccaggacatatgatgagatcacactgcacctacatagtaaatt tagttgctgtatcagagaagcacctgttgcggttcctttcgagctact
tgggggtgggtaccggaactgaggacagtgacctcaaataagtttatgta
tgatcctagccctgtatcggaggggagactttgcgagacttgacttagc tatcttcaagagttatgagcttaatctggagtcatatccacgataga

gctaataacattctttcaatatccagcgggaagttgattggccagtc tgtggtttcttatgatgaagataacctccataaagaatgacgccataat
agtgtagacaatacccgaaattggatcagtgaaagctcagaattcaga tgtgggccgctatttgaatatgcagcacttgaagtgtcctcgactg
ttcttaccactctattacctgagagtaagaggcctGgacaatatgt cttatatatgggtgattatacaagaatatgccaggaattctactttc
caacattgcagctacaatatctcatcccgtcattcattcaaggttaca
tgcagtgggcctgggtcaacatgacggatcacaccaacttgcagatac
ggattttatcgaaatgtctgcaaaactattagtagtcttgcacccgacg tgtgatctccggcttatattcaggaaataagtagatctgtgttccc
atctgtcttagatgataacctgaatgagaagatgcttcagctgatatc ccggttatgctgtctgtacacgggtactcttctgctacaacaagagaaat
cccgaaaataagaggcttaactgcagaagagaaatgttcaatactcac
tgagtatttactgtcggatgctgtgaaaccattacttagccccgatca agtgagctctatcatgtctcctaacataattacattcccagctaactt
gtactacatgtctcggagagcctcaatttgatcagggaaaggaggga
cagggtactatcctggcgtgtgttcccccaagagccattattaga gttcccttctgtgcaagatattgggtgctcgagtgaagatccattcac
ccgacaacctgcggcatttttgcaagagttagatttagtgctccagc aaggtatgacgcattcacacttagtcagattcatcctgaactcacatc
tccaaatccggaggaagactacttagtacgatacttgttcagagggat agggactgcattcttcttgggtataaggcatctcatctcttctgt
acccgaggtaagatgtgcaagacacgggaactccttatacttagctga
agggagcggagccatcatgagcttctcgaactgcatgtaccacatga
aactatctattacaatacgtctctttcaaagatgaacccccgca acgacatttccggccgaccccaactcagttttgaattcgggtgttta
taggaatctacaggcggaggttaacatgcaaagatggatttgcgaaga
gttccgtccattatggagagaaaatacagaggaaagCgacctgacctc
agataaagTagtgggggtatattacatctgcagtgcctacagatctgt atcattgctgcattgtgacattgaaattcctccagggtccaatcaaag
cttactagatcaactagctatcaatttatctctgattgccatgcattc tgaaggaggaggcggggtagtaatcatcaaagtgttgatgcaatggg
atactactttcatctactcatgaactgtttgctccgtgttcacaaa aggatatttctctctaatgggtatgcatgtcgaggagatatggagtg
ttacctggtatttgcagtggttacctgggcgggcctacattgtaca
tgagggtggtgaggatggcGaaaactctggtgcagcggcacggtacgt
Tttgtctaaatcagatgagatcacactgaccaggtattcacctcac gcggcagcgtgtgacagacatcctatccagtcctttaccaagattaat
aaagtacttgaggaagaatattgacactgcgctgattgaagccggggg
acagcccgtccgtccattctgtgcggagagctggtgagcacgctagc
gaacataactcagataaccagatCatcgctagtcacattgacacagt
tatccgggtctgtgatatatatggaagctgagggtgatctcgctgacac agtatttctatttacccttacaatctctctactgacgggaaaaagag
gacatcacttaAacagtgcacgagacagatcctagaggttacaatact
aggctcttagagtcgaaaatctcaataaaataggcgatataatcagcct
agtgtctaaaggcatgatctccatggaggaccttatcccactaaggac atacttgaagcatagtagctgcctaaatatttgaaggctgtcctagg
tattacaaactcaaagaaatgtttacagacactctgtatCtgtactt gactcgtgtcaacaaaaattctacatgaaaactataggcaatgcagt
caaaggatattacagtaactgtgactcttaacgaaaatcacatattaa taggctcctttttggccaattgtattctgttgatttaacatatta
tgttagaaaaaagtgaaccctgactccttaggactcgaattcgaact caaataaatgtcttaaaaaaagggtgcgcacaattattcttgagtga
gtctcgtcattcacaaatctttgtttggt Wild-type

ATGTCTTGGAAAGTGGTGATCATTTTTTTCATTGCTAATA 52 Nucleic acid
ACACCTCAACACGGTCTTAAAGAGAGCTACTTAGAAGA sequence
ATCATGTAGCACTATAACTGAGGGATATCTCAGTGTCT encoding hMPV
GAGGACAGGTTGGTATACCAACGTTTTTACATTAGAAG F protein of
TGGGTGATGTAGAAAACCTCACATGTGCTGATGGACCT strain
AGCCTAATAAAAACAGAATTAGATCTGACCAAAAGTGC HMPV/Homo
ACTAAGAGAGCTCAAAACAGTTTCTGCTGACCAATTGG sapiens/PER/FPP
CAAGAGAGGAACAGATTGAGAATCCCAGACAATCTAG 00726/2011/A
ATTTGTTCTAGGAGCAATAGCACTCGGTGTTGCGACAG
CAGCTGCAGTTACAGCAGGTGTTGCAATTGCCAAAAC
ATCCGACTTGAGAGTGAAGTTACAGCAATTAAGAATGC
CCTTAAAAAGACTAATGAAGCAGTGTCTACATTGGGGA
ATGGAGTTCGAGTGTTAGCAACTGCAGTGAGGGAACCTG
AAAGATTTTGTGAGCAAGAATTTAACTCGTGCAATCAA
CAAAAACAAGTGCGACATTGATGACCTAAAAATGGCTG

TTAGCTTCACTCAATCAGTAAGGTTTCTAAATGTTG
TGCGGCAATTTTCAGACAATGCTGGAATAACACCAGCA
ATATCTTTAGACTTAATGACAGATGCTGAACTAGCCAG
GGCCGTTTCCAACATGCCGACATCTGCAGGACAAATAA
AATTGATGTTGGAAAACCGTGCAATGGTGCGAAGGAAG
GGGTTTCGGAATCCTGATCGGGGTCTACGGGAGCTCCGT
AATTTACATGGTGCACTGCCAATCTTTGGCGTTATAGA
CACGCCTTGCTGGATAGTAAAAGCAGCCCCTTCTTGTTT
CGAAAAAAAGGGAAACTATGCTTGCCTCTTAAGAGAAG
ACCAAGGGTGGTATTGTCAGAATGCAGGGTCAACTGTT
TACTACCCAAATGAGAAGGACTGTGAAACAAGAGGAG
ACCATGTCTTTTGCGACACAGCAGCAGGAATTAATGTT
GCTGAGCAATCAAAGGAGTGCAACATCAATATATCCAC
CACAAATTACCCGTGCAAAGTCAGCACAGGAAGGCATC
CCATCAGTATGGTTGCACTGTCCCCTCTTGGGGCTCTGG
TTGCCTGTTACAAGGGAGTAAGCTGTTCCATTGGCAGC
AATAGAGTAGGGATCATCAAGCAGCTGAACAAAGGTTG
CTCTTATATAACCAACCAAGATGCAGACACAGTGACAA
TAGACAACACTGTATATCAGCTGAGCAAAGTTGAGGGT
GAACAGCATGTTATAAAAGGCAGACCAGTGTCAGCAG
CTTTGATCCAGTCAAGTTTCCTGAAGATCAATTCAATGT
TGCACTTGATCAAGTTTTTGAAAACATTGAAAACAGCC
AGGCCTTGGTAGATCAATCAAACAGAATCCTAAGCAGT
GCAGAGAAAGGGAACACTGGCTTCATCATTGTGATAAT
TCTAATTGCTGTCTTGGCTCTAGCATGATCCTGGTGAG
CGTCTTTATTATAATCAAGAAAACAAAGAAACCAACAG
GAGCACCTCCAGAGCTAAGCGGTGTCACAAACAATGGC TTCATACCGCACAGTTAG
Codon ATGAGCTGGAAGGTGGTGATCATCTTCAGCCTGCTGATCACCCC 53 optimized
CCAGCACGGCCTGAAGGAGAGCTACCTGGAGGAGAGCTGCAGCA nucleic acid
CCATCACCGAGGGCTACCTGAGCGTGCTGAGAACCGGCTGGTAC encoding hMPV
ACCAACGTGTTCAACCCTGGAGGTGGGCGACGTGGAGAACCTGAC F protein of
CTGCGCCGACGGCCCCAGCCTGATCAAGACCGAGCTGGACCTGA HMPV/Homo
CCAAGAGCGCCCTGAGAGAGCTGAAGACCGTGAGCGCCGACCAG sapiens/PER/
CTGGCCAGAGAGGAGCAGATCGAGAACCCCAGACAGAGCAGATT FPP00726/2011/A
CGTGCTGGGCGCCATCGCCCTGGGCGTGGCCACCGCCGCCGCCG
TGACCGCCGGCGTGGCCATCGCCAAGACCATCAGACTGGAGAGC
GAGGTGACCGCCATCAAGAACGCCCTGAAGAAGACCAACGAGGC
CGTGAGCACCTGGGCAACGGCGTGAGAGTGCTGGCCACCGCCG
TGAGAGAGCTGAAGGACTTCGTGAGCAAGAACCTGACCAGAGCC
ATCAACAAGAACAAGTGCGACATCGACGACCTGAAGATGGCCGT
GAGCTTCAGCCAGTTCAACAGAAGATTCCTGAACGTGGTGAGAC
AGTTCAGCGACAACGCCGGCATCACCCCCGCCATCAGCCTGGAC
CTGATGACCGACGCCGAGCTGGCCAGAGCCGTGAGCAACATGCC
CACCAGCGCCGGCCAGATCAAGCTGATGCTGGAGAACAGAGCCA
TGGTGAGAAGAAAGGGCTTCGGCATCCTGATCGGCGTGTACGGC
AGCAGCGTGATCTACATGGTGCACTGCCCATCTTCGGCGTGAT
CGACACCCCCTGCTGGATCGTGAAGGCCGCCCCAGCTGCAGCG
AGAAGAAGGGCAACTACGCCTGCCTGCTGAGAGAGGACCAGGGC
TGGTACTGCCAGAACGCCGGCAGCACCGTGTACTACCCCAACGA
GAAGGACTGCGAGACCAGAGGCGACCACGTGTTCTGCGACACCG

CCGCCGCGCATCGGACGATGACGAGCAAGGAGTGCAACATC
AACATCAGCACCACCAACTACCCCTGCAAGGTGAGCACCGGCAG
ACACCCCATCAGCATGGTGGCCCTGAGCCCCCTGGGCGCCCTGG
TGGCCTGCTACAAGGGCGTGAGCTGCAGCATCGGCAGCAACAGA
GTGGGCATCATCAAGCAGCTGAACAAGGGCTGCAGCTACATCAC
CAACCAGGACGCCGACACCGTGACCATCGACAACACCGTGTACC
AGCTGAGCAAGGTGGAGGGCGAGCAGCACGTGATCAAGGGCAGA
CCCGTGAGCAGCAGCTTCGACCCCGTGAAGTTCCCCGAGGACCA
GTTCAACGTGGCCCTGGACCAGGTGTTTCGAGAACATCGAGAACA
GCCAGGCCCTGGTGGACCAGAGCAACAGAATCCTGAGCAGCGCC
GAGAAGGGCAACACCGGCTTCATCATCGTGATCATCCTGATCGC
CGTGCTGGGCAGCAGCATGATCCTGGTGAGCGTGTTTCATCATCA
TCAAGAAGACCAAGAAGCCCACCGGCGCCCCCCCCCGAGCTGAGC
GGCGTGACCAACAACGGCTTCATCCCCCACAGCTGA Wild-type
ATGTCTTGGAAAGTGATGATCATCATTTTCGTTACTCATA 54 Nucleic acid
ACACCCCAGCACGGGCTAAAGGAGAGTTATTTGGAAGA sequence
ATCATGCAGTACTATAACTGAGGGATACCTCAGTGTTTT encoding hMPV
AAGAACAGGCTGGTACACCAATGTCTTCACATTAGAAG F protein of
TTGGTGATGTTGAAAATCTTACATGTACTGATGGACCTA strain
GCTTAATCAAAACAGAACTTGACCTAACAAAAAGTGCT HMPV/USA/C2-
TTAAGGGAACCTCAAAACAGTCTCTGCTGATCAGTTAGC 175/2005/B
GAGAGAGGAGCAAATTGAAAATCCCAGACAATCAAGA
TTTGTCCTAGGTGCAATAGCTCTCGGAGTTGCTACAGCA
GCAGCAGTCACAGCAGGCATTGCAATAGCCAAAACCAT
AAGGCTTGAGAGTGAGGTGAATGCAATTAAAGGTGCTC
TCAAACAACTAATGAAGCAGTATCCACATTAGGAAAT
GGTGTGCGGGTCCTAGCCACTGCAGTGAGAGAGCTAAA
AGAATTTGTGAGCAAAAATCTGACTAGTGCAATCAACA
GGAACAAATGTGACATTGCTGATCTGAAGATGGCTGTC
AGCTTCAGTCAATTCAACAGAAGATTTCTAAATGTTGTG
CGGCAGTTTTTCAGACAATGCAGGGATAACACCAGCAAT
ATCATTGGACCTAATGACTGATGCTGAATTGGCCAGAG
CTGTATCATACATGCCAACATCTGCAGGACAGATAAAA
CTGATGTTGGAGAACCGCGCAATGGTAAGGAGAAAAG
GATTTGGAATCCTAATAGGGGTCTACGGAAGCTCTGTG
ATTTACATGGTTCAATTGCCGATCTTTGGTGTCATAGAT
ACACCTTGTTGGATAATCAAGGCAGCTCCCTCTTGCTCA
GAAAAAAACGGGAATTATGCTTGCCTCCTAAGAGAGGA
TCAAGGGTGATTGTAAAAATGCAGGATCCACTGTTT
ACTACCCAAACGAAAAAGACTGTGAAACAAGAGGTGA
TCATGTTTTTTGTGACACAGCAGGGATCAATGTTGC
TGAGCAATCAAGAGAATGCAACATCAACATATCTACTA
CCAACCTACCCATGCAAAGTCAGCACAGGAAGACACCCT
ATAAGCATGGTTGCACTATCACCTCTCGGTGCTTTGGTG
GCTTGCTATAAAGGGGTAAAGCTGCTCGATTGGCAGCAA
TCGGGTTGGAATCATCAAACAATTACCTAAAGGCTGCT
CATAATACTAACCAGGATGCAGACACTGTAACAATT
GACAATACCGTGTATCAACTAAGCAAAGTTGAAGGTGA
ACAGCATGTAATAAAAGGGAGACCAGTTTCAAGCAGTT
TTGACCCAATCAGGTTTCCTGAGGATCAGTTCAATGTTG

CAGTTGATCTTCTGATTGAGAACAGTCAG
GCACTGGTGGAACAGTCAAACAAAATTCTAAACAGTGC
AGAAAAAGGAAACACTGGCTTCATTATTGTAATAATTT
TGGTTGCTGTTCTTGGTTTAACCATGATTTCAGTGAGCA
TCATCATCATAATCAAGAAAACAAGGAAGCCCACAGGA
GCACCTCCAGAGCTGAATGGTGTACCAACGGCGGTTT CATAACCACATAGTTAG Codon
ATGAGCTGGAAGGTGATGATCATCATCAGCCTGCTGATCACCCC 55 optimized
CCAGCACGGCCTGAAGGAGAGCTACCTGGAGGAGAGCTGCAGCA nucleic acid
CCATCACCGAGGGCTACCTGAGCGTGCTGAGAACCGGCTGGTAC sequence
ACCAACGTGTTACCCCTGGAGGTGGGCGACGTGGAGAACCTGAC encoding hMPV
CTGCACCGACGGCCCCAGCCTGATCAAGACCGAGCTGGACCTGA F protein of
CCAAGAGCGCCCTGAGAGAGCTGAAGACCGTGAGCGCCGACCAG strain
CTGGCCAGAGAGGAGCAGATCGAGAACCCCAGACAGAGCAGATT HMPV/USA/C2-
CGTGCTGGGCGCCATCGCCCTGGGCGTGGCCACCGCCGCGCCG 175/2005/B
TGACCGCCGGCATCGCCATCGCCAAGACCATCAGACTGGAGAGC
GAGGTGAACGCCATCAAGGGCGCCCTGAAGCAGACCAACGAGGC
CGTGAGCACCTGGGCAACGGCGTGAGAGTGCTGGCCACCGCCG
TGAGAGAGCTGAAGGAGTTCGTGAGCAAGAACCTGACCAGCGCC
ATCAACAGAAACAAGTGCGACATCGCCGACCTGAAGATGGCCGT
GAGCTTCAGCCAGTTCAACAGAAGATTCTGAACGTGGTGAGAC
AGTTCAGCGACAACGCCGGCATCACCCCCGCCATCAGCCTGGAC
CTGATGACCGACGCCGAGCTGGCCAGAGCCGTGAGCTACATGCC
CACCAGCGCCGGCCAGATCAAGCTGATGCTGGAGAACAGAGCCA
TGGTGAGAAGAAAGGGCTTCGGCATCCTGATCGGCGTGACGGC
AGCAGCGTGATCTACATGGTGCAGCTGCCCATCTTCGGCGTGAT
CGACACCCCCTGCTGGATCATCAAGGCCGCCCCAGCTGCAGCG
AGAAGAACGGCAACTACGCCTGCCTGCTGAGAGAGGACCAGGGC
TGGTACTGCAAGAACGCCGGCAGCACCGTGTACTACCCCAACGA
GAAGGACTGCGAGACCAGAGGGCGACCACGTGTTCTGCGACACCG
CCGCCGGCATCAACGTGGCCGAGCAGAGCAGAGAGTGCAACATC
AACATCAGCACCACTACCCCTGCAAGGTGAGCACCGGCAG
ACACCCCATCAGCATGGTGGCCCTGAGCCCCCTGGGCGCCCTGG
TGGCCTGCTACAAGGGCGTGAGCTGCAGCATCGGCAGCAACAGA
GTGGGCATCATCAAGCAGCTGCCCCAAGGGCTGCAGCTACATCAC
CAACCAGGACGCCGACACCGTGACCATCGACAACACCGTGTACC
AGCTGAGCAAGGTGGAGGGCGAGCAGCACGTGATCAAGGGCAGA
CCCGTGAGCAGCAGCTTCGACCCCATCAGATTCCCCGAGGACCA
GTTCAACGTGGCCCTGGACCAGGTGTTTCGAGAGCATCGAGAAC
GCCAGGCCCTGGTGGAGCAGAGCAACAAGATCCTGAACAGCGCC
GAGAAGGGCAACACCGGCTTCATCATCGTGATCATCCTGGTGGC
CGTGCTGGGCCTGACCATGATCAGCGTGAGCATCATCATCATCA
TCAAGAAGACCAGAAAGCCCACCGGCGCCCCCCCCGAGCTGAAC
GGCGTGACCAACGGCGGCTTCATCCCCACAGCTGA Wild-type
ATGTCTTGGAAAGTGGTGATCATTTTTTTCATTGCTAATAACACC 56 Nucleic acid
TCAACACGGTCTTAAAGAAAGCTACCTAGAAGAATCATGTAGCA sequence
CTATAACTGAGGGATATCTTAGTGTTCTGAGGACAGGTTGGTAT encoding hMPV
ACCAACGTTTTTACATTAGAGGTGGGTGATGTAGAAAACCTTAC F protein of
ATGTTCTGATGGACCTAGCCTAATAAAAACAGAATTAGATCTGA strain
CCAAAAGTGCATAAGAGAGCTCAAAACAGTCTCTGCTGACCAA HMPV/ARG/107/
TTGGCAAGAGAGGAACAAATTGAGAATCCCAGACAATCTAGGTT 2002/A

TGTGTTCTAGGAGCAATATAGCACTCGGTGTTGCAACACAGCAGCTGCAG
 TCACAGCAGGTGTTGCAATTGCCAAAACCATCCGGCTTGAGAGT
 GAAGTCACAGCAATTAAGAATGCCCTCAAACGACCAATGAAGC
 AGTATCTACATTGGGGAATGGAGTTCGAGTGTTAGCAACTGCAG
 TGAGAGAGCTGAAAGACTTTGTGAGCAAGAATTTAACCCGTGCA
 ATCAACAAAAACAAGTGTGACATTGATGACCTAAAAATGGCCGT
 TAGCTTCAGTCAATTCAACAGAAGGTTTCTAAATGTTGTGCGGC
 AATTTTCAGACAATGCTGGAATAACACCAGCAATATCTCTGGAC
 TTAATGACAGATGCTGAACTAGCCAGGGCCGTTTCTAACATGCC
 GACATCTGCAGGACAAATAAAATTGATGTTGGAGAACCGTGCGA
 TGGTGCGAAGAAAGGGGTTCGGAATCCTGATAGGGGTCTACGGG
 AGCTCCGTAATTTACATGGTGCAGCTGCCAATCTTTGGCGTTAT
 AGACACGCCTTGCTGGATAGTAAAAGCAGCCCCTTCTTGTTCCG
 AAAAAAAGGGAACTATGCTTGCCTCTTAAGAGAAGACCAAGGG
 TGGTATTGTCAGAATGCAGGGTCAACTGTTTACTACCCAAATGA
 GAAAGACTGTGAAACAAGAGGAGACCATGTCTTTTGCGACACAG
 CAGCAGGAATTAATGTTGCTGAGCAATCAAAGGAGTGCAACATC
 AACATATCCACTACAAATTACCCATGCAAAGTCAGCACAGGAAG
 ACATCCTATCAGTATGGTTGCACTGTCTCCTCTTGGGGCTCTGG
 TTGCTTGCTACAAAGGAGTAAGCTGTTCCATTGGCAGCAACAGA
 GTAGGGATCATCAAGCAGCTGAACAAAGGTTGCTCCTATATAAC
 CAACCAAGATGCAGACACAGTGACAATAGACAACACTGTATATC
 AGCTAAGCAAAGTTGAGGGTGAACAGCATGTTATAAAAGGCAGA
 CCAGTGTC AAGCAGCTTTGATCCAATCAAGTTTCCTGAAGATCA
 ATTCAATGTTGCACTTGACCAAGTTTTTTGAGAACATTGAAAACA
 GCCAGGCCTTGGTAGATCAATCAAACAGAATCCTAAGCAGTGCA
 GAGAAAGGGAACTGGCTTCATCATTGTAATAATTCTAATTGC
 TGTCTTTGGCTCTAGCATGATCCTAGTGAGCATCTTCATTATAA
 TCAAGAAAACAAAGAAACCAACGGGAGCACCTCCAGAGCTGAGT
 GGTGTCACAAACAATGGCTTCATACCACACAGTTAG Codon
 ATGAGCTGGAAGGTGGTGATCATCTTCAGCCTGCTGAT 57 optimized
 CACCCCCCAGCACGGCCTGAAGGAGAGCTACCTGGAGG nucleic acid
 AGAGCTGCAGCACCATCACCGAGGGCTACCTGAGCGTG sequence
 CTGAGAACCGGCTGGTACACCAACGTGTTACCCCTGGA encoding hMPV
 GGTGGGCGACGTGGAGAACCTGACCTGCAGCGACGGCC F protein of
 CCAGCCTGATCAAGACCGAGCTGGACCTGACCAAGAGC strain
 GCCCTGAGAGAGCTGAAGACCGTGAGCGCCGACCAGCT HMPV/ARG/107/
 GGCCAGAGAGGAGCAGATCGAGAACCCCAGACAGAGC 2002/A
 AGATTCGTGCTGGGCGCCATCGCCCTGGGCGTGGCCAC
 CGCCGCCGCCGTGACCGCCGGCGTGGCCATCGCCAAGA
 CCATCAGACTGGAGAGCGAGGTGACCGCCATCAAGAAC
 GCCCTGAAGACCACCAACGAGGCCGTGAGCACCCCTGGG
 CAACGGCGTGAGAGTGCTGGCCACCGCCGTGAGAGAGC
 TGAAGGACTTCGTGAGCAAGAACCTGACCAGAGCCATC
 AACAAGAACAAGTGCGACATCGACGACCTGAAGATGG
 CCGTGAGCTTCAGCCAGTTCAACAGAAGATTCCTGAAC
 GTGGTGAGACAGTTCAGCGACAACGCCGGCATCACCCC
 CGCCATCAGCCTGGACCTGATGACCGACGCCGAGCTGG
 CCAGAGCCGTGAGCAACATGCCACACGCGCCGCCAG
 ATCAAGCTGATGCTGGAGAACAGAGCCATGGTGAGAAG

AAAGGCTTCGGCTATCGGCGTGTACGGCAGCA
GCGTGATCTACATGGTGCAGCTGCCCATCTTCGGCGTGA
TCGACACCCCCTGCTGGATCGTGAAGGCCGCCCCCAGC
TGCAGCGAGAAGAAGGGCAACTACGCCTGCCTGCTGAG
AGAGGACCAGGGCTGGTACTGCCAGAACGCCGGCAGC
ACCGTGTACTACCCCAACGAGAAGGACTGCGAGACCAG
AGGCGACCACGTGTTCTGCGACACCGCCGCGGCATCA
ACGTGGCCGAGCAGAGCAAGGAGTGCAACATCAACATC
AGCACCACCAACTACCCCTGCAAGGTGAGCACCGGCAG
ACACCCCATCAGCATGGTGGCCCTGAGCCCCCTGGGCG
CCCTGGTGGCCTGCTACAAGGGCGTGAGCTGCAGCATC
GGCAGCAACAGAGTGGGCATCATCAAGCAGCTGAACA
AGGGCTGCAGCTACATCACCAACCAGGACGCCGACACC
GTGACCATCGACAACACCGTGTACCAGCTGAGCAAGGT
GGAGGGCGAGCAGCACGTGATCAAGGGCAGACCCGTG
AGCAGCAGCTTCGACCCCATCAAGTTCCCCGAGGACCA
GTTCAACGTGGCCCTGGACCAGGTGTTTCGAGAACATCG
AGAACAGCCAGGCCCTGGTGGACCAGAGCAACAGAAT
CCTGAGCAGCGCCGAGAAGGGCAACACCGGCTTCATCA
TCGTGATCATCCTGATCGCCGTGCTGGGCAGCAGCATG
ATCCTGGTGAGCATCTTCATCATCATCAAGAAGACCAA
GAAGCCCACCGGCGCCCCCCCCCGAGCTGAGCGGCGTGA
CCAACAACGGCTTCATCCCCACAGCTGA

(191) TABLE-US-00007 TABLE 5 Signal Sequences Human RSV F
MELLILKANAITTILTAVTFCFASG 23 signal sequence of the F protein of RSV
strain A2 (first 25 amino acids of SEQ ID NO: 6) hMPV F signal
MSWKVVIIFSLITPQHG 24 sequence of the F protein of hMPV strains CAN00-
16 (SEQ ID NO: 16), HMPV/*Homo sapiens*/PER/FPP 00726/2011/A (SEQ ID NO: 52)
and HMPV/ARG/10 7/2002/A (SEQ ID NO: 56) Bovine RSV F
MATTAMRMIISIIFISTYVTHITLC 60 signal sequence of the F protein of ATCC51908
strain (SEQ ID NO: 33)

(192) TABLE-US-00008 TABLE 6 cDNA of genome of NDV Strains cDNA of
accaaacagagaatccgtgagttacgataaaaaggcgaaggagcaattgaagtcgcacggg 47 genomic
tagaaggtgtgaatctcgagtgcgagcccgaagcacaactcgagaaagccttctgccaac sequence of
atgtcttccgtatttgatgagtacgaacagctcctcgcggtcagactcgcccaatggagct NDV strain
catggagggggagaaaaaggagtagctttaaagtagacgtcccgggtattcactcttaaca LaSota
gtgatgaccagaagatagatggagctttgtggtattctgcctccggattgctgtagcgaag
atgccaacaaaccactcaggcaaggtgctctcatatctctttatgctcccactcacaggtaat
gaggaacctgttgccCttgcagggaaacagaatgaagccacattggccgtgcttgagatt
gatggctttgccaacggcacgccccagttcaacaataggagtggagtgtctgaagagagag
cacagagatttgcgatgatagcaggatctctccctcgggcatgcagcaacggaacccccgttc
gtcacagccggggcCgaagatgatgcaccagaagacatcaccgataccctggagaggat
cctctctatccaggctcaagtatgggtcacagtagcaaaagccatgactgcgtatgagactg
cagatgagtcggaacaaggcgaatcaataagtatatgcagcaaggcaggggtccaaaaga
aatacatcctctaccccgtatgcaggagcacaatccaactcacgatcagacagtctcttgag
tccgcatcttttggtagcgagctcaagagaggccgcaacacggcaggtggtaccttactt
attataacctggtaggggacgtagactcatacatcaggaataaccgggcttactgcattcttctt
gacactcaagtacggaatcaacaccaagacatcagcccttgacttagtagcctctcaggcg
acatccagaagatgaagcagctcatgcgtttgtatcggtgaaaggagataatgcgccgtac
atgacattacttggtagatgtagcagatgagctttgcgctgccgagtagtcacaactttact

cctttgccatgggtatggcatcagtcctagataaaggctactgggaaataccaatttggccaggg
actttatgagcacatcattctggagacttggagtagagtacgctcaggctcagggaagtagc
attaacgaggatatggctgccgagctaaagctaaccacagcagcaaGgaGgggcctggc
agctgctgcccacgggtctccgaGgaGaccagcagcataGacatgcctactcaacaag
tcggagtcctcactgggcttagcaggggggggtcccaagctctacaaggcggatcgaata
gatcgcaagggcaaccagaagccgggggatggggagacccaattcctggatctgatgaga
gcggtagcaaatagcatgagggaggcgccaaactctgcacagggcactccccaatcggg
gcctccccaaactcctgggccatcccaagataacgacaccgactgggggtattgatggaca
aaaccagcctgcttccacaaaaacatcccaatgccctcacccgtagtcgacccctcgattg
cggctctatatgaccacaccctcaaacaacatccccctcttctcctccccctgctgtaca
actAcgTacgccctagataccacaggcacaatgcggctcactaacaatcaaaacagagc
cgagggaattagaaaaaagtacgggtagaagagggatattcagagatcagggcaagtctc
ccgagtctctgctctctcctctacctgatagaccaggacaaacatggccacctttacagatgc
agagatcgacgagctatttgagacaagtggaaactgtcattgacaacataattacagcccagg
gtaaaccagcagagactgttggaaggagtgaatcccacaaggcaagaccaaggtgctga
gcgcagcatgggagaagcatgggagcatccagccaccggccagtcaagacaacccgat
cgacaggacagatctgacaaacaacatccacacccgagcaaacgaccccgcatgacag
cccggccgacatccggccagcagccccccacccaggccacagacgaagccgctgcac
acacagCtcaggaccggagcaagcaactctctgctgttgatgcttgacaagctcagcaata
aatcgtccaatgctaaaaaggggcccatggctcgagccccaagagggggaatcaccaacgtc
cgactcaacagcaggggagtcacccagtcgcggaaacagtcaggaaagaccgcagaa
ccaagtcaaggccgcccctggaaaccagggcacagacgtgaacacagcatatcatggac
aatgggaggagtcacaactatcagctgggtgaacccctcatgctctccgatcaaggcagag
ccaagacaataccctgtatctgcggatcatgtccagccacctgtagacttgtgcaagcgat
gatgtctatgatggaggcgatatcacagagagtaagtaagggtgactatcagctagatctgtc
ttgaaacagacatcctccatccctatgatgcggtcggaaatccaacagctgaaaacatctgtt
gcagtcattggaagccaacttgggaatgatgaagattctggatcccgggtgtgccaacatttca
tctctgagtgatctacgggcagttgcccgatctcacccgggttttagtttcaggccctggagacc
cctctccctatgtgacacaaggaggcgaaatggcacttaataaaactttcgcaaccagtgcca
catccatctgaattgattaaacccgccactgcatgcgggcctgatataggagtggaaaagga
cactgtccgtgcattgatcatgtcacgccaatgcacccgagttcttcagccaagctcctaag
caagttagatgcagccgggtcgatcgaggaaatcaggaaaatcaagcgccttgctctaaat
ggctaattactactgccacacgtagcgggtccctgtccactcggcatcacacggaatctgca
ccgagttccccccgcGgaccaaggtccaactctccaagcggcaatcctctctcgcttct
cagccccactgaatgAtcgcgtaaccgtaattaatctagctacatttaagattaagaaaaat
acgggtagaattggagtgccccaattgtgccaagatggactcatctaggacaattgggctgt
actttgattctgcccattctttagcaacctgttagcatttccgatcgtcctacaagAcacagga
gatgggaagaagcaaatcgccccgcaatataggatccagcgccttgacttgaggactgata
gtaaggaggactcagttatcatcaccacctatggattcatctttcaagttgggaatgaagaagc
cacCgtcggcatgatcgatgataaaccgaagcgcgagttacttccgctgcgatgctctgcc
taggaagcgtcccaaataccgggagaccttattgagctggcaagggcctgtctcactatgata
gtcacatgcaagaagagtgaactaatactgagagaatggttttctcagtagtgaggcacc
ccaagtgtgcaaagctgtagggtgtggcaacaaatactcatcagtgaatgcagtcaagc
acgtgaaagcgccagagaagattcccgggagtggaaccctagaatacaaggtgaactttgt
ctccttgactgtgtaccgaagaGggatgtctacaagatcccagctgcagtattgaaggtttc
tggctcgagtctgtacaatcttgcgctcaatgtcactattaatgtggaggtagacccgaggagt
cctttgggttaaactCtgtctaagtctgacagcggatactatgctaacctcttctgcatattgga
cttatgaccacTgtagataggaaggggaagaaaagtacatttgacaagctggaaaagaaa
ataaggagccttgatctatctgtcgggctcagtgatgtgctcgggccttccgtgttggtaaaag
caagagggtgcacggactaagcttttggcacctttcttcttagcagtgaggacagcctgctatc

ccatagcaaatgcttctcctcaggtggccaagatactctggagtcaaaaccgcgtgcctgcgg
agcggtaaaaatcattatccaagcaggtacccaacgcgctgtcgcagtgaccgccgaccacg
aggttacctctactaagctggagaaggggcacacccttgccaaatacaatccttttaagaaat
aagctgcgtctctgagattgcgctccgcccactcaccagatcatcatgacacaaaaaacta
atctgtcttgattattacagtttagttacctgtctatcaagttagaaaaaacacgggtagaagatt
ctggatcccgggtggcgccctccaggtgcaagatgggctccagaccttctaccaagaaccc
agcacctatgatgctgactatccggggtgcgctgggtactgagttgcatctgtccggcaaactc
cattgatggcaggcctcttgcaagctgcaggaattgtggttacaggagacaaagccgtcaaca
tatacacctcatcccagacaggatcaatcatagttaagctcctcccgaatctgcccaaggata
aggaggcatgtgcgaaagcccccttgatgcatacaacaggacattgaccactttgctcacc
ccccttggtgactctatccgtaggatacaagagtctgtgactacatctggaggggggagaca
ggggcgcccttataggcgccattattggcggtgtggctcttgggggtgcaactgccgcacaaa
taacagcggccgcagctctgatacaagccaaacaaaatgctccaacatcctccgacttaa
gagagcattgccgcaaccaatgaggctgtgcatgaggctactgacggattatcgcaactag
cagtggcagttgggaagatgcagcagtttgttaatgaccaatttaaaaaacagctcaggaat
tagactgcatcaaaattgcacagcaagttgggtgtagagctcaacctgtacctaaccgaattga
ctacagtattcggaccacaaatcacctcacctgctttaacaagctgactattcaggcactttac
aatctagctgggtggaaatatggattacttattgactaagttagggttagggacaatcaactca
gctcattaatcggtagcggcttaatcaccgtaacctattctatacgactcacagactcaact
cttgggtatacaggttaactctaccttcagtcgggaacctaaataatatgctgccacctacttg
gaaaccttatccgtaagcacaccaggggatttgctcggcacttgtcccAaaagtgggtga
cacaggtcggttctgtgatagaagaactgacacctatactgtatagaactgacttagattta
tattgtacaagaatagtaacgttccctatgtcccctgggtatttattctgcttgagcggcaatacg
tcggcctgtatgtactcaaagaccgaaggcgcacttactacaccatacatgactatcaaaggt
tcagtcacgccaaactgcaagatgacaacatgtagatgtgtaaacccccgggtatcatatcg
caaaactatggagaagccgtgtctctaatagataaacaatcatgcaatgtttatccttaggcg
ggataactttaaggctcagtggggaattcgatgtaacttatcagaagaatatctcaatacaaga
ttctcaagtaataataacaggcaatcttgatatctcaactgagcttgggaatgtcaacaactcg
atcagtaatgcttgaataagttagaggaaagcaacagaaaactagacaaagtcaatgtcaa
actgactagcacatctgctctcattacctatatcgttttgactatcatactcttgttttggtatactt
agcctgattctagcatgctacctaattgtacaagcaaaaaggcgcaaaaaagaccttattatgg
cttgggaataatactctagatcagatgagagccactacaaaaatgtgaacacagatgaggaa
cgaaggtttccctaataagtaatttgtgtgaaagtcttggtagtctgtcagttcagagagttaaga
aaaaactaccgggttagatgaccaaaggacgatatacgggtagaacggtaagagaggcc
gcccccaattgcgagccaggcttcacaacctccgttctaccgcttcaccgacaacagtcctc
aatcatggaccgcgccgttagccaagttgcgttagagaatgatgaaagagaggcaaaaaat
acatggcgcttgatattccggattgcaatcttattcttaacagtagtgacctgggtatatctgta
gcctcccttttatatagcatgggggctagcacacctagcgatctttaggcataccgactagg
atttcagggcagaagaaaagattacatctacacttgggtccaatcaagatgtagtagatagg
atatataagcaagtggcccttgagtctccgttggcattgttaaatactgagaccacaattatgaa
cgcaataacatctctcttatcagattaatggagctgcaaaacaacagtgggtgggggggcac
ctatccatgaccagattatataggggggtagaggcaaagaactcattgtagatgatgctagt
atgtcacatcattctatccctctgcatttcaagaacatctgaattttatcccggcgctactacag
gatcaggttgcaactcgaataacctcatttgacatgagtgtacccattactgctacaccataat
gtaattattgtctggatgcagagatcactcacattcatatcagtatttagcacttgggtgtgctccg
gacatctgcaacaggaggagggtattcttttctactctgcgttccatcaacctggacgacacca
aaatcggaagtcttgagtgagtgcaactcccctggggtgtgatatgctgtgctcgaaagt
cacggagacagaggaagaagattataactcagctgtccctacgcggatggtacatgggag
gttaggggttcgacggccagtagcacgaaaaggacctaagatgtcacaacattattcggggact
gggtggccaactaccaggagtaggggggtggatcttttattgacagccgcgtatggttctca

gtctacggagggttaaaacattcaccagtgacactgtacaggaagggaatatgtgat
atacaagcgatacaatgacacatgccagatgagcaagactaccagattcgaatggccaag
tcttcgtataagcctggacgggttggtgggaaacgcatacagcaggctatcttatcaagg
tgtcaacatccttaggcgaagaccgggtactgactgtaccgccaacacagtcacactcatg
ggggccgaaggcagaattctcacagtagggacatctcatttctgtatcaacgagggcatca
tacttctctcccgcgttattatatcctatgacagtcagcaaaaaacagccactcttcatagtcct
tatacattcaatgccttcactcggccaggtagtatcccttgccaggcttcagcaagatgcccc
actcgtgtgttactggagtctatacagatccatatcccctaattcttctatagaaaccacacttg
cgaggggtattcgggacaatgcttgatgggtgtacaagcaagacttaaccctgcgtctgcagt
attcgatagcacatcccgcatcgcattactcgagtgaagtcaagcagtaacaaagcagcat
acacaacatcaactgttttaaagtggtcaagactaataagacctattgtctcagcattgctgaa
atatctaatactctcttcggagaattcagaatcggtcccgttactagttgagatcctcaaagatga
cgggggttagagaagccaggtctggctagttgagtcattataaaggagttggaaagatggca
ttgtatcacctatcttctgcgacatcaagaatcaaaccgaatgccggcgcggtgctcgaattcca
tgttgccagttgaccacaatcagccagtgctcatgcgatcagattaagccttgtaAaGtct
cttgattaagaaaaaatgtaagtggaatgagatacaaggcaaaacagctcatgggTaaCa
atacgggtaggacatggcgagctccggctcctgaaagggcagagcatcagattatcctacca
gagTcacacctgtcttcaccattgggtcaagcacaactactctattactggaaattaactggg
ctaccgcttcctgatgaatgtgacttcgaccacctcattctcagccgacaatggaaaaaata
cttgaatcggcctctcctgatactgagagaatgataaaactcgggaagggcagtacaccaaac
tcttaaccacaattccagaataaccggagtgctccaccccagggtgttagaaGaactggcta
atattgagggtccagattcaaccaaaaaatttcggaagattgagaagaagatccaaattcaca
acacgagatatggagaactgttcacaaggctgtgtacgcatatagagaagaaactgctggg
gtcatcttggtctaacaatgtccccgggtcagaggagttcagcagcattcgtacggatccggc
attctgggttcactcaaaatgggtccacagccaagtttgcattggctccatataaaacagatccag
aggcatctgatgggtggcagctaGgacaagggtctgcggccaacaaattgggtgatgtaacc
ataaggtaggccaagtctttgtcactcctgaactgtcgttgtagcgatacgaatgagaacaa
gttcacatgtcttaccaggaactgtattgatgtatgcagatatgatggagggcagagatatg
gtcaacataatatcaaccacggcggtgcatctcagaagcttatcagagaaaattgatgacattt
tgcggttaatagacgctctggcaaaaagacttgggtaatcaagtctacgatgttgatcactaat
ggaggggatttgcatacggagctgtccagctactcagccgtcaggtacatttgaggagattt
cttcgattcaacctgcaggagcttaaagacattctaattggcctcctcccaatgatatagca
gaatccgtgactcatgcaatcgctactgtattctctggtttagaacagaatcaagcagctgaga
tgttggtgtgttgctgtgtgggtcacccactgcttgagtccttgattgcagcaaaaggcagt
caggagccaaatgtgcgcaccgaaaatggtagactttgatatgatccttcaggtactgtcttc
ttcaagggaacaatcatcaacgggtacagaaagaagaatgcagggtgtgtggccgcgagtc
aaagtggatacaatatatgggaagggtcattgggcaactacatgcagattcagcagagatttca
cacgatatcatgttgagagagtataagagttatctgcacttgaatttgagccatgtatagaatat
gacctgtcaccaacctgagcatgttcctaaaagacaaggcaatcgcacaccccaacgata
attggcttgccctgtttaggcggaaccttctctccgaagaccagaagaaacatgtaaaagaa
gcaacttcgactaatcgctccttgatagagtttttagagtcaaattgatccatataaaagag
atggaatatctgacgacccttgagtaccttagagatgacaatgtggcagtatcatactcgctca
aggagaagggaagtgaaagttaatggacggatcttcgctaagctgacaaagaagttaaggaa
ctgtcaggtgatggcggaagggtacctagccgatcagattgcacctttcttcagggaatgg
agtcattcaggatagcatatccttgaccaagagtatgctagcgatgagtcaactgtctttaaca
gcaataagaaacgtatcactgactgtaaagaaagagtatctcaaaccgcaatcatgatccga
aaagcaagaaccgtcggagagttgcaaccttcatacaactgacctgcaaaagtactgtctt
aattggagatatcagacaatcaaattgttcgctcatgccatcaatcagttgatgggcctacctc
acttcttcgaatggattcacctaagactgatggacactacgatgttcgtaggagacctttcaat
cctccaagtgaccctactgactgtgacctctcaagagtcctaattgatgacatatatattgtca

gtgccagagggggtatcgaaaggattatgccagaagctatggacaatgatctcaattgctgca
atccaacttgctgcagctagatcgattgtcgtgttgctgtatggtaggggtgataatcaag
taatagcagtaacgagagaggtaagatcagacgactctccggagatgggtgtgacacagttg
catcaagccagtataatttctcaaggaattaattcatgtcaatcatttgattggccataatttga
aggatcgtgaaaccatcagggtcagacacattcttcatatacagcaaacgaatcttcaaagatg
gagcaatcctcagtcaagtcctcaaaaattcatctaaattagtgtctagtgtcaggtgatctcagt
gaaaacaccgtaatgtcctgtgccaacattgccttactgtagcacggctatgcgagaacgg
gcttcccaaagacttctgttactatttaaactatataatgagttgtgtgcagacatacttgactct
gagtttccatcaccaacaattcgcacccccgatcttaatcagtcgtggattgaggacatctcttt
tgtgcactcatatgttctgactcctgcccaattaggggggactgagtaaccttcaatactcaagg
ctctacactagaaatatcggtgacccgggggactactgcttttgagagatcaagcgactaga
agcagtgggattactgagtcctaacattatgactaatatcttaactaggccgcctgggaatgg
agattggggccagtctgtgcaacgaccatactctttcaattttgagactgttgcaagcccaa
attgttcttaagaaacatacgcaaagagtcctatttgaacctgttcaaacccttattgtctgga
gtgcacacagaggataatgaggcagaagagaaggcattggctgaattcttgcttaatcaaga
gggtgattcatccccgcgttgccgatgccatcatggaggcaagctctgtaggtaggagaaag
caaattcaagggctgttgacacaacaaacaccgtaattaagattgcgcttactaggaggcca
ttaggcataagaggctgatgcggatagtcattattctagcatgcatgcaatgctgtttagag
acgatgttttctccagtagatccaaccaccccttagtctcttctaatatgtgttctctgacact
ggcagactatgcacggaatagaagctgggtcacctttgacgggaggcaggaaaatactgggt
gtatctaactctgatacgatagaactcgtagagggtgagattcttagtgtaagcggagggtgt
acaagatgtgacagcggagatgaacaatttacttgggtccatcttccaagcaatatagaattga
ccgatgacaccagcaagaatcctccgatgagggtaccatatctcgggtcaaagacacagga
gaggagagctgcctcacttgcaaaaatagctcatatgtcgccacatgtaaaggctgccctaa
gggcatcatccgtgttgatctgggcttatggggataatgaagtaaattggactgctgctcttac
gattgcaaaatctcgggtgaatgtaaacttagagtatcttcgggttactgtcccctttaccacgg
ctgggaatcttcaacatagactagatgatggtataactcagatgacattcacccctgcatctct
ctacaggGgtgcaccttacattcacatatccaatgattctcaaaggctgttactgaagaagg
agtcaaagagggggaatgtggtttaccaacagatcatgctcttgggtttatcttaatcgaatcg
atctttccaatgacaacaaccaggacatatgatgagatcacactgcacctacatagtaaattta
gttgctgtatcagagaagcacctgttcgggttccttcgagctacttgggggtgtaccggaact
gaggacagtgaacctaaataagtttatgtatgatcctagccctgtatcgaggaggagacttgc
gagacttgacttagctatcttcaagagttatgagcttaatctggagtcatatccacgatagag
ctaataaacattcttcaatatccagcgggaagtgtgattggccagtctgtggtttcttatgatgaa
gatacctcataaagaatgacgccataatagtgtatgacaatacccgaaattggatcagtga
gctcagaattcagatgtgggtccgcctatttgaatatgcagcacttgaagtgtcctcgactgttc
ttaccaactctattacctgagagtaagaggcctGgacaatattgtcttatatatgggtgattata
caagaatatgccaggaattctactttccaacattgcagctacaatatctcatcccgctcattcattc
aagggttatcagtgagtgggcctgggtcaaccatgacggatcacaccaacttcagatacggatt
ttatcgaaatgtctgcaaaactattagatcttgcacccgacgtgtgatctccggcttatattcag
gaaataagtatgatctgctgttcccatctgtcttagatgataacctgaatgagaagatgcttcag
ctgataccccggttatgctgtctgtacacgggtactctttgctacaacaagagaaatccccgaaa
taagaggcttaactgcagaagagaaatgttcaatactcactgagtatttactgtcggatgctgt
gaaaccattacttagccccgatcaagtgagctctatcatgtctcctaacataattacattccag
ctaattctgtactacatgtctcgggaagagcctcaatttgatcagggaaggaggagacagggat
actatcctggcgttgtgttcccccaagagccattattagagttcccttctgtgcaagatattggt
gctcgagtgaagatccattcacccgacaacctgcggcatttttgcaagagttagattgagt
gctccagcaagggtatgacgcattcacacttagtcagattcatcctgaactcacatctccaaatc
cggaggaagactacttagtacgatacttgttcagagggatagggactgcatcttctcttggtta
taaggcatctcatctcctttctgtacccgaggtaagatgtgcaagacacgggaactccttatac

ttagctgaaggaggagcgagccatcatgagctcttctcgaactgcatgtaccacatgaaactatc
tattacaatacgcctcttttcaaagatgagatgaacccccgcaacgacatttcgggcccgaaccca
actcagtttttgaattcgggtgtttataggaatctacaggcgagggaacatgcaaagatggatt
tgtccaagagttccgtccattatggagagaaaatacagaggaaagCgacctgacctcagat
aaagTagtgggggtatattacatctgcagtgccctacagatctgtatcattgctgcattgtgaca
ttgaaattcctccagggtccaatcaaagcttactagatcaactagctatcaatttatctctgattg
ccatgcattctgtaaggggagggcggggtagtaatcatcaaagtgttgatgcaatgggatact
actttcatctactcatgaacttggttgcctcgtgtccacaaaaggatatattctctctaattgggtat
gcatgtcgaggagatattggagtgtacctgggtatttgcattgggttacctggggcgggcctaca
tttgatcatgaggtggtgaggatggcGaaaactctgggtgcagcggcacggtacgctTttgt
ctaaatcagatgagatcacactgaccaggtattcacctcacagcggcagcgtgtgacagac
atcctatccagtcctttaccaagattaataaagtacttgaggaagaatattgacactgcgctgat
tgaagccgggggacagcccgtccgtccattctgtgcggagagtgctggtgagcacgctagc
gaacataactcagataaccagatCatcgctagtcacattgacacagttatccggtctgtgat
atatatggaagctgaggggtgatctcgctgacacagttatttctatttacccttacaatctctctac
tgacgggaaaaagaggacatcacttaAacagtgcacgagacagatcctagagggtacaat
actaggtcttagagtcgaaaatctcaataaaataggcgatataatcagcctagtgcctaaagg
catgatctccatggaggaccttatcccactaaggacatactgaagcatagtacctgccctaa
atatttgaaggctgtcctaggtattacaaactcaaagaaatgtttacagacacttctgtaCtgt
acttgactcgtgctcaacaaaaattctacatgaaaactataggcaatgcagtcaaaggatatta
cagtaactgtgactcttaacgaaaatcacatattaataggctcctttttggccaattgtattctgt
tgatttaatcatattatgttagaaaaaagttgaacctgactccttaggactcgaattcgaactca
aataaatgtcttaaaaaaaggttgcgacaattattcttgagtgtagctctgcctaccacaaatc tttgtttggt cDNA of
ACCAAACAGAGAATCCGTAAGTTACGATAAAAGGCGA 48 genomic
AGGAGCAATTGAAGTCGCACGGGTAGAAGGTGTGAATC sequence of
TCGAGTGCGAGCCCGAAGCACAAACTCGAGGAAGCCTT NDV strain
CTGCCAACATGTCTTCCGTATTCGACGAGTACGAACAG Hitchner B1
CTCCTCGCGGCTCAGACTCGCCCCAATGGAGCTCATGG
AGGGGGGGGAGAAAGGGAGTACCTTAAAAGTAGACGTC
CCGGTATTCACTCTTAACAGTGATGACCCAGAAGATAG
GTGGAGCTTTGTGGTATTCTGCCTCCGGATTGCTGTTAG
CGAAGATGCCAACAAACCACTCAGGCAAGGTGCTCTCA
TATCTCTTTTATGCTCCCACTCACAGGTAATGAGGAACC
ATGTTGCCCTTGCAGGGAAACAGAATGAAGCCACATTG
GCCGTGCTTGAGATTGATGGCTTTGCCAACGGGCACGCC
CCAGTTCAACAATAGGAGTGGAGTGTCTGAAGAGAGAG
CACAGAGATTTGCGATGATAGCAGGATCTCTCCCTCGG
GCATGCAGCAACGGCACCCCGTTTCGTCACAGCCGGGGC
TGAAGATGATGCACCAGAAGACATCACCGATACCCTGG
AGAGGATCCTCTCTATCCAGGCTCAAGTATGGGTCACA
GTAGCAAAAGCCATGACTGCGTATGAGACTGCAGATGA
GTCGGAAACAAGGCGAATCAATAAGTATATGCAGCAAG
GCAGGGTCCAAAAGAAATACATCCTCTACCCCGTATGC
AGGAGCACAACTCAACTCACGATCAGACAGTCTCTTGC
AGTCCGCATCTTTTTTGGTTAGCGAGCTCAAGAGAGGCC
GCAACACGGCAGGTGGTACCTCTACTTATTATAACCTA
GTAGGGGACGTAGACTCATATATCAGGAATACCGGGCT
TACTGCATTCTTCTTGACACTCAAGTACGGAATCAACAC
CAAGACATCAGCCCTTGCACTTAGTAGCCTCTCAGGCG
ACATCCAGAAGATGAAGCAGCTCATGCGTTTGTATCGG

ATGAAAAGCTACCGCTGACATGACATTACTTGG
TGATAGTGACCAGATGAGCTTTGCGCCTGCCGAGTATG
CACAACTTTACTCCTTTGCCATGGGTATGGCATCAGTCC
TAGATAAAGGTACTGGGAAATACCAATTTGCCAGGGAC
TTTATGAGCACATCATTCTGGAGACTTGGAGTAGAGTA
CGCTCAGGCTCAGGGAAGTAGCATTAAACGAGGATATGG
CTGCCGAGCTAAAGCTAACCCCGGCAGCAAGGAGGGGGC
CTGGCAGCTGCTGCCCAACGAGTCTCCGAGGTGACCAG
CAGCATAGACATGCCTACTCAACAAGTCGGAGTCCTCA
CTGGGGCTTAGCGAGGGGGGGATCCCAAGCCCTACAAGGC
GGATCGAATAGATCGCAAGGGCAACCAGAAGCCGGGG
ATGGGGGAGACCCAATTCCTGGATCTGATGAGAGCGGTA
GCAAATAGCATGAGGGAGGGCGCCAAACTCTGCACAGG
GCACTCCCCAATCGGGGGCCTCCCCCAACTCCTGGGGCCAT
CCCAAGATAACGACACCGACTGGGGGTATTGATTGACA
AAACCCAGCCTGCTTCTACAAGAACATCCCAATGCTCTC
ACCCGTAGTCGACCCCTCGATTTGCGGGCTCTATATGACC
ACACCCTCAAACAAACATCCCCCTCTTTCCTCCCTCCCC
CTGCTGTACAACTCCGCACGCCCTAGATAACCACAGGCA
CACCGCGGCTCACTAACAATCAAAACAGAGCCGAGGGA
ATTAGAAAAAAGTACGGGTAGAAGAGGGATATTCAGA
GATCAGGGCAAGTCTCCCGAGTCTCTGCTCTCTCCTCTA
CCTGATAGACCAGGACAAACATGGCCACCTTTACAGAT
GCAGAGATCGACGAGCTATTTGAGACAAGTGGAAGTGT
CATTGACAACATAATTACAGCCCAGGGTAAACCAGCAG
AGACTGTTGGAAGGAGTGCAATCCCACAGGGCAAGACC
AAGGTGCTGAGCGCAGCATGGGAGAAGCATGGGAGCA
TCCAGCCACCGGCCAGTCAAGACAACCTCGATCGACAG
GACAGATCTGACAAACAACCATCCACACCCGAGCAAAC
GACCCCGCACGACAGCCCGCCGGCCACATCCGCTGACC
AGCCCCCCACCCAGGCCACAGACGAAGCCGTCGACACA
CAGCTCAGGACCGGAGCAAGCAACTCTCTGCTGTTGAT
GCTTGACAAGCTCAGCAATAAATCGTCCAATGCTAAAA
AGGGCCCATGGTCGAGCCCCCAAGAGGGGAATCACCAA
CGTCCGACTCAACAGCAGGGGAGTCAACCCAGTCGCGG
AAACAGCCAGGAAAGACTGCAGAACCAAGTCAAGGCC
GCCCCTGGAAACCAGGGCACAGACGTGAACACAGCATA
TCATGGACAATGGGAGGAGTCACAACTATCAGCTGGTG
CAACCCCTCATGCTCTCCGATCAAGGCAGAGCCAAGAC
AATACCCTTGTATCTGCGGATCATGTCCAGCCACCTGTA
GACTTTGTGCAAGCGATGATGTCTATGATGGGGGCGAT
ATCACAGAGAGTAAGTAAGGTTGACTATCAGCTAGATC
TTGTCTTGAAACAGACATCCTCCATCCCTATGATGCGGT
CCGAAATCCAACAGCTGAAAACATCTGTTGCAGTCATG
GAAGCCAACTTGGGAATGATGAAGATTCTGGATCCCGG
TTGTGCCAACATTTTCATCTCTGAGTGATCTACGGGCAGT
TGCCCGATCTACCCGGTTTTAGTTTTAGGCCCTGGAGA
CCCATCTCCCTATGTGATACAAGGAGGCGAAATGGCAC
TTAATAAACTTTTCGCAACCAGTGCCACATCCATCTGAAT
TGATTAAACCCGCCACTGCATGCGGGCCTGATATAGGA

GTGGAGTGAGCGGTGCGATTGATCATGTCACG
CCCAATGCACCCGAGTTCTTCAGCCAAGCTCCTAAGCA
AGTTAGATGCAGCCGGGTCGATCGAGGAAATCAGGAAA
ATCAAGCGCCTTGCTCTAAATGGCTAATTACTACTGCCA
CACGTAGCGGGTCCCTGTCCACTCGGCATCACACGGAA
TCTGCACCGAGTTCCCCCCCCGCAGACCCAAGGTCCAAC
TCTAGAAGCGGCAATCCTCTCTCGCTTCCTCAGCCCCAC
TGAATGATCGCGTAACCGTAATTAATCTAGCTACATTAA
GGATTAAGAAAAAATACGGGTAGAATTGGAGTGCCCCA
ATTGTGCCAAGATGGACTCATCTAGGACAATTGGGCTG
TACTTTGATTCTGCCCATTCTTCTAGCAACCTGTTAGCA
TTTCCGATCGTCCTACAAGACACAGGAGATGGGAAGAA
GCAAATCGCCCCGCAATATAGGATCCAGCGCCTTGACT
CGTGGACTGATAGTAAGGAAGACTCAGTATTCATCACC
ACCTATGGATTCATCTTTCAAGTTGGGAATGAGGAAGC
CACTGTCGGCATGATCGATGATAAACCCAAGCGCGAGT
TACTTTCCGCTGCGATGCTCTGCCTAGGAAGCGTCCCAA
ATACCGGAGACCTTGTTGAGCTGGCAAGGGCCTGTCTC
ACTATGATGGTCACATGCAAGAAGAGTGCAACTAATAC
TGAGAGAATGGTTTTTCTCAGTAGTGCAAGCACCCCAAG
TGCTGCAAAGCTGTAGGGTTGTGGCAAATAAATACTCA
TCAGTGAATGCAGTCAAGCACGTGAAAGCGCCAGAGAA
GATCCCCGGGAGTGGAACCCTAGAATAACAAGGTGAACT
TTGTCTCCTTGACTGTGGTACCGAAGAAGGATGTCTACA
AGATCCCAGCTGCAGTATTGAAGATTTCTGGCTCGAGTC
TGTACAATCTTGCGCTCAATGTCACTATTAATGTGGAGG
TAGACCCGAGGAGTCCTTTGGTTAAATCTCTGTCTAAGT
CTGACAGCGGATACTATGCTAACCTCTTCTTGTCATATTG
GACTTATGACCACCGTAGATAGGAAGGGGAAGAAAGT
GACATTTGACAAGCTGGAAAAGAAAATAAGGAGCCTTG
ATCTATCTGTCGGGCTCAGTGATGTGCTCGGGCCTTCCG
TGTTGGTAAAAGCAAGAGGTGCACGGACTAAGCTTTTG
GCACCTTTCTTCTCTAGCAGTGGGACAGCCTGCTATCCC
ATAGCAAATGCTTCTCCTCAGGTGGCCAAGATACTCTG
GAGTCAAACCGCGTGCCTGCGGAGCGTTAAAATCATT
TCCAAGCAGGTACCCAACGCGCTGTCGCAGTGACCGCT
GACCACGAGGTTACCTCTACTAAGCTGGAGAAGGGGCA
CACCTTGCCAAATAACAATCCTTTTAAGAAATAAGCTGC
GTCTCTGAGATTGCGCTCCGCCCACTCACCCAGATCATC
ATGACACAAAAAACTAATCTGTCTTGATTATTTACAGTT
AGTTTACCTGTCCATCAAGTTAGAAAAAACACGGGTAG
AAGATTCTGGATCCCGGTTGGCGCCCTCCAGGTGCAGG
ATGGGCTCCAGACCTTCTACCAAGAACCCAGCACCTAT
GATGCTGACTATCCGGGTCGCGCTGGTACTGAGTTGCAT
CTGCCCGGCAAACCTCCATTGATGGCAGGCCTCTTGCA
CTGCAGGAATTGTGGTTACAGGAGACAAAGCAGTCAAC
ATATACACCTCATCCCAGACAGGATCAATCATAGTTAA
GCTCCTCCCGAATCTGCCCAAGGATAAGGAGGCATGTG
CGAAAGCCCCCTTGATGCATACAACAGGACATTGACC
ACTTTGCTCACCCCCCTTGGTGACTCTATCCGTAGGATA

CAAGTGTCTGATCTGAGGGGGGAGACAGGG
GCGCCTTATAGGCGCCATTATTGGCGGTGTGGCTCTTGG
GGTTGCAACTGCCGCACAAATAACAGCGGCCGCAGCTC
TGATACAAGCCAAACAAAATGCTGCCAACATCCTCCGA
CTTAAAGAGAGCATTGCCGCAACCAATGAGGCTGTGCA
TGAGGTCACTGACGGATTATCGCAACTAGCAGTGGCAG
TTGGGAAGATGCAGCAGTTTGTTAATGACCAATTTAAT
AAAACAGCTCAGGAATTAGACTGCATCAAAATTGCACA
GCAAGTTGGTGTAGAGCTCAACCTGTACCTAACCGAAT
TGACTACAGTATTCGGACCACAAATCACTTCACCTGCCT
TAAACAAGCTGACTATTCAGGCACTTTACAATCTAGCTG
GTGGGAATATGGATTACTTATTGACTAAGTTAGGTATA
GGGAACAATCAACTCAGCTCATTAAATCGGTAGCGGCTT
AATCACCGGTAACCCTATTCTATACGACTCACAGACTCA
ACTCTTGGGTATACAGGTAACCTCTACCTTCAGTCGGGAA
CCTAAATAATATGCGTGCCACCTACTTGGAAACCTTATC
CGTAAGCACAACCAGGGGATTTGCCTCGGCACTTGTCC
CAAAAGTGGTGACACAGGTCGGTTCTGTGATAGAAGAA
CTTGACACCTCATACTGTATAGAACTGACTTAGATTTA
TATTGTACAAGAATAGTAACGTTCCCTATGTCCCCTGGT
ATTTACTCCTGCTTGAGCGGCAATACATCGGCCTGTATG
TACTCAAAGACCGAAGGCGCACTTACTACACCATATAT
GACTATCAAAGGCTCAGTCATCGCTAACTGCAAGATGA
CAACATGTAGATGTGTAAACCCCCCGGGTATCATATCG
CAAACTATGGAGAAGCCGTGTCTCTAATAGATAAACA
ATCATGCAATGTTTTATCCTTAGGCGGGATAACTTTAAG
GCTCAGTGGGGAATTCGATGTAACCTTATCAGAAGAATA
TCTCAATACAAGATTCTCAAGTAATAATAACAGGCAAT
CTTGATATCTCAACTGAGCTTGGGAATGTCAACAACTCG
ATCAGTAATGCTTTGAATAAGTTAGAGGAAAGCAACAG
AAAAC TAGACAAAGTCAATGTCAAACCTGACCAGCACAT
CTGCTCTCATTACCTATATCGTTTTGACTATCATATCTCT
TGTTTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCT
AATGTACAAGCAAAAGGCGCAACAAAAGACCTTATTAT
GGCTTGGGAATAATACCCTAGATCAGATGAGAGCCACT
ACAAAAATGTGAACACAGATGAGGAACGAAGGTTTCCC
TAATAGTAATTTGTGTGAAAGTTCTGGTAGTCTGTCAGT
TCGGAGAGTTAAGAAAAAACTACCGGTTGTAGATGACC
AAAGGACGATATACGGGTAGAACGGTAAGAGAGGCCG
CCCCTCAATTGCGAGCCAGACTTCACAACCTCCGTTCTA
CCGCTTCACCGACAACAGTCCTCAATCATGGACCGCGC
CGTTAGCCAAGTTGCGTTAGAGAATGATGAAAGAGAGG
CAAAAAATACATGGCGCTTGATATTCCGGATTGCAATC
TTATTCTTAACAGTAGTGACCTTGGCTATATCTGTAGCC
TCCCTTTTATATAGCATGGGGGCTAGCACACCTAGCGAT
CTTGTAGGCATACCGACTAGGATTTCCAGGGCAGAAGA
AAAGATTACATCTACACTTGGTTCCAATCAAGATGTAGT
AGATAGGATATATAAGCAAGTGGCCCTTGAGTCTCCAT
TGGCATTGTAAATACTGAGACCACAATTATGAACGCA
ATAACATCTCTCTCTTATCAGATTAATGGAGCTGCAAAC

ACACGCGGGTGGCGACCGCTATTTCATGACCCAGATTATATAGGGGGGATAGGCCAAAGAACTCATTGTAGATGATGCTAGTGATGTCACATCATTCTATCCCTCTGCATTTCAAGAACATCTGAATTTTATCCCGGCGCCTACTACAGGATCAGGTTGCACTCGAATAACCCTCATTTGACATGAGTGCTACCCATTACTGCTACACCCATAATGTAATATTGTCTGGATGCA GAGATCACTCACACTCACATCAGTATTTAGCACTTGGTG TGCTCCGGACATCTGCAACAGGGAGGGTATTCTTTTCTA CTCTGCGTTCCATCAACCTGGACGACACCCAAAATCGG AAGTCTTGCAGTGTGAGTGCAACTCCCCTGGGTTGTGAT ATGCTGTGCTCGAAAGCCACGGAGACAGAGGAAGAAG ATTATAACTCAGCTGTCCCTACGCGGATGGTACATGGG AGGTTAGGGTTCGACGGCCAATATCACGAAAAGGACCT AGATGTCACAACATTATTCGGGGACTGGGTGGCCAACT ACCCAGGAGTAGGGGGTGGATCTTTTATTGACAGCCGC GTATGGTTCTCAGTCTACGGAGGGTTAAAACCCAATAC ACCCAGTGACACTGTACAGGAAGGGAAATATGTGATAT ACAAGCGATACAATGACACATGCCCAGATGAGCAAGAC TACCAGATTCTGAATGGCCAAGTCTTCGTATAAGCCTGG ACGGTTTGGTGGGAAACGCATACAGCAGGCTATCTTAT CTATCAAAGTGTC AACATCCTTAGGCGAAGACCCGGTA CTGACTGTACCGCCCAACACAGTCACACTCATGGGGGC CGAAGGCAGAATTCTCACAGTAGGGACATCCCATTTCT TGTATCAGCGAGGGTCATCATACTTCTCTCCCGCGTTAT TATATCCTATGACAGTCAGCGACAAAACAGCCACTCTT CATAGTCCTTATACATTCAATGCCTTCACTCGGCCAGGT AGTATCCCTTGCCAGGCTTCAGCAAGATGCCCCAACTC GTGTGTTACTGGAGTCTATACAGATCCATATCCCCTAAT CTTCTATAGAAACCACACCTTGCGAGGGGTATTCGGGA CAATGCTTGATGGTGAACAAGCAAGACTTAACCCTGCG TCTGCAGTATTCGATAGCACATCCCGCAGTCGCATAACT CGAGTGAGTTCAAGCAGCATCAAAGCAGCATAACAAAC ATCAACTTGTTTTAAAGTGGTCAAGACCAATAAGACCT ATTGTCTCAGCATTGCTGAAATATCTAATACTCTCTTCG GAGAATTCAGAATCGTCCCGTACTAGTTGAGATCCTCA AAGATGACGGGGTTAGAGAAGCCAGGTCTGGCTAGTTG AGTCAACTATGAAAGAGTTGGAAAGATGGCATTGTATC ACCTATCTTCTGCGACATCAAGAATCAAACCGAATGCC GGCGCGTGCTCGAATTCCATGTCGCCAGTTGACCACAA TCAGCCAGTGCTCATGCGATCAGATTAAGCCTTGTCAAT AGTCTCTTGATTAAGAAAAAATGTAAGTGGCAATGAGA TACAAGGCAAAACAGCTCACGGTAAATAATACGGGTAG GACATGGCGAGCTCCGGTCCTGAAAGGGCAGAGCATCA GATTATCCTACCAGAGTCACACCTGTCTTCACCATTGGT CAAGCACAACTACTCTATTATTGGAAATTA ACTGGGC TACCGCTTCCTGATGAATGTGACTTCGACCACCTCATTC TCAGCCGACAATGGAAAAAATACTTGAATCGGCCTCT CCTGATACTGAGAGAATGATAAAACTCGGAAGGGCAGT ACACCAA ACTCTTAACCACAATTCCAGAATAACCGGAG TACTCCACCCCAGGTGTTTAGAAGAACTGGCTAATATTG

AGGTCCTCCGCTTCAACCAAAATTTTCGGAAAGATTGAG
AAGAAGATCCAAATTCACAACACGAGATATGGAGAACT
GTTCAACAAGGCTGTGTACGCATATAGAGAAGAACTGC
TGGGGTCATCTTGGTCTAACAATGTCCCCCGGTCAGAG
GAGTTCAGCAGCATTCGTACGGATCCGGCATTCTGGTTT
CACTCAAAATGGTCCACAGCCAAGTTTGCATGGCTCCA
TATAAAACAGATCCAGAGGCATCTGATTGTGGCAGCTA
GGACAAGGTCTGCGGCCAACAAATTGGTGATGCTAACC
CATAAGGTAGGCCAAGTCTTTGTCACTCCTGAACTTGTT
GTTGTGACGCATACGAATGAGAACAAGTTCACATGTCT
TACCCAGGAACTTGTATTGATGTATGCAGATATGATGG
AGGGCAGAGATATGGTCAACATAATATCAACCACGGCG
GTGCATCTCAGAAGCTTATCAGAGAAAATTGATGACAT
TTTGCGGTTAATAGACGCTCTGGCAAAGACTTGGGTA
ATCAAGTCTACGATGTTGTATCACTAATGGAGGGATTTG
CATACGGAGCTGTCCAGCTACTCGAGCCGTCAGGTACA
TTTGCGGGAGATTTCTTCGCATTCAACCTGCAGGAGCTT
AAAGACATTCTAATTGGCCTCCTCCCCAATGATATAGCA
GAATCCGTGACTCATGCAATCGCTACTGTATTCTCTGGT
TTAGAACAGAATCAAGCAGCTGAGATGTTGTGCCTGTT
GCGTCTGTGGGGTCACCCACTGCTTGAGTCCCGTATTGC
AGCAAAGGCAGTCAGGAGCCAAATGTGCGCACCGAAA
ATGGTAGACTTTGATATGATCCTTCAGGTACTGTCTTTC
TTCAAGGGAACAATCATCAACGGATACAGAAAGAAGA
ATGCAGGTGTGTGGCCGCGAGTCAAAGTGGATACAATA
TATGGGAAGGTCATTGGGCAACTACATGCAGATTCAGC
AGAGATTTACACGATATCATGTTGAGAGAGTATAAGA
GTTTATCTGCACTTGAATTTGAGCCATGTATAGAATACG
ACCCTGTCTACTAACCTGAGCATGTTCCCTAAAAGACAAG
GCAATCGCACACCCCCAACGATAATTGGCTTGCCTCGTTT
AGGCGGAACCTTCTCTCCGAAGACCAGAAGAAACATGT
AAAGGAAGCGACTTCGACTAACCGCCTCTTGATAGAGT
TTTTAGAGTCAAATGATTTTGATCCATATAAAGAGATGG
AATATCTGACGACCCTTGAGTACCTTAGAGATGACAAT
GTGGCAGTATCATACTCGCTCAAAGAGAAGGAAGTGAA
AGTTAATGGACGGATCTTCGCTAAGCTGACAAAGAAGT
TAAGGAACTGTCAGGTGATGGCGGAAGGGATCCTAGCC
GATCAGATTGCACCTTTCTTTCAGGGGAAATGGAGTCATT
CAGGATAGCATATCCTTGACCAAGAGTATGCTAGCGAT
GAGTCAACTGTCTTTTAACAGCAATAAGAAACGTATCA
CTGACTGTAAAGAAAGAGTATGTTCAAACCGCAATCAT
GATCCGAAAAGCAAGAACCGTCGGAGAGTTGCAACCTT
CATAACAACCTGACCTGCAAAGTACTGTCTTAATTGGA
GATATCAGACGATCAAATTGTTTCGCTCATGCCATCAATC
AGTTGATGGGCCTACCTCATTTCCTTCGAGTGGATTCACC
TAAGACTGATGGACACTACGATGTTTCGTAGGAGACCCT
TTCAATCCTCCAAGTGACCCTACTGACTGTGACCTCTCA
AGAGTCCCTAATGATGACATATATATTGTCAGTGCCAG
AGGGGGTATCGAAGGATTATGCCAGAAGCTATGGACAA
TGATCTCAATTGCTGCAATCCAACCTTGCTGCAGCTAGAT

CGCATTTGTTATGGTACAGGGTGATAATC
AAGTAATAGCAGTAACGAGAGAGGTAAGATCAGATGA
CTCTCCGGAGATGGTGTGACACAGTTGCATCAAGCCA
GTGATAATTTCTTCAAGGAATTAATCCATGTCAATCATT
TGATTGGCCATAATTTGAAGGATCGTGAAACCATCAGG
TCAGACACATTCTTCATATACAGCAAACGAATCTTCAA
AGATGGAGCAATCCTCAGTCAAGTCCTCAAAAATTCAT
CTAAATTAGTGCTAGTGTCAGGTGATCTCAGTGAAAAC
ACCGTAATGTCCTGTGCCAACATTGCCTCTACTGTAGCA
CGGCTATGCGAGAACGGGCTTCCCAAAGACTTCTGTTA
CTATTTAAACTATATAATGAGTTGTGTGCAGACATACTT
TGACTCTGAGTTCTCCATCACCAACAATTCGCACCCCGA
TCTTAATCAGTCGTGGATTGAGGACATCTCTTTTGTGCA
CTCATATGTTCTGACTCCTGCCCAATTAGGGGGACTGAG
TAACCTTCAATACTCAAGGCTCTACACTAGAAATATCG
GTGACCCGGGGACTACTGCTTTTGCAGAGATCAAGCGA
CTAGAAGCAGTGGGACTACTGAGTCCTAACATTAGGAC
TAATATCTTAACTAGGCCGCCTGGGAATGGAGATTGGG
CCAGTCTGTGCAACGACCCATACTCTTTC AATTTTGAGA
CTGTTGCAAGCCCAAACATTGTTCTTAAGAAACATACG
CAAAGAGTCCTATTTGAAACTTGTTCAAATCCCTTATTG
TCTGGAGTGCACACAGAGGATAATGAGGCAGAAGAGA
AGGCATTGGCTGAATTCTTGCTTAATCAAGAGGTGATTC
ATCCCCGCGTTGCGCATGCCATCATGGAGGCAAGCTCT
GTAGGTAGGAGAAAGCAAATTCAAGGGCTTGTTGACAC
AACAAACACTGTAATTAAGATTGCGCTTACTAGGAGGC
CATTAGGCATCAAGAGGCTGATGCGGATAGTCAATTAT
TCTAGCATGCATGCAATGCTGTTTAGAGACGATGTTTTT
TCCTCTAGTAGATCCAACCACCCCTTAGTCTCTTCTAAT
ATGTGTTCTCTGACACTGGCAGACTATGCACGGAATAG
AAGCTGGTCACCTTTGACGGGAGGCAGGAAAATACTGG
GTGTATCTAATCCTGATACGATAGAACTCGTAGAGGGT
GAGATTCTTAGTGTAAGCGGAGGGTGTACAAGATGTGA
CAGCGGAGATGAACAATTTACTTGGTTCCATCTTCCAAG
CAATATAGAATTGACCGATGACACCAGCAAGAATCCTC
CGATGAGGGTACCATATCTCGGGTCAAAGACACAGGAG
AGGAGAGCTGCCTCACTTGCGAAAATAGCTCATATGTC
GCCACATGTGAAGGCTGCCCTAAGGGCATCATCCGTGT
TGATCTGGGCTTATGGGGATAATGAAGTAAATTGGACT
GCTGCTCTTACGATTGCAAAATCTCGGTGTAATGTAAAC
TTAGAGTATCTTCGGTTACTGTCCCCTTTACCCACGGCT
GGGAATCTTCAACATAGACTAGATGATGGTATAACTCA
GATGACATTCACCCCTGCATCTCTCTACAGGGTGTCACC
TTACATTACATATCCAATGATTCTCAAAGGCTGTTTAC
TGAAGAAGGAGTCAAAGAGGGGAATGTGGTTTACCAAC
AGATCATGCTCTTGGGTTTATCTCTAATCGAATCGATCT
TTCCAATGACAACAACCAGAACATATGATGAGATCACA
CTGCACCTACATAGTAAATTTAGTTGCTGTATCAGGGAA
GCACCTGTTGCGGTTCTTTTCGAGCTACTTGGGGTGGCA
CCGGA ACTGAGGACAGTGACCTCAAATAAGTTTATGTA

TGATCTCTAGCTATCTTCAAGAGTTATGAGCTTAATCTGG
TTGACTTAGCTATCTTCAAGAGTTATGAGCTTAATCTGG
AGTCATATCCCACGATAGAGCTAATGAACATTCTTTCAA
TATCCAGCGGGAAGTTGATTGGCCAGTCTGTGGTTTCTT
ATGATGAAGATACCTCCATAAAGAATGATGCCATAATA
GTGTATGACAATACCCGAAATTGGATCAGTGAAGCTCA
GAATTCAGATGTGGTCCGCCTATTTGAATATGCAGCACT
TGAAGTGCTCCTCGACTGTTCTTACCAACTCTATTACCT
GAGAGTAAGAGACCTAGACAATATTGTCTTATATATGG
GTGATTTATACAAGAATATGCCAGGAATTCTACTTTCCA
ACATTGCAGCTACAATATCTCATCCTGTCATTCAATCAA
GGTTACATGCAGTGGGCCTGGTCAACCATGACGGATCA
CACCAACTTGCAGATACGGATTTTATCGAAATGTCTGCA
AAACTGTTAGTATCTTGCACCCGACGTGTGATCTCCGGC
TTATATTCAGGAAATAAGTATGATCTGCTGTTCCCATCT
GTCTTAGATGATAACCTGAATGAGAAGATGCTTCAGCT
GATATCCCGGTTATGCTGTCTGTACACGGTACTCTTTGC
TACAACAAGAGAAATCCCGAAAATAAGAGGCTTAACTG
CAGAAGAGAAATGTTCAATACTCACTGAGTATTTACTG
TCGGATGCTGTGAAACCATTACTTAGCCCCGATCAAGT
GAGCTCTATCATGTCTCCTAACATAATTACATTCCCAGC
TAATCTGTACTACATGTCTCGGAAGAGCCTCAATTTGAT
CAGGGAAAGGGAGGACAGGGATACTATCCTGGCGTTGT
TGTTCCCCCAAGAGCCATTATTAGAGTTCCCTTCTGTGC
AAGATATTGGTGCTCGAGTGAAAGATCCATTCACCCGA
CAACCTGCGGCATTTTTTGCAAGAGTTAGATTTGAGTGCT
CCAGCAAGGTATGACGCATTCACACTTAGTCAGATTCA
TCCTGAACTCACATCTCCAAATCCGGAGGAAGACTACT
TAGTACGATACTTGTTTACAGAGGGATAGGGACTGCATCT
TCCTCTTGGTATAAGGCATCCCATCTCCTTTCTGTACCC
GAGGTAAGATGTGCAAGACACGGGAACTCCTTATACTT
GGCTGAAGGAAGCGGAGCCATCATGAGTCTTCTTGAAC
TGCATGTACCACATGAAACTATCTATTACAATACGCTCT
TTTCAAATGAGATGAACCCCCCGCAACGACATTTCTGGG
CCGACCCCAACTCAGTTTTTTGAATTCGGTTGTTTATAGG
AATCTACAGGCGGAGGTAACATGCAAGGATGGATTTGT
CCAAGAGTTCCGTCCATTATGGAGAGAAAATACAGAGG
AAAGTGACCTGACCTCAGATAAAGCAGTGGGGTATATT
ACATCTGCAGTACCCTACAGATCTGTATCATTGCTGCAT
TGTGACATTGAAATTCCTCCAGGGTCCAATCAAAGCTTA
CTAGATCAACTAGCTATCAATTTATCTCTGATTGCCATG
CATTCTGTAAGGGAGGGCGGGGTAGTAATCATCAAAGT
GTTGTATGCAATGGGATACTACTTTCATCTACTCATGAA
CTTGTTTGCTCCGTGTTCCACAAAAGGATATATTCTCTC
TAATGGTTATGCATGTGAGGGGATATGGAGTGTTACC
TGGTATTTGTCATGGGTTACCTGGGCGGGCCTACATTTG
TACATGAGGTGGTGAGGATGGCAAAAACCTCTGGTGCAG
CGGCACGGTACGCTTTTGTCTAAATCAGATGAGATCAC
ACTGACCAGGTTATTCACCTCACAGCGGCAGCGTGTGA
CAGACATCCTATCCAGTCCTTTACCAAGATTAATAAAGT

ACTTGAGGAAGAATATTGACACTGCGCTGATTGAAGCC
 GGGGGACAGCCCGTCCGTCCATTCTGTGCGGAGAGTCT
 GGTGAGCACGCTAGCGAACATAACTCAGATAACCCAGA
 TCATCGCTAGTCACATTGACACAGTCATCCGGTCTGTGA
 TATATATGGAAGCTGAGGGTGATCTCGCTGACACAGTA
 TTTCTATTTACCCCTTACAATCTCTCTACTGACGGGAAA
 AAGAGGACATCACTTAAACAGTGCACGAGACAGATCCT
 AGAGGTTACAATACTAGGTCTTAGAGTCGAAAATCTCA
 ATAAAATAGGCGATATAATCAGCCTAGTGCTTAAAGGC
 ATGATCTCCATGGAGGACCTTATCCCCTAAGGACATA
 CTTGAAGCATAGTACCTGCCCTAAATATTTGAAGGCTGT
 CCTAGGTATTACCAAACCTCAAAGAAATGTTTACAGACA
 CTTCTGTACTGTACTTGACTCGTGCTCAACAAAAATTCT
 ACATGAAAACCTATAGGCAATGCAGTCAAAGGATATTAC
 AGTAACTGTGACTCCTAACGAAAATCACATATTAATAG
 GCTCCTTTTTTTGGCCAATTGTATTCTTGTTGATTTAATTA
 TATTATGTTAGAAAAAAGTTGAACTCTGACTCCTTAGGA
 CTCGAATTCGAACTCAAATAAATGTCTTTAAAAAAGGT
 TGCGCACAAATTATTCTTGAGTGTAGTCTCGTCATTACACC AAATCTTTGTTTGGT

6. EXAMPLES

6.1 Example 1: Viral Vectored RSV Vaccine Induces Long-Lived Humoral Immunity in Cotton Rats

(193) This example demonstrates that vaccination with a recombinant Newcastle disease virus-vectored vaccine that expresses the F glycoprotein of RSV (rNDV-F) protects cotton rats from RSV challenge and induces long-lived neutralizing antibody production, even in RSV immune animals. In addition, this example demonstrates that the pulmonary eosinophilia induced by RSV infection of unvaccinated cotton rats is prevented by rNDV-F vaccination. Overall, these example demonstrates enhanced protective immunity to RSV F when this protein is presented in the context of an abortive NDV infection.

(194) 6.1.1 Materials & Methods

(195) 6.1.1.1 Animals

(196) 6-10 week old female cotton rats (*Sigmodon hispidus*) were used in this study (Sigmovir Biosystems, Inc., Bethesda, MD, USA). Animals were housed in groups of 3 in a BSL 2 facility and were offered a commercial pelleted rat chow and water ad libitum. Animals were acclimated for 7 to 10 days prior to vaccination or viral challenge. All procedures used here were conducted humanely. Data were collected from 3-5 cotton rats per cohort per time point, unless otherwise specified. All cotton rat studies were approved by the Institutional Animal Care and Use Committees of New York University School of Medicine (protocol 100504-02) and Rutgers-New Jersey Medical School.

(197) 6.1.1.2 Generation rNDV-F Construct

(198) rNDV-F was constructed as previously described [6]. Briefly, the sequence coding for the F protein of RSV (strain Long; SEQ ID NO:1) was inserted in the cDNA of the full length genome of NDV strain Hitchner B1 (a non-virulent vaccine strain of NDV), between the P and M genes. The inserted sequence contained NDV gene end and gene start sequences to make it a functional transcription unit, and a Kozak sequence for efficient translation. The cDNA sequence of the full length genome of NDV strain Hitchner B1 with the nucleic acid sequence encoding the RSV F protein inserted.

(199) 6.1.1.3 Vaccination

(200) rNDV-F was constructed as previously described [6] then amplified in 10-day-old embryonated chicken eggs. While under isoflurane anesthesia, cotton rats were intranasally (i.n.)

vaccinated with $1 \times 10^{6.6}$ pfu rNDV-F, $1 \times 10^{6.6}$ pfu NDV vector alone, or allantoic fluid as a mock control in a 100 μ l total volume divided equally between nares. Twenty-eight days after the priming vaccination, cotton rats were boosted in the same manner.

(201) 6.1.1.4 RSV Challenge

(202) Human RSV strain A2, originally obtained from ATCC (VR-1540), was passaged on murine STAT1^{sup.-/-} fibroblast monolayers as previously described [12]. The challenge dose of $1 \times 10^{6.6}$ pfu RSV was instilled into the nasal cavities of cotton rats under isoflurane anesthesia in a 50 μ l total volume, divided equally between nares. The inoculum was delivered slowly, over approximately 15 seconds, to restrict initial virus infection to the upper airway.

(203) 6.1.1.5 Plaque Assay

(204) Cotton rats were euthanized by CO₂ asphyxiation, and nasal wash fluids and lungs were collected immediately after euthanasia and stored at -80° C. until use in viral plaque assay. Viral plaque assay was performed on murine STAT1^{sup.-/-} fibroblast monolayers as previously described [13].

(205) 6.1.1.6 Histology and Immunohistochemistry

(206) Nasal cavities and lung were collected immediately following CO₂ asphyxiation. Nasal cavities were fixed in neutral buffered formalin then decalcified with 0.35M EDTA in 0.1M Tris (pH 6.95). The lung and decalcified nasal cavities were processed routinely, paraffin-embedded and sectioned at 5 μ m. Tissue sections were stained with hematoxylin and eosin (H&E) or were left unstained for immunohistochemistry (IHC). For IHC, tissue sections were incubated with goat polyclonal RSV antiserum (Biodesign, Saco, Maine, USA) diluted 1:500 followed by incubation with biotinylated rabbit anti-goat IgG antibody then HRP (ScyTek, Logan, UT, USA). Virus detection was accomplished with the streptavidin link and AEC chromagen (Scytek, Logan, UT, USA). IHC-labeled tissue sections were counterstained with hematoxylin.

(207) 6.1.1.7 RSV F-Specific ELISA

(208) Purified RSV F glycoprotein, whose transmembrane anchor was replaced with a 6His tag (SEQ ID NO:61), was produced by transfection of HEK293F cells and purified as described [14]. For RSV F-specific ELISA, Nunc maxisorb immunoplates (ThermoScientific, Waltham, MA, USA) were coated overnight at 4° C. with 0.3 μ g purified F protein diluted in 0.1M sodium carbonate buffer. Plates were washed with PBS containing 0.05% Tween 20 (PBS-T) then blocked with PBS containing 10% fetal calf serum (PBS-F). Plates were washed with PBS-T then samples, serially diluted in PBS-F, were added followed by incubation at 25° C. for 60 minutes. Wells were washed with PBS-T and secondary antibody diluted in PBS-F was added (chicken anti-cotton rat IgG, Immunology Consultants Laboratory, Inc., Portland, OR, USA). Plates were incubated at 25° C. for 60 minutes. After washing with PBS-T, tertiary antibody diluted in PBS-F was added (goat anti-chicken IgG conjugated to HRP, Southern Biotech, Birmingham, AL, USA). Following a second incubation at 25° C. for 60 minutes, plates were again washed with PBS-T followed by the addition of tetramethylbenzidine peroxidase substrate (ScyTek, Logan, UT, USA). After incubation at 25° C. for 30 minutes, the reaction was stopped by addition of 2N H₂SO₄. Absorbance at 450 nm (A₄₅₀) was determined using an Epoch Microplate Spectrophotometer (BioTek, Winooki, VT, USA). Titers were defined as the highest dilution of the sample resulting in A₄₅₀ four-fold greater than background (PBS-F alone).

(209) 6.1.1.8 Serum Neutralization Assay

(210) Two different methods were used for detection of RSV neutralizing antibody. Data shown in FIG. 4D were collected using a standard plaque reduction assay. Serum dilutions pre-incubated with 200 pfu RSV were used as the inoculum for plaque assay on murine STAT1^{sup.-/-} fibroblast monolayers as previously described [13].

(211) For a more quantitative approach to determining neutralizing antibody titer, a flow cytometric assay was devised. For data shown in FIG. 7A, serum samples at a 1/800 dilution were incubated with rg-RSV [8] at a concentration of $3 \times 10^{5.5}$ pfu/ml. in DMEM+5% fetal calf serum for 1 hour

at 37° C. The serum-virus mixture was then used to inoculate, in duplicate, monolayers of Vero cells (10^{sup}.5 cells plated in 2 cm^{sup}.2 wells). After a 20 hour incubation, cells were resuspended and analyzed by flow cytometry. Only infected cells will express gfp. Percent neutralization was calculated as $100 (1 - f_{sub.s}/f_{sub.0})$ where $f_{sub.s}$ =fraction of gfp⁺ cells in the sample, and $f_{sub.0}$ =the fraction of gfp⁺ cells in the no-serum control.

(212) 6.1.1.9 Enumeration of RSV F-Specific B Cells

(213) For T cell depletion, mouse anti-cotton rat CD8 and mouse anti-cotton rat CD4 monoclonal antibodies (r&d systems) were labeled with DSB-X according to the manufacturer's protocol (Dynabeads FlowComp Flexi, Invitrogen Life Technologies, Grand Island, NY, USA). Single cell suspensions of cervical lymph nodes, pooled by cohort, were incubated with 0.5 µg each of CD8 and CD4 DSB-X labeled antibody per sample, then T cells were depleted according to the manufacturer's protocol (Dynabeads FlowComp Flexi, Invitrogen Life Technologies, Grand Island, NY, USA). Cervical lymph node single cell suspensions, depleted of T cells and pooled by cohort, were used in ELISpot assays.

(214) For F-specific ELISpot, multiscreen HTS HA plates (Millipore, Jaffrey, NH, USA) were coated overnight at 4° C. with 0.3 µg purified F protein diluted in 0.1M sodium carbonate buffer. Plates were washed with RPMI supplemented with 10% fetal calf serum (cRPMI) and then were blocked with cRPMI for 1 hr at 37° C. in a 5% CO₂ incubator. Plates were washed with RPMI then cervical lymph node single cell suspensions were added in duplicate. Plates were incubated 4-6 hrs at 37° C. in 5% CO₂. Plates were then washed with PBS containing 0.05% Tween-20 (PBS-T). Secondary antibody diluted in PBS-T (chicken anti-cotton rat IgG, Immunology Consultants Laboratory, Inc., Portland, OR, USA) was added and plates were incubated overnight at 4° C. Plates were washed with PBS-T. Tertiary antibody diluted in PBS-T was added (goat anti-chicken AP, Southern Biotech, Birmingham, AL, USA) and plates were incubated at 25° C. for 60 minutes. Plates were again washed with PBS-T. NBT/BCIP substrate (Millipore, Jaffrey, NH, USA) was added and plates were incubated in the dark for 30 minutes at 25° C. Plates were washed with tap water. Spots were counted visually.

(215) 6.1.1.10 BAL Fluid Analysis

(216) BAL fluids were collected from cotton rats immediately after CO₂ asphyxiation by washing the lung with 2 ml of sterile saline. BAL fluid cell counts were determined by hemocytometer and BAL differentials were determined on Wright-Giemsa stained CytoSpin preparations (ThermoScientific, Waltham, MA, USA). For the 1 month time point, BAL fluid data was available for only 2 NDV vaccinated animals as BAL fluid collected from one NDV vaccinated animals was of poor quality and thus excluded. For the 2 month time point, BAL fluid data was available for only 2 mock vaccinated animals because one of the mock vaccinated animals died prior to sampling. For the 4 days after secondary RSV infection time point, BAL fluid data was available for only 2 mock vaccinated animals as necropsy of a third mock-vaccinated animal revealed pyometra, an abdominal mass and fibrinous pericarditis. BAL fluid analysis was not performed on that animal.

(217) 6.1.1.11 Statistical Analysis

(218) SigmaPlot 12.0 (Systat Software, Inc., San Jose, CA, USA) and GraphPad (GraphPad Software, Inc., La Jolla, CA USA) were used to perform ANOVA and t-test where appropriate. A p-value<0.05 was considered statistically significant.

(219) 6.1.2 Results

(220) 6.1.2.1 rNDV-F Vaccination Induces a Mild Inflammatory Response

(221) The immunization regimen involved intranasal delivery of 10^{sup}.6 pfu of rNDV-F into the nasal cavity of cotton rats. To determine the effect of inoculation, the nasal cavity and lung tissues were examined following each of the two i.n. vaccine doses. Histologic sections from the upper airway taken 24 hours, 48 hours, 72 hours and 7 days following intranasal instillation of 10^{sup}.6 pfu of rNDV-F showed minimal signs of inflammation, indistinguishable from similar sections

taken from untreated cotton rats (Data not shown).

(222) The nasal cavities of inoculated animals were also examined 28 days following priming, and 28 days following boost, for any evidence of disease secondary to immunization alone. No histologic changes were observed at this time point following delivery of allantoic fluid (FIGS. 3A and 3B) or the NDV vector alone (FIGS. 3E and 3F), but nasal cavity tissues collected from rNDV-F vaccinated animals showed submucosal lymphoplasmacytic infiltrates (FIGS. 1C and 1D), indicative of an acquired immune response. A more pronounced infiltrate was observed following rNDV-F boosting (FIGS. 1G and 1H).

(223) No histologic changes were observed in the lung following delivery of allantoic fluid or a priming dose of NDV vector alone (FIGS. 1I, 1L, and 1J). However, in response to rNDV-F priming, submucosal lymphoplasmacytic infiltrates similar to those observed in the nasal cavity mucosa, were observed surrounding bronchioles (FIG. 1K), with an increased number of lymphocytes and plasma cells surrounding bronchioles following rNDV-F boosting (FIG. 1N). Similar peribronchiolar infiltrates were observed in response to a boosting dose of NDV vector alone (FIG. 1M), but these were less cellular than those that accompanied rNDV-F vaccination. Additionally, following a second dose with NDV vector alone, or either dose of rNDV-F, rare eosinophils were observed intermixed with the lymphocytic and plasma cell infiltrates surrounding bronchioles (FIGS. 1K, 1M and 1N). Based on these studies it appears that intranasal immunization with rNDV-F causes no acute injury to the upper or lower airway, but does induce a mucosal lymphocytic response. The basis for the enhanced, though still mild, inflammatory response to rNDV-F when compared with the NDV vector alone is not clear, but is consistent with the ability to boost immunity with a second, boosting vaccine dose.

(224) 6.1.2.2 rNDV-F Vaccination Protects Against RSV Infection and Reinfection

(225) Tissues were harvested 4 days after RSV challenge of animals that had been immunized with two i.n. doses of allantoic fluid, NDV vector alone, or rNDV-F. In the nasal cavity and lung tissue of animals receiving allantoic fluid or NDV vector alone prior to RSV challenge, primary RSV infection was associated with mucosal and submucosal inflammatory infiltrates composed primarily of lymphocytes (FIGS. 2A and 2B). The lungs from these mock-immunized cotton rats had sparse peribronchiolar lymphohistiocytic infiltrates, with mucus and inflammatory cells in small airways (FIGS. 2D and 2E). In contrast, both lung and nasal cavity tissues collected from RSV-challenged, rNDV-F vaccinated cotton rats exhibited changes similar to those observed in response to vaccination alone (FIGS. 2C and 2F), with a more pronounced lymphocytic infiltrate, but lacking the mucus production and mixed inflammatory infiltrate.

(226) Viral load following challenge was assayed by immunohistochemistry (IHC) and by plaque assay. IHC staining with a polyclonal antiviral serum revealed diffuse RSV antigen staining in the nasal cavities and small foci of infected cells in the lower airways of mock and NDV vaccinated animals. In contrast, only rare RSV antigen positive cells were detected in the nasal cavities of the rNDV-F vaccinated cotton rats, and none in the lower airways of the immunized animals (FIGS. 3A-3F). In support of IHC findings, viable virions were not detected by plaque assay of nasal mucosa and lung collected from animals that had received rNDV-F vaccination prior to RSV challenge (FIG. 3G).

(227) Following a second RSV challenge delivered 6 months after the first RSV challenge of mock-vaccinated or rNDV-F vaccinated animals, lung and nasal tissues were collected. IHC of nasal cavity and lung collected 4 days after secondary RSV challenge revealed rare, scattered antigen-positive cells in the nasal mucosa of cohorts primed with allantoic fluid or NDV vector only. No RSV antigen was detected in the lungs or nasal cavities of rNDV-F vaccinated animals. Plaque assay of nasal cavity mucosa and lung tissues collected after secondary RSV challenge was negative in all animals (not shown).

(228) 6.1.2.3 rNDV-F Vaccination Induces a Long-Lived Mucosal Antibody Response

(229) Serum, bronchoalveolar lavage (BAL) fluid and nasal wash (NW) fluid were collected at

multiple time points after vaccination and RSV challenge for determination of RSV F-specific IgG levels by ELISA (FIG. 4). rNDV-F vaccination induced robust, F-specific IgG responses that were boosted following RSV challenge and then maintained at high levels. While serum antibody levels were similar amongst cohorts 1-2 months after RSV challenge, mucosal F-specific antibody responses remained significantly elevated in BAL and NW fluid samples collected from rNDV-F vaccinated animals compared to controls up to 6 months after RSV infection, the latest time point examined. Serum neutralization, assayed by plaque reduction, demonstrated that antibodies produced in response to rNDV-F vaccination were neutralizing (FIG. 4D).

(230) Cervical lymph nodes were collected 4 days or 5 months after RSV challenge of mock and rNDV-F primed and boosted animals, and RSV F-specific IgG antibody secreting cells (ASCs) were enumerated by ELISpot (FIG. 4E). While a response was not detected, or was minimal, by cells harvested from cervical lymph nodes of mock or NDV vaccinated animals, robust F-specific ASC responses were observed in cervical lymph nodes collected from rNDV-F vaccinated animals at both the 4 day and 5 month time points. This is a significant finding given the short-lived mucosal protection that follows natural infection, and the importance of augmenting this protection by any RSV vaccine candidate [5]. Reagents to assay IgA in the cotton rat model are not yet available.

(231) 6.1.2.4 rNDV-F Vaccination Prevents Eosinophilic Lung Inflammation

(232) The cellularity of BAL fluid collected at multiple time points after vaccination and RSV challenge was determined by cell counts, and evaluated for cell type by examination of Wright-Giemsa stained cytospin preparations (FIGS. 5A and 5B). While vaccination alone did not increase total BAL fluid cellularity, a small, but not statistically significant, increase in total BAL fluid cellularity was detected in all cohorts after primary RSV challenge. After secondary RSV infection this increase in cellularity was again detected in BAL fluid collected from animals that had been vaccinated with allantoic fluid or NDV vector alone, but not present in BAL fluids collected from rNDV-F vaccinated animals.

(233) As previously described [7], eosinophils are present in the lungs of naïve cotton rats, in the absence of pulmonary pathology, and the composition of the BAL fluid was not altered by vaccination alone. Consistent with these prior observations, eosinophils were significantly increased one month after primary RSV challenge of control animals [7], but this increase did not occur in animals vaccinated with rNDV-F (FIG. 5). The number of pulmonary eosinophils decreased nearly to baseline 2 months after primary RSV infection, but rose again in response to secondary RSV challenge of control animals that had received only allantoic fluid or NDV vector. This increase in BAL eosinophils following secondary RSV infection, while striking, did not reach statistical significance ($p=0.06$). Thus, in the cotton rat model, rNDV-F vaccination confers long-lasting protection, but also inhibits eosinophilic inflammation in response to RSV infection and reinfection.

(234) 6.1.2.5 Immunization of RSV-Immune Animals

(235) Based on the vaccination strategy wherein the RSV F protein is expressed only in rNDV-F infected cells, it was hypothesized that it might be possible to immunize animals previously exposed to RSV. To test this possibility, 20 cotton rats were infected with RSV, and half of these were immunized with 10^{sup}.6 pfu of rNDV-F administered i.n. at 2 months, and again at 3 months, after primary RSV infection (FIG. 6). The 2 month time point for vaccination was chosen to ensure that RSV F protein specific antibodies would be present in serum and at the mucosal surface of the respiratory tree (see FIG. 4). Serum antibody titers were determined at 4 months, and again at 9 months after primary RSV infection as outlined in FIG. 6. All remaining animals were re-challenged with RSV at the 9 month time point, and no virus was detected in the lungs or noses of animals in either cohort at 2 or 4 days post virus instillation.

(236) RSV F-specific IgG levels in all animals were measured by an F-protein specific ELISA assay which demonstrated an approximately 3-fold increase in animals that had received rNDV-F

(1/2500 versus 1/7500), a level maintained for 6 months after the boosting dose of vaccine was administered (FIG. 7B). Neutralizing antibody was assessed by incubating serum samples with rg-RSV, a recombinant virus expressing eGFP [8], and determining % of Vero cell infection by flow cytometry. The result of this assay is shown in FIG. 7A, demonstrating an enhancement of neutralizing antibody in animals immunized with rNDV-F in the presence of RSV-neutralizing antibody, an enhancement which remained stable at the 9 month time point. This study suggests that, in addition to the ability of this vaccine to give long-lasting protection, rNDV-F is also efficacious in those with pre-existing antibody.

(237) 6.1.3 Discussion

(238) The data presented in this example demonstrates that a Newcastle disease virus vectored RSV vaccine, designed to stimulate robust IFN production simultaneous with RSV F protein presentation to the immune system, protects against RSV infection and induces a long-lived humoral immune response. Delivery of a protective dose of rNDV-F was not associated with adverse histopathology. Lymphoplasmacytic infiltrates observed in immunized animals in response to the boosting rNDV-F dose and subsequent challenge are consistent with an appropriate adaptive immune response (FIG. 1). Additionally, the increase in lung eosinophils detected in cotton rats that been mock vaccinated prior to primary and secondary RSV challenge was not detected in the lungs of rNDV-F vaccinated cotton rats. This finding is of particular interest when considered in light of mounting evidence that severe RSV lung disease during infancy is associated with asthma and recurrent wheezing later in childhood [1-4].

(239) As shown here, the eosinophilic response to RSV infection in the cotton rat was abrogated by rNDV-F vaccination. Without being bound by any theory, the adjuvant effect of the type I IFN induction, in addition to the prolongation of antibody responses by the rNDV-F vaccines described in this example, may also decrease T.sub.H2 responses to inhaled allergens encountered during RSV infections in young children.

(240) RSV-specific secretory and serum antibodies have been correlated with protection of the upper and lower airways, respectively, against re-infection with RSV [9, 10]. A recent study suggests that the ability of this virus to re-infect immunocompetent adults is correlated with a defect in B cell memory. Neutralizing serum and nasal antibody levels in experimentally infected healthy adult volunteers increased post-infection, but returned to baseline levels 6 months later [5]. This differs markedly from the lifelong persistence of influenza specific IgG that follows this infection [11]. Thus, it is essential that any RSV vaccine candidate induce both systemic and mucosal RSV-specific antibodies, but do so in a manner that induces a B cell memory response superior to that induced by RSV infection. It is this requirement that recommends consideration of a vectored RSV vaccine.

(241) In this example, rNDV-F vaccination induced serum and mucosal F-specific antibodies that were maintained for at least 7 months. Importantly, the mucosal F-specific antibody response, which correlates most closely with protection against reinfection [5], was more robust and longer lived in rNDV-F immunized cotton rats than that induced by RSV infection alone. The ability of NDV-F to boost neutralizing antibody titers in RSV seropositive animals indicates that vaccination of older children and adults may provide an avenue for protecting a population while avoiding the difficulties associated with vaccination of young infants.

6.1.4 REFERENCES CITED IN THIS EXAMPLE 6.1

(242) 1. Korppi M, Piippo-Savolainen E, Korhonen K, Remes S. Respiratory morbidity 20 years after RSV infection in infancy. *Pediatric pulmonology*. 2004; 38(2):155-60. Epub 2004/06/24. doi: 10.1002/ppul.20058. PubMed PMID: 15211700. 2. Pelaia G, Vatrella A, Gallelli L, Renda T, Cazzola M, Maselli R, et al. Respiratory infections and asthma. *Respiratory medicine*. 2006; 100(5):775-84. Epub 2005/11/18. doi: 10.1016/j.rmed.2005.08.025. PubMed PMID: 16289785. 3. Sigurs N, Bjarnason R, Sigurbergsson F, Kjellman B. Respiratory syncytial virus bronchiolitis in infancy is an important risk factor for asthma and allergy at age 7. *American journal of respiratory*

and critical care medicine. 2000; 161(5):1501-7. Epub 2000/05/12. PubMed PMID: 10806145. 4. Sigurs N, Gustafsson P M, Bjarnason R, Lundberg F, Schmidt S, Sigurbergsson F, et al. Severe respiratory syncytial virus bronchiolitis in infancy and asthma and allergy at age 13. American journal of respiratory and critical care medicine. 2005; 171(2):137-41. Epub 2004/11/02. doi: 10.1164/rccm.200406-7300C. PubMed PMID: 15516534. 5. Habibi M S, Jozwik A, Makris S, Dunning J, Paras A, DeVincenzo J P, et al. Impaired Antibody-mediated Protection and Defective IgA B-Cell Memory in Experimental Infection of Adults with Respiratory Syncytial Virus. American journal of respiratory and critical care medicine. 2015; 191(9):1040-9. doi: 10.1164/rccm.201412-22560C. PubMed PMID: 25730467; PubMed Central PMCID: PMC4435460. 6. Martinez-Sobrido L, Gitiban N, Fernandez-Sesma A, Cros J, Mertz S E, Jewell N A, et al. Protection against respiratory syncytial virus by a recombinant Newcastle disease virus vector. Journal of virology. 2006; 80(3):1130-9. Epub 2006/01/18. doi: 10.1128/JVI.80.3.1130-1139.2006. PubMed PMID: 16414990; PubMed Central PMCID: PMC1346968. 7. Grieves J L, Yin Z, Durbin R K, Durbin J E. Acute and Chronic Airway Disease After Human Respiratory Syncytial Virus Infection in Cotton Rats (*Sigmodon hispidus*). Comp Med. 2015; 65(4):315-26. PubMed PMID: 26310461; PubMed Central PMCID: PMC4549677. 8. Hallak L K, Collins P L, Knudson W, Peebles M E. Iduronic acid-containing glycosaminoglycans on target cells are required for efficient respiratory syncytial virus infection. Virology. 2000; 271(2):264-75. doi: 10.1006/viro.2000.0293. PubMed PMID: 10860881. 9. Shinoff J J, O'Brien K L, Thumar B, Shaw J B, Reid R, Hua W, et al. Young infants can develop protective levels of neutralizing antibody after infection with respiratory syncytial virus. The Journal of infectious diseases. 2008; 198(7):1007-15. Epub 2008/08/16. doi: 10.1086/591460. PubMed PMID: 18702606. 10. Walsh E E, Falsey A R. Humoral and mucosal immunity in protection from natural respiratory syncytial virus infection in adults. The Journal of infectious diseases. 2004; 190(2):373-8. Epub 2004/06/25. doi: 10.1086/421524. PubMed PMID: 15216475. 11. Fujimoto C, Takeda N, Matsunaga A, Sawada A, Tanaka T, Kimoto T, et al. Induction and maintenance of anti-influenza antigen-specific nasal secretory IgA levels and serum IgG levels after influenza infection in adults. Influenza Other Respir Viruses. 2012; 6(6):396-403. doi: 10.1111/j.1750-2659.2011.00330.x. PubMed PMID: 22226319; PubMed Central PMCID: PMC4941696. 12. Durbin J E, Hackenmiller R, Simon M C, Levy D E. Targeted disruption of the mouse Stat1 gene results in compromised innate immunity to viral disease. Cell. 1996; 84(3):443-50. Epub 1996/02/09. PubMed PMID: 8608598. 13. Gitiban N, Jurcisek J A, Harris R H, Mertz S E, Durbin R K, Bakaletz L O, et al. Chinchilla and murine models of upper respiratory tract infections with respiratory syncytial virus. Journal of virology. 2005; 79(10):6035-42. Epub 2005/04/29. doi: 10.1128/JVI.79.10.6035-6042.2005. PubMed PMID: 15857989; PubMed Central PMCID: PMC1091680. 14. Chaiwatpongsakorn S, Epand R F, Collins P L, Epand R M, Peebles M E. Soluble respiratory syncytial virus fusion protein in the fully cleaved, pretriggered state is triggered by exposure to low-molarity buffer. Journal of virology. 2011; 85(8):3968-77. doi: 10.1128/JVI.01813-10. PubMed PMID: 21307202; PubMed Central PMCID: PMC3126149.

6.2 Example 2: Optimized Recombinant Ndv Constructs

(243) This example demonstrates the production of optimized recombinant NDV constructs. In particular, the example demonstrates the enhanced expression of RSV F protein by the optimized recombinant NDV constructs.

(244) 6.2.1 Materials & Methods

(245) 6.2.1.1 Generation of Recombinant Newcastle Disease Virus (rNDV) Expressing the F Protein of Human Respiratory Syncytial Virus.

(246) The generation and characterization of the first version of rNDV-RSV-F has been described in detail in reference Martinez-Sobrido et al. (Journal of virology. 2006; 80(3):1130-9). This first version will be referred here as rNDV-F. Briefly, the sequence coding for the F protein of RSV (strain Long) was inserted in the cDNA of the full length genome of NDV strain Hitchner B1 (a

non-virulent vaccine strain of NDV), between the P and M genes. The inserted sequence contained NDV gene end and gene start sequences to make it a functional transcription unit, and a Kozak sequence for efficient translation.

(247) The plasmid obtained was used to rescue an infectious rNDV-F following a well-established protocol that has been recently described in Ayllon et al. (J Vis Exp. 80:50830 (2013)). The next versions of rNDV—RSV-F were prepared in a similar way, but using a full length cDNA of NDV vaccine strain LaSota and a codon optimized sequence of the RSV-F protein. The codon optimized sequence was purchased from Genewiz (see the website for Genewiz) and used to generate 2 different versions of rNDV-RSV-F that will be referred here as rNDV-Fopt and rNDV-F.sub.chim and are described in detail in the following paragraphs. NDV-Fopt was generated in a similar way to the original rNDV-F. The synthetic, codon optimized, sequence of RSV-F was used as template to PCR amplify an insert that contained the F open reading frame flanked by the necessary restriction sites, gene end and gene start regulatory sequences, and a Kozak sequence for efficient translation. The codon-optimized sequence of RSV-F sequence is shown in SEQ ID NO:2. The PCR product was directly cloned into the plasmid containing the full length NDV cDNA, between the P and M genes, and used to rescue an infectious rNDV-Fopt as previously described. The full length NDV cDNA sequence of the NDV strain LaSota with the codon-optimized RSV-F sequence inserted is shown in SEQ ID NO:3.

(248) rNDV-F.sub.chim was designed to express a chimeric F protein with the transmembrane and cytoplasmic domains of the RSV F protein replaced by the corresponding sequences of the NDV F protein. The rationale for this was that the chimeric F protein is predicted to be efficiently incorporated in the envelope of the viral particle and result in better immunogenicity. To obtain the rescue plasmid for rNDV-F.sub.chim we proceeded in two steps, summarized in FIG. 8. First, the codon optimized synthetic sequence was used as template to PCR amplify the ectodomain, flanked by the appropriate restriction sites and Kozak sequence. The PCR product was cloned into plasmid pShuttle LaSota that contained the required gene end and gene start sequences, as well as the sequence coding for the transmembrane and cytoplasmic domains of the NDV F protein. The fusion in frame of the RSV-F ectodomain and rNDV-F transmembrane and cytoplasmic domains resulted in the chimeric F protein. In the second step, after sequence confirmation, the insert in plasmid pShuttle LaSota was excised and cloned into the plasmid containing the full length NDV cDNA, between the P and M genes, and used to rescue the infectious rNDV-chimF. The chimeric RSV-F sequence is shown in SEQ ID NO:4. The full length NDV cDNA sequence of the NDV strain LaSota with the codon-optimized RSV-F sequence inserted is shown in SEQ ID NO:5. A schematic representation of the different rNDV-F, rNDV-F.sub.opt, rNDV-F.sub.chim are shown in FIG. 9.

(249) 6.2.1.2 Animals

(250) 6-10 week old female cotton rats (*Sigmodon hispidus*) were used in this study (Sigmovir Biosystems, Inc., Bethesda, MD, USA). Animals were housed in groups of 3 in a BSL 2 facility and were offered a commercial pelleted rat chow and water ad libitum. Animals were acclimated for 7 to 10 days prior to vaccination or viral challenge. Animal housing and experimental procedures were approved by the Institutional Animal Care and Use Committee at the New York University School of Medicine and the Rutgers—New Jersey Medical School. All procedures used here were conducted humanely. Data were collected from 3-5 cotton rats per cohort per time point, unless otherwise specified.

(251) 6.2.1.3 Vaccination

(252) NDV-F was constructed as previously described (33) then amplified in 10-day-old embryonated chicken eggs. While under isoflurane anesthesia, cotton rats were intranasally (i.n.) vaccinated with $1 \times 10^{5.6}$ pfu NDV-F, $1 \times 10^{5.6}$ pfu NDV vector alone, or allantoic fluid as a mock control in a 100 μ l total volume divided equally between nares. Twenty-eight days after the priming vaccination, cotton rats were boosted in the same manner.

(253) 6.2.1.4 RSV Challenge

(254) Human RSV strain A2, originally obtained from ATCC (VR-1540), was passaged on murine STAT1.sup.-/- fibroblast monolayers as previously described (54). The challenge dose of $1 \times 10^{6.6}$ pfu RSV was instilled into the nasal cavities of cotton rats under isoflurane anesthesia in a 50 μ l total volume, divided equally between nares. The inoculum was delivered slowly, over approximately 15 seconds, to restrict initial virus infection to the upper airway.

(255) 6.2.1.5 Plaque Assay

(256) Cotton rats were euthanized by CO.sub.2 asphyxiation, and lungs were collected immediately after euthanasia and stored at -80° C. until use in viral plaque assay. Viral plaque assay was performed on murine STAT1-/-fibroblast monolayers as previously described (Gitiban et al., Journal of Virology 2005; 79(10):6035-42).

(257) 6.2.1.6 Flow Cytometry Assay

(258) With respect to the data shown in FIG. 18: Vero cells growing in 6 well plates were infected with dilutions of the viral stocks by adding 250 μ l of virus diluted in PBS per well. After 1 h adsorption, 2 ml of complete medium (DMEM supplemented with 1% antibiotics and 10% fetal bovine serum) was added per well and cells were incubated at 37° C. in a CO2 incubator. At 24 h post infection cells were resuspended by trypsin treatment and fixed with 4% formaldehyde for 30 min. at room temperature. After fixation, approximately $10^{5.5}$ cells (1/10 of the sample) were labelled with RSV F specific humanized monoclonal antibody Synagis followed by fluorescently labelled anti-human secondary antibody. Fluorescently labelled cells were detected using a flow cytometer Canto II.

(259) With respect to the data shown in FIGS. 9A-9B: Serum samples at a 1/800 dilution were incubated with rg-RSV [37] at a concentration of $3 \times 10^{5.5}$ pfu/ml. in DMEM+5% fetal calf serum for 1 hour at 37° C. The serum-virus mixture was then used to inoculate, in duplicate, monolayers of Vero cells ($10^{5.5}$ cells plated in 2 cm² wells). After a 20 hour incubation, cells were resuspended and analyzed by flow cytometry. Only infected cells will express gfp. Percent neutralization was calculated as $100 (1 - f_s/f_0)$ where f_s =fraction of gfp+ cells in the sample, and f_0 =the fraction of gfp+ cells in the no-serum control.

(260) 6.2.1.7 Rt-PCR and Immunofluorescence Assay

(261) RT-PCR: Viral RNA was purified from the viral stocks using the kit E.Z.N.A. viral RNA kit (OMEGA bio-tek) and conditions recommended by the manufacturer. 5 μ l of purified RNA were used as template in a 50 μ l RT-PCR reaction using the kit Superscript III high-fidelity RT-PCR kit (Invitrogen, 12574-035) and the following primers specific for NDV genome sequences flanking the insert site. pNDV/3102+5'-CTGTCCACTCGGCATCACAC-3' (SEQ ID NO:62) and pNDV/3231-5'-CTAGATTAATTACGGTTACGC-3' (SEQ ID NO:63). The amplified band was purified from an agarose gel using the kit E.Z.N.A. gel extraction kit (OMEGA bio-tek) and sequenced using the same primers.

(262) Immunofluorescence Microscopy: Vero cells growing in 96 well plates were infected with dilutions of the viral stocks by adding 50 μ l of virus diluted in PBS per well. After 1 h adsorption, 200 μ l of complete medium (DMEM supplemented with 1% antibiotics and 10% fetal bovine serum) was added per well and cells were incubated at 37° C. in a CO2 incubator. At 24 h post infection cells were fixed with 4% formaldehyde for 30 min. at room temperature and then labelled with RSV F specific humanized monoclonal antibody Synagis and/or NDV specific mouse serum, followed by fluorescently labelled anti-human and/or anti-mouse secondary antibodies. Fluorescently labelled cells were observed and photographed in a fluorescent microscope.

(263) 6.2.1.8 RSV-F Specific ELISA

(264) Purified RSV F glycoprotein, whose transmembrane anchor was replaced with a 6His tag (SEQ ID NO:61), was produced by transfection of HEK293F cells and purified as described (56). For RSV F-specific ELISA, Nunc maxisorb immunoplates (ThermoScientific, Waltham, MA, USA) were coated overnight at 4° C. with 0.3 μ g purified F protein diluted in 0.1M sodium

carbonate buffer. Plates were washed with PBS containing 0.05% Tween 20 (PBS-T) then blocked with PBS containing 10% fetal calf serum (PBS-F). Plates were washed with PBS-T then samples, serially diluted in PBS-F, were added followed by incubation at 25° C. for 60 minutes. Wells were washed with PBS-T and secondary antibody diluted in PBS-F was added (chicken anti-cotton rat IgG, Immunology Consultants Laboratory, Inc., Portland, OR, USA). Plates were incubated at 25° C. for 60 minutes. After washing with PBS-T, tertiary antibody diluted in PBS-F was added (goat anti-chicken IgG conjugated to HRP, Southern Biotech, Birmingham, AL, USA). Following a second incubation at 25° C. for 60 minutes, plates were again washed with PBS-T followed by the addition of tetramethylbenzidine peroxidase substrate (ScyTek, Logan, UT, USA). After incubation at 25° C. for 30 minutes, the reaction was stopped by addition of 2N H.sub.2SO.sub.4. Absorbance at 450 nm (A.sub.450) was determined using an Epoch Microplate Spectrophotometer (BioTek, Winooki, VT, USA). Titers were defined as the highest dilution of the sample resulting in A.sub.450 four-fold greater than background (PBS-F alone).

(265) 6.2.1.9 RSV Neutralization Assay

(266) The potent RSV neutralizing monoclonal antibody Synagis® (concentration 100 mg/ml) was serially diluted in tissue culture medium (DMEM supplemented with 1% fetal bovine serum and 1% antibiotics). Dilutions tested were 1:1000, 1:5000, 1:25000, 1:125000, 1:625000, as indicated in FIGS. 14 and 15.

(267) Next, the antibody dilutions were distributed in 96 well plates, (50 µl per well), and the indicated viruses were added, in duplicate rows (see table below). Viral stocks were diluted in tissue culture medium to give an approximate titer of 10²-10³ infectious particles per 50 µl. Virus and antibody dilutions were mixed by pipetting up and down and incubated at 37° C. for 2 h. After the 2 h incubation, 100 µl of medium containing 10⁴ cells was added to each well and the plates were incubated at 37° C. for 24 h. For the RSV-GFP virus cells were the HEp2 cell line, for all the recombinant NDVs we used Vero cells.

(268) At 24 h post infection cells were fixed with 10% paraformaldehyde in PBS for 15 minutes at room temperature and then overlaid with 1×PBS.

(269) Cells infected with GFP expressing viruses (RSV-GFP and NDV-GFP) were visualized directly by fluorescence microscopy. For the recombinant NDV viruses, infected cells were visualized by immuno-fluorescence using a commercial monoclonal antibody that recognizes the NDV HN protein followed by a fluorescent anti-mouse secondary antibody. Representative microscopic fields were photographed to prepare FIGS. 14 and 15.

(270) TABLE-US-00009 TABLE 2 distribution of virus and antibody dilutions in the plate. Each virus was added to two replicate rows.

| | dilution | dilution | dilution | dilution | dilution |
|------------------|-------------------|-------------|----------------|-----------------|---------------------|
| 1:1000 | 1:5000 | 1:25000 | 1:125000 | 1:625000 | no antibody |
| RSV-GFP.sup.1 | NDV-GFP.sup.2 | NDV-F.sup.3 | NDV-Fopt.sup.4 | NDV-Fchim.sup.5 | NDV-LaSota wt.sup.6 |
| NDV-boFopt.sup.7 | NDV-boFchim.sup.8 | | | | |

.sup.1SV-GFP: recombinant RSV expressing the Green Fluorescent Protein, kindly provided by Dr. Megan Shaw (Mount Sinai). Positive control for neutralization. .sup.2NDV-GFP: recombinant NDV (strain LaSota) expressing the Green Fluorescent Protein. Negative control for neutralization. .sup.3NDV-F: original recombinant NDV (strain Hitchner B1) expressing the wild type version of the human RSV F protein. .sup.4NDV-Fopt: recombinant NDV (strain LaSota) expressing a codon optimized full length version of the human RSV F protein. .sup.5NDV-Fchim: recombinant NDV (strain LaSota) expressing a codon optimized chimeric version of the human RSV F protein ectodomain fused to the transmembrane and cytoplasmic domains of the NDV F protein. .sup.6NDV-LaSota wt: recombinant NDV with no insert. Negative control for neutralization. .sup.7NDV-boFopt: recombinant NDV (strain LaSota) expressing a codon optimized full length version of the bovine RSV F protein. .sup.8NDV-boFchim: recombinant NDV (strain LaSota) expressing a codon optimized chimeric version of the bovine RSV F protein ectodomain fused to the transmembrane and cytoplasmic domains of the NDV F protein.

6.2.2 Results

(271) Optimized versions of the rNDV-F construct have been produced as described in Section 6.2.1.1. These constructs are named rNDV-F.sub.opt and rNDV-F.sub.chim. FIG. 9 summarizes the main differences in the 3 recombinant NDVs expressing RSV-F. After rescue, all of the recombinant NDVs were amplified by inoculation in 8-10 old embryonated chicken eggs and characterized by RT-PCR and immunofluorescence (FIG. 13).

(272) While inflammation is a necessary adjuvant in any vaccination strategy, it is important to limit inflammation in the respiratory tree when the vaccine is administered by the intranasal route. Both primary immunization and challenge with the optimized constructs, rNDV-F.sub.opt and rNDV-F.sub.chim, showed a decreased acute inflammatory response when compared with the original rNDV-F construct. In addition, the optimized constructs, rNDV-F.sub.opt and rNDV-F.sub.chim, show enhanced RSV F protein expression (FIG. 10). Expression of the RSV F protein is enhanced in both NDV-F.sub.opt and NDV-F.sub.chim when compared with the original NDV-F construct. With being bound by any theory, it is possible that the relative reduction in RSV-F protein expression in the original NDV-F is related to the instability of the original rNDV-F construct. Initial observations indicate that the optimized constructs, NDV-F.sub.opt and NDV-F.sub.chim, appear to be much more stable than the original rNDV-F construct. The initial observations indicate that the original rNDV-F construct loses the RSV sequence upon passage.

(273) Cotton rats were mock treated or given priming and boosting doses of NDV, rNDV-F, rNDV-F.sub.opt or rNDV-F.sub.chim. Following RSV challenge, no RSV was detected in lung homogenates from animals receiving either rNDV-F.sub.opt or rNDV-F.sub.chim. Thus, the administration of rNDV-F.sub.opt and rNDV-F.sub.chim prior to challenge with RSV infection inhibited RSV replication. The administration of rNDV-Fopt to cotton rats resulted in a better protection against RSV challenge than administration rNDV-F to cotton rats.

(274) Another important measure of efficacy is the longevity of antibody protection. To determine whether rNDV-F.sub.opt and rNDV-F.sub.chim could also induce long-lived RSV-F specific antibody, animals were given allantoic fluid, rNDV vector, rNDV-F.sub.opt or rNDV-F.sub.chim, boosted one month later, and challenged with RSV on day 0. Serum samples were taken at day—31, and days 6, 13, 91 and 147 post-RSV challenge. The data in FIG. 12 demonstrates that the optimized constructs, unlike natural infection, can induce high titers of RSV F specific antibody that persist for 6 months after immunization. Titers are boosted by virus challenge, while protecting the host from infection, in the absence of disease.

(275) To assess the ability of Synagis®, a monoclonal antibody that binds to RSV F protein, to neutralize the recombination NDV constructs. As shown in FIG. 14, the recombinant RSV-GFP was neutralized by Synagis®, but NDV-GFP was not inhibited. Similarly, the recombinant NDV constructs, rNDV-F, rNDV-Fopt, and rNDV-Fchim as well as wild-type NDV were not neutralized by Synagis® (FIG. 15). For all the recombinant NDVs, no reduction in the percentage of infected cells at any of the dilutions tested was observed.

6.3 Example 3: Stability Assays for Recombinant NDV Constructs

(276) Stability of the recombinant NDVs expressing the F protein of RSV is evaluated by the ability to retain expression of the inserted gene after passage in eggs. The percentage of viruses retaining RSV-F expression is measured by simultaneous detection of the NDV protein HN and the RSV F protein by double immunofluorescence in cells infected with the different passages.

(277) Based on initial observations, the optimized versions (rNDV-F.sub.opt and rNDV-F.sub.chim) will retain a high percentage of cells with double labelling, while the original version (rNDV-F) will lose expression of the F protein in a high percentage of the infected cells after a few passages.

(278) Viruses rNDV-F, rNDV-F.sub.opt and rNDV-F.sub.chim are inoculated in triplicate at a 1:10000 dilution in chicken embryonated eggs (100 µl per egg). At 3 days post-infection the allantoic fluid is harvested, evaluated for hemagglutination and used to inoculate new eggs at 1:10000 dilution, as described above. This strategy is repeated for several passages.

(279) Serial dilutions of the harvested allantoic fluids is used to infect Vero cells in 96 well plates.

At 24 hours post infection cells are fixed and co-stained with a commercial mouse monoclonal antibody that recognizes the NDV HN protein and a humanized monoclonal antibody that recognize the RSV-F protein. Single and doubly labelled cells are quantified using a fluorescence plate reader (Celigo).

6.4 Example 4: Recombinant NDV Encoding Bovine RSV F Protein

(280) 6.4.1 Materials & Methods

(281) 6.4.1.1 Generation of Recombinant Newcastle Disease Virus (rNDV) Expressing the F Protein of Human Respiratory Syncytial Virus.

(282) 6.4.1.2 RSV Neutralization Assay

(283) This assay was performed as described in Section 6.2.1.9.

(284) 6.4.1.3 Generation of the Recombinant NDVs Expressing Bovine RSV F Proteins.

(285) The generation of the recombinant NDVs (LaSota strain) expressing a full length codon optimized F protein from bovine RSV (strain ATCC51908) (see SEQ ID NO:11 for the codon optimized nucleic acid sequence encoding bovine RSV F protein) and a chimeric F protein with the codon optimized ectodomain from bovine RSV (strain ATCC51908) fused to the transmembrane and cytoplasmic domains of NDV (strain LaSota) (see SEQ ID NO:14 for the nucleic acid sequence encoding chimeric F protein) was performed as described for the construction of the recombinant NDVs expressing the full length and chimeric F proteins from human RSV, respectively.

(286) 6.4.1.4 Characterization of the Recombinant NDVs Expressing Bovine RSV F Proteins.

(287) The presence of the inserted gene in the genome of the recombinant NDVs was confirmed by RT-PCR using viral RNA as template and primers flanking the insertion site. The amplified product was confirmed by sequencing.

(288) Expression of the recombinant bovine RSV F proteins was confirmed by double immunofluorescence in Vero cells infected with the recombinant NDVs as described for the human RSV F expressing NDVs, using the same antibodies (monoclonal antibodies Synagis recognizes the F protein from both human and bovine RSV).

(289) Absence of neutralization of the recombinant NDVs expressing bovine RSV F proteins by the potent RSV neutralizing Synagis antibody was performed as described for the recombinant NDVs expressing human RSV F proteins.

(290) 6.4.2 Results

(291) To assess the ability of Synagis®, a monoclonal antibody that binds to RSV F protein, to neutralize the recombination NDV constructs. As shown in FIG. 14, the recombinant RSV-GFP was neutralized by Synagis®, but NDV-GFP was not inhibited. Similarly, the recombinant NDV constructs, rNDV-bovine RSV-Fopt, and rNDV-bovine RSV-Fchim as well as wild-type NDV were not neutralized by Synagis® (FIGS. 14 and 18). For all the recombinant NDVs, no reduction in the percentage of infected cells at any of the dilutions tested was observed.

6.5 Example 5: Recombinant NDV Encoding Human Metapneumovirus F Protein

(292) 6.5.1 Generation of the Recombinant NDVs Expressing Human Metapneumovirus F Proteins.

(293) The generation of the recombinant NDVs expressing a full length codon optimized F protein from human MPV (strain CAN00-16) (see SEQ ID NO:18 for the codon optimized nucleic acid sequence encoding hMPV F protein) and a chimeric F protein with the codon optimized ectodomain from human MPV (strain CAN00-16) fused to the transmembrane and cytoplasmic domains of NDV (strain LaSota) (see SEQ ID NO:9 for the nucleic acid sequence encoding chimeric F protein) was performed as described for the construction of the recombinant NDVs expressing the full length and chimeric F proteins from human RSV, respectively.

(294) 6.5.2 Characterization of the Recombinant NDVs Expressing Human Metapneumovirus F Proteins.

(295) The presence of the inserted gene in the genome of the recombinant NDVs was confirmed by

RT-PCR using viral RNA as template and primers flanking the insertion site. The amplified product was confirmed by sequencing.

(296) Expression of the human MPV F protein by the recombinant NDVs was confirmed by double immunofluorescence in Vero cells infected with the recombinant NDVs as described for the human RSV F expressing NDVs, using human monoclonal antibody MPE8 to detect huMPV F protein and a commercial mouse monoclonal antibody to detect the NDV HN protein.

7. Embodiments

(297) Provided herein are the following exemplary embodiments:

(298) 1. A recombinant Newcastle disease virus (NDV) comprising a packaged genome comprising a transgene encoding a human respiratory syncytial virus ("RSV") F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:2, 26, 28, or 30.

(299) 2. A recombinant NDV comprising a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a human RSV F protein ectodomain and an NDV F protein transmembrane and cytoplasmic domains, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:4, 44, 45, or 46.

(300) 3. The recombinant NDV of embodiment 1, wherein the RSV F protein comprises the amino acid sequence set forth in SEQ ID NO:6.

(301) 4. The recombinant NDV of embodiment 2, wherein the chimeric F protein comprises the ectodomain of the amino acid sequence set forth in SEQ ID NO:7.

(302) 5. The recombinant NDV of embodiment 2 or 4, wherein the NDV virion comprises the chimeric F protein.

(303) 6. The recombinant NDV of embodiment 1 or 3, wherein the transgene is inserted between two transcription units of the packaged genome.

(304) 7. The recombinant NDV of embodiment 6, wherein the two transcription units of the packaged genome are the transcription units for the NDV P gene and the NDV M gene.

(305) 8. The recombinant NDV of embodiment 2, 4 or 5, wherein the transgene is inserted between two transcription units of the packaged genome.

(306) 9. The recombinant NDV of embodiment 8, wherein the two transcription units of the packaged genome are the transcription units for the NDV P gene and the NDV M gene.

(307) 10. The recombinant NDV of any one of embodiments 1 to 9 which comprises an NDV backbone which is lentogenic.

(308) 11. The recombinant NDV of any one of embodiments 1 to 9 which comprises an NDV backbone of LaSota strain.

(309) 12. The recombinant NDV of any one of embodiments 1 to 9 which comprises an NDV backbone of Hitchner B1 strain.

(310) 13. The recombinant NDV of embodiment 1 or 3, wherein the genome comprises a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO:3.

(311) 14. The recombinant NDV of embodiment 2, 4 or 5, wherein the genome comprises a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO:5.

(312) 15. An immunogenic composition comprising the recombinant NDV of any one of embodiments 1 to 14.

(313) 16. A method for inducing an immune response to RSV F protein, comprising administering the immunogenic composition of embodiment 15 to a subject.

(314) 17. A method for preventing an RSV disease, comprising administering the immunogenic composition of embodiment 15 to a subject.

(315) 18. A method for immunizing a subject against RSV, comprising administering the immunogenic composition of embodiment 15 to the subject.

(316) 19. The method of any one of embodiment 16 to 18, wherein the composition is administered

to the subject intranasally.

(317) 20. The method of any one of embodiments 16 to 19, wherein the subject is seropositive for anti-RSV F antibodies.

(318) 21. The method of any one of embodiments 16 to 19, wherein the subject is seronegative for anti-RSV F antibodies.

(319) 22. The method of any one of embodiments 16 to 21, wherein the subject is a human.

(320) 23. The method of any one of embodiments 16 to 21, wherein the subject is a human infant six months old or older.

(321) 24. The method of any one of embodiments 16 to 21, wherein the subject is a human child.

(322) 25. The method of any one of embodiments 16 to 21, wherein the subject is a human adult or an elderly human.

(323) 26. A recombinant NDV comprising a packaged genome comprising a transgene encoding a bovine respiratory syncytial virus ("RSV") F protein.

(324) 27. The recombinant NDV of embodiment 26, wherein bovine RSV F protein comprises the amino acid sequence of SEQ ID NO:10.

(325) 28. The recombinant NDV of embodiment 26 or 27, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:9 or 11.

(326) 29. The recombinant NDV of embodiment 26, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:40, 41, 42, or 43.

(327) 30. A recombinant NDV comprising a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises a bovine RSV F protein ectodomain and an NDV F protein transmembrane and cytoplasmic domains.

(328) 31. The recombinant NDV of embodiment 30, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:14, 31, 38 or 39.

(329) 32. The recombinant NDV of embodiment 30, wherein the chimeric F protein comprises the ectodomain of the amino acid sequence set forth in SEQ ID NO:10.

(330) 33. The recombinant NDV of embodiment 30, wherein the chimeric F protein comprises the amino acid sequence set forth in SEQ ID NO:33.

(331) 34. The recombinant NDV of any one of embodiments 30 to 33, wherein the NDV virion comprises the chimeric F protein.

(332) 35. The recombinant NDV of any one of embodiments 26 to 29, wherein the transgene is inserted between two transcription units of the packaged genome.

(333) 36. The recombinant NDV of embodiment 35, wherein the two transcription units of the packaged genome are the transcription units for the NDV P gene and the NDV M gene.

(334) 37. The recombinant NDV of any one of embodiments 30 to 34, wherein the transgene is inserted between two transcription units of the packaged genome.

(335) 38. The recombinant NDV of embodiment 37, wherein the two transcription units of the packaged genome are the transcription units for the NDV P gene and the NDV M gene.

(336) 39. The recombinant NDV of any one of embodiments 26 to 38 which comprises an NDV backbone which is lentogenic.

(337) 40. The recombinant NDV of any one of embodiments 26 to 39 which comprises an NDV backbone of LaSota strain.

(338) 41. The recombinant NDV of any one of embodiments 26 to 39 which comprises an NDV backbone of Hitchner B1 strain.

(339) 42. The recombinant NDV of embodiment 26, wherein the genome comprises a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO:12 or 13.

(340) 43. The recombinant NDV of embodiment 30, wherein the genome comprises a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO: 37 or 59.

- (341) 44. An immunogenic composition comprising the recombinant NDV of any one of embodiments 26 to 43.
- (342) 45. A method for inducing an immune response to RSV F protein, comprising administering the immunogenic composition of embodiment 44 to a bovine subject.
- (343) 46. A method for preventing an RSV disease, comprising administering the immunogenic composition of embodiment 44 to a bovine subject.
- (344) 47. A method for immunizing a subject against RSV, comprising administering the immunogenic composition of embodiment 44 to the bovine subject.
- (345) 48. The method of any one of embodiment 45 to 47, wherein the composition is administered to the subject intranasally.
- (346) 49. The method of any one of embodiments 45 to 48, wherein the subject is seropositive for anti-RSV F antibodies.
- (347) 50. The method of any one of embodiments 45 to 48, wherein the subject is seronegative for anti-RSV F antibodies.
- (348) 51. A method for inducing an immune response to RSV F protein in a human subject seropositive for anti-RSV F antibodies, comprising administering to the subject a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, wherein the transgene encodes a human RSV F protein.
- (349) 52. A method for preventing an RSV disease in a human subject seropositive for anti-RSV F antibodies, comprising administering to the subject a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, wherein the transgene encodes a human RSV F protein.
- (350) 53. A method for immunizing a subject against RSV in a human subject seropositive for anti-RSV F antibodies, comprising administering to the subject a recombinant NDV, wherein the recombinant NDV comprises a packaged genome comprising a transgene, wherein the transgene encodes a human RSV F protein.
- (351) 54. The method of any one of embodiments 51 to 53, wherein the human RSV F protein comprises the amino acid sequence set forth in SEQ ID NO:6, 49, 50 or 58.
- (352) 55. The method of embodiment 54, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO:1 or 2.
- (353) 56. The method of embodiment 54, wherein the transgene comprises a codon optimized nucleic acid sequence encoding the RSV F protein.
- (354) 57. The method of any one of embodiments 51 to 53, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence set forth in SEQ ID NO: 25, 26, 27, 28, 29 or 30.
- (355) 58. The method of any one of embodiments 53 to 57, wherein the transgene is inserted between two transcription units of the packaged genome.
- (356) 59. The method of embodiment 58, wherein the two transcription units of the packaged genome are the transcription units for the NDV P gene and the NDV M gene.
- (357) 60. The method of any one of embodiments 51 to 59, wherein the recombinant NDV comprises an NDV backbone which is lentogenic.
- (358) 61. The method of any one of embodiments 51 to 59, wherein the recombinant NDV comprises an NDV backbone of LaSota strain.
- (359) 62. The method of any one of embodiments 51 to 59, wherein the recombinant NDV comprises an NDV backbone of Hitchner B1 strain.
- (360) 63. The method of any one of embodiments 51 to 54, wherein the genome comprises a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO:3.
- (361) 64. The method of any one of embodiments 51 to 63, wherein the composition is administered to the subject intranasally.
- (362) 65. The method of any one of embodiments 51 to 64, wherein the human subject is an elderly

human.

(363) 66. The method of any one of embodiments 51 to 64, wherein the human subject is a human adult.

(364) 67. The method of any one of embodiments 51 to 64, wherein the human subject is a human child.

(365) 68. A recombinant NDV comprising a packaged genome comprising a transgene encoding a human metapneumovirus ("hMPV") F protein.

(366) 69. The recombinant NDV of embodiment 68, wherein the hMPV F protein comprises the amino acid sequence set forth in SEQ ID NO: 17.

(367) 70. The recombinant NDV of embodiment 68 or 69, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:16 or 18.

(368) 71. The recombinant NDV of embodiment 68, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO:52, 53, 54, 55, 56, or 57.

(369) 72. A recombinant Newcastle disease virus (NDV) comprising a packaged genome comprising a transgene encoding a chimeric F protein, wherein the chimeric F protein comprises an hMPV F protein ectodomain and an NDV F protein transmembrane and cytoplasmic domains.

(370) 73. The recombinant NDV of embodiment 72, wherein the chimeric F protein comprises the amino acid sequence of SEQ ID NO:15.

(371) 74. The recombinant NDV of embodiment 72, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 19, 32, 34, 35 or 36.

(372) 75. The recombinant NDV of embodiment 72, 73 or 74, wherein the NDV virion comprises the chimeric F protein.

(373) 76. The recombinant NDV of any one of embodiments 68 to 71, wherein the transgene is inserted between two transcription units of the packaged genome.

(374) 77. The recombinant NDV of embodiment 76, wherein the two transcription units of the packaged genome are the transcription units for the NDV P gene and the NDV M gene.

(375) 78. The recombinant NDV of any one of embodiments 72 to 75, wherein the transgene is inserted between two transcription units of the packaged genome.

(376) 79. The recombinant NDV of embodiment 78, wherein the two transcription units of the packaged genome are the transcription units for the NDV P gene and the NDV M gene.

(377) 80. The recombinant NDV of any one of embodiments 68 to 79 which comprises an NDV backbone which is lentogenic.

(378) 81. The recombinant NDV of any one of embodiments 68 to 79 which comprises an NDV backbone of LaSota strain.

(379) 82. The recombinant NDV of any one of embodiments 68 to 78 which comprises an NDV backbone of Hitchner B1 strain.

(380) 83. The recombinant NDV of embodiment 68, wherein the genome comprises a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO:20 or 21.

(381) 84. The recombination NDV of embodiment 72, wherein the genome comprises a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO: 22.

(382) 85. An immunogenic composition comprising the recombinant NDV of any one of embodiments 68 to 84.

(383) 86. A method for inducing an immune response to hMPV F protein, comprising administering the immunogenic composition of embodiment 85 to a human subject.

(384) 87. A method for preventing an hMPV disease, comprising administering the immunogenic composition of embodiment 85 to a human subject.

(385) 88. A method for immunizing a subject against hMPV, comprising administering the immunogenic composition of embodiment 85 to a human subject.

- (386) 89. The method of any one of embodiments 86 to 88, wherein the composition is administered to the subject intranasally.
- (387) 90. The method of any one of embodiments 86 to 89, wherein the subject is seropositive for anti-hMPV F antibodies.
- (388) 91. The method of any one of embodiments 86 to 89, wherein the subject is seronegative for anti-hMPV F antibodies.
- (389) 92. The method of any one of embodiments 86 to 91, wherein the subject is a human infant.
- (390) 93. The method of any one of embodiments 86 to 91, wherein the subject is an elderly human.
- (391) 94. A kit comprising the recombinant NDV of any one of embodiments 1 to 14, 26 to 43, or 68 to 84.
- (392) 95. A cell line or chicken embryonated egg comprising the propagating the recombinant NDV of any one of embodiments 1 to 14, 26 to 43, or 68 to 84.
- (393) 96. A method for propagating the recombinant NDV of any one of embodiments 1 to 14, 26 to 43, or 68 to 84, the method comprising culturing the cell or embryonated egg of embodiment 95.
- (394) 97. The method of embodiment 96, wherein the method further comprises isolating the recombinant NDV from the egg or embryonated egg.
- (395) The invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described will become apparent to those skilled in the art from the foregoing description and accompanying Figures. Such modifications are intended to fall within the scope of the appended claims.
- (396) All references cited herein are incorporated herein by reference in their entirety and for all purposes to the same extent as if each individual publication or patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety for all purposes.

Claims

1. A recombinant Newcastle disease virus (NDV) comprising a packaged genome comprising (a) a transgene encoding a fusion (F) protein of a human virus of the family Pneumoviridae, wherein the transgene comprises (i) a ribonucleic acid (RNA) sequence corresponding to the negative sense of the complementary deoxyribonucleic acid (cDNA) sequence of SEQ ID NO: 2, 26, 28, or 30, or (ii) an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 18, 53, 55, or 57; (b) a transgene encoding a chimeric protein comprising an ectodomain of a F protein of a human virus of the family Pneumoviridae and a transmembrane domain and a cytoplasmic domain of a NDV F protein, wherein: (i) the human virus of the family Pneumoviridae is human respiratory syncytial virus ("RSV"), and the chimeric protein comprises the ectodomain of the amino acid sequence set forth in SEQ ID NO:7, (ii) the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 19, 32, 34, 35, or 36, or (iii) the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 4, 44, 45, or 46; or (c) a transgene encoding a bovine respiratory syncytial virus ("RSV") F protein, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 11, 41, or 43; or (d) a transgene encoding a chimeric protein comprising a bovine Respiratory Syncytial Virus (RSV) F protein ectodomain and a transmembrane domain and a cytoplasmic domain of a NDV F protein, wherein: (i) the chimeric protein comprises the amino acid sequence set forth in SEQ ID NO:33, or (ii) the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 14, 31, 38, or 39.
2. The recombinant NDV of claim 1, wherein the human virus of the family Pneumoviridae is the RSV, and the chimeric protein comprises the ectodomain of the amino acid sequence set forth in

SEQ ID NO:7.

3. The recombinant NDV of claim 1, wherein the genome comprises a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO: 3, 5, 13, 37 or 59.

4. An immunogenic composition comprising the recombinant NDV of claim 1.

5. A method for inducing an immune response to an F protein of a human virus of the family Pneumoviridae in a subject, preventing a human virus of the family Pneumoviridae disease in a subject, or immunizing a subject against a human virus of the family Pneumoviridae, comprising administering the immunogenic composition of claim 4 to the subject.

6. The recombinant NDV of claim 1, wherein the chimeric protein comprises the amino acid sequence set forth in SEQ ID NO:33.

7. The recombinant NDV of claim 1, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 11, 14, 31, 38, 39, 41, or 43.

8. A method for inducing an immune response to RSV F protein to a bovine subject, preventing an RSV disease in a bovine subject, or immunizing a bovine subject against RSV, the method comprising administering the immunogenic composition of claim 1 to the bovine subject.

9. A method for inducing an immune response to RSV F protein in a human subject seropositive for anti-RSV F antibodies, preventing an RSV disease in a human subject seropositive for anti-RSV F antibodies, or immunizing a subject against RSV in a human subject seropositive for anti-RSV F antibodies, the method comprising administering the immunogenic composition of claim 1 to the human subject.

10. The recombinant NDV of claim 1, wherein (a) the F protein of a human virus of the family Pneumoviridae is a human metapneumovirus ("hMPV") F protein; or (b) the ectodomain of a fusion (F) protein of a human virus of the family Pneumoviridae is a human metapneumovirus (hMPV) fusion (F) protein ectodomain.

11. The recombinant NDV of claim 10, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 18, 53, 55, 57, 19, 32, 34, 35 or 36.

12. The recombinant NDV of claim 10, wherein the genome comprises a negative sense RNA sequence transcribed from the cDNA sequence set forth in SEQ ID NO: 21, or 22.

13. An immunogenic composition comprising the recombinant NDV of claim 10.

14. A method for inducing an immune response to hMPV F protein, preventing an hMPV disease, or immunizing a subject against hMPV, the method comprising administering the immunogenic composition of claim 13 to a human subject.

15. A cell line or chicken embryonated egg comprising the propagating the recombinant NDV of claim 1.

16. A method for propagating the recombinant NDV of claim 1, the method comprising culturing the cell or embryonated egg of claim 15.

17. The recombinant NDV of claim 1, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 2, 26, 28, or 30.

18. The recombinant NDV of claim 1, wherein the transgene comprises an RNA sequence corresponding to the negative sense of the cDNA sequence of SEQ ID NO: 4, 44, 45, or 46.
