



US012390525B2

(12) **United States Patent**  
**Ertl et al.**

(10) **Patent No.:** US 12,390,525 B2  
(45) **Date of Patent:** \*Aug. 19, 2025

(54) **CONSTRUCTS FOR ENHANCING IMMUNE RESPONSES**

(71) Applicant: **THE WISTAR INSTITUTE OF ANATOMY AND BIOLOGY,**  
Philadelphia, PA (US)

(72) Inventors: **Hildegund C. J. Ertl**, Villanova, PA (US); **Marcio O. Lasaro**, Maple Shade, NJ (US); **Luis C. S. Ferreira**, Sao Paulo (BR)

(73) Assignee: **THE WISTAR INSTITUTE OF ANATOMY AND BIOLOGY,**  
Philadelphia, PA (US)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 682 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **17/529,665**

(22) Filed: **Nov. 18, 2021**

(65) **Prior Publication Data**

US 2022/0331423 A1 Oct. 20, 2022

**Related U.S. Application Data**

(63) Continuation of application No. 16/408,730, filed on May 10, 2019, now Pat. No. 11,207,402, which is a continuation of application No. 15/639,296, filed on Jun. 30, 2017, now Pat. No. 10,328,146, which is a continuation of application No. 14/628,784, filed on Feb. 23, 2015, now Pat. No. 9,724,406, which is a continuation of application No. 13/239,771, filed on Sep. 22, 2011, now Pat. No. 8,962,816, which is a continuation of application No. 12/438,889, filed as application No. PCT/US2007/018939 on Aug. 28, 2007, now abandoned.

(60) Provisional application No. 60/840,526, filed on Aug. 28, 2006.

(51) **Int. Cl.**

**A61K 39/245** (2006.01)  
**A61K 39/12** (2006.01)  
**C07K 14/005** (2006.01)  
**C12N 7/00** (2006.01)  
**A61K 39/00** (2006.01)

(52) **U.S. Cl.**

CPC ..... **A61K 39/245** (2013.01); **A61K 39/12** (2013.01); **C07K 14/005** (2013.01); **C12N 7/00** (2013.01); **A61K 2039/53** (2013.01); **A61K 2039/57** (2013.01); **C07K 2319/00** (2013.01); **C07K 2319/10** (2013.01); **C12N 2710/10043** (2013.01); **C12N 2710/16622** (2013.01); **C12N 2710/16634** (2013.01); **Y02A 50/30** (2018.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

## U.S. PATENT DOCUMENTS

5,814,486 A	9/1998	Cohen et al.
6,936,255 B1	8/2005	Wettendorff et al.
8,962,816 B2	2/2015	Ertl et al.
9,624,510 B2	4/2017	Ertl et al.
9,724,406 B2 *	8/2017	Ertl ..... A61K 39/245
10,328,146 B2	6/2019	Ertl et al.
11,207,402 B2 *	12/2021	Ertl ..... C07K 14/005
2004/0253210 A1	12/2004	Robert-Guroff et al.
2005/0095270 A1	5/2005	Staecker et al.
2013/0315871 A1	11/2013	Roy et al.
2014/0065105 A1	3/2014	Wilson et al.
2014/0248305 A1	9/2014	Ertl et al.
2019/0167813 A1	6/2019	Ertl et al.

## FOREIGN PATENT DOCUMENTS

EP	1336619 A2	8/2003
WO	2006120034 A1	11/2006
WO	2007071997 A2	6/2007

## OTHER PUBLICATIONS

European Search Report, EP17837398, date of mailing Apr. 20, 2020.

International Search Report and Written Opinion, PCT/US17/43315, date of mailing Nov. 30, 2017.

International Search Report, PCT/US2007/018939, date of mailing May 16, 2008.

Singapore Search Report, SG Application No. 11201900808S, mailing date Apr. 29, 2020.

Altstein , et al., "Immunization with influenza A NP-Expressing Vaccinia Virus Recombinant Protects Mice Against Experimental Infection with Hyuman and Avian Influenza Viruses", Archives of Virology, vol. 151, No. 5, May 2006, pp. 921-931.

Alves , et al., "Antibody Response in Mice Immunized with a Plasmid DNA Encoding the Colonization Factor Antigen I of Enterotoxigenic *Escherichia coli*", FEMS Immunology Medical Microbiology, vol. 23, No. 4, Apr. 1999, pp. 321-330.

Bayer , et al., "Improved vaccine protection against retrovirus infection after co-administration of adenoviral vectors encoding viral antigens and type I interferon subtypes.", 2011 Retrovirology 8:75 (15 pages).

Casimiro , et al., "Attenuation of simian immunodeficiency virus SIVmac239 infection by prophylactic immunization with DNA and recombinant adenoviral vaccine vectors expressing Gag.", 2005, J Virol 79(24):15547-15555.

(Continued)

*Primary Examiner* — Janet L Andres

*Assistant Examiner* — Myron G Hill

(74) *Attorney, Agent, or Firm* — Saul Ewing LLP;  
Kathryn Doyle; Lukas Pfannenstiel

(57) **ABSTRACT**

Chimeric protein constructs including a herpesvirus glycoprotein D (gD) and a heterologous polypeptide that interact with herpes virus entry mediator (HVEM) and enhance and enhance an immune response against the heterologous polypeptide and methods for their use are provided.

**23 Claims, 38 Drawing Sheets**

**Specification includes a Sequence Listing.**

(56)

**References Cited****OTHER PUBLICATIONS**

- Casimiro , et al., "Comparative Immunogenicity in Rhesus Monkeys of DNA Plasmid, Recombinant Vaccinia Virus, and Replication-Defective Adenovirus Vectors Expressing a Human Immunodeficiency Virus Type 1 gag Gene", 2003, J Virol 77(11):6305-5313.
- Cervasi , et al., "Immunological and Virological Analyses of Rhesus Macaques Immunized with Chimpanzee Adenoviruses Expressing the Simian Immunodeficiency Virus Gag/Tat Fusion Protein and Challenged Intrarectally with Repeated Low Doses of SIVmac", Journal of Virology, 87(17), Sep. 2013, 9420-9430.
- Chawla , et al., "Adenovirus-vectorized vaccines", Dev Biol Stand, Mar. 16, 2008, vol. 18, No. 3, pp. 293-307, Table 1.
- Chen , et al., "Adenovirus-Based Vaccines: Comparison of Vectors from Three Species of Adenoviridae.", 2010, Journal of Virology 84(20):10522-32.
- Chen , et al., "Direct Observation of Xe and Kr Adsorption in a Xe-Selective Microporous Metal-Organic Framework.", J. Am. Chem. Soc., May 2015, 137, 7007-7010.
- Engram , et al., "Vaccine-induced, simian immunodeficiency virus-specific CD8+ T cells reduce virus replication but do not protect from simian immunodeficiency virus disease progression.", 2009, J Immunol 183:706-717.
- Hazama , et al., "Adjuvant-independent enhanced immune responses to recombinant Herpes Simplex Virus Type 1 Glycoprotein D by fusion biologically active interleukin-2", Vaccine, vol. 11, No. 6, 1993, pp. 629-636. (Abstract Only).
- He , et al., "A simplified system for generating recombinant adenoviruses.", 1998, PNAS 95:2509-14.
- Hinuma , et al., "A novel strategy for converting recombinant viral protein into high immunogenic antigen", FEBS Letters, vol. 288, No. 1/2, Aug. 1991, pp. 138-142 (Abstract Only).
- Horwitz , "Function of adenovirus E3 proteins and their interactions with immunoregulatory cell proteins.", 2004, J Gene Med 6:S172-S183 (Abstract only).
- Lasaro , et al., "Antibody-inducing properties of a prototype bivalent Herpes Simplex virus/Enterotoxigenic *Escherichia coli* DNA Vaccine", FEMS Immunology and Medical Microbiology, vol. 35, No. 1, Jan. 21, 2003, pp. 25-31.
- Lasaro , et al., "Anti-Tumor DNA Vaccines based on the Expression of Human Papillomavirus-16 E6/E7 Oncoproteins Genetically Fused With The Glycoprotein D from Herpes Simplex Virus-1", Microbes and Infection, vol. 7, No. 15, Dec. 2005, pp. 1541-1550. (Abstract Only).
- Lasaro , et al., "Human papillomavirus-associated cervical cancer: Prophylactic and therapeutic vaccines.", Gene Therapy Molecular Biology, 2004, vol. 8, pp. 291-306.
- Lasaro , et al., "New insights on adenovirus as vaccine vectors", Molecular Therapy, vol. 17, No. 8, Aug. 1, 2009, pp. 1333-1339.
- Lasaro , et al., "Vaccine-induced T cells Provide Partial Protection Against High-dose Rectal SIVmac239 Challenge of Rhesus Macaques", Molecular Therapy, 19(2), Feb. 2011, 417-426.
- Lewis , et al., "Response of a simian immunodeficiency virus (SIVmac251) to raltegravir: a basis for a new treatment for simian AIDS and an animal model for studying lentiviral persistence during antiretroviral therapy", Retrovirology, 7(21), 2010, 1-19.
- Lichtenstein , et al., "Functions and mechanisms of action of the adenovirus E3 proteins.", 2004, International Reviews of Immunology 23: 75-111.
- McCoy , et al., "Effect of Preexisting Immunity to Adenovirus Human Serotype 5 Antigens on the Immune Responses of Nonhuman Primates to Vaccine Regimens Based on Human- or Chimpanzee-Derived Adenovirus Vectors", Journal of Virology, 81(12), Jun. 2007, 6594-6604.
- Michel , et al., "Enhanced Immunogenicity of HPV 16 E7 Fusion Proteins in DNA Vaccination", Virology, 2002, vol. 294, pp. 47-59.
- Patel , et al., "DNA and virus particle vaccination protects against acquisition and confers control of viremia upon heterologous simian immunodeficiency virus challenge", PNAS, 110(8), Feb. 2013, 2975-2980.
- Plonka , et al., "Light Hydrocarbon Adsorption Mechanisms in Two Calcium-Based Microporous Metal Organic Frameworks.", Chem. Mater. 2016, 28, 1636-1646 (Abstract Only).
- Saha , et al., "A Fused Gene of Nucleoprotein (NP) and Herpes Simplex Virus Genes (VP22) Induces Highly Protective Immunity Against Different Subtypes of Influenza Virus", Virology, vol. 354, No. 1, Oct. 10, 2006, pp. 48-57.
- Santra , et al., "Heterologous Prime/Boost Immunizations of Rhesus Monkeys Using Chimpanzee Adenovirus Vectors", Vaccine, 27(42), Sep. 2009, 5837-5845.
- Shiver , et al., "Replication-incompetent adenoviral vaccine vector elicits effective anti-immunodeficiency-virus immunity.", 2002, Nature 415:331-335.
- Small , et al., "Viruses—From Pathogens to Vaccine Carriers", Curr. Opin. Virol., 1(4), Oct. 2011, 241-245.
- Tatsis , et al., "Adenovirus Vector-Induced Immune Responses in Nonhuman Primates: Responses to Prime Boost Regimens", Journal of Immunology, 182, 2009, 6587-6599.
- Tatsis , et al., "Chimpanzee-origin adenovirus vectors as vaccine", 2006, Gene Therapy 13:421-429.
- Tatsis , et al., "Multiple Immunizations with Adenovirus and MVA vectors improve CD8+ T cell functionality and mucosal homing", Virology, 367(1), Oct. 2007, 156-167.
- Watson , et al., "Herpes Simplex Virus Type=1 Glycoprotein D Gene: Nucleotide Sequence and Expression in *Escherichia coli*", Science, vol. 218, Oct. 22, 1982, pp. 381-384 (Abstract Only).
- Xiang , "Chimpanzee Adenovirus Antibodies in Humans, Sub-Saharan Africa", 2006, Emerging Infectious Diseases 12(10):1596-1599.
- Zago , et al., "Use of herpes simplex virus and pseudorabies virus chimeric glycoprotein D molecules to identify regions critical for membrane fusion.", PNAS, 2004, vol. 101, No. 50, pp. 17498-17503.
- Zhou , et al., "A Universal Influenza A Vaccine Based on Adenovirus Expressing Matrix-2 Ectodomain and Nucleoprotein Protects Mice from Lethal Challenge", Molecular Therapy, 18(12), Dec. 2010, 2182-2189.
- Zhou , et al., "An efficient method of directly cloning chimpanzee adenovirus as a vaccine vector", Natl. Protoc., 5(11), Nov. 2010, 1775-1785.
- Zolla-Pazner , et al., "Analysis of V2 Antibody Responses Induced in Vaccinees in the ALVAC/AIDSVAX HIV-1 Vaccine Efficacy Trial", PLOS ONE, 8(1), Jan. 2013, 1-11.

\* cited by examiner

FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 4




FIG. 5




FIG. 6







FIG. 7



**FIG. 8A**



**FIG. 8B**







FIG. 9



**FIG. 10A**




FIG. 10B





FIG. 11

FIG. 12






FIG. 13




FIG. 14




FIG. 15




FIG. 16

FIG. 17




FIG. 18A




FIG. 18C






FIG. 18B

FIG. 18D

FIG. 19




FIG. 20





FIG. 21






FIG. 22




FIG. 23A




FIG. 23C

FIG. 23B

FIG. 24A





FIG. 24B






FIG. 25




FIG. 26A

FIG. 26B

FIG. 26C




FIG. 27A




FIG. 27B

FIG. 28A




FIG. 28B




FIG. 28C




FIG. 29




FIG. 30A




FIG. 30B




FIG. 31A




FIG. 31B




FIG. 32A




FIG. 32B




FIG. 32C




FIG. 33A





FIG. 33B






FIG. 33C

## CONSTRUCTS FOR ENHANCING IMMUNE RESPONSES

### CROSS-REFERENCE TO RELATED APPLICATIONS

The present application is a continuation of U.S. patent application Ser. No. 16/408,730, filed on May 10, 2019, which is a continuation of U.S. patent application Ser. No. 15/639,296, filed on Jun. 30, 2017, now U.S. Pat. No. 10,328,146, which is a continuation of U.S. patent application Ser. No. 14/628,784, filed on Feb. 23, 2015, now U.S. Pat. No. 9,724,406, which is a continuation of U.S. patent application Ser. No. 13/239,771, filed on Sep. 22, 2011, now U.S. Pat. No. 8,962,816, which is a continuation of U.S. patent application Ser. No. 12/438,889, filed on Feb. 25, 2009, which is a National Stage application of PCT/US2007/018939, filed on Aug. 28, 2007, which claims priority under 35 U.S.C. § 119(e) to U.S. Provisional Patent Application No. 60/840,526, filed on Aug. 28, 2006, each of which application is incorporated by reference in its entirety herein.

### STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

This invention was made with government support under 5 P01 AI052271, awarded by the National Institutes of Health (NIH). The government has certain rights in the invention.

### FIELD OF THE INVENTION

Embodiments of the present invention relate in general to chimeric (fusion) protein constructs including a herpesvirus glycoprotein D (gD) and a heterologous polypeptide (e.g., antigen) that enhance the immune response against the heterologous polypeptide (e.g., antigen) in a subject.

### BACKGROUND OF THE INVENTION

gD is the receptor-binding glycoprotein of herpesviruses (Fusco et al. (2005) *Proc. Natl. Acad. Sci. U.S.A.* 102:9323). The gD ectodomain is organized in two structurally and functionally differentiated regions. The amino-terminus includes the signal sequence and receptor-binding sites, and the carboxy-terminus includes the pro-fusion domain and the transmembrane domain. gD interacts with two alternative receptors belonging to unrelated protein families, the herpesvirus entry mediator (HVEM) and the nectins (Geraghty et al. (1998) *Science* 280:1618; Montgomery et al. (1996) *Cell* 87:427; Cocchi et al. (1998) *J. Virol.* 72:9992; Warner et al. (1998) *Virology* 246:179; Lopez et al. (2000) *J. Virol.* 74:1267). HVEM is expressed on dendritic cells and the B and T lymphocyte attenuator (BTLA) is expressed on activated T and B lymphocytes. The interaction between HVEM and BTLA results in the down-regulation of immune responses.

### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 depicts a schematic representation of the chimeric gene gDE7E6E5. The HPV-16 E5, E6 and E7 genes without respective start and stop codons were linked in tandem and incorporated into the HSV-1 gD gene ApaI site, which corresponds to amino acid 244 in the gD mature form.

FIG. 2 depicts a schematic representation of chimeric gene gDgag. The codon-optimized truncated form of gag from HIV-1 clade B was fused into the HSV-1 gD gene Nan site, which corresponds to amino acid 289 in the gD mature form.

FIGS. 3A-3B depict the gag-specific CD8<sup>+</sup> IFN- $\gamma$  response in mice immunized with vaccine constructs carrying the gDgag chimeric gene. FIG. 3A depicts a FACS analysis of the gag-specific CD8<sup>+</sup>/IFN- $\gamma$ <sup>+</sup> response in peripheral blood mononuclear cells (PBMC) from mice immunized with DNA vaccine expressing either HIV-1 gag or HIV-1 gag fused to HSV-1 gD (gDgag). Numbers on the right corner represent percentage CD8<sup>+</sup>/IFN- $\gamma$ <sup>+</sup> cells over total of CD8<sup>+</sup> cells. FIG. 3B graphically depicts PBMC from mice immunized with AdC68 vectors carrying either genes encoding gag or gDgag, inoculated with different amounts of virus particles per mouse.

FIG. 4 graphically depicts the effect of pre-existing immunity to the AdHu5 adenovirus vector on the transgene product-specific CD8<sup>+</sup> T cell response to the AdC68 vector. AdC68 vectors carrying gag, gDgag or gDE7E65 were inoculated into naïve mice and mice previously immunized with an AdHu5 expressing an unrelated antigen (rabies glycoprotein, AdHu5rab(gp)). The percentage of reduction of CD8<sup>+</sup> T cell response was defined as a percentage of CD8<sup>+</sup>/IFN- $\gamma$ <sup>+</sup> frequency in mice previously immunized with AdHu5 over the frequency found in mice that did not receive AdHu5 vector.

FIG. 5 depicts the E7-specific CD8<sup>+</sup> IFN- $\gamma$  response in mice immunized with DNA vaccines expressing non-mutated or mutated gDE7E6E5. The mutation was designed to disrupt the HVEM binding site of gD. Mice were immunized with non-mutated gDE7E6E5 (pgDE7E6E5) or mutated gDE7E6E5 (pNBEGfDE7E6E5) and 14 days later peripheral blood mononuclear cells were investigated by intracellular cytokine staining for E7-specific CD8<sup>+</sup> IFN- $\gamma$  responses. Numbers on the right corners represent percentage CD8<sup>+</sup>/IFN- $\gamma$ <sup>+</sup> cells over total of CD8<sup>+</sup> cells.

FIG. 6 depicts the E7-specific CD8<sup>+</sup> IFN- $\gamma$  response in mice immunized with DNA vaccines expressing non-mutated (gDE7) or mutated gD (SgDE7). This mutation (SgD) was designed to increase binding between gD and HVEM. Mice were immunized with pgDE7 or pSgDE7 and 14 days later peripheral blood mononuclear cells were investigated by intracellular cytokine staining for E7-specific CD8<sup>+</sup> IFN- $\gamma$  response. Numbers in the corners represent percentage CD8<sup>+</sup>/IFN- $\gamma$ <sup>+</sup> cells over total of CD8<sup>+</sup> cells.

FIG. 7 graphically depicts an in vitro HVEM binding assay. CHO-CAR cells (Chinese hamster ovary-coxsackie-adenovirus receptor cells) were infected with either AdC68gD or AdC68gDgag and, after 48 hours, total protein was extracted. The amount of gD in each sample was quantified by capture ELISA and the protein extracts were diluted in extraction buffer to normalized levels of gD. Equalized extracts were diluted and added to 96-well plates coated with purified HVEM. The amount of gD bound to HVEM was detected by using anti-gD polyclonal antisera and anti-Rabbit IgG horseradish peroxidase. Data shown is one representative experiment from two performed.

FIGS. 8A-8B depict confocal microscopy for localization of gDgag and HVEM on AdC68gDgag infected cells. B78H1/3E5 cells, which expressed HVEM fused to enhanced green fluorescence protein (HVEM-EGFP), were infected with AdC68gDgag. After 48 hours, cells were either directly stained (FIG. 8A) or permeabilized and stained (FIG. 8B) with an anti-gD monoclonal antibody (DL-6) and

anti-mouse IgG conjugated with Texas Red. Cells were examined with a Leica TCS SP2 Confocal Microscope at 400 $\times$  final magnification.

FIG. 9 graphically depicts FACS analysis of gDgag expression on the surface of AdC68gDgag infected cells. B78H1/3E5 cells (darker line), which express HVEM-EGFP on the surface, and B78H1 cells (lighter line), which do not express HVEM, were infected with AdC68gDgag. Cells were cultivated for 48 hours, and then labeled with an anti-gD monoclonal antibody (DL-6) and anti-mouse IgG conjugated to phycoerythrin (PE). Cell suspensions were analyzed using an EPICS XL (Beckman-Coulter, Inc., Miami, FL) to determine presence of gDgag.

FIGS. 10A-10B depict that gDgag expressed by infected cells bound to HVEM expressed on non-infected cells. Confocal microscopy (FIG. 10A) and FACS analysis (FIG. 10B) were performed to localize gDgag on non-infected cells. CHO-CAR cells were infected with AdC68gDgag. After 48 hours, cells were harvested and washed extensively with cold PBS. AdC68gDgag-infected CHO-CAR cells were cultured with B78H1/3E5 cells, which expresses HVEM-EGFP on the surface, at 4:1 ratio. After 48 hours, cells were stained using the anti-gD monoclonal antibody DL-6 and anti-mouse IgG conjugated to Texas Red (microscopy) or PE (FACS). (FIG. 10A) Microscopy was performed with a Leica TCS SP2 Confocal Microscope at 400 $\times$  final magnification. (FIG. 10B) Cell suspensions were analyzed using an EPICS XL (Beckman-Coulter, Inc., Miami, FL). B78H1/3E5 cells were cultured with either AdC68gDgag-infected CHO-CAR cells (darker line) or non-infected CHO-CAR cells (lighter line). Data on graph show cells which are positive for GFP.

FIG. 11 depicts the gag-specific CD8 $^{+}$  T cell response in mice immunized with AdC68 vectors. PBMC and splenocytes from mice immunized with 1 $\times$ 10 $^{9}$  vp of AdC68 vectors carrying either gag, gDgag or gD were tested. Percentage represents CD8 $^{+}$ /IFN- $\gamma$  $^{+}$  cells over total of CD8 $^{+}$  cells. CD8 $^{+}$ /IFN- $\gamma$  $^{+}$  frequencies in all groups stimulated with an unrelated control peptide were below 0.20%. Data shown are representative of two performed experiments.

FIG. 12 graphically depicts the gag-specific IFN- $\gamma$  response of CD8 $^{+}$  T cells from mice immunized with AdC68 vectors. PBMC are from mice immunized with different amounts of AdC68 vectors carrying either gag, gDgag or gD. Percentage represents CD8 $^{+}$ /IFN- $\gamma$  $^{+}$  cells over total of CD8 $^{+}$  cells. CD8 $^{+}$ /IFN- $\gamma$  $^{+}$  frequencies in all groups stimulated with an unrelated control peptide were 0.20%. Data shown are representative of two performed experiments.

FIG. 13 depicts the phenotypic profile of CD8 $^{+}$  cells activated by AdC68 vaccination. PBMC from naïve mice and mice immunized with AdC68 carrying either gag or gDgag were stained with gag-tetramer-APC and anti-CD8-PerCP, in combination with anti-CD25-PE, anti-CD-122-PE, anti-CD127-PE, anti-CD27-PE, anti-CD62L-FITC, anti-CD69-PE, anti-CD103-PE, anti-CD43-PE, anti-CD44-FITC, anti-CD54-PE, anti-Bcl2-PE, anti-BTLA-PE, anti-CTLA4-PE and anti-PD1-PE. Graphs show data from CD8 $^{+}$ /gag-tet $^{+}$  cells for AdC68gag (gray line) and AdC68gDgag (black line), and total CD8 $^{+}$  for naïve (black dotted line). Data were analyzed on Flowjo software (Tree Star Inc.).

FIG. 14. Intracellular IFN- $\gamma$  staining of gag-specific and AAV-specific CD8 $^{+}$  T cells from mice immunized with AAV vectors expressing gD, gag or gDgag. Detection of gag-specific and AAV-specific CD8 $^{+}$  T cells was carried out after stimulation of peripheral blood mononuclear cells (PBMCs) with either MHC class I restricted gag peptide or AAV-

capsid peptide and cell surface staining for CD8 and intracellular staining for IFN- $\gamma$ . Unspecific peptide was used as control. The numbers in the right upper corners show the frequencies of peptide-specific CD8 $^{+}$  T cells, as percentages of IFN- $\gamma$ -producing CD8 $^{+}$  T cells over all detected CD8 $^{+}$  T cells.

FIG. 15. Schematic representation of chimeric gene gD-NP. Nucleoprotein P (NP) from Influenza virus A/PR8 without its start and stop codons was incorporated into the HSV-1 gD Apal site, which corresponds to amino acid 244 in the mature form of gD.

FIG. 16. NP-tetramer staining of CD8 $^{+}$  T cells isolated from blood of mice immunized with either pgD-NP or pNP DNA vaccine. Mice were immunized with 100  $\mu$ g of each DNA vaccine vector. Fourteen days after immunization peripheral blood mononuclear cells (PBMCs) were isolated and cell surface stained with the NP-tetramer and a labeled antibody to CD8. Naïve mice were used as negative control. Data represent percentages of NP-tetramer $^{+}$  CD8 $^{+}$  T cells over all detected CD8 $^{+}$  T cells.

FIG. 17. Schematic representation of chimeric gene gDTRAPTB. Thrombospondin-related anonymous protein (TRAP) from parasite *Plasmodium falciparum* and *Mycobacterium tuberculosis* epitope string (TB) without their start and stop codons are incorporated into the HSV-1 gD Nan site, which corresponds to amino acid 288 in the gD mature form.

FIGS. 18A-18D. Molecular modeling of the gD-gag chimeric protein. Ribbon representations of gD-gag in the unligated (FIGS. 18A, 18B) and HVEM ligated (FIGS. 18C, 18D) conformations. FIG. 18A, the gag insertion is connected to the gD ectodomain core by a long flexible linker. FIG. 18B, a superposition of the native gD X-ray structure (2C36) (darker ribbon) and that of the gD-gag chimera model (lighter ribbon) shows that the gag insert repositions the C-terminus of native gD away from the HVEM binding pocket. The dashed line indicates an 11 residue gD loop segment that like the first 22 N-terminal residues (not shown) is unresolved in the X-ray structure and presumed to be highly flexible (Krummenacher et al., *EMBO J.* 24, 4144-53, 2005). FIG. 18C, gD-gag chimera model in the HVEM ligated conformation with HVEM positioned as observed in the gD-HVEM complex X-ray structure (1JMA). The gD N-terminus changes conformation upon formation of the HVEM complex. FIG. 18D, a superposition of the native gD X-ray structure (2C36) (darker ribbon) with the gD-gag chimera model (lighter ribbon) in the HVEM-bound conformation shows that the gag insert does not disrupt the gD core domain.

FIG. 19. Confocal microscopy was carried out with B78-H1/3E5 cells, which express HVEM fused to Enhanced Green Fluorescence Protein (HVEM-EGFP). B78-H1/3E5 cells were infected with AdC68gag, AdC68gD or AdC68gD-gag, then stained anti-gD DL-6 MAb and anti-mouse IgG conjugated with Texas Red. AdC68gD-gag-infected cells were permeabilized then stained as above. Cells were examined with a Leica TCS SP2 Confocal Microscope at 400 $\times$  final magnification.

FIG. 20. Comparison of gD expression on the surface of B78-H1 (black line) and B78-H1/3E5 (gray) cells infected with AdC68 vectors carrying gD-gag, gD or gag.

FIG. 21. Presence of gD on the surface of non-infected HVEM $^{+}$  cells co-cultivated with HVEM $^{-}$  cells infected with either AdC68gD-gag, AdC68gD or AdC68gag. Non-infected cells were used as negative control.

FIG. 22. AdC68gD induces enhanced expansion of CD8 $^{+}$  T cells in vitro. Irradiated lymph nodes cells from naïve mice

and mice immunized with either AdC68E7E6E5 or AdC68gD were incubated with CFSE-labeled CD8<sup>+</sup> OT-1 (V $\alpha$ 2<sup>+</sup>) cells for 72 hrs. Total live cells (left) were analyzed for expression of CD8<sup>+</sup>V $\alpha$ 2<sup>+</sup>. CSFE expression by these double positive populations (highlighted by the squares on the right graphs) is shown on the left graphs. The bars and numbers show from right to left the percentages of the population that underwent no replication, 1, 2, 3 or  $\leq$ 4 cycles of replication. Graphs show data from one representative experiment of two performed.

FIGS. 23A-23C. CD8<sup>+</sup> T cell responses to vectors expressing antigens fused to gD. FIG. 23A, intracellular cytokine staining of E7- and gag-specific CD8<sup>+</sup> T cells were carried out on PBMCs from mice i.m. immunized with DNA vaccines (upper graphs) or AdC68 vectors (lower graphs) expressing either gD, E7E6E5, gD-E7E6E5, gag or gD-gag, after stimulation with E7 or gag peptide and cell surface staining for CD8 (FITC) and intracellular staining for IFN- $\gamma$  (PE). PBMCs were isolated from animals 14 days after DNA vaccination or 10 days after application of AdC68 vector. The numbers in the right upper corners show frequencies of IFN- $\gamma$ -producing CD8<sup>+</sup> T cells as a percentage of all CD8<sup>+</sup> T cells. Frequencies of IFN- $\gamma$ <sup>+</sup>/CD8<sup>+</sup> T cells stimulated with an unrelated control peptide were below 0.2% in all groups. FIG. 23B, Gag-specific CD8<sup>+</sup> T cell frequencies were determined 10 days after immunization of mice with decreasing doses of either AdC68gag (open bars) or AdC68gD-gag (black bars) vectors. FIG. 23C, The kinetics of E7-specific CD8<sup>+</sup> T cell responses induced by the AdC68gD-E7E6E5 vector were analyzed from BPMCs of mice immunized with either AdC68E7E6E5 (squares), AdC68gD (diamonds) or AdC68gD-E7E6E5 (triangles) vectors at different days after a single dose of  $10^{10}$  vps of the vaccines.

FIGS. 24A-24B. The enhancement of CD8<sup>+</sup> T cell responses requires binding of gD to HVEM. FIG. 24A, E7-specific IFN- $\gamma$ <sup>+</sup>CD8<sup>+</sup> responses were evaluated with splenocytes from mice immunized with one dose of DNA vaccines expressing the E7E6E5 polypeptide either within wild-type gD (pgD-E7E6E5), a mutated gD that shows loss of binding to HVEM (NBEGd-E7E6E5) or that shows enhanced binding to HVEM (SgD-E7E6E5). FIG. 24B, splenocytes from mice immunized with one dose of DNA vaccines carrying E7 fused to either wild type gD (gD-E7) or mutated gD with high affinity to HVEM (SgD-E7) were evaluated for E7-specific IFN- $\gamma$ <sup>+</sup>CD8<sup>+</sup> response. PBMCs were isolated 14 days after DNA vaccine immunizations.

FIG. 25. Phenotypes of gag-specific CD8<sup>+</sup> T cells were analyzed on PBMCs from mice immunized with either AdC68gD-gag or AdC68gag. PBMCs were isolated 10 days after immunization. Naïve mice were used as controls (black dotted line). The graphs shown reflect expression levels of total CD8<sup>+</sup> T cells from naïve mice (black dotted line) and gag-tet<sup>+</sup>CD8<sup>+</sup> T cells from mice immunized with either AdC68gag (grey line) or AdC68gD-gag (black line).

FIGS. 26A-26C. CD8<sup>+</sup> T cells induced by gD-antigen chimeric protein are functional in vivo. Protection against TC-1 tumor challenge was evaluated in mice vaccinated with DNA (FIG. 26A) or AdC68 vectors (FIG. 26B) expressing either gD (circles), E7E6E5 (diamonds) or gD-E7E6E5 (squares). FIG. 26C, Protection to TC-1 tumor challenge in mice vaccinated with DNA vaccine expressing either NBEGd-E7E6E5 (diamonds), SgD-E7E6E5 (squares), gD-E7 (circles) or SgD-E7 (triangles) chimeric genes. Mice were challenged 14 and 10 days after vaccination with DNA and AdC68 vectors, respectively. Tumor development was followed for up to 60 days after challenge.

FIGS. 27A-27B. Quantification of specific mRNA copies and protein expression by cells infected in vitro with AdC68 vectors. FIG. 27A, RNA isolated from non-infected cells and from cells infected with AdC68gD (white bars), AdC68E7E6E5 (gray bars) or AdC68gD-E7E6E5 (black bars) were reverse transcribed and quantified by Real-Time PCR. After quantification of GAPDH mRNA copies, all samples were normalized to  $10^9$  GAPDH mRNA copies. Specific mRNA copies were quantified using gD, E7, E6, and E5 specific primers. Neither E7, E6 nor E5 mRNA were detected in cells infected with AdC68gD, and no gD specific mRNA was detected in cells infected with AdC68E7E6E5. mRNA levels were assessed in three independent experiments and each sample was investigated in triplicates. p values from two-tail student's t test are shown on top of the bars. FIG. 27B, confocal microscopy was carried out with CHO/CAR cells infected with AdC68 expressing either gD, E7E6E5 or gD-E7E6E5, then permeabilized and stained with anti-gD DL-6 MAb and anti-mouse IgG conjugated with FITC. Immunofluorescence is shown on the top panel while differential interference contrast (DIC) microscopy is shown on the bottom panel. Cells were examined with a Leica TCS SP2 Confocal Microscope at 400 $\times$  magnification.

FIGS. 28A-28C. Intracellular IFN- $\gamma$  staining of E7-specific CD8<sup>+</sup> T cells from mice immunized with AdC68 and DNA vectors. FIG. 28A, frequencies of E7-specific CD8<sup>+</sup> T cells in PBMC (top) or spleens (bottom) from naïve mice or mice immunized with AdC68 vector expressing either gD, E7E6E5 or gD-E7E6E5 were determined 10 days after immunization. FIG. 28B, frequencies of E7-specific CD8<sup>+</sup> T cells in PBMC (top) or spleens (bottom) from naïve mice or mice immunized with DNA vaccines expressing either gD, E7E6E5 or gD-E7E6E5 were determined 14 days after immunization. FIG. 28C, prime and boost regimens with pgD-E7E6E5 and AdC68gD-E7E6E5 vectors. Mice immunized with one dose of pgD-E7E6E5 vector were boosted after 90 days with AdC68gD-E7E6E5 (open bars), while mice immunized with AdC68gD-E7E6E5 were boosted after 90 days with pgD-E7E6E5 (black bars). Detection of E7-specific CD8<sup>+</sup> T response was carried out after stimulation with a MHC class I restricted E7 peptide and cell surface staining for CD8 (FITC) and intracellular staining for IFN- $\gamma$  (PE). The numbers in the right upper corners show the frequencies of E7-specific CD8<sup>+</sup> T cells, as percentages of IFN- $\gamma$ -producing CD8<sup>+</sup> T cells over all detected CD8<sup>+</sup> T cells. IFN- $\gamma$ -producing CD8<sup>+</sup> cell frequencies in all groups stimulated with an unrelated peptide or in the absence of stimulus were below 0.2%. The data shown in a and b are from one representative experiment of four performed.

FIG. 29. Dose-response of the CD8<sup>+</sup> T cell response to AdC68gD-E7E6E5. Frequencies of IFN- $\gamma$ -producing E7-specific CD8<sup>+</sup> T cells in spleens and PBMCs induced by  $5 \times 10^{10}$  to  $1 \times 10^8$  vp/animal of AdC68gD-E7E6E5 vector were determined as described on legend to FIGS. 28A-28C.

FIGS. 30A-30B. Anti-tumor effects of AdC68 vectors against TC-1 cell challenge. FIG. 30A, for post-challenge vaccination, groups of 10 mice were s.c. inoculated with TC-1, then 5 days later immunized with either AdC68gD (triangles), AdC68E7E6E5 (circles) or AdC68gD-E7E6E5 (squares). FIG. 30B, for pre-challenge vaccination, groups of 10 mice vaccinated one year earlier with AdC68 vectors carrying either gD (triangles), E7E6E5 (circles) or gD-E7E6E5 (squares) were s.c. challenged with TC-1. For all TC-1 challenge experiments animals were monitored 3 times per week for evidence of tumor growth over a period of 60 days.

FIGS. 31A-31B. CD8<sup>+</sup> T cell response after challenge with TC-1 in mice vaccinated one year earlier. One year after vaccination with AdC68 vectors expressing either gD, E7E6E5 or gd-E7E6E5, mice were challenged with TC-1 cells and 10 days later E7-specific frequencies of E7-specific CD8<sup>+</sup> T cells were determined. FIG. 31A, ICS of E7-specific CD8<sup>+</sup> T cells in spleen from non-challenged mice (top) or mice challenged with TC-1 cells (bottom). Frequencies of IFN-γ-producing E7-specific CD8<sup>+</sup> T cells in spleen were determined as in FIG. 2 legend. FIG. 31B, E7-tetramer staining of CD8<sup>+</sup> T cells isolated from spleen, blood and liver of mice challenged (white bars) or not (black bars) with TC-1 cells. Data represent percentages of E7-tetramer<sup>+</sup> CD8<sup>+</sup> T cells over all detected CD8<sup>+</sup> T cells. E7-tetramer<sup>+</sup> CD8<sup>+</sup> cells were not detected in mice immunized with either AdC68gD or AdC68E7E6E5.

FIGS. 32A-32C. CD8<sup>+</sup> T cell response and phenotypic profile in mice immunized with AdC68 vectors and subsequently challenged with TC-1 cells. Lymphocytes were isolated 3 days after TC-1 challenge from animals immunized with AdC68gD-E7E6E5 and 7 days after challenge from animals immunized with either AdC68gD or AdC68E7E6E5. E7-specific CD8<sup>+</sup> T response were determined in spleen, PBMC and TIL by ICS (FIG. 32A) and E7-tetramer staining (FIG. 32B). ICS data was determined as in FIG. 2 legend, while E7-tetramer staining data represent percentages of E7-tetramer<sup>+</sup> CD8<sup>+</sup> T cells over all detected CD8<sup>+</sup> T cells. E7-specific IFN-γ<sup>+</sup>CD8<sup>+</sup> cells and E7-tetramer<sup>+</sup>CD8<sup>+</sup> cells were not detected in mice immunized with either AdC68gD or AdC68E7E6E5. FIG. 32C, phenotype analysis of splenocytes, PBMCs and TILs were determined with cells isolated from mice immunized with AdC68gD-E7E6E5 (black line) or AdC68gD (dotted black line) then challenged with TC-1. Cells were stained with E7-tetramer-APC and anti-CD8-PerCP, in combination with antibodies to CD44, CD62L, CD27, Bcl2, BTLA, CTLA-4 and PD-1. CD8<sup>+</sup> T cells isolated from either naïve mice or mice immunized with AdC68E7E6E5 showed similar phenotype profiles as CD8<sup>+</sup> T cells isolated from mice immunized with AdC68gD.

FIGS. 33A-33C. Comparison of CD8<sup>+</sup> T cell responses and phenotype profiles induced by AdC68 vectors in wild-type and HPV-16 E6/E7-tg mice. One-year old E6/E7-tg mice were vaccinated with AdC68 vectors expressing either gD, E7E6E5 or gd-E7E6E5, and 1-year old C57Bl/6 mice were vaccinated with AdC68gD-E7E6E5 vector. Ten days later frequencies and phenotypes of E7-specific CD8<sup>+</sup> T cells were determined. FIG. 33A, E7-specific CD8<sup>+</sup> T cells isolated from spleens of E6/E7-tg mice immunized with AdC68gD, AdC68E7E6E5 or AdC68gD-E7E6E5, or from spleen of wild-type mice immunized with AdC68gD-E7E6E5 were tested by ICS. Frequencies of IFN-γ-producing E7-specific CD8<sup>+</sup> T cells in spleen were determined as described in legend to FIG. 2. FIG. 33B, E7-tetramer staining of CD8<sup>+</sup> T cells was performed with splenocytes and PBMC from wild-type and E6/E7-tg mice, and with lymphocytes from thyroids of E6/E7-tg mice. Data show percentages of E7-tetramer<sup>+</sup> CD8<sup>+</sup> T cells over all detected CD8<sup>+</sup> T cells. ND, not determined. FIG. 33C, phenotypic profile of E7-specific CD8<sup>+</sup> T cells were determined using cells isolated from spleen and blood of AdC68gD-E7E6E5 vaccinated E6/E7 tg (filled gray) and wild-type (black line) mice, and from cells isolated from thyroid of E6/E7 tg mice vaccinated with AdC68gD-E7E6E5. CD8<sup>+</sup> T cells isolated from E6/E7 tg naïve mice were used as controls (black dotted line).

## DETAILED DESCRIPTION

The invention provides chimeric, or fusion, proteins in which one or more antigens is inserted into the C terminal region of a mature HSV gD protein. Such fusion proteins enhance the immune response of a host against the antigen(s) to a much greater degree than is observed without the gD. The gD chimeric proteins of the present invention are particularly suitable for use as genetic vaccines (e.g., DNA vaccines or viral vector vaccines) to therapeutically or prophylactically treat a subject. Thus, the invention also provides nucleic acid molecules which encode fusion proteins of the invention.

## Glycoprotein D

Glycoprotein D (gD) is an envelope glycoprotein found on Herpes simplex viruses such as HSV-1 or HSV-2 and is expressed in cells infected by the viruses. An HSV gD has a 25-amino acid amino-terminal signal sequence and a carboxy-terminal transmembrane domain. The signal sequence is typically cleaved in the mature form of the protein. The amino acid sequence of HSV-1 gD is shown in SEQ ID NO:27 (amino acids 1-25 are the signal sequence; amino acids 26-394 are mature HSV-1 gD); a coding sequence for SEQ ID NO:27 is shown in SEQ ID NO:26. The amino acid sequence of HSV-2 gD is shown in SEQ ID NO:29 (amino acids 1-25 are the signal sequence; amino acids 26-393 are mature HSV-2 gD); a coding sequence for SEQ ID NO:25 is shown in SEQ ID NO:28.

An HSV gD or mutant thereof which is useful in the present invention has the ability to interact with HVEM and, in addition, may have one or more of the following properties: 1) ability to stimulate a CD8<sup>+</sup> T cell response to the fusion partner; 2) ability to disrupt an HVEM-BTLA pathway activity; 3) ability to interact with nectin-1; 4) ability to mediate cell entry by an HSV-1 and/or HSV-2 virus; and 5) ability to mediate cell-to-cell spread of HSV-1 and/or HSV-2. Thus, as used herein, a “gD” or an “immunostimulatory portion of a gD” refers to a polypeptide having an amino acid sequence of a wild-type gD or a mutant thereof which retains one or more gD activities.

gD chimeric (fusion) proteins of the invention comprise at least two, preferably three polypeptide segments. The first polypeptide segment comprises at least amino acids 1-240 of a mature Herpes simplex virus (HSV) glycoprotein D; in preferred embodiments the first polypeptide segment does not comprise a full length mature glycoprotein D; in this case a third polypeptide segment is included. The second polypeptide segment, the N terminus of which is linked to the C terminus of the first polypeptide segment, comprises at least one antigen which is not an HSV glycoprotein D antigen. The third polypeptide comprises a C terminal portion of the HSV glycoprotein D; the N terminus of the third polypeptide segment is linked to the C terminus of the second polypeptide segment. Thus, in certain embodiments an antigen is fused to the carboxy-terminal region of gD. In other aspects, an antigen is inserted within the carboxy-terminal amino acid sequence of gD such that the amino-terminal end of the gD chimeric protein is an amino-terminal gD amino acid sequence, fused to an internal antigenic sequence, fused to a carboxy-terminal amino acid sequence of gD.

In certain embodiments, the first polypeptide segment of a gD chimeric protein of the present invention includes the entire gD amino acid sequence (e.g., amino acids 1-394) or the mature gD amino acid sequence (e.g., amino acids 26-394, the carboxy-terminal 369 amino acids). In other embodiments, the first polypeptide segment includes less

than full-length mature gD but includes 250, 260, 270, 280, 290, 300, 305, 310, 315, 320, 325, 330, 335, 340, 345, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 360, 361, 362, 363, 362, 365, 366, 367 or 368 amino acids of the mature gD sequence.

In certain aspects, an antigenic amino acid sequence is inserted within a region of a gD that is between amino acids 230 and 300, between amino acids 235 and 295, or between amino acids 240 and 290 of a mature gD amino acid sequence. In other aspects an antigenic amino acid sequence is inserted at a position carboxy-terminal to amino acid 230, 235, 240, 245, 250, 255, 260, 265, 270, 275, 280, 285 of a mature gD amino acid sequence. In certain aspects, an antigenic amino acid sequence is inserted immediately adjacent to amino acid 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 282, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, or 300 of a mature gD amino acid sequence.

In certain embodiments, a chimeric gD protein of the invention has a structure that is similar to the structure of the wild-type protein, that is, the chimeric gD protein has the ability to interact with HVEM and, in addition, may have one or more of the following activities: 1) stimulating a CD8<sup>+</sup> T cell response to the fusion partner; 2) disrupting an HVEM-BTLA pathway activity; 3) interacting with nectin-1; 4) mediating cell entry by an HSV-1 and/or HSV-2 virus; and/or 5) mediating cell-to-cell spread of HSV-1 and/or HSV-2.

#### Antigens

As used herein, the term "antigen" (also termed "fusion partner" in relation to gD) is intended to include, but is not limited to, a substance that an immune response is specifically mounted against, such as a protein or a polypeptide. An antigen of the invention can be of any length, ranging in size from a few amino acids in length to hundreds of amino acids in length, provided that the chimeric gD maintains a structure that is similar to that of the wild-type gD, e.g., the chimeric gD retains one or more activities of the wild-type gD, particularly the activities of interacting with HVEM and/or stimulating a CD8<sup>+</sup> T cell response to the fusion partner in the host. In certain aspects of the invention, the antigen is 1000, 900, 800, 700, 600, 500, 400, 350, 340, 330, 320, 310, 300, 290, 280, 270, 260, 250, 240, 230, 220, 210, 200, 190, 180, 170, 160, 150, 140, 130, 120, 110, 100, 90, 80, 70, 60, 50, 40, 30, 20, 10 or less amino acids in length.

An antigen of the present invention includes a heterologous protein and/or polypeptide, such as a viral, bacterial, fungal, or parasite protein or polypeptide, and/or a host polypeptide and/or protein e.g., a tumor cell polypeptide or protein (such as an oncoprotein or a portion thereof) or a polypeptide or protein associated with inflammation. Antigens of particular interest include, but are not limited to, influenza virus antigens, such as nucleoprotein P (NP; see FIG. 16), matrixprotein (M), and hemagglutinin (HA); *Plasmodium* antigens such as thrombospondin-related anonymous protein (TRAP; see FIG. 17), ring-infected erythrocyte surface antigen (RESA), merozoite surface protein 1 (MSP1), merozoite surface protein 2 (MSP2), merozoite surface protein 3 (MSP3), and glutamate-rich antigen (GLURP); human papilloma virus (HPV) antigens, particularly HPV-16 antigens, such as E5 protein, E6 protein, and E7 protein; and HIV antigens, such as gag, pol, nef, tet, and env.

Viruses include DNA or RNA animal virus. As used herein, RNA viruses include, but are not limited to, virus families such as picornaviridae (e.g., polioviruses), reoviridae (e.g., rotaviruses), togaviridae (e.g., encephalitis viruses, yellow fever virus, rubella virus), orthomyxoviridae (e.g., influenza viruses), paramyxoviridae (e.g., respiratory syncytial virus (RSV), measles virus (MV), mumps virus (MuV), parainfluenza virus (PIV)), rhabdoviridae (e.g., rabies virus (RV)), coronaviridae, bunyaviridae, flaviviridae (e.g., hepatitis C virus (HCV)), filoviridae, arenaviridae, bunyaviridae, and retroviridae (e.g., human T-cell lymphotropic viruses (HTLV), human immunodeficiency viruses (HIV)). As used herein, DNA viruses include, but are not limited to, virus families such as papovaviridae (e.g., papilloma viruses), adenoviridae (e.g., adenovirus), herpesviridae (e.g., herpes simplex viruses, e.g., HSV-1, HSV-2; varicella zoster virus (VZV); Epstein-Barr virus (EBV); cytomegalovirus (CMV); human herpesviruses, e.g., HHV-6 and HHV-7; Kaposi's sarcoma-associated herpesvirus (KSHV) and the like), and poxviridae (e.g., variola viruses). These and other viruses and viral proteins are included in the present invention and are described further in Knipe et al., *Field's Virology*, 4<sup>th</sup> ed., Lippincott Williams & Wilkins, 2001, incorporated herein by reference in its entirety for all purposes.

Bacteria include, but are not limited to, gram positive bacteria, gram negative bacteria, acid-fast bacteria and the like. As used herein, gram positive bacteria include, but are not limited to, Actinomycetaceae, *Actinomyces israelii*, *Bacillus anthracis*, *Bacillus cereus*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium*, *Enterococcus faecalis*, *Listeria monocytogenes*, *Nocardia*, *Propionibacterium acnes*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Streptococcus mutans*, *Streptococcus pneumoniae* and the like. As used herein, gram negative bacteria include, but are not limited to, *Afipia felis*, *Bacteroides*, *Bartonella bacilliformis*, *Bordetella pertussis*, *Borrelia burgdorferi*, *Borrelia recurrentis*, *Brucella*, *Calymmatobacterium granulomatis*, *Campylobacter*, *Escherichia coli*, *Francisella tularensis*, *Gardnerella vaginalis*, *Haemophilus aegyptius*, *Haemophilus ducreyi*, *Haemophilus influenzae*, *Helicobacter pylori*, *Legionella pneumophila*, *Leptospira interrogans*, *Neisseria meningitidis*, *Porphyromonas gingivalis*, *Providencia sturtii*, *Pseudomonas aeruginosa*, *Salmonella enteridis*, *Salmonella typhi*, *Serratia marcescens*, *Shigella boydii*, *Streptococcus moniliformis*, *Streptococcus pyogenes*, *Treponema pallidum*, *Vibrio cholerae*, *Yersinia enterocolitica*, *Yersinia pestis* and the like. As used herein, acid-fast bacteria include, but are not limited to, *Mycobacterium avium*, *Mycobacterium leprae*, *Mycobacterium tuberculosis* and the like.

Other bacteria not falling into the other three categories include, but are not limited to, *Bartonella henseiae*, *Chlamydia psittaci*, *Chlamydia trachomatis*, *Coxiella burnetii*, *Mycoplasma pneumoniae*, *Rickettsia akari*, *Rickettsia prowazekii*, *Rickettsia rickettsii*, *Rickettsia tsutsugamushi*, *Rickettsia typhi*, *Ureaplasma urealyticum*, *Diplococcus pneumoniae*, *Ehrlichia chafensis*, *Enterococcus faecium*, *Meningococci* and the like.

Fungi include, but are not limited to, Aspergilli, Candidae, *Candida albicans*, *Coccidioides immitis*, Cryptococci, and combinations thereof.

Parasites include, but are not limited to, *Balantidium coli*, *Cryptosporidium parvum*, *Cyclospora cayetanensis*, *Encephalitozoa*, *Entamoeba histolytica*, *Enterocytozoon*

## 11

*bieneusi*, *Giardia lamblia*, *Leishmaniae*, *Plasmodii*, *Toxoplasma gondii*, *Trypanosoma*, trapezoidal amoeba and the like.

Oncoproteins are intended, without limitation, to refer to proteins and/or peptides that are capable of inducing cell transformation. Oncoproteins include, but are not limited to, cellular proteins such as PDGF, ERB-B, ERB-B2, K-RAS, N-RAS, C-MYC, N-MYC, L-MYC, BCL-2, BCL-1, MDM2 and the like. Oncoproteins also include, but are not limited to, viral proteins from RNA and/or DNA tumor viruses such as hepatitis B viruses, SV40 viruses, polyomaviruses, adenoviruses, herpes viruses, retroviruses and the like. Tumor suppressor proteins are intended, without limitation, to refer to proteins or polypeptides that can suppress or block aberrant cellular proliferation, as well as tumor suppressor proteins that have been mutated and, accordingly, no longer suppress or block aberrant cellular proliferation. Tumor suppressor proteins include, but are not limited to, cellular proteins such as APC, DPC4, NF-1, NF-2, MTS1, RB, p53 and the like.

gD chimeric proteins of the present invention are useful for modulating disorders associated with aberrant cellular proliferation mediated by oncoproteins and/or tumor suppressor proteins, such as cancer. Aberrant cellular proliferation is intended to include, but is not limited to, inhibition of proliferation including rapid proliferation. As used herein, the term "disorder associated with aberrant cellular proliferation" includes, but is not limited to, disorders characterized by undesirable or inappropriate proliferation of one or more subset(s) of cells in a multicellular organism. The term "cancer" refers to various types of malignant neoplasms, most of which can invade surrounding tissues, and may metastasize to different sites (*PDR Medical Dictionary* 1st edition (1995)). The terms "neoplasm" and "tumor" refer to an abnormal tissue that grows by cellular proliferation more rapidly than normal and continues to grow after the stimuli that initiated proliferation is removed (*PDR Medical Dictionary* 1st edition (1995)). Such abnormal tissue shows partial or complete lack of structural organization and functional coordination with the normal tissue which may be either benign (benign tumor) or malignant (malignant tumor).

Polypeptides and proteins associated with inflammation include those that modulate a disease or disorder characterized by, caused by, resulting from, or becoming affected by inflammation. Examples of inflammatory diseases or disorders include, but not limited to, acute and chronic inflammation disorders such as asthma, psoriasis, rheumatoid arthritis, osteoarthritis, psoriatic arthritis, inflammatory bowel disease (Crohn's disease, ulcerative colitis), sepsis, vasculitis, and bursitis; autoimmune diseases such as lupus, polymyalgia, rheumatica, scleroderma, Wegener's granulomatosis, temporal arteritis, cryoglobulinemia, and multiple sclerosis; transplant rejection; reperfusion injury in strokes or myocardial infarction; osteoporosis; cancer, including solid tumors (e.g., lung, CNS, colon, kidney, and pancreas); Alzheimer's disease; atherosclerosis; viral (e.g., HIV or influenza) infections; chronic viral (e.g., Epstein-Barr, cytomegalovirus, herpes simplex virus) infection; and ataxia telangiectasia.

## Nucleic Acid Molecules

The invention also provides nucleic acid molecules which encode fusion proteins of the invention. In certain embodiments of the invention, the nucleic acid molecule is a vector. As used herein, the term "vector" refers to a nucleic acid molecule, a protein, or a liquid structure capable of transporting another nucleic acid. One type of vector is a "plas-

## 12

mid," which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated.

Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors are replication-defective and remain in the nucleus as episomes. Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors." In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. The terms "plasmid" and "vector" can be used interchangeably as the plasmid is a commonly used form of vector.

In one embodiment, a recombinant virus is provided for eliciting an immune response in a host infected by the virus. In certain aspects, the recombinant virus is replication-incompetent. A recombinant virus may be constructed from any virus using methods known in the art, provided that the native progenitor is rendered replication incompetent. For example, replication-incompetent adenovirus, adeno-associated virus, SV40 virus, retrovirus, herpes simplex virus or vaccinia virus may be used to generate the recombinant virus by inserting the viral antigen into a region that is non-essential to the infectivity of the recombinant virus. A recombinant virus does not have the pathologic regions of the native progenitor of the benign virus but retains its infectivity to the host.

In a certain embodiment, the recombinant virus is a replication-incompetent chimpanzee-derived adenovirus. Chimpanzee-derived adenovirus vectors have distinct advantages over previously used adenoviral recombinants of the human serotype 5 that is typically used in the art. Most importantly, the efficacy of simian adenoviral vaccine carriers is not impaired by pre-existing neutralizing antibodies to human adenovirus serotype 5 that can be detected in up to 45% of the adult human population in the United States. Furthermore, simian adenoviral recombinants have interactions with cells of the innate immune system, most notably dendritic cells, which sponsor development of strongly biased Th1 responses suited to induce potent responses of CD8<sup>+</sup> T cells, a subset of immunocytes that is particularly important to control the spread of HIV-1. For a review of replication-incompetent chimpanzee-derived adenovirus, see U.S. Pat. No. 6,019,978, incorporated herein by reference in its entirety for all purposes.

A number of viral vectors suitable for in vivo expression of the gD chimeric proteins described herein are known. Such vectors include retroviral vectors (see, e.g., Miller (1992) *Curr. Top. Microbiol. Immunol.* 158:1; Salmons and Gunzburg (1993) *Human Gene Therapy* 4:129; Miller et al. (1994) *Meth. Enz.* 217:581) and adeno-associated vectors (reviewed in Carter (1992) *Curr. Opinion Biotech.* 3:533; Muzyczka (1992) *Curr. Top. Microbiol. Immunol.* 158:97). Other viral vectors that are used include adenoviral vectors, alphavirus replicons, herpes virus vectors, pox virus vectors, and rhabdovirus vectors, as generally described in, e.g., Jolly (1994) *Cancer Gene Therapy* 1:51; Latchman (1994) *Molec. Biotechnol.* 2:179; Johanning et al. (1995) *Nucl. Acids Res.* 23:1495; Berencsi et al. (2001) *J. Infect. Dis.* 183:1171; Rosenwirth et al. (2001) *Vaccine* February 19:1661; Kittlesen et al. (2000) *J. Immunol.* 164:4204;

## 13

Brown et al. (2000) *Gene Ther.* 7:1680; Kanessa-thasan et al. (2000) *Vaccine* 19:483; and Sten (2000) *Drug* 60:249. Compositions comprising vectors and an acceptable excipient are provided herein.

## Nucleic Acid and Protein Variants

In certain aspects, gD nucleic acid molecules and polypeptides are "naturally occurring." As used herein, a "naturally-occurring" molecule refers to a gD molecule having a nucleotide sequence that occurs in nature (e.g., encodes a gD polypeptide sequence found in a herpes simplex virus, e.g., HSV-1 or HSV-2). In addition, naturally or non-naturally occurring variants of these polypeptides and nucleic acid molecules which retain the same functional activity, e.g., the ability to bind HVEM and/or stimulate an immune response to a fusion partner in a host. Such variants can be made, e.g., by mutation using techniques that are known in the art. Alternatively, variants can be chemically synthesized.

As used herein, the term "variant" is intended to include, but is not limited to, nucleic acid molecules or polypeptides that differ in sequence from a reference nucleic acid molecule or polypeptide, but retains its essential properties, that is, it retains the ability to interact with HVEM and, in addition, it may have one or more of the following activities: 1) stimulating a CD8<sup>+</sup> T cell response to a fusion partner 2) disrupting an HVEM-BTLA pathway activity; 3) interacting with nectin-1; 4) mediating cell entry by an HSV-1 and/or HSV-2 virus; and/or 5) mediating cell-to-cell spread of HSV-1 and/or HSV-2. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference nucleic acid molecule. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence.

Variants can be made using mutagenesis techniques that are known in the art. Alternatively, variants can be chemically synthesized. Mutations can include one or more point mutations, deletions and/or insertions. In certain aspects of the invention, a mutant gD chimeric polypeptide has the ability to bind HVEM at a level that is the same as or greater than the ability of a wild-type gD protein to bind HVEM. In certain aspects of the invention, amino acid W 294 of the mature gD sequence is mutated to alanine.

## Construction of Fusion Proteins

Fusion proteins of the invention typically are prepared recombinantly, as described in Example 1. Many kits for constructing fusion proteins are available from companies such as Promega Corporation (Madison, WI), Stratagene (La Jolla, CA), CLONTECH (Mountain View, CA), Santa Cruz Biotechnology (Santa Cruz, CA), MBL International Corporation (MIC; Watertown, MA), and Quantum Biotechnologies (Montreal, Canada; 1-888-DNA-KITS). Alternatively, a fusion protein can be synthesized chemically, for example, using solid phase techniques. See, e.g., Merrifield, *J. Am. Chem. Soc.* 85, 2149-54, 1963; Roberge et al., *Science* 269, 202-04, 1995. Protein synthesis can be performed using manual techniques or by automation. Automated synthesis can be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Optionally, fragments of a fusion protein can be separately synthesized and combined using chemical methods to produce a full-length molecule.

## Methods of Using Fusion Proteins and Nucleic Acids of the Invention Pharmaceutical Compositions

## Dosage Regimens

Certain embodiments of the invention are directed to prophylactically treating an individual in need thereof. As used herein, the term "prophylactic treatment" includes, but

## 14

is not limited to, the administration of a nucleic acid sequence encoding a gD chimeric protein to a subject who does not display signs or symptoms of a disease, pathology, or medical disorder, or displays only early signs or symptoms of a disease, pathology, or disorder, such that treatment is administered for the purpose of diminishing, preventing, or decreasing the risk of developing the disease, pathology, or medical disorder. A prophylactic treatment functions as a preventative treatment against a disease or disorder.

10 Certain embodiments of the invention are directed to therapeutically treating an individual in need thereof. As used herein, the term "therapeutically" includes, but is not limited to, the administration of a nucleic acid sequence encoding a gD chimeric protein to a subject who displays 15 symptoms or signs of pathology, disease, or disorder, in which treatment is administered to the subject for the purpose of diminishing or eliminating those signs or symptoms of pathology, disease, or disorder.

10 Embodiments of the present invention are directed to 20 compositions and methods for enhancing the immune response of a subject to one or more antigens. As used herein, the terms "subject" and "host" are intended to include living organisms such as mammals. Examples of subjects or hosts include, but are not limited to, horses, 25 cows, sheep, pigs, goats, dogs, cats, rabbits, guinea pigs, rats, mice, gerbils, non-human primates, humans and the like, non-mammals, including, e.g., non-mammalian vertebrates, such as birds (e.g., chickens or ducks) fish or frogs (e.g., *Xenopus*), and a non-mammalian invertebrates, as well 30 as transgenic species thereof.

As used herein, the term "immune response" is intended to include, but is not limited to, T and/or B cell responses, that is, cellular and/or humoral immune responses. In one embodiment, the claimed methods can be used to stimulate 35 cytotoxic T cell responses. The claimed methods can be used to stimulate both primary and secondary immune responses. The immune response of a subject can be determined by, for example, assaying antibody production, immune cell proliferation, the release of cytokines, the expression of cell surface markers, cytotoxicity, and the like. In certain 40 aspects, the claimed gD chimeric proteins increase the immune response in a subject when compared to the immune response by an untreated subject or a subject who receives a vaccine containing the same antigen but without 45 gD.

As used herein, the term "enhancing an immune response" includes increasing T and/or B cell responses, that is, cellular and/or humoral immune responses, by treatment 50 of a subject using the claimed gD chimeric proteins and/or methods. In one embodiment, the claimed gD chimeric proteins and/or methods can be used to enhance cytotoxic T cell responses to the antigen (fusion partner). In another embodiment, the claimed compounds and methods can be used to inhibit the ability of the HVEM to interact with B 55 and T lymphocyte attenuator (BTLA), thus enhancing an immune response to the antigen (fusion partner). In another embodiment, the claimed gD chimeric protein interacts with HVEM.

As used herein, the term "immune cell" is intended to 60 include, but is not limited to, cells that are of hematopoietic origin and play a role in an immune response. Immune cells include, but are not limited to, lymphocytes, such as B cells and T cells; natural killer cells; myeloid cells, such as monocytes, macrophages, eosinophils, mast cells, basophils, and granulocytes.

As used herein, the term "adjuvant" includes, but is not limited to, agents which potentiate the immune response to

## 15

an antigen. Adjuvants can be administered in conjunction with a nucleic acid sequence encoding a gD chimeric protein of the invention to additionally augment the immune response.

Nucleic acid sequences encoding the gD chimeric proteins described herein can be administered to subjects in whom it is desirable to promote an immune response. In one embodiment, a nucleic acid sequence encoding a gD chimeric protein described herein is administered prophylactically, e.g., prior to infection with a pathogen or to a subject who is free of cancer or free of an autoimmune disease. In another embodiment, a nucleic acid sequence encoding a gD chimeric protein described herein is administered therapeutically, e.g., to a subjects who has a preexisting condition, e.g., a subject who is infected with a pathogen, who has cancer, or who suffers from an autoimmune disease.

In one embodiment, the gD chimeric protein is administered by "genetic immunization." In this embodiment, a DNA expression vector encoding the gD chimeric protein is injected into the subject animal, e.g., into the skin or into a muscle of the subject. The gene products are correctly synthesized, glycosylated, folded, and expressed by the subject to elicit the desired immune response. In one embodiment, DNA is injected into muscles or delivered into the skin coated onto gold microparticles by a particle bombardment device, a "gene gun." Genetic immunization has been shown to induce specific humoral responses and cellular immune responses (See, e.g., Mori et al. (1995) *J. Immunol.* 155:2039; Xu and Liew (1995) *Immunology* 84:173; Davis et al. (1994) *Vaccine* 12:1503).

A dosage regimen of administration of a gD chimeric protein or a nucleic acid sequence encoding a gD chimeric protein may be adjusted to provide the optimum therapeutic response for each subject without undue experimentation. For example, antibody titers to an antigen or cellular immune responses to an antigen can be measured to determine whether or not the subject is developing an immune response or is manifesting an enhanced immune response to the antigen and the dosage regimen can be adjusted accordingly.

The composition including a gD chimeric protein or a nucleic acid sequence encoding a gD chimeric protein may also be administered parenterally or intraperitoneally. The agent can be administered, for example, intranasally, orally, intravenously, intramuscularly, subcutaneously or mucosally. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms.

Pharmaceutical compositions suitable for injection include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. A pharmaceutical composition of the invention can be formulated to be suitable for a particular route of administration. For example, in various embodiments, a pharmaceutical composition of the invention can be suitable for injection, inhalation or insufflation (either through the mouth or the nose), or for intranasal, mucosal, oral, buccal, parenteral, rectal, intramuscular, intravenous, intraperitoneal, and subcutaneous delivery.

The composition including a gD chimeric protein or a nucleic acid sequence encoding a gD chimeric protein will be sterile. In addition, it will be stable under the conditions of manufacture and storage and preserved against the contaminating action of microorganisms such as bacteria and

## 16

fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be desirable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile, injectable solutions can be prepared by incorporating a gD chimeric protein or a nucleic acid sequence encoding a gD chimeric protein in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filter sterilization. Generally, dispersions are prepared by incorporating the gD chimeric protein or the nucleic acid sequence (with or without a carrier) encoding the gD chimeric protein into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient (e.g., agent or composition) plus any additional desired ingredient from a previously sterile-filtered solution thereof. The agent or composition can be administered in a form suitable for use with a needle-less injector device (such devices are known in the art (see, e.g., U.S. Pat. Nos. 5,383,851; 5,581,198; 5,846,233) for example as described in *Mol. Med.* (1998) 4:109.

When the composition including a gD chimeric protein or a nucleic acid sequence encoding a gD chimeric protein is suitably protected, as described above, the composition may be orally administered, for example, with an inert diluent or an assimilable edible carrier. As used herein "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the composition including a gD chimeric protein or a nucleic acid sequence encoding a gD chimeric protein, use thereof in the therapeutic compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active agent or composition for the treatment of individuals.

The composition including a gD chimeric protein or a nucleic acid sequence encoding a gD chimeric protein of the invention is administered to subjects in a biologically compatible form suitable for pharmaceutical administration in vivo to enhance immune responses. By "biologically compatible form suitable for administration in vivo" is meant a form of the gD chimeric protein or a nucleic acid sequence encoding a gD chimeric protein to be administered in which any toxic effects are outweighed by the therapeutic effects of the agent.

Administration of a therapeutically or prophylactically active amount of the compositions of a gD chimeric protein or a nucleic acid sequence encoding a gD chimeric protein is defined as an amount effective, at dosages and for periods of time necessary to achieve the desired result. The administration of a gD chimeric protein or a nucleic acid sequence encoding a gD chimeric protein can result in an enhanced immune response (e.g., a stimulation of CD8<sup>+</sup> T cells) to an antigen (e.g., a viral or a tumor cell antigen).

As defined herein, a therapeutically or prophylactically effective amount of a composition of a gD chimeric protein or a nucleic acid sequence encoding a gD chimeric protein (an effective dosage) ranges from about 0.001 to 30 mg/kg body weight, about 0.01 to 25 mg/kg body weight, about 0.1 to 20 mg/kg body weight, or from about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight. The skilled artisan will appreciate that certain factors may influence the dosage required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of an inhibitor can include a single treatment or can include a series of treatments. It will also be appreciated that the effective dosage of inhibitor used for treatment may increase or decrease over the course of a particular treatment. Changes in dosage may result from the results of diagnostic assays as described herein.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of cell biology, cell culture, molecular biology, microbiology, recombinant DNA and immunology, which are within the skill of the art. Such techniques are explained fully in the literature. See, for example, *Molecular Cloning A Laboratory Manual*, 2nd Ed., ed. by Sambrook, J. et al. (Cold Spring Harbor Laboratory Press (1989)); *Short Protocols in Molecular Biology*, 3rd Ed., ed. by Ausubel, F. et al. (Wiley, NY (1995)); *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., 1998; *DNA Cloning*, Volumes I and II (D. N. Glover ed., 1985); Oligonucleotide Synthesis (M. J. Gait ed. (1984)); Mullis et al. U.S. Pat. No. 4,683,195; Nucleic Acid Hybridization (B. D. Hames & S. J. Higgins eds. (1984)); the treatise, *Methods In Enzymology* (Academic Press, Inc., N.Y.); *Immunochemical Methods in Cell and Molecular Biology* (Mayer and Walker, eds., Academic Press, London (1987)); *Handbook of Experimental Immunology*, Volumes I IV (D. M. Weir and C. C. Blackwell, eds. (1986)); and Miller, J. *Experiments in Molecular Genetics* (Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1972)).

The following examples are set forth as being representative of the present invention. These examples are not to be construed as limiting the scope of the invention as these and other equivalent embodiments will be apparent in view of the present disclosure, figures, tables, and accompanying claims. The contents of all references, patents and published

patent applications cited throughout this application are hereby incorporated by reference in their entirety for all purposes.

5 Example 1

Materials and Methods

Construction of HSV-1 gD Fused Genes

A number of DNA and adenovirus vector vaccines were 10 constructed and tested (Table 1). The chimeric gene gDE7E6E5 was constructed based on the fusion of the HPV-16 E7, E6 and E5 oncoproteins and the HSV-1 gD protein. Although in this example the HPV proteins are in the order E7, E6, and E5, they can be used in any order. The 15 E7, E6 and E5 genes, without their respective stop codons, were amplified by PCR using the HPV-16 complete genome as a template. The gD gene is from HSV-1. E7, E6, and E5 genes are from HPV-16. Gag is a codon-optimized truncated 20 form of gag from HIV-1 clade B. SgD is a mutated form (W294A) of HSV-1 gD, which shows high affinity to HVEM (Krummenacher et al., 2005). NBGF is a mutated HSV-1 gD, which contains mutations that has been described to prevent 25 gD-HVEM interaction (Connelly et al., 2003); see SEQ ID NO:37. pRE4 was provided by Drs. Gary Cohen and Roselyn Eisenberg (Cohen et al., 1988). AdC68gag was previously described by Fitzgerald and collaborators (2003).

TABLE 1

List of vaccine vectors used.		
Vector name	Genes encoded	Vaccine carrier
pRE4 (pgD)	gD	DNA vaccine
pE7E6E5	E7, E6, and E5	DNA vaccine
pgDE7E6E5	gD, E7, E6, and E5	DNA vaccine
pgDE7	gD and E7	DNA vaccine
pSgDE7	SgD (gDW294A) and E7	DNA vaccine
pSgDE7E6E5	SgD, E7, E6, and E5	DNA vaccine
pNBGFgDE7E6E5	NBGFgD, E7, E6 and E5	DNA vaccine
pgag	gag	DNA vaccine
pgDgag	gD and gag	DNA vaccine
40 AdC68gD	gD	E1-deleted adenovirus vector, chimpanzee serotype 68
AdC68E7	E7	E1-deleted adenovirus vector, chimpanzee serotype 68
45 AdC68E7E6E5	E7, E6, and E5	E1-deleted adenovirus vector, chimpanzee serotype 68
AdC68gDE7E6E5	gD, E7, E6, and E5	E1-deleted adenovirus vector, chimpanzee serotype 68
50 AdC68gag	gag	E1-deleted adenovirus vector, chimpanzee serotype 68
AdC68gDgag	gD and gag	E1-deleted adenovirus vector, chimpanzee serotype 68

55 Separate amplification reactions were carried out with the following primers: E7FwApal and E7RvNarI, E6FwNarI and E6RvNotI, and E5FwNotI and E5RvApal (Table 2). The DNA fragment of the E7 gene was cleaved with Apal and NotI. The E6 DNA fragment was cleaved with NotI and NarI, and the E5 was cleaved with NarI and Apal. All DNA fragments were cloned into the Apal site in the pRE4 vector, provided by Drs. Gary Cohen and Roselyn Eisenberg (University of Pennsylvania, USA) (Cohen et al., 1988). The correct in-frame cloning of E7-, E6- and E5-encoding genes was confirmed after nucleotide sequencing at Wistar Sequencing Facility. Control vectors pE7E6E5 and

AdC68E7E6E5 were generated by PCR using pgDE7E6E5 as template and primers E7FwHindIII and E5RvHindIII (Table 2). AdC68gD control vector was generated using pRE4 as template and primers gDFwXbaI and gDRvXbaI (Table 2).

designed to mutate the amino acid residue 294 of gD, were used to PCR amplify the entire pgDE7 vector. The reaction products were then treated with DpnI and used to transform DH5a *E. coli* cells. The NBEFgDE7E6E5 gene (see SEQ ID NO:37) was generated by mutation of residues crucial for

TABLE 2

List of primers used	
Primer	Sequence (5'-3')
E7FwApal	GCTGTAGGGCCCCATGGAGATAACACCTAC (SEQ ID NO: 1)
E7RvNarI	CATGGTGGCGCCCTGGTTCTGAGAACAG (SEQ ID NO: 2)
E6FwNarI	AGACATGGCGCCCACCAAAAGAGAACTGC (SEQ ID NO: 3)
E6RvNotI	CTCCATCGGGCCGCCAGCTGGGTTCTCTACG (SEQ ID NO: 4)
E5FwNotI	GACAAAGCGGCCCTGCATCCACAAACATTAC (SEQ ID NO: 5)
E5RvApal	ACATATGGGCCCTGTAATTAAAAGCGTGC (SEQ ID NO: 6)
E7FwHindIII	GGGTGGAAGCTTATGGGAGATAACACCTAC (SEQ ID NO: 7)
E5RvHindIII	TGGGGCAAGCTTTAATTAAAAGCGTGC (SEQ ID NO: 8)
gDFwXbaI	CCCTAGTCTAGAATGGGGGGCTGCCGCC (SEQ ID NO: 9)
gDRvXbaI	CCCTAGTCTAGACTAGTAAACAAAGGGCTGGTG (SEQ ID NO: 10)
gagFwNarI	AAGAAGGGCGCCGGTGCAGAGCGTCAG (SEQ ID NO: 11)
gagRvNarI	AAGGGTGGCGCCAAAACTCTGCCTTATGGC (SEQ ID NO: 12)
gDFwHindIII	AAGCCCAAGCTTATGGGGGGCTGCCGCC (SEQ ID NO: 13)
gDRvHindIII	AAGCCCAAGCTCTAGTAAACAAAGGGCTGGTG (SEQ ID NO: 14)
NBEFgDRv	GACCGGAAGGTCTTGCCCGAAAGCGAGCGGGCTGGCCCTTGAG (SEQ ID NO: 15)
NBEFgDFw	CGCTTTCGCGCAAAGACCTTCCGGTCGCGACGCGCGGCC (SEQ ID NO: 16)
SgDFw	CAAATCCAACAAAACGCGCACATAGGCTCGATCC (SEQ ID NO: 17)
SgDRv	GATCGACGGTATGTGCGCGTTGGTGGGATTG (SEQ ID NO: 18)

To construct the AdC68gDE7E6E5 vector, the gDE7E6E5 chimeric gene was amplified by PCR using the pgDE7E6E5 vector as a template. The PCR reaction was carried out with gDFwXbaI and gDRvXbaI primers (Table 2). The DNA fragment of the gDE7E6E5 chimeric gene was cleaved with XbaI and cloned into XbaI site on the shuttle vector (BD PharMingen, San Diego, CA). The pShuttlegDE7E6E5 clone was confirmed by restriction analysis and sub-cloned into El-deleted chimpanzee-derived adenovirus vector serotype 68 using PI-SceI and I-CeuI sites as described (Fitzgerald et al. 2003).

The gDgag chimeric gene was generated by insertion of the codon-optimized truncated form of gag from HIV-1 clade B into the HSV-1 gD NarI site. The gag gene was amplified by PCR using the pCMVgag vector as a template and primers gagFwNarI and gagRvNarI (Table 2). The DNA fragment corresponding to gag gene was cleaved with NarI, cloned into pShuttlegD, and then sub-cloned into AdC68 vector as described above.

#### Construction of gD Mutants

The SgDE7 mutated gene was constructed using QUICK-CHANGE® site-directed mutagenesis kit (Stratagene Cloning Systems, La Jolla, CA) as recommended by the manufacturer. Briefly, SgDFw and SgDRv primers (Table 2),

HVEM-gD interaction. HSV-1 gD residues 11, 15, 25, 27, 45 28, 29, and 30, were mutated to alanine by gene splicing by overlap extension (i.e., gene SOEing). Briefly, two PCR reactions were carried out using set of primers, (i) with gDFwHindIII and NBEFgDRv, and (ii) with NBEFgDFw and gDRvHindIII (Table 2). Vector pgDE7E6E5 was used as a template in both PCR reactions. Two amplified fragments were used as template to PCR reaction with gDFwHindIII and gDRvHindIII primers (Table 2). The NBEFgDE7E6E5 DNA fragment was cloned into the same pgDE7E6E5 backbone vector. Both mutant gD sequences were confirmed by sequencing the entire gene at Wistar Sequencing Facility. DNA Vaccine and El-Deleted Chimpanzee-Derived Adenovirus Purification

DNA vaccines were propagated in *E. coli* K12 DH5α cells in LB medium supplemented with ampicillin and purified with the Maxi Prep Kit (QIAGEN®, Valencia, CA). The DNA concentration was determined by spectrophotometry at 260 nm and confirmed by visual inspection of ethidium bromide-stained 1% agarose gels in comparison to DNA fragments of known concentration (Invitrogen, Carlsbad, CA). Plasmids were kept at -20° C. until use, at which time the DNA concentration was adjusted to 1 µg/µl in phosphate-buffered saline (PBS). AdC68 vectors were propa-

gated using El-transfected HEK 293 cells and purified by CsCl gradient centrifugation as previously described (Fitzgerald et al., 2003). Upon purification, the concentration of each virus vector batch was determined by measuring virus particles (vp) by spectrophotometry at 260 nm.

#### Cell Lines

TC-1 tumor cells, derived from C57BL/6 origin lung epithelial cells transformed with v-Ha-ras and HPV-16 E6 and E7 genes, were provided by Dr. T. C. Wu, Johns Hopkins University, USA (Lin et al., 1996). Mouse melanoma cells B78H1 and B78H1/3E5, which express HVEM fused to EGFP, were provided by Drs. Gary Cohen and Roselyn Eisenberg (University of Pennsylvania, USA). El-transfected HEK 293 cells were used to propagate El-deleted chimpanzee derived adenovirus vectors. All cells were propagated in DMEM supplemented with glutamine, sodium pyruvate, nonessential amino acids, HEPES buffer, antibiotic, and 10% FBS (TC-1, CHO/CAR and El-transfected HEK 293 cells) or 5% FBS (B78H1 and B78H1/3E5 cells). CHO cells stably transfected to express the coxsackie adenovirus receptor (CHO/CAR) were obtained from J. Bergelson (Childrens Hospital of Philadelphia).

#### Animals and Immunization

Female BALB/c and C57Bl/6 mice at 6-8 weeks of age were purchased from Jackson Laboratory (Bar Harbor, ME) or Charles River Laboratories (Boston, MA) and housed at the Animal Facility of the Wistar Institute. All procedures involving handling and sacrifice of animals were performed according to approved protocols in accordance with recommendations for the proper use and care of laboratory animals at the Wistar Institute. Groups of 5 to 10 BALB/c and C57Bl/6 mice were intramuscularly (i.m.) vaccinated with the DNA vaccines or El-deleted chimpanzee derived adenovirus vectors into the tibialis anterior muscle of each hind limb. DNA vaccine was given at 100 µg divided in two 50 µl aliquots. El-deleted chimpanzee-derived adenovirus vectors (AdC68) were inoculated from  $1\times 10^8$  to at  $5\times 10^{10}$  vp per mouse. For most experiments AdC68 vectors were inoculated at  $10^{10}$  viral particles per mouse.

#### E7 Transgenic Mouse

The E7 transgenic mouse was based on a similar mouse where E7 is expressed in the thymus under the control of the thyroglobulin promoter (Ledent et al., Oncogene 1995; 10:1789-97). The plasmid used to create the E7 mouse was a very generous gift from Dr. Catherine Ledent, Université Libre de Bruxelles. The plasmid, constructed in the pSG5 vector, contained the bovine thyroglobulin promoter, a rabbit β-intron, the E7 gene, and a polyadenylation signal in a gene cassette. The bovine thyroglobulin promoter was used as it has been shown to be tightly regulated and expressed in thyrocytes (Ledent et al., *Proc Natl Acad Sci USA* 87:6176, 1990). The rabbit β-intron was used to increase the expression of the transgenes (Palmiter et al., *Proc Natl Acad Sci USA* 88:478, 1991). This cassette was removed and purified by gel electrophoresis followed by a Geneclean kit (Q-BioGen, Morgan Irvine, CA). DNA was concentrated by ethanol precipitation. The cut DNA was then microinjected by the University of Pennsylvania School of Medicine Transgenic Facility under the direction of Dr. Jean Richa. The founder mouse strain was C57BL/6. The founder mice were thus mated to wild type C57BL/6, and the progeny back crossed and screened by the ΔCT real time PCR method for homozygosity. The E7 tg mice were bred at the Animal Facility of the Wistar Institute from a pair provided by Dr. Y. Patterson (University of Pennsylvania). All animal procedures were performed in accordance with recommendations for the proper use and care of laboratory animals at the Wistar

Institute. Groups of 3 to 10 animals were vaccinated i.m. with the El-deleted AdC68 vectors into the tibialis anterior muscle of each hind limb. AdC68 vectors were inoculated at  $5\times 10^{10}$  vp per mouse unless stated otherwise.

#### 5 Intracellular Cytokine Staining

Intracellular IFN-γ staining was performed using peripheral blood mononuclear cells (PBMC) and cells from the spleen two weeks after the DNA vaccine dose or 10 days after El-deleted chimpanzee-derived adenoviral vector 10 administration unless stated otherwise. Samples were washed twice with L-15 medium. Cells were then treated for 5 minutes on ice with ACK lysis buffer (Invitrogen) to rupture red blood cells, washed, and suspended in DMEM supplement with 2% FBS. Samples were cultured at a 15 concentration of  $10^6$  cells/well for 5 hours at 37° C. in a 96-well round bottom microtiter plate (Costar) in 200 µl of DMEM supplemented with 2% FBS and  $10^{-6}$  M 2-mercaptoethanol. Brefeldin A (GolgiPlug; BD PharMingen) was added at 1 µl/ml. The E7-specific RAHYNIVTF (SEQ ID NO:19) peptide, which carries the immunodominant epitope 20 of E7 for mice of the H-2b haplotype, or the AMQMLKETI (SEQ ID NO:20) peptide, which carries the immunodominant MHC class I epitope of gag for mice of the H-2d haplotype, were used for peptide stimulation at a concentration of 3 µg/ml. The V3 control peptide delineated from the sequence of the envelope protein of HIV-1 clade B (VVEDEGCTNLSGF; SEQ ID NO:21) and the SIINFEKL peptide (SEQ ID NO:30) were used as control peptides. After washing, cells were incubated for 30 min at 4° C. with 25 100 µl of a 1:100 dilution of a fluorescein (FITC)-conjugated monoclonal antibody to mouse CD8a (BD PharMingen). Cells were washed once with PBS followed by permeabilization with Cytofix/Cytoperm (BD PharMingen) for 20 min at 4° C., washed twice with Perm/Wash buffer (BD 30 PharMingen) and incubated in the same buffer for 30 min at 4° C. with 50 50 µl of a 1:100 dilution of a phycoerythrin (PE)-labeled monoclonal antibody to mouse IFN-γ (BD PharMingen). After washing, cells were suspended in PBS and examined by two-color flow cytometry using an EPICS 35 Elite XL (Beckman Coulter). Data were analyzed by 40 WinMDi software. The percentages of antigen specific CD8+ T cells that stained positive for IFN-γ over all CD8+ T cells were determined.

#### TC-1 Challenge

C57Bl/6 Mice were challenged subcutaneously (s.c.) with 45  $1\times 10^5$  TC-1 cells suspended in 100 µl of serum-free media, and injected at one rear flank. To determine the protection of pre-challenge vaccination mice were challenged 10 and 14 days after vaccination with DNA vaccine or El-deleted chimpanzee derived adenovirus vector, respectively. Post-challenge vaccination was evaluated with mice challenged five days before vaccination (Example 3 and 15). Tumor growth in pre- and post-challenge vaccinated mice was monitored by visual inspection and palpation three times a 50 week. Animals were scored as tumor-bearing when tumors attained sizes of approximately 1-2 mm in diameter. Mice were euthanized once tumors exceeded a diameter of 1 cm. Tumor growth was followed for a period of 60 days after the challenge.

#### 60 Statistical Analysis

Experiments were conducted using 3-10 mice per group. Samples tested by ELISA were assayed in triplicates. Results show the means±standard deviation (SD). Intracellular cytokine staining was conducted with PBMC from individual mice, while tetramer and markers staining were performed with pooled samples. Significances between two groups were analyzed by one-tailed student's t-test.

## HVEM Binding Assay

CHO/CAR cells were infected with either AdC68gD, AdC68gag, or AdC68gDgag. After 72 hrs, cells were harvested, suspend in 1 ml of extraction buffer (10 mM Tris, 150 mM NaCl, 10 mM EDTA, 1% NP-40, 0.5% sodium deoxycholate, 1 mM PMSF [pH 8.0]) supplement with Complete Protease inhibitor (Roche, Basel, Switzerland), and incubated at 4° C. for 1 hour. After spinning at 12,000 g for 15 min at 4° C., protein extracts were kept at -80° C. until use. A capture enzyme-linked immunosorbent assay (ELISA) was used to normalize the amount of gD in the extracts. ELISA plates were coated with 50 µl of a 10 µg/ml concentration of ID3 monoclonal antibody (MAb) diluted in PBS/well. After an overnight incubation at 4° C., plates were exposed to blocking solution for 1 hour and then to extracts diluted in blocking solution for 2 hours at room temperature. Captured gD was detected by adding 50 µl of a 1 µg/ml concentration of Pab R7/well followed by goat anti-rabbit antibody coupled to horseradish peroxidase. Plates were rinsed with 20 mM citrate buffer (pH 4.5), ABTS peroxidase substrate was added, and the absorbance at 405 nm was recorded by using a microtiter plate reader. The level of gD in each extract was normalized by dilution in extraction buffer. To assess receptor binding of the gD mutants, ELISA plates were coated overnight with 50 µl of human-HVEM (5 µg/ml), exposed to blocking solution, and incubated with normalized cell extracts diluted in blocking solution for 2 hours at room temperature. Bound gD was detected as described above. soluble gD306 (Nicola et al., *J. Virol.* 71, 2940-46, 1997; Sisk et al., *J. Virol.* 68, 766-75, 1994) and gD285 (Whitbeck et al., *J. Virol.* 71, 6083-93, 1997), purified as described previously, were used at 1 µM.

## Detection of Transcripts by Real-Time PCR

Total RNA was isolated from CHO/CAR cells 48 hrs after infection with the Ultraspec RNA solution system kit (Biotex). The mRNA was reversed transcribed in vitro using MEGAscript transcription kit (Ambion). Remaining DNA was removed by treatment with DNase I (Ambion) for 1 h at 37° C. Quantification of the housekeeping gene glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was performed to normalize the amount of cDNA in each sample. Normalized cDNA samples were used for amplification with E5-, E6-, E7- and gD-specific primers. Quantitative real-time PCR reactions were conducted using the LightCycler-RNA amplification kit SYBR Green 1 (Boehringer Mannheim), following manufacturer's conditions. Samples were quantified in triplicate by three independent experiments.

## Confocal Microscopy and FACS Analysis

B78H1/3E5 cells, which express HVEM fused to Enhanced Green Fluorescence Protein (HVEM-EGFP), or B78H1 cells (both obtained from Drs. Gary Cohen and Roselyn Eisenberg, University of Pennsylvania) were infected with either AdC68gD, AdC68gag, or AdC68gDgag. CHO/CAR cells were infected with AdC68gD, AdC68E7E6E5, or AdC68gD-E7E6E5. After 48 hours, cells were directly stained or permeabilized with Cytofix/Cytoperm (BD PharMingen) and then stained with anti-gD MAb (DL-6) followed by anti-mouse IgG conjugated with Texas Red or PE. Confocal microscopy was performed with a Leica TCS SP2 Confocal Microscope at 400× final magnification. Cell suspensions were analyzed using an EPICS XL (Beckman-Coulter, Inc., Miami, FL) to determine presence of gDgag. Data were analyzed by Flowjo software (Tree Star, Inc.).

## Molecular Modeling of gD-Gag

The 3-dimensional models of the gD-gag were constructed with the MODELLER package<sup>44,45</sup> by combining

the structures of individual protein domains as determined by X-ray crystallography. The receptor bound form of gD-gag model was based upon the HSV-1 gD HVEM complex (1JMA)<sup>12</sup>, chain A, residues 1:259; SIV gag (1ECW)<sup>46</sup>, residues 1:119; and HIV-1 gag (1E6J)<sup>47</sup>, chain P, residues 11:220. The gD-gag unligated form was based upon the Cyclophilin A/HIV-1Chimera Complex (1M9D)<sup>48</sup>, chain A, residues 1:15; HSV-1 gD (2C36)<sup>15</sup>, chain A, residues 23:256; SIV gag (1ECW), residues 1:119; and HIV-1 gag (1E6J), chain P, residues 11:220. Ribbon representations were prepared within the Swiss-PdbViewer program<sup>49</sup> and rendered with the Persistence of Vision Ray Tracer program (POV-Ray 2004, version 3.6).

## In Vitro T Cell Proliferation Assay

Cells were harvested from draining popliteal lymph node of naïve and mice i.m. immunized 24 hrs earlier with 10<sup>11</sup> vps of AdC68gD or AdC68E7E6E5 then irradiated with 2000 RADs. CD8<sup>+</sup> cells were isolated from the spleens of OT-1 mice by negative selection using magnetic beads (Miltenyi Biotec) and labeled with 204 CFSE (Molecular Probes). A total of 1×10<sup>6</sup> irradiated lymph node cells were cultured with 1×10<sup>5</sup> CD8<sup>+</sup> CF SE-labeled OT-1 cells in presence of either SIINFEKL peptide or control peptide AMQMLKETI (both at 10<sup>-8</sup>M, Alpha Diagnostic International) in 96-well plate wells for 72 hours. Cells were stained with anti-CD8-PerCP and anti-Vα2-PE (both BD Pharmingen) for 30 min on ice. Cells were examined on a FACSCalibur using CellQuest software (BD Biosciences Pharmingen) and were analyzed using FlowJo software version 7.1.2 (TriStar, Inc.).

## ELISA for Antibodies to Gag

Sera from the vaccinated or naïve mice were tested on plates coated with purified gag protein. Briefly, 96-well round-bottom Maxisorb (Nunc) plate wells were coated overnight with 0.2 µg of gag p24 HIV-1 (Immuno Diagnostics, Inc.) diluted in 100 µl of coating buffer (15 mM Na<sub>2</sub>CO<sub>3</sub>, 35 mM NaHCO<sub>3</sub> and 3 mM NaN<sub>3</sub>, pH 9.6). The next day plates were blocked with 200 µl of PBS containing 3% BSA for 2 hours. Serum samples were serially diluted in PBS supplemented with 3% BSA and in triplicates at 10 µl/well on the gag-coated plates for 1 h at room temperature. Plates were washed, and a 1:200 dilution of alkaline phosphatase-conjugated goat anti-mouse IgGs (Cappel) was added to each well for 1 h at room temperature. After washing, plates were incubated with substrate (10 mg d-nitrophenyl phosphate disodium dissolved in 10 ml of 1 mM MgCl<sub>2</sub>, 3 mM NaN<sub>3</sub>, and 0.9 M diethanolamine, pH 9.8), and then read in an automated ELISA reader at 405 nm (model EL311, Bio-Tek Instruments).

## Isolation of Lymphocytes

Peripheral blood mononuclear cells (PBMC), spleen and livers were harvested as described (Lasaro et al., *Microbes Infect* 2005; 7:1541-50; Lin et al., *Cancer Res* 1996; 56:21-6). Tumor-infiltrating lymphocytes (TILs) were isolated from matrigel-tumors 3 or 10 days after challenge as described (22). TILs from the thyroid of E7-tg mice were harvested upon treatment of thyroid tissue fragments with 2 mg/ml collagenase P (Roche) and 1 mg/ml DNase I (Invitrogen). After 1 hour, the thyroid was homogenized and filtrated through a 70-mm cell strainer. Cells were washed with RPMI (Cellgro) media supplemented with 5% FBS, treated for 5 min on ice with Ack lysing buffer (Invitrogen) to rupture red blood cells, then suspended in 7 mL 40% percoll (Amersham Bioscience, Piscataway, NJ), and applied on top of 3 mL 70% percoll. After centrifugation at 2200 rpm for 20 min at room temperature, the cells at the interface were harvested and resuspended in media.

## Tetramer and Lymphocyte Markers Staining

Antigen-specific CD8<sup>+</sup> T cells were detected by APC-labeled MHC class I tetramers carrying the AMQMLKETI peptide (SEQ ID NO:20) of gag or the RAHYNIVTF peptide of E7 (E7tet; SEQ ID NO:19) (MHC Tetramer Core Facility, Emory University Vaccine Center, Atlanta, GA). PBMC and splenocytes isolated 10 days after immunization with AdC68 vectors were treated as described for intracellular cytokine staining. Samples were stained for 30 minutes at room temperature with gag-tet and anti-CD8a-PerCP in combination with the following antibodies: CD25-PE, CD122-PE, CD127-PE, CD27-PE, BTLA-PE (eBioscience), PD1-PE, CD62L-FITC, CD69-FITC, CD103-FITC, CD43-FITC, CD44-FITC and CD54-FITC (all from BD Biosciences Pharmingen, unless indicated otherwise). For Bcl2 and CTLA-4 staining, cells were washed, permeabilized for 30 min at 4° C. with Cytofix/Cytoperm (BD Biosciences Pharmingen), and then stained with antibodies Bcl2-PE or CTLA-PE (BD Biosciences Pharmingen).

Lymphocytes were stained for 30 minutes at room temperature with E7-tet and anti-CD8a-PerCP together with the following antibodies: BTLA-PE (eBioscience), CD44-FITC, CD62L-PE, CD27-PE, CD127-PE, CD122-PE and PD1-PE (BD Biosciences Pharmingen). For Bcl2 and CTLA-4 staining, cells were washed, permeabilized for 30 min at 4° C. with Cytofix/Cytoperm (BD Biosciences Pharmingen), and then stained with antibodies Bcl2-PE or CTLA-PE (BD Biosciences Pharmingen). Flow cytometry analyses were performed with at least 100,000 viable cells live-gated on FACSCalibur using CellQuest software (BD Biosciences Pharmingen) and were analyzed using FlowJo software version 7.1.2 (TriStar, Inc.).

## Example 2

## Fusion Protein Constructs

The HPV-16/HSV-1 glycoprotein D (gD) chimeric gene, called gDE7E6E5, was composed of the complete open reading frame of gD, which had incorporated into the Apal site a fusion gene composed of HPV-16 E5, E6 and E7 genes without respective start and stop codons (FIG. 1). The HIV-1/gD chimeric gene, named gDgag, was composed of the complete open reading frame of gD which had incorporated into the Nan site a codon-optimized truncated form of gag HIV-1 clade B (FIG. 2).

## Example 3

## Effects of Fusion Protein Constructs on Host Immune System

Mice were immunized with 100 µg/mouse of DNA vaccines or 5×10<sup>10</sup> virus particles/mouse of El-deleted adenovirus vectors. After 14 days (DNA vaccines) or 10 days (El-deleted adenovirus vaccines) peripheral blood mononuclear cells (PBMC) were stained for CD8 and IFN-γ by intracellular cytokine staining. Percentage represents number of CD8<sup>+</sup>/IFN-γ<sup>+</sup> cells over total number of CD8<sup>+</sup> cells.

Pre-challenge vaccination: ten days after immunization mice were challenged with 5×10<sup>5</sup> TC-1 cells (10 mice per group), in vitro transformed syngeneic cells which express E7 and E6 and induce tumors in C57Bl/6 mice, and tumor growth was followed per 60 days. Percentage represents number of tumor-free mice over total mice challenged at day 60.

Post-challenge vaccination: Mice (10 per group) were vaccinated 5 days after challenged with 5×10<sup>5</sup> TC-1 cells. Tumor growth was followed for 60 days.

The results are shown in Table 3. Percentage represents number of tumor-free mice over total mice challenged at day 60. The fusion gene gDE7E6E5 expressed by a DNA vaccine (pgDE7E6E5) or an El-deleted chimpanzee-derived adenovirus vector (AdC68gDE7E6E5) induced high frequencies of E7-specific CD8<sup>+</sup> T cells and complete protection against tumor cell challenge after a single dose. However, neither E7-specific CD8<sup>+</sup> T cell responses nor protection to challenge were developed when gD was not present, indicating that this fusion had dramatically improved the efficacy of the vaccines.

TABLE 3

Immunization	E7-specific CD8 <sup>+</sup> T cell response <sup>b</sup>	Protection upon pre-challenge vaccination <sup>c</sup>	Protection upon post-challenge vaccination <sup>d</sup>
pRE4 (gD only)	0.11%	0%	0%
pE7E6E5	0.18%	0%	0%
pgDE7E6E5	5.53%	100%	70%
AdC68gD	0.05%	0%	0%
AdC68E7E6E5	0.05%	0%	0%
AdC68gDE7E6E5	21.94%	100%	100%

An El-deleted chimpanzee-derived adenovirus carrying a codon-optimized, truncated form of gag HIV-1 clade B (AdC68gag) has been shown to induce a strong specific CD8<sup>+</sup> T cell response (Fitzgerald et al. (2003) *J. Immunol.* 170:1416). However, gag-specific CD8<sup>+</sup> T cell frequencies induced by the DNA vaccine (pgDgag) and the El-deleted chimpanzee-derived adenovirus vector (AdC68gDgag) were higher when gag was fused into gD (FIGS. 3A-3B). The effect of gD fusion was more remarkable when lower amounts of adenovirus vector were used for vaccination (FIG. 3B).

Although chimpanzee-derived adenovirus vectors circumvented the expected negative effect of pre-existing immunity to common human serotypes of adenovirus, such as serotype 5 (AdHu5), the efficacy of AdC68 vaccination was decreased by approximately 50% when neutralizing antibodies to AdHu5 were present. Although not wishing to be bound by this explanation, this reduction likely is caused by pre-existing T cell that cross-react between adenovirus of the human serotype 5 and the chimpanzee adenoviruses. However, AdC68 vaccinations carrying antigens fused with HSV-1 gD were only weakly influenced by those antibodies (FIG. 4).

## Example 4

## Interaction of Constructs which Encode Fusion Proteins with HVEM

Efficacy of the vaccine was related to the localization of the transgene in relationship the N-terminus of gD. The specific region localized in the N-terminal portion of gD, which interacts with the herpes virus entry mediator (HVEM), was crucial in enhancing stimulation of specific CD8<sup>+</sup> T cell mediated immune response. A mutated form of gD chimeric protein gDE7E6E5, called NBEFgDE7E6E5, was generated altering specific gD amino acids (M11A, N15A, L25, Q27A, L28A, T29A, D30A), which promote gD-HVEM interaction. The mutated form of gDE7E6E5 expressed by DNA vaccine was not able to induce an E7-specific CD8<sup>+</sup> T cell response, although mutated and non-mutated gDE7E6E5 genes were transcribed at the same level (FIG. 5). On the other hand, a single-amino acid

modification on HSV-1 gD (W294A), was able to enhance the efficacy of a gD/E7 fusion DNA vaccine (FIG. 6).

The interaction of HSV-1 gD with HVEM interferes with down-regulation of immune responses associated with HVEM-BTLA (B and T lymphocyte attenuator) pathway. Without intending to be bound by theory, maintenance of gD interference in HVEM-BTLA pathway is an important key to enhance a specific T cell mediated immune response to one or more heterologous antigens. To this end, it was demonstrated that a fusion of an antigen, such as HIV-1 gag, into the C-terminal region of HSV-1 gD did not decrease its affinity to bind to HVEM (FIG. 7). The ability of HSV-1 gD to bind HVEM was not affected by fusion with HIV-1 gag. In fact, the affinity to HVEM of HIV-1 gag/HSV-1 gD chimeric protein was slight higher than non-fused gD.

Confocal microscopy and FACS analyses of cells infected in vitro with AdC68gDgag provided additional evidence that gDgag bound HVEM (FIGS. 8A-8B and FIG. 9). Confocal microscopy indicated that gDgag was co-localized with HVEM on the membrane (FIG. 8A) and inside (FIG. 8B) cells infected with AdC68gDgag. Also, cell surface gD staining on AdC68gDgag-infected cells demonstrated that the amount of gD was greater on cells that expressed HVEM than on cells that did not express HVEM (FIG. 9). These results indicate that gDgag bound HVEM, that part of gDgag expressed by the infected cell bound HVEM inside the cell, and that the complex of gDgag-HVEM was transported from the inside the cell to the cell surface. In addition, confocal microscopy and gD staining indicate that gDgag encoded by AdC68gDgag-infected cell was able to bind to HVEM expressed on the surface of other non-infected cells (FIGS. 10A-10B).

Human derived serotype 2 adeno-associated virus (AAV) vectors encoding the gDgag chimeric protein induced higher CD8+ T cell response against gag than AAV vector encoding only gag (FIG. 14). In addition, immunization with AAVgD-gag enhanced CD8+ T cell response against AAV capsid.

#### Example 5

##### CD8<sup>+</sup> T Cell Response Against NP

The chimeric protein gDNP is composed of the complete open reading frame of HSV-1 gD, into which we incorporated into the ApaI site the nucleoprotein P (NP) gene from Influenza virus A/PR8 without respective start and stop codons (FIG. 15).

Mice were immunized with 100 µg of DNA vaccine vector pgDNP or pNP. Fourteen days after immunization peripheral blood mononuclear cells (PBMCs) were isolated and cell surface stained with the NP-tetramer and a labeled antibody to CD8. Naïve mice were used as negative control. Data are shown in FIG. 16 and represent percentages of NP-tetramer<sup>+</sup> CD8<sup>+</sup> T cells over all detected CD8<sup>+</sup> T cells. This experiment demonstrated that the gDNP chimeric protein expressed by DNA vaccine vector induced NP-specific CD8<sup>+</sup> T cell response in mice after a single immunization.

#### Example 6

##### Molecular Modeling

To examine how insertion of a foreign sequence within the C terminus may modify folding of gD we modeled the structure of unligated gD-gag (FIGS. 18A, 18B) and of gD-gag bound to HVEM (FIGS. 18C, 18D). The C-terminus of the native unligated gD structure is anchored near the N-terminal region and thus masks the HVEM binding site (Krummenacher et al., *EMBO J.* 24, 4144-53, 2005; FIG.

19). However, computational modeling of the gD-gag structure predicts that the C-terminus would be shifted away from the N-terminal portion without altering the core structure or the N terminus, which is required to form the HVEM binding site (FIGS. 18A, 18B). A superposition of X-ray structure of gD upon the gD-gag model indicates that insertion of gag into gD does not disrupt the integrity of the HVEM binding surface (FIGS. 18B, 18D).

10

#### Example 7

##### Cellular Localization of Chimeric gD Proteins

Binding of gD to HVEM presumably depends on secretion or cell surface expression of gD. To determine the cellular localization of the chimeric gD-gag protein, B78-H1/3E5 cells, which express human HVEM-EGFP on their surface, were infected with AdC68gag, AdC68gD or AdC68gD-gag, stained with a monoclonal antibody (MAb) to gD and analyzed by confocal microscopy (FIG. 19). Both gD and gD-gag co-localize with HVEM on the surface of infected cells as well as within infected cells. Similar results were obtained upon staining of cells with antibodies to gag. Absolute levels of gD-gag expressed on the surface of infected cells were below those of native gD (FIG. 20). Analyses of mRNA levels showed that all vectors transcribed the chimeric genes at equal amounts to their corresponding non-chimeric versions. Thus the lower levels of cell surface expression of the chimeric protein suggest inefficient secretion, rapid re-internalization of gD-gag or accelerated proteolytic degradation. The presence of HVEM on the cell surface increased the amount of gD-gag (FIG. 20) on the cell surface suggesting that intracellular binding of HVEM to gD-gag stabilized gD-gag or facilitated export of the chimeric protein.

When AdC68gD or AdC68gD-gag infected HVEM negative (HVEM<sup>-</sup>) cells were mixed with uninfected cells expressing HVEM (HVEM<sup>+</sup>), both gD and gD-gag co-localized with HVEM on the uninfected HVEM<sup>+</sup> cells indicating that some of the protein was released and then bound to HVEM. This was more pronounced with gD-gag than with gD, and may reflect either increased secretion of the chimeric protein or its superior binding to HVEM (FIG. 21). To ensure that the observed transfer of gD or gD-gag from HVEM<sup>-</sup> to HVEM<sup>+</sup> cells was not a result of cell fusion, cells were stained for an additional marker, the coxsackie adenovirus receptor (CAR), expressed only by the HVEM<sup>-</sup> cell line. Uninfected HVEM cells which co-stained for gD failed to express CAR thus ruling out cell fusion.

30

#### Example 8

##### Expression of Antigens within gD Augments Antigen-Specific CD8<sup>+</sup> T Cell Responses

To determine if expression of an antigen within gD affects stimulation of CD8<sup>+</sup> T cells in general to epitopes presented on gD transduced antigen presenting cells we initially performed proliferation assays. Irradiated cells from draining lymph nodes of mice intramuscularly (i.m.) immunized with AdC68 carrying either gD or E7E6E5 were pulsed with low amounts of the SIINFEKL peptide and served as antigen presenting cells for carboxyfluorescein diacetate succinimidyl ester (CFSE)-labeled CD8<sup>+</sup> T cells from OT-1 mice, which carry CD8<sup>+</sup> T cells with a transgenic receptor for SIINFEKL in the context of Kb. We had shown previously that i.m. application of an AdC68 vector causes an accumulation of vector transduced mature dendritic cells within draining lymph nodes<sup>26</sup>. We thus expected that upon appli-

cation of the AdC68gD vector some of the mature dendritic cells would express gD, which in turn may modulate the response of OT-I derived CD8<sup>+</sup> T cells to their cognate antigen. As shown in FIG. 22, OT-I CD8<sup>+</sup> T cells proliferated more vigorously upon co-culture with lymph node cells from AdC68gD injected mice than upon co-culture with cells from AdC68E7E6E5 injected mice.

To evaluate if blockade of the HVEM inhibitory pathway enhances adaptive immune responses *in vivo*, we vaccinated mice with DNA and AdC68 vectors expressing the gD-antigen chimeric proteins and tested T and B cell responses in comparison to those of mice injected with vectors expressing E7, E6 and E5 or gag without gD. Mice fail to mount detectable E7-specific CD8<sup>+</sup> T cell responses following vaccination with DNA or AdC68 vectors encoding either E7E6E5 (FIG. 23A) or E7 alone. In contrast, the DNA vaccine and the AdC68 vector expressing E7 or the fusion polypeptide of E7, E6 and E5 within gD induce robust E7-specific CD8<sup>+</sup> T cell responses. Similarly, the DNA vaccine expressing the gD-gag chimeric protein stimulate more potent gag-specific CD8<sup>+</sup> T cell responses compared to vectors expressing gag only (FIG. 23A). The AdC68 vector expressing the gD-gag chimeric protein also elicits stronger gag-specific CD8<sup>+</sup> T cells than those expressing gag only, which was especially pronounced at low vector doses (FIGS. 23A, 23B). To determine the longevity of vaccine-induced responses, E7-specific CD8<sup>+</sup> T cells were monitored for over a year following vaccination with the AdC68gD-E7E6E5 vectors. CD8<sup>+</sup> T cell responses were maintained at stable frequencies indicating that the enhancement of the initial primary T cell response resulted in an increase of the memory T cell population (FIG. 23C).

#### Example 9

##### Expression of Antigens within gD Increases Antigen-Specific Antibody Responses

It is well established that neutralizing antibodies are the primary correlate of vaccine-induced protection against most virus infections. LIGHT stimulation enhances both B cell proliferation and immunoglobulin production<sup>27</sup> and might balance inhibition exerted by ligation of BTLA to HVEM. To analyze whether antibody responses can be enhanced by the gD-antigen chimeric proteins, sera of mice immunized with the AdC68 vectors expressing gag or gD-gag were analyzed for antibodies to gag (Table 4).

TABLE 4

Gag-specific antibody response after immunization with AdC68 vectors expressing gD, gag, or gD-gag.

Immunization	Anti-gag antibody titer ± SD <sup>a</sup> (p value) <sup>b</sup>			
	10 days	7 weeks	20 weeks	49 weeks
AdC68gag	26 ± 8	35 ± 8	152 ± 19	6 ± 4
AdC68gD	3 ± 4 (0.007)	9 ± 4 (0.0037)	28 ± 15 (0.0004)	8 ± 2 (0.2573)
AdC68gD-gag	287 ± 7 (2.6E-7)	740 ± 2 (6.8E-9)	1257 ± 21 (4.8E-6)	294 ± 8 (3.3E-7)

<sup>a</sup>Anti-gag Ig titers were established as the reciprocal of the serum dilution that gave an absorbance twice above that of the sera from naïve mice. SD-standard deviation.

<sup>b</sup>p-values were determined using one-tailed Student t test comparing titers either from mice immunized with AdC68gD or AdC68gD-gag with titers from mice immunized with AdC68gag.

The AdC68gag vector induces only marginal levels of gag-specific antibodies while AdC68gD-gag vector elicits a significantly higher response that remained detectable for at least 11 months. Antibody responses to gD, measured in

parallel, were comparable in mice vaccinated with the AdC68gD or the AdC68gD-gag vector.

#### Example 10

##### gD-HVEM Interaction is Needed to Augment Immune Responses

To determine if enhancement of the CD8<sup>+</sup> T cell responses by expression of an antigen within gD requires binding of gD to HVEM, we constructed DNA vaccines expressing the E7E6E5 sequence within two modified versions of gD; In one construct, termed NBEFgD-E7E6E5, 7 amino acids at the N-terminus of gD, i.e., M11, N15, L25, Q27, L28, T29 and D30, were replaced with alanine residues. Previous studies have shown that these 7 amino acids on wild type gD are crucial to gD-HVEM interaction<sup>13</sup>. In a second construct, termed SgD-E7E6E5, the tryptophan (W) in position 294 of gD was changed to alanine (A). This modification, which is thought to destabilize the native conformation of the gD C-terminus, was shown to increase binding of gD306 to HVEM<sup>15</sup>. Mice immunized with one dose of a DNA vaccine expressing NBEFgD-E7E6E5 failed to mount detectable frequencies of E7-specific CD8<sup>+</sup> T cells (FIG. 24A). In contrast, frequencies of E7-specific IFN-γ<sup>+</sup>CD8<sup>+</sup> T cells induced by pSgD-E7E6E5 were higher compared to those induced by pgD-E7E6E5 (FIG. 24A), although this difference did not reach statistical significance. To confirm this observation we also modified the pgd-E7 DNA vaccine<sup>28</sup>, which carries only E7 inserted into gD, by changing the tryptophan in position 294 to alanine. This new vector, termed pSgD-E7, stimulated significantly higher frequencies of E7-specific CD8<sup>+</sup> T cells compared to a plasmid vector expressing E7 within the wild-type form of gD (FIG. 24B). Taken together, these results demonstrate that binding to HVEM is essential for the immunopotentiating effect of gD and that gD mutations that increase its binding to HVEM can under some conditions further augment CD8<sup>+</sup> T cell responses.

#### Example 11

##### Functionality of CD8<sup>+</sup> T Cells Induced by gD Chimeric Proteins

Modulation of regulatory pathways may affect the functionality of the resultant T cell responses. Phenotypes of vaccine-induced CD8<sup>+</sup> T cells were analyzed from mice immunized with AdC68 vectors expressing gD-gag or gag (FIG. 25). We measured expression of the following phe-

notypic markers, i.e., CD25, CD122, CD127, CD27, CD62L, CD69, CD103, CD43, CD44, CD54, Bcl2, BTLA, CTLA-4 and PD-1 on gag-specific CD8<sup>+</sup> T cells (FIG. 25). Most of the markers tested (CD122, CD127, CD27, CD62L,

31

CD69, CD103, CD43, CD44, CD54, Bcl2, BTLA, CTLA-4 and PD-1) are modulated on antigen-specific CD8<sup>+</sup> T cells as compared to naive CD8<sup>+</sup> T cells and for most markers expression levels on CD8<sup>+</sup> T cells induced by gag or gD-gag are identical. CD27 is increased on a subpopulation of gD-gag induced gag-specific CD8<sup>+</sup> T cells, while CTLA expression is marginally lower when compared to CD8<sup>+</sup> T cells induced by gag alone. Overall, although AdC68gD-gag elicits higher frequencies of gag-specific CD8<sup>+</sup> T cells than AdC68gag, the phenotypic profiles of the resultant effector cells are very similar. IFN- $\gamma$ <sup>+</sup>CD8<sup>+</sup> T cells induced by gD-gag or gag respond to the same epitope when tested against a panel of gag peptides, indicating that expression of gag within gD does not increase the breadth of the CD8<sup>+</sup> response.

T cell functionally was further accessed by testing if mice vaccinated with either DNA (FIG. 26A) or AdC68 vectors (FIG. 26B) expressing E7E6E5 with or without gD were protected against challenge with TC-1 cells, which are lung epithelial cells derived from C57Bl/6 mice that are transformed with v-Ha-ras and the E6 and E7 oncoproteins of HPV-16<sup>26</sup>. Animals immunized with pgD-E7E6E5 or AdC68gD-E7E6E5 were completely protected against TC-1 tumor progression, and protection was only seen upon vaccination with constructs carrying gD fusion proteins. Mice vaccinated with DNA vaccines expressing E7 or E7E6E5 within mutant forms of gD were also tested for protection against TC-1 tumor formation (FIG. 26C). Protection correlated with CD8<sup>+</sup> T cell responses; mice immunized with a vector expressing E7E6E5 within the form of gD that can not bind to HVEM were not protected while mice immunized with the same antigen expressed within the gDW294A variant were fully protected. Additionally, animals immunized once with a DNA vaccine expressing only E7 within gD developed tumors while those that expressed E7 within the gDW294A variant were completely protected.

#### Discussion of Examples 2-11

Incorporation of the antigens into the extracellular C-terminal domain of gD markedly increases vaccine-induced CD8<sup>+</sup> T and B cell responses. As demonstrated in the specific examples, above, this is especially pronounced for CD8<sup>+</sup> T cell responses to E7, which expresses a T cell epitope with low to moderate affinity to H-2 K<sup>b</sup> (He et al., *virol.* 270, 146-61, 2000). A single dose of DNA vaccines or AdC68 vectors expressing the E7E6E5 polypeptide fails to elicit detectable CD8<sup>+</sup> T cell responses or protective immunity to challenge with an E7-expressing tumor cell line, while the same antigen expressed within gD by either vector results in high and sustained frequencies of E7-specific CD8<sup>+</sup> T cells and complete protection against tumor cell challenge. Gag of HIV-1 carries a high affinity epitope for mice of the H-2<sup>d</sup> haplotype and DNA vaccines or AdC68 vectors (Fitzgerald et al., *J. Immunol.* 170, 1416-22, 2003) expressing gag induce readily detectable CD8<sup>+</sup> T cell responses. Responses to gag are also increased upon expression of gag within gD.

The immunopotentiating effect of gD on the response to gag is more impressive when the gD-gag chimeric protein is delivered by a DNA vaccine rather than by the highly immunogenic AdC68 vector. Upon dose reduction of the AdC68 vector, results clearly demonstrate that an induction of gag-specific CD8<sup>+</sup> T cells can be achieved with an approximately 100 fold lower vaccine dose than needed for the AdC68 vector expressing gag only. This is important because Ad vectors of the common human serotype 5

32

(AdHu5) when tested in clinical trials as vaccine carriers for antigens of HIV-1 encountered dose limiting toxicity (Kresge, *LAVI rep.* 9, 18-20, 2005). Reactogenicity at high doses in humans is also anticipated with the chimpanzee-origin Ad vectors such as AdC68, which was developed to circumvent the effect of neutralizing antibodies to common human serotypes of adenovirus. Thus, further improvement of the immunogenicity of Ad vector vaccines that allows for a substantial dose reduction while maintaining efficacy could lower vaccine-related side effects, reduce the overall cost of the vaccine and facilitate production for mass vaccination. Increased efficacy may also lessen the need for complex prime boost regimens that are currently being tested in clinical HIV-1 vaccine trials but that may be unmanageable and too costly for developing countries.

Insertion of antigens into gD does not affect the functionality of antigen-specific CD8<sup>+</sup> T cells; T cells induced by the gD chimeric proteins protect against tumor challenge in the E7 model, are phenotypically similar to those induced in absence of gD, and efficiently differentiate into memory cells as shown with gag-expressing AdC68 vaccines, confirming a previous study with BTLA-deficient T cells (Krieg et al., *Nat. Immunol.* 8, 162-71, 2007).

Antibody responses are also augmented by expressing the antigen within gD. LIGHT participates in B cell expansion as a co-stimulus of CD40 and induces antibody production (Duhen et al., *Eur. J. Immunol.* 34, 3534-41, 2004). The control of LIGHT-induced B cell activation appears to be provided by down-regulation of HVEM following its engagement by LIGHT (Duhen et al., 2004). BTLA is constitutively expressed on B cells and BTLA deficient mice mount higher antibody responses compared to wild-type mice although this effect was shown previously to be modest (Hurchla et al., *J. Immunol.* 174, 3377-85, 2005). In contrast, in our vaccine model, antibody responses were markedly enhanced upon expressing gag within gD. While not wishing to be bound by the explanation, this may reflect that binding of gD to HVEM augments CD4<sup>+</sup> T cell responses, which in turn promote activation of B cells. Alternative pathways such as an enhancement of the co-stimulatory HVEM-LIGHT pathway may also have contributed.

Binding of gD to HVEM is essential for augmentation of CD8<sup>+</sup> T cell responses to the fusion partner as vaccines expressing antigen within a modified gD in which the HVEM binding site had been obliterated fail to induce enhanced CD8<sup>+</sup> T cell responses. In the native unligated structure form of gD, the C-terminus largely obstructs movement of the N-terminus into its HVEM-binding conformation (Krummenacher et al., *EMBO J.* 24, 4144-53, 2005). According to our molecular model, insertion of foreign sequences of certain lengths such as gag or the E7E6E5 polypeptide may effect a structural change of gD which improves its binding to HVEM. This may not be achieved by short sequences such as E7 alone, for which the immunogenicity and efficacy of the gD chimeric vaccines can be improved further through a single amino acid exchange in position 294 of gD which had been described previously to improve binding of gD to HVEM (Krummenacher et al., 2005).

Manipulation or disruption of negative regulatory pathways has been shown previously to augment T cell responses. Recent reports showed that antibody-mediated interruption of the PD1-PD-L1 pathway reverses T cell exhaustion caused by chronic infections and allows for increased T cell proliferation (Barber et al., *Nature* 439, 682-87, 2006; Day et al., *Nature* 443, 350-54, 2006). Manipulation of this pathway may not readily affect primary

T cell responses as PD1 requires induction by activation and is not expressed on naïve T cells (Barber et al., 2006). In contrast, low levels of BTLA are expressed on naïve T cells; the levels rapidly increase upon T cell activation and then decline (Hurchla et al., 2005; Han et al., *J. Immunol.* 172, 5931-39, 2004).

Medicinal targeting of immunoregulatory pathways can result in immunopathology such as the devastating cytokine storms observed upon application of anti-CD28 antibodies to human volunteers (Suntharalingam et al., *NEJM* 355, 1018-28, 2006) or auto-immunity commonly seen in genetically modified mice (Watanabe et al., *Nat. Immunol.* 4, 670-79, 2003). The use of a gD-antigen chimeric protein to enhance the immunogenicity of vaccines has the advantage that it does not involve systemic interruption of an inhibitory pathway but rather exerts its effects locally to the site of antigen presentation. We confirmed the spatially limited effect of gD experimentally by injecting two Ad vectors expressing either gD or E7E6E5 into distant anatomical sites (left leg versus right leg) and failed to observe enhancement of antigen-specific CD8<sup>+</sup> T cell responses. It has been suggested previously that targeting of BTLA through inhibitory antibody or small molecules may enhance vaccine immunogenicity. Results shown here suggest that such novel adjuvants may indeed be useful; nevertheless, their effect, unlike that of gD chimeric antigens, would be systemic and thus carry a higher likelihood of unwanted side effects.

#### Example 12

##### In Vitro Characterization of the Vaccine Vectors

We compared levels of transgene in cells infected in vitro with AdC68 vectors. Total RNA isolated from CHO/CAR cells infected with 10<sup>9</sup> virus particles (vps) of AdC68gD, AdC68E7E6E5 and AdC68gD-E7E6E5 vectors was reversed transcribed and gD, E7, E6 and E5 specific mRNA were quantified by real-time PCR (FIG. 27A). To minimize inherent differences in RNA isolation procedures, mRNA of a housekeeping gene (GAPDH) was used to normalize the amount of cDNA in each sample. The amount of gD-specific mRNA transcripts in cells infected with AdC68gD-E7E6E5 was not statistically different from that in cells infected with AdC68gD. The mRNA levels of E7, E6 and E5 were also similar in samples infected with AdC68gD-E7E6E5 and AdC68E7E6E5. Protein expression in cells infected with AdC68 vectors was evaluated by immunofluorescence with a monoclonal antibody (MAb) against gD (FIG. 27B). CHO/CAR cells infected with AdC68gD expressed gD mainly on their surface although some of the protein could be detected within the cells. Cells infected with AdC68E7E6E5 did not stain with the gD-specific MAb. In cells infected with AdC68gD-E7E6E5, gD was mainly detected within the cells; levels of gD on the cell surface were markedly reduced when compared to those on cells infected with AdC68gD. The lower levels of cell surface expression of the chimeric protein might be due to inefficient secretion, rapid re-internalization of gD-E7E6E5 or accelerated proteolytic degradation.

#### Example 13

##### E7-Specific CD8<sup>+</sup> T Cell Responses Induced by AdC68 Vectors

The induction of immune responses by the gD-E7E6E5 expressed by AdC68 and DNA vectors were described above. Here, we confirmed our studies by testing splenocytes and PBMCs from mice immunized with either AdC68

or DNA by ICS for frequencies of CD8<sup>+</sup> T cells producing IFN-γ in response to a peptide expressing the immunodominant epitope of E7 (FIGS. 28A and 28B, respectively). We confirm that immunization with AdC68gD-E7E6E5 induced a potent E7-specific CD8<sup>+</sup> T cell response detectable from spleens and blood while no specific CD8<sup>+</sup> T cell response was found upon immunization with AdC68gD or AdC68E7E6E5. Also, DNA vaccines carrying either the E7E6E5 fusion protein alone or gD failed to induce CD8<sup>+</sup> T cells to E7, while such cells were elicited by the DNA vaccine expressing E7E6E5 within gD.

To investigate whether the CD8<sup>+</sup> T cell response induced by AdC68gD-E7E6E5 could be enhanced further, AdC68gD-E7E6E5 was tested in a prime and boost regimen with the DNA vaccine expressing the same transgene product (FIG. 28C). Groups of mice were primed with the DNA vaccine and then boosted with the AdC68 vector 90 days later. In other animals the order of the vaccines was reversed. T cell frequencies were tested from blood 14, 30 and 90 days after priming and then on days 7, 10 and 14 after the boost. After priming, the AdC68 vector induced higher frequencies of E7-specific CD8<sup>+</sup> T cells compared to the DNA vaccine. Upon booster immunization the AdC68 prime DNA boost regimen performed poorly, and although CD8<sup>+</sup> T cell frequencies increased they remained below the peak frequencies seen upon AdC68 priming. In contrast, in the group primed with the DNA vaccine, a boost with the AdC68 vector induced a pronounced increase in E7-specific CD8<sup>+</sup> T cell frequencies.

Once we defined the most efficient prime boost regimen, we tested whether the DNA vaccine or the AdC68E7E6E5 vector could prime or boost a CD8<sup>+</sup> T cell response to E7. Mice primed with the DNA vaccines carrying E7E6E5 with or without gD were boosted 90 days later with either AdC68E7E6E5 or AdC68gD-E7E6E5, respectively. See Table 5, below.

TABLE 5

AdC68 vectors boost after prime with DNA vaccines.			
Prime <sup>a</sup>	Percentage of IFN-γ CD8 <sup>+</sup> cells over total CD8 <sup>+</sup> cells <sup>b</sup>		
	No Boost	AdC68E7E6E5	AdC68gD-E7E6E5
No prime	0.1	0.1	21.9
pE7E6E5	0.1	0.1	21.2
pgD-E7E6E5	0.3	2.5	47.4

<sup>a</sup> Mice were primed i.m. immunized with 100 µg of DNA vaccine.

<sup>b</sup> Frequencies of E7-specific CD8<sup>+</sup> T cells over all CD8<sup>+</sup> cells were determined 10 days after boost.

<sup>c</sup> Mice were boosted with 5 × 10<sup>10</sup> vps of AdC68 vectors 60 days after prime.

Ten days after immunization with AdC68gD-E7E6E5, E7-specific CD8<sup>+</sup> T-cell frequencies were similar in mice primed with pE7E6E5 and unprimed control mice, indicating that the DNA vaccine expressing the oncoproteins of HPV-16 was non-immunogenic. Again, mice primed with pgDE7E6E5 and boosted with AdC68gD-E7E6E5 developed high frequencies of E7-specific IFN-γ-producing CD8<sup>+</sup> T cells. A low but significant increase in E7-specific CD8<sup>+</sup> T cells was observed in mice primed with the DNA vaccine expressing gD-E7E6E5 and then boosted with AdC68E7E6E5, indicating that within this more immunogenic vaccine vehicle, the oncoproteins could trigger expansion of a memory response.

To test whether the effect of gD required that the antigen was expressed within gD or if concomitant presence of gD and the antigen sufficed, we immunized mice with a mixture

35

of equal doses of AdC68gD and AdC68E7E6E5. This mixture failed to elicit a detectable E7-specific CD8<sup>+</sup> T cell response, suggesting that the immunopotentiating effect of gD requires that the antigen is expressed within gD or that the same cells that express gD have to express the antigen.

## Example 14

Dose Response Curve of the CD8<sup>+</sup> T Cell Response to AdC68gD-E7E6E5

In humans, Ad vectors cause dose limiting toxicity, as has been established for El-deleted AdHu5 vector tested as vaccine carriers for antigens of HIV-1 in clinical trials (Kresge, K. J et al., 2005). Although the AdC68 vector has not yet undergone clinical testing, we anticipate that this vector would also cause significant toxicity if used at high doses. We therefore tested the E7-specific CD8<sup>+</sup> T cell response elicited by varied doses of AdC68gD-E7E6E5 (FIG. 29A).

Mice were immunized i.m. with  $1 \times 10^8$  to  $5 \times 10^{10}$  vps of AdC68gD-E7E6E5. Frequencies of E7-specific CD8<sup>+</sup> T cells were measured 10 days later from blood and spleens. The specific CD8<sup>+</sup> T response declined with decreasing doses of adenovirus vector. A response could still be detected upon immunization with  $5 \times 10^8$  vps of the vaccine but then became undetectable at  $1 \times 10^8$  vps. Comparable results were obtained with other immunogens, as described above.

## Example 15

## AdC68gD-E7E6E5 Protects Against Challenge with an E7-Expressing Tumor Cell Line

To formally demonstrate efficacy of AdC68gD-E7E6E5 we conducted challenge experiments with the TC-1 cell line. To determine the effectiveness of the AdC68 vector-induced immune response in causing regression of already established tumors, groups of C57Bl/6 mice were first injected with TC-1 cells then vaccinated 5 days later with AdC68gD, AdC68E7E6E5, or AdC68gD-E7E6E5 (FIG. 30A). Mice immunized with AdC68gD-E7E6E5 rejected the tumors and remained tumor-free for at least 60 days after challenge. In contrast, mice immunized with AdC68gD and AdC68E7E6E5 showed progressive growth of the TC-1 tumors.

Induction of long-term memory by a therapeutic vaccine is advantageous, since longevity of T cells may prevent resurgence of virus-infected cells that escape the initial wave of the immune response. Recently, we showed that E7-specific CD8<sup>+</sup> T response in mice immunized with a single dose of AdC68gD-E7E6E5 was detected over a year after immunization. We challenged mice that had been vaccinated 1 year earlier with AdC68gD-E7E6E5, AdC68gD or AdC68E7E6E5 with TC-1 cells (FIG. 30B). Mice immunized with AdC68gD-E7E6E5 were protected against TC-1 challenge given one year later, while mice injected with AdC68gD or AdC68E7E6E5 developed tumors after challenge.

To determine if tumor cell challenge boosted the vaccine-induced E7-specific CD8<sup>+</sup> T cell response, we investigated E7-specific CD8<sup>+</sup> T cell response after challenge (FIGS. 31A-31B). Frequencies of E7-specific CD8<sup>+</sup> T cells as detected by E7/D<sup>b</sup> tetramers increased slightly in mice that were challenged compared to those that had only been vaccinated (FIGS. 31A and 31B).

36

## Example 16

Phenotypic Profile of E7-Specific CD8<sup>+</sup> T Cells

We analyzed CD8<sup>+</sup> T cells from blood, spleens and tumors of vaccinated mice. Mice were immunized with AdC68gD, AdC68E7E6E5 or AdC68gD-E7E6E5. They were challenged 10 days later with TC-1 cells in matrigel. Lymphocytes were isolated 3 days later from spleens, blood and tumors. A pronounced cellular infiltrate was seen in tumors from AdC68gD-E7E6E5 vaccinated animals while comparatively few cells could be isolated from tumors of the other groups. We therefore isolated cells from additional mice on day 10 after challenge. At this time point there was a pronounced infiltrate in the tumors of AdC68gD and AdC68E7E6E5 vaccinated mice. In mice vaccinated with AdC68gD-E7E6E5, tumors had resolved and only a few cells could be recovered. We therefore compared cells isolated from day 3 tumors from AdC68gD-E7E6E5 vaccinated mice with those isolated on day 10 from tumors of AdC68gD and AdC68E7E6E5 vaccinated mice. In addition we analyzed PBMCs and splenocytes harvested on the corresponding days.

Tumors from mice vaccinated with AdC68gD-E7E6E5 had frequencies of E7-specific CD8<sup>+</sup> T cells that exceeded those in spleens or blood of the same mice, indicating a rapid recruitment or retention of E7-specific CD8<sup>+</sup> T cells within the tumors. E7-specific CD8<sup>+</sup> T cells from spleens, blood and tumors were analyzed phenotypically for CD44, CD62L, CD27, Bcl2, BTLA, CTLA-4 and PD-1 in comparison to tetramer negative (tet-) T cells from naive mice or mice immunized with AdC68gD or AdC68E7E6E5. The phenotypic profiles of tet-T cells from either of these groups were identical. E7-specific CD8<sup>+</sup> T cells from spleens and blood up-regulated CD44 and down-regulated CD62L, CD27, Bcl2 and BTLA. There was no change of CTLA-4 or PD1 expression compared to tet-CD8<sup>+</sup> T cells. CD8<sup>+</sup> T cells isolated from tumors showed a distinct phenotype. Tee CD8<sup>+</sup> cells up-regulated CD44 and down-regulated CD62L, CD27 and BTLA. Unlike cells from blood and spleens, they failed to down-regulate Bcl2 and strongly upregulated CTLA-4 and PD1, two molecules involved in negative immunoregulation.

## Example 17

AdC68gD-E7E6E5 Induces an E7-Specific CD8<sup>+</sup> T Cell Response in E7-Transgenic Mice

Women with HPV-16-associated cancers are expected to respond poorly to E7 as a progressing tumor would impair the adaptive immune response directed against its antigen. To determine if AdC68gD-E7E6E5 induces an E7-specific CD8<sup>+</sup> T cell response in mice that are tolerant to E7 we tested transgenic (tg) mice that constitutively express E7 under a tissue-specific promoter in the thyroid. These mice develop with age large goiters and thyroid carcinomas. E7-tg mice as well as age-matched control mice were vaccinated at one year of age with  $5 \times 10^{10}$  vps of AdC68gD-E7E6E5 or as a control with AdC68gD or AdC68E7E6E5. Ten days later lymphocytes were isolated from blood and spleens.

E7-tg mice had markedly enlarged thyroids and lymphocytes were also isolated from their thyroids. The thyroids of wild-type mice were comparatively small and we were unsuccessful to obtain lymphocytes from them. T cells from individual animals were tested for frequencies of E7-specific CD8<sup>+</sup> T cells. Mice failed to develop E7-specific CD8<sup>+</sup> T cells upon vaccination with AdC68gD or AdC68E7E6E5 (FIG. 33A). Both E7-tg and wild-type mice developed vigorous responses in blood and spleens to E7 upon vaccination with AdC68gD-E7E6E5. Responses were slightly higher in samples from wild-type mice compared to those

from E7-tg mice. This difference was statistically not significant by ICS although it was significant upon analysis by tetramer staining. E7-specific CD8<sup>+</sup> T cells could also be detected at high frequencies in the thyroid of AdC68gD-E7E6E5-vaccinated E7-tg mice. E7-specific CD8<sup>+</sup> T cells isolated from blood and spleen of wild-type and E7-tg mice as well as from thyroids of E7-tg mice were analyzed for expression of phenotypic markers.

CD8<sup>+</sup> T cells from naïve E7-tg mice isolated from spleens, blood or thyroids were analyzed for comparison. In spleen and blood expression of CD62, CD127, Bcl2, BTLA, and CTLA-4 on E7-specific CD8<sup>+</sup> T cells from E6/E7-tg and wild-type mice were indistinguishable. CD27, CD44 and PD-1 were slightly higher on E7-specific CD8<sup>+</sup> T cells isolated from blood and to a lesser degree from spleens of E7-tg mice. E7-specific CD8<sup>+</sup> T cells isolated from the thyroids expressed markedly higher levels of CD44, CD27 and PD-1 in comparison to CD8<sup>+</sup> T cells isolated from the thyroids of naïve E7-tg mice. The other markers were similar to those seen on E7-specific T cells from blood and spleens. None of the CD8<sup>+</sup> T cells isolated from the thyroids of naïve E7-tg mice stained with the tetramer to E7, phenotypes of the vaccine-induced E7-specific CD8<sup>+</sup> T cells could thus not be compared to phenotypes of T cells induced by the transgenic E7 protein.

#### Discussion of Examples 12-17

A preventative vaccine for HPV-16 based on L1 virus like particles, which induces neutralizing antibodies, has recently been licensed by FDA and has been recommended for use in teenage girls and young women. Once an infection has occurred, however, neutralizing antibodies do not affect viral clearance nor do they inhibit development of malignancies. Consequently, women with humoral immunodeficiency do not have an increased susceptibility to cervical cancer, while women with cell-mediated immunodeficiency, such as HIV-1-infected patients, renal transplant patients or patients with genetic T cell deficiencies, have increased incidence rates (Moscicki et al., J Infect Dis 2004; 190:37-45; Matas et al., Lancet 1975; 1:883-6; Frisch et al., J Natl Cancer Inst 2000; 92:1500-10; Lowy et al., J Natl Cancer Inst 2003; 95:1648-50). This together with extensive studies in animal models and clinical trials (reviewed in Galloway, Lancet Infect Dis 2003; 3:469-75) implicates a crucial role for T cells in eliminating cells persistently infected with oncogenic types of HPVs.

The vaccine described here expressed three of the onco-proteins of HPV-16, i.e., E7, E6 and E5 to broaden responses in human outbred populations. Mice of the H-2<sup>b</sup> haplotype respond to a low affinity epitope of E7, but according to our results fail to develop CD8<sup>+</sup> T cells to E6. Our CD8<sup>+</sup> T cell analyses thus focused on responses to E7. To enhance immune responses we incorporated the oncoproteins into gD of HSV-1 which binds to the herpes virus entry mediator (HVEM) (Montgomery et al., Cell 1996; 87:427-36; Whittle et al., J Virol 1997; 71:6083-93). HVEM, which is expressed on dendritic cells, is a member of the tumor-necrosis factor receptor (TNFR) family and interacts with LIGHT (Marsters et al., J Biol Chem 1997; 272:14029-32; Granger & Rickert, Cytokine Growth Factor Rev 2003; 14:289-96) and lymphotoxin- $\alpha$  (LT- $\alpha$ ) (Mauri et al., Immunity 1998; 8:21-30; Sarrias et al., Mol Immunol 2000; 37:665-73). Also, HVEM binds to B and T lymphocyte attenuator (BTLA), a recently described member of the B7-family (Gonzalez et al., Proc Natl Acad Sci USA 2005; 102:1116-21; Sedy et al., Nat Immunol 2005; 6:90-8). The

HVEM-BTLA interaction inhibits T cell activation in vitro thus defining these molecules as part of an inhibitory pathway (Sedy et al., 2005). Expression of BTLA is upregulated on tumor-infiltrating T cells as was shown in cancer patients (Wang et al., Tissue Antigens 2007; 69:62-72). HSV-1 gD competes with BTLA for binding to HVEM (Compaan et al., J Biol Chem 2005; 280:39553-61) and would thus be expected to enhance activation of naïve T cells by blockade of this negative immunoregulatory pathway. Recently, we showed that viral antigens expressed within gD induced CD8<sup>+</sup> T and B cell responses to the antigens that are far more potent than those elicited by the same antigen expressed without gD.

Our data confirm a very strong increase of CD8<sup>+</sup> T cell responses to E7 expressed within HSV-1 gD. Responses to E7 expressed by the DNA vaccine or the AdC68 vector without gD were below the level of detection, while E7 expressed within gD induced frequencies of E7-specific CD8<sup>+</sup> T cells of 1-3% upon DNA vaccination and 10-24% upon Ad vector immunization. As expected, responses were markedly higher upon vaccination with the AdC68 vector than the DNA vaccine. Most of our studies therefore focused on the AdC68 vector vaccine. Mice vaccinated with AdC68gD-E7E6E5 were completely protected against challenge with the E7 and E6 expressing TC-1 tumor cells given shortly after vaccination or one year later. More importantly, mice with pre-existing TC-1 tumors rejected the tumors upon vaccination and then remained disease-free. Vaccine-induced E7-specific CD8<sup>+</sup> T cells rapidly enriched within the TC-1 tumors. Phenotypically E7-specific CD8<sup>+</sup> T cells isolated from TC-1 tumors upregulated CD27, CTLA-4 and PD-1 and down-regulated Bcl2, which may have been a consequence of the engagement of their receptors by the tumor cells. Expression of BTLA, which was previously reported to become upregulated on TILs cells in humans, was not increased on E7-specific CD8<sup>+</sup> T cells isolated from TC-1 tumors.

Numerous studies have shown efficacy of HPV-16 E7 vaccines against TC-1 tumors. Transplantable tumors grow very rapidly in mice and vaccines are thus applied before or shortly after challenge. At this early stage, T cells to the tumor antigens are probably not yet compromised. We therefore tested AdC68gD-E7E6E5 in E7-tg mice, which express the oncoprotein under a tissue specific promoter within their thyroid. *Listeria* based E7 vaccines have been tested in E6/E7-tg mice in which the transgenes were similarly expressed with a bovine thyroglobulin promoter (Souders et al., Cancer Immun 2007; 7:2). The *Listeria* vaccine was shown to induce lower frequencies of E7-specific CD8<sup>+</sup> T cells compared to wild-type mice and the average avidity of CD8<sup>+</sup> T cells that were induced was ten fold lower than those isolated from wild-type mice. Nevertheless the *Listeria* vaccine could eradicate 7 day established (5 mm) transplanted tumors in some E6/E7 transgenic mice, albeit at lower frequency than in wild-type mice. We tested AdC68gD-E7E6E5 in one-year-old E7-tg with thyroid hyperplasia. AdC68gD-E7E6E5 induced an E7-specific CD8<sup>+</sup> T cell response in the E7-tg mice that was only slightly below that induced in age-matched wild-type mice. The vaccine-induced E7-specific CD8<sup>+</sup> T cells infiltrated the thyroid. Phenotypically, the thyroid-infiltrating vaccine-induced E7-specific CD8<sup>+</sup> T cells in E7-tg mice showed only minor difference to those isolated from TC-1 tumors and these differences, i.e., an overall decrease in CD44, CD27, Bcl2, and CTLA4 may be a reflection of difference in sampling time rather than T cell functionality. Accordingly

39

E7-specific CD8<sup>+</sup> T cells isolated from blood or spleens of vaccinated wild-type or E7-tg mice showed virtually identical phenotypes.

Our studies show that E7 of HPV-16 expressed as a fusion protein together with E6 and E5 within gD induces a robust CD8<sup>+</sup> T cell response even in animals with developing E7-associated malignancies. Without further studies directly comparing the effect of similar numbers of E7-specific CD8<sup>+</sup> T cells induced by a vaccine expressing E7 without gD to those induced by a vaccine expressing E7 within gD we can only speculate that the high efficacy of the latter is linked to blockade of an immunoinhibitory pathway. The Listeria vector used previously induces frequencies of E7-specific CD8<sup>+</sup> T cells that in wild-type mice are comparable in magnitude to those induced by AdC68gD-E7E6E5. In E6/E7-tg mice the Listeria vector induced markedly lower frequencies of T cells (Souders et al., 2007), while the response to AdC68gD-E7E6E5 is only slightly reduced. Although it is tempting to speculate that this is caused by an immunopotentiating effect of gD, other differences in the vaccine delivery vehicles such as their interactions with antigen presenting cells, in addition to differences between the E6/E7 and E7 transgenic mice such as transgene copy number and E7 expression may also have contributed to these results.

Blockade of immunoinhibitory pathways that are upregulated in cancer patients such as regulatory T cells or PD-1 expression on T cells has been shown pre-clinically to increase T cell responses to tumor antigens. In these systems regulatory T cells were depleted or rendered dysfunctional by antibodies and using a similar approach, the PD-1 pathway was blocked by antibodies to PD-1 or its ligand. These interventions, which have shown promise in animal models, exert a global effect on the immune system, which poses the risk of augmenting auto-immune reactivity. In contrast, blockade of the BTLA-HVEM pathway only exerts a local effect on T cells that are being activated to the antigen expressed within gD and should thus not subject patients to unwanted immune responses.

#### Example 18

##### Immunization of Rhesus Macaques

Two groups of rhesus macaques (4 per group) are enrolled into the study. Animals are screened for antibodies to

40

AdC68. Only seronegative animals are used. Sera and peripheral PBMCs are harvested 4 and 2 weeks before vaccination and preserved to serve as controls.

Animals are vaccinated once with 500 µg of pgag (group 1) or pgD-gag (group 2). They are boosted 2 months later with 5x10<sup>10</sup> vps of purified and quality controlled AdC68gag (group 1) or AdC68gD-gag (group 2) vector given i.m. in saline. Animals are bled 2, 4, and 8 weeks after priming and 2, 6 and 12 weeks after booster vaccination. PBMCs are tested for T cell responses to a pool of gag peptides by ELISpot for IFN-γ and IL-2 and by ICS for CD3, CD8, CD4, IFN-γ and IL-2 as described (Reyes-Sandoval, et al. J. Virol. 78:7392-7399).

Sera are tested for antibodies to gag (ELISA) and neutralizing antibodies to the vaccine carrier. The experiments are controlled by samples collected prior to vaccination. Animals are euthanized ~4 months after the boost, and lymphocytes are isolated from various compartments (spleen, blood, lymph nodes, liver, intestine) and tested by ELISpot for T cell responses to gag including analyses for IFN-γ, IL-2, TNF-α and MIP-113. In addition they are analyzed by ICS for secretion of IFN-γ and expression of T cell markers (CD3, CD4, CD8, included into each panel), activation (CD69, CD25, CD95, CD71) and proliferation (Ki67) markers, markers that identify T cell subsets (naive, central memory T cells, effector memory T cells and effector T cells) (CD28, CD95, CD45RA, CD62L, CCR7, CD27, CD127); chemokine and homing receptors (CCR5, CCR9, CXCR4, C11a-c, a4b7, CD103, CD49d); and markers indicative for lytic potential (CD107, perforin, granzyme B).

Immunophenotypical studies are performed by multicolor (7-8 colors) flow cytometry on mononuclear cells. The analyses are restricted to compartments that allow for isolation of sufficient numbers of lymphocytes. The experiment is controlled using cryopreserved splenocytes from sham-vaccinated rhesus.

It is to be understood that the embodiments of the present invention which have been described are merely illustrative of some of the applications of the principles of the present invention. Numerous modifications may be made by those skilled in the art based upon the teachings presented herein without departing from the true spirit and scope of the invention.

---

#### SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 37

```

<210> SEQ ID NO 1
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 1

```

gctgttagggc cccatggaga tacacctac

29

```

<210> SEQ ID NO 2
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

```

&lt;400&gt; SEQUENCE: 2

catgggtggcg cctgggttct gagaacag

28

<210> SEQ ID NO 3  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: primer

&lt;400&gt; SEQUENCE: 3

agacatggcg cccaccaaaa gagaactgc

29

<210> SEQ ID NO 4  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: primer

&lt;400&gt; SEQUENCE: 4

ctccatgcgg ccgcccagct gggtttctct acg

33

<210> SEQ ID NO 5  
<211> LENGTH: 32  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: primer

&lt;400&gt; SEQUENCE: 5

gacaaagcgg ccgcctgcat ccacaacatt ac

32

<210> SEQ ID NO 6  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: primer

&lt;400&gt; SEQUENCE: 6

acatatggc cctgtaatta aaaagcgtgc

30

<210> SEQ ID NO 7  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: primer

&lt;400&gt; SEQUENCE: 7

gggtggaagc ttatggaga tacacctac

29

<210> SEQ ID NO 8  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: primer

&lt;400&gt; SEQUENCE: 8

tggggcaagc ttttaaatta aaaagcgtgc

30

<210> SEQ ID NO 9  
<211> LENGTH: 30

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 9
ccctagtcta gaatggggg ggctgccccc                                30

<210> SEQ ID NO 10
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 10
ccctagtcta gactagtaaa acaaggcgtg gtg                                33

<210> SEQ ID NO 11
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 11
aagaagggcg ccgggtgcgag agcgtcag                                28

<210> SEQ ID NO 12
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 12
aagggtggcg cccaaaactc ttgccttatgc                                32

<210> SEQ ID NO 13
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 13
aagcccaagc ttatgggggg ggctgccccc                                30

<210> SEQ ID NO 14
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 14
aagcccaagc ttcttagtaaa acaaggcgtg gtg                                33

<210> SEQ ID NO 15
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 15

```

## US 12,390,525 B2

**45**

-continued

gaccggaagg tctttgccgc gaaagcgago ggggtcgccc gcctttag

48

<210> SEQ ID NO 16  
<211> LENGTH: 47  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: primer

&lt;400&gt; SEQUENCE: 16

cgctttcgcg gcaaagacct tccggtcgag gacgccccgg ccgcccc

47

<210> SEQ ID NO 17  
<211> LENGTH: 34  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: primer

&lt;400&gt; SEQUENCE: 17

caaatccaac aaaacgcgca cataggctcg atcc

34

<210> SEQ ID NO 18  
<211> LENGTH: 34  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: primer

&lt;400&gt; SEQUENCE: 18

gatcgacggt atgtgcgcgt ttggggat ttgc

34

<210> SEQ ID NO 19  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: HPV

&lt;400&gt; SEQUENCE: 19

Arg Ala His Tyr Asn Ile Val Thr Phe  
1 5

<210> SEQ ID NO 20  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: HIV-1

&lt;400&gt; SEQUENCE: 20

Ala Met Gln Met Leu Lys Glu Thr Ile  
1 5

<210> SEQ ID NO 21  
<211> LENGTH: 13  
<212> TYPE: PRT  
<213> ORGANISM: HIV-1

&lt;400&gt; SEQUENCE: 21

Val Val Glu Asp Glu Gly Cys Thr Asn Leu Ser Gly Phe  
1 5 10

<210> SEQ ID NO 22  
<211> LENGTH: 758  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: gDgag

**46**

## US 12,390,525 B2

**47**

-continued

&lt;400&gt; SEQUENCE: 22

Met Gly Gly Ala Ala Ala Arg Leu Gly Ala Val Ile Leu Phe Val Val  
 1 5 10 15

Ile Val Gly Leu His Gly Val Arg Gly Lys Tyr Ala Leu Ala Asp Ala  
 20 25 30

Ser Leu Lys Met Ala Asp Pro Asn Arg Phe Arg Gly Lys Asp Leu Pro  
 35 40 45

Val Leu Asp Gln Leu Thr Asp Pro Pro Gly Val Arg Arg Val Tyr His  
 50 55 60

Ile Gln Ala Gly Leu Pro Asp Pro Phe Gln Pro Pro Ser Leu Pro Ile  
 65 70 75 80

Thr Val Tyr Tyr Ala Val Leu Glu Arg Ala Cys Arg Ser Val Leu Leu  
 85 90 95

Asn Ala Pro Ser Glu Ala Pro Gln Ile Val Arg Gly Ala Ser Glu Asp  
 100 105 110

Val Arg Lys Gln Pro Tyr Asn Leu Thr Ile Ala Trp Phe Arg Met Gly  
 115 120 125

Gly Asn Cys Ala Ile Pro Ile Thr Val Met Glu Tyr Thr Glu Cys Ser  
 130 135 140

Tyr Asn Lys Ser Leu Gly Ala Cys Pro Ile Arg Thr Gln Pro Arg Trp  
 145 150 155 160

Asn Tyr Tyr Asp Ser Phe Ser Ala Val Ser Glu Asp Asn Leu Gly Phe  
 165 170 175

Leu Met His Ala Pro Ala Phe Glu Thr Ala Gly Thr Tyr Leu Arg Leu  
 180 185 190

Val Lys Ile Asn Asp Trp Thr Glu Ile Thr Gln Phe Ile Leu Glu His  
 195 200 205

Arg Ala Lys Gly Ser Cys Lys Tyr Ala Leu Pro Leu Arg Ile Pro Pro  
 210 215 220

Ser Ala Cys Leu Ser Pro Gln Ala Tyr Gln Gln Gly Val Thr Val Asp  
 225 230 235 240

Ser Ile Gly Met Leu Pro Arg Phe Ile Pro Glu Asn Gln Arg Thr Val  
 245 250 255

Ala Val Tyr Ser Leu Lys Ile Ala Gly Trp His Gly Pro Lys Ala Pro  
 260 265 270

Tyr Thr Ser Thr Leu Leu Pro Pro Glu Leu Ser Glu Thr Pro Asn Ala  
 275 280 285

Thr Gln Pro Glu Leu Ala Pro Glu Asp Pro Glu Asp Ser Ala Leu Leu  
 290 295 300

Glu Asp Pro Val Gly Thr Val Ala Pro Gly Ala Arg Ala Ser Val Leu  
 305 310 315 320

Ser Gly Gly Glu Leu Asp Arg Trp Glu Lys Ile Arg Leu Arg Pro Gly  
 325 330 335

Gly Lys Lys Tyr Lys Leu Lys His Ile Val Trp Ala Ser Arg Glu  
 340 345 350

Leu Glu Arg Phe Ala Val Asn Pro Gly Leu Leu Glu Thr Ser Glu Gly  
 355 360 365

Cys Arg Gln Ile Leu Gly Gln Leu Gln Pro Ser Leu Gln Thr Gly Ser  
 370 375 380

Glu Glu Leu Arg Ser Leu Tyr Asn Thr Val Ala Thr Leu Tyr Cys Val  
 385 390 395 400

His Gln Arg Ile Glu Ile Lys Asp Thr Lys Glu Ala Leu Asp Lys Ile  
 405 410 415

**48**

## US 12,390,525 B2

**49****50**

-continued

Glu Glu Glu Gln Asn Lys Ser Lys Lys Ala Gln Gln Ala Ala Ala  
 420 425 430  
 Asp Thr Gly His Ser Asn Gln Val Ser Gln Asn Tyr Pro Ile Val Gln  
 435 440 445  
 Asn Ile Gln Gly Gln Met Val His Gln Ala Ile Ser Pro Arg Thr Leu  
 450 455 460  
 Asn Ala Trp Val Lys Val Val Glu Lys Ala Phe Ser Pro Glu Val  
 465 470 475 480  
 Ile Pro Met Phe Ser Ala Leu Ser Glu Gly Ala Thr Pro Gln Asp Leu  
 485 490 495  
 Asn Thr Met Leu Asn Thr Val Gly Gly His Gln Ala Ala Met Gln Met  
 500 505 510  
 Leu Lys Glu Thr Ile Asn Glu Glu Ala Ala Asp Trp Asp Arg Val His  
 515 520 525  
 Pro Val His Ala Gly Pro Ile Ala Pro Gly Gln Met Arg Glu Pro Arg  
 530 535 540  
 Gly Ser Asp Ile Ala Gly Thr Thr Ser Thr Leu Gln Glu Gln Ile Gly  
 545 550 555 560  
 Trp Met Thr Asn Asn Pro Pro Ile Pro Val Gly Glu Ile Tyr Lys Arg  
 565 570 575  
 Trp Ile Ile Leu Gly Leu Asn Lys Ile Val Arg Met Tyr Ser Pro Thr  
 580 585 590  
 Ser Ile Leu Asp Ile Arg Gln Gly Pro Lys Glu Pro Phe Arg Asp Tyr  
 595 600 605  
 Val Asp Arg Phe Tyr Lys Thr Leu Arg Ala Glu Gln Ala Ser Gln Glu  
 610 615 620  
 Val Lys Asn Trp Met Thr Glu Thr Leu Leu Val Gln Asn Ala Asn Pro  
 625 630 635 640  
 Asp Cys Lys Thr Ile Leu Lys Ala Leu Gly Pro Ala Ala Thr Leu Glu  
 645 650 655  
 Glu Met Met Thr Ala Cys Gln Gly Val Gly Gly Pro Gly His Lys Ala  
 660 665 670  
 Arg Val Leu Ala Pro Gln Ile Pro Pro Asn Trp His Ile Pro Ser Ile  
 675 680 685  
 Gln Asp Ala Ala Thr Pro Tyr His Pro Pro Ala Thr Pro Asn Asn Met  
 690 695 700  
 Gly Leu Ile Ala Gly Ala Val Gly Gly Ser Leu Leu Ala Ala Leu Val  
 705 710 715 720  
 Ile Cys Gly Ile Val Tyr Trp Met Arg Arg Arg Thr Gln Lys Ala Pro  
 725 730 735  
 Lys Arg Ile Arg Leu Pro His Ile Arg Glu Asp Asp Gln Pro Ser Ser  
 740 745 750  
 His Gln Pro Leu Phe Tyr  
 755

<210> SEQ ID NO 23  
 <211> LENGTH: 732  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: gDE7E6E5

<400> SEQUENCE: 23

Met Gly Gly Ala Ala Ala Arg Leu Gly Ala Val Ile Leu Phe Val Val  
 1 5 10 15

-continued

Ile Val Gly Leu His Gly Val Arg Gly Lys Tyr Ala Leu Ala Asp Ala  
20 25 30

Ser Leu Lys Met Ala Asp Pro Asn Arg Phe Arg Gly Lys Asp Leu Pro  
35 40 45

Val Leu Asp Gln Leu Thr Asp Pro Pro Gly Val Arg Arg Val Tyr His  
50 55 60

Ile Gln Ala Gly Leu Pro Asp Pro Phe Gln Pro Pro Ser Leu Pro Ile  
65 70 75 80

Thr Val Tyr Tyr Ala Val Leu Glu Arg Ala Cys Arg Ser Val Leu Leu  
85 90 95

Asn Ala Pro Ser Glu Ala Pro Gln Ile Val Arg Gly Ala Ser Glu Asp  
100 105 110

Val Arg Lys Gln Pro Tyr Asn Leu Thr Ile Ala Trp Phe Arg Met Gly  
115 120 125

Gly Asn Cys Ala Ile Pro Ile Thr Val Met Glu Tyr Thr Glu Cys Ser  
130 135 140

Tyr Asn Lys Ser Leu Gly Ala Cys Pro Ile Arg Thr Gln Pro Arg Trp  
145 150 155 160

Asn Tyr Tyr Asp Ser Phe Ser Ala Val Ser Glu Asp Asn Leu Gly Phe  
165 170 175

Leu Met His Ala Pro Ala Phe Glu Thr Ala Gly Thr Tyr Leu Arg Leu  
180 185 190

Val Lys Ile Asn Asp Trp Thr Glu Ile Thr Gln Phe Ile Leu Glu His  
195 200 205

Arg Ala Lys Gly Ser Cys Lys Tyr Ala Leu Pro Leu Arg Ile Pro Pro  
210 215 220

Ser Ala Cys Leu Ser Pro Gln Ala Tyr Gln Gln Gly Val Thr Val Asp  
225 230 235 240

Ser Ile Gly Met Leu Pro Arg Phe Ile Pro Glu Asn Gln Arg Thr Val  
245 250 255

Ala Val Tyr Ser Leu Lys Ile Ala Gly Trp His Gly Pro His Gly Asp  
260 265 270

Thr Pro Thr Leu His Glu Tyr Met Leu Asp Leu Gln Pro Glu Thr Thr  
275 280 285

Asp Leu Tyr Cys Tyr Glu Gln Leu Asn Asp Ser Ser Glu Glu Glu Asp  
290 295 300

Glu Ile Asp Gly Pro Ala Gly Gln Ala Glu Pro Asp Arg Ala His Tyr  
305 310 315 320

Asn Ile Val Thr Phe Cys Cys Lys Cys Asp Ser Thr Leu Arg Leu Cys  
325 330 335

Val Gln Ser Thr His Val Asp Ile Arg Thr Leu Glu Asp Leu Leu Met  
340 345 350

Gly Thr Leu Gly Ile Val Cys Pro Ile Cys Ser Gln Lys Pro Gly Ala  
355 360 365

His Gln Lys Arg Thr Ala Met Phe Gln Asp Pro Gln Glu Arg Pro Arg  
370 375 380

Lys Leu Pro Gln Leu Cys Thr Glu Leu Gln Thr Thr Ile His Asp Ile  
385 390 395 400

Ile Leu Glu Cys Val Tyr Cys Lys Gln Gln Leu Leu Arg Arg Glu Val  
405 410 415

Tyr Asp Phe Ala Phe Arg Asp Leu Cys Ile Val Tyr Arg Asp Gly Asn  
420 425 430

## US 12,390,525 B2

**53****54**

-continued

Pro Tyr Ala Val Cys Asp Lys Cys Leu Lys Phe Tyr Ser Lys Ile Ser  
435 440 445

Glu Tyr Arg His Tyr Cys Tyr Ser Leu Tyr Gly Thr Thr Leu Glu Gln  
450 455 460

Gln Tyr Asn Lys Pro Leu Cys Asp Leu Leu Ile Arg Cys Ile Asn Cys  
465 470 475 480

Gln Lys Pro Leu Cys Pro Glu Glu Lys Gln Arg His Leu Asp Lys Lys  
485 490 495

Gln Arg Phe His Asn Ile Arg Gly Arg Trp Thr Gly Arg Cys Met Ser  
500 505 510

Cys Cys Arg Ser Ser Arg Thr Arg Arg Glu Thr Gln Leu Ala Ala Ala  
515 520 525

Cys Ile His Asn Ile Thr Gly Val Leu Phe Ala Leu Leu Cys Val Leu  
530 535 540

Leu Cys Val Cys Leu Leu Ile Arg Pro Leu Leu Ser Val Ser Thr  
545 550 555 560

Tyr Thr Ser Leu Ile Ile Leu Val Leu Leu Leu Trp Ile Thr Ala Ala  
565 570 575

Ser Ala Phe Arg Cys Phe Ile Val Tyr Ile Ile Phe Val Tyr Ile Pro  
580 585 590

Leu Phe Leu Ile His Thr His Ala Arg Phe Leu Ile Thr Gly Pro Lys  
595 600 605

Ala Pro Tyr Thr Ser Thr Leu Leu Pro Pro Glu Leu Ser Glu Thr Pro  
610 615 620

Asn Ala Thr Gln Pro Glu Leu Ala Pro Glu Asp Pro Glu Asp Ser Ala  
625 630 635 640

Leu Leu Glu Asp Pro Val Gly Thr Val Ala Pro Gln Ile Pro Pro Asn  
645 650 655

Trp His Ile Pro Ser Ile Gln Asp Ala Ala Thr Pro Tyr His Pro Pro  
660 665 670

Ala Thr Pro Asn Asn Met Gly Leu Ile Ala Gly Ala Val Gly Ser  
675 680 685

Leu Leu Ala Ala Leu Val Ile Cys Gly Ile Val Tyr Trp Met Arg Arg  
690 695 700

Arg Thr Gln Lys Ala Pro Lys Arg Ile Arg Leu Pro His Ile Arg Glu  
705 710 715 720

Asp Asp Gln Pro Ser Ser His Gln Pro Leu Phe Tyr  
725 730

&lt;210&gt; SEQ ID NO 24

&lt;211&gt; LENGTH: 732

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: gDE7E6E5 W294A

&lt;400&gt; SEQUENCE: 24

Met Gly Gly Ala Ala Ala Arg Leu Gly Ala Val Ile Leu Phe Val Val  
1 5 10 15

Ile Val Gly Leu His Gly Val Arg Gly Lys Tyr Ala Leu Ala Asp Ala  
20 25 30

Ser Leu Lys Met Ala Asp Pro Asn Arg Phe Arg Gly Lys Asp Leu Pro  
35 40 45

Val Leu Asp Gln Leu Thr Asp Pro Pro Gly Val Arg Arg Val Tyr His  
50 55 60

## US 12,390,525 B2

**55****56**

-continued

---

Ile Gln Ala Gly Leu Pro Asp Pro Phe Gln Pro Pro Ser Leu Pro Ile  
65 70 75 80

Thr Val Tyr Tyr Ala Val Leu Glu Arg Ala Cys Arg Ser Val Leu Leu  
85 90 95

Asn Ala Pro Ser Glu Ala Pro Gln Ile Val Arg Gly Ala Ser Glu Asp  
100 105 110

Val Arg Lys Gln Pro Tyr Asn Leu Thr Ile Ala Trp Phe Arg Met Gly  
115 120 125

Gly Asn Cys Ala Ile Pro Ile Thr Val Met Glu Tyr Thr Glu Cys Ser  
130 135 140

Tyr Asn Lys Ser Leu Gly Ala Cys Pro Ile Arg Thr Gln Pro Arg Trp  
145 150 155 160

Asn Tyr Tyr Asp Ser Phe Ser Ala Val Ser Glu Asp Asn Leu Gly Phe  
165 170 175

Leu Met His Ala Pro Ala Phe Glu Thr Ala Gly Thr Tyr Leu Arg Leu  
180 185 190

Val Lys Ile Asn Asp Trp Thr Glu Ile Thr Gln Phe Ile Leu Glu His  
195 200 205

Arg Ala Lys Gly Ser Cys Lys Tyr Ala Leu Pro Leu Arg Ile Pro Pro  
210 215 220

Ser Ala Cys Leu Ser Pro Gln Ala Tyr Gln Gln Gly Val Thr Val Asp  
225 230 235 240

Ser Ile Gly Met Leu Pro Arg Phe Ile Pro Glu Asn Gln Arg Thr Val  
245 250 255

Ala Val Tyr Ser Leu Lys Ile Ala Gly Trp His Gly Pro His Gly Asp  
260 265 270

Thr Pro Thr Leu His Glu Tyr Met Leu Asp Leu Gln Pro Glu Thr Thr  
275 280 285

Asp Leu Tyr Cys Tyr Glu Gln Leu Asn Asp Ser Ser Glu Glu Glu Asp  
290 295 300

Glu Ile Asp Gly Pro Ala Gly Gln Ala Glu Pro Asp Arg Ala His Tyr  
305 310 315 320

Asn Ile Val Thr Phe Cys Cys Lys Cys Asp Ser Thr Leu Arg Leu Cys  
325 330 335

Val Gln Ser Thr His Val Asp Ile Arg Thr Leu Glu Asp Leu Leu Met  
340 345 350

Gly Thr Leu Gly Ile Val Cys Pro Ile Cys Ser Gln Lys Pro Gly Ala  
355 360 365

His Gln Lys Arg Thr Ala Met Phe Gln Asp Pro Gln Glu Arg Pro Arg  
370 375 380

Lys Leu Pro Gln Leu Cys Thr Glu Leu Gln Thr Thr Ile His Asp Ile  
385 390 395 400

Ile Leu Glu Cys Val Tyr Cys Lys Gln Gln Leu Leu Arg Arg Glu Val  
405 410 415

Tyr Asp Phe Ala Phe Arg Asp Leu Cys Ile Val Tyr Arg Asp Gly Asn  
420 425 430

Pro Tyr Ala Val Cys Asp Lys Cys Leu Lys Phe Tyr Ser Lys Ile Ser  
435 440 445

Glu Tyr Arg His Tyr Cys Tyr Ser Leu Tyr Gly Thr Thr Leu Glu Gln  
450 455 460

Gln Tyr Asn Lys Pro Leu Cys Asp Leu Leu Ile Arg Cys Ile Asn Cys  
465 470 475 480

Gln Lys Pro Leu Cys Pro Glu Glu Lys Gln Arg His Leu Asp Lys Lys

## US 12,390,525 B2

**57**

-continued

**58**

485	490	495
Gln Arg Phe His Asn Ile Arg Gly Arg Trp Thr Gly Arg Cys Met Ser		
500	505	510
Cys Cys Arg Ser Ser Arg Thr Arg Arg Glu Thr Gln Leu Ala Ala Ala		
515	520	525
Cys Ile His Asn Ile Thr Gly Val Leu Phe Ala Leu Leu Cys Val Leu		
530	535	540
Leu Cys Val Cys Leu Leu Ile Arg Pro Leu Leu Leu Ser Val Ser Thr		
545	550	555
Tyr Thr Ser Leu Ile Ile Leu Val Leu Leu Trp Ile Thr Ala Ala		
565	570	575
Ser Ala Phe Arg Cys Phe Ile Val Tyr Ile Ile Phe Val Tyr Ile Pro		
580	585	590
Leu Phe Leu Ile His Thr His Ala Arg Phe Leu Ile Thr Gly Pro Lys		
595	600	605
Ala Pro Tyr Thr Ser Thr Leu Leu Pro Pro Glu Leu Ser Glu Thr Pro		
610	615	620
Asn Ala Thr Gln Pro Glu Leu Ala Pro Glu Asp Pro Glu Asp Ser Ala		
625	630	635
Leu Leu Glu Asp Pro Val Gly Thr Val Ala Pro Gln Ile Pro Pro Asn		
645	650	655
Ala His Ile Pro Ser Ile Gln Asp Ala Ala Thr Pro Tyr His Pro Pro		
660	665	670
Ala Thr Pro Asn Asn Met Gly Leu Ile Ala Gly Ala Val Gly Ser		
675	680	685
Leu Leu Ala Ala Leu Val Ile Cys Gly Ile Val Tyr Trp Met Arg Arg		
690	695	700
Arg Thr Gln Lys Ala Pro Lys Arg Ile Arg Leu Pro His Ile Arg Glu		
705	710	720
Asp Asp Gln Pro Ser Ser His Gln Pro Leu Phe Tyr		
725	730	

&lt;210&gt; SEQ ID NO 25

&lt;211&gt; LENGTH: 394

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: gD W294A

&lt;400&gt; SEQUENCE: 25

Met Gly Gly Ala Ala Ala Arg Leu Gly Ala Val Ile Leu Phe Val Val		
1	5	10
Ile Val Gly Leu His Gly Val Arg Gly Lys Tyr Ala Leu Ala Asp Ala		
20	25	30
Ser Leu Lys Met Ala Asp Pro Asn Arg Phe Arg Gly Lys Asp Leu Pro		
35	40	45
Val Leu Asp Gln Leu Thr Asp Pro Pro Gly Val Arg Arg Val Tyr His		
50	55	60
Ile Gln Ala Gly Leu Pro Asp Pro Phe Gln Pro Pro Ser Leu Pro Ile		
65	70	75
Thr Val Tyr Tyr Ala Val Leu Glu Arg Ala Cys Arg Ser Val Leu Leu		
85	90	95
Asn Ala Pro Ser Glu Ala Pro Gln Ile Val Arg Gly Ala Ser Glu Asp		
100	105	110
Val Arg Lys Gln Pro Tyr Asn Leu Thr Ile Ala Trp Phe Arg Met Gly		

## US 12,390,525 B2

**59****60**

-continued

115 120 125

Gly Asn Cys Ala Ile Pro Ile Thr Val Met Glu Tyr Thr Glu Cys Ser  
 130 135 140

Tyr Asn Lys Ser Leu Gly Ala Cys Pro Ile Arg Thr Gln Pro Arg Trp  
 145 150 155 160

Asn Tyr Tyr Asp Ser Phe Ser Ala Val Ser Glu Asp Asn Leu Gly Phe  
 165 170 175

Leu Met His Ala Pro Ala Phe Glu Thr Ala Gly Thr Tyr Leu Arg Leu  
 180 185 190

Val Lys Ile Asn Asp Trp Thr Glu Ile Thr Gln Phe Ile Leu Glu His  
 195 200 205

Arg Ala Lys Gly Ser Cys Lys Tyr Ala Leu Pro Leu Arg Ile Pro Pro  
 210 215 220

Ser Ala Cys Leu Ser Pro Gln Ala Tyr Gln Gln Gly Val Thr Val Asp  
 225 230 235 240

Ser Ile Gly Met Leu Pro Arg Phe Ile Pro Glu Asn Gln Arg Thr Val  
 245 250 255

Ala Val Tyr Ser Leu Lys Ile Ala Gly Trp His Gly Pro Lys Ala Pro  
 260 265 270

Tyr Thr Ser Thr Leu Leu Pro Pro Glu Leu Ser Glu Thr Pro Asn Ala  
 275 280 285

Thr Gln Pro Glu Leu Ala Pro Glu Asp Pro Glu Asp Ser Ala Leu Leu  
 290 295 300

Glu Asp Pro Val Gly Thr Val Ala Pro Gln Ile Pro Pro Asn Ala His  
 305 310 315 320

Ile Pro Ser Ile Gln Asp Ala Ala Thr Pro Tyr His Pro Pro Ala Thr  
 325 330 335

Pro Asn Asn Met Gly Leu Ile Ala Gly Ala Val Gly Gly Ser Leu Leu  
 340 345 350

Ala Ala Leu Val Ile Cys Gly Ile Val Tyr Trp Met Arg Arg Arg Thr  
 355 360 365

Gln Lys Ala Pro Lys Arg Ile Arg Leu Pro His Ile Arg Glu Asp Asp  
 370 375 380

Gln Pro Ser Ser His Gln Pro Leu Phe Tyr  
 385 390

&lt;210&gt; SEQ ID NO 26

&lt;211&gt; LENGTH: 1206

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Herpes simplex 1

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: CDS

&lt;222&gt; LOCATION: (1)...(1185)

&lt;400&gt; SEQUENCE: 26

atg ggg ggg gct gcc gcc agg ttg ggg gcc gtg att ttg ttt gtc gtc	48
Met Gly Gly Ala Ala Ala Arg Leu Gly Ala Val Ile Leu Phe Val Val	
1 5 10 15	

ata gtg ggc ctc cat ggg gtc cgc ggc aaa tat gcc ttg gcg gat gcc	96
Ile Val Gly Leu His Gly Val Arg Gly Lys Tyr Ala Leu Ala Asp Ala	
20 25 30	

tct ctc aag atg gcc gac ccc aat cgc ttt cgc ggc aaa gac ctt ccg	144
Ser Leu Lys Met Ala Asp Pro Asn Arg Phe Arg Gly Lys Asp Leu Pro	
35 40 45	

gtc ctg gac cag ctg acc gac cct ccg ggg gtc cgg cgc gtg tac cac	192
Val Leu Asp Gln Leu Thr Asp Pro Pro Gly Val Arg Arg Val Tyr His	
50 55 60	

## US 12,390,525 B2

**61****62**

-continued

---

atc cag ggc cta ccg gac ccg ttc cag ccc ccc agc ctc ccg atc Ile Gln Ala Gly Leu Pro Asp Pro Phe Gln Pro Pro Ser Leu Pro Ile 65 70 75 80	240
acg gtt tac tac gcc gtg ttg gag cgc gcc tgc cgc agc gtg ctc cta Thr Val Tyr Tyr Ala Val Leu Glu Arg Ala Cys Arg Ser Val Leu Leu 85 90 95	288
aac gca ccg tcg gag gcc ccc cag att gtc cgc ggg gcc tcc gaa gac Asn Ala Pro Ser Glu Ala Pro Gln Ile Val Arg Gly Ala Ser Glu Asp 100 105 110	336
gtc cgg aaa caa ccc tac aac ctg acc atc gct tgg ttt cgg atg gga Val Arg Lys Gln Pro Tyr Asn Leu Thr Ile Ala Trp Phe Arg Met Gly 115 120 125	384
ggc aac tgt gct atc ccc atc acg gtc atg gag tac acc gaa tgc tcc Gly Asn Cys Ala Ile Pro Ile Thr Val Met Glu Tyr Thr Glu Cys Ser 130 135 140	432
tac aac aag tct ctg ggg gcc tgt ccc atc cga acg cag ccc cgc tgg Tyr Asn Lys Ser Leu Gly Ala Cys Pro Ile Arg Thr Gln Pro Arg Trp 145 150 155 160	480
aac tac tat gac agc ttc agc gcc gtc agc gag gat aac ctg ggg ttc Asn Tyr Tyr Asp Ser Phe Ser Ala Val Ser Glu Asp Asn Leu Gly Phe 165 170 175	528
ctg atg cac gcc ccc ggc ttt gag acc gcc ggc acg tac ctg cgg ctc Leu Met His Ala Pro Ala Phe Glu Thr Ala Gly Thr Tyr Leu Arg Leu 180 185 190	576
gtg aag ata aac gac tgg acg gag att aca cag ttt atc ctg gag cac Val Lys Ile Asn Asp Trp Thr Glu Ile Thr Gln Phe Ile Leu Glu His 195 200 205	624
cga gcc aag ggc tcc tgt aag tac gcc ctc ccg ctg cgc atc ccc ccc Arg Ala Lys Gly Ser Cys Lys Tyr Ala Leu Pro Leu Arg Ile Pro Pro 210 215 220	672
tca gcc tgc ctg tcc ccc cag gcc tac cag cag ggg gtg acg gtg gac Ser Ala Cys Leu Ser Pro Gln Ala Tyr Gln Gln Gly Val Thr Val Asp 225 230 235 240	720
agc atc ggg atg ctg ccc cgc ttc atc ccc gag aac cag cgc acc gtc Ser Ile Gly Met Leu Pro Arg Phe Ile Pro Glu Asn Gln Arg Thr Val 245 250 255	768
gcc gta tac agc ttg aag atc gcc ggg tgg cac ggg ccc aag gcc cca Ala Val Tyr Ser Leu Lys Ile Ala Gly Trp His Gly Pro Lys Ala Pro 260 265 270	816
tac acg agc acc ctg ctg ccc ccg gag ctg tcc gag acc ccc aac gcc Tyr Thr Ser Thr Leu Leu Pro Pro Glu Leu Ser Glu Thr Pro Asn Ala 275 280 285	864
acg cag cca gaa ctc gcc ccg gaa gac ccc gag gat tcg gcc ctc ttg Thr Gln Pro Glu Leu Ala Pro Glu Asp Pro Glu Asp Ser Ala Leu Leu 290 295 300	912
gag gac ccc gtg ggg acg gtg gcg ccg caa atc cca cca aac tgg cac Glu Asp Pro Val Gly Thr Val Ala Pro Gln Ile Pro Pro Asn Trp His 305 310 315 320	960
ata ccg tcg atc cag gac gcc gcg acg cct tac cat ccc ccg gcc acc Ile Pro Ser Ile Gln Asp Ala Ala Thr Pro Tyr His Pro Pro Ala Thr 325 330 335	1008
ccg aac aac atg ggc ctg atc gcc ggc gcg gtg ggc ggc agt ctc ctg Pro Asn Asn Met Gly Leu Ile Ala Gly Ala Val Gly Gly Ser Leu Leu 340 345 350	1056
gca gcc ctg gtc att tgc gga att gtg tac tgg atg cgc cgc cgc act Ala Ala Leu Val Ile Cys Gly Ile Val Tyr Trp Met Arg Arg Arg Thr 355 360 365	1104
caa aaa gcc cca aag cgc ata cgc ctc ccc cac atc cgg gaa gac gac Gln Lys Ala Pro Lys Arg Ile Arg Leu Pro His Ile Arg Glu Asp Asp	1152

## US 12,390,525 B2

**63****64**

-continued

370

375

380

cag ccg tcc tcg cac cac ccc ttg ttt tac tag atacccccc ttaatgggtg 1205  
 Gln Pro Ser Ser His Gln Pro Leu Phe Tyr \*  
 385 390

c 1206

&lt;210&gt; SEQ ID NO 27

&lt;211&gt; LENGTH: 394

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Herpes simplex 1

&lt;400&gt; SEQUENCE: 27

Met	Gly	Gly	Ala	Ala	Ala	Arg	Leu	Gly	Ala	Val	Ile	Leu	Phe	Val	Val
1															
							5			10			15		

Ile	Val	Gly	Leu	His	Gly	Val	Arg	Gly	Lys	Tyr	Ala	Leu	Ala	Asp	Ala
							20		25			30			

Ser	Leu	Lys	Met	Ala	Asp	Pro	Asn	Arg	Phe	Arg	Gly	Lys	Asp	Leu	Pro
							35		40		45				

Val	Leu	Asp	Gln	Leu	Thr	Asp	Pro	Pro	Gly	Val	Arg	Arg	Val	Tyr	His
							50		55		60				

Ile	Gln	Ala	Gly	Leu	Pro	Asp	Pro	Phe	Gln	Pro	Pro	Ser	Leu	Pro	Ile
							65		70		75		80		

Thr	Val	Tyr	Tyr	Ala	Val	Leu	Glu	Arg	Ala	Cys	Arg	Ser	Val	Leu	Leu
							85		90		95				

Asn	Ala	Pro	Ser	Glu	Ala	Pro	Gln	Ile	Val	Arg	Gly	Ala	Ser	Glu	Asp
							100		105		110				

Val	Arg	Lys	Gln	Pro	Tyr	Asn	Leu	Thr	Ile	Ala	Trp	Phe	Arg	Met	Gly
							115		120		125				

Gly	Asn	Cys	Ala	Ile	Pro	Ile	Thr	Val	Met	Glu	Tyr	Thr	Glu	Cys	Ser
							130		135		140				

Tyr	Asn	Lys	Ser	Leu	Gly	Ala	Cys	Pro	Ile	Arg	Thr	Gln	Pro	Arg	Trp
							145		150		155		160		

Asn	Tyr	Tyr	Asp	Ser	Phe	Ser	Ala	Val	Ser	Glu	Asp	Asn	Leu	Gly	Phe
							165		170		175				

Leu	Met	His	Ala	Pro	Ala	Phe	Glu	Thr	Ala	Gly	Thr	Tyr	Leu	Arg	Leu
							180		185		190				

Val	Lys	Ile	Asn	Asp	Trp	Thr	Glu	Ile	Thr	Gln	Phe	Ile	Leu	Glu	His
							195		200		205				

Arg	Ala	Lys	Gly	Ser	Cys	Lys	Tyr	Ala	Leu	Pro	Leu	Arg	Ile	Pro	Pro
							210		215		220				

Ser	Ala	Cys	Leu	Ser	Pro	Gln	Ala	Tyr	Gln	Gln	Gly	Val	Thr	Val	Asp
							225		230		235		240		

Ser	Ile	Gly	Met	Leu	Pro	Arg	Phe	Ile	Pro	Glu	Asn	Gln	Arg	Thr	Val
							245		250		255				

Ala	Val	Tyr	Ser	Leu	Lys	Ile	Ala	Gly	Trp	His	Gly	Pro	Lys	Ala	Pro
							260		265		270				

Tyr	Thr	Ser	Thr	Leu	Leu	Pro	Pro	Glu	Leu	Ser	Glu	Thr	Pro	Asn	Ala
							275		280		285				

Thr	Gln	Pro	Glu	Leu	Ala	Pro	Glu	Asp	Pro	Glu	Asp	Ser	Ala	Leu	Leu
							290		295		300				

Glu	Asp	Pro	Val	Gly	Thr	Val	Ala	Pro	Gln	Ile	Pro	Pro	Asn	Trp	His
							305		310		315		320		

Ile	Pro	Ser	Ile	Gln	Asp	Ala	Ala	Thr	Pro	Tyr	His	Pro	Pro	Ala	Thr
								325		330		335			

## US 12,390,525 B2

**65****66**

-continued

---

Pro Asn Asn Met Gly Leu Ile Ala Gly Ala Val Gly Gly Ser Leu Leu  
340 345 350

Ala Ala Leu Val Ile Cys Gly Ile Val Tyr Trp Met Arg Arg Arg Thr  
355 360 365

Gln Lys Ala Pro Lys Arg Ile Arg Leu Pro His Ile Arg Glu Asp Asp  
370 375 380

Gln Pro Ser Ser His Gln Pro Leu Phe Tyr  
385 390

&lt;210&gt; SEQ ID NO 28

&lt;211&gt; LENGTH: 1730

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Herpes simplex 2

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: CDS

&lt;222&gt; LOCATION: (275) ... (1456)

&lt;400&gt; SEQUENCE: 28

agagcggtgg	gggggggggg	gggaagaaac	taaaaacaca	tcaagccac	aaccctccac	60	
aaggggggtt	atggcgacc	caccgcacca	ccatactccg	attcgaccac	atatgcaacc	120	
aaatcacccc	cagaggggaa	ggttccattt	ttacgaggag	gaggagtata	atagagtctt	180	
tgtgtttaaa	accgggggtc	ggtgtggtgt	tcggtcataa	gctgcattgc	gaacgactag	240	
tcggccgttt	ttcgtgtgca	tcgctatca	cggc	atg ggg cgt ttg acc tcc ggc	Met Gly Arg Leu Thr Ser Gly	295	
				1	5		
gtc	ggg acg	gcc ctg	cta gtt	gtc gcg	gtg gga ctc cgc	gtc gtc	343
Val	Gly	Thr	Ala	Leu	Leu	Val	
10	15	20					
tgc	gcc aaa tac	gcc tta	gca gac	ccc tcg	ctt aag atg	gcc gat	391
Cys	Ala	Lys	Tyr	Ala	Leu	Ala	
25	30	35					
aat	cga ttt	cgc ggg	aag aac	ctt ccg	gtt ttg	gac cag	439
Asn	Arg	Phe	Arg	Gly	Lys	Asn	
40	45	50	55				
ccc	ccc ggg	gtg aag	cgt gtt	tac cac	att cag	ccg agc	487
Pro	Pro	Gly	Val	Lys	Arg	Val	
60	65	70					
ccg	tcc cag	ccc agc	atc ccg	atc act	gtg tac	tac gca	535
Pro	Phe	Gln	Pro	Pro	Ser	Ile	
75	80	85					
gaa	cgt	gcc	tgc	cgc	agc	gtg	583
Glu	Arg	Ala	Cys	Arg	Ser	Val	
90	95	100					
cag	atc	gtc	ccc	ggg	gtc	gac	631
Gln	Ile	Val	Arg	Gly	Ala	Ser	
105	110	115					
ctg	acc atc	gcc tgg	tat	cgc atg	gga gac	aat tgc	679
Leu	Thr	Ile	Ala	Trp	Tyr	Arg	
120	125	130					
acg	gtt	atg	gaa	tac	acc gag	tgc ccc tac	727
Thr	Val	Met	Glu	Tyr	Thr	Glu	
140	145	150					
tgc	ccc atc	cga acg	cag	ccc cgc	tgg agc	tac tat	775
Cys	Pro	Ile	Arg	Thr	Gln	Pro	
155	160	165					
gcc	gtc	agc	gag	gat	aac ctg	gga ttc	823
Ala	Val	Ser	Glu	Asp	Asn	Leu	
170	175	180					

## US 12,390,525 B2

**67****68**

-continued

---

gag acc gcg ggt acg tac ctg cggtt cta gtg aag ata aac gac tgg acg Glu Thr Ala Gly Thr Tyr Leu Arg Leu Val Lys Ile Asn Asp Trp Thr 185 190 195	871
gag atc aca caa ttt atc ctg gag cac cggtt gcc ggc tcc tgc aag Glu Ile Thr Gln Phe Ile Leu Glu His Arg Ala Arg Ala Ser Cys Lys 200 205 210 215	919
tac gct ctc ccc ctg cgc atc ccc ccgtt gca ggc tgc ctc acc tcg aag Tyr Ala Leu Pro Leu Arg Ile Pro Pro Ala Ala Cys Leu Thr Ser Lys 220 225 230	967
gcc tac caa cag ggc gtg acg gtc gac agc atc ggg atg tta ccc cgc Ala Tyr Gln Gln Gly Val Thr Val Asp Ser Ile Gly Met Leu Pro Arg 235 240 245	1015
ttt act ccc gaa aac cag cgc acc gtc gcc cta tac agc tta aaa atc Phe Thr Pro Glu Asn Gln Arg Thr Val Ala Leu Tyr Ser Leu Lys Ile 250 255 260	1063
gcc ggg tgg cac ggc ccc aag ccc ccgtt tac acc agc acc ctg ctg ccc Ala Gly Trp His Gly Pro Lys Pro Pro Tyr Thr Ser Thr Leu Leu Pro 265 270 275	1111
ccg gag ctg tcc gac acc acc aac gcc acg caa ccc gaa ctc gtt ccg Pro Glu Leu Ser Asp Thr Thr Asn Ala Thr Gln Pro Glu Leu Val Pro 280 285 290 295	1159
gaa gac ccc gag gac tcg gcc ctc tta gag gat ccc gcc ggg acg gtg Glu Asp Pro Glu Asp Ser Ala Leu Leu Glu Asp Pro Ala Gly Thr Val 300 305 310	1207
tct tcg cag atc ccc cca aac tgg cac atc ccgtt tcg atc cag gac gtc Ser Ser Gln Ile Pro Pro Asn Trp His Ile Pro Ser Ile Gln Asp Val 315 320 325	1255
gcg ccg cac cac ggc ccc gcc gcc cca gcc aac ccgtt ggc ctg atc atc Ala Pro His His Ala Pro Ala Ala Pro Ala Asn Pro Gly Leu Ile Ile 330 335 340	1303
ggc gcg ctg gcc ggc agt acc ctg gcg gctt gtc atc ggc ggt att Gly Ala Leu Ala Gly Ser Thr Leu Ala Ala Leu Val Ile Gly Gly Ile 345 350 355	1351
gcg ttt tgg gta cgc cgc ccgtt tca gtg gcc ccc aag cgc cta cgt Ala Phe Trp Val Arg Arg Arg Ser Val Ala Pro Lys Arg Leu Arg 360 365 370 375	1399
ctc ccc cac atc ccgtt gat gac gac ggc ccc ccgtt tcg cac cag cca ttg Leu Pro His Ile Arg Asp Asp Asp Ala Pro Pro Ser His Gln Pro Leu 380 385 390	1447
ttt tac tag aggagttcc ccgttccgtt gtacctctga ggccgttgt Phe Tyr *	1496
gagggtggctt ggggtattta ggggtggact tggactccgc ataaaggagt ctgcaggag ggaaaactagg acagttcata ggccgggagc gtggggcgcg cgccgcgtgt cccgacgatt agccagccgc gccccacagcc acctcgaccc ggtccgatcc gcggtatgcc cggccgcctcg ctgcaggggcc tggcgatccct gggccgttgtt gtctgcgcaca ccggcccttgtt ccgt	1556 1616 1676 1730
<210> SEQ ID NO 29	
<211> LENGTH: 393	
<212> TYPE: PRT	
<213> ORGANISM: Herpes simplex 2	
<400> SEQUENCE: 29	
Met Gly Arg Leu Thr Ser Gly Val Gly Thr Ala Ala Leu Leu Val Val 1 5 10 15	
Ala Val Gly Leu Arg Val Val Cys Ala Lys Tyr Ala Leu Ala Asp Pro 20 25 30	
Ser Leu Lys Met Ala Asp Pro Asn Arg Phe Arg Gly Lys Asn Leu Pro 35 40 45	

## US 12,390,525 B2

**69****70**

-continued

Val Leu Asp Gln Leu Thr Asp Pro Pro Gly Val Lys Arg Val Tyr His  
50 55 60

Ile Gln Pro Ser Leu Glu Asp Pro Phe Gln Pro Pro Ser Ile Pro Ile  
65 70 75 80

Thr Val Tyr Tyr Ala Val Leu Glu Arg Ala Cys Arg Ser Val Leu Leu  
85 90 95

His Ala Pro Ser Glu Ala Pro Gln Ile Val Arg Gly Ala Ser Asp Glu  
100 105 110

Ala Arg Lys His Thr Tyr Asn Leu Thr Ile Ala Trp Tyr Arg Met Gly  
115 120 125

Asp Asn Cys Ala Ile Pro Ile Thr Val Met Glu Tyr Thr Glu Cys Pro  
130 135 140

Tyr Asn Lys Ser Leu Gly Val Cys Pro Ile Arg Thr Gln Pro Arg Trp  
145 150 155 160

Ser Tyr Tyr Asp Ser Phe Ser Ala Val Ser Glu Asp Asn Leu Gly Phe  
165 170 175

Leu Met His Ala Pro Ala Phe Glu Thr Ala Gly Thr Tyr Leu Arg Leu  
180 185 190

Val Lys Ile Asn Asp Trp Thr Glu Ile Thr Gln Phe Ile Leu Glu His  
195 200 205

Arg Ala Arg Ala Ser Cys Lys Tyr Ala Leu Pro Leu Arg Ile Pro Pro  
210 215 220

Ala Ala Cys Leu Thr Ser Lys Ala Tyr Gln Gln Gly Val Thr Val Asp  
225 230 235 240

Ser Ile Gly Met Leu Pro Arg Phe Thr Pro Glu Asn Gln Arg Thr Val  
245 250 255

Ala Leu Tyr Ser Leu Lys Ile Ala Gly Trp His Gly Pro Lys Pro Pro  
260 265 270

Tyr Thr Ser Thr Leu Leu Pro Pro Glu Leu Ser Asp Thr Thr Asn Ala  
275 280 285

Thr Gln Pro Glu Leu Val Pro Glu Asp Pro Glu Asp Ser Ala Leu Leu  
290 295 300

Glu Asp Pro Ala Gly Thr Val Ser Ser Gln Ile Pro Pro Asn Trp His  
305 310 315 320

Ile Pro Ser Ile Gln Asp Val Ala Pro His His Ala Pro Ala Ala Pro  
325 330 335

Ala Asn Pro Gly Leu Ile Ile Gly Ala Leu Ala Gly Ser Thr Leu Ala  
340 345 350

Ala Leu Val Ile Gly Gly Ile Ala Phe Trp Val Arg Arg Arg Arg Ser  
355 360 365

Val Ala Pro Lys Arg Leu Arg Leu Pro His Ile Arg Asp Asp Asp Ala  
370 375 380

Pro Pro Ser His Gln Pro Leu Phe Tyr  
385 390

&lt;210&gt; SEQ ID NO 30

&lt;211&gt; LENGTH: 8

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: HIV-1

&lt;400&gt; SEQUENCE: 30

Ser Ile Ile Asn Phe Glu Lys Leu  
1 5

-continued

---

<210> SEQ ID NO 31  
 <211> LENGTH: 2199  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: nucleotide sequence encoding gDE7E6E5

<400> SEQUENCE: 31

atgggggggg	ctgccgcccag	gttggggggc	gtgattttgt	ttgtcgcat	agtgggcctc	60
catgggttcc	cgggcaata	tgccttggcg	gatgcctctc	tcaagatggc	cgaccaccaat	120
cgctttcgcg	gcaaagacct	tccggtctcg	gaccagctga	ccgaccctcc	gggggttcgg	180
cgcgtgtacc	acatccaggc	gggccttaccg	gaccggttcc	agccccccag	cctcccgatc	240
acggtttact	acggcgtgtt	ggagcgcgcc	tgcgcagcg	tgctcctaaa	cgcaccgtcg	300
gaggcccccc	agattgtccg	cggggcctcc	gaagacgtcc	ggaaacaacc	ctacaacctg	360
accatcgctt	ggtttccgat	gggaggcaac	tgtgctatcc	ccatcacggt	catggagtag	420
acccaatgt	cctacaacaa	gtctctgggg	gcctgtccca	tccgaacgca	gccccgtgg	480
aactactatg	acagcttcag	cgcgtcago	gaggataacc	tggggttcct	gatgcacgccc	540
cccgcggttgc	agaccgcccc	cacgtacctg	cggctcgta	agataaacgca	ctggacggag	600
attacacagt	ttatcttggaa	gcacccgagcc	aagggtctcc	gtaagtacgc	cctcccgctg	660
cgcacccccc	cgtcagccctg	cctgtccccc	caggccctacc	agcagggggt	gacgggtggac	720
agcatcggga	tgctgccccg	cttcatcccc	gagaaccaggc	gcaccgtcgc	cgtatacagc	780
ttgaagatcg	cgggggtggca	cggggcccat	ggagatacac	ctacattgca	tgaatatatg	840
tttagatttgc	aaccagagac	aactgatctc	tactgttatg	agcaattaaa	tgacagctca	900
gaggaggagg	atgaaaataga	tggtccagct	ggacaaggcag	aaccggacag	agccattac	960
aatattgtaa	cctttgttg	caagtgtgac	tctacgcttc	ggttgtgcgt	acaaaggcaca	1020
cacgttagaca	ttcgtacttt	ggaagacctg	ttaatggca	cactaggaat	tgtgtgcccc	1080
atctgttctc	agaaaaccagg	cgcacccacaa	aagagaactg	caatgttca	ggaccacacag	1140
gagcgcacca	gaaagtacc	acagttatgc	acagagctgc	aaacaactat	acatgatata	1200
atattagaat	gtgtgtactg	caagcaacag	ttactgcfgac	gtgagggtata	tgactttgt	1260
tttcgggatt	tatgcatagt	atataagat	ggaaatccat	atgctgtatg	tgataaatgt	1320
ttaaagttt	attctaaaat	tagtgagtat	agacattatt	gttataatgtt	gtatggaca	1380
acatttagaac	agcaataacaa	caaaccgttg	tgtgatttg	taatttagtg	tattaactgt	1440
caaaagccac	tgtgtctga	agaaaagcaa	agacatctgg	acaaaaagca	aagattccat	1500
aatataaggg	gtcgggtggac	cggtcgtatgt	atgtcttgtt	gcagatcatc	aagaacacgt	1560
agagaaaacc	agctggcgcc	cgcctgcata	cacaacatta	ctggcgctgt	ttttgttgc	1620
ctttgtgtgc	ttttgtgtgt	ctgccttatta	atacgtccgc	tgcgtttgtc	tgtgtctaca	1680
tacacatcat	taataatatt	ggtattacta	ttgtggataa	cagcagccctc	tgcgtttagg	1740
tgttttattg	tatataattat	atttgtttat	ataccattat	tttaatacaca	tacacatgca	1800
cgctttttaa	ttacaggggcc	caaggccccca	tacacgagca	cctgtgtcc	cccgaggctg	1860
tccgagaccc	ccaacgcccac	gcagccagaa	ctggcccccgg	aagaccccgaa	ggattcggcc	1920
ctcttggagg	accccggtgg	gacggtggcg	ccgcaaatcc	caccaaactg	gcacataccg	1980
tcgatccagg	acgcccgcac	gccttaccat	ccccggcca	ccccgaacaa	catgggcctg	2040
atcgccggcg	cggtggcg	cagtctcctg	gcagccctgg	tcatttgccg	aattgtgtac	2100

-continued

---

tggatgcgcc	gcccactca	aaaagccca	aagcgcatac	gcctccccca	catccggaa	2160
gacgaccaggc	cgtctcgca	ccagcccttg	ttttactag			2199
<hr/>						
<210> SEQ ID NO 32						
<211> LENGTH: 2277						
<212> TYPE: DNA						
<213> ORGANISM: Artificial Sequence						
<220> FEATURE:						
<223> OTHER INFORMATION: nucleotide sequence encoding gDgag						
<400> SEQUENCE: 32						
atgggggggg	ctggcccaag	gttgggggc	gtgattttgt	ttgtcgcat	agtgggcctc	60
catggggtcc	gcccataata	tgccttggcg	gatgcctctc	tcaagatggc	cgacccaaat	120
cggttcgcg	gcaaagacct	tccggtcctg	gaccagctga	ccgaccctcc	gggggtccgg	180
cgcggttacc	acatccaggc	gggccttaccg	gaccgcgtcc	agccccccag	cctcccgatc	240
acggtttact	acgcgtgtt	ggagcgcgc	tgccgcagcg	tgctctaaa	cgcaccgtcg	300
gaggcccccc	agattgtccg	cggggcctcc	gaagacgtcc	ggaaacaacc	ctacaacctg	360
accatcgctt	ggtttccggat	gggaggcaac	tgtgctatcc	ccatcacggt	catggagtagc	420
accgaatgt	cctacaacaa	gtctctgggg	gcctgtccca	tccgaacgca	gccccgctgg	480
aactactatg	acagcttcag	cgccgtcago	gaggataacc	tggggttcct	gatgcacgcc	540
cccgcgtttgc	agaccgcgg	cacgtacctg	cggtctcgta	agataaacga	ctggacggag	600
attacacagt	ttatcctgga	gcaccggagcc	aagggtctct	gtaagtacgc	cctcccgctg	660
cgcacatcccc	cgtcagccctg	cctgtccccc	caggcctacc	agcaggggggt	gacgggtggac	720
agcatcggga	tgctgccccg	cttcatcccc	gagaaccagc	gcaccgtcgc	cgtatacagc	780
ttgaagatcg	ccgggtggca	cggggccaag	gccccataca	cgagcacccct	gctgcccccg	840
gagctgtccg	agaccccaa	cgccacgcag	ccagaactcg	ccccggaaaga	ccccgaggat	900
tcggccctct	tggaggaccc	cgtggggacg	gtggcgccgg	gtgcgagagc	gtcagtattta	960
agcgggggag	aattagatcg	atggaaaaaa	attcggttaa	ggccaggggg	aaagaagaag	1020
tacaagctaa	agcacatcg	atgggcaagc	agggagctag	aacgattcgc	agttaatcct	1080
ggcctgttag	aaacatcaga	aggctgtaga	caaatactgg	gacagctaca	accatccctt	1140
cacagaggat	cagaggagct	tcgatcaacta	tacaacacag	tagcaaccct	ctattgtgt	1200
caccagcgga	tcgagatcaa	ggacaccaag	gaagctttag	acaagataga	ggaagagcaa	1260
aacaagtcca	agaagaaggc	ccagcaggca	gcagctgaca	caggacacag	caatcaggtc	1320
agccaaaatt	accctatagt	gcagaacatc	cagggcaaa	tggtacatca	ggccatatca	1380
cctagaactt	taaatgcatt	ggtaaaagta	gtagaagaga	aggcttcag	cccagaagt	1440
atacccatgt	tttcagcatt	atcagaagga	gccacccac	aggacctgaa	cacgatgtt	1500
acacccgtgg	ggggacatca	agcagccatg	caaatgttaa	aagagaccat	caatgaggaa	1560
gctgcagatt	gggatagagt	gcatccagt	catgcaggc	ctattgcacc	aggccagatg	1620
agagaaccaa	gggaaagtga	catagcagg	actactagta	cccttcagga	acaaatagga	1680
tggatgacaa	ataatccacc	tatcccagta	ggagagatct	acaagagg	gataatcctg	1740
ggattgaaca	agatcgtgag	gatgtatagc	cctaccagca	ttctggacat	aagacaagga	1800
ccaaaggaa	cctttagaga	ctatgttagac	cggttctata	aaactctaag	agctgagcaa	1860
gcttcacagg	aggtaaaaaa	ttggatgaca	gaaaccttg	tggtccaaa	tgcgaaccca	1920
gattgtaa	ccatcctgaa	ggctctcg	ccagcggcta	cactagaaga	aatgtgaca	1980

-continued

```

gcatgtcagg gagtaggagg acccgccat aaggcaagag tttggcgcc gcaaatccca 2040
ccaaactggc acataccgtc gatccaggac gcccgcacgc cttaccatcc cccggccacc 2100
ccgaacaaca tgggcctgat cgccggcgcg gtgggcggca gtctcctggc agccctggc 2160
atttgcgaa ttgtgtactg gatgcgccgc cgcaactcaa aagccccaaa gcgcatacgc 2220
ctcccccaca tccggaaaga cgaccagccg tcctcgacc agcccttgtt ttactag 2277

```

<210> SEQ ID NO 33  
<211> LENGTH: 758  
<212> TYPE: PRT  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: amino acid sequence of gDgagW294A

&lt;400&gt; SEQUENCE: 33

```

Met Gly Ala Ala Ala Arg Leu Gly Ala Val Ile Leu Phe Val Val
1           5          10          15

Ile Val Gly Leu His Gly Val Arg Gly Lys Tyr Ala Leu Ala Asp Ala
20          25          30

Ser Leu Lys Met Ala Asp Pro Asn Arg Phe Arg Gly Lys Asp Leu Pro
35          40          45

Val Leu Asp Gln Leu Thr Asp Pro Pro Gly Val Arg Arg Val Tyr His
50          55          60

Ile Gln Ala Gly Leu Pro Asp Pro Phe Gln Pro Pro Ser Leu Pro Ile
65          70          75          80

Thr Val Tyr Tyr Ala Val Leu Glu Arg Ala Cys Arg Ser Val Leu Leu
85          90          95

Asn Ala Pro Ser Glu Ala Pro Gln Ile Val Arg Gly Ala Ser Glu Asp
100         105         110

Val Arg Lys Gln Pro Tyr Asn Leu Thr Ile Ala Trp Phe Arg Met Gly
115         120         125

Gly Asn Cys Ala Ile Pro Ile Thr Val Met Glu Tyr Thr Glu Cys Ser
130         135         140

Tyr Asn Lys Ser Leu Gly Ala Cys Pro Ile Arg Thr Gln Pro Arg Trp
145         150         155         160

Asn Tyr Tyr Asp Ser Phe Ser Ala Val Ser Glu Asp Asn Leu Gly Phe
165         170         175

Leu Met His Ala Pro Ala Phe Glu Thr Ala Gly Thr Tyr Leu Arg Leu
180         185         190

Val Lys Ile Asn Asp Trp Thr Glu Ile Thr Gln Phe Ile Leu Glu His
195         200         205

Arg Ala Lys Gly Ser Cys Lys Tyr Ala Leu Pro Leu Arg Ile Pro Pro
210         215         220

Ser Ala Cys Leu Ser Pro Gln Ala Tyr Gln Gln Gly Val Thr Val Asp
225         230         235         240

Ser Ile Gly Met Leu Pro Arg Phe Ile Pro Glu Asn Gln Arg Thr Val
245         250         255

Ala Val Tyr Ser Leu Lys Ile Ala Gly Trp His Gly Pro Lys Ala Pro
260         265         270

Tyr Thr Ser Thr Leu Leu Pro Pro Glu Leu Ser Glu Thr Pro Asn Ala
275         280         285

Thr Gln Pro Glu Leu Ala Pro Glu Asp Pro Glu Asp Ser Ala Leu Leu
290         295         300

Glu Asp Pro Val Gly Thr Val Ala Pro Gly Ala Arg Ala Ser Val Leu

```

## US 12,390,525 B2

77

-continued

78

---

305	310	315	320													
Ser	Gly	Glu	Leu	Asp	Arg	Trp	Glu	Lys	Ile	Arg	Leu	Arg	Pro	Gly		
325							330							335		
Gly	Lys	Lys	Tyr	Lys	Leu	Lys	His	Ile	Val	Trp	Ala	Ser	Arg	Glu		
340							345							350		
Leu	Glu	Arg	Phe	Ala	Val	Asn	Pro	Gly	Leu	Leu	Glu	Thr	Ser	Glu	Gly	
355							360							365		
Cys	Arg	Gln	Ile	Leu	Gly	Gln	Leu	Gln	Pro	Ser	Leu	Gln	Thr	Gly	Ser	
370							375							380		
Glu	Glu	Leu	Arg	Ser	Leu	Tyr	Asn	Thr	Val	Ala	Thr	Leu	Tyr	Cys	Val	
385							390							400		
His	Gln	Arg	Ile	Glu	Ile	Lys	Asp	Thr	Lys	Glu	Ala	Leu	Asp	Lys	Ile	
405							410							415		
Glu	Glu	Glu	Gln	Asn	Lys	Ser	Lys	Lys	Ala	Gln	Gln	Ala	Ala	Ala		
420							425							430		
Asp	Thr	Gly	His	Ser	Asn	Gln	Val	Ser	Gln	Asn	Tyr	Pro	Ile	Val	Gln	
435							440							445		
Asn	Ile	Gln	Gly	Gln	Met	Val	His	Gln	Ala	Ile	Ser	Pro	Arg	Thr	Leu	
450							455							460		
Asn	Ala	Trp	Val	Lys	Val	Val	Glu	Glu	Lys	Ala	Phe	Ser	Pro	Glu	Val	
465							470							480		
Ile	Pro	Met	Phe	Ser	Ala	Leu	Ser	Glu	Gly	Ala	Thr	Pro	Gln	Asp	Leu	
485							490							495		
Asn	Thr	Met	Leu	Asn	Thr	Val	Gly	Gly	His	Gln	Ala	Ala	Met	Gln	Met	
500							505							510		
Leu	Lys	Glu	Thr	Ile	Asn	Glu	Glu	Ala	Ala	Asp	Trp	Asp	Arg	Val	His	
515							520							525		
Pro	Val	His	Ala	Gly	Pro	Ile	Ala	Pro	Gly	Gln	Met	Arg	Glu	Pro	Arg	
530							535							540		
Gly	Ser	Asp	Ile	Ala	Gly	Thr	Thr	Ser	Thr	Leu	Gln	Glu	Gln	Ile	Gly	
545							550							560		
Trp	Met	Thr	Asn	Asn	Pro	Pro	Ile	Pro	Val	Gly	Glu	Ile	Tyr	Lys	Arg	
565							570							575		
Trp	Ile	Ile	Leu	Gly	Leu	Asn	Lys	Ile	Val	Arg	Met	Tyr	Ser	Pro	Thr	
580							585							590		
Ser	Ile	Leu	Asp	Ile	Arg	Gln	Gly	Pro	Lys	Glu	Pro	Phe	Arg	Asp	Tyr	
595							600							605		
Val	Asp	Arg	Phe	Tyr	Lys	Thr	Leu	Arg	Ala	Glu	Gln	Ala	Ser	Gln	Glu	
610							615							620		
Val	Lys	Asn	Trp	Met	Thr	Glu	Thr	Leu	Leu	Val	Gln	Asn	Ala	Asn	Pro	
625							630							640		
Asp	Cys	Lys	Thr	Ile	Leu	Lys	Ala	Leu	Gly	Pro	Ala	Ala	Thr	Leu	Glu	
645							650							655		
Glu	Met	Met	Thr	Ala	Cys	Gln	Gly	Val	Gly	Gly	Pro	Gly	His	Lys	Ala	
660							665							670		
Arg	Val	Leu	Ala	Pro	Gln	Ile	Pro	Pro	Asn	Ala	His	Ile	Pro	Ser	Ile	
675							680							685		
Gln	Asp	Ala	Ala	Thr	Pro	Tyr	His	Pro	Pro	Ala	Thr	Pro	Asn	Asn	Met	
690							695							700		
Gly	Leu	Ile	Ala	Gly	Ala	Val	Gly	Gly	Ser	Leu	Leu	Ala	Ala	Leu	Val	
705							710							720		
Ile	Cys	Gly	Ile	Val	Tyr	Trp	Met	Arg	Arg	Arg	Arg	Thr	Gln	Lys	Ala	Pro
725							730							735		

-continued

Lys Arg Ile Arg Leu Pro His Ile Arg Glu Asp Asp Gln Pro Ser Ser  
 740 745 750

His Gln Pro Leu Phe Tyr  
 755

<210> SEQ ID NO 34  
 <211> LENGTH: 2277  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: nucleotide sequence encoding gDgagW294A

<400> SEQUENCE: 34

atgggggggg	ctgccgcac	gttggggggc	gtgattttgt	ttgtcgcat	agtgggcctc	60
catggggtcc	cgggcaata	tgcctggcg	gatgcctctc	tcaagatggc	cgaccggaaat	120
cgcgttcgcg	gcaaagacct	tccggtcctg	gaccagctga	ccgaccctcc	gggggtccgg	180
cgcgtgtacc	acatccaggc	gggccttaccg	gaccggttcc	agccccccag	cctcccgatc	240
acggtttaact	acgcccgtgtt	ggagcgcgcc	tgcgcgcagcg	tgctcctaaa	cgcaccgtcg	300
gaggcccccc	agattgtccg	cggggctcc	gaagacgtcc	ggaaacaacc	ctacaacctg	360
accatcgott	ggtttccgat	ggggaggcaac	tgtgtatacc	ccatcacggt	catggagtagc	420
accgaatgt	cctacaacaa	gtctctgggg	gcctgtccca	tccgaacgca	gccccgctgg	480
aactactatg	acagcttcag	cgcgcgtcago	gaggataacc	tggggttcct	gatgcacgcc	540
cccgcggtt	agaccgcccc	cacgtacctg	cggctcgta	agataaacga	ctggacggag	600
attacacagt	ttatcctgga	gcaccgaged	aagggtctcc	gtaagtacgc	cctcccgctg	660
cgcattcccc	cgtcagccctg	cctgtcccc	caggcctacc	agcaggggggt	gacgggtggac	720
agcatcgaaa	tgctgecccg	cttcattcccc	gagaaccaggc	gcaccgtcgc	cgtatacagc	780
ttgaagatcg	cgggggtggca	cggggccaag	gccccataca	cgagcaccc	gctgeccccc	840
gagctgtccg	agaccccaa	cgcacgcag	ccagaactcg	ccccggaaaga	ccccgaggat	900
tcggccctct	tggaggaccc	cgtggggacg	gtggcgccgg	gtgcgagagc	gtcagtattta	960
agcggggggag	aattagatcg	atggaaaaaa	attcggttaa	ggccaggggg	aaagaagaag	1020
tacaagctaa	agcacatcgt	atgggcaago	agggagctag	aacgattcgc	agttaatcct	1080
ggcctgttag	aaacatcaga	aggctgtaga	caaatactgg	gacagctaca	accatccctt	1140
cacacaggat	cagaggagct	tcgatacta	tacaacacag	tagcaaccct	ctattgttg	1200
caccagcgga	tcgagatcaa	ggacaccaag	gaagctttag	acaagataga	ggaagagcaa	1260
aacaagtcca	agaagaaggc	ccagcaggca	gcagctgaca	caggacacag	caatcaggtc	1320
agccaaaatt	accctatagt	gcagaacatc	caggggcaaa	tggtacatca	ggccatatca	1380
cctagaacctt	taaatgcatt	ggtaaaagta	gtagaagaga	aggctttcag	cccagaagtg	1440
ataccccatgt	tttcagcatt	atcagaagga	gccacccac	aggacctgaa	cacgatgtt	1500
aacaccgtgg	ggggacatca	agcagccatg	caaatgttaa	aagagaccat	caatgaggaa	1560
gctgcagatt	gggatagagt	gcatccatgt	catgcaggcc	ctattgcacc	aggccagatg	1620
agagaaccaa	ggggaaagtga	catagcagga	actactagta	cccttcagga	acaaataagga	1680
tggatgacaa	ataatccacc	tatcccgat	ggagagatct	acaagagggt	gataatcctg	1740
ggattgaaca	agatcgtgag	gatgtatago	cctaccagca	ttctggacat	aagacaagga	1800
ccaaaggaac	ccttagaga	ctatgttagac	cggttctata	aaactctaag	agctgagcaa	1860

-continued

---

gcttcacagg aggtaaaaaa ttggatgaca gaaaccttg tggtccaaaa tgcgAACCCA	1920
gattgttAAGA ccatcctgaa ggctctcgcc ccagcggcta cactagaaga aatgtatgaca	1980
gcattgtcagg gagtaggagg acccggccat aaggcaagag ttttggcgcc gcaaATCCCA	2040
ccaaACGCGC acataccgtc gatccaggac gcccgcacgc ctTaccatcc cccggccacc	2100
ccgaacaaca tgggcctgtat cgccggcgcg gtgggcggca gtctctggc agccctggc	2160
atTTGCGGAA ttgtgtactg gatgcggcgc cgcactcaaa aagccccaaa ggcatacgc	2220
ctcccccaaca tccgggaaga cgaccagccg tcctcgacc accccttgtt ttactag	2277
<210> SEQ ID NO 35	
<211> LENGTH: 2682	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: nucleotide sequence encoding gDNP	
<400> SEQUENCE: 35	
atgggggggg ctggcccaag gttggggggcc gtgattttgt ttgtcgcat agtggggctc	60
catgggggtcc ggggcaataa tgccttggcg gatgcctctc tcaagatggc cgaccccaat	120
cgttttcgat gcaaagacat tccgggtcctg gaccagctga ccgaccctcc ggggggtccgg	180
cgcgtgtacc acatccaggc gggcctaccg gaccgcgtcc agccccccag cctcccgatc	240
acgggtttact acggcggttt ggagcgccgc tgccgcagcg tgctcctaaa cgcaccgtcg	300
gaggcccccc agattgtccg cggggcctcc gaagacgtcc ggaaacaacc ctacaactg	360
accatcgctt ggtttccggat gggaggcaac tggctatcc ccatcacggt catggagtag	420
accgaatgtc cctacaacaa gtctctgggg gcctgtccca tccgaacgca gccccgtgg	480
aactactatg acagcttcag cgccgtcago gaggataacc tgggggtcct gatgcacgcc	540
cccgcggttg agaccgcccgg cacgtacctg cggctcgat gataaacgca ctggacggag	600
attacacagt ttatctggaa gcaccgagcc aagggtctt gtaagtacgc cctcccgctg	660
cgcacatcccc cgtcagccctg cctgtcccc caggccatacc agcaggggggt gacgggtggac	720
agcatcgatggaa tgctgtcccc cttcatcccc gagaaccaggc gcaccgtcgc cgtatacagc	780
ttgaagatcg cgggggtggca cggggcccgcc tcccaaggca ccaaacggtc ttacgaacag	840
atggagactg atggagaacg ccagaatgcc actgaaatca gaggcatccgt cggaaaaatg	900
attgggtggaa ttggacgatt ctacatccaa atgtgcacccg aactcaaact cagtgattat	960
gaggggacggt tgatccaaaa cagcttaaca atagagagaa tggtgctctc tgctttgac	1020
gaaaggagaa ataaataacct ggaagaacat cccagtgcgg gggaaagatcc taagaaaact	1080
ggaggaccta tatacaggag agtaaacggaa aagtggatga gagaactcat cctttatgac	1140
aaagaagaaa taaggcgaat ctggcgccaa actaataatg gtgacgtatgc aacggcttgt	1200
ctgactcaca tgatgtatcg gcattccat ttgaatgtatc caacttatca gaggacaaga	1260
gctcttgttc gcacccgaat ggatcccagg atgtgccttc tgatgcagg ttcaactctc	1320
ccttagggat ctggagccgc aggtgctgc gtcggatgggg ttggaaacaat ggtgtggaa	1380
ttgggtcagga tgatcaaaccg tgggtatcaat gatcggaaact tctggggggg cgagaatgg	1440
cgaaaaacaa gaattgctta tgaaagaatg tgcaacattc tcaaaggaa atttcaaact	1500
gctgcacaaa aagcaatgtatggatcaat ggtcaatgt agagagagcc ggaacccagg gaatgtcgat	1560
tgcgaagatc tcaactttctt acgtacggatc gcactcatat tgagagggc ggttgctcac	1620
aagtccgtcc tgcctgcctg tggatgtatggaa cctggccgtat ccagtgggtt ccactttgaa	1680

-continued

agagagggat	actctctagt	cggaatagac	ccttcagac	tgcttcaaaa	cagccaagtg	1740
tacagcctaa	tcagacccaaa	tgagaatcca	gcacacaaga	gtcaacttgt	gtggatggca	1800
tgcattctg	ccgcatttga	agatctaaga	gtattaagct	tcatcaaagg	gacgaagggt	1860
ctccccaaag	ggaagcttc	cactagagga	gttcaaattg	cttccaaatga	aaatatggat	1920
gttatggaa	taagtacact	tgaactgaga	agcaggtact	gggccataag	gaccagaagt	1980
gggggaaaca	ccaatcaaca	gagggcatct	gcgggccaaa	ttagcataca	acctacgttc	2040
tcagtacaga	gaaatctccc	tttgacaga	acaaccatta	tggcagcatt	caatggaaat	2100
acagaggaa	gaacatctga	catgaggacc	gaaatcataa	ggatgtatgg	aagtgcaga	2160
ccagaagatg	tgtcttcca	ggggggggga	gtcttcgagc	tctcgacga	aaaggcagcg	2220
agcccgatcg	tgccttcctt	tgacatgagt	aatgaaggat	cttattttctt	cgagacaat	2280
gcagaggagt	acgacaatgg	gcccaaggcc	ccatacacga	gcaccctgt	gccccggag	2340
ctgtccgaga	cccccaacgc	cacgcagcca	gaactcgccc	cggaagaccc	cgaggattcg	2400
gcctcttgg	aggacccgt	ggggacggtg	gcgcgcgaaa	tcccacccaa	ctggcacata	2460
cgtcgatcc	aggacgcccgc	gacgccttac	catccccgg	ccaccccgaa	caacatggc	2520
ctgatcgccg	ggcggtgggg	cggcagtc	ctggcagccc	tggcatttg	cggaatttg	2580
tactggatgc	gccgcggcac	tcaaaagcc	ccaaagcgca	tacgcctccc	ccacatccgg	2640
gaagacgacc	agccgttctc	gcaccagccc	ttgttttact	ag		2682

<210> SEQ ID NO 36  
 <211> LENGTH: 1185  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: nucleotide sequence encoding gDW294A

<400> SEQUENCE: 36

atgggggggg	ctgcccacag	gttgggggccc	gtgattttgt	ttgtcgcat	agtgggcctc	60
catgggttcc	cgggcaata	tgccttgggg	gatgcctctc	tcaagatggc	cgaccccaat	120
cgttttgcg	gaaaagacct	tccggctctg	gaccagctga	ccgaccctcc	gggggtccgg	180
cgcgtgtacc	acatccaggc	gggectacccg	gaccgcgtcc	agccccccag	cctcccgatc	240
acggtttact	acgcegtgtt	ggagcgcgc	tgcgcagcg	tgtctctaaa	cgcaccgtcg	300
gaggcccccc	agattgtccg	cggggcctcc	gaagacgtcc	ggaaacaacc	ctacaacctg	360
accatcgctt	ggtttcggat	gggaggcaac	tgtgctatcc	ccatcacggt	catggagtag	420
accgaatgct	cctacaacaa	gtctctgggg	gcctgtccca	tccgaacgca	gccccgtgg	480
aactactatg	acagcttcag	cgccgtcago	gaggataacc	tggggctct	gatgcacgccc	540
cccgcggtt	agaccgcgg	cacgtacctg	cggctcgatg	agataaacgca	ctggacggag	600
attacacagt	ttatccttgg	gcaccgagcc	aagggtctct	gtaagtacgc	cctcccgctg	660
cgcacatcccc	cgtcagccctg	cctgtcccc	caggccatacc	agcaggggggt	gacgggtggac	720
agcatcgaaa	tgctgcccc	cttcatcccc	gagaaccaggc	gcaccgtcg	cgtatacagc	780
ttgaagatcg	ccgggtggca	cgggcccaag	gccccataca	cgagcaccct	gctgcccccg	840
gagctgtccg	agaccccaa	cgccacgcag	ccagaactcg	ccccggaa	ccccgaggat	900
tccggccctct	tggaggaccc	cgtggggac	gtggcgccgc	aaatcccacc	aaacgcgcac	960
ataccgtcga	tccaggacgc	cgcgacgcct	taccatcccc	cggccacccc	gaacaacatg	1020

-continued

---

ggcctgatcg	ccggcgccgt	ggggggcagt	ctccctggcag	ccctggtcat	ttgcggatt	1080
gtgtactgga	tgcgcgcgg	cactaaaaaa	gccccaaagc	gcataacgcct	cccccacatc	1140
cgggaaagacg	accagecgtc	ctcgaccag	cccttgtttt	actag		1185
 <b>&lt;210&gt; SEQ ID NO 37</b>						
<b>&lt;211&gt; LENGTH: 2199</b>						
<b>&lt;212&gt; TYPE: DNA</b>						
<b>&lt;213&gt; ORGANISM: Artificial Sequence</b>						
<b>&lt;220&gt; FEATURE:</b>						
<b>&lt;223&gt; OTHER INFORMATION: nucleotide sequence encoding NBEF gDE7E6E5</b>						
 <b>&lt;400&gt; SEQUENCE: 37</b>						
atgggggggg	ctgcccggcag	gttgggggcc	gtgatttgt	ttgtcgatc	agtgggcctc	60
catggggtcc	cgggcaaaaata	tgccttggcg	gatgcctctc	tcaaggcgcc	cgaccccgct	120
cgctttcgcg	gcaaagacct	tccggtcgag	gacgcggcgg	ccgccccctcc	gggggtccgg	180
cgcgtgtacc	acatccaggc	gggccttaccg	gaccgcgtcc	agccccccag	cctcccgatc	240
acgggtttact	acgcggcggtt	ggagcgcgcgc	tgccgcgcgc	tgctccataaa	cgcaccgcgt	300
gaggcccccc	agattgtccg	cggggcctcc	gaagacgtcc	ggaaacaacc	ctacaacctg	360
accatcgctt	ggtttcggat	gggaggcaac	tgtgctatcc	ccatcacggt	catggagtag	420
accgaatgct	cctacaacaa	gtctctgggg	gcctgtccca	tccgaacgc	gcggcgctgg	480
aactactatg	acagcttcag	cgccgtcago	gaggataacc	tgggggtcct	gatgcacgccc	540
cccgcggtt	agaccgcgg	cacgtacctg	cggctcgta	agataaacga	ctggacggag	600
attacacagt	ttatcctgga	gcaccgagcc	aagggctcct	gtaagtacgc	cctcccgctg	660
cgcacatcccc	cgtcagccctg	cctgtccccc	caggcctacc	agcaggggg	gacgggtggac	720
agcatcgaaa	tgctgcccc	cttcatcccc	gagaaccagg	gcaccgtcgc	cgtatacagc	780
ttgaagatcg	ccgggtggca	cggggccat	ggagatacac	ctacattgca	tgaatatatg	840
tttagattgc	aaccagagac	aactgatctc	tactgttatg	agcaattaaa	tgacagctca	900
gaggaggagg	atgaaataga	tggccagct	ggacaaggcag	aaccggacag	agccattac	960
aatattgtaa	ccttttgtt	caagtgtac	tctacgttcc	ggttgtgcgt	acaaagcaca	1020
cacgtagaca	ttcgtacttt	ggaagacctg	ttaatggca	cactaggaat	tgtgtgcccc	1080
atctgttctc	agaaaccagg	cgcacccacaa	aagagaactg	caatgttca	ggacccacag	1140
gagcgaccca	gaaagttacc	acagttatgc	acagagctgc	aaacaactat	acatgtatata	1200
atattagaat	gtgtgtactg	caagcaacag	ttactgcgc	gtgaggtata	tgactttgt	1260
tttcgggatt	tatgcatagt	atataagat	gggaatccat	atgctgtatg	tgataaaatgt	1320
ttaaagtttt	attctaaaat	tagtgagat	agacattatt	gttataatgtt	gtatggaaaca	1380
acattagaac	agcaataacaa	caaaccgtt	tgtgatttgt	taattaggtg	tattaactgt	1440
caaaggccac	tgtgtctga	agaaaagcaa	agacatctgg	acaaaaagca	aagattccat	1500
aatataaggg	gtcggtggac	cggtcgtatgt	atgtcttgc	gcagatcatc	aagaacacgt	1560
agagaaaaccc	agctggcgcc	cgcctgcata	cacaacatta	ctggcgtgct	ttttgcttg	1620
ctttgtgtgc	ttttgtgtgt	ctgccttatta	atacgtccgc	tgctttgtc	tgtgtctaca	1680
tacacatcat	taataatatt	ggttacta	ttgtggataa	cagcagccctc	tgcgtttagg	1740
tgttttattt	tatataattat	atttgtttat	ataccattat	tttaataaca	tacacatgca	1800
cgcttttaa	ttacagggcc	caaggccccaa	tacacgagca	ccctgctgcc	cccgagctg	1860
tccgagaccc	ccaacgcccac	gcagccagaa	ctcgccccgg	aagaccccgaa	ggattcgcc	1920

---

ctcttggagg accccgtggg gacggtgccg cgc当地atcc caccataactg gcacataccg	1980
tgc当地tccagg acgccc当地cgc当地 gc当地taccat cccccggccca ccccgaaacaa catgggc当地g	2040
atcgccggcg cgg当地ggccgg cag当地tc当地ctg gc当地gccc当地gg tcat当地tggccg aattgtgtac	2100
tggatgccc gccc当地actca aaaagcccca aaggcc当地atac gc当地tccccc当地 catccggaa	2160
gacgaccagg cgtcc当地cgca ccagcccttg tttactag	2199

---

## What is claimed:

1. A method of inducing an immune response comprising providing to a subject in need thereof a vaccine comprising a nucleic acid molecule which encodes a fusion protein, wherein the fusion protein comprises:
  - a. a first polypeptide segment comprising at least amino acids 1-240 of a mature Herpes simplex virus (HSV) glycoprotein D, wherein the first polypeptide segment does not comprise a full length mature glycoprotein D;
  - b. a second polypeptide segment comprising at least one antigen, wherein the at least one antigen is not an HSV glycoprotein D antigen, wherein the N terminus of the second polypeptide segment is linked to the C terminus of the first polypeptide segment; and
  - c. a third polypeptide segment comprising a C terminal portion of the HSV glycoprotein D, wherein the N terminus of the third polypeptide segment is linked to the C terminus of the second polypeptide segment, wherein the at least one antigen is selected from the group consisting of: an influenza virus antigen; a nucleoprotein P influenza virus antigen, a *Plasmodium* antigen, a human papilloma virus (HPV) antigen, a human papilloma virus HPV16 antigen, an HPV E5 protein, an HPV E6 protein, an HPV E7 protein, a human immunodeficiency virus (HIV) antigen, and an HIV gag antigen.
2. The method of claim 1, wherein the HSV is selected from the group consisting of HSV-1 and HSV-2.
3. The method of claim 2, wherein the first polypeptide segment comprises an amino acid sequence selected from the group consisting of: amino acids 26-265 of SEQ ID NO: 27; amino acids 26-265 of SEQ ID NO: 29; amino acids 26-269 of SEQ ID NO: 27; amino acids 26-269 of SEQ ID NO: 29; amino acids 26-313 of SEQ ID NO: 27; amino acids 26-313 of SEQ ID NO: 29; amino acids 26-319 of SEQ ID NO: 27 with the exception that amino acid 319 is alanine instead of tryptophan; and amino acids 26-319 of SEQ ID NO: 29 with the exception that amino acid 319 is alanine instead of tryptophan.
4. The method of claim 3, wherein the first polypeptide segment is encoded by a nucleic acid sequence comprising nucleotides 76-795 of SEQ ID NO:26; or nucleotides 350-1069 of SEQ ID NO:28.
5. The method of claim 1, wherein the nucleic acid molecule encodes the amino acid sequence encoded SEQ ID NO:35.
6. The method of claim 1, wherein the nucleic acid molecule comprises a nucleotide sequence selected from the group consisting of SEQ ID NO:31; SEQ ID NO:32; SEQ ID NO: 34; SEQ ID NO:35; SEQ ID NO:36; and SEQ ID NO:37.
7. The method of claim 1, wherein the fusion protein comprises an amino acid sequence selected from the group consisting of SEQ ID NO:22; SEQ ID NO:23; SEQ ID NO:24; and SEQ ID NO:33.
8. The method of claim 1, wherein the nucleic acid molecule is in a viral vector.
9. The method of claim 1, wherein the nucleic acid molecule is naked DNA.
10. The method of claim 1, wherein the nucleic acid molecule is in a bacterial vector.
11. The method of claim 1, wherein the second polypeptide segment comprises the HPV E5 protein, the HPV E6 protein, and the HPV E7 protein.
12. The method of claim 1, wherein the third polypeptide segment comprises the transmembrane domain of the HSV glycoprotein D.
13. A method of inducing an immune response comprising providing to a subject in need thereof a vaccine comprising a fusion protein, wherein the fusion protein comprises:
  - a. a first polypeptide segment comprising at least amino acids 1-240 of a mature Herpes simplex virus (HSV) glycoprotein D, wherein the first polypeptide segment does not comprise a full length glycoprotein D;
  - b. a second polypeptide segment comprising at least one antigen, wherein the at least one antigen is not an HSV glycoprotein D antigen, wherein the N terminus of the second polypeptide segment is linked to the C terminus of the first polypeptide segment; and
  - c. a third polypeptide segment comprising a C terminal portion of the HSV glycoprotein D, wherein the N terminus of the third polypeptide segment is linked to the C terminus of the second polypeptide segment.
14. The method of claim 13, wherein the HSV is selected from the group consisting of HSV-1 and HSV-2.
15. The method of claim 14, wherein the first polypeptide segment comprises an amino acid sequence selected from the group consisting of amino acids 26-265 of SEQ ID NO: 27, amino acids 26-265 of SEQ ID NO: 29; amino acids 26-269 of SEQ ID NO: 27; amino acids 26-269 of SEQ ID NO: 29; amino acids 26-313 of SEQ ID NO: 27; amino acids 26-313 of SEQ ID NO: 29; amino acids 26-319 of SEQ ID NO: 27 with the exception that amino acid 319 is alanine instead of tryptophan; and amino acids 26-319 of SEQ ID NO: 29 with the exception that amino acid 319 is alanine instead of tryptophan.
16. The method of claim 15, wherein the first polypeptide segment is encoded by a nucleic acid sequence comprising nucleotides 76-795 of SEQ ID NO:26; or nucleotides 350-1069 of SEQ ID NO:28.
17. The method of claim 13, wherein the at least one antigen is selected from the group consisting of an influenza virus antigen, a nucleoprotein P influenza virus antigen, a *Plasmodium* antigen, a human papilloma virus (HPV) antigen, human papilloma virus HPV16 antigen, HPV E5 protein, HPV E6 protein, HPV E7 protein, a human immunodeficiency virus (HIV) antigen, and an HIV gag antigen.

18. The method of claim 13, wherein the fusion protein comprises an amino acid sequence selected from the group consisting of SEQ ID NO:22; SEQ ID NO:23; SEQ ID NO:24; and SEQ ID NO:33.

19. The method of claim 13, wherein the fusion protein is encoded by a nucleic acid sequence selected from the group consisting of SEQ ID NO:31; SEQ ID NO:32; SEQ ID NO:34; SEQ ID NO:35; SEQ ID NO:36; and SEQ ID NO:37.

20. The method of claim 17, wherein the second polypeptide segment comprises the HPV E5 protein, the HPV E6 protein, and the HPV E7 protein.

21. The method of claim 13, wherein the third polypeptide segment comprises the transmembrane domain of the HSV glycoprotein D.

22. The method of claim 1, wherein the *Plasmodium* antigen selected from the group consisting of thrombospondin-related anonymous protein (TRAP), ring-infected erythrocyte surface antigen (RESA), merozoite surface protein 1 (MSP1), merozoite surface protein 2 (MSP2), merozoite surface protein 3 (MSP3), and glutamate-rich antigen (GLURP).

23. The method of claim 17, wherein a *Plasmodium* antigen selected from the group consisting of thrombospondin-related anonymous protein (TRAP), ring-infected erythrocyte surface antigen (RESA), merozoite surface protein 1 (MSP1), merozoite surface protein 2 (MSP2), merozoite surface protein 3 (MSP3), and glutamate-rich antigen (GLURP).

\* \* \* \* \*