

(12) United States Plant Patent

Danziger

US PP36.885 P2 (10) Patent No.:

(45) Date of Patent: Aug. 12, 2025

(54) IMPATIENS PLANT NAMED 'DIMPSLPRRD'

Latin Name: Impatiens hawkeri Varietal Denomination: **DIMPSLPRRD**

(71) Applicant: Danziger "DAN" Flower Farm, Beit Dagan (IL)

Inventor: Gavriel Danziger, Beit Dagan (IL)

Assignee: Danziger "DAN" Flower Farm (IL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

(21) Appl. No.: 19/026,197

(22) Filed: Jan. 16, 2025

(51) **Int. Cl.**

A01H 5/02 (2018.01)A01H 6/16 (2018.01)

(52) U.S. Cl.

USPC Plt./318.7

(58) Field of Classification Search

USPC Plt./263.1, 317, 318.1, 318.7 See application file for complete search history.

(56)References Cited

U.S. PATENT DOCUMENTS

PP25,015 P3 * 10/2014 Sato A01H 5/02 Plt./318.7

* cited by examiner

Primary Examiner — Karen M Redden (74) Attorney, Agent, or Firm — Cassandra Bright

ABSTRACT

A new and distinct cultivar of Impatiens plant named 'DIMPSLPRRD' is disclosed, characterized by large red flowers, and dark olive-green foliage. Plants are sun-resistant, have mounded habits and are well-branched. The new variety is an Impatiens, typically produced as an ornamental

2 Drawing Sheets

1

Latin name of the genus and species: Impatiens hawkeri. Variety denomination: 'DIMPSLPRRD'.

BACKGROUND OF THE INVENTION

The new Impatiens cultivar is a product of a planned breeding program conducted by the inventor, Gavriel Danziger. The objective of the breeding program was to produce useful new Impatiens varieties for ornamental horticultural production. The new variety is the result of the July 2020 crossing of the seed parent, an unpatented, proprietary variety referred to as Impatiens 'BM-18-1792', and the pollen parent, an unpatented, proprietary variety referred to as Impatiens 'BM-17-1438'. The new variety was discovered by the inventor in July of 2020 at a nursery in Moshav Mishmar Hashiva, Israel.

Asexual propagation of Impatiens 'DIMPSLPRRD' was first performed by leaf cuttings in July of 2020, at the same nursery in Moshav Mishmar Hashiva, Israel. This and sub- 20 sequent propagation has shown that the unique features of this cultivar are stable and reproduced true to type in multiple successive generations.

SUMMARY OF THE INVENTION

The cultivar 'DIMPSLPRRD' has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature, day length, and light intensity, without, how- 30 ever, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'DIMPSL-PRRD'. These characteristics in combination distinguish 'DIMPSLPRRD' as a new and distinct *Impatiens* cultivar:

2

- 1. Large, red flowers.
- 2. Well branched plants.
- 3. Sun-resistant plants.

PARENT COMPARISON

Plants of 'DIMPSLPRRD' are similar in some horticultural characteristics to the seed parent. The new variety, however, differs in the following:

- 1. Flower color of the new variety is darker than the flower color of the seed parent.
- 2. Flowers of the new variety are larger than flowers of the seed parent.
- 3. Plants of the new variety are larger than plants of the seed parent.
- 4. Flower color of the new variety is more stable than the flower color of the seed parent.

Plants of 'DIMPSLPRRD' are similar in some horticultural characteristics to the pollen parent. The new variety, however, differs in the following:

- 1. Flowers of the new variety are larger than flowers of the pollen parent.
- 2. Plants of the new variety are larger than plants of the pollen parent.
- 3. Plants of the new variety has more branching than plants of the pollen parent.

COMMERCIAL COMPARISONS

Plants of 'DIMPSLPRRD' are similar in some horticultural characteristics to the variety *Impatiens* 'SAKIMP030', U.S. Plant Pat. No. 25,015. The new variety, however, differs in the following:

1. Leaf color of the new variety is lighter than the leaf color of this comparator.

3

- 2. Flowers of the new variety are smaller than flowers of this comparator.
- 3. Plants of the new variety are larger than plants of this comparator.

Plants of 'DIMPSLPRRD' are similar in some horticultural characteristics to the variety *Impatiens* 'SAKIMP060', U.S. Pat. No. 11,076,549. The new variety, however, differs in the following:

- 1. Leaf color of the new variety is lighter than the leaf color of this comparator.
- Flowers of the new variety are larger than flowers of this comparator.
- 3. Plants of the new variety are larger than plants of this comparator.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photograph in FIG. 1 illustrates in full color a typical plant of 'DIMPSLPRRD' at about 13 weeks of age, grown in a 15 cm pot.

FIG. 2 illustrates a close-up view of a typical flower of the new variety.

The photographs were taken using conventional techniques and although colors may appear different from actual 25 colors due to light reflectance it is as accurate as possible by conventional photographic techniques.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Mini Colour Chart 2005 except where general terms of ordinary dictionary significance are used. The following observations and measurements describe 'DIMPSLPRRD' plants grown in a greenhouse in Moshav Mishmar Hashiva, Israel, under natural lighting. Measurements were taken during autumn. The growing temperature ranged from approximately 20° C. to 30° C. during the days, 15° C. to 25° C. during the nights. Measurements and numerical values represent averages of typical plant types.

Botanical classification: *Impatiens hawkeri* 'DIMPSL-PRRD'.

PROPAGATION

Time to initiate rooting: 10 to 14 days in summer; 14 to 20 days in winter.

Age of plant described: 12 weeks from a rooted cutting. Root description: Fibrous, adventitious roots.

PLANT

Growth habit: Mounded.

Pot size of plant described: 12 cm.

Height: 20 to 25 cm. Plant spread: 30 to 35 cm.

Growth rate: Medium.
Branching characteristics: Medium branched.

Length of lateral branches: 15 to 20 cm. Quantity of lateral branches: 5 to 8. Characteristics of primary lateral branches:

Shape.—Oval.

Diameter.—1 cm.

Texture.—Smooth.

Strength.—Strong, very stable. Color.—RHS Yellow-Green 147C. Internode length: 4 to 5 cm.

FOLIAGE

Leaf:

15

Arrangement.—Alternate.

Quantity.—Approximately 25 to 30 per branch.

Average length.—7 cm. Average width.—4 cm.

Pubescence.—None.

Shape.—Lanceolate. Apex.—Apiculate.

Base.—Acuminate.

Margin.—Ciliate.

Texture.—Smooth.

Aspect.-45°.

Color.—Young Foliage, Upper Side: RHS Yellow-Green 147A. Young Foliage, Under Side: RHS Yellow-Green 147C. Mature Foliage, Upper Side: RHS Yellow-Green 147A. Mature Foliage, Under Side: RHS Yellow-Green 147C.

Venation.—Type: Pinnate. Color, Upper Side: RHS Yellow-Green 147D. Color, Under Side: RHS Yellow-Green 147C.

Petiole:

30

Length.—0.5 to 1 cm.

Diameter.—0.2 to 0.3 cm.

Color.—RHS Yellow-Green 145B.

Texture.—Smooth.

FLOWER

Natural flowering season: Flowers at any day length. All year round in a greenhouse environment. Flowering is continuous from spring until fall in an outdoor garden.

Days to flowering from rooted cutting: 10 weeks.

Inflorescence and flower type and habit: Single flowers raised above the foliage. Corolla consists of 5 petals.

Rate of flower opening: 10 to 15 days from bud to fully opened flower.

Flower longevity on plant: 5 to 8 days.

Persistent or self-cleaning: Self-cleaning.

45 Bud:

Shape.—Ovate.

Length.—1.5 cm.

Diameter.—1 cm.

Color.—RHS Red-Orange N34A.

50 Flower size:

Diameter.—6 cm.

Depth.—6.5 cm including spur.

Petals:

65

Arrangement.—Overlapping.

Length.—3.5 cm.

Width.—4.5 cm.

Quantity.—5.

Texture.—Smooth.

Apex.—Heart-shaped.

Apex.—neart-snape

Base.—Acute.

Shape.—Obcordate.

Margin.—Entire.

Color.—When Opening, Upper Surface: RHS Red 45B. When Opening, Lower Surface: RHS Red 47C. Fully Opened, Upper Surface: RHS Red 45B. Fully Opened, Lower Surface: RHS Red 47C.

6 5

| Spurs: | | Quantity.—5, with anthers that are more or less fused, |
|---|----|---|
| Quantity.—1 per flower. | | forming a cap over the ovary. |
| Length.—5.5 cm. | | Anthers: |
| Diameter.—0.2 cm. | | Shape.—Oblong. |
| Aspect.—Spur curvature 40° to 45° from peduncle. | 5 | Length.—About 0.5 cm. |
| Color.—RHS Red 45C towards base; Red 45C at apex. | | Color.—RHS Red-Orange 43B. |
| Sepals: | | Pollen.—None. |
| Quantity.—2 per flower. | | Gynoecium: |
| Shape.—Oblong. | | Pistil: |
| Length.—1 cm. | 10 | Quantity.—1. |
| <i>Width.</i> —0.4 cm. | | Length.—0.5 cm. |
| Apex.—Caudate. | | Style.—Length: 0.5 cm. Color: RHS Yellow-Green |
| Base.—Rounded. | | 141B. |
| Margin.—Entire. | | Stigma.—Shape: Round. Color: RHS Yellow-Green |
| Color.—RHS Yellow-Green 145B. | 15 | 149D. Ovary Color: RHS Yellow-Green 144D. |
| Peduncles: | | |
| Length.—5 to 5.5 cm. | | OTHER CHARACTERISTICS |
| <i>Width.</i> —0.2 cm. | | |
| Angle.—About 45° to lateral branch. | | Seeds and fruits: 5 to 10 seeds per fruit. |
| Strength.—Stable, flexible. | 20 | Disease/pest resistance: Neither resistance nor susceptibility |
| Texture.—Smooth. | | of diseases common to <i>Impatiens</i> observed. |
| Color.—RHS Yellow-Green 145B. | | Temperature tolerance: Tolerant at 30° C. and 5° C. Grows |
| Fragrance: None. | | best at 10° C. to 28° C. |
| REPRODUCTIVE ORGANS | 25 | What is claimed is: 1. A new and distinct cultivar of <i>Impatiens</i> plant named |
| Androecium: | | 'DIMPSLPRRD' as herein illustrated and described. |

Androecium:

Stamens:






FIG. 2