

# US Patent & Trademark Office

## Patent Public Search | Text View

---

United States Patent Application Publication	20250262187
Kind Code	A1
Publication Date	August 21, 2025
Inventor(s)	LAYZELL; Marie Claire et al.

---

### INJECTABLE FORMULATIONS

---

#### Abstract

Provided herein are pharmaceutical formulations, methods for their production, and uses thereof. The pharmaceutical formulations comprise a salt of an optionally substituted dimethyltryptamine compound, a buffer, which is separate to the salt, and water. The formulations have pH values of from about 3.5 to about 6.5 and osmolalities of about 250 to about 350 mOsm/Kg. Such formulations are suitable for injection, being both stable and clinically acceptable, and have potential uses in the treatment of psychiatric or neurological disorders.

---

**Inventors:** LAYZELL; Marie Claire (London, GB), RENNIE; James Maxwell (London, GB)

**Applicant:** CYBIN UK LTD (London, GB)

**Family ID:** 1000008586655

**Assignee:** CYBIN UK LTD (London, GB)

**Appl. No.:** 19/202059

**Filed:** May 08, 2025

#### Foreign Application Priority Data

GB	2013571.1	Aug. 28, 2020
----	-----------	---------------

#### Related U.S. Application Data

parent US continuation 17806526 20220613 parent-grant-document US 12343327 child US 19202059

parent US continuation 17459284 20210827 parent-grant-document US 11406619 child US 17806526

parent US continuation-in-part 17006115 20200828 ABANDONED child US 17459284

---

## Publication Classification

**Int. Cl.:** A61K31/4045 (20060101); A61K47/02 (20060101); A61K47/12 (20060101)

**U.S. Cl.:**

**CPC** A61K31/4045 (20130101); A61K47/02 (20130101); A61K47/12 (20130101);

---

## Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS [0001] This application is a continuation of U.S. patent application Ser. No. 17/806,526 filed Jun. 13, 2022, which is a continuation of U.S. patent application Ser. No. 17/459,284 filed Aug. 27, 2021, which is a continuation in part of U.S. patent application Ser. No. 17/006,115, filed Aug. 28, 2020, and claims benefit to United Kingdom Application No. 2013571.1, filed Aug. 28, 2020, the entire disclosures of each of which are incorporated by reference herein.

### BACKGROUND

[0002] Classical psychedelics have shown preclinical and clinical promise in treating psychiatric disorders (Carhart-Harris and Goodwin, *Neuropsychopharmacology* 42, 2105-2113 (2017)). In particular, psilocybin has demonstrated significant improvement in a range of depression and anxiety rating scales in randomised double blind studies (Griffiths et al. *Journal of Psychopharmacology*, 30 (12), 1181-1197 (2016)).

[0003] N,N-dimethyltryptamine (DMT) is also understood to hold therapeutic value as a short-acting psychedelic. A review of research into the biosynthesis and metabolism of DMT in the brain and peripheral tissues, methods and results for DMT detection in body fluids and the brain, new sites of action for DMT, and new data regarding the possible physiological and therapeutic roles of DMT is provided by S. A. Barker in *Front. Neurosci.*, 12, 536, 1-17 (2018). In this review, DMT is described as having a possible therapeutic role in the treatment of depression, obsessive-compulsive disorder, and substance abuse disorders.

[0004] The injection of saline solutions of DMT fumarate salts into human volunteers is described in C. Timmermann et al., *Sci. Rep.*, 9, 16324 (2019). The effect of DMT fumarate on the power spectrum and signal diversity of human brain activity was recorded via multivariate EEG and compared with the results obtained on injection of a placebo (saline solution). It was found that, relative to the results obtained with the placebo, DMT fumarate suppressed alpha power and normalized/increased delta and theta power. Alpha power has been linked with high-level psychological functioning, top-down predictive processing and related feedback connectivity, whilst theta and delta power is classically associated with REM sleep dreaming and related 'visionary' states. It is described that these results relate injection of DMT fumarate to the experience of feeling profoundly immersed in an entirely other world.

[0005] According to the Human Metabolome Database (HMDB), dimethyltryptamine degrades relatively quickly in solution (see specifically <http://www.hmdb.ca/metabolites/HMDB0005973>). Consequently, there is a need in the art for injectable solutions of DMT that are stable over longer periods of time, and are clinically acceptable. The present invention addresses this need.

### SUMMARY

[0006] The present invention relates to pharmaceutical formulations, methods for their production, and uses thereof. The pharmaceutical formulations comprise a salt of an optionally substituted dimethyltryptamine compound, a buffer, which is separate to the salt, and water. The formulations have pH values of from about 3.5 to about 6.5 and osmolalities of about 250 to about 350

mOsm/Kg. Such formulations are suitable for injection, being both stable and clinically acceptable, and have potential uses in the treatment of psychiatric or neurological disorders. Certain aspects and embodiments are described herein and further aspects and embodiments will be evident from the discussion that follows below.

---

## Description

### DETAILED DESCRIPTION

[0007] Throughout this specification, one or more aspects of the invention may be combined with one or more features described in the specification to define distinct embodiments of the invention.

[0008] In the discussion that follows, reference is made to a number of terms, which are to be understood to have the meanings provided below, unless a context expressly indicates to the contrary. The nomenclature used herein for defining compounds, in particular the compounds described herein, is intended to be in accordance with the rules of the International Union of Pure and Applied Chemistry (IUPAC) for chemical compounds, specifically the “IUPAC Compendium of Chemical Terminology (Gold Book)” (see A. D. Jenkins et al., Pure & Appl. Chem., 1996, 68, 2287-2311). For the avoidance of doubt, if a rule of the IUPAC organisation is contrary to a definition provided herein, the definition herein is to prevail.

[0009] References herein to a singular of a noun encompass the plural of the noun, and vice-versa, unless the context implies otherwise.

[0010] Throughout this specification the word “comprise”, or variations such as “comprises” or “comprising”, will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

[0011] The term “consisting” or variants thereof is to be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, and the exclusion of any other element, integer or step or group of elements, integers or steps.

[0012] The term “about” herein, when qualifying a number or value, is used to refer to values that lie within  $\pm 5\%$  of the value specified. For example, if a pH range is specified to be about 3.5 to about 6.5, pH values of 3.3 to 6.8 are included.

[0013] The formulations of the invention are useful in therapy and may be administered to a patient in need thereof. As used herein, the term ‘patient’ preferably refers to a mammal. Typically the mammal is a human, but may also refer to a domestic mammal. The term does not encompass laboratory mammals.

[0014] The term “treatment” defines the therapeutic treatment of a patient, in order to reduce or halt the rate of progression of a disorder, or to ameliorate or cure the disorder. Prophylaxis of a disorder as a result of treatment is also included. References to prophylaxis are intended herein not to require complete prevention of a disorder: its development may instead be hindered through treatment in accordance with the invention. Typically, treatment is not prophylactic, and the formulation is administered to a patient having a diagnosed or suspected disorder.

[0015] As is understood in the art, psychiatric or neurological disorders are disorders which may be associated with one or more cognitive impairment. As used herein, the term ‘psychiatric disorder’ is a clinically significant behavioural or psychological syndrome or pattern that occurs in an individual and that is associated with present distress (e.g., a painful symptom) or disability (i.e., impairment in one or more important areas of functioning) or with a significantly increased risk of suffering death, pain, disability, or an important loss of freedom.

[0016] Diagnostic criteria for psychiatric or neurological disorders referred to herein are provided in the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, (DSM-5).

[0017] As used herein the term ‘obsessive-compulsive disorder’ (OCD) is defined by the presence

of either obsessions or compulsions, but commonly both. The symptoms can cause significant functional impairment and/or distress. An obsession is defined as an unwanted intrusive thought, image or urge that repeatedly enters the person's mind. Compulsions are repetitive behaviours or mental acts that the person feels driven to perform. Typically, OCD manifests as one or more obsessions, which drive adoption of a compulsion. For example, an obsession with germs may drive a compulsion to clean or an obsession with food may drive a compulsion to overeat, eat too little or throw up after eating (i.e. an obsession with food may manifest itself as an eating disorder). A compulsion can either be overt and observable by others, such as checking that a door is locked, or a covert mental act that cannot be observed, such as repeating a certain phrase in one's mind.

[0018] The present invention relates to pharmaceutical formulations suitable for injection, comprising a salt of an optionally substituted dimethyltryptamine compound, a buffer, which is separate to the salt, and water, wherein the formulations have pH values of from about 3.5 to about 6.5 and typical osmolalities of about 250 to about 350 mOsm/Kg.

[0019] Human blood serum has a pH of about 7.4 (typically ranging between 7.35 to 7.45, see G. K. Shwalfenberg, *J. Environ. Public Health*, 2012; 2012:727630), and the obvious go-to formulation of salts of optionally substituted dimethyltryptamine compounds is isotonic with a pH of 7.4. It has now been found that formulations described in the prior art or adapted therefrom have non-optimal shelf-life when stored under ambient conditions. The present invention addresses this problem of providing pharmaceutical formulations suitable for injection with substantially reduced degradation products compared with known formulations when stored under stressed conditions. This is indicative of improved shelf-life over such pharmaceutical formulations described in the prior art.

[0020] The invention provides a formulation or kit according to the first and second aspect of the invention for use in a method of treating an eating disorder. The term “eating disorder” includes anorexia nervosa, bulimia and binge eating disorder (BED). The symptoms of anorexia nervosa include eating too little and/or exercising too much in order to keep weight as low as possible. The symptoms of bulimia include eating a lot of food in a very short amount of time (i.e. bingeing) and then being deliberately sick, using laxatives, eating too little and/or exercising too much to prevent weight gain. The symptoms of BED include regularly eating large portions of food until uncomfortably full, and consequently feeling upset or guilty.

[0021] As used herein the term ‘depressive disorder’ includes major depressive disorder, persistent depressive disorder, bipolar disorder, bipolar depression, and depression in terminally ill patients.

[0022] As used herein the term ‘major depressive disorder’ (MDD, also referred to as major depression or clinical depression) is defined as the presence of five or more of the following symptoms over a period of two-weeks or more (also referred to herein as a ‘major depressive episode’), most of the day, nearly every day: [0023] depressed mood, such as feeling sad, empty or tearful (in children and teens, depressed mood can appear as constant irritability); [0024] significantly reduced interest or feeling no pleasure in all or most activities; [0025] significant weight loss when not dieting, weight gain, or decrease or increase in appetite (in children, failure to gain weight as expected); [0026] insomnia or increased desire to sleep; [0027] either restlessness or slowed behaviour that can be observed by others; [0028] fatigue or loss of energy; [0029] feelings of worthlessness, or excessive or inappropriate guilt; [0030] trouble making decisions, or trouble thinking or concentrating; [0031] recurrent thoughts of death or suicide, or a suicide attempt. [0032] At least one of the symptoms must be either a depressed mood or a loss of interest or pleasure.

[0033] Persistent depressive disorder, also known as dysthymia, is defined as a patient exhibiting the following two features: [0034] A. has depressed mood for most the time almost every day for at least two years. Children and adolescents may have irritable mood, and the time frame is at least one year. [0035] B. While depressed, a person experiences at least two of the following symptoms: [0036] Either overeating or lack of appetite. [0037] Sleeping too much or having difficulty

sleeping. [0038] Fatigue, lack of energy. [0039] Poor self-esteem. [0040] Difficulty with concentration or decision-making.

[0041] As used herein the term ‘treatment resistant major depressive disorder’ describes MDD that fails to achieve an adequate response to an adequate treatment with standard of care therapy.

[0042] As used herein, ‘bipolar disorder’, also known as manic-depressive illness, is a disorder that causes unusual shifts in mood, energy, activity levels, and the ability to carry out day-to-day tasks.

[0043] There are two defined sub-categories of bipolar disorder; all of them involve clear changes in mood, energy, and activity levels. These moods range from periods of extremely “up,” elated, and energised behaviour (known as manic episodes, and defined further below) to very sad, “down,” or hopeless periods (known as depressive episodes). Less severe manic periods are known as hypomanic episodes.

[0044] Bipolar I Disorder—defined by manic episodes that last at least 7 days, or by manic symptoms that are so severe that the person needs immediate hospital care. Usually, depressive episodes occur as well, typically lasting at least 2 weeks. Episodes of depression with mixed features (having depression and manic symptoms at the same time) are also possible.

[0045] Bipolar II Disorder—defined by a pattern of depressive episodes and hypomanic episodes, but not the full-blown manic episodes described above.

[0046] As used herein ‘bipolar depression’ is defined as an individual who is experiencing depressive symptoms with a previous or coexisting episode of manic symptoms, but does not fit the clinical criteria for bipolar disorder.

[0047] As used herein, the term ‘anxiety disorder’ includes generalised anxiety disorder, phobia, panic disorder, social anxiety disorder, and post-traumatic stress disorder.

[0048] ‘Generalised anxiety disorder’ (GAD) as used herein means a chronic disorder characterised by long-lasting anxiety that is not focused on any one object or situation. Those suffering from GAD experience non-specific persistent fear and worry, and become overly concerned with everyday matters. GAD is characterised by chronic excessive worry accompanied by three or more of the following symptoms: restlessness, fatigue, concentration problems, irritability, muscle tension, and sleep disturbance.

[0049] ‘Phobia’ is defined as a persistent fear of an object or situation the affected person will go to great lengths to avoid, typically disproportional to the actual danger posed. If the feared object or situation cannot be avoided entirely, the affected person will endure it with marked distress and significant interference in social or occupational activities.

[0050] A patient suffering from a ‘panic disorder’ is defined as one who experiences one or more brief attack (also referred to as a panic attack) of intense terror and apprehension, often marked by trembling, shaking, confusion, dizziness, nausea, and/or difficulty breathing. A panic attack is defined as a fear or discomfort that abruptly arises and peaks in less than ten minutes.

[0051] ‘Social anxiety disorder’ is defined as an intense fear and avoidance of negative public scrutiny, public embarrassment, humiliation, or social interaction. Social anxiety often manifests specific physical symptoms, including blushing, sweating, and difficulty speaking.

[0052] ‘Post-traumatic stress disorder’ (PTSD) is an anxiety disorder that results from a traumatic experience. Post-traumatic stress can result from an extreme situation, such as combat, natural disaster, rape, hostage situations, child abuse, bullying, or even a serious accident. Common symptoms include hypervigilance, flashbacks, avoidant behaviours, anxiety, anger and depression.

[0053] As used herein, the term “post-partum depression” (PPD, also known as postnatal depression) is a form of depression experienced by either parent of a newborn baby. Symptoms typically develop within 4 weeks of delivery of the baby and often include extreme sadness, fatigue, anxiety, loss of interest or pleasure in hobbies and activities, irritability, and changes in sleeping or eating patterns.

[0054] As used herein, the term ‘substance abuse’ means a patterned use of a drug in which the user consumes the substance in amounts or with methods that are harmful to themselves or others.

[0055] As used herein, the term ‘an avolition disorder’ refers to a disorder that includes as a symptom the decrease in motivation to initiate and perform self-directed purposeful activities.

[0056] The invention provides a pharmaceutical formulation suitable for injection, comprising a salt of a dimethyltryptamine (DMT) compound optionally substituted with deuterium and optionally substituted at position 4 or 5 with acetoxy or methoxy or position 4 with monohydrogen phosphate; a buffer which is separate to the salt; and water, wherein the formulation has a pH of about 3.5 to about 6.5 and an osmolality of about 250 to about 350 mOsm/Kg.

[0057] Owing to the instability of dimethyltryptamine in solution, solutions comprising dimethyltryptamine are generally prepared immediately before or close to the time of use, i.e. storage of solutions of dimethyltryptamine is avoided. Alternatively, solutions of dimethyltryptamine are frozen. The inventors have found that when a buffer, which is separate to the salt, is used, the resultant formulations are more stable than formulations prepared without a buffer separate to the salt. In addition, when a container adapted to prevent penetration of ultraviolet light is used, the resultant formulations are more stable than those stored in containers that allow for ultraviolet light penetration.



[0058] The inventors have found that the formulation is surprisingly more stable than formulations prepared at higher pH (specifically those prepared at a pH matching human blood serum, i.e. at a pH of about 7.4). The greater stability of the formulation of the invention relative to the go-to formulation is discussed in more detail in the Example section.

[0059] Osmolality is formally defined as the quotient of the negative natural logarithm of the rational activity of water and the molar mass of water, as represented by formula:

$$[00001] \text{osmolality} = \frac{-\ln a_w}{18.015}; a_w = \frac{p}{p^*}$$

where p is the partial vapour pressure of water in the solution and p\* is the partial vapour pressure of pure water. In simpler terms, osmolality is the number of osmotically active particles (the number of solute particles) in 1 kg of a solution. Thus, osmolality is a function only of the number of particles, and is not related to particle molecular weight, size, shape, or charge (see D. K. Faria et al., M. E. Mendes and N. M. Sumita, *J. Bras. Patol. Med. Lab.*, 53, 1, 38-45 (2017) for a review of the measurement of serum osmolality). For example, one mole of a nondissociating substance (e.g. DMT as a free base) dissolved in 1 kg of water has an osmolality of 1 Osm/kg (1000 mOsm/kg), whilst one mole of a substance that dissociates into two separate species in solution (e.g. DMT fumarate) dissolved in 1 kg of water has an osmolality of 2 Osm/kg (2000 mOsm/kg).

[0060] Where a first solution is defined herein to be isotonic with a second solution, the solutions have the same osmolality. For example, where a formulation is defined to be isotonic with human blood serum, the formulation has the same osmolality as human blood serum. Human blood serum typically has an osmolality of about 275 to about 300 mOsm/Kg (L. Hooper et al., *BMJ Open*, 2015; 5 (10): e008846).

[0061] The formulation (i.e. of the invention) is suitable for injection, by which is meant that it is in accordance with Pharmacopeial requirements of sterility, contaminants, and pyrogens (see for example The United States Pharmacopeial Convention, General Requirements/custom-character1 custom-character Injections, page 33). Sometimes, the formulation contains inhibitors of the growth of microorganisms (e.g. antimicrobial preservatives) and/or anti-oxidants.

[0062] Formulations suitable for injection have a pH of about 3 to 9 and an osmolality of about 250 to about 600 mOsm/Kg. pH values above 9 are reported by I. Usach et al. in *Adv. Ther.*, 36, 2986-2996 (2019) to relate to tissue necrosis (death of cells within the tissue), whereas values lower than 3 are reported to cause pain and phlebitis (inflammation of veins). Osmolality values greater than 600 mOsm/Kg are also reported to cause pain. The pH and osmolality of the formulation of the invention lie within the ranges reported to be suitable for injection.

[0063] The formulation comprises a salt of a DMT compound optionally substituted with deuterium and optionally substituted at position 4 or 5 with acetoxy or methoxy or position 4 with

monohydrogen phosphate, referred to herein as “the DMT compound”. Formulations in accordance with this invention may comprise one or more than one DMT compound. For the avoidance of doubt, formulations comprise an optionally substituted DMT salt when they comprise ions of optionally substituted DMT and ions that counter the charge of the optionally substituted DMT ions (counterions). Accordingly, the optionally substituted DMT salt within the formulation may be formed, for example, by contacting optionally substituted DMT as a free base with an aqueous solution comprising an excess of buffer relative to the molar quantity of optionally substituted DMT.

[0064] The DMT compound is optionally substituted with deuterium, wherein a deuterium atom is a hydrogen atom with an additional neutron. The DMT compound is also optionally substituted at position 4 or 5 with acetoxy or methoxy or position 4 with monohydrogen phosphate. The term “acetoxy” (often abbreviated to OAc) defines a univalent group derived from acetic acid by removal of a hydrogen atom from the OH moiety. The term “methoxy” (often abbreviated to OMe) defines a univalent group derived from methanol by removal of a hydrogen atom from the OH moiety. The term monohydrogen phosphate defines a diivalent group of formula  $\text{HPO}_4^{\text{sub.4}}$ , derived from phosphoric acid by removal of a proton from two of the three OH moieties, and thus denotes a substituent of formula  $\text{—OP(O)(OH)O—}$ .

[0065] In some embodiments, the dimethyltryptamine compound is optionally substituted at position 5 with methoxy or position 4 with acetoxy or monohydrogen phosphate.

[0066] Where the DMT compound is substituted at position 4 with monohydrogen phosphate, this is to reflect that psilocybin (also known as [3-(2-Dimethylaminoethyl)-1H-indol-4-yl] dihydrogen phosphate) in water generally has monohydrogen phosphate at the 4-position, this generally being understood to be the predominant form owing to the pKa values of the two terminal phosphate oxygen atoms being estimated as 1.3 and 6.5. It is further understood that the monohydrogen phosphate-containing form of psilocybin exists as a zwitterion (i.e. an internal salt) in which the nitrogen atom of the dimethylamino moiety is protonated. This form is thus, and psilocybin is to be regarded as, a salt of a DMT compound substituted at position 4 with monohydrogen phosphate.

[0067] In some embodiments, the dimethyltryptamine compound is optionally substituted with deuterium at one or more positions selected from the  $\alpha$ ,  $\beta$  and dimethyl carbon atoms. In further embodiments, the dimethyltryptamine compound is optionally substituted at one or more positions selected from the  $\alpha$  and  $\beta$  carbon atoms, such as the  $\alpha$  carbon.

[0068] For the avoidance of doubt, positions 4, 5,  $\alpha$  and  $\beta$  of the optionally substituted DMT salt refer to the positions labelled in the structure below (substitution not shown).

##STR00001##

[0069] The formulation comprises a buffer, which is separate to the salt, i.e. the buffer is not merely a counterion to the optionally substituted DMT. For example, where the salt is dimethyltryptamine fumarate (i.e. the fumaric acid salt of dimethyltryptamine), an amount of buffer is required over and above the buffer provided by the fumarate. The term “buffer” is well known in the art and refers to a chemical which, on inclusion within a formulation, resists a change in pH on addition of acid or base to the formulation. Within a formulation, a buffer comprises a weak acid and its conjugate base. A suitable buffer comprises an acid with a pKa value that lies within +1 of the desired pH of the formulation. For example, if the desired pH of the formulation is about 4.0, a suitable buffer comprises a weak acid with a pKa value of from about 3.0 to about 5.0. If the acid of a buffer has more than one pKa value (i.e. each molecule of the acid is able to donate more than one proton), in order for the buffer to be suitable, at least one of the pKa values lies within the desired pH range.

[0070] The weak acid and conjugate base of the buffer are in equilibrium with one another. In accordance with Le Chatelier's principle (if a constraint (such as a change in concentration of a reactant) is applied to a system in equilibrium, the equilibrium will shift so as to counteract the effect of the constraint), addition of acid or base to the formulation shifts the position of

equilibrium in favour of the conjugate base or weak acid, respectively. Consequently, the concentration of free protons in the formulation (and thus the pH) is relatively unchanged.

[0071] As described above, the formulation of the invention has a pH of from about 3.5 to about 6.5. In some embodiments, the buffer comprises an acetate salt and acetic acid ( $pK_a=4.75$ ); a citrate salt and citric acid ( $pK_a=3.13, 4.76$  and  $6.40$ ); an ascorbate salt and ascorbic acid ( $pK_a=4.17$  and  $11.6$ ); a benzoate salt and benzoic acid ( $pK_a=4.20$ ); a phosphate salt and phosphoric acid ( $pK_a=2.14, 7.20$  and  $12.37$ ); an oxalate salt and oxalic acid ( $pK_a=1.25$  and  $4.14$ ); or a formate salt and formic acid ( $pK_a=3.75$ ). The  $pK_a$  values cited herein are those reported at  $25^\circ\text{C}$ . in water. Typically, the buffer comprises only one of the pairs listed above, i.e. one acid and its conjugate base.

[0072] In some embodiments, the buffer comprises an acetate salt and acetic acid; a citrate salt and citric acid; an ascorbate salt and ascorbic acid; a benzoate salt and benzoic acid; or a phosphate salt and phosphoric acid.

[0073] In some embodiments, the pH of the formulation is from about 3.75 to about 6.5, such as from about 3.75 to about 5.75. Often, the pH of the formulation is from about 3.75 to about 4.25, typically about 4.0. In such embodiments, the buffer often comprises an acetate salt and acetic acid; a citrate salt and citric acid; an ascorbate salt and ascorbic acid; a benzoate salt and benzoic acid; an oxalate salt and oxalic acid; or a formate salt and formic acid. Sometimes, the buffer comprises an acetate salt and acetic acid; a citrate salt and citric acid; an ascorbate salt and ascorbic acid; or a benzoate salt and benzoic acid.

[0074] In some embodiments, the buffer comprises an acetate salt and acetic acid, often sodium acetate and acetic acid, or potassium acetate and acetic acid.

[0075] The concentration of buffer within the formulation is typically great enough to resist significant pH change of the formulation on storage of the formulation for two weeks (i.e. the pH typically fluctuates less than about 0.1 pH unit), and is small enough so that the osmolality of the formulation lies within the desired range. The skilled person is able to assess suitable buffer concentrations and to achieve this. Often, the concentration of buffer is from about 15 mM to about 75 mM, such as about 20 mM to about 30 mM. In some embodiments, the concentration of the buffer is about 25 mM.

[0076] As described above, the formulation comprises a salt of a DMT compound optionally substituted with deuterium and optionally substituted at position 4 or 5 with acetoxy or methoxy or position 4 with monohydrogen phosphate.

[0077] The salt comprises an acid and the DMT compound, or the salt comprises a DMT compound substituted at position 4 with monohydrogen phosphate. An example of a salt comprising an acid and DMT compound is dimethyltryptamine fumarate, which is the fumaric acid salt of dimethyltryptamine. P. H. Stahl and C. G. Wermuth provide an overview of pharmaceutical salts and the acids comprised therein in *Handbook of Pharmaceutical Salts: Properties, Selection and Use*, Weinheim/Zürich:Wiley-VCH/VHCA, 2002. The acids described in this review are suitable acids for inclusion within the salt of the formulation.

[0078] The salt may comprise an acid selected from the group consisting of fumaric acid, tartaric acid, citric acid, acetic acid, lactic acid, gluconic acid, 1-hydroxy-2-naphthoic acid, 2,2-dichloroacetic acid, 2-hydroxyethanesulfonic acid, 2-oxoglutaric acid, 4-acetamidobenzoic acid, 4-aminosalicylic acid, adipic acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, camphoric acid, camphor-10-sulfonic acid, decanoic acid, hexanoic acid, octanoic acid, carbonic acid, cinnamic acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, formic acid, galactaric acid, gentisic acid, glucoheptonic acid, glucuronic acid, glutamic acid, glutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, hydrobromic acid, hydrochloric acid, isobutyric acid, lactobionic acid, lauric acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, nicotinic acid, nitric acid, oleic acid, oxalic acid, palmitic acid, pamoic acid, phosphoric acid, propionic



acid, pyroglutamic acid (-L), salicylic acid, sebacic acid, stearic acid, succinic acid, sulfuric acid, thiocyanic acid, toluenesulfonic acid and undecylenic acid.

[0079] In some embodiments, where the salt comprises an acid and the DMT compound, the acid is a Brønsted acid having a pKa at 25° C. in water of from about 3 to about 5. In these embodiments, the Brønsted acid may act both as a counterion to the DMT compound and as a buffer. Thus, the formulation may be stabilised to a greater extent, i.e. degradation of the DMT compound may be further ameliorated, when the salt comprises such an acid.

[0080] In some embodiments, the salt comprises a Brønsted acid having a pKa at 25° C. of from about 3 to about 5, and a compound of Formula I

##STR00002## [0081] wherein: [0082] R<sup>sup.4</sup> and R<sup>sup.5</sup> are both H and each .sup.XH and each .sup.YH is independently selected from H and D, or [0083] one of R<sup>sup.4</sup> and R<sup>sup.5</sup> is H and the other is acetoxy or methoxy, each .sup.YH is H and each .sup.XH is independently selected from H and D, or [0084] the salt comprises a compound of formula I wherein R<sup>sup.4</sup> is monohydrogen phosphate, R<sup>sup.5</sup> is H and each .sup.YH and each .sup.XH is H.

[0085] In some embodiments, R<sup>sup.4</sup> and R<sup>sup.5</sup> are both H. In these embodiments, the DMT compound is any one or a combination of N,N-dimethyltryptamine, α-monodeutero-N,N-dimethyltryptamine, α,α-dideutero-N,N-dimethyltryptamine, α,β-dideutero-N,N-dimethyltryptamine, α,α,β-trideutero-N,N-dimethyltryptamine, α,β,β-trideutero-N,N-dimethyltryptamine and α,α,β,β-tetradeutero-N,N-dimethyltryptamine. Often, the DMT compound is N,N-dimethyltryptamine.

[0086] In some embodiments, R<sup>sup.4</sup>, R<sup>sup.5</sup> and each .sup.YH are H and each .sup.XH is independently selected from H and D. In these embodiments, the DMT compound is any one or a combination of N,N-dimethyltryptamine, α-monodeutero-N,N-dimethyltryptamine, and α,α-dideutero-N,N-dimethyltryptamine.

[0087] Partially deuterated and deuterated N,N-dimethyltryptamine compounds can be synthesised following the reaction schemes (synthetic schemes) provided in Schemes 1 and 2 below. The chemistry depicted in the schemes was reported by PE Morris and C Chiao (*Journal of Labelled Compounds And Radiopharmaceuticals*, Vol. XXXIII, No. 6, 455-465 (1993)). Partially deuterated and deuterated N,N-dimethyltryptamine compounds can also be synthesised following the synthetic scheme depicted in Scheme 3.

[0088] Herein, the terms α,α-dideutero-N,N-dimethyltryptamine compounds and α-protio, α-deutero-N,N-dimethyltryptamine compounds are referred to as deuterated (or fully deuterated) N,N-dimethyltryptamine and partially deuterated N,N-dimethyltryptamine respectively. A deuterated (or fully deuterated) N,N-dimethyltryptamine compound thus refers strictly to an N,N-dimethyltryptamine compound with both protons at the a position substituted with deuterium atoms. The term partially deuterated N,N-dimethyltryptamine compound strictly refers to an N,N-dimethyltryptamine compound in which one of the two protons at the a position is substituted with a deuterium atom. A deuterated N,N-dimethyltryptamine compound herein is any N,N-dimethyltryptamine compound substituted with two deuterium atoms at the a position, and a partially deuterated N,N-dimethyltryptamine compound any N,N-dimethyltryptamine compound with one hydrogen atom and one deuterium atom at the a position.

[0089] If desired, compositions comprising amounts of N,N-dimethyltryptamine and deuterated N,N-dimethyltryptamine compounds, with the relative proportions of N,N-dimethyltryptamine against deuterated N,N-dimethyltryptamine compounds and partially deuterated N,N-dimethyltryptamine compounds may be controlled by varying the ratio of lithium aluminium hydride and lithium aluminium deuteride in the reducing agent. It is to be understood that in such compositions, R<sup>sup.4</sup>, R<sup>sup.5</sup> and each .sup.YH are H and each .sup.XH is independently selected from H and D, i.e. the DMT compound is any one or a combination of N,N-dimethyltryptamine, α-monodeutero-N,N-dimethyltryptamine, and α,α-dideutero-N,N-dimethyltryptamine. Relative proportions may further be varied by adding one or more of N,N-dimethyltryptamine, α,α-

dideutero-N,N-dimethyltryptamine and  $\alpha,\alpha,\beta,\beta$ -tetra deutero-N,N-dimethyltryptamine to the compositions described hereinabove.

##STR00003##

##STR00004##

##STR00005##

[0090] Identification of the compositions resultant from the reduction step in Schemes 1 and 2 may be achieved, if desired, by chromatographic separation of the components of the mixtures by conventional means at the disposal of the skilled person in combination with spectroscopic and/or mass spectrometric analysis.

[0091] Alternative compositions are obtainable by mixing N,N-dimethyltryptamine, obtainable by Scheme 1 or Scheme 2 when the reducing agent is exclusively lithium aluminium hydride, with a deuterated N,N-dimethyltryptamine compound obtainable from Scheme 1 or Scheme 2 when the reducing agent is exclusively lithium aluminium deuteride.

[0092] The compositions described hereinabove may be further modified by adding one or more deuterated or partially deuterated N,N-dimethyltryptamine compounds. Stocks of such deuterated or partially deuterated N,N-dimethyltryptamine compounds may be obtained, for example, from the chromatographic separation described above.

[0093] In some embodiments, R.sup.4 is acetoxy and R.sup.5 is H, or R.sup.5 is acetoxy and R.sup.4 is H. According to some embodiments, R.sup.4 is acetoxy and R.sup.5 is H, thus the DMT compound is any one or a combination of 4-acetoxy-N,N-dimethyltryptamine, 4-acetoxy- $\alpha$ -monodeutero-N,N-dimethyltryptamine and 4-acetoxy- $\alpha,\alpha$ -dideutero-N,N-dimethyltryptamine. For example, the DMT compound is 4-acetoxy-N,N-dimethyltryptamine.

[0094] In some embodiments, R.sup.4 is H and R.sup.5 is methoxy, or R.sup.5 is H and R.sup.4 is methoxy. According to some embodiments, R.sup.4 is H and R.sup.5 is methoxy, thus the DMT compound is any one or a combination of 5-methoxy-N,N-dimethyltryptamine, 5-methoxy- $\alpha$ -monodeutero-N,N-dimethyltryptamine and 5-methoxy- $\alpha,\alpha$ -dideutero-N,N-dimethyltryptamine. For example, the DMT compound is 5-methoxy-N,N-dimethyltryptamine.

[0095] Scheme 4 represents schemes known in the art to synthesise DMT compounds, in which substituent R.sup.1 denotes hydrogen or the substituent R.sup.4 or R.sup.5, when other than hydrogen, as defined in Formula I; each R.sub.2 is methyl and HX refers to the acids described herein with which the DMT compounds described herein may form salts.

##STR00006##

[0096] Mixtures of compounds of Formula I comprising controllable proportions of optionally R.sup.4- or R.sup.5-substituted DMT and the same optionally R.sup.4- or R.sup.5-substituted DMT but with  $\alpha$ -mono- and/or  $\alpha,\alpha$ -di-deuteration may if desired be prepared by reducing 2-(3-indolyl)-N,N-dimethyl acetamide with a desired ratio of lithium aluminium hydride and lithium aluminium deuteride.

[0097] For more detail on the synthesis of DMT compounds, see the Example section herein.

[0098] In some embodiments, the salt is of an optionally substituted dimethyltryptamine compound and an acid selected from the group consisting of fumaric acid, tartaric acid, citric acid, acetic acid, lactic acid and gluconic acid, typically fumaric acid.

[0099] Accordingly, the salt may comprise: [0100] any one or a combination of N,N-dimethyltryptamine,  $\alpha$ -monodeutero-N,N-dimethyltryptamine,  $\alpha,\alpha$ -dideutero-N,N-dimethyltryptamine,  $\alpha,\beta$ -dideutero-N,N-dimethyltryptamine,  $\alpha,\alpha,\beta$ -trideutero-N,N-dimethyltryptamine,  $\alpha,\beta,\beta$ -trideutero-N,N-dimethyltryptamine and  $\alpha,\alpha,\beta,\beta$ -tetra deutero-N,N-dimethyltryptamine; or [0101] any one or a combination of 4-acetoxy-N,N-dimethyltryptamine, 4-acetoxy- $\alpha$ -monodeutero-N,N-dimethyltryptamine and 4-acetoxy- $\alpha,\alpha$ -dideutero-N,N-dimethyltryptamine; or [0102] any one or a combination of 5-methoxy-N,N-dimethyltryptamine, 5-methoxy- $\alpha$ -monodeutero-N,N-dimethyltryptamine and 5-methoxy- $\alpha,\alpha$ -dideutero-N,N-dimethyltryptamine; and [0103] an acid selected from the group consisting of fumaric acid, tartaric

acid, citric acid, acetic acid, lactic acid and gluconic acid, typically fumaric acid.

[0104] In some embodiments, the salt is DMT fumarate, i.e. it comprises DMT and fumaric acid.

[0105] The DMT compound may have a purity of about 80 to 100%. Sometimes, the purity is about 90 to 100%, such as from about 95 to 100%. Typically, the DMT compound has a purity of from about 99 to 100%, i.e. a purity greater than or equal to 99%. Percentages of purity herein are as determined by HPLC.

[0106] It is particularly advantageous to prepare the formulations of the present invention with a drug substance comprising the optionally substituted DMT compound or salt thereof with a purity of greater than 99%. By drug substance is meant, as is understood in the art, an active ingredient intended to furnish pharmacological activity or other direct effect in the diagnosis, cure, mitigation, treatment, or prevention of disease or to affect the structure or any function of the patient concerned, but does not include intermediates used in the synthesis of such ingredient. It will be understood that the drug substance may comprise one or more such active ingredients.

[0107] Formulations made with less pure drug substance show greater rates of related substances, which is indicative of inferior shelf-life. Accordingly, preferred embodiments of any aspect of the present invention comprise a drug substance comprising an optionally substituted dimethyltryptamine compound or salt thereof having a purity of greater than or equal to 99% when measured by HPLC. Particularly preferred embodiments comprise a drug substance comprising an optionally substituted dimethyltryptamine compound or salt thereof having a purity of greater than or equal to 99.5%, even more preferably 99.7%, and even more preferably 99.9%, when measured by HPLC. The concentration of the DMT compound within the formulation may be any desired concentration, provided that the osmolality of the formulation is about 250 to about 350 mOsm/Kg. The DMT compound may be at a concentration of about 0.001 to about 28 mg/mL, such as about 2.5 mg/mL to about 28 mg/mL. Accordingly, in some embodiments, the formulation of the invention comprises the DMT compound at a concentration of from 2.5 mg/ml to 28 mg/mL, including any integer value within this range. A concentration of 28 mg/mL of DMT provides approximately 148 mOsm/kg (approximately 296 mOsm/kg with counterions taken into account). This allows for the provision of a further 54 mOsm/kg by other components of the formulation, such as the buffer.

[0108] In some embodiments, the concentration of the DMT compound within the formulation is about 2.5 mg/mL, which provides approximately 13.2 mOsm/kg (approximately 26.4 mOsm/kg with counterions taken into account).

[0109] As described above, the formulation of the invention has an osmolality of about 250 to about 350 mOsm/Kg. As described above, to be injectable, a formulation may have an osmolality of about 250 to about 600 mOsm/Kg. In some embodiments of the invention, the osmolality of the formulation is about 250 to about 500 mOsm/Kg or about 250 to about 400 mOsm/Kg. In some embodiments, the osmolality of the formulation of the invention is about 275 to about 325 mOsm/Kg, such as about 280 to about 310 mOsm/Kg. Typically, the osmolality of the formulation is about 295 to about 305 mOsm/Kg. In some embodiments, the formulation is isotonic with human blood serum.

[0110] Sometimes, the concentration of optionally substituted DMT salt and buffer in the formulation gives rise to the desired osmolality. Alternatively, the desired osmolality may be achieved by inclusion of one or more tonicity agents in the formulation. Thus, in some embodiments, the formulation further comprises a tonicity agent. A tonicity agent is defined herein as a chemical that, on inclusion within a formulation, increases the osmolality of the formulation. As described above, the osmolality is the number of osmotically active particles (the number of solute particles) in 1 kg of a solution. Thus, a chemical that acts as a solute when incorporated into the formulation lies within the definition of a tonicity agent.

[0111] If the formulation further comprises a tonicity agent, the concentration of tonicity agent depends on the concentration of other components within the formulation, such as the optionally

substituted DMT and buffer. For example, where the formulation without tonicity agent has an osmolality of about 60 mOsm/kg, at least about 190 mOsm/kg would be provided by a tonicity agent (e.g. 95 mM of sodium chloride). Inclusion of a tonicity agent is often preferable in low concentration formulations useful for intravenous administration, for example in formulations comprising about 2.5 mg/ml of the DMT compound. In higher concentration formulations, for example those comprising the DMT compound in concentrations greater than about 5 mg/mL, the tonicity agent may be less preferable or absent.

[0112] M. F. Powell, T. Nguyen and L. Baloian provide a review of excipients suitable for parenteral administration (administration other than by the mouth or alimentary canal) in *PDA J. Pharm. Sci. Technol.*, 52, 238-311 (1998). All soluble excipients listed in this review article that can be given by the intravenous route will, when added to the formulation, contribute to the osmolality and thus can be considered tonicity agents.

[0113] In some embodiments, the tonicity agent is any one or a combination selected from the group consisting of sodium chloride; potassium chloride; dextrose; glucose; mannitol; phosphoric acid; lactose; sorbitol; sucrose; a phosphate salt such as sodium phosphate or potassium phosphate; acetic acid; an acetate salt such as sodium acetate, potassium acetate or ammonium acetate; alanine; ethanol; citric acid; a citrate salt such as sodium citrate or potassium citrate; arginine; ascorbic acid; an ascorbate salt such as potassium ascorbate or sodium ascorbate; benzyl alcohol; calcium chloride; creatinine; edetic acid; an edetate salt such as sodium edetate or calcium edetate; glycine; glycerol; histidine; lactic acid; magnesium chloride; polyethylene glycol; propylene glycol; sodium bicarbonate; sodium hydroxide; hydrochloric acid; lactic acid; lactate salts such as potassium lactate or sodium lactate; tartaric acid and tartrate salts such as sodium tartrate or potassium tartrate.

[0114] Some of the tonicity agents listed above may be used to buffer the formulation (e.g. acetate salt, acetic acid, citrate salt, citric acid, ascorbate salt, ascorbic acid, phosphate salt, phosphoric acid). For the avoidance of doubt, where one of the tonicity agents listed above is used as the buffer, it is not also the defined tonicity agent, i.e. where the formulation further comprises a tonicity agent, the tonicity agent is different from the buffer.

[0115] Often, the tonicity agent is any one or a combination selected from the group consisting of sodium chloride, potassium chloride, dextrose, glucose, mannitol, lactose, sorbitol and sucrose. Typically, the tonicity agent is sodium chloride.

[0116] In some embodiments, the formulation comprises sodium chloride at a concentration of about 120 mM to about 140 mM, such as about 125 mM to about 135 mM. Sometimes, the concentration of sodium chloride within the formulation is about 130 mM.

[0117] In some embodiments, the formulation consists essentially of the optionally substituted DMT salt, the buffer, water, and optionally a tonicity agent. By this is meant, for example, that the presence of additional components within the formulation is permitted, provided the amounts of such additional components do not materially affect, in a detrimental manner, the essential characteristics of the formulation. Given that the intention behind including the optionally substituted DMT salt, the buffer, water, and optional tonicity agent in the formulation is to produce a pharmaceutical formulation of optionally substituted DMT suitable for injection, and stable for at least several weeks when stored, it will be understood that the inclusion of components that materially affect, in a detrimental manner, the stability of the formulation or its suitability for injection (e.g. its osmolality or pH), are excluded from the formulation. On the other hand, it will be understood that the presence of any components that do not materially affect, in a detrimental manner, the stability of the formulation or its suitability for injection, is included. Such components include anti-oxidants and antimicrobial preservatives. For an overview of pharmaceutical excipients and their properties, including those with anti-oxidant and antimicrobial properties, see P. J. Sheskey, W G Cook and C G Cable, *Handbook of Pharmaceutical Excipients*, Eighth Edition, Pharmaceutical Press, London 2017.

[0118] Anti-oxidants commonly used in aqueous injectable formulations include ascorbic acid, citric acid, tartaric acid, sodium metabisulfite and thiol derivatives.

[0119] Antimicrobial preservatives commonly used in injectable formulations include methylparaben (methyl parahydroxybenzoate), ethylparaben (ethyl parahydroxybenzoate) and propylparaben (n-propyl parahydroxybenzoate) benzoic acid, benzyl alcohol, chlorobutanol, phenol and sodium benzoate.

[0120] In specific embodiments, the formulation consists of the salt, the buffer, water, and optionally a tonicity agent, i.e. the presence of any other components is excluded.

[0121] Often, the formulation has an oxygen content of less than 2 ppm, such as between 0.1 ppm and 2 ppm. The skilled person is able to determine the oxygen content of the formulation using any technique known in the art to be suitable, such as using a dissolved oxygen meter (e.g. a Jenway 970 Enterprise Dissolved Oxygen Meter, available from Keison Products:

<http://www.keison.co.uk/products/jenway/970.pdf>).

[0122] The formulation may be stored in any suitable container. In some embodiments, to ameliorate degradation of the formulation further, the formulation is stored in a container adapted to prevent penetration of ultraviolet light, such as amber glass vial. In others, the container within which the formulation is stored is not so adapted (and may be, for example, made of clear glass) with protection against ultraviolet light, if desired, provided by secondary packaging (for example packaging within which the receptacle containing the formulation may be placed). Often, the container is airtight and the formulation is stored under an inert atmosphere, such as under nitrogen or argon, typically nitrogen. The formulation may be stored at room temperature, e.g. at about 20 to about 30° C. or at cooler temperatures, for example at about 2 to about 8° C. Alternatively, to ameliorate degradation of the formulation further, it may be stored in a freezer.

[0123] Viewed from a second aspect, the invention provides a kit suitable for preparing a formulation of the first aspect, said kit comprising a salt of a DMT compound optionally substituted with deuterium and optionally substituted at position 4 or 5 with acetoxy or methoxy or position 4 with monohydrogen phosphate; and a buffer, which is separate to the salt.

[0124] Also provided is a kit to generate a formulation of the first aspect, the kit comprising:

[0125] a first composition comprising a salt of a DMT compound optionally substituted with deuterium and optionally substituted at position 4 or 5 with acetoxy or methoxy or position 4 with monohydrogen phosphate; and [0126] a second composition comprising a buffer, which is separate to the salt, [0127] wherein the first and second compositions are mixed with water and optionally a tonicity agent, and the resulting mixture generates the formulation of the first aspect.

[0128] For the avoidance of doubt, embodiments related to the optionally substituted DMT salt and the buffer of the first aspect of the invention as defined herein apply mutatis mutandis to the second aspect. For example, the optionally substituted DMT salt of the kit may comprise a Brønsted acid having a pKa at 25° C. of from about 3 to about 5, and a compound of Formula I and/or the buffer may comprise an acetate salt and acetic acid.

[0129] The optionally substituted DMT salt within the kit may be a solid, e.g. in a powder or crystalline form. To ameliorate degradation of the optionally substituted DMT salt in the solid form, the salt may be lyophilised (freeze-dried) before incorporation into the kit. Lyophilising the salt comprises freezing it in the presence of solvent (typically water) and separating the solvent from the salt by sublimation.

[0130] The kit may further comprise a tonicity agent. When the kit further comprises a tonicity agent, the embodiments related to the optional tonicity agent of the first aspect of the invention as defined herein apply mutatis mutandis to the second aspect. For example, the tonicity agent may be any one or a combination selected from the group consisting of sodium chloride, potassium chloride, dextrose, glucose, mannitol, lactose, sorbitol and sucrose.

[0131] Viewed from a third aspect, the invention provides a method of preparing a pharmaceutical formulation of the first aspect, which is typically a solution. The method comprises contacting the

optionally substituted DMT salt, buffer, water and optionally a tonicity agent. For the avoidance of doubt, the embodiments of the first aspect of the invention apply mutatis mutandis to the third aspect. For example, the salt may be DMT, the buffer may comprise acetic acid and an acetate salt, and/or sodium chloride may be used as a tonicity agent.

[0132] It will be understood that the contacting of the method may be achieved in a variety of ways. Often, the optionally substituted DMT salt is dissolved in water to form a first solution to which the buffer is added and dissolved, forming a second solution. If a tonicity agent is used, it is often added to and dissolved in the second solution.

[0133] In some embodiments, an aqueous solution of the buffer is contacted with the salt, wherein the aqueous solution has a pH of about 3.5 to about 6.5, such as a pH from about 3.75 to about 6.5. Sometimes, the aqueous solution has a pH of about 3.75 to about 5.75, such as a pH from about 3.75 to about 4.25. In some embodiments, the aqueous solution has a pH of about 4.0.

[0134] The optionally substituted DMT salt within the formulation may be formed by contacting optionally substituted DMT as a free base with an aqueous solution comprising a quantity of buffer suitable to stabilise the pH and act as counterion to the optionally substituted DMT when protonated. Accordingly, the method of the invention may comprise contacting the optionally substituted dimethyltryptamine in free base form with the buffer, water and optionally a tonicity agent.

[0135] In some embodiments, the method further comprises adjusting the pH of the solution resultant from the contacting. Since the pH of the solution resultant from the contacting is usually low, pH adjustment often comprises contacting the solution with a suitable base. The skilled person is able to assess which bases are suitable to adjust the pH of the solution resultant from the contacting without risk of degradation of the optionally substituted DMT salt.

[0136] Often, the pH of the solution resultant from the contacting is adjusted with any one selected from the group consisting of sodium hydroxide, potassium hydroxide, sodium carbonate, ammonium hydroxide, calcium hydroxide and magnesium hydroxide. In some embodiments, the pH is adjusted with sodium hydroxide or potassium hydroxide.

[0137] As described above, to ameliorate degradation of the formulation further, it may be desirable to minimise the total oxygen content within the container in which the formulation is stored, the oxygen within the container equilibrating between the formulation and the headspace (if any) within the container. Accordingly, it may be desirable to store the formulation under an inert atmosphere for example by purging the headspace to reduce its oxygen content from about 20% typically found in air, to less than, for example, 0.5%. Additionally or alternatively, in some embodiments, the method further comprises sparging the solution resultant from the contacting with an inert gas, such as nitrogen or argon, typically nitrogen.

[0138] Viewed from a fourth aspect, there is provided use of a buffer to ameliorate degradation of an injectable pharmaceutical formulation of a salt of a dimethyltryptamine compound optionally substituted with deuterium and optionally substituted at position 4 or 5 with acetoxy or methoxy or position 4 with monohydrogen phosphate.

[0139] For the avoidance of doubt, the embodiments of the first aspect of the invention apply mutatis mutandis to the fourth aspect. Specifically, the embodiments of the first aspect relating to the buffer and the optionally substituted DMT salt apply mutatis mutandis to the fourth aspect. For example, the buffer of the fourth aspect may comprise an acetate salt and acetic acid; a phosphate salt and phosphoric acid; a citrate salt and citric acid; an ascorbate salt and ascorbic acid; or a benzoate salt and benzoic acid; and/or the optionally substituted DMT salt of the fourth aspect may comprise a Brønsted acid having a pKa at 25° C. of from about 3 to about 5.

[0140] As described above, DMT has a possible therapeutic role in the treatment of depression, obsessive-compulsive disorder, and substance abuse disorders (S. A. Barker, 2018, supra). Viewed from a fifth aspect, therefore, the invention provides a formulation of the first aspect for use in therapy.

[0141] Viewed from a sixth aspect, the invention provides a formulation of the first aspect for use in a method of treating a psychiatric or neurological disorder in a patient. Often, the psychiatric or neurological disorder is selected from the group consisting of (i) an obsessive compulsive disorder, (ii) a depressive disorder, (iii) an anxiety disorder, (iv) substance abuse, and (v) an avolition disorder. Often, the disorder is selected from the group consisting of major depressive disorder, treatment resistant major depressive disorder, post-partum depression, an obsessive compulsive disorder and an eating disorder such as a compulsive eating disorder.

[0142] Viewed from a seventh aspect, the invention provides a method of treating a psychiatric or neurological disorder comprising administering to a patient in need thereof a formulation of the first aspect. The psychiatric or neurological disorder may be any of those described in relation to the sixth aspect. For example, the disorder may be selected from the group consisting of major depressive disorder, treatment resistant major depressive disorder, post-partum depression, an obsessive compulsive disorder and an eating disorder such as a compulsive eating disorder.

[0143] In order to treat the disorder, the formulation comprises an effective amount of the DMT compound, i.e. an amount that is sufficient to reduce or halt the rate of progression of the disorder, or to ameliorate or cure the disorder and thus produce the desired therapeutic or inhibitory effect.

[0144] The formulation is suitable for injection, thus its administration in therapy typically comprises injection of the formulation.

[0145] The formulation may be suitable for bolus injection, in which a discrete amount of an optionally substituted DMT salt is administered in one injection such that the concentration of DMT in the body quickly increases. Bolus injections are typically administered intravenously (directly into the vein), intramuscularly (within the muscle), intradermally (beneath the skin) or subcutaneously (within the fat or skin).

[0146] Each and every reference referred to herein is hereby incorporated by reference in its entirety, as if the entire content of each reference was set forth herein in its entirety.

[0147] The invention may be further understood with reference to the following non-limiting clauses and examples following thereafter:

[0148] A pharmaceutical formulation suitable for injection is described, comprising a salt of a dimethyltryptamine compound optionally substituted with deuterium and optionally substituted at position 4 or 5 with acetoxy or methoxy or position 4 with monohydrogen phosphate; a buffer which is separate to the salt; and water, wherein the formulation has a pH of about 3.5 to about 6.5 and an osmolality of about 250 to about 350 mOsm/Kg. The pH is from about 3.75 to about 6.5, from about 3.75 to about 5.75, from about 3.75 to about 4.25, or about 4.0. The formulation has an osmolality of about 275 to about 325 mOsm/Kg, about 280 to about 310 mOsm/Kg, or about 295 to about 305 mOsm/Kg.

[0149] The dimethyltryptamine compound is optionally substituted at position 5 with methoxy or position 4 with acetoxy or monohydrogen phosphate. The dimethyltryptamine compound is optionally substituted with deuterium at one or more positions selected from the a, B and dimethyl carbon atoms. The dimethyltryptamine compound is optionally substituted with deuterium at one or more positions selected from the  $\alpha$  and  $\beta$  carbon atoms. The dimethyltryptamine compound is optionally substituted with deuterium once or twice at the a carbon atom.

[0150] In some embodiments, the salt comprises a Brønsted acid having a pKa of from about 3 to about 5 and a compound of Formula I

##STR00007## [0151] wherein: [0152] R.sup.4 and R.sup.5 are both H and each .sup.XH and each .sup.YH is independently selected from H and D, or one of R.sup.4 and R.sup.5 is H and the other is acetoxy or methoxy, each .sup.YH is H and each .sup.XH is independently selected from H and D, or [0153] the salt comprises a compound of formula I wherein R.sup.4 is monohydrogen phosphate, R.sup.5 is H and each .sup.YH and each .sup.XH is H.

[0154] In some embodiments, R.sup.4 and R.sup.5 are both H. In some embodiments, each .sup.YH is H and each .sup.XH is independently selected from H and D. In some embodiments,

R.sup.4 is acetoxy and R.sup.5 is H. In some embodiments, R.sup.4 is H and R.sup.5 is methoxy. [0155] In any of the foregoing, the optionally substituted dimethyltryptamine compound may be dimethyltryptamine.

[0156] The salt may be of an optionally substituted dimethyltryptamine compound and an acid selected from the group consisting of fumaric acid, tartaric acid, citric acid, acetic acid, lactic acid and gluconic acid. Preferably, the acid is fumaric acid.

[0157] The optionally substituted dimethyltryptamine compound may have a purity of greater than 99% by HPLC.

[0158] The optionally substituted dimethyltryptamine compound may be at a concentration of about 0.001 to about 28 mg/mL, about 2.5 to about 28 mg/mL, or about 2.5 mg/mL.

[0159] In some embodiments, the buffer comprises an acetate salt and acetic acid; a phosphate salt and phosphoric acid; a citrate salt and citric acid; an ascorbate salt and ascorbic acid; a benzoate salt and benzoic acid; an oxalate salt and oxalic acid; or a formate salt and formic acid. In some embodiments, the buffer comprises an acetate salt and acetic acid; a phosphate salt and phosphoric acid; a citrate salt and citric acid; an ascorbate salt and ascorbic acid; or a benzoate salt and benzoic acid. In some embodiments, the buffer comprises an acetate salt and acetic acid. In some embodiments, the buffer comprises sodium acetate and acetic acid, or potassium acetate and acetic acid.

[0160] In some embodiments, the formulation comprises the buffer in a concentration of about 15 mM to about 75 mM, about 20 mM to about 30 mM, or about 25 mM.

[0161] In some embodiments, the formulation consists essentially of the salt, the buffer, water, and optionally, a tonicity agent. The tonicity agent may be sodium chloride. In some embodiments, the formulation consists of the salt, the buffer, water, and optionally, a tonicity agent.

[0162] In some embodiments, the formulation comprises sodium chloride at a concentration of about 120 mM to about 140 mM, about 125 mM to about 135 mM, or about 130 mM.

[0163] In some embodiments, the formulation has an oxygen content of less than 2 ppm or between 0.1 ppm and 2 ppm.

[0164] In some embodiments, the formulation is stored in a container adapted to prevent penetration of ultraviolet light, for example, in an amber glass vial.

[0165] In some embodiments, a kit suitable for preparing a formulation as above is provided, said kit comprising a salt of a dimethyltryptamine compound optionally substituted with deuterium and optionally substituted at position 4 or 5 with acetoxy or methoxy or position 4 with monohydrogen phosphate; and a buffer which is separate to the salt.

[0166] In some embodiments, a method of preparing a pharmaceutical formulation as defined as above comprises contacting the salt, buffer, water and optionally a tonicity agent. In some embodiments, an aqueous solution of the buffer is contacted with the salt, wherein the aqueous solution has a pH as defined in any one of clauses 1 to 5. In some embodiments, the method comprises contacting the optionally substituted dimethyltryptamine in free base form with the buffer, water and optionally a tonicity agent. In some embodiments, the method comprises adjusting the pH of the solution resultant from the contacting. In some embodiments, the pH is adjusted with sodium hydroxide or potassium hydroxide. In some embodiments, the method comprises sparging the solution resultant from the contacting with an inert gas (such as, for example nitrogen).

[0167] In some embodiments, a use of a buffer to ameliorate degradation of an injectable pharmaceutical formulation of a salt of a dimethyltryptamine optionally substituted with deuterium and optionally substituted at position 4 or 5 with acetoxy or methoxy or position 4 with monohydrogen phosphate is provided.

[0168] In some embodiments, a method of treating a psychiatric or neurological disorder in a patient with the above formulations is provided. The psychiatric or neurological disorder may be selected from the group consisting of (i) an obsessive compulsive disorder, (ii) a depressive



disorder, (iii) an anxiety disorder, (iv) substance abuse, and (v) an avolition disorder. The disorder may be major depressive disorder. The disorder may be treatment resistant major depressive disorder. The disorder may be post-partum depression. The disorder may be an obsessive compulsive disorder. The disorder may be an eating disorder (such as, for example, a compulsive eating disorder).

## EXAMPLES

### Example 1

[0169] N,N-DMT 220.9 g (as free base) was prepared as N,N-DMT fumarate, using the chemistry depicted in Scheme 3 above. An additional 4-6 g of six partially deuterated mixtures were also produced using modified conditions.

[0170] In Scheme 3, the carbodiimide coupling agent EDC.Math.HCl and additive coupling agent (which enhance the reactivity of the coupling agent) HOBt are used. More generally, the combination of two or more coupling agents comprises an agent selected from (i) a phosphonium coupling agent and a carbodiimide coupling agent selected from DCC, EDC, and DIC; and (ii) an additive coupling agent selected from HOBt, HOOBt, HOSu, HOAt, Ethyl 2-cyano-2-(hydroximino)acetate and DMAP. Often, as exemplified below, EDC is used, preferably as the HCl salt. Often, as exemplified below, the additive coupling agent HOBt. Often, as exemplified below, EDC is used, preferably as the HCl salt in combination with the additive coupling agent HOBt.

### Stage 1: Coupling of indole-3-acetic acid and Dimethylamine

[0171] To a 5 L vessel under N.sub.2 was charged indole-3-acetic acid (257.0 g, 1.467 mol), hydroxybenzotriazole (HOBt, ~20% wet) (297.3 g, 1.760 mol) and dichloromethane (2313 mL) to give a milky white suspension. 1-Ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (EDC.Math.HCl, 337.5 g, 1.760 mol) was then charged portion-wise over 5 minutes at 16-22° C. The reaction mixture was stirred for 2 hours at ambient temperature before 2M dimethylamine in THF (1100 mL, 2.200 mol) was charged dropwise over 20 minutes at 20-30° C. The resultant solution was stirred at ambient temperature for 1 hour where HPLC indicated 1.1% indole-3-acetic acid and 98.1% target product referred to as Stage 1). The reaction mixture was then charged with 10% K.sub.2CO.sub.3 (1285 mL) and stirred for 5 minutes. The layers were separated, and the upper aqueous layer extracted with dichloromethane (643 mL×2). The organic extracts were combined and washed with saturated brine (643 mL). The organic extracts were then dried over MgSO.sub.4, filtered and concentrated in vacuo at 45° C. This provided 303.1 g of crude Stage 1 as an off-white sticky solid. The crude material was then subjected to a slurry in tert-butyl methyl ether (TBME, 2570 mL) at 50° C. for 2 hours before being cooled to ambient temperature, filtered and washed with TBME (514 mL×2). The filter-cake was then dried in vacuo at 50° C. to afford Stage 1 266.2 g (yield=90%) as an off-white solid in a purity of 98.5% by HPLC and >95% by NMR.

### Stage 2: Preparation of DMT

[0172] To a 5 L vessel under N.sub.2 was charged Stage 1 (272.5 g, 1.347 mol) and tetrahydrofuran (THF, 1363 mL) to give an off-white suspension. 2.4 M LiH.sub.4 in THF (505.3 mL, 1.213 mol) was then charged dropwise over 35 minutes at 20-56° C. to give an amber solution. The solution was heated to 60° C. for 2 hours where HPLC indicated Stage 1 ND, target product bracket referred to as Stage 2, 92.5%), Impurity 1 (2.6%), Impurity 2 (1.9%). The complete reaction mixture was cooled to ambient temperature and then charged to a solution of 25% Rochelle's salts (aq) (2725 mL) dropwise over 30 minutes at 20-30° C. The resultant milky white suspension was allowed to stir at 20-25° C. for 1 hour after which the layers were separated and the upper organic layer washed with saturated brine (681 mL). The organic layer was then dried over MgSO.sub.4, filtered and concentrated in vacuo at 45° C. The resultant crude oil was subjected to an azeotrope from ethanol (545 mL×2). This provided 234.6 g (yield=92%) of Stage 2 in a purity of 95.0% by HPLC and >95% by NMR.

### Stage 3a (i)-(iii): Preparation of Seed Crystals of DMT Fumarate

[0173] (i) Stage 2 (100 mg) was taken up in 8 volumes of isopropyl acetate and warmed to 50° C. before charging fumaric acid (1 equivalent) as a solution in ethanol. The flask was then allowed to mature at 50° C. for 1 hour before cooling to room temperature and stirring overnight, resulting in a white suspension. The solids were isolated by filtration and dried for 4 hours at 50° C. to provide 161 mg of product (>99% yield). Purity by HPLC was determined to be 99.5% and by NMR to be >95%.

[0174] (ii) Substitution of isopropyl acetate for isopropyl alcohol in method (i) afforded a white suspension after stirring overnight. The solids were isolated by filtration and dried for 4 hours at 50° C. to provide 168 mg of product (>99% yield). Purity by HPLC was determined to be 99.8% and by NMR to be >95%.

[0175] Substitution of isopropyl acetate for tetrahydrofuran in method (i) afforded a white suspension after stirring overnight. The solids were isolated by filtration and dried for 4 hours at 50° C. to provide 161 mg of product (>99% yield). Purity by HPLC was determined to be 99.4% and by NMR to be >95%.

[0176] Analysis by x-ray powder diffraction, showed the products of each of methods 9i) to (iii) to be the same, which was labelled Pattern A.

#### Stage 3b: Preparation of DMT Fumarate

[0177] To a 5 L flange flask under N.sub.2 was charged fumaric acid (152.7 g, 1.315 mol) and Stage 2 (248.2 g, 1.315 mol) as a solution in ethanol (2928 mL). The mixture was heated to 75° C. to give a dark brown solution. The solution was polish filtered into a preheated (80° C.) 5 L jacketed vessel. The solution was then cooled to 70° C. and seeded with Pattern A (0.1 wt %), the seed was allowed to mature for 30 minutes before cooling to 0° C. at a rate of 5° C./hour. After stirring for an additional 4 hours at 0° C., the batch was filtered and washed with cold ethanol (496 mL×2) and then dried at 50° C. overnight. This provided 312.4 g (yield=78%) of Stage 3 in a purity of 99.9% by HPLC and >95% by NMR. XRPD: Pattern A.

[0178] 5-methoxy-DMT fumarate was prepared analogously to the DMT fumarate described immediately above except for the use of 5-methoxyindole-3-acetic acid.

#### Synthesis of Deuterated Mixtures of DMT Compounds

[0179] A modified synthesis at stage 2 using solid LiAlH.sub.4/LiAlD.sub.4 mixtures was adopted, using 1.8 equivalents of LiAlH.sub.4/LiAlD.sub.4 versus 0.9 equivalents using the process described above for undeuterated DMT.

[0180] Six deuteration reactions were performed.

#### Representative Synthesis of a Deuterated Mixture (Using 1:1 LiAlH.sub.4:LiAlD.sub.4) of DMT Compounds

[0181] To a 250 mL 3-neck flask under N.sub.2 was charged LiAlH.sub.4 (1.013 g, 26.7 mmol), LiAlD.sub.4 (1.120 g, 26.7 mmol) and THF (100 mL). The resultant suspension was stirred for 30 minutes before stage 1 (6 g, 29.666 mmol) was charged portion-wise over 15 minutes at 20-40° C. The reaction mixture was then heated to reflux (66° C.) for 2 hours where HPLC indicated no stage 1 remained. The mixture was cooled to 0° C. and quenched with 25% Rochelle's salts (aq) (120 mL) over 30 minutes at <30° C. The resultant milky suspension was stirred for 1 hour and then allowed to separate. The lower aqueous layer was removed and the upper organic layer washed with saturated brine (30 mL). The organics were then dried over MgSO.sub.4, filtered and concentrated in vacuo. This provided 4.3 g of crude material. The crude was then taken up in ethanol (52 mL) and charged with fumaric acid (2.66 g, 22.917 mmol) before heating to 75° C. The resultant solution was allowed to cool to ambient temperature overnight before further cooling to 0-5° C. for 1 hour. The solids were isolated by filtration and washed with cold ethanol (6.5 mL×2). The filtercake was dried at 50° C. overnight to provided 5.7 g (yield=63%) of product in a purity of 99.9% by HPLC and >95% by NMR.

#### Assessment of Extents of Deuteration

[0182] This was achieved by LCMS-SIM (SIM=single ion monitoring), the analysis giving a

separate ion counts for each mass for the three deuterated N,N-dimethyltryptamine compounds (N,N-dimethyltryptamine (D0),  $\alpha$ -deutero-N,N-dimethyltryptamine (D1) and  $\alpha,\alpha$ -dideutero-N,N-dimethyltryptamine (D2) at the retention time for N,N-dimethyltryptamine. The percentage of each component was then calculated from these ion counts.

[0183] For example, % D0=[D0/(D0+D1+D2)] $\times$ 100.

#### HPLC Parameters

TABLE-US-00001 System: Agilent 1100/1200 series liquid chromatograph or equivalent Column: Triart Phenyl; 150  $\times$  4.6 mm, 3.0  $\mu$ m particle size (Ex: YMC, Part number: TPH12S03-1546PTH) Mobile phase A: Water:Trifluoroacetic acid (100:0.05%) Mobile phase B: Acetonitrile:Trifluoroacetic acid (100:0.05%) Time % A % B Gradient: 0 95 5 13 62 38 26 5 95 30.5 5 95 31 95 5 Flow rate: 1.0 mL/min Stop time: 31 minutes Post runtime: 4 minutes Injection volume: 5  $\mu$ L Wash vial: N/A Column temperature: 30° C. combined Wavelength: 200 nm, (4 nm) Reference: N/A

#### Mass Spectrometry Parameters

TABLE-US-00002 System: Agilent 6100 series Quadrupole LC-MS or equivalent Drying gas flow: 12.0 L/min Drying gas temp.: 350° C. Nebuliser 35 psig pressure: Fragmentor: 110 Gain: 1.00 Cpd RT RRT Conc Diluent Detection Mass D0 10.64 1.00 0.30 mg/ml CH.sub.3CN:H.sub.2O (50:50) (+) SIM 189.10 m/z D1 10.64 1.00 0.30 mg/ml CH.sub.3CN:H.sub.2O (50:50) (+) SIM 190.10 m/z D2 10.64 1.00 0.30 mg/ml CH.sub.3CN:H.sub.2O (50:50) (+) SIM 191.10 m/z MS-SIM range is the target mass  $\pm$ 0.1 m/z

[0184] The data for the six deuterated reactions are tabulated in Table A below:

TABLE-US-00003 TABLE A Mixture No. Input Output Purity Purity (LiAlH.sub.4:LiAlD.sub.4 (stage stage 3 by by Deuteration % ratio) 1) (yield) HPLC NMR D.sub.0 D.sub.1 D.sub.2 1 (SPL028) (0:1) 5 g 5.3 g (65%) 99.7% >95% 0.7% 2.7% 96.6% 2 (1:1) 6 g 5.699 g (63%) 99.9% >95% 30.0% 48.3% 21.7% 3 (1:2) 5 g 4.206 g (52%) 99.9% >95% 16.5% 46.8% 36.8% 4 (1:3) 5 g 5.558 g (68% 99.8% >95% 9.3% 41.5% 49.2% 5 (2:1) 5 g 4.218 g (52%) 99.9% >95% 47.5% 41.3% 11.2% 6 (3:1) 5 g 5.0 g (62%) 99.4% >95% 57.5% 35.3% 7.4%

[0185] To synthesise 5-methoxy-N,N-dimethyltryptamine or 4-methoxy-N,N-dimethyltryptamine, 3-indoleacetic acid (see Scheme 3) may be replaced with 5-methoxyindole-3-acetic acid (see synthesis of  $\alpha,\alpha$ -dideutero-5-methoxydimethyltryptamine described below) or 4-methoxyindole-3-acetic acid respectively, both of which are commercially available (for 5-methoxyindole-3-acetic acid, for example from Sigma-Aldrich (code M14935-1G), for 4-methoxyindole-3-acetic acid see for example Aaron chemicals (code AR00VTP1)).

[0186] 5-methoxy-N,N-dimethyltryptamine (see Sigma-Aldrich code M-168-1ML), 4-methoxy-N,N-dimethyltryptamine (see Cayman Chemical code 9000895), 4-acetoxy-N,N-dimethyltryptamine (see Cayman Chemical code 14056) and 3-[2-(Dimethylamino)ethyl]-1H-indol-4-yl phosphate (psilocybin, see Sigma-Aldrich CAS Number 520-52-5) are also commercially available.

#### Synthesis of $\alpha,\alpha$ -dideutero-5-methoxydimethyltryptamine

##### Stage 1:

[0187] To a 100 mL 3-neck flask under N.sub.2 was charged 5-methoxyindole-3-acetic acid (3.978 g, 19.385 mmol), HOBt (~20% wet) (3.927 g, 23.261 mmol) and DCM (40 mL). EDC.Math.HCl (4.459 g, 23.261 mmol) was then charged in portions over 15 minutes at <30° C. The reaction mixture was stirred at ambient temperature for 1 hour before being charged with 2 M dimethylamine (14.54 mL, 29.078 mmol) dropwise over 15 minutes at <25° C. After stirring for 1 hour HPLC indicated no SM remained. The reaction mixture was then charged with 10% K.sub.2CO.sub.3 (20 mL), stirred for 5 minutes then allowed to separate. The lower aqueous layer was removed and back extracted with DCM (10 mL $\times$ 2). The organic extracts were combined, washed with saturated brine (10 mL) then dried over MgSO.sub.4 and filtered. The filtrate was concentrated in vacuo at 45° C. to provide 3.898 g active (yield=87%) of product in a purity of

95.7% by HPLC.

#### Stage 2

[0188] To a 100 mL 3-neck flask under N.sub.2 was charged Stage 1 methoxy derivative (3.85 g, 16.586 mmol) and THF (19.25 mL). 2.4 M LiAlD.sub.4 in THF (6.22 mL, 14.927 mmol) was then charged dropwise over 30 minutes at <40° C. The reaction mixture was heated to 60° C. for 1 hour where HPLC indicated 0.1% SM remained. The reaction mixture was then cooled to ambient temperature and quenched into 25% Rochelle's salts (38.5 mL) dropwise over 30 minutes at <30° C. The resultant suspension was stirred for 1 hour before being allowed to separate. The lower aqueous layer was then removed, and the upper organic layer washed with saturated brine (9.6 mL). The organics were then dried over MgSO.sub.4, filtered and concentrated in vacuo before being subjected to an azeotrope from EtOH (10 mL×2). This provided 3.196 g active (yield=88%) of product in a purity of 91.5% by HPLC.

#### Stage 3

[0189] To a 50 mL 3-neck flask under N.sub.2 was charged fumaric acid (1.675 g, 14.430 mmol) and a solution of Stage 2 methoxy derivative (3.15 g, 14.299 mmol) in EtOH (37.8 mL). The mixture was then heated to 75° C. for 1 hour, this did not produce a solution as expected, the mixture was further heated to reflux (78° C.) which still failed to provide a solution. The suspension was therefore cooled to 0-5° C., filtered and washed with EtOH (8 mL×2) before being dried at 50° C. overnight. This provided 3.165 g (yield=65%) of material in a purity of 99.9% by HPLC.

#### Development of Formulation

[0190] A stable formulation isotonic with human blood serum and suitable for intravenous (IV) bolus administration of DMT fumarate was developed. A suitable process for preparation of such a formulation comprising DMT fumarate at a concentration of 2.5 mg/ml was also developed. These formulations were prepared and placed under accelerated storage to assess stability over several weeks.

[0191] All stated concentrations below are expressed in terms of the free base (i.e. in the absence of fumarate counterion). To do so, a correction factor of 1.59 has been applied to the specific batch of drug substance as supplied.

#### Experimental Details

##### Initial Tests

[0192] Solubility of DMT fumarate was assessed at a concentration of 10 mg/ml in a small selection of aqueous vehicles (water, saline, 20 mM phosphate buffer and a combination of buffer and saline).

[0193] Phosphate buffer (100 mL) was prepared using 219.53 mg of the dibasic form [HPO.sub.4] [Na].sub.2 with 183.7 mg of the monobasic form [H.sub.2PO.sub.4]Na, both dihydrate salts. The solution was adjusted to pH 7.0 with addition of NaOH (1 M) and then made to volume. 10 ml of a 10 mg/mL formulation was prepared.

[0194] A phosphate buffer combined with saline was initially tested (20 mM phosphate buffer in 0.45% w/v saline) as a good starting point for a physiologically acceptable formulation with no solubility issues. To prepare this, sodium chloride was first dissolved in water to produce the saline solution (100 mL, 0.45% w/v). The phosphate salts, in the quantities described above, were then dissolved in the saline solution and the pH was adjusted using NaOH (1 M).

[0195] DMT fumarate was readily soluble in each aqueous vehicle. In terms of appearance, each solution was a clear beige colour, which on filtration (using 0.2 µm filters) was removed to produce a clear colourless solution. The pH of these solutions was in the range 3-4.

[0196] Buffer strength at 30 and 50 mM (as phosphate buffer, pH 7.4, prepared in 0.45% w/v sodium chloride) was tested to assess the effect of the buffer on pH control of formulations comprising concentrations of 2 or 2.5 mg/mL DMT fumarate. This buffer strength range was chosen in order to determine the required buffer strength so as to fix the pH of the formulation to

about pH 7.4. When developing formulations for injection, it is typical to match the pH of the formulation with those of the patient's blood serum. Human blood serum has a pH of about 7.4. The buffers were prepared as follows. Saline solution was prepared by dissolving 9 g of sodium chloride in 2 litres of water. The phosphate salts (e.g. 30 mM=dibasic dihydrate (4.29 g), monobasic dihydrate (1.43 g), 50 mM=dibasic dihydrate (7.28 g), monobasic dihydrate (2.25 g)) were dissolved into the saline solution the pH was adjusted to 7.4 using NaOH (1 M). 9 g of NaCl in 2 litres of water. pH adjustment to pH 7.4 with 1M NaOH.

[0197] The initial pH of each solution following preparation was less than pH 7.4. At 20 mM the initial pH value of 5.9 continued to drop on storage of the solution overnight in the laboratory, indicating that the buffering capacity of the buffer at 20 mM concentration was insufficient. At both 30 mM and 50 mM buffer strengths, the initial pH values were >6.5 and remained stable.

[0198] A short-term stability assessment was performed and data obtained for the pH, osmolality and assay are presented in Tables 1 and 2. The data in Table 2 were obtained on storage of the formulations at between 40 to 50° C.

TABLE-US-00004 TABLE 1 Short-term formulation stability assessment pH Osmolality (mOsm/kg) Assay (mg/ML) 24 Day Day 24 Day Day 24 Day Day Sample Initial hr 5 7 Initial hr 5 7											
2.00	2.02	—	2.09	2 mg .Math. mL.sup.-1/30 mM buffer/light.sup.a	6.74	6.74	—	6.71	239	238	—
253	—	1.99	—	2.05 2 mg .Math. mL.sup.-1/30 mM buffer/dark.sup.b	6.74	6.73	—	6.72	239	235	—
237	1.99	1.96	—	2.09 2 mg .Math. mL.sup.-1/30 mM buffer/2-8° C.	6.74	6.73	—	6.72	239	238	—
—	289	1.95	1.99	—	2.04 2 mg .Math. mL.sup.-1/50 mM buffer/light.sup.a	6.95	6.96	—	6.95	285	284
—	289	1.95	1.99	—	2.04 2 mg .Math. mL.sup.-1/50 mM buffer/dark.sup.b	6.85	6.98	—	6.95	285	—
285	—	286	—	—	2.04 2 mg .Math. mL.sup.-1/50 mM buffer/2-8° C.	6.95	6.97	—	6.95	285	284
—	283	1.98	1.98	—	2.08 2.5 mg .Math. mL.sup.-1/50 mM buffer/light.sup.a	6.87	—	6.86	—	288	—
—	286	—	2.46	—	2.51 — 2.5 mg .Math. mL.sup.-1/50 mM buffer/dark.sup.b	6.87	—	6.86	—	288	—
—	287	—	2.43	—	2.60 — 2.5 mg .Math. mL.sup.-1/50 mM	6.87	—	6.86	—	288	—
2.57	—	—	—	—	buffer/2-8° C. .sup.a,b	Light and dark at laboratory temperature storage (15-25° C.)					

TABLE-US-00005 TABLE 2 Short-term formulation stability assessment pH Assay (mg/mL) Related Substances (%) Sample Initial Day 7 Initial Day 7 Initial Day 7 2.5 mg/mL/50 mM buffer											
6.88	6.87	2.49	2.38	N.D	<0.02	1.13	2.5 mg/mL/50 mM buffer (N.sub.2 sparged)	6.88	6.87	2.49	—
2.55	N.D	<0.02	ND	<0.02	2.5 mg/mL/50 mM buffer (2-8° C.)	6.88	6.89	2.49	2.53	N.D	<0.02
<0.02	2.5	mg/mL/50 mM buffer	6.83	—	2.10 — 3.57 — (UV light exposure)	2.5	mg/mL/50 mM buffer	6.84	—	2.50	—
—	—	—	—	—	N.D <0.02 — (UV light exposure Control)						

[0199] Noticeably, the go-to buffers stored under ambient conditions (1 week at 40° C.), without N.sub.2 sparging or control of UV light exposure, contained 1.13% of related substances after 1 week of storage. This is compared to the related substances formed under the same conditions for Britton-Robinson buffered formulations below.

## Development of Formulation

### Materials

[0200] Details of the DMT fumarate employed for stability purposes are provided in Table 3 and excipients used are listed in Table 4.

TABLE-US-00006 TABLE 3 DMT fumarate used for the stability study Material Batch Number Supplier DMT fumarate SPL026 Onyx Scientific, Sunderland											
---	--	--	--	--	--	--	--	--	--	--	--

TABLE-US-00007 TABLE 4 Excipients used for the formulation development study Material Batch Number Supplier Purified water Not applicable Elga dispenser, asset number ARC37642 Sodium chloride 17D194102 VWR di-Sodium hydrogen 1997160 Fisher Chemicals orthophosphate dihydrate Sodium dihydrogen 1724808 Fisher Chemicals orthophosphate dihydrate Volumetric 1M sodium 726144 Scientific Laboratory hydroxide solution Supplies Glacial acetic acid 1727841 Fisher Chemicals											
---	--	--	--	--	--	--	--	--	--	--	--

### Equipment

[0201] Equipment, excluding standard laboratory glassware, used throughout the studies is listed in

Table 5. Calibration and verification of equipment were performed in accordance with standard operating procedures for all measurements, as required.

TABLE-US-00008 TABLE 5 Typical equipment used during the formulation development study

Item	Make and Model	Asset Number	Balance	Mettler Toledo, MX5	32721	Balance Sartorius, ME215S	31476	Single Stir Plate Bibby HB502	20234	pH Meter Mettler Toledo, MP225	20322
Osmolality	Advanced Instruments Osmo 1	38564	Filters Millex MP PES 0.22 µm	n/a	Light Box Heraeus SunTest 28 694						

#### Osmolality Readings

[0202] Osmolality readings were obtained using an Advanced Instruments Osmo1 instrument. A single sample syringe was used to introduce the sample into the osmometer, which employed the industry-preferred principles of freezing point depression to determine osmolality accurately and precisely.

[0203] Instrument verification was performed using 50, 850 and 2000 mOsm/kg H.sub.2O calibration standards prior to analysis, for confirmation of accuracy.

#### pH Readings

[0204] pH readings were obtained using a Mettler Toledo MP225 pH meter. The electrode probe was inserted into the test solutions, contained in a glass vial, with brief stirring at ambient temperature.

[0205] Instrument verification was performed prior and post each use using, as supplied, pH buffer solutions over the range pH 1.68 to 10.01 for confirmation of accuracy.

#### High Performance Liquid Chromatography (HPLC)

[0206] The following HPLC parameters were employed to assess assays and the quantity of related substances (substances resulting from DMT fumarate degradation) of solutions of DMT fumarate that were prepared as part of the formulation development. [0207] Column: YMC-Triart Phenyl; 150×4.6 mm, 3 µm, [0208] Mobile phase A: Water:Trifluoroacetic acid (100:0.05 v/v) [0209] Mobile phase B: Acetonitrile:Trifluoroacetic acid (100:0.05 v/v) [0210] Diluent: Acetonitrile:Water (50:50)

TABLE-US-00009 Time (min) % A % B Gradient timetable: 0.0 95 5 13.0 62 38 26.0 5 95 30.5 5 95 31.0 95 5 [0211] Flow rate: 1.0 mL.Math.min.sup.-1 [0212] Column temperature: 30° C. [0213] Injection volume: 7.5 UL [0214] Needle wash: Water:Acetonitrile (50:50) [0215] Seal Wash: Water:Acetonitrile (50:50) [0216] Run time: 35 minutes [0217] Detection Wavelength: 220 nm

#### Formulation Development

[0218] The solubility of DMT fumarate was initially assessed over a range of different pH values, from pH 4 to pH 10. Formulations were then prepared at the target concentration of 2.5 mg/mL of DMT fumarate over a pH range of pH 4 to pH 9.

#### Solubility of DMT Fumarate at Different pH Values

[0219] Seven solutions, each containing a concentration of 20 mg/ml of DMT fumarate were prepared in Britton-Robinson (B-R) buffer solution. On dissolution of DMT fumarate in each test formulation (DMT fumarate was very soluble, needing only swirling and shaking in each), the pH of each test formulation was then adjusted to pH 4, 5, 6, 7, 8 and 9 using sodium hydroxide solution.

[0220] Solubility of a concentration of 20 mg/ml of DMT fumarate was confirmed at pH 4, 5, 6 and 7—these solutions were clear and colourless. The sample at pH 8 was hazy and the samples at pH 9 and pH 10 contained a precipitate. Following overnight storage under ambient conditions, the pH of each solution was measured and the results showed no changes from the initial pH values. Each sample was then filtered and analysed for content. Each solution, including the high pH solutions where precipitate was present, contained approximately the same content of DMT fumarate.

#### pH Stability

[0221] pH-stability of DMT fumarate at a concentration of 2.5 mg/ml was assessed in 40 mM Britton-Robinson buffer solution over the buffer solution range pH 4 to 9 (nominal). The pH of

each formulation was measured at preparation, following 7 days storage at 40° C. and then further storage over an additional 3 days at 40° C. and 7 days at 50° C. (so a total further storage of 10 days). Analysis of these formulations was performed on preparation, and then after 7 and 17 days storage for content (assay) and related substances.

[illegible]



these both buffer well at the optimal pH for stability, and assess both sodium chloride and dextrose as tonicity agents.

#### Formulation Preparation

[0230] Details of each individual formulation (numbered 1 to 8) are presented in Table 12 and Table 13. For each formulation, the requisite acid and tonicity agent was dissolved in 80 ml of water. The pH of this solution was then adjusted to pH 4 (+0.5) with 1 M sodium hydroxide solution. The drug substance was then dissolved, the pH adjusted to pH 4 (+0.1) with more 1 M sodium hydroxide solution and then made to volume with water and final pH check and adjusted as required. For each formulation, the volume of sodium hydroxide used was documented. The composition of each formulation is presented below in Table 12 (saline) and Table 13 (dextrose).

[0231] An aliquot of each formulation was taken for assay/related substances and osmolality check. The remainder of each formulation was filtered (filter size 0.2 µm) into a clear glass multi-dose vial, sparged with nitrogen, capped and placed into storage (60° C.) for 14 days. The 40 mM phosphate/dextrose formulation (formulation 8) was split into two aliquots with one aliquot stored in an amber glass multi-dose vial and one aliquot in a clear glass multi-dose vial.

TABLE-US-00016 TABLE 12 Candidate SPL026 formulation preparations (saline) Saline Formulations

Formulation Number	Ingredient 1	2	3	4	SPL026	398 mg	398 mg	398 mg	398 mg		
Acetic acid	120 mg	240 mg	—	—	(20 mM)	(40 mM)	Ortho-phosphoric	—	—	231 mg	461 mg
acid (85%)	(20 mM)	(40 mM)			Sodium chloride	780 mg	720 mg	780 mg	720 mg	Sodium hydroxide	38.8 mg
	60.8 mg	103.2 mg	185.6 mg	Volume prepared	100	100	100	100	(mL)		

TABLE-US-00017 TABLE 13 Candidate SPL026 formulation preparations (dextrose) Dextrose Formulations

Formulation Number	Ingredient 5	6	7	8	SPL026	398 mg	398 mg	398 mg	398 mg		
Acetic acid	120 mg	240 mg	—	—	(20 mM)	(40 mM)	Ortho-phosphoric	—	—	231 mg	461 mg
acid (85%)	(20 mM)	(40 mM)			Dextrose	4300 mg	3900 mg	4300 mg	3900 mg	Sodium hydroxide	27.6 mg
	47.2	96.4	175.2	mg	Volume prepared	100	100	100	100	(mL)	

#### Results

[0232] Concentration, osmolality and pH results for formulations at preparation and following storage at 60° C. are presented in Table 14. Related substances results following storage are presented in Table 15.

[0233] All formulations on preparation were clear colourless solutions. No related substances were present in any of the formulations following preparation.

[0234] Following removal from storage all formulations in terms of their appearance were no longer colourless but had acquired to varying degrees a hint of beige but all remained clear, colour was most pronounced in formulation 5 (20 mM acetate buffer/dextrose) which had the greatest concentration of related substances.

[0235] Osmolality and pH were confirmed as stable for each tested formulation with no significant changes.

[0236] Total related substances as peaks of more than 0.05% of total peak area ranged between 0.07% up to 0.52%. These data would suggest that for SPL026 saline is the preferred tonicity agent over dextrose.

[0237] All Day 14 results for the 40 mM phosphate/dextrose formulation stored in amber glass mirrored the clear glass results confirming that clear glass/amber glass storage has no impact on stability in terms of these storage conditions but given the previously noted light instability amber glass should be used as the primary pack.

TABLE-US-00018 TABLE 14 Candidate SPL026 formulation results, assay, osmolality and pH

Concentration	Osmolality (mg .Math. mL.sup.-1)	(mOsm/kg)	pH 14 days	14 days	14 days	No. Vehicle composition
Day 0	60° C.	Day 0	60° C.	Day 0	60° C.	1
20 mM acetate/saline	2.50	2.52	305	300	3.94	3.94
2	40 mM acetate/saline	2.54	2.54	307	318	3.98
4.01	3	20 mM phosphate/saline	2.55	2.54	315	319
4.01	4.02	4	40 mM phosphate/saline	2.53	2.46	330
347	4.00	3.99	5	20 mM acetate/dextrose	2.50	2.46
300	308	3.97	4.05	6	40 mM acetate/dextrose	2.51
2.47	304	310	4.02			

4.06 7 20 mM phosphate/dextrose 2.55 2.51 320 320 4.02 4.04 8 40 mM phosphate/dextrose 2.58 2.48 339 336 4.01 4.04 8.sup.a 40 mM phosphate/dextrose — 2.49 — 334 — 4.02 .sup.astored in amber glass

TABLE-US-00019 TABLE 15 Related substances assay for candidate SPL026 formulations following storage at 60° C. for 14 days RRT and percentage area of total peak area of peaks >0.05% of total peak area No. 0.62 0.72 0.80 0.91 1.60 1.61 Total 1 — — — — — 0.11 0.11 2 — — — — — 0.08 0.08 3 — — — — — 0.09 0.09 4 — — — — — 0.07 0.07 5 0.05 0.05 0.07 0.07 0.08 0.20 0.52 6 — — — — — 0.13 0.13 7 — — — — — 0.05 — 0.16 0.21 8 — — — — — 0.05 — 0.15 0.20 8.sup.a — — — — — 0.07 — 0.23 0.30 .sup.astored in amber glass

## CONCLUSION

[0238] All the candidate formulations following the 14-day stability assessment were stable in terms of osmolality and pH. For formulations 1 to 3 there was no change in achieved concentration and for formulations 5-8 changes (losses on storage) were small (<0.1 mg.Math.mL.sup.-1). For each candidate formulation, related substances were very low at between 0.07 to 0.52% of the total peak area. More related substances were observed in the dextrose formulations than in the saline formulations.

[0239] Following review of the data the following SPL026 formulation was chosen (herein referred to as SPL026 formulation 10). This is based on formulation 1 but the acetate buffer content is increased slightly to ensure robust buffering in the formulation over the shelf life but at a low enough level to ensure that blood buffering effects are minimal. Sodium chloride levels were dropped slightly to compensate for the additional acetate but to maintain an iso-osmotic solution. This formulation was prepared and analysed for assay, pH and osmolality on preparation. Preparation details and achieved results are presented below in Table 16.

TABLE-US-00020 TABLE 16 SPL026, 2.5 mg/mL formulation 10 preparation and results  
Preparation SPL026 398 mg\* Acetic acid 150 mg Sodium chloride 760 mg Sodium hydroxide q.s. to pH 4.0 ≈ 40 mg Volume prepared 100 (mL) Result Appearance Clear colourless solution Osmolality 299 (mOsm/kg) pH 3.96 Assay (mg.mL.sup.-1) 2.49 \*398 mg SPL026 (DMT fumarate) is equivalent to 250 mg of free base (DMT)

Recommended Process Overview for Formulation Preparation (Batch Size 100 mL)

## Preparation of Vehicles

### Preparation of 1M Sodium Hydroxide Solution (100 mL)

[0240] 1) Weigh 4 g of sodium hydroxide pellets into a suitably sized beaker. [0241] 2) Dispense 80 ml of Water for injection (WFI) water into the beaker. [0242] 3) Magnetically stir to achieve dissolution and allow the solution to cool to laboratory temperature. [0243] 4) Add purified water to make up to 100 mL. [0244] 5) Transfer into a type 1 borosilicate glass container.

Preparation of Formulation 10 (2.5 mg/mL) [0245] 1) Accurately weigh 760 mg of sodium chloride into a suitable container [0246] 2) Weigh by difference the required amount of drug substance into a suitable container (glass weigh boat). Ensure that mass of drug substance taken includes correction for salt and purity. [0247] 3) Carefully transfer the weighed drug substance into a beaker. Rinse out the weighing container with WFI ensuring no solids remain. Add further WFI to the drug substance up to ¾ of the required total volume and magnetically stir to dissolve. Add batch quantity of acetic acid (note acetic acid is volatile and so this step must be performed immediately after weighing). [0248] 4) Add the pre-weighed sodium chloride. [0249] 5) Once dissolution is complete, adjust the pH of the formulation to pH 4 (+0.1) with dropwise addition of the freshly prepared 1M sodium hydroxide solution whilst continually stirring. [0250] 6) Make to volume in a suitable container and lastly check the pH is pH 4 (+0.1) and adjust if required. [0251] 7) The drug product solution is clear with a very slight hint of a beige colour. This colour is removed on filtration (step 6) to leave a colourless solution filtration. [0252] 8) Bubble nitrogen through the formulation until the measured dissolved oxygen content is below 2 ppm. [0253] 9) Syringe filter the solution either 0.22 µm or 0.2 µm into an amber glass multi-dose vial.

[0254] Below, reference is made to both technical and clinical batches of Formulation 10. Each of these have identical composition, but the technical batch was not made to GMP.

#### Stability of Formulation 10 (Technical Batch)

##### Appearance and pH

[0255] Appearance and pH results for formulation 10 (technical batch) at preparation and following storage at 2-8° C., 25° C./60% relative humidity (RH) and 40°/75% RH are presented in Table 17. At T=0 (the initial analysis), the pH met the specification of 3.8-4.2 and the osmolality met the specification of 270-330 mOsm/kg (296 mOsm/kg). By T=9 months, there are no significant changes in appearance on storage at 2-8° C. or at 25° C./60% RH. However, at 40° C./75% RH there is an obvious colour change in the product at T=6 months. There was an indication of colour change at the 2 and 3 month points on storage at 40° C./75% RH when compared to the other storage conditions, but at 6 months the colour change was obvious. There was no change in pH by the 9 month time point on storage at 2-8° C. or 25° C./60% RH or by the 6 month time point on storage at 40° C./75% RH.

TABLE-US-00021 TABLE 17 Appearance and pH of formulation 10 after storage under the conditions specified

Time-point/ months	Appearance	pH
Initial (T = 0)	Clear colourless liquid, free from visible particulates	4.0
2-8° C. T = 1	Clear colourless liquid, free from visible particulates	4.0
T = 2	Clear colourless liquid, free from visible particulates	4.0
T = 3	Clear colourless liquid, free from visible particulates	4.0
T = 6	Clear colourless liquid, free from visible particulates	4.0
T = 9	Clear colourless liquid, free from visible particulates	4.0
25° C./ T = 1	Clear colourless liquid, free from visible particulates	4.0
60% T = 2	Clear colourless liquid, free from visible particulates	4.0
RH T = 3	Clear colourless liquid, free from visible particulates	4.0
T = 6	Clear colourless liquid, free from visible particulates	4.0
T = 9	Clear colourless liquid, free from visible particulates	4.0
40° C./ T = 1	Clear colourless liquid, free from visible particulates	4.0
75% T = 2	Clear colourless liquid, free from visible particulates	4.0
RH T = 3	Clear colourless liquid, free from visible particulates	4.0
T = 6	Clear light yellow liquid, free from visible	4.0

##### Sub-Visible Particulates

[0256] The number of sub-visible particles within formulation 10 (technical batch) at preparation and following storage at 2-8° C., 25° C./60% relative humidity (RH) and 40°/75% RH is presented in Table 18. After 6 months storage at 2-8° C., 25° C./60% RH and 40° C./75% RH there is no significant change in sub-visible particulates (see Table 18).

TABLE-US-00022 TABLE 18 Sub-visible particulates of formulation 10 after storage under the conditions specified

Time-point/ Count/vial	≥10 µm	≥25 µm	Pass/fail
Initial (T = 0)	5	1	Pass
5° C. T = 6	3	0	Pass
25° C./60% RH T = 1	9	0	Pass
T = 6	3	0	Pass
40° C./75% RH T = 1	4	0	Pass
T = 6	14	1	Pass

.sup.1Where fill volume is 10.5 mL .sup.2Number particles with diameter ≥10 µm per vial <6000 = PASS and number particles with diameter ≥25 µm per vial <600 = PASS

##### Extractable Volume

[0257] The extractable volume from 6 vials of formulation 10 (technical batch) is presented in Table 19. The density used to calculate the volume was taken from the placebo data used in batch manufacture (1.008 g/cm<sup>3</sup>). The extractable volume meets the specification of NLT 10.0 mL.

TABLE-US-00023 TABLE 19 Extractable volume of formulation 10

Sample	Sample weight (g)	Extractable volume (mL)
1	10.2791	10.1975
2	10.4192	10.3365
3	10.2732	10.1917
4	10.3162	10.2343
5	10.2696	10.1881
6	10.2516	10.1702
Mean	10.3	10.2

##### Percentage Recovery

[0258] The percentage recovery of SPL026 (free base, mg/mL) within formulation 10 (technical batch) at preparation and following storage at 2-8° C., 25° C./60% relative humidity (RH) and 40°/75% RH is presented in Table 20. There were no significant changes in the purity of the formulation after 9 months storage at 2-8° C. or 25° C./60% RH and 6 months storage at 40° C./75% RH. All recoveries versus the theoretical concentration and T=0 are within the specification

of 90.0-105.0%

TABLE-US-00024 TABLE 20 Percentage recovery of SPL026 from formulation 10 after storage under the conditions specified Time-point/months 1 2 Mean Rec./theory.sup.1 (%) Rec./T = 0.sup.2 (%) Pre-filtration 2.518 2.506 2.51 100.5 — Post-filtration 2.516 2.504 2.51 100.4 99.9.sup.3 Initial (T = 0) 2.503 2.501 2.50 100.1 — 5° C. T = 1 2.508 2.508 2.51 100.2 100.3 T = 2 2.538 2.539 2.54 101.5 101.5 T = 3 2.512 2.523 2.52 100.8 100.7 T = 6 2.515 2.525 2.52 100.8 100.7 T = 9 2.511 2.504 2.51 100.3 100.2 25° C./60% RH T = 1 2.508 2.509 2.51 100.3 100.3 T = 2 2.541 2.535 2.54 101.5 101.4 T = 3 2.525 2.524 2.52 101.0 100.9 T = 6 2.497 2.525 2.51 100.4 100.4 T = 9 2.475 2.496 2.49 99.4 99.3 40° C./75% RH T = 1 2.510 2.499 2.50 100.1 100.2 T = 2 2.538 2.536 2.54 101.5 101.4 T = 3 2.505 2.501 2.50 100.1 100.0 T = 6 2.494 2.494 2.49 99.8 99.7 .sup.1As percentage of theoretical 2.5 mg/mL. .sup.2As percentage of initial (T = 0) result. .sup.3As percentage of pre-filtration result.

#### Purity/Related Substances

[0259] The purity of SPL026 (free base, mg/mL) and the amount of impurities observed at different retention times within formulation 10 (technical batch) at preparation and following storage at 2-8° C., 25° C./60% relative humidity (RH) and 40°/75% RH are presented in Table 21, Table 22 and Table 23, respectively. The total impurities observed in formulations after 9 months storage at 2-8° C. and 25° C./60% RH have increased slightly due to the impurity at RRT 1.04 being observed for the first time and above the LOQ limit at both conditions. Increases in total impurities were noted for the sample stored at 40° C./75% RH from 2 months. An increase in the impurity observed at RRT 1.60 is noted from T=2 months at 40° C./75% RH in comparison to the other storage conditions, which may be attributed to the higher storage temperature of 40° C.

TABLE-US-00025 TABLE 21 Purity/related substances of SPL026 from formulation 10 after storage at 2-8° C. Time-point/months and amount (% w/w) RRT Initial (T = 0) 1 2 3 6 9 SPL026 (free base).sup.1 100.0 100.0 100.0 100.0 100.0 100.0 (99.95) 1.04 ND ND ND ND ND 0.05 1.60 ND ND <LOQ <LOQ <LOQ <LOQ Total.sup.2 ND ND <LOQ <LOQ <LOQ 0.05 .sup.1SPL026 purity calculated as 100 – total related substances. .sup.2Sum of related substances ≥0.05%

TABLE-US-00026 TABLE 22 Purity/related substances of SPL026 from formulation 10 after storage at 25° C./60% RH Time-point/months and amount (% w/w) Initial RRT (T = 0) 1 2 3 6 9 SPL026 (free base) .sup.1 100.0 100.0 100.0 100.0 100.0 100.0 (99.95) 1.04 ND ND ND ND ND 0.05 1.18 ND ND <LOQ ND ND ND 1.60 ND ND <LOQ <LOQ <LOQ <LOQ Total.sup.2 ND ND <LOQ <LOQ <LOQ 0.05 .sup.1 SPL026 purity calculated as 100 – total related substances. .sup.2Sum of related substances ≥0.05%

TABLE-US-00027 TABLE 23 Purity/related substances of SPL026 from formulation 10 after storage at 40° C./75% RH Time-point/months and amount (% w/w) Initial RRT (T = 0) 1 2 3 6 0.49 ND ND ND ND 0.05 0.70 ND ND <LOQ <LOQ 0.05 0.77 ND ND ND <LOQ <LOQ 0.81 ND ND <LOQ <LOQ 0.05 0.92 ND ND <LOQ <LOQ 0.06 SPL026 100.0 100.0 99.9 99.9 99.7 (free base) .sup.1 1.19 ND ND ND ND <LOQ 1.22 ND ND ND ND <LOQ 1.60 ND ND 0.05 0.07 0.10 Total.sup.2 ND ND 0.05 0.07 0.31 .sup.1 SPL026 purity calculated as 100 – total related substances. .sup.2Sum of related substances ≥ 0.05%

#### Stability of Formulation 10 (Clinical Batch)

[0260] At T=0 (the initial analysis), the pH met the specification of 3.8-4.2, the osmolality met the specification of 270-330 mOsm/kg (being 306 mOsm/kg), and the extractable volume met the NLT of 10.0 mL (being 10.4 mL). In addition, at T=0, the number of sub visible particulates was 4 particles/vial of a size ≥10 µm and 0 particles/vial of a size ≥25 µm, which is well within the specification of not more than 6000 particles/vial of a size ≥10 µm and not more than 600 particles/vial of a size ≥25 µm. At T=0, the UV-vis spectrum of the formulation conformed to reference spectra (λ.sub.max at 221±3 nm and 279±3 nm), and the signal was observed at ±2% retention time of the reference standard. Finally, at T=0, container closure integrity testing showed no dye ingress, and the formulation was sterile, comprising <0.01 EU/mL of bacterial endotoxins,

which is well within the specification of 20.5 EU/mL.

#### Appearance and pH

[0261] Appearance and pH results for formulation 10 (clinical batch) at preparation and following storage at 2-8° C., 25° C./60% relative humidity (RH) and 40°/75% RH are presented in Table 24. By T=3 months, there are no significant changes in appearance on storage at 2-8° C., at 25° C./60% RH or at 40° C./75% RH. There was an indication of colour change at the 3 month point. There was no change in pH.

TABLE-US-00028 TABLE 24 Appearance and pH of formulation 10 after storage under the conditions specified Time-point/months Appearance pH Initial (T = 0) Clear colourless liquid, practically free from 4.0 particulates 2-8° C. T = 1 Clear colourless liquid, practically free from 3.9 particulates T = 3 Clear pale yellow liquid, practically free from 4.0 particulates 25° C./60% RH T = 1 Clear colourless liquid, practically free from 4.0 particulates T = 3 Clear pale yellow liquid, practically free from 4.0 particulates 40° C./75% RH T = 1 Clear colourless liquid, practically free from 3.9 particulates T = 3 Clear pale yellow liquid, practically free from 4.0 particulates

#### Percentage Recovery

[0262] The percentage recovery of SPL026 (free base, mg/mL) within formulation 10 (clinical batch) at preparation and following storage at 2-8° C., 25° C./60% relative humidity (RH) and 40°/75% RH is presented in Table 25. There were no significant changes in the purity of the formulation after 3 months storage at 2-8° C. or 25° C./60% RH and 40° C./75% RH. All recoveries versus the theoretical concentration and T=0 are within the specification of 95.0-105.0%

TABLE-US-00029 TABLE 25 Percentage recovery of SPL026 from formulation 10 after storage under the conditions specified Time-point/months Percentage recover of SPL026 (%) Initial (T = 0) 103.6 2-8° C. T = 1 102.9 T = 3 103.2 25° C./60% RH T = 1 102.7 T = 3 103.0 40° C./75% RH T = 1 102.6 T = 3 102.5

#### Purity/Related Substances

[0263] The purity of SPL026 (free base, mg/mL) and the amount of impurities observed at different retention times within formulation 10 (clinical batch) at preparation and following storage at 2-8° C., 25° C./60% relative humidity (RH) and 40°/75% RH are presented in Table 26, Table 27 and Table 28, respectively.

TABLE-US-00030 TABLE 26 Purity/related substances of SPL026 from formulation 10 after storage at 2-8° C. Time-point/months and amount (Area %) Initial RRT (T = 0) 1 3 1.11 <LOQ <LOQ <LOQ 1.59-1.60 0.07 0.08 0.07 Total.sup.2 0.07 0.08 0.07 .sup.2Sum of related substances ≥ 0.05%

TABLE-US-00031 TABLE 27 Purity/related substances of SPL026 from formulation 10 after storage at 25° C./60% RH Time-point/months and amount (Area %) Initial RRT (T = 0) 1 3 1.11 <LOQ <LOQ <LOQ 1.59-1.60 0.07 0.08 0.07 Total.sup.2 0.07 0.08 0.07 .sup.2Sum of related substances ≥ 0.05%

TABLE-US-00032 TABLE 28 Purity/related substances of SPL026 from formulation 10 after storage at 40° C./75% RH Time-point/months and amount (Area %) Initial RRT (T = 0) 1 3 0.56 <LOQ <LOQ <LOQ 0.70 <LOQ <LOQ <LOQ 0.82 <LOQ <LOQ <LOQ 0.94 <LOQ <LOQ <LOQ 1.11 <LOQ <LOQ <LOQ 1.59-1.60 0.07 0.08 0.09 Total.sup.2 0.07 0.08 0.09 .sup.2Sum of related substances ≥ 0.05%

## Claims

1. A pharmaceutical formulation suitable for injection, comprising: a salt of a dimethyltryptamine compound substituted with deuterium and an acid selected from the group consisting of fumaric acid, tartaric acid, citric acid, acetic acid, lactic acid, gluconic acid, benzoic acid, salicylic acid, and succinic acid; a buffer which is separate to the salt; and water, wherein the formulation has a pH of about 3.5 to about 6.5 and an osmolality of about 250 to about 350 mOsm/Kg.

2. The formulation of claim 1, wherein the pH is from about 3.75 to about 5.75.
3. The formulation of claim 1, wherein the pH is from about 3.75 to about 4.25.
4. The formulation of claim 1, wherein the dimethyltryptamine compound is substituted with deuterium at one or more positions selected from the  $\alpha$ ,  $\beta$  and dimethyl carbon atoms.
5. The formulation of claim 1, wherein the dimethyltryptamine compound is a compound of Formula I: ##STR00008## wherein: R.sup.4 and R.sup.5 are both H; and each .sup.XH and each .sup.YH is independently selected from H and D, wherein at least one of .sup.XH and .sup.YH is D.
6. The formulation of claim 1, comprising any one of or a combination of  $\alpha$ -monodeutero-N,N-dimethyltryptamine,  $\alpha,\alpha$ -dideutero-N,N-dimethyltryptamine,  $\alpha,\beta$ -dideutero-N,N-dimethyltryptamine,  $\alpha,\alpha,\beta$ -trideutero-N,N-dimethyltryptamine,  $\alpha,\beta,\beta$ -trideutero-N,N-dimethyltryptamine and  $\alpha,\alpha,\beta,\beta$ -tetra-deutero-N,N-dimethyltryptamine.
7. The formulation of claim 6, wherein the formulation comprises  $\alpha,\alpha$ -dideutero-N,N-dimethyltryptamine.
8. The formulation of claim 6, wherein the formulation comprises  $\alpha$ -monodeutero-N,N-dimethyltryptamine.
9. The formulation of claim 6, wherein the formulation comprises  $\alpha,\alpha$ -dideutero-N,N-dimethyltryptamine and  $\alpha$ -monodeutero-N,N-dimethyltryptamine.
10. The formulation of claim 1, wherein the dimethyltryptamine compound is substituted with deuterium at one or more positions on the dimethyl carbon atoms.
11. The formulation of claim 10, wherein the dimethyltryptamine compound is substituted with deuterium at all positions on the dimethyl carbon atoms.
12. The formulation of claim 10, wherein the dimethyltryptamine compound is substituted with deuterium at one or more positions on the dimethyl carbon atoms and at one or both  $\alpha$ -carbon positions.
13. The formulation of claim 10, wherein the dimethyltryptamine compound is substituted with deuterium at one or more positions on the dimethyl carbon atoms, at one or both  $\alpha$ -carbon positions ( $\alpha$ -monodeutero or  $\alpha,\alpha$ -dideutero), and at one or both  $\beta$ -carbon positions ( $\beta$ -monodeutero or  $\beta,\beta$ -dideutero).
14. The formulation of claim 10, wherein the dimethyltryptamine compound is substituted with deuterium at all positions on the dimethyl carbon atoms and at one or both  $\alpha$ -carbon positions.
15. The formulation of claim 10, wherein the dimethyltryptamine compound is substituted with deuterium at all positions on the dimethyl carbon atoms and at both  $\alpha$ -carbon positions.
16. The formulation of claim 10, wherein the dimethyltryptamine compound is substituted with deuterium at all positions on the dimethyl carbon atoms, at both  $\alpha$ -carbon positions, and at both  $\beta$ -carbon positions.
17. The formulation of claim 1, wherein the acid is fumaric acid.
18. The formulation of claim 1, wherein the buffer comprises sodium acetate and acetic acid, or potassium acetate and acetic acid; or the buffer comprises acetate salt, acetic acid, citrate salt, citric acid, ascorbate salt, ascorbic acid, phosphate salt, or phosphoric acid.
19. The formulation of claim 1, further comprising a tonicity agent.
20. A method of treating a psychiatric or neurological disorder comprising administering to a patient in need thereof a formulation as defined in claim 1.
21. The method of claim 20, wherein the psychiatric or neurological disorder is an obsessive compulsive disorder, a depressive disorder, an anxiety disorder, substance abuse, or an avolition disorder.
22. The method of claim 21, wherein the depressive disorder is major depressive disorder or treatment resistant depression.
23. The method of claim 21, wherein the anxiety disorder is generalised anxiety disorder, phobia, panic disorder, social anxiety disorder, or post-traumatic stress disorder.

**24.** The formulation of claim 1, comprising any one of or a combination of  $\alpha$ -monodeutero-N,N-dimethyltryptamine,  $\alpha,\alpha$ -dideutero-N,N-dimethyltryptamine,  $\alpha,\beta$ -dideutero-N,N-dimethyltryptamine,  $\alpha,\alpha,\beta$ -trideutero-N,N-dimethyltryptamine,  $\alpha,\beta,\beta$ -trideutero-N,N-dimethyltryptamine and  $\alpha,\alpha,\beta,\beta$ -tetra-deutero-N,N-dimethyltryptamine; and wherein the buffer comprises sodium acetate and acetic acid, or potassium acetate and acetic acid; or the buffer comprises acetate salt, acetic acid, citrate salt, citric acid, ascorbate salt, ascorbic acid, phosphate salt, or phosphoric acid.

---