



US 20250261651A1

(19) **United States**(12) **Patent Application Publication****Dow et al.**(10) **Pub. No.: US 2025/0261651 A1**(43) **Pub. Date: Aug. 21, 2025**(54) **INSECT NEUROPEPTIDES 3****Publication Classification**(71) Applicant: **Solasta Bio Limited**, Glasgow (GB)(72) Inventors: **Julian A.T. Dow**, East Dunbartonshire (GB); **Shireen A. Davies**, East Dunbartonshire (GB); **Yousef Abul-Haija**, East Dunbartonshire (GB); **Lewis Archibald**, Glasgow (GB)(73) Assignee: **Solasta Bio Limited**, Glasgow (GB)(21) Appl. No.: **19/090,002**(22) Filed: **Mar. 25, 2025****Related U.S. Application Data**

(62) Division of application No. 18/492,006, filed on Oct. 23, 2023, now Pat. No. 12,279,621.

(51) **Int. Cl.****A01N 63/50** (2020.01)**A01P 7/04** (2006.01)(52) **U.S. Cl.**CPC **A01N 63/50** (2020.01); **A01P 7/04** (2021.08)


(57)

ABSTRACT

The present invention relates to natural or natural-like analogues of insect pyrokinin neuropeptide [Hy]-NAD-EDQQSVDFTPRL-[NH₂] (SEQ ID NO:2) having activity against insects, for example hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insects, such as aphids, moths and fruit flies, and their use as insect control agents (e.g. insecticides) and plant protection agents.

Specification includes a Sequence Listing.

SEQ ID NO: 2



SEQ ID NO: 2




FIG. 1




FIG. 3

PYROKININ VS *SPODOPTERA FRUGIPERDA*
SB-P-80




FIG. 4

INSECT NEUROPEPTIDES 3

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a divisional of U.S. application Ser. No. 18/492,006, filed Oct. 23, 2023, the entirety of which is hereby incorporated by reference.

REFERENCE TO SEQUENCE LISTING

[0002] This application contains a Sequence Listing submitted as a computer readable form named "P365404US_corr.xml", having a size in bytes of 25,793 bytes, and created on Jul. 9, 2024 ("production date"). The information contained in this computer readable form is hereby incorporated by reference in its entirety.

FIELD OF THE INVENTION

[0003] The present invention relates to natural or natural-like analogues of insect pyrokinin neuropeptides having activity against insects, for example hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insects, such as aphids, moths and fruit flies, and their use as insect control agents (e.g. insecticides) and plant protection agents.

BACKGROUND

[0004] With a global dependence on broad-spectrum insecticides, the damaging effects of which are well documented, there is increasing need for the development of greener, target-specific insecticides with which to protect valuable crop plants. The development and employment of neuropeptides, and their synthetic analogues offers a promising avenue in the drive for greener and target-specific insecticidal agents.

[0005] Within insects, neuropeptides are regulatory peptides with functional roles in growth and development, behaviour and reproduction, metabolism and homeostasis, and muscle movement. Insect neuropeptide families are large and include the insect kinins and CAPA (CAPA, CAP2b, CAPA3) neuropeptides, AKH peptides, and the pyrokinin peptides.

[0006] Due to their high specificity, insect neuropeptides and their cognate receptors (G-protein coupled receptors, GPCRs) may be developed towards insecticidal agents to selectively reduce the fitness of target pest insects, whilst minimising detrimental environmental impacts, and minimising harm to other non-target species such as pollinators, which have seen dangerously declining population numbers due to pesticide/insecticide use.

[0007] There have been attempts in the art to provided modified synthetic analogues of such insect neuropeptides for use as targeted insecticides, which show promising results. However, whilst desirable and effective, these modified peptides may be challenging to bring to market in many territories in which the use of such modified peptides is heavily regulated.

[0008] The present invention aims to solve one or more of the above-mentioned problems, and provides novel natural pyrokinin neuropeptides, and natural-like variants or analogues thereof for use as insect control agents and plant protection agents.

SUMMARY OF THE INVENTION


[0009] The inventors have discovered new pyrokinin peptides, and natural-like variants or analogues thereof, having insecticidal activity against insects such as hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insects, and so potentially finding use as pest control agents or insecticides, while having little or no effect against important pollinator species such as bees.

[0010] Thus, in a first aspect, the invention provides an insecticidal compound having the formula (I) below, or a salt or solvate thereof:



wherein:

[0011] R^1 is hydrogen (which may be designated "H-" or "Hy-"), C_{1-4} alkyl (e.g. methyl, ethyl, propyl, butyl), formyl, acyl, fatty acyl, a sugar moiety, phosphate or sulphate, or R^1 is a pyroglutamate group of the formula:



[0012] and any alkyl, formyl, acyl or fatty acyl may optionally be substituted with one or more groups selected from oxo or C_{1-6} alkyl or a sugar moiety, phosphate or sulphate;

[0013] Y^1 is absent or is a peptide comprising from 1 to 2 amino acids;

[0014] Z is a peptide according to formula



[0015] Y^2 is absent or is a peptide comprising from 1 to 2 amino acids;

[0016] R^2 is NH_2 , NR^{2a}H , $\text{NR}^{2a}\text{R}^{2b}$, OH or OR^{2a} , wherein each of R^{2a} and R^{2b} if present are independently C_{1-6} alkyl (e.g. methyl, ethyl, propyl, butyl, pentyl or hexyl).

[0017] In another aspect, there is provided a composition comprising a compound as defined herein, or a salt thereof, in admixture with one or more solvents, carriers, diluents, adjuvants, preservatives, dispersants, emulsifying agents, or synergists. Suitably, the composition is an agricultural composition, an insect control composition or plant protection composition.

[0018] In another aspect, there is provided the use of a compound as defined herein, or a salt thereof, or a composition as defined herein, as an insect control agent. The use may be as an insecticide against insects that encode the pyrokinin peptide. The use may be as an insecticide against hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insects, preferably hemipteran, dipteran and/or lepidopteran insects, most preferably lepidopteran insects such as moths.

[0019] In another aspect, there is provided a method of increasing insect mortality, the method comprising contacting an insect population with a compound as defined herein, or a salt thereof, or a composition as defined herein. Suitably, the insect is an insect that encodes the pyrokinin peptide, such as hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insects, preferably hemipteran, dip-

teran and/or lepidopteran insects, most preferably lepidopteran insects such as moths.

[0020] In another aspect, there is provided a method of increasing hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insect mortality, the method comprising contacting a hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insect population with a compound as defined herein, or a salt thereof, or a composition as defined herein. Preferably the insect is a lepidopteran insect such as moths.

[0021] In another aspect, there is provided the use of a compound as defined herein, or a salt thereof, or a composition as defined herein, as a plant protection agent, for protecting a plant against insects that encode the pyrokinin peptide. Preferably the insect is a lepidopteran insect such as moths.

[0022] In another aspect, there is provided the use of a compound as defined herein, or a salt thereof, or a composition as defined herein, as a plant protection agent, for protecting a plant against hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insects, preferably hemipteran, dipteran and/or lepidopteran insects, most preferably lepidopteran insects such as moths.

[0023] In another aspect, there is provided a method of inhibiting infestation of a plant by insects that encode the pyrokinin peptide, the method comprising contacting the plant with a compound as defined herein, or a salt thereof, or a composition as defined herein. Suitably, the compound or composition is applied to the plant while the plant is free or substantially free of insects that encode the pyrokinin peptide.

[0024] In another aspect, there is provided a method of inhibiting infestation of a plant by hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects comprising contacting the plant with a compound as defined herein, or a salt thereof, or a composition as defined herein. Suitably, the compound or composition is applied to the plant while the plant is free or substantially free of hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insects. Preferably the insects are selected from hemipteran, dipteran and/or lepidopteran insects, most preferably lepidopteran insects such as moths.

[0025] In another aspect, there is provided a method of reducing infestation of a plant by insects that encode the pyrokinin peptide, or of reducing load of insects that encode the pyrokinin peptide on a plant, the method comprising contacting the plant with a compound as defined herein, or a salt thereof, or a composition as defined herein.

[0026] In another aspect, there is provided a method of reducing hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insect infestation of a plant, or of reducing hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insect load on a plant, the method comprising contacting the plant with a compound as defined herein, or a salt thereof, or a composition as defined herein. Preferably the insects are selected from hemipteran, dipteran and/or lepidopteran insects, most preferably lepidopteran insects such as moths. In one embodiment, the compounds, compositions, and agents of the invention suitably have activity against the moth species *Plutella xylostella*. In one embodiment, the compounds, compositions, and agents of the invention suitably have activity against the moth species *Spodoptera frugiperda*.

[0027] Suitable insects which encode the pyrokinin peptide include hemipteran, dipteran, lepidopteran, blattodean and/or coleopteran insects, preferably hemipteran, dipteran and/or lepidopteran insects. In a preferred embodiment, the insects which encode the pyrokinin peptide are lepidopteran insects such as moths.

[0028] In another aspect, there is provided a method of producing an insecticidal compound according to the first aspect, the method comprising:

[0029] (a) chemically synthesising the insecticidal compound of formula (I) or a precursor thereof.

[0030] In one embodiment, the chemical synthesis comprises the method described in the examples. In one embodiment, the chemical synthesis comprises the method and steps shown in FIG. 2.

[0031] Further features of the aspects and embodiments of the invention will now be defined under the following headed sections. Any feature in any section may be combined with any aspect of embodiment in any workable combination.


DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0032] Throughout the present description and claims the conventional three-letter and one-letter codes for naturally occurring amino acids are used, i.e. A (Ala), G (Gly), L (Leu), I (Ile), V (Val), F (Phe), W (Trp), S (Ser), T (Thr), Y (Tyr), N (Asn), Q (Gln), D (Asp), E (Glu), K (Lys), R (Arg), H (His), M (Met), C (Cys) and P (Pro).

[0033] ‘Amino acid’ as referred to herein may refer to a naturally occurring amino acid or any other amino acid including synthetic amino acids, and non-proteinogenic amino acids. By “naturally occurring” in this context is meant the 20 amino acids encoded by the standard genetic code, sometimes referred to as proteinogenic amino acids.

[0034] The term amino acid is short for α -amino [alpha-amino]carboxylic acid. Each molecule contains a central carbon atom, called the α -carbon, to which both an amino and a carboxyl group are attached. The remaining two bonds of the α -carbon atom are generally satisfied by a hydrogen (H) atom and the side chain, shown as R below:



[0035] Unless otherwise specified, amino acid residues in peptides of the invention are of the L-configuration. As used herein, the terms “polypeptide”, “protein”, “peptide”, are used interchangeably, and refer to a polymeric form of amino acids of any length, which can include coded and non-coded amino acids. As used herein, amino acid residues will be indicated as above either by their full name or according to the standard three-letter or one-letter amino acid code.

[0036] The notation C_{x-xx} refers to the number of carbon atoms in a functional group. The number in the ‘x’ position is the lowest number of carbon atoms and the number in the ‘xx’ position denotes the highest number of carbon atoms.

For example, C₁₋₆-alkyl refers to alkyl groups as defined herein having from 1 to 6 carbon atoms.

[0037] The notation l, n or t are used herein in relation to various alkyl groups in the normal way. Specifically, the suffixes refer to the arrangement of atoms and denote straight chain ('n') or branched ('i' or 't') alkyl groups.

[0038] The term "alkyl" as used herein refers to a saturated linear or branched-chain monovalent hydrocarbon radical, wherein the alkyl radical may be optionally substituted. The number of carbon atoms in the alkyl group may be specified using the above notation, for example, when there are from 1 to 8 carbon atoms the term "C₁₋₈-alkyl" may be used. Examples of alkyl groups include methyl (Me, —CH₃), ethyl (Et, —CH₂CH₃), 1-propyl (n-Pr, n-propyl, —CH₂CH₂CH₃), 2-propyl (i-Pr, i-propyl, —CH(CH₃)₂), and 1-butyl (n-Bu, n-butyl, —CH₂CH₂CH₂CH₃).

[0039] An "alkylene," "alkenylene," or "alkynylene" group is an alkyl, alkenyl, or alkynyl group that is positioned between and serves to connect two other chemical groups. Thus, "C₁₋₆alkylene" means a linear saturated divalent hydrocarbon radical of one to six carbon atoms or a branched saturated divalent hydrocarbon radical of three to six carbon atoms. Example alkylene groups include methylene (—CH₂—), 1,1-ethylene (—CH(CH₃)—), 1,2-ethylene (—CH₂CH₂—), 1,1-propylene (—CH(CH₂CH₃)—), and 2,2-propylene (—C(CH₃)₂—).

[0040] The term "alkenyl" as used herein refers to a linear or branched-chain monovalent hydrocarbon radical with at least one site of unsaturation, i.e., a carbon-carbon double bond. The alkenyl radical may be optionally substituted, and includes radicals having "cis" and "trans" orientations, or alternatively, "E" and "Z" orientations. The number of carbon atoms in the alkenyl group may be specified using the above notation, for example, when there are from 2 to 8 carbon atoms the term "C₂₋₈-alkenyl" may be used. Example alkenyl groups include, but are not limited to, ethenyl (—CH=CH₂), and prop-1-enyl (—CH=CHCH₃).

[0041] In the chemical structures drawn herein, the presence of "⌵" or "⌵" denotes a point of attachment or a radical for example, a radical as discussed in relation to various functional groups.

[0042] The term "comprising" is used in the present description and claims, it does not exclude other elements or steps. Where an indefinite or definite article is used when referring to a singular noun e.g. "a" or "an", "the", this includes a plural of that noun unless something else is specifically stated.

[0043] The term "about" as used herein when referring to a measurable value such as a parameter, an amount, a temporal duration, and the like, is meant to encompass variations of +/−10% or less, preferably +/−5% or less, more preferably +/−1% or less, and still more preferably +/−0.1% or less of and from the specified value, insofar such variations are appropriate to perform in the disclosed invention. It is to be understood that the value to which the modifier 'about' refers is itself also specifically, and preferably, disclosed.

[0044] "Plant" as used herein, means an entire plant or a part thereof, including fresh fruit, vegetables and seeds. The plant or plant part may be a live plant or part thereof. Also, the term "plant" as used herein encompasses whole plants, ancestors and progeny of the plants and plant parts, including seeds, shoots, stems, leaves, roots (including tubers), flowers, and tissues and organs, wherein each of the afore-

mentioned comprise the gene/nucleic acid of interest. The term "plant" also encompasses plant cells, suspension cultures, callus tissue, embryos, meristematic regions, gametophytes, sporophytes, pollen and microspores, again wherein each of the aforementioned comprises the gene/nucleic acid of interest.

[0045] "Crop" as used herein means a plant species or variety that is grown to be harvested as food, livestock fodder, fuel raw material, or for any other economic purpose. As a non-limiting example, said crops can be maize, cereals, such as wheat, rye, barley and oats, sorghum, rice, sugar beet and fodder beet, fruit, such as pome fruit (e.g. apples and pears), citrus fruit (e.g. oranges, lemons, limes, grapefruit, or mandarins), stone fruit (e.g. peaches, nectarines or plums), nuts (e.g. almonds or walnuts), soft fruit (e.g. cherries, strawberries, blackberries or raspberries), the plantain family or grapevines, leguminous crops, such as beans, lentils, peas and soya, oil crops, such as sunflower, safflower, rapeseed, canola, castor or olives, cucurbits, such as cucumbers, melons or pumpkins, fibre plants, such as cotton, flax or hemp, fuel crops, such as sugarcane, miscanthus or switchgrass, vegetables, such as potatoes, tomatoes, peppers, lettuce, spinach, onions, carrots, egg-plants, asparagus or cabbage, ornamentals, such as flowers (e.g. petunias, pelargoniums, roses, tulips, lilies, or chrysanthemums), shrubs, broadleaved trees (e.g. poplars or willows) and evergreens (e.g. conifers), grasses, such as lawn, turf or forage grass or other useful plants, such as coffee, tea, tobacco, hops, pepper, rubber or latex plants.

[0046] A "pest", as used here, is an organism that is harmful to plants, animals, humans or human concerns, and includes, but is not limited to crop pests such as insects (as later defined), household pests or insects, such as cockroaches, ants, etc., and disease vectors, such as malaria mosquitoes.

[0047] "Pest infection" or "pest disease" as used herein refers to any inflammatory condition, disease or disorder in a living organism, such as a plant, animal or human, which is caused by a pest.

[0048] "Active substance", "active ingredient" or "active principle", as used interchangeably herein, means any biological, biochemical or chemical element and its derivatives, fragments or compounds based thereon, including microorganisms, having general or specific action against harmful organisms on a subject, and in particular on insect pests of plants, parts of plants or on plant products, as they occur naturally or by manufacture, including any impurity inevitably resulting from the manufacturing process.

[0049] The terms "effective amount" and "effective dose", as used herein, mean the amount needed to achieve the desired result or results.

[0050] An "insect", as used here, is used in the broad popular sense and includes all species of the superphylum Panarthropoda (classification Systema Naturae, Brands, S. (comp.) 1989-2005. Systema Naturae 2000. Amsterdam, The Netherlands, [http://sn2000.taxonomy.nl/]), including the phyla Arthropoda, Tardigrada and Onychophora; it includes all the different phases of the life cycle, such as, but not limited to eggs, larvae, nymphs, pupae and adults. Suitable insect pests are defined elsewhere herein.

[0051] An "insecticidal compound" as used herein refers to compounds having biological activity on insects (as defined above), including but not limited to compounds capable of killing the insect, larvaecides, insect growth

regulators, behaviour modifying compounds, attractants, repellents, pheromones, kairomones, allomones and entomopathogenic fungi, viruses and proteins. The insecticide exerts its biological activity preferably by the contact of the compound with the insect, without the need of being ingested by the insect. It includes not only compounds or compound formulations that are ready to use, but also precursors in an inactive form, which may be activated by outside factors. Possibly, the insecticide may be combined with materials used in conjunction, such as synergists or safeners, flavour or odour compositions. Preferably, said compound is comprised in a carrier as defined above. "Comprised in a carrier" as used herein means bound on or contained in by means such as but not limited to embedding, encapsulation and adsorption.


[0052] Throughout this specification, including the claims which follow, unless the context requires otherwise, the word "comprise" and "include", and variations such as "comprises", "comprising", and "including" will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

[0053] It must be noted that, as used in the specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. Ranges may be expressed herein as from "about" one particular value, and/or to "about" another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by the use of the antecedent "about," it will be understood that the particular value forms another embodiment. The term "about" in relation to a numerical value is optional and means for example $\pm 10\%$.

COMPOUNDS OF THE INVENTION

[0054] Particular compounds of the invention include, for example, compounds of the formula (I), wherein, unless otherwise stated, each of R^1 , Y^1 , Y^2 , R^2 and any associated substituent group has any of the meanings defined hereinbefore or in any of paragraphs (1) to (12) hereinafter:—


[0055] (1) R^1 is selected from hydrogen, acyl or fatty acyl, or phosphate or sulphate, or R^1 is a pyroglutamate group of the formula:



[0056] wherein acyl or fatty acyl are optionally substituted by a sugar moiety, phosphate or sulphate.

[0057] (2) R^1 is selected from hydrogen, formyl, acetyl (Ac), propanoyl, butanoyl, palmitoyl [palm], butyryl, cerotoyl, decanoyl, docosenoyl, dodecanoyl, eleostearoyl, heptanoyl, hexanoyl, icosanoyl, icosenoyl, lignoceroyl, linoleoyl, lipoyl, myristoleoyl, nonanoyl, octadecanoyl, ocatanoyl, palmitoleoyl, stearoyl, undecanoyl, and valeryl.


[0058] (3) R^1 is selected from hydrogen, acetyl or palmitoyl, or R^1 is a pyroglutamate group of the formula:



[0059] (4) R^1 is selected from hydrogen or palmitoyl.

[0060] (5) R^1 is hydrogen.

[0061] (6) R^1 is a pyroglutamate group of the formula:



[0062] (7) Y^1 is absent or a peptide comprising 1 or 2 amino acid residues.

[0063] (8) Y^1 is absent.

[0064] (9) Y^2 is absent or a peptide comprising 1 or 2 amino acid residues.

[0065] (10) Y^2 is absent.

[0066] (11) R^2 is NH_2 , $NR^{2a}H$, $NR^{2a}R^{2b}$ or OH wherein each of R^{2a} and R^{2b} if present is independently C_{1-6} -alkyl (e.g. methyl, ethyl, propyl, butyl, pentyl or hexyl).

[0067] (12) R^2 is NH_2 .

[0068] Suitably, R^1 is as defined in any one of paragraphs (1) to (6) above. More suitably, R^1 is as defined in any one of paragraphs (3) to (6) above. Most suitably, R^1 is as defined in paragraph (4) or (5) above.

[0069] Suitably, Y^1 or Y^2 is as defined in any one of paragraphs (7) to (10) above. Most suitably, Y^1 or Y^2 is as defined in paragraphs (8) or (10) above.

[0070] Suitably, R^2 is as defined in paragraph (11) or (12) above. Most suitably, R^2 is as defined in paragraph (12) above.

[0071] In a preferred embodiment, R^1 is hydrogen, Y^1 and Y^2 are absent, and R^2 is NH_2 .

[0072] In a preferred embodiment, R^1 is hydrogen, Y^2 is absent, Y^1 is one or two amino acids, and R^2 is NH_2 .

[0073] In a preferred embodiment, R^1 is hydrogen, Y^1 is absent, Y^2 is one or two amino acids, and R^2 is NH_2 .

FURTHER EXPLANATION OF COMPOUNDS OF THE INVENTION


R^1 and R^2

[0074] The terminal groups present at the N- and C-termini of the peptide backbone are designated R^1 and R^2 respectively. Thus, R^1 is bonded to the nitrogen atom of the N-terminal amino group of Y^1 and R^2 is bonded to the C-terminal carbonyl carbon atom of Y^2 .


[0075] The compounds of the invention may include further functionalisation, suitably at the N or C terminus. Suitably, the compounds may be functionalised to increase cuticle permeability or to increase stability. Suitably, the compound may be functionalised with an aromatic, aliphatic or lipophilic group. Suitably therefore, R^1 may be an aromatic, heteroaromatic, aliphatic or lipophilic group.

[0076] In certain embodiments, the compound may be functionalised with a lipophilic group such as a fatty acyl group. Fatty acyl groups include but are not limited to

palmitoyl, butyryl, cerotoyl, decanoyl, docosenoyl, dodecanoyl, eleostearoyl, heptanoyl, hexanoyl, icosanoyl, icosenoyl, lignoceroyl, linoleoyl, lipoyl, myristoleoyl, nonanoyl, octadecanoyl, ocatanoyl, palmitoleoyl, stearoyl, undecanoyl, and valeryl. Suitably therefore the R^1 group may be palmitoyl ([Palm]), i.e.:




[0077] In one embodiment, the compound may be functionalised with an acyl group. An “acyl” group is a group of the formula $R^{3a}-C(O)-$ group wherein R^{3a} is a C_{1-6} alkyl, for example formyl, acetyl (Ac), propanoyl, butanoyl, or wherein R^{3a} is benzoyl. Suitably, the R^{3a} group may be $R^{1b}-C(O)-$, such as acetyl (Ac), i.e.:



[0078] In certain embodiments, the R^1 group may be substituted with a sugar moiety.

[0079] In certain embodiments, one or more amino acid residues in peptides Y and Z may be naturally modified with a sugar moiety, i.e. the amino acid residue may be a “sugar



modified analogue”. For example, a sugar may be part of a Ser or Thr side chain modification (glycosylation) or a Lys or Arg side chain modification (glycation).

[0080] The sugar moieties discussed herein may be a monosaccharide or disaccharide. Examples of monosaccharides include glucose, 6-deoxyglucose, mannose, galactose, glucosamine, galactosamine, N-acetylglucosamine, N-acetylgalactosamine, glucuronic acid, allose, altrose, gulose, idose, fucose, talose, ribose, deoxyribose, arabinose, xylose, lyxose, ribulose, xylulose, fructose, psicose, sorbose or tagatose. Examples of disaccharides include sucrose, lactulose, allolactose, maltose, isomaltose, isomaltulose, trehalose, cellobiose, kojibiose, nigerose, sophorose, laminaribiose, gentiobiose, thiomaltose, mannobiose or their N-, C- or S-interglycosidic derivative. Most suitably, the sugar moiety is selected from glucosamine or galactosamine. The sugar can be present as an N-terminus modification or as part of Ser sidechain modification. In certain embodiments, the R^1 group may be substituted with a phosphate or a sulphate.

[0081] In certain embodiments, one or more amino acid residues in peptides Y and Z may be naturally modified with

a phosphate or sulphate. For example the phosphate or sulphate may be part of a side chain modification i.e. the amino acid may be phosphorylated or sulphated.

[0082] R^1 is selected from hydrogen (which may not be noted in the specific peptide sequence, or may instead be designated “H-” or “Hy-”), C_{1-4} alkyl (e.g. methyl, ethyl, propyl, butyl), $-N(R^{1a})-C(=N^+(R^{1b})(R^{1c}))NR^{1d}R^{1e}$, or $-C(=N^+(R^{1b})(R^{1c}))NR^{1d}R^{1e}$; wherein each of R^{1a} , R^{1b} , R^{1c} , R^{1d} and R^{1e} is independently selected from hydrogen or C_{1-4} alkyl (e.g. methyl, ethyl, propyl, butyl), preferably hydrogen or methyl.


[0083] In some embodiments, if R^1 is $-C(=N^+(R^{1b})(R^{1c}))NR^{1d}R^{1e}$; each of R^{1a} , R^{1b} , R^{1c} , R^{1d} and R^{1e} are methyl, i.e. R^1 is $-C(=N^+Me_2)NMe_2$.

[0084] When $R^1=“H”$ (or “Hy”), it typically indicates a free primary amino group at the N-terminus. The other hydrogen atom of the N-terminal amino group is typically invariant, regardless of the nature of R^1 . Exceptionally, when the residue at the N-terminus is N-methylated, R^1 may still be indicated as H even though the N-terminal residue has a secondary amine group. Thus an N-methylated leucine residue at the N-terminus may be indicated as $R^1-[n-me-L]$ - where R^1 is H. However, it could also be shown as simply R^1-L - where R^1 is methyl and the other hydrogen atom is not shown.

[0085] In some embodiments, an N-terminal glutamine (Gln or Q) or N-terminal glutamic acid (Glu or E) residue may undergo a conversion to form the pyroglutamate terminal group, [pyr]:

An N-terminal glutamine (Q) or glutamic acid (E) residue may undergo such a conversion in biological systems via reaction with certain enzymes. The conversion may also be achieved synthetically.

[0086] The Y^1 peptide, if present, or the Z peptide may therefore also include an N-terminal [pyr] group:



An N-terminal [pyr] group may be formed as a result of cyclisation of an N-terminal glutamine (Gln or Q) or N-terminal glutamic acid (Glu or E) residue to form the pyroglutamate terminal group. Since this conversion may occur under certain biological conditions, it is considered to be a naturally occurring modification of an N-terminal glutamine (Gln or Q) or N-terminal glutamic acid (Glu or E) residue.

[0087] Suitably, R^2 is NH_2 , $NR^{2a}H$, $NR^{2a}R^{2b}$, or OR^{2a} , wherein each of R^{2a} and R^{2b} are as defined herein. Preferably R^2 is NH_2 .

[0088] A particular example of a pyrokinin peptide according to the invention includes:

SB-P-80 [Hy]-NADEDQQQSVDFTPRL-[NH₂] (SEQ
ID NO: 2)

[0089] or a salt or solvate thereof.

[0090] In some embodiments, the compounds of the invention may be in the form of a salt or solvate (e.g. a hydrate).

[0091] The compounds of the invention may be provided in combination with one or more additional active insecticides, such as those described herein.

Y^1 and Y^2

[0092] The peptide may comprise one or two further peptide groups located between the terminal R^1 and R^2 groups at the N and C terminus respectively. Suitably these groups are the Y^1 and Y^2 group. If present, these groups are located on either side of the Z peptide group.

[0093] In one embodiment, the Y^1 and Y^2 groups are absent.

[0094] In one embodiment, the compound comprises a Y^1 group only. In one embodiment, the compound comprises a Y^2 group only.

[0095] In one embodiment, the compound comprises a Y^1 and Y^2 group.

[0096] Suitably the Y groups when present comprise one or two amino acids. Suitably the Y groups may comprise any amino acid, suitably selected from: M, R, G, and K.

[0097] In one embodiment, the Y groups are selected from: GR, RG, KR, RK, K, R, M, and G. In one embodiment, the Y groups are selected from: RG, KR, and K.

[0098] In one embodiment, the compound comprises a Y^2 group only, wherein the Y^2 group is selected from RG, KR, and K.

Modified Amino Acid or Non-Natural Amino Acid Analogue

[0099] In preferred embodiments, the insecticidal compound does not comprise any modified amino acid or non-natural amino acid analogues.

[0100] In particular, preferably peptide Y^1 , if present does not comprise any a modified amino acid or non-natural amino acid analogues, and peptide Y^2 if present does not comprise any modified amino acid or non-natural amino acid analogues. Thus, in certain embodiments, the peptides Y^1 and Y^2 comprise only unmodified amino acid residues, suitably only natural amino acid residues.

[0101] In particular, peptide Z does not comprise any modified amino acid or non-natural amino acid analogues. Thus, in certain embodiments, the peptide Z comprises only unmodified amino acid residues, suitably only natural amino acid residues.

Activity

[0102] Suitably, the compounds of the invention have activity against insects. Suitably the activity is against hemipteran insects, dipteran insects, lepidopteran insects, blattodean insects and/or coleopteran insects.

[0103] In a preferred embodiment, the compounds of the invention have activity against lepidopteran insects. The compounds of the invention may therefore find particular use against lepidopteran insects, and find particular application in the associated uses and methods described herein. Particularly preferred lepidopteran insects are moths.

[0104] The compounds of the invention typically increase insect mortality, for example when contacted topically to a suitable insect, or ingested by a suitable insect. Thus, the compounds of the invention described herein (and compositions containing them) may be regarded as insecticides, and may be referred to as “insect control agents”.

[0105] Suitably the compounds of the invention thereby indirectly increase plant health, including increasing plant growth, yields, and the like, due to preventing insect infestations of plants and reducing insect load on plants. Thus, the compounds of the invention described herein (and compositions containing them) may also be regarded as “plant protection agents”.

[0106] Without wishing to be bound by theory, any or all of the effects described may be mediated by agonist activity at the pyrokinin receptor of the target insects. The pyrokinin receptors believed to be present in target insects include the GPCRs pyrokinin receptor 1 and pyrokinin receptor 2, PK1-R and PK2-R. Suitably the compounds of the invention bind to one or more pyrokinin receptors of target insects. Suitably the compounds of the invention have agonist activity when bound to a pyrokinin receptor of a target insect. Suitably the compounds of the invention bind to pyrokinin receptor 1 and/or pyrokinin receptor 2. Suitably the compounds of the invention are agonists of pyrokinin receptor 1 and/or pyrokinin receptor 2.

Insect Control Agent

[0107] The term “insect control agent” refers to agents used to increase insect mortality (i.e. as insecticides). Any compound or composition thereof of the invention may be considered as an insect control agent, and may be used as an insect control agent. Suitably therefore the invention provides the use of a compound or composition of the invention as an insect control agent, for example against diptera insects, hemiptera insects, coleoptera insects, blattodea insects, and/or lepidoptera insects. Thus an insect control agent may be administered to increase mortality of a given insect or insect population.

[0108] An increase in mortality used herein is intended to refer to an increase in the percentage of dead insects, as compared to the percentage of dead insects of an otherwise identical insect population which have not been exposed to the insect control agent of the invention.

[0109] Suitably, insect mortality may be calculated as number of dead insects/total number of insects per treated area. Suitably the treated area may be a well of a plate, or may be one or more leaves, or an entire plant. Suitably insect mortality may be measured by performing a lepidoptera artificial diet assay as described herein in the examples.

[0110] An insect control agent may be used to reduce the size of an insect population, or inhibit growth of an insect population or inhibit feeding of an insect population (e.g. as compared to an otherwise identical insect population not exposed to the agent).

[0111] An insect control composition is a composition comprising an insect control agent i.e. a compound of the invention as described.

Plant Protection Agent

[0112] The term “plant protection agent” refers to agents used to protect a plant or plant part against insects, e.g. against infestation or colonisation, or being used as a food source by such insects (e.g. by the draining of sap). Any compound or composition thereof of the invention may be considered as a plant protection agent, and may be used as a plant protection agent. Suitably therefore the invention provides the use of a compound or composition of the invention as a plant protection agent, for example to protect a plant against diptera insects, hemiptera insects, coleoptera insects, blattodea insects, and/or lepidoptera insects.

[0113] Infestation or colonisation may be by larvae (or nymphs), by adult insects, or may be by being used as a host or repository for eggs. The terms “infestation” and “colonisation” should not be construed as requiring the presence of the insects to be deleterious to the plant, however.

[0114] A plant protection agent may be applied inter alia for reducing insect load on a plant or plant part, for inhibiting or reducing infestation of a plant by insects, for inhibiting (e.g. reducing the rate of) an increase in insect load on a plant or plant part, or for maintaining a plant in an insect-free state, as compared to an otherwise identical plant having an insect population not exposed to the agent. Thus, the plant protection agent may be applied to a plant or plant part which already carries hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects, or to a plant or plant part which is free or substantially free of hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects.

[0115] A plant protection composition is a composition comprising a plant protection agent i.e. a compound of the invention as described.

Plants

[0116] By ‘plant or plant part’, or ‘plant or part thereof’ referred to herein it is meant any part of a plant including but not limited to; the leaf, stem, root, flower, bud, bulb, and seed.

[0117] Suitable plants or parts thereof which may be protected by the compounds or compositions thereof of the invention, or by the agents of the present invention include crops and plants of agricultural, horticultural, or economic significance. Suitable plants may include any of the following or parts thereof: *Musa textilis*, *Medicago sativa*, *Prunus dulcis*, *Pimpinella anisum*, *Malus sylvestris*, *Prunus armeniaca*, *Areca catechu*, *Arracacia xanthorrhiza*, *Maranta arundinacea*, *Cynara scolymus*, *Helianthus tuberosus*, *Asparagus officinalis*, *Persea americana*, *Pennisetum americanum*, *Vigna subterranean*, *Musa paradisiaca*, *Hordeum vulgare*, *Phaseolus vulgaris*, *Phaseolus vigna* spp., *Beta vulgaris*, *Citrus bergamia*, *Rubus* spp., *Piper nigrum*, *Acacia mearnsii*, *Vaccinium* spp., *Bertholletia excelsa*, *Artocarpus altilis*, *Vicia faba*, *Brassica oleracea botrytis*, *Sorghum bicolor*, *Brassica oleracea gemmifera*, *Fagopyrum esculentum*, *Brassica oleracea capitata*, *Brassica rapa*, *Brassica* spp., *Theobroma cacao*, *Cucumis melo*, *Carum carvi*, *Elettaria cardamomum*, *Cynara cardunculus*, *Cerantonia siliqua*, *Daucus carota*, *Anacardium occidentale*, *Manihot esculenta*, *Ricinus communis*, *Brassica oleracea botrytis*, *Apium graveolens*, *Sechium edule*, *Prunus* spp., *Castanea sativa*, *Cicer arietinum*, *Cichorium intybus*, *Cichorium intybus*, *Capsicum* spp., *Cinnamomum verum*,

Cymbopogon nardus, *Citrus medica*, *Citrus reticulata*, *Trifolium* spp., *Syzygium aromaticum*, *Cocos nucifera*, *Colocasia* spp., *Xanthosoma* spp., *Coffee* spp., *Cola* spp., *Brassica napus*, *Zea mays*, *Valerianella locusta*, *Gossypium* spp., *Vigna unguiculata*, *Vaccinium* spp., *Lepidium sativum*, *Cucumis sativus*, *Ribes* spp., *Annona reticulata*, *Colocasia esculenta*, *Phoenix dactylifera*, *Moringa oleifera*, *Phaseolus* spp., *Allium sativum*, *Allium cepa*, *Pisum sativum*, *Triticum durum*, *Xanthosoma* spp., *Colocasia* spp., *Solanum melongena*, *Cichorium endivia*, *Lygeum spartum*, *Foeniculum vulgare*, *Trigonella foenumgraecum*, *Ficus carica*, *Corylus avellane*, *Furcraea macrophylla*, *Linum usitatissimum*, *Phormium tenax*, *Pelargonium* spp.; *Geranium* spp., *Zingiber officinalis*, *Langenaria* spp.; *Cucurbita* spp., *Cicer arietinum*, *Citrus paradise*, *Vitis vinifera*, *Lygeum spartum*, *Dactylis glomerata*, *Arachis hypogaea*, *Psidium guajava*, *Corylus avellane*, *Cannabis sativa*, *Crotalaria juncea*, *Agave fourcroydes*, *Lawsonia inermis*, *Humulus lupulus*, *Armoracia rusticana*, *Indigofera tinctoria*, *Jasminum* spp., *Corchorus* spp., *Brassica oleracea acephala*, *Ceiba pentandra*, *Hibiscus cannabinus*, *Brassica oleracea gongyloides*, *Lavandula* spp., *Allium ampeloprasum*, *Citrus limon*, *Cymbopogon citratus*, *Lens culinaris*, *Lependeza* spp., *Lactuca sativa*, *Glycyrrhiza glabra*, *Citrus aurantifolia*, *Citrus limetta*, *Linum usitatissimum*, *Litchi chinensis*, *Eriobotrya japonica*, *Lupinus* spp., *Macadamia* spp., *Myristica fragrans*, *Agave atrovirens*, *Citrus reticulata*, *Mangifera indica*, *Manihot esculenta*, *Secale cereal*, *Mespilus germanica*, *Cucumis melo*, *Penicum miliaceum*, *Eleusine coracana*, *Setaria italica*, *Echinochloa crusgalli*, *Eleusine coracana*, *Mentha* spp., *Morus* spp., *Morus alba*, *Agaricus* spp.; *Pleurotus* spp., *Volvariella*, *Brassica nigra*, *Sinapis alba*, *Prunus persica*, *Phormium tenax*, *Guizotia abyssinica*, *Myristica fragrans*, *Avena* spp., *Elaeis guineensis*, *Abelmoschus esculentus*, *Olea europea*, *Papaver somniferum*, *Citrus sinensis*, *Citrus aurantium*, *Dactylis glomerata*, *Metroxylon* spp., *Borassus flabellifer*, *Carica papaya*, *Pastinaca sativa*, *Pyrus communis*, *Pisum sativum*, *Carya illinoensis*, *Capsicum annuum*, *Diospyros kaki*, *Diospyros virginiana*, *Cajanus cajan*, *Ananas comosus*, *Pistacia* spp., *Prunus domestica*, *Punica granatum*, *Citrus grandis*, *Solanum tuberosum*, *Ipomoea batatas*, *Cucurbita* spp., *Chrysanthemum cinerariaefolium*, *Aspidosperma* spp., *Cydonia oblonga*, *Cinchona* spp., *Chenopodium quinoa*, *Raphanus sativus* (including *Cochlearia armoracia*), *Boehmeria nivea*, *Agrostis* spp., *Boehmeria nivea*, *Rheum* spp., *Oryza sativa*; *Oryza glaberrima*, *Rose* spp., *Hevea brasiliensis*, *Secale cereal*, *Lolium* spp., *Carthamus tinctorius*, *Metroxylon* spp., *Onobrychis viciifolia*, *Valerianella locusta*, *Tragopogon porrifolius*, *Achras sapota*, *Citrus reticulata*, *Brassica ileracea capitata*, *Scorzonera hispanica*, *Sesamum indicum*, *Butyrospermum paradoxum*, *Agave sisilana*, *Citrus aurantifolia*, *Glycine max*, *Triticum spelta*, *Spinacia oleracea*, *Secale cereal*, *Cucurbita* spp., *Fragaria* spp., *Sorghum bicolor* Sudanense, *Saccharum officinarum*, *Helianthus annuus*, *Crotalaria juncea*, *Citrus limetta*, *Ipomoea batatas*, *Citrus reticulata*, *Xanthosoma sagittifolium*, *Manihot esculenta*, *Colocasia esculenta*, *Camellia sinensis*, *Eragrostis abyssinica*, *Phleum pratense*, *Nicotiana tabacum*, *Lycopersicon esculentum*, *Lotus* spp., *Aleurites* spp., *Brassica rapa*, *Urena lobata*, *Vanilla planifolia*, *Vicia sativa*, *Juglans* spp., *Citrullus lanatus*, *Acacia mearnsii*, *Triticum* spp., *Hordeum* spp., *Dioscorea* spp., and *Ilex paraguariensis*.

[0118] Suitably, the plant or part thereof which may be protected by the compounds, compositions, or agents of the present invention is selected from a plant or part thereof which suffers from hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insect infestations, or which attracts hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects. Suitably, the plant or part thereof which suffers from hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insect infestations, or which attracts hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects is any of those listed above.

[0119] Suitably, the plant or part thereof which suffers from hemipteran insect infestations, or which attracts hemipteran insects, is any of those listed above.

[0120] Suitably, the plant or part thereof which suffers from dipteran insect infestations, or which attracts dipteran insects, is any of those listed above.

[0121] Suitably, the plant or part thereof which suffers from lepidopteran insect infestations, or which attracts lepidopteran insects, is any of those listed above.

[0122] Suitably, the plant or part thereof which suffers from coleopteran insect infestations, or which attracts coleopteran insects, is any of those listed above.

[0123] Suitably, the plant or part thereof which suffers from blattodean insect infestations, or which attracts blattodean insects, is any of those listed above.

[0124] In one embodiment, the plant or part thereof is selected from a plant or part thereof which suffers from or attracts hemipteran insect infestations, for example: cereal crops such as wheat (*Triticum* spp.), oats (*Avena* spp.), rye (*Secale* spp.), barley (*Hordeum* spp.), rice (*Oryza* spp.) and corn (*Zea* spp.); fruit and vegetable crops including apples (*Malus* spp.); pears (*Pyrus* spp.); strawberry (*Fragaria* spp.), blueberry (*Vaccinium* spp.), blackberry (*Rubus* spp.), raspberry (*Rubus* spp.), citrus (*Citrus* spp.), olive (*Olea* spp.), durian (*Durio* spp.), longan (*Dimocarpus* spp.), litchi (*L. chinensis*), persimmon (*Diospyros* spp.); beans and peas (including but not limited to *Phaseolus*, *Vigna*, *Pisum*, *Lens*, *Glycine*, *Cicer*, *Cajanus*, *Arachis* spp), sugar beet (*Beta vulgaris*), sugar cane (*Saccharum* spp.), lettuce (*Lactuca* spp.), brassicas (*Brassica* spp.) including oil seed rape, alliums (*Allium* spp.), tomato (*Solanum* spp.), pepper (*Capsicum* spp.), asparagus (*A. officinalis*), melon, squash, pumpkins (*Cucumis* spp.), and tubers (potato)(*Solanum* spp.), or a part thereof.

[0125] In one embodiment, the plant or part thereof is selected from a plant or part thereof which suffers from or attracts aphid insect infestations, suitably *M. persicae* insect infestations, including Solanaceae, Cruciferae, and Leguminosae for example: cereal crops such as wheat (*Triticum* spp) including winter wheat *Triticum aestivum* L); fruit and vegetable crops including peach (*Prunus* spp.), strawberry (*Fragaria* spp.), blueberry (*Vaccinium* spp.), blackberry (*Rubus* spp.), raspberry (*Rubus* spp.), brassicas (*Brassica* spp.) such as oil seed rape, lettuce (*Lactuca* spp.), tomato (*Solanum* spp.), pepper (*Capsicum* spp.), beans and peas (including but not limited to *Vigna*, *Pisum* spp), melon, squash, pumpkins (*Cucumis* spp.), citrus (*Citrus* spp.), and tubers (potato)(*Solanum* spp.), or a part thereof. In one embodiment, the plant is a vegetable crop, suitably a brassica spp.

[0126] In one embodiment, the plant or part thereof is selected from a plant or part thereof which suffers from or attracts dipteran insect infestations, for example: cereals (*Triticum* spp.); oats (*Avena* spp); rye (*Secale* spp.); barley

(*Hordeum* spp.), rice (*Oryza* spp.) and corn (*Zea* spp.); beans and peas (including but not limited to *Phaseolus*, *Vigna*, *Pisum*, *Lens*, *Glycine*, *Cicer*, *Cajanus*, *Arachis* spp); fruit crops including apples (*Malus* spp), pears (*Pyrus* spp.), strawberry (*Fragaria* spp.), blueberry (*Vaccinium* spp.), blackberry (*Rubus* spp.), raspberry (*Rubus* spp.), cherry, plum, apricot, peach, nectarine (*Prunus* spp.), blackcurrant, redcurrant, whitecurrant, gooseberry (*Ribes* spp.), kiwi fruit (*Actinidia* spp), papaya (*Carica* spp.), avocado (*Persea* spp.), mango (*Mangifera indica* L), longan (*Dimocarpus* spp.), litchi (*L. chinensis*), grapes (*Vitis* spp.), fig (*Ficus* spp.), passionfruit (*Passiflora* spp.), Asian pears (*Pyrus* spp), citrus (*Citrus* spp.), and olive (*Olea* spp.); vegetable crops including alliums (*Allium* spp.), aubergine, tomato (*Solanum* spp.) and peppers (*Capsicum* spp.), lettuce (*Lactuca* spp.), brassicas (*Brassica* spp.) and courgette, melon, squash, pumpkins (*Cucumis* spp.); Apiaceae root crops including carrot (*Daucus* spp.), parsnip (*Pastinaca* spp.), or a part thereof.

[0127] In one embodiment, the plant or part thereof is selected from a plant of part thereof which suffers from or attracts lepidoptera insect infestations, for example: cereal crops such as wheat (*Triticum* spp.), oats (*Avena* spp), rye (*Secale* spp.), barley (*Hordeum* spp.), rice (*Oryza* spp.) and corn (*Zea* spp.); fruit and vegetable crops including apples (*Malus* spp); pears (*Pyrus* spp); tree nuts (including for example almonds (*P. amygdalus*), pistachio (*Pistacia vera*), walnuts (Juglandaceae), hazlenuts (*Corylus*)); avocado, including *Persea americana* (Lauraceae), blueberry (*Vaccinium* spp.), citrus (*Citrus* spp.), olive (*Olea* spp.), durian (*Durio* spp.), longan (*Dimocarpus* spp.), litchi (*L. chinensis*), persimmon (*Diospyros* spp.); beans and peas (including but not limited to *Phaseolus*, *Vigna*, *Pisum*, *Lens*, *Glycine*, *Cicer*, *Cajanus*, *Arachis* spp), sugar beet (*Beta vulgaris*), sugar cane (*Saccharum* spp.), lettuce (*Lactuca* spp.), brassicas (*Brassica* spp.) including oil seed rape, alliums (*Allium* spp.), tomato (*Solanum* spp.), pepper (*Capsicum* spp.), asparagus (*A. officinalis*), melon, squash, pumpkins (*Cucumis* spp.), and tubers (potato)(*Solanum* spp.), or a part thereof.

Insects

[0128] The compounds, compositions, and agents of the invention suitably have activity against insects as described above.

[0129] The compounds, compositions, and agents of the invention may be effective against any insects. Such as Arthropoda, in particular from the class of the arachnids, for example *Acarus* spp., *Aceria sheldoni*, *Aculops* spp., *Aculus* spp., *Amblyomma* spp., *Amphitetranychus viennensis*, *Argas* spp., *Boophilus* spp., *Brevipalpus* spp., *Bryobia praetiosa*, *Centruroides* spp., *Choriotptes* spp., *Dermanyssus gallinae*, *Dermatophagoides pteronyssius*, *Dermatophagoides farinae*, *Dermacentor* spp., *Eotetranychus* spp., *Epirimerus pyri*, *Eutetranychus* spp., *Eriophyes* spp., *Halotydeus destructor*, *Hemitarsonemus* spp., *Hyalomma* spp., *Ixodes* spp., *Latrodectus* spp., *Loxosceles* spp., *Metatetranychus* spp., *Nuphessa* spp., *Oligonychus* spp., *Ornithodoros* spp., *Ornithonyssus* spp., *Panonychus* spp., *Phyllocoptruta oleivora*, *Polyphagotarsonemus latus*, *Psoroptes* spp., *Rhipicephalus* spp., *Rhizoglyphus* spp., *Sarcoptes* spp., *Scorpio maurus*, *Stenotarsonemus* spp., *Tarsonemus* spp., *Tetranychus* spp., *Vaejovis* spp., *Vasates lycopersici*.

[0130] Still other examples are from the order of the Anoplura (Phthiraptera), for example, *Damalinea* spp., *Haematopinus* spp., *Linognathus* spp., *Pediculus* spp., *Ptirus pubis*, *Trichodectes* spp.

[0131] Still other examples are from the order of the Chilopoda, for example, *Geophilus* spp., *Scutigera* spp.

[0132] Still other examples are from the order of the Coleoptera, for example, *Acalymma vittatum*, *Acanthoscelides obtectus*, *Adoretus* spp., *Agelastica alni*, *Agriotes* spp., *Alphitobius diaperinus*, *Amphimallon solstitialis*, *Anobium punctatum*, *Anoplophora* spp., *Anthonomus* spp., *Anthrenus* spp., *Apion* spp., *Apogonia* spp., *Atomaria* spp., *Attagenus* spp., *Bruchidius obtectus*, *Bruchus* spp., *Cassida* spp., *Ceratomya trifurcata*, *Ceutorhynchus* spp., *Chaetocnema* spp., *Cleonus mendicus*, *Conoderus* spp., *Cosmopolites* spp., *Costelytra zealandica*, *Ctenicera* spp., *Curculio* spp., *Cryptorhynchus lapathi*, *Cylindrocopturus* spp., *Dermestes* spp., *Diabrotica* spp., *Dichocrocis* spp., *Diloboderus* spp., *Epilachna* spp., *Epitrix* spp., *Faustinus* spp., *Gibbium psyllioides*, *Hellula undalis*, *Heteronychia arator*, *Heteronyx* spp., *Hylamorphia elegans*, *Hylotrupes bajulus*, *Hypera postica*, *Hypothenemus* spp., *Lachnosterna consanguinea*, *Lema* spp., *Leptinotarsa decemlineata*, *Leucoptera* spp., *Lissorhynchus oryzophilus*, *Lixus* spp., *Luperodes* spp., *Lycus* spp., *Megascelis* spp., *Melanotus* spp., *Meligethes aeneus*, *Melolontha* spp., *Migdolus* spp., *Monochamus* spp., *Naupactus xanthographus*, *Niptus hololeucus*, *Oryctes rhinoceros*, *Oryzaephilus surinamensis*, *Oryzaphagus oryzae*, *Otiorynchus* spp., *Oxyctonia jucunda*, *Phaedon cochleariae*, *Phyllorhiza* spp., *Phyllotreta* spp., *Popillia japonica*, *Premnotrypes* spp., *Prostephanus truncatus*, *Psylliodes* spp., *Ptinus* spp., *Rhizobius ventralis*, *Rhizopertha dominica*, *Sitophilus* spp., *Sphenophorus* spp., *Stegobium paniceum*, *Sternechus* spp., *Symphyletes* spp., *Tanymecus* spp., *Tenebrio molitor*, *Tribolium* spp., *Trogoderma* spp., *Tychius* spp., *Xylotrechus* spp., *Zabrus* spp. Suitably compounds, compositions, and agents of the invention have activity against Coleoptera species selected from, for example: Bruchid beetle (*Bruchus rufimanus*), cabbage seed weevil (*Ceutorhynchus obstrictus*), cabbage stem flea beetle (*Psylliodes chrysocephalus*), cabbage stem weevil (*Ceutorhynchus pallidactylus*), Chafer grubs (*Melolontha melolontha*), Colorado potato beetle (*Leptinotarsa decemlineata*), pea and bean weevils (*Sitona lineatus*), pollen beetle (*Meligethes* spp.), pygmy beetle (*Atomaria linearis*), rape winter stem weevil (*Ceutorhynchus picipitarsis*), and wireworms (*Agriotes* spp.).

[0133] Still other examples are from the order of the Collembola, for example, *Onychiurus armatus*.

[0134] Still other examples are from the order of the Diplopoda, for example, *Blaniulus guttulatus*.

[0135] Still other examples are from the order of the Diptera, for example, *Aedes* spp., *Agromyza* spp., *Anastrepha* spp., *Anopheles* spp., *Asphondylia* spp., *Bactrocera* spp., *Bibio hortulanus*, *Calliphora erythrocephala*, *Ceratitis capitata*, *Chironomus* spp., *Chrysomya* spp., *Chrysops* spp., *Cochliomyia* spp., *Contarinia* spp., *Cordylobia anthropophaga*, *Culex* spp., *Culicoides* spp., *Culiseta* spp., *Cuterebra* spp., *Dacus oleae*, *Dasyneura* spp., *Delia* spp., *Dermatobia hominis*, *Drosophila* spp., *Echinocneme* spp., *Fannia* spp., *Gasterophilus* spp., *Glossina* spp., *Haematopota* spp., *Hydrellia* spp., *Hylemyia* spp., *Hyppobosca* spp., *Hypoderma* spp., *Liriomyza* spp., *Lucilia* spp., *Lutzomyia* spp., *Mansonella* spp., *Musca* spp., *Nezara* spp., *Oestrus* spp.,

Oscinella frit, *Pegomyia* spp., *Phlebotomus* spp., *Phorbia* spp., *Phormia* spp., *Prodiptosis* spp., *Psila rosae*, *Rhagoletis* spp., *Sarcophaga* spp., *Simulium* spp., *Stomoxys* spp., *Tabanus* spp., *Tannia* spp., *Tetanops* spp., *Tipula* spp. Still other examples are from the order of the Heteroptera, for example, *Anasa tristis*, *Antestiopsis* spp., *Boisea* spp., *Blissus* spp., *Calocoris* spp., *Campylomma livida*, *Cavelerius* spp., *Cimex* spp., *Collaria* spp., *Creontiades dilutus*, *Dasyneura piperis*, *Dichelops furcatus*, *Diconocoris hewetti*, *Dysdercus* spp., *Euschistus* spp., *Eurygaster* spp., *Heliopeltis* spp., *Horcias nobilellus*, *Leptocoris* spp., *Leptoglossus phyllopus*, *Lygus* spp., *Macropes excavatus*, *Miridae*, *Monalion atratum*, *Nezara* spp., *Oebalus* spp., *Pentomidae*, *Piesma quadrata*, *Piezodorus* spp., *Psallus* spp., *Pseudocysta perseae*, *Rhodnius* spp., *Sahlbergella singularis*, *Scaptocoris castanea*, *Scotinophora* spp., *Stephanitis nashi*, *Tibraca* spp., *Triatoma* spp. Still other examples are from the order of the Homoptera, for example, *Acyrtosiphon* spp., *Acrogonia* spp., *Aeneolamia* spp., *Agonosceles* spp., *Aleurodes* spp., *Aleurolobus barodensis*, *Aleurothrixus* spp., *Amrasca* spp., *Anuraphis cardui*, *Aonidiella* spp., *Aphanostigma pin*, *Aphis* spp., *Arboridia apicalis*, *Aspidiella* spp., *Aspidiotus* spp., *Atanus* spp., *Aulacorthum solani*, *Bemisia* spp., *Brachycaudus helichrysi*, *Brachycolus* spp., *Brevicoryne brassicae*, *Calligypona marginata*, *Carneiocephala fulgida*, *Ceratovacuna lanigera*, *Cercopidae*, *Ceroplastes* spp., *Chaetosiphon fragaefolii*, *Chionaspis tegalensis*, *Chlorita onukii*, *Chromaphis juglandicola*, *Chrysomphalus ficus*, *Cicadulina mbila*, *Cocomytilus halli*, *Coccus* spp., *Cryptomyzus ribis*, *Dalbulus* spp., *Dialeurodes* spp., *Diaphorina* spp., *Diaspis* spp., *Drosicha* spp., *Dysaphis* spp., *Dysmicoccus* spp., *Empoasca* spp., *Eriosoma* spp., *Erythroneura* spp., *Euscelis bilobatus*, *Ferrisia* spp., *Geococcus coffeae*, *Hieroglyphus* spp., *Homalodisca coagulata*, *Hyalopteris arundinis*, *Icerya* spp., *Idiocerus* spp., *Idioscopus* spp., *Laodelphax striatellus*, *Lecanium* spp., *Lepidosaphes* spp., *Lipaphis erysimi*, *Macrosiphum* spp., *Mahanarva* spp., *Melanaphis sacchari*, *Metcalfiella* spp., *Metopolophium dirhodum*, *Monellia costalis*, *Monelliopsis pecanis*, *Myzus* spp., *Nasonovia ribisnigri*, *Nephotettix* spp., *Nilaparvata lugens*, *Oncometopia* spp., *Orthezia praelonga*, *Parabemisia myricae*, *Paratrioza* spp., *Parlatoria* spp., *Pemphigus* spp., *Peregrinus maidis*, *Phenacoccus* spp., *Phloeomyzus passerinii*, *Phorodon humuli*, *Phylloxera* spp., *Pinnaspis aspidistrae*, *Planococcus* spp., *Protopulvinaria pyriformis*, *Pseudaulacaspis pentagona*, *Pseudococcus* spp., *Psylla* spp., *Pteromalus* spp., *Pyrilla* spp., *Quadrastipidiotus* spp., *Quesada gigas*, *Rastrococcus* spp., *Rhopalosiphum* spp., *Saissetia* spp., *Scaphoides titanus*, *Schizaphis graminum*, *Selenaspis articulatus*, *Sogata* spp., *Sogatella furcifera*, *Sogatodes* spp., *Stictocephala festina*, *Tenalaphara malayensis*, *Tinocallis caryaefoliae*, *Tomaspis* spp., *Toxoptera* spp., *Trialeurodes* spp., *Trioza* spp., *Typhlocyba* spp., *Unaspis* spp., *Viteus vitifolii*, *Zygina* spp.

[0136] Still other examples are from the order of the Hymenoptera, for example, *Acromyrmex* spp., *Athalia* spp., *Atta* spp., *Diprion* spp., *Hoplocampa* spp., *Lasius* spp., *Monomorium pharaonis*, *Solenopsis invicta*, *Tapinoma* spp., *Vespa* spp.

[0137] Still other examples are from the order of the Isopoda, for example, *Armadillidium vulgare*, *Oniscus asellus*, *Porcellio scaber*.

[0138] Still other examples are from the order of the Isoptera, for example, *Coptotermes* spp., *Cornitermes cumu-*

lans, *Cryptotermes* spp., *Incisitermes* spp., *Microtermes* *obesi*, *Odontotermes* spp., *Reticulitermes* spp.

[0139] Still other examples are from the order of the Lepidoptera, for example, *Acrionicta major*, *Adoxophyes* spp., *Aedia leucomelas*, *Agrotis* spp., *Alabama* spp., *Amyelois transitella*, *Anarsia* spp., *Anticarsia* spp., *Argyroplece* spp., *Barathra brassicae*, *Borbo cinnara*, *Bucculatrix thurberii*, *Bupalus piniarius*, *Busseola* spp., *Cacoecia* spp., *Caloptilia theivora*, *Capua reticulana*, *Carpocapsa pomonella*, *Carposina niponensis*, *Chematobia brumata*, *Chilo* spp., *Choristoneura* spp., *Clysia ambiguella*, *Cnaphalocerus* spp., *Cnephasia* spp., *Conopomorpha* spp., *Conotrachelus* spp., *Copitarsia* spp., *Cydia* spp., *Dalaca noctuides*, *Diaphania* spp., *Diatraea saccharalis*, *Earias* spp., *Ecdytoplopha aurantium*, *Elasmopalpus lignosellus*, *Eldana saccharina*, *Ephestia* spp., *Epinotia* spp., *Epiphyas postvittana*, *Etiella* spp., *Eulia* spp., *Eupoecilia ambiguella*, *Euproctis* spp., *Euxoa* spp., *Feltia* spp., *Galleria mellonella*, *Gracilaria* spp., *Grapholitha* spp., *Hedylepta* spp., *Helicoverpa* spp., *Heliothis* spp., *Hofmannophila pseudospretella*, *Homoeosoma* spp., *Homona* spp., *Hyponomeuta padella*, *Kakivoria flavofasciata*, *Laphygma* spp., *Laspeyresia molesta*, *Leucinodes orbonalis*, *Leucoptera* spp., *Lithocolletis* spp., *Lithophane antennata*, *Lobesia* spp., *Loxagrotis albicosta*, *Lymantria* spp., *Lyonetia* spp., *Malacosoma neustria*, *Maruca testulalis*, *Mamestra brassicae*, *Moda* spp., *Mythimna separata*, *Nymphula* spp., *Oiketicus* spp., *Oria* spp., *Orthaga* spp., *Ostrinia* spp., *Oulema oryzae*, *Panolis flammea*, *Parnara* spp., *Pectinophora* spp., *Perileucoptera* spp., *Phthorimaea* spp., *Phyllocnistis citrella*, *Phyllonorycter* spp., *Pieris* spp., *Platynota stultana*, *Plodia interpunctella*, *Plusia* spp., *Plutella xylostella*, *Prays* spp., *Prodenia* spp., *Protoparce* spp., *Pseudaletia* spp., *Pseudoplusia includens*, *Pyrausta nubilalis*, *Rachiplusia nu*, *Schoenobius* spp., *Scirpophaga* spp., *Scotia segetum*, *Sesamia* spp., *Sparganothis* spp., *Spodoptera* spp., *Stathmopoda* spp., *Stomopteryx subsecivella*, *Synanthedon* spp., *Tecia solanivora*, *Thermesia gemmatilis*, *Tinea pellionella*, *Tineola bisselliella*, *Tortrix* spp., *Trichophaga tapetzella*, *Trichoplusia* spp., *Tuta absoluta*, *Virachola* spp.

[0140] Still other examples are from the order of the Orthoptera, for example, *Acheta domesticus*, *Dichroplus* spp., *Gryllotalpa* spp., *Leucophaea maderae*, *Locusta* spp., *Melanoplus* spp., *Pulex irritans*, *Schistocerca gregaria*.

[0141] Still other examples are from the order of the Blattodea, for example, *Blatta orientalis*, *Blattella germanica*, *Periplaneta americana*, *Periplaneta* spp., *Supella longipalpa*, termites of the infraorder Isoptera such as those of the family Termitidae.

[0142] Still other examples are from the order of the Siphonaptera, for example, *Ceratophyllus* spp., *Ctenocephalides* spp., *Tunga penetrans*, *Xenopsylla cheopis*.

[0143] Still other examples are from the order of the Symphyla, for example, *Scutigera* spp.

[0144] Still other examples are from the order of the Thysanoptera, for example, *Anaphothrips obscurus*, *Baliothrips bififormis*, *Drepanothrips reuteri*, *Enneothrips flavens*, *Frankliniella* spp., *Heliothrips* spp., *Hercinothrips femoralis*, *Rhipiphorothrips cruentatus*, *Scirtothrips* spp., *Taeniothrips cardamoni*, *Thrips* spp. Still other examples are from the order of the Zygentoma (=Thysanura), for example, *Lepisma saccharina*, *Thermobia domestica*.

[0145] In one preferred embodiment, the compounds, compositions, and agents of the invention suitably have activity against insects of the order hemiptera, diptera, coleoptera, blattodea and/or lepidoptera. In one preferred embodiment, the compounds, compositions, and agents of the invention suitably have activity against insects of the order Lepidoptera. In one preferred embodiment, the compounds, compositions, and agents of the invention suitably have activity against moths. In one embodiment, the compounds, compositions, and agents of the invention suitably have activity against the moth species *Plutella xylostella*. In one embodiment, the compounds, compositions, and agents of the invention suitably have activity against the moth species *Spodoptera frugiperda*.

Hemipteran Insects

[0146] The compounds, compositions and agents of the invention suitably have activity against insects of the order Hemiptera, which comprises groups including aphids, planthoppers, leafhoppers, stink bugs, shield bugs and cicadas. Suitably, the compounds, compositions and agents of the invention suitably have activity against aphids.

[0147] Hemipterans are defined by distinctive mouthparts in the form of a “beak”, comprising modified mandibles and maxillae which form a “stylet”, sheathed within a modified labium.

[0148] Many insects within these groups have endogenous neuropeptides with sequence homology to the peptides described herein, suggesting that the compounds of the present invention may have activity against those insects.

[0149] The insects may belong to the sub-order Sternorrhyncha, e.g. to the super-family of Aphidoidea (aphid superfamily), Aleyrodoidea (whiteflies), Coccoidea (scale insects), Phylloxeroidea (including Phylloxeridae or “phylloxerans”, and Adelgidae or woolly conifer aphids) or Psylloidea (jumping plant lice etc.).

[0150] Thus, the insects may be aphids, i.e. members of the aphid superfamily (Aphidoidea). Aphids (Hemiptera: Aphididae) are one of the most significant groups of agricultural pests and are vectors in the transmission of approximately 50% of all insect transmitted plant viruses. Within that superfamily, the aphids may be part of the family Aphididae, which contains sub-families Aiceoninae, Anoeciinae, Aphidinae, Baltichaitophorinae, Calaphidinae, Chaitophorinae, Drepanosiphinae, Eriosomatinae, Greenideinae, Hormaphidinae, Israelaphidinae, Lachninae, Lizeriinae, Macropodaphidinae, Mindarinae, Neophyllaphidinae, Phloeomyzinae, Phyllaphidinae, Pterastheniinae, Saltusaphidinae, Spicaphidinae, Taiwanaphidinae, Tamaliinae and Thelaxinae.

[0151] The aphids may, for example, be of the genus *Acyrtosiphon* (e.g. *Acyrtosiphon pisum*), *Aphis* (e.g. *Aphis gossypii*, *Aphis glycines*, *Aphis fabae*), *Diuraphis* (e.g. *Diuraphis noxia*) *Macrosiphum* (e.g. *Macrosiphum rosae*, *Macrosiphum euphorbiae*), *Myzus* (e.g. *Myzus persicae*), *Rhopalosiphum* (e.g. *Rhopalosiphum padi*), or *Sitobion* (e.g. *Sitobion avenae*).

[0152] *Myzus persicae* (peach potato aphid) is the most economically important aphid crop pest worldwide, with a global distribution and host range encompassing more than 400 species in 40 different plant families. For example, it is a major pest of agricultural crops including fruit and potatoes, and act as a vector for viruses.

[0153] *Macrosiphum rosae*, (rose aphid) is an important horticultural pest, especially of cultivated species of *Rosa*, and is a vector in the transmission of 12 plant viruses including the strawberry mild yellow edge virus.

[0154] *Aphis gossypii* (cotton or melon aphid) is a pest of Curcibitae and cotton.

[0155] Other than aphids, the insects may, for example, be of the Adelgidae family, e.g. of the genus *Adelges* (e.g. *Adelges tsugae*).

[0156] The insects may be of the Aleyrodidae family, e.g. of the genus *Bemisia* (e.g. *Bemisia tabaci*) or *Trialeurodes* (e.g. *Trialeurodes vaporariorum*).

[0157] The insects may be of the Psylloidea family, e.g. of the genus *Pachypsylla* (e.g. *Pachypsylla venusta*).

[0158] As examples of hemipteran insects outside the sub-order Sternorrhyncha, the insects may be of the Cimicidae family, e.g. of the genus *Cimex* (bed bugs), e.g. *Cimex lectularius*.

[0159] The insects may be of the Cicadellidae family, e.g. of the genus *Cuerna* (e.g. *Cuerna arida*), *Graminella* (e.g. *Graminella nigrifrons*) or *Homalodisca* (e.g. *Homalodisca vitripennis*). Also included in the Cicadellidae family are *Ammrasca biguttula*. The insects may be part of the Delphacidae family, e.g. of the genus *Nilaparvata* (e.g. *Nilaparvata lugens*) or *Sogatella* (e.g. *Sogatella furcifera*). For example, *Nilaparvata lugens* (brown planthopper) is a pest of rice crops, especially in Asia.

[0160] The insects may be of the Liviidae family, e.g. of the genus *Diaphorina* (e.g. *Diaphorina citri*).

[0161] The insects may be part of the Miridae family, e.g. of the genus *Pseudatomoscelis* (e.g. *Pseudatomoscelis seriatus*), *Lygus* (e.g. *Lygus hesperus*) or *Tupiocoris* (e.g. *Tupiocoris notatus*). For example, *Pseudatomoscelis seriatus* (cotton fleahopper) is a pest of cotton.

[0162] The insects may be of the Pentatomidae family, e.g. of the genus *Acrosternum* (e.g. *Acrosternum hilare*), *Banasa* (e.g. *Banasa dimiata*), *Euschistus* (e.g. *Euschistus servus*, *Euschistus heroes*), *Halyomorpha* (e.g. *Halyomorpha halys*), *Murgantia* (e.g. *Murgantia histrionica*), *Nezara* (e.g. *Nezara viridula*), *Plautia* (e.g. *Plautia stali*), or *Podisus* (e.g. *Podisus maculiventris*). For example, *Acrosternum hilare* (green stink bug) is a significant pest of cotton. *Euschistus servus* (brown stink bug) is a pest of many agricultural crops including seeds, grains, nuts and fruits, especially in the southern USA. *Nezara viridula* is a pest of grain and soybean crops, especially in Brazil.

[0163] The insects may be of the Pyrrhocoridae family, e.g. of the genus *Pyrrhocoris* (e.g. *Pyrrhocoris apterus*).

[0164] The insects may be of the Reduviidae family, e.g. of the genus *Rhodnius* (e.g. *Rhodnius prolixus*), or *Triatoma* (e.g. *Triatoma infestans*). *Rhodnius prolixus* is a vector of human disease (Chagas disease).

[0165] The insects may be of the Triozidae family, e.g. of the genus *Acanthocasuarina* (e.g. *Acanthocasuarina muel-lerianae*).

[0166] In one embodiment, the insect may be selected from the following species: *H. halys*, *E. heroes*, *A. hilare*, *A. gossypii*, *E. servus*, *M. persicae*, *N. viridula*, *N. lugens*, *P. seriatus*, and *R. prolixus*.

[0167] In one embodiment, the insect is of the species *M. persicae*.

Dipteran Insects

[0168] The compounds, compositions and agents of the invention may have activity against insects of the order Diptera.

[0169] In particular, they may have activity against insects of the family Drosophilidae, such as fruit flies, including those of genus *Drosophila*, such as *Drosophila suzukii*. They may also have activity against insects of the family Tephritidae, including those of the genera *Anastrepha* (*Anastrepha* spp.); *Bactrocera* (*Bactrocera* spp.); *Ceratitis* (*Ceratitis* spp.); *Dacus* (*Dacus* spp.); *Rhagoletis* (*Rhagoletis* spp.); *Tephritis* (*Tephritis* spp.).

[0170] The families Drosophilidae and Tephritidae together are commonly referred to as fruit flies.

[0171] The compounds, compositions, and agents may also have activity against other important dipteran pests, such as flies of the family Chloropidae (chloropid flies) and those of the genera:

[0172] *Phytomyza* (e.g. *Phytomyza angelicastris*);

[0173] *Melani* (e.g. *Melani agromyza*);

[0174] *Antherigona* (e.g. *Antherigona* spp);

[0175] *Delia* (e.g. *Delia radicum*);

[0176] *Contarinia* (e.g. *Contarinia sorghicola*);

[0177] For more detail on these, and other examples, see *Developing the Arsenal Against Pest and Vector Dipterans: Inputs of Transgenic and Paratransgenic Biotechnologies*, Ogaugwu and Durvasula, *IntechOpen*, 2017; DOI: 10.5772/66440

Lepidopteran Insects

[0178] The compounds, compositions and agents of the invention may have activity against insects of the order Lepidoptera.

[0179] In particular, they may have activity against insects of the genera: *Heliothis*, *Plutella*, *Spodoptera*, and *Cydia*. Suitably the compounds, compositions and agents of the invention may have activity against the species: *Heliothis peltigera*, *H. virescens*, *Spodoptera* spp., and *Cydia pomonella* (Codling Moth), Larvae of *Heliothis* spp., including *peltigera* and *virescens* *Spodoptera littoralis* (which represent a large variety of Heliothinae and *Spodoptera* moth species and are world-wide agricultural pests), and *Plutella xylostella* (diamondback moth, most important world-wide pest of Brassicas).

[0180] In one embodiment, the compounds, compositions and agents of the invention may have activity against *Plutella xylostella*. In one embodiment, the compounds, compositions and agents of the invention may have activity against *Spodoptera frugiperda*.

Coleopteran Insects

[0181] The compounds, compositions and agents of the invention may have activity against insects of the order Coleoptera.

[0182] In particular, they may have activity against insects of the genera: *Bruchus*, *Ceutorhynchus*, *Psylliodes*, *Leptinotarsa*, *Sitona*, *Meligethes*, *Atomaria*, and *Agriotes*. Suitably the compounds, compositions and agents of the invention may have activity against the species: Bruchid beetle (*Bruchus rufimanus*), cabbage seed weevil (*Ceutorhynchus obstrictus*), cabbage stem flea beetle (*Psylliodes chrysocephalus*), cabbage stem weevil (*Ceutorhynchus pallidactylus*), Chafer grubs (*Melolontha melolontha*), Colorado

potato beetle (*Leptinotarsa decemlineata*), pea and bean weevils (*Sitona lineatus*), pollen beetle (*Meligethes* spp.), pygmy beetle (*Atomaria linearis*), rape winter stem weevil (*Ceutorhynchus picipitarsis*), and wireworms (*Agriotes* spp.)

Blattodean Insects

[0183] The compounds, compositions and agents of the invention may have activity against insects of the order Blattodea.

[0184] In particular, they may have activity against insects of the infraorder Isoptera. Suitably the compounds, compositions and agents of the invention may have activity against the species of the family Termitidae.

[0185] Suitably the compounds, compositions and agents of the invention may have activity against the species: *Blatta orientalis*, *Blattella germanica*, *Periplaneta* America, *Periplaneta* spp., *Supella longipalpa*.

Household Pests

[0186] The compounds, compositions, and agents of the invention may also have activity against domestic pests, such as cockroaches and termites (such as those of the family Termitidae). Of more than 3000 species, these may include the German cockroach (*Blattella germanica*), the Oriental cockroach (*Blatta orientalis*), the American cockroach (*Periplaneta americana*) and the brown banded cockroach (*Supella longipalpa*). Cockroaches are common domestic pests worldwide, and may carry various diseases. Control of these and other domestic pests (e.g. ants) is envisaged by treating surfaces where the insects run, but particularly with food baits containing compounds, compositions and agents of the invention, and with direct spray application of compounds, compositions and agents of the invention.

[0187] Therefore further aspects of the invention may include a bait comprising a compound or composition of the invention, suitably which may be a bait for domestic pests. Further aspects of the invention may include a method of controlling domestic pests, reducing domestic pest populations, inhibiting domestic pest populations, increasing domestic pest mortality, the method comprising treating a surface (e.g. wood) which is contacted by the domestic pest with a compound or composition of the invention, or contacting the domestic pest with a compound or composition of the invention. Further details of such methods are described hereinbelow. Optionally treating or contacting may comprise a suitable means of application as described elsewhere herein, such as by spraying. Suitably therefore a sprayable formulation comprising a compound or composition of the invention is also envisaged. Suitable formulations are described elsewhere herein.

Methods and Uses of the Invention

Methods of Increasing Insect Mortality

[0188] The invention provides a method of increasing insect mortality, comprising contacting an insect or insect population with a compound, composition or combinations as described herein. The insect or insect population may be hemipteran, dipteran, coleopteran, blattodean, and/or lepidopteran insects.

[0189] In one embodiment, there is provided a method of increasing dipteran insect mortality, comprising contacting a

dipteran insect or dipteran insect population with a compound or composition or combinations of the invention.

[0190] In one particular embodiment, there is provided a method of increasing *Drosophila suzukii* mortality, comprising contacting a *Drosophila suzukii* insect or insect population with a compound or composition or combinations of the invention.

[0191] In one embodiment, there is provided a method of increasing hemipteran insect mortality, comprising contacting a hemipteran insect or hemipteran insect population with a compound or composition or combinations as defined herein.

[0192] In one particular embodiment, there is provided a method of increasing aphid mortality, comprising contacting an aphid insect or insect population with a compound or composition or combinations as defined herein.

[0193] In one particular embodiment, there is provided a method of increasing *Aphis fabae*, *Aphis gossypii*, *Acyrtosiphon pisum*, *Myzus persicae*, *Amrasca biguttula* or *Rhopalosiphum padi* mortality, comprising contacting an *Aphis fabae*, *Aphis gossypii*, *Acyrtosiphon pisum*, *Myzus persicae*, *Amrasca biguttula* or *Rhopalosiphum padi* insect or insect population with a compound or composition or combinations as defined herein.

[0194] In one particular embodiment, there is provided a method of increasing *Myzus persicae* mortality, comprising contacting a *Myzus persicae* insect or insect population with a compound or composition or combinations as defined herein.

[0195] In one particular embodiment, there is provided a method of increasing *Rhopalosiphum padi* mortality, comprising contacting a *Rhopalosiphum padi* insect or insect population with a compound or composition or combinations as defined herein.

[0196] In one embodiment, there is provided a method of increasing lepidoptera insect mortality, comprising contacting a lepidopteran insect or lepidopteran insect population with a compound or composition or combinations as defined herein.

[0197] In one particular embodiment, there is provided a method of increasing *Plutella xylostella* mortality, comprising contacting a *Plutella xylostella* insect or insect population with a compound or composition or combinations as defined herein.

[0198] In one particular embodiment, there is provided a method of increasing *Spodoptera frugiperda* mortality, comprising contacting a *Spodoptera frugiperda* insect or insect population with a compound or composition or combinations as defined herein.

[0199] In one embodiment, there is provided a method of increasing coleopteran insect mortality, comprising contacting a coleopteran insect or coleopteran insect population with a compound or composition or combinations as defined herein.

[0200] In one embodiment, there is provided a method of increasing blattodean insect mortality, comprising contacting a blattodean insect or blattodean insect population with a compound or composition or combinations as defined herein.

Methods of Inhibiting or Reducing Infestation of a Plant or Site

[0201] The invention further provides a method of inhibiting infestation of a plant or plant part by insects, suitably

by hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects comprising contacting the plant or plant part with a compound, composition or combinations as described.

[0202] In one embodiment, there is provided a method of inhibiting infestation of a plant by dipteran insects comprising contacting the plant with a compound or composition or combinations of the invention.

[0203] In one particular embodiment, there is provided a method of inhibiting infestation of a plant by *Drosophila suzukii* comprising contacting the plant with a compound or composition or combinations of the invention.

[0204] In one embodiment, there is provided a method of inhibiting infestation of a plant by hemipteran insects comprising contacting the plant with a compound or composition or combinations as defined herein.

[0205] In one embodiment, there is provided a method of inhibiting infestation of a plant by aphids comprising contacting the plant with a compound or composition or combinations as defined herein.

[0206] In one embodiment, there is provided a method of inhibiting infestation of a plant by *Aphis fabae*, *Aphis gossypii*, *Acyrtosiphon pisum*, *Myzus persicae* or *Rhopalosiphum padi* comprising contacting the plant with a compound or composition or combinations as defined herein.

[0207] In one particular embodiment, there is provided a method of inhibiting infestation of a plant by *Rhopalosiphum padi* comprising contacting the plant with a compound or composition or combinations as defined herein.

[0208] In one embodiment, there is provided a method of inhibiting infestation of a plant by *Myzus persicae* comprising contacting the plant with a compound or composition or combinations as defined herein.

[0209] In one embodiment, there is provided a method of inhibiting infestation of a plant by *Halyomorpha halys* comprising contacting the plant with a compound or composition or combinations as defined herein.

[0210] In one embodiment, there is provided a method of inhibiting infestation of a plant by lepidopteran insects comprising contacting the plant with a compound or composition or combinations as defined herein.

[0211] In one embodiment, there is provided a method of inhibiting infestation of a plant by *Plutella xylostella* comprising contacting the plant with a compound or composition or combinations as defined herein.

[0212] In one embodiment, there is provided a method of inhibiting infestation of a plant by *Spodoptera frugiperda* comprising contacting the plant with a compound or composition or combinations as defined herein.

[0213] In one embodiment, there is provided a method of inhibiting infestation of a plant by coleopteran insects comprising contacting the plant with a compound or composition or combinations as defined herein.

[0214] In one embodiment, there is provided a method of inhibiting infestation of a plant by blattodean insects comprising contacting the plant with a compound or composition or combinations as defined herein.

[0215] The invention further provides a method of inhibiting infestation of a site on which plants are growing or intended to be grown by insects, suitably by hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects comprising contacting the site with a compound, composition, or combinations as described.

[0216] The method may be prophylactic. Thus, for example, the compound may be applied to the plant or plant part, or site, while the plant or part or field is free or substantially free of insects.

[0217] Suitably the site may be any agricultural site suitable for growing plants. Suitably the site may be any area or locale which may be suitable for growing plants, or in which plants are grown. Suitable sites may include farmland, brownfield sites, fields, greenhouses, conservatories, containers, aquaponics and hydroponics systems etc. In one embodiment, the site is a field.

[0218] Alternatively, the plant or plant part or site may already be colonised or infested by insects, suitably by hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects.

[0219] Thus, the invention further provides a method of reducing insect infestation of a plant or plant part, or of reducing insect load on a plant or plant part, the method comprising contacting the plant or plant part with a compound, composition, or combinations as described herein. The invention also provides a method of reducing insect infestation of a field, or of reducing an insect load in a field, the method comprising contacting the field with a compound, composition, or combinations as described herein. Suitably of hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects.

Methods of Decreasing Insect Feeding

[0220] The invention further provides a method of decreasing insect feeding, comprising contacting an insect or insect population with a compound, composition or combinations as described herein. Suitably decreasing insect feeding on a plant or plant part. The insect or insect population may be hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects.

Methods of Protecting a Plant from Insects

[0221] The invention further provides a method of protecting a plant or part thereof from insects, or from insect infestation, and specifically a method of protecting a plant or plant part against hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects or infestation thereof, wherein the method comprises the step of applying directly or indirectly to the plant or to a part of the plant, an insecticidal compound or composition, or combinations of the invention. Suitably applying indirectly may comprise applying the compound or composition or combinations of the invention to the field or locale in which the plant is growing, or in which it is intended to be grown.

[0222] The present invention also provides post-harvest treatment methods for protecting or treating a harvested plant or a harvested part of the plant against insects, or insect infestation, specifically against hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects or infestation thereof, the method comprising the step of applying directly or indirectly to the harvested plant or to a harvested part of the plant, an insecticidal compound or composition or combinations of the invention, under conditions effective to protect or treat the harvested plant or a harvested part of the plant against the insects. Suitably applying indirectly may comprise applying the compound or composition or combinations of the invention to the location in which the plant or a harvested part of the plant is stored, or in which it is intended to be stored.

[0223] Suitably the insect may be of the order hemiptera, diptera, coleoptera, blattodea and/or lepidoptera.

Methods of Contacting an Insect

[0224] In one embodiment, there is provided a method of contacting an insect with a compound, preferably an insect control agent, even more preferably an insecticide, said method comprising applying to or on sites frequented by the insect a compound or composition or combinations according to the invention.

[0225] The site, frequented by insects may be a natural habitat for insects, or a place regularly visited by insects. This site can be treated then with the compound or composition as described above: as a non-limiting example mosquito nets, impregnated with encapsulated insecticide, can be used as application method. Alternatively, said site is created by application of a visual lure or of an attractant for the insects. Visual lures are known to the person skilled in the art, and include but are not limited to light sources, coloured object and shapes or silhouettes that stand out of a contrasting background. As mentioned above, insect attractants include but are not limited to pheromones, kairomones and allomones. The attractant may be present in the composition or it may be applied separately from the compound or composition or combinations, to ensure that the insects are attracted to the site where the compound or composition or combinations is applied.

[0226] Suitably the insect may be of the order hemiptera, diptera, coleoptera, blattodea and/or lepidoptera. More suitably, the insect is of the order lepidoptera. Most suitably, the insect is a moth, such as *Plutella xylostella* or *Spodoptera frugiperda*.

Application of the Compounds or Compositions

[0227] The compound or composition or combinations of the invention may be contacted with the insect or insect population, suitably this may comprise applying the compound or composition or combinations directly to an insect or insect population. For example, it may be applied topically. Alternatively, the compound or composition or combinations may be applied indirectly. For example, it may be applied to a substrate, site, or locale likely to come into contact with an insect or insect population. The substrate may be a plant or plant part, especially for Hemiptera or Diptera or Coleoptera or Lepidoptera or Blattodea which represent pests of plants (whether crops or horticultural plants), or may be a field, locale, or area suitably in which the plants are grown. Suitably therefore the compound or composition or combinations may be applied to the plant or plant part. Suitably therefore the compound or composition or combinations may be applied to the site in which the plant is grown.

[0228] However, for insects which represent pests to humans, such as the Cimicidae family (e.g. bedbugs of the genus *Cimex*, such as *Cimex lectularius*) or the Reduviidae family (e.g. of the genus *Rhodnius* such as *Rhodnius prolixus*, or *Triatoma* such as *Triatoma infestans*) which can be vectors of human disease, the substrate may be a domestic surface or article, such as bedding, a mattress, or any other suitable domestic surface. The compound or composition may be applied to the substrate in a form suitable for ingestion by an insect.

[0229] Suitably contacting may comprise feeding or spraying, for example. Suitably feeding may be encouraged via bait attractants, which may be comprised in a composition of the invention, as explained below.

[0230] The methods may comprise contacting or applying directly or indirectly to the plant or to a part of the plant a compound or composition or combinations as disclosed herein, for example at an application rate higher than 5 g of the compound per hectare, such as but not limited to an application rate higher than 10 g of the compound per hectare, such as an application rate higher than 24 g of the compound per hectare, such as an application rate higher than 50 g of the compound per hectare, such as an application rate higher than 75 g of the compound per hectare, such as an application rate higher than 100 g of the compound per hectare, or in particular an application rate higher than 200 g of the compound per hectare. These methods may comprise applying directly or indirectly to the plant or to a part of the plant a compound, composition or combinations as disclosed herein, for example at an application rate between 5 g and 100 g of the compound composition or combinations per hectare, such as but not limited to an application rate of between 5 g and 200 g of the compound composition or combinations per hectare, in particular an application rate of between 5 g and 50 g of the compound composition or combinations per hectare, such as between 5 g and 30 g of the compound composition or combinations per hectare or between 10 g and 25 g per hectare.

[0231] The compounds or compositions or combinations as disclosed herein may be directly or indirectly contacted with/applied to the plant or to a part of the plant by spraying, atomizing, foaming, fogging, culturing in hydroculture, culturing in hydroponics, coating, submerging, injecting and/or encrusting, optionally post-harvest. Suitably contacting may comprise feeding or spraying, for example. In some embodiments, the contacting is by feeding. Contacting may comprise injecting a plant (e.g. a tree) or part thereof with the composition as defined herein, using systems and methods such as those described in WO2020/021041, WO2023/161802, WO2022/264053, WO2022/189386, WO2022/165248, WO2021/152093, WO2020/212612 and WO2020/021041, the entire contents of which being incorporated by reference.

[0232] Suitably the compound may be contacted with the insect or insect population, or plant or plant part, at any suitable concentration which is effective. Suitably the compositions and combinations may comprise such an effective concentration of a compound. Suitably the concentration of the compound is between 10⁻³ to 10⁻⁹ M, suitably between 10⁻⁴ to 10⁻⁶ M, suitably between 10⁻⁴ to 10⁻⁵ M.

Use as a Plant Protection Agent

[0233] The invention further provides the use of a compound, composition or combination as described herein as a plant protection agent, and specifically for protecting a plant or plant part against insects, suitably against hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects. Plant protection agents are described further hereinabove.

Use as an Insect Control Agent

[0234] The present invention provides the use of a compound or composition or combination thereof as described

herein as an insect control agent, specifically in methods of increasing mortality in insects, or a method of inhibiting infestation of a plant by insects.

[0235] The present invention also provides the use of a compound or composition or combination thereof as described herein as an insect control agent, specifically in methods of increasing hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insect mortality, or in a method of inhibiting or reducing infestation, or reducing insect load of a plant by insects, suitably by hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects.

[0236] The present invention also provides the use of a compound or composition or combination thereof as described herein as an insect control agent, specifically by having a biostatic effect on hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects, a biocidal effect on hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects, and/or a pesticidal effect hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insects.

[0237] “Biostatic (effect)” or “biostatic use”, as used herein, includes any effect or use of a compound or composition or combination as described herein (optionally comprised in a biostatic, biocidal, fungicidal or fungistatic composition as defined herein) for controlling, modulating or interfering with the harmful activity of a pest, such as a plant pest or a plant pathogen. Suitably the pest is of hemipteran, dipteran, coleopteran, blattodean and/or lepidopteran insect orders. Suitably including but not limited to inhibiting the growth or activity of the insect, altering the behaviour of the insect, and repelling or attracting the insect in plants, plant parts or in other agro-related settings, such as for example for household uses or in soil.

[0238] “Pesticidal activity” or “biocidal activity”, as used interchangeably herein, means killing the pest or severely disabling the pest. Suitably which may be the same as insecticidal activity wherein the pest is an insect. Suitably, the compound, composition or combination may be for use as an insect control agent wherein the insect encodes a pyrokinin peptide.

[0239] Suitably, the compound or composition or combination may be for use as an insect control agent wherein the insect is of the order diptera. Suitably, the compound may be for use as an insect control agent wherein the insect is of the genus *Drosophila*. Suitably, the compound may be for use as an insect control agent wherein the insect is *Drosophila suzukii*.

[0240] Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is of the order hemiptera. Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is an aphid. Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is of the genus *Myzus*. Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is *Myzus persicae*.

[0241] Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is of the order hemiptera. Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the

insect is of the genus *Halyomorpha*. Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is *Halyomorpha halys*.

[0242] Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is of the order lepidoptera. Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is of the genus *Plutella*. Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is *Plutella xylostella*.

[0243] Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is of the order lepidoptera. Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is of the genus *Spodoptera*. Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is *Spodoptera frugiperda*.

[0244] Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is of the order coleoptera.

[0245] Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is of the order blattodea. Suitably, the compound or composition or combination as defined herein may be for use as an insect control agent wherein the insect is a cockroach or a termite.

Methods of Controlling Insect Fecundity

[0246] The present invention also provides insecticidal compounds, compositions and combinations as described herein for controlling fecundity of insect species. Suitably, the insect is selected from hemipteran, dipteran, coleopteran or lepidopteran insects, such as those described anywhere herein. Suitably wherein in some embodiments the hemipteran insect is *Myzus persicae*. Suitably wherein in some embodiments the dipteran insect is *D. suzukii*. Suitably wherein in some embodiments the lepidopteran insect is *Plutella xylostella*. Suitably wherein in some embodiments the lepidopteran insect is *Spodoptera frugiperda*.

[0247] There is also provided a method of controlling fecundity of a hemipteran insect, the method comprising contacting a compound of the invention, or a composition of the invention to a substrate, e.g. a plant or soil. Suitably, the hemipteran insect is any of those disclosed herein, e.g., an aphid.

[0248] There is also provided a method of controlling fecundity of a dipteran insect, the method comprising contacting a compound of the invention, or a composition of the invention to a substrate, e.g. a plant or soil. Suitably, the dipteran insect is any of those disclosed herein.

[0249] There is also provided a method of controlling fecundity of a lepidopteran insect, the method comprising contacting a compound of the invention, or a composition of the invention to a substrate, e.g. a plant or soil. Suitably, the lepidopteran insect is any of those disclosed herein.

[0250] There is also provided a method of controlling fecundity of an coleopteran insect, the method comprising contacting a compound of the invention, or a composition of

the invention to a substrate, e.g. a plant or soil. Suitably, the coleopteran insect is any of those disclosed herein.

[0251] There is also provided a method of controlling fecundity of a blattodean insect, the method comprising contacting a compound of the invention, or a composition of the invention to a substrate, e.g. a plant or soil. Suitably, the blattodean insect is any of those disclosed herein.

[0252] There is also provided a method of controlling fecundity of a hemipteran insect, the method comprising contacting a combination as described herein, to a substrate, e.g. a plant or soil. Suitably, the hemipteran insect is any of those disclosed herein. Suitably, the combination comprises a compound of the invention, and a kinin peptide, or an analogue thereof.

[0253] There is also provided a method of controlling fecundity of a dipteran insect, the method comprising contacting a combination as described herein, to a substrate, e.g. a plant or soil. Suitably, the dipteran insect is any of those disclosed herein. Suitably, the combination comprises a compound of the invention, and a kinin peptide, or an analogue thereof.

[0254] There is also provided a method of controlling fecundity of a lepidopteran insect, the method comprising contacting a combination as described herein, or a composition of the invention to a substrate, e.g. a plant or soil. Suitably, the lepidopteran insect is any of those disclosed herein. Suitably, the combination comprises a compound of the invention, and a kinin peptide, or an analogue thereof.

[0255] There is also provided a method of controlling fecundity of a coleopteran insect, the method comprising contacting a combination as described herein, or a composition of the invention to a substrate, e.g. a plant or soil. Suitably, the coleopteran insect is any of those disclosed herein. Suitably, the combination comprises a compound of the invention, and a kinin peptide, or an analogue thereof.

[0256] There is also provided a method of controlling fecundity of a blattodean insect, the method comprising contacting a combination as described herein, or a composition of the invention to a substrate, e.g. a plant or soil. Suitably, the blattodean insect is any of those disclosed herein. Suitably, the combination comprises a compound of the invention, and a kinin peptide, or an analogue thereof.

[0257] There is also provided a method of controlling fecundity of a hemipteran insect or insect population, the method comprising contacting a compound of the invention, or a composition of the invention to a hemipteran insect or insect population. Suitably, the hemipteran insect is any of those disclosed herein.

[0258] There is also provided a method of controlling fecundity of a dipteran insect or insect population, the method comprising contacting a compound of the invention, or a composition of the invention to a dipteran insect or insect population. Suitably, the dipteran insect is any of those disclosed herein.

[0259] There is also provided a method of controlling fecundity of a lepidopteran insect or insect population, the method comprising contacting a compound of the invention, or a composition of the invention to a lepidopteran insect or insect population. Suitably, the lepidopteran insect is any of those disclosed herein.

[0260] There is also provided a method of controlling fecundity of a coleopteran insect or insect population, the method comprising contacting a compound of the invention,

or a composition of the invention to a coleopteran insect or insect population. Suitably, the coleopteran insect is any of those disclosed herein.

[0261] There is also provided a method of controlling fecundity of a blattodean insect or insect population, the method comprising contacting a compound of the invention, or a composition of the invention to a blattodean insect or insect population. Suitably, the blattodean insect is any of those disclosed herein.

[0262] There is also provided a method of controlling fecundity of a hemipteran insect or insect population, the method comprising contacting a combination as described herein, to a hemipteran insect or insect population. Suitably, the hemipteran insect is any of those disclosed herein. Suitably, the combination comprises a compound of the invention, and a kinin peptide, or an analogue thereof.

[0263] There is also provided a method of controlling fecundity of a dipteran insect or insect population, the method comprising contacting a combination as described herein, to a dipteran insect or insect population. Suitably, the dipteran insect is any of those disclosed herein. Suitably, the combination comprises a compound of the invention, and a kinin peptide, or an analogue thereof.

[0264] There is also provided a method of controlling fecundity of a lepidopteran insect or insect population, the method comprising contacting a combination as described herein, or a composition of the invention to a lepidopteran insect or insect population. Suitably, the lepidopteran insect is any of those disclosed herein. Suitably, the combination comprises a compound of the invention, and a kinin peptide, or an analogue thereof.

[0265] There is also provided a method of controlling fecundity of a coleopteran insect or insect population, the method comprising contacting a combination as described herein, or a composition of the invention to a coleopteran insect or insect population. Suitably, the coleopteran insect is any of those disclosed herein. Suitably, the combination comprises a compound of the invention, and a kinin peptide, or an analogue thereof.

[0266] There is also provided a method of controlling fecundity of a blattodean insect or insect population, the method comprising contacting a combination as described herein, or a composition of the invention to a blattodean insect or insect population. Suitably, the blattodean insect is any of those disclosed herein. Suitably, the combination comprises a compound of the invention, and a kinin peptide, or an analogue thereof.

[0267] In the context of the present invention, “controlling fecundity” refers to reducing, inhibiting, or eliminating the presence of an insect species during one or more of its growth stages. For example, the compositions of the present invention may be used for controlling the growth of mites at any stage such as egg, larva, nymph, and adult form.

Combinations with Further Insecticides

[0268] The compound of the invention may be used in combination with one or more further insecticides or insecticidal compounds, such as those as described herein. There is also provided a combination comprising a compound of the invention, or a salt or solvate thereof, in combination with one or more additional active insecticides, or insecticidal compounds.

[0269] The composition of the invention may further comprise one or more additional active insecticides, or active insecticidal compounds.

[0270] Suitably, the insecticide may be selected from an insect neuropeptide or analogues thereof such as a kinin peptide, an AKH peptide, a DH31 peptide, a DH44 peptide, a pyrokinin peptide or a CAPA peptide (e.g. a CAPA-1, CAPA-2 or CAPA-3 analogue). In a particular embodiment, the further insecticide is a kinin peptide, for example SB-P-69 ([Hy]-NFSPWG-[NH₂], SEQ ID NO:14). Additionally or alternatively, the insecticide may be selected from a chemical insecticide, such as: Pyrethroids (Permethrin, Cypermethrin, Deltamethrin); Organophosphates (Malathion, Chlorpyrifos, Diazinon); Neonicotinoids (Imidacloprid, Clothianidin, Thiamethoxam); Carbamates (Carbaryl, Methomyl, Propoxur); Botanical Insecticides (Pyrethrin (derived from chrysanthemum flowers), Rotenone (derived from certain plant roots)); Biopesticides (*Bacillus thuringiensis* (Bt) products, *Beauveria bassiana* (fungus), *Metarhizium anisopliae* (fungus)); Insect Growth Regulators (IGRs) (Methoprene, Pyriproxyfen); Fipronil; Spinosad; Avermectins (including abamectin and ivermectin); Chitin Synthesis Inhibitors (Diflubenzuron, Hexaflumuron); Piperonyl Butoxide; Flonicamid; Indoxacarb; and Sulfoxaflor.

[0271] Suitably said further insecticides or insecticidal compounds may be comprised in a composition of the invention.

[0272] The methods of the invention may further comprise contacting the insect population or plant or part thereof with a further insecticide agent, or insecticidal compound, such as those described herein. Suitably, the further insecticide or insecticidal compound is a kinin peptide.

[0273] Suitably any references herein to the compound of the invention or a composition thereof may equally refer to a combination of a compound of the invention with one or more insecticides or insecticidal compounds, or a composition containing such a combination.

[0274] Suitably, there is provided a combination comprising a compound of the invention, or a salt or solvate thereof, in combination with one or more of the peptides listed in the table below:

Family	Peptide	Sequence	SEQ ID NO.
Pyrokinin	SB-P-46	[Hy]-SPPYSPPFSPRL-[NH ₂]	6
	SB-P-47	[Pyr]-AIMARPQVPRL-[NH ₂]	3
	SB-P-48	Hy-EQNVQSNGEPAAYRVTPRL-[NH ₂]	15
	SB-P-49	[Hy]-SVPFKPRL-[NH ₂]	4
	SB-P-51	[Hy]-LRQLQSNGEPAAYRVTPRL-[NH ₂]	5
	SB-P-81	[Hy]-GGSMTFSPRL-[NH ₂]	7
Kinin	SB-P-70	[Hy]-KVKFSAWG-[NH ₂]	8
	SB-P-65	[Hy]-RQKTVFSSWG-[NH ₂]	9
	SB-P-66	[Hy]-PAFSSWG-[NH ₂]	10
	SB-P-69	[Hy]-NFSPWG-[NH ₂]	14
AKH	SB-P-86	[Pyr]-LTFTSSWGG-[NH ₂]	11
	SB-P-39	[Palm]-QLTFSPDW-[NH ₂]	12
	SB-P-41	Hy-QLTFSPDW-[NH ₂]	16
	SB-P-42	[Pyr]-LTTFSPDW-[NH ₂]	13

[0275] In one embodiment, the compound of the invention is utilised in combination with a further pyrokinin peptide, e.g.

Family	Peptide	Sequence	SEQ ID NO.
Pyrokinin	SB-P-46	[Hy]-SPPYSPPFSPRL-[NH ₂]	6
	SB-P-47	[Pyr]-AIMARPQVPRL-[NH ₂]	3
	SB-P-48	Hy-EQNVQSNGEPAAYRVTPRL-[NH ₂]	15
	SB-P-49	[Hy]-SVPFKPRL-[NH ₂]	4
	SB-P-51	[Hy]-LRQLQSNGEPAAYRVTPRL-[NH ₂]	5
	SB-P-81	[Hy]-GGSMTFSPRL-[NH ₂]	7

[0276] In one embodiment, the compound of the invention is utilised in combination with a kinin peptide, e.g.

Family	Peptide	Sequence	SEQ ID NO.
Kinin	SB-P-70	[Hy]-KVKFSAWG-[NH ₂]	8
	SB-P-65	[Hy]-RQKTVFSSWG-[NH ₂]	9
	SB-P-66	[Hy]-PAFSSWG-[NH ₂]	10
	SB-P-69	[Hy]-NFSPWG-[NH ₂]	14

[0277] In one embodiment, the compound of the invention is utilised in combination with a CAPA peptide (e.g. a CAPA2 peptide).

[0278] In one embodiment, the compound of the invention is utilised in combination with an AKH peptide, e.g.

Family	Peptide	Sequence	SEQ ID NO.
AKH	SB-P-86	[Pyr]-LTFTSSWGG-[NH ₂]	11
	SB-P-39	[Palm]-QLTFSPDW-[NH ₂]	12
	SB-P-41	Hy-QLTFSPDW-[NH ₂]	16
	SB-P-42	[Pyr]-LTTFSPDW-[NH ₂]	13

[0279] Suitably any of the above combinations of compounds may be comprised within a composition of the invention.

[0280] The choice of ancillary or additional insecticides will typically depend on the particular target species.

Beneficial Insect Species

[0281] The compounds and compositions and combinations of the invention may be substantially non-toxic to beneficial insect species, including species which prey on pests and pollinator species. Important pollinator species, such as insects of the superfamily Apoidea, including bees, such as the Apidae, e.g. those of the genus *Bombus*, such as *Bombus terrestris*. Important predatory species include Coccinellidae (lady bugs) such as *Adalia bipunctata*.

[0282] By substantially non-toxic, it is meant that the compounds and compositions and combinations of the invention do not cause death of the beneficial insect species (e.g. pollinator species), suitably that they do not cause premature death of the beneficial insect species (e.g. pollinator species). It is also meant that the compounds and compositions and combinations of the invention do not cause any detrimental side effects to the beneficial insect

species (e.g. pollinator species), for example they do not have a negative effect on feeding behaviour, or ability to move.

Compositions

[0283] The present inventors have provided compositions comprising at least one insecticidal compound or combination thereof of the invention, which can specifically bind to an insect. Importantly, through this interaction with a specific molecular structure of the insect, such as a receptor, suitably a neurological receptor, the compositions disclosed herein are capable of inhibiting, preventing or reducing one or more biological activities of the insect, such that the growth or fecundity of the insect is inhibited, prevented or reduced. In certain embodiments, the compositions as disclosed herein are capable of killing insects through the specific interaction of at least one insecticidal compound, which can specifically bind to an insect receptor, and which is comprised in the compositions.

[0284] Compositions of the invention, or for use in accordance with the invention, typically comprise a compound as described in combination with one or more ancillary component such as solvents, carriers, diluents, adjuvants, preservatives, dispersants, emulsifying agents, or synergists. Suitably the composition may be an agricultural composition, an insect control composition (e.g. insecticide composition), or a plant protection composition.

[0285] In any of these embodiments, the compound or combination of compounds of the invention may be provided as part of a composition, such as an agricultural composition, an insect control composition (e.g. insecticide composition) or a plant protection composition. Reference to application or use of a compound or combination herein should therefore be construed as encompassing application or use of a suitable composition thereof, unless the context demands otherwise.

[0286] The composition typically comprises a compound as described herein in admixture with one or more ancillary component such as solvents, carriers, diluents, adjuvants, preservatives, dispersants, emulsifying agents, or synergists.

[0287] The composition may further comprise a combination with one or more additional active insecticides as described herein.

[0288] The invention further provides a composition, e.g. an agricultural composition, an insect control composition or plant protection composition, comprising a compound of the invention, or a combination thereof, in admixture with one or more solvents, carriers, diluents, adjuvants, preservatives, dispersants, emulsifying agents, or synergists. The composition may be an aqueous composition. Further details of which are explained hereinbelow.

[0289] “Agricultural”, as used herein, means suitable for use in the agricultural or agrochemical industry, including horticulture, floriculture and home and garden uses, but also products intended for non-crop related uses such as public health/pest control operator uses to control undesirable insects and rodents, household uses, such as household fungicides and insecticides and agents, for protecting plants or parts of plants, crops, bulbs, tubers, fruits (e.g. from harmful organisms, diseases or pests); for controlling, preferably promoting or increasing, the growth of plants; and/or for promoting the yield of plants, crops or the parts of plants that are harvested (e.g. its fruits, flowers, seeds etc.). Examples of such substances will be clear to the skilled

person and preferably in the context of the present invention, include compounds that are active as insecticides (e.g. contact insecticides or systemic insecticides, including insecticides for household use). Other such agrochemicals may be pesticides, growth regulators, nutrients/fertilizers, repellants, defoliants etc.

[0290] “Agricultural use”, as used herein, not only includes the use of the insecticidal compounds of the invention, combinations thereof or compositions, and optionally the agrochemicals as defined above (for example, pesticides, growth regulators, nutrients/fertilizers, repellants, defoliants etc.), that are suitable and/or intended for use in field grown crops (e.g., agriculture), but also includes the use of compounds of the invention combinations thereof or compositions and said agrochemicals as defined above that are meant for use in greenhouse grown crops (e.g. horticulture/floriculture) or hydroponic culture systems and even the use of compounds combinations thereof or compositions of the invention and agrochemicals as defined above that are suitable and/or intended for non-crop uses such as uses in private gardens, household uses (for example, herbicides or insecticides for household use), or uses by pest control operators (for example, weed control etc.).

[0291] Suitably an insect control composition according to the invention is for controlling insect populations. Formulations of such compositions for controlling insect populations are known to the person skilled in the art and include, but are not limited to liquid emulsifiable concentrates, wettable powders, solutions, suspension concentrates, emulsions, suspoemulsions, granules and water dispersible granules (Mulqueen, 2003). Preferably, said insect control compositions according to the invention comprise an insecticidal compound of the invention, and optionally a combination of further insecticidal compounds. Suitably said insecticidal compound is comprised in a carrier as described hereinbelow.

[0292] Compositions of the invention may comprise a pesticide formulation or an agrochemical formulation. A “pesticide formulation” as used herein means any composition comprising a compound or combination of compounds intended for preventing, destroying, repelling, attracting or mitigating any pest. An “agrochemical formulation” as used herein means a composition for agricultural use, comprising a biologically active agent, optionally with one or more additives favoring optimal dispersion, atomization, distribution, retention and/or activity of agrochemicals. As a non-limiting example such additives are diluents, solvents, adjuvants, surfactants, wetting agents, spreading agents, oils, stickers, penetrants, buffering agents, acidifiers, defoaming agents or drift control agents.

[0293] A composition as used herein means a composition comprising at least one active substance, suitably an insecticidal compound of the invention, optionally with one or more additives favouring optimal dispersion, atomization, deposition, leaf wetting, distribution, retention and/or uptake of said active substance. It will become clear from the further description herein that a composition as used herein includes biological insect control agents or biological insecticides, and these terms will be interchangeably used in the present application. Accordingly, a composition as used herein includes compositions comprising at least one biological molecule as an active ingredient, substance or principle for controlling pests in plants or in other agro-related settings (such for example in soil). Suitably wherein the at

least one biological molecule comprises an insecticidal compound of the invention or a combination thereof. Suitably wherein the pest is an insect. As a non-limiting example, the additives in the compositions disclosed herein may include but are not limited to diluents, solvents, adjuvants, surfactants, wetting agents, spreading agents, oils, stickers, thickeners, penetrants, buffering agents, acidifiers, antisetling agents, anti-freeze agents, photo-protectors, defoaming agents, biocides and/or drift control agents.

[0294] Suitably, the compositions of the invention are aqueous compositions.

[0295] The compound content of the composition can vary within wide limits. The compound concentration of the composition is suitably an effective amount, and can be from 0.0000001 to 95% by weight of the compound, preferably between 0.0001 and 1% by weight. The terms “effective amount” and “effective dose”, as used herein, mean the amount needed to achieve the desired result or results.

[0296] In a specific embodiment the concentration of the compound of the invention contained in the composition may be at least 0.0001% by weight. In a specific embodiment the concentration of the compound of the invention contained in the composition may be up to 50% by weight. In a specific embodiment the concentration of the compound of the invention contained in the composition may be from 0.0001% to 50% by weight. In particular embodiments, the present invention provides compositions comprising at least one insecticidal compound of the invention, wherein the concentration of the at least one compound of the invention in the composition ranges from 0.001% to 50% by weight. In yet another specific embodiment the concentration of the at least one compound of the invention contained in the composition may be from 0.001% to 50% by weight. In yet another specific embodiment the concentration of the at least one compound of the invention contained in the composition may be from 0.01% to 50% by weight. In yet another specific embodiment the concentration of the at least one compound of the invention contained in the composition may be from 0.1% to 50% by weight.

[0297] The compositions of the invention, or for use in accordance with the invention, may comprise more than one compound of the invention in combination, and/or with other insecticidal compounds as described herein in combination. Therefore the compositions of the invention may comprise a first compound of the invention and a second compound of the invention, or a first compound of the invention and a second insecticidal compound, for example. Suitably the first and second compound may be any of those described herein, and may be present in the composition in any relative proportion.

[0298] The composition may be an aqueous composition, e.g. a saline composition. The aqueous composition may contain one or more buffers, such as a phosphate buffer (e.g. phosphate buffered saline) or a Tris buffer. Alternatively the composition may be an oil dispersion or an emulsion, e.g. an oil and water emulsion. Alternatively the composition may be a suspension, powder, foam, paste, granule, aerosol, impregnated natural and synthetic substance, or encapsulated in polymeric substance for example. A suitable form of the composition may be chosen for the intended use having regard to the target insect, and to its habitat.

[0299] Adjuvants may enhance product performance, for example, by increasing the efficiency of the delivery of

active ingredients, reducing the level of active ingredient required, or extending the spectrum of effectiveness.

[0300] Different types of adjuvants offer various benefits and advantages, which are achieved by modulating properties such as spray formation, spray retention, wetting, deposit formation or uptake.

[0301] Adjuvants modulating spray formation may influence spray quality by reducing spray drift and wastage, allowing more of the product to reach the target. This can reduce use rates, leading to a better environmental profile and a potentially more cost effective solution. Such adjuvants include non-ionic surfactants and emulsifier blends.

[0302] Adjuvants modulating spray retention may dissipate the kinetic energy of the droplet during impact, meaning the likelihood of bounce or run-off is reduced. Such adjuvants include alkyl polyglucosides, alkoxyated alcohols, and polyoxyethylene monobranched alcohols (e.g. polyoxyethylene (8) monobranched alcohol).

[0303] Adjuvants modulating wetting properties (i.e. wetting agents) may reduce surface tension and contact angle, leading to enhanced coverage. Such adjuvants include polyoxyethylene sorbitan monolaurate (e.g. polyoxyethylene (8) sorbitan monolaurate), surfactant blends, and alkyl polyglucosides.

[0304] Adjuvants modulating deposit formation may influence evaporation of water from the droplet and thus provide a more homogeneous distribution. Such adjuvants include alkoxyated polyol esters, polyoxyethylene sorbitan monolaurate (e.g. polyoxyethylene (12) sorbitan monolaurate), and alkyl polyglucoside.

[0305] Adjuvants modulating uptake can improve penetration and uptake of active ingredients, e.g. through the insect cuticle, resulting in increased bioavailability. Such adjuvants include alkoxyated polyol esters and polyoxyethylene sorbitan monolaurate (e.g. polyoxyethylene (12) sorbitan monolaurate and polyoxyethylene (16) sorbitan monolaurate).

[0306] Dispersants may be aqueous or non-aqueous. An oil dispersion (OD) formulation typically comprises a solid active ingredient dispersed in oil. The oil can vary from paraffinic to aromatic solvent types and vegetable oil or methylated seed oils. Typically the active ingredient is uniformly suspended in the oil phase. Although primarily used for water sensitive active ingredients, OD formulations have extended to other active ingredients due to their better spray retention, spreading, foliar uptake, and penetration enhancement (e.g. across the insect cuticle) as the carrier oil often acts as an adjuvant.

[0307] Oils suitable for use in OD dispersions include linseed, rapeseed and soyabean oils.

[0308] Aqueous dispersants may be used, for example, to improve stability in the spray tank after dilution in water, and may include modified styrene acrylic polymers, and polymeric amphoteric dispersants and adjuvants.

[0309] An emulsifier may be employed to emulsify a continuous oil phase into water when an OD formulation is diluted prior to being sprayed. The emulsifier may be selected based upon its ability to spontaneously form the emulsion. Their performance is primarily dictated by the nature of the surfactant and their collective effect on how they arrange themselves at the oil/water interface. Examples include polyoxyethylene sorbitol hexaoate (e.g. polyoxyethylene (40) sorbitol hexaoate), emulsifier blends, and calcium alkylaryl sulphonate.

[0310] The composition may further comprise an adhesive or a dye.

[0311] The compound may be provided in the form of a concentrate, for dilution prior to application. Alternatively the compound may be provided in a solid form to be suspended or dissolved prior to formulation.

[0312] The compositions as disclosed herein are themselves in fairly diverse, solid or liquid, forms. As solid composition forms, there may be mentioned dustable powders (content of active substance which may be up to 100%) and granules, in particular those obtained by extrusion, by compacting, by impregnation of a granulated carrier, by granulation using a powder as starting material (the content of active substance in these granules being between 0.5 and 80% for these latter cases). Such solid compositions may be optionally used in the form of a liquid which is viscous to a greater or lesser degree, depending on the type of application desired, for example by diluting in water. As liquid composition forms or forms intended to constitute liquid compositions during application, there may be mentioned solutions, in particular water-soluble concentrates, emulsions, suspension concentrates, wettable powders (or spraying powder), oils and waxes. The suspension concentrates, which can be applied by spraying, are prepared so as to obtain a stable fluid product which does not form a deposit and they usually contain from 10 to 75% of active substance, from 0.5 to 15% of surfactants, from 0.1 to 10% of thixotropic agents, from 0 to 10% of appropriate additives, such as antifoams, corrosion inhibitors, stabilizers, penetrating agents and adhesives and, as carrier, water or an organic liquid in which the active substance is not or not very soluble: some organic solids or inorganic salts may be dissolved in the carrier to help prevent sedimentation or as antigels for water.

[0313] A "carrier", as used herein, means any solid, semi-solid or liquid carrier in or on (to) which an active substance, such as the insecticidal compounds or combinations of the invention, can be suitably incorporated, included, immobilized, adsorbed, absorbed, bound, encapsulated, embedded, attached, or comprised. Non-limiting examples of such carriers include nanocapsules, microcapsules, nanospheres, microspheres, nanoparticles, microparticles, liposomes, vesicles, beads, a gel, weak ionic resin particles, liposomes, cochleate delivery vehicles, small granules, granulates, nano-tubes, bucky-balls, water droplets that are part of an water-in-oil emulsion, oil droplets that are part of an oil-in-water emulsion, organic materials such as cork, wood or other plant-derived materials (e.g. in the form of seed shells, wood chips, pulp, spheres, beads, sheets or any other suitable form), paper or cardboard, inorganic materials such as talc, clay, microcrystalline cellulose, silica, alumina, silicates and zeolites, or even microbial cells (such as yeast cells) or suitable fractions or fragments thereof.

[0314] In one embodiment, the carrier is a liposome. In one embodiment, the composition of the invention comprises one more liposomes suitably wherein each liposome comprises a compound of the invention, or combination thereof. Suitably therefore the invention provides a composition or formulation comprising a plurality of liposomes, said liposomes comprising a compound of the invention, or a combination thereof. Suitably the liposomes comprise a lipid component, suitably which may be soy lecithin and glycerol. Suitably the liposomes are formed of soy lecithin and glycerol. Suitably the liposomes are formed of 25 mg/ml

of glycerol and 3% glycerol. Suitably the liposomes comprise an effective concentration of the compound of the invention. Suitably the liposomes comprise around 10-4 M of a compound of the invention, or a combination thereof. Suitably therefore the invention provides a composition or formulation comprising a plurality of liposomes, said liposomes comprising a compound of the invention, or a combination thereof, soy lecithin (preferably 25 mg/ml) and glycerol (preferably 3%).

[0315] Suitably the liposomes may be manufactured by the Mozafari Heating Method (Mozafari, M. R. "Nanoliposomes: preparation and analysis." Liposomes: Methods and Protocols, Volume 1: Pharmaceutical Nanocarriers (2010): 29-50. Suitably such a method comprises (a) admixing the compound of the invention with glycerol, suitably with a 50% glycerol solution, (b) admixing the mixture of step (a) with soy lecithin, suitably with 350 mg of soy lecithin in an aqueous solution, (c) heating the mixture of step (b) whilst stirring to form liposomes, suitably at around 60° C. and at around 800 RPM for around 40 minutes, (d) annealing the liposomes, suitably by placing in 40° C. water for around 1-2 hours, and (e) sonicating the liposomes, suitably for around 30 minutes in a sonicator bath.

[0316] Suitably the liposomes have an average diameter of around 150 to 250 nm, suitably around 175 nm to 225 nm, suitably around 190 nm to 210 nm, suitably around 200 nm. Suitably the size of the liposomes is determined by using Dynamic light scattering (DLS), for example using a Malvern Zetasizer Nano. Suitably the liposomes have a polydispersity index of between 0.25 to 0.35, suitably 0.26 to 0.33, suitably 0.26 to 0.31, suitably 0.26 to 0.30, suitably 0.26 to 0.29, suitably 0.26 to 0.28, suitably around 0.26.

[0317] In the compositions of the invention, the carrier with the one or more insecticidal compounds may for example be maintained as a wettable powder, wettable granule, emulsifiable concentrate, suspension concentrate, microemulsion, capsule suspension, dry microcapsule, tablet or gel or be suspended, dispersed, emulsified or otherwise brought in a suitable liquid medium (such as water or another suitable aqueous, organic or oily medium) so as to provide a (concentrated) liquid composition of the invention that has a stability that allows the composition of the invention to be suitably stored or (where necessary after further dilution) applied to the intended site of action. Suitably the composition of the invention can be transported and/or stored prior to final use, optionally (and usually preferably) as a suitable liquid concentrate, dry powder, tablet, capsule suspension, slurry or "wet cake", which can be suitably diluted, dispersed, suspended, emulsified or otherwise suitably reconstituted by the end user prior to final use. The composition of the invention allows to be applied to the intended site of action using any suitable or desired manual or mechanical technique such as spraying, pouring, dripping, brushing, coating, dripcoating, applying as small droplets, a mist or an aerosol or any other suitable technique. In one embodiment, said intended site of action is an intact living insect, even more preferably an insect surface.

[0318] The composition may be a bait composition for ingestion by the target insect. A bait composition may comprise one or more phagostimulants, i.e. a substance which will entice the insect to ingest the compound. Phagostimulants may include artificial sweeteners, amino acids, other peptides or proteins and carbohydrates (e.g. glucose,

fructose, sucrose, maltose) etc. Examples include honey, syrups and aqueous solutions of sucrose.

[0319] Commercially available base formulations may also be suitable for use in formulating the compounds described in this specification, such as Armid® FMPC (Akzo Nobel).

[0320] The composition may comprise one or more synergists, i.e. compounds which increase the efficacy of insecticides against their targets, often by inhibiting an insect's ability to metabolise the active agent. Common synergists include piperonyl butoxide and MGK-264 (n-octyl bicycloheptane dicarboximide), or peptidase inhibitors.

[0321] The composition may comprise one or more agents which promote stability of the insecticidal compounds of the invention. Suitably the one or more agents which promote stability may prevent degradation of the insecticidal compounds of the invention. Suitable agents which prevent degradation of the insecticidal compounds of the invention may inhibit or reduce the activity of enzymes, suitably of enzymes which act to degrade proteins, suitably proteases, for example. Suitably therefore the composition may comprise one or more protease inhibitors, suitably which may be selected from: Bowman-Birk Inhibitors, Kunitz Inhibitors, Cystatin, Trypsin Inhibitors, Serpins, Tannins, Proteinase Inhibitor-II (PI-II), and Alpha-AI1.

[0322] The composition may further comprise one or more additional, attractants, sterilizing agents, acaricides, nematocides, fungicides, growth-regulating substances or herbicides.

[0323] In some embodiments, the compositions may comprise one or more further agricultural chemicals or agrochemicals such as herbicides (e.g. contact herbicides or systemic herbicides, including herbicides for household use), fungicides (e.g. contact fungicides or systemic fungicides, including fungicides for household use), nematocides (e.g. contact nematocides or systemic nematocides, including nematocides for household use) and other pesticides or biocides (for example agents for killing insects or snails); as well as fertilizers; growth regulators such as plant hormones; micro-nutrients, safeners, pheromones; repellants; insect baits; and/or active principles that are used to modulate (i.e. increase, decrease, inhibit, enhance and/or trigger) gene expression (and/or other biological or biochemical processes) in or by the targeted plant (e.g. the plant to be protected or the plant to be controlled), such as nucleic acids (e.g., single stranded or double stranded RNA, as for example used in the context of RNAi technology) and other factors, proteins, chemicals, etc. known per se for this purpose, etc.

[0324] Examples of such agrochemicals will be clear to the skilled person; and for example include, without limitation: glyphosate, paraquat, metolachlor, acetochlor, mesotrione, 2,4-D, atrazine, glufosinate, sulfosate, fenoxaprop, pendimethalin, picloram, trifluralin, bromoxynil, clodinafop, fluroxypyr, nicosulfuron, bensulfuron, imazetapyr, dicamba, imidacloprid, thiamethoxam, fipronil, chlorpyrifos, deltamethrin, lambda-cyhalothrin, endosulfan, methamidophos, carbofuran, clothianidin, cypermethrin, abamectin, diflufenican, spinosad, indoxacarb, bifenthrin, tefluthrin, azoxystrobin, thiamethoxam, tebuconazole, mancozeb, cyazofamid, fluazinam, pyraclostrobin, epoxiconazole, chlorothalonil, copper fungicides, trifloxystrobin, prothioconazole, difenoconazole, carbendazim, propiconazole,

thiophanate, sulphur, boscalid, chlorantraniliprole and other known agrochemicals or any suitable combination(s) thereof.

Methods of Producing the Insecticidal Compounds

[0325] The insecticidal compounds described herein may be produced by any method known in the art for the production of peptides. In one embodiment, the insecticidal compounds may be produced by a chemical method, suitably a chemical synthesis method.

[0326] Any suitable chemical method may be used to synthesize the insecticidal compounds of the invention.

[0327] Suitably the insecticidal compounds are peptides. Suitably therefore they are synthesised by solid-state peptide synthesis. Suitably such synthesis may be carried out by using commercially available machines such as the Biotage Initiator+ Alstra microwave-assisted peptide synthesiser or the CEM Liberty Prime microwave-assisted peptide synthesiser.

[0328] In one embodiment of the present invention, the insecticidal compound is a peptide of the formula: [Hy]-NADEDQQSVDFTPRL-[NH₂] (SEQ ID NO:2). Suitably this peptide may be synthesised by the method defined in the examples. Suitably by the method defined in FIG. 2.

[0329] In a further aspect of the invention, there is provided an isolated insecticidal compound produced by a process of the invention. Suitably an isolated insecticidal compound of formula (I) produced by a process of the invention.

[0330] In a further aspect of the invention there is provided a method of producing a composition as disclosed herein, at least comprising the steps of: (a) obtaining at least one insecticidal compound of formula (I), and (b) formulating the insecticidal compound into a composition.

[0331] In a further aspect of the invention there is provided a method of producing and/or manufacturing a variant of the insecticidal compound of formula (I). Such may comprise the steps of: (i) modifying the peptide Z of formula (I) by adding, replacing or deleting at least one amino acid; (ii) assessing the so created variant for its insecticidal activity and—optionally—at least one property selected from the group consisting of biostability, chemical stability, bioavailability, solubility (including the ability to form stable formulations), producibility, and production costs. Where the insecticidal activity is reduced in comparison to the unmodified peptide Z, repeating the process of steps (i) and (ii) until a variant with improved insecticidal activity is obtained. Where the insecticidal activity is improved in comparison to the unmodified peptide Z, the method may include further steps of manufacturing, isolating and purifying the variant. Screening for insecticidal activity and the other above described properties can be conducted as described below or generally known to the person skilled in the art.

[0332] In one embodiment is provided a method for manufacturing a variant of the insecticidal compound of the invention, comprising the steps of:

[0333] a) modifying the peptide Z by adding, replacing or deleting at least one amino acid;

[0334] b) assessing the ability of the variant for its insecticidal activity and;

[0335] c) where the insecticidal activity is reduced in comparison to the insecticidal compound comprising

the unmodified peptide Z, repeating the process of steps (a) and (b) until a variant with improved insecticidal activity is obtained;

[0336] d) where the insecticidal activity is improved in comparison to the insecticidal compound comprising the unmodified peptide Z, further manufacturing, isolating and purifying the variant.

[0337] Suitably the step of obtaining at least one insecticidal compound comprises (a) chemically synthesising the insecticidal compound.

[0338] Suitable compositions are described hereinabove. Suitable manufacturing methods for formulating the composition are known in the art and include, but are not limited to, high or low shear mixing, wet or dry milling, drip-casting, encapsulating, emulsifying, coating, encrusting, pilling, extrusion granulation, fluid bed granulation, co-extrusion, spray drying, spray chilling, atomization, addition or condensation polymerization, interfacial polymerization, in situ polymerization, coacervation, spray encapsulation, cooling melted dispersions, solvent evaporation, phase separation, solvent extraction, sol-gel polymerization, fluid bed coating, pan coating, melting, passive or active absorption or adsorption.

[0339] The features disclosed in the foregoing description, or in the following claims, or in the accompanying drawings, expressed in their specific forms or in terms of a means for performing the disclosed function, or a method or process for obtaining the disclosed results, as appropriate, may, separately, or in any combination of such features, be utilised for realising the invention in diverse forms thereof.

[0340] While the invention has been described in conjunction with the exemplary embodiments described above, many equivalent modifications and variations will be apparent to those skilled in the art when given this disclosure. Accordingly, the exemplary embodiments of the invention set forth above are considered to be illustrative and not limiting. Various changes to the described embodiments may be made without departing from the spirit and scope of the invention.

[0341] For the avoidance of any doubt, any theoretical explanations provided herein are provided for the purposes of improving the understanding of a reader. The inventors do not wish to be bound by any of these theoretical explanations.

[0342] Any section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

BRIEF DESCRIPTION OF THE FIGURES

[0343] Aspects and embodiments of the present invention will now be discussed with reference to the accompanying figures in which:

[0344] FIG. 1 shows the chemical structure of SB-P-80;

[0345] FIG. 2 shows the chemical synthesis route for synthesising SB-P-80;

[0346] FIG. 3 shows the lepidoptera artificial diet assay experimental set up;

[0347] FIG. 4 shows activity data of unformulated SB-P-80 against *Spodoptera frugiperda* after 168 hours. One larva was assayed per cell. (n=10).

EXAMPLES

[0348] The invention will now be demonstrated and further described by way of the following non-limiting examples.

General Procedures

[0349] All amino acids are of L-configuration unless otherwise stated. Standard Fmoc-protected amino acids were purchased from CEM Corporation or Pepceuticals. Suppliers of specialist amino acids are indicated as and when appropriate, as is peptide synthesis resin supplier. Peptide-grade DMF was purchased from Rathburn.

[0350] Peptides were synthesised on a Biotage Initiator+Alstra microwave-assisted peptide synthesiser or a CEM Liberty Prime microwave-assisted peptide synthesiser as specified.

[0351] High-resolution mass spectrometry (HRMS) was performed on a Bruker microTOF-Q II (ESI+).

[0352] Peptides were purified on a reverse-phase Dionex HPLC system equipped with Dionex P680 pumps and a Dionex UVD170U UV-vis detector (monitoring at 214 nm and 280 nm), using a Phenomenex, Gemini, C18, 5 μ m, 250 \times 21.2 mm column. Gradients were performed using solvents consisting of A (H₂O+0.1% TFA) and B (MeCN+0.1% TFA) and fractions were lyophilised on a Christ Alpha 2-4 LO plus freeze dryer.

[0353] Pure peptides were analysed on a Shimadzu reverse-phase HPLC (RP-HPLC) system equipped with Shimadzu LC-20AT pumps, a SIL-20A autosampler and a SPD-20A UV-vis detector (monitoring at 214 nm and 280 nm) using a Phenomenex, Aeris, 5 μ m, peptide XB-C18, 150 \times 4.6 mm column at a flow rate of 1 mL/min. RP-HPLC gradients were run using a solvent system consisting of solution A (100% H₂O+0.1% TFA) and B (100% MeCN+0.1% TFA). Typically, two gradients were used to characterise each peptide; a gradient from 5% to 95% solution B over 20 min (incorporating a 2 min hold at 5% solution B and a 5 min wash at 95% solution B at the start and end of the gradient respectively) and a gradient from 5-95% solution B over 50 min (incorporating a 5 min hold at 5% solution B and a 5 min wash at 95% solution B at the start and end of the gradient respectively). In some cases, specialised gradients were used and this is indicated where appropriate. Analytical RP-HPLC data is reported as column retention time (tR) in minutes (min). Analytical columns were maintained at ambient temperature.

[0354] LC-MS analysis was performed on a Thermo Scientific LCQ Fleet quadrupole mass spectrometer of m/z range 50-2000 Da with an ESI source coupled to a Dionex Ultimate 3000 LC. Analyses were performed on a ReprosilGold 120 C18, 3 μ m 150 \times 4 mm column using a linear gradient of buffer A (95/5 H₂O/MeCN with 0.1% v/v TFA) to buffer B (95/5 MeCN/H₂O with 0.1% v/v TFA) over 20 min (incorporating a 2 min hold at 0% solution B and a 5 min wash at 100% solution B at the start and end of the gradient respectively). In cases where a specialised analytical RP-HPLC gradient was used to characterise a compound, an analogous LC-MS gradient was used. Analytical RP-HPLC and LC-MS samples were injected as 25 μ L of a stock with concentration of 1 mg/ml in H₂O/MeCN with 0.1% v/v TFA. LC-MS column oven temperature was maintained at 30° C.

[0355] High-resolution mass spectrometry (HRMS) of pure peptides was performed on a Bruker microTOF-Q II (ESI+).

[0356] Proton nuclear magnetic resonance spectra (^1H NMR) for the purpose of peptide content calculations were recorded on an AVANCE III 400 Bruker (400 MHz). Proton chemical shifts are expressed in parts per million (ppm, δ scale) and are referenced to residual protium in the NMR solvent (CDCl_3 , δ 7.26; CD_3OD , δ 3.31 and D_2O , δ 4.79). The following abbreviations were used to describe peak patterns when appropriate: br=broad, s=singlet, d=doublet, t=triplet, q=quadruplet, m=multiplet. Coupling constants, J, are reported in Hertz unit (Hz).

General Procedure for Automated Peptide Synthesis

Biotage Initiator+ Alstra Synthesiser:

[0357] Fmoc-protected amino acids were prepared as a 0.2 M (0.1 mmol syntheses), 0.5 M (0.2 mmol syntheses) or 0.7 M (0.5 mmol syntheses) solution in DMF. 5 equivalents of amino acid (relative to the resin loading) were used during coupling cycles. Oxyma and diisopropyl carbodiimide (DIC) were prepared as 0.2 M (0.1 mmol syntheses), 0.5 M (0.2 mmol syntheses) or 0.7 M (0.5 mmol syntheses) solutions in DMF. 5 equivalents of Oxyma and 5 equivalents of DIC (relative to resin loading) were used during coupling cycles. For Fmoc-deprotections, a solution of 20% morpholine (with 5% formic acid) in DMF was used. Coupling reactions were performed under microwave heating at 90° C. for 2 min with the exception of Fmoc-Cys (Trt)-OH, Fmoc-His (Trt)-OH and Fmoc-Arg (Pbf)-OH. Coupling of Fmoc-Cys (Trt) and Fmoc-His (Trt)-OH was performed for 10 min at 50° C. Coupling of Fmoc-Arg (Pbf)-OH was performed for 2 successive cycles (double-coupled) for 2 min at 90° C. Microwave-assisted Fmoc deprotections were carried out at 90° C. for 1 min.

CEM Liberty Blue Synthesiser

[0358] Fmoc-protected amino acids were prepared as a 0.2 M solution in NBP (Tamisolve). 5 equivalents of amino acid (relative to the resin loading) were used during coupling cycles. Oxyma was prepared as a 0.5 M solution in NBP. DIC was prepared as 5 M solution in NBP. 5 equivalents of Oxyma and 5 equivalents of DIC (relative to resin loading) were used during coupling cycles. For Fmoc-deprotections, a solution of 20% pyrrolidine was used. Coupling reactions and Fmoc-deprotections were performed under microwave heating at 90° C. for 2 min and 1 min respectively with the exception of Fmoc-Cys (Trt)-OH, Fmoc-His (Trt)-OH and Fmoc-Arg (Pbf)-OH. Coupling of Fmoc-Cys (Trt) and Fmoc-His (Trt)-OH was performed for 5 min at 50° C. Coupling of Fmoc-Arg (Pbf)-OH was performed for 2 successive cycles of 5 min at 75° C.

General Procedure for TFA Cleavage of Peptides

[0359] Typically, cleavage tests of peptides were performed by taking ~3 mg of dried resin beads and treating them with TFA/TIS/water (95:2.5:2.5) for 3 h. The filtrate was drained, concentrated and then triturated in cold diethyl ether (Et_2O). The triturate was dissolved in acetonitrile/water, then analysed by RP-HPLC/LC-MS.

[0360] Peptides were typically cleaved from the resin in bulk by gently rocking the resin at rt in a cleavage cocktail

of TFA/TIS/ H_2O (95:2.5:2.5) for 3 h before being drained and the TFA blown off with a steady stream of N_2 gas. Peptides containing cysteine or tryptophan residues were cleaved from the resin using a cleavage cocktail of TFA/TIS/ H_2O /DODT (94:2.5:1:2.5) for 3 h. In all cases the crude peptide was triturated with cold Et_2O . Et_2O was removed from the resulting crude peptide pellet under a steady stream of nitrogen. The crude peptide was then redissolved in H_2O /MeCN and purified by RP-HPLC.

[0361] A specific synthesis route for SB-P-80 is shown in FIG. 2. For the synthesis of SB-P-80, double coupling methods are used for 9Ser, 11Asp, 13Thr. Extended 6 min couplings are used for 3Asp, 4Glu, 5Asp, 6Gln and 7Gln, and a 7 min arginine coupling are used for 15Arg.

[0362] Unless otherwise stated herein, peptides were synthesised according to the general procedures above, in line with the exemplary procedures below, or were ordered directly.

Mortality Assessment Methods

Lepidoptera Artificial Diet Assay

[0363] Frontier Lepidoptera artificial diet was made up by preparing agar at 1% and then mixing it with artificial diet powder. Peptide solution was aliquoted in 10 cells and diet poured straight onto it. The final concentration of the peptide was 1×10^{-4} M, and each peptide was aliquoted in 10 cells. Once the diet solidified and cooled down, one L1 larvae was infested in each cell. The trays were sealed with heat and let larva were allowed to develop for 168 hrs. Finally, pupation and mortality were scored for. The lepidoptera artificial diet assay set up is shown in FIG. 3. Lepidoptera artificial diet assays were performed with SB-P-80, and compared to a negative control vehicle and the insecticide dichlorodiphenyltrichloroethane (DDT). The activity of SB-P-80 is shown in FIG. 4; SB-P-80 has much higher activity whilst also being eco-friendly and targeted to specific pests unlike broad spectrum chemical insecticides such as DDT.

REFERENCES

- [0364]** A number of publications are cited in order to more fully describe and disclose the invention and the state of the art to which the invention pertains. Full citations for these references are provided below. The entirety of each of these references is incorporated herein by reference.
- [0365]** Koyama T, Terhzaz S, Naseem M T, Nagy S, Rewitz K, Dow J A T, Davies S A, Halberg K V. A nutrient-responsive hormonal circuit mediates an inter-tissue program regulating metabolic homeostasis in adult *Drosophila*. Nat Commun. 2021
- [0366]** Kean L, Cazenave W, Costes L, Broderick K E, Graham S, Pollock V P, Davies S A, Veenstra J A, Dow J A. Two nitridergic peptides are encoded by the gene capability in *Drosophila melanogaster*. Am J Physiol Regul Integr Comp Physiol. 2002
- [0367]** Yeoh J G C, Pandit A A, Zandawala M, Nässel D R, Davies S A, Dow J A T. DInER: Database for Insect Neuropeptide Research. Insect Biochem Mol Biol. 2017
- [0368]** Ahn S J, Corcoran J A, Vander Meer R K, Choi M Y. Identification and Characterization of GPCRs for Pyrokinin and CAPA Peptides in the Brown Marmorated Stink Bug, *Halyomorpha halys* (Hemiptera: Pentatomidae). Front Physiol. 2020

- [0369] Audsley N and Down R E, G protein coupled receptors as targets for next generation pesticides. *Insect Biochem Molec* 67:27-37 (2015).
- [0370] Halberg K A, Terhzaz S, Cabrero P, Davies S A and Dow J A T, Tracing the evolutionary origins of insect renal function. *Nat Commun* 6 (2015).
- [0371] Dow J A, Insights into the Malpighian tubule from functional genomics. *J Exp Biol* 212:435-445 (2009).
- [0372] Huesmann G R, Cheung C C, Loi P K, Lee T D, Swiderek K M and Tublitz N J, Amino acid sequence of CAP2b, an insect cardioacceleratory peptide from the tobacco hawkmoth *Manduca sexta*. *FEBS Lett* 371:311-314 (1995).
- [0373] Davies S A, Cabrero P, Povsic M, Johnston N R, Terhzaz S and Dow J A T, Signaling by *drosophila* capa neuropeptides. *Gen Comp Endocr* 188:60-66 (2013).
- [0374] Terhzaz S, Teets N M, Cabrero P, Henderson L, Ritchie M G, Nachman R J, Dow J A T, Denlinger D L and Davies S A, Insect capa neuropeptides impact desiccation and cold tolerance. *Proc Natl Acad Sci* 201501518 (2015).
- [0375] Terhzaz S, Alford L, Yeoh J G C, Marley R, Dornan A T, Dow J A T and Davies S A, Renal neuroendocrine control of desiccation and cold tolerance by *Drosophila* suzukii. *Pest Manag Sci* 74:800-810 (2017).
- [0376] Predel R, Wegener C, Biology of the CAPA peptides in insects. *Cell Mol Life Sci* 63:2477-2490 (2006).
- [0377] Lamango N S, Nachman R J, Hayes T K, Strey A and Isaac R E, Hydrolysis of insect neuropeptides by an angiotensin converting enzyme from the housefly, *M. domestica*. *Peptides* 18:47-52 (1997).
- [0378] Terhzaz S, Cabrero P, Robben J H, Radford J C, Hudson B D, Milligan G, Dow J A and Davies S A, Mechanism and function of *Drosophila* capa GPCR: a desiccation stress-responsive receptor with functional homology to human neuromedinU receptor. *PLOS One* 7 (1): e29897 (2012).
- [0379] Blackman R L and Eastop V F, Aphids on the World's Crops: An Identification Guide, John Wiley & Sons Ltd, Chichester, UK. (2000).
- [0380] Dow J A T, Maddrell S H P, Gortz A, Skaer N J V, Brogan S and Kaiser K, The Malpighian tubules of *Drosophila melanogaster*—a novel phenotype for studies of fluid secretion and its control. *J Exp Biol* 197:421-428 (1994).
- [0381] Davies S A, Huesmann G R, Maddrell S H, O'Donnell M J, Skaer N J, Dow J A T and Tublitz N J, CAP2b, a cardioacceleratory peptide, is present in *Drosophila* and stimulates tubule fluid secretion via cGMP. *Am J Physiol* 269: R1321-R1326 (1995).
- [0382] Beyenbach K W, Skaer H and Dow J A, The developmental, molecular, and transport biology of Malpighian tubules. *Annu Rev Entomol* 55:351-374 (2010).
- [0383] Sadeghi A., Van Damme, E. J. M. and Smagghe G. (2009) Evaluation of the susceptibility of the pea aphid, *Acyrtosiphon pisum*, to a selection of novel biorational insecticides using an artificial diet. *Journal of Insect Science* 9:65.
- [0384] Van Emden F (2009). Artificial diet for aphids—thirty years' experience. *REDIA*, XCII: 163-167

SEQUENCE LISTING

```

Sequence total quantity: 17
SEQ ID NO: 1          moltype = AA  length = 16
FEATURE              Location/Qualifiers
source                1..16
                     mol_type = protein
                     note = Z peptide
                     organism = synthetic construct

SEQUENCE: 1
NADEDQQQSV DFTPLR                                           16

SEQ ID NO: 2          moltype = AA  length = 16
FEATURE              Location/Qualifiers
source                1..16
                     mol_type = protein
                     note = SB-P-80
                     organism = synthetic construct

MOD_RES              16
                     note = leucinamide

SEQUENCE: 2
NADEDQQQSV DFTPLR                                           16

SEQ ID NO: 3          moltype = AA  length = 11
FEATURE              Location/Qualifiers
source                1..11
                     mol_type = protein
                     note = SB-P-47
                     organism = synthetic construct

MOD_RES              11
                     note = leucinamide

MOD_RES              1
                     note = pyroglutamyl-alanine

SEQUENCE: 3
AIMARPQVPR L                                                11

SEQ ID NO: 4          moltype = AA  length = 8
FEATURE              Location/Qualifiers
source                1..8

```

-continued

	mol_type = protein note = SB-P-49 organism = synthetic construct	
MOD_RES	8 note = leucinamide	
SEQUENCE: 4 SVPFKPRL		8
SEQ ID NO: 5 FEATURE source	moltype = AA length = 19 Location/Qualifiers 1..19 mol_type = protein note = SB-P-51 organism = synthetic construct	
MOD_RES	19 note = leucinamide	
SEQUENCE: 5 LRQLQSNQEP AYRVTPRL		19
SEQ ID NO: 6 FEATURE source	moltype = AA length = 12 Location/Qualifiers 1..12 mol_type = protein note = SB-P-46 organism = synthetic construct	
MOD_RES	12 note = leucinamide	
SEQUENCE: 6 SPPYSPPFSP RL		12
SEQ ID NO: 7 FEATURE source	moltype = AA length = 10 Location/Qualifiers 1..10 mol_type = protein note = SB-P-81 organism = synthetic construct	
MOD_RES	10 note = leucinamide	
SEQUENCE: 7 GGSMTFSPRL		10
SEQ ID NO: 8 FEATURE source	moltype = AA length = 8 Location/Qualifiers 1..8 mol_type = protein note = SB-P-70 organism = synthetic construct	
MOD_RES	8 note = glycinamide	
SEQUENCE: 8 KVKFSAWG		8
SEQ ID NO: 9 FEATURE source	moltype = AA length = 10 Location/Qualifiers 1..10 mol_type = protein note = SB-P-65 organism = synthetic construct	
MOD_RES	10 note = glycinamide	
SEQUENCE: 9 RQKTVFSSWG		10
SEQ ID NO: 10 FEATURE source	moltype = AA length = 7 Location/Qualifiers 1..7 mol_type = protein note = SB-P-66 organism = synthetic construct	
MOD_RES	7 note = glycinamide	
SEQUENCE: 10 PAFSSWG		7
SEQ ID NO: 11 FEATURE	moltype = AA length = 9 Location/Qualifiers	

-continued

source	1..9	
	mol_type = protein	
	note = SB-P-86	
	organism = synthetic construct	
MOD_RES	9	
	note = glycineamide	
MOD_RES	1	
	note = pyroglutamyl-leucine	
SEQUENCE: 11		
LTFTSSWGG		9
SEQ ID NO: 12	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
source	1..8	
	mol_type = protein	
	note = SB-P-039	
	organism = synthetic construct	
MOD_RES	1	
	note = palmitoyl-glutamine	
MOD_RES	8	
	note = tryptophanamide	
SEQUENCE: 12		
QLTFSPDW		8
SEQ ID NO: 13	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
source	1..7	
	mol_type = protein	
	note = SB-P-042	
	organism = synthetic construct	
MOD_RES	1	
	note = pyroglutamyl-leucine	
MOD_RES	7	
	note = tryptophanamide	
SEQUENCE: 13		
LTFSPDW		7
SEQ ID NO: 14	moltype = AA length = 6	
FEATURE	Location/Qualifiers	
source	1..6	
	mol_type = protein	
	note = SB-P-69	
	organism = synthetic construct	
MOD_RES	6	
	note = glycineamide	
SEQUENCE: 14		
NFSPWG		6
SEQ ID NO: 15	moltype = AA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = protein	
	note = SB-P-48	
	organism = synthetic construct	
MOD_RES	19	
	note = leucinamide	
SEQUENCE: 15		
EQNVQSNQEP AYRVTPRL		19
SEQ ID NO: 16	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
source	1..8	
	mol_type = protein	
	note = SB-P-41	
	organism = synthetic construct	
MOD_RES	8	
	note = tryptophanamide	
SEQUENCE: 16		
QLTFSPDW		8

-continued

```

SEQ ID NO: 17      moltype = AA  length = 16
FEATURE            Location/Qualifiers
source              1..16
                    mol_type = protein
                    note = Fig2, middle
                    organism = synthetic construct
MOD_RES            16
                    note = Modified with Rink Amide resin
SEQUENCE: 17
NADEDQQQSV DFTPLR

```


16

1. An insecticidal compound having the formula (I) below, or a salt or solvate thereof:



wherein:

R^1 is hydrogen (which may be designated "H-" or "Hy-"), C_{1-4} alkyl (e.g. methyl, ethyl, propyl, butyl), formyl, acyl, fatty acyl, a sugar moiety, phosphate or sulphate, or R^1 is a pyroglutamate group of the formula:



and any alkyl, formyl, acyl or fatty acyl may optionally be substituted with one or more groups selected from oxo or C_{1-6} alkyl, or a sugar moiety, phosphate or sulphate; Y^1 is absent or is a peptide comprising from 1 to 2 amino acids;


Z is a peptide according to formula:



Y^2 is absent or is a peptide comprising from 1 to 2 amino acids;


R^2 is NH_2 , $NR^{2a}H$, $NR^{2a}R^{2b}$, OH or OR^{2a} ; wherein each of R^{2a} and R^{2b} if present are independently C_{1-6} alkyl (e.g. methyl, ethyl, propyl, butyl, pentyl or hexyl).

2. The insecticidal compound according to claim 1, wherein R^1 is selected from hydrogen, acyl or fatty acyl, or phosphate or sulphate, or R^1 is a pyroglutamate group of the formula:



wherein acyl or fatty acyl are optionally substituted by a sugar moiety, phosphate or sulphate.

3. The insecticidal compound according to claim 1, wherein R^1 is selected from hydrogen, formyl, acetyl (Ac), propanoyl, butanoyl, palmitoyl, butyryl, cerotoyl, decanoyl, docosenoyl, dodecanoyl, eleostearoyl, heptanoyl, hexanoyl, icosanoyl, icosenoyl, lignoceroyl, linoleoyl, lipoyl, myristoleoyl, nonanoyl, octadecanoyl, ocatanoyl, palmitoleoyl, stearoyl, undecanoyl, valeryl, and a pyroglutamate group of the formula:



optionally wherein R^1 is selected from hydrogen, or palmitoyl, optionally wherein R^1 is hydrogen.


4. The insecticidal compound according to claim 1, wherein:

Y^1 is absent; and/or

wherein Y^2 is present and is selected from: GR, RG, KR, RK, K, R, and G; and/or

wherein R^2 is NH_2 .

5. The insecticidal compound according to claim 1, having the formula:



6. A composition comprising a compound according to claim 1 in admixture with one or more solvents, carriers, diluents, adjuvants, preservatives, dispersants, emulsifying agents, or synergists; optionally wherein the composition is an aqueous composition, optionally wherein the composition comprises an effective amount of the compound.

7. The composition according to claim 6, wherein the composition is an agricultural composition, optionally wherein the composition is an insect control composition or a plant protection composition.

* * * * *