

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12390498
Kind Code	B2
Date of Patent	August 19, 2025
Inventor(s)	Schneider; Jessica et al.

Treatment of clostridium difficile infection

Abstract

Provided herein are compositions and methods for the treatment or prevention of pathogenic infections.

Inventors: Schneider; Jessica (Cambridge, MA), Kim; Yun-Gi (Watertown, MA), Olle; Bernat (Cambridge, MA), Reddy; Shilpa (Watertown, MA), Norman; Jason (North Weymouth, MA), Patarroyo; Juan (Lexington, MA)

Applicant: Vedanta Biosciences, Inc. (Cambridge, MA)

Family ID: 1000008766535

Assignee: Vedanta Biosciences, Inc. (Cambridge, MA)

Appl. No.: 18/326511

Filed: May 31, 2023

Prior Publication Data

Document Identifier	Publication Date
US 20240123000 A1	Apr. 18, 2024

Related U.S. Application Data

continuation parent-doc US 16702659 20191204 US 11701396 child-doc US 18326511
continuation parent-doc US 16423487 20190528 US 10555980 child-doc US 16702659
continuation parent-doc US 16157640 20181011 US 10456431 child-doc US 16423487
continuation parent-doc US 15993037 20180530 US 10350250 child-doc US 16157640
continuation parent-doc US 15630088 20170622 US 9999641 child-doc US 15993037
continuation parent-doc WO PCT/US2017/037498 20170614 PENDING child-doc US 15630088
us-provisional-application US 62349914 20160614

Publication Classification

Int. Cl.: A61K35/747 (20150101); A23L5/00 (20160101); A23L33/135 (20160101); A61K9/00 (20060101); A61K9/48 (20060101); A61K35/74 (20150101); A61K35/742 (20150101); A61K38/14 (20060101); C12N1/20 (20060101); A23K10/18 (20160101); A23K50/30 (20160101)

U.S. Cl.:

CPC A61K35/747 (20130101); A23L5/00 (20160801); A23L33/135 (20160801); A61K9/0053 (20130101); A61K9/48 (20130101); A61K35/74 (20130101); A61K35/742 (20130101); A61K38/14 (20130101); C12N1/20 (20130101); A23K10/18 (20160501); A23K50/30 (20160501); A61K9/0031 (20130101)

Field of Classification Search

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
6635260	12/2002	Gerding	N/A	N/A
8460648	12/2012	Borody	N/A	N/A
9386793	12/2015	Blaser et al.	N/A	N/A
9642881	12/2016	Honda et al.	N/A	N/A
9649345	12/2016	Honda et al.	N/A	N/A
9999641	12/2017	Schneider	N/A	A23L 5/00
10064904	12/2017	Schneider	N/A	C12N 1/20
10350250	12/2018	Schneider	N/A	A23L 5/00
10456431	12/2018	Schneider	N/A	A61K 35/74
10555980	12/2019	Schneider	N/A	A61K 38/14
11701396	12/2022	Schneider	424/93.3	A23L 33/135
2004/0028689	12/2003	Borody	N/A	N/A
2009/0269321	12/2008	Sashihara et al.	N/A	N/A
2010/0074872	12/2009	Blaser et al.	N/A	N/A
2013/0045274	12/2012	Hlavka	N/A	N/A
2013/0195804	12/2012	Borody	N/A	N/A
2014/0199281	12/2013	Henn et al.	N/A	N/A
2014/0328803	12/2013	McKenzie et al.	N/A	N/A
2014/0341921	12/2013	Honda et al.	N/A	N/A
2015/0037476	12/2014	Dhingra et al.	N/A	N/A
2015/0079209	12/2014	Kameyama et al.	N/A	N/A
2016/0022745	12/2015	Wang	N/A	N/A
2016/0022746	12/2015	Lawley et al.	N/A	N/A
2016/0040215	12/2015	Henn et al.	N/A	N/A
2016/0113971	12/2015	Kaplan et al.	N/A	N/A
2016/0193256	12/2015	Honda et al.	N/A	N/A
2016/0193257	12/2015	Honda et al.	N/A	N/A
2016/0228476	12/2015	Cutcliffe et al.	N/A	N/A
2017/0143772	12/2016	Mulder et al.	N/A	N/A
2017/0209502	12/2016	Honda et al.	N/A	N/A
2017/0216378	12/2016	Honda et al.	N/A	N/A
2017/0290889	12/2016	Loke et al.	N/A	N/A
2017/0354697	12/2016	Schneider et al.	N/A	N/A
2018/0169153	12/2017	Berry et al.	N/A	N/A
2018/0169157	12/2017	Schneider et al.	N/A	N/A
2018/0221286	12/2017	Kabadi et al.	N/A	N/A
2018/0264056	12/2017	Schneider et al.	N/A	N/A
2019/0030098	12/2018	Schneider et al.	N/A	N/A
2019/0134106	12/2018	Borody	N/A	N/A
2019/0275090	12/2018	Schneider et al.	N/A	N/A
2020/0206284	12/2019	Schneider et al.	N/A	N/A
2022/0143108	12/2021	Norman et al.	N/A	N/A
2024/0100103	12/2023	Olle et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
3052111	12/2019	EP	N/A
2015-500792	12/2014	JP	N/A
2016-509002	12/2015	JP	N/A
WO 2002/007741	12/2001	WO	N/A
WO 2006/050479	12/2005	WO	N/A
WO 2011/033310	12/2010	WO	N/A
WO 2011/152566	12/2010	WO	N/A
WO 2013/037067	12/2012	WO	N/A
WO 2013/037068	12/2012	WO	N/A
WO 2013/080561	12/2012	WO	N/A

WO 2013/182038	12/2012	WO	N/A
WO 2014/082050	12/2013	WO	N/A
WO 2014/121298	12/2013	WO	N/A
WO 2014/121301	12/2013	WO	N/A
WO 2014/121302	12/2013	WO	N/A
WO 2014/145958	12/2013	WO	N/A
WO 2014/153194	12/2013	WO	N/A
WO 2015/006355	12/2014	WO	N/A
WO 2015/051323	12/2014	WO	N/A
WO 2015/077794	12/2014	WO	N/A
WO 2015/095241	12/2014	WO	N/A
WO 2015/156419	12/2014	WO	N/A
WO 2015/164555	12/2014	WO	N/A
WO 2015/179437	12/2014	WO	N/A
WO 2016/086161	12/2015	WO	N/A
WO 2016/086205	12/2015	WO	N/A
WO 2016/086206	12/2015	WO	N/A
WO 2016/086208	12/2015	WO	N/A
WO 2016/086209	12/2015	WO	N/A
WO 2016/086210	12/2015	WO	N/A
WO 2016/185469	12/2015	WO	N/A
WO 2016/201053	12/2015	WO	N/A
WO 2016/203217	12/2015	WO	N/A
WO 2016/203218	12/2015	WO	N/A
WO 2016/203220	12/2015	WO	N/A
WO 2016/203221	12/2015	WO	N/A
WO 2016/203223	12/2015	WO	N/A
WO 2016/209806	12/2015	WO	N/A
WO 2017/008026	12/2016	WO	N/A
WO 2017/035188	12/2016	WO	N/A
WO 2017/075098	12/2016	WO	N/A
WO 2017/085518	12/2016	WO	N/A
WO 2017/085520	12/2016	WO	N/A
WO 2017/089794	12/2016	WO	N/A
WO 2017/089795	12/2016	WO	N/A
WO 2017/091783	12/2016	WO	N/A
WO 2017/148596	12/2016	WO	N/A
WO 2017/218680	12/2016	WO	N/A
WO 2018/005606	12/2017	WO	N/A
WO 2018/080477	12/2017	WO	N/A
WO 2019/227085	12/2018	WO	N/A
WO 2020/037271	12/2019	WO	N/A

OTHER PUBLICATIONS

Drancourt et al. 2000 (16S Ribosomal DNA Sequence Analysis of a Large Collection of Environmental and Clinical Unidentifiable Bacterial Isolates; Journal of Clinical Microbiology, vol. 38, No. 10: 3623-3630; (Year: 2000). cited by examiner

Reigadas et al. 2021 (How to: prophylactic interventions for prevention of Clostridiolides difficile infection; Clinical Microbiology and Infection 27: 1777-1783) (Year: 2021). cited by examiner

Smits et al. 2016 (Clostridium difficile infection; Nature Reviews Disease Primers, vol. 2, p. 1-20) (Year: 2016). cited by examiner

Mounsey et al. 2020 (Clostridioides difficile Infection: Update on Management; <https://familydoctor.org/condition/clostridium-difficile-cdiff-infection>) (Year: 2020). cited by examiner

Dsouza et al., Colonization of the live biotherapeutic product VE303 and modulation of the microbiota and metabolites in healthy volunteers. Cell Host Microbe. Apr. 13, 2022;30(4):583-598.e8. doi: 10.1016/j.chom.2022.03.016. cited by applicant

Martiny et al., Phylogenetic conservatism of functional traits in microorganisms. ISME J. Apr. 2013;7(4):830-8. doi: 10.1038/ismej.2012.160. Epub Dec. 13, 2012. cited by applicant

Stackebrandt et al., Authors need to be prudent when assigning names to microbial isolates. Antonie Van Leeuwenhoek. Jan. 2022;115(1):1-5. doi: 10.1007/s10482-021-01675-8. cited by applicant

[No Author Listed], [Clostridium] innocuum strain I46 16S ribosomal RNA gene, partial sequence. GenBank Accession No. KR364751.1. Nov. 28, 2016. 2 pages. cited by applicant

[No Author Listed], [Clostridium] symbiosum gene for 16S ribosomal RNA, partial sequence, strain: JCM 1297. GenBank Accession No. LC036311.1. Mar. 20, 2015. 1 page. cited by applicant

Alang et al., Weight gain after fecal microbiota transplantation. Open Forum Infect Dis. Feb. 4, 2015;2(1):ofv004. doi: 10.1093/ofid/ofv004. eCollection Jan. 2015. cited by applicant

Apisarnthanarak et al., Adjunctive intracolonic vancomycin for severe *Clostridium difficile* colitis: case series and review of the literature. Clin Infect Dis. Sep. 15, 2002;35(6):690-6. Epub Aug. 26, 2002. cited by applicant

Bajaj et al., Fecal microbiota transplant from a rational stool donor improves hepatic encephalopathy: A randomized clinical trial. Hepatology. Dec. 2017;66(6):1727-1738. doi: 10.1002/hep.29306. Epub Oct. 30, 2017. cited by applicant

Blaser, The microbiome revolution. J Clin Invest. Oct. 2014;124(10):4162-5. doi: 10.1172/JCI78366. Epub Oct. 1, 2014. cited by applicant

Bloom et al., Microbiome therapeutics for hepatic encephalopathy. J Hepatol. Dec. 2021;75(6):1452-1464. doi: 10.1016/j.jhep.2021.08.004. Epub Aug. 25, 2021. cited by applicant

Bobilev et al., 1953. VE303, a Rationally Designed Bacterial Consortium for Prevention of Recurrent *Clostridioides difficile* (*C. difficile*) infection (rCDI), Stably Restores the Gut Microbiota After Vancomycin (vanco)-Induced Dysbiosis in Adult Healthy Volunteers (HV). Open Forum Infect Dis. Oct. 2019; 6(Suppl 2): S60. Epub Oct. 23, 2019. doi: 10.1093/ofid/ofz359.130. cited by applicant

Borody et al., Therapeutic faecal microbiota transplantation: current status and future developments. Curr Opin Gastroenterol. Jan. 2014;30(1):97-105. doi: 10.1097/MOG.0000000000000027. cited by applicant

Bucci et al., MDSINE: Microbial Dynamical Systems INference Engine for microbiome time-series analyses. Genome Biol. Jun. 3, 2016;17(1):121. doi: 10.1186/s13059-016-0980-6. cited by applicant

Buffie et al., Precision microbiome reconstitution restores bile acid mediated resistance to *Clostridium difficile*. Nature. Jan. 8, 2015;517(7533):205-8. doi: 10.1038/nature13828. Epub Oct. 22, 2014. cited by applicant

Burns et al., Donor Recruitment and Eligibility for Fecal Microbiota Transplantation: Results From an International Public Stool Bank. Gastro. Apr. 2015;148(4):S96-S97. cited by applicant

Calfee, *Clostridium difficile*: a reemerging pathogen. Geriatrics. Sep. 1, 2008;63(9):10-21. cited by applicant

Cammarota et al., Randomised clinical trial: faecal microbiota transplantation by colonoscopy vs. vancomycin for the treatment of recurrent *Clostridium difficile* infection. Aliment Pharmacol Ther. May 2015;41(9):835-43. doi: 10.1111/apt.13144. Epub Mar. 1, 2015. cited by applicant

Cash et al., Current concepts in the assessment and treatment of hepatic encephalopathy. QJM. Jan. 2010;103(1):9-16. doi: 10.1093/qjmed/hcp152. Epub Nov. 10, 2009. cited by applicant

Demorrow, Bile Acids in Hepatic Encephalopathy. J Clin Exp Hepatol. Jan.-Feb. 2019;9(1):117-124. doi: 10.1016/j.jceh.2018.04.011. Epub May 4, 2018. cited by applicant

Drancourt et al., 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. J Clin Microbiol. Oct. 2000;38(10):3623-30. doi: 10.1128/JCM.38.10.3623-3630.2000. cited by applicant

Eyre et al., Whole-genome sequencing demonstrates that fidaxomicin is superior to vancomycin for preventing reinfection and relapse of infection with *Clostridium difficile*. J Infect Dis. May 1, 2014;209(9):1446-51. doi:10.1093/infdis/jit598. Epub Nov. 11, 2013. cited by applicant

Ferenci et al., Hepatic encephalopathy—definition, nomenclature, diagnosis, and quantification: final report of the working party at the 11th World Congresses of Gastroenterology, Vienna, 1998. Hepatology. Mar. 2002;35(3):716-21. doi: 10.1053/jhep.2002.31250. cited by applicant

Genbank Accession No. NR_104687.1. NCBI. Sakamoto. Feb. 3, 2015. cited by applicant

Hooper et al., Interactions between the microbiota and the immune system. Science. Jun. 8, 2012;336(6086):1268-73. doi: 10.1126/science.1223490. Epub Jun. 6, 2012. cited by applicant

Hughes et al., Immune activation in irritable bowel syndrome: can neuroimmune interactions explain symptoms? Am J Gastroenterol. Jul. 2013;108(7):1066-74. doi: 10.1038/ajg.2013.120. Epub May 7, 2013. cited by applicant

Janda et al., 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: pluses, perils, and pitfalls. J Clin Microbiol. Sep. 2007;45(9):2761-4. doi: 10.1128/JCM.01228-07. Epub Jul. 11, 2007. cited by applicant

Kakihana et al., Fecal microbiota transplantation for patients with steroid-resistant acute graft-versus-host disease of the gut. Blood. Oct. 20, 2016;128(16):2083-2088. doi: 10.1182/blood-2016-05-717652. Epub Jul. 26, 2016. cited by applicant

Kassam et al., Fecal microbiota transplantation for *Clostridium difficile* infection: systematic review and meta-analysis. Am J Gastroenterol. Apr. 2013;108(4):500-8. doi: 10.1038/ajg.2013.59. Epub Mar. 19, 2013. cited by applicant

Khoruts et al., Emergence of fecal microbiota transplantation as an approach to repair disrupted microbial gut ecology. Immunol Lett. Dec. 2014;162(2 Pt A):77-81. doi: 10.1016/j.imlet.2014.07.016. Epub Aug. 10, 2014. cited by applicant

Leblanc et al., Bacteria as vitamin suppliers to their host: a gut microbiota perspective. Curr Opin Biotechnol. Apr. 2013;24(2):160-8. doi: 10.1016/j.copbio.2012.08.005. Epub; Aug. 30, 2012. cited by applicant

Lessa et al., Burden of *Clostridium difficile* infection in the United States. N Engl J Med. Feb. 26, 2015;372(9):825-34. doi:10.1056/NEJMoa1408913. cited by applicant

Louie et al., Fidaxomicin preserves the intestinal microbiome during and after treatment of *Clostridium difficile* infection (CDI) and reduces both toxin reexpression and recurrence of CDI. Clin Infect Dis. Aug. 2012;55 Suppl 2:S132-42. doi: 10.1093/cid/cis338. cited by applicant

Marvola et al., Enteric polymers as binders and coating materials in multiple-unit site-specific drug delivery systems. Eur J Pharm Sci. Feb. 1999;7(3):259-67. cited by applicant

Miller, Fidaxomicin (OPT-80) for the treatment of *Clostridium difficile* infection. Expert Opin Pharmacother. Jun. 2010;11(9):1569-78. doi:10.1517/14656566.2010.485614. cited by applicant

Mullane, Fidaxomicin in *Clostridium difficile* infection: latest evidence and clinical guidance. Ther Adv Chronic Dis. Mar.

2014;5(2):69-84. doi:10.1177/2040622313511285. cited by applicant

Narushima et al., Characterization of the 17 strains of regulatory T cell-inducing human-derived *Clostridia*. *Gut Microbes*. May-Jun. 2014;5(3):333-9. doi: 10.4161/gmic.28572. Epub Mar. 18, 2014. cited by applicant

Paramsothy et al., Donor Recruitment for Fecal Microbiota Transplantation. *Inflamm Bowel Dis*. Jul. 2015;21(7):1600-6. doi: 10.1097/MIB.0000000000000405. cited by applicant

Rose et al., Hepatic encephalopathy: Novel insights into classification, pathophysiology and therapy. *J Hepatol*. Dec. 2020;73(6):1526-1547. doi: 10.1016/j.jhep.2020.07.013. Epub Oct. 21, 2020. cited by applicant

Rossi-Tamisier et al., Cautionary tale of using 16S rRNA gene sequence similarity values in identification of human-associated bacterial species. *Int J Syst Evol Microbiol*. Jun. 2015;65(Pt 6):1929-34. doi:10.1099/ij.s.0.000161. Epub Mar. 3, 2015. cited by applicant

Shannon-Lowe et al., Prevention and medical management of *Clostridium difficile* infection. *BMJ*. Mar. 12, 2010;340:c1296. doi:10.1136/bmj.c1296. cited by applicant

Surawicz, Fecal microbiota transplantation: what we know and what we need to know. *Ann Intern Med*. May 5, 2015;162(9):662-3. doi: 10.7326/M15-0609. cited by applicant

Tannock et al., A new macrocyclic antibiotic, fidaxomicin (OPT-80), causes less alteration to the bowel microbiota of *Clostridium difficile*-infected patients than does vancomycin. *Microbiology*. Nov. 2010;156(Pt 11):3354-9. doi: 10.1099/mic.0.042010-0. Epub Aug. 19, 2010. cited by applicant

Van Nood et al., Duodenal infusion of donor feces for recurrent *Clostridium difficile*. *N Engl J Med*. Jan. 31, 2013;368(5):407-15. doi: 10.1056/NEJMoa1205037. Epub Jan. 16, 2013. cited by applicant

Wang et al., Microbiota-derived butyrate dynamically regulates intestinal homeostasis through regulation of actin-associated protein synaptopodin. *Proc Natl Acad Sci U S A*. May 26, 2020;117(21):11648-11657. doi: 10.1073/pnas.1917597117. Epub May 12, 2020. cited by applicant

Wei et al., Fecal microbiota transplantation restores dysbiosis in patients with methicillin resistant *Staphylococcus aureus* enterocolitis. *BMC Infect Dis*. Jul. 11, 2015;15:265. doi: 10.1186/s12879-015-0973-1. cited by applicant

Xiao et al., Bacterial diversity and community structure of supragingival plaques in adults with dental health or caries revealed by 16S pyrosequencing. *Frontiers in microbiology*. Jul. 22, 2016;7:1145. 15 pages. cited by applicant

Xie et al., Fecal Microbiota Transplantation for Treating Hepatic Encephalopathy: Experimental and Clinical Evidence and Possible Underlying Mechanisms. *Journal of Exploratory Research in Pharmacology* 2018;3(4):119-124. doi: 10.14218/JERP.2018.00017. cited by applicant

Youngster et al., Fecal microbiota transplant for relapsing *Clostridium difficile* infection using a frozen inoculum from unrelated donors: a randomized, open-label, controlled pilot study. *Clin Infect Dis*. Jun. 2014;58(11):1515-22. doi: 10.1093/cid/ciu135. Epub Apr. 23, 2014. cited by applicant

Primary Examiner: Lyons; Mary Maille

Attorney, Agent or Firm: Wolf, Greenfield & Sacks, P.C.

Background/Summary

RELATED APPLICATIONS (1) This application is a continuation of U.S. application Ser. No. 16/702,659, filed Dec. 4, 2019, which is a continuation of U.S. application Ser. No. 16/423,487, filed May 28, 2019, now issued as U.S. Pat. No. 10,555,980, which is a continuation of U.S. application Ser. No. 16/157,640, filed Oct. 11, 2018, now issued as U.S. Pat. No. 10,456,431, which is a continuation of U.S. application Ser. No. 15/993,037, filed May 30, 2018, now issued as U.S. Pat. No. 10,350,250, which is a continuation of U.S. application Ser. No. 15/630,088, filed Jun. 22, 2017, now issued as U.S. Pat. No. 9,999,641, which is a continuation of international application number PCT/US2017/037498, filed Jun. 14, 2017, which claims the benefit under 35 U.S.C. § 119(e) of U.S. provisional application No. 62/349,914, filed Jun. 14, 2016, each of which is incorporated by reference herein in its entirety.

REFERENCE TO AN ELECTRONIC SEQUENCE LISTING

(1) The contents of the electronic sequence listing (P074570006US07-SEQ-NTJ.xml; Size: 340,528 bytes; and Date of Creation: May 15, 2023) is herein incorporated by reference in its entirety.

FIELD OF INVENTION

(2) The disclosure relates to compositions of purified bacterial strains, and methods for treating pathogenic infections, such as *Clostridium difficile* infections, by administering the compositions to a subject having a pathogenic infection.

BACKGROUND OF THE INVENTION

(3) The collection of bacterial, viral, and fungal commensal microorganisms that reside within and on the human body are collectively known as the human microbiome. The bacterial subset of the human microbiome plays an important role in host nutrient acquisition, development, immunological homeostasis, neurological health, and protection against pathogens (LeBlanc et al. *Curr. Opin. Biotechnol.* (2013) 24(2): 160-168; Hooper et al. *Science* (2012) 336(6086): 1268-1273; Hughes et al. *Am. J. Gastroenterol.* (2013) 108(7): 1066-1074). As the largest reservoir of mammalian commensals, bacteria residing in the gastrointestinal (GI) tract influence nearly all of these aspects of human biology (Blaser *J. Clin. Invest.* (2014) 124(10): 4162-4165). Consequently, perturbation of the normal bacterial populations within the GI niche, a state known as

dysbiosis, can predispose humans to a variety of diseases.

(4) *Clostridium difficile* infection (CDI) arises after intestinal colonization by the anaerobic spore-forming Gram-positive pathogen *Clostridium difficile*. Upon colonization of the GI tract, *C. difficile* produces toxins which causes diarrhea and may ultimately lead to death. This illness is the most common identifiable cause of nosocomial diarrhea and is thought to arise as a direct result of dysbiosis (Calfee *Geriatrics* (2008) 63: 10-21; Shannon-Lowe et al *BMJ* (2010) 340: c1296). Not surprisingly, usage of nearly all classes of antibiotics has been associated with CDI, presumably by inducing dysbiosis in the GI tract and thereby enabling *C. difficile* outgrowth. The Center for Disease Control currently classifies CDI as a public health threat requiring immediate and aggressive action because of its natural resistance to many drugs and the emergence of a fluoroquinolone-resistant strain that is now prevalent throughout North America and Europe. *C. difficile* was responsible for almost half a million infections and was associated with approximately 29,000 deaths in 2011 (Lessa et al. *NEJM* 2015 372: 825-834).

(5) The antibiotics metronidazole, vancomycin, and fidaxomicin are the current therapeutic options for treatment of CDI. However, metronidazole is inadequate because of decreased response rates and neither metronidazole nor vancomycin prevent disease recurrence, with up to 30% of patients initially responding experiencing a clinical recurrence after antibiotic cessation (Miller *Expert Opin. Pharmacother.* (2010) 11: 1569-1578). Fidaxomicin has been shown to be superior to vancomycin in preventing recurrent CDI (Mullane *Ther. Adv. Chronic Dis.* (2014) 5(2): 69-84). Because of its narrow spectrum of activity, fidaxomicin is thought to enable normal microbiome repopulation of the gut following dysbiosis and CDI, thereby lowering the likelihood of recurrent disease (Tannock et al. *Microbiology* (2010) 156 (Pt 11): 3354-3359; Louie et al. *Clin. Infect. Dis.* (2012) 55 Suppl. 2: S132-142). Nonetheless, 14% of fidaxomicin-treated patients experience CDI relapse and mutations conferring reduced sensitivity have already been reported (Eyre et al. *J. Infect. Dis.* (2014) 209(9): 1446-1451).

(6) Because the risk of recurrent CDI is heightened by antibiotic use and *C. difficile* spores are inherently recalcitrant to the available chemotherapeutic arsenal, alternative therapeutic modalities are being pursued for the treatment of CDI. Fecal microbiota transplantation (FMT) is one such modality that has shown efficacy against CDI (Khoruts et al. *Immunol. Lett.* (2014) 162(2): 77-81; van Nood et al. *N. Engl. J. Med.* (2013) 368(5): 407-415). To date, results of FMT studies for the treatment of CDI, have reported cure rates up to 90% in three randomized controlled studies (Cammarota et al. *Alimen. Pharmacol. Therap.* (2015) 41(9): 835-843; Kassam et al. *Am. J. Gastroenterol.* (2013) 108(4): 500-508; van Nood et al. *N. Engl. J. Med.* (2013) 368(5): 407-415; Youngster et al. *Infect. Dis. Soc. Am.* (2014) 58(11): 1515-1522).

(7) Despite the success of FMT, this therapeutic approach is not without risks and logistical concerns. Selection of FMT donors is critical and challenging. When FMT donor recruitment is performed with stringent screening and standardization protocols, most prospective donors fail this process. Only 6-10% of prospective FMT donors qualify, with the majority of failures arising from asymptomatic carriage of GI pathogens (Paramsothy et al. *Inflamm. Bowel Dis.* (2015) 21(7): 1600-1606; Borody et al. *Curr. Opin. Gastroenterol.* (2014) 30(10): 97-105; Burns et al. *Gastroenterology* (2015) 148: S96-S97; Surawicz *Ann. Intern. Med.* (2015) 162(9): 662-663). Furthermore, variation between donors may lead to variation in FMT efficacy. In addition, the risk of transmission of even non-infectious illnesses may be heightened by FMT. Indeed, significant weight gain has been reported in a patient who received an FMT from an overweight stool donor (Alang et al. *Open Forum Infect. Dis.* (Winter 2015) 2(1)).

SUMMARY OF THE INVENTION

(8) Provided herein are compositions and methods for the treatment or prevention of pathogenic infections including *C. difficile*.

(9) In one aspect, the disclosure provides compositions comprising two or more purified bacterial strains of species selected from the group consisting of: *Clostridium hathewayi*, *Blautia hansenii*, *Blautia producta*, *Blautia producta* ATCC 27340, *Clostridium bacterium* UC5.1-1D4, *Blautia coccoides*, *Eubacterium contortum*, *Eubacterium fissicatena*, *Sellimonas intestinalis*, *Dracourtella massiliensis*, *Dracourtella massilinesis* GD1, *Ruminococcus torques*, *Anaerostipes caccae*, *Clostridium scindens*, *Marvinbryanta formatexigens*, *Eisenbergiella tayi*, *Flavinofractor plautii*, *Clostridium orbiscindens* 1_3_50 AFAA, *Lachnospiraceae bacterium* 7_1_58 FAA, *Subdoligranulum*, *Anaerotruncus colihominis*, *Anaerotruncus colihominis* DSM 17241, *Clostridium symbiosum*, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae*, *Clostridium bolteae* 90A9, *Dorea longicatena*, *Dorea longicatena* CAG:42, *Clostridium innocuum*, *Erysipelotrichaceae bacterium* 21-3, *Blautia wexlerae*, *Clostridium disporicum*, *Erysipelatoclostridium ramosum*, *Pseudoflavinofractor capillosus*, *Turicibacter sanguinis*, *Lactobacillus mucosae*, *Ruminococcus obeum*, *Megasphaera elsdenii*, *Acidaminococcus fermentans*, *Acidaminococcus intestine*, *Ruminococcus faecis*, *Bacteroides cellulosilyticus*, *Anaerostipes hadrus*, *Eubacterium rectale*, *Ruminococcus champanellensis*, *Ruminococcus albus*, *Bifidobacterium bifidum*, *Blautia luti*, *Roseburia faecis*, *Fusicatenibacter saccharivorans*, *Roseburia faecis*, *Blautia faecis*, *Dorea formicigenerans* and *Bacteroides ovatus*.

(10) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Clostridium hathewayi*, *Blautia hansenii*, *Blautia producta*, *Blautia producta* ATCC 27340, *Clostridium bacterium* UC5.1-1D4, *Blautia coccoides*, *Eubacterium contortum*, *Eubacterium fissicatena*, *Sellimonas intestinalis*, *Dracourtella massiliensis*, *Dracourtella massilinesis* GD1, *Ruminococcus torques*, *Anaerostipes caccae*, *Clostridium scindens*, *Marvinbryanta formatexigens*, *Eisenbergiella tayi*, *Flavinofractor plautii*, *Clostridium orbiscindens* 1_3_50 AFAA, *Lachnospiraceae bacterium* 7_1_58 FAA, *Subdoligranulum*, *Anaerotruncus colihominis*, *Anaerotruncus colihominis* DSM 17241, *Clostridium symbiosum*, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae*, *Clostridium bolteae* 90A9, *Dorea longicatena*, *Dorea longicatena* CAG:42, *Clostridium innocuum*, *Erysipelotrichaceae bacterium* 21-3, *Blautia wexlerae*, *Turicibacter sanguinis*, *Lactobacillus mucosae*, and *Bacteroides ovatus*.

(11) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Clostridium hathewayi*, *Blautia hansenii*, *Blautia producta*, *Blautia coccoides*, *Eubacterium contortum*, *Eubacterium fissicatena*, *Anaerostipes caccae*, *Clostridium scindens*, *Marvinbryanta formatexigens*, and *Eisenbergiella tayi*.

(12) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Flavinofractor plautii*, *Clostridium orbiscindens* 1_3_50 AFAA, *Lachnospiraceae bacterium* 7_1_58 FAA, *Subdoligranulum*, *Anaerotruncus colihominis*, *Anaerotruncus colihominis* DSM 17241, *Eubacterium fissicatena*, *Sellimonas intestinalis*, *Dracourtella massiliensis*, *Dracourtella massilinesis* GD1, *Ruminococcus torques*, *Clostridium symbiosum*, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae*, *Clostridium bolteae* 90A9, *Dorea longicatena*, *Dorea longicatena* CAG:42, *Blautia producta*, *Blautia producta* ATCC 27340, *Clostridium bacterium* UC5.1-1D4, *Clostridium innocuum*, and *Erysipelotrichaceae bacterium* 21-3.

(13) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Clostridium orbiscindens* 1_3_50 AFAA, *Anaerotruncus colihominis* DSM 17241, *Dracourtella massilinesis* GD1, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae* 90A9, *Dorea longicatena* CAG:42, *Clostridium bacterium* UC5.1-1D4, and *Erysipelotrichaceae bacterium* 21-3.

(14) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Clostridium orbiscindens* 1_3_50 AFAA, *Anaerotruncus colihominis* DSM 17241, *Sellimonas intestinalis*, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae* 90A9, *Dorea longicatena* CAG:42, *Clostridium bacterium* UC5.1-1D4, and *Erysipelotrichaceae bacterium* 21_3.

(15) In some embodiments of the compositions provided herein, the composition comprises purified bacterial strains *Clostridium orbiscindens* 1_3_50 AFAA, *Anaerotruncus colihominis* DSM 17241, *Dracourtella massilinesis* GD1, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae* 90A9, *Dorea longicatena* CAG:42, *Clostridium bacterium* UC5.1-1D4, and *Erysipelotrichaceae bacterium* 21-3.

(16) In some embodiments of the compositions provided herein, the composition comprises purified bacterial strains *Clostridium orbiscindens* 1_3_50 AFAA, *Anaerotruncus colihominis* DSM 17241, *Sellimonas intestinalis* GD1, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae* 90A9, *Dorea longicatena* CAG:42, *Clostridium bacterium* UC5.1-1D4, and *Erysipelotrichaceae bacterium* 21_3.

(17) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Flavinofractor plautii*, *Anaerotruncus colihominis*, *Dracourtella massiliensis*, *Clostridium symbiosum*, *Clostridium bolteae*, *Dorea longicatena*, *Blautia producta*, and *Clostridium innocuum*.

(18) In some embodiments of the compositions provided herein, the composition comprises purified bacterial strains *Flavinofractor plautii*, *Anaerotruncus colihominis*, *Dracourtella massiliensis*, *Clostridium symbiosum*, *Clostridium bolteae*, *Dorea longicatena*, *Blautia producta*, and *Clostridium innocuum*.

(19) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Flavinofractor plautii*, *Anaerotruncus colihominis*, *Eubacterium fissicatena*, *Clostridium symbiosum*, *Clostridium bolteae*, *Dorea longicatena*, *Blautia producta*, and *Clostridium innocuum*.

(20) In some embodiments of the compositions provided herein, the composition comprises purified bacterial strains *Flavinofractor plautii*, *Anaerotruncus colihominis*, *Eubacterium fissicatena*, *Clostridium symbiosum*, *Clostridium bolteae*, *Dorea longicatena*, *Blautia producta*, and *Clostridium innocuum*.

(21) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Flavinofractor plautii*, *Lachnospiraceae bacterium* 7_1_58 FAA, *Subdoligranulum*, *Anaerotruncus colihominis*, *Eubacterium fissicatena*, *Ruminococcus torques*, *Clostridium symbiosum*, *Clostridium bolteae*, *Dorea longicatena*, *Blautia producta*, *Clostridium innocuum*, *Erysipelotrichaceae bacterium* 21-3, and *Bacteroides ovatus*.

(22) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Clostridium orbiscindens* 1_3_50 AFAA, *Anaerotruncus colihominis* DSM 17241, *Dracourtella massiliensis* GD1, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae* 90A9, *Dorea longicatena* CAG:42, *Clostridium bacterium* UC5.1-1D4, *Erysipelotrichaceae bacterium* 21-3, and *Bacteroides ovatus*.

(23) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Clostridium orbiscindens* 1_3_50 AFAA, *Anaerotruncus colihominis* DSM 17241, *Sellimonas intestinalis*, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae* 90A9, *Dorea longicatena* CAG:42, *Clostridium bacterium* UC5.1-1D4, *Erysipelotrichaceae bacterium* 21-3, and *Bacteroides ovatus*.

(24) In some embodiments of the compositions provided herein, the composition does not include a bacterial strain of the species *Flavinofractor plautii*, *Subdoligranulum*, or *Lachnospiraceae bacterium* 7_1_58 FAA. In some embodiments of the compositions provided herein, the composition does not include a bacterial strain of the species *Bacteroides ovatus*. The composition of any one of claims 4-12, wherein the composition does not include a bacterial strain of the species *Flavinofractor plautii*, *Subdoligranulum*, *Clostridium orbiscindens* 1_3_50 AFAA, or *Lachnospiraceae bacterium* 7_1_58 FAA.

(25) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Clostridium scindens*, *Clostridium hathewayi*, *Blautia hansenii*, *Blautia wexlerae*, *Blautia producta*, *Blautia coccoides*, *Dorea longicatena*, *Clostridium innocuum*, *Erysipelotrichaceae bacterium* 21-3, *Flavinofractor plautii*, *Lachnospiraceae bacterium* 7_1_58 FAA, *Subdoligranulum*, *Anaerotruncus colihominis*, and *Clostridium symbiosum*. In some embodiments of the compositions provided herein, the

composition does not include a bacterial strain of the species *Flavinofractor plautii*, *Subdoligranulum*, or *Lachnospiraceae bacterium 7_1_58* FAA.

(26) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Clostridium scindens*, *Clostridium hathewayi*, *Blautia hansenii*, *Blautia wexlerae*, *Anaerotruncus colihominis*, *Dorea longicatena*, *Clostridium innocuum*, *Erysipelotrichaceae bacterium 21-3*, *Flavinofractor plautii*, *Lachnospiraceae bacterium 7-1_58* FAA, *Subdoligranulum*, *Turicibacter sanguinis*, and *Lactobacillus mucosae*. In some embodiments of the compositions provided herein, the composition does not include a bacterial strain of the species *Flavinofractor plautii*, *Subdoligranulum* or *Lachnospiraceae bacterium 7_1_58* FAA.

(27) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Dorea longicatena*, *Ruminococcus obeum*, *Megasphaera elsdenii*, *Acidaminococcus fermentans*, *Acidaminococcus intestine*, *Ruminococcus faecis*, *Bacteroides cellulosilyticus*, *Anaerostipes hadrus*, *Flavinofractor plautii*, *Eubacterium rectale*, *Ruminococcus champanellensis*, *Ruminococcus albus*, *Bifidobacterium bifidum*, *Ruminococcus faecis*, *Blautia luti*, *Roseburia faecis*, *Fusicatenibacter saccharivorans*, *Blautia faecis*, *Dorea formicigenerans*, and *Blautia hansenii*.

(28) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains of species selected from the group consisting of: *Acidaminococcus fermentans*, *Acidaminococcus intestine*, *Anaerostipes hadrus*, *Blautia faecis*, *Blautia hansenii*, *Dorea formicigenerans*, *Dorea longicatena*, *Eubacterium rectale*, *Flavinofractor plautii*, *Fusicatenibacter saccharivorans*, *Megasphaera elsdenii*, *Roseburia faecis*, *Ruminococcus champanellensis*, *Ruminococcus albus*, *Ruminococcus faecis*, and *Ruminococcus obeum*.

(29) In one aspect the disclosure provides compositions comprising two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs:1-83 and 124-159.

(30) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs:1-23, SEQ ID NO:83, SEQ ID NOs: 124-159.

(31) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23.

(32) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NOs: 124-159. In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21. In some embodiments of the compositions provided herein, the composition comprises purified bacterial strains that comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21. In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs: 124-159. In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO: 124, SEQ ID NO: 129, SEQ ID NO: 132, SEQ ID NO: 137, SEQ ID NO: 141, SEQ ID NO: 146, SEQ ID NO: 152, and SEQ ID NO: 157. In some embodiments of the compositions provided herein, the composition comprises purified bacterial strains that comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO: 124, SEQ ID NO: 129, SEQ ID NO: 132, SEQ ID NO: 137, SEQ ID NO: 141, SEQ ID NO: 146, SEQ ID NO: 152, and SEQ ID NO: 157.

(33) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20 and SEQ ID NO:21, and wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NO:10. In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs:124-156, and wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence selected from the group consisting of SEQ ID NOs:157-159.

(34) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with

nucleic acid sequences selected from the group consisting of SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:106, SEQ ID NO:110, and SEQ ID NO:122.

(55) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:106, SEQ ID NO:110, and SEQ ID NO:122, and wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NO:93.

(56) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:87, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:106, and SEQ ID NO:122.

(57) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:87, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, and SEQ ID NO:105.

(58) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:104, SEQ ID NO:107, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, SEQ ID NO:118, SEQ ID NO:119, and SEQ ID NO:120.

(59) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:104, SEQ ID NO:107, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, and SEQ ID NO:119.

(60) In some embodiments of the compositions provided herein, the composition comprises two or more purified bacterial strains, wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:86, SEQ ID NO:95, SEQ ID NO:98, SEQ ID NO:110, SEQ ID NO:122, and SEQ ID NO:123.

(61) In some embodiments of the compositions provided herein, the composition comprises at least one bacterial strain from *Clostridium* cluster XIVa and at least one bacterial strain from *Clostridium* cluster XVII. In some embodiments of the compositions provided herein, the composition comprises at least one bacterial strain from *Clostridium* cluster IV and at least one bacterial strain from *Clostridium* cluster XVII. In some embodiments of the compositions provided herein, the composition comprises at least one bacterial strain from *Clostridium* cluster XIVa, at least one strain from *Clostridium* cluster IV and at least one bacterial strain from *Clostridium* cluster XVII.

(62) In some embodiments of the compositions provided herein, the composition comprises at least one *Bacteroides* strain. In some embodiments of the compositions provided herein, the composition does not include *Clostridium scindens*.

(63) In some embodiments of the compositions provided herein, the composition comprises at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, or at least 20 purified bacterial strains.

(64) In some embodiments of the compositions provided herein, one or more of the bacterial strains are spore formers. In some embodiments of the compositions provided herein, one or more of the bacterial strains are in spore form. In some embodiments of the compositions provided herein, each of the bacterial strains is in spore form.

(65) In some embodiments of the compositions provided herein, one or more of the bacterial strains is in vegetative form. In some embodiments of the compositions provided herein, each of the bacterial strains is in vegetative form.

(66) In some embodiments of the compositions provided herein, the composition comprises only obligate anaerobic bacterial strains. In some embodiments of the compositions provided herein, the composition comprises bacterial strains that originate from more than one human donor.

(67) In some embodiments of the compositions provided herein, one or more of the bacterial strains are baiCD⁻. In some embodiments of the compositions provided herein, each of the bacterial strains is baiCD⁻. In some embodiments of the compositions provided herein, the composition does not mediate bile acid 7- α -dehydroxylation. In some embodiments of the compositions provided herein, the composition inhibits *C. difficile* toxin production. In some embodiments of the compositions provided herein, the composition inhibits *C. difficile* replication and/or survival.

(68) In some embodiments of the compositions provided herein, the bacterial strains are lyophilized.

(69) In some embodiments of the compositions provided herein, the composition induces the proliferation and/or accumulation of regulatory T cells (Tregs).

(70) In one aspect, the disclosure provides compositions comprising two or more purified bacterial strains, wherein the composition comprises at least one bacterial strain from *Clostridium* cluster XIVa and at least one bacterial strain from *Clostridium* cluster XVII. In one aspect, the disclosure provides compositions comprising two or more purified bacterial strains, wherein the composition comprises at least one bacterial strain from *Clostridium* cluster IV and at least one bacterial

strain from *Clostridium* cluster XVII. In one aspect, the disclosure provides compositions comprising two or more purified bacterial strains, wherein the composition comprises at least one bacterial strain from *Clostridium* cluster IV, at least one bacterial strain from *Clostridium* cluster XIVa and at least one bacterial strain from *Clostridium* cluster XVII.

(71) In some embodiments of the compositions provided herein, the composition comprises at least one *Bacteroides* strain. In some embodiments of the compositions provided herein, the composition does not include *Clostridium scindens*.

(72) In some embodiments of the compositions provided herein, the composition comprises at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, or at least 20 purified bacterial strains.

(73) In some embodiments of the compositions provided herein, one or more of the bacterial strains are spore formers. In some embodiments of the compositions provided herein, one or more of the bacterial strains are in spore form. In some embodiments of the compositions provided herein, each of the bacterial strains is in spore form.

(74) In some embodiments of the compositions provided herein, one or more of the bacterial strains is in vegetative form. In some embodiments of the compositions provided herein, each of the bacterial strains is in vegetative form.

(75) In some embodiments of the compositions provided herein, the composition comprises only obligate anaerobic bacterial strains.

(76) In some embodiments of the compositions provided herein, the composition comprises bacterial strains that originate from more than one human donor.

(77) In some embodiments of the compositions provided herein, one or more of the bacterial strains are baiCD⁻. In some embodiments of the compositions provided herein, each of the bacterial strains is baiCD⁻. In some embodiments of the compositions provided herein, the composition does not mediate bile acid 7- α -dehydroxylation. In some embodiments of the compositions provided herein, the composition inhibits *C. difficile* toxin production. In some embodiments of the compositions provided herein, the composition inhibits *C. difficile* replication and/or survival.

(78) In some embodiments of the compositions provided herein, the bacterial strains are lyophilized.

(79) In some embodiments of the compositions provided herein, the composition induces the proliferation and/or accumulation of regulatory T cells (Tregs).

(80) In one aspect, the disclosure provides a pharmaceutical composition comprising any of the compositions provided herein further comprising a pharmaceutically acceptable excipient. In some embodiments of the pharmaceutical compositions provided herein, the pharmaceutical composition is formulated for oral delivery. In some embodiments of the pharmaceutical compositions provided herein, the pharmaceutical composition is formulated for rectal delivery. In some embodiments of the pharmaceutical compositions provided herein, the pharmaceutical composition is formulated for delivery to the intestine. In some embodiments of the pharmaceutical compositions provided herein, the pharmaceutical composition is formulated for delivery to the colon. In one aspect, the disclosure provides a food product comprising any of the compositions provided herein further comprising a nutrient.

(81) In one aspect, the disclosure provides a method of treating a pathogenic infection in a subject, comprising administering to the subject a therapeutically effective amount of any of the compositions or food products provided herein to treat the pathogenic infection.

(82) In some embodiments of the methods provided herein, the pathogenic infection is *C. difficile*, Vancomycin Resistant Enterococci (VRE), Carbapenem Resistant Enterobacteriaceae (CRE), *Neisseria gonorrhoeae*, Multidrug Resistant *Acinetobacter*, *Campylobacter*, Extended spectrum beta-lactamase (ESBL) producing Enterobacteriaceae, Multidrug Resistant *Pseudomonas aeruginosa*, *Salmonella*, Drug resistant non-typhoid *Salmonella*, Drug resistant *Salmonella typhi*, Drug resistant *Shigella*, Methicillin Resistant *Staphylococcus aureus*, Drug resistant *Streptococcus pneumoniae*, Drug resistant Tuberculosis, Vancomycin resistant *Staphylococcus aureus*, Erythromycin Resistant Group A *Streptococcus*, Clindamycin resistant Group B *Streptococcus*, and combinations thereof. In some embodiments of the methods provided herein, the pathogenic infection is *C. difficile*. In some embodiments of the methods provided herein, the pathogenic infection is Vancomycin-Resistant Enterococci.

(83) In some embodiments of the methods provided herein, the subject is human. In some embodiments of the methods provided herein, the subject is an asymptomatic carrier.

(84) In some embodiments of the methods provided herein, the subject is administered a dose of an antibiotic prior to administration of the composition. In some embodiments of the methods provided herein, the subject is administered more than one dose of the antibiotic prior to administration of the composition. In some embodiments of the methods provided herein, the subject has not been administered an antibiotic prior to administration of the composition.

(85) In some embodiments of the methods provided herein, the composition is administered to the subject by oral administration. In some embodiments of the methods provided herein, the composition is administered to the subject by rectal administration.

(86) In some embodiments of the methods provided herein, the administering results in proliferation and/or accumulation of regulatory T cells (Tregs).

(87) Each of the limitations of the invention can encompass various embodiments of the invention. It is, therefore, anticipated that each of the limitations of the invention involving any one element or combinations of elements can be included in each aspect of the invention. This invention is not limited in its application to the details of construction and the arrangement of components set forth in the following description or illustrated in the drawings. The invention is capable of other embodiments and of being practiced or of being carried out in various ways.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

(1) The accompanying drawings are not intended to be drawn to scale. The figures are illustrative only and are not required for enablement of the disclosure. For purposes of clarity, not every component may be labeled in every drawing. In the drawings:

(2) FIG. 1 shows the strains of Compositions A-D. Each entry includes the SEQ ID NO of the 16S rDNA sequence of the strain, a strain identifier, and the species with the closest known homology (can be more than one species). The bracketed roman numeral indicates the *Clostridium* cluster classification of each strain based on the closest species homology. Strains that are not classified in Cluster XIVa are highlighted in bold. The two non-clostridial strains (SEQ ID NO:2, closest known species *Turicibacter sanguinis*, and SEQ ID NO:6, closest known species *Lactobacillus mucosae*) do not belong to the *Clostridium* genus.

(3) FIG. 2 shows various *Clostridium difficile* infection models. Timelines indicate antibiotic type, duration of treatment, as well as exposure to *C. difficile* spores. The top panel shows an antibiotic cocktail treatment model in which the antibiotic cocktail is provided in the drinking water from day -10 to day -3 followed by intraperitoneal clindamycin on day -1. The middle panel shows a clindamycin IP injection model, in which clindamycin is administered by intraperitoneal injection on day -1. The bottom panel shows the cefoperazone treatment model, in which cefoperazone is provided in the drinking water from day -12 to day -2, followed by administration of a live biotherapeutic product (LBP) on day -1.

(4) FIG. 3 shows the experimental conditions described in Example 1. The groups of mice were divided based on the antibiotic regimen received prior to administration of the indicated amount of *C. difficile* spores. "Abx" refers to treatment with any of the antibiotic regimens.

(5) FIGS. 4A-4L show data obtained in Example 1. FIGS. 4A-4D show survival of mice that received no treatment (FIG. 4A), antibiotic cocktail (FIG. 4B), clindamycin (FIG. 4C), or cefoperazone (FIG. 4D) prior to *C. difficile* infection. FIGS. 4E-4H show body weight of mice that received no treatment (FIG. 4E), antibiotic cocktail (FIG. 4F), clindamycin (FIG. 4G), or cefoperazone (FIG. 4H) prior to *C. difficile* infection. FIGS. 4I-4L show *C. difficile* burden (CFU) per gram of feces from mice that received no treatment (FIG. 4I), antibiotic cocktail (FIG. 4J), clindamycin (FIG. 4K), or cefoperazone (FIG. 4L) prior to *C. difficile* infection. Open circles indicate infection with 10 *C. difficile* spores; closed squares indicate infection with 10,000 *C. difficile* spores. Black triangles in FIG. 4J indicate an additional experimental arm in which mice were treated with vancomycin following *C. difficile* infection.

(6) FIG. 5 shows experimental conditions evaluated in Example 2, the results for which are presented in FIGS. 7-9. Composition E corresponds to a mixture of 17 bacterial strains (See e.g., Narushima et al., Gut Microbes 5: 3, 333-339). Composition I corresponds to a mixture of *Clostridium scindens*, *Pseudoflavonifractor capillosus*, and *Blautia hansenii*. "Abx" refers to treatment with any of the antibiotic regimens.

(7) FIG. 6 shows survival of mice over time post infection with *C. difficile* spores, according to the experimental conditions shown in FIG. 5. Mice losing >20% body weight of baseline were included in mortality numbers in survival curves.

(8) FIGS. 7A-7I show weight of the mice at various times post infection with *C. difficile* spores. Groups of mice received cefoperazone (Abx) treatment followed by the indicated composition, or no cefoperazone (no Abx), then were administered *C. difficile* spores. FIG. 7A shows weight of the mice that received no antibiotic treatment. FIG. 7B shows weight of the mice that received cefoperazone treatment. FIG. 7C shows weight of the mice that received cefoperazone treatment followed by vancomycin. FIG. 7D shows weight of the mice that received cefoperazone treatment followed by Composition I. FIG. 7E shows weight of the mice that received cefoperazone treatment followed by Composition E. FIG. 7F shows weight of the mice that received cefoperazone treatment followed by composition A. FIG. 7G shows weight of the mice that received cefoperazone treatment followed by composition B. FIG. 7H shows weight of the mice that received cefoperazone treatment followed by composition C. FIG. 7I shows weight of the mice that received cefoperazone treatment followed by composition D.

(9) FIGS. 8A-8C show the load of *C. difficile* in colony forming units (CFUs) in fecal pellets at various times post infection with *C. difficile*. FIG. 8A shows *C. difficile* CFU/g feces one-day post infection. FIG. 8B shows *C. difficile* CFU/g feces 3 days post infection. FIG. 8C shows *C. difficile* CFU/g feces 8 days post infection.

(10) FIG. 9 shows experimental conditions evaluated in Example 3, the results for which are presented in FIGS. 10-12.

(11) FIG. 10 shows survival of the mice over time post infection with *C. difficile* spores, according to the experimental conditions shown in FIG. 9. Mice losing >20% body weight of baseline were included in mortality numbers in survival curves.

(12) FIG. 11 shows weight of the mice at various times post infection with *C. difficile* spores.

(13) FIG. 12 shows the *C. difficile* burden in colony forming units (CFUs) in fecal pellets collected from mice 1, 3, and 8 days post infection with *C. difficile*.

(14) FIG. 13 shows the strains of Composition F. The genus-species notation indicates the closest species based on the sequence of the isolated strain.

(15) FIG. 14 shows the classification by *Clostridium* cluster of the strains in Composition F and their short-chain fatty acid producing abilities.

(16) FIG. 15 shows experimental conditions evaluated in Example 4, the results for which are presented in FIGS. 16-18. The dosing days are relative to *C. difficile* infection. FMT refers to Fecal Matter Transplant with fecal matter isolated from mice or from humans.

(17) FIG. 16 shows survival of the mice over time post infection with *C. difficile* spores, according to the experimental conditions shown in FIG. 15. Mice losing >20% body weight of baseline were included in mortality numbers in survival curves.

(18) FIGS. 17A-17H show weight of the mice at various times post infection with *C. difficile* spores. Groups of mice received cefoperazone (Abx) treatment followed by the indicated composition, then were administered *C. difficile* spores. FIG. 17A shows weight of the mice that received cefoperazone treatment. FIG. 17B shows weight of the mice that received cefoperazone treatment followed by FMT with fecal matter from a human. FIG. 17C shows weight of the mice that received cefoperazone treatment followed by FMT with fecal matter from a mouse. FIG. 17D shows weight of the mice that received cefoperazone treatment followed by Composition B on day -1. FIG. 17E shows weight of the mice that received cefoperazone treatment followed by Composition B on days -2 and -1. FIG. 17F shows weight of the mice that received cefoperazone treatment followed by Composition B on days -2, -1, 1, 2, and 3. FIG. 17G shows weight of the mice that received cefoperazone treatment followed by Composition F on day -1. FIG. 17H shows weight of the mice that received cefoperazone treatment followed by Composition F on days -2, -1, 1, 2, and 3.

(19) FIGS. 18A-18B show the load of *C. difficile* in colony forming units (CFUs) in fecal pellets at various times post infection with *C. difficile*. FIG. 18A shows *C. difficile* CFU/g feces 8 days post infection. FIG. 18B shows *C. difficile* CFU/g feces 17 days post infection.

(20) FIG. 19 shows the strains of Composition G. The genus-species notation indicates the closest species based on the sequence of the isolated strain.

(21) FIG. 20 shows experimental conditions evaluated in Example 5, the results for which are presented in FIGS. 21-23. Composition B1=Composition B with *Bacteroides*; Composition B2=Composition B with *Bacteroides* but without *Flavonifractor plautii*.

(22) FIG. 21 shows survival of the mice over time post infection with *C. difficile* spores, according to the experimental conditions shown in FIG. 20. Mice losing >20% body weight of baseline were included in mortality numbers in survival curves.

(23) FIGS. 22A-22J show weight of the mice at various times post infection with *C. difficile* spores. FIG. 22A shows weight of the mice that received vehicle control. FIG. 22B shows weight of the mice that received Composition F. FIG. 22C shows weight of the mice that received Composition G. FIG. 22D shows weight of the mice that received cefoperazone treatment followed by Composition B. FIG. 22E shows weight of the mice that received cefoperazone treatment followed by Composition B2 (=Composition B without *Flavonifractor plautii* and with added *Bacteroides*). FIG. 22F shows weight of the mice that received cefoperazone treatment followed by Composition B1 (=Composition B with *Bacteroides* added). FIG. 22G shows weight of the mice that received cefoperazone treatment followed by frozen Composition B. FIG. 22H shows weight of the mice that received cefoperazone treatment followed by ethanol treated human fecal samples. FIG. 22I shows weight of the mice that received cefoperazone treatment followed by ethanol treated Composition B. FIG. 22J shows weight of the mice that received cefoperazone treatment followed by Composition J.

(24) FIG. 23 shows the load of *C. difficile* in colony forming units (CFUs) in fecal pellets at various times post infection with *C. difficile*.

(25) FIG. 24 shows weight of the indicated groups of mice at various times post infection with *C. difficile* spores.

(26) FIG. 25 shows experimental conditions evaluated in Example 6, the results of which are presented in FIGS. 27-29.

(27) FIG. 26 shows the strains in Composition H (SEQ ID NO:14—VE202-13—*Anaerotruncus colihominis* (Cluster IV); SEQ ID NO:16—VE202-16—*Clostridium symbiosum* (Cluster XIVa); SEQ ID NO:21-189—*Clostridium innocuum* (Cluster XVII); SEQ ID NO:82—PE9—*Clostridium disporicum* (Cluster I); SEQ ID NO:81—PE5—*Clostridium bolteae* (Cluster XIVa); SEQ ID NO:80—VE202-18—*Erysipelatoclostridium ramosum* (Cluster XVIII)).

(28) FIGS. 27A and 27B shows survival and weight loss of the mice over time post infection with *C. difficile* spores, according to the experimental conditions shown in FIG. 25. Mice losing >20% body weight of baseline were included in mortality numbers in survival curves. FIG. 29A shows survival/mortality of mice that received the indicated treatment prior to *C. difficile* infection. FIG. 29B shows the weight over time of mice that received the indicated treatment prior to *C. difficile* infection.

(29) FIGS. 28A and 28B show results from the experimental conditions shown in FIG. 25. FIG. 28A shows survival/mortality of mice that received the indicated treatment prior to *C. difficile* infection. FIG. 28B shows the weight over time of mice that received the indicated treatment prior to *C. difficile* infection.

(30) FIGS. 29A and 29B show the *C. difficile* burden in CFU/gram feces collected from mice that received the indicated treatment prior to *C. difficile*. FIG. 29A shows *C. difficile* burden at one-day post *C. difficile* infection. FIG. 29B shows *C. difficile* burden at 4 days post *C. difficile* infection. FIG. 29C shows *C. difficile* burden at 19 days post *C. difficile* infection.

(31) FIG. 30 shows that Composition B reduced the amount of *C. difficile* Toxin B compared to no treatment controls: “2-1 (Cdiff)” and “2-4 (Cdiff)” and FMT. In addition, Composition B reduced the amount of *C. difficile* Toxin B compared to Composition B with additional spores.

(32) FIG. 31 shows Composition B reduced *C. difficile* growth in in vitro competition experiments. Cultures of *C. difficile* were incubated in the presence of *B. thetaiotaomicron*, *C. bifermentans*, or Composition B, or in the absence of a competing strain(s) (*C. diff* only). The quantity of *C. difficile* is presented as the percentage of the control (*C. diff* only).

(33) FIG. 32 shows that inoculation with Composition B induced the percentage of FoxP3+CD4+ cells (regulatory T cells) in the intestine of germ-free mice as compared to control mice (“GF”).

DETAILED DESCRIPTION OF THE INVENTION

(34) Disclosed herein are compositions comprising purified bacterial strains and pharmaceutical compositions and food

products containing such compositions and bacterial strains. Also disclosed are methods of treating a pathogenic infection, such as *Clostridium difficile* (*C. difficile*) infection, in a subject by administering said compositions to the subject.

(35) Various factors including antibiotic usage can induce dysbiosis of the gastrointestinal tract, which may allow for colonization by pathogenic microorganisms, such as *C. difficile*. Such colonization or pathogenic infection can lead to a variety of adverse effects in the subject including diarrhea, which is one of the primary symptoms characteristic of *C. difficile* infection (CDI). In the case of CDI, diarrhea is thought to be a result of *C. difficile* production of Toxin B (also referred to as cytotoxin TcdB), which results in opening of the tight junctions between intestinal epithelial cells, increasing vascular permeability, hemorrhage, and inflammation.

(36) The compositions described herein are effective in the treatment of *C. difficile* infection. As shown herein, the disclosed compositions are effective in suppressing the pathogenic effects of *C. difficile* infection. The compositions provided herein reduce the amount of *C. difficile* after infection and thereby provide an effective method for eliminating *C. difficile* from the body (e.g., the gut). The compositions provided herein induce the proliferation and/or accumulation of regulatory T cells (Tregs), for example when administered to a subject. Remarkably, the compositions disclosed herein have been found to reduce or inhibit production or activity of *C. difficile* Toxin B and thereby represent effective compositions for the treatment or prevention of CDI. The compositions disclosed herein have also been found to inhibit the growth and/or survival of *C. difficile*.

(37) The present disclosure provides compositions comprising purified bacterial strains that can be administered to subjects experiencing or having experienced a pathogenic infection to treat the infection. In some embodiments, the compositions may be administered to subjects who may be at risk for a pathogenic infection. Such subjects include subjects who previously had pathogenic infections, subjects who have been treated with antibiotics and subjects who will undergo a procedure that will put them at an increased risk for a pathogenic infection (e.g., surgery and/or hospitalization). In some embodiments, the pathogenic infection, is infection by a pathogen that is present predominantly in the gut or the intestine. In some embodiments, the pathogen that is present predominantly in the gut or the intestine is *Clostridium difficile*.

(38) In some embodiments, the one or more of the bacterial strains of the compositions provided herein colonize or recolonize the intestinal tract or parts of the intestinal tract (e.g., the colon or the cecum) of the subject. Such colonization or recolonization may also be referred to as grafting. In some embodiments, the one or more of the bacterial strains of the compositions recolonize the intestinal tract (e.g., the colon or the cecum) of the subject after the naturally present microbiome has been partially or completely removed, e.g., because of administration of antibiotics. In some embodiments, the one or more of the bacterial strains of the compositions colonize a dysbiotic gastrointestinal tract.

(39) In some embodiments, the one or more of the bacterial strains of the compositions can “outgrow” a pathogen, such as *C. difficile*. Thus, in some embodiments, if a pathogen (e.g., *C. difficile*) and one or more bacteria of compositions provided herein are both present in the intestinal tract (e.g., the colon or the cecum), the one or more bacteria of compositions provided herein grow faster (e.g., have a shorter doubling time) than the pathogen, thereby preventing the pathogen from accumulating in the intestinal tract (e.g., the colon or the cecum). In some embodiments, the faster growth results because the one or more bacteria of the compositions provided herein are better at grafting in the intestinal tract (e.g., the colon or the cecum). In some embodiments, the faster growth results because the one or more bacteria of the compositions provided herein are better at metabolizing nutrients present in the intestinal tract (e.g., the colon or the cecum). In some embodiments, the compositions of bacterial strains provided herein prevent or inhibit production of bacterial toxins by the pathogenic infection, or prevent or inhibit the cytopathic or cytotoxic effects of such bacterial toxins. In some embodiments, the bacterial strains of the compositions provided herein can treat pathogenic infections, because of the synergy between the bacterial strains. Thus, without being limiting, in some embodiments, the combination of the bacterial strains of the compositions provided herein act synergistically because the combination of the strains is particularly well-suited to use nutrients in the intestinal tract (e.g., the colon or the cecum), or instance through metabolic interactions, and/or because the combination is superior in grafting (e.g., by providing a favorable microenvironment).

(40) In some embodiments, a pathogenic infection such as *C. difficile* is treated because the combination of bacterial strains of the compositions provided herein is superior in the use of nutrients when compared to the pathogen such as *C. difficile*, thereby suppressing the growth of the pathogen such as *C. difficile*. In some embodiments, a pathogenic infection such as *C. difficile* is treated because the combination of bacterial strains of the compositions provided herein is superior in grafting when compared to the pathogen such as *C. difficile*, thereby suppressing the growth of the pathogen such as *C. difficile*. In some embodiments, a pathogenic infection such as *C. difficile* is treated because the combination of bacterial strains of the compositions provided herein is superior in the use of nutrients and in grafting when compared to the pathogen such as *C. difficile*, thereby suppressing the growth of the pathogen such as *C. difficile*. In some embodiments, a pathogenic infection such as *C. difficile* is treated because the combination of bacterial strains of the compositions provided herein inhibits the growth and/or survival of the pathogen such as *C. difficile*. In some embodiments, a pathogenic infection such as *C. difficile* is treated because the combination of bacterial strains of the compositions provided herein induces regulatory T cells (Tregs) in the subject that results in reduction or elimination of the pathogen such as *C. difficile*. In some embodiments, a pathogenic infection such as *C. difficile* is treated because the combination of bacterial strains of the compositions provided herein inhibits the growth and/or survival of the pathogen and induces regulatory T cells (Tregs) in the subject that results in reduction or elimination of the pathogen such as *C. difficile*.

(41) In some embodiments, the synergistic effect is provided by the capacity of the combination to colonize specific niches in the intestinal tract (e.g., the colon or the cecum). In some embodiments, the synergistic effect is provided by the capacity of the combination to metabolize specific nutrients. In some embodiments, the synergistic effect is provided by the capacity of the combination to provide specific metabolites to the environment. Such specific metabolites may suppress growth of the

pathogen and/or stimulate growth of non-pathogens. In some embodiments, the synergistic effect is provided by the capacity of the combination to provide short-chain fatty acids to the environment. In some embodiments, the synergistic effect is provided by the capacity of the combination to provide specific short-chain fatty acids to the environment. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce butyrate. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce acetate. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce lactate. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce propionate. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce succinate. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce multiple metabolites. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce multiple short-chain fatty acids. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce both butyrate and acetate. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce both butyrate and lactate. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce both butyrate and propionate. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce both butyrate and succinate. In some embodiments, the synergistic effect is provided by the capacity of the combination to produce butyrate, acetate and additional short-chain fatty acids.

(42) The bacterial strains used in the compositions provided herein generally are isolated from the microbiome of healthy individuals. In some embodiments, the compositions include strains originating from a single individual. In some embodiments, the compositions include strains originating from multiple individuals. In some embodiments, the bacterial strains are obtained from multiple individuals, isolated and grown up individually. The bacterial compositions that are grown up individually may subsequently be combined to provide the compositions of the disclosure. It should be appreciated that the origin of the bacterial strains of the compositions provided herein is not limited to the human microbiome from a healthy individual. In some embodiments, the bacterial strains originate from a human with a microbiome in dysbiosis. In some embodiments, the bacterial strains originate from non-human animals or the environment (e.g., soil or surface water). In some embodiments, the combinations of bacterial strains provided herein originate from multiple sources (e.g., human and non-human animals).

(43) In some embodiments, the bacteria of the compositions provided herein are anaerobic bacteria. In some embodiments, the bacteria of the compositions provided herein are obligate anaerobic bacteria. In some embodiments, the bacteria of the compositions provided herein are clostridia. Clostridia may be classified into phylogenetic clusters with other closely related strains and species. (See e.g., Rajilic-Stojanovic, M., and de Vos, W. M. *FEMS Microbiol Rev* 38, (2014) 996-1047). In general, clostridia are classified as belonging to a specific cluster based on their 16S rRNA (or 16S rDNA) nucleic acid sequence. Methods for determining the identity of specific bacterial species based on their 16S rRNA (or 16S rDNA) nucleic acid sequence are well known in the art (See e.g., *Jumpstart Consortium Human Microbiome Project Data Generation Working*, G. *PLoS One* (2012) 7, e39315).

(44) Provided herein are compositions comprising bacterial strains belonging to specific *Clostridium* clusters that have been found to be effective in treating and/or preventing pathogenic infection (e.g., *C. difficile* infection). In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster IV. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster XIVa. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster XVII. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster I. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster IX. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster XIVa and at least one of the bacterial strains belongs to *Clostridium* cluster XVII. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster IV and at least one of the bacterial strains belongs to *Clostridium* cluster XVII. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster IV, at least one of the bacterial strains belongs to *Clostridium* cluster XIVa, and at least one of the bacterial strains belongs to *Clostridium* cluster XVII.

(45) In some embodiments, the composition has at least twice as many bacterial strains that belong to *Clostridium* cluster XIVa when compared to the bacterial strains that belong to *Clostridium* cluster IV. In some embodiments, at least two of the bacterial strains of the composition belong to *Clostridium* cluster IV and at least five of the bacterial strains belong to *Clostridium* cluster XIVa. In some embodiments, the composition has at least twice as many bacterial strains that belong to *Clostridium* cluster XIVa when compared to the bacterial strains that belong to *Clostridium* cluster IV, and the composition has at least one strain that belongs to *Clostridium* cluster XVII. In some embodiments, at least two of the bacterial strains of the composition belong to *Clostridium* cluster IV, at least five of the bacterial strains belongs to *Clostridium* cluster XIVa, and at least one of the bacterial strains belongs to *Clostridium* cluster XVII.

(46) In some embodiments, the compositions provided herein do not include bacterial strains belonging to *Clostridium* cluster XVIII. In some embodiments, the compositions provided herein do not include bacterial strains belonging to *Clostridium* cluster XVI. In some embodiments, the compositions provided herein do not include bacterial strains belonging to *Clostridium* cluster XI. In some embodiments, the compositions provided herein do not include bacterial strains belonging to *Clostridium* cluster I.

(47) In one aspect, the disclosure provides bacterial strains comprising a 16S rDNA sequence with a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1-83 and 124-159. It should be appreciated that SEQ ID NOs: 1-83 and 124-159 may include both full length and partial 16S rDNA sequences.

(48) In one aspect, the disclosure provides compositions comprising a bacterial strain comprising a 16S rDNA sequence with

a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1-83 and 124-159. In one aspect, the disclosure provides compositions comprising as an active ingredient a bacterial strain comprising a 16S rDNA sequence with a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1-83 and 124-159. It should be appreciated that for all compositions provided herein, in some embodiments, the bacterial strain or the bacterial strains are the active ingredient of the composition.

(49) It should be appreciated that for all compositions provided herein, in some embodiments, the bacterial strains are purified. Thus, for example the disclosure provides purified bacterial strains comprising a 16S rDNA sequence with a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1-83 and 124-159. In addition, for example, the disclosure provides compositions comprising purified bacterial strains comprising a 16S rDNA sequence with a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 1-83 and 124-159. The bacterial strains disclosed herein originally may have been obtained and purified from the microbiota of one or more human individuals or obtained from sources other than the human microbiota, including soil and non-human microbiota. As provided herein, in some embodiments, bacteria isolated from the human microbiota, non-human microbiota, soil, or any alternative source are purified prior to use in the compositions and methods provided herein.

(50) In one aspect, the disclosure provides compositions comprising one or more bacterial strains, wherein the one or more bacterial strains comprise a 16S rDNA sequence with a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1-83 and 124-159. In one aspect, the disclosure provides compositions comprising one or more bacterial strains wherein the one or more bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs:1-83 and 124-159. As discussed previously, in some embodiments, the bacterial strains are purified. Thus, in one aspect, the disclosure provides compositions comprising one or more purified bacterial strains wherein the one or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs:1-83 and 124-159.

(51) In one aspect, the disclosure provides compositions comprising two or more purified bacterial strains wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs:1-83 and 124-159. As discussed above, in some embodiments, the bacterial strains are the active ingredient of the composition. Thus, in some embodiments, the disclosure provides compositions comprising as an active ingredient two or more purified bacterial strains wherein the two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs:1-83 and 124-159.

(52) In one aspect, the disclosure provides bacterial strains and combinations of bacterial strains that are homologous or have a high percent of homology with bacterial strains comprising 16S rDNA sequences selected from the group consisting of SEQ ID NOs:1-83 and 124-159. As discussed previously, in some embodiments, the bacterial strains are purified. The bacterial strains disclosed herein that have a 16S rDNA sequence with a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1-83 and 124-159 have a high percent of homology (e.g., greater than 90%) with 16S rDNA sequences of bacterial strains that have been described in various databases (See e.g., the National Center for Biotechnology Information). Table 1 and Table 3 provides the closest known species by homology when the 16S rDNA sequences comprising SEQ ID NOs:1-83 and 124-159 are compared to 16S rDNA sequences of bacterial species available in public databases. By way of example, the bacterial strain comprising a 16S rDNA sequence with SEQ ID NO:1 (also referred to herein as “Strain 71”) disclosed herein has the highest homology with a bacterial strain of the species *Blautia wexlerae* as defined by Accession #NR_044054 (having 16S rDNA sequence SEQ ID NO:94). While the bacterial strain with SEQ ID NO:1 has homology with other published bacterial strains as well, the highest homology is with a bacterial strain of the species *Blautia wexlerae* as defined by Accession #NR_044054. In this particular example the homology of SEQ ID NO:1 is 96.6% with SEQ ID NO:94 (corresponding to *Blautia wexlerae*). It should be appreciated that multiple bacterial strains disclosed herein may have the highest homology with the same species. (e.g., both SEQ ID NO:4 and SEQ ID NO:5 have the highest homology with a 16S rDNA sequence of a strain of the species *Blautia hansenii*).

(53) It should further be appreciated that the bacterial strains disclosed herein that have a 16S rDNA sequence with a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1-83 and 124-159, are also homologous to other strains based on their whole genome sequence, or subset of their whole genome sequence. Homologies based on whole genome analysis are provided in Table 2 and Table 3.

(54) In one aspect, the disclosure provides compositions comprising one or more bacterial strains wherein the one or more bacterial strains are of species selected from the group consisting of *Clostridium hathewayi*, *Blautia hansenii*, *Blautia producta*, *Blautia producta* ATCC 27340, *Clostridium bacterium* UC5.1-1D4, *Blautia coccoides*, *Eubacterium contortum*, *Eubacterium fissicatena*, *Sellimona intestinalis*, *Dracourtella massiliensis*, *Dracourtella massiliensis* GD1, *Ruminococcus torques*, *Anaerostipes caccae*, *Clostridium scindens*, *Marvinbryanta formatexigens*, *Eisenbergiella tayi*, *Flavinofractor plautii*, *Clostridium orbiscindens* 1_3_50 AFAA, *Lachnospiraceae bacterium* 7_1_58 FAA, *Subdoligranulum*, *Anaerotruncus colihominis*, *Anaerotruncus colihominis* DSM 17241, *Clostridium symbiosum*, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae*, *Clostridium bolteae* 90A9, *Dorea longicatena*, *Dorea longicatena* CAG:42, *Clostridium innocuum*, *Erysipelotrichaceae bacterium* 21-3, *Blautia wexlerae*, *Clostridium disporicum*, *Erysipelatoclostridium ramosum*, *Pseudoflavinofractor capillosus*, *Turicibacter sanguinis*, *Lactobacillus mucosae*, *Ruminococcus obeum*, *Megasphaera elsdenii*, *Acidaminococcus fermentans*, *Acidaminococcus intestine*, *Ruminococcus faecis*, *Bacteroides cellulosilyticus*, *Anaerostipes hadrus*, *Eubacterium rectale*, *Ruminococcus champanellensis*, *Ruminococcus albus*, *Bifidobacterium bifidum*, *Blautia luti*, *Roseburia faecis*, *Fusicatenibacter saccharivorans*, *Roseburia faecis*, *Blautia faecis*, *Dorea formicigenerans* and *Bacteroides ovatus*.

(55) In some embodiments, the disclosure provides compositions comprising two or more bacterial strains, wherein the two or more bacterial strains are of species selected from the group consisting of *Clostridium hathewayi*, *Blautia hansenii*, *Blautia producta*, *Blautia producta* ATCC 27340, *Clostridium bacterium* UC5.1-1D4, *Blautia coccoides*, *Eubacterium contortum*, *Eubacterium fissicatena*, *Sellimona intestinalis*, *Dracourtella massiliensis*, *Dracourtella massiliensis* GD1, *Ruminococcus torques*, *Anaerostipes caccae*, *Clostridium scindens*, *Marvinbryanta formatexigens*, *Eisenbergiella tayi*, *Flavinofractor plautii*, *Clostridium orbiscindens* 1_3_50 AFAA, *Lachnospiraceae bacterium* 7_1_58 FAA, *Subdoligranulum*, *Anaerotruncus colihominis*, *Anaerotruncus colihominis* DSM 17241, *Clostridium symbiosum*, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae*, *Clostridium bolteae* 90A9, *Dorea longicatena*, *Dorea longicatena* CAG:42, *Clostridium innocuum*, *Erysipelotrichaceae_bacterium_21-3*, *Blautia wexlerae*, *Clostridium disporicum*, *Erysipelatoclostridium ramosum*, *Pseudoflavinofractor capillosus*, *Turicibacter sanguinis*, *Lactobacillus mucosae*, *Ruminococcus obeum*, *Megasphaera elsdenii*, *Acidaminococcus fermentans*, *Acidaminococcus intestine*, *Ruminococcus faecis*, *Bacteroides cellulosilyticus*, *Anaerostipes hadrus*, *Eubacterium rectale*, *Ruminococcus champanellensis*, *Ruminococcus albus*, *Bifidobacterium bifidum*, *Blautia luti*, *Roseburia faecis*, *Fusicatenibacter saccharivorans*, *Roseburia faecis*, *Blautia faecis*, *Dorea formicigenerans* and *Bacteroides ovatus*.

(56) It should be appreciated that the compositions may include multiple strains of a particular species. Thus, for illustration, a non-limiting example of the compositions disclosed herein, comprises one strain of *Clostridium hathewayi* and two strains of *Blautia hansenii*.

(57) The invention also encompasses compositions comprising bacterial strains that are close in homology to and/or fall within the species *Clostridium hathewayi*, *Blautia hansenii*, *Blautia producta*, *Blautia producta* ATCC 27340, *Clostridia* bacteria UC5.1-1D4, *Blautia coccoides*, *Eubacterium contortum*, *Eubacterium fissicatena*, *Sellimona intestinalis*, *Dracourtella massiliensis*, *Dracourtella massiliensis* GD1, *Ruminococcus torques*, *Anaerostipes caccae*, *Clostridium scindens*, *Marvinbryanta formatexigens*, *Eisenbergiella tayi*, *Flavinofractor plautii*, *Clostridium orbiscindens* 1_3_50 AFAA, *Lachnospiraceae bacterium* 7_1_58 FAA, *Subdoligranulum*, *Anaerotruncus colihominis*, *Anaerotruncus colihominis* DSM 17241, *Clostridium symbiosum*, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae*, *Clostridium bolteae* 90A9, *Dorea longicatena*, *Dorea longicatena* CAG:42, *Clostridium innocuum*, *Erysipelotrichaceae_bacterium_21-3*, *Blautia wexlerae*, *Clostridium disporicum*, *Erysipelatoclostridium ramosum*, *Pseudoflavinofractor capillosus*, *Turicibacter sanguinis*, *Lactobacillus mucosae*, *Ruminococcus obeum*, *Megasphaera elsdenii*, *Acidaminococcus fermentans*, *Acidaminococcus intestine*, *Ruminococcus faecis*, *Bacteroides cellulosilyticus*, *Anaerostipes hadrus*, *Eubacterium rectale*, *Ruminococcus champanellensis*, *Ruminococcus albus*, *Bifidobacterium bifidum*, *Blautia luti*, *Roseburia faecis*, *Fusicatenibacter saccharivorans*, *Roseburia faecis*, *Blautia faecis*, *Dorea formicigenerans* and *Bacteroides ovatus*. Thus, in one embodiment, the compositions of the disclosure include one or more bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs: 84-123. In some embodiments, the compositions of the disclosure include two or more bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs:84-123.

(58) In one aspect, the compositions of the disclosure include two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs:1-23 and 124-159. In some embodiments, the compositions of the disclosure include two or more bacterial strains of species selected from the group consisting of *Clostridium hathewayi*, *Blautia hansenii*, *Blautia producta*, *Blautia producta* ATCC 27340, *Clostridia* bacteria UC5.1-1D4, *Blautia coccoides*, *Eubacterium contortum*, *Eubacterium fissicatena*, *Sellimona intestinalis*, *Dracourtella massiliensis*, *Dracourtella massiliensis* GD1, *Ruminococcus torques*, *Anaerostipes caccae*, *Clostridium scindens*, *Marvinbryanta formatexigens*, *Eisenbergiella tayi*, *Flavinofractor plautii*, *Clostridium orbiscindens* 1_3_50 AFAA, *Lachnospiraceae bacterium* 7_1_58 FAA, *Subdoligranulum*, *Anaerotruncus colihominis*, *Anaerotruncus colihominis* DSM 17241, *Clostridium symbiosum*, *Clostridium symbiosum* WAL-14163, *Clostridium bolteae*, *Clostridium bolteae* 90A9, *Dorea longicatena*, *Dorea longicatena* CAG:42, *Clostridium innocuum*, *Erysipelotrichaceae_bacterium_21-3*, *Blautia wexlerae*, *Turicibacter sanguinis*, *Lactobacillus mucosae*, and *Bacteroides ovatus*. In some embodiments, the compositions of the disclosure include two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:109, SEQ ID NO:110, SEQ ID NO:121, and SEQ ID NO:122.

(59) In one aspect, the disclosure provides Composition A (See e.g., FIG. 1, Table A). As shown in FIG. 1, Composition A contains bacterial strains that comprise 16S rDNA sequences with nucleic acid sequences: SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23. In some embodiments, the disclosure provides compositions with two or more purified bacterial strains that comprise 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23. In some embodiments, the disclosure provides compositions with five or more purified bacterial strains that comprise 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23. In some embodiments, the disclosure provides compositions with at least ten purified bacterial strains, wherein the bacterial strains comprise 16S rDNA sequences with nucleic acid sequences SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23, respectively. In some embodiments, the disclosure provides a composition consisting of ten purified bacterial strains, wherein

the bacterial strains comprise 16S rDNA sequences with nucleic acid sequences SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23, respectively. In some embodiments, the disclosure provides a composition essentially consisting of ten purified bacterial strains, wherein the bacterial strains comprise 16S rDNA sequences with nucleic acid sequences SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23, respectively. As used herein, essentially consisting of refers to a composition that includes no additional bacterial strains.

(60) In some embodiments, the disclosure provides compositions with bacterial strains that comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of: SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23. In some embodiments, the disclosure provides compositions with two or more purified bacterial strains that comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23. In some embodiments, the disclosure provides compositions with five or more purified bacterial strains that comprise 16S rDNA having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23. In some embodiments, the disclosure provides compositions with at least ten purified bacterial strains, wherein the bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23, respectively. In some embodiments, the disclosure provides a composition consisting of ten purified bacterial strains, wherein the bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23, respectively. In some embodiments, the disclosure provides a composition essentially consisting of ten purified bacterial strains, wherein the bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23, respectively.

(61) The bacterial strains in Composition A are related to the following bacterial species: *Clostridium hathewayi*, *Blautia hansenii*, *Blautia producta*, *Blautia coccooides*, *Eubacterium contortum*, *Eubacterium fissicatena*, *Anaerostipes caccae*, *Clostridium scindens*, *Marvinbryanta formatexigens*, and *Eisenbergiella tayi* (See e.g., Table 1). It should be appreciated that multiple bacterial strains of the compositions disclosed herein can have the same related bacterial species. For instance, the bacterial strains having 16S rDNA sequences with nucleic acid sequences SEQ ID NO:4, SEQ ID NO:5 and SEQ ID NO:7 all have *Blautia hansenii* as related species. In some embodiments, the disclosure provides compositions with two or more bacteria of species selected from the group consisting of *Clostridium hathewayi*, *Blautia hansenii*, *Blautia producta*, *Blautia coccooides*, *Eubacterium contortum*, *Eubacterium fissicatena*, *Anaerostipes caccae*, *Clostridium scindens*, *Marvinbryanta formatexigens*, and *Eisenbergiella tayi*. In some embodiments, the disclosure provides compositions with two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:87, SEQ ID NO:88, SEQ ID NO:89, SEQ ID NO:99, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:106, SEQ ID NO:108, SEQ ID NO:109, and SEQ ID NO:121.

(62) In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5; SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:13.

(63) Each of the bacterial strains of Composition A are BaiCD⁺, meaning that the bacterial strains encode, or are predicted to encode, the bile inducible operon gene BaiCD and/or a protein with stereospecific NAD(H)-dependent 3-oxo- Δ .sup.4-cholenoic acid oxidoreductase activity. The BaiCD status of a bacterial strain can be determined for instance by PCR (See

e.g., Wells et al. *Clin Chim Acta* (2003) May; 331(1-2):127-34). Furthermore, each of the strains of Composition A are classified as belonging to *Clostridium* cluster XIVa. In some embodiments, the disclosure provides compositions comprising two or more bacterial strains, wherein the bacterial strains are BaiCD⁺ strains. In some embodiments, the disclosure provides compositions comprising two or more bacterial strains, wherein the bacterial strains are BaiCD⁺ and belong to *Clostridium* cluster XIVa. In some embodiments of the compositions comprising two or more bacterial strains that are BaiCD⁺ strains and that belong to *Clostridium* cluster XIVa, the compositions do not include bacterial strains that belong to *Clostridium* cluster IV.

(64) In some embodiments, the disclosure provides compositions with two or more purified bacterial strains that comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:23, wherein all the bacterial strains belong to *Clostridium* cluster XIVa.

(65) TABLE-US-00001 TABLE A Composition A SEQ_03 - 5 - *Clostridium_hathewayi* (XIVa)* SEQ_04 - 7 - *Blautia_hansenii* (XIVa)* SEQ_05 - 10 - *Blautia_hansenii* (XIVa)* SEQ_07 - 59 - *Blautia_producta/Blautia_coccoides* (XIVa) SEQ_08 - 79 - *Blautia_hansenii* (XIVa)* SEQ_09 - VE202-21 - *Eubacterium_contortum/Eubacterium_fissicatena* (XIVa)* SEQ_11 - VE202-9 - *Anaerostipes_caccae* (XIVa)* SEQ_12 - VE202-26 - *Clostridium_scindens* (XIVa)* SEQ_13 - 136 - *Marvinbryantia_formatexigens* (XIVa)* SEQ_23 - VE202-29 - *Eisenbergiella_tayi* (XIVa)* *= BaiCD^{sup.}+

(66) In one aspect, the disclosure provides Composition B (See e.g., FIG. 1, Table B). As shown in FIG. 1, Composition B contains bacterial strains that comprise 16S rDNA sequences with nucleic acid sequences: SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NOs: 124-159. In some embodiments, the compositions include two or more purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NOs: 124-159. In some embodiments, the compositions include two or more purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21. In some embodiments, the compositions include two or more purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NOs: 124-159. In some embodiments, the compositions include two or more purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO: 124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ ID NO:152 and SEQ ID NO:157.

(67) In some embodiments, the compositions include five or more purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NOs: 124-159. In some embodiments, the compositions include five or more purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21. In some embodiments, the compositions include five or more purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NOs: 124-159. In some embodiments, the compositions include five or more purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO: 124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ ID NO:152 and SEQ ID NO:157.

(68) In some embodiments, the compositions include at least eight purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NOs: 124-159, respectively. In some embodiments, the compositions include at least eight purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21, respectively. In some embodiments, the compositions include at least eight purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NOs: 124-159, respectively. In some embodiments, the compositions include at least eight purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO: 124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ ID NO:152 and SEQ ID NO:157.

(69) In some embodiments, the composition consists of at least eight purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NOs: 124-159, respectively. In some embodiments, the composition consists of at least eight purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21, respectively. In some embodiments, the composition consists of eight purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NOs:124-159, respectively. In some embodiments, the compositions consists of at least eight purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences SEQ ID NO: 124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ ID NO:152 and SEQ ID NO:157.

(70) In some embodiments, the composition essentially consists of at least eight purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NOs: 124-159,

SEQ ID NO:22, and SEQ ID NOs: 124-159, respectively. In some embodiments, the compositions include at least ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NO:22, respectively. In some embodiments, the compositions include at least ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO: 18, SEQ ID NO: 22, and SEQ ID NOs: 124-159.

(96) In some embodiments, the composition consists of ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, and SEQ ID NO: 124-159, respectively. In some embodiments, the composition consists of ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NO:22, respectively. In some embodiments, the composition consists of ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:18, SEQ ID NO:22, SEQ ID NOs: 124-159, respectively.

(97) In some embodiments, the composition essentially consists of ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, and SEQ ID NO: 124-159, respectively. In some embodiments, the composition essentially consists of ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NO:22, respectively. In some embodiments, the composition essentially consists of ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:18, SEQ ID NO:22, and SEQ ID NO: 124-159, respectively.

(98) The bacterial strains in Composition B are related to the following bacterial species: *Flavinofractor plautii*, *Lachnospiraceae, bacterium 7_1_58* FAA, *Subdoligranulum Anaerotruncus colihominis*, *Eubacterium fissicatena*, *Ruminococcus torques Clostridium symbiosum*, *Clostridium bolteae*, *Dorea longicatena*, *Blautia producta*, *Clostridium innocuum*, and *Erysipelotrichaceae bacterium 21-3* (See e.g., Table 2).

(99) Selected strains were subjected to whole genome sequencing using a PacBio Biosciences platform (Menlo Park, CA) and sequences were assembled into whole genomes (Table 3). The 16S rDNA sequences were identified using Prokka and Barrnap. It was found that several strains contained more than one 16S sequence. All identified 16S rRNA gene nucleotide sequences for each strain were then clustered at 97% identity using the usearch (v 5.2.236) algorithm and the cluster seed sequence was selected as the representative sequence for each Composition B strain (The Consensus 16S sequence: column labeled “*Consensus SEQ ID # of 16S region as determined by WGS” in Table 3). Table 3 provides identification of the indicated strains included in Composition B based on Sanger sequencing of the 16S region as well as on whole genome sequencing (WGS). The closest species of the bacterial strains were identified both by comparison to a 16S database (column labeled: “Closest species based on Consensus SEQ ID # of 16S region as compared with 16S database”) and to whole genome databases (column labeled: “Closest species based on WGS compared versus WG databases”).

(100) Based on identification of 16S sequences through whole genome sequencing, and by comparing these sequences with 16S databases, the bacterial strains in Composition B are related to the following bacterial species: *Clostridium bolteae*, *Anaerotruncus colihominis*, *Dracourtella massiliensis*, *Clostridium symbiosum* *Blautia producta*, *Dorea longicatena* *Clostridium innocuum* and *Flavinofractor plautii* (see, e.g., Table 3).

(101) Based on whole genome sequencing and comparing of the whole genome to whole genome databases, the bacterial strains in Composition B are most closely related to the following bacterial species: *Clostridium bolteae* 90A9, *Anaerotruncus colihominis* DSM 17241, *Dracourtella massiliensis* GD1, *Clostridium symbiosum* WAL-14163, *Clostridium bacterium* UC5.1-1D4, *Dorea longicatena* CAG:42, *Erysipelotrichaceae bacterium 21_3*, and *Clostridium orbiscindens 1_3_50* AFAA (see, e.g., Table 3).

(102) It should be appreciated that multiple strains of the compositions disclosed herein can have the same related bacterial species. For instance, the bacterial strains comprising 16S rDNA sequences with nucleic acid sequences SEQ ID NO 18, SEQ ID NO:20 and SEQ ID NO:22 all have *Dorea longicatena* as related bacterial species. In some embodiments, the disclosure provides compositions with two or more bacteria selected from the group consisting of *Flavinofractor plautii*, *Lachnospiraceae, bacterium 7_1_58* FAA, *Subdoligranulum Anaerotruncus colihominis*, *Eubacterium fissicatena*, *Ruminococcus torques Clostridium symbiosum*, *Clostridium bolteae*, *Dorea longicatena*, *Blautia producta*, *Clostridium innocuum* and *Erysipelotrichaceae bacterium 21-3*. In some embodiments, the disclosure provides compositions with two or more bacteria selected from the group consisting of *Flavinofractor plautii*, *Anaerotruncus colihominis*, *Eubacterium fissicatena*, *Clostridium symbiosum*, *Clostridium bolteae*, *Dorea longicatena*, *Blautia producta*, and *Clostridium innocuum*. In some embodiments, the disclosure provides compositions that include two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:102, SEQ ID NO:106, SEQ ID NO:110, and SEQ ID NO:122.

(103) In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10,

consisting of SEQ ID NO: 18, SEQ ID NOs: 124-145, and SEQ ID NOs: 152-159.

(109) In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NOs: 157-159, and SEQ ID NOs:141-156. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NO:22. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:18, SEQ ID NO:22, SEQ ID NOs: 157-159, and SEQ ID NO:141-156.

(110) Each of the bacteria of Composition B are BaiCD⁻ strains, meaning that the strains do not encode and/or are not predicted to encode the bile inducible operon gene baiCD and/or a protein with stereospecific NAD(H)-dependent 3-oxo- Δ .sup.4-cholenoic acid oxidoreductase activity. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the bacteria are BaiCD⁻ strains. The strains of Composition B are classified as belonging to *Clostridium* clusters IV, XIVa, and XVII. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and belong to *Clostridium* clusters IV, XIVa, or XVII. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and belong to *Clostridium* clusters IV or XVII. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and belong to *Clostridium* clusters XIVa or XVII. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and belong to *Clostridium* cluster IV. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and belong to *Clostridium* cluster XIVa. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and belong to *Clostridium* clusters XVII. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and belong to *Clostridium* clusters IV, XIVa, and XVII and do not belong to *Clostridium* clusters XVI or XVIII.

(111) In some embodiments, the disclosure provides two or more bacterial strains wherein the bacterial strains are spore forming bacterial strains. In some embodiments, the disclosure provides two or more bacterial strains wherein the bacteria are spore formers and wherein the bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:21, SEQ ID NOs 124-140, and SEQ ID NO: 152-156. In some embodiments, the disclosure provides two or more bacterial strains wherein the bacteria are spore formers and wherein the bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, and SEQ ID NO:21. In some embodiments, the disclosure provides two or more bacterial strains wherein the bacteria are spore formers and wherein the bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs: 124-140, and SEQ ID NOs: 152-156.

(112) In some embodiments, the disclosure provides two or more bacterial strains wherein the bacteria include both spore formers and non-spore formers. In some embodiments, the disclosure provides two or more bacterial strains wherein the bacteria include both spore formers and non-spore formers, and wherein the spore forming bacterial strains comprise two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:21, SEQ ID NOs: 124-140, and SEQ ID NOs: 152-156. In some embodiments, the disclosure provides two or more bacterial strains wherein the bacteria include both spore formers and non-spore formers, and wherein the spore forming bacterial strains comprise two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:17, and SEQ ID NO:21. In some embodiments, the disclosure provides two or more bacterial strains wherein the bacteria include both spore formers and non-spore formers, and wherein the spore forming bacterial strains comprise two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs: 124-140 and SEQ ID NO: 152-156.

(113) TABLE-US-00002 TABLE B Composition B SEQ_10 - 211 - *Flavonifractor plautii* (IV) SEQ_14 - VE202-13 - *Anaerotruncus colihominis* (IV) SEQ_15 - VE202-14 - *Eubacterium fissicatena* (XIVa) SEQ_16 - VE202-16 - *Clostridium symbiosum* (XIVa) SEQ_17 - VE202-7 - *Clostridium bolteae* (XIVa) SEQ_19 - 16 - *Blautia producta* (XIVa) SEQ_20 - 170 - *Dorea longicatena* (XIVa) SEQ_21 - 189 - *Clostridium innocuum* (XVII)

(114) In some embodiments, the compositions include one or more bacterial species from the *Bacteroides* genus. In some embodiments, the compositions include one or more bacterial species selected from the group consisting of *B. acidifaciens*, *B. caccae*, *B. coprocola*, *B. coprosuis*, *B. eggerthii*, *B. finegoldii*, *B. fragilis*, *B. helcogenes*, *B. intestinalis*, *B. massiliensis*, *B. nordii*, *B. ovatus*, *B. thetaiotaomicron*, *B. vulgatus*, *B. plebeius*, *B. uniformis*, *B. salyersai*, *B. pyogenes*, *B. goldsteinii*, *B. dorei*, and *B. johnsonii*. In some embodiments, the compositions include *Bacteroides ovatus*. In some embodiments, the *Bacteroides ovatus* has a 16S rDNA sequence comprising SEQ ID NO:83. In some embodiments, the *Bacteroides ovatus* has a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence comprising SEQ ID NO:83. In some embodiments, the *Bacteroides ovatus* has a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence comprising SEQ ID NO:101.

(115) While not being limited to a specific mechanism it is thought that the inclusion of a *Bacteroides* species in the bacterial compositions disclosed herein increases the ability to sense and adapt to nutrient availability or influence the host immune system so that it becomes more effective in fighting pathogens (e.g., *C. difficile*). In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO: 124-159, and SEQ ID NO:83. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NO:83. (Composition B1, See e.g., Table B1). In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs: 124-159, and SEQ ID NO: 83.

(116) In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO: 124-145, SEQ ID NO: 152-159, and SEQ ID NO:83. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, and SEQ ID NO:83. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs: 124-145, SEQ ID NO: 152-159, and SEQ ID NO: 83.

(117) In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO: 124-159, and SEQ ID NO:83. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22 and SEQ ID NO:83. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs: 124-159, SEQ ID NO: 18, SEQ ID NO: 22, and SEQ ID NO: 83.

(118) In some embodiments, the composition comprises two or more purified bacterial strains comprising 16s rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:93, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:106, SEQ ID NO:110, and SEQ ID NO:122. It should also be appreciated that in some embodiments, the compositions disclosed herein do not include bacterial species from the *Bacteroides* genus.

(119) TABLE-US-00003 TABLE B1 Composition B1 SEQ_10 - 211 - *Flavonifractor plautii* (IV) SEQ_14 - VE202-13 - *Anaerotruncus colihominis* (IV) SEQ_15 - VE202-14 - *Eubacterium fissicatena* (XIVa) SEQ_16 - VE202-16 - *Clostridium symbiosum* (XIVa) SEQ_17 - VE202-7 - *Clostridium bolteae* (XIVa) SEQ_20 - 170 - *Dorea longicatena* (XIVa) SEQ_19 - 16 - *Blautia producta* (XIVa) SEQ_21 - 189 - *Clostridium innocuum* (XVII) SEQ_83 *Bacteroides ovatus*

(120) In some embodiments, the compositions disclosed herein do not include *Clostridium orbiscindens* 1_3_50 AFAA, *Flavonifractor plautii*, *Subdoligranulum* or *Lachnospiraceae bacterium* 7_1_58 FAA. In some embodiments, the compositions disclosed herein do not include *Clostridium orbiscindens* 1_3_50 AFAA. In some embodiments, the compositions disclosed herein do not include *Flavonifractor plautii*. In some embodiments, the compositions disclosed herein do not include *Subdoligranulum*. In some embodiments, the compositions disclosed herein do not include *Lachnospiraceae bacterium* 7_1_58 FAA.

(121) In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO: 15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NO: 124-156, wherein the composition does not include a bacterial strain comprising a 16s rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NO:10 and SEQ ID NOs: 157-159. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO: 15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21, wherein the composition does not include a bacterial strain comprising a 16s rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NO:10. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NOs: 124-156, wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NOs:157-159.

(122) In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:21, SEQ ID NO:22, and SEQ ID NOs: 124-146 and SEQ ID NO: 152-156, wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence

SEQ ID NO:83, and SEQ ID NO: 124-156, wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NO:10 and SEQ ID NOs: 157-159. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22 and SEQ ID NO:83, wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NO:10. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:22, SEQ ID NO:83, and SEQ ID NO: 124-156, wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NOs: 157-159.

(129) In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:101, SEQ ID NO:102, SEQ ID NO:106, SEQ ID NO:110, and SEQ ID NO:122, wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NO:93.

(130) TABLE-US-00004 TABLE B2 Composition B2 SEQ_14 - VE202-13 - *Anaerotruncus colihominis* (IV) SEQ_15 - VE202-14 - *Eubacterium fissicatena* (XIVa) SEQ_16 - VE202-16 - *Clostridium symbiosum* (XIVa) SEQ_17 - VE202-7 - *Clostridium bolteae* (XIVa) SEQ_20 - 170 - *Dorea longicatena* (XIVa) SEQ_19 - 16 - *Blautia producta* (XIVa) SEQ_21 - 189 - *Clostridium innocuum* (XVII) SEQ_83 *Bacteroides ovatus*

(131) In one aspect, the disclosure provides Composition C (See e.g., FIG. 1, Table C). As shown in FIG. 1, Composition C contains bacteria that have the following 16S rDNA sequences: SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16. In some embodiments, the disclosure provides compositions with two or more purified bacterial strains that have 16S rDNA sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16. In some embodiments, the compositions include four or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16. In some embodiments, the compositions include at least ten purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16. In some embodiments, the composition consists of ten purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16, respectively. In some embodiments, the composition consists essentially of ten purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16, respectively.

(132) In some embodiments, the compositions include two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16. In some embodiments, the compositions include four or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16. In some embodiments, the compositions include at least ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16. In some embodiments, the composition consists of ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16, respectively. In some embodiments, the composition essentially consists of ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16, respectively.

(133) The bacterial strains in Composition C are related to the following species: *Clostridium scindens*, *Clostridium hathewayi*, *Blautia hansenii*, *Blautia wexlerae*, *Blautia producta*, *Blautia coccoides*, *Dorea longicatena*, *Clostridium innocuum*, *Flavonifractor plautii*, *Lachnospiraceae bacterium 7_1_58* FAA, *Subdoligranulum*, *Anaerotruncus colihominis*, and *Clostridium symbiosum*. In some embodiments, the disclosure provides compositions with two or more bacterial strains of species selected from the group consisting of *Clostridium scindens*, *Clostridium hathewayi*, *Blautia hansenii*, *Blautia wexlerae*, *Blautia producta*, *Blautia coccoides*, *Dorea longicatena*, *Clostridium innocuum*, *Flavonifractor plautii*, *Lachnospiraceae bacterium 7_1_58* FAA, *Subdoligranulum*, *Anaerotruncus colihominis*, and *Clostridium symbiosum*. In some embodiments, the disclosure provides compositions that include two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID

SEQ ID NO:87, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:106, and SEQ ID NO:122.

(134) In some embodiments, the compositions disclosed herein do not include *Flavinofractor plautii*, *Subdoligranulum* or *Lachnospiraceae bacterium 7_1_58* FAA. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:14, and SEQ ID NO:16, wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NO:10. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:87, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:103, SEQ ID NO:105, SEQ ID NO:106, and SEQ ID NO:122, wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NO:93.

(135) The strains of Composition C include both BaiCD^{sup}.+ strains and BaiCD⁻ strains. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein one or more bacteria are BaiCD⁺ strains and one or more bacteria are BaiCD⁻ strains. In some embodiments of the one or more bacteria that are BaiCD⁺ strains are selected from bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, and SEQ ID NO:7. In some embodiments the one or more bacteria that are BaiCD⁻ strains are selected from bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:14, and SEQ ID NO:16. In some embodiments of the one or more bacteria that are BaiCD⁺ strains are selected from the bacterial species *Clostridium scindens*, *Clostridium hathewayi*, *Blautia hansenii*, *Blautia wexlerae*, *Blautia producta*, and *Blautia coccoides*. In some embodiments of the one or more bacteria that are BaiCD⁻ strains are selected from the bacterial species *Dorea longicatena*, *Clostridium innocuum*, *Flavinofractor plautii*, or *Lachnospiraceae bacterium 7_1_58* FAA, *Anaerotruncus colihominis*, and *Clostridium symbiosum*. The clostridial strains of Composition C are classified as belonging to *Clostridium* clusters IV, XIVa, and XVII. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and BaiCD⁺ strains and belong to *Clostridium* clusters IV, XIVa, or XVII. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and BaiCD⁺ strains and belong to *Clostridium* clusters XIVa or XVII. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and BaiCD⁺ strains and belong to *Clostridium* clusters IV or XIVa.

(136) TABLE-US-00005 TABLE C Composition C SEQ_12 - VE202-26 - *Clostridium scindens* (XIVa)* SEQ_03 - 5 - *Clostridium hathewayi* (XIVa)* SEQ_05 - 10 - *Blautia hansenii* (XIVa)* SEQ_01 - 71 - *Blautia wexlerae* (XIVa)* SEQ_07 - 59 - *Blautia producta/Blautia coccoides* (XIVa)* SEQ_18 - 148 - *Dorea longicatena* (XIVa) SEQ_21 - 189 - *Clostridium innocuum* (XVII) SEQ_10 - 211 - *Flavinofractor plautii* (IV) SEQ_14 - VE202-13 - *Anaerotruncus colihominis* (IV) SEQ_16 - VE202-16 - *Clostridium symbiosum* (XIVa) *= BaiCD⁺

(137) In one aspect, the disclosure provides Composition D (See e.g., FIG. 1, Table D). As shown in FIG. 1, Composition D contains bacteria that have the following 16S rDNA sequences: SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:2, and SEQ ID NO:6. In some embodiments, the disclosure provides compositions with two or more purified bacterial strains that have 16S rDNA sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:2, and SEQ ID NO:6. In some embodiments, the disclosure provides compositions that include three or more purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:2, and SEQ ID NO:6. In some embodiments, the disclosure provides compositions that include at least ten purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:2, and SEQ ID NO:6. In some embodiments, the disclosure provides a composition that consists of ten purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:2, and SEQ ID NO:6, respectively. In some embodiments, the disclosure provides a composition that consists essentially of ten purified bacterial strains comprising 16S rDNA sequences with nucleic acid sequences SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:2, and SEQ ID NO:6, respectively. In some embodiments, the disclosure provides compositions that include two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:2, and SEQ ID NO:6. In some embodiments, the disclosure provides compositions that include three or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:2, and SEQ ID NO:6. In some embodiments, the disclosure provides compositions that include at least ten more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID

NO:21, SEQ ID NO:10, SEQ ID NO:2, and SEQ ID NO:6. In some embodiments, the disclosure provides a composition that consists of ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:2, and SEQ ID NO:6, respectively. In some embodiments, the disclosure provides a composition that consists essentially of ten purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, SEQ ID NO:2, and SEQ ID NO:6, respectively.

(138) The bacterial strains in Composition D are related to the following bacteria: *Clostridium scindens*, *Clostridium hathewayi*, *Blautia hansenii*, *Blautia wexlerae*, *Anaerotruncus colihominis*, *Dorea longicatena*, *Clostridium innocuum*, *Flavonifractor plautii*, *Lachnospiraceae bacterium 7_1_58* FAA, *Subdoligranulum*, *Turicibacter sanguinis*, and *Lactobacillus mucosae*. In some embodiments, the disclosure provides compositions with two or more bacterial strains of species selected from the group consisting of *Clostridium scindens*, *Clostridium hathewayi*, *Blautia hansenii*, *Blautia wexlerae*, *Anaerotruncus colihominis*, *Dorea longicatena*, *Clostridium innocuum*, *Erysipelotrichaceae bacterium 21-3*, *Flavonifractor plautii*, *Lachnospiraceae bacterium 7_1_58* FAA, *Turicibacter sanguinis*, and *Lactobacillus mucosae*. In some embodiments, the disclosure provides compositions that include two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:87, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, and SEQ ID NO:105.

(139) In some embodiments, the compositions disclosed herein do not include *Flavonifractor plautii*, *Subdoligranulum* or *Lachnospiraceae bacterium 7_1_58* FAA. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:1, SEQ ID NO:14, SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:2, and SEQ ID NO:6, wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NO:10. In some embodiments, the composition comprises two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:87, SEQ ID NO:90, SEQ ID NO:91, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:97, SEQ ID NO:98, SEQ ID NO:99, and SEQ ID NO:105, wherein the composition does not include a bacterial strain comprising a 16S rDNA sequence having at least 97% homology with a nucleic acid sequence of SEQ ID NO:93.

(140) The strains of Composition D include both BaiCD⁺ strains and BaiCD⁻ strains. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein one or more bacteria are BaiCD⁺ strains and one or more bacteria are BaiCD⁻ strains. In some embodiments of the one or more bacteria that are BaiCD⁺ strains are selected from bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:12, SEQ ID NO:3, SEQ ID NO:5, and SEQ ID NO:1. In some embodiments the one or more bacteria that are BaiCD⁻ strains are selected from bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:18, SEQ ID NO:21, SEQ ID NO:10, and SEQ ID NO:14. In some embodiments of the one or more bacteria that are BaiCD⁺ strains are selected from the bacterial species *Clostridium scindens*, *Clostridium hathewayi*, *Blautia hansenii*, and *Blautia wexlerae*. In some embodiments of the one or more bacteria that are BaiCD⁻ strains are selected from the bacterial species *Dorea longicatena*, *Clostridium innocuum*, *Flavonifractor plautii*, and *Anaerotruncus colihominis*. The Clostridial strains of Composition D are classified as belonging to *Clostridium* clusters IV, XIVa, and XVII. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and BaiCD⁺ strains and belong to *Clostridium* clusters IV, XIVa, or XVII. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and BaiCD⁺ strains and belong to *Clostridium* clusters XIVa or XVII. In some embodiments, the disclosure provides two or more bacterial strains, wherein the bacteria are BaiCD⁻ strains and BaiCD⁺ strains and belong to *Clostridium* clusters IV or XIVa.

(141) Composition D includes the non-*Clostridium* strains *Turicibacter sanguinis* and *Lactobacillus mucosae*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition includes both *Clostridium* strains and non-*Clostridium* strains. In some embodiments, the non-*Clostridium* strains are the members of the genus *Lactobacillus*. Members of the genus *Lactobacillus* include, without limitation *L. acetotolerans*, *L. acidifarinae*, *L. acidipiscis*, *L. acidophilus*, *L. agilis*, *L. algidus*, *L. alimentarius*, *L. amylolyticus*, *L. amylophilus*, *L. amylophobicus*, *L. amylovorus*, *L. animalis*, *L. antri*, *L. apodemi*, *L. aviarius*, *L. bifermentans*, *L. brevis*, *L. buchneri*, *L. camelliae*, *L. casei*, *L. cateniformis*, *L. ceti*, *L. coleohominis*, *L. collinoides*, *L. composti*, *L. concavus*, *L. coryniformis*, *L. crispatus*, *L. crustorum*, *L. curvatus*, *L. delbrueckii* subsp. *bulgaricus*, *L. delbrueckii* subsp. *delbrueckii*, *L. delbrueckii* subsp. *lactis*, *L. dextrinicus*, *L. diolivorans*, *L. equi*, *L. equigenerosi*, *L. farraginis*, *L. farciminis*, *L. fermentum*, *L. fornicalis*, *L. fructivorans*, *L. frumenti*, *L. fuchuensis*, *L. gallinarum*, *L. gasseri*, *L. gastricus*, *L. ghanensis*, *L. graminis*, *L. hammesii*, *L. hamsteri*, *L. harbinensis*, *L. hayakitensis*, *L. helveticus*, *L. hilgardii*, *L. homohiochii*, *L. iners*, *L. ingluviei*, *L. intestinalis*, *L. jensenii*, *L. johnsonii*, *L. kalixensis*, *L. kefiranoferiens*, *L. kefiri*, *L. kimchii*, *L. kitasatonis*, *L. kunkeei*, *L. leichmannii*, *L. lindneri*, *L. malefermentans*, *L. mali*, *L. manihotivorans*, *L. mindensis*, *L. mucosae*, *L. murinus*, *L. nagelii*, *L. namurensis*, *L. nantensis*, *L. oligofermentans*, *L. oris*, *L. panis*, *L. pantheris*, *L. parabrevis*, *L. parabuchneri*, *L. paracasei*, *L. paracollinoides*, *L. parafarraginis*, *L. parakefiri*, *L. paralimentarius*, *L. paraplantarum*, *L. pentosus*, *L. perolens*, *L. plantarum*, *L. pontis*, *L. protectus*, *L. psittaci*, *L. rennini*, *L. reuteri*, *L. rhamnosus*, *L. rimae*, *L. rogosa*, *L. rossiae*, *L. ruminis*, *L. saerimneri*, *L. sakei*, *L. salivarius*, *L. sanfranciscensis*, *L. satsumensis*, *L. secaliphilus*, *L. sharpeae*, *L. siliginis*, *L. spicheri*, *L. suebicus*, *L. thailandensis*, *L.*

ultunensis, *L. vaginalis*, *L. versmoldensis*, *L. vini*, *L. vitulinus*, *L. zeae*, and *L. zymae*. In some embodiments, the non-*Clostridium* strain is *Lactobacillus mucosae*. In some embodiments, the non-*Clostridium* strain is *Lactobacillus mucosae*. In some embodiments, the *Lactobacillus mucosae* has a 16S rDNA sequence comprising SEQ ID NO:2. In some embodiments, the *Lactobacillus mucosae* has a 16S rDNA sequence having at least 97% homology with a nucleic acid comprising SEQ ID NO:2. In some embodiments, the *Lactobacillus mucosae* has a 16S rDNA sequence having at least 97% homology with a nucleic acid comprising SEQ ID NO:91.

(142) In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition includes both *Clostridium* strains and non-*Clostridium* strains. In some embodiments, the non-*Clostridium* strains are members of the genus *Turicibacter*. In some embodiments, the non-*Clostridium* strain is *Turicibacter sanguinis*. In some embodiments, the *Turicibacter sanguinis* has a 16S rDNA sequence comprising SEQ ID NO:6. In some embodiments, the *Turicibacter sanguinis* has a 16S rDNA sequence having at least 97% homology with a nucleic acid comprising SEQ ID NO:6. In some embodiments, the *Turicibacter sanguinis* has a 16S rDNA sequence having at least 97% homology with a nucleic acid comprising SEQ ID NO:90.

(143) In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition includes both *Clostridium* strains and non-*Clostridium* strains. In some embodiments, the non-*Clostridium* strains are *Lactobacillus mucosae* and *Turicibacter sanguinis*.

(144) In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Lactobacillus*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Turicibacter*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Lactobacillus* or *Turicibacter*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition only includes clostridia strains. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition only includes clostridia strains belonging to *Clostridium* cluster IV, XIVa or XVII strains. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Clostridium* cluster XI strains.

(145) In some embodiments, the disclosure provides compositions comprising two or more purified bacterial strains selected from the group consisting of: *Clostridium scindens*, *Pseudoflavonifractor capillosus*, and *Blautia hansenii*. In some embodiments, the compositions disclosed herein do not include *Clostridium scindens*, *Pseudoflavonifractor capillosus*, or *Blautia hansenii*.

(146) TABLE-US-00006 TABLE D Composition D SEQ_12 - VE202-26 - *Clostridium scindens* (XIVa)* SEQ_03 - 5 - *Clostridium hathewayi* (XIVa)* SEQ_05 - 10 - *Blautia hansenii* (XIVa)* SEQ_01 - 71 - *Blautia wexlerae* (XIVa)* SEQ_14 - VE202-13 - *Anaerotruncus colihominis* (IV) SEQ_18 - 148 - *Dorea longicatena* (XIVa) SEQ_21 - 189 - *Clostridium innocuum* (XVII) SEQ_10 - 211 - *Flavonifractor plautii* (IV) SEQ_02 - 102 - *Turicibacter sanguinis* (non-*Clostridium*) SEQ_06 - 40 - *Lactobacillus mucosae* (non-*Clostridium*) *= BaiCD+

(147) In one aspect, the disclosure provides Composition F (See e.g., FIGS. 13 and 14, and Tables F1 and F2). As shown in FIG. 13, Composition F contains bacteria that have the following 16S rDNA sequences: SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, and SEQ ID NO:79.

(148) In some embodiments, the disclosure provides compositions with two or more purified bacterial strains that have 16S rDNA sequences selected from the group consisting of SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, and SEQ ID NO:79.

(149) In some embodiments, the compositions include two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, and SEQ ID NO:79.

(150) The bacterial strains in Composition F are related to the following bacteria: *Dorea longicatena*, *Ruminococcus obeum*, *Megasphaera elsdenii*, *Acidaminococcus fermentans*, *Acidaminococcus intestine*, *Megasphaera elsdenii*, *Ruminococcus faecis*, *Bacteroides cellulosilyticus*, *Anaerostipes hadrus*, *Ruminococcus obeum*, *Flavonifractor plautii*, *Eubacterium rectale*, *Flavonifractor plautii*, *Megasphaera elsdenii*, *Eubacterium rectale*, *Ruminococcus champanellensis*, *Ruminococcus albus*, *Ruminococcus champanellensis*, *Ruminococcus faecis*, *Bifidobacterium bifidum*, *Anaerostipes hadrus*, *Anaerostipes hadrus*, *Anaerostipes hadrus*, *Eubacterium rectale*, *Ruminococcus faecis*, *Blautia luti*, *Ruminococcus faecis*, *Anaerostipes hadrus*, *Anaerostipes hadrus*, *Ruminococcus faecis*, *Eubacterium rectale*, *Eubacterium rectale*, *Anaerostipes hadrus*, *Ruminococcus faecis*, *Ruminococcus faecis*, *Dorea longicatena*, *Roseburia faecis*, *Blautia luti*, *Fusicatenibacter saccharivorans*, *Fusicatenibacter saccharivorans*, *Roseburia faecis*, *Megasphaera elsdenii*, *Eubacterium rectale*, *Eubacterium rectale*, *Roseburia faecis*, *Blautia faecis*, *Fusicatenibacter saccharivorans*, and *Dorea formicigenerans*.

(151) In some embodiments, the disclosure provides compositions with two or more bacterial strains of species selected from the group consisting of *Dorea longicatena*, *Ruminococcus obeum*, *Megasphaera elsdenii*, *Acidaminococcus fermentans*, *Acidaminococcus intestine*, *Megasphaera elsdenii*, *Ruminococcus faecis*, *Bacteroides cellulosilyticus*, *Anaerostipes hadrus*, *Ruminococcus obeum*, *Flavonifractor plautii*, *Eubacterium rectale*, *Flavonifractor plautii*, *Megasphaera elsdenii*, *Eubacterium rectale*, *Ruminococcus champanellensis*, *Ruminococcus albus*, *Ruminococcus champanellensis*, *Ruminococcus faecis*, *Bifidobacterium bifidum*, *Anaerostipes hadrus*, *Anaerostipes hadrus*, *Anaerostipes hadrus*, *Eubacterium rectale*, *Ruminococcus faecis*, *Blautia luti*, *Ruminococcus faecis*, *Anaerostipes hadrus*, *Anaerostipes hadrus*, *Ruminococcus faecis*, *Eubacterium rectale*, *Eubacterium rectale*, *Anaerostipes hadrus*, *Ruminococcus faecis*, *Ruminococcus faecis*, *Dorea longicatena*, *Roseburia faecis*, *Blautia luti*, *Fusicatenibacter saccharivorans*, *Fusicatenibacter saccharivorans*, *Roseburia faecis*, *Megasphaera elsdenii*, *Eubacterium rectale*, *Eubacterium rectale*, *Roseburia faecis*, *Blautia faecis*, *Fusicatenibacter saccharivorans*, and *Dorea formicigenerans*.

(152) In some embodiments, the disclosure provides compositions that include two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:100, SEQ ID NO:104, SEQ ID NO:107, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, SEQ ID NO:118, SEQ ID NO:119, and SEQ ID NO:120. It should be appreciated that multiple strains of the compositions disclosed herein can have the same related bacterial species. For instance, Composition F includes 12 strains that have *Eubacterium rectale* as the closest related species.

(153) In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster IV. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster XIVa. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster IX. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster IV. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster XIVa and at least one of the bacterial strains belongs to *Clostridium* cluster IX. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster IV and at least one of the bacterial strains belongs to *Clostridium* cluster IX. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster IV, at least one of the bacterial strains belongs to *Clostridium* cluster XIVa, and at least one of the bacterial strains belongs to *Clostridium* cluster IX. In some embodiments, the compositions provided herein do not include bacterial strains belonging to *Clostridium* cluster XVIII. In some embodiments, the compositions provided herein do not include bacterial strains belonging to *Clostridium* cluster XVI or XVIII.

(154) Composition F includes non-*Clostridium* bacterial strains. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition includes both *Clostridium* strains and non-*Clostridium* strains. In some embodiments, the non-*Clostridium* strains are the members of the genus *Bacteroides*. In some embodiments, the non-*Clostridium* strain is *Bacteroides cellulosilyticus*. In some embodiments, the non-*Clostridium* strains are the members of the genus *Bifidobacterium*. In some embodiments, the non-*Clostridium* strain is *Bifidobacterium bifidum*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition includes both *Clostridium* strains and non-*Clostridium* strains, and wherein the non-*Clostridium* strains are *Bacteroides cellulosilyticus* and *Bifidobacterium bifidum*.

(155) In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Bacteroides*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Bifidobacterium*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Bacteroides* and does not include *Bifidobacterium*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include non-*Clostridium* strains. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition only includes clostridia strains belonging to *Clostridium* cluster IV, XIVa or XVII strains. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Clostridium* cluster XI strains.

(156) TABLE-US-00007
TABLE F1
Composition F
SEQ_NO StrainID Genus_species SEQ_NO StrainID Genus_species
SEQ_24 YK96 *Dorea longicatena* SEQ_52 YK51 *Eubacterium rectale* SEQ_25 YK101 *Ruminococcus obeum* SEQ_53 YK52 *Eubacterium rectale* SEQ_26 YK110 *Megasphaera elsdenii* SEQ_54 YK54 *Anaerostipes hadrus* SEQ_27 YK149 *Acidaminococcus fermentans* / SEQ_55 YK56 *Ruminococcus faecis* *Acidaminococcus intestini* SEQ_28 YK154 *Megasphaera elsdenii* SEQ_56 YK57 *Ruminococcus faecis* SEQ_29 YK36 *Ruminococcus faecis* SEQ_57 YK58 *Dorea longicatena* SEQ_30 YK95 *Bacteroides cellulosilyticus* SEQ_58 YK65 *Roseburia faecis* SEQ_31 YK32

Anaerostipes SEQ_59 YK67 *Blautia_luti* SEQ_32 YK64 *Ruminococcus_obeum* SEQ_60 YK69
Fusicatenibacter_saccharivorans SEQ_33 YK73 *Flavonifractor_plautii* SEQ_61 YK70 *Fusicatenibacter_saccharivorans*
SEQ_34 YK87 *Eubacterium_rectale* SEQ_62 YK71 *Roseburia_faecis* SEQ_35 YK105 *Flavonifractor_plautii* SEQ_63
YK74 *Megasphaera_elsdenii* SEQ_36 YK153 *Megasphaera_elsdenii* SEQ_64 YK88 *Eubacterium_rectale* SEQ_37 YK163
Eubacterium_rectale SEQ_65 YK89 *Eubacterium_rectale* SEQ_38 YK191 *Ruminococcus_champanellensis*/ SEQ_66 YK97
Roseburia_faecis *Ruminococcus_albus* SEQ_39 YK99 *Ruminococcus_champanellensis* SEQ_67 YK98 *Blautia_faecis*
SEQ_40 YK55 *Ruminococcus_faecis* SEQ_68 YK139 *Fusicatenibacter_saccharivorans* SEQ_41 YK75
Bifidobacterium_bifidum SEQ_69 YK141 *Dorea_formicigenerans* SEQ_42 YK90 *Anaerostipes_hadrus* SEQ_70 YK142
Ruminococcus_faecis SEQ_43 YK30 *Anaerostipes_hadrus* SEQ_71 YK152 *Blautia_hansenii* SEQ_44 YK31
Anaerostipes_hadrus SEQ_72 YK155 *Blautia_hansenii* SEQ_45 YK12 *Eubacterium_rectale* SEQ_73 YK157
Eubacterium_rectale SEQ_46 YK27 *Ruminococcus_faecis* SEQ_74 YK160 *Roseburia_faecis* SEQ_47 YK28 *Blautia_luti*
SEQ_75 YK166 *Eubacterium_rectale* SEQ_48 YK29 *Ruminococcus_faecis* SEQ_76 YK168 *Eubacterium_rectale* SEQ_49
YK33 *Anaerostipes_hadrus* SEQ_77 YK169 *Eubacterium_rectale* SEQ_50 YK34 *Anaerostipes_hadrus* SEQ_78 YK171
Eubacterium_rectale SEQ_51 YK35 *Ruminococcus_faecis* SEQ_79 YK192 *Roseburia_faecis*

(157) TABLE-US-00008 TABLE F2 Composition F, strain groupings Cluster Composition F *SCFAs XIVa *Eubacterium_rectale* 12 A, B, L *Ruminococcus_faecis* 8 A, L *Ruminococcus_obeum* 2 A, L *Blautia_faecis* 1 A, L *Blautia_hansenii* 2 A, L *Blautia_luti* 2 A, L *Anaerostipes_hadrus* 7 B *Roseburia_faecis* 5 A, B *Fusicatenibacter* A, L *saccharivorans* 3 *Dorea_formicigenerans* 1 A *Dorea_longicatena* 2 A IV *Flavomfractor_plautii* 2 A, B *Ruminococcus* A *champanellensis* 2 IX *Acidaminococcus_fermentans* A, B, P 1 *Megasphaera_elsdenii* 4 P other *Bacteroides_cellulosilyticus* A, S 1 *Bifidobacterium_bifidum* L, A *Short chain fatty acid legend: A, acetate; B, Butyrate; L, lactate; P, propionate; S, succinate

(158) In one aspect, the disclosure provides Composition G (See e.g., FIG. 19; Table G). As shown in FIG. 19, Composition G contains bacteria that have the following 16S rDNA sequences: SEQ ID NO:27, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:51, SEQ ID NO:55, SEQ ID NO:68, SEQ ID NO:72, SEQ ID NO:70, SEQ ID NO:24, SEQ ID NO:34, SEQ ID NO:37, SEQ ID NO:46, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:35, SEQ ID NO:62, SEQ ID NO:26, SEQ ID NO:63, SEQ ID NO:67, SEQ ID NO:40, SEQ ID NO:38, SEQ ID NO:47, SEQ ID NO:56, SEQ ID NO:25, and SEQ ID NO: 32.

(159) In some embodiments, the disclosure provides compositions with two or more purified bacterial strains that have 16S rDNA sequences selected from the group consisting of SEQ ID NO:27, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:51, SEQ ID NO:55, SEQ ID NO:68, SEQ ID NO:72, SEQ ID NO:70, SEQ ID NO:24, SEQ ID NO:34, SEQ ID NO:37, SEQ ID NO:46, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:35, SEQ ID NO:62, SEQ ID NO:26, SEQ ID NO:63, SEQ ID NO:67, SEQ ID NO:40, SEQ ID NO:38, SEQ ID NO:47, SEQ ID NO:56, SEQ ID NO:25, and SEQ ID NO: 32.

(160) In some embodiments, the compositions include two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:27, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:51, SEQ ID NO:55, SEQ ID NO:68, SEQ ID NO:72, SEQ ID NO:70, SEQ ID NO:24, SEQ ID NO:34, SEQ ID NO:37, SEQ ID NO:46, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:35, SEQ ID NO:62, SEQ ID NO:26, SEQ ID NO:63, SEQ ID NO:67, SEQ ID NO:40, SEQ ID NO:38, SEQ ID NO:47, SEQ ID NO:56, SEQ ID NO:25, and SEQ ID NO: 32.

(161) The bacterial strains in Composition G are related to the following bacteria: *Acidaminococcus_fermentans*, *Acidaminococcus_intestine*, *Anaerostipes_hadrus*, *Blautia_faecis*, *Blautia_hansenii*, *Dorea_formicigenerans*, *Dorea_longicatena*, *Eubacterium_rectale*, *Flavonifractor_plautii*, *Fusicatenibacter_saccharivorans*, *Megasphaera_elsdenii*, *Roseburia_faecis*, *Ruminococcus_champanellensis*, *Ruminococcus_albus*, *Ruminococcus_faecis*, and *Ruminococcus_obeum*.

(162) In some embodiments, the disclosure provides compositions with two or more bacterial strains of species selected from the group consisting of *Acidaminococcus_fermentans*, *Acidaminococcus_intestine*, *Anaerostipes_hadrus*, *Blautia_faecis*, *Blautia_hansenii*, *Dorea_formicigenerans*, *Dorea_longicatena*, *Eubacterium_rectale*, *Flavonifractor_plautii*, *Fusicatenibacter_saccharivorans*, *Megasphaera_elsdenii*, *Roseburia_faecis*, *Ruminococcus_champanellensis*, *Ruminococcus_albus*, *Ruminococcus_faecis*, and *Ruminococcus_obeum*.

(163) In some embodiments, the disclosure provides compositions that include two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:96, SEQ ID NO:97, SEQ ID NO:99, SEQ ID NO:104, SEQ ID NO:107, SEQ ID NO:111, SEQ ID NO:112, SEQ ID NO:113, SEQ ID NO:114, SEQ ID NO:115, SEQ ID NO:116, SEQ ID NO:117, and SEQ ID NO:119.

(164) TABLE-US-00009 TABLE G Composition G SEQ_27 YK149 *Acidaminococcus_fermentans*/*Acidaminococcus_intestini* SEQ_43 YK90 *Anaerostipes_hadrus* SEQ_44 YK30 *Anaerostipes_hadrus* SEQ_51 YK34 *Anaerostipes_hadrus* SEQ_55 YK54 *Anaerostipes_hadrus* SEQ_68 YK98 *Blautia_faecis* SEQ_72 YK152 *Blautia_hansenii* SEQ_70 YK141 *Dorea_formicigenerans* SEQ_24 YK96 *Dorea_longicatena* SEQ_34 YK87 *Eubacterium_rectale* SEQ_37 YK163 *Eubacterium_rectale* SEQ_46 YK12 *Eubacterium_rectale* SEQ_76 YK166 *Eubacterium_rectale* SEQ_77 YK168 *Eubacterium_rectale* SEQ_35 YK105 *Flavonifractor_plautii* SEQ_62 YK70 *Fusicatenibacter_saccharivorans* SEQ_26 YK110 *Megasphaera_elsdenii* SEQ_63 YK71 *Roseburia_faecis* SEQ_67 YK97 *Roseburia_faecis* SEQ_40 YK99 *Ruminococcus_champanellensis* SEQ_38 YK191 *Ruminococcus_champanellensis*/*Ruminococcus_albus* SEQ_47 YK27 *Ruminococcus_faecis* SEQ_56 YK56 *Ruminococcus_faecis* SEQ_25 YK101 *Ruminococcus_obeum* SEQ_32 YK64 *Ruminococcus_obeum*

(165) In one aspect, the disclosure provides Composition H (See e.g., FIG. 26, Table H). As shown in FIG. 26, Composition H contains bacteria that have the following 16S rDNA sequences: SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:21, SEQ ID NO:82, SEQ ID NO:81, and SEQ ID NO:80. In some embodiments, the disclosure provides compositions with two or more

purified bacterial strains that have 16S rDNA sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:21, SEQ ID NO:82, SEQ ID NO:81, and SEQ ID NO:80.

(166) In some embodiments, the compositions include two or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:21, SEQ ID NO:82, SEQ ID NO:81, and SEQ ID NO:80. In some embodiments, the compositions include four or more purified bacterial strains comprising 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:21, SEQ ID NO:82, SEQ ID NO:81, and SEQ ID NO:80.

(167) The bacterial strains in Composition H are related to the following bacteria: *Anaerotruncus colihominis*, *Clostridium symbiosum*, *Clostridium innocuum*, *Erysipelotrichaceae_bacterium_21-3*, *Clostridium disporicum*, *Clostridium bolteae*, and *Erysipelatoclostridium ramosum*. In some embodiments, the disclosure provides compositions with two or more bacterial strains selected from the group consisting of *Anaerotruncus colihominis*, *Clostridium symbiosum*, *Clostridium innocuum*, *Erysipelotrichaceae_bacterium_21-3*, *Clostridium disporicum*, *Clostridium bolteae*, and *Erysipelatoclostridium ramosum*.

(168) In some embodiments, the disclosure provides compositions that include two or more purified bacterial strains comprise 16S rDNA sequences having at least 97% homology with nucleic acid sequences selected from the group consisting of SEQ ID NO:86, SEQ ID NO:95, SEQ ID NO:98, SEQ ID NO:110, SEQ ID NO:122, and SEQ ID NO:123.

(169) Composition H includes bacteria from *Clostridium* cluster I, IV, XIVa, XVII and XVIII. In some embodiments, the disclosure provides compositions that include two or more purified bacterial strains from *Clostridium* cluster I, IV, XIVa, XVII and XVIII. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster IV. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster XIVa. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster XVII. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster I. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster XVIII. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster XIVa and at least one of the bacterial strains belongs to *Clostridium* cluster IV. In some embodiments, at least one of the bacterial strains of the composition belongs to *Clostridium* cluster XIVa and at least one of the bacterial strains belongs to *Clostridium* cluster XVII.

(170) TABLE-US-00010 TABLE H Composition H SEQ ID NO Strain Closest species Cluster SEQ ID NO: 14 VE202-13 *Anaerotruncus colihominis* Cluster IV SEQ ID NO: 16 VE202-16 *Clostridium symbiosum* Cluster XIVa WAL-14163 SEQ ID NO: 21 189 *Clostridium innocuum* Cluster XVII SEQ ID NO: 82 PE9 *Clostridium disporicum* Cluster I SEQ ID NO: 81 PE5 *Clostridium bolteae* Cluster XIVa SEQ ID NO: 80 VE202-18 *Erysipelatoclostridium* Cluster XVIII *ramosum*

(171) In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include bacteria from *Clostridium* cluster I. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include bacteria from *Clostridium* cluster XVIII. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include bacteria from *Clostridium* cluster I and does not include bacteria from *Clostridium* cluster XVIII.

(172) In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein all the bacteria are anaerobic bacteria. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein all the bacteria are obligate anaerobic bacteria.

(173) In some embodiments, the disclosure provides compositions comprising two or more bacteria (e.g., purified bacterial strains), wherein the composition does not include *Clostridium scindens*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Flavonifractor plautii*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include Parabacteroides. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Lactobacillus*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include Colinsella. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include Dialister. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include Raoultella. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Streptococcus*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Staphylococcus*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include *Microbacterium*. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include Proteobacteria. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include Peptostreptococcaceae. In some embodiments, the disclosure provides compositions comprising two or more bacteria, wherein the composition does not include Oscillospiraceae.

(174) In one aspect, the disclosure provides bacterial strains with 16S rDNA sequences that have homology to a nucleic acid sequence of any one of the sequences of the bacterial strains or species described herein. In some embodiments, the bacterial strain has at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.6%, 99.7%, 99.8% or 99.9% homology relative to any of the strains or bacterial species described herein over a specified region or over the entire sequence. It would be appreciated by one of skill in the art that the term "homology" or "percent homology," in the context of two or more nucleic acid sequences or amino acid sequences, refers to a measure of similarity between two or more sequences or portion(s) thereof. The homology may exist over a region of a sequence that is at least about 50 nucleotides in length, or more preferably over a region that is 100 to 500 or 1000 or more

nucleotides in length. In some embodiments, the homology exists over the length the 16S rRNA or 16S rDNA sequence, or a portion thereof.

(175) Additionally, or alternatively, two or more sequences may be assessed for the identity between the sequences. The terms “identical” or percent “identity” in the context of two or more nucleic acids or amino acid sequences, refer to two or more sequences or subsequences that are the same. Two sequences are “substantially identical” if two sequences have a specified percentage of amino acid residues or nucleotides that are the same (e.g., at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.6%, 99.7%, 99.8% or 99.9% identical) over a specified region or over the entire sequence, when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. Optionally, the identity exists over a region that is at least about 50 nucleotides in length, or more preferably over a region that is 100 to 500 or 1000 or more nucleotides in length. In some embodiments, the identity exists over the length the 16S rRNA or 16S rDNA sequence.

(176) Additionally, or alternatively, two or more sequences may be assessed for the alignment between the sequences. The terms “alignment” or percent “alignment” in the context of two or more nucleic acids or amino acid sequences, refer to two or more sequences or subsequences that are the same. Two sequences are “substantially aligned” if two sequences have a specified percentage of amino acid residues or nucleotides that are the same (e.g., at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, 99.5%, 99.6%, 99.7%, 99.8% or 99.9% identical) over a specified region or over the entire sequence, when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. Optionally, the alignment exists over a region that is at least about 50 nucleotides in length, or more preferably over a region that is 100 to 500 or 1000 or more nucleotides in length. In some embodiments, the identity exists over the length the 16S rRNA or 16S rDNA sequence.

(177) For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. Methods of alignment of sequences for comparison are well known in the art. See, e.g., by the local homology algorithm of Smith and Waterman (1970) *Adv. Appl. Math.* 2:482c, by the homology alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* (1970) 48:443, by the search for similarity method of Pearson and Lipman. *Proc. Natl. Acad. Sci. USA* (1998) 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, Madison, WI), or by manual alignment and visual inspection (see, e.g., Brent et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (Ringbom ed., 2003)). Two examples of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., *Nuc. Acids Res.* (1977) 25:3389-3402, and Altschul et al., *J. Mol. Biol.* (1990) 215:403-410, respectively.

(178) In one aspect, the disclosure provides compositions comprising multiple purified bacterial strains (e.g., Compositions A-J). For instance, FIGS. 1, 13, 19, and 26 present several example compositions comprising multiple bacterial strains. In one aspect, the 16S rDNA sequences of purified bacterial strains of the compositions were compared to 16S rDNA sequences of known bacterial species/strains in a bacterial genome database to identify the closest known related bacterial species to the bacterial strains disclosed herein (See e.g., Table 1). It should be appreciated that multiple bacterial strains of the compositions disclosed herein may have the same closest related bacterial species. In one aspect, the disclosure provides compositions comprising one or more bacterial strains or species with 16S rDNA sequences that have homology to a nucleic acid sequence of any one of the sequences provided by SEQ ID NOs:1-83 and 124-159. In some embodiments, the species with 16S rDNA sequences with homology to a nucleic acid sequence of any one of the closest related species to any of the strains described herein, correspond to bacterial strains with 16S rDNA sequences provided by SEQ ID NOs:84-123.

(179) In some embodiments, the compositions disclosed herein provide at least one of the bacterial strains (e.g., purified bacterial strains) described herein. In some embodiments, the compositions that comprise at least one bacterial strain, comprise at least one bacterial strain with a 16S rDNA sequence selected from any one of SEQ ID NOs:1-122 and 124-159. In some embodiments, the compositions that comprise at least one bacterial strain, comprise at least one bacterial strain with a 97% homology to 16S rDNA sequence selected from any one of SEQ ID NOs:1-122 and 124-159.

(180) In some embodiments, the compositions disclosed herein comprise two or more bacterial strains. In some embodiments, the compositions described herein comprise at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, or at least 20 or more bacterial strains (e.g., purified bacterial strains).

(181) It should be appreciated the compositions and methods provided herein can be distinguished from compositions and methods associated with the treatment of *C. difficile* infection that are available. For instance, it has been proposed that non-toxigenic *C. difficile* strains, i.e., strains that do not produce *C. difficile* toxins, may be used to treat *C. difficile* infection (See, e.g., U.S. Pat. No. 6,635,260). The compositions disclosed herein can be distinguished at least because the compositions described herein do not comprise non-toxigenic strains of *C. difficile*. Thus, in some embodiments, the compositions herein do not include comprise non-toxigenic strains of *C. difficile*. *C. difficile* belongs to *Clostridium* cluster XI. In some embodiments, the compositions herein do not include bacterial strains belonging to *Clostridium* cluster XI.

(182) It is also considered in the art that bacterial strains expressing a bile inducible 7 α / β -dehydroxylation operon can be used in the treatment of *C. difficile* (see, e.g., Buffie et al. *Nature* (2015) 517:205-208). The catalysis of bile acid 7 α dihydroxylation is mediated by a stereo-specific NAD(H)-dependent 3-oxo- Δ .sup.4-cholenoic acid oxidoreductase encoded by the gene baiCD. In some embodiments, the compositions provided herein do not mediate bile acid 7- α -dehydroxylation.

(183) In contrast to the findings in the art, in some embodiments, as shown herein, combinations of bacterial strains that do not encode baiCD (or a homolog thereof), or encode a baiCD that comprises one or more mutations that result in a non-functional BaiCD protein (“baiCD-”), are more effective at treating *C. difficile* infection and/or reducing or inhibiting production of Toxin B by *C. difficile* than combinations of bacterial strains that have a functional BaiCD protein (“baiCD+”). Thus, in some embodiments, the compositions of bacterial strains provided herein are baiCD- (i.e., the combination of the bacteria has no effective baiCD+ function). In some embodiments, all of bacterial strains in the compositions provided herein are baiCD-. In some embodiments, the majority (i.e., 50% or greater) of the bacterial strains in the compositions are baiCD-. In some embodiments, the majority (i.e., 50% or greater) of the bacterial strains in the compositions are baiCD- and the composition has no effective BaiCD function. In some embodiments, the minority (i.e., 50% or less) of the bacterial strains in the compositions are baiCD- and the composition has no effective BaiCD function. In some embodiments, bacterial strains for the compositions are selected based on the absence (or presence) of a baiCD gene or a predicted baiCD gene. In some embodiments, bacterial strains may be modified (e.g., genetically engineered) to prevent or reduce expression of a baiCD gene and/or to reduce or eliminate NAD(H)-dependent 3-oxo- Δ .sup.4-cholenoic acid oxidoreductase activity of BaiCD protein. The NAD(H)-dependent 3-oxo- Δ .sup.4-cholenoic acid oxidoreductase activity of a bacterial strain may be assessed by methods such as measuring the amount of 7 α -dehydroxylated bile acid. In some embodiments, the compositions described herein comprise bacterial strains without the baiCD operon (baiCD.sup.-) or baiCD function.

(184) In some embodiments, the compositions described herein do not include *Clostridium scindens*. In some embodiments, the compositions described herein do not include *Barnesiella intestihominis*. In some embodiments, the compositions described herein do not include *Blautia hansenii*. In some embodiments, the compositions described herein do not include *Pseudoflavinofractor capillosus*. In some embodiments, the compositions described herein do not include *Clostridium scindens*, *Barnesiella intestihominis*, *Blautia hansenii* or *Pseudoflavinofractor capillosus*.

(185) In some embodiments, the compositions provided herein do not include *Colinsella aerofaciens*. In some embodiments, the compositions provided herein do not include *Acetovibrio ethanoligignens*. In some embodiments, the compositions provided herein do not include bacterial strains belonging to *Clostridium* cluster I. In some embodiments, the compositions provided herein do not include *Clostridium butyricum*. In some embodiments, the compositions provided herein do not include *Clostridium disporicum*. In some embodiments, the compositions provided herein do not include strains belonging to *Clostridium* cluster XI. In some embodiments, the compositions provided herein do not include *Clostridium glycolicum*. In some embodiments, the compositions provided herein do not include *Faecalibacterium prausnitzii*. In some embodiments, the compositions provided herein do not include *Turicibacter sanguinis*. In some embodiments, the compositions provided herein do not include *Eubacterium rectale*. In some embodiments, the compositions provided herein do not include *Eubacterium ventriosum*. In some embodiments, the compositions provided herein do not include *Ruminococcus obeum*. In some embodiments, the compositions provided herein do not include *Pseudobutyrvibrio*. In some embodiments, the compositions provided herein do not include *Christensenellaceae*. In some embodiments, the compositions do not comprise gram-negative bacteria. In some embodiments, the compositions do not comprise *E. coli*. In some embodiments, the compositions do not comprise fungi, such as *Monilla* species.

(186) In some embodiments of the compositions provided herein, the compositions do not include bacterial strains that are resistant to one or more antibiotics. It should be appreciated that it may be desirable to have a mechanism to remove the bacterial compositions provided herein from the body of the subject after administration. One such mechanism is to remove the bacterial compositions by antibiotic treatment. Thus, in some embodiments, the compositions do not include bacterial strains that are resistant to one or more antibiotics. In some embodiments, the compositions do not include bacterial strains that are resistant to one or more antibiotics selected from the group consisting of penicillin, benzylpenicillin, ampicillin, sulbactam, amoxicillin, clavulanate, tazobactam, piperacillin, cefmetazole, vancomycin, imipenem, meropenem, metronidazole and clindamycin. In some embodiments, the compositions do not include bacterial strains that are resistant to vancomycin.

(187) In some embodiments, the compositions include bacterial strains that are susceptible to at least four antibiotics that are efficacious in humans. In some embodiments, the compositions include bacterial strains that are susceptible to at least three antibiotics that are efficacious in humans. In some embodiments, the compositions include bacterial strains that are susceptible to at least two antibiotics that are efficacious in humans. In some embodiments, the compositions include bacterial strains that are susceptible to at least one antibiotic that is efficacious in humans. In some embodiments, the compositions include only bacterial strains that are susceptible to at least four antibiotics that are efficacious in humans. In some embodiments, the compositions include only bacterial strains that are susceptible to at least three antibiotics that are efficacious in humans. In some embodiments, the compositions include only bacterial strains that are susceptible to at least two antibiotics that are efficacious in humans. In some embodiments, the compositions include bacterial strains that are susceptible to at least one antibiotic that is efficacious in humans. As used herein, an “antibiotic that is efficacious in a human” refers to an antibiotic that has been used to successfully treat bacterial infections in a human.

(188) In some embodiments, the compositions described herein comprise spore forming and non-spore forming bacterial strains. In some embodiments, the compositions described herein comprise spore forming bacterial strains. In some embodiments, the compositions described herein comprise only spore forming bacterial strains. In some embodiments, the compositions described herein comprise only non-spore forming bacterial strains. The spore-forming bacteria can be in spore form (i.e., as spores) or in vegetative form (i.e., as vegetative cells). In spore form, bacteria are generally more resistant to environmental conditions, such as heat, acid, radiation, oxygen, chemicals, and antibiotics. In contrast, in the vegetative state or actively growing state, bacteria are more susceptible to such environmental conditions, compared to in the spore form. In general, bacterial spores are able to germinate from the spore form into a vegetative/actively growing state, under appropriate

conditions. For instance, in spore form format may germinate when they are introduced in the intestine.

(189) In some embodiments, at least one (e.g., 1, 2, 3, 4, 5, or more) of the bacterial strains in the composition is a spore former. In some embodiments, at least one (e.g., 1, 2, 3, 4, 5, or more) of the bacterial strains in the composition is in spore form. In some embodiments, at least one (e.g., 1, 2, 3, 4, 5, or more) of the bacterial strains in the composition is a non-spore former. In some embodiments, at least one (e.g., 1, 2, 3, 4, 5, or more) of the bacterial strains in the composition is in vegetative form (As discussed above, spore forming bacteria can also be in vegetative form). In some embodiments, at least one (e.g., 1, 2, 3, 4, 5, or more) of the bacterial strains in the composition is in spore form and at least one (e.g., 1, 2, 3, 4, 5, or more) of the bacterial strains in the composition is in vegetative form. In some embodiments, at least one bacterial strain that is considered able to form spores (i.e., a spore-former) but is present in the composition in vegetative form. In some embodiments, at least one bacterial strain that is considered able to form spores is present in the composition both in spore form and in vegetative form.

(190) In some embodiments, the disclosure provides compositions wherein the compositions comprise bacterial strains that are spore forming bacterial strains. In some embodiments, the disclosure provides compositions wherein the compositions comprise bacterial strains that are non-spore forming bacterial strains. In some embodiments, the disclosure provides compositions wherein the compositions comprise bacterial strains that are spore forming bacterial strains and bacterial strains that are non-spore forming bacterial strains. In some embodiments, the disclosure provides compositions, wherein the compositions comprise a mixture of bacterial strains wherein at least 10% of the bacterial strains are spore forming bacterial strains, at least 20% of the bacterial strains are spore forming bacterial strains, at least 30% of the bacterial strains are spore forming bacterial strains, at least 40% of the bacterial strains are spore forming bacterial strains, at least 50% of the bacterial strains are spore forming bacterial strains, at least 60% of the bacterial strains are spore forming bacterial strains, at least 70% of the bacterial strains are spore forming bacterial strains, at least 80% of the bacterial strains are spore forming bacterial strains, at least 90% of the bacterial strains are spore forming bacterial strains bacteria up to 100% spore forming bacterial strains. Whether a bacterial strain is a spore forming strain can be determined for instance by evaluating the genome of the bacterial strain for the presence of sporulation genes. However, it should be appreciated that not all bacteria that are predicted to encode spore forming genes can be made to sporulate. In addition, whether a bacterial strain is a spore forming strain can be determined by exposing the bacterial strain to stress conditions, e.g., heat or exposure to chemicals (e.g., ethanol or chloroform), that are known to induce sporulation.

(191) It should be appreciated that spore forming bacteria can be in spore form or in vegetative form. In some embodiments of the compositions provided herein, the spore forming bacteria are in spore form. In some embodiments of the compositions provided herein, the spore forming bacteria are in vegetative form. In some embodiments of the compositions provided herein, the spore forming bacteria are both present in spore form and in vegetative form. In some embodiments, the disclosure provides compositions, wherein the compositions comprise spore forming bacteria at least 10% of the spore forming bacteria are in spore format, at least 20% of the spore forming bacteria are in spore format, at least 30% of the spore forming bacteria are in spore format, at least 40% of the spore forming bacteria are in spore format, at least 50% of the spore forming bacteria are in spore format, at least 60% of the spore forming bacteria are in spore format, at least 70% of the spore forming bacteria are in spore format, at least 80% of the spore forming bacteria are in spore format, at least 90% of the spore forming bacteria are in spore format, up to 100% in spore format.

(192) It is envisioned that the bacterial strains of the compositions provided herein are alive and will be alive when they reach the target area (e.g., the intestines). Bacterial spores are considered to be alive in this regards. In some embodiments, bacteria that are administered as spores may germinate in the target area (e.g., the intestines). It should further be appreciated that not all of the bacteria are alive and the compositions can include a percentage (e.g., by weight) that is not alive. In addition, in some embodiments, the compositions include bacterial strains that are not alive when administered or at the time when the composition reaches the target area (e.g., the intestines). It is envisioned that non-living bacteria may still be useful by providing some nutrients and metabolites for the other bacterial strains in the composition.

(193) Methods of inducing sporulation of spore-forming bacterial strains are well known in the art (See e.g., Paredes-Sabja et al., *Trends Microbiol.* (2011) 19(2):85-94). Generally, bacterial strains that are spore-formers can be made to go into spore form by stressing the bacterial strains. Non-limiting examples of stresses that can induce sporulation are an increase in temperature, change in the nutrients available and/or exposure to chemicals (e.g., ethanol or chloroform). It should be noted that bacteria that are non-spore formers, for instance because they are missing sporulation genes, cannot be made to sporulate by stress. To prepare compositions in which all the bacterial strains are in the spore form, the composition or bacterial cultures used to prepare the composition may be subjected to treatment to kill any bacteria not in spore form (e.g., in vegetative form), for example by exposing the composition to heat and are chemically breaking down the non-spore bacteria. The bacteria in spore format can subsequently be separated from the non-spore bacteria for instance by filtration.

(194) The amount of spores can be quantified using techniques know in the art. These techniques include phase contrast microscopy for enumerating spores using a hemocytometer. In addition, the viability of spores can be determined by plating the spores and growing the spores. For instance, spores can be plated in appropriate media and incubated in the anaerobic chamber for a period of time (e.g., 48-96 hrs.). Viability can subsequently be determined by quantifying the colony forming units which correspond to spores that germinated. For instance, spores can be plated on TCCFA plates (Taurocholate, cycloserine, cefoxitin, fructose agar plates), in which taurocholate helps the spores to germinate. In addition, spores can be quantified using the dipicolinic assay (DPA assay). DPA is an agent that allows for spore selection and is a clear indicator of endospores. When complexed with terbium, bright green luminescence is observed.

(195) In any of the compositions provided herein, in some embodiments, the bacterial strains are purified. In any of the compositions provided herein, in some embodiments, the bacterial strains are isolated. Any of the bacterial strains described

herein may be isolated and/or purified, for example, from a source such as a culture or a microbiota sample (e.g., fecal matter). The bacterial strains used in the compositions provided herein generally are isolated from the microbiome of healthy individuals. However, bacterial strains can also be isolated from individuals that are considered not to be healthy. In some embodiments, the compositions include strains originating from multiple individuals.

(196) As used herein, the term “isolated” bacteria that have been separated from one or more undesired component, such as another bacterium or bacterial strain, one or more component of a growth medium, and/or one or more component of a sample, such as a fecal sample. In some embodiments, the bacteria are substantially isolated from a source such that other components of the source are not detected.

(197) As also used herein, the term “purified” refers to a bacterial strain or composition comprising such that has been separated from one or more components, such as contaminants. In some embodiments, the bacterial strain is substantially free of contaminants. In some embodiments, one or more bacterial strains of a composition may be independently purified from one or more other bacteria produced and/or present in a culture or a sample containing the bacterial strain. In some embodiments, a bacterial strain is isolated or purified from a sample and then cultured under the appropriate conditions for bacterial replication, e.g., under anaerobic culture conditions. The bacteria that is grown under appropriate conditions for bacterial replication can subsequently be isolated/purified from the culture in which it is grown.

(198) In some embodiments, the bacterial strains of the compositions provided herein are obligate anaerobes. In some embodiments, the bacterial strains of the compositions provided herein are facultative anaerobes.

(199) Aspects of the present disclosure are related to methods for treating a pathogenic infection in a subject by administering a therapeutically effective amount of any of the compositions described herein. In some embodiments, the subject is a mammalian subject, such as a human, non-human primate, rodent, rabbit, sheep, pig, dog, cat, horse, or cow. In some embodiments, the subject is a human subject. In some embodiments, the subject is a pig.

(200) In some embodiments, the subject is a carrier of a pathogenic organism and is suffering from the effects of the infection (e.g., diarrhea caused by *C. difficile* toxins). In some embodiments the subject is an asymptomatic carrier of a pathogen. In some embodiments, the subject is a carrier of *C. difficile*. In some embodiments the subject is an asymptomatic *C. difficile* carrier. In some embodiments, the subject has experienced recurrent or chronic pathogenic infections. In some embodiments, the subject is suffering from a first occurrence of a particular pathogenic infection. In some embodiments, the subject has been treated with antibiotics which resulted in the recurrence of the pathogenic infection. In some embodiments, the subject has been treated with antibiotics which resulted in a first occurrence of a pathogenic infection. In some embodiments, the subject is to undergo a procedure that puts the subject at a higher risk of infection. In some embodiments, the compositions provided herein are administered to a subject to lower the risk of becoming infected by a pathogen.

(201) In some embodiments, the compositions provided herein are administered to a subject if the subject has a dysbiosis (e.g., has a microbiome associated with a disease state). In some embodiments, treatment with the compositions provided herein results in the change in the microbiome of the subject. In some embodiments, treatment with the compositions provided herein removes the dysbiosis in the subject resulting in a healthy microbiome. In some embodiments, treatment with the compositions provided herein removes the dysbiosis in the subject resulting in microbiome refractory or less susceptible to infection by a pathogen.

(202) As used herein, the term “pathogen” in regard to a pathogenic infection refers to a microorganism (e.g., a *bacterium*) that causes a disease or a disease state in a subject. In some embodiments, the disease or disease state of the subject may include symptoms such as colitis, diarrhea, watery diarrhea, abdominal cramping, fever, blood or pus in the stool, nausea, dehydration, loss of appetite, chills, weight loss, and/or kidney failure. In some embodiments, the pathogenic infection may be diagnosed, for example, by detecting a pathogen (or protein or nucleic acid associated with a pathogen) in a fecal sample collected from the subject. In some embodiments, the pathogenic infection may be diagnosed, for example, by comparing the microbiota of a fecal sample of the subject with the microbiota in a fecal sample of a healthy subject.

(203) In some embodiments, the pathogenic infection is *C. difficile*; *Clostridium perfringens*; *Clostridium botulinum*; *Clostridium tributrycum*; *Clostridium sporogenes*; *Escherichia coli*; *Pseudomonas aeruginosa*, such as Multidrug Resistant *Pseudomonas aeruginosa*; Vancomycin Resistant Enterococci (VRE); Carbapenem Resistant Enterobacteriaceae (CRE); *Neisseria gonorrhoeae*; *Acinetobacter*; Multidrug Resistant *Acinetobacter*; *Campylobacter*; Multi-drug resistant *Campylobacter*; *Candida*; Fluconazole-resistant *Candida*; Extended spectrum beta-lactamase (ESBL) producing Enterobacteriaceae; *Salmonella*, *Salmonella Typhimurium*, Drug resistant non-typhoid *Salmonella* spp.; Drug resistant *Salmonella Typhi*; Drug resistant *Shigella*; *Staphylococcus aureus*, such as Methicillin Resistant *S. aureus* or vancomycin resistant *S. aureus*; Drug resistant *Streptococcus pneumoniae*; Drug resistant Tuberculosis; Erythromycin Resistant Group A *Streptococcus*; Clindamycin resistant Group B *Streptococcus*, and any combinations thereof. In some embodiments, the pathogenic infection is *C. difficile*. In some embodiments, the *C. difficile* is an antibiotic-resistant *C. difficile*, e.g., fluoroquinolone resistant *C. difficile*. In some embodiments, the pathogenic infection is vancomycin-resistant Enterococci.

(204) Additional non-limiting examples of pathogens responsible for pathogenic infection that can be treated according to the methods provided herein are *Leishmania*, *Staphylococcus epidermis*, *Staphylococcus saprophyticus*, *Streptococcus pyogenes*, *Streptococcus pneumoniae*, *Streptococcus agalactiae*, *Enterococcus faecalis*, *Corynebacterium diphtheriae*, *Bacillus anthracis*, *Listeria monocytogenes*, *Clostridium perfringens*, *Clostridium tetanus*, *Clostridium botulinum*, *Clostridium difficile*, *Neisseria meningitidis*, *Neisseria gonorrhoeae*, *Escherichia coli*, *Salmonella typhimurium*, *Salmonella choleraesuis*, *Salmonella enterica*, *Salmonella enteritidis*, *Yersinia pestis*, *Yersinia pseudotuberculosis*, *Yersinia enterocolitica*, *Vibrio cholerae*, *Campylobacter jejuni*, *Campylobacter fetus*, *Helicobacter pylori*, *Pseudomonas aeruginosa*, *Pseudomonas mallei*, *Haemophilus influenzae*, *Bordetella pertussis*, *Mycoplasma pneumoniae*, *Ureaplasma urealyticum*, *Legionella pneumophila*, *Treponema pallidum*, *Leptospira interrogans*, *Borrelia burgdorferi*, *Mycobacterium tuberculosis*, *Mycobacterium leprae*,

Chlamydia psittaci, *Chlamydia pneumoniae*, *Rickettsia rickettsii*, *Rickettsia akari*, *Rickettsia prowazekii*, *Brucella abortus*, *Brucella melitens*, *Brucella suis*, and *Francisella tularensis*. In general, any *bacterium* that is capable of inducing a disease in a subject and/or that is not present in healthy individual is considered a pathogen herein. It should be appreciated that a subject may carry multiple pathogens and/or have multiple pathogenic infections.

(205) Any of the compositions described herein may be administered to a subject in a therapeutically effective amount or a dose of a therapeutically effective amount to treat or prevent a pathogenic infection (e.g., one or more pathogenic infections). The terms “treat” or “treatment” refer to reducing or alleviating one or more of the symptoms associated with a pathogenic infection, reducing the amount of bacterial toxin produced by the pathogenic infection, and/or reducing the bacterial load of the pathogenic infection. The terms “prevent” or “prevention” encompass prophylactic administration and may reduce the incidence or likelihood of pathogenic infection or a recurrent or chronic pathogenic infection. For instance, in some embodiments, administration of the compositions provided herein result in a healthy microbiome that is refractory to pathogenic infection, thereby preventing the pathogenic infection.

(206) As used herein, a “therapeutically effective amount” of composition, such as a pharmaceutical composition, is any amount that results in a desired response or outcome in a subject, such as those described herein, including but not limited to prevention of infection, an immune response or an enhanced immune response to the pathogenic infection, prevention or reduction of symptoms associated with pathogenic infection, and/or a reduction or inhibition of toxin production by the pathogenic infection. It should be appreciated that the term effective amount may be expressed in number of bacteria or bacterial spores to be administered. It should further be appreciated that the bacteria can multiply once administered. Thus, administration of even a relatively small amount of bacteria may have therapeutic effects.

(207) In some embodiments, the therapeutically effective amount of any of the compositions described herein is an amount sufficient to enhance survival of the subject, reduce the bacterial burden of the pathogenic infection in the subject, and/or reduce or inhibit toxin production by the pathogenic infection. In some embodiments, the therapeutically effective amount is an amount sufficient to reduce the bacterial burden of the pathogenic infection in a fecal sample from the subject by at least 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 20-fold, 30-fold, 40-fold, 50-fold, 100-fold, 1000-fold, 10.sup.4-fold, 10.sup.5-fold or more, as compared to the bacterial burden in a subject with a pathogenic infection that has not received any of the compositions described herein, or as compared to a fecal sample from the same subject that was collected prior to administration of any of the compositions.

(208) In some embodiments, the compositions provided herein inhibit the production of a bacterial toxin, e.g., *C. difficile* Toxin B. In some embodiments, the therapeutically effective amount is an amount sufficient to reduce or inhibit the amount of bacterial toxin (e.g., *C. difficile* Toxin B) produced by pathogenic infection in a fecal sample from the subject by at least 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 20-fold, 30-fold, 40-fold, 50-fold, 100-fold, 150-fold, 200-fold, 500-fold or more, as compared to the amount of the bacterial toxin in a subject with a pathogenic infection that has not received any of the compositions described herein or as compared to a fecal sample from the same subject that was collected prior to administration of any of the compositions.

(209) In some embodiments, the compositions provided herein induce the proliferation and/or accumulation of regulatory T cells in the subject. As will be evident to one of ordinary skill in the art, regulatory T cells, also referred to as “Tregs,” are a subset of T lymphocytes that are generally thought to suppress an abnormal or excessive immune response and play a role in immune tolerance. Regulatory T cells may be identified based expression of the markers Foxp3 and CD4 (Foxp3+ CD4+). The term regulatory T cells may also include Foxp3-negative regulatory T cells that are IL-10-producing CD4-positive T cells.

(210) In some embodiments, the therapeutically effective amount is an amount sufficient to induce the proliferation and/or accumulation of Tregs in the subject (or in a sample obtained from a subject) by at least 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 20-fold, 30-fold, 40-fold, 50-fold, 100-fold, 150-fold, 200-fold, 500-fold or more, as compared to the amount of Tregs in a subject (e.g., a subject with a pathogenic infection) that has not received any of the compositions described herein or as compared to a fecal sample from the same subject that was collected prior to administration of any of the compositions.

(211) As used herein, the phrase “induces proliferation and/or accumulation of regulatory T cells” refers to an effect of inducing the differentiation of immature T cells into regulatory T cells, which differentiation leads to the proliferation and/or the accumulation of regulatory T cells. Further, the meaning of “induces proliferation and/or accumulation of regulatory T cells” includes in vivo effects, in vitro effects, and ex vivo effects. In some embodiments, the proliferation and/or accumulation of regulatory T cells may be assessed by detecting and/or quantifying the number of cells that express markers of regulatory T cells (e.g., Foxp3 and CD4), for example by flow cytometry. In some embodiments, the proliferation and/or accumulation of regulatory T cells may be assessed by determining the activity of the regulatory T cells, such as the production of cytokines (e.g., IL-10).

(212) In some embodiments, the therapeutically effective amount is an amount sufficient to recolonize or repopulate the gastrointestinal tract of the subject with non-pathogenic bacteria. In some embodiments, the therapeutically effective amount is an amount sufficient to graft one or more of the bacterial strains of the composition in the gastrointestinal tract of the subject. In some embodiments, a fecal sample is obtained from the subject to assess the bacterial burden of the pathogenic infection and/or evaluate the efficacy of administration of the bacterial compositions described herein. In some embodiments, the microbiota of the subject (e.g., the identity and abundance of strains and/or species of the microbiota) may be assessed to determine a disease state of the subject and/or assess progress of the treatment. In some embodiments, the microbiota of the subject having a pathogenic infection is compared to the microbiota of a healthy subject, such as a subject that is not experiencing or has not experienced the pathogenic infection. In some embodiments, the microbiota of the subject having a

pathogenic infection is compared to the microbiota of the same subject from a fecal sample obtained from the subject prior to the pathogenic infection.

(213) Any of the compositions described herein, including the pharmaceutical compositions and food products comprising the compositions, may contain bacterial strains in any form, for example in an aqueous form, such as a solution or a suspension, embedded in a semi-solid form, in a powdered form or freeze dried form. In some embodiments, the composition or the bacterial strains of the composition are lyophilized. In some embodiments, a subset of the bacterial strains in a composition is lyophilized. Methods of lyophilizing compositions, specifically compositions comprising bacteria, are well known in the art. See, e.g., U.S. Pat. Nos. 3,261,761; 4,205,132; PCT Publications WO 2014/029578 and WO 2012/098358, herein incorporated by reference in their entirety. The bacteria may be lyophilized as a combination and/or the bacteria may be lyophilized separately and combined prior to administration. A bacterial strain may be combined with a pharmaceutical excipient prior to combining it with the other bacterial strain or multiple lyophilized bacteria may be combined while in lyophilized form and the mixture of bacteria, once combined may be subsequently be combined with a pharmaceutical excipient. In some embodiments, the bacterial strain is a lyophilized cake. In some embodiments, the compositions comprising the one or more bacterial strains are a lyophilized cake.

(214) The bacterial strains of the composition can be manufactured using fermentation techniques well known in the art. In some embodiments, the active ingredients are manufactured using anaerobic fermenters, which can support the rapid growth of anaerobic bacterial species. The anaerobic fermenters may be, for example, stirred tank reactors or disposable wave bioreactors. Culture media such as BL media and EG media, or similar versions of these media devoid of animal components, can be used to support the growth of the bacterial species. The bacterial product can be purified and concentrated from the fermentation broth by traditional techniques, such as centrifugation and filtration, and can optionally be dried and lyophilized by techniques well known in the art.

(215) In some embodiments, the composition of bacterial strains may be formulated for administration as a pharmaceutical composition. The term “pharmaceutical composition” as used herein means a product that results from the mixing or combining of at least one active ingredient, such as any two or more purified bacterial strains described herein, and one or more inactive ingredients, which may include one or more pharmaceutically acceptable excipient.

(216) An “acceptable” excipient refers to an excipient that must be compatible with the active ingredient and not deleterious to the subject to which it is administered. In some embodiments, the pharmaceutically acceptable excipient is selected based on the intended route of administration of the composition, for example a composition for oral or nasal administration may comprise a different pharmaceutically acceptable excipient than a composition for rectal administration. Examples of excipients include sterile water, physiological saline, solvent, a base material, an emulsifier, a suspending agent, a surfactant, a stabilizer, a flavoring agent, an aromatic, an excipient, a vehicle, a preservative, a binder, a diluent, a tonicity adjusting agent, a soothing agent, a bulking agent, a disintegrating agent, a buffer agent, a coating agent, a lubricant, a colorant, a sweetener, a thickening agent, and a solubilizer.

(217) Pharmaceutical compositions of the invention can be prepared in accordance with methods well known and routinely practiced in the art (see e.g., Remington: The Science and Practice of Pharmacy, Mack Publishing Co. 20th ed. 2000). The pharmaceutical compositions described herein may further comprise any carriers or stabilizers in the form of a lyophilized formulation or an aqueous solution. Acceptable excipients, carriers, or stabilizers may include, for example, buffers, antioxidants, preservatives, polymers, chelating reagents, and/or surfactants. Pharmaceutical compositions are preferably manufactured under GMP conditions. The pharmaceutical compositions can be used orally, nasally or parenterally, for instance, in the form of capsules, tablets, pills, sachets, liquids, powders, granules, fine granules, film-coated preparations, pellets, troches, sublingual preparations, chewables, buccal preparations, pastes, syrups, suspensions, elixirs, emulsions, liniments, ointments, plasters, cataplasms, transdermal absorption systems, lotions, inhalations, aerosols, injections, suppositories, and the like.

(218) In some embodiments, the bacteria are formulated for delivery to the intestines (e.g., the small intestine and/or the colon). In some embodiments, the bacteria are formulated with an enteric coating that increases the survival of the bacteria through the harsh environment in the stomach. The enteric coating is one which resists the action of gastric juices in the stomach so that the bacteria which are incorporated therein will pass through the stomach and into the intestines. The enteric coating may readily dissolve when in contact with intestinal fluids, so that the bacteria enclosed in the coating will be released in the intestinal tract. Enteric coatings may consist of polymer and copolymers well known in the art, such as commercially available EUDRAGIT (Evonik Industries). (See e.g., Zhang, *AAPS PharmSciTech*, (2016) 17 (1), 56-67).

(219) The bacteria may also be formulated for rectal delivery to the intestine (e.g., the colon). Thus, in some embodiments, the bacterial compositions may be formulated for delivery by suppository, colonoscopy, endoscopy, sigmoidoscopy or enema. A pharmaceutical preparation or formulation and particularly a pharmaceutical preparation for oral administration, may include an additional component that enables efficient delivery of the compositions of the disclosure to the intestine (e.g., the colon). A variety of pharmaceutical preparations that allow for the delivery of the compositions to the intestine (e.g., the colon) can be used. Examples thereof include pH sensitive compositions, more specifically, buffered sachet formulations or enteric polymers that release their contents when the pH becomes alkaline after the enteric polymers pass through the stomach. When a pH sensitive composition is used for formulating the pharmaceutical preparation, the pH sensitive composition is preferably a polymer whose pH threshold of the decomposition of the composition is between about 6.8 and about 7.5. Such a numeric value range is a range in which the pH shifts toward the alkaline side at a distal portion of the stomach, and hence is a suitable range for use in the delivery to the colon. It should further be appreciated that each part of the intestine (e.g., the duodenum, jejunum, ileum, cecum, colon and rectum), has different biochemical and chemical environment. For instance, parts of the intestines have different pHs, allowing for targeted delivery by compositions that

have a specific pH sensitivity. Thus, the compositions provided herein may be formulated for delivery to the intestine or specific parts of the intestine (e.g., the duodenum, jejunum, ileum, cecum, colon and rectum) by providing formulations with the appropriate pH sensitivity. (See e.g., Villena et al., *Int J Pharm* 2015, 487 (1-2): 314-9).

(220) Another embodiment of a pharmaceutical preparation useful for delivery of the compositions to the intestine (e.g., the colon) is one that ensures the delivery to the colon by delaying the release of the contents (e.g., the bacterial strains) by approximately 3 to 5 hours, which corresponds to the small intestinal transit time. In one embodiment of a pharmaceutical preparation for delayed release, a hydrogel is used as a shell. The hydrogel is hydrated and swells upon contact with gastrointestinal fluid, with the result that the contents are effectively released (released predominantly in the colon). Delayed release dosage units include drug-containing compositions having a material which coats or selectively coats a drug or active ingredient to be administered. Examples of such a selective coating material include in vivo degradable polymers, gradually hydrolyzable polymers, gradually water-soluble polymers, and/or enzyme degradable polymers. A wide variety of coating materials for efficiently delaying the release is available and includes, for example, cellulose-based polymers such as hydroxypropyl cellulose, acrylic acid polymers and copolymers such as methacrylic acid polymers and copolymers, and vinyl polymers and copolymers such as polyvinylpyrrolidone.

(221) Additional examples of pharmaceutical compositions that allow for the delivery to the intestine (e.g., the colon) include bioadhesive compositions which specifically adhere to the colonic mucosal membrane (for example, a polymer described in the specification of U.S. Pat. No. 6,368,586) and compositions into which a protease inhibitor is incorporated for protecting particularly a biopharmaceutical preparation in the gastrointestinal tracts from decomposition due to an activity of a protease.

(222) Another example of a system enabling the delivery to the intestine (e.g., the colon) is a system of delivering a composition to the colon by pressure change in such a way that the contents are released by utilizing pressure change caused by generation of gas in bacterial fermentation at a distal portion of the stomach. Such a system is not particularly limited, and a more specific example thereof is a capsule which has contents dispersed in a suppository base and which is coated with a hydrophobic polymer (for example, ethyl cellulose).

(223) A further example of a system enabling the delivery of a composition to the intestine (e.g., the colon), is a composition that includes a coating that can be removed by an enzyme present in the gut (e.g., the colon), such as, for example, a carbohydrate hydrolase or a carbohydrate reductase. Such a system is not particularly limited, and more specific examples thereof include systems which use food components such as non-starch polysaccharides, amylose, xanthan gum, and azopolymers.

(224) The compositions provided herein can also be delivered to specific target areas, such as the intestine, by delivery through an orifice (e.g., a nasal tube) or through surgery. In addition, the compositions provided herein that are formulated for delivery to a specific area (e.g., the cecum or the colon), may be administered by a tube (e.g., directly into the small intestine). Combining mechanical delivery methods such as tubes with chemical delivery methods such as pH specific coatings, allow for the delivery of the compositions provided herein to a desired target area (e.g., the cecum or the colon).

(225) The compositions comprising bacterial strains are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art. Dosage regimens are adjusted to provide the optimum desired response (e.g., the prophylactic or therapeutic effect). In some embodiments, the dosage form of the composition is a tablet, pill, capsule, powder, granules, solution, or suppository. In some embodiments, the pharmaceutical composition is formulated for oral administration. In some embodiments, the pharmaceutical composition is formulated such that the bacteria of the composition, or a portion thereof, remain viable after passage through the stomach of the subject. In some embodiments, the pharmaceutical composition is formulated for rectal administration, e.g. as a suppository. In some embodiments, the pharmaceutical composition is formulated for delivery to the intestine or a specific area of the intestine (e.g., the colon) by providing an appropriate coating (e.g., a pH specific coating, a coating that can be degraded by target area specific enzymes, or a coating that can bind to receptors that are present in a target area).

(226) Dosages of the active ingredients in the pharmaceutical compositions of the present invention can be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired pharmaceutical response for a particular subject, composition, and mode of administration, without being toxic or having an adverse effect on the subject. The selected dosage level depends upon a variety of factors including the activity of the particular compositions of the present invention employed, the route of administration, the time of administration, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the subject being treated, and like factors.

(227) A physician, veterinarian or other trained practitioner, can start doses of the pharmaceutical composition at levels lower than that required to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect (e.g., treatment of a pathogenic infection, reduction of bacterial burden of pathogenic infection, reduction or inhibition of toxin production) is achieved. In general, effective doses of the compositions of the present invention, for the prophylactic treatment of groups of people as described herein vary depending upon many different factors, including routes of administration, physiological state of the subject, whether the subject is human or an animal, other medications administered, and the therapeutic effect desired. Dosages need to be titrated to optimize safety and efficacy. In some embodiments, the dosing regimen entails oral administration of a dose of any of the compositions described herein. In some embodiments, the dosing regimen entails oral administration of multiple doses of any of the compositions described herein. In some embodiments, the composition is administered orally the subject once, twice, 3 times, 4 times, 5 times, 6 times, 7 times, 8 times, 9 times, or at least 10 times.

(228) The compositions, including the pharmaceutical compositions disclosed herein, include compositions with a range of active ingredients (e.g., live bacteria, bacteria in spore format). The amount of bacteria in the compositions may be expressed

and nutrient products are, in general, intended for the consumption of a human or an animal. Any of the bacterial strains described herein may be formulated as a food product. In some embodiments, the bacterial strains are formulated as a food product in spore form. In some embodiments, the bacterial strains are formulated as a food product in vegetative form. In some embodiments, the food product comprises both vegetative bacteria and bacteria in spore form. The compositions disclosed herein can be used in a food or beverage, such as a health food or beverage, a food or beverage for infants, a food or beverage for pregnant women, athletes, senior citizens or other specified group, a functional food, a beverage, a food or beverage for specified health use, a dietary supplement, a food or beverage for patients, or an animal feed. Non-limiting examples of the foods and beverages include various beverages such as juices, refreshing beverages, tea beverages, drink preparations, jelly beverages, and functional beverages; alcoholic beverages such as beers; carbohydrate-containing foods such as rice food products, noodles, breads, and pastas; paste products such as fish hams, sausages, paste products of seafood; retort pouch products such as curries, food dressed with a thick starchy sauces, soups; dairy products such as milk, dairy beverages, ice creams, cheeses, and yogurts; fermented products such as fermented soybean pastes, yogurts, fermented beverages, and pickles; bean products; various confectionery products such as Western confectionery products including biscuits, cookies, and the like, Japanese confectionery products including steamed bean-jam buns, soft adzuki-bean jellies, and the like, candies, chewing gums, gummies, cold desserts including jellies, cream caramels, and frozen desserts; instant foods such as instant soups and instant soy-bean soups; microwavable foods; and the like. Further, the examples also include health foods and beverages prepared in the forms of powders, granules, tablets, capsules, liquids, pastes, and jellies.

(233) Food products containing bacterial strains described herein may be produced using methods known in the art and may contain the same amount of bacteria (e.g., by weight, amount or CFU) as the pharmaceutical compositions provided herein. Selection of an appropriate amount of bacteria in the food product may depend on various factors, including for example, the serving size of the food product, the frequency of consumption of the food product, the specific bacterial strains contained in the food product, the amount of water in the food product, and/or additional conditions for survival of the bacteria in the food product.

(234) Examples of food products which may be formulated to contain any of the bacterial strains described herein include, without limitation, a beverage, a drink, a bar, a snack, a dairy product, a confectionery product, a cereal product, a ready-to-eat product, a nutritional formula, such as a nutritional supplementary formulation, a food or beverage additive.

(235) In some embodiments, the subject has not received a dose of an antibiotic prior to administration of the bacterial composition. In some embodiments, the subject has not been administered an antibiotic at least 1, at least 2, at least 3, at least 5, at least 10, at least 15, at least 20, at least 25, at least 30, at least 60, at least 90, at least 120, at least 180 or at least 360 days prior to administration of the compositions provided herein. In some embodiments, the person has not been administered and antibiotic to treat the pathogenic infection. In some embodiments, the compositions provided herein comprise the first treatment of the pathogenic infection.

(236) In some embodiments, the subject may be administered one or more doses of an antibiotic prior to or concurrently with a bacterial composition. Generally, the first line of defense in the treatment of a pathogenic infection is the administration of an antibiotic. In some embodiments, the subject is administered a single dose of an antibiotic prior to the bacterial composition. In some embodiments, the subject is administered multiple doses of an antibiotic prior to the bacterial composition. In some embodiments, the subject is administered at least 2, 3, 4, 5 or more doses of an antibiotic prior to the bacterial composition. In some embodiments, the subject is administered a dose of an antibiotic at substantially the same time as the bacterial composition. Examples of antibiotics that can be administered include, without limitation, kanamycin, gentamicin, colistin, metronidazole, vancomycin, clindamycin, fidaxomicin, and cefoperazone.

(237) Table 1 below provides sequence identifier numbers (SEQ ID NOs) used in the compositions of the experiments disclosed herein, along with the accompanying strain identification number (Strain ID). The closest bacterial species to the indicated strain is presented by genus-species. The 16S rDNA sequence associated with each genus species identified as the closest related genus species is also provided. The percent alignment presents the percent identity between the sequence of the indicated strain with the sequence from the closest genus species and the length of the alignment. The GenBank Accession Number of the closest related species is provided in the last column.

(238) TABLE-US-00011 TABLE 1 Closest bacterial species to the strains described herein SEQ ID NO. Accession # of closest Percent Alignment of closest SEQ ID Strain ID Closest Genus_species species alignment length species SEQ ID 71
Blautia_wexlerae SEQ_94 96.62 207 NR_044054 NO: 01 SEQ ID 102 *Turicibacter_sanguinis* SEQ_91 97.81 183 NR_028816 NO: 02 SEQ ID 5 *Clostridium_hathewayi* SEQ_105 92.42 198 NR_036928 NO: 03 SEQ ID 7
Blautia_hansenii SEQ_99 96.62 207 NR_104687 NO: 04 SEQ ID 10 *Blautia_hansenii* SEQ_99 98.06 206 NR_104687 NO: 05 SEQ ID 40 *Lactobacillus_mucosae* SEQ_90 87.57 185 NR_024994 NO: 06 SEQ ID 59 *Blautia_producta* SEQ_106 98.54 206 NR_113270 NO: 07 SEQ ID 59 *Blautia_coccoides* SEQ_103 98.54 206 NR_104700 NO: 07 SEQ ID 79
Blautia_hansenii SEQ_99 100 194 NR_104687 NO: 08 SEQ ID VE202- *Eubacterium_contortum* SEQ_109 94.59 296 NR_117147 NO: 09 21 SEQ ID VE202- *Eubacterium_fissicatena* SEQ_108 94.59 296 NR_117142 NO: 09 21 SEQ ID 211
Flavonifractor_plautii SEQ_93 98.49 199 NR_043142 NO: 10 SEQ ID VE202-9 *Anaerostipes_caccae* SEQ_88 99.5 399 NR_028915 NO: 11 SEQ ID VE202- *Clostridium_scindens* SEQ_87 95.76 354 NR_028785 NO: 12 26 SEQ ID 136
Marvinbryantia_formatexigens SEQ_89 94.66 131 NR_042152 NO: 13 SEQ ID VE202- *Anaerotruncus_colihominis* SEQ_95 99.34 1365 NR_027558 NO: 14 13 SEQ ID VE202- *Eubacterium_fissicatena* SEQ_102 93.33 1530 NR_117563 NO: 15 14 SEQ ID VE202- *Clostridium_symbiosum* SEQ_122 98.43 1469 NR_118730 NO: 16 16 SEQ ID VE202-7
Clostridiumbolteae SEQ_110 99.86 1390 NR_113410 NO: 17 SEQ ID 148 *Dorea_longicatena* SEQ_97 99.7 1318 NR_028883 NO: 18 SEQ ID 16 *Blautia_producta* SEQ_106 98.33 1493 NR_113270 NO: 19 SEQ ID 170
Dorea_longicatena SEQ_97 99.7 1318 NR_028883 NO: 20 SEQ ID 189 *Clostridium_innocuum* SEQ_98 98.64 1476

NR_029164 NO: 21 Dorea_longicatena SEQ_97 99.58 475 NR_028883 NO: 22 SEQ ID VE202-
Eisenbergiella_tayi SEQ_121 100 354 NR_118643 NO: 23 29 SEQ ID YK96 Dorea_longicatena SEQ_97 99.48 191
NR_028883 NO: 24 SEQ ID YK101 Ruminococcus_obeum SEQ_85 96.81 188 NR_118692 NO: 25 SEQ ID YK110
Megasphaera_elsdenii SEQ_119 96.62 207 NR_102980 NO: 26 SEQ ID YK149 Acidaminococcus_fermentans SEQ_115
99.48 192 NR_074928 NO: 27 SEQ ID YK149 Acidaminococcus_intestini SEQ_112 99.48 192 NR_074306 NO: 27 SEQ ID
YK154 Megasphaera_elsdenii SEQ_119 96.12 206 NR_102980 NO: 28 SEQ ID YK36 Ruminococcus_faecis SEQ_96 99.29
425 NR_116747 NO: 29 SEQ ID YK95 Bacteroides_cellulosilyticus SEQ_100 99.54 437 NR_112933 NO: 30 SEQ ID YK32
Anaerostipes_hadrus SEQ_107 98.8 415 NR_104799 NO: 31 SEQ ID YK64 Ruminococcus_obeum SEQ_84 99.04 415
NR_119185 NO: 32 SEQ ID YK73 Flavonifractor_plautii SEQ_93 98.56 418 NR_043142 NO: 33 SEQ ID YK87
Eubacterium_rectale SEQ_114 99.52 416 NR_074634 NO: 34 SEQ ID YK105 Flavonifractor_plautii SEQ_93 99.26 407
NR_043142 NO: 35 SEQ ID YK153 Megasphaera_elsdenii SEQ_119 96.04 429 NR_102980 NO: 36 SEQ ID YK163
Eubacterium_rectale SEQ_114 99.76 415 NR_074634 NO: 37 SEQ ID YK191 Ruminococcus_champanellensis SEQ_117
94.47 416 NR_102884 NO: 38 SEQ ID YK191 Ruminococcus_albus SEQ_113 94.47 416 NR_074399 NO: 38 SEQ ID
YK99 Ruminococcus_champanellensis SEQ_117 97.28 184 NR_102884 NO: 39 SEQ ID YK55 Ruminococcus_faecis
SEQ_96 99.02 408 NR_116747 NO: 40 SEQ ID YK75 Bifidobacterium_bifidum SEQ_118 99.45 183 NR_102971 NO: 41
SEQ ID YK90 Anaerostipes_hadrus SEQ_107 98.97 194 NR_104799 NO: 42 SEQ ID YK30 Anaerostipes_hadrus
SEQ_107 99.48 191 NR_104799 NO: 43 SEQ ID YK31 Anaerostipes_hadrus SEQ_107 98.97 194 NR_104799 NO: 44 SEQ
ID YK12 Eubacterium_rectale SEQ_114 99.27 412 NR_074634 NO: 45 SEQ ID YK27 Ruminococcus_faecis SEQ_96 99.51
412 NR_116747 NO: 46 SEQ ID YK28 Blautia_luti SEQ_111 99.5 400 NR_041960 NO: 47 SEQ ID YK29
Ruminococcus_faecis SEQ_96 99.03 413 NR_116747 NO: 48 SEQ ID YK33 Anaerostipes_hadrus SEQ_107 99.27 413
NR_104799 NO: 49 SEQ ID YK34 Anaerostipes_hadrus SEQ_107 99.51 410 NR_104799 NO: 50 SEQ ID YK35
Ruminococcus_faecis SEQ_96 99.51 409 NR_116747 NO: 51 SEQ ID YK51 Eubacterium_rectale SEQ_114 99.27 413
NR_074634 NO: 52 SEQ ID YK52 Eubacterium_rectale SEQ_114 99.03 413 NR_074634 NO: 53 SEQ ID YK54
Anaerostipes_hadrus SEQ_107 85.82 409 NR_104799 NO: 54 SEQ ID YK56 Ruminococcus_faecis SEQ_96 99.03 413
NR_116747 NO: 55 SEQ ID YK57 Ruminococcus_faecis SEQ_96 98.79 413 NR_116747 NO: 56 SEQ ID YK58
Dorea_longicatena SEQ_97 98.8 417 NR_028883 NO: 57 SEQ ID YK65 Roseburia_faecis SEQ_92 99.27 413 NR_042832
NO: 58 SEQ ID YK67 Blautia_luti SEQ_111 98.57 419 NR_041960 NO: 59 SEQ ID YK69
Fusicatenibacter_saccharivorans SEQ_116 99.27 413 NR_114326 NO: 60 SEQ ID YK70 Fusicatenibacter_saccharivorans
SEQ_116 98.79 414 NR_114326 NO: 61 SEQ ID YK71 Roseburia_faecis SEQ_92 99.28 414 NR_042832 NO: 62 SEQ ID
YK74 Megasphaera_elsdenii SEQ_119 96.06 431 NR_102980 NO: 63 SEQ ID YK88 Eubacterium_rectale SEQ_114 99.28
415 NR_074634 NO: 64 SEQ ID YK89 Eubacterium_rectale SEQ_114 99.27 413 NR_074634 NO: 65 SEQ ID YK97
Roseburia_faecis SEQ_92 99.28 414 NR_042832 NO: 66 SEQ ID YK98 Blautia_faecis SEQ_104 98.02 405 NR_109014
NO: 67 SEQ ID YK139 Fusicatenibacter_saccharivorans SEQ_116 99.03 412 NR_114326 NO: 68 SEQ ID YK141
Dorea_formicigenerans SEQ_120 98.51 402 NR_044645 NO: 69 SEQ ID YK142 Ruminococcus_faecis SEQ_96 98.79 413
NR_116747 NO: 70 SEQ ID YK152 Blautia_hansenii SEQ_99 99.5 401 NR_104687 NO: 71 SEQ ID YK155
Blautia_hansenii SEQ_99 98.79 413 NR_104687 NO: 72 SEQ ID YK157 Eubacterium_rectale SEQ_114 99.27 413
NR_074634 NO: 73 SEQ ID YK160 Roseburia_faecis SEQ_92 99.03 414 NR_042832 NO: 74 SEQ ID YK166
Eubacterium_rectale SEQ_114 99.27 409 NR_074634 NO: 75 SEQ ID YK168 Eubacterium_rectale SEQ_114 99.27 413
NR_074634 NO: 76 SEQ ID YK169 Eubacterium_rectale SEQ_114 99.28 416 NR_074634 NO: 77 SEQ ID YK171
Eubacterium_rectale SEQ_114 97.87 188 NR_074634 NO: 78 SEQ ID YK192 Roseburia_faecis SEQ_92 99.03 414
NR_042832 NO: 79 SEQ ID VE202- Erysipelatoclostridium_ramosum SEQ_123 100 1485 NR_113243 NO: 80 18 SEQ ID
PE5 Clostridium_bolteae SEQ_110 100 1385 NR_113410 NO: 81 SEQ ID PE9 Clostridium_disporicum SEQ_86 99.21 382
NR_026491 NO: 82 SEQ ID 211-B Bacteroides_ovatus SEQ_101 95.64 436 NR_112940 NO: 83

(239) TABLE-US-00012 TABLE 2 Bacterial species with a high degree of homology based on whole genome analysis:
Strain Whole genome homology SEQ_10 - 211 Lachnospiraceae bacterium 7_1_58FAA Subdoligranulum Flavinoifractor
plautii SEQ_14 - VE202-13 Anaerotruncus_colihominis SEQ_15 - VE202-14 Eubacterium_fissicatena Ruminococcus
torques SEQ_16 - VE202-16 Clostridium_symbiosum SEQ_17 - VE202-7 Clostridium_bolteae SEQ_22 - 169/SEQ_20 - 170
Dorea_longicatena SEQ_19 - 16 Blautia_producta SEQ_21 - 189 Clostridium_innocuum
Erysipelotrichaceae bacterium_21_3

(240) TABLE-US-00013 TABLE 3 Bacterial species with highest degree of homology based on whole genome analysis SEQ
ID # of Closest *Consensus 16S region as species based SEQ ID # of SEQ ID # of determined on Sanger 16S regions as 16S
region as Composition B Strain by Sanger sequencing of determined determined strain number identifier sequencing 16S
region by WGS{circumflex over ()} by WGS 1 VE202-7 17 Clostridium 124, 125, 124 bolteae 126, 127, 128 2 VE202-13
14 Anaerotruncus 129,130, 129 colihominis 131 3 VE202-14 15 Eubacterium 132, 133, 132 fissicatena 134, 135, 136 4
VE202-16 16 Clostridium 137, 138, 137 symbiosum 139, 140 5 strain #16 19 Blautia 141, 142, 141 producta 143, 144, 145 6
strain #170 20 Dorea 146, 147, 146 longicatena 148, 149, 150, 151 7 strain #189 21 Clostridium 152, 153, 152 innocuum
154, 155, 156 8 strain #211 10 Flavinoifractor 157, 158, 157 plautii 159 Closest species based on Consensus Closest SEQ ID
of species based 16S region as on WGS Additional Composition B compared with compared versus closely related
Clostridium strain number 16S database WG databases sequences cluster 1 Clostridium Clostridium XIVa bolteae bolteae
90A9 2 Anaerotruncus Anaerotruncus IV colihominis colihominis DSM 17241 3 Dracourtella Dracourtella Ruminococcus
XIVa massiliensis massiliensis GD1 torques; Sellimonas intestinalis 4 Clostridium Clostridium XIVa symbiosum symbiosum
WAL- 14163 5 Blautia Clostridium Blautia product XIVa producta bacterium UC5.1- ATCC 27340 1D4 6 Dorea Dorea
XIVa longicatena longicatena CAG: 42 7 Clostridium Erysipelotrichaceae XVII innocuum bacterium 21_3 8 Flavinoifractor

Clostridium Subdoligranulum *IV plautii orbiscindens* 1_3_50AFAA {circumflex over ()}WGS refers to Whole Genome Sequencing performed on a PacBio Biosciences platform (Menlo Park, CA). *Consensus sequence is defined as the 16S sequence that has the most overlap with all other identified 16S sequences.

(241) In some embodiments, in any of the compositions described herein, *Clostridium bolteae* can be replaced with *Clostridium bolteae* 90A9. In some embodiments, in any of the compositions described herein, *Anaerotruncus colihominis* can be replaced with *Anaerotruncus colihominis* DSM 17241. In some embodiments, in any of the compositions described herein, *Eubacterium fissicatena* can be replaced with *Sellimonas instestinalis*, *Drancourtella massiliensis* or *Drancourtella massiliensis* GPI. In some embodiments, in any of the compositions described herein, *Clostridium symbiosum* can be replaced with *Clostridium symbiosum* WAL-14163. In some embodiments, in any of the compositions described herein, *Blautia producta* can be replaced with *Clostridium bacterium* CD5.1-1D4 or *Blautia product* ATCC27340. In some embodiments, in any of the compositions described herein, *Dorea longicatena* can be replaced with *Dorea longicatena* CAG:42. In some embodiments, in any of the compositions described herein, *Clostridium innocuum* can be replaced with *Erysipelotrichaceae bacterium* 21_3. In some embodiments, in any of the compositions described herein, *Flavonifractor plautii* can be replaced with *Clostridium orbiscindens* 1_3_50 AFAA.

(242) Aspects described herein provide pharmaceutical composition comprising a purified bacterial mixture consisting of bacterial strains comprising 16S rDNA sequences of at least 95% homology to SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21. In some aspects, the bacterial strains have at least 97% homology to SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21. In some aspects, the bacterial strains have at least 98% homology to SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21. In some aspects, the bacterial strains have at least 99% homology to SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21.

(243) In some aspects, at least a portion of the bacteria of the pharmaceutical composition are in spore-form. In some aspects, the pharmaceutical composition further comprises a pharmaceutically acceptable excipient.

(244) In some aspects, the pharmaceutical composition is formulated for oral administration. In some aspects, the pharmaceutical composition is in the form of a capsule. In some aspects, the pharmaceutical composition is formulated for delivery to the colon. In some aspects, the pharmaceutical composition further comprises a pH sensitive composition comprising one or more enteric polymers.

(245) Aspects described herein provide pharmaceutical compositions comprising a purified bacterial mixture consisting of bacterial strains comprising 16S rDNA sequences of at least 95% homology to SEQ ID NO:124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ ID NO:152, and SEQ ID NO:157. In some aspects, the bacterial strains have at least 97% homology to SEQ ID NO:124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ ID NO:152, and SEQ ID NO:157. In some aspects, the bacterial strains have at least 98% homology to SEQ ID NO:124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ ID NO:152, and SEQ ID NO:157. In some aspects, the bacterial strains have at least 99% homology to SEQ ID NO:124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ ID NO:152, and SEQ ID NO:157.

(246) In some aspects, at least a portion of the bacterial strains are in spore-form. In some aspects, the pharmaceutical composition further comprises a pharmaceutically acceptable excipient.

(247) In some aspects, the pharmaceutical composition is formulated for oral administration. In some aspects, the pharmaceutical composition is in the form of a capsule. In some aspects, the pharmaceutical composition is formulated for delivery to the colon. In some aspects, the pharmaceutical composition further comprises a pH sensitive composition comprising one or more enteric polymers.

(248) Aspects described herein provide pharmaceutical compositions comprising a purified bacterial mixture comprising bacterial strains comprising 16S rDNA sequences of at least 95% homology to SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21. In some aspects, the bacterial strains have at least 97% homology to SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21. In some aspects, the bacterial strains have at least 98% homology to SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21. In some aspects, the bacterial strains have at least 99% homology to SEQ ID NO:10, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:19, SEQ ID NO:20, and SEQ ID NO:21.

(249) In some aspects, at least a portion of the bacterial strains are in spore-form. In some aspects, the pharmaceutical composition further comprises a pharmaceutically acceptable excipient.

(250) In some aspects, the pharmaceutical composition is formulated for oral administration. In some aspects, the pharmaceutical composition is in the form of a capsule. In some aspects, the pharmaceutical composition is formulated for delivery to the colon. In some aspects, the pharmaceutical composition further comprises a pH sensitive composition comprising one or more enteric polymers.

(251) Aspects described herein provide pharmaceutical compositions comprising a purified bacterial mixture comprising bacterial strains comprising 16S rDNA sequences of at least 95% homology to SEQ ID NO:124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ ID NO:152, and SEQ ID NO:157. In some aspects, the bacterial strains have at least 97% homology to SEQ ID NO:124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ ID NO:152, and SEQ ID NO:157. In some aspects, the bacterial strains have at least 98% homology to SEQ ID NO:124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ

ID NO:152, and SEQ ID NO:157. In some aspects, the bacterial strains have at least 99% homology to SEQ ID NO:124, SEQ ID NO:129, SEQ ID NO:132, SEQ ID NO:137, SEQ ID NO:141, SEQ ID NO:146, SEQ ID NO:152, and SEQ ID NO:157. In some aspects, at least a portion of the bacterial strains are in spore-form. In some aspects, the pharmaceutical composition further comprises a pharmaceutically acceptable excipient.

(252) In some aspects, the pharmaceutical composition is formulated for oral administration. In some aspects, the pharmaceutical composition is in the form of a capsule. In some aspects, the pharmaceutical composition is formulated for delivery to the colon. In some aspects, the pharmaceutical composition further comprises a pH sensitive composition comprising one or more enteric polymers.

(253) Aspects described herein provide pharmaceutical compositions comprising a purified bacterial mixture consisting of the following bacterial strains: *Clostridium bolteae*, *Anaerotruncus colihominis*, *Sellimonas intestinalis*, *Clostridium symbiosum*, *Blautia producta*, *Dorea Longicatena*, *Erysipelotrichaceae bacterium*, and *Clostridium orbiscindens*.

(254) In some aspects, at least a portion of the bacterial strains are in spore-form. In some aspects, the pharmaceutical composition further comprises a pharmaceutically acceptable excipient.

(255) In some aspects, the pharmaceutical composition is formulated for oral administration. In some aspects, the pharmaceutical composition is in the form of a capsule. In some aspects, the pharmaceutical composition is formulated for delivery to the colon. In some aspects, the pharmaceutical composition further comprises a pH sensitive composition comprising one or more enteric polymers.

(256) Aspects described herein provide pharmaceutical compositions comprising a purified bacterial mixture comprising the following bacterial strains: *Clostridium bolteae*, *Anaerotruncus colihominis*, *Sellimonas intestinalis*, *Clostridium symbiosum*, *Blautia producta*, *Dorea Longicatena*, *Erysipelotrichaceae bacterium*, and *Clostridium orbiscindens*.

(257) In some aspects, at least a portion of the bacterial strains are in spore-form. In some aspects, the pharmaceutical composition further comprises a pharmaceutically acceptable excipient.

(258) In some aspects, the pharmaceutical composition is formulated for oral administration. In some aspects, the pharmaceutical composition is in the form of a capsule. In some aspects, the pharmaceutical composition is formulated for delivery to the colon. In some aspects, the pharmaceutical composition further comprises a pH sensitive composition comprising one or more enteric polymers.

(259) Aspects described herein provide methods of treating an infectious disease in a subject, the method comprising administering the pharmaceutical composition of any of the aspects described herein to the subject in an amount sufficient to treat the infectious disease. In some aspects, the infectious disease is *Clostridium difficile* infection.

(260) The nucleic acid sequences of the 16S rDNA, or portion thereof, for the bacterial strains described herein are provided below:

(261) TABLE-US-00014 >SEQ ID NO: 01|71|

GCCCCGAGCAGTTGATGTGAAGGATGGGTCACCTGTGGACTGCATTGGAAGTGCATACTTGAGTGCCGGAGGGTAA
GCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAGTGCGGAAGGCGGCTTACTGGACGGT
AACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTAA >SEQ ID NO: 02|102|
CTAACCGTGGAGGTCATTGGAAACTGGTCAACTTGAGTGCAGAAGAGGGGAAGTGAATTCATGTGTAGCGGTGAAA
TGCGTAGAGATATGGAGGAACACCAGTGCGGAAGGCGGCTTCCTGGTCTGTAAGTACACTGAGGCGCGAAAGCGTG
GGGGCAAACAGGATTAGATCCCCCGGTAA >SEQ ID NO: 03|5|

ATGAAAGCCGGGGCTCAACCCCGGTACTGCTTTGGAACTGTTTGACTTGAGTGCTTGAGAGGTAAGTGAATTCCT
AGTGTAGCGGGAAATGTTTAGATATTAGGAGGACACCAGTGCGGAAGGCGGCTTACTGGACTGTAAGTACGTTGTG
GCTCGATTGTGGGGAGCAAACAGGATTATATCCCCTGGTAA >SEQ ID NO: 04|7|

CGGAAGGTCTGATGTGAAGGTTGGGGCTTACCCCGGACTGCATTGGAACTGTTTTTCTAGAGTGCCCGAGAGGTAA
GCGGAATTCCTAGTGTAGCGGTGAAATGCTTTAGATATTAGGAGGAACACCAGTGCGGAAGGCGGCTTACTGGACGG
TAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTAA >SEQ ID NO: 05|10|
CGATGTCTGAGTGAAGGCTGGGGCTTACCCAGGACTGCATTGGAAGTGTCTTCTAGAGTGCCGGAGAGGTAAGCG
GAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAGTGCGGAAGGCGGCTTACTGGACGGTAAC
TGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTAA >SEQ ID NO: 06|40|

TTAACCAAGAAGTGCATTGGAAGTGCAGACTTGGGGGAAAAAAGACAGTGCAACTCCATGTGTAGCGGTGGAATG
CTCCATATATATGGAAGAACACCAGTGCGGAAGGCGGCTGTCTGGTCTGCAACTGACGCTGAGGCTCGAATTCATGG
GTAAGAAAGTATTAGTCCCTTGTA >SEQ ID NO: 07|59|

ACCCGCTTGGTCTGAGGTGAGGCTGGGGCTTAACCCAGGACTGCATTGGAACTGTTGTTCTAGAGTGCCGGAGAG
GTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAGTGCGGAAGGCGGCTTACTGGA
CGGTAAGTACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTAA >SEQ ID NO:
08|79|

TAGGCTGGGGCTTAACCCAGGACTGCATTGGAACTGTTTTTCTAGAGTGCCGGAGAGGTAAGCGGAATTCCTAGT
GTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAGTGCGGAAGGCGGCTTACTGGACGGTAAGTACGTTGAGG
CTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTAA >SEQ ID NO: 09|VE202-21|

TTGCATTGGACACTATGTCAGCTGAGTGTGCGAGAGGTAAGTGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAGTGCGGAAGGCGGCTTACTGCACGTTTTCTGACGTTGAGGCTCGAAATCGTGGGGAGCAAACA
AAAATAGATACCCTGGTAGTCCACGCCGTAAACGATGCATACTAGGTGTGCGGTGGCAAAGCCATTCGGTGCCGCAG
CAAACGCAATAAGTATGCCACCTGGGGAGTACGTTGCAAGAATGAACTCAAATAAATTGACGGA >SEQ
ID NO: 10|211|

CCCGTCTGATGTGAAGTGGGGGCTCACCTCCAGCCTGCATTTGAACTGTAGTTCTTGAGTGCTGGAGAGGCAAT

CGGAATTCCGTTGCTGAGTACGTAAGTACGTTAGGAGCAACAGGATTAGATACCTCATAA >SEQ ID NO: 11|VE202-9|
ACCTGATGCAGCGACGCCGCTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGAAGAAAAAGACG
GTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCCGGAAT
TACTGGGTGTAAAGGGTGCCTAGGTGGCATGGTAAGTCAGAAGTGAAAGCCCCGGGGCTTAACCCCGGGACTGCTTTT
GAAACTGTCATGCTGGAGTGCAGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGA
ACACCAGTGGCGAAGGCGGCTTACTGGACTGTCACTGACACTGATGCACGAAAGCGTGGGGAGCAAACAGGATTAGA
TACCCTGGAAGTCCAT >SEQ ID NO: 12|VE202-26|
ATGGGAGCGTAGATGGCGACTGGGCCATATGTGACAGCCCTGGTCTCAACCCCTTAACTGCATTTGGAAGTGAAGTGG
CTGGAGTGTCCGAGAGGCAGGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAGTGGCG
AAGGCGGCCTGCTGGACGATGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCTGGTAGT
CCACGCCGTAAACGATGACTACTAGGTGTCCGGTGGCAAGGACATTCGGTGCCGCAGCAAACGCAATAAGTAGTCCA
CCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAAATTGACGGA >SEQ ID NO: 13|136|
CGCAGCGGAGTGTATCCTAGGCTCACCTGGCTGCTTTTCAACTGGTTTTCTAGATCGTGTAGAGGGGGAGATTCCTG
GTGTAGCGTGAAATGCGTAGATATCTGGAGGAACACCAGTGGCGAAGGCGGCCTCCTGGACGGCAACTGACGTTGAG
GCTCGAAAGTGTGGGGAGCAAACAGGATTAGATACCTGGTAA >SEQ ID NO: 14|VE202-13|
TCAAAGAGTTTGATCCTGGCTCAGGACGAACGCTGGCGGCGCGCCTAACACATGCAAGTCGAACGGAGCTTACGTTT
TGAAGTTTTTCGGATGGACGAATGTAAGCTTAGTGGCGGACGGGTGAGTAACACGTGAGCAACCTGCCTTTCAGAGGG
GGATAACAGCCGGAACCGGCTGCTAATACCGCATGATGTTGCGGGGGCACATGCCCTGCAACCAAAGGAGCAATCC
GCTGAAAGATGGGCTCGCGTCCGATTAGCCAGTTGGCGGGGTAACGGCCACCAAAGCGACGATCGGTAGCCGGACT
GAGAGGTTGAACGGCCACATTGGGACTGAGACACGGCCAGACTCCTACGGGAGGCAGCAGTGGGGGATATTGCACA
ATGGGCGAAAGCCTGATGCAGCGACGCCGCTGAGGGAAGACGGTCTTCGGATTGTAAACCTCTGTCTTTGGGGAAG
AAAATGACGGTACCCAAAGAGGAAGCTCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGAGCAAGCGTT
GTCCGGAATTACTGGGTGTAAAGGGAGCGTAGGCGGGATGGCAAGTAGAATGTAAATCCATCGGCTCAACCGGTGG
CTGCGTTCTAAACTGCCGTTCTTGAGTGAAGTAGAGGCAGGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAGTGGCGAAGGCGGCCTGCTGGGCTTTAACTGACGCTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCTGGTAGTCCACGCCGTAAACGATGATTACTAGGTGTGGGGGGACTGACCCCTTCCGTGCCGCAG
TTAACACAATAAGTAATCCACCTGGGGAGTACGGCCGCAAGGTTGAAACTCAAAGGAATTGACGGGGGCCCGCACAA
GCAGTGGAGTATGTGGTTTAATTGCAAGCAACGCGAAGAACCTTACCAGGTCTTGACATCGGATGCATAGCCTAGAG
ATAGGTGAAGCCCTTCGGGGCATCCAGACAGGTGGTGCATGGTTGTCTCAGCTCGTGTCTGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCCTTATTATTAGTTGCTACGCAAGAGCACTCTAATGAGACTGCCGTTGACAAAACGGA
GGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGACCTGGGCTACACACGTACTACAATGGCACTAAAACAG
AGGGCGGCGACACCGCGAGGTGAAGCGAATCCCGAAAAAGTGTCTCAGTTCAGATTGCAGGCTGCAACCCGCTGCA
TGAAGTCGGAATTGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCCCGGGCCTTGACACACCGCCCGT
CACACCATGGGAGTCGGTAACACCCGAAGCCAGTAGCCTAACCGCAAGGGGGGCGCTGTGCAAGGTGGGATTGATGA
CTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID NO: 15|VE202-14|
TACGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAGCGAAGCGCTGTTTT
CAGAATCTTCGGAGGAAGAGGACAGTGAAGTGAAGCGGTGAGTAACGCGTGGGCAACCTGCCTCATACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGGACCGCATGGTGTAGTGTGAAAACTCCGGTG
GTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAAAGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAA
TGGGGGAAACCCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATTTTCGGTATGTAAACTTCTATCAGCAGGGAAGA
AGATGACGGTACCTGAGTAAGAAGCACCGGCTAAATACGTGCCAGCAGCCGCGGTAATACGTATGGTGCAAGCGTTA
TCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGATAGGCAAGTCTGGAGTGAAAACCCAGGGCTCAACCCCTGGGA
CTGCTTTGGAACTGCAGATCTGGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGGTGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCTGGTAGTCCACGCCGTAAACGATGACTACTAGGTGTCTGGTGTGCAAAGCACATCGGTGCCGCAG
CAAACGCAATAAGTAGTCCACCTGGGGAGTACGTTGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAA
GCGGTGGAGCATGTGGTTTAATTGCAAGCAACGCGAAGAACCTTACCTGGTCTTGACATCCGGATGACGGGCGAGTA
ATGTCGCCGTCCTTTCGGGGCATCCGAGACAGGTGGTGCATGGTTGTCTCAGCTCGTGTCTGTGAGATGTTGGGTAA
AGTCCCGCAACGAGCGCAACCCCTTATCTTCAGTAGCCAGCATATAAGGTGGGCACTCTGGAGAGACTGCCAGGGAGA
ACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGGCCAGGGCTACACACGTGCTACAATGGCGTA
AACAAAGGGAAGCGAGAGGGTGACCTGGAGCGAATCCCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCG
ACTACATGAAGCTGGAATCGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCCCGGGTCTTGACACACC
GCCCGTCACACCATGGGAGTCAGTAACGCCCGAAGCCAGTGACCCAACCTTAGAGGAGGGAGCTGTGCAAGGCGGGA
CGGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID NO: 16|VE202-16|
ATGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCGATTAAACG
GAAGTTTTTCGGATGGAAGTTGAATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGGTAACTGCCTTGTACTGGG
GGACAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTATCGCATGATACAGTGTGAAAACTCCGGTGG
TACAAGATGGACCCGCGTCTGATTAGCTAGTTGGTAAGGTAACGGCTTACCAAGGCGACGATCAGTAGCCGACCTGA

GAGGTCGACCGGACCATTTGGGACCTAGAGACAGGCCCAAACCTACCTACGGGAGGCGACGAGTGGGGGGAATATTGCACAAT
GGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGGAAGAA
AATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGGCAAGCGTTAT
CCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGTAAAGCAAGTCTGAAGTGAAAGCCCCGCGGCTCAACTGCGGGAC
TGCTTTGGAACTGTTTAACTGGAGTGTTCGGAGAGGTAAGTGGAAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCAGTGGCGAAGGCGACTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAC
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTTGGGGAGCAAAGCTCTTCGGTGCCGTTCG
AAACGCAGTAAGTATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAGGTCTTGACATCGATCCGACGGGGGAGTAA
CGTCCCCCTTCCCTTCGGGGCGGAGAAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCTTATTCTAAGTAGCCAGCGGTTTCGGCCGGGAACCTCTTGGGAGACTGCCAGGGATAA
CCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCCTTATGATCTGGGCTACACACGTGCTACAATGGCGTAA
ACAAAGAGAAGCAAGACCGCGAGGTGGAGCAAATCTCAAAAATAACGTCTCAGTTCGGACTGCAGGCTGCAACTCGC
CTGCACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCG
CCCGTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCGCAAGGAGGGAGCTGCCGAAGGCGGGAC
GATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 17|VE202-7|

ATGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAACGAAGCAATTAAAT
GAAGTTTTTCGGATGGATTTTTGATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGATAACCTGCCTCACACTGGG
GGATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTACCGCATGGTACGGTGTGAAAACTCCGGTGG
TGTGAGATGGATCCGCGTCTGATTAGCCAGTTGGCGGGGTAACGGCCACCAAAGCGACGATCAGTAGCCGACCTGA
GAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCTACGGGAGGCAGCAGTGGGGAATATTGCACAAT
GGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGGAAGAA
AATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGGCAAGCGTTAT
CCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCGAAGCAAGTCTGAAGTGAAACCCAGGGCTCAACCCTGGGAC
TGCTTTGGAACTGTTTTGCTAGAGTGTTCGGAGAGGTAAGTGGAAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAC
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATGCTAGGTGTTGGGGGGCAAAGCCCTTCGGTGCCGTTCG
AAACGCAGTAAGCATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCTCTTGACCGGCGTGTA
CGGCGCCTTCCCTTCGGGGCAAAGAGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCTTATCCTTAGTAGCCAGCAGGTAAAGCTGGGCACTCTAGGGAGACTGCCAGGGATA
ACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTA
AACAAAGGAAGCAAGACAGTGATGTGGAGCAAATCCCAAAAATAACGTCCCAGTTCGGACTGTAGTCTGCAACCCG
ACTACACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTACACACC
GCCCGTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCTCGCAAGAGAGGGAGCTGCCGAAGGCGGG
GCAGGTAACCTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 18|148|

AACGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTTTGGA
AGATTCTTCGGATGAAGACTTTTTGTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATACAGGG
GGATAACAGTTAGAAATGACTGCTAATACCGCATAAGACCACGGTACCGCATGGTACAGTGGTAAAACTCCGGTGG
TATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAACGGCCTACCAAGCCGACGATCAGTAGCCGACCTGA
GAGGGTGACCGGCCACATTGGGACTGAGACACGGCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAAT
GGAGGAAACTCTGATGCAGCGACGCCGCGTGAAGGATGAAGTATTTTCGGTATGTAAACTTCTATCAGCAGGGGAAGAA
AATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGGCAAGCGTTAT
CCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCACGGCAAGCCAGATGTGAAAGCCCCGGGGCTCAACCCCGGGAC
TGCATTTGGAACCTGCTGAGCTAGAGTGTTCGGAGAGGCAAGTGGAAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCAGTGGCGAAGGCGGCTTGCTGGACGATGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAC
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGACTGCTAGGTGTCGGGTGGCAAAGCCATTTCGGTGCCGCAGC
TAACGCAATAAGCAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCTGATCTTGACATCCCGATGACCGCTTCGTAA
TGGAAGCTTTTCTTCGGAACATCGGTGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCTTATCTTCAGTAGCCAGCAGGTAAAGCTGGGCACTCTGGAGAGACTGCCAGGGATA
ACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCCTTATGACCAGGGCTACACACGTGCTACAATGGCGTA
AACAAAGAGAAGCGAAGTTCGCGAGGGTAAGCAAATCTCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCG
ACTACATGAAGCTGGAATCGCTAGTAATCGCAGATCAGAATGCTGCGGTGAATACGTTCCCGGGTCTTGTACACACC
GCCCGTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCGTAAGGAGGGAGCTGCCGAAGGTGGGAC
CGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 19|16|

ATCAGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTTAAG
TGATCTCTTCGGATTGAACTTATTTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATACAG
GGGGATAACAGTTAGAAATGGCTGCTAATACCGCATAAGCGCACAGGACCGCATGGTCTGGTGTGAAAACTCCGGT
GGTATGAGATGGACCCGCGTCTGATTAGCTAGTTGGAGGGGTAACGGCCACCAAAGGCGACGATCAGTAGCCGGCCT

TCGGTATGTTAAACCTTCTATCAGGTAAGGGAAGAAATGACGGTACCTGACTGACGAAGACCCGGGCTAACTACGTGCCAGC
AGCCGCGGTAATACGTAGGGGGGCAAGCGTTATCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCACGGCAAGCC
AGATGTGAAAGCCC >SEQ ID NO: 23|VE202-29|
CAGGCTGGAGTGCAGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCCAGT
GGCGAAGGCGGCTTACTGGACTGTAAGTACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGG
TAGTCCACGCGGTAAACGATGATTGCTAGGTGTAGGTGGGTATGGACCCATCGGTGCCGAGCTAACGCAATAAGCA
ATCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGCGGTGGAGCATGTG
GTTTAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCC >SEQ ID NO: 24|YK96|
CCGGGGCTCACCCCGGGACTGCATTTGGAAGTCTGAGCTAGAGTGTTCGGAGAGGCAAGTGAATTCCTAGTGTAGC
GGTGAAATGCGTAGATATTAGGAGGAACACCCAGTGGCGAAGGCGGCTTGCTGGACGATGACTGACGTTGAGGCTCGA
AAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTA >SEQ ID NO: 25|YK101|
AGGGTCAACCCCTGGACTGCATTGGAACTGTCAGGCTGGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGG
TGAAATGCGTAGATATTAGGAGGAACACCCAGTGGCGAAGGCGGCTTACTGGACGGTAAGTACGTTGATGCTCGAAA
GCGTGGGGAGCAAACAGGATTAGATAACCTGGTAAA >SEQ ID NO: 26|YK110|
GGGAAGTCGGTCTTAAGTGCGGGGCTTAACCCCGTGAGGGGACCGAAACTGTGAAGCTCGAGTGTTCGGAGAGGAAAG
CGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCCAGTGGCGAAAGCGGCTTTCTGGACGACA
ACTGACGCTGAGGCGCGAAAGCCAGGGGAGCAAACGGGATTAGATACCCCAAGTAA >SEQ ID NO:
27|YK149|
TAGTCTGAGTGATGCGGGGCTTAACCCCGTATGGCGTTGGATACTGGAAGTCTTGAGTGCAGGAGAGGAAAGGGGAA
TTCCCACTGTAGCGGTGAAATGCGTAGATATTGGGAGGAACACCCAGTGGCGAAGGCGCCTTTCTGGACTGTGTCTGA
CGCTGAGATGCGAAAGCCAGGGTAGCAAACGGGATTAGATACCACGGTA >SEQ ID NO: 28|YK154|
GATAGTCGGTCTTAAGTGCGGGGCTTACCCCGTGAGGGGACCGAAACTGTGAAGCTCGAGTGTTCGGAGAGGAAAGCG
GAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCCAGTGGCGAAAGCGGCTTTCTGGACGACAAC
TGACGCTGAGGCGCGAAAGCCAGGGGAGCAAACGGGATTAGATACCACGGTAA >SEQ ID NO: 29|YK36|
CGTTTGCTCCACGCTTTTCGAGCCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCCTCCTAATA
TCTACGCATTTACCCGCTACACTAGGAATTCGCTTACCTCTCCGGTACTCTAGATTGACAGTTTCCAATGCAGTCC
CGGGGTTGAGCCCCGGGTTTTACATCAGACTTGCCACTCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAA
CGCTTGACCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTTTCT
TCCCTGCTGATAGAGCTTTACATACCGAAATACTTCATCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTG
TGCAATATTCCCCACTGCTGCCTCCCGTAGGAGTTTGGA >SEQ ID NO: 30|YK95|
TGTCACACTTTTCGAGCATCAGCGTCAGTTACAGTCCAGTAAGCTGCCTTCGCAATCGGAGTTCTTCGTGATATCTAA
GCATTTACCCGCTACACCACGAATTCGCGCTACCTCTACTGCACTCAAGACGACCAGTATCAACTGCAATTTTACGG
TTGAGCCGCAAACCTTTCACAGCTGACTTAATAGTCCGCTACGCTCCCTTTAAACCCAATAAATCCGGATAACGCTT
GGATCCTCCGTATTACCGCGGCTGCTGGCACGGAGTTAGCCGATCCTTATTCGTATGGTACATACAAAAGCCACAC
GTGGCTCACTTTATTCCCATATAAAAGAAGTTTACAACCCATAGGGCAGTCATCCTTCACGCTACTTGGCTGGTTCA
GACTCTCGTCCATTGACCAATATTCTCACTGCTGCCTCCCGTAGGTAGTTTGGA >SEQ ID NO: 31|YK32|
CCGTTGTCACGCTTTTCGTGCTCAGTGTCAGTTTCAGTCCAGTAAGCCGCCTTCGCCACTGATGTTCTCCTAATATC
TACGCATTTACCCGCTACACTAGGAATTCGCTTACCTCTCCTGCACTCCAGTCTGACAGTTTCAAAGCAGTCCCA
GAGTTAAGCCCTGGGTTTTCACTTCTGACTTGCCATACCACCTACGCACCCTTTACACCCAGTAAATCCGGATAACG
CTTGCCCCCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTTTCTTC
CCTGCTGATAGAGCTTTACATACCGAAATACTTCTTCACTCACGCGGCGTCGCTGCATCAGGGTTCCCCCATTGTG
CAATATTCCCCACTGCTGCCTCCCGTGGAAGTTTGGA >SEQ ID NO: 32|YK64|
GCGAATGTCACGCATTCGAGCCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCTCCTAATAT
CTACGCATTTACCCGCTACACTAGGAATTCGCTTACCTCTCCGGCACTCAAGACTAACAGTTTCCAATGCAGTCCA
GGGGTTGAGCCCCCGCCTTTCACATCAGACTTGCCAGTCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAAC
GCTTGCCCCCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCACTATCTT
CCCTGCTGATAGAAGTTTACATACCGAGATACTTCTTCTTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGT
GCAATATTCCCCACTGCTGCCTCCCGTGGGAGTTTGGA >SEQ ID NO: 33|YK73|
TGCTCACGCTTTTCGCGCTCAGCGTCAGTTACTGTCCAGCAATCCGCCTTCGCCACTGGTGTTCTCCTCGTATATCTAC
GCATTTACCCGCTACACACGGAATTCGATTGCCTCTCCAGCACTCAAGAACTACAGTTTCAAATGCAGGCTGGAGG
TTGAGCCCCCAGTTTTACATCTGACTTGCAATCCCGCCTACACGCCCTTTACACCCAGTAAATCCGGATAACGCTT
GCCACCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGTGGCTTATTCGTCAGGTACCGTCATTTGTTTCGTC
CCTGACAAAAGAAGTTTACAACCCGAAAGCCTTCTTCTTCACGCGGCGTTGCTGGGTCAGGCTTGCGCCCATTGCC
CAATATTCCCCACTGCTGCCTCCCGTGGTAGTTTGGA >SEQ ID NO: 34|YK87|
TGTCACGCTTTTCGAGCTCAGCGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCTCCTAATATCTAC
GCATTTACCCGCTACACTAGGAATTCGCTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGG
TTGAGCCCCGGGCTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTT
GCACCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTTCTTCCCT
GCTGATAGAGCTTTACATACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGCTTTGCGCCATTGTGCAA
TATTCCCCACTGCTGACTCCCGTAGGAGTTTGGA >SEQ ID NO: 35|YK105|
CGTTTCTCCACGCTTCGCGCTCAGCGTCAGTTACTGTCCAGCAATCCGCCTTCGCCACTGGTGTTCTCCTCGTATATC
TACGCATTTACCCGCTACACACGGAATTCGATTGCCTCTCCAGCACTCAAGAACTACAGTTTCAAATGCAGGCTGG
AGGTTGAGCCCCCAGTTTTACATCTGACTTGCAATCCCGCCTACACGCCCTTTACACCCAGTAAATCCGGATAACG

CTTGCCACTACGTATTACCGCGGCTGCTGCCAGTATGACCGTGGCTTATTCTCAGGTACCGTCATTTGTTTC
GTCCCCGACAAAAGAAGTTTACAACCCGAAAGCCTTCTTCTTCACGCGGCGTTGCTGGGTCAGGCTTGCGCCCAT
GCCCAATATTCCCCACTGCTGCCTCCCTGGGAAGTTTGG >SEQ ID NO: 36|YK153|
ATGTCCTGACTTCGCGCCTCAGCGTCAGTTGTCTGCCAGAAAGCCGCTTTCGCCACTGGTGTTCCTCCTAATATCTA
CGCATTTACACGCTACACTAGGAATTCCGCTTTCCTCTCCGACACTCGAGCTTACAGTTTCGGTCCCCTCACGGGG
TTAAGCCCCGCACTTTTAAGACCGACTTGCGATGCCGCTGCGCGCCCTTTACGCCCAATAATTCCGGACAACGCTT
GCCACCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGTGGCTTCTCTTACGGTACCGTCAGGGATAACGGG
TATTGACCGCTATCCTGTTTCGTCCCATAAACAGAACTTTACAACCCGAAGGCCGTCATCGTTCACGCGGCGTTGCT
CCGTCAGACTTTCGTCCATTGCGGAAGATTCCCCACTGCTGCCTCCCTGGGAAGTTTGGA >SEQ ID NO:
37|YK163|
GTTTGCTCACGCTTTCGAGCTCAGCGTCAGTTATCGTCCAGTAAGCCGCTTTCGCCACTGGTGTTCCTCCTAATATC
TACGCATTTACACGCTACACTAGGAATTCCGCTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCG
GGGTTGAGCCCCGGGCTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACG
CTTGCAACCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTATCTTC
CCTGCTGATAGAGCTTTACATAACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGCTTTCGCCCATTTGTG
CAATATTCCCCACTGCTGCCTCCCGTAGGAGTTTGG >SEQ ID NO: 38|YK191|
CGTTGCTCACGCATTCGAGCCTCAGCGTCAGTTAAGCCCAGTAAGCCGCTTTCGCCACTGATGTTCCTCCTAATATC
TACGCATTTACACGCTACACTAGGAATTCCGCTTACCTCTACTTCACTCAAGAACCACAGTTTCAAATGCAGTTTAT
GGGTTAAGCCCATAGTTTTCACATCTGACTTGCGATCCCGCCTACGCTCCCTTTACACCCAGTAAATCCGGACAACG
CTCGCTCCCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGGAGCTTCCCTCCTCAGGTACCGTCTTTTTTCGT
CCCTGAAGACAGAGGTTTACAATCCTAAAACCTTCTTCCCTCACGCGGCATCGCTGCATCAGAGTTTCTCCTCATTGT
GCAATATTCCCCACTGCTGCCTCCCGTAGGAGTTTGGA >SEQ ID NO: 39|YK99|
TGGGCTTACCCATAAACTGCATTTGAACTGTGGTTCTTGAGTGAAGTAGAGGTAAGCGGAATTCCTAGTGTAGCGG
TGAAATGCGTAGATATTAGGAGGAACATCAGTGCGCAAGGCGGCTTACTGGGCTTTAACTGACGCTGAGGCTCGAAA
GCGTGGGGAGCAAACAGGATTAGATACCCAAGTAA >SEQ ID NO: 40|YK55|
GTCAGCATCGAGCTCACGTCAGTTACCGTCCAGTAAGCCGCTTTCGCCACTGGTGTTCCTCCTAATATCTACGCATT
TCACCGCTACACTAGGAATTCCGCTTACCTCTCCGGTACTCTAGATTGACAGTTTCCAATGCAGTCCCGGGGTTGAG
CCCCGGGTTTTACATCAGACTTGCCACTCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCACC
ATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGCTGA
TAGAGCTTTACATAACCGAAATACTTCATCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAATATC
CCCCTGCTGCCTCCCGAGGGAGTTTGGA >SEQ ID NO: 41|YK75|
TCATCGCTTACGGTGGATCTGCGCCGGGTACGGGCGGGCTGGAGTGCGGTAGGGGAGACTGGAATTCCTGGTGTAAC
GGTGGAATGTGTAGATATCGGGAAGAACACCGATGGCGAAGGCAGGTCTCTGGGCCGTCCTGACGCTGAGGAGCGA
AAGCGTGGGGAGCGAACAGGATTAGATACAACGGTAA >SEQ ID NO: 42|YK90|
TGAACCCAGGGCTTAACTCTGGGACTGCTTTTGAAGTGTGAGTGCAGGAGAGGTAAGCGGAATTCCTAGT
GTAGCGGTGAAATGCGTAGATATTAGGAGGAACATCAGTGCGCAAGGCGGCTTACTGGACTGAAACTGACACTGAGG
CACGAAAGCGTGGGGAGCAAACAGGATTAGATACCATGGTAA >SEQ ID NO: 43|YK30|
ACCAGGGCTTAACTCTGGGACTGCTTTTGAAGTGTGAGTGCAGGAGAGGTAAGCGGAATTCCTAGTGTAG
CGGTGAAATGCGTAGATATTAGGAGGAACATCAGTGCGCAAGGCGGCTTACTGGACTGAAACTGACACTGAGGCACG
AAAGCGTGGGGAGCAAACAGGATTAGATACCTGGTAA >SEQ ID NO: 44|YK31|
GAACCCAGGGCTTAACTCTGGGACTGCTTTTGAAGTGTGAGTGCAGGAGAGGTAAGCGGAATTCCTAGTG
TAGCGGTGAAATGCGTAGATATTAGGAGGAACATCAGTGCGCAAGGCGGCTTACTGGACTGAAACTGACACTGAGGC
ACGAAAGCGTGGGGAGCAAACAGGATTAGATACCCCGGTAA >SEQ ID NO: 45|YK12|
GAGTCAGCTTTCGAGCTCAGCGTCAGTTATCGTCCAGTAAGCCGCTTTCGCCACTGGTGTTCCTCCTAATATCTACG
CATTTACCGCTACACTAGGAATTCCGCTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGT
TGAGCCCCGGGCTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTG
CACCATACTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTG
CTGATAGAGCTTTACATAACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGCTTTCGCCCATTTGTGCAAT
ATTCCCCACTGCTGCCTCCCGAGGGAGTTTGGA >SEQ ID NO: 46|YK27|
TGTCAGCTTTCGAGCTCACGTCAGTTACCGTCCAGTAAGCCGCTTTCGCCACTGGTGTTCCTCCTAATATCTACGCA
TTTCACCGCTACACTAGGAATTCCGCTTACCTCTCCGGTACTCTAGATTGACAGTTTCCAATGCAGTCCCGGGGTTG
AGCCCCGGGTTTTACATCAGACTTGCCACTCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCA
CCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGCT
GATAGAGCTTTACATAACCGAAATACTTCATCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAATAT
TCCCCACTGCTGCCTCCCGTAGGAGTTTGGA >SEQ ID NO: 47|YK28|
CACGTCAGTTACCGTCCAGTAAGCCGCTTTCGCCACTGGTGTTCCTCCTAATATCTACGCATTTACACCGCTACACTA
GGAATTCGCTTACCTCTCCGGCACTCAAGACGGGCAGTTTCCAATGCAGTCCCGGGGTTGAGCCCCAGCCTTTAC
ATCAGACTTGTCCATCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCCCCCTACGTATTACCGC
GGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGCTGATAGAAGTTTACATA
CCGAGATACTTCTTCTTACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAATATTCCCCACTGCTGCCTC
CCGTAGGAGTTTGGG >SEQ ID NO: 48|YK29|
GTCAGCTTTCGAGCTCACGTCAGTTACCGTCCAGTAAGCCGCTTTCGCCACTGGTGTTCCTCCTAATATCTACGCAT
TTCACCGCTACACTAGGAATTCCGCTTACCTCTCCGGTACTCTAGATTGACAGTTTCCAATGCAGTCCCGGGGTTGA

GCCCGCGGTTTTTACATACAGACTTGCCACTCCGCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCA
 CATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGCTG
 ATAGAGCTTTACATACCGAAATACTTCATCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAATATT
 CCCCCTGCTGCCTCCCGTGGGGAGTTTGGA>SEQ ID NO: 49|YK33|
 GATGCTCAGCTTTTCGTGCTCAGTGTACGTTTCAGTCCAGTAAGCCGCCTTCGCCACTGATGTTTCTCCTAATATCTA
 CGCATTTACCGCTACACTAGGAATTCCGCTTACCTCTCCTGCACTCCAGTCTGACAGTTTCAAAGCAGTCCCAGA
 GTTAAGCCCTGGGTTTTCACTTCTGACTTGCCATACCACCTACGCACCCTTTACACCCAGTAATTCCGGATAACGCT
 TGCCCCCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTTTCTTCCC
 TGCTGATAGAGCTTTACATACCGAGATACTTCTTCACTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCA
 ATATTCCCCACTGCTGCCTCCCGAAGGAAGTTTGGA>SEQ ID NO: 50|YK34|70A_009_YK34_A1_A02
 GTGTCAGCTTCGTGCTCAGTGTACGTTTCAGTCCAGTAAGCCGCCTTCGCCACTGATGTTTCTCCTAATATCTACGC
 ATTTACCGCTACACTAGGAATTCCGCTTACCTCTCCTGCACTCCAGTCTGACAGTTTCAAAGCAGTCCCAGAGTT
 AAGCCCTGGGTTTTCACTTCTGACTTGCCATACCACCTACGCACCCTTTACACCCAGTAATTCCGGATAACGCTTGC
 CCCCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGC
 TGATAGAGCTTTACATACCGAGATACTTCTTCACTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAATA
 TTCCCCACTGCTGCCTCCCGTAGGGAGTTTGGA>SEQ ID NO: 51|YK35|
 GTCAGCTTCGAGCCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTTCTCCTAATATCTACGCAT
 TTCACCGCTACACTAGGAATTCCGCTTACCTCTCCGGTACTCTAGATTGACAGTTTCCAATGCAGTCCCAGGGGTTGA
 GCCCCGGGTTTTACATCAGACTTGCCACTCCGCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCAC
 CATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGCTG
 ATAGAGCTTTACATACCGAAATACTTCATCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAATATT
 CCCCCTGCTGCCTCCCGTAGGGAGTTTGGA>SEQ ID NO: 52|YK51|
 TGTCAGCTTTTCGAGCTCAGCGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTTCTCCTAATATCTACGC
 ATTTACCGCTACACTAGGAATTCCGCTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGTT
 GAGCCCCGGGCTTTACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGC
 ACCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTGC
 TGATAGAGCTTTACATACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGCTTTCGCCCATTTGTGCAATA
 TTCCCCACTGCTGCCTCCCGAAGGGAGTTTGGA>SEQ ID NO: 53|YK52|
 TTCAGCTTTTCGAGCTCAGCGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTTCTCCTAATATCTACGCA
 TTTACCGCTACACTAGGAATTCCGCTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGTTG
 AGCCCCGGGCTTTACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCAC
 CCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTGCT
 GATAGAGCTTTACATACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGCTTTCGCCCATTTGTGCAATAT
 TCCCCACTGCTGCCTCCCGAGGGGAGTTTG>SEQ ID NO: 54|YK54|
 TTCGGTCTGCTTTCCCTTCTCGCGCCTCAGTGTACGTTTCTGTCTAGTAAGCCGCCTTCGCCACTGATGTTTCTCTC
 TAATATCTACGCACTTCACCGCTCCACAATGAATTCCGCTTACCCCTCCCGCGCTCTAGTCTGACAGTTTAAAAAA
 ACTCCCCGAGAGAAACCCTGGGTTTTTTCTTCTGACATGCGATATCCCACCCCCACCCTTTATACACCCAAAAATCG
 GATAAAAGGTGCGACCTACGTATTATACCGGCTGCTGGGGCGTAGATAGCCGGGGGTTCTTATACAGGGACCGTCAT
 TTTCTTTCCCGCTGATACAGCTTTACATACCGAAATACTTCTTTCTCACGCGGCGTCGCTGCATCAGGGTTTCCCC
 ATTGTGCAATATTCCCCACTGCTGCCTCCCGAAGGGGAAGTTGGGGGAA>SEQ ID NO: 55|YK56|
 GTTCAGCTTTTCGAGCCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTTCTCCTAATATCTACGC
 ATTTACCGCTACACTAGGAATTCCGCTTACCTCTCCGGTACTCTAGATTGACAGTTTCCAATGCAGTCCCAGGGGTT
 GAGCCCCGGGTTTTACATCAGACTTGCCACTCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGC
 ACCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGC
 TGATAGAGCTTTACATACCGAAATACTTCATCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAATA
 TTCCCCACTGCTGCCTCCCGAGGGGAGTTTGGA>SEQ ID NO: 56|YK57|
 GTCAGCTTTTCGAGCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTTCTCCTAATATCTACGCAT
 TTCACCGCTACACTAGGAATTCCGCTTACCTCTCCGGTACTCTAGATTGACAGTTTCCAATGCAGTCCCAGGGGTTGA
 GCCCCGGGTTTTACATCAGACTTGCCACTCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCAC
 CATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGCTG
 ATAGAGCTTTACATACCGAAATACTTCATCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAATATT
 CCCCCTGCTGCCTCCCGAGGGGAGTTTGGA>SEQ ID NO: 57|YK58|
 TCTCACGCTTTTCGAGCTCACGTCAGTCATCGTCCAGCAAGCCGCCTTCGCCACTGGTGTTTCTCCTAATATCTACGC
 ATTTACCGCTACACTAGGAATTCCACTTGCCTCTCCGACACTCTAGCTCAGCAGTTCCAAATGCAGTCCCAGGGGTT
 GAGCCCCGGGCTTTACATCTGGCTTGCCGTGCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGC
 CCCCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGC
 TGATAGAAGTTTACATACCGAAATACTTCATCCTTCACGCGGCGTCGCTGCATCAGAGTTTCTCCTCATTGTGCAATA
 TTCCCCACTGCTGCCTCCCGTAGGGAGTTTG>SEQ ID NO: 58|YK65|
 GTCAGCTTTTCGAGCTCAGCGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTTCTCCTAATATCTACGCA
 TTTACCGCTACACTAGGAATTCCACTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGTTG
 AGCCCCGGGCTTTACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCAC
 CCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTCTTCTTCCCTGC
 TGATAGAGCTTTACATACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAATA

TTCCCTCCGCTGCTGCTCCCGAGGAGTTTGGGA>SEQ ID NO: 59|YK67|
AGCCCCGCTTTTCGAGCCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCTCCTAATATCTACG
CATTTACCGCTACACTAGGAATTCCGCTTACCTCTCCGGCACTCAAGACGGGCAGTTTCCAATGCAGTCCCGGGGT
TGAGCCCCAGCCTTTACATCAGACTTGTCCATCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTG
CCCCCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTG
CTGATAGAAGTTTACATACCGAGATACTTCTTCCCTTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAAT
ATTCCCCACTGCTGCCTCCCGAAGGAAGTTTGGGA>SEQ ID NO: 60|YK69|
TGCTCAGCTTTTCGAGCCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCTCCTAATATCTACG
CATTTACCGCTACACTAGGAATTCCGCTTACCTCTCCGGCACTCGAGCCAGACAGTTTCCAATGCAGTCCCAGGGT
TAAGCCCTGGGTTTTTCACATCAGACTTGCCTTGCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTG
CCCCCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTG
CTGATAGAGCTTTACATACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAAT
ATTCCCCACTGCTGCCTCCCGAAGGGAGTTTGGGA>SEQ ID NO: 61|YK70|
GTTGCTCAGCTTTTCGAGCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCTCCTAATATCTAC
GCATTTACCGCTACACTAGGAATTCCGCTTACCTCTCCGGCACTCGAGCCAGACAGTTTCCAATGCAGTCCCAGGG
TTAAGCCCTGGGTTTTTCACATCAGACTTGCCTTGCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTT
GCCCCCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTATCTTCCCT
GCTGATAGAGCTTTACATACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAA
TATTCCCCACTGCTGCCTCCCGAAGGAAGTTTGGGA>SEQ ID NO: 62|YK71|
TGCTCAGCTTTTCGAGCTCAGCGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCTCCTAATATCTACG
CATTTACCGCTACACTAGGAATTCCACTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGT
TGAGCCCCGGGCTTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTG
CACCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTCTTCTTCCCT
GCTGATAGAGCTTTACATACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAA
TATTCCCCACTGCTGCCTCCCGAAGGGAGTTTGGGA>SEQ ID NO: 63|YK74|
GATGCCCTGGCTTCGCGCTCAGCGTCAGTTGTCTGCCAGAAAGCCGCTTTTCGCCACTGGTGTTCCTCCTAATATCTA
CGCATTTACCGCTACACTAGGAATTCCGCTTTCTCTCCGACACTCGAGCTTCACAGTTTCGGTCCCCTCACGGGG
TTAAGCCCCGCACTTTTTAAGACCGACTTGCGATGCCGCTTGCAGCGCCCTTTACGCCCAATAATTCCGGACAACGCTT
GCCACCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGTGGCTTCTCTTACGGTACCGTCAGGGATAACGGG
TATTGACCGCTATCCTGTTTCGTCCCATATAACAGAACTTTACAACCCGAAGGCCGTCATCGTTCACGCGGCGTTGCT
CCGTCAGACTTTCGTCCATTGCGGAAGATTCCCCACTGCTGCCTCCCGGGGGGAGTTTGGGA>SEQ ID NO:
64|YK88|
GTCCCGCTTTTCGAGCCTCAGCGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCTCCTAATATCTACG
CATTTACCGCTACACTAGGAATTCCGCTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGT
TGAGCCCCGGGCTTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTG
CACCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTG
CTGATAGAGCTTTACATACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGCTTTTCGCCATTGTGCAAT
ATTCCCCACTGCTGCCTCCCGAAGGGAAGTTTGG>SEQ ID NO: 65|YK89|
TGTCAGCTTTTCGAGCTCAGCGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCTCCTAATATCTACGC
ATTTACCGCTACACTAGGAATTCCGCTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGT
GAGCCCCGGGCTTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTG
ACCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTGC
TGATAGAGCTTTACATACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGCTTTTCGCCATTGTGCAATA
TTCCCCACTGCTGCCTCCCGAAGGGAGTTTGGGA>SEQ ID NO: 66|YK97|
TGCTCAGCTTTTCGAGCTCAGCGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCTCCTAATATCTACG
CATTTACCGCTACACTAGGAATTCCACTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGT
TGAGCCCCGGGCTTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTG
CACCATACGTATTACCGCGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTCTTCTTCCCT
GCTGATAGAGCTTTACATACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAA
TATTCCCCACTGCTGCCTCCCGAAGGGAGTTTGGGA>SEQ ID NO: 67|YK98|
ATTAGCTTTTCGAGCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCTCCTAATATCTACGCA
TTTCACCGCTACACTAGGAATTCCGCTTACCCCTCCGGCACTCAAGCATAACAGTTTCCAATGCAGTCCAGGGGT
AGCCCTGCCTTTTCACATCAGACTTGATACGCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTCGCC
CCCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTGCT
GATAGAAGTTTACATACCGAGATACTTCTTCCCTTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAAT
TCCCCACTGCTGCCTCCCGAGGGAAGTTTGGGA>SEQ ID NO: 68|YK139|
GTGTCAGCTTTTCGAGCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCTCCTAATATCTACGC
ATTTACCGCTACACTAGGAATTCCGCTTACCTCTCCGGCACTCGAGCCAGACAGTTTCCAATGCAGTCCCAGGGT
AAGCCCTGGGTTTTTCACATCAGACTTGCCTTGCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTG
CCCCTACGTATTACCGCGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTGC
TGATAGAGCTTTACATACCGAAATACTTCTTCGCTCACGCGGCGTCGCTGCATCAGGGTTTCCCCCATTGTGCAATA
TTCCCCACTGCTGCCTCCCGAGGGAGTTTGG>SEQ ID NO: 69|YK141|
GCCAGCTTCGAGCCTCACGTCAGTCATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCTCCTAATATCTACGCAT

TTACACCTACCAATGGAATTTCCACTTCACTTACCTTCCGACACTCTAGCTGACAGTCTTCCAAAGAGGATCCACAGTCACGATGCCCCATGCCTTTTCACTTCAGACTTGCCACAGCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCCCCCTACGTATTACCGCGGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGCTGATAGAAGTTTACATAACCGAAATACTTTCATCCTTCACGCGGGCTCGCTGCATCAGGCTTTCGCCCCATTGTGCAATATCCCCACTGCTGCCTCCCGAGGGGAAGTTTGGA>SEQ ID NO: 70|YK142|

TGATCAGCTTTCGAGCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCCTCCTAATATCTACGCATTTACACCGCTACACTAGGAATTCCGCTTACCTCTCCGGTACTCTAGATTGACAGTTTCCAATGCAGTCCCGGGGTTGAGCCCCGGGTTTTACATCAGACTTGCCACTCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCAACATACGTATTACCGCGGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGCTGATAGAGCTTTACATAACCGAAATACTTTCATCGCTCACGCGGGCTCGCTGCATCAGGCTTTCGCCCCATTGTGCAATAATCCCCACTGCTGCCTCCCGGGGGAGTTTGGA>SEQ ID NO: 71|YK152|

GATGATCAGCTTTCGAGCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCCTCCTAATATCTACGCATTTACACCGCTACACTAGGAATTCCGCTTACCTCTCCGGCACTCTAGAAAAACAGTTTCCAATGCAGTCCCTGGGGTTAAGCCCCAGCCTTTCACATCAGACTTGCTCTTCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCCCCCTACGTATTACCGCGGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGCTGATAGAAGTTTACATAACCGAGATACTTCTTCCCTTCACGCGGGCTCGCTGCATCAGGCTTTCGCCCCATTGTGCAATATTTCCCCACTGCTGCCTCCCGGGGGAGTTTGGA>SEQ ID NO: 72|YK155|

TTGATCAGCTTTCGAGCTCACGTCAGTTACCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCCTCCTAATATCTACGCATTTACACCGCTACACTAGGAATTCCGCTTACCTCTCCGGCACTCTAGAAAAACAGTTTCCAATGCAGTCCCTGGGGTTAAGCCCCAGCCTTTCACATCAGACTTGCTCTTCCGTCTACGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCCCCCTACGTATTACCGCGGGCTGCTGGCACGTAGTTAGCCGGGGCTTCTTAGTCAGGTACCGTCATTTTCTTCCCTGCTGATAGAAGTTTACATAACCGAGATACTTCTTCCCTTCACGCGGGCTCGCTGCATCAGGCTTTCGCCCCATTGTGCAATATTCCCCACTGCTGCCTCCCGAAGGGAGTTTGGA>SEQ ID NO: 73|YK157|

GTATTTACAGCTTTCGAGCTCACGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCCTCCTAATATCTACCGCATTTACACCGCTACACTAGGAATTCCGCTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGTTGAGCCCCGGGCTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGACACCATAACGTATTACCGCGGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTGCTGATAGAGCTTTACATAACCGAAATACTTCTTCGCTCACGCGGGCTCGCTGCATCAGGCTTTCGCCCCATTGTGCAATATATTCCCCACTGCTGCCTCCCGAAGGGAGTTTGGA>SEQ ID NO: 74|YK160|

GCTCAGCTTTCGAGCTCACGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCCTCCTAATATCTACGCATTTACACCGCTACACTAGGAATTCCACTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGTTGAGCCCCGGGCTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCAACATACGTATTACCGCGGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTCTTCTTCCCTGCTGATAGAGCTTTACATAACCGAAATACTTCTTCGCTCACGCGGGCTCGCTGCATCAGGCTTTCGCCCCATTGTGCAATATTCCCCACTGCTGCCTCCCGAGGGGAGTTTGGA>SEQ ID NO: 75|YK166|

TTTCAGCTTCGAGCCTCAGCGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCCTCCTAATATCTACGCATTTACACCGCTACACTAGGAATTCCGCTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGTTGAGCCCCGGGCTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCAACATACGTATTACCGCGGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTGCTGATAGAGCTTTACATAACCGAAATACTTCTTCGCTCACGCGGGCTCGCTGCATCAGGCTTTCGCCCCATTGTGCAATAATCCCCACTGCTAGCTCCCGAAGGAGTTTGGA>SEQ ID NO: 76|YK168|

AGCTCAGCTTTCGAGCTCACGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCCTCCTAATATCTACGCATTTACACCGCTACACTAGGAATTCCGCTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGTTGAGCCCCGGGCTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCAACATACGTATTACCGCGGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTGCTGATAGAGCTTTACATAACCGAAATACTTCTTCGCTCACGCGGGCTCGCTGCATCAGGCTTTCGCCCCATTGTGCAATATTCCCCACTGCTGCCTCCCGAAGGGAGTTTGGA>SEQ ID NO: 77|YK169|

GTCCAGCTTTCGAGCCTCAGCGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCCTCCTAATATCTACGCATTTACACCGCTACACTAGGAATTCCGCTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGTTGAGCCCCGGGCTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCAACATACGTATTACCGCGGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTATCTTCCCTGCTGATAGAGCTTTACATAACCGAAATACTTCTTCGCTCACGCGGGCTCGCTGCATCAGGCTTTCGCCCCATTGTGCAATATTCCCCACTGCTGCCTCCCGAGGGAGTTTGGA>SEQ ID NO: 78|YK171|

TGAGCCGGGCTCACCCCGGTACTGCATTGGAAGTGTCTGACTAGAGTGTGCGAGGGGTAAGCGGAATTCCTAGTGTAACGGTGAAATGCGTAGATATTAGGAGGAACACCAAGTGGCGAAGGCGGCTTACTGGACGATAACTGACGCTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACACCGGTAA>SEQ ID NO: 79|YK192|

CACGATGTCAGCTTTCGAGCTCAGCGTCAGTTATCGTCCAGTAAGCCGCCTTCGCCACTGGTGTTCCCTCCTAATATCTACGCATTTACACCGCTACACTAGGAATTCCACTTACCCCTCCGACACTCTAGTACGACAGTTTCCAATGCAGTACCGGGGGTTGAGCCCCGGGCTTTCACATCAGACTTGCCGCACCGCCTGCGCTCCCTTTACACCCAGTAAATCCGGATAACGCTTGCAACATACGTATTACCGCGGGCTGCTGGCACGTATTTAGCCGGTGCTTCTTAGTCAGGTACCGTCATTCTTCTTCCCTGCTGATAGAGCTTTACATAACCGAAATACTTCTTCGCTCACGCGGGCTCGCTGCATCAGGCTTTCGCCCCATTGTGCAATATTCCCCACTGCTGCCTCCCGAGGGGAGTTTGGA>SEQ ID NO: 80|VE202-18|

ATGGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGGCTGCCTAATACATGCAAGTCGAACGCGAGCACTTGTG

CTCGAGTGGCGAAGCAACGGGTGAGTAATACATAAAGTAACCTGCCCCTAGACACAGGGGGATAAAGTATTTGGAAACCGATAGCTAA
GACCGCATAGGTACGGACACTGCATGGTGACCGTATTAATAAAGTGCCTCAAAGCACTGGTAGAGGATGGACTTATGG
CGCATTAGCTGGTTGGCGGGGTAACGGCCCAACAGGCGACGATGCGTAGCCGACCTGAGAGGGTGACCGGCCACAC
TGGGACTGAGACACGGCCAGACTCCTACGGGAGGCAGCAGTAGGGAATTTTCGGCAATGGGGGAAACCCTGACCGA
GCAACGCCGCGTGAAGGAAGAAGGTTTTTCGGATTGTAAACTTCTGTTATAAAGGAAGAACGGCGGCTACAGGAAAT
GGTAGCCGAGTGACGGTACTTTATTAGAAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGTGGCA
AGCGTTATCCGGAATTATTGGGCGTAAAGAGGGAGCAGGCGGCAGCAAGGGTCTGTGGTGAAAGCCTGAAGCTTAAC
TTCAGTAAGCCATAGAAACCAGGCAGCTAGAGTGCAGGAGAGGATCGTGGAATTCCATGTGTAGCGGTGAAATGCG
TAGATATATGGAGGAACACCAGTGGCGAAGGCGACGATCTGGCCTGCAACTGACGCTCAGTCCCGAAAGCGTGCGGA
GCAAATAGGATTAGATACCCTAGTAGTCCACGCCGTAAACGATGAGTACTAAGTGTTGGATGTCAAAGTTCAGTGCT
GCAGTTAACGCAATAAGTACTCCGCCTGAGTAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGGCCCG
CACAAGCGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAGGTCTTGACATACTCATAAAGGCT
CCAGAGATGGAGAGATAGCTATATGAGATACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTT
AAGTCCCGCAACGAGCGCAACCCTTATCGTTAGTTACCATCATTAAAGTTGGGGACTCTAGCGAGACTGCCAGTGAC
AAGCTGGAGGAAGGCGGGGATGACGTCAAATCATCATGCCCTTATGACCTGGGCTACACACGTGCTACAATGGATG
GTGCAGAGGGAAGCGAAGCCGCGAGGTGAAGCAAAACCCATAAAACCATTTCTCAGTTTCGGATTGTAGTCTGCAACTC
GACTACATGAAGTTGGAATCGCTAGTAATCGCGAATCAGCATGTGCGCGGTGAATACGTTCTCGGGCCTTGTAACA
CCGCCCCGTACACCACGAGAGTTGATAACACCCGAAGCCGGTGGCCTAACCGCAAGGAAGGAGCTGTCTAAGGTGGG
ATTGATGATTGGGGTGAAGTCGTAACAAGGTATCCCTACGGGAACGTGGGGATGGATCACCTCCTTT >SEQ

ID NO: 81|PE5|

ATGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCAATTGAAGG
AAGTTTTTCGGATGGAATTTCGATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGAATAACCTGCCTCACACTGGGGG
ATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTACCGCATGGTACAGTGTGAAAACTCCGGTGGT
GTGAGATGGATCCGCGTCTGATTAGCCAGTTGGCGGGGTAACGGCCCAACAAAGCGACGATCAGTAGCCGACCTGAG
AGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATG
GGCGAAAGCCTGATGCAGCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGAAGAA
AATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTAT
CCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCGAAGCAAGTCTGAAGTGAAAACCCAGGGCTCAACCCTGGGAC
TGCTTTGGAAACTGTTTTGCTAGAGTGTGCGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATGCTAGGTGTTGGGGGGCAAAGCCCTTCGGTGCCGTCG
CAAACGCAGTAAGCATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACA
AGCGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCTCTTGACCGGCGTGT
AACGGCGCCTTCCCTTCGGGGCAAGAGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTT
AAGTCCCGCAACGAGCGCAACCCTTATCCTTAGTAGCCAGCAGGTAAAGCTGGGCACTCTAGGGAGACTGCCAGGG
ATAACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGC
GTAAACAAAGGGAAGCAAGACAGTGATGTGGAGCAAATCCCAAAAATAACGTCCCAGTTTCGGACTGTAGTCTGCAAC
CCGACTACACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTAACA
CACCGCCCCGTACACCATGGGAGTCAGCAACGCCCGAAGTCAGTGACCCAACTCGCAAGAGAGGGAGCTGCCGAAGC
CGGGGCAGGTAACCTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT

>SEQ ID NO: 82|PE9|

AATTCGACGTTGTCCGGATTACTGGGCGTAAAGGGAGCGTAGGCGGACTTTTAAGTGAGATGTGAAATACCCGGGCT
CAACTTGGGTGCTGCATTTCAAACCTGGAAGTCTAGAGTGCAGGAGAGGAGAATGGAATTCCTAGTGTAGCGGTGAAA
TGCGTAGAGATTAGGAAGAACACCAGTGGCGAAGGCGATTCTCTGGACTGTAAGTACGCTGAGGCTCGAAAGCGTG
GGGAGCAAACAGGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTAGGGGTTGTATGACCT
CTGTGCCGCCGCTAACGCATTAAGTATTCGCCTGGGGAGTACGGTTCGCAAGATTAAAACTCAAAGAAATTGACGGA

>SEQ ID NO: 83|211-B|

ACGAGCGTATCGGATTATTGGGTTTAAGGGAGCGTAGGTGGATTGTAAAGTCAGTTGTGAAAGTTTGCGGCTCAACC
GTAAATTTGCAGTTGAAACTGGCAGTCTTGAGTACAGTAGAGGTGGGCGGAATTCGTGGTGATGCGGTGAAATGCTT
AGATATCACGAAGAACTCCGATTGCGAAGGCAGCTCACTAGACTGTCACTGACACTGATGCTCGAAAGTGTGGGTAT
CAAACAGGATTAGATACCCTGGTAGTCCACACAGTAAACGATGAATACTCGCTGTTTTCGATATACAGTAAGCGGCC
AAGCGAAAGCATTAAAGTATTCACCTGGGGAGTACGCCGGCAACGGTGAAACTCAAAGAAATTGACGGAAGCCCGCC
CAGGGGGGAAAAACATGGGGTTTATGTTGGATGATACGGGGAGGAACCTC >SEQ ID NO:

84|NR_119185.1|Ruminococcus obeum 16S ribosomal RNA gene, complete sequence

GGCGGCGTGCTTAACACATGCAAGTCGAACGGGAAACCTTTTCATTGAAGCTTCGGCAGATTGNNNTGTTTCTAGTG
GCGGACGGGTGAGTAACGCGTGGGTAACTGCCTTATACAGGGGGGATAACAACCAGAAATGGTTGCTAATACCGCAT
AAGCGCACAGGACCGCATGGTCTGGTGTGAAAACTCCGGTGGTATAAGATGGACCCGCGTTGGATTAGCTAGTTGG
CAGGGTAACGGCCTACCAAGGCGACGATCCATAGCCGGCCTGAGAGGGTGAACGGCCACATTGGGACTGAGACACGG
CCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGGGAAACCCTGATGCAGCGACGCCGCGTGAAG
GAAGAAGTATCTCGGTATGTAAACTTCTATCAGCAGGGAAGATAGTGACGGTACCTGACTAAGAAGCCCCGGCTAAC
TACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGG
ACTGGCAAGTCTGATGTGAAAGGCGGGGGCTCAACCCCTGGACTGCATTGGAAACTGTAGTCTTGAGTGCCGGAGA
GGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAGTGGCGAAGGCGGCTTACTGG

GATTAGATACCCTGGTAGTCCACGCCGTA AACGATGACTACTAGGTGTCTGGGTGGCAAGGCCATTTCGGTGCCGCAGC
AAACGCAATAAGTAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCTGATCTTGACATCCCGATGCCAAAGCGCGTA
ACGCGCTCTTTCTTCGGAACATCGGTGACAGGTGGTGCATGGTTGTCTGCAGCTCGTGTCTGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCCTATCTTCAGTAGCCAGCATTTTGGATGGGCACTCTGGAGAGACTGCCAGGGAGAA
CCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGACCAGGGCTACACACGTGCTACAATGGCGTAA
ACAAAGGGAGGCGAACCCGCGAGGGTGGGCAAATCCCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCGA
CTACATGAAGTTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCG
CCCGTCACACCATGGGAGTCAGTAACGCCCGAAGCCGGTGACCCAACCCGTAAGGGAGGGAGCCGTCGAAGGTGGGA
CCGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTC >SEQ

>SEQ ID NO: 89|NR_042152.1|*Marvinbryantia formatexigens* strain I-52 16S ribosomal RNA gene, partial sequence >gi|636558750|ref|NR_114807.1| *Marvinbryantia formatexigens* strain I-52 16S ribosomal RNA gene, complete sequence

>SEQ ID NO: 90|NR_024994.1|*Lactobacillus mucosae* strain S32 16S ribosomal RNA gene, complete sequence

AGAGTTTTCATCCTGGCTCAGGATGAACGCCGGCGGTGTGCCTAATACATGCAAGTCGAACGCGTTGGCCCAACTGAT
TGAACGTGCTTGCACGGACTTGACGTTGGTTTACCAGCGAGTGGCGGACGGGTGAGTAACACGTAGGTAACCTGCCC
CAAAGCGGGGGATAACATTTGGAACAGATGCTAATACCGCATAACAATTTGAATCGCATGATTCAAATTTAAAAGA
TGGCTTCGGCTATCACTTTGGGATGGACCTGCGGCGCATTAGCTTGTGGTAGGGTAACGGCCTACCAAGGCTGTGA
TGCGTAGCCGAGTTGAGAGACTGATCGGCCACAATGGAAGTGAAGACACGGTCCATACTCCTACGGGAGGCAGCAGTA
GGGAATCTTCCACAATGGGCGCAAGCCTGATGGAGCAACACCGCGTGAGTGAAGAAGGGTTTCGGCTCGTAAAGCTC
TGTTGTTAGAGAAGAACGTGCGTGAGAGCAACTGTTACGCAGTGACGGTATCTAACCAGAAAGTCACGGCTAACTA
CGTGCCAGCAGCCGCGGTAATACGTAGGTGGCAAGCGTTATCCGGATTTATTGGGCGTAAAGCGAGCGCAGGCGGTT
TGATAAGTCTGATGTGAAAGCCTTTGGCTTAACCAAAGAAGTGCATCGGAAACTGTCAGACTTGAGTGCAGAAGAGG

ACAGTGAAGTAAGTGTGATGTGATGTGGAAGTGTGGAAGAACACCATGTCGCGAAGGCGGCTGTCTGGTC
TGCAACTGACGCTGAGGCTCGAAAGCATGGGTAGCGAACAGGATTAGATACCCTGGTAGTCCATGCCGTAAACGATG
AGTGCTAGGTGTTGGAGGGTTTCCGCCCTTCAGTGCCGCAGCTAACGCATTAAGCACTCCGCTGGGGAGTACGACC
GCAAGGTTGAAACTCAAAGGAATTGACGGGGGCCCCGCACAAGCGGTGGAGCATGTGGTTTAATTCTGAAGCTACGCGA
AGAACCTTACCAGGTCTTGACATCTTGCGCCAACCCTAGAGATAGGGCGTTTTCCTTCGGGAACGCAATGACAGGTGG
TGCATGGTCGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCAACCCTTGTTACTAGTTGC
CAGCATTGAGTTGGGCACTCTAGTGAGACTGCCGGTGACAAACCGGAGGAAGGTGGGGACGACGTCAGATCATCATG
CCCCTTATGACCTGGGCTACACACGTGCTACAATGGACGGGTACAACGAGTCGCGAACTCGCGAGGGCAAGCTAATCT
CTTAAAACCGTTCTCAGTTCGGACTGCAGGCTGCAACTCGCCTGCACGAAGTCGGAATCGCTAGTAATCGCGGATCA
GCATGCCGCGGTGAATACGTTCCCGGGCCTTGTACACACCGCCCGTCACACCATGAGAGTTTGCAACACCCAAAGTC
GGTGGGGTAACCTTCGGGGAGCTAGCCGCCTAAGGTGGGGCAGATGATTAGGGTGAAGTCGTAACAAGGTAGCCGT
AGGAGAACCTGCGGCTGGATCACCTCCT>SEQ ID NO: 91|NR_028816.1|*Turicibacter sanguinis* strain

MOL361 16S ribosomal RNA gene, complete sequence

AGAGTTTGATCATGGCTCAGGATGAACGCTGGCGGCGTGCCTAATACATGCAAGTCGAGCGAACCCTTCGGTGGTG
AGCGGCGAACGGGTGAGTAACACGTAGGTTATCTGCCCATCAGACGGGGACAACGATTGGAACGATCGCTAATACC
GGATAGGACGAAAGTTTAAAGGTGCTTCGGCACCCTGATGGATGAGCCTGCGGCGCATTAGCTAGTTGGTAGGGTA
AAGGCCTACCAAGGCGACGATGCGTAGCCGACCTGAGAGGGTGAACGGCCACACTGGGACTGAGACACGGCCCAGAC
TCCTACGGGAGGCAGCAGTAGGGAATCTTCGGCAATGGGCGAAAGCCTGACCGAGCAACGCCGCGTGAATGATGAAG
GCCTTCGGGTTGTAAAATTCTGTTATAAGGGAAGAATGGCTCTAGTAGGAAATGGCTAGAGTGTGACGGTACCTTAT
GAGAAAGCCACGGCTAACTACGTGCCAGCAGCCGCGTAATACGTAGGTGGCGAGCGTTATCCGGAATTATTGGGCG
TAAAGAGCGCGCAGGTGGTTGATTAAAGTCTGATGTGAAAGCCCACGGCTTAACCGTGGAGGGTCATTGGAACTGGT
CAACTTGAGTGCAGAAGAGGGAAGTGAATTCATGTGTAGCGGTGAAATGCGTAGAGATATGGAGGAACACCAGTG
GCGAAGGCGGCTTCCTGGTCTGTAAGTGTGACTGAGGCGCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGT
AGTCCACGCCGTAAACGATGAGTGCTAAGTGTGGGGGTCGAACCTCAGTGCTGAAGTTAACGCATTAAGCACTCCG
CCTGGGGAGTACGGTCGCAAGACTGAACTCAAAGGAATTGACGGGGACCCGCACAAGCGGTGGAGCATGTGGTTTA
ATTCGAAGCAACGCGAAGAACCCTTACCAGGTCTTGACATACCAGTGACCGTCCTAGAGATAGGATTTTCCCTTCGGG
GACAATGGATACAGGTGGTGCATGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCA
ACCCCTGTCGTTAGTTGCCAGCATTGAGTTGGGGACTCTAACGAGACTGCCAGTGACAACTGGAGGAAGGTGGGGA
TGACGTCAAATCATCATGCCCCCTTATGACCTGGGCTACACACGTGCTACAATGGTTGGTACAAAGAGAAGCGAAGCG
GTGACGTGGAGCAAACCTCATAAAGCCAATCTCAGTTCGGATTGTAGGCTGCAACTCGCCTACATGAAGTTGGAATC
GCTAGTAATCGCGAATCAGCATGTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCGCCCGTCACACCACGAGAG
TTTACAACACCCGAAGTCAGTGGCCTAACCGCAAGGAGGGAGCTGCCTAAGGTGGGGTAGATGATTGGGGTGAAGTC
GTAACAAGGTATCCCTACCGGAAGGTGGGGTTGGATCACCTCCTT>SEQ ID NO:

92|NR_042832.1|*Roseburia faecis* strain M72/1 16S ribosomal RNA gene, partial sequence

GATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAACGAAGCACTCTATTTGATTTTCTTCGGAAATGAAGATT
TTGTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGGTAACTGCCTCATAACAGGGGGATAACAGTTGGAAACGACT
GCTAATACCGCATAAGCGCACAGGATCGCATGATCCGGTGTGAAAACTCCGGTGGTATGAGATGGACCCGCGTCTG
ATTAGCCAGTTGGCAGGGTAACGGCCTACCAAAGCGACGATCAGTAGCCGACCTGAGAGGGTGACCGGCCACATTGG
GACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGGGAACCCCTGATGCAGCG
ACGCCGCGTGAGCGAAGAAGTATTTCCGTATGTAAAGCTCTATCAGCAGGGAAGAAGAATGACGGTACCTGACTAAG
AAGCACCGGCTAAATACGTGCCAGCAGCCGCGGTAATACGTATGGTGCAAGCGTTATCCGGATTTACTGGGTGTAAA
GGGAGCGCAGGCGGTGCGGCAAGTCTGATGTGAAAGCCCGGGGCTCAACCCCGGTACTGCATTGGAACTGTGCTAC
TAGAGTGTCCGAGGGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAGTGGCGA
AGGCGGCTTACTGGACGATAACTGACGCTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTAGTC
CACGCCGTAAACGATGAATACTAGGTGTCGGGGAGCATTGCTCTTCGGTGCCGCAGCAAACGCAATAAGTATTCCAC
CTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCCGCACAAGCGGTGGAGCATGTGGTTTAA
TTCGAAGCAACGCGAAGAACCCTTACCAAGTCTTGACATCCCGATGACAGAGTATGTAATGTACYTTCTCTTCGGAGC
ATCGGTGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCAACC
CCTGTCTTAGTAGCCAGCGGTTTCGGCCGGGCACTCTAGGGAGACTGCCAGGGATAACCTGGAGGAAGCGGGGGATG
ACGTCAAATCATCATGCCCCCTTATGACTTGGGCTACACACGTGCTACAATGGCGTAAACAAAGGGAAGCGGAGCCGT
GAGGCCGAGCAAATCTCAAAAATAACGTCTCAGTTCGGACTGTAGTCTGCAACCCGACTACACGAAGCTGGAATCGC
TAGTAATCGCAGATCAGAATGCTGCGGTGAATACGTTCCCGGGTCTTGTACACACCGCCCGTCACACCATGGGAGTT
GGAAATGCCCGAAGTCAGTGACCCAACCGCAAGGAGGGAGCTGCCGAAGGCAGGTTGATAACTGGGGTG

>SEQ ID NO: 93|NR_043142.1|*Flavonifractor plautii* strain Prevot S1 16S ribosomal RNA gene, partial sequence

CGCTGGCGGCGTGCTTAACACATGCAAGTCGAACGGGGTGCTCATGACGGAGGATTTCGTCCAATGGATTGAGTTACC
TAGTGGCGGACGGGTGAGTAACGCGTGAGGAACCTGCCTTGGAGAGGGGAATAACACTCCGAAAGGAGTGCTAATAC
CGCATGAAGCAGTTGGGTGCGATGGCTCTGACTGCCAAAGATTTATCGTCTGAGATGGCCTCGCGTCTGATTAGCT
AGTAGGCGGGGTAAACGGCCACCTAGGCGACGATCAGTAGCCGACTGAGAGGTTGACCGGCCACATTGGGACTGAG
ACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGGGCAATGGGCGCAAGCCTGACCCAGCAACGCCCG
GTGAAGGAAGAAGGCTTTCGGGTTGTAACTTCTTTTGTGCGGGACGAAACAAATGACGGTACCCGACGAATAAGCC
ACGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGTGGCAAGCGTTATCCGGATTTACTGGGTGTAAAGGGCG
GTAGGCGGGATTGCAAGTCAGATGTGAAACTGGGGGCTCAACCTCCAGCCTGCATTTGAAACTGTAGTTCTTGAG

AAAGGAATTGACGGACCCGACCAAGCGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCTGCTC
CTTGACATCTCCCTGACCGGCAAGTAATGTTGCCCTTTCCTTCGGGACAGGGATGACAGGTGGTGCATGGTTGTCGTC
AGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCAACCCCTATCTTTAGTAGCCAGCGGTTTGGCCGG
GCACTCTAGAGAGACTGCCAGGGATAACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGAGCAG
GGCTACACACGTGCTACAATGGCGTAAACAAAGGGAGGCAGAACCGCGAGGTTCGAGCAAATCCCAAAAATAACGTCT
CAGTTCGGATTGTAGTCTGCAACTCGACTACATGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGA
ATACGTTCCCGGGTCTTGTACACACCGCCCGTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCGT
AAGGAGGAGCTGCCGAAG >SEQ ID NO: 97|NR_028883.1|*Dorea longicatena* strain 111-35 16S

ribosomal RNA gene, partial sequence

TAACGCGTGGGTAACTGCCTCATAACAGGGGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGACCACGTACC
GCATGGTACAGTGGTAAAACTCCGGTGGTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAACGGCCT
ACCAAGCCGACGATCAGTAGCCGACCTGAGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACG
GGAGGCAGCAGTGGGGAATATTGCACAATGGAGGAACTCTGATGCAGCGACGCCGCGTGAAGGATGAAGTATTTCTG
GTATGTAAACTTCTATCAGCAGGGAAGAAAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGC
CGCGGTAATACGTAGGGGGCAAGCGTTATCCGGATTACTGGGTGTAAAGGGAGCGTAGACGGCACGGCAAGCCAGA
TGTGAAAAGCCCCGGGGCTCAACCCCCGGGACTGCATTTGGAAGTCTGAGCTAGAGTGTTCGGAGAGGCAAGTGAATT
CCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAGTGGCGAAGGCGGCTTGCTGGACGATGACTGACG
TTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGACTGCTAGGTG
TCGGGTGGCAAAGCCATTCCGGTGGCGCAGCTAACGCAATAAGCAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAAA
CTCAAAGGAATTGACGGGGACCCGCACAAGCGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCT
GATCTTGACATCCCGATGACCGCTTCGTAATGGAAGTTTTTCTTCGGAACATCGGTGACAGGTGGTGCATGGTTGTC
GTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCAACCCCTATCTTCAGTAGCCAGCAGGTTAAG
CTGGGCACTCTGGAGAGACTGCCAGGGATAACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGA
CCAGGGCTACACACGTGCTACAATGGCGTAAACAAAGAGAAGCGAACTCGCGAGGGTAAGCAAATCTCAAAAATAAC
GTCTCAGTTCGGATTGTAGTCTGCAACTCGACTACATGAAGCTGGAATCGCTAGTAATCGCAGATCAGAATGCTGCG
GTGAATACGTTCCCGGGTCTTGTACACACCGCCCGTCACACCATGGGAGTCATAACGCCCGAAGTCAGTGACCCAAC
CGTAAGG >SEQ ID NO: 98|NR_029164.1|*Clostridium innocuum* strain B-3 16S ribosomal RNA gene,

partial sequence

ATGGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCATGCCTAATACATGCAAGTCGAACGAAGTCTTCAGGA
AGCTTGCTTCCAAAAAGACTTAGTGCGAACGGGTGAGTAACACGTAGGTAACCTGCCCATGTGTCCGGGATAACTG
CTGGAAACGGTAGCTAAAACCGGATAGGTATACGGAGCGCATGCTCTGTATATTAAAGCGCCCTTCAAGGCGTGAAC
ATGGATGGACCTGCGACGCATTAGCTAGTTGGTGAGGTAACGGCCCCACCAAGGCGATGATGCGTAGCCGGCCTGAGA
GGGTAAACGGCCACATTGGGACTGAGACACGGCCCCAACTCCTACGGGAGGCAGCAGTAGGGAATTTTCGTCAATGG
GGGAAACCTGAACGAGCAATGCCGCGTGAGTGAAGAAGGTCTTCGGATCGTAAAGCTCTGTTGTAAGTGAAGAACC
GCTCATAGAGGAAATGCTATGGGAGTGACGGTAGCTTACCAGAAAGCCACGGCTAACTACGTGCCAGYAGCCGCGGT
AATACGTAGGTGGCAAGCGTTATCCGGAATCATTGGGCGTAAAGGGTTCGTAGGTGGCGTACTAAGTCTGTAGTAAA
AGGCAATGGCTCAACCATTTGTAAGCTATGGAAGTGGTATGCTGGAGTGCAGAAGAGGGCGATGGAATTCATGTGT
AGCGGTAAAATGCGTAGATATATGGAGGAACACCAGTGGCGAAGGCGGTTCGCTGTTGTAAGTGAAGGCA
CGAAAGCGTGGGGAGCAAATAGGATTAGATACCCTAGTAGTCCACGCCGTAAACGATGAGAACTAAGTGTGGAGAA
ATTCAGTGCTGCAGTTAACGCAATAAGTTCTCCGCTGGGGAGTATGCACGCAAGTTNGAACTCAAAGGAATTGAC
GGGGGCCCGCACAAAGCGNTGGAGTATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAGGCCTTGACATGGAA
ACAAATACCCTAGAGATAGGGGGATAATTATGGATCACACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGA
TGTTGGGTAAAGTCCCGCAACGAGCGCAACCCCTTGTGCGCATGTTACCAGCATCAAGTTGGGGACTCATGCGAGACTG
CCGGTGACAAACCGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGGCCTGGGCTACACACGTACTAC
AATGGCGACCACAAAGAGCAGCGACTTGGTGACAAGAAGCGAATCTCATAAAGATCGTCTCAGTTCGGATTGAAGTC
TGCAACTCGACTTCATGAAGTCGGAATCGCTAGTAATCGCAGATCAGCATGCTGCGGTGAATACGTTCTCGGGCCTT
GTACACACCGCCCGTCAAACCATGGGAGTCAGTAATACCCGAAGCCGGTGGCATAACCGTAAGGAGTGAGCCGTGCA
AGGTAGGACCGA >SEQ ID NO: 99|NR_104687.1|*Blautia hansenii* strain JCM 14655 16S ribosomal

RNA gene, partial sequence

AGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTTATCATTGA
CTCTTCGGAAGATTTGATAATTTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATAACAGGGGAA
TAACAGTTAGAAATGGCTGCTAATGCCGCATAAGCGCACAGGACCGCATGGTCTGGTGTGAAAAACTGAGGTGGTAT
GAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAACGGCCTACCAAGCCGACGATCAGTAGCCGGCCTGAGAG
GGTGAACGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGG
GGAACCCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATCTCGGTATGTAACTTCTATCAGCAGGGAAGAAAAT
GACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCCG
GATTTACTGGGTGTAAAGGGAGCGTAGACGGAAGAGCAAGTCTGATGTGAAAGGCTGGGGCTTAACCCCAGGACTGC
ATTGGAAACTGTTTTTCTAGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGG
AGGAACACCAGTGGCGAAGGCGGCTTACTGGACGGTAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGAT
TAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTCGGGGTGCAAAGCAGTTCGGTGGCGCAGCAAA
CGCAATAAGTATTCCACCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCCGCACAAGCGG
TGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCTGCCTGACCGTTCCTTAACCG
GAGCTTTCCTTCGGGACAGGCAAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTC

CCGCAACGACCGACCTAGCCATCTTAGTACCGAGAGCTGCGGAGACTGCCGGGGATAACCC
GGAGGAAGGCGGGGACGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTAAACA
AAGGGAAGCGAAGCGGTGACGCTTAGCAAATCTCAAAAATAACGTCCCAGTTCGGACTGCAGTCTGCAACTCGACTG
CACGAAGCTGGAATCGTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCGCCC
GTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCTTATGGAGGGAGCTGCCGAAGGCGGGACCGAT
AACTGGGGTGAAGTCGTAACAAGGTAACC >SEQ ID NO: 100|NR_112933.1|*Bacteroides cellulosilyticus*
strain JCM 15632 16S ribosomal RNA gene, partial sequence
AGAGTTTGATCCTGGCTCAGGATGAACGCTAGCTACAGGCTTAACACATGCAAGTCGAGGGGCAGCATGACCTAGCA
ATAGGTTGATGGCGACCGGCGCACGGGTGAGTAACACGTATCCAACCTACCGGTTATTCCGGGATAGCCTTTTCGAAA
GAAAGATTAATACCGGATAGTATAACGAGAAGGCATCTTTTTTGTATTAAAGAATTTTCGATAACCGATGGGGATGCG
TTCCATTAGTTTGTGCGGGGTAACGGCCCCACCAAGACATCGATGGATAGGGGTTCTGAGAGGAAGGTCCCCCACA
TTGGAAGTGAAGACACGGTCCAACTCCTACGGGAGGCAGCAGTGAGGAATATTGGTCAATGGACGAGAGTCTGAACC
AGCCAAGTAGCGTGAAGGATGACTGCCCTATGGGTTGTAAACTTCTTTTATATGGGAATAAAGTGAGCCACGTGTGG
CTTTTTGTATGTACCATACGAATAAGGATCGGCTAACTCCGTGCCAGCAGCCGCGGTAATACGGAGGATCCGAGCGT
TATCCGGATTTATTGGGTTTAAAGGGAGCGTAGGCGGACTATTAAGTCAGCTGTGAAAGTTTGCGGCTCAACCGTAA
AATTGCAGTTGATACTGGTCGTCTTGAGTGCAGTAGAGGTAGGCGGAATTCGTGGTGTAGCGGTGAAATGCTTAGAT
ATCACGAAGAACTCCGATTGCGAAGGCAGCTTACTGGACTGTAACCTGACGCTGATGCTCGAAAGTGTGGGTATCAA
CAGGATTAGATACCCTGGTAGTCCACACAGTAAACGATGAATACTCGCTGTTTGCATATACAGCAAGCGGCCAAGC
GAAAGCATTAAAGTATTCCACCTGGGGAGTACGCCGGCAACGGTGAAACTCAAAGGAATTGACGGGGGCCCCGCACAAG
CGGAGGAACATGTGGTTTAATTCGATGATACGCGAGGAACCTTACCCGGGCTTAAATTGCATCTGAATAATTTGGAA
ACAGATTAGCCGCAAGGCAGATGTGAAGGTGCTGCATGGTTGTCGTCAGCTCGTGCCGTGAGGTGTGCGCTTAAGTG
CCATAACGAGCGCAACCCTTATCTTTAGTTACTAACAGGTCATGCTGAGGACTCTAGAGAGACTGCCGTCGTAAGAT
GTGAGGAAGGTGGGGATGACGTCAAATCAGCACGGCCCTTACGTCCGGGGCTACACACGTGTTACAATGGGGGGTAC
AGAAGGCAGCTACACAGCGATGTGATGCTAATCCCAAAGCCTCTCTCAGTTTCGGATTGGAGTCTGCAACCCGACTC
CATGAAGCTGGATTGCTAGTAATCGCGCATCAGCCACGGCGCGGTGAATACGTTCCCGGGCCTTGTACACACCGCC
CGTCAAGCCATGAAAGCCGGGGGTACCTGAAGTCCGTAACCGCAAGGAGCGGCCTAGGGTAAAACTGGTAATTGGGG
CTAAGTCGTA >SEQ ID NO: 101|NR_112940.1|*Bacteroides ovatus* strain JCM 5824 16S ribosomal
RNA gene, partial sequence
GGCTCAGGATGAACGCTAGCTACAGGCTTAACACATGCAAGTCGAGGGGCAGCATTTTATGTTTGCTTGCAAACCTGAA
GATGGCGACCGGCGCACGGGTGAGTAACACGTATCCAACCTGCCGATAACTCCGGAATAGCCTTTTCGAAAGAAAGAT
TAATACCGGATAGCATAACGAATATCGCATGATATTTTTATTAAAGAATTTTCGGTTATCGATGGGGATGCGTTCCATT
AGTTTGTGCGGGGTAACGGCCCCACCAAGACTACGATGGATAGGGGTTCTGAGAGGAAGGTCCCCCACATTGGAAC
TGAGACACGGTCCAACTCCTACGGGAGGCAGCAGTGAGGAATATTGGTCAATGGGCGAGAGCCTGAACCAGCCAAG
TAGCGTGAAGGATGAAGGCTCTATGGGTCGTAAACTTCTTTTATATGGGAATAAAGTTTTCCACGTGTGGAATTTG
TATGTACCATATGAATAAGGATCGGCTAACTCCGTGCCAGCAGCCGCGGTAATACGGAGGATCCGAGCGTTATCCGG
ATTTATTGGGTTTAAAGGGAGCGTAGGTGGATTGTTAAGTCAGTTGTGAAAGTTTGCGGCTCAACCGTAAATTTGCA
GTTGAAACTGGCAGTCTTGAGTACAGTAGAGGTGGGCGGAATTCGTGGTGTAGCGGTGAAATGCTTAGATATCACGA
AGAACTCCGATTGCGAAGGCAGCTCACTAGACTGTTACTGACACTGATGCTCGAAAGTGTGGGTATCAAACAGGATT
AGATACCCTGGTAGTCCACACAGTAAACGATGAATACTCGCTGTTTGCATATACAGTAAGCGGCCAAGCGAAAGCA
TTAAGTATTCCACCTGGGGAGTACGCCGGCAACGGTGAAACTCAAAGGAATTGACGGGGGCCCCGCACAAGCGGAGGA
ACATGTGGTTTAATTCGATGATACGCGAGGAACCTTACCCGGGCTTAAATTGCAACAGAATATATTGGAAACAGTAT
AGCCGTAAGGCTGTTGTGAAGGTGCTGCATGGTTGTCGTCAGCTCGTGCCGTGAGGTGTGCGCTTAAGTGCCATAAC
GAGCGCAACCCTTATCTTTAGTTACTAACAGGTTATGCTGAGGACTCTAGAGAGACTGCCGTCGTAAGATGTGAGGA
AGGTGGGGATGACGTCAAATCAGCACGGCCCTTACGTCCGGGGCTACACACGTGTTACAATGGGGGGTACAGAAGGC
AGCTACCTGGCGACAGGATGCTAATCCCAAACCTCTCTCAGTTTCGGATCGAAGTCTGCAACCCGACTTTCGTGAAG
CTGGATTGCTAGTAATCGCGCATCAGCCATGGCGCGGTGAATACGTTCCCGGGCCTTGTACACACCGCCCGTCAAG
CCATGAAAGCCGGGGGTACCTGAAGTACGTAACCGCAAGGAGCGTCCTAGGGTAAAACTGGTAATTGGGGCTA
>SEQ ID NO: 102|NR_117563.1|*Eubacterium fissicatena* 16S ribosomal RNA gene, partial sequence
TAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCGCTTTACTTAG
ATTTCTTCGGATTGAAGAGTTTTTGCGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACTGCCTCATAACAGGG
GGATAACAGTTAGAAATGACTGCTAATACCGCATAAGACCACAGTACCGCATGGTACAGTGGGAAAAACTCCGGTGG
TATGAGATGGACCCGCGTCTGATTAGCTAGTTGGTAAGGTAACGGCTTACCAAGGCGACGATCAGTAGCCGACCTGA
GAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAAT
GGGGGAAACCCTGATGCAGCGACGCCGCGTGAAGGATGAAGTATTTCCGTATGTAAACTTCTATCAGCAGGGAAGAA
AATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTAT
CCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGTTATGTAAGTCTGATGTGAAAACCCGGGGCTCAACCCCGGGAC
TGCATTGGAACCTATGTAAGTGTGCGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGATCACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAC
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTGCGGTGGCAAAGCCATTCGGTGCCGCAGC
AAACGCAATAAGTATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCTGCTCTTGACATCCCACTGACCGGCGTGTAA
TGGCGCCTTCCCTTCGGGGCAGTGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCTTATCTTTAGTAGCCAGCGTTTTGGCCGGGCACTCTAGAGAGACTGCCAGGGATAA

CCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGAGCAGGGCTACACACGTGCTACAATGGCGTAA
 ACAAAGGGAGGCAATACCGCGAGGTTGAGCAAATCCCCAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCGA
 CTACATGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGACACACCG
 CCCGTCACACCATGGGAGTTGGTAACGCCCCGAAGTCAGTGACCCAACCGTAAGGAGGGAGCTGCCGAAGGCGGGATC
 GATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTT >SEQ ID
 NO: 103|NR_104700.1|*Blautia coccoides* strain JCM 1395 16S ribosomal RNA gene, partial sequence
 AGAGTTTGTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCGCTAAGACAGAT
 TTCTTCGGATTGAAGTCTTTGTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATACAGGGGGA
 TAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGGACCGCATGGTCTGGTGTGAAAAACTCCGGTGGTAT
 GAGATGGACCCGCGTCTGATTAGCTAGTTGGAGGGGTAACGGCCCAACAAGGCGACGATCAGTAGCCGGCCTGAGAG
 GGTGAACGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGG
 GGAAACCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATCTCGGTATGTAACTTCTATCAGCAGGGAAGAAAAT
 GACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCCG
 GATTTACTGGGTGTAAAGGGAGCGTAGACGGAAGAGCAAGTCTGATGTGAAAGGCTGGGGCTTAACCCCAAGGACTGC
 ATTGGAACTGTTGTTCTAGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGG
 AGGAACACCAAGTGGCGAAGGCGGCTTACTGGACGGTAAGTACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGAT
 TAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTGCGGTGGCAAAGCCATTTCGGTGCCGCAGCAAA
 CGCAATAAGTATTCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGCGG
 TGGAGCATGTGGTTTAAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCCTCTGACCGTCCCGTAACGG
 GGGCTTCCCTTCGGGGCAGAGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTGCTGAGATGTTGGGTAAAGTC
 CCGCAACGAGCGCAACCCTTATCCTTAGTAGCCAGCACATGATGGTGGGCACTCTAGGGAGACTGCCGGGGATAACC
 CGGAGGAAGGCGGGGACGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTAAAC
 AAAGGGAAGCGAGACAGCGATGTTGAGCGAATCCCCAAAATAACGTCCCAGTTCGGACTGCAGTCTGCAACTCGACT
 GCACGAAGCTGGAATCGCTAGTAATCGCGGATCAGAATGCCGCGGTGAATACGTTCCCGGGTCTTGACACACCCGCC
 CGTCACACCATGGGAGTCAGTAACGCCCCGAAGTCAGTGACCTAACCAGAAAGGAAGGAGCTGCCGAAGGCGGGACCGA
 TAACTGGGGTGAAGTCGTAACAAGGTAACC >SEQ ID NO: 104|NR_109014.1|*Blautia faecis* strain M25
 16S ribosomal RNA gene, partial sequence

strain 1313 16S ribosomal RNA gene, partial sequence
CTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCGGTTTCAATGAAGTTTTCGGATGGATT
TGAAATTGACTTAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTTACACTGGGGGATAACAGTTAGAAATG
ACTGCTAATACCGCATAAGCGCACAGGGCCGCATGGNCTGGTGTGAAAACTCCGNGGTGTAAGATGGACCCGCGT
CTGATTAGGTAGTTGGNGGGGTAACGGCCCCACCAAGCCGACGATCAGTAGCCGACCTGAGAGGGTGACCGGCCACAT
TGGGACTGAGACACGGCCCCAACTCCTACGGGAGGCAGCAGTGGGGAATATTGGACAATGGGCGAAAGCCTGATCCA
GCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGAAGAAAATGACGGTACCTGACTA
AGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCCGGATTTACTGGGTGTA
AAGGGAGCGTAGACGTTTTAGCAAGTCTGAAGTGAAAGCCCGGGGCTCAACCCCGGTACTGCTTTGGAACTGTTAG
ACTTGAGTGCAGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAAGTGGC
GAAGGCGGCTTACTGGACTGTAAGTACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTAG
TCCACGCCGTAAACGATGAATACTAGGTGTCGGGGGGCAAAGCCCTTCGGTGCCGCCGCAAACGCAATAAGTATTCC
ACCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCCGCACAAGCGGTGGAGCATGTGGTTT
AATTTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCCACTGAAAACACNTTAACCGTGATCCCTCTTCGGA
GCAGTGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCAA
CCCTTATCCTTAGTAGCCAGCGAGTAGAGTCGGGCACTCTGGGGAGACTGCCAGGGATAACCTGGAGGAAGGTGGGG
ATGACGTCAAATCATCATGCCCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTAAACAAAGGGAGGCAAAGG
AGCGATCTGGAGCAAACCCCCAAAAATAACGTCTCAGTTCGGATTGCAGGCTGCAACTCGCCTGCATGAAGCTGGAAT
CGTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTAACACACCGCCCGTCACACCATGGGA

106|NR_113270.1|*Blautia producta* strain JCM 1471 16S ribosomal RNA gene, partial sequence
AGAGTTTGTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTAAGACGGAT
TTCTTCGGATTGAAGTCTTTGTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATACAGGGGGA
TAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGGACCGCATGGTCTGGTGTGAAAACTCCGGTGGTAT
GAGATGGACCCGCGTCTGATTAGCTAGTTGGAGGGGTAAACGGCCACCAAGGCGACGATCAGTAGCCGGCCTGAGAG
GGTGAACGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGG
GGAAACCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATCTCGGTATGTAACTTCTATCAGCAGGGAAGAAAAT
GACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCCG
GATTTACTGGGTGTAAAGGGAGCGTAGACGGAAGAGCAAGTCTGATGTGAAAGGCTGGGGCTTAACCCAGGACTGC
ATTGGAACTGTTGTTCTAGAGTGCCGGAGAGGTAAGCGGAATTCTAGTGTAGCGGTGAAATGCGTAGATATTAGG
AGGAACACCAGTGGCGAAGGCGGCTTACTGGACGGTAAGTACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGAT
TAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTCTGGGTGGCAAAGCCATTCCGGTGCCGCAGCAA
CGCAATAAGTATTCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGCGG
TGGAGCATGTGGTTTAAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCCTCTGACCGTCCCGTAACGG
GGACTTCCCTTCGGGGCAGAGGAGACAGGTGGTGCATGGTTGTCGTACAGTCTGTGTCGTGAGATGTTGGGTAAAGTC
CCGCAACGAGCGCAACCCTTATCCTTAGTAGCCAGCACATGATGGTGGGCACTCTAGGGAGACTGCCGGGGATAACC
CGGAGGAAGGCGGGGACGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTAAAC
AAAGGGAAGCGAGACAGCGATGTTGAGCGAATCCCAAAAATAACGTCCAGTTCGGACTGCAGTCTGCAACTCGACT
GCACGAAGCTGGAATCGCTAGTAATCGCGGATCAGAATGCCGCGGTGAATACGTTCCCGGGTCTTGTACACACCGCC
CGTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCTAACCGAAAGGAAGGAGCTGCCGAAGGCGGGACCGA
TAACTGGGGTGAAGTCGTAACAAGGTAACC >SEQ ID NO: 107|NR_104799.1|*Anaerostipes hadrus* strain

DSM 3319 16S ribosomal RNA gene, partial sequence

TGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTGAACGAAGCTGCTTAACCTGATCTTCTTCGGAAT
TGACGTTTTGTAGACTGAGTGGCGGACGGGTGAGTAACGCGTGGGCAACCTGCCCTGTACAGGGGGATAACAGTCAG
AAATGACTGCTAATACCGCATAAGACCACAGCACCGCATGGTGCAGGGGTAAAACTCCGGTGGTACAGGATGGACC
CGCGTCTGATTAGCTGTTTGGTGAGGTAACGGCTCACCAAGGCGACGATCAGTAGCCGGCTTGAGAGAGTGAACGGC
CACATTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGGGAAACCCTG
ATGCAGCGACGCCGCGTGAGTGAAGAAGTATCTCGGTATGTAAAGCTCTATCAGCAGGGAAGAAAATGACGGTACCT
GACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCCGGAATTACTGG
GTGTAAAGGGTGCCTAGGTGGTATGGCAAGTCAGAAAGTGAACCCAGGGCTTAACTCTGGGACTGCTTTTGAACT
GTCAGACTGGAGTGCAGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACATCA
GTGGCGAAGGCGGCTTACTGGACTGAAACTGACACTGAGGCACGAAAGCGTGGGGAGCAAACAGGATTAGATACCCT
GGTAGTCCACGCCGTAAACGATGAATACTAGGTGTCGGGGCCGTAGAGGCTTCGGTGCCGCAGCCAACGCAGTAAGT
ATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGCGGTGGAGCATGT
GGTTTAATTTCGAAGCAACGCGAAGAACCCTTACCTGGTCTTGACATCCTTCTGACCGGTCCTTAACCGGACCTTTCT
TCGGGACAGGAGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGAGATGTTGGGTAAAGTCCCGCAACGAG
CGCAACCCCTATCTTTAGTAGCCAGCATTTCAGGTGGGCACTCTAGAGAGACTGCCAGGGATAACCTGGAGGAAGGT
GGGGACGACGTCAAATCATCATGCCCTTATGACCAGGGCTACACACGTGCTACAATGGCGTAAACAGAGGGAAGCA
GCCTCGTGAGAGTGAGCAAATCCCCAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCGACTACATGAAGCTG
GAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCGCCCGTCACACCAT
GGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCGTAAGGAGGGAGCTGCCGAAGGCGGGACCGATAACTGGGGT
AAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTTC >SEQ ID NO:

108|NR_117142.1|*Eubacterium fissicatena* strain DSM 3598 16S ribosomal RNA gene, partial sequence

GTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCGCTTTACTTAGATTT
 CTTCCGATTGAAGAGTTTTTGC GACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATA CAGGGGGAT
 AACAGTTAGAAATGACTGCTAATACCGCATAAGACCACAGTACCGCATGGTACAGTGGGAAAACTCCGGTGGTATG
 AGATGGACCCGCGTCTGATTAGCTAGTTGGTAAGGTAACGGCTTACCAAGGCAACGATCAGTAGCCGACCTGAGAGG
 GTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGG
 GAAACCCCTGATGCAGCGACGCCGCGTGAAGGATGAAGTATTTCCGTATGTAACTTCTATCAGCAGGGAAGAAAATG
 ACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAAACGTAGGGGGCAAGCGTTATCCGG
 ATTTACTGGGTGTAAAGGGAGCGTAGACGGTTATGTAAGTCTGATGTGAAAACCCGGGGCTCAACCCCGGGACTGCA
 TTGGAAACTATGTAACTAGAGTGTCCGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGA
 GGAACACCAGTGGCGAAGGCGGCTTACTGGACGATCACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATT
 AGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTCGGGTGGCAAAGCCATTCGGTGCCGCAGCAAAC
 GCAATAAGTATTCCACCTGGGGAGTACGTTCCGAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGCGGT
 GGAGCATGTGGTTTAATTCTGAAGCAACGCGAAGAACCTTACCTGCTCTTGACATCCCACTGACCGGCGTGTAATGGC
 GCCTTCCCTTCGGGGCAGTGGAGACAGGTGGTGCATGGTTGTCTGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCC
 CGCAACGAGCGCAACCCTTATCTTTAGTAGCCAGCGGTTTGCCCGGGCACTCTAGAGAGACTGCCAGGGATAACCTG
 GAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGAGCAGGGCTACACACGTGCTACAATGGCGTAAACAA
 AGGGAGGCAATACCGCGAGGTTGAGCAAATCCCAAAAATAACGTCTCAGTTCCGATTGTAGTCTGCAACTCGACTAC
 ATGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTTACACACCGCCCG
 TCACACCATGGGAGTTGGTAACGCCCGAAGTCAGTGACCCAACCGTAAGGAGGGAGCTGCCGAAGGCGGGATCGATA

CTGGGGTGAAGTCGTAACAAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT>SEQ ID NO: 109|NR_117147.1|*Eubacterium contortum* strain DSM 3982 16S ribosomal RNA gene, partial sequence
TTTGATCTGGCTCAGGATGAACGCTGGCGACGTGCTTAACACATGCAAGTCGAGCGAAGCACTTTACTTTGATTTC
TTCGGAATGAAAGGTTTTGTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATACAGGGGGATA
ACAGTTAGAAATGACTGCTAATACCGCATAAGACCACAGTACCGCATGGTACAGTGGGAAAAACTCCGGTGGTATGA
GATGGACCCGCGTCTGATTAGCTAGTTGGTAAGGTAACGGCTTACCAAGGCGACGATCAGTAGCCGACCTGAGAGGG
TGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGGG
AAACCCTGATGCAGCGACGCCGCGTGAAGGATGAAGTATTTCCGGTATGTAACTTCTATCAGCAGGGAAGAAAATGA
CGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCCGGA
TTTACTGGGTGTAAAGGGAGCGTAGACGGTTATGTAAGTCTGATGTGAAAACCCGGGGCTCAACCCCGGGACTGCAT
TGAAACTATGTAAGTAGAGTGTTCGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAG
GAACACCAGTGGCGAAGGCGGCTTACTGGACGATGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTA
GATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTCGGGTGGCAAAGCCATTCCGTGCCGCAGCAAACG
CAATAAGTATTCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGCGGTG
GAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCTGCTCTTGACATCCCCCTGACCGGCGTGTAATGGTG
CCTTTCCTTCGGGACAGGGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCC
GCAACGAGCGCAACCCTTATCTTTAGTAGCCAGCGGTTTGGCCGGGCACTCTAGAGAGACTGCCAGGGATAACCTGG
AGGAAGGTGGGGATGACGTCAAATCATCATGCCCCCTTATGAGCAGGGCTACACACGTGCTACAATGGCGTAAACAAA
GGGAGGCGAAGCCGTGAGGTGGAGCAAATCCCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCGACTACA
TGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCGCCCCG
CACACCATGGGAGTTGGTAACGCCCGAAGTCAGTGACCCAACCGCAAGGAGGGAGCTGCCGAGGGTGGGACCGATAA
CTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTTCT>SEQ ID NO:

110|NR_113410.1|*Clostridium bolteae* strain JCM 12243 16S ribosomal RNA gene, partial sequence
TTTTAATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGATAACCTGCCTCACACTGGGGGATAACAGTTAGAAAT
GACTGCTAATACCGCATAAGCGCACAGTACCGCATGGTACAGTGTGAAAACTCCGGTGGTGTGAGATGGATCCGCG
TCTGATTAGCCAGTTGGCGGGGTAAACGGCCACCAAAGCGACGATCAGTAGCCGACCTGAGAGGGTGACCGGCCACA
TTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGCGAAAGCCTGATGC
AGCGACGCCGCGTGAGTGAAGAAGTATTTCCGGTATGTAAAGCTCTATCAGCAGGGAAGAAAATGACGGTACCTGACT
AAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCCGGATTTACTGGGTGT
AAAGGGAGCGTAGACGGCGAAGCAAGTCTGAAGTGAAAACCCAGGGCTCAACCCTGGGACTGCTTTGGAAACTGTTT
TGCTAGAGTGTTCGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAAGTGG
CGAAGGCGGCTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTA
GTCCACGCCGTAAACGATGAATGCTAGGTGTTGGGGGGCAAAGCCCTTCGGTGCCGTCGCAAACGCAGTAAGCATT
CACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGCGGTGGAGCATGTGGTT
TAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCTCTTGACCGGCGTGTAACGGCGCCTTCCCTTCGG
GGCAAGAGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCA
ACCCTTATCCTTAGTAGCCAGCAGGTAAAGCTGGGCACTCTAGGGAGACTGCCAGGGATAACCTGGAGGAAGGTGGG
GATGACGTCAAATCATCATGCCCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTAAACAAAGGGAAGCAAGA
CAGTGATGTGGAGCAAATCCCAAAAATAACGTCCCAGTTTCGGACTGTAGTCTGCAACCCGACTACACGAAGCTGGAA
TCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCGCCCCGTCACACCATGGG
AGTCAGCAACGCCCGAAGTCAGTGACCCAACTCGCAAGAGAGGGAGCTGCCGAAGGCGGGGCAGGTAACCTGGGGTG
AGTC>SEQ ID NO: 111|NR_041960.1|*Blautia luti* strain BInIX 16S ribosomal RNA gene, complete
sequence

GTGGGTAACCTGCCTTATACAGGGGGATAACAGTCAGAAATGACTGCTAATACCGCATAAGCGCACAGAGCTGCATG
GCTCCGGTGTGAAAAACTCCGGTGGTATAAGATGGACCCGCGTTGGATTAGCTAGTTGGTGAGGTAACGGCCACCA
AGGCGACGATCCATAGCCGGCCTGAGAGGGTGAACGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAG
GCAGCAGTGGGGAATATTGCACAATGGGGGAAACCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATCTCGGTAT
GTAACTTCTATCAGCAGGGAAGAAAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCG
GTAATACGTAGGGGGCAAGCGTTATCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCATGGACAAGTCTGATGT
GAAAGGCTGGGGCTCAACCCCGGGACTGCATTGGAAACTGCCCGTCTTGAGTGCCGGAGAGGTAAGCGGAATTCCTA
GTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAAGTGGCGAAGGCGGCTTACTGGACGGTAACCTGACGTTGA
GGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTAGTCCACGCGGTAAACGATGAATCCTAGGTGTTCGG
GGAGCAAANNNTTCGGTGCCGCGCAAACGCATTAAGCATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCA
AAGGAATTGACGGGGACCCGCACAAGCGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAAGTC
TTGACATCCCTCTGACCGAGTATGTATGGTACTTTTCCTTCGGGAGAGAGAGGAGACAGGTGGTGCATGGTTGTCGT
CAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCAACCCCTATCCCCAGTAGCCAGCGGTTTCGGCCG
GGCACTCTGAGGAGACTGCCAGGGATAACCTGGAGGAAGGCGGGGATGACGTCAAATCATCATGCCCCCTTATGATTT
GGGCTACACACGTGCTACAATGGCGTAAACAAAGGGAAGCAAGCCTGCGAGGGTGGGCAAATCCCAAAAATAACGTC
CCAGTTCGGACTGTAGTCTGCAACCCGACTACACGAAGCTGGAATCGCTAGTAATCGCGGATCAGAATGCCGCGGTG
AATACGTTCCCGGGTCTTGTACACACCGCCCCGTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCTAACT
>SEQ ID NO: 112|NR_074306.1|*Acidaminococcus intestini* RyC-MR95 strain RyC-MR95 16S ribosomal
RNA, complete sequence
CTGGCGGCGTGCTTAACACATGCAAGTCGAACGGAGAACTTATTTCCGGTAAGTTCTTAGTGGCGAACGGGTGAGTAA

CGCGTGGGCAACCTGCCCTCCAGTTGGGGACAACATTCCGAAAGGGATGCTAATACCGAATGTCCTCCCTCCTCCGC
ATGGAGGAGGGAGGAAAGATGGCCTCTGCTTGCAAGCTATCGCTGGAAGATGGGCCCGCTCTGATTAGCTAGTTGG
TGGGGTAACGGCTACCAAGGCGATGATCAGTAGCCGGTCTGAGAGGATGAACGGCCACATTGGGACTGAGACACGG
CCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATCTTCCGCAATGGACGAAAGTCTGACGGAGCAACGCCGCGTGAGT
GATGAAGGTCTTTCGGATTGTAAAACTCTGTTGTTAGGGACGAAAGCACCGTGTTTCAACAGGTCATGGTGTTGACGG
TACCTAACGAGGAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGTGGCAAGCGTTGTCCGGAATT
ATTGGGCGTAAAGAGCATGTAGGCGGGCTTTTAAGTCTGACGTGAAAATGCGGGGCTTAACCCCGTATGGCGTTGGA
TACTGGAAGTCTTGAGTGCAGGAGAGGAAAGGGGAATTCCCAGTGTAGCGGTGAAATGCGTAGATATTGGGAGGAAC
ACCAGTGGCGAAGGCGCCTTTCTGGACTGTGTCTGACGCTGAGATGCGAAAGCCAGGGTAGCAAACGGGATTAGATA
CCCCGGTAGTCCTGGCCGTAAACGATGGATACTAGGTGTAGGAGGTATCGACCCCTTCTGTGCCGGAGTTAACGCAA
TAAGTATCCCGCCTGGGGACTACGATCGCAAGATTGAAACTCAAAGGAATTGACGGGGGCCCCGCACAAGCGGTGGAG
TATGTGGTTTAATTCGACGCAACGCGAAGAACCTTACCAAGGCTTGACATTGAGTGAAAGACCTAGAGATAGGTCCC
TCCCTTCGGGGACACGAAAACAGGTGGTGCATGGCTGTCGTCAGCTCGTGTCGTGAGATGTTGGGTAAAGTCCCGCA
ACGAGCGCAACCCCTATCCTATGTTACCAGCGCGTAAAGGCGGGGACTCATAGGAGACTGCCAGGGATAACTTGGAG
GAAGGCGGGGATGACGTCAAGTCATCATGCCCTTATGTCTTGGGCTACACACGTACTACAATGGTCGGCAACAAAG
GGCAGCGAAACCGCGAGGTGGAGCAAATCCAGAAACCCGACCCAGTTCGGATCGTAGGCTGCAACCCGCCTACGT
GAAGTTGGAATCGCTAGTAATCGCAGGTCAGCATACTGCGGTGAATACGTTCCCGGGCCTTGACACACCGCCCGTC
ACACCACGAAAGTTGGTAACACCCGAAGCCGGTGAGATAACCTTTTAGGAGTCAGCTGTCTAAGGTGGGGCCGATGA
TTGGGGTGAAGTCGTAACAAGGTAGC >SEQ ID NO: 113|NR_074399.1|*Ruminococcus albus* strain 7

AGAGTTGATTCGATCGCTGAGGCTGAGGACGCTGAGGCTGCTTAACACATGCAAGTCGAACGGGAGAACTTCTTCGGA
ATGTTCTTAGTGGCGAACGGGTGAGTAACGCGTAGGCAACCTGCCCTCTGGTTGGGGACAACATTCCGAAAGGGATG
CTAATACCGAATGAGATCCTCTTTCCGCATGGAGAGAGGATGAAAGATGGCCTCTACTTGTAAAGCTATCGCCAGAAG
ATGGGCCTGCGTCTGATTAGCTAGTAGGTGAGGTAACGGCTCACCTAGGCGATGATCAGTAGCCGGTCTGAGAGGAT
GAACGGCCACATTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATCTTCCGCAATGGACGA
AAGTCTGACGGAGCAACGCCGCGTGAGTGATGAAGGCCTTCGGGTTGTAAAACCTCTGTTGTCAGGGACGAAAGCACC
GATCTATAATACATTTTGGTGTGACGGTACCTGACGAGGAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGTAAT
ACGTAGGTGGCAAGCGTTGTCCGGAATTATTGGGCGTAAAGAGCATGTAGGCGGGCTTTTAAGTCCGACGTGAAAAT
GCGGGGCTTAACCCCGTATGGCGTTGGATACTGGAAGTCTTGAGTGCAGGAGAGGAAAGGGGAATTCCCACTGTAGC
GGTGAAATGCGTAGATATTGGGAGGAACACCAGTGGCGAAGGCGCCTTTCTGGACTGTGTCTGACGCTGAGATGCGA
AAGCCAGGGTAGCAAACGGGATTAGATACCCCGGTAGTCCTGGCCGTAAACGATGGGTACTAGGTGTAGGAGGTATC
GACCCCTTCTGTGCCGGAGTTAACGCAATAAGTACCCCGCCTGGGGACTACGATCGCAAGATTGAAACTCAAAGGAA
TTGACGGGGGCCCCGCACAAGCGGTGGAGTATGTGGTTTAATTTCGACGCAACGCGAAGAACCTTACCAAGGCTTGACA
TTGAGTGAAAGACCCAGAGATGGGTCCCCTTCTTCGGAAGCACGAAACAGGTGGTGCATGGCTGTCGTCAGCTCGT
GTCGTGAGATGTTGGGTTAAGTCCCGCAACGAGCGCAACCCTTATCCTATGTTACCAGCACGTAATGGTGGGGACTC
ATAGGAGACTGCCAGGGATAACCTGGAGGAAGGCGGGGATGACGTCAAGTCATCATGCCCTTATGTCTTGGGCTAC
ACACGTACTACAATGGTCGGCAACAAAGGGCAGCGAAGCCGCGAGGCGGAGCCAATCCAGAAACCCGACCCCACT
CGGATCGCAGGCTGCAACCCGCGCTGCGTGAAGTTGGAATCGCTAGTAATCGCAGGTCAGCATACTGCGGTGAATACG
TTCCCGGGCCTTGTACACACCCGCCGTCACACCACGAAAGTTGGTAACACCCGAAGCCGGTGAGATAACCTTTTAGG
AGTCAGCTGTCTAAGGTGGGGCCGATGATTGGGGTGAAGTCGTAACAAGGTAGCCGTTTCGAGAACGAGCGGCTGGAT
CACCT>SEQ ID NO: 116|NR_114326.1|*Fusicatenibacter saccharivorans* strain HT03-11 16S ribosomal

RNA gene, partial sequence

TGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCAGTTAAGAAGATTYTTCCGGATGAT
TCTTGACTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTGACCTGCCCCATACCGGGGGATAACAGCTGGAAAC
GGCTGCTAATACCGCATAAGCGCACAGAGCTGCATGGCTCGGTGTGAAAACTCCGGTGGTATGGGATGGGCCCCGCG
TCTGATTAGGCAGTTGGCGGGGTAAACGGCCCCACAAACCGACGATCAGTAGCCGGCCTGAGAGGGCGACCGGCCACA
TTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGGGAAACCCTGATGC
AGCGACGCCGCGTGAGCGAAGAAGTATTTCCGGTATGTAAAGCTCTATCAGCAGGGAAGATAATGACGGTACCTGACT
AAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCCGGATTTACTGGGTGT
AAAGGGAGCGTAGACGGCAAGGCAAGTCTGATGTGAAAACCCAGGGCTTAACCCTGGGACTGCATTGGAAACTGTCT
GGCTCGAGTGCCGAGAGGTAAGCGGAATTTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAAGAACACCAGTGG
CGAAGGCGGCTTACTGGACGGTAACCTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTA
GTCCACGCCGTAAACGATGAATGCTAGGTGTTGGGGAGCAAAGCTCTTCGGTGCCGCCGCAACGCATTAAGCATT
CACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGCGGTGGAGCATGTGGTT
TAATTGCAAGCAACGCGAAGAACCTTACCAGGTCTTGACATCCCGATGACCGGGCCCGTAACGGGGCCTTCTCTTCGG
AGCATTGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTGTCGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCA
ACCCTTATCCTCAGTAGCCAGCAGGTAAAGCTGGGCACTCTGTGGAGACTGCCAGGGATAACCTGGAGGAAGGTGGG
GATGACGTCAAATCATCATGCCCTTATGATCTGGGCTACACACGTGCTACAATGGCGTAAACAAAGGGAGGCAAAG
CCGCGAGGTGGAGCAAATCCCAAAAATAACGTCTCAGTTCGGACTGCAGTCTGCAACTCGACTGCACGAAGCTGGAA
TCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTAACACCCGCCGTCACACCATGGG
AGTTGGTAACGCCCGAAGTCAGTGACCCAACCTTTTA>SEQ ID NO: 117|NR_102884.1|*Ruminococcus*

champanellensis strain 18P13 16S ribosomal RNA gene, complete sequence

AGAGTTTGATCCTGGCTCAGGACGAACGCTGGCGGCACGCCTAACACATGCAAGTCGAACGGAGATAAAGACTTCGG
TTTTTATCTTAGTGGCGGACGGGTGAGTAACACGTGAGCAACCTGCCTCTGAGAGAGGGATAGCTTCTGGAAACGGA
TGTAATACCTCATAACATAGCGGTACCGCATGATACTGCTATCAAAGATTTATCGCTCAGAGATGGGCTCGCGTCT
GATTAGCTAGATGGTGAGGTAACGGCTCACCATGGCGACGATCAGTAGCCGGACTGAGAGGTTGAACGGCCACATTG
GGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGCGCAAGCCTGATGCAGC
GATGCCGCGTGAGGAAGAAGGTTTTTCGGATTGTAAACTCCTGTCTTAAGGGACGATAATGACGGTACCTTAGGAGG
AAGCTCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGAGCGAGCGTTGTCCGGAATTACTGGGTGTAAA
GGGAGCGTAGGCGGGATTGCAAGTCAGATGTGAAAACCTATGGGCTTAACCCATAGACTGCATTTGAAACTGTAGTTC
TTGAGTGAAGTAGAGGTAAGCGGAATTTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACATCGGTGGCGA
AGGCGGCTTACTGGGCTTTTACTGACGCTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCCTGGTAGTC
CACGCTGTAAACGATGATTACTAGGTGTGGGGGGACTGACCCCTTCCGTGCCGCAGTTAACACAATAAGTAATCCAC
CTGGGGAGTACGGCCGCAAGGTTGAAACTCAAAGGAATTGACGGGGGCCCCGCACAAGCAGTGAGATATGTGGTTTAA
TTCGAAGCAACGCGA AAAACCTTACCAGGTCTTGACATCGAGTGAATGATCTAGAGATAGATCAGTCCTTCGGGACA
CAAAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTGTCGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCAACCCT
TACCTTTAGTTGCTACGCAAGAGCACTCTAGAGGGACTGCCGTTGACAAAACGGAGGAAGGTGGGGATGACGTCAA
TCATCATGCCCTTATGACCTGGGCTACACACGTACTACAATGGCAATGAACAGAGGGAAGCAATACAGTGATGTGG
AGCAAATCCCCAAAATTGTCCAGTTTCAGATTGTAGGCTGCAACTCGCCTACATGAAGTCGGAATTGCTAGTAATC
GCAGATCAGCATGCTGCGGTGAATACGTTCCCGGGCCTTGTAACACCCGCCGTCACACCATGGGAGTCGGTAACAC
CCGAAGCCAGTAGCCTAACCGCAAGGAGGGCGCTGTGCAAGGTGGGATTGATGACTGGGGTGAAGTCGTAACAAGGT
AGCCGTATCGGAAGGTGCGGCTGGATCACCT>SEQ ID NO: 118|NR_102971.1|*Bifidobacterium bifidum*

S17 strain S17 16S ribosomal RNA, complete sequence

ribosomal RNA gene, partial sequence

GGTATAACTTAGTGGCGGACGGGTGAGTAACGCGTGGGAAACCTGCCCTGTACCGGGGGATAACACTTAGAAATAGG
TGCTAATACCGCATAAGCGCACGGAACCGCATGGTTCCGTGTGAAAACTCCGGTGGTACAGGATGGTCCCGCGTCT
GATTAGCCAGTTGGCAGGGTAACGGCCTACCAAAGCGACGATCAGTAGCCGGCCTGAGAGGGTGAACGGCCACATTG
GGACTGAGACACGGCCCCAACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGGGGGAAACCTGATGCAGC
GACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGAAGAAAATGACGGTACCTGACTAAG
AAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCCGGATTTACTGGGTGTAAA
GGGAGCGTAGACGGCATGGCAAGCCAGATGTGAAAACCCAGGGCTCAACCTTGGGATTGCATTTGGAAGTGCCAGGC
TGGAGTGCAGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAGGAGGAACACCAGTGGCGA
AGGCGGCTTACTGGACTGTAAGTACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGATTAGATACCTTGGTAGTC
CACGCGGTAAACGATGATTGCTAGGTGTAGGTGGGTATGGACCCATCGGTGCCGCAGCTAACGCAATAAGCAATCCA
CCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCCGCACAAGCGGTGGAGCATGTGGTTTA
ATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCCAATGACGCACCTGTAAAGAGGTGTTCCCTTCGGGG
CATTGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCGCAACGAGCGCAAC
CCTTATTCTTAGTAGCCAGCAGGTAAAGCTGGGCACTCTAAGGAGACTGCCGGGGATAACCCGGAGGAAGGCGGGGA
TGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTAAACAAAGGGAAGCGAGACA
GTGATGTGGAGCAAATCYCAGAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCGACTACATGAAGCTGGAATC
GCTAGTAATCGCGAATCAGCATGTGCGCGGTGAATACGTTCCCGGGTCTTGTACACACCGCCCCGTCACACCATGGGAG
TTGGAATGCCCGAAGTCTGTGACCTAACCGAAAGGGAGGAGCAGCCGAAGGCAGGTCTGATAACTGGGGTGAAGTC
GTAA>SEQ ID NO: 122|NR_118730.1|*Clostridium symbiosum* strain ATCC 14940 16S ribosomal

RNA gene, partial sequence

AAACATGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCGATTT
AACGGAAGTTTTTCGGATGGAAGTTGAATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGGTAACTGCCTTGTA
TGGGGGACAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTATTGCATGATACAGTGTGAAAACTCCG
GTGGTACAAGATGGACCCGCGTCTGATTAGCTAGTTGGTAAGGTAACGGCTTACCAAGGCGACGATCAGTAGCCGAC
CTGAGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCNNACTCCTACGGGAGGCAGCAGTGGGGAATATTGCA
CAATGGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGA
AGAAAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGNAGCG
TTATCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGTAAAGCAAGTCTGAAGTGAAAGCCCGCGNCTCAACTGCG
GNNCTGCTTTGGAACTGTTTAACTGGAGTGTGCGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGA
TATTAGGAGGAACACNAGTGGCGAAGGCGACTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAA
ACAGGATTAGATACCTTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTTGGGGAGCAAAGCTCTTCGGTGCCG
TCGCAAACGCAGTAAGTATTCACCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCNCA
CAAGCGGTGGAGCATGTGGTTTAATTCTGAANNAACGCGAAGAACCTTACCAGGTCTTGACATCGACTCGACGGGGGA
GTAACGTCCCNNTNCCTTCGGGGCGGAGAAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGG
TTNAGTCCCGCAACGAGCGCAACCCTTATTCTAAGTAGCCAGCGGTTTCGGCCGGGAACTCTTGGGAGACTGCCAGGG
ATAACCTGGAGGAAGGTGGGGATGACGTCNAATCATCATGCCCTTATGATCTGGGCTACACACGTGCTACAATGGC
GTAAACANAGAGAAGCAAGACCGCGAGGTGGAGCAAATCTCAAAAATAACGTCTCAGTTCGGACTGCAGGCTGCAAC
TCGCCTGCACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTACAC
ACCGNNGCTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCGCAAGGAGGGAGCTGCCGAAGGCGC
GACCGANAACNNGG>SEQ ID NO: 123|NR_113243.1|*Erysipelatoclostridium ramosum* strain JCM 1298

16S ribosomal RNA gene, partial sequence

AGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAATACATGCAAGTCGAACGCGAGCACTTGTGCTCG
AGTGGCGAACGGGTGAGTAATACATAAGTAACCTGCCCTAGACAGGGGGATAACTATTGGAAACGATAGCTAAGACC
GCATAGGTACGGACACTGCATGGTGACCGTATTAAGAGTGCCTCAAAGCACTGGTAGAGGATGGACTTATGGCGCAT
TAGCTGGTTGGCGGGGTAAACGGCCACCAAGGCGACGATGCGTAGCCGACCTGAGAGGGTGACCGGCCACACTGGGA
CTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTAGGGAATTTTCGGCAATGGGGGAAACCTGACCGAGCAAC
GCCGCGTGAAGGAAGAAGGTTTTTCGGATTGTAACTTCTGTTATAAAGGAAGAACGGCGGCTACAGGAAATGGTAGC
CGAGTGACGGTACTTTATTAGAAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGTGGCAAGCGTT
ATCCGGAATTATTGGGCGTAAAGAGGGAGCAGGCGGCAGCAAGGGTCTGTGGTGAAAGCCTGAAGCTTAACTTCAGT
AAGCCATAGAAACCAGGCAGCTAGAGTGCAGGAGAGGATCGTGGAATTCATGTGTAGCGGTGAAATGCGTAGATAT
ATGGAGGAACACCAGTGGCGAAGGCGACGATCTGGCCTGCAACTGACGCTCAGTCCCGAAAGCGTGGGGAGCAAATA
GGATTAGATACCTAGTAGTCCACGCCGTAAACGATGAGTACTAAGTGTGGATGTCAAAGTTCAGTGCTGCAGTTA
ACGCAATAAGTACTCCGCTGAGTAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGGCCCCGCACAAGCG
GTGGAGCATGTGGTTTAATTCTGAAGCAACGCGAAGAACCTTACCAGGTCTTGACATACTCATAAAGGCTCCAGAGAT
GGAGAGATAGCTATATGAGATACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGTCCCG
CAACGAGCGCAACCCTTATCGTTAGTTACCATCATTAAAGTTGGGGACTCTAGCGAGACTGCCAGTGACAAGCTGGAG
GAAGGCGGGGATGACGTCAAATCATCATGCCCTTATGACCTGGGCTACACACGTGCTACAATGGATGGTGCAGAGG
GAAGCGAAGCCGCGAGGTGAAGCAAAACCCATAAAACCATTCAGTTCGGATTGTAGTCTGCAACTCGACTACATG
AAGTTGGAATCGCTAGTAATCGCGAATCAGCATGTGCGCGGTGAATACGTTCTCGGGCCTTGTACACACCGCCCCGTC
CACCACGAGAGTTGATAACACCCGAAGCCGGTGGCCTAACCGCAAGGAAGGAGCTGTCTAAGGTGGGATTGATGATT
GGGGTGAAGTCGTAACAAGGTAACC>SEQ ID NO: 124 |PROKKA_00507|16S ribosomal RNA

gene | VE202-7

ATGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCAATTAAGG
AAGTTTTTCGGATGGAATTTGATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGATAACCTGCCTCACACTGGGGG
ATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTACCGCATGGTACGGTGTGAAAACTCCGGTGGTG
TGAGATGGATCCGCGTCTGATTAGCCAGTTGGCGGGGTAACGGCCCAACAAAGCGACGATCAGTAGCCGACCTGAGA
GGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCTACGGGAGGCAGCAGTGGGGAATATTGCACAATGG
GCGAAAGCCTGATGCAGCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGAAGAAAA
TGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATCC
GGATTCACTGGGTGTAAAGGGAGCGTAGACGGCGAAGCAAGTCTGAAGTGAAAACCCAGGGCTCAACCCTGGGACTG
CTTTGGAAACTGTTTTGCTAGAGTGTTCGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTAG
GAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGGA
TTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATGCTAGGTGTTGGGGGGCAAAGCCCTTCGGTGCCGTCGCAA
ACGCAGTAAGCATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGCG
GTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCTCTTGACCGGCGTGTAACG
GCGCCTTCCCTTCGGGGCAAGAGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGT
CCCGCAACGAGCGCAACCCTTATCCTTAGTAGCCAGCAGGTAGAGCTGGGCACTCTAGGGAGACTGCCAGGGATAAC
CTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTAAA
CAAAGGGAAGCAAGACAGTGATGTGGAGCAAATCCCAAAAATAACGTCCCAGTTCGGACTGTAGTCTGCAACCCGAC
TACACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCCG
CCGTCACACCATGGGAGTCAGCAACGCCCGAAGTCAGTGACCCAACTCGCAAGAGAGGGAGCTGCCGAAGGCGGGG
AGGTAAGTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 125 |PROKKA_00709 16S ribosomal RNA gene | VE202-7

ATGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCAATTAAT
GAAGTTTTTCGGATGGATTTTTGATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGATAACCTGCCTCACACTGGG
GGATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTACCGCATGGTACGGTGTGAAAACTCCGGTGG
TGTGGGATGGATCCGCGTCTGATTAGCCAGTTGGCGGGGTAACGGCCCAACAAAGCGACGATCAGTAGCCGACCTGA
GAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCTACGGGAGGCAGCAGTGGGGAATATTGCACAAT
GGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGAAGAA
AATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTAT
CCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCGAAGCAAGTCTGAAGTGAAAACCCAGGGCTCAACCCTGGGAC
TGCTTTGGAAACTGTTTTGCTAGAGTGTTCGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAG
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATGCTAGGTGTTGGGGGGCAAAGCCCTTCGGTGCCGTCGC
AAACGCAGTAAGCATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCTCTTGACCGGCGTGTA
CGGCGCCTTCCCTTCGGGGCAAGAGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCTTATCCTTAGTAGCCAGCAGGTAAAGCTGGGCACTCTAGGGAGACTGCCAGGGATA
ACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTA
AACAAAGGGAAGCAAGACAGTGATGTGGAGCAAATCCCAAAAATAACGTCCCAGTTCGGACTGTAGTCTGCAACCCG
ACTACACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTACACACC
GCCCGTCACACCATGGGAGTCAGCAACGCCCGAAGTCAGTGACCCAACTCGCAAGAGAGGGAGCTGCCGAAGGCGGG
GCAGGTAAGTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 126 |PROKKA_01766 16S ribosomal RNA gene | VE202-7

ATGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCAATTAAT
GAAGTTTTTCGGATGGATTTTTGATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGATAACCTGCCTCACACTGGG
GGATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTACCGCATGGTACGGTGTGAAAACTCCGGTGG
TGTGAGATGGATCCGCGTCTGATTAGCCAGTTGGCGGGGTAACGGCCCAACAAAGCGACGATCAGTAGCCGACCTGA
GAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCTACGGGAGGCAGCAGTGGGGAATATTGCACAAT
GGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGAAGAA
ATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATC
CGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCGAAGCAAGTCTGAAGTGAAAACCCAGGGCTCAACCCTGGGACT
GCTTTGGAAACTGTTTTGCTAGAGTGTTCGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTA
GGAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAG
ATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATGCTAGGTGTTGGGGGGCAAAGCCCTTCGGTGCCGTCGCAA
ACGCAGTAAGCATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGCG
GTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCTCTTGACCGGCGTGTAACG
GCGCCTTCCCTTCGGGGCAAGAGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAAAGT
CCCGCAACGAGCGCAACCCTTATCCTTAGTAGCCAGCAGGTAAAGCTGGGCACTCTAGGGAGACTGCCAGGGATAAC
CTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTAAA
CAAAGGGAAGCAAGACAGTGATGTGGAGCAAATCCCAAAAATAACGTCCCAGTTCGGACTGTAGTCTGCAACCCGAC
TACACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCCG
CCGTCACACCATGGGAGTCAGCAACGCCCGAAGTCAGTGACCCAACTCGCAAGAGAGGGAGCTGCCGAAGGCGGGG
AGGTAAGTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 127 |PROKKA_01779 16S ribosomal RNA gene | VE202-7

ATGAGAGTTTGTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCAATTAATAAAT
GAAGTTTTTCGGATGGATTTTTTATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGATAACCTGCCTCACACTGGG
GGATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTACCGCATGGTACGGTGTGAAAAACTCCGGTGG
TGTGAGATGGATCCGCGTCTGATTAGCCAGTTGGCGGGGTAACGGCCCAACAAAGCGACGATCAGTAGCCGACCTGA
GAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCTACGGGAGGCAGCAGTGGGGAATATTGCACAAT
GGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGAAGAA
AATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGGCAAGCGTTAT
CCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCGAAGCAAGTCTGAAGTGAAAACCCAGGGCTCAACCCTGGGAC
TGCTTTGGAAACTGTTTTGCTAGAGTGTTCGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAC
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATGCTAGGTGTTGGGGGCAAAGCCCTTCGGTGCCGTCGCA
AACGCAGTAAGCATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGC
GGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCTCTTGACCGGCGTGTAAC
GGCGCCTTCCCTTCGGGGCAGGAGAGACAGGTGGTGCATGGTTGTCTCAGCTCGTGTCTGTGAGATGTTGGGTAAAG
TCCCGCAACGAGCGCAACCCTTATCCTTAGTAGCCAGCAGGTAGAGCTGGGCACTCTAGGGAGACTGCCAGGGATAA
CCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTAA
ACAAAGGGAAGCAAGACAGTGATGTGGAGCAAATCCCAAATAACGTCCCAGTTTCGGACTGTAGTCTGCAACCCGAC
TACACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCCG
CCGTCACACCATGGGAGTCAGCAACGCCCGAAGTCAGTGACCCAACTCGCAAGAGAGGGAGCTGCCGAAGGCGGGG
AGGTAACCTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 128 |PROKKA_05926 16S ribosomal RNA gene | VE202-7

ATGAGAGTTTGTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCAATTAATAAAT
GAAGTTTTTCGGATGGATTTTAATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGATAACCTGCCTCACACTGGGG
GATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTACCGCATGGTACGGTGTGAAAAACTCCGGTGGT
GTGAGATGGATCCGCGTCTGATTAGCCAGTTGGCGGGGTAACGGCCCAACAAAGCGACGATCAGTAGCCGACCTGAG
AGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCTACGGGAGGCAGCAGTGGGGAATATTGCACAATG
GGCGAAAGCCTGATGCAGCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGAAGAAA
ATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTATC
CGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCGAAGCAAGTCTGAAGTGAAAACCCAGGGCTCAACCCTGGGACT
GCTTTGGAAACTGTTTTGCTAGAGTGTTCGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATTA
GGAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAGC
ATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATGCTAGGTGTTGGGGGGCAAAGCCCTTCGGTGCCGTCGCA
AACGCAGTAAGCATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAGC
GGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCTCTTGACCGGCGTGTAAC
GGCGCCTTCCCTTCGGGGCAAAGAGAGACAGGTGGTGCATGGTTGTCTCAGCTCGTGTCTGTGAGATGTTGGGTAAAG
TCCCGCAACGAGCGCAACCCTTATCCTTAGTAGCCAGCAGGTAAAGCTGGGCACTCTAGGGAGACTGCCAGGGATAA
CCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCGTAA
ACAAAGGGAAGCAAGACAGTGATGTGGAGCAAATCCCAAATAACGTCCCAGTTTCGGACTGTAGTCTGCAACCCGAC
TACACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCCG
CCGTCACACCATGGGAGTCAGCAACGCCCGAAGTCAGTGACCCAACTCGCAAGAGAGGGAGCTGCCGAAGGCGGGG
AGGTAACCTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 129 | PROKKA_01784 16S ribosomal RNA gene

TCAAAGAGTTTGTATCCTGGCTCAGGACGAACGCTGGCGGCGCGCCTAACACATGCAAGTCGAACGGAGCTTACGTTT
TGAAGTTTTTCGGATGGATGAATGTAAGCTTAGTGGCGGACGGGTGAGTAACACGTGAGCAACCTGCCTTTCAGAGGG
GGATAACAGCCGGAACGGCTGCTAATACCGCATGATGTTGCGGGGGCACATGCCCTGCAACCAAAGGAGCAATCC
GCTGAAAGATGGGCTCGCGTCCGATTAGCCAGTTGGCGGGGTAACGGCCCAACAAAGCGACGATCGGTAGCCGGACT
GAGAGGTTGAACGGCCACATTGGGACTGAGACACGGCCAGACTCCTACGGGAGGCAGCAGTGGGGGATATTGCACA
ATGGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGGGAAGACGGTCTTCGGATTGTAAACCTCTGTCTTTGGGGAAG
AAAATGACGGTACCCAAAGAGGAAGCTCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGAGCAAGCGTT
GTCCGGAATTACTGGGTGTAAAGGGAGCGTAGGCGGGATGGCAAGTAGAATGTAAATCCATCGGCTCAACCGGTGG
CTGCGTTCTAAACTGCCGTTCTTGAGTGAAGTAGAGGCAGGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAGTGGCGAAGGCGGCCTGCTGGGCTTTAACTGACGCTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGATTACTAGGTGTGGGGGGACTGACCCCTTCCGTGCCGCAG
TTAACACAATAAGTAATCCACCTGGGGAGTACGGCCGCAAGGTTGAAACTCAAAGGAATTGACGGGGGCCCGCACA
GCAGTGGAGTATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAGGTCTTGACATCGGATGCATAGCCTAGAG
ATAGGTGAAGCCCTTCGGGGCATCCAGACAGGTGGTGCATGGTTGTCTCAGCTCGTGTCTGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCTTATTATTAGTTGCTACGCAAGAGCACTCTAATGAGACTGCCGTTGACAAAACGGA
GGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGACCTGGGCTACACACGTACTACAATGGCACTAAAACAG
AGGGCGGCGACACCGCGAGGTGAAGCGAATCCCGAAAAAGTGCTCTCAGTTCAGATTGCAGGCTGCAACCCGCTGCA
TGAAGTCGGAATTGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCCCGGGCCTTGTACACACCGCCCGT
CACACCATGGGAGTCGGTAACACCCGAAGCCAGTAGCCTAACCGCAAGGGGGGCGCTGTCTGAAGGTGGGATTGATGA
CTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID NO:

130 | PROKKA_01864 16S ribosomal RNA gene

TCAAAGAGTTTATCCTGGCTCAGGACGAACGCTGGCGGCGCGCCTAACACATGCAAGTCGAACGGAGCTTACGTTT
TGAAGTTTTTCGGATGGACGAATGTAAGCTTAGTGCGGACGGGTGAGTAACACGTGAGCAACCTGCCTTTCAGAGGG
GATAACAGCCGGAACGGCTGCTAATACCGCATGATGTTGCGGGGGCACATGCCCTGCAACCAAAGGAGCAATCCG
CTGAAAGATGGGCTCGCGTCCGATTAGCCAGTTGGCGGGGTAACGGCCACCAAAGCGACGATCGGTAGCCGGACTG
AGAGGTTGAACGGCCACATTGGGACTGAGACACGGCCAGACTCCTACGGGAGGCAGCAGTGGGGGATATTGCACAA
TGGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGGGAAGACGGTCTTCGGATTGTAAACCTCTGTCTTTGGGGAAGA
AAATGACGGTACCCAAAGAGGAAGCTCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGAGCAAGCGTTG
TCCGGAATTACTGGGTGTAAAGGGAGCGTAGGCGGGATGGCAAGTAGAATGTTAAATCCATCGGCTCAACCGGTGGC
TGCGTTCTAAACTGCCGTTCTTGAGTGAAGTAGAGGCAGGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCAAGTGGCGAAGGCGGCCTGCTGGGCTTTAACTGACGCTGAGGCTCGAAAGCGTGGGGAGCAAACA
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGATTACTAGGTGTGGGGGGACTGACCCCTTCCGTGCCGCAGT
TAACACAATAAGTAATCCACCTGGGGAGTACGGCCGCAAGGTTGAAACTCAAAGGAATTGACGGGGGGCCCGCACAAG
CAGTGGAGTATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAGGTCTTGACATCGGATGCATAGCCTAGAGA
TAGGTGAAGCCCTTCGGGGCATCCAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTGTCGTGAGATGTTGGGTAA
TCCCGCAACGAGCGCAACCCCTTATTATTAGTTGCTACGCAAGAGCACTCTAATGAGACTGCCGTTGACAAAACGGAG
GAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGACCTGGGCTACACACGTACTACAATGGCACTAAAACAGA
GGGCGGCGACACCGCGAGGTGAAGCGAATCCCGAAAAAGTGTCTCAGTTCAGATTGCAGGCTGCAACCCGCGCTGCAT
GAAGTCGGAATTGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCCCGGGCCTTGTACACACCGCCCGTC
ACACCATGGGAGTCGGTAACACCCGAAGCCAGTAGCCTAACCGCAAGGGGGGCGCTGTCTGAAGGTGGGATTGATGAC
TGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID NO:

131 | PROKKA_02671 16S ribosomal RNA gene

TCAAAGAGTTTATCCTGGCTCAGGACGAACGCTGGCGGCGCGCCTAACACATGCAAGTCGAACGGAGCTTACGTTT
TGAAGTTTTTCGGATGGATGAATGTAAGCTTAGTGCGGACGGGTGAGTAACACGTGAGCAACCTGCCTTTCAGAGGG
GGATAACAGCCGGAACGGCTGCTAATACCGCATGATGTTGCGGGGGCACATGCCCTGCAACCAAAGGAGCAATCC
GCTGAAAGATGGGCTCGCGTCCGATTAGCCAGTTGGCGGGGTAACGGCCACCAAAGCGACGATCGGTAGCCGGACT
GAGAGGTTGAACGGCCACATTGGGACTGAGACACGGCCAGACTCCTACGGGAGGCAGCAGTGGGGGATATTGCACA
ATGGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGGGAAGACGGTCTTCGGATTGTAAACCTCTGTCTTTGGGGAAG
AAAATGACGGTACCCAAAGAGGAAGCTCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGAGCAAGCGTT
GTCCGGAATTACTGGGTGTAAAGGGAGCGTAGGCGGGATGGCAAGTAGAATGTTAAATCCATCGGCTCAACCGGTGG
CTGCGTTCTAAACTGCCGTTCTTGAGTGAAGTAGAGGCAGGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAAGTGGCGAAGGCGGCCTGCTGGGCTTTAACTGACGCTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGATTACTAGGTGTGGGGGGACTGACCCCTTCCGTGCCGCAG
TTAACACAATAAGTAATCCACCTGGGGAGTACGGCCGCAAGGTTGAAACTCAAAGGAATTGACGGGGGGCCCGCACA
GCAGTGGAGTATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAGGTCTTGACATCGGATGCATAGCCTAGAG
ATAGGTGAAGCCCTTCGGGGCATCCAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTGTCGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCCTTATTATTAGTTGCTACGCAAGAGCACTCTAATGAGACTGCCGTTGACAAAACGGA
GGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGACCTGGGCTACACACGTACTACAATGGCACTAAAACAG
AGGGCGGCGACACCGCGAGGTGAAGCGAATCCCGAAAAAGTGTCTCAGTTCAGATTGCAGGCTGCAACCCGCGCTGCA
TGAAGTCGGAATTGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCCCGGGCCTTGTACACACCGCCCGT
CACACCATGGGAGTCGGTAACACCCGAAGCCAGTAGCCTAACCGCAAGGGGGGCGCTGTCTGAAGGTGGGATTGATGA
CTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID NO:

132 | PROKKA_00690 16S ribosomal RNA gene

TACGAGAGTTTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTAACACATGCAAGTCGAGCGAAGCGCTGTTTT
CAGAATCTTCGGAGGAAGAGGACAGTGAAGTGAAGCGGCGGACGGGTGAGTAACGCGTGGGCAACCTGCCTCATACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGGACCGCATGGTGTAGTGTGAAAACTCCGGTG
GTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAAAGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGGAATATTGCACAA
TGGGGGAACCCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATTTCCGGTATGTAAACTTCTATCAGCAGGGAAGA
AAATGACGGTACCTGAGTAAGAAGCACCGGCTAAATACGTGCCAGCAGCCGCGGTAATACGTATGGTGCAAGCGTTA
TCCGGATTACTGGGTGTAAAGGGAGCGTAGACGGATAGGCAAGTCTGGAGTGAAAACCCAGGGCTCAACCCCTGGGA
CTGCTTTGGAAACTGCAGATCTGGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAAGTGGCGAAGGCGGCTTACTGGACGGTGAAGTGAAGTGTAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGACTACTAGGTGTGCGGTGTGCAAAGCACATCGGTGCCGCAG
CAAACGCAATAAGTAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAA
GCGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCTGGTCTTGACATCCGGATGACGGGGCGAGTA
ATGTCGCCGTCCTTCGGGGCGTCCGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTGTCGTGAGATGTTGGGTAA
AGTCCCGCAACGAGCGCAACCCCTTATCTTCAGTAGCCAGCATATAAGGTGGGCACTCTGGAGAGACTGCCAGGGAGA
ACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGGCCAGGGCTACACACGTGCTACAATGGCGTA
AACAAAGGGAAGCGAGAGGGTGACCTGGAGCGAATCCCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCG
ACTACATGAAGCTGGAATCGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCCCGGGTCTTGTACACACC
GCCCGTCACACCATGGGAGTCAGTAACGCCCGAAGCCAGTGACCCAACCTTAGAGGAGGGAGCTGTCTGAAGGCGGGA
CGGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 131 | PROKKA_00991 16S ribosomal RNA gene

TACGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAGCGAAGCGCTGTTTT
CAGAATCTTCGGAGGAAGAGGACAGTGAAGTGAAGCGGCGGACGGGTGAGTAACGCGTGGGCAACCTGCCTCATAACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGGACCGCATGGTGTAGTGTGAAAACTCCGGTG
GTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAAAGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCCAACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAA
TGGGGGAAACCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATTTCCGGTATGTAAACTTCTATCAGCAGGGAAGA
AGATGACGGTACCTGAGTAAGAAGCACCGGCTAAATACGTGCCAGCAGCCGCGGTAATACGTATGGTGCAAGCGTTA
TCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGATAGGCAAGTCTGGAGTGAAAACCCAGGGCTCAACCCTGGGA
CTGCTTTGGAAACTGCAGATCTGGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAAGTGGCGAAGGCGGCTTACTGGACGGTGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGACTACTAGGTGTTCGGTGTGCAAAGCACATCGGTGCCGCAG
CAAACGCAATAAGTAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAA
GCGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCTGGTCTTGACATCCGGATGACGGGGCGAGTA
ATGTCGCCGTCCCTTCGGGGCATCCGAGACAGGTGGTGCATGGTTGTCTGTCAGCTCGTGTCTGTGAGATGTTGGGTTA
AGTCCCGCAACGAGCGCAACCCTTATCTTCAGTAGCCAGCATATAAGGTGGGCACTCTGGAGAGACTGCCAGGGAGA
ACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCCTTATGGCCAGGGCTACACACGTGCTACAATGGCGTA
AACAAAGGGAAGCGAGAGGGTGACCTGAAGCGAATCCCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCG
ACTACATGAAGCTGGAATCGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCGCCGGGTCTTGACACACC
GCCCCTCACACCATGGGAGTCAGTAACGCCCCGAAGCCAGTGACCCAACCTTAGAGGAGGGAGCTGTCTGAAGGCGGGA
CGGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 134 | PROKKA_01948 16S ribosomal RNA gene

TACGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAGCGAAGCGCTGTTTT
CAGAATCTTCGGAGGAAGAGGACAGTGAAGTGAAGCGGCGGACGGGTGAGTAACGCGTGGGCAACCTGCCTCATAACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGGACCGCATGGTGTAGTGTGAAAACTCCGGTG
GTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAAAGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCCAACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAA
TGGGGGAAACCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATTTCCGGTATGTAAACTTCTATCAGCAGGGAAGA
AGATGACGGTACCTGAGTAAGAAGCACCGGCTAAATACGTGCCAGCAGCCGCGGTAATACGTATGGTGCAAGCGTTA
TCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGATAGGCAAGTCTGGAGTGAAAACCCAGGGCTCAACCCTGGGA
CTGCTTTGGAAACTGCAGATCTGGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAAGTGGCGAAGGCGGCTTACTGGACGGTGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGACTACTAGGTGTTCGGTGTGCAAAGCACATCGGTGCCGCAG
CAAACGCAATAAGTAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAA
GCGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCTGGTCTTGACATCCGGATGACGGGGCGAGTA
ATGTCGCCGTCCCTTCGGGGCATCCGAGACAGGTGGTGCATGGTTGTCTGTCAGCTCGTGTCTGTGAGATGTTGGGTTA
AGTCCCGCAACGAGCGCAACCCTTATCTTCAGTAGCCAGCATATAAGGTGGGCACTCTGGAGAGACTGCCAGGGAGA
ACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCCTTATGGCCAGGGCTACACACGTGCTACAATGGCGTA
AACAAAGGGAAGCGAGAGGGTGACCTGGAGCGAATCCCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCG
ACTACATGAAGCTGGAATCGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCGCCGGGTCTTGACACACC
GCCCCTCACACCATGGGAGTCAGTAACGCCCCGAAGCCAGTGACCCAACCTTAGAGGAGGGAGCTGTCTGAAGGCGGGA
CGGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 135 | PROKKA_02310 16S ribosomal RNA gene

TACGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAGCGAAGCGCTGTTTT
CAGAATCTTCGGAGGAAGAGGACAGTGAAGTGAAGCGGCGGACGGGTGAGTAACGCGTGGGCAACCTGCCTCATAACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGGACCGCATGGTGTAGTGTGAAAACTCCGGTG
GTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAAAGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCCAACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAA
TGGGGGAAACCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATTTCCGGTATGTAAACTTCTATCAGCAGGGAAGA
AGATGACGGTACCTGAGTAAGAAGCACCGGCTAAATACGTGCCAGCAGCCGCGGTAATACGTATGGTGCAAGCGTTA
TCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGATAGGCAAGTCTGGAGTGAAAACCCAGGGCTCAACCCTGGGA
CTGCTTTGGAAACTGCAGATCTGGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAAGTGGCGAAGGCGGCTTACTGGACGGTGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGACTACTAGGTGTTCGGTGTGCAAAGCACATCGGTGCCGCAG
CAAACGCAATAAGTAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAA
GCGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCTGGTCTTGACATCCGGATGACGGGGCGAGTA
ATGTCGCCGTCCCTTCGGGGCATCCGAGACAGGTGGTGCATGGTTGTCTGTCAGCTCGTGTCTGTGAGATGTTGGGTTA
AGTCCCGCAACGAGCGCAACCCTTATCTTCAGTAGCCAGCATATAAGGTGGGCACTCTGGAGAGACTGCCAGGGAGA
ACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCCTTATGGCCAGGGCTACACACGTGCTACAATGGCGTA
AACAAAGGGAAGCGAGAGGGTGACCTGAAGCGAATCCCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCG
ACTACATGAAGCTGGAATCGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCGCCGGGTCTTGACACACC
GCCCCTCACACCATGGGAGTCAGTAACGCCCCGAAGCCAGTGACCCAACCTTAGAGGAGGGAGCTGTCTGAAGGCGGGA
CGGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 136 | PROKKA_02993 16S ribosomal RNA gene

TACGAGAGTTTGTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAGCGAAGCGCTGTTTT
CAGAATCTTCGGAGGAAGAGGACAGTGAAGTGAAGCGGCGGACGGGTGAGTAACGCGTGGGCAACCTGCCTCATACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGGACCGCATGGTGTAGTGTGAAAACTCCGGTG
GTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAAAGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAA
TGGGGGAAACCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATTTCCGGTATGTAAACTTCTATCAGCAGGGAAGA
AGATGACGGTACCTGAGTAAGAAGCACCGGCTAAATACGTGCCAGCAGCCGCGGTAATACGTATGGTGCAGCGTTA
TCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGATAGGCAAGTCTGGAGTGAAAACCCAGGGCTCAACCCTGGGA
CTGCTTTGGAACTGCAGATCTGGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCACTGGCGAAGGCGGCTTACTGGACGGTGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGACTACTAGGTGTGCGGTGTGCAAAGCACATCGGTGCCGCAG
CAAACGCAATAAGTAGTCCACCTGGGGAGTACGTTCCGAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAA
GCGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCTGGTCTTGACATCCGGATGACGGGGCGAGTA
ATGTCGCCGTCCCTTCGGGGCATCCGAGACAGGTGGTGCATGGTTGTGTCGTCAGCTCGTGTGTCGTGAGATGTTGGGT
AGTCCCGCAACGAGCGCAACCCTTATCTTCAGTAGCCAGCATATAAGGTGGGCACTCTGGAGAGACTGCCAGGGAGA
ACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGGCCAGGGCTACACACGTGCTACAATGGCGTA
AACAAAGGGAAGCGAGAGGGTGACCTGGAGCGAATCCCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCG
ACTACATGAAGCTGGAATCGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCGCCGGGTCTTGACACACC
GCCCCGTACACCATGGGAGTCAGTAACGCCCGAAGCCAGTGACCCAACCTTAGAGGAGGGAGCTGTGGAAGGCGGGA
CGGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 137 | PROKKA_00436 16S ribosomal RNA gene

ATGAGAGTTTGTATCCTAGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCAATTTAACG
GAAGTTTTTCGGATGGAAGTTGAATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTTGACTGGG
GGACAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTATCGCATGATACAGTGTGAAAACTCCGGTG
TACAAGATGGACCCGCGTCTGATTAGCTAGTTGGTAAGGTAACGGCTTACCAAGGCGACGATCAGTAGCCGACCTGA
GAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAAT
GGGCGAAAGCCTGATGCAGCGACGCCGCGTGAAGTAAGAAGTATTTCCGGTATGTAAAGCTCTATCAGCAGGGAAGAA
AATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTAT
CCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGTAAAGCAAGTCTGAAGTGAAAGCCCCGCGGCTCAACTGCGGGAC
TGCTTTGGAACTGTTTAACTGGAGTGTGCGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCACTGGCGAAGGCGACTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAC
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTTGGGGAGCAAAGCTCTTCGGTGCCGTCCG
AAACGCAGTAAGTATTCCACCTGGGGAGTACGTTCCGAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCAGGTCTTGACATCGATCCGACGGGGGAGTAA
CGTCCCCCTTCCCTTCGGGGCGGAGAAGACAGGTGGTGCATGGTTGTGTCGTCAGCTCGTGTGTCGTGAGATGTTGGGT
GTCCCGCAACGAGCGCAACCCTTATTCTAAGTAGCCAGCGGTTCCGGCCGGGAACCTTGGGAGACTGCCAGGGATAA
CCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGATCTGGGCTACACACGTGCTACAATGGCGTAA
ACAAAGAGAAGCAAGACCGCGAGGTGGAGCAAATCTCAAAAATAACGTCTCAGTTCGGACTGCAGGCTGCAACTCGC
CTGCACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTGCGCGGTGAATACGTTCGCCGGGTCTTGACACACCG
CCCGTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCGCAAGGAGGGAGCTGCCGAAGGCGGGAC
GATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 138 | PROKKA_00685 16S ribosomal RNA gene

ATGAGAGTTTGTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCGATTTAACG
GAAATTTTCGGATGGAAGTTGAATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTTGACTGGG
GGACAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTATCGCATGATACAGTGTGAAAACTCCGGTG
TACAAGATGGACCCGCGTCTGATTAGCTAGTTGGTAAGGTAACGGCTTACCAAGGCGACGATCAGTAGCCGACCTGA
GAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAAT
GGGCGAAAGCCTGATGCAGCGACGCCGCGTGAAGTAAGAAGTATTTCCGGTATGTAAAGCTCTATCAGCAGGGAAGAA
AATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTAT
CCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGTAAAGCAAGTCTGAAGTGAAAGCCCCGCGGCTCAACTGCGGGAC
TGCTTTGGAACTGTTTAACTGGAGTGTGCGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCACTGGCGAAGGCGACTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACAC
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTTGGGGAGCAAAGCTCTTCGGTGCCGTCCG
AAACGCAGTAAGTATTCCACCTGGGGAGTACGTTCCGAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCAGGTCTTGACATCGATCCGACGGGGGAGTAA
CGTCCCCCTTCCCTTCGGGGCGGAGAAGACAGGTGGTGCATGGTTGTGTCGTCAGCTCGTGTGTCGTGAGATGTTGGGT
GTCCCGCAACGAGCGCAACCCTTATTCTAAGTAGCCAGCGGTTCCGGCCGGGAACCTTGGGAGACTGCCAGGGATAA
CCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGATCTGGGCTACACACGTGCTACAATGGCGTAA
ACAAAGAGAAGCAAGACCGCGAGGTGGAGCAAATCTCAAAAATAACGTCTCAGTTCGGACTGCAGGCTGCAACTCGC
CTGCACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTGCGCGGTGAATACGTTCGCCGGGTCTTGACACACCG
CCCGTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCGCAAGGAGGGAGCTGCCGAAGGCGGGAC
GATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

NO: 130 | PROKKA_01171 16S ribosomal RNA gene

ATGAGAGTTTGTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCGATTTAACG
GAAGTTTTTCGGATGGAAGTTGAATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTTGTACTGGG
GGACAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTATCGCATGATACAGTGTGAAAAACTCCGGTGG
TACAAGATGGACCCGCGTCTGATTAGCTAGTTGGTAAGGTAACGGCTTACCAAGGCGACGATCAGTAGCCGACCTGA
GAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCTACGGGAGGCAGCAGTGGGGAATATTGCACAAT
GGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGAAGAA
AATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGGCAAGCGTTAT
CCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGTAAAGCAAGTCTGAAGTGAAAGCCCCGCGGCTCAACTGCGGGAC
TGCTTTTGAAACTGTTTAACTGGAGTGTCTGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCAAGTGGCGAAGGCGACTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTTGGGGAGCAAAGCTCTTCGGTGCCGTCTGC
AAACGCAGTAAGTATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAAATTCGAAGCAACGCGAAGAACCTTACCAGGTCTTGACATCGATCCGACGGGGGAGTAA
CGTCCCCCTTCCCTTCGGGGCGGAGAAGACAGGTGGTGCATGGTTGTCTGTCAGCTCGTGTCTGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCCTTATTCTAAGTAGCCAGCGGTTTCGGCCGGGAACCTCTTGGGAGACTGCCAGGGATAA
CCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGATCTGGGCTACACACGTGCTACAATGGCGTAA
ACAAAGAGAAGCAAGACCCGCGAGGTGGAGCAAATCTCAAAAATAACGTCTCAGTTCGGACTGCAGGCTGCAACTCGC
CTGCACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCG
CCCGTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCGCAAGGAGGGAGCTGCCGAAGGCGGGAC
GATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 140 | PROKKA_05969 16S ribosomal RNA gene

ATGAGAGTTTGTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAACGAAGCGATTTAACG
GAAGTTTTTCGGATGGAAGTTGGATTGACTGAGTGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTTGTACTGGG
GGACAACAGTTAGAAATGACTGCTAATACCGCATAAGCGCACAGTATCGCATGATACAGTGTGAAAAACTCCGGTGG
TACAAGATGGACCCGCGTCTGATTAGCTAGTTGGTAAGGTAACGGCTTACCAAGGCGACGATCAGTAGCCGACCTGA
GAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAAACCTCTACGGGAGGCAGCAGTGGGGAATATTGCACAAT
GGGCGAAAGCCTGATGCAGCGACGCCGCGTGAGTGAAGAAGTATTTTCGGTATGTAAAGCTCTATCAGCAGGGAAGAA
AATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGGCAAGCGTTAT
CCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGTAAAGCAAGTCTGAAGTGAAAGCCCCGCGGCTCAACTGCGGGAC
TGCTTTTGAAACTGTTTAACTGGAGTGTCTGGAGAGGTAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCAAGTGGCGAAGGCGACTTACTGGACGATAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTTGGGGAGCAAAGCTCTTCGGTGCCGTCTGC
AAACGCAGTAAGTATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAAATTCGAAGCAACGCGAAGAACCTTACCAGGTCTTGACATCGATCCGACGGGGGAGTAA
CGTCCCCCTTCCCTTCGGGGCGGAGAAGACAGGTGGTGCATGGTTGTCTGTCAGCTCGTGTCTGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCCTTATTCTAAGTAGCCAGCGGTTTCGGCCGGGAACCTCTTGGGAGACTGCCAGGGATAA
CCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGATCTGGGCTACACACGTGCTACAATGGCGTAA
ACAAAGAGAAGCAAGACCCGCGAGGTGGAGCAAATCTCAAAAATAACGTCTCAGTTCGGACTGCAGGCTGCAACTCGC
CTGCACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCTCGCGGTGAATACGTTCCCGGGTCTTGTACACACCG
CCCGTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCGCAAGGAGGGAGCTGCCGAAGGCGGGAC
GATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 141 | PROKKA_00279 16S ribosomal RNA gene

ATCAGAGAGTTTGTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCCTAACACATGCAAGTCGAGCGAAGCACTTAAG
TGGATCTCTTCGGATTGAAACTTATTTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATAACG
GGGGATAACAGTTAGAAATGGCTGCTAATACCGCATAAGCGCACAGGACCGCATGGTCTGGTGTGAAAAACTCCGGT
GGTATGAGATGGACCCGCGTCTGATTAGCTAGTTGGAGGGGTAACGGCCCACCAAGGCGACGATCAGTAGCCGGCCT
GAGAGGGTGAACGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACA
ATGGGGGAAACCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATCTCGGTATGTAAACTTCTATCAGCAGGGAAG
AAAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGGCAAGCGTT
ATCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGAAGAGCAAGTCTGATGTGAAAGGCTGGGGCTTAACCCCAAG
ACTGCATTGGAAACTGTTTTTCTAGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATA
TTAGGAGGAACACCAAGTGGCGAAGGCGGCTTACTGGACGGTAACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAAC
AGGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTCTGGGTGGCAAAGCCATTTCGGTGCCGCA
GCAAACGCAATAAGTATTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACA
AGCGGTGGAGCATGTGGTTTAAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCCTCTGACCGGCCCGT
AACGGGGCCTTCCCTTCGGGGCAGAGGAGACAGGTGGTGCATGGTTGTCTGTCAGCTCGTGTCTGTGAGATGTTGGGT
AAGTCCCGCAACGAGCGCAACCCCTATCCTTAGTAGCCAGCAGGTGAAGCTGGGCACTCTAGGGAGACTGCCGGGGA
TAACCCGGAGGAAGGCGGGGACGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCG
TAAACAAAGGGAAGCGAGACAGCGATGTTGAGCAAATCCCAAAAATAACGTCCCAGTTCGGACTGCAGTCTGCAACT
CGACTGCACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCTCGCGGTGAATACGTTCCCGGGTCTTGTACACA
CCGCCCGTCACACCATGGGAGTCAGTAACGCCCGAAGTCAGTGACCCAACCTTTTAGGAGGGAGCTGCCGAAGGCGG
GACCGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 142 | PROKKA_01221 16S ribosomal RNA gene

ATCAGAGAGTTTGGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTTAAG
TGGATCTCTTCGGATTGAACTTATTTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATAACAG
GGGGATAACAGTTAGAAATGGCTGCTAATACCGCATAAGCGCACAGGACCGCATGGTCTGGTGTGAAAAACTCCGGT
GGTATGAGATGGACCCGCGTCTGATTAGCTAGTTGGAGGGGTAACGGCCCACCAAGGCGACGATCAGTAGCCGGCCT
GAGAGGGTGAACGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACA
ATGGGGGAAACCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATCTCGGTATGTAACTTCTATCAGCAGGGAAG
AAAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTT
ATCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGAAGAGCAAGTCTGATGTGAAAGGCTGGGGCTTAACCCCAAGG
ACTGCATTGGAACTGTTTTTCTAGAGTGCCGGAGAGGTAAGTGGAAATTCCTAGTGTAGCGGTGAAATGCGTAGATA
TTAGGAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGGTAAGTACGTTGAGGCTCGAAAGCGTGGGGAGCAAAC
AGGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTCGGGTGGCAAAGCCATTTCGGTGCCGCA
GCAAACGCAATAAGTATTCCACCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCCGCACA
AGCGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCCTCTGACCGGCCCGT
AACGGGGCCTTCCCTTCGGGGCAGAGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTT
AAGTCCCGCAACGAGCGCAACCCCTATCCTTAGTAGCCAGCAGGTGAAGCTGGGCACTCTAGGGAGACTGCCGGGGA
TAACCCGGAGGAAGGCGGGGACGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCG
TAAACAAAGGGAAGCGGGACAGCGATGTTGAGCAAATCCCAAAAATAACGTCCCAGTTCGGACTGCAGTCTGCAACT
CGACTGCACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGACACA
CCGCCCCGTACACCATGGGAGTCAGTAACGCCCCGAAGTCAGTGACCCAACCTTACAGGAGGGAGCTGCCGAAGGCGG
GACCGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 143 | PROKKA_02318 16S ribosomal RNA gene

ATCAGAGAGTTTGGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTTAAG
TGGATCTCTTCGGATTGAAGCTTATTTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATAACAG
GGGGATAACAGTTAGAAATGGCTGCTAATACCGCATAAGCGCACAGGACCGCATGGTCTGGTGTGAAAAACTCCGGT
GGTATGAGATGGACCCGCGTCTGATTAGCTAGTTGGAGGGGTAACGGCCCACCAAGGCGACGATCAGTAGCCGGCCT
GAGAGGGTGAACGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACA
ATGGGGGAAACCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATCTCGGTATGTAACTTCTATCAGCAGGGAAG
AAAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTT
ATCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGAAGAGCAAGTCTGATGTGAAAGGCTGGGGCTTAACCCCAAGG
ACTGCATTGGAACTGTTTTTCTAGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATA
TTAGGAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGGTAAGTACGTTGAGGCTCGAAAGCGTGGGGAGCAAAC
AGGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTCGGGTGGCAAAGCCATTTCGGTGCCGCA
GCAAACGCAATAAGTATTCCACCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCCGCACA
AGCGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCCTCTGACCGGCCCGT
AACGGGGCCTTCCCTTCGGGGCAGAGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTT
AAGTCCCGCAACGAGCGCAACCCCTATCCTTAGTAGCCAGCAGGTGAAGCTGGGCACTCTAGGGAGACTGCCGGGGA
TAACCCGGAGGAAGGCGGGGACGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCG
TAAACAAAGGGAAGCGAGACAGCGATGTTGAGCAAATCCCAAAAATAACGTCCCAGTTCGGACTGCAGTCTGCAACT
CGACTGCACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGACACA
CCGCCCCGTACACCATGGGAGTCAGTAACGCCCCGAAGTCAGTGACCCAACCTTACAGGAGGGAGCTGCCGAAGGCGG
GACCGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 144 | PROKKA_02336 16S ribosomal RNA gene

ATCAGAGAGTTTGGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTTAAG
CGGATCTCTTCGGATTGAACTTATTTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATAACAG
GGGGATAACAGTTAGAAATGGCTGCTAATACCGCATAAGCGCACAGGACCGCATGGTCTGGTGTGAAAAACTCCGGT
GGTATGAGATGGACCCGCGTCTGATTAGCTAGTTGGAGGGGTAACGGCCCACCAAGGCGACGATCAGTAGCCGGCCT
GAGAGGGTGAACGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACA
ATGGGGGAAACCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATCTCGGTATGTAACTTCTATCAGCAGGGAAG
AAAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTT
ATCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGAAGAGCAAGTCTGATGTGAAAGGCTGGGGCTTAACCCCAAGG
ACTGCATTGGAACTGTTTTTCTAGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATA
TTAGGAGGAACACCAGTGGCGAAGGCGGCTTACTGGACGGTAAGTACGTTGAGGCTCGAAAGCGTGGGGAGCAAAC
AGGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTCGGGTGGCAAAGCCATTTCGGTGCCGCA
GCAAACGCAATAAGTATTCCACCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCCGCACA
AGCGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCCTCTGACCGGCCCGT
AACGGGGCCTTCCCTTCGGGGCAGAGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTT
AAGTCCCGCAACGAGCGCAACCCCTATCCTTAGTAGCCAGCAGGTGAAGCTGGGCACTCTAGGGAGACTGCCGGGGA
TAACCCGGAGGAAGGCGGGGACGACGTCAAATCATCATGCCCTTATGATTTGGGCTACACACGTGCTACAATGGCG
TAAACAAAGGGAAGCGAGACAGCGATGTTGAGCAAATCCCAAAAATAACGTCCCAGTTCGGACTGCAGTCTGCAACT
CGACTGCACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGACACA
CCGCCCCGTACACCATGGGAGTCAGTAACGCCCCGAAGTCAGTGACCCAACCTTACAGGAGGGAGCTGCCGAAGGCGG
GACCGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 145 | PROKKA_04947 16S ribosomal RNA gene

ATCAGAGAGTTTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTTAAG
TGGATCTCTTCGGATTGAACTTATTTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATAACAG
GGGATAACAGTTAGAAATGGCTGCTAATACCGCATAAGCGCACAGGACCGCATGGTCTGGTGTGAAAAACTCCGGT
GGTATGAGATGGACCCGCGTCTGATTAGCTAGTTGGAGGGGTAACGGCCCACCAAGGCGACGATCAGTAGCCGGCCT
GAGAGGGTGAACGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACA
ATGGGGGAAACCCTGATGCAGCGACGCCGCGTGAAGGAAGAAGTATCTCGGTATGTAACTTCTATCAGCAGGGAAG
AAAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTT
ATCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGAAGAGCAAGTCTGATGTGAAAGGCTGGGGCTTAACCCCAAGG
ACTGCATTGGAACTGTTTTTCTAGAGTGCCGGAGAGGTAAGCGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATA
TTAGGAGGAACATCAGTGGCGAAGGCGGCTTACTGGACGGTAAGTACGTTGAGGCTCGAAAGCGTGGGGAGCAAAC
AGGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGAATACTAGGTGTCGGGTGGCAAAGCCATTTCGGTGCCGCA
GCAAACGCAATAAGTATTCCACCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCCGCACA
AGCGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAAGTCTTGACATCCCTCTGACCGGCCCGT
AACGGGGCCCTTCCCTTCGGGGCAGAGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTT
AAGTCCCGCAACGAGCGCAACCCCTATCCTTAGTAGCCAGCAGGTGAAGCTGGGCACTCTAGGGAGACTGCCGGGGA
TAACCCCGAGGAAGGCGGGGACGACGTCAAATCATCATGCCCCCTTATGATTTGGGCTACACACGTGCTACAATGGCG
TAAACAAAGGGAAGCGAGACAGCGATGTTGAGCAAATCCCAAAAATAACGTCCTAGTTCGGACTGCAGTCTGCAACT
CGACTGCACGAAGCTGGAATCGCTAGTAATCGCGAATCAGAATGTCGCGGTGAATACGTTCCCGGGTCTTGACACA
CCGCCCCGTACACCATGGGAGTCAGTAACGCCCCGAAGTCAGTGACCCAACCTTACAGGAGGGAGCTGCCGAAGGCGG
GACCGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 146 | PROKKA_00208 16S ribosomal RNA gene

AACGAGAGTTTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTTAAGT
TTGATTCTTCGGATGAAGACTTTTGTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATAACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGACCACGGTACCGCATGGTACAGTGGTAAAAACTCCGGTG
GTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAACGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAA
TGGAGGAAACTCTGATGCAGCGACGCCGCGTGAAGGATGAAGTATTTTCGGTATGTAACTTCTATCAGCAGGGAAGA
AAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTA
TCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCACGGCAAGCCAGATGTGAAAGCCCCGGGGCTCAACCCCGGGA
CTGCATTTGGAAGTCTGCTGAGCTAGAGTGTTCGGAGAGGCAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAGTGGCGAAGGCGGCTTGCTGGACGATGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGACTGCTAGGTGTCGGGTGGCAAAGCCATTTCGGTGCCGCAG
CTAACGCAATAAGCAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCCGCACAA
GCGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCTGATCTTGACATCCCGATGACCGCTTCGTA
ATGGAAGTTTTTCTTCGGAACATCGGTGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTTA
AGTCCCGCAACGAGCGCAACCCCTATCTTCAGTAGCCAGCAGGTAAAGCTGGGCACTCTGGAGAGACTGCCAGGGAT
AACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCCCTTATGACCAGGGCTACACACGTGCTACAATGGCGT
AAACAAAGAGAAGCGAACTCGCGAGGGTAAGCAAATCTCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTC
GACTACATGAAGCTGGAATCGCTAGTAATCGCAGATCAGAATGCTGCGGTGAATACGTTCCCGGGTCTTGACACAC
CGCCCGTCACACCATGGGAGTCAGTAACGCCCCGAAGTCAGTGACCCAACCGTAAGGAGGGAGCTGCCGAAGGTGGGA
CCGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 147 | PROKKA_00340 16S ribosomal RNA gene

AACGAGAGTTTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTTTGGA
AAGATTCTTCGGATGATTTCCTTTGTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATAACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGACCACGGTACCGCATGGTACAGTGGTAAAAACTCCGGTG
GTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAACGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAA
TGGAGGAAACTCTGATGCAGCGACGCCGCGTGAAGGATGAAGTATTTTCGGTATGTAACTTCTATCAGCAGGGAAGA
AAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTA
TCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCACGGCAAGCCAGATGTGAAAGCCCCGGGGCTCAACCCCGGAC
TGCATTTGGAAGTCTGCTGAGCTAGAGTGTTCGGAGAGGCAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCAGTGGCGAAGGCGGCTTGCTGGACGATGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GATTAGATACCCTGGTAGTCCACGCCGTAAACGATGACTGCTAGGTGTCGGGTGGCAAAGCCATTTCGGTGCCGCAGC
TAACGCAATAAGCAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCTGATCTTGACATCCCGATGACCGCTTCGTAA
TGGAAGCTTTTCTTCGGAACATCGGTGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCCTATCTTCAGTAGCCAGCAGGTAAAGCTGGGCACTCTGGAGAGACTGCCAGGGATA
ACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCCCTTATGACCAGGGCTACACACGTGCTACAATGGCGTA
AACAAAGAGAAGCGAACTCGCGAGGGTAAGCAAATCTCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCG
ACTACATGAAGCTGGAATCGCTAGTAATCGCAGATCAGAATGCTGCGGTGAATACGTTCCCGGGTCTTGACACACC
GCCCGTCACACCATGGGAGTCAGTAACGCCCCGAAGTCAGTGACCCAACCGTAAGGAGGGAGCTGCCGAAGGTGGGAC
CGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID

NO: 148 | PROKKA_01031 16S ribosomal RNA gene

AACGAGAGTTTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTTAAGT
TTGATTCTTCGGATGAAGACTTTTGTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATAACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGACCACGGTACCGCATGGTACAGTGGTAAAAACTCCGGTG
GTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAACGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAA
TGGAGGAAACTCTGATGCAGCGACGCCGCGTGAAGGATGAAGTATTTTCGGTATGTAAACTTCTATCAGCAGGGAAGA
AAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTA
TCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCACGGCAAGCCAGATGTGAAAGCCCCGGGGCTCAACCCCGGGA
CTGCATTTGGAAGTCTGCTGAGCTAGAGTGTTCGGAGAGGCAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAAGTGGCGAAGGCGGCTTGCTGGACGATGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATAACCTGGTAGTCCACGCCGTAAACGATGACTGCTAGGTGTTCGGGTGGCAAAGCCATTTCGGTGCCGCAG
CTAACGCAATAAGCAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAA
GCGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCTGATCTTGACATCCCGATGACCGCTTCGTA
ATGGAAGCTTTTCTTCGGAACATCGGTGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTTCGTGAGATGTTGGGTTA
AGTCCCGCAACGAGCGCAACCCCTATCTTCAGTAGCCAGCAGGTAAAGCTGGGCACTCTGGAGAGACTGCCAGGGAT
AACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGACCAGGGGTACACACGTGCTACAATGGCGT
AAACAAAGAGAAGCGAACTCGCGAGGGTAAGCAAATCTCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTC
GACTACATGAAGCTGGAATCGCTAGTAATCGCAGATCAGAATGCTGCGGTGAATACGTTCCCGGGTCTTGTACACAC
CGCCCGTCACACCATGGGAGTCAGTAACGCCCCGAAGTCAGTGACCCAACCGTAAGGAGGGGAGCTGCCGAAGGTGGGA
CCGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 149 | PROKKA_01840 16S ribosomal RNA gene

AACGAGAGTTTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCACTTTGGA
AAGATTCTTCGGATGATTTCCTTTGTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATAACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGACCACGGTACCGCATGGTACAGTGGTAAAAACTCCGGTG
GTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAACGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAA
TGGAGGAAACTCTGATGCAGCGACGCCGCGTGAAGGATGAAGTATTTTCGGTATGTAAACTTCTATCAGCAGGGAAGA
AAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTA
TCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCACGGCAAGCCAGATGTGAAAGCCCCGGGGCTCAACCCCGGGA
CTGCATTTGGAAGTCTGCTGAGCTAGAGTGTTCGGAGAGGCAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAAGTGGCGAAGGCGGCTTGCTGGACGATGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATAACCTGGTAGTCCACGCCGTAAACGATGACTGCTAGGTGTTCGGGTGGCAAAGCCATTTCGGTGCCGCAG
CTAACGCAATAAGCAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAA
GCGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCTGATCTTGACATCCCGATGACCGCTTCGTA
ATGGAAGCTTTTCTTCGGAACATCGGTGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTTCGTGAGATGTTGGGTTA
AGTCCCGCAACGAGCGCAACCCCTATCTTCAGTAGCCAGCAGGTAAAGCTGGGCACTCTGGAGAGACTGCCAGGGAT
AACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGACCAGGGGTACACACGTGCTACAATGGCGT
AAACAAAGAGAAGCGAACTCGCGAGGGTAAGCAAATCTCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTC
GACTACATGAAGCTGGAATCGCTAGTAATCGCAGATCAGAATGCTGCGGTGAATACGTTCCCGGGTCTTGTACACAC
CGCCCGTCACACCATGGGAGTCAGTAACGCCCCGAAGTCAGTGACCCAACCGTAAGGAGGGGAGCTGCCGAAGGTGGGA
CCGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 150 | PROKKA_02944 16S ribosomal RNA gene

AACGAGAGTTTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCGCTTAAGT
TTGATTCTTCGGATGAAGACTTTTGTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATAACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGACCACGGTACCGCATGGTACAGTGGTAAAAACTCCGGTG
GTATGAGATGGACCCGCGTCTGATTAGGTAGTTGGTGGGGTAACGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAA
TGGAGGAAACTCTGATGCAGCGACGCCGCGTGAAGGATGAAGTATTTTCGGTATGTAAACTTCTATCAGCAGGGAAGA
AAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTA
TCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCACGGCAAGCCAGATGTGAAAGCCCCGGGGCTCAACCCCGGGAC
TGCATTTGGAAGTCTGCTGAGCTAGAGTGTTCGGAGAGGCAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATATT
AGGAGGAACACCAAGTGGCGAAGGCGGCTTGCTGGACGATGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GATTAGATAACCTGGTAGTCCACGCCGTAAACGATGACTGCTAGGTGTTCGGGTGGCAAAGCCATTTCGGTGCCGCAGC
TAACGCAATAAGCAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAAG
CGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCTGATCTTGACATCCCGATGACCGCTTCGTAA
TGGAAGTTTTTCTTCGGAACATCGGTGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTTCGTGAGATGTTGGGTAA
GTCCCGCAACGAGCGCAACCCCTATCTTCAGTAGCCAGCAGGTAAAGCTGGGCACTCTGGAGAGACTGCCAGGGATA
ACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGACCAGGGGTACACACGTGCTACAATGGCGTA
AACAAAGAGAGGCAAACCTCGCGAGGGTAAGCAAATCTCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTCG
ACTACATGAAGCTGGAATCGCTAGTAATCGCAGATCAGAATGCTGCGGTGAATACGTTCCCGGGTCTTGTACACACC
GCCCGTCACACCATGGGAGTCAGTAACGCCCCGAAGTCAGTGACCCAACCGTAAGGAGGGGAGCTGCCGAAGGTGGGAC
CGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 151 | PROKKA_04036 16S ribosomal RNA gene

AACGAGAGTTTATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAGCGAAGCGCTTTGGG
AAGATTCTTCGGATGATTTCCTTTGTGACTGAGCGGCGGACGGGTGAGTAACGCGTGGGTAACCTGCCTCATACAGG
GGGATAACAGTTAGAAATGACTGCTAATACCGCATAAGACCACGGTACCGCATGGTACAGTGGTAAAACTCCGGTG
GTATGAGATGGACCCGCGTTTGATTAGGTAGTTGGTGGGGTAACGGCCTACCAAGCCGACGATCAGTAGCCGACCTG
AGAGGGTGACCGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGCACAA
TGGAGGAAACTCTGATGCAGCGACGCCGCGTGAAGGATGAAGTATTTTCGGTATGTAAACTTCTATCAGCAGGGAAGA
AAATGACGGTACCTGACTAAGAAGCCCCGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGGGGCAAGCGTTA
TCCGGATTTACTGGGTGTAAAGGGAGCGTAGACGGCACGGCAAGCCAGATGTGAAAGCCCCGGGGCTCAACCCCGGGA
CTGCATTTGGAAGTCTGAGCTAGAGTGTTCGGAGAGGCAAGTGGAATTCCTAGTGTAGCGGTGAAATGCGTAGATAT
TAGGAGGAACACCAAGTGGCGAAGGCGGCTTGCTGGACGATGACTGACGTTGAGGCTCGAAAGCGTGGGGAGCAAACA
GGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGACTGCTAGGTGTTCGGGTGGCAAAGCCATTCGGTGCCGCAG
CTAACGCAATAAGCAGTCCACCTGGGGAGTACGTTTCGCAAGAATGAAACTCAAAGGAATTGACGGGGACCCGCACAA
GCGGTGGAGCATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCTGATCTTGACATCCCGATGACTGCTTCGTA
ATGGAAGTTTTTCTTCGGAACATCGGTGACAGGTGGTGCATGGTTGTCTGTCAGCTCGTGTCTGTGAGATGTTGGGTTA
AGTCCCGCAACGAGCGCAACCCCTATCTTCAGTAGCCAGCAGGTAAAGCTGGGCACTCTGGAGAGACTGCCAGGGAT
AACCTGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGACCAGGGCTACACACGTGCTACAATGGCGT
AAACAAAGAGAAGCGAACTCGCGAGGGTAAGCAAATCTCAAAAATAACGTCTCAGTTCGGATTGTAGTCTGCAACTC
GACTACATGAAGCTGGAATCGCTAGTAATCGCAGATCAGAATGCTGCGGTGAATACGTTCCCGGGTCTTGTACACAC
CGCCCGTCACACCATGGGAGTCAGTAACGCCCCGAAGTCAGTGACCCAACCGTAAGGAGGGAGCTGCCGAAGGTGGGA
CCGATAACTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 152 | PROKKA_00437 16S ribosomal RNA gene

ATGGAGAGTTTATCCTGGCTCAGGATGAACGCTGGCGGCATGCCTAATACATGCAAGTCGAACGAAGTTTTCGAGGA
AGCTTGCTTCCAAAGAGACTTAGTGCGCAACGGGTGAGTAACACGTAGGTAACTGCCCATGTGTCCGGGATAACTG
CTGGAACCGGTAGCTAAAACCGGATAGGTATACAGAGCGCATGCTCAGTATATTAAAGCGCCCATCAAGGCGTGAAC
ATGGATGGACCTGCGGCGCATTAGCTAGTTGGTGAGGTAAACGGCTACCAAGGCGATGATGCGTAGCCGGCCTGAGA
GGGTAAACGGGCCACATTGGGACTGAGACACGGCCCCAACTCCTACGGGAGGCAGCAGTAGGGAATTTTCGTCAATGG
GGGAAACCTGAACGAGCAATGCCGCGTGAGTGAAGAAGGTCTTCGGATCGTAAAGCTCTGTTGTAAGTGAAGAACC
GCTCATAGAGGAAATGCTATGGGAGTGACGGTAGCTTACCAGAAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGT
AATACGTAGGTGGCAAGCGTTATCCGGAATCATTGGGCGTAAAGGGTGCGTAGGTGGCGTACTAAGTCTGTAGTAAA
AGGCAATGGCTCAACCATTTGTAAGCTATGGAAGTGGTATGCTGGAGTGCAGAAGAGGGCGATGGAATTCATGTGT
AGCGGTAAAATGCGTAGATATATGGAGGAACACCAAGTGGCGAAGGCGGTTCGCTGTAAGTGAAGTGAAGGCA
CGAAAGCGTGGGGAGCAAATAGGATTAGATACCCTAGTAGTCCACGCCGTAAACGATGAGAACTAAGTGTGGAGGA
ATTCAGTGCTGCAGTTAACGCAATAAGTTCTCCGCTGGGGAGTATGCACGCAAGTGTGAAACTCAAAGGAATTGAC
GGGGCCCCGCACAAGCGGTGGAGTATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCAGGCCTTGACATGGAA
ACAAATACCCTAGAGATAGGGGGATAATTATGGATCACACAGGTGGTGCATGGTTGTCTGTCAGCTCGTGTCTGTGAGA
TGTTGGGTAAAGTCCCGCAACGAGCGCAACCCCTGTCTGCATGTTACCAGCATCAAGTTGGGGACTCATGCGAGACTG
CCGGTGACAAACCGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGGCTGGGCTACACACGTACTAC
AATGGCGGCCACAAAGAGCAGCGACACAGTGATGTGAAGCGAATCTCATAAAGGTCTGCTCAGTTCGGATTGAAGTC
TGCAACTCGACTTCATGAAGTCGGAATCGCTAGTAATCGCAGATCAGCATGCTGCGGTGAATACGTTCTCGGGCCTT
GTACACACCGCCCGTCAAACCATGGGAGTCAGTAATACCCGAAGCCGGTGGCATAACCGTAAGGAGTGAGCCGTCTGA
AGGTAGGACCGATGACTGGGGTTAAGTCGTAACAAGGTATCCCTACGGGAACGTGGGGATGGATCACCTCCTTT

>SEQ ID NO: 153 | PROKKA_00896 16S ribosomal RNA gene

ATGGAGAGTTTATCCTGGCTCAGGATGAACGCTGGCGGCATGCCTAATACATGCAAGTCGAACGAAGTTTTCGAGGA
AGCTTGCTTCCAAAGAGACTTAGTGCGCAACGGGTGAGTAACACGTAGGTAACTGCCCATGTGTCCGGGATAACTG
CTGGAACCGGTAGCTAAAACCGGATAGGTATACAGAGCGCATGCTCAGTATATTAAAGCGCCCATCAAGGCGTGAAC
ATGGATGGACCTGCGGCGCATTAGCTAGTTGGTGAGGTAAACGGCCACCAAGGCGATGATGCGTAGCCGGCCTGAGA
GGGTAAACGGGCCACATTGGGACTGAGACACGGCCCCAACTCCTACGGGAGGCAGCAGTAGGGAATTTTCGTCAATGG
GGGAAACCTGAACGAGCAATGCCGCGTGAGTGAAGAAGGTCTTCGGATCGTAAAGCTCTGTTGTAAGTGAAGAACC
GCTCATAGAGGAAATGCTATGGGAGTGACGGTAGCTTACCAGAAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGT
AATACGTAGGTGGCAAGCGTTATCCGGAATCATTGGGCGTAAAGGGTGCGTAGGTGGCGTACTAAGTCTGTAGTAAA
AGGCAATGGCTCAACCATTTGTAAGCTATGGAAGTGGTATGCTGGAGTGCAGAAGAGGGCGATGGAATTCATGTGT
AGCGGTAAAATGCGTAGATATATGGAGGAACACCAAGTGGCGAAGGCGGTTCGCTGTAAGTGAAGTGAAGGCA
CGAAAGCGTGGGGAGCAAATAGGATTAGATACCCTAGTAGTCCACGCCGTAAACGATGAGAACTAAGTGTGGAGGA
ATTCAGTGCTGCAGTTAACGCAATAAGTTCTCCGCTGGGGAGTATGCACGCAAGTGTGAAACTCAAAGGAATTGAC
GGGGCCCCGCACAAGCGGTGGAGTATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCAGGCCTTGACATGGAA
ACAAATACCCTAGAGATAGGGGGATAATTATGGATCACACAGGTGGTGCATGGTTGTCTGTCAGCTCGTGTCTGTGAGA
TGTTGGGTAAAGTCCCGCAACGAGCGCAACCCCTGTCTGCATGTTACCAGCATCAAGTTGGGGACTCATGCGAGACTG
CCGGTGACAAACCGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGGCTGGGCTACACACGTACTAC
AATGGCGGCCACAAAGAGCAGCGACACAGTGATGTGAAGCGAATCTCATAAAGGTCTGCTCAGTTCGGATTGAAGTC
TGCAACTCGACTTCATGAAGTCGGAATCGCTAGTAATCGCAGATCAGCATGCTGCGGTGAATACGTTCTCGGGCCTT
GTACACACCGCCCGTCAAACCATGGGAGTCAGTAATACCCGAAGCCGGTGGCATAACCGTAAGGAGTGAGCCGTCTGA
AGGTAGGACCGATGACTGGGGTTAAGTCGTAACAAGGTATCCCTACGGGAACGTGGGGATGGATCACCTCCTTT

>SEQ ID NO: 154 | PROKKA_02845 16S ribosomal RNA gene

ATGGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCATGCCTAATACATGCAAGTCGAACGAAGTTTCGAGGA
AGCTTGCTTCCAAAGAGACTTAGTGCGGAACGGGTGAGTAACACGTAGGTAACCTGCCCATGTGTCCGGGATAACTG
CTGGAACCGGTAGCTAAAACCGGATAGGTATACAGAGCGCATGCTCAGTATATTAAAGCGCCCATCAAGGCGTGAAC
ATGGATGGACCTGCGGCGCATTAGCTAGTTGGTGAGGTAACGGCTCACCAAGGCGATGATGCGTAGCCGGCCTGAGA
GGGTAAACGGCCACATTGGGACTGAGACACGGCCCCAACTCCTACGGGAGGCAGCAGTAGGGAATTTTCGTCAATGG
GGGAAACCTTGAACGAGCAATGCCGCGTGAGTGAAGAAGGTCTTCGGATCGTAAAGCTCTGTTGTAAGTGAAGAACC
GCTCATAGAGGAAATGCTATGGGAGTGACGGTAGCTTACCAGAAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGT
AATACGTAGGTGGCAAGCGTTATCCGGAATCATTGGGCGTAAAGGGTGCGTAGGTGGCGTACTAAGTCTGTAGTAAA
AGGCAATGGCTCAACCATTTGTAAGCTATGGAACCTGGTATGCTGGAGTGCAGAAGAGGGCGATGGAATTCCATGTGT
AGCGGTAAAATGCGTAGATATATGGAGGAACACCAAGTGGCGAAGGCGGTTCGCCTGGTCTGTAAGTACACTGAGGCA
CGAAAGCGTGGGGAGCAAATAGGATTAGATACCCTAGTAGTCCACGCCGTAAACGATGAGAACTAAGTGTGGAGGA
ATTCAGTGCTGCAGTTAACGCAATAAGTTCTCCGCTGGGGAGTATGCACGCAAGTGTGAAACTCAAAGGAATTGAC
GGGGGCCCCGCACAAGCGGTGGAGTATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCAGGCCTTGACATGGAA
ACAAATACCCTAGAGATAGGGGGATAATTATGGATCACACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGA
TGTTGGGTAAAGTCCCGCAACGAGCGCAACCCTTGTTCGCATGTTACCAGCATCAAGTTGGGGACTCATGCGAGACTG
CCGGTGACAAACCGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGGCCTGGGCTACACACGTACTAC
AATGGCGGCCACAAAGAGCAGCGACACAGTGATGTGAAGCGAATCTCATAAAGGTCGTCTCAGTTCGGATTGAAGTC
TGCAACTCGACTTCATGAAGTCGGAATCGCTAGTAATCGCAGATCAGCATGCTGCGGTGAATACGTTCTCGGGCCTT
GTACACACCGCCCCGTCAAACCATGGGAGTCAGTAATACCCGAAGCCGGTGGCATAACCGTAAGGAGTGAGCCGTCTGA
AGGTAGGACCGATGACTGGGGTTAAGTCGTAACAAGGTATCCCTACGGGAACGTGGGGATGGATCACCTCCTTT

>SEQ ID NO: 155 | PROKKA_04164 16S ribosomal RNA gene

ATGGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCATGCCTAATACATGCAAGTCGAACGAAGTTTCGAGGA
AGCTTGCTTCCAAAGAGACTTAGTGCGGAACGGGTGAGTAACACGTAGGTAACCTGCCCATGTGTCCGGGATAACTG
CTGGAACCGGTAGCTAAAACCGGATAGGTATACAGAGCGCATGCTCAGTATATTAAAGCGCCCATCAAGGCGTGAAC
ATGGATGGACCTGCGGCGCATTAGCTAGTTGGTGAGGTAACGGCCCCACCAAGGCGATGATGCGTAGCCGGCCTGAGA
GGGTAAACGGCCACATTGGGACTGAGACACGGCCCCAACTCCTACGGGAGGCAGCAGTAGGGAATTTTCGTCAATGG
GGGAAACCTTGAACGAGCAATGCCGCGTGAGTGAAGAAGGTCTTCGGATCGTAAAGCTCTGTTGTAAGTGAAGAACC
GCTCATAGAGGAAATGCTATGGGAGTGACGGTAGCTTACCAGAAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGT
AATACGTAGGTGGCAAGCGTTATCCGGAATCATTGGGCGTAAAGGGTGCGTAGGTGGCGTACTAAGTCTGTAGTAAA
AGGCAATGGCTCAACCATTTGTAAGCTATGGAACCTGGTATGCTGGAGTGCAGAAGAGGGCGATGGAATTCCATGTGT
AGCGGTAAAATGCGTAGATATATGGAGGAACACCAAGTGGCGAAGGCGGTTCGCCTGGTCTGTAAGTACACTGAGGCA
CGAAAGCGTGGGGAGCAAATAGGATTAGATACCCTAGTAGTCCACGCCGTAAACGATGAGAACTAAGTGTGGAGGA
ATTCAGTGCTGCAGTTAACGCAATAAGTTCTCCGCTGGGGAGTATGCACGCAAGTGTGAAACTCAAAGGAATTGAC
GGGGGCCCCGCACAAGCGGTGGAGTATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCAGGCCTTGACATGGAA
ACAAATACCCTAGAGATAGGGGGATAATTATGGATCACACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGA
TGTTGGGTAAAGTCCCGCAACGAGCGCAACCCTTGTTCGCATGTTACCAGCATCAAGTTGGGGACTCATGCGAGACTG
CCGGTGACAAACCGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGGCCTGGGCTACACACGTACTAC
AATGGCGACCACAAAGAGCAGCGACACAGTGATGTGAAGCGAATCTCATAAAGGTCGTCTCAGTTCGGATTGAAGTC
TGCAACTCGACTTCATGAAGTCGGAATCGCTAGTAATCGCAGATCAGCATGCTGCGGTGAATACGTTCTCGGGCCTT
GTACACACCGCCCCGTCAAACCATGGGAGTCAGTAATACCCGAAGCCGGTGGCATAACCGTAAGGAGTGAGCCGTCTGA
AGGTAGGACCGATGACTGGGGTTAAGTCGTAACAAGGTATCCCTACGGGAACGTGGGGATGGATCACCTCCTTT

>SEQ ID NO: 156 | PROKKA_04921 16S ribosomal RNA gene

ATGGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCATGCCTAATACATGCAAGTCGAACGAAGTTTCGAGGA
AGCTTGCTTCCAAAGAGACTTAGTGCGGAACGGGTGAGTAACACGTAGGTAACCTGCCCATGTGTCCGGGATAACTG
CTGGAACCGGTAGCTAAAACCGGATAGGTATACAGAGCGCATGCTCAGTATATTAAAGCGCCCATCAAGGCGTGAAC
ATGGATGGACCTGCGGCGCATTAGCTAGTTGGTGAGGTAACGGCCCCACCAAGGCGATGATGCGTAGCCGGCCTGAGA
GGGTAAACGGCCACATTGGGACTGAGACACGGCCCCAACTCCTACGGGAGGCAGCAGTAGGGAATTTTCGTCAATGG
GGGAAACCTTGAACGAGCAATGCCGCGTGAGTGAAGAAGGTCTTCGGATCGTAAAGCTCTGTTGTAAGTGAAGAACC
GCTCATAGAGGAAATGCTATGGGAGTGACGGTAGCTTACCAGAAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGT
AATACGTAGGTGGCAAGCGTTATCCGGAATCATTGGGCGTAAAGGGTGCGTAGGTGGCGTACTAAGTCTGTAGTAAA
AGGCAATGGCTCAACCATTTGTAAGCTATGGAACCTGGTATGCTGGAGTGCAGAAGAGGGCGATGGAATTCCATGTGT
AGCGGTAAAATGCGTAGATATATGGAGGAACACCAAGTGGCGAAGGCGGTTCGCCTGGTCTGTAAGTACACTGAGGCA
CGAAAGCGTGGGGAGCAAATAGGATTAGATACCCTAGTAGTCCACGCCGTAAACGATGAGAACTAAGTGTGGAGGA
ATTCAGTGCTGCAGTTAACGCAATAAGTTCTCCGCTGGGGAGTATGCACGCAAGTGTGAAACTCAAAGGAATTGAC
GGGGGCCCCGCACAAGCGGTGGAGTATGTGGTTTAATTTCGAAGCAACGCGAAGAACCTTACCAGGCCTTGACATGGAA
ACAAATACCCTAGAGATAGGGGGATAATTATGGATCACACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGA
TGTTGGGTAAAGTCCCGCAACGAGCGCAACCCTTGTTCGCATGTTACCAGCATCAAGTTGGGGACTCATGCGAGACTG
CCGGTGACAAACCGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGGCCTGGGCTACACACGTACTAC
AATGGCGGCCACAAAGAGCAGCGACACAGTGATGTGAAGCGAATCTCATAAAGGTCGTCTCAGTTCGGATTGAAGTC
TGCAACTCGACTTCATGAAGTCGGAATCGCTAGTAATCGCAGATCAGCATGCTGCGGTGAATACGTTCTCGGGCCTT
GTACACACCGCCCCGTCAAACCATGGGAGTCAGTAATACCCGAAGCCGGTGGCATAACCGTAAGGAGTGAGCCGTCTGA
AGGTAGGACCGATGACTGGGGTTAAGTCGTAACAAGGTATCCCTACGGGAACGTGGGGATGGATCACCTCCTTT

>SEQ ID NO: 157 | PROKKA_00199 16S ribosomal RNA gene

TATTGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAACGGGGTGCTCATG
ACGGAGGATTCGTCCAACGGATTGAGTTACCTAGTGGCGGACGGGTGAGTAACGCGTGAGGAACCTGCCTTGGAGAG
GGGAATAACACTCCGAAAGGAGTGCTAATACCGCATGATGCAGTTGGGTTCGCATGGCTCTGACTGCCAAAGATTTAT
CGCTCTGAGATGGCCTCGCGTCTGATTAGCTAGTAGGCGGGGTAACGGCCACCTAGGCGACGATCAGTAGCCGGAC
TGAGAGGTTGACCGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGGGC
AATGGGCGCAAGCCTGACCCAGCAACGCCGCGTGAAGGAAGAAGGCTTTCGGGTTGTAAACTTCTTTTGTCTGGGGAC
GAAACAAATGACGGTACCCGACGAATAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGTGGCAAG
CGTTATCCGGATTTACTGGGTGTAAAGGGCGTGTAGGCGGGATTGCAAGTCAGATGTGAAAACCTGGGGGCTCAACCT
CCAGCCTGCATTTGAAACTGTAGTTCTTGAGTGCTGGAGAGGCAATCGGAATTCCGTGTGTAGCGGTGAAATGCGTA
GATATACGGAGGAACACCACTGGCGAAGGCGGATTGCTGGACAGTAACTGACGCTGAGGCGCGAAAGCGTGGGGAGC
AAACAGGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGGATACTAGGTGTGGGGGTCTGACCCCTCCGTGCC
GCAGTTAACACAATAAGTATCCACCTGGGGAGTACGATCGCAAGGTTGAAACTCAAAGGAATTGACGGGGGCCCCG
ACAAGCGGTGGAGTATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAGGGCTTGACATCCCACTAACGAAGC
AGAGATGCATTAGGTGCCCTTCGGGGAAAGTGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGT
TGGGTAAAGTCCCGCAACGAGCGCAACCCTTATTGTTAGTTGCTACGCAAGAGCACTCTAGCGAGACTGCCGTTGAC
AAAACGGAGGAAGGTGGGGACGACGTCAAATCATCATGCCCCTTATGTCCTGGGCCACACACGTACTACAATGGTGG
TTAACAGAGGGAGGCAATACCGCGAGGTGGAGCAAATCCCTAAAAGCCATCCAGTTCGGATTGCAGGCTGAAACCC
GCCTGTATGAAGTTGGAATCGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCCCGGGCCTTGTACACAC
CGCCCGTCACACCATGAGAGTCGGGAACACCCGAAGTCCGTAGCCTAACCGCAAGGAGGGCGCGGCCGAAGGTGGGT
TCGATAATTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ ID

NO: 158 | PROKKA_00208 16S ribosomal RNA gene

TATTGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAACGGGGTGCTCATG
ACGGAGGATTCGTCCAACGGATTGAGTTACCTAGTGGCGGACGGGTGAGTAACGCGTGAGGAACCTGCCTTGGAGAG
GGGAATAACACTCCGAAAGGAGTGCTAATACCGCATGATGCAGTTGGGTTCGCATGGCTCTGACTGCCAAAGATTTAT
CGCTCTGAGATGGCCTCGCGTCTGATTAGCTAGTAGGCGGGGTAACGGCCACCTAGGCGACGATCAGTAGCCGGAC
TGAGAGGTTGACCGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGGGC
AATGGGCGCAAGCCTGACCCAGCAACGCCGCGTGAAGGAAGAAGGCTTTCGGGTTGTAAACTTCTTTTGTCTGGGGAC
GAAACAAATGACGGTACCCGACGAATAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGTGGCAAG
CGTTATCCGGATTTACTGGGTGTAAAGGGCGTGTAGGCGGGATTGCAAGTCAGATGTGAAAACCTGGGGGCTCAACCT
CCAGCCTGCATTTGAAACTGTAGTTCTTGAGTGCTGGAGAGGCAATCGGAATTCCGTGTGTAGCGGTGAAATGCGTA
GATATACGGAGGAACACCACTGGCGAAGGCGGATTGCTGGACAGTAACTGACGCTGAGGCGCGAAAGCGTGGGGAGC
AAACAGGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGGATACTAGGTGTGGGGGGTCTGACCCCTCCGTGC
CGCAGTTAACACAATAAGTATCCACCTGGGGAGTACGATCGCAAGGTTGAAACTCAAAGGAATTGACGGGGGCCCCG
CACAAGCGGTGGAGTATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAGGGCTTGACATCCCACTAACGAAG
CAGAGATGCATTAGGTGCCCTTCGGGGAAAGTGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATG
TTGGGTAAAGTCCCGCAACGAGCGCAACCCTTATTGTTAGTTGCTACGCAAGAGCACTCTAGCGAGACTGCCGTTGA
CAAAACGGAGGAAGGTGGGGACGACGTCAAATCATCATGCCCCTTATGTCCTGGGCCACACACGTACTACAATGGTG
GTTAACAGAGGGAGGCAATACCGCGAGGTGGAGCAAATCCCTAAAAGCCATCCAGTTCGGATTGCAGGCTGAAACC
CGCCTGTATGAAGTTGGAATCGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCCCGGGCCTTGTACACA
CCGCCCCGTACACCATGAGAGTCGGGAACACCCGAAGTCCGTAGCCTAACCGCAAGGAGGGCGCGGCCGAAGGTGGG
TTCGATAATTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT >SEQ

ID NO: 159 | PROKKA_04460 16S ribosomal RNA gene

TATTGAGAGTTTGATCCTGGCTCAGGATGAACGCTGGCGGCGTGCTTAACACATGCAAGTCGAACGGGGTGCTCATG
ACGGAGGATTCGTCCAACGGATTGAGTTACCTAGTGGCGGACGGGTGAGTAACGCGTGAGGAACCTGCCTTGGAGAG
GGGAATAACACTCCGAAAGGAGTGCTAATACCGCATAATGCAGTTGGGTTCGCATGGCTCTGACTGCCAAAGATTTAT
CGCTCTGAGATGGCCTCGCGTCTGATTAGCTAGTAGGCGGGGTAACGGCCACCTAGGCGACGATCAGTAGCCGGAC
TGAGAGGTTGACCGGCCACATTGGGACTGAGACACGGCCCAGACTCCTACGGGAGGCAGCAGTGGGGAATATTGGGC
AATGGGCGCAAGCCTGACCCAGCAACGCCGCGTGAAGGAAGAAGGCTTTCGGGTTGTAAACTTCTTTTGTCTAGGGAC
GAAACAAATGACGGTACCTGACGAATAAGCCACGGCTAACTACGTGCCAGCAGCCGCGGTAATACGTAGGTGGCAAG
CGTTATCCGGATTTACTGGGTGTAAAGGGCGTGTAGGCGGGATTGCAAGTCAGATGTGAAAACCTGGGGGCTCAACCT
CCAGCCTGCATTTGAAACTGTAGTTCTTGAGTGCTGGAGAGGCAATCGGAATTCCGTGTGTAGCGGTGAAATGCGTA
GATATACGGAGGAACACCACTGGCGAAGGCGGATTGCTGGACAGTAACTGACGCTGAGGCGCGAAAGCGTGGGGAGC
AAACAGGATTAGATACCCTGGTAGTCCACGCCGTAAACGATGGATACTAGGTGTGGGGGGTCTGACCCCTCCGTGCC
GCAGTTAACACAATAAGTATCCACCTGGGGAGTACGATCGCAAGGTTGAAACTCAAAGGAATTGACGGGGGCCCCG
ACAAGCGGTGGAGTATGTGGTTTAATTCGAAGCAACGCGAAGAACCTTACCAGGGCTTGACATCCCACTAACGAAGC
AGAGATGCATTAGGTGCCCTTCGGGGAAAGTGGAGACAGGTGGTGCATGGTTGTCGTCAGCTCGTGTCTGTGAGATGT
TGGGTAAAGTCCCGCAACGAGCGCAACCCTTATTGTTAGTTGCTACGCAAGAGCACTCTAGCGAGACTGCCGTTGAC
AAAACGGAGGAAGGTGGGGACGACGTCAAATCATCATGCCCCTTATGTCCTGGGCCACACACGTACTACAATGGTGG
TTAACAGAGGGAGGCAATACCGCGAGGTGGAGCAAATCCCTAAAAGCCATCCAGTTCGGATTGCAGGCTGAAACCC
GCCTGTATGAAGTTGGAATCGCTAGTAATCGCGGATCAGCATGCCGCGGTGAATACGTTCCCGGGCCTTGTACACAC
CGCCCGTCACACCATGAGAGTCGGGAACACCCGAAGTCCGTAGCCTAACCGCAAGGAGGGCGCGGCCGAAGGTGGGT
TCGATAATTGGGGTGAAGTCGTAACAAGGTAGCCGTATCGGAAGGTGCGGCTGGATCACCTCCTTT

(262) This invention is not limited to its application to the details of construction and the arrangement of components set forth in the following description or illustrated in the drawings. The invention is capable of other embodiments and of being practiced or of being carried out in various ways. Also, the phraseology and terminology used herein is for the purpose of description and should not be regarded as limiting. The use of “including,” “comprising,” or “having,” “containing,” “involving,” and variations thereof herein, is meant to encompass the items listed thereafter and equivalents thereof as well as additional items.

(263) Unless otherwise defined herein, scientific and technical terms used in connection with the present disclosure shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. The methods and techniques of the present disclosure are generally performed according to conventional methods well-known in the art. Generally, nomenclatures used in connection with, and techniques of biochemistry, enzymology, molecular and cellular biology, microbiology, virology, cell or tissue culture, genetics and protein and nucleic chemistry described herein are those well-known and commonly used in the art. The methods and techniques of the present disclosure are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated.

(264) The present invention is further illustrated by the following Examples, which in no way should be construed as further limiting. The entire contents of all of the references (including literature references, issued patents, published patent applications, and co-pending patent applications) cited throughout this application are hereby expressly incorporated by reference, in particular for the teaching that is referenced hereinabove. However, the citation of any reference is not intended to be an admission that the reference is prior art.

EXAMPLES

Example 1: Mouse Model of *C. difficile* Infection

(265) Mouse Husbandry

(266) Experiments were performed using C57BL/6J female mice purchased from Jackson Laboratories (Bar Harbor, ME) and housed in ventilated sterile cages. All animals were maintained in a specific-pathogen-free facility. Animals were acclimated to the vivarium for at least 3 days prior to study (i.e., commencing antibiotic courses). For experiments involving *C. difficile* infection, mice were administered 10-10^{sup.4} *C. difficile* VPI 10463 spores in 200 µl PBS by oral gavage. Experiments were performed in compliance with institutional guidelines and approved by the institution's Institutional Animal Care and Use Committee. Sterile food and drinking water were provided to the animals

(267) Live Biotherapeutic Product (LBP) Preparation

(268) Individual bacterial strains were isolated from fecal material obtained from healthy donors. The individual strains were struck out from 15% glycerol freezer stocks onto EG (Eggerth Gagnon) agar plates containing 5% horse blood in an anaerobic chamber and incubated for 24-48 hours at 37° C. Colonies were inoculated into pre-reduced liquid Peptone Yeast Glucose (PYG) media and grown for 24-48 hours until dense (static in the anaerobic chamber). Optical density (OD_{sub.600}) of the cultures was assessed and live biotherapeutic product (LBP) cocktails were prepared inside an anaerobic chamber adjusting inputs based upon OD_{sub.600} for equal CFU ratio cocktails in PBS (sterile, pre-treated).

(269) *C. difficile* Colony Forming Unit (CFU) Determination

(270) Fecal pellets were collected, transported to an anaerobic chamber (<2 hours), and manually homogenized in 500 µL of pre-reduced PBS using a pipette tip and through repeated pipetting. Serial dilutions of fecal homogenates were prepared in pre-reduced PBS, 100 µL of which was spread onto cycloserine-cefoxitin-fructose agar with sodium taurocholate (TCCFA) plates, and incubated anaerobically at 37° C. *C. difficile* CFUs were enumerated at 48 hours.

(271) Murine Susceptibility to *C. difficile* Infection

(272) Groups of mice were evaluated for susceptibility to *C. difficile* using three antibiotic regimen protocols: (1) an antibiotic cocktail, (2) clindamycin administration, or (3) cefoperazone administration (FIGS. 2 and 3). The antibiotic cocktail consisted of kanamycin (0.4 mg/ml), gentamicin (0.035 mg/ml), colistin (0.056 mg/ml), metronidazole (0.215 mg/ml), vancomycin (0.045 mg/ml) in the drinking water from day -10 to day -3, followed by a single intraperitoneal clindamycin injection (200 µg/mouse). The clindamycin administration involved a single intraperitoneal injection of clindamycin (200 µg/mouse) on day -1. The notation of days is relative to day 0, the day of *C. difficile* infection.

(273) Mice were treated with the indicated antibiotic regimen as described above and then infected with either 10 or 10^{sup.4} *C. difficile* spores by oral gavage on day 0 (FIGS. 2 and 3). An additional experimental arm was added to the antibiotic treatment model in which mice were treated with vancomycin after *C. difficile* infection (FIG. 4J; black triangles).

(274) Mice were monitored daily following infection for mortality/survival (FIGS. 4A-4D) and weight (FIGS. 4E-4H). Fecal pellets were also collected daily and used for *C. difficile* CFU enumeration, presented as CFU/gram feces (FIGS. 4I-4L).

(275) The groups of mice that received cefoperazone treatment had a significant change in weight (FIG. 4H) and substantial *C. difficile* bacterial load in the fecal pellets (FIG. 4L), even following administration with 10 *C. difficile* spores. These results indicated that the cefoperazone pre-treatment regimen provided a good model for *C. difficile* infection and for evaluating protection and/or treatment of *C. difficile* infection. In the absence of antibiotic treatment prior to infection, *C. difficile* infection was not established (FIG. 4I) and all mice survived (FIG. 4A) without significant change in body weight (FIG. 4E).

Example 2: Live Biotherapeutic Product (LBP) Preparations Protect Against *C. difficile* Infection

(276) The following LBP compositions were evaluated for their capacity to protect and/or treat *C. difficile* infection:

Composition A, Composition B, Composition C, Composition D, Composition E (See e.g., Narushima et al., Gut Microbes (2014) 5(3) 333-339), and Composition I: a mixture of *Clostridium scindens*, *Pseudoflavonifractor capillosus* and *Blautia*

hansenii (FIG. 5).

(277) In general, LBP cocktails were mixed in PYG media, and each mouse was administered a dose by oral gavage in 250 μ L pre-reduced PBS (media-free). For composition E, bacteria were mixed in equal volumes (not equal ratios/CFUs) and administered in a 250 μ L dose. Each LBP of Compositions A-D contained 10^{sup.8} CFUs total in a 250 μ L dose, comprised of 10^{sup.7} CFU of each of the bacterial strains (FIG. 1), for a total of 10^{sup.8} CFU administered to each animal. Composition I contained a total of 10^{sup.6} CFUs in a 250 μ L dose (approximately 333,000 of each of the 3 bacteria mixed). (278) Groups of mice were subjected to cefoperazone treatment, as described in Example 1, and were administered the indicated composition by oral gavage 2 days after the cessation of cefoperazone treatment. Twenty-four hours later, the mice were subjected to infection with 10^{sup.4} *C. difficile* spores (FIG. 5). Mice evaluated for survival/mortality (FIG. 6), weight (FIGS. 7A-7I, and *C. difficile* CFUs (FIGS. 8A-8C). The results show that administration of Composition B prior to *C. difficile* infection is an effective protection and/or treatment against *C. difficile* infection.

Example 3: Composition B Protects Against and/or Treats *C. difficile* Infection

(279) Groups of 10-12 week old mice were used in the *C. difficile* mouse model (FIG. 9). Mice were subjected to cefoperazone treatment as described in Example 1. One group of mice was then administered Composition B (10^{sup.8} CFU per mouse) administered by oral gavage, as described in Examples 1 and 2, 2 days after the cessation of cefoperazone treatment. The other group of mice did not receive a live biotherapeutic product after cefoperazone treatment (control). Twenty-four hours later, the mice were subjected to *C. difficile* infection (10^{sup.4} *C. difficile* spores) and then evaluated for survival/mortality (FIG. 10), weight (FIG. 11), and *C. difficile* burden (CFUs per gram feces; FIG. 12). These results confirm the results of Example 2 that demonstrate treatment with Composition B prior to *C. difficile* infection is an effective protection and/or treatment against *C. difficile* infection.

Example 4: LBP Composition F Protects Against and/or Treat *C. difficile* Infection

(280) FIG. 13 shows the strains of live biotherapeutic product (LBP) Composition F. The genus-species classification indicates the closest species based on the sequence of the isolated strain. FIG. 14 shows the classification by *Clostridium* cluster of the strains in Composition F.

(281) Groups of mice were administered cefoperazone, as described in the Examples above, then administered LBPs or fecal matter transplant (FMT) from mice or human (FIG. 15). Composition B was administered to the indicated groups on day -1; days -2 and -1; or on days -2, -1, 1, 2, and 3, relative to infection with 10^{sup.4} *C. difficile* spores. Composition F was administered to the indicated groups on day -1 or on days -2, -1, 1, 2, and 3, relative to administration of *C. difficile* spores. Additional groups received FMT from mice or from humans (200 μ L of a 10% fecal sample s per mouse). Mice were then evaluated for survival/mortality (FIG. 16), weight (FIGS. 17A-17H), and *C. difficile* burden (CFU/gram feces) on days 1, 3, 8 and 17 after infection (FIGS. 18A and 18B). The data demonstrate that Composition B, Composition F, and FMT protect against and/or treat *C. difficile* infection.

Example 5: LBP Compositions Protect Against and/or Treat *C. difficile* Infection

(282) FIG. 19 shows the strains of LBP Composition G. The genus-species notation indicates the closest species based on the sequence of the isolated strain. Composition G includes a subset of the strains of Composition F. Groups of mice were administered cefoperazone, as described in the Examples above, then administered the LBP: Composition B; Composition B-1 (Composition B with *Bacteroides* added); Composition B-2 (Composition B from which *Flavonifractor plautii* was removed and *Bacteroides* added); Composition F; Composition G; Human fecal samples subjected to ethanol treatment; Composition B subjected to ethanol treatment; Composition B that had been frozen; or Composition J: *Clostridium innocuum*, *Clostridium bolteae* and *Clostridium symbiosum* subjected to ethanol treatment; (See also FIG. 20).

(283) The *Bacteroides* strain used in Composition B-1 and B-2 was *Bacteroides ovatus* (strain identifier 211-B; SEQ ID NO: 83).

(284) Mice were challenged with *C. difficile* VPI 10463 spores (10^{sup.4}) and monitored daily (Day 0 to Day 7 post *C. difficile* infection) for survival/mortality (FIGS. 21 and 23) and change in weight (FIGS. 22A-22J and 24). These data show that the compositions protect against and/or treat *C. difficile* infection.

Example 6: LBP Compositions Protect Against and/or Treat *C. difficile* Infection

(285) Groups of mice were subjected to cefoperazone treatment, as described above, then administered human fecal matter transplant, Composition B, Composition B+4 spores, or Composition H (FIG. 25). "Composition B+4 spores" refers to Composition B plus the following four strains in spore form: *Clostridium bolteae*, *Anaerotruncus colihominis*, *Clostridium symbiosum* and *Clostridium innocuum*. Composition H contains the following six strains in spore form: *Clostridium bolteae*, *Anaerotruncus colihominis*, *Clostridium symbiosum*, *Clostridium innocuum*, *Clostridium disporicum* and *Erysipelatoclostridium ramosum* (FIG. 26).

(286) Mice were then challenged with *C. difficile* infection with 10^{sup.4} *C. difficile* VPI 10463 spores and monitored for survival/mortality (FIGS. 27A and 28A), weight (FIGS. 27B and 28B). Mice that lost more than 20% body weight relative to baseline were included in mortality numbers in survival curves. The *C. difficile* burden was assessed by CFU in fecal pellets on days 1, 4 and 19 after infection (FIGS. 29A-29C).

(287) These data indicate that Composition B as well as other compositions can improve survival in the cefoperazone-induced *C. difficile* mouse model and protect against and/or treat *C. difficile* infection.

Example 7: *C. difficile* Toxin Experiment

(288) Vero cells, epithelial cells derived from African Green Monkey kidney epithelium, are sensitive to a variety of bacterial toxins, including *C. difficile* Toxin B. Exposure of cells to *C. difficile* Toxin B results in inhibition of the function of Rho, Rac, and Cdc42 leading to a decline in F-actin, a change in cell morphology (e.g., cell rounding), and eventually apoptosis.

(289) To determine whether administration of bacterial compositions described herein has an effect on the production or

activity of *C. difficile* Toxin B, a cellular assay was performed. Briefly, groups of mice were treated with cefoperazone, as described above, and administered human fecal matter transplant (FMT) (“4-3”); Composition B (“5-3”); Composition B plus four strains in spore form: *Clostridium bolteae*, *Anaerotruncus colihominis*, *Clostridium symbiosum* and *Clostridium innocuum* (“7-4”), or no treatment. Each of the groups of mice were then exposed to *C. difficile* infection with 10^{sup.4} *C. difficile* spores. The groups of mice that did not receive a treatment after cefoperazone administration and prior to *C. difficile* infection are referred to as “2-1 (Cdiff)” and “2-4 (Cdiff).” An additional group of mice was not exposed to *C. difficile* as indicated by “N3 (Healthy)”.

(290) Fecal pellets were collected from each of the groups of mice, weighed, and homogenized in PBS and normalized to a fixed concentration (~25 mg/mL). The samples were centrifuged to prepare a clarified supernatant, which was then diluted in 10-fold serial dilutions to produce a range from 1:10 to 1:10⁶ dilutions of clarified pellet supernatant. Vero cell cultures were exposed to the diluted samples for approximately 18 hours, then visualized by phase contract microscopy to assess morphological changes (i.e., cell rounding) associated with *C. difficile* toxin exposure. The cells were scored based on the highest concentration of supernatant that did not yield a change in morphology (FIG. 30). The samples from mice that had been treated with Composition B prior to *C. difficile* infection had reduced amounts of *C. difficile* Toxin B, as compared to samples from control mice that did not receive a treatment after cefoperazone administration and prior to *C. difficile* infection (“2-1 (Cdiff)” and “2-4 (Cdiff)”) as well as compared to samples from mice that received FMT. Notably, the samples from mice that had been treated with Composition B also had reduced amounts of *C. difficile* Toxin B, as compared to samples from mice that had been treated with Composition B with additional spores.

Example 8: In Vitro Competition Between Compositions B and *C. difficile*

(291) Composition B was assessed for its ability to suppress *Clostridium difficile* growth by an in vitro mixed culture competition assay. From glycerol freezer stocks, individual strains of Composition B, *C. difficile* (Cdiff), *Clostridium bifermentans*, and *Bacteroides thetaiotaomicron* were struck out onto Eggerth-Gagnon agar plates with horse blood (EG+HB). Single colonies of each of the strains were subsequently inoculated into brain heart infusion (BHI) liquid media and allowed to grow in pure culture for 24-48 hours. Turbid cultures were sub-cultured then grown to exponential phase and finally diluted and combined to prepare a mixed culture with an optical density (OD₆₀₀) of 0.1. Exponential phase Cdiff culture was added to the mixed culture at a final concentration with an OD of 0.1. After the cultures were combined and incubated for 2-3 hours, samples were collected, serially diluted, and plated on Taurocholate-Cycloserine-Cefoxitin-Fructose Agar (TCCFA) plates to select for Cdiff growth. After 48-72 hours, the colony forming units (CFUs) of Cdiff in each competition experiment were determined by manual colony counting.

(292) EG+HB agar plates were prepared according to standard procedures and reduced in an anaerobic environment for at least 6-8 hours prior to use. Liquid BHI medium was obtained from BD Biosciences (Catalog #211059, San Jose, CA), prepared according to the manufacturer's instructions, and reduced in an anaerobic environment for at least 18-24 hours prior to use. TCCFA plates were prepared according to standard procedures and reduced in an anaerobic environment for at least 6-8 hours prior to use. *Clostridium difficile* strain used in the experiments: American Type Culture Collection (ATCC) 43255.

(293) TABLE-US-00015 TABLE 4 Composition B strains Composition B VE202-7 VE202-13 VE202-14 VE202-16 Strain #16 Strain #170 Strain #189 Strain #211

(294) Strains were struck out onto EG+HB agar plates from frozen glycerol stocks inside an anaerobic chamber for 48-72 hours. Single colonies were inoculated into 10 mL of BHI media and grown 24-48 hours at 37° C. in the anaerobic chamber. Turbid cultures were then diluted to an OD of 0.1 and grown for 2-3 hours at 37° C. in the anaerobic chamber. Exponential phase cultures were diluted and combined at equivalent ODs. For the competition assay, each of the strains of Combination B (Table 4) were combined in equal parts, based on OD_{sub.600}, to reach a final consortium OD_{sub.600} of 0.1. *C. bifermentans* and *B. thetaiotaomicron* were setup to compete with Cdiff individually at an OD of 0.1. The OD_{sub.600} for Cdiff in each of the mixed culture competition experiments was 0.1. After combination, the cultures were incubated for 2-3 hours at 37° C. in the anaerobic chamber, then prepared for enumerations on Cdiff selective plates.

(295) TCCFA plates are selective for Cdiff growth, and none of the Combination B strains, nor either of the control strains (*C. bifermentans* and *B. thetaiotaomicron*), grow on these plates. Inside an anaerobic chamber, a 100 µL sample of each competition culture was collected and serially diluted 1:10 to reach a final dilution of 1×10^{sup.-6}. Plates for CFU enumeration were prepared by spreading 100 µL of each of the 1×10^{sup.-4} through 1×10^{sup.-6} dilutions on TCCFA plates using sterile spreading loops. CFU plates were incubated for 48-72 hours at 37° C. in the anaerobic chamber. CFU enumeration was completed by manually counting colonies.

(296) To determine the effect of competition, the ratio of CFUs determined for the competition samples and Cdiff alone was calculated and expressed as a percentage. Inhibition of Cdiff growth by the Composition B cocktail was compared to the responses of *B. thetaiotaomicron* (negative control) and *C. bifermentans* (positive control). The results are shown in Table 5 and FIG. 31.

(297) TABLE-US-00016 TABLE 5 Summary Results for In Vitro Competition Competition with Exper- No Competition
Composi- iment Competing Competition with with tion Number Strain(s) *B. thetaiotaomicron* *C. bifermentans* B n = 1 100
33.8 n = 2 100 9.90 0.1 0.5 n = 3 100 115 39.5 33.1 n = 4 100 41.3 0.7 0.7 n = 5 100 105 14.1 20.9 n = 6 100 57.4 4.1 1.6
Mean 100 65.6 11.7 15.1 Std. Dev. 0 43.8 16.5 16.2 Total N 6 5 5 6

(298) Data is expressed as Cdiff CFU as a percentage of control. Each n is representative of a single biological replicate, independent of other measurements.

(299) In in vitro competition, Composition B inhibited Cdiff growth to 15.1±16.2% of control (absence of competing strain(s)). This result is consistent with the inhibition observed by the positive control, *C. bifermentans*, of 11.7±16.5% of control. *B. thetaiotaomicron*, a negative control, yielded a negligible effect on Cdiff growth at 65.6±43.8% of control. Given

the variability between the assessment of CFU, inhibition of growth to <25% of control is considered to be significant inhibition and both the positive control and Composition B cocktail meet this threshold of activity. The Composition B consortium attenuated Cdiff growth in vitro comparable to the direct competition observed by *C. bifermentans*. Direct competition with *B. thetaiotaomicron* did not significantly inhibit Cdiff growth.

Example 9: Determination of In Vitro Short-Chain Fatty Acid Production

(300) Each strain of Composition B was assessed for individual short-chain fatty acid (SCFA) production in vitro.

Composition B strains were grown in pure cultures inside an anaerobic chamber. Spent supernatant from liquid media cultures was harvested by centrifugation, filter sterilized, and then stored at <-70° C. Frozen clarified supernatant specimens were analyzed for short-chain fatty acids (SCFAs).

(301) EG+HB agar plates (Eggerth-Gagnon agar plates with horse blood) were prepared according to standard methods and reduced in an anaerobic environment for at least 6-8 hours prior to use. Liquid PYG medium (pre-formulated, pre-reduced) was obtained from Anaerobe Systems (Catalog #AS-822; Morgan Hill, CA).

(302) Strains were struck out onto EG+HB agar plates from frozen 15% glycerol stocks inside an anaerobic chamber for 48-72 hours. Single colonies were inoculated into 7 mL PYG media and grown 24-48 hours at 37° C. in the anaerobic chamber. Unless otherwise noted, when the optical density (OD) was ≥0.2, samples were collected for CFU enumeration and filtration. Inside an anaerobic chamber, a 100 µL sample of turbid culture was collected and serially diluted 1:10 to reach a final dilution of 1×10^{sup}.-6. Plates for CFU enumeration were prepared by spreading 100 µL/dilution for the 1×10^{sup}.-4 through 1×10^{sup}.-6 dilutions on EG+HB agar plates using sterile glass beads. CFU plates were incubated for 48-72 hours in the anaerobic chamber. CFU enumeration was completed using the EasyCount 2 (bioMérieux SA, Marcy-l'Étoile, France). Immediately after samples of turbid cultures were collected for CFU enumeration, the remaining turbid cultures were centrifuged at approximately 1000 RCF for 10 minutes to pellet cellular debris. The clarified supernatants were transferred to a 0.2 µm plate filter and vacuum filtered to remove any remaining particulates prior to bioanalysis. In the event of blockage in the filter plate, clarified supernatants were manually filtered using 0.2 µm syringe filters. Filtered supernatants were aliquoted and stored at <-70° C. prior to bioanalysis of SCFAs.

(303) To facilitate easier comparisons between samples, raw SCFA data (µg/mL) was normalized by the log of corresponding determined/estimated CFU for the culture. The results are depicted in Table 6 and Table 7 below.

(304) TABLE-US-00017 TABLE 6 Enumerated CFUs for Composition B Strains Enumerated CFU Sample ID OD600 (CFU/mL) VE202-7 >2 6.11E+08 VE202-13 0.8 4.00E+08 VE202-14 >2 1.60E+09 VE202-16 1.92 1.28E+09 #16 1.97 1.69E+08 # 170 1.8 1.08E+08 # 189 1.03 1.74E+09 # 211 0.35 3.71E+08

(305) TABLE-US-00018 TABLE 7 SCFAs produced by individual Composition B strains Normalized (µg/Log(CFU/mL)*mL) 2- Sample Iso- Methyl- Iso- ID Acetate Propionate butyrate Butyrate butyrate valerate Valerate Hexanoate VE202-7 123.7 0.077 0.102 0.208 0.015 0.056 BLOQ 0.031 VE202-13 30.1 0.545 0.116 34.452 0.288 0.188 0.097 0.034 VE202-14 110.5 0.054 0.022 0.248 0.011 0.014 BLOQ 0.009 VE202-16 313.2 0.000 0.000 0.280 0.004 0.000 BLOQ 0.009 #16 104.0 0.005 0.000 50.988 0.014 0.033 BLOQ 0.009 # 170 87.1 0.055 0.025 0.215 0.011 0.039 BLOQ 0.016 # 189 0.0 BLOQ 0.000 35.751 0.005 0.019 0.359 0.587 # 211 57.6 5.289 0.000 78.227 0.028 0.050 0.053 0.095

(306) Seven strains of Composition B were found to produce significant quantities (>1 µg/Log(CFU/mL)*mL) of the 2-carbon SCFA, acetate. One strain, (#211), produced substantial quantities of the 3-carbon SCFA, propionate. Four strains of Composition B produced substantial quantities of the 4-carbon SCFA, butyrate. Trace quantities (<1 µg/Log(CFU/mL)*mL) of other SCFAs were also produced by the Composition B strains.

Example 10: Composition B Induces Regulatory T Cells (Tregs)

(307) Each of the bacterial strains of Composition B were grown to log phase, combined to a total dose of ~10^{sup}.8 cfu per mouse. Germ-free mice were inoculated with Composition B or a negative control by oral gavage and sacrificed following four weeks of colonization. *Lamina propria* leukocytes were isolated from colonic tissue of individual mice by standard procedures and assessed by flow cytometry. The regulatory T cell content was evaluated as the percentage of Foxp3-positive cells among CD4+ T cells.

(308) As shown in FIG. 32, mice that were inoculated with Composition B were found to have significantly more regulatory T cells as compared to mice that were inoculated with the control.

Claims

1. A pharmaceutical composition comprising a purified bacterial mixture, wherein the purified bacterial mixture consists of 7 to 10 bacterial strains, wherein the purified bacterial mixture comprises at least 7 of: (i) a bacterial strain comprising a 16S rDNA sequence having at least 97% sequence identity to the nucleotide sequence of SEQ ID NO: 157; (ii) a bacterial strain comprising a 16S rDNA sequence having at least 97% sequence identity to the nucleotide sequence of SEQ ID NO: 129; (iii) a bacterial strain comprising a 16S rDNA sequence having at least 97% sequence identity to the nucleotide sequence of SEQ ID NO: 132; (iv) a bacterial strain comprising a 16S rDNA sequence having at least 97% sequence identity to the nucleotide sequence of SEQ ID NO: 137; (v) a bacterial strain comprising a 16S rDNA sequence having at least 97% sequence identity to the nucleotide sequence of SEQ ID NO: 124; (vi) a bacterial strain comprising a 16S rDNA sequence having at least 97% sequence identity to the nucleotide sequence of SEQ ID NO: 141; (vii) a bacterial strain comprising a 16S rDNA sequence having at least 97% sequence identity to the nucleotide sequence of SEQ ID NO: 146; and (viii) a bacterial strain comprising a 16S rDNA sequence having at least 97% sequence identity to the nucleotide sequence of SEQ ID NO: 152, wherein the bacterial strains are lyophilized.

2. The pharmaceutical composition of claim 1, wherein the purified bacterial mixture comprises: (i) a bacterial strain

comprising a 16S rDNA sequence having at least 99% sequence identity to the nucleotide sequence of SEQ ID NO: 157; (ii) a bacterial strain comprising a 16S rDNA sequence having at least 99% sequence identity to the nucleotide sequence of SEQ ID NO: 129; (iii) a bacterial strain comprising a 16S rDNA sequence having at least 99% sequence identity to the nucleotide sequence of SEQ ID NO: 132; (iv) a bacterial strain comprising a 16S rDNA sequence having at least 99% sequence identity to the nucleotide sequence of SEQ ID NO: 137; (v) a bacterial strain comprising a 16S rDNA sequence having at least 99% sequence identity to the nucleotide sequence of SEQ ID NO: 124; (vi) a bacterial strain comprising a 16S rDNA sequence having at least 99% sequence identity to the nucleotide sequence of SEQ ID NO: 141; (vii) a bacterial strain comprising a 16S rDNA sequence having at least 99% sequence identity to the nucleotide sequence of SEQ ID NO: 146; and (viii) a bacterial strain comprising a 16S rDNA sequence having at least 99% sequence identity to the nucleotide sequence of SEQ ID NO: 152.

3. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises a pH-sensitive composition comprising one or more enteric polymers.
 4. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition is in the form of a capsule.
 5. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition comprises between 1×10^7 and 1×10^{10} colony forming units (CFUs) per bacterial strain.
 6. The pharmaceutical composition of claim 1, further comprising a pharmaceutically acceptable excipient.
 7. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition is formulated for oral administration.
 8. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition is formulated for rectal administration.
 9. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition is formulated for delivery to the intestine.
 10. The pharmaceutical composition of claim 1, wherein the pharmaceutical composition is formulated for delivery to the colon.
 11. A method of reducing the likelihood of a *Clostridium difficile* infection in a subject, the method comprising administering the pharmaceutical composition of claim 1 to the subject in a therapeutically effective amount to reduce the likelihood of the *Clostridium difficile* infection.
 12. The method of claim 11, wherein the *Clostridium difficile* infection is a first occurrence of a *Clostridium difficile* infection.
 13. The method of claim 11, wherein the *Clostridium difficile* infection is a recurrence of *Clostridium difficile* infection.
 14. The method of claim 11, wherein the subject is administered one or more doses of an antibiotic prior to the pharmaceutical composition.
 15. The method of claim 14, wherein the antibiotic is vancomycin, kanamycin, gentamicin, colistin, metronidazole, clindamycin, fidaxomicin, or cefoperazone.
 16. The method of claim 14, wherein the antibiotic is vancomycin.
 17. A method to suppress an abnormal or excessive immune response in a subject comprising administering the pharmaceutical composition of claim 1 to the subject in a therapeutically effective amount to suppress the abnormal or excessive immune response.
 18. The method of claim 17, wherein the abnormal or excessive immune response is suppressed by inducing proliferation and/or accumulation of regulatory T cells.
 19. The method of claim 17, wherein the subject is administered one or more doses of an antibiotic prior to the pharmaceutical composition.
 20. The method of claim 19, wherein the antibiotic is vancomycin, kanamycin, gentamicin, colistin, metronidazole, clindamycin, fidaxomicin, or cefoperazone.
-