



US 20250257318A1

(19) United States

(12) Patent Application Publication (10) Pub. No.: US 2025/0257318 A1

Noggle et al.

(43) Pub. Date: Aug. 14, 2025

(54) METHOD AND COMPOSITION FOR
GENERATING BASAL FOREBRAIN
CHOLINERGIC NEURONS (BFCNS)(71) Applicants: New York Stem Cell Foundation, Inc.,
New York, NY (US); Icahn School of
Medicine at Mount Sinai, New York,
NY (US)(72) Inventors: Scott Noggle, New Rochelle, NY (US);
Maitane Ortiz-Virumbrales, New
York, NY (US); Sam Gandy, New
York, NY (US); Ilya Kruglikov, New
York, NY (US); Michelle Ehrlich, New
York, NY (US)

(21) Appl. No.: 18/925,851

(22) Filed: Oct. 24, 2024

Related U.S. Application Data

- (60) Division of application No. 17/146,336, filed on Jan. 11, 2021, now abandoned, which is a continuation of application No. 15/990,208, filed on May 25, 2018, now Pat. No. 10,889,800.
- (60) Provisional application No. 62/586,571, filed on Nov. 15, 2017, provisional application No. 62/574,639, filed on Oct. 19, 2017, provisional application No. 62/571,741, filed on Oct. 12, 2017, provisional application No. 62/511,271, filed on May 25, 2017.





Publication Classification

(51)	Int. Cl.		
	C12N 5/0793	(2010.01)	
	A61K 35/30	(2015.01)	
	C12N 5/074	(2010.01)	
	C12N 9/22	(2006.01)	
	C12N 15/11	(2006.01)	
	G01N 33/50	(2006.01)	
(52)	U.S. Cl.		
	CPC	C12N 5/0619 (2013.01); A61K 35/30 (2013.01); C12N 5/0696 (2013.01); C12N 9/22 (2013.01); C12N 15/11 (2013.01); G01N 33/5058 (2013.01); C12N 2310/20 (2017.05); C12N 2501/15 (2013.01); C12N 2501/41 (2013.01); C12N 2501/999 (2013.01); C12N 2506/02 (2013.01); C12N 2506/45 (2013.01); C12N 2510/00 (2013.01); C12N 2800/80 (2013.01); G01N 2800/2814 (2013.01)	

ABSTRACT





The invention relates to methods and compositions for developing basal forebrain cholinergic neurons (BFCNs) from stem cells, and in particular, BFCNs having repaired electrophysiological defects relating to one or more mutations in PSEN2, and to the use of such BFCNs in cell-based therapies to treat Alzheimer's disease.



Specification includes a Sequence Listing.



IPSC Lines	Gender	Age	PSEN2 genotype	ApoE	Family Related
948 (AD1)	F	40	x1419WT	e3e4	Yes
949 (Control)	F	33	WT/WT	e3e3	Yes
960 (AD2)	F	37	x1419WT	e3e4	Yes
000843 (Control)	M	26	WT/WT	e3e3	No

FIG. 2A









FIG. 4A

FIG. 4B

AP-40 AP-40




FIG. 4C

FIG. 4D




FIG. 5A




FIG. 5B




FIG. 6A
AS.89




FIG. 6B




FIG. 6C




FIG. 6D
BFCNs




FIG. 6E





FIG. 6F






FIG. 8A




FIG. 8B




FIG. 8C




FIG. 9A




FIG. 9B




FIG. 9C




FIG. 9D




FIG. 10A




FIG. 10B




FIG. 11B




FIG. 11A

METHOD AND COMPOSITION FOR GENERATING BASAL FOREBRAIN CHOLINERGIC NEURONS (BFCNS)**CROSS REFERENCE TO RELATED APPLICATIONS**

[0001] This application is a continuation application of U.S. application Ser. No. 15/990,208 filed May 25, 2018, now issued as U.S. Pat. No. 10,889,800; which claims the benefit under 35 USC § 119(e) to U.S. Application Ser. No. 62/586,571 filed Nov. 15, 2017, U.S. Application Ser. No. 62/574,639 filed Oct. 19, 2017, U.S. Application Ser. No. 62/571,741 filed Oct. 12, 2017 and U.S. Application Ser. No. 62/511,271 filed May 25, 2017, all now expired. The disclosure of each of the prior applications is considered part of and is incorporated by reference in the disclosure of this application.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0002] This invention was made with government support under Grant Nos. AG005138, AG046170 and AG042965 awarded by the National Institutes of Health. The government has certain rights in this invention.

INCORPORATION OF SEQUENCE LISTING

[0003] The material in the accompanying sequence listing is hereby incorporated by reference into this application. The accompanying sequence listing text file, named NYSC1390-5_ST25.txt, was created on Jan. 8, 2021 and is 53 kB in size. The file can be accessed using Microsoft Word on a computer that uses Windows OS.

BACKGROUND OF THE INVENTION**Field of the Invention**

[0004] The present invention relates generally to the field of medicine, and more specifically to methods and compositions for developing basal forebrain cholinergic neurons (BFCNs) from stem cells, and in particular, BFCNs comprising repaired electrophysiological defects relating to one or more mutations in the presenilin 2 gene (PSEN2), and the use of such BFCNs in cell-based therapies to treat Alzheimer's disease.

Background Information

[0005] Alzheimer's disease (AD) is a progressive disease resulting in senile dementia. Broadly speaking the disease falls into two categories: late onset, which occurs in old age (65+ years) and early onset, which develops well before the senile period, i.e., between 35 and 60 years. In both types of disease, the pathology is the same but the abnormalities tend to be more severe and widespread in cases beginning at an earlier age. The disease is characterized by at least two types of lesions in the brain, senile plaques and neurofibrillary tangles. Senile plaques are areas of disorganized neuropil up to 150 m across with extracellular amyloid deposits at the center visible by microscopic analysis of sections of brain tissue. Neurofibrillary tangles are intracellular deposits of microtubule associated tau protein consisting of two filaments twisted about each other in pairs.

[0006] The principal constituent of the plaques is a peptide termed A β or β -amyloid peptide. A β peptide is an internal fragment of 39-43 amino acids of a precursor protein termed amyloid precursor protein (APP). Several mutations within the APP protein have been correlated with the presence of Alzheimer's disease. Such mutations are thought to cause Alzheimer's disease by increased or altered processing of APP to A β , particularly processing of APP to increased amounts of the long form of A β (i.e., A β 1-42 and A β 1-43). Mutations in other genes, such as the presenilin genes, PSEN1 and PSEN2, are thought indirectly to affect processing of APP to generate increased amounts of long form A β . These observations suggest that A β , and particularly its long form, is a causative element in Alzheimer's disease.

[0007] There are 5 million people currently affected by Alzheimer's disease in the US and, according to the Alzheimer's Association, this number will increase to 16 million by the year 2050. Unfortunately, we only have direct evidence for genetic causation that accounts for 3-5% of these patients. This percentage encompasses the autosomal dominant early onset familial Alzheimer's disease (EOFAD) variants caused by inherited fully penetrant autosomal dominant mutations in the APP, or PSEN1, PSEN2 that constitute the γ -secretase apparatus [87], and changes in their function increases the production of A β 42 oligomers and/or deposition of amyloid plaques.

[0008] After decades studying murine models of AD that do not fully recapitulate the pathophysiology of this disease in the human brain [5, 57, 58], a complementary new concept of AD modeling in vitro has emerged upon the breakthrough by [81] allowing adult human tissue reprogramming into iPSC using defined factors, and their subsequent in vitro differentiation into specific brain cell types.

[0009] BFCNs are one of the most vulnerable neuronal populations whose deterioration explains, in part, the cognitive decline in AD patients. Apart from the evidence for BFCN failure and atrophy, other studies have revealed that human embryonic stem cell-derived BFCNs transplanted into AD mouse models can be associated with improvement in the learning behavior of the implanted mouse [94]. These findings highlight the relevance of iPSC- and ESC-derived BFCNs as not only early clinical indicators but also as a potential strategy for subtype-specific cell-based therapy for AD [39]. In order to move this cell-based therapeutic strategy forward, there has been an urgent need for a refined differentiation protocol to generate human ESC- and/or iPSC-derived BFCNs.

SUMMARY OF THE INVENTION

[0010] The present invention provides a highly reproducible protocol to efficiently derive BFCNs from pluripotent stem cells (PSCs), including embryonic stem cells (ESCs) and induced pluripotent stem cells (iPSCs).

[0011] Accordingly, in one embodiment, the invention provides a method of generating BFCNs. The method includes culturing PSCs in a basal media comprising an inhibitor of transforming growth factor beta (TGF- β) signaling and an activator of sonic hedgehog (Shh) signaling to induce neuroectodermal differentiation. In some aspects, the basal media is a modified mTeSR1 formulation that lacks factors that support pluripotency including basic fibroblast growth factor (bFGF), TGF- β , lithium chloride (Li—Cl), GABA and pipecolic acid. In some aspects, culturing is performed in the presence of dual SMAD inhibitors, such as

SB431542 and LDN193189 along with one or more agonists of smoothened protein, such as smoothened agonist (SAG) and purmorphamine. After about 9, 10, 11 or 12 days of culturing, CD271+ cells are selected and in a neuronal basal medium, such as Brainphys™ to generate neuronal embryoid bodies (NEBs). The neuronal basal medium is optionally supplemented with one or more of B27 supplement, an inhibitor of rho-associated protein kinase (ROCK), nerve growth factor (NGF) and brain derived neurotrophic factor (BDNF). After about 7, 8, 9 or 10 days of culturing the CD271+ cells, the formed NEBs are harvested, dissociated, and plated as monolayer cultures and further cultured in a neuronal basal medium optionally supplemented with B27 supplement, NGF and BDNF. To ensure differentiation into BFCNs, the cultured cells are analyzed for positive expression of Tuj1, MAP2, BF1, Nkx2.1 and p75.

[0012] In another embodiment, the method utilizes iPSCs which may be treated with a gene editing system to repair one or more mutations, such as a mutation of presenilin 1 (PSEN1) or presenilin 2 (PSEN2). In one aspect, the mutation is PSEN2^{N141I}, repair of which restores neuronal excitability in BFCNs.

[0013] Accordingly, in another embodiment, the invention provides a method of treating a disease or disorder in a subject. The method includes administering to a subject a BFCN generated using the culturing method described herein. In certain aspects, the disease or disorder is an amyloidogenic disease, such as systemic amyloidosis, Alzheimer's disease, mature onset diabetes, Parkinson's disease, Huntington's disease, fronto-temporal dementia, and prion-related transmissible spongiform encephalopathies. In embodiments, a BFCN having a mutation that impairs neuronal excitability, such as PSEN2^{N141I}, may be obtained from a subject and used to generate an iPSC, which in turn may be treated with a gene editing system to repair the mutation. The gene edited iPSC is then cultured as described herein to produce BFCNs having restored neuronal excitability which are administered to the subject to treat the disease or disorder. In certain aspects, the disease or disorder is AD.

[0014] In a related embodiment, the invention provides a method of restoring neuronal excitability in BFCNs in a subject. The method includes: a) isolating a BFCN from the subject, wherein the BFCN has a mutation in PSEN2 resulting in impaired neuronal excitability of the BFCN; b) generating an iPSC using the BFCN of (a); c) repairing the PSEN2 mutation in the iPSC; d) culturing the iPSC of (c) using the differentiation protocol described herein to generate a BFCN having the repaired mutation; and e) administering the iPSC of (d) to the subject, thereby restoring neuronal excitability in BFCNs in the subject.

[0015] Also provided is a method of identifying a compound for treatment or prevention of a disease or disorder associated with diminished neuronal excitability in BFCNs. The method includes: a) contacting a BFCN or neuronal embryoid body (NEB) generated using the differentiation protocol described herein with a candidate compound, wherein the BFCN comprises a mutation in PSEN2 resulting in impaired neuronal excitability of the BFCN; and b) detecting neuronal excitability of the BFCN after contact with the candidate compound. An increase in neuronal excitability of the BFCN after contact with the candidate compound identifies the compound as a compound potentially capable of restoring neuronal excitability in BFCNs.

[0016] The invention further provides a BFCN generated using the differentiation protocol as described herein. The BFCN may include a gene edited repair of PSEN2, as well as a detectable marker recombinantly introduced into the BFCN genome.

[0017] The invention also provides a kit for generating a BFCN. The kit includes a culture media having an inhibitor of TGF-β signaling and an activator of Shh signaling. In embodiments, the culture media is a modified mTeSR1 formulation that lacks factors that support pluripotency including bFGF, TGF-β, lithium chloride (Li—Cl), GABA and pipecolic acid. In embodiments, the culture media includes dual SMAD inhibitors, such as SB431542 and LDN193189, along with one or more agonists of smoothened protein, such as smoothened agonist (SAG) and purmorphamine. The kit may also include a neuronal basal medium, such as Brainphys™ optionally supplemented with one or more of B27 supplement, an inhibitor of ROCK, NGF and BDNF. In embodiments, the kit includes reagents for detection of CD271+ cells as well as cells which positively express Tuj1, MAP2, BF1, Nkx2.1 and p75.

BRIEF DESCRIPTION OF THE DRAWINGS

[0018] FIGS. 1A-1E. Overview schematic of basal cholinergic differentiation protocol. (FIG. 1A) Cells are plated and allowed to reach 100% confluence (Day 0), before the initiation of dual SMAD inhibition and the subsequent introduction of ventralizing agents (Day 2). At day 10 the monolayer is dissociated, sorted for p75+ cells, and kept as NEBs until day 19. Then the culture is dissociated again into a monolayer (See Methods for more details). (FIG. 1B) Left panel shows sustained EGFP expression driven by Nkx2.1 induction in NKx2.1-EGFP hESCs upon SHH plus purmorphamine or SAG plus purmorphamine treatment, maintained at Day 14, after removal of treatment at Day 8. Right panel shows Nkx2.1, Lhx8 and BF1 relative gene expression to GAPDH measured by qPCR, in NKx2.1-EGFP cell line in the presence of the indicated ventralizing agents, or unpatterned (UNP) at Day 12. n=3, in technical triplicates. (FIG. 1C) Confocal microscope images of Nestin, Sox2 and DRAQ5 immunostaining in fControl and control lines at Day 11, showing typical neural rosettes (left panel), or Tuj1, Nkx2.1 right and DRAQ5 in the right panel. Images representative of 3 independent experiments. (FIG. 1D) Fluorescence microscope images of immunostained NEB cryosections or dissociated NEBs into a monolayer with the BFCN markers Nkx2.1/Tuj1/p75/BF1/MAP2/ChAT. (FIG. 1E) Dissociated NEBs into a monolayer immunostained at Day 50 with MAP2, ChAT and Hoechst. Fluorescence microscope images the effect of NGF addition to SAG plus purmorphamine treatment alone. Images are representative of at least 3 independent experiments.

[0019] FIGS. 2A-2F. Basal cholinergic markers in PSEN2^{N141I}, neuroprecursors. (FIG. 2A) Table showing the cell lines used. Four iPS lines reprogrammed from fibroblasts were used; two controls (949 and 050643, labelled as fControl and Control, respectively) that do not carry the PSEN2^{N141I} mutation nor the £4 allele; and two AD patients (948 and 950, labelled as AD1 and AD2, respectively) who carry the mutation and the £4 allele. Three of the four iPS lines were family related (fControl, AD1, and AD2). (FIG. 2B) Representative Sanger sequencing chromatograms showing a fragment of exon 5 of PSEN2. Arrow marks site of the missense point mutation Chr1:227,073,304 A>T.

(FIG. 2C) Immunocytochemistry and RT-PCR for early neuronal and basal forebrain markers. n 3, 3 independent experiments with technical triplicates. (FIG. 2D) RTPCR fold changes for TUJ1 and BF1. n 3, 3 independent experiments with technical triplicates. (FIG. 2E) Representative histograms for P75 staining. n>6. (F) A β 40 and A β 42 ELISA quantifications. n 3, 3 independent experiments with technical triplicates. ***, p<0.001. *, p<0.05.

[0020] FIGS. 3A-3B. Neuronal and basal cholinergic markers by immunocytochemistry. (FIG. 3A) Immunostaining for TrkA on DIV 21. (FIG. 3B) Immunostainings for ChAT and vAChT at different magnifications at DIV65; and Tuj1 and MAP2. Images are representative of at least 3 independent experiments.

[0021] FIGS. 4A-4D. CRISPR/Cas9-mediated correction of PSEN2^{N141I} iPS lines. (FIG. 4A) Schematic showing guide RNAs used in the targeting of CRISPR/Cas9, as well as donor ssODNs utilized to introduce wild-type genotype. Sequence identifiers from top to bottom: SEQ ID NOS: 26-31. (FIG. 4B) Left 2 panels show GFP positive HEK293T cells indicating Cas9 system with guide RNA expression, NT refers to non-transfected; right 2 panels show sample of GFP positive iPSCs after lipofection with pCas9-gN141I-GFP vector. (FIG. 4C) Sanger sequencing results from iPSC lines, showing corrections in the N141I mutation. (FIG. 4D) A β 42/40 ratio detected by ELISA in 72 h conditioned media from mutant, control or Cispr-Cas9 corrected BFCNs (DIV 34). n 4, 4 independent experiments with technical triplicates. *, p<0.05; **, p<0.01 Student T-test.

[0022] FIGS. 5A-5B. BFCNs carrying various PSEN mutations are not consistently more susceptible to A β 42 oligomer toxicity. (FIG. 5A) Sample images of BFCNs from the indicated genotypes treated with propidium iodide to visualize cell death in response to 72-h exposure to A β 42 oligomers (5 μ M). (FIG. 5B) % LDH Release recorded from media collected after 72-h exposure. n 3, 3 independent experiments with technical triplicates. *, p<0.05; **, p<0.01 as detected by 2-Way ANOVA Bonferroni post hoc tests.

[0023] FIGS. 6A-6F. NLRP2 inflammasome mRNA levels are over-expressed in some PSEN2^{N141I} cells, but it is not driven by mutation. RT-PCR expression of (FIG. 6A) NLRP2, (FIG. 6B) NLRP3, and (FIG. 6C) ASB9 in cholinergic neuroprecursors. (FIG. 6D) Western blot showing NLRP2, PSEN2 and β -Actin. RT-PCR expression of NLRP2 in Neuroprecursors (FIG. 6E) and BFCNs (F). n 3, 3 independent experiments with technical triplicates, for all panels. ***, p<0.001.

[0024] FIGS. 7A-7B. Electrophysiological and morphological features of BFCN. (FIG. 7A) Top row—compound sodium and potassium currents produced by a voltage protocol shown in bottom row. Current trace produced by a voltage step to -20 mV shown in red. Inset shows first 25 ms of a current produced by a voltage step to -20 mV (scale bars 200 pA, 5 ms). (FIG. 7B) Differential interference contrast image of a patched BFCN recorded in (A). Ninety-four neurons (22 wild-type control, 21 familial control, 18 AD1, 28 AD2 and 5 iAD1_control). Scale bar is 30 μ m.

[0025] FIGS. 8A-8C. Electrophysiological deficits in BFCNs from AD lines. (FIG. 8A) Co-localization of biocytin-labelled neurons with cholinergic markers ChAT and VACHT. Arrows indicate positions of recorded neurons somas, scale bar is 50 μ m. (FIG. 8B) Representative firing patterns of BFCNs produced by a 1 sec negative and positive

square current injection are depicted. A grand total of 94 individual neurons were studied electrophysiologically: 22 wild-type control neurons, 21 familial control neurons, 18 AD1 neurons, 28 AD2 neurons, and 5 iAD1_(CRISPR-corrected) neurons. The experiments on the 94 neurons required days to weeks. On each experimental day, representatives from each genotype were included, with at least three samples from each genotype studied each day. (FIG. 8C) Summary data on maximum number of action potentials that neurons are capable of sustaining (left) and height of a single action potential at rheobase (right) across all conditions. Individual data points are shown as circles, group means are shown as bars. **, p<0.01 Tukey HSD test.

[0026] FIGS. 9A-9D. Intrinsic electrophysiological properties of BFCNs. Summary data on all recorded BFCNs from five groups. Ninety-four neurons (22 wild-type control, 21 familial control, 18 AD1, 28 AD2 and 5 iAD1_control). Histograms show individual values from each neuron (circle) and group means (bars) for membrane resistance (FIG. 9A), capacitance (FIG. 9B), resting potential (FIG. 9C) and rheobase current (FIG. 9D). Statistical significance was tested with ANOVA and Tukey's post hoc comparisons.

[0027] FIGS. 10A-10B. Quality control of iPSC lines. (FIG. 10A) Immunofluorescence shows expression of pluripotency markers SSEA4, Nanog, Tra160 and in 7889 (S)B iPSC line. (FIG. 10B) Three germ layers (endoderm, mesoderm, and ectoderm) from teratomas generated by 7889(S)B iPSC line.

[0028] FIGS. 11A-11B. Amyloid 3 levels in mature BFCNs. (FIG. 11A) Levels of A β 40 on BFCNs (DIV 34). *, P<0.01 vs. other lines in study according to One-Way ANOVA Bonferroni Post-hoc test. (FIG. 11B) Levels of A β 42 on BFCNs (DIV 34). n=3, 3 independent experiments with technical triplicates. *, P<0.01 based on Student's T-test.

DETAILED DESCRIPTION OF THE INVENTION

[0029] The present invention is based on the discovery of a robust, fast, and reproducible differentiation protocol to generate BFCNs from PSCs using a chemically defined medium.

[0030] The following is a detailed description of the invention provided to aid those skilled in the art in practicing the present invention. Those of ordinary skill in the art may make modifications and variations in the embodiments described herein without departing from the spirit or scope of the present invention. Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The terminology used in the description of the invention herein is for describing particular embodiments only and is not intended to be limiting of the invention. All publications, patent applications, patents, figures and other references mentioned herein are expressly incorporated by reference in their entirety.

[0031] Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and described the methods and/or materials in connection with which the publications are cited.

[0032] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The following references, the entire disclosures of which are incorporated herein by reference, provide one of skill with a general definition of many of the terms (unless defined otherwise herein) used in this invention: Singleton et al., Dictionary of Microbiology and Molecular Biology (2nd ed. 1994); The Cambridge Dictionary of Science and Technology (Walker ed., 1988); The Glossary of Genetics, 5th Ed., R. Rieger et al. (eds.), Springer Verlag (1991); and Hale & Marham, the Harper Collins Dictionary of Biology (1991). Generally, the procedures of molecular biology methods described or inherent herein and the like are common methods used in the art. Such standard techniques can be found in reference manuals such as for example Sambrook et al. (2000, Molecular Cloning—A Laboratory Manual, Third Edition, Cold Spring Harbor Laboratories); and Ausubel et al. (1994, Current Protocols in Molecular Biology, John Wiley & Sons, New-York).

[0033] The terminology used in the description is for describing particular embodiments only and is not intended to be limiting of the invention. Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise (such as in the case of a group containing a number of carbon atoms in which case each carbon atom number falling within the range is provided), between the upper and lower limit of that range and any other stated or intervening value in that stated range is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges is also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either both of those included limits are also included in the invention.

[0034] The following terms are used to describe the present invention. In instances where a term is not specifically defined herein, that term is given an art-recognized meaning by those of ordinary skill applying that term in context to its use in describing the present invention.

[0035] The articles “a” and “an” as used herein and in the appended claims are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article unless the context clearly indicates otherwise. By way of example, “an element” means one element or more than one element.

[0036] The phrase “and/or,” as used herein in the specification and in the claims, should be understood to mean “either or both” of the elements so conjoined, i.e., elements that are conjunctively present in some cases and disjunctively present in other cases. Multiple elements listed with “and/or” should be construed in the same fashion, i.e., “one or more” of the elements so conjoined. Other elements may optionally be present other than the elements specifically identified by the “and/or” clause, whether related or unrelated to those elements specifically identified. Thus, as a non-limiting example, a reference to “A and/or B”, when used in conjunction with open-ended language such as “comprising” can refer, in one embodiment, to A only (optionally including elements other than B); in another embodiment, to B only (optionally including elements other

than A); in yet another embodiment, to both A and B (optionally including other elements).

[0037] As used herein in the specification and in the claims, “or” should be understood to have the same meaning as “and/or” as defined above. For example, when separating items in a list, “or” or “and/or” shall be interpreted as being inclusive, i.e., the inclusion of at least one, but also including more than one, of a number or list of elements, and, optionally, additional unlisted items. Only terms clearly indicated to the contrary, such as “only one of” or “exactly one of,” or, when used in the claims, “consisting of,” will refer to the inclusion of exactly one element of a number or list of elements. In general, the term “or” as used herein shall only be interpreted as indicating exclusive alternatives (i.e., “one or the other but not both”) when preceded by terms of exclusivity, such as “either,” “one of,” “only one of,” or “exactly one of.”

[0038] As used herein in the specification and in the claims, the phrase “at least one,” in reference to a list of one or more elements, should be understood to mean at least one element selected from anyone or more of the elements in the list of elements, but not necessarily including at least one of each and every element specifically listed within the list of elements and not excluding any combinations of elements in the list of elements. This definition also allows that elements may optionally be present other than the elements specifically identified within the list of elements to which the phrase “at least one” refers, whether related or unrelated to those elements specifically identified. Thus, as a nonlimiting example, “at least one of A and B” (or, equivalently, “at least one of A or B,” or, equivalently “at least one of A and/or B”) can refer, in one embodiment, to at least one, optionally including more than one, A, with no B present (and optionally including elements other than B); in another embodiment, to at least one, optionally including more than one, B, with no A present (and optionally including elements other than A); in yet another embodiment, to at least one, optionally including more than one, A, and at least one, optionally including more than one, B (and optionally including other elements).

[0039] It should also be understood that, in certain methods described herein that include more than one step or act, the order of the steps or acts of the method is not necessarily limited to the order in which the steps or acts of the method are recited unless the context indicates otherwise.

[0040] The term “PSEN2 gene” refers herein to a gene that encodes a PSEN2 polypeptide. The PSEN2 gene is represented by NCBI Reference Sequence: NC_000001.11 (SEQ ID NO: 1) as well as known orthologs. The term “PSEN2 polypeptide” refers herein to a polypeptide that is represented by NCBI Reference Sequence: NP_000438.2 (SEQ ID NO: 2) as well as known orthologs.

[0041] The term “amyloidogenic disease” includes any disease associated with (or caused by) the formation or deposition of insoluble amyloid fibrils. Exemplary amyloidogenic diseases include, but are not limited to systemic amyloidosis, Alzheimer’s disease, mature onset diabetes, Parkinson’s disease, Huntington’s disease, fronto-temporal dementia, and the prion-related transmissible spongiform encephalopathies (kuru and Creutzfeldt-Jacob disease in humans and scrapie and BSE in sheep and cattle, respectively). Different amyloidogenic diseases are defined or characterized by the nature of the polypeptide component of the fibrils deposited. For example, in subjects or patients

having Alzheimer's disease, β -amyloid protein (e.g., wild-type, variant, or truncated β -amyloid protein) is the characterizing polypeptide component of the amyloid deposit. Accordingly, Alzheimer's disease is an example of a "disease characterized by deposits of $A\beta$ " or a "disease associated with deposits of $A\beta$ ", e.g., in the brain of a subject or patient. The terms " β -amyloid protein", " β -amyloid peptide", " β -amyloid", " $A\beta$ " and " $A\beta$ peptide" are used interchangeably herein.

[0042] The term "patient" or "subject" is used throughout the specification to describe an animal, preferably a human or a domesticated animal, to whom treatment, including prophylactic treatment, with the compositions according to the present disclosure is provided. For treatment of conditions or disease states which are specific for a specific animal such as a human patient, the term patient refers to that specific animal, including a domesticated animal such as a dog or cat or a farm animal such as a horse, cow, sheep, etc. In general, in the present disclosure, the term patient refers to a human patient unless otherwise stated or implied from the context of the use of the term.

[0043] The terms " β -amyloid protein", " β -amyloid peptide", " β -amyloid", " $A\beta$ " and "AD peptide" are used interchangeably herein. $A\beta$ peptide (e.g., $A\beta$ 39, $A\beta$ 40, $A\beta$ 41, $A\beta$ 42 and $A\beta$ 43) is about 4-kDa internal fragment of 39-43 amino acids of APP. $A\beta$ 40, for example, consists of residues 672-711 of APP and $A\beta$ 42 consists of residues 672-713 of APP. $A\beta$ peptides include peptides resulting from secretase cleavage of APP and synthetic peptides having the same or essentially the same sequence as the cleavage products. $A\beta$ peptides can be derived from a variety of sources, for example, tissues, cell lines, or body fluids (e.g., sera or cerebrospinal fluid). For example, an $A\beta$ can be derived from APP-expressing cells such as Chinese hamster ovary (CHO) cells stably transfected with APP_{717V}F, as described, for example, in Walsh et al. (2002), Nature, 416, pp 535-539. An $A\beta$ preparation can be derived from tissue sources using methods previously described (see, e.g., Johnson-Wood et al. (1997), Proc. Natl. Acad. Sci. USA 94:1550). Alternatively, $A\beta$ peptides can be synthesized using methods which are well known to those in the art. See, for example, Fields et al., Synthetic Peptides: A User's Guide, ed. Grant, W. H. Freeman & Co., New York, N.Y., 1992, p 77). Hence, peptides can be synthesized using the automated Merrifield techniques of solid phase synthesis with the α -amino group protected by either t-Boc or F-moc chemistry using side chain protected amino acids on, for example, an Applied Biosystems Peptide Synthesizer Model 430A or 431. Longer peptide antigens can be synthesized using well known recombinant DNA techniques. For example, a polynucleotide encoding the peptide or fusion peptide can be synthesized or molecularly cloned and inserted in a suitable expression vector for the transfection and heterologous expression by a suitable host cell. A B peptide also refers to related $A\beta$ sequences that results from mutations in the $A\beta$ region of the normal gene.

[0044] As used herein the phrase "substantially pure" refers to a population of cells wherein at least 95% of the cells have the recited phenotype. In all embodiments that refer to a "substantially pure" cell population, alternative embodiments in which the cell populations have a lower or higher level of purity are also contemplated. For example, in some embodiments, instead of a given cell population being "substantially pure" the cell population may be one in which

at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% of the cells, or 100% of the cells, have the recited phenotype.

[0045] The terms "co-administration", "co-administered" and "co-administering" or "combination therapy" refer to both concurrent administration (administration of two or more agents at the same time) and time varied administration (administration of one or more agents at a time different from that of the administration of an additional agent or agents), as long as the agents are present in the area to be treated to some extent, preferably at effective amounts, at the same time.

[0046] The term "therapeutically effective amount" means the amount required to achieve a therapeutic effect. The therapeutic effect could be any therapeutic effect ranging from prevention, symptom amelioration, symptom treatment, to disease termination or cure, e.g., the treatment of Alzheimer's disease or an associated condition.

[0047] As used herein, the term "administering" is meant to refer to a means of providing the composition to the subject in a manner that results in the composition being inside the subject's body. Such an administration can be by any route including, without limitation, subcutaneous, intradermal, intravenous, intra-arterial, intraperitoneal, and intramuscular.

[0048] The term "effective" is used to describe an amount of a compound, composition or component which, when used within the context of its intended use, effects an intended result. The term effective subsumes all other effective amount or effective concentration terms, which are otherwise described or used in the present application.

[0049] As used herein, the term "comprising" is intended to mean that the compositions and methods include the recited elements, but do not exclude other elements. "Consisting essentially of", when used to define compositions and methods, shall mean excluding other elements of any essential significance to the combination. Thus, a composition consisting essentially of the elements as defined herein would not exclude trace contaminants from the isolation and purification method and pharmaceutically acceptable carriers, such as phosphate buffered saline, preservatives, and the like. "Consisting of" shall mean excluding more than trace elements of other ingredients and substantial method steps for administering the compositions of this invention. Embodiments defined by each of these transition terms are within the scope of this invention.

[0050] Ranges provided herein are understood to be shorthand for all of the values within the range. For example, a range of 1 to 50 is understood to include any number, combination of numbers, or sub-range from the group consisting of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, and 50.

[0051] As used herein, "kits" are understood to contain at least the non-standard laboratory reagents of the invention and one or more non-standard laboratory reagents for use in the methods of the invention.

[0052] The term "obtaining" is understood herein as manufacturing, purchasing, or otherwise coming into possession of.

[0053] As used herein, the term "treated," "treating" or "treatment" includes the diminishment or alleviation of at least one symptom associated or caused by the state, disor-

der or disease being treated. A subject that has been treated can exhibit a partial or total alleviation of symptoms (for example, Alzheimer's disease or associated condition), or symptoms can remain static following treatment according to the invention. The term "treatment" is intended to encompass prophylaxis, therapy and cure.

[0054] As used herein, the term "control" refers to a sample or standard used for comparison with an experimental sample. In some embodiments, the control is a sample obtained from a healthy patient. In other embodiments, the control is a historical control or standard reference value or range of values (such as a previously tested sample, subject, or group of samples or subjects).

Methods

[0055] BFCNs are believed to be one of the first cell types to be affected in all forms of AD, and their dysfunction is clinically correlated with impaired short-term memory formation and retrieval. As detailed in the Example of this disclosure, the inventors present an optimized *in vitro* protocol to generate human BFCNs from iPSCs, using cell lines from PSEN2 mutation carriers and controls. Cell lines harboring the PSEN2^{N141T} mutation displayed an increase in the Aβ42/40 in iPSC-derived BFCNs. Neurons derived from PSEN2^{N141T}, lines generated fewer maximum number of spikes in response to a square depolarizing current injection. The height of the first action potential at rheobase current injection was also significantly decreased in PSEN2^{N141T} BFCNs. CRISPR/Cas9 correction of the PSEN2 point mutation abolished the electrophysiological deficit, restoring both the maximal number of spikes and spike height to the levels recorded in controls. Increased Aβ42/40 was also normalized following CRISPR/Cas-mediated correction of the PSEN2^{N141T} mutation. The genome editing data set forth herein confirms the robust consistency of mutation-related changes in Aβ42/40 ratio while also showing a PSEN2-mutation-related alteration in electrophysiology.

[0056] Accordingly, in one embodiment, the invention provides a method of generating BFCNs. The method may first include preparing PSC colonies. PSCs are seeded (plated) at low density and grown in an adherent culture for about 1-2 days. "Low density" means about 8,000 to about 11,000 cells/cm². Cells are preferably seeded at about 9,500 to about 10,500 cells/cm², more preferably at about 10,000 cells/cm². After about 1-2 days (or greater, i.e., 3, 4, 5, 6, 7, 8, 9, 10 or more), the PSCs form colonies, which are preferably about 75 μm to about 300 μm in diameter, more preferably about 100 μm to about 250 μm in diameter.

[0057] The term "PSCs" has its usual meaning in the art, i.e., self-replicating cells that have the ability to develop into endoderm, ectoderm, and mesoderm cells. Preferably, PSCs are hPSCs. PSCs include ESCs and iPSCs, preferably hESCs and hiPSCs. PSCs can be seeded on a surface comprising a matrix, such as a gel or basement membrane matrix. A preferable matrix is the protein mixture secreted by Engelbreth-Holm-Swarm (EHS) mouse sarcoma cells, sold under trade names including MATRIGEL®, CULTREX®, and GELTREX®. Other suitable matrices include, without limitation, collagen, fibronectin, gelatin, laminin, poly-lysine, vitronectin, and combinations thereof.

[0058] In some embodiments, media suitable for use in maintaining pluripotent stem cells is used. In embodiments such a medium is mTeSR1 medium from Stem Cell Technologies. However, one of skill in the art will recognize that

there are several other types of media that are equivalent to mTeSR medium in terms of their suitability for use in maintaining pluripotent stem cells, any of which could be used. Typically such media will contain one or more pluripotency factors to facilitate the maintenance of cells in a pluripotent state. The composition of mTeSR1 medium is known in the art and described in, for example, Ludwig et al., 2006 (Nat Methods. 2006 August; 3(8):637-46; "Feeder-Independent Culture of Human Embryonic Stem Cells"), the contents of which are hereby incorporated by reference.

[0059] The pluripotent stem cells used in the method of the invention can be any suitable type of pluripotent stem cells. Where iPSCs are used, such cells may have been "reprogrammed" to the pluripotent state from a non-pluripotent state using any suitable means known in the art, including, but not limited to, modified RNA-based methods, Sendai virus based methods, and the like. Furthermore, such cells may have been reprogrammed to the pluripotent state using any suitable cocktail of reprogramming factors known in the art.

[0060] In one embodiment, after PSCs are prepared and grown to confluence in, for example mTeSR1 media, the method includes culturing the PSCs in a basal media comprising an inhibitor of transforming growth factor beta (TGF-β) signaling and an activator of sonic hedgehog (Shh) signaling to induce neuroectodermal differentiation. The basal media utilized is a modified mTeSR1 medium, which is a variant of mTeSR1 medium (sometimes referred to herein as "mTeSR1 Custom" medium) that does not comprise lithium chloride, GABA, pipecolic acid, bFGF or TGFβ1. Inhibitors of TGFβ signaling include, for example, one or more of SB431542, GW788388, LDN193189, LY2109761, LY2157299, and LY364947. Activators of Shh signaling include agonists of Smoothened, such as Smoothened Agonist (SAG; 3-chloro-N-[(1r,4r)-4-(methylamino)cyclohexyl]-N-[3-(pyridin-4-yl)benzyl]benzo[b]thiophene-2-carboxamide) and purmorphamine.

[0061] After about 6, 7, 8, 9, 10, 11 or 12 days (or greater, i.e., 15, 16, 17, 18, 19, 20, 25, 30 or more) of culturing, CD271+ cells are selected and cultured in a neuronal basal medium, such as Brainphys™ to generate neuronal embryoid bodies (NEBs). The neuronal basal medium is optionally supplemented with one or more of B27 supplement, an inhibitor of rho-associated protein kinase (ROCK), nerve growth factor (NGF) and brain derived neurotrophic factor (BDNF). Inhibitors of ROCK include, for example, GSK269962, GSK429286, H-1152, HA-1077, RKI-1447, thiazovivin, Y-27632, or derivatives thereof.

[0062] To select for CD271+ cells, overconfluent cells are lifted from the culture surface and purified by FACS and re-plated. This process allows for the formation of cell aggregates or spheres, also referred to herein as NEBs. For purposes of the present invention, the terms "NEB," "aggregate" and "sphere" are used interchangeably and refer to a multicellular three-dimensional structure, preferably, but not necessarily, of at least about 100 cells.

[0063] Lifting can be performed mechanically, with a cell scraper or other suitable implement, or chemically. Chemical lifting can be achieved using a proteolytic enzyme, for example, collagenase, trypsin, trypsin-like proteinase, recombinant enzymes, such as that sold under the trade name TRYPLE™, naturally derived enzymes, such as that sold under the trade name ACCUTASE™, and combinations thereof. Chemical lifting can also be done using a chelator,

such as EDTA, or a compound such as urea. Mechanical lifting or detachment offers the advantage of minimal cell death, however it produces aggregates of variable size, thus suitable spheres need to be selected through a manual picking process. Good spheres are defined as those having a round-shape, golden/brown color, with darker core and with a diameter between about 300 μm and about 800 μm . Detaching the cells using chemical methods, such as enzymatic digestion predominantly produces spheres that are appropriate for further culture. Therefore manual picking of spheres is not required, and the detachment steps can be adapted for automation and used in high throughput studies. However, enzymatic digestion increases cell death, resulting in a lower number of spheres.

[0064] After about 5, 6, 7, 8, 9 or 10 days (or greater, i.e., 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30 or more) of culturing selected CD271+ cells, the formed NEBs are harvested, dissociated, and plated as monolayer cultures and further cultured in a neuronal basal medium, such as BrainphysTM, optionally supplemented with B27 supplement, NGF and BDNF. The surface on which the cells are plated and cultured can comprise an extracellular matrix protein (e.g., collagen, fibronectin, laminin) and/or a positively charged poly-amino acid (e.g., poly-arginine, poly-lysine, poly-ornithine). Preferably the surface comprises laminin and/or poly-omithine.

[0065] To ensure differentiation into BFCNs, the cultured cells are analyzed for positive expression of Tuj1, MAP2, BF1, Nkx2.1 and p75.

[0066] Many of the embodiments of the present invention involve certain factors to be used in (or excluded from) the compositions and methods described herein, for example as media supplements. These include, but are not limited to, bFGF, GABA, pipecolic acid, lithium chloride, TGF- β , NGF and BDNF. Each of these factors is well known in the art, including the full names of each of these factors in the cases where acronyms or other abbreviations are used. Furthermore, all of these factors are available to the public from multiple sources, including commercial sources. Exemplary amounts/concentrations for use of each of these factors in the methods and compositions of the present invention are provided in the Examples section of this patent disclosure. For all embodiments where specified amounts are referred to, amounts that are “about” the specified amount are also intended. Furthermore, one of skill in the art will recognize that in some situations further deviations from the specified amounts can be used, and will be able to determine how much of each factor to use by performing routine testing, optimization, dose-response studies, and the like, for example to reduce or increase the specified amounts, as long as the amounts used still achieve the stated effect, e.g., the stated differentiation effect. For example, in some embodiments specified amounts of the specified agents may be reduced to 10%, or 20%, or 30%, or 40%, or 50%, or 60%, or 70%, or 80%, or 90% of the stated amounts. Similarly, in some embodiments the specified amounts of the specified agents may be increased by 10%, by 20%, by 30%, by 40%, by 50%, by 60%, by 70%, by 80%, by 90%, by 100%, by 150%, by 200%, by 300%, by 400%, or by 500% of the stated amounts. Similarly, where specified factors are referred to, one of skill in the art will recognize that analogs, variants, or derivatives of such factors can also be used as long as the analogs, variants, or derivatives have the same general function/activity as the specified factors.

[0067] As discussed herein, the inventors have discovered that mutation of PSEN2 results in impaired neuronal excitability in BFCNs. As discussed in the Example herein the inventors observed significant mutation-related, editing-reversible differences in excitability of BFCNs via repair of PSEN2^{N141T} mutation which restored neuronal excitability. Accordingly, the method of the invention may utilize iPSCs which may be treated with a gene editing system to repair one or more mutations, such as a mutation of presenilin 1 (PSEN1) or presenilin 2 (PSEN2). In one embodiment, the mutation is PSEN2^{N141T}, repair of which restores neuronal excitability in BFCNs.

[0068] As used herein the term “gene editing” or “genome editing” refers to a type of genetic engineering in which DNA is inserted, replaced, or removed from a target DNA, e.g., the genome of a cell, using one or more nucleases and/or nickases. The nucleases create specific double-strand breaks (DSBs) at desired locations in the genome, and harness the cell’s endogenous mechanisms to repair the induced break by homology-directed repair (HDR) (e.g., homologous recombination) or by nonhomologous end joining (NHEJ). The nickases create specific single-strand breaks at desired locations in the genome. In one non-limiting example, two nickases can be used to create two single-strand breaks on opposite strands of a target DNA, thereby generating a blunt or a sticky end. Any suitable nuclease can be introduced into a cell to induce genome editing of a target DNA sequence including, but not limited to, CRISPR-associated protein (Cas) nucleases, zinc finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs), meganucleases, other endo- or exo-nucleases, variants thereof, fragments thereof, and combinations thereof. In particular embodiments, nuclease-mediated genome editing of a target DNA sequence (e.g., a safe harbor gene) by homology-directed repair (HDR) (e.g., homologous recombination) is used for generating a genetically modified human neural stem cell in accordance with the methods described herein.

[0069] The term “DNA nuclease” refers to an enzyme capable of cleaving the phosphodiester bonds between the nucleotide subunits of DNA, and may be an endonuclease or an exonuclease. According to the invention, the DNA nuclease may be an engineered (e.g., programmable or targetable) DNA nuclease which can be used to induce genome editing of a target DNA sequence such as a safe harbor gene. Any suitable DNA nuclease can be used including, but not limited to, CRISPR-associated protein (Cas) nucleases, zinc finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs), meganucleases, other endo- or exo-nucleases, variants thereof, fragments thereof, and combinations thereof.

[0070] In various embodiments of the invention, a gene editing system utilizes a DNA nuclease to edit a gene and repair a mutation. In specific embodiments, a mutation is repaired using one or more of the following gene editing system: CRISPR/Cas system, Cre/Lox system, TALEN system and homologous recombination.

[0071] The differentiation protocol of the present invention may be utilized to treat a disease or disorder in a subject. The method includes administering to a subject a BFCN generated using the culturing method described herein. In various embodiments, the disease or disorder is an amyloidogenic disease, such as systemic amyloidosis, Alzheimer’s disease, mature onset diabetes, Parkinson’s disease,

Huntington's disease, fronto-temporal dementia, and prion-related transmissible spongiform encephalopathies. In embodiments, a BFCN having a mutation that impairs neuronal excitability, such PSEN2^{N141I}, may be obtained from a subject and used to generate an iPSC, which in turn may be treated with a gene editing system to repair the mutation. The gene edited iPSC is then cultured as described herein to produce BFCNs having restored neuronal excitability which are administered to the subject to treat the disease or disorder.

[0072] In a related manner, neuronal excitability in BFCNs in a subject may be restored. This method includes: a) isolating a BFCN from the subject, wherein the BFCN has a genetic mutation resulting in impaired neuronal excitability of the BFCN; b) generating an iPSC using the BFCN of (a); c) repairing the mutation in the iPSC; d) culturing the iPSC of (c) using the differentiation protocol described herein to generate a BFCN having the repaired mutation; and e) administering the iPSC of (d) to the subject, thereby restoring neuronal excitability in BFCNs in the subject. In embodiments, the mutations is of PSEN1 or PSEN2, such as PSEN2^{N141I}.

[0073] The invention also encompasses BFCNs generated using the differentiation protocol as described herein. The BFCN may include a gene edited repair of PSEN2, as well as a detectable marker recombinantly introduced into the BFCN genome. In some embodiments, BFCNs are differentiated from PSCs, and in such embodiments, the PSCs can be iPSCs. The iPSCs can be derived from a somatic cell of a subject. In one aspect, the subject has an amyloidogenic disease or disorder.

[0074] Alongside its potential for autologous cell transplantation, iPSC technology is emerging as a tool for developing new drugs and gaining insight into disease pathogenesis. Han, S. S. W. et al., *Neuron*. 70:626-644 (2011). The methods and cells of the invention can aid the development of high-throughput in vitro screens for compounds that promote restoration of neuronal excitability. To that end, the disclosure provides a method of identifying a compound that can be used for treatment or prevention of a disease or disorder associated with diminished neuronal excitability in BFCNs. The method includes: a) contacting a BFCN or neuronal embryoid body (NEB) generated using the differentiation protocol described herein with a candidate compound, wherein the BFCN comprises a mutation in PSEN2 resulting in impaired neuronal excitability of the BFCN; and b) detecting neuronal excitability of the BFCN after contact with the candidate compound. A beneficial effect on neuronal excitability is evident in partial or complete restoration of neuronal excitability, thereby being indicative of a candidate therapeutic agent for treating a disease or disorder associated with diminished neuronal excitability in BFCNs, such as an amyloidogenic disease or disorder. Preferably, the method is conducted in a high-throughput format.

[0075] The invention also provides a model system for a neurological disease, preferably a disease or disorder associated with diminished neuronal excitability in BFCNs, such as an amyloidogenic disease or disorder. In one aspect, the model system comprises a BFCN differentiated from an iPSC derived from a subject having a disease or disorder associated with diminished neuronal excitability in BFCNs, such as an amyloidogenic disease or disorder. The model system can further comprise a non-human mammal into which the myelin-producing cell has been transplanted. In

one embodiment, the non-human mammal is a mouse or a rat. Model systems provided by the invention can be used to study diseases or disorders, including understanding underlying mechanisms and defining therapeutic targets.

[0076] In some embodiments the present invention provides tissue culture media, tissue culture media supplements, and various kits useful in performing the various methods described herein.

[0077] In one embodiment, the invention provides a kit for generating a BFCN via the differentiation protocol of the invention. The kit includes a culture media having an inhibitor of TGF-β signaling and an activator of Shh signaling. In embodiments, the culture media is a modified mTeSR1 formulation that lacks factors that support pluripotency including bFGF, TGF-β, lithium chloride (Li—Cl), GABA and pipecolic acid. In embodiments, the culture media includes dual SMAD inhibitors, such as SB431542 and LDN193189, along with one or more agonists of smoothened protein, such as smoothened agonist (SAG) and purmorphamine.

[0078] The kit may also include an additional neuronal basal medium, such as Brainphys™ optionally supplemented with one or more of B27 supplement, an inhibitor of ROCK, NGF and BDNF.

[0079] The kit may also include reagents for detection and isolation of CD271+ cells via FACS as well as detection of expression of TuJ1, MAP2, BF1, Nkx2.1 and p75.

[0080] The kit may optionally comprise instructions for use, one or more containers, one or more antibodies, or any combination thereof. A label typically accompanies the kit, and includes any writing or recorded material, which may be electronic or computer readable form (e.g., disk, optical disc, memory chip, or tape) providing instructions or other information for use of the kit contents.

[0081] The following example is provided to further illustrate the advantages and features of the present invention, but it is not intended to limit the scope of the invention. While this example is typical of those that might be used, other procedures, methodologies, or techniques known to those skilled in the art may alternatively be used.

Example I

CRISPR/Cas9-Correctable Mutation-Related Molecular and Physiological Phenotypes in iPSC-Derived Alzheimer's PSEN2^{N141I} Mutation

[0082] Basal forebrain cholinergic neurons (BFCNs) are believed to be one of the first cell types to be affected in all forms of AD, and their dysfunction is clinically correlated with impaired short-term memory formation and retrieval. We present an optimized in vitro protocol to generate human BFCNs from iPSCs, using cell lines from presenilin 2 (PSEN2) mutation carriers and controls. As expected, cell lines harboring the PSEN2^{N141I} mutation displayed an increase in the Aβ42/40 in iPSC-derived BFCNs. Neurons derived from PSEN2^{N141I} lines generated fewer maximum number of spikes in response to a square depolarizing current injection. The height of the first action potential at rheobase current injection was also significantly decreased in PSEN2^{N141I} BFCNs. CRISPR/Cas9 correction of the PSEN2 point mutation abolished the electrophysiological deficit, restoring both the maximal number of spikes and spike height to the levels recorded in controls. Increased Aβ42/40 was also normalized following CRISPR/Cas-me-

diated correction of the PSEN2^{N141T} mutation. The genome editing data confirms the robust consistency of mutation-related changes in Aβ42/40 ratio while also showing a PSEN2-mutation-related alteration in electrophysiology.

[0083] The “amyloid hypothesis” is one of the most popular formulations for the pathogenesis of Alzheimer’s disease (AD). Recent examples of clinicopathological and/or clinico-radiological dissociation have led to the consideration of alternative models in order to explain, respectively, why neuropathological AD is not always associated with dementia [24], and why about one-third of patients with clinical AD have negative amyloid brain scans [40]. It has been proposed that clinical AD can be caused by one of several “feed-forward” scenarios linking amyloidosis, tauopathy, neuroinflammation, and neurodegeneration [22]. Mutations in the gene encoding presenilin 2 (PSEN2) are associated with autosomal dominant early onset familial Alzheimer’s disease (EOFAD). The linkage of a locus on human chromosome 1q31-42 linked to EOFAD led to the identification of the PSEN2^{N141T} point mutation in the Volga German kindreds in 1995 [43]. This mutation causes elevation in the Aβ42/43/40 ratio, thereby promoting assembly of Aβ oligomers and fibrils [83].

[0084] In considering the progression of AD, human basal forebrain cholinergic neurons (BFCNs) are one of the first cell types whose dysfunction underlies the early loss of short-term memory recall in all forms of AD. The “cholinergic hypothesis of AD” was formulated in the mid-1970s [6, 20, 61], and the discoveries of reduced acetylcholine release from neurons of the nucleus basalis of Meynert confirmed the presence of a presynaptic cholinergic deficit in the basal forebrain of AD patients [1, 71]. Based on those observations, acetylcholinesterase inhibitors were developed and continue as the most widely used symptomatic treatments for AD [21, 28, 33, 82]. Eventually, post-mortem brain biochemical and volumetric studies at different stages of the disease identified several other regions of the brain that were also affected early in the course of AD [63]. These studies have traditionally focused on the hippocampus and cortex, but more recently, attention has begun shifting back to the basal forebrain and adding other areas, such as the striatum [27, 62]. The latest analyses suggest that cholinergic basal forebrain volume measurement may be a better predictor of the transition from MCI to AD than the previous standard, hippocampal volume [10].

[0085] We previously reported the generation of iPSC-derived neurons from banked fibroblasts from subjects harboring PSEN1^{A246E} and PSEN1^{M146L} mutations [77]. In characterizing the gene expression profiles from these iPSC-derived neurons, we observed an unexpected association of elevated expression of the inflammasome gene NLRP2 in undifferentiated PSEN1 mutant iPSCs and their and neuronally differentiated progeny [77]. This led us to examine NLRP2 expression in our PSEN2 mutant lines and employ CRISPR/Cas9 [15] to investigate if activation of the inflammasome was tightly linked to the pathogenic mutation in PSEN2. While we did not find altered expression of NLRP2 in gene-corrected PSEN2 lines, we observed significant mutation-related, editing-reversible differences in excitability of BFCNs.

Materials and Methods

Generation and Maintenance of iPSC Lines

[0086] 7889(s)B, 050643 (Control), 948 (AD1), 949 (fControl), and 950 (AD2) iPSC lines were obtained via the NYSCF Repository following the guidelines from [60]. The derivation and characterization of Nkx2.1-GFP ESC line was previously published [30]. ES and iPS cell lines were expanded and maintained in serum-free mTeSR1™ media (Stem Cell Technologies). Cells were lifted using Strem-Pro™ Accutase (ThermoFisher) and media was supplemented with 10 µM ROCK inhibitor (Y27632, Stemgent) during cell passaging.

[0087] For all studies in this paper, cell lines underwent at least 3 independent differentiations from the iPSC stage to the mature neuron stage. Data were routinely compared across these independently derived genotype-identical neurons (or in some cases neuronal precursors), and if comparable results were obtained across independently genotype-identical derived cells, they were considered to be qualified representatives of their genotype and so were passed along for genotype-specific experimentation.

Aβ42 Oligomer Preparation

[0088] Aβ42 oligomers were prepared as previously reported [23, 78]. Briefly, we dissolved 1 mg of Aβ42 (American Peptide Company) in 1,1,1,3,3-hexafluoro-2-propanol (HFIP) (Sigma). This preparation was aliquoted and dried using a SpeedVac™ centrifuge. The pellet was then resuspended in DMSO to obtain a 5 mM solution which was sonicated in a water bath for 10 min. From here aliquots were stored at -20 C and used within 2 weeks by diluting with 100 µl of PBS and leaving for 12 h at 4° C. in order for oligomerization to proceed. This final solution was diluted 1:16 in cell media for studies, allowing cells to be exposed to 5 µM of Aβ42 oligomers. Control wells were diluted with 1:16 PBS. Cells were exposed to oligomers or PBS without media change for a period of 3 days.

Cell Death Assays

[0089] Cells were assayed in a 96-well plate format. Oligomer or vehicle solutions were added to media and allowed to incubate for a period of 3 days. Media was then collected and assayed using a lactate dehydrogenase toxicity assay (Thermo Fisher Scientific). 50 µl of media and an equal amount of reaction mix buffer were incubated for a period of 30 min. An additional set of wells per experiment were treated with 2% Triton™ X-100 for a 5-min period in order to lyse all cells, and media from these wells was also collected and incubated as described. After incubation absorbance was recorded at 490 nm and 680 nm, signal and background absorbance, respectively. Signal values were subtracted from background, and values were adapted to the total LDH content as determined by Triton X-100 treated wells. Propidium iodide (Thermo Fisher Scientific) was added to cell media for a 1 µM final concentration and allowed to incubate for 5 min. Cells were then washed twice with media and imaged. Images were captured using CELIGOTM image cytometer and accompanying software (Nexcelom Bioscience). Each biological variable was assessed in technical triplicates within each designated “Experiment”, and each designated “Experiment” was performed in at least three complete “start to finish” iterations.

Differentiation of Basal Forebrain Cholinergic Neurons from iPS and ES Cells

[0090] Human ES or iPSC were plated as single cells after chemical dissociation using Accutase™ (Sigma-Aldrich) into Cul-Trex™ (Trevigen) coated plates, at a density of 4-8×10⁵ cells per well in 6-well plates or petri dishes and adapting cell numbers. Cells were initially maintained in mTeSR1™ media (Stem Cell Technologies) until reaching full confluence. On “day 0” of differentiation, media was replaced by Custom mTeSR1™ media (Stem Cell Technologies) lacking factors promoting pluripotency i.e., bFGF, TGF-β, Li—Cl, GABA and pipecolic acid. The addition of dual SMAD inhibitors (SB431542 10 μM plus LDN193189 250 nM, Selleckchem) at day 0 drives cells towards neuroectoderm specification. At day 2 of differentiation, media was replaced by Custom mTeSR1 containing dual SMAD inhibitors plus two ventralizing agents: SAG at 500 nM (R&D) and Purmorphamine at 2 μM (Stemgent™). Cells were fed every 2 days with this media until day 9, when media was progressively switched to Brainphys™ media (Stemcell Technologies) supplemented with B27 (Life Technologies) [3]. Neural progenitors were harvested at day 11 using Accutase, p75+(CD271) NPCs were purified by FACS and plated at a density of 80,000 cells per well into non-adherent 96 well V-bottom plates in Brainphys™+B27 supplemented with 10 μM ROCK inhibitor (Y27632, Stemgent), Nerve Growth Factor, NGF (Alamone labs, 50 ng/mL) and Brain Derived Neurotrophic Factor, BDNF (R&D, 50 ng/mL). Cells were allowed to aggregate and form Neuronal Embryoid Bodies (NEBs) and were fed every other day until day 19. At day 19 NEBs were dissociated using Accutase (Sigma-Aldrich) and were plated as monolayer cultures on plates coated with branched polyethylenimine (0.1%, Sigma-Aldrich) and laminin (10 mg/mL, Life Technology) in Brainphys™ media+B27 supplement with BDNF and NGF. The media was changed every 2 days until analysis. As an alternative, 3D NEBs were dissected manually into 3-4 pieces for expansion and further grown, or were cryopreserved. Initial versions of the protocol used Neurobasal™ as a base media instead of Brainphys™.

Genomic DNA Isolation and Sequencing

[0091] Genomic DNA was isolated from PSEN2 mutant or control iPSC lines using High Pure™ PCR Template Preparation Kit (Roche) following manufacturer instructions. Genomic samples were treated with RNase (QIAGEN) prior to amplification. A fragment from exon 5 of PSEN2 containing PSEN2^{N141I} mutation was amplified using the following primers: Forward 5'-CATCAGCCCTTGCCCTCT-3' (SEQ ID NO: 3), Reverse: 5'-CTCACCTTGTAGCAGCGGTA-3' (SEQ ID NO: 4), generating a 173 bp fragment, regardless of the genotype. For detection of ApoE allelic variants, a fragment of 244 bp was amplified prior to sequencing using the primers: Forward: 5'-ACAGAAT-TCGCCCGGCCTGGTACAC-3' (SEQ ID NO: 5), Reverse: 5'-TAAGCTTGGCACGGCTGTCCAAGGA-3' (SEQ ID NO: 6). Both PCR were performed with the following settings: 10 min 94 C, 40 cycles (30 s 94 C, 20 s 62 C, 10 s 72 C) 7 min 72 C. PCR products were run in a 2% agarose gel to check the size of the amplified fragment. After amplification, samples were cleaned using EXOSAP-It™ (Thermo Fisher Scientific) and then sequenced using the following primers: PSEN2 (Forward: 5'-TCAGCATC-

TACACGCCATT-3' (SEQ ID NO: 7), Reverse: 5'-AGCACCACCAAGAAGATGGT-3') (SEQ ID NO: 8), from [53]; ApoE (Forward: 5'-AT-TCGCCCCGGCCTGGTACAC TGCCA-3' (SEQ ID NO: 9), Reverse: 5'-CTGTCCAAGGAGCTGC AGGCGGCGCAG-3' (SEQ ID NO: 10)), from [36].

CRISPR/Cas9 Gene Correction

[0092] iAD1 Control and iAD2 Control lines were originated from 948 (AD1) and 950 (AD2) iPSC lines by CRISPR/Cas9-mediated correction of the PSEN2^{N141I/WT} heterozygous point mutation to PSEN2^{WT/WT}. g1N141I single guide RNA (sgRNA) was cloned into pSpCas9(BB)-2A-GFP (PX458) vector, generating pSpCas9-g1N141I-GFP vector to direct gene editing to the sequence in exon 5 of PSEN2 where the Volga mutation is located. Single stranded oligonucleotides (ssODN) are efficient templates for the CRISPR/Cas9 correction [13, 66]. ssODN #A-N141I (sequence detail below) was used as donor sequence for gene correction. We designed asymmetric ssODN sequences with a long homology arm of 91 bp, and a short homology arm of 36 bp since asymmetrical ssODNs showed a higher efficiency of homology-directed repair using CRISPR/Cas9

TABLE 1

Sequence Name	Bases	Sequence	SEQ ID NO
g1N141I guide RNA F	25	/5Phos/CACCGCATCATGATC AGCGTCATCG	11
g1N141I guide RNA R	25	/5Phos/AACCGATGACGCTG ATCATGATGC	12
Donor ssODN#A N141I	127	GAGAGAACGCGTGGCTGGAGGGC AGGGCCAGGGCCTCACCTTGTA GCAGCGGTACTTGTAGAGCACC ACCAAGAAAGATGGTCATAACCA CGATGACGCTGATCATGATGAG GGTGT <u>C</u> AGCACGGAGT	13

Underline = Base of the ssODN that Corrects the Point Mutation

[0093] The donor sequence and pSpCas9-g1N141I-GFP vector were transduced in the AD1 and AD2 iPSC lines, plated at 50-70% confluence, using Amaxa Human Stem Cell Nucleofector™ kit (Lonza VPH-5002) and re-plated for recovery. GFP+ cells were sorted in a BD FACSAria IIu Cell Sorter™ and were seeded at 30-50 cells per well in 96-well format to detect and pick single clones. Positive clones were expanded, qDNA was extracted and analyzed for successful HDR was determined using a custom designed TaqMan™ genotyping assay with a probe specific for the SNP (dbSNP ID: rs63750215) located in Chr1:227,073,304 A>T. Selected clones were analyzed by Sanger sequencing to confirm correction of Chr1:227,073,304 location and discard possible insertions or deletions in the surrounding areas.

Fluorescence-Activated Cell Sorting (FACS)

[0094] Neural progenitors at day 12 of differentiation were dissociated with Accutase (Sigma-Aldrich) for 5 min at 37 C and inactivated in Neurobasal media. Cells were spun at 1000 rpm for 4 min and the pellets were resuspended in FACS buffer (DPBS, 0.5% BSA Fraction V-Solution, 100 U/mL Penicillin-Streptomycin, 0.5% EDTA and 20 mM Glucose) with PE Mouse anti-human CD271 antibody

(clone C40-1457, BD) at 1:100, also known as p75 or NGFR, and incubated for 20 min at room temperature (RT) in the dark. After the incubation time, cells were washed with FACS buffer and the pellet was resuspended in 2 mL of FACS buffer with 10 μ M ROCK inhibitor (Y27632, Stemgent). p75 positive cells were purified in a BD FACSAria IIu Cell Sorter™ and data was analyzed using FlowJo™ software.

Real-Time Quantitative Polymerase Chain Reaction (RT-qPCR)

[0095] Human iPSC from PSEN2 mutants or control patients were grown in a monolayer and lysed directly in the cell culture wells with RLT buffer. Total RNA purification was performed with the RNeasy™ Micro kit (Qiagen), and was carried out according to the manufacturer's instructions. cDNA was synthesized using SuperScript™ III Reverse Transcriptase (RT) (Invitrogen, Carlsbad, CA). Semi-quantitative real-time PCR was performed on StepOnePlus™ Real-Time PCR System (Applied Biosystems, Foster City, CA) using the primers listed in Table 2 below. We normalized expression levels to GAPDH. The PCR cycling parameters were: 50° C. for 2 min, 95° C. for 10 min, followed by 40 cycles of 95° C. for 15 s and 60° C. for 1 min. Each biological variable was assessed in technical triplicates within each designated "Experiment", and each designated "Experiment" was performed in at least three complete "start to finish" iterations. Expression levels were normalized to the control line, and results were expressed as AVG \pm SEM.

TABLE 2

Gene	Forward Primer 5'-3'	SEQ ID NO	Reverse Primer 5'-3'	SEQ ID NO
BDNF	TAACGGCGGCAGAC AAAAAGA	14	GAAGTATTGCTTCAG TTGGCCT	15
BF1	AGAAGAACGGCAAG TACCGAGA	16	TGTTGAGGGACAGAT TGTGGC	17
Nkx2.1	TAACGGCGGCAGAC AAAAAGA	18	GAAGTATTGCTTCAG TTGGCCT	19
NLRP2, ASB9	From [77]	20	From [77]	21
NLRP3	ACGAATCTCCGACC ACCACT	22	CCATGGCCACAACAA CTGAC	23
Tuj1	GAAGTGTCCCAGGA CATGATAA	24	CTCTTGAGTAGCTGG GATTGAG	25

A β Assays

[0096] Cells were conditioned for 3 days after day 8 of dual SMAD inhibition to measure secretion of A3 by neural progenitors in vitro. A3 levels were quantified using human/rat 3 amyloid 40 ELISA Kit and 3 amyloid 42 ELISA Kit high sensitive (Wako). Each biological variable was assessed using technical triplicates within each designated "Experiment", and each designated "Experiment" was performed in at least three complete "start to finish" iterations.

Immunostaining/ICC

[0097] Cells were fixed with PFA 4% directly on the wells of 12, 48 or 96 well plates for 20 min, washed 3 times with

DPBS 1×(ThermoFisher). For the staining, cells were incubated in blocking solution (DPBS 1× with 0.1% Triton™ X-100 plus 5% Donkey serum) for two hours at room temperature. The corresponding primary antibodies were diluted at suitable concentration in blocking solution, and incubated overnight at 4 C. The primary antibodies used are represented in the table below. Cells were washed three times with DPBST (DPBS 1×+0.1% Triton™ X-100) and suitable secondary antibody was added in blocking solution for 1 h at room temperature. Then cells were washed three times with DPBST and incubated with DRAQ5 or Hoechst 33,342 (1 μ g/mL, diluted in DPBS 1×) for 10 min at room temperature for nuclear counterstain. Cells were visualized using an inverted fluorescence microscope (Olympus™ IX71 microscope) or a confocal microscope (Zeiss™ LSM5 Pascal microscope) under 10×, 20× or 63× magnification.

Western Blots

[0098] Human iPSC from PSEN2 mutants or control patients were grown in a monolayer and lysed directly in the cell culture wells with RIPA buffer (Thermo Scientific) with protease and phosphatase inhibitors. The protein concentration was measured using the BCA protein assay kit (Thermo Scientific). After protein estimation, 20 μ g of cell lysate were separated by SDS-PAGE electrophoresis on a 4-12% Bis-Tris gel (Bolt™ protein gels) and transferred onto nitrocellulose membranes by electrophoresis blotting. The membranes were blocked with blocking buffer 1×TBST (tris-buffered saline+0.10% Tween) plus 5% nonfat dry milk for 1 h in agitation at room temperature and washed three times with TBST. After washing, membranes were incubated at 4° C. overnight in agitation, with the primary antibodies against NLRP2 (1:1000), PSEN2 (1:200) or 3-actin (1:1000). After rinsing, the membranes were incubated with horseradish peroxidase (HRP)-conjugated suitable secondary antibodies for 1 h at room temperature. Finally, protein bands were visualized with a chemiluminescent reagent according to the manufacturer's instructions. 3-actin was used as loading control. Electrophysiology

[0099] Whole cell patch-clamp recordings were obtained from single neurons between differentiation days 38 and 55. Cells were seeded at low density onto plastic coverslips which were placed in a perfusion based enclosed recording chamber. Neurons were localized using differential interference contrast optics under an Olympus BX61WI microscope fitted with a Hamamatsu Orca™ R² CCD camera. Recordings were carried out at room temperature using Multi-Clamp™ 700B amplifier (Molecular Devices, Sunnyvale, CA, USA). Signals were sampled at 10 kHz and filtered at 6 kHz using a Digidata™ 1440A analog to digital converter (Molecular Devices). Amplifier control and data acquisition was done using pClamp™ 10.0 software (Molecular Devices).

[0100] During recordings neurons were perfused with oxygenated Brainphys' media (StemCell Technologies Inc). Medium resistance recording pipettes (4-6 M Ω) were filled with an intracellular solution consisting of (in mM) 130 K-gluconate, 10 KCl, 2 Mg-ATP, 0.2 Na-GTP, 0.6 CaCl₂, 2 MgCl₂, 0.6 EGTA, and 5 HEPES titrated to pH 7.1 and osmolarity of 310 mOsm. In some experiments, the intracellular solution also contained 4 mg/mL biocytin (Sigma-Aldrich) for post-hoc identification of individual neurons, which were visualized with streptavidin-conjugated Alexa 488 (Life Sciences) as described elsewhere [42]. After initial

break-in, access resistance (R_s) was constantly monitored and recordings were discarded if R_s exceeded 20 M Ω or changed more than 30%. The voltage protocol for compound Na $^+$ and K $^+$ currents characterization was as follows: cells were held at -80 mV potential followed by 500 ms steps from -100 mV to 30 mV with 10 mV increment at a frequency of 0.1 Hz. Following transition to current-clamp mode, resting membrane potential was recorded and cells were hyperpolarized by a negative DC current injection to -70 mV to ensure consistency of excitability measurements. Action potentials were evoked with square 1 s current steps from -10 pA to 40 pA with 1 pA steps.

[0101] Electrophysiological recordings were analyzed using ClampFit™ software (Molecular Devices, Sunnyvale, CA, USA) and statistical significance of the results was measured using ANOVA test with Tukey's post-hoc comparison of means. Salts and other reagents were purchased from Sigma-Aldrich (St. Louis, MO, USA).

Statistical Analysis

[0102] qPCR gene expression experiments and A β 42/40 ELISAs were analyzed for statistical significance using Student t-test. LDH Release assays were analyzed by 2-Way ANOVA Bonferroni post hoc tests. ANOVA test with Tukey's post hoc comparisons were used for analysis of electrophysiology results. The experiments needed to study each of the 94 neurons recorded for electrophysiology analyses required days to weeks. On each experimental day, representatives from each genotype were included, with at least three samples from each genotype studied on each day.
*, p<0.05; **, p<0.01; ***, p<0.001.

Results

Optimization of Protocol for BFCN Differentiation

[0103] The scheme of BFCN differentiation is described in FIG. 1A. iPSCs from control subjects or AD patients were plated in feeder-free conditions and allowed to reach 100% con-fluency prior to differentiation using mTeSR1 basal media. Both branches of TGF- β signaling were inhibited (dual SMAD inhibition) to induce neuroectodermal fate on "day 0" [12]. Differentiations (day 2-10) were performed using a modified mTeSR1 formulation, lacking factors that support pluripotency (bFGF, TGF- β , Li—Cl, GABA and pipecolic acid). To specify these cells to basal forebrain cholinergic neurons, ventralization for medial ganglionic eminences (MGE) induction is required [19, 85, 91]. As such cells were treated with the Sonic Hedgehog (Shh) analog (SAG) at 500 nM and Purmorphamine at 2 μ M from days 2 to 8. SAG is a suitable substitute to activate Shh signaling, as demonstrated during differentiation of ChAT $^+$ motor neurons and glutamatergic interneurons [91], with lower cost than recombinant Shh and some advantages in neuronal survival properties over Shh itself [7, 35]. We used the Nkx2.1-GFP embryonic stem cell (ESC) reporter line as a tool to adjust the combination, dosage and timing of ventralizing agents more beneficial for specification of BFCNs from induced Nkx2.1 basal forebrain precursors.

[0104] However, given the potential of Nkx2.1 intermediate neural precursors to generate multiple neuronal subtypes, such as TH $^+$ and GABA $^+$ hypothalamic neurons, we analyzed the expression of the downstream cholinergic specification factor Lhx8 over expression of the GABAergic

interneuron specific transcription factor Lhx6 [26] under different specification conditions (FIG. 1B). These data agree with those from [50] supporting the existence of a synergistic effect of SAG and purmorphamine on Nkx2.1 induction although an effect that is less than the effect of Shh plus purmorphamine (FIG. 1B). Nkx2.1-driven GFP levels were maintained after Day 14, even after withdrawal of SAG+purmorphamine at day 8 (FIG. 1B). We observed higher Lhx8 induction upon SAG plus purmorphamine treatment than SAG alone, or even Shh plus purmorphamine (FIG. 1B). Interestingly, intermediate Nkx2.1 levels driven by SAG plus purmorphamine correlate with higher induction of Lhx8 and BFT gene expression (FIG. 1B). Our choice of starting SHH pathway-driven ventralization at day 2 was based on reports demonstrating other MGE-derived populations being generated by earlier (e.g., hypothalamic neurons) or later (e.g., GABAergic interneurons.) ventralization in the context of dual smad inhibition protocols.

[0105] Following the patterning stage, we gradually switched from Custom mTESR1™ media to Brainphys™ media with B-27 supplement to support neuronal survival and growth [3]. At day 11, we observed neural rosettes positive for Nestin and Sox2 markers (FIG. 1C); also, we observed Tuj1+ neurites as early as day 11 (FIG. 1C). To obtain cholinergic populations of a higher purity, we developed a P75+ FACS strategy to isolate progenitors specific for cholinergic neurons due to the fact that BFCNs are the only CNS neuron type to express robust levels of P75 under non-pathogenic conditions in the adult brain). Support for this strategy includes a previously published protocol using FACS to isolate high expressing P75+ cells from the embryonic murine septum [65]. This population correlated with best expression of cholinergic-related markers.

[0106] At day 11/12, we lifted the cells using chemical dissociation (Accutase) and purified day 11-12 p75+ (CD271) neural progenitors and generated 3D neuronal embryoid bodies (NEBs) by spinning down neural progenitors in V-bottom 96 well plates. On day 19 NEBs were dissociated and re-plated as a monolayer on plates coated with branched polyethylenimine (Aldrich catalog number 408727) and laminin. Monolayer cultures were maintained with the addition of growth factors BDNF, NGF and DAPT until day 26, when cultures no longer had DAPT added. Immunostaining of both cryosections of NEB structures and fixed monolayers, resulting from chemical dissociation of NEBs from several control iPSC and H9 hESC lines, demonstrated expression of BFCN lineage markers Tuj1, MAP2, BF1, Nkx2.1 and p75, at final stages of the differentiation protocol (FIG. 1D). NGF addition to neuronal cultures showed an advantageous effect on maturation, neurite outgrowth and presence of ChAT (FIG. 1E).

Generation and QC of PSEN2 N141I iPSC Lines

[0107] PSEN2 N141I , mutant iPSC and control lines were generated from fresh skin biopsies. Established fibroblast lines were grown from skin punches donated by a kindred of 2 carriers for presenilin 2 Volga familial AD mutation (PSEN2 N141I) and one non-affected member. Additionally, we included a non-family related control. Fibroblasts were reprogrammed using modified RNA method to introduce the Yamanaka factors (Oct4, KLF4, SOX2 and c-Myc), and the iPSC lines obtained were subject to several quality control processes to ensure robust cell-renewal and pluripotency, including alkaline-phosphatase (AP) enzymatic activity, gene expression analysis and immunostaining for pluripo-

tency markers, as well as karyotyping for detection of chromosome abnormalities, following the automated iPSC reprogramming and QC methods developed by [60]. A summary of the genotypes, sex and age of the subjects included in the study is shown in FIG. 2A. The two PSEN2^{N141I}, iPSC lines were also heterozygous for APOE ε4 (ε3/ε4), whereas the control iPSC lines were homozygous ε3/ε3. The characterization of the iPSC lines, expression of pluripotency markers and quality control results are shown in FIGS. 10A-10B. Briefly, all iPSC clones selected demonstrated pluripotency by embryoid body formation and differentiation into the three germ layers (FIG. 10A), incorporated herein by reference). Finally, the lines were finger-printed (Cell Line Genetics) to ensure they matched the parental fibroblast lines (data not shown). All the parental fibroblast lines and the iPSC lines were subject to Sanger sequencing to determine PSEN2 and APOE genotypes. A 173 bp fragment from the exon 5 of PSEN2, surrounding the area where the PSEN2^{N141I} point mutation is located (Chr1: 227,073,304 A>T), was amplified by PCR and sequenced using the primers published in [53]; similarly a fragment of 244 bp from APOE locus that contains two SNPs which determine the three allelic variants was amplified by PCR from genomic DNA, and subsequently sequenced to distinguish between 82/83/84 variants, using the primers from [36]. Sample chromatograms showing the presence of PSEN2N141I point mutation are shown in FIG. 2B, and all genotypes are summarized in FIG. 2A.

Characterization of PSEN2^{N141I} Neural Progenitors

[0108] To study the effect of the PSEN2^{N141I} mutation in early stages of the differentiation of cholinergic neurons, we analyzed the neural progenitors (NPCs) obtained at DIV 11-16 along the BFCN differentiation protocol. The analysis of this intermediate immature population allows us to detect possible early alterations in the generation of BFCNs that would otherwise not be detected in terminally differentiated cholinergic neurons. Such defects could potentially play roles in mature neurons and contribute to the pathophysiology of AD. We analyzed the expression of early neuronal markers in PSEN2^{N141I} mutant and control NPCs by gene expression and immunofluorescence methods. Although, we found a lower RNA expression of Tuj1 (βIII-Tubulin), a general neuronal marker, in mutant NPCs at day 11 of differentiation, we did not detect quantifiable differences by immunocytochemistry circa days 16-21 (FIGS. 2C and 2D). NPC monolayer cultures at day 11 were also immunostained for typical NPC markers: Sox2, and Pax6; with Pax6 levels dropping as expected along with Nkx2.1 induction (not shown). We observed comparable expression of Sox2 and Nestin in PSEN2^{N141I}, cultures at day 11 (FIG. 2C, top panel). At day 21, mutant NPCs expressed comparable levels of Nkx2.1 (MGE marker), but reduced levels of BF1 (forebrain marker) by qPCR; however, BF1 protein expression did not seem affected by immunostaining at this differentiation stage (FIG. 2C bottom panel, and D). We did not observe differences in the surface expression of NGFR (p75/CD271) in DIV11-12 PSEN2^{N141I} cells, in terms of percentage of positive cells or fluorescence mean peak value (FIG. 2E).

[0109] As previously published by [59, 73], the expression of mutant PSEN2^{N141I} causes an increase in the Aβ42/40 ratio in the brains of transgenic mice; additionally, this enhanced Aβ42 production was observed in neural cell lines

upon induced overexpression of mutant PSEN2^{N141I} protein [83] and in iPSC derived from PSEN2^{N141I} mutant patients [93]. Consistently, we observed a 2-fold increase in the Aβ42/40 ratio, a 50% increase in the amount of secreted Aβ40 and 2.5-fold increase in Aβ42 species in the conditioned media from PSEN2N141I neural progenitors at DIV 11 (**p<0.001) (FIG. 2E). The levels of secreted Aβ40 and 42 observed in our study and the levels found in [93], using a different neuronal differentiation method applied to FAD1/PS2 iPSC lines derived from fibroblasts from the Coriell repository are very similar in both absolute number and in fold-increase.

Characterization of Mature BFCNs from PSEN2^{N141I} iPSC Lines and Controls

[0110] With the aim of determining the impact of PSEN2^{N141I} mutation on the differentiation, gene expression, function, and communication of BFCNs, we characterized cells at later time points for appropriate expression markers; our goal was to explore whether PSEN2^{N141I} iPSC were able to complete BFCN maturation process and if so, if any abnormalities along later stages of BFCN differentiation may account for the pathophysiology of EOFAD (FIGS. 3A-3B). In addition to p75, which preferentially binds pro-NGF, we analyzed the expression of TrkB, the primary mature NGF receptor, was also expressed in PSEN2^{N141I} BFCNs and control (FIG. 3A). This suggested that PSEN2^{N141I} BFCNs are susceptible to receiving and benefiting from NGF pro-survival and differentiation signals as expected and further confirms their proper identity. We observed comparable expression of additional cholinergic neuron specific markers choline acetyltransferase (ChAT) and vesicular acetylcholine transporter (vAChT) in PSEN2^{N141I} BFCNs and controls (FIG. 3B). Other general neuronal markers such as Tuj1, and the mature marker microtubule-associated protein 2 (MAP2) showed no apparent differences by immunofluorescence (FIG. 3B).

CRISPR/Cas9-Mediated Correction of PSEN2^{N141I} Mutation and Effect on Aβ 42/40 Ratio

[0111] To determine if the molecular alterations in the processing and cleavage of APP and/or the exacerbated activation of NLRP2 inflammasome, as previously observed in PSEN1 mutants [77], can be attributed to PSEN2^{N141I} mutation only, we modified the PSEN2 locus in our iPSC lines employing CRISPR/Cas9 technology. We did this by correcting the PSEN2^{N141I} point mutation in the two PSEN2 mutant iPSC lines (AD1, AD2). For this purpose, a specific guide RNA (g1N141I) was designed using an online tool (tools.genome-engineering.org) to direct Cas9 to the region of PSEN2 exon 5 surrounding PSEN2^{N141I} mutation (23 bp upstream of Chr1:227,073,304 A>T). g1N141I was cloned into pSpCas9(BB)-2A-GFP (PX458) vector. Expression was assessed by GFP fluorescence upon transfection of pSp-Cas9-g1N141I-GFP in HEK293T (FIG. 4A).

[0112] In order to correct the mutation, we designed an asymmetric ssODN HDR (homology directed repair) template, ssODN #A-N141I, with a long homology arm of 91 bp, and a short homology arm of 36 bp since asymmetrical donor sequences with a shorter arm oriented to the area closer to the PAM side demonstrated a superior efficiency of homology-directed repair using CRISPR/Cas9 system [13]. We then proceeded to transduce pSpCas9-g1N141I-GFP and ssODN #A-N141I into the iPSC lines using Amaxa™ nucleofection (FIG. 4A). Forty-eight hours post-nucleofec-

tion cells were dissociated and the GFP⁺ population was purified by FACS and replated at low density feeder free for isolation of single gene-corrected clones (FIG. 4B). Subsequently, clones were grown and gDNA extracted post expansion. The screening of positive clones that demonstrated successful HDR was determined by qPCR using a custom designed TaqMan™ genotyping assay with a probe specific for the SNP (dbSNP ID: rs63750215) located in Chr1:227,073,304 A>T. We were able to distinguish by this method between homozygous PSEN2^{N141I}, heterozygous PSEN2^{N141I} and PSEN2^{WT} single clones derived from the original iPSC lines, and pre-selected clones were subjected to Sanger sequencing to confirm Chr1:227,073,304 location and detect possible insertions, deletions or mismatches introduced by CRISPR/Cas9 modification in the surrounding area and corroborate successful HDR (FIG. 4C).

[0113] Successfully corrected clones were expanded and subjected to the BFCN differentiation protocol in parallel to the other 4 lines used in the study. We collected media from BFCNs (DIV 34) and re-tested for amyloid beta production. In support of our previous finding in NPCs at DIV 11-12 (FIG. 2F), we observed that mature BFCNs also display significant increases in Aβ42/40 ratio (FIG. 4D) and overall A3 production (FIGS. 11A-11B). Importantly, these results also showed a normalization of Aβ42/40 ratio to control levels in corrected lines (iAD1 Control and iAD2 Control, are corrected clones of AD1 and AD2, respectively) (FIG. 4D). These results also strengthen previous findings linking the PSEN2^{N141I} mutation to abnormal APP processing and reinforcing that presenilins contains the catalytic site of γ-secretase [90].

Assessment of Sensitivity to Aβ42 Oligomer Toxicity in iPSC-Derived PSEN2^{N141I} Neurons

[0114] Previous reports have shown that iPSC lines carrying FAD mutations may display an enhanced susceptibility to noxious stimuli, such as high concentrations of Aβ42 oligomers [2]. We therefore tested whether our BFCNs from PSEN2^{N141I} mutants would display enhanced toxicity to Aβ42 oligomers in the media (FIGS. 5A-5B). We assessed neurotoxicity by measuring the percentage of lactate dehydrogenase (LDH) released by dead cells, thus providing an indirect measurement for toxicity. Using this methodology by 2-way ANOVA we detected a significant effect in toxicity driven by 5 μM Aβ42 oligomer addition to the culture media, after 72-h exposure (***, p<0.01). Post hoc Bonferroni analysis revealed significant differences between the AD2 line and its corrected isogenic control (iAD2 Control). However, this apparent enhanced sensitivity to Aβ42 oligomer toxicity was not observed in the AD1 line and its corresponding control. These results indicate that differences in susceptibility to Aβ42 are not exclusively linked to mutant PSEN2 genotype, and that likely additional genetic factors different between AD1 and AD2 subjects affect susceptibility to this stress, further emphasizing the importance of multiple isogenic models.

Assessment of NLRP2 mRNA in iPSC-Derived PSEN2^{N141I} Neurons

[0115] We previously reported that NLRP2 mRNA was elevated in PSEN1 mutant iPSC and NPCs, [77] which was also the case for PSEN1 mutant cortical neurons (unpublished observation). Therefore, we wanted to analyze the status of the components of the inflammasome in the context of PSEN2^{N141I} mutation. When we assayed by qPCR the mRNA levels of NLRP2 in NPCs at DIV12, we observed an

increase over 100-fold in AD1 and AD2 lines, as compared to control lines (FIG. 6A). This correlated with a notable increase in NLRP2 protein, as observed by SDS-PAGE in whole cell lysates from day 11 PSEN2 mutants (FIG. 6D). Noticeably, however we did not detect bands for NLRP2 by immuno-blot in the AD2 line lysates. Further, we were unable to corroborate some other transcriptional events previously seen in PSEN1 mutant iPS neural precursors, such as the elevated ASB9 that encodes an E3 ligase that directs mitochondrial creatine kinase for degradation. Instead, we observed a trend toward decreased levels in PSEN2 mutation carriers by 20-30%.

Assessment of Excitability of iPSC-Derived PSEN2N141I BFCNs

[0116] Using BFCN differentiation protocol, we were able to generate electrophysiologically active cholinergic neurons in a dish from two PSEN2^{N141I} mutant AD patients, wild-type and familial controls starting from differentiation day 35. We were initially unable to obtain mature action potential waveforms from BFCNs grown in neurobasal media at this stage, but switching to Brainphys' media significantly improved electrophysiological properties of cultured neurons [3]. These findings are in line with electrophysiological characterization of other iPSC generated neurons used to compare both media [3]. The benefits of the protocol containing Brainphys media in two additional cell lines (including the H9 embryonic stem cell line) with comparable endpoint expression of ChAT and VACHT as well as electrophysiological responses were repeated. In order to investigate the electrophysiological properties of BFCN, we recorded from a total of 94 neurons (22 wild-type control, 21 familial control, 18 AD1, 28 AD2 and 5 iAD1_control) using whole cell patch-clamp method. In all experimental groups, recorded neurons displayed voltage-activated currents through sodium and potassium ion channels, ability to generate action potentials and displayed classical neuronal morphologies (FIGS. 7A-7B). In subset of experiments, recorded neurons were labeled with biocytin through a patch pipette, which allowed for post hoc cell identification and ICH characterization. We found that all biocytin-labelled cells were also immuno-positive for ChAT and VACHT (n=12, FIG. 8A).

[0117] Significant differences between the groups in terms of neuronal membrane resistance and capacitance were not observed, membrane resting potential and the minimum current required for generation of a single action potential (FIGS. 9A-9D). However, it was observed that there were significant mutation-related, editing-reversible differences in excitability of BFCNs.

[0118] Neurons derived from AD1 and AD2 lines (as compared to WT and familial controls) were able to generate fewer maximum number of spikes in response to a square depolarizing current injection (ANOVA test with Tukey's post hoc comparisons, FIGS. 8B, 8C). Height of the first action potential at rheobase current injection was also significantly decreased in AD1 and AD2 BFCNs (FIG. 1C). Importantly, CRISPR/Cas9 correction of the PSEN2 point mutation in the AD1 mutant iPSC line abolished the observed electro-physiological deficit, restoring both the maximal number of spikes and spike height to the levels recorded in wild-type and familial controls (ANOVA test with Tukey's post hoc comparisons, FIGS. 8A-8C).

Discussion

[0119] There are 5 million people currently affected by Alzheimer's disease in the US and, according to the Alzheimer's Association, this number will increase to 16 million by the year 2050. Unfortunately, we only have direct evidence for genetic causation that accounts for 3-5% of these patients. This percentage encompasses the EOFAD variants, caused by inherited fully penetrant autosomal dominant mutations in the amyloid protein precursor (APP), or PSEN1, PSEN2 that constitute the γ -secretase apparatus [87], and changes in their function increases the production of A β 42 oligomers and/or deposition of amyloid plaques.

[0120] After decades studying murine models of AD that do not fully recapitulate the pathophysiology of this disease in the human brain [5, 57, 58], a complementary new concept of AD modeling in vitro has emerged upon the breakthrough by [81] allowing adult human tissue reprogramming into iPSC using defined factors, and their subsequent in vitro differentiation into specific brain cell types.

[0121] BFCNs are one of the most vulnerable neuronal populations whose deterioration explains, in part, the cognitive decline in AD patients. Apart from the evidence for BFCN failure and atrophy, other studies have revealed that human embryonic stem cell-derived BFCNs transplanted into AD mouse models can be associated with improvement in the learning behavior of the implanted mouse [94]. These findings highlight the relevance of iPSC- and ESC-derived BFCNs as not only early clinical indicators but also as a potential strategy for subtype-specific cell-based therapy for AD [39]. In order to move this cell-based therapeutic strategy forward, there has been an urgent need for a refined differentiation protocol to generate human ESC- and/or iPSC-derived BFCNs.

[0122] Our first goal was to develop an improved protocol for the generation of BFCNs and intermediate neural progenitors (NPCs), followed by the use of these methods when differentiating cell lines from both control subjects and those harboring the PSEN2^{N141T} mutation. Using fibroblasts isolated from 3 sisters, 2 carrying the PSEN2 mutation and displaying cognitive decline, with the third wild-type for the mutation, iPSCs were developed [60]. In order to approach the dissection of the fidelity of linkage of various phenotypes to the pathogenic mutation, we began by optimizing published BFCNs protocols [4, 17, 46, 50, 89] including the purification of an intermediate CD271+(p75) forebrain progenitor population by Fluorescence Activated Cell Sorting (FACS) to generate 3D ventralized neural embryoid bodies (vNEBs), which can be later dissociated to look at neuronal populations in a monolayer.

[0123] After induction of BFCN differentiation in these cell lines, we have analyzed: (1) capacity to generate Tuj1+/BF1+/ChAT+ neurons in vitro; (2) expression of genes/proteins of interest related to neuronal differentiation or inflammation; (3) generation of soluble and oligomeric A β 40 and 42; (4) electrophysiological (ePhys) properties; and (5) selective vulnerability of BFCNs to one or more innate or microenvironmental factors within or in close approximation to those cells.

[0124] Several studies in AD mouse models highlight electrophysiological defects associated to late stages of AD pathology. Synaptic function in the hippocampus was reduced in APP23 mouse models [70]. Similarly, cholinergic neurons from the prefrontal cortex of TgCRND8 mice are unable to sustain cholinergic excitation as compared to

control mice [64]. Here we report deficient electrophysiological properties in PSEN2^{N141T} iPSC-derived BFCNs in vitro. Notably, correction of this point mutation re-established neuronal excitability to the level of the control iPSC-derived neurons.

[0125] We have optimized an in vitro BFCN differentiation protocol from human iPSC, focusing on generating a homogeneous population of electrophysiologically active ChAT+/VACHT+ neurons in a reproducible and fast way. The innovations introduced to the protocol granted a homogeneous expression of Nkx2.1, a transcriptional marker for MGE subregions, as soon as day 8 and very robust by day 11, compared to day 20 suggested in previously published protocols [38]; in defined serum-free media conditions and without forcing overexpression of factors implicated in cholinergic fate. We were able to record mature action potentials in neurons from day 38 in culture, accompanying co-expression of cholinergic specific markers, which is an earlier time point as compared to other existing protocols using ES or iPSC [4, 17, 46, 50, 89]. Therefore, our protocol has potential application to high-throughput drug screening in homogeneous cholinergic cultures. In addition, the 3D structure of NEBs themselves if left undissociated organoid form would also allow mechanistic analysis in a more physiological setting.

[0126] After applying this optimized protocol to PSEN2^{N141T} mutant iPSC lines, we found an increase in A β 42/40 ratio in the conditioned media. We did not observe any evident defects in the neuronal differentiation process and expression of BFCN markers. Interestingly, we observed a decrease on BDNF gene expression in PSEN2^{N141T} NPCs, similar to results described in a report [18] wherein BDNF changes were observed in homozygous and heterozygous APP^{sws}/PSEN1^{M146V} mice. The two mutant lines are also carriers of one APOE E4 allele. The presence of this allelic variant, the most common and well characterized risk factor polymorphism for LOAD [16], may modulate the age of onset and severity of the phenotype [49]. Therefore, these iPSC lines combining both the EOFAD PSEN2 Volga mutation (or CRISPR/Cas9 corrected) and APOE E4 allele constitute a tremendously useful tool to study the pathophysiology of early onset AD in vitro, especially when apoE-secreting iPSC-derived astrocytes are also present.

[0127] Searching for adjacent mechanisms or events that may be a cause or a consequence of elevated β -amyloid production, researchers have found overactivated inflammation and electrophysiological defects associated with AD mutations. The concept of these defects being independent from β -amyloid deposition and their demonstration using CRISPR/Cas9 technology to correct EOFAD mutations would open the debate to the need of combined AD treatments not only targeting β -amyloid plaques (Gandy et al., in press), but also to overcome parallel inflammatory processes or excitotoxicity/defective neuronal firing.

[0128] NLRPs are components of the inflammasome, which induces the secretion of mature pro-inflammatory cytokine IL-1 β in response to pathogens and toxic stimuli [11, 41]. NLRP2 appears dysregulated in astrocytes [45, 51], and NLRP3 in microglia [34] in the context of Alzheimer's disease as well as in other neurological diseases like Parkinson's disease [14, 32]; additionally, NLRP2/3 are altered in pathologies that show comorbidity with AD: obesity, type-2 diabetes. We previously reported an unexpected

association of elevated expression of the inflammasome gene NLRP2 in iPSC-derived neurons from banked fibroblasts from subjects harboring PSEN1^{A246E} and PSEN1^{M146L} mutations [77]. This association reminded us of the association of the inflammatory skin disease acne inversa (AI) with mutations in presenilin 1, nicastrin, APH-1 and PEN-2, raising in our minds the question of whether some γ -secretase component mutations might be associated not only with proamyloidogenic actions but also with proinflammatory mechanisms.

[0129] Despite our observations PSEN2^{N141I} mutant cells had elevated NLRP2 compared to controls, we were not able to attribute this upregulation to the familial PSEN2 mutation, as gene correction did not significantly reduce NLRP2 levels. Our results suggest that, although inflammasome dysregulation may occur in the brains of EOFAD patients, there may be factors triggering this event apart from any effect of PSENs on inflammasome biology that are reflected in reprogrammed PSEN2 mutant cell lines. Some potential explanations for this PSEN2-independent NLRP2 upregulation include effects of the apoE4 allele present in both PSEN2 subjects (not present in controls) or epigenetic effects on fibroblasts collected from the EOFAD subjects that are maintained through the reprogramming process.

[0130] Electrophysiological defects in neurons have been associated with PSEN1 and PSEN2 mutations. Some of these defects are attributed to altered function of voltage-gated K⁺ channels, potentially through the cleavage of channel components mediated by the PS/ γ -secretase apparatus [44, 72]. Presenilin mutations also disrupt calcium signaling by increasing the levels of calcium stored in the endoplasmic reticulum that result in increased stimulus-induced release into the cytosol, rather than altered influx of calcium. One of the mechanisms behind neuronal calcium dysregulation was described in cortical neurons from PSEN1^{M146V} mice, mediated by inositol triphosphate (IP3) [79]; and, more directly, the formation of dual function protein-ion channels by unprocessed PSEN1 and PSEN2 themselves, modulating the exit of calcium from the endoplasmic reticulum [29, 55, 80, 84]. Given the important role of presenilins on potassium and calcium flux and neuronal excitability, mutations in PSEN1 and PSEN2 may lead to reduced neuronal excitability and neurotoxicity. Mice carrying mutant forms of APP exhibited aberrant action potentials associated to a decrease in sodium currents with no alteration in potassium currents, only after plaque burden was considerable [9]. There is evidence that APP overexpression causes hyperexcitability in mouse cortical neurons [75, 86, 92].

[0131] Mucke and Selkoe [52] have highlighted a toxic effect of A3 resulting in synaptic and network dysfunction. In fibroblasts and neural cell lines, A3-mediated accumulation of mitochondrial Ca²⁺ was elevated when mutant forms of PS1 were expressed [31]. Neuronal firing patterns in mouse hippocampal neurons were altered by exposure to A3 [67, 69]. A3 exposure was also associated with altered K⁺ channel conductance in pyramidal neurons [54]. PSEN1 mutations have been observed to associate with altered Ca²⁺ mitochondrial channels in the cerebellum, apparently causing reduced spike activity in Purkinje cells in the absence of amyloid plaque deposition [74]. A β 42 may accentuate the defects present in Ca²⁺ homeostasis by modulation of additional voltage-dependent ion channels [8, 25, 76, 88].

[0132] Apart from mouse data and immortalized neuronal cell lines, electrophysiological defects in iPSC-derived neurons upon exposure to A3 have been shown: hiPSC-derived cortical pyramidal neurons and GABAergic interneurons have deficient action potentials upon exposure to A3 [56], and neurons differentiated from hiPSC harboring PS1^{A426E} mutation also showed deficient firing patterns [47]. However, there are no previously published data on characterization of electrophysiological properties of PSEN2 mutant iPSC-derived BFCNs.

[0133] Hyper- or hypoexcitatory effects and differences in firing frequency vary with the gene mutation and are highly dependent on the neuronal subtype [37, 48]. All these events may contribute to the progressive neurodegeneration present in the pathogenesis of AD, and we specifically document events that may account for the neuronal defects associated to early stages of EOFAD human pathogenesis. Here we report defective electrophysiological properties in iPSC-derived BFCNs that are specifically associated with the PSEN2N141I familial mutation. Interestingly, although some of the previous studies attribute this impairment in neuronal activity to the build-up of plaques in the brain of AD mice, we found a substantial impairment in the induced action potentials in the absence of amyloid plaques, solely in the presence of a discrete excess of A β 42 oligomers in the culture media, consistent with other reports [18]. Correction of this point mutation re-established the firing patterns to those of the wildtype iPSC-derived neurons.

[0134] Modulators of potassium channels in neurons have proven efficacy in memory improvement in AD mouse models [44]. Modulation of Ca²⁺ channels and excitotoxicity may open a new wave of AD drugs. Understanding the mechanism through which PSEN2 mutations affect the electrophysiological activity in different subsets of neuronal populations and unraveling the connection between PSEN2, other genetic modulatory factors and inflammation will potentially lead to, not only alternative symptomatic treatments, but also to novel drugs decreasing the Ca²⁺-mediated vulnerability to ROS and potentially stopping the neuronal loss and progression of the disease, if administered at early stages.

[0135] It is clear that mutant presenilins alter neuronal excitability even before the formation of A3 plaques [18, 74]. One plausible hypothesis is that APP and presenilins may exert effects that modulate neuronal excitability through currently unrecognized mechanisms acting separate from their roles in the biogenesis of A3. Accumulation of A3 could synergize with the altered electrophysiological mechanisms in a pathway leading to AD. With the wealth of data supporting neuronal excitotoxicity as a key mechanism implicated in AD, further studies focusing on clarifying the possible role(s) of PSENs and/or A3 in physiological or pathological events are warranted.

Conclusions

[0136] We have optimized an in vitro protocol to generate human BFCNs from iPSCs from presenilin 2 (PSEN2) mutation carriers and controls. As expected, PSEN2^{N141I} was associated with an increase in the A β 42/40 in iPSC-derived BFCNs, and this was reversed by CRISPR/Cas9-mediated gene editing. Unexpectedly, iPSC-derived BFCNs or cortical neurons from PSEN2^{N141I} carriers showed diminished basal excitability as quantified by a reduction of both spike frequency and spike amplitude. This electrophysi-

ological phenotype was also abolished following CRISPR/Cas9 correction of the PSEN2^{N141T} mutation. The gene editing data confirm that there was a robust consistency of mutation-related changes that characterized all the expected findings and genotypes from all cells.

Abbreviations

- [0137] AD: Alzheimer's Disease;
- [0138] ApoE: Apolipoprotein E;
- [0139] APP: Amyloid Protein Precursor;
- [0140] AVG: Average;
- [0141] A3: Amyloid Beta;
- [0142] BDNF: Brain Derived Neurotrophic Factor;
- [0143] BF1: Brain Factor 1;
- [0144] BFCNs: Basal Forebrain Cholinergic Neurons;
- [0145] ChAT: Acetylcholine Transferase;
- [0146] DAPT: (N-[N-(3,5-difluorophenacetyl)-L-alanyl]-S-phenylglycine t-butyl ester);
- [0147] DIV: Days in vitro;
- [0148] DNA: Deoxyribonucleic Acid;
- [0149] DPBS: Dulbecco's Phosphate-Buffered Saline;
- [0150] DPBST: Dulbecco's Phosphate-Buffered Saline+0.1% Triton X-100;
- [0151] EGTA: Ethylene-bis(oxyethylenenitrilo)tetraacetic acid;
- [0152] EOFAD: Early Onset Familial Alzheimer's Disease;
- [0153] ESC: Embryonic Stem Cells;
- [0154] FACS: Fluorescence-Activated Cell Sorting;
- [0155] GAPDH: Glyceraldehyde-3-Phosphate Dehydrogenase;
- [0156] GFP: Green Fluorescent Protein;
- [0157] HDR: Homology Directed Repair;
- [0158] HFIP: 1,1,1,3,3-hexafluoro-2-propanol;
- [0159] HRP: Horseradish Peroxidase;
- [0160] iPSCs: Induced Pluripotent Stem Cells;
- [0161] LDH: Lactate Dehydrogenase;
- [0162] MAP2: Microtubule-Associated Protein 2;
- [0163] MGE: Medial Ganglionic Eminences;
- [0164] NEBs: Neuronal Embryoid Bodies;
- [0165] NGF: Nerve Growth Factor;
- [0166] NLRP2: NLR Family Pyrin Domain Containing 2;
- [0167] NPC: Neural Progenitor Cells;
- [0168] PFA: Paraformaldehyde;
- [0169] PSEN: Presenilin;
- [0170] RNA: Ribonucleic Acid;
- [0171] Rock: Rho-Associated, coiled-coil containing protein kinase;
- [0172] RT: Reverse Transcriptase;
- [0173] RT-qPCR: Real-time quantitative polymerase chain reaction;
- [0174] SAG: Smoothened Agonist;
- [0175] SDS-PAGE: Sodium Dodecyl Sulfate PolyAcrylamide Gel Electrophoresis;
- [0176] SEM: Standard Error of the Mean;
- [0177] sgRNA: Single Guide RNA;
- [0178] Shh: Sonic Hedgehog;
- [0179] SNP: Single Nucleotide Polymorphism;
- [0180] ssODN: Single Stranded Oligonucleotides;
- [0181] TBST: Tris-Buffered Saline+0.1% Tween;
- [0182] VACht: Vesicular Acetylcholine Transporter;
- [0183] WT: Wild Type

REFERENCES (EACH ARE INCORPORATED HEREIN BY REFERENCE)

- [0184] 1. Arendt T, Bigl V, Arendt A, Tennstedt A (1983) Loss of neurons in the nucleus basalis of Meynert in Alzheimer's disease, paralysis agitans and Korsakoff's disease. *Acta Neuropathol* 61:101-108
- [0185] 2. Armijo E, Gonzalez C, Shahnawaz M, Flores A, Davis B, Soto C (2017) Increased susceptibility to A3 toxicity in neuronal cultures derived from familial Alzheimer's disease (PSEN1^{A246E}) induced pluripotent stem cells. *Neurosci Lett* 639:74-81. doi:10.1016/j.neulet.2016.12.060
- [0186] 3. Bardy C, van den Hurk M, Eames T, Marchand C, Hernandez R V, Kellogg M, Gorris M, Galet B, Palomares V, Brown J, Bang A G, Mertens J, Böhnke L, Boyer L, Simon S, Gage F H (2015) Neuronal medium that supports basic synaptic functions and activity of human neurons in vitro. *Proc Natl Acad Sci USA* 112: E2725-E2734. doi:10.1073/pnas.1504393112
- [0187] 4. Bissonnette C J, Lyass L, Bhattacharyya B J, Belmadani A, Miller R J, Kessler J A (2011) The controlled generation of functional basal forebrain cholinergic neurons from human embryonic stem cells. *Stem Cells* 29:802-811. doi:10.1002/stem.626
- [0188] 5. Borchelt D R, Ratovitski T, van Lare J, Lee M K, Gonzales V, Jenkins N A, Copeland N G, Price D L, Sisodia S S (1997) Accelerated amyloid deposition in the brains of transgenic mice coexpressing mutant presenilin 1 and amyloid precursor proteins. *Neuron* 19:939-945. doi:10.1016/S0896-6273(00)80974-5
- [0189] 6. Bowen D M, Smith C B, White P, Davison A N (1976) Neurotransmitter-related enzymes and indices of hypoxia in senile dementia and other abiotrophies. *Brain* 99:459-496
- [0190] 7. Bragina O, Sergeieva S, Serg M, Zarkovsky T, Maloverjan A, Kogerman P, Zarkovsky A (2010) Smoothened agonist augments proliferation and survival of neural cells. *Neurosci Lett* 482:81-85. doi:10.1016/j.neulet.2010.06.068
- [0191] 8. Briggs C A, Schneider C, Richardson J C, Stutzmann G E (2013) Beta amyloid peptide plaques fail to alter evoked neuronal calcium signals in APP/PS1 Alzheimer's disease mice. *Neurobiol Aging* 34:1632-1643. doi:10.1016/j.neurobiolaging.2012.12.013
- [0192] 9. Brown J T, Chin J, Leiser S C, Pangalos M N, Randall A D (2011) Altered intrinsic neuronal excitability and reduced Na⁺ currents in a mouse model of Alzheimer's disease. *Neurobiol Aging* 32:2109.e1-2109.e14. doi:10.1016/j.neurobiolaging.2011.05.025
- [0193] 10. Brueggen K, Dyrba M, Barkhof F, Hausner L, Filippi M, Nestor P J, Hauenstein K, Klöppel S, Grothe M J, Kasper E, Teipel S J (2015) Basal forebrain and hippocampus as predictors of conversion to Alzheimer's disease in patients with mild cognitive impairment—a multicenter DTI and Volumetry study. *J Alzheimers Dis* 48:197-204. doi:10.3233/JAD-150063
- [0194] 11. Bruey J M, Bruey-Sedano N, Newman R, Chandler S, Stehlík C, Reed J C (2004) PAN1/NALP2/PYPAF2, an inducible inflammatory mediator that regulates NF-κappaB and caspase-1 activation in macrophages. *J Biol Chem* 279:51897-51907. doi:10.1074/jbc.M406741200
- [0195] 12. Chambers S M, Fasano C A, Papapetrou E P, Tomishima M, Sadelain M, Studer L (2009) Highly

- efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nat Biotechnol* 27:275-280. doi: 10.1038/nbt.1529
- [0196] 13. Chu V T, Weber T, Wefers B, Wurst W, Sander S, Rajewsky K, Kühn R (2015) Increasing the efficiency of homology-directed repair for CRISPR-Cas9-induced precise gene editing in mammalian cells. *Nat Biotechnol* 33:543-548. doi:10.1038/nbt.3198
- [0197] 14. Codolo G, Plotegher N, Pozzobon T, Brucale M, Tessari I, Bubacco L, de Bernard M (2013) Triggering of inflammasome by aggregated α -synuclein, an inflammatory response in synucleinopathies. *PLoS One* 8:e55375. doi:10.1371/journal.pone.0055375
- [0198] 15. Cong L, Ran F A, Cox D, Lin S, Barretto R, Habib N, Hsu P D, Wu X, Jiang W, Marraffini L A, Zhang F (2013) Multiplex genome engineering using CRISPR/Cas systems. *Science* 339:819-823. doi:10.1126/science.1231143
- [0199] 16. Corder E H, Saunders A M, Strittmatter W J, Schmeichel D E, Gaskell P C, Small G W, Roses A D, Haines J L, Pericak-Vance M A (1993) Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science* 261:921-923
- [0200] 17. Crompton L A, Byrne M L, Taylor H, Kerrigan T L, Bru-Mercier G, Badger J L, Barbuti P A, Jo J, Tyler S J, Allen S J, Kunath T, Cho K, Caldwell M A (2013) Stepwise, non-adherent differentiation of human pluripotent stem cells to generate basal forebrain cholinergic neurons via hedgehog signaling. *Stem Cell Res* 11:1206-1221. doi:10.1016/j.scr.2013.08.002
- [0201] 18. Cummings D M, Liu W, Portelius E, Bayram S, Yasvoina M, Ho S-H, Smits H, Ali S S, Steinberg R, Pegasiou C-M, James O T, Matarin M, Richardson J C, Zetterberg H, Blennow K, Hardy J A, Salih D A, Edwards F A (2015) First effects of rising amyloid- β in transgenic mouse brain: synaptic transmission and gene expression. *Brain* 138:1992-2004. doi:10.1093/brain/awv127
- [0202] 19. Danjo T, Eiraku M, Muguruma K, Watanabe K, Kawada M, Yanagawa Y, Rubenstein J L R, Sasai Y (2011) Subregional specification of embryonic stem cell-derived ventral Telencephalic tissues by timed and combinatorial treatment with extrinsic signals. *J Neurosci* 31:1919-1933. doi:10.1523/JNEUROSCI.5128-10.2011
- [0203] 20. Davies P, Maloney A J (1976) Selective loss of central cholinergic neurons in Alzheimer's disease. *Lancet* (London, England). 2:1403
- [0204] 21. Davis K L, Mohs R C, Davis B M, Horvath T B, Greenwald B S, Rosen W G, Levy M I, Johns C A (1983) Oral physostigmine in Alzheimer's disease. *Psychopharmacol Bull* 19:451-453
- [0205] 22. De Strooper B After Solanezumab: Where Should Alzheimer's Research Go? ALZFORUM. 2017
- [0206] 23. Echeverria V, Berman D E, Arancio O (2007) Oligomers of beta-amyloid peptide inhibit BDNF-induced arc expression in cultured cortical neurons. *Curr Alzheimer Res* 4:518-521
- [0207] 24. Epelbaum S, Genton R, Cavedo E, Habert M O, Lamari F, Gagliardi G, Lista S, Teichmann M, Bakardjian H, Hampel H, Dubois B (2017) Preclinical Alzheimer's disease: a systematic review of the cohorts underlying the concept. *Alzheimers Dement*. doi:10.1016/j.jalz.2016.12.003
- [0208] 25. Etcheberrigaray R, Ito E, Kim C S, Alkon D L (1994) Soluble beta-amyloid induction of Alzheimer's phenotype for human fibroblast K $^{+}$ channels. *Science* 264:276-279
- [0209] 26. Flandin P, Zhao Y, Vogt D, Jeong J, Long J, Potter G, Westphal H, Rubenstein J L R (2011) Lhx6 and Lhx8 coordinately induce neuronal expression of Shh that controls the generation of interneuron progenitors. *Neuron* 70:939-950. doi:10.1016/j.neuron.2011.04.020
- [0210] 27. Francis P T, Palmer A M, Snape M, Wilcock G K (1999) The cholinergic hypothesis of Alzheimer's disease: a review of progress. *J Neurol Neurosurg Psychiatry* 66:137-147. doi:10.1136/JNNP.66.2.137
- [0211] 28. Gamzu E R, Thal L J, Davis K L (1990) Therapeutic trials using tacrine and other cholinesterase inhibitors. *Adv Neurol* 51:241-245
- [0212] 29. Gandy S, Doeuvre M K, Poolman B (2006) Alzheimer disease: presenilin springs a leak. *Nat Med* 12:1121-1123. doi:10.1038/nm1006-1121
- [0213] 30. Goulbum A L, Alden D, Davis R P, Micallef S J, Ng E S, Yu Q C, Lim S M, Soh C-L, Elliott D A, Hatzistavrou T, Bourke J, Watmuff B, Lang R J, Haynes J M, Pouton C W, Giudice A, Trounson A O, Anderson S A, Stanley E G, Elefanty A G (2011a) A targeted NKX2.1 human embryonic stem cell reporter line enables identification of human basal forebrain derivatives. *Stem Cells* 29:462-473. doi:10.1002/stem.587
- [0214] 31. Guo Q, Fu W, Holtsberg F W, Steiner S M, Mattson M P (1999) Superoxide mediates the cell-death-enhancing action of presenilin-1 mutations. *J Neurosci Res* 56:457-470. doi:10.1002/(SICI)1097-4547(19990601)56:5<457::AID-JNR2>3.0.CO;2-P
- [0215] 32. Gustot A, Gallea J I, Sarroukh R, Celej M S, Ruysschaert J-M, Raussens V (2015) Amyloid fibrils are the molecular trigger of inflammation in Parkinson's disease. *Biochem J* 471:323-333. doi:10.1042/BJ20150617
- [0216] 33. Hager K, Baseman A S, Nye J S, Brashear H R, Han J, Sano M, Davis B, Richards H M (2014) Effects of galantamine in a 2-year, randomized, placebo-controlled study in Alzheimer's disease. *Neuropsychiatr Dis Treat* 10:391-401. doi:10.2147/NDT.557909
- [0217] 34. Halle A, Hornung V, Petzold G C, Stewart C R, Monks B G, Reinheinkel T, Fitzgerald K A, Latz E, Moore K J, Golenbock D T (2008) The NALP3 inflammasome is involved in the innate immune response to amyloid-. *Nat Immunol* 9:857-865. doi:10.1038/ni.1636
- [0218] 35. Heine V M, Griveau A, Chapin C, Ballard P L, Chen J K, Rowitch D H (2011) A small-molecule smoothed agonist prevents Glucocorticoid-induced neonatal Cerebellar injury. *Sci Transl Med* 3:105ra104. doi:10.1126/scitranslmed.3002731
- [0219] 36. Hixson J E, Vernier D T (1990) Restriction isotyping of human apolipoprotein E by gene amplification and cleavage with HhaI. *J Lipid Res* 31:545-548. doi:10.0000/PMID2341813
- [0220] 37. Hoxha E, Boda E, Montarolo F, Parolisi R, Tempia F (2012) Excitability and synaptic alterations in the cerebellum of APP/PS1 mice. *PLoS One* 7:e34726. doi:10.1371/journal.pone.0034726
- [0221] 38. Hu Y, Qu Z, Cao S, Li Q, Ma L, Krencik R, Xu M, Liu Y (2016) Directed differentiation of basal fore-

- brain cholinergic neurons from human pluripotent stem cells. *J Neurosci Methods* 266:42-49. doi:10.1016/j.jneumeth.2016.03.017
- [0222] 39. Hunsberger J G, Rao M, Kurtzberg J, Bulte J W M, Atala A, LaFerla F M, Greely H T, Sawa A, Gandy S, Schneider L S, Doraiswamy P M (2016) Accelerating stem cell trials for Alzheimer's disease. *Lancet Neurol* 15:219-230. doi:10.1016/S1474-4422(15)00332-4
- [0223] 40. Jack C R, Wiste H J, Weigand S D, Knopman D S, Lowe V, Vemuri P, Mielke M M, Jones D T, Senjem M L, Gunter J L, Gregg B E, Pankratz V S, Petersen R C (2013) Amyloid-first and neurodegeneration-first profiles characterize incident amyloid PET positivity. *Neurology* 81:1732-1740. doi:10.1212/01.wnl.0000435556.21319.e4
- [0224] 41. Kinoshita T, Wang Y, Hasegawa M, Imamura R, Suda T (2005) PYPAF3, a PYRIN-containing APAF-1-like protein, is a feedback regulator of caspase-1-dependent interleukin-1beta secretion. *J Biol Chem* 280: 21720-21725. doi:10.1074/jbc.M410057200
- [0225] 42. Kruglikov I, Rudy B (2008) Perisomatic GABA release and thalamocortical integration onto neocortical excitatory cells are regulated by neuromodulators. *Neuron* 58:911-924. doi:10.1016/j.neuron.2008.04.024
- [0226] 43. Levy-Lahad E, Wasco W, Poorkaj P, Romano D, Oshima J, Pettingell W, Yu C, Jondro P, Schmidt S, Wang K et al (1995) Candidate gene for the chromosome 1 familial Alzheimer's disease locus. *Science* 269:973-977. doi:10.1126/science.7638622
- [0227] 44. Liu D, Pitta M, Lee J-H, Ray B, Lahiri D K, Furukawa K, Mughal M, Jiang H, Villarreal J, Cutler R G, Greig N H, Mattson M P (2010) The KATP channel activator diazoxide ameliorates amyloid- and tau pathologies and improves memory in the 3xTgAD mouse model of Alzheimer's disease. *J Alzheimers Dis* 22:443-457. doi:10.3233/JAD-2010-101017
- [0228] 45. Liu L, Chan C (2014) The role of inflammasome in Alzheimer's disease. *Ageing Res Rev* 15:6-15. doi:10.1016/j.arr.2013.12.007
- [0229] 46. Liu Y, Weick J P, Liu H, Krencik R, Zhang X, Ma L, Zhou G, Ayala M, Zhang S-C (2013) Medial ganglionic eminence-like cells derived from human embryonic stem cells correct learning and memory deficits. *Nat Biotechnol* 31:440-447. doi:10.1038/nbt.256
- [0230] 47. Mahairaki V, Ryu J, Peters A, Chang Q, Li T, Park T S, Burridge P W, Talbot C C, Asnaghi L, Martin L J, Zambidis E T, Koliatsos V E, Koliatsos V E (2014) Induced pluripotent stem cells from familial Alzheimer's disease patients differentiate into mature neurons with amyloidogenic properties. *Stem Cells Dev* 23:2996-3010. doi:10.1089/scd.2013.0511
- [0231] 48. Marcantonio A, Raymond E F, Carbone E, Marie H (2014) Firing properties of entorhinal cortex neurons and early alterations in an Alzheimer's disease transgenic model. *Pflugers Arch* 466:1437-1450. doi:10.1007/s00424-013-1368-z
- [0232] 49. Marchani E E, Bird T D, Steinbart E J, Rosenthal E, Yu C-E, Schellenberg G D, Wijsman E M (2010) Evidence for three loci modifying age-at-onset of Alzheimer's disease in early-onset PSEN2 families. *Am J Med Genet B Neuropsychiatr Genet* 153B:1031-1041. doi:10.1002/ajmg.b.31072
- [0233] 50. Maroof A M, Keros S, Tyson J A, Ying S-W, Ganat Y M, Merkle F T, Liu B, Goulburn A, Stanley E G, Elefanti A G, Widmer H R, Eggan K, Goldstein P A, Anderson S A, Studer L (2013) Directed differentiation and functional maturation of cortical interneurons from human embryonic stem cells. *Cell Stem Cell* 12:559-572. doi:10.1016/j.stem.2013.04.008
- [0234] 51. Minkiewicz J, de Rivero Vaccari J P, Keane R W (2013) Human astrocytes express a novel NLRP2 inflammasome. *Glia* 61:1113-1121. doi:10.1002/glia.22499
- [0235] 52. Mucke L, Selkoe D J (2012) Neurotoxicity of amyloid-protein: synaptic and network dysfunction. *Cold Spring Harb Perspect Med* 2:a006338. doi:10.1101/cshperspect.a006338
- [0236] 53. Müller U, Winter P, Graeber M B (2011) Alois Alzheimer's case, Auguste D., did not carry the N141I mutation in <emph type="quot; italic" > PSEN2</emph> characteristic of Alzheimer disease in Volga Germans. *Arch Neurol* 68:1210. doi:10.1001/archneurol.2011.218
- [0237] 54. Nava-Mesa M O, Jimenez-Diaz L, Yajeya J, Navarro-Lopez J D (2013) Amyloid-induces synaptic dysfunction through G protein-gated inwardly rectifying potassium channels in the fimbria-CA3 hippocampal synapse. *Front Cell Neurosci* 7:117. doi:10.3389/fncel.2013.00117
- [0238] 55. Nelson O, Supnet C, Tolia A, Horre K, De Strooper B, Bezprozvanny I (2011) Mutagenesis mapping of the Presenilin 1 calcium leak conductance pore. *J Biol Chem* 286:22339-22347. doi:10.1074/jbc.M111.243063
- [0239] 56. Nieweg K, Andreyeva A, van Stegen B, Tanrıöver G, Gottmann K (2015) Alzheimer's disease-related amyloid-induces synaptotoxicity in human iPS cell-derived neurons. *Cell Death Dis* 6:e1 709. doi:10.1038/cddis.2015.72
- [0240] 57. Oddo S, Caccamo A, Shepherd J D, Murphy M P, Golde T E, Kayed R, Metherate R, Mattson M P, Akbari Y, LaFerla F M (2003) Triple-transgenic model of Alzheimer's disease with plaques and tangles: intracellular Abeta and synaptic dysfunction. *Neuron* 39:409-421
- [0241] 58. Onos K D, Sukoff Rizzo S J, Howell G R, Sasner M (2016) Toward more predictive genetic mouse models of Alzheimer's disease. *Brain Res Bull* 122:1-11. doi:10.1016/j.brainresbull.2015.12.003
- [0242] 59. Oyama F, Sawamura N, Kobayashi K, Morishima-Kawashima M, Kuramochi T, Ito M, Tomita T, Maruyama K, Saido T C, Iwatsubo T, Capell A, Walter J, Grünberg J, Ueyama Y, Haass C, Ihara Y (1998) Mutant presenilin 2 transgenic mouse: effect on an age-dependent increase of amyloid beta-protein 42 in the brain. *J Neurochem* 71:313-322
- [0243] 60. Paull D, Sevilla A, Zhou H, Hahn A K, Kim H, Napolitano C, Tsankov A, Shang L, Krumholz K, Jagadeesan P, Woodard C M, Sun B, Vilboux T, Zimmer M, Forero E, Moroziewicz D N, Martinez H, Malicdan M C V, Weiss K A, Vensand L B, Dusenberry C R, Polus H, Sy K T L, Kahler D J, Gahl W A, Solomon S L, Chang S, Meissner A, Eggan K, Noggle S A (2015) Automated, high-throughput derivation, characterization and differentiation of induced pluripotent stem cells. *Nat Methods* 12:885-892. doi: 10.1038/nmeth.3507
- [0244] 61. Perry E K, Gibson P H, Blessed G, Perry R H, Tomlinson B E (1977) Neurotransmitter enzyme abnormalities in senile dementia. Choline acetyltransferase and

- glutamic acid decarboxylase activities in necropsy brain tissue. *J Neurol Sci* 34:247-265
- [0245] 62. Pini L, Pievani M, Bocchetta M, Altomare D, Bosco P, Cavedo E, Galluzzi S, Marizzoni M, Frisoni G B (2016) Brain atrophy in Alzheimer's disease and aging. *Ageing Res Rev* 30:25-48. doi:10.1016/j.arr.2016.01.002
- [0246] 63. Price D L, Tanzi R E, Borchelt D R, Sisodia S S (1998) Alzheimer's disease: genetic studies and transgenic models. *Annu Rev Genet* 32:461-493. doi:10.1146/annurev.genet.32.1.461
- [0247] 64. Proulx É, Fraser P, McLaurin J, Lambe E K (2015) Impaired cholinergic excitation of prefrontal attention circuitry in the TgCRND8 model of Alzheimer's disease. *J Neurosci* 35:12779-12791. doi:10.1523/JNEUROSCI.4501-14.2015
- [0248] 65. Pruszak J, Sonntag K C, Aung M H, Sanchez-Pernaute R, Isaacson O (2007) Markers and methods for cell sorting of human embryonic stem cell-derived neural cell populations. *Stem Cells* 25(9):2257-2268 Epub 2007 Jun. 21. PubMed PMID: 17588935; PubMed Central PMCID: PMC2238728
- [0249] 66. Ran F A, Hsu P D, Wright J, Agarwala V, Scott D A, Zhang F (2013) Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* 8:2281-2308. doi:10.1038/nprot.2013.143
- [0250] 67. Ren S-C, Shao H, Ji W-G, Jiang H-H, Xu F, Chen P-Z, Mi Z, Wen B, Zhu G-X, Zhu Z-R (2015) Riluzole prevents soluble α_{1-42} oligomers-induced perturbation of spontaneous discharge in the hippocampal CA1 region of rats. *Amyloid* 22:36-44. doi:10.3109/13506129.2014.990558
- [0251] 68. Richardson C D, Ray G J, DeWitt M A, Curie G L, Com J E (2016) Enhancing homology-directed genome editing by catalytically active and inactive CRISPR-Cas9 using asymmetric donor DNA. *Nat Biotechnol* 34:339-344. doi:10.1038/nbt.3481
- [0252] 69. Ripoli C, Cocco S, Li Puma D D, Piacentini R, Mastrodonato A, Scala F, Puzzo D, D'Ascenzo M, Grassi C (2014) Intracellular accumulation of Amyloid-(a) protein plays a major role in a-induced alterations of Glutamatergic synaptic transmission and plasticity. *J Neurosci* 34:12893-12903. doi:10.1523/JNEUROSCI.1201-14.2014
- [0253] 70. Roder S, Danober L, Pozza M, Lingenoehlk K, Wiederhold K—H, Olpe H-R (2003) Electrophysiological studies on the hippocampus and prefrontal cortex assessing the effects of amyloidosis in amyloid precursor protein 23 transgenic mice. *Neuroscience* 120:705-720. doi:10.1016/50306-4522(03)00381-6
- [0254] 71. Rylett R J, Ball M J, Colhoun E H (1983) Evidence for high affinity choline transport in synaptosomes prepared from hippocampus and neocortex of patients with Alzheimer's disease. *Brain Res* 289:169-175
- [0255] 72. Sachse C C, Kim Y H, Agsten M, Huth T, Alzheimer C, Kovacs D M, Kim D Y (2013) BACE1 and presenilin/-secretase regulate proteolytic processing of KCNE1 and 2, auxiliary subunits of voltage-gated potassium channels. *FASEB J* 27:2458-2467. doi:10.1096/fj.12-214056
- [0256] 73. Sawamura N, Morishima-Kawashima M, Waki H, Kobayashi K, Kuramochi T, Frosch M P, Ding K, Ito M, Kim T W, Tanzi R E, Oyama F, Tabira T, Ando S, Ihara Y (2000) Mutant-presenilin 2-transgenic mice: a large increase in the levels of a beta 42 is presumably associated with the low-density membrane domain that contains decreased levels of glycerophospholipids and sphingomyelin. *J Biol Chem* 275:27901-27908. doi: 10.1074/jbc.M004308200
- [0257] 74. Sepulveda-Falla D, Barrera-Ocampo A, Hagel C, Korwitz A, Vinuela-Veloz M F, Zhou K, Schoneville M, Zhou H, Velazquez-Perez L, Rodriguez-Labrador R, Villegas A, Ferrer I, Lopera F, Langer T, De Zeeuw C I, Glatzel M (2014) Familial Alzheimer's disease-associated presenilin-1 alters cerebellar activity and calcium homeostasis. *J Clin Invest* 124:1552-1567. doi:10.1172/JC166407
- [0258] 75. Šišková Z, Justus D, Kaneko H, Friedrichs D, Henneberg N, Beutel T, Pitsch J, Schoch S, Becker A, von der Kammer H, Remy S (2014) Dendritic structural degeneration is functionally linked to cellular Hyperexcitability in a mouse model of Alzheimer's disease. *Neuron* 84:1023-1033. doi:10.1016/j.neuron.2014.10.024
- [0259] 76. Smilansky A, Dangoor L, Nakdimon I, Ben-Hail D, Mizrahi D, Shoshan-Barmatz V (2015) The voltage-dependent Anion Channel 1 mediates Amyloid-toxicity and represents a potential target for Alzheimer disease therapy. *J Biol Chem* 290:30670-30683. doi:10.1074/jbc.M115.69149
- [0260] 77. Sproul A A, Jacob S, Pre D, Kim S H, Nestor M W, Navarro-Sobrino M, Santa-Maria I, Zimmer M, Aubry S, Steele J W, Kahler D J, Dranovsky A, Arancio O, Crary J F, Gandy S, Noggle S A (2014) Characterization and molecular profiling of PSEN1 familial Alzheimer's disease iPSC-derived neural progenitors. *PLoS One* 9:e84547. doi:10.1371/journal.pone.0084547
- [0261] 78. Stine W B, Dahlgren K N, Kraft G A, LaDu M J (2003) In vitro characterization of conditions for amyloid-beta peptide oligomerization and fibrillogenesis. *J Biol Chem* 278:11612-11622. doi:10.1074/jbc.M210207200
- [0262] 79. Stutzmann G E, Caccamo A, LaFerla F M, Parker I (2004) Dysregulated IP3 signaling in cortical neurons of knock-in mice expressing an Alzheimer's-linked mutation in Presenilin1 results in exaggerated Ca²⁺ signals and altered membrane excitability. *J Neurosci* 24:508-513. doi:10.1523/JNEUROSCI.4386-03.2004
- [0263] 80. Supnet C, Bezprozvanny I (2011) Presenilins function in E R calcium leak and Alzheimer's disease pathogenesis. *Cell Calcium* 50:303-309. doi:10.1016/j.ceca.2011.05.013
- [0264] 81. Takahashi K, Yamanaka S (2006) Induction of Pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 126(4):663-676. doi:10.1016/j.cell.2006.07.024
- [0265] 82. Thal L J, Fuld P A, Masur D M, Sharpless N S (1983) Oral physostigmine and lecithin improve memory in Alzheimer disease. *Ann Neurol* 13:491-496. doi:10.1002/ana.410130504
- [0266] 83. Tomita T, Maruyama K, Saido T C, Kume H, Shinohaki K, Tokuhiro S, Capell A, Walter J, Grunberg J, Haass C, Iwatsubo T, Obata K (1997) The presenilin 2 mutation (N141I) linked to familial Alzheimer disease (Volga German families) increases the secretion of amyloid beta protein ending at the 42nd (or 43rd) residue. *Proc Natl Acad Sci USA* 94:2025-2030
- [0267] 84. Tu H, Nelson O, Bezprozvanny A, Wang Z, Lee S—F, Hao Y—H, Semeels L, De Strooper B, Yu G,

- Bezprozvanny I (2006) Presenilins form ER Ca²⁺ leak channels, a function disrupted by familial Alzheimer's disease-linked mutations. *Cell* 126:981-993. doi:10.1016/j.cell.2006.06.059
- [0268] 85. Tucker E S, Segall S, Gopalakrishna D, Wu Y, Vernon M, Polleux F, Lamantia A-S (2008) Molecular specification and patterning of progenitor cells in the lateral and medial ganglionic eminences. *J Neurosci* 28:9504-9518. doi:10.1523/JNEUROSCI.2341-08.2008
- [0269] 86. Varga E, Juhász G, Bozsó Z, Penke B, Fülop L, Szegedi V (2014) Abeta(1-42) enhances neuronal excitability in the CA1 via NR2B subunit-containing NMDA receptors. *Neural Plast* 2014:1-12. doi:10.1155/2014/584314
- [0270] 87. Verdile G, Gandy S E, Martins R N (2007) The role of presenilin and its interacting proteins in the biogenesis of Alzheimer's beta amyloid. *Neurochem Res* 32:609-623. doi:10.1007/s11064-006-9131-x
- [0271] 88. Wang X, Zhang X-G, Zhou T-T, Li N, Jang C—Y, Xiao Z—C, Ma Q-H, Li S (2016) Elevated neuronal excitability due to modulation of the voltage-gated Sodium Channel Nav1.6 by A 1-42. *Front Neurosci* 10:94. doi:10.3389/fnins.2016.00094
- [0272] 89. Wicklund L, Leão R N, Stromberg A-M, Mousavi M, Hovatta O, Nordberg A, Marutle A (2010) β-amyloid 1-42 oligomers impair function of human embryonic stem cell-derived forebrain cholinergic neurons. *PLoS One* 5:e15600. doi:10.1371/journal.pone.0015600
- [0273] 90. Wolfe M S, Selkoe D J, Xia W, Ostaszewski B L, Diehl T S, Kimberly W T (1999) Two transmembrane aspartates in presenilin-1 required for presenilin endoproteolysis and [gamma]-secretase activity. *Nature* 398:513-517. doi:10.1038/19077
- [0274] 91. Xu H, Sakiyama-Elbert S E (2015) Directed differentiation of V3 Interneurons from mouse embryonic stem cells. *Stem Cells Dev* 24:2723-2732. doi:10.1089/scd.2015.0122
- [0275] 92. Xu W, Fitzgerald S, Nixon R A, Levy E, Wilson D A (2015) Early hyperactivity in lateral entorhinal cortex is associated with elevated levels of A PP metabolites in the Tg2576 mouse model of Alzheimer's disease. *Exp Neurol* 264:82-91. doi:10.1016/j.expneurol.2014.12.008
- [0276] 93. Yagi T, Ito D, Okada Y, Akamatsu W, Nihei Y, Yoshizaki T, Yamanaka S, Okano H, Suzuki N (2011) Modeling familial Alzheimer's disease with induced pluripotent stem cells. *Hum Mol Genet* 20:4530-4539. doi:10.1093/hmg/ddr394
- [0277] 94. Yue W, Li Y, Zhang T, Jiang M, Qian Y, Zhang M, Sheng N, Feng S, Tang K, Yu X, Shu Y, Yue C, Jing N (2015) ESC-derived basal forebrain cholinergic neurons ameliorate the cognitive symptoms associated with Alzheimer's disease in mouse models. *Stem Cell Rep* 5:776-790. doi:10.1016/j.stemcr.2015.09.01.
- [0278] Although the invention has been described with reference to the examples herein, it will be understood that modifications and variations are encompassed within the spirit and scope of the invention. Accordingly, the invention is limited only by the following claims.

SEQUENCE LISTING

```

Sequence total quantity: 31
SEQ ID NO: 1      moltype = DNA length = 33258
FEATURE          Location/Qualifiers
misc_feature     1..33258
                  note = chromosome 1, GRCh38.p12 Primary Assembly
source           1..33258
                  mol_type = genomic DNA
                  organism = Homo sapiens
SEQUENCE: 1
ggggcctggg cccgcggccgg gtcggccgg ggcgtcaagcc agctgcgtaa actccgcgtgg 60
agcgcggccgg cagagcagggt gagcgccgggg tgccgggggg tgcccgaggcc aggccctgt 120
cgccctgcggc gctgaggggcc cgggggtgggg ctgcgcctgc aggccctgc cctgcctcc 180
gcacgcctct ggccacggtc ctttccccgg ctgtgggtct gcccgcctgt cgtgcgcagc 240
gtccctggcc tctgcggcca gcgcgggggc ggagagaggc gagtgcccgag caggccgggg 300
ctggggccgc cccgaactgg gtcgtggaa gatcgcgggg agccgccttc aggcccttcgg 360
cctcaactgcg tccccacttc cttgcgcggc cttgcgcggc agcccccgtt ggggggtggc 420
gccccgcgcgag cggtaaagg gcccgtcat ttaaaggagc ggtgcacgtg ggtctctgag 480
gcgtgttagca ggcggggggc ttttgggtt cttctcttc gccggagacc tccgttgcgc 540
cgagtccatt cggcccttag cacccgggttc tgggcatgtt ttccccggga aggaggcgcc 600
cgggggctct gcccccatgt gaggggcagg gccgcaggct caagcttaga gccgggtttct 660
gttagcagcg gtgtttggc gttttatcag gcatttccag cagtggagg acagccagaa 720
gcaagctttt ggagctgaa gaacctgaga cagaagtagc tccccctctt gaattttact 780
gatgaagaaa ctggggccac agagctaata gtagtttcc caaggctcgcc caggtaacgt 840
atacgacgcg caggcttcg ccccaagtgc ctggcttcta gatctgtctgt ccatccctcc 900
gagcagaccc caccctgtt tattgcctta ataagtattc ccttggaaag gtagaaccgg 960
tggtttagtgc agtaactgc tccttcttcaaatggaga acctgcggagg attccataga 1020
gacgattgtta gactaactta actcagaagc gacagcgttc ggttgcggaa gctgtctacg 1080
aagtaacttg attaggaccc accccaggctt ccagtaaggc agccctctgtat gcctctgtat 1140
ccaaattctgc agacacactga gcctccaagg cttcagccca agacctttgg cggttaattgg 1200
agtctcgggta taagctgttgc cagggtgttg agcctcaggat tcttctcttc tgaatgtgg 1260
tgcggccaggc cggctactgg cgcagggtgcga aaaggggggctt ggttctggc cccacactcag 1320
agctgcgtcc tcacgacgcc caccgttgcac cttgggttcc agggcagaga ctggaggatgg 1380
ggcttgggggg catgttgcattt tgaagtggga tggatgtatc aggtttttgg gaaaaactct 1440
gtaccctttt gtgtttagtgc gcccattgtc caagtctgtca gtccagcatg ttcacatgtg 1500
gggagtgagt ggcttggccc tgcgttattt aaagagcagg aaggaggagg aggagcaagg 1560
gctaggggctt gctgtgggg tgcctggagc tgcgttgcatt aatgtcacac ctgtctcccc 1620
tccgttagtgc ctcaccgtcc ccccaagggg ggtttgccttc ttgcctactt tggccctttc 1680

```

-continued

ctgttatcga	tgttaataat	gacatatata	tcgttatcga	gttggcata	ataaaaagct	1740
atcttgata	gaatattaga	atttaaatgc	ttaagaattt	caatgacact	gaaataggtt	1800
atttacact	ttttacaga	gaggaaaca	gttcaggag	ttaagcagct	agtccagtt	1860
tgtggcttc	gtgccttaag	caccaggatt	tgaacatagc	tggctacact	gtcttatct	1920
cttgagttt	tgcgcaggag	gttctgtat	tcaactctca	cccggtctc	tccactactg	1980
ctggaaagt	tttggggat	ccccatgago	aacttctgta	caaacaaca	aaatttttt	2040
aaagaaaacca	aagcagtgt	tgttaggtca	atgcagtgt	tctaatgaaa	acatctctgg	2100
cgggtttca	gtgttgtctt	tgacttcegg	acactgttt	gttggggact	gataagacag	2160
caaattattc	tgcaagtatt	cccacctgtt	ctattccca	ctggcacagc	tgccggaaagg	2220
cgggggtag	gtgagggag	cccggaggaa	acatttccca	ctgggtctca	atccggaga	2280
tggatgacc	atcatgtta	tgttggaga	aaaatgtat	ttcaggctgg	gtgtgtggc	2340
tcatgtctgt	aatccccaga	cttggggagg	ccggagggtgg	ttgatcacct	gaggtcgaa	2400
gtttgagacc	agcctgacca	acatggggaa	accccatctt	tactaaaaat	acaaaatagg	2460
cggggcgctgg	ttgcacatgc	ctgtaatccc	agctactca	gaggctgggg	caggagaatc	2520
gcttgaaccc	aggaggcgga	gggttcgtgt	agccgagatc	gggcacatggc	actccagcct	2580
gggcaacaag	agcaaaactc	catctccaaa	aaaaaaaaat	ggttcacat	cagtcctcag	2640
gaaaagatcg	atgtcgtga	ggggagatcat	ttcttgagag	cctcttca	gagtgggaga	2700
atgggctgt	ttttttgtct	tgttaaaaat	ctagaacggg	aagaacaatt	caaagggtgt	2760
ccaccattt	gtctgtaccc	aaccatggaa	ttactggact	ctttttaaaat	taaaaggtaat	2820
tcatgtttat	tctagaaaaat	tagggaaaaaa	aaatttttt	ttagatggag	tttactctt	2880
gttgcggcagg	ctggaggcgt	atggatgtat	ctcagctaa	tgcaacccct	gcttgcgggg	2940
ttttagcgac	tccctctgc	cageccctcg	ggtagctggg	attacagggt	cctccaccca	3000
ctcccgatct	atttttgtat	tttttttaga	ggcgggggtt	caccatgtt	gccaggctgg	3060
tctccaaactc	ctgacccctcg	gtgtccccc	aaccttggcc	tcccaaaaat	ctggattttac	3120
aagcataaag	cattggcgct	ggctgagata	accactteta	acatgcattt	cctccacac	3180
tgttcacata	tgcatattta	tctataaaaa	caataggggg	agggaaacct	aggtgaagtt	3240
tgtgtatct	ttttttttct	ttatgcctt	agaattttgc	tttttttagg	tactatgg	3300
aaagcaggatt	tgtatggctgt	tgtaggattt	tgaatatttt	atcccttac	tccccacagca	3360
ggacactttc	ccacccctccc	ttttggccca	gagaaggcgt	gccctctgtc	ctcccccattc	3420
ccatatgtgg	gcaactccccc	ccatggggcc	aaaccttactt	gggcaagttag	cagaggggaga	3480
gcagagtgtag	ccctgggggg	aggagggagaa	cttggagatgt	tttgggttgc	agatggactgt	3540
gtgtgtgtgt	gatggggag	catttttgc	catacatcttgc	cccggtgtgt	aggcatgtgt	3600
cttggatgt	tgctcccgaa	aagccctgtgt	atgtgtgtgt	gggccttgct	gtgtgaccaa	3660
accctggcca	ctgggtacgt	gaccctcaca	agtgtctact	gggctgtggaa	gagctccctt	3720
atgggcgtt	ttggagactt	atgtgtact	gttgccttttgc	gccatggggac	attaactatgt	3780
tacttttgc	cctcttaggt	tcagttgtcc	ttaattataaa	tacggggac	tgacttaggt	3840
gcgttgcgtt	cctttacgg	ccttgcacat	ttcttgcatttgc	ttctatgtat	atgtccttcc	3900
atttgataat	atgtttacgt	ggccatagtg	cctggggctgt	ggccggggat	ggaaaacttga	3960
tctctggggc	ctggggcgtt	aaggccatgt	atgtgtctgg	tgttgcagca	gatccgttaac	4020
tttccaaagag	gacatccat	aggctaccgt	gtccttttcc	actgtgtccc	tcctccattt	4080
catcttcttt	ataactacga	cttattgaac	atctactgttgc	tgctggacac	tttacaggtt	4140
atctcttaggt	tttacgataa	tcttgcagg	tatgcctgtt	ctgttttttgc	cagcagagga	4200
aatgagctgt	gtcagatgtt	actgtctgt	gcctcttggc	caggagatgt	gtgggttcaaa	4260
teacataggc	aggcgtatgt	aaaccttgc	gtctccaaatgt	cetctgttttgc	tgaccgttgc	4320
cttgcgtctgt	tttgggtttaa	aataaatgttgc	tttttggggc	ctactccaga	ggggcgtgttgc	4380
agggggctccc	tctcccaactt	cccccaaaac	cacccttttgc	cctgggtgttgc	tcaggaaatgt	4440
agagaactt	gctggggccccc	caggacttgc	tgagtggggat	aggggctgttgc	gaggtaagtgc	4500
tagagcctgg	cccaaaaattc	aggaggcccc	ccctggggcc	tgtgggttgc	4560	
gagggtgtt	ggggcgttcc	tcacttccctt	tttggacttca	ggccccagct	ctgggtttagg	4620
ccagggagaa	ccatccccaa	gttggatgttgc	ttactatatgc	agctgagatg	gatgggttgc	4680
tggaccaat	acatagtgttgc	gttaccgggg	ccaggggggat	gagggttgc	agggaaactgt	4740
tggggcaatgt	tctgggttgc	acctgtttttgc	tttttttttttttttgc	tttttttttttttttgc	4800	
gttggacttgc	tcagacgttgc	ggaggtgtat	gttggggatgc	tgtgttgcacca	tagaaatgt	4860
cgtgtttaaa	accaggcgttgc	cccttttgc	aaggccaggaa	gcatcattca	tttagccgttgc	4920
tgagaagaag	aaaccaatgttgc	tccgggttgc	agaccccttcttgc	ggggccccaa	gttggctgttgc	4980
taatgtgttgc	gactccaaatgttgc	ctgttttttgc	acccttttttgc	tttttttttttttttgc	5040	
ggaggccatgt	ttccatgggttgc	gttttttttgc	gttatttttttgc	atctgttttttgc	taccttttttgc	5100
cattggcctt	aatatgttaatgttgc	gccaaggcttgc	ttacatttttgc	ctgtatgttgc	aagaaggcttgc	5160
agggaaatgg	gtcatgttttttgc	tttccataatgttgc	attttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5220
ggggagatgttgc	tcttaggttgc	actcccttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5280
agcttggat	ttgggttgc	cttgggttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5340
aagcagctat	cagecccttgc	acttcccttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5400
ggacacttcc	ttctgccttgc	ctgcatttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5460
gggtggggaga	ggggggaaatgttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5520
ccaccccccac	ctggggcgttgc	gtcagatgttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5580
aatgtgttgc	gacacccatgttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5640
ctccaaaggat	ttcagatgttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5700
gcccttcaggaa	attcagatgttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5760
gtctggat	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5820
ctgtcttcc	ttggccatgttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5880
taaatgttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	5940
atacagtctgttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	6000
aaccacttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	6060
tccgttataat	acatacatgttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	6120
ttcgggtttaaa	tttaacatgttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	6180
tgggtgttgc	gtcagatgttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	tttttttttttttttgc	6240

-continued

taaaagtgg attcttactc cttcaggtt aaggcagtgc agtttgccca ggttagtgact	6300
gagacccagg tttcacac tctcccccgc agtggggatt gttttggggcc tttttcagcc	6360
caagagctct ctctccca tgccgctctg ctggctcgat attttccac tectctccct	6420
cctctggc tctctgacca gactcttagt attcaggaga aagtgttcat tgctctactc	6480
tctcatgtgg caatcaagta gtgccaagga gtgagagggt gaaggtgggt gggtgaggga	6540
cactcacccctt gotgagaaaag ggccccaggc tgttcggggtt attttaaagc agagacactg	6600
ccaggaaaaag ttctgacactg getgagaatc accccggggac caaccatccc gaatgcggat	6660
ccttgacact gggtgaggat ggagttgaa gatctgcattt gtaataaagc agcctagcag	6720
agtggtaag agtccagaca cactacatcg gtccaagggtt aacccgttgc taattactt	6780
tttggacttctt ttggacttctt cagttttttttt gggctttttttt gggctttttttt	6840
tttggacttctt ttggacttctt cagttttttttt gggctttttttt gggctttttttt	6900
ttggccaccc ttggacttctt gggattttttt gggattttttt gggattttttt gggattttttt	6960
ccacactgtca ccccccaccc accaggctag gagggttggattt attaggggtt gcccctgt	7020
tgacagctga gactgtcttcc tccctgttca ttccctttaatg acagtcttcc cttttccctgc	7080
tttcttggat tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7140
ttttttaaagg tggatgaggaa gaagcaggtt gttttttttt tttttttttt tttttttttt	7200
aggccgcattt tatgtatgac gttttttttt tttttttttt tttttttttt tttttttttt	7260
ttttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7320
tgcaactgtttt gatgtttttt tttttttttt tttttttttt tttttttttt tttttttttt	7380
tagcgcagggtt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7440
tcacatgtttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7500
ttttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7560
tgcagggtttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7620
gttacatgtttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7680
gtatttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7740
tcagggtttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7800
tattttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7860
ccttgggttcc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7920
gcaccctttgc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	7980
cacaacatcc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8040
atttggaggctt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8100
catagcagat tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8160
ttttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8220
aagagccacc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8280
attttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8340
ttttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8400
ggattttatct tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8460
ttttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8520
gaactcttgg tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8580
gtgagccgg tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8640
ttttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8700
ggatgttgg tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8760
ccttggggctt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8820
caaagggtttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8880
ggggggcttc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	8940
ggcatggccc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9000
aggatagggtt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9060
aatggagccc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9120
ttttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9180
ttttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9240
gttggatgtt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9300
agttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9360
tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9420
ctgttcttgc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9480
gggttggatgtt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9540
actctgttat tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9600
tgagactatc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9660
ccttagtatgg tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9720
ctgttcttgc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9780
aaagaagggtt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9840
gggttggatgtt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9900
gtgaagctca tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	9960
tcagcctctg tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10020
gacagacagcc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10080
gcagggtttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10140
tgttgatgtt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10200
cactgtttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10260
cggcccttgc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10320
taaacccatgg tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10380
gacactcaaa tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10440
cttagcccccc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10500
tttgacccctt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10560
cagctctctt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10620
caactgtttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10680
gtgtgtctcc tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10740
tgaggccctt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt	10800

-continued

ccatcagata ggctttctc tagaaacat tcgtgatctc tgagatttg ttccacttt	10860
gtgcttctgc acctaccatc aaacaccgg attgtatcat ttgtcacatt agatgattt	10920
tgttttgtt taagacaagg gtgtttcta ctcatcttt tateccccaga gcccagecatg	10980
atcttttgtt catabatagat gcaccacaga tglttgcga ttgaatgaat gagcacactg	11040
acagtttggaa gtcgcctga cttdcggtgo tatgcgtttt gccccctggg atgtgagtc	11100
cetcaggcoca gcccaggca aggccgtgc tgctccatc gtaactctca aggctcttg	11160
ttttatggca gtctttgtat tgacaggcat ctcttggaaagg ctttggggc aggacttgc	11220
tccaagtctc caggtgcctc ccagccacc cctgagtcctc caactgcctc tgctccatc	11280
gaaaatggaa caaggcttctt gtgtcttctt ttccagggtgc ttccagaggc agggctatgc	11340
tcacatttcat gccttgcgtac aagttgtgtga tgagcgaccc tccctaattgt	11400
cggctgtggaa cccacccgcg cgcttcgtgc aggaggccgac gcaaggccca gaggatggag	11460
agaacactgc caatgggtt ggtccccca gcaagctgggg gccttcaaac aggtccctgc	11520
ggctactgtt ctttacatgtt gaaaaccaga catttcatttc ctgtatgggg agggagaagg	11580
gaagatgtatg tggggatgg ccggaaagggt ggggtggcttccatgttgc cttttccat	11640
gcagggttcc ataggactgc gtcatttcac ccagatgggg actttggcactt gggctggaa	11700
acgctgttca ctcttgcacc ttgggtttac ctcttcatc caggttacttgc ttccactgt	11760
aataggagag ttgtttggta tgccctgggtg cttagacagg taacacagaa gctttaggt	11820
gtacgggggg aacgtttt tggcagatgg ccagacatgg taagtgtggagg agggactctgc	11880
ctgtatcacgc atggatgtt gatgtttttt tattttgggttgc tcaactgtatcattc	11940
ccagggggggg agattgttac gtttagaaaga gttagatattt gttttggggag ccacttagt	12000
gtgtcccttc tctcccgatc agggcagaac atctgaattt gctgaaccc tttttctgt	12060
tttgcattt atagaattttaaaaatgttgc ttttttttttgc agggactctt	12120
atccctgtt gctgaaattt gatgttttttgc ttttttttttgc cggaaaccatc	12180
gtcccttaacgc ctttgcgttac atacacccgg ctttgcgttcc actcttcagc	12240
ctttttttgtt ttgttatctc cggggagttt ttttttttttgc ttttttttttgc	12300
atgtatgttgc ttcttttttgc gtcgttttttgc tagattttggta ctcttcgttgc agaaacccgg	12360
tacttgcgttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	12420
cttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	12480
agatcggtt gtttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	12540
tgcctgcctt ctttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	12600
cttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	12660
ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	12720
acttcccttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	12780
cccatatccat ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	12840
aggggaaaatc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	12900
gccttgcacac ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	12960
tttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13020
tatctggaaa gcaacattca aacttctcat ttctgggttcc aaaaatccgt gcaatccat	13080
gtataggctgc ctttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13140
aggagaacgca gggggggggc ttttttttttgc ttttttttttgc ttttttttttgc	13200
ggcggccggcc aggccgggg gggggggggc ttttttttttgc ttttttttttgc	13260
tgtgttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13320
gcttcttacac agagaagaatggc gggggggggc ttttttttttgc ttttttttttgc	13380
gtgggggggtt gtttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13440
caagtttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13500
gtctacttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13560
cttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13620
ttagtgcgttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13680
cttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13740
aaacaaatca ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13800
caagtttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13860
tttagtgcgttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13920
ccccctcccc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	13980
tttaaagtttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14040
tcttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14100
gggggggggtt gtttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14160
ggcggggggcc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14220
tagtctcgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14280
caagtggccca ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14340
aaaatttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14400
gtgggggggtt gtttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14460
gtgggggggtt gtttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14520
gagcttgcgttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14580
tttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14640
acttggatgggg ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14700
gtgggggggtt gtttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14760
gtgggggggtt gtttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14820
ggatgggttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14880
cccttagccat ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	14940
catcgttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	15000
gtgggggggtt gtttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	15060
gttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	15120
cccttgccttc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	15180
aacagccccc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	15240
tcccttgccttc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	15300
tttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc ttttttttttgc	15360

-continued

cgatcatgac tcactgcagc	ctcaacctcc	tgggctcaag	ttctccacc	cagccctca	15420
agttagtgg actcacagg	tgcacccaca	tacctggcta	attaaaaaat	tttttttgt	15480
gcaggctaga ttcacatgt	ttggccagg	ttgtctcaa	ttccctggact	caagtgtatct	15540
cccacccctgg cttccaaag	ttctgggatt	acatgtgtga	gccatgtcat	ccagccctgt	15600
gtcttttaaa ttacacatt	atcccacttg	agttcctcat	tgcaagtgtc	caagcatcat	15660
tttcatttataa tcaaagtta	ttttgttttgc	tttcttttca	tttttaggtc	tttttaggtc	15720
cccteacccc gatactccc	ctgaagattt	atttttatgt	ttccctttcc	ttttcgcc	15780
aggatgtca gagggccatgc	tgagggtctg	cagccctgg	agacttttg	tttgtactgt	15840
cctatagctg ccgagtagcc	ccaggaggta	gtggaaagg	agatcccatc	tggccagaat	15900
catgggcaact gctgtcccc	aaagatgcca	taagctttt	gacageggct	tcaggcttt	15960
ctcccaggta aggggttggaa	ccccataacg	tggaaaggaa	attaagctgg	gcattacta	16020
ttttaaaaact gtttacacac	agggtcctca	caggatttt	ttttcaggccc	gtgtccatcc	16080
atggagcagg tagatagaag	tgcagagtgc	ccaggctaga	gggatgggac	aggggacatg	16140
caaggggggg gctgagcccc	cttccagccg	gggcagcaga	ggggaaaggcc	atgggagggg	16200
ctgcaggatg tgctgtgac	tgaaagctt	caacaagtaa	tgatgtaccag	ctgggcattt	16260
tggtgcacgc ctgtgttccc	aactacttg	gaqactgagg	caggaggatc	gcctgacccc	16320
aggagttcaa gtctagctg	ggcaatgtaa	gaccctgtct	ctaaaaaaat	aataataaaa	16380
taagtaacaa ttacatgtgt	aactgtgacg	aggcagggtt	tgaacatgtc	cgctgggagg	16440
ttggcagatg tggtggaa	gggtgggggg	ctgtgtgggt	ggagcagagg	atacagattt	16500
catggggtca agctagaaat	tgctggcag	atgtaaag	ctggcccccac	tgcgggca	16560
aggtgtctgg tggccagttc	cagaggctgt	gaagaggggc	tcagccatct	gtctgttagg	16620
gtttccctgg aggttccac	atacaggcag	atgggtgggg	ccggggcage	cagggtgggg	16680
ctggggatgaa aggggttggc	agggtccaga	ggcagccccct	ttcccttttgc	gtctgtgtgt	16740
cagcaggccg gtggggctg	cttttagtc	aggtagacca	ggggccacgt	gagggtcccg	16800
tgggctgagc tggtgactgt	tgagttggc	ctcaggggtt	aggctgggtt	gaagtgtatgt	16860
cactgtcccc ccgtatggca	gctaaggaa	tgggtagga	tcagccccc	ttgtccccc	16920
actctccat ctttggccag	gagaaggagg	acagggtttt	ctgaggact	gttgttagag	16980
ctttgggtag ggggggtt	cccggttct	ctgttggagg	cactctatct	aaaatagcac	17040
cccagtgtatccatc	tgtatctaa	cattatttt	ttccatggccc	tcatcattac	17100
ctgctgtatgtatgtatgt	tgtctatata	tcatctaa	ccccctcacac	tggaaacacaa	17160
tggccgtgggg cagagatctt	gctagcctt	gttccagagg	ctagaacagt	gcctggcaag	17220
taggagacac ccagcattac	cttttcaatgt	gaaccatgt	agatgggggg	agacccaaag	17280
gctatgcggg cagacccgt	ggagtcctgt	ctgcatgc	tgcaggatga	cctggggga	17340
actccttggc cttctgtgcc	ctctttatct	gtaaggtggc	cacccgtatcc	tttccagcgt	17400
aggcatgtaa tagccatgt	aaggccatcc	aggcttgggt	atcgttccca	ggatcttggg	17460
ggcccttagaa ttgtggcgc	ttggggacac	cttggatgtc	tgcataatttt	ttgtctatgt	17520
tcatccatgg ctgggttgc	atgttttca	tgatgtgtgt	tttccttcc	acctataat	17580
accttgggtt agtgcacat	aaggcagg	gtccctgggg	gcccccttcc	atgtggcaca	17640
agtggacatg ggcacatgg	cctggggcc	gaaaatgac	catcgac	cagtcttccc	17700
cagtggccagc ctgtttgggg	accggccct	ccgtccgcct	ctctcatgtc	cttgacacac	17760
ggggatgtggaa gtggggctgc	atggggac	acatgtttt	gtctgttcc	tgatgttttt	17820
tgaacccttc atggagaagg	ctctctgtg	accccagggg	gatagaaacc	ccccaaattt	17880
taacattctgat tttttgggt	aggctgggtt	actttctgggt	ttgtggggaa	attatctgt	17940
tctatcgccc ttgtttttgg	gatatcago	tgacccagg	gccccaaagg	acttggggga	18000
caagagaaaa cactttccca	aggacccat	ctatgtcaca	gggttccca	ggtcatgccc	18060
atgcacatctt ctgtgtatgt	ttccaagcat	ccccaccc	tttttagaaaa	tgctgcaaat	18120
gtgtttatgtt aaggacatgt	aaggatgggg	aaggaaatgt	tagttaaagg	ggccagggtt	18180
ggactgtatgtt ggttttttttt	gttccatccaa	gttccatccaa	gttccatccaa	gttccatccaa	18240
ccacccctgtt ctgcagggtt	agtgtcaag	acccatcaat	tttccatccaa	tttccatccaa	18300
ctcttgcgtca ctgtctggaa	cttcggggca	gtgggcattt	tgtgcattca	cttggaggcc	18360
cctctgtgc tgcacccagg	cttccatccaa	atgtatgtt	ctgcatttgc	cttgcatttgc	18420
atcaagatcc tccccaggat	gtcccggtt	gttccatccaa	gttccatccaa	gttccatccaa	18480
agggtggggc caagggtgtt	ggggccatgt	ggggccatgt	ccaggcccaa	atcgccccca	18540
gtgtgcaca aggaggccag	gttccatccaa	gttccatccaa	gttccatccaa	gttccatccaa	18600
gggatccctt ctgcaggatgt	ccgccttgc	aaacagagg	gggtccacta	tttctggaa	18660
actcctgtgtt gtcataat	aacgcgtat	tcactgtatgt	cctccatccaa	tttttttttt	18720
tttggatgtt agtcttgc	tgtcccccac	gttccatccaa	gttccatccaa	gttccatccaa	18780
ctgcacaccc cgcctccccc	gttcaagtga	tttccatccaa	tcagccctt	tttttagatgt	18840
gattataggc atgtggccacc	acgttgggtt	atttttgc	tttttagatgt	agatgggggt	18900
tcaccatgtt ggccagggtt	atctogaact	cctgcacttgc	tttgcatttgc	tttgcatttgc	18960
cccaaatgtt tggggatata	ggcatgtat	actacatccca	gttccatccaa	ccattatccat	19020
tgctatgtgtt attgatgtt	tgcaatgtt	agcaatgtt	gttccatccaa	tttttttttt	19080
atcaagacatcc tccctttgg	accatgtat	tacataacc	tttttagatgt	tttttagatgt	19140
atattatgtat gtttgcattt	ggccatccat	tttttagatgt	tttttagatgt	tttttagatgt	19200
ggccatccatc agtggggact	cccacccat	tttttagatgt	tttttagatgt	tttttagatgt	19260
tgcttgcgtt tgcctgcgt	ggccatccat	tttttagatgt	tttttagatgt	tttttagatgt	19320
acttggggat gagaatttgg	gttccatccaa	tttttagatgt	tttttagatgt	tttttagatgt	19380
ccggggctcc tgcgtacat	ggcaggctt	tttttagatgt	tttttagatgt	tttttagatgt	19440
caaggatgtc tctgtttttt	tagatgtatgt	tttttagatgt	tttttagatgt	tttttagatgt	19500
aatgtgtt gaaactgtcc	tttttagatgt	tttttagatgt	tttttagatgt	tttttagatgt	19560
atgtgtatgt gccccccgtt	tttttagatgt	tttttagatgt	tttttagatgt	tttttagatgt	19620
acaggggctt gtttcttggc	tttttagatgt	tttttagatgt	tttttagatgt	tttttagatgt	19680
ccacttttcg aagccaggga	tttttagatgt	tttttagatgt	tttttagatgt	tttttagatgt	19740
ttaggacagg aagcaggat	tttttagatgt	tttttagatgt	tttttagatgt	tttttagatgt	19800
tcctccagcg gcatgtgtcg	tttttagatgt	tttttagatgt	tttttagatgt	tttttagatgt	19860
cttcctccaggc tcttgcata	tttttagatgt	tttttagatgt	tttttagatgt	tttttagatgt	19920

-continued

gcaaggagct	tctgggcttc	cttgacagca	gcgtggctga	ttggcattaa	tcctaactga	19980
agggaaggca	cacgggatgg	ccccctggct	cggggtaaat	gtgttagat	ttggacttac	20040
acatgcagtc	aacaaggca	catcaagtcc	ccattttgg	acaggcactg	tgcttaggcata	20100
tgggggaccc	agcaggaaag	aagaccacag	ggtcccgaggc	ctcatggagc	tcacggccct	20160
gtgattgtga	tgccctcggt	ctgttgcattgg	cggggcttaa	atagctgaa	tttctggagc	20220
tctggcgtct	gcaagggtggc	ctgggaaaga	gtttatggaa	cagctacaga	gttcttaggt	20280
ccttcatgca	gttgaggatt	cgagcccgta	gaggagaata	gcctgcaggc	tgccccccacg	20340
ggaagcaca	ttccaggcgc	atcccgagg	tgacggagaa	ccatgtatgg	aaaggtagtg	20400
ccaggactgt	catgagtgtc	ccagggtcg	ggggattcac	ccgtgaactg	tgagggtttg	20460
getctgtatg	actctgttct	tatgttttag	gaggggagaa	aaacgtaac	agaatagaca	20520
aatgtcaagag	agagtgtact	tggaccctc	ccacaacggc	ctccaaaca	tggagcatga	20580
gcagataacct	cgaggatgg	gggtctgtg	caggcttct	gggacgcaga	ctggccacct	20640
cccccaggcc	ctgcaggcag	ccactgttag	caccgcctga	gatgtgaacc	tttctctc	20700
cccccagtcg	catgggtgt	acgggtggca	tggcgaagct	ggacccccc	tctcagggtg	20760
ccctccagct	ccccatcgc	ccggagatgt	gtgagtagat	tggggagatc	acagectctc	20820
atcaactgggg	ggcagtcctc	tacccgtacc	cagctctgt	ccgcctggc	tccctqaqag	20880
gcatgagttc	aggaggggca	gaggggaaagg	tcggtaaaa	accagccga	cacatgcggc	20940
ttgaagatc	accaagtgtt	ggacccttgc	tcctctgc	gctctgtgt	categttctg	21000
ctggggctgg	tggggtggag	tggggggaa	cctgggtgt	ggtgtgtgt	ctcaggggga	21060
cccccttctt	gagttttgtt	ccctgttaad	actctgtacc	gtgtttgtt	ctctctttt	21120
ttgtccccctc	ctcacggtga	tgacggacat	tttctttcc	tggacaccca	gaagaagact	21180
cctatgacag	ttttggggag	ccttccatacc	ccgaagtctt	tgagccccc	ttgactggct	21240
acccagggg	ggagctggag	gaagaggagg	aaagtaggt	gcccattgttc	acacggctcg	21300
cttcagccct	ccgcggggc	ggagacagg	gggtggggct	ccctgcaggc	tgggtggagg	21360
agggcatgag	ggggagggcc	ccttttccca	tcaaggccat	ctctgtgaaa	gtagaagatg	21420
cctgcagcgc	tggggttctt	tcagcaggcc	ccatgtatgt	gtccggatgt	tattgagat	21480
ggcccacgtg	ccctgtgtgt	gtctgggtgt	gcccaggccc	gggtggggcc	acaggcttaag	21540
gagacacggg	cgtaaatcac	atagactgag	aaggccaaaga	ctatgaaagg	ggccatgggg	21600
tttggggagg	ggggcaggag	agcatgccc	ggggcttctt	gacctgtgt	gcaggggatg	21660
gagaaagtca	gttgaggaaag	taactgtgt	gtctggatct	gaagggttag	tcacagcagt	21720
caactaggaa	gaggagcaca	gggtgggggg	catttcttgc	cagacagact	caggaaatcag	21780
aggaaggccgg	ggccggatgc	agagacaga	agtgtgggg	agccctgcaa	acaggccctgg	21840
agacatcgca	agataggagt	tcatcttgc	gtcagtagac	ggtgccctgg	taacacccaa	21900
tgccagccca	ctgctcggt	ccaggcagca	ccctggagca	gggagatgt	gcactgtcgt	21960
aacagccct	gcctctggag	gtgccttacg	ggagcagctg	ggtgacatgt	gttggcata	22020
caggactcca	tgacacacggg	agggggaa	tagggaaaga	tcaactctgc	gtgggtctgg	22080
aaggaggagc	aggtggccac	cctccaggca	ggcttggggg	aggtatttat	tccaaggccaa	22140
actgggtgtc	tcagagaccag	gagttagcac	agatcccacg	gggcccgcag	gactggcttc	22200
cctccagacca	ccagccacaa	gctctagagg	gtcttagat	cacttgtgt	tctgaccggc	22260
tgccaaat	gggcttccat	gaccccccata	gttcaataaa	tttgcataaa	tgactcacag	22320
aactcaggaa	agcaactacac	ttaaaatgtc	agttgtttt	ttgttgtgt	tttgttttgg	22380
agacagggtc	tcgtctgtt	gcccaggctg	gagtgcagta	gcacgattgt	ggctactgc	22440
aaccttgcact	tcctggctc	aatgtatcc	cccccattttg	cctctgtatgt	agctgggtat	22500
acaggcacgt	gttaccacac	ctggcttatt	tatattttt	gtagagacaa	gttttgtact	22560
tgttgcgggg	gtgtttttcg	aactcttggg	ctcaagttgt	ccacctgcct	tggcttccaa	22620
aagtactggg	attatagggtc	tgagccacag	caccggccaa	aaattagttt	tattataaga	22680
gatgcacact	aggaggacac	aaatgtggaa	acagtgtggaa	agtaatgtgt	atggatcaca	22740
cctgggtgggg	gaggaggagc	agctggggcc	aggaggagg	gggacacccgt	cagggtgtgg	22800
agggcagggg	agggtggccct	ccatgttttg	tgttttatgc	ataaccattt	ttatgtctta	22860
cagtggagca	agtttaccta	taaacaagtgt	tcagggacc	ttgcactaaa	gaaaacaac	22920
gagggcattt	tggaaggtct	aattttcttgc	tcagttgtgt	gtgactaaat	tcccagtat	22980
atttacctgt	tgtaagggtg	aaaggtttgc	aggaggatctt	tgtttttgtt	ttatatggc	23040
tttgaatgt	actgtaaaa	aattttcttca	aaatctgttgc	ttcaqactt	atactaaat	23100
gtacagcagt	gccccagccca	aggecttgc	tttcttatttt	ttgttttttt	tactctctaa	23160
gtgccccaa	ctgggtttac	ctgagttca	gaactgcggc	ttttttcttg	cccagggtgt	23220
aagtccaccc	gtccacagggt	gttccctgt	tttccctactgt	ccactgtt	ggggaggccag	23280
ctgtccatgt	ccccaggcc	catcttagct	tctagaggcc	agggtgggtt	ggctgggtt	23340
ggcaagagca	gttggccctt	ctggggccaga	gtttcttctt	ttttttccatt	ctgtgcacgc	23400
cttttca	cggttactt	tcttccctca	cacaggggcc	gtgaagctg	gcctcgggga	23460
cttcatctt	taacgtgtc	tggtggccaa	ggggctgtcc	acgggcaggc	ggggactggaa	23520
taccacgtc	gtctgttct	tggccatctt	catttgtgt	gtctggggat	ggctccaggct	23580
gcctcggtt	ggggggccccc	agggtccatca	ttgtgtgtgg	ggcagggttcc	aggatccata	23640
gggatttttc	atttcttctc	ttccctcttgc	gggacaagag	caggggaggcc	ggcttggaaagg	23700
gtcagcttgc	gaccaaggct	cacagggtt	gtgttgcggc	ctagggtggc	tccaggctgt	23760
ggaggacagt	gggggggggg	gtgaggatgt	tacccggcccc	agcgttggct	agcacacacg	23820
ctccaggcc	aggaccaccc	tgacagctt	gcccgtgt	gatacttctg	aggtgtgtgt	23880
gatgacatca	gatttgcaga	aaagaaaatt	gtttaagggtc	cttgcctat	ggcgcacaaac	23940
tagtggagac	catgttttcc	ccctccctca	tgccatttgg	acaccacagg	gtctgaatct	24000
ggggcactag	gggtggccccc	gttactgtgt	accacagca	tgaaatgtgg	aggccctgtt	24060
gtcagttaac	gtgaccat	acacataat	ggggacacgt	ctgcctgt	ttcatctcag	24120
agatttcgct	gtcacgttag	aggaggatgt	gtgttgcggc	ctgttgcgtt	gcatcttccc	24180
cttagtgaaa	accctggggc	ttggatgtt	aagggttgcatt	ctccagtgtc	cagaagggttt	24240
tcttttttgc	cacaagtata	tcaggatgtt	gatgttgcac	ccaggctctt	ccaccacccag	24300
actgccttac	ctgagccctg	ctggcccaa	agatataat	ggcaccctgg	ttccctgtc	24360
tcacccatgc	cactgcctc	atcagctggg	tcaggggagg	atggggc	cccacac	24420
cttccca	ggaggatgtt	gttcccttgc	gttcccttgc	gttcccttgc	gttcccttgc	24480

-continued

tactgccatt	tcagttgagc	ccaccttagct	ctcatataaaa	tacatgttcc	ctgaggggcat	24540
cttaccatcc	catgtgacca	ctccagccag	acaggggagg	cagcacggcc	tcggggcacaca	24600
gcactgtccc	aggagtccagg	aggectgcct	tctgggttca	tcactaacag	gtgagggtgat	24660
ctaatggggg	tgagaacttc	tgcccttaac	acctaagag	ctgttgagg	accaggaaag	24720
ataatggggt	gtctagcgcc	gttatccgac	tggtccctcg	acaagctct	gtgcccaggg	24780
actagaccat	gactcacagc	tcctgtccac	accagggatc	accacgctca	ccctccctc	24840
catgtcttc	agggcttgc	tetgaccctc	ctgctgttgc	ctgtgttcaa	gaaggcgctg	24900
ccccccctcc	ccatctccat	cacgtcggg	ctcatcttt	atctctccac	ggacaactcg	24960
gtgcggcgt	tcatggacac	cctggctcc	catcagctct	acatctgagg	gacatgttgt	25020
gccccggct	gcaagctgca	ggggatttt	atggatgca	gttgtatagt	tttacactc	25080
agtgcctat	attttttaaga	ctttttttc	ctaaaaaaat	aaagtacgt	tttacttgg	25140
gaggaggagg	cagaaccage	tctttttgtc	cagctgtttc	atcaccagac	tttggctccc	25200
gctttgggg	gcgcctcgct	tcacggacag	gaagcacage	agggttattcc	agatgaactg	25260
agaagggtcg	attaggccgg	ggagaagago	atccggcatg	agggctgaga	tgccaaaga	25320
gtgtgtctgg	gagtggcc	tggcacctt	gtgctctgg	tggagaggaa	aaggccatc	25380
cctacagagga	gtgttccca	tgtttgtcc	atgtgtct	tgttattttt	ttgcctttag	25440
aaactgagtc	ctgttcttgc	tacggcagtc	acactgtgg	gaagtggctt	aatagtaata	25500
tcaataataa	gatgagtctc	gttagaatct	tggagtttg	tccgtgttaa	atgttgaccc	25560
ctctccctgc	atcttggca	ccccctggat	aacttgcgt	gtgagccacag	gatggaggca	25620
gtttgccttg	tttggaggaa	ctttatagt	tctccctct	ctgcacacat	ttcttaact	25680
agaaagttc	ctaagcaaag	gagtttaggag	agcaggggtgg	cctgacatct	gccagccctg	25740
agctgttaagg	ctgtgtatgc	tgagcagggt	cctggactca	gttgtca	gtggcacaga	25800
caactgccc	tgggtggccaa	aacatccagt	ggttcccttc	goaagtgttc	accctctgca	25860
gaagcctgt	aggggctctg	ctcagaaacc	actctctt	ccttcttgg	ctttggccct	25920
gggactgt	gtggggaggt	ggagacattt	gcttgcctt	ctctgtacat	caatcatggg	25980
ttgcaaaagag	aatctcagaa	gtgcctctt	ctgagcacag	tggctcacac	ctgtaatccc	26040
aataacttgg	gagggttgcgt	cggagaggat	acttgagocca	aggagtttg	gaccagccgg	26100
ggcacaatcag	tgagacttgc	tacaaaaaaa	aattttaaaa	tttagcca	atgggtggcat	26160
gcacatgttag	tctcagctac	tctggaggt	gagggtggag	gatcacttaa	gccccaggagg	26220
ctgaggctgtc	aatgagccga	gatcaagcag	gtgttagtta	tatcagcacag	ctgagaagac	26280
gcaagtgtgc	ctgggggttc	aaactgtgtt	ccctgtctcc	ctgttccagg	aataacatga	26340
gtggggggac	aatgcattt	tattatgaga	ggaatgagaa	ttgtgtatct	tgacatttga	26400
caaggagttt	ctttccccca	ggctgtttga	ggaagggcag	aggaaaaatgt	ggtgcctaa	26460
gaaggaagga	cagaggaggc	cgaacactgg	cgggtggaa	cccactgatt	agttagtgcag	26520
gtcagagacc	tggggatgggg	ggcattgtcc	tcatggaaac	cacagcgggg	agcgggtaaa	26580
gcagacaggg	atggtccctg	atggtacaca	ctcgcaagag	gtaaggggaa	aagaaaaact	26640
gaaaaggctt	ttaattttgg	caattatgg	agtgtttatc	ttcagaaagag	cagttttagg	26700
gtgggggttc	caaagatggg	attggacata	tattttgaat	cattaagctt	gaggtcttc	26760
aaaggccctgg	ccaaagggttgc	ctgggtggag	accacattca	gaggtaaagg	cagaatattgg	26820
ggggcccttaa	tgtagacagcg	agggagggaa	aatgaaagg	gcctgggtat	ggttaggggt	26880
aagtgttaag	actgagaaaa	caaggacatg	tgagaagacg	agggaaagac	attggagaga	26940
acaaagacac	tggaggagat	gtctatttgc	ggtccccaga	gagcaggggag	acaaatgaac	27000
ccagaacaca	aatggcaaag	aagaaaaatgt	agagaatttg	taaaagacag	cattcgaaca	27060
tgcgcgaatc	gaggcaggta	ctgggttca	aacacctgtt	tctcccccgt	taaaccgcgtc	27120
aactaatatc	tttccatatt	tgcctcagat	ttgtctttag	aaataaaaaacc	cacgttctga	27180
agtctgtttt	gtatgtggcc	ccagtcctgt	tgctccgc	tcctgtctt	aagtgcattt	27240
ctgccccttc	catctatgtt	tagttttgt	ttgtatgtt	gtatgtttt	tttaactttac	27300
agaaatgtt	tcataactgt	catattgtat	aattttttaa	aattttgtat	tctggaggca	27360
tgataataatg	tagtcctagt	tcatttttt	tattttttt	ttgagatgga	gttttgcct	27420
tgtcaccctcg	gttagatgtc	aatggcgtga	tgttggctca	ctgcaaccc	tgcctcttgg	27480
gttcaagcaa	tttccctgtc	tcaattttct	gatgtatgtt	gattacagg	gccccccacc	27540
atgcctgtgt	aattttgtat	tttagtagat	acggggttt	accacgtttag	ccaggctgtt	27600
ctcaaacttc	tgcacttcgagg	tgatccacgc	accttggcc	ccaaaatgtc	tgggattaca	27660
ggcgtgagcc	accgtcccc	gcccaggat	ttaactatt	gtatagtgtt	ccattgtatg	27720
agtttcaatc	tttatatgtt	attgtatgc	ctgttgggg	tttgcagtgt	ttctgttattt	27780
cagctgtgt	cgagtgcag	teccatcaca	ttgtgtggat	ttgaggaaat	atttggaaatc	27840
ccccatgtt	ctggacattc	ccaaatacc	tccaaatgtt	tgctgttta	tccttccatc	27900
cgcaatctga	gagttcccc	acttataat	acttgggttc	atcagactt	tcatcttgc	27960
tgttggatg	gtgttcat	cttttaggtt	tttatttt	tttttcatat	gtgttattgg	28020
tgtacaagg	tccttcctgt	tttattttt	tttattttt	tagacagat	ctcgcgcgtgt	28080
cgcccgaggct	ggagggtcgc	agegtgtat	ttggctactt	taagctccgc	ctccgggtt	28140
catgcccc	tcctgttca	gccttcttgc	tagtggggat	tacagggttgc	tgccatca	28200
cccgccgtat	ttttttgtat	ttttagtata	gttgggtttt	caccgtgtt	gccaggagg	28260
tctcgatcc	ctgacccgt	gatccaccc	cctccggcc	ccaaatgtt	gggattacag	28320
gtgtgtatgc	tcgcggcc	cccaatgtt	tttcttctgt	attcttctat	atccctgtcc	28380
catttttcc	tttggattttt	tgttcttacgg	atattttaa	tcctttaat	atatatttt	28440
gagagatgt	aatcttgtat	taattttat	tatttgcata	gtctgttaca	tttgggttt	28500
cctctcttcc	ctgcctttag	gagtttttt	ctggacccaa	gtatttttt	aatgttata	28560
tttattttat	tatcaat	tttgcgtt	tttgcgtt	tttgcgtt	tttgcgtt	28620
atcccttcct	accctcaagg	ttttttttat	tttttttttc	ataataat	ttttttttt	28680
tctgtttatgt	atttttatgt	tttgcgtt	tttgcgtt	tttgcgtt	tttgcgtt	28740
cagggtgtcc	agcaactgttt	actgaacatgt	cttccttcc	tcgttgcgtt	gtagaactct	28800
cctgacat	accaatgtt	cataagtggg	tggatgggtt	cctgagctt	ctactgttta	28860
tagaacttgc	tctctcgca	gccaatgtt	caccagggtt	tttgcgtt	tttgcgtt	28920
tggtggaaagg	agaagatggg	ggctgttctg	agagttttct	tttgcgtt	tttgcgtt	28980
gagggtatgt	aatatctgaag	tttacacttgg	tctccaaaaa	cagtataaa	ctacagaatgt	29040

-continued

ataatagtgt	ggaatggtgg	tgggagtcag	taagggttag	gtcaactgcag	tggttaaac	29100
aagatggct	agaatccctt	cacaggcaca	ggcagcttg	agaggggtca	atagtgcatt	29160
gtatcagggg	tcagatgcct	cttttccctt	tgagatcagt	aagtggctt	caccctcatga	29220
ccttaggtgg	ctgctgtgt	ctagccgtca	agtcacactc	catccacat	gaaaggaggt	29280
tagaaaagg	tgcatattcc	cttcttaaaa	acatgtctca	aagtgcaca	cagcactttt	29340
gcctatattc	aattggccat	tagtcccac	gccatatacc	tctgagactg	agagactggg	29400
aaatgtctt	atttcaagtg	gccccatata	ccatctaa	agataaggga	tacgtgggta	29460
tggcgtgtct	tttggtttac	caatgcagat	aatgaagttt	ccaaaaacat	gagaaaatgg	29520
ggtcgtgagg	gatcatgtga	atcacaagct	gatgtcttca	aagacgggtt	aatggggccc	29580
cgggaggcag	cagatgcacag	cgtggggat	taaggtagac	ctccatctg	gggttaaaat	29640
gaggggaaagg	tgtggagat	ggaccacac	tcagaatgg	cagtggttt	gagaccctt	29700
gccccccacc	gtgcaccca	ttggctctt	acagaatgcc	tgcgagttgc	tttagagtac	29760
caaggatgag	gtgcagatcc	atgtgcaccc	ccctggccccc	tctgtggaca	attttcatgc	29820
ctgacagcac	agtctatgtg	gattgcacag	cgtatggaaat	atgcacaaat	gaagcatgc	29880
tgcagtttgt	tgtcggtga	tgtgtttat	gcttatgtga	gtcgaatggg	gcggcagggt	29940
cctgtgttca	cccgctgaga	aggagggtt	ctgtacacc	tgcctttt	tcagctactt	30000
gagaaaagg	ttgtgggg	ccgtggattt	tgggacagct	ttgaatgg	gtagggagga	30060
agggtcccg	ctgagtgt	ggccagaaag	ctgtggggaa	gtttttaga	cattggccaa	30120
gagctccctg	aaggcagcca	ggggatgtt	tgtgactaa	ggccaactgt	30180	
ataatacgac	aagtgtgt	ttgtgtgt	caacactgtt	caccttaga	aggacctcg	30240
gacagtggtt	gtggactctg	tagagagta	cagtgcacgt	agcaaacctt	tacccagtc	30300
caacccctgt	ctaggctcgc	actaaatgt	tttaccttca	attctcgtaa	caataggagg	30360
taactactat	tctaaattcc	attttata	tgaggaaat	aaggcacaga	gatcactgac	30420
ttggccaaaa	tcaagcagg	agtagttgt	atataagccc	acggatgtt	gtttgttagaa	30480
taggtgetct	tgactagcag	aaataggtt	tccctgcagt	gtgtatgt	taacaagcat	30540
gggctgccc	cttcctgtcg	aggccacta	aaacacccaa	caggctacgc	acggtggct	30600
acacccctgt	tcggcggact	gtggggggcc	gagggtggcc	ggtcacccgt	ggtcaggagt	30660
tcgagaccc	cttggccgc	atggtaaaac	tccgtctcta	ctaagtcgt	aaaaatttagc	30720
tggcgtgtt	ggccggccacc	tgtaaatcc	gctactcagg	aggctgagac	agaagaatca	30780
cttgaaccag	ggaggcagag	gttgcagtga	gacaagatca	cgccatttgc	ctccagcct	30840
tgtgcacaaa	ggggaaactgt	ctcaaaaaaa	aagtatgtt	ttataatccc	30900	
agcacccctt	gggggtgt	gtcgaaatc	acttgagcc	aggagttaa	gaccaatcta	30960
ggaacacatgg	caagacccca	tctctgcca	aaataaaaaa	tagtctaatt	ttagtttac	31020
atgtgtgt	gaaatgggt	cttccctgtt	cttgcatttgc	cattttccca	atgctgacta	31080
atgacgtttt	gcacccttgc	atgtgcattt	tggtcagat	tctttttt	gttacattt	31140
attaagtttt	aaaataaaaa	gtcaagattt	tccctatgt	aatgactttt	aaaatgacca	31200
aaaaggggaa	gataacatta	attcttgc	agaaggcc	ttgagaaaaat	acagttgt	31260
caagctgtt	ctttgcataat	gacccatgt	ttttatattt	ccccatagga	aggccaagga	31320
agagtcttt	caccttcagg	caggatgtt	aggggacttgg	gtcattttat	aagagtgtt	31380
ggtttggaaa	ctcaacccca	gaagacttct	tagatttt	cccaggatgt	agggaaagggg	31440
ccgcacatcc	ggagagagga	ggatgtact	tagagcattt	gtccccatc	ttttagggacc	31500
agggacttgt	gtcatggtag	acagttttt	cacagatagg	gttgggggg	atgatttgg	31560
gtctgaaactt	ctccacactt	ggtcatcagg	cattagattt	tcatgtgg	tgtccactt	31620
agatccctgg	ctgtcagact	tcacatgg	gttgcagg	ctatgagaat	ccgtgcac	31680
tgatttgaca	gggggggg	ctcagttgtt	atgtctatc	tccacccgtt	accacctgt	31740
gtgcagctt	gttcctata	ggctatagat	tggtaactgtt	ccatggctt	gggggtgggg	31800
accctctttt	tagaggtttt	aagggtatgg	cttaccgttgc	ggccctgggtt	gttctgggaa	31860
tgggggggg	gggggggg	gaggatgtt	ggaagaccc	cccttgc	ccatttggg	31920
tttggggaga	aaatcggtt	tcaggcttca	cagtacccgt	tcctgttacca	tcttccaaag	31980
ggaagtgttt	gggggtgg	ggttgcagg	gttgcattt	tctgagcc	ggcacaagac	32040
agaatgtttt	caccatcc	gaggggggcc	gttgcatttgc	cccaagcaga	gagccatgc	32100
tccctctctg	aggcccttgc	tggatgggg	tttttttttt	actgagccaa	gggtctgtca	32160
gtgagatgtt	gggggggtt	gttgcatttgc	aaggccatgt	tataacccaa	ccaggaaat	32220
aggttccgg	tcttgcac	aaggccctac	agttgcac	agagatggac	agccaggaga	32280
ccccaatgg	ccacccatgg	ccacccctt	ctgccttacc	caccctcc	tactccat	32340
cctactcggt	ggggggcc	aaatcgat	gttgcatttgc	cccaagcaga	gagccatgc	32400
tagaaaaata	tttggggat	caggaaat	tatttttact	tttccatgtt	tataatccata	32460
ggctgtttat	tatcaaata	attttttaaa	attgtataa	acatttctat	actaaaacac	32520
tgagttttt	tcttttattt	tttttttttt	tccagatgt	tttacttctt	32580	
caatgggtt	tctcgat	tttataat	tttccatgtt	ccatccatgtt	32640	
cttgtgtttt	tttttttttt	tttttttttt	tcagcccttt	tttctgtttt	32700	
taagaaaacc	ccccatgtt	tgtgttctt	cagacatgtt	tttgcatttgc	32760	
cagagcaactt	aggcccttta	tttttttttt	tttttttttt	tttttttttt	32820	
gacccttccaa	tttccatgtt	tttttttttt	tttttttttt	tttttttttt	32880	
gattaatcg	tttttttttt	tttttttttt	tttttttttt	tttttttttt	32940	
tttccctgttt	tttttttttt	tttttttttt	tttttttttt	tttttttttt	33000	
ctgatctacc	tttttttttt	tttttttttt	tttttttttt	tttttttttt	33060	
gtgtttactt	tttttttttt	tttttttttt	tttttttttt	tttttttttt	33120	
tctggaaat	tttttttttt	tttttttttt	tttttttttt	tttttttttt	33180	
tttccatgtt	tttttttttt	tttttttttt	tttttttttt	tttttttttt	33240	
aaattaat	tttttagag				33258	

SEQ ID NO: 2 moltype = AA length = 448
 FEATURE Location/Qualifiers
 REGION 1..448
 note = misc_feature - presenilin-2 isoform 1

-continued

```

source          1..448
               mol_type = protein
               organism = Homo sapiens
SEQUENCE: 2
MLTFMNASDSE EEVCDERTSL MSAESPTPRS CQEGRQGPED GENTAOWRSQ ENEEDGEEDP 60
DRYVCSGVPG RPPGLEEEELT LYKGAHKVIM LFVFPVTLCEMI VVATIKSVR FYTEKNGQLI 120
YTPFTEDTPS VGQRLLNSVL NTLIMISIV VMTIFLVVLY KYRCYKFIHG WLIMSSLL 180
FLFTYIYLGE VLKTYNVAMD YPTLLLTVWN FGAVGMVCIH WKGPLVLLQQA YLIMISALMA 240
LVFIKYLPWE SAWVILGAIS VYDLVAVLCP KGPLRMLVET AQERNEPIFP ALIYSSAMW 300
TVGMAKLDPS SQGALQLPYD PEMEEDSYDS FGEPSPYPEVF EPPLTGYPGR ELEEEERGV 360
KLGLGDFIFY SVLVGKAAAT GSGDWNTTLA CFVAILIGLC LTLLLLAVFK KALPALPISI 420
TFGLIFYFST DNLVRPFMDT LASHQLYI 448

SEQ ID NO: 3          moltype = DNA  length = 19
FEATURE
misc_feature          Location/Qualifiers
1..19
note = Forward 5'-3' Primer
source                1..19
mol_type = other DNA
organism = Homo sapiens
SEQUENCE: 3
catcagccct ttgccttct 19

SEQ ID NO: 4          moltype = DNA  length = 20
FEATURE
misc_feature          Location/Qualifiers
1..20
note = Reverse 3'-5' Primer
source                1..20
mol_type = other DNA
organism = Homo sapiens
SEQUENCE: 4
ctcaccttgt agcagcggtat 20

SEQ ID NO: 5          moltype = DNA  length = 26
FEATURE
misc_feature          Location/Qualifiers
1..26
note = Forward 5'-3' Primer
source                1..26
mol_type = other DNA
organism = Homo sapiens
SEQUENCE: 5
acagaatttcg ccccgccctg gtacac 26

SEQ ID NO: 6          moltype = DNA  length = 25
FEATURE
misc_feature          Location/Qualifiers
1..25
note = Reverse 3'-5' Primer
source                1..25
mol_type = other DNA
organism = Homo sapiens
SEQUENCE: 6
taagcttggc acggctgtcc aagga 25

SEQ ID NO: 7          moltype = DNA  length = 20
FEATURE
misc_feature          Location/Qualifiers
1..20
note = Forward 5'-3' Primer
source                1..20
mol_type = other DNA
organism = Homo sapiens
SEQUENCE: 7
tcagcatcta cacgccattc 20

SEQ ID NO: 8          moltype = DNA  length = 20
FEATURE
misc_feature          Location/Qualifiers
1..20
note = Reverse 3'-5' Primer
source                1..20
mol_type = other DNA
organism = Homo sapiens
SEQUENCE: 8
agcaccacca agaagatgg 20

SEQ ID NO: 9          moltype = DNA  length = 26
FEATURE
misc_feature          Location/Qualifiers
1..26

```

-continued

```

source          note = Forward 5'-3' Primer
                1..26
                mol_type = other DNA
                organism = Homo sapiens
SEQUENCE: 9      attcgccccg gcctggata ctgcca                               26
SEQ ID NO: 10     moltype = DNA  length = 27
FEATURE          Location/Qualifiers
misc_feature     1..27
note = Reverse 3'-5' Primer
source          1..27
                mol_type = other DNA
                organism = Homo sapiens
SEQUENCE: 10      ctgtccaaagg agctgcaggc ggcgcag                                27
SEQ ID NO: 11     moltype = DNA  length = 25
FEATURE          Location/Qualifiers
misc_feature     1..25
note = gLN141I guide RNA F
source          1..25
                mol_type = other DNA
                organism = Homo sapiens
SEQUENCE: 11      caccgcatca tgatcagcgt catcg                                25
SEQ ID NO: 12     moltype = DNA  length = 25
FEATURE          Location/Qualifiers
misc_feature     1..25
note = gLN141I guide RNA R
source          1..25
                mol_type = other DNA
                organism = Homo sapiens
SEQUENCE: 12      aaaccgatga cgctgatcat gatgc                                25
SEQ ID NO: 13     moltype = DNA  length = 100
FEATURE          Location/Qualifiers
misc_feature    1..100
note = Donor ssODN#A N141I
source          1..100
                mol_type = other DNA
                organism = Homo sapiens
SEQUENCE: 13      gagagaagcg tggctggagg gcagggccag gcacctcacct tgttagcagcg gtactttag 60
                  agcaccacca agaagatggc cagggtgttc agcacggagt                                100
SEQ ID NO: 14     moltype = DNA  length = 21
FEATURE          Location/Qualifiers
misc_feature     1..21
note = Forward 5' Primer
source          1..21
                mol_type = other DNA
                organism = Homo sapiens
SEQUENCE: 14      taacggcggc agacaaaaag a                                         21
SEQ ID NO: 15     moltype = DNA  length = 22
FEATURE          Location/Qualifiers
misc_feature     1..22
note = Reverse 5'-3' Primer
source          1..22
                mol_type = other DNA
                organism = Homo sapiens
SEQUENCE: 15      gaagtattgc ttcaagtggc ct                                         22
SEQ ID NO: 16     moltype = DNA  length = 21
FEATURE          Location/Qualifiers
misc_feature     1..21
note = Forward 5' Primer
source          1..21
                mol_type = other DNA
                organism = Homo sapiens
SEQUENCE: 16

```

-continued

agaagaacgg caagtacgag a	21
SEQ ID NO: 17 moltype = DNA length = 21 FEATURE Location/Qualifiers misc_feature 1..21 note = Reverse 5'-3' Primer source 1..21 mol_type = other DNA organism = Homo sapiens	
SEQUENCE: 17 tgttgagggc cagattgtgg c	21
SEQ ID NO: 18 moltype = DNA length = 21 FEATURE Location/Qualifiers misc_feature 1..21 note = Forward 5' Primer source 1..21 mol_type = other DNA organism = Homo sapiens	
SEQUENCE: 18 taacggcgcc agacaaaaag a	21
SEQ ID NO: 19 moltype = DNA length = 22 FEATURE Location/Qualifiers misc_feature 1..22 note = Reverse 5'-3' Primer source 1..22 mol_type = other DNA organism = Homo sapiens	
SEQUENCE: 19 gaagtattgc ttcaagttggc ct	22
SEQ ID NO: 20 moltype = length = SEQUENCE: 20 000	
SEQ ID NO: 21 moltype = length = SEQUENCE: 21 000	
SEQ ID NO: 22 moltype = DNA length = 20 FEATURE Location/Qualifiers misc_feature 1..20 note = Forward 5' Primer source 1..20 mol_type = other DNA organism = Homo sapiens	
SEQUENCE: 22 acgaatctcc gaccaccact	20
SEQ ID NO: 23 moltype = DNA length = 20 FEATURE Location/Qualifiers misc_feature 1..20 note = Reverse 5'-3' Primer source 1..20 mol_type = other DNA organism = Homo sapiens	
SEQUENCE: 23 ccatggccac aacaactgac	20
SEQ ID NO: 24 moltype = DNA length = 22 FEATURE Location/Qualifiers misc_feature 1..22 note = Forward 5' Primer source 1..22 mol_type = other DNA organism = Homo sapiens	
SEQUENCE: 24 gaagtgtccc aggacatgat aa	22
SEQ ID NO: 25 moltype = DNA length = 22 FEATURE Location/Qualifiers misc_feature 1..22 note = Reverse 5'-3' Primer source 1..22 mol_type = other DNA	

-continued

```

SEQUENCE: 25          organism = Homo sapiens
ctcttgagta gctgggattg ag                                22

SEQ ID NO: 26          moltype = DNA  length = 46
FEATURE
misc_feature           Location/Qualifiers
1..46
note = Forward 5'-3' Primer
source
1..46
mol_type = other DNA
organism = Homo sapiens

SEQUENCE: 26
ctccgtgctg atcaccctca tcatgatcg cgtcatcggt tatgac      46

SEQ ID NO: 27          moltype = DNA  length = 48
FEATURE
misc_feature           Location/Qualifiers
1..48
note = Reverse 3'-5' Primer
source
1..48
mol_type = other DNA
organism = Homo sapiens

SEQUENCE: 27
gaggcacgac tagtgggagt agtacttagtc gcagtagcac caatactg      48

SEQ ID NO: 28          moltype = DNA  length = 20
FEATURE
misc_feature           Location/Qualifiers
1..20
note = Forward 5'-3' Primer
source
1..20
mol_type = other DNA
organism = Homo sapiens

SEQUENCE: 28
catcatgatc agcgtcatcg                                20

SEQ ID NO: 29          moltype = DNA  length = 25
FEATURE
misc_feature           Location/Qualifiers
1..25
note = gLN2411 guide RNA F
source
1..25
mol_type = other DNA
organism = Homo sapiens

SEQUENCE: 29
caccgcatca tgatcagcgt catcg                                25

SEQ ID NO: 30          moltype = DNA  length = 25
FEATURE
misc_feature           Location/Qualifiers
1..25
note = gLN2411 guide RNA R
source
1..25
mol_type = other DNA
organism = Homo sapiens

SEQUENCE: 30
aaaccgatga cgctgtatcat gatgc                                25

SEQ ID NO: 31          moltype = DNA  length = 118
FEATURE
misc_feature           Location/Qualifiers
1..118
note = Donor ssODN#A N141I
source
1..118
mol_type = other DNA
organism = Homo sapiens

SEQUENCE: 31
gagagaagcg tggctggagg gcagggccag ggcttcacct tgttagcagcg gtacttgtag  60
agcaccacca agaagatggc cataaccacg atgacgtcta tcatgttag ggtgttca   118

```

What is claimed is:

1. A method of generating basal forebrain cholinergic neurons (BFCNs) comprising: culturing pluripotent stem cells (PSCs) in a basal media comprising an inhibitor of transforming growth factor beta (TGF- β) signaling and an activator of sonic hedgehog (Shh) signaling to induce neuroectodermal differentiation, wherein the basal media lacks

basic fibroblast growth factor (bFGF), TGF- β , lithium chloride (Li—Cl), GABA and pipecolic acid, thereby generating BFCNs.

2. A method of treating a disease or disorder in a subject comprising administering a BFCN generated using the method of claim 1 to the subject, thereby treating the disease or disorder.

3. The method of claim **2**, wherein the disease or disorder is an amyloidogenic disease.
4. The method of claim **2**, wherein the disease or disorder is associated with diminished neuronal excitability in BFCNs of the subject.
5. The method of claim **3**, wherein the disease or disorder is selected from the group consisting of systemic amyloidosis, Alzheimer's disease, mature onset diabetes, Parkinson's disease, Huntington's disease, fronto-temporal dementia, and prion-related transmissible spongiform encephalopathies.
6. The method of claim **2**, wherein the BFCN comprises a genome repaired PSEN2 mutation.
7. The method of claim **6**, wherein the PSEN2 mutation is PSEN2^{N141I}.
8. A method of restoring neuronal excitability in basal forebrain cholinergic neurons (BFCNs) in a subject, comprising:
 - a) isolating a BFCN from the subject, wherein the BFCN has a mutation in presenilin 2 (PSEN2) resulting in impaired neuronal excitability of the BFCN;
 - b) generating an induced pluripotent stem cell (iPSC) using the BFCN of (a);
 - c) repairing the PSEN2 mutation in the iPSC;
 - d) culturing the iPSC of (c) using the method of claim **1** to generate a BFCN having the repaired mutation; and
 - e) administering the iPSC of (d) to the subject, thereby restoring neuronal excitability in BFCNs in the subject.
9. The method of claim **8**, wherein the mutation is PSEN2^{N141I}.
10. The method of claim **8**, wherein the subject has, or is at risk of having Alzheimer's disease.
11. A method of identifying a compound for treatment or prevention of a disease or disorder associated with diminished neuronal excitability in basal forebrain cholinergic neurons (BFCNs), the method comprising:
 - a) contacting a BFCN or neuronal embryoid body (NEB) generated by the method of claim **1** with a candidate compound, wherein the BFCN comprises a mutation in presenilin 2 (PSEN2) resulting in impaired neuronal excitability of the BFCN; and
 - b) detecting neuronal excitability of the BFCN after contact with the candidate compound.
12. The method of claim **11**, wherein the mutation is PSEN2^{N141I}.
13. The method of claim **11**, wherein the disease or disorder is Alzheimer's disease.
14. A basal forebrain cholinergic neuron (BFCN) generated using the method of claim **1**.
15. The BFCN of claim **14**, wherein the genome has a recombinantly introduced marker.
16. A kit for generating a basal forebrain cholinergic neuron (BFCN) comprising a culture media having an inhibitor of transforming growth factor beta (TGF-β) signaling and an activator of sonic hedgehog (Shh) signaling, wherein the basal media lacks basic fibroblast growth factor (bFGF), TGF-β, lithium chloride (Li—Cl), GABA and pipecolic acid, wherein the inhibitor comprises SB431542 and LDN193189, and wherein the activator comprises Smoothened Agonist (SAG) and purmorphamine.
17. The kit of claim **16**, wherein the kit further comprises reagents for generating an induced pluripotent stem cell (iPSC).
18. The kit of claim **16**, wherein the kit further comprises a gene-editing reagent.
19. The kit of claim **16**, wherein the kit further comprises a reagent for detecting a genetic mutation.
20. The kit of claim **19**, wherein the mutation is PSEN2^{N141I}.

* * * * *