



US012391980B2

(12) **United States Patent**  
**Chee**

(10) **Patent No.:** US 12,391,980 B2  
(45) **Date of Patent:** \*Aug. 19, 2025

(54) **SPATIALLY ENCODED BIOLOGICAL ASSAYS**(71) Applicant: **Prognosys Biosciences, Inc.**, San Diego, CA (US)(72) Inventor: **Mark S. Chee**, San Diego, CA (US)(73) Assignee: **Prognosys Biosciences, Inc.**, San Diego, CA (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **19/074,001**(22) Filed: **Mar. 7, 2025**(65) **Prior Publication Data**

US 2025/0207183 A1 Jun. 26, 2025

**Related U.S. Application Data**

(63) Continuation of application No. 18/972,148, filed on Dec. 6, 2024, now Pat. No. 12,297,488, which is a continuation of application No. 18/793,359, filed on Aug. 2, 2024, now Pat. No. 12,234,505, which is a continuation of application No. 18/100,127, filed on Jan. 23, 2023, which is a continuation of application No. 17/878,519, filed on Aug. 1, 2022, now Pat. No. 11,560,587, which is a continuation of application No. 17/556,588, filed on Dec. 20, 2021, now Pat. No. 11,401,545, which is a continuation of application No. 17/223,669, filed on Apr. 6, 2021, now Pat. No. 11,208,684, which is a continuation of application No. 17/030,230, filed on Sep. 23, 2020, now Pat. No. 11,384,386, which is a continuation of application No. 16/988,284, filed on Aug. 7, 2020, now Pat. No. 10,961,566, which is a continuation of application No. 16/414,213, filed on May 16, 2019, now Pat. No. (Continued)

(51) **Int. Cl.**

<i>CI2Q 1/6837</i>	(2018.01)
<i>B01L 3/00</i>	(2006.01)
<i>CI2Q 1/68</i>	(2018.01)
<i>CI2Q 1/6804</i>	(2018.01)
<i>CI2Q 1/6809</i>	(2018.01)
<i>CI2Q 1/6834</i>	(2018.01)
<i>CI2Q 1/6841</i>	(2018.01)
<i>CI2Q 1/6869</i>	(2018.01)
<i>CI2Q 1/6874</i>	(2018.01)
<i>CI2Q 1/6876</i>	(2018.01)
<i>C40B 30/04</i>	(2006.01)
<i>C40B 60/04</i>	(2006.01)
<i>G01N 33/53</i>	(2006.01)
<i>G01N 33/543</i>	(2006.01)
<i>G01N 33/68</i>	(2006.01)
<i>CI2Q 1/6858</i>	(2018.01)

(52) **U.S. Cl.**

CPC ..... *CI2Q 1/6837* (2013.01); *B01L 3/502715* (2013.01); *CI2Q 1/68* (2013.01); *CI2Q 1/6804* (2013.01); *CI2Q 1/6809* (2013.01); *CI2Q 1/6834* (2013.01); *CI2Q 1/6841* (2013.01); *CI2Q 1/6869* (2013.01); *CI2Q 1/6874* (2013.01); *CI2Q 1/6876* (2013.01); *C40B 30/04* (2013.01); *C40B 60/04* (2013.01); *G01N 33/5308* (2013.01); *G01N 33/543* (2013.01); *G01N 33/54366* (2013.01); *G01N 33/6845* (2013.01); *G01N 33/6848* (2013.01); *CI2Q 1/6858* (2013.01); *CI2Q 2600/156* (2013.01); *CI2Q 2600/158* (2013.01); *G01N 2458/10* (2013.01); *G01N 2458/40* (2013.01)

(58) **Field of Classification Search**

CPC ..... *CI2Q 1/6837*; *CI2Q 1/68*; *CI2Q 1/6804*; *CI2Q 1/6809*; *CI2Q 1/6834*; *CI2Q 1/6841*; *CI2Q 1/6869*; *CI2Q 1/6874*; *CI2Q 1/6876*; *CI2Q 1/6858*; *CI2Q 2600/156*; *CI2Q 2600/158*; *B01L 3/502715*; *C40B 30/04*; *C40B 60/04*; *G01N 33/5308*; *G01N 33/543*; *G01N 33/54366*; *G01N 33/6845*; *G01N 33/6848*; *G01N 2458/10*; *G01N 2458/40*

See application file for complete search history.

## (56)

**References Cited**

## U.S. PATENT DOCUMENTS

4,514,388 A	4/1985	Psaledakis
4,557,903 A	12/1985	McCormick

(Continued)

## FOREIGN PATENT DOCUMENTS

AU	2003200718	10/2006
CA	2169928	4/2007

(Continued)

## OTHER PUBLICATIONS

U.S. Appl. No. 16/353,937, filed Mar. 14, 2019, Frisen et al.  
(Continued)

Primary Examiner — Narayan K Bhat

(74) Attorney, Agent, or Firm — Fish &amp; Richardson P.C.

## (57)

**ABSTRACT**

The present invention provides assays and assay systems for use in spatially encoded biological assays. The invention provides an assay system comprising an assay capable of high levels of multiplexing where reagents are provided to a biological sample in defined spatial patterns; instrumentation capable of controlled delivery of reagents according to the spatial patterns; and a decoding scheme providing a readout that is digital in nature.

**Related U.S. Application Data**

- 10,787,701, which is a continuation of application No. 16/402,098, filed on May 2, 2019, now Pat. No. 10,472,669, which is a continuation of application No. 16/276,235, filed on Feb. 14, 2019, now Pat. No. 10,480,022, which is a continuation of application No. 15/187,661, filed on Jun. 20, 2016, now Pat. No. 10,308,982, which is a continuation of application No. 13/080,616, filed on Apr. 5, 2011, now Pat. No. 9,371,598.
- (60) Provisional application No. 61/321,124, filed on Apr. 5, 2010.

**(56) References Cited****U.S. PATENT DOCUMENTS**

4,574,729 A	3/1986	Wells	5,962,271 A	10/1999	Chenchik et al.
4,683,195 A	7/1987	Mullis	5,962,272 A	10/1999	Chenchik et al.
4,683,202 A	7/1987	Mullis	5,965,443 A	10/1999	Reznikoff et al.
4,731,335 A	3/1988	Brigati	5,994,136 A	11/1999	Naldini et al.
4,800,159 A	1/1989	Mullis	6,013,440 A	1/2000	Lipshutz
4,829,012 A	5/1989	Cambiaso et al.	6,013,516 A	1/2000	Verma et al.
4,883,867 A	11/1989	Lee	6,027,889 A	2/2000	Barany et al.
4,965,188 A	10/1990	Mullis	6,054,274 A	4/2000	Sampson et al.
4,968,601 A	11/1990	Jacobson et al.	6,060,240 A	5/2000	Kamb et al.
4,988,617 A	1/1991	Landegren et al.	6,083,761 A	7/2000	Kedar et al.
5,002,882 A	3/1991	Lunnen	6,130,073 A	10/2000	Eggerding
5,061,049 A	10/1991	Hornbeck	6,133,436 A	10/2000	Koester et al.
5,130,238 A	7/1992	Malek	6,136,592 A	10/2000	Leighton
5,183,053 A	2/1993	Yeh et al.	6,143,496 A	11/2000	Brown
5,308,751 A	5/1994	Ohkawa	6,153,389 A	11/2000	Haarer
5,321,130 A	6/1994	Yue	6,153,408 A	11/2000	Abastado et al.
5,410,030 A	4/1995	Yue	6,157,432 A	12/2000	Helbing
5,436,129 A	7/1995	Stapleton	6,159,736 A	12/2000	Reznikoff et al.
5,436,134 A	7/1995	Haugland	6,165,714 A	12/2000	Lane et al.
5,455,166 A	10/1995	Walker	6,172,218 B1	1/2001	Brenner
5,472,881 A	12/1995	Beebe et al.	6,210,891 B1	4/2001	Nyren
5,487,972 A	1/1996	Gelfand et al.	6,210,894 B1	4/2001	Brennan
5,494,810 A	2/1996	Barany et al.	6,214,587 B1	4/2001	Dattagupta
5,496,518 A	3/1996	Arai et al.	6,221,591 B1	4/2001	Aerts
5,503,980 A	4/1996	Cantor	6,221,654 B1	4/2001	Quake
5,512,439 A	4/1996	Hornes	6,251,639 B1	6/2001	Kurn
5,512,462 A	4/1996	Cheng	6,258,558 B1	7/2001	Szostak
5,512,478 A	4/1996	Orser et al.	6,258,568 B1	7/2001	Nyren
5,559,032 A	9/1996	Porneroy	6,261,804 B1	7/2001	Szostak
5,582,977 A	12/1996	Yue	6,265,552 B1	7/2001	Schatz
5,589,173 A	12/1996	O'Brien	6,266,459 B1	7/2001	Walt
5,599,675 A	2/1997	Brenner	6,268,148 B1	7/2001	Barany et al.
5,610,287 A	3/1997	Nikiforov et al.	6,274,320 B1	8/2001	Rothberg
5,641,658 A	6/1997	Adams	6,281,804 B1	8/2001	Haller
5,648,245 A	7/1997	Fire et al.	6,291,180 B1	9/2001	Chu
5,658,751 A	8/1997	Yue	6,291,187 B1	9/2001	Kingsmore et al.
5,667,976 A	9/1997	Van Ness et al.	6,300,063 B1	10/2001	Lipshutz et al.
5,672,344 A	9/1997	Kelley et al.	6,300,093 B1	10/2001	Kindsvogel et al.
5,695,940 A	12/1997	Drmanac et al.	6,306,597 B1	10/2001	Macevicz
5,716,825 A	2/1998	Hancock et al.	6,309,824 B1	10/2001	Drmanac
5,750,341 A	5/1998	Macevicz	6,323,009 B1	11/2001	Lasken et al.
5,752,982 A	5/1998	Lang et al.	6,337,472 B1	1/2002	Garner et al.
5,763,175 A	6/1998	Brenner	6,344,316 B1	2/2002	Lockhart
5,807,522 A	9/1998	Brown et al.	6,344,329 B1	2/2002	Lizardi et al.
5,817,783 A	10/1998	Calabretta et al.	6,348,990 B1	2/2002	Igasaki et al.
5,830,711 A	11/1998	Barany et al.	6,355,431 B1	3/2002	Choe
5,837,832 A	11/1998	Chee et al.	6,358,508 B1	3/2002	Ni et al.
5,837,860 A	11/1998	Anderson et al.	6,368,801 B1	4/2002	Faruqi
5,854,033 A	12/1998	Lizardi	6,391,937 B1	5/2002	Beuhler et al.
5,863,753 A	1/1999	Haugland	6,401,267 B1	6/2002	Drmanac
5,866,377 A	2/1999	Kim et al.	6,404,907 B1	6/2002	Gilchrist
5,871,921 A	2/1999	Landegren et al.	6,416,950 B1	7/2002	Lohse
5,875,258 A	2/1999	Ortyn et al.	6,426,215 B1	7/2002	Sandell
5,912,148 A	6/1999	Eggerding	6,432,360 B1	8/2002	Church et al.
5,919,626 A	7/1999	Shi et al.	6,444,661 B1	9/2002	Barton et al.
5,925,545 A	7/1999	Reznikoff et al.	6,485,982 B1	11/2002	Charlton
5,928,906 A	7/1999	Koester et al.	6,503,713 B1	1/2003	Rana
5,958,775 A	9/1999	Wickstrom et al.	6,506,561 B1	1/2003	Cheval et al.
			6,518,018 B1	2/2003	Szostak
			6,534,266 B1	3/2003	Singer
			6,544,732 B1	4/2003	Choe
			6,544,790 B1	4/2003	Sabatini
			6,565,727 B1	5/2003	Shenderov
			6,573,043 B1	6/2003	Cohen et al.
			6,579,695 B1	6/2003	Lambalot
			6,620,584 B1 *	9/2003	Chee ..... C12Q 1/6837 435/7.1
			6,632,641 B1	10/2003	Brennan
			6,673,620 B1	1/2004	Loeffler
			6,677,160 B1	1/2004	Stockman et al.
			6,699,710 B1	3/2004	Kononen
			6,737,236 B1	5/2004	Pieken et al.
			6,770,441 B2	8/2004	Dickinson
			6,773,566 B2	8/2004	Shenderov
			6,773,886 B2	8/2004	Kaufman
			6,787,308 B2	9/2004	Balasubramanian
			6,797,470 B2	9/2004	Barany et al.
			6,800,453 B2	10/2004	Labaer
			6,812,005 B2	11/2004	Fan et al.

# US 12,391,980 B2

Page 3

---

(56)

## References Cited

### U.S. PATENT DOCUMENTS

6,828,100 B1	12/2004	Ronaghi	7,635,566 B2	12/2009	Brenner
6,833,246 B2	12/2004	Balasubramanian	7,641,779 B2	1/2010	Becker
6,852,487 B1	2/2005	Barany et al.	7,655,898 B2	2/2010	Miller
6,859,570 B2	2/2005	Walt	7,666,612 B2	2/2010	Johnsson
6,864,052 B1	3/2005	Drmanac	7,674,589 B2	3/2010	Cohen et al.
6,867,028 B2	3/2005	Janulaitis	7,674,752 B2	3/2010	He
6,872,816 B1	3/2005	Hall et al.	7,700,286 B2	4/2010	Stroun et al.
6,875,572 B2	4/2005	Prudent et al.	7,709,198 B2	5/2010	Luo et al.
6,878,515 B1	4/2005	Landgren	7,741,106 B2	6/2010	Moyle et al.
6,890,741 B2	5/2005	Fan et al.	7,754,429 B2	7/2010	Rigatti
6,897,023 B2	5/2005	Fu	7,776,547 B2	8/2010	Roth
6,911,132 B2	6/2005	Pamula	7,776,567 B2	8/2010	Mao
6,911,345 B2	6/2005	Quake	7,785,869 B2	8/2010	Belgrader et al.
6,913,881 B1	7/2005	Aizenstein et al.	7,803,943 B2	9/2010	Mao
6,913,921 B2	7/2005	Fischer	7,844,940 B2	11/2010	Shin et al.
6,942,968 B1	9/2005	Dickinson et al.	7,848,553 B2	12/2010	Hertel et al.
6,969,488 B2	11/2005	Bridgeman	7,858,321 B2	12/2010	Glezer
6,969,589 B2	11/2005	Patil	7,888,009 B2	2/2011	Barany et al.
6,977,033 B2	12/2005	Becker	7,892,747 B2	2/2011	Barany et al.
7,001,792 B2	2/2006	Sauer et al.	7,910,304 B2	3/2011	Drmanac
7,011,944 B2	3/2006	Prudent et al.	7,914,981 B2	3/2011	Barany et al.
7,052,244 B2	5/2006	Fouillet	7,941,279 B2	5/2011	Hwang et al.
7,057,026 B2	6/2006	Barnes	7,955,794 B2	6/2011	Shen et al.
7,060,431 B2	6/2006	Chee et al.	7,960,119 B2	6/2011	Chee
7,083,980 B2	8/2006	Reznikoff et al.	7,960,120 B2	6/2011	Rigatti
7,098,041 B2	8/2006	Kaylor et al.	7,985,565 B2	7/2011	Mayer et al.
7,115,400 B1	10/2006	Adessi	8,003,354 B2	8/2011	Shen et al.
7,118,883 B2	10/2006	Inoue	8,030,466 B2	10/2011	Shin et al.
7,128,893 B2	10/2006	Leamon et al.	8,030,477 B2	10/2011	Cerrina et al.
7,163,612 B2	1/2007	Sterling	8,046,043 B2	10/2011	Asano et al.
7,166,431 B2	1/2007	Chee et al.	8,076,063 B2	12/2011	Fan
7,192,735 B2	3/2007	Lambalot	8,092,784 B2	1/2012	Mao
7,211,414 B2	5/2007	Hardin	8,124,751 B2	2/2012	Pierce et al.
7,223,371 B2	5/2007	Hayenga et al.	8,131,476 B2	3/2012	Cline et al.
7,229,769 B2	6/2007	Kozlov	8,148,068 B2	4/2012	Brenner
7,244,559 B2	7/2007	Rothberg	8,148,518 B2	4/2012	Buchanan
7,255,994 B2	8/2007	Lao	8,198,028 B2	6/2012	Rigatti et al.
7,258,976 B2	8/2007	Mitsuhashi	8,199,999 B2	6/2012	Hoyt et al.
7,259,258 B2	8/2007	Kozlov et al.	8,206,917 B2	6/2012	Chee
7,264,929 B2	9/2007	Rothberg	8,207,093 B2	6/2012	Szostak
7,270,950 B2	9/2007	Szostak	8,268,554 B2	9/2012	Schallmeiner
7,282,328 B2	10/2007	Kong et al.	8,278,034 B2	10/2012	Muraca
7,297,518 B2	11/2007	Quake	8,288,103 B2	10/2012	Olyphant
7,315,019 B2	1/2008	Turner	8,288,122 B2	10/2012	O'Leary et al.
7,328,979 B2	2/2008	Decre	8,330,087 B2	12/2012	Domenicali
7,329,492 B2	2/2008	Hardin	8,337,851 B2	12/2012	Aukerman
7,358,047 B2	4/2008	Hafner et al.	8,343,500 B2	1/2013	Wraith
7,361,488 B2	4/2008	Fan et al.	8,383,338 B2	2/2013	Kitzman
7,375,234 B2	5/2008	Sharpless et al.	8,415,102 B2	4/2013	Geiss et al.
7,378,242 B2	5/2008	Hurt	8,431,691 B2	4/2013	McKernan et al.
7,393,665 B2	7/2008	Brenner	8,460,865 B2	6/2013	Cheee
7,405,281 B2	7/2008	Xu	8,462,981 B2	6/2013	Determan et al.
7,407,757 B2	8/2008	Brenner	8,481,257 B2	7/2013	Van Eijk
7,427,678 B2	9/2008	Pieken et al.	8,481,258 B2	7/2013	Church et al.
7,456,012 B2	11/2008	Ryttsen et al.	8,481,292 B2	7/2013	Casbor
7,462,449 B2	12/2008	Quake	8,481,698 B2	7/2013	Lieberman et al.
7,473,767 B2	1/2009	Dimitrov	8,486,625 B2	7/2013	Gunderson
7,499,806 B2	3/2009	Kermani et al.	8,507,204 B2	8/2013	Pierce et al.
7,500,637 B2	3/2009	Marimon et al.	8,519,115 B2	8/2013	Webster et al.
7,501,245 B2	3/2009	Quake	8,551,710 B2	10/2013	Bernitz et al.
7,510,637 B2	3/2009	Barlow et al.	8,568,979 B2	10/2013	Stuelpnagel et al.
7,534,991 B2	5/2009	Miller et al.	RE44,596 E	11/2013	Stroun et al.
7,537,897 B2	5/2009	Brenner	8,586,310 B2	11/2013	Mitra
7,544,473 B2	6/2009	Brennan	8,597,891 B2	12/2013	Barany et al.
7,547,380 B2	6/2009	Velev	8,603,743 B2	12/2013	Liu et al.
7,555,155 B2	6/2009	Levenson et al.	8,604,182 B2	12/2013	Luo et al.
7,561,336 B2	7/2009	Osaka et al.	8,614,073 B2	12/2013	Van Eijk
7,563,576 B2	7/2009	Chee	8,624,016 B2	1/2014	Barany et al.
7,579,153 B2	8/2009	Brenner	8,637,242 B2	1/2014	Shen
7,582,420 B2	9/2009	Oliphant et al.	8,658,361 B2	2/2014	Wu et al.
7,595,883 B1	9/2009	El Gamal	8,685,889 B2	4/2014	Van Eijk
7,601,492 B2	10/2009	Fu et al.	8,741,564 B2	6/2014	Seligmann
7,601,498 B2	10/2009	Mao	8,741,606 B2	6/2014	Casbon
7,608,434 B2	10/2009	Reznikoff et al.	8,748,103 B2	6/2014	Faham et al.
7,611,869 B2	11/2009	Fan	8,771,950 B2	7/2014	Church et al.
			8,778,849 B2	7/2014	Bowen
			8,785,353 B2	7/2014	Van Eijk
			8,790,873 B2	7/2014	Namsaraev et al.
			8,809,238 B2	8/2014	Livak et al.

(56)	References Cited					
U.S. PATENT DOCUMENTS						
8,815,512 B2	8/2014	Van Eijk	10,266,876 B2	4/2019	Cai et al.	
8,835,358 B2	9/2014	Fodor	10,267,808 B2	4/2019	Cai	
8,865,410 B2	10/2014	Shendure	10,357,771 B2	7/2019	Bharadwaj	
8,906,626 B2	12/2014	Oliphant et al.	10,391,467 B2	8/2019	Zhou et al.	
8,911,945 B2	12/2014	Van Eijk	10,428,326 B2	10/2019	Belhocine et al.	
8,936,912 B2	1/2015	Mitra	10,465,235 B2	11/2019	Gullberg et al.	
8,951,726 B2	2/2015	Luo et al.	10,501,791 B2	12/2019	Church et al.	
8,951,728 B2	2/2015	Rasmussen	10,633,648 B2	4/2020	Seelig et al.	
8,951,781 B2	2/2015	Reed	10,669,569 B2	6/2020	Gullberg et al.	
8,986,926 B2	3/2015	Ferree et al.	10,685,210 B2	6/2020	Wimberger-Friedl et al.	
9,005,891 B2	4/2015	Sinicropi et al.	10,697,013 B1	6/2020	Brenner et al.	
9,005,935 B2	4/2015	Belyaev	10,725,027 B2	7/2020	Bell	
9,023,768 B2	5/2015	Van Eijk	10,767,223 B1	9/2020	Brenner et al.	
9,062,348 B1	6/2015	Van Eijk	10,774,372 B2	9/2020	Chee et al.	
9,080,210 B2	7/2015	Van Eijk	10,774,374 B2	9/2020	Frisen et al.	
9,085,798 B2	7/2015	Chee	10,829,803 B2	11/2020	Terbrueggen et al.	
9,121,069 B2	9/2015	Lo	10,858,702 B2	12/2020	Lucero et al.	
9,163,283 B2	10/2015	Chee et al.	10,927,403 B2	2/2021	Chee et al.	
9,194,001 B2	11/2015	Brenner	10,995,362 B2	5/2021	Dallett et al.	
9,201,063 B2	12/2015	Sood et al.	11,046,996 B1	6/2021	Chee et al.	
9,217,176 B2	12/2015	Faham et al.	11,162,132 B2	11/2021	Frisen et al.	
9,290,808 B2	3/2016	Fodor	11,214,796 B2	1/2022	Shirai et al.	
9,290,809 B2	3/2016	Fodor	11,286,515 B2	3/2022	Chee et al.	
9,309,556 B2	4/2016	Myllykangas et al.	11,299,774 B2	4/2022	Frisen et al.	
9,328,383 B2	5/2016	Van Eijk	11,332,790 B2	5/2022	Chell et al.	
9,334,536 B2	5/2016	Van Eijk	11,352,659 B2	6/2022	Frisen et al.	
9,340,830 B2	5/2016	Lipson	11,359,228 B2	6/2022	Chee et al.	
9,371,563 B2	6/2016	Geiss et al.	11,390,912 B2	7/2022	Frisen et al.	
9,371,598 B2	6/2016	Chee	11,407,992 B2	8/2022	Dadhwal	
9,376,716 B2	6/2016	Van Eijk	11,408,029 B2	8/2022	Katiraei et al.	
9,376,717 B2	6/2016	Gao et al.	11,434,524 B2	9/2022	Ramachandran Iyer et al.	
9,376,719 B2	6/2016	Van Eijk	11,459,607 B1	10/2022	Terry et al.	
9,404,156 B2	8/2016	Hicks	11,479,809 B2	10/2022	Frisen et al.	
9,416,409 B2	8/2016	Hayden	11,492,612 B1	11/2022	Dadhwal	
9,447,459 B2	9/2016	Van Eijk	11,501,440 B2	11/2022	Weisenfeld et al.	
9,453,256 B2	9/2016	Van Eijk	11,505,828 B2	11/2022	Chell et al.	
9,493,820 B2	11/2016	Van Eijk	11,512,308 B2	11/2022	Gallant et al.	
9,494,588 B2	11/2016	Springer	11,519,033 B2	12/2022	Schnall-Levin et al.	
9,506,061 B2	11/2016	Brown	11,535,887 B2	12/2022	Gallant et al.	
9,512,487 B2	12/2016	Faham et al.	11,560,592 B2	1/2023	Chew et al.	
9,541,504 B2	1/2017	Hoyt	11,560,593 B2	1/2023	Chell et al.	
9,557,330 B2	1/2017	Siciliano et al.	11,592,447 B2	2/2023	Uytingco et al.	
9,574,230 B2	2/2017	Van Eijk	11,624,063 B2	4/2023	Dadhwal	
9,593,365 B2	3/2017	Frisen et al.	11,624,086 B2	4/2023	Uytingco et al.	
9,598,728 B2	3/2017	Barany et al.	11,649,485 B2	5/2023	Yin et al.	
9,624,538 B2	4/2017	Church et al.	11,661,626 B2	5/2023	Katiraei et al.	
9,657,335 B2	5/2017	Van Eijk	11,680,260 B2	6/2023	Kim et al.	
9,670,542 B2	6/2017	Van Eijk	11,692,218 B2	7/2023	Engblom et al.	
9,671,344 B2	6/2017	Staker	11,702,693 B2	7/2023	Bharadwaj	
9,694,361 B2	7/2017	Bharadwaj	11,702,698 B2	7/2023	Stoeckius	
9,702,004 B2	7/2017	Van Eijk	11,732,299 B2	8/2023	Ramachandran Iyer	
9,714,446 B2	7/2017	Webster et al.	11,732,300 B2	8/2023	Bava	
9,714,937 B2	7/2017	Dunaway	11,739,372 B2	8/2023	Frisen et al.	
9,745,627 B2	8/2017	Van Eijk	11,739,381 B2	8/2023	Chew et al.	
9,777,324 B2	10/2017	Van Eijk	11,753,673 B2	9/2023	Chew et al.	
9,816,134 B2	11/2017	Namsaraev	11,753,674 B2	9/2023	Chee et al.	
9,850,536 B2	12/2017	Oliphant et al.	11,753,675 B2	9/2023	Ramachandran Iyer	
9,868,979 B2	1/2018	Chee et al.	11,761,038 B1	9/2023	Stoeckius	
9,879,313 B2	1/2018	Chee et al.	11,768,175 B1	9/2023	Kim et al.	
9,889,422 B2	2/2018	Smith et al.	11,773,433 B2	10/2023	Gallant et al.	
9,896,721 B2	2/2018	Van Eijk	11,781,130 B2	10/2023	Dadhwal	
9,898,576 B2	2/2018	Van Eijk	11,788,122 B2	10/2023	Frisen et al.	
9,902,950 B2	2/2018	Church et al.	11,795,498 B2	10/2023	Frisen et al.	
9,902,991 B2	2/2018	Sinicropi et al.	11,795,507 B2	10/2023	Chell et al.	
9,958,454 B2	5/2018	Kozlov et al.	11,808,769 B2	11/2023	Uytingco et al.	
10,023,907 B2	7/2018	Van Eijk	11,821,024 B2	11/2023	Chee et al.	
10,030,261 B2	7/2018	Frisen et al.	11,821,035 B1	11/2023	Bent et al.	
10,072,104 B2	9/2018	Winnik et al.	11,827,935 B1	11/2023	Ramachandran Iyer et al.	
10,077,478 B2	9/2018	Faham et al.	11,835,462 B2	12/2023	Bava	
10,095,832 B2	10/2018	Van Eijk	11,840,687 B2	12/2023	Gallant et al.	
10,144,966 B2	12/2018	Cantor	11,840,724 B2	12/2023	Chew et al.	
10,196,691 B2	2/2019	Harkin et al.	11,845,979 B2	12/2023	Engblom et al.	
10,227,639 B2	3/2019	Levner et al.	11,859,178 B2	1/2024	Gallant et al.	
10,246,752 B2	4/2019	Faham et al.	11,866,767 B2	1/2024	Uytingco et al.	

# US 12,391,980 B2

Page 5

---

(56)

## References Cited

### U.S. PATENT DOCUMENTS

11,873,482 B2	1/2024	Kim et al.	2003/0108726 A1	6/2003	Schembri et al.
11,891,654 B2	2/2024	Alvarado Martinez et al.	2003/0113713 A1	6/2003	Glezer
11,898,205 B2	2/2024	Bava	2003/0152927 A1	7/2003	Lizardi
11,926,822 B1	3/2024	Gohil et al.	2003/0153850 A1	8/2003	Davis et al.
11,926,863 B1	3/2024	Boutet	2003/0162216 A1	8/2003	Gold
11,926,867 B2	3/2024	Yin et al.	2003/0165948 A1	9/2003	Alsmadi et al.
11,933,957 B1	3/2024	Tentori et al.	2003/0170637 A1	9/2003	Pirrung et al.
11,952,627 B2	4/2024	Stoeckius	2003/0175947 A1	9/2003	Liu et al.
11,959,076 B2	4/2024	Kim et al.	2003/0190744 A1	10/2003	McGarry et al.
11,959,130 B2	4/2024	Galonska et al.	2003/0205632 A1	11/2003	Kim et al.
11,965,213 B2	4/2024	Williams	2003/0211489 A1	11/2003	Shen et al.
11,970,739 B2	4/2024	Chew et al.	2003/0215936 A1	11/2003	Kallionиеми et al.
11,981,958 B1	5/2024	Galonska	2003/0224419 A1	12/2003	Corcoran
11,981,960 B1	5/2024	Lin et al.	2003/0232348 A1	12/2003	Jones et al.
11,981,965 B2	5/2024	Chell et al.	2003/0232382 A1	12/2003	Brennan
RE50,065 E	7/2024	Frisen et al.	2003/0235535 A1	12/2003	Zhou
12,024,741 B2	7/2024	Tentori et al.	2003/0235852 A1	12/2003	Roberts
12,031,177 B1	7/2024	Tentori et al.	2003/0235854 A1	12/2003	Chan et al.
12,060,604 B2	8/2024	Katiraei et al.	2004/0002090 A1	1/2004	Mayer et al.
12,071,655 B2	8/2024	Sukovich et al.	2004/0019005 A1	1/2004	Van Ness
12,076,701 B2	9/2024	Bava	2004/0023320 A1	2/2004	Steiner et al.
12,098,417 B2	9/2024	Engblom et al.	2004/0033499 A1	2/2004	Ilsley et al.
12,098,985 B2	9/2024	Cox et al.	2004/0050699 A1	3/2004	Goncalves
12,110,541 B2	10/2024	Bava	2004/0067492 A1	4/2004	Peng et al.
12,117,439 B2	10/2024	Delaney et al.	2004/0067493 A1	4/2004	Matsuzaki
12,128,403 B2	10/2024	Kim et al.	2004/0082058 A1	4/2004	Schleifer et al.
12,129,516 B2	10/2024	Tentori et al.	2004/0082059 A1	4/2004	Webb
12,157,124 B2	12/2024	Cox et al.	2004/0096853 A1	5/2004	Mayer
12,180,543 B2	12/2024	Uytingco et al.	2004/0106110 A1	6/2004	Balasubramanian
12,195,790 B2	1/2025	Sukovich et al.	2004/0112442 A1	6/2004	Maerkl
12,203,134 B2	1/2025	Nagendran et al.	2004/0121456 A1	6/2004	Fischer
12,209,280 B1	1/2025	Mignardi et al.	2004/0175822 A1	9/2004	Timperman et al.
D1,064,308 S	2/2025	Alimsijah et al.	2004/0219588 A1	11/2004	Furuta
12,223,751 B2	2/2025	Li et al.	2004/0224326 A1	11/2004	Kim et al.
12,228,544 B2	2/2025	Kim et al.	2004/0235103 A1	11/2004	Reznikoff et al.
12,241,060 B2	3/2025	Kim et al.	2004/0241660 A1	12/2004	Wojtowicz et al.
12,241,890 B2	3/2025	Delaney et al.	2004/0248287 A1	12/2004	Hu et al.
12,249,085 B2	3/2025	Tentori et al.	2004/0248325 A1	12/2004	Bukusoglu et al.
12,265,079 B1	4/2025	Bent	2004/0259105 A1	12/2004	Fan et al.
12,270,077 B2	4/2025	Schnall-Levin et al.	2005/0003431 A1	1/2005	Wucherpfennig
12,275,988 B2	4/2025	Galonska et al.	2005/0014203 A1	1/2005	Darfler et al.
12,281,357 B1	4/2025	Tentori et al.	2005/0019776 A1	1/2005	Callow et al.
12,286,673 B2	4/2025	Bava	2005/0019842 A1	1/2005	Prober et al.
12,287,264 B2	4/2025	Cox et al.	2005/0026188 A1	2/2005	Van Kessel
12,297,486 B2	5/2025	Patterson et al.	2005/0037362 A1	2/2005	Remacle et al.
12,344,892 B2	7/2025	Schnall-Levin et al.	2005/0037393 A1	2/2005	Gunderson et al.
2001/0029017 A1	10/2001	Yasuda et al.	2005/0042695 A1	2/2005	Meares et al.
2001/0039029 A1	11/2001	Nemori et al.	2005/0048535 A1	3/2005	Santin
2001/0053519 A1	12/2001	Fodor et al.	2005/0048580 A1	3/2005	Labaer
2001/0055764 A1	12/2001	Empedocles et al.	2005/0064432 A1	3/2005	Huang et al.
2002/0006477 A1	1/2002	Shishido et al.	2005/0064460 A1	3/2005	Holliger et al.
2002/0022261 A1	2/2002	Anderson et al.	2005/0079518 A1	4/2005	Baker et al.
2002/0040275 A1	4/2002	Cravatt	2005/0079520 A1	4/2005	Wu
2002/0045169 A1	4/2002	Shoemaker et al.	2005/0095627 A1	5/2005	Kolman et al.
2002/0045272 A1	4/2002	McDevitt et al.	2005/0100900 A1	5/2005	Kawashima et al.
2002/0048766 A1	4/2002	Doyle et al.	2005/0106617 A1	5/2005	Besemer et al.
2002/0051986 A1	5/2002	Baez et al.	2005/0118602 A1	6/2005	Li et al.
2002/0055100 A1	5/2002	Kawashima	2005/0130173 A1	6/2005	Leamon et al.
2002/0058250 A1	5/2002	Firth	2005/0130188 A1	6/2005	Walt
2002/0064779 A1	5/2002	Landegren	2005/0136414 A1	6/2005	Gunderson et al.
2002/0086441 A1	7/2002	Baranov et al.	2005/0147537 A1	7/2005	Sangha
2002/0132246 A1	9/2002	Kallionиеми et al.	2005/0164292 A1	7/2005	Farooqui
2002/0137031 A1	9/2002	Wolber	2005/0170373 A1	8/2005	Monforte
2002/0150909 A1	10/2002	Stuelpnagel	2005/0179746 A1	8/2005	Roux et al.
2002/0164611 A1	11/2002	Bamdad	2005/0191656 A1	9/2005	Drmanac et al.
2002/0168645 A1	11/2002	Taylor	2005/0191698 A1	9/2005	Chee et al.
2003/0017451 A1	1/2003	Wang et al.	2005/0196786 A1	9/2005	Levy
2003/0022207 A1	1/2003	Balasubramanian	2005/0202433 A1	9/2005	Van Beuningen
2003/0027342 A1	2/2003	Sheridan et al.	2005/0226780 A1	10/2005	Sandell et al.
2003/0040035 A1	2/2003	Slamon	2005/0227271 A1	10/2005	Kwon
2003/0064398 A1	4/2003	Barnes	2005/0239119 A1	10/2005	Tsukada et al.
2003/0073086 A1	4/2003	Guire et al.	2005/0239192 A1	10/2005	Nasarabadi et al.
2003/0087232 A1	5/2003	Christians	2005/0244850 A1	11/2005	Huang
2003/0092624 A1	5/2003	Wang et al.	2005/0255548 A1	11/2005	Lipovsek et al.
2003/0096323 A1	5/2003	James	2005/0257284 A1	11/2005	Nakajima et al.

(56)

**References Cited****U.S. PATENT DOCUMENTS**

2005/0260653 A1	11/2005	LaBaer	2008/0032310 A1	2/2008	Shannon et al.
2005/0266417 A1	12/2005	Barany et al.	2008/0033159 A1	2/2008	Kadushin et al.
2006/0003394 A1	1/2006	Song	2008/0038734 A1	2/2008	Sorge et al.
2006/0039823 A1	2/2006	Yamakawa et al.	2008/0043235 A1	2/2008	Oldham et al.
2006/0041384 A1	2/2006	Kermani et al.	2008/0047835 A1	2/2008	MacConnell
2006/0041385 A1	2/2006	Bauer et al.	2008/0071071 A1	3/2008	LaBaer et al.
2006/0046313 A1	3/2006	Roth	2008/0108082 A1	5/2008	Rank et al.
2006/0063160 A1	3/2006	West et al.	2008/0108804 A1	5/2008	Hayashizaki et al.
2006/0073506 A1	4/2006	Christians et al.	2008/0124252 A1	5/2008	Marchand et al.
2006/0079453 A1	4/2006	Sidney et al.	2008/0124810 A1	5/2008	Terbrueggen et al.
2006/0084078 A1	4/2006	Zhao	2008/0128627 A1	6/2008	Lundquist et al.
2006/0105352 A1	5/2006	Qiao et al.	2008/0132429 A1	6/2008	Perov et al.
2006/0110739 A1	5/2006	Heyduk	2008/0145378 A1	6/2008	Ovaa et al.
2006/0127946 A1	6/2006	Montagu et al.	2008/0145616 A1	6/2008	Gharib et al.
2006/0134669 A1	6/2006	Casasanta	2008/0153086 A1	6/2008	Wong
2006/0154286 A1	7/2006	Kong et al.	2008/0160580 A1	7/2008	Adessi et al.
2006/0164490 A1	7/2006	Kim et al.	2008/0199929 A1	8/2008	Yeung et al.
2006/0180489 A1	8/2006	Guiney et al.	2008/0218838 A1	9/2008	Rey-Mermet
2006/0183150 A1	8/2006	Cohen et al.	2008/0220434 A1	9/2008	Thomas
2006/0188875 A1	8/2006	Cox et al.	2008/0220981 A1	9/2008	McGregor
2006/0188901 A1	8/2006	Barnes et al.	2008/0261204 A1	10/2008	Lexow
2006/0188906 A1	8/2006	Kim et al.	2008/0280773 A1	11/2008	Fedurco et al.
2006/0194331 A1	8/2006	Pamula et al.	2008/0286795 A1	11/2008	Kawashima et al.
2006/0199183 A1	9/2006	Valat et al.	2008/0293046 A1	11/2008	Allawi et al.
2006/0199207 A1	9/2006	Matysiak	2008/0293591 A1	11/2008	Taussig et al.
2006/0211001 A1	9/2006	Yu et al.	2008/0312103 A1	12/2008	Nemoto et al.
2006/0216721 A1	9/2006	Kozlov et al.	2009/0005252 A1	1/2009	Drmanac et al.
2006/0216775 A1	9/2006	Burkart et al.	2009/0006002 A1	1/2009	Honisich et al.
2006/0228758 A1	10/2006	Muchhal et al.	2009/0011943 A1	1/2009	Drmanac et al.
2006/0240439 A1	10/2006	Smith et al.	2009/0018024 A1	1/2009	Church et al.
2006/0246475 A1	11/2006	Peterson et al.	2009/0023148 A1	1/2009	Moyle et al.
2006/0263789 A1	11/2006	Kincaid	2009/0026082 A1	1/2009	Rothberg et al.
2006/0275782 A1	12/2006	Gunderson et al.	2009/0036323 A1	2/2009	van Eijk et al.
2006/0275799 A1	12/2006	Banerjee et al.	2009/0048510 A1	2/2009	Miller et al.
2006/0281109 A1	12/2006	Barr Ost et al.	2009/0060866 A1	3/2009	Dousson et al.
2006/0292559 A1	12/2006	Reddy et al.	2009/0062148 A1	3/2009	Goldberg
2007/0003954 A1	1/2007	Kodadek et al.	2009/0068667 A1	3/2009	Meisner et al.
2007/0014810 A1	1/2007	Baker et al.	2009/0082212 A1	3/2009	Williams
2007/0020625 A1	1/2007	Duchaud et al.	2009/0099041 A1	4/2009	Church et al.
2007/0020640 A1	1/2007	McCloskey et al.	2009/0105959 A1	4/2009	Braverman et al.
2007/0020669 A1	1/2007	Ericsson	2009/0117573 A1	5/2009	Fu et al.
2007/0023292 A1	2/2007	Kim et al.	2009/0127589 A1	5/2009	Rothberg et al.
2007/0026430 A1	2/2007	Andersen et al.	2009/0152116 A1	6/2009	Boles et al.
2007/0036511 A1	2/2007	Lundquist et al.	2009/0155781 A1	6/2009	Drmanac et al.
2007/0048812 A1	3/2007	Moravec et al.	2009/0169089 A1	7/2009	Hunt et al.
2007/0054288 A1	3/2007	Su et al.	2009/0170713 A1	7/2009	van Eijk et al.
2007/0087360 A1	4/2007	Boyd	2009/0181375 A1	7/2009	Peter et al.
2007/0099208 A1	5/2007	Drmanac et al.	2009/0192044 A1	7/2009	Fouillet
2007/0116612 A1	5/2007	Williamson	2009/0197326 A1	8/2009	El Gamal et al.
2007/0128071 A1	6/2007	Shea et al.	2009/0202998 A1	8/2009	Schlumpberger et al.
2007/0128624 A1	6/2007	Gormley et al.	2009/0215633 A1	8/2009	van Eijk et al.
2007/0128656 A1	6/2007	Agrawal	2009/0226911 A1	9/2009	Mauk et al.
2007/0134723 A1	6/2007	Kozlov et al.	2009/0233802 A1	9/2009	Bignell et al.
2007/0141718 A1	6/2007	Bui et al.	2009/0239232 A1	9/2009	Kurn
2007/0161020 A1	7/2007	Luo et al.	2009/0253163 A1	10/2009	Xie et al.
2007/0161029 A1	7/2007	Li et al.	2009/0253581 A1	10/2009	van Eijk et al.
2007/0166705 A1	7/2007	Milton et al.	2009/0253582 A1	10/2009	Pena et al.
2007/0166725 A1	7/2007	McBride et al.	2009/0264299 A1	10/2009	Drmanac et al.
2007/0172873 A1	7/2007	Brenner et al.	2009/0270273 A1	10/2009	Burns et al.
2007/0178503 A1	8/2007	Jiang	2009/0280487 A1	11/2009	Hung et al.
2007/0184456 A1	8/2007	Chee et al.	2009/0283407 A1	11/2009	Shah et al.
2007/0207482 A1	9/2007	Church et al.	2009/0286249 A1	11/2009	Becker et al.
2007/0215466 A1	9/2007	Okada	2009/0289184 A1	11/2009	Deininger
2007/0231823 A1	10/2007	McKernan	2009/0291854 A1	11/2009	Weisinger-Mayr et al.
2007/0231824 A1	10/2007	Chee et al.	2009/0305237 A1	12/2009	Cantor et al.
2007/0243634 A1	10/2007	Pamula et al.	2009/0312193 A1	12/2009	Kim et al.
2007/0251824 A1	11/2007	Patton	2009/0321262 A1	12/2009	Adachi et al.
2007/0254305 A1	11/2007	Paik et al.	2010/0009871 A1	1/2010	Reed et al.
2007/0264656 A1	11/2007	Kawamura	2010/0014537 A1	1/2010	Jacquet et al.
2007/0269805 A1	11/2007	Hogers	2010/0031757 A1	2/2010	Hoyer
2007/0280517 A1	12/2007	De La Torre-Bueno et al.	2010/0035249 A1	2/2010	Hayashizaki et al.
2008/0003586 A1	1/2008	Hyde et al.	2010/0047790 A1	2/2010	Southern et al.
2008/0009071 A1	1/2008	Sogard	2010/0055733 A1	3/2010	Lutolf et al.
2008/0009420 A1	1/2008	Schroth et al.	2010/0069263 A1	3/2010	Shendure et al.
2008/0032301 A1	2/2008	Rank et al.	2010/0096266 A1	4/2010	Kim et al.
			2010/0099103 A1	4/2010	Hsieh et al.
			2010/0105021 A1	4/2010	Drmanac et al.
			2010/0105112 A1	4/2010	Heltze et al.
			2010/0108577 A1	5/2010	Wang et al.

(56)

**References Cited****U.S. PATENT DOCUMENTS**

2010/0111768 A1	5/2010	Banerjee et al.	2012/0279954 A1	11/2012	Ceremony et al.
2010/0113302 A1	5/2010	Williams	2012/0283106 A1	11/2012	Wang et al.
2010/0120043 A1	5/2010	Sood et al.	2012/0289414 A1	11/2012	Mitra et al.
2010/0120097 A1	5/2010	Matz et al.	2012/0301925 A1	11/2012	Belyaev
2010/0120098 A1	5/2010	Grunenwald et al.	2012/0308445 A1	12/2012	Roper et al.
2010/0126862 A1	5/2010	Sabin et al.	2012/0316086 A1	12/2012	Lin et al.
2010/0129874 A1	5/2010	Mitra et al.	2012/0322099 A1	12/2012	Lapen et al.
2010/0137143 A1	6/2010	Rothberg et al.	2013/0005594 A1	1/2013	Terbrueggen et al.
2010/0145037 A1	6/2010	Makarov et al.	2013/0005600 A1	1/2013	Olek
2010/0151447 A1	6/2010	Ely	2013/0023433 A1	1/2013	Luo et al.
2010/0151464 A1	6/2010	Stuelpnagel et al.	2013/0035239 A1	2/2013	Kong et al.
2010/0151511 A1	6/2010	Gereenizer et al.	2013/0040842 A1	2/2013	Lim et al.
2010/0159446 A1	6/2010	Haff et al.	2013/0052331 A1	2/2013	Kram et al.
2010/0173384 A1	7/2010	Johnsson et al.	2013/0053273 A1	2/2013	Juncker et al.
2010/0184614 A1	7/2010	Ye et al.	2013/0065768 A1	3/2013	Zheng et al.
2010/0184618 A1	7/2010	Namsaraev et al.	2013/0079232 A1	3/2013	Kain et al.
2010/0201809 A1	8/2010	Oyama et al.	2013/0096033 A1	4/2013	Routenberg
2010/0210475 A1	8/2010	Lee et al.	2013/0109595 A1	5/2013	Routenberg
2010/0216137 A1	8/2010	Bankaitis-Davis et al.	2013/0122516 A1	5/2013	Hong et al.
2010/0227329 A1	9/2010	Cuppens	2013/0146459 A1	6/2013	Bazant et al.
2010/0267590 A1	10/2010	Grudzien et al.	2013/0171621 A1	7/2013	Luo et al.
2010/0273219 A1	10/2010	May et al.	2013/0195963 A1	8/2013	Serda et al.
2010/0273679 A1	10/2010	Cuppoletti et al.	2013/0202718 A1	8/2013	Pepin et al.
2010/0282617 A1	11/2010	Rothberg et al.	2013/0203100 A1	8/2013	Otter et al.
2010/0303722 A1	12/2010	Jin et al.	2013/0211249 A1	8/2013	Barnett et al.
2011/0015494 A1	1/2011	Spaulding	2013/0244884 A1	9/2013	Jacobson et al.
2011/0024511 A1	2/2011	Rietzler et al.	2013/0244895 A1	9/2013	Voros et al.
2011/0027772 A1	2/2011	Ahn et al.	2013/0252847 A1	9/2013	McKenna et al.
2011/0028685 A1	2/2011	Purkayastha et al.	2013/0260372 A1	10/2013	Buermann et al.
2011/0033854 A1	2/2011	Drmanac et al.	2013/0261019 A1	10/2013	Lin et al.
2011/0045462 A1	2/2011	Fu et al.	2013/0302801 A1	11/2013	Asbury et al.
2011/0048951 A1	3/2011	Wu	2014/0011289 A1	1/2014	Smith et al.
2011/0059436 A1	3/2011	Hardin et al.	2014/0011707 A1	1/2014	Ye et al.
2011/0059865 A1	3/2011	Smith et al.	2014/0065609 A1	3/2014	Hicks et al.
2011/0086774 A1	4/2011	Dunaway	2014/0066318 A1	3/2014	Friesen et al.
2011/0090563 A1	4/2011	Krasov	2014/0121118 A1	5/2014	Warner
2011/0111409 A1	5/2011	Sinicropi et al.	2014/0155274 A1	6/2014	Xie et al.
2011/0151451 A1	6/2011	Lemaire et al.	2014/0155297 A1	6/2014	Heinz
2011/0152111 A1	6/2011	Fan et al.	2014/0162293 A1	6/2014	Springer et al.
2011/0165178 A1	7/2011	Wasyluk et al.	2014/0227705 A1	8/2014	Vogelstein et al.
2011/0172115 A1	7/2011	Thompson et al.	2014/0342921 A1	11/2014	Weiner
2011/0177518 A1	7/2011	Kartalov et al.	2014/0378350 A1	12/2014	Hindson et al.
2011/0201515 A1	8/2011	Webster et al.	2015/0010860 A1	1/2015	Kataoka et al.
2011/0207134 A1	8/2011	Faham et al.	2015/0051085 A1	2/2015	Vogelstein et al.
2011/0223613 A1	9/2011	Gut	2015/0065371 A1	3/2015	Seppo et al.
2011/0237449 A1	9/2011	McMaster et al.	2015/0072867 A1	3/2015	Soldatov
2011/0244448 A1	10/2011	Shirai et al.	2015/0087027 A1	3/2015	Makarov et al.
2011/0245101 A1	10/2011	Chee et al.	2015/0344942 A1	12/2015	Friesen et al.
2011/0245111 A1	10/2011	Chee	2016/0003812 A1	1/2016	Porreca et al.
2011/0269155 A1	11/2011	Reker-Hadrup et al.	2016/0024576 A1	1/2016	Chee
2011/0269647 A1	11/2011	Ule et al.	2016/0138091 A1	5/2016	Chee et al.
2011/0275077 A1	11/2011	James	2016/0145677 A1	5/2016	Chee et al.
2011/0287435 A1	11/2011	Grunenwald et al.	2016/0201125 A1	7/2016	Samuels et al.
2012/0010091 A1	1/2012	Linnarson	2016/0291006 A1	10/2016	Trau et al.
2012/0021930 A1	1/2012	Schoen et al.	2016/0298180 A1	10/2016	Chee
2012/0046175 A1	2/2012	Rodesch et al.	2016/0304952 A1	10/2016	Boyden et al.
2012/0046178 A1	2/2012	Van Den Boom et al.	2016/0333403 A1	11/2016	Chee
2012/0065081 A1	3/2012	Chee	2017/0058339 A1	3/2017	Chee
2012/0077693 A1	3/2012	Cazalis et al.	2017/0058340 A1	3/2017	Chee
2012/0129248 A1	5/2012	Chee et al.	2017/0058345 A1	3/2017	Chee
2012/0135871 A1	5/2012	van Eijk et al.	2017/0088881 A1	3/2017	Chee
2012/0142014 A1	6/2012	Cai	2017/0166962 A1	6/2017	van Eijk et al.
2012/0157322 A1	6/2012	Myllykangas	2017/0266667 A1	9/2017	Mortillaro et al.
2012/0160683 A1	6/2012	Yé et al.	2018/0094316 A1	4/2018	Scott et al.
2012/0177543 A1	7/2012	Battrell	2018/0105808 A1	4/2018	Mikkelsen et al.
2012/0195810 A1	8/2012	Cohen et al.	2018/0112209 A1	4/2018	Eshoo
2012/0196297 A1	8/2012	Yost et al.	2018/0179590 A1	6/2018	Belgrader et al.
2012/0202698 A1	8/2012	van Eijk et al.	2018/0201980 A1	7/2018	Chee et al.
2012/0202704 A1	8/2012	Fan et al.	2018/0216162 A1	8/2018	Belhocine et al.
2012/0220479 A1	8/2012	Ericsson et al.	2018/0247017 A1	8/2018	van Eijk et al.
2012/0245053 A1	9/2012	Shirai et al.	2018/0291439 A1	10/2018	van Eijk et al.
2012/0252702 A1	10/2012	Muratani et al.	2018/0334670 A1	11/2018	Bharadwaj et al.
2012/0258871 A1	10/2012	Kozlov et al.	2019/0002971 A1	1/2019	Koslover et al.
2012/0270305 A1	10/2012	Reed et al.	2019/0017106 A1	1/2019	Friesen et al.
2012/0270748 A1	10/2012	Chee et al.	2019/0024153 A1	1/2019	Friesen et al.
			2019/0024154 A1	1/2019	Friesen et al.
			2019/0064173 A1	2/2019	Bharadwaj et al.
			2019/0177800 A1	6/2019	Boutet et al.
			2019/0203275 A1	7/2019	Friesen et al.

(56)	References Cited					
U.S. PATENT DOCUMENTS						
2019/0264268 A1	8/2019	Frisen et al.	2022/0267844 A1	8/2022	Ramachandran Iyer et al.	
2019/0360034 A1	11/2019	Zhou et al.	2022/0275444 A1	9/2022	Salmanzadeh	
2019/0367982 A1	12/2019	Belhocine et al.	2022/0282329 A1	9/2022	Chell et al.	
2019/0367997 A1	12/2019	Bent et al.	2022/0290217 A1	9/2022	Frenz et al.	
2020/0109443 A1	4/2020	Chee	2022/0298560 A1	9/2022	Frisen et al.	
2020/0277663 A1	9/2020	Ramachandran Iyer et al.	2022/0325325 A1	10/2022	Chee et al.	
2020/0277664 A1	9/2020	Frenz	2022/0326251 A1	10/2022	Uytingco et al.	
2020/0283852 A1	9/2020	Oliphant et al.	2022/0333191 A1	10/2022	Mikkelsen et al.	
2020/0354774 A1	11/2020	Church et al.	2022/0333192 A1	10/2022	Uytingco	
2020/0399687 A1	12/2020	Frisen et al.	2022/0333195 A1	10/2022	Schnall-Levin et al.	
2020/0407781 A1	12/2020	Schnall-Levin et al.	2022/0334031 A1	10/2022	Delaney et al.	
2021/0010068 A1	1/2021	Chee et al.	2022/0348905 A1	11/2022	Dadhwal	
2021/0010070 A1	1/2021	Schnall-Levin et al.	2022/0348992 A1	11/2022	Stoeckius et al.	
2021/0017583 A1	1/2021	Chee et al.	2022/0356464 A1	11/2022	Kim et al.	
2021/0140982 A1	5/2021	Uytingco	2022/0364163 A1	11/2022	Stahl et al.	
2021/0172007 A1	6/2021	Chee et al.	2022/0389503 A1	12/2022	Mikkelsen et al.	
2021/0189475 A1	6/2021	Tentori et al.	2022/0389504 A1	12/2022	Chee et al.	
2021/0190770 A1	6/2021	Delaney et al.	2022/0403455 A1	12/2022	Ramachandran Iyer et al.	
2021/0198741 A1	7/2021	Williams	2022/0404245 A1	12/2022	Chell et al.	
2021/0199660 A1	7/2021	Williams et al.	2023/0002812 A1	1/2023	Stoeckius et al.	
2021/0214785 A1	7/2021	Stoeckius	2023/0014008 A1	1/2023	Shastry	
2021/0222241 A1	7/2021	Bharadwaj	2023/0017773 A1	1/2023	Kim et al.	
2021/0222242 A1	7/2021	Ramachandran Iyer	2023/0031305 A1	2/2023	Hernandez Neuta et al.	
2021/0222253 A1	7/2021	Uytingco	2023/0033960 A1	2/2023	Gallant et al.	
2021/0223227 A1	7/2021	Stoeckius	2023/0034039 A1	2/2023	Shahjamali	
2021/0230681 A1	7/2021	Patterson et al.	2023/0034216 A1	2/2023	Bava	
2021/0230692 A1	7/2021	Daugharty et al.	2023/0042817 A1	2/2023	Mignardi	
2021/0237022 A1	8/2021	Bava	2023/0047782 A1	2/2023	Tentori et al.	
2021/0238581 A1	8/2021	Mikkelsen et al.	2023/0056549 A1	2/2023	Dadhwal	
2021/0238664 A1	8/2021	Bava	2023/0064372 A1	3/2023	Chell et al.	
2021/0238675 A1	8/2021	Bava	2023/0069046 A1	3/2023	Chee et al.	
2021/0238680 A1	8/2021	Bava	2023/0077364 A1	3/2023	Patterson et al.	
2021/0247316 A1	8/2021	Bava	2023/0080543 A1	3/2023	Katiraei et al.	
2021/0262019 A1	8/2021	Alvarado Martinez et al.	2023/0081381 A1	3/2023	Chee et al.	
2021/0277460 A1	9/2021	Bava	2023/0100497 A1	3/2023	Frisen et al.	
2021/0285036 A1	9/2021	Yin et al.	2023/0111225 A1	4/2023	Chew et al.	
2021/0285046 A1	9/2021	Chell et al.	2023/0113230 A1	4/2023	Kim et al.	
2021/0292748 A1	9/2021	Frisen et al.	2023/0126825 A1	4/2023	Nagendran et al.	
2021/0292822 A1	9/2021	Frisen et al.	2023/0129552 A1	4/2023	Ramachandran Iyer	
2021/0317510 A1	10/2021	Chee et al.	2023/0135010 A1	5/2023	Tentori et al.	
2021/0317524 A1	10/2021	Lucero et al.	2023/0143569 A1	5/2023	Iyer et al.	
2021/0324457 A1	10/2021	Ramachandran Iyer et al.	2023/0145575 A1	5/2023	Gallant et al.	
2021/0332424 A1	10/2021	Schnall-Levin	2023/0147726 A1	5/2023	Hadrup et al.	
2021/0332425 A1	10/2021	Pfeiffer et al.	2023/0159995 A1	5/2023	Iyer et al.	
2021/0348221 A1	11/2021	Chell et al.	2023/0160008 A1	5/2023	Chell et al.	
2022/0002791 A1	1/2022	Frisen et al.	2023/0175045 A1	6/2023	Katsori et al.	
2022/0010367 A1	1/2022	Ramachandran Iyer et al.	2023/0183684 A1	6/2023	Gallant et al.	
2022/0017951 A1	1/2022	Ramachandran Iyer et al.	2023/0183785 A1	6/2023	Frisen et al.	
2022/0025446 A1	1/2022	Shah	2023/0194469 A1	6/2023	Tentori et al.	
2022/0025447 A1	1/2022	Tentori et al.	2023/0194470 A1	6/2023	Kim et al.	
2022/0033888 A1	2/2022	Schnall-Levin et al.	2023/0203478 A1	6/2023	Bent et al.	
2022/0049293 A1	2/2022	Frenz et al.	2023/02020454 A1	7/2023	Galonska et al.	
2022/0049294 A1	2/2022	Uytingco et al.	2023/0227811 A1	7/2023	Dadhwal	
2022/0064630 A1	3/2022	Bent et al.	2023/0228762 A1	7/2023	Uytingco et al.	
2022/0081728 A1	3/2022	Williams	2023/0242973 A1	8/2023	Frisen et al.	
2022/0088561 A1	3/2022	Brenan et al.	2023/0242976 A1	8/2023	Tentori et al.	
2022/0090058 A1	3/2022	Frisen et al.	2023/0265488 A1	8/2023	Gohil et al.	
2022/0090175 A1	3/2022	Uytingco et al.	2023/0265489 A1	8/2023	Uytingco et al.	
2022/0090181 A1	3/2022	Gallant et al.	2023/0265491 A1	8/2023	Tentori et al.	
2022/0098576 A1	3/2022	Dadhwal	2023/0267625 A1	8/2023	Tentori et al.	
2022/0098661 A1	3/2022	Chee et al.	2023/0279474 A1	9/2023	Katiraei	
2022/0106632 A1	4/2022	Galonska et al.	2023/0279477 A1	9/2023	Kvastad et al.	
2022/0106633 A1	4/2022	Engblom et al.	2023/0279481 A1	9/2023	Marrache et al.	
2022/0112486 A1	4/2022	Ramachandran Iyer et al.	2023/0287399 A1	9/2023	Gallant et al.	
2022/0119869 A1	4/2022	Ramachandran Iyer et al.	2023/0287475 A1	9/2023	Chell et al.	
2022/0127659 A1	4/2022	Frisen et al.	2023/0287481 A1	9/2023	Katsori et al.	
2022/0127660 A1	4/2022	Katiraei et al.	2023/0295699 A1	9/2023	Sukovich et al.	
2022/0127672 A1	4/2022	Stoeckius	2023/0295722 A1	9/2023	Bharadwaj	
2022/0145361 A1	5/2022	Frenz et al.	2023/0304072 A1	9/2023	Gohil et al.	
2022/0154255 A1	5/2022	Chee et al.	2023/0304074 A1	9/2023	Chee et al.	
2022/0170083 A1	6/2022	Khaled et al.	2023/0304078 A1	9/2023	Frisen et al.	
2022/0195422 A1	6/2022	Gallant et al.	2023/0313279 A1	10/2023	Giacomello et al.	
2022/0195505 A1	6/2022	Frisen et al.	2023/0323340 A1	10/2023	Dadhwal	
2022/0213526 A1	7/2022	Frisen et al.	2023/0323434 A1	10/2023	Yin et al.	
2022/0241780 A1	8/2022	Tentori et al.	2023/0323447 A1	10/2023	Schnall-Levin et al.	
			2023/0323453 A1	10/2023	Stoeckius	

# US 12,391,980 B2

Page 9

---

(56)

## References Cited

### U.S. PATENT DOCUMENTS

2023/0332138 A1	10/2023	Kim et al.	2024/0417784 A1	12/2024	Sukovich et al.
2023/0332212 A1	10/2023	Chew et al.	2025/0002980 A1	1/2025	Tentori et al.
2023/0332227 A1	10/2023	Ramachandran Iyer	2025/0002982 A1	1/2025	Stoeckius et al.
2023/0332247 A1	10/2023	Singh et al.	2025/0003956 A1	1/2025	Delaney et al.
2023/0351619 A1	11/2023	Tentori et al.	2025/0019689 A1	1/2025	Galonska et al.
2023/0358733 A1	11/2023	Chee	2025/0019749 A1	1/2025	Katirae et al.
2023/0366008 A1	11/2023	Chew et al.	2025/0066762 A1	2/2025	Man et al.
2023/0383285 A1	11/2023	Kim et al.	2025/0066770 A1	2/2025	Costa
2023/0383344 A1	11/2023	Stoeckius	2025/0073719 A1	3/2025	Cox et al.
2023/0392204 A1	12/2023	Chell et al.	2025/0075261 A1	3/2025	Kim
2023/0393071 A1	12/2023	Bava	2025/0101504 A1	3/2025	Nagendran et al.
2023/0407404 A1	12/2023	Baumgartner et al.	2025/0122564 A1	4/2025	Mignardi et al.
2023/0416808 A1	12/2023	Sukovich et al.	2025/0122565 A1	4/2025	Schnall-Levin et al.
2023/0416850 A1	12/2023	Singh et al.	2025/0129412 A1	4/2025	Uytingco et al.
2024/0002931 A1	1/2024	Bava	2025/0129421 A1	4/2025	Schnall-Levin et al.
2024/0011090 A1	1/2024	Chew et al.	2025/0137043 A1	5/2025	Tentori
2024/0018572 A1	1/2024	Mignardi	2025/0145984 A1	5/2025	Ma et al.
2024/0018575 A1	1/2024	Gallant et al.	2025/0146057 A1	5/2025	Schnall-Levin et al.
2024/0018589 A1	1/2024	Schnall-Levin et al.	2025/0154569 A1	5/2025	Frisen et al.
2024/0026445 A1	1/2024	Ramachandran Iyer et al.	2025/0154571 A1	5/2025	Stoeckius et al.
2024/0033743 A1	2/2024	Tentori et al.	2025/0154588 A1	5/2025	Ramachandran Iyer et al.
2024/0035937 A1	2/2024	Cox et al.	2025/0155446 A1	5/2025	Uytingco et al.
2024/0043908 A1	2/2024	Chew et al.	2025/0163501 A1	5/2025	Singh et al.
2024/0043925 A1	2/2024	Bent et al.	2025/0163509 A1	5/2025	Daugharthy et al.
2024/0052343 A1	2/2024	Gallant et al.	2025/0171833 A1	5/2025	Frisen et al.
2024/0053351 A1	2/2024	Uytingco et al.	2025/0171848 A1	5/2025	Chell et al.
2024/0060115 A1	2/2024	Chee et al.	2025/0179475 A1	6/2025	Borgstrom et al.
2024/0067953 A1	2/2024	Mikkelsen et al.	2025/0182305 A1	6/2025	Tentori et al.
2024/0068016 A1	2/2024	Frisen et al.	2025/0182503 A1	6/2025	Li et al.
2024/0068017 A1	2/2024	Lundeberg et al.	2025/0188526 A1	6/2025	Sukovich et al.
2024/0076723 A1	3/2024	Mignardi	2025/0189483 A1	6/2025	Kim et al.
2024/0080346 A1	3/2024	Engblom et al.	2025/0197847 A1	6/2025	Kim et al.
2024/0084365 A1	3/2024	Frisen et al.	2025/0197938 A1	6/2025	Bjorninen
2024/0084366 A1	3/2024	Chee	2025/0207125 A1	6/2025	Gupta et al.
2024/0084383 A1	3/2024	Ramachandran Iyer et al.	2025/0207195 A1	6/2025	Chell et al.
2024/0093274 A1	3/2024	Frisen et al.	2025/0208115 A1	6/2025	Bent
2024/0093290 A1	3/2024	Stahl et al.			
2024/0110228 A1	4/2024	Uytingco et al.			
2024/0124933 A1	4/2024	Chell et al.			
2024/0125772 A1	4/2024	Delaney et al.			
2024/0141327 A1	5/2024	Kim et al.			
2024/0158838 A1	5/2024	Alvarado Martinez et al.			
2024/0175080 A1	5/2024	Galonska et al.			
2024/0182968 A1	6/2024	Bava			
2024/0191286 A1	6/2024	Boutet et al.			
2024/0200121 A1	6/2024	Boutet			
2024/0209425 A1	6/2024	Yin et al.			
2024/0218427 A1	7/2024	Sukovich et al.			
2024/0218432 A1	7/2024	Mielinis			
2024/0219701 A1	7/2024	Tentori et al.			
2024/0253036 A1	8/2024	Kim et al.			
2024/0263218 A1	8/2024	Katirae et al.			
2024/0271190 A1	8/2024	Stoeckius et al.			
2024/0271195 A1	8/2024	Mikhail et al.			
2024/0279747 A1	8/2024	Williams			
2024/0287600 A1	8/2024	Iyer et al.			
2024/0294971 A1	9/2024	Galonska			
2024/0294974 A1	9/2024	Galonska et al.			
2024/0294975 A1	9/2024	Lin et al.			
2024/0301488 A1	9/2024	Stoeckius			
2024/0301489 A1	9/2024	Chee et al.			
2024/0360494 A1	10/2024	Costa et al.			
2024/0368711 A1	11/2024	Giacomello et al.			
2024/0377297 A1	11/2024	Cox et al.			
2024/0385088 A1	11/2024	Kim et al.			
2024/0392349 A1	11/2024	Frisen et al.			
2024/0392352 A1	11/2024	Stahl et al.			
2024/0392353 A1	11/2024	Engblom et al.			
2024/0401109 A1	12/2024	Kim et al.			
2024/0401117 A1	12/2024	Bava			
2024/0401118 A1	12/2024	Tentori et al.			
2024/0404301 A1	12/2024	Li et al.			
2024/0408593 A1	12/2024	Kim et al.			
2024/0416315 A1	12/2024	Bava			
2024/0417783 A1	12/2024	Chee et al.			

### FOREIGN PATENT DOCUMENTS

CN	1273609	11/2000
CN	1425133	6/2003
CN	1537953	10/2004
CN	1680604	10/2005
CN	1749752	3/2006
CN	1813059	8/2006
CN	1898398	1/2007
CN	1934452	3/2007
CN	1981188	6/2007
CN	101142325	3/2008
CN	101205560	6/2008
CN	101221182	7/2008
CN	101405400	4/2009
CN	101460633	6/2009
CN	101522915	9/2009
CN	202548048	11/2012
CN	102851369	1/2013
CN	102947330	2/2013
CN	105441549	3/2016
EP	0901631	3/1999
EP	0961110	12/1999
EP	1782737	5/2007
EP	1878502	1/2008
EP	1923471	5/2008
EP	1929039	6/2008
EP	1966393	9/2008
EP	2002017	12/2008
EP	2130913	12/2009
EP	2161336	3/2010
EP	1910562	12/2010
EP	2292788	3/2011
EP	2302070	3/2011
EP	2350648	8/2011
EP	2363504	9/2011
EP	2580351	4/2013
EP	2789696	10/2014

(56)	References Cited					
FOREIGN PATENT DOCUMENTS						
EP	2963127	1/2016	WO	WO 2006/065597	6/2006	
EP	3045544	7/2016	WO	WO 2006/074351	7/2006	
EP	3239304	11/2017	WO	WO 2006/081021	8/2006	
EP	1846164	5/2018	WO	WO 2006/081222	8/2006	
JP	2007-014297	1/2007	WO	WO 2006/084130	8/2006	
JP	2007-074967	3/2007	WO	WO 2006/117541	11/2006	
JP	2009-036694	2/2009	WO	WO 2006/124771	11/2006	
JP	2011-182702	9/2011	WO	WO 2006/137733	12/2006	
JP	2013-544498	12/2013	WO	WO 2007/000669	1/2007	
JP	2014-217381	11/2014	WO	WO 2007/010251	1/2007	
KR	10-2009-0000812	1/2009	WO	WO 2007/030373	3/2007	
KR	10-2009-0081260	7/2009	WO	WO 2007/037678	4/2007	
RU	2145635	2/2000	WO	WO 2007/041689	4/2007	
RU	2270254	2/2006	WO	WO 2007/053719	5/2007	
RU	2410439 C1	1/2011	WO	WO 2007/060599	5/2007	
WO	WO 1989/010977	11/1989	WO	WO 2007/073165	6/2007	
WO	WO 1991/006678	5/1991	WO	WO 2007/073171	6/2007	
WO	WO 1993/004199	3/1993	WO	WO 2007/073271	6/2007	
WO	WO 1995/023875	9/1995	WO	WO 2007/076128	7/2007	
WO	WO 1995/025116	9/1995	WO	WO 2007/076726	7/2007	
WO	WO 1995/035505	12/1995	WO	WO 2007/114693	10/2007	
WO	WO 1996/007669	3/1996	WO	WO 2007/120241	10/2007	
WO	WO 1997/031256	8/1997	WO	WO 2007/139766	12/2007	
WO	WO 1997/047640	12/1997	WO	WO 2007/145612	12/2007	
WO	WO 1998/010277	3/1998	WO	WO 2008/005673	1/2008	
WO	WO 1998/044151	10/1998	WO	WO 2008/022332	2/2008	
WO	WO 1999/032654	7/1999	WO	WO 2008/069906	6/2008	
WO	WO 1999/044062	9/1999	WO	WO 2008/075086	6/2008	
WO	WO 1999/044063	9/1999	WO	WO 2008/075086 A1 *	6/2008	..... G01N 33/554
WO	WO 1999/049082	9/1999	WO	WO 2008/157801	12/2008	
WO	WO 1999/063385	12/1999	WO	WO 2009/032167	3/2009	
WO	WO 1999/067641	12/1999	WO	WO 2009/036525	3/2009	
WO	WO 2000/017390	3/2000	WO	WO 2009/086487	7/2009	
WO	WO 2000/018957	4/2000	WO	WO 2009/137521	11/2009	
WO	WO 2000/024940	5/2000	WO	WO 2009/152928	12/2009	
WO	WO 2000/075373	12/2000	WO	WO 2009/156725	12/2009	
WO	WO 2001/006012	1/2001	WO	WO 2010/019826	2/2010	
WO	WO 2001/007915	2/2001	WO	WO 2010/027870	3/2010	
WO	WO 2001/009363	2/2001	WO	WO 2010/053587	5/2010	
WO	WO 2001/012862	2/2001	WO	WO 2010/060439	6/2010	
WO	WO 2001/042796	6/2001	WO	WO 2010/088517	8/2010	
WO	WO 2001/046402	6/2001	WO	WO 2010/100265	9/2010	
WO	WO 2001/059161	8/2001	WO	WO 2010/110929	9/2010	
WO	WO 2001/090415	11/2001	WO	WO 2010/126614	11/2010	
WO	WO 2001/096608	12/2001	WO	WO 2010/127186	11/2010	
WO	WO 2002/024952	3/2002	WO	WO 2011/008502	1/2011	
WO	WO 2002/040874	5/2002	WO	WO 2011/014879	2/2011	
WO	WO 2002/059355	8/2002	WO	WO 2011/019964	2/2011	
WO	WO 2002/059364	8/2002	WO	WO 2011/062933	5/2011	
WO	WO 2002/077283	10/2002	WO	WO 2011/071943	6/2011	
WO	WO 2002/088396	11/2002	WO	WO 2011/102903	8/2011	
WO	WO 2003/002979	1/2003	WO	WO 2011/127006	10/2011	
WO	WO 2003/003810	1/2003	WO	WO 2011/127099	10/2011	
WO	WO 2003/008538	1/2003	WO	WO 2011/143583	11/2011	
WO	WO 2003/010176	2/2003	WO	WO 2011/155833	12/2011	
WO	WO 2003/020261	3/2003	WO	WO 2012/022975	2/2012	
WO	WO 2003/077851	9/2003	WO	WO 2012/049316	4/2012	
WO	WO 2003/102233	12/2003	WO	WO 2012/058096	5/2012	
WO	WO 2003/106973	12/2003	WO	WO 2012/061832	5/2012	
WO	WO 2004/015080	2/2004	WO	WO 2012/071428	5/2012	
WO	WO 2004/028955	4/2004	WO	WO 2012/083225	6/2012	
WO	WO 2004/055159	7/2004	WO	WO 2012/129242	9/2012	
WO	WO 2004/067759	8/2004	WO	WO 2012/139110	10/2012	
WO	WO 2004/081225	9/2004	WO	WO 2012/140224	10/2012	
WO	WO 2004/108268	12/2004	WO	WO 2012/142213	10/2012	
WO	WO 2005/007814	1/2005	WO	WO 2012/148477	11/2012	
WO	WO 2005/010145	2/2005	WO	WO 2012/148497	11/2012	
WO	WO 2005/026387	3/2005	WO	WO 2012/159089	11/2012	
WO	WO 2005/042759	5/2005	WO	WO 2012/168003	12/2012	
WO	WO 2005/067648	7/2005	WO	WO 2013/022807	2/2013	
WO	WO 2005/084367	9/2005	WO	WO 2013/033271	3/2013	
WO	WO 2005/113804	12/2005	WO	WO 2013/040257	3/2013	
WO	WO 2006/020515	2/2006	WO	WO 2013/090390	6/2013	
WO	WO 2006/056861	6/2006	WO	WO 2013/123442	8/2013	
WO	WO 2006/064199	6/2006	WO	WO 2013/131962	9/2013	
			WO	WO 2013/138510	9/2013	
			WO	WO 2013/142389	9/2013	
			WO	WO 2013/150083	10/2013	
			WO	WO 2013/025952	12/2013	

(56)	References Cited		WO	WO 2024/035844	2/2024	
FOREIGN PATENT DOCUMENTS						
WO	WO 2014/060483	4/2014	WO	WO 2024/081212	4/2024	
WO	WO 2014/210223	12/2014	WO	WO 2024/086167	4/2024	
WO	WO 2014/210225	12/2014	WO	WO 2024/086776	4/2024	
WO	WO 2014/210353	12/2014	WO	WO 2024/102809	5/2024	
WO	WO 2017/013170	1/2017	WO	WO 2024/137826	6/2024	
WO	WO 2020/076979	4/2020	WO	WO 2024/145224	7/2024	
WO	WO 2020/123305	6/2020	WO	WO 2024/145441	7/2024	
WO	WO 2020/123309	6/2020	WO	WO 2024/145445	7/2024	
WO	WO 2020/123311	6/2020	WO	WO 2024/145491	7/2024	
WO	WO 2020/123316	6/2020	WO	WO 2024/206603	10/2024	
WO	WO 2020/123317	6/2020	WO	WO 2024/220882	10/2024	
WO	WO 2020/123318	6/2020	WO	WO 2024/238900	11/2024	
WO	WO 2020/123319	6/2020	WO	WO 2024/254316	12/2024	
WO	WO 2020/167862	8/2020	WO	WO 2025/029605	2/2025	
WO	WO 2020/176882	9/2020	WO	WO 2025/029627	2/2025	
WO	WO 2020/190509	9/2020	WO	WO 2025/043076	2/2025	
WO	WO 2020/198071	10/2020	WO	WO 2025/072119	4/2025	
WO	WO 2020/219901	10/2020	WO	WO 2025/090912	5/2025	
WO	WO 2021/091611	5/2021	WO	WO 2025/096581	5/2025	
WO	WO 2021/092433	5/2021	WO	WO 2025/101864	5/2025	
WO	WO 2021/097255	5/2021	OTHER PUBLICATIONS			
WO	WO 2021/133842	7/2021	U.S. Appl. No. 17/707,189, filed Mar. 29, 2022, Chell et al.			
WO	WO 2021/133849	7/2021	[No Author Listed], "HuSNP Mapping Assay User's Manual," Affymetrix Part No. 90094 (Affymetrix, Santa Clara, Calif.), GeneChip, 2000, 104 pages.			
WO	WO 2021/142233	7/2021	[No Author Listed], "Microarray technologies have excellent possibilities in genomics-related researches," Science Tools From Amersham Pharmacia Biotech, 1998, 3(4): 8 pages (with English Translation).			
WO	WO 2021/168261	8/2021	Adessi et al., "Solid phase DNA amplification: characterisation of primer attachment and amplification mechanisms," Nucl. Acids Res., Oct. 2000, 28(20):E87, 8 pages.			
WO	WO 2021/168278	8/2021	Affymetrix, "GeneChip Human Genome U133 Set," retrieved from the Internet: on the World Wide Web at <a href="http://affymetrix.com/support/technical/datasheets/hgu133_datasheet.pdf">affymetrix.com/support/technical/datasheets/hgu133_datasheet.pdf</a> , retrieved on Feb. 26, 2003.			
WO	WO 2021/207610	10/2021	Affymetrix, "Human Genome U95Av2," Internet Citation, retrieved from the internet: on the World Wide Web <a href="http://affymetrix.com">affymetrix.com</a> , retrieved on Oct. 2, 2002.			
WO	WO 2021/216708	10/2021	Agbavwe et al., "Efficiency, error and yield in light-directed maskless synthesis of DNA microarrays," Journal of Nanobiotechnology, Dec. 2011, 9:57, 17 pages.			
WO	WO 2021/225900	11/2021	Ahern et al., "Biochemical, Reagents Kits Offer Scientists Good Return On Investment," The Scientist, 1995, 9(15):20, 7 pages.			
WO	WO 2021/236625	11/2021	Ahlfen et al., "Determinants of RNA quality from FFPE samples," PLoS One, Dec. 2007, 2(12):e1261, 7 pages.			
WO	WO 2021/236929	11/2021	AJCC, "25 Lungs and 26 Pleural Mesothelioma," AJCC Cancer Staging Manual 7th Ed., Springer, 2010, pp. 253-278, 38 pages.			
WO	WO 2021/237056	11/2021	Akatsuka et al., "Rapid screening of T-cell receptor (TCR) variable gene usage by multiplex PCR: Application for assessment of clonal composition," Tissue Antigens, Jan. 5, 1999, 53:122-134.			
WO	WO 2021/237087	11/2021	Akatsuka et al., "T cell receptor clonal diversity following allogeneic marrow grafting," Human Immunology, Jun.-Jul. 1996, 48:125-134.			
WO	WO 2021/242834	12/2021	Akeroyd, "Click chemistry for the preparation of advanced macromolecular architectures," Stellenbosch University, PhD Dissertation, Mar. 2010, 138 pages.			
WO	WO 2021/247543	12/2021	Albretsen et al., "Applications of magnetic beads with covalently attached oligonucleotides in hybridization: Isolation and detection of specific measles virus mRNA from a crude cell lysate", Anal. Biochem. 189: 40-50, 1990.			
WO	WO 2021/247568	12/2021	Albretsen et al., "Optimal conditions for hybridization with oligonucleotides: a study with myc-oncogene DNA probes," Anal Biochem., Apr. 1988, 170(1):193-202.			
WO	WO 2021/252499	12/2021	Allawi et al., "Thermodynamics and NMR of Internal G.T Mismatches in DNA," Biochemistry, 1996, 36(34):10581-10594.			
WO	WO 2021/252576	12/2021	Almog et al., "The crystal structures of the psychrophilic subtilisin S41 and the mesophilic subtilisin Sph reveal the same calcium-loaded state," Proteins, Feb. 1, 2009, 74(2):489-496.			
WO	WO 2021/252591	12/2021	Altaras et al., "Production and formulation of adenovirus vectors," Adv Biochem Eng Biotechnol., Nov. 2005, 99:193-260.			
WO	WO 2021/263111	12/2021	Altman et al., "Phenotypic Analysis of Antigen-Specific T Lymphocytes," Science, Oct. 4, 1996, 274(5284):94-96.			
WO	WO 2022/025965	2/2022				
WO	WO 2022/060798	3/2022				
WO	WO 2022/060953	3/2022				
WO	WO 2022/061150	3/2022				
WO	WO 2022/087273	4/2022				
WO	WO 2022/098810	5/2022				
WO	WO 2022/099037	5/2022				
WO	WO 2022/103712	5/2022				
WO	WO 2022/109181	5/2022				
WO	WO 2022/140028	6/2022				
WO	WO 2022/147005	7/2022				
WO	WO 2022/147296	7/2022				
WO	WO 2022/164615	8/2022				
WO	WO 2022/178267	8/2022				
WO	WO 2022/198068	9/2022				
WO	WO 2022/221425	10/2022				
WO	WO 2022/226057	10/2022				
WO	WO 2022/236054	11/2022				
WO	WO 2022/256503	12/2022				
WO	WO 2022/271820	12/2022				
WO	WO 2023/018799	2/2023				
WO	WO 2023/034489	3/2023				
WO	WO 2023/076345	5/2023				
WO	WO 2023/086880	5/2023				
WO	WO 2023/102118	6/2023				
WO	WO 2023/122033	6/2023				
WO	WO 2023/287765	6/2023				
WO	WO 2023/150098	8/2023				
WO	WO 2023/150163	8/2023				
WO	WO 2023/150171	8/2023				
WO	WO 2023/215552	11/2023				
WO	WO 2023/225519	11/2023				
WO	WO 2023/229988	11/2023				
WO	WO 2023/250077	12/2023				
WO	WO 2024/015578	1/2024				

(56)

**References Cited****OTHER PUBLICATIONS**

- Altschul et al., "Basic local alignment search tool," *J. Mol. Biol.*, Oct. 5, 1990, 215(3):403-410.
- Anderson et al., "Microarrayed Compound Screening to Identify Activators and Inhibitors of AMP-Activated Protein Kinase," *J. of Biomolecular Screening*, 2004, 9:112.
- Andersson et al., "Analysis of protein expression in cell microarrays: a tool for antibody-based proteomics..," *J Histochem Cytochem.*, 4(12): 1413-1423, 2006.
- Andresen et al., "Deciphering the Antibodyome—Peptide Arrays for Serum Antibody Biomarker Diagnostics," *Current Proteomics*, 6(1), 1-12, 2009.
- Andresen et al., "Helicase-dependent amplification: use in OnChip amplification and potential for point-of-care diagnostics," *Expert Rev Mol Diagn.*, Oct. 2009, 9(7):645-650.
- Angenendt et al., "Cell-free Protein expression and functional assay in a nanowell chip format," *Analytical Chemistry*, 2004, 76(7):1844-49.
- Angenendt et al., "Generation of High Density Protein Microarrays by Cell-free in Situ Expression of Unpurified PCR Products," *Molecular and Cellular Proteomics*, (2006) Ch. 5.9, pp. 1658-1666.
- Appella, "Non-natural nucleic acids for synthetic biology," *Current Opinion in Chemical Biology*, Dec. 2009, 13(5-6): 687-696.
- Armani et al., "2D-PCR: a method of mapping DNA in tissue sections," *Lab on a Chip*, 2009, 9(24):3526-34.
- Atkinson, Overview of Translation: Lecture Manuscript, U of Texas (2000) DD. 6.1-6.8.
- Azioune et al., "Simple and rapid process for single cell micropatterning," *Lab Chip*, Jun. 2009, 9(11):1640-1642.
- Baerwald et al., "Discovery of genes implicated in whirling disease infection and resistance in rainbow trout using genome-wide expression profiling," *BMC Genomics*, Jan. 24, 2008, 9(1):37, 11 pages.
- Bains et al, "A Novel Method for Nucleic Acid Sequence Determination", *Journal of Theoretical Biology*, 1988, 135(3), 303-7.
- Baird et al., "Rapid SNP Discovery and Genetic Mapping Using Sequenced RAD markers," *PLoS One*, 2008, 3(10):e3376.
- Bakker et al., "Conditional MHC class I ligands and peptide exchange technology for the human MHC gene products HLA-A1, -A3, -All, and -B7," *Proc Natl Acad Sci USA*, Mar. 11, 2008, 105(10):3825-3830.
- Bandiera et al., "Nuclear Outsourcing of RNA Interference Components to Human Mitochondria," *PLoS One*, Jun. 2011, 6(6):e20746, 1-16.
- Baner et al., "Signal amplification of padlock probes by rolling circle replication," *Nucleic Acids Res.*, 1998, 26(22):5073-5078.
- Barbie et al., "Systematic RNA interference reveals that oncogenic KRAS-driven cancers require TBK1," *Nature*, Nov. 2009, 462(7269):108-12.
- Barnes, "PCR amplification of up to 35-kb DNA with high fidelity and high yield from lambda bacteriophage templates..," *Proc. Natl. Acad. Sci USA*, 91: 2216-2220, 1994.
- Bates et al., "Block copolymers-designer soft materials," *Phys Today*, Feb. 2000, 52:32-8.
- Baugh et al., "Quantitative analysis of mRNA amplification by in vitro transcription," *Nucleic Acids Res.*, 2001, 29:5:e29.
- Beattie et al., "Advances in genosensor research," *Clin Chem.*, May 1995, 41(5):700-6.
- Beier et al., "Versatile derivatization of solid support media for covalent bonding on DNA-microchips," *Nucleic Acids Res.*, May 1999, 27(9):1970-7.
- Bell, "A Simple Way to Treat PCR Products Prior to Sequencing Using ExoSAP-IT," *Biotechniques*, 2008, vol. 44, No. 6.
- Bentley et al, "Accurate whole human genome sequencing using reversible terminator chemistry", *Nature*, 2008, 456:53-59.
- Berent et al., "Comparison of oligonucleotide and long DNA fragments as probes in DNA and RNA dot, Southern, Northern, colony and plaque hybridizations," *Biotechniques*, 1985, 3(3):208-220 (Abstract Only).
- Berger et al., "Universal bases for hybridization, replication and chain termination," *Nucleic Acid Res.*, Aug. 2000, 28(15):2911-2914.
- Bessmerlykh et al., "Efficient Palladium-Catalyzed Synthesis of Aminopyridyl Phosphonates from Bromopyridines and Diethyl Phosphite," *Synthesis*, 2008, 10:1575-1579.
- Bibikova et al., "Quantitative gene expression profiling in formalin-fixed paraffin-embedded tissues using universal bead arrays," *The American Journal of Pathology*, Nov. 1, 2004, 165(5):1799-1807.
- Bielas et al., "Human cancers express a mutator phenotype," *Proc. Natl. Acad. Sci. USA*, 2006, 103(48): 18238-18242.
- Bielas et al., "Quantification of random genomic mutations," *Nat. Methods*, 2005, 2(4):285-290.
- Biol.wwu.edu [online], "Principles of Di-Base Sequencing and the Advantages of Color Space Analysis in the SOLiD System," 2008, retrieved on Mar. 11, 2022, retrieved from URL<<https://biol.wwu.edu/young/470/stuff/abi-solid.pdf>>, 4 pages.
- Birney, et al., "Identification and analysis of functional elements in 1% of the human genome by the Encode pilot project," *Nature*, 2007, 447:799-816.
- Blair et al., "Microarray temperature optimization using hybridization kinetics," *Methods Mol Biol.*, 2009, 529:171-96.
- Blanchard et al., "High-density oligonucleotide arrays," *Biosensors & Bioelectronics*, 1996, 11(6-7):687-690.
- Blanco et al., "A practical approach to FRET-based PNA fluorescence in situ hybridization," *Methods*, Dec. 2010, 52(4):343-51.
- Blandini et al., "Animal models of Parkinson's disease," *FEBS J.*, Apr. 2012, 279(7):1156-66.
- Blokzijl et al., "Profiling protein expression and interactions: proximity ligation as a tool for personalized medicine," *J Intern. Med.*, 2010, 268:232-245.
- Blow, "Tissue Issues," *Nature*, 448(23), 959-962, 2007.
- Boeke et al., "Transcription and reverse transcription of retrotransposons," *Annu Rev Microbiol*, 1989, 43:403-34.
- Bonfield et al., "The application of numerical estimates of base calling accuracy to DNA sequencing projects," *Nucleic Acids Research*, 1995, 23(8):1406-1410.
- Bos et al., "In Vitro Evaluation of DNA-DNA Hybridization as a Two-Step Approach in Radioimmunotherapy of Cancer," *Cancer Res.*, Jul. 1, 1994, 54(13):3479-3486.
- Boulé et al., "Terminal deoxynucleotidyl transferase indiscriminately incorporates ribonucleotides and deoxyribonucleotides," *J Biol Chem.*, Aug. 2001, 276(33):31388-93.
- Boutros et al., "The art and design of genetic screens: RNA interference," *Nat Rev Genet.*, Jul. 2008, 9(7):554-66.
- Bowtell, "The genesis and evolution of high-grade serous ovarian cancer," *Nat. Rev. Cancer*, 2010, (11 ):803-808 Abstract.
- Brandon et al., "Mitochondrial mutations in cancer," *Oncogene*, 2006, 25(34):4647-4662.
- Brenner et al., "In vitro cloning of complex mixtures of DNA on microbeads: physical separation of differentially expressed cDNAs," *Proc. Natl. Acad. Sci. USA*, 2000, 97, 1665-1670.
- Brenner et al., "Gene expression analysis by massively parallel signature sequencing (MPSS) on microbead arrays", *Nat. Biotech.* 18: 630-634, 2000.
- Brockman et al., "Quality scores and SNP detection in sequencing-by-synthesis systems," *Methods*, 2008, 18:763-770.
- Brow, "35—The Cleavase I enzyme for mutation and polymorphism scanning," *PCR Applications Protocols for Functional Genomics*, 1999, pp. 537-550.
- Brown et al., "Multiplex Three-Dimensional Brain Gene Expression Mapping in a Mouse Model of Parkinson's Disease," *Genome Research*, 2002, 12:868-884.
- Brown et al., "Retroviral integration: structure of the initial covalent product and its precursor, and a role for the viral IN protein," *Proc Natl Acad Sci USA*, Apr. 1989, 86(8):2525-9.
- Bullard et al., "Direct comparison of nick-joining activity of the nucleic acid ligases from bacteriophage T4," *Biochem. J.* 2006, 398:135-144.
- Burns et al., "Well-less, gel-permeation formats for ultra-HTS," *DDT*, 2001, 6(12):S40-S47.

(56)

**References Cited****OTHER PUBLICATIONS**

- Cai et al., "Glutathione-mediated shedding of PEG layers based on disulfide-linked cationomers for DNA delivery," *J. Mater. Chem.*, Sep. 20, 2011, 21(38):14639-14645.
- Calvert, "Materials science. Printing cells," *Science*, Oct. 2007, 318(5848):208-209.
- Cardona et al., "TrakEM2 0.9a User Manual," Sep. 8, 2011, retrieved on Jul. 29, 2022, retrieved from URL <[https://www.ini.uzh.ch/~acardona/trakem2\\_manual.html](https://www.ini.uzh.ch/~acardona/trakem2_manual.html)>, 38 pages.
- Cardullo et al., "Detection of nucleic acid hybridization by nonradiative fluorescence resonance energy transfer," *PNAS*, Dec. 1, 1988, 85:8790-8794.
- Carlson et al., "Function and Structure of a Prokaryotic Formylglycine-generating Enzyme," *J. of Biological Chemistry*, 2008, 283(29):20117-125.
- Carter et al., "Stabilization of an optical microscope to 0.1 nm in three dimensions," *Applied Optics*, 2007, 46:421-427.
- Cerritelli et al., "Ribonuclease H: the enzymes in eukaryotes," *FEBS Journal*, Mar. 2009, 276(6):1494-505.
- Cerutti et al., "Generation of sequence-specific, high affinity anti-DNA antibodies," *Journal of Biological Chemistry*, 2001, 276(16):12769-12773.
- Cha et al., "Specificity, Efficiency and Fidelity of PCR," *Genome Res.*, 1993, 3:518-29.
- Chandra et al., "Cell-free synthesis-based protein microarrays and their applications," *Proteomics*, 2009, 5(6):717-30.
- Chapin et al., "Rapid microRNA Profiling on Encoded Gel Microparticles," *Angew Chem Int Ed Engl.*, 2011, 50(10):2289-2293.
- Chatterjee et al., "Protein Microarray On-Demand: A Novel Protein Microarray System," *PLoS One*, 2008, 3(9):e3265.
- Chatterjee, et al., "Mitochondrial DNA mutations in human cancer. Oncogene," 2006, 25(34):4663-4674.
- Chen et al., "A Homogeneous, Ligase-mediated DNA diagnostic test," *Genome research*, 1998, 8(5):549-556.
- Chen et al., "DNA hybridization detection in a microfluidic Channel using two fluorescently labelled nucleic acid probes," *Biosensors and Bioelectronics*, 2008, 23:1878-1882.
- Chen et al., "Geometric control of cell life and death," *Science*, May 1997, 276(5317):1425-1428.
- Chen et al., "Gray-scale photolithography using microfluidic photomasks," *PNAS*, Feb. 2003, 100(4):1499-1504.
- Chen et al., "Parallel single nucleotide polymorphism genotyping by surface invasive cleavage with universal detection," *Anal Chem.*, Apr. 2005, 77(8):2400-5.
- Cheng et al., "Sensitive Detection of Small Molecules by Competitive Immunomagnetic-Proximity Ligation Assay," *Anal Chem*, 2012, 84:2129-2132.
- Cheng, "The Contrast Formation in Optical Microscopy," *Handbook Of Biological Confocal Microscopy*, 2006, Chapter 8, pp. 162-206.
- Chester et al., "Dimethyl sulfoxide-mediated primer Tm reduction: a method for analyzing the role of renaturation temperature in the polymerase chain reaction," *Anal Biochem*, Mar. 1993, 209(2):284-90.
- Chial, "DNA Sequencing Technologies Key to the Human Genome Project," *Nature Education*, 2008, 1(1):219, 7 pages.
- Chiang et al., "NFκappaB translocation in human microvessel endothelial cells using a four-compartment subcellular protein redistribution assay," *J Biochem Biophys Methods*, Nov. 2000, 46(1-2):53-68.
- Chrisey et al., "Covalent attachment of synthetic DNA to self-assembled monolayer films," *Nucleic Acids Res.*, Aug. 1996, 24(15):3031-9.
- Chu et al., "SV40 DNA transfection of cells in suspension: analysis of the efficiency of transcription and translation of T-antigen," *Gene*, Mar. 1981, 13(2):197-202, 1 page (Abstract Only).
- Chung et al., "Imaging single-cell signaling dynamics with a deterministic high-density single-cell trap array," *Anal Chem*, Sep. 2011, 83(18):7044-7052.
- Ciaccio et al., "Systems analysis of EGF receptor signaling dynamics with microwestern arrays," *Nat Methods*, Feb. 2010, 7(2):148-55.
- Cockcroft et al., "A single-molecule nanopore device detects DNA polymerase activity with single-nucleotide resolution," *J Am Chem Soc.*, Jan. 2008, 130(3):818-20.
- Colegio et al., "In vitro transposition system for efficient generation of random mutants of *Campylobacter jejuni*," *J Bacteriol.*, Apr. 2001, 183(7):2384-8.
- Condina et al., "A sensitive magnetic bead method for the detection and identification of tyrosine phosphorylation in proteins by MALDI-TOF/TOF MS," *Proteomics*, 2009, 9:3047-3057.
- Constantine et al., "Use of genechip high-density oligonucleotide arrays for gene expression monitoring," *Life Science News*, Amersham Life Science; 11-14, 1998.
- Cook et al., "The effects of secondary structure and O<sub>2</sub> on the formation of direct strand breaks upon UV irradiation of 5-bromodeoxyuridine-containing oligonucleotides," *Chem Biol.*, Jul. 1999, 6(7):451-9.
- Copeland et al., "Mitochondrial DNA Alterations in Cancer," *Cancer Invest.*, 2002, 20:57-59.
- Cornett et al., "MALDI imaging mass spectrometry: molecular snapshots of biochemical systems," *Nature Methods*, 2007, 4(10):828-833.
- Cox et al., "Tissue subcellular fractionation and protein extraction for use in mass-spectrometry-based proteomics," *Nat Protoc.*, 2006, 1(4):1872-8.
- Craig, "Transposon Tn7," *Curr Top Microbiol Immunol.*, 1996, 204:27-48.
- Craig, "V(D)J recombination and transposition: closer than expected," *Science*, Mar. 1996, 271(5255):1512, 1 page.
- Crisalli et al., "Multi-Path Quenchers: Efficient Quenching of Common Fluorophores," *Bioconjug Chem.*, Oct. 28, 2011, 22(11): 2345-2354.
- Crisalli et al., "Water-soluble Organocatalysts for Hydrazine and Oxime Formation," *J Org Chem*, Feb. 1, 2013, 78(3):1184-1189, 20 pages (Author Manuscript).
- Cujec et al. "Selection of v-abl tyrosine kinase substrate sequences from randomized peptide and cellular proteomic libraries using mRNA display," *Chemistry and Biology*, 2002, 9:253-264.
- Curtis et al., "Adhesion of cells to polystyrene surfaces," *J Cell Biol.*, Nov. 1983, 97(5):1500-1506.
- Czarnik, "Encoding methods for combinatorial chemistry," *Curr Opin Chem Biol.*, Jun. 1997, 1(1):60-6.
- Dahl et al., "Circle-to-circle amplification for precise and sensitive DNA analysis," *Proc. Natl. Acad. Sci.*, 2004, 101:4548-4553.
- Dalma-Weiszhausz et al., "The Affymetrix GeneChip platform: an overview," *Methods Enzymol.*, 2006, 410:3-28.
- Darmanis, et al., "ProteinSeq: High-Performance Proteomic Analyses by Proximity, Ligation and Next Generation Sequencing," *PLoS One*, 2011, 6(9):e25583.
- Daubendiek et al., "Rolling-Circle RNA Synthesis: Circular Oligonucleotides as Efficient Substrates for T7 RNA Polymerase," *J. Am. Chem. Soc.*, 1995, 117:77818-7819.
- Dawson et al., "Genetic animal models of Parkinson's disease," *Neuron*, Jun. 2010, 66(5):646-661.
- De Clercq, "A 40-year journey in search of selective antiviral chemotherapy," *Annu Rev Pharmacol Toxicol.*, 2011, 51:1-24.
- Deamer et al., "Characterization of nucleic acids by nanopore analysis," *Acc Chem Res.*, Oct. 2002, 35(10):817-25.
- Deamer et al., "Nanopores and nucleic acids: prospects for ultrarapid sequencing," *Trends Biotechnol.*, Apr. 2000, 18(4):147-51.
- Dean et al., "Comprehensive human genome amplification using multiple displacement amplification," *Proc Natl. Acad. Sci. USA* 99:5261-66, 2002.
- Dean et al., "Rapid Amplification Of Plasmid And Phage DNA Using Phi29 DNA Polymerase And Multiply-Primed Rolling Circle Amplification," *Genome Research*, Jun. 2001, 11:1095-1099.
- Dedeoglu et al., "On the Source of Asymmetry in Image Registration Problems," Technical Report, 2005, retrieve from URL <[https://www.ri.cmu.edu/pub\\_files/pub\\_dedeoglu\\_goksel\\_2005\\_1/dedeoglu\\_goksel\\_2005\\_1.pdf](https://www.ri.cmu.edu/pub_files/pub_dedeoglu_goksel_2005_1/dedeoglu_goksel_2005_1.pdf)>, 18 pages.

(56)

**References Cited****OTHER PUBLICATIONS**

- Deibel et al., "Biochemical properties of purified human terminal deoxynucleotidyltransferase," *J Biol Chem.*, May 1980, 255(9):4206-12.
- Deininger et al., "Allograft inflammatory factor-1 defines a distinct subset of infiltrating macrophages/microglial cells in rat and human gliomas," *Acta Neuropathol.*, Dec. 2000, 100(6):673-680.
- Deo et al., "Detection of mammalian microRNA expression by *in situ* hybridization with RNA oligonucleotides," *Dev Dyn.*, Sep. 2006, 235(9):2538-48.
- Devereux et al., "A comprehensive set of sequence analysis programs for the VAX," *Nucleic Acids Res.*, 1984, 12:387-395.
- Devine et al., "Efficient integration of artificial transposons into plasmid targets *in vitro*: a useful tool for DNA mapping, sequencing and genetic analysis," *Nucleic Acids Res.*, Sep. 1994, 22(18):3765-72.
- Dhindsa et al., "Virtual Electrowetting Channels: Electronic Liquid Transport with Continuous Channel Functionality," *Lab Chip*, 2010, 10:832-836.
- Diez-Roux et al., "A high-resolution anatomical atlas of the transcriptome in the mouse embryo," *PLoS Biol.*, Jan. 2011, 9(1):e1000582, 14 pages.
- Doddridge et al., "UV-induced strand break damage in single stranded bromodeoxyuridine-containing DNA oligonucleotides," *Chem Commun.*, 1998, p. 1997-1998.
- Dressman et al., "Transforming single DNA molecules into fluorescent magnetic particles for detection and enumeration of genetic variations," *Proc. Natl. Acad. Sci. USA*, 2003, 100:8817-8822.
- Drmanac et al., "Accurate sequencing by hybridization for DNA diagnostics and individual genomics," *Nature Biotechnology*, 16:54-58, 1998.
- Druley et al., "Quantification of rare allelic variants from pooled genomic DNA," *Nat. Methods* 6: 263-65, 2009.
- Duhr et al., "Why molecules move along a temperature gradient," *Proc Natl Acad Sci USA*, Dec. 2006, 103(52):19678-19682.
- Duncan et al., "Affinity chromatography of a sequence-specific DNA binding protein using teflon-linked oligonucleotides", *Anal. Biochem.* 169: 104-108, 1988.
- Eberwine et al., "Analysis of gene expression in single live neurons," *Proc. Natl. Acad. Sci., USA* 89, 3010-3014, 1992.
- Eberwine et al., "Complementary DNA Synthesis *In Situ*: Methods and Applications," *Methods in Enzymology*, 1992, 216:80-100.
- Eberwine, "Amplification of mRNA Populations Using aRNA Generated from Immobilized Oligo(dT)-T7 Primed cDNA," *BioTechniques* 20 (4), 584, 1996.
- Ebihara et al., "Molecular detection of dermatophytes and nondermatophytes in onychomycosis by nested polymerase chain reaction based on 28S ribosomal RNA gene sequences," *Br J Dermatol.*, Nov. 2009, 161(5):1038-44.
- Eguiluz et al., "Multitissue array review: A chronological description of tissue array techniques, applications and procedures," *Pathology Research and Practice*, 2006, 202:561-568.
- Ekins et al., "Microarrays: their origins and applications," *Trends in Biotechnology*, Jun. 1999, 17(6):217-218.
- Eldridge et al. "An *in vitro* selection strategy for conferring protease resistance to ligand binding peptides," *Protein Eng Des Sel.*, 22(11): 691-698, 2009.
- Ellington et al., "Antibody-based protein multiplex platforms: technical and operational challenges," *Clin Chem* 56(2): 186-193, 2010.
- Emmert-Buck et al., "Laser capture microdissection," *Science*, Nov. 1996, 274(5289):998-1001.
- Ergin et al., "Proteomic Analysis of PAXgene-Fixed Tissues," *J Proteome Res.*, 2010, 9(10):5188-96.
- Ertsey et al., "Coverslip Mounted-Immersion Cycled *In Situ* RT-PCR for the Localization of mRNA in Tissue Sections," *Biotechniques*, 1998, 24; pp. 92-100.
- Escholarship.org [online], "Methods and devices for fabricating and assembling DNA and protein arrays for high-throughput analyses [electronic resource]," 2010, retrieved on Jun. 8, 2022, retrieved from URL<<https://escholarship.org/uc/item/6tf7p46s>>, 155 pages.
- Espina et al., "Laser-capture microdissection," *Nat Protoc*, 2006, 1(2):586-603.
- Evers et al., "The effect of formaldehyde fixation on RNA: optimization of formaldehyde adduct removal," *J Mol Diagn.*, May 2011, 13(3):282-8.
- Extended European Search Report in European Appln. No. 11766613. 1, dated Jan. 15, 2014, 4 pages.
- Fahy et al., "Design and synthesis of polyacrylamide-based oligonucleotide supports for use in nucleic acid diagnostics," *Nucleic Acids Res.*, Apr. 1993, 21(8):1819-26.
- Falconnet et al., "Rapid, Sensitive and Real-Time Multiplexing Platform for the Analysis of Protein and Nucleic-Acid Biomarkers," *Anal. Chem.*, Jan. 7, 2015, 87:1582-1589.
- Falconnet et al., "Surface engineering approaches to micropattern surfaces for cell-based assays," *Biomaterials*, Jun. 2006, 27(16):3044-3063.
- Fan et al., "A versatile assay for high-throughput gene expression profiling on universal array matrices," *Genome Research*, May 1, 2004, 14(5):878-885.
- Fan et al., "Highly parallel SNP genotyping," *Cold Spring Symp. Quant. Biol.*, 68: 69-78, 2003.
- Fan et al., "Illumina Universal Bead Arrays," *Methods in Enzymology*, 2006, 410:57-73.
- Fang et al., "Fluoride-cleavable biotinylation phosphoramidite for 5'-end-labeling and affinity purification of synthetic oligonucleotides," *Nucleic Acids Res.*, Jan. 2003, 31(2):708-715.
- Faruqi et al., "High-throughput genotyping of single nucleotide polymorphisms with rolling circle amplification," *BMC Genomics*, Aug. 2001, 2:4, 10 pages.
- Ferreira et al., "Photocrosslinkable Polymers for Biomedical Applications," *Biomedical Engineering-Frontiers and Challenges*, Prof. Reza, 2011, 22 pages.
- Fiandaca et al., "Self-Reporting PNA/DNA Primers for PCR Analysis," *Genome Research*, Apr. 2001, 11:609-613.
- Fire and Xu, "Rolling replication of short DNA circles," *Proc. Natl. Acad. Sci.*, 92: 4641-4645, 1995.
- Fischer et al., "Hematoxylin and eosin staining of tissue and cell sections," *CSH Protoc.*, May 2008, 3(5):1-3.
- Fodor et al., "Light-Directed, Spatially Addressable Parallel Chemical Synthesis," *Science*, 251(4995), 767-773, 1995.
- Folch et al., "Microfabricated elastomeric stencils for micropatterning cell cultures," *J Biomed Mater Res*, Nov. 2000, 52(2):346-353.
- Fox, "Applications of Ultra-high-Throughput Sequencing," *Plant Systems Biology, Methods in Molecular Biology*, Belostotsky (ed.), Humana Press, 2009, 553:79-108.
- Fredriksson et al., "Multiplexed protein detection by proximity ligation for cancer detection," *Nature Methods*, 4(4): 327-29, 2007.
- Fredriksson et al., "Multiplexed proximity ligation assays to profile putative plasma biomarkers relevant to pancreatic and ovarian cancer," *Clin. Chem.*, 53(3): 582-89, 2008.
- Fredriksson et al., "Protein detection using proximity-dependent DNA ligation assays," *Nature Biotech.*, 20: 473-77, 2002.
- Frese et al., "Formylglycine Aldehyde Tag-Protein Engineering through a Novel Posttranslational Modification," *ChemBioChem*, 10: 425-27, 2009.
- Friedman et al., "The synthesis of high yields of full-length reverse transcripts of globin mRNA," *Nucleic Acids Res.*, Oct. 1, 1977, 4(10):3455-3471.
- Fu et al., "Counting individual DNA molecules by the stochastic attachment of diverse labels," *PNAS*, 108: 9026-9031, 2011.
- Fu et al., "Repeat subtraction-mediated sequence capture from a complex genome," *Plant J.*, Jun. 2010, 62(5):898-909.
- Fullwood et al., "Next-generation DNA sequencing of paired-end tags (PET) for transcriptome and genome analyses," *Genome Res.*, 19: 521-532, 2009.
- Galon et al., "The immune score as a new possible approach for the classification of cancer," *J Transl Med.*, Jan. 2012, 10:1, 4 pages.
- Gamper et al., "Gene expression profile of bladder tissue of patients with ulcerative interstitial cystitis," *BMC Genomics*, Apr. 28, 2009, 10(199):1-17.
- Gans et al., "Inkjet Printing of Polymers: State of the Art and Future Developments," *Advanced Materials*, Feb. 2004, 16(3):203-213.

(56)

**References Cited****OTHER PUBLICATIONS**

- Gao et al., "High density peptide microarrays. In situ synthesis and applications," *Molecular Diversity*, 8, 177-187, 2004.
- Geiss et al., "Direct multiplexed measurement of gene expression with color-coded probe pairs," *nature biotechnology*, 2008, 26(3):317-325.
- Genbank Accession No. AC009495.1, "*Homo sapiens* clone NH0490102, \*\*\* Sequencing in Progress \*\*\* , 12 unordered pieces," Aug. 24, 1999, 53 pages.
- Genbank Accession No. AC009495.5, "*Homo sapiens* BAC clone RP11-49012 from 2, complete sequence," Apr. 21, 2005, 32 pages.
- Genbank Accession No. AC037198.2, "*Homo sapiens* chromosome 15 clone CTD-2033D15 map 15q14, \*\*\* Sequencing in Progress \*\*\* , 62 unordered pieces," Apr. 25, 2000, 39 pages.
- Genbank Accession No. AC087379.2, "*Homo sapiens* chromosome 11 clone RP11-396020 map 11, \*\*\* Sequencing in Progress \*\*\* , 5 ordered pieces," Jul. 6, 2002, 47 pages.
- Genbank Accession No. AC087741.1, "*Homo sapiens* chromosome 17 clone RP11-334C17 map 17, Low-Pass Sequence Sampling," Jan. 22, 2001, 18 pages.
- Genbank Accession No. AC100826.1, "*Homo sapiens* chromosome 15 clone RP11-279F6 map 15, Low-Pass Sequence Sampling," Nov. 22, 2001, 21 pages.
- Genbank Accession No. AL445524.1, "*Homo sapiens* chromosome 1 clone RP11-295G20, Working Draft Sequence, 19 unordered pieces," Oct. 14, 2000, 47 pages.
- Genome.ucsc.edu, [online], "Genome Browser Gateway," 2000, retrieved on Jun. 11, 2021, retrieved from URL<<https://genome.ucsc.edu/cgi-bin/hgGateway>>, 3 pages.
- Gerard et al., "Excess dNTPs minimize RNA hydrolysis during reverse transcription," *Biotechniques*, Nov. 2002, 33(5):984, 986, 988, 990.
- Gerry et al., "Universal DNA Microarray Method for Multiplex Detection of Low Abundance Point Mutations," *J. Mol. Biol.*, 1999, 292:251-262.
- Giam et al., "Scanning probe-enabled nanocombinatorics define the relationship between fibronectin feature size and stem cell fate," *PNAS*, Mar. 2012, 109(12):4377-4382.
- Gibson et al., "Enzymatic assembly of DNA molecules up to several hundred kilobases," *Nat Methods*, May 2009, 6(5):343-5.
- Gilar et al., "Study of phosphorothioate-modified oligonucleotide resistance to 3'-exonuclease using capillary electrophoresis," *J Chromatogr B Biomed Sci Appl.*, Aug. 28, 1998, 714(1):13-20.
- Gill et al., "Nucleic acid isothermal amplification technologies: a review," *Nucleosides Nucleotides Nucleic Acids*, Mar. 2008, 27(3):224-43.
- Gilles et al., "Single nucleotide polymorphic discrimination by an electronic dot blot assay on semiconductor microchips," *Nat Biotechnol.*, Apr. 1999, 17(4):365-70.
- Glass et al., "SIMPLE: a sequential immunoperoxidase labeling and erasing method," *J. Histochem. Cytochem.*, Oct. 2009, 57(10):899-905.
- Gloor, "Gene targeting in *Drosophila*," *Methods Mol Biol.*, 2004, 260:97-114.
- Glorioso et al., "Development and Application of Herpes Simplex Virus Vectors for Human Gene Therapy," *Annu. Rev. Microbiol.*, 1995; 49:675-710, 1 page (Abstract Only).
- Gnanapragasam, "Unlocking the molecular archive: the emerging use of formalin-fixed paraffin-embedded tissue for biomarker research in urological cancer," *BJU International*, 105, 274-278, 2009.
- Goebl et al., "Development of a sensitive and specific *in situ* hybridization technique for the cellular localization of antisense oligodeoxynucleotide drugs in tissue sections," *Toxicologic Pathology*, Jun. 2007, 35(4):541-548.
- Goldkom and Prokrop, "A simple and efficient enzymatic method for covalent attachment of DNA to cellulose. Application for hybridization-restriction analysis and for *in vitro* synthesis of DNA probes.", *Nucleic Acids Res.* 14:9171-9191, 1986.
- Goldmeyer et al., "Development of a novel one-tube isothermal reverse transcription thermophilic helicase-dependent amplification platform for rapid RNA detection," *Journal of Molecular Diagnostics*, American Society for Investigative Pathology and the Association for Molecular Pathology, Nov. 1, 2007, 9(5):639-644.
- Goransson et al., "A single molecule array for digital targeted molecular analyses," *Nucleic Acids Res.*, Nov. 25, 2009, 37(1):e7, 9 pages.
- Goryshin et al., "Tn5 *in vitro* transposition," *J Biol Chem.*, Mar. 1998, 273(13):7367-74.
- Gotz et al., "Animal models of Alzheimer's disease and frontotemporal dementia," *Nat Rev Neurosci.*, Jul. 2008, 9(7):532-44.
- Graham et al., "A new technique for the assay of infectivity of human adenovirus 5 DNA," *Virology*, Apr. 1973, 52(2):456-467, 3 pages (Abstract Only).
- Grant et al., "Pathways and mechanisms of endocytic recycling," *Nat. Rev. Mol. Cell Biol.*, Sep. 2009, 10(9):597-608.
- Grigoryev, "How DNA microarrays are built," Bitesize Bio, first published Jul. 13, 2011, updated Oct. 2021, retrieved from URL <<https://bitesizebio.com/7206/introduction-to-dna-microarrays/#:~:text=Microarrays%20evolved%20from%20a%20technique%20known%20as%20Southern,filter%20paper>>, 11 pages.
- Grokhovsky, "Specificity of DNA cleavage by ultrasound," *Molecular Biology*, 2006, 40(2):276-283.
- Grünweller et al., "Locked Nucleic Acid Oligonucleotides," *BioDrugs*, Jul. 2007, 21(4): 235-243.
- Gruttadaira et al., "Supported proline and proline-derivatives as recyclable organocatalysts," *Chemical Society Reviews*, Aug. 1, 2008, 37(8):1666-1688.
- Gudjonsson et al., "Myoepithelial cells: their origin and function in breast morphogenesis and neoplasia," *J Mammary Gland Biol Neoplasia*, Jul. 2005, 10(3):261-72.
- Gunderson et al., "Decoding Randomly Ordered DNA Arrays," *Genome Research* 14: 870-877, 2004.
- Guo et al., "Direct fluorescence analysis of genetic polymorphisms by hybridization with oligonucleotide arrays on glass supports," *Nucleic Acids Res.*, Dec. 1994, 22(24):5456-65.
- Ha et al., "Self-assembly hollow nanosphere for enzyme encapsulation," *Soft Matter*, Feb. 11, 2010, 6, 1405-1408, 10 pages.
- Hadrup et al., "Parallel detection of antigen-specific T-cell responses by multidimensional encoding of MHC multimers," *Nat. Methods.*, Jul. 2009, 6(7): 520-526.
- Hafner et al., "Identification of microRNAs and other small regulatory RNAs using cDNA library sequencing," *Methods*, Jan. 2008, 44(1):3-12.
- Hahnke et al., "Striptease on glass: validation of an improved stripping procedure for *in situ* microarrays," *J Biotechnol.*, Jan. 2007, 128(1):1-13.
- Hajduk et al., "Drug discovery: A question of library design," *Nature*, Feb. 2011, 470(7332):42-43.
- Hamaguchi et al., "Direct reverse transcription-PCR on oligo(dT)-immobilized polypropylene microplates after capturing total mRNA from crude cell lysates," *Clin Chem.*, Nov. 1998, 44(11):2256-63.
- Hamers-Casterman et al., "Naturally occurring antibodies devoid of light chains," *Nature*, 1993, 363:446-448.
- Hammond et al., "Profiling cellular protein complexes by proximity ligation with dual tag microarray readout," *PLoS One*, 2012, 7(7):e40405.
- Han et al., "3C and 3C-based techniques: the powerful tools for spatial genome organization deciphering", *Molecular Cytogenetics* (2018) 11:21, 10 pages, 2018.
- Hanauer et al., "Separation of nanoparticles by gel electrophoresis according to size and shape," *Nano Lett.*, Sep. 2007, 7(9):2881-5.
- Hardenbol et al., "Highly multiplexed molecular inversion probe genotyping: over 10,000 targeted SNPs genotyped in a single tube assay," *Genome Res.*, Feb. 2005, 15(2):269-75.
- Hardenbol et al., "Multiplexed genotyping with sequence-tagged molecular inversion probes," *Nature Biotechnol.*, Jun. 2003, 21(6):673-678.
- Harold, "Molecules into Cells: Specifying Spatial Architecture," *Microbiology and Molecular Biology Reviews*, Dec. 2005, 69(4):544-564.
- Harris et al., "Chloroplast ribosomes and protein synthesis," *Microbiol. Mol. Biol. Rev.*, Dec. 1, 1994, 58(4): 700-754.

(56)

**References Cited****OTHER PUBLICATIONS**

- Harris et al., "The design and application of target-focused compound libraries," *Comb Chem High Throughput Screen*, Jul. 2011, 14(6):521-531.
- Hattersley et al., "Development of a microfluidic device for the maintenance and interrogation of viable tissue biopsies," *Lab Chip*, Nov. 2008, 8(11):1842-6.
- Hayes et al., "Electrophoresis of proteins and nucleic acids: I-Theory," *BMJ*, Sep. 1989, 299(6703):843-6.
- He et al., "In situ synthesis of protein arrays," *Current Opinion in Biotechnology*, 19:4-9, 2008.
- He et al., "Printing protein arrays from DNA arrays," *Nature Methods*, 5:175-77, 2008.
- He et al., "Ribosome display: Cell-free protein display technology," *Briefings in Functional Genomics and Proteomics*, Jul. 2002, 204-212.
- He, "Cell-free protein synthesis: applications in proteomics and biotechnology," *New Biotechnology* 25: 126-132, 2008.
- Healy, "Nanopore-based single-molecule DNA analysis," *Nanomedicine (Lond)*, Aug. 2007, 2(4):459-81.
- Hedskog et al., "Dynamics of HIV-1 Quasispecies during Antiviral Treatment Dissected using Ultra-Deep Pyrosequencing," *PLoS One*, 5(7):e11345, 2010.
- Hein et al., "Click Chemistry, A Powerful Tool for Pharmaceutical Sciences", *Pharm Res.*, 25(10):2216-2230, 2008.
- Heiter et al., "Site-Specific DNA-nicking Mutants of the Heterodimeric Restriction Endonuclease R.BbvC," *J. Mol. Biol.*, 2005, 348:631-40.
- Hejatkova et al., "In Situ Hybridization Techniques for mRNA Detection in Whole Mount *Arabidopsis* Samples," *Nature Protocols*, 2006, 1(4):1939-1946.
- Hendrickson et al., "High sensitivity multianalyte immunoassay using covalent DNA-labeled antibodies and polymerase chain reaction," *Nucleic Acid Research*, Feb. 11, 1995, 23(3):522-529.
- Hessner et al., "Genotyping of factor V G1691A (Leiden) without the use of PCR by invasive cleavage of oligonucleotide probes," *Clin Chem.*, Aug. 2000, 46(8 Pt 1):1051-6.
- Hewitt et al., "Tissue Handling and Specimen Preparation in Surgical Pathology," *Arch. Pathol. Lab. Med.*, 2008, 132:1929-1935.
- Hiatt et al., "Parallel, tag-directed assembly of locally-derived short sequence reads," *Nature Methods*, 7(2): 119-25, 2010.
- Higgins et al., "The nicking endonuclease N.BstNBI is closely related to Type II restriction endonucleases MlyI and PstI," *Nucleic Acids Res.*, 2001, 29:2492-2501.
- Hlubek et al., "Heterogeneous expression of Wnt/beta-catenin target genes within colorectal cancer," *Int J Cancer*, Nov. 2007, 121(9):1941-8.
- Ho et al., "Bacteriophage T4 RNA ligase 2 (gp24.1) exemplifies a family of RNA ligases found in all phylogenetic domains," *PNAS*, Oct. 2002, 99(20):12709-14.
- Ho et al., "Characterization of an ATP-Dependent DNA Ligase Encoded by Chlorella Virus PBCV-1," *Journal of Virology*, Mar. 1997, 71(3):1931-1937.
- Hofer et al., "Human protein atlas and the use of microarray technologies," *Curr Opin Biotechnol.*, Feb. 2008, 19(1):30-35.
- Holmstrom et al., "A highly sensitive and fast nonradioactive method for detection of polymerase chain reaction products," *Anal Biochem*, Mar. 1993, 209(2):278-83.
- Holscher et al., "Application of Laser-Assisted Microdissection for Tissue and Cell-Specific Analysis of RNA," *Progress in Botany*, Jan. 2008, 69(3):141-167.
- Hong et al., "Background-Free Detection of Single 5 nm Nanoparticles through Interferometric Cross-Polarization Microscopy," *Nano Letters*, Jan. 4, 2011, 11(2):541-547.
- Howell et al., "Glycosylases and AP-cleaving enzymes as a general tool for probe-directed cleavage of ssDNA targets," *Nucleic Acids Research*, Jan. 15, 2010, 38(7):e99, 10 pages.
- Howell et al., "iFRET: An Improved Fluorescence System for DNA-Melting Analysis," *Genome Research*, 2002, 12:1401-1407.
- Hoyer et al., "Electrostatic spraying: a novel technique for preparation of polymer coatings on electrodes," *Anal Chem*, Nov. 1996, 68(21):3840-4.
- Hsuih et al., "Novel, Ligation-Dependent PCR Assay for Detection of Hepatitis C Virus in Serum," *Journal of Clinical Microbiology*, Mar. 1996, 34(3):501-507.
- Hu et al., "High reproducibility using sodium hydroxide-stripped long oligonucleotide DNA microarrays," *Biotechniques*, Jan. 2005, 38(1):121-4.
- Hycultbiotech.com, [online], "Immunohistochemistry, Paraffin" Apr. 2010, retrieved on Apr. 16, 2020, retrieved from URL<[https://www.hycultbiotech.com/media/wysiwyg/Protocol\\_Immunohistochemistry\\_Paraffin\\_2.pdf](https://www.hycultbiotech.com/media/wysiwyg/Protocol_Immunohistochemistry_Paraffin_2.pdf)>, 3 pages.
- Hytönen et al., "Design and construction of highly stable, protease-resistant chimeric avidins," *J Biol Chem*, Mar. 2005, 280(11):10228-33.
- Ichikawa et al., "In vitro transposition of transposon Tn3," *J Biol. Chem.*, Nov. 1990, 265(31):18829-32, Abstract.
- Illumina Gene Expression Profiling, "Whole-Genome Expression Analysis Using the Sentrix Human-6 and HumanRef-8 Expression BeadChips," Illumina, 2005, pp. 1-8.
- Illumina, "M1\_SamplePrepSlides," Slides of Broad/Illumina Genome Analyzer Boot Camp, Feb. 2010, 76 pages.
- Illumina, "M2\_ClusterGeneration," Slides of Broad/Illumina Genome Analyzer (GA) Boot Camp, Feb. 2010, 45 pages.
- Illumina, "M3\_Sequencing," Slides of Broad/Illumina Genome Analyzer (GA) Boot Camp, Feb. 2010, 73 pages.
- Illumina.com [online], "Array-Based Gene Expression Analysis," 2011, retrieved on Dec. 13, 2021, retrieved from URL<[https://www.illumina.com/documents/products/datasheets/datasheet\\_gene\\_exp\\_analysis.pdf](https://www.illumina.com/documents/products/datasheets/datasheet_gene_exp_analysis.pdf)>, 5 pages.
- Im et al., "An Introduction to Performing Immunofluorescence Staining," *Biobanking: Methods and Protocols*, Method in Molecular Biology, Yong (ed.), 2019, 1897, Chapter 26, 299-311.
- Imbeaud et al., "Towards standardization of RNA quality assessment using user-independent classifiers of microcapillary electrophoresis traces," *Nucleic Acids Res.*, Mar. 2005, 33(6):e56, 12 pages.
- Inoue and Wittbrodt, "One for All—A Highly Efficient and Versatile Method for Fluorescent Immunostaining in Fish Embryos," *PLoS One* 6, e19713, 2011.
- Invitrogen, Immune Response Biomarker Profiling Service Report, Invitrogen, 2009, 1-33.
- Jabara et al., Accurate sampling and deep sequencing of the HIV-1 protease gene using a Primer ID. *PNAS* 108(50); 20166-20171, 2011.
- Jain, "Transport of molecules, particles, and cells in solid tumors," *Annu. Rev. Biomed. Eng.*, 1999, 1:241-263.
- Jamur and Oliver, "Permeabilization of cell membranes," *Method Mol. Biol.*, 588: 63-66, 2010.
- Janda et al., "16S rRNA Gene Sequencing for Bacterial Identification in the Diagnostic Laboratory: Pluses, Perils, and Pitfalls," *Journal of Clinical Microbiology*, 2007, 45(9):2761-2764.
- Jawhar et al., "Tissue Microarray: A rapidly evolving diagnostic and research tool," *Annals of Saudi Medicine*, Mar. 2009, 29(2):123-7.
- Jeffers, "A Basic Subroutine for Geary's Contiguity Ratio," *J. Royal Stat. Society, Series D*, Dec. 1973, 22(4):299-302.
- Jennane et al., "Photolithography of self-assembled monolayers: optimization of protecting groups by an electroanalytical method," *Can. J Chem.*, Dec. 1996, 74(12):2509-2517.
- Jensen et al., "Zinc fixation preserves flow cytometry scatter and fluorescence parameters and allows simultaneous analysis of DNA content and synthesis, and intracellular and surface epitopes," *Cytometry A*, Aug. 2010, 77(8):798-804.
- Jones et al., Comparative lesion sequencing provides insights into tumor evolution. *Proc. Natl. Acad. Sci. USA* 105(11): 4283-4288, 2008.
- Joos et al., "Covalent attachment of hybridizable oligonucleotides to glass supports," *Anal Biochem*, Apr. 1997, 247(1):96-101.
- Ju et al., "Supramolecular dendrimer capsules by cooperative binding," *Chem. Commun.*, Jan. 7, 2011, 47(1):268-270, 8 pages.
- Jucá et al., "Effect of dimethyl sulfoxide on reverse transcriptase activity," *Braz. J. Med. Biol. Res.*, Mar. 1995, 28(3):285-90.

- (56) **References Cited**
- OTHER PUBLICATIONS**
- Kainkaryam et al., "Pooling in high-throughput drug screening" *Curr Opin Drug Discov Devel.*, May 2009, 12(3):339-50.
- Kanehisa, "Use of statistical criteria for screening potential homologies in nucleic acid sequences", *Nucleic Acids Res.* 12: 203-213, 1984.
- Kap et al., "Histological Assessment of PAXgene Tissue Fixation and Stabilization Reagents," *PLoS One* 6, e27704, 10 pages, 2011.
- Kapteyn et al., "Incorporation of Non-Natural Nucleotides Into Template-Switching Oligonucleotides Reduces Background and Improves cDNA Synthesis From Very Small RNA Samples," *BMC Genomics*, 2010, 11(413): 1-9.
- Karlin et al., "Applications and statistics for multiple high-scoring segments in molecular sequences," *Proc. Natl. Acad. Sci.*, Jun. 15, 1993, 90:5873-7.
- Kelleher et al., "Characterization of RNA Strand Displacement Synthesis by Moloney Murine Leukemia Virus Reverse Transcriptase," *J Biol Chem*, Apr. 1998, 273(16):9976-86.
- Kibbe, "OligoCalc: an online oligonucleotide properties calculator," *Nucleic Acids Res.*, Jul. 2007, 35:W43-6.
- Kim et al., "Replication of DNA Microarrays Prepared by In Situ Oligonucleotide Polymerization and Mechanical Transfer," *Anal Chem.*, 2007, 79:7267-7274.
- Kim, "Development of Microdevices for Applications to Bioanalysis," Dissertation for the degree of Doctor of Philosophy, University of Texas at Austin, Aug. 2007, 176 pages.
- Kirby et al., "Cryptic plasmids of Mycobacterium avium: Tn552 to the rescue," *Mol Microbiol*, Jan. 2002, 43(1):173-86.
- Kleckner et al., "Tn10 and IS10 transposition and chromosome rearrangements: mechanism and regulation in vivo and in vitro," *Curr Top Microbiol Immunol.*, 1996, 204:49-82.
- Koch et al., "Photochemical immobilization of anthraquinone conjugated oligonucleotides and PCR amplicons on solid surfaces," *Bioconjugate Chem.*, Jul. 2000, 11(4):474-483.
- Kolb et al., "Click Chemistry: Diverse Chemical Function from a Few Good Reactions," *Angew. Chem. Int. Ed.*, 40(11): 2004-2021, 2001.
- Kolbert et al., "Ribosomal DNA sequencing as a tool for identification of bacterial pathogens," *Curr Opin Microbiol*, Jun. 1999, 2(3):299-305.
- Kong et al., "Duplex probes: a new approach for the detection of specific nucleic acids in homogenous assays," *Analytics Chimica Acta*, Sep. 2003, 491:135-143, 16 pages.
- König et al., "iCLIP reveals the function of hnRNP particles in splicing at individual nucleotide resolution," *Nat Struct Mol Biol*, Jul. 2010, 17(7):909-915.
- Korbel et al., "Paired-End Mapping Reveals Extensive Structural Variation in the Human Genome," *Science*, 318(5849): 420-426, 2007.
- Korlach et al., "Selective aluminum passivation for targeted immobilization of single DNA polymerase molecules in zero-mode waveguide nanostructures," *Proc. Natl. Acad. Sci. USA* 105, 1176-1181, 2008.
- Kozlov et al., "A High-Complexity Multiplexed Solution-Phase Assay for Profiling Protease Activity on Microarrays," *Comb Chem High Throughput Screen*, 11: 24-35, 2008.
- Kozlov et al., "A Highly Scalable Peptide-Based Assay System for Proteomics," *PLoS One*, 7(6):e37441, 2012.
- Kozlov et al., "A Method for Rapid Protease Substrate Evaluation and Optimization," *Comb Chem High Throughput Screen*, 9: 481-87, 2006.
- Kristensen et al., "High-Throughput Methods for Detection of Genetic Variation," *BioTechniques*, Feb. 2001, 30(2):318-332.
- Kuhn et al., "A novel, high-performance random array platform for quantitative gene expression profiling," *Genome Res*, 2004, 14:2347-2356.
- Kuhn et al., "Poly(A) Tail Length Is Controlled by the Nuclear Poly(A)-binding Protein Regulating the Interaction between Poly(A)
- Polymerase and the Cleavage and Poladenylation Specificity Factor," *The Journal of Biologic Chemistry*, Aug. 21, 2009, 284(34):22803-22814.
- Kuijpers et al. "Specific recognition of antibody-oligonucleotide conjugates by radiolabeled antisense nucleotides: a novel approach for two-step radioimmunotherapy of cancer," *Bioconjugate Chem.*, Jan. 1, 1993, 4(1):94-102.
- Kuiper et al., "Enzymes containing porous polymersomes as nano reaction vessels for cascade reactions," *Org. Biomol. Chem.*, Oct. 15, 2008, 6(23):4315-4318.
- Kumar et al., "Template-directed oligonucleotide strand ligation, covalent intramolecular DNA circularization and catenation using click chemistry," *J Am Chem Soc.*, May 2007, 129(21):6859-64.
- Kurz et al., "cDNA-Protein Fusions: Covalent Protein-Gene Conjugates for the In Vitro Selection of Peptides and Proteins," *ChemBioChem.*, 2: 666-72, 2001.
- Kwok, "High-throughput genotyping assay approaches," *Pharmacogenomics*, Feb. 2000, 1(1):95-100.
- Kwon et al., *Polyelectrolyte Gels-Fundamentals and Applications*, Nov. 10, 2006, *Polymer Journal*, 38, pp. 1211-1219.
- Lage et al., "Whole Genome Analysis of Genetic Alterations in Small DNA Samples Using Hyperbranched Strand Displacement Amplification and Array-CGH," *Genome Research* 13: 294-307, 2003.
- Lampe et al., "A purified mariner transposase is sufficient to mediate transposition in vitro," *EMBO J.*, Oct. 1996, 15(19):5470-9.
- Lamture et al., "Direct detection of nucleic acid hybridization on the surface of a charge coupled device," *Nucleic Acid Res.*, Jun. 1994, 22(11):2121-5.
- Landegren et al., "A Ligase-Mediated Gene Detection Technique," *Science*, 1988, 241(4869):1077-1080.
- Landegren et al., "Reading bits of genetic information: methods for single-nucleotide polymorphism analysis," *Genome Res.*, Aug. 1998, 8(8):769-76.
- Langdale et al., "A rapid method of gene detection using DNA bound to Sephadryl," *Gene* 36: 201-210, 1985.
- Larman et al., "Autoantigen discovery with a synthetic human peptidome," *Nature Biotechnology*, doi:10.1038/nbt.1856, vol. 29, No. 6, pp. 535-541, 2011.
- Larsen et al., "Characterization of a recombinantly expressed proteinase K-like enzyme from a psychrotrophic *Serratia* sp," *FEBS J.*, Jan. 2006, 273(1):47-60.
- Larsson et al., "In situ detection and genotyping of individual mRNA molecules," *Nat Methods*, May 2010, 7(5):395-7.
- Larsson et al., "In situ genotyping individual DNA molecules by target-primer rolling-circle amplification of padlock probes," *Nat Methods*, Dec. 2004, 1(3):227-32.
- Lassmann et al., "A Novel Approach For Reliable Microarray Analysis of Microdissected Tumor Cells From Formalin-Fixed and Paraffin-Embedded Colorectal Cancer Resection Specimens," *J Mol Med*, 87, 211-224, 2009.
- Laurell et al., "Chip integrated strategies for acoustic separation and manipulation of cells and particles," *Chem. Soc. Rev.*, Mar. 2007, 36(3):492-506.
- Lee et al., "A novel COL3A1 gene mutation in patient with aortic dissected aneurysm and cervical artery dissections," *Heart Vessels*, Mar. 2008, 23(2):144-8.
- Lee et al., "Cytokines in cancer immunotherapy," *Cancers (Basel)*, Oct. 2011, 3(4):3856-3893.
- Lee et al., "Hydrogels for Tissue Engineering," *Chemical Reviews*, 2001, 101(7):1869-1879.
- Lee et al., "Improving the efficiency of genomic loci capture using oligonucleotide arrays for high throughput resequencing," *BMC Genomics*, Dec. 2009, 10:646, 12 pages.
- Lee et al., "Protein nanoarrays generated by dip-pen nanolithography," *Science*, Mar. 2002, 295(5560):1702-1705.
- Lenard, "Viral Membranes," *Encyclopedia of Virology*, Jul. 2008, pp. 308-314.
- Leriche et al., "Cleavable linkers in chemical biology.", *Bioorganic & Medicinal Chemistry*, 20: 571-582, 2012.
- Levene et al., "Zero-Mode Waveguides for Single-Molecule Analyses at High Concentrations," *Science* 299, 682-686, 2003.

(56)

**References Cited****OTHER PUBLICATIONS**

- Li et al., "A photocleavable fluorescent nucleotide for DNA sequencing and analysis," *Proc. Natl. Acad. Sci.*, 100: 414-419, 2003.
- Li et al., "Beyond Moran's I: Testing for Spatial Dependence Based on the Spatial Autoregressive Model," *Geographical Analysis*, Sep. 18, 2007, 39(4):357-375.
- Li et al., "DNA molecules and configurations in a solid-state nanopore microscope," *Nat Mater.*, Sep. 2003, 2(9):611-5.
- Li et al., "RASL-seq for Massively Parallel and Quantitative Analysis of Gene Expression," *Curr Protoc Mol Biol.*, Apr. 2012, 4(13):1-10.
- Life Technologies, "Illumina TotalPrep RNA Amplification Kit," Ambion, 2011, 17 pages.
- Ligasová et al., "In situ reverse transcription: the magic of strength and anonymity," *Nucleic Acids Research*, 2010, 38(16):e167.
- Lin et al., "Replication of DNA microarrays from zip code masters," *J. Am. Chem. Soc.*, 2006, 128(10):3268-3272.
- Linnarsson, "Recent advances in DNA sequencing methods—general principles of sample preparation," *Experimental Cell Research*, 316: 1339-1343, 2010.
- Liu et al., "Barcoded oligonucleotides ligated on RNA amplified for multiplexed and parallel *in situ* analyses," *Nucleic Acids Res.*, Mar. 8, 2021, 49(10):e58, 15 pages.
- Liu et al., "Method for Quantitative Proteomics Research by Using Metal Element Chelated Tags Coupled with Mass Spectrometry," *Analytical Chemistry*, 2006, 78:6614-6621.
- Liu et al., "Preparation and Characterization of Temperature-Sensitive Poly(N-isopropylacrylamide)-b-poly(d,L-lactide) Microspheres for Protein Delivery," *Biomacromolecules*, 2003, 4(6):1784-1793.
- Liu et al., "Surface and interface control on photochemically initiated immobilization," *J Am Chem Soc.*, Nov. 2006, 128(43):14067-72.
- Liu et al., An integrated and sensitive detection platform for biosensing application based on Fe@Au magnetic nanoparticles as bead array carries Biosensors and Bioelectronics, 2010, 26(4):1442-1448.
- Lizardi et al., "Mutation detection and single-molecule counting using isothermal rolling-circle amplification," *Nat. Genet.* 19: 225-232, 1998.
- Lopez-Otin et al., "Protease degradomics: a new challenge for proteomics," *Nat Rev Mol Cell Biol.*, Jul. 2002, 3(7):509-19.
- Lu et al., "A microfluidic electroporation device for cell lysis," *Lab Chip.*, Jan. 2005, 5(1):23-29.
- Lund et al., "Assessment of methods for covalent binding of nucleic acids to magnetic beads, Dynabeads, and the characteristics of the bound nucleic acids in hybridization reactions," *Nucleic Acids Res.*, 16: 10861-80, 1988.
- Lundberg et al., "High-fidelity amplification using a thermostable DNA polymerase isolated from Pyrococcus furiosus," *Gene.*, 108(1): 1-6, 1991.
- Lundberg et al., "Homogeneous antibody-based proximity extension assays provide sensitive and specific detection of low-abundant proteins in human blood," *Nucleic Acids Res.*, 39(15): e102, 2011.
- Lundberg et al., "Multiplexed homogeneous proximity ligation assays for high-throughput protein biomarker research in serological material," *Mol Cell Proteomics*, 10(4): M110.004978, 2011.
- Lundin et al., "Increased throughput by parallelization of library preparation for massive sequencing," *PLoS One*, Apr. 2010, 5(4):e10029, 7 pages.
- Lundquist et al., "Parallel confocal detection of single molecules in real time," *Opt. Lett.* 33, 1026-1028, 2008.
- Lyamichev et al., "Invader assay for SNP genotyping," *Methods Mol Biol.*, 2003, 212:229-40.
- Lyamichev et al., "Polymorphism identification and quantitative detection of genomic DNA by invasive cleavage of oligonucleotide probes," *Nat Biotechnol.*, Mar. 1999, 17(3):292-6.
- Lyck, et al., "Immunohistochemical Markers for Quantitative Studies of Neurons and Glia in Human Neocortex," *J Histochem Cytochem* 56, 201-21, 2008.
- Lykidis et al., "Novel zinc-based fixative for high quality DNA, RNA and protein analysis," *Nucleic Acids Res.*, Jun. 2007, 35(12):e85, 10 pages.
- Mabruk et al., "In situ hybridization: detecting viral nucleic acid in formalin-fixed, paraffin-embedded tissue samples," *Expert Rev. Mol. Diagn.*, 2004, 4(5):653-661.
- MacBeath et al., "Printing proteins as microarrays for high-throughput function determination," *Science*, Sep. 2000, 289(5485):1760-1763.
- MacIntyre, "Unmasking antigens for immunohistochemistry," *Br J Biomed Sci.* 58, 190-6, 2001.
- Magaki et al., "An introduction to Performance of Immunohistochemistry," *Biobanking: Methods and Protocols*, Method in Molecular Biology, Yong (ed.), 2019, 1897, Chapter 25, 289-298.
- Makaryus et al., "Coronary venous angioplasty and stenting for biventricular pacemaker left ventricular lead implantation," *Journal of Invasive Cardiology*, 2008, 19(5), 3 pages.
- Malkov et al., "Multiplexed measurements of gene signatures in different analytes using the Nanostring nCounter™ Assay System," *BMC research notes.*, 2009, 2:80.
- Manz et al., "Phylogenetic Composition, Spatial Structure, and Dynamics of Lotic Bacterial Biofilms Investigated by Fluorescent *in situ* Hybridization and Confocal Laser Scanning Microscopy," *Microb Ecol*, May 1999, 37(4):225-237.
- Marras, "Selection of fluorophore and quencher pairs for fluorescent nucleic acid hybridization probes," *Methods Mol Biol.*, 2006, 335:3-16.
- Marsden et al., "3D small-molecule microarrays," *Chem. Commun.*, 2009, pp. 7107-7109.
- Martin, "Cutadapt removes adapter sequences from high-throughput sequencing reads," *EMBnet Journal*, 2011, 17(1):10-12.
- Massey et al., "Fluorescence resonance energy transfer (FRET) for DNA biosensors: FRET pairs and Förster distances for various dye-DNA conjugates," *Anal Chim Acta.*, May 2006, 568(1-2):181-9.
- Masuda et al., "Analysis of chemical modification of RNA from formalin-fixed samples and optimization of molecular biology applications for such samples," *Nucleic Acids Research*, Nov. 1999, 27(22):4436-4443.
- Materna et al., "High accuracy, high-resolution prevalence measurement for the majority of locally expressed regulatory genes in early sea urchin development," *Gene Expr Patterns.*, 2010, 10(4-5):177-184.
- Mattheyes et al., "Imaging with total internal reflection fluorescence microscopy for the cell biologist," *J Cell Sci.*, Nov. 2010, 123(Pt 21):3621-3628.
- McCloskey et al., "Encoding PCR Products with Batch-stamps and Barcodes," *Biochem. Genet.* 45:761-767, 2007.
- McGee, "Structure and Analysis of Affymetrix Arrays," UTSW Microarray Analysis Course, Oct. 28, 2005, 68 pages.
- McKernan et al., "Sequence and structural variation in a human genome uncovered by short-read, massively parallel ligation sequencing using two-base encoding," *Genome Res.*, 19: 1527-41, 2009.
- Megason et al., "Imaging in Systems Biology," *Cell* 130, Sep. 7, 2007, pp. 784-795.
- Metzker "Sequencing technologies—the next generation," *Nature Reviews Genetics*, 11: 31-46, 2010.
- Meyer et al., "Fast evolving 18S rRNA sequences from Solenogastres (Mollusca) resist standard PCR amplification and give new insights into mollusk substitution rate heterogeneity," *BMC Evol. Biol.*, Mar. 2010, 10:70, 12 pages.
- Michael et al., "Randomly Ordered Addressable High-Density Optical Sensor Arrays," *Analytical Chemistry*, American Chemical Society, Apr. 1998, 70:1242-1248.
- Mické et al., "Biobanking of fresh frozen tissue: RNA is stable in nonfixed surgical specimens," *Lab Invest.*, Feb. 2006, 86(2):202-11.
- Miele et al., "Mapping cis- and trans-chromatin interaction networks using chromosome conformation capture (3C)," *Methods Mol Biol.*, 2009, 464:105-21.
- Miller et al., "Basic Concepts of Microarrays and Potential Applications in Clinical Microbiology," *Clinical Microbiology Reviews*, vol. 22, No. 4, pp. 611-633, 2009.

(56)

**References Cited****OTHER PUBLICATIONS**

- Miller et al., "Rapid and Efficient Enzyme Encapsulation in a Dendrimer Silica Nanocomposite," *Macromolecular Bioscience*, Oct. 25, 2006, 6(10):839-845.
- Miner et al., "Molecular barcodes detect redundancy and contamination in hairpin-bisulfite PCR," *Nucleic Acids Res.*, Sep. 2004, 32(17):e135, 4 pages.
- Mir et al., "Sequencing by cyclic ligation and cleavage (CycliC) directly on a microarray captured template," *Nucleic Acids Research*, 37(1)e5, 8 pages, 2009.
- Mitra et al., "Digital genotyping and haplotyping with polymerase colonies," *Proc. Natl. Acad. Sci. USA*, May 2003, 100(10):5926-5931.
- Mitra et al., "Fluorescent in situ sequencing on polymerase colonies," *Anal Biochem*, Sep. 2003, 320(1):55-65.
- Mitra et al., "In situ localized amplification and contact replication of many individual DNA molecules," *Nucleic Acids Res.*, Dec. 1999, 27(24):e34, 6 pages.
- Mitsuhashi et al., "Gene manipulation on plastic plates," *Nature* 357: 519-520, 1992.
- Mizusawa et al., "A bacteriophage lambda vector for cloning with BamHI and Sau3A," *Gene*, 20: 317-322, 1982.
- Mlecinik et al., "Histopathologic-based prognostic factors of colorectal cancers are associated with the state of the local immune reaction," *J Clin Oncol.*, Feb. 2011, 29(6):610-8.
- Mohanty et al., "Bacterial/archaeal/organellar poly adenylation," *Wiley Interdiscip Rev RNA*, Mar.-Apr. 2011, 2(2):256-76, 36 pages (Author Manuscript).
- Morgan et al., "Characterization of the specific DNA nicking activity of restriction endonuclease N. BstNBI," *Biol. Chem.*, 2000, 381:1123-1125.
- Mortazavi et al., "Mapping and quantifying mammalian transcriptomes by RNA-Seq," *Nature Methods*, 5(7): 621-8, 2008.
- Moses et al., "Museum of spatial transcriptomics," *Nature Methods*, May 2022, 19:534-546.
- Moshrefzadeh et al., "Nonuniform photobleaching of dyed polymers for optical waveguides," *Applied Physics Letters*, 1993, 62:16-18.
- Motea et al., "Terminal deoxynucleotidyl transferase: the story of a misguided DNA polymerase," *Biochim Biophys Acta.*, May 2010, 1804(5):1151-66.
- Mueller et al., "RNA Integrity Number (RIN)—Standardization of RNA Quality Control," *Agilent Technologies*, 2004, 8 pages.
- Nadji et al., "Immunohistochemistry of tissue prepared by a molecular-friendly fixation and processing system," *Appl Immunohistochem Mol Morphol.*, Sep. 2005, 13(3):277-82.
- Nagahara et al., "Neuroprotective effects of brain-derived neurotrophic factor in rodent and primate models of Alzheimer's disease," *Nat Med.*, Mar. 2009, 15(3):331-337.
- Nagai et al., "Site-specific DNA cleavage by antisense oligonucleotides covalently linked to phenazine di-N-oxide," *J Biol. Chem.*, Dec. 1991, 266(35):23994-4002.
- Nakamura et al., "Biocompatible inkjet printing technique for designed seeding of individual living cells," *Tissue Eng.*, Nov. 2005, 11(11-12):1658-1666.
- Nakao et al., "Myosin heavy chain gene expression in human heart failure," *J Clin Invest.*, Nov. 1997, 100(9):2362-70.
- Nallur et al., "Signal amplification by rolling circle amplification on DNA microarrays," *Nucleic Acids Res.*, Dec. 1, 2001, 29(23):e118, 9 pages.
- Nam et al., "Nanoparticle-Based Bio-Bar Codes for the Ultrasensitive Detection of Proteins," *Science*, Sep. 26, 2003, 301(5641):1884-1886.
- Nandakumar et al., "How an RNA Ligase Discriminates RNA versus DNA Damage," *Molecular Cell*, 2004, 16(2):211-221.
- Nandakumar et al., "RNA Substrate Specificity and Structure-guided Mutational Analysis of Bacteriophage T4 RNA Ligase 2," *Journal of Biological Chemistry*, Jul. 2004, 279(30):31337-31347.
- Ng et al., "Gene identification signature (GIS) analysis for transcriptome characterization and genome annotation," *Nature Methods*, 2(2): 105-111, 2005.
- Ng et al., "Massively parallel sequencing and rare disease," *Human Malec. Genetics*, 19(2): R119-R124, 2010.
- Ng et al., "Multiplex sequencing of paired-end ditags (MS-PET): a strategy for the ultra-high-throughput analysis of transcriptomes and genomes," *Nucleic Acids Research*, Jul. 2006, 34(12): e84, 10 pages.
- Nichols et al., "RNA Ligases," *Curr Protoc Mol Biol.*, Oct. 2008, 84(1):3.15.1-3.15.4.
- Nicholson, "Diffusion and related transport mechanisms in brain tissue," *Rep. Prog. Phys.*, Jun. 2001, 64(7):815-884.
- Niedringhaus et al., "Landscape of next-generation sequencing technologies," *Anal Chem.*, Jun. 2011, 83(12):4327-41.
- Niemeyer, "The developments of semisynthetic DNA-protein conjugates," *Trends Biotechnol.*, Sep. 2002, 20(9): 395-401.
- Nikiforov et al., "The use of 96-well polystyrene plates for DNA hybridization-based assays: an evaluation of different approaches to oligonucleotide immobilization," *Anal Biochem*, May 1995, 227(1):201-9.
- Niklas et al., "Selective permeabilization for the high-throughput measurement of compartmentalized enzyme activities in mammalian cells," *Anal Biochem*, Sep. 2011, 416(2):218-27.
- Nilsson et al., "Padlock Probes: Circularizing Oligonucleotides for Localized DNA Detection," *Science*, Sep. 30, 1994, 265(5181):2085-2088.
- Nilsson et al., "RNA-templated DNA ligation for transcript analysis," *Nucleic Acids Res.*, Jan. 2001, 29(2):578-81.
- Nuovo, "In situ detection of microRNAs in paraffin embedded, formalin fixed tissues and the co-localization of their putative targets," *Methods*, 2010, 52(4):307-315.
- Nuovo, "In situ PCR: protocols and applications," *Genome Res*, Feb. 1995, 4 (4):151-167.
- Ohtsubo et al., "Bacterial insertion sequences," *Curr Top Microbiol Immunol.*, 1996, 204:1-26.
- Oleinikov et al., "Self-assembling protein arrays using electronic semiconductor microchips and in vitro translation," *J Proteome Res*, May-Jun. 2003, 2(3): 313-319.
- Olivier, "The Invader assay for SNP genotyping," *Mutat. Res.*, Jun. 2005, 573(1-2):103-110.
- Oren et al., "Selective lysis of bacteria but not mammalian cells by diastereomers of melittin: structure-function study," *Biochemistry*, Feb. 1997, 36(7):1826-35.
- Osada et al., "Epitope mapping using ribosome display in a reconstituted cell-free protein synthesis system," *J Biochem*, May 2009, 145(5): 693-700.
- O-Shannessy et al., "Detection and quantitation of hexa-histidine-tagged recombinant proteins on western blots and by a surface plasmon resonance biosensor technique," *Anal Biochem*, 229(1): 119-124, 1995.
- Ostuni et al., "Patterning Mammalian Cells Using Elastomeric Membranes," *Langmuir*, Aug. 2000, 16(20):7811-7819.
- Ozsolak et al., "Digital transcriptome profiling from attomole-level RNA samples," *Genome Res.*, Apr. 2010, 20(4):519-25.
- Palamanda et al., "Evaluation of CYP1A1 and CYP2B1/2 m-RNA Induction in Rat Liver Slices Using the NanoString® Technology: A Novel Tool for Drug Discovery Lead Optimization," *Drug metabolism letters*, Nov. 3, 2009, 3(3):171-175.
- Pandey et al., "Inhibition of terminal deoxynucleotidyl transferase by adenine dinucleotides. Unique inhibitory action of Ap5A," *FEBS Lett.*, Mar. 1987, 213(1):204-8.
- Parameswaran et al., "A pyrosequencing-tailored nucleotide barcode design unveils opportunities for large-scale sample multiplexing," *Nucleic Acids Research*, 2007, 35(19):e130.
- Park et al., "Cancer gene therapy using adeno-associated virus vectors," *Front Biosci.*, Jan. 2008, 13:2653-59.
- Park et al., "Detection of Hepatitis C Virus RNA Using Ligation-Dependent Polymerase Chain Reaction in Formalin-Fixed, Paraffin-Embedded Liver Tissues," *American Journal of Pathology*, Nov. 5, 1996, 149(5):1485-1491.

(56)

**References Cited****OTHER PUBLICATIONS**

- Park et al., "The Estimation of Breast Cancer Disease-Probability by Difference of Individual Susceptibility," *Cancer Res. Treat.*, Feb. 2003, 35(1):35-51, Abstract.
- Patil et al., "DNA-based therapeutics and DNA delivery systems: a comprehensive review," *AAPS J.*, Apr. 2005, 7(1):E61-77.
- Patton et al., "Rainbow's end: the quest for multiplexed fluorescence quantitative analysis in proteomics," *Current Opinion in Chemical Biology*, Feb. 1, 2002, 6(1):63-69.
- Pawlowski, "Photolithographic synthesis of high-density DNA probe arrays: Challenges and opportunities," *J. Vac. Sci. Technol. B*, 2007, 25:2537-2546.
- PCT International Preliminary Report on Patentability in International Appln. No. PCT/US2011/031308, dated Oct. 9, 2012, 7 pages.
- PCT International Search Report and Written Opinion in International Appln. No. PCT/US2011/031308, dated May 25, 2011, 8 pages.
- Pearson et al., "Improved tools for biological sequence comparison," *Proc. Natl. Acad. Sci.*, May 1988, 85:2444-2448.
- Pellestor et al., "The peptide nucleic acids (PNAs), powerful tools for molecular genetics and cytogenetics," *Eur J Hum Genet.*, Sep. 2004, 12(9):694-700.
- Pemov et al., "DNA analysis with multiplex microarray-enhanced PCR," *Nucl. Acids Res.*, Jan. 2005, 33(2):e11, 9 pages.
- Penland et al., "RNA expression analysis of formalin-fixed paraffin-embedded tumors," *Laboratory Investigation*, Apr. 2007, 87(4):383-391.
- Perler et al., "Intervening sequences in an Archaea DNA polymerase gene," *Pnas USA*, Jun. 1992, 89(12): 5577-5581.
- Perocchi et al., "Antisense artifacts in transcriptome microarray experiments are resolved by actinomycin D," *Nucleic Acids Res.*, 2007, 35(19):e128, 7 pages.
- Petterson et al., "Generations of sequencing technologies," *Genomics*, 2009, 105-111.
- Pipenburg et al., "DNA detection using recombination proteins," *PLoS Biol.*, Jul. 2006, 4(7):e204, 7 pages.
- Pirici et al., "Antibody elution method for multiple immunohistochemistry on primary antibodies raised in the same species and of the same subtypem," *J. Histochem. Cytochem.*, Jun. 2009, 57(6):567-75.
- Piston et al., "Fluorescent protein FRET: the good, the bad and the ugly," *Trends Biochem Sci.*, Sep. 2007, 32(9):407-14.
- Plasterk, "The Tcl/mariner transposon family," *Curr Top Microbiol Immunol.*, 1996, 204:125-43.
- Pluen et al., "Diffusion of macromolecules in agarose gels: comparison of linear and globular configurations," *Biophys J.*, Jul. 1999, 77(1):542-552.
- Polsky-Cynkin et al., "Use of DNA Immobilizedon Plastic and Agarose Supports to Detect DNA by Sandwich Hybridization," *Clin. Chem.* 31: 1438-1443, 1985.
- Porreca et al., "Polony DNA sequencing," *Curr Protoc Mol Biol.*, Nov. 2006, Chapter 7, Unit 7.8, pp. 7.8.1-7.8.22.
- Pringle et al., "In situ hybridization demonstration of polyadenylated RNA sequences in formalin-fixed paraffin sections using a biotinylated oligonucleotide poly d(T) probe," *Journal of Pathology*, Aug. 1989, 158:279-286.
- Proudfoot et al., "Integrating mRNA Processing with Transcription," *Cell*, Feb. 22, 2002, 108:501-512.
- U.S. Appl. No. 61/267,363, filed Dec. 7, 2009 (Year: 2009).
- Punwaney et al., "Human papillomavirus may be common within nasopharyngeal carcinoma of Caucasian Americans: investigation of Epstein-Barr virus and human papillomavirus in eastern and western nasopharyngeal carcinoma using ligation-dependent polymerase chain reaction," *Head & Neck*, Jan. 1999, 21(1):21-29.
- Qiu et al., "Mutation detection using Surveyor nuclease," *Biotechniques*, Apr. 2004, 36(4):702-707.
- Raab et al., "Human tRNA genes function as chromatin insulators," *EMBO J.*, Jan. 2012, 31(2):330-50.
- Rahimi et al., "Synthesis and Characterization of Thermo-Sensitive Nanoparticles for Drug Delivery Applications," *J. Biomed. Nanotechnol.* Dec. 2008, 4(4):482-490, 19 pages.
- Raj et al., "Imaging individual mRNA molecules using multiple singly labeled probes," *Nature Methods*, Oct. 2008, 5(10):877-879, 9 pages.
- Ramachandran et al., "Next-generation high-density self-assembling functional protein arrays," *Nature Methods*, Jun. 2008, 5(6):535-538.
- Ramanujan et al., "Diffusion and convection in collagen gels: implications for transport in the tumor interstitium," *Biophys. J.*, Sep. 2002, 83(3):1650-1660.
- Ranki et al., "Sandwich hybridization as a convenient method for the detection of nucleic acids in crude samples", *Gene* 21: 77-85, cellulose, 1983.
- Rasila et al., "Flexibility in MuA transposase family protein structures: functional mapping with scanning mutagenesis and sequence alignment of protein homologues," *PLoS One*, 2012, 7(5):e37922, 14 pages.
- Razonable, "Antiviral drugs for viruses other than human immunodeficiency virus," *Mayo Clinic Proceedings*, Oct. 2011, 86(10):1009-26.
- Reijenga et al., "Buffer Capacity, Ionic Strength and Heat Dissipation in Capillary Electrophoresis," *Journal of Chromatography A*, Sep. 13, 1996, 744(1-2):147-153.
- Reinartz et al., "Massively parallel signature sequencing (MPSS) as a tool for in-depth quantitative gene expression profiling in all organisms," *Brief Funct Genomic Proteomic*, Feb. 2002, 1(1):95-104.
- Rettig et al., "Large-scale single-cell trapping and imaging using microwell arrays," *Anal Chem*, Sep. 2005, 77(17):5628-5634.
- Reznikoff, "Tn5 as a model for understanding DNA transposition," *Mol Microbiol.*, Mar. 2003, 47(5):1199-206.
- Ristova et al., "Study of hydrogenated amorphous silicon thin films as a potential sensor for He—Ne laser light detection," *Applied Surface Science*, Sep. 2003, 218(1-4):44-53.
- Roberts et al., "RNA-peptide fusions for the in vitro selection of peptides and proteins," *PNAS USA*, Nov. 1997, 94: 12297-12302.
- Robins et al., "Comprehensive assessment of T-cell receptor β-chain diversity in αβ T cells," *Blood*, Nov. 5, 2009, 114(19):4099-4107.
- Robinson et al., "Small-sample estimation of negative binomial dispersion, with applications to SAGE data," *Biostatistics*, Apr. 2008, 9(2):321-332.
- Rogers et al., "Immobilization of oligonucleotides onto a glass support via disulfide bonds: A method for preparation of DNA microarrays," *Anal Biochem.* Jan. 1999, 266(1):23-30.
- Rogers et al., "Use of a novel cross-linking method to modify adenovirus tropism," *Gene Ther.*, Dec. 1997, 4(12):1387-92.
- Ronaghi et al., "A sequencing method based on real-time pyrophosphate," *Science*, Jul. 1998, 281(5375): 363-365.
- Ronaghi et al., "Real-time DNA sequencing using detection of pyrophosphate release," *Analytical Biochemistry*, Nov. 1996, 242(1):84-89.
- Ronaghi, "Pyrosequencing sheds light on DNA sequencing," *Genome Res.* Jan. 2001, 11(1):3-11.
- Rosenthal et al., "Cell patterning chip for controlling the stem cell microenvironment," *Biomaterials*, Jul. 2007, 28(21):3208-3216.
- Rouillard et al., "OligoArray 2.0: design of oligonucleotide probes for DNA microarrays using a thermodynamic approach," *Nuc. Acid Research*, Jun. 2003, 31(12): 3057-3062.
- Rountenberg et al., "Microfluidic probe: a new tool for integrating microfluidic environments and electronic wafer-probing," *Lab Chip*, Oct. 2009, 10(1):123-127.
- Rubin et al., "Whole-genome resequencing reveals loci under selection during chicken domestication," *Nature*, Mar. 2010, 464: 587-591.
- Rubina et al., "Hydrogel-based protein microchips: manufacturing, properties, and applications," *Biotechniques*, May 2003, 34(5):1008-14.
- Rush et al., "New Aldehyde Tag Sequences Identified by Screening Formylglycine Generating Enzymes in Vitro and in Vivo," *J. of American Chemical Society*, Aug. 2008, 130(37): 12240-12241.

(56)

**References Cited****OTHER PUBLICATIONS**

- Rusk, "Tracing Cell Lineage with 5hmC," *Nature Methods*, 2016, 13:710-711.
- Russell et al., "Molecular mechanisms of late endosome morphology, identity and sorting," *Curr. Opin. Cell Bio.*, Aug. 2006, 18(4):422-428.
- Samuelson et al., "The isolation of strand-specific nicking endonucleases from a randomized SapI expression library," *Nucleic Acids Res.*, 2004, 32:3661-3671.
- San Paulo et al., "High-resolution imaging of antibodies by tapping-mode atomic force microscopy: attractive and repulsive tip-sample interaction regimes," *Biophys J.*, Mar. 2000, 78(3):1599-1605.
- Sano et al., "Immuno-PCR: Very Sensitive Antigen Detection by Means of Specific Antibody-DNA Conjugates," *Science*, Oct. 2, 1992, 258(5079):120-122.
- Schellings et al., "Absence of SPARC results in increased cardiac rupture and dysfunction after acute myocardial infarction," *J Exp Med.*, Jan. 2009, 206(1):113-23.
- Schena et al., "Quantitative monitoring of gene expression patterns with a complementary DNA microarray," *Science*, Oct. 1995, 270(5235):467-470.
- Schena et al., "Entering the Postgenome Era," *Science*, 1995, 270:368-9, 371.
- Schlupak et al., "Glass surfaces grafted with high-density poly(ethylene glycol) as substrates for DNA oligonucleotide microarrays," *Langmuir*, Jan. 2006, 22: 277-285.
- Schmitt et al., "Detection of ultra-rare mutations by next-generation sequencing," *PNAS* (2012) 109:14508-14523.
- Scholz et al., "The Molecular Chaperone Hsp90 Is Required for Signal Transduction by Wild-Type Hck and Maintenance of Its Constitutively Active Counterpart1," *Cell Growth Differ.*, Aug. 2001, 12(8):409-417.
- Schouten et al., "Relative quantification of 40 nucleic acid sequences by multiplex ligation-dependent probe amplification," *Nucleic Acids Res.*, Jun. 2002, 30(12):e57, 13 pages.
- Schroeder et al., "The RIN: an RNA integrity No. for assigning integrity values to RNA measurements," *BMC Molecular Biology*, Jan. 2006, 7:3, 14 pages.
- Schweitzer et al., "Immunoassays with rolling circle DNA amplification: A versatile platform for ultrasensitive antigen detection," *Proc. Natl Acad. Sci. USA*, May 22, 2000, 97:10113-119.
- Schweitzer et al., "Multiplexed protein profiling on microarrays by rolling-circle amplification," *Nature Biotechnology*, Apr. 2002, 20(4):359-365.
- Schwers et al., "A high-sensitivity, medium-density, and target amplification-free planar waveguide microarray system for gene expression analysis of formalin-fixed and paraffin-embedded tissue," *Clin. Chem.*, Nov. 2009, 55(11):1995-2003.
- Scicchitano et al., "Preliminary Comparison of Quantity, Quality, and Microarray Performance of RNA Extracted From Formalin-fixed, Paraffin-embedded, and Unfixed Frozen Tissue Samples," *J. Histochemistry & Cytochemistry*, 2006, 54(11):1229-1237.
- ScienceDirect.com [online]. "Plant Fibers." Definition, 2011, retrieved on Apr. 13, 2022, retrieved from URL<<https://www.sciencedirect.com/topics/agricultural-and-biological-sciences/plant-fibers>>, 9 pages.
- Sekar et al., "Fluorescence resonance energy transfer (FRET) microscopy imaging of live cell protein localizations," *J Cell Biol.*, Mar. 2003, 160(5):629-33.
- Sel et al., "Feasibility of Employing Model-Based Optimization of Pulse Amplitude and Electrode Distance for Effective Tumor Electroporation," *IEEE Transactions on Biomedical Engineering*, May 2007, 54(5):773-781.
- Sergeeva et al., "Display technologies: Application for the discovery of drug and gene delivery agents," *Advanced Drug Delivery Reviews* (2006) 58(15):1622-1654.
- Seurnick-Servoss et al., "Evaluation of Surface Chemistries for Antibody Microarrays," *Anal Biochem*, 371(1): 105-115, 2007.
- Shalon et al., "A DNA microarray system for analyzing complex DNA samples using two-color fluorescent probe hybridization," *Genome Res.*, Jul. 1996, 6(7):639-45.
- Shastry, "SNPs in disease gene mapping, medicinal drug development and evolution," *J. Hum. Genet.*, 2007, 52:871-880.
- Shelbourne et al., "Fast copper-free click DNA ligation by the ring-strain promoted alkyne-azide cycloaddition reaction," *Chem. Commun.*, 47: 6257-6259, 2011.
- Shendure et al., "Accurate Multiplex Polony Sequencing of an Evolved Bacterial Genome," *Science*, 2005, 309:1728-1732.
- Shi et al., "The MicroArray Quality Control (MAQC) project shows inter- and intraplatform reproducibility of gene expression measurements," *Nature Biotechnology*, 2006, 24(9):1151-61.
- Shi, "Enabling large-scale pharmacogenetic studies by high-throughput mutation detection and genotyping technologies," *Clin. Chem.*, Feb. 2001, 47(2):164-172.
- Shibata et al., "Detection of human papilloma virus in paraffin-embedded tissue using the polymerase chain reaction," *J Exp Med.*, Jan. 1988, 167(1):225-30.
- Shirai et al., "Novel Tools for Analyzing Gene Expressions in Single Cells," The 5th International Workshop on Approaches to Single-Cell Analysis, The University of Tokyo, Mar. 3-4, 2011, 1 page.
- Shoemaker et al., "Quantitative phenotypic analysis of yeast deletion mutants using a highly parallel molecular bar-coding strategy," *Nature genetics* (1996) 14:450-456.
- Shults et al., "A multiplexed protein kinase assay," *Chem Bio Chem* (2007) 8:933-942.
- Sieverzon et al., "Transcriptome analysis in primary neural stem cells using a tag cDNA amplification method," *BMC Neuroscience*, Dec. 2005, 6: 28.
- Simonis et al., "Nuclear organization of active and inactive chromatin domains uncovered by chromosome conformation capture-on-chip (4C)," *Nat Genet.*, Nov. 2006, 38(11):1348-54.
- Slomovic et al., "Addition of poly(A) and poly(A)-rich tails during RNA degradation in the cytoplasm of human cells," *Proc Natl Acad Sci USA*, Apr. 2010, 107(16):7407-12.
- Slonim and Yanai, "Getting started in gene expression microarray analysis," *Plos Computational Biology*, 2009, 5(10):e1000543.
- Soderberg et al. "Characterizing proteins and their interactions in cells and tissues using the *in situ* proximity ligation assay," *Methods*, Jul. 2008, 45(3):227-232.
- Soderberg et al., "Direct observation of individual endogenous protein complexes *in situ* by proximity ligation," *Nature Methods*, 2006, 3:995-1000.
- Soen et al., "Detection and Characterization of Cellular Immune Responses Using Peptide-MHC Microarrays," *PLoS Biology*, Dec. 22, 2003, 1(3):429-438.
- Son et al., "A platform for ultrasensitive and selective multiplexed marker protein assay toward early-stage cancer diagnosis," *Nanomedicine*, Feb. 7, 2007, 2(1):79-82.
- Soni and Meller, "Progress toward ultrafast DNA sequencing using solid-state nanopores," *Clin Chem.*, 2007, 53: 1996-2001.
- Spiess et al., "A highly efficient method for long-chain cDNA synthesis using trehalose and betaine," *Anal. Biochem.*, Feb. 2002, 301(2):168-74.
- Spurgeon et al., "High Throughput Gene Expression Measurement with Real Time PCR in a Microfluidic Dynamic Array," *Plos ONE*, 2008, 3(2):e1662.
- Stahl et al., "Visualization and analysis of gene expression in tissue sections by spatial transcriptomics," *Science*, Jul. 2016, 353(6294):78-82.
- Stahl et al., "Visualization and analysis of gene expression in tissue sections by spatial transcriptomics," *Science*, Jun. 2016, Supplementary Materials, 353(6294):78-82, 41 pages.
- Stanton et al., "Altered patterns of gene expression in response to myocardial infarction," *Circulation research*, May 12, 2000, 86(9), 939-945.
- Stevens Jr. et al., "Enhancement of phosphoprotein analysis using a fluorescent affinity tag and mass spectrometry," *Rapid Commun Mass Spectrom*, 2005, 19(15):2157-62.
- Stimpson et al., "Real-time detection of DNA hybridization and melting on oligonucleotide arrays by using optical wave guides," *Proc Natl Acad Sci USA*, Jul. 1995, 92(14):6379-83.
- Stoddart et al., "Single-nucleotide discrimination in immobilized DNA oligonucleotides with a biological nanopore," *PNAS U S A*, May 2009, 106(19):7702-7707.

(56)

**References Cited****OTHER PUBLICATIONS**

- Stougaard et al., "In situ detection of non-polyadenylated RNA molecules using Turtle Probes and target primed rolling circle PRINS," *BMC Biotechnology*, Dec. 18, 2007, 7(69):1-10.
- Stroh et al., "Quantum dots spectrally distinguish multiple species within the tumor milieu *in vivo*," *Nat Med.*, Jun. 2005, 11(6):678-82.
- Subramanian et al., "Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles," *PNAS*, Oct. 2005, 102(43):15545-15550.
- Suh et al., "A simple soft lithographic route to fabrication of poly(ethylene glycol) microstructures for protein and cell patterning," *Biomaterials*, Feb. 2004, 25(3):557-563.
- Sumitomo et al., "Ca<sup>2+</sup> ion transport through channels formed by—hemolysin analyzed using a microwell array on a Si substrate," *Biosensors and Bioelectronics*, 2012, 31(1):445-450.
- Summersgill et al., "Fluorescence In Situ Hybridization Analysis of Formalin Fixed Paraffin Embedded Tissues, Including Tissue Microarrays," Chapter 4, Bridger, J. Ed., *Methods in Molecular Biology* 659, 2010, 51-70, 2010.
- Sun et al., "Direct immobilization of DNA probes on non-modified plastics by UV irradiation and integration in microfluidic devices for rapid bioassay," *Anal. Bio. Chem.*, 402: 741-748, 2012.
- Surzhik et al., "Template-dependent biosynthesis of poly(G)xpoly(C) and its antiviral activity *in vitro* and *in vivo*," *Antiviral Res.*, May 1988, 38(2):131-40.
- Sutherland et al., "Utility of formaldehyde cross-linking and mass spectrometry in the study of protein-protein interactions," *J. Mass Spectrom.*, Jun. 2008, 43(6):699-715.
- Swartz et al., "Interstitial flow and its effects in soft tissues," *Annu Rev Biomed Eng.*, 2007, 9:229-56.
- Syková et al., "Diffusion in brain extracellular space," *Physiol Rev.*, Oct. 2008, 88(4):1277-340.
- Tai et al., "Replication-competent retrovirus vectors for cancer gene therapy," *Front Biosci.*, Jan. 2008, 13:3083-95.
- Takahashi et al., "In Vitro Selection of Protein and Peptide Libraries Using mRNA Display," *Nucleic Acid and Peptide Aptamers: Methods and Protocols* (2009) 535:293-314 (Ch. 17).
- Tan et al., "Parylene peel-off arrays to probe the role of cell-cell interactions in tumour angiogenesis," *Integr Biol (Camb)*, Oct. 2009, 1(10):587-594.
- Tang et al., "mRNA-Seq whole-transcriptome analysis of a single cell," *Nat Methods*, 2009, 6:377-382.
- Tang et al., "RNA-Seq analysis to capture the transcriptome landscape of a single cell," *Nat Protoc.*, 5:516-35, 2010.
- Taniguchi et al., "Quantitative analysis of gene expression in a single cell by qPCR," *Nature Methods*, 6, pp. 503-506, 2009.
- Tawfik et al., "Man-made cell-like compartments for molecular evolution," *Nat Biotechnol.*, Jul. 1998, 16(7):652-6.
- Taylor et al., "Microfluidic local perfusion chambers for the visualization and manipulation of synapses," *Neuron*, Apr. 2010, 66(1):57-68, 25 pages.
- Taylor et al., "Mitochondrial DNA mutations in human disease," *Nature Reviews Genetics*. May 2005, 6(5):389-402.
- Tecott et al., "In Situ Transcription: Specific Synthesis of Complementary DNA in Fixed Tissue Sections," *Science*, 1988, 240:1661-1664.
- Tegtmeier et al., "Alternative Interactions of the SV40 A Protein with DNA," *Virology*, 1981, 115:75-87.
- Thacker et al., "Alkaline Hydrolysis—Carcass Disposal: A Comprehensive Review," National Agriculture Biosecurity Center, Aug. 2004, Chapter 6, pp. 1-12.
- Thiery et al., "Multiplex target protein imaging in tissue sections by mass spectrometry—TAMSIM," *Rapid Commun. Mass Spectrom.*, 2007, 21:823-829.
- Thorne et al., "In vivo diffusion analysis with quantum dots and dextrans predicts the width of brain extracellular space," *Proc Natl Acad Sci USA*, Apr. 2006, 103(14):5567-5572.
- Thornton, "High rate thick film growth." Annual review of materials science, Aug. 1977, 7(1):239-60.
- Tian et al., "Antigen peptide-based immunosensors for rapid detection of antibodies and antigens," *Anal Chem* May 26, 2009, 81 (13):5218-5225.
- Tijssen et al., "Overview of principles of hybridization and the strategy of nucleic acid assays" in *Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, 1993, 24(Chapter 2), 65 pages.
- Timofeev et al., "Regioselective immobilization of short oligonucleotides to acrylic copolymer gels," *Nucleic Acids Res.*, Aug. 1996, 24(16):3142-8.
- Tolbert et al., "New Methods for Proteomic Research: Preparation of Proteins with N-Terminal Cysteines for Labeling and Conjugation," *Angewandte Chemie International Edition*, Jun. 17, 2002, 41(12):2171-4.
- Totet et al., "Immunocompetent infants as a human reservoir for *Pneumocystis jirovecii*: rapid screening by non-invasive sampling and real-time PCR at the mitochondrial large subunit rRNA gene," *J Eukaryot Microbiol.*, 2003, pp. 668-669.
- Toubanaki et al., "Dry-reagent disposable biosensor for visual genotyping of single nucleotide polymorphisms by oligonucleotide ligation reaction: application to pharmacogenetic analysis," *Hum Mutat.*, Aug. 2008, 29(8):1071-8.
- Toy et al., "A Simple Plastic Perfusion Chamber for Continuous Maintenance and Cinematography of Tissue Cultures," *Experimental Cell Research*, 1958, 14:97-103.
- Trevino et al., "DNA Microarrays: a Powerful Genomic Tool for Biomedical and Clinical Research," *Mol Med*, 2007, 13(9-10):527-541.
- Twyman et al., "Techniques Patents for SNP Genotyping," *Pharmacogenomics*, Jan. 2003, 4(1):67-79.
- Tzanetakis et al., "The use of reverse transcriptase for efficient first- and second-strand cDNA synthesis from single- and double-stranded RNA templates," *J Virol Methods*, Mar. 2005, 24(1-2):73-7.
- Ueno et al., "cDNA Display: Rapid Stabilization of mRNA Display," *Antibody-Drug Conjugates, Methods in Molecular Biology*, Jan. 2012, pp. 113-135.
- Ulery et al., "Biomedical Applications of Biodegradable Polymers," *J Polym Sci B Polym Phys.*, Jun. 2011, 49(12):832-864.
- U.S. Appl. No. 60/416,118 Fan et al., Multiplex Nucleic Acid Analysis Using Archived or Fixed Samples, filed Oct. 3, 2002, 22 pages.
- Valencia et al., "mRNA-Display-Based Selections for Proteins with Desired Functions: A Protease-Substrate Case Study." *Biotechnology progress*, May 2008, 24(3):561-9.
- Valley et al., "Optoelectronic tweezers as a tool for parallel single-cell manipulation and stimulation," *IEEE Trans Biomed Circuits Syst*, Dec. 2009, 3(6):424-31.
- Van Gelder et al., "Amplified RNA synthesized from limited quantities of heterogeneous cDNA," *Proc. Natl. Acad. Sci. USA* 87, 1663-1667, 1990.
- Van Ness et al., "A versatile solid support system for oligodeoxynucleotide probe-based hybridization assays," *Nucleic Acids Res.* 19: 3345-3350, 1991.
- Vandenbroucke et al., "Quantification of splice variants using real-time PCR," *Nucleic Acids Research*, 2001, 29(13):e68, 7 pages.
- Vasiliskov et al., "Fabrication of microarray of gel-immobilized compounds on a chip by copolymerization," *Biotechniques*, Sep. 1999, 27(3):592-606.
- Velculescu et al., "Serial analysis of gene expression." *Science*, Oct. 20, 1995, 270(5235):484-7.
- Verma et al., "Modified Oligonucleotides: Synthesis and Strategy for Users," *Annual Review of Biochemistry*, 1998, 67(1):99-134.
- Vermesh et al., "High-density, multiplexed patterning of cells at single-cell resolution for tissue engineering and other applications," *Angew Chem Int Ed Engl*, Aug. 2011, 50(32):7378-7380.
- Villa et al., "Partial V(D)J Recombination Activity Leads to Omenn Syndrome," *Cell*, May 29, 1998, 93:885-896.
- Villemejane et al., "Physical methods of nucleic acid transfer: general concepts and applications," *British Journal of Pharmacology*, 2009, 157:207-219.

(56)

**References Cited****OTHER PUBLICATIONS**

- Vincent et al., "Helicase-dependent isothermal DNA amplification," *EMBO Rep.*, Aug. 2004, 5(8):795-800.
- Viollet et al., "T4 RNA ligase 2 truncated active site mutants: improved tools for RNA analysis," *BMC Biotechnol.*, Jul. 2011, 11:72, 14 pages.
- viralzone.com, "Coronaviridae," viralzone.com, available on or before May 21, 2017, retrieved on Mar. 20, 2025, retrieved from URL <<https://web.archive.org/web/20200525133654/https://viralzone.expasy.org/30/>>, 2 pages.
- Voelkerding et al., "Next-Generation Sequencing: From Basic Research to Diagnostics," *Clinical Chemistry*, 2009, 55(4):641-658.
- Vogelstein et al., "Digital PCR," *Proceedings of the National Academy of Sciences*, Aug. 3, 1999, 96:9236-9241.
- Wade et al., "Genome sequence, comparative analysis, and population genetics of the domestic horse," *Science*, 326: 865-7, 2009.
- Waichman et al., "Functional immobilization and patterning of proteins by an enzymatic transfer reaction." *Analytical chemistry*, Jan. 21, 2010, 82(4):1478-85.
- Walker et al., "Isothermal in vitro amplification of DNA by a restriction enzyme/DNA polymerase system," *Proc. Natl. Acad. Sci. USA*, 1992, 89:392-396.
- Walker et al., "Strand displacement amplification—an isothermal, in vitro DNA amplification technique." *Nucleic acids research*. Apr. 11, 1992, 1992, 20(7):1691-1696.
- Walker et al., Ed., "Chapter 1: Basic Techniques in Molecular Biology," Medical Biotechnology Handbook, Humana Press, Totowa, New Jersey, 2005, 19 pages.
- Wang "Preparation of DNA substrates for in vitro mismatch repair," *Mol. Biotechnol.*, 2000, 15:97-104.
- Wang et al., "Mutations in NEXN, a Z-disc gene, are associated with hypertrophic cardiomyopathy," *Am J Hum Genet.*, Nov. 2010, 87(5):687-93.
- Wang et al., "Paramagnetic microspheres with core-shell-ed structures," *Journal of Materials Science*, Apr. 2012, 47(16):5946-54.
- Wang et al., "Single cell analysis: the new frontier in 'omics,'" *Trends Biotechnol.*, 28: 281-90, 2010.
- Wang et al., "High-fidelity mRNA amplification for gene profiling." *Nature biotechnology*. Apr. 2000, 18(4):457-459.
- Wang, "RNA amplification for successful gene profiling analysis," *J Transl Med.*, Jul. 2005, 3:28, 11 pages.
- Watanabe et al., "Cellular networks involved in the influenza virus life cycle," *Cell Host & Microbe*, Jun. 2010, 7(6):427-39.
- Waxman et al., "De-regulation of common housekeeping genes in hepatocellular carcinoma," *BMC Genomics*, 2007, 1-9.
- Weichhart et al., "Functional selection of vaccine candidate peptides from *Staphylococcus aureus* whole-genome expression libraries in vitro," *Infection and Immunity*, 2003, 71 (8):4333-4641.
- Weinreich et al., "Evidence that the cis Preference of the Tn5 Transposase is Caused by Nonproductive Multimerization," *Genes and Development*, Oct. 1994, 8(19):2363-2374.
- Wheeler et al., "Microfluidic device for single-cell analysis," *Analytical Chemistry*, Jul. 2003, 75(14):3581-3586.
- Wiedmann et al., "Ligase chain reaction (LCR)—overview and applications," *PCR Methods Appl.*, Feb. 1994, 3(4):S51-64.
- Wikipedia.org [online]. "Random hexamer," Jan. 2012, Retrieved on Jan. 21, 2022, retrieved from URL <[https://en.wikipedia.org/w/index.php?title=Random\\_hexamer&oldid=47304223](https://en.wikipedia.org/w/index.php?title=Random_hexamer&oldid=47304223)>, 1 page.
- Williams et al., "Disc electrophoresis in polyacrylamide gels: extension to new conditions of pH and buffer," *Annals of the New York Academy of Sciences*, Dec. 1964, 121(2):373-381.
- Williams, "RAC reviews serious adverse event associated with AAV therapy trial," *Mol Ther.*, Dec. 2007, 15(12):2053-54.
- Willi-Monnerat et al., "Comprehensive spatiotemporal transcriptomic analyses of the ganglionic eminences demonstrate the uniqueness of its caudal subdivision," *Molecular and Cellular Neuroscience* 37(4):845-856, 2008.
- Wilson et al., "New transposon delivery plasmids for insertional mutagenesis in *Bacillus anthracis*," *J Microbiol Methods*, Dec. 2007, 71(3):332-5.
- Wolf et al., "Rapid hybridization kinetics of DNA attached to submicron latex particles", *Nucleic Acids Res.* 15: 2911-2926, 1987.
- Wolf et al., "tadA, an essential tRNA-specific adenosine deaminase from *Escherichia coli*," *EMBO J.*, Jul. 15, 2002, 21(14):3841-3851.
- Wong et al., "Direct Site-Selective Covalent Protein Immobilization Catalyzed by a Phosphopantetheinyl Transferase," *J. Am. Chem Soc.*, 2008, 130:12456-64.
- Woo et al., "A Comparison of cDNA, Oligonucleotide, and Affymetrix GeneChip Gene Expression Microarray Platforms," *Journal of Biomolecular Techniques*, 2004, 15(4), 276-284.
- Wood et al., "Single cell trapping and DNA damage analysis using microwell arrays," *PNAS*, Jun. 2010, 107(22):10008-10013.
- Worthington et al., "Cloning of random oligonucleotides to create single-insert plasmid libraries," *Analyt. Biochem*, 2001, 294:169-175.
- Wright et al., "Reusable, reversibly sealable parylene membranes for cell and protein patterning," *J Biomed Mater Res A*, May 2008, 85(2):530-538.
- Wu et al., "Detection DNA Point Mutation with Rolling-Circle Amplification Chip," *Bioinformatics & Biomed Eng Conference*, Piscataway, NJ, 2010, pp. 1-4.
- Wu et al., "Detection DNA Point Mutation with Rolling-Circle Amplification Chip," *Bioinformatics and Biomedical Engineering (ICBBE)*, 2010 4th International Conference on IEEE, Piscataway, NJ, USA, Jun. 18, 2010, 1-4 pages.
- Xiao et al., "Direct determination of haplotypes from single DNA molecules," *Nature Methods*, 2009, 6(3):199-201.
- Xie et al., "CryoFISH: Fluorescence In Situ Hybridization on Ultrathin Cryosections," *Fluorescence in situ Hybridization (FISH)*, Jul. 2010, pp. 221-230.
- Xu et al., "Engineering a nicking endonuclease N.AlwI by domain swapping," *Proc. Natl. Acad. Sci. USA*, 2001, 98:12990-12995.
- Yamamoto et al., "Generation of stable co-cultures of vascular cells in a honeycomb alginate scaffold," *Tissue Eng Part A*, Jan. 1, 2010, 16(1):299-308.
- Yan et al., "Decorin gene delivery inhibits cardiac fibrosis in spontaneously hypertensive rats by modulation of transforming growth factor-beta/Smad and p38 mitogen-activated protein kinase signaling pathways," *Hum Gene Ther.*, Oct. 2009, 20(10):1190-200.
- Yang et al., "Nucleoside alpha-Thiotriphosphates, Polymerases and the Exonuclease III Analysis of Oligonucleotides Containing Phosphorothioate Linkages," *Nucleic Acids Research*, Apr. 22, 2007, 35(9):3118-3127.
- Yao et al., "Influence of laser parameters on nanoparticle-induced membrane permeabilization," *Journal of Biomedical Optics*, 2009, 14(5):054034, 7 pages.
- Yasukawa et al., "Effects of organic solvents on the reverse transcription reaction catalyzed by reverse transcriptases from avian myeloblastosis virus and Moloney murine leukemia virus," *Biosci Biotechnol Biochem.*, 2010, 74(9):1925-30.
- Yeakley et al., "Profiling alternative splicing on fiber-optic arrays," *Nature Biotechnology*, Apr. 2002, 20(4):353-358.
- Yershov et al., "DNA analysis and diagnostics on oligonucleotide microchips," *Proc. Natl. Acad. Sci. USA*, May 1996, 93(10):4913-4918.
- Yet et al., "Cardiac-specific expression of heme oxygenase-1 protects against ischemia and reperfusion injury in transgenic mice," *Circ Res.*, Jul. 2001, 89(2):168-73.
- Yin et al., "Genetically encoded short peptide tag for versatile protein labeling by Sfp phosphopantetheinyl transferase," *PNAS*, 2005, 102(44):15815-20.
- Yonezawa et al., "DNA display for in vitro selection of diverse peptide libraries," *Nucleic Acids Research*, 2003, 31 (19):e118.
- Yusof et al., "Inkjet-like printing of single-cells," *Lab Chip*, Jul. 2011, 11(14):2447-2454.
- Zhang et al., "A novel mechanism of transposon-mediated gene activation," *PLoS Genet.*, Oct. 2009, 5(10):e1000689, 10 pages.
- Zhang et al., "A self-assembly pathway to aligned monodomain gels," *Nat Mater*, Jul. 2010, 9(7):594-601, 12 pages (Author Manuscript).

(56)

**References Cited****OTHER PUBLICATIONS**

- Zhang et al., "Binding-induced DNA assembly and its application to octomole detection of proteins," *Anal Chem* (2012) 84(2):877-884.
- Zhang et al., "Single-base mutational analysis of cancer and genetic diseases using membrane bound modified oligonucleotides," *Nucleic Acids Res.*, Jul. 1991, 19(14):3929-33.
- Zhang et al., "Stripping custom microRNA microarrays and the lessons learned about probe-slide interactions," *Anal Biochem*, Mar. 2009, 386(2):222-7.
- Zhao et al., "Ultrasensitive DNA detection using highly fluorescent bioconjugated nanoparticles," *Journal American Chemical Society*, 2003, 125:11474-11475.
- Zheng et al., Origins of human mitochondrial point mutations as DNA polymerase mediated errors. *Mutat. Res.* 599(1-2): 11-20, 2006.
- Zheng, "Spectroscopy-based quantitative fluorescence resonance energy transfer analysis," *Methods Mol Biol.*, 2006, 337:65-77.
- Zhou et al., "Analysis of the expression profile of Dickkopf-1 gene in human glioma and the associate with tumor malignancy," *Journal of Experimental & Clinical Cancer Research*, Oct. 28, 2010, 29(138):1-7.
- Zhou et al., "Genetically encoded short peptide tags for orthogonal protein labeling by Sfp and AcpS phosphopantetheinyl transferases," *ACS Chemical Biol.*, 2007 2 5 : 337-346.
- Zhu et al., "Engineering strand-specific DNA nicking enzymes from the type IIS restriction endonucleases BsAI, BsmBI, and BsmAI," *J. Mol. Biol.*, 2004, 337:573-583.
- Zhu et al., "Reverse Transcriptase Template Switching: A SMART Approach for Full-Length cDNA Library Construction," *BioTechniques*, 2001, 30(4): 892-897.
- Zieba et al., "Bright-field microscopy visualization of proteins and protein complexes by in situ proximity ligation with peroxidase detection," *Clin Chem*, Jan. 2010, 56(1):99-110.
- Zilberman et al., "Genome-wide analysis of DNA methylation patterns," *Development* (2007) 134:3959-3965.
- Zimmerman et al., "Chapter 13, Imaging of Cells and Tissue with Mass Spectrometry," *Methods in Cell Biology, Biophysical Tools for Biologists*, vol. Two: *In Vivo Techniques*, Jan. 2008, 89:361-390.
- Zuker, "Mfold web server for nucleic acid folding and hybridization prediction," *Nucleic Acids Res.*, Jul. 2003, 31(13):3406-15.
- Cellosaurus, "Cellosaurus Hs 742.Sk (CVCL\_0887)," cellosaurus.org, Apr. 4, 2012, retrieved on May 23, 2025, retrieved from URL <[https://www.cellosaurus.org/CVCL\\_0887](https://www.cellosaurus.org/CVCL_0887)>, 1 page.
- Cellosaurus, "Cellosaurus MDA-MB-231 (CVCL\_0062)," cellosaurus.org, Apr. 4, 2012, retrieved on May 23, 2025, retrieved from URL <[https://www.cellosaurus.org/CVCL\\_0062](https://www.cellosaurus.org/CVCL_0062)>, 13 pages.
- Litosh et al., "Improved nucleotide sensitivity and termination of 3'-OH unblocked reversible terminators by molecular tuning of 2-nitrobenzyl alkylated HOMedU triphosphates," *Nucleic Acids Research*, Jan. 11, 2011, 339(6):E39, 13 pages.
- Manning et al., "Benefits and pitfalls of secondary antibodies: why choosing the right secondary is of primary importance," *PLoS One*, 2012, 7(6):e38313, 11 pages.
- Pray, "Eukaryotic Genome Complexity," *Nature Education*, 2008, 1(1):96, pp. 1-4.
- Simon et al., "Immunohistochemical analysis of tissue microarrays," *Methods Mol Biol.*, 2010, 664:113-26.
- Webb et al., "Chapter 2: Epi-Fluorescence Microscopy," *Cell Imaging Techniques, Methods in Molecular Biology*, 2012, 931:29-59.

\* cited by examiner




FIG. 1






200

FIG. 2



**FIG. 4A****FIG. 4B****FIG. 4C**




FIG. 5

**1****SPATIALLY ENCODED BIOLOGICAL ASSAYS****CROSS-REFERENCE TO RELATED APPLICATIONS**

This application is a continuation of U.S. patent application Ser. No. 18/972,148, filed Dec. 6, 2024, which is a continuation of U.S. patent application Ser. No. 18/793,359, filed on Aug. 2, 2024, which is a continuation of U.S. patent application Ser. No. 18/100,127, filed on Jan. 23, 2023, which is a continuation of U.S. patent application Ser. No. 17/878,519, filed on Aug. 1, 2022, now U.S. Pat. No. 11,560,587, which is a continuation of U.S. patent application Ser. No. 17/556,588, filed Dec. 20, 2021, now U.S. Pat. No. 11,401,545, which is a continuation of U.S. patent application Ser. No. 17/223,669, filed Apr. 6, 2021, now U.S. Pat. No. 11,208,684, which is a continuation of U.S. patent application Ser. No. 17/030,230, filed Sep. 23, 2020, now U.S. Pat. No. 11,384,386, which is a continuation of U.S. patent application Ser. No. 16/988,284, filed Aug. 7, 2020, now U.S. Pat. No. 10,961,566, which is a continuation of U.S. patent application Ser. No. 16/414,213, filed May 16, 2019, now U.S. Pat. No. 10,787,701, which is a continuation of U.S. patent application Ser. No. 16/402,098, filed May 2, 2019, now U.S. Pat. No. 10,472,669, which is a continuation of U.S. patent application Ser. No. 16/276,235, filed Feb. 14, 2019, now U.S. Pat. No. 10,480,022, which is a continuation application of U.S. patent application Ser. No. 15/187,661, filed Jun. 20, 2016, now U.S. Pat. No. 10,308,982, which is a continuation of U.S. patent application Ser. No. 13/080,616, filed Apr. 5, 2011, now U.S. Pat. No. 9,371,598, which claims the benefit of U.S. Provisional Patent Application No. 61/321,124, filed Apr. 5, 2010, each of which are herein incorporated by reference.

**FIELD OF THE INVENTION**

This invention relates to assays of biological molecules, and more particularly to assays for determining spatial distributions of a large number of biological molecules in a solid sample simultaneously.

**BACKGROUND OF THE INVENTION**

In the following discussion certain articles and methods will be described for background and introductory purposes. Nothing contained herein is to be construed as an "admission" of prior art. Applicant expressly reserves the right to demonstrate, where appropriate, that the articles and methods referenced herein do not constitute prior art under the applicable statutory provisions.

Comprehensive gene expression analysis and protein analysis have been useful tools in understanding mechanisms of biology. Use of these tools has allowed the identification of genes and proteins involved in development and in various diseases such as cancer and autoimmune disease. Conventional methods such as *in situ* hybridization and other multiplexed detection of different transcripts have revealed spatial patterns of gene expression and have helped shed light on the molecular basis of development and disease. Other technologies that have enabled the quantitative analysis of many RNA sequences per sample include microarrays (see Shi, et al., *Nature Biotechnology*, 24(9): 1151-61 (2006); and Slonim and Yanai, *Plos Computational Biology*, 5(10):e1000543 (2009)); serial analysis of gene expression (SAGE) (see Velculescu, et al., *Science*, 270

**2**

(5235):484-87 (1995)), high-throughput implementations of qPCR (see Spurgeon, et al., *Plos ONE*, 3(2):e1662 (2008)) and *in situ* PCR (see Nuovo, *Genome Res.*, 4:151-67 (1995)). As useful as these methods are, however, they do not enable simultaneous measurement of the expression of many genes or the presence and/or activity of multiple proteins at many spatial locations in a sample. Laser capture microdissection has permitted the analysis of many genes at a small number of locations, but it is very expensive, 10 laborious, and does not scale well. Certain PCR assays in a 2D format preserve spatial information (see Armani, et al., *Lab on a Chip*, 9(24):3526-34 (2009)), but these methods have low spatial resolution because they rely on physical transference of tissue into wells, which also prevents random access to tissue samples and high levels of multiplexing.

At present, no practical method exists to analyze at high resolution the spatial expression patterns of large numbers of genes, proteins, or other biologically active molecules 20 simultaneously. There is thus a need for reproducible, high-resolution spatial maps of biological molecules in tissues. The present invention addresses this need.

**SUMMARY OF THE INVENTION**

This Summary is provided to introduce a selection of concepts in a simplified form that are further described below in the Detailed Description. This Summary is not intended to identify key or essential features of the claimed subject matter, nor is it intended to be used to limit the scope 30 of the claimed subject matter. Other features, details, utilities, and advantages of the claimed subject matter will be apparent from the following written Detailed Description including those aspects illustrated in the accompanying drawings and defined in the appended claims.

The invention encompasses assay systems that provide high-resolution spatial maps of biological activity in tissues. The assay system comprises an assay capable of high levels of multiplexing where encoded probes are provided to a 40 biological sample in defined spatial patterns; instrumentation capable of controlled delivery of reagents according to the spatial patterns; and a decoding scheme providing a readout that is digital in nature. In short, the present invention provides the ability to look at many biological targets in many locations, providing the resolution of *in situ* hybridization with the highly-parallel data analysis of sequencing.

Thus, in some embodiments, the invention provides an assay system to determine spatial patterns of abundance or activity or both of multiple biological targets at multiple sites in a sample, where the assay system performs the following steps: providing a sample affixed to a support; delivering encoded probes for the multiple biological targets to the multiple sites in the sample in a known spatial pattern, where each encoded probe comprises a probe region that 50 may interact with the biological targets and a coding tag that identifies a location of the site to which the encoded probe was delivered; allowing the encoded probes to interact with the biological targets; separating encoded probes that interact with the biological targets from encoded probes that do not interact with the biological targets; determining all or a portion of a sequence of the encoded probes, and associating the abundance or activity or both of the multiple biological targets to the locations of the sites in the sample.

In particular aspects of the invention the biological targets 55 comprise nucleic acids and the encoded probes are oligonucleotides, and in some aspects, there are two encoded probes for each of the multiple nucleic acid targets. In some

aspects, the multiple biological targets comprise proteins, the probe regions of the encoding probes are proteins and the coding tags comprise oligonucleotides. In some aspects the multiple biological targets comprise enzymes. In some aspects the probe regions of the encoded probes comprise antibodies, aptamers or small molecules.

Some aspects of the assay system further comprise an amplification step between the separating step and the determining step. In some aspects, the determining step is performed by nucleic acid sequencing, and in preferred aspects, the sequencing is high-throughput digital nucleic acid sequencing.

In some aspects of the invention, the product of the multiple biological targets being assayed and the multiple sites in the sample is greater than 20, in some aspects product of the multiple biological targets being assayed and the multiple sites in the sample is greater than 50, in some aspects the product of the multiple biological targets being assayed and the multiple sites in the sample is greater than 75, 100, 150, 500, 750, 1,000, 5,000, 10,000, 25,000, 50,000, 100,000, 500,000, or 1,000,000 or more. In other aspects, the sequence of at least fifty thousand encoding probes are determined in parallel, in other aspects the sequence of at least one hundred thousand encoding probes are determined in parallel, in some aspects the sequence of at least five hundred thousand encoding probes are determined in parallel, and in some aspects the sequence of at least one million, ten million, one hundred million, one billion, ten billion, one hundred billion or more encoding probes are determined in parallel.

In some aspects, the known spatial pattern is determined by histological features of the sample. Also in some aspects, software programmed hardware performs at least two steps of the delivering step, the separation step, the determining step and the associating step.

In some aspects, the probe regions of the encoded probes are proteins and the separating step is accomplished by encoded probes that interact with the biological targets being captured by an affinity capture agent. In some aspects the probe regions of the encoding probes are nucleic acids and the separating step is accomplished by a washing of the sample.

In other embodiments there is provided an assay system to determine spatial patterns of abundance or activity or both of multiple nucleic acid targets at multiple sites in a sample, where the assay system performs the following steps: providing a sample affixed to a support; delivering oligonucleotide probes for multiple nucleic acid targets to the multiple sites in the sample in a known spatial pattern; allowing the oligonucleotide probes to hybridize with the nucleic acid targets; washing unhybridized encoded oligonucleotide probes from the sample; delivering one or more encoding agents to locations of the multiple sites in the sample according to a known spatial pattern, where the combination of encoding agents delivered to each site is different; coupling the encoding agents and the oligonucleotide probes to form encoded probes; determining all or a portion of a sequence of the encoded probes using high-throughput sequencing, and associating the abundance or activity or both of multiple biological targets to the locations of multiple sites in the sample.

Other embodiments of the invention provide an assay system to determine spatial patterns of abundance or activity or both of multiple protein targets at multiple sites in a sample, where the assay system performs the following steps: providing a sample affixed to a support; delivering encoded probes for the multiple protein targets to the mul-

multiple sites in the sample in a known spatial pattern, where each encoded probe comprises a protein probe region that may interact with the protein targets and a coding tag that identifies a location of the site to which the encoded probe was delivered and the protein probe region of the encoding probe of which the coding tag is part; allowing the encoded probes to interact with the protein targets; separating encoded probes that interact with the protein targets from encoded probes that do not interact with the protein targets; determining all or a portion of a sequence of the encoded probes by high throughput sequencing, and associating the abundance or activity or both of the multiple protein targets to the locations of the multiple sites in the sample.

Other embodiments provide an assay system to determine spatial patterns of abundance or activity or both of multiple biological targets at multiple sites in a sample, where the assay system performs the following steps: providing a sample affixed to a support; delivering encoded probes for the multiple biological targets to the multiple sites in the sample in a known spatial pattern, where each encoded probe comprises a probe region that may interact with the biological targets and a coding tag that identifies a location of the site to which the encoded probe was delivered and identifies the biological target; allowing the encoded probes to interact with the biological targets; determining all or a portion of a sequence of the encoded probes, and associating the abundance or activity or both of the multiple biological targets to the locations of the sites in the sample.

The assay system of the invention can utilize various detection mechanisms, based on the molecules to be detected and the reagents needed for such detection system. Exemplary methods that can be used with the assay systems of the invention are described in more detail below.

## DESCRIPTION OF THE FIGURES

FIG. 1 provides a simplified overview of the assay system of the present invention.

FIG. 2 provides a simplified overview of one embodiment of the assay system of the present invention for detecting nucleic acids.

FIG. 3 is a representational depiction of one embodiment of the assay overviewed in FIG. 2.

FIG. 4A-C illustrates a general mechanism for one embodiment of a combinatorial encoding scheme of the assay systems of the invention. FIG. 4A shows two target-specific/encoding oligonucleotide constructs specifically bound to a target nucleic acid of interest in a sample. FIG. 4B shows a scheme for delivering twenty different coding tags, a1 through a10 and b1 through b10, to a sample to form a 10×10 coding tag grid. FIG. 4C shows a tissue section sample to which the coding tags are delivered, forming the coding tag grid in the sample.

FIG. 5 provides a simplified, specific example of the embodiment of a combinatorial encoding scheme shown in FIG. 4.

## DEFINITIONS

The terms used herein are intended to have the plain and ordinary meaning as understood by those of ordinary skill in the art. The following definitions are intended to aid the reader in understanding the present invention, but are not intended to vary or otherwise limit the meaning of such terms unless specifically indicated.

The term “antibody” as used herein is intended to refer to an entire immunoglobulin or antibody or any functional

fragment of an immunoglobulin molecule which is capable of specific binding to an antigen (antibodies and antigens are "binding partners" as defined herein). "Antibody" as used herein is meant to include the entire antibody as well as any antibody fragments capable of binding the antigen or antigenic fragment of interest. Examples of such peptides include complete antibody molecules, antibody fragments, such as Fab, F(ab')<sub>2</sub>, CDRS, VL, VH, and any other portion of an antibody which is capable of specifically binding to an antigen. Antibodies for assays of the invention are immunoactive or immunospecific for, and therefore specifically and selectively bind to, proteins either detected (i.e., biological targets) or used for detection (i.e., probes) in the assays of the invention.

The term "binding agent" as used herein refers to any agent that specifically binds to a biological molecule of interest

"Complementary" or "substantially complementary" refers to the hybridization or base pairing or the formation of a duplex between nucleotides or nucleic acids, such as, for instance, between the two strands of a double-stranded DNA molecule or between an oligonucleotide primer and a primer binding site on a single-stranded nucleic acid. Complementary nucleotides are, generally, A and T (or A and U), or C and G. Two single-stranded RNA or DNA molecules are said to be substantially complementary when the nucleotides of one strand, optimally aligned and compared and with appropriate nucleotide insertions or deletions, pair with at least about 80% of the other strand, usually at least about 90% to about 95%, and even about 98% to about 100%.

"Hybridization" refers to the process in which two single-stranded polynucleotides bind non-covalently to form a stable double-stranded polynucleotide. The resulting (usually) double-stranded polynucleotide is a "hybrid" or "duplex." "Hybridization conditions" will typically include salt concentrations of approximately less than 1M, often less than about 500 mM and may be less than about 200 mM. A "hybridization buffer" is a buffered salt solution such as 5% SSPE, or other such buffers known in the art. Hybridization temperatures can be as low as 5° C., but are typically greater than 22° C., and more typically greater than about 30° C., and typically in excess of 37° C. Hybridizations are often performed under stringent conditions, i.e., conditions under which a primer will hybridize to its target subsequence but will not hybridize to the other, non-complementary sequences. Stringent conditions are sequence-dependent and are different in different circumstances. For example, longer fragments may require higher hybridization temperatures for specific hybridization than short fragments. As other factors may affect the stringency of hybridization, including base composition and length of the complementary strands, presence of organic solvents, and the extent of base mismatching, the combination of parameters is more important than the absolute measure of any one parameter alone. Generally stringent conditions are selected to be about 5° C. lower than the  $T_m$  for the specific sequence at a defined ionic strength and pH. Exemplary stringent conditions include a salt concentration of at least 0.01 M to no more than 1M sodium ion concentration (or other salt) at a pH of about 7.0 to about 8.3 and a temperature of at least 25° C. For example, conditions of 5xSSPE (750 mM NaCl, 50 mM sodium phosphate, 5 mM EDTA at pH 7.4) and a temperature of approximately 30° C. are suitable for allele-specific hybridizations, though a suitable temperature depends on the length and/or GC content of the region hybridized.

"Ligation" means to form a covalent bond or linkage between the termini of two or more nucleic acids, e.g.,

oligonucleotides and/or polynucleotides, in a template-driven reaction. The nature of the bond or linkage may vary widely and the ligation may be carried out enzymatically or chemically. As used herein, ligations are usually carried out enzymatically to form a phosphodiester linkage between a 5' carbon terminal nucleotide of one oligonucleotide with a 3' carbon of another nucleotide.

"Nucleic acid", "oligonucleotide", "oligo" or grammatical equivalents used herein refers generally to at least two nucleotides covalently linked together. A nucleic acid generally will contain phosphodiester bonds, although in some cases nucleic acid analogs may be included that have alternative backbones such as phosphoramidite, phosphordithioate, or methylphosphoroamidite linkages; or peptide nucleic acid backbones and linkages. Other analog nucleic acids include those with bicyclic structures including locked nucleic acids, positive backbones, non-ionic backbones and non-ribose backbones. Modifications of the ribose-phosphate backbone may be done to increase the stability of the molecules; for example, PNA:DNA hybrids can exhibit higher stability in some environments.

"Primer" means an oligonucleotide, either natural or synthetic, that is capable, upon forming a duplex with a polynucleotide template, of acting as a point of initiation of nucleic acid synthesis and being extended from its 3' end along the template so that an extended duplex is formed. The sequence of nucleotides added during the extension process is determined by the sequence of the template polynucleotide. Primers usually are extended by a DNA polymerase.

The term "SNP" or "single nucleotide polymorphism" refers to a genetic variation between individuals; e.g., a single nitrogenous base position in the DNA of organisms that is variable. SNPs are found across the genome; much of the genetic variation between individuals is due to variation at SNP loci, and often this genetic variation results in phenotypic variation between individuals. SNPs for use in the present invention and their respective alleles may be derived from any number of sources, such as public databases (U.C. Santa Cruz Human Genome Browser Gateway or the NCBI dbSNP website, or may be experimentally determined as described in U.S. Pat No. 6,969,589; and US Pub. No. 2006/0188875 entitled "Human Genomic Polymorphisms." Although the use of SNPs is described in some of the embodiments presented herein, it will be understood that other biallelic or multi-allelic genetic markers may also be used. A biallelic genetic marker is one that has two polymorphic forms, or alleles. As mentioned above, for a biallelic genetic marker that is associated with a trait, the allele that is more abundant in the genetic composition of a case group as compared to a control group is termed the "associated allele," and the other allele may be referred to as the "unassociated allele." Thus, for each biallelic polymorphism that is associated with a given trait (e.g., a disease or drug response), there is a corresponding associated allele. Other biallelic polymorphisms that may be used with the methods presented herein include, but are not limited to multinucleotide changes, insertions, deletions, and translocations. It will be further appreciated that references to DNA herein may include genomic DNA, mitochondrial DNA, episomal DNA, and/or derivatives of DNA such as amplicons, RNA transcripts, cDNA, DNA analogs, etc. The polymorphic loci that are screened in an association study may be in a diploid or a haploid state and, ideally, would be from sites across the genome.

The term "selectively binds", "selective binding" and the like as used herein, when referring to a binding partner (e.g. protein, nucleic acid, antibody or other affinity capture

agent, etc.), refers to a binding reaction of two or more binding partners with high affinity and/or complementarity to ensure selective hybridization under designated assay conditions. Typically, specific binding will be at least three times the standard deviation of the background signal. Thus, under designated conditions the binding partner binds to its particular “target” molecule and does not bind in a significant amount to other molecules present in the sample.

“Sequencing”, “sequence determination” and the like means determination of information relating to the nucleotide base sequence of a nucleic acid. Such information may include the identification or determination of partial as well as full sequence information of the nucleic acid. Sequence information may be determined “with varying degrees of statistical reliability or confidence. In one aspect, the term includes the determination of the identity and ordering of a plurality of contiguous nucleotides in a nucleic acid, “High throughput digital sequencing” or “next generation sequencing” means sequence determination using methods that determine many (typically thousands to billions) of nucleic acid sequences in an intrinsically parallel manner, i.e. where DNA templates are prepared for sequencing not one at a time, but in a bulk process, and where many sequences are read out preferably in parallel, or alternatively using an ultra-high throughput serial process that itself may be parallelized. Such methods include but are not limited to pyrosequencing (for example, as commercialized by 454 Life Sciences, Inc., Branford, CT); sequencing by ligation (for example, as commercialized in the SOLiD™ technology, Life Technology, Inc., Carlsbad, CA); sequencing by synthesis using modified nucleotides (such as commercialized in TruSeq™ and HiSeq™ technology by Illumina, Inc., San Diego, CA, HeliScope™ by Helicos Biosciences Corporation, Cambridge, MA, and PacBio RS by Pacific Biosciences of California, Inc., Menlo Park, CA), sequencing by ion detection technologies (Ion Torrent, Inc., South San Francisco, CA); sequencing of DNA nanoballs (Complete Genomics, Inc., Mountain View, CA); nanopore-based sequencing technologies (for example, as developed by Oxford Nanopore Technologies, LTD, Oxford, UK), and like highly parallelized sequencing methods.

The term “ $T_m$ ” is used in reference to the “melting temperature.” The melting temperature is the temperature at which a population of double-stranded nucleic acid molecules becomes half dissociated into single strands. Several equations for calculating the  $T_m$  of nucleic acids are well known in the art. As indicated by standard references, a simple estimate of the  $T_m$  value may be calculated by the equation,  $T_m = 81.5 + 0.41 (\% G+C)$ , when a nucleic acid is in aqueous solution at 1M NaCl (see e.g., Anderson and Young, Quantitative Filter Hybridization, in *Nucleic Acid Hybridization* (1985)). Other references (e.g., Allawi and SantaLucia, Jr., Biochemistry, 36:10581-94 (1997)) include alternative methods of computation which take structural and environmental, as well as sequence characteristics into account for the calculation of  $T_m$ .

#### DETAILED DESCRIPTION OF THE INVENTION

The practice of the techniques described herein may employ, unless otherwise indicated, conventional techniques and descriptions of organic chemistry, polymer technology, molecular biology (including recombinant techniques), cell biology, biochemistry, and sequencing technology, which are within the skill of those who practice in the art. Such conventional techniques include polymer array synthesis,

hybridization and ligation of polynucleotides, and detection of hybridization using a label. Specific illustrations of suitable techniques can be had by reference to the examples herein. However, other equivalent conventional procedures can, of course, also be used. Such conventional techniques and descriptions can be found in standard laboratory manuals such as Green, et al., Eds., *Genome Analysis: A Laboratory Manual Series* (Vols. I-IV) (1999); Weiner, Gabriel, Stephens, Eds., *Genetic Variation: A Laboratory Manual* (2007); Dieffenbach, Dveksler, Eds., *PCR Primer: A Laboratory Manual* (2003); Bowtell and Sambrook, *DNA Microarrays: A Molecular Cloning Manual* (2003); Mount, *Bioinformatics: Sequence and Genome Analysis* (2004); Sambrook and Russell, *Condensed Protocols from Molecular Cloning: A Laboratory Manual* (2006); and Sambrook and Russell, *Molecular Cloning: A Laboratory Manual* (2002) (all from Cold Spring Harbor Laboratory Press); Stryer, *Biochemistry* (4th Ed.) (1995) W.H. Freeman, New York N.Y.; Gait, “*Oligonucleotide Synthesis: A Practical Approach*” (2002) IRL Press, London; Nelson and Cox, Lehninger, *Principles of Biochemistry* (2000) 3<sup>rd</sup> Ed., W. H. Freeman Pub., New York, N.Y.; and Berg, et al., *Biochemistry* (2002) 5<sup>th</sup> Ed., W. H. Freeman Pub., New York, N.Y., all of which are herein incorporated in their entirety by reference for all purposes.

Note that as used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a nucleic acid” refers to one or more nucleic acids, and reference to “the assay” includes reference to equivalent steps and methods known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. All publications mentioned herein are incorporated by reference for the purpose of describing and disclosing devices, formulations and methodologies that may be used in connection with the presently described invention.

Where a range of values is provided, it is understood that each intervening value, between the upper and lower limit of that range and any other stated or intervening value in that stated range is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either both of those included limits are also included in the invention.

In the following description, numerous specific details are set forth to provide a more thorough understanding of the present invention. However, it will be apparent to one of skill in the art that the present invention may be practiced without one or more of these specific details. In other instances, well-known features and procedures well known to those skilled in the art have not been described in order to avoid obscuring the invention.

#### The Invention in General

The assay systems of the invention provide spatially-encoded, multiplexed assays comprising 1) an assay capable of high levels of multiplexing with an efficient spatial encoding scheme; 2) instrumentation capable of delivering reagents according to a spatial pattern; and 3) decoding determined by a readout that is digital in nature. The assay

systems of the invention detect the presence or absence and relative amount of a biological target or biological activity indicative of a biological target, as well as the location of the biological target or activity in a biological sample, e.g., a tissue section or other biological structure disposed upon a support such as a microscope slide or culture dish.

The assay system further provides instrumentation with an ability to deliver reagents in a spatially-defined pattern. This instrumentation, together "with software, reagents and protocols, provides a key component of the highly innovative assay system of the invention, allowing for measurement of numerous biological targets or activities in a meaningful spatial environment, including gene expression and peptide localization. An encoding scheme used in these assay systems allows one to determine the location of biological targets or activity (or lack thereof) in the biological samples after the products of the multiplexed assay are removed from the biological sample and pooled for analysis. Decoding of the encoding scheme can be performed by, e.g., next-generation sequencing, which easily provides millions to trillions of data points at low cost. The assay results such as the amount or activity of biological targets can then be mapped back to specific location in the biological sample. The assay systems open a new analytical window into the complex spatial patterns of cellular function and regulation in biological samples.

A simplified overview of the assay system 100 of the present invention is provided at FIG. 1. At step 110, a biological sample affixed to a support is provided. The biological sample contains biological targets of interest. Biological targets can include any molecule of interest, such as nucleic acids (including, e.g., RNA transcripts, genomic DNA sequences, cDNAs, amplicons, or other nucleic acid sequences) and proteins, enzymes and the like. At step 120, encoded probes are delivered to the biological sample according to a known spatial pattern. Encoded probes comprise probes, which can interact "with biological targets of interest, and coding tags, which identify the positions in the sample of the biological targets being assayed, and thus can be used to link assay results back to locations in the sample. Coding tags in most embodiments are oligonucleotides. However, coding tags may also be mass tags, fluorescent labels, or other moieties.

In some embodiments, the probe and coding tag portions of the encoded probe are pre-coupled before being delivered to the biological sample. For example, in the case where the encoded probes are oligonucleotides, both the probe and coding tag sequence can be synthesized as a single oligonucleotide. Alternatively, the probe and coding tag portions of the encoding probes can be synthesized or obtained separately and combined before delivery to the biological sample (e.g., two separate oligonucleotides can be synthesized and coupled by, e.g., ligation; or an antibody and an oligonucleotide can be prepared separately and conjugated before delivery to the biological sample). Also, as is described in FIGS. 2-5, the probes and the coding tags (in encoding oligonucleotides) are synthesized separately, and are delivered to the biological sample at different steps (e.g., probes first and coding tags thereafter, or vice versa) in the assay.

At step 130, the encoded probes are allowed to react or interact with the biological targets, i.e., conditions are provided to allow e.g., oligonucleotides to hybridize to nucleic acid targets, enzymes to catalyze reactions with protein targets, antibodies to bind epitopes, etc. In the case where the biological targets are nucleic acids, the encoded probes are typically oligonucleotides and hybridize to the target nucleic

acids. In the case that the biological targets are proteins, the encoded probes typically are aptamers, small molecules, or oligonucleotide-conjugated proteins that interact with target proteins by binding to them or by reacting with them (that is, one of the proteins is a substrate for the other). Encoding oligonucleotides may be coupled to the probes (proteins) by conjugation, chemical or photo-crosslinking via suitable groups and the like.

Once encoded probes interact with the biological targets, 10 the encoded probes that interacted with the biological targets must be separated from the encoded probes that did not interact with the biological targets at step 140. In the case where the biological targets are nucleic acids and the encoded probes are oligonucleotides, the separation can be 15 accomplished by, e.g., washing the unhybridized encoded probes from the sample. Similarly, for other assays that are based on affinity binding, including those using aptamer, small molecule, and protein probes, washing steps can be used to remove low affinity binders. In the case where the 20 probe is transformed via interaction with the target, e.g., in the case of a peptide, e.g., via cleavage by a protease or phosphorylation by a kinase, it is convenient to collect, all encoded probes—both encoded probes that interacted with the biological targets and were transformed and encoded 25 probes that were not transformed. After collection or pooling, an antibody or other affinity capture agent can be used to capture probes that were transformed by addition of a moiety (e.g., a phosphate group). In cases where probes have been transformed via cleavage, the transformed probes can 30 be separated, e.g., by capturing the non-transformed probes via a tag that is removed from the transformed probes during the transformation (e.g., by cleavage), or by adding a new tag at the site of cleavage.

Once the reacted (transformed) or interacted encoded 35 probes are separated from the unreacted or un-interacted encoded probes, the sequence of the reacted and/or interacted encoded probes is determined at step 150 by, preferably, sequencing. The sequence of the encoded probes allows the mapping of the assay results at step 160 back to 40 locations in the biological sample.

FIG. 2 provides a simplified overview of an assay system 200 of the present invention embodying an efficient implementation of a combinatorial coding scheme for the encoding of spatial information. For purposes of this overview, the 45 probes are oligonucleotides, but as explained elsewhere, other types of probes can also be used. In step 210, a biological sample affixed to a support, e.g., a tissue sample or other biological structure, is provided. In step 220, one or more oligonucleotide probes are delivered to the biological 50 sample, where the oligonucleotide probes are capable of hybridizing with biological targets in the biological sample. In step 230, the oligonucleotide probes are allowed to interact with (hybridize to) the nucleic acid targets; that is, appropriate conditions are provided where oligonucleotide 55 probes can hybridize to the target nucleic acids.

In step 240, the oligonucleotide probes that did not hybridize to target nucleic acids are removed, and thereby 60 separated from oligonucleotide probes that did hybridize to target nucleic acids. In this embodiment, separation can be accomplished by, e.g., washing the sample to remove unhybridized oligonucleotide probes. Next, in step 250, encoding 65 oligonucleotides (the encoding agents) are delivered to the biological sample according to a chosen spatial pattern, where the encoding oligonucleotides comprise coding tags that are used to encode the location of biological targets in the biological sample. Note that in contrast to the assay system of FIG. 1, here the probes and encoding agents

(encoding oligonucleotides) are delivered in separate steps. In step 260, the encoding oligonucleotides are coupled to the oligonucleotide probes to create encoded probes. In this case where the probes are oligonucleotides, the encoding oligonucleotides may be coupled to the oligonucleotides probes by, e.g., ligation. Alternatively, the information in the encoding oligonucleotides can be transferred by using a DNA polymerase to extend a probe oligonucleotide that acts as a primer, and thereby copy and incorporate the sequence of the encoding oligonucleotides.

In step 270, the sequence of the coding tags in the encoded probes as well as the sequence or a portion of the sequence of the probe itself is determined, and in step 280, the target nucleic acids are mapped back to the biological sample. In some embodiments, the abundance of sequences reveals the relative quantity of biological targets at the location. Although this embodiment shows the individual steps in a particular order, so as to better explain the invention, the precise order of the steps can be varied. For example, steps 220 and 250 can be combined, so that a mixture of the probes and encoding oligonucleotides is delivered according to a chosen spatial pattern. Coupling step 260 can then be carried out immediately after the combined steps 220 and 250, or concomitantly with them. In this case, step 240 would then occur after step 260. It can therefore be appreciated that the two key results of this series of steps, i.e., the location-specific encoding of probe molecules and the separation of probe molecules based on their ability to interact with corresponding target molecules, can be accomplished with some flexibility in the implementation of the particular steps. Similarly, there is considerable flexibility in the design of the coding scheme. As described infra, the assays of the invention are particularly amenable to combinatorial methods.

Thus, the present invention provides an ability to look at many different biological targets in many locations, providing the resolution of *in situ* hybridization with the highly-parallel data analysis of sequencing. In some embodiments, the sum of the multiple biological targets being assayed and the multiple sites in the biological sample is greater than 20, in other embodiments, the sum of the multiple biological targets being assayed and the multiple sites in the biological sample is greater than 50, in other embodiments, the sum of the multiple biological targets being assayed and the multiple sites in the biological sample is greater than 100, greater than 500, 1,000, 10,000, 25,000, 100,000, 500,000, 1,000,000. It will be appreciated that, due to the spatial encoding dimension of the invention, even much larger numbers can be contemplated. For example, assaying 10,000 targets per location×10,000 locations would generate 10<sup>8</sup> different assays, and even larger numbers than these can easily be contemplated, particularly if spatial locations with resolution on the order of that of single cells are utilized. Further, in embodiments where high-throughput digital sequencing is employed, the sequences of at least 1,000 encoding probes are typically determined in parallel. More typically, using a digital readout, it is desirable to obtain multiple sequence reads for each assay (defined by a probe and a spatial location code). It is desirable to obtain an average of at least 3 copies per assay, and more typically at least 10 or at least 30 copies per assay, depending on the design of the experiment and requirements of the assay. For a quantitative readout with suitable dynamic range, it may be desirable to obtain at least 1,000 reads per assay. Therefore, if 1,000,000 assays are carried out, the number of sequence reads may be 1 billion or more. With high-throughput digital sequencing, and allowing for redundancy, the sequence of at

least 10,000 encoding probes are determined in parallel, or the sequence of at least 100,000, 500,000, 1,000,000, 10,000,000, 100,000,000, 1,000,000,000 or more encoding probes are determined in parallel.

5

### Assays

The assay portion of the assay systems of the present invention comprise the following general steps: delivering 10 probes and encoding agents where the encoding agents (in some embodiments pre-coupled to the probes) are delivered to the sample according to a known spatial pattern, allowing the probes to interact or react with biological targets in the sample, and, if the probes and encoding agents have not been 15 pre-coupled, coupling the encoding agents to probes.

The samples of the present invention include virtually any 20 biological sample or samples that can be affixed to a support or provided essentially in a two-dimensional manner, where the ability to tie an assayed biological target or activity back 25 to the location within the biological sample is important. Exemplary biological samples include tissue sections (e.g., including whole animal sectioning and tissue biopsies), cell populations on slides or culture dishes, and the like. The assay systems of the invention are particularly advantageous 30 in that they are compatible with numerous biological sample types, including fresh samples, such as primary tissue sections, and preserved samples including but not limited to frozen samples and paraformalin-fixed, paraffin-embedded (FFPE) samples. An important aspect of the assay systems 35 of the invention is that the biological samples are immobilized on a substrate surface having discrete, independently measurable areas.

The biological targets to be detected can be any biological 40 molecules including but not limited to proteins, nucleic acids, lipids, carbohydrates, ions, or multicomponent complexes containing any of the above. Examples of subcellular targets include organelles, e.g., mitochondria, Golgi apparatus, endoplasmic reticulum, chloroplasts, endocytic vesicles, exocytic vesicles, vacuoles, lysosomes, etc.

In some particular embodiments, the assay system is used 45 to analyze nucleic acids, e.g., by genotyping, quantitation of DNA copy number or RNA transcripts, localization of particular transcripts within samples, and the like. FIG. 3 illustrates an overall scheme for an exemplary assay for, e.g., detecting single nucleotide polymorphisms (SNPs) that can be used with the assay system of the invention. In FIG. 3, two oligonucleotide probes are provided. Each oligonucleotide probe comprises a target-specific region (located on either side of the SNP to be analyzed) seen at 305 and 307, and ligation regions, seen at 301 and 303. The oligonucleotide probes are allowed to hybridize to a target nucleic acid 50 (not shown) in the biological sample. At step 302, one of the oligonucleotide probes is extended to incorporate the SNP sequence and ligated to the other probe to form an extended 55 probe comprising target nucleic acid region 309 and ligation regions 301 and 303.

Two encoding agents, both comprising a coding tag (seen 60 at 315 and 317), a ligation region (seen at 311 and 313), and a primer region (seen at 319 and 321) are combined with and ligated to the extended probe at step 304 to form an encoded target-specific oligonucleotide. Again, in contrast with FIG. 1, the probes and encoding agents are delivered at separate steps. Doing so allows use of the combinatorial embodiments described infra. In preferred embodiments, the encoding oligonucleotides within a pair of encoding oligonucleotides ligate specifically to one side of the target sequence or the other (i.e., 5' or 3' of the target sequence) in step 306.

Also, typically, the ligation and primer regions of the encoding oligonucleotides and probes are universal; that is, the set of ligation and primer regions used in constructing the probes and encoding oligonucleotides are constant, and only the target-specific regions of the probes and the coding tags of the encoding oligonucleotides differ. However, again in alternative embodiments, the ligation and primer regions are not universal and differ between probes and encoding agents.

Following ligation, the encoded probes are eluted, pooled, and, optionally, sequencing adapters are added to the encoded probes via PCR. In alternative embodiments, sequencing primers may be ligated to the encoding oligonucleotides, or sequencing primer sequences can be included as part of the encoding oligonucleotide. As seen in FIG. 3, each sequencing adapter comprises primer region 319 or 321, compatible with the primer regions 319 and 321 on the encoded probes. The final construct comprising first adapter 327, first primer region 319, first coding tag 315, ligation regions 311 and 301, target region 309, ligation regions 313 and 303, second coding tag 317, second primer region 325 and second adapter 329 is now ready for input into a digital high-throughput sequencing process.

A combination of extension and ligation reactions are exemplified in FIG. 3, but it should be appreciated that a variety of reactions may be used to couple the encoding oligonucleotides to the target-specific oligonucleotides, including ligation only (e.g., for oligonucleotides that hybridize to contiguous portions of the target nucleic acid sequence). Alternatively, an assay utilizing an additional oligonucleotide, such as in the GOLDEN-GATE® assay (see Fan, et al., Cold Spring Symp. Quant. Biol., 68:69-78 (2003); (Illumina, Inc., San Diego, CA)), may be employed.

In other embodiments, the assay system of the invention also can be used to analyze peptides or proteins, the presence of antibodies, enzymatic and other protein activities, post-translational modifications, active and non-active forms of peptides, as well as peptide isoforms in a biological sample. Accordingly, the probes may comprise an active region of an enzyme, a binding domain of an immunoglobulin, defined domains of proteins, whole proteins, synthetic peptides, peptides with introduced mutations, aptamers and the like.

In certain aspects, the probes are substrates for enzymes or proenzymes, e.g., kinases, phosphatases, zymogens, proteases, or fragments thereof. In certain aspects, the probes are phosphorylation substrates used to detect proteins involved in one or more signal transduction pathways, e.g., a kinase or a phosphatase. In another specific aspect of the invention, the probes are specific protease substrates that associate only with individual proteases or classes of proteases. In other aspects, the probes are different processed forms, isoforms and/or domains of an enzyme. Protein-based probes are typically conjugated or otherwise linked to oligonucleotide encoding agents. The oligonucleotide encoding agents in this case would also include a nucleotide sequence component that allows for identification of the protein probe.

In certain aspects, the present invention provides assays for evaluating differences in the amount and/or activity of biological targets between different locations in a sample and/or between samples. The method includes determining a plurality of encoded results from the biological sample and evaluating the differences in quantity of the biological targets at each location in the biological sample.

#### Combinatorial Embodiments

To maximize the efficiency of encoding, a combinatorial approach using pairs of coding tags in the encoding oligo-

nucleotides can be used. By de-coupling the target-specific information and the coding tags, the number of oligonucleotides required is dramatically reduced, with a concomitant decrease in cost.

FIG. 4 illustrates a general mechanism for one embodiment of a combinatorial encoding scheme of the assay systems of the invention, where nucleic acids in a representative tissue section (shown at 416) are assayed. FIG. 4 at A shows two target-specific/encoding oligonucleotide constructs 420 and 422 (e.g., formed between steps 302 and 304 of FIG. 3) specifically bound to a target nucleic acid 402 of interest. The first encoded probe 420 comprises coding tag 408, associated with, e.g., a universal priming site for amplification of the assay products or an adapter to enable identification of the coding identifiers using sequencing technologies 404. The second encoded probe 422 comprises coding tag 406, associated with, e.g., a universal priming site for amplification of the assay products or an adapter to enable identification of the coding identifiers using sequencing technologies 410.

FIG. 4 at B shows the spatial pattern that may be used for twenty different coding tags, a1 through a10 (coding tag 406 on encoded probe 420) and b1 through b10 (coding tag 408 encoded probe 422). Coding tag a1, for example, is deposited on the biological sample in ten discrete areas or spots (shown as the first horizontal line of spots in 412). Coding tag a2 is deposited on the biological sample in ten spots on the second horizontal line in 412. Coding tag a3 is deposited on the biological sample in ten spots on the third horizontal line in 412, and so on. Whereas the "a" tags are deposited in ten horizontal rows, the "b" tags are deposited in ten vertical rows as shown in 414. For example, coding tag b1 is deposited on the biological sample in ten discrete spots in the first vertical row of 414, coding tag b2 is deposited on the biological sample in ten discrete spots in the second vertical row of 414, and so on. Using such a configuration allows for twenty coding tags to uniquely define 100 different locations on the biological sample.

FIG. 4 at C shows a representative tissue section 416 coincident with coding tag grid 418. The arrows show how the "a" coding tags and the "b" coding tags are deposited on grid 418 that is coincident with tissue section 416. If, once sequenced, coding tags a1 and b4, e.g., are associated with a target nucleic acid sequence, then that target nucleic acid sequence (i.e., biological target) was present in the tissue section at location a1, b4.

FIG. 5 provides a simplified, specific example of the encoding scheme of the assay systems of the invention. FIG. 5 shows encoding oligonucleotides 510, comprising a1, a2, a3, a4 and b1, b2, b3 and b4. Target-specific oligonucleotides (TSOs) (probes) 1 and 2 are shown at 520. A deposit or dispensing scheme is shown at 530. Like the grid exemplified in FIG. 4, encoding oligonucleotides a1 through a4 are deposited in spots in a pattern (here, in a vertical pattern), and encoding oligonucleotides b1 through b4 are deposited in spots in a pattern (here, a horizontal pattern). The grid though shown as a square with spots is actually a deposition pattern on a biological sample (not shown) such as tissue section 416 shown in FIG. 4.

The target-specific oligonucleotides are delivered to the biological sample, where the target-specific oligonucleotides hybridize to target nucleic acids in the biological sample if target nucleic acids are present. Unhybridized target-specific oligonucleotides are then removed, e.g., by washing. The encoding oligonucleotides are then delivered to the biological sample according to the spatial pattern shown at 530. The encoding oligonucleotides are ligated (or, e.g., extended and

15

ligated) to any target-specific oligonucleotides that hybridized to the target nucleic acid in the biological sample, the ligated constructs are then eluted from the biological sample, pooled, and sequencing adapters are added through, e.g., PCR or ligation, if the sequences were not previously included in the encoding oligonucleotides. The ligated constructs are sequenced by, e.g., high throughput or “next generation” sequencing.

The pool of resulting sequences is shown at 540. A sequence readout was obtained for target-specific oligonucleotide 1 only at a4b1, a4b2, a1b3, a2b3, a3b3, a4b3 and a4b4 (positions shown with horizontal lines). A sequence readout was obtained for target-specific oligonucleotide 2 only at a1b1 (position shown with vertical lines). A sequence readout was obtained for both target-specific oligonucleotides 1 and 2 at positions a2b1, a3b1, a1b2, a2b2, and a3b2 (positions shown with cross-hatching). No sequence readout was obtained for either target-specific oligonucleotides at a1b4, a2b4 or a3b4 (positions shown without shading). Thus, in the biological sample on which the assay took place the first target nucleic acid was detected in a large portion of the left side and at the bottom of the biological sample, the second target nucleic acid was detected only in the upper left portion of the biological sample, and neither target nucleic acid was detected in the upper right portion of the biological sample. The differential expression of the two target nucleic acids now can be mapped back to the biological sample and to the biological structures or cell types in these locations in the biological sample, as shown in 550.

In addition to location information, information relating to relative abundance of the encoded tags can be obtained. For example, if it is found that there are ten times as many a4T1b1 sequences occurring in the data set as compared to a4T1b2 sequences, this would indicate that target nucleic acid sequence 1 is ten times more abundant at the a4T1b1 location than at the a4T1b2 location.

In the case of nucleotide analysis as shown in FIG. 3, by ligating the coding tags directly to target-specific oligonucleotides, only  $2n$  target-specific oligonucleotides are needed for  $n$  targets. For example, using the combinatorial approach outlined in FIG. 2, assaying 100 different targets at 10,000 spatial locations would require  $2 \times 100$  target-specific oligonucleotides and  $2 \times 100$  encoding oligonucleotides. The total count of assay oligonucleotides would be only 400 (200 target-specific and 200 encoding), not counting universal primers. In contrast, if the coding oligonucleotides were not decoupled from the target-specific oligonucleotides, ( $n \times X$  positional codes) + ( $n \times Y$  positional codes) would be needed, or in the above example, 20,000 oligonucleotides, not counting universal primer sequences. Moreover, though the embodiments shown in FIGS. 2-5 depict a combinatorial scheme using two encoding agents (coding tags), three, four or more encoding agents and coding tags may be used, and attached to the probe or one another by varying means and in varying combinations of steps.

Due to the spatial encoding aspect of the assay system of the invention, a large amount of information can be generated with even a modest number of assays. For example, five or more biological targets assayed at five or more positions in the sample generates 25 or more combinations. Using digital sequencing as a readout, the optimum number of sequence reads per combination depends on the sensitivity and dynamic range required, and can be adjusted. For example, if for each combination on average 100 reads are sampled, the total for 25 combination is 25,000 reads. If 1,000 targets are assayed at 1,000 locations with an average sampling depth of 1,000, then  $10^9$  reads are required. These

16

numbers, although large, are within the capacity of intrinsically parallel digital sequencing methods, which can generate datasets of billions or even trillions of reads in a reasonable timeframe and at a very low cost per read. Therefore, by varying the numbers of positions interrogated or biological targets assayed, or both, and using digital sequencing, large amounts of information can be obtained. In specific aspects, multiple locations are interrogated for two or more biological molecules.

### Reagent Delivery Systems

The reagent delivery system of the invention includes instrumentation that allows the delivery of reagents to discrete portions of the biological sample, maintaining the integrity of the spatial patterns of the encoding scheme. Reagent delivery systems of the assay systems of the invention comprise optional imaging means, reagent delivery hardware and control software. Reagent delivery can be achieved in a number of different ways. It should be noted that reagent delivery may be to many different biological samples at one time. A single tissue section has been exemplified herein; however, multiple biological samples may be affixed and analyzed simultaneously. For example, pions of a tissue sample can be analyzed in parallel and the data combined to build a 3D map.

Integral to the assay system of the invention is instrumentation that allows for spatial patterning of reagents onto the biological sample. Technologies for formulating and delivering both biological molecules (e.g. oligonucleotides or antibodies) and chemical reagents (e.g., small molecules or dNTPs) are known in the art, and uses of these instrument systems are known to one skilled in the art and easily adaptable to the assay systems of the invention. One example of a suitable reagent delivery system is the Labcyte<sup>TM</sup> Echo acoustic liquid handier, which can be used to deliver nanoliter scale droplets containing biological molecules with high precision and reproducibility. One skilled in the art could incorporate this reagent delivery device into the overall system, using software to specify the locations to which reagents should be delivered.

Other instruments that can be used for the deposition of agents and/or coding identifiers onto biological samples include, but are not limited to, ink jet spotting; mechanical spotting by means of pin, pen or capillary; micro contact printing; photochemical or photolithographic methods; and the like. For several applications, it may be preferred to segment or sequester certain areas of the biological samples into one or more assay areas for different reagent distributions and/or biological target determination. The assay areas may be physically separated using barriers or channels.

In one exemplary aspect, the reagent delivery system may be a flow-based system. The flow-based systems for reagent delivery in the present invention can include instrumentation such as one or more pumps, valves, fluid reservoirs, channels, and/or reagent storage cells. Reagent delivery systems are configured to move fluid to contact a discrete section of the biological sample. Movement of the reagents can be driven by a pump disposed, for example, downstream of the fluid reagents. The pump can drive each fluid reagent to (and past) the reaction compartment. Alternatively, reagents may be driven through the fluid by gravity. US Pub. Nos. 20070166725 and 20050239192 disclose certain general-purpose fluidics tools that can be used with the assay systems of the invention, allowing for the precise manipu-

17

lation of gases, liquids and solids to accomplish very complex analytical manipulations with relatively simple hardware.

In a more specific example, one or more flow-cells can be attached to the substrate-affixed biological sample from above. The flow-cell can include inlet and outlet tubes connected thereto and optionally an external pump is used to deliver reagents to the flow-cell and across the biological sample. The flow cells are configured to deliver reagents only to certain portions of the biological sample, restricting the amount and type of reagent delivered to any specific section of the biological sample.

In another aspect, a microfluidic system can be integrated into the substrate upon which the biological sample is disposed or externally attached on top of the substrate. Microfluidic passages for holding and carrying fluid may be formed on and/or above the planar substrate by a fluidics layer abutted to the substrate. Fluid reagents can be selected and delivered according to selective opening and closing of valves disposed between reagent reservoirs.

Pumps generally include any mechanism for moving fluid and/or reagents disposed in fluid. In some examples, the pump can be configured to move fluid and/or reagents through passages with small volumes (i.e., microfluidic structures). The pump can operate mechanically by exerting a positive or negative pressure on fluid and/or on a structure carrying fluid, electrically by appropriate application of an electric field(s), or both, among other means. Exemplary mechanical pumps may include syringe pumps, peristaltic pumps, rotary pumps, pressurized gas, pipettors, etc. Mechanical pumps may be micromachined, molded, etc. Exemplary electrical pumps may include electrodes and may operate by electrophoresis, electroendoosmosis, electrocapillarity, dielectrophoresis (including traveling wave forms thereof), and/or the like.

Valves generally include any mechanism for regulating the passage of fluid through a channel. Valves can include, for example, deformable members that can be selectively deformed to partially or completely close a channel, a movable projection that can be selectively extended into a channel to partially or completely block a channel, an electrocapillary structure, and/or the like.

An open gasket can be attached to the top of the biological sample and the sample and reagents can be injected into the gasket. Suitable gasket materials include, but are not limited to, neoprene, nitrile, and silicone rubber. Alternatively, a watertight reaction chamber may be formed by a gasket sandwiched between the biological sample on the substrate and a chemically inert, water resistant material such as, but not limited to, black-anodized aluminum, thermoplastics (e.g., polystyrene, polycarbonate, etc), glass, etc.

In an optional embodiment, the assay system comprises imaging means to determine features and organization of the biological sample of interest. The images obtained, e.g., may be used to design the deposition pattern of the reagents. Imaging means are optional, as an individual can instead view the biological sample using, e.g., a microscope, analyze the organization of the biological sample, and specify a spatial pattern for delivery assay reagents. If included, the delivery system can comprise a microcircuit arrangement including an imager, such as a CCD or IGFET-based (e.g., CMOS-based) imager and an ultrasonic sprayer for reagent delivery such as described in US Pub. No. 20090197326, which is incorporated herein by reference. Also, it should be noted that although FIGS. 4 and 5 illustrate using a x,y grid configuration, other configurations can be used, such as, e.g.,

18

following the topology of a tissue sample; targeting certain groups of cells, cell layers and/or cell types in a tissue, and the like.

In yet another alternative, the reagent delivery system controls the delivery of reagents to specific patterns on a biological sample surface using semiconductor techniques such as masking and spraying. Specific areas of a biological sample can be protected from exposure to reagents through use of a mask to protect specific areas from exposure. The reagents may be introduced to the biological sample using conventional techniques such as spraying or fluid flow. The use of masked delivery results in a patterned delivery scheme on the substrate surface.

In a preferred aspect of the invention, the reagent delivery instrumentation is based on inkjet printing technology. There are a variety of different ink-jetting mechanisms (e.g., thermal, piezoelectric) and compatibility has been shown with aqueous and organic ink formulations. Sets of independently actuated nozzles can be used to deliver multiple reagents at the same time, and very high resolutions are be achieved.

In order to target specific sites of interest, an informative image of the biological sample to be assayed may be used to assist in the reagent delivery methods and associated encoding scheme. Sample regions of the biological sample can be identified using image processing (e.g., images of cell types differentiated by immunohistochemistry or other staining chemistries) integrated with other features of the assay system. In some aspects, software is used to automatically translate image information into a reagent delivery pattern. A mechanism to register and align very precisely the biological sample for reagent delivery is thus an important component of the assay systems of the invention. Mechanisms such as the use of fiducial markers on slides and/or other very accurate physical positioning systems can be adapted to this purpose.

The invention preferably comprises a complete suite of software tailored to the assay system. Optionally, oligonucleotide design software is used to design the encoding nucleotides (and in embodiments where nucleic acids are assayed, the target-specific oligonucleotides) for the specific assay to be run, and may be integrated as a part of the system. Also optionally, algorithms and software for reagent delivery and data analysis (i.e., sequence analysis) may be integrated to determine assay results. Integrated data analysis is particularly useful, as the type of dataset that is generated may be massive as a consequence of scale. Algorithms and software tools that are specifically designed for analysis of the spatially-associated data generated by the assay systems, including pattern-analysis software and visualization tools, enhance the value of the data generated by the assay systems.

In certain aspects, the assay system comprises processes for making and carrying out the quality control of reagents, e.g., the integrity and sequence fidelity of oligonucleotide pools. In particular, reagents are formulated according to factors such as volatility, stability at key temperatures, and chemical compatibility for compatibility with the reagent delivery instrumentation and may be analyzed by instrumentation integrated within the assay system.

### Sequencing

Numerous methods can be used to identify the coding tags and probe sequences in the encoded probes of the assay systems of the invention. The coding tags can be detected using techniques such as mass spectroscopy (e.g., Maldi-T

of, LC-MS/MS), nuclear magnetic resonance imaging, or, preferably, nucleic acid sequencing. Examples of techniques for decoding the coding tags of the present invention can be found, for example, in US Pub. No. 20080220434, which is incorporated herein by reference. For example, the coding tags may be oligonucleotide mass tags (OMTs or massTags). Such tags are described, e.g., in US Pub. No. 20090305237, which is incorporated by reference in its entirety. In yet another alternative, the encoded probes can be amplified and hybridized to a microarray. This would require separate amplification reactions to be carried out, in which each amplification is specific to a particular spatial code or subset of codes, accomplished by using code-specific primers. Each amplification would also incorporate a different resolvable label (e.g. fluorophor). Following hybridization, the relative amounts of a particular target mapping to different spatial locations in the sample can be determined by the relative abundances of the resolvable labels.

In one particularly preferred aspect, the resulting coding tags according to the assay system are substrates for high-throughput, next-generation sequencing, and highly parallel next-generation sequencing methods are used to confirm the sequence of the coding tags, for example, with SOLID™ technology (Life Technologies, Inc.) or Genome Analyzer (Illumina, Inc.). Such next-generation sequencing methods can be carried out, for example, using a one pass sequencing method or using paired-end sequencing. Next generation sequencing methods include, but are not limited to, hybridization-based methods, such as disclosed in e.g., Drmanac, U.S. Pat. Nos. 6,864,052; 6,309,824; and 6,401,267; and Drmanac et al., U.S. patent publication 2005/0191656; sequencing-by-synthesis methods, e.g., U.S. Pat. Nos. 6,210,891; 6,828,100; 6,969,488; 6,897,023; 6,833,246; 6,911,345; 6,787,308; 7,297,518; 7,462,449 and 7,501,245; US Publication application Ser. Nos. 20/110,059436; 20040106110; 20030064398; and 20030022207; Ronaghi, et al., *Science*, 281:363-365 (1998); and Li, et al., *Proc. Natl. Acad. Sci.*, 100:414-419 (2003); ligation-based methods, e.g., U.S. Pat. Nos. 5,912,148 and 6,130,073; and U.S. Pat. Appln Nos. 20100105052, 20070207482 and 20090018024; nanopore sequencing e.g., U.S. Pat. Appln Nos. 20070036511; 20080032301; 20080128627; 20090082212; and Soni and Meller, *Clin Chem* 53:1996-2001 (2007)), as well as other methods, e.g., U.S. Pat. Appln Nos. 20110033854; 20090264299; 20090155781; and 20090005252; also, see, McKernan, et al., *Genome Res.*, 19:1527-41 (2009) and Bentley, et al., *Nature* 456:53-59 (2008), all of which are incorporated herein in their entirety for all purposes.

#### Applications of Assay System

It will be apparent to one skilled in the art upon reading the present disclosure that there are numerous important areas of biological research, diagnostics, and drug development that will benefit from a high throughput multiplexed assay system that can measure simultaneously the amount and spatial location of a biological target in a biological sample. For example, combining the ability to estimate the relative abundance of different RNA transcripts with the ability to reconstruct an image of spatial patterns of abundance across many locations, which may be as small as or even smaller than individual cells, in a tissue enables many different areas of basic research. The following are exemplary uses and are by no means meant to be limiting in scope.

In one example, 3-dimensional patterns of gene expression are determined by analyzing a series of tissue sections, in a manner analogous to image reconstruction in CT scanning. Such a method can be used to measure changes in gene expression in disease pathology, e.g., in cancerous tissue and/or a tissue upon injury, inflammation or infection. With the assay systems of the invention, more detailed information on gene expression and protein localization in complex tissues is obtained, leading to new insights into the function and regulation both in normal and diseased states, and provides new hypotheses that can be tested. For example, an assay system of the invention may enable some of the insights gained from many individual studies and larger programs like ENCODE (Birney, et al., *Nature*, 447: 799-816 (2007)) and modENCODE to be integrated at the tissue level. The assay systems also aid computational efforts to model interacting networks of gene expression in the field of systems biology.

The assay systems also provide a novel approach to analysis of somatic variation, e.g., somatic mutations in cancer or variability in response to infectious organisms. For example, tumors are typically highly heterogeneous, containing cancer cells as well as genetically normal cells in an abnormal local environment. Cancer cells undergo mutation and selection, and in this process it is not unusual for local clones to develop. Identifying relatively rare somatic mutations in the context of tumors may enable the study of the role of key mutations in the selection of clonal variants. Transcriptional patterns associated with angiogenesis, inflammation, or other cancer-related processes in both cancer and genetically normal cells can be analyzed for insights into cancer biology and assist in the development of new therapeutic agents for the treatment of cancers. In another example, individuals have varying susceptibility to infectious organisms, and the assay systems of the invention can be used to study the interaction between microbes and tissues or the various cell types within the tissue.

Importantly, in addition to providing spatially-associated information, the invention allows a great increase in the sensitivity of detecting rare mutations, as signal to noise can be dramatically increased since only a small location is assayed in any given reaction. In a typical assay for rare mutations in a mixed sample, the sample is treated in bulk, i.e., nucleic acids are extracted from many cells into a single pool. Thus, if a mutation is present in one cell in 10,000, it must be detected against a background of normal DNA from ~10,000 cells. In contrast, with the assay systems of the invention many cells can be analyzed, but individual cells or small groups of cells would be identified by the spatial coding system. Therefore, in the assay systems of the present invention, background is reduced by orders of magnitude, greatly increasing sensitivity. Furthermore, the spatial organization of mutant cells can be observed, which may be particularly important in detecting key mutations in tissue sections in cancer. Already molecular histological analyses are yielding insights into cancer biology and may have potential for use in diagnostics. The technology of the invention promises to greatly increase the power of such approaches.

The present invention provides assays, assay systems, and methods of using such assays in spatially encoded biological assays. The invention provides an assay system comprising one or more agents provided in defined spatial patterns on a substrate surface, and a detection system for identifying the presence or absence, relative amount, and location of a biological molecule. Such biological molecules include, but are not limited to, nucleic acids, peptides, carbohydrates,

cellular components, and the like. The assay system is a novel multiplexing approach, as it allows multiple molecules and their respective multiple locations to be identified in a single system using a unique encoding scheme. This encoding scheme uses both molecule-specific binding agents and coding identifiers to provide a practical and cost-effective determination of information on multiple biological molecules, including specific positional information of such molecules in a biological sample, e.g., a tissue section. The single molecule detection analysis using the encoding system also allows relative amounts of biological molecules to be detected, thus providing information on expression levels, sequestering in specific locales, and the like.

The assay systems detect the presence or absence, and relative amount, of a biological molecule at more than one spatial location in a sample. In addition, the assays provide methods for doing this for multiple biological molecules simultaneously. The assay systems utilize one or more binding agents that specifically bind to the biological molecule of interest and unique coding identifiers associated with specific binding agents. The detection system utilizes a method for identifying the presence and spatial address of the agent binding based on the positive and/or negative results that are obtained using detection of the agent and identifier and the encoding scheme of the spatial patterns on the substrate surface. In a specific aspect, the encoding scheme employs limited reagent delivery to the spatial patterns on the substrate surface, and access of the coding identifiers and/or binding agents to portions of the sample is controlled through such limited delivery.

In one aspect, the assay system detects the presence or absence and spatial location of a biological molecule based on the positive and/or negative results that are obtained using limited reagent delivery and the encoding scheme of the spatial patterns on the substrate surface.

The assay system and methods of the invention are based on relational, solid-state substrates with positions that represent specific spatial locations within a biological sample, e.g., a cell, organelle or tissue. The ability to use encoding features to represent locations allows high-throughput analysis of the presence or absence, and relative amount, of a biological molecule at more than one spatial location in a sample. The encoding features also allow provide assaying of multiple biological molecules at these multiple locations simultaneously.

A primary feature of the invention is the preservation of the spatial organization of elements in a sample of interest through the use of an encoding scheme. For example, the assay may be designed to preserve the relative position of cells in a tissue, and the assay may interrogate the individual cells for genomic DNA variation (including epigenetic modifications), and RNA and protein expression.

In one specific aspect, the encoding scheme of the assay system comprises the use of two or more coding patterns, each comprising regions defined by spatial patterns on the substrate surface. For example, the assay system can utilize an encoding scheme that comprises a 2-dimensional grid format based on the discrete positioning of the binding agents in the substrate surfaces. In another example, the spatial patterns may be based on more randomized cell locations, e.g., the patterns on the substrate surface follow an underlying biological structure rather than a strict, x,y grid pattern. This aspect includes systems with two or more substantially identical spatial patterns using different binding agents and/or coding identifiers, as well as systems having different patterns for different agents and/or coding

identifiers. The encoding scheme of the systems can be controlled by delivery of different reagents to discrete regions on the substrate surfaces, which allows different reactions to take place on substantially similar agents of known location on the substrate surfaces.

In one specific aspect, the invention provides high resolution, high-throughput analysis of nucleic acids and/or expression levels that provides both detection and spatial identification of large numbers of nucleic acids, e.g., DNA or RNA.

In another specific aspect, the invention provides high resolution, high-throughput analysis of proteins that provides both detection and spatial identification of large numbers of such proteins, e.g., kinases or proteases.

Numerous reagent delivery systems can be used with the assay system of the invention. The primary criteria of such reagent delivery systems is the ability to direct delivery of specific agents based on spatial patterns on the substrate surface.

In one preferred aspect, the encoding scheme utilizes a reagent delivery system based on printing and informatics technologies to implement the spatial patterns used for identification and localization of the biological materials. For example, the patterns found in the encoding scheme may be created using ink jet printing technology to provide reagents at specific locations on one or more substrate surfaces. The desired patterns are set out in specific coding patterns on the substrate surface.

In certain aspects of the invention, the binding agents are immobilized directly to the substrate surface, and the location of the binding agents is known or determined prior to use of the substrate surface in the assay system. In another aspect, the binding agents are immobilized onto beads or other separate structural elements that are then provided in known locations on the substrate surface. In yet another aspect, the binding agents may be provided in or on features of the substrate surface, e.g., provided in wells or channels.

In specific aspects of the invention, the binding agents are nucleic acids immobilized directly or indirectly to the substrate surface, e.g., directly through the use of amino groups on the substrate surface or indirectly through the use of a linker. The location of the nucleic acid sequences is known or determined prior to use of the substrate surface in the assay system. In another specific aspect, the nucleic acids may be immobilized directly or indirectly onto beads that are then provided in known locations on the substrate surface. In yet another aspect, the nucleic acids may be provided in or on features of the substrate surface, e.g., provided in wells.

In these aspects involving nucleic acid agents, any methods of sequence determination can be used, e.g., sequencing, hybridization and the like. In a preferred aspect, nucleic acid sequencing, and preferably next-generation sequencing, is used to decode the spatial encoding scheme in the assay system of the invention. This provides a very wide dynamic range for very large numbers of assays, allowing for efficient multiplexing.

In some aspects, the assay utilizes two or more oligonucleotides, the oligonucleotides comprising a universal primer region and a region that correlates specifically to a single spatial pattern within the spatial encoding scheme. In a specific aspect, the assay comprises two allele specific oligonucleotides and one locus specific oligonucleotides. These oligonucleotides allow the identification of specific SNPs, indels or mutations within an allele. This is useful in the identification of genetic changes in somatic cells, genotyping of tissues, and the like.

In other specific aspects of the invention, the binding agents are peptides. In one aspect, these peptides are associated directly or indirectly to known locations on a substrate surface, e.g., using binding protein pairs or through oligonucleotide linkers complementary to oligonucleotides on the substrate surface. In another aspect, the binding agents are peptides are immobilized directly or indirectly onto beads or other separate structural elements that are then provided in known locations on the substrate surface. In yet another aspect, the peptides may be provided in or on features of the substrate surface, e.g., provided in wells.

In yet other specific aspects of the invention, the binding agents are chemical entities (e.g., small molecules) that are coded, e.g. using sequence tags or mass spectroscopy tags as coding identifiers. In one aspect, these chemical entities can be immobilized directly to the substrate surface. In another aspect, the binding agents are immobilized onto beads or other separate structural elements that are then provided in known locations on the substrate surface. In yet another aspect, the binding agents may be provided in or on features of the substrate surface, e.g., provided in wells.

The assay system of the invention can utilize various detection mechanisms, based on the molecules to be detected and the reagents needed for such detection system. Exemplary methods that can be used with the assay systems of the invention are described in more detail below.

#### The Invention in General

The assay system and methods of the invention are based on relational methods that allow extraction of data to detect the presence or absence and relative amount of a biological molecule, and the location of this molecule in a sample having a distinct structure, e.g., a tissue section or other biological structure with distinct locations of specific biological molecules. The encoding scheme used in these systems corresponds to the structural elements of the sample, and the information obtained using a two-dimensional coding system is indicative of the spatial addresses of these molecules in a sample of interest.

Integral to the assay system of the invention is a method for spatial patterning of reagents. Technologies for formulating and delivering both biological molecules (e.g. DNA or antibodies) and chemical reagents (e.g., small molecules or dNTPs) have already been demonstrated, and use of these systems will be available to one skilled in the art and easily adaptable upon reading this specification.

The assay design of the invention provides an accurate and easily scalable spatial encoding system. The ability to deliver reagents in a spatially defined pattern together with software, reagents and protocols comprises a novel and highly innovative assay system for spatial analysis of various biological molecules and activities. This allows the assays to measure numerous biological functions in a meaningful spatial environment, including functions such as gene expression and peptide localization. The systems provide the potential to open a new analytical window into the complex spatial patterns of cellular function and regulation in biological systems.

The biological molecules to be detected can be any biological molecules such as proteins, nucleic acids, lipids, carbohydrates, ions, or multicomponent complexes containing any of the above. Further examples of subcellular objects include organelles, e.g., mitochondria, Golgi apparatus, endoplasmic reticulum, chloroplast, endocytic vesicle, exocytic vesicles, vacuole, lysosome, etc.

FIG. 4 illustrates such a target-specific assay system for identification of nucleic acid sequences in a sample. In this system, two reagents 420, 422 that specifically bind to a biological molecule of interest are associated with coding identifiers 406, 408 that encode for a spatial location in the sample. These coding identifiers 406, 408 are optionally associated with sites that assist in their identification in the assay format, e.g., universal priming sites 404, 410 for amplification of the assay products or adapters to enable identification of the coding identifiers and the binding agents using sequencing technologies. The sample that is tested, here shown as a tissue section 416 is encoded using the combination of the patterns 412, 414 created using the separate coding identifiers 406, 408 which provide a two dimensional code 418 that shows the location of any positive detection of the biological molecule 402 as well as quantifying the biological molecule 402 at each location assayed in the tissue.

The assay systems of the invention are particularly advantageous in that they are compatible with numerous samples types, such as fresh samples, such as primary tissue sections, and preserved samples including but not limited to frozen samples and paraformalin-fixed, paraffin-embedded (FFPE) samples. An important aspect of the assay systems of the invention is that the binding agents are immobilized on a substrate surface in discrete, independently measurable areas. These discrete areas can be formed by spatially selective deposition of the binding agents on the substrate surface. Numerous methods can be used for the deposition of the agent and the coding identifiers associates with the agent. For example, the coding identifiers can be delivered together or separately from the agent. If delivered together they can be attached (e.g., synthesized as a single molecule or attached through ligation or a chemical coupling mechanism) or simply mixed together to be attached after delivery to the substrate. In a preferred aspect, the agent and the coding identifier are made separately, mixed together for attachment, and delivered either attached or as a mixture to be attached on the surface. In a specific aspect the binding agents are delivered generally over the substrate surface and the coding identifiers are delivered in a pattern-specific manner.

Examples of methods that can be used for deposition of agents and/or coding identifiers onto the substrate surface include, but are not limited to, ink jet spotting, mechanical spotting by means of pin, pen or capillary, micro contact printing, fluidically contacting the measurement areas with the biological or biochemical or synthetic recognition elements upon their supply in parallel or crossed micro channels, upon exposure to pressure differences or to electric or electromagnetic potentials, and photochemical or photolithographic immobilization methods.

For several applications, it may be preferred to arrange the substrates into segments of one or more measurement areas for reagent distribution and agent determination. These regions may be physically separated using barriers or channels. They may still comprise several additional discrete measurement areas with agents that are different or in different combination from each other.

In certain aspects, the present invention provides a method, e.g., a machine-based method, for evaluating changes in the presence and/or location of a biological molecule over time. The method includes providing a plurality of encoded array results representative of the biologi-

cal molecule over time and evaluating the differences in detection and/or localization of the biological molecules.

#### Nucleic Acid Detection and Localization

In a particular aspect, the assay system is used to analyze nucleic acids, e.g. genotyping, gene expression analysis, localization of particular transcripts within samples, and the like.

Genotyping may be performed using any technique known to those of skill in the art. Preferred techniques permit rapid, accurate determination of multiple variations with a minimum of sample handling. Some examples of suitable techniques involve but are not limited to direct DNA sequencing, capillary electrophoresis, hybridization, allele-specific probes or primers, single-strand conformation polymorphism analysis, nucleic acid arrays, bead arrays, restriction fragment length polymorphism analysis, cleavage fragment length polymorphism analysis, random amplified polymorphic DNA, ligase detection reaction, heteroduplex or fragment analysis, differential sequencing with mass spectrometry, atomic force microscopy, pyrosequencing, FRET (e.g., TaqMan (Applied Biosystems, Inc., Foster City, Calif.) and Molecular Beacon (Stratagene, La Jolla, Calif.) assays), and other related techniques. Several methods for DNA sequencing are well known and generally available in the art. See, for example, Sambrook, et al., *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory, New York) (2001); Ausubel, et al., *Current Protocols in Molecular Biology* (John Wiley and Sons, New York) (1997), Twyman, et al. (2003) "Techniques Patents for SNP Genotyping", *Pharmacogenomics* 4 (1): 67-79; and Kristensen, et al. (2001) "High-Throughput Methods for Detection of Genetic Variation", *BioTechniques* 30 (2): 318-332. For details on the use of nucleic acid arrays (DNA chips) for the detection of, for example, SNPs, see U.S. Pat. No. 6,300,063 issued to Lipshultz, et al., and U.S. Pat. No. 5,837,832 to Chee, et al., *HuSNP Mapping Assay*, reagent kit and user manual, Affymetrix Part No. 90094 (Affymetrix, Santa Clara, Calif.). The molecular inversion probe (MIP) assay format (Hardenbol et al., 2003) is another example of a highly multiplexable assay that may be used with the assay systems of the invention.

In one exemplary and preferred method for analyzing nucleic acids using the assay system of the invention, the detection of nucleic acids uses two allele-specific oligonucleotides and a locus specific oligonucleotide. The assay methods are carried out according to the strategy outlined in FIG. 2 using next-generation sequencing or another highly parallel nucleic acid assay technology. In this assay, a set of two oligonucleotides is designed to hybridize to each target sequence, with a common oligonucleotide and two unique coding identifiers. The allele can be determined, e.g., by primer extension of the locus specific oligonucleotide. Following primer extension and ligation, an amplifiable template is formed with universal primer sequences at either end. Assay oligonucleotides are annealed to a template and enzymatic reactions are used to join the two oligonucleotides only when both are correctly annealed. The detection techniques and read out parameters used in this system of the invention include a much shorter tag than the oligonucleotides used in the assays that are based on capture by hybridization. These shorter tags are designed to be read out by sequencing or, preferably, used to ligate codes onto both ends of the fragment as illustrated in FIG. 2.

In FIG. 3, two target-specific assay oligonucleotides are ligated together 302 following *in situ* hybridization to target

sequences. At the same time, encoding oligonucleotides containing tag sequence sets X and Y are ligated 304 to the target specific oligonucleotides. Oligonucleotides containing X ligate specifically to one side of the targeting construct 5 and oligonucleotides containing Y ligate to the other. The oligonucleotides contain universal primer sites P1 and P2. Following ligation, the constructs are eluted and, optionally, sequencing adapters can be attached 306, e.g., via PCR.

In one preferred aspect, the final construct created from 10 the assay method is a substrate for next-generation sequencing, and highly parallel next-generation sequencing methods are used to confirm the sequence of constructs. Such sequencing methods can be carried out, for example, using a one pass sequencing method or using paired-end sequencing. Next generation sequencing methods include, but are 15 not limited to, hybridization-based methods, such as disclosed in Drmanac, U.S. Pat. Nos. 6,864,052; 6,309,824; and 6,401,267; and Drmanac et al, U.S. patent publication 2005/0191656, and sequencing by synthesis methods, e.g., 20 Nyren et al, U.S. Pat. No. 6,210,891; Ronaghi, U.S. Pat. No. 6,828,100; Ronaghi et al (1998), *Science*, 281:363-365; Balasubramanian, U.S. Pat. No. 6,833,246; Quake, U.S. Pat. No. 6,911,345; Li et al, *Proc. Natl. Acad. Sci.*, 100:414-419 (2003); Smith et al, PCT publication WO 2006/074351; use 25 of reversible extension terminators, e.g., Turner, U.S. Pat. No. 6,833,246 and Turner, U.S. Pat. No. 6,833,246 and ligation-based methods, e.g., Shendure et al (2005), *Science*, 309:1728-1739, Macevicz, U.S. Pat. No. 6,306,597; which references are incorporated by reference. Soddart et al., 30 PNAS USA, 2009 Apr. 20; Xiao et al., *Nat Methods*. 2009 March; 6(3):199-201. Epub 2009 Feb. 8.

To maximize the efficiency of encoding, a combinatorial approach using pairs of oligonucleotides can be used. For example, with only two sets of 100 codes, a substrate can 35 theoretically encode up to 10,000 locations. The number of assay oligonucleotides required is dramatically reduced, the cost decreased, and the robustness of the approach increased by decoupling the coding sequences from the genome-specific sequences. Alternative assay formats can also be 40 used (e.g. ligation or primer extension followed by ligation).

By ligating the codes on separately, only  $2n$  target-specific 45 assay oligonucleotides are needed for  $n$  targets. For example, assaying 100 different targets at 10,000 spatial locations would require  $2 \times 100$  targeting oligonucleotides and  $2 \times 100$  encoding oligonucleotides, using a combinatorial 50 approach outlined in FIG. 2. The total count of assay oligonucleotides would be only 400 (200 target-specific and 200 encoding), not counting universal primers. In contrast, if the coding oligonucleotides were not decoupled,  $(nxX)$  positional codes)+ $(nxY)$  positional codes) would be needed, or in the above example, 20,000 oligonucleotides, not counting 55 universal primer sequences.

Due to the matrix system of the invention, a large amount 60 of information can be obtained even using five or more positions interrogated for five or more biological molecules. By varying one or the other of these, large amounts of information can be obtained, both in terms of locations and/or specific biological In specific aspects, multiple locations are interrogated for two or more biological molecules. As an example, for each datapoint ~1,000 reads may be sampled, for a total of ~10E9 reads for 10E6 datapoints.

#### Peptide Detection Systems

The assay system of the invention can be used to analyze 65 biological molecules using peptide agents that are associated with the substrate surface in a spatial pattern. Such peptides

may comprise an active region of an enzyme, a binding domain of an immunoglobulin, defined domains of proteins, whole proteins, synthetic peptides, peptides with introduced mutations, etc.

The assay system of the invention allows the identification and spatial location of various forms of peptides, including isoforms and peptides that have undergone post-translational modification. Importantly, certain aspects of the invention allow the identification of active versus non-active forms of such peptides in a sample. This allows the identification of the presence or absence of specific peptide isoforms, and also acts as a proxy for identification of peptide activity in a sample.

In certain aspects of the invention, the binding agents associated with the substrate surfaces of the assay system include substrates for enzymes or proenzymes, e.g., a kinase, a phosphatase, a zymogen, a protease, or a fragment thereof. In certain aspects, the binding agents associated with the substrate surfaces are phosphorylation substrates used to detect proteins involved with one or more signal transduction pathways, e.g., a kinase or a phosphatase. In another specific aspect of the invention, the binding agents are specific protease substrates that associate only with individual or classes of proteases. In other aspects, the binding agents on the substrate surface are different processed forms, isoforms and/or domains of an enzyme.

#### Reagent Delivery

The reagent delivery system of the invention can be any system that allows the delivery of reagents to discrete portions of the array in order to keep the integrity of the defined spatial patterns of the encoding scheme. Such discrete delivery can be achieved in a number of different ways.

In one exemplary aspect, the reagent delivery system can be a flow-based system. The flow-based systems for reagent delivery in the present invention can include one or more pumps, valves, fluid reservoirs, channels, and/or reagent storage cells. Such a reagent delivery system is configured to move fluid in contact with a discrete section of the substrate surface. Movement of the reagents can be driven through a fluid by a pump disposed, for example, downstream of the fluid reagents. The pump can drive each fluid reagent to (and past) the reaction compartment. Alternatively, the reagents may be driven through the fluid by gravity.

US Appln Nos. 20070166725 and 20050239192 disclose certain general-purpose fluidics tools that can be used with the assay systems of the invention. These allow the precise manipulation of gases, liquids and solids to accomplish very complex analytical manipulations with relatively simple hardware.

In a more specific example, one or more flow-cells can be attached to the substrate from above. The flow-cell can include inlet and outlet tubes connected thereto and optionally an external pump can be used to deliver the sample or reagents to the flow-cell and across the substrate. The flow cell is configured to deliver reagents only to certain portions of the array, restricting the amount and type of reagent delivered to any specific section of the array.

In another aspect, a microfluidic system can be integrated into the substrate or externally attached on top of the substrate. Microfluidic passages for holding and carrying fluid can be formed on and/or above the planar substrate by a fluidics layer abutted to the substrate. Fluid reagents can be selected according to selective opening and closing of valves disposed between reagent reservoirs.

Pumps generally include any mechanism for moving fluid and/or reagents disposed in fluid. In some examples, the pump can be configured to move fluid and/or reagents through passages with small volumes (i.e., microfluidic structures). The pump can operate mechanically by exerting a positive or negative pressure on fluid and/or on a structure carrying fluid, electrically by appropriate application of an electric field(s), or both, among others. Exemplary mechanical pumps may include syringe pumps, peristaltic pumps, rotary pumps, pressurized gas, pipettors, etc. The mechanical pumps may be micromachined, molded, etc. Exemplary electrical pumps can include electrodes and may operate by electrophoresis, electroendoosmosis, electrocapillarity, dielectrophoresis (including traveling wave forms thereof), and/or the like.

Valves generally include any mechanism for regulating the passage of fluid through a channel. The valves can include, for example, deformable members that can be selectively deformed to partially or completely close a channel, a movable projection that can be selectively extended into the channel to partially or completely block the channel, an electrocapillary structure, and/or the like.

In yet another aspect, an open gasket can be attached to the top of the substrate and the sample and reagents can be injected into the gasket. Suitable gasket materials include, but are not limited to, neoprene, nitrile, and silicone rubber. Alternatively, a watertight reaction chamber formed by a gasket sandwiched between the substrate and a chemically inert, water resistant material such as, but not limited to, black-anodized aluminum, thermoplastics (e.g., polystyrene, polycarbonate, etc), glass, etc.

In a specific aspect of the present invention, the delivery system can compose a microcircuit arrangement including an imager, such as a CCD or IGFET-based (e.g., CMOS-based) imager and an ultrasonic sprayer for reagent delivery such as described in US Appln No. 20090197326, which is incorporated herein by reference.

In yet another aspect of the invention, the reagent delivery system controls the delivery of reagents to specific patterns on a substrate surface using semiconductor techniques such as masking and spraying. Specific areas of a substrate surface can be protected from exposure to reagents through use of a mask to protect specific areas from exposure. The reagents may be introduced to the substrate using conventional techniques such as spraying or fluid flow. The use of the masked substrate delivery results in a patterned delivery scheme on the substrate surface.

In a preferred aspect of the invention, the reagent delivery instrumentation is based on inkjet printing technology. There are a variety of different ink-jetting mechanisms (e.g., thermal, piezoelectric) and compatibility has been shown with aqueous and organic ink formulations. Sets of independently actuated nozzles can be used to deliver multiple reagents at the same time, and very high resolutions can be achieved.

#### Software for Use in the Assay System

In order to target specific sites of interest, an informative image of the biological section to be analyzed can be used to assist in the reagent delivery methods and associated encoding scheme. Sample regions can be identified using image processing (e.g., images of cell types differentiated by immunohistochemistry or other staining chemistries) integrated with the other features of the assay system. In some aspects, software is used to automatically translate this information into a reagent delivery pattern. A mechanism to

register and align very precisely the biological sample in a targeting system is thus a preferred component of the assay systems of the invention. Mechanisms such as the use of fiducial markers on slides and other very accurate physical positioning systems can be adapted to this purpose.

Additional software components will also be key components that will be part of a complete assay system. The invention thus preferably comprises a complete suite of software tailored to the assay system. Optionally, oligonucleotide design software will be customized for the specific assay to be run, and may be integrated as a part of the system. Also optionally, algorithms and software for data analysis may be integrated to assist in determination of results of the assays. This can be especially useful, as the type of dataset that will be generated will be novel, particularly as a consequence of scale. The ability to provide algorithms and software tools that are specifically designed for analysis of spatially-associated data for significant patterns, including pattern-analysis software and visualization tools, is a novel feature that will enhance the value of the data generated by the assay systems.

In certain aspects, the assay system will comprise processes for making and carrying out quality control of reagents, e.g., the integrity and sequence fidelity of oligonucleotide pools. In particular, reagents will need to be formulated for compatibility with the reagent delivery instrumentation. Factors such as volatility, stability at key temperatures, and chemical compatibility can be optimized by those skilled in the art upon reading the present disclosure.

#### Applications of Assay System

It will be apparent to one skilled in the art upon reading the present disclosure that there are numerous very important areas of biological research, diagnostics, and drug development that will benefit from a high throughput means to simultaneously measure the presence or absence and spatial location of a biological molecule in a sample. For example, this technology combining the ability to analyze semi-quantitatively the expression of many different genes with the ability to image the spatial organization of expression across many cells in a tissue is enabling for many different areas of basic research. The following are exemplary uses and are by no means meant to be limiting in scope.

In one example, 3-dimensional patterns of expression can be determined by analyzing a series of tissue sections, in a manner analogous to image reconstruction in CT scanning. This can be used to measure changes in gene expression in disease pathology, e.g., in cancerous tissue and/or a tissue upon injury, inflammation or infection. With the assay systems of the invention, more detailed information on gene expression and protein localization in complex tissues can be obtained. This may lead to new insights into the function and regulation both in normal and diseased states, and is likely to provide new hypotheses that can be tested. For example, a system of the invention may enable some of the insights gained from many individual studies and larger programs like ENCODE (Birney et al., 2007) and modENCODE to be integrated at the tissue level. The assay systems will also aid in computational efforts to model interacting networks of gene expression in the field of systems biology.

The assay systems also provide a novel approach that enables the analysis of somatic variation, e.g., somatic mutations in cancer or variability in response to infectious organisms. For example, tumors are typically highly hetero-

geneous, containing cancer cells as well as genetically normal cells in an abnormal local environment. Cancer cells undergo mutation and selection, and in this process it is not unusual for local clones to develop. Identifying relatively rare somatic mutations in the context of tumors may enable the study of the role of key mutations in the selection of clonal variants. Transcriptional patterns associated with angiogenesis, inflammation, or other cancer related processes in both cancer and genetically normal cells can be analyzed for insights into cancer biology and assist in the development of new therapeutic agents for the treatment of cancers.

In another example, different people have varying susceptibility to infectious organisms, and much of this may be to underlying genetic differences in individuals and/or populations. Identifying these differences will aid in an understanding of the underlying disease pathologies and assist in the development of vaccines or therapeutics to prevent or ameliorate these disease states.

Importantly, in addition to providing spatially associated information, the technology of the invention will allow a great increase in the sensitivity of detecting rare mutations. The reason is that signal to noise can be dramatically increased because the approach of the invention assays a small location in any given reaction. In a typical assay for rare mutations in a mixed sample, the sample is treated in bulk, i.e. nucleic acids are extracted from many cells into a single pool. Thus, if a mutation is present in 1 cell in 10,000, it must be detected against a background of normal DNA from ~10,000 cells. In contrast, with the systems of the invention many cells can be analyzed, but individual cells or small groups of cells would be identified by the spatial coding system. Therefore, the background can be reduced by orders of magnitude, greatly increasing sensitivity. Furthermore, the spatial organization of mutant cells can be observed. This may be particularly important in detecting key mutations in tissue sections in cancer. Already, molecular histological analyses are yielding insights into cancer biology and may have potential for use in diagnostics (Choe et al., 2003). The technology of the invention promises to greatly increase the power of such approaches.

#### EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present invention, and are not intended to limit the scope of what the inventor regards as his invention, nor are they intended to represent or imply that the experiments below are all of or the only experiments performed. It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees centigrade, and pressure is at or near atmospheric.

##### Example 1: Initial Proof of Concept of Encoding Scheme

As an initial proof of concept, a model system is developed using a microarray to demonstrate a working single-

31

plex assay. The basic design validates the concept of the assay, and establishes a working assay prior to addressing issues related to the analysis of a more complicated biological sample. Conventional sequencing is used as a readout for this proof of concept.

A microarray is used as a proxy for a tissue section. The target sequences of the microarray are fully specified, so that the composition of the targets are known and can be varied systematically. Synthetic oligonucleotide templates are attached to a glass slide via a 5' amino modification. Each slide has a single oligonucleotide template sequence, and the assays that are carried out may employ either ligation, or extension followed by ligation as this may be useful in determining certain polymorphisms.

Once the *in situ* part of the assay is complete, the reaction products are eluted and analyzed by qPCR to determine presence or absence of a product and estimate yield, and by conventional sequencing to determine the structure of the assay products. The single plex assays that are tested include appropriate positive and negative controls, and a single nucleotide variant (SNV) to check ability to discriminate single base variants.

#### Example 2: Scalability

The complexity of the assay system is increased to establish scalability of the assay for use in high throughput studies. Scalability of both the spatial encoding and assay systems is demonstrated by carrying out a 24-plex×24-site assay using a microarray model system.

The amount of biological target, here a DNA target sequence, at each assay location is systematically varied on microarray substrate. For example, in a microarray with 50 micron spot size (center to center), a 1 mm<sup>2</sup> area contains ~400 spots. The region around each site is optionally occupied by a region that is devoid of these spots to allow individual resolvability of the target sequences. Alternatively, the spots may be clustered, with two or more directly adjacent spots surrounded by or adjacent to a region that is devoid of target sequences.

In order to demonstrate that spatial encoding is accurate, the sites comprise different target compositions to show that the assay readout matches the expected composition of each site. With 24 target sequences, a simple digital pattern is made with each site having a different set of 12 targets present and 12 targets absent, to make a binary code (0=absent, 1=present). The assay readout is then determined to show that the detected regions match the expected signal after spatial decoding. In this particular example, the code space is large enough ( $2^{24}$ ) so that even a few errors would not result in different codes being mixed up. Moreover, this design allows identification of errors and allows an estimation not only of accuracy of spatial encoding but also of accuracy calling the presence or absence of target sequences.

In an exemplary aspect, a 4×4 arrangement of 16 sequences is used for the array configuration. A white square indicates that the sequence is absent and a black square that it is present, i.e. 8 of the 16 possible sequences are present in this sample. In a different sample, a different pattern of absent and present sequences can be constructed. In this way, unique patterns are associated with spatial locations so that the accuracy of spatial encoding can be measured.

The ability to detect quantitative differences is evaluated by generating dose-response curves for each of the 24 assays that are carried out at each site in a 24-site assay. This allows estimation of the limit of detection, dynamic range, and power to detect a given fold-change across the range.

32

In one aspect, a latin square design is used to represent individual targets at different ratios by varying the number of features for each target. In other words, with multiple spots in a site, the number of spots allocated to each of the 24 target sequences can be varied and each of the 24 sites can have a different composition. A 1×3 inch microarray is sufficiently large to permit multiple replicates. This larger set of 24 sequences will require deconvolution, and this is accomplished using high throughput techniques such as next-generation sequencing technologies (e.g., SOLiD™ technology (Life Technologies, Inc., Carlsbad, CA) or Genome Analyzer (Illumina, Inc., San Diego, CA)). The use of the 24-plex assay demonstrates both the accuracy of spatial encoding and decoding, and the quantitative response of the assay system.

#### Example 3: Adaptation of the Assay to Preserved Samples

Genomic DNA is assayed as a proof of concept for assaying RNA, as it provides a way to establish a single-copy reference signal. Once a working assay is developed for FFPE samples, it is adapted to an RNA assay. To this end, assay oligonucleotide concentrations are assayed to ensure compatibility with high multiplexing. Assuming a cell diameter of 10 microns, and delivery of a 10 micron diameter reagent droplet to an individual cell, the volume of the droplet will be ~500 µl and can contain ~3×10<sup>11</sup> molecules at a 1 µM concentration. To assay 1,000 target sequences in 10,000 cells, ~2,000 targeting oligonucleotides would be required in a droplet. Therefore, each droplet could contain ~160 million copies of each assay oligo, a vast excess over the few thousand target sequences in a cell.

The handling of small absolute numbers of product molecules generated from very small or compromised samples are enhanced to counter the issue of low recovery efficiency; that is, elution is efficient and losses resulting from adsorption of molecules to surfaces are prevented. An approach to addressing the latter issue is to include a carrier material, such as glycogen or carrier nucleic acids.

#### Example 4: Adapting the Assay to a Biological Sample

A control RNA template is immobilized to a solid support in order to create an artificial system. The assay is performed using T4 DNA ligase, which can repair nicks in DNA/RNA hybrids. Assays are carried out on matched slides, or different sections of the same slide, where in one case gDNA is assayed and in the other RNA is assayed. When assaying gDNA the slide can be pretreated with RNase, and when assaying RNA the slide is pretreated with DNase. Results of the assay are confirmed by extracting gDNA or RNA and quantitating the relative amounts by qPCR or RT-qPCR respectively.

In order to make the tissue section RNA assays as informative as possible, pre-existing information on expression levels in specific tissues to target transcripts across a range of abundances are used in the assay design. Both high abundance transcripts, as well as some medium and low abundance transcripts, are targeted to enable an initial assessment of the quantitative performance characteristics of the assay.

The preceding merely illustrates the principles of the invention. It will be appreciated that those skilled in the art will be able to devise various arrangements which, although not explicitly described or shown herein, embody the prin-

**33**

ciples of the invention and are included within its spirit and scope. Furthermore, all examples and conditional language recited herein are principally intended to aid the reader in understanding the principles of the invention and the concepts contributed by the inventors to furthering the art, and are to be construed as being without limitation to such specifically recited examples and conditions. Moreover, all statements herein reciting principles, aspects, and embodiments of the invention as well as specific examples thereof, are intended to encompass both structural and functional equivalents thereof. Additionally, it is intended that such equivalents include both currently known equivalents and equivalents developed in the future, i.e., any elements developed that perform the same function, regardless of structure. The scope of the present invention, therefore, is not intended to be limited to the exemplary embodiments shown and described herein. Rather, the scope and spirit of present invention is embodied by the appended claims. In the claims that follow, unless the term "means" is used, none of the features or elements recited therein should be construed as means-plus-function limitations pursuant to 35 U.S.C. § 112, ¶6.

The invention claimed is:

**1. A method comprising:**

- (a) providing a substrate comprising a plurality of nucleic acid molecules each having (i) a first sequence configured to bind a target biological molecule in a tissue section, and (ii) a second sequence corresponding to a location on the substrate;
- (b) contacting the tissue section with the substrate;
- (c) capturing the target biological molecule from the tissue section onto a nucleic acid molecule of the plurality of nucleic acid molecules using the first sequence; and
- (d) generating a nucleic acid molecule comprising (i) a sequence of the captured target biological molecule, or a complement thereof, and (ii) the second sequence corresponding to the location on the substrate, or a complement thereof.

**2. The method of claim 1, wherein the tissue section is a fresh-frozen or formalin-fixed paraffin-embedded tissue sample.**

**3. The method of claim 1, wherein the target biological molecule is an RNA molecule.**

**34**

**4. The method of claim 3, wherein the RNA molecule is an mRNA molecule.**

**5. The method of claim 1, wherein the target biological molecule is a DNA molecule.**

**6. The method of claim 1, wherein (c) comprises hybridizing the target biological molecule to the first sequence of the nucleic acid molecule.**

**7. The method of claim 1, wherein (c) comprises coupling the target biological molecule to the nucleic acid molecule.**

**8. The method of claim 7, wherein the coupling comprises ligation.**

**9. The method of claim 8, wherein the ligation comprises enzymatic ligation.**

**10. The method of claim 1, wherein (d) comprises extending the nucleic acid molecule using the target biological molecule as a template.**

**11. The method of claim 10, wherein the extending is performed by a polymerase.**

**12. The method of claim 1, wherein the substrate is a flow cell.**

**13. The method of claim 1, wherein the plurality of nucleic acid molecules further comprises a universal priming site.**

**14. The method of claim 1, further comprising staining the tissue section.**

**15. The method of claim 14, wherein the staining is selected from immunohistochemistry or hematoxylin and eosin staining.**

**16. The method of claim 14, further comprising obtaining an image of the stained tissue section.**

**17. The method of claim 1, further comprising amplifying the generated nucleic acid molecule.**

**18. The method of claim 17, further comprising sequencing all or a portion of the amplified, generated nucleic acid molecule.**

**19. The method of claim 1, further comprising determining the location on the substrate where the target biological molecule was captured.**

**20. The method of claim 19, wherein the method further comprises mapping the location of the target biological molecule to a location in the tissue section.**

\* \* \* \* \*