

### (19) United States

### (12) Patent Application Publication (10) Pub. No.: US 2025/0264475 A1 zeng et al.

#### Aug. 21, 2025 (43) **Pub. Date:**

#### (54) METHODS FOR IDENTIFYING BINDING PARTNERS OF PROGRANULIN

(71) Applicant: Janssen Pharmaceutica NV, Beerse

(72) Inventors: Wei-zheng zeng, San Diego, CA (US); Joseph Chang, La Jolla, CA (US); Pascal Bonaventure, La Jolla, CA (US); Changlu Liu, San Diego, CA (US); David S Bredt, La Jolla, CA

(US)

(21) Appl. No.: 18/288,025 Apr. 22, 2022 (22) PCT Filed:

(86) PCT No.: PCT/EP2022/060724

§ 371 (c)(1),

Oct. 23, 2023 (2) Date:

#### Related U.S. Application Data

(60) Provisional application No. 63/178,831, filed on Apr. 23, 2021.

#### **Publication Classification**

(51) Int. Cl. (2006.01)G01N 33/68 G01N 1/30 (2006.01)G01N 21/64 (2006.01)G01N 33/58 (2006.01)


(52) U.S. Cl. CPC ...... G01N 33/68 (2013.01); G01N 1/30 (2013.01); G01N 21/6428 (2013.01); G01N 33/582 (2013.01); G01N 2021/6439 (2013.01)

#### (57)ABSTRACT

Disclosed herein are screening methods for identifying cell surface receptors for progranulin. Also disclosed herein are screening methods for identifying intracellular proteins bind to progranulin.

Specification includes a Sequence Listing.

HEK293T cells




~3000 CNS ~6000 Broad cDNA libraries




Protein expression

**Fixation** Permeabilization 100 nM Flag-GRN **Fixation Immunostaining** 

High content imaging








Figure 2




Figure 3A-3B




Figure 3C




Figure 4

# METHODS FOR IDENTIFYING BINDING PARTNERS OF PROGRANULIN

# CROSS REFERENCE TO RELATED APPLICATION

[0001] This application claims the benefit of U.S. Provisional Application 63/178,831, filed on Apr. 23, 2021, which is incorporated by reference herein in its entirety.

#### FIELD OF INVENTION

[0002] This invention relates to high-throughput screening methods for identifying binding partners of programulin.

# REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

[0003] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Feb. 28, 2022, is named PRD4129WOPCT1\_SL.txt and is 16,384 bytes in size.

#### BACKGROUND OF INVENTION

[0004] Neurodegenerative diseases occur when nervous system cells (neurons) in the brain and spinal cord begin to deteriorate. Today, 5.4 million Americans suffer from Alzheimer's disease; 500,000 from Parkinson's disease; and 50,000-60,000 from frontotemporal dementia (FTD). Because neurodegenerative diseases strike primarily in midto late-life, the incidence is expected to soar as the population ages. By 2030, as many as 1 in 5 Americans will be over the age of 65. Finding treatments and cures for neurodegenerative diseases is a goal of increasing urgency.

[0005] Progranulin (PGRN) is a cysteine-rich, secreted protein encoded by the GRN gene and known to be involved in neurodegeneration and other diseases. A reduction of PGRN levels in the brain correlates with familial forms of frontotemporal lobar degeneration (FTLD). Despite the strong correlation, its biological mechanism that causes neurodegeneration remains unclear. Though there have been proposed cell surface receptors that bind to PGRN, the link between those receptors and neurodegeneration remains murky, suggesting other binding partners may be at play. There is a need for methods to identify those interaction partners in a high-throughput and unbiased manner.

#### BRIEF SUMMARY OF THE INVENTION

[0006] Provided herein is a high-throughput screening method for identifying transmembrane proteins that bind progranulin (PGRN), comprising: a) providing a cDNA collection comprising a plurality of expression vectors, each expression vector encoding a distinct transmembrane protein; b) providing a plate comprising a plurality of wells, wherein one or more wells thereof each contains host cells; c) into each of the one or more wells, introducing a distinct expression vector from said collection, thereby producing distinct transfected cells in each well; d) culturing the transfected cells under a condition to allow expression of a distinct transmembrane protein in each well; e) contacting the transfected cells in each of the one or more wells with a target protein comprising PGRN attached to a detectable tag; f) fixing the transfected cells with a fixing agent after contact with the target protein; g) washing the fixed cells; and h) after washing the fixed cells, determining the presence of the detectable tag in each of the one or more wells, wherein if the detectable tag is determined to be present in a well, the transmembrane protein expressed by the transfected cells in the well is identified as a transmembrane protein that binds PGRN.

[0007] In one embodiment of the high-throughput screening method, the host cells are selected from the group consisting of human embryonic kidney 293T (HEK293T) cell, HEK293F cells, Hela cells, Chinese hamster ovary (CHO) cells, NIH 3T3 cells, MCF-7 cells, Hep G2 cells, baby hamster kidney (BHK) cells, BV-2 (mouse, C57BL/6, brain, microglial) cells, Neuro-2a (N2a) cells and Cos7 cells. [0008] In a further embodiment of the high-throughput screening, the fixing agent is selected from glutaraldehyde, methanol and paraformaldehyde.

[0009] In a yet further embodiment of the high-throughput screening method, the detectable tag is selected from the group consisting of green fluorescent protein (GFP), myc, myc-pyruvate kinase (myc-PK), His6, maltose binding protein (MBP), human influenza virus hemagglutinin (HA) tag, flag tag (FLAG), and glutathione-S-transferase (GST) tag. [0010] In a yet further embodiment of the high-throughput screening method, the target protein is a FLAG tagged PGRN. In one aspect, step h) comprises h1) adding an anti-FLAG antibody that is covalently attached to a fluorescent dye into the wells containing the transfected cells to immunostain the FLAG tagged PGRN, if present, and h2) detecting the fluorescent dye using a fluorometer or an imager in a high-throughput manner. In another aspect, the fluorescent dye is cyanine dye.

[0011] In a yet further embodiment of the high-throughput screening method, the target protein is an HA tagged PGRN. In one aspect, step h) comprises h1) exposing the cells to an anti-HA antibody that is covalently attached to a fluorescent dye to immunostain the HA tagged PGRN, if present, and h2) detecting the fluorescent dye using a fluorometer or an imager. In another aspect, the fluorescent dye is Dylight 650. [0012] In a yet further embodiment of the high-throughput screening method, between steps g) and h), the method further comprises permeabilizing the fixed cells with a permeabilizing agent and washing the permeabilized cells. [0013] In a yet further embodiment of the high-throughput screening method, the permeabilizing agent is selected from the group consisting of Saponin, Triton X-100, lysolecithin, n-Octyl-B-D-glucopyranoside, and Tween 20.

[0014] Further provided herein is a method for identifying intracellular proteins that bind PGRN, comprising: a) introducing into host cells an expression vector encoding a heterologous protein, thereby producing transfected cells; b) culturing the transfected cells under a condition to allow expression of the heterologous protein; c) permeabilizing the transfected cells with a permeabilizing agent; d) contacting the permeabilized cells with a target protein comprising PGRN attached to a detectable tag; e) fixing the transfected cells with a fixing agent after contact with the target protein; f) washing the fixed cells; and g) after washing the fixed cells, determining the presence of the detectable tag on the washed cells, wherein if the detectable tag is determined to be present, the heterologous protein expressed by the cells is identified as an intracellular protein that binds PGRN.

[0015] In one embodiment of the method, the host cells are selected from the group consisting of human embryonic kidney 293T (HEK293T) cell, HEK293F cells, HeLa cells,

Chinese hamster ovary (CHO) cells, NIH 3T3 cells, MCF-7 cells, Hep G2 cells, baby hamster kidney (BHK) cells, BV-2 (mouse, C57BL/6, brain, microglial) cells, Neuro-2a (N2a) cells and Cos7 cells.

[0016] In a further embodiment of the method, the detectable tag is selected from the group consisting of green fluorescent protein (GFP), myc, myc-pyruvate kinase (myc-PK), His6, maltose binding protein (MBP), human influenza hemagglutinin (HA) tag, flag tag (FLAG), and glutathione-S-transferase (GST) tag.

[0017] In a yet further embodiment of the method, the target protein is a FLAG tagged PGRN. In one aspect, step g) comprises g1) exposing the cells to an anti-FLAG antibody that is covalently attached to a fluorescent dye to immunostain the FLAG tagged PGRN, if present, and g2) detecting the fluorescent dye using a fluorometer or an imager. In another aspect, the fluorescent dye is cyanine dye. [0018] In a yet further embodiment of the high-throughput screening method, the target protein is an HA tagged PGRN. In one aspect, step g) comprises g1) exposing the cells to an anti-HA antibody that is covalently attached to a fluorescent dye to immunostain the HA tagged PGRN, if present, and g2) detecting the fluorescent dye using a fluorometer or an imager. In another aspect, the fluorescent dye is Dylight 650. [0019] In a yet further embodiment of the method, between steps b) and c), the method further comprises, fixing the cultured cells with the fixing agent and washing the fixed cells.

[0020] In a yet further embodiment of the method, the fixing agent is selected from glutaraldehyde, methanol and paraformaldehyde.

[0021] In a yet further embodiment of the method, between steps f) and g), the method further comprises permeabilizing the fixed cells with a permeabilizing agent and washing the permeabilized cells.

[0022] In a yet further embodiment of the method, the permeabilizing agent is selected from the group consisting of Saponin, Triton X-100, lysolecithin, n-Octyl-B-D-glucopyranoside, and Tween 20.

[0023] Yet further provided herein is a high-throughput screening method for identifying intracellular proteins that bind PGRN, comprising: a) providing a cDNA collection comprising a plurality of expression vectors, each expression vector encoding a distinct heterologous protein; b) providing a plate comprising a plurality of wells, wherein one or more wells thereof each contains host cells; c) into each of the one or more wells, introducing a distinct expression vector from said collection, thereby producing distinct transfected cells in each well; d) culturing the transfected cells under a condition to allow expression of a distinct heterologous protein in each well; e) permeabilizing the transfected cells with a permeabilizing agent; f) contacting the permeabilized cells in each of the one or more wells with a target protein comprising PGRN attached to a detectable tag; g) fixing the transfected cells with a fixing agent after contact with the target protein; h) washing the fixed cells; and i) after washing the fixed cells, determining the presence of the detectable tag in each of the one or more wells, wherein if the detectable tag is determined to be present in a well, the heterologous protein expressed by the transfected cells in the well is identified as a intracellular protein that binds PGRN.

[0024] In one embodiment of the high-throughput screening method, the host cells are selected from the group

consisting of human embryonic kidney 293T (HEK293T) cell, HEK293F cells, Hela cells, Chinese hamster ovary (CHO) cells, NIH 3T3 cells, MCF-7 cells, Hep G2 cells, baby hamster kidney (BHK) cells, BV-2 (mouse, C57BL/6, brain, microglial) cells, Neuro-2a (N2a) cells and Cos7 cells. [0025] In a further embodiment of the high-throughput screening method, the detectable tag is selected from the group consisting of green fluorescent protein (GFP), myc, myc-pyruvate kinase (myc-PK), His6, maltose binding protein (MBP), human influenza hemagglutinin (HA) tag, flag tag (FLAG), and glutathione-S-transferase (GST) tag.

[0026] In a yet further embodiment of the high-throughput screening method, the target protein is a FLAG tagged PGRN. In one aspect, step i) comprises i1) adding an anti-FLAG antibody that is covalently attached to a fluorescent dye into the wells containing the transfected cells to immunostain the FLAG tagged PGRN, if present, and i2) detecting the fluorescent dye using a fluorometer or an imager in a high-throughput manner. In another aspect, the fluorescent dye is cyanine dye.

[0027] In a yet further embodiment of the high-throughput screening method, the target protein is an HA tagged PGRN. In one aspect, step i) comprises i1) exposing the cells to an anti-HA antibody that is covalently attached to a fluorescent dye to immunostain the HA tagged PGRN, if present, and i2) detecting the fluorescent dye using a fluorometer or an imager. In another aspect, the fluorescent dye is Dylight 650. [0028] In a yet further embodiment of the high-throughput screening method, between steps d) and e), the method further comprises, fixing the cultured cells with the fixing agent and washing the fixed cells.

[0029] In a yet further embodiment of the high-throughput screening method, the fixing agent is selected from glutar-aldehyde, methanol and paraformaldehyde.

[0030] In a yet further embodiment of the high-throughput screening method, between steps h) and i), the method further comprises permeabilizing the fixed cells with a permeabilizing agent and washing the permeabilized cells. [0031] In a yet further embodiment of the high-throughput screening method, the permeabilizing agent is selected from the group consisting of Saponin, Triton X-100, lysolecithin, n-Octyl-B-D-glucopyranoside, and Tween 20.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0032] FIG. 1 is a schematic depicting a screening strategy for detection of intracellular proteins that bind PGRN.

[0033] FIG. 2 shows the Cy3 intensity of cells transfected with pcDNA3 vector (negative control), cells transfected with an expression vector carrying coding sequence of sortilin (positive control), and cells transfected with an expression plasmid carrying the coding sequence of a hit intracellular protein.

[0034] FIGS. 3A and 3C shows the Cy3 intensity or Dylight 650 intensity of the transfected cells (negative control, positive control, and Hit #1) that were exposed to FLAG tagged PGRN or HA tagged PGRN, followed by immunostaining with anti-Flag M2-Cy3 antibody or Dylight 650 anti-HA antibody, respectively. FIG. 3B shows the Cy3 signal and Dylight 650 signal for cells transfected with Hit #1 plasmid but not exposed to tagged PGRN prior to immunostaining.

[0035] FIG. 4 shows the Cy3 intensity or Dylight 650 intensity of the transfected cells (negative control, positive

control, and Hit #1) that were not permeabilized prior to exposure to FLAG tagged PGRN or HA tagged PGRN and immunostaining.

# DETAILED DESCRIPTION OF THE INVENTION

[0036] Various publications, articles and patents are cited or described in the background and throughout the specification; each of these references is herein incorporated by reference in its entirety. Discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is for the purpose of providing context for the invention. Such discussion is not an admission that any or all of these matters form part of the prior art with respect to any inventions disclosed or claimed. [0037] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention pertains. Otherwise, certain terms used herein have the meanings as set forth in the specification.

[0038] It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise.

[0039] Unless otherwise stated, any numerical values, such as a concentration or a concentration range described herein, are to be understood as being modified in all instances by the term "about." Thus, a numerical value typically includes ±10% of the recited value. For example, a concentration of 1 mg/ml includes 0.9 mg/mL to 1.1 mg/mL. Likewise, a concentration range of 1% to 10% (w/v) includes 0.9% (w/v) to 11% (w/v). As used herein, the use of a numerical range expressly includes all possible subranges, all individual numerical values within that range, including integers within such ranges and fractions of the values unless the context clearly indicates otherwise.

[0040] Unless otherwise indicated, the term "at least" preceding a series of elements is to be understood to refer to every element in the series. Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the invention.

[0041] As used herein, the terms "comprises," "comprising," "includes," "including," "has," "having," "contains" or "containing," or any other variation thereof, will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers and are intended to be non-exclusive or openended. For example, a composition, a mixture, a process, a method, an article, or an apparatus that comprises a list of elements is not necessarily limited to only those elements but can include other elements not expressly listed or inherent to such composition, mixture, process, method, article, or apparatus. Further, unless expressly stated to the contrary, "or" refers to an inclusive or and not to an exclusive or. For example, a condition A or B is satisfied by any one of the following: A is true (or present) and B is false (or not present), A is false (or not present) and B is true (or present), and both A and B are true (or present).

[0042] As used herein, the conjunctive term "and/or" between multiple recited elements is understood as encompassing both individual and combined options. For instance, where two elements are conjoined by "and/or," a first option

refers to the applicability of the first element without the second. A second option refers to the applicability of the second element without the first. A third option refers to the applicability of the first and second elements together. Any one of these options is understood to fall within the meaning, and therefore satisfy the requirement of the term "and/or" as used herein. Concurrent applicability of more than one of the options is also understood to fall within the meaning, and therefore satisfy the requirement of the term "and/or."

[0043] As used herein, the term "consists of," or variations such as "consist of" or "consisting of," as used throughout the specification and claims, indicate the inclusion of any recited integer or group of integers, but that no additional integer or group of integers can be added to the specified method, structure, or composition.

[0044] As used herein, the term "consists essentially of," or variations such as "consist essentially of" or "consisting essentially of," as used throughout the specification and claims, indicate the inclusion of any recited integer or group of integers, and the optional inclusion of any recited integer or group of integers that do not materially change the basic or novel properties of the specified method, structure or composition. See M.P.E.P. § 2 111.03.

[0045] It should also be understood that the terms "about," "approximately," "generally," "substantially," and like terms, used herein when referring to a dimension or characteristic of a component of the preferred invention, indicate that the described dimension/characteristic is not a strict boundary or parameter and does not exclude minor variations therefrom that are functionally the same or similar, as would be understood by one having ordinary skill in the art. At a minimum, such references that include a numerical parameter would include variations that, using mathematical and industrial principles accepted in the art (e.g., rounding, measurement or other systematic errors, manufacturing tolerances, etc.), would not vary the least significant digit.

[0046] As used herein, the term "polynucleotide," synonymously referred to as "nucleic acid molecule," "nucleotides" or "nucleic acids," refers to any polyribonucleotide or polydeoxyribonucleotide, which can be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that can be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short nucleic acid chains, often referred to as oligonucleotides.

[0047] As used herein, the terms "peptide," "polypeptide," or "protein" can refer to a molecule comprised of amino acids and can be recognized as a protein by those of skill in

the art. The conventional one-letter or three-letter code for amino acid residues is used herein. The terms "peptide," "polypeptide," and "protein" can be used interchangeably herein to refer to polymers of amino acids of any length. The polymer can be linear or branched, it can comprise modified amino acids, and it can be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification, such as conjugation with a labeling component. Also included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, etc.), as well as other modifications known in the art.

[0048] The peptide sequences described herein are written according to the usual convention whereby the N-terminal region of the peptide is on the left and the C-terminal region is on the right. Although isomeric forms of the amino acids are known, it is the L-form of the amino acid that is represented unless otherwise expressly indicated.

Identification of Transmembrane Proteins that Bind PGRN [0049] In one aspect, the invention disclosed herein are methods for screening for transmembrane proteins that bind to progranulin (PGRN) in a high throughput (HTP) manner. [0050] In particular, a target protein comprised of PGRN fused with a detectable tag, a cDNA library or collection, and host cells are utilized in the screening methods.

[0051] PGRN is a cysteine-rich, secreted protein encoded by the GRN gene (NCBI Accession #NM\_002087.3). In forming the target protein, PGRN is fused with a detectable tag peptide. Such tag peptides are well known in the art and include, without limitation, green fluorescent protein (GFP), myc, myc-pyruvate kinase (myc-PK), polyhistidine (His6) tag, maltose binding protein (MBP), human influenza virus hemagglutinin (HA) tag (YPYDVPDYA, SEQ ID NO: 1), flag tag (FLAG) (DYKDDDDK, SEQ ID NO: 2), and glutathione-S-transferase (GST) tag. However, the invention should in no way be construed to be limited to the abovelisted tags. Rather, any peptide or polypeptide which may function in a manner substantially similar to these tags should be construed to be included in the present invention. [0052] The cDNA library comprises a plurality of expression vectors each encoding a distinct transmembrane protein. In one embodiment, each of the expression vectors encodes a human transmembrane protein. The term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular doublestranded DNA loop into which additional DNA segments can be inserted. Another type of vector is a viral vector wherein additional DNA segments can be inserted. Expression vectors are those vectors capable of directing the expression of genes to which they are operably linked. The expression vectors used herein comprise a nucleic acid encoding a protein sequence in a form suitable for expression of the nucleic acid in a host cell. Thus, the expression vectors can include one or more regulatory sequences, such as a promoter, selected on the basis of the host cells to be used for expression, operably linked to the nucleic acid sequence to be expressed. When used in reference to a expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner allowing for expression of the nucleotide sequence (e.g., in an in vitro transcription/ translation system or in a host cell when the vector is introduced into the host cell). It will be appreciated by those of ordinary skill in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed and the level of expression of protein desired as well as the intended use of the vector. Exemplary cDNA libraries suitable herein may include, without limitation, Ultimate® ORF Clones from ThermoFisher, Human cDNA Clone Sets for Pathway Study or HTP Screening from OriGene, etc.

[0053] Any suitable host cells may be used herein. In one embodiment, the host cells used herein are mammalian cells. Suitable mammalian cells may be selected from human embryonic kidney 293T (HEK293T) cell, HEK293F cells, Hela cells, Chinese hamster ovary (CHO) cells, NIH 3T3 cells, MCF-7 cells, Hep G2 cells, baby hamster kidney (BHK) cells, BV-2 (mouse, C57BL/6, brain, microglial) cells, Neuro-2a (N2a) cells and Cos7 cells.

[0054] The high-throughput screening method for identifying transmembrane proteins that bind progranulin (PGRN) uses multi-well plates, such as 96-well, 384-well, or 1536well plates, and wherein the host cells in culture media are placed in at least one well of the plate. In a high-throughput manner, expression vectors from the cDNA library are introduced into each of the wells containing host cells and the multi-well plate is maintained under suitable culturing conditions to allow transfection and expression of the transmembrane proteins encoded by the expression vectors. The culture media used herein are not particularly limited. Exemplary culture media include, without limitation, Dulbecco's Modified Eagle Medium (DMEM), Roswell Park Memorial Institute 1640 Medium (RPMI 1640), Ham's F-12 Nutrient Mixture (F-12), Dulbecco's Modified Eagle Medium: Nutrient Mixture F-12 (DMEM/F-12) and Minimum Essential Medium (MEM). Suitable culturing conditions may be with 5% CO<sub>2</sub> and at about 30-37° C. for 1-3 days.

[0055] After the culturing media is removed, in a highthroughput manner, the target protein (i.e., PGRN fused with detectable tag) is dispensed into each of the wells containing the transfected host cells and the multi-well plate is maintained under a suitable condition to allow potential bond between the transmembrane proteins and the target protein. In one embodiment, the transfected cells are maintained in contact with the target protein at about 20-37° C. for 1-3 hours. Then, fixing agent is added into each well of the multi-well plate to fix the transfected cells followed by washing with washing buffer to remove the un-bond target protein. Examples of suitable fixing agents include glutaraldehyde, methanol and paraformaldehyde. The range of concentrations of these fixatives is generally 0.01 to 8% for paraformaldehyde, and 0.001 to 2% for glutaraldehyde. Combinations of these two reagents may also be employed. Preferred concentrations of paraformaldehyde are 0.1 to 6% with the most preferred concentrations of 1 to 5%. Preferred concentrations of glutaraldehyde are 0.01 to 1% with the most preferred concentrations of 0.05 to 0.5%. The particular concentrations of fixatives are adjusted to optimally fix a particular cell type. Examples of suitable washing buffer include HEPES Balanced Salt Solution (HBSS), Phosphate-Buffered Saline (PBS), Tris-Buffered Saline (TBS), etc.

[0056] Finally, the plate is screened for the presence of the detectable tag in each well, wherein if the detectable tag is determined to be present in a well, the transmembrane

protein expressed by the transfected cells contained in the well is identified as a transmembrane protein binds PGRN. Preferably the screening for the detectable tag is an automated process.

[0057] Optionally, prior to the final screening for the detectable tag, the cells in each well are permeabilized with a permeabilization agent, followed by washing with washing buffer. Permeabilization agents generally are detergent or detergent-like compounds and may include, without limitation, Saponin, Triton X-100, lysolecithin, n-Octyl-B-D-glucopyranoside, and Tween 20.

[0058] Starting from placing host cells in the one or more wells until screening for the presence of the detectable tag, each step of the process may be performed in a high-throughput manner, such as by a robotic arm.

[0059] In one embodiment, the detectable tag is FLAG tag and the target protein is a N-terminal FLAG tagged PGRN. The final screening includes, adding an anti-FLAG antibody that is covalently attached to a fluorescent dye (such as a cyanine dye, e.g., Cy3 or Cy5) into the wells containing the transfected cells to immunostain the FLAG tagged PGRN, if present, and detecting the fluorescent dye using, for example, fluorometers or imagers.

N-terminal FLAG tagged PGRN: SEO ID NO: 3 ATGTGGACTCTGGTCTCATGGGTGGCTCTGACTGCTGGACTGGTG GCTGGAACCGACTACAAGGACGACGACGACAAACTCGCTGCCCTG ACGGCCAGTTCTGCGACAAATGGCCCACAACACTGAGCAGGCATC TGGGTGGCCCTGCCAGGTTGATGCCCACTGCTCTGCCGGCCACT CCTGCATCTTTACCGTCTCAGGGACTTCCAGTTGCTGCCCCTTCC CAGAGGCCGTGGCATGCGGGGATGGCCATCACTGCTGCCCACGGG GCTTCCACTGCAGTGCAGACGGGCGATCCTGCTTCCAAAGATCAG GTAACAACTCCGTGGGTGCCATCCAGTGCCCTGATAGTCAGTTCG AATGCCCGGACTTCTCCACGTGCTGTTTTTTGGTCGATGGCTCCT GGGGGTGCTGCCCCATGCCCCAGGCTTCCTGCTGTGAAGACAGGG  $\tt TGCACTGCTGTCCGCACGGTGCCTTCTGCGACCTGGTTCACACCC$ GCTGCATCACACCCACGGGCACCCCACCCCTGGCAAAGAAGCTCC CTGCCCAGAGGACTAACAGGGCAGTGGCCTTGTCCAGCTCGGTCA  ${\tt TGTGTCCGGACGCACGGTCCCGGTGCCCTGATGGTTCTACCTGCT}$ GTGAGCTGCCCAGTGGGAAGTATGGCTGCTGCCCAATGCCCAACG  $\tt CCACCTGCTGCTCCGATCACCTGCACTGCTGCCCCCAAGACACTG$ TGTGTGACCTGATCCAGAGTAAGTGCCTCTCCAAGGAGAACGCTA  $\tt CCACGGACCTCCTCACTAAGCTGCCTGCGCACACAGTGGGGGATG$ TGAAATGTGACATGGAGGTGAGCTGCCCAGATGGCTATACCTGCT GCCGTCTACAGTCGGGGGCCTGGGGCTGCTGCCCTTTTACCCAGG  $\tt CTGTGTGCTGTGAGGACCACATACACTGCTGTCCCGCGGGGTTTA$ 

#### -continued

CGTGTGACACGCAGAAGGGTACCTGTGAACAGGGGCCCCACCAGG TGCCCTGGATGGAGAAGGCCCCAGCTCACCTCAGCCTGCCAGACC  ${\tt CACAAGCCTTGAAGAGAGATGTCCCCTGTGATAATGTCAGCAGCT}$ GTCCCTCCGATACCTGCTGCCAACTCACGTCTGGGGAGTGGG GCTGCTGTCCAATCCCAGAGGCTGTCTGCTGCTCGGACCACCAGC ACTGCTGCCCCCAGGGCTACACGTGTGTAGCTGAGGGGCAGTGTC GCCGGGCTTCCTTATCCCACCCCAGAGACATCGGCTGTGACCAGC ACACCAGCTGCCCGGTGGGGCAGACCTGCTGCCCGAGCCTGGGTG GGAGCTGGGCCTGCCAGTTGCCCCATGCTGTGTGCTGCGAGG ATCGCCAGCACTGCTGCCCGGCTGGCTACACCTGCAACGTGAAGG CTCGATCCTGCGAGAAGGAAGTGGTCTCTGCCCAGCCTGCCACCT TCCTGGCCCGTAGCCCTCACGTGGGTGTGAAGGACGTGGAGTGTG GGGAAGGACACTTCTGCCATGATAACCAGACCTGCTGCCGAGACA ACCGACAGGGCTGGGCCTGCTGTCCCTACCGCCAGGGCGTCTGTT  $\tt GTGCTGATCGGCGCCACTGCTGTCCTGCTGGCTTCCGCTGCGCAG$  $\tt CCAGGGGTACCAAGTGTTTGCGCAGGGAGGCCCCGCGCTGGGACG$  $\tt CCCCTTTGAGGGACCCAGCCTTGAGACAGCTGCTGTGAGGGACAG$  ${\tt TACTGAAGACTCTGCAGCCCTCGGGACCCCACTCGGAGGGTGCCC}$ TCTGCTCA

SEQ ID NO: 4

MWTLVSWVALTAGLVAGTDYKDDDDKRCPDGQFCPVACCLDPGGA

SYSCCRPLLDKWPTTLSRHLGGPCQVDAHCSAGHSCIFTVSGTSS

CCPFPEAVACGDGHHCCPRGFHCSADGRSCFQRSGNNSVGAIQCP

DSQFECPDFSTCCVMVDGSWGCCPMPQASCCEDRVHCCPHGAFCD

LVHTRCITPTGTHPLAKKLPAQRTNRAVALSSSVMCPDARSRCPD

GSTCCELPSGKYGCCPMPNATCCSDHLHCCPQDTVCDLIQSKCLS

KENATTDLLTKLPAHTVGDVKCDMEVSCPDGYTCCRLQSGAWGCC

PFTQAVCCEDHIHCCPAGFTCDTQKGTCEQGPHQVPWMEKAPAHL

SLPDPQALKRDVPCDNVSSCPSSDTCCQLTSGEWGCCPIPEAVCC

SDHQHCCPQGYTCVAEGQCQRGSEIVAGLEKMPARRASLSHPRDI

GCDQHTSCPVGQTCCPSLGGSWACCQLPHAVCCEDRQHCCPAGYT

CNVKARSCEKEVVSAQPATFLARSPHVGVKDVECGEGHFCHDNQT

CCRDNRQGWACCPYRQGVCCADRRHCCPAGFRCAARGTKCLRREA

PRWDAPLRDPALROLL

[0060] In one embodiment, the detectable tag is HA tag and the target protein is a N-terminal HA tagged PGRN. The final screening includes, adding an anti-HA antibody that is covalently attached to a dye (such as Dylight 650® anti-HA antibody) into the wells containing the transfected cells to immunostain the HA tagged PGRN, if present, and detecting the dye.

N-terminal HA tagged PGRN: SEQ ID NO: 5 ATGTGGACTCTGGTCTCATGGGTGGCTCTGACTGCTGGACTGGTG GACGGCCAGTTCTGCGACAAATGGCCCACAACACTGAGCAGGCAT  $\tt CTGGGTGGCCCTGCCAGGTTGATGCCCACTGCTCTGCCGGCCAC$ TCCTGCATCTTTACCGTCTCAGGGACTTCCAGTTGCTGCCCCTTC CCAGAGGCCGTGGCATGCGGGGATGGCCATCACTGCTGCCCACGG GGCTTCCACTGCAGTGCAGACGGGCGATCCTGCTTCCAAAGATCA GGTAACAACTCCGTGGGTGCCATCCAGTGCCCTGATAGTCAGTTC GAATGCCCGGACTTCTCCACGTGCTGTTTATGGTCGATGGCTCC TGGGGGTGCTGCCCCATGCCCCAGGCTTCCTGCTGTGAAGACAGG GTGCACTGCTGTCCGCACGGTGCCTTCTGCGACCTGGTTCACACC CGCTGCATCACACCCACGGGCACCCACCCCCTGGCAAAGAAGCTC CCTGCCCAGAGGACTAACAGGGCAGTGGCCTTGTCCAGCTCGGTC ATGTGTCCGGACGCACGGTCCCGGTGCCCTGATGGTTCTACCTGC TGTGAGCTGCCCAGTGGGAAGTATGGCTGCTGCCCAATGCCCAAC GCCACCTGCTGCTCCGATCACCTGCACTGCTGCCCCCAAGACACT GTGTGTGACCTGATCCAGAGTAAGTGCCTCTCCAAGGAGAACGCT ACCACGGACCTCCTCACTAAGCTGCCTGCGCACACAGTGGGGGAT  $\tt GTGAAATGTGACATGGAGGTGAGCTGCCCAGATGGCTATACCTGC$  $\tt TGCCGTCTACAGTCGGGGGCCTGGGGCTGCTGCCCTTTTACCCAG$ GCTGTGTGCTGTGAGGACCACATACACTGCTGTCCCGCGGGGTTT ACGTGTGACACGCAGAAGGGTACCTGTGAACAGGGGCCCCACCAG GTGCCTGGATGGAGAGGCCCCAGCTCACCTCAGCCTGCCAGAC CCACAAGCCTTGAAGAGAGATGTCCCCTGTGATAATGTCAGCAGC  ${\tt TGTCCCTCCGATACCTGCTGCCAACTCACGTCTGGGGAGTGG}$ GGCTGCTGTCCAATCCCAGAGGCTGTCTGCTGCTCGGACCACCAG CACTGCTGCCCCCAGGGCTACACGTGTGTAGCTGAGGGGCAGTGT CGCCGGGCTTCCTTATCCCACCCCAGAGACATCGGCTGTGACCAG CACACCAGCTGCCCGGTGGGGCAGACCTGCTGCCCGAGCCTGGGT GGGAGCTGGGCCTGCCAGTTGCCCCATGCTGTGTGCTGCGAG GATCGCCAGCACTGCTGCCCGGCTGGCTACACCTGCAACGTGAAG GCTCGATCCTGCGAGAAGGAAGTGGTCTCTGCCCAGCCTGCCACC TTCCTGGCCCGTAGCCCTCACGTGGGTGTGAAGGACGTGGAGTGT GGGGAAGGACACTTCTGCCATGATAACCAGACCTGCTGCCGAGAC AACCGACAGGGCTGGGCCTGCTGTCCCTACCGCCAGGGCGTCTGT TGTGCTGATCGGCGCCACTGCTGTCCTGCTGGCTTCCGCTGCGCA GCCAGGGGTACCAAGTGTTTGCGCAGGGAGGCCCCGCGCTGGGAC

-continued
GCCCCTTTGAGGGACCCAGCCTTGAGACAGCTGCTGTGAGGGACA
GTACTGAAGACTCTGCAGCCCTCGGGACCCCACTCGGAGGGTGCC
CTCTGCTCA

SEQ ID NO: 6
MWTLVSWVALTAGLVAGTYPYDVPDYARCPDGQFCPVACCLDPGG
ASYSCCRPLLDKWPTTLSRHLGGPCQVDAHCSAGHSCIFTVSGTS
SCCPFPEAVACGDGHHCCPRGFHCSADGRSCFQRSGNNSVGAIQC
PDSQFECPDFSTCCVMVDGSWGCCPMPQASCCEDRVHCCPHGAFC
DLVHTRCITPTGTHPLAKKLPAQRTNRAVALSSSVMCPDARSRCP
DGSTCCELPSGKYGCCPMPNATCCSDHLHCCPQDTVCDLIQSKCL
SKENATTDLLTKLPAHTVGDVKCDMEVSCPDGYTCCRLQSGAWGC
CPFTQAVCCEDHIHCCPAGFTCDTQKGTCEQGPHQVPWMEKAPAH
LSLPDPQALKRDVPCDNVSSCPSSDTCCQLTSGEWGCCPIPEAVC
CSDHQHCCPQGYTCVAEGQCQRGSEIVAGLEKMPARRASLSHPRD
IGCDQHTSCPVGQTCCPSLGGSWACCQLPHAVCCEDRQHCCPAGY
TCNVKARSCEKEVVSAQPATFLARSPHVGVKDVECGEGHFCHDNQ
TCCRDNRQGWACCPYRQGVCCADRRHCCPAGFRCAARGTKCLRRE
APRWDAPLRDPALROL

[0061] Preferably, a high-content throughput imager is used to detect the tagged PGRN.

Identification of Intracellular Proteins that Bind PGRN

[0062] In another aspect, the invention disclosed herein are methods for screening for intracellular proteins that bind to progranulin (PGRN). A schematic outline of the method is shown in FIG. 1.

[0063] In this method the target protein comprised of PGRN fused with a detectable tag (as described above), a cDNA library or collection comprising a plurality of expression vectors each encoding a distinct protein (preferably human proteins, more preferably human intracellular proteins), and host cells (as described above), are utilized.

[0064] The method includes introducing an expression vector from the cDNA library into a host cell and maintaining the expression vector and the host cell in a suitable culture media under suitable culturing conditions to allow transfection and expression of the protein encoded by the expression vector. Suitable culturing conditions are the same as those described above.

[0065] After removing the culture media, the transfected cells are permeabilized with a permeabilizing agent (as described above) followed by washing with wash buffer. The permeabilizing agent can sufficiently disrupt the cellular membrane of the cells to permit target proteins to enter the cell. The permeabilized cells are then brought into contact with the target protein and maintained under a suitable condition (as described above) to allow potential bond between the intracellular proteins and the target protein. Optionally, prior to permeabilization, the transfected cells are fixed with a fixing agent followed by washing with washing buffer.

[0066] After contacting the target protein, the cells are fixed with a fixing agent (as described above) and washed with washing buffer to remove the un-bond target protein.

[0067] Finally, the presence of the detectable tag is determined, wherein if the detectable tag is determined to be present, the protein encoded by the expression vector is identified as an intracellular protein binds PGRN.

[0068] Optionally, prior to detecting the detectable tag, the fixed cells are again permeabilized with a permeabilization agent, followed by washing with washing buffer.

[0069] In one embodiment, the detectable tag is FLAG tag and the target protein is a N-terminal FLAG tagged PGRN. To determine the presence of the FLAG tag, an anti-FLAG antibody that is covalently attached to a fluorescent dye (such as a cyanine dye, e.g., Cy3 or Cy5) is added to immunostain the FLAG tagged PGRN, if present, and an imager or fluorometer is used to detect the presence of the fluorescent dye on the cells.

[0070] In one embodiment, the detectable tag is HA tag and the target protein is a N-terminal HA tagged PGRN. The final screening includes, adding an anti-HA antibody that is covalently attached to a dye (such as Dylight 650® anti-HA antibody) into the wells containing the transfected cells to immunostain the HA tagged PGRN, if present, and detecting the dye.

[0071] In one embodiment, the screening for intracellular proteins that bind to progranulin (PGRN) is conducted in a high-throughput manner. In the high-throughput process, multi-well plates, such as 96-well, 384-well, or 1536-well plates, are used. Host cells in a suitable culture media are placed in one or more wells of the plate. And starting from placing host cells in the one or more wells until screening for the presence of the detectable tag, each step of the process is performed in a high-throughput manner, such as by a robotic arm.

### **EXAMPLE**

[0072] It will be appreciated by those skilled in the art that changes could be made to the embodiments described above without departing from the broad inventive concept thereof. It is understood, therefore, that this invention is not limited to the particular embodiments disclosed, but it is intended to cover modifications within the spirit and scope of the present invention as defined by the present description.

Identification of Cell Surface Receptor for Progranulin

[0073] In this study, a cDNA collection containing over 4,000 transfection-ready cDNA plasmids for both human and mouse proteins with a transmembrane domain (obtained from OriGene) was used. Human open reading frames (ORFs) are located on pCMV6-AC pCMV6-XL4, pCMV6-XL5, pCMV6-XL6, or pCMV6-NEO backbones and mouse ORFs are located on a pCMV6-Kan/Neo skeleton. The cDNA plasmids were transfected individually into HEK293T cells (ATCC) in 384-well poly-D-lysine coated plates (Perkin Elmer) using Fugene 6 (Promega). Each well was seeded with 10,000 cells. Using Fugene 6 according to the manufacturer's instructions, each well was transfected with 60 ng of plasmid DNA, 0.3 ng of which was GFP plasmid (green fluorescent protein (GFP) gene expressed on pcDNA3.1) as a transfection efficiency control. The cells were incubated for 1 day at 37° C., after which they were moved to 30° C. and incubated for 2 days. Three (3) days after transfection, cells in each well were incubated with 100 nM N-terminal flag tagged PGRN at room temperature for 2 hours. Following PGRN incubation, the media was removed and the cells were fixed with 4% paraformaldehyde for 40 minutes at room temperature. Afterwards, the cells were washed with HEPES Balanced Salt Solution (HBSS) 3 times, and permeabilized with 0.3% Triton X-100 for 30 minutes at room temperature. After permeabilization, cells were washed 3 times again, and then immunostained with 400 ng/mL of Anti-Flag M2-Cy3 antibody (Sigma) at 4° C. overnight. The following day, the cells were washed 3 times again and then stained with NucBlue Live ReadyProbes Reagent (Invitrogen) according to manufacturer's instructions. Subsequently, the plate was imaged using the Opera Phenix high-content throughput imaging machine (Perkin Elmer).

Identification of Intracellular Binding Partner for Progranulin

[0074] Two separate cDNA collections of around 9.000 total plasmids were screened. The first cDNA collection includes about 3000 transfection-ready cDNA plasmids for both human and mouse proteins expressed in the central nervous system (CNS). Human ORFs are similar to the above, located on pCMV6-AC pCMV6-XL4, pCMV6-XL5, pCMV6-XL6, or pCMV6-NEO backbones. Mouse ORFs in this collection are located on a pCMV6-Entry backbone. Some of the proteins in this collection were tagged with a Myc-DDK tag, in which case HA-tagged PGRN was used for screening. The second cDNA collection containing about 6000 transfection-ready cDNA plasmids selected from the hORFeome V8.1 library (obtained from the Broad Institute). The plasmids were transfected individually into HEK293T cells (ATCC) in 384-well poly-D-lysine coated plates (Perkin Elmer) using Fugene 6 (Promega). Each well was seeded with 10,000 cells. Using Fugene 6 according to the manufacturer's instructions, each well was transfected with 60 ng of plasmid DNA, 3 ng of which was GFP plasmid as a transfection control. The cells were incubated for 1 day at 37° C., after which they were moved to 30° C. and incubated for 2 days. Three (3) days after transfection the media was removed and the cells were fixed with 4% paraformaldehyde for 40 minutes at room temperature. Afterwards, the cells were washed with HBSS 3 times, and permeabilized with 0.3% Triton X-100 for 30 minutes at room temperature. After permeabilization, cells were washed 3 times again, and incubated in Dulbecco's Modified Eagle Medium (DMEM) containing 100 nM N-terminal flag tagged PGRN for 2 hours at room temperature. After incubation with PGRN, the DMEM was removed and the cells were fixed again with 4% paraformaldehyde for 40 minutes at room temperature. After the 2nd fixation, the cells were washed again with HBSS for 3 times and then immunostained with 400 ng/ml of Anti-Flag M2-Cy3 antibody (Sigma) in 4° C. overnight. The following day, the cells were washed 3 times again and then stained with NucBlue Live ReadyProbes Reagent (Invitrogen) according to manufacturer's instructions. Subsequently, the plate was imaged using the Opera Phenix high-content throughput imaging machine (Perkin Elmer). FIG. 2 show the Cy3 intensity for cells transfected with pcDNA3 vector (negative control) and cells transfected with an expression vector carrying the coding sequence of sortilin (positive control). Also shown in FIG. 2 is the Cy3 intensity for cells transfected with one of the 9000 expression plasmids, which is comparable to the positive control. Thus, the intracellular protein encoded by the expression plasmid was identified as Hit #1.

#### Confirmation of Hit #1

[0075] The plasmid for Hit #1, along with pcDNA3 vector (negative control) and an expression vector carrying the coding sequence of sortilin (positive control) were individually transfected into HEK293 cells. Following the same process described under "Identification of Intracellular Binding Partner for Progranulin", the transfected cells were exposed to N-terminal FLAG tagged PGRN or N-terminal HA tagged PGRN, followed by immunostaining with anti-Flag M2-Cy3 antibody and Dylight 650 anti-HA antibody, respectively. As shown in FIGS. 3A and 3C, the Cy3 and Dylight intensity for cells transfected with Hit #1 plasmid was comparable to that for the positive control. In addition, the cells transfected with Hit #1 plasmid was also immunostained with anti-Flag M2-Cy3 antibody and Dylight 650 anti-HA antibody without prior exposure to the FLAG

tagged PGRN or HA tagged PGRN. And as shown in FIG. 3B, in both cases, the fluorescent signal was the same as the negative control, confirming that the identification of Hit #1 was not due to non-specific antibody binding to the cells transfected with the Hit #1 plasmid.

[0076] Further, following the same process described under "Identification of Cell Surface Receptor for Progranulin", HEK293 cells transfected with pcDNA3 vector (negative control), expression vector carrying the coding sequence of sortilin (positive control), and Hit #1 plasmid were exposed to FLAG tagged PGRN or HA tagged PGRN without being permeabilized, followed by immunostaining with anti-Flag M2-Cy3 antibody and Dylight 650 anti-HA antibody, respectively. As shown, in FIG. 4, the fluorescent signal intensity for Hit #1 was the same as the negative control, confirming that Hit #1 is not a surface receptor for PGRN.

SEQUENCE LISTING

```
<160> NUMBER OF SEQ ID NOS: 6
<210> SEQ ID NO 1
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide
<400> SEQUENCE: 1
Tyr Pro Tyr Asp Val Pro Asp Tyr Ala
<210> SEQ ID NO 2
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic peptide
<400> SEQUENCE: 2
Asp Tyr Lys Asp Asp Asp Lys
<210> SEQ ID NO 3
<211> LENGTH: 1808
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<400> SEQUENCE: 3
atgtggactc tggtctcatg ggtggctctg actgctggac tggtggctgg aaccgactac
aaggacgacg acgacaaact cgctgccctg acggccagtt ctgcgacaaa tggcccacaa
                                                                      120
cactgagcag gcatctgggt ggcccctgcc aggttgatgc ccactgctct gccggccact
cctgcatctt taccgtctca gggacttcca gttgctgccc cttcccagag gccgtggcat
geggggatgg ceateactge tgeceaeggg gettecaetg cagtgeagae gggegateet
gcttccaaag atcaggtaac aactccgtgg gtgccatcca gtgccctgat agtcagttcg
                                                                      360
aatgcccgga cttctccacg tgctgtgtta tggtcgatgg ctcctggggg tgctgcccca
                                                                      420
tgccccaggc ttcctgctgt gaagacaggg tgcactgctg tccgcacggt gccttctgcg
```

acctggttca cacccgctgc atcacaccca cgggcaccca ccccctggca aaga	agetee 540
ctgcccagag gactaacagg gcagtggcct tgtccagctc ggtcatgtgt ccg	gacgcac 600
ggtcccggtg ccctgatggt tctacctgct gtgagctgcc cagtgggaag tatg	ggctgct 660
geccaatgee caaegecaee tgetgeteeg ateaeetgea etgetgeeee caae	gacactg 720
tgtgtgacct gatccagagt aagtgcctct ccaaggagaa cgctaccacg gacc	etectea 780
ctaagctgcc tgcgcacaca gtgggggatg tgaaatgtga catggaggtg agct	gcccag 840
atggctatac ctgctgccgt ctacagtcgg gggcctgggg ctgctgccct ttta	acccagg 900
ctgtgtgctg tgaggaccac atacactgct gtcccgcggg gtttacgtgt gaca	acgcaga 960
agggtacctg tgaacagggg ccccaccagg tgccctggat ggagaaggcc cca	geteace 1020
teageetgee agaceeacaa geettgaaga gagatgteee etgtgataat gtea	agcagct 1080
gtccctcctc cgatacctgc tgccaactca cgtctgggga gtggggctgc tgtc	caatcc 1140
cagaggctgt ctgctgctcg gaccaccagc actgctgccc ccagggctac acgt	gtgtag 1200
ctgaggggca gtgtcagcga ggaagcgaga tcgtggctgg actggagaag atg	ectgece 1260
gccgggcttc cttatcccac cccagagaca tcggctgtga ccagcacacc agct	gcccgg 1320
tggggcagac ctgctgcccg agcctgggtg ggagctgggc ctgctgccag ttgc	cccatg 1380
ctgtgtgctg cgaggatcgc cagcactgct gcccggctgg ctacacctgc aac	gtgaagg 1440
ctcgatcctg cgagaaggaa gtggtctctg cccagcctgc caccttcctg gccc	gtagcc 1500
ctcacgtggg tgtgaaggac gtggagtgtg gggaaggaca cttctgccat gata	aaccaga 1560
cctgctgccg agacaaccga cagggctggg cctgctgtcc ctaccgccag ggc	gtctgtt 1620
gtgctgatcg gcgccactgc tgtcctgctg gcttccgctg cgcagccagg ggta	accaagt 1680
gtttgegeag ggaggeeeeg egetgggaeg eeeetttgag ggaeeeagee ttga	agacagc 1740
tgctgtgagg gacagtactg aagactctgc agccctcggg accccactcg gagg	ggtgeee 1800
tctgctca	1808
<210> SEQ ID NO 4	
<pre>&lt;211&gt; LENGTH: 601 &lt;212&gt; TYPE: PRT</pre>	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE: <223> OTHER INFORMATION: Synthetic peptide	
<400> SEQUENCE: 4	
Met Trp Thr Leu Val Ser Trp Val Ala Leu Thr Ala Gly Leu Val 1 5 10 15	Ala
Gly Thr Asp Tyr Lys Asp Asp Asp Lys Arg Cys Pro Asp Gly 20 25 30	/ Gln
Phe Cys Pro Val Ala Cys Cys Leu Asp Pro Gly Gly Ala Ser Tys 35 40 45	: Ser
Cys Cys Arg Pro Leu Leu Asp Lys Trp Pro Thr Thr Leu Ser Arg	, His
Leu Gly Gly Pro Cys Gln Val Asp Ala His Cys Ser Ala Gly His	; Ser
65 70 75	80

Cys Ile Phe Thr Val Ser Gly Thr Ser Ser Cys Cys Pro Phe Pro Glu

Ala Val Ala Cys Gly Asp Gly His His Cys Cys Pro Arg Gly Phe His 100  $$105\$ 

85 90

Cys	Ser	Ala 115	Asp	Gly	Arg	Ser	Cys 120	Phe	Gln	Arg	Ser	Gly 125	Asn	Asn	Ser
Val	Gly 130	Ala	Ile	Gln	Cys	Pro 135	Asp	Ser	Gln	Phe	Glu 140	Cys	Pro	Asp	Phe
Ser 145	Thr	Сув	Сув	Val	Met 150	Val	Asp	Gly	Ser	Trp 155	Gly	Сув	Сув	Pro	Met 160
Pro	Gln	Ala	Ser	Cys 165	СЛа	Glu	Asp	Arg	Val 170	His	СЛа	СЛа	Pro	His 175	Gly
Ala	Phe	Сла	Asp 180	Leu	Val	His	Thr	Arg 185	Сув	Ile	Thr	Pro	Thr 190	Gly	Thr
His	Pro	Leu 195	Ala	Lys	Lys	Leu	Pro 200	Ala	Gln	Arg	Thr	Asn 205	Arg	Ala	Val
Ala	Leu 210	Ser	Ser	Ser	Val	Met 215	Cya	Pro	Asp	Ala	Arg 220	Ser	Arg	Cya	Pro
Asp 225	Gly	Ser	Thr	CÀa	Cys 230	Glu	Leu	Pro	Ser	Gly 235	Lys	Tyr	Gly	Cys	Сув 240
Pro	Met	Pro	Asn	Ala 245	Thr	Cys	Cys	Ser	Asp 250	His	Leu	His	Сув	Сув 255	Pro
Gln	Asp	Thr	Val 260	CAa	Asp	Leu	Ile	Gln 265	Ser	Lys	CAa	Leu	Ser 270	Lys	Glu
Asn	Ala	Thr 275	Thr	Asp	Leu	Leu	Thr 280	Lys	Leu	Pro	Ala	His 285	Thr	Val	Gly
Asp	Val 290	Lys	Cys	Asp	Met	Glu 295	Val	Ser	Cys	Pro	300	Gly	Tyr	Thr	Cya
Сув 305	Arg	Leu	Gln	Ser	Gly 310	Ala	Trp	Gly	Cys	Суз 315	Pro	Phe	Thr	Gln	Ala 320
Val	CAa	Сла	Glu	Asp 325	His	Ile	His	CAa	330 230	Pro	Ala	Gly	Phe	Thr 335	Сув
Asp	Thr	Gln	Lys 340	Gly	Thr	CÀa	Glu	Gln 345	Gly	Pro	His	Gln	Val 350	Pro	Trp
Met	Glu	155 355	Ala	Pro	Ala	His	Leu 360	Ser	Leu	Pro	Asp	Pro 365	Gln	Ala	Leu
ГЛа	Arg 370	Asp	Val	Pro	CÀa	Asp 375	Asn	Val	Ser	Ser	380 CÀa	Pro	Ser	Ser	Asp
Thr 385	CÀa	Cya	Gln	Leu	Thr 390	Ser	Gly	Glu	Trp	Gly 395	CÀa	СЛа	Pro	Ile	Pro 400
Glu	Ala	Val	СЛа	Cys 405	Ser	Asp	His	Gln	His 410	CÀa	CÀa	Pro	Gln	Gly 415	Tyr
Thr	CÀa	Val	Ala 420	Glu	Gly	Gln	CÀa	Gln 425	Arg	Gly	Ser	Glu	Ile 430	Val	Ala
Gly	Leu	Glu 435	Lys	Met	Pro	Ala	Arg 440	Arg	Ala	Ser	Leu	Ser 445	His	Pro	Arg
Asp	Ile 450	Gly	Cys	Asp	Gln	His 455	Thr	Ser	Cys	Pro	Val 460	Gly	Gln	Thr	Cys
Cys 465	Pro	Ser	Leu	Gly	Gly 470	Ser	Trp	Ala	Сув	Cys 475	Gln	Leu	Pro	His	Ala 480
Val	Сла	СЛа	Glu	Asp 485	Arg	Gln	His	СЛа	Сув 490	Pro	Ala	Gly	Tyr	Thr 495	Сув
Asn	Val	Lys	Ala 500	Arg	Ser	CÀa	Glu	Lys 505	Glu	Val	Val	Ser	Ala 510	Gln	Pro

Ala Thr Phe Leu Ala A	rg Ser Pro His Val Gly 520	Val Lys Asp Val Glu 525	
Cys Gly Glu Gly His P	he Cys His Asp Asn Gln 535	Thr Cys Cys Arg Asp	
9 1 1	la Cys Cys Pro Tyr Arg	Gln Gly Val Cys Cys	
	555 ys Cys Pro Ala Gly Phe		
565	ys cys PIO AIA GIY PNE 570	575	
Gly Thr Lys Cys Leu A 580	rg Arg Glu Ala Pro Arg 585	Trp Asp Ala Pro Leu 590	
Arg Asp Pro Ala Leu A	-		
595	600		
<210> SEQ ID NO 5 <211> LENGTH: 1809 <212> TYPE: DNA <213> ORGANISM: Artif <220> FEATURE: <223> OTHER INFORMATION	icial Sequence ON: Synthetic polynucl	eotide	
<400> SEQUENCE: 5			
		tggtggctgg aacctaccca	60
		tctgcgacaa atggcccaca	120
		cccactgctc tgccggccac	180
		gcagtgcaga cgggcgatcc	300
		agtgccctga tagtcagttc	360
		gctcctgggg gtgctgcccc	420
		gtccgcacgg tgccttctgc	480
gacctggttc acacccgctg	catcacaccc acgggcaccc	acccctggc aaagaagctc	540
cctgcccaga ggactaacag	ggcagtggcc ttgtccagct	cggtcatgtg tccggacgca	600
cggtcccggt gccctgatgg	ttctacctgc tgtgagctgc	ccagtgggaa gtatggctgc	660
tgcccaatgc ccaacgccac	ctgctgctcc gatcacctgc	actgctgccc ccaagacact	720
gtgtgtgacc tgatccagag	taagtgcctc tccaaggaga	acgctaccac ggacctcctc	780
actaagctgc ctgcgcacac	agtgggggat gtgaaatgtg	acatggaggt gagctgccca	840
		gctgctgccc ttttacccag	900
		ggtttacgtg tgacacgcag	960
		tggagaaggc cccagctcac	1020
		cctgtgataa tgtcagcagc	1080
		agtggggctg ctgtccaatc	1140
		cccagggcta cacgtgtgta	1200
		gactggagaa gatgcctgcc	1260
		accagcacac cagctgcccg	1320
		cctgctgcca gttgccccat	1380
gctgtgtgct gcgaggatcg	ccagcactgc tgcccggctg	gctacacctg caacgtgaag	1440

_												COII		ucu			
gctc	gato	cct (	gcga	gaag	ga a	gtgg	tete	gco	ccago	cctg	ccad	cctt	cct (	ggcc	cgtagc	1500	
cctc	acgt	gg (	gtgt	gaag	ga c	gtgg.	agtgi	gg	ggaaq	ggac	acti	tctg	cca 1	tgata	aaccag	1560	
acct	gct	gcc (	gaga	caac	cg a	cagg	gctg	g gc	ctgct	gtc	ccta	accg	cca 🤅	gggcg	gtctgt	1620	
tgtg	ctga	atc q	ggcg	ccact	tg c	tgtc	ctgc	gg(	cttco	eget	gcg	cage	cag q	gggta	accaag	1680	
tgtt	tgc	gca q	ggga	ggcc	cc g	cgct	ggga	e ge	ccctt	tga	ggga	accca	agc (	cttga	agacag	1740	
ctgc	tgt	gag g	ggac	agta	ct g	aaga	ctct	g caq	gadat	cgg	gaco	ccca	ctc (	ggagg	ggtgcc	1800	
ctct	gcto	ca														1809	
<220	> LI > T? > OI > FI	ENGTI YPE : RGAN: EATUI	H: 6 PRT ISM: RE:	01 Art:			Seque nthe		pept:	ide							
< 400	> SI	EQUEI	NCE:	6													
Met 1	Trp	Thr	Leu	Val 5	Ser	Trp	Val	Ala	Leu 10	Thr	Ala	Gly	Leu	Val 15	Ala		
Gly	Thr	Tyr	Pro 20	Tyr	Asp	Val	Pro	Asp 25	Tyr	Ala	Arg	Cys	Pro 30	Asp	Gly		
Gln	Phe	Сув 35	Pro	Val	Ala	CAa	Cys 40	Leu	Asp	Pro	Gly	Gly 45	Ala	Ser	Tyr		
Ser	Сув 50	Сув	Arg	Pro	Leu	Leu 55	Asp	Lys	Trp	Pro	Thr	Thr	Leu	Ser	Arg		
His 65	Leu	Gly	Gly	Pro	Cys	Gln	Val	Asp	Ala	His 75	СЛа	Ser	Ala	Gly	His 80		
Ser	Cys	Ile	Phe	Thr	Val	Ser	Gly	Thr	Ser 90	Ser	Càa	CAa	Pro	Phe 95	Pro		
Glu	Ala	Val	Ala 100	СЛа	Gly	Asp	Gly	His	His	СЛа	СЛа	Pro	Arg 110	Gly	Phe		
His	Cys	Ser 115	Ala	Asp	Gly	Arg	Ser 120	Сув	Phe	Gln	Arg	Ser 125	Gly	Asn	Asn		
Ser	Val 130	Gly	Ala	Ile	Gln	Cys 135	Pro	Asp	Ser	Gln	Phe	Glu	CÀa	Pro	Asp		
Phe 145	Ser	Thr	Cys	Cys	Val 150	Met	Val	Asp	Gly	Ser 155	Trp	Gly	Cys	CAa	Pro 160		
	Pro	Gln	Ala	Ser 165		Cys	Glu	Asp	Arg 170		His	Сув	Сув	Pro 175			
Gly	Ala	Phe	Cys 180	Asp	Leu	Val	His	Thr	Arg	Cys	Ile	Thr	Pro	Thr	Gly		
Thr	His	Pro			Lys	Lys	Leu 200			Gln	Arg	Thr 205		Arg	Ala		
Val	Ala 210		Ser	Ser	Ser	Val 215	Met	Cys	Pro	Asp	Ala 220		Ser	Arg	Cys		
		Gly	Ser	Thr	-			Leu	Pro			Lys	Tyr	Gly	_		
225					230					235					240		
CÀa	Pro	Met	Pro	Asn 245	Ala	Thr	Сув	CAa	Ser 250	Asp	His	Leu	His	Сув 255	Cys		
Pro	Gln	Asp	Thr 260		CAa	Asp	Leu	Ile 265	Gln	Ser	Lys	CAa	Leu 270	Ser	Lys		

Glu Asn Ala Thr Thr Asp Leu Leu Thr Lys Leu Pro Ala His Thr Val

_																
			275					280					285			
G:	_	Asp 290	Val	Lys	CAa	Asp	Met 295	Glu	Val	Ser	Сув	Pro 300	Asp	Gly	Tyr	Thr
	/s 05	Cys	Arg	Leu	Gln	Ser 310	Gly	Ala	Trp	Gly	Сув 315	CAa	Pro	Phe	Thr	Gln 320
A.	la	Val	Сув	Сув	Glu 325	Asp	His	Ile	His	Cys	Сув	Pro	Ala	Gly	Phe 335	Thr
C	Ys	Asp	Thr	Gln 340	Lys	Gly	Thr	Сув	Glu 345	Gln	Gly	Pro	His	Gln 350	Val	Pro
T	rp	Met	Glu 355	Lys	Ala	Pro	Ala	His 360	Leu	Ser	Leu	Pro	Asp 365	Pro	Gln	Ala
Le		Lys 370	Arg	Asp	Val	Pro	Cys 375	Asp	Asn	Val	Ser	Ser 380	Сув	Pro	Ser	Ser
	sp 85	Thr	Сув	Cys	Gln	Leu 390	Thr	Ser	Gly	Glu	Trp 395	Gly	Сув	Сув	Pro	Ile 400
Pi	ro	Glu	Ala	Val	Сув 405	CÀa	Ser	Asp	His	Gln 410	His	Cys	Cys	Pro	Gln 415	Gly
T	γr	Thr	Сув	Val 420	Ala	Glu	Gly	Gln	Cys 425	Gln	Arg	Gly	Ser	Glu 430	Ile	Val
Α:	la	Gly	Leu 435	Glu	ГÀв	Met	Pro	Ala 440	Arg	Arg	Ala	Ser	Leu 445	Ser	His	Pro
A		Asp 450	Ile	Gly	Cys	Asp	Gln 455	His	Thr	Ser	Cys	Pro 460	Val	Gly	Gln	Thr
	/s 55	Cys	Pro	Ser	Leu	Gly 470	Gly	Ser	Trp	Ala	Cys 475	СЛв	Gln	Leu	Pro	His 480
A.	la	Val	Сув	Сув	Glu 485	Asp	Arg	Gln	His	Сув 490	Сув	Pro	Ala	Gly	Tyr 495	Thr
C.	/s	Asn	Val	Lys 500	Ala	Arg	Ser	Сув	Glu 505	Lys	Glu	Val	Val	Ser 510	Ala	Gln
P	ro	Ala	Thr 515	Phe	Leu	Ala	Arg	Ser 520	Pro	His	Val	Gly	Val 525	Lys	Asp	Val
G:		Cys 530	Gly	Glu	Gly	His	Phe 535	Сув	His	Asp	Asn	Gln 540	Thr	Сув	Сув	Arg
	sp 15	Asn	Arg	Gln	Gly	Trp 550	Ala	Сув	Сув	Pro	Tyr 555	Arg	Gln	Gly	Val	Сув 560
C	Ys	Ala	Asp	Arg	Arg 565	His	CÀa	CÀa	Pro	Ala 570	Gly	Phe	Arg	CÀa	Ala 575	Ala
A	rg	Gly	Thr	580	CÀa	Leu	Arg	Arg	Glu 585	Ala	Pro	Arg	Trp	Asp 590	Ala	Pro
Le	∋u	Arg	Asp 595	Pro	Ala	Leu	Arg	Gln 600	Leu							

### What is claimed is:

- 1. A high-throughput screening method for identifying transmembrane proteins that bind progranulin (PGRN), comprising:
  - a) providing a cDNA collection comprising a plurality of expression vectors, each expression vector encoding a distinct transmembrane protein;
  - b) providing a plate comprising a plurality of wells, wherein one or more wells thereof each contains host cells;
- c) into each of the one or more wells, introducing a distinct expression vector from said collection, thereby producing distinct transfected cells in each well;
- d) culturing the transfected cells under a condition to allow expression of a distinct transmembrane protein in each well;
- e) contacting the transfected cells in each of the one or more wells with a target protein comprising PGRN attached to a detectable tag;
- f) fixing the transfected cells with a fixing agent after contact with the target protein;

- g) washing the fixed cells; and
- h) after washing the fixed cells, determining the presence of the detectable tag in each of the one or more wells, wherein if the detectable tag is determined to be present in a well, the transmembrane protein expressed by the transfected cells in the well is identified as a transmembrane protein that binds PGRN.
- 2. The high-throughput screening method of claim 1, wherein the host cells are selected from the group consisting of human embryonic kidney 293T (HEK293T) cell, HEK293F cells, Hela cells, Chinese hamster ovary (CHO) cells, NIH 3T3 cells, MCF-7 cells, Hep G2 cells, baby hamster kidney (BHK) cells, BV-2 (mouse, C57BL/6, brain, microglial) cells, Neuro-2a (N2a) cells and Cos7 cells.
- 3. The high-throughput screening method of claim 1 or 2, wherein the fixing agent is selected from glutaraldehyde, methanol and paraformaldehyde.
- **4**. The high-throughput screening method of any one of claims 1-3, wherein, the detectable tag is selected from the group consisting of green fluorescent protein (GFP), myc, myc-pyruvate kinase (myc-PK), His6, maltose binding protein (MBP), human influenza virus hemagglutinin (HA) tag, flag tag (FLAG), and glutathione-S-transferase (GST) tag.
- 5. The high-throughput screening method of claim 4, wherein, the target protein is a FLAG tagged PGRN.
- 6. The high-throughput screening method of claim 5, wherein, step h) comprises h1) adding an anti-FLAG antibody that is covalently attached to a fluorescent dye into the wells containing the transfected cells to immunostain the FLAG tagged PGRN, if present, and h2) detecting the fluorescent dye using a fluorometer or an imager in a high-throughput manner.
- 7. The high-throughput screening method of claim 6, wherein the fluorescent dye is cyanine dye.
- **8**. The high-throughput screening method of claim **4**, wherein, the target protein is a HA tagged PGRN.
- 9. The high-throughput screening method of claim 8, wherein, step h) comprises h1) adding an anti-HA antibody that is covalently attached to a fluorescent dye into the wells containing the transfected cells to immunostain the HA tagged PGRN, if present, and h2) detecting the fluorescent dye using a fluorometer or an imager in a high-throughput manner.
- 10. The high-throughput screening method of claim 6, wherein the fluorescent dye is Dylight 650.
- 11. The high-throughput screening method of any one of claims 1-10, wherein between steps g) and h), the method further comprises permeabilizing the fixed cells with a permeabilizing agent and washing the permeabilized cells.
- 12. The high-throughput screening method of claim 11, wherein the permeabilizing agent is selected from the group consisting of Saponin, Triton X-100, lysolecithin, n-Octyl-B-D-glucopyranoside, and Tween 20.
- 13. A method for identifying intracellular proteins that bind PGRN, comprising:
  - a) introducing into host cells an expression vector encoding a heterologous protein, thereby producing transfected cells:
  - b) culturing the transfected cells under a condition to allow expression of the heterologous protein;
  - c) permeabilizing the transfected cells with a permeabilizing agent;
  - d) contacting the permeabilized cells with a target protein comprising PGRN attached to a detectable tag;

- e) fixing the transfected cells with a fixing agent after contact with the target protein;
- f) washing the fixed cells; and
- g) after washing the fixed cells, determining the presence of the detectable tag on the washed cells, wherein if the detectable tag is determined to be present, the heterologous protein expressed by the cells is identified as an intracellular protein that binds PGRN.
- 14. The method of claim 13, wherein the host cells are selected from the group consisting of human embryonic kidney 293T (HEK293T) cell, HEK293F cells, HeLa cells, Chinese hamster ovary (CHO) cells, NIH 3T3 cells, MCF-7 cells, Hep G2 cells, baby hamster kidney (BHK) cells, BV-2 (mouse, C57BL/6, brain, microglial) cells, Neuro-2a (N2a) cells and Cos7 cells.
- 15. The method of 13 or 14, wherein, the detectable tag is selected from the group consisting of green fluorescent protein (GFP), myc, myc-pyruvate kinase (myc-PK), His6, maltose binding protein (MBP), human influenza virus hemagglutinin (HA) tag, flag tag (FLAG), and glutathione-S-transferase (GST) tag.
- 16. The method of claim 15, wherein, the target protein is a FLAG tagged PGRN.
- 17. The method of claim 16, wherein, step g) comprises g1) exposing the cells to an anti-FLAG antibody that is covalently attached to a fluorescent dye to immunostain the FLAG tagged PGRN, if present, and g2) detecting the fluorescent dye using a fluorometer or an imager.
- 18. The method of claim 17, wherein the fluorescent dye is cyanine dye.
- 19. The method of claim 15, wherein, the target protein is an HA tagged PGRN.
- **20**. The method of claim **19**, wherein, step g) comprises g1) exposing the cells to an anti-HA antibody that is covalently attached to a fluorescent dye to immunostain the HA tagged PGRN, if present, and g2) detecting the fluorescent dye using a fluorometer or an imager.
- 21. The method of claim 20, wherein the fluorescent dye is Dylight 650.
- 22. The method of any one of claims 13-21, wherein between steps b) and c), the method further comprises, fixing the cultured cells with the fixing agent and washing the fixed cells.
- 23. The method of any one of claims 13-22, wherein the fixing agent is selected from glutaraldehyde, methanol and paraformaldehyde.
- 24. The method of any one of claims 13-23, wherein between steps f) and g), the method further comprises permeabilizing the fixed cells with a permeabilizing agent and washing the permeabilized cells.
- 25. The method of any one of claims 13-24, wherein the permeabilizing agent is selected from the group consisting of Saponin, Triton X-100, lysolecithin, n-Octyl-B-D-glucopyranoside, and Tween 20.
- **26**. A high-throughput screening method for identifying intracellular proteins that bind PGRN, comprising:
  - a) providing a cDNA collection comprising a plurality of expression vectors, each expression vector encoding a distinct heterologous protein;
  - b) providing a plate comprising a plurality of wells, wherein one or more wells thereof each contains host cells;

- c) into each of the one or more wells, introducing a distinct expression vector from said collection, thereby producing distinct transfected cells in each well;
- d) culturing the transfected cells under a condition to allow expression of a distinct heterologous protein in each well;
- e) permeabilizing the transfected cells with a permeabilizing agent;
- f) contacting the permeabilized cells in each of the one or more wells with a target protein comprising PGRN attached to a detectable tag;
- g) fixing the transfected cells with a fixing agent after contact with the target protein;
- h) washing the fixed cells; and
- i) after washing the fixed cells, determining the presence of the detectable tag in each of the one or more wells, wherein if the detectable tag is determined to be present in a well, the heterologous protein expressed by the transfected cells in the well is identified as a intracellular protein that binds PGRN.
- 27. The high-throughput screening method of claim 26, wherein the host cells are selected from the group consisting of human embryonic kidney 293T (HEK293T) cell, HEK293F cells, Hela cells, Chinese hamster ovary (CHO) cells, NIH 3T3 cells, MCF-7 cells, Hep G2 cells, baby hamster kidney (BHK) cells, BV-2 (mouse, C57BL/6, brain, microglial) cells, Neuro-2a (N2a) cells and Cos7 cells.
- 28. The high-throughput screening method of 26 or 27, wherein, the detectable tag is selected from the group consisting of green fluorescent protein (GFP), myc, myc-pyruvate kinase (myc-PK), His6, maltose binding protein (MBP), influenza virus hemagglutinin tag, flag tag (FLAG), and glutathione-S-transferase (GST) tag.
- 29. The high-throughput screening method of claim 28, wherein, the target protein is a FLAG tagged PGRN.

- **30**. The high-throughput screening method of claim **29**, wherein, step i) comprises i1) adding an anti-FLAG antibody that is covalently attached to a fluorescent dye into the wells containing the transfected cells to immunostain the FLAG tagged PGRN, if present, and i2) detecting the fluorescent dye using a fluorometer or an imager in a high-throughput manner.
- 31. The high-throughput screening method of claim 30, wherein the fluorescent dye is cyanine dye.
- 32. The high-throughput screening method of claim 28, wherein, the target protein is an HA tagged PGRN.
- 33. The high-throughput screening method of claim 32, wherein, step i) comprises i1) adding an anti-HA antibody that is covalently attached to a fluorescent dye into the wells containing the transfected cells to immunostain the HA tagged PGRN, if present, and i2) detecting the fluorescent dye using a fluorometer or an imager in a high-throughput manner.
- **34**. The high-throughput screening method of claim **33**, wherein the fluorescent dye is Dylight 650.
- **35**. The high-throughput screening method of any one of claims **26-34**, wherein between steps d) and e), the method further comprises, fixing the cultured cells with the fixing agent and washing the fixed cells.
- **36**. The high-throughput screening method of any one of claims **26-35**, wherein the fixing agent is selected from glutaraldehyde, methanol and paraformaldehyde.
- 37. The high-throughput screening method of any one of claims 26-36, wherein between steps h) and i), the method further comprises permeabilizing the fixed cells with a permeabilizing agent and washing the permeabilized cells.
- **38**. The high-throughput screening method of any one of claims **26-37**, wherein the permeabilizing agent is selected from the group consisting of Saponin, Triton X-100, lysolecithin, n-Octyl-B-D-glucopyranoside, and Tween 20.

\* \* \* \* \*