



US 20250263740A1

(19) **United States**

(12) **Patent Application Publication**
AIDA et al.

(10) **Pub. No.: US 2025/0263740 A1**

(43) **Pub. Date: Aug. 21, 2025**

(54) **NON-REPLICATING BOVINE INFECTIOUS
LYMPHOMA VIRUS (BLV) AND CELLS FOR
PRODUCING SAME**

Publication Classification

(51) **Int. Cl.**
C12N 15/86 (2006.01)
A61K 39/00 (2006.01)
A61K 39/21 (2006.01)
(52) **U.S. Cl.**
CPC *C12N 15/86* (2013.01); *A61K 39/21*
(2013.01); *A61K 2039/5254* (2013.01); *C12N*
2740/14021 (2013.01); *C12N 2740/14034*
(2013.01); *C12N 2740/14043* (2013.01)

(71) Applicant: **THE UNIVERSITY OF TOKYO,**
Tokyo (JP)

(72) Inventors: **Yoko AIDA,** Tokyo (JP); **Samy Gamal
Alsaied Mohamed METWALLY,**
Tokyo (JP); **Ryosuke MATSUURA,**
Tokyo (JP)

(21) Appl. No.: **18/857,565**

(22) PCT Filed: **Apr. 19, 2023**

(86) PCT No.: **PCT/JP2023/015633**

§ 371 (c)(1),

(2) Date: **Mar. 14, 2025**


(30) **Foreign Application Priority Data**

Apr. 20, 2022 (JP) 2022-069710

(57) **ABSTRACT**

An object of the present invention is to provide a novel non-replicating bovine leukemia virus (BLV) and a producing cell thereof. According to the present invention, there is provided a bovine leukemia virus (BLV) in which at least a part of the function of a pol gene is deficient. Also, according to the present invention, there is provided a non-replicating BLV-producing cell comprising a gene of a BLV in which at least a part of the function of a pol gene is deficient. The present invention is advantageous in that it can provide a BLV vaccine which is highly immunogenic, and is highly safe without replicating in an infected subject.

Specification includes a Sequence Listing.






FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14

NON-REPLICATING BOVINE INFECTIOUS LYMPHOMA VIRUS (BLV) AND CELLS FOR PRODUCING SAME

CROSS-REFERENCE TO RELATED APPLICATION

[0001] The present application enjoys the benefit of priority from the prior Japanese Patent Application No. 2022-69710 filed on Apr. 20, 2022, the entire disclosure of which is incorporated herein by reference.

TECHNICAL FIELD

[0002] The present invention relates to a non-replicating bovine leukemia virus (BLV) and a producing cell thereof.

BACKGROUND ART

[0003] The bovine leukemia virus (BLV) is a causative virus of enzootic bovine leukosis (EBL) which is a malignant B lymphoma, and is a retrovirus which is integrated as a provirus into the DNA of a host cell. About 70% of BLV-infected cattle remain presymptomatic and healthy, while about 30% thereof have persistent lymphocytosis, and, after a long period of latency, about a few percent thereof develop bovine leukosis (Non-patent Documents 1 and 2).

[0004] Although EBL has been increasing in recent years, a vaccine effective in preventing EBL has not yet been developed. One of the main factors for this is that the amount of the virus to be produced from BLV-infected cells is extremely small, about $1/1,000$ to $1/10,000$ of the amount of AIDS virus to be produced, which belongs to the same Retrovirus family, thereby making it difficult to develop an inactivated vaccine or a cell-derived vaccine. Attempts have been made so far, for example, to develop expression vectors for increasing the amount of BLV to be produced (Patent Document 1). However, there is a risk that, after infection with BLV which is a retrovirus, the viral gene will be integrated into the genome of a host, and therefore problems with practical use of a live vaccine still remain.

REFERENCE LIST

Patent Documents

[0005] Patent Document 1: JP 2019-24351 A

Non-Patent Documents

[0006] Non-Patent Document 1: Gillet NA, et al., *Retrovirology*, 2007, 4:18.

[0007] Non-Patent Document 2: Aida Y, et al., *Frontiers in Microbiology*, 2013, 4:328.

SUMMARY OF THE INVENTION

[0008] An object of the present invention is to provide a novel non-replicating bovine leukemia virus (BLV) and a producing cell thereof.

[0009] The present inventors have now succeeded in establishing a virus-producing cell line that produces an infectious but replication-incompetent bovine leukemia virus (BLV). As a result of analysis of the gene sequence of the virus produced by the cell line, the present inventors have also found that a part of the pol gene is deleted. Further, the present inventors have found that a cell having introduced thereto a BLV gene in which a part of the pol gene

is deleted (pBLV-416ΔRT-introduced cell) has increased expression level and release amount of the virus produced and also has enhanced syncytium formation ability, as compared with a control cell (pBLV-416-introduced cell). The present inventors have also found that, upon inoculation of mice with the virus, an anti-p24 antibody increased over time, whereas no BLV gene was detected. As a result of studies using a non-replicating BLV-producing cell line (PK15-BLVΔRT cell line), the present inventors have also found that the virus produced from the cell line can be useful as a non-replicating BLV vaccine. The present invention is based on the findings.

[0010] According to the present invention, the following inventions are provided.

[0011] [1] A bovine leukemia virus (BLV) in which at least a part of the function of a pol gene is deficient.

[0012] [2] The virus according to [1], wherein the function of the pol gene is the function of a reverse transcriptase and/or an integrase.

[0013] [3] The virus according to [1] or [2], wherein at least a part of the pol gene is mutated.

[0014] [4] The virus according to [3], wherein the mutation is a deletion of 1 to 1643 bases in a reverse transcriptase region of the pol gene and/or 1 to 894 bases in an integrase region of the pol gene.

[0015] [5] The virus according to [3], wherein the mutation is a mutation in bases corresponding to 1st to 1643rd bases and/or bases corresponding to 1644th to 2537th bases from the 5' side in a base sequence of the pol gene as shown in SEQ ID NO: 2.

[0016] [6] The virus according to any one of [1] to [5], which does not replicate in an infected subject.

[0017] [7] A non-replicating bovine leukemia virus (BLV)-producing cell, comprising a gene of a BLV in which at least a part of the function of a pol gene is deficient.

[0018] [8] A method for producing a non-replicating BLV, comprising the step of culturing the cell according to [7].

[0019] [9] A BLV vaccine comprising the virus according to any one of [1] to [6].

[0020] [10] A method for preventing or treating a BLV, comprising the step of inoculating a subject excluding a human with the virus according to any one of [1] to [6] or the vaccine according to [9].

[0021] [11] A method for producing a non-replicating bovine leukemia virus (BLV)-producing cell, comprising the step of introducing, into a host cell, a gene of a BLV in which at least a part of the function of a pol gene is deficient.

[0022] The present invention is advantageous in that it can provide a bovine leukemia virus (BLV) vaccine which is highly immunogenic, and is highly safe without replicating in an infected subject.

BRIEF DESCRIPTION OF THE DRAWINGS

[0023] FIG. 1 shows the genetic constitution of a wild-type BLV, and shows a deleted portion of a pol gene of a CMVΔU3-pBLV-416ΔRT plasmid (pBLV-416ΔRT) in comparison with a CMVΔU3-pBLV-416 plasmid (pBLV-416).

[0024] FIG. 2A shows results of PCR amplification of pol gene regions of pBLV-416ΔRT and pBLV-416, respectively. The negative control used was a pBluescript II KS(-) plasmid which was an empty vector, and the positive control

used was an FLK-BLV cell which was a persistently BLV-infected cell. FIG. 2B shows results of Western blotting using a BLV-infected bovine serum (left) and a non-BLV-infected bovine serum (right) for expression of viral proteins produced by a pBLV-416 Δ ART-introduced cell and a pBLV-416-introduced cell, respectively. The negative control used was a cell introduced with a pBluescript II KS(-) plasmid which was an empty vector, and the positive control used was an FLK-BLV cell which was a persistently BLV-infected cell.

[0025] FIG. 3 shows results of an immunofluorescent antibody method for intracellular localization of the viral proteins in the pBLV-416 Δ ART-introduced cell and the pBLV-416-introduced cell, respectively. The negative control used was a cell introduced with a pBluescript II KS(-) plasmid which was an empty vector, and the positive control used was an FLK-BLV cell which was a persistently BLV-infected cell.

[0026] FIG. 4 shows results of Western blotting for expression levels of the viral proteins in the pBLV-416 Δ ART-introduced cell and the pBLV-416-introduced cell, respectively. The negative control used was a cell introduced with a pBluescript II KS(-) plasmid which was an empty vector, and the positive control used was an FLK-BLV cell which was a persistently BLV-infected cell. As a result of a Student's t-test, * indicates $p < 0.05$ and ** indicates $p < 0.01$.

[0027] FIG. 5 shows results for syncytium formation ability in the pBLV-416 Δ ART-introduced cell and the pBLV-416-introduced cell, respectively. The negative control used was a cell introduced with a pBluescript II KS(-) plasmid which was an empty vector. As a result of a Student's t-test, ** indicates $p < 0.01$ and *** indicates $p < 0.001$.

[0028] FIG. 6 shows results for cell-to-cell infection ability in the pBLV-416 Δ ART-introduced cell and the pBLV-416-introduced cell, respectively. The negative control used was a cell introduced with a pBluescript II KS(-) plasmid which was an empty vector, and the positive control used was an FLK-BLV cell which was a persistently BLV-infected cell.

[0029] FIG. 7A shows release amounts of the viral protein produced by the pBLV-416 Δ ART-introduced cell and the pBLV-416-introduced cell, respectively. FIG. 7B shows the reverse transcriptase activities of the viral protein produced by the pBLV-416 Δ ART-introduced cell and the pBLV-416-introduced cell, respectively. FIG. 7C shows results for syncytium formation ability in the pBLV-416 Δ ART-introduced cell and the pBLV-416-introduced cell, respectively. FIG. 7D shows results of Western blotting for the viral protein released from the pBLV-416 Δ ART-introduced cell and the pBLV-416-introduced cell, respectively. The negative control used was a cell introduced with a pBluescript II KS(-) plasmid which was an empty vector, and the positive control used was an FLK-BLV cell which was a persistently BLV-infected cell. As a result of a Student's t-test, *** indicates $p < 0.001$.

[0030] FIG. 8 shows the genetic constitution of a wild-type BLV, and shows a deleted portion of a pol gene of a CMV Δ U3-pBLV-IFART plasmid (pBLV-IFART) in comparison with a CMV Δ U3-pBLV-IF plasmid (pBLV-IF).

[0031] FIG. 9A shows results of Western blotting for expression levels of the viral proteins in the pBLV-IFART-introduced cell and the pBLV-IF-introduced cell, respectively. FIG. 9B shows results for syncytium formation ability in the pBLV-IFART-introduced cell and the pBLV-IF-introduced cell, respectively. FIG. 9C shows results of

measuring amounts of the virus in culture supernatants of the pBLV-IFART-introduced cell and the pBLV-IF-introduced cell, respectively, by Capture ELISA. The negative control used was a cell introduced with a pBluescript II KS(-) plasmid which was an empty vector, and the positive control used was an FLK-BLV cell which was a persistently BLV-infected cell. As a result of a Student's t-test, * indicates $p < 0.05$, ** indicates $p < 0.01$, and *** indicates $p < 0.001$.

[0032] FIG. 10A shows a schedule of an experiment in which mice were inoculated with the viral protein produced by the pBLV-416 Δ ART-introduced cell and the pBLV-416-introduced cell. FIG. 10B shows results of a Nested PCR method for BLV genes in blood cells 3 weeks after inoculation. FIG. 10C shows results of co-culturing a spleen cell extracted from mice and a CC81-GREMG cell. FIG. 10D shows results of quantifying anti-p24 antibodies from 0 to 6 weeks after inoculation by ELISA. The negative control used was PBS, and the positive control used was an FLK-BLV cell which was a persistently BLV-infected cell. As a result of a Student's t-test, * indicates $p < 0.05$.

[0033] FIG. 11 shows the genetic constitution of a wild-type BLV, and shows a deleted portion of a pol gene of a BLV in a PK15-BLV Δ ART cell line in comparison with a CMV Δ U3-pBLV-416 plasmid (pBLV-416).

[0034] FIG. 12 shows results of Western blotting for expression levels of the viral proteins in the PK15-BLV Δ ART cell line. The positive control used was an FLK-BLV cell which was a persistently BLV-infected cell.

[0035] FIG. 13 shows results of an immunofluorescent antibody method for intracellular localization of the viral protein in the PK15-BLV Δ ART cell line. The positive control used was an FLK-BLV cell which was a persistently BLV-infected cell.

[0036] FIG. 14A shows a release amount of the viral protein produced by the PK15-BLV Δ ART cell line. FIG. 14B shows a standard curve created using His-p24 antigen as a standard.

DETAILED DESCRIPTION OF THE INVENTION

Virus and Method for Producing Same

[0037] The "bovine leukemia virus (BLV)" is a causative virus of enzootic bovine leukosis (EBL), which is a malignant B lymphoma, and is a retrovirus which is integrated as a provirus into the DNA of a host cell. The genome of a wild-type BLV has a full length of about 8,720 bp, and is composed of gag, pro, pol and env genes that encode structural proteins; rex and tax genes that encode regulatory proteins; and R3 and G4 genes that encode accessory proteins, in a form in which the genes are sandwiched between two identical long terminal repeat (LTR) sequences (see FIG. 1). The gag gene encodes a protein involved in viral particle formation, the pol gene encodes a reverse transcriptase and an integrase, and the env gene encodes a coat protein involved in adsorption onto and entry into a host cell. An example of the gene sequence of the wild-type BLV is shown in Table 1 (the gene sequence of the wild-type BLV shows a sequence from the 5' LTR to the 3' LTR, and the underlined part shows a sequence encoding the pol gene). In the present invention, a wild-type BLV sequence other than the gene sequence indicated in Table 1 can be used.

TABLE 1

Gene sequence of wild-type BLV (SEQ ID NO: 1)

TGTATGAAGATCATGCCGACCTAGGCGCCGCCACCGCCCCGTAAACC
 AGACAGAGACGTCAGCTGCCAGAAAAGCTGGTGACGGCAGCTGGTGGC
 TAGAATCCCCGTACCTCCCCAATTCCCTTTCCGAAAATCCACAC
 CTTGAGCTGCTGACCTCACCTGCTGATAAACTAATAAATGCCGGCCC
 TCTCGAGTTAGCGGCACCCAGAGCTTCTTCTCCTGAGACCTCGTGC
 TCAGTCTCGGTCTGAGCTCTCTTGCTCCCGAGACCTTCTGGTCCGC
 TATCCGGCAGCGGTGAGTAAGGCAAAACACGGTTTGGAGGGTGGTTC
 TCGGCTGAGACCACCGCAGCTCTATCTCCGGTCTCTGACCGTCC
 ACGTGGACTCTCTCTTTGCCCTCTGACCCCGCGCTCCAAGGCGCTCT
 GGCTTGACCCCGCTTTGTTTCTGCTTACTTTCTGTTTCTCGCGGC
 CCGCGCTCTCTCTTCCGGCGCCACTAGCGGCCAGGAGAGACCGCAA
 ACAATTTGGGGGCTCGTCCGGGATTGATCACCCCGGAACCTTAACAAT
 CTCTGGACCCACCCCTCGGCGGCATTTTGGGTCTCTCTTCAAAATTA
 TATCATGGGAAATCCCCCTCTATTAACCCCGCGCTGGTATCTCCCC
 CTCAGACTGGCTCAACCTTCTGCAAGCGCGCAAAGGCTCAATCCGCG
 ACCCTCTCTAGCGATTTTACCGATTTAAAGAAATACATCCATTGGTT
 ATATAAGACCCAGAAAAACCATGGACTTTCACCTCTGGTGGCCAC
 CTCATGTCCACCCGGGAGGTTCCGCGGGTTCCTTGTCTTGGCCAC
 CCTAAACGAAGTGCTCTCAACGAAGGGGCGCCCGGGTGTCATCGGC
 CCAGAAGAACACCCCGCTTATGACCCCGCGCGTTTGGCCAAT
 CATATCTGAGGGAATCGCAACCGGCATCGTCTTGGGCACCTCGAGA
 ATTACAAGATATCAAAAAGGAAATGAAAAAAGGCACCGGTTTCGCA
 AGTATGGATACAAACACTACGACTTGCATCTGCGAGGCCACCTAC
 TCCGGCTGACTAGAACAACTTTCGCAATATATTGCTTCCCGGTGCA
 CCAACCGGCCATATGACCGAGCTTAACGGCAGCAATTGCCCGCGTCA
 CGCGCCCAACACCCCTTAAACCCCAAAACGGGACCTTAAC
 CCAACAATCAGCTCAGCCCAACCGCGGGGATTTAGAACTCAATATCA
 AAACCTCTGGCTTCAGGCTCGAAAAATCTCCCTACTCGTCTTAGT
 ACAACCTTGGTCCACCTCTGCAAGGCCCGCGCAAGCTATGTAGA
 GTTTGTCAACCGGTACAAATTTTATTAGCTGACAACTTCCCGACGG
 AGTCCTTAAGGAACCCATTATGACTCCCTTAGTTATGCAATGCTAA
 CAAAGAGTGTCAAGAAATTTGCAAGGGCGAGGCTAGTGGCCGGCC
 GGTGGGCAAAAACATGCGAGGTTGCGCACATTGGGCCCAAGATGAA
 ACAGCTCGCAATTTCTGCTCACACCCCGAGGCCAAGATGCCCCGGCC
 TCGGCACACCGGCTCCCGGAACAGCTTCCCGAGGACCTGCTATCGAT
 CCTCAAGAGAGCCATTGGGCCCGGGATTGTCTTACCAAGGCCACCGG
 CCCCCCTCGGGACCTTCCCCCATATGTAAAGATCTTCCCATTTGAA
 AGGACTGTCCAAACCTCAATCAAAACCTTAATAGGGGGGACTT
 AGCGCCCCCAACCATTAACCCATAACAGATTCTCTAGTGAGGCC
 GAATTAGAATGCTTACTTTCTTCTCTGCTCGCAGCCGCTCCCTCC
 GTGGCTGTATACCTGTCTGGCCCTTGGCTGCGAGCCTCTCAGAATCAA
 GCGCTCATGCTTGTGGACACCGGGGCTGAAATACGGTTCTCCACAA
 AATTGGCTGGTTCCGGATTACCCAGGATCCCCCGCGAGTGCTCGGA
 GCAGGGGAGTCTCCCGGAACAGATCAATTTGGCTTACAGGCCCTCTG
 ACCCTGGCTCTTAAACAGAGGGTCCCTTTATCACCATCCCAAAAT
 TTAGTTGACACTTTTCGACAAATGGCAAAATTTTAGGACGGGACGCTCT
 TTCGCTTACAGGCTCTTATCTTCTTACCTGAGGAAGTACGCCCTCT
 GTGGTAGGGCTTGGATGCCCCCGAGCCACATTGGATTAGAATAT
 CTGCCCCCTCCACCTGAAGTGCCCTCAATTTCCCTTTAACTAGAAGCC
 TCCAGGCTCTCAAGACCTTGCTTACCTGCTCTGAGGCGAGTTATA
 TCTCCCCCTGGGACGGGCCAGGCAATAATCAGTCTTCCCGGTACGGA
 AACCAATGGCGCTGGAGGTTTGTGATGACCTACGAGCTACAAATG
 CTCTTACAAGGCCATTCCGGCAGTCTCCCGGACCGCCAGACCTTA
 CCGCTATCCCTACACACTTCCACATATCATTGCTTAGATCTCAAG
 ACGCTTCTTCCAGATTCCAGTCGAAGACCGCTTCCGCTCTACTTTG
 CTTTACCCTCCCTACCCCGGGGACTCCAACCTCATAGACGCTTTG
 CTTGGCGGGTCTTACCTCAAGGCTTCAATTAACAGCCAGCTCTTTTCG
 AACGAGCACTACAGGACCTCTTTCGCCAAGTTTTCGCGCCTTTTCCC
 AGTCTCTTCTGGTGATCTTATGGAGCATATCTTATGCTTCGCTCA
 CAGAAGAACAGCGGTACAAATGTTATCAAGCCCTGGCTGCCCGCTCC
 GGGACCTAGGTTTTCAGGTGGCATCCGAAAAGACTCGCCAGACGCTT
 CCGCGTCCCTTCTTGGGCAAACTTGCACCTTCTACTCTTCCCTTAAAGCA
 TCGATGACCTTAGGGCCATCATCCAGCTTTCGCCGAACAGCTGCAAG
 GCATTGACAGCTTCGACAGCCCTGTCCATAACGCAAGATCTAGAT
 ATACAGAGCAAGAACCCCTGAGCTAGCTACCTTACCTTAAAGGCA
 TCGATGACCTTAGGGCCATCATCCAGCTTTCGCCGAACAGCTGCAAG
 GCATTGACAGCTTCGACAGCCCTGTCCATAACGCAAGATCTAGAT
 ATACAGAGCAAGAACCCCTGAGCTAGCTACCTTACCTTAAAGGCA
 GGTCCACCTTGGTACTCTTCCAAAAGGGCGCTCAATTTCCCTTGGCT
 ACTTCCAGACCCCTTACTGACAAACCAAGCTTACCTTGGGCGCTCT
 TTCTCTGCTGGGATGCCCAATACCTGCGAGCTCAGGCTTAAAGCTCGT
 ATGCAAGCCCATCTTAAATATTATCAAACTTCTTCTTAAACTTCTCT
 TAGACAAATTTGATTTCAATCATCTGAGGACCTCGAGTCCAGGAGTTGC
 TGCAATTTGGGCCAGATTTCTCTCAGGGAATACAGCCCCGGGCT
 CTTGGAAGACCTTAATCACCAGGGCAGAGGTTTTTTTACGCCCCAGT

TABLE 1-continued

Gene sequence of wild-type BLV (SEQ ID NO: 1)

TCTCCCCGTATCCGATTCTCGGGCCCTTTGCCTCTTTAGTGACGGGG
 CTAAGGACGAGGAGCATATTGCTTGTGGAAGGACCCACTTTTAGACT
 TTCAGGCGGTTCGGCTCCAGAACTCCGCTCAAAAGGGAGAACTAGCAG
 GTCTCTTGGCGGGCTTAGCAGCCCGCCCGCTGAACCTGTAATATAT
 GGGTAGATTCCAAATACCTGTACTCTTGTCTCAGAACCTTAGTCTGG
 GAGCTTGGCTTCAACCTGACCCCGTACCTCTACGCCCTCTATATA
 AAAGCCTCTCCGACATCCAGCAATCTTGTGTGGCATGTCCGGAGCC
 ACTCTTCAGCATCCACCTTATGCTTCCCTGAACAATATGTAGATC
 AACTGCTTCCCTTAGAACTCCAGAGCAATGGCATAAGCTCACCACAT
 GCAACTCTCGGGCCTTGTCTCGATGGCCGAACCCACGTATCTCTGCCT
 GGGACCCCGTTCGCCCGCTACGCTGTGTGAACTGTCAAAAGCTTA
 ATCCAACTGGAGGAGGAAAGATGGCAACTATTTCAGAGAGGGTGGGCC
 CGAACCATATTGGCAGGCGCATATAACCATTATAAATACAAACAGT
 TCACCTACGCTCTGCATGTGTTTGTAGATCTACTCTGGAGCTACTC
 ATGCTCTCGGCAAGCGTGGGCTCACCACTCAAAAGACCATTTAGGGGCC
 TTCTTGGGCCATAGTGATCTGGGTCTGCCAAAAAGCTAAACACTG
 ACCAAGGTGCAAACTACACCTCCAAACCTTTGTCAAGTTTGTGCCAGC
 AGTTTCGAGTTTCCCTTTCTCATCATGTTCTCTCAACCCACAAAGTT
 CGGGGTAGTAGAACGGACAAATGGACTGTCTAAACTTCTTCTATCTA
 AATATCACCTAGACGAACCCACCTTCCATGACTCAGGCGCTTCTC
 GAGCCCTCTGAGCTCACAACTAGATTAACTCTTACCAATTTCTAAAGA
 CCAGATGGGAGCTACACCATTCACCCCACTTGTCTGTCATTTACAGG
 GCGGAGAAACACCCCAAGGGCTCTGATAAACTCTTTTGTACAAGCTCC
 CCGGGCAAAACAATCGTGGTGGCTAGGACCACTCCCGGCCCTAGTCT
 AAGCCTCGGGAGGCGCTCTCTGGCTACTGACCCCGCGTGTGGGTTCT
 CTTGGCGTTTGTCTGAAAGCCTTCAATGCTTAAAGAACGAGCTCCCG
 AAGACGCCCAACCCGATCATCAGATGGGTAACTCTCACTCTCACTCT
 CCTCGCTCTCTCTCGGCCATCCAGACTTGGAGATGCTCCCTGTCCCT
 AGGAACCAACAATGGATGACAGCATATAACCAAGAGCAAAATTTTC
 CATCTCATTGACCAAACTAGAGGCTCATAATCAGTCACCTTTCTG
 TGCCAAGTCTCCAGATACACCTTGGACTCTGTAATGGCTATCCCTAA
 GATCTACTGGCCCCCCCCCAAGGGCGCGCGCTTGGAGCCAGGGC
 CATGGTCACATATGATTGCGAGCCCCGATGCCCTTATGTGGGGCAGA
 TCGCTTCGACTGCCCCCACTGGGCAATGCTCCAGGCCGATCAAGG
 ATCTTTTATGTCAATCATCAGATTATTCTCTGATCTCAAAACAATG
 TCATGGAATTTTCACTTTAACTGGGAGATATGGGATATGATCCCCCT
 CATCACCTTTTCTTACATAAGATCCCTGATCCCCCTCAACCCGACTT
 TCCCGAGTTGAACAGTGACTGGGTCTCCCTCTGTCAGATCTGGGCCCT
 GCTTTTAAATCAACAGCACGGGCTTCCAGACTGTGCTATATGTTG
 GGAACCTTCCCTCCCTGGGCTCCGAAATATTAGTATATAACAAAA
 CATCTCCAGCTCTGGACCCGGCCCTCGCCCTCCGGACCGCCAAATCTT
 CTGGGTCAACACGCTCTCGTTTAAACACCCCAAGGATGGACACACCC
 TTCCAGAGGTTGTTGTTCAATGTTTCTCAAGGCAACGCTTGTATT
 ACCTCTATCTCCCTGGTTAATCTCTCTACGGCTTCTCCGCCCTCTC
 TACCCGGGTGAGAGTAGTCCCGTTCGAGCCCTGACCTTAGGCTTAGC
 CCTGTAGTGGGGCTCACTGGAATTAATGTGGCCGTGTCTGCCCTTAG
 CCATCAGAGACTCACCTCCCTGATCCAGCTTCTGGAGCAAGATCAGCA
 ACGCTTGATCAGCAATTAACAGACCCACTATAATTTGCTTAATGT
 GGCTCTGTGGTTGCCGGAACCGAGGGGGCTTGGATTGTTATACAT
 CCGCTGGGTTTCAAGCCTATGTCCCACTAATTAAGCAAGCTCTGCTG
 TTTCTGCGCATTAATAAGTACTCATTATCCGCTCGGTGATCTCCA
 GCCTCTCTCGCAAGAGTCTCTACAGACTGGCAGTGCCCTGGAATTTG
 GGATCTGGGGCTCACTGCTGGGTGCGAGAAACCATCTTCTGTCTT
 AAGCTGTCTTATAGCCCTTTTGTCTCTCTGGCCCCCTGCTCT
 GATAAATGCTTACCTCTCGCTTTTAAAGCTCCTCCGGCAGGCTCC
 CCATCTCCCTGAAATCTCTTAAACCCCAACCCGATCTGATTATCA
 GGCTTGTCTACCATCTGCACAGAGATCTACTCTCACTCTCCCCCGC
 CAAACCCGATTACATCAACCTCCGACCTGTCTTGATACCCCGCGCT
 TTCACGACCCCGAGGCTGTGGTGGTGCAGTGGGTAGGCTTAAAGATG
 AGTGTACCATCACAGCTCTTCTTGTGCGCGCACCGAGTTGGAACA
 CAGCCCTACCTGAGCTCTCTGAGTGCATGACTGAGTGTAGCGCAGA
 GAGATTGTGCTTCTGCTGTGCTGCTGCTGCTGCTGCTGCTGCTGCTG
 GGGTTGCGCCCTTCCGTTGCTGTGACACAGATAAGACCTCTCTCAC
 TTCTGCTTACCATCCCCCTGCCAGCTTGGTCTAGTGGAAAGAACTA
 ACGCTGACGGGGCGATTCTTGCAGCTGTGCTGCTGCTGCTGCTGCTG
 TGCTGGGGATAAGGTGTGGCCCTTAGCACACAGTCTCTGCGCTTTT
 GGGTTCGAATCTTCCACAGCAGCTTCCGCTTTTACGCCCTGTGTGCA
 CACCTTTCTAGAGATACCTGAAATCTCAGCTGCGACCTGAGGAGG
 GTTGTGGCTCAGAGGTTAAATAGCTCGGGCCGAACCTCTCTTCTT
 TTTATTCACCTCGCAAGGCCCGGGTTCTGAGCCCCCTAACGGAGG
 TTAATAATTTCTCTACTAGGGGATGCTCGGGTCCAGGTGTCACAAAT
 ATCTCTTCAAAAGGCTCTGATGAACGCTTTCCTATGTAACAGCCCC
 AGCAGAGACATTCAGGCACATCCAGCAGCATTTGGGCCGCTTTTCT
 AACAGTGCCATAAAGCTCCCTTCCGTTTCCCAACAGGCTGCTGCT
 TCTTCTATTTCACCTCGGCACCGACTCCCCCGCGAGCCTTCGAGC

TABLE 1-continued

Gene sequence of wild-type BLV (SEQ ID NO: 1)
<p>TCTTCGGGATCCATTACCTGATAACGACAAAATTATTTCTTGCTTTT AAGCAAGTGTGTGGTTGGGGGCCCCACTCTCTACATGCTGCCCGG CCCTGGTTTGTCCATGATGTCACCATCGATGCTGGTGCCCCCTCT GCGGGCCCCATGAGCGACTCCAATTGCGAAGGATCGACACCACTGCA CCTGCGAGACCCACCGTATCAACTGGACCGCGATGGACGACCTTGGC GCCTCAATGGAACGTGTGTTCCCTCGACTGCATGTCTCCGAGACCCGCC CCCAAGGGCCCCGACGACTCTGGATCAACTGCCCCCTTCCGGCCGTTT GGCTCAGCCCCGCGGCTTCACTTTCCTCCCTTCGAGCGGTCCCTT TCCAGCCCTACCAATGCGCAATTGCGCTCGGCTCTAGCGACGGTTGCC CCATTATCGGGCAGCGCTTCTTCCCTGGAACAACCTAGTAACGCATC CTGTCTCGGAAAAGTCTTATATTAATCAAATGGCAATTTTTCCT TACTCCCTCCTTCGATACCTCCTTGTGGACCCCTCCGGCTGTCGG TCTTTGCCCCAGACACAGGGGAGCCATACGTTATCTCTCCACCTTT TGACGCTATGCCCGAGCTACTTGTATTCTACCCCTAGGCGAGCCCTCT CTCCTAATGTCCCATATGCGCCTTCCCGGGACTCTAATGAACCCC CCCTTTCAGAATTTCGAGCTGCCCTTATCCAAACGCCCGCTGTCT GGTCTGTCCCCGCGATCGACCTATTCTAACCGGTCCCTTCCCAT GCGACCGGTTACACGTATGGTCCAGTCTCAGGCTTACAGCGCTTCC TTCATGACCTTACGCTAACCTGGTCCGAATTGGTTGCTAGCAGAAAA TAAGACTTGATTCCCTTAAATTAACAACTGCTAGAAAAATGAATGGC TCTCCGCTTTTTCGAGGGGAGTCATTGTATGAAAGATCATGCCG ACCTAGGCGCGCCACCGCCCGTAAACAGACAGAGACGTACAGTGC CAGAAAAGCTGGTGACGCGAGCTGGTGGCTAGAATCCCGTACCTCCC CAACTTCCCTTTCGCGAAAATCCACACCTGAGCTGCTGACCTCAC CTGCTGATAAACTAATAAATGCCGGCCTGTGCGAGTTAGCGGCACCA GAAGCGTTCTTCTCTGAGACCTCGTGCTCAGCTCTCGGTCTGAGC TCTCTGTCTCCGAGACCTTCTGGTCCGCTATCCGCGAGCGGTGAGGT AAGGCAAAACCGGTTTGGAGGGTGGTTCTCGGCTGAGACCAACCGCA GCTCTATCTCCGGTCTCTGACCGCTCCACGCTGAGCTCTCTCTTTG CCTCTGACCCCGCTCCAAGGGCGTCTGGCTTGACACCCGCTTGT TTCTGTCTTACTTCTGTTTCTCGCGCCCGCTCTCTCTCGCG GCCACTAGCGCGCAGGAGAGACCGGCAACA</p>

[0038] According to the present invention, there is provided a bovine leukemia virus (BLV) in which at least a part of the function of a pol gene is deficient. In the present invention, the sentence that “at least a part of the function of a pol gene is deficient” means that at least a part of the function of the pol gene is deficient in comparison with the wild-type BLV, and specifically means that at least a part of the function of a reverse transcriptase and/or an integrase is deficient. That is, the virus of the present invention does not express the reverse transcriptase and/or the integrase, or, even if it expresses these enzymes, the enzymes do not function normally. The virus does not replicate in an infected subject. For this reason, the virus of the present invention is sometimes referred to as “non-replicating BLV” in the present specification.

[0039] Whether or not at least a part of the function of the pol gene is deficient may be determined by preparing a BLV in which at least a part of the pol gene is mutated, and determining whether or not the prepared virus expresses proteins encoded by the pol gene or whether or not the proteins normally function due to the presence or absence of the enzymatic activities of the proteins. The presence or absence of the expression of the proteins can be determined, for example, by a known method such as an immunological assay using an antibody against each of the reverse transcriptase and integrase encoded by the pol gene. The presence or absence of the enzymatic activities of the proteins can be determined, for example, by a known method such as a method for measuring reverse transcriptase activity using colorimetric analysis or a method for measuring integrase activity using the strand transfer ability as an index.

[0040] The virus of the present invention is a virus in which at least a part of a pol gene is mutated. In the present invention, the sentence that “at least a part of a pol gene is

mutated” means that at least a part of the base sequence of the pol gene is mutated in comparison with the wild-type BLV, and means a mutation that brings a state where at least a part of the function of the pol gene is deficient. The mutation of the pol gene means a mutation in which a deletion, a substitution, an insertion and/or an addition occur/occurs in at least a part of the base sequence of the pol gene, resulting in a state where at least a part of the function of the pol gene is deficient. The function of the pol gene that is deficient in the virus of the present invention is either or both of the reverse transcriptase function and the integrase function, and is preferably the reverse transcriptase function. That is, the virus of the present invention can be said to be a BLV in which a reverse transcriptase region (RT) and/or an integrase region (IN) encoded by the pol gene are/is mutated, or a BLV having a pol gene in which the reverse transcriptase region (RT) and/or the integrase region (IN) are/is mutated. Typical examples of the virus of the present invention include BLVs having a pol gene in which a part of RT (e.g., 1 to 1643 bases, 1 to 1600 bases, 1 to 1500 bases, 1 to 1400 bases, 1 to 1300 bases, 1 to 1200 bases, 1 to 1100 bases, 1 to 1000 bases, 1 to 900 bases, 1 to 850 bases, 1 to 800 bases, 10 to 750 bases, 20 to 700 bases, or 40 to 650 bases) and/or a part of IN (e.g., 1 to 894 bases, 1 to 850 bases, 1 to 800 bases, 10 to 750 bases, 20 to 700 bases, or 40 to 650 bases) is deleted. The deletion of bases in a part of RT and/or IN may be a deletion of continuous bases or a deletion of discontinuous bases.

[0041] In the virus of the present invention, the mutation of a part of the base sequence of the pol gene is not limited as long as at least a part of the function of the pol gene is deficient. When the pol gene is a wild-type pol gene of SEQ ID NO: 2 (Table 2), the mutation of RT can be, for example, a mutation in 1st to 1643rd, 1st to 1600th, 1st to 1500th, 1st to 1400th, 1st to 1300th, 1st to 1200th, 1st to 1100th, 1st to 1000th, 10th to 900th, 20th to 800th, 30th to 700th, or 40th to 650th bases from the 5' side of the base sequence of the pol gene as shown in SEQ ID NO: 2 in Table 2, and the mutation of IN can be, for example, a mutation in 1644th to 2537th, 1650th to 2500th, 1700th to 2400th, 1800th to 2300th, 1700th to 2200th, 1650th to 2100th, 1650th to 2000th, 1650th to 1900th, or 1650th to 1800th bases from the 5' side of the base sequence of the pol gene as shown in SEQ ID NO: 2. When the pol gene is composed of a base sequence other than that of the wild-type pol gene as shown in SEQ ID NO: 2 (Table 2), the mutation can be a mutation in bases corresponding to the bases of SEQ ID NO: 2. For example, the mutation site can be identified by appropriately aligning the base sequence of the pol gene to be mutated with the base sequence of SEQ ID NO: 2. That is, the mutation site can be identified by appropriately aligning the base sequence of the pol gene to be mutated with the base sequence of SEQ ID NO: 2 using a publicly available homology search software or program such as BLAST (Basic local alignment search tool) (Altschul et al., J. Mol. Biol. 215:403-410 (1990)), FASTA (Pearson et al., Methods in Enzymology 183:63-69 (1990)), or Smith-Waterman (Meth. Enzym., 164, 765 (1988)).

[0042] The base sequence of the pol gene of SEQ ID NO: 2 in Table 2 is shown in the 5' to 3' direction and corresponds to the underlined base sequence in SEQ ID NO: 1 in Table 1. The number of mutations of RT or the number of mutations of IN is not particularly limited as long as at least a part of the function of the pol gene is deficient. The lower limit value of the number of the mutations can be, for

example, 1, 10, 20, 30, 40, 50, 60, 70, 80, or 90, and the upper limit value of the number of the mutations can be 1,000, 900, 800, 700, 600, or 500. These lower limit values and upper limit values can be combined arbitrarily, and the numerical range thereof can be, for example, 1 to 1,000, 10 to 900, 20 to 800, 30 to 700, or 40 to 600.

[0043] In the virus of the present invention, the mutation of the base sequence of the pol gene may be a mutation spanning RT and IN. When the pol gene is the wild-type pol gene of SEQ ID NO: 2 (Table 2), such a mutation can be, for example, a mutation in 1st to 2537th, 10th to 2500th, 20th to 2400th, 30th to 2300th, 40th to 2200th, 50th to 2100th, 60th to 2000th, 70th to 1900th, 80th to 1850th, 90th to 1800th, 100th to 1750th, 200th to 1700th, 300th to 1650th, 400th to 1650th, 500th to 1650th, 600th to 1650th, 700th to 1650th, 800th to 1650th, 900th to 1650th, or 1000th to 1650th bases from the 5' side of the base sequence of the pol gene as shown in SEQ ID NO: 2 in Table 2. When the pol gene is composed of a base sequence other than that of the wild-type pol gene as shown in SEQ ID NO: 2 (Table 2), the mutation can be a mutation in bases corresponding to the bases of SEQ ID NO: 2. For example, the mutation site can be identified by appropriately aligning the base sequence of the pol gene to be mutated with the base sequence of SEQ ID NO: 2, as described above.

[0044] The number of the mutations spanning RT and IN is not particularly limited as long as at least a part of the function of the pol gene is deficient. The lower limit value of the number of the mutations can be, for example, 1, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400 or 500. The upper limit value of the number of the mutations can be, for example, 2,200, 2,000, 1,800, 1,600, 1,400, 1,200, 1,000, 900, 800, 700, 600 or 500. These lower limit values and upper limit values can be combined arbitrarily, and the numerical range thereof can be, for example, 1 to 2,200, 10 to 2,000, 20 to 1,800, 30 to 1,600, 40 to 1,400, 50 to 1,200, 60 to 1,000, 70 to 900, 80 to 800, 90 to 700, or 100 to 600.

TABLE 2

pol Gene of wild-type BLV (SEQ ID NO: 2)
AAGTGAAGCGCTCCAGGCGCTTCAAGACCTGGTCCATCGCTCTCTGGA GGCAGGTTATATCTCCCGCTGGGACGGCCAGGCAATAATCCAGCTCTC CCGGTACGGAACCAATGGCGCTTGGAGGTTTGTGCATGACCTACGAG CTACAAATGCTCTTACAAAGCCCATTCGGGCACTCTCCCGGACCGCC AGACCTTACCGCTATCCCTACACACCTTCCACATATCATTTGCTTAGAT CTCAAAGACGCTTCTTCCAGATTCAGTCGAAGACCGCTTCCGCTCCT ACTTTGCTTTTACCCTCCCTACCCCGGGGAGCTCCAACCTCATAGAG CTTTGCTGGCGGCTTACCTCAAGGCTTCATTAACAGCCAGCTCTT TTCGAACGAGCACTACAGGAACCTTTCGCCAAGTTTCGCCGCTTTT CCCAGTCTCTTCTGGTGTCTATATGGACGATATCTTATCGCTTCGCC TACAGAAGAAGACGCGGTACCAATGTTATCAAGCCCTGGCTGCCGCTC CGGGACCTAGGTTTTCAGGTGGCATCCGAAAAGACTCGCCAGACGCTT CGCCCGTCCCTTCTTGGGACAAATGGTCCATGAGCAGATTGTACCTA CGAGTCCCTACCTATCTGCGAGATCTCATCCCAATTTCTTACCAA TTACAGGCGGTCTTAGGAGACCTCAATGGGTCTTAGGGGCACACCA CTACCCGCGGCGCTTCAACTTCTTACTCTTCCCTTAAAGGCATCGA TGACCTTAGGGCCATCATCCAGCTTTCGCCGAACAGCTGCAAGGCATT GCAGAGCTTCGAACAGCCCTGTCCTAAGCAAGATAGATATAACG AGCAAGAAGCCCTGTAGCTACGTACACCTAACCCGGGCGGGTCCAC CTGTGTACTCTTCCAAAGGGCGCTCAATTTCCCTGGCTTACTCCAG ACCCCTTGTACTGACAAACAGCCTCACCTTGGGCGCTCTTCTCTCTGC TGGGATGCCAATACCTGCACTCAGGCTTAAAGCTCGTATGCCAAGCC CATACTTAAATATATCACAATCTTCTTAAACTCTCTAGACAATTGG ATTCAATCATCTGAGGACCTCGAGTCCAGGAGTTGCTGCAATTGTGCG CCCAGATTCTCTCAGGGAATACAGCCCCGGGCGCTTGGAAAGACCTT AATCACCAGGGCAGAGGTTTTTTGACGCCCGAGTTCTCCCTGATCCG ATTCTCGCGGCCCTTTCCTCTTTAGTGACGGGCTTAAAGACGAGGAG

TABLE 2-continued

pol Gene of wild-type BLV (SEQ ID NO: 2)
CATATTGCTTGTGGAAGGACCACTTTTAGACTTTTCAGGCCGTTCCGGC TCCAGAATCCGCTCAAAAGGGAGAACTAGCAGGTCTCTGGCGGGCTTA GCAGCCGCCCGCTGAACCTGTAAATATATGGGTAGATTCCAATACC TGTAATCTTGTCTCAGAACCCTAGTTCTGGGAGCTTGGCTTCAACCTGA CCCCGTACCTCTACGCCCTCTATATAAAGCTCTCCGACATCCA GCAATCTTGTGTGGCATGTCCGGAGCCACTCTTCAGCATCCACCCTA TTGCTTCCCTGAACAATTATGTAGATCAACTGCTTCCCTTAGAACTCC AGAGCAATGGCATAAGCTCACCCACTGCAACTCTCGGCCCTGTCTCGA TGGCGGAACCCACGTATCTCTGCCGAGGACCCCGTCCCCCGCTACGC TGTGTGAACCTGTCAAAGCTTAATCCAAGTGGAGGAGGAAGATGCG AACTATTGAGAGAGGTTGGGCCGGAACCATATTGGCAGGCCGATATA ACCCATTATAAATACAAACAGTTACCTACGCTCTGATGTGTTGTAG ATACTTACTCTGGAGCTACTCATGCCCTGGCGAAGCGTGGGCTCACCAC TCAATGACCATTTGAGGGCTTCTTGGGCGCATAGTGCATCTGGGTGCT CCAAAAAGCTAAACACTGACCAAGGTGCAAACTACACCTCAAACCT TTGTGAGTTTTTCCAGCAGTTCGGAGTTTCCCTTTCTCATCATGTTCC CTACAACCCCAAGTTTCGGGTTAGTAGAACGGACAAATGGATGCTCGA AACTTCTTCTATCTAAATATCACCTAGACGAACCCACCTTCCCATGA CTCAGGCCCTTTCTCGAGCCCTCTGGACTCACAATCAGATTAACCTCCT ACCAATTTCTAAAGACGAGTGGGAGCTACACCATTCACCCCGCATGATA GTCTATTGAGAGGCGGAGAACACCCAGGGCTCTGATAAATCTTTT TGTAACAGCTCCCGGGCAAAACATCGTCGGTGGCTAGGACCACTCCC GGCCCTAGTCGAAGCCTCGGGAGGCGCTCTCTGGCTACTGACCCCGC GTGTGGTTCCTTGGCGTTTGTCTGAAAGCCTTCAATGCCTAAGAACG ACGGTCCCGAAGACGCCCAACCGATCATCAGATGGG

[0045] The virus of the present invention may have a region other than the pol gene as it is, in the full length of the sequence between the 5' LTR and 3' LTR of the wild-type BLV as shown in SEQ ID NO: 1, as long as at least a part of the function of the pol gene is deficient and the functions of the BLV genes other than the pol gene are retained. Alternatively, the virus may have a sequence having 80% or more (preferably 85% or more, more preferably 90% or more, even more preferably 95% or more, and particularly preferably 96% or more, 97% or more, 98% or more, 99% or more, 99.5% or more, or 99.8% or more) identity to that of the region (wild type) other than the pol gene. The term "identity" as used herein refers to the degree of identity, for example, when sequences to be compared are appropriately aligned, and means an appearance rate (%) of exactly matched amino acids between the sequences. In the calculation of the identity, for example, the presence of gaps in the sequence and the nature of amino acids are taken into consideration (Wilbur, Natl. Acad. Sci. U.S.A. 80:726-730 (1983)). The alignment can be performed, for example, by using an arbitrary algorithm. Specifically, publicly available homology search software such as BLAST (Basic local alignment search tool) (Altschul et al., J. Mol. Biol. 215: 403-410 (1990)), FASTA (Pearson et al., Methods in Enzymology 183:63-69 (1990)), or Smith-Waterman (Meth. Enzym., 164, 765 (1988)) can be used. The identity can be calculated, for example, using the publicly available homology search program as described above, and can be calculated, for example, using a default parameter in the homology algorithm BLAST (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) of the National Center for Biotechnology Information (NCBI).

[0046] The virus of the present invention can be obtained by allowing a cell having a BLV gene in which at least a part of the function of the pol gene is deficient to produce it. The origin of the cell is not limited, and the cell may be a naturally occurring cell, a cell obtained by self-cloning, or a cell prepared by a gene recombination technique. However,

from the viewpoint that the virus is used as a vaccine, a naturally occurring cell or a cell obtained by self-cloning can be preferably used.

Cell and Method for Producing Same

[0047] According to the present invention, there is also provided a cell comprising a gene of a bovine leukemia virus (BLV) in which at least a part of the function of a pol gene is deficient. The phrase “comprising a gene” as used herein means that the gene is contained in the cell in an expressible manner, and typically means that the gene has been introduced into the cell and is in a transformed state. The cell of the present invention may be a naturally occurring cell, a cell obtained by self-cloning, or a cell prepared by a gene recombination technique, as long as the cell comprises a polynucleotide encoding a BLV in which at least a part of the function of the pol gene is deficient. However, from the viewpoint that the virus is used as a vaccine, a naturally occurring cell or a cell obtained by self-cloning can be preferably used.

[0048] According to another aspect of the present invention, there is provided a method for producing the cell of the present invention. When the cell of the present invention is produced by a gene recombination technique, it can be prepared, for example, by introducing into a host cell an expression vector having incorporated therein a polynucleotide encoding a BLV in which at least a part of the function of the pol gene is deficient. To make the function of the BLV pol gene deficient, the entire or a part of RT and/or IN encoded by the pol gene may be deleted. Also, the pol gene may be mutated by substitution, deletion, insertion and/or addition of bases to prevent expression of a normal reverse transcriptase or integrase. Alternatively, a foreign gene may be inserted into RT or IN. The deletion or mutation of a gene and the insertion of a foreign gene can be carried out by known homologous recombination or site-directed mutagenesis.

[0049] The host cell for producing the BLV of the present invention is not particularly limited as long as it is capable of transcription and translation of an expression vector. Examples of the host cell include insect cells (e.g., silkworm cells), amphibian cells, reptilian cells, avian cells, fish cells, and mammalian cells (e.g., PK15 cells, HEK293 cells, HeLa cells, COS cells, BHK cells, CHL cells, and CHO cells). The host cell is preferably a mammalian cell.

[0050] The transformation method of the host cell may be selected depending on the type of host cell or the like, and the transformation can be performed, for example, by an electroporation method, a lithium acetate method, a calcium phosphate method, a lipofection method, a particle gun method, or the like.

[0051] The cell of the invention can be implemented according to the descriptions of the virus of the invention, in addition to the above description.

Method for Producing Virus

[0052] According to the present invention, there is also provided a method for producing a virus. The method for producing a virus according to the present invention comprises the step of culturing the cell of the present invention. The method for producing a virus according to the present invention may further comprise the step of recovering a virus produced by the cell of the present invention. The

recovered virus may be purified as necessary. The purification method may be appropriately selected from known methods depending on the virus to be produced. In the case of a virus that is accumulated within the host cell, the virus may be purified after disruption of the host cell. In the case of a virus that is released outside the host cell, the virus can be purified and recovered, for example, from a culture supernatant of the host cell. The method for producing a virus according to the present invention can be implemented according to the descriptions of the virus of the present invention, and the cell of the present invention and the method for producing the same, in addition to the above description.

BLV Vaccine

[0053] According to the present invention, there is also provided a BLV vaccine. The vaccine of the present invention is characterized by comprising the virus of the present invention. That is, the vaccine of the present invention can be used to prevent infection with a BLV or to treat (or ameliorate symptoms of) EBL, because administration of the vaccine into a subject induces an immune response against the BLV (for example, production of a specific antibody or proliferation of a cytotoxic T cell) and promotes production of a neutralizing antibody against the BLV and production of an interferon or the like in the subject.

[0054] The virus of the present invention, which is contained in the vaccine of the present invention, preferably has at least a gene encoding a structural protein, more preferably has at least any one of p24 of the gag gene and gp51 and gp30 of the env gene, and even more preferably has at least gp51, from the viewpoint of effectively inducing an immune response against the BLV in the subject to be administered with the vaccine. Without being bound by the following theory, it is believed that, in the infection with the BLV, when gp51 binds to a receptor on the surface of the cell membrane of the host (subject), the virus and the cell membrane of the host fuse by virtue of the membrane fusion activity of gp30, thereby allowing entry of the core of the BLV into the host cell (Bai L, et al., FASEB Journal, 2019, fj201901528R).

[0055] In the vaccine of the present invention, the content of the virus of the present invention can be appropriately set depending on the subject to be administered with the vaccine, the administration method, and the form of administration. For example, when the subject is cattle, the content can be 0.01 mg to 10 mg (preferably 0.5 mg to 5 mg) per cattle. The vaccine of the present invention may be administered once, twice, three or more times depending on the subject, and the number of administrations may be added whenever the antibody titer decreases in the subject.

[0056] The vaccine of the present invention can comprise a pharmaceutically acceptable carrier, a lubricant, a preservative, a stabilizer, a wetting agent, an emulsifier, a salt for adjusting an osmotic pressure, a buffer, a colorant, an antioxidant, a viscosity adjusting agent, an activator (a concept which also includes carbonate apatite, sodium hydroxide, alum, an incomplete/complete Freund's adjuvant and the like, as immunostimulants), nanoparticles or the like. Examples of the pharmaceutically acceptable carrier include water, various salt solutions, alcohols, vegetable oils, and mineral oils.

[0057] Examples of the form of the vaccine of the present invention include an injection, a liquid medicine, a suspen-

sion, an emulsion, a powder, a granule, and a capsule. The form of administration of the vaccine may be oral administration or parenteral administration (subcutaneous administration, transnasal administration, intraperitoneal administration, intracapsular administration, intramuscular administration, or an intravenous administration).

[0058] The vaccine of the present invention can be used in the prevention or treatment of bovine leukosis. According to another aspect of the present invention, there is provided a method for preventing or treating bovine leukosis, comprising the step of administering the vaccine of the present invention to a subject.

[0059] In the present invention, the “subject” is a non-human animal or a non-human mammal, and examples thereof include mammals such as cattle (*Bos taurus*), zebu (*Bos indicus*), water buffalo (*Bubalus bubalis*), sheep, goat, pig, mouse, rat, rabbit, cat, and monkey. Examples of the cattle include dairy breeds, meat breeds, dual-purpose breeds for dairy and meat, draft breeds, and dual-purpose breeds for draft and meat. Specific examples of the cattle include Japanese cattle such as Japanese Black cattle and Japanese Shorthorn cattle, Holstein cattle, Jersey cattle, and local breeds of various countries. From the viewpoint of preventing or treating bovine leukosis, the subject is preferably cattle, zebu and water buffalo, more preferably cattle and water buffalo, and even more preferably cattle.

[0060] The vaccine of the present invention can be implemented according to the descriptions of the virus of the present invention and the method for producing the same, and the cell of the present invention and the method for producing the same, in addition to the above description.

EXAMPLES

[0061] The present invention will be described in more detail by way of the following examples, but is not limited to these examples.

Example 1: Establishment of Non-Replicating BLV-Producing Cell Line

[0062] The present inventors previously constructed a CMVΔU3-pBLV-IF plasmid having a high virus production amount of BLV as disclosed in JP 2019-24351 A, but constructed a CMVΔU3-pBLV-416 plasmid using, as a based plasmid, a pBLV-416 plasmid cloned as a wild-type strain having a higher virus production amount than that of the CMVΔU3-pBLV-IF plasmid. The CMVΔU3-pBLV-416 plasmid was prepared in the same manner as in the preparation of the CMVΔU3-pBLV-IF plasmid described in JP 2019-24351 A. This CMVΔU3-pBLV-416 plasmid was introduced into PK15 cells to attempt to establish a cell line having a high BLV production amount. As a result of repeating cell subculture and seven runs of cloning, the inventors succeeded in creating a CMVΔU3-pBLV-416 plasmid-introduced mutated stable cell line (sometimes referred to as “PK15-BLVΔRT cell line” in the present specification) in which the virus produced had lost its replicating ability and integrating ability while retaining infectivity. As a result of analysis of the entire base sequence of the provirus integrated into this cell line, it was confirmed that 46 bp to 696 bp in the RT region of the pol gene were deleted (see FIG. 1).

Example 2: Study (1) Using pBLV-416ΔRT-Introduced Cell

[0063] In Example 2, a CMVΔU3-pBLV-416ΔRT plasmid introduced with a deletion of 46 bp to 696 bp in the RT region of the pol gene was prepared, and viral proteins produced from cells (pBLV-416ΔRT-introduced cells) introduced with this molecular clone were analyzed.

(1) Method

a. Plasmid

[0064] The CMVΔU3-pBLV-416 plasmid was prepared in the same manner as in the preparation of the CMVΔU3-pBLV-IF plasmid described in JP 2019-24351 A. CMVΔU3-pBLV416ART was prepared by site-directed mutagenesis using the CMVΔU3-pBLV-416 plasmid as a template (Bai L, et al., Retrovirology, 2015; 12(1):106, Inabe K, et al., J. Virol., 1999; 73: 1293-1301., Matsuura R, et al., 2019; 11(12):1140). The deletion of the RT region of the pol gene in pBLV-416ΔRT was confirmed by PCR.

b. Cell

[0065] African green monkey kidney-derived cells (COS-1) were used as the cells into which the plasmid was to be introduced. The COS-1 cells were maintained on a DMEM medium added with 10% fetal bovine serum (FBS) and 1×Penicillin-Streptomycin-Glutamine (PSG) at 37° C. in a CO₂ incubator.

c. Transfection

[0066] The COS-1 cells (5×10⁵ cells/60 mm-dish) were subjected to transfection using the reagents and plasmids indicated in Tables 3 to 5, and the cells and culture supernatants were recovered 48 hours after transfection. In the present specification, the cells subjected to transfection are sometimes referred to as “introduced cells.”

TABLE 3

Transfection of pBLV-416ART	
FuGENE HD	32 μL
pBLV-416ART	8 μg
Opti-MEM	400 μL

TABLE 4

Transfection of pBLV-416	
FuGENE HD	32 μL
pBLV-416	8 μg
Opti-MEM	400 μL

TABLE 5

Transfection of pBluescript II KS(-)	
FuGENE HD	32 μL
pBluescript II KS(-)	8 μg
Opti-MEM	400 μL

d. Western Blotting

[0067] The expression of the viral proteins was confirmed by Western blotting using a BLV-infected bovine serum or a non-BLV-infected bovine serum.

(2) Result

[0068] The results were as shown in FIG. 2. As a result of amplifying the pol gene region by PCR, it was confirmed that, in the pBLV-416 Δ RT plasmid, a fragment shorter by 650 bp than that in the pBLV-416 was amplified, and that a region corresponding to 46 bp to 696 bp in the RT region of the pol gene was deleted (FIG. 2A). As a result of the Western blotting, it was confirmed that, in the case of using the BLV-infected bovine serum, the expression levels of structural proteins such as Gag protein (p24) and Env proteins (gp51 and gp30) were not affected by the deletion of the pol gene in the pBLV-416 Δ RT-introduced cells as compared with the pBLV-416-introduced cells (FIG. 2B, left), whereas no viral protein was detected in the case of using the non-BLV-infected bovine serum (FIG. 2B, right).

Example 3: Study (2) Using
pBLV-416 Δ RT-Introduced Cell

[0069] In Example 3, the influences of the deletion of the pol gene on the intracellular localization and expression levels of the viral proteins were studied using pBLV-416 Δ RT-introduced cells.

(1) Method

a. Plasmid

[0070] A plasmid similar to that in Example 2 (1) a was used.

b. Cell

[0071] COS-1 cells were used similarly as in Example 2 (1) b.

c. Western Blotting

[0072] The viral proteins were confirmed by Western blotting using an anti-Gag antibody and an anti-Env antibody.

d. Immunofluorescent Antibody Method

[0073] The intracellular localization of the viral proteins was confirmed by an immunofluorescent antibody method using an anti-Gag antibody and an anti-Env antibody.

(2) Result

[0074] The results were as shown in FIGS. 3 and 4. In the pBLV-416 Δ RT-introduced cells, as compared with the pBLV-416-introduced cells, the influence of the deletion of the pol gene on the intracellular localization of the Gag protein (p24) and the Env protein (gp51) was not observed (FIG. 3), but increases in expression levels of both p24 and gp51 were confirmed (FIGS. 4A and 4B).

Example 4: Study (3) Using
pBLV-416 Δ RT-Introduced Cell

[0075] In Example 4, the syncytium formation ability and cell-to-cell infection ability of the virus produced from the pBLV-416 Δ RT-introduced cells were studied.

(1) Method

a. Syncytium Formation Ability

[0076] The pBLV-416 Δ RT plasmid or pBLV-416 plasmid was introduced into 293T cells together with a pEGFP-N1 plasmid (EGFP expression plasmid) to evaluate the syncytium formation ability. The negative control used was 293T cells into which a pBluescript II KS(−) plasmid, as an empty vector, was introduced together with an EGFP-N1 plasmid.

b. Evaluation of Cell-to-Cell Infection Ability

[0077] To evaluate the cell-to-cell infection ability, the pBLV-416 Δ RT plasmid or pBLV-416 plasmid was introduced into COS-1 cells, which were cocultured with CC81-GREMG cells (reporter cells for the BLV). The CC81-GREMG cells express EGFP in a BLV Tax-dependent manner. When the BLV-producing cells and the CC81-GREMG cells fuse due to the Env protein of the BLV to form syncytia, syncytia having expressed EGFP are observed. Therefore, the cell-to-cell infection ability can be evaluated.

(2) Result

[0078] The results were as shown in FIGS. 5 and 6. It was confirmed that both the pBLV-416 Δ RT-introduced cells and the pBLV-416-introduced cells formed syncytia (FIG. 5A). It was confirmed that the number and size of syncytia were larger in the pBLV-416 Δ RT-introduced cells than those in the pBLV-416-introduced cells (FIG. 5B). The results indicated that the deletion of the pol gene enhances the syncytium formation ability in the pBLV-416 Δ RT-introduced cells.

[0079] Regarding the cell-to-cell infection ability, it was also confirmed that the number and size of syncytia formed were larger in the pBLV-416 Δ RT-introduced cells than those in the pBLV-416-introduced cells (FIGS. 6A and B). The results suggested the possibility that the deletion of the pol gene may enhance the cell-to-cell infection.

Example 5: Study (4) Using
pBLV-416 Δ RT-Introduced Cell

[0080] In Example 5, the production amount of the virus produced from the pBLV-416 Δ RT-infected cells and the reverse transcriptase activity were analyzed.

(1) Method

a. Plasmid

[0081] A plasmid similar to that in Example 2 (1) a was used.

b. Cell

[0082] COS-1 cells were used similarly as in Example 2 (1) b.

c. Western Blotting

[0083] The viral proteins were detected by Western blotting using an anti-Gag antibody and an anti-Env antibody.

d. Virus Release Amount

[0084] To confirm the virus production amounts of the pBLV-416 Δ RT-introduced cells and the pBLV-416-introduced cells, the amount of p24 in the culture supernatant was measured by Capture ELISA. Specific procedures are as follows. A BLV-positive serum was diluted with a carbonate buffer solution adjusted to a pH of 9.0 and immobilized on a 96-well plate. Next, the culture supernatant of the pBLV-416 Δ RT-introduced cells or the pBLV-416-introduced cells and Tween 20 were added to the plate on which the serum was immobilized to capture p24. The amount of captured p24 was then measured by a colorimetric method using an anti-BLV p24 antibody (BLV3, VMRD) and an HRP-labeled anti-mouse IgG antibody.

e. Reverse Transcriptase Activity

[0085] The virus released into the culture supernatant was recovered by ultracentrifugation, and the reverse transcriptase activity in the viral particles was measured using a Reverse Transcriptase Assay, Colorimetric kit (Roche).

f. Viral Infectivity

[0086] To evaluate the viral infectivity, the virus-containing culture supernatant was added to CC81-GREMG cells to confirm the formation of syncytia. Using 2 µg of the virus, p24 and gp51 in the virus were quantified by Western blotting.

(2) Result

[0087] The results were as shown in FIG. 7. There was no difference in virus release amount due to the deletion of the pol gene between the pBLV-416ΔRT-introduced cells and the pBLV-416-introduced cells (FIG. 7A). Furthermore, the virus released from the pBLV-416ΔRT-introduced cells did not exhibit reverse transcriptase activity (FIG. 7B), nor did it form syncytia (FIG. 7C). Furthermore, the Gag protein (p24) and the Env protein (gp51) were detected in the pBLV-416ΔRT-introduced cells, indicating that the deletion of the pol gene does not affect viral maturation (FIG. 7D).

Example 6: Study Using pBLV-IFΔRT-Introduced Cell

[0088] In Example 6, a CMVΔU3-pBLV-IFΔRT plasmid introduced with a deletion of 46 bp to 696 bp in the RT region of the pol gene was prepared, and viral protein produced from cells (pBLV-IFΔRT-introduced cells) introduced with this molecular clone were analyzed.

(1) Method

a. Plasmid

[0089] A CMVΔU3-pBLV-IF2 plasmid was prepared according to the description of JP 2019-24351 A. Similarly as in Example 2 (1) a, a CMVΔU3-pBLV-IF2ΔRT plasmid was prepared by site-directed mutagenesis using the CMVΔU3-pBLV-IF plasmid as a template (Tajima S, et al., J Virol. 2000; 74 (23):10939-10949) (FIG. 8).

b. Cell

[0090] COS-1 cells and 293T cells were used as the cells into which the plasmid was to be introduced.

c. Western Blotting

[0091] The viral proteins were confirmed by Western blotting using an anti-Gag antibody and an anti-Env antibody.

d. Syncytium Formation Ability

[0092] The same procedures as in Example 4 (1) a were conducted.

(2) Result

[0093] The results were as shown in FIG. 9. It was confirmed that the expression levels of the Gag protein (p24) and the Env protein (gp51) increased in the pBLV-IF2ΔRT-introduced cells (293T cells) as compared with the pBLV-IF2-introduced cells (293T cells) (FIG. 9A). Also, it was confirmed that the number and size of syncytia increased in the pBLV-IF2ΔRT-introduced cells as compared with the pBLV-IF2-introduced cells (FIG. 9B).

[0094] In the pBLV-IF2ΔRT-introduced cells (COS-1 cells), an equivalent amount of Gag protein (p24) to that in the pBLV-IF2-introduced cells (COS-1 cells) was detected by Capture ELISA, and no influence of the deletion of the pol gene was observed (FIG. 9C).

Example 7: Study on Bovine Leukosis Vaccine

[0095] In Example 7, the virus produced from the pBLV-416ΔRT-introduced cells was inoculated into mice to study the effect of the virus as a vaccine.

(1) Method

a. Plasmid

[0096] A plasmid to similar that in Example 1 (1) a was used.

b. Cell

[0097] COS-1 cells were used as the cells into which the plasmid was to be introduced.

c. Virus

[0098] The culture supernatant of the pBLV-416ΔRT-introduced cells or the pBLV-416-introduced cells was centrifuged at 3,000 rpm for 10 minutes to remove cell debris, and then ultracentrifuged at 141,118×g and 4° C. for 2 hours to remove the culture supernatant. Then, the precipitate was resuspended in PBS to purify the virus.

d. Inoculation Into Mouse

[0099] According to the schedule shown in FIG. 10A, 100 µg of the virus was inoculated into mice in the 0th and 2nd weeks, and PBS was inoculated as the negative control (n=6 per group). Blood sampling was conducted from 0 to 6 weeks after inoculation. Following euthanasia 6 weeks after inoculation, the spleen was extracted. To confirm viral proliferation within the mouse bodies, the BLV genes in the blood cells 3 weeks after inoculation were detected using a Nested PCR method. In addition, spleen cells were co-cultured with CC81-GREMG cells to evaluate the syncytium formation ability. In addition, anti-p24 antibodies from 0 to 6 weeks after inoculation were quantified by ELISA.

(2) Result

[0100] The results were as shown in FIG. 10. In the mice inoculated with the virus produced by the pBLV-416ΔRT-introduced cells, no BLV gene was detected (FIG. 10B), and no syncytium was observed even when the spleen cells extracted from the mice were co-cultured with CC81-GREMG (FIG. 10C). On the other hand, it was confirmed that the anti-p24 antibody increased over time in the mice inoculated with the virus produced by the pBLV-416ΔRT-introduced cells (FIG. 10D). These results suggested the possibility that the virus produced from the pBLV-416ΔRT-introduced cells may be useful as a non-replicating BLV vaccine.

Example 8: Study (1) Using Non-Replicating BLV-Producing Cell Line

[0101] In Example 8, the entire base sequence of the provirus integrated into the PK15-BLVΔRT cell line successfully created in Example 1 was analyzed again. As a result, it was confirmed that 43 bp to 696 bp in the RT region of the pol gene were deleted (see FIG. 11). The results showed a difference of three bases in number of bases deleted in the RT region of the pol gene from the analysis results in Example 1. This difference is considered to be due to addition of further mutations during subculture after establishment of the PK15-BLVΔRT cell line. In addition, as compared with the base sequence of the wild-type BLV, a mutation of two bases and a deletion of one base were confirmed after 696 bp in the pol gene, but these mutations did not affect the protein expression.

Example 9: Study (2) Using Non-Replicating
BLV-Producing Cell Line

[0102] In Example 9, the expression and localization of the viral proteins in cells, and the amount of virus released were studied using the PK15-BLVΔRT cell line.

(1) Method

a. Western Blotting

[0103] The viral proteins were confirmed by Western blotting using an anti-Gag antibody and an anti-Env antibody.

b. Immunofluorescent Antibody Method

[0104] The intracellular localization of the viral proteins was confirmed by the immunofluorescent antibody method using an anti-Gag antibody and an anti-Env antibody.

c. Virus Release Amount

[0105] The virus release amount of the PK15-BLVΔRT cell line was measured by the Capture ELISA method. Specific procedures are as follows. The culture supernatant of the cells was centrifuged at 141,118×g for 2 hours to remove the supernatant. The precipitate was then resuspended in an amount of PBS one-hundredth the amount of the culture supernatant to concentrate the virus, and the

concentration of p24 was measured. The concentration of viral p24 was calculated from a standard curve created using, as a standard, His-p24 antigen expressed using *E. coli* and purified (FIG. 14B).

(2) Result

[0106] The results were as shown in FIGS. 12 to 14. The results in FIG. 12 confirmed that the PK15-BLVΔRT cell line expressed the viral structural proteins p24 and gp51. Further, as compared with the positive control, persistently BLV-infected cell FLK-BLV, no changes in molecular weights of p24 and gp51 due to the deletion of the pol gene were observed. From the results in FIG. 13, spotted accumulation was observed within the cells, similarly as in the positive control, persistently BLV-infected cell FLK-BLV. The results confirmed that the deletion of the pol gene does not affect the intracellular localization of p24. From the results in FIG. 14A, the concentration of p24 was 25.1±1.2 μg, confirming that a sufficient amount of virus was released from the PK15-BLVΔRT cell line. These results suggested the possibility that the virus produced from the PL15-BLVΔRT cell line may be useful as a non-replicating BLV vaccine.

SEQUENCE LISTING

Sequence total quantity: 2
SEQ ID NO: 1 moltype = DNA length = 8720
FEATURE Location/Qualifiers
source 1..8720
mol_type = other DNA
organism = Bovine leukemia virus

SEQUENCE: 1

tgtatgaaag	atcatgccga	cctaggcgcc	gccaccgccc	cgtaaaccag	acagagacgt	60
cagctgccag	aaaagctggt	gacggcagct	ggtggctaga	atccccgtac	ctccccaact	120
tcccctttcc	cgaaaaatcc	acaccctgag	ctgctgacct	cacctgctga	taaacataata	180
aaatgccggc	cctgtcgagt	tagcggcacc	agaagcgctc	ttctcctgag	accctcgtgc	240
tcagctctcg	gtcctgagct	ctcttgctcc	cgagaccttc	tggtcggcta	tcgggcagcg	300
gtcaggtaag	gcaaacccag	gtttggaggg	tggttctcgg	ctgagaccac	cgcgagctct	360
atctccggtc	ctctgaccgt	ctccacgtgg	actctctcct	ttgcctctcg	accccgcgct	420
ccaaggcgct	ctggcttgca	cccgcgcttg	tttctgtctc	tactttctgt	ttctcggcgg	480
ccgcgctctc	tccttcggcg	cccactagcg	gccaggagag	accggcaaac	aattgggggc	540
tcgtccggga	ttgatcaccc	cggaacccta	acaactctct	ggacccaccc	cctcggcgcc	600
attttgggtc	ttctcttcaa	attatatcat	gggaaattcc	ccctcctata	acccccccgc	660
tggtatctcc	ccctcagact	ggctcaacct	tctgcaaaag	gcgcaaaagg	tcaatccgcg	720
accctctcct	agcgatttta	ccgattttaa	gaattacatc	cattggtttc	ataagaccga	780
gaaaaaacca	tggagtttca	cttctgggtg	ccccacctca	tgctccaccg	ggagggttcg	840
ccgggttccc	cttgtcttgg	ccaccctaaa	cgaaagtctc	tcaaacgaag	ggggcgcccc	900
gggtgcatcg	gccccagaag	aacaaccccc	cccttatgac	ccccccgcg	ttttgccaat	960
cataatctga	gggaatcgca	accgccatcg	tgcttgggca	ctccgagaat	tacaagatat	1020
caaaaaggaa	attgaaaata	aggcaccggg	ttcgcaagta	tggtatacaa	cactacgact	1080
tgcaatctcg	caggccgacc	ctactccggc	tgacctagaa	caactttgcc	aatatattgc	1140
ttccccggtc	gaccaaaccg	cccatatgac	cagcctaacg	gcagcaattg	ccgccgctga	1200
agcggccaac	accctccagg	gttttaaccc	ccaaaacggg	accctaacc	aacaatcagc	1260
tcagcccaac	gccggggatc	ttagaagtca	atatcaaaac	ctctggcttc	aggcctggaa	1320
aaatctcctc	actcgtcctt	tagtacaacc	ttgggtccacc	atcgctccaa	gccccgcgca	1380
aagctatgta	gagtttgcta	accggttaca	aatttcatta	gctgacaacc	ttcccagcgg	1440
agtccttaag	gaaccattta	ttgactcctc	tagttatgca	aatgctaaca	aagagtgtca	1500
gcaaatcttg	caagggcgag	gcctagtggc	cgccccgggtg	gggcaaaaac	tgacgggttg	1560
cgcacattgg	gcccccaaga	tgaacagacc	tgcaattctc	gtccacaccc	caggggcccaa	1620
gatgccgggg	cctcggaac	cggcccccaa	aaggcctccc	ccaggaccat	gctatcgatg	1680
cctcaaaaga	ggccatttgg	cccgggattg	tcttaccaa	gccaccggcc	cccctccggg	1740
accttgcccc	atatgtaaag	atccttccca	ttggaaacga	gactgtccaa	ccctcaaatc	1800
aaaaaactaa	tagagggggg	acttagcgcc	ccccaaacca	taacacccat	aacagattct	1860
cttagtgagg	ccgaattaga	atgcttactt	tctattcctc	tggtctcgag	cgtccctccc	1920
gtggctgtat	acctgtctgg	cccttggtcg	cagccctctc	agaatcaagc	cctcatgctt	1980
gtggacaccc	gggctgaaaa	tacggttctc	ccacaaaatt	ggctgggttc	ggattaccca	2040
cggatccccg	cgcagtgct	cggagcaggg	ggagtctccc	ggaacagata	caattggcta	2100
caaggccctc	tgaccctggc	tctaaaaacca	gagggtccct	ttatcccat	cccaaaaatt	2160
ttagttgaca	ctttcgacaa	atggcaaat	ttaggacggg	acgtcctctc	ccgcctacag	2220

-continued

gcctctatct	ccatacctga	ggaagtacgc	ccccctgtgg	taggcgtctt	ggatgcccc	2280
ccgagccaca	ttggattaga	acatctgccc	cctccacctg	aagtgcctca	attcccttta	2340
aactagaacg	cctccaggcc	cttcaagacc	tggtccatcg	ctctctggag	gcaggttata	2400
ttccccctg	ggaaggccca	ggcaataatc	cagtcttccc	ggtacggaaa	ccaaatggcg	2460
ectggaggtt	tgtgcattgac	ctacagagcta	caaatgctct	tacaaagccc	attccggcac	2520
ttccccccgg	accgccagac	cttaccgcta	ttccctacaca	ccttccacat	atcatttgcc	2580
tagatctcaa	agacgccttc	ttccagattc	cagtcgaaga	ccgcttccgc	tcctactttg	2640
cttttacct	ccctaccccc	gggggactcc	aacctcatag	acgctttgcc	tggcggtgcc	2700
taacctcaagg	cttcattaac	agcccagctc	ttttcgaacg	agcactacag	gaacctcttc	2760
gccaagtctc	cgccgccttt	ttccagttct	ttctgggtgc	ctatatggac	gatataccta	2820
tcgcttccgc	tacagaagaa	cagcgggtcac	aatgttatca	agccctggct	gcccgcctcc	2880
gggacctag	gtttcagggtg	gcctccgaaa	agactcgcca	gacgccttcg	cccgctccct	2940
ttttgggaca	aatgggtccat	gagcagattg	tcacctaacc	gtccctacct	accttgccga	3000
ttctatcccc	aatctctctt	caccaattac	aggcggtctt	aggagacctc	caatgggtct	3060
ctagggggac	accactacc	cgccggcccc	tgcaacttct	ctactcttcc	cttaaaaggca	3120
tcgatgcccc	tagggccatc	atccagcttt	ccccggaaca	gctgcaaggc	attgcagagc	3180
ttcgacaagc	cctgtcccat	aacgcaagat	ctagatataa	cgagcaagaa	ccccgtctag	3240
cctacgtaca	cctaaccogg	gcggggtcca	ccctgggtact	cttccaaaag	ggcgctcaat	3300
ttccctgggc	ctacttccag	acccccttga	ctgacaacca	agcctcacct	tggggcctcc	3360
ttctcctgct	gggatgccaa	tacctgcaga	ctcaggcctt	aagctcgtat	gccaaagccc	3420
tacttaataa	ttatcacaat	cttccataaa	cttctctaga	caattggatt	caatcatctg	3480
aggacctcg	agtcaggag	ttgctgcaat	tgtggcccca	gatttctctc	cagggaaatac	3540
agcccccg	cccttggaag	accttaatac	ccagggcaga	ggtttttttg	acgccccagt	3600
ttccccctg	tcggattcct	gcgccctttt	gcctctttag	tgacggggct	aaaggacgag	3660
gagcatattg	cttgtggaag	gaccaccttt	tagactttca	ggcggttccg	gctccagaat	3720
ccgctcaaaa	gggagaacta	cgagggtctc	tggcgggctt	agcagccgcc	ccgcctgaac	3780
ctgtaaatat	atgggttagat	ttcaaatacc	tgtactcttt	gctcagaacc	ctagtctctg	3840
gagcttggtc	tcaacctgac	cccgtaacct	cctacgcctc	cctatataaa	agcctctccc	3900
gacatccagc	aatctttggt	ggccatgtcc	ggagccactc	ttcagcatcc	cacctatttg	3960
cttccctgaa	caattatgta	gatcaactgc	ttcccttaga	aaactccagag	caatggcata	4020
agctcaccca	ctgcaactct	cgggccttgt	ctcgatggcc	gaaccacagt	atctctgcct	4080
gggacccccg	ttcccccgct	acgctgtgtg	aaacctgtca	aaagcttaat	ccaactggag	4140
gaggaagagt	gcgaactatt	cagagagggt	gggccccgaa	ccatatttgg	caggccgata	4200
taaccattta	taaatacaaa	cagttcacct	acgctctgca	tgtgtttgta	gatacttact	4260
ctggagctac	tcctgcctcg	gcgaagcgtg	ggctcaccac	tcaaatgacc	attgaggggc	4320
ttcttgaggc	catagtgcat	ctgggtcgct	caaaaaagct	aaacactgac	caagggtgca	4380
actacacctc	caaaaacctt	gtcaggtttt	gccagcagtt	cggagtttcc	ctttctcatc	4440
atgttcccta	caacccccca	agtttcgggg	tagtagaacg	gacaaatgga	ctgctcaaac	4500
ttcttctatc	taaatatcac	ctagacgaac	cccaccttcc	catgactcag	gcccctttct	4560
gagccctctg	gactccaact	cagattaacc	tctaaccaat	tctaaagacc	agatggggagc	4620
tacaccattc	acccccactt	gctgtcattt	cagagggcgg	agaaacaccc	aagggtctctg	4680
ataaactctt	ttgttacaa	ctccccgggc	aaaacaatcg	tcggtggcta	ggaccactcc	4740
cgccctagtg	ggaagcctcg	ggaggcgctc	tcctgggtac	tgaccccccc	gtgtgggttc	4800
cctggcggtt	gctgaaagcc	ttcaaatgcc	taaaagaacga	cggtcccgaa	gacgcccaca	4860
accgatcatc	agatgggtga	gtctcactct	cactctctct	gctctctctc	ggcccatcca	4920
gacttgagga	gtctccctgt	cctagggaag	ccaacaatgg	atgacagcat	ataaccaaga	4980
ggcaaaaatt	ttcatctcca	ttgaccaaat	actagaggct	cataatcagt	cacctttctg	5040
tgccaaagtct	cccagataca	ccttggaact	tgtaaatggc	tatcctaaga	tctactggcc	5100
ccccccacaa	ggggcgccgc	ggtttggagc	caggggccatg	gtcacatatg	attgcgagcc	5160
ccgatgcctc	tatgtggggg	cagatcgctt	cgactgcccc	cactggggaca	atgcctccca	5220
ggccgatcaa	ggatcccttt	atgtcaatca	tcagatttta	ttctctgcat	tcaaaccaatg	5280
tcatggaatt	ttcaacttta	cctggggagat	atgggggatat	gatccccctga	tcaccttttc	5340
tttacataag	atccctgatc	ccctcaacc	cgactttccc	cagttgaaca	gtgactgggt	5400
ttccctctgt	agatcatggg	ccctgctttt	aaatcaaaac	gcacggggct	tccagactgc	5460
tgctatatgt	tgggaacctt	ccctccctcg	ggctccccga	atattagtat	ataacaaaac	5520
catctccagc	tttggaccgc	gcctcgccct	cccggaacgc	caaatcttct	gggtcaaacac	5580
gtcctcggtt	aacaccaccc	aaggatggca	ccaccttccc	cagaggttgt	tgttcaatgt	5640
ttctcaaggc	aacgccttgt	tattacctcc	tatctccctg	gttaactctc	ctacggcttc	5700
ctccgccccct	cctaccgggg	tcagacgtag	ttccgtcgca	gcctgacct	taggcctagc	5760
cctgtcagtg	gggctcactg	gaattaatgt	ggccgtgtct	gccttagacc	atcagagact	5820
caactccctg	atccacgttc	tgaggcaaga	tcagcaacgc	ttgatcacag	caattaaaca	5880
gaccactat	aatgtgttta	atgtggcttc	tgtggttgcc	cagaaccgac	gggggcttga	5940
ttgggttatc	atccggctgg	gttttcaaag	cctatgtccc	acaattaatg	agccttgctg	6000
tttctgcgc	attcaaaaat	actccattat	ccgctcggt	gatctccagc	ctctctcgca	6060
aagagtctct	acagactggc	agtggccctg	gaattgggat	ctggggctca	ctgcctgggt	6120
gcgagaaacc	attcattctg	ttctaagcct	gttccctatta	gccccttttt	tgctcttctc	6180
ggcccccctg	ctgataaaat	gcttgacctc	tcgcctttta	aagctctctc	ggcaggctcc	6240
ccacttccct	gaaatctctc	taacccctaa	accgattctc	gattatcagg	ccttgctacc	6300
atctgcacca	gagatctact	ctcacctctc	ccccgccaaa	cccgattaca	tcaacctccg	6360
accctgtcct	tgataccccc	gcgtttcacg	cacccccagg	ctgtgggtgt	gcactgggtt	6420
agtggaaatg	tcagttgtac	atcaaaagcc	ttctcttgct	gccggcaccc	agttcgaaaca	6480
cagccctacc	ctgagctctc	ctgagtgcac	gactgagtg	agcgcagaga	gattgtcgct	6540
ttctgcgtgc	gctcagttat	tttttatagc	cgattggggg	tcgcgccctt	ccgttgccctg	6600
tgacacagat	aagacctctc	tcacttctgc	ttcaccatcc	ccctgccagc	gttggtctag	6660
tggaagaac	taacgctgac	gggggcgatt	ttctgcagct	gtgctagcgg	gaggctctgg	6720
tgctggggat	aaggtgtggc	ccttagcacc	acagctctctg	cgcttttttg	gttcgaatct	6780

-continued

```

tccccacgca gcttccgctt tttacgcct gttgcacacc ctttctagag atacctgaaa 6840
atctcagctc gcacctgag gaagggtgtg gctcagaggt taaaatagct cgggccgcaa 6900
cctcccttto tttttattcc accctcgcaa ggccccgggt tctgagcccc ctaacggagg 6960
ttcaaaatth cctctactag gggatgctcg ggtccaagtg tgcacaatat ctcttccaaa 7020
aggctctgat gaacgtcttc ccatgtaaca agccccagca gagacatttc agccacatcc 7080
agcagcattt gggcgcctt ttttaacagt gcccataaag tcccttcctg ttcacacaag 7140
gctgcctctg cttctctat ttcacctcg gcaccgactc ccccgccgag cccttcgagc 7200
tcttcgggat ccattacctg ataacgacaa aattatttct tgtcttttaa gcaagtgttg 7260
ttgggttggg gccccactct ctacatgcct gcccgccctt ggttttctgc aatgatgtca 7320
ccatcgatgc ctggtgcccc cctgcgggac cccatgagcg actccaattc gaaaggatcg 7380
acaccacgct cacttgagag acccacgcta tcaactggac cgcgatgga cgaccttgcg 7440
gcctcaatgg aacgttgttc cctcgactgc atgtctccga gaccgcccc caagggcccc 7500
gacgactctg gatcaactgc ccccttcctg ccgttcgcgc tcagcccgcc ccggtttcac 7560
tttccccctt cgagcgggtc ccccttcagc cctaccaatg ccaattgccc tcggcctcta 7620
gcgacgggtg cccattatc gggcagcgcc ttcttcctcg gaacaactta gtaacgcatc 7680
ctgtcctcgg aaaagtcctt atattaaatc aaatggccaa ttttctta ctccctcct 7740
tcgatccctt ccttgtggac cccctccggc tgtccgtctt tgccccagac accaggggag 7800
ccatacgcta tctctccacc cttttgacgc tatgcccagc tacttgtatt ctacccttag 7860
gcgagccctt ctctcctaat gtcccataat gccgttttcc cgggactct aatgaacccc 7920
ccctttcaga attcgagctg ccccttatcc aaacgcccgg cctgtcctgg tctgtccccg 7980
cgatcgacct attctaaccc gggtccctct ccccatgcca cgggttacac gtatggtcca 8040
gtctcaggc cttacagcgc ttcttcctat accctacgct aacctggtcc gaattgggtg 8100
ctagcagaaa aataagactt gattccccct taaaattaca actgctagaa aatgaatggc 8160
tctcccgctt tttttgaggg ggagtcattt gtatgaaaga tcatgcccag ctaggcgccg 8220
ccaccgcccc gtaaacccaga cagagacgtc agctgccaga aaagctgggt acggcagctg 8280
gtggctagaa tcccgtacc tccccaaact cccctttccc gaaaaatcca caccctgagc 8340
tgctgacctc acctgctgat aaactaataa aatgcgggcc ctgtcgagtt agcggcacca 8400
gaagcgttct tctcctgaga cctcgtgctc cagctctcgg tctgagctc tcttgcctcc 8460
gagaccttct ggtcggttat ccggcagcgg tcaggttaagg caaacccagg ttggagggtg 8520
ggttctcggc tgagaccacc gcgagctcta tctccggtcc tctgaccgct tccacgtgga 8580
ctctctcctt tgcctctgta cccgcgcgtc caaggcgctc tggcttgac ccgcgtttgt 8640
ttcctgtctt accttctgtt tctcgcggcc cgcgtctctt ccttcggcgc ccactagcgg 8700
ccaggagaga ccgcaaaaaa 8720

```

```

SEQ ID NO: 2      moltype = DNA length = 2537
FEATURE          Location/Qualifiers
source           1..2537
                 mol_type = other DNA
                 organism = Bovine leukemia virus

```

```

SEQUENCE: 2
aactagaacg cctccaggcc cttcaagacc tgggtccatcg ctctctggag gcaggttata 60
tctccccctg ggaaggccca ggcaataatc cagtcttccc ggtacggaaa ccaaatggcg 120
cctggagggt tgtgcatgac ctacgagcta caaatgctct tacaagccc attccggcac 180
tctccccctg accgccagac cttaccgcta tccctacaca ccttccacat atcatttgcc 240
tagatctcaa agacgccttc ttccagattc cagtccaaga ccgttccgc tctactttg 300
cttttacctc ccttaccccc gggggactcc aacctcatag acgctttgcc tggcgggtcc 360
tacctcaagg cttcattaac agcccagctc ttttcgaacg agcactacag gaacctcttc 420
gccaaagttc cgcgccttt ttccagttct tcttggtgtc ctatatggac gatatcctta 480
tcgcttcgcc tacagaagaa cagcgggtac aatgttatca agcctgggtt gcccgctccc 540
gggacctagg gtttcagggt gcatccgaaa agactcgcca gacgccttcg ccggtccctc 600
tcttgggaca aatgggtcat gagcagattg tcacctacca gtccctacct accttgacga 660
tctcatcccc aatttctctt caccattac aggcggtctt agggagacct caatgggtct 720
ctagggggac acccactacc cgcgggcccc tgcacttctt ctactcttc cttaaaggca 780
tcgatgaccc tagggccatc atccagcttt ccccggaaca gctgcaaggc attgcagagc 840
ttcgacaagc cctgtcccat aacgcaagat ctagatataa cgagcaagaa cccctgctag 900
cctacgtaca cctaaccctg gcgggggtcca ccttggtact cttccaaaag ggcgctcaat 960
ttcccttgcc ctacttccag accccttga ctgacaacca agcctcactt tggggcctcc 1020
ttctcctgct gggatgccaa tacctgcaga ctcaggcctt aagctcgtat gccaaagcca 1080
tacttaataa ttatcacaat cttcctaaaa cttctctaga caattggatt caatcatctg 1140
aggacctctg agtccaggag ttgctgcaat tgtggcccca gatttctct cagggaatac 1200
agcccccggt ccttgggaag accttaatca ccagggcaga ggtttttttg acgccccagt 1260
tctccccctg tccgattcct gcggcccttt gcctctttg tgacgggggt aaaggacgag 1320
gagcatattg cttgtggaag gaccaccttt tagactttca ggccgttccg gctccagaat 1380
ccgctcaaaa gggagaacta gcaggtctct tggcgggctt agcagccgac ccgctgaac 1440
ctgtaaatat atgggtagat tccaaatacc tgtactcttt gctcagaacc ctagtctggt 1500
gagcttggct tcaacctgac ccggtaccct cctacgcccct cctatataaa agcctcctcc 1560
gacatccagc aatcttgggt ggccatgtcc ggagccactc ttcagcatcc caccctattg 1620
ctccctgaa caattatgta gatcaactgc ttcccttaga aactccagag caatggcata 1680
agctcaccia ctgcaactct cgggccttgt ctcgatggcc gaacccacgt atctctgct 1740
gggacccccg ttcccccgct acgctgtgtg aaacctgtca aaagcttaat ccaactggag 1800
gaggaaagat tgcgaactat cagagagggt gggcccccga ccatatttgg caggccgata 1860
taaccattta taataacaaa cagttcacct acgctctgca tgtgtttgta gatacttact 1920
ctggagctac tcatgctctg gcgaagcgtg ggctcaccac tcaaatgacc attgagggcc 1980
ttcttgaggg catagtgcct cttgggtcgtc caaaaaagct aaacactgac caaggtgcaa 2040
actacacctc caaacctttt gtcaggtttt gccagcagtt cggagtttcc ctttctcatc 2100
atgttcccta caaccccaca agttcgggggt tagtagaacg gacaatgga ctgctcaaac 2160

```

-continued

tttttttato	taaatatcac	ctagacgaac	cccaccttcc	catgactcag	gccctttctc	2220
gagccctctg	gactcacaat	cagattaacc	tcctaccaat	tctaaagacc	agatgggagc	2280
tacaccatto	acccccactt	gctgtcattt	cagagggcgg	agaaacaccc	aagggtctctg	2340
ataaactctt	tttgtacaag	ctccccgggc	aaaacaatcg	tcggtggcta	ggaccactcc	2400
cggccctagt	cgaagcctcg	ggaggcgctc	tcctgggtac	tgaccccccc	gtgtgggttc	2460
cctggcggtt	gctgaaagcc	ttcaaatgcc	taaagaacga	cggccccgaa	gacgcccaca	2520
accgatcatc	agatggg					2537

1. A bovine leukemia virus (BLV) in which at least a part of the function of a pol gene is deficient.

2. The virus according to claim 1, wherein the function of the pol gene is the function of a reverse transcriptase and/or an integrase.

3. The virus according to claim 1, wherein at least a part of the pol gene is mutated.

4. The virus according to claim 3, wherein the mutation is a deletion of 1 to 1643 bases in a reverse transcriptase region of the pol gene and/or 1 to 894 bases in an integrase region of the pol gene.

5. The virus according to claim 1, which does not replicate in an infected subject.

6. A non-replicating bovine leukemia virus (BLV)-producing cell, comprising a gene of a BLV in which at least a part of the function of a pol gene is deficient.

7. A method for producing a non-replicating BLV, comprising the step of culturing the cell according to claim 6.

8. A method for preventing or treating enzootic bovine leukosis (EBL), comprising the step of inoculating a subject excluding a human with the virus according to claim 1.

9. A method for producing a non-replicating bovine leukemia virus (BLV)-producing cell, comprising the step of introducing, into a host cell, a gene of a BLV in which at least a part of the function of a pol gene is deficient.

* * * * *