

# US Patent & Trademark Office

## Patent Public Search | Text View

United States Patent	12390514
Kind Code	B2
Date of Patent	August 19, 2025
Inventor(s)	Maianti; Juan Pablo et al.

### Cancer vaccine

#### Abstract

Provided herein are systems, compositions, and methods for generating immunogenic peptides or epitopes from tumor associated antigens (e.g., in vivo or ex vivo). Polynucleotides (e.g., genes) encoding the tumor associated antigens may be edited at selected target sites by nucleobase editors comprising a catalytically-inactive Cas9 and a cytosine deaminase, leading to the expression of heteroclitic or cryptic peptides that are more immunogenic than the native peptide derived from the tumor associated antigens. The heteroclitic or cryptic peptide elicit strong tumor-specific immune response (e.g., T-cell response or B-cell response), which inhibits tumor growth and metastasis.

<b>Inventors:</b>	<b>Maianti; Juan Pablo (Revere, MA), Liu; David R. (Lexington, MA)</b>
<b>Applicant:</b>	<b>President and Fellows of Harvard College (Cambridge, MA)</b>
<b>Family ID:</b>	<b>1000008767269</b>
<b>Assignee:</b>	<b>President and Fellows of Harvard College (Cambridge, MA)</b>
<b>Appl. No.:</b>	<b>16/492534</b>
<b>Filed (or PCT Filed):</b>	<b>March 09, 2018</b>
<b>PCT No.:</b>	<b>PCT/US2018/021880</b>
<b>PCT Pub. No.:</b>	<b>WO2018/165631</b>
<b>PCT Pub. Date:</b>	<b>September 13, 2018</b>

#### Prior Publication Data

Document Identifier	Publication Date
US 20210196809 A1	Jul. 01, 2021

#### Related U.S. Application Data

us-provisional-application US 62469219 20170309

#### Publication Classification

**Int. Cl.:** A61K38/50 (20060101); A61K31/7088 (20060101); A61K38/46 (20060101); A61K39/00 (20060101); A61K45/06 (20060101)

#### U.S. Cl.:

**CPC** A61K38/50 (20130101); A61K31/7088 (20130101); A61K38/465 (20130101); A61K39/0011 (20130101); A61K39/001106 (20180801); A61K39/00113 (20180801); A61K39/00115 (20180801); A61K39/001153 (20180801); A61K39/001156 (20180801); A61K39/001157 (20180801); A61K39/00117 (20180801); A61K39/001182 (20180801); A61K39/001188 (20180801); A61K39/001191 (20180801); A61K39/001192

## Field of Classification Search

USPC: None

## References Cited

### U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
4182449	12/1979	Kozlow	N/A	N/A
4186183	12/1979	Steck et al.	N/A	N/A
4217344	12/1979	Vanlerberghe et al.	N/A	N/A
4235871	12/1979	Papahadjopoulos et al.	N/A	N/A
4261975	12/1980	Fullerton et al.	N/A	N/A
4485054	12/1983	Mezei et al.	N/A	N/A
4501728	12/1984	Geho et al.	N/A	N/A
4663290	12/1986	Weis et al.	N/A	N/A
4737323	12/1987	Martin et al.	N/A	N/A
4774085	12/1987	Fidler	N/A	N/A
4797368	12/1988	Carter et al.	N/A	N/A
4837028	12/1988	Allen	N/A	N/A
4873316	12/1988	Meade et al.	N/A	N/A
4880635	12/1988	Janoff et al.	N/A	N/A
4889818	12/1988	Gelfand et al.	N/A	N/A
4897355	12/1989	Eppstein et al.	N/A	N/A
4906477	12/1989	Kurono et al.	N/A	N/A
4911928	12/1989	Wallach	N/A	N/A
4917951	12/1989	Wallach	N/A	N/A
4920016	12/1989	Allen et al.	N/A	N/A
4921757	12/1989	Wheatley et al.	N/A	N/A
4946787	12/1989	Eppstein et al.	N/A	N/A
4965185	12/1989	Grischenko et al.	N/A	N/A
5017492	12/1990	Kotewicz et al.	N/A	N/A
5047342	12/1990	Chatterjee	N/A	N/A
5049386	12/1990	Eppstein et al.	N/A	N/A
5079352	12/1991	Gelfand et al.	N/A	N/A
5139941	12/1991	Muzyczka et al.	N/A	N/A
5173414	12/1991	Lebkowski et al.	N/A	N/A
5223409	12/1992	Ladner et al.	N/A	N/A
5244797	12/1992	Kotewicz et al.	N/A	N/A
5270179	12/1992	Chatterjee	N/A	N/A
5374553	12/1993	Gelfand et al.	N/A	N/A
5405776	12/1994	Kotewicz et al.	N/A	N/A
5436149	12/1994	Barnes	N/A	N/A
5449639	12/1994	Wei et al.	N/A	N/A
5496714	12/1995	Comb et al.	N/A	N/A
5512462	12/1995	Cheng	N/A	N/A
5580737	12/1995	Polisky et al.	N/A	N/A
5614365	12/1996	Tabor et al.	N/A	N/A
5652094	12/1996	Usman et al.	N/A	N/A
5658727	12/1996	Barbas et al.	N/A	N/A
5668005	12/1996	Kotewicz et al.	N/A	N/A
5677152	12/1996	Birch et al.	N/A	N/A
5767099	12/1997	Harris et al.	N/A	N/A
5780053	12/1997	Ashley et al.	N/A	N/A
5830430	12/1997	Unger et al.	N/A	N/A
5834247	12/1997	Comb et al.	N/A	N/A
5835699	12/1997	Kimura	N/A	N/A
5844075	12/1997	Kawakami	530/328	A61P 35/00

5849548	12/1997	Haseloff et al.	N/A	N/A
5851548	12/1997	Dattagupta et al.	N/A	N/A
5855910	12/1998	Ashley et al.	N/A	N/A
5856463	12/1998	Blankenborg et al.	N/A	N/A
5962313	12/1998	Podsakoff et al.	N/A	N/A
5981182	12/1998	Jacobs, Jr. et al.	N/A	N/A
6015794	12/1999	Haseloff et al.	N/A	N/A
6057153	12/1999	George et al.	N/A	N/A
6063608	12/1999	Kotewicz et al.	N/A	N/A
6077705	12/1999	Duan et al.	N/A	N/A
6099857	12/1999	Gross	N/A	N/A
6156509	12/1999	Schellenberger	N/A	N/A
6183998	12/2000	Ivanov et al.	N/A	N/A
6355415	12/2001	Wagner et al.	N/A	N/A
6416997	12/2001	Mir-Shekari et al.	N/A	N/A
6429298	12/2001	Ellington et al.	N/A	N/A
6453242	12/2001	Eisenberg et al.	N/A	N/A
6479264	12/2001	Louwrier	N/A	N/A
6503717	12/2002	Case et al.	N/A	N/A
6534261	12/2002	Cox, III et al.	N/A	N/A
6558671	12/2002	Slingluff et al.	N/A	N/A
6589768	12/2002	Kotewicz et al.	N/A	N/A
6599692	12/2002	Case et al.	N/A	N/A
6607882	12/2002	Cox, III et al.	N/A	N/A
6610522	12/2002	Kotewicz et al.	N/A	N/A
6689558	12/2003	Case	N/A	N/A
6716973	12/2003	Baskerville et al.	N/A	N/A
6824978	12/2003	Cox, III et al.	N/A	N/A
6933113	12/2004	Case et al.	N/A	N/A
6979539	12/2004	Cox, III et al.	N/A	N/A
7013219	12/2005	Case et al.	N/A	N/A
7045337	12/2005	Schultz et al.	N/A	N/A
7067650	12/2005	Tanaka	N/A	N/A
7070928	12/2005	Liu et al.	N/A	N/A
7078208	12/2005	Smith et al.	N/A	N/A
7083970	12/2005	Schultz et al.	N/A	N/A
7163824	12/2006	Cox, III et al.	N/A	N/A
7192739	12/2006	Liu et al.	N/A	N/A
7223545	12/2006	Liu et al.	N/A	N/A
7329807	12/2007	Vadrucci et al.	N/A	N/A
7354761	12/2007	Schultz et al.	N/A	N/A
7368275	12/2007	Schultz et al.	N/A	N/A
7419669	12/2007	Kosmatopoulos et al.	N/A	N/A
7442160	12/2007	Liu et al.	N/A	N/A
7476500	12/2008	Liu et al.	N/A	N/A
7476734	12/2008	Liu	N/A	N/A
7479573	12/2008	Chu et al.	N/A	N/A
7488718	12/2008	Scheinberg et al.	N/A	N/A
7491494	12/2008	Liu et al.	N/A	N/A
7510706	12/2008	Yonemitsu et al.	N/A	N/A
7541450	12/2008	Liu et al.	N/A	N/A
7556940	12/2008	Galarza et al.	N/A	N/A
7557068	12/2008	Liu et al.	N/A	N/A
7595179	12/2008	Chen et al.	N/A	N/A
7638300	12/2008	Schultz et al.	N/A	N/A
7670807	12/2009	Lampson et al.	N/A	N/A
7678554	12/2009	Liu et al.	N/A	N/A
7713721	12/2009	Schultz et al.	N/A	N/A
7771935	12/2009	Liu et al.	N/A	N/A
7794931	12/2009	Breaker et al.	N/A	N/A
7807408	12/2009	Liu et al.	N/A	N/A

7851658	12/2009	Liu et al.	N/A	N/A
7915025	12/2010	Schultz et al.	N/A	N/A
7919277	12/2010	Russell et al.	N/A	N/A
7993672	12/2010	Huang et al.	N/A	N/A
7998904	12/2010	Liu et al.	N/A	N/A
7999071	12/2010	Schlom et al.	N/A	N/A
8012739	12/2010	Schultz et al.	N/A	N/A
8017323	12/2010	Liu et al.	N/A	N/A
8017755	12/2010	Liu et al.	N/A	N/A
8030074	12/2010	Schultz et al.	N/A	N/A
8067556	12/2010	Hogrefe et al.	N/A	N/A
8114648	12/2011	Schultz et al.	N/A	N/A
8173364	12/2011	Schultz et al.	N/A	N/A
8173392	12/2011	Schultz et al.	N/A	N/A
8183012	12/2011	Schultz et al.	N/A	N/A
8183178	12/2011	Liu et al.	N/A	N/A
8206914	12/2011	Liu et al.	N/A	N/A
8354380	12/2012	Liu et al.	N/A	N/A
8361725	12/2012	Russell et al.	N/A	N/A
8394604	12/2012	Liu et al.	N/A	N/A
8420104	12/2012	Charneau et al.	N/A	N/A
8440431	12/2012	Voytas et al.	N/A	N/A
8440432	12/2012	Voytas et al.	N/A	N/A
8450471	12/2012	Voytas et al.	N/A	N/A
8492082	12/2012	De Franciscis et al.	N/A	N/A
8546553	12/2012	Terns et al.	N/A	N/A
8569256	12/2012	Heyes et al.	N/A	N/A
8586363	12/2012	Voytas et al.	N/A	N/A
8673612	12/2013	Klatzmann et al.	N/A	N/A
8680069	12/2013	de Fougerolles et al.	N/A	N/A
8691729	12/2013	Liu et al.	N/A	N/A
8691750	12/2013	Constien et al.	N/A	N/A
8697359	12/2013	Zhang	N/A	N/A
8697439	12/2013	Mangeot et al.	N/A	N/A
8697853	12/2013	Voytas et al.	N/A	N/A
8709466	12/2013	Coady et al.	N/A	N/A
8728526	12/2013	Heller	N/A	N/A
8729038	12/2013	Gruber et al.	N/A	N/A
8741279	12/2013	Kasahara et al.	N/A	N/A
8748667	12/2013	Budzik et al.	N/A	N/A
8758810	12/2013	Okada et al.	N/A	N/A
8759103	12/2013	Kim et al.	N/A	N/A
8759104	12/2013	Unciti-Broceta et al.	N/A	N/A
8771728	12/2013	Huang et al.	N/A	N/A
8790664	12/2013	Pitard et al.	N/A	N/A
8795965	12/2013	Zhang	N/A	N/A
8822663	12/2013	Schrum et al.	N/A	N/A
8835148	12/2013	Janulaitis et al.	N/A	N/A
8846578	12/2013	McCray et al.	N/A	N/A
8871445	12/2013	Cong et al.	N/A	N/A
8889418	12/2013	Zhang et al.	N/A	N/A
8900814	12/2013	Yasukawa et al.	N/A	N/A
8945839	12/2014	Zhang	N/A	N/A
8975232	12/2014	Liu et al.	N/A	N/A
8993233	12/2014	Zhang et al.	N/A	N/A
8999641	12/2014	Zhang et al.	N/A	N/A
9023594	12/2014	Liu et al.	N/A	N/A
9023649	12/2014	Mali et al.	N/A	N/A
9034650	12/2014	Padidam	N/A	N/A
9068179	12/2014	Liu et al.	N/A	N/A
9150626	12/2014	Liu et al.	N/A	N/A

9163271	12/2014	Schultz et al.	N/A	N/A
9163284	12/2014	Liu et al.	N/A	N/A
9181535	12/2014	Liu et al.	N/A	N/A
9200045	12/2014	Liu et al.	N/A	N/A
9221886	12/2014	Liu et al.	N/A	N/A
9228207	12/2015	Liu et al.	N/A	N/A
9234213	12/2015	Wu	N/A	N/A
9243038	12/2015	Liu et al.	N/A	N/A
9267127	12/2015	Liu et al.	N/A	N/A
9290773	12/2015	Edgerton	N/A	N/A
9296790	12/2015	Chatterjee et al.	N/A	N/A
9322006	12/2015	Liu et al.	N/A	N/A
9322037	12/2015	Liu et al.	N/A	N/A
9340799	12/2015	Liu et al.	N/A	N/A
9340800	12/2015	Liu et al.	N/A	N/A
9359599	12/2015	Liu et al.	N/A	N/A
9388430	12/2015	Liu et al.	N/A	N/A
9394537	12/2015	Liu et al.	N/A	N/A
9434774	12/2015	Liu et al.	N/A	N/A
9458484	12/2015	Ma et al.	N/A	N/A
9512446	12/2015	Joung et al.	N/A	N/A
9526784	12/2015	Liu et al.	N/A	N/A
9534210	12/2016	Park et al.	N/A	N/A
9580698	12/2016	Xu et al.	N/A	N/A
9593356	12/2016	Haugwitz et al.	N/A	N/A
9610322	12/2016	Liu et al.	N/A	N/A
9637739	12/2016	Siksnys et al.	N/A	N/A
9663770	12/2016	Rogers et al.	N/A	N/A
9695446	12/2016	Mangeot et al.	N/A	N/A
9737604	12/2016	Liu et al.	N/A	N/A
9738693	12/2016	Telford et al.	N/A	N/A
9753340	12/2016	Saitou	N/A	N/A
9765304	12/2016	Klatzmann et al.	N/A	N/A
9771574	12/2016	Liu et al.	N/A	N/A
9777043	12/2016	Anderson et al.	N/A	N/A
9783791	12/2016	Hogrefe et al.	N/A	N/A
9816093	12/2016	Donohoue et al.	N/A	N/A
9840538	12/2016	Telford et al.	N/A	N/A
9840690	12/2016	Karli et al.	N/A	N/A
9840699	12/2016	Liu et al.	N/A	N/A
9840702	12/2016	Collingwood et al.	N/A	N/A
9850521	12/2016	Braman et al.	N/A	N/A
9873907	12/2017	Zeiner et al.	N/A	N/A
9879270	12/2017	Hittinger et al.	N/A	N/A
9914939	12/2017	Church et al.	N/A	N/A
9932567	12/2017	Xu et al.	N/A	N/A
9938288	12/2017	Kishi et al.	N/A	N/A
9944933	12/2017	Storici et al.	N/A	N/A
9982279	12/2017	Gill et al.	N/A	N/A
9999671	12/2017	Liu et al.	N/A	N/A
10011868	12/2017	Liu et al.	N/A	N/A
10040830	12/2017	Chatterjee et al.	N/A	N/A
10053725	12/2017	Liu et al.	N/A	N/A
10059940	12/2017	Zhong	N/A	N/A
10077453	12/2017	Liu et al.	N/A	N/A
10113163	12/2017	Liu et al.	N/A	N/A
10150955	12/2017	Lambowitz et al.	N/A	N/A
10167457	12/2018	Liu et al.	N/A	N/A
10179911	12/2018	Liu et al.	N/A	N/A
10189831	12/2018	Arrington et al.	N/A	N/A
10202593	12/2018	Liu et al.	N/A	N/A

10202658	12/2018	Parkin et al.	N/A	N/A
10227581	12/2018	Liu et al.	N/A	N/A
10323236	12/2018	Liu et al.	N/A	N/A
10336997	12/2018	Liu et al.	N/A	N/A
10358670	12/2018	Janulaitis et al.	N/A	N/A
10392674	12/2018	Liu et al.	N/A	N/A
10407474	12/2018	Liu et al.	N/A	N/A
10407695	12/2018	Charneau et al.	N/A	N/A
10407697	12/2018	Doudna et al.	N/A	N/A
10465176	12/2018	Liu et al.	N/A	N/A
10508298	12/2018	Liu et al.	N/A	N/A
10583201	12/2019	Chen et al.	N/A	N/A
10597679	12/2019	Liu et al.	N/A	N/A
10612011	12/2019	Liu et al.	N/A	N/A
10640767	12/2019	Maianti et al.	N/A	N/A
10682410	12/2019	Liu et al.	N/A	N/A
10704062	12/2019	Liu et al.	N/A	N/A
10745677	12/2019	Maianti et al.	N/A	N/A
10858639	12/2019	Liu et al.	N/A	N/A
10912833	12/2020	Liu et al.	N/A	N/A
10930367	12/2020	Zhang et al.	N/A	N/A
10947530	12/2020	Liu et al.	N/A	N/A
10954548	12/2020	Liu et al.	N/A	N/A
11046948	12/2020	Liu et al.	N/A	N/A
11053481	12/2020	Liu et al.	N/A	N/A
11124782	12/2020	Liu et al.	N/A	N/A
11214780	12/2021	Liu	N/A	A61P 17/00
11319532	12/2021	Liu et al.	N/A	N/A
11421016	12/2021	Nguyen et al.	N/A	N/A
11447770	12/2021	Liu et al.	N/A	N/A
11542496	12/2022	Liu et al.	N/A	N/A
11542509	12/2022	Maianti et al.	N/A	N/A
11560566	12/2022	Liu et al.	N/A	N/A
11578343	12/2022	Liu et al.	N/A	N/A
11643652	12/2022	Liu et al.	N/A	N/A
11661590	12/2022	Liu et al.	N/A	N/A
11702651	12/2022	Liu et al.	N/A	N/A
11732274	12/2022	Liu et al.	N/A	N/A
11795443	12/2022	Liu et al.	N/A	N/A
11795452	12/2022	Liu et al.	N/A	N/A
11820969	12/2022	Maianti et al.	N/A	N/A
11898179	12/2023	Maianti et al.	N/A	N/A
11912985	12/2023	Liu et al.	N/A	N/A
11920181	12/2023	Liu et al.	N/A	N/A
11932884	12/2023	Liu et al.	N/A	N/A
11999947	12/2023	Liu et al.	N/A	N/A
12006520	12/2023	Liu et al.	N/A	N/A
12031126	12/2023	Liu et al.	N/A	N/A
12043852	12/2023	Liu et al.	N/A	N/A
12084663	12/2023	Maianti et al.	N/A	N/A
12157760	12/2023	Liu et al.	N/A	N/A
12215365	12/2024	Liu et al.	N/A	N/A
2003/0082575	12/2002	Schultz et al.	N/A	N/A
2003/0087817	12/2002	Cox et al.	N/A	N/A
2003/0096337	12/2002	Hillman et al.	N/A	N/A
2003/0108885	12/2002	Schultz et al.	N/A	N/A
2003/0119764	12/2002	Loeb et al.	N/A	N/A
2003/0167533	12/2002	Yadav et al.	N/A	N/A
2003/0203480	12/2002	Kovesdi et al.	N/A	N/A
2004/0003420	12/2003	Kuhn et al.	N/A	N/A
2004/0028687	12/2003	Waelti	N/A	N/A

2004/0115184	12/2003	Smith et al.	N/A	N/A
2004/0156861	12/2003	Figdor et al.	N/A	N/A
2004/0197892	12/2003	Moore et al.	N/A	N/A
2004/0203109	12/2003	Lal et al.	N/A	N/A
2005/0136429	12/2004	Guarente et al.	N/A	N/A
2005/0222030	12/2004	Allison	N/A	N/A
2005/0260626	12/2004	Lorens et al.	N/A	N/A
2006/0088864	12/2005	Smolke et al.	N/A	N/A
2006/0104984	12/2005	Littlefield et al.	N/A	N/A
2006/0216702	12/2005	Compans et al.	N/A	N/A
2006/0246568	12/2005	Honjo et al.	N/A	N/A
2007/0015238	12/2006	Snyder et al.	N/A	N/A
2007/0049533	12/2006	Liu et al.	N/A	N/A
2007/0264692	12/2006	Liu et al.	N/A	N/A
2007/0269817	12/2006	Shapero	N/A	N/A
2007/0298118	12/2006	Lotvall et al.	N/A	N/A
2008/0008697	12/2007	Mintier et al.	N/A	N/A
2008/0051317	12/2007	Church et al.	N/A	N/A
2008/0124725	12/2007	Barrangou et al.	N/A	N/A
2008/0182254	12/2007	Hall et al.	N/A	N/A
2008/0220502	12/2007	Schellenberger et al.	N/A	N/A
2008/0241917	12/2007	Akita et al.	N/A	N/A
2008/0268516	12/2007	Perreault et al.	N/A	N/A
2009/0111119	12/2008	Doyon et al.	N/A	N/A
2009/0130718	12/2008	Short	N/A	N/A
2009/0202622	12/2008	Fleury et al.	N/A	N/A
2009/0215878	12/2008	Tan et al.	N/A	N/A
2009/0234109	12/2008	Han et al.	N/A	N/A
2010/0076057	12/2009	Sontheimer et al.	N/A	N/A
2010/0093617	12/2009	Barrangou et al.	N/A	N/A
2010/0104690	12/2009	Barrangou et al.	N/A	N/A
2010/0105134	12/2009	Quay et al.	N/A	N/A
2010/0273857	12/2009	Thakker et al.	N/A	N/A
2010/0305197	12/2009	Che	N/A	N/A
2010/0316643	12/2009	Eckert et al.	N/A	N/A
2011/0016540	12/2010	Weinstein et al.	N/A	N/A
2011/0059160	12/2010	Essner et al.	N/A	N/A
2011/0059502	12/2010	Chalasani	N/A	N/A
2011/0104787	12/2010	Church et al.	N/A	N/A
2011/0123509	12/2010	Jantz et al.	N/A	N/A
2011/0177495	12/2010	Liu et al.	N/A	N/A
2011/0189775	12/2010	Ainley et al.	N/A	N/A
2011/0189776	12/2010	Terns et al.	N/A	N/A
2011/0206672	12/2010	Little	N/A	N/A
2011/0217739	12/2010	Terns et al.	N/A	N/A
2011/0301073	12/2010	Gregory et al.	N/A	N/A
2012/0129759	12/2011	Liu et al.	N/A	N/A
2012/0141523	12/2011	Castado et al.	N/A	N/A
2012/0159653	12/2011	Weinstein et al.	N/A	N/A
2012/0244601	12/2011	Bertozzi et al.	N/A	N/A
2012/0270273	12/2011	Zhang et al.	N/A	N/A
2012/0322861	12/2011	Byrne et al.	N/A	N/A
2013/0022980	12/2012	Nelson et al.	N/A	N/A
2013/0053426	12/2012	Seow et al.	N/A	N/A
2013/0059931	12/2012	Petersen-Mahrt et al.	N/A	N/A
2013/0108657	12/2012	Yee et al.	N/A	N/A
2013/0117869	12/2012	Duchateau et al.	N/A	N/A
2013/0130248	12/2012	Haurwitz et al.	N/A	N/A
2013/0158245	12/2012	Russell et al.	N/A	N/A
2013/0165389	12/2012	Schellenberger et al.	N/A	N/A
2013/0212725	12/2012	Kuhn et al.	N/A	N/A

2013/0309720	12/2012	Schultz et al.	N/A	N/A
2013/0344117	12/2012	Mirosevich et al.	N/A	N/A
2013/0345064	12/2012	Liu et al.	N/A	N/A
2014/0004280	12/2013	Loomis	N/A	N/A
2014/0005269	12/2013	Ngwuluka et al.	N/A	N/A
2014/0017214	12/2013	Cost	N/A	N/A
2014/0018404	12/2013	Chen et al.	N/A	N/A
2014/0044793	12/2013	Goll et al.	N/A	N/A
2014/0065711	12/2013	Liu et al.	N/A	N/A
2014/0068797	12/2013	Doudna et al.	N/A	N/A
2014/0127752	12/2013	Zhou et al.	N/A	N/A
2014/0128449	12/2013	Liu et al.	N/A	N/A
2014/0141094	12/2013	Smyth et al.	N/A	N/A
2014/0141487	12/2013	Feldman et al.	N/A	N/A
2014/0179770	12/2013	Zhang et al.	N/A	N/A
2014/0186843	12/2013	Zhang et al.	N/A	N/A
2014/0186919	12/2013	Zhang et al.	N/A	N/A
2014/0186958	12/2013	Zhang et al.	N/A	N/A
2014/0201858	12/2013	Ostertag et al.	N/A	N/A
2014/0234289	12/2013	Liu et al.	N/A	N/A
2014/0248702	12/2013	Zhang et al.	N/A	N/A
2014/0273037	12/2013	Wu	N/A	N/A
2014/0273226	12/2013	Wu	N/A	N/A
2014/0273230	12/2013	Chen et al.	N/A	N/A
2014/0273234	12/2013	Zhang et al.	N/A	N/A
2014/0283156	12/2013	Zador et al.	N/A	N/A
2014/0295556	12/2013	Joung et al.	N/A	N/A
2014/0295557	12/2013	Joung et al.	N/A	N/A
2014/0342456	12/2013	Mali et al.	N/A	N/A
2014/0342457	12/2013	Mali et al.	N/A	N/A
2014/0342458	12/2013	Mali et al.	N/A	N/A
2014/0349400	12/2013	Jakimo et al.	N/A	N/A
2014/0356867	12/2013	Peter et al.	N/A	N/A
2014/0356956	12/2013	Church et al.	N/A	N/A
2014/0356958	12/2013	Mali et al.	N/A	N/A
2014/0356959	12/2013	Church et al.	N/A	N/A
2014/0357523	12/2013	Zeiner et al.	N/A	N/A
2014/0377868	12/2013	Joung et al.	N/A	N/A
2015/0010526	12/2014	Liu et al.	N/A	N/A
2015/0031089	12/2014	Lindstrom	N/A	N/A
2015/0031132	12/2014	Church et al.	N/A	N/A
2015/0031133	12/2014	Church et al.	N/A	N/A
2015/0044191	12/2014	Liu et al.	N/A	N/A
2015/0044192	12/2014	Liu et al.	N/A	N/A
2015/0044772	12/2014	Zhao	N/A	N/A
2015/0050699	12/2014	Siksnys et al.	N/A	N/A
2015/0056177	12/2014	Liu et al.	N/A	N/A
2015/0056629	12/2014	Guthrie-Honea	N/A	N/A
2015/0064138	12/2014	Lu et al.	N/A	N/A
2015/0064789	12/2014	Paschon et al.	N/A	N/A
2015/0071898	12/2014	Liu et al.	N/A	N/A
2015/0071899	12/2014	Liu et al.	N/A	N/A
2015/0071900	12/2014	Liu et al.	N/A	N/A
2015/0071901	12/2014	Liu et al.	N/A	N/A
2015/0071902	12/2014	Liu et al.	N/A	N/A
2015/0071903	12/2014	Liu et al.	N/A	N/A
2015/0071906	12/2014	Liu et al.	N/A	N/A
2015/0079680	12/2014	Bradley et al.	N/A	N/A
2015/0079681	12/2014	Zhang	N/A	N/A
2015/0098954	12/2014	Hyde et al.	N/A	N/A
2015/0118216	12/2014	Liu et al.	N/A	N/A



2015/0128300	12/2014	Warming et al.	N/A	N/A
2015/0132269	12/2014	Orkin et al.	N/A	N/A
2015/0140664	12/2014	Byrne et al.	N/A	N/A
2015/0159172	12/2014	Miller et al.	N/A	N/A
2015/0165054	12/2014	Liu et al.	N/A	N/A
2015/0166980	12/2014	Liu et al.	N/A	N/A
2015/0166981	12/2014	Liu et al.	N/A	N/A
2015/0166982	12/2014	Liu et al.	N/A	N/A
2015/0166983	12/2014	Liu et al.	N/A	N/A
2015/0166984	12/2014	Liu et al.	N/A	N/A
2015/0166985	12/2014	Liu et al.	N/A	N/A
2015/0191744	12/2014	Wolfe et al.	N/A	N/A
2015/0197759	12/2014	Xu et al.	N/A	N/A
2015/0211058	12/2014	Carstens	N/A	N/A
2015/0218573	12/2014	Loque et al.	N/A	N/A
2015/0225773	12/2014	Farmer et al.	N/A	N/A
2015/0241440	12/2014	Fasan et al.	N/A	N/A
2015/0252358	12/2014	Maeder et al.	N/A	N/A
2015/0275202	12/2014	Liu et al.	N/A	N/A
2015/0291965	12/2014	Zhang et al.	N/A	N/A
2015/0307889	12/2014	Petolino et al.	N/A	N/A
2015/0315252	12/2014	Haugwitz et al.	N/A	N/A
2015/0344549	12/2014	Muir et al.	N/A	N/A
2016/0017393	12/2015	Jacobson et al.	N/A	N/A
2016/0017396	12/2015	Cann et al.	N/A	N/A
2016/0032292	12/2015	Storici et al.	N/A	N/A
2016/0032353	12/2015	Braman et al.	N/A	N/A
2016/0040155	12/2015	Maizels et al.	N/A	N/A
2016/0046952	12/2015	Hittinger et al.	N/A	N/A
2016/0046961	12/2015	Jinek et al.	N/A	N/A
2016/0046962	12/2015	May et al.	N/A	N/A
2016/0053272	12/2015	Wurtzel et al.	N/A	N/A
2016/0053304	12/2015	Wurtzel et al.	N/A	N/A
2016/0074535	12/2015	Ranganathan et al.	N/A	N/A
2016/0076093	12/2015	Shendure et al.	N/A	N/A
2016/0090603	12/2015	Carnes et al.	N/A	N/A
2016/0090622	12/2015	Liu et al.	N/A	N/A
2016/0115488	12/2015	Zhang et al.	N/A	N/A
2016/0137716	12/2015	El Andaloussi et al.	N/A	N/A
2016/0138046	12/2015	Wu	N/A	N/A
2016/0153003	12/2015	Joung et al.	N/A	N/A
2016/0186214	12/2015	Brouns et al.	N/A	N/A
2016/0200779	12/2015	Liu et al.	N/A	N/A
2016/0201040	12/2015	Liu et al.	N/A	N/A
2016/0201089	12/2015	Gersbach et al.	N/A	N/A
2016/0206566	12/2015	Lu et al.	N/A	N/A
2016/0208243	12/2015	Zhang et al.	N/A	N/A
2016/0208288	12/2015	Liu et al.	N/A	N/A
2016/0215275	12/2015	Zhong	N/A	N/A
2016/0215276	12/2015	Liu et al.	N/A	N/A
2016/0215300	12/2015	May et al.	N/A	N/A
2016/0244784	12/2015	Jacobson et al.	N/A	N/A
2016/0244829	12/2015	Bang et al.	N/A	N/A
2016/0264934	12/2015	Giallourakis et al.	N/A	N/A
2016/0272593	12/2015	Ritter et al.	N/A	N/A
2016/0272965	12/2015	Zhang et al.	N/A	N/A
2016/0281072	12/2015	Zhang	N/A	N/A
2016/0298136	12/2015	Chen et al.	N/A	N/A
2016/0304846	12/2015	Liu et al.	N/A	N/A
2016/0304855	12/2015	Stark et al.	N/A	N/A
2016/0312304	12/2015	Sorrentino et al.	N/A	N/A

2016/0319262	12/2015	Doudna et al.	N/A	N/A
2016/0333389	12/2015	Liu et al.	N/A	N/A
2016/0340622	12/2015	Abdou	N/A	N/A
2016/0340661	12/2015	Cong et al.	N/A	N/A
2016/0340662	12/2015	Zhang et al.	N/A	N/A
2016/0345578	12/2015	Barrangou et al.	N/A	N/A
2016/0346360	12/2015	Quake et al.	N/A	N/A
2016/0346361	12/2015	Quake et al.	N/A	N/A
2016/0346362	12/2015	Quake et al.	N/A	N/A
2016/0348074	12/2015	Quake et al.	N/A	N/A
2016/0348096	12/2015	Liu et al.	N/A	N/A
2016/0350476	12/2015	Quake et al.	N/A	N/A
2016/0355796	12/2015	Davidson et al.	N/A	N/A
2016/0369262	12/2015	Reik et al.	N/A	N/A
2017/0009224	12/2016	Liu et al.	N/A	N/A
2017/0009242	12/2016	McKinley et al.	N/A	N/A
2017/0014449	12/2016	Bangera et al.	N/A	N/A
2017/0020922	12/2016	Wagner et al.	N/A	N/A
2017/0022251	12/2016	Rammensee et al.	N/A	N/A
2017/0037432	12/2016	Donohoue et al.	N/A	N/A
2017/0044520	12/2016	Liu et al.	N/A	N/A
2017/0044592	12/2016	Peter et al.	N/A	N/A
2017/0053729	12/2016	Kotani et al.	N/A	N/A
2017/0058271	12/2016	Joung et al.	N/A	N/A
2017/0058272	12/2016	Carter et al.	N/A	N/A
2017/0058298	12/2016	Kennedy et al.	N/A	N/A
2017/0073663	12/2016	Wang et al.	N/A	N/A
2017/0073670	12/2016	Nishida et al.	N/A	N/A
2017/0087224	12/2016	Quake	N/A	N/A
2017/0087225	12/2016	Quake	N/A	N/A
2017/0088587	12/2016	Quake	N/A	N/A
2017/0088828	12/2016	Quake	N/A	N/A
2017/0107536	12/2016	Zhang et al.	N/A	N/A
2017/0107560	12/2016	Peter et al.	N/A	N/A
2017/0112773	12/2016	Stachowiak et al.	N/A	N/A
2017/0114367	12/2016	Hu et al.	N/A	N/A
2017/0121693	12/2016	Liu et al.	N/A	N/A
2017/0145394	12/2016	Yeo et al.	N/A	N/A
2017/0145405	12/2016	Tang et al.	N/A	N/A
2017/0145438	12/2016	Kantor	N/A	N/A
2017/0152528	12/2016	Zhang	N/A	N/A
2017/0152787	12/2016	Kubo et al.	N/A	N/A
2017/0159033	12/2016	Kamtekar et al.	N/A	N/A
2017/0166928	12/2016	Vyas et al.	N/A	N/A
2017/0173113	12/2016	Besner et al.	N/A	N/A
2017/0175086	12/2016	Schmitt et al.	N/A	N/A
2017/0175104	12/2016	Doudna et al.	N/A	N/A
2017/0175142	12/2016	Zhang et al.	N/A	N/A
2017/0191047	12/2016	Terns et al.	N/A	N/A
2017/0191078	12/2016	Zhang et al.	N/A	N/A
2017/0198269	12/2016	Zhang et al.	N/A	N/A
2017/0198277	12/2016	Kmiec et al.	N/A	N/A
2017/0198302	12/2016	Feng et al.	N/A	N/A
2017/0211061	12/2016	Weiss et al.	N/A	N/A
2017/0224843	12/2016	Deglon et al.	N/A	N/A
2017/0226522	12/2016	Hu et al.	N/A	N/A
2017/0233703	12/2016	Xie et al.	N/A	N/A
2017/0233708	12/2016	Liu et al.	N/A	N/A
2017/0233756	12/2016	Begemann et al.	N/A	N/A
2017/0247671	12/2016	Yung et al.	N/A	N/A
2017/0247703	12/2016	Sloan et al.	N/A	N/A

2017/0268022	12/2016	Liu et al.	N/A	N/A
2017/0275648	12/2016	Barrangou et al.	N/A	N/A
2017/0275665	12/2016	Silas et al.	N/A	N/A
2017/0283797	12/2016	Robb et al.	N/A	N/A
2017/0283831	12/2016	Zhang et al.	N/A	N/A
2017/0306306	12/2016	Potter et al.	N/A	N/A
2017/0314016	12/2016	Kim et al.	N/A	N/A
2017/0362635	12/2016	Chamberlain et al.	N/A	N/A
2018/0023062	12/2017	Lamb et al.	N/A	N/A
2018/0037877	12/2017	Gao et al.	N/A	N/A
2018/0064077	12/2017	Dunham et al.	N/A	N/A
2018/0066258	12/2017	Powell	N/A	N/A
2018/0068062	12/2017	Zhang et al.	N/A	N/A
2018/0073012	12/2017	Liu et al.	N/A	N/A
2018/0080051	12/2017	Sheikh et al.	N/A	N/A
2018/0087046	12/2017	Badran et al.	N/A	N/A
2018/0100147	12/2017	Yates et al.	N/A	N/A
2018/0105867	12/2017	Xiao et al.	N/A	N/A
2018/0119118	12/2017	Lu et al.	N/A	N/A
2018/0127759	12/2017	Lu et al.	N/A	N/A
2018/0127780	12/2017	Liu et al.	N/A	N/A
2018/0155708	12/2017	Church et al.	N/A	N/A
2018/0155720	12/2017	Donohoue et al.	N/A	N/A
2018/0163213	12/2017	Aneja et al.	N/A	N/A
2018/0170984	12/2017	Harris et al.	N/A	N/A
2018/0177727	12/2017	Kalluri et al.	N/A	N/A
2018/0179503	12/2017	Maianti et al.	N/A	N/A
2018/0179547	12/2017	Zhang et al.	N/A	N/A
2018/0201921	12/2017	Malcolm	N/A	N/A
2018/0230464	12/2017	Zhong	N/A	N/A
2018/0230471	12/2017	Storici et al.	N/A	N/A
2018/0236081	12/2017	Liu et al.	N/A	N/A
2018/0237787	12/2017	Maianti et al.	N/A	N/A
2018/0245066	12/2017	Yao et al.	N/A	N/A
2018/0258418	12/2017	Kim	N/A	N/A
2018/0265864	12/2017	Li et al.	N/A	N/A
2018/0273935	12/2017	Lane et al.	N/A	N/A
2018/0273939	12/2017	Yu et al.	N/A	N/A
2018/0273976	12/2017	Ümit et al.	N/A	N/A
2018/0282722	12/2017	Jakimo et al.	N/A	N/A
2018/0298391	12/2017	Jakimo et al.	N/A	N/A
2018/0305688	12/2017	Zhong	N/A	N/A
2018/0305704	12/2017	Zhang	N/A	N/A
2018/0312822	12/2017	Lee et al.	N/A	N/A
2018/0312825	12/2017	Liu et al.	N/A	N/A
2018/0312828	12/2017	Liu et al.	N/A	N/A
2018/0312835	12/2017	Yao et al.	N/A	N/A
2018/0327756	12/2017	Zhang et al.	N/A	N/A
2018/0346927	12/2017	Doudna et al.	N/A	N/A
2018/0371497	12/2017	Gill et al.	N/A	N/A
2019/0010481	12/2018	Joung et al.	N/A	N/A
2019/0032053	12/2018	Ji et al.	N/A	N/A
2019/0055543	12/2018	Tran et al.	N/A	N/A
2019/0055549	12/2018	Capurso et al.	N/A	N/A
2019/0062734	12/2018	Cotta-Ramusino et al.	N/A	N/A
2019/0093099	12/2018	Liu et al.	N/A	N/A
2019/0135869	12/2018	Chatterjee et al.	N/A	N/A
2019/0167810	12/2018	Hean et al.	N/A	N/A
2019/0185883	12/2018	Liu et al.	N/A	N/A
2019/0203228	12/2018	Bouille et al.	N/A	N/A
2019/0218547	12/2018	Lee et al.	N/A	N/A

2019/0224331	12/2018	Wiklander	N/A	N/A
2019/0225955	12/2018	Liu et al.	N/A	N/A
2019/0233847	12/2018	Savage et al.	N/A	N/A
2019/0241633	12/2018	Fotin-Mleczek et al.	N/A	N/A
2019/0256842	12/2018	Liu et al.	N/A	N/A
2019/0264202	12/2018	Church et al.	N/A	N/A
2019/0276816	12/2018	Liu et al.	N/A	N/A
2019/0309290	12/2018	Neuteboom et al.	N/A	N/A
2019/0322992	12/2018	Liu et al.	N/A	N/A
2019/0330619	12/2018	Smith et al.	N/A	N/A
2019/0352632	12/2018	Liu et al.	N/A	N/A
2019/0367891	12/2018	Liu et al.	N/A	N/A
2019/0388347	12/2018	Wiklander et al.	N/A	N/A
2020/0010818	12/2019	Liu et al.	N/A	N/A
2020/0010835	12/2019	Maianti et al.	N/A	N/A
2020/0023012	12/2019	Joseph et al.	N/A	N/A
2020/0056206	12/2019	Tremblay et al.	N/A	N/A
2020/0060980	12/2019	Von Maltzahn et al.	N/A	N/A
2020/0062813	12/2019	Nordin et al.	N/A	N/A
2020/0063127	12/2019	Lu et al.	N/A	N/A
2020/0071722	12/2019	Liu et al.	N/A	N/A
2020/0109398	12/2019	Rubens et al.	N/A	N/A
2020/0172931	12/2019	Liu et al.	N/A	N/A
2020/0181619	12/2019	Tang et al.	N/A	N/A
2020/0190493	12/2019	Liu et al.	N/A	N/A
2020/0255868	12/2019	Liu et al.	N/A	N/A
2020/0277587	12/2019	Liu et al.	N/A	N/A
2020/0323984	12/2019	Liu et al.	N/A	N/A
2020/0399619	12/2019	Maianti et al.	N/A	N/A
2020/0399626	12/2019	Liu et al.	N/A	N/A
2021/0054416	12/2020	Liu et al.	N/A	N/A
2021/0115428	12/2020	Maianti et al.	N/A	N/A
2021/0198330	12/2020	Liu et al.	N/A	N/A
2021/0214698	12/2020	Liu et al.	N/A	N/A
2021/0230577	12/2020	Liu et al.	N/A	N/A
2021/0254127	12/2020	Liu et al.	N/A	N/A
2021/0315994	12/2020	Liu et al.	N/A	N/A
2021/0317440	12/2020	Liu et al.	N/A	N/A
2022/0033785	12/2021	Liu et al.	N/A	N/A
2022/0170013	12/2021	Liu et al.	N/A	N/A
2022/0177877	12/2021	Church et al.	N/A	N/A
2022/0204975	12/2021	Liu et al.	N/A	N/A
2022/0213507	12/2021	Liu et al.	N/A	N/A
2022/0220462	12/2021	Liu et al.	N/A	N/A
2022/0238182	12/2021	Shen et al.	N/A	N/A
2022/0249697	12/2021	Liu et al.	N/A	N/A
2022/0282275	12/2021	Liu et al.	N/A	N/A
2022/0290115	12/2021	Liu et al.	N/A	N/A
2022/0307001	12/2021	Liu et al.	N/A	N/A
2022/0307003	12/2021	Liu et al.	N/A	N/A
2022/0315906	12/2021	Liu et al.	N/A	N/A
2022/0356469	12/2021	Liu et al.	N/A	N/A
2022/0380740	12/2021	Liu et al.	N/A	N/A
2022/0389395	12/2021	Liu et al.	N/A	N/A
2023/0002745	12/2022	Liu et al.	N/A	N/A
2023/0021641	12/2022	Liu et al.	N/A	N/A
2023/0056852	12/2022	Liu et al.	N/A	N/A
2023/0058176	12/2022	Liu et al.	N/A	N/A
2023/0078265	12/2022	Liu et al.	N/A	N/A
2023/0086199	12/2022	Liu et al.	N/A	N/A
2023/0090221	12/2022	Liu et al.	N/A	N/A

2023/0108687	12/2022	Liu et al.	N/A	N/A
2023/0123669	12/2022	Liu et al.	N/A	N/A
2023/0127008	12/2022	Liu et al.	N/A	N/A
2023/0159913	12/2022	Liu et al.	N/A	N/A
2023/0193295	12/2022	Maianti et al.	N/A	N/A
2023/0220374	12/2022	Liu et al.	N/A	N/A
2023/0272425	12/2022	Liu et al.	N/A	N/A
2023/0279443	12/2022	Liu et al.	N/A	N/A
2023/0332144	12/2022	Liu et al.	N/A	N/A
2023/0340465	12/2022	Liu et al.	N/A	N/A
2023/0340466	12/2022	Liu et al.	N/A	N/A
2023/0340467	12/2022	Liu et al.	N/A	N/A
2023/0348883	12/2022	Liu et al.	N/A	N/A
2023/0357766	12/2022	Liu et al.	N/A	N/A
2023/0383289	12/2022	Liu et al.	N/A	N/A
2024/0035017	12/2023	Liu et al.	N/A	N/A
2024/0076652	12/2023	Liu et al.	N/A	N/A
2024/0110166	12/2023	Maianti et al.	N/A	N/A
2024/0124866	12/2023	Liu et al.	N/A	N/A
2024/0173430	12/2023	Liu et al.	N/A	N/A
2024/0209329	12/2023	Liu et al.	N/A	N/A
2024/0229077	12/2023	Liu et al.	N/A	N/A
2024/0271116	12/2023	Maianti et al.	N/A	N/A
2024/0287487	12/2023	Liu et al.	N/A	N/A
2024/0327872	12/2023	Liu et al.	N/A	N/A
2024/0401018	12/2023	Liu et al.	N/A	N/A
2024/0417715	12/2023	Liu et al.	N/A	N/A
2024/0417719	12/2023	Liu et al.	N/A	N/A
2024/0417753	12/2023	Liu et al.	N/A	N/A
2025/0011748	12/2024	Liu et al.	N/A	N/A
2025/0027114	12/2024	Liu et al.	N/A	N/A
2025/0034549	12/2024	Liu et al.	N/A	N/A
2025/0059244	12/2024	Liu et al.	N/A	N/A
2025/0064979	12/2024	Liu et al.	N/A	N/A
2025/0064981	12/2024	Liu et al.	N/A	N/A

**FOREIGN PATENT DOCUMENTS**

Patent No.	Application Date	Country	CPC
2012244264	12/2011	AU	N/A
2012354062	12/2013	AU	N/A
2015252023	12/2014	AU	N/A
2015101792	12/2015	AU	N/A
112015013786	12/2016	BR	N/A
2480696	12/2002	CA	N/A
2894668	12/2013	CA	N/A
2894681	12/2013	CA	N/A
2894684	12/2013	CA	N/A
2852593	12/2014	CA	N/A
1069962	12/1992	CN	N/A
101460619	12/2008	CN	N/A
101873862	12/2009	CN	N/A
104342457	12/2010	CN	N/A
102057039	12/2010	CN	N/A
102892777	12/2012	CN	N/A
103224947	12/2012	CN	N/A
103233028	12/2012	CN	N/A
103388006	12/2012	CN	N/A
103614415	12/2013	CN	N/A
103642836	12/2013	CN	N/A
103668472	12/2013	CN	N/A
103820441	12/2013	CN	N/A
103820454	12/2013	CN	N/A

103911376	12/2013	CN	N/A
103923911	12/2013	CN	N/A
103088008	12/2013	CN	N/A
103981211	12/2013	CN	N/A
103981212	12/2013	CN	N/A
104004778	12/2013	CN	N/A
104004782	12/2013	CN	N/A
104017821	12/2013	CN	N/A
104109687	12/2013	CN	N/A
104178461	12/2013	CN	N/A
104404036	12/2014	CN	N/A
104450774	12/2014	CN	N/A
104480144	12/2014	CN	N/A
104498493	12/2014	CN	N/A
104504304	12/2014	CN	N/A
104531704	12/2014	CN	N/A
104531705	12/2014	CN	N/A
104560864	12/2014	CN	N/A
104561095	12/2014	CN	N/A
104593418	12/2014	CN	N/A
104593422	12/2014	CN	N/A
104611370	12/2014	CN	N/A
104651392	12/2014	CN	N/A
104651398	12/2014	CN	N/A
104651399	12/2014	CN	N/A
104651401	12/2014	CN	N/A
104673816	12/2014	CN	N/A
104725626	12/2014	CN	N/A
104726449	12/2014	CN	N/A
104726494	12/2014	CN	N/A
104745626	12/2014	CN	N/A
104762321	12/2014	CN	N/A
104805078	12/2014	CN	N/A
104805099	12/2014	CN	N/A
104805118	12/2014	CN	N/A
104846010	12/2014	CN	N/A
104894068	12/2014	CN	N/A
104894075	12/2014	CN	N/A
104928321	12/2014	CN	N/A
105039339	12/2014	CN	N/A
105039399	12/2014	CN	N/A
105063061	12/2014	CN	N/A
105087620	12/2014	CN	N/A
105112422	12/2014	CN	N/A
105112445	12/2014	CN	N/A
105112519	12/2014	CN	N/A
105121648	12/2014	CN	N/A
105132427	12/2014	CN	N/A
105132451	12/2014	CN	N/A
105177038	12/2014	CN	N/A
105177126	12/2014	CN	N/A
105210981	12/2015	CN	N/A
105219799	12/2015	CN	N/A
105238806	12/2015	CN	N/A
105255937	12/2015	CN	N/A
105274144	12/2015	CN	N/A
105296518	12/2015	CN	N/A
105296537	12/2015	CN	N/A
105316324	12/2015	CN	N/A
105316327	12/2015	CN	N/A
105316337	12/2015	CN	N/A

105331607	12/2015	CN	N/A
105331608	12/2015	CN	N/A
105331609	12/2015	CN	N/A
105331627	12/2015	CN	N/A
105400773	12/2015	CN	N/A
105400779	12/2015	CN	N/A
105400810	12/2015	CN	N/A
105441451	12/2015	CN	N/A
105462968	12/2015	CN	N/A
105463003	12/2015	CN	N/A
105463027	12/2015	CN	N/A
105492608	12/2015	CN	N/A
105492609	12/2015	CN	N/A
105505976	12/2015	CN	N/A
105505979	12/2015	CN	N/A
105518134	12/2015	CN	N/A
105518135	12/2015	CN	N/A
105518137	12/2015	CN	N/A
105518138	12/2015	CN	N/A
105518139	12/2015	CN	N/A
105518140	12/2015	CN	N/A
105543228	12/2015	CN	N/A
105543266	12/2015	CN	N/A
105543270	12/2015	CN	N/A
105567688	12/2015	CN	N/A
105567689	12/2015	CN	N/A
105567734	12/2015	CN	N/A
105567735	12/2015	CN	N/A
105567738	12/2015	CN	N/A
105593367	12/2015	CN	N/A
105594664	12/2015	CN	N/A
105602987	12/2015	CN	N/A
105624146	12/2015	CN	N/A
105624187	12/2015	CN	N/A
105646719	12/2015	CN	N/A
105647922	12/2015	CN	N/A
105647962	12/2015	CN	N/A
105647968	12/2015	CN	N/A
105647969	12/2015	CN	N/A
105671070	12/2015	CN	N/A
105671083	12/2015	CN	N/A
105695485	12/2015	CN	N/A
105779448	12/2015	CN	N/A
105779449	12/2015	CN	N/A
105802980	12/2015	CN	N/A
105821039	12/2015	CN	N/A
105821040	12/2015	CN	N/A
105821049	12/2015	CN	N/A
105821072	12/2015	CN	N/A
105821075	12/2015	CN	N/A
105821116	12/2015	CN	N/A
105838733	12/2015	CN	N/A
105861547	12/2015	CN	N/A
105861552	12/2015	CN	N/A
105861554	12/2015	CN	N/A
105886498	12/2015	CN	N/A
105886534	12/2015	CN	N/A
105886616	12/2015	CN	N/A
105907758	12/2015	CN	N/A
105907785	12/2015	CN	N/A
105925608	12/2015	CN	N/A

105934516	12/2015	CN	N/A
105950560	12/2015	CN	N/A
105950626	12/2015	CN	N/A
105950633	12/2015	CN	N/A
105950639	12/2015	CN	N/A
105985985	12/2015	CN	N/A
106011104	12/2015	CN	N/A
106011150	12/2015	CN	N/A
106011167	12/2015	CN	N/A
106011171	12/2015	CN	N/A
106032540	12/2015	CN	N/A
106047803	12/2015	CN	N/A
106047877	12/2015	CN	N/A
106047930	12/2015	CN	N/A
106086008	12/2015	CN	N/A
106086028	12/2015	CN	N/A
106086061	12/2015	CN	N/A
106086062	12/2015	CN	N/A
106103475	12/2015	CN	N/A
106109417	12/2015	CN	N/A
106119275	12/2015	CN	N/A
106119283	12/2015	CN	N/A
106148286	12/2015	CN	N/A
106148370	12/2015	CN	N/A
106148416	12/2015	CN	N/A
106167525	12/2015	CN	N/A
106167808	12/2015	CN	N/A
106167810	12/2015	CN	N/A
106167821	12/2015	CN	N/A
106172238	12/2015	CN	N/A
106190903	12/2015	CN	N/A
106191057	12/2015	CN	N/A
106191061	12/2015	CN	N/A
106191062	12/2015	CN	N/A
106191064	12/2015	CN	N/A
106191071	12/2015	CN	N/A
106191099	12/2015	CN	N/A
106191107	12/2015	CN	N/A
106191113	12/2015	CN	N/A
106191114	12/2015	CN	N/A
106191116	12/2015	CN	N/A
106191124	12/2015	CN	N/A
106222177	12/2015	CN	N/A
106222193	12/2015	CN	N/A
106222203	12/2015	CN	N/A
106232823	12/2015	CN	N/A
106244555	12/2015	CN	N/A
106244557	12/2015	CN	N/A
106244591	12/2015	CN	N/A
106244609	12/2015	CN	N/A
106282241	12/2016	CN	N/A
106318934	12/2016	CN	N/A
106318973	12/2016	CN	N/A
106350540	12/2016	CN	N/A
106367435	12/2016	CN	N/A
106399306	12/2016	CN	N/A
106399311	12/2016	CN	N/A
106399360	12/2016	CN	N/A
106399367	12/2016	CN	N/A
106399375	12/2016	CN	N/A
106399377	12/2016	CN	N/A



106434651	12/2016	CN	N/A
106434663	12/2016	CN	N/A
106434688	12/2016	CN	N/A
106434737	12/2016	CN	N/A
106434748	12/2016	CN	N/A
106434752	12/2016	CN	N/A
106434782	12/2016	CN	N/A
106446600	12/2016	CN	N/A
106479985	12/2016	CN	N/A
106480027	12/2016	CN	N/A
106480036	12/2016	CN	N/A
106480067	12/2016	CN	N/A
106480080	12/2016	CN	N/A
106480083	12/2016	CN	N/A
106480097	12/2016	CN	N/A
106544351	12/2016	CN	N/A
106544353	12/2016	CN	N/A
106544357	12/2016	CN	N/A
106554969	12/2016	CN	N/A
106566838	12/2016	CN	N/A
106701763	12/2016	CN	N/A
106701808	12/2016	CN	N/A
106701818	12/2016	CN	N/A
106701823	12/2016	CN	N/A
106701830	12/2016	CN	N/A
106754912	12/2016	CN	N/A
106755026	12/2016	CN	N/A
106755077	12/2016	CN	N/A
106755088	12/2016	CN	N/A
106755091	12/2016	CN	N/A
106755097	12/2016	CN	N/A
106755424	12/2016	CN	N/A
106801056	12/2016	CN	N/A
106834323	12/2016	CN	N/A
106834341	12/2016	CN	N/A
106834347	12/2016	CN	N/A
106845151	12/2016	CN	N/A
106868008	12/2016	CN	N/A
106868031	12/2016	CN	N/A
106906240	12/2016	CN	N/A
106906242	12/2016	CN	N/A
106916820	12/2016	CN	N/A
106916852	12/2016	CN	N/A
106939303	12/2016	CN	N/A
106947750	12/2016	CN	N/A
106947780	12/2016	CN	N/A
106957830	12/2016	CN	N/A
106957831	12/2016	CN	N/A
106957844	12/2016	CN	N/A
106957855	12/2016	CN	N/A
106957858	12/2016	CN	N/A
106967697	12/2016	CN	N/A
106967726	12/2016	CN	N/A
106978428	12/2016	CN	N/A
106987570	12/2016	CN	N/A
106987757	12/2016	CN	N/A
107012164	12/2016	CN	N/A
107012174	12/2016	CN	N/A
107012213	12/2016	CN	N/A
107012250	12/2016	CN	N/A
107022562	12/2016	CN	N/A

107034188	12/2016	CN	N/A
107034218	12/2016	CN	N/A
107034229	12/2016	CN	N/A
107043775	12/2016	CN	N/A
107043779	12/2016	CN	N/A
107043787	12/2016	CN	N/A
107058320	12/2016	CN	N/A
107058328	12/2016	CN	N/A
107058358	12/2016	CN	N/A
107058372	12/2016	CN	N/A
107083392	12/2016	CN	N/A
107099533	12/2016	CN	N/A
107099850	12/2016	CN	N/A
107119053	12/2016	CN	N/A
107119071	12/2016	CN	N/A
107129999	12/2016	CN	N/A
107130000	12/2016	CN	N/A
107142272	12/2016	CN	N/A
107142282	12/2016	CN	N/A
107177591	12/2016	CN	N/A
107177595	12/2016	CN	N/A
107177625	12/2016	CN	N/A
107177631	12/2016	CN	N/A
107190006	12/2016	CN	N/A
107190008	12/2016	CN	N/A
107217042	12/2016	CN	N/A
107217075	12/2016	CN	N/A
107227307	12/2016	CN	N/A
107227352	12/2016	CN	N/A
107236737	12/2016	CN	N/A
107236739	12/2016	CN	N/A
107236741	12/2016	CN	N/A
107245502	12/2016	CN	N/A
107254485	12/2016	CN	N/A
107266541	12/2016	CN	N/A
107267515	12/2016	CN	N/A
107287245	12/2016	CN	N/A
107298701	12/2016	CN	N/A
107299114	12/2016	CN	N/A
107304435	12/2016	CN	N/A
107312785	12/2016	CN	N/A
107312793	12/2016	CN	N/A
107312795	12/2016	CN	N/A
107312798	12/2016	CN	N/A
107326042	12/2016	CN	N/A
107326046	12/2016	CN	N/A
107354156	12/2016	CN	N/A
107354173	12/2016	CN	N/A
107356793	12/2016	CN	N/A
107362372	12/2016	CN	N/A
107365786	12/2016	CN	N/A
107365804	12/2016	CN	N/A
107384894	12/2016	CN	N/A
107384922	12/2016	CN	N/A
107400677	12/2016	CN	N/A
107418974	12/2016	CN	N/A
107435051	12/2016	CN	N/A
107435069	12/2016	CN	N/A
107446922	12/2016	CN	N/A
107446923	12/2016	CN	N/A
107446924	12/2016	CN	N/A

107446932	12/2016	CN	N/A
107446951	12/2016	CN	N/A
107446954	12/2016	CN	N/A
107460196	12/2016	CN	N/A
107474129	12/2016	CN	N/A
107475300	12/2016	CN	N/A
107488649	12/2016	CN	N/A
107502608	12/2016	CN	N/A
107502618	12/2016	CN	N/A
107513531	12/2016	CN	N/A
107519492	12/2016	CN	N/A
107523567	12/2016	CN	N/A
107523583	12/2016	CN	N/A
107541525	12/2017	CN	N/A
107557373	12/2017	CN	N/A
107557378	12/2017	CN	N/A
107557381	12/2017	CN	N/A
107557390	12/2017	CN	N/A
107557393	12/2017	CN	N/A
107557394	12/2017	CN	N/A
107557455	12/2017	CN	N/A
107574179	12/2017	CN	N/A
107586777	12/2017	CN	N/A
107586779	12/2017	CN	N/A
107604003	12/2017	CN	N/A
107619829	12/2017	CN	N/A
107619837	12/2017	CN	N/A
107630006	12/2017	CN	N/A
107630041	12/2017	CN	N/A
107630042	12/2017	CN	N/A
107630043	12/2017	CN	N/A
107641631	12/2017	CN	N/A
107653256	12/2017	CN	N/A
107686848	12/2017	CN	N/A
206970581	12/2017	CN	N/A
107760652	12/2017	CN	N/A
107760663	12/2017	CN	N/A
107760684	12/2017	CN	N/A
107760715	12/2017	CN	N/A
107784200	12/2017	CN	N/A
107794272	12/2017	CN	N/A
107794276	12/2017	CN	N/A
107815463	12/2017	CN	N/A
107828738	12/2017	CN	N/A
107828794	12/2017	CN	N/A
107828826	12/2017	CN	N/A
107828874	12/2017	CN	N/A
107858346	12/2017	CN	N/A
107858373	12/2017	CN	N/A
107880132	12/2017	CN	N/A
107881184	12/2017	CN	N/A
107893074	12/2017	CN	N/A
107893075	12/2017	CN	N/A
107893076	12/2017	CN	N/A
107893080	12/2017	CN	N/A
107893086	12/2017	CN	N/A
107904261	12/2017	CN	N/A
107937427	12/2017	CN	N/A
107937432	12/2017	CN	N/A
107937501	12/2017	CN	N/A
107974466	12/2017	CN	N/A

107988229	12/2017	CN	N/A
107988246	12/2017	CN	N/A
107988256	12/2017	CN	N/A
107988268	12/2017	CN	N/A
108018316	12/2017	CN	N/A
108034656	12/2017	CN	N/A
108048466	12/2017	CN	N/A
108102940	12/2017	CN	N/A
108103090	12/2017	CN	N/A
108103092	12/2017	CN	N/A
108103098	12/2017	CN	N/A
108103586	12/2017	CN	N/A
108148835	12/2017	CN	N/A
108148837	12/2017	CN	N/A
108148873	12/2017	CN	N/A
108192956	12/2017	CN	N/A
108243575	12/2017	CN	N/A
108251423	12/2017	CN	N/A
108251451	12/2017	CN	N/A
108251452	12/2017	CN	N/A
108342480	12/2017	CN	N/A
108359691	12/2017	CN	N/A
108359712	12/2017	CN	N/A
108384784	12/2017	CN	N/A
108396027	12/2017	CN	N/A
108410877	12/2017	CN	N/A
108410906	12/2017	CN	N/A
108410907	12/2017	CN	N/A
108410911	12/2017	CN	N/A
108424931	12/2017	CN	N/A
108441519	12/2017	CN	N/A
108441520	12/2017	CN	N/A
108472314	12/2017	CN	N/A
108486108	12/2017	CN	N/A
108486111	12/2017	CN	N/A
108486145	12/2017	CN	N/A
108486146	12/2017	CN	N/A
108486154	12/2017	CN	N/A
108486159	12/2017	CN	N/A
108486234	12/2017	CN	N/A
108504657	12/2017	CN	N/A
108504685	12/2017	CN	N/A
108504693	12/2017	CN	N/A
108513575	12/2017	CN	N/A
108546712	12/2017	CN	N/A
108546717	12/2017	CN	N/A
108546718	12/2017	CN	N/A
108559730	12/2017	CN	N/A
108559732	12/2017	CN	N/A
108559745	12/2017	CN	N/A
108559760	12/2017	CN	N/A
108570479	12/2017	CN	N/A
108588071	12/2017	CN	N/A
108588123	12/2017	CN	N/A
108588128	12/2017	CN	N/A
108588182	12/2017	CN	N/A
108610399	12/2017	CN	N/A
108611364	12/2017	CN	N/A
108624622	12/2017	CN	N/A
108642053	12/2017	CN	N/A
108642055	12/2017	CN	N/A

108642077	12/2017	CN	N/A
108642078	12/2017	CN	N/A
108642090	12/2017	CN	N/A
108690844	12/2017	CN	N/A
108699542	12/2017	CN	N/A
108707604	12/2017	CN	N/A
108707620	12/2017	CN	N/A
108707621	12/2017	CN	N/A
108707628	12/2017	CN	N/A
108707629	12/2017	CN	N/A
108715850	12/2017	CN	N/A
108728476	12/2017	CN	N/A
108728486	12/2017	CN	N/A
108753772	12/2017	CN	N/A
108753783	12/2017	CN	N/A
108753813	12/2017	CN	N/A
108753817	12/2017	CN	N/A
108753832	12/2017	CN	N/A
108753835	12/2017	CN	N/A
108753836	12/2017	CN	N/A
108795902	12/2017	CN	N/A
108822217	12/2017	CN	N/A
108823248	12/2017	CN	N/A
108823249	12/2017	CN	N/A
108823291	12/2017	CN	N/A
108841845	12/2017	CN	N/A
108853133	12/2017	CN	N/A
108866093	12/2017	CN	N/A
108893529	12/2017	CN	N/A
108913664	12/2017	CN	N/A
108913691	12/2017	CN	N/A
108913714	12/2017	CN	N/A
108913717	12/2017	CN	N/A
208034188	12/2017	CN	N/A
109517841	12/2018	CN	N/A
0264166	12/1987	EP	N/A
0321201	12/1988	EP	N/A
0519463	12/1991	EP	N/A
1085892	12/2000	EP	N/A
1092770	12/2000	EP	N/A
2350295	12/2012	EP	N/A
2604255	12/2012	EP	N/A
2840140	12/2014	EP	N/A
2877490	12/2014	EP	N/A
2966170	12/2015	EP	N/A
3009511	12/2015	EP	N/A
3031921	12/2015	EP	N/A
3045537	12/2015	EP	N/A
3115457	12/2016	EP	N/A
3144390	12/2016	EP	N/A
2583974	12/2016	EP	N/A
3199632	12/2016	EP	N/A
3216867	12/2016	EP	N/A
3235828	12/2016	EP	N/A
3252160	12/2016	EP	N/A
2498823	12/2017	EP	N/A
3365437	12/2017	EP	N/A
3454889	12/2018	EP	N/A
3008192	12/2018	EP	N/A
3079725	12/2018	EP	N/A
3450553	12/2018	EP	N/A

2740248	12/2019	ES	N/A
2528177	12/2015	GB	N/A
2531454	12/2015	GB	N/A
2542653	12/2016	GB	N/A
1208045	12/2015	HK	N/A
2007-501626	12/2006	JP	N/A
2008-515405	12/2007	JP	N/A
2010-033344	12/2009	JP	N/A
2010-535744	12/2009	JP	N/A
2010-539929	12/2009	JP	N/A
2011-081011	12/2010	JP	N/A
2011-523353	12/2010	JP	N/A
2012-525146	12/2011	JP	N/A
2012-210172	12/2011	JP	N/A
2012-531909	12/2011	JP	N/A
2015-523856	12/2014	JP	N/A
2015-532654	12/2014	JP	N/A
2016-525888	12/2015	JP	N/A
2016-534132	12/2015	JP	N/A
2017-500035	12/2016	JP	N/A
2018-521045	12/2017	JP	N/A
2019-506123	12/2018	JP	N/A
6629734	12/2019	JP	N/A
6633524	12/2019	JP	N/A
6830517	12/2020	JP	N/A
7324523	12/2022	JP	N/A
101584933	12/2015	KR	N/A
2016-0050069	12/2015	KR	N/A
20160133380	12/2015	KR	N/A
20170037025	12/2016	KR	N/A
20170037028	12/2016	KR	N/A
101748575	12/2016	KR	N/A
20170128137	12/2016	KR	N/A
2018-0022465	12/2017	KR	N/A
2016104674	12/2016	RU	N/A
2634395	12/2016	RU	N/A
2652899	12/2017	RU	N/A
2015128057	12/2018	RU	N/A
2015128098	12/2018	RU	N/A
2687451	12/2018	RU	N/A
2019112514	12/2018	RU	N/A
2019127300	12/2018	RU	N/A
2701850	12/2018	RU	N/A
10201707569	12/2016	SG	N/A
10201710486X	12/2017	SG	N/A
10201710487V	12/2017	SG	N/A
10201710488	12/2017	SG	N/A
I608100	12/2016	TW	N/A
201809272	12/2017	TW	N/A
2018-29773	12/2017	TW	N/A
WO 1990/002809	12/1989	WO	N/A
WO 1991/003162	12/1990	WO	N/A
WO 1991/016024	12/1990	WO	N/A
WO 1991/017271	12/1990	WO	N/A
WO 1991/017424	12/1990	WO	N/A
WO 1992/006188	12/1991	WO	N/A
WO 1992/006200	12/1991	WO	N/A
WO 1992/007065	12/1991	WO	N/A
WO 1993/015187	12/1992	WO	N/A
WO 1993/024641	12/1992	WO	N/A
WO 1994/018316	12/1993	WO	N/A

WO 1994/026877	12/1993	WO	N/A
WO 1996/004403	12/1995	WO	N/A
WO 1996/010640	12/1995	WO	N/A
WO 1997/025416	12/1996	WO	N/A
WO 1998/032845	12/1997	WO	N/A
WO 1998/050538	12/1997	WO	N/A
WO 2001/036452	12/2000	WO	N/A
WO 2001/038547	12/2000	WO	N/A
WO 2002/059296	12/2001	WO	N/A
WO 2002/068676	12/2001	WO	N/A
WO 2002/103028	12/2001	WO	N/A
WO 2003/004608	12/2002	WO	N/A
WO 2004/007684	12/2003	WO	N/A
WO 2005/014791	12/2004	WO	N/A
WO 2005/019415	12/2004	WO	N/A
WO 2006/002547	12/2005	WO	N/A
WO 2006/042112	12/2005	WO	N/A
WO 2007/025097	12/2006	WO	N/A
WO 2007/037444	12/2006	WO	N/A
WO 2007/066923	12/2006	WO	N/A
WO 2007/136815	12/2006	WO	N/A
WO 2007/143574	12/2006	WO	N/A
WO 2008/005529	12/2007	WO	N/A
WO 2008/108989	12/2007	WO	N/A
WO 2009/002418	12/2007	WO	N/A
WO 2009/019317	12/2008	WO	N/A
WO 2009/098290	12/2008	WO	N/A
WO 2009/134808	12/2008	WO	N/A
WO 2010/011961	12/2009	WO	N/A
WO 2010/012902	12/2009	WO	N/A
WO 2010/028347	12/2009	WO	N/A
WO 2010/054108	12/2009	WO	N/A
WO 2010/054154	12/2009	WO	N/A
WO 2010/068289	12/2009	WO	N/A
WO 2010/075424	12/2009	WO	N/A
WO 2010/091122	12/2009	WO	N/A
WO 2010/102257	12/2009	WO	N/A
WO 2010/104749	12/2009	WO	C07K 19/00
WO 2010/129019	12/2009	WO	N/A
WO 2010/129023	12/2009	WO	N/A
WO 2010/132092	12/2009	WO	N/A
WO 2010/144150	12/2009	WO	N/A
WO 2011/002503	12/2010	WO	N/A
WO 2011/017293	12/2010	WO	N/A
WO 2011/053868	12/2010	WO	N/A
WO 2011/053982	12/2010	WO	N/A
WO 2011/068810	12/2010	WO	N/A
WO 2011/075627	12/2010	WO	N/A
WO 2011/091311	12/2010	WO	N/A
WO 2011/091396	12/2010	WO	N/A
WO 2011/109031	12/2010	WO	N/A
WO 2011/143124	12/2010	WO	N/A
WO 2011/147590	12/2010	WO	N/A
WO 2011/159369	12/2010	WO	N/A
WO 2012/054726	12/2011	WO	N/A
WO 2012/061815	12/2011	WO	N/A
WO 2012/065043	12/2011	WO	N/A
WO 2012/088381	12/2011	WO	N/A
WO 2012/125445	12/2011	WO	N/A
WO 2012/138927	12/2011	WO	N/A
WO 2012/149470	12/2011	WO	N/A

WO 2012/158985	12/2011	WO	N/A
WO 2012/158986	12/2011	WO	N/A
WO 2012/164565	12/2011	WO	N/A
WO 2012/170930	12/2011	WO	N/A
WO 2013/012674	12/2012	WO	N/A
WO 2013/013105	12/2012	WO	N/A
WO 2013/039857	12/2012	WO	N/A
WO 2013/039861	12/2012	WO	N/A
WO 2013/040093	12/2012	WO	N/A
WO 2013/045632	12/2012	WO	N/A
WO 2013/047844	12/2012	WO	N/A
WO 2013/066438	12/2012	WO	N/A
WO 2013/086441	12/2012	WO	N/A
WO 2013/086444	12/2012	WO	N/A
WO 2013/098244	12/2012	WO	N/A
WO 2013/119602	12/2012	WO	N/A
WO 2013/120022	12/2012	WO	N/A
WO 2013/122617	12/2012	WO	N/A
WO 2013/126794	12/2012	WO	N/A
WO 2013/130683	12/2012	WO	N/A
WO 2013/130824	12/2012	WO	N/A
WO 2013/141680	12/2012	WO	N/A
WO 2013/142578	12/2012	WO	N/A
WO 2013/152359	12/2012	WO	N/A
WO 2013/160230	12/2012	WO	N/A
WO 2013/166315	12/2012	WO	N/A
WO 2013/169398	12/2012	WO	N/A
WO 2013/169802	12/2012	WO	N/A
WO 2013/176772	12/2012	WO	N/A
WO 2013/176915	12/2012	WO	N/A
WO 2013/176916	12/2012	WO	N/A
WO 2013/181440	12/2012	WO	N/A
WO 2013/186754	12/2012	WO	N/A
WO 2013/188037	12/2012	WO	N/A
WO 2013/188522	12/2012	WO	N/A
WO 2013/188638	12/2012	WO	N/A
WO 2013/192278	12/2012	WO	N/A
WO 2013/142378	12/2013	WO	N/A
WO 2014/004336	12/2013	WO	N/A
WO 2014/005042	12/2013	WO	N/A
WO 2014/011237	12/2013	WO	N/A
WO 2014/011901	12/2013	WO	N/A
WO 2014/018423	12/2013	WO	N/A
WO 2014/020608	12/2013	WO	N/A
WO 2014/022120	12/2013	WO	N/A
WO 2014/022702	12/2013	WO	N/A
WO 2014/036219	12/2013	WO	N/A
WO 2014/039513	12/2013	WO	N/A
WO 2014/039523	12/2013	WO	N/A
WO 2014/039585	12/2013	WO	N/A
WO 2014/039684	12/2013	WO	N/A
WO 2014/039692	12/2013	WO	N/A
WO 2014/039702	12/2013	WO	N/A
WO 2014/039872	12/2013	WO	N/A
WO 2014/039970	12/2013	WO	N/A
WO 2014/041327	12/2013	WO	N/A
WO 2014/043143	12/2013	WO	N/A
WO 2014/047103	12/2013	WO	N/A
WO 2014/055782	12/2013	WO	N/A
WO 2014/059173	12/2013	WO	N/A
WO 2014/059255	12/2013	WO	N/A



WO 2014/065596	12/2013	WO	N/A
WO 2014/066505	12/2013	WO	N/A
WO 2014/068346	12/2013	WO	N/A
WO 2014/070887	12/2013	WO	N/A
WO 2014/071006	12/2013	WO	N/A
WO 2014/071219	12/2013	WO	N/A
WO 2014/071235	12/2013	WO	N/A
WO 2014/072941	12/2013	WO	N/A
WO 2014/081729	12/2013	WO	N/A
WO 2014/081730	12/2013	WO	N/A
WO 2014/081855	12/2013	WO	N/A
WO 2014/082644	12/2013	WO	N/A
WO 2014/085261	12/2013	WO	N/A
WO 2014/085593	12/2013	WO	N/A
WO 2014/085830	12/2013	WO	N/A
WO 2014/089212	12/2013	WO	N/A
WO 2014/089290	12/2013	WO	N/A
WO 2014/089348	12/2013	WO	N/A
WO 2014/089513	12/2013	WO	N/A
WO 2014/089533	12/2013	WO	N/A
WO 2014/089541	12/2013	WO	N/A
WO 2014/093479	12/2013	WO	N/A
WO 2014/093595	12/2013	WO	N/A
WO 2014/093622	12/2013	WO	N/A
WO 2014/093635	12/2013	WO	N/A
WO 2014/093655	12/2013	WO	N/A
WO 2014/093661	12/2013	WO	N/A
WO 2014/093694	12/2013	WO	N/A
WO 2014/093701	12/2013	WO	N/A
WO 2014/093709	12/2013	WO	N/A
WO 2014/093712	12/2013	WO	N/A
WO 2014/093718	12/2013	WO	N/A
WO 2014/093736	12/2013	WO	N/A
WO 2014/093768	12/2013	WO	N/A
WO 2014/093852	12/2013	WO	N/A
WO 2014/096972	12/2013	WO	N/A
WO 2014/099744	12/2013	WO	N/A
WO 2014/099750	12/2013	WO	N/A
WO 2014/104878	12/2013	WO	N/A
WO 2014/110006	12/2013	WO	N/A
WO 2014/110552	12/2013	WO	N/A
WO 2014/113493	12/2013	WO	N/A
WO 2014/123967	12/2013	WO	N/A
WO 2014/124226	12/2013	WO	N/A
WO 2014/125668	12/2013	WO	N/A
WO 2014/127287	12/2013	WO	N/A
WO 2014/128324	12/2013	WO	N/A
WO 2014/128659	12/2013	WO	N/A
WO 2014/130706	12/2013	WO	N/A
WO 2014/130955	12/2013	WO	N/A
WO 2014/131833	12/2013	WO	N/A
WO 2014/138379	12/2013	WO	N/A
WO 2014/143381	12/2013	WO	N/A
WO 2014/144094	12/2013	WO	N/A
WO 2014/144155	12/2013	WO	N/A
WO 2014/144288	12/2013	WO	N/A
WO 2014/144592	12/2013	WO	N/A
WO 2014/144761	12/2013	WO	N/A
WO 2014/144951	12/2013	WO	N/A
WO 2014/145599	12/2013	WO	N/A
WO 2014/145736	12/2013	WO	N/A

WO 2014/150624	12/2013	WO	N/A
WO 2014/152432	12/2013	WO	N/A
WO 2014/152940	12/2013	WO	N/A
WO 2014/153118	12/2013	WO	N/A
WO 2014/153470	12/2013	WO	N/A
WO 2014/158593	12/2013	WO	N/A
WO 2014/161821	12/2013	WO	N/A
WO 2014/164466	12/2013	WO	N/A
WO 2014/165177	12/2013	WO	N/A
WO 2014/165349	12/2013	WO	N/A
WO 2014/165612	12/2013	WO	N/A
WO 2014/165707	12/2013	WO	N/A
WO 2014/165825	12/2013	WO	N/A
WO 2014/172458	12/2013	WO	N/A
WO 2014/172470	12/2013	WO	N/A
WO 2014/172489	12/2013	WO	N/A
WO 2014/173955	12/2013	WO	N/A
WO 2014/182700	12/2013	WO	N/A
WO 2014/183071	12/2013	WO	N/A
WO 2014/184143	12/2013	WO	N/A
WO 2014/184741	12/2013	WO	N/A
WO 2014/184744	12/2013	WO	N/A
WO 2014/186585	12/2013	WO	N/A
WO 2014/186686	12/2013	WO	N/A
WO 2014/190181	12/2013	WO	N/A
WO 2014/191128	12/2013	WO	N/A
WO 2014/191518	12/2013	WO	N/A
WO 2014/191521	12/2013	WO	N/A
WO 2014/191525	12/2013	WO	N/A
WO 2014/191527	12/2013	WO	N/A
WO 2014/193583	12/2013	WO	N/A
WO 2014/194190	12/2013	WO	N/A
WO 2014/197568	12/2013	WO	N/A
WO 2014/197748	12/2013	WO	N/A
WO 2014/199358	12/2013	WO	N/A
WO 2014/200659	12/2013	WO	N/A
WO 2014/201015	12/2013	WO	N/A
WO 2014/204578	12/2013	WO	N/A
WO 2014/204723	12/2013	WO	N/A
WO 2014/204724	12/2013	WO	N/A
WO 2014/204725	12/2013	WO	N/A
WO 2014/204726	12/2013	WO	N/A
WO 2014/204727	12/2013	WO	N/A
WO 2014/204728	12/2013	WO	N/A
WO 2014/204729	12/2013	WO	N/A
WO 2014/205192	12/2013	WO	N/A
WO 2014/207043	12/2013	WO	N/A
WO 2015/002780	12/2014	WO	N/A
WO 2015/004241	12/2014	WO	N/A
WO 2015/006290	12/2014	WO	N/A
WO 2015/006294	12/2014	WO	N/A
WO 2015/006437	12/2014	WO	N/A
WO 2015/006498	12/2014	WO	N/A
WO 2015/006747	12/2014	WO	N/A
WO 2015/007194	12/2014	WO	N/A
WO 2015/010114	12/2014	WO	N/A
WO 2015/011483	12/2014	WO	N/A
WO 2015/013583	12/2014	WO	N/A
WO 2015/017866	12/2014	WO	N/A
WO 2015/018503	12/2014	WO	N/A
WO 2015/021353	12/2014	WO	N/A

WO 2015/021426	12/2014	WO	N/A
WO 2015/021990	12/2014	WO	N/A
WO 2015/024017	12/2014	WO	N/A
WO 2015/024986	12/2014	WO	N/A
WO 2015/026883	12/2014	WO	N/A
WO 2015/026885	12/2014	WO	N/A
WO 2015/026886	12/2014	WO	N/A
WO 2015/026887	12/2014	WO	N/A
WO 2015/027134	12/2014	WO	N/A
WO 2015/028969	12/2014	WO	N/A
WO 2015/030881	12/2014	WO	N/A
WO 2015/031619	12/2014	WO	N/A
WO 2015/031775	12/2014	WO	N/A
WO 2015/032494	12/2014	WO	N/A
WO 2015/033293	12/2014	WO	N/A
WO 2015/034872	12/2014	WO	N/A
WO 2015/034885	12/2014	WO	N/A
WO 2015/035136	12/2014	WO	N/A
WO 2015/035139	12/2014	WO	N/A
WO 2015/035162	12/2014	WO	N/A
WO 2015/040075	12/2014	WO	N/A
WO 2015/040402	12/2014	WO	N/A
WO 2015/042393	12/2014	WO	N/A
WO 2015/042585	12/2014	WO	N/A
WO 2015/048577	12/2014	WO	N/A
WO 2015/048690	12/2014	WO	N/A
WO 2015/048707	12/2014	WO	N/A
WO 2015/048801	12/2014	WO	N/A
WO 2015/049897	12/2014	WO	N/A
WO 2015/051191	12/2014	WO	N/A
WO 2015/052133	12/2014	WO	N/A
WO 2015/052231	12/2014	WO	N/A
WO 2015/052335	12/2014	WO	N/A
WO 2015/053995	12/2014	WO	N/A
WO 2015/054253	12/2014	WO	N/A
WO 2015/054315	12/2014	WO	N/A
WO 2015/057671	12/2014	WO	N/A
WO 2015/057834	12/2014	WO	N/A
WO 2015/057852	12/2014	WO	N/A
WO 2015/057976	12/2014	WO	N/A
WO 2015/057980	12/2014	WO	N/A
WO 2015/059265	12/2014	WO	N/A
WO 2015/065964	12/2014	WO	N/A
WO 2015/066119	12/2014	WO	N/A
WO 2015/066634	12/2014	WO	N/A
WO 2015/066636	12/2014	WO	N/A
WO 2015/066637	12/2014	WO	N/A
WO 2015/066638	12/2014	WO	N/A
WO 2015/066643	12/2014	WO	N/A
WO 2015/069682	12/2014	WO	N/A
WO 2015/070083	12/2014	WO	N/A
WO 2015/070193	12/2014	WO	N/A
WO 2015/070212	12/2014	WO	N/A
WO 2015/071474	12/2014	WO	N/A
WO 2015/073683	12/2014	WO	N/A
WO 2015/073867	12/2014	WO	N/A
WO 2015/073990	12/2014	WO	N/A
WO 2015/075056	12/2014	WO	N/A
WO 2015/075154	12/2014	WO	N/A
WO 2015/075175	12/2014	WO	N/A
WO 2015/075195	12/2014	WO	N/A

WO 2015/075557	12/2014	WO	N/A
WO 2015/077058	12/2014	WO	N/A
WO 2015/077290	12/2014	WO	N/A
WO 2015/077318	12/2014	WO	N/A
WO 2015/079056	12/2014	WO	N/A
WO 2015/079057	12/2014	WO	N/A
WO 2015/086795	12/2014	WO	N/A
WO 2015/086798	12/2014	WO	N/A
WO 2015/088643	12/2014	WO	N/A
WO 2015/089046	12/2014	WO	N/A
WO 2015/089077	12/2014	WO	N/A
WO 2015/089277	12/2014	WO	N/A
WO 2015/089351	12/2014	WO	N/A
WO 2015/089354	12/2014	WO	N/A
WO 2015/089364	12/2014	WO	N/A
WO 2015/089419	12/2014	WO	N/A
WO 2015/089427	12/2014	WO	N/A
WO 2015/089462	12/2014	WO	N/A
WO 2015/089465	12/2014	WO	N/A
WO 2015/089473	12/2014	WO	N/A
WO 2015/089486	12/2014	WO	N/A
WO 2015/095804	12/2014	WO	N/A
WO-2015089406	12/2014	WO	A61K 38/465
WO 2015/099850	12/2014	WO	N/A
WO 2015/100929	12/2014	WO	N/A
WO 2015/103057	12/2014	WO	N/A
WO 2015/103153	12/2014	WO	N/A
WO 2015/105928	12/2014	WO	N/A
WO 2015/108993	12/2014	WO	N/A
WO 2015/109752	12/2014	WO	N/A
WO 2015/110474	12/2014	WO	N/A
WO 2015/112790	12/2014	WO	N/A
WO 2015/112896	12/2014	WO	N/A
WO 2015/113063	12/2014	WO	N/A
WO 2015/114365	12/2014	WO	N/A
WO 2015/115903	12/2014	WO	N/A
WO 2015/116686	12/2014	WO	N/A
WO 2015/116969	12/2014	WO	N/A
WO 2015/117021	12/2014	WO	N/A
WO 2015/117041	12/2014	WO	N/A
WO 2015/117081	12/2014	WO	N/A
WO 2015/118156	12/2014	WO	N/A
WO 2015/119941	12/2014	WO	N/A
WO 2015/121454	12/2014	WO	N/A
WO 2015/122967	12/2014	WO	N/A
WO 2015/123339	12/2014	WO	N/A
WO 2015/124715	12/2014	WO	N/A
WO 2015/124718	12/2014	WO	N/A
WO 2015/126927	12/2014	WO	N/A
WO 2015/127428	12/2014	WO	N/A
WO 2015/127439	12/2014	WO	N/A
WO 2015/129686	12/2014	WO	N/A
WO 2015/131101	12/2014	WO	N/A
WO 2015/133554	12/2014	WO	N/A
WO 2015/134121	12/2014	WO	N/A
WO 2015/134812	12/2014	WO	N/A
WO 2015/136001	12/2014	WO	N/A
WO 2015/138510	12/2014	WO	N/A
WO 2015/138739	12/2014	WO	N/A
WO 2015/138855	12/2014	WO	N/A
WO 2015/138870	12/2014	WO	N/A

WO 2015/139008	12/2014	WO	N/A
WO 2015/139139	12/2014	WO	N/A
WO 2015/143046	12/2014	WO	N/A
WO 2015/143177	12/2014	WO	N/A
WO 2015/145417	12/2014	WO	N/A
WO 2015/148431	12/2014	WO	N/A
WO 2015/148670	12/2014	WO	N/A
WO 2015/148680	12/2014	WO	N/A
WO 2015/148760	12/2014	WO	N/A
WO 2015/148761	12/2014	WO	N/A
WO 2015/148860	12/2014	WO	N/A
WO 2015/148863	12/2014	WO	N/A
WO 2015/153760	12/2014	WO	N/A
WO 2015/153780	12/2014	WO	N/A
WO 2015/153789	12/2014	WO	N/A
WO 2015/153791	12/2014	WO	N/A
WO 2015/153889	12/2014	WO	N/A
WO 2015/153940	12/2014	WO	N/A
WO 2015/155341	12/2014	WO	N/A
WO 2015/155686	12/2014	WO	N/A
WO 2015/157070	12/2014	WO	N/A
WO 2015/157534	12/2014	WO	N/A
WO 2015/159068	12/2014	WO	N/A
WO 2015/159086	12/2014	WO	N/A
WO 2015/159087	12/2014	WO	N/A
WO 2015/160683	12/2014	WO	N/A
WO 2015/161276	12/2014	WO	N/A
WO 2015/163733	12/2014	WO	N/A
WO 2015/164740	12/2014	WO	N/A
WO 2015/164748	12/2014	WO	N/A
WO 2015/165274	12/2014	WO	N/A
WO 2015/165275	12/2014	WO	N/A
WO 2015/165276	12/2014	WO	N/A
WO 2015/166272	12/2014	WO	N/A
WO 2015/167766	12/2014	WO	N/A
WO 2015/167956	12/2014	WO	N/A
WO 2015/168125	12/2014	WO	N/A
WO 2015/168158	12/2014	WO	N/A
WO 2015/168404	12/2014	WO	N/A
WO 2015/168547	12/2014	WO	N/A
WO 2015/168800	12/2014	WO	N/A
WO 2015/171603	12/2014	WO	N/A
WO 2015/171894	12/2014	WO	N/A
WO 2015/171932	12/2014	WO	N/A
WO 2015/172128	12/2014	WO	N/A
WO 2015/173436	12/2014	WO	N/A
WO 2015/175642	12/2014	WO	N/A
WO 2015/179540	12/2014	WO	N/A
WO 2015/183025	12/2014	WO	N/A
WO 2015/183026	12/2014	WO	N/A
WO 2015/183885	12/2014	WO	N/A
WO 2015/184259	12/2014	WO	N/A
WO 2015/184262	12/2014	WO	N/A
WO 2015/184268	12/2014	WO	N/A
WO 2015/188056	12/2014	WO	N/A
WO 2015/188065	12/2014	WO	N/A
WO 2015/188094	12/2014	WO	N/A
WO 2015/188109	12/2014	WO	N/A
WO 2015/188132	12/2014	WO	N/A
WO 2015/188135	12/2014	WO	N/A
WO 2015/188191	12/2014	WO	N/A

WO 2015/189693	12/2014	WO	N/A
WO 2015/191693	12/2014	WO	N/A
WO 2015/191899	12/2014	WO	N/A
WO 2015/191911	12/2014	WO	N/A
WO 2015/193858	12/2014	WO	N/A
WO 2015/195547	12/2014	WO	N/A
WO 2015/195621	12/2014	WO	N/A
WO 2015/195798	12/2014	WO	N/A
WO 2015/198020	12/2014	WO	N/A
WO 2015/200334	12/2014	WO	N/A
WO 2015/200378	12/2014	WO	N/A
WO 2015/200555	12/2014	WO	N/A
WO 2015/200805	12/2014	WO	N/A
WO 2016/001978	12/2015	WO	N/A
WO 2016/004010	12/2015	WO	N/A
WO 2016/004318	12/2015	WO	N/A
WO 2016/007347	12/2015	WO	N/A
WO 2016/007604	12/2015	WO	N/A
WO 2016/007948	12/2015	WO	N/A
WO 2016/011080	12/2015	WO	N/A
WO 2016/011210	12/2015	WO	N/A
WO 2016/011428	12/2015	WO	N/A
WO 2016/012544	12/2015	WO	N/A
WO 2016/012552	12/2015	WO	N/A
WO 2016/014409	12/2015	WO	N/A
WO 2016/014565	12/2015	WO	N/A
WO 2016/014794	12/2015	WO	N/A
WO 2016/014837	12/2015	WO	N/A
WO 2016/016119	12/2015	WO	N/A
WO 2016/016358	12/2015	WO	N/A
WO 2016/019144	12/2015	WO	N/A
WO 2016/020399	12/2015	WO	N/A
WO 2016/021972	12/2015	WO	N/A
WO 2016/021973	12/2015	WO	N/A
WO 2016/022363	12/2015	WO	N/A
WO 2016/022866	12/2015	WO	N/A
WO 2016/022931	12/2015	WO	N/A
WO 2016/025131	12/2015	WO	N/A
WO 2016/025469	12/2015	WO	N/A
WO 2016/025759	12/2015	WO	N/A
WO 2016/026444	12/2015	WO	N/A
WO 2016/028682	12/2015	WO	N/A
WO 2016/028843	12/2015	WO	N/A
WO 2016/028887	12/2015	WO	N/A
WO 2016/033088	12/2015	WO	N/A
WO 2016/033230	12/2015	WO	N/A
WO 2016/033246	12/2015	WO	N/A
WO 2016/033298	12/2015	WO	N/A
WO 2016/035044	12/2015	WO	N/A
WO 2016/035918	12/2015	WO	N/A
WO 2016/036754	12/2015	WO	N/A
WO 2016/037157	12/2015	WO	N/A
WO 2016/040030	12/2015	WO	N/A
WO 2016/040594	12/2015	WO	N/A
WO 2016/044182	12/2015	WO	N/A
WO 2016/044416	12/2015	WO	N/A
WO 2016/046635	12/2015	WO	N/A
WO 2016/049024	12/2015	WO	N/A
WO 2016/049163	12/2015	WO	N/A
WO 2016/049230	12/2015	WO	N/A
WO 2016/049251	12/2015	WO	N/A

WO 2016/049258	12/2015	WO	N/A
WO 2016/053397	12/2015	WO	N/A
WO 2016/054326	12/2015	WO	N/A
WO 2016/057061	12/2015	WO	N/A
WO 2016/057821	12/2015	WO	N/A
WO 2016/057835	12/2015	WO	N/A
WO 2016/057850	12/2015	WO	N/A
WO 2016/057951	12/2015	WO	N/A
WO 2016/057961	12/2015	WO	N/A
WO 2016/061073	12/2015	WO	N/A
WO 2016/061374	12/2015	WO	N/A
WO 2016/061481	12/2015	WO	N/A
WO 2016/061523	12/2015	WO	N/A
WO 2016/064894	12/2015	WO	N/A
WO 2016/065364	12/2015	WO	N/A
WO 2016/069282	12/2015	WO	N/A
WO 2016/069283	12/2015	WO	N/A
WO 2016/069591	12/2015	WO	N/A
WO 2016/069774	12/2015	WO	N/A
WO 2016/069910	12/2015	WO	N/A
WO 2016/069912	12/2015	WO	N/A
WO 2016/070037	12/2015	WO	N/A
WO 2016/070070	12/2015	WO	N/A
WO 2016/070129	12/2015	WO	N/A
WO 2016/072399	12/2015	WO	N/A
WO 2016/072936	12/2015	WO	N/A
WO 2016/073433	12/2015	WO	N/A
WO 2016/073559	12/2015	WO	N/A
WO 2016/073990	12/2015	WO	N/A
WO 2016/075662	12/2015	WO	N/A
WO 2016/076672	12/2015	WO	N/A
WO 2016/077273	12/2015	WO	N/A
WO 2016/077350	12/2015	WO	N/A
WO 2016/080097	12/2015	WO	N/A
WO 2016/080795	12/2015	WO	N/A
WO 2016/081923	12/2015	WO	N/A
WO 2016/081924	12/2015	WO	N/A
WO 2016/082135	12/2015	WO	N/A
WO 2016/083811	12/2015	WO	N/A
WO 2016/084084	12/2015	WO	N/A
WO 2016/084088	12/2015	WO	N/A
WO 2016/086177	12/2015	WO	N/A
WO 2016/089433	12/2015	WO	N/A
WO 2016/089866	12/2015	WO	N/A
WO 2016/089883	12/2015	WO	N/A
WO 2016/090385	12/2015	WO	N/A
WO 2016/094679	12/2015	WO	N/A
WO 2016/094845	12/2015	WO	N/A
WO 2016/094867	12/2015	WO	N/A
WO 2016/094872	12/2015	WO	N/A
WO 2016/094874	12/2015	WO	N/A
WO 2016/094880	12/2015	WO	N/A
WO 2016/094888	12/2015	WO	N/A
WO 2016/097212	12/2015	WO	N/A
WO 2016/097231	12/2015	WO	N/A
WO 2016/097751	12/2015	WO	N/A
WO 2016/099887	12/2015	WO	N/A
WO 2016/100272	12/2015	WO	N/A
WO 2016/100389	12/2015	WO	N/A
WO 2016/100568	12/2015	WO	N/A
WO 2016/100571	12/2015	WO	N/A

WO 2016/100951	12/2015	WO	N/A
WO 2016/100955	12/2015	WO	N/A
WO 2016/100974	12/2015	WO	N/A
WO 2016/103233	12/2015	WO	N/A
WO 2016/104716	12/2015	WO	N/A
WO 2016/106236	12/2015	WO	N/A
WO 2016/106239	12/2015	WO	N/A
WO 2016/106244	12/2015	WO	N/A
WO 2016/106338	12/2015	WO	N/A
WO 2016/108926	12/2015	WO	N/A
WO 2016/109255	12/2015	WO	N/A
WO 2016/109840	12/2015	WO	N/A
WO 2016/110214	12/2015	WO	N/A
WO 2016/110453	12/2015	WO	N/A
WO 2016/110511	12/2015	WO	N/A
WO 2016/110512	12/2015	WO	N/A
WO 2016/111546	12/2015	WO	N/A
WO 2016/112242	12/2015	WO	N/A
WO 2016/112351	12/2015	WO	N/A
WO 2016/112963	12/2015	WO	N/A
WO 2016/113357	12/2015	WO	N/A
WO 2016/114972	12/2015	WO	N/A
WO 2016/115179	12/2015	WO	N/A
WO 2016/115326	12/2015	WO	N/A
WO 2016/115355	12/2015	WO	N/A
WO 2016/116032	12/2015	WO	N/A
WO 2016/120480	12/2015	WO	N/A
WO 2016/123071	12/2015	WO	N/A
WO 2016/123230	12/2015	WO	N/A
WO 2016/123243	12/2015	WO	N/A
WO 2016/123578	12/2015	WO	N/A
WO 2016/126747	12/2015	WO	N/A
WO 2016/130600	12/2015	WO	N/A
WO 2016/130697	12/2015	WO	N/A
WO 2016/131009	12/2015	WO	N/A
WO 2016/132122	12/2015	WO	N/A
WO 2016/133165	12/2015	WO	N/A
WO 2016/135507	12/2015	WO	N/A
WO 2016/135557	12/2015	WO	N/A
WO 2016/135558	12/2015	WO	N/A
WO 2016/135559	12/2015	WO	N/A
WO 2016/137774	12/2015	WO	N/A
WO 2016/137949	12/2015	WO	N/A
WO 2016/141224	12/2015	WO	N/A
WO 2016/141893	12/2015	WO	N/A
WO 2016/142719	12/2015	WO	N/A
WO 2016/145150	12/2015	WO	N/A
WO 2016/148994	12/2015	WO	N/A
WO 2016/149484	12/2015	WO	N/A
WO 2016/149547	12/2015	WO	N/A
WO 2016/150336	12/2015	WO	N/A
WO 2016/150855	12/2015	WO	N/A
WO 2016/154016	12/2015	WO	N/A
WO 2016/154579	12/2015	WO	N/A
WO 2016/154596	12/2015	WO	N/A
WO 2016/155482	12/2015	WO	N/A
WO 2016/161004	12/2015	WO	N/A
WO 2016/161207	12/2015	WO	N/A
WO 2016/161260	12/2015	WO	N/A
WO 2016/161380	12/2015	WO	N/A
WO 2016/161446	12/2015	WO	N/A



WO 2016/164305	12/2015	WO	N/A
WO 2016/164356	12/2015	WO	N/A
WO 2016/164797	12/2015	WO	N/A
WO 2016/166340	12/2015	WO	N/A
WO 2016/167300	12/2015	WO	N/A
WO 2016/168631	12/2015	WO	N/A
WO 2016/170484	12/2015	WO	N/A
WO 2016/172359	12/2015	WO	N/A
WO 2016/172727	12/2015	WO	N/A
WO 2016/174056	12/2015	WO	N/A
WO 2016/174151	12/2015	WO	N/A
WO 2016/174250	12/2015	WO	N/A
WO 2016/176191	12/2015	WO	N/A
WO 2016/176404	12/2015	WO	N/A
WO 2016/176690	12/2015	WO	N/A
WO 2016/177682	12/2015	WO	N/A
WO 2016/178207	12/2015	WO	N/A
WO 2016/179038	12/2015	WO	N/A
WO 2016/179112	12/2015	WO	N/A
WO 2016/181357	12/2015	WO	N/A
WO 2016/182893	12/2015	WO	N/A
WO 2016/182917	12/2015	WO	N/A
WO 2016/182959	12/2015	WO	N/A
WO 2016/183236	12/2015	WO	N/A
WO 2016/183298	12/2015	WO	N/A
WO 2016/183345	12/2015	WO	N/A
WO 2016/183402	12/2015	WO	N/A
WO 2016/183438	12/2015	WO	N/A
WO 2016/183448	12/2015	WO	N/A
WO 2016/184955	12/2015	WO	N/A
WO 2016/184989	12/2015	WO	N/A
WO 2016/185411	12/2015	WO	N/A
WO 2016/186745	12/2015	WO	N/A
WO 2016/186772	12/2015	WO	N/A
WO 2016/186946	12/2015	WO	N/A
WO 2016/186953	12/2015	WO	N/A
WO 2016/187717	12/2015	WO	N/A
WO 2016/187904	12/2015	WO	N/A
WO 2016/191684	12/2015	WO	N/A
WO 2016/191869	12/2015	WO	N/A
WO 2016/196273	12/2015	WO	N/A
WO 2016/196282	12/2015	WO	N/A
WO 2016/196308	12/2015	WO	N/A
WO 2016/196361	12/2015	WO	N/A
WO 2016/196499	12/2015	WO	N/A
WO 2016/196539	12/2015	WO	N/A
WO 2016/196655	12/2015	WO	N/A
WO 2016/196805	12/2015	WO	N/A
WO 2016/196887	12/2015	WO	N/A
WO 2016/197132	12/2015	WO	N/A
WO 2016/197133	12/2015	WO	N/A
WO 2016/197354	12/2015	WO	N/A
WO 2016/197355	12/2015	WO	N/A
WO 2016/197356	12/2015	WO	N/A
WO 2016/197357	12/2015	WO	N/A
WO 2016/197358	12/2015	WO	N/A
WO 2016/197359	12/2015	WO	N/A
WO 2016/197360	12/2015	WO	N/A
WO 2016/197361	12/2015	WO	N/A
WO 2016/197362	12/2015	WO	N/A
WO 2016/198361	12/2015	WO	N/A

WO 2016/198500	12/2015	WO	N/A
WO 2016/200263	12/2015	WO	N/A
WO 2016/201047	12/2015	WO	N/A
WO 2016/201138	12/2015	WO	N/A
WO 2016/201152	12/2015	WO	N/A
WO 2016/201153	12/2015	WO	N/A
WO 2016/201155	12/2015	WO	N/A
WO 2016/205276	12/2015	WO	N/A
WO 2016/205613	12/2015	WO	N/A
WO 2016/205623	12/2015	WO	N/A
WO 2016/205680	12/2015	WO	N/A
WO 2016/205688	12/2015	WO	N/A
WO 2016/205703	12/2015	WO	N/A
WO 2016/205711	12/2015	WO	N/A
WO 2016/205728	12/2015	WO	N/A
WO 2016/205745	12/2015	WO	N/A
WO 2016/205749	12/2015	WO	N/A
WO 2016/205759	12/2015	WO	N/A
WO 2016/205764	12/2015	WO	N/A
WO 2017/001572	12/2016	WO	N/A
WO 2017/001988	12/2016	WO	N/A
WO 2017/004261	12/2016	WO	N/A
WO 2017/004279	12/2016	WO	N/A
WO 2017/004616	12/2016	WO	N/A
WO 2017/005807	12/2016	WO	N/A
WO 2017/009399	12/2016	WO	N/A
WO 2017/010556	12/2016	WO	N/A
WO 2017/011519	12/2016	WO	N/A
WO 2017/011721	12/2016	WO	N/A
WO 2017/011804	12/2016	WO	N/A
WO 2017/015015	12/2016	WO	N/A
WO 2017/015101	12/2016	WO	N/A
WO 2017/015545	12/2016	WO	N/A
WO 2017/015567	12/2016	WO	N/A
WO 2017/015637	12/2016	WO	N/A
WO 2017/017016	12/2016	WO	N/A
WO 2017/019867	12/2016	WO	N/A
WO 2017/019895	12/2016	WO	N/A
WO 2017/023803	12/2016	WO	N/A
WO 2017/023974	12/2016	WO	N/A
WO 2017/024047	12/2016	WO	N/A
WO 2017/024319	12/2016	WO	N/A
WO 2017/024343	12/2016	WO	N/A
WO 2017/024602	12/2016	WO	N/A
WO 2017/025323	12/2016	WO	N/A
WO 2017/027423	12/2016	WO	N/A
WO 2017/028768	12/2016	WO	N/A
WO 2017/029664	12/2016	WO	N/A
WO 2017/031360	12/2016	WO	N/A
WO 2017/031483	12/2016	WO	N/A
WO 2017/035416	12/2016	WO	N/A
WO 2017/040348	12/2016	WO	N/A
WO 2017/040511	12/2016	WO	N/A
WO 2017/040709	12/2016	WO	N/A
WO 2017/040786	12/2016	WO	N/A
WO 2017/040793	12/2016	WO	N/A
WO 2017/040813	12/2016	WO	N/A
WO 2017/043573	12/2016	WO	N/A
WO 2017/043656	12/2016	WO	N/A
WO 2017/044419	12/2016	WO	N/A
WO 2017/044776	12/2016	WO	N/A

WO 2017/044857	12/2016	WO	N/A
WO 2017/048390	12/2016	WO	N/A
WO 2017/049129	12/2016	WO	N/A
WO 2017/050963	12/2016	WO	N/A
WO 2017/053312	12/2016	WO	N/A
WO 2017/053431	12/2016	WO	N/A
WO 2017/053713	12/2016	WO	N/A
WO 2017/053729	12/2016	WO	N/A
WO 2017/053753	12/2016	WO	N/A
WO 2017/053762	12/2016	WO	N/A
WO 2017/053879	12/2016	WO	N/A
WO 2017/054721	12/2016	WO	N/A
WO 2017/058658	12/2016	WO	N/A
WO 2017/059241	12/2016	WO	N/A
WO 2017/062605	12/2016	WO	N/A
WO 2017/062723	12/2016	WO	N/A
WO 2017/062754	12/2016	WO	N/A
WO 2017/062855	12/2016	WO	N/A
WO 2017/062886	12/2016	WO	N/A
WO 2017/062983	12/2016	WO	N/A
WO 2017/064439	12/2016	WO	N/A
WO 2017/064546	12/2016	WO	N/A
WO 2017/064566	12/2016	WO	N/A
WO 2017/066175	12/2016	WO	N/A
WO 2017/066497	12/2016	WO	N/A
WO 2017/066588	12/2016	WO	N/A
WO 2017/066707	12/2016	WO	N/A
WO 2017/066781	12/2016	WO	N/A
WO 2017/068077	12/2016	WO	N/A
WO 2017/068377	12/2016	WO	N/A
WO 2017/069829	12/2016	WO	N/A
WO 2017/070029	12/2016	WO	N/A
WO 2017/070032	12/2016	WO	N/A
WO 2017/070169	12/2016	WO	N/A
WO 2017/070284	12/2016	WO	N/A
WO 2017/070598	12/2016	WO	N/A
WO 2017/070605	12/2016	WO	N/A
WO 2017/070632	12/2016	WO	N/A
WO 2017/070633	12/2016	WO	N/A
WO 2017/072590	12/2016	WO	N/A
WO 2017/074526	12/2016	WO	N/A
WO 2017/074962	12/2016	WO	N/A
WO 2017/075261	12/2016	WO	N/A
WO 2017/075335	12/2016	WO	N/A
WO 2017/075475	12/2016	WO	N/A
WO 2017/077135	12/2016	WO	N/A
WO 2017/077329	12/2016	WO	N/A
WO 2017/078751	12/2016	WO	N/A
WO 2017/079400	12/2016	WO	N/A
WO 2017/079428	12/2016	WO	N/A
WO 2017/079673	12/2016	WO	N/A
WO 2017/079724	12/2016	WO	N/A
WO 2017/081097	12/2016	WO	N/A
WO 2017/081288	12/2016	WO	N/A
WO 2017/083368	12/2016	WO	N/A
WO 2017/083722	12/2016	WO	N/A
WO 2017/083766	12/2016	WO	N/A
WO 2017/087395	12/2016	WO	N/A
WO 2017/090724	12/2016	WO	N/A
WO 2017/091510	12/2016	WO	N/A
WO 2017/091630	12/2016	WO	N/A

WO 2017/092201	12/2016	WO	N/A
WO 2017/093370	12/2016	WO	N/A
WO 2017/093969	12/2016	WO	N/A
WO 2017/095111	12/2016	WO	N/A
WO 2017/096041	12/2016	WO	N/A
WO 2017/096237	12/2016	WO	N/A
WO 2017/100158	12/2016	WO	N/A
WO 2017/100431	12/2016	WO	N/A
WO 2017/104404	12/2016	WO	N/A
WO 2017/105251	12/2016	WO	N/A
WO 2017/105350	12/2016	WO	N/A
WO 2017/105991	12/2016	WO	N/A
WO 2017/106414	12/2016	WO	N/A
WO 2017/106528	12/2016	WO	N/A
WO 2017/106537	12/2016	WO	N/A
WO 2017/106569	12/2016	WO	N/A
WO 2017/106616	12/2016	WO	N/A
WO 2017/106657	12/2016	WO	N/A
WO 2017/106767	12/2016	WO	N/A
WO 2017/109134	12/2016	WO	N/A
WO 2017/109757	12/2016	WO	N/A
WO 2017/112620	12/2016	WO	N/A
WO 2017/115268	12/2016	WO	N/A
WO 2017/117395	12/2016	WO	N/A
WO 2017/118598	12/2016	WO	N/A
WO 2017/118720	12/2016	WO	N/A
WO 2017/123609	12/2016	WO	N/A
WO 2017/123910	12/2016	WO	N/A
WO 2017/124086	12/2016	WO	N/A
WO 2017/124100	12/2016	WO	N/A
WO 2017/124652	12/2016	WO	N/A
WO 2017/126987	12/2016	WO	N/A
WO 2017/127807	12/2016	WO	N/A
WO 2017/131237	12/2016	WO	N/A
WO 2017/132112	12/2016	WO	N/A
WO 2017/132580	12/2016	WO	N/A
WO 2017/136520	12/2016	WO	N/A
WO 2017/136629	12/2016	WO	N/A
WO 2017/136794	12/2016	WO	N/A
WO 2017/139264	12/2016	WO	N/A
WO 2017/139505	12/2016	WO	N/A
WO 2017/141173	12/2016	WO	N/A
WO 2017/142835	12/2016	WO	N/A
WO 2017/142923	12/2016	WO	N/A
WO 2017/142999	12/2016	WO	N/A
WO 2017/143042	12/2016	WO	N/A
WO 2017/147056	12/2016	WO	N/A
WO 2017/147278	12/2016	WO	N/A
WO 2017/147432	12/2016	WO	N/A
WO 2017/147446	12/2016	WO	N/A
WO 2017/147555	12/2016	WO	N/A
WO 2017/151444	12/2016	WO	N/A
WO 2017/151719	12/2016	WO	N/A
WO 2017/152015	12/2016	WO	N/A
WO 2017/155717	12/2016	WO	N/A
WO 2017/157422	12/2016	WO	N/A
WO 2017/158153	12/2016	WO	N/A
WO 2017/160689	12/2016	WO	N/A
WO 2017/160752	12/2016	WO	N/A
WO 2017/160890	12/2016	WO	N/A
WO 2017/161068	12/2016	WO	N/A

WO 2017/165741	12/2016	WO	N/A
WO 2017/165826	12/2016	WO	N/A
WO 2017/165862	12/2016	WO	N/A
WO 2017/167712	12/2016	WO	N/A
WO 2017/172644	12/2016	WO	N/A
WO 2017/172645	12/2016	WO	N/A
WO 2017/172860	12/2016	WO	N/A
WO 2017/173004	12/2016	WO	N/A
WO 2017/173054	12/2016	WO	N/A
WO 2017/173092	12/2016	WO	N/A
WO 2017/174329	12/2016	WO	N/A
WO 2017/176529	12/2016	WO	N/A
WO 2017/176806	12/2016	WO	N/A
WO 2017/178590	12/2016	WO	N/A
WO 2017/180694	12/2016	WO	N/A
WO 2017/180711	12/2016	WO	N/A
WO 2017/180915	12/2016	WO	N/A
WO 2017/180926	12/2016	WO	N/A
WO 2017/181107	12/2016	WO	N/A
WO 2017/181735	12/2016	WO	N/A
WO 2017/182468	12/2016	WO	N/A
WO 2017/182585	12/2016	WO	N/A
WO 2017/182607	12/2016	WO	N/A
WO 2017/184334	12/2016	WO	N/A
WO 2017/184768	12/2016	WO	N/A
WO 2017/184786	12/2016	WO	N/A
WO 2017/186550	12/2016	WO	N/A
WO 2017/189308	12/2016	WO	N/A
WO 2017/189336	12/2016	WO	N/A
WO 2017/190041	12/2016	WO	N/A
WO 2017/190257	12/2016	WO	N/A
WO 2017/190664	12/2016	WO	N/A
WO 2017/191210	12/2016	WO	N/A
WO 2017/191274	12/2016	WO	N/A
WO 2017/192172	12/2016	WO	N/A
WO 2017/192512	12/2016	WO	N/A
WO 2017/192544	12/2016	WO	N/A
WO 2017/192573	12/2016	WO	N/A
WO 2017/193029	12/2016	WO	N/A
WO 2017/193053	12/2016	WO	N/A
WO 2017/196768	12/2016	WO	N/A
WO 2017/197038	12/2016	WO	N/A
WO 2017/197238	12/2016	WO	N/A
WO 2017/197301	12/2016	WO	N/A
WO 2017/201476	12/2016	WO	N/A
WO 2017/205290	12/2016	WO	N/A
WO 2017/205423	12/2016	WO	N/A
WO 2017/207589	12/2016	WO	N/A
WO 2017/208247	12/2016	WO	N/A
WO 2017/209809	12/2016	WO	N/A
WO 2017/213896	12/2016	WO	N/A
WO 2017/213898	12/2016	WO	N/A
WO 2017/214460	12/2016	WO	N/A
WO 2017/216392	12/2016	WO	N/A
WO 2017/216771	12/2016	WO	N/A
WO 2017/218185	12/2016	WO	N/A
WO 2017/219027	12/2016	WO	N/A
WO 2017/219033	12/2016	WO	N/A
WO 2017/220751	12/2016	WO	N/A
WO 2017/222370	12/2016	WO	N/A
WO 2017/222773	12/2016	WO	N/A

WO 2017/222834	12/2016	WO	N/A
WO 2017/223107	12/2016	WO	N/A
WO 2017/223330	12/2016	WO	N/A
WO 2018/000657	12/2017	WO	N/A
WO 2018/002719	12/2017	WO	N/A
WO 2018/005117	12/2017	WO	N/A
WO 2018/005289	12/2017	WO	N/A
WO 2018/005691	12/2017	WO	N/A
WO 2018/005782	12/2017	WO	N/A
WO 2018/005873	12/2017	WO	N/A
WO 2018/06693	12/2017	WO	N/A
WO 2018/009520	12/2017	WO	N/A
WO 2018/009562	12/2017	WO	N/A
WO 2018/009822	12/2017	WO	N/A
WO 2018/013821	12/2017	WO	N/A
WO 2018/013932	12/2017	WO	N/A
WO 2018/013990	12/2017	WO	N/A
WO 2018/014384	12/2017	WO	N/A
WO 2018/015444	12/2017	WO	N/A
WO 2018/015936	12/2017	WO	N/A
WO 2018/017754	12/2017	WO	N/A
WO 2018/018979	12/2017	WO	N/A
WO 2018/020248	12/2017	WO	N/A
WO 2018/021878	12/2017	WO	N/A
WO 2018/022480	12/2017	WO	N/A
WO 2018/022634	12/2017	WO	N/A
WO 2018/025206	12/2017	WO	N/A
WO 2018/026723	12/2017	WO	N/A
WO 2018/026976	12/2017	WO	N/A
WO 2018/027078	12/2017	WO	N/A
WO 2018/030608	12/2017	WO	N/A
WO 2018/031683	12/2017	WO	N/A
WO 2018/035250	12/2017	WO	N/A
WO 2018/035300	12/2017	WO	N/A
WO 2018/035423	12/2017	WO	N/A
WO 2018/035503	12/2017	WO	N/A
WO 2018/039145	12/2017	WO	N/A
WO 2018/039438	12/2017	WO	N/A
WO 2018/039440	12/2017	WO	N/A
WO 2018/039448	12/2017	WO	N/A
WO 2018/045630	12/2017	WO	N/A
WO 2018/048827	12/2017	WO	N/A
WO 2018/049073	12/2017	WO	N/A
WO 2018/049168	12/2017	WO	N/A
WO 2018/051347	12/2017	WO	N/A
WO 2018/058064	12/2017	WO	N/A
WO 2018/062866	12/2017	WO	N/A
WO 2018/064352	12/2017	WO	N/A
WO 2018/064371	12/2017	WO	N/A
WO 2018/064516	12/2017	WO	N/A
WO 2018/067546	12/2017	WO	N/A
WO 2018/067846	12/2017	WO	N/A
WO 2018/068053	12/2017	WO	N/A
WO 2018/069474	12/2017	WO	N/A
WO 2018/071623	12/2017	WO	N/A
WO 2018/071663	12/2017	WO	N/A
WO 2018/071868	12/2017	WO	N/A
WO 2018/071892	12/2017	WO	N/A
WO 2018/074979	12/2017	WO	N/A
WO 2018/079134	12/2017	WO	N/A
WO 2018/080573	12/2017	WO	N/A

WO 2018/081504	12/2017	WO	N/A
WO 2018/081535	12/2017	WO	N/A
WO 2018/081728	12/2017	WO	N/A
WO 2018/083128	12/2017	WO	N/A
WO 2018/083606	12/2017	WO	N/A
WO 2018/085288	12/2017	WO	N/A
WO 2018/085414	12/2017	WO	N/A
WO 2018/085842	12/2017	WO	N/A
WO 2018/086623	12/2017	WO	N/A
WO 2018/089664	12/2017	WO	N/A
WO 2018/093990	12/2017	WO	N/A
WO 2018/098383	12/2017	WO	N/A
WO 2018/098480	12/2017	WO	N/A
WO 2018/098587	12/2017	WO	N/A
WO 2018/099256	12/2017	WO	N/A
WO 2018/103686	12/2017	WO	N/A
WO 2018/106268	12/2017	WO	N/A
WO 2018/107028	12/2017	WO	N/A
WO 2018/107103	12/2017	WO	N/A
WO 2018/107129	12/2017	WO	N/A
WO 2018/108272	12/2017	WO	N/A
WO 2018/109101	12/2017	WO	N/A
WO 2018/111946	12/2017	WO	N/A
WO 2018/111947	12/2017	WO	N/A
WO 2018/112336	12/2017	WO	N/A
WO 2018/112446	12/2017	WO	N/A
WO 2018/119354	12/2017	WO	N/A
WO 2018/119359	12/2017	WO	N/A
WO 2018/120283	12/2017	WO	N/A
WO 2018/130830	12/2017	WO	N/A
WO 2018/135838	12/2017	WO	N/A
WO 2018/136396	12/2017	WO	N/A
WO 2018/138385	12/2017	WO	N/A
WO 2018/142364	12/2017	WO	N/A
WO 2018/148246	12/2017	WO	N/A
WO 2018/148256	12/2017	WO	N/A
WO 2018/148647	12/2017	WO	N/A
WO 2018/149418	12/2017	WO	N/A
WO 2018/149888	12/2017	WO	N/A
WO 2018/149915	12/2017	WO	N/A
WO 2018/152197	12/2017	WO	N/A
WO 2018/152418	12/2017	WO	N/A
WO 2018/154380	12/2017	WO	N/A
WO 2018/154387	12/2017	WO	N/A
WO 2018/154412	12/2017	WO	N/A
WO 2018/154413	12/2017	WO	N/A
WO 2018/154418	12/2017	WO	N/A
WO 2018/154439	12/2017	WO	N/A
WO 2018/154459	12/2017	WO	N/A
WO 2018/154462	12/2017	WO	N/A
WO 2018/156372	12/2017	WO	N/A
WO 2018/156824	12/2017	WO	N/A
WO 2018/161009	12/2017	WO	N/A
WO 2018/161032	12/2017	WO	N/A
WO 2018/165504	12/2017	WO	N/A
WO 2018/165629	12/2017	WO	N/A
WO 2018/170015	12/2017	WO	N/A
WO 2018/170340	12/2017	WO	N/A
WO 2018/175502	12/2017	WO	N/A
WO 2018/176009	12/2017	WO	N/A
WO 2018/177351	12/2017	WO	N/A

WO 2018/179578	12/2017	WO	N/A
WO 2018/183403	12/2017	WO	N/A
WO 2018/189184	12/2017	WO	N/A
WO 2018/191388	12/2017	WO	N/A
WO 2018/195402	12/2017	WO	N/A
WO 2018/195545	12/2017	WO	N/A
WO 2018/195555	12/2017	WO	N/A
WO 2018/197020	12/2017	WO	N/A
WO 2018/197495	12/2017	WO	N/A
WO 2018/200597	12/2017	WO	N/A
WO 2018/202800	12/2017	WO	N/A
WO 2018/204493	12/2017	WO	N/A
WO 2018/208755	12/2017	WO	N/A
WO 2018/208998	12/2017	WO	N/A
WO 2018/209158	12/2017	WO	N/A
WO 2018/209320	12/2017	WO	N/A
WO 2018/213351	12/2017	WO	N/A
WO 2018/213708	12/2017	WO	N/A
WO 2018/213726	12/2017	WO	N/A
WO 2018/213771	12/2017	WO	N/A
WO 2018/213791	12/2017	WO	N/A
WO 2018/217852	12/2017	WO	N/A
WO 2018/217981	12/2017	WO	N/A
WO 2018/218166	12/2017	WO	N/A
WO 2018/218188	12/2017	WO	N/A
WO 2018/218206	12/2017	WO	N/A
WO 2018/226855	12/2017	WO	N/A
WO 2019/005884	12/2018	WO	N/A
WO 2019/005886	12/2018	WO	N/A
WO 2019/010384	12/2018	WO	N/A
WO 2019/023680	12/2018	WO	N/A
WO 2019/042284	12/2018	WO	N/A
WO 2019/051097	12/2018	WO	N/A
WO 2019/067992	12/2018	WO	N/A
WO 2019/075357	12/2018	WO	N/A
WO 2019/079347	12/2018	WO	N/A
WO 2019/084062	12/2018	WO	N/A
WO 2019/090169	12/2018	WO	N/A
WO 2019/090367	12/2018	WO	N/A
WO 2019/092042	12/2018	WO	N/A
WO 2019/118497	12/2018	WO	N/A
WO 2019/118935	12/2018	WO	N/A
WO 2019/118949	12/2018	WO	N/A
WO 2019/123430	12/2018	WO	N/A
WO 2019/126709	12/2018	WO	N/A
WO 2019/139645	12/2018	WO	N/A
WO 2019/139951	12/2018	WO	N/A
WO 2019/147014	12/2018	WO	N/A
WO 2019/161251	12/2018	WO	N/A
WO 2019/168953	12/2018	WO	N/A
WO 2019/183641	12/2018	WO	N/A
WO 2019/204369	12/2018	WO	N/A
WO 2019/213257	12/2018	WO	N/A
WO 2019/217942	12/2018	WO	N/A
WO 2019/226593	12/2018	WO	N/A
WO 2019/226953	12/2018	WO	N/A
WO 2019/236566	12/2018	WO	N/A
WO 2019/241649	12/2018	WO	N/A
WO 2020/014261	12/2019	WO	N/A
WO 2020/028555	12/2019	WO	N/A
WO 2020/028823	12/2019	WO	N/A



WO 2020/041751	12/2019	WO	N/A
WO 2020/047124	12/2019	WO	N/A
WO 2020/051360	12/2019	WO	N/A
WO 2020/086908	12/2019	WO	N/A
WO 2020/092453	12/2019	WO	N/A
WO 2020/102659	12/2019	WO	N/A
WO 2020/154500	12/2019	WO	N/A
WO 2020/180975	12/2019	WO	N/A
WO 2020/181178	12/2019	WO	N/A
WO 2020/181180	12/2019	WO	N/A
WO 2020/181193	12/2019	WO	N/A
WO 2020/181195	12/2019	WO	N/A
WO 2020/181202	12/2019	WO	N/A
WO 2020/191153	12/2019	WO	N/A
WO 2020/191171	12/2019	WO	N/A
WO 2020/191233	12/2019	WO	N/A
WO 2020/191234	12/2019	WO	N/A
WO 2020/191239	12/2019	WO	N/A
WO 2020/191241	12/2019	WO	N/A
WO 2020/191242	12/2019	WO	N/A
WO 2020/191243	12/2019	WO	N/A
WO 2020/191245	12/2019	WO	N/A
WO 2020/191246	12/2019	WO	N/A
WO 2020/191248	12/2019	WO	N/A
WO 2020/191249	12/2019	WO	N/A
WO 2020/210751	12/2019	WO	N/A
WO 2020/214842	12/2019	WO	N/A
WO 2020/236982	12/2019	WO	N/A
WO 2021/025750	12/2020	WO	N/A
WO 2021/030666	12/2020	WO	N/A
WO 2021/072328	12/2020	WO	N/A
WO 2021/108717	12/2020	WO	N/A
WO 2021/138469	12/2020	WO	N/A
WO 2021/155065	12/2020	WO	N/A
WO 2021/158921	12/2020	WO	N/A
WO 2021/158995	12/2020	WO	N/A
WO 2021/158999	12/2020	WO	N/A
WO 2021/222318	12/2020	WO	N/A
WO 2021/226558	12/2020	WO	N/A
WO 2021/252924	12/2020	WO	N/A
WO 2022/067130	12/2021	WO	N/A
WO 2022/150790	12/2021	WO	N/A
WO 2022/165262	12/2021	WO	N/A
WO 2023/015309	12/2022	WO	N/A
WO 2023/102537	12/2022	WO	N/A
WO 2023/102538	12/2022	WO	N/A
WO 2023/102550	12/2022	WO	N/A
WO 2023/173140	12/2022	WO	N/A
WO 2024/155741	12/2023	WO	N/A
WO 2024/155745	12/2023	WO	N/A
WO 2024/215652	12/2023	WO	N/A

## OTHER PUBLICATIONS

Mahoney et al., “The Next Immune-Checkpoint Inhibitors: PD-1/PD-L1 Blockade in Melanoma” 37(4) Clinical Therapeutics 764-782 (Year: 2015). cited by examiner

Bae et al., “Heteroclitic CD33 Peptide With Enhanced Anti-Acute Myeloid Leukemic Immunogenicity” 10 Clinical Cancer Research 7043-7052 (Year: 2004). cited by examiner

Fikes et al., “Design of multi-epitope, analogue-based cancer vaccines” 3(6) Expert Opinion On Biological Therapy 985-993 (Year: 2003). cited by examiner

Houghton et al., “Immunological validation of the EpiOptimizer program for streamlined design of heteroclitic epitopes” 25 Vaccine 5330-5342 (Year: 2007). cited by examiner

U.S. Appl. No. 61/716,256, filed Oct. 19, 2012, Jinek et al. cited by applicant

U.S. Appl. No. 61/717,324, filed Oct. 23, 2012, Cho et al. cited by applicant

U.S. Appl. No. 61/734,256, filed Dec. 6, 2012, Chen et al. cited by applicant

U.S. Appl. No. 61/758,624, filed Jan. 30, 2013, Chen et al. cited by applicant

U.S. Appl. No. 61/761,046, filed Feb. 5, 2013, Knight et al. cited by applicant

U.S. Appl. No. 61/794,422, filed Mar. 15, 2013, Knight et al. cited by applicant

U.S. Appl. No. 61/803,599, filed Mar. 20, 2013, Kim et al. cited by applicant

U.S. Appl. No. 61/837,481, filed Jun. 20, 2013, Cho et al. cited by applicant

U.S. Appl. No. 61/838,178, filed Jun. 21, 2013, Joung et al. cited by applicant

U.S. Appl. No. 61/874,682, filed Sep. 6, 2013, Liu et al. cited by applicant

U.S. Appl. No. 61/874,746, filed Sep. 6, 2013, Liu et al. cited by applicant

U.S. Appl. No. 62/288,661, filed Jan. 29, 2016, Muir et al. cited by applicant

U.S. Appl. No. 62/357,332, filed Jun. 30, 2016, Liu et al. cited by applicant

International Search Report for PCT/US2018/021880, mailed Jun. 20, 2018. cited by applicant

International Preliminary Report on Patentability for PCT/US2018/021880, mailed on Sep. 19, 2019. cited by applicant

[No Author Listed] HyPhy—Hypothesis testing using Phylogenies. Last modified Apr. 21, 2017. Accessed online via [http://hyphy.org/w/index.php/Main\\_Page](http://hyphy.org/w/index.php/Main_Page) on Apr. 28, 2021. cited by applicant

[No Author Listed] NCBI Accession No. XP\_015843220.1. C →U editing enzyme APOBEC-1 [*Peromyscus maniculatus bairdii*], XP002793540. Mar. 21, 2016. cited by applicant

[No Author Listed] NCBI Accession No. XP\_021505673.1. C →U editing enzyme APOBEC-1 [*Meriones unguiculatus*], XP002793541. Jun. 27, 2017. cited by applicant

[No Author Listed] NCBI Reference Sequence: WP\_00087959824.1. Oct. 9, 2019. 2 pages. cited by applicant

[No Author Listed] Score result for SEQ 355 to W02017032580. Muir et al. 2016. cited by applicant

[No Author Listed], “Human genome.” Encyclopedia Britannica. Encyclopedia Britannica, Inc. Published Feb. 15, 2019. Last accessed online via <https://www.britannica.com/science/human-genome> on Mar. 19, 2021. 2 pages. cited by applicant

[No Author Listed], EMBL Accession No. Q99ZW2. Nov. 2012. 2 pages. cited by applicant

[No Author Listed], Invitrogen Lipofectamine™ 2000 product sheets, 2002. 2 pages. cited by applicant

[No Author Listed], Invitrogen Lipofectamine™ 2000 product sheets, 2005. 3 pages. cited by applicant

[No Author Listed], Invitrogen Lipofectamine™ LTX product sheets, 2011. 4 pages. cited by applicant

[No Author Listed], *Mus musculus* (Mouse). UniProtKB Accession No. P51908 (ABEC1\_MOUSE). Oct. 1, 1996. 10 pages. cited by applicant

[No Author Listed], Thermo Fisher Scientific—How Cationic Lipid Mediated Transfection Works, retrieved from the internet Aug. 27, 2015. 2 pages. cited by applicant

Abremski et al., Bacteriophage P1 site-specific recombination. Purification and properties of the Cre recombinase protein. J Biol Chem. Feb. 10, 1984;259(3):1509-14. cited by applicant

Abudayyeh et al., C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector. Science Aug. 2016;353(6299):aaf5573. DOI: 10.1126/science.aaf5573. cited by applicant

Abudayyeh et al., A cytosine deaminase for programmable single-base RNA editing. Science. Jul. 26, 2019;365(6451):382-386. doi: 10.1126/science.aax7063. Epub Jul. 11, 2019. cited by applicant

Abudayyeh et al., RNA targeting with CRISPR-Cas13. Nature. Oct. 12, 2017;550(7675):280-284. doi: 10.1038/nature24049. Epub Oct. 4, 2017. cited by applicant

Ada et al., Carbohydrate-protein conjugate vaccines. Clin Microbiol Infect. Feb. 2003;9(2):79-85. doi: 10.1046/j.1469-0691.2003.00530.x. cited by applicant

Adamala et al., Programmable RNA-binding protein composed of repeats of a single modular unit. Proc Natl Acad Sci U S A. May 10, 2016;113(19):E2579-88. doi: 10.1073/pnas.1519368113. Epub Apr. 26, 2016. cited by applicant

Adams et al., New biarsenical ligands and tetracysteine motifs for protein labeling in vitro and in vivo: synthesis and biological applications. J Am Chem Soc. May 29, 2002;124(21):6063-76. doi: 10.1021/ja017687n. cited by applicant

Addgene Plasmid # 44246. pCas9-humanized, 2017, Stanley Qi. cited by applicant

Addgene Plasmid # 73021. PCMV-BE3, 2017, David Liu. cited by applicant

Addgene Plasmid # 79620. pcDNA3.1\_pCMV-nCas-PmCDA1-ugi pH1-gRNA(HPRT), 2017, Akihiko Kondo. cited by applicant

Adli, The CRISPR tool kit for genome editing and beyond. Nat Commun. May 15, 2018;9(1):1911. doi: 10.1038/s41467-018-04252-2. cited by applicant

Aguilo et al., Coordination of m(6)A mRNA Methylation and Gene Transcription by ZFP217 Regulates Pluripotency and Reprogramming. Cell Stem Cell. Dec. 3, 2015;17(6):689-704. doi: 10.1016/j.stem.2015.09.005. Epub Oct. 29, 2015. cited by applicant

Ahmad et al., Antibody-mediated specific binding and cytotoxicity of liposome-entrapped doxorubicin to lung cancer cells in vitro. Cancer Res. Sep. 1, 1992;52(17):4817-20. cited by applicant

Aihara et al., A conformational switch controls the DNA cleavage activity of lambda integrase. Mol Cell. Jul. 2003;12(1):187-98. cited by applicant

Aik et al., Structure of human RNA 7-methyladenine demethylase ALKBH5 provides insights into its mechanisms of

nucleic acid recognition and demethylation. *Nucleic Acids Res.* Apr. 2014;42(7):4741-54. doi: 10.1093/nar/gku085. Epub Jan. 30, 2014. cited by applicant

Aird et al., Increasing Cas9-mediated homology-directed repair efficiency through covalent tethering of DNA repair template. *Commun Biol.* May 31, 2018;1:54. doi: 10.1038/s42003-018-0054-2. cited by applicant

Akcakaya et al., In vivo CRISPR editing with no detectable genome-wide off-target mutations. *Nature.* Sep. 2018;561(7723):416-419. doi: 10.1038/s41586-018-0500-9. Epub Sep. 12, 2018. PMID: 30209390; PMCID: PMC6194229. cited by applicant

Akins et al., Mitochondrial plasmids of *Neurospora*: integration into mitochondrial DNA and evidence for reverse transcription in mitochondria. *Cell.* Nov. 21, 1986;47(4):505-16. doi: 10.1016/0092-8674(86)90615-x. cited by applicant

Akinsheye et al., Fetal hemoglobin in sickle cell anemia. *Blood.* Jul. 7, 2011;118(1):19-27. doi: 10.1182/blood-2011-03-325258. Epub Apr. 13, 2011. cited by applicant

Akopian et al., Chimeric recombinases with designed DNA sequence recognition. *Proc Natl Acad Sci U S A.* Jul. 22, 2003;100(15):8688-91. Epub Jul. 1, 2003. cited by applicant

Alarcón et al., HNRNPA2B1 Is a Mediator of m(6)A-Dependent Nuclear RNA Processing Events. *Cell.* Sep. 10, 2015;162(6):1299-308. doi: 10.1016/j.cell.2015.08.011. Epub Aug. 27, 2015. cited by applicant

Alarcón et al., N6-methyladenosine marks primary microRNAs for processing. *Nature.* Mar. 26, 2015;519(7544):482-5. doi: 10.1038/nature14281. Epub Mar. 18, 2015. cited by applicant

Alexander, HFE-associated hereditary hemochromatosis. *Genet Med.* May 2009;11(5):307-13. doi: 10.1097/GIM.0b013e31819d30f2. cited by applicant

Alexandrov et al., Signatures of mutational processes in human cancer. *Nature.* Aug. 22, 2013;500(7463):415-21. doi: 10.1038/nature12477. Epub Aug. 14, 2013. cited by applicant

Ali et al., Novel genetic abnormalities in Bernard-Soulier syndrome in India. *Ann Hematol.* Mar. 2014;93(3):381-4. doi: 10.1007/s00277-013-1895-x. Epub Sep. 1, 2013. cited by applicant

Altschul et al., Basic local alignment search tool. *J Mol Biol.* Oct. 5, 1990;215(3):403-10. doi: 10.1016/S0022-2836(05)80360-2. cited by applicant

Amato et al., Interpreting elevated fetal hemoglobin in pathology and health at the basic laboratory level: new and known  $\gamma$ -gene mutations associated with hereditary persistence of fetal hemoglobin. *Int J Lab Hematol.* Feb. 2014;36(1):13-9. doi: 10.1111/ijlh.12094. Epub Apr. 29, 2013. cited by applicant

Ames et al., A eubacterial riboswitch class that senses the coenzyme tetrahydrofolate. *Chem Biol.* Jul. 30, 2010;17(7):681-5. doi: 10.1016/j.chembiol.2010.05.020. cited by applicant

Amrann et al., Tightly regulated tac promoter vectors useful for the expression of unfused and fused proteins in *Escherichia coli*. *Gene.* Sep. 30, 1988;69(2):301-15. cited by applicant

Anders et al., Chapter One: In Vitro Enzymology of Cas9. in *Methods in Enzymology*, eds Doudna et al. 2014: 546:1-20. cited by applicant

Anders et al., Structural basis of PAM-dependent target DNA recognition by the Cas9 endonuclease. *Nature.* Sep. 25, 2014;513(7519):569-73. doi: 10.1038/nature13579. Epub Jul. 27, 2014. cited by applicant

Anderson, Human gene therapy. *Science.* May 8, 1992;256(5058):808-13. doi: 10.1126/science.1589762. cited by applicant

André et al., Axotomy-induced expression of calcium-activated chloride current in subpopulations of mouse dorsal root ganglion neurons. *J Neurophysiol.* Dec. 2003;90(6):3764-73. doi: 10.1152/jn.00449.2003. Epub Aug. 27, 2003. cited by applicant

Anzalone et al., Reprogramming eukaryotic translation with ligand-responsive synthetic RNA switches. *Nat Methods.* May 2016;13(5):453-8. doi: 10.1038/nmeth.3807. Epub Mar. 21, 2016. cited by applicant

Anzalone et al., Search-and-replace genome editing without double-strand breaks or donor DNA. *Nature.* Dec. 2019;576(7785):149-157. doi: 10.1038/s41586-019-1711-4. Epub Oct. 21, 2019. cited by applicant

Aplan, Causes of oncogenic chromosomal translocation. *Trends Genet.* Jan. 2006;22(1):46-55. doi: 10.1016/j.tig.2005.10.002. Epub Oct. 28, 2005. cited by applicant

Arakawa et al., A method to convert mRNA into a gRNA library for CRISPR/Cas9 editing of any organism. *Sci Adv.* Aug. 24, 2016;2(8):e1600699. doi: 10.1126/sciadv.1600699. cited by applicant

Araki et al., Comparative analysis of right element mutant lox sites on recombination efficiency in embryonic stem cells. *BMC Biotechnol.* Mar. 31, 2010;10:29. doi: 10.1186/1472-6750-10-29. cited by applicant

Araki et al., Site-specific recombinase, R, encoded by yeast plasmid pSR1. *J Mol Biol.* May 5, 1992;225(1):25-37. doi: 10.1016/0022-2836(92)91023-i. cited by applicant

Araki et al., Targeted integration of DNA using mutant lox sites in embryonic stem cells. *Nucleic Acids Res.* Feb. 15, 1997;25(4):868-72. doi: 10.1093/nar/25.4.868. cited by applicant

Arambula et al., Surface display of a massively variable lipoprotein by a *Legionella* diversity-generating retroelement. *Proc Natl Acad Sci U S A.* May 14, 2013;110(20):8212-7. doi: 10.1073/pnas.1301366110. Epub Apr. 30, 2013. cited by applicant

Arazoe et al., Targeted Nucleotide Editing Technologies for Microbial Metabolic Engineering. *Biotechnol J.* Sep. 2018;13(9):e1700596. doi: 10.1002/biot.201700596. Epub Jun. 19, 2018. cited by applicant

Arbab et al., Cloning-free CRISPR. *Stem Cell Reports.* Nov. 10, 2015;5(5):908-917. doi: 10.1016/j.stemcr.2015.09.022.

Epub Oct. 29, 2015. cited by applicant

Arezi et al., Novel mutations in Moloney Murine Leukemia Virus reverse transcriptase increase thermostability through tighter binding to template-primer. *Nucleic Acids Res.* Feb. 2009;37(2):473-81. doi: 10.1093/nar/gkn952. Epub Dec. 4, 2008. cited by applicant

Arnold et al., Mutants of Tn3 resolvase which do not require accessory binding sites for recombination activity. *EMBO J.* Mar. 1, 1999;18(5):1407-14. cited by applicant

Asante et al., A naturally occurring variant of the human prion protein completely prevents prion disease. *Nature.* Jun. 25, 2015;522(7557):478-81. doi: 10.1038/nature14510. Epub Jun. 10, 2015. cited by applicant

Asokan et al., The AAV vector toolkit: poised at the clinical crossroads. *Mol Ther.* Apr. 2012;20(4):699-708. doi: 10.1038/mt.2011.287. Epub Jan. 24, 2012. cited by applicant

Atkins et al., Ribosomal frameshifting and transcriptional slippage: From genetic steganography and cryptography to adventitious use. *Nucleic Acids Res.* Sep. 6, 2016;44(15):7007-78. doi: 10.1093/nar/gkw530. Epub Jul. 19, 2016. cited by applicant

Auer et al., Highly efficient CRISPR/Cas9-mediated knock-in in zebrafish by homology-independent DNA repair. *Genome Res.* Jan. 2014;24(1):142-53. doi: 10.1101/gr.161638.113. Epub Oct. 31, 2013. cited by applicant

Auricchio et al., Exchange of surface proteins impacts on viral vector cellular specificity and transduction characteristics: the retina as a model. *Hum Mol Genet.* Dec. 15, 2001;10(26):3075-81. doi: 10.1093/hmg/10.26.3075. cited by applicant

Autieri et al., IRT-1, a novel interferon-gamma-responsive transcript encoding a growth-suppressing basic leucine zipper protein. *J Biol Chem.* Jun. 12, 1998;273(24):14731-7. doi: 10.1074/jbc.273.24.14731. cited by applicant

Avidan et al., The processivity and fidelity of DNA synthesis exhibited by the reverse transcriptase of bovine leukemia virus. *Eur J Biochem.* Feb. 2002;269(3):859-67. doi: 10.1046/j.0014-2956.2001.02719.x. cited by applicant

Babacic et al., CRISPR-cas gene-editing as plausible treatment of neuromuscular and nucleotide-repeat-expansion diseases: A systematic review. *PLoS One.* Feb. 22, 2019;14(2):e0212198. doi: 10.1371/journal.pone.0212198. cited by applicant

Bacman et al., Specific elimination of mutant mitochondrial genomes in patient-derived cells by mitoTALENs. *Nat Med.* Sep. 2013;19(9):1111-3. doi: 10.1038/nm.3261. Epub Aug. 4, 2013. cited by applicant

Badran et al., Continuous evolution of *Bacillus thuringiensis* toxins overcomes insect resistance. *Nature.* May 5, 2016;533(7601):58-63. doi: 10.1038/nature17938. Epub Apr. 27, 2016. cited by applicant

Badran et al., Development of potent in vivo mutagenesis plasmids with broad mutational spectra. *Nat Commun.* Oct. 7, 2015;6:8425. doi: 10.1038/ncomms9425. cited by applicant

Badran et al., In vivo continuous directed evolution. *Curr Opin Chem Biol.* Feb. 2015;24:1-10. doi: 10.1016/j.cbpa.2014.09.040. Epub Nov. 7, 2014. cited by applicant

Bae et al., Cas-OFFinder: a fast and versatile algorithm that searches for potential off-target sites of Cas9 RNA-guided endonucleases. *Bioinformatics.* May 15, 2014;30(10):1473-5. doi: 10.1093/bioinformatics/btu048. Epub Jan. 24, 2014. cited by applicant

Bae et al., Microhomology-based choice of Cas9 nuclease target sites. *Nat Methods.* Jul. 2014;11(7):705-6. doi: 10.1038/nmeth.3015. cited by applicant

Bagal et al., Recent progress in sodium channel modulators for pain. *Bioorg Med Chem Lett.* Aug. 15, 2014;24(16):3690-9. doi: 10.1016/j.bmcl.2014.06.038. Epub Jun. 21, 2014. cited by applicant

Bagyinszky et al., Characterization of mutations in PRNP (prion) gene and their possible roles in neurodegenerative diseases. *Neuropsychiatr Dis Treat.* Aug. 14, 2018;14:2067-2085. doi: 10.2147/NDT.S165445. cited by applicant

Balakrishnan et al., Flap endonuclease 1. *Annu Rev Biochem.* 2013;82:119-38. doi: 10.1146/annurev-biochem-072511-122603. Epub Feb. 28, 2013. cited by applicant

Baldari et al., A novel leader peptide which allows efficient secretion of a fragment of human interleukin 1 beta in *Saccharomyces cerevisiae*. *EMBO J.* Jan. 1987;6(1):229-34. cited by applicant

Banerjee et al., Cadmium inhibits mismatch repair by blocking the ATPase activity of the MSH2-MSH6 complex [published correction appears in *Nucleic Acids Res.* 2005;33(5):1738]. *Nucleic Acids Res.* 2005;33(4):1410-1419. Published Mar. 3, 2005. doi:10.1093/nar/gki291. cited by applicant

Banerji et al., A lymphocyte-specific cellular enhancer is located downstream of the joining region in immunoglobulin heavy chain genes. *Cell.* Jul. 1983;33(3):729-40. doi: 10.1016/0092-8674(83)90015-6. cited by applicant

Bannert et al., Retroelements and the human genome: new perspectives on an old relation. *Proc Natl Acad Sci U S A.* Oct. 5, 2004;101 Suppl 2(Suppl 2):14572-9. doi: 10.1073/pnas.0404838101. Epub Aug. 13, 2004. cited by applicant

Banno et al., Deaminase-mediated multiplex genome editing in *Escherichia coli*. *Nat Microbiol.* Apr. 2018;3(4):423-429. doi: 10.1038/s41564-017-0102-6. Epub Feb. 5, 2018. cited by applicant

Barmania et al., C—C chemokine receptor type five (CCR5): An emerging target for the control of HIV infection. *Appl Transl Genom.* May 26, 2013;2:3-16. doi: 10.1016/j.atg.2013.05.004. cited by applicant

Barnes et al., Repair and genetic consequences of endogenous DNA base damage in mammalian cells. *Annu Rev Genet.* 2004;38:445-76. cited by applicant

Barnes et al., The fidelity of Taq polymerase catalyzing PCR is improved by an -terminal deletion. *Gene.* Mar. 1, 1992;112(1):29-35. doi: 10.1016/0378-1119(92)90299-5. cited by applicant

Barrangou et al., CRISPR provides acquired resistance against viruses in prokaryotes. *Science.* Mar. 23,

2007;315(5819):1709-12. cited by applicant

Barrangou, RNA-mediated programmable DNA cleavage. *Nat Biotechnol.* Sep. 2012;30(9):836-8. doi: 10.1038/nbt.2357. cited by applicant

Bartlett et al., Efficient expression of protein coding genes from the murine U1 small nuclear RNA promoters. *Proc Natl Acad Sci U S A.* Aug. 20, 1996;93(17):8852-7. doi: 10.1073/pnas.93.17.8852. cited by applicant

Bartosovic et al., N6-methyladenosine demethylase FTO targets pre-mRNAs and regulates alternative splicing and 3'-end processing. *Nucleic Acids Res.* Nov. 2, 2017;45(19):11356-11370. doi: 10.1093/nar/gkx778. cited by applicant

Basha et al., Influence of cationic lipid composition on gene silencing properties of lipid nanoparticle formulations of siRNA in antigen-presenting cells. *Mol Ther.* Dec. 2011;19(12):2186-200. doi: 10.1038/mt.2011.190. Epub Oct. 4, 2011. cited by applicant

Basturea et al., Substrate specificity and properties of the *Escherichia coli* 16S rRNA methyltransferase, RsmE. *RNA.* Nov. 2007;13(11):1969-76. doi: 10.1261/rna.700507. Epub Sep. 13, 2007. cited by applicant

Batey et al., Structure of a natural guanine-responsive riboswitch complexed with the metabolite hypoxanthine. *Nature.* Nov. 18, 2004;432(7015):411-5. cited by applicant

Beale et al., Comparison of the differential context-dependence of DNA deamination by APOBEC enzymes: correlation with mutation spectra in vivo. *J Mol Biol.* Mar. 26, 2004;337(3):585-96. cited by applicant

Beaudry et al., Directed evolution of an RNA enzyme. *Science.* Jul. 31, 1992;257(5070):635-41. doi: 10.1126/science.1496376. cited by applicant

Bebenek et al., Error-prone polymerization by HIV-1 reverse transcriptase. Contribution of template-primer misalignment, miscoding, and termination probability to mutational hot spots. *J Biol Chem.* May 15, 1993;268(14):10324-34. cited by applicant

Bedell et al., In vivo genome editing using a high-efficiency TALEN system. *Nature.* Nov. 1, 2012;491(7422):114-8. Doi: 10.1038/nature11537. Epub Sep. 23, 2012. cited by applicant

Begley, Scientists unveil the 'most clever CRISPR gadget' so far. *STAT*, Apr. 20, 2016. <https://www.statnews.com/2016/04/20/clever-crispr-advance-unveiled/>. cited by applicant

Behr, Gene transfer with synthetic cationic amphiphiles: prospects for gene therapy. *Bioconjug Chem.* Sep.-Oct. 1994;5(5):382-9. doi: 10.1021/bc00029a002. cited by applicant

Bell et al., Ribozyme-catalyzed excision of targeted sequences from within RNAs. *Biochemistry.* Dec. 24, 2002;41(51):15327-33. doi: 10.1021/bi0267386. cited by applicant

Belshaw et al., Controlling programmed cell death with a cyclophilin-cyclosporin-based chemical inducer of dimerization. *Chem Biol.* Sep. 1996;3(9):731-8. doi: 10.1016/s1074-5521(96)90249-5. cited by applicant

Belshaw et al., Controlling protein association and subcellular localization with a synthetic ligand that induces heterodimerization of proteins. *Proc Natl Acad Sci U S A.* May 14, 1996;93(10):4604-7. doi: 10.1073/pnas.93.10.4604. cited by applicant

Benarroch, HCN channels: function and clinical implications. *Neurology.* Jan. 15, 2013;80(3):304-10. doi: 10.1212/WNL.0b013e31827dec42. cited by applicant

Bennett et al., Painful and painless channelopathies. *Lancet Neurol.* Jun. 2014;13(6):587-99. doi: 10.1016/S1474-4422(14)70024-9. Epub May 6, 2014. cited by applicant

Bentin, T., A ribozyme transcribed by a ribozyme. *Artif DNA PNA XNA.* Apr. 2011;2(2):40-42. doi: 10.4161/adna.2.2.16852. cited by applicant

Berger et al., Reverse transcriptase and its associated ribonuclease H: interplay of two enzyme activities controls the yield of single-stranded complementary deoxyribonucleic acid. *Biochemistry.* May 10, 1983;22(10):2365-72. doi: 10.1021/bi00279a010. cited by applicant

Berges et al., Transduction of brain by herpes simplex virus vectors. *Mol Ther.* Jan. 2007;15(1):20-9. doi: 10.1038/sj.mt.6300018. cited by applicant

Berkhout et al., Identification of an active reverse transcriptase enzyme encoded by a human endogenous HERV-K retrovirus. *J Virol.* Mar. 1999;73(3):2365-75. doi: 10.1128/JVI.73.3.2365-2375.1999. cited by applicant

Bernhart et al., Local RNA base pairing probabilities in large sequences. *Bioinformatics.* Mar. 1, 2006;22(5):614-5. doi: 10.1093/bioinformatics/btk014. Epub Dec. 20, 2005. cited by applicant

Bernstein et al., Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature.* Jan. 18, 2001;409(6818):363-6. doi: 10.1038/35053110. cited by applicant

Bershtein et al., Advances in laboratory evolution of enzymes. *Curr Opin; Chem Biol.* Apr. 2008;12(2):151-8. doi: 10.1016/j.cbpa.2008.01.027. Epub Mar. 7, 2008. Review. cited by applicant

Bertolotti et al., Toward genosafe endonuclease-boosted gene targeting using breakthrough CRISP/Cas9 for next generation stem cell gene therapy culminating in efficient ex VIVO in VIVO gene repair/genomic editing. *Molecular Therapy.* May 2015;23(Suppl1):S139. Abstract 350. 18th Ann Meeting of the American Society of Gene and Cell Therapy. ASGCT 2015. New Orleans, LA. May 13, 2015-May 16, 2015. cited by applicant

Bertrand et al., Localization of ASHI mRNA particles in living yeast. *Mol Cell.* Oct. 1998;2(4):437-45. doi: 10.1016/s1097-2765(00)80143-4. cited by applicant

Beumer et al., Efficient gene targeting in *Drosophila* with zinc-finger nucleases. *Genetics.* Apr. 2006;172(4):2391-403. Epub Feb. 1, 2006. cited by applicant

Bhagwat, DNA-cytosine deaminases: from antibody maturation to antiviral defense. *DNA Repair (Amst)*. Jan. 5, 2004;3(1):85-9. cited by applicant

Bi et al., Pseudo attP sites in favor of transgene integration and expression in cultured porcine cells identified by *Streptomyces* phage phiC31 integrase. *BMC Mol Biol*. Sep. 8, 2013;14:20. doi: 10.1186/1471-2199-14-20. cited by applicant

Bibb et al., Integration and excision by the large serine recombinase phiRv1 integrase. *Mol Microbiol*. Mar. 2005;55(6):1896-910. doi: 10.1111/j.1365-2958.2005.04517.x. cited by applicant

Biehs et al., DNA Double-Strand Break Resection Occurs during Non-homologous End Joining in G1 but Is Distinct from Resection during Homologous Recombination. *Mol Cell*. Feb. 16, 2017;65(4):671-684.e5. doi: 10.1016/j.molcel.2016.12.016. Epub Jan. 26, 2017. cited by applicant

Billon et al., CRISPR-Mediated Base Editing Enables Efficient Disruption of Eukaryotic Genes through Induction of STOP Codons. *Mol Cell*. Sep. 21, 2017;67(6):1068-1079.e4. doi: 10.1016/j.molcel.2017.08.008. Epub Sep. 7, 2017. cited by applicant

Birling et al., Site-specific recombinases for manipulation of the mouse genome. *Methods Mol Biol*. 2009;561:245-63. doi: 10.1007/978-1-60327-019-9\_16. cited by applicant

Biswas et al., A structural basis for allosteric control of DNA recombination by lambda integrase. *Nature*. Jun. 23, 2005;435(7045):1059-66. doi: 10.1038/nature03657. cited by applicant

Bitinaite et al., FokI dimerization is required for DNA cleavage. *Proc Natl Acad Sci U S A*. Sep. 1, 1998;95(18):10570-5. cited by applicant

Blaese et al., Vectors in cancer therapy: how will they deliver? *Cancer Gene Ther*. Dec. 1995;2(4):291-7. cited by applicant

Blain et al., Nuclease activities of Moloney murine leukemia virus reverse transcriptase. Mutants with altered substrate specificities. *J Biol Chem*. Nov. 5, 1993;268(31):23585-92. cited by applicant

Blaisonneau et al., A circular plasmid from the yeast *Torulaspora delbrueckii*. *Plasmid*. 1997;38(3):202-9. doi: 10.1006/plas.1997.1315. cited by applicant

Blau et al., A proliferation switch for genetically modified cells. *PNAS* Apr. 1, 1997 94 (7) 3076-3081; <https://doi.org/10.1073/pnas.94.7.3076>. cited by applicant

Bloom et al., Evolving strategies for enzyme engineering. *Curr Opin Struct Biol*. Aug. 2005;15(4):447-52. cited by applicant

Boch, TALEs of genome targeting. *Nat Biotechnol*. Feb. 2011;29(2):135-6. Doi: 10.1038/nbt.1767. cited by applicant

Bodi et al., Yeast m6A Methylated mRNAs Are Enriched on Translating Ribosomes during Meiosis, and under Rapamycin Treatment. *PLoS One*. Jul. 17, 2015;10(7):e0132090. doi: 10.1371/journal.pone.0132090. cited by applicant

Boeckle et al., Melittin analogs with high lytic activity at endosomal pH enhance transfection with purified targeted PEI polyplexes. *J Control Release*. May 15, 2006;112(2):240-8. Epub Mar. 20, 2006. cited by applicant

Boersma et al., Selection strategies for improved biocatalysts. *FEBS J*. May 2007;274(9):2181-95. cited by applicant

Bogdanove et al., Engineering altered protein-DNA recognition specificity. *Nucleic Acids Res*. Jun. 1, 2018;46(10):4845-4871. doi: 10.1093/nar/gky289. cited by applicant

Bogdanove et al., TAL effectors: customizable proteins for DNA targeting. *Science*. Sep. 30, 2011;333(6051):1843-6. doi: 10.1126/science.1204094. cited by applicant

Bohlke et al., Sense codon emancipation for proteome-wide incorporation of noncanonical amino acids: rare isoleucine codon AUA as a target for genetic code expansion. *FEMS Microbiol Lett*. Feb. 2014;351(2):133-44. doi: 10.1111/1574-6968.12371. Epub Jan. 27, 2014. cited by applicant

Bolotin et al., Clustered regularly interspaced short palindrome repeats (CRISPRs) have spacers of extrachromosomal origin. *Microbiology*. Aug. 2005;151(Pt 8):2551-61. cited by applicant

Bolusani et al., Evolution of variants of yeast site-specific recombinase F1p that utilize native genomic sequences as recombination target sites. *Nucleic Acids Res*. 2006;34(18):5259-69. Epub Sep. 26, 2006. cited by applicant

Bondeson et al., Inversion of the IDS gene resulting from recombination with IDS-related sequences is a common cause of the Hunter syndrome. *Hum Mol Genet*. Apr. 1995;4(4):615-21. doi: 10.1093/hmg/4.4.615. cited by applicant

Borchardt et al., Controlling mRNA stability and translation with the CRISPR endoribonuclease Csy4. *RNA*. Nov. 2015;21(11):1921-30. doi: 10.1261/rna.051227.115. Epub Sep. 9, 2015. cited by applicant

Borman, Improved route to single-base genome editing. *Chemical & Engineering News*, Apr. 25, 2016;94(17)p5. <http://cen.acs.org/articles/94/i17/Improved-route-single-base-genome.html>. cited by applicant

Bourinet et al., Silencing of the Cav3.2 T-type calcium channel gene in sensory neurons demonstrates its major role in nociception. *EMBO J*. Jan. 26, 2005;24(2):315-24. doi: 10.1038/sj.emboj.7600515. Epub Dec. 16, 2004. cited by applicant

Boutabout et al., DNA synthesis fidelity by the reverse transcriptase of the yeast retrotransposon Ty1. *Nucleic Acids Res*. Jun. 1, 2001;29(11):2217-22. doi: 10.1093/nar/29.11.2217. cited by applicant

Box et al., A multi-domain protein system based on the HC fragment of tetanus toxin for targeting DNA to neuronal cells. *J Drug Target*. Jul. 2003;11(6):333-43. doi: 10.1080/1061186310001634667. cited by applicant

Branden and Tooze, Introduction to Protein Structure. 1999; 2nd edition. Garland Science Publisher: 3-12. cited by applicant

Braun et al., Immunogenic duplex nucleic acids are nuclease resistant. *J Immunol.* Sep. 15, 1988;141(6):2084-9. cited by applicant

Brierley et al., Viral RNA pseudoknots: versatile motifs in gene expression and replication. *Nat Rev Microbiol.* Aug. 2007;5(8):598-610. doi: 10.1038/nrmicro1704. cited by applicant

Briner et al., Guide RNA functional modules direct Cas9 activity and orthogonality. *Mol Cell.* Oct. 23, 2014;56(2):333-339. doi: 10.1016/j.molcel.2014.09.019. cited by applicant

Britt et al., Re-engineering plant gene targeting. *Trends Plant Sci.* Feb. 2003;8(2):90-5. cited by applicant

Brouns et al., Small CRISPR RNAs guide antiviral defense in prokaryotes. *Science.* Aug. 15, 2008;321(5891):960-4. doi: 10.1126/science.1159689. cited by applicant

Brown et al., A mammalian protein targeted by G1-arresting rapamycin-receptor complex. *Nature.* Jun. 30, 1994;369(6483):756-8. doi: 10.1038/369756a0. cited by applicant

Brown et al., Characterization of the genetic elements required for site-specific integration of plasmid pSE211 in *Saccharopolyspora erythraea*. *J Bacteriol.* Apr. 1990;172(4):1877-88. doi: 10.1128/jb.172.4.1877-1888.1990. cited by applicant

Brown et al., Serine recombinases as tools for genome engineering. *Methods.* Apr. 2011;53(4):372-9. doi: 10.1016/j.ymeth.2010.12.031. Epub Dec. 30, 2010. cited by applicant

Brown et al., Structural insights into the stabilization of MALAT1 noncoding RNA by a bipartite triple helix. *Nat Struct Mol Biol.* Jul. 2014;21(7):633-40. doi: 10.1038/nsmb.2844. Epub Jun. 22, 2014. cited by applicant

Brusse et al., Spinocerebellar ataxia associated with a mutation in the fibroblast growth factor 14 gene (SCA27): A new phenotype. *Mov Disord.* Mar. 2006;21(3):396-401. cited by applicant

Brzezicha et al., Identification of human tRNA:m5C methyltransferase catalysing intron-dependent m5C formation in the first position of the anticodon of the pre-tRNA Leu (CAA). *Nucleic Acids Res.* 2006;34(20):6034-43. doi: 10.1093/nar/gk1765. Epub Oct. 27, 2006. cited by applicant

Buchholz et al., Alteration of Cre recombinase site specificity by substrate-linked protein evolution. *Nat Biotechnol.* Nov. 2001;19(11):1047-52. cited by applicant

Buchsachner et al., Human immunodeficiency virus vectors for inducible expression of foreign genes. *J Virol.* May 1992;66(5):2731-9. doi: 10.1128/JVI.66.5.2731-2739.1992. cited by applicant

Buchwald et al., Long-term, continuous intravenous heparin administration by an implantable infusion pump in ambulatory patients with recurrent venous thrombosis. *Surgery.* Oct. 1980;88(4):507-16. cited by applicant

Buckley et al., Targeting the von Hippel-Lindau E3 ubiquitin ligase using small molecules to disrupt the VHL/HIF-1? interaction. *J Am Chem Soc.* Mar. 14, 2012;134(10):4465-8. doi: 10.1021/ja209924v. Epub Feb. 27, 2012. cited by applicant

Budisa et al., Residue-specific bioincorporation of non-natural, biologically active amino acids into proteins as possible drug carriers: structure and stability of the per-thiaproline mutant of annexin V. *Proc Natl Acad Sci U S A.* Jan. 20, 1998;95(2):455-9. cited by applicant

Budker et al., Protein/amphipathic polyamine complexes enable highly efficient transfection with minimal toxicity. *Biotechniques.* Jul. 1997;23(1):139, 142-7. doi: 10.2144/97231rr02. cited by applicant

Budworth et al., A brief history of triplet repeat diseases. *Methods Mol Biol.* 2013;1010:3-17. doi: 10.1007/978-1-62703-411-1\_1. cited by applicant

Bulow et al., Multienzyme systems obtained by gene fusion. *Trends Biotechnol.* Jul. 1991;9(7):226-31. cited by applicant

Burke et al., Activating mutations of Tn3 resolvase marking interfaces important in recombination catalysis and its regulation. *Mol Microbiol.* Feb. 2004;51(4):937-48. cited by applicant

Burke et al., RNA Aptamers to the Adenosine Moiety of S-adenosyl Methionine: Structural Inferences From Variations on a Theme and the Reproducibility of SELEX. *Nucleic Acids Res.* May 15, 1997;25(10):2020-4. doi: 10.1093/nar/25.10.2020. cited by applicant

Burstein et al., New CRISPR-Cas systems from uncultivated microbes. *Nature* Feb. 2017;542(7640):237-240. cited by applicant

Burton et al., Gene delivery using herpes simplex virus vectors. *DNA Cell Biol.* Dec. 2002;21(12):915-36. doi: 10.1089/104454902762053864. cited by applicant

Buskirk et al., Directed evolution of ligand dependence: small-molecule-activated protein splicing. *Proc Natl Acad Sci U S A.* Jul. 20, 2004;101(29):10505-10. Epub Jul. 9, 2004. cited by applicant

Buskirk et al., In vivo evolution of an RNA-based transcriptional activator. *Chem Biol.* Jun. 2003;10(6):533-40. doi: 10.1016/s1074-5521(03)00109-1. cited by applicant

Butt et al., Efficient CRISPR/Cas9-Mediated Genome Editing Using a Chimeric Single-Guide RNA Molecule. *Front Plant Sci.* Aug. 24, 2017;8:1441(1-8). doi: 10.3389/fpls.2017.01441. cited by applicant

Byrne et al., Multiplex gene regulation: a two-tiered approach to transgene regulation in transgenic mice. *Proc Natl Acad Sci U S A.* Jul. 1989;86(14):5473-7. doi: 10.1073/pnas.86.14.5473. cited by applicant

Böck et al., Selenocysteine: the 21st amino acid. *Mol Microbiol.* Mar. 1991;5(3):515-20. cited by applicant

Cade et al., Highly efficient generation of heritable zebrafish gene mutations using homo- and heterodimeric TALENs. *Nucleic Acids Res.* Sep. 2012;40(16):8001-10. Doi: 10.1093/nar/gks518. Epub Jun. 7, 2012. cited by applicant

Cadwell et al., Randomization of genes by PCR mutagenesis. *PCR Methods Appl.* Aug. 1992;2(1):28-33. doi:

10.1101/gr.2.1.28. cited by applicant

Cai et al., Reconstruction of ancestral protein sequences and its applications. BMC Evol Biol. Sep. 17, 2004;4:33. doi: 10.1186/1471-2148-4-33. cited by applicant

Calame et al., Transcriptional controlling elements in the immunoglobulin and T cell receptor loci. Adv Immunol. 1988;43:235-75. doi: 10.1016/s0065-2776(08)60367-3. cited by applicant

Caldecott et al., Single-strand break repair and genetic disease. Nat Rev Genet. Aug. 2008;9(8):619-31. doi: 10.1038/nrg2380. cited by applicant

Camarero et al., Biosynthesis of a Head-to-Tail Cyclized Protein with Improved Biological Activity. J. Am. Chem. Soc. May 29, 1999; 121(23):5597-5598. <https://doi.org/10.1021/ja990929n>. cited by applicant

Cameron, Recent advances in transgenic technology. Mol Biotechnol. Jun. 1997;7(3):253-65. cited by applicant

Camper et al., Postnatal repression of the alpha-fetoprotein gene is enhancer independent. Genes Dev. Apr. 1989;3(4):537-46. doi: 10.1101/gad.3.4.537. cited by applicant

Camps et al., Targeted gene evolution in *Escherichia coli* using a highly error-prone DNA polymerase I. Proc Natl Acad Sci U S A. Aug. 19, 2003;100(17):9727-32. Epub Aug. 8, 2003. cited by applicant

Canchaya et al., Genome analysis of an inducible prophage and prophage remnants integrated in the *Streptococcus pyogenes* strain SF370. Virology. Oct. 25, 2002;302(2):245-58. doi: 10.1006/viro.2002.1570. cited by applicant

Canver et al., Customizing the genome as therapy for the  $\beta$ -hemoglobinopathies. Blood. May 26, 2016;127(21):2536-45. doi: 10.1182/blood-2016-01-678128. Epub Apr. 6, 2016. cited by applicant

Cargill et al., Characterization of single-nucleotide polymorphisms in coding regions of human genes. Nat Genet. Jul. 1999;22(3):231-8. cited by applicant

Carlier et al., Burkholderia cenocepacia H111 Rhy-family protein. Apr. 16, 2015. Retrieved from the Internet via <https://www.ebi.ac.uk/ena/browser/api/embl/CDN65395.1?lineLimit=1000>. Last retrieved Apr. 26, 2021. cited by applicant

Carlson et al., Negative selection and stringency modulation in phage-assisted continuous evolution. Nat Chem Biol. Mar. 2014;10(3):216-22. doi: 10.1038/nchembio.1453. Epub Feb. 2, 2014. With Supplementary Results. cited by applicant

Caron et al., Intracellular delivery of a Tat-eGFP fusion protein into muscle cells. Mol Ther. Mar. 2001;3(3):310-8. cited by applicant

Carr et al., Genome engineering. Nat Biotechnol. Dec. 2009;27(12):1151-62. doi: 10.1038/nbt.1590. cited by applicant

Carroll et al., Gene targeting in *Drosophila* and *Caenorhabditis elegans* with zinc-finger nucleases. Methods Mol Biol. 2008;435:63-77. doi: 10.1007/978-1-59745-232-8\_5. cited by applicant

Carroll et al., Progress and prospects: zinc-finger nucleases as gene therapy agents. Gene Ther. Nov. 2008;15(22):1463-8. doi: 10.1038/gt.2008.145. Epub Sep. 11, 2008. cited by applicant

Carroll, A CRISPR approach to gene targeting. Mol Ther. Sep. 2012;20(9):1658-60. doi: 10.1038/mt.2012.171. cited by applicant

Carroll, Genome engineering with zinc-finger nucleases. Genetics. Aug. 2011;188(4):773-82. doi: 10.1534/genetics.111.131433. Review. cited by applicant

Carvalho et al., Evolution in health and medicine Sackler colloquium: Genomic disorders: a window into human gene and genome evolution. Proc Natl Acad Sci U S A. Jan. 26, 2010;107 Suppl 1(Suppl 1):1765-71. doi: 10.1073/pnas.0906222107. Epub Jan. 13, 2010. cited by applicant

Caspi et al., Distribution of split DnaE inteins in cyanobacteria. Mol Microbiol. Dec. 2003;50(5):1569-77. doi: 10.1046/j.1365-2958.2003.03825.x. cited by applicant

Cattaneo et al., SELIL affects human pancreatic cancer cell cycle and invasiveness through modulation of PTEN and genes related to cell-matrix interactions. Neoplasia. 2005;7(11):1030-1038. cited by applicant

Ceccaldi et al., Repair Pathway Choices and Consequences at the Double-Strand Break. Trends Cell Biol. Jan. 2016;26(1):52-64. doi: 10.1016/j.tcb.2015.07.009. Epub Oct. 1, 2015. cited by applicant

Cermak et al., Efficient design and assembly of custom TALEN and other TAL effector-based constructs for DNA targeting. Nucleic Acids Res. Jul. 2011;39(12):e82. Doi: 10.1093/nar/gkr218. Epub Apr. 14, 2011. cited by applicant

Chadalavada et al., Wild-type is the optimal sequence of the HDV ribozyme under cotranscriptional conditions. RNA. Dec. 2007;13(12):2189-201. doi: 10.1261/rna.778107. Epub Oct. 23, 2007. cited by applicant

Chadwick et al., In Vivo Base Editing of PCSK9 (Proprotein Convertase Subtilisin/Kexin Type 9) as a Therapeutic Alternative to Genome Editing. Arterioscler Thromb Vasc Biol. Sep. 2017;37(9):1741-1747. doi: 10.1161/ATVBAHA.117.309881. Epub Jul. 27, 2017. cited by applicant

Chaikind et al., A programmable Cas9-serine recombinase fusion protein that operates on DNA sequences in mammalian cells. Nucleic Acids Res. Nov. 16, 2016;44(20):9758-9770. Epub Aug. 11, 2016. cited by applicant

Chalberg et al., Integration specificity of phage phiC31 integrase in the human genome. J Mol Biol. Mar. 17, 2006;357(1):28-48. doi: 10.1016/j.jmb.2005.11.098. Epub Dec. 22, 2005. cited by applicant

Chalberg et al., phiC31 integrase confers genomic integration and long-term transgene expression in rat retina. Invest Ophthalmol Vis Sci. Jun. 2005;46(6):2140-6. doi: 10.1167/iovs.04-1252. cited by applicant

Chan et al., Molecular recording of mammalian embryogenesis. Nature. Jun. 2019;570(7759):77-82. doi: 10.1038/s41586-019-1184-5. Epub May 13, 2019. cited by applicant

Chan et al., Novel selection methods for DNA-encoded chemical libraries. Curr Opin Chem Biol. 2015;26:55-61.



doi:10.1016/j.cbpa.2015.02.010. cited by applicant

Chan et al., The choice of nucleotide inserted opposite abasic sites formed within chromosomal DNA reveals the polymerase activities participating in translesion DNA synthesis. *DNA Repair (Amst)*. Nov. 2013;12(11):878-89. doi: 10.1016/j.dnarep.2013.07.008. Epub Aug. 26, 2013. cited by applicant

Chapman et al., Playing the end game: DNA double-strand break repair pathway choice. *Mol Cell*. Aug. 24, 2012;47(4):497-510. doi: 10.1016/j.molcel.2012.07.029. cited by applicant

Chari et al., Unraveling CRISPR-Cas9 genome engineering parameters via a library-on-library approach. *Nat Methods*. Sep. 2015;12(9):823-6. doi: 10.1038/nmeth.3473. Epub Jul. 13, 2015. cited by applicant

Charpentier et al., Biotechnology: Rewriting a genome. *Nature*. Mar. 7, 2013;495(7439):50-1. doi: 10.1038/495050a. cited by applicant

Chaturvedi et al., Stabilization of triple-stranded oligonucleotide complexes: use of probes containing alternating phosphodiester and stereo-uniform cationic phosphoramidate linkages. *Nucleic Acids Res*. Jun. 15, 1996;24(12):2318-23. cited by applicant

Chavez et al., Highly efficient Cas9-mediated transcriptional programming. *Nat Methods*. Apr. 2015;12(4):326-8. doi: 10.1038/nmeth.3312. Epub Mar. 2, 2015. cited by applicant

Chavez et al., Precise Cas9 targeting enables genomic mutation prevention. *bioRxiv*. Jun. 14, 2016; <http://dx.doi.org/10.1101/058974>. 6 pages. bioRxiv preprint first posted online Jun. 14, 2016. cited by applicant

Chavez et al., Therapeutic applications of the PhiC31 integrase system. *Curr Gene Ther*. Oct. 2011;11(5):375-81. Review. cited by applicant

Chawla et al., An atlas of RNA base pairs involving modified nucleobases with optimal geometries and accurate energies. *Nucleic Acids Res*. Aug. 18, 2015;43(14):6714-29. doi: 10.1093/nar/gkv606. Epub Jun. 27, 2015. cited by applicant

Chelico et al., Biochemical basis of immunological and retroviral responses to DNA-targeted cytosine deamination by activation-induced cytidine deaminase and APOBEC3G. *J Biol Chem*. Oct. 9, 2009;284(41):27761-5. doi: 10.1074/jbc.R109.052449. Epub Aug. 13, 2009. cited by applicant

Chelico et al., Stochastic properties of processive cytidine DNA deaminases AID and APOBEC3G. *Philos Trans R Soc Lond B Biol Sci*. Mar. 12, 2009;(1517):583-93. doi: 10.1098/rstb.2008.0195. cited by applicant

Chen et al., Enhanced proofreading governs CRISPR-Cas9 targeting accuracy. *Nature*. Oct. 19, 2017;550(7676):407-410. doi: 10.1038/nature24268. Epub Sep. 20, 2017. cited by applicant

Chen et al., A general strategy for the evolution of bond-forming enzymes using yeast display. *Proc Natl Acad Sci U S A*. Jul. 12, 2011;108(28):11399-404. doi: 10.1073/pnas.1101046108. Epub Jun. 22, 2011. cited by applicant

Chen et al., Alterations in PMS2, MSH2 and MLH1 expression in human prostate cancer. *Int J Oncol*. May 2003;22(5):1033-43. cited by applicant

Chen et al., Fusion protein linkers: property, design and functionality. *Adv Drug Deliv Rev*. Oct. 2013;65(10):1357-69. doi: 10.1016/j.addr.2012.09.039. Epub Sep. 29, 2012. cited by applicant

Chen et al., Genome-wide CRISPR screen in a mouse model of tumor growth and metastasis. *Cell*. Mar. 12, 2015;160(6):1246-60. doi: 10.1016/j.cell.2015.02.038. Epub Mar. 5, 2015. cited by applicant

Chen et al., Highly Efficient Mouse Genome Editing by CRISPR Ribonucleoprotein Electroporation of Zygotes. *J Biol Chem*. Jul. 8, 2016;291(28):14457-67. doi: 10.1074/jbc.M116.733154. Epub May 5, 2016. cited by applicant

Chen et al., m(6)A RNA methylation is regulated by microRNAs and promotes reprogramming to pluripotency. *Cell Stem Cell*. Mar. 5, 2015;16(3):289-301. doi: 10.1016/j.stem.2015.01.016. Epub Feb. 12, 2015. cited by applicant

Chen et al., Structure of the DNA deaminase domain of the HIV-1 restriction factor APOBEC3G. *Nature*. Mar. 6, 2008;452(7183):116-9. doi: 10.1038/nature06638. Epub Feb. 20, 2008. cited by applicant

Chen et al., Targeting genomic rearrangements in tumor cells through Cas9-mediated insertion of a suicide gene. *Nat Biotechnol*. Jun. 2017;35(6):543-550. doi: 10.1038/nbt.3843. Epub May 1, 2017. cited by applicant

Cheng et al., Multiplexed activation of endogenous genes by CRISPR-on, an RNA-guided transcriptional activator system. *Cell Res*. Oct. 2013;23(10):1163-71. doi: 10.1038/cr.2013.122. Epub Aug. 27, 2013. cited by applicant

Chesnoy et al., Structure and function of lipid-DNA complexes for gene delivery. *Annu Rev Biophys Biomol Struct*. 2000;29:27-47. cited by applicant

Chester et al., The apolipoprotein B mRNA editing complex performs a multifunctional cycle and suppresses nonsense-mediated decay. *EMBO J*. Aug. 1, 2003;22(15):3971-82. doi: 10.1093/emboj/cdg369. cited by applicant

Chew et al., A multifunctional AAV-CRISPR-Cas9 and its host response. *Nat Methods*. Oct. 2016;13(10):868-74. doi: 10.1038/nmeth.3993. Epub Sep. 5, 2016. cited by applicant

Chew et al., A multifunctional AAV-CRISPR-Cas9 and its host response. *Nat Methods*. Oct. 2016;13(10):868-74. doi: 10.1038/nmeth.3993. Epub Sep. 5, 2016. Supplementary Information. cited by applicant

Chichili et al., Linkers in the structural biology of protein-protein interactions. *Protein Science*. 2013;22:153-67. cited by applicant

Chin, Expanding and reprogramming the genetic code of cells and animals. *Annu Rev Biochem*. 2014;83:379-408. doi: 10.1146/annurev-biochem-060713-035737. Epub Feb. 10, 2014. cited by applicant

Chipev et al., A leucine----proline mutation in the H1 subdomain of keratin 1 causes epidermolytic hyperkeratosis. *Cell*. Sep. 4, 1992;70(5):821-8. cited by applicant

Cho et al., Analysis of off-target effects of CRISPR/Cas-derived RNA-guided endonucleases and nickases. *Genome Res*.

Jan. 2012;14(1):132-41. doi: 10.1101/gr.162339.113. Epub Nov. 19, 2013. cited by applicant

Cho et al., Site-specific recombination of bacteriophage P22 does not require integration host factor. *J Bacteriol.* Jul. 1999;181(14):4245-9. doi: 10.1128/JB.181.14.4245-4249.1999. cited by applicant

Cho et al., Targeted genome engineering in human cells with the Cas9 RNA-guided endonuclease. *Nat Biotechnol.* Mar. 2013;31(3):230-2. doi: 10.1038/nbt.2507. Epub Jan. 29, 2013. cited by applicant

Cho et al., The calcium-activated chloride channel anoctamin 1 acts as a heat sensor in nociceptive neurons. *Nat Neurosci.* May 27, 2012;15(7):1015-21. doi: 10.1038/nn.3111. cited by applicant

Choe et al., Forging Ahead through Darkness: PCNA, Still the Principal Conductor at the Replication Fork. *Mol Cell.* Feb. 2, 2017;65(3):380-392. doi: 10.1016/j.molcel.2016.12.020. cited by applicant

Choi et al., (6)-methyladenosine in mRNA disrupts tRNA selection and translation-elongation dynamics. *Nat Struct Mol Biol.* Feb. 2016;23(2):110-5. doi: 10.1038/nsmb.3148. Epub Jan. 11, 2016. cited by applicant

Choi et al., Protein trans-splicing and characterization of a split family B-type DNA polymerase from the hyperthermophilic archaeal parasite *Nanoarchaeum equitans*. *J Mol Biol.* Mar. 10, 2006;356(5):1093-106. doi: 10.1016/j.jmb.2005.12.036. Epub Dec. 27, 2005. cited by applicant

Choi et al., Translesion synthesis across abasic lesions by human B-family and Y-family DNA polymerases  $\eta$ ,  $\theta$ ,  $\kappa$ ,  $\nu$ , and REV1. *J Mol Biol.* Nov. 19, 2010;404(1):34-44. doi: 10.1016/j.jmb.2010.09.015. Epub Oct. 1, 2010. cited by applicant

Chong et al., Modulation of protein splicing of the *Saccharomyces cerevisiae* vacuolar membrane ATPase intein. *J Biol Chem.* Apr. 24, 1998;273(17):10567-77. doi: 10.1074/jbc.273.17.10567. cited by applicant

Chong et al., Utilizing the C-terminal cleavage activity of a protein splicing element to purify recombinant proteins in a single chromatographic step. *Nucleic Acids Res.* Nov. 15, 1998;26(22):5109-15. doi: 10.1093/nar/26.22.5109. cited by applicant

Chong et al., Protein splicing involving the *Saccharomyces cerevisiae* VMA intein. The steps in the splicing pathway, side reactions leading to protein cleavage, and establishment of an in vitro splicing system. *J Biol Chem.* Sep. 6, 1996;271(36):22159-68. doi: 10.1074/jbc.271.36.22159. cited by applicant

Chong et al., Protein splicing of the *Saccharomyces cerevisiae* VMA intein without the endonuclease motifs. *J Biol Chem.* Jun. 20, 1997;272(25):15587-90. doi: 10.1074/jbc.272.25.15587. cited by applicant

Chong et al., Single-column purification of free recombinant proteins using a self-cleavable affinity tag derived from a protein splicing element. *Gene.* Jun. 19, 1997;192(2):271-81. doi: 10.1016/s0378-1119(97)00105-4. cited by applicant

Choudhury et al., CRISPR-dCas9 mediated TET1 targeting for selective DNA demethylation at BRCA1 promoter. *Oncotarget.* Jul. 19, 2016;7(29):46545-46556. doi: 10.18632/oncotarget.10234. cited by applicant

Choudhury et al., CRISPR/Cas9 recombineering-mediated deep mutational scanning of essential genes in *Escherichia coli*. *Mol Syst Biol.* Mar. 2020;16(3):e9265. doi: 10.15252/msb.20199265. cited by applicant

Choudhury et al., Engineering RNA endonucleases with customized sequence specificities. *Nat Commun.* 2012;3:1147. doi: 10.1038/ncomms2154. cited by applicant

Chouluka et al., Induction of homologous recombination in mammalian chromosomes by using the I-SceI system of *Saccharomyces cerevisiae*. *Mol Cell Biol.* Apr. 1995;15(4):1968-73. doi: 10.1128/MCB.15.4.1968. cited by applicant

Christian et al., Targeting G with TAL effectors: a comparison of activities of TALENs constructed with NN and NK repeat variable di-residues. *PLoS One.* 2012;7(9):e45383. doi: 10.1371/journal.pone.0045383. Epub Sep. 24, 2012. cited by applicant

Christian et al., Targeting DNA double-strand breaks with TAL effector nucleases. *Genetics.* Oct. 2010;186(2):757-61. Doi: 10.1534/genetics.110.120717. Epub Jul. 26, 2010. cited by applicant

Christiansen et al., Characterization of the lactococcal temperate phage TP901-1 and its site-specific integration. *J Bacteriol.* Feb. 1994;176(4):1069-76. doi: 10.1128/jb.176.4.1069-1076.1994. cited by applicant

Chu et al., Increasing the efficiency of homology-directed repair for CRISPR-Cas9-induced precise gene editing in mammalian cells. *Nat Biotech.* Feb. 13, 2015;33:543-8. doi: 10.1038/nbt.3198. Epub Mar. 24, 2015. cited by applicant

Chuai et al., DeepCRISPR: optimized CRISPR guide RNA design by deep learning. *Genome Biol.* Jun. 26, 2018;19(1):80. doi: 10.1186/s13059-018-1459-4. cited by applicant

Chuai et al., In Silico Meets In Vivo: Towards Computational CRISPR-Based sgRNA Design. *Trends Biotechnol.* Jan. 2017;35(1):12-21. doi: 10.1016/j.tibtech.2016.06.008. Epub Jul. 11, 2016. cited by applicant

Chuang et al., Novel Heterotypic Rox Sites for Combinatorial Dre Recombination Strategies. *G3 (Bethesda).* Dec. 29, 2015;6(3):559-71. doi: 10.1534/g3.115.025841. cited by applicant

Chujo et al., Trmt61B is a methyltransferase responsible for 1-methyladenosine at position 58 of human mitochondrial tRNAs. *RNA.* Dec. 2012;18(12):2269-76. doi: 10.1261/rna.035600.112. Epub Oct. 24, 2012. cited by applicant

Chung-II et al., Artificial control of gene expression in mammalian cells by modulating RNA interference through aptamer-small molecule interaction. *RNA.* May 2006;12(5):710-6. Epub Apr. 10, 2006. cited by applicant

Chylinski et al., The tracrRNA and Cas9 families of type II CRISPR-Cas immunity systems. *RNA Biol.* May 2013;10(5):726-37. doi: 10.4161/rna.24321. Epub Apr. 5, 2013. cited by applicant

Clackson et al., Redesigning an FKBP-ligand interface to generate chemical dimerizers with novel specificity. *Proc Natl Acad Sci U S A.* Sep. 1, 1998;95(18):10437-42. doi: 10.1073/pnas.95.18.10437. cited by applicant

Clement et al., CRISPResso2 provides accurate and rapid genome editing sequence analysis. *Nat Biotechnol.* Mar.

2019;37(3):224-226. doi: 10.1038/s41587-019-0032-3. cited by applicant

Cobb et al., Directed evolution as a powerful synthetic biology tool. *Methods*. Mar. 15, 2013;60(1):81-90. doi: 10.1016/j.ymeth.2012.03.009. Epub Mar. 23, 2012. cited by applicant

Coffey et al., The Economic Impact of BSE on the U.S. Beef Industry: Product Value Losses, Regulatory Costs, and Consumer Reactions. Kansas State University Agricultural Experiment Station and Cooperative Extension Service. MF-2678. May 2005. 68 pages. Accessed via <https://bookstore.ksre.ksu.edu/pubs/MF2678.pdf>. cited by applicant

Cokol et al., Finding nuclear localization signals. *EMBO Rep*. Nov. 2000;1(5):411-5. doi: 10.1093/embo-reports/kvd092. cited by applicant

Cole et al., Reconstructing evolutionary adaptive paths for protein engineering. *Methods Mol Biol*. 2013;978:115-25. doi: 10.1007/978-1-62703-293-3\_8. cited by applicant

Cole-Strauss et al., Correction of the mutation responsible for sickle cell anemia by an RNA-DNA oligonucleotide. *Science*. Sep. 6, 1996;273(5280):1386-9. cited by applicant

Collinge, Prion diseases of humans and animals: their causes and molecular basis. *Annu Rev Neurosci*. 2001;24:519-50. doi: 10.1146/annurev.neuro.24.1.519. cited by applicant

Cong et al., Multiplex genome engineering using CRISPR/Cas systems. *Science*. Feb. 15, 2013;339(6121):819-23. doi: 10.1126/science.1231143. Epub Jan. 3, 2013. cited by applicant

Conrad et al., A Kaposi's sarcoma virus RNA element that increases the nuclear abundance of intronless transcripts. *EMBO J*. May 18, 2005;24(10):1831-41. doi: 10.1038/sj.emboj.7600662. Epub Apr. 28, 2005. cited by applicant

Conticello, The AID/APOBEC family of nucleic acid mutators. *Genome Biol*. 2008;9(6):229. doi: 10.1186/GB-2008-9-6-229. Epub Jun. 17, 2008. cited by applicant

Cornu et al., Refining strategies to translate genome editing to the clinic. *Nat Med*. Apr. 3, 2017;23(4):415-423. doi: 10.1038/nm.4313. cited by applicant

Costa et al., Frequent use of the same tertiary motif by self-folding RNAs. *EMBO J*. Mar. 15, 1995;14(6):1276-85. cited by applicant

Cotton et al., Insertion of a Synthetic Peptide into a Recombinant Protein Framework: A Protein Biosensor. *J. Am. Chem. Soc*. Jan. 22, 1999; 121(5):1100-1. <https://doi.org/10.1021/ja983804b>. cited by applicant

Covino et al., The CCL2/CCR2 Axis in the Pathogenesis of HIV-1 Infection: A New Cellular Target for Therapy? *Current Drug Targets* Dec. 2016;17(1):76-110. DOI : 10.2174/138945011701151217110917. cited by applicant

Cox et al., An SCN9A channelopathy causes congenital inability to experience pain. *Nature*. Dec. 14, 2006;444(7121):894-8. doi: 10.1038/nature05413. cited by applicant

Cox et al., Conditional gene expression in the mouse inner ear using Cre-loxP. *J Assoc Res Otolaryngol*. Jun. 2012;13(3):295-322. doi: 10.1007/s10162-012-0324-5. Epub Apr. 24, 2012. cited by applicant

Cox et al., Congenital insensitivity to pain: novel SCN9A missense and in-frame deletion mutations. *Hum Mutat*. Sep. 2010;31(9):E1670-86. doi: 10.1002/humu.21325. cited by applicant

Cox et al., RNA editing with CRISPR-Cas13. *Science*. Nov. 24, 2017;358(6366):1019-1027. doi: 10.1126/science.aag0180. Epub Oct. 25, 2017. cited by applicant

Cox et al., Therapeutic genome editing: prospects and challenges. *Nat Med*. Feb. 2015;21(2):121-31. doi: 10.1038/nm.3793. cited by applicant

Cox, Proteins pinpoint double strand breaks. *Elife*. Oct. 29, 2013;2:e01561. doi: 10.7554/eLife.01561. cited by applicant

Crabtree et al., Three-part inventions: intracellular signaling and induced proximity. *Trends Biochem Sci*. Nov. 1996;21(11):418-22. doi: 10.1016/s0968-0004(96)20027-1. cited by applicant

Cradick et al., CRISPR/Cas9 systems targeting  $\beta$ -globin and CCR5 genes have substantial off-target activity. *Nucleic Acids Res*. Nov. 1, 2013;41(20):9584-92. doi: 10.1093/nar/gkt714. Epub Aug. 11, 2013. cited by applicant

Cradick et al., ZFN-site searches genomes for zinc finger nuclease target sites and off-target sites. *BMC Bioinformatics*. May 13, 2011;12:152. doi: 10.1186/1471-2105-12-152. cited by applicant

Cradick et al., Zinc-finger nucleases as a novel therapeutic strategy for targeting hepatitis B virus DNAs. *Mol Ther*. May 2010;18(5):947-54. Doi: 10.1038/mt.2010.20. Epub Feb. 16, 2010. cited by applicant

Crick, On protein synthesis. *Symp Soc Exp Biol*. 1958;12:138-63. cited by applicant

Cronican et al., A class of human proteins that deliver functional proteins into mammalian cells in vitro and in vivo. *Chem Biol*. Jul. 29, 2011;18(7):833-8. doi: 10.1016/j.chembiol.2011.07.003. cited by applicant

Cronican et al., Potent delivery of functional proteins into Mammalian cells in vitro and in vivo using a supercharged protein. *ACS Chem Biol*. Aug. 20, 2010;5(8):747-52. doi:10.1021/cb1001153. cited by applicant

Crystal, Transfer of genes to humans: early lessons and obstacles to success. *Science*. Oct. 20, 1995;270(5235):404-10. doi: 10.1126/science.270.5235.404. cited by applicant

Cui et al., Consequences of Cas9 cleavage in the chromosome of *Escherichia coli*. *Nucleic Acids Res*. May 19, 2016;44(9):4243-51. doi: 10.1093/nar/gkw223. Epub Apr. 8, 2016. cited by applicant

Cui et al., m6A RNA Methylation Regulates the Self-Renewal and Tumorigenesis of Glioblastoma Stem Cells. *Cell Rep*. Mar. 14, 2017;18(11):2622-2634. doi: 10.1016/j.celrep.2017.02.059. cited by applicant

Cui et al., Review of CRISPR/Cas9 sgRNA Design Tools. *Interdiscip Sci*. Jun. 2018;10(2):455-465. doi: 10.1007/s12539-018-0298-z. Epub Apr. 11, 2018. cited by applicant

Cui et al., Targeted integration in rat and mouse embryos with zinc-finger nucleases. *Nat Biotechnol*. Jan. 2011;29(1):64-

7. Doi: 10.1038/nbt.1731. Epub Dec. 12, 2010. cited by applicant

Cunningham et al., Ensembl 2015. Nucleic Acids Res. Jan. 2015;43(Database issue):D662-9. doi: 10.1093/nar/gku1010. Epub Oct. 28, 2014. cited by applicant

Cupples et al., A set of lacZ mutations in *Escherichia coli* that allow rapid detection of each of the six base substitutions. Proc Natl Acad Sci U S A. Jul. 1989;86(14):5345-9. cited by applicant

D'Adda di Fagagna et al., The Gam protein of bacteriophage Mu is an orthologue of eukaryotic Ku. EMBO Rep. Jan. 2003;4(1):47-52. cited by applicant

Dahlem et al., Simple methods for generating and detecting locus-specific mutations induced with TALENs in the zebrafish genome. PLoS Genet. 2012;8(8):e1002861. doi: 10.1371/journal.pgen.1002861. Epub Aug. 16, 2012. cited by applicant

Dahlgren et al., A novel mutation in ribosomal protein S4 that affects the function of a mutated RF1. Biochimie. Aug. 2000;82(8):683-91. cited by applicant

Dahlman et al., Orthogonal gene knockout and activation with a catalytically active Cas9 nuclease. Nat Biotechnol. Nov. 2015;33(11):1159-61. doi: 10.1038/nbt.3390. cited by applicant

Dandage et al., beditor: A Computational Workflow for Designing Libraries of Guide RNAs for CRISPR-Mediated Base Editing. Genetics. Jun. 2019;212(2):377-385. doi: 10.1534/genetics.119.302089. Epub Apr. 1, 2019. cited by applicant

Dang et al., Optimizing sgRNA structure to improve CRISPR-Cas9 knockout efficiency. Genome Biol. Dec. 15, 2015;16:280. doi: 10.1186/s13059-015-0846-3. cited by applicant

Das et al., The crystal structure of the monomeric reverse transcriptase from Moloney murine leukemia virus. Structure. May 2004;12(5):819-29. doi: 10.1016/j.str.2004.02.032. cited by applicant

Dassa et al., Fractured genes: a novel genomic arrangement involving new split inteins and a new homing endonuclease family. Nucleic Acids Res. May 2009;37(8):2560-73. doi: 10.1093/nar/gkp095. Epub Mar. 5, 2009. cited by applicant

Dassa et al., Trans protein splicing of cyanobacterial split inteins in endogenous and exogenous combinations. Biochemistry. Jan. 9, 2007;46(1):322-30. doi: 10.1021/bi0611762. cited by applicant

Database EBI Accession No. ADE34233 Jan. 29, 2004. cited by applicant

Database EBI Accession No. BFF09785. May 31, 2018. 2 pages. cited by applicant

Database EBI Accession No. BGE38086. Jul. 25, 2019. 2 pages. cited by applicant

Database UniProt Accession No. G8I3E0. Jan. 14, 2012. cited by applicant

Datsenko et al., One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. Proc Natl Acad Sci U S A. Jun. 6, 2000;97(12):6640-5. cited by applicant

Davidson et al., Viral vectors for gene delivery to the nervous system. Nat Rev Neurosci. May 2003;4(5):353-64. doi: 10.1038/nrn1104. cited by applicant

Davis et al., DNA double strand break repair via non-homologous end-joining. Transl Cancer Res. Jun. 2013;2(3):130-143. cited by applicant

Davis et al., Small molecule-triggered Cas9 protein with improved genome-editing specificity. Nat Chem Biol. May 2015;11(5):316-8. doi: 10.1038/nchembio.1793. Epub Apr. 6, 2015. cited by applicant

De Felipe et al., Co-translational, intraribosomal cleavage of polypeptides by the foot-and-mouth disease virus 2A peptide. J Biol Chem. Mar. 28, 2003;278(13):11441-8. doi: 10.1074/jbc.M211644200. Epub Jan. 8, 2003. cited by applicant

De La Peña et al., The Hammerhead Ribozyme: A Long History for a Short RNA. Molecules. Jan. 4, 2017;22(1):78. doi: 10.3390/molecules22010078. cited by applicant

De Souza, Primer: genome editing with engineered nucleases. Nat Methods. Jan. 2012;9(1):27. cited by applicant

De Wit et al., The Human CD4+ T Cell Response against Mumps Virus Targets a Broadly Recognized Nucleoprotein Epitope. J Virol. Mar. 5, 2019;93(6):e01883-18. doi: 10.1128/JVI.01883-18. cited by applicant

Dean et al., Genetic restriction of HIV-1 infection and progression to AIDS by a deletion allele of the CKR5 structural gene. Hemophilia Growth and Development Study, Multicenter AIDS Cohort Study, Multicenter Hemophilia Cohort Study, San Francisco City Cohort, ALIVE Study. Science. Sep. 27, 1996;273(5283):1856-62. doi: 10.1126/science.273.5283.1856. cited by applicant

DeKosky et al., Large-scale sequence and structural comparisons of human naive and antigen-experienced antibody repertoires. Proc Natl Acad Sci U S A. May 10, 2016;113(19):E2636-45. doi: 10.1073/pnas.1525510113. Epub Apr. 25, 2016. cited by applicant

Delebecque et al., Organization of intracellular reactions with rationally designed RNA assemblies. Science. Jul. 22, 2011;333(6041):470-4. doi: 10.1126/science.1206938. Epub Jun. 23, 2011. cited by applicant

Deltcheva et al., CRISPR RNA maturation by trans-encoded small RNA and host factor RNase III. Nature. Mar. 31, 2011;471(7340):602-7. doi: 10.1038/nature09886. cited by applicant

Deng et al., Widespread occurrence of N6-methyladenosine in bacterial mRNA. Nucleic Acids Res. Jul. 27, 2015;43(13):6557-67. doi: 10.1093/nar/gkv596. Epub Jun. 11, 2015. cited by applicant

Denizio et al., Harnessing natural DNA modifying activities for editing of the genome and epigenome. Curr Opin Chem Biol. Aug. 2018;45:10-17. doi: 10.1016/j.cbpa.2018.01.016. Epub Feb. 13, 2018. cited by applicant

Deriano et al., Modernizing the nonhomologous end-joining repertoire: alternative and classical NHEJ share the stage. Annu Rev Genet. 2013;47:433-55. doi: 10.1146/annurev-genet-110711-155540. Epub Sep. 11, 2013. cited by applicant

Deussing, Targeted mutagenesis tools for modelling psychiatric disorders. *Cell Tissue Res.* Oct. 2013;354(1):9-25. doi: 10.1007/s00441-013-1708-5. Epub Sep. 10, 2013. cited by applicant

Dever et al., CRISPR/Cas9  $\beta$ -globin gene targeting in human haematopoietic stem cells. *Nature.* Nov. 17, 2016;539(7629):384-389. doi: 10.1038/nature20134. Epub Nov. 7, 2016. cited by applicant

Deverman et al., Cre-dependent selection yields AAV variants for widespread gene transfer to the adult brain. *Nat Biotechnol.* Feb. 2016;34(2):204-9. doi: 10.1038/nbt.3440. Epub Feb. 1, 2016. cited by applicant

Devigili et al., Paroxysmal itch caused by gain-of-function Nav1.7 mutation. *Pain.* Sep. 2014;155(9):1702-1707. doi: 10.1016/j.pain.2014.05.006. Epub May 10, 2014. cited by applicant

Dianov et al., Mammalian base excision repair: the forgotten archangel. *Nucleic Acids Res.* Apr. 1, 2013;41(6):3483-90. doi: 10.1093/nar/gkt076. Epub Feb. 13, 2013. cited by applicant

Dicarolo et al., Genome engineering in *Saccharomyces cerevisiae* using CRISPR-Cas systems. *Nucleic Acids Research* Apr. 2013;41(7):4336-43. cited by applicant

Dicarolo et al., Genome engineering in *Saccharomyces cerevisiae* using CRISPR-Cas systems. *Nucleic Acids Res.* Apr. 2013;41(7):4336-43. doi: 10.1093/nar/gkt135. Epub Mar. 4, 2013. cited by applicant

Dicarolo et al., Safeguarding CRISPR-Cas9 gene drives in yeast. *Nat Biotechnol.* Dec. 2015;33(12):1250-1255. doi: 10.1038/nbt.3412. Epub Nov. 16, 2015. cited by applicant

Dickey et al., Single-stranded DNA-binding proteins: multiple domains for multiple functions. *Structure.* Jul. 2, 2013;21(7):1074-84. doi: 10.1016/j.str.2013.05.013. cited by applicant

Dickinson et al., Experimental interrogation of the path dependence and stochasticity of protein evolution using phage-assisted continuous evolution. *Proc Natl Acad Sci USA.* May 2013;110(22):9007-12. cited by applicant

Dillon, Regulating gene expression in gene therapy. *Trends Biotechnol.* May 1993;11(5):167-73. doi: 10.1016/0167-7799(93)90109-M. cited by applicant

Ding et al., A TALEN genome-editing system for generating human stem cell-based disease models. *Cell Stem Cell.* Feb. 7, 2013;12(2):238-51. Doi: 10.1016/j.stem.2012.11.011. Epub Dec. 13, 2012. cited by applicant

Ding et al., Permanent alteration of PCSK9 with in vivo CRISPR-Cas9 genome editing. *Circ Res.* Aug. 15, 2014;115(5):488-92. doi: 10.1161/CIRCRESAHA.115.304351. Epub Jun. 10, 2014. cited by applicant

Dingwall et al., Nuclear targeting sequences—a consensus? *Trends Biochem Sci.* Dec. 1991;16(12):478-81. doi: 10.1016/0968-0004(91)90184-w. cited by applicant

Diver et al., Single-Step Synthesis of Cell-Permeable Protein Dimerizers That Activate Signal Transduction and Gene Expression. *J. Am. Chem. Soc.* Jun. 4, 1997;119(22):5106-5109. <https://doi.org/10.1021/ja963891c>. cited by applicant

Dixon et al., Reengineering orthogonally selective riboswitches. *Proc Natl Acad Sci U S A.* Feb. 16, 2010;107(7):2830-5. doi: 10.1073/pnas.0911209107. Epub Jan. 26, 2010. cited by applicant

Doench et al., Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. *Nat Biotechnol.* Feb. 2016;34(2):184-191. doi: 10.1038/nbt.3437. cited by applicant

Doench et al., Rational design of highly active sgRNAs for CRISPR-Cas9-mediated gene inactivation. *Nat Biotechnol.* Dec. 2014;32(12):1262-7. doi: 10.1038/nbt.3026. Epub Sep. 3, 2014. cited by applicant

Dolan et al., Trans-splicing with the group I intron ribozyme from *Azoarcus*. *RNA.* Feb. 2014;20(2):202-13. doi: 10.1261/rna.041012.113. Epub Dec. 16, 2013. cited by applicant

Doman et al., Evaluation and minimization of Cas9-independent off-target DNA editing by cytosine base editors. *Nat Biotechnol.* May 2020;38(5):620-628. doi: 10.1038/s41587-020-0414-6. Epub Feb. 10, 2020. cited by applicant

Dominissini et al., Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. *Nature.* Apr. 29, 2012;485(7397):201-6. doi: 10.1038/nature11112. cited by applicant

Dorgan et al., An enzyme-coupled continuous spectrophotometric assay for S-adenosylmethionine-dependent methyltransferases. *Anal Biochem.* Mar. 15, 2006;350(2):249-55. doi: 10.1016/j.ab.2006.01.004. Epub Feb. 7, 2006. cited by applicant

Dormiani et al., Long-term and efficient expression of human  $\beta$ -globin gene in a hematopoietic cell line using a new site-specific integrating non-viral system. *Gene Ther.* Aug. 2015;22(8):663-74. doi: 10.1038/gt.2015.30. Epub Apr. 1, 2015. cited by applicant

Dorr et al., Reprogramming the specificity of sortase enzymes. *Proc Natl Acad Sci U S A.* Sep. 16, 2014;111(37):13343-8. doi: 10.1073/pnas.1411179111. Epub Sep. 3, 2014. cited by applicant

Doudna et al., Genome editing. The new frontier of genome engineering with CRISPR-Cas9. *Science.* Nov. 28, 2014;346(6213):1258096. doi: 10.1126/science.1258096. cited by applicant

Doudna, The promise and challenge of therapeutic genome editing. *Nature.* Feb. 2020;578(7794):229-236. doi: 10.1038/s41586-020-1978-5. Epub Feb. 12, 2020. cited by applicant

Dove et al., Conversion of the omega subunit of *Escherichia coli* RNA polymerase into a transcriptional activator or an activation target. *Genes Dev.* Mar. 1, 1998;12(5):745-54. cited by applicant

Doyon et al., Directed evolution and substrate specificity profile of homing endonuclease I-SceI. *J Am Chem Soc.* Feb. 22, 2006;128(7):2477-84. cited by applicant

Doyon et al., Heritable targeted gene disruption in zebrafish using designed zinc-finger nucleases. *Nat Biotechnol.* Jun. 2008;26(6):702-8. Doi: 10.1038/nbt1409. Epub May 25, 2008. cited by applicant

Drake, A constant rate of spontaneous mutation in DNA-based microbes. *Proc Natl Acad Sci USA.* Aug. 15,

1991;88(16):7160-4. cited by applicant

Drost et al., Inactivation of DNA mismatch repair by variants of uncertain significance in the PMS2 gene. *Hum Mutat*. Nov. 2013;34(11):1477-80. doi: 10.1002/humu.22426. Epub Sep. 11, 2013. cited by applicant

Duan et al., Enhancement of muscle gene delivery with pseudotyped adeno-associated virus type 5 correlates with myoblast differentiation. *J Virol*. Aug. 2001;75(16):7662-71. doi: 10.1128/JVI.75.16.7662-7671.2001. cited by applicant

Dubois et al., Retroviral RNA Dimerization: From Structure to Functions. *Front Microbiol*. Mar. 22, 2018;9:527. doi: 10.3389/fmicb.2018.00527. cited by applicant

Dumas et al., Designing logical codon reassignment—Expanding the chemistry in biology. *Chem Sci*. Jan. 1, 2015;6(1):50-69. doi: 10.1039/c4sc01534g. Epub Jul. 14, 2014. Review. cited by applicant

Dunaime, Breakthrough method means CRISPR just got a lot more relevant to human health. *The Verge*. Apr. 20, 2016. <http://www.theverge.com/2016/4/20/11450262/crispr-base-editing-single-nucleotides-dna-gene-liu-harvard>. cited by applicant

Dunbar et al., Gene therapy comes of age. *Science*. Jan. 12, 2018;359(6372):eaan4672. doi: 10.1126/science.aan4672. cited by applicant

Dupuy et al., Le syndrome de De La Chapelle [De La Chapelle syndrome]. *Presse Med*. Mar. 3, 2001;30(8):369-72. French. cited by applicant

Durai et al., A bacterial one-hybrid selection system for interrogating zinc finger-DNA interactions. *Comb Chem High Throughput Screen*. May 2006;9(4):301-11. cited by applicant

Durai et al., Zinc finger nucleases: custom-designed molecular scissors for genome engineering of plant and mammalian cells. *Nucleic Acids Res*. Oct. 26, 2005;33(18):5978-90. doi: 10.1093/nar/gki912. cited by applicant

During et al., Controlled release of dopamine from a polymeric brain implant: in vivo characterization. *Ann Neurol*. Apr. 1989;25(4):351-6. cited by applicant

East-Seletsky et al., Two distinct RNase activities of CRISPR-C2c2 enable guide-RNA processing and RNA detection. *Nature* Oct. 2016;538(7624):270-3. cited by applicant

Edlund et al., Cell-specific expression of the rat insulin gene: evidence for role of two distinct 5' flanking elements. *Science*. Nov. 22, 1985;230(4728):912-6. doi: 10.1126/science.3904002. cited by applicant

Edwards et al., An *Escherichia coli* tyrosine transfer RNA is a leucine-specific transfer RNA in the yeast *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A*. Feb. 15, 1991;88(4):1153-6. cited by applicant

Edwards et al., Crystal structures of the thi-box riboswitch bound to thiamine pyrophosphate analogs reveal adaptive RNA-small molecule recognition. *Structure*. Sep. 2006;14(9):1459-68. cited by applicant

Eick et al., Robustness of Reconstructed Ancestral Protein Functions to Statistical Uncertainty. *Mol Biol Evol*. Feb. 1, 2017;34(2):247-261. doi: 10.1093/molbev/msw223. cited by applicant

Eiler et al., Structural Basis for the Fast Self-Cleavage Reaction Catalyzed by the Twister Ribozyme. *Proc Natl Acad Sci U S A*. Sep. 9, 2014;111(36):13028-33. doi: 10.1073/pnas.1414571111. Epub Aug. 25, 2014. cited by applicant

Eltoukhy et al., Nucleic acid-mediated intracellular protein delivery by lipid-like nanoparticles. *Biomaterials*. Aug. 2014;35(24):6454-61. doi: 10.1016/j.biomaterials.2014.04.014. Epub May 13, 2014. cited by applicant

Emery et al., HCN2 ion channels play a central role in inflammatory and neuropathic pain. *Science*. Sep. 9, 2011;333(6048):1462-6. doi: 10.1126/science.1206243. cited by applicant

Endo et al., Toward establishing an efficient and versatile gene targeting system in higher plants. *Biocatalysis and Agricultural Biotechnology* 2014;3(1):2-6. cited by applicant

Engel et al., The emerging role of mRNA methylation in normal and pathological behavior. *Genes Brain Behav*. Mar. 2018;17(3):e12428. doi: 10.1111/gbb.12428. Epub Nov. 17, 2017. cited by applicant

Engelward et al., Base excision repair deficient mice lacking the Aag alkyladenine DNA glycosylase. *Proc Natl Acad Sci U S A*. Nov. 25, 1997;94(24):13087-92. cited by applicant

England, Unnatural amino acid mutagenesis: a precise tool for probing protein structure and function. *Biochemistry*. Sep. 21, 2004;43(37):11623-9. cited by applicant

Enyeart et al., Biotechnological applications of mobile group II introns and their reverse transcriptases: gene targeting, RNA-seq, and non-coding RNA analysis. *Mobile DNA* 5, 2 (2014). <https://doi.org/10.1186/1759-8753-5-2>. <https://doi.org/10.1186/1759-8753-5-2>. cited by applicant

Epstein, HSV-1-based amplicon vectors: design and applications. *Gene Ther*. Oct. 2005;12 Suppl 1:S154-8. doi: 10.1038/sj.gt.3302617. cited by applicant

Eriksson et al., Recurrent de novo point mutations in lamin A cause Hutchinson-Gilford progeria syndrome. *Nature*. May 15, 2003;423(6937):293-8. doi: 10.1038/nature01629. Epub Apr. 25, 2003. PMID: 12714972. cited by applicant

Estacion et al., A sodium channel gene SCN9A polymorphism that increases nociceptor excitability. *Ann Neurol*. Dec. 2009;66(6):862-6. doi: 10.1002/ana.21895. cited by applicant

Esvelt et al., A system for the continuous directed evolution of biomolecules. *Nature*. Apr. 28, 2011;472(7344):499-503. doi: 10.1038/nature09929. Epub Apr. 10, 2011. cited by applicant

Esvelt et al., Genome-scale engineering for systems and synthetic biology. *Mol Syst Biol*. 2013;9:641. doi: 10.1038/msb.2012.66. cited by applicant

Esvelt et al., Orthogonal Cas9 proteins for RNA-guided gene regulation and editing. *Nat Methods*. Nov. 2013;10(11):1116-21. doi: 10.1038/nmeth.2681. Epub Sep. 29, 2013. cited by applicant

Evans et al., Protein trans-splicing and cyclization by a naturally split intein from the dnaE gene of *Synechocystis* species PCC6803. *J Biol Chem.* Mar. 31, 2000;275(13):9091-4. doi: 10.1074/jbc.275.13.9091. cited by applicant

Evans et al., Semisynthesis of cytotoxic proteins using a modified protein splicing element. *Protein Sci.* Nov. 1998;7(11):2256-64. doi: 10.1002/pro.5560071103. cited by applicant

Evans et al., The cyclization and polymerization of bacterially expressed proteins using modified self-splicing inteins. *J Biol Chem.* Jun. 25, 1999;274(26):18359-63. doi: 10.1074/jbc.274.26.18359. cited by applicant

Evans et al., The in vitro ligation of bacterially expressed proteins using an intein from *Methanobacterium thermoautotrophicum*. *J Biol Chem.* Feb. 12, 1999;274(7):3923-6. doi: 10.1074/jbc.274.7.3923. cited by applicant

Evers et al., CRISPR knockout screening outperforms shRNA and CRISPRi in identifying essential genes. *Nat Biotechnol.* Jun. 2016;34(6):631-3. doi: 10.1038/nbt.3536. Epub Apr. 25, 2016. cited by applicant

Fagerlund et al., The Cpf1 CRISPR-Cas protein expands genome-editing tools. *Genome Biology* Nov. 17, 2015;16:251. <https://doi.org/10.1186/s13059-015-0824-9>. cited by applicant

Falnes et al., DNA repair by bacterial AlkB proteins. *Res Microbiol.* Oct. 2003;154(8):531-8. doi: 10.1016/S0923-2508(03)00150-5. cited by applicant

Falnes et al., Repair of methyl lesions in DNA and RNA by oxidative demethylation. *Neuroscience.* Apr. 14, 2007;145(4):1222-32. doi: 10.1016/j.neuroscience.2006.11.018. Epub Dec. 18, 2006. cited by applicant

Fang et al., Synthetic Studies Towards Halichondrins: Synthesis of the Left Halves of Norhalichondrins and Homohalichondrins. *Tetrahedron Letters* 1992;33(12):1557-1560. cited by applicant

Farboud et al., Dramatic enhancement of genome editing by CRISPR/Cas9 through improved guide RNA design. *Genetics.* Apr. 2015;199(4):959-71. doi: 10.1534/genetics.115.175166. Epub Feb. 18, 2015. cited by applicant

Farhood et al., Codelivery to mammalian cells of a transcriptional factor with cis-acting element using cationic liposomes. *Anal Biochem.* Feb. 10, 1995;225(1):89-93. cited by applicant

Fawcett et al., Transposable elements controlling I-R hybrid dysgenesis in *D. melanogaster* are similar to mammalian LINEs. *Cell.* Dec. 26, 1986;47(6):1007-15. doi: 10.1016/0092-8674(86)90815-9. cited by applicant

Feldstein et al., Two sequences participating in the autolytic processing of satellite tobacco ringspot virus complementary RNA. *Gene.* Oct. 15, 1989;82(1):53-61. doi: 10.1016/0378-1119(89)90029-2. cited by applicant

Felletti et al., Twister Ribozymes as Highly Versatile Expression Platforms for Artificial Riboswitches. *Nat Commun.* Sep. 27, 2016;7:12834. doi: 10.1038/ncomms12834. cited by applicant

Feng et al., Crystal structures of the human RNA demethylase Alkbh5 reveal basis for substrate recognition. *J Biol Chem.* Apr. 25, 2014;289(17):11571-11583. doi: 10.1074/jbc.M113.546168. Epub Mar. 10, 2014. cited by applicant

Feng et al., Human L1 retrotransposon encodes a conserved endonuclease required for retrotransposition. *Cell.* Nov. 29, 1996;87(5):905-16. doi: 10.1016/s0092-8674(00)81997-2. cited by applicant

Ferretti et al., Complete genome sequence of an M1 strain of *Streptococcus pyogenes*. *Proc Natl Acad Sci U S A.* Apr. 10, 2001;98(8):4658-63. cited by applicant

Ferry et al., Rational design of inducible CRISPR guide RNAs for de novo assembly of transcriptional programs. *Nat Commun.* Mar. 3, 2017;8:14633. doi: 10.1038/ncomms14633. cited by applicant

Feuk, Inversion variants in the human genome: role in disease and genome architecture. *Genome Med.* Feb. 12, 2010;2(2):11. doi: 10.1186/gm132. cited by applicant

Filippov et al., A novel type of RNase III family proteins in eukaryotes. *Gene.* Mar. 7, 2000;245(1):213-21. doi: 10.1016/s0378-1119(99)00571-5. cited by applicant

Filippova et al., Guide RNA modification as a way to improve CRISPR/Cas9-based genome-editing systems. *Biochimie.* Dec. 2019;167:49-60. doi: 10.1016/j.biochi.2019.09.003. Epub Sep. 4, 2019. cited by applicant

Fine et al., Trans-spliced Cas9 allows cleavage of HBB and CCR5 genes in human cells using compact expression cassettes. *Scientific Reports* 2015;5(1):Article No. 10777. doi:10.1038/srep10777. With Supplementary Information. cited by applicant

Fire et al., Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature.* Feb. 19, 1998;391(6669):806-11. doi: 10.1038/35888. cited by applicant

Fischbach et al., Directed evolution can rapidly improve the activity of chimeric assembly-line enzymes. *Proc Natl Acad Sci U S A.* Jul. 17, 2007;104(29):11951-6. doi: 10.1073/pnas.0705348104. Epub Jul. 9, 2007. cited by applicant

Fischer et al., Cryptic epitopes induce high-titer humoral immune response in patients with cancer. *J Immunol.* Sep. 1, 2010;185(5):3095-102. doi: 10.4049/jimmunol.0902166. Epub Jul. 26, 2010. cited by applicant

Fitzjohn, Diversitree: comparative phylogenetic analyses of diversification in R. *Methods in Evology and Evolution.* Dec. 2012;3(6):1084-92 .doi: 10.1111/j.2041-210X.2012.00234.x. cited by applicant

Flajolet et al., Woodchuck hepatitis virus enhancer I and enhancer II are both involved in -myc2 activation in woodchuck liver tumors. *J Virol.* Jul. 1998;72(7):6175-80. doi: 10.1128/JVI.72.7.6175-6180.1998. cited by applicant

Flaman et al., A rapid PCR fidelity assay. *Nucleic Acids Res.* Aug. 11, 1994;22(15):3259-60. doi: 10.1093/nar/22.15.3259. cited by applicant

Flynn et al., CRISPR-mediated genotypic and phenotypic correction of a chronic granulomatous disease mutation in human iPS cells. *Exp Hematol.* Oct. 2015;43(10):838-848.e3. doi: 10.1016/j.exphem.2015.06.002. Epub Jun. 19, 2015. Including supplementary figures and data. cited by applicant

Fogg et al., New applications for phage integrases. *J Mol Biol.* Jul. 29, 2014;426(15):2703-16. doi:

10.1016/j.jmb.2014.05.011. Epub May 22, 2014. cited by applicant

Fogg et al., Genome Integration and Excision by a New Streptomyces Bacteriophage, ?Joe. Appl Environ Microbiol. Feb. 15, 2017;83(5):e02767-16. doi: 10.1128/AEM.02767-16. cited by applicant

Fonfara et al., Phylogeny of Cas9 determines functional exchangeability of dual-RNA and Cas9 among orthologous type II CRISPR-Cas systems. Nucleic Acids Res. Feb. 2014;42(4):2577-90. doi: 10.1093/nar/gkt1074. Epub Nov. 22, 2013. Including Supplementary Information. cited by applicant

Forster et al., Self-cleavage of virusoid RNA is performed by the proposed 55-nucleotide active site. Cell. Jul. 3, 1987;50(1):9-16. doi: 10.1016/0092-8674(87)90657-x. cited by applicant

Fortini et al., Different DNA polymerases are involved in the short- and long-patch base excision repair in mammalian cells. Biochemistry. Mar. 17, 1998;37(11):3575-80. doi: 10.1021/bi972999h. cited by applicant

Fouts et al., Sequencing Bacillus anthracis typing phages gamma and cherry reveals a common ancestry. J Bacteriol. May 2006;188(9):3402-8. doi: 10.1128/JB.188.9.3402-3408.2006. cited by applicant

Freitas et al., Mechanisms and signals for the nuclear import of proteins. Curr Genomics. Dec. 2009;10(8):550-7. doi: 10.2174/138920209789503941. cited by applicant

Freshney, Culture of Animal Cells. A Manual of Basic Technique. Alan R. Liss, Inc. New York. 1983;4. cited by applicant

Fu et al., Improving CRISPR-Cas nuclease specificity using truncated guide RNAs. Nat Biotechnol. Mar. 2014;32(3):279-84. doi: 10.1038/nbt.2808. Epub Jan. 26, 2014. cited by applicant

Fu et al., High-frequency off-target mutagenesis induced by CRISPR-Cas nucleases in human cells. Nat Biotechnol. Sep. 2013;31(9):822-6. doi: 10.1038/nbt.2623. Epub Jun. 23, 2013. cited by applicant

Fu et al., Promises and Pitfalls of Intracellular Delivery of Proteins. Bioconjugate Chemistry. Aug. 2014;25:1602-8. cited by applicant

Fuchs et al., Polyarginine as a multifunctional fusion tag. Protein Sci. Jun. 2005;14(6):1538-44. cited by applicant

Fujisawa et al., Disease-associated mutations in CIAS1 induce cathepsin B-dependent rapid cell death of human THP-1 monocytic cells. Blood. Apr. 1, 2007;109(7):2903-11. cited by applicant

Fukui et al., DNA Mismatch Repair in Eukaryotes and Bacteria. J Nucleic Acids. Jul. 27, 2010;2010. pii: 260512. doi: 10.4061/2010/260512. cited by applicant

Fung et al., Repair at single targeted DNA double-strand breaks in pluripotent and differentiated human cells. PLoS One. 2011;6(5):e20514. doi: 10.1371/journal.pone.0020514. Epub May 25, 2011. cited by applicant

Furukawa et al., In vitro selection of allosteric ribozymes that sense the bacterial second messenger c-di-GMP. Methods Mol Biol. 2014;1111:209-20. doi: 10.1007/978-1-62703-755-6\_15. cited by applicant

Fusi et al., In Silico Predictive Modeling of CRISPR/Cas9 guide efficiency. Jun. 26, 2015; bioRxiv. <http://dx.doi.org/10.1101/021568>. cited by applicant

Gaj et al., 3rd. Genome engineering with custom recombinases. Methods Enzymol. 2014;546:79-91. doi: 10.1016/B978-0-12-801185-0.00004-0. cited by applicant

Gaj et al., A comprehensive approach to zinc-finger recombinase customization enables genomic targeting in human cells. Nucleic Acids Res. Feb. 6, 2013;41(6):3937-46. cited by applicant

Gaj et al., Enhancing the specificity of recombinase-mediated genome engineering through dimer interface redesign. J Am Chem Soc. Apr. 2, 2014;136(13):5047-56. doi: 10.1021/ja4130059. Epub Mar. 20, 2014. cited by applicant

Gaj et al., Expanding the scope of site-specific recombinases for genetic and metabolic engineering. Biotechnol Bioeng. Jan. 2014;111(1):1-15. doi: 10.1002/bit.25096. Epub Sep. 13, 2013. cited by applicant

Gaj et al., Structure-guided reprogramming of serine recombinase DNA sequence specificity. Proc Natl Acad Sci U S A. Jan. 11, 2011;108(2):498-503. doi: 10.1073/pnas.1014214108. Epub Dec. 27, 2010. cited by applicant

Gaj et al., ZFN, TALEN, and CRISPR/Cas-based methods for genome engineering. Trends Biotechnol. Jul. 2013;31(7):397-405. doi: 10.1016/j.tibtech.2013.04.004. Epub May 9, 2013. cited by applicant

Gajula, Designing an Elusive C•G?G•C Crispr Base Editor. Trends Biochem Sci. Feb. 2019;44(2):91-94. doi: 10.1016/j.tibs.2018.10.004. Epub Nov. 13, 2018. cited by applicant

Gallo et al., A novel pathogenic PSEN1 mutation in a family with Alzheimer's disease: phenotypical and neuropathological features. J Alzheimers Dis. 2011;25(3):425-31. doi: 10.3233/JAD-2011-110185. cited by applicant

Gangopadhyay et al., Precision Control of CRISPR-Cas9 Using Small Molecules and Light. Biochemistry. Jan. 29, 2019;58(4):234-244. doi: 10.1021/acs.biochem.8b01202. Epub Jan. 22, 2019. cited by applicant

Gao et al., Cationic liposome-mediated gene transfer. Gene Ther. Dec. 1995;2(10):710-22. cited by applicant

Gao et al., DNA-guided genome editing using the Natronobacterium gregoryi Argonaute. Nat Biotechnol. Jul. 2016;34(7):768-73. doi: 10.1038/nbt.3547. Epub May 2, 2016. cited by applicant

Gao et al., Self-processing of ribozyme-flanked RNAs into guide RNAs in vitro and in vivo for CRISPR-mediated genome editing. J Integr Plant Biol. Apr. 2014;56(4):343-9. doi: 10.1111/jipb.12152. Epub Mar. 6, 2014. cited by applicant

Gao et al., Treatment of autosomal dominant hearing loss by in vivo delivery of genome editing agents. Nature. Jan. 11, 2018;553(7687):217-221. doi: 10.1038/nature25164. Epub Dec. 20, 2017. cited by applicant

Gapinske et al., CRISPR-SKIP: programmable gene splicing with single base editors. Genome Biol. Aug. 15, 2018;19(1):107. doi: 10.1186/s13059-018-1482-5. cited by applicant

Garcia et al., Transglycosylation: a mechanism for RNA modification (and editing?). Bioorg Chem. Jun. 2005;33(3):229-



51. doi: 10.1016/j.bioorg.2005.01.001. Epub Feb. 23, 2005. cited by applicant

Gardlik et al., Vectors and delivery systems in gene therapy. Med Sci Monit. Apr. 2005;11(4):RA110-21. Epub Mar. 24, 2005. cited by applicant

Gariyban et al., Use of the rpoB gene to determine the specificity of base substitution mutations on the *Escherichia coli* chromosome. DNA Repair (Amst). May 13, 2003;2(5):593-608. cited by applicant

Garneau et al., The CRISPR/Cas bacterial immune system cleaves bacteriophage and plasmid DNA. Nature. Nov. 4, 2010;468(7320):67-71. doi: 10.1038/nature09523. cited by applicant

Gasiunas et al., Cas9-crRNA ribonucleoprotein complex mediates specific DNA cleavage for adaptive immunity in bacteria. Proc Natl Acad Sci U S A. Sep. 25, 2012;109(39):E2579-86. Epub Sep. 4, 2012. Supplementary materials included. cited by applicant

Gasiunas et al., RNA-dependent DNA endonuclease Cas9 of the CRISPR system: Holy Grail of genome editing? Trends Microbiol. Nov. 2013;21(11):562-7. doi: 10.1016/j.tim.2013.09.001. Epub Oct. 1, 2013. cited by applicant

Gaudelli et al., Programmable base editing of A to T to G to C in genomic DNA without DNA cleavage. Nature. Nov. 23, 2017;551(7681):464-471. doi: 10.1038/nature24644. Epub Oct. 25, 2017. Erratum in: Nature. May 2, 2018. cited by applicant

Gearing, Addgene blog. CRISPR 101: Cas9 nickase design and homology directed repair. 2018. pp. 1-12. <https://blog.addgene.org/crispr-101-cas9-nickase-design-and-homology-directed-repair>. Last retrieved online Jun. 25, 2021. cited by applicant

Gehrke et al., An APOBEC3A-Cas9 base editor with minimized bystander and off-target activities. Nat Biotechnol. Nov. 2018;36(10):977-982. doi: 10.1038/nbt.4199. Epub Jul. 30, 2018. cited by applicant

GenBank Accession No. J01600.1. Brooks et al., *E.coli* dam gene coding for DNA adenine methylase. Apr. 26, 1993. cited by applicant

GenBank Accession No. U07651.1. Lu, *Escherichia coli* K12 negative regulator of replication initiation (seqA) gene, complete cds. Jul. 19, 1994. cited by applicant

GenBank Submission; NIH/NCBI Accession No. 4UN5\_B. Anders et al., Jul. 23, 2014. 5 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. AAA66622.1. Martinelli et al., May 18, 1995. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. AGT42196. Farzadfar et al., Nov. 2, 2013. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. AIT42264.1. Hyun et al., Oct. 15, 2014. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. AKA60242.1. Tong et al., Apr. 5, 2015. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. AKQ21048.1. Gilles et al., Jul. 19, 2015. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. AKS40380.1. Nodvig et al., Aug. 2, 2015. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. APG80656.1. Burstein et al., Dec. 10, 2016. 1 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. AYD60528.1. Ram et al., Oct. 2, 2018. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. BDB43378. Zhang et al., Aug. 11, 2016. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. J04623. Kita et al., Apr. 26, 1993. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. KR710351.1. Sahni et al., Jun. 1, 2015. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_002737.1. Ferretti et al., Jun. 27, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_015683.1. Trost et al., Jul. 6, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_016782.1. Trost et al., Jun. 11, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_016786.1. Trost et al., Aug. 28, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_017053.1. Fittipaldi et al., Jul. 6, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_017317.1. Trost et al., Jun. 11, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_017861.1. Heidelberg et al., Jun. 11, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_018010.1. Lucas et al., Jun. 11, 2013. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_018721.1. Feng et al., Jun. 11, 2013. 1 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_021284.1. Ku et al., Jul. 12, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_021314.1. Zhang et al., Jul. 15, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_021846.1. Lo et al., Jul. 22, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NM\_002945.3. Weiser et al., Sep. 3, 2017. 5 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NM\_002947.4. Xiao et al., May 1, 2019. 4 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NM\_174936. Guo et al., Oct. 28, 2015. 6 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_358988.1. Hoskins et al., Jan. 11, 2017. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_472073.1. Glaser et al., Jun. 27, 2013. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_628093.1. Hsiao et al., Aug. 3, 2016. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_955579.1. Chen et al., Aug. 13, 2018. 5 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. P42212. Prasher et al., Mar. 19, 2014. 7 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. RFF81513.1. Zhou et al., Aug. 21, 2018. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. SNX31424.1. Weckx, S., Feb. 16, 2018. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. TGH57013. Xu et al., Apr. 9, 2019. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_002989955.1. No Author Listed, May 6, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_010922251.1. No Author Listed, May 15, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_011054416.1. No Author Listed, May 15, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_011284745.1. No Author Listed, May 16, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_011285506.1. No Author Listed, May 16, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_011527619.1. No Author Listed, May 16, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_012560673.1. No Author Listed, May 17, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_014407541.1. No Author Listed, May 18, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_020905136.1. No Author Listed, Jul. 25, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_023080005.1. No Author Listed, Oct. 27, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_023610282.1. No Author Listed, Nov. 27, 2013. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_030125963.1. No Author Listed, Jul. 9, 2014. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_030126706.1. No Author Listed, Jul. 9, 2014. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_031386437. No Author Listed., Sep. 23, 2019. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_031488318.1. No Author Listed., Aug. 5, 2014. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_031589969.1. Haft et al., Oct. 9, 2019. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_032460140.1. No Author Listed, Oct. 4, 2014. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_032461047.1. No Author Listed, Oct. 4, 2014. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_032462016.1. Haft et al., Oct. 4, 2014. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_032462936.1. No Author Listed, Oct. 4, 2014. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_032464890.1. No Author Listed, Oct. 4, 2014. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_038431314.1. No Author Listed, Dec. 26, 2014. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_038432938.1. No Author Listed, Dec. 26, 2014. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_038434062.1. No Author Listed, Dec. 26, 2014. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_044924278.1. Haft et al., Oct. 9, 2019. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_047338501.1. Haft et al., Oct. 9, 2019. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_048327215.1. No Author Listed, Jun. 26, 2015. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_049519324.1. No Author Listed, Jul. 20, 2015. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_060798984.1. Haft et al., Oct. 9, 2019. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_062913273.1. Haft et al., Oct. 9, 2019. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_072754838. No Author Listed., Sep. 23, 2019. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_095142515.1. No Author Listed., Sep. 23, 2019. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_118538418.1. No Author Listed., Oct. 13, 2019. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_119223642.1. No Author Listed., Oct. 13, 2019. 1 page. cited by applicant

applicant  
 GenBank Submission; NIH/NCBI, Accession No. WP\_119227726.1. No Author Listed., Oct. 13, 2019. 1 page. cited by applicant  
 GenBank Submission; NIH/NCBI, Accession No. WP\_119623382.1. No Author Listed., Oct. 13, 2019. 1 page. cited by applicant  
 GenBank Submission; NIH/NCBI, Accession No. WP\_132221894.1. No Author Listed., Sep. 23, 2019. 1 page. cited by applicant  
 GenBank Submission; NIH/NCBI, Accession No. WP\_133478044.1. Haft et al., Oct. 9, 2019. 2 pages. cited by applicant  
 GenBank Submission; NIH/NCBI, Accession No. YP\_002342100.1. Bernardini et al., Jun. 10, 2013. 2 pages. cited by applicant  
 GenBank Submission; NIH/NCBI, Accession No. YP\_002344900.1. Gundogdu et al., Mar. 19, 2014. 2 pages. cited by applicant  
 GenBank Submission; NIH/NCBI, Accession No. YP\_009137104.1. Davison, Aug. 13, 2018. 2 pages. cited by applicant  
 GenBank Submission; NIH/NCBI, Accession No. YP\_009283008.1. Bernardini et al., Sep. 23, 2016. 2 pages. cited by applicant  
 GenBank Submission; NIH/NCBI, Accession No. YP\_820832.1. Makarova et al., Aug. 27, 2013. 2 pages. cited by applicant  
 George et al., Adenosine deaminases acting on RNA, RNA editing, and interferon action. J Interferon Cytokine Res. Jan. 2011;31(1):99-117. doi: 10.1089/jir.2010.0097. Epub Dec. 23, 2010. PMID: 21182352; PMCID: PMC3034097. cited by applicant  
 Gerard et al., Influence on stability in *Escherichia coli* of the carboxy-terminal structure of cloned Moloney murine leukemia virus reverse transcriptase. DNA. Aug. 1986;5(4):271-9. doi: 10.1089/dna.1986.5.271. cited by applicant  
 Gerard et al., Purification and characterization of the DNA polymerase and RNase H activities in Moloney murine sarcoma-leukemia virus. J Virol. Apr. 1975;15(4):785-97. doi: 10.1128/JVI.15.4.785-797.1975. cited by applicant  
 Gerard et al., The role of template-primer in protection of reverse transcriptase from thermal inactivation. Nucleic Acids Res. Jul. 15, 2002;30(14):3118-29. doi: 10.1093/nar/gkf417. cited by applicant  
 Gerber et al., An adenosine deaminase that generates inosine at the wobble position of tRNAs. Science. Nov. 5, 1999;286(5442):1146-9. doi: 10.1126/science.286.5442.1146. cited by applicant  
 Gerber et al., RNA editing by base deamination: more enzymes, more targets, new mysteries. Trends Biochem Sci. Jun. 2001;26(6):376-84. cited by applicant  
 Gersbach et al., Directed evolution of recombinase specificity by split gene reassembly. Nucleic Acids Res. Jul. 2010;38(12):4198-206. doi: 10.1093/nar/gkq125. Epub Mar. 1, 2010. cited by applicant  
 Gersbach et al., Targeted plasmid integration into the human genome by an engineered zinc-finger recombinase. Nucleic Acids Res. Sep. 1, 2011;39(17):7868-78. doi: 10.1093/nar/gkr421. Epub Jun. 7, 2011. cited by applicant  
 Ghahfarokhi et al., Blastocyst Formation Rate and Transgene Expression are Associated with Gene Insertion into Safe and Non-Safe Harbors in the Cattle Genome. Sci Rep. Nov. 13, 2017;7(1):15432. doi: 10.1038/s41598-017-15648-3. cited by applicant  
 Gibson et al., Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat Methods. May 2009;6(5):343-5. doi: 10.1038/nmeth.1318. Epub Apr. 12, 2009. cited by applicant  
 Gil, Position-dependent sequence elements downstream of AAUAAA are required for efficient rabbit beta-globin mRNA 3' end formation. Cell. May 8, 1987;49(3):399-406. doi: 10.1016/0092-8674(87)90292-3. cited by applicant  
 Gilbert et al., CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes. Cell. 2013 154(2):442-51. cited by applicant  
 Gilleron et al., Image-based analysis of lipid nanoparticle-mediated siRNA delivery, intracellular trafficking and endosomal escape. Nat Biotechnol. Jul. 2013;31(7):638-46. doi: 10.1038/nbt.2612. Epub Jun. 23, 2013. cited by applicant  
 Glasgow et al., DNA-binding properties of the Hin recombinase. J Biol Chem. Jun. 15, 1989;264(17):10072-82. cited by applicant  
 Glassner et al., Generation of a strong mutator phenotype in yeast by imbalanced base excision repair. Proc Natl Acad Sci U S A. Aug. 18, 1998;95(17):9997-10002. cited by applicant  
 Goldberg et al., Epigenetics: a landscape takes shape. Cell. Feb. 23, 2007;128(4):635-8. doi: 10.1016/j.cell.2007.02.006. cited by applicant  
 Goldberg et al., Loss-of-function mutations in the Nav1.7 gene underlie congenital indifference to pain in multiple human populations. Clin Genet. Apr. 2007;71(4):311-9. doi: 10.1111/j.1399-0004.2007.00790.x. cited by applicant  
 Gong et al., Active DNA demethylation by oxidation and repair. Cell Res. Dec. 2011;21(12):1649-51. doi: 10.1038/cr.2011.140. Epub Aug. 23, 2011. cited by applicant  
 Gonzalez et al., An iCRISPR platform for rapid, multiplexable, and inducible genome editing in human pluripotent stem cells. Cell Stem Cell. Aug. 7, 2014;15(2):215-26. doi: 10.1016/j.stem.2014.05.018. Epub Jun. 12, 2014. cited by applicant  
 Goodnough et al., Development of a delivery vehicle for intracellular transport of botulinum neurotoxin antagonists. FEBS Lett. Feb. 27, 2002;513(2-3):163-8. cited by applicant  
 Gordley et al., Evolution of programmable zinc finger-recombinases with activity in human cells. J Mol Biol. Mar. 30, 2007;367(3):802-13. Epub Jan. 12, 2007. cited by applicant

Gordley et al., Comprehensive integrases. *Proc Natl Acad Sci U S A*. Mar. 31, 2009;106(13):5053-8. doi: 10.1073/pnas.0812502106. Epub Mar. 12, 2009. cited by applicant

Gou et al., Designing single guide RNA for CRISPR-Cas9 base editor by deep learning. Peer reviewed Thesis/Dissertation. UCLA Electronic Theses and Dissertations. Jan. 1, 2019. Retrieved from the Internet via <https://escholarship.org/uc/item/7vf9z54t>. Last accessed on Apr. 29, 2021. cited by applicant

Grainge et al., The integrase family of recombinase: organization and function of the active site. *Mol Microbiol*. Aug. 1999;33(3):449-56. cited by applicant

Gregory et al., Integration site for *Streptomyces* phage phiBT1 and development of site-specific integrating vectors. *J Bacteriol*. Sep. 2003;185(17):5320-3. doi: 10.1128/jb.185.17.5320-5323.2003. cited by applicant

Griffiths, Endogenous retroviruses in the human genome sequence. *Genome Biol*. 2001;2(6):Reviews1017. doi: 10.1186/GB-2001-2-6-reviews1017. Epub Jun. 5, 2001. cited by applicant

Grindley et al., Mechanisms of site-specific recombination. *Annu Rev Biochem*. 2006;75:567-605. doi: 10.1146/annurev.biochem.73.011303.073908. cited by applicant

Grishok et al., Genes and Mechanisms Related to RNA Interference Regulate Expression of the Small Temporal RNAs that Control *C. elegans* Developmental Timing. *Jul. 13, 2001*;106(1):P23-4. cited by applicant

Groher et al., Synthetic riboswitches—A tool comes of age. *Biochim Biophys Acta*. Oct. 2014;1839(10):964-973. doi: 10.1016/j.bbagr.2014.05.005. Epub May 17, 2014. cited by applicant

Groth et al., Construction of transgenic *Drosophila* by using the site-specific integrase from phage phiC31. *Genetics*. Apr. 2004;166(4):1775-82. doi: 10.1534/genetics.166.4.1775. cited by applicant

Groth et al., Phage integrases: biology and applications. *J Mol Biol*. Jan. 16, 2004;335(3):667-78. cited by applicant

Gruber et al., Strategies for measuring evolutionary conservation of RNA secondary structures. *BMC Bioinformatics*. Feb. 26, 2008;9:122. doi: 10.1186/1471-2105-9-122. cited by applicant

Gruber et al., The Vienna RNA websuite. *Nucleic Acids Res*. Jul. 1, 2008;36(Web Server issue):W70-4. doi: 10.1093/nar/gkn188. Epub Apr. 19, 2008. cited by applicant

Grunebaum et al., Recent advances in understanding and managing adenosine deaminase and purine nucleoside phosphorylase deficiencies. *Curr Opin Allergy Clin Immunol*. Dec. 2013;13(6):630-8. doi: 10.1097/ACI.0000000000000006. cited by applicant

Grünwald et al., Transcriptome-wide off-target RNA editing induced by CRISPR-guided DNA base editors. *Nature*. May 2019;569(7756):433-437. doi: 10.1038/s41586-019-1161-z. Epub Apr. 17, 2019. cited by applicant

Guedon et al., Current gene therapy using viral vectors for chronic pain. *Mol Pain*. May 13, 2015;11:27. doi: 10.1186/s12990-015-0018-1. cited by applicant

Guilinger et al., Broad specificity profiling of TALENs results in engineered nucleases with improved DNA-cleavage specificity. *Nat Methods*. Apr. 2014;11(4):429-35. doi: 10.1038/nmeth.2845. Epub Feb. 16, 2014. cited by applicant

Guilinger et al., Fusion of catalytically inactive Cas9 to FokI nuclease improves the specificity of genome modification. *Nat Biotechnol*. Jun. 2014;32(6):577-82. doi: 10.1038/nbt.2909. Epub Apr. 25, 2014. cited by applicant

Gumulya et al., Exploring the past and the future of protein evolution with ancestral sequence reconstruction: the 'retro' approach to protein engineering. *Biochem J*. Jan. 1, 2017;474(1):1-19. doi: 10.1042/BCJ20160507. cited by applicant

Guo et al., Evolution of Tetrahymena ribozyme mutants with increased structural stability. *Nat Struct Biol*. Nov. 2002;9(11):855-61. doi: 10.1038/nsb850. cited by applicant

Guo et al., Facile functionalization of FK506 for biological studies by the thiol-ene 'click' reaction. *RSC Advances*. 2014;22:11400-3. cited by applicant

Guo et al., Protein tolerance to random amino acid change. *Proc Natl Acad Sci U S A*. Jun. 22, 2004;101(25):9205-10. Epub Jun. 14, 2004. cited by applicant

Guo et al., Structure of Cre recombinase complexed with DNA in a site-specific recombination synapse. *Nature*. Sep. 4, 1997;389(6646):40-6. cited by applicant

Gupta et al., Cross-talk between cognate and noncognate RpoE sigma factors and Zn(2+)-binding anti-sigma factors regulates photooxidative stress response in *Azospirillum brasilense*. *Antioxid Redox Signal*. Jan. 1, 2014;20(1):42-59. doi: 10.1089/ars.2013.5314. Epub Jul. 19, 2013. cited by applicant

Gupta et al., Sequences in attB that affect the ability of phiC31 integrase to synapse and to activate DNA cleavage. *Nucleic Acids Res*. 2007;35(10):3407-19. doi: 10.1093/nar/gkm206. Epub May 3, 2007. cited by applicant

Guzman et al., Tight regulation, modulation, and high-level expression by vectors containing the arabinose PBAD promoter. *J Bacteriol*. 1995;177(14):4121-4130. cited by applicant

Haapaniemi et al., CRISPR-Cas9 genome editing induces a p53-mediated DNA damage response. *Nat Med*. Jul. 2018;24(7):927-930. doi: 10.1038/s41591-018-0049-z. Epub Jun. 11, 2018. cited by applicant

Haddada et al., Gene therapy using adenovirus vectors. *Curr Top Microbiol Immunol*. 1995;199 ( Pt 3):297-306. doi: 10.1007/978-3-642-79586-2\_14. cited by applicant

Haeussler et al., Evaluation of off-target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. *Genome Biol*. Jul. 5, 2016;17(1):148. doi: 10.1186/s13059-016-1012-2. cited by applicant

Halbert et al., Repeat transduction in the mouse lung by using adeno-associated virus vectors with different serotypes. *J Virol*. Feb. 2000;74(3):1524-32. doi: 10.1128/jvi.74.3.1524-1532.2000. cited by applicant

Hale et al., RNA-guided RNA cleavage by a CRISPR RNA-Cas protein complex. *Cell*. Nov. 25, 2009;139(5):945-56. doi:

10.1016/j.cell.2009.07.040. cited by applicant

Halmai et al., Targeted CRISPR/dCas9-mediated reactivation of epigenetically silenced genes suggests limited escape from the inactive X chromosome. 2nd Intl Conf on Epigenetics and Bioengineering. Oct. 4, 2018; Retrieved from the Internet: <https://aiche.confex.com/aiche/epibio18/webprogram/paper544785.html>. Retrieved Jun. 29, 2020. cited by applicant

Halperin et al., CRISPR-guided DNA polymerases enable diversification of all nucleotides in a tunable window. *Nature*. Aug. 2018;560(7717):248-252. doi: 10.1038/s41586-018-0384-8. Epub Aug. 1, 2018. cited by applicant

Halvas et al., Role of murine leukemia virus reverse transcriptase deoxyribonucleoside triphosphate-binding site in retroviral replication and in vivo fidelity. *J Virol*. Nov. 2000;74(22):10349-58. doi: 10.1128/jvi.74.22.10349-10358.2000. cited by applicant

Hamano-Takaku et al., A mutant *Escherichia coli* tyrosyl-tRNA synthetase utilizes the unnatural amino acid azatyrosine more efficiently than tyrosine. *J Biol Chem*. Dec. 22, 2000;275(51):40324-8. cited by applicant

Han, New CRISPR/Cas9-based Tech Edits Single Nucleotides Without Breaking DNA. *Genome Web*, Apr. 20, 2016. <https://www.genomeweb.com/gene-silencing/gene-editing/new-crisprcas9-based-tech-edits-single-nucleotides-without-breaking-dna>. cited by applicant

Handa et al., Template-assisted synthesis of adenine-mutagenized cDNA by a retroelement protein complex. *Nucleic Acids Res*. Oct. 12, 2018;46(18):9711-9725. doi: 10.1093/nar/gky620. cited by applicant

Hanson et al., Codon optimality, bias and usage in translation and mRNA decay. *Nat Rev Mol Cell Biol*. Jan. 2018;19(1):20-30. doi: 10.1038/nrm.2017.91. Epub Oct. 11, 2017. cited by applicant

Hardt et al., Missense variants in hMLH1 identified in patients from the German HNPCC consortium and functional studies. *Fam Cancer*. Jun. 2011;10(2):273-84. doi: 10.1007/s10689-011-9431-4. cited by applicant

Harms et al., Evolutionary biochemistry: revealing the historical and physical causes of protein properties. *Nat Rev Genet*. Aug. 2013;14(8):559-71. doi: 10.1038/nrg3540. cited by applicant

Harmsen et al., DNA mismatch repair and oligonucleotide end-protection promote base-pair substitution distal from a CRISPR/Cas9-induced DNA break. *Nucleic Acids Res*. Apr. 6, 2018;46(6):2945-2955. doi: 10.1093/nar/gky076. cited by applicant

Harrington et al., A thermostable Cas9 with increased lifetime in human plasma. *Nat Commun*. Nov. 10, 2017;8(1):1424. doi: 10.1038/s41467-017-01408-4. Posted May 16, 2017 as bioRxiv preprint. [Doi.org/10.1101/138867](https://doi.org/10.1101/138867). cited by applicant

Harris et al., RNA Editing Enzyme APOBEC1 and Some of Its Homologs Can Act as DNA Mutators. *Mol Cell*. Nov. 2002;10(5):1247-53. cited by applicant

Hartung et al., Correction of metabolic, craniofacial, and neurologic abnormalities in MPS I mice treated at birth with adeno-associated virus vector transducing the human alpha-L-iduronidase gene. *Mol Ther*. Jun. 2004;9(6):866-75. cited by applicant

Hartung et al., Cre mutants with altered DNA binding properties. *J Biol Chem*. Sep. 4, 1998;273(36):22884-91. cited by applicant

Hasadsri et al., Functional protein delivery into neurons using polymeric nanoparticles. *J Biol Chem*. Mar. 13, 2009;284(11):6972-81. doi: 10.1074/jbc.M805956200. Epub Jan. 7, 2009. cited by applicant

Hasegawa et al., Spontaneous mutagenesis associated with nucleotide excision repair in *Escherichia coli*. *Genes Cells*. May 2008;13(5):459-69. doi: 10.1111/j.1365-2443.2008.01185.x. cited by applicant

Hector et al., CDKL5 variants: Improving our understanding of a rare neurologic disorder. *Neurol Genet*. Dec. 15, 2017;3(6):e200. doi: 10.1212/NXG.0000000000000200. cited by applicant

Heidenreich et al., Non-homologous end joining as an important mutagenic process in cell cycle-arrested cells. *EMBO J*. May 1, 2003;22(9):2274-83. doi: 10.1093/emboj/cdg203. cited by applicant

Held et al., In vivo correction of murine hereditary tyrosinemia type I by phiC31 integrase-mediated gene delivery. *Mol Ther*. Mar. 2005;11(3):399-408. doi: 10.1016/j.ymthe.2004.11.001. cited by applicant

Heller et al., Replisome assembly and the direct restart of stalled replication forks. *Nat Rev Mol Cell Biol*. Dec. 2006;7(12):932-43. Epub Nov. 8, 2006. cited by applicant

Hendricks et al., The *S. cerevisiae* Mag1 3-methyladenine DNA glycosylase modulates susceptibility to homologous recombination. *DNA Repair (Amst)*. 2002;1(8):645-659. cited by applicant

Hermonat et al., Use of adeno-associated virus as a mammalian DNA cloning vector: transduction of neomycin resistance into mammalian tissue culture cells. *Proc Natl Acad Sci U S A*. Oct. 1984;81(20):6466-70. doi: 10.1073/pnas.81.20.6466. cited by applicant

Herschhorn et al., Retroviral reverse transcriptases. *Cell Mol Life Sci*. Aug. 2010;67(16):2717-47. doi: 10.1007/s00018-010-0346-2. Epub Apr. 1, 2010. cited by applicant

Herzig et al., A Novel Leu92 Mutant of HIV-1 Reverse Transcriptase with a Selective Deficiency in Strand Transfer Causes a Loss of Viral Replication. *J Virol*. Aug. 2015;89(16):8119-29. doi: 10.1128/JVI.00809-15. Epub May 20, 2015. cited by applicant

Hess et al., Directed evolution using dCas9-targeted somatic hypermutation in mammalian cells. *Nat Methods*. Dec. 2016;13(12):1036-1042. doi: 10.1038/nmeth.4038. Epub Oct. 31, 2016. cited by applicant

Hickford et al., Antitumour polyether macrolides: four new halichondrins from the New Zealand deep-water marine sponge *Lissodendoryx* sp. *Bioorg Med Chem*. Mar. 15, 2009;17(6):2199-203. doi: 10.1016/j.bmc.2008.10.093. Epub Nov.

19, 2008. cited by applicant

Hida et al., Directed evolution for drug and nucleic acid; delivery. *Adv Drug Deliv Rev.* Dec. 22, 2007;59(15):1562-78. Epub Aug. 28, 2007.; Review. cited by applicant

Higgs et al., Genetic complexity in sickle cell disease. *Proc Natl Acad Sci U S A.* Aug. 19, 2008;105(33):11595-6. doi: 10.1073/pnas.0806633105. Epub Aug. 11, 2008. cited by applicant

Hilbers et al., New developments in structure determination of pseudoknots. *Biopolymers.* 1998;48(2-3):137-53. doi: 10.1002(SICI)1097-0282(1998)48:2<137::AID-BIP4>3.0.CO;2-H. cited by applicant

Hill et al., Functional analysis of conserved histidines in ADP-glucose pyrophosphorylase from *Escherichia coli*. *Biochem Biophys Res Commun.* Mar. 17, 1998;244(2):573-7. cited by applicant

Hille et al., The Biology of CRISPR-Cas: Backward and Forward. *Cell.* Mar. 8, 2018;172(6):1239-1259. doi: 10.1016/j.cell.2017.11.032. cited by applicant

Hilton et al., Enabling functional genomics with genome engineering. *Genome Res.* Oct. 2015;25(10):1442-55. doi: 10.1101/gr.190124.115. cited by applicant

Hirano et al., Site-specific recombinases as tools for heterologous gene integration. *Appl Microbiol Biotechnol.* Oct. 2011;92(2):227-39. doi: 10.1007/s00253-011-3519-5. Epub Aug. 7, 2011. Review. cited by applicant

Hirano et al., Structural Basis for the Altered PAM Specificities of Engineered CRISPR-Cas9. *Mol Cell.* Mar. 17, 2016;61(6):886-94. doi: 10.1016/j.molcel.2016.02.018. cited by applicant

Hoang et al., UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Mol Biol Evol.* Feb. 1, 2018;35(2):518-522. doi: 10.1093/molbev/msx281. cited by applicant

Hockemeyer et al., Efficient targeting of expressed and silent genes in human ESCs and iPSCs using zinc-finger nucleases. *Nat Biotechnol.* Sep. 2009;27(9):851-7. doi: 10.1038/nbt.1562. Epub Aug. 13, 2009. cited by applicant

Hockemeyer et al., Genetic engineering of human pluripotent cells using TALE nucleases. *Nat Biotechnol.* Jul. 7, 2011;29(8):731-4. doi: 10.1038/nbt.1927. cited by applicant

Hoernes et al., Translating the epitranscriptome. *Wiley Interdiscip Rev RNA.* Jan. 2017;8(1):e1375. doi: 10.1002/wrna.1375. Epub Jun. 27, 2016. cited by applicant

Hoess et al., DNA specificity of the Cre recombinase resides in the 25 kDa carboxyl domain of the protein. *J Mol Biol.* Dec. 20, 1990;216(4):873-82. doi: 10.1016/S0022-2836(99)80007-2. cited by applicant

Holden et al., Crystal structure of the anti-viral APOBEC3G catalytic domain and functional implications. *Nature.* Nov. 6, 2008;456(7218):121-4. doi: 10.1038/nature07357. Epub Oct. 12, 2008. cited by applicant

Hollis et al., Phage integrases for the construction and manipulation of transgenic mammals. *Reprod Biol Endocrinol.* Nov. 7, 2003;1:79. doi: 10.1186/1477-7827-1-79. cited by applicant

Holsinger et al., Signal transduction in T lymphocytes using a conditional allele of Sos. *Proc Natl Acad Sci U S A.* Oct. 10, 1995;92(21):9810-4. doi: 10.1073/pnas.92.21.9810. cited by applicant

Holt et al., Human hematopoietic stem/progenitor cells modified by zinc-finger nucleases targeted to CCR5 control HIV-1 in vivo. *Nat Biotechnol.* Aug. 2010;28(8):839-47. doi: 10.1038/nbt.1663. Epub Jul. 2, 2010. cited by applicant

Hondares et al., Peroxisome Proliferator-activated Receptor ? (PPAR?) Induces PPAR? Coactivator 1? (PGC-1?) Gene Expression and Contributes to Thermogenic Activation of Brown Fat. *J Biol. Chem.* Oct. 2011; 286(50):43112-22. doi: 10.1074/jbc.M111.252775. cited by applicant

Hoogenboom et al., Natural and designer binding sites made by phage display technology. *Immunol Today.* Aug. 2000;21(8):371-8. cited by applicant

Horvath et al., CRISPR/Cas, the immune system of bacteria and archaea. *Science.* Jan. 8, 2010;327(5962):167-70. doi: 10.1126/science.1179555. cited by applicant

Horvath et al., Diversity, Activity, and Evolution of CRISPR Loci in *Streptococcus thermophilus*. *J Bacteriol.* Feb. 2008;190(4):1401-12. doi: 10.1128/JB.01415-07. Epub Dec. 7, 2007. cited by applicant

Hotta et al., [Neurotropic viruses—classification, structure and characteristics]. *Nihon Rinsho.* Apr. 1997;55(4):777-82. Japanese. cited by applicant

Hou et al., Efficient genome engineering in human pluripotent stem cells using Cas9 from *Neisseria meningitidis*. *Proc Natl Acad Sci U S A.* Sep. 24, 2013;110(39):15644-9. doi: 10.1073/pnas.1313587110. Epub Aug. 12, 2013. cited by applicant

Houdebine, The methods to generate transgenic animals and to control transgene expression. *J Biotechnol.* Sep. 25, 2002;98(2-3):145-60. cited by applicant

Housden et al., Identification of potential drug targets for tuberous sclerosis complex by synthetic screens combining CRISPR-based knockouts with RNAi. *Sci Signal.* Sep. 8, 2015;8(393):rs9. doi: 10.1126/scisignal.aab3729. cited by applicant

Howard et al., Intracerebral drug delivery in rats with lesion-induced memory deficits. *J Neurosurg.* Jul. 1989;71(1):105-12. cited by applicant

Hower et al., Shape-based peak identification for ChIP-Seq. *BMC Bioinformatics.* Jan. 12, 2011;12:15. doi: 10.1186/1471-2105-12-15. cited by applicant

Hsu et al., DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat Biotechnol.* Sep. 2013;31(9):827-32. doi: 10.1038/nbt.2647. Epub Jul. 21, 2013. cited by applicant

Hsu et al., DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat Biotechnol.* Sep. 2013;31(9):827-32. doi:

10.1038/nbt.2647. Epub Jul. 21, 2013. Supplementary Information. 27 pages. cited by applicant

Hu et al., Chemical Biology Approaches to Genome Editing: Understanding, Controlling, and Delivering Programmable Nucleases. *Cell Chem Biol.* Jan. 21, 2016;23(1):57-73. doi: 10.1016/j.chembiol.2015.12.009. cited by applicant

Hu et al., Evolved Cas9 variants with broad PAM compatibility and high DNA specificity. *Nature.* Apr. 5, 2018;556(7699):57-63 and Extended/Supplementary Data. doi: 10.1038/nature26155. Epub Feb. 28, 2018. 21 pages. cited by applicant

Hu et al., Evolved Cas9 variants with broad PAM compatibility and high DNA specificity. *Nature.* Apr. 5, 2018;556(7699):57-63. doi: 10.1038/nature26155. Epub Feb. 28, 2018. cited by applicant

Hua et al., Expanding the base editing scope in rice by using Cas9 variants. *Plant Biotechnol J.* Feb. 2019;17(2):499-504. doi: 10.1111/pbi.12993. Epub Oct. 5, 2018. cited by applicant

Hua et al., Precise A·T to G·C Base Editing in the Rice Genome. *Mol Plant.* Apr. 2, 2018;11(4):627-630. doi: 10.1016/j.molp.2018.02.007. Epub Feb. 21, 2018. cited by applicant

Huang et al., Circularly permuted and PAM-modified Cas9 variants broaden the targeting scope of base editors. *Nat Biotechnol.* Jun. 2019;37(6):626-631. doi: 10.1038/s41587-019-0134-y. Epub May 20, 2019. Including Supplementary Information. cited by applicant

Huang et al., Heritable gene targeting in zebrafish using customized TALENs. *Nat Biotechnol.* Aug. 5, 2011;29(8):699-700. doi: 10.1038/nbt.1939. cited by applicant

Huggins et al., Flap endonuclease 1 efficiently cleaves base excision repair and DNA replication intermediates assembled into nucleosomes. *Mol Cell.* Nov. 2002;10(5):1201-11. doi: 10.1016/s1097-2765(02)00736-0. cited by applicant

Humbert et al., Targeted gene therapies: tools, applications, optimization. *Crit Rev Biochem Mol Biol.* May-Jun. 2012;47(3):264-81. doi: 10.3109/10409238.2012.658112. cited by applicant

Hung et al., Protein localization in disease and therapy. *J Cell Sci.* Oct. 15, 2011;124(Pt 20):3381-92. doi: 10.1242/jcs.089110. cited by applicant

Hurt et al., Highly specific zinc finger proteins obtained by directed domain shuffling and cell-based selection. *Proc Natl Acad Sci U S A.* Oct. 14, 2003;100(21):12271-6. Epub Oct. 3, 2003. cited by applicant

Husimi, Selection and evolution of bacteriophages in cellstat. *Adv Biophys.* ; 1989;25:1-43. Review. cited by applicant

Hwang et al., Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat Biotechnol.* Mar. 2013;31(3):227-9. doi: 10.1038/nbt.2501. Epub Jan. 29, 2013. cited by applicant

Hwang et al., Efficient In Vivo Genome Editing Using RNA-Guided Nucleases. *Nat Biotechnol.* Mar. 2013; 31(3): 227-229. doi: 10.1038/nbt.2501. Epub Jan. 29, 2013. cited by applicant

Hwang et al., Web-based design and analysis tools for CRISPR base editing. *BMC Bioinformatics.* Dec. 27, 2018;19(1):542. doi: 10.1186/s12859-018-2585-4. cited by applicant

Ibba et al., Relaxing the substrate specificity of an aminoacyl-tRNA synthetase allows in vitro and in vivo synthesis of proteins containing unnatural amino acids. *FEBS Lett.* May 15, 1995;364(3):272-5. cited by applicant

Ibba et al., Substrate specificity is determined by amino acid binding pocket size in *Escherichia coli* phenylalanyl-tRNA synthetase. *Biochemistry.* Jun. 14, 1994;33(23):7107-12. cited by applicant

Ihry et al., p53 inhibits CRISPR-Cas9 engineering in human pluripotent stem cells. *Nat Med.* Jul. 2018;24(7):939-946. doi: 10.1038/s41591-018-0050-6. Epub Jun. 11, 2018. cited by applicant

Iida et al., A site-specific, conservative recombination system carried by bacteriophage P1. Mapping the recombinase gene *cin* and the cross-over sites *cix* for the inversion of the C segment. *EMBO J.* 1982;1(11):1445-53. cited by applicant

Iida et al., The Min DNA inversion enzyme of plasmid p15B of *Escherichia coli* 15T-: a new member of the Din family of site-specific recombinases. *Mol Microbiol.* Jun. 1990;4(6):991-7. doi: 10.1111/j.1365-2958.1990.tb00671.x. cited by applicant

Ikediodi et al., Mutation analysis of 24 known cancer genes in the NCI-60 cell line set. *Mol Cancer Ther.* Nov. 2006;5(11):2606-12. Epub Nov. 6, 2006. cited by applicant

Imanishi et al., Detection of N6-methyladenosine based on the methyl-sensitivity of MazF RNA endonuclease. *Chem Commun (Camb).* Nov. 30, 2017;53(96):12930-12933. doi: 10.1039/c7cc07699a. cited by applicant

Imburgio et al., Studies of promoter recognition and start site selection by T7 RNA polymerase using a comprehensive collection of promoter variants. *Biochemistry.* Aug. 29, 2000;39(34):10419-30. cited by applicant

Ingram, A specific chemical difference between the globins of normal human and sickle-cell anaemia haemoglobin. *Nature.* Oct. 13, 1956;178(4537):792-4. doi: 10.1038/178792a0. cited by applicant

Irion et al., Identification and targeting of the ROSA26 locus in human embryonic stem cells. *Nat Biotechnol.* Dec. 2007;25(12):1477-82. doi: 10.1038/nbt1362. Epub Nov. 25, 2007. cited by applicant

Irrthum et al., Congenital hereditary lymphedema caused by a mutation that inactivates VEGFR3 tyrosine kinase. *Am J Hum Genet.* Aug. 2000;67(2):295-301. Epub Jun. 9, 2000. cited by applicant

Isaacs et al., Engineered riboregulators enable post-transcriptional control of gene expression. *Nat Biotechnol.* Jul. 2004;22(7):841-7. doi: 10.1038/nbt986. Epub Jun. 20, 2004. cited by applicant

Ishino et al., Nucleotide sequence of the *iap* gene, responsible for alkaline phosphatase isozyme conversion in *Escherichia coli*, and identification of the gene product. *J Bacteriol.* Dec. 1987;169(12):5429-33. cited by applicant

Iwai et al., Circular beta-lactamase: stability enhancement by cyclizing the backbone. *FEBS Lett.* Oct. 8, 1999;459(2):166-72. doi: 10.1016/s0014-5793(99)01220-x. cited by applicant

Iwai et al., Highly efficient protein trans-splicing by a naturally split DnaE intein from *Nostoc punctiforme*. *FEBS Lett.* Mar. 20, 2006;580(7):1853-8. doi: 10.1016/j.febslet.2006.02.045. Epub Feb. 24, 2006. cited by applicant

Jaffrey et al., Emerging links between m6A and misregulated mRNA methylation in cancer. *Genome Med.* Jan. 12, 2017;9(1):2. doi: 10.1186/s13073-016-0395-8. cited by applicant

Jamieson et al., Drug discovery with engineered zinc-finger proteins. *Nat Rev Drug Discov.* May 2003;2(5):361-8. cited by applicant

Jansen et al., Backbone and nucleobase contacts to glucosamine-6-phosphate in the glmS ribozyme. *Nat Struct Mol Biol.* Jun. 2006;13(6):517-23. Epub May 14, 2006. cited by applicant

Jansen et al., Identification of genes that are associated with DNA repeats in prokaryotes. *Mol Microbiol.* Mar. 2002;43(6):1565-75. cited by applicant

Jardine et al., HIV-1 Vaccines. Priming a broadly neutralizing antibody response to HIV-1 using a germline-targeting immunogen. *Science.* Jul. 10, 2015;349(6244):156-61. doi: 10.1126/science.aac5894. Epub Jun. 18, 2015. cited by applicant

Jasin et al., Repair of strand breaks by homologous recombination. *Cold Spring Harb Perspect Biol.* Nov. 1, 2013;5(11):a012740. doi: 10.1101/cshperspect.a012740. cited by applicant

Jeggo, DNA breakage and repair. *Adv Genet.* 1998;38:185-218. doi: 10.1016/s0065-2660(08)60144-3. cited by applicant

Jemielity et al., Novel “anti-reverse” cap analogs with superior translational properties. *RNA.* Sep. 2003;9(9):1108-22. doi: 10.1261/rna.5430403. cited by applicant

Jenkins et al., Comparison of a preQ1 riboswitch aptamer in metabolite-bound and free states with implications for gene regulation. *J Biol Chem.* Jul. 15, 2011;286(28):24626-37. doi: 10.1074/jbc.M111.230375. Epub May 18, 2011. cited by applicant

Jeong et al., Measurement of deoxyinosine adduct: Can it be a reliable tool to assess oxidative or nitrosative DNA damage? *Toxicol Lett.* Oct. 17, 2012;214(2):226-33. doi: 10.1016/j.toxlet.2012.08.013. Epub Aug. 23, 2012. cited by applicant

Jia et al., The MLH1 ATPase domain is needed for suppressing aberrant formation of interstitial telomeric sequences. *DNA Repair (Amst).* May 2018;65:20-25. doi: 10.1016/j.dnarep.2018.03.002. Epub Mar. 7, 2018. cited by applicant

Jiang et al., CRISPR-Cas9 Structures and Mechanisms. *Annu Rev Biophys.* May 22, 2017;46:505-529. doi: 10.1146/annurev-biophys-062215-010822. Epub Mar. 30, 2017. cited by applicant

Jiang et al., RNA-guided editing of bacterial genomes using CRISPR-Cas systems. *Nat Biotechnol.* Mar. 2013;31(3):233-9. doi: 10.1038/nbt.2508. Epub Jan. 29, 2013. cited by applicant

Jiang et al., Structural Biology. A Cas9-guide RNA Complex Preorganized for Target DNA Recognition. *Science.* Jun. 26, 2015;348(6242):1477-81. doi: 10.1126/science.aab1452. cited by applicant

Jiang et al., Structures of a CRISPR-Cas9 R-loop complex primed for DNA cleavage. *Science.* Feb. 19, 2016;351(6275):867-71. doi: 10.1126/science.aad8282. Epub Jan. 14, 2016. cited by applicant

Jin et al., Cytosine, but not adenine, base editors induce genome-wide off-target mutations in rice. *Science.* Apr. 19, 2019;364(6437):292-295. doi: 10.1126/science.aaw7166. Epub Feb. 28, 2019. cited by applicant

Jinek et al., A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science.* Aug. 17, 2012;337(6096):816-21. doi: 10.1126/science.1225829. Epub Jun. 28, 2012. cited by applicant

Jinek et al., RNA-programmed genome editing in human cells. *Elife.* Jan. 29, 2013;2:e00471. doi: 10.7554/eLife.00471. cited by applicant

Jinek et al., Structures of Cas9 endonucleases reveal RNA-mediated conformational activation. *Science.* Mar. 14, 2014;343(6176):1247997. doi: 10.1126/science.1247997. Epub Feb. 6, 2014. cited by applicant

Jiricny, The multifaceted mismatch-repair system. *Nat Rev Mol Cell Biol.* May 2006;7(5):335-46. doi: 10.1038/nrm1907. cited by applicant

Johann et al., GLVR1, a receptor for gibbon ape leukemia virus, is homologous to a phosphate permease of *Neurospora crassa* and is expressed at high levels in the brain and thymus. *J Virol.* Mar. 1992;66(3):1635-40. doi: 10.1128/JVI.66.3.1635-1640.1992. cited by applicant

Johansson et al., RNA Recognition by the MS2 Phage Coat Protein. *Seminars in Virology.* 1997;8(3):176-85. <https://doi.org/10.1006/smvy.1997.0120>. cited by applicant

Johansson et al., Selenocysteine in proteins-properties and biotechnological use. *Biochim Biophys Acta.* Oct. 30, 2005;1726(1):1-13. Epub Jun. 1, 2005. cited by applicant

Johns et al., The promise and peril of continuous in vitro evolution. *J Mol Evol.* Aug. 2005;61(2):253-63. Epub Jun. 27, 2005. cited by applicant

Johnson et al., Trans insertion-splicing: ribozyme-catalyzed insertion of targeted sequences into RNAs. *Biochemistry.* Aug. 9, 2005;44(31):10702-10. doi: 10.1021/bi0504815. cited by applicant

Joho et al., Identification of a region of the bacteriophage T3 and T7 RNA polymerases that determines promoter specificity. *J Mol Biol.* Sep. 5, 1990;215(1):31-9. cited by applicant

Jore et al., Structural basis for CRISPR RNA-guided DNA recognition by Cascade. *Nat Struct Mol Biol.* May 2011;18(5):529-36. doi: 10.1038/nsmb.2019. Epub Apr. 3, 2011. cited by applicant

Joung et al., TALENs: a widely applicable technology for targeted genome editing. *Nat Rev Mol Cell Biol.* Jan. 2013;14(1):49-55. doi: 10.1038/nrm3486. Epub Nov. 21, 2012. cited by applicant



Joyce et al., Amplification, mutation and selection of catalytic RNA. *Gene*. Oct. 15, 1989;82(1):83-7. doi: 10.1016/0378-1119(89)90033-4. cited by applicant

Jusiak et al., Comparison of Integrases Identifies Bxb1-GA Mutant as the Most Efficient Site-Specific Integrase System in Mammalian Cells. *ACS Synth Biol*. Jan. 18, 2019;8(1):16-24. doi: 10.1021/acssynbio.8b00089. Epub Jan. 9, 2019. cited by applicant

Jyothy et al., Translocation Down syndrome. *Indian J Med Sci*. Mar. 2002;56(3):122-6. cited by applicant

Kacian et al., Purification of the DNA polymerase of avian myeloblastosis virus. *Biochim Biophys Acta*. Sep. 24, 1971;246(3):365-83. doi: 10.1016/0005-2787(71)90773-8. cited by applicant

Kaczmarczyk et al., Manipulating the Prion Protein Gene Sequence and Expression Levels with CRISPR/Cas9. *PLoS One*. Apr. 29, 2016;11(4):e0154604. doi: 10.1371/journal.pone.0154604. cited by applicant

Kadoch et al., Reversible disruption of mSWI/SNF (BAF) complexes by the SS18-SSX oncogenic fusion in synovial sarcoma. *Cell*. Mar. 28, 2013;153(1):71-85. doi: 10.1016/j.cell.2013.02.036. cited by applicant

Kahmann et al., G inversion in bacteriophage Mu DNA is stimulated by a site within the invertase gene and a host factor. *Cell*. Jul. 1985;41(3):771-80. doi: 10.1016/s0092-8674(85)80058-1. cited by applicant

Kaiser et al., Gene therapy. Putting the fingers on gene repair. *Science*. Dec. 23, 2005;310(5756):1894-6. cited by applicant

Kakiyama et al., A peptide release system using a photo-cleavable linker in a cell array format for cell-toxicity analysis. *Polymer J*. Feb. 27, 2013;45:535-9. cited by applicant

Kalyaanamoorthy et al., ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods*. Jun. 2017;14(6):587-589. doi: 10.1038/nmeth.4285. Epub May 8, 2017. cited by applicant

Kandavelou et al., Targeted manipulation of mammalian genomes using designed zinc finger nucleases. *Biochem Biophys Res Commun*. Oct. 9, 2009;388(1):56-61. doi: 10.1016/j.bbrc.2009.07.112. Epub Jul. 25, 2009. cited by applicant

Kang et al., Structural Insights into riboswitch control of the biosynthesis of queuosine, a modified nucleotide found in the anticodon of tRNA. *Mol Cell*. Mar. 27, 2009;33(6):784-90. doi: 10.1016/j.molcel.2009.02.019. Epub Mar. 12, 2009. cited by applicant

Kang et al., Precision genome engineering through adenine base editing in plants. *Nat Plants*. Jul. 2018;4(7):427-431. doi: 10.1038/s41477-018-0178-x. Epub Jun. 4, 2018. Erratum in: *Nat Plants*. Sep. 2018;4(9):730. cited by applicant

Kao et al., Cleavage specificity of *Saccharomyces cerevisiae* flap endonuclease 1 suggests a double-flap structure as the cellular substrate. *J Biol Chem*. Apr. 26, 2002;277(17):14379-89. doi: 10.1074/jbc.M110662200. Epub Feb. 1, 2002. cited by applicant

Kappel et al., Regulating gene expression in transgenic animals. *Curr Opin Biotechnol*. Oct. 1992;3(5):548-53. cited by applicant

Karimova et al., Discovery of Nigri/nox and Panto/pox site-specific recombinase systems facilitates advanced genome engineering. *Sci Rep*. Jul. 22, 2016;6:30130. doi: 10.1038/srep30130. cited by applicant

Karimova et al., Vika/vox, a novel efficient and specific Cre/loxP-like site-specific recombination system. *Nucleic Acids Res*. Jan. 2013;41(2):e37. doi: 10.1093/nar/gks1037. Epub Nov. 9, 2012. cited by applicant

Karpenshif et al., From yeast to mammals: recent advances in genetic control of homologous recombination. *DNA Repair (Amst)*. Oct. 1, 2012;11(10):781-8. doi: 10.1016/j.dnarep.2012.07.001. Epub Aug. 11, 2012. Review. cited by applicant

Karpinsky et al., Directed evolution of a recombinase that excises the provirus of most HIV- 1 primary isolates with high specificity. *Nat Biotechnol*. Apr. 2016;34(4):401-9. doi: 10.1038/nbt.3467. Epub Feb. 22, 2016. cited by applicant

Katafuchi et al., DNA polymerases involved in the incorporation of oxidized nucleotides into DNA: their efficiency and template base preference. *Mutat Res*. Nov. 28, 2010;703(1):24-31. doi: 10.1016/j.mrgentox.2010.06.004. Epub Jun. 11, 2010. cited by applicant

Kato et al., Improved purification and enzymatic properties of three forms of reverse transcriptase from avian myeloblastosis virus. *J Virol Methods*. Dec. 1984;9(4):325-39. doi: 10.1016/0166-0934(84)90058-2. cited by applicant

Katoh et al., MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol*. Apr. 2013;30(4):772-80. doi: 10.1093/molbev/mst010. Epub Jan. 16, 2013. cited by applicant

Kaufman et al., Translational efficiency of polycistronic mRNAs and their utilization to express heterologous genes in mammalian cells. *EMBO J*. Jan. 1987;6(1):187-93. cited by applicant

Kavli et al., Excision of cytosine and thymine from DNA by mutants of human uracil-DNA glycosylase. *EMBO J*. Jul. 1, 1996;15(13):3442-7. cited by applicant

Kawarasaki et al., Enhanced crossover SCRATCHY: construction and high-throughput screening of a combinatorial library containing multiple non-homologous crossovers. *Nucleic Acids Res*. Nov. 1, 2003;31(21):e126. cited by applicant

Kay et al., Viral vectors for gene therapy: the art of turning infectious agents into vehicles of therapeutics. *Nat Med*. Jan. 2001;7(1):33-40. cited by applicant

Kaya et al., A bacterial Argonaute with noncanonical guide RNA specificity. *Proc. Natl. Acad. Sci. USA*. Apr. 2016;113(15):4057-62. cited by applicant

Keijzers et al., Human exonuclease 1 (EXO1) activity characterization and its function on flap structures. *Biosci Rep*. Apr. 25, 2015;35(3):e00206. doi: 10.1042/BSR20150058. cited by applicant

Kellendonk et al., Regulation of Cre recombinase activity by the synthetic steroid RU 486. *Nucleic Acids Res*. Apr. 15, 1996;24(8):1404-11. cited by applicant

Kelman, PCNA: structure, functions and interactions. *Oncogene*. Feb. 13, 1997;14(6):629-40. doi: 10.1038/sj.onc.1200886. cited by applicant

Keravala et al., A diversity of serine phage integrases mediate site-specific recombination in mammalian cells. *Mol Genet Genomics*. Aug. 2006;276(2):135-46. doi: 10.1007/s00438-006-0129-5. Epub May 13, 2006. cited by applicant

Kessel et al., Murine developmental control genes. *Science*. Jul. 27, 1990;249(4967):374-9. doi: 10.1126/science.1974085. cited by applicant

Kessler et al., Gene delivery to skeletal muscle results in sustained expression and systemic delivery of a therapeutic protein. *Proc Natl Acad Sci U S A*. Nov. 26, 1996;93(24):14082-7. doi: 10.1073/pnas.93.24.14082. cited by applicant

Ketha et al., Application of bioinformatics-coupled experimental analysis reveals a new transport-competent nuclear localization signal in the nucleoprotein of Influenza A virus strain. *BMC Cell Biol*. Apr. 28, 2008; 9:22. <https://doi.org/10.1186/1471-2121-9-22>. cited by applicant

Kiga et al., An engineered *Escherichia coli* tyrosyl-tRNA synthetase for site-specific incorporation of an unnatural amino acid into proteins in eukaryotic translation and its application in a wheat germ cell-free system. *Proc Natl Acad Sci U S A*. Jul. 23, 2002;99(15):9715-20. Epub Jul. 3, 2002. cited by applicant

Kilbride et al., Determinants of product topology in a hybrid Cre-Tn3 resolvase site-specific recombination system. *J Mol Biol*. Jan. 13, 2006;355(2):185-95. Epub Nov. 9, 2005. cited by applicant

Kilcher et al., Brochothrix thermosphacta bacteriophages feature heterogeneous and highly mosaic genomes and utilize unique prophage insertion sites. *J Bacteriol*. Oct. 2010;192(20):5441-53. doi: 10.1128/JB.00709-10. Epub Aug. 13, 2010. cited by applicant

Kim et al., DJ-1, a novel regulator of the tumor suppressor PTEN. *Cancer Cell*. 2005;7(3):263-273. cited by applicant

Kim et al., Genome-wide target specificity of CRISPR RNA-guided adenine base editors. *Nat Biotechnol*. Apr. 2019;37(4):430-435. doi: 10.1038/s41587-019-0050-1. Epub Mar. 4, 2019. cited by applicant

Kim et al., A library of TAL effector nucleases spanning the human genome. *Nat Biotechnol*. Mar. 2013;31(3):251-8. Doi: 10.1038/nbt.2517. Epub Feb. 17, 2013. cited by applicant

Kim et al., An anionic human protein mediates cationic liposome delivery of genome editing proteins into mammalian cells. *Nat Commun*. Jul. 2, 2019;10(1):2905. doi: 10.1038/s41467-019-10828-3. cited by applicant

Kim et al., Evaluating and Enhancing Target Specificity of Gene-Editing Nucleases and Deaminases. *Annu Rev Biochem*. Jun. 20, 2019;88:191-220. doi: 10.1146/annurev-biochem-013118-111730. Epub Mar. 18, 2019. cited by applicant

Kim et al., Genome-wide target specificities of CRISPR RNA-guided programmable deaminases. *Nat Biotechnol*. May 2017;35(5):475-480. doi: 10.1038/nbt.3852. Epub Apr. 10, 2017. cited by applicant

Kim et al., High cleavage efficiency of a 2A peptide derived from porcine teschovirus-1 in human cell lines, zebrafish and mice. *PLoS One*. 2011;6(4):e18556. doi: 10.1371/journal.pone.0018556. Epub Apr. 29, 2011. cited by applicant

Kim et al., High-throughput analysis of the activities of xCas9, SpCas9-NG and SpCas9 at matched and mismatched target sequences in human cells. *Nat Biomed Eng*. Jan. 2020;4(1):111-124. doi: 10.1038/s41551-019-0505-1. Epub Jan. 14, 2020. cited by applicant

Kim et al., Highly efficient RNA-guided base editing in mouse embryos. *Nat Biotechnol*. May 2017;35(5):435-437. doi: 10.1038/nbt.3816. Epub Feb. 27, 2017. cited by applicant

Kim et al., Highly efficient RNA-guided genome editing in human cells via delivery of purified Cas9 ribonucleoproteins. *Genome Res*. Jun. 2014;24(6):1012-9. doi: 10.1101/gr.171322.113. Epub Apr. 2, 2014. cited by applicant

Kim et al., In vivo genome editing with a small Cas9 orthologue derived from *Campylobacter jejuni*. *Nat Commun*. Feb. 21, 2017;8:14500. doi: 10.1038/ncomms14500. PMID: 28220790; PMCID: PMC5473640. cited by applicant

Kim et al., In vivo high-throughput profiling of CRISPR-Cpf1 activity. *Nat Methods*. Feb. 2017;14(2):153-159. doi: 10.1038/nmeth.4104. Epub Dec. 19, 2016. cited by applicant

Kim et al., Increasing the genome-targeting scope and precision of base editing with engineered Cas9-cytidine deaminase fusions. *Nat Biotechnol*. Apr. 2017;35(4):371-376. doi: 10.1038/nbt.3803. Epub Feb. 13, 2017. cited by applicant

Kim et al., Mycobacteriophage Bxb1 integrates into the *Mycobacterium smegmatis* groEL1 gene. *Mol Microbiol*. Oct. 2003;50(2):463-73. doi: 10.1046/j.1365-2958.2003.03723.x. cited by applicant

Kim et al., Rescue of high-specificity Cas9 variants using sgRNAs with matched 5' nucleotides. *Genome Biol*. Nov. 15, 2017;18(1):218. doi: 10.1186/s13059-017-1355-3. cited by applicant

Kim et al., Structural and kinetic characterization of *Escherichia coli* TadA, the wobble-specific tRNA deaminase. *Biochemistry*. May 23, 2006;45(20):6407-16. doi: 10.1021/bi0522394. PMID: 16700551. cited by applicant

Kim et al., TALENs and ZFNs are associated with different mutationsignatures. *Nat Methods*. Mar. 2013;10(3):185. doi: 10.1038/nmeth.2364. Epub Feb. 10, 2013. cited by applicant

Kim et al., Targeted genome editing in human cells with zinc finger nucleases constructed via modular assembly. *Genome Res*. Jul. 2009;19(7):1279-88. doi: 10.1101/gr.089417.108. Epub May 21, 2009. cited by applicant

Kim et al., The role of apolipoprotein E in Alzheimer's disease. *Neuron*. Aug. 13, 2009;63(3):287-303. doi: 10.1016/j.neuron.2009.06.026. cited by applicant

Kim et al., Transcriptional repression by zinc finger peptides. Exploring the potential for applications in gene therapy. *J Biol Chem*. Nov. 21, 1997;272(47):29795-800. cited by applicant

King et al., No gain, no pain: Nav1.7 as an analgesic target. *ACS Chem Neurosci*. Sep. 17, 2014;5(9):749-51. doi: 10.1021/cn500171p. Epub Aug. 11, 2014. cited by applicant

Kitamura et al., Uracil DNA glycosylase counteracts APOBEC3G-induced hypermutation of hepatitis B viral genomes: excision repair of covalently closed circular DNA. *PLoS Pathog.* 2013;9(5):e1003361. doi: 10.1371/journal.ppat.1003361. Epub May 16, 2013. cited by applicant

Klapacz et al., Frameshift mutagenesis and microsatellite instability induced by human alkyladenine DNA glycosylase. *Mol Cell. Mar.* 26, 2010;37(6):843-53. doi: 10.1016/j.molcel.2010.01.038. cited by applicant

Klauser et al., An engineered small RNA-mediated genetic switch based on a ribozyme expression platform. *Nucleic Acids Res.* May 1, 2013;41(10):5542-52. doi: 10.1093/nar/gkt253. Epub Apr. 12, 2013. cited by applicant

Klein et al., Cocystal structure of a class I preQ1 riboswitch reveals a pseudoknot recognizing an essential hypermodified nucleobase. *Nat Struct Mol Biol.* Mar. 2009;16(3):343-4. doi: 10.1038/nsmb.1563. Epub Feb. 22, 2009. cited by applicant

Kleiner et al., In vitro selection of a DNA-templated small-molecule library reveals a class of macrocyclic kinase inhibitors. *J Am Chem Soc.* Aug. 25, 2010;132(33):11779-91. doi: 10.1021/ja104903x. cited by applicant

Kleinstiver et al., Broadening the targeting range of *Staphylococcus aureus* CRISPR-Cas9 by modifying PAM recognition. *Nat Biotechnol.* Dec. 2015;33(12):1293-1298. doi: 10.1038/nbt.3404. Epub Nov. 2, 2015. cited by applicant

Kleinstiver et al., Engineered CRISPR-Cas9 nucleases with altered PAM specificities. *Nature.* Jul. 23, 2015;523(7561):481-5 and Supplementary Materials. doi: 10.1038/nature14592. Epub Jun. 22, 2015. 27 pages. cited by applicant

Kleinstiver et al., Engineered CRISPR-Cas9 nucleases with altered PAM specificities. *Nature.* Jul. 23, 2015;523(7561):481-5. doi: 10.1038/nature14592. Epub Jun. 22, 2015. cited by applicant

Kleinstiver et al., High-fidelity CRISPR-Cas9 nucleases with no detectable genome-wide off-target effects. *Nature.* Jan. 28, 2016;529(7587):490-5. doi: 10.1038/nature16526. Epub Jan. 6, 2016. cited by applicant

Kleinstiver et al., Monomeric site-specific nucleases for genome editing. *Proc Natl Acad Sci U S A.* May 22, 2012;109(21):8061-6. doi: 10.1073/pnas.1117984109. Epub May 7, 2012. cited by applicant

Klement et al., Discrimination between bacteriophage T3 and T7 promoters by the T3 and T7 RNA polymerases depends primarily upon a three base-pair region located 10 to 12 base-pairs upstream from the start site. *J Mol Biol.* Sep. 5, 1990;215(1):21-9. cited by applicant

Klippel et al., Isolation and characterization of unusual gin mutants. *EMBO J.* Dec. 1, 1988;7(12):3983-9. cited by applicant

Klippel et al., The DNA invertase Gin of phage Mu: formation of a covalent complex with DNA via a phosphoserine at amino acid position 9. *EMBO J.* Apr. 1988;7(4):1229-37. cited by applicant

Klompe et al., Transposon-encoded CRISPR-Cas systems direct RNA-guided DNA integration. *Nature.* Jul. 2019;571(7764):219-225. doi: 10.1038/s41586-019-1323-z. Epub Jun. 12, 2019. cited by applicant

Knott et al., Guide-bound structures of an RNA-targeting A-cleaving CRISPR-Cas13a enzyme. *Nat Struct Mol Biol.* Oct. 2017;24(10):825-833. doi: 10.1038/nsmb.3466. Epub Sep. 11, 2017. cited by applicant

Koblan et al., Improving cytidine and adenine base editors by expression optimization and ancestral reconstruction. *Nat Biotechnol.* Oct. 2018;36(9):843-846. doi: 10.1038/nbt.4172. Epub May 29, 2018. cited by applicant

Kobori et al., Deep Sequencing Analysis of Aptazyme Variants Based on a Pistol Ribozyme. *ACS Synth Biol.* Jul. 21, 2017;6(7):1283-1288. doi: 10.1021/acssynbio.7b00057. Epub Apr. 14, 2017. cited by applicant

Kohli et al., A portable hot spot recognition loop transfers sequence preferences from APOBEC family members to activation-induced cytidine deaminase. *J Biol Chem.* Aug. 21, 2009;284(34):22898-904. doi: 10.1074/jbc.M109.025536. Epub Jun. 26, 2009. cited by applicant

Kohli et al., Local sequence targeting in the AID/APOBEC family differentially impacts retroviral restriction and antibody diversification. *J Biol Chem.* Dec. 24, 2010;285(52):40956-64. doi: 10.1074/jbc.M110.177402. Epub Oct. 6, 2010. cited by applicant

Koike-Yusa et al., Genome-wide recessive genetic screening in mammalian cells with a lentiviral CRISPR-guide RNA library. *Nat Biotechnol.* Mar. 2014;32(3):267-73. doi: 10.1038/nbt.2800. Epub Dec. 23, 2013. cited by applicant

Kolot et al., Site promiscuity of coliphage HK022 integrase as a tool for gene therapy. *Gene Ther.* Jul. 2015;22(7):521-7. doi: 10.1038/gt.2015.9. Epub Mar. 12, 2015. cited by applicant

Kolot et al., Site-specific recombination in mammalian cells expressing the Int recombinase of bacteriophage HK022. *Mol Biol Rep.* Aug. 1999;26(3):207-13. doi: 10.1023/a:1007096701720. cited by applicant

Komor et al., CRISPR-Based Technologies for the Manipulation of Eukaryotic Genomes. *Cell.* Jan. 12, 2017;168(1-2):20-36. doi: 10.1016/j.cell.2016.10.044. cited by applicant

Komor et al., Improved base excision repair inhibition and bacteriophage Mu Gam protein yields C:G-to-T:A base editors with higher efficiency and product purity. *Sci Adv.* Aug. 30, 2017;3(8):eaao4774. doi: 10.1126/sciadv.aao4774. eCollection Aug. 2017. cited by applicant

Komor et al., Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage. *Nature.* Apr. 20, 2016;533(7603):420-4. doi: 10.1038/nature17946. cited by applicant

Komor, Editing the Genome Without Double-Stranded DNA Breaks. *ACS Chem Biol.* Feb. 16, 2018;13(2):383-388. doi: 10.1021/acschembio.7b00710. Epub Oct. 9, 2017. cited by applicant

Konermann et al., Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature.* Jan. 29, 2015;517(7536):583-8. doi: 10.1038/nature14136. Epub Dec. 10, 2014. cited by applicant

Koonin et al., Diversity, classification and evolution of CRISPR-Cas systems. *Curr Opin Microbiol.* 2017;37:67-78.

doi:10.1016/j.mib.2017.05.008. cited by applicant

Kosicki et al., Repair of double-strand breaks induced by CRISPR-Cas9 leads to large deletions and complex rearrangements. *Nat Biotechnol.* Sep. 2018;36(8):765-771. doi: 10.1038/nbt.4192. Epub Jul. 16, 2018. cited by applicant

Kotewicz et al., Cloning and overexpression of Moloney murine leukemia virus reverse transcriptase in *Escherichia coli*. *Gene.* 1985;35(3):249-58. doi: 10.1016/0378-1119(85)90003-4. cited by applicant

Kotewicz et al., Isolation of cloned Moloney murine leukemia virus reverse transcriptase lacking ribonuclease H activity. *Nucleic Acids Res.* Jan. 11, 1988;16(1):265-77. doi: 10.1093/nar/16.1.265. cited by applicant

Kotin, Prospects for the use of adeno-associated virus as a vector for human gene therapy. *Hum Gene Ther.* Jul. 1994;5(7):793-801. doi: 10.1089/hum.1994.5.7-793. cited by applicant

Kouzminova et al., Patterns of chromosomal fragmentation due to uracil-DNA incorporation reveal a novel mechanism of replication-dependent double-stranded breaks. *Mol Microbiol.* Apr. 2008;68(1):202-15. doi: 10.1111/j.1365-2958.2008.06149.x. cited by applicant

Kowal et al., Exploiting unassigned codons in *Micrococcus luteus* for tRNA-based amino acid mutagenesis. *Nucleic Acids Res.* Nov. 15, 1997;25(22):4685-9. cited by applicant

Kowalski et al., Delivering the Messenger: Advances in Technologies for Therapeutic mRNA Delivery. *Mol Ther.* Apr. 10, 2019;27(4):710-728. doi: 10.1016/j.ymthe.2019.02.012. Epub Feb. 19, 2019. cited by applicant

Kozak, An analysis of 5'-noncoding sequences from 699 vertebrate messenger RNAs. *Nucleic Acids Res.* Oct. 26, 1987;15(20):8125-48. doi: 10.1093/nar/15.20.8125. cited by applicant

Kraft et al., Deletions, Inversions, Duplications: Engineering of Structural Variants using CRISPR/Cas in Mice. *Cell Rep.* Feb. 10, 2015;10(5):833-839. doi: 10.1016/j.celrep.2015.01.016. Epub Feb. 7, 2015. cited by applicant

Kremer et al., Adenovirus and adeno-associated virus mediated gene transfer. *Br Med Bull.* Jan. 1995;51(1):31-44. doi: 10.1093/oxfordjournals.bmb.a072951. cited by applicant

Krokan et al., Uracil in DNA—occurrence, consequences and repair. *Oncogene.* Dec. 16, 2002;21(58):8935-48. doi: 10.1038/sj.onc.1205996. cited by applicant

Krokan et al., Base excision repair. *Cold Spring Harb Perspect Biol.* Apr. 1, 2013;5(4):a012583. doi: 10.1101/cshperspect.a012583. cited by applicant

Krzywkowski et al., Limited reverse transcriptase activity of phi29 DNA polymerase. *Nucleic Acids Res.* Apr. 20, 2018;46(7):3625-3632. doi: 10.1093/nar/gky190. cited by applicant

Kumar et al., Gene therapy for chronic neuropathic pain: how does it work and where do we stand today? *Pain Med.* May 2011;12(5):808-22. doi: 10.1111/j.1526-4637.2011.01120.x. cited by applicant

Kumar et al., Structural and functional consequences of the mutation of a conserved arginine residue in alphaA and alphaB crystallins. *J Biol Chem.* Aug. 20, 1999;274(34):24137-41. cited by applicant

Kundu et al., Leucine to proline substitution by SNP at position 197 in Caspase-9 gene expression leads to neuroblastoma: a bioinformatics analysis. *3 Biotech.* 2013; 3:225-34. cited by applicant

Kunkel et al., Eukaryotic Mismatch Repair in Relation to DNA Replication. *Annu Rev Genet.* 2015;49:291-313. doi: 10.1146/annurev-genet-112414-054722. cited by applicant

Kunz et al., DNA Repair in mammalian cells: Mismatched repair: variations on a theme. *Cell Mol Life Sci.* Mar. 2009;66(6):1021-38. doi: 10.1007/s00018-009-8739-9. cited by applicant

Kurjan et al., Structure of a yeast pheromone gene (MF alpha): a putative alpha-factor precursor contains four tandem copies of mature alpha-factor. *Cell.* Oct. 1982;30(3):933-43. doi: 10.1016/0092-8674(82)90298-7. cited by applicant

Kury et al., De Novo Disruption of the Proteasome Regulatory Subunit PSMD12 Causes a Syndromic Neurodevelopmental Disorder. *Am J Hum Genet.* Feb. 2, 2017;100(2):352-363. doi: 10.1016/j.ajhg.2017.01.003. Epub Jan. 26, 2017. cited by applicant

Kuscu et al., CRISPR-Cas9-AID base editor is a powerful gain-of-function screening tool. *Nat Methods.* Nov. 29, 2016;13(12):983-984. doi: 10.1038/nmeth.4076. cited by applicant

Kuscu et al., CRISPR-STOP: gene silencing through base-editing-induced nonsense mutations. *Nat Methods.* Jul. 2017;14(7):710-712. doi: 10.1038/nmeth.4327. Epub Jun. 5, 2017. cited by applicant

Kuscu et al., Genome-wide analysis reveals characteristics of off-target sites bound by the Cas9 endonuclease. *Nat Biotechnol.* Jul. 2014;32(7):677-83. doi: 10.1038/nbt.2916. Epub May 18, 2014. cited by applicant

Kwart et al., Precise and efficient scarless genome editing in stem cells using CORRECT. *Nat Protoc.* Feb. 2017;12(2):329-354. doi: 10.1038/nprot.2016.171. Epub Jan. 19, 2017. cited by applicant

Kweon et al., Fusion guide RNAs for orthogonal gene manipulation with Cas9 and Cpf1. *Nat Commun.* Nov. 23, 2017;8(1):1723. doi: 10.1038/s41467-017-01650-w. Erratum in: *Nat Commun.* Jan. 16, 2018;9(1):303. cited by applicant

Kwon et al., Chemical basis of glycine riboswitch cooperativity. *RNA.* Jan. 2008;14(1):25-34. Epub Nov. 27, 2007. cited by applicant

Köhler et al., A possible approach to site-specific insertion of two different unnatural amino acids into proteins in mammalian cells via nonsense suppression. *Chem Biol.* Nov. 2003;10(11):1095-102. cited by applicant

Köhler et al., Complete set of orthogonal 21st aminoacyl-tRNA synthetase-amber, ochre and opal suppressor tRNA pairs: concomitant suppression of three different termination codons in an mRNA in mammalian cells. *Nucleic Acids Res.* Dec. 1, 2004;32(21):6200-11. Print 2004. cited by applicant

Kügler et al., Human synapsin 1 gene promoter confers highly neuron-specific long-term transgene expression from an

adenoviral vector in the adult rat brain depending on the transduced area. *Gene Ther.* Feb. 2003;10(4):337-47. doi: 10.1038/sj.gt.3301905. cited by applicant

Lada et al., Mutator effects and mutation signatures of editing deaminases produced in bacteria and yeast. *Biochemistry (Mosc).* Jan. 2011;76(1):131-46. cited by applicant

Lakich et al., Inversions disrupting the factor VIII gene are a common cause of severe haemophilia A. *Nat Genet.* Nov. 1993;5(3):236-41. doi: 10.1038/ng1193-236. cited by applicant

Lancaster et al., Limited trafficking of a neurotropic virus through inefficient retrograde axonal transport and the type I interferon response. *PLoS Pathog.* Mar. 5, 2010;6(3):e1000791. doi: 10.1371/journal.ppat.1000791. cited by applicant

Landrum et al., ClinVar: public archive of interpretations of clinically relevant variants. *Nucleic Acids Res.* Jan. 4, 2016;44(D1):D862-8. doi: 10.1093/nar/gkv1222. Epub Nov. 17, 2015. cited by applicant

Landrum et al., ClinVar: public archive of relationships among sequence variation and human phenotype. *Nucleic Acids Res.* Jan. 2014;42(Database issue):D980-5. doi: 10.1093/nar/gkt1113. Epub Nov. 14, 2013. cited by applicant

Langer et al., Chemical and Physical Structure of Polymers as Carriers for Controlled Release of Bioactive Agents: A Review. *Journal of Macromolecular Science*, 2006;23(1):61-126. DOI: 10.1080/07366578308079439. cited by applicant

Langer et al., New methods of drug delivery. *Science.* Sep. 28, 1990;249(4976):1527-33. cited by applicant

Larson et al., CRISPR interference (CRISPRi) for sequence-specific control of gene expression. *Nat Protoc.* Nov. 2013;8(11):2180-96. doi: 10.1038/nprot.2013.132. Epub Oct. 17, 2013. cited by applicant

Lau et al., Molecular basis for discriminating between normal and damaged bases by the human alkyladenine glycosylase, AAG. *Proc Natl Acad Sci U S A.* Dec. 5, 2000;97(25):13573-8. cited by applicant

Lauer et al., Construction, characterization, and use of two *Listeria monocytogenes* site-specific phage integration vectors. *J Bacteriol.* Aug. 2002;184(15):4177-86. doi: 10.1128/jb.184.15.4177-4186.2002. cited by applicant

Lavergne et al., Defects in type IIA von Willebrand disease: a cysteine 509 to arginine substitution in the mature von Willebrand factor disrupts a disulphide loop involved in the interaction with platelet glycoprotein Ib-IX. *Br J Haematol.* Sep. 1992;82(1):66-72. cited by applicant

Lawrence et al., Supercharging proteins can impart unusual resilience. *J Am Chem Soc.* Aug. 22, 2007;129(33):10110-2. Epub Aug. 1, 2007. cited by applicant

Lawyer et al., High-level expression, purification, and enzymatic characterization of full-length *Thermus aquaticus* DNA polymerase and a truncated form deficient in 5' to 3' exonuclease activity. *PCR Methods Appl.* May 1993;2(4):275-87. doi: 10.1101/gr.2.4.275. cited by applicant

Lazar et al., Transforming growth factor alpha: mutation of aspartic acid 47 and leucine 48 results in different biological activities. *Mol Cell Biol.* Mar. 1988;8(3):1247-52. cited by applicant

Lazarevic et al., Nucleotide sequence of the *Bacillus subtilis* temperate bacteriophage SPbetac2. *Microbiology (Reading).* May 1999;145 ( Pt 5):1055-1067. doi: 10.1099/13500872-145-5-1055. cited by applicant

Le Grice et al., Purification and characterization of recombinant equine infectious anemia virus reverse transcriptase. *J Virol.* Dec. 1991;65(12):7004-7. doi: 10.1128/JVI.65.12.7004-7007.1991. cited by applicant

Leaver-Fay et al., ROSETTA3: an object-oriented software suite for the simulation and design of macromolecules. *Methods Enzymol.* 2011;487:545-74. doi: 10.1016/B978-0-12-381270-4.00019-6. cited by applicant

Leconte et al., A population-based experimental model for protein evolution: effects of mutation rate and selection stringency on evolutionary outcomes. *Biochemistry.* Feb. 26, 2013;52(8): 1490-9. doi: 10.1021/bi3016185. Epub Feb. 14, 2013. cited by applicant

Ledford, Gene-editing hack yields pinpoint precision. *Nature*, Apr. 20, 2016. <http://www.nature.com/news/gene-editing-hack-yields-pinpoint-precision-1.19773>. cited by applicant

Lee et al., A chimeric thyroid hormone receptor constitutively bound to DNA requires retinoid X receptor for hormone-dependent transcriptional activation in yeast. *Mol Endocrinol.* Sep. 1994;8(9):1245-52. cited by applicant

Lee et al., A monoclonal antibody that targets a NaV1.7 channel voltage sensor for pain and itch relief. *Cell.* Jun. 5, 2014;157(6):1393-1404. doi: 10.1016/j.cell.2014.03.064. Epub May 22, 2014. Retraction in: *Cell.* Jun. 25, 2020;181(7):1695. cited by applicant

Lee et al., An allosteric self-splicing ribozyme triggered by a bacterial second messenger. *Science.* Aug. 13, 2010;329(5993):845-8. doi: 10.1126/science.1190713. cited by applicant

Lee et al., Failure to detect DNA-guided genome editing using *Natronobacterium gregoryi* Argonaute. *Nat Biotechnol.* Nov. 28, 2016;35(1):17-18. doi: 10.1038/nbt.3753. cited by applicant

Lee et al., Group I Intron-Based Therapeutics Through Trans-Splicing Reaction. *Prog Mol Biol Transl Sci.* 2018;159:79-100. doi: 10.1016/bs.pmbts.2018.07.001. Epub Aug. 9, 2018. cited by applicant

Lee et al., PIK3CA gene is frequently mutated in breast carcinomas and hepatocellular carcinomas. *Oncogene.* Feb. 17, 2005;24(8):1477-80. cited by applicant

Lee et al., Recognition of liposomes by cells: in vitro binding and endocytosis mediated by specific lipid headgroups and surface charge density. *Biochim Biophys Acta.* Jan. 31, 1992;1103(2):185-97. cited by applicant

Lee et al., Ribozyme Mediated gRNA Generation for In Vitro and In Vivo CRISPR/Cas9 Mutagenesis. *PLoS One.* Nov. 10, 2016;11(11):e0166020. doi: 10.1371/journal.pone.0166020. eCollection 2016. cited by applicant

Lee et al., Simultaneous targeting of linked loci in mouse embryos using base editing. *Sci Rep.* Feb. 7, 2019;9(1):1662. doi: 10.1038/s41598-018-33533-5. cited by applicant

Lee et al., Site-specific integration of mycobacteriophage L5: integration-proficient vectors for *Mycobacterium smegmatis*, *Mycobacterium tuberculosis*, and bacille Calmette-Guérin. *Proc Natl Acad Sci U S A*. Apr. 15, 1991;88(8):3111-5. doi: 10.1073/pnas.88.8.3111. cited by applicant

Lee et al., Synthetically modified guide RNA and donor DNA are a versatile platform for CRISPR-Cas9 engineering. *Elife*. May 2, 2017;6:e25312. doi: 10.7554/eLife.25312. cited by applicant

Lee et al., Targeted chromosomal deletions in human cells using zinc finger nucleases. *Genome Res*. Jan. 20, 2010: 81-89; Published in Advance Dec. 1, 2009, doi:10.1101/gr.099747.109. cited by applicant

Lee et al., Targeting fidelity of adenine and cytosine base editors in mouse embryos. *Nat Commun*. Nov. 15, 2018;9(1):4804. doi: 10.1038/s41467-018-07322-7. cited by applicant

Lee et al., Transcriptional regulation and its misregulation in disease. *Cell*. Mar. 14, 2013;152(6):1237-51. doi: 10.1016/j.cell.2013.02.014. cited by applicant

Lei et al., Efficient targeted gene disruption in *Xenopus* embryos using engineered transcription activator-like effector nucleases (TALENs). *Proc Natl Acad Sci U S A*. Oct. 23, 2012;109(43):17484-9. Doi: 10.1073/pnas.1215421109. Epub Oct. 8, 2012. cited by applicant

Lei et al., Site-specificity of serine integrase demonstrated by the attB sequence preference of  $\phi$ BT1 integrase. *FEBS Lett*. Apr. 2018;592(8):1389-1399. doi: 10.1002/1873-3468.13023. Epub Mar. 25, 2018. cited by applicant

Leipold et al., A de novo gain-of-function mutation in SCN11A causes loss of pain perception. *Nat Genet*. Nov. 2013;45(11):1399-404. doi: 10.1038/ng.2767. Epub Sep. 15, 2013. cited by applicant

Lemos et al., CRISPR/Cas9 cleavages in budding yeast reveal templated insertions and strand-specific insertion/deletion profiles. *Proc Natl Acad Sci U S A*. Feb. 27, 2018;115(9):E2040-E2047. doi: 10.1073/pnas.1716855115. Epub Feb. 13, 2018. cited by applicant

Lenk et al., Pathogenic mechanism of the FIG4 mutation responsible for Charcot-Marie-Tooth disease CMT4J. *PLoS Genet*. Jun. 2011;7(6):e1002104. doi: 10.1371/journal.pgen.1002104. Epub Jun. 2, 2011. cited by applicant

Levy et al., Cytosine and adenine base editing of the brain, liver, retina, heart and skeletal muscle of mice via adeno-associated viruses. *Nat Biomed Eng*. 2020;4(1):97-110. doi:10.1038/s41551-019-0501-5. cited by applicant

Levy et al., Inhibition of calcification of bioprosthetic heart valves by local controlled-release diphosphonate. *Science*. Apr. 12, 1985;228(4696):190-2. cited by applicant

Levy et al., Membrane-associated guanylate kinase dynamics reveal regional and developmental specificity of synapse stability. *J Physiol*. Mar. 1, 2017;595(5):1699-1709. doi: 10.1113/JP273147. Epub Jan. 18, 2017. cited by applicant

Lew et al., Protein splicing in vitro with a semisynthetic two-component minimal intein. *J Biol Chem*. Jun. 26, 1998;273(26):15887-90. doi: 10.1074/jbc.273.26.15887. cited by applicant

Lewis et al., A serum-resistant cytofectin for cellular delivery of antisense oligodeoxynucleotides and plasmid DNA. *Proc Natl Acad Sci U S A*. Apr. 16, 1996;93(8):3176-81. cited by applicant

Lewis et al., Building the Class 2 CRISPR-Cas Arsenal. *Mol Cell* 2017;65(3):377-379. cited by applicant

Lewis et al., Codon 129 polymorphism of the human prion protein influences the kinetics of amyloid formation. *J Gen Virol*. Aug. 2006;87(Pt 8):2443-9. cited by applicant

Lewis et al., Cytosine deamination and the precipitous decline of spontaneous mutation during Earth's history. *Proc Natl Acad Sci U S A*. Jul. 19, 2016;113(29):8194-9. doi: 10.1073/pnas.1607580113. Epub Jul. 5, 2016. cited by applicant

Lewis et al., RNA modifications and structures cooperate to guide RNA-protein interactions. *Nat Rev Mol Cell Biol*. Mar. 2017;18(3):202-210. doi: 10.1038/nrm.2016.163. Epub Feb. 1, 2017. cited by applicant

Li et al., A Radioactivity-Based Assay for Screening Human m6A-RNA Methyltransferase, METTL3-METTL14 Complex, and Demethylase ALKBH5. *J Biomol Screen*. Mar. 2016;21(3):290-7. doi: 10.1177/1087057115623264. Epub Dec. 23, 2015. cited by applicant

Li et al., Base editing with a Cpf1-cytidine deaminase fusion. *Nat Biotechnol*. Apr. 2018;36(4):324-327. doi: 10.1038/nbt.4102. Epub Mar. 19, 2018. cited by applicant

Li et al., Current approaches for engineering proteins with diverse biological properties. *Adv Exp Med Biol*. 2007;620:18-33. cited by applicant

Li et al., Disruption of splicing-regulatory elements using CRISPR/Cas9 to rescue spinal muscular atrophy in human iPSCs and mice. *National Science Review*. Jan. 1, 2020:92-101. DOI: 10.1093/nsr/nwz131. Retrieved from the Internet via <https://academic.oup.com/nsr/article-pdf/7/1/92/33321439/nwz131.pdf>. Last accessed Apr. 28, 2021. cited by applicant

Li et al., Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*. Jul. 15, 2009;25(14):1754-60. doi: 10.1093/bioinformatics/btp324. Epub May 18, 2009. cited by applicant

Li et al., Generation of Targeted Point Mutations in Rice by a Modified CRISPR/Cas9 System. *Mol Plant*. Mar. 6, 2017;10(3):526-529. doi: 10.1016/j.molp.2016.12.001. Epub Dec. 8, 2016. cited by applicant

Li et al., Highly efficient and precise base editing in discarded human tripronuclear embryos. *Protein Cell*. Aug. 19, 2017. doi: 10.1007/s13238-017-0458-7. [Epub ahead of print]. cited by applicant

Li et al., Lagging strand DNA synthesis at the eukaryotic replication fork involves binding and stimulation of FEN-1 by proliferating cell nuclear antigen. *J Biol Chem*. Sep. 22, 1995;270(38):22109-12. doi: 10.1074/jbc.270.38.22109. cited by applicant

Li et al., Loss of post-translational modification sites in disease. *Pac Symp Biocomput*. 2010:337-47. doi:

10.1142/9789814295291\_0036. cited by applicant

Li et al., Modularly assembled designer TAL effector nucleases for targeted gene knockout and gene replacement in eukaryotes. *Nucleic Acids Res.* Aug. 2011;39(14):6315-25. doi: 10.1093/nar/gkr188. Epub Mar. 31, 2011. cited by applicant

Li et al., Multiplex and homologous recombination-mediated genome editing in *Arabidopsis* and *Nicotiana benthamiana* using guide RNA and Cas9. *Nat Biotechnol.* Aug. 2013;31(8):688-91. doi: 10.1038/nbt.2654. cited by applicant

Li et al., Programmable Single and Multiplex Base-Editing in *Bombyx mori* Using RNA-Guided Cytidine Deaminases. *G3 (Bethesda)*. May 4, 2018;8(5):1701-1709. doi: 10.1534/g3.118.200134. cited by applicant

Li et al., Protein trans-splicing as a means for viral vector-mediated in vivo gene therapy. *Hum Gene Ther.* Sep. 2008;19(9):958-64. doi: 10.1089/hum.2008.009. cited by applicant

Li et al., RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics.* Aug. 4, 2011;12:323. doi: 10.1186/1471-2105-12-323. cited by applicant

Li et al., TAL nucleases (TALNs): hybrid proteins composed of TAL effectors and FokI DNA-cleavage domain. *Nucleic Acids Res.* Jan. 2011;39(1):359-72. doi: 10.1093/nar/gkq704. Epub Aug. 10, 2010. cited by applicant

Li, Mechanisms and functions of DNA mismatch repair. *Cell Res.* Jan. 2008;18(1):85-98. doi: 10.1038/cr.2007.115. cited by applicant

Liang et al., Correction of  $\beta$ -thalassemia mutant by base editor in human embryos. *Protein Cell.* Nov. 2017;8(11):811-822. doi: 10.1007/s13238-017-0475-6. Epub Sep. 23, 2017. cited by applicant

Liang et al., Homology-directed repair is a major double-strand break repair pathway in mammalian cells. *Proc Natl Acad Sci U S A.* Apr. 28, 1998;95(9):5172-7. doi: 10.1073/pnas.95.9.5172. cited by applicant

Liang et al., Rapid and highly efficient mammalian cell engineering via Cas9 protein transfection. *Send to; J Biotechnol.* Aug. 20, 2015;208:44-53. doi: 10.1016/j.jbiotec.2015.04.024. cited by applicant

Liao et al., One-step assembly of large CRISPR arrays enables multi-functional targeting and reveals constraints on array design. *bioRxiv.* May 2, 2018. doi: 10.1101/312421. 45 pages. cited by applicant

Lieber et al., Mechanism and regulation of human non-homologous DNA end-joining. *Nat Rev Mol Cell Biol.* Sep. 2003;4(9):712-20. cited by applicant

Liefke et al., The oxidative demethylase ALKBH3 marks hyperactive gene promoters in human cancer cells. *Genome Med.* Jun. 30, 2015;7(1):66. doi: 10.1186/s13073-015-0180-0. cited by applicant

Lienert et al., Two- and three-input TALE-based AND logic computation in embryonic stem cells. *Nucleic Acids Res.* Nov. 2013;41(21):9967-75. doi: 10.1093/nar/gkt758. Epub Aug. 27, 2013. cited by applicant

Lilley, D.M. The Varkud Satellite Ribozyme. *RNA.* Feb. 2004;10(2):151-8. doi: 10.1261/rna.5217104. cited by applicant

Lim et al., Crystal structure of the moloney murine leukemia virus RNase H domain. *J Virol.* Sep. 2006;80(17):8379-89. doi: 10.1128/JVI.00750-06. cited by applicant

Lim et al., Viral vectors for neurotrophic factor delivery: a gene therapy approach for neurodegenerative diseases of the CNS. *Pharmacol Res.* Jan. 2010;61(1):14-26. doi: 10.1016/j.phrs.2009.10.002. Epub Oct. 17, 2009. cited by applicant

Lin et al., Enhanced homology-directed human genome engineering by controlled timing of CRISPR/Cas9 delivery. *Elife.* Dec. 15, 2014;3:e04766. doi: 10.7554/eLife.04766. cited by applicant

Lin et al., The human REV1 gene codes for a DNA template-dependent dCMP transferase. *Nucleic Acids Res.* Nov. 15, 1999;27(22):4468-75. doi: 10.1093/nar/27.22.4468. cited by applicant

Link et al., Engineering ligand-responsive gene-control elements: lessons learned from natural riboswitches. *Gene Ther.* Oct. 2009;16(10):1189-201. doi: 10.1038/gt.2009.81. Epub Jul. 9, 2009. Review. cited by applicant

Liu et al., C2c1-sgRNA Complex Structure Reveals RNA-Guided DNA Cleavage Mechanism. *Molecular Cell* Jan. 2017;65(2):310-22. cited by applicant

Liu et al., Split dnaE genes encoding multiple novel inteins in *Trichodesmium erythraeum*. *J Biol Chem.* Jul. 18, 2003;278(29):26315-8. doi: 10.1074/jbc.C300202200. Epub May 24, 2003. cited by applicant

Liu et al., A METTL3-METTL14 complex mediates mammalian nuclear RNA N6-adenosine methylation. *Nat Chem Biol.* Feb. 2014;10(2):93-5. doi: 10.1038/nchembio.1432. Epub Dec. 6, 2013. cited by applicant

Liu et al., Adding new chemistries to the genetic code. *Annu Rev Biochem.* 2010;79:413-44. doi: 10.1146/annurev.biochem.052308.105824. cited by applicant

Liu et al., Apolipoprotein E and Alzheimer disease: risk, mechanisms and therapy. *Nat Rev Neurol.* Feb. 2013;9(2):106-18. doi: 10.1038/nrneurol.2012.263. Epub Jan. 8, 2013. cited by applicant

Liu et al., Balancing AID and DNA repair during somatic hypermutation. *Trends Immunol.* Apr. 2009;30(4):173-81. doi: 10.1016/j.it.2009.01.007. cited by applicant

Liu et al., Calcineurin is a common target of cyclophilin-cyclosporin A and FKBP-FK506 complexes. *Cell.* Aug. 23, 1991;66(4):807-15. doi: 10.1016/0092-8674(91)90124-h. cited by applicant

Liu et al., CasX enzymes comprise a distinct family of RNA-guided genome editors. *Nature.* Feb. 2019;566(7743):218-223. doi: 10.1038/s41586-019-0908-x. Epub Feb. 4, 2019. Author manuscript entitled CRISPR-CasX is an RNA-dominated enzyme active for human genome editing. cited by applicant

Liu et al., Cell-penetrating peptide-mediated delivery of TALEN proteins via bioconjugation for genome engineering. *PLoS One.* Jan. 20, 2014;9(1):e85755. doi: 10.1371/journal.pone.0085755. eCollection 2014. cited by applicant

Liu et al., Computational approaches for effective CRISPR guide RNA design and evaluation. *Comput Struct Biotechnol*

J. Nov. 29, 2019;18:35-44. doi: 10.1016/j.csbj.2019.11.006. cited by applicant

Liu et al., Design of polydactyl zinc-finger proteins for unique addressing within complex genomes. Proc Natl Acad Sci U S A. May 27, 1997;94(11):5525-30. cited by applicant

Liu et al., Direct Promoter Repression by BCL11A Controls the Fetal to Adult Hemoglobin Switch. Cell. Apr. 5, 2018;173(2):430-442.e17. doi: 10.1016/j.cell.2018.03.016. Epub Mar. 29, 2018. cited by applicant

Liu et al., Distance determination by GIY-YIG intron endonucleases: discrimination between repression and cleavage functions. Nucleic Acids Res. Mar. 31, 2006;34(6):1755-64. Print 2006. cited by applicant

Liu et al., Editing DNA Methylation in the Mammalian Genome. Cell. Sep. 22, 2016;167(1):233-247.e17. doi: 10.1016/j.cell.2016.08.056. cited by applicant

Liu et al., Engineering a tRNA and aminoacyl-tRNA synthetase for the site-specific incorporation of unnatural amino acids into proteins in vivo. Proc Natl Acad Sci U S A. Sep. 16, 1997;94(19):10092-7. cited by applicant

Liu et al., Fast Colorimetric Sensing of Adenosine and Cocaine Based on a General Sensor Design Involving Aptamers and Nanoparticles. Angew Chem. Dec. 16, 2006;45(1):90-4. DOI: 10.1002/anie.200502589. cited by applicant

Liu et al., Fast Colorimetric Sensing of Adenosine and Cocaine Based on a General Sensor Design Involving Aptamers and Nanoparticles. Angew Chem. 2006;118(1):96-100. cited by applicant

Liu et al., Flap endonuclease 1: a central component of DNA metabolism. Annu Rev Biochem. 2004;73:589-615. doi:10.1146/annurev.biochem.73.012803.092453. cited by applicant

Liu et al., Functional Nucleic Acid Sensors. Chem Rev. May 2009;109(5):1948-98. doi: 10.1021/cr030183i. cited by applicant

Liu et al., Genetic incorporation of unnatural amino acids into proteins in mammalian cells. Nat Methods. Mar. 2007;4(3):239-44. Epub Feb. 25, 2007. cited by applicant

Liu et al., Highly efficient RNA-guided base editing in rabbit. Nat Commun. Jul. 13, 2018;9(1):2717. doi: 10.1038/s41467-018-05232-2. cited by applicant

Liu et al., (6)-methyladenosine-dependent RNA structural switches regulate RNA-protein interactions. Nature. Feb. 26, 2015;518(7540):560-4. doi: 10.1038/nature14234. cited by applicant

Liu et al., Probing N6-methyladenosine RNA modification status at single nucleotide resolution in mRNA and long noncoding RNA. RNA. Dec. 2013;19(12):1848-56. doi: 10.1261/rna.041178.113. Epub Oct. 18, 2013. cited by applicant

Liu et al., Reverse transcriptase of foamy virus. Purification of the enzymes and immunological identification. Arch Virol. 1977;55(3):187-200. doi: 10.1007/BF01319905. cited by applicant

Liu et al., Reverse transcriptase-mediated tropism switching in Bordetella bacteriophage. Science. Mar. 15, 2002;295(5562):2091-4. doi: 10.1126/science.1067467. cited by applicant

Liu et al., *Saccharomyces cerevisiae* flap endonuclease 1 uses flap equilibration to maintain triplet repeat stability. Mol Cell Biol. May 2004;24(9):4049-64. doi: 10.1128/MCB.24.9.4049-4064.2004. cited by applicant

Liu et al., The Molecular Architecture for RNA-Guided RNA Cleavage by Cas13a. Cell. Aug. 10, 2017;170(4):714-726.e10. doi: 10.1016/j.cell.2017.06.050. Epub Jul. 27, 2017. cited by applicant

Loessner et al., Complete nucleotide sequence, molecular analysis and genome structure of bacteriophage A118 of *Listeria monocytogenes*: implications for phage evolution. Mol Microbiol. Jan. 2000;35(2):324-40. doi: 10.1046/j.1365-2958.2000.01720.x. cited by applicant

Lombardo et al., Gene editing in human stem cells using zinc finger nucleases and integrase-defective lentiviral vector delivery. Nat Biotechnol. Nov. 2007;25(11):1298-306. Epub Oct. 28, 2007. cited by applicant

Long et al., Postnatal genome editing partially restores dystrophin expression in a mouse model of muscular dystrophy. Science. Jan. 22, 2016;351(6271):400-3. doi: 10.1126/science.aad5725. Epub Dec. 31, 2015. cited by applicant

Lopez-Girona et al., Cereblon is a direct protein target for immunomodulatory and antiproliferative activities of lenalidomide and pomalidomide. Leukemia. Nov. 2012;26(11):2326-35. doi: 10.1038/leu.2012.119. Epub May 3, 2012. cited by applicant

Lorenz et al., ViennaRNA Package 2.0. Algorithms Mol Biol. Nov. 24, 2011;6:26. doi: 10.1186/1748-7188-6-26. cited by applicant

Losey et al., Crystal structure of *Staphylococcus aureus* tRNA adenosine deaminase tadA in complex with RNA. Nature Struct. Mol. Biol. Feb. 2006;13(2):153-9. cited by applicant

Lu et al., Precise Editing of a Target Base in the Rice Genome Using a Modified CRISPR/Cas9 System. Mol Plant. Mar. 6, 2017;10(3):523-525. doi: 10.1016/j.molp.2016.11.013. Epub Dec. 6, 2016. cited by applicant

Luan et al., Reverse transcription of R2Bm RNA is primed by a nick at the chromosomal target site: a mechanism for non-LTR retrotransposition. Cell. Feb. 26, 1993;72(4):595-605. doi: 10.1016/0092-8674(93)90078-5. cited by applicant

Luckow et al., High level expression of nonfused foreign genes with Autographa californica nuclear polyhedrosis virus expression vectors. Virology. May 1989;170(1):31-9. doi: 10.1016/0042-6822(89)90348-6. cited by applicant

Lukacsovich et al., Repair of a specific double-strand break generated within a mammalian chromosome by yeast endonuclease I-SceI. Nucleic Acids Res. Dec. 25, 1994;22(25):5649-57. doi: 10.1093/nar/22.25.5649. cited by applicant

Lundberg et al., Delivery of short interfering RNA using endosomolytic cell-penetrating peptides. FASEB J. Sep. 2007;21(11):2664-71. Epub Apr. 26, 2007. cited by applicant

Lundquist et al., Site-directed mutagenesis and characterization of uracil-DNA glycosylase inhibitor protein. Role of specific carboxylic amino acids in complex formation with *Escherichia coli* uracil-DNA glycosylase. J Biol Chem. Aug.



22, 1997;272(34):21408-19. cited by applicant

Lynch, Evolution of the mutation rate. Trends Genet. Aug. 2010;26(8):345-52. doi: 10.1016/j.tig.2010.05.003. Epub Jun. 30, 2010. cited by applicant

Lyons et al., Efficient Recognition of an Unpaired Lesion by a DNA Repair Glycosylase. J. Am. Chem. Soc., 2009;131(49):17742-3. DOI: 10.1021/ja908378y. cited by applicant

Lüke et al., Partial purification and characterization of the reverse transcriptase of the simian immunodeficiency virus TYO-7 isolated from an African green monkey. Biochemistry. Feb. 20, 1990;29(7):1764-9. doi: 10.1021/bi00459a015. cited by applicant

Ma et al., Identification of pseudo attP sites for phage phiC31 integrase in bovine genome. Biochem Biophys Res Commun. Jul. 7, 2006;345(3):984-8. doi: 10.1016/j.bbrc.2006.04.145. Epub May 3, 2006. cited by applicant

Ma et al., In vitro protein engineering using synthetic tRNA(Ala) with different anticodons. Biochemistry. Aug. 10, 1993;32(31):7939-45. cited by applicant

Ma et al., PhiC31 integrase induces efficient site-specific recombination in the Capra hircus genome. DNA Cell Biol. Aug. 2014;33(8):484-91. doi: 10.1089/dna.2013.2124. Epub Apr. 22, 2014. cited by applicant

Ma et al., Single-Stranded DNA Cleavage by Divergent CRISPR-Cas9 Enzymes. Mol Cell. Nov. 5, 2015;60(3):398-407. doi: 10.1016/j.molcel.2015.10.030. cited by applicant

Ma et al., Targeted AID-mediated mutagenesis (TAM) enables efficient genomic diversification in mammalian cells. Nature Methods. Oct. 2016;13:1029-35. doi:10.1038/nmeth.4027. cited by applicant

Maas et al., Identification and characterization of a human tRNA-specific adenosine deaminase related to the ADAR family of pre-mRNA editing enzymes. Proc Natl Acad Sci U S A. Aug. 3, 1999;96(16):8895-900. doi: 10.1073/pnas.96.16.8895. cited by applicant

Macbeth et al., Inositol hexakisphosphate is bound in the ADAR2 core and required for RNA editing. Science. Sep. 2, 2005;309(5740):1534-9. doi: 10.1126/science.1113150. cited by applicant

Macrae et al., Ribonuclease revisited: structural insights into ribonuclease III family enzymes. Curr Opin Struct Biol. Feb. 2007;17(1):138-45. doi: 10.1016/j.sbi.2006.12.002. Epub Dec. 27, 2006. cited by applicant

Madura et al., Structural basis for ineffective T-cell responses to MHC anchor residue-improved “heteroclitic” peptides. Eur J Immunol. Feb. 2015;45(2):584-91. doi: 10.1002/eji.201445114. Epub Dec. 28, 2014. cited by applicant

Maeder et al., CRISPR RNA-guided activation of endogenous human genes. Nat Methods. Oct. 2013;10(10):977-9. doi: 10.1038/nmeth.2598. Epub Jul. 25, 2013. cited by applicant

Maeder et al., Rapid “open-source” engineering of customized zinc-finger nucleases for highly efficient gene modification. Mol Cell. Jul. 25, 2008;31(2):294-301. doi:10.1016/j.molcel.2008.06.016. cited by applicant

Maeder et al., Robust, synergistic regulation of human gene expression using TALE activators. Nat Methods. Mar. 2013;10(3):243-5. doi: 10.1038/nmeth.2366. Epub Feb. 10, 2013. cited by applicant

Magin et al., Corf, the Rev/Rex homologue of HTDV/HERV-K, encodes an arginine-rich nuclear localization signal that exerts a trans-dominant phenotype when mutated. Virology. Aug. 15, 2000;274(1):11-6. doi: 10.1006/viro.2000.0438. cited by applicant

Mahfouz et al., De novo-engineered transcription activator-like effector (TALE) hybrid nuclease with novel DNA binding specificity creates double-strand breaks. Proc Natl Acad Sci U S A. Feb. 8, 2011;108(6):2623-8. doi: 10.1073/pnas.1019533108. Epub Jan. 24, 2011. cited by applicant

Maizels et al., Initiation of homologous recombination at DNA nicks. Nucleic Acids Res. Aug. 21, 2018;46(14):6962-6973. doi: 10.1093/nar/gky588. cited by applicant

Maji et al., A High-Throughput Platform to Identify Small-Molecule Inhibitors of CRISPR-Cas9. Cell. May 2, 2019;177(4):1067-1079.e19. doi: 10.1016/j.cell.2019.04.009. cited by applicant

Makarova et al., An updated evolutionary classification of CRISPR-Cas systems. Nat Rev Microbiol. Nov. 2015;13(11):722-36. doi: 10.1038/nrmicro3569. Epub Sep. 28, 2015. cited by applicant

Makarova et al., Classification and Nomenclature of CRISPR-Cas Systems: Where from Here? CRISPR J. Oct. 2018;1(5):325-336. doi: 10.1089/crispr.2018.0033. cited by applicant

Makarova et al., Evolution and classification of the CRISPR-Cas systems. Nat Rev Microbiol. Jun. 2011;9(6):467-77. doi: 10.1038/nrmicro2577. Epub May 9, 2011. cited by applicant

Makarova et al., Prokaryotic homologs of Argonaute proteins are predicted to function as key components of a novel system of defense against mobile genetic elements. Biology Direct 2009;4:29. doi: 10.1186/1745-6150-4-29. cited by applicant

Makeyev et al., Evolutionary potential of an RNA virus. J Virol. Feb. 2004;78(4):2114-20. cited by applicant

Malashkevich et al., Crystal structure of tRNA adenosine deaminase TadA from *Escherichia coli*. Deposited: Mar. 10, 2005 Released: Feb. 21, 2006 doi:10.2210/pdb1z3a/pdb (2006). cited by applicant

Mali et al., Cas9 as a versatile tool for engineering biology. Nat Methods. Oct. 2013;10(10):957-63. doi: 10.1038/nmeth.2649. cited by applicant

Mali et al., CAS9 transcriptional activators for target specificity screening and paired nickases for cooperative genome engineering. Nat Biotechnol. Sep. 2013;31(9):833-8. doi: 10.1038/nbt.2675. Epub Aug. 1, 2013. cited by applicant

Mali et al., RNA-guided human genome engineering via Cas9. Science. Feb. 15, 2013;339(6121):823-6. doi: 10.1126/science.1232033. Epub Jan. 3, 2013. cited by applicant

Malito et al., Structural basis for lack of the diphtheria toxin mutant CRM197. *Proc Natl Acad Sci U S A*. Apr. 3, 2012;109(14):5229-34. doi: 10.1073/pnas.1201964109. Epub Mar. 19, 2012. cited by applicant

Mandal et al., Efficient ablation of genes in human hematopoietic stem and effector cells using CRISPR/Cas9. *Cell Stem Cell*. Nov. 6, 2014;15(5):643-52. doi: 10.1016/j.stem.2014.10.004. Epub Nov. 6, 2014. cited by applicant

Mandal et al., Riboswitches Control Fundamental Biochemical Pathways in *Bacillus Subtilis* and Other Bacteria. *Cell*. May 30, 2003;113(5):577-86. doi: 10.1016/s0092-8674(03)00391-x. cited by applicant

Mani et al., Design, engineering, and characterization of zinc finger nucleases. *Biochem Biophys Res Commun*. Sep. 23, 2005;335(2):447-57. cited by applicant

Marceau, Functions of single-strand DNA-binding proteins in DNA replication, recombination, and repair. *Methods Mol Biol*. 2012;922:1-21. doi: 10.1007/978-1-62703-032-8\_1. cited by applicant

Maresca et al., Obligate ligation-gated recombination (ObLiGaRe): custom-designed nuclease-mediated targeted integration through nonhomologous end joining. *Genome Res*. Mar. 2013;23(3):539-46. Doi: 10.1101/gr.145441.112. Epub Nov. 14, 2012. cited by applicant

Marioni et al., DNA methylation age of blood predicts all-cause mortality in later life. *Genome Biol*. Jan. 30, 2015;16:25. doi: 10.1186/s13059-015-0584-6. cited by applicant

Marraffini et al., CRISPR interference limits horizontal gene transfer in staphylococci by targeting DNA. *Science*. Dec. 19, 2008;322(5909):1843-5. doi: 10.1126/science.1165771. cited by applicant

Martinez et al., Hypermutagenesis of RNA using human immunodeficiency virus type 1 reverse transcriptase and biased dNTP concentrations. *Proc Natl Acad Sci U S A*. Dec. 6, 1994;91(25):11787-91. doi: 10.1073/pnas.91.25.11787. cited by applicant

Martsof et al., Complete trisomy 17p a relatively new syndrome. *Ann Genet*. 1988;31(3):172-4. cited by applicant

Martz, L., Nav-i-gating antibodies for pain. *Science-Business eXchange*. Jun. 12, 2014;7(662):1-2. doi: 10.1038/scibx.2014.662. cited by applicant

Maruyama et al., Increasing the efficiency of precise genome editing with CRISPR-Cas9 by inhibition of nonhomologous end joining. *Nat Biotechnol*. May 2015;33(5):538-42. doi: 10.1038/nbt.3190. Epub Mar. 23, 2015. cited by applicant

Marzec et al., Prime Editing: A New Way for Genome Editing. *Trends Cell Biol*. Apr. 2020;30(4):257-259. doi: 10.1016/j.tcb.2020.01.004. Epub Jan. 27, 2020. cited by applicant

Mascola et al., HIV-1 neutralizing antibodies: understanding nature's pathways. *Immunol Rev*. Jul. 2013;254(1):225-44. doi: 10.1111/immr.12075. cited by applicant

Mathys et al., Characterization of a self-splicing mini-intein and its conversion into autocatalytic—and C-terminal cleavage elements: facile production of protein building blocks for protein ligation. *Gene*. Apr. 2, 1999;231(1-2):1-13. doi: 10.1016/s0378-1119(99)00103-1. cited by applicant

Matsuura et al., A gene essential for the site-specific excision of actinophage r4 prophage genome from the chromosome of a lysogen. *J Gen Appl Microbiol*. 1995;41(1):53-61. cited by applicant

Matthews, Structures of human ADAR2 bound to dsRNA reveal base-flipping mechanism and basis for site selectivity. *Nat Struct Mol Biol*. May 2016;23(5):426-33. doi: 10.1038/nsmb.3203. Epub Apr. 11, 2016. cited by applicant

May et al., Emergent lineages of mumps virus suggest the need for a polyvalent vaccine. *Int J Infect Dis*. Jan. 2018;66:1-4. doi: 10.1016/j.ijid.2017.09.024. Epub Oct. 4, 2017. cited by applicant

McCarroll et al., Copy-number variation and association studies of human disease. *Nat Genet*. Jul. 2007;39(7 Suppl):S37-42. doi: 10.1038/ng2080. cited by applicant

McDonald et al., Characterization of mutations at the mouse phenylalanine hydroxylase locus. *Genomics*. Feb. 1, 1997;39(3):402-5. doi: 10.1006/geno.1996.4508. cited by applicant

McInerney et al., Error Rate Comparison during Polymerase Chain Reaction by DNA Polymerase. *Mol Biol Int*. 2014;2014:287430. doi: 10.1155/2014/287430. Epub Aug. 17, 2014. cited by applicant

McKenna et al., Recording development with single cell dynamic lineage tracing. *Development*. Jun. 27, 2019;146(12):dev169730. doi: 10.1242/dev.169730. cited by applicant

McKenna et al., Whole-organism lineage tracing by combinatorial and cumulative genome editing. *Science*. Jul. 29, 2016;353(6298):aaf7907. doi: 10.1126/science.aaf7907. Epub May 26, 2016. cited by applicant

McNaughton et al., Mammalian cell penetration, siRNA transfection, and DNA transfection by supercharged proteins. *Proc Natl Acad Sci U S A*. Apr. 14, 2009;106(15):6111-6. doi: 10.1073/pnas.0807883106. Epub Mar. 23, 2009. cited by applicant

McVey et al., MMEJ repair of double-strand breaks (director's cut): deleted sequences and alternative endings. *Trends Genet*. Nov. 2008;24(11):529-38. doi: 10.1016/j.tig.2008.08.007. Epub Sep. 21, 2008. cited by applicant

Mead et al., A novel protective prion protein variant that colocalizes with kuru exposure. *Engl J Med*. Nov. 19, 2009;361(21):2056-65. doi: 10.1056/NEJMoa0809716. cited by applicant

Mei et al., Recent Progress in CRISPR/Cas9 Technology. *J Genet Genomics*. Feb. 20, 2016;43(2):63-75. doi: 10.1016/j.jgg.2016.01.001. Epub Jan. 18, 2016. cited by applicant

Meinke et al., Cre Recombinase and Other Tyrosine Recombinases. *Chem Rev*. Oct. 26, 2016;116(20):12785-12820. doi: 10.1021/acs.chemrev.6b00077. Epub May 10, 2016. cited by applicant

Meng et al., Targeted gene inactivation in zebrafish using engineered zinc-finger nucleases. *Nat Biotechnol*. Jun. 2008;26(6):695-701. doi: 10.1038/nbt1398. Epub May 25, 2008. cited by applicant

Menéndez-Arias, Mutation rates and intrinsic fidelity of retroviral reverse transcriptases. *Viruses*. Dec. 2009;1(3):1137-65. doi: 10.3390/v1031137. Epub Dec. 4, 2009. cited by applicant

Mercer et al., Chimeric TALE recombinases with programmable DNA sequence specificity. *Nucleic Acids Res*. Nov. 2012;40(21):11163-72. doi: 10.1093/nar/gks875. Epub Sep. 26, 2012. cited by applicant

Mertens et al., Site-specific recombination in bacteriophage Mu: characterization of binding sites for the DNA invertase Gin. *EMBO J*. Apr. 1988;7(4):1219-27. cited by applicant

Meyer et al., Breathing life into polycations: functionalization with pH-responsive endosomolytic peptides and polyethylene glycol enables siRNA delivery. *J Am Chem Soc*. Mar. 19, 2008;130(11):3272-3. doi: 10.1021/ja710344v. Epub Feb. 21, 2008. cited by applicant

Meyer et al., Comprehensive analysis of mRNA methylation reveals enrichment in 3' UTRs and near stop codons. *Cell*. Jun. 22, 2012;149(7):1635-46. doi: 10.1016/j.cell.2012.05.003. Epub May 17, 2012. cited by applicant

Meyer et al., Confirmation of a second natural preQ1 aptamer class in Streptococcaceae bacteria. *RNA*. Apr. 2008;14(4):685-95. doi: 10.1261/rna.937308. Epub Feb. 27, 2008. cited by applicant

Meyer et al., Library generation by gene shuffling. *Curr Protoc Mol Biol*. Jan. 6, 2014;105: Unit 15.12 . . . doi: 10.1002/0471142727.mb1512s105. cited by applicant

Meyer et al., Ribosome biogenesis factor Tsr3 is the aminocarboxypropyl transferase responsible for 18S rRNA hypermodification in yeast and humans. *Nucleic Acids Res*. May 19, 2016;44(9):4304-16. doi: 10.1093/nar/gkw244. Epub Apr. 15, 2016. cited by applicant

Meyer et al., The dynamic epitranscriptome: N6-methyladenosine and gene expression control. *Nat Rev Mol Cell Biol*. May 2014;15(5):313-26. doi: 10.1038/nrm3785. Epub Apr. 9, 2014. cited by applicant

Michel et al., Mitochondrial class II introns encode proteins related to the reverse transcriptases of retroviruses. *Nature*. Aug. 15-21, 1985;316(6029):641-3. doi: 10.1038/316641a0. cited by applicant

Midoux et al., Chemical vectors for gene delivery: a current review on polymers, peptides and lipids containing histidine or imidazole as nucleic acids carriers. *Br J Pharmacol*. May 2009;157(2):166-78. doi: 10.1111/j.1476-5381.2009.00288.x. cited by applicant

Mihai et al., PTEN inhibition improves wound healing in lung epithelia through changes in cellular mechanics that enhance migration. *Am J Physiol Lung Cell Mol Physiol*. 2012;302(3):L287-L299. cited by applicant

Mijakovic et al., Bacterial single-stranded DNA-binding proteins are phosphorylated on tyrosine. *Nucleic Acids Res*. Mar. 20, 2006;34(5):1588-96. doi: 10.1093/nar/gkj514. cited by applicant

Miller et al., A TALE nuclease architecture for efficient genome editing. *Nat Biotechnol*. Feb. 2011;29(2):143-8. doi:10.1038/nbt.1755. Epub Dec. 22, 2010. cited by applicant

Miller et al., An improved zinc-finger nuclease architecture for highly specific genome editing. *Nat Biotechnol*. Jul. 2007;25(7):778-85. Epub Jul. 1, 2007. cited by applicant

Miller et al., Construction and properties of retrovirus packaging cells based on gibbon ape leukemia virus. *J Virol*. May 1991;65(5):2220-4. doi: 10.1128/JVI.65.5.2220-2224.1991. cited by applicant

Miller et al., Continuous evolution of SpCas9 variants compatible with non-G PAMs. *Nat Biotechnol*. Apr. 2020;38(4):471-481. doi: 10.1038/s41587-020-0412-8. Epub Feb. 10, 2020. cited by applicant

Miller, Human gene therapy comes of age. *Nature*. Jun. 11, 1992;357(6378):455-60. doi: 10.1038/357455a0. cited by applicant

Mills et al., Protein splicing in trans by purified—and C-terminal fragments of the *Mycobacterium tuberculosis* RecA intein. *Proc Natl Acad Sci U S A*. Mar. 31, 1998;95(7):3543-8. doi: 10.1073/pnas.95.7.3543. cited by applicant

Minoche et al., Evaluation of genomic high-throughput sequencing data generated on Illumina HiSeq and genome analyzer systems. *Genome Biol*. Nov. 8, 2011;12(11):R112. doi: 10.1186/GB-2011-12-11-r112. cited by applicant

Minoretto et al., A W148R mutation in the human FOXD4 gene segregating with dilated cardiomyopathy, obsessive-compulsive disorder, and suicidality. *Int J Mol Med*. Mar. 2007;19(3):369-72. cited by applicant

Mir et al., Two Active Site Divalent Ions in the Crystal Structure of the Hammerhead Ribozyme Bound to a Transition State Analogue. *Biochemistry* . . . Feb. 2, 2016;55(4):633-6. doi: 10.1021/acs.biochem.5b01139. Epub Jan. 19, 2016. cited by applicant

Mir et al., Type II-C CRISPR-Cas9 Biology, Mechanism, and Application. *ACS Chem Biol*. Feb. 16, 2018;13(2):357-365. doi: 10.1021/acscchembio.7b00855. Epub Dec. 20, 2017. cited by applicant

Mishina et al., Conditional gene targeting on the pure C57BL/6 genetic background. *Neurosci Res*. Jun. 2007;58(2):105-12. doi: 10.1016/j.neures.2007.01.004. Epub Jan. 18, 2007. cited by applicant

Mitani et al., Delivering therapeutic genes—matching approach and application. *Trends Biotechnol*. May 1993;11(5):162-6. doi: 10.1016/0167-7799(93)90108-L. cited by applicant

Mitton-Fry et al., Poly(A) tail recognition by a viral RNA element through assembly of a triple helix. *Science*. Nov. 26, 2010;330(6008):1244-7. doi: 10.1126/science.1195858. cited by applicant

Miyaoka et al., Systematic quantification of HDR and NHEJ reveals effects of locus, nuclease, and cell type on genome-editing. *Sci Rep*. Mar. 31, 2016;6:23549. doi: 10.1038/srep23549. cited by applicant

Moede et al., Identification of a nuclear localization signal, RRMKWKK, in the homeodomain transcription factor PDX-1. *FEBS Lett*. Nov. 19, 1999;461(3):229-34. doi: 10.1016/s0014-5793(99)01446-5. cited by applicant

Mohr et al., A Reverse Transcriptase-Cas1 Fusion Protein Contains a Cas6 Domain Required for Both CRISPR RNA

Biogenesis and RNA Spacer Acquisition. *Mol Cell*. Nov. 15, 2018;72(4):700-714.e8. doi: 10.1016/j.molcel.2018.09.013. Epub Oct. 18, 2018. Including Supplemental Information. cited by applicant

Mohr et al., Thermostable group II intron reverse transcriptase fusion proteins and their use in cDNA synthesis and next-generation RNA sequencing. *RNA*. Jul. 2013;19(7):958-70. doi: 10.1261/rna.039743.113. Epub May 22, 2013. cited by applicant

Mojica et al., Intervening sequences of regularly spaced prokaryotic repeats derive from foreign genetic elements. *J Mol Evol*. Feb. 2005;60(2):174-82. cited by applicant

Mol et al., Crystal structure and mutational analysis of human uracil-DNA glycosylase: structural basis for specificity and catalysis. *Cell*. Mar. 24, 1995;80(6):869-78. doi: 10.1016/0092-8674(95)90290-2. cited by applicant

Mol et al., Crystal structure of human uracil-DNA glycosylase in complex with a protein inhibitor: protein mimicry of DNA. *Cell*. Sep. 8, 1995;82(5):701-8. cited by applicant

Molla et al., CRISPR/Cas-Mediated Base Editing: Technical Considerations and Practical Applications. *Trends Biotechnol*. Oct. 2019;37(10):1121-1142. doi: 10.1016/j.tibtech.2019.03.008. Epub Apr. 14, 2019. cited by applicant

Monahan et al., Site-specific incorporation of unnatural amino acids into receptors expressed in Mammalian cells. *Chem Biol*. Jun. 2003;10(6):573-80. cited by applicant

Monot et al., The specificity and flexibility of 11 reverse transcription priming at imperfect T-tracts. *PLoS Genet*. May 2013;9(5):e1003499. doi: 10.1371/journal.pgen.1003499. Epub May 9, 2013. cited by applicant

Montange et al., Structure of the S-adenosylmethionine riboswitch regulatory mRNA element. *Nature*. Jun. 29, 2006;441(7097):1172-5. cited by applicant

Moore et al., Improved somatic mutagenesis in zebrafish using transcription activator-like effector nucleases (TALENs). *PloS One*. 2012;7(5):e37877. Doi: 10.1371/journal.pone.0037877. Epub May 24, 2012. cited by applicant

Mootz et al., Conditional protein splicing: a new tool to control protein structure and function in vitro and in vivo. *J Am Chem Soc*. Sep. 3, 2003;125(35):10561-9. cited by applicant

Mootz et al., Protein splicing triggered by a small molecule. *J Am Chem Soc*. Aug. 7, 2002;124(31):9044-5 and Supporting Information. doi: 10.1021/ja0267690. 4 pages. cited by applicant

Mootz et al., Protein splicing triggered by a small molecule. *J Am Chem Soc*. Aug. 7, 2002;124(31):9044-5. cited by applicant

Morbitzer et al., Assembly of custom TALE-type DNA binding domains by modular cloning. *Nucleic Acids Res*. Jul. 2011;39(13):5790-9. doi: 10.1093/nar/gkr151. Epub Mar. 18, 2011. cited by applicant

Moreno-Mateos et al., CRISPRscan: designing highly efficient sgRNAs for CRISPR-Cas9 targeting in vivo. *Nat Methods*. Oct. 2015;12(10):982-8. doi: 10.1038/nmeth.3543. Epub Aug. 31, 2015. cited by applicant

Morita et al., The site-specific recombination system of actinophage TG1. *FEMS Microbiol Lett*. Aug. 2009;297(2):234-40. doi: 10.1111/j.1574-6968.2009.01683.x. cited by applicant

Morris et al., A peptide carrier for the delivery of biologically active proteins into mammalian cells. *Nat Biotechnol*. Dec. 2001;19(12):1173-6. cited by applicant

Moscou et al., A simple cipher governs DNA recognition by TAL effectors. *Science*. Dec. 11, 2009;326(5959):1501. doi: 10.1126/science.1178817. cited by applicant

Mougiakos et al., Characterizing a thermostable Cas9 for bacterial genome editing and silencing. *Nat Commun*. Nov. 21, 2017;8(1):1647. doi: 10.1038/s41467-017-01591-4. cited by applicant

Muir et al., Expressed protein ligation: a general method for protein engineering. *Proc Natl Acad Sci U S A*. Jun. 9, 1998;95(12):6705-10. doi: 10.1073/pnas.95.12.6705. cited by applicant

Muller et al., Nucleotide exchange and excision technology (NExT) DNA shuffling: a robust method for DNA fragmentation and directed evolution. *Nucleic Acids Res*. Aug. 1, 2005;33(13):e117. doi: 10.1093/nar/gni116. PMID: 16061932; PMCID: PMC1182171. cited by applicant

Muller, U.F., Design and Experimental Evolution of trans-Splicing Group I Intron Ribozymes. *Molecules*. Jan. 2, 2017;22(1):75. doi: 10.3390/molecules22010075. cited by applicant

Mullins et al., Transgenesis in nonmurine species. *Hypertension*. Oct. 1993;22(4):630-3. cited by applicant

Mumtsidu et al., Structural features of the single-stranded DNA-binding protein of Epstein-Barr virus. *J Struct Biol*. Feb. 2008;161(2):172-87. doi: 10.1016/j.jsb.2007.10.014. Epub Nov. 1, 2007. cited by applicant

Murphy, Phage recombinases and their applications. *Adv Virus Res*. 2012;83:367-414. doi: 10.1016/B978-0-12-394438-2.00008-6. Review. cited by applicant

Mussolino et al., A novel TALE nuclease scaffold enables high genome editing activity in combination with low toxicity. *Nucleic Acids Res*. Nov. 2011;39(21):9283-93. Doi: 10.1093/nar/gkr597. Epub Aug. 3, 2011. cited by applicant

Mussolino et al., TALE nucleases: tailored genome engineering made easy. *Curr Opin Biotechnol*. Oct. 2012;23(5):644-50. doi: 10.1016/j.copbio.2012.01.013. Epub Feb. 17, 2012. cited by applicant

Muzyczka et al., Adeno-associated virus (AAV) vectors: will they work? *J Clin Invest*. Oct. 1994;94(4):1351. doi: 10.1172/JCI117468. cited by applicant

Myerowitz et al., The major defect in Ashkenazi Jews with Tay-Sachs disease is an insertion in the gene for the alpha-chain of beta-hexosaminidase. *J Biol Chem*. Dec. 15, 1988;263(35):18587-9. cited by applicant

Myers et al., Insulin signal transduction and the IRS proteins. *Annu Rev Pharmacol Toxicol*. 1996;36:615-58. doi: 10.1146/annurev.pa.36.040196.003151. cited by applicant

Nabel et al., Direct gene transfer from immunotherapy and immunization. Trends Biotechnol. May 1993;11(5):211-5. doi: 10.1016/0167-7799(93)90117-R. cited by applicant

Nahar et al., A G-quadruplex motif at the 3' end of sgRNAs improves CRISPR-Cas9 based genome editing efficiency. Chem Commun (Camb). Mar. 7, 2018;54(19):2377-2380. doi: 10.1039/c7cc08893k. Epub Feb. 16, 2018. cited by applicant

Nahvi et al., Coenzyme B12 riboswitches are widespread genetic control elements in prokaryotes. Nucleic Acids Res. Jan. 2, 2004;32(1):143-50. cited by applicant

Nakade et al., Microhomology-mediated end-joining-dependent integration of donor DNA in cells and animals using TALENs and CRISPR/Cas9. Nat Commun. Nov. 20, 2014;5:5560. doi: 10.1038/ncomms6560. cited by applicant

Nakamura et al., Codon usage tabulated from international DNA sequence databases: status for the year 2000. Nucleic Acids Res. Jan. 1, 2000;28(1):292. doi: 10.1093/nar/28.1.292. cited by applicant

Naorem et al., DGR mutagenic transposition occurs via hypermutagenic reverse transcription primed by nicked template RNA. Proc Natl Acad Sci U S A. Nov. 21, 2017;114(47):E10187-E10195. doi: 10.1073/pnas.1715952114. Epub Nov. 6, 2017. cited by applicant

Narayanan et al., Clamping down on weak terminal base pairs: oligonucleotides with molecular caps as fidelity-enhancing elements at the 5'- and 3'-terminal residues. Nucleic Acids Res. May 20, 2004;32(9):2901-11. Print 2004. cited by applicant

Navaratnam et al., An overview of cytidine deaminases. Int J Hematol. Apr. 2006;83(3):195-200. cited by applicant

NCBI Reference Sequence: NM\_002427.3. Wu et al., May 3, 2014. 5 pages. cited by applicant

Neel et al., Riboswitches: Classification, function and in silico approach, International Journal of Pharma Sciences and Research. 2010;1(9):409-420. cited by applicant

Nelson et al., Filamentous phage DNA cloning vectors: a noninfective mutant with a nonpolar deletion in gene III. Virology. 1981; 108(2): 338-50. cited by applicant

Nelson et al., The unstable repeats—three evolving faces of neurological disease. Neuron. Mar. 6, 2013;77(5):825-43. doi: 10.1016/j.neuron.2013.02.022. cited by applicant

Nern et al., Multiple new site-specific recombinases for use in manipulating animal genomes. Proc Natl Acad Sci U S A. Aug. 23, 2011;108(34):14198-203. doi: 10.1073/pnas.1111704108. Epub Aug. 9, 2011. cited by applicant

Nguyen et al., Evolutionary drivers of thermoadaptation in enzyme catalysis. Science. Jan. 20, 2017;355(6322):289-294. doi: 10.1126/science.aah3717. Epub Dec. 22, 2016. cited by applicant

Nguyen et al., IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. Jan. 2015;32(1):268-74. doi: 10.1093/molbev/msu300. Epub Nov. 3, 2014. cited by applicant

Ni et al., A PCSK9-binding antibody that structurally mimics the EGF(A) domain of LDL-receptor reduces LDL cholesterol in vivo. J Lipid Res. 2011;52:76-86. cited by applicant

Ni et al., Nucleic acid aptamers: clinical applications and promising new horizons. Curr Med Chem. 2011;18(27):4206-14. Review. cited by applicant

Nishida et al., Targeted nucleotide editing using hybrid prokaryotic and vertebrate adaptive immune systems. Science. Sep. 16, 2016;353(6305):1248. pii: aaf8729. doi: 10.1126/science.aaf8729. Epub Aug. 4, 2016. cited by applicant

Nishikura, Functions and regulation of RNA editing by ADAR deaminases. Annu Rev Biochem. 2010;79:321-349. doi:10.1146/annurev-biochem-060208-105251. cited by applicant

Nishimasu et al., Crystal structure of Cas9 in complex with guide RNA and target DNA. Cell. Feb. 27, 2014;156(5):935-49. doi: 10.1016/j.cell.2014.02.001. Epub Feb. 13, 2014. cited by applicant

Nishimasu et al., Crystal Structure of *Staphylococcus aureus* Cas9. Cell. Aug. 27, 2015;162(5):1113-26. doi: 10.1016/j.cell.2015.08.007. cited by applicant

Nishimasu et al., Engineered CRISPR-Cas9 nuclease with expanded targeting space. Science. Sep. 21, 2018;361(6408):1259-1262. doi: 10.1126/science.aas9129. Epub Aug. 30, 2018. cited by applicant

Noack et al., Epitranscriptomics: A New Regulatory Mechanism of Brain Development and Function. Front Neurosci. Feb. 20, 2018;12:85. doi: 10.3389/fnins.2018.00085. 9 pages. cited by applicant

Nomura et al., Controlling Mammalian Gene Expression by Allosteric Hepatitis Delta Virus Ribozymes. ACS Synth Biol. Dec. 20, 2013;2(12):684-9. doi: 10.1021/sb400037a. Epub May 22, 2013. cited by applicant

Nomura et al., Synthetic mammalian riboswitches based on guanine aptazyme. Chem Commun (Camb). Jul. 21, 2012;48(57):7215-7. doi: 10.1039/c2cc33140c. Epub Jun. 13, 2012. cited by applicant

Noris et al., A phenylalanine-55 to serine amino-acid substitution in the human glycoprotein IX leucine-rich repeat is associated with Bernard-Soulier syndrome. Br J Haematol. May 1997;97(2):312-20. cited by applicant

Nottingham et al., RNA-seq of human reference RNA samples using a thermostable group II intron reverse transcriptase. RNA. Apr. 2016;22(4):597-613. doi: 10.1261/rna.055558.115. Epub Jan. 29, 2016. cited by applicant

Nowak et al., Characterization of single-stranded DNA-binding proteins from the psychrophilic bacteria *Desulfotalea psychrophila*, *Flavobacterium psychrophilum*, *Psychrobacter arcticus*, *Psychrobacter cryohalolentis*, *Psychromonas ingrahamii*, *Psychroflexus torquis*, and *Photobacterium profundum*. BMC Microbiol. Apr. 14, 2014;14:91. doi: 10.1186/1471-2180-14-91. cited by applicant

Nowak et al., Guide RNA Engineering for Versatile Cas9 Functionality. Nucleic Acids Res. Nov. 16, 2016;44(20):9555-9564. doi: 10.1093/nar/gkw908. Epub Oct. 12, 2016. cited by applicant

Nowak et al., Structural analysis of monomeric retroviral reverse transcriptase in complex with an RNA/DNA hybrid. *Nucleic Acids Res.* Apr. 1, 2013;41(6):3874-87. doi: 10.1093/nar/gkt053. Epub Feb. 4, 2013. cited by applicant

Numrych et al., A comparison of the effects of single-base and triple-base changes in the integrase arm-type binding sites on the site-specific recombination of bacteriophage lambda. *Nucleic Acids Res.* Jul. 11, 1990;18(13):3953-9. doi: 10.1093/nar/18.13.3953. cited by applicant

Nyerges et al., A highly precise and portable genome engineering method allows comparison of mutational effects across bacterial species. *Proc Natl Acad Sci U S A.* Mar. 1, 2016;113(9):2502-7. doi: 10.1073/pnas.1520040113. Epub Feb. 16, 2016. cited by applicant

O'Connell et al., Programmable RNA recognition and cleavage by CRISPR/Cas9. *Nature.* Dec. 11, 2014;516(7530):263-6. doi: 10.1038/nature13769. Epub Sep. 28, 2014. cited by applicant

O'Maille et al., Structure-based combinatorial protein engineering (Scope). *J Mol Biol.* Aug. 23, 2002;321(4):677-91. cited by applicant

Oakes et al., CRISPR-Cas9 Circular Permutants as Programmable Scaffolds for Genome Modification. *Cell.* Jan. 10, 2019;176(1-2):254-267.e16. doi: 10.1016/j.cell.2018.11.052. cited by applicant

Oakes et al., Profiling of engineering hotspots identifies an allosteric CRISPR-Cas9 switch. *Nat Biotechnol.* Jun. 2016;34(6):646-51. doi: 10.1038/nbt.3528. Epub May 2, 2016. cited by applicant

Oakes et al., Protein engineering of Cas9 for enhanced function. *Methods Enzymol.* 2014;546:491-511. cited by applicant

Odsbu et al., Specific-terminal interactions of the *Escherichia coli* SeqA protein are required to form multimers that restrain negative supercoils and form foci. *Genes Cells.* Nov. 2005;10(11):1039-49. cited by applicant

Oeemig et al., Solution structure of DnaE intein from *Nostoc punctiforme*: structural basis for the design of a new split intein suitable for site-specific chemical modification. *FEBS Lett.* May 6, 2009;583(9):1451-6. cited by applicant

Offord, *Advances in Genome Editing. The Scientist*, Apr. 20, 2016. <http://www.the-scientist.com/?articles.view/articleNo/45903/title/Advances-in-Genome-Editing/>. cited by applicant

Oh et al., Positional cloning of a gene for Hermansky-Pudlak syndrome, a disorder of cytoplasmic organelles. *Nat Genet.* Nov. 1996;14(3):300-6. doi: 10.1038/ng1196-300. cited by applicant

Ohe et al., Purification and properties of xanthine dehydrogenase from *Streptomyces cyanogenus*. *J Biochem.* Jul. 1979;86(1):45-53. cited by applicant

Olivares et al., Site-specific genomic integration produces therapeutic Factor IX levels in mice. *Nat Biotechnol.* Nov. 2002;20(11):1124-8. doi: 10.1038/nbt753. Epub Oct. 15, 2002. cited by applicant

Olorunniji et al., Purification and In Vitro Characterization of Zinc Finger Recombinases. *Methods Mol Biol.* 2017;1642:229-245. doi: 10.1007/978-1-4939-7169-5\_15. cited by applicant

Olorunniji et al., Site-specific recombinases: molecular machines for the Genetic Revolution. *Biochem J.* Mar. 15, 2016;473(6):673-84. doi: 10.1042/BJ20151112. cited by applicant

Olorunniji et al., Synapsis and catalysis by activated Tn3 resolvase mutants. *Nucleic Acids Res.* Dec. 2008;36(22):7181-91. doi: 10.1093/nar/gkn885. Epub Nov. 10, 2008. cited by applicant

Orlando et al., Zinc-finger nuclease-driven targeted integration into mammalian genomes using donors with limited chromosomal homology. *Nucleic Acids Res.* Aug. 2010;38(15):e152. doi: 10.1093/nar/gkq512. Epub Jun. 8, 2010. cited by applicant

Orthwein et al., A mechanism for the suppression of homologous recombination in G1 cells. *Nature.* Dec. 17, 2015;528(7582):422-6. doi: 10.1038/nature16142. Epub Dec. 9, 2015. cited by applicant

Ortiz-Urda et al., Stable nonviral genetic correction of inherited human skin disease. *Nat Med.* Oct. 2002;8(10):1166-70. doi: 10.1038/nm766. Epub Sep. 16, 2002. Erratum in: *Nat Med.* Feb. 2003;9(2):237. cited by applicant

Osborn et al., Base Editor Correction of COL7A1 in Recessive Dystrophic Epidermolysis Bullosa Patient-Derived Fibroblasts and iPSCs. *J Invest Dermatol.* Feb. 2020;140(2):338-347.e5. doi: 10.1016/j.jid.2019.07.701. Epub Aug. 19, 2019. cited by applicant

Osborn et al., TALEN-based gene correction for epidermolysis bullosa. *Mol Ther.* Jun. 2013;21(6):1151-9. doi: 10.1038/mt.2013.56. Epub Apr. 2, 2013. cited by applicant

Ostermeier et al., A combinatorial approach to hybrid enzymes independent of DNA homology. *Nat Biotechnol.* Dec. 1999;17(12):1205-9. cited by applicant

Ostertag et al., Biology of mammalian L1 retrotransposons. *Annu Rev Genet.* 2001;35:501-38. doi: 10.1146/annurev.genet.35.102401.091032. cited by applicant

Otomo et al., Improved segmental isotope labeling of proteins and application to a larger protein. *J Biomol NMR.* Jun. 1999;14(2):105-14. doi: 10.1023/a:1008308128050. cited by applicant

Otomo et al., NMR observation of selected segments in a larger protein: central-segment isotope labeling through intein-mediated ligation. *Biochemistry.* Dec. 7, 1999;38(49):16040-4. doi: 10.1021/bi991902j. cited by applicant

Otto et al., The probability of fixation in populations of changing size. *Genetics.* Jun. 1997;146(2):723-33. cited by applicant

Packer et al., Methods for the directed evolution of proteins. *Nat Rev Genet.* Jul. 2015;16(7):379-94. doi: 10.1038/nrg3927. Epub Jun. 9, 2015. cited by applicant

Packer et al., Phage-assisted continuous evolution of proteases with altered substrate specificity. *Nat Commun.* Oct. 16, 2017;8(1):956. doi: 10.1038/s41467-017-01055-9. cited by applicant

Paige et al., RNA mimics of green fluorescent protein. *Science*. Jul. 29, 2011;333(6042):642-6. doi:10.1126/science.1207339. cited by applicant

Paiva et al., Targeted protein degradation: elements of PROTAC design. *Curr Opin Chem Biol*. Jun. 2019;50:111-119. doi: 10.1016/j.cbpa.2019.02.022. Epub Apr. 17, 2019. cited by applicant

Pan et al., Biological and biomedical applications of engineered nucleases. *Mol Biotechnol*. Sep. 2013;55(1):54-62. doi: 10.1007/s12033-012-9613-9. cited by applicant

Paquet et al., Efficient introduction of specific homozygous and heterozygous mutations using CRISPR/Cas9. *Nature*. May 5, 2016;533(7601):125-9. doi: 10.1038/nature17664. Epub Apr. 27, 2016. cited by applicant

Park et al., Digenome-seq web tool for profiling CRISPR specificity. *Nat Methods*. May 30, 2017;14(6):548-549. doi: 10.1038/nmeth.4262. cited by applicant

Park et al., Highly efficient editing of the  $\beta$ -globin gene in patient-derived hematopoietic stem and progenitor cells to treat sickle cell disease. *Nucleic Acids Res*. Sep. 5, 2019;47(15):7955-7972. doi: 10.1093/nar/gkz475. cited by applicant

Park et al., Sendai virus, an RNA virus with no risk of genomic integration, delivers CRISPR/Cas9 for efficient gene editing. *Mol Ther Methods Clin Dev*. Aug. 24, 2016;3:16057. doi: 10.1038/mtm.2016.57. cited by applicant

Parker et al., Admixture mapping identifies a quantitative trait locus associated with FEV1/FVC in the COPD Gene Study. *Genet Epidemiol*. Nov. 2014;38(7):652-9. doi: 10.1002/gepi.21847. Epub Aug. 11, 2014. cited by applicant

Patel et al., Flap endonucleases pass 5'-flaps through a flexible arch using a disorder-thread-order mechanism to confer specificity for free 5'-ends. *Nucleic Acids Res*. May 2012;40(10):4507-19. doi: 10.1093/nar/gks051. Epub Feb. 8, 2012. cited by applicant

Pattanayak et al., Determining the specificities of TALENs, Cas9, and other genome-editing enzymes. *Methods Enzymol*. 2014;546:47-78. doi: 10.1016/B978-0-12-801185-0.00003-9. cited by applicant

Pattanayak et al., High-throughput profiling of off-target DNA cleavage reveals RNA-programmed Cas9 nuclease specificity. *Nat Biotechnol*. Sep. 2013;31(9):839-43. doi: 10.1038/nbt.2673. Epub Aug. 11, 2013. cited by applicant

Pattanayak et al., Revealing off-target cleavage specificities of zinc-finger nucleases by in vitro selection. *Nat Methods*. Aug. 7, 2011;8(9):765-70. doi: 10.1038/nmeth.1670. cited by applicant

Pavletich et al., Zinc finger-DNA recognition: crystal structure of a Zif268-DNA complex at 2.1 Å. *Science*. May 10, 1991;252(5007):809-17. cited by applicant

Pawson et al., Protein phosphorylation in signaling—50 years and counting. *Trends Biochem Sci*. Jun. 2005;30(6):286-90. doi: 10.1016/j.tibs.2005.04.013. cited by applicant

Pearl, Structure and function in the uracil-DNA glycosylase superfamily. *Mutat Res*. Aug. 30, 2000;460(3-4):165-81. cited by applicant

Peck et al., Directed evolution of a small-molecule-triggered intein with improved splicing properties in mammalian cells. *Chem Biol*. May 27, 2011;18(5):619-30. doi: 10.1016/j.chembiol.2011.02.014. cited by applicant

Pellenz et al., New human chromosomal safe harbor sites for genome engineering with CRISPR/Cas9, TAL effector and homing endonucleases. Aug. 20, 2018. bioRxiv doi: <https://doi.org/10.1101/396390>. cited by applicant

Pelletier, CRISPR-Cas systems for the study of the immune function. Nov. 15, 2016. <https://doi.org/10.1002/9780470015902.a0026896>. cited by applicant

Pennisi et al., The CRISPR craze. *Science*. Aug. 23, 2013;341(6148):833-6. doi: 10.1126/science.341.6148.833. cited by applicant

Pennisi et al., The tale of the TALEs. *Science*. Dec. 14, 2012;338(6113):1408-11. doi: 10.1126/science.338.6113.1408. cited by applicant

Perach et al., Catalytic features of the recombinant reverse transcriptase of bovine leukemia virus expressed in bacteria. *Virology*. Jun. 20, 1999;259(1):176-89. doi: 10.1006/viro.1999.9761. cited by applicant

Perez et al., Establishment of HIV-1 resistance in CD4<sup>+</sup> T cells by genome editing using zinc-finger nucleases. *Nat Biotechnol*. Jul. 2008;26(7):808-16. Doi: 10.1038/nbt1410. Epub Jun. 29, 2008. cited by applicant

Perez-Pinera et al., Advances in targeted genome editing. *Curr Opin Chem Biol*. Aug. 2012;16(3-4):268-77. doi: 10.1016/j.cbpa.2012.06.007. Epub Jul. 20, 2012. cited by applicant

Perez-Pinera et al., RNA-guided gene activation by CRISPR-Cas9-based transcription factors. *Nat Methods*. Oct. 2013;10(10):973-6. doi: 10.1038/nmeth.2600. Epub Jul. 25, 2013. cited by applicant

Perler et al., Protein splicing and autoprolysis mechanisms. *Curr Opin Chem Biol*. Oct. 1997;1(3):292-9. doi: 10.1016/s1367-5931(97)80065-8. cited by applicant

Perler et al., Protein splicing elements: inteins and exteins—a definition of terms and recommended nomenclature. *Nucleic Acids Res*. Apr. 11, 1994;22(7):1125-7. doi: 10.1093/nar/22.7.1125. cited by applicant

Perler, InBase, the New England Biolabs Intein Database. *Nucleic Acids Res*. Jan. 1, 1999;27(1):346-7. doi: 10.1093/nar/27.1.346. cited by applicant

Perler, Protein splicing of inteins and hedgehog autoprolysis: structure, function, and evolution. *Cell*. Jan. 9, 1998;92(1):1-4. doi: 10.1016/s0092-8674(00)80892-2. cited by applicant

Perreault et al., Mixed deoxyribo- and ribo-oligonucleotides with catalytic activity. *Nature*. Apr. 5, 1990;344(6266):565-7. doi: 10.1038/344565a0. cited by applicant

Petek et al., Frequent endonuclease cleavage at off-target locations in vivo. *Mol Ther*. May 2010;18(5):983-6. Doi: 10.1038/mt.2010.35. Epub Mar. 9, 2010. cited by applicant

Petersen-Mahrt et al., AID mutates *E. coli* suggesting a DNA deamination mechanism for antibody diversification. *Nature*. Jul. 4, 2002;418(6893):99-103. cited by applicant

Petolino et al., Editing Plant Genomes: a new era of crop improvement. *Plant Biotechnol J*. Feb. 2016;14(2):435-6. doi: 10.1111/pbi.12542. cited by applicant

Peyrottes et al., Oligodeoxynucleoside phosphoramidates (P-NH<sub>2</sub>): synthesis and thermal stability of duplexes with DNA and RNA targets. *Nucleic Acids Res*. May 15, 1996;24(10):1841-8. cited by applicant

Pfeiffer et al., Mechanisms of DNA double-strand break repair and their potential to induce chromosomal aberrations. *Mutagenesis*. Jul. 2000;15(4):289-302. doi: 10.1093/mutage/15.4.289. cited by applicant

Phillips, The challenge of gene therapy and DNA delivery. *J Pharm Pharmacol*. Sep. 2001;53(9):1169-74. cited by applicant

Pickart et al., Ubiquitin: structures, functions, mechanisms. *Biochim Biophys Acta*. Nov. 29, 2004;1695(1-3):55-72. doi: 10.1016/j.bbamcr.2004.09.019. cited by applicant

Pieken et al., Kinetic characterization of ribonuclease-resistant 2'-modified hammerhead ribozymes. *Science*. Jul. 19, 1991;253(5017):314-7. doi: 10.1126/science.1857967. cited by applicant

Pinkert et al., An albumin enhancer located 10 kb upstream functions along with its promoter to direct efficient, liver-specific expression in transgenic mice. *Genes Dev*. May 1987;1(3):268-76. doi: 10.1101/gad.1.3.268. cited by applicant

Pirakitikulr et al., PCRless library mutagenesis via oligonucleotide recombination in yeast. *Protein Sci*. Dec. 2010;19(12):2336-46. doi: 10.1002/pro.513. cited by applicant

Plasterk et al., DNA inversions in the chromosome of *Escherichia coli* and in bacteriophage Mu: relationship to other site-specific recombination systems. *Proc Natl Acad Sci U S A*. Sep. 1983;80(17):5355-8. cited by applicant

Plosky et al., CRISPR-Mediated Base Editing without DNA Double-Strand Breaks. *Mol Cell*. May 19, 2016;62(4):477-8. doi: 10.1016/j.molcel.2016.05.006. cited by applicant

Pluciennik et al., PCNA function in the activation and strand direction of MutL $\alpha$  endonuclease in mismatch repair. *Proc Natl Acad Sci U S A*. Sep. 14, 2010;107(37):16066-71. doi: 10.1073/pnas.1010662107. Epub Aug. 16, 2010. cited by applicant

Poller et al., A leucine-to-proline substitution causes a defective alpha 1-antichymotrypsin allele associated with familial obstructive lung disease. *Genomics*. Sep. 1993;17(3):740-3. cited by applicant

Popp et al., Sortagging: a versatile method for protein labeling. *Nat Chem Biol*. Nov. 2007;3(11):707-8. doi: 10.1038/nchembio.2007.31. Epub Sep. 23, 2007. cited by applicant

Porteus, Design and testing of zinc finger nucleases for use in mammalian cells. *Methods Mol Biol*. 2008;435:47-61. doi: 10.1007/978-1-59745-232-8\_4. cited by applicant

Posnick et al., Imbalanced base excision repair increases spontaneous mutation and alkylation sensitivity in *Escherichia coli*. *J Bacteriol*. Nov. 1999;181(21):6763-71. cited by applicant

Pospíšilová et al., Hydrolytic cleavage of N6-substituted adenine derivatives by eukaryotic adenine and adenosine deaminases. *Biosci Rep*. 2008;28(6):335-347. doi:10.1042/BSR20080081. cited by applicant

Pourcel et al., CRISPR elements in *Yersinia pestis* acquire new repeats by preferential uptake of bacteriophage DNA, and provide additional tools for evolutionary studies. *Microbiology*. Mar. 2005;151(Pt 3):653-63. cited by applicant

Prasad et al., Rev1 is a base excision repair enzyme with 5'-deoxyribose phosphate lyase activity. *Nucleic Acids Res*. Dec. 15, 2016;44(22):10824-10833. doi: 10.1093/nar/gkw869. Epub Sep. 28, 2016. cited by applicant

Prashant et al., CAS9 transcriptional activators for target specificity screening and paired nickases for cooperative genome engineering. *Nature Biotechnology* 2013;31(9):833-8. cited by applicant

Prorocic et al., Zinc-finger recombinase activities in vitro. *Nucleic Acids Res*. Nov. 2011;39(21):9316-28. doi: 10.1093/nar/gkr652. Epub Aug. 17, 2011. cited by applicant

Proudfoot et al., Zinc finger recombinases with adaptable DNA sequence specificity. *PLoS One*. Apr. 29, 2011;6(4):e19537. doi: 10.1371/journal.pone.0019537. cited by applicant

Pruschy et al., Mechanistic studies of a signaling pathway activated by the organic dimerizer FK1012. *Chem Biol*. Nov. 1994;1(3):163-72. doi: 10.1016/1074-5521(94)90006-x. cited by applicant

Prykhodzhiy et al., CRISPR multitargeter: a web tool to find common and unique CRISPR single guide RNA targets in a set of similar sequences. *PLoS One*. Mar. 5, 2015;10(3):e0119372. doi: 10.1371/journal.pone.0119372. eCollection 2015. cited by applicant

Pu et al., Evolution of a split RNA polymerase as a versatile biosensor platform. *Nat Chem Biol*. Apr. 2017;13(4):432-438. doi: 10.1038/nchembio.2299. Epub Feb. 13, 2017. cited by applicant

Putnam et al., Protein mimicry of DNA from crystal structures of the uracil-DNA glycosylase inhibitor protein and its complex with *Escherichia coli* uracil-DNA glycosylase. *J Mol Biol*. Mar. 26, 1999;287(2):331-46. cited by applicant

Qi et al., Engineering naturally occurring trans-acting non-coding RNAs to sense molecular signals. *Nucleic Acids Res*. Jul. 2012;40(12):5775-86. doi: 10.1093/nar/gks168. Epub Mar. 1, 2012. cited by applicant

Qi et al., Repurposing CRISPR as an RNA-guided platform for sequence-specific control of gene expression. *Cell*. Feb. 28, 2013;152(5):1173-83. doi: 10.1016/j.cell.2013.02.022. cited by applicant

Qu et al., Global mapping of binding sites for phic31 integrase in transgenic maden-darby bovine kidney cells using ChIP-seq. *Hereditas*. Jan. 14, 2019;156:3. doi: 10.1186/s41065-018-0079-z. cited by applicant

Queen et al., Immunoglobulin gene transcription is activated by downstream sequence elements. *Cell*. Jul.



1983;33(3):741-8. doi: 10.1016/0092-8674(83)90016-8. cited by applicant

Radany et al., Increased spontaneous mutation frequency in human cells expressing the phage PBS2-encoded inhibitor of uracil-DNA glycosylase. *Mutat Res. Sep. 15, 2000;461(1):41-58.* doi: 10.1016/s0921-8777(00)00040-9. cited by applicant

Raghavan et al., Abstract 27: Therapeutic Targeting of Human Lipid Genes with in vivo CRISPR-Cas9 Genome Editing. Oral Abstract Presentations: Lipoprotein Metabolism and Therapeutic Targets. *Arterioscler THromb Vasc Biol.* 2015;35(Suppl. 1):Abstract 27. 5 pages. cited by applicant

Raillard et al., Targeting sites within HIV-1 cDNA with a DNA-cleaving ribozyme. *Biochemistry. Sep. 10, 1996;35(36):11693-701.* doi: 10.1021/bi960845g. cited by applicant

Raina et al., PROTAC-induced BET protein degradation as a therapy for castration-resistant prostate cancer. *Proc Natl Acad Sci U S A. Jun. 28, 2016;113(26):7124-9.* doi: 10.1073/pnas.1521738113. Epub Jun. 6, 2016. cited by applicant

Rakonjac et al., Roles of PIII in filamentous phage assembly. *J Mol Biol.* 1998; 282(1)25-41. cited by applicant

Ramakrishna et al., Gene disruption by cell-penetrating peptide-mediated delivery of Cas9 protein and guide RNA. *Genome Res. Jun. 2014;24(6):1020-7.* doi: 10.1101/gr.171264.113. Epub Apr. 2, 2014. cited by applicant

Ramamurthy et al., Identification of immunogenic B-cell epitope peptides of rubella virus E1 glycoprotein towards development of highly specific immunoassays and/or vaccine. Conference Abstract. 2019. cited by applicant

Ramirez et al., Engineered zinc finger nickases induce homology-directed repair with reduced mutagenic effects. *Nucleic Acids Res. Jul. 2012;40(12):5560-8.* doi: 10.1093/nar/gks179. Epub Feb. 28, 2012. cited by applicant

Ramirez et al., Unexpected failure rates for modular assembly of engineered zinc fingers. *Nat Methods.* May 2008;5(5):374-5. Doi: 10.1038/nmeth0508-374. cited by applicant

Ran et al., Double Nicking by RNA-guided CRISPR Cas9 for Enhanced Genome Editing Specificity. *Cell. Sep. 12, 2013;154(6):1380-9.* doi: 10.1016/j.cell.2013.08.021. Epub Aug. 29, 2013. cited by applicant

Ran et al., Genome engineering using the CRISPR-Cas9 system. *Nat Protoc. Nov. 2013;8(11):2281-308.* doi: 10.1038/nprot.2013.143. Epub Oct. 24, 2013. cited by applicant

Ran et al., In vivo genome editing using *Staphylococcus aureus* Cas9. *Nature.* Apr. 9, 2015;520(7546):186-91. doi: 10.1038/nature14299. Epub Apr. 1, 2015. cited by applicant

Ranzau et al., Genome, Epigenome, and Transcriptome Editing via Chemical Modification of Nucleobases in Living Cells. *Biochemistry.* Feb. 5, 2019;58(5):330-335. doi: 10.1021/acs.biochem.8b00958. Epub Dec. 12, 2018. cited by applicant

Rashel et al., A novel site-specific recombination system derived from bacteriophage phiMR11. *Biochem Biophys Res Commun.* Apr. 4, 2008;368(2):192-8. doi: 10.1016/j.bbrc.2008.01.045. Epub Jan. 22, 2008. cited by applicant

Rasila et al., Critical evaluation of random mutagenesis by error-prone polymerase chain reaction protocols, *Escherichia coli* mutator strain, and hydroxylamine treatment. *Anal Biochem.* May 1, 2009;388(1):71-80. doi: 10.1016/j.ab.2009.02.008. Epub Feb. 10, 2009. cited by applicant

Raskin et al., Substitution of a single bacteriophage T3 residue in bacteriophage T7 RNA polymerase at position 748 results in a switch in promoter specificity. *J Mol Biol.* Nov. 20, 1992;228(2):506-15. cited by applicant

Raskin et al., T7 RNA polymerase mutants with altered promoter specificities. *Proc Natl Acad Sci U S A.* Apr. 15, 1993;90(8):3147-51. cited by applicant

Rath et al., Fidelity of end joining in mammalian episomes and the impact of Metnase on joint processing. *BMC Mol Biol.* Mar. 22, 2014;15:6. doi: 10.1186/1471-2199-15-6. cited by applicant

Rauch et al., Programmable RNA Binding Proteins for Imaging and Therapeutics. *Biochemistry.* Jan. 30, 2018;57(4):363-364. doi: 10.1021/acs.biochem.7b01101. Epub Nov. 17, 2017. cited by applicant

Ravishankar et al., X-ray analysis of a complex of *Escherichia coli* uracil DNA glycosylase (EcUDG) with a proteinaceous inhibitor. The structure elucidation of a prokaryotic UDG. *Nucleic Acids Res.* 26 (21): 4880-4887 (1998). cited by applicant

Ray et al., A compendium of RNA-binding motifs for decoding gene regulation. *Nature.* Jul. 11, 2013;499(7457):172-7. doi: 10.1038/nature12311. cited by applicant

Ray et al., Homologous recombination: ends as the means. *Trends Plant Sci.* Oct. 2002;7(10):435-40. cited by applicant

Rebar et al., Phage display methods for selecting zinc finger proteins with novel DNA-binding specificities. *Methods Enzymol.* 1996;267:129-49. cited by applicant

Rebuzzini et al., New mammalian cellular systems to study mutations introduced at the break site by non-homologous end-joining. *DNA Repair (Amst).* May 2, 2005;4(5):546-55. cited by applicant

Rees et al., Analysis and minimization of cellular RNA editing by DNA adenine base editors. *Sci Adv.* May 8, 2019;5(5):eaax5717. doi: 10.1126/sciadv.aax5717. cited by applicant

Rees et al., Base editing: precision chemistry on the genome and transcriptome of living cells. *Nat Rev Genet.* Dec. 2018;19(12):770-788. doi: 10.1038/s41576-018-0059-1. cited by applicant

Rees et al., Development of hRad51-Cas9 nickase fusions that mediate HDR without double-stranded breaks. *Nat Commun.* May 17, 2019;10(1):2212. doi: 10.1038/s41467-019-09983-4. cited by applicant

Rees et al., Improving the DNA specificity and applicability of base editing through protein engineering and protein delivery. *Nat Commun.* Jun. 6, 2017;8:15790. doi: 10.1038/ncomms15790. cited by applicant

Relph et al., Recent developments and current status of gene therapy using viral vectors in the United Kingdom. *BMJ.* 2004;329(7470):839-842. doi: 10.1136/bmj.329.7470.839. cited by applicant

Remy et al., Gene transfer with a series of lipophilic DNA-binding molecules. *Bioconjug Chem.* Nov.-Dec. 1994;5(6):647-54. doi: 10.1021/bc00030a021. cited by applicant

Ren et al., In-line Alignment and Mg.SUP.2.? Coordination at the Cleavage Site of the env22 Twister Ribozyme. *Nat Commun.* Nov. 20, 2014;5:5534. doi: 10.1038/ncomms6534. cited by applicant

Ren et al., Pistol Ribozyme Adopts a Pseudoknot Fold Facilitating Site-Specific In-Line Cleavage. *Nat Chem Biol.* Sep. 2016;12(9):702-8. doi: 10.1038/nchembio.2125. Epub Jul. 11, 2016. cited by applicant

Reynaud et al., What role for AID: mutator, or assembler of the immunoglobulin mutasome? *Nat Immunol.* Jul. 2003;4(7):631-8. cited by applicant

Reyon et al., FLASH assembly of TALENs for high-throughput genome editing. *Nat Biotechnol.* May 2012;30(5):460-5. doi: 10.1038/nbt.2170. cited by applicant

Ribeiro et al., Protein Engineering Strategies to Expand CRISPR-Cas9 Applications. *Int J Genomics.* Aug. 2, 2018;2018:1652567. doi: 10.1155/2018/1652567. cited by applicant

Richardson et al., Enhancing homology-directed genome editing by catalytically active and inactive CRISPR-Cas9 using asymmetric donor DNA. *Nat Biotechnol.* Mar. 2016;34(3):339-44. doi: 10.1038/nbt.3481. Epub Jan. 20, 2016. cited by applicant

Richter et al., Function and regulation of clustered regularly interspaced short palindromic repeats (CRISPR) / CRISPR associated (Cas) systems. *Viruses.* Oct. 19, 2012;4(10):2291-311. doi: 10.3390/v4102291. cited by applicant

Riechmann et al., The C-terminal domain of TolA is the coreceptor for filamentous phage infection of *E. coli*. *Cell.* 1997;90(2):351-60. PMID:9244308. cited by applicant

Ringrose et al., The Kw recombinase, an integrase from *Kluyveromyces waltii*. *Eur J Biochem.* Sep. 15, 1997;248(3):903-12. doi: 10.1111/j.1432-1033.1997.00903.x. cited by applicant

Risso et al., Hyperstability and substrate promiscuity in laboratory resurrections of Precambrian  $\beta$ -lactamases. *J Am Chem Soc.* Feb. 27, 2013;135(8):2899-902. doi: 10.1021/ja311630a. Epub Feb. 14, 2013. cited by applicant

Ritchie et al., limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res.* Apr. 20, 2015;43(7):e47. doi: 10.1093/nar/gkv007. Epub Jan. 20, 2015. cited by applicant

Robertson et al., DNA repair in mammalian cells: Base excision repair: the long and short of it. *Cell Mol Life Sci.* Mar. 2009;66(6):981-93. doi: 10.1007/s00018-009-8736-z. cited by applicant

Robertson et al., Selection in vitro of an RNA enzyme that specifically cleaves single-stranded DNA. *Nature.* Mar. 29, 1990;344(6265):467-8. doi: 10.1038/344467a0. cited by applicant

Robinson et al., The protein tyrosine kinase family of the human genome. *Oncogene.* Nov. 20, 2000;19(49):5548-57. doi: 10.1038/sj.onc.1203957. cited by applicant

Rogozin et al., Evolution and diversification of lamprey antigen receptors: evidence for involvement of an AID-APOBEC family cytosine deaminase. *Nat Immunol.* Jun. 2007;8(6):647-56. doi: 10.1038/ni1463. Epub Apr. 29, 2007. cited by applicant

Rong et al., Homologous recombination in human embryonic stem cells using CRISPR/Cas9 nickase and a long DNA donor template. *Protein Cell.* Apr. 2014;5(4):258-60. doi: 10.1007/s13238-014-0032-5. cited by applicant

Rongrong et al., Effect of deletion mutation on the recombination activity of Cre recombinase. *Acta Biochim Pol.* 2005;52(2):541-4. Epub May 15, 2005. cited by applicant

Roth et al., A widespread self-cleaving ribozyme class is revealed by bioinformatics. *Nat Chem Biol.* Jan. 2014;10(1):56-60. doi: 10.1038/nchembio.1386. Epub Nov. 17, 2013. cited by applicant

Roth et al., Purification and characterization of murine retroviral reverse transcriptase expressed in *Escherichia coli*. *J Biol Chem.* Aug. 5, 1985;260(16):9326-35. cited by applicant

Rouet et al., Expression of a site-specific endonuclease stimulates homologous recombination in mammalian cells. *Proc Natl Acad Sci U S A.* Jun. 21, 1994;91(13):6064-8. doi: 10.1073/pnas.91.13.6064. cited by applicant

Rouet et al., Introduction of double-strand breaks into the genome of mouse cells by expression of a rare-cutting endonuclease. *Mol Cell Biol.* Dec. 1994;14(12):8096-106. doi: 10.1128/mcb.14.12.8096. cited by applicant

Rouet et al., Receptor-Mediated Delivery of CRISPR-Cas9 Endonuclease for Cell-Type-Specific Gene Editing. *J Am Chem Soc.* May 30, 2018;140(21):6596-6603. doi: 10.1021/jacs.8b01551. Epub May 18, 2018. cited by applicant

Roundtree et al., YTHDC1 mediates nuclear export of N6-methyladenosine methylated mRNAs. *Elife.* Oct. 6, 2017;6:e31311. doi: 10.7554/eLife.31311. cited by applicant

Rowland et al., Regulatory mutations in Sin recombinase support a structure-based model of the synaptosome. *Mol Microbiol.* Oct. 2009;74(2):282-98. doi: 10.1111/j.1365-2958.2009.06756.x. Epub Jun. 8, 2009. cited by applicant

Rowland et al., Sin recombinase from *Staphylococcus aureus*: synaptic complex architecture and transposon targeting. *Mol Microbiol.* May 2002;44(3):607-19. doi: 10.1046/j.1365-2958.2002.02897.x. cited by applicant

Rowley, Chromosome translocations: dangerous liaisons revisited. *Nat Rev Cancer.* Dec. 2001;1(3):245-50. doi: 10.1038/35106108. cited by applicant

Rubio et al., An adenosine-to-inosine tRNA-editing enzyme that can perform C-to-U deamination of DNA. *Proc Natl Acad Sci U S A.* May 8, 2007;104(19):7821-6. doi: 10.1073/pnas.0702394104. Epub May 1, 2007. PMID: 17483465; PMCID: PMC1876531. cited by applicant

Rubio et al., Transfer RNA travels from the cytoplasm to organelles. *Wiley Interdiscip Rev RNA.* Nov.-Dec. 2011;2(6):802-17. doi: 10.1002/wrna.93. Epub Jul. 11, 2011. cited by applicant

Rudolph et al., Synthetic ribwitches for the conditional control of gene expression in *Streptomyces coelicolor*. *Microbiology*. Jul. 2013;159(Pt 7):1416-22. doi: 10.1099/mic.0.067322-0. Epub May 15, 2013. cited by applicant

Rutherford et al., Attachment site recognition and regulation of directionality by the serine integrases. *Nucleic Acids Res*. Sep. 2013;41(17):8341-56. doi: 10.1093/nar/gkt580. Epub Jul. 2, 2013. cited by applicant

Ryu et al., Adenine base editing in mouse embryos and an adult mouse model of Duchenne muscular dystrophy. *Nat Biotechnol*. Jul. 2018;36(6):536-539. doi: 10.1038/nbt.4148. Epub Apr. 27, 2018. cited by applicant

Rüfer et al., Non-contact positions impose site selectivity on Cre recombinase. *Nucleic Acids Res*. Jul. 1, 2002;30(13):2764-71. doi: 10.1093/nar/gkf399. cited by applicant

Sadelain et al., Safe harbours for the integration of new DNA in the human genome. *Nat Rev Cancer*. Dec. 1, 2011;12(1):51-8. doi: 10.1038/nrc3179. cited by applicant

Sadowski, The FLP recombinase of the 2-microns plasmid of *Saccharomyces cerevisiae*. *Prog Nucleic Acid Res Mol Biol*. 1995;51:53-91. cited by applicant

Safari et al., CRISPR Cpf1 proteins: structure, function and implications for genome editing. *Cell Biosci*. May 9, 2019;9:36. doi: 10.1186/s13578-019-0298-7. cited by applicant

Sage et al., Proliferation of functional hair cells in vivo in the absence of the retinoblastoma protein. *Science*. Feb. 18, 2005;307(5712):1114-8. Epub Jan. 13, 2005. cited by applicant

Sakuma et al., MMEJ-assisted gene knock-in using TALENs and CRISPR-Cas9 with the PITCh systems. *Nat Protoc*. Jan. 2016;11(1):118-33. doi: 10.1038/nprot.2015.140. Epub Dec. 17, 2015. cited by applicant

Sale et al., Y-family DNA polymerases and their role in tolerance of cellular DNA damage. *Nat Rev Mol Cell Biol*. Feb. 23, 2012;13(3):141-52. doi: 10.1038/nrm3289. cited by applicant

Saleh-Gohari et al., Conservative homologous recombination preferentially repairs DNA double-strand breaks in the S phase of the cell cycle in human cells. *Nucleic Acids Res*. Jul. 13, 2004;32(12):3683-8. Print 2004. cited by applicant

Samal et al., Cationic polymers and their therapeutic potential. *Chem Soc Rev*. Nov. 7, 2012;41(21):7147-94. doi: 10.1039/c2cs35094g. Epub Aug. 10, 2012. cited by applicant

Samanta et al., A reverse transcriptase ribozyme. *Elife*. Sep. 26, 2017;6:e31153. doi: 10.7554/eLife.31153. cited by applicant

Samulski et al., Helper-free stocks of recombinant adeno-associated viruses: normal integration does not require viral gene expression. *J Virol*. Sep. 1989;63(9):3822-8. doi: 10.1128/JVI.63.9.3822-3828.1989. cited by applicant

Sander et al., CRISPR-Cas systems for editing, regulating and targeting genomes. *Nat Biotechnol*. Apr. 2014;32(4):347-55. doi: 10.1038/nbt.2842. Epub Mar. 2, 2014. cited by applicant

Sander et al., In silico abstraction of zinc finger nuclease cleavage profiles reveals an expanded landscape of off-target sites. *Nucleic Acids Res*. Oct. 2013;41(19):e181. doi: 10.1093/nar/gkt716. Epub Aug. 14, 2013. cited by applicant

Sander et al., Targeted gene disruption in somatic zebrafish cells using engineered TALENs. *Nat Biotechnol*. Aug. 5, 2011;29(8):697-8. doi: 10.1038/nbt.1934. cited by applicant

Sang et al., A unique uracil-DNA binding protein of the uracil DNA glycosylase superfamily. *Nucleic Acids Res*. Sep. 30, 2015;43(17):8452-63. doi: 10.1093/nar/gkv854. Epub Aug. 24, 2015. cited by applicant

Sang, Prospects for transgenesis in the chick. *Mech Dev*. Sep. 2004;121(9):1179-86. cited by applicant

Sanjana et al., A transcription activator-like effector toolbox for genome engineering. *Nat Protoc*. Jan. 5, 2012;7(1):171-92. doi: 10.1038/nprot.2011.431. cited by applicant

Santiago et al., Targeted gene knockout in mammalian cells by using engineered zinc-finger nucleases. *Proc Natl Acad Sci U S A*. Apr. 15, 2008;105(15):5809-14. doi: 10.1073/pnas.0800940105. Epub Mar. 21, 2008. cited by applicant

Santoro et al., Directed evolution of the site specificity of Cre recombinase. *Proc Natl Acad Sci U S A*. Apr. 2, 2002;99(7):4185-90. Epub Mar. 19, 2002. cited by applicant

Saparbaev et al., Excision of hypoxanthine from DNA containing dIMP residues by the *Escherichia coli*, yeast, rat, and human alkylpurine DNA glycosylases. *Proc Natl Acad Sci U S A*. Jun. 21, 1994;91(13):5873-7. doi: 10.1073/pnas.91.13.5873. cited by applicant

Sapranaukas et al., The *Streptococcus thermophilus* CRISPR/Cas system provides immunity in *Escherichia coli*. *Nucleic Acids Res*. Nov. 2011;39(21):9275-82. doi: 10.1093/nar/gkr606. Epub Aug. 3, 2011. cited by applicant

Sapunar et al., Dorsal root ganglion—a potential new therapeutic target for neuropathic pain. *J Pain Res*. 2012;5:31-8. doi: 10.2147/JPR.S26603. Epub Feb. 16, 2012. cited by applicant

Saraconi et al., The RNA editing enzyme APOBEC1 induces somatic mutations and a compatible mutational signature is present in esophageal adenocarcinomas. *Genome Biol*. Jul. 31, 2014;15(7):417. doi: 10.1186/s13059-014-0417-z. cited by applicant

Sarkar et al., HIV-1 proviral DNA excision using an evolved recombinase. *Science*. Jun. 29, 2007;316(5833):1912-5. doi: 10.1126/science.1141453. cited by applicant

Sashital et al., Mechanism of foreign DNA selection in a bacterial adaptive immune system. *Mol Cell*. Jun. 8, 2012;46(5):606-15. doi: 10.1016/j.molcel.2012.03.020. Epub Apr. 19, 2012. cited by applicant

Sasidharan et al., The selection of acceptable protein mutations. *PNAS*; Jun. 12, 2007;104(24):10080-5. [www.pnas.org/cgi/doi/10.1073.pnas.0703737104](http://www.pnas.org/cgi/doi/10.1073.pnas.0703737104). cited by applicant

Satomura et al., Precise genome-wide base editing by the CRISPR Nickase system in yeast. *Sci Rep*. May 18, 2017;7(1):2095. doi: 10.1038/s41598-017-02013-7. cited by applicant

Saudek et al., A preliminary trial of the programmable implantable medication system for insulin delivery. *Engl J Med*. Aug. 31, 1989;321(9):574-9. cited by applicant

Sauer et al., DNA recombination with a heterospecific Cre homolog identified from comparison of the pac-cl regions of P1-related phages. *Nucleic Acids Res*. Nov. 18, 2004;32(20):6086-95. doi: 10.1093/nar/gkh941. cited by applicant

Savic et al., Covalent linkage of the DNA repair template to the CRISPR-Cas9 nuclease enhances homology-directed repair. *Elife*. May 29, 2018;7:e33761. doi: 10.7554/eLife.33761. cited by applicant

Saville et al., A site-specific self-cleavage reaction performed by a novel RNA in *Neurospora* mitochondria. *Cell*. May 18, 1990;61(4):685-96. doi: 10.1016/0092-8674(90)90480-3. cited by applicant

Savva et al., The structural basis of specific base-excision repair by uracil-DNA glycosylase. *Nature*. Feb. 9, 1995;373(6514):487-93. doi: 10.1038/373487a0. cited by applicant

Schaaper et al., Base selection, proofreading, and mismatch repair during DNA replication in *Escherichia coli*. *J Biol Chem*. Nov. 15, 1993;268(32):23762-5. cited by applicant

Schaaper et al., Spectra of spontaneous mutations in *Escherichia coli* strains defective in mismatch correction: the nature of in vivo DNA replication errors. *Proc Natl Acad Sci U S A*. Sep. 1987;84(17):6220-4. cited by applicant

Schaefer et al., Understanding RNA modifications: the promises and technological bottlenecks of the 'epitranscriptome'. *Open Biol*. May 2017;7(5):170077. doi: 10.1098/rsob.170077. cited by applicant

Schechner et al., Multiplexable, locus-specific targeting of long RNAs with CRISPR-Display. *Nat Methods*. Jul. 2015;12(7):664-70. doi: 10.1038/nmeth.3433. Epub Jun. 1, 2015. Author manuscript entitled CRISPR Display: A modular method for locus-specific targeting of long noncoding RNAs and synthetic RNA devices in vivo. cited by applicant

Schek et al., Definition of the upstream efficiency element of the simian virus 40 late polyadenylation signal by using in vitro analyses. *Mol Cell Biol*. Dec. 1992;12(12):5386-93. doi: 10.1128/mcb.12.12.5386. cited by applicant

Schenk et al., MPDU1 mutations underlie a novel human congenital disorder of glycosylation, designated type If. *J Clin Invest*. Dec. 2001;108(11):1687-95. doi: 10.1172/JCI13419. cited by applicant

Schmitz et al., Behavioral abnormalities in prion protein knockout mice and the potential relevance of PrP(C) for the cytoskeleton. *Prion*. 2014;8(6):381-6. doi: 10.4161/19336896.2014.983746. cited by applicant

Schriefer et al., Low pressure DNA shearing: a method for random DNA sequence analysis. *Nucleic Acids Res*. Dec. 25, 1990;18(24):7455-6. cited by applicant

Schultz et al., Expression and secretion in yeast of a 400-kDa envelope glycoprotein derived from Epstein-Barr virus. *Gene*. 1987;54(1):113-23. doi: 10.1016/0378-1119(87)90353-2. cited by applicant

Schultz et al., Oligo-2'-fluoro-2'-deoxynucleotide N3'-->P5' phosphoramidates: synthesis and properties. *Nucleic Acids Res*. Aug. 1, 1996;24(15):2966-73. cited by applicant

Schwank et al., Functional repair of CFTR by CRISPR/Cas9 in intestinal stem cell organoids of cystic fibrosis patients. *Cell Stem Cell*. Dec. 5, 2013;13(6):653-8. doi:10.1016/j.stem.2013.11.002. cited by applicant

Schwartz et al., Post-translational enzyme activation in an animal via optimized conditional protein splicing. *Nat Chem Biol*. Jan. 2007;3(1):50-4. Epub Nov. 26, 2006. cited by applicant

Schwarze et al., In vivo protein transduction: delivery of a biologically active protein into the mouse. *Science*. Sep. 3, 1999;285(5433):1569-72. cited by applicant

Schöller et al., Interactions, localization, and phosphorylation of the m6A generating METTL3-METTL14-WTAP complex. *RNA*. Apr. 2018;24(4):499-512. doi: 10.1261/rna.064063.117. Epub Jan. 18, 2018. cited by applicant

Scimienti et al., Directed evolution of a recombinase for improved genomic integration at a native human sequence. *Nucleic Acids Res*. Dec. 15, 2001;29(24):5044-51. cited by applicant

SCORE Results for Luetticken et al., Complete genome sequence of a *Streptococcus dysgalactiae* subsp. RT equisimilis strain possessing Lancefield's group A antigen. RL Submitted to the EMBL/GenBank/DDBJ databases. May 2012. 3 pages. cited by applicant

SCORE Results for Okumura et al., Evolutionary paths of streptococcal and staphylococcal superantigens. RL *BMC Genomics*. 2012;13:404-404. 3 pages. cited by applicant

SCORE Results for Shimomura et al., Complete Genome Sequencing and Analysis of a Lancefield Group G RT *Streptococcus dysgalactiae* Subsp. Equisimilis Strain Causing Streptococcal RT Toxic Shock Syndrome (STSS). RL *BMC Genomics*. 2011;12:17-17. 3 pages. cited by applicant

Scott et al., Production of cyclic peptides and proteins in vivo. *Proc Natl Acad Sci U S A*. Nov. 23, 1999;96(24):13638-43. doi: 10.1073/pnas.96.24.13638. cited by applicant

Sebastián-Martín et al., Transcriptional inaccuracy threshold attenuates differences in RNA-dependent DNA synthesis fidelity between retroviral reverse transcriptases. *Sci Rep*. Jan. 12, 2018;8(1):627. doi: 10.1038/s41598-017-18974-8. cited by applicant

Seed, An LFA-3 cDNA encodes a phospholipid-linked membrane protein homologous to its receptor CD2. *Nature*. Oct. 29-Nov. 4, 1987;329(6142):840-2. doi: 10.1038/329840a0. cited by applicant

Sefton et al., Implantable pumps. *Crit Rev Biomed Eng*. 1987;14(3):201-40. cited by applicant

Segal et al., Toward controlling gene expression at will: selection and design of zinc finger domains recognizing each of the 5'-GNN-3' DNA target sequences. *Proc Natl Acad Sci U S A*. Mar. 16, 1999;96(6):2758-63. cited by applicant

Sells et al., Delivery of protein into cells using polycationic liposomes. *Biotechniques*. Jul. 1995;19(1):72-6, 78. cited by applicant

Semenova et al., Interference by clustered regularly spaced short palindromic repeat (CRISPR) RNA is governed by a seed sequence. *Proc Natl Acad Sci U S A*. Jun. 21, 2011;108(25):10098-103. doi: 10.1073/pnas.1104144108. Epub Jun. 6, 2011. cited by applicant

Semple et al., Rational design of cationic lipids for siRNA delivery. *Nat Biotechnol*. Feb. 2010;28(2):172-6. doi: 10.1038/nbt.1602. Epub Jan. 17, 2010. cited by applicant

Serganov et al., Coenzyme recognition and gene regulation by a flavin mononucleotide riboswitch. *Nature*. Mar. 12, 2009;458(7235):233-7. doi: 10.1038/nature07642. Epub Jan. 25, 2009. cited by applicant

Serganov et al., Structural basis for discriminative regulation of gene expression by adenine- and guanine-sensing mRNAs. *Chem Biol*. Dec. 2004;11(12):1729-41. cited by applicant

Serganov et al., Structural basis for gene regulation by a thiamine pyrophosphate-sensing riboswitch. *Nature*. Jun. 29, 2006;441(7097):1167-71. Epub May 21, 2006. cited by applicant

Seripa et al., The missing ApoE allele. *Ann Hum Genet*. Jul. 2007;71(Pt 4):496-500. Epub Jan. 22, 2007. cited by applicant

Serrano-Heras et al., Protein p56 from the *Bacillus subtilis* phage phi29 inhibits DNA-binding ability of uracil-DNA glycosylase. *Nucleic Acids Res*. 2007;35(16):5393-401. Epub Aug. 13, 2007. cited by applicant

Setten et al., The current state and future directions of RNAi-based therapeutics. *Nat Rev Drug Discov*. Jun. 2019;18(6):421-446. doi: 10.1038/s41573-019-0017-4. cited by applicant

Severinov et al., Expressed protein ligation, a novel method for studying protein-protein interactions in transcription. *J Biol Chem*. Jun. 26, 1998;273(26):16205-9. doi: 10.1074/jbc.273.26.16205. cited by applicant

Sha et al., Monobodies and other synthetic binding proteins for expanding protein science. *Protein Sci*. May 2017;26(5):910-924. doi: 10.1002/pro.3148. Epub Mar. 24, 2017. cited by applicant

Shah et al., Inteins: nature's gift to protein chemists. *Chem Sci*. 2014;5(1):446-461. cited by applicant

Shah et al., Kinetic control of one-pot trans-splicing reactions by using a wild-type and designed split intein. *Angew Chem Int Ed Engl*. Jul. 11, 2011;50(29):6511-5. doi: 10.1002/anie.201102909. Epub Jun. 8, 2011. cited by applicant

Shah et al., Protospacer recognition motifs: mixed identities and functional diversity. *RNA Biol*. May 2013;10(5):891-9. doi: 10.4161/rna.23764. Epub Feb. 12, 2013. cited by applicant

Shah et al., Target-specific variants of Flp recombinase mediate genome engineering reactions in mammalian cells. *FEBS J*. Sep. 2015;282(17):3323-33. doi: 10.1111/febs.13345. Epub Jul. 1, 2015. cited by applicant

Shaikh et al., Chimeras of the Flp and Cre recombinases: tests of the mode of cleavage by Flp and Cre. *J Mol Biol*. Sep. 8, 2000;302(1):27-48. cited by applicant

Shalem et al., Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science*. Jan. 3, 2014;343(6166):84-7. doi: 10.1126/science.1247005. Epub Dec. 12, 2013. cited by applicant

Shalem et al., High-throughput functional genomics using CRISPR-Cas9. *Nat Rev Genet*. May 2015;16(5):299-311. doi: 10.1038/nrg3899. Epub Apr. 9, 2015. cited by applicant

Sharbeen et al., Ectopic restriction of DNA repair reveals that UNG2 excises AID-induced uracils predominantly or exclusively during G1 phase. *J Exp Med*. May 7, 2012;209(5):965-74. doi: 10.1084/jem.20112379. Epub Apr. 23, 2012. cited by applicant

Sharer et al., The ARF-like 2 (ARL2)-binding protein, BART. Purification, cloning, and initial characterization. *J Biol Chem*. Sep. 24, 1999;274(39):27553-61. doi: 10.1074/jbc.274.39.27553. cited by applicant

Sharma et al., Efficient introduction of aryl bromide functionality into proteins in vivo. *FEBS Lett*. Feb. 4, 2000;467(1):37-40. cited by applicant

Sharma et al., Identification of novel methyltransferases, Bmt5 and Bmt6, responsible for the m3U methylations of 25S rRNA in *Saccharomyces cerevisiae*. *Nucleic Acids Res*. Mar. 2014;42(5):3246-60. doi: 10.1093/nar/gkt1281. Epub Dec. 11, 2013. cited by applicant

Sharon et al., Functional Genetic Variants Revealed by Massively Parallel Precise Genome Editing. *Cell*. Oct. 4, 2018;175(2):544-557.e16. doi: 10.1016/j.cell.2018.08.057. Epub Sep. 20, 2018. cited by applicant

Shaw et al., Implications of human genome architecture for rearrangement-based disorders: the genomic basis of disease. *Hum Mol Genet*. Apr. 1, 2004;13 Spec No. 1:R57-64. doi: 10.1093/hmg/ddh073. Epub Feb. 5, 2004. cited by applicant

Shcherbakova et al., Near-infrared fluorescent proteins for multicolor in vivo imaging. *Nat Methods*. Aug. 2013;10(8):751-4. doi: 10.1038/nmeth.2521. Epub Jun. 16, 2013. cited by applicant

Shechner et al., Multiplexable, locus-specific targeting of long RNAs with CRISPR-Display. *Nat Methods*. Jul. 2015;12(7):664-70. doi: 10.1038/nmeth.3433. Epub Jun. 1, 2015. cited by applicant

Shee et al., Engineered proteins detect spontaneous DNA breakage in human and bacterial cells. *Elife*. Oct. 29, 2013;2:e01222. doi: 10.7554/eLife.01222. cited by applicant

Shen et al., Herpes simplex virus 1 (HSV-1) for cancer treatment. *Cancer Gene Ther*. Nov. 2006;13(11):975-92. doi: 10.1038/sj.cgt.7700946. Epub Apr. 7, 2006. cited by applicant

Shen et al., Predictable and precise template-free CRISPR editing of pathogenic variants. *Nature*. Nov. 2018;563(7733):646-651. doi: 10.1038/s41586-018-0686-x. Epub Nov. 7, 2018. cited by applicant

Shen, Data processing, Modeling and Analysis scripts for CRISPR-inDelphi. GitHub—maxwshen/indelphi-dataprocessinganalysis at 6b68e3cec73c9358fef6e5f178a935f3c2a4118f. Apr. 10, 2018. Retrieved online via <https://github.com/maxwshen/indelphi-sataprocessinganalysis/tree/6b68e3cec73c9358fef6e5f178a935f3c2a4118f> Last

retrieved on Jul. 26, 2021. 2 pages. cited by applicant

Sheridan, First CRISPR-Cas patent opens race to stake out intellectual property. *Nat Biotechnol.* 2014;32(7):599-601. cited by applicant

Sheridan, Gene therapy finds its niche. *Nat Biotechnol.* Feb. 2011;29(2):121-8. doi: 10.1038/nbt.1769. cited by applicant

Sherwood et al., Discovery of directional and nondirectional pioneer transcription factors by modeling DNase profile magnitude and shape. *Nat Biotechnol.* Feb. 2014;32(2):171-178. doi: 10.1038/nbt.2798. Epub Jan. 19, 2014. cited by applicant

Shi et al., Structural basis for targeted DNA cytosine deamination and mutagenesis by APOBEC3A and APOBEC3B. *Nat Struct Mol Biol.* Feb. 2017;24(2):131-139. doi: 10.1038/nsmb.3344. Epub Dec. 19, 2016. cited by applicant

Shi et al., YTHDF3 facilitates translation and decay of N6-methyladenosine-modified RNA. *Cell Res.* Mar. 2017;27(3):315-328. doi: 10.1038/cr.2017.15. Epub Jan. 20, 2017. cited by applicant

Shimantani et al., Targeted base editing in rice and tomato using a CRISPR-Cas9 cytidine deaminase fusion. *Nat Biotechnol.* May 2017;35(5):441-443. doi: 10.1038/nbt.3833. Epub Mar. 27, 2017. cited by applicant

Shimojima et al., Spinocerebellar ataxias type 27 derived from a disruption of the fibroblast growth factor 14 gene with mimicking phenotype of paroxysmal non-kinesigenic dyskinesia. *Brain Dev.* Mar. 2012;34(3):230-3. doi: 10.1016/j.braindev.2011.04.014. Epub May 19, 2011. cited by applicant

Shin et al., CRISPR/Cas9 targeting events cause complex deletions and insertions at 17 sites in the mouse genome. *Nat Commun.* May 31, 2017;8:15464. doi: 10.1038/ncomms15464. cited by applicant

Shindo et al., A Comparison of Two Single-Stranded DNA Binding Models by Mutational Analysis of APOBEC3G. *Biology (Basel).* Aug. 2, 2012;1(2):260-76. doi: 10.3390/biology1020260. cited by applicant

Shingledecker et al., Molecular dissection of the Mycobacterium tuberculosis RecA intein: design of a minimal intein and of a trans-splicing system involving two intein fragments. *Gene.* Jan. 30, 1998;207(2):187-95. doi: 10.1016/s0378-1119(97)00624-0. cited by applicant

Shmakov et al., Discovery and Functional Characterization of Diverse Class 2 CRISPR Cas Systems. *Molecular Cell Nov.* 2015;60(3):385-97. cited by applicant

Shmakov et al., Diversity and evolution of class 2 CRISPR-Cas systems. *Nat Rev Microbiol.* Mar. 2017;15(3):169-182. doi: 10.1038/nrmicro.2016.184. Epub Jan. 23, 2017. cited by applicant

Shultz et al., A genome-wide analysis of FRT-like sequences in the human genome. *PLoS One.* Mar. 23, 2011;6(3):e18077. doi: 10.1371/journal.pone.0018077. cited by applicant

Siebert et al., An improved PCR method for walking in uncloned genomic DNA. *Nucleic Acids Res.* Mar. 25, 1995;23(6):1087-8. cited by applicant

Silas et al., Direct CRISPR spacer acquisition from RNA by a natural reverse transcriptase-Cas1 fusion protein. *Science.* Feb. 26, 2016;351(6276):aad4234. doi: 10.1126/science.aad4234. cited by applicant

Silva et al., Selective disruption of the DNA polymerase III  $\alpha$ - $\beta$  complex by the umuD gene products. *Nucleic Acids Res.* Jul. 2012;40(12):5511-22. doi: 10.1093/nar/gks229. Epub Mar. 9, 2012. cited by applicant

Simonelli et al., Base excision repair intermediates are mutagenic in mammalian cells. *Nucleic Acids Res.* Aug. 2, 2005;33(14):4404-11. Print 2005. cited by applicant

Singh et al., Cross-talk between diverse serine integrases. *J Mol Biol.* Jan. 23, 2014;426(2):318-31. doi: 10.1016/j.jmb.2013.10.013. Epub Oct. 22, 2013. cited by applicant

Singh et al., Real-time observation of DNA recognition and rejection by the RNA-guided endonuclease Cas9. *Nat Commun.* Sep. 14, 2016;7:12778. doi: 10.1038/ncomms12778. cited by applicant

Singh et al., Real-time observation of DNA target interrogation and product release by the RNA-guided endonuclease CRISPR Cpf1 (Cas12a). *Proc Natl Acad Sci U S A.* May 22, 2018;115(21):5444-5449. doi: 10.1073/pnas.1718686115. Epub May 7, 2018. cited by applicant

Sirk et al., Expanding the zinc-finger recombinase repertoire: directed evolution and mutational analysis of serine recombinase specificity determinants. *Nucleic Acids Res.* Apr. 2014;42(7):4755-66. doi: 10.1093/nar/gkt1389. Epub Jan. 21, 2014. cited by applicant

Siu et al., Riboregulated toehold-gated gRNA for programmable CRISPR-Cas9 function. *Nat Chem Biol.* Mar. 2019;15(3):217-220. doi: 10.1038/s41589-018-0186-1. Epub Dec. 10, 2018. cited by applicant

Sivalingam et al., Biosafety assessment of site-directed transgene integration in human umbilical cord-lining cells. *Mol Ther.* Jul. 2010;18(7):1346-56. doi: 10.1038/mt.2010.61. Epub Apr. 27, 2010. cited by applicant

Sjoblom et al., The consensus coding sequences of human breast and colorectal cancers. *Science.* Oct. 13, 2006;314(5797):268-74. Epub Sep. 7, 2006. cited by applicant

Skretas et al., Regulation of protein activity with small-molecule-controlled inteins. *Protein Sci.* Feb. 2005;14(2):523-32. Epub Jan. 4, 2005. cited by applicant

Slaymaker et al., Rationally engineered Cas9 nucleases with improved specificity. *Science.* Jan. 1, 2016;351(6268):84-8. doi: 10.1126/science.aad5227. Epub Dec. 1, 2015. cited by applicant

Sledz et al., Structural insights into the molecular mechanism of the m(6)A writer complex. *Elife.* Sep. 14, 2016;5:e18434. doi: 10.7554/eLife.18434. cited by applicant

Slupphaug et al., A nucleotide-flipping mechanism from the structure of human uracil-DNA glycosylase bound to DNA. *Nature.* Nov. 7, 1996;384(6604):87-92. doi: 10.1038/384087a0. cited by applicant

Smargon et al., Cas13b Is a Type VI-B CRISPR-Associated RNA-Guided RNase Differentially Regulated by Accessory Proteins Csx27 and Csx28. *Mol Cell*. Feb. 16, 2017;65(4):618-630.e7. doi: 10.1016/j.molcel.2016.12.023. Epub Jan. 5, 2017. cited by applicant

Smith et al., Diversity in the serine recombinases. *Mol Microbiol*. Apr. 2002;44(2):299-307. Review. cited by applicant

Smith et al., Expression of a dominant negative retinoic acid receptor  $\gamma$  in *Xenopus* embryos leads to partial resistance to retinoic acid. *Roux Arch Dev Biol*. Mar. 1994;203(5):254-265. doi: 10.1007/BF00360521. cited by applicant

Smith et al., Herpesvirus transport to the nervous system and back again. *Annu Rev Microbiol*. 2012;66:153-76. doi: 10.1146/annurev-micro-092611-150051. Epub Jun. 15, 2012. cited by applicant

Smith et al., Production of human beta interferon in insect cells infected with a baculovirus expression vector. *Mol Cell Biol*. Dec. 1983;3(12):2156-65. doi: 10.1128/mcb.3.12.2156. cited by applicant

Smith et al., Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione S-transferase. *Gene*. Jul. 15, 1988;67(1):31-40. doi: 10.1016/0378-1119(88)90005-4. cited by applicant

Smith, Filamentous fusion phage: novel expression vectors that display cloned antigens on the virion surface. *Science*. Jun. 14, 1985;228(4705):1315-7. cited by applicant

Smith, Phage-encoded Serine Integrases and Other Large Serine Recombinases. *Microbiol Spectr*. Aug. 2015;3(4). doi: 10.1128/microbiolspec.MDNA3-0059-2014. cited by applicant

Somanathan et al., AAV vectors expressing LDLR gain-of-function variants demonstrate increased efficacy in mouse models of familial hypercholesterolemia. *Circ Res*. Aug. 29, 2014;115(6):591-9. doi: 10.1161/CIRCRESAHA.115.304008. Epub Jul. 14, 2014. cited by applicant

Sommerfelt et al., Receptor interference groups of 20 retroviruses plating on human cells. *Virology*. May 1990;176(1):58-69. doi: 10.1016/0042-6822(90)90230-o. cited by applicant

Song et al., Adenine base editing in an adult mouse model of tyrosinaemia. *Nat Biomed Eng*. Jan. 2020;4(1):125-130. doi: 10.1038/s41551-019-0357-8. Epub Feb. 25, 2019. cited by applicant

Southworth et al., Control of protein splicing by intein fragment reassembly. *EMBO J*. Feb. 16, 1998;17(4):918-26. doi: 10.1093/emboj/17.4.918. cited by applicant

Southworth et al., Purification of proteins fused to either the amino or carboxy terminus of the *Mycobacterium xenopi* gyrase A intein. *Biotechniques*. Jul. 1999;27(1):110-4, 116, 118-20. doi: 10.2144/99271st04. cited by applicant

Spencer et al., A general strategy for producing conditional alleles of Src-like tyrosine kinases. *Proc Natl Acad Sci U S A*. Oct. 10, 1995;92(21):9805-9. doi: 10.1073/pnas.92.21.9805. cited by applicant

Spencer et al., Controlling signal transduction with synthetic ligands. *Science*. Nov. 12, 1993;262(5136):1019-24. doi: 10.1126/science.7694365. cited by applicant

Spencer et al., Functional analysis of Fas signaling in vivo using synthetic inducers of dimerization. *Curr Biol*. Jul. 1, 1996;6(7):839-47. doi: 10.1016/s0960-9822(02)00607-3. cited by applicant

Srivastava et al., An inhibitor of nonhomologous end-joining abrogates double-strand break repair and impedes cancer progression. *Cell*. Dec. 21, 2012;151(7):1474-87. doi: 10.1016/j.cell.2012.11.054. cited by applicant

Stadtman, Selenocysteine. *Annu Rev Biochem*. 1996;65:83-100. cited by applicant

Stamos et al., Structure of a Thermostable Group II Intron Reverse Transcriptase with Template-Primer and Its Functional and Evolutionary Implications. *Mol Cell*. Dec. 7, 2017;68(5):926-939.e4. doi: 10.1016/j.molcel.2017.10.024. Epub Nov. 16, 2017. cited by applicant

Steele et al., The prion protein knockout mouse: a phenotype under challenge. *Prion*. Apr.-Jun. 2007;1(2):83-93. doi: 10.4161/pri.1.2.4346. Epub Apr. 25, 2007. cited by applicant

Steiner et al., The neurotropic herpes viruses: herpes simplex and varicella-zoster. *Lancet Neurol*. Nov. 2007;6(11):1015-28. doi: 10.1016/S1474-4422(07)70267-3. cited by applicant

Stella et al., Structure of the Cpf1 endonuclease R-loop complex after target DNA cleavage. *Nature*. Jun. 22, 2017;546(7659):559-563. doi: 10.1038/nature22398. Epub May 31, 2017. cited by applicant

Stenglein et al., APOBEC3 proteins mediate the clearance of foreign DNA from human cells. *Nat Struct Mol Biol*. Feb. 2010;17(2):222-9. doi: 10.1038/nsmb.1744. Epub Jan. 10, 2010. cited by applicant

Stenson et al., The Human Gene Mutation Database: towards a comprehensive repository of inherited mutation data for medical research, genetic diagnosis and next-generation sequencing studies. *Hum Genet*. Jun. 2017;136(6):665-677. doi: 10.1007/s00439-017-1779-6. Epub Mar. 27, 2017. cited by applicant

Stephens et al., The landscape of cancer genes and mutational processes in breast cancer. *Nature*. Jun. 2012;486:400-404. doi: 10.1038/nature11017. cited by applicant

Sternberg et al., Conformational control of DNA target cleavage by CRISPR-Cas9. *Nature*. Nov. 5, 2015;527(7576):110-3. doi: 10.1038/nature15544. Epub Oct. 28, 2015. cited by applicant

Sternberg et al., DNA interrogation by the CRISPR RNA-guided endonuclease Cas9. *Nature*. Mar. 6, 2014;507(7490):62-7. doi: 10.1038/nature13011. Epub Jan. 29, 2014. cited by applicant

Sterne-Weiler et al., Exon identity crisis: disease-causing mutations that disrupt the splicing code. *Genome Biol*. Jan. 23, 2014;15(1):201. doi: 10.1186/gb4150. cited by applicant

Stevens et al., A promiscuous split intein with expanded protein engineering applications. *Proc Natl Acad Sci U S A*. Aug. 8, 2017;114(32):8538-8543. doi: 10.1073/pnas.1701083114. Epub Jul. 24, 2017. cited by applicant

Stockwell et al., Probing the role of homomeric and heteromeric receptor interactions in TGF-beta signaling using small

molecular dimerizers. *Curr Biol.* Jun. 18, 1998;8(13):761-70. doi: 10.1016/s0960-9822(98)70299-4. cited by applicant

Strecker et al., Engineering of CRISPR-Cas12b for human genome editing. *Nat Commun.* Jan. 22, 2019;10(1):212. doi: 10.1038/s41467-018-08224-4. cited by applicant

Strecker et al., RNA-guided DNA insertion with CRISPR-associated transposases. *Science.* Jul. 5, 2019;365(6448):48-53. doi: 10.1126/science.aax9181. Epub Jun. 6, 2019. cited by applicant

Strutt et al., RNA-dependent RNA targeting by CRISPR-Cas9. *Elife.* Jan. 5, 2018;7:e32724. doi: 10.7554/eLife.32724. cited by applicant

Su et al., Human DNA polymerase  $\gamma$  has reverse transcriptase activity in cellular environments. *J Biol Chem.* Apr. 12, 2019;294(15):6073-6081. doi: 10.1074/jbc.RA119.007925. Epub Mar. 6, 2019. cited by applicant

Sudarsan et al., An mRNA structure in bacteria that controls gene expression by binding lysine. *Genes Dev.* Nov. 1, 2003;17(21):2688-97. cited by applicant

Sudarsan et al., Riboswitches in eubacteria sense the second messenger cyclic di-GMP. *Science.* Jul. 18, 2008;321(5887):411-3. doi: 10.1126/science.1159519. cited by applicant

Suess et al., A theophylline responsive riboswitch based on helix slipping controls gene expression in vivo. *Nucleic Acids Res.* Mar. 5, 2004;32(4):1610-4. cited by applicant

Sullenger et al., Ribozyme-mediated repair of defective mRNA by targeted, trans-splicing. *Nature.* Oct. 13, 1994;371(6498):619-22. doi: 10.1038/371619a0. cited by applicant

Sun et al., Optimized TAL effector nucleases (TALENs) for use in treatment of sickle cell disease. *Mol Biosyst.* Apr. 2012;8(4):1255-63. doi: 10.1039/c2mb05461b. Epub Feb. 3, 2012. cited by applicant

Sun et al., The CRISPR/Cas9 system for gene editing and its potential application in pain research. *Transl Periop & Pain Med.* Aug. 3, 2016;1(3):22-33. cited by applicant

Surun et al., High Efficiency Gene Correction in Hematopoietic Cells by Donor-Template-Free CRISPR/Cas9 Genome Editing. *Mol Ther Nucleic Acids.* Mar. 2, 2018;10:1-8. doi: 10.1016/j.omtn.2017.11.001. Epub Nov. 10, 2017. cited by applicant

Suzuki et al., Crystal structures reveal an elusive functional domain of pyrrolysyl-tRNA synthetase. *Nat Chem Biol.* Dec. 2017;13(12):1261-1266. doi: 10.1038/nchembio.2497. Epub Oct. 16, 2017. cited by applicant

Suzuki et al., In vivo genome editing via CRISPR/Cas9 mediated homology-independent targeted integration. *Nature.* Dec. 1, 2016;540(7631):144-149. doi: 10.1038/nature20565. Epub Nov. 16, 2016. cited by applicant

Suzuki et al., VCre/VloxP and SCre/SloxP: new site-specific recombination systems for genome engineering. *Nucleic Acids Res.* Apr. 2011;39(8):e49. doi: 10.1093/nar/gkq1280. Epub Feb. 1, 2011. cited by applicant

Swarts et al., Argonaute of the archaeon *Pyrococcus furiosus* is a DNA-guided nuclease that targets cognate DNA. *Nucleic Acids Res.* May 26, 2015;43(10):5120-9. doi: 10.1093/nar/gkv415. Epub Apr. 29, 2015. cited by applicant

Swarts et al., DNA-guided DNA interference by a prokaryotic Argonaute. *Nature.* Mar. 13, 2014;507(7491):258-61. doi: 10.1038/nature12971. Epub Feb. 16, 2014. cited by applicant

Swarts et al., The evolutionary journey of Argonaute proteins. *Nat Struct Mol Biol.* Sep. 2014;21(9):743-53. doi: 10.1038/nsmb.2879. cited by applicant

Szczeppek et al., Structure-based redesign of the dimerization interface reduces the toxicity of zinc-finger nucleases. *Nat Biotechnol.* Jul. 2007;25(7):786-93. Epub Jul. 1, 2007. cited by applicant

Tabebordbar et al., In vivo gene editing in dystrophic mouse muscle and muscle stem cells. *Science.* Jan. 22, 2016;351(6271):407-411. doi: 10.1126/science.aad5177. Epub Dec. 31, 2015. cited by applicant

Tagalakakis et al., Lack of RNA-DNA oligonucleotide (chimeraplast) mutagenic activity in mouse embryos. *Mol Reprod Dev.* Jun. 2005;71(2):140-4. cited by applicant

Tahara et al., Potent and Selective Inhibitors of 8-Oxoguanine DNA Glycosylase. *J Am Chem Soc.* Feb. 14, 2018;140(6):2105-2114. doi: 10.1021/jacs.7b09316. Epub Feb. 5, 2018. cited by applicant

Tajiri et al., Functional cooperation of MutT, MutM and MutY proteins in preventing mutations caused by spontaneous oxidation of guanine nucleotide in *Escherichia coli*. *Mutat Res.* May 1995;336(3):257-67. doi: 10.1016/0921-8777(94)00062-b. cited by applicant

Takimoto et al., Stereochemical basis for engineered pyrrolysyl-tRNA synthetase and the efficient in vivo incorporation of structurally divergent non-native amino acids. *ACS Chem Biol.* Jul. 15, 2011;6(7):733-43. doi: 10.1021/cb200057a. Epub May 5, 2011. cited by applicant

Tambunan et al., Vaccine Design for H5N1 Based on B- and T-cell Epitope Predictions. *Bioinform Biol Insights.* Apr. 28, 2016;10:27-35. doi: 10.4137/BBI.S38378. cited by applicant

Tanenbaum et al., A protein-tagging system for signal amplification in gene expression and fluorescence imaging. *Cell.* Oct. 23, 2014;159(3):635-46. doi: 10.1016/j.cell.2014.09.039. Epub Oct. 9, 2014. cited by applicant

Tanese et al., Expression of enzymatically active reverse transcriptase in *Escherichia coli*. *Proc Natl Acad Sci U S A.* Aug. 1985;82(15):4944-8. doi: 10.1073/pnas.82.15.4944. cited by applicant

Tang et al., Aptazyme-embedded guide RNAs enable ligand-responsive genome editing and transcriptional activation. *Nat Commun.* Jun. 28, 2017;8:15939. doi: 10.1038/ncomms15939. cited by applicant

Tang et al., Evaluation of Bioinformatic Programmes for the Analysis of Variants within Splice Site Consensus Regions. *Adv Bioinformatics.* 2016;2016:5614058. doi: 10.1155/2016/5614058. Epub May 24, 2016. cited by applicant

Tang et al., Rewritable multi-event analog recording in bacterial and mammalian cells. *Science.* Apr. 13,



2018;360(eaap8992):eaap8992. doi: 10.1126/science.aap8992. Epub Feb. 15, 2018. cited by applicant

Tassabehji, Williams-Beuren syndrome: a challenge for genotype-phenotype correlations. *Hum Mol Genet.* Oct. 15, 2003;12 Spec No. 2:R229-37. doi: 10.1093/hmg/ddg299. Epub Sep. 2, 2003. cited by applicant

Taube et al., Reverse transcriptase of mouse mammary tumour virus: expression in bacteria, purification and biochemical characterization. *Biochem J.* Feb. 1, 1998;329 ( Pt 3)(Pt 3):579-87. doi: 10.1042/bj3290579. Erratum in: *Biochem J* Jun. 15, 1998;332(Pt 3):808. cited by applicant

Tebas et al., Gene editing of CCR5 in autologous CD4 T cells of persons infected with HIV. *Engl J Med.* Mar. 6, 2014;370(10):901-10. doi: 10.1056/NEJMoa1300662. cited by applicant

Tee et al., Polishing the craft of genetic diversity creation in directed evolution. *Biotechnol Adv.* Dec. 2013;31(8):1707-21. doi: 10.1016/j.biotechadv.2013.08.021. Epub Sep. 6, 2013. cited by applicant

Telenti et al., The *Mycobacterium xenopi* GyrA protein splicing element: characterization of a minimal intein. *J Bacteriol.* Oct. 1997;179(20):6378-82. doi: 10.1128/jb.179.20.6378-6382.1997. cited by applicant

Telesnitsky et al., RNase H domain mutations affect the interaction between Moloney murine leukemia virus reverse transcriptase and its primer-template. *Proc Natl Acad Sci U S A.* Feb. 15, 1993;90(4):1276-80. doi: 10.1073/pnas.90.4.1276. cited by applicant

Teng et al., Mutational analysis of apolipoprotein B mRNA editing enzyme (APOBEC1). structure-function relationships of RNA editing and dimerization. *J Lipid Res.* Apr. 1999;40(4):623-35. cited by applicant

Tessarollo et al., Targeted mutation in the neurotrophin-3 gene results in loss of muscle sensory neurons. *Proc Natl Acad Sci U S A.* Dec. 6, 1994;91(25):11844-8. cited by applicant

Tesson et al., Knockout rats generated by embryo microinjection of TALENs. *Nat Biotechnol.* Aug. 5, 2011;29(8):695-6. doi: 10.1038/nbt.1940. cited by applicant

Thompson et al., Cellular uptake mechanisms and endosomal trafficking of supercharged proteins. *Chem Biol.* Jul. 27, 2012;19(7):831-43. doi: 10.1016/j.chembiol.2012.06.014. cited by applicant

Thompson et al., Engineering and identifying supercharged proteins for macromolecule delivery into mammalian cells. *Methods Enzymol.* 2012;503:293-319. doi: 10.1016/B978-0-12396962-0.00012-4. cited by applicant

Thompson et al., The Future of Multiplexed Eukaryotic Genome Engineering. *ACS Chem Biol.* Feb. 16, 2018;13(2):313-325. doi: 10.1021/acscchembio.7b00842. Epub Dec. 28, 2017. cited by applicant

Thomson et al., Mutational analysis of loxP sites for efficient Cre-mediated insertion into genomic DNA. *Genesis.* Jul. 2003;36(3):162-7. doi: 10.1002/gene.10211. cited by applicant

Thorpe et al., Functional correction of episomal mutations with short DNA fragments and RNA-DNA oligonucleotides. *J Gene Med.* Mar.-Apr. 2002;4(2):195-204. cited by applicant

Thuronyi et al., Continuous evolution of base editors with expanded target compatibility and improved activity. *Nat Biotechnol.* Sep. 2019;37(9):1070-1079. doi: 10.1038/s41587-019-0193-0. Epub Jul. 22, 2019. cited by applicant

Thyagarajan et al., Creation of engineered human embryonic stem cell lines using phiC31 integrase. *Stem Cells.* Jan. 2008;26(1):119-26. doi: 10.1634/stemcells.2007-0283. Epub Oct. 25, 2007. cited by applicant

Thyagarajan et al., Mammalian genomes contain active recombinase recognition sites. *Gene.* Feb. 22, 2000;244(1-2):47-54. cited by applicant

Thyagarajan et al., Site-specific genomic integration in mammalian cells mediated by phage phiC31 integrase. *Mol Cell Biol.* Jun. 2001;21(12):3926-34. cited by applicant

Tinland et al., The T-DNA-linked VirD2 protein contains two distinct functional nuclear localization signals. *Proc Natl Acad Sci U S A.* Aug. 15, 1992;89(16):7442-6. doi: 10.1073/pnas.89.16.7442. cited by applicant

Tirumalai et al., Recognition of core-type DNA sites by lambda integrase. *J Mol Biol.* Jun. 12, 1998;279(3):513-27. cited by applicant

Tom et al., Mechanism whereby proliferating cell nuclear antigen stimulates flap endonuclease 1. *J Biol Chem.* Apr. 7, 2000;275(14):10498-505. doi: 10.1074/jbc.275.14.10498. cited by applicant

Tone et al., Single-stranded DNA binding protein Gp5 of Bacillus subtilis phage ?29 is required for viral DNA replication in growth-temperature dependent fashion. *Biosci Biotechnol Biochem.* 2012;76(12):2351-3. doi: 10.1271/bbb.120587. Epub Dec. 7, 2012. cited by applicant

Toor et al., Crystal structure of a self-spliced group II intron. *Science.* Apr. 4, 2008;320(5872):77-82. doi: 10.1126/science.1153803. cited by applicant

Toro et al., On the Origin and Evolutionary Relationships of the Reverse Transcriptases Associated With Type III CRISPR-Cas Systems. *Front Microbiol.* Jun. 15, 2018;9:1317. doi: 10.3389/fmicb.2018.01317. cited by applicant

Toro et al., The Reverse Transcriptases Associated with CRISPR-Cas Systems. *Sci Rep.* Aug. 2, 2017;7(1):7089. doi: 10.1038/s41598-017-07828-y. cited by applicant

Torres et al., Non-integrative lentivirus drives high-frequency cre-mediated cassette exchange in human cells. *PLoS One.* 2011;6(5):e19794. doi: 10.1371/journal.pone.0019794. Epub May 23, 2011. cited by applicant

Tourdout et al., A general strategy to enhance immunogenicity of low-affinity HLA-A2. 1-associated peptides: implication in the identification of cryptic tumor epitopes. *Eur J Immunol.* Dec. 2000;30(12):3411-21. cited by applicant

Townsend et al., Role of HFE in iron metabolism, hereditary haemochromatosis, anaemia of chronic disease, and secondary iron overload. *Lancet.* Mar. 2, 2002;359(9308):786-90. doi: 10.1016/S0140-6736(02)07885-6. cited by applicant

Tracewell et al., Directed evolution: climbing fitness peaks one amino acid at a time. *Curr Opin Chem Biol.* Feb. 2009;13(1):3-9. doi: 10.1016/j.cbpa.2009.01.017. Epub Feb. 25, 2009. cited by applicant

Tratschin et al., A human parvovirus, adeno-associated virus, as a eucaryotic vector: transient expression and encapsidation of the procaryotic gene for chloramphenicol acetyltransferase. *Mol Cell Biol.* Oct. 1984;4(10):2072-81. doi: 10.1128/mcb.4.10.2072. cited by applicant

Tratschin et al., Adeno-associated virus vector for high-frequency integration, expression, and rescue of genes in mammalian cells. *Mol Cell Biol.* Nov. 1985;5(11):3251-60. doi: 10.1128/mcb.5.11.3251. cited by applicant

Trausch et al., The structure of a tetrahydrofolate-sensing riboswitch reveals two ligand binding sites in a single aptamer. *Structure.* Oct. 12, 2011;19(10):1413-23. doi: 10.1016/j.str.2011.06.019. Epub Sep. 8, 2011. cited by applicant

Traxler et al., A genome-editing strategy to treat  $\beta$ -hemoglobinopathies that recapitulates a mutation associated with a benign genetic condition. *Nat Med.* Sep. 2016;22(9):987-90. doi: 10.1038/nm.4170. Epub Aug. 15, 2016. cited by applicant

Trojan et al., Functional analysis of hMLH1 variants and HNPCC-related mutations using a human expression system. *Gastroenterology.* Jan. 2002;122(1):211-9. doi: 10.1053/gast.2002.30296. cited by applicant

Trudeau et al., On the Potential Origins of the High Stability of Reconstructed Ancestral Proteins. *Mol Biol Evol.* Oct. 2016;33(10):2633-41. doi: 10.1093/molbev/msw138. Epub Jul. 12, 2016. cited by applicant

Truong et al., Development of an intein-mediated split-Cas9 system for gene therapy. *Nucleic Acids Res.* Jul. 27, 2015;43(13):6450-8. doi: 10.1093/nar/gkv601. Epub Jun. 16, 2015. With Supplementary Data. cited by applicant

Tsai et al., CIRCLE-seq: a highly sensitive in vitro screen for genome-wide CRISPR-Cas9 nuclease off-targets. *Nat Methods.* Jun. 2017;14(6):607-614. doi: 10.1038/nmeth.4278. Epub May 1, 2017. cited by applicant

Tsai et al., Dimeric CRISPR RNA-guided FokI nucleases for highly specific genome editing. *Nat Biotechnol.* Jun. 2014;32(6):569-76. doi: 10.1038/nbt.2908. Epub Apr. 25, 2014. cited by applicant

Tsai et al., GUIDE-seq enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases. *Nat Biotechnol.* Feb. 2015;33(2):187-97. doi: 10.1038/nbt.3117. Epub Dec. 16, 2014. cited by applicant

Tsang et al., Specialization of the DNA-cleaving activity of a group I ribozyme through in vitro evolution. *J Mol Biol.* Sep. 13, 1996;262(1):31-42. doi: 10.1006/jmbi.1996.0496. cited by applicant

Tsutakawa et al., Human flap endonuclease structures, DNA double-base flipping, and a unified understanding of the FEN1 superfamily. *Cell.* Apr. 15, 2011;145(2):198-211. doi: 10.1016/j.cell.2011.03.004. cited by applicant

Turan et al., Recombinase-mediated cassette exchange (RMCE)—a rapidly-expanding toolbox for targeted genomic modifications. *Gene.* Feb. 15, 2013;515(1):1-27. doi: 10.1016/j.gene.2012.11.016. Epub Nov. 29, 2012. cited by applicant

Turan et al., Recombinase-mediated cassette exchange (RMCE): traditional concepts and current challenges. *J Mol Biol.* Mar. 25, 2011;407(2):193-221. doi: 10.1016/j.jmb.2011.01.004. Epub Jan. 15, 2011. cited by applicant

Turan et al., Site-specific recombinases: from tag-and-target- to tag-and-exchange-based genomic modifications. *Faseb J.* Dec. 2011;25(12):4088-107. doi: 10.1096/fj.11-186940. Epub Sep. 2, 2011. Review. cited by applicant

Tycko et al., Pairwise library screen systematically interrogates *Staphylococcus aureus* Cas9 specificity in human cells. *bioRxiv.* doi: <https://doi.org/10.1101/269399> Posted Feb. 22, 2018. cited by applicant

UniProt Consortium, UniProt: the universal protein knowledgebase. *Nucleic Acids Res.* Mar. 16, 2018;46(5):2699. doi: 10.1093/nar/gky092. cited by applicant

UniProt Submission; UniProt, Accession No. P01011. Last modified Jun. 11, 2014, version 2. 15 pages. cited by applicant

UniProt Submission; UniProt, Accession No. P01011. Last modified Sep. 18, 2013, version 2. 15 pages. cited by applicant

UniProt Submission; UniProt, Accession No. P04264. Last modified Jun. 11, 2014, version 6. 15 pages. cited by applicant

UniProt Submission; UniProt, Accession No. P04275. Last modified Jul. 9, 2014, version 107. 29 pages. cited by applicant

UniProtKB A0A1V6. Dec. 11, 2019. cited by applicant

UniProtKB Submission; Accession No. F0NH53. May 3, 2011. 4 pages. cited by applicant

UniProtKB Submission; Accession No. F0NN87. May 3, 2011. 4 pages. cited by applicant

UniProtKB Submission; Accession No. P0DOC6. No Author Listed., Oct. 5, 2016. 5 pages. cited by applicant

UniProtKB Submission; Accession No. T0D7A2. Oct. 16, 2013. 10 pages. cited by applicant

Urasaki et al., Functional dissection of the Tol2 transposable element identified the minimal cis-sequence and a highly repetitive sequence in the subterminal region essential for transposition. *Genetics.* Oct. 2006;174(2):639-49. doi: 10.1534/genetics.106.060244. Epub Sep. 7, 2006. cited by applicant

Urnov et al., Genome editing with engineered zinc finger nucleases. *Nat Rev Genet.* Sep. 2010;11(9):636-46. doi: 10.1038/nrg2842. cited by applicant

Urnov et al., Highly efficient endogenous human gene correction using designed zinc-finger nucleases. *Nature.* Jun. 2, 2005;435(7042):646-51. Epub Apr. 3, 2005. cited by applicant

Usman et al., Exploiting the chemical synthesis of RNA. *Trends Biochem Sci.* Sep. 1992;17(9):334-9. doi: 10.1016/0968-0004(92)90306-t. cited by applicant

Vagner et al., Efficiency of homologous DNA recombination varies along the *Bacillus subtilis* chromosome. *J Bacteriol.* Sep. 1988;170(9):3978-82. cited by applicant

Van Brunt et al., Genetically Encoded Azide Containing Amino Acid in Mammalian Cells Enables Site-Specific Antibody-Drug Conjugates Using Click Cycloaddition Chemistry. *Bioconjug Chem.* Nov. 18, 2015;26(11):2249-60. doi:

10.1021/acs.biochem.5b00359. Epub Sep. 11, 2015. cited by applicant

Van Brunt et al., Molecular Farming: Transgenic Animals as Bioreactors. *Biotechnology (Y)*. 1988;6(10):1149-1154. doi: 10.1038/nbt1088-1149. cited by applicant

Van Duyne et al., Teaching Cre to follow directions. *Proc Natl Acad Sci U S A*. Jan. 6, 2009;106(1):4-5. doi: 10.1073/pnas.0811624106. Epub Dec. 31, 2008. cited by applicant

Van Overbeek et al., DNA Repair Profiling Reveals Nonrandom Outcomes at Cas9-Mediated Breaks. *Mol Cell*. Aug. 18, 2016;63(4):633-646. doi: 10.1016/j.molcel.2016.06.037. Epub Aug. 4, 2016. cited by applicant

Van Swieten et al., A mutation in the fibroblast growth factor 14 gene is associated with autosomal dominant cerebellar ataxia [corrected]. *Am J Hum Genet*. Jan. 2003;72(1):191-9. Epub Dec. 13, 2002. cited by applicant

Van Wijk et al., Identification of 51 novel exons of the Usher syndrome type 2A (USH2A) gene that encode multiple conserved functional domains and that are mutated in patients with Usher syndrome type II. *Am J Hum Genet*. Apr. 2004;74(4):738-44. doi: 10.1086/383096. Epub Mar. 10, 2004. cited by applicant

Vanamee et al., FokI requires two specific DNA sites for cleavage. *J Mol Biol*. May 25, 2001;309(1):69-78. cited by applicant

Varga et al., Progressive vascular smooth muscle cell defects in a mouse model of Hutchinson-Gilford progeria syndrome. *Proc Natl Acad Sci U S A*. Feb. 28, 2006;103(9):3250-5. doi: 10.1073/pnas.0600012103. Epub Feb. 21, 2006. cited by applicant

Vellore et al., A group II intron-type open reading frame from the thermophile *Bacillus (Geobacillus) stearothermophilus* encodes a heat-stable reverse transcriptase. *Appl Environ Microbiol*. Dec. 2004;70(12):7140-7. doi: 10.1128/AEM.70.12.7140-7147.2004. cited by applicant

Venken et al., Genome-wide manipulations of *Drosophila melanogaster* with transposons, FLP recombinase, and ΦC31 integrase. *Methods Mol Biol*. 2012;859:203-28. doi: 10.1007/978-1-61779-603-6\_12. cited by applicant

Verma, The reverse transcriptase. *Biochim Biophys Acta*. Mar. 21, 1977;473(1):1-38. doi: 10.1016/0304-419x(77)90005-1. cited by applicant

Vigne et al., Third-generation adenovectors for gene therapy. *Restor Neurol Neurosci*. Jan. 1, 1995;8(1):35-6. doi: 10.3233/RNN-1995-81208. cited by applicant

Vik et al., Endonuclease V cleaves at inosines in RNA. *Nat Commun*. 2013;4:2271. doi: 10.1038/ncomms3271. cited by applicant

Vilenchik et al., Endogenous DNA double-strand breaks: production, fidelity of repair, and induction of cancer. *Proc Natl Acad Sci U S A*. Oct. 28, 2003;100(22):12871-6. doi: 10.1073/pnas.2135498100. Epub Oct. 17, 2003. cited by applicant

Villiger et al., Treatment of a metabolic liver disease by in vivo genome base editing in adult mice. *Nat Med*. Oct. 2018;24(10):1519-1525. doi: 10.1038/s41591-018-0209-1. Epub Oct. 8, 2018. cited by applicant

Vitreschak et al., Regulation of the vitamin B12 metabolism and transport in bacteria by a conserved RNA structural element. *RNA*. Sep. 2003;9(9):1084-97. cited by applicant

Voigt et al., Rational evolutionary design: the theory of in vitro protein evolution. *Adv Protein Chem*. 2000;55:79-160. cited by applicant

Vriend et al., Nick-initiated homologous recombination: Protecting the genome, one strand at a time. *DNA Repair (Amst)*. Feb. 2017;50:1-13. doi: 10.1016/j.dnarep.2016.12.005. Epub Dec. 29, 2016. cited by applicant

Wacey et al., Disentangling the perturbational effects of amino acid substitutions in the DNA-binding domain of p53. *Hum Genet*. Jan. 1999;104(1):15-22. cited by applicant

Wadia et al., Modulation of cellular function by TAT mediated transduction of full length proteins. *Curr Protein Pept Sci*. Apr. 2003;4(2):97-104. cited by applicant

Wadia et al., Transducible TAT-HA fusogenic peptide enhances escape of TAT-fusion proteins after lipid raft macropinocytosis. *Nat Med*. Mar. 2004;10(3):310-5. Epub Feb. 8, 2004. cited by applicant

Wah et al., Structure of FokI has implications for DNA cleavage. *Proc Natl Acad Sci U S A*. Sep. 1, 1998;95(18):10564-9. cited by applicant

Wals et al., Unnatural amino acid incorporation in *E. coli*: current and future applications in the design of therapeutic proteins. *Front Chem*. Apr. 1, 2014;2:15. doi: 10.3389/fchem.2014.00015. eCollection 2014. cited by applicant

Wan et al., Material solutions for delivery of CRISPR/Cas-based genome editing tools: Current status and future outlook. *Materials Today*. Jun. 2019;26:40-66. doi: 10.1016/j.mattod.2018.12.003. cited by applicant

Wang et al. CRISPR-Cas9 and CRISPR-Assisted Cytidine Deaminase Enable Precise and Efficient Genome Editing in *Klebsiella pneumoniae*. *Appl Environ Microbiol*. 2018;84(23):e01834-18. Published Nov. 15, 2018. doi:10.1128/AEM.01834-18. cited by applicant

Wang et al., AID upmutants isolated using a high-throughput screen highlight the immunity/cancer balance limiting DNA deaminase activity. *Nat Struct Mol Biol*. Jul. 2009;16(7):769-76. doi: 10.1038/nsmb.1623. Epub Jun. 21, 2009. cited by applicant

Wang et al., Continuous directed evolutions of proteins with improved soluble expression. *Nature Chemical Biology*. Nat Publishing Group. Aug. 20, 2018; 14(10):972-980. cited by applicant

Wang et al., CRISPR-Cas9 Targeting of PCSK9 in Human Hepatocytes In Vivo-Brief Report. *Arterioscler Thromb Vasc Biol*. May 2016;36(5):783-6. doi: 10.1161/ATVBAHA.116.307227. Epub Mar. 3, 2016. cited by applicant

Wang et al., Efficient delivery of genome-editing proteins using bio-reducible lipid nanoparticles. *Proc Natl Acad Sci U S A*

A. Feb. 2016; pii: 201520244. [Epub ahead of print]. cited by applicant

Wang et al., Enhanced base editing by co-expression of free uracil DNA glycosylase inhibitor. *Cell Res.* Oct. 2017;27(1):1289-92. doi: 10.1038/cr.2017.111. Epub Aug. 29, 2017. cited by applicant

Wang et al., Evolution of new nonantibody proteins via iterative somatic hypermutation. *Proc Natl Acad Sci U S A.* Nov. 30, 2004;101(48):16745-9. Epub Nov. 19, 2004. cited by applicant

Wang et al., Expanding the genetic code. *Annu Rev Biophys Biomol Struct.* 2006;35:225-49. Review. cited by applicant

Wang et al., Genetic screens in human cells using the CRISPR-Cas9 system. *Science.* Jan. 3, 2014;343(6166):80-4. doi: 10.1126/science.1246981. Epub Dec. 12, 2013. cited by applicant

Wang et al., Highly efficient CRISPR/HDR-mediated knock-in for mouse embryonic stem cells and zygotes. *Biotechniques.* 2015;59,201-2;204;206-8. cited by applicant

Wang et al., (6)-methyladenosine Modulates Messenger RNA Translation Efficiency. *Cell.* Jun. 4, 2015;161(6):1388-99. doi: 10.1016/j.cell.2015.05.014. cited by applicant

Wang et al., N6-methyladenosine-dependent regulation of messenger RNA stability. *Nature.* Jan. 2, 2014;505(7481):117-20. doi: 10.1038/nature12730. Epub Nov. 27, 2013. cited by applicant

Wang et al., Nucleation, propagation and cleavage of target RNAs in Ago silencing complexes. *Nature.* Oct. 8, 2009;461(7265):754-61. doi: 10.1038/nature08434. cited by applicant

Wang et al., One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. *Cell.* May 9, 2013;153(4):910-8. doi: 10.1016/j.cell.2013.04.025. Epub May 2, 2013. cited by applicant

Wang et al., Optimized paired-sgRNA/Cas9 cloning and expression cassette triggers high-efficiency multiplex genome editing in kiwifruit. *Plant Biotechnol J.* Aug. 2018;16(8):1424-1433. doi: 10.1111/pbi.12884. Epub Feb. 6, 2018. cited by applicant

Wang et al., Programming cells by multiplex genome engineering and accelerated evolution. *Nature.* Aug. 13, 2009;460(7257):894-8. Epub Jul. 26, 2009. cited by applicant

Wang et al., Reading RNA methylation codes through methyl-specific binding proteins. *RNA Biol.* 2014;11(6):669-72. doi: 10.4161/rna.28829. Epub Apr. 24, 2014. cited by applicant

Wang et al., Recombinase technology: applications and possibilities. *Plant Cell Rep.* Mar. 2011;30(3):267-85. doi: 10.1007/s00299-010-0938-1. Epub Oct. 24, 2010. cited by applicant

Wang et al., Riboswitches that sense S-adenosylhomocysteine and activate genes involved in coenzyme recycling. *Mol Cell.* Mar. 28, 2008;29(6):691-702. doi: 10.1016/j.molcel.2008.01.012. cited by applicant

Wang et al., *Staphylococcus aureus* protein SAUGI acts as a uracil-DNA glycosylase inhibitor. *Nucleic Acids Res.* Jan. 2014;42(2):1354-64. doi: 10.1093/nar/gkt964. Epub Oct. 22, 2013. cited by applicant

Wang et al., Structural basis of (6)-adenosine methylation by the METTL3-METTL14 complex. *Nature.* Jun. 23, 2016;534(7608):575-8. doi: 10.1038/nature18298. Epub May 25, 2016. cited by applicant

Wang et al., Targeted gene addition to a predetermined site in the human genome using a ZFN-based nicking enzyme. *Genome Res.* Jul. 2012;22(7):1316-26. doi: 10.1101/gr.122879.111. Epub Mar. 20, 2012. cited by applicant

Wang et al., Uracil-DNA glycosylase inhibitor gene of bacteriophage PBS2 encodes a binding protein specific for uracil-DNA glycosylase. *J Biol Chem.* Jan. 15, 1989;264(2):1163-71. cited by applicant

Warren et al., A chimeric Cre recombinase with regulated directionality. *Proc Natl Acad Sci USA.* Nov. 25, 2008;105(47):18278-83. doi: 10.1073/pnas.0809949105. Epub Nov. 14, 2008. cited by applicant

Warren et al., Mutations in the amino-terminal domain of lambda-integrase have differential effects on integrative and excisive recombination. *Mol Microbiol.* Feb. 2005;55(4):1104-12. cited by applicant

Watowich, The erythropoietin receptor: molecular structure and hematopoietic signaling pathways. *J Investig Med.* Oct. 2011;59(7):1067-72. doi: 10.2310/JIM.0b013e31820fb28c. cited by applicant

Waxman et al., Regulating excitability of peripheral afferents: emerging ion channel targets. *Nat Neurosci.* Feb. 2014;17(2):153-63. doi: 10.1038/nn.3602. Epub Jan. 28, 2014. cited by applicant

Weber et al., Assembly of designer TAL effectors by Golden Gate cloning. *PLoS One.* 2011;6(5):e19722. doi: 10.1371/journal.pone.0019722. Epub May 19, 2011. cited by applicant

Weill et al., DNA polymerases in adaptive immunity. *Nat Rev Immunol.* Apr. 2008;8(4):302-12. doi: 10.1038/nri2281. Epub Mar. 14, 2008. cited by applicant

Weinberg et al., New Classes of Self-Cleaving Ribozymes Revealed by Comparative Genomics Analysis. *Nat Chem Biol.* Aug. 2015;11(8):606-10. doi: 10.1038/nchembio.1846. Epub Jul. 13, 2015. cited by applicant

Weinberg et al., The aptamer core of SAM-IV riboswitches mimics the ligand-binding site of SAM-I riboswitches. *RNA.* May 2008;14(5):822-8. doi: 10.1261/rna.988608. Epub Mar. 27, 2008. cited by applicant

Weinberger et al., Disease-causing mutations C277R and C277Y modify gating of human CIC-1 chloride channels in myotonia congenita. *J Physiol.* Aug. 1, 2012;590(Pt 15):3449-64. doi: 10.1113/jphysiol.2012.232785. Epub May 28, 2012. cited by applicant

Weinert et al., Unbiased detection of CRISPR off-targets in vivo using DISCOVER-Seq. *Science.* Apr. 19, 2019;364(6437):286-289. doi: 10.1126/science.aav9023. Epub Apr. 18, 2019. cited by applicant

Weiss et al., Loss-of-function mutations in sodium channel Nav1.7 cause anosmia. *Nature.* Apr. 14, 2011;472(7342):186-90. doi: 10.1038/nature09975. Epub Mar. 23, 2011. cited by applicant

Wen et al., Inclusion of a universal tetanus toxoid CD4(+) T cell epitope P2 significantly enhanced the immunogenicity of

recombinant rotavirus ?VP8\* subunit parental vaccines. *Vaccine*. Jul. 31, 2014;32(35):4420-4427. doi: 10.1016/j.vaccine.2014.06.060. Epub Jun. 21, 2014. cited by applicant

West et al., Gene expression in adeno-associated virus vectors: the effects of chimeric mRNA structure, helper virus, and adenovirus VA1 RNA. *Virology*. Sep. 1987;160(1):38-47. doi: 10.1016/0042-6822(87)90041-9. cited by applicant

Wharton et al., A new-specificity mutant of 434 repressor that defines an amino acid-base pair contact. *Nature*. Apr. 30-May 6, 1987;326(6116):888-91. cited by applicant

Wharton et al., Changing the binding specificity of a repressor by redesigning an alpha-helix. *Nature*. Aug. 15-21, 1985;316(6029):601-5. cited by applicant

Wheeler et al., The thermostability and specificity of ancient proteins. *Curr Opin Struct Biol*. Jun. 2016;38:37-43. doi: 10.1016/j.sbi.2016.05.015. Epub Jun. 9, 2016. cited by applicant

Wiedenheft et al., RNA-guided genetic silencing systems in bacteria and archaea. *Nature*. Feb. 15, 2012;482(7385):331-8. doi: 10.1038/nature10886. Review. cited by applicant

Wienert et al., KLF1 drives the expression of fetal hemoglobin in British HPFH. *Blood*. Aug. 10, 2017;130(6):803-807. doi: 10.1182/blood-2017-02-767400. Epub Jun. 28, 2017. cited by applicant

Wijesinghe et al., Efficient deamination of 5-methylcytosines in DNA by human APOBEC3A, but not by AID or APOBEC3G. *Nucleic Acids Res*. Oct. 2012;40(18):9206-17. doi: 10.1093/nar/gks685. Epub Jul. 13, 2012. cited by applicant

Wijnker et al., Managing meiotic recombination in plant breeding. *Trends Plant Sci*. Dec. 2008;13(12):640-6. doi: 10.1016/j.tplants.2008.09.004. Epub Oct. 22, 2008. cited by applicant

Williams et al., Assessing the accuracy of ancestral protein reconstruction methods. *PLoS Comput Biol*. Jun. 23, 2006;2(6):e69. doi: 10.1371/journal.pcbi.0020069. Epub Jun. 23, 2006. cited by applicant

Wills et al., Pseudoknot-dependent read-through of retroviral gag termination codons: importance of sequences in the spacer and loop 2. *EMBO J*. Sep. 1, 1994;13(17):4137-44. doi: 10.1002/j.1460-2075.1994.tb06731.x. cited by applicant

Wilson et al., Assessing annotation transfer for genomics: quantifying the relations between protein sequence, structure and function through traditional and probabilistic scores. *J Mol Biol* 2000;297:233-49. cited by applicant

Wilson et al., Formation of infectious hybrid virions with gibbon ape leukemia virus and human T-cell leukemia virus retroviral envelope glycoproteins and the gag and pol proteins of Moloney murine leukemia virus. *J Virol*. May 1989;63(5):2374-8. doi: 10.1128/JVI.63.5.2374-2378.1989. cited by applicant

Wilson et al., In Vitro Selection of Functional Nucleic Acids. *Annu Rev Biochem*. 1999;68:611-47. doi: 10.1146/annurev.biochem.68.1.611. cited by applicant

Wilson et al., Kinase dynamics. Using ancient protein kinases to unravel a modern cancer drug's mechanism. *Science*. Feb. 20, 2015;347(6224):882-6. doi: 10.1126/science.aaa1823. cited by applicant

Winkler et al., An mRNA structure that controls gene expression by binding FMN. *Proc Natl Acad Sci U S A*. Dec. 10, 2002;99(25):15908-13. Epub Nov. 27, 2002. cited by applicant

Winkler et al., Control of gene expression by a natural metabolite-responsive ribozyme. *Nature*. Mar. 18, 2004;428(6980):281-6. cited by applicant

Winkler et al., Thiamine derivatives bind messenger RNAs directly to regulate bacterial gene expression. *Nature*. Oct. 31, 2002;419(6910):952-6. Epub Oct. 16, 2002. cited by applicant

Winoto et al., A novel, inducible and T cell-specific enhancer located at the 3' end of the T cell receptor alpha locus. *EMBO J*. Mar. 1989;8(3):729-33. cited by applicant

Winter et al., Drug Development. Phthalimide conjugation as a strategy for in vivo target protein degradation. *Science*. Jun. 19, 2015;348(6241):1376-81. doi: 10.1126/science.aab1433. Epub May 21, 2015. cited by applicant

Winter et al., Targeted exon skipping with AAV-mediated split adenine base editors. *Cell Discov*. Aug. 20, 2019;5:41. doi: 10.1038/s41421-019-0109-7. cited by applicant

Wold, Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism. *Annu Rev Biochem*. 1997;66:61-92. doi: 10.1146/annurev.biochem.66.1.61. cited by applicant

Wolf et al., tadA, an essential tRNA-specific adenosine deaminase from *Escherichia coli*. *EMBO J*. Jul. 15, 2002;21(14):3841-51. cited by applicant

Wolfe et al., Analysis of zinc fingers optimized via phage display: evaluating the utility of a recognition code. *J Mol Biol*. Feb. 5, 1999;285(5):1917-34. cited by applicant

Wong et al., A statistical analysis of random mutagenesis methods used for directed protein evolution. *J Mol Biol*. Jan. 27, 2006;355(4):858-71. Epub Nov. 17, 2005. cited by applicant

Wong et al., The Diversity Challenge in Directed Protein Evolution. *Comb Chem High Throughput Screen*. May 2006;9(4):271-88. cited by applicant

Wood et al., A genetic system yields self-cleaving inteins for bioseparations. *Nat Biotechnol*. Sep. 1999;17(9):889-92. doi: 10.1038/12879. cited by applicant

Wood et al., Targeted genome editing across species using ZFNs and TALENs. *Science*. Jul. 15, 2011;333(6040):307. doi: 10.1126/science.1207773. Epub Jun. 23, 2011. cited by applicant

Woods et al., The phenotype of congenital insensitivity to pain due to the NaV1.9 variant p.L811P. *Eur J Hum Genet*. May 2015;23(5):561-3. doi: 10.1038/ejhg.2014.166. Epub Aug. 13, 2014. cited by applicant

Wright et al., Continuous in vitro evolution of catalytic function. *Science*. Apr. 25, 1997;276(5312):614-7. cited by

applicant

- Wright et al., Rational design of a split-Cas9 enzyme complex. *Proc Natl Acad Sci U S A*. Mar. 10, 2015;112(10):2984-9. doi: 10.1073/pnas.1501698112. Epub Feb. 23, 2015. cited by applicant
- Wu et al., Correction of a genetic disease in mouse via use of CRISPR-Cas9. *Cell Stem Cell*. Dec. 5, 2013;13(6):659-62. doi: 10.1016/j.stem.2013.10.016. cited by applicant
- Wu et al., Genome-wide binding of the CRISPR endonuclease Cas9 in mammalian cells. *Nat Biotechnol*. Jul. 2014;32(7):670-6. doi: 10.1038/nbt.2889. Epub Apr. 20, 2014. cited by applicant
- Wu et al., Human single-stranded DNA binding proteins: guardians of genome stability. *Acta Biochim Biophys Sin (Shanghai)*. Jul. 2016;48(7):671-7. doi: 10.1093/abbs/gmw044. Epub May 23, 2016. cited by applicant
- Wu et al., Protein trans-splicing and functional mini-inteins of a cyanobacterial dnaB intein. *Biochim Biophys Acta*. Sep. 8, 1998;1387(1-2):422-32. doi: 10.1016/s0167-4838(98)00157-5. cited by applicant
- Wu et al., Protein trans-splicing by a split intein encoded in a split DnaE gene of *Synechocystis* sp. PCC6803. *Proc Natl Acad Sci U S A*. Aug. 4, 1998;95(16):9226-31. doi: 10.1073/pnas.95.16.9226. cited by applicant
- Wu et al., Readers, writers and erasers of N6-methylated adenosine modification. *Curr Opin Struct Biol*. Dec. 2017;47:67-76. doi: 10.1016/j.sbi.2017.05.011. Epub Jun. 16, 2017. cited by applicant
- Xiang et al., RNA m6A methylation regulates the ultraviolet-induced DNA damage response. *Nature*. Mar. 23, 2017;543(7646):573-576. doi: 10.1038/nature21671. Epub Mar. 15, 2017. cited by applicant
- Xiao et al., Genetic incorporation of multiple unnatural amino acids into proteins in mammalian cells. *Angew Chem Int Ed Engl*. Dec. 23, 2013;52(52):14080-3. doi: 10.1002/anie.201308137. Epub Nov. 8, 2013. cited by applicant
- Xiao et al., Nuclear m(6)A Reader YTHDC1 Regulates mRNA Splicing. *Mol Cell*. Feb. 18, 2016;61(4):507-519. doi: 10.1016/j.molcel.2016.01.012. Epub Feb. 11, 2016. cited by applicant
- Xie et al., Adjusting the attB site in donor plasmid improves the efficiency of  $\phi$ C31 integrase system. *DNA Cell Biol*. Jul. 2012;31(7):1335-40. doi: 10.1089/dna.2011.1590. Epub Apr. 10, 2012. cited by applicant
- Xiong et al., Origin and evolution of retroelements based upon their reverse transcriptase sequences. *EMBO J*. Oct. 1990;9(10):3353-62. cited by applicant
- Xu et al., Accuracy and efficiency define Bxb1 integrase as the best of fifteen candidate serine recombinases for the integration of DNA into the human genome. *BMC Biotechnol*. Oct. 20, 2013;13:87. doi: 10.1186/1472-6750-13-87. cited by applicant
- Xu et al., Chemical ligation of folded recombinant proteins: segmental isotopic labeling of domains for NMR studies. *Proc Natl Acad Sci U S A*. Jan. 19, 1999;96(2):388-93. doi: 10.1073/pnas.96.2.388. cited by applicant
- Xu et al., Multiplex nucleotide editing by high-fidelity Cas9 variants with improved efficiency in rice. *BMC Plant Biol*. 2019;19(1):511. Published Nov. 2, 2019. doi:10.1186/s12870-019-2131-1. Includes supplementary data and materials. cited by applicant
- Xu et al., Protein splicing: an analysis of the branched intermediate and its resolution by succinimide formation. *EMBO J*. Dec. 1, 1994;13(23):5517-22. cited by applicant
- Xu et al., PTMD: A Database of Human Disease-associated Post-translational Modifications. *Genomics Proteomics Bioinformatics*. Aug. 2018;16(4):244-251. doi: 10.1016/j.gpb.2018.06.004. Epub Sep. 21, 2018. cited by applicant
- Xu et al., Sequence determinants of improved CRISPR sgRNA design. *Genome Res*. Aug. 2015;25(8):1147-57. doi: 10.1101/gr.191452.115. Epub Jun. 10, 2015. cited by applicant
- Xu et al., Structures of human ALKBH5 demethylase reveal a unique binding mode for specific single-stranded N6-methyladenosine RNA demethylation. *J Biol Chem*. Jun. 20, 2014;289(25):17299-311. doi: 10.1074/jbc.M114.550350. Epub Apr. 28, 2014. cited by applicant
- Xu et al., The mechanism of protein splicing and its modulation by mutation. *EMBO J*. Oct. 1, 1996;15(19):5146-53. cited by applicant
- Yahata et al., Unified, Efficient, and Scalable Synthesis of Halichondrins: Zirconium/Nickel-Mediated One-Pot Ketone Synthesis as the Final Coupling Reaction. *Angew Chem Int Ed Engl*. Aug. 28, 2017;56(36):10796-10800. doi: 10.1002/anie.201705523. Epub Jul. 28, 2017. cited by applicant
- Yamada et al., Crystal Structure of the Minimal Cas9 from *Campylobacter jejuni* Reveals the Molecular Diversity in the CRISPR-Cas9 Systems. *Mol Cell*. Mar. 16, 2017;65(6):P1109-1121. /doi.org/10.1016/j.molcel.2017.02.007. cited by applicant
- Yamamoto et al., The ons and offs of inducible transgenic technology: a review. *Neurobiol Dis*. Dec. 2001;8(6):923-32. cited by applicant
- Yamamoto et al., Virological and immunological bases for HIV-1 vaccine design. *Uirusu* 2007;57(2):133-139. <https://doi.org/10.2222/jsv.57.133>. cited by applicant
- Yamano et al., Crystal Structure of Cpf1 in Complex with Guide RNA and Target DNA. *Cell*. May 5, 2016;165(4):949-62 and Supplemental Info. doi: 10.1016/j.cell.2016.04.003. Epub Apr. 21, 2016. cited by applicant
- Yamano et al., Crystal Structure of Cpf1 in Complex with Guide RNA and Target DNA. *Cell*. May 5, 2016;165(4):949-62. doi: 10.1016/j.cell.2016.04.003. Epub Apr. 21, 2016. cited by applicant
- Yamazaki et al., Segmental Isotope Labeling for Protein NMR Using Peptide Splicing. *J. Am. Chem. Soc*. May 22, 1998;120(22):5591-2. <https://doi.org/10.1021/ja9807760>. cited by applicant
- Yan et al., Cas13d Is a Compact RNA-Targeting Type VI CRISPR Effector Positively Modulated by a WYL-Domain-

Containing Accessory Protein. *Mol Cell*. Apr. 19, 2018;70(2):327-339.e5. doi: 10.1016/j.molcel.2018.02.028. Epub Mar. 15, 2018. cited by applicant

Yan et al., Functionally diverse type V CRISPR-Cas systems. *Science*. Jan. 4, 2019;363(6422):88-91. doi: 10.1126/science.aav7271. Epub Dec. 6, 2018. cited by applicant

Yan et al., Highly Efficient A·T to G·C Base Editing by Cas9n-Guided tRNA Adenosine Deaminase in Rice. *Mol Plant*. Apr. 2, 2018;11(4):631-634. doi: 10.1016/j.molp.2018.02.008. Epub Feb. 22, 2018. cited by applicant

Yang et al., APOBEC: From mutator to editor. *J Genet Genomics*. Sep. 20, 2017;44(9):423-437. doi: 10.1016/j.jgg.2017.04.009. Epub Aug. 7, 2017. cited by applicant

Yang et al., Construction of an integration-proficient vector based on the site-specific recombination mechanism of enterococcal temperate phage phiFC1. *J Bacteriol*. Apr. 2002;184(7):1859-64. doi: 10.1128/jb.184.7.1859-1864.2002. cited by applicant

Yang et al., Engineering and optimising deaminase fusions for genome editing. *Nat Commun*. Nov. 2, 2016;7:13330. doi: 10.1038/ncomms13330. cited by applicant

Yang et al., Genome editing with targeted deaminases. *BioRxiv*. Preprint. First posted online Jul. 28, 2016. cited by applicant

Yang et al., Genome-wide inactivation of porcine endogenous retroviruses (PERVs). *Science*. Nov. 27, 2015;350(6264):1101-4. doi: 10.1126/science.aad1191. Epub Oct. 11, 2015. cited by applicant

Yang et al., Increasing targeting scope of adenosine base editors in mouse and rat embryos through fusion of TadA deaminase with Cas9 variants. *Protein Cell*. Sep. 2018;9(9):814-819. doi: 10.1007/s13238-018-0568-x. cited by applicant

Yang et al., Mutations in SCN9A, encoding a sodium channel alpha subunit, in patients with primary erythralgia. *J Med Genet*. Mar. 2004;41(3):171-4. doi: 10.1136/jmg.2003.012153. cited by applicant

Yang et al., New CRISPR-Cas systems discovered. *Cell Res*. Mar. 2017;27(3):313-314. doi: 10.1038/cr.2017.21. Epub Feb. 21, 2017. cited by applicant

Yang et al., One Prime for All Editing. *Cell*. Dec. 12, 2019;179(7):1448-1450. doi: 10.1016/j.cell.2019.11.030. cited by applicant

Yang et al., One-step generation of mice carrying reporter and conditional alleles by CRISPR/Cas-mediated genome engineering. *Cell*. Sep. 12, 2013;154(6):1370-9. doi: 10.1016/j.cell.2013.08.022. Epub Aug. 29, 2013. cited by applicant

Yang et al., PAM-dependent Target DNA Recognition and Cleavage by C2C1 CRISPR-Cas endonuclease. *Cell*. Dec. 2016;167(7):1814-28. cited by applicant

Yang et al., Permanent genetic memory with >1-byte capacity. *Nat Methods*. Dec. 2014;11(12):1261-6. doi: 10.1038/nmeth.3147. Epub Oct. 26, 2014. cited by applicant

Yang et al., Preparation of RNA-directed DNA polymerase from spleens of Balb-c mice infected with Rauscher leukemia virus. *Biochem Biophys Res Commun*. Apr. 28, 1972;47(2):505-11. doi: 10.1016/0006-291x(72)90743-7. cited by applicant

Yang et al., Small-molecule control of insulin and PDGF receptor signaling and the role of membrane attachment. *Curr Biol*. Jan. 1, 1998;8(1):11-8. doi: 10.1016/s0960-9822(98)70015-6. cited by applicant

Yang, Development of Human Genome Editing Tools for the Study of Genetic Variations and Gene Therapies. Doctoral Dissertation. Harvard University. 2013. Accessible via nrs.harvard.edu/urn-3:HUL.InstRepos:11181072. 277 pages. cited by applicant

Yang, Nucleases: diversity of structure, function and mechanism. *Q Rev Biophys*. Feb. 2011;44(1):1-93. doi: 10.1017/S0033583510000181. Epub Sep. 21, 2010. cited by applicant

Yang, PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol*. Aug. 2007;24(8):1586-91. doi: 10.1093/molbev/msm088. Epub May 4, 2007. cited by applicant

Yanover et al., Extensive protein and DNA backbone sampling improves structure-based specificity prediction for C2H2 zinc fingers. *Nucleic Acids Res*. Jun. 2011;39(11):4564-76. doi: 10.1093/nar/gkr048. Epub Feb. 22, 2011. cited by applicant

Yasui et al., Miscoding Properties of 2'-Deoxyinosine, a Nitric Oxide-Derived DNA Adduct, during Translesion Synthesis Catalyzed by Human DNA Polymerases. *J Molec Biol*. Apr. 4, 2008;377(4):1015-23. cited by applicant

Yasui, Alternative excision repair pathways. *Cold Spring Harb Perspect Biol*. Jun. 1, 2013;5(6):a012617. doi: 10.1101/cshperspect.a012617. cited by applicant

Yasukawa et al., Characterization of Moloney murine leukaemia virus/avian myeloblastosis virus chimeric reverse transcriptases. *J Biochem*. Mar. 2009;145(3):315-24. doi: 10.1093/jb/mvn166. Epub Dec. 6, 2008. cited by applicant

Yazaki et al., Hereditary systemic amyloidosis associated with a new apolipoprotein AII stop codon mutation Stop78Arg. *Kidney Int*. Jul. 2003;64(1):11-6. cited by applicant

Yeh et al., In vivo base editing of post-mitotic sensory cells. *Nat Commun*. Jun. 5, 2018;9(1):2184. doi: 10.1038/s41467-018-04580-3. cited by applicant

Yin et al., Genome editing with Cas9 in adult mice corrects a disease mutation and phenotype. *Nat Biotechnol*. Jun. 2014;32(6):551-3. doi: 10.1038/nbt.2884. Epub Mar. 30, 2014. cited by applicant

Yokoe et al., Spatial dynamics of GFP-tagged proteins investigated by local fluorescence enhancement. *Nat Biotechnol*. Oct. 1996;14(10):1252-6. doi: 10.1038/nbt1096-1252. cited by applicant

Young et al., Beyond the canonical 20 amino acids: expanding the genetic lexicon. *J Biol Chem*. Apr. 9,

2010;285(15):11079-44. doi: 10.1074/jbc.R109.091306. Epub Feb. 10, 2010. cited by applicant

Yu et al., Circular permutation: a different way to engineer enzyme structure and function. *Trends Biotechnol.* Jan. 2011;29(1):18-25. doi: 10.1016/j.tibtech.2010.10.004. Epub Nov. 17, 2010. cited by applicant

Yu et al., Liposome-mediated in vivo ELA gene transfer suppressed dissemination of ovarian cancer cells that overexpress HER-2/neu. *Oncogene.* Oct. 5, 1995;11(7):1383-8. cited by applicant

Yu et al., Progress towards gene therapy for HIV infection. *Gene Ther.* Jan. 1994;1(1):13-26. cited by applicant

Yu et al., Small molecules enhance CRISPR genome editing in pluripotent stem cells. *Cell Stem Cell.* Feb. 5, 2015;16(2):142-7. doi: 10.1016/j.stem.2015.01.003. cited by applicant

Yu et al., Synthesis-dependent microhomology-mediated end joining accounts for multiple types of repair junctions. *Nucleic Acids Res.* Sep. 2010;38(17):5706-17. doi: 10.1093/nar/gkq379. Epub May 11, 2010. cited by applicant

Yuan et al., Laboratory-directed protein evolution. *Microbiol Mol Biol Rev.* 2005; 69(3):373-92. PMID: 16148303. cited by applicant

Yuan et al., Tetrameric structure of a serine integrase catalytic domain. *Structure.* Aug. 6, 2008;16(8):1275-86. doi: 10.1016/j.str.2008.04.018. cited by applicant

Yuen et al., Control of transcription factor activity and osteoblast differentiation in mammalian cells using an evolved small-molecule-dependent intein. *J Am Chem Soc.* Jul. 12, 2006;128(27):8939-46. cited by applicant

Zakas et al., Enhancing the pharmaceutical properties of protein drugs by ancestral sequence reconstruction. *Nat Biotechnol.* Jan. 2017;35(1):35-37. doi: 10.1038/nbt.3677. Epub Sep. 26, 2016. cited by applicant

Zalatan et al., Engineering complex synthetic transcriptional programs with CRISPR RNA scaffolds. *Cell.* Jan. 15, 2015;160(1-2):339-50. doi: 10.1016/j.cell.2014.11.052. Epub Dec. 18, 2014. cited by applicant

Zelphati et al., Intracellular delivery of proteins with a new lipid-mediated delivery system. *J Biol Chem.* Sep. 14, 2001;276(37):35103-10. Epub Jul. 10, 2001. cited by applicant

Zeng et al., Correction of the Marfan Syndrome Pathogenic FBN1 Mutation by Base Editing in Human Cells and Heterozygous Embryos. *Mol Ther.* Nov. 7, 2018;26(11):2631-2637. doi: 10.1016/j.ymthe.2018.08.007. Epub Aug. 14, 2018. cited by applicant

Zetsche et al., A split-Cas9 architecture for inducible genome editing and transcription modulation. *Nat Biotechnol.* Feb. 2015;33(2):139-42. doi: 10.1038/nbt.3149. cited by applicant

Zetsche et al., Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. *Cell.* Oct. 22, 2015;163(3):759-71 and Supplemental Info. doi: 10.1016/j.cell.2015.09.038. Epub Sep. 25, 2015. cited by applicant

Zetsche et al., Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. *Cell.* Oct. 22, 2015;163(3):759-71. doi: 10.1016/j.cell.2015.09.038. Epub Sep. 25, 2015. cited by applicant

Zettler et al., The naturally split Npu DnaE intein exhibits an extraordinarily high rate in the protein trans-splicing reaction. *FEBS Lett.* Mar. 4, 2009;583(5):909-14. doi: 10.1016/j.febslet.2009.02.003. Epub Feb. 10, 2009. cited by applicant

Zhang et al., II-Clamp-mediated cysteine conjugation. *Nat Chem.* Feb. 2016;8(2):120-8. doi: 10.1038/nchem.2413. Epub Dec. 21, 2015. cited by applicant

Zhang et al., A new strategy for the site-specific modification of proteins in vivo. *Biochemistry.* Jun. 10, 2003;42(22):6735-46. cited by applicant

Zhang et al., Circular intronic long noncoding RNAs. *Mol Cell.* Sep. 26, 2013;51(6):792-806. doi: 10.1016/j.molcel.2013.08.017. Epub Sep. 12, 2013. cited by applicant

Zhang et al., Comparison of non-canonical PAMs for CRISPR/Cas9-mediated DNA cleavage in human cells. *Sci Rep.* Jun. 2014;4:5405. cited by applicant

Zhang et al., Conditional gene manipulation: Cre-ating a new biological era. *J Zhejiang Univ Sci B.* Jul. 2012;13(7):511-24. doi: 10.1631/jzus.B1200042. Review. cited by applicant

Zhang et al., Copy number variation in human health, disease, and evolution. *Annu Rev Genomics Hum Genet.* 2009;10:451-81. doi: 10.1146/annurev.genom.9.081307.164217. cited by applicant

Zhang et al., CRISPR/Cas9 for genome editing: progress, implications and challenges. *Hum Mol Genet.* Sep. 15, 2014;23(R1):R40-6. doi: 10.1093/hmg/ddu125. Epub Mar. 20, 2014. cited by applicant

Zhang et al., Efficient construction of sequence-specific TAL effectors for modulating mammalian transcription. *Nat Biotechnol.* Feb. 2011;29(2):149-53. doi: 10.1038/nbt.1775. Epub Jan. 19, 2011. cited by applicant

Zhang et al., Global analysis of small RNA and mRNA targets of Hfq. *Mol Microbiol.* Nov. 2003;50(4):1111-24. doi: 10.1046/j.1365-2958.2003.03734.x. cited by applicant

Zhang et al., Myoediting: Toward Prevention of Muscular Dystrophy by Therapeutic Genome Editing. *Physiol Rev.* Jul. 1, 2018;98(3):1205-1240. doi: 10.1152/physrev.00046.2017. cited by applicant

Zhang et al., Programmable base editing of zebrafish genome using a modified CRISPR-Cas9 system. *Nat Commun.* Jul. 25, 2017;8(1):118. doi: 10.1038/s41467-017-00175-6. cited by applicant

Zhang et al., Reversible RNA Modification N1-methyladenosine (m1A) in mRNA and tRNA. *Genomics Proteomics Bioinformatics.* Jun. 2018;16(3):155-161. doi: 10.1016/j.gpb.2018.03.003. Epub Jun. 14, 2018. cited by applicant

Zhang et al., Ribozymes and Riboswitches: Modulation of RNA Function by Small Molecules. *Biochemistry.* Nov. 2, 2010;49(43):9123-31. doi: 10.1021/bi1012645. cited by applicant

Zhang et al., Stabilized plasmid-lipid particles for regional gene therapy: formulation and transfection properties. *Gene*



Ther. Aug. 1999;6(8):1438-47. cited by applicant  
 Zhao et al., An ultraprocessive, accurate reverse transcriptase encoded by a metazoan group II intron. RNA. Feb. 2018;24(2):183-195. doi: 10.1261/rna.063479.117. Epub Nov. 6, 2017. cited by applicant  
 Zhao et al., Crystal structures of a group II intron maturase reveal a missing link in spliceosome evolution. Nat Struct Mol Biol. Jun. 2016;23(6):558-65. doi: 10.1038/nsmb.3224. Epub May 2, 2016. cited by applicant  
 Zhao et al., Post-transcriptional gene regulation by mRNA modifications. Nat Rev Mol Cell Biol. Jan. 2017;18(1):31-42. doi: 10.1038/nrm.2016.132. Epub Nov. 3, 2016. cited by applicant  
 Zheng et al., ALKBH5 is a mammalian RNA demethylase that impacts RNA metabolism and mouse fertility. Mol Cell. Jan. 10, 2013;49(1):18-29. doi: 10.1016/j.molcel.2012.10.015. Epub Nov. 21, 2012. cited by applicant  
 Zheng et al., DNA editing in DNA/RNA hybrids by adenosine deaminases that act on RNA. Nucleic Acids Res. Apr. 7, 2017;45(6):3369-3377. doi: 10.1093/nar/gkx050. cited by applicant  
 Zheng et al., Highly efficient base editing in bacteria using a Cas9-cytidine deaminase fusion. Commun Biol. Apr. 19, 2018;1:32. doi: 10.1038/s42003-018-0035-5. cited by applicant  
 Zheng et al., Structural basis for the complete resistance of the human prion protein mutant G127V to prion disease. Sci Rep. Sep. 4, 2018;8(1):13211. doi: 10.1038/s41598-018-31394-6. cited by applicant  
 Zhong et al., Rational Design of Aptazyme Riboswitches for Efficient Control of Gene Expression in Mammalian Cells. Elife. Nov. 2, 2016;5:e18858. doi: 10.7554/eLife.18858. cited by applicant  
 Zhou et al., Dynamic m(6)A mRNA methylation directs translational control of heat shock response. Nature. Oct. 22, 2015;526(7574):591-4. doi: 10.1038/nature15377. Epub Oct. 12, 2015. cited by applicant  
 Zhou et al., GISSD: Group I Intron Sequence and Structure Database. Nucleic Acids Res. Jan. 2008;36(Database issue):D31-7. doi: 10.1093/nar/gkm766. Epub Oct. 16, 2007. cited by applicant  
 Zhou et al., Off-target RNA mutation induced by DNA base editing and its elimination by mutagenesis. Nature. Jul. 2019;571(7764):275-278. doi: 10.1038/s41586-019-1314-0. Epub Jun. 10, 2019. cited by applicant  
 Zhou et al., Protective V127 prion variant prevents prion disease by interrupting the formation of dimer and fibril from molecular dynamics simulations. Sci Rep. Feb. 24, 2016;6:21804. doi: 10.1038/srep21804. cited by applicant  
 Zhou et al., Seamless Genetic Conversion of SMN2 to SMN1 via CRISPR/Cpf1 and Single-Stranded Oligodeoxynucleotides in Spinal Muscular Atrophy Patient-Specific Induced Pluripotent Stem Cells. Hum Gene Ther. Nov. 2018;29(11):1252-1263. doi: 10.1089/hum.2017.255. Epub May 9, 2018. cited by applicant  
 Zielenski, Genotype and phenotype in cystic fibrosis. Respiration. 2000;67(2):117-33. doi: 10.1159/000029497. cited by applicant  
 Zimmerly et al., An Unexplored Diversity of Reverse Transcriptases in Bacteria. Microbiol Spectr. Apr. 2015;3(2):MDNA3-0058-2014. doi: 10.1128/microbiolspec.MDNA3-0058-2014. cited by applicant  
 Zimmerly et al., Group II intron mobility occurs by target DNA-primed reverse transcription. Cell. Aug. 25, 1995;82(4):545-54. doi: 10.1016/0092-8674(95)90027-6. cited by applicant  
 Zimmermann et al., Molecular interactions and metal binding in the theophylline-binding core of an RNA aptamer. RNA. May 2000;6(5):659-67. cited by applicant  
 Zolotukhin et al., Production and purification of serotype 1, 2, and 5 recombinant adeno-associated viral vectors. Methods. Oct. 2002;28(2):158-67. doi: 10.1016/s1046-2023(02)00220-7. cited by applicant  
 Zong et al., Precise base editing in rice, wheat and maize with a Cas9-cytidine deaminase fusion. Nat Biotechnol. May 2017;35(5):438-440. doi: 10.1038/nbt.3811. Epub Feb. 27, 2017. cited by applicant  
 Zorko et al., Cell-penetrating peptides: mechanism and kinetics of cargo delivery. Adv Drug Deliv Rev. Feb. 28, 2005;57(4):529-45. Epub Jan. 22, 2005. cited by applicant  
 Zou et al., Gene targeting of a disease-related gene in human induced pluripotent stem and embryonic stem cells. Cell Stem Cell. Jul. 2, 2009;5(1):97-110. doi: 10.1016/j.stem.2009.05.023. Epub Jun. 18, 2009. cited by applicant  
 Zufferey et al., Woodchuck hepatitis virus posttranscriptional regulatory element enhances expression of transgenes delivered by retroviral vectors. J Virol. Apr. 1999;73(4):2886-92. doi: 10.1128/JVI.73.4.2886-2892.1999. cited by applicant  
 Zuker et al., Optimal computer folding of large RNA sequences using thermodynamics and auxiliary information. Nucleic Acids Res. Jan. 10, 1981;9(1):133-48. doi: 10.1093/nar/9.1.133. cited by applicant  
 Zuo et al., Cytosine base editor generates substantial off-target single-nucleotide variants in mouse embryos. Science. Apr. 19, 2019;364(6437):289-292. doi: 10.1126/science.aav9973. Epub Feb. 28, 2019. cited by applicant  
 Zuris et al., Cationic lipid-mediated delivery of proteins enables efficient protein-based genome editing in vitro and in vivo. Nat Biotechnol. 2015;33:73-80. cited by applicant  
 [No Author Listed], "Lambda DNA" from Catalog & Technical Reference. New England Biolabs Inc. 2002/2003. pp. 133 and 270-273. cited by applicant  
 [No Author Listed], Gag-Pol polyprotein. UniProtKB/Swiss-Prot No. P03355.5. Sep. 18, 2019. 18 pages. cited by applicant  
 [No Author Listed], *Homo sapiens* signal transducer and activator of transcription 3 (STAT3), transcript variant 1, mRNA. NCBI Ref Seq No. NM\_139276.2. Retrived from [https://www.ncbi.nlm.nih.gov/nuccore/nm\\_139276.2](https://www.ncbi.nlm.nih.gov/nuccore/nm_139276.2). Feb. 26, 2020. 8 pages. cited by applicant  
 Ai et al., C-terminal Loop Mutations Determine Folding and Secretion Properties of PCSK9. iMedPub J: Biochem Mol

Biol J. Nov. 5, 2016;2(3):17. doi: 10.21767/2471-8084.100026. 12 pages. cited by applicant

Andersen et al., High frequency of T cells specific for cryptic epitopes in melanoma patients. *Oncoimmunology*. Jul. 1, 2013;2(7):e25374. doi: 10.4161/onci.25374. 7 pages. cited by applicant

Anzalone et al., Search-and-replace genome editing without double-strand breaks or donor DNA. *Nature*. Dec. 2019;576(7785):149-157 and Suppl Info. doi: 10.1038/s41586-019-1711-4. Epub Oct. 21, 2019. 72 pages. cited by applicant

Baba et al., Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol Syst Biol*. 2006;2:2006.0008. doi: 10.1038/msb4100050. Epub Feb. 21, 2006. cited by applicant

Bass, B.L., RNA editing by adenosine deaminases that act on RNA. *Annu Rev Biochem*. 2002;71:817-46. doi: 10.1146/annurev.biochem.71.110601.135501. Epub Nov. 9, 2001. cited by applicant

Bibikova et al., Targeted chromosomal cleavage and mutagenesis in *Drosophila* using zinc-finger nucleases. *Genetics*. Jul. 2002;161(3):1169-75. doi: 10.1093/genetics/161.3.1169. cited by applicant

Blauw et al., SMN1 gene duplications are associated with sporadic ALS. *Neurology*. Mar. 13, 2012;78(11):776-80. doi: 10.1212/WNL.0b013e318249f697. Epub Feb. 8, 2012. cited by applicant

Bothmer et al., Characterization of the interplay between DNA repair and CRISPR/Cas9-induced DNA lesions at an endogenous locus. *Nat Commun*. Jan. 9, 2017;8:13905. doi: 10.1038/ncomms13905. cited by applicant

Brutlag et al., Improved sensitivity of biological sequence database searches. *Comput Appl Biosci*. Jul. 1990;6(3):237-45. doi: 10.1093/bioinformatics/6.3.237. cited by applicant

Buonaguro et al., Translating tumor antigens into cancer vaccines. *Clin Vaccine Immunol*. Jan. 2011;18(1):23-34. doi: 10.1128/CVI.00286-10. Epub Nov. 3, 2010. cited by applicant

Campos-Perez et al., DNA fusion vaccine designs to induce tumor-lytic CD8+ T-cell attack via the immunodominant cysteine-containing epitope of NY-ESO 1. *Int J Cancer*. Sep. 15, 2013;133(6):1400-7. doi: 10.1002/ijc.28156. Epub Apr. 11, 2013. cited by applicant

Canny et al., Inhibition of 53BP1 Favors Homology-Dependent DNA Repair and Increases CRISPR-Cas9 Genome-Editing Efficiency. *Nat Biotechnol*. Jan. 2018;36(1):95-102. doi: 10.1038/nbt.4021. Epub Nov. 27, 2017. cited by applicant

Cao et al., Rapamycin reverses cellular phenotypes and enhances mutant protein clearance in Hutchinson-Gilford progeria syndrome cells. *Sci Transl Med*. Jun. 29, 2011;3(89):89ra58. doi: 10.1126/scitranslmed.3002346. cited by applicant

Carlier et al., Genome Sequence of Burkholderia cenocepacia H111, a Cystic Fibrosis Airway Isolate. *Genome Announc*. Apr. 10, 2014;2(2):e00298-14. doi: 10.1128/genomeA.00298-14. cited by applicant

Cartegni et al., Determinants of exon 7 splicing in the spinal muscular atrophy genes, SMN1 and SMN2. *Am J Hum Genet*. Jan. 2006;78(1):63-77. doi: 10.1086/498853. Epub Nov. 16, 2005. cited by applicant

Chang et al., Degradation of survival motor neuron (SMN) protein is mediated via the ubiquitin/proteasome pathway. *Neurochem Int*. Dec. 2004;45(7):1107-12. doi: 10.1016/j.neuint.2004.04.005. cited by applicant

Chatterjee et al., Robust Genome Editing of Single-Base PAM Targets; with Engineered ScCas9 Variants. *bioRxiv*. doi: 10.1101/620351. Posted Apr. 26, 2019. cited by applicant

Cheng et al., [Cloning, expression and activity identification of human innate immune protein apolipoprotein B mRNA editing enzyme catalytic subunit 3A(APOBEC3A)]. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi*. Chinese Journal of Cellular and Molecular Immunology, Feb. 2017;33(2):179-84. Chinese. cited by applicant

Cho et al., A degron created by SMN2 exon 7 skipping is a principal contributor to spinal muscular atrophy severity. *Genes Dev*. Mar. 1, 2010;24(5):438-42. doi: 10.1101/gad.1884910. cited by applicant

Chowell et al., TCR contact residue hydrophobicity is a hallmark of immunogenic CD8+ T cell epitopes. *Proc Natl Acad Sci U S A*. Apr. 7, 2015;112(14):E1754-62. doi: 10.1073/pnas.1500973112. Epub Mar. 23, 2015. cited by applicant

Corcia et al., The importance of the SMN genes in the genetics of sporadic ALS. *Amyotroph Lateral Scler*. Oct.-Dec. 2009;10(5-6):436-40. doi: 10.3109/17482960902759162. cited by applicant

Corti et al., Genetic correction of human induced pluripotent stem cells from patients with spinal muscular atrophy. *Sci Transl Med*. Dec. 19, 2012;4(165):165ra162. doi: 10.1126/scitranslmed.3004108. cited by applicant

Cucchiari et al., Enhanced expression of the central survival of motor neuron (SMN) protein during the pathogenesis of osteoarthritis. *J Cell Mol Med*. Jan. 2014;18(1):115-24. doi: 10.1111/jcmm.12170. Epub Nov. 17, 2013. cited by applicant

Davis et al., Assaying Repair at DNA Nicks. *Methods Enzymol*. 2018;601:71-89. doi: 10.1016/bs.mie.2017.12.001. Epub Feb. 1, 2018. cited by applicant

Davis et al., Homology-directed repair of DNA nicks via pathways distinct from canonical double-strand break repair. *Proc Natl Acad Sci U S A*. Mar. 11, 2014;111(10):E924-32. doi: 10.1073/pnas.1400236111. Epub Feb. 20, 2014. cited by applicant

Davis et al., Two Distinct Pathways Support Gene Correction by Single-Stranded Donors at DNA Nicks. *Cell Rep*. Nov. 8, 2016;17(7):1872-1881. doi: 10.1016/j.celrep.2016.10.049. cited by applicant

De Sandre-Giovannoli et al., Lamin a truncation in Hutchinson-Gilford progeria. *Science*. Jun. 27, 2003;300(5628):2055. doi: 10.1126/science.1084125. Epub Apr. 17, 2003. cited by applicant

Dickinson et al., A system for the continuous directed evolution of proteases rapidly reveals drug-resistance mutations. *Nat Commun*. Oct. 30, 2014;5:5352. doi: 10.1038/ncomms6352. cited by applicant

Drenth et al., Mutations in sodium-channel gene SCN9A cause a spectrum of human genetic pain disorders. *J Clin Invest*.

Dec. 2007;117(12):3603-9. doi: 10.1172/JCI33297. cited by applicant

Dugar et al., CRISPR RNA-Dependent Binding and Cleavage of Endogenous RNAs by the *Campylobacter jejuni* Cas9. *Mol Cell*. Mar. 1, 2018;69(5):893-905.e7. doi: 10.1016/j.molcel.2018.01.032. cited by applicant

D'Ydewalle et al., The Antisense Transcript SMN-AS1 Regulates SMN Expression and Is a Novel Therapeutic Target for Spinal Muscular Atrophy. *Neuron*. Jan. 4, 2017;93(1):66-79 and Supplemental Information. doi: 10.1016/j.neuron.2016.11.033. Epub Dec. 22, 2016. cited by applicant

Eisenberg et al., A-to-I RNA editing—immune protector and transcriptome diversifier. *Nat Rev Genet*. Aug. 2018;19(8):473-490. doi: 10.1038/s41576-018-0006-1. cited by applicant

Ekstrand et al., Frequent alterations of the PI3K/AKT/mTOR pathways in hereditary nonpolyposis colorectal cancer. *Fam Cancer*. Jun. 2010;9(2):125-9. doi: 10.1007/s10689-009-9293-1. cited by applicant

Entin-Meer et al., The role of phenylalanine-119 of the reverse transcriptase of mouse mammary tumour virus in DNA synthesis, ribose selection and drug resistance. *Biochem J*. Oct. 15, 2002;367(Pt 2):381-91. doi: 10.1042/BJ20020712. cited by applicant

Friedman, J. H., Greedy function approximation: A gradient boosting machine. *Ann. Statist*. Oct. 2001;29(5):1189-232. doi: 10.1214/aos/1013203451. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NC\_000001.11. Gregory et al., Jun. 6, 2016. 3 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NG\_008692.2. McClintock et al., Aug. 27, 2018. 33 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NM\_206933.2. Khalaileh et al., Sep. 16, 2018. 12 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_001075493.1. Schiaffella et al., Jun. 24, 2018. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_001157741.1. Zeng et al., Sep. 17, 2018. 3 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_001157742.1. Zeng et al., Oct. 21, 2018. 3 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_033040.2. Liu et al., Jun. 23, 2018. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_996816.2. Fu et al., Sep. 22, 2019. 9 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. XP\_003314669.1. No Author Listed, Mar. 20, 2018. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. XP\_026671085.1. No Author Listed, Oct. 17, 2018. 1 page. cited by applicant

Grati et al., Localization of PDZD7 to the stereocilia ankle-link associates this scaffolding protein with the Usher syndrome protein network. *J Neurosci*. Oct. 10, 2012;32(41):14288-93. doi: 10.1523/JNEUROSCI.3071-12.2012. cited by applicant

Gutschner et al., Post-translational Regulation of Cas9 during G1 Enhances Homology-Directed Repair. *Cell Rep*. Feb. 16, 2016;14(6):1555-1566. doi: 10.1016/j.celrep.2016.01.019. Epub Feb. 4, 2016. cited by applicant

Hagen et al., A high rate of polymerization during synthesis of mouse mammary tumor virus DNA alleviates hypermutation by APOBEC3 proteins. *PLoS Pathog*. Feb. 15, 2019;15(2):e1007533. doi: 10.1371/journal.ppat.1007533. cited by applicant

Harrington et al., Programmed DNA destruction by miniature CRISPR-Cas14 enzymes. *Science*. Nov. 16, 2018;362(6416):839-842. doi: 10.1126/science.aav4294. Epub Oct. 18, 2018. cited by applicant

Hart et al., High-Resolution CRISPR Screens Reveal Fitness Genes and Genotype-Specific Cancer Liabilities. *Cell*. Dec. 3, 2015;163(6):1515-26. doi: 10.1016/j.cell.2015.11.015. Epub Nov. 25, 2015. cited by applicant

Hawley-Nelson et al., Transfection of Cultured Eukaryotic Cells Using Cationic Lipid Reagents. *Curr Prot Mol Biol*. Jan. 2008;9.4.1-9.4.17. doi: 10.102/0471142727.mb0904s81. 17 pages. cited by applicant

Hendel et al., Chemically modified guide RNAs enhance CRISPR-Cas genome editing in human primary cells. *Nat Biotechnol*. Sep. 2015;33(9):985-989. doi: 10.1038/nbt.3290. Epub Jun. 29, 2015. Author Manuscript. 14 pages. cited by applicant

Heyer et al., Regulation of homologous recombination in eukaryotes. *Annu Rev Genet*. 2010;44:113-39. doi: 10.1146/annurev-genet-051710-150955. Author Manuscript. 33 pages. cited by applicant

Huang et al., Gain-of-function mutations in sodium channel Na(v)1.9 in painful neuropathy. *Brain*. Jun. 2014;137(Pt 6):1627-42. doi: 10.1093/brain/awu079. Epub Apr. 27, 2014. cited by applicant

Ishizuka et al., Loss of ADAR1 in tumours overcomes resistance to immune checkpoint blockade. *Nature*. Jan. 2019;565(7737):43-48. doi: 10.1038/s41586-018-0768-9. Epub Dec. 17, 2018. cited by applicant

Iyama et al., DNA repair mechanisms in dividing and non-dividing cells. *DNA Repair (Amst)*. Aug. 2013;12(8):620-36. doi: 10.1016/j.dnarep.2013.04.015. Epub May 16, 2013. cited by applicant

Jakimo et al., A Cas9 with Complete PAM Recognition for Adenine Dinucleotides. *bioRxiv preprint*. Sep. 27, 2018. doi.org/10.1101/429654. 29 pages. cited by applicant

Kan et al., Mechanisms of precise genome editing using oligonucleotide donors. *Genome Res*. Jul. 2017;27(7):1099-1111. doi: 10.1101/gr.214775.116. Epub Mar. 29, 2017. cited by applicant

Kim et al., Adenine base editors catalyze cytosine conversions in human cells. *Nat Biotechnol*. Oct. 2019;37(10):1145-

1148. doi: 10.1038/s41587-019-0254-4. Epub Sep. 23, 2019. cited by applicant

Kim et al., RAD51 mutants cause replication defects and chromosomal instability. *Mol Cell Biol.* Sep. 2012;32(18):3663-80. doi: 10.1128/MCB.00406-12. Epub Jul. 9, 2012. cited by applicant

Knott et al., CRISPR-Cas guides the future of genetic engineering. *Science.* Aug. 31, 2018;361(6405):866-869. doi: 10.1126/science.aat5011. cited by applicant

Konishi et al., Amino acid substitutions away from the RNase H catalytic site increase the thermal stability of Moloney murine leukemia virus reverse transcriptase through RNase H inactivation. *Biochem Biophys Res Commun.* Nov. 14, 2014;454(2):269-74. doi: 10.1016/j.bbrc.2014.10.044. Epub Oct. 17, 2014. cited by applicant

Kreiter et al., Mutant MHC class II epitopes drive therapeutic immune responses to cancer. *Nature.* Apr. 30, 2015;520(7549):692-6. doi: 10.1038/nature14426. Epub Apr. 22, 2015. Erratum in: *Nature.* Jul. 16, 2015;523(7560):370. cited by applicant

Kuan et al., A systematic evaluation of nucleotide properties for CRISPR sgRNA design. *BMC Bioinformatics.* Jun. 6, 2017;18(1):297. doi: 10.1186/s12859-017-1697-6. cited by applicant

Kweon et al., A CRISPR-based base-editing screen for the functional assessment of BRCA1 variants. *Oncogene.* Jan. 2020;39(1):30-35. doi: 10.1038/s41388-019-0968-2. Epub Aug. 29, 2019. cited by applicant

Langer et al., Chemical and Physical Structure of Polymers as Carriers for Controlled Release of Bioactive Agents: A Review. *J Macromol Sci, Part C,* 1983;23(1):61-126. doi: 10.1080/07366578308079439. cited by applicant

Le et al., SMNDelta7, the major product of the centromeric survival motor neuron (SMN2) gene, extends survival in mice with spinal muscular atrophy and associates with full-length SMN. *Hum Mol Genet.* Mar. 15, 2005;14(6):845-57. doi: 10.1093/hmg/ddi078. Epub Feb. 9, 2005. cited by applicant

Lefebvre et al., Identification and characterization of a spinal muscular atrophy-determining gene. *Cell.* Jan. 13, 1995;80(1):155-65. doi: 10.1016/0092-8674(95)90460-3. cited by applicant

Lesinski et al., The potential for targeting the STAT3 pathway as a novel therapy for melanoma. *Future Oncol.* Jul. 2013;9(7):925-7. doi: 10.2217/fon.13.83. Author Manuscript. 4 pages. cited by applicant

Lin et al., [Construction and evaluation of DnaB split intein high expression vector and a six amino acids cyclic peptide library]. *Sheng Wu Gong Cheng Xue Bao.* Nov. 2008;24(11):1924-30. Chinese. cited by applicant

Lindahl, T., Instability and decay of the primary structure of DNA. *Nature.* Apr. 22, 1993;362(6422):709-15. doi: 10.1038/362709a0. cited by applicant

Liu et al., Human BRCA2 protein promotes RAD51 filament formation on RPA-covered single-stranded DNA. *Nat Struct Mol Biol.* Oct. 2010; 17(10):1260-2. doi: 10.1038/nsmb.1904. Epub Aug. 22, 2010. cited by applicant

Liu et al., Improving Editing Efficiency for the Sequences with NGH PAM Using xCas9-Derived Base Editors. *Mol Ther Nucleic Acids.* Sep. 6, 2019;17:626-635. doi: 10.1016/j.omtn.2019.06.024. Epub Jul. 12, 2019. cited by applicant

Liu et al., Intrinsic Nucleotide Preference of Diversifying Base Editors Guides Antibody Ex Vivo Affinity Maturation. *Cell Rep.* Oct. 23, 2018;25(4):884-892.e3. doi: 10.1016/j.celrep.2018.09.090. cited by applicant

Liu et al., Usherin is required for maintenance of retinal photoreceptors and normal development of cochlear hair cells. *Proc Natl Acad Sci U S A.* Mar. 13, 2007;104(11):4413-8. doi: 10.1073/pnas.0610950104. Epub Mar. 5, 2007. cited by applicant

Lorson et al., A single nucleotide in the SMN gene regulates splicing and is responsible for spinal muscular atrophy. *Proc Natl Acad Sci U S A.* May 25, 1999;96(11):6307-11. doi: 10.1073/pnas.96.11.6307. cited by applicant

Lutz et al., Postsymptomatic restoration of SMN rescues the disease phenotype in a mouse model of severe spinal muscular atrophy. *J Clin Invest.* Aug. 2011;121(8):3029-41. doi: 10.1172/JCI57291. Epub Jul. 25, 2011. cited by applicant

Ma et al., Human RAD52 interactions with replication protein A and the RAD51 presynaptic complex. *J Biol Chem.* Jul. 14, 2017;292(28):11702-11713. doi: 10.1074/jbc.M117.794545. Epub May 27, 2017. cited by applicant

Maerker et al., A novel Usher protein network at the periciliary reloading point between molecular transport machineries in vertebrate photoreceptor cells. *Hum Mol Genet.* Jan. 1, 2008;17(1):71-86. doi: 10.1093/hmg/ddm285. Epub Sep. 28, 2007. cited by applicant

Mali et al., CAS9 transcriptional activators for target specificity screening and paired nickases for cooperative genome engineering. *Nat Biotechnol.* Sep. 2013;31(9):Supplemental Info. doi: 10.1038/nbt.2675. Epub Aug. 1, 2013. cited by applicant

Marcovitz et al., Frustration in protein-DNA binding influences conformational switching and target search kinetics. *Proc Natl Acad Sci U S A.* Nov. 1, 2011;108(44):17957-62. doi: 10.1073/pnas.1109594108. Epub Oct. 14, 2011. cited by applicant

Marsden et al., The Tumor-Associated Variant RAD51 G151D Induces a Hyper-Recombination Phenotype. *PLOS Genet.* Aug. 11, 2016;12(8):e1006208. doi: 10.1371/journal.pgen.1006208. cited by applicant

Mason et al., Non-enzymatic roles of human RAD51 at stalled replication forks. *bioRxiv.* Jul. 31, 2019; doi.org/10.1101/359380. 36 pages. bioRxiv preprint first posted online Jul. 31, 2019. cited by applicant

Melero et al., Therapeutic vaccines for cancer: an overview of clinical trials. *Nat Rev Clin Oncol.* Sep. 2014;11(9):509-24. doi: 10.1038/nrclinonc.2014.111. Epub Jul. 8, 2014. cited by applicant

Mendell et al., Single-Dose Gene-Replacement Therapy for Spinal Muscular Atrophy. *N Engl J Med.* Nov. 2, 2017;377(18):1713-1722. doi: 10.1056/NEJMoa1706198. cited by applicant

Min et al., Deep learning in bioinformatics. *Brief Bioinform.* Sep. 1, 2017;18(5):851-869. doi: 10.1093/bib/bbw068. cited

by applicant

Monani et al., A single nucleotide difference that alters splicing patterns distinguishes the SMA gene SMN1 from the copy gene SMN2. *Hum Mol Genet.* Jul. 1999;8(7):1177-83. doi: 10.1093/hmg/8.7.1177. cited by applicant

Murray et al., Selective vulnerability of motor neurons and dissociation of pre- and post-synaptic pathology at the neuromuscular junction in mouse models of spinal muscular atrophy. *Hum Mol Genet.* Apr. 1, 2008;17(7):949-62. doi: 10.1093/hmg/ddm367. Epub Dec. 8, 2007. cited by applicant

Murugan et al., The Revolution Continues: Newly Discovered Systems Expand the CRISPR-Cas Toolkit. *Mol Cell.* Oct. 5, 2017;68(1):15-25. doi: 10.1016/j.molcel.2017.09.007. cited by applicant

Nelson et al., In vivo genome editing improves muscle function in a mouse model of Duchenne muscular dystrophy. *Science.* Jan. 22, 2016;351(6271):403-7. doi: 10.1126/science.aad5143. Epub Dec. 31, 2015. cited by applicant

Ottesen, ISS-N1 makes the First FDA-approved Drug for Spinal Muscular Atrophy. *Transl Neurosci.* Jan. 26, 2017;8:1-6. doi: 10.1515/tnsci-2017-0001. cited by applicant

Ousterout et al., Multiplex CRISPR/Cas9-based genome editing for correction of dystrophin mutations that cause Duchenne muscular dystrophy. *Nat Commun.* Feb. 18, 2015;6:6244. doi: 10.1038/ncomms7244. cited by applicant

Parente et al., Advances in spinal muscular atrophy therapeutics. *Ther Adv Neurol Disord.* Feb. 5, 2018;11:1756285618754501. doi: 10.1177/1756285618754501. 13 pages. cited by applicant

Passini et al., Antisense oligonucleotides delivered to the mouse CNS ameliorate symptoms of severe spinal muscular atrophy. *Sci Transl Med.* Mar. 2, 2011;3(72):72ra18. doi: 10.1126/scitranslmed.3001777. cited by applicant

Pellegrini et al., Insights into DNA recombination from the structure of a RAD51-BRCA2 complex. *Nature.* Nov. 21, 2002;420(6913):287-93. doi: 10.1038/nature01230. Epub Nov. 10, 2002. cited by applicant

Pendse et al., Exon 13-skipped USH2A protein retains functional integrity in mice, suggesting an exon-skipping therapeutic approach to treat USH2A-associated disease. *bioRxiv preprint.* Feb. 4, 2020. Retrieved from [www.biorxiv.org](http://www.biorxiv.org). doi: 10.1101/2020.02.04.934240. 34 pages. cited by applicant

Pendse et al., In Vivo Assessment of Potential Therapeutic Approaches for USH2A-Associated Diseases. *Adv Exp Med Biol.* 2019;1185:91-96. doi: 10.1007/978-3-030-27378-1\_15. cited by applicant

Perez-Palma et al., Simple ClinVar: an interactive web server to explore and retrieve gene and disease variants aggregated in ClinVar database. *Nucleic Acids Res.* Jul. 2, 2019;47(W1):W99-W105. doi: 10.1093/nar/gkz411. cited by applicant

Porensky et al., A single administration of morpholino antisense oligomer rescues spinal muscular atrophy in mouse. *Hum Mol Genet.* Apr. 1, 2012;21(7):1625-38. doi: 10.1093/hmg/ddr600. Epub Dec. 20, 2011. cited by applicant

Prasad et al., Visualizing the assembly of human Rad51 filaments on double-stranded DNA. *J Mol Biol.* Oct. 27, 2006;363(3):713-28. doi: 10.1016/j.jmb.2006.08.046. Epub Aug. 22, 2006. cited by applicant

Purcell et al., More than one reason to rethink the use of peptides in vaccine design. *Nat Rev Drug Discov.* May 2007;6(5):404-14. doi: 10.1038/nrd2224. cited by applicant

Rajagopal et al., High-throughput mapping of regulatory DNA. *Nat Biotechnol.* Feb. 2016;34(2):167-74. doi: 10.1038/nbt.3468. Epub Jan. 25, 2016. cited by applicant

Ramos et al., Age-dependent SMN expression in disease-relevant tissue and implications for SMA treatment. *J Clin Invest.* Nov. 1, 2019;129(11):4817-4831. doi: 10.1172/JCI124120. cited by applicant

Reiners et al., Scaffold protein harmonin (USH1C) provides molecular links between Usher syndrome type 1 and type 2. *Hum Mol Genet.* Dec. 15, 2005;14(24):3933-43. doi: 10.1093/hmg/ddi417. Epub Nov. 21, 2005. cited by applicant

Richardson et al., CRISPR-Cas9 genome editing in human cells occurs via the Fanconi anemia pathway. *Nat Genet.* Aug. 2018;50(8):1132-1139. doi: 10.1038/s41588-018-0174-0. Epub Jul. 27, 2018. cited by applicant

Richardson et al., Frequent chromosomal translocations induced by DNA double-strand breaks. *Nature.* Jun. 8, 2000;405(6787):697-700. doi: 10.1038/35015097. cited by applicant

Rivoltini et al., A superagonist variant of peptide MART1/Melan A27-35 elicits anti-melanoma CD8+ T cells with enhanced functional characteristics: implication for more effective immunotherapy. *Cancer Res.* Jan. 15, 1999;59(2):301-6. 18 pages. cited by applicant

Rodriguez-Muela et al., Single-Cell Analysis of SMN Reveals Its Broader Role in Neuromuscular Disease. *Cell Rep.* Feb. 7, 2017;18(6):1484-1498 and Supplemental Information. doi: 10.1016/j.celrep.2017.01.035. cited by applicant

Saayman et al., The therapeutic application of CRISPR/Cas9 technologies for HIV. *Expert Opin Biol Ther.* Jun. 2015;15(6):819-30. doi: 10.1517/14712598.2015.1036736. Epub Apr. 12, 2015. cited by applicant

San Filippo et al., Mechanism of eukaryotic homologous recombination. *Annu Rev Biochem.* 2008;77:229-57. doi: 10.1146/annurev.biochem.77.061306.125255. cited by applicant

Sanjurjo-Soriano et al., Genome Editing in Patient iPSCs Corrects the Most Prevalent USH2A Mutations and Reveals Intriguing Mutant mRNA Expression Profiles. *Mol Ther Methods Clin Dev.* Nov. 27, 2019;17:156-173. doi: 10.1016/j.omtm.2019.11.016. cited by applicant

Schlacher et al., Double-strand break repair-independent role for BRCA2 in blocking stalled replication fork degradation by MRE11. *Cell.* May 13, 2011;145(4):529-42. doi: 10.1016/j.cell.2011.03.041. Erratum in: *Cell.* Jun. 10, 2011;145(6):993. cited by applicant

Schrank et al., Inactivation of the survival motor neuron gene, a candidate gene for human spinal muscular atrophy, leads to massive cell death in early mouse embryos. *Proc Natl Acad Sci USA.* Sep. 2, 1997;94(18):9920-5. doi: 10.1073/pnas.94.18.9920. cited by applicant

Shen et al., Efficient genome modification by CRISPR-Cas9 nickase with minimal off-target effects. *Nat Methods*. Apr. 2014;11(4):399-402. doi: 10.1038/nmeth.2857. Epub Mar. 2, 2014. cited by applicant

Singh et al., Splicing of a critical exon of human Survival Motor Neuron is regulated by a unique silencer element located in the last intron. *Mol Cell Biol*. Feb. 2006;26(4):1333-46. doi: 10.1128/MCB.26.4.1333-1346.2006. cited by applicant

Song et al., RS-1 enhances CRISPR/Cas9- and TALEN-mediated knock-in efficiency. *Nat Commun*. Jan. 28, 2016;7:10548. doi: 10.1038/ncomms10548. cited by applicant

Sorusch et al., Characterization of the ternary Usher syndrome SANS/ush2a/whirlin protein complex. *Hum Mol Genet*. Mar. 15, 2017;26(6):1157-1172. doi: 10.1093/hmg/ddx027. cited by applicant

Stark et al., ATP hydrolysis by mammalian RAD51 has a key role during homology-directed DNA repair. *J Biol Chem*. Jun. 7, 2002;277(23):20185-94. doi: 10.1074/jbc.M112132200. Epub Mar. 28, 2002. cited by applicant

Sumner et al., Two breakthrough gene-targeted treatments for spinal muscular atrophy: challenges remain. *J Clin Invest*. Aug. 1, 2018;128(8):3219-3227. doi: 10.1172/JCI121658. Epub Jul. 9, 2018. cited by applicant

Talbot et al., Spinal muscular atrophy. *Semin Neurol*. Jun. 2001;21(2):189-97. doi: 10.1055/s-2001-15264. cited by applicant

Tan et al., Engineering of high-precision base editors for site-specific single nucleotide replacement. *Nat Commun*. Jan. 25, 2019;10(1):439. doi: 10.1038/s41467-018-08034-8. Erratum in: *Nat Commun*. May 1, 2019;10(1):2019. cited by applicant

Vakulskas et al., A high-fidelity Cas9 mutant delivered as a ribonucleoprotein complex enables efficient gene editing in human hematopoietic stem and progenitor cells. *Nat Med*. Aug. 2018;24(8):1216-1224. doi: 10.1038/s41591-018-0137-0. Epub Aug. 6, 2018. cited by applicant

Van Den Oord et al., Pixel Recurrent Neural Networks. *Proceedings of the 33rd International Conference on Machine Learning. Journal of Machine Learning Research*. Aug. 19, 2016. vol. 48. 11 pages. cited by applicant

Vidal et al., Yeast forward and reverse 'n'-hybrid systems. *Nucleic Acids Res*. Feb. 15, 1999;27(4):919-29. doi: 10.1093/nar/27.4.919. cited by applicant

Vigneron et al., Database of T cell-defined human tumor antigens: the 2013 update. *Cancer Immun*. Jul. 15, 2013;13:15. 6 pages. cited by applicant

Webb et al., Functional and structural characteristics of NY-ESO-1-related HLA A2-restricted epitopes and the design of a novel immunogenic analogue. *J Biol Chem*. May 28, 2004;279(22):23438-46. doi: 10.1074/jbc.M314066200. Epub Mar. 5, 2004. cited by applicant

Wirth et al., Mildly affected patients with spinal muscular atrophy are partially protected by an increased SMN2 copy number. *Hum Genet*. May 2006;119(4):422-8. doi: 10.1007/s00439-006-0156-7. Epub Mar. 1, 2006. cited by applicant

Woo et al., Gene activation of SMN by selective disruption of lncRNA-mediated recruitment of PRC2 for the treatment of spinal muscular atrophy. *Proc Natl Acad Sci U S A*. Feb. 21, 2017;114(8):E1509-E1518. doi:10.1073/pnas.1616521114. Epub Feb. 13, 2017. cited by applicant

Wu et al., A novel SCN9A mutation responsible for primary erythromelalgia and is resistant to the treatment of sodium channel blockers. *PLoS One*. 2013;8(1):e55212. doi: 10.1371/journal.pone.0055212. Epub Jan. 31, 2013. 15 pages. cited by applicant

Yamane et al., Deep-sequencing identification of the genomic targets of the cytidine deaminase AID and its cofactor RPA in B lymphocytes. *Nat Immunol*. Jan. 2011;12(1):62-9. doi: 10.1038/ni.1964. Epub Nov. 28, 2010. cited by applicant

Yang et al., BRCA2 function in DNA binding and recombination from a BRCA2-DSS1-ssDNA structure. *Science*. Sep. 13, 2002;297(5588):1837-48. doi: 10.1126/science.297.5588.1837. cited by applicant

Yang et al., The BRCA2 homologue Brh2 nucleates RAD51 filament formation at a dsDNA-ssDNA junction. *Nature*. Feb. 10, 2005;433(7026):653-7. doi: 10.1038/nature03234. cited by applicant

Yu et al., Dynamic control of Rad51 recombinase by self-association and interaction with BRCA2. *Mol Cell*. Oct. 2003;12(4):1029-41. doi: 10.1016/s1097-2765(03)00394-0. cited by applicant

Zaremba et al., Identification of an enhancer agonist cytotoxic T lymphocyte peptide from human carcinoembryonic antigen. *Cancer Res*. Oct. 15, 1997;57(20):4570-7. cited by applicant

Zhang et al., Efficient precise knockin with a double cut HDR donor after CRISPR/Cas9-mediated double-stranded DNA cleavage. *Genome Biol*. Feb. 20, 2017;18(1):35. doi: 10.1186/s13059-017-1164-8. cited by applicant

Zhang et al., Large genomic fragment deletions and insertions in mouse using CRISPR/Cas9. *PLoS One*. Mar. 24, 2015;10(3):e0120396. doi: 10.1371/journal.pone.0120396. 14 pages. cited by applicant

Zhu et al., Novel Thrombotic Function of a Human SNP in STXBP5 Revealed by CRISPR/Cas9 Gene Editing in Mice. *Arterioscler Thromb Vasc Biol*. Feb. 2017;37(2):264-270. doi: 10.1161/ATVBAHA.116.308614. Epub Dec. 29, 2016. cited by applicant

[No Author Listed], MutL homolog 1. UniProtKB Acc. No. F1MPG0. May 3, 2011. Accessible at <https://rest.uniprot.org/unisave/F1MPG0?format=txt&versions=1>. 1 page. cited by applicant

Acharya et al., hMSH2 forms specific mispair-binding complexes with hMSH3 and hMSH6. *Proc Natl Acad Sci U S A*. Nov. 26, 1996;93(24):13629-34. doi: 10.1073/pnas.93.24.13629. cited by applicant

Basila et al., Minimal 2'-O-methyl phosphorothioate linkage modification pattern of synthetic guide RNAs for increased stability and efficient CRISPR-Cas9 gene editing avoiding cellular toxicity. *PLoS One*. Nov. 27, 2017;12(11):e0188593. doi: 10.1371/journal.pone.0188593. cited by applicant

Bertsimas et al., Simulated annealing. *Statistical Science*. Feb. 1993;8(1):10-15. doi: 10.1214/ss/1177011077. cited by applicant

Chen et al., Dynamic imaging of genomic loci in living human cells by an optimized CRISPR/Cas system. *Cell*. Dec. 19, 2013;155(7):1479-91. doi: 10.1016/j.cell.2013.12.001. Erratum in: *Cell*. Jan. 16, 2014;156(1-2):373. cited by applicant

Edraki et al., A Compact, High-Accuracy Cas9 with a Dinucleotide PAM for In Vivo Genome Editing. *Mol Cell*. Feb. 21, 2019;73(4):714-726.e4 and Supplemental Info. doi: 10.1016/j.molcel.2018.12.003. Epub Dec. 20, 2018. cited by applicant

Fang et al., Human strand-specific mismatch repair occurs by a bidirectional mechanism similar to that of the bacterial reaction. *J Biol Chem*. Jun. 5, 1993;268(16):11838-44. cited by applicant

Fang et al., The Menu of Features that Define Primary MicroRNAs and Enable De Novo Design of MicroRNA Genes. *Mol Cell*. Oct. 1, 2015;60(1):131-45. doi: 10.1016/j.molcel.2015.08.015. Epub Sep. 24, 2015. cited by applicant

Feng et al., Efficient genome editing in plants using a CRISPR/Cas system. *Cell Res*. Oct. 2013;23(10):1229-32. doi: 10.1038/cr.2013.114. Epub Aug. 20, 2013. cited by applicant

Fishel et al., The human mutator gene homolog MSH2 and its association with hereditary nonpolyposis colon cancer. *Cell*. Dec. 3, 1993;75(5):1027-38. doi: 10.1016/0092-8674(93)90546-3. Erratum in: *Cell*. Apr. 8, 1994;77(1):1 p following 166. cited by applicant

Fu et al., Targeted genome editing in human cells using CRISPR/Cas nucleases and truncated guide RNAs. *Methods Enzymol*. 2014;546:21-45. doi: 10.1016/B978-0-12-801185-0.00002-7. cited by applicant

Geisberg et al., Global analysis of mRNA isoform half-lives reveals stabilizing and destabilizing elements in yeast. *Cell*. Feb. 13, 2014;156(4):812-24. doi: 10.1016/j.cell.2013.12.026. cited by applicant

Geng et al., In vitro studies of DNA mismatch repair proteins. *Anal Biochem*. Jun. 15, 2011;413(2):179-84. doi: 10.1016/j.ab.2011.02.017. Epub Feb. 15, 2011. cited by applicant

Genschel et al., Human exonuclease I is required for 5' and 3' mismatch repair. *J Biol Chem*. Apr. 12, 2002;277(15):13302-11. doi: 10.1074/jbc.M111854200. Epub Jan. 24, 2002. cited by applicant

Genschel et al., Isolation of MutSbeta from human cells and comparison of the mismatch repair specificities of MutSbeta and MutSalpha. *J Biol Chem*. Jul. 31, 1998;273(31):19895-901. doi: 10.1074/jbc.273.31.19895. Erratum in: *J Biol Chem* Oct. 9, 1998;273(41):27034. cited by applicant

Green et al., Characterization of the mechanical unfolding of RNA pseudoknots. *J Mol Biol*. Jan. 11, 2008;375(2):511-28. doi: 10.1016/j.jmb.2007.05.058. Epub May 26, 2007. cited by applicant

Gueneau et al., Structure of the MutLalpha C-terminal domain reveals how Mlh1 contributes to Pms1 endonuclease site. *Nat Struct Mol Biol*. Apr. 2013;20(4):461-8. doi: 10.1038/nsmb.2511. Epub Feb. 24, 2013. cited by applicant

Guerrette et al., The interaction of the human MutL homologues in hereditary nonpolyposis colon cancer. *J Biol Chem*. Mar. 5, 1999;274(10):6336-41. doi: 10.1074/jbc.274.10.6336. cited by applicant

Gupta et al., Mechanism of mismatch recognition revealed by human MutSbeta bound to unpaired DNA loops. *Nat Struct Mol Biol*. Dec. 18, 2011;19(1):72-8. doi: 10.1038/nsmb.2175. cited by applicant

Hänsel-Hertsch et al., DNA G-quadruplexes in the human genome: detection, functions and therapeutic potential. *Nat Rev Mol Cell Biol*. May 2017;18(5):279-284. doi: 10.1038/nrm.2017.3. Epub Feb. 22, 2017. cited by applicant

Houck-Loomis et al., An equilibrium-dependent retroviral mRNA switch regulates translational recoding. *Nature*. Nov. 27, 2011;480(7378):561-4. doi: 10.1038/nature 10657. cited by applicant

Houseley et al., The many pathways of RNA degradation. *Cell*. Feb. 20, 2009;136(4):763-76. doi: 10.1016/j.cell.2009.01.019. cited by applicant

Iaccarino et al., hMSH2 and hMSH6 play distinct roles in mismatch binding and contribute differently to the ATPase activity of hMutSalpha. *EMBO J*. May 1, 1998;17(9):2677-86. doi: 10.1093/emboj/17.9.2677. cited by applicant

Ibrahim et al., RNA recognition by 3'-to-5' exonucleases: the substrate perspective. *Biochim Biophys Acta*. Apr. 2008;1779(4):256-65. doi: 10.1016/j.bbagr.2007.11.004. Epub Dec. 3, 2007. cited by applicant

Iyer et al., DNA mismatch repair: functions and mechanisms. *Chem Rev*. Feb. 2006;106(2):302-23. doi: 10.1021/cr0404794. cited by applicant

Jost et al., Titrating gene expression using libraries of systematically attenuated CRISPR guide RNAs. *Nat Biotechnol*. Mar. 2020;38(3):355-364. doi: 10.1038/s41587-019-0387-5. Epub Jan. 13, 2020. cited by applicant

Kadyrov et al., Endonucleolytic function of MutLalpha in human mismatch repair. *Cell*. Jul. 28, 2006;126(2):297-308. doi: 10.1016/j.cell.2006.05.039. cited by applicant

Ku et al., Nucleic Acid Aptamers: An Emerging Tool for Biotechnology and Biomedical Sensing. *Sensors (Basel)*. Jul. 6, 2015;15(7):16281-313. doi: 10.3390/s150716281. cited by applicant

Kunkel et al., DNA mismatch repair. *Annu Rev Biochem*. 2005;74:681-710. doi: 10.1146/annurev.biochem.74.082803.133243. cited by applicant

Kwok et al., G-Quadruplexes: Prediction, Characterization, and Biological Application. *Trends Biotechnol*. Oct. 2017;35(10):997-1013. doi: 10.1016/j.tibtech.2017.06.012. Epub Jul. 26, 2017. cited by applicant

Lahue et al., DNA mismatch correction in a defined system. *Science*. Jul. 14, 1989;245(4914):160-4. doi: 10.1126/science.2665076. cited by applicant

Leach et al., Mutations of a mutS homolog in hereditary nonpolyposis colorectal cancer. *Cell*. Dec. 17, 1993;75(6):1215-25. doi: 10.1016/0092-8674(93)90330-s. cited by applicant

Longworth, Expanding the Enzymatic Activity of the Programmable Endonuclease Cas9 in Zebrafish. Thesis. Rice

University. Houston, TX. May 17, 2019. 41 pages. cited by applicant

Lujan et al., Heterogeneous polymerase fidelity and mismatch repair bias genome variation and composition. *Genome Res.* Nov. 2014;24(11):1751-64. doi: 10.1101/gr.178335.114. Epub Sep. 12, 2014. cited by applicant

MacFadden et al., Mechanism and structural diversity of exoribonuclease-resistant RNA structures in flaviviral RNAs. *Nat Commun.* Jan. 9, 2018;9(1):119. doi: 10.1038/s41467-017-02604-y. cited by applicant

Mangeot et al., Genome editing in primary cells and in vivo using viral-derived Nanoblades loaded with Cas9-sgRNA ribonucleoproteins. *Nat Commun.* Jan. 3, 2019;10(1):45. doi: 10.1038/s41467-018-07845-z. cited by applicant

Micozzi et al., Human cytidine deaminase: a biochemical characterization of its naturally occurring variants. *Int J Biol Macromol.* Feb. 2014;63:64-74. doi: 10.1016/j.ijbiomac.2013.10.029. Epub Oct. 29, 2013. Erratum in: *Int J Biol Macromol.* Feb. 2014;63:262. cited by applicant

Millevoi et al., G-quadruplexes in RNA biology. *Wiley Interdiscip Rev RNA.* Jul.-Aug. 2012;3(4):495-507. doi: 10.1002/wrna.1113. Epub Apr. 4, 2012. cited by applicant

Pandey et al., Effect of loops and G-quartets on the stability of RNA G-quadruplexes. *J Phys Chem B.* Jun. 13, 2013;117(23):6896-905. doi: 10.1021/jp401739m. Epub May 29, 2013. Supplementary Information, 21 pages. cited by applicant

Parsons et al., Hypermutability and mismatch repair deficiency in RER+ tumor cells. *Cell.* Dec. 17, 1993;75(6):1227-36. doi: 10.1016/0092-8674(93)90331-j. cited by applicant

Petit et al., Powerful mutators lurking in the genome. *Philos Trans R Soc Lond B Biol Sci.* Mar. 12, 2009;364(1517):705-15. doi: 10.1098/rstb.2008.0272. cited by applicant

Pijlman et al., A highly structured, nuclease-resistant, noncoding RNA produced by flaviviruses is required for pathogenicity. *Cell Host Microbe.* Dec. 11, 2008;4(6):579-91. doi: 10.1016/j.chom.2008.10.007. cited by applicant

Piotukh et al., Directed evolution of sortase A mutants with altered substrate selectivity profiles. *J Am Chem Soc.* Nov. 9, 2011;133(44):17536-9. doi: 10.1021/ja205630g. Epub Oct. 13, 2011. cited by applicant

Plotz et al., N-terminus of hMLH1 confers interaction of hMutLalpha and hMutLbeta with hMutSalpha. *Nucleic Acids Res.* Jun. 15, 2003;31(12):3217-26. doi: 10.1093/nar/gkg420. cited by applicant

Räschle et al., Mutations within the hMLH1 and hPMS2 subunits of the human MutLalpha mismatch repair factor affect its ATPase activity, but not its ability to interact with hMutSalpha. *J Biol Chem.* Jun. 14, 2002;277(24):21810-20. doi: 10.1074/jbc.M108787200. Epub Apr. 10, 2002. cited by applicant

Robert et al., Virus-Like Particles Derived from HIV-1 for Delivery of Nuclear Proteins: Improvement of Production and Activity by Protein Engineering. *Mol Biotechnol.* Jan. 2017;59(1):9-23. doi: 10.1007/s12033-016-9987-1. cited by applicant

Shcherbakova et al., Mutator phenotypes conferred by MLH1 overexpression and by heterozygosity for mlh1 mutations. *Mol Cell Biol.* Apr. 1999;19(4):3177-83. doi: 10.1128/MCB.19.4.3177. cited by applicant

Steckelberg et al., A folded viral noncoding RNA blocks host cell exoribonucleases through a conformationally dynamic RNA structure. *Proc Natl Acad Sci U S A.* Jun. 19, 2018;115(25):6404-6409. doi: 10.1073/pnas.1802429115. Epub Jun. 4, 2018. cited by applicant

Strand et al., Destabilization of tracts of simple repetitive DNA in yeast by mutations affecting DNA mismatch repair. *Nature.* Sep. 16, 1993;365(6443):274-6. doi: 10.1038/365274a0. Erratum in: *Nature* Apr. 7, 1994;368(6471):569. cited by applicant

Su et al., Mispair specificity of methyl-directed DNA mismatch correction in vitro. *J Biol Chem.* May 15, 1988;263(14):6829-35. Erratum in: *J Biol Chem* Aug. 5, 1988;263(22):11015. cited by applicant

Sugawara et al., Heteroduplex rejection during single-strand annealing requires Sgs1 helicase and mismatch repair proteins Msh2 and Msh6 but not Pms1. *Proc Natl Acad Sci U S A.* Jun. 22, 2004;101(25):9315-20. doi: 10.1073/pnas.0305749101. Epub Jun. 15, 2004. cited by applicant

Supek et al., Differential DNA mismatch repair underlies mutation rate variation across the human genome. *Nature.* May 7, 2015;521(7550):81-4. doi: 10.1038/nature14173. Epub Feb. 23, 2015. cited by applicant

Svitashev et al., Targeted Mutagenesis, Precise Gene Editing, and Site-Specific Gene Insertion in Maize Using Cas9 and Guide RNA. *Plant Physiol.* Oct. 2015;169(2):931-45. doi: 10.1104/pp15.00793. Epub Aug. 12, 2015. cited by applicant

Thomas et al., Heteroduplex repair in extracts of human HeLa cells. *J Biol Chem.* Feb. 25, 1991;266(6):3744-51. cited by applicant

Tomer et al., Contribution of human mlh1 and pms2 ATPase activities to DNA mismatch repair. *J Biol Chem.* Jun. 14, 2002;277(24):21801-9. doi: 10.1074/jbc.M111342200. Epub Mar. 15, 2002. cited by applicant

Tran et al., Hypermutability of homonucleotide runs in mismatch repair and DNA polymerase proofreading yeast mutants. *Mol Cell Biol.* May 1997;17(5):2859-65. doi: 10.1128/MCB.17.5.2859. cited by applicant

Umar et al., DNA loop repair by human cell extracts. *Science.* Nov. 4, 1994;266(5186):814-6. doi: 10.1126/science.7973637. cited by applicant

Warren et al., Structure of the human MutSalpha DNA lesion recognition complex. *Mol Cell.* May 25, 2007;26(4):579-92. doi: 10.1016/j.molcel.2007.04.018. cited by applicant

Wu et al., MLV based viral-like-particles for delivery of toxic proteins and nuclear transcription factors. *Biomaterials.* Sep. 2014;35(29):8416-26. doi: 10.1016/j.biomaterials.2014.06.006. Epub Jul. 3, 2014. cited by applicant

Wu et al., Widespread Influence of 3'-End Structures on Mammalian mRNA Processing and Stability. *Cell.* May 18,



2017;169(5):905-917.e11. doi: 10.1016/j.cell.2017.04.036. cited by applicant

Xi et al., C-terminal Loop Mutations Determine Folding and Secretion Properties of PCSK9. *Biochem Mol Biol J*. 2016;2(3):17. doi: 10.21767/2471-8084.100026. 12 pages. cited by applicant

Yi et al., Engineering of TEV protease variants by yeast ER sequestration screening (YESS) of combinatorial libraries. *Proc Natl Acad Sci U S A*. Apr. 30, 2013;110(18):7229-34. doi: 10.1073/pnas.1215994110. Epub Apr. 15, 2013. cited by applicant

Zhang et al., Reconstitution of 5'-directed human mismatch repair in a purified system. *Cell*. Sep. 9, 2005;122(5):693-705. doi: 10.1016/j.cell.2005.06.027. cited by applicant

[No Author Listed] Ncbi Reference Sequence: WP\_032188360.1. Apr. 6, 2015. 1 page. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase [*Escherichia coli*]. GenBank Acc. No. CTS26096.1. Accessible at <https://www.ncbi.nlm.nih.gov/protein/CTS26096.1>. Aug. 22, 2015. 1 page. cited by applicant

[No Author Listed], tumor-specific antigen. Retrieved from <http://www.cancer.gov/publications/dictionaries/cancer-terms/def/tumor-specific-antigen>. Retrieved on Oct. 7, 2022. 1 page. cited by applicant

Alizadeh et al., HR9: An Important Cell Penetrating Peptide for Delivery of HCV NS3 DNA into HEK-293T Cells. *Avicenna J Med Biotechnol*. Jan.-Mar. 2020;12(1):44-51. cited by applicant

Avidan et al., Expression and characterization of a recombinant novel reverse transcriptase of a porcine endogenous retrovirus. *Virology*. Mar. 15, 2003;307(2):341-57. doi: 10.1016/s0042-6822(02)00131-9. cited by applicant

Bae et al., Heteroclitic CD33 peptide with enhanced anti-acute myeloid leukemic immunogenicity. *Clin Cancer Res*. Oct. 15, 2004;10(20):7043-52. doi: 10.1158/1078-0432.CCR-04-0322. cited by applicant

Baños-Sanz et al., Crystal structure and functional insights into uracil-DNA glycosylase inhibition by phage Φ29 DNA mimic protein p56. *Nucleic Acids Res*. Jul. 2013;41(13):6761-73. doi: 10.1093/nar/gkt395. Epub May 13, 2013. cited by applicant

Celluzzi et al., Peptide-pulsed dendritic cells induce antigen-specific CTL-mediated protective tumor immunity. *J Exp Med*. Jan. 1, 1996;183(1):283-7. doi: 10.1084/jem.183.1.283. cited by applicant

Cheriyian et al., Faster protein splicing with the Nostoc punctiforme DnaE intein using non-native extein residues. *J Biol Chem*. Mar. 1, 2013;288(9):6202-11. doi: 10.1074/jbc.M112.433094. Epub Jan. 10, 2013. cited by applicant

Choi et al., Optimization of AAV expression cassettes to improve packaging capacity and transgene expression in neurons. *Mol Brain*. Mar. 11, 2014;7:17. doi: 10.1186/1756-6606-7-17. cited by applicant

Damdindorj et al., A comparative analysis of constitutive promoters located in adeno-associated viral vectors. *PLoS One*. Aug. 29, 2014;9(8):e106472. doi: 10.1371/journal.pone.0106472. cited by applicant

Ding et al., Gene therapy for cardiovascular disease. *Journal of Shanghai University (Natural Science Edition)*. 2016;3:270-9. DOI: 10.3969/j.issn.1007-2861.2016.03.013. cited by applicant

Ekman et al., CRISPR-Cas9-Mediated Genome Editing Increases Lifespan and Improves Motor Deficits in a Huntington's Disease Mouse Model. *Mol Ther Nucleic Acids*. Sep. 6, 2019;17:829-839. doi: 10.1016/j.omtn.2019.07.009. Epub Jul. 26, 2019. cited by applicant

Eriksen et al., Occlusion of the Ribosome Binding Site Connects the Translational Initiation Frequency, mRNA Stability and Premature Transcription Termination. *Front Microbiol*. Mar. 14, 2017;8:362. doi: 10.3389/fmicb.2017.00362. cited by applicant

Fikes et al., Design of multi-epitope, analogue-based cancer vaccines. *Expert Opin Biol Ther*. Sep. 2003;3(6):985-93. doi: 10.1517/14712598.3.6.985. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_060228.2. Bi et al., Dec. 21, 2005. 1 page. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_062826.2. Bokar et al., Sep. 18, 2004. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. NP\_066012.1. Ota et al., Apr. 3, 2005. 2 pages. cited by applicant

GenBank Submission; NIH/NCBI, Accession No. WP\_042518169.1. No Author, Feb. 10, 2015. 1 page. cited by applicant

Hizi et al., Retroviral reverse transcriptases (other than those of HIV-1 and murine leukemia virus): a comparison of their molecular and biochemical properties. *Virus Res*. Jun. 2008;134(1-2):203-20. doi: 10.1016/j.virusres.2007.12.008. Epub Mar. 3, 2008. cited by applicant

Houghton et al., Immunological validation of the EpiOptimizer program for streamlined design of heteroclitic epitopes. *Vaccine*. Jul. 20, 2007;25(29):5330-42. doi: 10.1016/j.vaccine.2007.05.008. Epub Jun. 4, 2007. cited by applicant

Hwang et al., Heritable and precise zebrafish genome editing using a CRISPR-Cas system. *PLoS One*. Jul. 9, 2013;8(7):e68708. doi: 10.1371/journal.pone.0068708. cited by applicant

Kirshenboim et al., Expression and characterization of a novel reverse transcriptase of the LTR retrotransposon Tf1. *Virology*. Sep. 30, 2007;366(2):263-76. doi: 10.1016/j.virol.2007.04.002. Epub May 23, 2007. cited by applicant

Kwon et al., Precision targeting tumor cells using cancer-specific InDel mutations with CRISPR-Cas9. *Proc Natl Acad Sci U S A*. Mar. 1, 2022;119(9):e2103532119. doi: 10.1073/pnas.2103532119. cited by applicant

Men et al., Assessment of immunogenicity of human Melan-A peptide analogues in HLA-A\*0201/Kb transgenic mice. *J Immunol*. Mar. 15, 1999;162(6):3566-73. cited by applicant

Misra et al., An enzymatically active chimeric HIV-1 reverse transcriptase (RT) with the RNase-H domain of murine leukemia virus RT exists as a monomer. *J Biol Chem*. Apr. 17, 1998;273(16):9785-9. doi: 10.1074/jbc.273.16.9785. cited by applicant

Nicholson et al., Tuning T cell activation threshold and effector function with cross-reactive peptide ligands. *Int Immunol*.

Feb. 2000;12(2):205-13. doi: 10.1093/intimm/12.2.205. cited by applicant

Niemeyer, C.M., Semisynthetic DNA-protein conjugates for biosensing and nanofabrication. *Angew Chem Int Ed Engl.* Feb. 8, 2010;49(7):1200-16. doi: 10.1002/anie.200904930. cited by applicant

Nowak et al., Ty3 reverse transcriptase complexed with an RNA-DNA hybrid shows structural and functional asymmetry. *Nat Struct Mol Biol.* Apr. 2014;21(4):389-96. doi: 10.1038/nsmb.2785. Epub Mar. 9, 2014. Author Manuscript, 22 pages. cited by applicant

Pfeiffer et al., Altered peptide ligands can control CD4 T lymphocyte differentiation in vivo. *J Exp Med.* Apr. 1, 1995;181(4):1569-74. doi: 10.1084/jem.181.4.1569. cited by applicant

Raaijmakers et al., CRISPR/Cas Applications in Myotonic Dystrophy: Expanding Opportunities. *Int J Mol Sci.* Jul. 27, 2019;20(15):3689. doi: 10.3390/ijms20153689. cited by applicant

Riddle et al., Frameshift suppression: a nucleotide addition in the anticodon of a glycine transfer RNA. *Nat New Biol.* Apr. 25, 1973;242(121):230-4. doi: 10.1038/newbio242230a0. cited by applicant

Riddle et al., Frameshift suppressors. II. Genetic mapping and dominance studies. *J Mol Biol.* May 28, 1972;66(3):483-93. doi: 10.1016/0022-2836(72)90428-7. cited by applicant

Riddle et al., Suppressors of frameshift mutations in *Salmonella typhimurium*. *J Mol Biol.* Nov. 28, 1970;54(1):131-44. doi: 10.1016/0022-2836(70)90451-1. cited by applicant

Romagnani, Lymphokine production by human T cells in disease states. *Annu Rev Immunol.* 1994;12:227-57. doi: 10.1146/annurev.iy.12.040194.001303. cited by applicant

Sadowski et al., The sequence-structure relationship and protein function prediction. *Curr Opin Struct Biol.* Jun. 2009;19(3):357-62. doi: 10.1016/j.sbi.2009.03.008. Epub May 4, 2009. cited by applicant

Salazar et al., Agonist peptide from a cytotoxic t-lymphocyte epitope of human carcinoembryonic antigen stimulates production of tc1-type cytokines and increases tyrosine phosphorylation more efficiently than cognate peptide. *Int J Cancer.* Mar. 15, 2000;85(6):829-38. doi: 10.1002/(sici)1097-0215(20000315)85:6<829::aid-ijc16>3.0.co;2-k. cited by applicant

SCORE Results for US 2014-0186919 A1 to Zhang et al. Aug. 28, 2014. 3 pages. cited by applicant

Seffernick et al., Melamine deaminase and atrazine chlorohydrolase: 98 percent identical but functionally different. *J Bacteriol.* Apr. 2001;183(8):2405-10. doi: 10.1128/JB.183.8.2405-2410.2001. cited by applicant

Selby et al., Hepatitis C virus envelope glycoprotein E1 originates in the endoplasmic reticulum and requires cytoplasmic processing for presentation by class I MHC molecules. *J Immunol.* Jan. 15, 1999;162(2):669-76. cited by applicant

Simon et al., Retrons and their applications in genome engineering. *Nucleic Acids Res.* Dec. 2, 2019;47(21):11007-11019. doi: 10.1093/nar/gkz865. cited by applicant

Singh et al., Protein Engineering Approaches in the Post-Genomic Era. *Curr Protein Pept Sci.* 2018;19(1):5-15. doi: 10.2174/1389203718666161117114243. cited by applicant

Studebaker et al., Depletion of uracil-DNA glycosylase activity is associated with decreased cell proliferation. *Biochem Biophys Res Commun.* Aug. 26, 2005;334(2):509-15. doi: 10.1016/j.bbrc.2005.06.118. cited by applicant

Tang et al., Identification of Dehalobacter reductive dehalogenases that catalyse dechlorination of chloroform, 1,1,1-trichloroethane and 1,1-dichloroethane. *Philos Trans R Soc Lond B Biol Sci.* Mar. 11, 2013;368(1616):20120318. doi: 10.1098/rstb.2012.0318. cited by applicant

Tao et al., Induction of IL-4-producing CD4<sup>+</sup> T cells by antigenic peptides altered for TCR binding. *J Immunol.* May 1, 1997;158(9):4237-44. cited by applicant

Toro et al., Comprehensive phylogenetic analysis of bacterial reverse transcriptases. *PLoS One.* Nov. 25, 2014;9(11):e114083. doi: 10.1371/journal.pone.0114083. cited by applicant

Wang et al., The stimulation of low-affinity, nontolerized clones by heteroclitic antigen analogues causes the breaking of tolerance established to an immunodominant T cell epitope. *J Exp Med.* Oct. 4, 1999;190(7):983-94. doi: 10.1084/jem.190.7.983. cited by applicant

Witkowski et al., Conversion of a beta-ketoacyl synthase to a malonyl decarboxylase by replacement of the active-site cysteine with glutamine. *Biochemistry.* Sep. 7, 1999;38(36):11643-50. doi: 10.1021/bi990993h. cited by applicant

Yin et al., Optimizing genome editing strategy by primer-extension-mediated sequencing. *Cell Discov.* Mar. 26, 2019;5:18. doi: 10.1038/s41421-019-0088-8. cited by applicant

Zhang et al., Propagated Perturbations from a Peripheral Mutation Show Interactions Supporting WW Domain Thermostability. *Structure.* Nov. 6, 2018;26(11):1474-1485.e5. doi: 10.1016/j.str.2018.07.014. Epub Sep. 6, 2018. cited by applicant

Zitvogel et al., Therapy of murine tumors with tumor peptide-pulsed dendritic cells: dependence on T cells, B7 costimulation, and T helper cell 1-associated cytokines. *J Exp Med.* Jan. 1, 1996;183(1):87-97. doi: 10.1084/jem.183.1.87. cited by applicant

Zügel et al., Termination of peripheral tolerance to a T cell epitope by heteroclitic antigen analogues. *J Immunol.* Aug. 15, 1998;161(4):1705-9. cited by applicant

Abed et al., The Gag protein PEG10 binds to RNA and regulates trophoblast stem cell lineage specification. *PLoS One.* Apr. 5, 2019;14(4):e0214110. doi: 10.1371/journal.pone.0214110. cited by applicant

Abifadel et al., Mutations in PCSK9 cause autosomal dominant hypercholesterolemia. *Nat Genet.* Jun. 2003;34(2):154-6. doi: 10.1038/ng1161. cited by applicant

Adikusuma et al., Versatile single-step assembly of CRISPR/Cas9 vectors for dual gRNA expression. *PLoS One*. Dec. 6, 2017;12(12):e0187236. doi: 10.1371/journal.pone.0187236. cited by applicant

Addgene Plasmid #42234. pMJ920. 2013. Retrieved Jan. 22, 2025. 3 pages. cited by applicant

Ashley et al., Retrovirus-like Gag Protein Arc1 Binds RNA and Traffics across Synaptic Boutons. *Cell*. Jan. 11, 2018;172(1-2):262-274.e11. doi: 10.1016/j.cell.2017.12.022. cited by applicant

Ayala-Ramirez et al., A new autosomal recessive syndrome consisting of posterior microphthalmos, retinitis pigmentosa, foveoschisis, and optic disc drusen is caused by a MFRP gene mutation. *Mol Vis*. Dec. 4, 2006;12:1483-9. cited by applicant

Bender et al., Receptor-Targeted Nipah Virus Glycoproteins Improve Cell-Type Selective Gene Delivery and Reveal a Preference for Membrane-Proximal Cell Attachment. *PLoS Pathog*. Jun. 9, 2016;12(6):e1005641. doi: 10.1371/journal.ppat.1005641. cited by applicant

Bernardi et al., Nucleotide sequence at the binding site for coat protein on RNA of bacteriophage R17. *Proc Natl Acad Sci U S A*. Oct. 1972;69(10):3033-7. doi: 10.1073/pnas.69.10.3033. cited by applicant

Bikard et al., Programmable repression and activation of bacterial gene expression using an engineered CRISPR-Cas system. *Nucleic Acids Res*. Aug. 2013;41(15):7429-37. doi: 10.1093/nar/gkt520. Epub Jun. 12, 2013. cited by applicant

Bolukbasi et al., DNA-binding-domain fusions enhance the targeting range and precision of Cas9. *Nat Methods*. Dec. 2015;12(12):1150-6. doi: 10.1038/nmeth.3624. Epub Oct. 19, 2015. cited by applicant

Cai et al. Targeted genome editing by lentiviral protein transduction of zinc-finger and TAL-effector nucleases. *Elife*. Apr. 24, 2014;3:e01911. doi: 10.7554/eLife.01911. cited by applicant

Cai et al., Abstract OR021: Targeted Genome Editing by Lentiviral Protein Transduction of ZFN and Cas9 Proteins Abstract, Presented at Proceedings of the ESGCT and NVGCT Collaborative Congress: The Hague. *Human Gene Therapy*. 2014. 15 pages. cited by applicant

Cai, Protein Transduction Using Lentiviral Vectors for Transposition and Site-directed Gene Editing. Thesis for the degree of Doctor of Philosophy, Aarhus University, Department of Biomedicine. 2014. 74 pages. cited by applicant

Cameron et al., Mapping the genomic landscape of CRISPR-Cas9 cleavage. *Nat Methods*. Jun. 2017;14(6):600-606 with Erratum. doi: 10.1038/nmeth.4284. Epub May 1, 2017. Erratum in: *Nat Methods*. Dec. 2023;20(12):2068. doi: 10.1038/s41592-023-02114-4. 8 pages. cited by applicant

Chandler et al., Recombinant Adeno-Associated Viral Integration and Genotoxicity: Insights from Animal Models. *Hum Gene Ther*. Apr. 2017;28(4):314-322. doi: 10.1089/hum.2017.009. cited by applicant

Chang et al., Functional characterization of the placental fusogenic membrane protein syncytin. *Biol Reprod*. Dec. 2004;71(6):1956-62. doi: 10.1095/biolreprod.104.033340. Epub Jul. 21, 2004. cited by applicant

Chen et al., DNA methylation and demethylation in mammals. *J Biol Chem*. May 27, 2011;286(21):18347-53. doi: 10.1074/jbc.R110.205286. Epub Mar. 24, 2011. cited by applicant

Cho et al., Heritable gene knockout in *Caenorhabditis elegans* by direct injection of Cas9-sgRNA ribonucleoproteins. *Genetics*. Nov. 2013;195(3):1177-80. doi: 10.1534/genetics.113.155853. Epub Aug. 26, 2013. cited by applicant

Chylinski et al., The tracrRNA and Cas9 families of type II CRISPR-Cas immunity systems. *RNA Biol*. May 2013;10(5):Supplementary Material. doi: 10.4161/rna.24321. Epub Apr. 5, 2013. 12 pages. cited by applicant

Cideciyan, Leber congenital amaurosis due to RPE65 mutations and its treatment with gene therapy. *Prog Retin Eye Res*. Sep. 2010;29(5):398-427. doi: 10.1016/j.preteyeres.2010.04.002. Epub Apr. 24, 2010. cited by applicant

Coey, Sumoylation of thymine DNA glycosylase occurs efficiently and weakens DNA binding but does not regulate enzymatic turnover. Dissertation. 2017. 178 pages. cited by applicant

Cohen et al., Low LDL cholesterol in individuals of African descent resulting from frequent nonsense mutations in PCSK9. *Nat Genet*. Feb. 2005;37(2):161-5. doi: 10.1038/ng1509. Epub Jan. 16, 2005. Erratum in: *Nat Genet*. Mar. 2005;37(3):328. cited by applicant

Cohen et al., Sequence variations in PCSK9, low LDL, and protection against coronary heart disease. *N Engl J Med*. Mar. 23, 2006;354(12):1264-72. doi: 10.1056/NEJMoa054013. cited by applicant

Contreras-Galindo et al., Human Endogenous Retrovirus Type K (HERV-K) Particles Package and Transmit HERV-K-Related Sequences. *J Virol*. Jul. 2015;89(14):7187-201. doi: 10.1128/JVI.00544-15. Epub Apr. 29, 2015. cited by applicant

Coquin, Characterization of lentiviral vectors pseudotyped with murine syncytins and their cellular targets in vitro and in vivo. Thesis. University of Évry Val d'Essonne. Defense on Dec. 10, 2019. 238 pages. cited by applicant

Den Hollander et al., Leber congenital amaurosis: genes, proteins and disease mechanisms. *Prog Retin Eye Res*. Jul. 2008;27(4):391-419. doi: 10.1016/j.preteyeres.2008.05.003. Epub Jun. 1, 2008. cited by applicant

Farhy-Tselnicker et al., Astrocytes, neurons, synapses: a tripartite view on cortical circuit development. *Neural Dev*. May 1, 2018;13(1):7. doi: 10.1186/s13064-018-0104-y. cited by applicant

Fehér et al., Characterization of the murine leukemia virus protease and its comparison with the human immunodeficiency virus type 1 protease. *J Gen Virol*. May 2006;87(Pt 5):1321-1330. doi: 10.1099/vir.0.81382-0. cited by applicant

Fitzgerald et al., Effect of an RNA interference drug on the synthesis of proprotein convertase subtilisin/kexin type 9 (PCSK9) and the concentration of serum LDL cholesterol in healthy volunteers: a randomised, single-blind, placebo-controlled, phase 1 trial. *Lancet*. Jan. 4, 2014;383(9911):60-68. doi: 10.1016/S0140-6736(13)61914-5. Epub Oct. 3, 2013. cited by applicant

Fonfara et al., Phylogeny of Cas9 determines functional exchangeability of dual-RNA and Cas9 among orthologous type II CRISPR-Cas systems. *Nucleic Acids Res.* Feb. 2014;42(4):2577-90. doi: 10.1093/nar/gkt1074. Epub Nov. 22, 2013. Supplementary Information. 67 pages. cited by applicant

Gaidukov et al., A multi-landing pad DNA integration platform for mammalian cell engineering. *Nucleic Acids Res.* May 4, 2018;46(8):4072-4086. doi: 10.1093/nar/gky216. cited by applicant

Gao et al., Delineation of the Exact Transcription Termination Signal for Type 3 Polymerase III. *Mol Ther Nucleic Acids.* Mar. 2, 2018;10:36-44. doi: 10.1016/j.omtn.2017.11.006. Epub Nov. 21, 2017. cited by applicant

Garnier et al., WW domains and retrovirus budding. *Nature.* Jun. 27, 1996;381(6585):744-5. doi: 10.1038/381744a0. cited by applicant

GenBank Accession No. AAH57574.1 2009. 2 pages. cited by applicant

Giannoukos et al., UDiTaS™, a genome editing detection method for indels and genome rearrangements. *BMC Genomics.* Mar. 21, 2018;19(1):212. doi: 10.1186/s12864-018-4561-9. cited by applicant

Golczak et al., Importance of membrane structural integrity for RPE65 retinoid isomerization activity. *J Biol Chem.* Mar. 26, 2010;285(13):9667-9682. doi: 10.1074/jbc.M109.063941. Epub Jan. 25, 2010. cited by applicant

Gusel'Nikova et al., NeuN As a Neuronal Nuclear Antigen and Neuron Differentiation Marker. *Acta Naturae.* Apr.-Jun. 2015;7(2):42-7. cited by applicant

Hamilton et al., Knocking out barriers to engineered cell activity. *Science.* Feb. 28, 2020;367(6481):976-977. doi: 10.1126/science.aba9844. Epub Feb. 6, 2020. cited by applicant

Heins et al., Designing Automated, High-throughput, Continuous Cell Growth Experiments Using e Volver. *J Vis Exp.* May 19, 2019;(147):10.3791/59652. doi: 10.3791/59652. cited by applicant

Heintze et al., A CRISPR CAsE for high-throughput silencing. *Front Genet.* Oct. 7, 2013;4:193. doi: 10.3389/fgene.2013.00193. cited by applicant

Hooper et al., The C679X mutation in PCSK9 is present and lowers blood cholesterol in a Southern African population. *Atherosclerosis.* Aug. 2007;193(2):445-8. doi: 10.1016/j.atherosclerosis.2006.08.039. Epub Sep. 20, 2006. cited by applicant

Jacobs et al., DNA glycosylases: in DNA repair and beyond. *Chromosoma.* Feb. 2012;121(1):1-20. doi: 10.1007/s00412-011-0347-4. Epub Nov. 3, 2011. 20 pages. cited by applicant

Jinek et al., A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science.* Aug. 17, 2012;337(6096):Supplementary Material. doi: 10.1126/science.1225829. Epub Jun. 28, 2012. 37 pages. cited by applicant

Johnson et al., Mass spectrometry analysis reveals differences in the host cell protein species found in pseudotyped lentiviral vectors. *Biologicals.* Mar. 2018;52:59-66. doi: 10.1016/j.biologicals.2017.12.005. Epub Feb. 1, 2018. cited by applicant

Johnson, Origins and evolutionary consequences of ancient endogenous retroviruses. *Nat Rev Microbiol.* Jun. 2019;17(6):355-370. doi: 10.1038/s41579-019-0189-2. cited by applicant

Kameya et al., Mfrp, a gene encoding a frizzled related protein, is mutated in the mouse retinal degeneration 6. *Hum Mol Genet.* Aug. 1, 2002;11(16):1879-86. doi: 10.1093/hmg/11.16.1879. cited by applicant

Katoh et al., Exploitation of the interaction of measles virus fusogenic envelope proteins with the surface receptor CD46 on human cells for microcell-mediated chromosome transfer. *BMC Biotechnol.* May 6, 2010;10:37. doi: 10.1186/1472-6750-10-37. cited by applicant

Kleinstiver et al., Broadening the targeting range of *Staphylococcus aureus* CRISPR-Cas9 by modifying PAM recognition. *Nat Biotechnol.* Dec. 2015;33(12):1293-1298. doi: 10.1038/nbt.3404. Epub Nov. 2, 2015. Author Manuscript, 14 pages. cited by applicant

Kneissl et al., Measles virus glycoprotein-based lentiviral targeting vectors that avoid neutralizing antibodies. *PLoS One.* 2012;7(10):e46667. doi: 10.1371/journal.pone.0046667. Epub Oct. 10, 2012. cited by applicant

Kulcsár et al., Blackjack mutations improve the on-target activities of increased fidelity variants of SpCas9 with 5'-G-extended sgRNAs. *Nat Commun.* Mar. 6, 2020;11(1):1223. doi: 10.1038/s41467-020-15021-5. cited by applicant

Lau et al., In vivo epigenome editing and transcriptional modulation using CRISPR technology. *Transgenic Res.* Dec. 2018;27(6):489-509. doi: 10.1007/s11248-018-0096-8. Epub Oct. 4, 2018. cited by applicant

Lebar et al., A tunable orthogonal coiled-coil interaction toolbox for engineering mammalian cells. *Nat Chem Biol.* May 2020;16(5):513-519. doi: 10.1038/s41589-019-0443-y. Epub Jan. 6, 2020. cited by applicant

Lee et al., Reconstitution of an infectious human endogenous retrovirus. *PLoS Pathog.* Jan. 2007;3(1):e10. doi: 10.1371/journal.ppat.0030010. cited by applicant

Leenay et al., Identifying and Visualizing Functional PAM Diversity across CRISPR-Cas Systems. *Mol Cell.* Apr. 7, 2016;62(1):137-47. doi: 10.1016/j.molcel.2016.02.031. Epub Mar. 31, 2016. cited by applicant

Leibundgut-Landmann et al., Mini-review: Specificity and expression of CIITA, the master regulator of MHC class II genes. *Eur J Immunol.* Jun. 2004;34(6):1513-25. doi: 10.1002/eji.200424964. cited by applicant

Li et al., A dominant-negative form of mouse SOX2 induces trophectoderm differentiation and progressive polyploidy in mouse embryonic stem cells. *J Biol Chem.* Jul. 6, 2007;282(27):19481-92. doi: 10.1074/jbc.M702056200. Epub May 15, 2007. cited by applicant

Li et al., Base-Resolution Mapping Reveals Distinct m1A Methylome in Nuclear- and Mitochondrial-Encoded Transcripts. *Mol Cell.* Dec. 7, 2017;68(5):993-1005.e9. doi: 10.1016/j.molcel.2017.10.019. Epub Nov. 5, 2017. cited by

applicant

- Lim et al., Specific insertions of zinc finger domains into Gag-Pol yield engineered retroviral vectors with selective integration properties. *Proc Natl Acad Sci U S A*. Jul. 13, 2010;107(28):12475-80. doi: 10.1073/pnas.1001402107. Epub Jun. 28, 2010. cited by applicant
- Liu et al., Delivery methods for site-specific nucleases: Achieving the full potential of therapeutic gene editing. *J Control Release*. Dec. 28, 2016;244(Pt A):83-97. doi: 10.1016/j.jconrel.2016.11.014. Epub Nov. 16, 2016. cited by applicant
- Ma et al., Pol III Promoters to Express Small RNAs: Delineation of Transcription Initiation. *Mol Ther Nucleic Acids*. May 6, 2014;3(5):e161. doi: 10.1038/mtna.2014.12. cited by applicant
- Martín et al., Envelope-targeted retrovirus vectors transduce melanoma xenografts but not spleen or liver. *Mol Ther*. Mar. 2002;5(3):269-74. doi: 10.1006/mthe.2002.0550. cited by applicant
- Mason et al., Coiled coil domains: stability, specificity, and biological implications. *Chembiochem*. Feb. 6, 2004;5(2):170-6. doi: 10.1002/cbic.200300781. cited by applicant
- Mercuri et al., Nusinersen versus Sham Control in Later-Onset Spinal Muscular Atrophy. *N Engl J Med*. Feb. 15, 2018;378(7):625-635. doi: 10.1056/NEJMoa1710504. cited by applicant
- Meunier et al., Drug-Induced Liver Injury: Biomarkers, Requirements, Candidates, and Validation. *Front Pharmacol*. Dec. 11, 2019;10:1482. doi: 10.3389/fphar.2019.01482. cited by applicant
- Nawaz et al., Extracellular Vesicles, Tunneling Nanotubes, and Cellular Interplay: Synergies and Missing Links. *Front Mol Biosci*. Jul. 18, 2017;4:50. doi: 10.3389/fmolb.2017.00050. cited by applicant
- Nesbitt, Targeted Intracellular Therapeutic Delivery Using Liposomes Formulated with Multifunctional FAST proteins. Electronic Thesis and Dissertation Repository. The University of Western Ontario. 2012. 126 pages. cited by applicant
- Pan et al., Biodistribution and toxicity studies of VSVG-pseudotyped lentiviral vector after intravenous administration in mice with the observation of in vivo transduction of bone marrow. *Mol Ther*. Jul. 2002;6(1):19-29. doi: 10.1006/mthe.2002.0630. cited by applicant
- Pan et al., Identification of a nuclear localization signal in OCT4 and generation of a dominant negative mutant by its ablation. *J Biol Chem*. Aug. 27, 2004;279(35):37013-20. doi: 10.1074/jbc.M405117200. Epub Jun. 24, 2004. cited by applicant
- Pang et al., Retinal degeneration 12 (rd12): a new, spontaneously arising mouse model for human Leber congenital amaurosis (LCA). *Mol Vis*. Feb. 28, 2005;11:152-62. cited by applicant
- Parr-Brownlie et al., Lentiviral vectors as tools to understand central nervous system biology in mammalian model organisms. *Front Mol Neurosci*. May 18, 2015;8:14. doi: 10.3389/fnmol.2015.00014. cited by applicant
- Pastuzyn et al., The Neuronal Gene Arc Encodes a Repurposed Retrotransposon Gag Protein that Mediates Intercellular RNA Transfer. *Cell*. Jan. 11, 2018;172(1-2):275-288.e18. doi: 10.1016/j.cell.2017.12.024. Erratum in: *Cell*. Mar. 22, 2018;173(1):275. doi: 10.1016/j.cell.2018.03.024. cited by applicant
- Pavlov et al., Roles of DNA polymerases in replication, repair, and recombination in eukaryotes. *Int Rev Cytol*. 2006;255:41-132. doi: 10.1016/S0074-7696(06)55002-8. cited by applicant
- Podbilewicz, Virus and cell fusion mechanisms. *Annu Rev Cell Dev Biol*. 2014;30:111-39. doi: 10.1146/annurev-cellbio-101512-122422. Epub Jun. 27, 2014. cited by applicant
- Puppo et al., Retinal transduction profiles by high-capacity viral vectors. *Gene Ther*. Oct. 2014;21(10):855-65. doi: 10.1038/gt.2014.57. Epub Jul. 3, 2014. cited by applicant
- Qi et al., Repurposing CRISPR as an RNA-guided platform for sequence-specific control of gene expression. *Cell*. Feb. 28, 2013;152(5):Supplementary Material. doi: 10.1016/j.cell.2013.02.022. 4 pages. cited by applicant
- Ramiro et al., Transcription enhances AID-mediated cytidine deamination by exposing single-stranded DNA on the nontemplate strand. *Nat Immunol*. May 2003;4(5):452-6. doi: 10.1038/ni920. cited by applicant
- Rao et al., Large-Scale Phenome-Wide Association Study of PCSK9 Variants Demonstrates Protection Against Ischemic Stroke. *Circ Genom Precis Med*. Jul. 2018;11(7):e002162. doi: 10.1161/CIRCGEN.118.002162. cited by applicant
- Remington et al., Complete nucleotide sequence of a neuropathogenic variant of Friend murine leukemia virus PVC-211. *Nucleic Acids Res*. Jun. 25, 1992;20(12):3249. doi: 10.1093/nar/20.12.3249. cited by applicant
- Romero et al., Exploring protein fitness landscapes by directed evolution. *Nat Rev Mol Cell Biol*. Dec. 2009;10(12):866-76. doi: 10.1038/nrm2805. cited by applicant
- Sanjana et al., Improved vectors and genome-wide libraries for CRISPR screening. *Nat Methods*. Aug. 2014;11(8):783-784. doi: 10.1038/nmeth.3047. cited by applicant
- Sapir et al., Viral and developmental cell fusion mechanisms: conservation and divergence. *Dev Cell*. Jan. 2008;14(1):11-21. doi: 10.1016/j.devcel.2007.12.008. cited by applicant
- Schellekens, Bioequivalence and the immunogenicity of biopharmaceuticals. *Nat Rev Drug Discov*. Jun. 2002;1(6):457-62. doi: 10.1038/nrd818. cited by applicant
- Schneider et al., MuLV IN mutants responsive to HDAC inhibitors enhance transcription from unintegrated retroviral DNA. *Virology*. May 10, 2012;426(2):188-96. doi: 10.1016/j.virol.2012.01.034. Epub Feb. 23, 2012. cited by applicant
- Semple et al., Brain development in rodents and humans: Identifying benchmarks of maturation and vulnerability to injury across species. *Prog Neurobiol*. Jul.-Aug. 2013;106-107:1-16. doi: 10.1016/j.pneurobio.2013.04.001. Epub Apr. 11, 2013. cited by applicant
- Serreze et al., Major histocompatibility complex class I-deficient NOD-B2mnull mice are diabetes and insulinitis resistant.

Diabetes. Mar. 1994;43(3):505-9. doi: 10.2337/diab.43.3.505. cited by applicant

Shen et al., Activation-induced cytidine deaminase (AID) can target both DNA strands when the DNA is supercoiled. *Proc Natl Acad Sci U S A*. Aug. 31, 2004;101(35):12997-3002. doi: 10.1073/pnas.0404974101. Epub Aug. 24, 2004. cited by applicant

Skipper et al., Delivering the Goods for Genome Engineering and Editing. *Hum Gene Ther*. Aug. 2015;26(8):486-97. doi: 10.1089/hum.2015.063. cited by applicant

Stadtmauer et al., CRISPR-engineered T cells in patients with refractory cancer. *Science*. Feb. 28, 2020;367(6481):eaba7365. doi: 10.1126/science.aba7365. Epub Feb. 6, 2020. cited by applicant

Stevens et al., Design of a Split Intein with Exceptional Protein Splicing Activity. *J Am Chem Soc*. Feb. 24, 2016;138(7):2162-5. doi: 10.1021/jacs.5b13528. Epub Feb. 8, 2016. Abstract Only. 1 page. cited by applicant

Swiech et al., In vivo interrogation of gene function in the mammalian brain using CRISPR-Cas9. *Nat Biotechnol*. Jan. 2015;33(1):102-6. doi: 10.1038/nbt.3055. Epub Oct. 19, 2014. Author Manuscript. 22 pages. cited by applicant

Taylor, Ocular immune privilege. *Eye (Lond)*. Oct. 2009;23(10):1885-9. doi: 10.1038/eye.2008.382. Epub Jan. 9, 2009. cited by applicant

Thakore et al., Highly specific epigenome editing by CRISPR-Cas9 repressors for silencing of distal regulatory elements. *Nat Methods*. Dec. 2015;12(12):1143-9. doi: 10.1038/nmeth.3630. Epub Oct. 26, 2015. cited by applicant

Thorne et al., In vivo diffusion analysis with quantum dots and dextrans predicts the width of brain extracellular space. *Proc Natl Acad Sci U S A*. Apr. 4, 2006;103(14):5567-72. doi: 10.1073/pnas.0509425103. Epub Mar. 27, 2006. cited by applicant

Tokuriki et al., Stability effects of mutations and protein evolvability. *Curr Opin Struct Biol*. Oct. 2009;19(5):596-604. doi: 10.1016/j.sbi.2009.08.003. Epub Sep. 16, 2009. cited by applicant

Tözsér, Comparative studies on retroviral proteases: substrate specificity. *Viruses*. Jan. 2010;2(1):147-165. doi: 10.3390/v2010147. Epub Jan. 14, 2010. cited by applicant

Urano et al., Substitution of the myristoylation signal of human immunodeficiency virus type 1 Pr55Gag with the phospholipase C-delta1 pleckstrin homology domain results in infectious pseudovirion production. *J Gen Virol*. Dec. 2008;89(Pt 12):3144-3149. doi: 10.1099/vir.0.2008/004820-0. cited by applicant

Voisset et al., Phylogeny of a novel family of human endogenous retrovirus sequences, HERV-W, in humans and other primates. *AIDS Res Hum Retroviruses*. Nov. 20, 1999;15(17):1529-33. doi: 10.1089/088922299309810. cited by applicant

Wang et al., Characterization of an Mps I-H knock-in mouse that carries a nonsense mutation analogous to the human IDUA-W402X mutation. *Mol Genet Metab*. Jan. 2010;99(1):62-71. doi: 10.1016/j.ymgme.2009.08.002. Erratum in: *Mol Genet Metab*. Apr. 2010;99(4):439. cited by applicant

Wang et al., Influence of the polyanion on the physico-chemical properties and biological activities of polyanion/DNA/polycation ternary polyplexes. *Acta Biomater*. Aug. 2012;8(8):3014-26. doi: 10.1016/j.actbio.2012.04.034. Epub Apr. 27, 2012. cited by applicant

Webber et al., Highly efficient multiplex human T cell engineering without double-strand breaks using Cas9 base editors. *Nat Commun*. Nov. 19, 2019;10(1):5222. doi: 10.1038/s41467-019-13007-6. Erratum in: *Nat Commun*. Dec. 6, 2019;10(1):5659. doi: 10.1038/s41467-019-13778-y. cited by applicant

Wheeler et al., Proteomics analysis of cellular components in lentiviral vector production using Gel-LC-MS/MS. *Proteomics Clin Appl*. Feb. 2007;1(2):224-30. doi: 10.1002/prca.200600522. Epub Jan. 22, 2007. cited by applicant

Wu et al., Effect of genome size on AAV vector packaging. *Mol Ther*. Jan. 2010;18(1):80-6. doi: 10.1038/mt.2009.255. Epub Nov. 10, 2009. cited by applicant

Xu et al., Cas9-based tools for targeted genome editing and transcriptional control. *Appl Environ Microbiol*. Mar. 2014;80(5):1544-52. doi: 10.1128/AEM.03786-13. Epub Jan. 3, 2014. cited by applicant

Xu et al., Sequence and structural analyses of nuclear export signals in the NESdb database. *Mol Biol Cell*. Sep. 2012;23(18):3677-93. doi: 10.1091/mbc.E12-01-0046. Epub Jul. 25, 2012. cited by applicant

Zhang et al., Morphology and ultrastructure of retrovirus particles. *AIMS Biophys*. 2015;2(3):343-369. doi: 10.3934/biophy.2015.3.343. Epub Aug. 18, 2015. Author Manuscript. 33 pages. cited by applicant

Zhong et al., Seven novel variants expand the spectrum of RPE65-related Leber congenital amaurosis in the Chinese population. *Mol Vis*. Mar. 18, 2019;25:204-214. cited by applicant

[No Author Listed], CMP/dCMP-type deaminase domain-containing protein. Uniprot Accession No. A0A2Z6RZE9. Oct. 10, 2018. Accessible at <https://www.uniprot.org/uniprotkb/A0A2Z6RZE9/entry>. 8 pages. cited by applicant

[No Author Listed], dCas9-5xPlat2AfID-P2A-scFvGCN4sfGFPTET1CD [Cloning vector pPlatTET-grNA2]. GenBank No. BAV54124. Apr. 18, 2017. 5 pages. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase [Candidatus Moranella endobia PCVAL]. GenBank Acc. No. AGJ61179.1. Accessible at <https://www.ncbi.nlm.nih.gov/protein/AGJ61179>. Jan. 3, 20140. 3 pages. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase 2 [Terrapene triunguis]. GenBank Acc. No. XP\_024075810.1. Accessible at [https://www.ncbi.nlm.nih.gov/protein/XP\\_024075810](https://www.ncbi.nlm.nih.gov/protein/XP_024075810). Jul. 15, 2019. 2 pages. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase TAD2 isoform X2 [Panicum hallii]. GenBank Acc. No. XP\_025793740.1. Accessible at <https://www.ncbi.nlm.nih.gov/protein/025793740>. Jul. 27, 2018. 1 page. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase TAD2 isoform X1 [*Oryza sativa* Japonica Group]. GenBank Acc. No. XP\_15631651.1. Accessible at <https://www.ncbi.nlm.nih.gov/protein/1002254769?sat=58&satkey=133677684>. Aug. 7, 2018. 2 pages. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase. Uniprot Accession No. A0A1U7M801. May 10, 2017. Accessible at <https://www.uniprot.org/uniprotkb/A0A1U7M801/history>. 3 pages. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase. Uniprot Accession No. A0A1Z4VPW4. Sep. 27, 2017. Accessible at <https://www.uniprot.org/uniprotkb/A0A1Z4VPW4/history>. 3 pages. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase. Uniprot Accession No. A0A1Z9LYI9. Oct. 25, 2017. Accessible at <https://www.uniprot.org/uniprotkb/A0A1Z9LYI9/entry>. 12 pages. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase. Uniprot Accession No. A0A2P5TOZ9. May 23, 2018. Accessible at <https://www.uniprot.org/uniprotkb/A0A2P5TOZ9/entry>. 10 pages. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase. Uniprot Accession No. A0A4P6PH16. Jul. 31, 2019. Accessible at <https://www.uniprot.org/uniprotkb/A0A4P6PH16/entry>. 12 pages. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase. Uniprot Accession No. A0A520SVM3. Oct. 16, 2019. Accessible at <https://www.uniprot.org/uniprotkb/A0A520SVM3/entry>. 10 pages. cited by applicant

[No Author Listed], tRNA-specific adenosine deaminase. Uniprot Accession No. U2JUU0. Nov. 13, 2013. Accessible at <https://www.uniprot.org/uniprotkb/U2JUU0/entry>. 11 pages. cited by applicant

Alves et al., Immunogenicity of the carcinoembryonic antigen derived peptide 694 in HLA-A2 healthy donors and colorectal carcinoma patients. *Cancer Immunol Immunother.* Nov. 2007;56(11):1795-805. doi: 10.1007/s00262-007-0323-2. Epub Apr. 20, 2007. cited by applicant

Asemissen et al., Identification of a highly immunogenic HLA-A\*01-binding T cell epitope of WT1. *Clin Cancer Res.* Dec. 15, 2006;12(24):7476-82. doi: 10.1158/1078-0432.CCR-06-1337. cited by applicant

Attia et al., Autoimmunity correlates with tumor regression in patients with metastatic melanoma treated with anti-cytotoxic T-lymphocyte antigen-4. *J Clin Oncol.* Sep. 1, 2005;23(25):6043-53. doi: 10.1200/JCO.2005.06.205. Epub Aug. 8, 2005. cited by applicant

Aurischio et al., A novel minigene scaffold for therapeutic cancer vaccines. *Oncoimmunology.* Jan. 1, 2014;3(1):e27529. doi: 10.4161/onci.27529. Epub Jan. 16, 2014. cited by applicant

Bae et al., Identification of novel CD33 antigen-specific peptides for the generation of cytotoxic T lymphocytes against acute myeloid leukemia. *Cell Immunol.* Jan. 2004;227(1):38-50. doi: 10.1016/j.cellimm.2004.01.002. cited by applicant

Bakker et al., Analogues of CTL epitopes with improved MHC class-I binding capacity elicit anti-melanoma CTL recognizing the wild-type epitope. *Int J Cancer.* Jan. 27, 1997;70(3):302-9. doi: 10.1002/(sici)1097-0215(19970127)70:3<302::aid-ijc10>3.0.co;2-h. cited by applicant

Banerjee et al., Viral glycoproteins: biological role and application in diagnosis. *Virusdisease.* Mar. 2016;27(1):1-11. doi: 10.1007/s13337-015-0293-5. Epub Jan. 18, 2016. cited by applicant

Barve et al., Induction of immune responses and clinical efficacy in a phase II trial of IDM-2101, a 10-epitope cytotoxic T-lymphocyte vaccine, in metastatic non-small-cell lung cancer. *J Clin Oncol.* Sep. 20, 2008;26(27):4418-25. doi: 10.1200/JCO.2008.16.6462. cited by applicant

Benlalam et al., Identification of five new HLA-B\*3501-restricted epitopes derived from common melanoma-associated antigens, spontaneously recognized by tumor-infiltrating lymphocytes. *J Immunol.* Dec. 1, 2003;171(11):6283-9. doi: 10.4049/jimmunol.171.11.6283. cited by applicant

Bernatchez et al., Altered decamer and nonamer from an HLA-A0201-restricted epitope of Survivin differentially stimulate T-cell responses in different individuals. *Vaccine.* Apr. 5, 2011;29(16):3021-30. doi: 10.1016/j.vaccine.2011.01.115. Epub Feb. 12, 2011. cited by applicant

Bioley et al., Melan-A/MART-1-specific CD4 T cells in melanoma patients: identification of new epitopes and ex vivo visualization of specific T cells by MHC class II tetramers. *J Immunol.* Nov. 15, 2006;177(10):6769-79. doi: 10.4049/jimmunol.177.10.6769. cited by applicant

Blanchet et al., A new generation of Melan-A/MART-1 peptides that fulfill both increased immunogenicity and high resistance to biodegradation: implication for molecular anti-melanoma immunotherapy. *J Immunol.* Nov. 15, 2001;167(10):5852-61. doi: 10.4049/jimmunol.167.10.5852. cited by applicant

Borbulevych et al., Increased immunogenicity of an anchor-modified tumor-associated antigen is due to the enhanced stability of the peptide/MHC complex: implications for vaccine design. *J Immunol.* Apr. 15, 2005;174(8):4812-20. doi: 10.4049/jimmunol.174.8.4812. cited by applicant

Brichard et al., A tyrosinase nonapeptide presented by HLA-B44 is recognized on a human melanoma by autologous cytolytic T lymphocytes. *Eur J Immunol.* Jan. 1996;26(1):224-30. doi: 10.1002/eji.1830260135. cited by applicant

Cacabelos et al., Chapter 1—The Epigenetic Machinery in the Life Cycle and Pharmacoeugenetics. *Pharmacoeugenetics.* vol. 10 in *Translational Epigenetics.* 2019:1-100. doi: <https://doi.org/10.1016/B978-0-12-813939-4.00001-2>. 7 pages. cited by applicant

Campbell et al., Gesicle-Mediated Delivery of CRISPR/Cas9 Ribonucleoprotein Complex for Inactivating the HIV Provirus. *Mol Ther.* Jan. 2, 2019;27(1):151-163. doi: 10.1016/j.ymthe.2018.10.002. Epub Oct. 11, 2018. cited by applicant

Campi et al., CD4(+) T cells from healthy subjects and colon cancer patients recognize a carcinoembryonic antigen-specific immunodominant epitope. *Cancer Res.* Dec. 1, 2003;63(23):8481-6. cited by applicant

Casnici et al., Immunologic evaluation of peptides derived from BCR/ABL-out-of-frame fusion protein in HLA A2.1 transgenic mice. *J Immunother.* May 2012;35(4):321-8. doi: 10.1097/CJI.0b013e3182562d37. cited by applicant

Casnici et al., Out of frame peptides from BCR/ABL alternative splicing are immunogenic in HLA A2.1 transgenic mice. *Cancer Lett.* Apr. 8, 2009;276(1):61-7. doi: 10.1016/j.canlet.2008.10.032. Epub Dec. 4, 2008. cited by applicant

Castelli et al., Mass spectrometric identification of a naturally processed melanoma peptide recognized by CD8+ cytotoxic T lymphocytes. *J Exp Med.* Jan. 1, 1995;181(1):363-8. doi: 10.1084/jem.181.1.363. cited by applicant

Castelli et al., Novel HLA-Cw8-restricted T cell epitopes derived from tyrosinase-related protein-2 and gp100 melanoma antigens. *J Immunol.* Feb. 1, 1999;162(3):1739-48. cited by applicant

Castle et al., Exploiting the mutanome for tumor vaccination. *Cancer Res.* Mar. 1, 2012;72(5):1081-91. doi: 10.1158/0008-5472.CAN-11-3722. Epub Jan. 11, 2012. cited by applicant

Cervera et al., Generation of HIV-1 Gag VLPs by transient transfection of HEK 293 suspension cell cultures using an optimized animal-derived component free medium. *J Biotechnol.* Jul. 20, 2013;166(4):152-65. doi: 10.1016/j.jbiotec.2013.05.001. Epub May 17, 2013. cited by applicant

Chen et al., Identification of NY-ESO-1 peptide analogues capable of improved stimulation of tumor-reactive Ctl. *J Immunol.* Jul. 15, 2000;165(2):948-55. doi: 10.4049/jimmunol.165.2.948. cited by applicant

Cho et al., Optimized peptide vaccines eliciting extensive CD8 T-cell responses with therapeutic antitumor effects. *Cancer Res.* Dec. 1, 2009;69(23):9012-9. doi: 10.1158/0008-5472.CAN-09-2019. Epub Nov. 10, 2009. cited by applicant

Choi et al., Lentivirus pre-packed with Cas9 protein for safer gene editing. *Gene Ther.* Jul. 2016;23(7):627-33. doi: 10.1038/gt.2016.27. Epub Apr. 7, 2016. cited by applicant

Christensen et al., Melan-A/MART1 analog peptide triggers anti-myeloma T-cells through crossreactivity with HM1.24. *J Immunother.* Jul.-Aug. 2009;32(6):613-21. doi: 10.1097/CJI.0b013e3181a95198. cited by applicant

Correale et al., In vitro generation of human cytotoxic T lymphocytes specific for peptides derived from prostate-specific antigen. *J Natl Cancer Inst.* Feb. 19, 1997;89(4):293-300. doi: 10.1093/jnci/89.4.293. cited by applicant

Courtney et al., CRISPR/Cas9 DNA cleavage at SNP-derived PAM enables both in vitro and in vivo KRT12 mutation-specific targeting. *Gene Ther.* Jan. 2016;23(1):108-12. doi: 10.1038/gt.2015.82. Epub Aug. 20, 2015. cited by applicant

Cox et al., Identification of a peptide recognized by five melanoma-specific human cytotoxic T cell lines. *Science.* Apr. 29, 1994;264(5159):716-9. doi: 10.1126/science.7513441. cited by applicant

Cronin et al., Altering the tropism of lentiviral vectors through pseudotyping. *Curr Gene Ther.* Aug. 2005;5(4):387-98. doi: 10.2174/1566523054546224. Erratum in: *Curr Gene Ther.* Oct. 2005;5(5):531. Author Manuscript, 19 pages. cited by applicant

Crosti et al., Identification of novel subdominant epitopes on the carcinoembryonic antigen recognized by CD4+ T cells of lung cancer patients. *J Immunol.* Apr. 15, 2006;176(8):5093-9. doi: 10.4049/jimmunol.176.8.5093. cited by applicant

Dalet et al., An antigenic peptide produced by reverse splicing and double asparagine deamidation. *Proc Natl Acad Sci U S A.* Jul. 19, 2011;108(29):E323-31. doi: 10.1073/pnas.1101892108. Epub Jun. 13, 2011. cited by applicant

David et al., Viral Vectors: The Road to Reducing Genotoxicity. *Toxicol Sci.* Feb. 2017;155(2):315-325. doi: 10.1093/toxsci/kfw220. Epub Nov. 1, 2016. cited by applicant

Depontieu et al., Identification of tumor-associated, MHC class II-restricted phosphopeptides as targets for immunotherapy. *Proc Natl Acad Sci U S A.* Jul. 21, 2009;106(29):12073-8. doi: 10.1073/pnas.0903852106. Epub Jul. 6, 2009. cited by applicant

Di Stasi et al., Review of the Results of WT1 Peptide Vaccination Strategies for Myelodysplastic Syndromes and Acute Myeloid Leukemia from Nine Different Studies. *Front Immunol.* Feb. 4, 2015;6:36. doi: 10.3389/fimmu.2015.00036. cited by applicant

Duan et al., Immune rejection of mouse tumors expressing mutated self. *Cancer Res.* Apr. 15, 2009;69(8):3545-53. doi: 10.1158/0008-5472.CAN-08-2779. Epub Apr. 7, 2009. Author Manuscript. 18 pages. cited by applicant

Duportet et al., A platform for rapid prototyping of synthetic gene networks in mammalian cells. *Nucleic Acids Res.* Dec. 1, 2014;42(21):13440-51. doi: 10.1093/nar/gku1082. Epub Nov. 5, 2014. cited by applicant

Fontana et al., Rabies virus-like particles expressed in HEK293 cells. *Vaccine.* May 19, 2014;32(24):2799-804. doi: 10.1016/j.vaccine.2014.02.031. Epub Mar. 12, 2014. cited by applicant

Fonteneau et al., The Tumor Antigen NY-ESO-1 Mediates Direct Recognition of Melanoma Cells by CD4+ T Cells after Intercellular Antigen Transfer. *J Immunol.* Jan. 1, 2016;196(1):64-71. doi: 10.4049/jimmunol.1402664. Epub Nov. 25, 2015. cited by applicant

Fourcade et al., PD-1 and Tim-3 regulate the expansion of tumor antigen-specific CD8? T cells induced by melanoma vaccines. *Cancer Res.* Feb. 15, 2014;74(4):1045-55. doi: 10.1158/0008-5472.CAN-13-2908. Epub Dec. 16, 2013. cited by applicant

Fridman et al., An efficient T-cell epitope discovery strategy using in silico prediction and the iTopia assay platform. *Oncoimmunology.* Nov. 1, 2012;1(8):1258-1270. doi: 10.4161/onci.21355. cited by applicant

Fujiki et al., Identification and characterization of a WT1 (Wilms Tumor Gene) protein-derived HLA-DRB1\*0405-restricted 16-mer helper peptide that promotes the induction and activation of WT1-specific cytotoxic T lymphocytes. *J Immunother.* Apr. 2007;30(3):282-93. doi: 10.1097/01.cji.0000211337.91513.94. cited by applicant

Gee et al., Extracellular nanovesicles for packaging of CRISPR-Cas9 protein and sgRNA to induce therapeutic exon skipping. *Nat Commun.* Mar. 13, 2020;11(1):1334. doi: 10.1038/s41467-020-14957-y. cited by applicant



GenBank Access No. BAP64357. Aug. 1, 2013. 1 page. cited by applicant

Geynisman et al., A randomized pilot phase I study of modified carcinoembryonic antigen (CEA) peptide (CAP1-6D)/montanide/GM-CSF-vaccine in patients with pancreatic adenocarcinoma. *J Immunother Cancer*. Jun. 27, 2013;1:8. doi: 10.1186/2051-1426-1-8. cited by applicant

Ghosh et al., Synapsis in phage Bxb1 integration: selection mechanism for the correct pair of recombination sites. *J Mol Biol*. Jun. 3, 2005;349(2):331-48. doi: 10.1016/j.jmb.2005.03.043. Epub Apr. 7, 2005. cited by applicant

Girard-Gagnepain et al., Baboon envelope pseudotyped LVs outperform VSV-G-LVs for gene transfer into early-cytokine-stimulated and resting HSCs. *Blood*. Aug. 21, 2014;124(8):1221-31. doi: 10.1182/blood-2014-02-558163. Epub Jun. 20, 2014. cited by applicant

Godefroy et al., Identification of two Melan-A CD4+ T cell epitopes presented by frequently expressed MHC class II alleles. *Clin Immunol*. Oct. 2006;121(1):54-62. doi: 10.1016/j.clim.2006.05.007. Epub Jun. 30, 2006. cited by applicant

Graff-Dubois et al., Generation of CTL recognizing an HLA-A\*0201-restricted epitope shared by MAGE-A1, -A2, -A3, -A4, -A6, -A10, and -A12 tumor antigens: implication in a broad-spectrum tumor immunotherapy. *J Immunol*. Jul. 1, 2002;169(1):575-80. doi: 10.4049/jimmunol.169.1.575. cited by applicant

Gross et al., High vaccination efficiency of low-affinity epitopes in antitumor immunotherapy. *J Clin Invest*. Feb. 2004;113(3):425-33. doi: 10.1172/JCI19418. cited by applicant

Guevara-Patiño et al., Optimization of a self antigen for presentation of multiple epitopes in cancer immunity. *J Clin Invest*. May 2006;116(5):1382-90. doi: 10.1172/JCI25591. Epub Apr. 13, 2006. cited by applicant

Guibinga et al., Cell surface heparan sulfate is a receptor for attachment of envelope protein-free retrovirus-like particles and VSV-G pseudotyped MLV-derived retrovirus vectors to target cells. *Mol Ther*. May 2002;5(5 Pt 1):538-46. doi: 10.1006/mthe.2002.0578. cited by applicant

Gulley et al., Combining a Recombinant Cancer Vaccine with Standard Definitive Radiotherapy in Patients with Localized Prostate Cancer. *Clin Cancer Res*. May 2, 2005;11(9):3353-62. doi: 10.1158/1078-0432.CCR-04-2062. cited by applicant

Guo et al., Direct recognition and lysis of leukemia cells by WT1-specific CD4+ T lymphocytes in an HLA class II-restricted manner. *Blood*. Aug. 15, 2005;106(4):1415-8. doi: 10.1182/blood-2005-01-0413. Epub Apr. 21, 2005. cited by applicant

Haeussler et al., Genome Editing with CRISPR-Cas9: Can It Get Any Better? *J Genet Genomics*. May 20, 2016;43(5):239-50. doi: 10.1016/j.jgg.2016.04.008. Epub Apr. 24, 2016. Author Manuscript. 22 pages. cited by applicant

Herbst-Kralovetz et al., Norwalk virus-like particles as vaccines. *Expert Rev Vaccines*. Mar. 2010;9(3):299-307. doi: 10.1586/erv.09.163. Author Manuscript, 16 pages. cited by applicant

Hirohashi et al., An HLA-A24-restricted cytotoxic T lymphocyte epitope of a tumor-associated protein, survivin. *Clin Cancer Res*. Jun. 2002;8(6):1731-9. cited by applicant

Hong et al., Novel recombinant hepatitis B virus vectors efficiently deliver protein and RNA encoding genes into primary hepatocytes. *J Virol*. Jun. 2013;87(12):6615-24. doi: 10.1128/JVI.03328-12. Epub Apr. 3, 2013. cited by applicant

Jalaguier et al., Efficient production of HIV-1 virus-like particles from a mammalian expression vector requires the N-terminal capsid domain. *PLoS One*. 2011;6(11):e28314. doi: 10.1371/journal.pone.0028314. Epub Nov. 30, 2011. cited by applicant

Jaramillo et al., Identification of HLA-A3-restricted CD8+ T cell epitopes derived from mammaglobin-A, a tumor-associated antigen of human breast cancer. *Int J Cancer*. Dec. 10, 2002;102(5):499-506. doi: 10.1002/ijc.10736. cited by applicant

Joglekar et al., Pseudotyped Lentiviral Vectors: One Vector, Many Guises. *Hum Gene Ther Methods*. Dec. 2017;28(6):291-301. doi: 10.1089/hgtb.2017.084. Epub Sep. 4, 2017. cited by applicant

Kaczmarczyk et al., Protein delivery using engineered virus-like particles. *Proc Natl Acad Sci US A*. Oct. 11, 2011;108(41):16998-7003. doi: 10.1073/pnas.1101874108. Epub Sep. 26, 2011. cited by applicant

Kang et al., Chimeric rabies virus-like particles containing membrane-anchored GM-CSF enhances the immune response against rabies virus. *Viruses*. Mar. 11, 2015;7(3):1134-52. doi: 10.3390/v7031134. cited by applicant

Kang et al., Identification of a tyrosinase epitope recognized by HLA-A24-restricted, tumor-infiltrating lymphocytes. *J Immunol*. Aug. 1, 1995;155(3):1343-8. cited by applicant

Karbach et al., Long-term complete remission following radiosurgery and immunotherapy in a melanoma patient with brain metastasis: immunologic correlates. *Cancer Immunol Res*. May 2014;2(5):404-9. doi: 10.1158/2326-6066.CIR-13-0200. Epub Feb. 5, 2014. cited by applicant

Kato et al., A lentiviral strategy for highly efficient retrograde gene transfer by pseudotyping with fusion envelope glycoprotein. *Hum Gene Ther*. Feb. 2011;22(2):197-206. doi: 10.1089/hum.2009.179. Epub Jan. 27, 2011. cited by applicant

Kato et al., Selective neural pathway targeting reveals key roles of thalamostriatal projection in the control of visual discrimination. *J Neurosci*. Nov. 23, 2011;31(47):17169-79. doi: 10.1523/JNEUROSCI.4005-11.2011. cited by applicant

Kawakami et al., Identification of a human melanoma antigen recognized by tumor-infiltrating lymphocytes associated with in vivo tumor rejection. *Proc Natl Acad Sci U S A*. Jul. 5, 1994;91(14):6458-62. doi: 10.1073/pnas.91.14.6458. cited by applicant

Kawakami et al., Identification of new melanoma epitopes on melanosomal proteins recognized by tumor infiltrating T

lymphocytes restricted by HLA-A1, -A2, and -A3 alleles. *J Immunol.* Dec. 15, 1998;161(12):6985-92. cited by applicant

Kawakami et al., Identification of the immunodominant peptides of the MART-1 human melanoma antigen recognized by the majority of HLA-A2-restricted tumor infiltrating lymphocytes. *J Exp Med.* Jul. 1, 1994;180(1):347-52. doi: 10.1084/jem.180.1.347. cited by applicant

Kawakami et al., Recognition of multiple epitopes in the human melanoma antigen gp100 by tumor-infiltrating T lymphocytes associated with in vivo tumor regression. *J Immunol.* Apr. 15, 1995;154(8):3961-8. cited by applicant

Kawashima et al., Identification of gp100-derived, melanoma-specific cytotoxic T-lymphocyte epitopes restricted by HLA-A3 supertype molecules by primary in vitro immunization with peptide-pulsed dendritic cells. *Int J Cancer.* Nov. 9, 1998;78(4):518-24. doi: 10.1002/(sici)1097-0215(19981109)78:4<518::aid-ijc20>3.0.co;2-0. cited by applicant

Kawashima et al., Identification of HLA-A3-restricted cytotoxic T lymphocyte epitopes from carcinoembryonic antigen and HER-2/neu by primary in vitro immunization with peptide-pulsed dendritic cells. *Cancer Res.* Jan. 15, 1999;59(2):431-5. cited by applicant

Kawashima et al., The multi-epitope approach for immunotherapy for cancer: identification of several CTL epitopes from various tumor-associated antigens expressed on solid epithelial tumors. *Hum Immunol.* Jan. 1998;59(1):1-14. doi: 10.1016/s0198-8859(97)00255-3. cited by applicant

Kemmler et al., Elevated tumor-associated antigen expression suppresses variant peptide vaccine responses. *J Immunol.* Nov. 1, 2011;187(9):4431-9. doi: 10.4049/jimmunol.1101555. Epub Sep. 21, 2011. cited by applicant

Kittlesen et al., Human melanoma patients recognize an HLA-A1-restricted CTL epitope from tyrosinase containing two cysteine residues: implications for tumor vaccine development. *J Immunol.* Mar. 1, 1998;160(5):2099-106. Erratum in: *J Immunol* Mar. 1, 1999;162(5):3106. Shabanowitz JA [corrected to Shabanowitz J]. cited by applicant

Kizer et al., Application of functional genomics to pathway optimization for increased isoprenoid production. *Appl Environ Microbiol.* May 2008;74(10):3229-41. doi: 10.1128/AEM.02750-07. Epub Mar. 14, 2008. cited by applicant

Kobayashi et al., CD4+ T cells from peripheral blood of a melanoma patient recognize peptides derived from nonmutated tyrosinase. *Cancer Res.* Jan. 15, 1998;58(2):296-301. cited by applicant

Kobayashi et al., Identification of an antigenic epitope for helper T lymphocytes from carcinoembryonic antigen. *Clin Cancer Res.* Oct. 2002;8(10):3219-25. cited by applicant

Kobayashi et al., Identification of helper T-cell epitopes that encompass or lie proximal to cytotoxic T-cell epitopes in the gp100 melanoma tumor antigen. *Cancer Res.* Oct. 15, 2001;61(20):7577-84. cited by applicant

Kotterman et al., Engineering adeno-associated viruses for clinical gene therapy. *Nat Rev Genet.* Jul. 2014;15(7):445-51. doi: 10.1038/nrg3742. Epub May 20, 2014. cited by applicant

Kueh et al., The new editor-targeted genome engineering in the absence of homology-directed repair. *Cell Death Discov.* Jun. 13, 2016;2:16042. doi: 10.1038/cddiscovery.2016.42. cited by applicant

Kushnir et al., Virus-like particles as a highly efficient vaccine platform: diversity of targets and production systems and advances in clinical development. *Vaccine.* Dec. 17, 2012;31(1):58- 83. doi: 10.1016/j.vaccine.2012.10.083. Epub Nov. 6, 2012. cited by applicant

Lally et al., Unmasking cryptic epitopes after loss of immunodominant tumor antigen expression through epitope spreading. *Int J Cancer.* Sep. 2001;93(6):841-7. doi: 10.1002/ijc.1420. cited by applicant

Lapointe et al., Retrovirally transduced human dendritic cells can generate T cells recognizing multiple MHC class I and class II epitopes from the melanoma antigen glycoprotein 100. *J Immunol.* Oct. 15, 2001;167(8):4758-64. doi: 10.4049/jimmunol.167.8.4758. cited by applicant

Larrieu et al., A HLA-Cw\*0701 restricted Melan-A/MART1 epitope presented by melanoma tumor cells to CD8+ tumor infiltrating lymphocytes. *Cancer Immunol Immunother.* May 2008;57(5):745-52. doi: 10.1007/s00262-007-0436-7. Epub Dec. 21, 2007. cited by applicant

Latham et al., Formation of wild-type and chimeric influenza virus-like particles following simultaneous expression of only four structural proteins. *J Virol.* Jul. 2001;75(13):6154-65. doi: 10.1128/JVI.75.13.6154-6165.2001. cited by applicant

Lennerz et al., The response of autologous T cells to a human melanoma is dominated by mutated neoantigens. *Proc Natl Acad Sci U S A.* Nov. 1, 2005;102(44):16013-8. doi: 10.1073/pnas.0500090102. Epub Oct. 24, 2005. cited by applicant

Li et al., Expression and self-assembly of empty virus-like particles of hepatitis E virus. *J Virol.* Oct. 1997;71(10):7207-13. doi: 10.1128/JVI.71.10.7207-7213.1997. cited by applicant

Lin et al., HLA-DPB1\*05: 01-restricted WT1332-specific TCR-transduced CD4+ T lymphocytes display a helper activity for WT1-specific CTL induction and a cytotoxicity against leukemia cells. *J Immunother.* Apr. 2013;36(3):159-70. doi: 10.1097/CJI.0b013e3182873581. cited by applicant

Lu, Periodic Chart of Amino Acid PDF. Accessed on the internet at [https://figshare.com/articles/figure/periodic\\_chart\\_of\\_amino\\_acid\\_pdf/3445001/1](https://figshare.com/articles/figure/periodic_chart_of_amino_acid_pdf/3445001/1). Posted Jun. 21, 2016. [www.bachem.com](http://www.bachem.com). 1 page. cited by applicant

Ludwig et al., Virus-like particles-universal molecular toolboxes. *Curr Opin Biotechnol.* Dec. 2007;18(6):537-45. doi: 10.1016/j.copbio.2007.10.013. cited by applicant

Lueck et al., Engineered transfer RNAs for suppression of premature termination codons. *Nat Commun.* Feb. 18, 2019;10(1):822. doi: 10.1038/s41467-019-08329-4. cited by applicant

Lupetti et al., Translation of a retained intron in tyrosinase-related protein (TRP) 2 mRNA generates a new cytotoxic T

lymphocyte (CTL)-defined and shared human melanoma antigen not expressed in normal cells of the melanocytic lineage. *J Exp Med.* Sep. 21, 1998;188(6):1005- 16. doi: 10.1084/jem.188.6.1005. cited by applicant

Lyu et al., Delivering Cas9/sgRNA ribonucleoprotein (RNP) by lentiviral capsid-based bionanoparticles for efficient 'hit-and-run' genome editing. *Nucleic Acids Res.* Sep. 26, 2019;47(17):e99. doi: 10.1093/nar/gkz605. cited by applicant

Maetzig et al., Retroviral protein transfer: falling apart to make an impact. *Curr Gene Ther.* Oct. 2012;12(5):389-409. doi: 10.2174/156652312802762581. cited by applicant

Mandic et al., The alternative open reading frame of LAGE-1 gives rise to multiple promiscuous HLA-DR-restricted epitopes recognized by T-helper 1-type tumor-reactive CD4+ T cells. *Cancer Res.* Oct. 1, 2003;63(19):6506-15. cited by applicant

Mangeot et al., A universal transgene silencing method based on RNA interference. *Nucleic Acids Res.* Jul. 12, 2004;32(12):e102. doi: 10.1093/nar/gnh105. cited by applicant

Mangeot et al., Development of minimal lentivirus vectors derived from simian immunodeficiency virus (SIVmac251) and their use for gene transfer into human dendritic cells. *J Virol.* Sep. 2000;74(18):8307-15. doi: 10.1128/jvi.74.18.8307-8315.2000. cited by applicant

Mangeot et al., Protein transfer into human cells by VSV-G-induced nanovesicles. *Mol Ther.* Sep. 2011;19(9):1656-66. doi: 10.1038/mt.2011.138. Epub Jul. 12, 2011. cited by applicant

Mariani et al., Species-specific exclusion of APOBEC3G from HIV-1 virions by Vif. *Cell.* Jul. 11, 2003;114(1):21-31. doi: 10.1016/s0092-8674(03)00515-4. cited by applicant

Meng et al., Identification of an HLA-DPB1\*0501 restricted Melan-A/MART-1 epitope recognized by CD4+ T lymphocytes: prevalence for immunotherapy in Asian populations. *J Immunother.* Sep. 2011;34(7):525-34. doi: 10.1097/CJI.0b013e318226bd45. Author Manuscript. 16 pages. cited by applicant

Michaux et al., A spliced antigenic peptide comprising a single spliced amino acid is produced in the proteasome by reverse splicing of a longer peptide fragment followed by trimming. *J Immunol.* Feb. 15, 2014;192(4):1962-71. doi: 10.4049/jimmunol.1302032. Epub Jan. 22, 2014. cited by applicant

Milone et al., Clinical use of lentiviral vectors. *Leukemia.* Jul. 2018;32(7):1529-1541. doi: 10.1038/s41375-018-0106-0. Epub Mar. 22, 2018. cited by applicant

Momose et al., Diving into marine genomics with CRISPR/Cas9 systems. *Mar Genomics.* Dec. 2016;30:55-65. doi: 10.1016/j.margen.2016.10.003. Epub Oct. 12, 2016. cited by applicant

Morel et al., A tyrosinase peptide presented by HLA-B35 is recognized on a human melanoma by autologous cytotoxic T lymphocytes. *Int J Cancer.* Dec. 10, 1999;83(6):755-9. doi: 10.1002/(sici)1097-0215(19991210)83:6<755::aid-ijc10>3.0.co;2-s. cited by applicant

Mselli-Lakhal et al., Gene transfer system derived from the caprine arthritis-encephalitis lentivirus. *J Virol Methods.* Sep. 2006;136(1-2):177-84. doi: 10.1016/j.jviromet.2006.05.006. Epub Jun. 21, 2006. cited by applicant

Murawski et al., Newcastle disease virus-like particles containing respiratory syncytial virus G protein induced protection in BALB/c mice, with no evidence of immunopathology. *J Virol.* Jan. 2010;84(2):1110-23. doi: 10.1128/JVI.01709-09. Epub Nov. 4, 2009. cited by applicant

Naskalska et al., Virus Like Particles as Immunogens and Universal Nanocarriers. *Pol J Microbiol.* 2015;64(1):3-13. cited by applicant

Negre et al., Characterization of novel safe lentiviral vectors derived from simian immunodeficiency virus (SIVmac251) that efficiently transduce mature human dendritic cells. *Gene Ther.* Oct. 2000;7(19):1613-23. doi: 10.1038/sj.gt.3301292. cited by applicant

Noppen et al., Naturally processed and concealed HLA-A2.1-restricted epitopes from tumor-associated antigen tyrosinase-related protein-2. *Int J Cancer.* Jul. 15, 2000;87(2):241-6. cited by applicant

Nukaya et al., Identification of HLA-A24 epitope peptides of carcinoembryonic antigen which induce tumor-reactive cytotoxic T lymphocyte. *Int J Cancer.* Jan. 5, 1999;80(1):92-7. doi: 10.1002/(sici)1097-0215(19990105)80:1<92::aid-ijc18>3.0.co;2-m. cited by applicant

Ogasawara et al., Recombinant viral-like particles of parvovirus B19 as antigen carriers of anthrax protective antigen. *In Vivo.* May-Jun. 2006;20(3):319-24. cited by applicant

Ohminami et al., HLA class I-restricted lysis of leukemia cells by a CD8(+) cytotoxic T-lymphocyte clone specific for WT1 peptide. *Blood.* Jan. 1, 2000;95(1):286-93. cited by applicant

Oka et al., WT1 peptide vaccine for the treatment of cancer. *Curr Opin Immunol.* Apr. 2008;20(2):211-20. doi: 10.1016/j.coi.2008.04.009. Epub May 24, 2008. cited by applicant

Olsen, J.C., Gene transfer vectors derived from equine infectious anemia virus. *Gene Ther.* Nov. 1998;5(11):1481-7. doi: 10.1038/sj.gt.3300768. cited by applicant

Olson et al., HLA-A2-restricted T-cell epitopes specific for prostatic acid phosphatase. *Cancer Immunol Immunother.* Jun. 2010;59(6):943-53. doi: 10.1007/s00262-010-0820-6. Epub Feb. 6, 2010. cited by applicant

Osen et al., Screening of human tumor antigens for CD4 T cell epitopes by combination of HLA-transgenic mice, recombinant adenovirus and antigen peptide libraries. *PLoS One.* Nov. 30, 2010;5(11):e14137. doi: 10.1371/journal.pone.0014137. cited by applicant

Parkhurst et al., Identification of a shared HLA-A\*0201-restricted T-cell epitope from the melanoma antigen tyrosinase-related protein 2 (TRP2). *Cancer Res.* Nov. 1, 1998;58(21):4895-901. cited by applicant

Parkhurst et al., Induction of CD4+ Th1 lymphocytes that recognize known and novel class II MHC restricted epitopes from the melanoma antigen gp100 by stimulation with recombinant protein. *J Immunother.* Mar.-Apr. 2004;27(2):79-91. doi: 10.1097/00002371-200403000-00001. Author Manuscript. 22 pages. cited by applicant

Paschen et al., Detection of spontaneous CD4+ T-cell responses in melanoma patients against a tyrosinase-related protein-2-derived epitope identified in HLA-DRB1\*0301 transgenic mice. *Clin Cancer Res.* Jul. 15, 2005;11(14):5241-7. doi: 10.1158/1078-0432.CCR-05-0170. cited by applicant

Pinilla et al., Combinatorial peptide libraries as an alternative approach to the identification of ligands for tumor-reactive cytolytic T lymphocytes. *Cancer Res.* Jul. 1, 2001;61(13):5153-60. cited by applicant

Pinilla-Ibarz et al., Improved human T-cell responses against synthetic HLA-0201 analog peptides derived from the WT1 oncoprotein. *Leukemia.* Nov. 2006;20(11):2025-33. doi: 10.1038/sj.leu.2404380. Epub Aug. 31, 2006. cited by applicant

Prather et al., De novo biosynthetic pathways: rational design of microbial chemical factories. *Curr Opin Biotechnol.* Oct. 2008;19(5):468-74. doi: 10.1016/j.copbio.2008.07.009. Epub Sep. 5, 2008. cited by applicant

Quan et al., Influenza M1 VLPs containing neuraminidase induce heterosubtypic cross-protection. *Virology.* Sep. 1, 2012;430(2):127-35. doi: 10.1016/j.virol.2012.05.006. Epub Jun. 2, 2012. cited by applicant

Rasmussen et al., Characterization of virus-like particles produced by a recombinant baculovirus containing the gag gene of the bovine immunodeficiency-like virus. *Virology.* Oct. 1990;178(2):435-51. doi: 10.1016/0042-6822(90)90341-n. cited by applicant

Renner et al., Intact Viral Particle Counts Measured by Flow Virometry Provide Insight into the Infectivity and Genome Packaging Efficiency of Moloney Murine Leukemia Virus. *J Virol.* Jan. 6, 2020;94(2):e01600-19. doi: 10.1128/JVI.01600-19. cited by applicant

Riley et al., Identification of a new shared HLA-A2.1 restricted epitope from the melanoma antigen tyrosinase. *J Immunother.* May-Jun. 2001;24(3):212-20. cited by applicant

Rimoldi et al., Efficient simultaneous presentation of NY-ESO-1/LAGE-1 primary and nonprimary open reading frame-derived CTL epitopes in melanoma. *J Immunol.* Dec. 15, 2000;165(12):7253-61. doi: 10.4049/jimmunol.165.12.7253. cited by applicant

Robbins et al., Multiple HLA class II-restricted melanocyte differentiation antigens are recognized by tumor-infiltrating lymphocytes from a patient with melanoma. *J Immunol.* Nov. 15, 2002;169(10):6036-47. doi: 10.4049/jimmunol.169.10.6036. cited by applicant

Robbins et al., The intronic region of an incompletely spliced gp100 gene transcript encodes an epitope recognized by melanoma-reactive tumor-infiltrating lymphocytes. *J Immunol.* Jul. 1, 1997;159(1):303-8. cited by applicant

Rohovie et al., Virus-like particles: Next-generation nanoparticles for targeted therapeutic delivery. *Bioeng Transl Med.* Jan. 19, 2017;2(1):43-57. doi: 10.1002/btm2.10049. cited by applicant

Rosenberg et al., Immunologic and therapeutic evaluation of a synthetic peptide vaccine for the treatment of patients with metastatic melanoma. *Nat Med.* Mar. 1998;4(3):321-7. doi: 10.1038/nm0398-321. cited by applicant

Rubio-Godoy et al., Toward synthetic combinatorial peptide libraries in positional scanning format (PS-SCL)-based identification of CD8+ Tumor-reactive T-Cell Ligands: a comparative analysis of PS-SCL recognition by a single tumor-reactive CD8+ cytolytic T-lymphocyte clone. *Cancer Res.* Apr. 1, 2002;62(7):2058-63. cited by applicant

Ruiz et al., Identification and characterization of a T-helper peptide from carcinoembryonic antigen. *Clin Cancer Res.* Apr. 15, 2004;10(8):2860-7. doi: 10.1158/1078-0432.ccr-03-0476. cited by applicant

Rusk, Cas9 and the importance of asymmetry. *Nat Methods.* Apr. 2016;13(4):286-7. doi: 10.1038/nmeth.3826. cited by applicant

Saenger et al., Improved tumor immunity using anti-tyrosinase related protein-1 monoclonal antibody combined with DNA vaccines in murine melanoma. *Cancer Res.* Dec. 1, 2008;68(23):9884-91. doi: 10.1158/0008-5472.CAN-08-2233. Author Manuscript. 19 pages. cited by applicant

Saenz et al., Feline immunodeficiency virus-based lentiviral vectors. *Cold Spring Harb Protoc.* Jan. 1, 2012;2012(1):71-6. doi: 10.1101/pdb.ip067579. cited by applicant

Saenz et al., Production, harvest, and concentration of feline immunodeficiency virus-based lentiviral vector from cells grown in CF10 or CF2 devices. *Cold Spring Harb Protoc.* Jan. 1, 2012;2012(1):118-23. doi: 10.1101/pdb.prot067546. cited by applicant

Sakuma et al., Multiplex genome engineering in human cells using all-in-one CRISPR/Cas9 vector system. *Sci Rep.* Jun. 23, 2014;4:5400. doi: 10.1038/srep05400. cited by applicant

Schneider et al., Overlapping peptides of melanocyte differentiation antigen Melan-A/MART-1 recognized by autologous cytolytic T lymphocytes in association with HLA-B45.1 and HLA-A2.1. *Int J Cancer.* Jan. 30, 1998;75(3):451-8. doi: 10.1002/(sici)1097-0215(19980130)75:3<451::aid-ijc20>3.0.co;2-a. cited by applicant

Sensi et al., Identification of a novel gp100/pMel17 peptide presented by HLA-A\*6801 and recognized on human melanoma by cytolytic T cell clones. *Tissue Antigens.* Apr. 2002;59(4):273-9. doi: 10.1034/j.1399-0039.2002.590404.x. cited by applicant

Shang et al., The spontaneous CD8+ T-cell response to HLA-A2-restricted NY-ESO-1b peptide in hepatocellular carcinoma patients. *Clin Cancer Res.* Oct. 15, 2004;10(20):6946-55. doi: 10.1158/1078-0432.CCR-04-0502. cited by applicant

Sharma et al., Noninfectious virus-like particles produced by Moloney murine leukemia virus-based retrovirus packaging

cells deficient in viral envelope become infectious in the presence of lipofection reagents. *Proc Natl Acad Sci U S A*. Sep. 30, 1997;94(20):10803-8. doi: 10.1073/pnas.94.20.10803. cited by applicant

Shellenberger et al., A recombinant polypeptide extends the in vivo half-life of peptides and proteins in a tunable manner. *Nat Biotechnol*. Dec. 2009;27(12):1186-90. doi: 10.1038/nbt.1588. cited by applicant

Shen et al., Identification of a MHC class-II restricted epitope in carcinoembryonic antigen. *Cancer Immunol Immunother*. May 2004;53(5):391-403. doi: 10.1007/s00262-003-0455-y. Epub Nov. 18, 2003. cited by applicant

Shim et al., Nonviral Delivery Systems for Cancer Gene Therapy: Strategies and Challenges. *Curr Gene Ther*. 2018;18(1):3-20. doi: 10.2174/1566523218666180119121949. cited by applicant

Skipper et al., An HLA-A2-restricted tyrosinase antigen on melanoma cells results from posttranslational modification and suggests a novel pathway for processing of membrane proteins. *J Exp Med*. Feb. 1, 1996;183(2):527-34. doi: 10.1084/jem.183.2.527. cited by applicant

Skipper et al., Shared epitopes for HLA-A3-restricted melanoma-reactive human CTL include a naturally processed epitope from Pmel-17/gp100. *J Immunol*. Dec. 1, 1996;157(11):5027-33. cited by applicant

Slansky et al., Enhanced antigen-specific antitumor immunity with altered peptide ligands that stabilize the MHC-peptide-TCR complex. *Immunity*. Oct. 2000;13(4):529-38. doi: 10.1016/s1074-7613(00)00052-2. cited by applicant

Slingluff et al., Clinical and immunologic results of a randomized phase II trial of vaccination using four melanoma peptides either administered in granulocyte-macrophage colony-stimulating factor in adjuvant or pulsed on dendritic cells. *J Clin Oncol*. Nov. 1, 2003;21(21):4016-26. doi: 10.1200/JCO.2003.10.005. cited by applicant

Slingluff et al., Immunologic and clinical outcomes of vaccination with a multiepitope melanoma peptide vaccine plus low-dose interleukin-2 administered either concurrently or on a delayed schedule. *J Clin Oncol*. Nov. 15, 2004;22(22):4474-85. doi: 10.1200/JCO.2004.10.212. cited by applicant

Tangri et al., Structural features of peptide analogs of human histocompatibility leukocyte antigen class I epitopes that are more potent and immunogenic than wild-type peptide. *J Exp Med*. Sep. 17, 2001;194(6):833-46. doi: 10.1084/jem.194.6.833. cited by applicant

Tomé-Amat et al., Secreted production of assembled Norovirus virus-like particles from *Pichia pastoris*. *Microb Cell Fact*. Sep. 10, 2014;13:134. doi: 10.1186/s12934-014-0134-z. cited by applicant

Topalian et al., Melanoma-specific CD4+ T cells recognize nonmutated HLA-DR-restricted tyrosinase epitopes. *J Exp Med*. May 1, 1996;183(5):1965-71. doi: 10.1084/jem.183.5.1965. cited by applicant

Touloukian et al., Expression of a "self-"antigen by human tumor cells enhances tumor antigen-specific CD4(+) T-cell function. *Cancer Res*. Sep. 15, 2002;62(18):5144-7. Author Manuscript. 11 pages. cited by applicant

Touloukian et al., Identification of a MHC class II-restricted human gp100 epitope using DR4-IE transgenic mice. *J Immunol*. Apr. 1, 2000;164(7):3535-42. doi: 10.4049/jimmunol.164.7.3535. cited by applicant

Touloukian et al., Normal tissue depresses while tumor tissue enhances human T cell responses in vivo to a novel self/tumor melanoma antigen, OA1. *J Immunol*. Feb. 1, 2003;170(3): 1579-85. doi: 10.4049/jimmunol.170.3.1579. cited by applicant

Trojan et al., Generation of cytotoxic T lymphocytes against native and altered peptides of human leukocyte antigen-A\*0201 restricted epitopes from the human epithelial cell adhesion molecule. *Cancer Res*. Jun. 15, 2001;61(12):4761-5. cited by applicant

Tsai et al., Identification of subdominant CTL epitopes of the GP100 melanoma-associated tumor antigen by primary in vitro immunization with peptide-pulsed dendritic cells. *J Immunol*. Feb. 15, 1997;158(4):1796-802. cited by applicant

Tsang et al., A human cytotoxic T-lymphocyte epitope and its agonist epitope from the nonvariable No. of tandem repeat sequence of MUC-1. *Clin Cancer Res*. Mar. 15, 2004;10(6):2139-49. doi: 10.1158/1078-0432.ccr-1011-03. cited by applicant

Tsang et al., Generation of human cytotoxic T cells specific for human carcinoembryonic antigen epitopes from patients immunized with recombinant vaccinia-CEA vaccine. *J Natl Cancer Inst*. Jul. 5, 1995;87(13):982-90. doi: 10.1093/jnci/87.13.982. cited by applicant

Tsuboi et al., Enhanced induction of human WT1-specific cytotoxic T lymphocytes with a 9-mer WT1 peptide modified at HLA-A\*2402-binding residues. *Cancer Immunol Immunother*. Dec. 2002;51(11-12):614-20. doi: 10.1007/s00262-002-0328-9. Epub Oct. 18, 2002. cited by applicant

Tuorto et al., Genome recoding by tRNA modifications. *Open Biol*. Dec. 2016;6(12):160287. doi: 10.1098/rsob.160287. cited by applicant

Tycko et al., Methods for Optimizing CRISPR-Cas9 Genome Editing Specificity. *Mol Cell*. Aug. 4, 2016;63(3):355-70. doi: 10.1016/j.molcel.2016.07.004. cited by applicant

Valmori et al., Analysis of the cytolytic T lymphocyte response of melanoma patients to the naturally HLA-A\*0201-associated tyrosinase peptide 368-376. *Cancer Res*. Aug. 15, 1999;59(16):4050-5. cited by applicant

Valmori et al., Enhanced generation of specific tumor-reactive CTL in vitro by selected Melan-A/MART-1 immunodominant peptide analogues. *J Immunol*. Feb. 15, 1998;160(4):1750-8. cited by applicant

Valmori et al., Naturally occurring human lymphocyte antigen-A2 restricted CD8+ T-cell response to the cancer testis antigen NY-ESO-1 in melanoma patients. *Cancer Res*. Aug. 15, 2000;60(16):4499-506. cited by applicant

Vigneron et al., A peptide derived from melanocytic protein gp100 and presented by HLA-B35 is recognized by autologous cytolytic T lymphocytes on melanoma cells. *Tissue Antigens*. Feb. 2005;65(2):156-62. doi: 10.1111/j.1399-

0039.2005.00365.x. cited by applicant

Vigneron et al., An antigenic peptide produced by peptide splicing in the proteasome. *Science*. Apr. 23, 2004;304(5670):587-90. doi: 10.1126/science.1095522. cited by applicant

Visseren et al., Affinity, specificity and T-cell-receptor diversity of melanoma-specific CTL generated in vitro against a single tyrosinase epitope. *Int J Cancer*. Sep. 17, 1997;72(6):1122-8. doi: 10.1002/(sici)1097-0215(19970917)72:6<1122::aid-ijc30>3.0.co;2-3. cited by applicant

Voelkel et al., Protein transduction from retroviral Gag precursors. *Proc Natl Acad Sci U S A*. Apr. 27, 2010;107(17):7805-10. doi: 10.1073/pnas.0914517107. Epub Apr. 12, 2010. cited by applicant

Volpe et al., Alternative BCR/ABL splice variants in Philadelphia chromosome-positive leukemias result in novel tumor-specific fusion proteins that may represent potential targets for immunotherapy approaches. *Cancer Res*. Jun. 1, 2007;67(11):5300-7. doi: 10.1158/0008-5472.CAN-06-3737. cited by applicant

Voutev et al., Bxb1 phage recombinase assists genome engineering in *Drosophila melanogaster*. *Biotechniques*. Jan. 1, 2017;62(1):37-38. doi: 10.2144/000114494. cited by applicant

Walpita et al., Mammalian Cell-Derived Respiratory Syncytial Virus-Like Particles Protect the Lower as well as the Upper Respiratory Tract. *PLoS One*. Jul. 14, 2015;10(7):e0130755. doi: 10.1371/journal.pone.0130755. cited by applicant

Walton et al., Spontaneous CD8 T cell responses against the melanocyte differentiation antigen RAB38/NY-MEL-1 in melanoma patients. *J Immunol*. Dec. 1, 2006;177(11):8212-8. doi: 10.4049/jimmunol.177.11.8212. cited by applicant

Wang et al., CRISPR/Cas9 in Genome Editing and Beyond. *Annu Rev Biochem*. Jun. 2, 2016;85:227-64. doi: 10.1146/annurev-biochem-060815-014607. Epub Apr. 25, 2016. cited by applicant

Wang et al., Recognition of an antigenic peptide derived from tyrosinase-related protein-2 by CTL in the context of HLA-A31 and -A33. *J Immunol*. Jan. 15, 1998;160(2):890-7. cited by applicant

Wang et al., Recognition of breast cancer cells by CD8+ cytotoxic T-cell clones specific for NY-BR-1. *Cancer Res*. Jul. 1, 2006;66(13):6826-33. doi: 10.1158/0008-5472.CAN-05-3529. cited by applicant

Wang et al., Utilization of an alternative open reading frame of a normal gene in generating a novel human cancer antigen. *J Exp Med*. Mar. 1, 1996;183(3):1131-40. doi: 10.1084/jem.183.3.1131. cited by applicant

Wang et al., Virus-like particles for the prevention of human papillomavirus-associated malignancies. *Expert Rev Vaccines*. Feb. 2013;12(2):129-41. doi: 10.1586/erv.12.151. Author Manuscript, 22 pages. cited by applicant

Wölfel et al., Two tyrosinase nonapeptides recognized on HLA-A2 melanomas by autologous cytolytic T lymphocytes. *Eur J Immunol*. Mar. 1994;24(3):759-64. doi: 10.1002/eji.1830240340. cited by applicant

Yang et al., HIV-1 virus-like particles produced by stably transfected *Drosophila* S2 cells: a desirable vaccine component. *J Virol*. Jul. 2012;86(14):7662-76. doi: 10.1128/JVI.07164-11. Epub May 2, 2012. cited by applicant

Yee et al., A general method for the generation of high-titer, pantropic retroviral vectors: highly efficient infection of primary hepatocytes. *Proc Natl Acad Sci U S A*. Sep. 27, 1994;91(20):9564-8. doi: 10.1073/pnas.91.20.9564. cited by applicant

Yu et al., Poor immunogenicity of a self/tumor antigen derives from peptide-MHC-I instability and is independent of tolerance. *J Clin Invest*. Aug. 2004;114(4):551-9. doi: 10.1172/JCI21695. cited by applicant

Zarour et al., Melan-A/MART-1(51-73) represents an immunogenic HLA-DR4-restricted epitope recognized by melanoma-reactive CD4(+) T cells. *Proc Natl Acad Sci U S A*. Jan. 4, 2000;97(1):400-5. doi: 10.1073/pnas.97.1.400. cited by applicant

Zeltins, A., Construction and characterization of virus-like particles: a review. *Mol Biotechnol*. Jan. 2013;53(1):92-107. doi: 10.1007/s12033-012-9598-4. cited by applicant

Zhang et al., Cell-specific targeting of lentiviral vectors mediated by fusion proteins derived from Sindbis virus, vesicular stomatitis virus, or avian sarcoma/leukosis virus. *Retrovirology*. Jan. 25, 2010;7:3. doi: 10.1186/1742-4690-7-3. cited by applicant

Zhang et al., CRISPR-Cpf1 correction of muscular dystrophy mutations in human cardiomyocytes and mice. *Sci Adv*. Apr. 12, 2017;3(4):e1602814. doi: 10.1126/sciadv.1602814. cited by applicant

Zhao et al., Study on p21 gene knock out in G401 cell line by using CRISPR/Cas9 system. *Tianjin Med J*. Oct. 2016;44(10):1190-1194. cited by applicant

---

*Primary Examiner:* Leith; Nancy J

*Attorney, Agent or Firm:* Wolf, Greenfield & Sacks, P.C.

---

## Background/Summary

RELATED APPLICATIONS (1) The present application is a national stage filing under 35 U.S.C. § 371 of international PCT application, PCT/US2018/021880, filed Mar. 9, 2018, which claims priority under 35 U.S.C. § 119(e) to U.S. provisional patent application, U.S. Ser. No. 62/469,219, filed Mar. 9, 2017, the entire contents of each of which are incorporated herein by reference.

## REFERENCE TO A SEQUENCE LISTING SUBMITTED AS A TEXT FILE VIA EFS-WEB

(1) The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Oct. 27, 2023, is named H082470241US01-SUBSEQ-AZW and is 3,827,234 bytes in size.

## BACKGROUND OF THE INVENTION

(2) Tumor-specific immune responses may be elicited by peptides generated from proteins expressed in tumor cells or on tumor cell surface (e.g., tumor-specific antigens). Native peptides derived from tumor-specific antigens are tolerated as “self” by the immune system and do not elicit strong immune response against the tumor-specific antigen. Altered versions of the native peptides derived from tumor-specific antigens (e.g., heteroclitic peptides or cryptic peptides) may be engineered to elicit potent immune reactions through the MHC-I and MHC-II antigen presentation pathways, which also produce cross-reactive responses towards the native tumor-specific antigen sequences.

(3) It is well established that the immune system can function to kill tumor cells, including both primary and metastatic cancer cells. Indeed, evidence that the immune system recognizes the presence of neoplastic cancerous cells is supported by the existence of infiltrating lymphocytes in tumor tissues (Haskill et al., 1978, *Contemp. Top. Immunobiol.* 8: 107-170; Vose and Moore, 1985, *Semin. Hematol.* 22: 27-40). Yet, for reasons that are not completely clear, despite the presence of immune cells, tumors often prevail and not only survive but metastasize to distant sites with unrestricted growth. Recent advances in the understanding of T cell activation and recognition of target cells have begun to permit some progress in development of T cell mediated cancer immunotherapy (Schwartz, 1992, *Cell* 71: 1065-1068; Pardoll, 1992, *Curr. Opin. Immunol.* 4: 619-623).

## SUMMARY OF THE INVENTION

(4) Described herein are systems, methods, compositions, and kits for producing immunogenic peptides derived from tumor specific antigens (e.g., heteroclitic epitopes or cryptic epitopes) that may be used as cancer vaccines in vivo or ex vivo. Targeted mutations are introduced into tumor-specific antigens using gene editing agents, e.g., a nucleobase editor comprising a programmable DNA binding domain (e.g., catalytically-inactive Cas9 or a Cas9 nickase) fused to a cytosine deaminase, to generate altered versions of peptides arising from the tumor-specific antigens (heteroclitic epitopes) or peptides arising from normally untranslated regions of the tumor-specific antigen genes (cryptic peptides). The heteroclitic peptides or cryptic peptides may be generated in vivo in a subject (e.g., a subject who has cancer) and presented to the adaptive immune system via the MHC class I or MHC class II pathway, which in turn induces a strong adaptive immune response, e.g., T cell response and B cell response. Such an adaptive immune response is antigen specific and is effective in reducing tumor growth and preventing metastasis.

(5) Some aspects of the present disclosure provide methods of eliciting a tumor-specific immune response in a subject in need thereof, the methods including administering to the subject a therapeutically effective amount of a composition comprising: (i) a fusion protein comprising (a) a guide nucleotide sequence-programmable DNA-binding protein domain; and (b) a cytosine deaminase domain; and (ii) a guide nucleotide sequence, wherein the guide nucleotide sequence of (ii) targets the fusion protein of (i) to a polynucleotide encoding a tumor-specific antigen in a tumor cell, wherein the fusion protein changes a target cytosine (C) base to a thymine (T) base via deamination.

(6) In some embodiments, the polynucleotide comprises a coding strand and a complementary strand. In some embodiments, the polynucleotide comprises a coding region and a non-coding region. In some embodiments, the polynucleotide encoding the tumor-specific antigen is located in the genome of the tumor cell. In some embodiments, deamination of the target C base results in a C-G base-pair to thymine-adenine (T-A) base-pair change.

(7) In some embodiments, the guide nucleotide sequence-programmable DNA binding protein domain is selected from the group consisting of: nuclease inactive Cas9 (dCas9) domains, nuclease inactive Cpf1 domains, nuclease inactive Argonaute domains, and variants thereof.

(8) In some embodiments, the guide nucleotide sequence-programmable DNA-binding protein domain is a nuclease inactive Cas9 (dCas9) domain. In some embodiments, the amino acid sequence of the dCas9 domain comprises mutations corresponding to a D10A and/or H840A mutation in SEQ ID NO: 1. In some embodiments, the amino acid sequence of the dCas9 domain comprises a mutation corresponding to a D10A mutation in SEQ ID NO: 1, and wherein the dCas9 domain comprises a histidine at the position corresponding to amino acid 840 of SEQ ID NO: 1.

(9) In some embodiments, the guide nucleotide sequence-programmable DNA-binding protein domain comprises a nuclease inactive Cpf1 (dCpf1) domain. In some embodiments, the dCpf1 domain is from a species of *Acidaminococcus* or *Lachnospiraceae*. In some embodiments, the guide nucleotide sequence-programmable DNA-binding protein domain comprises a nuclease inactive Argonaute (dAgo) domain. In some embodiments, the (dAgo) domain is from *Natronobacterium gregoryi* (dNgAgo).

(10) In some embodiments, the cytosine deaminase domain comprises an apolipoprotein B mRNA-editing complex (APOBEC) family deaminase. In some embodiments, the cytosine deaminase is selected from the group consisting of APOBEC1, APOBEC2, APOBEC3A, APOBEC3B, APOBEC3C, APOBEC3D, APOBEC3F, APOBEC3G deaminase, APOBEC3H deaminase, APOBEC4 deaminase, and activation-induced deaminase (AID). In some embodiments, the cytosine deaminase comprises an amino acid sequence of any of SEQ ID NOS: 27-292, 303, and 1072-1083.

(11) In some embodiments, the fusion protein of (a) further comprises a uracil glycosylase inhibitor (UGI) domain. In some embodiments, the cytosine deaminase domain is fused to the N-terminus of the guide nucleotide sequence-programmable DNA-binding protein domain. In some embodiments, the UGI domain is fused to the C-terminus of the

guide nucleotide sequence-programmable DNA-binding protein domain. In some embodiments, the cytosine deaminase and the guide nucleotide sequence-programmable DNA-binding protein domain is fused via an optional linker. In some embodiments, the UGI domain is fused to the guide nucleotide sequence-programmable DNA-binding protein domain via an optional linker.

(12) In some embodiments, the fusion protein comprises the structure NH.sub.2-[cytosine deaminase domain]-[optional linker sequence]-[guide nucleotide sequence-programmable DNA-binding protein domain]-[optional linker sequence]-[UGI domain]-COOH.

(13) In some embodiments, the optional linker comprises (GGGS).sub.n, (SEQ ID NO: 337) (GGGGS).sub.n (SEQ ID NO: 308), (G).sub.n (SEQ ID NO: 783), (EAAAK).sub.n (SEQ ID NO: 309), (GGS).sub.n (SEQ ID NO: 784), SGSETPGTSESATPES (SEQ ID NO: 310), or (XP).sub.n (SEQ ID NO: 785) motif, or a combination of any of these, wherein n is independently an integer between 1 and 30 and wherein X is any amino acid. In some embodiments, the linker comprises the amino acid sequence of SGSETPGTSESATPES (SEQ ID NO: 310). In some embodiments, the linker is (GGS).sub.n (SEQ ID NO: 784), and wherein n is 1, 3, or 7.

(14) In some embodiments, the fusion protein comprises the amino acid sequence of any one of SEQ ID NOs: 293-302, 1071, and 1084.

(15) In some embodiments, the tumor specific antigen is selected from the group consisting of: CEA; gp100; Pmel17; mammaglobin-A; Melan-A; MART-1; NY-BR-1; ERBB2; OA1; PAP; PSA; RAB38; NY-MEL-1; TRP-1; gp75; TRP-2; tyrosinase; WT1; CD33; BAGE-1; D393-CD20n; Cyclin-A1; GAGE-1,2,8; GAGE-3,4,5,6,7; GnTVf; HERV-K-MEL; KK-LC-1; KM-HN-1; LAGE-1; LY6K; MAGE-A1; MAGE-A2; MAGE-A3; MAGE-A4; MAGE-A6; MAGE-A9; MAGE-A10; MAGE-A12m; MAGE-C1; MAGE-C2; mucin; NA88-A; NY-ESO-1; LAGE-2; SAGE; Sp17; SSX-2; SSX-4; survivin; BIRC5; TAG-1; TAG-2; TRAG-3; TRP2-INT2g; XAGE-1b; GAGED2a; BCR-ABL (b3a2); adipophilin; AIM-2; ALDH1A1; BCLX(L); BING-4; CALCA; CD45; CD274; CPSE; cyclin D1; DKK1; ENAH (hMena); EpCAM; EphA3; EZH2; FGF5; glypican-3; G250; MN; CAIX; HER-2; neu; HLA-DOB; Hepsin; IDO1; IGF2B3; IL13Ralpha2; Intestinal carboxyl esterase; alpha-foetoprotein; Kallikrein 4; KIF20A; Lengsin; M-CSF; MCSP; mdm-2; Meloe; Midkine; MMP-2; MMP-7; MUC1; MUC5AC; p53; PAX5; PBF; PRAME; PSMA; RAGE-1; RGS5; RhoC; RNF43; RU2AS; secernin 1; SOX10; STEAP1; Telomerase; TPBG; and VEGF.

(16) In some embodiments, the target C base is in a target codon located in a coding region of the polynucleotide encoding the tumor-specific antigen. In some embodiments, the target codon is any one of the target codons in Tables 4 and 8.

(17) In some embodiments, the target codon is converted to a modified codon selected from any one of the modified codons in Table 4. In some embodiments, the target C base is located in a non-coding region of the polynucleotide encoding the tumor specific antigen. In some embodiments, the target C base is located in an intron in the polynucleotide encoding the tumor specific antigen.

(18) In some embodiments, the methods described herein further comprising generating an immunogenic peptide from the tumor-specific antigen. In some embodiments, the immunogenic peptide is a heteroclitic epitope. In some embodiments, the heteroclitic epitope is at least 2 fold, at least 5 fold, at least 10 fold, at least 20 fold, at least 30 fold, at least 40 fold, at least 50 fold, at least 60 fold, at least 70 fold, at least 80 fold, at least 90 fold, at least 100 fold, or more immunogenic than a native epitope from the tumor specific antigen. In some embodiments, the immunogenic peptide is a cryptic epitope. In some embodiments, the cryptic epitope is at least 2 fold, at least 5 fold, at least 10 fold, at least 20 fold, at least 30 fold, at least 40 fold, at least 50 fold, at least 60 fold, at least 70 fold, at least 80 fold, at least 90 fold, at least 100 fold, or more immunogenic than a native epitope from the tumor specific antigen.

(19) In some embodiments, the immunogenic peptide is displayed on the surface of the tumor cell via the MHC class I antigen presentation pathway. In some embodiments, the immunogenic peptide is displayed on the surface of an antigen presenting cell (APC) via the MHC class II antigen presentation pathway.

(20) In some embodiments, the method is carried out in vivo. In some embodiments, the method is carried out ex vivo.

(21) In some embodiments, the APC is selected from the group consisting of: tumor cells, dendritic cells, mononuclear phagocytes, thymic epithelial cells, and B cells.

(22) In some embodiments, the immunogenic peptide elicits adaptive immune response against the tumor-specific antigen. In some embodiments, the adaptive immune response comprises promoting the maturation of dendritic cells, activation of CD4+ T lymphocytes, activation of CD8+ T lymphocytes, activation and maturation of B lymphocytes, and/or production of tumor antigen-specific antibodies. In some embodiments, the adaptive immune response kills tumor cells, reduces tumor size, and/or prevents metastasis.

(23) In some embodiments, the guide nucleotide sequence is an RNA. In some embodiments, the RNA is chemically modified.

(24) In some embodiments, the guide nucleotide sequence is a single strand DNA (ssDNA).

(25) In some embodiments, the tumor specific antigen is gp100. In some embodiments, the gp100 is from melanoma. In some embodiments, the deamination of the target C base in codon T210 of gp100 results in a T210I mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of IIDQVPFSV (SEQ ID NO: 786) is generated, and wherein the I at position 2 corresponds to the T210I mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 724 and 870-888.

(26) In some embodiments, the deamination of the target C base in codon A288 of gp100 results in a A288V mutation. In



some embodiments, a heteroclitic epitope comprising the amino acid sequence of YLEPGPVTV (SEQ ID NO: 818) is generated, and wherein the V at position 7 corresponds to the A288V mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 725 and 889.

(27) In some embodiments, the deamination of the target C base in codon T155 of gp100 results in a T155I mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of KIWGQYWQV (SEQ ID NO: 787) is generated, and wherein the I at position 2 corresponds to the T155I mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 726 and 890-892.

(28) In some embodiments, the tumor specific antigen is melanoma antigen recognized by T cells 1 (MART-1). In some embodiments, the MART-1 antigen is from melanoma. In some embodiments, the deamination of the target C base in codon A27 of MART-1 results in a A27V mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of EVAGIGILTV (SEQ ID NO: 819) is generated, and wherein the V at position 2 corresponds to the A27V mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 727 and 893-896.

(29) In some embodiments, the tumor specific antigen is cancer/testis antigen 1B (NY-ESO-1). In some embodiments, the NY-ESO-1 antigen is from melanoma or breast cancer. In some embodiments, the deamination of the target C base in codon C165 of NY-ESO-1 results in a C165Y mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of SLLMWITQY (SEQ ID NO: 788) is generated, and wherein the C at position 9 corresponds to the C165Y mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 728 and 897.

(30) In some embodiments, the tumor specific antigen is Tyrosinase (TYR). In some embodiments, the TYR antigen is from melanoma. In some embodiments, the deamination of the target C base in codon T373 of TYR results in a T373I mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of YMNGIMSQV (SEQ ID NO: 789) is generated, and wherein the I at position 5 corresponds to the T373I mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 729 and 898-901.

(31) In some embodiments, the tumor specific antigen is tyrosinase-related protein 1 (TyRP1). In some embodiments, the TyRP1 antigen is from melanoma. In some embodiments, the deamination of the target C base in codon C244 of TyRP1 results in a C244Y mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of DAEKYDICTDEY (SEQ ID NO: 790) is generated, and wherein the Y at position 5 corresponds to the C244Y mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 730 and 902.

(32) In some embodiments, the tumor specific antigen is Survivin. In some embodiments, the Survivin is from melanoma, breast cancer, or leukemia. In some embodiments, the deamination of the target C base in codon T97 of Survivin results in a T97I mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of ELILGEFLKL (SEQ ID NO: 791) is generated, and wherein the I at position 3 corresponds to the T97I mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 731 and 903.

(33) In some embodiments, the tumor specific antigen is telomerase reverse transcriptase (hTERT). In some embodiments, the hTERT is from breast cancer. In some embodiments, the deamination of the target C base in codon M549 of hTERT results in a M549I mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of ILAKFLHWLI (SEQ ID NO: 792) is generated, and wherein the I at position 10 corresponds to the M549I mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 735 and 916-920.

(34) In some embodiments, the tumor specific antigen is human epidermal growth factor receptor 2 (HER2). In some embodiments, the HER2 is from breast cancer. In some embodiments, the deamination of the target C base in codon V658 of HER2 results in a V658M mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of AMVGILLVVV (SEQ ID NO: 793) is generated, and wherein the M at position 2 corresponds to the V658M mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 732 and 904-909.

(35) In some embodiments, the deamination of the target C base in codon T912 of HER2 results in a T912I mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of IIWELMTFGA (SEQ ID NO: 794) is generated, and wherein the V at position 2 corresponds to the T912I mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 733 and 910-912.

(36) In some embodiments, the deamination of the target C base in codon A920 of HER2 results in a A920V mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of ITWELMTFGV (SEQ ID NO: 795) is generated, and wherein the V at position 10 corresponds to the A920V mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 734 and 913-915.

(37) In some embodiments, the tumor specific antigen is CD33. In some embodiments, the CD33 is from leukemia. In some embodiments, the deamination of the target C base in codon A65 of CD33 results in a A65V mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of VIISGDSPV (SEQ ID NO: 796) is generated,

and wherein the V at position 1 corresponds to the A65V mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 740 and 930-932.

(38) In some embodiments, the tumor specific antigen is Synovial Sarcoma X Breakpoint 2 (SSX2). In some embodiments, the deamination of the target C base in codon A42 of SSX2 results in a A42V mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of KVSEKIFYV (SEQ ID NO: 797) is generated, and wherein the V at position 2 corresponds to the A42V mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 737 and 921.

(39) In some embodiments, the tumor specific antigen is Wilm's tumor 1 (WT1) protein. In some embodiments, the WT1 is from leukemia. In some embodiments, the deamination of the target C base in codon C235 of WT1 results in a C235Y mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of YMTWNQMNL (SEQ ID NO: 798) is generated, and wherein the Y at position 1 corresponds to the C235Y mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 738 and 922-925.

(40) In some embodiments, the deamination of the target C base in codon M236 of WT1 results in a M236I mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of CITWNQMNL (SEQ ID NO: 799) is generated, and wherein the I at position 2 corresponds to the M236I mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 739 and 926-929.

(41) In some embodiments, the tumor specific antigen is Epithelial cell adhesion molecule precursor (EpCAM). In some embodiments, the deamination of the target C base in codon T192 of EpCAM results in a T192I mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of ILYENNVII (SEQ ID NO: 800) is generated, and wherein the I at position 9 corresponds to the T192I mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 741 and 933-934.

(42) In some embodiments, the tumor specific antigen is carcinoembryonic antigen-related cell adhesion molecules (CEA-CAM). In some embodiments, the CEA-CAM is from colorectal cancer, lung cancer, or breast cancer. In some embodiments, the deamination of the target C base in codon T314 of CEA-CAM results in a T314I mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of LLTFWNPPPI (SEQ ID NO: 801) is generated, and wherein the I at position 9 corresponds to the T314I mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 742 and 935-936.

(43) In some embodiments, the deamination of the target C base in codon T311 of CEA-CAM results in a T311I mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of RITVTTITV (SEQ ID NO: 802) is generated, and wherein the V at position 2 corresponds to the T311I mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 743 and 937-940.

(44) In some embodiments, the deamination of the target C base in codon T688 of CEA-CAM results in a T688V mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of AVVGIMIGV (SEQ ID NO: 803) is generated, and wherein the V at position 2 corresponds to the T688V mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 744 and 941-947.

(45) In some embodiments, the deamination of the target C base in codon V695 of CEA-CAM results in a V695M mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of IMIGMLVGV (SEQ ID NO: 804) is generated, and wherein the M at position 5 corresponds to the V695M mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 745 and 948-953.

(46) In some embodiments, the tumor specific antigen is melanoma-associated antigen A3 (MAGEA3). In some embodiments, the deamination of the target C base in codon H118 of MAGEA3 results in a H118Y mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of KVAELVYFL (SEQ ID NO: 805) is generated, and wherein the Y at position 7 corresponds to the H118Y mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 746 and 954.

(47) In some embodiments, the tumor specific antigen is melanoma-associated antigen (MAGE) common antigen A3, A1, A4, A2, or A12. In some embodiments, the deamination of the target C base in codon C181 of MAGE common antigen A3, A1, A4, A2, or A12 results in a C181Y mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of YLGLSYDGLL (SEQ ID NO: 806) is generated, and wherein the Y at position 1 corresponds to the C181Y mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 747-750 and 955-983.

(48) In some embodiments, the tumor specific antigen is MUC-1. In some embodiments, the deamination of the target C base in codon T93 of MUC-1 results in a T93I mutation. In some embodiments, a heteroclitic epitope comprising the amino acid sequence of AIWGQDVTSV (SEQ ID NO: 807) is generated, and wherein the I at position 2 corresponds to the T93I mutation. In some embodiments, the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 751 and 984-985.

(49) In some embodiments, the target C base is located in intron 4 of the premelanosome protein (PMEL) gene. In some embodiments, the deamination of the target C base results in a cryptic peptide comprising the amino acid sequence of

VYFFLPDHL (SEQ ID NO 808). In some embodiments, the guide nucleotide sequence comprises a nucleotide selected from the group consisting of SEQ ID NOs: 752-753 and 986-998.

(50) In some embodiments, the target C base is located on the complementary strand of open reading frame 1 (ORF1) of TYRP1 gene. In some embodiments, the target C base is located in the complementary strand of the first start codon (ATG) of ORF1 of the TYRP1 gene. In some embodiments, the deamination of the target C base results in a cryptic peptide comprising the amino acid sequence of MSLQRQFLR (SEQ ID NO: 809). In some embodiments, the guide nucleotide sequence comprises a nucleotide selected from the group consisting of SEQ ID NOs: 754 and 999-1005.

(51) In some embodiments, the target C base is located on the complementary strand of the last base of intron 2 of the mannosyl (alpha-1,6-)-glycoprotein beta-1,6-N-acetyl glucosaminyltransferase (MGAT5) gene. In some embodiments, the deamination of the target C base results in a cryptic peptide comprising the amino acid sequence of VLPDVFIRCV (SEQ ID NO: 810). In some embodiments, the cryptic peptide is translated from exon 3 of the MGAT5 gene. In some embodiments, the guide nucleotide sequence comprises a nucleotide selected from the group consisting of SEQ ID NOs: 755 and 1006-1008.

(52) In some embodiments, the target C base is located in open reading frame 1 (ORF1) of cancer/testis antigen 2 (LAGE-1) gene. In some embodiments, the target C base is located in the complementary strand of the first start codon of ORF1 of the LAGE-1 gene. In some embodiments, the deamination of the target C base results in a cryptic peptide comprising the amino acid sequence of selected from the group consisting of: MLMAQEALAF (SEQ ID NO: 811), LAAQERRVPR (SEQ ID NO: 812), APRGVRMAV (SEQ ID NO: 813), QGAMLAAQERRVPRAAEVPR (SEQ ID NO: 814), and CLSRPWPWRSWSAGSCPGMPHL (SEQ ID NO: 815). In some embodiments, the guide nucleotide sequence comprises a nucleotide selected from the group consisting of SEQ ID NOs: 756 and 1009-1014.

(53) In some embodiments, the target C base is located in intron 2 of tyrosinase-related protein 2 (TRP-2) gene. In some embodiments, the target C base is located on the complementary strand of the first base of intron 2 of the TRP-2 gene. In some embodiments, the deamination of the target C base results in a cryptic peptide comprising the amino acid sequence of EVISCKLIKR (SEQ ID NO: 816). In some embodiments, the guide nucleotide sequence comprises a nucleotide selected from the group consisting of SEQ ID NOs: 757-758 and 1015-1023.

(54) In some embodiments, the target C base is located in intron 2 of baculoviral IAP repeat containing 5 (BIRC5) gene. In some embodiments, the target C base is located on the spliceosome branch site of intron 2 of the BIRC5 gene. In some embodiments, the target C base is located in the complementary strand of the last base of intron 2 of the BIRC5 gene. In some embodiments, the deamination of the target C base results in a cryptic peptide comprising the amino acid sequence of AYACNTSTL (SEQ ID NO: 817). In some embodiments, the guide nucleotide sequence comprises a nucleotide selected from the group consisting of SEQ ID NOs: 759 and 1024-1029.

(55) In some embodiments, the target C base is located in intron 1 acceptor site of BCR/ABL fusion proteins (BCR/ABL-OOF) gene. In some embodiments, the target C base is located in intron 2 acceptor site of BCR/ABL fusion proteins (BCR/ABL-OOF) gene. In some embodiments, the deamination of the target C base results in a cryptic peptide comprising the amino acid sequence of any one of SSKALQRPV (SEQ ID NO: 603), GFKQSSKAL (SEQ ID NO: 604), and ATGFKQSSKALQRPVAS (SEQ ID NO: 605). In some embodiments, the guide nucleotide sequence comprises a nucleotide selected from the group consisting of SEQ ID NOs: 761 and 1032-1045. In some embodiments, the guide nucleotide sequence comprises a nucleotide selected from the group consisting of SEQ ID NOs: 762 and 1046-1056.

(56) In some embodiments, the methods further comprising administering to the subject a therapeutically effective amount of an immune checkpoint inhibitor. In some embodiments, the immune checkpoint inhibitor inhibits binding of CTLA-4, PD-1, PD-L1, TIM3, LAG3, B7-H3, B7-H4, BTLA, GAL9, Chk1, or A2aR to a cognate binding partner. In some embodiments, the immune checkpoint inhibitor is an antibody or a fragment thereof. In some embodiments, the antibody is selected from anti-CTLA-4 antibodies, anti-PD-1 antibodies, anti-PD-L1 antibodies, anti-TIM3 antibodies, anti-LAG3 antibodies, anti-B7-H3 antibodies, anti-B7-H4 antibodies, anti-BTLA antibodies, anti-GAL9 antibodies, anti-Chk1 antibodies, and anti-A2aR antibodies. In some embodiments, the antibody is selected from pembrolizumab, nivolumab, and ipilimumab.

(57) In some embodiments, the immune checkpoint inhibitor is a small molecule.

(58) In some embodiments, the immune checkpoint inhibitor is a recombinant protein.

(59) In some embodiments, the immune checkpoint inhibitor is a nucleic acid aptamer.

(60) In some embodiments, the immune checkpoint inhibition is performed by genome editing of a gene selected from the group consisting of: CTLA-4, PD-1, PD-L1, TIM3, LAG3, B7-H3, B7-H4, BTLA, GAL9, Chk1, or A2aR.

(61) Other aspects of the present disclosure provide methods of treating cancer, the methods including administering to a subject in need thereof a therapeutically effective amount of a composition comprising: (i) a fusion protein comprising (a) a guide nucleotide sequence-programmable DNA-binding protein domain; and (b) a cytosine deaminase domain; and (ii) a guide nucleotide sequence; wherein the fusion protein of (i) and the guide nucleotide sequence of (ii) enters a tumor cell, and wherein the guide nucleotide sequence targets the fusion protein of (i) to a polynucleotide encoding a tumor-specific antigen, wherein the fusion protein changes a target cytosine (C) residue to a (T) residue in the polynucleotide.

(62) In some embodiments, the methods include administering to the subject a therapeutically effective amount of an immune checkpoint inhibitor.

(63) Further provided herein are methods of inducing a tumor-specific immune response in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a composition comprising: (i) a

fusion protein comprising (a) a guide nucleotide sequence-programmable DNA-binding protein domain; and (b) a nuclease domain; and (ii) a guide nucleotide sequence; wherein the fusion protein of (i) and the guide nucleotide sequence of (ii) enters the tumor cell, and wherein the guide nucleotide sequence targets the fusion protein of (i) to a polynucleotide encoding a tumor-specific antigen, wherein the fusion protein introduces an indel in the polynucleotide. In some embodiments, the nuclease is a FokI nuclease.

(64) Further provided herein are methods of inducing a tumor-specific immune response in a subject in need thereof, the methods including administering to the subject a therapeutically effective amount of a composition comprising: (i) a guide nucleotide sequence-programmable nuclease; and (ii) a guide nucleotide sequence; wherein the fusion protein of (i) and the guide nucleotide sequence of (ii) enters the tumor cell, and wherein the guide nucleotide sequence targets the fusion protein of (i) to a polynucleotide encoding a tumor-specific antigen, wherein the guide nucleotide sequence-programmable nuclease introduces an indel in the polynucleotide.

(65) In some embodiments, the guide nucleotide sequence-programmable nuclease comprises a Cas9, a Cpf1, an Argonaute, or a variant thereof. In some embodiments, the indel causes a mutation or frame shift.

(66) Method of inducing a tumor-specific immune response in a subject in need thereof are also provided, the methods including administering to a subject in need thereof a therapeutically effective amount of a composition comprising a fusion protein comprising (a) a programmable DNA-binding protein domain; and (b) a deaminase domain; wherein the fusion protein enters the tumor cell and changes a target base in the polynucleotide via deamination.

(67) In some embodiments, the deaminase domain comprises a cytosine deaminase and the target base is a cytosine (C) base. In some embodiments, the programmable DNA-binding domain comprises a zinc finger nuclease (ZFN). In some embodiments, the programmable DNA-binding domain comprises a transcription activator-like effector (TALE).

(68) In some embodiments, the programmable DNA-binding domain is a guide nucleotide sequence-programmable DNA binding protein domain. In some embodiments, the programmable DNA-binding domain is selected from the group consisting of: nuclease-inactive Cas9 domains, nuclease inactive Cpf1 domains, nuclease inactive Argonaute domains, and variants thereof. In some embodiments, the programmable DNA-binding domain is associated with a guide nucleotide sequence. In some embodiments, the deamination of the target C base results in a C to thymine (T) change. In some embodiments, the deamination of the target C base results in a C-G base pair to thymine-adenine (T-A) change in a translated codon, resulting in the incorporation of a different amino acid in an immunogenic or heteroclitic peptide. In some embodiments, the deamination of the target C base results in a C-G basepair to thymine-adenine (T-A) change in a non-coding intron region of a gene, resulting in alternative splicing and translation of immunogenic or cryptic peptide sequences. In some embodiments, the deamination of the target C base results in a C-G basepair to thymine-adenine (T-A) change in the start (Met) codon of the open reading frame of a gene, resulting in the translation of an alternative open reading frame comprising immunogenic or cryptic peptide sequences.

(69) Other aspects of the present disclosure provide compositions comprising: (i) a fusion protein comprising (a) a guide nucleotide sequence-programmable DNA-binding protein domain; and (b) a cytosine deaminase domain; and (ii) a guide nucleotide sequence targeting the fusion protein of (i) to a polynucleotide encoding a tumor specific antigen.

(70) Yet other aspects of the present disclosure provide compositions comprising a polynucleotide encoding a fusion protein and a guide nucleotide sequence, wherein the fusion protein comprises (a) a guide nucleotide sequence-programmable DNA-binding protein domain; and (b) a cytosine deaminase domain, and wherein the guide nucleotide sequence targets the fusion protein to a polynucleotide encoding a tumor specific antigen.

(71) Yet other aspects of the present disclosure provide cancer vaccines comprising: (i) a fusion protein comprising (a) a guide nucleotide sequence-programmable DNA-binding protein domain; and (b) a cytosine deaminase domain; and (ii) a guide nucleotide sequence targeting the fusion protein of (i) to a polynucleotide encoding a tumor specific antigen.

(72) Further provided herein are cancer vaccine comprising a polynucleotide encoding a fusion protein and a guide nucleotide sequence, wherein the fusion protein comprises (a) a guide nucleotide sequence-programmable DNA-binding protein domain; and (b) a cytosine deaminase domain, and wherein the guide nucleotide sequence targets the fusion protein to a polynucleotide encoding a tumor specific antigen.

(73) Kits comprising the cancer vaccines described herein are also provided.

(74) The details of certain embodiments of the disclosure are set forth in the Detailed Description of Certain Embodiments, as described below. Other features, objects, and advantages of the disclosure will be apparent from the Definitions, Examples, Figures, and Claims.

---

## Description

### BRIEF DESCRIPTION OF THE DRAWINGS

- (1) The accompanying drawings, which constitute a part of this specification, illustrate several embodiments of the disclosure and together with the description, serve to explain the principles of the disclosure.
- (2) FIG. 1 shows strategies to engineer heteroclitic and cryptic epitopes using genome base editing.
- (3) FIG. 2A shows strategies to introduce immunogenic heteroclitic epitopes by editing conservative anchor residues to match the binding preference of the main HLA allele supertypes. The example shows a base-editing reaction that turns an alanine residue at anchor position 9 of a weakly immunogenic peptide epitope into a preferred valine residue for binding HLA-A2.

(4) FIG. 2B shows amino-residue binding preference and population coverage of the main HLA allele supertypes (MHC-I pathway). The peptides in FIG. 2B are as follows: HLA A1, A2, A3, and A24 (SEQ ID NOs: 878-881) and HLA B7, B27, B44, B58, and B62 (SEQ ID NOs: 882-886).

(5) FIG. 3 shows a proposed mechanism for anti-cancer vaccination by heteroclitic/cryptic epitopes introduced by genome base-editing reactions programmed by guide-RNAs. The edited tumor cells produce heteroclitic and cryptic epitopes in cancer-specific genes, which chemotactically attract immature dendritic cells (DCs) (1a), inducing DC maturation (1b). Edited tumor cells produce apoptotic bodies (2a) that are taken up by DCs (2b), contributing to maturation of DCs (1b) and B cells (2b'). Mature DCs activate CD4<sup>+</sup> T lymphocytes (1c) and CD8<sup>+</sup> T lymphocytes (1c'). Activated CD4<sup>+</sup> T lymphocytes further stimulate B-lymphocyte activation (1d) and provide IL-2 for CD8<sup>+</sup> T lymphocytes (1d'). B lymphocytes produce TAA-specific antibodies to cell-surface proteins that result in antibody-dependent cell-mediated cytotoxicity or complement-mediated tumor cell death (1e). Activated CD8<sup>+</sup> T lymphocytes then kill tumor cells via recognition of MHC class I molecules in association with TAA epitopes (1e').

(6) FIG. 4 shows comparison of cancer lineages that display high frequency of mutagenesis, which may harbor non-synonymous hitchhiker mutations and “neo-epitopes”.

#### DEFINITIONS

(7) As used herein and in the claims, the singular forms “a,” “an,” and “the” include the singular and the plural reference unless the context clearly indicates otherwise. Thus, for example, a reference to “an agent” includes a single agent and a plurality of such agents.

(8) An “immunogenic peptide” or “antigenic peptide” is a peptide or epitope that can be recognized by the immune system and elicit an immune response. Immunogenic peptides or antigenic peptide may comprise a motif such that the peptide will bind an MHC molecule and induce a T cell response, or can be recognized by the B cell receptor on the B cell to induce antibody production. These terms are used interchangeably herein.

(9) An “immunogenic epitope” or “antigenic epitope” refers to a part of an antigen is recognized by the immune system, e.g., by antibodies, B cells, or T cells. In some embodiments, the epitope is the specific piece of the antigen to which an antibody binds. Although epitopes are usually non-self proteins, sequences derived from the host can, in some instances, be recognized.

(10) “Immune response” is how your body recognizes and defends itself against bacteria, viruses, and substances that appear foreign and harmful to the body. In its general form, the immune response begins with the sensitization of helper (TH, CD4<sup>+</sup>) and cytotoxic (CD8<sup>+</sup>) T cell subsets through their interaction with antigen presenting cells (APC) that express major histocompatibility (MHC)-class I or class II molecules associated with antigenic fragments (i.e., specific amino acid sequences derived from the antigen which bind to MHC I and/or MHC II for presentation on the cell surface). The sensitized or primed CD4<sup>+</sup> T cells produce lymphokines that participate in the activation of B cells as well as various T cell subsets. The sensitized CD8<sup>+</sup> T cells increase in numbers in response to lymphokines and are capable of destroying any cells that express the specific antigenic fragments associated with matching MHC-encoded class I molecules. Thus, in the course of a cancerous tumor, CTL eradicate cells expressing cancer associated or cancer specific antigens, thereby limiting the progression of tumor spread and disease development.

(11) The “adaptive immune system,” also known as the acquired immune system, is a subsystem of the overall immune system that is composed of highly specialized, systemic cells and processes that eliminate or prevent pathogen growth. The adaptive immune system is one of the two main immunity strategies found in vertebrates (the other being the innate immune system). Adaptive immunity creates immunological memory after an initial response to a specific pathogen, and leads to an enhanced response to subsequent encounters with that pathogen. This process of acquired immunity is the basis of vaccination. Like the innate system, the adaptive system includes both humoral immunity components and cell-mediated immunity components.

(12) Unlike the innate immune system, the adaptive immune system is highly specific to a particular pathogen or antigen. Adaptive immunity can also provide long-lasting protection. The adaptive system response destroys invading pathogens and any toxic molecules they produce. In accordance with the present disclosure, the adaptive immune system response destroys tumor or cancer cells. Sometimes the adaptive system is unable to distinguish harmful from harmless foreign molecules. The cells that carry out the adaptive immune response are white blood cells known as lymphocytes. Two main broad classes-antibody responses and cell mediated immune response—are also carried by two different lymphocytes (B cells and T cells). In antibody responses, B cells are activated to secrete antibodies, which are proteins also known as immunoglobulins. Antibodies travel through the bloodstream and bind to the foreign antigen causing it to inactivate, which does not allow the antigen to bind to the host.

(13) In adaptive immunity, pathogen-specific receptors are “acquired” during the lifetime of the organism (whereas in innate immunity pathogen-specific receptors are already encoded in the germline). The acquired response is called “adaptive” because it prepares the body's immune system for future challenges (though it can actually also be maladaptive when it results in autoimmunity).

(14) The immune system is highly adaptable because of somatic hypermutation (a process of accelerated somatic mutations), and V(D)J recombination (an irreversible genetic recombination of antigen receptor gene segments). This mechanism allows a small number of genes to generate a vast number of different antigen receptors, which are then uniquely expressed on each individual lymphocyte. Since the gene rearrangement leads to an irreversible change in the DNA of each cell, all progeny (offspring) of that cell inherit genes that encode the same receptor specificity, including the

memory B cells and memory T cells that are the keys to long-lived specific immunity.

(15) A “T cell” or “T lymphocyte” is a type of lymphocyte (a subtype of white blood cell) that plays a central role in cell-mediated immunity. T cells can be distinguished from other lymphocytes, such as B cells and natural killer cells, by the presence of a T-cell receptor on the cell surface. They are called T cells because they mature in the thymus from thymocytes. The several subsets of T cells each have a distinct function. The majority of human T cells rearrange their alpha and beta chains on the cell receptor and are termed alpha beta T cells ( $\alpha\beta$  T cells) and are part of the adaptive immune system. Specialized gamma delta T cells, (a small minority of T cells in the human body, more frequent in ruminants), have invariant T cell receptors with limited diversity, that can effectively present antigens to other T cells and are considered to be part of the innate immune system. Effector T cell broadly includes various T cell types that actively respond to a stimulus, such as co-stimulation. This includes helper, killer, regulatory, and potentially other T cell types. One skilled in the art is familiar with different types of T cells and their respective roles in adaptive immune response.

(16) A “human leukocyte antigen (HLA) system” is a gene complex encoding the major histocompatibility complex (MHC) proteins in humans. These cell-surface proteins are responsible for the regulation of the immune system in humans. The HLA gene complex resides on a 3 Mbp stretch within chromosome 6p21. HLA genes are highly polymorphic, which means that they have many different alleles, allowing them to fine-tune the adaptive immune system. The proteins encoded by certain genes are also known as antigens, as a result of their historic discovery as factors in organ transplants. Different classes have different functions:

(17) HLAs encoding major histocompatibility complex (MHC) class I MHC class I (A, B, and C) molecules, which present peptides from inside the cell. “Major histocompatibility complex (MHC) class I” or “MHC class I” molecules are found on the cell surface of all nucleated cells in the body. Their function is to display peptide fragments of antigens from within the cell to cytotoxic T cells; this will trigger an immediate response from the immune system against a particular non-self antigen displayed with the help of an MHC class I protein. Because MHC class I molecules present peptides derived from cytosolic proteins, the pathway of MHC class I presentation is often called cytosolic or endogenous pathway.

(18) Class I MHC molecules bind peptides generated mainly from degradation of cytosolic proteins by the proteasome. The MHC I peptide complex is then inserted via endoplasmic reticulum into the external plasma membrane of the cell. The epitope peptide is bound on extracellular parts of the class I MHC molecule. Thus, the function of the class I MHC is to display intracellular proteins to cytotoxic T cells (CTLs). However, class I MHC can also present peptides generated from exogenous proteins, in a process known as cross-presentation.

(19) A normal cell will display peptides from normal cellular protein turnover on its class I MHC, and CTLs will not be activated in response to them due to central and peripheral tolerance mechanisms. When a cell expresses foreign proteins, such as after viral infection, a fraction of the class I MHC will display these peptides on the cell surface. Consequently, CTLs specific for the MHC:peptide complex will recognize and kill presenting cells. Alternatively, class I MHC itself can serve as an inhibitory ligand for natural killer cells (NKs). Reduction in the normal levels of surface class I MHC, a mechanism employed by some viruses during immune evasion or in certain tumors, will activate NK cell killing. Antigens or antigenic epitopes presented by MHC class II molecules are recognized by cytotoxic T cells.

(20) HLAs encoding MHC class II (DP, DM, DOA, DOB, DQ, and DR) molecules, which present antigens from outside of the cell to T-lymphocytes. “Major histocompatibility complex class II” or “MHC class II” molecules are a family of molecules normally found only on antigen-presenting cells such as dendritic cells, mononuclear phagocytes, some endothelial cells, thymic epithelial cells, and B cells. The antigens presented by class II peptides are usually derived from extracellular proteins (not cytosolic as in class I); hence, the MHC class II-dependent pathway of antigen presentation is called the endocytic or exogenous pathway. Loading of a MHC class II molecule occurs by phagocytosis; extracellular proteins are endocytosed, digested in lysosomes, and the resulting epitopic peptide fragments are loaded onto MHC class II molecules prior to their migration to the cell surface. Antigens or antigenic epitopes presented by MHC class II molecules are recognized by T helper cells and stimulate the multiplication of T-helper cells, which in turn stimulate antibody-producing B-cells to produce antibodies to that specific antigen. Self-antigens are suppressed by regulatory T cells.

(21) An “antigen-presenting cell (APC)” is a cell that displays antigen complexed with major histocompatibility complexes (MHCs) on their surfaces; this process is known as antigen presentation. T cells may recognize these complexes using their T cell receptors (TCRs). These cells process antigens and present them to T-cells. Antigen-presenting cells fall into two categories: professional and non-professional. Those that express MHC class II molecules along with co-stimulatory molecules and pattern recognition receptors are often called professional antigen-presenting cells. The non-professional APCs express MHC class I molecules.

(22) Professional APCs specialize in presenting antigen to T cells. They are very efficient at internalizing antigens, either by phagocytosis (macrophages and dendritic cells) or by receptor-mediated endocytosis (B cells), processing the antigen into peptide fragments and then displaying those peptides, bound to a class II MHC molecule, on their membrane.[1] The T cell recognizes and interacts with the antigen-class II MHC molecule complex on the membrane of the antigen-presenting cell. An additional co-stimulatory signal is then produced by the antigen-presenting cell, leading to activation of the T cell. The expression of co-stimulatory molecules and MHC class II are defining features of professional APCs.

(23) Almost all cell types can serve as a non-professional APC. They are found in a variety of tissue types. Professional antigen-presenting cells, including dendritic cells, mononuclear phagocytes, thymic epithelial cells, and B cells, present

foreign antigens to helper T cells, while other cell types can present antigens originating inside the cell to cytotoxic T cells. In addition to the MHC family of proteins, antigen presentation relies on other specialized signaling molecules on the surfaces of both APCs and T cells.

(24) A “B lymphocyte” or “B cell” is a type of white blood cell of the lymphocyte subtype. B cells function in the humoral immunity component of the adaptive immune system by secreting antibodies. Additionally, B cells present antigen (they are also classified as professional antigen-presenting cells (APCs)) and secrete cytokines. In mammals, B cells mature in the bone marrow, which is at the core of most bones. B cells express B cell receptors (BCRs) on their cell membrane. BCRs allow the B cell to bind a specific antigen, against which it will initiate an antibody response.

(25) “Cancer immunotherapy” refers to a type of cancer treatment designed to boost the body's natural defenses to fight the cancer. It uses substances either made by the body or in a laboratory to improve or restore immune system function.

(26) “Tumor specific antigen (TSA)” or “tumor associated antigen (TAA)” refers to a protein that is specifically expressed or upregulated in cells of the respective tumor, as compared to non-cancerous cells of the same origin. A tumor specific antigen, or epitopes derived therefrom, can be recognized by the immune system to induce an immune response. Herein, the terms “tumor associated antigen” and “tumor specific antigen” are used interchangeably. The tumor specific antigen may be from all protein classes, e.g., enzymes, receptors, transcription factors, etc.

(27) A “heteroclitic epitope” or “heteroclitic analog” refers to an altered version of an endogenous peptide sequence (i.e., an analog) engineered to elicit potent immune reactions. Heteroclitic epitopes have increased stimulatory capacity or potency for a specific T cell, as measured by increased responses to a given dose, or by a requirement of lesser amounts to achieve the same response and therefore provide benefit as vaccine components since these epitopes induce T cell responses stronger than those induced by the native epitope.

(28) A “self-antigen” refers to an antigen that originates from within the body. The immune system usually does not react to self-antigens under normal homeostatic conditions. Epitopes from self-antigens (i.e., self-epitopes) are found in high concentration on the surface of Antigen-presenting cells (APC's) in association with its major histocompatibility complex (MHC) are known as dominant epitopes. These are stimulants of negative selection mechanisms to remove potentially self-destructing autoreactive T cells. Their “self” antigens are displayed to a developing T-cell and signal those “self-reactive” T-cells to die via programmed cell death (apoptosis) and thereby deletion from the T cell repertoire, preventing autoimmunity.

(29) A “cryptic epitope” refers to an epitope derived from a self-antigen that does not necessarily undergo antigen processing/presentation and are ‘hidden’ from immune recognition. Cryptic epitopes usually appear in very low concentration on APC and do not delete auto-reactive T cells. Cryptic epitopes are not presented for recognition by T cells unless they are produced in unusually large concentrations or unless they are freed from the configuration of their native antigen. Cryptic epitopes derived from tumor-specific antigens may be used to break the tolerance of T cells to the tumor and induce potent immune response against the tumor. Such principles have been described in Pardoll, et al., *PNAS*, Vol. 96, pp. 5340-5342 (1999), the entire contents of which are incorporated herein by reference.

(30) A “neoepitope” refers to an antigenic epitope generated via random somatic mutations occurring in tumor cells. Neoepitopes are usually derived from individually specific tumor antigens or unique antigens and is thus specific to the lineage of tumor cells it is derived from. Neoepitopes are regarded in the art to be responsible for the immunogenicity of tumors ((Srivastava et al., 1993, Duan et al., 2009; van der Bruggen et al., 2013), and mathematic modeling has predicted the existence of tens to hundreds of neoepitopes in individual human tumors (Srivastava 2009). The recent revolution in high-throughput DNA sequencing and accompanying bioinformatics approaches has finally made it possible to actually identify the individually specific neoepitopes in individual cancers.

(31) “Cancer vaccine,” as used herein, refers to a composition that induces tumor-specific immunoresponse against a tumor or a tumor-specific antigen. Such immunoresponse is effective in inhibiting tumor growth and/or preventing reoccurrence of tumor.

(32) An “intron” refers to any nucleotide sequence within a gene that is removed by RNA splicing during maturation of the final RNA product. The term intron refers to both the DNA sequence within a gene and the corresponding sequence in RNA transcripts. Sequences that are joined together in the final mature RNA after RNA splicing are exons. Introns are found in the genes of most organisms and many viruses, and can be located in a wide range of genes, including those that generate proteins, ribosomal RNA (rRNA), and transfer RNA (tRNA). When proteins are generated from intron-containing genes, RNA splicing takes place as part of the RNA processing pathway that follows transcription and precedes translation.

(33) An “exon” refers to any part of a gene that will become a part of the final mature RNA produced by that gene after introns have been removed by RNA splicing. The term exon refers to both the DNA sequence within a gene and to the corresponding sequence in RNA transcripts. In RNA splicing, introns are removed and exons are covalently joined to one another as part of generating the mature messenger RNA.

(34) “RNA splicing” refers to the processing of a newly synthesized messenger RNA transcript (also referred to as a primary mRNA transcript). After splicing, introns are removed and exons are joined together (ligated) for form mature mRNA molecule containing a complete open reading frame that is decoded and translated into a protein. For nuclear-encoded genes, splicing takes place within the nucleus either co-transcriptionally or immediately after transcription. The molecular mechanism of RNA splicing has been extensively described, e.g., in Pagani et al., *Nature Reviews Genetics* 5, 389-396, 2004; Clancy et al., *Nature Education* 1 (1): 31, 2011; Cheng et al., *Molecular Genetics and Genomics* 286 (5-

6): 395-414, 2011; Taggart et al., *Nature Structural & Molecular Biology* 19 (7): 719-2, 2012, the contents of each of which are incorporated herein by reference. One skilled in the art is familiar with the mechanism of RNA splicing.

(35) “Alternative splicing” refers to a regulated process during gene expression that results in a single gene coding for multiple proteins. In this process, particular exons of a gene may be included within or excluded from the final, processed messenger RNA (mRNA) produced from that gene. Consequently, the proteins translated from alternatively spliced mRNAs will contain differences in their amino acid sequence and, often, in their biological functions. Notably, alternative splicing allows the human genome to direct the synthesis of many more proteins than would be expected from its 20,000 protein-coding genes. Alternative splicing is sometimes also termed differential splicing. Alternative splicing occurs as a normal phenomenon in eukaryotes, where it greatly increases the biodiversity of proteins that can be encoded by the genome; in humans, ~95% of multi-exonic genes are alternatively spliced. There are numerous modes of alternative splicing observed, of which the most common is exon skipping. In this mode, a particular exon may be included in mRNAs under some conditions or in particular tissues, and omitted from the mRNA in others. Abnormal variations in splicing are also implicated in disease; a large proportion of human genetic disorders result from splicing variants. Abnormal splicing variants are also thought to contribute to the development of cancer, and splicing factor genes are frequently mutated in different types of cancer. The regulation of alternative splicing is also described in the art, e.g., in Douglas et al., *Annual Review of Biochemistry* 72 (1): 291-336, 2003; Pan et al., *Nature Genetics* 40 (12): 1413-1415, 2008; Martin et al., *Nature Reviews* 6 (5): 386-398, 2005; Skotheim et al., *The international journal of biochemistry & cell biology* 39 (7-8): 1432-49, 2007, the entire contents of each of which is incorporated herein by reference.

(36) A “coding frame” or “open reading frame” refers to a stretch of codons that encodes a polypeptide. Since DNA is interpreted in groups of three nucleotides (codons), a DNA strand has three distinct reading frames. The double helix of a DNA molecule has two anti-parallel strands so, with the two strands having three reading frames each, there are six possible frame translations. A functional protein may be produced when translation proceeds in the correct coding frame. An insertion or a deletion of one or two bases in the open reading frame causes a shift in the coding frame that is also referred to as a “frameshift mutation.” A frameshift mutation typical results in premature translation termination and/or truncated or non-functional protein.

(37) The term “proteome” refers to the entire set of proteins expressed by a genome, cell, tissue, or organism at a certain time. More specifically, it is the set of expressed proteins in a given type of cell or organism, at a given time, under certain conditions. The term is a blend of proteins and genome. “Proteome-wide” refers to each and every protein in the proteome without any bias.

(38) The term “genome” refers to the genetic material of a cell or organism. It typically includes DNA (or RNA in the case of RNA viruses). The genome includes both the genes, the coding regions, the noncoding DNA, and the genomes of the mitochondria and chloroplasts. A genome does not typically include genetic material that is artificially introduced into a cell or organism, e.g., a plasmid that is transformed into a bacteria is not a part of the bacterial genome.

(39) A “programmable DNA-binding protein,” as used herein, refers to DNA binding proteins that can be programmed to navigate to any desired target nucleotide sequence within the genome. To program the DNA-binding protein to bind a desired nucleotide sequence, the DNA binding protein may be modified to change its binding specificity, e.g., zinc finger nuclease (ZFN) or transcription activator-like effector proteins (TALE). ZFNs are artificial restriction enzymes generated by fusing a zinc finger DNA-binding domain to a DNA-cleavage domain. Zinc finger domains can be engineered to target specific desired DNA sequences and this enables zinc-finger nucleases to target unique sequences within complex genomes. Transcription activator-like effector nucleases (TALEN) are restriction enzymes that can be engineered to cut specific sequences of DNA. They are made by fusing a TAL effector DNA-binding domain to a DNA cleavage domain (a nuclease which cuts DNA strands). Transcription activator-like effectors (TALEs) can be engineered to bind practically any desired DNA sequence, so when combined with a nuclease, DNA can be cut at specific locations. The restriction enzymes can be introduced into cells, for use in gene editing or for genome editing in situ, Methods of programming ZFNs and TALEs are familiar to one skilled in the art. For example, such methods are described in Maeder, et al., *Mol. Cell* 31 (2): 294-301, 2008; Carroll et al., *Genetics Society of America*, 188 (4): 773-782, 2011; Miller et al., *Nature Biotechnology* 25 (7): 778-785, 2007; Christian et al., *Genetics* 186 (2): 757-61, 2008; Li et al., *Nucleic Acids Res* 39 (1): 359-372, 2010; and Moscou et al., *Science* 326 (5959): 1501, 2009, the entire contents of each of which are incorporated herein by reference.

(40) A “guide nucleotide sequence-programmable DNA-binding protein,” as used herein, refers to a protein, a polypeptide, or a domain that is able to bind DNA, and the binding to its target DNA sequence is mediated by a guide nucleotide sequence. Thus, it is appreciated that the guide nucleotide sequence-programmable DNA-binding protein binds to a guide nucleotide sequence. The “guide nucleotide” may be a RNA molecule or a DNA molecule (e.g., a single-stranded DNA or ssDNA molecule) that is complementary to the target sequence and can guide the DNA binding protein to the target sequence. In some embodiments, the guide nucleotide sequence is an oligonucleotide sequence. As such, a guide nucleotide sequence-programmable DNA-binding protein may be a RNA-programmable DNA-binding protein (e.g., a Cas9 protein), or an ssDNA-programmable DNA-binding protein (e.g., an Argonaute protein). “Programmable” means the DNA-binding protein may be programmed to bind any DNA sequence that the guide nucleotide targets.

(41) In some embodiments, the guide nucleotide sequence exists as a single nucleotide molecule and comprises comprise two domains: (1) a domain that shares homology to a target nucleic acid (e.g., and directs binding of a guide nucleotide sequence-programmable DNA-binding protein to the target); and (2) a domain that binds a guide nucleotide sequence-



programmable DNA-binding proteins, the guide nucleotide is a guide RNA (gRNA). In some embodiments, domain (2) of the gRNA corresponds to a sequence known as a tracrRNA, and comprises a stem-loop structure. For example, in some embodiments, domain (2) is identical or homologous to a tracrRNA as provided in Jinek et al., *Science* 337:816-821(2012), the entire contents of which is incorporated herein by reference. Other examples of gRNAs (e.g., those including domain 2) can be found in U.S. Provisional Patent Application, U.S. Ser. No. 61/874,682, filed Sep. 6, 2013, entitled "Switchable Cas9 Nucleases And Uses Thereof," and U.S. Provisional Patent Application, U.S. Ser. No. 61/874,746, filed Sep. 6, 2013, entitled "Delivery System For Functional Nucleases," the entire contents of each are hereby incorporated by reference in their entirety.

(42) Because the guide nucleotide sequence hybridizes to target DNA sequence, the guide nucleotide sequence-programmable DNA-binding proteins are able to be targeted, in principle, to any sequence specified by the guide nucleotide sequence. Methods of using guide nucleotide sequence-programmable DNA-binding protein, such as Cas9, for site-specific cleavage (e.g., to modify a genome) are known in the art (see e.g., Cong, L. et al. *Science* 339, 819-823 (2013); Mali, P. et al. *Science* 339, 823-826 (2013); Hwang, W. Y. et al. *Nature biotechnology* 31, 227-229 (2013); Jinek, M. et al. *eLife* 2, e00471 (2013); Dicarlo, J. E. et al. *Nucleic acids research* (2013); Jiang, W. et al. *Nature biotechnology* 31, 233-239 (2013); the entire contents of each of which are incorporated herein by reference).

(43) It is to be understood that any DNA binding domain that is programmable by a guide nucleotide sequence may be used in accordance with the present disclosure. For example, in some embodiments, the guide nucleotide sequence-programmable DNA binding protein may be a Cas9 protein, or a variant thereof. One skilled in the art would understand that the present disclosure is not limited to the use of Cas9 as the guide nucleotide sequence-programmable DNA binding protein, but that other DNA binding proteins that adopt similar mechanism of target sequence binding may also be used.

(44) As used herein, the term "Cas9" or "Cas9 nuclease" refers to an RNA-guided nuclease comprising a Cas9 protein, a fragment, or a variant thereof. A Cas9 nuclease is also referred to sometimes as a casnI nuclease or a CRISPR (clustered regularly interspaced short palindromic repeat)-associated nuclease. CRISPR is an adaptive immune system that provides protection against mobile genetic elements (viruses, transposable elements and conjugative plasmids). CRISPR clusters contain spacers, sequences complementary to antecedent mobile elements, and target invading nucleic acids. CRISPR clusters are transcribed and processed into CRISPR RNA (crRNA). In type II CRISPR systems correct processing of pre-crRNA requires a trans-encoded small RNA (tracrRNA), endogenous ribonuclease 3 (mc) and a Cas9 protein. The tracrRNA serves as a guide for ribonuclease 3-aided processing of pre-crRNA. Subsequently, Cas9/crRNA/tracrRNA endonucleolytically cleaves linear or circular dsDNA target complementary to the spacer. The target strand not complementary to crRNA is first cut endonucleolytically, then trimmed 3'-5' exonucleolytically. In nature, DNA-binding and cleavage typically requires protein and both RNAs. However, single guide RNAs ("sgRNA", or simply "gRNA") can be engineered so as to incorporate aspects of both the crRNA and tracrRNA into a single RNA species. See, e.g., Jinek et al., *Science* 337:816-821(2012), the entire contents of which is incorporated herein by reference.

(45) Cas9 nuclease sequences and structures are well known to those of skill in the art (see, e.g., Ferretti et al., *Proc. Natl. Acad. Sci.* 98:4658-4663(2001); Deltcheva E. et al., *Nature* 471:602-607(2011); and Jinek et al., *Science* 337:816-821(2012), the entire contents of each of which are incorporated herein by reference). Cas9 orthologs have been described in various species, including, but not limited to, *S. pyogenes* and *S. thermophilus*. Additional suitable Cas9 nucleases and sequences will be apparent to those of skill in the art based on this disclosure, and such Cas9 nucleases and sequences include Cas9 sequences from the organisms and loci disclosed in Chylinski et al., (2013) *RNA Biology* 10:5, 726-737; the entire contents of which are incorporated herein by reference. In some embodiments, wild type Cas9 corresponds to Cas9 from *Streptococcus pyogenes* (NCBI Reference Sequence: NC\_002737.2, SEQ ID NO: 4 (nucleotide); and Uniport Reference Sequence: Q99ZW2, SEQ ID NO: 1 (amino acid)).

(46) TABLE-US-00001 (SEQ ID NO: 4)

```
ATGGATAAGAAATACTCAATAGGCTTAGATATCGGCACAAATAGCGTCGGATGGGCGGTGATCAC
TGATGAATATAAGGTTCCGTCTAAAAAGTTCAAGGTTCTGGGAAATACAGACCGCCACAGTATCA
AAAAAAATCTTATAGGGGCTCTTTTATTTGACAGTGGAGAGACAGCGGAAGCGACTCGTCTCAAA
CGGACAGCTCGTAGAAGGTATACACGTCGGAAGAATCGTATTTGTTATCTACAGGAGATTTTTTCA
AATGAGATGGCGAAAGTAGATGATAGTTTCTTTCATCGACTTGAAGAGTCTTTTTTGGTGGAAGAA
GACAAGAAGCATGAACGTCATCCTATTTTTGGAAATATAGTAGATGAAGTTGCTTATCATGAGAAA
TATCCAACATCTATCATCTGCGAAAAAAATTGGTAGATTCTACTGATAAAGCGGATTTGCGCTTA
ATCTATTTGGCCTTAGCGCATATGATTAAGTTTCGTGGTCATTTTTTGATTGAGGGAGATTTAAATC
CTGATAATAGTGATGTGGACAAACTATTTATCCAGTTGGTACAAACCTACAATCAATTATTTGAAG
AAAACCCTATTAACGCAAGTGAGTAGATGCTAAAGCGATTCTTCTGCACGATTGAGTAAATCAA
GACGATTAGAAAATCTCATTGCTCAGCTCCCCGGTGAGAAGAAAAATGGCTTATTTGGGAATCTCA
TTGCTTTGTCAATTGGGTTTGACCCCTAATTTTAAATCAAATTTTGATTGCGAGAAGATGCTAAATT
ACAGCTTTCAAAGATACTTACGATGATGATTTAGATAATTTATTGGCGCAAATTGGAGATCAATA
TGCTGATTTGTTTTTGGCAGCTAAGAATTTATCAGATGCTATTTTACTTTCAGATATCCTAAGAGTA
AATACTGAAATAACTAAGGCTCCCCTATCAGCTTCAATGATTAACGCTACGATGAACATCATCAA
GACTTGACTCTTTTAAAAGCTTTAGTTTCGACAACAACCTCCAGAAAAGTATAAAGAAATCTTTTTT
GATCAATCAAAAAACGGATATGCAGGTTATATTGATGGGGGAGCTAGCCAAGAAGAATTTTATAA
ATTTATCAAACCAATTTTAGAAAAAATGGATGGTACTGAGGAATTATTGGTGAACTAAATCGTGA
```

AGATTTCGCGGCAAGCAACGACCTTTGACCGGATCTATTCCCATCCAAATTCACCTTGGGTGA  
GCTGCATGCTATTTTGAAGAACAAGAAGACTTTTATCCATTTTAAAAGACAATCGTGAGAAGAT  
TGAAAAAATCTTGACTTTTTCGAATTCCTTATTATGTTGGTCCATTGGCGCGTGGAATAGTCGTTTT  
GCATGGATGACTCGGAAGTCTGAAGAAACAATTACCCCATGGAATTTGAAGAAGTTGTCGATAA  
AGGTGCTTCAGCTCAATCATTTATTGAACGCATGACAACTTTGATAAAAAATCTTCCAAATGAAAA  
AGTACTACCAAACATAGTTTGCTTTATGAGTATTTTACGGTTTATAACGAATTGACAAAGGTCAA  
ATATGTTACTGAAGGAATGCGAAAACCAGCATTTCTTTCAGGTGAACAGAAGAAAGCCATTGTTG  
ATTTACTCTTCAAAACAAATCGAAAAGTAACCGTTAAGCAATTAAAAGAAGATTATTTCAAAAAA  
ATAGAATGTTTTGATAGTGTTGAAATTTTCAAGGAGTTGAAGATAGATTTAATGCTTCATTAGGTACC  
TACCATGATTTGCTAAAAATTATTAAAGATAAAGATTTTTTGGATAATGAAGAAAATGAAGATATC  
TTAGAGGATATTGTTTTAACATTGACCTTATTTGAAGATAGGGAGATGATTGAGGAAAGACTTAAA  
ACATATGCTCACCTCTTTGATGATAAGGTGATGAAACAGCTTAAACGTCGCCGTTATACTGGTTGG  
GGACGTTTGTCTCGAAAATTGATTAATGGTATTAGGGATAAGCAATCTGGCAAACAATATTAGAT  
TTTTTGAAATCAGATGGTTTTGCCAATCGCAATTTTATGCAGCTGATCCATGATGATAGTTTGACAT  
TTAAAGAAGACATTCAAAAAGCACAAAGTGTCTGGACAAGGCGATAGTTTACATGAACATATTGCA  
AATTTAGCTGGTAGCCCTGCTATTAAAAAAGGTATTTTACAGACTGTAAAAGTTGTTGATGAATTG  
GTCAAAGTAATGGGGCGGCATAAGCCAGAAAATATCGTTATTGAAATGGCACGTGAAAATCAGAC  
AACTCAAAAGGGCCAGAAAATTTCGCGAGAGCGTATGAAACGAATCGAAGAAGGTATCAAAGAA  
TTAGGAAGTCAGATTCTTAAAGAGCATCCTGTTGAAAATACTCAATTGCAAAATGAAAAGCTCTAT  
CTCTATTATCTCCAAATGGAAGAGACATGTATGTGGACCAAGAATTAGATATTAATCGTTTAAAGT  
GATTATGATGTCGATCACATTGTTCCACAAAGTTTCCTTAAAGACGATTCAATAGACAATAAGGTC  
TTAACGCGTTCTGATAAAAATCGTGGTAAATCGGATAACGTTCCAAGTGAAGAAGTAGTCAAAAA  
GATGAAAAACTATTGGAGACAACTTCTAAACGCCAAGTTAATCACTCAACGTAAGTTTGATAATTT  
AACGAAAGCTGAACGTGGAGGTTTGAGTGAACCTTGATAAAGCTGGTTTTATCAAACGCCAATTGG  
TTGAAACTCGCCAAATCACTAAGCATGTGGCACAAATTTTGGATAGTCGCATGAATACTAAATACG  
ATGAAAATGATAAACTTATTCGAGAGGTTAAAGTGATTACCTTAAAATCTAAATTAGTTTCTGACT  
TCCGAAAAGATTTCCAATTCTATAAAGTACGTGAGATTAACAATTACCATCATGCCCATGATGCGT  
ATCTAAATGCCGTCGTTGGAAGTCTTTGATTAAGAAATATCCAAACTTGAATCGGAGTTTGTCT  
ATGGTGATTATAAAGTTTATGATGTTCTGTAATAATGATTGCTAAGTCTGAGCAAGAAATAGGCAA  
GCAACCGCAAAATATTTCTTTTACTCTAATATCATGAACTTCTTCAAAACAGAAATTACACTTGCA  
AATGGAGAGATTCGCAAACGCCCTCTAATCGAACTAATGGGGAAACTGGAGAAATTGTCTGGGA  
TAAAGGGCGAGATTTTGCCACAGTGCGCAAAGTATTGTCCATGCCCCAAGTCAATATTGTCAAGAA  
AACAGAAGTACAGACAGGCGGATTCTCCAAGGAGTCAATTTTACCAAAAAGAAATTCGGACAAGC  
TTATTGCTCGTAAAAAAGACTGGGATCCAAAAAATATGGTGGTTTTGATAGTCCAACGGTAGCTT  
ATTCAGTCCTAGTGGTTGCTAAGGTGGAAAAAGGGAAATCGAAGAAGTTAAAATCCGTAAAGAG  
TTACTAGGGATCACAAATTATGGAAAGAAGTTCCTTTGAAAAAATCCGATTGACTTTTTAGAAAGCT  
AAAGGATATAAGGAAGTTAAAAAAGACTTAATCATTAACTACCTAAATATAGTCTTTTTGAGTTA  
GAAAACGGTCGTAAACGGATGCTGGCTAGTGCCGGAGAATTACAAAAGGAAATGAGCTGGCTCT  
GCCAAGCAAATATGTGAATTTTTTATATTTAGCTAGTCATTATGAAAAGTTGAAGGGTAGTCCAGA  
AGATAACGAACAAAAACAATTGTTTGTGGAGCAGCATAAGCATTATTTAGATGAGATTATTGAGC  
AAATCAGTGAATTTTCTAAGCGTGTTATTTTAGCAGATGCCAATTTAGATAAAGTTCTTAGTGCAT  
ATAACAAACATAGAGACAAACCAATACGTGAACAAGCAGAAAATATTATTCAATTTATTTACGTTG  
ACGAATCTTGAGCTCCCGCTGCTTTTAAATATTTTGATACAACAATTGATCGTAAACGATATACG  
TCTACAAAAGAAGTTTTAGATGCCACTCTTATCCATCAATCCATCACTGGTCTTTATGAAACACGC  
ATTGATTTGAGTCAGCTAGGAGGTGACTGA (SEQ ID NO: 1)

MDKKYSIGLDIGTNSVGWAVITDEYKVPSKFKVLGNTDRHSIKKNLIGALLEDSETAEATRLKRTAR  
RRYTRRKNRICYLQEIFSNEMAKVDDSFHRLSESFLEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLR  
KKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDA  
KAILSARLSKSRLENLIAQLPGEKKNGLFGNLIALSLGLTPNFKSNFDLAEDAKLQLSKDTYDDDLN  
LLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLKALVRQQLPEK  
YKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDNGSIPHQIHL  
GELHAILRRQEDFYPLKDNREKIEKILTFRIYYVGPLARGNSRFAWMTRKSEETITPWNFEEVVDKGA  
SAQSFIERMTNFDKNLPNEKVLPKHSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFK  
TNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEENEDILEDIVLTLT  
LFEDREMIEERLKTYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRN  
FMQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIKKGILOTVKVVDELVKVMGRHKPENIVI  
EMARENQTTQKGQKNSRERMKRIEELGSGILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQEL  
DINRLSDYDVDHIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKMKMKNYWRQLLNAKLITQRK  
FDNLTKAERGGELSELDKAGFIKROLVETROITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSD  
FRKDFQFYKVBREINNYHHAHDAYLNAVVGTAIIKKYPKLESEFVYGDYKVYDVRKMIKSEQEIGKA  
TAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVL SMPQVNIVKKTEVQ

IGGFSKESILPKRNSDKLIARKKDWDPKPKYGGFSDSPTVAYSVLVAKVEKGSKKLKSVKELLGITIME  
RSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELENGKRKMLASAGELQKGNELALPSKYVNFLYLA  
SHYEKLKGSPEDEQKQLFVEQHKHYLDEIIQISEFSKRVLADANLDKVL SAYNKHHRDKPIREQAENI  
IHLFTLTNLGAPAAFKYFDDTTIDRKRYTSTKEVLDATLIHQSTGLYETRIDLSQLGGD (single underline:  
HNH domain; double underline: RuvC domain) In some embodiments, wild type Cas9  
corresponds to Cas9 from *Streptococcus aureus*. *S. aureus* Cas9 wild type (SEQ ID NO: 6)  
MKRNYILGLDIGITSVGYGIIDYETRDVIDAGVRLFKEANVENNEGRRSKRGRARLRKRHRRIQRVKK  
LLFDYNLLTDHSELSGINPYEARVKGLSQKLSSEEFSAALLHLAKRRGVHNVNEVEEDTGNELSTKEQI  
SRNSKALEEKYVAELQLERLKKDGEVRGSINRFKTS DYVKEAKQLLKVQKAYHOLDQSFIDTYIDLLE  
TRRTYYEGPGEGSPFGWKDIKEWYEMLMGHCTYFPEELRSVKYAYNADLYNALNDLNNLVITRDENE  
KLEYYEKFQIENVFQKQKKKPTLKQIAKEILVNEEDIKGYRVTS TGKPEFTNLKVYHDIKDITARKEIEN  
AELLDDQIAKILTIYQSSEDIQEELTNLNSLTQEEIEQISNLKGYTGTHNLSLKAINLILDELWHTNDNQIA  
IFNRLKLVPKKVDLSQQKEIPTTLVDDFILSPVVKRSFIQSIKVINAIKKYGLPNDIIELAREKNSKDAQK  
MINEMQKRNRQTNERIEEIIRTTGKENAKYLIEKIKLHDMQEGKCLYSLEAIPLEDLLNPNFYEV DHIIP  
RSVSFDNSFNKVLVKQEENSKKGNRTPFQYLSSSDSKISYETFKKHILNLAKGKGRISKTKKEYLLEER  
DINRFSVQKDFINRNLVDTRYATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGY  
KHH AEDALIIANADFIKWKLDKAKVMENQMFE EKQAESMPEIETE QEYKEIFITPHQIKHIKDFK  
DYKYSHRVDKPNRELINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLKKLINKSPEKLLMYHHDP  
QTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDNGPVIKKIKYYGNKLNAHLDITDDYPNSR  
NKVVKLSLKP YRFDVYLDNGVYKFVTVKNLDVIKKENY YEVNSKCYEEAKKLKKISNQAEFIASFYNN  
DLIKINGEL YRVIGVNNDLLNRIEVNMIDITYREYLENMNDKRPPRIIKTIASKTQSIKKYSTDILGNLYE  
VKSKKHPPQIIKKG In some embodiments, wild type Cas9 corresponds to Cas9 from *Streptococcus*  
*thermophilus*. *Streptococcus thermophilus* wild type CRISPR3 Cas9 (St3Cas9) (SEQ ID NO: 7)  
MTKPYSIGLDIGTNSVGWAVITDNYKVPSKKMKVLGNTSKKYIKKNLLGVLLFDSGITAEGRRLKRTA  
RRRYTRRRNRILYLQEIFSTEMATLDDAFFQRLDDSFVPDDKRD SKYPIFGNLVEEKVYHDEFPTIYHL  
RKYLADSTKKADRLVYLALAHMIKYRGHFLIEGEFNSKNNDIQKNFQDFLDTYNAIFESDLSLENSKQ  
LEEIVKDKISKLEKKDRILKLFPG EKNSGIFSEFLKLIVGNQADFRKCFNLDEKASLHESKESYDEDLET  
LGYIGDDYSDVFLKAKKLYDAILSGFLTVDNETEAPLSSAMIKRYNEHKEDLALLKEYIRNISLKTYN  
EVFKDDTKNGYAGYIDGKTNQEDFYVYLKNLLAEFEGADYFLEKIDREDFLRKQRTFDNGSIPYQIHLQ  
EMRAILDKQAKFYPFLAKNKERIEKILTFRIPYYVGPLARGNSDFAWSIRKRNEKITPWNFEDVIDKESS  
AEAFINRMTSFDLYLP EKVLPKHSLLYETFN VYNELTKVRFIAESMRDYQFLDSKQKKDIVRLYFKDK  
RKVTDKDIIEYLHAIYGYDGI ELKGIEKQFNSSLSTYHDL LNIINDKEFLDDSSNEAIIIEIIHTLTFEDRE  
MIKQRLSKFENIFDKSVLKKLSRRHYTGWGKLSAKLINGIRDEKSGNTILDYLI DDGISNRNFMQLIHDD  
ALSFKKKIQKAQIIGDEDKGNIKEVVKSLPGSPA IKKGILQSIKIVDELVKVMGGRKPESIVVEMARENQ  
YTNQGKSNSQQLKRLEKSLKELGSKILKENIPAKLSKIDNNALQNDRLYLYLQNGKDMYTGD DLDI  
DRLSNYDIDHIIPQAF LKDNSIDNKVLVSSASN RGKSDDFPSLEVVKRKRTFWYQLLKS KLISQRKFDNL  
TKAERGGLLPEDKAGFIQRQLVETRQITKHVARLLDEKENNKKDENNR AVRTVKIITLKTSLVSQFRKD  
FELYK VREINDFHHAHDAYLNAVIA SALLKKYPKLEPEFVYGDYPKYNSFRERKSATEKVYFYSNIMNI  
FKKSISLADGRVIERPLIEVNEETGESVWNKESDLATVRRVLSYPQVNVVKKVEEQNHGLDRGKPKGL  
FNANLSSKPKPNSNENLVGAKEYLDPKKYGGYAGISNSFAVLVKGTIEKGAKKKITNVLEFQGISILDRI  
NYRKDKLNFLEKGYKDIELIIELPKYSLFELSDGSRRMLASILSTNNKRGEIHKGNQIFLSQKFVKLLYH  
AKRISNTINENHRKYVENHKKEFEELFYI LEFNENYVGAKKNGKLLNSAFQSWQNHSIDELCSSFIGPT  
GSEKGLFELTSRGSAADFEFLGVKIPRYRDYTPSSLLKDATLIHQSVTGLYETRIDLAKLGEG *Streptococcus*  
*thermophilus* CRISPR1 Cas9 wild type (St1Cas9) (SEQ ID NO: 8)  
MSDLVLGLDIGISVGVGILNKVTGEIIHKNSRIFPAAQAENNLVRRTRNRQGRRLTRRKKHRRVRLNRL  
FEESGLITDFTKISINLPYQLRVKGLTDELSNEELFIALKNMVKHRGISYLD DASDDGNSSIGDYAQIVK  
ENSKQLETKTPGQIQ LERYQTYGQLRGDFTVEKD GKKHRLINVFPTSAYRSEALRILQTQQEFNPQITDE  
FINRYLEILTGRKYYHGP GNEKSRTDYGRYRTSGETLDNIFGILIGKCTFY PDEFRAAKASYTAQEFNL  
LNDLNNLTVPTETKLSKEQKNQIINYVKNEKAMGPAKLFKYIAKLLSCDVADIKGYRIDKSGKAEIHT  
FEAYRKMKTLETLDIEQMDRETLDKLAYVLT LNTEREGIQEAL EHEFADGSFSQKQVDELVQFRKANS  
SIFGKGWHNFSVKLMMELIPELYETSEEQMTILTRLGKQKTTSSSNKTKYIDEKLLTEEIYNPVVAKSVR  
QAIKIVNAAIKEYGDFDNIV IEMARETNEDDEKKA IQKIQKANKDEKDAAMLKAANQYNGKAELPHSV  
FHGHKQLATKIRLWHQQGERCLYTGKTISI HDLINNSNQFEVDHILPLSITFDDSLANKVLVYATANQE  
KGQRTPYQALDSMDDAWSFRELKAFVRESKTL SNKKKEYLLTEEDISKFDVRKKFIERNLVDTRYASR  
VVLNALQEHFRAHKIDTKVSVVRGQFTSQLRRHWGIEKTRDTYHHH AVDALIAASSQLNLWKKQKN  
TLVSYSEDQLLDIETGELISDDEYKESVFKAPYQH FVDTLKSKEFEDSILFSYQVDSKFNRKISDATIYAT  
RQAKVGKDKADETYVLGKIKDIYTQDGYDAFMKIYKKDKSKFLMYRHDPQTFEKVIEPILENYPNKQI  
NEKGKEVPCNPFLKYKEEHGYIRKYSKKGNGPEIKSLKY YDSKLG NHIDITPKDSNNKVVLQSVSPWR  
ADVYFNKTTGKYEILGLKYADLQFEKGTGTYKISQEKYNDIKKKEGVDS DSEFKFTLYKNDLLLVKDT  
ETKEQQLFRFLSRTMPKQKH YVELKPYDKQKFEGGEALIKVLGNVANS GQCKGLGKSNISYKVRTD  
VLGNOHIIKNEGD KPKLDF

(47) In some embodiments, the Cas9 domain comprises any of the fusion proteins provided herein is a Cas9 from archaea (e.g. nanoarchaea), which constitute a domain and kingdom of single-celled prokaryotic microbes. In some embodiments, the Cas9 domain is CasX or CasY, which have been described in, for example, Burstein et al., "New CRISPR-Cas systems from uncultivated microbes." *Cell Res.* 2017 Feb. 21. doi: 10.1038/cr.2017.21, which is incorporated herein by reference. Using genome-resolved metagenomics, a number of CRISPR-Cas systems were identified, including the first reported Cas9 in the archaeal domain of life. This divergent Cas9 protein was found in nanoarchaea as part of an active CRISPR-Cas system. In bacteria, two previously unknown systems were discovered, CRISPR-CasX and CRISPR-CasY, which are among the most compact systems yet discovered. In some embodiments, Cas9 refers to CasX, or a variant of CasX. In some embodiments, Cas9 refers to a CasY, or a variant of CasY. It should be appreciated that other RNA-guided DNA binding proteins may be used as a nucleic acid programmable DNA binding protein (napDNAbp) and are within the scope of this disclosure.

(48) In some embodiments, the Cas9 domain comprises an amino acid sequence that is at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to a naturally-occurring CasX or CasY protein. In some embodiments, the Cas9 domain is a naturally-occurring CasX or CasY protein. In some embodiments, the Cas9 domain comprises an amino acid sequence that is at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to any one of SEQ ID NOs: 336-337 or 3000. In some embodiments, the Cas9 domain comprises an amino acid sequence of any one SEQ ID NOs: 336-337 or 3000. It should be appreciated that CasX and CasY from other bacterial species may also be used in accordance with the present disclosure.

(49) In some embodiments, wild-type Cas9 refers to CasX from *Sulfolobus islandicus* (strain REY15A).

(50) TABLE-US-00002 (SEQ ID NO: 338)

MEVPLYNIFGDNYIIQVATEAENSTIYNNKVEIDDEELRNVLNLAYKIAKNNEDAAAE  
RRGKAKKKKGEEGETTTSNIILPLSGNDKNPWTETLKCYNFPTTVALSEVFNFSQV  
KECEEVSAPSFVKPEFYKFGFRSPGMVERTRRVKLEVEPHYLMIAAAGWVLTRLGKA  
KVSEGDYVGVNVFTPTRGILYSLIQNVNGIVPGIKPETAFLGLWIARKVVSSVTNPVNS  
VVSITYTISDAVGQNPTTINGGFSIDLTKLLEKRDLLSERLEAIARNALSISSNMRERYIV  
LANIYIYEYLTGSKRLEDLLYFANRDLIMNLSDDGKVRDLKLISAYVNGELIRGEG In some embodiments,  
wild-type Cas9 refers to CasX from *Sulfolobus islandicus* (strain REY15A). (SEQ ID NO: 339)

MEVPLYNIFGDNYIIQVATEAENSTIYNNKVEIDDEELRNVLNLAYKIAKNNEDAAAE  
RRGKAKKKKGEEGETTTSNIILPLSGNDKNPWTETLKCYNFPTTVALSEVFNFSQV  
KECEEVSAPSFVKPEFYEFGRSPGMVERTRRVKLEVEPHYLIIAAAGWVLTRLGKAK  
VSEGDYVGVNVFTPTRGILYSLIQNVNGIVPGIKPETAFLGLWIARKVVSSVTNPVNSV  
VRIYTISDAVGQNPTTINGGFSIDLTKLLEKRYLLSERLEAIARNALSISSNMRERYIVL  
ANYIYIYEYLTGSKRLEDLLYFANRDLIMNLSDDGKVRDLKLISAYVNGELIRGEG In some embodiments,

wild-type Cas9 refers to CasY from a *Parcubacteria* group *bacterium*. CasY  
(ncbi.nlm.nih.gov/protein/APG80656.1) >APG80656.1 CRISPR-associated protein CasY [uncultured  
*Parcubacteria* group *bacterium*] (SEQ ID NO: 3000)

MSKRHPRISGVKGYRLHAQRLEYTGKSGAMRTIKYPLYSSPSGGRTVPREIVSAINDDYVGLYGLSNFD  
DLYNAEKRNEEKVYSVLDFWYDCVQYGAVFSYTAPGLLKNVAEVRGGSYELTKTLKGSHLYDELQID  
KVIKFLNKKEISRANGSLDKLKKDIIDCFKAEYRERHKDQCENKLADDIKNAKKDAGASLGERQKKLFR  
DFFGISEQSENDKPSFTNPLNLTCCLLPFDTVNNNRNRGEVLENKLKEYAQKLDKNEGSLEMWEYIGIG  
NSGTAFSNFLGEGFLGRLRENKITEKKAMMDITDAWRGQEQQEELEKRLRILAALTIKLREPKFNDHW  
GGYRSDINGKLSSWLQNYINQTVKIKEDLKGHKDLKKAKEMINRFGESDTKEEAVVSSLLESIEKIVP  
DDSADDEKPDIPAIAIYRRFLSDGRLTLNRFVQREDVQEALIKERLEAEKKKKPKKRKKKSDAEDEKETI  
DFKELFPHLAKPLKLPNFYGD SKRELYKKYKNAAIYTDALWKAVEKIYKSAFSSSLKNSFFD TDFDK  
DFFIKRLQKIFSVYRRFNTDKWKPIVKNSFAPYCDIVSLAENEVLYKPKQSRSRKSAIDKNRVRLPSTE  
NIAKAGIALARELSVAGFDWKDLLKKEEH EYIDLIELHKTALALLAVTETQLDISALDFVENGTVKD  
FMKTRDGNLVLEGRFLEMFSQSIVFSELRLAGLMSRKEFITRS AIQTMNGKQAELLYIPHEFQSAKITT  
PKEMSRAFLDLAPAEFATSLEPELSEKSLKLKQMRYYPHYFGYELTRTGQGIDGGVAENALRLEKSP  
VKKREIKCKQYKTLGRGQNKIVLYVRSSYYQTQFLEWFLHRPKNVQTDVAVSGSFLIDEKKVKTRWN  
YDALTVALEPVSGSERVFVSQPFTIFPEKSAEEEGQRYLGIDIGEYGIAYTALEITGDSAKILDQNFISDPQ  
LKTREEVKGLKLDQRRGTFAMPSTKIARIRESLVHSLRNRIHHLALKHKAKIVYELEVSRFEEGKQKIK  
KVYATLKKADVYSEIDADKNLQTTVWGKLAVASEISASYTSQFCGACKKLWRAEMQVDETITTQELIG  
TVRVIKGGTLIDAIKDFMRPPIFDENDTPFPKYRDFCDKHHISKMKMRGNSCLFICPFCRANADADIQASQ  
TIALLRVKEEKKVEDYFERFRKLKNIKVLGQMKKI

(51) In some embodiments, Cas9 refers to Cas9 from: *Corynebacterium ulcerans* (NCBI Refs: NC\_015683.1, NC\_017317.1); *Corynebacterium diphtheria* (NCBI Refs: NC\_016782.1, NC\_016786.1); *Spiroplasma syrphidicola* (NCBI Ref: NC\_021284.1); *Prevotella intermedia* (NCBI Ref: NC\_017861.1); *Spiroplasma taiwanense* (NCBI Ref: NC\_021846.1); *Streptococcus iniae* (NCBI Ref: NC\_021314.1); *Belliella baltica* (NCBI Ref: NC\_018010.1); *Psychroflexus torquus* (NCBI Ref: NC\_018721.1); *Listeria innocua* (NCBI Ref: NP\_472073.1), *Campylobacter jejuni*

(NCBI Ref: YP\_00234900.1) or *Neisseria meningitidis* (NCBI Ref: YP\_002342100.1) or to a Cas9 from any of the organisms listed in Example 1 (SEQ ID NOs: 11-260).

(52) To be used as in the fusion protein of the present disclosure as the guide nucleotide sequence-programmable DNA binding protein domain, a Cas9 protein needs to be nuclease inactive. A nuclease-inactive Cas9 protein may interchangeably be referred to as a “dCas9” protein (for nuclease-“dead” Cas9). Methods for generating a Cas9 protein (or a fragment thereof) having an inactive DNA cleavage domain are known (See, e.g., Jinek et al., *Science*. 337:816-821(2012); Qi et al., (2013) *Cell*. 28; 152(5):1173-83, the entire contents of each of which are incorporated herein by reference). For example, the DNA cleavage domain of Cas9 is known to include two subdomains, the HNH nuclease subdomain and the RuvC1 subdomain. The HNH subdomain cleaves the strand complementary to the gRNA, whereas the RuvC1 subdomain cleaves the non-complementary strand. Mutations within these subdomains can silence the nuclease activity of Cas9. For example, the mutations D10A and H840A completely inactivate the nuclease activity of *S. pyogenes* Cas9 (Jinek et al., *Science*. 337:816-821(2012); Qi et al., *Cell*. 28; 152(5):1173-83 (2013)). dCas9 (D10A and H840A)

(53) TABLE-US-00003 (SEQ ID NO: 2)

MDKKYSIGLAIGTNSVGWAVITDEYKVPSSKKFKVLGNTDRHSIKKNLIGALLEDSGETAEATRLKRTAR  
RRYTRRKNRICYLQEIFSNEMAKVDDSFHRLSEESFLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLR  
KKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDA  
KAILSARLSKSRLENLIAQLPGEKKNGLFGNLIALSLGLTPNFKSNFDLAEDAKLQLSKDTYDDDLDN  
LLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLKALVRQQLPEK  
YKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKNLNREDLLRKQRTFDNGSIPHQIHL  
GELHAILRRQEDFYFPFLKDNREKIEKILTFRIPIYVGPLARGNSRFAWMTRKSEETITPWNFEEVVDKGA  
SAQSFIERMTNFDKNLPNEKVLPKHSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFK  
TNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLTKIHKDKDFLDNEENEDILEDIVLTLT  
LFEDREMIEERLKTIAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRN  
FMQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIIKKGILQTVKVVDELVKVMGRHKPENIVI  
EMARENQTTQKGQKNSRERMKRIIEGKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQEL  
DINRLSDYDVEDAIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKMKMKNYWRQLLNAKLITQRK  
FDNLTKAERGGLSELDKAGFIKRLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSD  
FRKDFQFYKVINNYHHAHDAYLNAVVGTAIIKKYPKLESEFVYGDYKVYDVRKMIKSEQEIGKA  
TAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVLSPQVNVKKTVEVQ  
TGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVAKVEKGKSKKLKSVKELLGITIME  
RSSFENPIDFLEAKGYKEVKKDLIIKLPKYSLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLA  
SHYEKLKGSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVILADANLDKVL SAYNKHRRDKPIREQAENI  
IHLFTLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQSTGLYETRIDLSQLGGD (single underline:  
HNH domain; double underline: RuvC domain)

(54) The dCas9 of the present disclosure encompasses completely inactive Cas9 or partially inactive Cas9. For example, the dCas9 may have one of the two nuclease domain inactivated, while the other nuclease domain remains active. Such a partially active Cas9 may also be referred to as a Cas9 nickase, due to its ability to cleave one strand of the targeted DNA sequence. The Cas9 nickase suitable for use in accordance with the present disclosure has an active HNH domain and an inactive RuvC domain and is able to cleave only the strand of the target DNA that is bound by the sgRNA. The Cas9 nickase of the present disclosure may comprise mutations that inactivate the RuvC domain, e.g., a D10A mutation. It is to be understood that any mutation that inactivates the RuvC domain may be included in a Cas9 nickase, e.g., insertion, deletion, or single or multiple amino acid substitution in the RuvC domain. In a Cas9 nickase described herein, while the RuvC domain is inactivated, the HNH domain remains activate. Thus, while the Cas9 nickase may comprise mutations other than those that inactivate the RuvC domain (e.g., D10A), those mutations do not affect the activity of the HNH domain. In a non-limiting Cas9 nickase example, the histidine at position 840 remains unchanged. The sequence of an exemplary Cas9 nickase suitable for the present disclosure is provided below.

(55) TABLE-US-00004 Cas9 Nickase (D10A) (SEQ ID NO: 3)

MDKKYSIGLAIGTNSVGWAVITDEYKVPSSKKFKVLGNTDRHSIKKNLIGALLEDSGETAEATRLKRTAR  
RRYTRRKNRICYLQEIFSNEMAKVDDSFHRLSEESFLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLR  
KKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDA  
KAILSARLSKSRLENLIAQLPGEKKNGLFGNLIALSLGLTPNFKSNFDLAEDAKLQLSKDTYDDDLDN  
LLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLKALVRQQLPEK  
YKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKNLNREDLLRKQRTFDNGSIPHQIHL  
GELHAILRRQEDFYFPFLKDNREKIEKILTFRIPIYVGPLARGNSRFAWMTRKSEETITPWNFEEVVDKGA  
SAQSFIERMTNFDKNLPNEKVLPKHSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFK  
TNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLTKIHKDKDFLDNEENEDILEDIVLTLT  
LFEDREMIEERLKTIAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRN  
FMQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIIKKGILQTVKVVDELVKVMGRHKPENIVI  
EMARENQTTQKGQKNSRERMKRIIEGKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQEL  
DINRLSDYDVEDHIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKMKMKNYWRQLLNAKLITQRK  
FDNLTKAERGGLSELDKAGFIKRLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSD

FRKDFQFYKYNVLAHDAYLNAVVGTAALIKKYPKLESEFVYGDYKVDVVRKMIASEQKLEIGKA  
 TAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVL SMPQVNIVKKTEVQ  
 TGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVAKVEKGKSKKLKSVKELLGITIME  
 RSSFEKNPIDFLEAKGYKEVKKDLIILPKYSLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLA  
 SHYEKLKGSPEDEQKQLFVEQHKHYLDEIIEQISEFSKRVLADANLDKVL SAYNKH RDKPIREQAENI  
 IHLFTLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQ SITGLYETRIDLSQLGGD (single underline:  
 HNH domain; double underline: RuvC domain) *S. aureus* Cas9 Nickase (D10A) (SEQ ID NO: 5)  
 MKRNYILGLAIGITSVGYGIIDYETRDVIDAGVRLFKEANVENNEGRRSKRGARRLRRRRRHRIQRVKK  
 LLFDYNLLTDHSELSGINPYEARVKGLSQKLSEEEFSAALLHLAKRRGVHNVNEVEEDTGNELSTKEQI  
 SRNSKALEEKYVAELQLERLKKDGEVRGSINRFKTS DYVKEAKQLLKVQKAYHQLDQSFIDTYIDLL  
 TRRTYYEGPGEGSPFGWKDIKEWYEMLMGHCTYFPEELRSVKYAYNADLYNALNDLNNLVITRDENE  
 KLEYEYKFQIENVFKQKKKPTLTKQIAKEILVNEEDIKGYRVTSTGKPEFTNLKVYHDIKDITARKEIEN  
 AELLDQIAKILTIYQSSEDIQEELTNLSEL TQEEIEQISNLKGYTGTHNLSLKAINLILDELWHTNDNQIA  
 IFNRLKLVPKKVDLSQQKEIPTTLVDDFILSPVVKRSFIQSIKVINAIKKYGLPNDIIELAREKNSKDAQK  
 MINEMQKRNRQTNERIEEII RTTGKENAKYLIEKIKLHDMQEGKCLYSLEAIPLEDLLNPNFNYEVDHIIP  
 RSVSFDNSFNKVLVKQEENS KKG NRTPFQYLSSSDSKISYETFKKHILNLA KGKGRISKTKKEYLLEER  
 DINRFSVQKDFINRNLVDTRYATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGY  
 KHHAEDALIIANADFIFKEWKLDKAKKVMENQMFE EKQAESMPEIETE QEYKEIFITPHQIKHIKDFK  
 DYKYSHRVDKKPNRELINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLKLINKSPEKLLMYHHDP  
 QTYQKLKLIMEQYGDEKNPLYKYEEETGNYLTKYSKKDNGPVIKKIKYYGNKLNAHL DITDDYPNSR  
 NKVVKL SLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYYEVNSKCYEEAKKLKKISNQAEFIASFYNN  
 DLIKINGEL YRVIGVNNDLLNRIEVNMIDITYREYLENMNDKRPPRIIKTIASKTQSIKKYSTDILGNLYE  
 VKSKKHPQIIKKG

(56) It is appreciated that when the term “dCas9” or “nuclease-inactive Cas9” is used herein, it refers to Cas9 variants that are inactive in both HNH and RuvC domains as well as Cas9 nickases. For example, the dCas9 used in the present disclosure may include the amino acid sequence set forth in SEQ ID NO: 2 or SEQ ID NO: 3. In some embodiments, the dCas9 may comprise other mutations that inactivate RuvC or HNH domain. Additional suitable mutations that inactivate Cas9 will be apparent to those of skill in the art based on this disclosure and knowledge in the field, and are within the scope of this disclosure. Such additional exemplary suitable nuclease-inactive Cas9 domains include, but are not limited to, D839A and/or N863A (See, e.g., Prashant et al., *Nature Biotechnology*. 2013; 31(9): 833-838, the entire contents of which are incorporated herein by reference), or K603R (See, e.g., Chavez et al., *Nature Methods* 12, 326-328, 2015, the entire contents of which is incorporated herein by reference). The term Cas9, dCas9, or Cas9 variant also encompasses Cas9, dCas9, or Cas9 variant from any organism. Also appreciated is that dCas9, Cas9 nickase, or other appropriate Cas9 variants from any organisms may be used in accordance with the present disclosure.

(57) A “deaminase” refers to an enzyme that catalyzes the removal of an amine group from a molecule, or deamination. In some embodiments, the deaminase is a cytidine deaminase, catalyzing the hydrolytic deamination of cytidine or deoxycytidine to uridine or deoxyuridine, respectively. In some embodiments, the deaminase is a cytosine deaminase, catalyzing the hydrolytic deamination of cytosine to uracil (e.g., in RNA) or thymine (e.g., in DNA). In some embodiments, the deaminase is a naturally-occurring deaminase from an organism, such as a human, chimpanzee, gorilla, monkey, cow, dog, rat, or mouse. In some embodiments, the deaminase is a variant of a naturally-occurring deaminase from an organism, that does not occur in nature. For example, in some embodiments, the deaminase or deaminase domain is at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75% at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to a naturally-occurring deaminase from an organism.

(58) A “cytosine deaminase” refers to an enzyme that catalyzes the chemical reaction “cytosine+H.sub.2O.fwdarw.uracil+NH.sub.3” or “5-methyl-cytosine+H2O.fwdarw.thymine+NH.sub.3.” As it may be apparent from the reaction formula, such chemical reactions result in a C to U/T nucleobase change. In the context of a gene, such nucleotide change, or mutation, may in turn lead to an amino acid residue change in the protein, which may affect the protein function, e.g., loss-of-function or gain-of-function.

(59) One exemplary suitable class of cytosine deaminases is the apolipoprotein B mRNA-editing complex (APOBEC) family of cytosine deaminases encompassing eleven proteins that serve to initiate mutagenesis in a controlled and beneficial manner. The apolipoprotein B editing complex 3 (APOBEC3) enzyme provides protection to human cells against a certain HIV-1 strain via the deamination of cytosines in reverse-transcribed viral ssDNA. These cytosine deaminases all require a Zn.sup.2+-coordinating motif (His-X-Glu-X.sub.23-26-Pro-Cys-X.sub.24-Cys; SEQ ID NO: 820) and bound water molecule for catalytic activity. The Glu residue acts to activate the water molecule to a zinc hydroxide for nucleophilic attack in the deamination reaction. Each family member preferentially deaminates at its own particular “hotspot”, ranging from WRC (W is A or T, R is A or G) for hAID, to TTC for hAPOBEC3F. A recent crystal structure of the catalytic domain of APOBEC3G revealed a secondary structure comprised of a five-stranded 3-sheet core flanked by six  $\alpha$ -helices, which is believed to be conserved across the entire family. The active center loops have been shown to be responsible for both ssDNA binding and in determining “hotspot” identity. Overexpression of these enzymes has been linked to genomic instability and cancer, thus highlighting the importance of sequence-specific targeting.

Another suitable cytosine deaminase is the activation-induced cytidine deaminase (AID), which is responsible for the maturation of antibodies by converting cytosines in ssDNA to uracils in a transcription-dependent, strand-biased fashion. (60) Herein, a “nucleobase editor” refers to a protein that edits a nucleotide base. “Edit” refers to the conversion of one nucleotide base to another. For example, the nucleobase may target C bases in a nucleic acid sequence and convert the C to T base. In some embodiments, the C to T editing is carried out by a deaminase, e.g., a cytosine deaminase. Other types of base conversions are also contemplated. In some embodiments, the nucleobase editor comprises a DNA binding domain that directs it to a target sequence.

(61) As such, a base editor may be a cytosine deaminase-dCas9 fusion protein. In some embodiments, the base editor may be a deaminase-dCas9-UGI fusion protein. In some embodiments, the base editor may be a APOBEC1-dCas9-UGI fusion protein. In some embodiments, the base editor may be APOBEC1-Cas9 nickase-UGI fusion protein. In some embodiments, the base editor may be APOBEC1-dCpf1-UGI fusion protein. In some embodiments, the base editor may be APOBEC1-dNgAgo-UGI fusion protein. In some embodiments, the base editor may be a pmCDA1-Cas9 nickase-UGI fusion protein. In some embodiments, the base editor may be a human APOBEC3G-Cas9 nickase UGI fusion protein. In some embodiments, the base editor may comprise a second UGI domain. Non-limiting exemplary sequences of the nucleobase editors described herein are provided in Example 1, SEQ ID NOs: 293-302, 1071, and 1084. Such nucleobase editors and methods of using them for genome editing have been described in the art, e.g., in U.S. Pat. No. 9,068,179, US Patent Application Publications US 2015/0166980, US 2015/0166981, US 2015/0166982, US20150166984, and US20150165054, and US Provisional Applications, U.S. Ser. No. 62/245,828, filed Oct. 23, 2015; 62/279,346, filed Jan. 15, 2016; 62/311,763, filed Mar. 22, 2016; 62/322,178, filed Apr. 13, 2016; 62/357,352, filed Jun. 30, 2016, U.S. Pat. No. 62,370,700, filed Aug. 3, 2016; 62/398,490, filed Sep. 22, 2016; 62/408,686, filed Oct. 14, 2016; PCT Application PCT/US2016/058344, filed Oct. 22, 2016, U.S. patent application Ser. No. 15/311,852, filed Oct. 22, 2016; Komor et al. (2017) Improved Base Excision Repair Inhibition and Bacteriophage Mu Gam Protein Yields C:G-to-T:A base editors with higher efficiency and product purity. *Sci Adv*, 3: eaao4774; and in Komor et al., *Nature*, Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage, 533, 420-424 (2016), the entire contents of each of which are incorporated herein by reference.

(62) The term “target site” or “target sequence” refers to a sequence within a nucleic acid molecule (e.g., a DNA molecule) that is deaminated by the fusion protein provided herein. In some embodiments, the target sequence is a polynucleotide (e.g., a DNA), wherein the polynucleotide comprises a coding strand and a complementary strand. The meaning of a “coding strand” and “complementary strand” is the common meaning of the terms in the art. In some embodiments, the target sequence is a sequence in the genome of a mammal. In some embodiments, the target sequence is a sequence in the genome of a human. The term “target codon” refers to the amino acid codon that is edited by the base editor and converted to a different codon via deamination of C base. In some embodiments, the target codon is edited in the coding strand. In some embodiments, the target codon is edited in the complementary strand.

(63) The term “linker,” as used herein, refers to a chemical group or a molecule linking two molecules or moieties, e.g., two domains of a fusion protein, such as, for example, a nuclease-inactive Cas9 domain and a nucleic acid editing domain (e.g., a deaminase domain). Typically, the linker is positioned between, or flanked by, two groups, molecules, or other moieties and connected to each one via a covalent bond, thus connecting the two. In some embodiments, the linker is an amino acid or a plurality of amino acids (e.g., a peptide or protein). In some embodiments, the linker is an organic molecule, group, polymer, or chemical moiety. In some embodiments, the linker is 2-100 amino acids in length, for example, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 30-35, 35-40, 40-45, 45-50, 50-60, 60-70, 70-80, 80-90, 90-100, 100-150, or 150-200 amino acids in length. Longer or shorter linkers are also contemplated.

(64) The term “mutation,” as used herein, refers to a substitution of a residue within a sequence, e.g., a nucleic acid or amino acid sequence, with another residue, or a deletion or insertion of one or more residues within a sequence. Mutations are typically described herein by identifying the original residue followed by the position of the residue within the sequence and by the identity of the newly substituted residue. Various methods for making the amino acid substitutions (mutations) provided herein are well known in the art, and are provided by, for example, Green and Sambrook, *Molecular Cloning: A Laboratory Manual* (4<sup>sup</sup>.th ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2012)).

(65) The terms “nucleic acid” and “nucleic acid molecule,” as used herein, refer to a compound comprising a nucleobase and an acidic moiety, e.g., a nucleoside, a nucleotide, or a polymer of nucleotides. Typically, polymeric nucleic acids, e.g., nucleic acid molecules comprising three or more nucleotides are linear molecules, in which adjacent nucleotides are linked to each other via a phosphodiester linkage. In some embodiments, “nucleic acid” refers to individual nucleic acid residues (e.g. nucleotides and/or nucleosides). In some embodiments, “nucleic acid” refers to an oligonucleotide chain comprising three or more individual nucleotide residues. As used herein, the terms “oligonucleotide” and “polynucleotide” can be used interchangeably to refer to a polymer of nucleotides (e.g., a string of at least three nucleotides). In some embodiments, “nucleic acid” encompasses RNA as well as single and/or double-stranded DNA. Nucleic acids may be naturally occurring, for example, in the context of a genome, a transcript, an mRNA, tRNA, rRNA, siRNA, snRNA, a plasmid, cosmid, chromosome, chromatid, or other naturally occurring nucleic acid molecule. On the other hand, a nucleic acid molecule may be a non-naturally occurring molecule, e.g., a recombinant DNA or RNA, an artificial chromosome, an engineered genome, or fragment thereof, or a synthetic DNA, RNA, DNA/RNA hybrid, or including non-naturally occurring nucleotides or nucleosides. Furthermore, the terms “nucleic acid,” “DNA,” “RNA,”

and/or similar terms include nucleic acid analogs, e.g., analogs having other than a phosphodiester backbone. Nucleic acids can be purified from natural sources, produced using recombinant expression systems and optionally purified, chemically synthesized, etc. Where appropriate, e.g., in the case of chemically synthesized molecules, nucleic acids can comprise nucleoside analogs such as analogs having chemically modified bases or sugars, and backbone modifications. A nucleic acid sequence is presented in the 5' to 3' direction unless otherwise indicated. In some embodiments, a nucleic acid is or comprises natural nucleosides (e.g. adenosine, thymidine, guanosine, cytidine, uridine, deoxyadenosine, deoxythymidine, deoxyguanosine, and deoxycytidine); nucleoside analogs (e.g., 2-aminoadenosine, 2-thiothymidine, inosine, pyrrolo-pyrimidine, 3-methyl adenosine, 5-methylcytidine, 2-aminoadenosine, C5-bromouridine, C5-fluorouridine, C5-iodouridine, C5-propynyl-uridine, C5-propynyl-cytidine, C5-methylcytidine, 2-aminoadenosine, 7-deazaadenosine, 7-deazaguanosine, 8-oxoadenosine, 8-oxoguanosine, 0(6)-methylguanine, and 2-thiocytidine); chemically modified bases; biologically modified bases (e.g., methylated bases); intercalated bases; modified sugars (e.g., 2'-fluororibose, ribose, 2'-deoxyribose, arabinose, and hexose); and/or modified phosphate groups (e.g., phosphorothioates and 5'-N-phosphoramidite linkages).

(66) The terms “protein,” “peptide,” and “polypeptide” are used interchangeably herein, and refer to a polymer of amino acid residues linked together by peptide (amide) bonds. The terms refer to a protein, peptide, or polypeptide of any size, structure, or function. Typically, a protein, peptide, or polypeptide will be at least three amino acids long. A protein, peptide, or polypeptide may refer to an individual protein or a collection of proteins. One or more of the amino acids in a protein, peptide, or polypeptide may be modified, for example, by the addition of a chemical entity such as a carbohydrate group, a hydroxyl group, a phosphate group, a farnesyl group, an isofarnesyl group, a fatty acid group, a linker for conjugation, functionalization, or other modification, etc. A protein, peptide, or polypeptide may also be a single molecule or may be a multi-molecular complex. A protein, peptide, or polypeptide may be just a fragment of a naturally occurring protein or peptide. A protein, peptide, or polypeptide may be naturally occurring, recombinant, or synthetic, or any combination thereof. The term “fusion protein” as used herein refers to a hybrid polypeptide which comprises protein domains from at least two different proteins. One protein may be located at the amino-terminal (N-terminal) portion of the fusion protein or at the carboxy-terminal (C-terminal) protein thus forming an “amino-terminal fusion protein” or a “carboxy-terminal fusion protein,” respectively. A protein may comprise different domains, for example, a nucleic acid binding domain (e.g., the gRNA binding domain of Cas9 that directs the binding of the protein to a target site) and a nucleic acid cleavage domain or a catalytic domain of a nucleic-acid editing protein. In some embodiments, a protein comprises a proteinaceous part, e.g., an amino acid sequence constituting a nucleic acid binding domain, and an organic compound, e.g., a compound that can act as a nucleic acid cleavage agent. In some embodiments, a protein is in a complex with, or is in association with, a nucleic acid, e.g., RNA. Any of the proteins provided herein may be produced by any method known in the art. For example, the proteins provided herein may be produced via recombinant protein expression and purification, which is especially suited for fusion proteins comprising a peptide linker. Methods for recombinant protein expression and purification are well known, and include those described by Green and Sambrook, *Molecular Cloning: A Laboratory Manual* (4<sup>sup</sup>.th ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2012)), the entire contents of which are incorporated herein by reference.

(67) The term “subject,” as used herein, refers to an individual organism, for example, an individual mammal. In some embodiments, the subject is a human. In some embodiments, the subject is a non-human mammal. In some embodiments, the subject is a non-human primate. In some embodiments, the subject is a rodent. In some embodiments, the subject is a sheep, a goat, a cattle, a cat, or a dog. In some embodiments, the subject is a vertebrate, an amphibian, a reptile, a fish, an insect, a fly, or a nematode. In some embodiments, the subject is a research animal. In some embodiments, the subject is genetically engineered, e.g., a genetically engineered non-human subject. The subject may be of either sex and at any stage of development.

(68) The term “recombinant” as used herein in the context of proteins or nucleic acids refers to proteins or nucleic acids that do not occur in nature, but are the product of human engineering. For example, in some embodiments, a recombinant protein or nucleic acid molecule comprises an amino acid or nucleotide sequence that comprises at least one, at least two, at least three, at least four, at least five, at least six, or at least seven mutations as compared to any naturally occurring sequence.

#### DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS OF THE INVENTION

(69) The immune system is critical in preventing the outgrowth of cancers, and “immunosurveillance” exists to provide immunological resistance against cancer development (e.g., as described in Old et al., *Annu Rev Med* 1964; 15: 167-186; Burnet et al., *Prog Exp Tumor Res* 1970; 13: 1-27; and Graziano et al., *Cancer Treat Res* 2005; 123: 89-111, each of which is incorporated herein by reference).

(70) Despite the presence of immunosurveillance, cancers can develop in apparently immunocompetent animals and humans, due to the ability of cancer cells to evade immunosurveillance (e.g., as described in Hanahan et al., *Cell* 2000; 100: 57-70 and Zitvogel et al., *Nat Rev Immunol* 2006; 6: 715-727, each of which is incorporated herein by reference). The evasion of cancer cells from immunosurveillance occurs via various well-characterized mechanisms, including induction of T-cell tolerance by autochthonous tumors (e.g., as described in Willimsky et al., *Nature* 2005; 437: 141-146, incorporated herein by reference), cancer immunoediting (e.g., as described in Dunn et al., *Nat Immunol* 2002; 3: 991-998, incorporated herein by reference), and development of an immune suppressive cancer microenvironment (e.g., as described in Zou et al., *Nat Rev Cancer* 2005; 5: 263-274, incorporated herein by reference). Therapeutic cancer vaccines



or adoptive immunotherapy are being developed and tested as potential approaches to strengthen the immune responses after tumor arise in order to slow their progression and prevent their recurrence. Immunotherapeutic approaches, e.g., cancer vaccines have been described but are only partially successful (e.g., as described in Finn et al., *Nat Rev Immunol* 2003; 3: 630-641, incorporated herein by reference).

(71) Described herein are systems, methods, compositions, and kits for producing immunogenic peptides derived from tumor specific antigens (e.g., heteroclitic epitopes or cryptic epitopes) that may be used as cancer vaccines in vivo or ex vivo. Targeted mutations are introduced into tumor-specific antigens using a gene editing agent, e.g., a nucleobase editor comprising a programmable DNA binding domain (e.g., a catalytically-inactive Cas9 or Cas9 nickase) fused to a cytosine deaminase, to generate altered versions of peptides arising from the tumor-specific antigens (heteroclitic epitopes) or peptides arising from normally untranslated regions of the tumor-specific antigen genes (cryptic peptides). The heteroclitic peptides or cryptic peptides may be generated in vivo in a subject (e.g., a subject who has cancer) and presented to the adaptive immune system via the MHC class I or MHC class II pathway, which in turn induces a strong adaptive immune response, e.g., T cell response and B cell response. Such an adaptive immune response is antigen specific and is effective in reducing tumor growth and preventing metastasis.

(72) The advantage of the cancer vaccines of the present disclosure is that the vaccine (e.g., antigenic peptides derived from tumor-specific antigens) is generated from the genome and the proteome of the malignant cells in vivo and is highly personalized. The cancer vaccines described herein are also highly cancer-specific and do not induce unwanted immune response against “self,” since the immunogenic epitopes are derived from tumor-specific antigens. Further, the adaptive immune response induced by the cancer vaccine described herein confer “memory” to the immune system, promoting the immune system to efficiently recognize “neoepitopes” generated due to the highly mutagenic nature of the cancer genome, thus preventing metastasis and facilitate remission. To enhance the efficacy of the cancer vaccines described herein, combination therapies using an immune checkpoint inhibitor in conjunction with the cancer vaccine is also contemplated, aiming to enhance the tumor antigen specific immune response.

(73) The methods of producing endogenous cancer vaccines in vivo are enabled by the targeted nucleobase editing technology described herein. Such base editing technology is described in the art, e.g., in U.S. Pat. No. 9,068,179, US Patent Application Publications US 2015/0166980, US 2015/0166981, US 2015/0166982, US20150166984, and US20150165054, and US Provisional Applications, U.S. Ser. No. 62/245,828, filed Oct. 23, 2015; 62/279,346, filed Jan. 15, 2016; 62/311,763, filed Mar. 22, 2016; 62/322,178, filed Apr. 13, 2016, 62/357,352, filed Jun. 30, 2016, U.S. Pat. No. 62,370,700, filed Aug. 3, 2016; 62/398,490, filed Sep. 22, 2016; 62/408,686, filed Oct. 14, 2016; PCT Application PCT/US2016/058344, filed Oct. 22, 2016, U.S. patent application Ser. No. 15/311,852, filed Oct. 22, 2016; and in Komor et al., *Nature*, Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage, 533, 420-424 (2016), the entire contents of each of which are incorporated herein by reference.

(74) Immunogenic Peptides or Epitopes

(75) Some aspects of the present disclosure provide immunogenic peptides or epitopes derived from tumor-specific antigens and how these peptides or epitopes elicit tumor-specific immune response. A large number of proteins that specifically express in tumor cells or are upregulated in tumor cells have been identified (Hassane et al., *Holland-Frei Cancer Medicine*. 6th edition). The known tumor specific antigens are classified into different classes.

(76) a) Cancer-testis antigens: The first TAAs ever identified that can be recognized by T cells belong to this class, which was originally called cancer-testis (CT) antigens because of the expression of its members in histologically different human tumors and, among normal tissues, only in spermatocytes/spermatogonia of testis and, occasionally, in placenta. Since the cells of testis do not express class I and II HLA molecules, these antigens cannot be recognized by T cells in normal tissues and can therefore be considered as immunologically tumor-specific. Well-known examples of CT antigens are the MAGE family members or NY-ESO-1.

(77) b) Differentiation antigens: These TAAs are shared between tumors and normal tissue from which the tumor arose; most are found in melanomas and normal melanocytes. Many of these melanocyte lineage-related proteins are involved in the biosynthesis of melanin and are therefore not tumor specific but nevertheless are widely used for cancer immunotherapy. Examples include, but are not limited to, tyrosinase and Melan-A/MART-1 for melanoma, and PSA for prostate cancer.

(78) c) Overexpressed TAAs: Genes encoding widely expressed TSAs have been detected in histologically different types of tumors as well as in many normal tissues, generally with lower expression levels. It is possible that many of the epitopes processed and potentially presented by normal tissues are below the threshold level for T-cell recognition, while their overexpression in tumor cells can trigger an anticancer response by breaking previously established tolerance. Examples of this class of TAAs are Her-2/neu, Survivin, Telomerase and WT1.

(79) d) Tumor specific antigens: These unique TAAs arise from mutations of normal genes (such as  $\beta$ -catenin, CDK4, etc.). Some of these molecular changes are associated with neoplastic transformation and/or progression. Tumor specific antigens are generally able to induce strong immune responses without bearing the risk for autoimmune reactions against normal tissues. On the other hand, these TAAs are in most cases only relevant to the exact tumor on which they were identified and are usually not shared between many individual tumors.

(80) e) TAAs arising from abnormal post-translational modifications: Such TSAs may arise from proteins which are neither specific nor overexpressed in tumors but nevertheless become tumor associated by posttranslational processes primarily active in tumors. Examples for this class arise from altered glycosylation patterns leading to novel epitopes in

tumors (e.g., MUC1).

(81) f) Oncoviral proteins: These TSAs are viral proteins that may play a critical role in the oncogenic process, and because they are foreign (not of human origin), they can evoke a T-cell response. Examples of such proteins are the human papilloma type 16 virus proteins, E6 and E7, which are expressed in cervical carcinoma.

(82) TAAs are a starting point for the development of a tumor vaccine. The methods for identifying and characterizing the TAAs are based on the use of cytotoxic T lymphocytes (CTL) that can be isolated from patients or healthy subjects, or they are based on the generation of differential transcription profiles or differential peptide expression patterns between tumors and normal tissues.

(83) In some embodiments, the tumor-specific antigen is expressed in a broad range of different types of cancers. In some embodiments, the tumor-specific antigen is expressed only in one or a few types of cancers. The anti-cancer immune response described herein is antigen-specific. As such, an immune response induced by a tumor-specific antigen is specific to cancer types where the said antigen is expressed. Non-limiting, exemplary tumor-specific antigens that may be edited to generate immunogenic epitopes are provided in Tables 1-3. It is appreciated that the examples are for illustration purpose only, and the methods described herein may be applied to any tumor-specific antigen.

(84) In some embodiments, the immunogenic peptide or epitope is a portion of the tumor-specific antigen. For example, the immunogenic peptide or epitope may be a portion of the tumor-specific antigen that is 5-40 amino acids long. In some embodiments, the immunogenic peptide or epitope is 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 amino acids long. In some embodiments, the immunogenic peptide or epitope comprises modifications, e.g., amino acid substitutions (also termed “heteroclitic epitopes”), as compared to the native sequence in the tumor specific antigen. In some embodiments, the immunogenic peptide or epitope comprises more than one amino acid substitutions (e.g., 2, 3, 4, 5, or more) compared to the native sequence of the tumor-specific antigen it is derived from. In some embodiments, a heteroclitic peptide or epitope may be at least 60%, at least 70%, at least 80%, at least 90%, at least 98%, or at least 99% identical to the native sequence that it is derived from. In some embodiments, a heteroclitic peptide or epitope is 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99% identical to the native sequence that it is derived from.

(85) In some embodiments, a heteroclitic peptide or epitope is more immunogenic than a peptide of its native sequence. For example, a heteroclitic epitope may be at least 30% more immunogenic (i.e., induces a stronger immune response) than its corresponding native peptide. In some embodiments, a heteroclitic epitope may be at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 100%, at least 2-fold, at least 3-fold, at least 4-fold, at least 5-fold, at least 6-fold, at least 7-fold, at least 8-fold, at least 9-fold, at least 10-fold, at least 20-fold, at least 30-fold, at least 40-fold, at least 50-fold, at least 60-fold, at least 70-fold, at least 80-fold, at least 90-fold, at least 100-fold, or more immunogenic than its corresponding native peptide.

(86) In some embodiments, the immunogenic peptide or epitope is a cryptic peptide or epitope, e.g., generated from translation of a non-coding region of the tumor specific antigen gene or translation of a different reading frame of a coding region of the tumor specific antigen. A cryptic peptide or epitope may be more immunogenic (i.e., induces a stronger immune response) than any native peptide derived from the tumor associated antigen. For example, a cryptic peptide or epitope may be at least 30% more immunogenic than any native peptide derived from the tumor associated antigen. In some embodiments, a cryptic peptide or epitope may be at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 100%, at least 2-fold, at least 3-fold, at least 4-fold, at least 5-fold, at least 6-fold, at least 7-fold, at least 8-fold, at least 9-fold, at least 10-fold, at least 20-fold, at least 30-fold, at least 40-fold, at least 50-fold, at least 60-fold, at least 70-fold, at least 80-fold, at least 90-fold, at least 100-fold, or more immunogenic than any native peptide derived from the tumor associated antigen. One skilled in the art is familiar with how to assess the immune response induced by an antigen, e.g., measuring antibody titers.

(87) Tumor specific antigens from which antigenic epitopes (e.g., heteroclitic epitopes and cryptic epitopes) may be derived are provided in Tables 1-3. (HLA—human leukocyte antigen type)

(88) TABLE-US-00005 TABLE 1 Tumor associated antigen - Differentiation HLA SEQ Gene/ frequency ID  
protein Tumor HLA (%) Peptide NO Position CEA gut A2 44 YLSGANLNL 340 605-613 carcinoma A2 44  
IMIGVLVG 341 691-699 A2 44 GVLVGVALI 342 694-702 A2 44 LLTFWNPPT 343 24-33 A3 22 HLFYGSWYK 344  
61-69 A24 20 QYSWFVNGTF 345 268-277 A24 20 TYACFVSNL 346 652-660 DR3 21 AYVCGIQNSVSANRS 347  
568-582 DR4 24 DTGFYTLHVIKSDLVNEEA 348 116-140 TGQFRV DR4 24 YSWRINGIPQQHTQV 349 625-639  
DR7 25 TYYRPGVNL SLSC 350 425-437 DR7 25 EIIYPNASLLION 351 99-111 DR9 3 YACFVSNLATGRNNS 352  
653-667 DR11 25 LWVNNQSLPVSP 353 177-189, 355-367 DR13 19 LWVNNQSLPVSP 354 177-189, 355-367  
DR14 6 LWVNNQSLPVSP 355 177-189, 355-367 DR14 6 EIIYPNASLLIQN 356 99-111 DR14 6 NSIVKSITVSASG  
357 666-678 gp100/ melanoma A2 44 KTWGQYWQV 358 154-162 Pmel17 A2 44 (A)MLGTHTMEV 359  
177(8)-186 A2 44 ITDQVPFSV 360 209-217 A2 44 YLEPGPVT 361 280-288 A2 44 LLDGTATLRL 362 457-  
466 A2 44 VLYRYGSFSV 363 476-485 A2 44 SLADTNSLAV 364 570-579 A2 44 RLMKQDFSV 365 619-627 A2 44  
RLPRIFCSC 366 639-647 A3 22 LIYRRRLMK 367 614-622 A3 22 ALLAVGATK 368 17-25 A3 22 IALNFPQSQK 369  
86-95 A3 22 RSYVPLAHR 370 195-202 and 191 or 192 A3 22 ALNFPQSQK 371 87-95 A11 13 ALNFPQSQK 372  
87-95 A24 20 VYFFLPDHL 373 intron 4 A32 8 RTKQLYPEW 374 40-42 and 47-52 A68 8 HTMEVTVYHR 375  
182-191 B7 17 SSPGCQPPA 376 529-537 B35 20 VPLDCVLYRY 377 471-480 B35 20 LPHSSSHWL 378 630-638 Cw8  
— SNDGPTLI 379 71-78 DQ6 63 GRAMLGTHTMEVTVY 380 175-189 DR4 24 WNRQLYPEWTEAQRDL 381 44-59

DR7 25 TTEWVETTAPEIPE 382 420-437 DR7 25 TGRAMLGTHTMEVTYH 383 174-190 DR53 49  
 GRAMLGTHTMEVTYH 384 175-189 mammagl breast A3 22 PLENVISK 385 23-31 obin-A cancer Melan-A/  
 melanoma A2 44 (E)AAGIGILTV 386 26(27)-35 MART-1 A2 44 ILTVILGVL 387 32-40 B35 20  
 EAAGIGILTV 388 26-35 B45 2 AEAAAGIGIL(T) 389 24-33(34) Cw7 41 RNGYRALMDKS 390 51-61 DP5  
 3 YTTAEAAAGIGILTVILGV 391 21-50 LLLIGCWYCRR DQ6 63 EEAAGIGILTVI 392 25-36 DR1 18  
 AAGIGILTVILGVL 393 27-40 DR1 18 APPAYEKLpSAEQf 394 100-111 DR3 21 EEAAGIGILTVI 395 25-36 DR4 24  
 RNGYRALMDKSLHVGTC 396 51-73 ALTRR DR11 25 MPREDAHFIYGYPKKGHH 397 1-20 S DR52 41  
 KNCEPVVPNAPPAYEKL 398 91-110 E NY-BR-1 breast A2 44 SLSKILDTV 399 904-912 cancer ERBB2 breast  
 A2 44 IVWELMTFGA 400 911-921 cancer OA1 melanoma A24 20 LYSACFWWL 401 126-134 PAP prostate A2 44  
 FLFLFFWL 402 18-26 cancer A2 44 TLMSAMTNL 403 112-120 A2 44 ALDVYNGLL 404 299-307 PSA prostate A2  
 44 FLTPKKLQCV 405 165-174 carcinoma A2 44 VISNDVCAQV 406 178-187 RAB38/ melanoma A2 44  
 VLHWDPETV 407 50-58 NY-MEL-1 TRP-1/ melanoma A31 5 MSLQRQFLR 408 alt. ORF gp75 DR4 24  
 ISPNSVFSQWRVVCDSLED 409 277-297 YD DR15 20 SLPYWNFATG 410 245-254 DR17 21  
 SQWRVVCDSLEDYDT 411 284-298 TRP-2 melanoma A2 44 SVYDFFVWL 412 180-188 A2 44 TLDSQVMSL 413  
 360-368 A31 5 LLGPGRPYR 414 197-205 A33 5 LLGPGRPYR 415 197-205 Cw8 — ANDPIFVVL 416 387-395 DR3  
 21 QCTEVRADTRPWSGP 417 60-74 DR15 20 ALPYWNFATG 418 241-250 tyrosinase melanoma A1 26  
 KCDICTDEY 419 243-251 A1 26 SSDYVIPIGT 420 146-156 A2 44 MLLAVLYCL 421 1-9 A2 44 CLLWSFQ TSA  
 422 8-17 A2 44 YMDGTMSQV 423 369-377 A24 20 AFLPWHRFL 424 206-214 A24 20 IYMDGTADFSF 425 368-  
 373 and 336-340e A26 8 QCSGNFMGF 426 90-98 B35 20 TPRLPSSADVEF 427 309-320 B35 20 LPSSADVEF 428  
 312-320 B38 5 LHAFVDSIF 429 388-397 B44 21 SEIWRDIDFd 430 192-200 DR4 24 QNILLSNAPLGPQFP 431 56-  
 70 DR4 24 SYLQSDPDSFQD 432 450-462 DR15 20 FLLHAFVDSIFEQWLQRH 433 386-406 RP WT1 testis, A1  
 26 TSEKRPFMCAY 434 317-327 ovary, A24 20 CMTWNQMNL 435 235-243 bone DP5 3 LSHLQMHSRKH 436 337-  
 347 marrow, DP5 3 KRYFKLSHLQMHSRKH 437 332-347 spleen DR4 24 KRYFKLSHLQMHSRKH 438 332-347  
 CD33 leukemia A2 44 AIISGDSPV 439 65-73  
 (89) TABLE-US-00006 TABLE 2 Tumor specific antigen - Tumor specific HLA SEQ frequency ID  
 Gene/protein HLA (%) Peptide NO Position BAGE-1 Cw16 7 AARAVFLAL 440 1-10 D393-CD20n DR4 24  
 KPLFRRMSSLELVIA 441 28-42 Cyclin-A1 A2 44 FLDRFLSCM 442 227-235 A2 44 SLIAAAAFCLA 443 341-351  
 GAGE-1,2,8 Cw6 18 YRPRPRRY 444 9-16 GAGE-3,4,5,6,7 A29 6 YYWPRPRRY 445 10-18 GnTVf A2 44  
 VLPDVFIRC(V) 446 intron HERV-K-MEL A2 44 MLAVISCAV 447 1-9 KK-LC-1 B15 13 RQKRILVNL 448 76-84  
 KM-HN-1 A24 20 NYNNFYRFL 449 196-204 A24 20 EYSKECLKEF 450 499-508 A24 20 EYLSLSDKI 451 770-778  
 LAGE-1 A2 44 MLMAQEALAF 452 alt. ORF (1-11) A2 44 SLLMWITQC 453 157-165 A31 5 LAAQERRVPR  
 454 alt. ORF (18-27) A68 8 ELVRRILSR 455 103-111 B7 17 APRGVMAV 456 alt. ORF (46-54) DP4 75  
 SLLMWITQCFLPVF 457 157-170 DR3 21 QGAMLAAQERRVPRAAEVPR 458 alt. ORF (14-33) DR4 24  
 AADHRQLQLSISSCLQQL 459 139-156 DR11 25 CLSRRPWKRSWSAGSCPGMPHL 460 alt. ORFT (81-102)  
 DR12 5 CLSRRPWKRSWSAGSCPGMPHL 461 alt. ORFT (81-102) DR13 19 ILSRDAAPLPRPG 462 108-120  
 DR15 20 AGATGGRGPRGAGA 463 37-50 LY6K A24 20 RYCNLEGPI 464 119-128 DP5 3  
 KWTEPYCVIAAVKIFPRFFMVAKQ 465 61-84 DR15 20 KCCKIRYCNLEGPPINSSVF 466 114-133 MAGE-A1 A1  
 26 EADPTGHSY 467 161-169 A2 44 KVLEYVIKV 468 278-286 A3 22 SLFRAVITK 469 96-104 A68 8  
 EYDGREHSA 470 222-231 B7 17 RVRFFFPSL 471 289-298 B35 20 EADPTGHSY 472 161-169 B37 3  
 REPVTKAEML 473 120-129 B44 21 KEADPTGHSY 474 160-169 B53 2 DPARYEFLW 475 258-266 B57 8  
 ITKKVADLVGF 476 102-112 Cw2 10 SAFPTTINF 477 62-70 Cw3 17 SAYGEPRKL 478 230-238 Cw7 41  
 RVRFFFPSL 479 289-298 Cw16 7 SAYGEPRKL 480 230-238 DP4 75 TSCILESLFRAVITK 481 90-104 DP4 75  
 PRALAEYSYVKVLEY 482 268-282 DR13 19 FLLLYKRYAREPVTKAE 483 112-127 DR15 20 EYVIKVSARVRF 484  
 281-292 MAGE-A2 A2 44 YLQLVFGIEV 485 157-166 A24 20 EYLQLVFGI 486 156-164 B37 3 REPVTKAEML  
 487 127-136 Cw7 41 EGDCAPEEK 488 212-220 DR13 19 LLKYRAREPVTKAE 489 121-134 MAGE-A3 A1 26  
 EVDPIGHLY 490 168-176 A2 44 FLWGPRALVd 491 271-279 A2 44 KVAELVHFL 492 112-120 A24 20 TFPDLESEF  
 493 97-105 A24 20 VAELVHFL 494 113-121 B18 6 MEVDPIGHLY 495 167-176 B35 20 EVDPIGHLY 496 168-  
 176 B37 3 REPVTKAEML 497 127-136 B40 6 AELVHFL 498 114-122 B44 21 MEVDPIGHLY 499 167-176  
 B52 5 WQYFFPVIF 500 143-151 Cw7 41 EGDCAPEEK 501 212-220 DP4 75 KLLTQHVFVQENYLEY 502 243-258  
 DP4 75 RKVAELVHFLLLKYR 503 111-125 DQ6 63 KLLTQHVFVQENYLEY 504 243-258 DR1 18  
 ACYEFLWGPRALVETS 505 267-282 DR4 24 RKVAELVHFLLLKYR 506 111-125 DR4 24 VIFSKASSSLQL 507  
 149-160 DR7 25 VIFSKASSSLQL 508 149-160 DR7 25 VFGIELMEVDPIGHL 509 161-175 DR11 25  
 GDNQIMPKAGLLIIV 510 191-205 DR11 25 TSYVKVLHHMVKISG 511 281-295 DR13 19 RKVAELVHFLLLKYRA  
 512 111-126 DR13 19 FLLLYKRYAREPVTKAE 513 119-134 MAGE-A4 A1 26 EVDPASNTY 514 169-177 A2 44  
 GYDGREHTV 515 230-239 A24 20 NYKRCFPVI 516 143-151 B37 3 SESLKMIF 517 156-163 MAGE-A6 A34  
 1 MVKISGGPR 518 290-298 B35 20 EVDPIGHVY 519 168-176 B37 3 REPVTKAEML 520 127-136 Cw7 41  
 EGDCAPEEK 521 212-220 Cw16 7 ISGPRISY 522 293-301 DR13 19 LLKYRAREPVTKAE 523 121-134 MAGE-  
 A9 A2 44 ALSVMGVYV 524 223-231 MAGE-A10 A2 44 GLYDGMEHLI 525 254-262 B53 2 DPARYEFLW 526  
 290-298 MAGE-A12 m A2g 44 FLWGPRALV 527 271-279 Cw7 41 VRIGHLYIL 528 170-178 Cw7 41 EGDCAPEEK  
 529 212-220 DP4 75 REPVTKAEMLGSVIR 530 127-141 DR13 19 AELVHFLLLKYRAR 531 114-127 MAGE-C1 A2  
 44 ILFGISLREV 532 959-968 A2 44 KVVEFLAML 533 1083-1091 DQ6 63 SSALLSIFQSSPE 534 137-149 DQ6 63

SFSLTSL 535 450-458 DR15 20 VSSFFSYTL 536 779-787 MAGE-C2 A2 44 LLFGLALIEV 537 191-200 A2 44 ALKDVEERV 538 336-344 B44 21 SESIKKKVL 539 307-315 B57 8 ASSTLYLVF 540 42-50 DR15 20 SSTLYLVFSPSSFST 541 43-57 mucink PDTRPAPGSTAPPAHGVTS 542 NA88-A B13 6 QGQHFLQKV 543 A2 44 SLLMWITQC 544 157-165 NY-ESO-1/ A2 44 MLMAQEALAF 545 alt. ORF LAGE-2 (1-11) A24 20 YLAMPFATPME 546 91-101 A31 5 ASGPGGGAPR 547 53-62 A31 5 LAAQERRVPR 548 alt. ORF (18-27) A68 8 TVSGNILTIR 549 127-136 B7 17 APRGPHGGAASGL 550 60-72 B35 20 MPFATPMEAE 551 94-104 B49 KEFTVSGNILT 552 124-135 B51 12 MPFATPMEA 553 94-102 B52 5 FATPMEAE 554 96-104 C12 12 FATPMEAE 555 96-106 Cw3 17 LAMPFATPM 556 92-100 Cw6 18 ARGPE SRL 557 80-88 DP4 75 SLLMWITQCFLPVF 558 157-170 DP4 75 LLEFY LAMPFATPMEAE LARRSLAQ 559 87-111 DR1 18 LLEFY LAMPFATPMEAE LARRSLAQ 560 87-111 DR1 18 EFY LAMPFATPM 561 89-100 DR1 18 PGVLLKEFTVSGNILTIRLTAAADR 562 119-143 DR2 25 RLLEFY LAMPFA 563 86-97 DR3 21 QGAMLA AQERRVPRAAEVPR 564 alt. ORF (14-33) DR4 24 PFATPMEAE LARR 565 95-107 DR4 24 PGVLLKEFTVSGNILTIRLT 566 119-138 DR4 24 VLLKEFTVSG 567 121-130 DR4 24 AADRQLQLSISSCLQQL 568 139-156 DR4 24 LLEFY LAMPFATPMEAE LARRSLAQ 569 87-111 DR52b 25 LKEFTVSGNILTIRLT 570 123-137 DR7 25 PGVLLKEFTVSGNILTIRLTAAADR 571 119-143 DR7 25 LLEFY LAMPFATPMEAE LARRSLAQ 572 87-111 DR8 4 KEFTVSGNILT 573 124-134 DR9 3 LLEFY LAMPFATPM 574 87-100 DR15 20 AGATGGRGPRGAGA 575 37-50 SAGE A24 20 LYATVIHDI 576 715-723 Sp17 A1 26 ILDSSEEDK 577 103-111 SSX-2 A2 44 KASEKIFYV 578 41-49 DP1 14 EKIQKAFDDIAKYFSK 579 19-34 DR1 18 FGRLQGISP KI 580 101-111 DR3 21 WEKMKASEKIFYVYMKRK 581 37-54 DR4 24 KIFYVYMKRKYEAMT 582 45-59 DR11 25 KIFYVYMKRKYEAM 583 45-58 SSX-4 DP10 2 INKTSGPKRGKHAWTHRLRE 584 151-170 DR3 21 YFSKKEWEKMKSSSEKIVYVY 585 31-50 DR8 4 MKLNYEVMTKLGFKVTLPPF 586 51-70 DR8 4 KHAWTHRLRERKQLVVYEEI 587 161-180 DR11 25 LGFKVTLPPFMRSKRAADF 588 61-80 DR15 20 KSSEKIVYVYMKLNYEVMTK 589 41-60 DR52 41 KHAWTHRLRERKQLVVYEEI 590 161-180 survivin/ A2 44 ELTLGEFLKL 591 96-106 BIRC5 A24 20 AYACNTSTL 592 intron B2 TAG-1 A2 44 SLGWLFLL 593 78-86 B8 14 LSRLSNRLL 594 42-50 TAG-2 B8 14 LSRLSNRLL 595 42-50 TRAG-3 DR1 18 CEFHACWPAFTVLGE 596 34-48 DR4 24 CEFHACWPAFTVLGE 597 34-48 DR7 25 CEFHACWPAFTVLGE 598 34-48 TRP2-INT2g A68 8 EVISCKLIK 599 intron 2 XAGE-1b/ A2 44 RQKKIRIQL 600 21-29 GAGED2a DR4 24 HLGSRQKKIRIQLRSQ 601 17-32 DR9 3 CATWKVICKSCISQTPG 602 33-49 BCR-ABL A2 44 SSKALQRPV 603 926-934 (b3a2) B8 14 GFKQSSKAL 604 922-930 DR4 24 ATGFKQSSKALQRPVAS 605 920-936 DR9 3 ATGFKQSSKALQRPVAS 606 920-936

(90) TABLE-US-00007 TABLE 3 Overexpressed tumor specific antigen HLA SEQ Normal tissue Freq ID Gene/Protein expression HLAA (%) Peptide NO Position adipophilin adipocytes, macrophages A2 44 SVASTITGV 607 129-137 AIM-2 ubiquitous (low level) A1 26 RSDSGQQARY 608 intron ALDH1A1 mucosa, keratinocytes A2 44 LLYKLADLI 609 88-96 BCLX (L) ubiquitous (low level) A2 44 YLNDHLEPWI 610 173-182 BING-4 ubiquitous (low level) A2 44 CQWGRLWQL 611 ORF2 CALCA thyroid A2 44 VLLQAGSLHA 612 16-25 CD45 proliferating cells, A24 20 KFLDALISL 613 556-564 testis, multiple tissues (low level) CD274 multiple tissues A2 44 LLNAFTVT 614 15-23 (lung, heart, dendritic cells, etc.) and induced by IFN- $\gamma$  CPSF ubiquitous (low level) A2 44 KVHPVIWSL 615 250-258 A2 44 LMLQNALT 616 1360-1369 cyclin D1 ubiquitous (low level) A2 44 LLGATCMFV 617 101-109 DR4 24 NPPSMVAAGSVVAAV 618 198-212 DKK1 testis, prostate, A2 44 ALGGHPLLGV 619 20-29 mesenchymal stem cells ENAH (hMen a) breast, prostate stroma A2 44 TMNGSKSPV 620 502-510 and epithelium of colon-rectum, pancreas, endometrium EpCAM epithelial cells A24 20 RYQLDPKFI 621 173-181 EphA3 many DR11 25 DVTFNIICKKCG 622 356-367 EZH2 ubiquitous (low level) A2 44 FMVEDETVL 623 120-128 A2 44 FINDEIFVEL 624 165-174 A24 20 KYDCFLHPF 625 291-299 A24 20 KYVGIEREM 626 735-743 FGF5 brain, kidney A3 22 NTYASPRFKF 627 172-176 and 217-220 glypican-3 placental and A2 44 FVGEFFTDV 628 144-152 multiple tissues A24 20 EYILSLEEL 629 298-306 G250/MN/CAIX stomach, liver, A2 44 HLSTAFARV 630 254-262 pancreas HER-2/neu ubiquitous (low level) A2 44 KIFGSLAFL 631 369-377 A2 44 IISAVVGIL 632 654-662 A2 44 ALCRWGLLL 633 5-13 A2 44 ILHNGAYSL 634 435-443 A2 44 RLLQETELV 635 689-697 A2 44 VVLGVVFGI 636 665-673 A2 44 YMIMVKCWMI 637 952-961 A2 44 HLYQGCQVV 638 48-56 A2 44 YLVPQQGFFC 639 1023-1032 A2 44 PLQPEQLQV 640 391-399 A2 44 TLEEITGYL 641 402-410 A2 44 ALIHHNTHL 642 466-474 A2 44 PLTSIISAV 643 650-658 A3 22 VLRENTSPK 644 754-762 A24 20 TYLPTNASL 645 63-71 HLA-DOB B lymphocytes, A2 44 FLLGLIFLL 646 232-240 monocytes, blood cells, adrenals, . . . Hepsin kidney, liver, skin, A2 44 SLLSGDWVL 647 191-199 . . . A2 44 GLQLGVQAV 648 229-237 A2 44 PLTEYIQPV 649 268-276 IDO1 lymph nodes, placenta, A2 44 ALLEIASCL 650 199-207 and many cell types in the course of inflammatory response IGF2B3 ubiquitous (low level) A2 44 NLSSAEVVV 651 515-523 A3 44 RLLVPTQFV 652 199-207 IL13Ralpha2 A2 44 WLPFGFILI 653 345-353 Intestinal liver, intestine, B7 17 SPRWWPTCL 654 alt. ORF carboxyl kidney esterase alpha- liver A2 44 GVALQTMKQ 655 542-550 foetoprotein A2 44 FMNKFIYEI 656 158-166 DR13 19 QLAVSVILRV 657 364-373 Kallikrein 4 prostate and ovarian A2 44 FLGYLILGV 658 11-19 cancer DP4 75 SVSESDTIRSISIAS 659 125-139 cancer DR4 24 LLANGRMPTVLQCVN 660 155-169 DR7 25 RMPTVLQCVNVS 661 160-174 KIF20A ubiquitous (low level) A2 44 LLSDDDVVV 662 12-20 A2 44 AQPDTAPLPV 663 284-293 A2 44 CIAEQYHTV 664 809-817 Lengsin eye lens and low level A2 44 FLPEFGISSA 665 270-279 in multiple tissues M-CSF liver, kidney B35 20

LPVGLPEQGEY 666 alt. ORF MSCSP endothelial cells, DR11 25 VGQDVSVLFRVTGALQ 667 693-708 chondrocytes, smooth muscle cells mdm-2 ubiquitous (brain, A2 44 VLFYLGQY 668 53-60 muscle, lung) Meloe ubiquitous (low level) A2 44 TLNDECWPA 669 36-44 DQ2 41 ERISSTLNDECWPA 670 31-44 DQ6 63 FGRLQGISPKI 671 32-44 DR1 18 TSREQFLPSEGAA 672 11-23 DR11 25 CPPWHPSEIRISSTL 673 24-37 Midkine ubiquitous (low level) A2 44 ALLALTSAPV 674 13-21 A2 44 AQCQETIRV 675 114-122 DR4 24 LTLALLALTSAPVAK 676 9-23 MMP-2 ubiquitous A2 44 GLPPDVQVRVH 677 560-568 MMP-7 ubiquitous (low level) A3 22 SLFPNSPKWTSK 678 96-107 MUC1 glandular epithelia A2 44 STAPPVHNV 679 950-958 A2 44 LLLLTVLTV 680 12-20 DR3 21 PGSTAPPAHGVT 681 repeated region MUC5AC surface mucosal A24 20 TCQPTCRSL 682 716-724 cells, respiratory tract, and stomach epithelia p53 ubiquitous (low level) A2 44 LLGRNSFEV 683 264-272 A2 44 RMPEAAPPV 684 65-73 B46 0.1 SQKTYQGSY 685 99-107 DP5 3 PGTRVRAMAIYKQ 686 153-165 DR14 6 HLIRVEGNLRVE 687 193-204 PAX5 hemopoietic system A2 44 TLPGYPPHV 688 311-319 PBF ovary, pancreas, spleen, B55 4 CTACRWKKACQR 689 499-510 liver PRAME testis, ovary, A2 44 VLDGLDVLL 690 100-108 endometrium, A2 44 SLYSFPEPEA 691 142-151 adrenals A2 44 ALYVDSLFFL 692 300-309 A2 44 SLLQHLIGL 693 425-433 A24 20 LYVDSLFFLC 694 301-309 PSMA prostate, CNS, liver A24 20 NYARTEDFF 695 178-186 RAGE-1 retina A2 44 LKLSGVVRL 696 352-360 A2 44 PLPPARNGGLG 697 32-40 B7 17 SPSSNRIRNT 698 11-20 RGS5 heart, skeletal A2 44 LAALPHSCL 699 5-13 muscle, pericytes A3 22 GLASFKSFLK 700 74-83 RhoC ubiquitous (low level) A3 22 RAGLQVRKNK 701 176-185 RNF43 A2 44 ALWPWLLMA(T) 702 11-19(20) A24 20 NSQPVWLCL 703 721-729 RU2AS testis, kidney, bladder B7 17 LPRWPPPQL 704 antisense secernin 1 ubiquitous A2 44 KMDAEHPPEL 705 196-204 SOX10 ubiquitous (low level) A2 44 AWISKPPGV 706 332-340 A2 44 SAWISKPPGV 707 331-340 STEAP1 prostate A2 44 MIAVFLPIV 708 292-300 A2 44 HQQYFYKIPILVINK 709 102-116 survivin ubiquitous A2 44 ELTLGEFLKL 710 95-104 ubiquitous DR1 18 TLGEFLKLDREKAKN 711 97-111 Telomerase testis, thymus, A2 44 ILAKFLHWLE 712 540-548 bone marrow, A2 44 RLVDFFLLV 713 865-873 lymph nodes DR7 25 RPGLLGASVLGLDDI 714 672-686 DR11 25 LTDLQPYMRQFVAHL 715 766-780 hTERT WLFYRKS(R) 716 572-581 (572) TPBG multiple tissues A2 44 RLARLALVL 717 17-25 (esophagus, bladder, etc.) VEGF ubiquitous (low level) B27 7 SRFGGAVVR 718 -i WT1 testis, ovary, bone A1 26 TSEKRPFMCAY 719 317-327 marrow, spleen A24 20 CMTWNQMNL 720 235-243 DP5 3 LSHLQMHSRKH 721 337-347 DP5 3 KRYFKLSHLQMHSRKH 722 332-347 DR4 24 KRYFKLSHLQMHSRKH 723 332-347

(91) The identification of heteroclitic peptides are described in the art. For example, in previous studies (Selby, et al., *J. Immunol.*, 162(2):669 (1999), Skipper, et al., *J. Exp. Med.* 183:527 (1996), the entire contents of each of which are incorporated herein by reference), heteroclitic epitopes were fortuitously identified by eluting naturally occurring mutant peptides from melanoma cells, or by systematically screening a large number of epitopes consisting of substitutions at almost every position in the epitope (Zaremba, et al., *Cancer Research*, 57:4570 (1997), Loftus, et al., *Cancer Research* 58:2433 (1998), Blake, et al., *J. Exp. Med.* 18:121 (1996), the entire contents of each of which are incorporated herein by reference). Alternatively, heteroclitic epitopes were identified by screening random combinatorial peptide libraries which also has required the arduous synthesis and screening of large numbers of peptides (Pinilla, et al., *Current Opinion in Immunology* 11:193-202 (1999), the entire contents of each of which are incorporated herein by reference). Genetic approaches, such as screening of DNA expression libraries, have provided another method for generating CTL epitopes and analogs (Boon, et al., *Annu. Rev. Immunol.* 12:337-65 (1994), Gavin, et al., *Eur. J. Immunol.* 24(9):2124-33 (1994), the entire contents of each of which are incorporated herein by reference).

(92) Generating Cancer Vaccine in Tumor Cells

(93) Some aspects of the present disclosure provide systems, compositions, and methods of editing genes encoding tumor specific antigens in vivo (e.g., in tumor cells in a subject) or ex vivo (e.g., in isolated tumor cells) to introduce mutations in the genes encoding tumor specific antigens. In some embodiments, such mutations lead to the production of heteroclitic peptides that are more immunogenic than native peptides of the tumor-specific antigen. In some embodiments, such mutations lead to the translation of a non-coding region of the tumor specific antigen, which results in cryptic peptides that are more immunogenic than any native peptides from the tumor specific antigen.

(94) The gene editing methods described herein, rely on nucleobase editors as described in., in U.S. Pat. No. 9,068,179, US Patent Application Publications US 2015/0166980, US 2015/0166981, US 2015/0166982, US20150166984, and US20150165054, and US Provisional Applications, U.S. Ser. No. 62/245,828, filed Oct. 23, 2015; 62/279,346, filed Jan. 15, 2016; 62/311,763, filed Mar. 22, 2016; 62/322,178, filed Apr. 13, 2016, 62/357,352, filed Jun. 30, 2016, U.S. Pat. No. 62,370,700, filed Aug. 3, 2016; 62/398,490, filed Sep. 22, 2016; 62/408,686, filed Oct. 14, 2016; PCT Application PCT/US2016/058344, filed Oct. 22, 2016, U.S. patent application Ser. No. 15/311,852, filed Oct. 22, 2016; and in Komor et al., *Nature*, Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage, 533, 420-424 (2016), the entire contents of each of which are incorporated herein by reference.

(95) The nucleobase editors are highly efficient at precisely editing a target base in any of the tumor associated antigen genes described herein, and a DNA double strand break is not necessary for the gene editing, thus reducing genome instability and preventing possible oncogenic modifications that may be caused by other genome editing methods. The nucleobase editors described herein may be programmed to target and modify a single base. In some embodiments, the target base is a cytosine (C) base and may be converted to a thymine (T) base via deamination by the nucleobase editor.

(96) To edit the polynucleotide encoding a tumor associated antigen, the polynucleotide is contacted with a nucleobase

editor as described herein, the tumor-associated antigen encoding polynucleotide is contacted with a nucleobase editor and a guide nucleotide sequence, wherein the guide nucleotide sequence targets the nucleobase editor the target base (e.g., a C base) in the tumor-associated antigen encoding polynucleotide.

(97) In some embodiments, the tumor-associated antigen encoding polynucleotide is the tumor-associated antigen gene locus in the genomic DNA of a cell (e.g., a tumor cell). In some embodiments, the tumor cell is a cultured cell. In some embodiments, the tumor cell is in vivo. In some embodiments, the tumor cell is ex vivo. In some embodiments, the tumor cell is from a mammal. In some embodiments, the mammal is a human. In some embodiments, the mammal is a rodent. In some embodiments, the rodent is a mouse. In some embodiments, the rodent is a rat.

(98) In some embodiments, the tumor-associated antigen encoding polynucleotide may be a DNA molecule comprising a coding strand and a complementary strand, e.g., the tumor-associated antigen gene locus in the genome of a tumor cell. In some embodiments, the tumor-associated antigen encoding polynucleotide may also include coding regions (e.g., exons) and non-coding regions (e.g., introns or splicing sites). In some embodiments, the target base (e.g., a C base) is located in the coding region (e.g., an exon) of the tumor-associated antigen encoding polynucleotide. As such, the conversion of a base in the coding region may result in an amino acid change in the tumor-associated antigen protein sequence, i.e., a mutation. Tumor associated antigens comprising the desired mutation(s), once degraded (e.g., via any of the protein degradation pathways, such as degradation by the proteasome) results in immunogenic heteroclitic epitopes.

(99) In some embodiments, the target base is located in a non-coding region of the tumor-associated antigen gene, e.g., in an intron or a splice site. In some embodiments, a target base is located in a splice site, and the editing of such target base causes alternative splicing of the tumor-associated antigen mRNA. In some embodiments, the alternative splicing leads to translation of a non-coding region of the tumor-associated antigen gene, generating cryptic epitopes. The immunogenic epitopes (e.g., heteroclitic epitopes or cryptic epitopes) may be presented by the tumor cell, or a professional antigen presenting cell, and be recognized by the immune system, thus eliciting a tumor-specific immune response (e.g., T-cell response or B-cell response).

(100) To edit a tumor-associated antigen gene, the tumor-associated antigen gene (a polynucleotide molecule) may be contacted with the nucleobase editor, wherein the nucleobase editor binds to its target sequence and edits the desired base. For example, the nucleobase editor may be expressed in a cell where editing is desired (e.g., a tumor), allowing editing of the tumor-associated antigen gene by the nucleobase editor. In some embodiments, the binding of the nucleobase editor to its target sequence in the tumor-associated antigen gene is mediated by a guide nucleotide sequence, e.g., a nucleotide molecule comprising a nucleotide sequence that is complementary to one of the strands of the target sequence in the tumor-associated antigen gene. Thus, by designing the guide nucleotide sequence, the nucleobase editor may be programmed to edit any target base in any tumor associated antigen gene. In some embodiments, the guide nucleotide sequence is co-expressed with the nucleobase editor in a tumor cell where editing is desired. In some embodiments, a nucleobase editor/gRNA complex is delivered to the cell where editing is desired (e.g., a tumor cell).

(101) Provided herein are non-limiting, exemplary heteroclitic epitopes and cryptic epitopes that may be produced via base editing and strategies for making them.

(102) Codon Change

(103) Using the nucleobase editors described herein, several amino acid codons may be converted to a different codon via deamination of a target base within the codon. For example, in some embodiments, a cytosine (C) base is converted to a thymine (T) base via deamination by a nucleobase editor comprising a cytosine deaminase domain (e.g., APOBEC1 or AID). It is worth noting that during a C to T change via deamination (e.g., by a cytosine deaminase such as APOBEC1 or AID), the cytosine is first converted to a uridine (U), leading to a G:U mismatch. The G:U mismatch is then converted by DNA repair and replication pathways to T:A pair, thus introducing the thymine at the position of the original cytosine. As such, deamination of a C base results in a C-G base pair being replaced by a T-A base pair.

(104) As is familiar to one skilled in the art, conversion of a base in an amino acid codon may lead to a change of the amino acid the codon encodes. Cytosine deaminases are capable of converting a cytosine (C) base to a thymine (T) base via deamination. Thus, it is envisioned that, for amino acid codons containing a C base, the C base may be directly converted to T. For example, codon for leucine (CTC) may be changed to a TTC (phenylalanine) codon via the deamination of the first C on the coding strand. For amino acid codons that contain a guanine (G) base, a C base is present on the complementary strand; and the G base may be converted to an adenosine (A) via the deamination of the C on the complementary strand. For example, an ATG (Met/M) codon may be converted to a ATA (Ile/I) codon via the deamination of the third C on the complementary strand. In some embodiments, two C to T changes are required to convert a codon to a different codon. Non-limiting examples of possible mutations that may be made in any tumor associated antigens by the nucleobase editors of the present disclosure are summarized in Table 4.

(105) TABLE-US-00008 TABLE 4 Exemplary Codon Changes via Base Editing Target codon Base-editing reaction (s)  
Edited codon CTT (Leu/L) 1st base C to T on coding strand TTT (Phe/F) CTC (Leu/L) 1st base C to T on coding strand  
TTC (Phe/F) ATG (Met/M) 3rd base C to T on complementary strand ATA (Ile/I) GTT (Val/V) 1st base C to T on  
complementary strand ATT (Ile/I) GTA (Val/V) 1st base C to T on complementary strand ATA (Ile/I) GTC (Val/V) 1st base  
C to T on complementary strand ATC (Ile/I) GTG (Val/V) 1st base C to T on complementary strand ATG (Met/M) TCT  
(Ser/S) 2nd base C to T on coding strand TTT (Phe/F) TCC (Ser/S) 2nd base C to T on coding strand TTC (Phe/F) TCA  
(Ser/S) 2nd base C to T on coding strand TTA (Leu/L) TCG (Ser/S) 2nd base C to T on coding strand TTG (Leu/L) AGT  
(Ser/S) 2nd base C to T on complementary strand AAT (Asp/N) AGC (Ser/S) 2nd base C to T on complementary strand

AAC (Aps/N) CCT (Pro/P) 1st base C to T on coding strand TCT (Ser/S) CCC (Pro/P) 1st base C to T on coding strand TCC (Ser/S) CCA (Pro/P) 1st base C to T on coding strand TCA (Ser/S) CCG (Pro/P) 1st base C to T on coding strand TCG (Ser/S) CCT (Pro/P) 2nd base C to T on coding strand CTT (Leu/L) CCC (Pro/P) 2nd base C to T on coding strand CTC (Leu/L) CCA (Pro/P) 2nd base C to T on coding strand CTA (Leu/L) CCG (Pro/P) 2nd base C to T on coding strand CTG (Leu/L) ACT (Thr/T) 2nd base C to T on coding strand ATT (Leu/L) ACC (Thr/T) 2nd base C to T on coding strand ATC (Leu/L) ACA (Thr/T) 2nd base C to T on coding strand ATA (Leu/L) ACG (Thr/T) 2nd base C to T on coding strand ATG (Met/M) GCT (Ala/A) 2nd base C to T on coding strand GTT (Val/V) GCC (Ala/A) 2nd base C to T on coding strand GTC (Val/V) GCA (Ala/A) 2nd base C to T on coding strand GTA (Val/V) GCG (Ala/A) 2nd base C to T on coding strand GTG (Val/V) GCT (Ala/A) 1st base C to T on complementary stand ACT (Thr/T) GCC (Ala/A) 1st base C to T on complementary stand ACC (Thr/T) GCA (Ala/A) 1st base C to T on complementary stand ACA (Thr/T) GCG (Ala/A) 1st base C to T on complementary stand ACG (Thr/T) CAT (His/H) 1st base C to T on complementary stand TAT (Tyr/Y) CAC (His/H) 1st base C to T on complementary stand TAC (Tyr/Y) GAT (Asp/D) 1st base C to T on complementary stand AAT (Asp/N) GAC (Asp/D) 1st base C to T on complementary stand AAC (Asp/N) GAA (Glu/E) 1st base C to T on complementary stand AAA (Lys/K) GAG (Glu/E) 1st base C to T on complementary stand AAG (Lys/K) TGT (Cys/C) 2nd base C to T on complementary stand TAT (Tyr/Y) TGC (Cys/C) 2nd base C to T on complementary stand TAC (Tyr/Y) CGT (Arg/R) 1st base C to T on coding strand TGT (Cys/C) CGC (Arg/R) 1st base C to T on coding strand TGC (Cys/C) AGA (Arg/R) 2nd base C to T on complementary stand AAA (Lys/K) AGG (Arg/R) 2nd base C to T on complementary stand AAG (Lys/K) CGG (Arg/R) 2nd base C to T on complementary stand CAG (Gln/Q) CGG (Arg/R) 1st base C to T on coding strand TGG (Trp/W) GGT (Gly/G) 2nd base C to T on complementary stand GAT (Asp/D) GGC (Gly/G) 2nd base C to T on complementary stand GAC (Asp/D) GGA (Gly/G) 2nd base C to T on complementary stand GAA (Glu/E) GGG (Gly/G) 2nd base C to T on complementary stand GAG (Glu/E) GGT (Gly/G) 1st base C to T on complementary stand AGT (Ser/S) GGC (Gly/G) 1st base C to T on complementary stand AGC (Ser/S) GGA (Gly/G) 1st base C to T on complementary stand AGA (Arg/R) GGG (Gly/G) 1st base C to T on complementary stand AGG (Arg/R)

(106) Such amino acid substitutions introduced via base editing generate heteroclitic epitopes. Non-limiting examples of heteroclitic epitopes that may be generated from tumor associated antigens by nucleobase editors are summarized in Table 5.

(107) In some embodiments, to bind to its target sequence and edit the desired base, the nucleobase editor depends on its guide nucleotide sequence (e.g., a guide RNA). In some embodiments, the guide nucleotide sequence is a gRNA sequence. A gRNA typically comprises a tracrRNA framework allowing for Cas9 binding, and a guide sequence, which confers sequence specificity to fusion proteins disclosed herein. In some embodiments, the guide RNA comprises the structure 5'-[guide sequence]-

guuuuagagcuagaaauagcaaguuaaaauaaaggcuaguccguuaucacuugaaaaaguggcaccgagucggugcuu uuu-3' (SEQ ID NO: 336), wherein the guide sequence comprises a sequence that is complementary to the target sequence. Other suitable tracrRNA framework sequences are provided in Table 11. The guide sequence is typically about 20 nucleotides long. In certain embodiments, the guide sequence may be 15-25 nucleotides long. In some embodiments, the guide sequence is 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotides long. Such suitable guide RNA sequences typically comprise guide sequences that are complementary to a nucleic sequence within 50 (e.g., within 50, 45, 40, 35, 30, 35, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, or 10) nucleotides upstream or downstream of the target nucleotide to be edited.

(108) Guide sequences that may be used to target the nucleobase editor to its target sequence to induce specific mutations in tumor associated antigen genes are provided in Table 5. It is to be understood that the mutations and guide sequences presented herein are for illustration purpose only and are not meant to be limiting.

(109) TABLE-US-00009 TABLE 5 Heteroclitic epitopes Antigen Name/Epitope Heteroclitic amino acid epitope position (mutation), Programmable SEQ Genome (Exemplary Genomic guide-RNA ID gRNA size Editor condition) target region sequence NO (PAM) (C-edited) type.sup.a gp100/209-217 IIDQVPFSV

GCCUUCACCAUACUGGUAA 724 (GGG) 20 (C14) SpBE3 (melanoma) (T2101)  
CACCAUACUGGUAAAGGGUU 870 (TAG) 20 (C9) SpBE3 (SEQ ID ACCAUUACUGGUAAAGGGUUU 871 (AGG) 20 (C8) SpBE3 NO: 786) CCAUACUGGUAAAGGGUUUA 872 (GGAA) 20 (C7) VQR-SpBE3 ttcaccattaCt AUUACUGGUAAAGGGUUUAGG 873 (AAG) 20 (C5) SpBE3 ggtaagggttta  
UUACUGGUAAAGGGUUUAGGA 874 (AGG) 20 (C4) SpBE3 ggaaggggca UACUGGUAAAGGGUUUAGGAA 875 (GGG) 20 (C3) SpBE3 (SEQ ID ACUGGUAAAGGGUUUAGGAAG 876 (GGG) 20 (C2) SpBE3 NO: 831)  
CAGCCUUCACCAUACUGGU 877 (AAGGGT) 20 (C16) SaBE3 UUACUGGUAAAGGGUUUAGGA 888 (AGGGG) 20 (C5) St3BE3 gp100/280-288 YLEPGPVT UCACUGCCCAGGUGGUCCUG 725 (CAG) 20 (C7) SpBE3 (melanoma) V (A288V) CACUGCCCAGGUGGUCCUGC 889 (AGG) 20 (C6) SpBE3 (SEQ ID NO: 818) agtcactgCcc aggtgtgtctg caggctgccatt cc (SEQ ID NO: 832) gp100/154-162 KIWGQYW  
AUGUCUGGAAGACCUGGGGU 726 (GAG) 20 (C13) SpBE3 (melanoma) QV (T1551)  
UGUCUGGAAGACCUGGGGUG 890 (AGG) 20 (C12) SpBE3 (SEQ ID GUCUGGAAGACCUGGGGUGA 891 (GGG) 20 (C11) SpBE3 NO: 787) UCUGGAAGACCUGGGGUGAG 892 (GGAC) 20 (C10) VQR-SpBE3 tatgtctggaag aCctggggtg agggactccctt ct (SEQ ID NO: 833) MART-1/26-35 EVAGIGILT  
GGCCGCAGGGAUCGGCAUCC 727 (TGAT) 20 (C6) VQR-SpBE3 (melanoma) V (A27V)  
GCAGGGAUCGGCAUCCUGAU 893 (CGTG) 20 (C2) VQR-SpBE3 (SEQ ID UGUCCAGGGCCGCAGGGAU



894 (CGG) 20 (C14) SpBE3 (NO: 819) AGGCGGCGGCGGAU 895 (CCTGAT) 20 (C8) KKH-SaBE3  
 tgggtccagggc GCAGGGAUCGGCAUCCUGAU 896 (CGTGGT) 20 (C2) KKH-SaBE3 cgCagggatc ggcacatcctgatc  
 gtggctcctggg ga (SEQ ID NO: 834) NY-ESO-1/157- SLLMWITQ ACUGCGUGAUCCACAUAAC 728 (AGG)  
 20 (C-1) SpBE3 165 Y (C165Y) CACUGCGUGAUCCACAUAAC 897 (CAG) 20 (C1) SpBE3 (multiple (SEQ  
 ID tumors, e.g., NO: 788) melanoma and ctttcctgtgtat breast cancer) gtggatcacgc agtGctttctgc cc (SEQ ID  
 NO: 835) TYR/369-377 YMNGIMSQ AUGGAACAAUGUCCCAGGUA 729 (CAG) 20 (C7) SpBE3 (melanoma)  
 V (T3731) UGGAACAAUGUCCCAGGUAC 898 (AGG) 20 (C6) SpBE3 (SEQ ID  
 GGAACAAUGUCCCAGGUACA 899 (GGG) 20 (C5) SpBE3 NO: 789) GAACAAUGUCCCAGGUACAG 900  
 (GGAT) 20 (C4) VQR-SpBE3 tatgaatggaa UGGAACAAUGUCCCAGGUAC 901 (AGGGAT) 20 (C6) SaBE3  
Caatgtcccag gtacagggatc tgcc (SEQ ID NO: 836) TyRP-1/240- DAEKYDID CACACUUUUCUGCAUCCCGC 730  
 (CAG) 20 (C3) SpBE3 251 TDEY GUCACACUUUUCUGCAUCCC 902 (GCCAGT) 20 (C5) KKH-SaBE3  
 (melanoma) (C244Y) (SEQ ID NO: 790) tgggactggcg ggtatgcagaa aagtGtgacat ttgcacagatg agtacatggga (SEQ ID  
 NO: 837) Survivin/95-102 ELILGEFLK AUUAACCCUUGGUGAAUUUU 731 (TGAA) 20 (C6) VQR-SpBE3  
 (multiple L (T971) CCUUGGUGAAUUUUUGAAAC 903 (TGG) 20 (C-1) SpBE3 tumors, e.g., (SEQ ID  
 melanoma, NO: 791) breast cancer, ttgaagaatta and leukemia) aCcttggtga attttgaaactg gac (SEQ ID NO:  
 838) HER2/657-666 AMVIGILLV CGCAGAGAUGAUGGACGUA 732 (GAG) 20 (C-1) VQR-SpBE3 (breast  
 cancer) V (V658M) ACCGCAGAGAUGAUGGACGU 904 (CAG) 20 (C2) SpBE3 (SEQ ID  
 CCAACCACCGCAGAGAUGAU 905 (GGAC) 20 (C8) VQR-SpBE3 NO: 793) GCCAACCACCGCAGAGAUGA  
 906 (TGG) 20 (C9) SpBE3 gcttcgcggccc AAUGCCAACCACCGCAGAGA 907 (TGAT) 20 (C12) VQR-SpBE3  
 agccctctgac AGAAUGCCAACCACCGCAGA 908 (GATGAT) 20 (C14) KKH-SaBE3 gtccatcatctct  
 CCGCAGAGAUGAUGGACGUC 909 (AGAG) 20 (C1) EQR-SpBE3 gcgGtggttg cattctgctggc (SEQ ID NO:  
 839) HER2/911-920 IIWELMTFG AGGUGUGACUGUGUGGGAGC 733 (TGAT) 20 (C9) VQR-SpBE3 (breast  
 cancer) A (T9121) UGUGACUGUGUGGGAGCUGA 910 (TGAC) 20 (C6) VQR-SpBE3 (SEQ ID  
 UUAGGUGUGACUGUGUGGGA 911 (GCTGAT) 20 (C11) KKH-SaBE3 NO: 794)  
 GUGUGGGAGCUGAUGACUUU 912 (TGGGG) 20 (C-2) St3BE3 aggtgtgaCtg tgtgggagctg atgacttttggg gCcaaactta  
 cgtgggatcc cagccccgga gatccct (SEQ ID NO: 840) HER2/911-920 ITWELMTF ACUUUUGGGGCCAAACCUUA  
 734 (CGAT) 20 (C11) VQR-SpBE3 (breast cancer) GV (A920V) UUUGGGGCCAAACCUUACGA 913 (TGG)  
 20 (C8) SpBE3 (SEQ ID UUGGGGCCAAACCUUACGAU 914 (GGG) 20 (C7) SpBE3 NO: 795)  
 UUUGGGGCCAAACCUUACGA 915 (TGGGAT) 20 (C8) SaBE3 gacttttgggg Ccaaacttac gatgggatccc agccccggg  
 (SEQ ID NO: 841) hTERT/540- ILAKFLHWL CAUCAGCCAGUGCAGGAACU 735 (TGG) 20 (C1) SpBE3 549  
 I (M5491) CACACUCAUCAGCCAGUGCA 916 (GGAA) 20 (C7) VQR-SpBE3 (breast cancer) (SEQ ID  
 ACACACUCAUCAGCCAGUGC 917 (AGG) 20 (C8) SpBE3 NO: 792) UACACACUCAUCAGCCAGUG 918  
 (CAG) 20 (C9) SpBE3 cgtgaggagat GACGUACACACUCAUCAGCC 919 (AGTG) 20 (C12) VQR-SpBE3  
 cctggccaagtt GACGACGUACACACUCAUCA 920 (GCCAGT) 20 (C11) KKH-SaBE3 cctgcactggct gatGagtgtgt  
 acgtcgtcgag ctg (SEQ ID NO: 842) hTERT/572- WLFFYRKS UUCUGAUGCUCGGCUCUUCU 736 (TGG) 20  
 (C11) SpBE3 580 V (R572W) (breast cancer) (SEQ ID NO: 716) cccaagcctat cttttctgatct Cggctcttcttg  
 gtcacctctcg ttcca (SEQ ID NO: 843) SSX2/41-49 KVSEKIFYV GAAAAGAUGAAAGCCUCGGA 737  
 (GAAAAT) 20 (C14) KKH-SaBE3 (multiple (A42V) GCCUCGGAGAAAAUCUUCUA 921 (TGTG) 20 (C2) VQR-  
 SpBE3 tumors) (SEQ ID NO: 797) ggaaaagatg aaagCctcgg agaaaatcttct atgtgtatatga agagaaagat gaggctatgac t  
 (SEQ ID NO: 844) WT1/235-243 YMTWNQM CAGGUCAUGCAUUAAGCUG 738 (GGAT) 20 (C10) VQR-  
 SpBE3 (Leukemia) NL (C235Y) CCAGGUCAUGCAUUAAGCU 922 (GGG) 20 (C11) SpBE3 (SEQ ID  
 UCCAGGUCAUGCAUUAAGC 923 (TGG) 20 (C12) SpBE3 NO: 798) GCAUUAAGCUGGGAUGUCA 924  
 (TTTGGT) 20 (C2) KKH-SaBE3 atttataccaaa UCCAGGUCAUGCAUUAAGC 925 (TGGGAT) 20 (C12) SaBE3  
 tgacatcccag ctggaatGcatg acctgga (SEQ ID NO: 845) WT1/235-243 CITWNQMN UGAUUCAGGUCAUGCAUUC  
 739 (AAG) 20 (C12) SpBE3 (Leukemia) L (M2361) UCCAGGUCAUGCAUUAAGC 926 (TGGGAT) 20 (C8)  
 SaBE3 (SEQ ID CAGGUCAUGCAUUAAGCUG 927 (GGAT) 20 (C6) VQR-SpBE3 NO: 799)  
 CCAGGUCAUGCAUUAAGCU 928 (GGG) 20 (C7) SpBE3 atttataccaaa UCCAGGUCAUGCAUUAAGC 929  
 (TGG) 20 (C8) SpBE3 tgacatcccag ctggaatgcatG acctggaatca gat (SEQ ID NO: 846) CD33/65-73 YIISGDSPV  
 GGAAGGAGCCAUAUAUUAUCC 740 (AGG) 20 (C10) SpBE3 (Leukemia) (A65V)  
 GGAAGGAGCCAUAUAUUAUCCA 930 (GGG) 20 (C9) SpBE3 (SEQ ID GAAGGAGCCAUAUAUUAUCCAG 931  
 (GGAC) 20 (C8) VQR-SpBE3 NO: 796) GCCAUUAUAUCCAGGGACUC 932 (TCCAGT) 20 (C2) KKH-SaBE3  
 ctggttccggg aaggagCcat tatatccaggg actctccagt (SEQ ID NO: 847) EpCAM/184- ILYENNVII  
 UAAUGUUAUCACUAUUGAUC 741 (TGG) 20 (C12) SpBE3 192 (T1921) UUAUCACUAUUGAUCUGGUU 933  
 (CAAAAT) 20 (C7) KKH-SaBE3 (multiple (SEQ ID AAUAAUGUUAUCACUAUUGA 934 (TCTGGT) 20 (C14)  
 KKH-SaBE3 tumors) NO: 800) atgagaataat gttatcaCtattg atctggttcaaa attctctc (SEQ ID NO: 848) CEA-CAM/24-  
 LLTFWNPII ACCACUGCCAAGCUCACUAU 742 (TGA) 20 (C2) SpBE3 31 (T3141)  
 GGAACCCGCCCACCACUGCC 935 (AAG) 20 (C13) SpBE3 (multiple (SEQ ID ACCACUGCCAAGCUCACUAU  
 936 (TGAA) 20 (C2) VQR-SpBE3 tumors, e.g., NO: 801) colorectal taaccttctgga cancer, lung acccgcccaC  
 cancer, breast cactgccaagc cancer) tcactattgaatc cagccgt (SEQ ID NO: 849) CEA-CAM/310- RITVTITV  
 CCUCAUAGGACCACAGUCA 743 (CGAC) 20 (C12) VQR-SpBE3 318 (T3111)  
 CAAUAGGACCACAGUCACGA 937 (CGAT) 20 (C9) VQR-SpBE3 (multiple (SEQ ID



GACCACAGACACGACGAUCA 938 (CAG) 20 (C3) SpBE3 tumors, e.g., NO: 802) CUCAAUAGGACCACAGUCAC 939 (GACGAT) 20 (C11) KKH-SaBE3 colorectal gacactggcct AGGACCACAGUCACGACGAU 940 (CACAGT) 20 (C5) KKH-SaBE3 cancer, lung caataggaCc cancer, breast acagtacacgac cancer) gatcacagtct atggaagtgg atccacgaa (SEQ ID NO: 850) CEA-CAM/687- AVVGIMIGV UGGGGCCACUGUCGGCAUCA 744 (TGAT) 20 (C9) VQR-SpBE3 695 (T688V) GCCACUGUCGGCAUCAUGAU 941 (TGG) 20 (C5) SpBE3 (multiple (SEQ ID CCACUGUCGGCAUCAUGAUU 942 (GGAG) 20 (C4) EQR-SpBE3 tumors, e.g., NO: 803) CACUGUCGGCAUCAUGAUUG 943 (GAG) 20 (C3) SpBE3 colorectal tctctgtgtctt ACUGUCGGCAUCAUGAUUGG 944 (AGTG) 20 (C2) VQR-SpBE3 cancer, lung cagctggggcc GCUGGGGCCACUGUCGGCAU 945 (CATGAT) 20 (C11) KKH-SaBE3 cancer, breast aCgtgtcggcat GCCACUGUCGGCAUCAUGAU 946 (TGGAGT) 20 (C5) SaBE3 cancer) catgattggagt GCCACUGUCGGCAUCAUGAU 947 (TGGAG) 20 (C5) St3BE3 gctggttggggt t (SEQ ID NO: 851) CEA-CAM/691- IMIGMLVGV UCAAUCAUGAUGCCGACAG 745 (TGG) 20 (C-1) SpBE3 699 (V695M) ACUCCAAUCAUGAUGCCGAC 948 (AGTG) 20 (C2) VQR-SpBE3 (multiple (SEQ ID CACUCCAAUCAUGAUGCCGA 949 (CAG) 20 (C3) SpBE3 tumors, e.g., NO: 804) CAGCACUCCAAUCAUGAUGC 950 (CGAC) 20 (C6) VQR-SpBE3 colorectal tctcagctggg CCCAACCAGCACUCCAAUCA 951 (TGAT) 20 (C12) VQR-SpBE3 cancer, lung gccactgtcgg AGCACUCCAAUCAUGAUGCC 952 (GACAGT) 20 (C5) KKH-SaBE3 cancer, breast catcatgattgg ACCCCAACCAGCACUCCAAU 953 (CATGAT) 20 (C14) KKH-SaBE3 cancer) aGtgctggttg ggggtt (SEQ ID NO: 852) MAGEA3/112- KVAELVYF GUUGGUUCAUUUUCUGCUCC 746 (TCAAGT) 20 (C8) KKH-SaBE3 120 L (H118Y) UGGUUCAUUUUCUGCUCCUC 954 (AAG) 20 (C6) SpBE3 (multiple (SEQ ID tumors) NO: 805) aggtggccga gttggttCattttc tgctcctcaagt atcgagccag ggagccggtc ac (SEQ ID NO: 853) MAGEA3/181- YLGLSYDG AGGUGACAAGGACAUAGGAG 747 (TGG) 20 (C-1) SpBE3 190 (also in LL (C181Y) GCAGGUGACAAGGACAUAGG 955 (AGTG) 20 (C2) VQR-SpBE3 MAGE A1 (SEQ ID GGCAGGUGACAAGGACAUAG 956 (GAG) 20 (C3) SpBE3 (multiple NO: 806) AGGCAGGUGACAAGGACAU 957 (GGAG) 20 (C4) EQR-SpBE3 tumors) ctatgtccttgtc UAGGCAGGUGACAAGGACAU 958 (AGG) 20 (C5) SpBE3 acctGcctagg CUAGGCAGGUGACAAGGACA 959 (TAG) 20 (C6) SpBE3 tctctctatgat GAGACCUAGGCAGGUGACAA 960 (GGAC) 20 (C11) VQR-SpBE3 (SEQ ID AGAGACCUAGGCAGGUGACA 961 (AGG) 20 (C13) SpBE3 NO: 854) UAGGCAGGUGACAAGGACAU 962 (AGGAGT) 20 (C5) SaBE3 20 (C-1) SpBE3 MAGEA3/181- YLGLSYDG AGGUGACAAGGAUGUACAAG 748 (TGG) 20 (C-1) SpBE3 190 (also in LL (C181Y) GCAGGUGACAAGGAUGUACA 963 (AGTG) 20 (C2) VQR-SpBE3 MAGE A2/A12 (SEQ ID GGCAGGUGACAAGGAUGUAC 964 (AAG) 20 (C3) SpBE3 (multiple NO: 806) GAGGCCCAGGCAGGUGACAA 965 (GGAT) 20 (C11) VQR-SpBE3 tumors) tacatccttgtca AGAGGCCCAGGCAGGUGACA 966 (AGG) 20 (C13) SpBE3 cctGcctgggc CAGGCAGGUGACAAGGAUGU 967 (ACAAGT) 20 (C5) KKH-SaBE3 ctctcctacgat GAGAGGCCCCAGGCAGGUGAC 968 (AAGGAT) 20 (C14) SaBE3 (SEQ ID NO: 855) MAGEA3/181- YLGLSYDG AGGUGGCAAAGAUGUACAAG 749 (TGG) 20 (C-1) SpBE3 190 (also in LL (C181Y) GCAGGUGGCAAAGAUGUACA 969 (AGTG) 20 (C2) VQR-SpBE3 MAGE A3 (SEQ ID GGCAGGUGGCAAAGAUGUAC 970 (AAG) 20 (C3) SpBE3 (multiple NO: 806) GAGGCCCAGGCAGGUGGCAA 971 (AGAT) 20 (C11) VQR-SpBE3 tumors) gtacatccttggc AGAGGCCCAGGCAGGUGGCA 972 (AAG) 20 (C13) SpBE3 acctGcctggg CAGGCAGGUGGCAAAGAUGU 973 (ACAAGT) 20 (C6) KKH-SaBE3 cctctcctacga GAGAGGCCCCAGGCAGGUGGC 974 (AAAGAT) 20 (C12) KKH-SaBE3 (SEQ ID NO: 856) MAGEA3/181- YLGLSYDG AGGCAGGUGACAAGGGUGUA 750 (GGTG) 20 (C4) VQR-SpBE3 190 (also in LL (C181Y) CAGGCAGGUGACAAGGGUGU 975 (AGG) 20 (C5) SpBE3 MAGE A4 (SEQ ID CCAGGCAGGUGACAAGGGUG 976 (TAG) 20 (C6) SpBE3 (multiple NO: 806) AGGCCCAGGCAGGUGACAAG 977 (GGTG) 20 (C9) VQR-SpBE3 tumors) tacacccttgtc AAGGCCCAGGCAGGUGACAA 978 (GGG) 20 (C10) SpBE3 acctGcctggg AAAGGCCCCAGGCAGGUGACA 979 (AGG) 20 (C11) SpBE3 cctttctatgat GAAAGGCCCCAGGCAGGUGAC 980 (AAG) 20 (C12) SpBE3 (SEQ ID CCCAGGCAGGUGACAAGGGU 981 (GTAGGT) 20 (C7) KKH-SaBE3 NO: 857) CAGGCAGGUGACAAGGGUGU 982 (AGGTG) 20 (C5) St3BE3 AAGGCCCCAGGCAGGUGACAA 983 (GGGTG) 20 (C11) St3BE3 MUC-1/92-101 AIWGQDVT UCAGCUGCCACCUGGGGACA 751 (GGAT) 20 (C11) VQR-SpBE3 (multiple SV (T931) CUGGGGACAGGAUGUCACCU 984 (CGG) 20 (C-1) SpBE3 tumors) (SEQ ID ACCUGGGGACAGGAUGUCAC 985 (CTCGGT) 20 (C2) KKH-SaBE3 NO: 807) tcagctgccac ctggggacag gatgtcacctcg gtccagtcac caggcca (SEQ ID NO: 858).sup.aGenome-editor types abbreviations: SpBE3 = APOBEC1—SpCas9n—UGI; VQR-SpBE3 = APOBEC1—VQR-SpCas9n—UGI; EQR-SpBE3 = APOBEC1—EQR-SpCas9n—UGI; VRER-SpBE3 = APOBEC1—VRER-SpCas9n—UGI; SaBE3 = APOBEC1—SaCas9n—UGI; KKH-SaBE3 = APOBEC1—KKH-SaCas9n—UGI; St3BE3 = APOBEC1—St3Cas9n—UGI; St1BE3 = APOBEC1—St1Cas9n—UGI. Guide sequences (the portion of the guide RNA that targets the nucleobase editor to the target sequence) are provided. The guide sequences may be used with any tracrRNA framework sequences known in the art to generate the full guide RNA sequence.

#### Cryptic Epitopes

(110) Some aspects of the present disclosure provide strategies for generating cryptic epitopes in tumor cells. In some embodiments, such strategies involve alterations of splicing sites in a tumor associated antigen gene. Altered splicing site

may lead to altered splicing of an mRNA that encodes a tumor associated antigen. One outcome of altered splicing is the translation of an otherwise non-coding region of the gene, leading to otherwise “hidden peptides,” i.e., cryptic epitopes. The splicing site typically comprises an intron donor site, a Lariat branch point, and an intron acceptor site. The mechanism of splicing are familiar to those skilled in the art. The intron donor site has a consensus sequence of GGGTRAGT, and the C bases paired with the G bases in the intron donor site consensus sequence may be targeted by a nucleobase editor, thereby altering the intron donor site. The Lariat branch point also has a consensus sequence, e.g., YTRAC, wherein Y is a pyrimidine, and R is a purine. The C base in the Lariat branch point consensus sequence may be targeted by the nucleobase editors described herein, leading to skipping of the following exon. The intron acceptor site has a consensus sequence of YNCAGG, wherein Y is a pyrimidine, and N is any nucleotide. The C base of the consensus sequence of the intron acceptor site, and the C base paired with the G bases in the consensus sequence of the intron acceptor site may be targeted by the nucleobase editors described herein, thereby altering the intron acceptor site, in turn leading to skipping of an exon. General strategies of altering the splicing sites are described in Table 6.

(111) TABLE-US-00010 TABLE 6 Exemplary Alteration of Intron-Exon Junction via Base Editing Target Consensus Base-editing Edited site Sequence reaction (s) sequence Outcome Intron GGGTRAGT 2.sup.nd or 3.sup.rd base GAGTRAGT Intron sequence is donor (example) C to T on (example) translated as exon, in complementary frame premature STOP strand codon Lariat YTRAC 5.sup.th base C to T YTRAT The following exon is branch (example) on coding (example) skipped from the mature point strand mRNA, which may affect the coding frame Intron Y(rich)NCAGG 2.sup.nd to last base Y(rich)NCAAG The exon is skipped from acceptor (example) C to T on (example) the mature mRNA, which complementary may affect the coding strand frame Start ATG (Met/M) 3.sup.rd base C to T ATA (Ile/I) The next ATG is used as codon on start codon, which may complementary affect the coding frame strand (112) Non-limiting, exemplary cryptic epitopes that may be produced using the base editing methods described herein are provided in Table 7.

(113) TABLE-US-00011 TABLE 7 Cryptic Epitopes Antigen Name Cryptic and epitope(s) and Programmable SEQ gRNA Genome Exemplary Genomic target guide-RNA ID size (C- Editor Condition(s) region sequence NO (PAM) edited) type.sup.a gp100 (PMEL VYFFLPDHL UCCCUCACCCCAGGUCUUC 752 (AGAC) 20 (C9) VQR-SpBE3 gene) (SEQ ID NO: GUCCCUCACCCCAGGUCUUC 986 (CAG) 20 (C9/10) SpBE3 (melanoma) 808) AGAAGGGAGUCCCUCACCCC 987 (AGG) 20 (n.a.) WT Cas9 AAGCTTTGTTT UGUCUGGAAGACCUGGGGUG 988 (AGG) 20 (n.a.) WT Cas9 ATGTCTGGAAG GUCUGGAAGACCUGGGGUGA 989 (GGG) 20 (n.a.) WT Cas9 ACCTGGGggtgag AUGUCUGGAAGACCUGGGGU 990 (GAG) 20 (n.a.) WT Cas9 ggactcccttctcagcc GAGAAGGGAGUCCCUCACCCC 991 (CAG) 20 (n.a.) WT Cas9 tatcatccacac (SEQ ID NO: 859) intron 4, intron donor site gp100 (PMEL VYFFLPDHL CCUGAAGUUUUUGGAAUGAA 753 (AAG) 20 (C1/2) SpBE3 gene) (SEQ ID NO: UUGGCCUGAAGUUUUUGGAA 992 (TGAA) 20 (C4/5) VQR-SpBE3 (melanoma) 808) AGUAUUGGCCUGAAGUUUUU 993 (GGAA) 20 (C9/10) VQR-SpBE3 ccaaaaacttcagG CAGUAUUGGCCUGAAGUUUU 994 (TGG) 20 (C10/11) SpBE3 CCAATACTGGC CAGUAUUGGCCUGAAGUUUU 995 (TGGAAT) 20 (C10/11) SaBE3 AAGTTCT (SEQ CUUUUCAUUCCAAAAACUUC 996 (AGG) 20 (n.a.) WT Cas9 ID NO: 860) AACUUGCCAGUAUUGGCCUG 997 (AAG) 20 (n.a.) WT Cas9 intron 4, intron GCUUUUCAUUCCAAAAACUUC 998 (CAG) 20 (n.a.) WT Cas9 acceptor site TYRP1 MSLQRQFLR (S CUCAUUCUGCUUGAAAUAA 999 (GAG) 20 (C4) SpBE3 gcactctatttcaagc GCACUCAUUCUGCUUGAAAU 1000 (AAG) 20 (C6) SpBE3 agaatGagtgtctcta UUAGGAGCACUCAUUCUGCU 1001 (TGAA) 20 (C11) VQR-SpBE3 a (SEQ ID NO: GCACUCAUUCUGCUUGAAAU 1002 (AAGAGT) 20 (C6) SaBE3 861) UUAGGAGCACUCAUUCUGCU 1003 (TGAAAT) 20 (C11) KKH-SaBE3 ORF1, target CACUCAUUCUGCUUGAAAU 1004 (AGAG) 20 (C5) EQR-SpBE3 start codon ACUCUUAUUUCAAGCAGAAU 1005 (GAG) 20 (n.a.) WT Cas9 MGAT5 VLPDVFIRCV UGUAAGACAGAAAACCACAC 755 (AGCG) 20 (C-1) VRER-SpBE3 (melanoma) (SEQ ID NO: CUGUAAGACAGAAAACCACA 1006 (CAG) 20 (C1) SpBE3 810) CAAGUCCAACAAACAACUGU 1007 (AAG) 20 (n.a.) WT Cas9 tcatacgtgtgtgtgt UCUGUCUUACAGUUGUUUGU 1008 (TGG) 20 (n.a.) WT Cas9 tttctgtcttacagTT GTTTGTGGACTT GGGTTC (SEQ ID NO: 862) target intron2- exon3 junction, last base of intron 2 LAGE-1 MLMAQEALAFI CUUCGGCCUGCAUGGCUC 756 (GAG) 20 (C11) SpBE3 (multiple (SEQ ID NO: AUGGCUCGGAGCCUCUGCC 1009 (CGG) 20 (C -1) SpBE3 tumors, e.g., 811) GCCUUCGGCCUGCAUGGCUC 1010 (CGG) 20 (C13) SpBE3 melanoma LAAQERRVPR CCUUCGGCCUGCAUGGCUC 1011 (GGAG) 20 (C12) EQR-SpBE3 and breast (SEQ ID NO: CAGAGGCUCGGAGCCAUGC 1012 (AGG) 20 (n.a.) WT Cas9 cancer) 812) GCAGAGGCUCGGAGCCAUG 1013 (CAG) 20 (n.a.) WT Cas9 APRGVRMAV GCCUUCGGCCUGCAUGGCUC 1014 (CGG) 20 (n.a.) WT Cas9 (SEQ ID NO: 813) QGAMLAAQERR VPRAAEVPR (SEQ ID NO: 814) CLSRPWRKRS WSAGSCPMP HL (SEQ ID NO: 815) tctctgagagccgggc agaggctccggagcc atGcaggccgaagg c (SEQ ID NO: 863) target OFR1 start site TRP-2 EVISCKLIKR CCUAAUAAUGUAUCUCUAA 757 (AGAA) 20 (C1/2) VQR-SpBE3 (melanoma) (SEQ ID NO: ACCUAAUAAUGUAUCUCUAA 1015 (CAG) 20 (C2/3) SpBE3 816) ACCUAAUAAUGUAUCUCUAA 1016 (CAGAAT) 20 (C2/3) SaBE3 TATTCTGTAG

ACCAUAUACAUUUAUUCUA 1017 (ACAGAAT) 20 (C3/4) St1BE3 AGATACATTATT  
 UUAGAGAUACAUAUUAUAGGU 1018 (GGG) 20 (n.a.) WT Cas9 A**G**gtgggtttttcc  
 GUUAGAGAUACAUAUUAUAGG 1019 (TGG) 20 (n.a.) WT Cas9 (SEQ ID NO: 864) target intron 2  
 donor site TRP-2 EVISCKLIK R UCCUGGUCCUGAAACAAUUG 758 (GGAA) 20 (C8/9) VQR-SpBE3  
 (melanoma) (SEQ ID NO: GUCCUGGUCCUGAAACAAU 1020 (GGG) 20 (C9/10) SpBE3 816)  
 CGUCCUGGUCCUGAAACAAU 1021 (TGG) 20 (C10/11) SpBE3 tatgtttccaattgtttc  
 UUUCCCAAUUGUUUCAGGAC 1022 (CAG) 20 (n.a.) WT Cas9 ag**G**ACCAGGAC  
 UUCCCAAUUGUUUCAGGACC 1023 (AGG) 20 (n.a.) WT Cas9 GCCCCT (SEQ ID NO: 865) target  
 intron 2 acceptor site BIRC5/ AYACNTSTL AGCUGAAGGGAUUAAGCGG 759 (CAG) 20 (C3) SpBE3  
 Survivin (SEQ ID NO: GGCAGCUGAAGGGAUUAAG 1024 (CGG) 20 (C6) SpBE3 (multiple 817)  
 AAGGCAGCUGAAGGGAUUA 1025 (AGCG) 20 (C8) VRER-SpBE3 tumors, e.g., agtggactgccgttta  
 AAAGGCAGCUGAAGGGAUUA 1026 (AAG) 20 (C9) SpBE3 melanoma, atccctt**C**auctgcctt  
 GCAGCUGAAGGGAUUAAGC 1027 (GGCAGT) 20 (C5) KKH-SaBE3 breast cancer, tccgtgtt (SEQ ID  
 GACUGCCGCUUUAUCCCUU 1028 (CAG) 20 (n.a.) WT Cas9 and leukemia) NO: 866)  
 CUGCCGCUUUAUCCCUUCA 1029 (GCT) 20 (n.a.) WT Cas9 target intron 2 spliceosome branch site  
 BIRC5/ AYACNTSTL UCUAGAAAAAUCAAAACAAC 760 (AGCG) 20 (C2) VRER-SpBE3 Survivin (SEQ ID  
 NO: CUCUAGAAAAAUCAAAACA 1030 (CAG) 20 (C3) SpBE3 (multiple 817)  
 GAACAUAAAAAGCAUUCGUC 1031 (CGG) 20 (n.a.) WT Cas9 tumors, e.g., tccgtgtgttttgatttt melanoma,  
 ctagAGAGGAAC breast cancer, ATAA (SEQ ID and leukemia) NO: 867) target intron 2 acceptor site Bcr-  
 Abl-OOF SSKALQRPV UGGAAGAGAAAGGGGGGAAC 761 (AGAA) 20 (C-1) VQR-SpBE3 (Leukemia) (SEQ  
 ID CUGGAAGAGAAAGGGGGGA 1032 (CAG) 20 (C1) SpBE3 NO: GCUUCUGGAAGAGAAAGGGG 1033  
 (GGAA) 20 (C5) VQR-SpBE3 603) GGCUCUGGAAGAGAAAGGG 1034 (GGG) 20 (C6) SpBE3 GFKQSSKAL  
 GGGCUUCUGGAAGAGAAAGG 1035 (GGG) 20 (C7) SpBE3 (SEQ ID NO: AGGGCUUCUGGAAGAGAAAG  
 1036 (GGG) 20 (C8) SpBE3 604) AAGGGCUUCUGGAAGAGAAA 1037 (GGG) 20 (C9) SpBE3 ATGFKQSSKAL  
 GAAGGGCUUCUGGAAGAGAA 1038 (AGG) 20 (C10) SpBE3 QRPVAS (SEQ  
 UGAAGGGCUUCUGGAAGAGA 1039 (AAG) 20 (C11) SpBE3 ID NO: 605) UCUGGAAGAGAAAGGGGGGA  
 1040 (ACAGAAA) 20 (C2) St1BE3 ATGFKQSSKAL AGGGCUUCUGGAAGAGAAAG 1041 (GGGGG) 20 (C8)  
 St3BE3 QRPVAS (SEQ AAGGGCUUCUGGAAGAGAAA 1042 (GGGGG) 20 (C9) St3BE3 ID NO: 606)  
 GAAGGGCUUCUGGAAGAGAA 1043 (AGGGG) 20 (C10) St3BE3 tccccctttctctcca  
 UUUCCCCCUUUCUCUCCAG 1044 (AAG) 20 (n.a.) WT Cas9 **g**AAGCCCTTCA  
 CUGUCCCCCUUUCUCUUC 1045 (CAG) 20 (n.a.) WT Cas9 GC (SEQ ID NO: 868) target intron 1  
 acceptor site Bcr-Abl-OOF SSKALQRPV UGAGAAGAAAGGAACCAAU 762 (CAG) 20 (C-1) SpBE3  
 (Leukemia) (SEQ ID NO: GCUUUUCACCUGAGAAGAAA 1046 (GGAA) 20 (C9/10) VQR-SpBE3 603)  
 AGCUUUUCACCUGAGAAGAA 1047 (AGG) 20 (C10/11) SpBE3 GFKQSSKAL GAGCUUUUCACCUGAGAAGA  
 1048 (AAG) 20 (C11/12) SpBE3 (SEQ ID NO: UUCACCUGAGAAGAAAGGAA 1049 (CCAAAT) 20 (C5/6)  
 KKH-SaBE3 604) CCGGAGCUUUUCACCUGAGA 1050 (AGA) 20 (C7/8) SpBE3 ATGFKQSSKAL  
 CCCGGAGCUUUUCACCUGAG 1051 (AAG) 20 (C6/7) SpBE3 QRPVAS (SEQ ACCCGGAGCUUUUCACCUGA  
 1052 (GAAGAA) 20 (C8/9) KKH-SaBE3 ID NO: 605) CACCUGAGAAGAAAGGAACC 1053 (AAATC) 20  
 (C6/7) St3BE3 ATGFKQSSKAL GAUUUGGUUCCUUUCUUCUC 1054 (AGG) 20 (n.a.) WT Cas9 QRPVAS  
 (SEQ UUCUUUCUUCUUCAGGUGAA 1055 (AAG) 20 (n.a.) WT Cas9 ID NO: 606)  
 UGAUUUGGUUCCUUUCUUCU 1056 (CAG) 20 (n.a.) WT Cas9 tcctttctctcag**G**TG AAAAGCTCCGG GTCT  
 (SEQ ID NO: 869) target intron 2 acceptor site .sup.aGenome-editor types abbreviations: SpBE3 = APOBEC1  
 —SpCas9n—UGI; VQR-SpBE3 = APOBEC1—VQR-SpCas9n—UGI; EQR-SpBE3 = APOBEC1—EQR-SpCas9n—  
 UGI; VRER-SpBE3 = APOBEC1—VRER-SpCas9n—UGI; SaBE3 = APOBEC1—SaCas9n—UGI; KKH-SaBE3  
 = APOBEC1—KKH-SaCas9n—UGI; St3BE3 = APOBEC1—St3Cas9n—UGI; St1BE3 = APOBEC1—St1Cas9n—  
 UGI. Guide sequences (the portion of the guide RNA that targets the nucleobase editor to the target sequence) are  
 provided. The guide sequences may be used with any tracrRNA framework sequences known in the art to generate the full  
 guide RNA sequence.

(114) In some embodiments, the nucleobase editor may be used to introduce a premature stop codon (a stop codon that occurs upstream of the normal stop codon) into a tumor specific antigen gene (e.g., TAA, TAG, and TGA). In some embodiments, introduction of a premature stop codon destabilizes the tumor specific antigen. In some embodiments, destabilization of the tumor specific antigen leads to enhanced presentation of immunogenic epitopes (e.g., heteroclitic epitopes or cryptic epitopes).

(115) Premature stop codons are introduced by changing one or more bases in a target codon that encodes a target residue. For example, nucleobase editors including a cytosine deaminase domain are capable of converting a cytosine (C) base to a thymine (T) base via deamination. Thus, it is envisioned that, for amino acid codons containing a C base, the C base may be converted to T. For example, a CAG (Gln/Q) codon may be changed to a TAG (amber) codon via the deamination of the first C on the coding strand. For sense codons that contain a guanine (G) base, a C base is present on the complementary strand; and the G base may be converted to an adenosine (A) via the deamination of the C on the complementary strand. For example, a TGG (Trp/W) codon may be converted to a TAG (amber) codon via the deamination of the second C on the complementary strand. In some embodiments, two C to T changes are required to

convert a codon to a nonsense codon. For example, a CCG (R) codon is converted to a TAG (amber) codon via the deamination of the first C on the coding strand and the deamination of the second C on the complementary strand.

(116) In some embodiments, the target residue is located in a flexible loop region of the tumor specific antigen. In some embodiments, tandem premature stop codons are introduced. Non-limiting examples of codons that may be changed to stop codons via base editing are provided in Table 8.

(117) TABLE-US-00012 TABLE 8 Conversion to Stop Codon Target codon Base-editing process Edited codon CAG (Gln/Q) 1.sup.st base C to T on coding strand TAG (amber) TGG (Trp/W) 2.sup.nd base C to T on complementary strand TAG (amber) CGA (Arg/R) 1.sup.st base C to T on coding strand TGA (opal) CAA (Gln/Q) 1.sup.st base C to T on coding strand TAA (ochre) TGG (Trp/W) 3.sup.rd base C to T on complementary strand TGA (opal) CGG (Arg/R) 1.sup.st base C to T on coding strand and 2.sup.nd TAG (amber) base C to T on complementary strand CGA (Arg/R) 1.sup.st base C to T on coding strand and 2.sup.nd TAA (ochre) base C to T on complementary strand \*single underline: changes on the coding strand double underline: changes on the complementary strand

(118) In some embodiments, cryptic epitopes are generated by shifting the coding frame of a tumor specific antigen gene. In some embodiments, the coding frame is shifted by changing a start codon (ATG) to a sense codon that cannot be used as a start codon. As such, translation will start at the next start codon in the coding region, and the coding frame may be shifted. In some embodiments, a normal sense codon may be edited to generate a start codon to allow translation to start at the newly generated start codon, which may also lead to shifting of the coding frame. Alterations of start codons and the resulting shift in the coding frame generate peptides that would not otherwise be generated from the tumor specific antigen gene (i.e., cryptic epitopes). Non-limiting, exemplary start codon alterations that may be achieved by the nucleobase editors described herein are provided in Table 9.

(119) TABLE-US-00013 TABLE 9 Alteration of Start Codons via Base Editing Target codon Base-editing process Edited codon Cognate Start 3.sup.rd base C to T on complementary strand ATA (Ile/I, next ATG is codon ATG used as start codon) (Met/M) ACG (Thr/T) 2.sup.nd base C to T on coding strand ATG (Met/M) GTG (Val/V) 1.sup.st base C to T on complementary strand ATG (Met/M) GCG (Ala/A) 1.sup.st base C to T on complimentary strand and 2.sup.nd ATG (Met/M) base C to T on coding strand

(120) The tumor associated antigens listed in Table 5 and Table 6 and their respective gene sequences and protein sequences are known in the art. The amino acid sequence of the listed tumor specific antigens are listed in Table 10.

(121) TABLE-US-00014 TABLE 10 Amino acid sequences of human tumor specific antigens Name of tumor SEQ associated ID antigen Amino acid sequence NO

Melanocyte  
MDLVLRCLLHLAVIGALLAVGATKVPNRQDWLGVSRLRTKAWNRQLYPEW 763 protein PMEL  
TEAQRLLDCWRGGQVSLKVSNDGPTLIGANASFSIALNFPGSQKVLDPDGQVIW (gp100)  
VNNTIINGSQVWGGQPVYPQETDDACIFPDGGPCPSGSWSQKRSFVYVWKTW  
GQYWQVLGGPVSGLSIGTGRAMLGTHTMEVTYVYHRRGSRSYVPLAHSSSAFT  
ITDQVPFSVSVSQLRALDGGNKHFLRNQPLTFALQLHDPSTGYLAEADLSYTW  
DFGDSSGTLISRALVVTHTYLEPGPVTAQVVLQAAIPLTSCGSSPVPGTTDG  
HRPTAEAPNTTAGQVPTTEVVGTTPGQAPTAEPSTTSVQVPTTEVISTAPV  
QMPTAESTGMTPEKVPVSEVMGTTLAEMSTPEATGMTPAEVSIVVLSGTTAA  
QVTTTEWVETTARELPIPEPEGPDASSIMSTESITGSLGPLLDGTATLRLVK  
RQVPLDCVLYRYGSFSVTLDIVQGIESAEILQAVPSGEGDAFELTVSCQGGL  
PKEACMEISSPGCQPPAQRLCQPVLPSPACQLVLHQLKGGSGTYCLNVSLA  
DTNSLAVVSTQLIMPGQEAGLGQVPLIVGILLVLMMAVVLASLIYRRRLMKQD  
FSPVQLPHSSSHWLRLPRIFCSCPIGENSPLLSGQQV Melanoma  
MPREDAHFYGYPKKGHGHSYTTAEAAAGIGILTIVLGVLLIGCWYCRRRN 764 antigen  
GYRALMDKSLHVGTTQCALTRRCPPQEGFDHRDSKVSLEKNCPEVVPNAPPAY recognized by  
EKLAEQSPPPYSP T-cells 1 (MLANA/M ART-1) 5,6-  
MSAPKLLSLGCIFFPLLLFQQAQFPRQCATVEALRSGMCCPDLSPVSGPG 765 dihydroxyindole-  
TDRCGSSSGRGRCEAVTADSRPHSPQYPHDGRDDREVWPLRFFNRTCHCNGN 2-  
FSGHNCGTCTCRPGWRGAACDQVRLIVRRNLLDLSKEEKNHFVRALDMAKRTTH carboxylic  
PLFVIATRRSEEILGPDGNTQFENISIYNYFVWTHYYSVKKTFLGVGQESF acid oxidase  
GEVDFSHEGPAFLTWHRYHLLRLEKDMQEMLQEPSFSLPYWNFATGKNVCDI (TYRP1)  
CTDDLMSGSRNFDSTLISPNSVFSQWRVVCDSLEDYDTLGTLCNSTEDGPIR  
RNPAGNVARPMVQRLPEPQDVAQCLEVGFLDTPPFYSNSTNSFRNTVEGYSD  
PTGKYDPAVRSLHNLHLFLNGTGGQTHLSPNDPIFVLLHTFTDAVEDEWLR  
RYNADISTFPLENAPIGHNRQYNMVPFWPPVTNTEMFVTAPDNLGYTYEIQW  
PSREFSVPEIIAIAVVGALLLVALIFGTASYLIRARRSMDEANQPLLDQYQ CYAEEYEKLQNPNQSVV Alpha-1,6-  
MALFTPWKLSSQKLGFLLVTFGFIWGMMLLHFTIQQRTQPESSSMLREQILD 766 mannosyl-  
LSKRYIKALAEENRNVVDGPYAGVMTAYDLKKTAVLLDNILQRIGKLESKV glycoprotein 6-  
DNLVVNGTGTNSTNSTTAVPSLVALEKINVADIINGAQEKCVLPMDGYPHC beta-N-  
EGKIKWMKDMWRSDPCYADYGVDGSTCSFFIYLSEVENWCPLPWRANKNPYE acetylglucos-  
EADHNSLAEIRTDNFILYSMMKKHEEFRWMRLRIRRMADAWIQAIKSLAEKQ aminyltransferase  
NLEKRKRKKVLVHLGLLTKESGFKIAETAFFSGGPLGELVQWSDLITSLYLLG A

HDIRISALAEIIVGVGNRSGCTVGDRIYDIVGLAQFKKTLG (MGAT5)  
PSWVHYQCMLRVLDSFGTEPEFNHANYAQSKGHKTPWGKWNLNPPQQFYTMFP  
HTPDNSFLGFVVEQHLNSSDIHHINEIKRQNSLSVYGKVDSFWKNKKIYLDI  
IHTYMEVHATVYGSSTKNIPSYVKNHGLSGRDLQFLLRETKLFFVGLGFPYE  
GPAPLEAIANGCAFLNPKFNPPKSSKNTDFFIGKPTLRELTSQHPYAEVFIG  
RPHVWTVDLNNEEVEDAVKAILNQKIEPYMPYEFTCEGMLQRINAFIEKQD  
FCHGQVMWPPLSALQVKLAEPGQSCQVCQESQLICEPSFFQHLNKDKDMLK  
YKVTCQSSELAKDILVPSFDPKNKHCVFQGDLLLFSACAGHPRHQVPCPRD FIKGQVALCKDCL Cancer/testis  
MQAEGRTGGSTGDADGPGGPGIPDGPGGNAGGPGGEAGATGGRGPRGAGAAR 767 antigen 1  
ASGPGGGAPRGPHGGAASGLNGCCRCGARGPESRLLEFYLAMPFATPMEAE (CTAG1;  
ARRSLAQDAPPLPVPGVLLKEFTVSGNILTIRLTAADHRQLQLSISSCLQQL LAGE2/NY-  
SLLMWITQCFLPVFLAQPPSGQRR ESO-1) CTL-  
MLMAQEALAFMAQGAAMLAQERRVPRAAEVPGAQGGQQGPRGREEAPRGVRM 768 recognized  
AVPLLRMEGAPAGPGGRTAACFSCTSRCLSRRPWKRSWSAGSCPGMPHLSP antigen on DQGRF melanoma  
(CAMEL) L- MSPLWWGFLLSCLGCKILPGAQQGFPRVCMTVDSLVNKECCPRLGAESANVC 769 dopachrome  
GSQQGRGQCTEVRADTRPWSGPYILRNQDDRELWPRKFFHRTCKCTGNFAGY tautomerase  
NCGDCKFGWTGPNCERKKPPVIRQNIHSLSPQEREQFLGALDLAKKRVHPDY (DCT)  
VITTQHWLGLLGPNGTQPQFANCSVYDFFVWLHYYSVRDTLGPGRPYRAID  
FSHQGPAFVTWHRYHLLCLERDLQRLIGNESFALPYWNFATGRNECDVCTDQ  
LFGAARPDPTLISRNSRFSSWETVCDSLDDYNHLVTLNCGTYEGLLRNQM  
GRNSMKLPTLKDIRDCLSLQKFDNPPFFQNSTFSFRNALEGEDKADGTLDSQ  
VMSLHNLVHSFLNGTNALPHSAANDPIFVVLHSFTDAIFDEWMKRFNPPADA  
WPQELAPIGHNRMYNMVPFFPPVTNEELFLTSDQLGYSYIDLPSVEETPG  
WPTTLLVVMGTLVALVGLFVLLAFLQYRRLRKGYTPLMETHLSSKRYTEEA Tyrosinase  
MLLAVLYCLLWSFQTSAGHFPRACVSSKNLMEKECCPPWSGDRSPCGQLSGR 770 (TYR)  
GSCQNILLSNAPLGPQFPFTGVDDRESWPSVFYNRTCQCSGNFMGFNCGNCK  
FGFWGPNCTERRLLVRRNIFDLSAFEKDKFFAYLTLAKHTISSDYVIPICTY  
GQMKNGSTPMENDINIYDLFVWMHYVVSMDALLGGSEIWRDIDFAHEAPAF  
PWHRLFLLRWEQEIQLTGDENFTIPYWDWRDAEKCDICTDEYMGQHPNP  
NLLSPASFFSSWQIVCSRLEEYNHQSLCNGTPEGPLRRNPGNHDKSRTPL  
PSSADVEFCLSLTQYESGMDKAANFSFRNTLEGFASPLTGIADASQSSMHN  
ALHIYMNGTMSQVQGSANDPIFLLHHAFFVDSIFEQWLRRHRPLQEVYPEANA  
PIGHNRESYMPFIPLYRNGDFFISSKDLGYDYSYLQSDPDSFQDIKSYL  
EQASRIWSWLLGAAMVGAVLTALLAGLVSLLCRHKRKQLPEEKQPLMEKED YHSLYQSHL Baculoviral  
MGAPTLPPAWQPFLKDHRISTFKNWPFLGCACTPERMAEAGFIHCPTENEP 771 IAP repeat-  
DLAQCFFCFKELEGWEPDDPIEEHKKHSSGCAFLSVKKQFEELTLGEFLKL containing  
DRERAKNKIAKETNNKKKEFEETAKKVRRRAIEQLAAMD protein 5 (BIRC5) Receptor  
MELAAALCRWGLLLALLPPGAASTQVCTGTDMLRLPASPETHLDMRLHLYQG 772 tyrosine-  
CQVVQGNLELTYPNASLSFLQDIQEVQGYVLIHNVVRQVPLQRLRIVRG protein kinase  
TQLFEDNYALAVLDNGDPLNNTTPVTGASPGGLRELQLRSLTEILKGGVLIQ erbB-2  
RNPQLCYQDTILWKDIFHKNNQLALTIDTNRSRACHPCSPMCKGSRCWGES (ERBB2/HER  
SEDCQSLTRTVCAAGGCARCKGPLPTDCHEQCAAGCTGPKHSDCLACLHENH 2)  
SGICELHCPALVTYNTDTFESMPNPEGRYTFGASCVTACPYNYLSTDVGST  
LVCPLHNQEVTAEDGTQRCEKCSKPCARVCYGLGMEHLREVRAVTSANIQEF  
AGCKKIFGSLAFLPESEDGDPASNTAPLQPEQLQVFETLEEITGYLYISAWP  
DSLPLDSVFQNLQVIRGRILHNGAYSLTLQGLGISWLGLRSLRELGSGLALI  
HHNTHLCFVHTVPWDQLFRNPHQALLHTANRPEDECVGEGLACHQLCARGHC  
WGPAPTQCVNCSQFLRGQECVEECRVLQGLPREYVNARHCLPCHPECQPQNG  
SVTCFGPEADQCVACAHYKDPPFCVARCPSGVKPDLSYMPIWKFPDEEGACQ  
PCPINCTHSCVDLDDKGCPAEQRASPLTSIISAVVGILLVVVLGVVFGILIK  
RRQQKIRKYTMRRLLQETELVEPLTPSGAMPNQAQMRILKETELRKVKVLGS  
GAFGTVYKGIWIPDGENVKIPVAIKVLRENTSPKANKEILDEAYVMAGVGSP  
YVSRLLGICLTSTVQLVTQLMPYGCLLDHVRENRGRLGSQDLLNWCMQIAKG  
MSYLEDVRLVHRDLAARNVLVKSPNHVKITDFGLARLLDIDETEHADGGKV  
PIKWMALESILRRRFTHQSDVWSYGVTVWELMTFGAKPYDGIPAREIPDLE  
KGERLPQPPICTIDVYMIMVKCWMIDSECRPRFRELVSEFSRMARDPQRFV  
IQNEDLGASPLDSTFYRSLLEDDDMGDLVDAEEYLVPQQGFFCPDPAPGAG  
GMVHHRHRSSTRSGGDLTLGLEPSEEEAPRSPLAPSEGAGSDVEDGDLGM  
GAAKGLQSLPTHDPSPQLQRYSEDPTVPLPSETDGYVAPLTCSPQPEYVNQPD  
VRPQPPSPREGPLPAARPAGATLERPKTLSPGKNGVVKDVFAFGGAVENPEY  
LTPQGGAAPQPHPPAFSPAFDNLYYWDQDPPERGAPPSTFKGTPTAENPEY LGLDVPV Myeloid cell

MPLLLLPLLWALDMPDNPSTVQVEGLCVLVPCTFFHPIPYDK 773 surface  
NSPVHGYWFREGAIISRDSPVATNKLDEVQEETQGRFRLLGDP SRNNCSLS antigen CD33  
IVDARRRDNNGSYFFRMERGSTKY SYKSPQLSVHVTDLTHRPKILIPGTLEPG  
HSKNLTCSVSWACEQGTPIFSWLSAAPTSLGPRITHSSVLIITPRPQDHGT  
NLTCQVKFAGAGVTTERTIQLNVTYVPQNPTTGIFPGDGS GKQETRAGVVHG  
AIGGAGVTALLALCLCLIFFIVKTHRRKAARTAVGRNDTHPTTGSASPKHQK  
KSKLHGPTETSSCSGAAPT VEMDEELHYASLNFGHMNPSKDTSTEYSEVRTQ Telomerase  
MPRAPRCRAVRSLLRSHYREVLPLATFVRRLGPQGWRVLVQRGDPAAFRALVA 774 reverse  
QCLVCVPWDARPPPAAPSFRQV SCLKELVARVLQRLCERGAKNVLAFGFALL transcriptase  
DGARGGPPEAFTTSVRSYLPNTVTDALRGSGAWGLLLRRVGDDVLVHLLARC (TERT)  
ALFVLVAPSCAYQVCGPPPLYQLGAATQARPPPHASGP RRRLGCERAWNH SVR  
EAGVPLGLPAPGARRRGGSASRSLPLPKRP RRGA APEPERTPVGQGSWAHPG  
RTRGPSDRGFCVVS PARPAEEATSLEGALSGTRHSHPSVGRQHHAGPPSTSR  
PPRPWDTPCPPVYAETKHFLYSSGDKEQLRPSFLLSSLRPSLTGARRIVETI  
FLGSRPWMPGT PRRLPRLPQRYWQMRPLFLELLGNHAQCPYGVLLKTHCPLR  
AAVTPAAGVCAREKPQGSVA APEEEDTDPRRLVQLLRQHSSPWQVYGFVRAC  
LRLVPPGLWGS RHNERFLRNTKKFISLGKHAKLSLQELTWKMSVRDCAWL  
RRSPGVGCVPA AEHRLREEILAKFLHWLMSVYVVELLRSFFYVTETTFQKNR  
LFFYRKS VWSKLQSIGIRQHLKRVQLRELSEAEVRQHREARPALLTSRLRFI  
PKPDGLRP IVNMDYVVGARTFRREKRAERLT SRVKALFSVLNYERARRPGLL  
GASVLGLDDIHRAWRTFVL RVRAQDPPPELYFVKVDVTGAYDTIPQDRLTEV  
IASIIPQNTYCVRRYAVVQKAAHGHVRKA FKS HVSTLTDLQPYMRQFVAHL  
QETSPLRDAVVIEQSSSLNEASSGLEDVFLRFMCHHAVRIRGKSYVQCQGIP  
QGSILSTLLCSLCYGD MENKLFAGIRRDGLLLRLVDDFLLVTPHLTHAKTFL  
RTLVRGVPEYGC VVNLRKTVVNFVEDEALGGTAFVQMPAHGLFPWCGLLLD  
TRTLEVQSDYSSYARTSIRASLT FNRGFKAGRNMRRKLFGVLRLKCHSLFLD  
LQVNSLQTVCTNIYKILLQAYRFHACVLQLPFHQQVWKNPTFFLRVISDTA  
SLCYSILKAKNAGMSLGAKGAAGPLSEAVQWLCHQAFLCLKLTRHRVTYVPL  
LGLSRTAQTQLSRKLP GTTLTALEAAANPALPSDFKTILD Protein SSX2  
MNGDDAFARRPTVGAQIPEKIQKAFDDIAKYFSKEEWEKMKASEKIFYVYMK 775  
RKYEAMTKLGFKATLPPFMCNKRAEDFQGNLDNDPNRGNQVERPQM TFGRL  
QGISP KIMPKPAEEGNDSEEVPEASGPQNDGKELCPPGKPTTSEKIHESG  
PKRGEHAWTHRLRERKQLVIYEEISDPEEDDE Wilms tumor  
MGSDVRDLNALLPAVPSLGGGGGCALPVSGAAQWAPVLDFA PP GASAYGSLG 776 protein (WT1)  
GPAPPPAPPPPPPPPHSFIKQEPSWGAEPHEEQCLSAFTVHFSGQFTGTA  
GACRYGPF GPPPPSQASSGQARMFPNAPYLPSCLESQPAIRNQGYSTVTE DG  
TPSYGHTPSHHAAQFPNHSFKHEDPMGQQGSLGEQQYSVPPPVYGCHTPTDS  
CTGSQALLLRTPYSSDNL YQMTSQLECM TWNQMN LGATLKGVAAGSSSSVKW  
TEGQSNHSTGYESDNHTTPILCGAQYRIHTHGVERGIQDVRRVPGVAPT LVR  
SASETSEKRPFMCAYPGCNKRYFKLSHLQMHSRKHTGEKPYQCDFKDCERRF  
SRSDQLKRHQRRHTGVKPFQCKTCQRKFSRSDHLKTHTRTHTGKTSEKPFSC  
RWPSCQKKFARSDELVRHHNMHQ RNMTKLQLAL Sperm-  
MRQRLPSVTSLLLVALLPFGSSQARHVNHSATEALGELRERAPGQGINGFQ 777 associated  
LLRHAVKRDLLPRTPPYQVHISHQEARGPSFKICVGLGPRWARGCSTGNE antigen 11A  
KYHLPYAARDLQTFFLPFW (SPAG11A) BCR/ABL  
MVDPVGFAEAWKAQFPDSEPPRMELRSVGDIEQELERCKASIRRL EQEVNQE 778 fusion protein  
RFRMIYLQTL LAKEKKS YDRQRWGFRRAAQAPDGASEPRASASRPQAPADG isoform X3  
ADPPPAEEPEARPDGEGSPGKARPGTARRPGAAASGERDDRGPPASVAALRS  
NFERIRKGHGQPGADA EKPFYVNVEFHHERGLVKVNDKEVSDRISSLGSQAM  
QMERKKSQHGAGSSVGDASRPYRGRSSESSCGVDGDYEDAELNPRFLKDNL  
IDANGGSRPPWPPLEYQPYQSIYVGGMMEGEGKG PLLRSQSTSEQEKRLTWP  
RRSYSPRSFEDCGGGYTPDCSSNENLTSSEEDFSSGQSSRVSPSPTTYRMER  
DKSRSPSQNSQQSFDSSSPPTPQCHKRHRHCPVVVSEATIVGVRKTGQIWPN  
DGEGAFHGDADGSGFTPPGYGCAADRAEEQRRHQDGLPYIDDSPSSPHLSS  
KGRGSRDALVSGALESTKASELDLEKGLEMRKWVLSGILASEETYLSHLEAL  
LLPMKPLKAAATTSQPVLTSQQIETIFFKVPELYEIHKEFYDGLFPRVQQWS  
HQQRVGDLFQKLASQLGVYRVLGYNHNGEWCEAQTKNGQG WVPSNYITPVNS  
LEKHSWYHGPVSRNA AEYLLSSGINGSFLVRESESSPGQRSISLRYEGRVYH  
YRINTASDGKLYVSSESRENTLAELVHHHSTVADGLITTLHYPAPKR NKPTV  
YGVSPNYDKWEMERTDITMKHKLGGGQYGEVYEGVWKKYSLTVAVKTLKEDT  
MEVEEFLKEAAVMKEIKHPNLVQLLGVCTREPPFYIITEFMTYGNLLDYLRE

CNRQEVNAVVLVNAVLISSAMEYLEKKNFIHRDLAARNCLVGENHLVVKVAD  
FGLSRLMTGDTYTAHAGAKFPIKWTAPESLAYNKFSIKSDVWAFGVLLWEIA  
TYGMSPYPGIDLSQVYELLEKDYRMERPEGCEKVYELMRACWQWNPSPDRPS  
FAEIHQAFETMFQESSISDEVEKELGKQGVRGAVSTLLQAPELPTKTRTSRR  
AAEHRDTTDVPMPHSGKGQGESDPLDHEPAVSPLLPRKERGPPEGGLNEDER  
LLPKDKKTNLFSALIKKKKKKTAPTPPKRSSSFREMDGQPERRGAGEEEGRDI  
SNGALAFPTLDTADPAKSPKPSNGAGVPNGALRESGGSGFRSPHLWKKSSSTL Carcinoembry  
TSSRLATGEEEGGGSSSKRFLRSCSASCVPFHGAKDTEWRSVTLPRLDQSTGR 779 onic antigen-  
QFDSSTFGGHKSEKPALPRKRAGENRSDQVTRGTVTPPPRLVKKNEEAADDEV related cell  
FKDIMESSPGSSPPNLTpkPLRRQVTVAPASGLPHKEEAGKGSALGTPAAAE adhesion  
PVTPTSKAGSGAPGGTskGPAEESRVRHKKHSSSESPGRDKGKLSRLKPAPPP molecule 5  
PPAASAGKAGGKPSQSPSQEAAGEAVLGAKTKATSLVDAVNSDAAKPSQPGE (CEACAM5)  
GLKKPVLPATPKPQSAKPSGTPISPAPVPSTLPSASSALAGDQPSSTAFIPL  
ISTRVSLRKTQPPERIASGAITKGVVLDSTEALCLAISRNSEQMASHSAVL  
EAGKNLYTFCVSYVDSIQQMRNKFAFREAINKLENNLRELQICPATAGSGPA ATQDFSLLSSVKEISDIVQR  
MESPSAPPHRWCIWQRLLLLTASLLTFWNPPTTAKLTIESTPENVAEGKEVL  
LLVHNLQPQLFGYSWYKGERVDGNRQIIGYVIGTQQATPGPAYSGREIIPN  
ASLLIQNIIQNDTGfYTLHVIKSDLVNEEATGQFRVYPELPKPSISSNNSKP  
VEDKDAVAFTCEPETQDATYLWVWNNQSLPVSRLQLSNGNRTLTLNVNTRN  
DTASYKCETQNPVSARRSDSVILNVLYGPDAPTISPLNTSYRSGENLNLSCH  
AASNPPAQYSWFVNGTFQQSTQELFIPNITVNNSGsyTCQAHNSDTGLNRTT  
VTITITVYAEPKPFITSNNSNPVEDEDAVALTCEPEIQNTTYLWVWNNQSLP  
VSPRLQLSNDNRTLTLsvTRNDVGPYECGIQNKLSVDHSDPVILNVLYGPD  
DPTISPSYTYRPGVNLsLSCHAASNPPAQYSWLIDGNIQQHTQELFISNIT  
EKNSGLYTCQANNSASGHSRTTVKtITVSAELPKPSISSNNSKPVEDKDAVA  
FTCEPEAQNTTYLWVWNGQSLPVSRLQLSNGNRTLTLFNVNTRNDARAYVCG  
IQNSVSANRSDPVTLdVLYGPDTPHSPDSSYLsgANLNLSCHSASNPSPQ  
YSWRINGIPQQHTQVLfIAKITPNNNGTYACFVSNLATGRNNSIVKSITVSA  
SGTSPGLSAGATVGIMIGVLVGVALI Melanoma  
MPLEQRSQHCKPEEGLEARGEALGLVGAQAPATEEQEAASSSSTLVEVTLGE 780 antigen,  
VPAAESPDPPQSPQGASSLPTTMNYPLWSQSYEDSSNQEEEGPSTFPDLESE family A, 3  
FQAALSrkVAELVHfLLLKYRAREPVTKAEMLGsvVGNWQYFFPVIFSKAFS variant  
SLQLVFGIELMEVDPIGHLYIFATCLGLSYDGLLGDNQIMPKAGLLIIVLAI (MAGE-A3)  
IAREGDCAPEEKIWEELSVLEVFEGREDSILGDPKkLLTQHfVQENYLEYRQ  
VPGSDPACyEFLWGPRALVETSyVKVLHhMVKISGGPHISYPPLHEWVLREG EE Mucin-1  
MTPGTQSPFFLLLLLTvLTvVTGSGHASSTPGGEKETSATQRSSVPSSTEKN 781 (MUC1)  
AVSMTSSVLSSHSPGSGSSTTQGQDVTlAPATEPASGSAATWGQDVTsVPVT  
RPALGSTTPPAHDVTSAPDNKPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAH  
GVTSAPDTRPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTsAPDTRPAP  
GSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTs  
APDTRPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTsAPDTRPAPGSTA  
PPAHGVTSAPDTRPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTsAPDT  
RPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAH  
GVTSAPDTRPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTsAPDTRPAP  
GSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTs  
APDTRPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTsAPDTRPAPGSTA  
PPAHGVTSAPDTRPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTsAPDT  
RPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAHGVTsAPDTRPAPGSTAPPAH  
GVTSAPDNRPALGSTAPPVHNVISASGSASGSASTLVHNGTSARATTTpASK  
STPFsIPSHHSDTPtTLASHSTKTDASSTHSSVPPLTSSNHSTSPQLSTGV  
SFFFLSFHISNLQFNSSLEDpSTDYYQELQRDISEMFLQIYKQGgFLGLSNI  
KFRPGSVVVQLTLAFREGTINVHDVETQFNQYKTEAASRYNLtISDVSVSDV  
PFPFSAQSGAGVPGWGIALLVLCVLVALAIvYLIAlAVCQCRRKNYGQLDI  
FPARDTYHPMSEYPTYHthGRYVPPSSDRSPYEkVSAGNGGSSLSyTNPAV AATSANL Epithelial cell  
MAPPQVLAfGLLLAAATATFAAAQEECVcENYKLAVNCFVNNNRQCQCTSVG 782 adhesion  
AQNTVICSKLAAKCLVMKAEMNGSKLGRRAKPEGALQNNDGLYDPDCDESGL molecule

FKAKNGTSMCVNTAGVRRITDKDTEITCSERVRTYWIHILKHKAREKP precursor  
YDSKSLRTALQKEITTRYQLDPKFITSILYENNVITIDLQVNSSQKTQNDVD (EpCAM)  
IADVAYYFEKDVKGESLFHSSKMDLTVNGEQLDLDPGQTLIYYVDEKAPEFS  
MQGLKAGVIAVIVVVVIAVVAGIVVLVISRKKRMAKYEKAEIKEMGEMHREL NA

(122) In some embodiments, cancer vaccines containing immunogenic peptides from tumor specific antigens (e.g., heteroclitic epitopes and cryptic epitopes) are generated in vivo (e.g., in tumor cells in a subject) or ex vivo (e.g., in tumor cells isolated from a subject). In some embodiments, the tumor cells are treated with the nucleobase editors to generate the immunogenic peptides and are irradiated and administered to the subject as whole-cell cancer vaccines.

(123) To edit the genes encoding the tumor associated antigens, the nucleobase editor and/or the guide nucleotide sequence is introduced into the cell (e.g., a tumor cell) where the editing occurs. In some embodiments, nucleic acid molecules (e.g., expression vectors) encoding the nucleobase editors and/or the guide nucleotide sequences are delivered into the cell, resulting in co-expression of nucleobase editors and/or the guide nucleotide sequences in the cell. The nucleic acid molecules encoding the nucleobase editors and/or the guide nucleotide sequences may be delivered into the cell using any known methods in the art, e.g., transfection (e.g., transfection mediated by cationic liposomes), transduction (e.g., via viral infection) and electroporation. In some embodiments, an isolated nucleobase editor/gRNA complex is delivered. Methods of delivering an isolated protein to a cell is familiar to those skilled in the art. For example, the isolated nucleobase editor in complex with a gRNA be associated with a supercharged, cell-penetrating protein or peptide, which facilitates its entry into a cell (e.g., as described in PCT Application Publication WO2010129023 and US Patent Application Publication US20150071906, incorporated herein by reference). In some embodiments, the isolated nucleobase editor in complex with a gRNA may be delivered by a cationic transfection reagent, e.g., the Lipofectamine CRISPRMAX Cas9 Transfection Reagent from Thermofisher Scientific. In some embodiments, the nucleobase editor and the gRNA may be delivered separately. Other suitable delivery methods may also be used, e.g., AAV mediated gene transfer. Strategies for delivery a Cas9-based genome editing agent (e.g., the nucleobase editor described herein) using AAV have been described, e.g., in Zetsche et al., *Nature Biotechnology* 33, 139-142 (2015), incorporated herein by reference.

(124) In some embodiments, once generated, the immunogenic peptide is displayed on the surface of the tumor cell via the MHC class I antigen presentation pathway. In some embodiments, the immunogenic peptide is displayed on the surface of an antigen presenting cell (APC) via the MHC class II antigen presentation pathway. In some embodiments, the APC is selected from the group consisting of: tumor cells, dendritic cells, mononuclear phagocytes, thymic epithelial cells, and B cells. In some embodiments, the immunogenic peptide elicits an adaptive immune response against the tumor-specific antigen where the peptide is derived from. In some embodiments, the immunogenic peptide elicits an adaptive immune response against the tumor. In some embodiments, the adaptive immune response comprises promoting the maturation of dendritic cells, activation of CD4<sup>+</sup> T lymphocytes, (T helper cells) activation of CD8<sup>+</sup> T lymphocytes (cytotoxic T cells), activation and maturation of B lymphocytes, and/or production of tumor antigen-specific antibodies.

(125) T helper cells (TH cells) assist other white blood cells in immunologic processes, including maturation of B cells into plasma cells and memory B cells, and activation of cytotoxic T cells and macrophages. These cells are also known as CD4<sup>+</sup> T cells because they express the CD4 glycoprotein on their surfaces. Helper T cells become activated when they are presented with peptide antigens by MHC class II molecules, which are expressed on the surface of antigen-presenting cells (APCs). Once activated, they divide rapidly and secrete small proteins called cytokines that regulate or assist in the active immune response. These cells can differentiate into one of several subtypes, including TH1, TH2, TH3, TH17, TH9, or TFH, which secrete different cytokines to facilitate different types of immune responses. Signaling from the APC directs T cells into particular subtypes.

(126) Cytotoxic T cells (e.g., TC cells, CTLs, T-killer cells, killer T cells) destroy virus-infected cells and tumor cells, and are also implicated in transplant rejection. These cells are also known as CD8<sup>+</sup> T cells since they express the CD8 glycoprotein at their surfaces. These cells recognize their targets by binding to antigen associated with MHC class I molecules, which are present on the surface of all nucleated cells. Through IL-10, adenosine, and other molecules secreted by regulatory T cells, the CD8<sup>+</sup> cells can be inactivated to an anergic state, which prevents autoimmune diseases.

(127) Most cytotoxic T cells express T-cell receptors (TCRs) that can recognize a specific antigen. Antigens inside a cell are bound to class I MHC molecules, and brought to the surface of the cell by the class I MHC molecule, where they can be recognized by the T cell. If the TCR is specific for that antigen, it binds to the complex of the class I MHC molecule and the antigen, and the T cell destroys the cell, e.g., via inducing apoptosis. In order for the TCR to bind to the class I MHC molecule, the former must be accompanied by a glycoprotein called CD8, which binds to the constant portion of the class I MHC molecule. Therefore, these T cells are called CD8<sup>+</sup> T cells.

(128) Natural killer T cells (NKT cells—not to be confused with natural killer cells of the innate immune system) bridge the adaptive immune system with the innate immune system. Unlike conventional T cells that recognize peptide antigens presented by major histocompatibility complex (MHC) molecules, NKT cells recognize glycolipid antigen presented by a molecule called CD1d. Once activated, these cells can perform functions ascribed to both Th and Tc cells (i.e., cytokine production and release of cytolytic/cell killing molecules). They are also able to recognize and eliminate some tumor cells and cells infected with microorganisms, e.g., bacteria or virus.

(129) Memory T cells are a subset of antigen-specific T cells that persist long-term after an initial T cell response. They quickly expand to large numbers of effector T cells upon re-exposure to their cognate antigen, thus providing the immune



system with “memory” against past antigens. The cancer vaccine described herein provides the immune system with “memory” against the tumor specific antigen, thereby eliciting strong immune response against newly emerged cancer cells or metastasized cancer cells.

(130) Regulatory T cells (suppressor T cells) are crucial for the maintenance of immunological tolerance. Their major role is to shut down T cell-mediated immunity toward the end of an immune reaction and to suppress autoreactive T cells that escaped the process of negative selection in the thymus. Suppressor T cells along with Helper T cells can collectively be called Regulatory T cells due to their regulatory functions.

(131) B cell activation occurs in the secondary lymphoid organs (SLOs), such as the spleen and lymph nodes. After B cells mature in the bone marrow, they migrate through the blood to SLOs, which receive a constant supply of antigen through circulating lymph. At the SLO, B cell activation begins when the B cell binds to an antigen via its BCR. The antigen can either be free-floating or presented by APCs such as macrophages or dendritic cells (DCs), and include proteins, glycoproteins, polysaccharides, whole virus particles, and whole bacterial cells. Some subtypes of B cell preferentially undergo T cell-dependent activation while other subtypes of cells preferentially undergo T cell-independent activation.

(132) Antigens that activate B cells with the help of T-cell are known as T cell-dependent (TD) antigens and include foreign proteins. They are named as such because they are unable to induce a humoral response in organisms that lack T cells. B cell response to these antigens takes multiple days, though antibodies generated have a higher affinity and are more functionally versatile than those generated from T cell-independent activation.

(133) Once a BCR binds a TD antigen, the antigen is taken up into the B cell through receptor-mediated endocytosis, degraded, and presented to T cells as peptide pieces in complex with MHC-II molecules on the cell membrane. T helper (TH) cells, typically follicular T helper (TFH) cells, that were activated with the same antigen recognize and bind these MHC-II-peptide complexes through their T cell receptor (TCR). Following TCR-MHC-II-peptide binding, T cells express the surface protein CD40L as well as cytokines such as IL-4 and IL-21. CD40L serves as a necessary co-stimulatory factor for B cell activation by binding the B cell surface receptor CD40, which promotes B cell proliferation, immunoglobulin class switching, and somatic hypermutation as well as sustains T cell growth and differentiation. T cell-derived cytokines bound by B cell cytokine receptors also promote B cell proliferation, immunoglobulin class switching, and somatic hypermutation as well as guide differentiation. After B cells receive these signals, they are considered activated.

(134) Activated B cells participate in a two-step differentiation process that yields both short-lived plasmablasts for immediate protection and long-lived plasma cells and memory B cells for persistent protection. The first step, known as the extrafollicular response, occurs outside of lymphoid follicles but still in the SLO. During this step activated B cells proliferate, may undergo immunoglobulin class switching, and differentiate into plasmablasts that produce early, weak antibodies mostly of class IgM. The second step consists of activated B cells entering a lymphoid follicle and forming a germinal center (GC), which is a specialized microenvironment where B cells undergo extensive proliferation, immunoglobulin class switching, and affinity maturation directed by somatic hypermutation. These processes are facilitated by TFH cells within the GC and generate both high-affinity memory B cells and long-lived plasma cells. Resultant plasma cells secrete large amounts of antibody and either stay within the SLO or, more preferentially, migrate to bone marrow.

(135) Antigens that activate B cells without T cell help are known as T cell-independent (TI) antigens and include foreign polysaccharides and unmethylated CpG DNA. They are named as such because they are able to induce a humoral response in organisms that lack T cells. B cell response to these antigens is rapid, though antibodies generated tend to have lower affinity and are less functionally versatile than those generated from T cell-dependent activation.

(136) As with TD antigens, B cells activated by TI antigens need additional signals to complete activation, but instead of receiving them from T cells, they are provided either by recognition and binding of a common microbial constituent to toll-like receptors (TLRs) or by extensive crosslinking of BCRs to repeated epitopes on a bacterial cell. B cells activated by TI antigens go on to proliferate outside of lymphoid follicles but still in SLOs (GCs do not form), possibly undergo immunoglobulin class switching, and differentiate into short-lived plasmablasts that produce early, weak antibodies mostly of class IgM, but also some populations of long-lived plasma cells.

(137) Memory B cell activation begins with the detection and binding of their target antigen, which is shared by their parent B cell. Some memory B cells can be activated without T cell help, such as certain virus-specific memory B cells, but others need T cell help. Upon antigen binding, the memory B cell takes up the antigen through receptor-mediated endocytosis, degrades it, and presents it to T cells as peptide pieces in complex with MHC-II molecules on the cell membrane. Memory T helper (TH) cells, typically memory follicular T helper (TFH) cells, that were derived from T cells activated with the same antigen recognize and bind these MHC-II-peptide complexes through their TCR. Following TCR-MHC-II-peptide binding and the relay of other signals from the memory TFH cell, the memory B cell is activated and differentiates either into plasmablasts and plasma cells via an extrafollicular response or enter a germinal center reaction where they generate plasma cells and more memory B cells.

(138) In some embodiments, the adaptive immune response results in the killing tumor cells, reducing tumor burden, reducing tumor size, and/or preventing metastasis. In some embodiments, the adaptive immune response is active against neo-epitopes associated with spontaneous somatic mutations. In some embodiments, the neo-epitope is specific to the lineage of tumor cells.

(139) In some embodiments, the adaptive immune response elicited by the heteroclitic or cryptic epitopes is cross-reactive with the native tumor-specific antigen. In some embodiments, the adaptive immune response elicited by the heteroclitic or cryptic epitopes is cross-reactive with neoepitopes arising from spontaneous mutations occurring in the tumor specific antigen.

(140) There are advantages associated with using heteroclitic epitopes in clinical applications. For example, heteroclitic epitopes have the ability to break/overcome tolerance by reversing a state of T cell anergy, activating non-tolerized cross-reactive clones of T cells, or by mediating “immune deviation,” i.e., the type of CTL produced, such as Th1 or Th2. Recent studies indicate that heteroclitic epitopes are immunogenic (Zaremba, et al., *Cancer Research*, 57:4570 (1997); Rivoltini, et al., *Cancer Research*, 59:301 (1999); Selby, et al., *The Journal of Immunology* 162(2):669 (1999), the entire contents of each of which are incorporated herein by reference) in that they are capable of inducing CTLs that recognize endogenously processed epitopes. This is confirmed by studies in different immunological systems (Zugel, et al., *J. Immunol.*, 161:1705 (1998), Wang, et al., *J. Exp. Med.*, 190:983 (1999), Men, et al., *J. Immunol.*, 162:3566, (1999), the entire contents of each of which are incorporated herein by reference). For example, studies by Zugel et al. have shown that T cell tolerance to an immunodominant T cell epitope in adult mice can be overcome by immunization with heteroclitic cross-reactive peptide analogs of that peptide.

(141) In some embodiments, heteroclitic epitopes or cryptic epitopes modulate cytokine production from T cells (Pfeiffer, et al., *J. Exp. Med.*, 181:1569 (1995), Tao, et al., *J. Immunol.*, 158:4237 (1997), Salazar, et al., *Int. J. Cancer* 85(6):829-38 (2000), Nicholson, et al., *Int. Immunol.* 12(2):205-13 (2000), the entire contents of each of which are incorporated herein by reference). The immune deviation induced by such analogs has implications in several disease states, where generation of a specific subset of Th cell responses correlate with tumor regression (Zitvogel, et al., *J. Exp. Med.*, 183:87 (1996), Celluzzi, et al., *J. Exp. Med.* 183:283 (1996), the entire contents of each of which are incorporated herein by reference) or affected the clinical outcome of autoimmune or infectious disease (Romagnani, et al., *Annu. Rev. Immunol.*, 12:227-57 (1994), the entire contents of which are incorporated herein by reference). Thus, immunization with heteroclitic epitopes offers the capacity to modulate cytokine production by induction of specific subsets of effector T cells, thereby altering the course of disease.

(142) In some embodiments, heteroclitic epitopes offer an advantage in drug development since significantly smaller amounts of peptide are needed for treatment doses, due to their strong biological potency. This feature overcomes certain manufacturing and toxicity concerns. In this regard, it has been shown that a heteroclitic analog of a MART-1 peptide (Rivoltini, et al., *Cancer Research* 59:301 (1999), the entire contents of which are incorporated herein by reference), which generated antigen specific T cells in melanoma patients, was active at much lower concentrations than the native epitope. Similar results were reported by Schlom and colleagues (Zaremba, et al., *Cancer Research* 57:4570 (1997), the entire contents of which are incorporated herein by reference) regarding heteroclitic analog of the CEA derived CAP1 epitope.

(143) Nucleobase Editors

(144) The methods of generating immunogenic peptides or epitopes from tumor associated antigens as cancer vaccines described herein, are enabled by the use of the nucleobase editors. As described herein, a nucleobase editor is a fusion protein comprising: (i) a programmable DNA binding protein domain; and (ii) a deaminase domain. It is to be understood that any programmable DNA binding domain may be used in the based editors.

(145) In some embodiments, the programmable DNA binding protein domain comprises the DNA binding domain of a zinc finger nuclease (ZFN) or a transcription activator-like effector domain (TALE). In some embodiments, the programmable DNA binding protein domain may be programmed by a guide nucleotide sequence, and is thus referred as a “guide nucleotide sequence-programmable DNA binding-protein domain.” In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a nuclease inactive Cas9, or dCas9. A dCas9, as used herein, encompasses a Cas9 that is completely inactive in its nuclease activity, or partially inactive in its nuclease activity (e.g., a Cas9 nickase). Thus, in some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a Cas9 nickase. In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a nuclease inactive Cpf1. In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a nuclease inactive Argonaute. In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a nuclease inactive CasX or CasY, e.g., as described in Burstein et al., *New CRISPR-Cas systems from uncultivated microbes*, *Nature* 542, 237-241, 2017, incorporated herein by reference.

(146) In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a dCas9 domain. In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a Cas9 nickase. In some embodiments, the dCas9 domain comprises the amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 3. In some embodiments, the dCas9 domain comprises an amino acid sequence that is at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to any one of the Cas9 domains provided herein (e.g., SEQ ID NOs: 11-260), and comprises mutations corresponding to D10X (X is any amino acid except for D) and/or H840X (X is any amino acid except for H) in SEQ ID NO: 1. In some embodiments, the dCas9 domain comprises an amino acid sequence that is at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to any one of the Cas9 domains provided herein (e.g., SEQ ID NOs: 11-260), and

comprises mutations corresponding to D10A and/or H840A in SEQ ID NO: 1. In some embodiments, the Cas9 nickase comprises an amino acid sequence that is at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to any one of the Cas9 domains provided herein (e.g., SEQ ID NOs: 11-260), and comprises mutations corresponding to D10X (X is any amino acid except for D) in SEQ ID NO: 1 and a histidine at a position correspond to position 840 in SEQ ID NO: 1. In some embodiments, the Cas9 nickase comprises an amino acid sequence that is at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to any one of the Cas9 domains provided herein (e.g., SEQ ID NOs: 11-260), and comprises mutations corresponding to D10A in SEQ ID NO: 1 and a histidine at a position correspond to position 840 in SEQ ID NO: 1. In some embodiments, variants or homologues of dCas9 or Cas9 nickase (e.g., variants of SEQ ID NO: 2 or SEQ ID NO: 3, respectively) are provided which are at least about 70% identical, at least about 80% identical, at least about 90% identical, at least about 95% identical, at least about 98% identical, at least about 99% identical, at least about 99.5% identical, or at least about 99.9% to SEQ ID NO: 2 or SEQ ID NO: 3, respectively, and comprises mutations corresponding to D10A and/or H840A in SEQ ID NO: 1. In some embodiments, variants of Cas9 (e.g., variants of SEQ ID NO: 2) are provided having amino acid sequences which are shorter, or longer than SEQ ID NO: 2, by about 5 amino acids, by about 10 amino acids, by about 15 amino acids, by about 20 amino acids, by about 25 amino acids, by about 30 amino acids, by about 40 amino acids, by about 50 amino acids, by about 75 amino acids, by about 100 amino acids, or more, provided that the dCas9 variants comprise mutations corresponding to D10A and/or H840A in SEQ ID NO: 1. In some embodiments, variants of Cas9 nickase (e.g., variants of SEQ ID NO: 3) are provided having amino acid sequences which are shorter, or longer than SEQ ID NO: 3, by about 5 amino acids, by about 10 amino acids, by about 15 amino acids, by about 20 amino acids, by about 25 amino acids, by about 30 amino acids, by about 40 amino acids, by about 50 amino acids, by about 75 amino acids, by about 100 amino acids, or more, provided that the dCas9 variants comprise mutations corresponding to D10A and comprises a histidine at a position corresponding to position 840 in SEQ ID NO: 1. (147) Additional suitable nuclease-inactive dCas9 domains will be apparent to those of skill in the art based on this disclosure and knowledge in the field, and are within the scope of this disclosure. Such additional exemplary suitable nuclease-inactive Cas9 domains include, but are not limited to, D10A/H840A, D10A/D839A/H840A, D10A/D839A/H840A/N863A mutant domains (See, e.g., Prashant et al., *Nature Biotechnology*. 2013; 31(9): 833-838, which are incorporated herein by reference), or K603R (See, e.g., Chavez et al., *Nature Methods* 12, 326-328, 2015, which is incorporated herein by reference).

(148) In some embodiments, the nucleobase editors utilized in the present invention comprise a Cas9 domain with decreased electrostatic interactions between the Cas9 domain and a sugar-phosphate backbone of a DNA, as compared to a wild-type Cas9 domain. In some embodiments, a Cas9 domain comprises one or more mutations that decreases the association between the Cas9 domain and a sugar-phosphate backbone of a DNA. In some embodiments, the nucleobase editors described herein comprises a dCas9 (e.g., with D10A and H840A mutations) or a Cas9 nickase (e.g., with D10A mutation), wherein the dCas9 or the Cas9 nickase further comprises one or more of a N497X, R661X, Q695X, and/or Q926X mutation of the amino acid sequence provided in SEQ ID NO: 1, or a corresponding mutation in any of the amino acid sequences provided in SEQ ID NOs: 11-260, wherein X is any amino acid. In some embodiments, the nucleobase editors described herein comprises a dCas9 (e.g., with D10A and H840A mutations) or a Cas9 nickase (e.g., with D10A mutation), wherein the dCas9 or the Cas9 nickase further comprises one or more of a N497A, R661A, Q695A, and/or Q926A mutation of the amino acid sequence provided in SEQ ID NO: 1, or a corresponding mutation in any of the amino acid sequences provided in SEQ ID NOs: 11-260. In some embodiments, the dCas9 domain (e.g., of any of the nucleobase editors provided herein) comprises the amino acid sequence as set forth in any one of SEQ ID NOs: 2-9. In some embodiments, the nucleobase editor comprises the amino acid sequence as set forth in any one of SEQ ID NOs: 293-302 and 321.

(149) TABLE-US-00015 Cas9 variant with decreased electrostatic interactions between the Cas9 and DNA backbone

DKKYSIGLAIGTNSVGWAVITDEYKVPSKKFKVLGNTDRHSIKKNLIGALLFDSGETAEATRLKRTARR  
RYTRRKNRICYLQEIFSNEMAKVDDSFHRLSESLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLRK  
KLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDAK  
AILSARLSKSRLENLIAQLPGKKNGLFGNLIALSLGLTPNFKSNFDLAEDAKLQLSKDITYDDDLNLL  
AQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLKALVRQQLPEKYK  
EIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKNREDLLRKQRTFDNGSIPHQIHLGE  
LHAILRRQEDFYFPFLKDNREKIEKILTFRIPYYVGPLARGNSRFAWMTRKSEETITPWNFEEVVDKGASA  
QSFIERMTAFDKNLPNEKVLPKHSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFKTN  
RKVTVKQLKEDYFKKIECFDSVEISGVEDRNFNASLGTYHDLKIIKDKDFLDNEENEDILEDIVLTTLFE  
DREMIEERLKTYAHLFDDKVMKQLKRRRYTGWGALSRKLINGIRDKQSGKTILDFLKSDGFANRNF  
ALIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIKKGILQTVKVVDELVKVMGRHKPENIVIE  
ARENQTTQKGQKNSRERMKRIEKGELGSQILKEHPVENTQLQNEKLYLYLQNGRDMYVDQELDIN  
RLSDYDVDHIVPQSFLKDDSIDNKVLTRSDKNRGKSDNPSEEVVKKMKNYWRQLLNAKLITQRKFD  
NLTKAERGGLSELDKAGFIKRQLVETRAITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFR

KDFQFYKYNVYALNNAVVGTAALIKKYPKLESEFVYGDYVVRKMIAKSEQEIGKAT  
 KYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVLSPQVNVKKTEVQTG  
 GFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGKSKKLKSVKELLGITIMERS  
 SFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLASH  
 YEKLKGSPEDEQKQLFVEQHKHYLDEIIIEQISEFSKRVLADANLDKVL SAYNKH RD KPIREQAENIIH  
 LFTLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQ SITGLYETRIDLSQLGGD (SEQ ID NO: 9,  
 mutations relative to SEQ ID NO: 1 are bolded and underlined) High fidelity nucleobase editor  
 (SEQ ID NO: 321)

MSSETGPVAVDPTLRRRIEPHEFEVFFDPREL RKETCLLYEINWGGRH SIWRHTSQNTNKHVEVNFIEKF  
 TTERYFCPNTRCSITWFLSWSPCGECSRAITEFLSRYPHVTLFIYIARLYHHADPRNRQGLRDLISSGVTI  
 QIMTEQESGYCWRNFVNYSNEAHWPYPHLLWVRLYVLELYCIIILGLPPCLNLRKQPKLTFTTIALQ  
 SCHYQRLPPHILWATGLKSGSETPGTSESATPESDKKYSIGLAIGTNSVGWAVITDEYKVP SKKFKVLG  
 NTDRHSIKKNLIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIFS NEMAKVDDSFHRLEESFL  
 VEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLRKKLV DSTDKADLR LIYLALAHMIKFRGHFLIEGDLN  
 PDNSDVKLFIQLVQTYNQLFEENPINASGVDAKAILSARLSKSRLENLIAQLPGEKKNGLFGNLIALS  
 LGLTPNFKSNFDLAEDAKLQLSKD TYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITK  
 APLSASMIKRYDEHHQDLTLLKALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKM  
 DGTEELLVKLNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFY PFLKDNREKIEKILTRIPYYVGP  
 LARGNSRFAWMTRKSEETITPWNFEEVVDKGASAQSFIERMTAFDKNLPNEKVLPKHSLLYEYFTVYN  
 ELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASL  
 GTYHDLLKHKDKDFLDNEENEDILEDIVLTTLTFEDREMIEERLKYAHLFDDKVMKQLKRRRYTGWG  
 ALSRKLINGIRDKQSGKTILDFLKSDGFANRNFMALIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGS  
 PAIKKGILQTVKVVDDELVKVMGRHKPENIVIE MARENQTTQKGQKNSRERMKRIEEGIKELGSQILKEH  
 PVENTQLQNEKLYLYLQNGRDMYVDQELDINRLSDYDVDHIVPQSFLKDDSIDNKVLTRSDKNRGKS  
 DNV PSEEVVKMKKNYWRQLLNAKLITQRKFDNLTKAERGGLSELDKAGFIKRQLVETRAITKHVAQIL  
 DSRMNTKYDENDKLIREVKVITLKS KLVSDFRKDFQFYKVREINNYHHAHDAYLNAVVG TALIKKYPK  
 LESEFVYGDYK VYDVRKMIAKSEQEIGKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIV  
 WDKGRDFATVRKVLSPQVNVKKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVA  
 YSVLVVAKEKGKSKKLKSVKELLGITIMERS SFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELENGR  
 KRMLASAGELQKGNELALPSKYVNFLYLASHYEKLKGSPEDEQKQLFVEQHKHYLDEIIIEQISEFSKR  
 VILADANLDKVL SAYNKH RD KPIREQAENIIHLFTLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIH  
 QSITGLYETRIDLSQLGGD

(150) In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a single effector of a microbial CRISPR-Cas system. Single effectors of microbial CRISPR-Cas systems include, without limitation, Cas9, Cpf1, C2c1, C2c2, and C2c3. Typically, microbial CRISPR-Cas systems are divided into Class 1 and Class 2 systems. Class 1 systems have multisubunit effector complexes, while Class 2 systems have a single protein effector. Cas9 and Cpf1 are Class 2 effectors. In addition to Cas9 and Cpf1, three distinct Class 2 CRISPR-Cas systems (C2c1, C2c2, and C2c3) have been described by Shmakov et al., “Discovery and Functional Characterization of Diverse Class 2 CRISPR Cas Systems”, *Mol. Cell*, 2015 Nov. 5; 60(3): 385-397, the entire contents of which are herein incorporated by reference. Effectors of two of the systems, C2c1 and C2c3, contain RuvC-like endonuclease domains related to Cpf1. A third system, C2c2 contains an effector with two predicted HEPN RNase domains. Production of mature CRISPR RNA is tracrRNA-independent, unlike production of CRISPR RNA by C2c1. C2c1 depends on both CRISPR RNA and tracrRNA for DNA cleavage. Bacterial C2c2 has been shown to possess a unique RNase activity for CRISPR RNA maturation distinct from its RNA-activated single-stranded RNA degradation activity. These RNase functions are different from each other and from the CRISPR RNA-processing behavior of Cpf1. See, e.g., East-Seletsky, et al., “Two distinct RNase activities of CRISPR-C2c2 enable guide-RNA processing and RNA detection”, *Nature*, 2016 Oct. 13; 538(7624):270-273, the entire contents of which are hereby incorporated by reference. In vitro biochemical analysis of C2c2 in *Leptotrichia shahii* has shown that C2c2 is guided by a single CRISPR RNA and can be programmed to cleave ssRNA targets carrying complementary protospacers. Catalytic residues in the two conserved HEPN domains mediate cleavage. Mutations in the catalytic residues generate catalytically inactive RNA-binding proteins. See e.g., Abudayyeh et al., “C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector,” *Science*, 2016 Aug. 5; 353(6299), the entire contents of which are hereby incorporated by reference.

(151) The crystal structure of *Alicyclobacillus acidoterrastris* C2c1 (AacC2c1) has been reported in complex with a chimeric single-molecule guide RNA (sgRNA). See, e.g., Liu et al., “C2c1-sgRNA Complex Structure Reveals RNA-Guided DNA Cleavage Mechanism”, *Mol. Cell*, 2017 Jan. 19; 65(2):310-322, incorporated herein by reference. The crystal structure has also been reported for *Alicyclobacillus acidoterrestris* C2c1 bound to target DNAs as ternary complexes. See, e.g., Yang et al., “PAM-dependent Target DNA Recognition and Cleavage by C2C1 CRISPR-Cas endonuclease”, *Cell*, 2016 Dec. 15; 167(7):1814-1828, the entire contents of which are hereby incorporated by reference. Catalytically competent conformations of AacC2c1, both with target and non-target DNA strands, have been captured independently positioned within a single RuvC catalytic pocket, with C2c1-mediated cleavage resulting in a staggered seven-nucleotide break of target DNA. Structural comparisons between C2c1 ternary complexes and previously identified

Cas9 and Cpf1 counterparts demonstrate the diversity of mechanisms used by CRISPR-Cas9 systems. (152) In some embodiments, the nucleobase editors described herein comprise a C2c1, a C2c2, or a C2c3 protein. In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a C2c1 protein. In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a C2c2 protein. In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a C2c3 protein. In some embodiments, the guide nucleotide sequence-programmable DNA binding protein comprises an amino acid sequence that is at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to a naturally-occurring C2c1, C2c2, or C2c3 protein. In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a naturally-occurring C2c1, C2c2, or C2c3 protein. In some embodiments, the guide nucleotide sequence-programmable DNA binding protein comprises an amino acid sequence that is at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to any one of SEQ ID NOs: 1057-1059. In some embodiments, the guide nucleotide sequence-programmable DNA binding protein comprises an amino acid sequence of any one SEQ ID NOs: 1057-1059. It should be appreciated that C2c1, C2c2, or C2c3 from other bacterial species may also be used in accordance with the present disclosure.

(153) C2c1 (uniprot.org/uniprot/TOD7A2 #) sp|TOD7A2|C2C1\_ALIAG CRISPR-associated endonuclease C2c1 OS=*Alicyclobacillus acidoterrestris* (strain ATCC 49025/DSM 3922/CIP 106132/NCIMB 13137/GD3B) GN=c2c1 PE=1 SV=1

(154) TABLE-US-00016 C2c1 (uniprot.org/uniprot/TOD7A2#) sp|TOD7A2|C2C1\_ALIAG CRISPR-associated endonuclease C2c1 OS = *Alicyclobacillus acidoterrestris* (strain ATCC 49025/DSM 3922/CIP 106132/MCIMB 13137/GD3B) GN = c2c1 PE = 1 SV = 1 (SEQ ID NO: 1057)

MAVKSIVKLRLLDDMPEIRAGLWKLHKEVNAGVRRYYTEWLSLLRQENLYRRSPNGDGEQEC DKTAE  
ECKAELLERLRARQVENGHRRGPA GSDELLQLARQLYELLVPQAIGAKGDAQQIARKFLSPLADKDAV  
GGLGIAKAGNKPRWVRMREAGEPGWEEKEKAETRKSADRTADVLRALADFLGLKPLMRVYTDSEMS  
SVEWKPLRKQAVRTWDRDMFQQA IERMMSWESWNQRVGQEYAKLVEQKNRFEQKNFVGGQEHLV  
HLVNQLQQDMKEASPGLESKEQTAHYVTGRALRGSDKVF EKWGKLAPDAPFDLYDAEIKNVQRRNT  
RRFGSHDLFAKLAEPEYQALWREDASFLTRYAVYNSILRKLNHAKMFATFTLPDATAHPIWTRFDKLG  
GNLHQYTFLFNEFGERRHAIRFHKL LKVENGVA REVDVTVPI SMSEQLDNLLPRDPNEPIALYFRDYG  
AEQHFTGEFGGAKIQCR RDQLAHMHRRGARDVYLVNSVRVQSQSEARGERRPPYAAVFR LVGDNH  
RAVHFHDKLS DYLAHPDDGKLGSEGLLSGLRVMSVDLGLRTSASISVFRVARKDELKPN SKGRVPFFF  
PIKGNNDNLVAVHERS QLLKLPGETESKDLRAIREERQRTLRLQLRTQLAYLRLLVRC SEDVGRRRERSW  
AKLIEQPVDAANHMTPDWREAFENELQKLKSLHGICSDKEWMDAVYESVRRVWRHMGKQVRDWRK  
DVRSGERP KIRGYAKDVVGGNSIEQIEYLERQYKFLKSW SFFGKVSQVIRAEKGSRAITLREHIDHAK  
EDRLKKLADRIIMEALGYVYALDERGKGKWKVAKYPPCQLILLEELSEYQFNDRPPSENNQLMQWSH  
RGVFQELINQAQVHDL LVGTMYYAAFSSRFDARTGAPGIRCRRVPARCTQEHNPEPFPWWLNKFVVEHT  
LDACPLRADDLIPTGEGEIVSPFSAE EGDHFHQIHADLNAAQN LQQRLWSDFDISQIRLRCDWGEVDGE  
LVLIPRLTGKRTADSYSNKVFYTN TGVTYYERERGKKRRKVFAQEKLSEEEAELLVEADEAREKSVVL  
MRDPSGIINRGNWTRQKEFW SMVNQRIEGYLVKQIRSRVPLQDSACENTGDI C2c2

(uniprot.org/uniprot/P0DOC6) >sp|P0DOC6|C2C2\_LEPSD CRISPR-associated endonuclease C2c2 OS = *Leptotrichia shahii* (strain DSM 19757/CCUG 47503/CIP 107916/JCM 16776/LB37) GN = c2c2 PE = 1 SV = 1 (SEQ ID NO: 1058)

MGNLFGHKRWYEV RDKKDFKIKRKVKVKRNYDGNKYILNINENNNKEKIDNNKFIRKYINYKKNDNI  
LKEFTRKFHAGNILFKLKGKEGIIRIENDDFLETEE VVLYIEAYGKSEKLKALGITKKKIIDEAIRQGITK  
DDKKIEIKRQENEEIEIDIRDEYTNKTLNDCSILRIIENDELETKKSIYEIFKNINMSLYKIEKIENETEK  
VFENRYEEHLREKLLKDDKIDVILTNFMEIREKIKSNLEILGFVKFYLVN VGGDKKKSKNKKMLVEKIL  
NINVDLTVEDIAD FVIKELEFWNITKRIEKVKVNNEFLEKRRNR TYIKSYVLLDKHEKFKIERENKKDK  
IVKFFVENIKNSI KEKIEKILA EFKIDELIKLEKELKKGNC DTEIFGIFKKHYKVNFD SKKFSKKSDEEK  
ELYKIIYRYLKGRIEKILVNEQKVRLKKMEKIEIEKILNESILSEKILKRVKQY TLEHIMYLGKLRHNDID  
MTTVNTDDFSRLHAK EELDLELITFFASTNMELNKIFSRENINNDENIDFFGGDREKNYVLDKKILNSKI  
KIIRDLDFIDNKN NITNNFIRKFTKIGTNERNRILHAISKERDLQGTQDDY NKVINIIQNLKISDEEVSKAL  
NLDVVFKDKKNIITKINDIKISEENNDIKYLP SFSKVLPEILNLYRNNPKNEPFDTIETEKIVLNALIYVN  
KELYKKLILED DLEENESKNIFLQELKKT LGNIDEIDENIENYYKNAQISASKGNNAIKKYQKKVIECY  
IGYLRKNYEELFDFSDFKMNIQEIKKQIKDINDNKTYERITVKTSDKTIVINDDFEYIISIFALLNSNAVIN  
KIRNRFFATSVWLNTSEYQNIIDILDEIMQLN TLRNECITENWNLNLEEFIQKMKEIEKDFDDFKIQTKE  
IFNNYYEDIKNILTEFKDDINGCDVLEKKLEKIVIFDDETKFEIDKKSNI LQDEQRKLSNINKKDLKKKV  
DQYIKDKDQEIKSKILCRIIFNSDFLKKYKKEIDNLIEDMESENENKFQEIYYPKERKNELYTYKKNLFLNI  
GNPNFDKIYGLISNDIKMADAKFLFNIDGKNIRKNKISEIDAILKNLNDK LNGYSKEYKEYIKKLKEND  
DFFAKNIQNKNYKSFEKDYNRVSEYKKIRD LVEFN YLNKIESYLIDINWKLAIQMARFERDMHYIVNGL  
RELGIKLSGYNTGISRAYPKRNGSDGFYTTTAYYKFFDEESYKKFEKICYGFGID LSENSEINKPENESIR  
NYISHFYIVRNPFADYSIAEQIDRVSNLLSYSTRYNNSTYASVFEVFKKDVNLDYDELKKKFKLIGNNDI  
LERLMKPKKVS VLELESYNSDYIKNLIIELLTKIENTNDTL C2c3, translated from >CEPX01008730.1

marine metagenome genome assembly TARA\_037\_MES\_0.1-0.22, contig TARA\_037\_MES\_0.1-0.22\_scaffold22115\_1, whole genome shotgun sequence. (SEQ ID NO: 1059)

MRSNYHGGRNARQWRKQISGLARRTKETVFTYKFPLETDAAEIDFDKAVQTYGIAEGVGHGSLIGLVC  
AFHLSGFRLFSKAGEAMAFNRNRSRYPTDAFAEKL SAIMGIQLPTLSPEGLDLIFQSPPRS RDGIAPVWSE  
NEVRNRLYTNWTGRGPANKPDEHLLLEIAGEIAKQVFPKFGGWDDL ASDPKALAAADKYFQSQGD  
FIASLPAAIMLSPANSTVDFEGDYIAIDPAAETLLHQAVSRCAARLGRERPDLDQNKGPVSSLDALVS  
SQNNGLSWLFGVGFQHWKEKSPKELIDEYKVPADQHGAVTQVKSFVDAIPLNPLFDTTHYGEFRASVA  
GKVRSWVANYWKRLDLKSLLATTEFTLPESISDPKAVSLFSGLLVDPQGLKKVADSLPARLVSAEEAI  
DRLMGVGIPTAADIAQVERVADEIGAFIGVQVQFNNQVKQKLENLQDADDEEFLKGLKIELPSGDKEPP  
AINRISGGAPDAAAEISELEEKLQRLLDARSEHFQTISEWAEENAVTLDPIAAMVELERLRLAERGATGD  
PEEYALRLLLQRIGRLANRVSPVSAGSIRELLKPVFMEEREFNLFHNRLGSLYRSPYSTSRHQPFSDVG  
KAKAIDWIAGLDQISSDIEKALSGAGEALGDQLRDWINLAGFAISQRLRGLPDTVPNALAQVRCPPDVR  
IPLLAMLLEEDDIARDVCLKAFNLYVSAINGCLFGALREGFIVRTRFQRIGTDQIHYPKDKAW EYPDR  
LNTAKGPINAAVSSDWIEKDGAVIKPVETVRNLSSTGFAGAGVSEYLVQAPHDWYTPLDLRDVAHLVT  
GLPVEKNITKLKRLTNRTAFRMVGASSFKTHLDSVLLSDKIKLGDFTHIIDQHYRQSVTYGGKVKISYEP  
ERLQVEAAVPVVDTRDRTVPEPDTLFDHIVAIDLGERSVGFVFDIKSCLRTGEVKPIHDNNGNPVVG  
T VAVPSIRRLMKAVRSHRRRRQPQNQKVNQTYSTALQNYRENVIGDVCNRIDTLMERYNAFPVLEFQIKN  
FQAGAKQLEIVYGS

(155) Cas9 recognizes a short motif (PAM motif) in the CRISPR repeat sequences in the target DNA sequence. A “PAM motif,” or “protospacer adjacent motif,” as used herein, refers a DNA sequence immediately following the DNA sequence targeted by the Cas9 nuclease in the CRISPR bacterial adaptive immune system. PAM is a component of the invading virus or plasmid, but is not a component of the bacterial CRISPR locus. Naturally, Cas9 will not successfully bind to or cleave the target DNA sequence if it is not followed by the PAM sequence. PAM is an essential targeting component (not found in the bacterial genome) which distinguishes bacterial self from non-self DNA, thereby preventing the CRISPR locus from being targeted and destroyed by nuclease.

(156) Wild-type *Streptococcus pyogenes* Cas9 recognizes a canonical PAM sequence (5'-NGG-3'). Other Cas9 nucleases (e.g., Cas9 from *Streptococcus thermophiles*, *Staphylococcus aureus*, *Neisseria meningitidis*, or *Treponema denticola*) and Cas9 variants thereof have been described in the art to have different, or more relaxed PAM requirements. For example, in Kleinstiver et al., *Nature* 523, 481-485, 2015; Klenstiver et al., *Nature* 529, 490-495, 2016; Ran et al., *Nature*, April 9; 520(7546): 186-191, 2015; Kleinstiver et al., *Nat Biotechnol*, 33(12):1293-1298, 2015; Hou et al., *Proc Natl Acad Sci USA*, 110(39):15644-9, 2014; Prykhodzhiy et al., *PLoS One*, 10(3): e0119372, 2015; Zetsche et al., *Cell* 163, 759-771, 2015; Gao et al., *Nature Biotechnology*, doi:10.1038/nbt.3547, 2016; Want et al., *Nature* 461, 754-761, 2009; Chavez et al., doi: dx.doi.org/10.1101/058974; Fagerlund et al., *Genome Biol.* 2015; 16: 25, 2015; Zetsche et al., *Cell*, 163, 759-771, 2015; and Swarts et al., *Nat Struct Mol Biol*, 21(9):743-53, 2014, each of which is incorporated herein by reference.

(157) Thus, the guide nucleotide sequence-programmable DNA-binding protein of the present disclosure may recognize a variety of PAM sequences including, without limitation: NGG, NGAN, NGNG, NGAG, NGCG, NNGRRT, NNGRRN, NNNRRT, NNNGATT, NNAGAAW, NAAAC, TTN, TTTN, and YTN, wherein Y is a pyrimidine, and N is any nucleobase. In some embodiments, the PAM is located 5' of the target base. In some embodiments, the PAM is located 3' of the target base.

(158) One example of an RNA-programmable DNA-binding protein that has different PAM specificity is Clustered Regularly Interspaced Short Palindromic Repeats from *Prevotella* and *Francisella* 1 (Cpf1). Similar to Cas9, Cpf1 is also a class 2 CRISPR effector. It has been shown that Cpf1 mediates robust DNA interference with features distinct from Cas9. Cpf1 is a single RNA-guided endonuclease lacking tracrRNA, and it utilizes a T-rich protospacer-adjacent motif (TTN, TTTN, or YTN). Moreover, Cpf1 cleaves DNA via a staggered DNA double-stranded break. Out of 16 Cpf1-family proteins, two enzymes from *Acidaminococcus* and *Lachnospiraceae* are shown to have efficient genome-editing activity in human cells.

(159) Also useful in the present disclosure are nuclease-inactive Cpf1 (dCpf1) variants that may be used as a guide nucleotide sequence-programmable DNA-binding protein domain. The Cpf1 protein has a RuvC-like endonuclease domain that is similar to the RuvC domain of Cas9 but does not have a HNH endonuclease domain, and the N-terminal of Cpf1 does not have the alpha-helical recognition lobe of Cas9. It was shown in Zetsche et al., *Cell*, 163, 759-771, 2015 (which is incorporated herein by reference) that, the RuvC-like domain of Cpf1 is responsible for cleaving both DNA strands and inactivation of the RuvC-like domain inactivates Cpf1 nuclease activity. For example, mutations corresponding to D917A, E1006A, or D1255A in *Francisella novicida* Cpf1 (SEQ ID NO: 10) inactivates Cpf1 nuclease activity. In some embodiments, the dCpf1 of the present disclosure comprises mutations corresponding to D917A, E1006A, D1255A, D917A/E1006A, D917A/D1255A, E1006A/D1255A, or D917A/E1006A/D1255A in SEQ ID NO: 10. It is to be understood that any mutations, e.g., substitution mutations, deletions, or insertions that inactivates the RuvC domain of Cpf1 may be used in accordance with the present disclosure.

(160) Thus, in some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a nuclease inactive Cpf1 (dCpf1). In some embodiments, the dCpf1 comprises the amino acid sequence of any one SEQ ID NOs: 261-267. In some embodiments, the dCpf1 comprises an amino acid sequence that is at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at

ease 99.5% identity to SEQ ID NO: 10, and comprises mutations corresponding to D917A, E1006A, D1255A, D917A/E1006A, D917A/D1255A, E1006A/D1255A, or D917A/E1006A/D1255A in SEQ ID NO: 10. Cpf1 from other bacterial species may also be used in accordance with the present disclosure.

(161) Wild type *Francisella novicida* Cpf1 (SEQ ID NO: 10) (D917, E1006, and D1255 are bolded and underlined)

(162) TABLE-US-00017 Wild type *Francisella novicida* Cpf1 (SEQ ID NO: 10) (D917, E1006, and D1255 are bolded and underlined)

MSIYQEFVNKYSLSKTLRFELIPQGKTLENIKARGLILDDEKRAKDYKKAKQIIDKYHQFFIEEILSSVCIS  
EDLLQNYSDVYFKLKKSDDDNLQKDFKSAKDTIKKQISEYIKDSEKFKNLNQLIDAKKGQESDLILW  
LKQSKDNGIELFKANS DITDIDEALEIISFKGWTTYFKGFHENRKNVYSSNDIPTSIHYRIVDDNLPKFLE  
NKAKYESLKDKAPEAINYEQIKKDLAEELTFDIDYKTSEVNQRVFSLDEVFEIANFNYYLNQSGITKENT  
IIGGKFVNGENTKRKGINEYINLYSQQINDKTLKKYKMSVLFKQILSDTESKSFVIDKLEDDSDVVTMMQ  
SFYEQIAAFKTVEEKSIKETLSLLFDDDLKAQKLDLSKIYFKNDKSLTDLSSQVVEDDYSVIGTAVLEYITQ  
QIAPKNLDNPSKKEQELIAKKTEKAKYLSLETIKLALAEFNKHRDIDKQCRFEEILANFAAIPMIFDEIAQ  
NKDNLAQISIKYQNQGGKDLLQASAEDDVKAIKDLLDQTNLLHKLKIFHISQSEDKANILDKDEHFYL  
VFEECYFELANIVPLYNKIRNYITQKPYSDEKFKLNFENSTLANGWDKNKEPDNTAILFIKDDKYLYLV  
MNKKNKIFDDKAIKENKGEYKKIVYKLLPGANKMLPKVFFSAKSIKFYNPSEDILRIRNHSTHTKNG  
SPQKGYEKFENIEDCRKFIDFYKQSISKHPEWKDFGFRFSDTQRYNSIDEFYREVENQGYKLTFFENISES  
YIDSVVNQGGKLYLFQIYNKDFSAYSKGRPNLHTLYWKALFDERNLQDVVYKLNGEAELFYRKQSIPKK  
ITHPAKEAIAANKNDNPKKESVFEYDLIKDKRFTEDKFFHCPITINFKSSGANKENDEINLLLKEKAND  
VHILSI**DR**GERHLAYYTLVDGKGNIQKDTFNIIGNDRMKTNYHDKLAAIEKDRDSARKDWKKINNIKE  
MKEGYLSQVVHEIAKLVIEYNAIVVF**E**DLNFGFKRGRFKVEKQVYQKLEKMLIEKLNLYLVFKDNEFDK  
TGGVLRAYQLTAPFETFKKMGKQTGIYYVPAGFTSKICPVTGFVNQLYPKYESVSKSQEFFSKFDKICY  
NLDKGYFEFSFDYKNFGDKAAKGKWTIASFGSRLINFRNSDKNHNWDTREVPYPTKELEKLLKDYSIEY  
GHGECIKAACGESDCKKFFAKLTSLNLTILQMRNSKTGTLDYLISPVADVNGNFFDSRQAPKNMPQDA  
**D**ANGAYHIGLKGLMLLGRIKNNQEGKKLNLVIKNEEYFEFVQNRNN *Francisella novicida* Cpf1 D917A  
(SEQ ID NO: 261) (A917, E1006, and D1255 are bolded and underlined)

MSIYQEFVNKYSLSKTLRFELIPQGKTLENIKARGLILDDEKRAKDYKKAKQIIDKYHQFFIEEILSSVCIS  
EDLLQNYSDVYFKLKKSDDDNLQKDFKSAKDTIKKQISEYIKDSEKFKNLNQLIDAKKGQESDLILW  
LKQSKDNGIELFKANS DITDIDEALEIISFKGWTTYFKGFHENRKNVYSSNDIPTSIHYRIVDDNLPKFLE  
NKAKYESLKDKAPEAINYEQIKKDLAEELTFDIDYKTSEVNQRVFSLDEVFEIANFNYYLNQSGITKENT  
IIGGKFVNGENTKRKGINEYINLYSQQINDKTLKKYKMSVLFKQILSDTESKSFVIDKLEDDSDVVTMMQ  
SFYEQIAAFKTVEEKSIKETLSLLFDDDLKAQKLDLSKIYFKNDKSLTDLSSQVVEDDYSVIGTAVLEYITQ  
QIAPKNLDNPSKKEQELIAKKTEKAKYLSLETIKLALAEFNKHRDIDKQCRFEEILANFAAIPMIFDEIAQ  
NKDNLAQISIKYQNQGGKDLLQASAEDDVKAIKDLLDQTNLLHKLKIFHISQSEDKANILDKDEHFYL  
VFEECYFELANIVPLYNKIRNYITQKPYSDEKFKLNFENSTLANGWDKNKEPDNTAILFIKDDKYLYLV  
MNKKNKIFDDKAIKENKGEYKKIVYKLLPGANKMLPKVFFSAKSIKFYNPSEDILRIRNHSTHTKNG  
SPQKGYEKFENIEDCRKFIDFYKQSISKHPEWKDFGFRFSDTQRYNSIDEFYREVENQGYKLTFFENISES  
YIDSVVNQGGKLYLFQIYNKDFSAYSKGRPNLHTLYWKALFDERNLQDVVYKLNGEAELFYRKQSIPKK  
ITHPAKEAIAANKNDNPKKESVFEYDLIKDKRFTEDKFFHCPITINFKSSGANKENDEINLLLKEKAND  
VHILSI**AR**GERHLAYYTLVDGKGNIQKDTFNIIGNDRMKTNYHDKLAAIEKDRDSARKDWKKINNIKE  
MKEGYLSQVVHEIAKLVIEYNAIVVF**E**DLNFGFKRGRFKVEKQVYQKLEKMLIEKLNLYLVFKDNEFDK  
TGGVLRAYQLTAPFETFKKMGKQTGIYYVPAGFTSKICPVTGFVNQLYPKYESVSKSQEFFSKFDKICY  
NLDKGYFEFSFDYKNFGDKAAKGKWTIASFGSRLINFRNSDKNHNWDTREVPYPTKELEKLLKDYSIEY  
GHGECIKAACGESDCKKFFAKLTSLNLTILQMRNSKTGTLDYLISPVADVNGNFFDSRQAPKNMPQDA  
**D**ANGAYHIGLKGLMLLGRIKNNQEGKKLNLVIKNEEYFEFVQNRNN *Francisella novicida* Cpf1 E1006A  
(SEQ ID NO: 262) (D917, A1006, and D1255 are bolded and underlined)

MSIYQEFVNKYSLSKTLRFELIPQGKTLENIKARGLILDDEKRAKDYKKAKQIIDKYHQFFIEEILSSVCIS  
EDLLQNYSDVYFKLKKSDDDNLQKDFKSAKDTIKKQISEYIKDSEKFKNLNQLIDAKKGQESDLILW  
LKQSKDNGIELFKANS DITDIDEALEIISFKGWTTYFKGFHENRKNVYSSNDIPTSIHYRIVDDNLPKFLE  
NKAKYESLKDKAPEAINYEQIKKDLAEELTFDIDYKTSEVNQRVFSLDEVFEIANFNYYLNQSGITKENT  
IIGGKFVNGENTKRKGINEYINLYSQQINDKTLKKYKMSVLFKQILSDTESKSFVIDKLEDDSDVVTMMQ  
SFYEQIAAFKTVEEKSIKETLSLLFDDDLKAQKLDLSKIYFKNDKSLTDLSSQVVEDDYSVIGTAVLEYITQ  
QIAPKNLDNPSKKEQELIAKKTEKAKYLSLETIKLALAEFNKHRDIDKQCRFEEILANFAAIPMIFDEIAQ  
NKDNLAQISIKYQNQGGKDLLQASAEDDVKAIKDLLDQTNLLHKLKIFHISQSEDKANILDKDEHFYL  
VFEECYFELANIVPLYNKIRNYITQKPYSDEKFKLNFENSTLANGWDKNKEPDNTAILFIKDDKYLYLV  
MNKKNKIFDDKAIKENKGEYKKIVYKLLPGANKMLPKVFFSAKSIKFYNPSEDILRIRNHSTHTKNG  
SPQKGYEKFENIEDCRKFIDFYKQSISKHPEWKDFGFRFSDTQRYNSIDEFYREVENQGYKLTFFENISES  
YIDSVVNQGGKLYLFQIYNKDFSAYSKGRPNLHTLYWKALFDERNLQDVVYKLNGEAELFYRKQSIPKK  
ITHPAKEAIAANKNDNPKKESVFEYDLIKDKRFTEDKFFHCPITINFKSSGANKENDEINLLLKEKAND  
VHILSI**DR**GERHLAYYTLVDGKGNIQKDTFNIIGNDRMKTNYHDKLAAIEKDRDSARKDWKKINNIKE  
MKEGYLSQVVHEIAKLVIEYNAIVVF**A**DLNFGFKRGRFKVEKQVYQKLEKMLIEKLNLYLVFKDNEFDK

TGGVLRAYQLTAPFETFKKMGKQTGIYYVPAGFTSKICPVTGTFVNQLYPKYESVSKSQEFFSKFDKICY  
NLDKGYFEFSFDYKNFGDKAAKGKWTIASFGSRLINFRNSDKNHNWDTREVPYPTKELEKLLKDYSIEY  
GHGECIKAAICGESDCKKFFAKLTSLNLTILQMRNSKTGTELDYLISPVADVNGNFFDSRQAPKNMPQDA  
**D**ANGAYHIGLKGLMLLGRIKNNQEGKKLNLVIKNEEYFEFVQNRNN *Francisella novicida* Cpf1 D1255A  
(SEQ ID NO: 263) (D917, E1006, and A1255 are bolded and underlined)  
MSIYQEFVNKYSLSKTLRFELIPQGKTLENIKARGLILDDEKRAKDYKKAKQIIDKYHQFFIEILSSVCIS  
EDLLQNYSDVYFKLKKSDDDNLQKDFKSAKDTIKKQISEYIKDSEKFKNLFNQNLIDAKKGQESDLILW  
LKQSKDNGIELFKANS DITDIDEALEIISFKGWTTYFKGFHENRKNVYSSNDIPTSIHYRIVDDNLPKFLE  
NKAKYESLKDKAPEAINYEQIKKDLAEELTFDIDYKTSEVNQRVFSLDEVFEIANFNYYLNQSGITKENT  
IIGGKFVNGENTKRKGINEYINLYSQQINDKTLKKYKMSVLFKQILSDTESKSFVIDKLEDDSDVVTMMQ  
SFYEQIAAFKTVEEKSIKETLSLLFDDLKAQKLDLSKIYFKNDKSLTDLSSQVVEDDYSVIGTAVLEYITQ  
QIAPKNLDNPSKKEQELIAKKTEKAKYLSLETIKLALAEFNNKHRDIDKQCRFEEILANFAAIPMIFDEIAQ  
NKDNLAQISIKYQNQGKKDLLQASAEDDVKAIKDLLDQTNLLHKLKIFHISQSEDKANILDKDEHFYL  
VFEECYFELANIVPLYNKIRNYITQKPYSDEKFKLNFENSTLANGWDKNKEPDNTAILFIKDDKYLYLV  
MNKKNNKIFDDKAIKENKGEYKKIVYKLLPGANKMLPKVFFSAKSIKFYNPSEDILRIRNHSTHTKNG  
SPQKGYEKFENIEDCRKFIDFYKQSISKHPEWKDFGFRFSDTQRYNSIDEFYREVENQGYKLTFFENISES  
YIDSVVNQGKLYLFQIYNKDFSAYS KGRPNLHTLYWKALFDERNLQDVVYKLNGEAELFYRKQSIPKK  
ITHPAKEAIANKNDNPKKESVFEYDLIKDKRFTEDKFFHCPITINFKSSGANKENDEINLLLKEKAND  
VHIL**S**IRGERHLAYYTLVDGKGNIQKDTFNIIGNDRMKTNYHDKLAAIEKDRDSARKDWKKINNIKE  
MKEGYLSQVVHEIAKLVIEYNAIVVF**E**DLNFGFKRGRFKVEKQVYQKLEKMLIEKLNLYLVFKDNEFDK  
TGGVLRAYQLTAPFETFKKMGKQTGIYYVPAGFTSKICPVTGTFVNQLYPKYESVSKSQEFFSKFDKICY  
NLDKGYFEFSFDYKNFGDKAAKGKWTIASFGSRLINFRNSDKNHNWDTREVPYPTKELEKLLKDYSIEY  
GHGECIKAAICGESDCKKFFAKLTSLNLTILQMRNSKTGTELDYLISPVADVNGNFFDSRQAPKNMPQDA  
**A**ANGAYHIGLKGLMLLGRIKNNQEGKKLNLVIKNEEYFEFVQNRNN *Francisella novicida* Cpf1  
D917A/E1006A (SEQ ID NO: 264) (A917, A1006, and D1255 are bolded and underlined)  
MSIYQEFVNKYSLSKTLRFELIPQGKTLENIKARGLILDDEKRAKDYKKAKQIIDKYHQFFIEILSSVCIS  
EDLLQNYSDVYFKLKKSDDDNLQKDFKSAKDTIKKQISEYIKDSEKFKNLFNQNLIDAKKGQESDLILW  
LKQSKDNGIELFKANS DITDIDEALEIISFKGWTTYFKGFHENRKNVYSSNDIPTSIHYRIVDDNLPKFLE  
NKAKYESLKDKAPEAINYEQIKKDLAEELTFDIDYKTSEVNQRVFSLDEVFEIANFNYYLNQSGITKENT  
IIGGKFVNGENTKRKGINEYINLYSQQINDKTLKKYKMSVLFKQILSDTESKSFVIDKLEDDSDVVTMMQ  
SFYEQIAAFKTVEEKSIKETLSLLFDDLKAQKLDLSKIYFKNDKSLTDLSSQVVEDDYSVIGTAVLEYITQ  
QIAPKNLDNPSKKEQELIAKKTEKAKYLSLETIKLALAEFNNKHRDIDKQCRFEEILANFAAIPMIFDEIAQ  
NKDNLAQISIKYQNQGKKDLLQASAEDDVKAIKDLLDQTNLLHKLKIFHISQSEDKANILDKDEHFYL  
VFEECYFELANIVPLYNKIRNYITQKPYSDEKFKLNFENSTLANGWDKNKEPDNTAILFIKDDKYLYLV  
MNKKNNKIFDDKAIKENKGEYKKIVYKLLPGANKMLPKVFFSAKSIKFYNPSEDILRIRNHSTHTKNG  
SPQKGYEKFENIEDCRKFIDFYKQSISKHPEWKDFGFRFSDTQRYNSIDEFYREVENQGYKLTFFENISES  
YIDSVVNQGKLYLFQIYNKDFSAYS KGRPNLHTLYWKALFDERNLQDVVYKLNGEAELFYRKQSIPKK  
ITHPAKEAIANKNDNPKKESVFEYDLIKDKRFTEDKFFHCPITINFKSSGANKENDEINLLLKEKAND  
VHIL**S**IRGERHLAYYTLVDGKGNIQKDTFNIIGNDRMKTNYHDKLAAIEKDRDSARKDWKKINNIKE  
MKEGYLSQVVHEIAKLVIEYNAIVVF**A**DLNFGFKRGRFKVEKQVYQKLEKMLIEKLNLYLVFKDNEFDK  
TGGVLRAYQLTAPFETFKKMGKQTGIYYVPAGFTSKICPVTGTFVNQLYPKYESVSKSQEFFSKFDKICY  
NLDKGYFEFSFDYKNFGDKAAKGKWTIASFGSRLINFRNSDKNHNWDTREVPYPTKELEKLLKDYSIEY  
GHGECIKAAICGESDCKKFFAKLTSLNLTILQMRNSKTGTELDYLISPVADVNGNFFDSRQAPKNMPQDA  
**D**ANGAYHIGLKGLMLLGRIKNNQEGKKLNLVIKNEEYFEFVQNRNN *Francisella novicida* Cpf1  
D917A/D1255A (SEQ ID NO: 265) (A917, E1006, and A1255 are bolded and underlined)  
MSIYQEFVNKYSLSKTLRFELIPQGKTLENIKARGLILDDEKRAKDYKKAKQIIDKYHQFFIEILSSVCIS  
EDLLQNYSDVYFKLKKSDDDNLQKDFKSAKDTIKKQISEYIKDSEKFKNLFNQNLIDAKKGQESDLILW  
LKQSKDNGIELFKANS DITDIDEALEIISFKGWTTYFKGFHENRKNVYSSNDIPTSIHYRIVDDNLPKFLE  
NKAKYESLKDKAPEAINYEQIKKDLAEELTFDIDYKTSEVNQRVFSLDEVFEIANFNYYLNQSGITKENT  
IIGGKFVNGENTKRKGINEYINLYSQQINDKTLKKYKMSVLFKQILSDTESKSFVIDKLEDDSDVVTMMQ  
SFYEQIAAFKTVEEKSIKETLSLLFDDLKAQKLDLSKIYFKNDKSLTDLSSQVVEDDYSVIGTAVLEYITQ  
QIAPKNLDNPSKKEQELIAKKTEKAKYLSLETIKLALAEFNNKHRDIDKQCRFEEILANFAAIPMIFDEIAQ  
NKDNLAQISIKYQNQGKKDLLQASAEDDVKAIKDLLDQTNLLHKLKIFHISQSEDKANILDKDEHFYL  
VFEECYFELANIVPLYNKIRNYITQKPYSDEKFKLNFENSTLANGWDKNKEPDNTAILFIKDDKYLYLV  
MNKKNNKIFDDKAIKENKGEYKKIVYKLLPGANKMLPKVFFSAKSIKFYNPSEDILRIRNHSTHTKNG  
SPQKGYEKFENIEDCRKFIDFYKQSISKHPEWKDFGFRFSDTQRYNSIDEFYREVENQGYKLTFFENISES  
YIDSVVNQGKLYLFQIYNKDFSAYS KGRPNLHTLYWKALFDERNLQDVVYKLNGEAELFYRKQSIPKK  
ITHPAKEAIANKNDNPKKESVFEYDLIKDKRFTEDKFFHCPITINFKSSGANKENDEINLLLKEKAND  
VHIL**S**IRGERHLAYYTLVDGKGNIQKDTFNIIGNDRMKTNYHDKLAAIEKDRDSARKDWKKINNIKE  
MKEGYLSQVVHEIAKLVIEYNAIVVF**E**DLNFGFKRGRFKVEKQVYQKLEKMLIEKLNLYLVFKDNEFDK  
TGGVLRAYQLTAPFETFKKMGKQTGIYYVPAGFTSKICPVTGTFVNQLYPKYESVSKSQEFFSKFDKICY



NLDKGYFEFSFDYKNFGDKAAKGKWTIASFGSRLINFRNSDKNHNWDTREVPYPTKELEKLLKDYSIEY  
 GHGECIKAAICGESDCKKFFAKLTSLVNTILQMRNSKTGTELDYLISPVADVNGNFFDSRQAPKNMPQDA  
AANGAYHIGLKGLMLLGRIKNNQEGKKLNLVIKNEEYFEFVQNRNN *Francisella novicida* Cpf1  
 E1006A/D1255A (SEQ ID NO: 266) (D917, A1006, and A1255 are bolded and underlined)  
 MSIQEFVNKYSLSKTLRFELIPQGKTLENIKARGLILDDEKRAKDYKKAKQIIDKYHQFFIEILSSVCIS  
 EDLLQNYSDVYFKLKKSDDDNLQKDFKSAKDTIKKQISEYIKDSEKFKNLFNQNLIDAKKGQESDLILW  
 LKQSKDNGIELFKANSDITDIDEALEIISFKGWTTYFKGFHENRKNVYSSNDIPTSIHYRIVDDNLPKFLE  
 NKAKYESLKDKAPEAINYEQIKKDLAEELTFDIDYKTSEVNQRVFSLDEVFEIANFNNYLNQSGITKENT  
 IIGGKFVNGENTKRKGINEYINLYSQQINDKTLKKYKMSVLFKQILSDTESKSFVIDKLEDDSDVVTMMQ  
 SFYEQIAAFKTVEEKSIKETLSLLFDDDLKAQKLDLSKIYFKNDKSLTDLSSQQVFDDYSVIGTAVLEYITQ  
 QIAPKNLDNPSKKEQELIAKKTEKAKYLSLETIKLALAEFNKHRDIDKQCRFEEILANFAAIPMIFDEIAQ  
 NKDNLAQISIKYQNQGKKDLLQASAEDDVKAIKDLLDQTNNLLHKLKIFHISQSEDKANILDKDEHFYL  
 VFEECYFELANIVPLYNKIRNYITQKPYSDEKFKLNFENSTLANGWDKNKEPDNTAILFIKDDKYLYLV  
 MNKKNNKIFDDKAIKENKGEYKKIVYKLLPGANKMLPKVFFSAKSIKFYNPSEDILRIRNHSTHTKNG  
 SPQKGYEKFENIEDCRKFIDFYKQSISKHPEWKDFGFRFSDTQRYNSIDEFYREVENQGYKLTFFENISES  
 YIDSVVNQGKLYLFQIYNKDFSAYSKGRPNLHTLYWKALFDERNLQDVVYKLNGEAEFYRKQSIPKK  
 ITHPAKEAIANKNDNPKKESVFEYDLIKDKRFTEDKFFHCPITINFKSSGANKENDEINLLLKEKAND  
 VHILSI**D**RGERHLAYYTLVDGKGNIQKDTFNIIGNDRMKTNYHDKLAAIEKDRDSARKDWKKINNIKE  
 MKEGYLSQVVHEIAKLVIEYNAIVVF**A**DLNFGFKRGRFKVEKQVYQKLEKMLIEKLNLYLVFKDNEFDK  
 TGGVLRAYQLTAPFETFKKMKGKTGIIYYVPAGFTSKICPVTGTFVNQLYPKYESVSKSQEFFSKFDKICY  
 NLDKGYFEFSFDYKNFGDKAAKGKWTIASFGSRLINFRNSDKNHNWDTREVPYPTKELEKLLKDYSIEY  
 GHGECIKAAICGESDCKKFFAKLTSLVNTILQMRNSKTGTELDYLISPVADVNGNFFDSRQAPKNMPQDA  
AANGAYHIGLKGLMLLGRIKNNQEGKKLNLVIKNEEYFEFVQNRNN *Francisella novicida* Cpf1  
 D917A/E1006A/D1255A (SEQ ID NO: 267) (A917, A1006, and A1255 are bolded and  
 underlined)

MSIQEFVNKYSLSKTLRFELIPQGKTLENIKARGLILDDEKRAKDYKKAKQIIDKYHQFFIEILSSVCIS  
 EDLLQNYSDVYFKLKKSDDDNLQKDFKSAKDTIKKQISEYIKDSEKFKNLFNQNLIDAKKGQESDLILW  
 LKQSKDNGIELFKANSDITDIDEALEIISFKGWTTYFKGFHENRKNVYSSNDIPTSIHYRIVDDNLPKFLE  
 NKAKYESLKDKAPEAINYEQIKKDLAEELTFDIDYKTSEVNQRVFSLDEVFEIANFNNYLNQSGITKENT  
 IIGGKFVNGENTKRKGINEYINLYSQQINDKTLKKYKMSVLFKQILSDTESKSFVIDKLEDDSDVVTMMQ  
 SFYEQIAAFKTVEEKSIKETLSLLFDDDLKAQKLDLSKIYFKNDKSLTDLSSQQVFDDYSVIGTAVLEYITQ  
 QIAPKNLDNPSKKEQELIAKKTEKAKYLSLETIKLALAEFNKHRDIDKQCRFEEILANFAAIPMIFDEIAQ  
 NKDNLAQISIKYQNQGKKDLLQASAEDDVKAIKDLLDQTNNLLHKLKIFHISQSEDKANILDKDEHFYL  
 VFEECYFELANIVPLYNKIRNYITQKPYSDEKFKLNFENSTLANGWDKNKEPDNTAILFIKDDKYLYLV  
 MNKKNNKIFDDKAIKENKGEYKKIVYKLLPGANKMLPKVFFSAKSIKFYNPSEDILRIRNHSTHTKNG  
 SPQKGYEKFENIEDCRKFIDFYKQSISKHPEWKDFGFRFSDTQRYNSIDEFYREVENQGYKLTFFENISES  
 YIDSVVNQGKLYLFQIYNKDFSAYSKGRPNLHTLYWKALFDERNLQDVVYKLNGEAEFYRKQSIPKK  
 ITHPAKEAIANKNDNPKKESVFEYDLIKDKRFTEDKFFHCPITINFKSSGANKENDEINLLLKEKAND  
 VHILSI**A**RGERHLAYYTLVDGKGNIQKDTFNIIGNDRMKTNYHDKLAAIEKDRDSARKDWKKINNIKE  
 MKEGYLSQVVHEIAKLVIEYNAIVVF**A**DLNFGFKRGRFKVEKQVYQKLEKMLIEKLNLYLVFKDNEFDK  
 TGGVLRAYQLTAPFETFKKMKGKTGIIYYVPAGFTSKICPVTGTFVNQLYPKYESVSKSQEFFSKFDKICY  
 NLDKGYFEFSFDYKNFGDKAAKGKWTIASFGSRLINFRNSDKNHNWDTREVPYPTKELEKLLKDYSIEY  
 GHGECIKAAICGESDCKKFFAKLTSLVNTILQMRNSKTGTELDYLISPVADVNGNFFDSRQAPKNMPQDA  
AANGAYHIGLKGLMLLGRIKNNQEGKKLNLVIKNEEYFEFVQNRNN

(163) In some embodiments, the guide nucleotide sequence-programmable DNA binding protein is a Cpf1 protein from a *Acidaminococcus* species (AsCpf1). Cpf1 proteins from *Acidaminococcus* species have been described previously and would be apparent to the skilled artisan. Exemplary *Acidaminococcus* Cpf1 proteins (AsCpf1) include, without limitation, any of the AsCpf1 proteins provided herein.

(164) TABLE-US-00018 Wild-type AsCpf1 - Residue R912 is indicated in bold underlining and residues 661-667 are indicated in italics and underlining. (SEQ ID NO: 1060)

TQFEGFTNLYQVSKTLRFELIPQGKTLKHIQEQGFIEEDKARNDDHYKELKPIIDRIYKTYADQCLQLVQL  
 DWENLSAAIDSYRKEKTEETRNLALIEEQATYRNAIHDIYFIGRTDNLTDAINKRHAEIYKGLFKAELFNG  
 KVLKQLGTVTTTEHENALLRSFDKFTTYFSGFYENRKNVFSAEIDTAIPHRIVQDNFPKFKENCHIFTRL  
 ITAVPSLREHFENVKKAIGIFVSTSIEEVFSFPFYNQLLTQTQIDLYNQLLGGISREAGTEKIKGLNEVLNL  
 AIQKNDETAHIIASLPHRFIPLFKQILSDRNTLSFILEEFKSDEEVIQSFCYKTLRNENVLETAELFNL  
 NSIDLTHIFISHKKLETISSALCDHWDTLRNALYERRISELTGKITKSAKEKVQRSLKHEDINLQEIIAAG  
 KELSEAFKQKTSEILSHAAALDQPLPTTMLKKQEEKEILKSQDLSLLGLYHLLDWFVAVDESNEVDPEF  
 SARLTGIKLEMEPSLSFYNKARNYATKPYKVEKFKLNFQMPTLASGWDVNKEKNNGAILFVKNGLY  
 YLGIMPKQKGRYKALSFEPTSEKTFEGDKMYDYDFPDAKMIPKCSTQLKAVTAHFQTHHTPILLSNNF  
 IEPLEITKEIYDLNNPEKEPKFKQTAYAKKTGDQKGYREALCKWIDFTRDFLSKYTKTTSIDLSSLRPSSQ  
 YKDLGEYYAELNPLLYHISFQRIAEKEIMDAVETGKLYLFQIYNKDFAKGHHGKPNLHTLYWTGLFSPE

NLAQKNGQALFNGQPKSRMKRMAHRLGEKMLNKKLDQKTPIDTLYQELYDYVNHRLSHDLS  
DEARALLPNVITKEVSHEIHKDRRFTSDKFFFHVPITLNYQAANSPSKFNQRVNAYLKEHPETPIIGIDRGE  
RNLIYITVIDSTGKILEQRSLNTIQQFDYQKKLDNREKERVAAARQAWSVVGTIKDLKQGYLSQVIHEIVD  
LMIHYQAVVVLENLNFNGFYSKRTGIAEKAVYQQFEKMLIDKLNCLVLKDYPAEKVGGVLPYQLTDQ  
FTSFAKMGQTQSGFLFYVPAPYTSKIDPLTGFDVPFVWKTIKNHESRKHFLGEGDFLHYDVKTGDFILHFK  
MNRNLSFQRGLPGFMPAWDIVFEKNETQFDAQGTPFIAGKRIVPVIENTHRFTGRYRDLYPANELIALLE  
EKGIVFRDGSNILPKLLENDSDHAIDTMVALIRSVLQMRNSNAATGEDYINSPVRDLNGVCFDSRFQNP  
EWPMADANGAYHIALKGQLLLNHLKESKDLKLQNGISNQDWLAYIQELRN AsCpf1(R912A)- Residue  
A912 is indicated in bold underlining and residues 661- 667 are indicated in italics and  
underlining. (SEQ ID NO: 1061)

TQFEGFTNLYQVSKTLRFELIPQGKTLKHIQEQQFIEEDKARNNDHYKELKPIIDRIYKTYADQCLQLVQL  
DWENLSAAIDSYRKEKTEETRNLALIEEQATYRNAIHDFIGRTDNLTDANKRHAIEYKGLFKAELFNG  
KVLKQLGTVTTTEHENALLRSFDKFTTYFSGFYENRKNVFSADISTAIPHRIVQDNFPKFKENCHIFTRL  
ITAVPSLREHFENVKKAIGIFVSTSIEEVFSFPFYNQLLTQTQIDLYNQLLGGISREAGTEKIKGLNEVLNL  
AIQKNDETAHIIASLPHRFIPLFKQILSDRNTLSFILEEFKSDEEVIQSFCYKTLRNENVLETAELFNL  
NSIDLTHIFISHKKLETISSALCDHWDTLRNALYERRISELTGKITSAKEKVQRSLKHEDINLQEIIAAG  
KELSEAFKQKTSEILSHAAALDQPLPTTMLKKQEEKEILKSQDLSLLGLYHLLDWFVAVDESNEVDPEF  
SARLTGIKLEMEPSLSFYNKARNYATKKPYSVEKFKLNFQMPTLASGWDVNKEKNNGAILFVKNGLY  
YLGIMPKQKGRYKALSFEPTKTESEGFDKMYDYDFPDAAKMIPKCSTQLKAVTAHFQTHHTTPILLSNNF  
IEPLEITKEIYDLNNPEKEPKKFQTAYAKKTGDQKGYREALCKWIDFTRDFLSKYTKTTSIDLSSLRPSSQ  
YKDLGEYYAELNPLLYHISFQRIAEKEIMDAVETGKLYLFQIYNKDFAKGHHGKPNLHTLYWTGLFSPE  
NLAKTSIKLNGQAELFYRPKSRMKRMAHRLGEKMLNKKLDQKTPIDTLYQELYDYVNHRLSHDLS  
DEARALLPNVITKEVSHEIHKDRRFTSDKFFFHVPITLNYQAANSPSKFNQRVNAYLKEHPETPIIGIDRGE  
ANLIYITVIDSTGKILEQRSLNTIQQFDYQKKLDNREKERVAAARQAWSVVGTIKDLKQGYLSQVIHEIVD  
LMIHYQAVVVLENLNFNGFYSKRTGIAEKAVYQQFEKMLIDKLNCLVLKDYPAEKVGGVLPYQLTDQ  
FTSFAKMGQTQSGFLFYVPAPYTSKIDPLTGFDVPFVWKTIKNHESRKHFLGEGDFLHYDVKTGDFILHFK  
MNRNLSFQRGLPGFMPAWDIVFEKNETQFDAQGTPFIAGKRIVPVIENTHRFTGRYRDLYPANELIALLE  
EKGIVFRDGSNILPKLLENDSDHAIDTMVALIRSVLQMRNSNAATGEDYINSPVRDLNGVCFDSRFQNP  
EWPMADANGAYHIALKGQLLLNHLKESKDLKLQNGISNQDWLAYIQELRN

(165) In some embodiments, the nucleic acid programmable DNA binding protein is a Cpf1 protein from a  
Lachnospiraceae species (LbCpf1). Cpf1 proteins from Lachnospiraceae species have been described previously and  
would be apparent to the skilled artisan. Exemplary Lachnospiraceae Cpf1 proteins  
(LbCpf1) include, without limitation, any of the LbCpf1 proteins provided herein.

(166) TABLE-US-00019 Wild-type LbCpf1 (SEQ ID NO: 1062)

MSKLEKFTNCYSLSKTLRFKAIPVGKTQENIDNKRLLEVEDEKRAEDYKGVKKLLDRYYLSFINDVLHSI  
KLKNLNNYISLFRKKTRTEKENKELENLEINLRKEIAKAFKGNEGYKSLFKKDIETILPEFLDDKDEIAL  
VNSFNGFTTAFTGFFDNRENMFSEEAKSTSIAFRCINENLTRYISNMDIFEKVDAIFDKHEVQEIKEILN  
SDYDVEDFFEGEFFNFVLTQEGIDVYNAIIGGFVTESGEKIKGLNEYINLYNQKTKQKLPKFKPLYKQVL  
SDRESLSFYGEGYTSDEEVLEVFRNTLNKNSEIFFSIKKLEKLFKNFDEYSSAGIFVKNNGPAISTISKDIFG  
EWNVIRDKWNAEYDDIHLKKKAVVTEKYEDDRRSFKKIGSFSLEQLQEYADADLSVVEKLKEIIIQK  
VDEIYKVYGSSEKLFDAADFVLEKSLKKNDVAVVIMKDLLDSVKSFENYIKAFFGEGKETNRDESFYGD  
FVLAYDILLKVDHIYDAIRNYVTQKPYSKDKFKLYFQNPQFMGGWDKDKETDYRATILRYGSKYYLAI  
MDKKYAKCLQKIDKDDVNGNYEKINYKLLPGPNKMLPKVFFSKKWMAYYNPSEDIQKIYKNGTFKK  
GDMFNLNDCHKLIDFFKDSISRPKWSNAYDFNFSETEKYKDIAGFYREVEEQGYKVSFESASKKEVD  
KLVEEGKLYMFQIYNKDFSDKSHGTPNLHTMYFKLLFDENNHGQIRLSGGAELFMRRASLKKEELVVH  
PANSPIANKNPDNPKKTTTSLSYDVYKDKRFSEDQYELHIPIAINKCPKNIFKINTEVRVLLKHDDNPYVI  
GIDRGERNLLYIVVVDGKGNIVEQYSLNEIINNENGIRIKTDYHSLLDKKEKERFEARQNWTSIENIKELK  
AGYISQVVHKICELVEKYDAVIALEDLNSGFKNSRVKVEKQVYQKFEKMLIDKLNMYMVDKKSNPCAT  
GGALKGYQITNKFESFKSMSTQNGFIFYIPAWLTSKIDPSTGFVNLLKTKYTSIADSKKFISFDRIMYVP  
EEDLFEFALDYKNFSRTDADYIKWKLYSYGNRIRIFRNPKNVFDWEEVCLTSAYKELFNKYGINY  
QQGDIRALLCEQSDKAFYSSFMALMSLMLQMRNSITGRTDVDFLISPVKNSDGIFYDSRNYEAQENAIL  
PKNADANGAYNIARKVLWAIGQFKKAEDEKLDKVKIAISNKEWLEYAQTSVKH LbCpf1 (R836A) (SEQ ID  
NO: 1063)

MSKLEKFTNCYSLSKTLRFKAIPVGKTQENIDNKRLLEVEDEKRAEDYKGVKKLLDRYYLSFINDVLHSI  
KLKNLNNYISLFRKKTRTEKENKELENLEINLRKEIAKAFKGNEGYKSLFKKDIETILPEFLDDKDEIAL  
VNSFNGFTTAFTGFFDNRENMFSEEAKSTSIAFRCINENLTRYISNMDIFEKVDAIFDKHEVQEIKEILN  
SDYDVEDFFEGEFFNFVLTQEGIDVYNAIIGGFVTESGEKIKGLNEYINLYNQKTKQKLPKFKPLYKQVL  
SDRESLSFYGEGYTSDEEVLEVFRNTLNKNSEIFFSIKKLEKLFKNFDEYSSAGIFVKNNGPAISTISKDIFG  
EWNVIRDKWNAEYDDIHLKKKAVVTEKYEDDRRSFKKIGSFSLEQLQEYADADLSVVEKLKEIIIQK  
VDEIYKVYGSSEKLFDAADFVLEKSLKKNDVAVVIMKDLLDSVKSFENYIKAFFGEGKETNRDESFYGD  
FVLAYDILLKVDHIYDAIRNYVTQKPYSKDKFKLYFQNPQFMGGWDKDKETDYRATILRYGSKYYLAI

MDKKYAKCLQKIDKDDVNGNYEKINYKLLPGPNKMLPKVFFSKKWMAYYNPSEDIQKIYKNGTFFK  
GDMFNLNDCHKLIDFFKDSISRYPKWSNAYDFNFSETEKYKDIAGFYREVEEQGYKVSFESASKKEVD  
KLVEEGKLYMFQIYNKDFSDKSHGTPNLHTMYFKLLFDENNHGQIRLSGGAELFMRRASLKKEELVVH  
PANSPIANKNPDNPKKTTTSLSYDVYKDKRFSEDQYELHIPIAINKCPKNIFKINTEVRVLLKHDDNPYVI  
GIDRGEANLLYIVVVDGKGNIVEQYSLNEIINNFNNGIRIKTDYHSLLDKKEKERFEARQNWTSIENIKEL  
KAGYISQVVHKICELVEKYDAVIALEDLNSGFKNSRVKVEKQVYQKFEKMLIDKLNMYMDKKSNPCA  
TGGALKGYQITNKFESFKSMSTQNGFIFYIPAWLTSKIDPSTGFVNLLKTKYTSIADSKKFISFDRIMYV  
PEEDLFEFALDYKNFSRTDADYIKKWKLYSYGNRIRIFRNPKKNNVFDWEEVCLTSAYKELFNKYGIN  
YQQGDIRALLCEQSDKAFYSSFMALMSLMLQMRNSITGRTDVDFLISPVKNSDGIFYDSRNYEAQENAI  
LPKNADANGAYNIARKVLWAIGQFKKAEDEKLDKVKIAISNKEWLEYAQTSVKH LbCpf1 (R1138A) (SEQ  
ID NO: 1064)

MSKLEKFTNCYSLSKTLRFKAIPVGKTQENIDNKRLLEVEDEKRAEDYKGVKKLLDRYYLSFINDVLHSI  
KLKNLNNYISLFRKKTRTEKENKELENLEINLRKEIAKAFKGNEGYKSLFKKDIIETILPEFLDDKDEIAL  
VNSFNGFTTAFTGFFDNRENMFSEEAKSTSIAFRCINENLTRYISNMDIFEKVD AIFDKHEVQEIKEKILN  
SDYDVEDFFEGEFFNFVLTQEGIDVYNAIIGGFVTESGEKIKGLNEYINLYNQKTKQKLPKFKPLYKQVL  
SDRESLSFYGEGYTSDEEVLEVFRTNLNKNSEIFSSIKKLEKLFKNFDEYSSAGIFVKNGPAISTISKDIFG  
EWNVIRDKWNAEYDDIHLKKKAVVTEKYEDDRRSFKKIGSFSLEQLQEYADADLSVVEKLKEIIIQK  
VDEIYKVYGSSEKLFDAADFVLEKSLKKNDAVVAIMKDLLDSVKSFENYIKAFFGEGKETNRDESFYGD  
FVLAYDILLKVDHIYDAIRNYVTQKPYSKDKFKLYFQNPQFMGGWDKDKETDYRATILRYGSKYYLAI  
MDKKYAKCLQKIDKDDVNGNYEKINYKLLPGPNKMLPKVFFSKKWMAYYNPSEDIQKIYKNGTFFK  
GDMFNLNDCHKLIDFFKDSISRYPKWSNAYDFNFSETEKYKDIAGFYREVEEQGYKVSFESASKKEVD  
KLVEEGKLYMFQIYNKDFSDKSHGTPNLHTMYFKLLFDENNHGQIRLSGGAELFMRRASLKKEELVVH  
PANSPIANKNPDNPKKTTTSLSYDVYKDKRFSEDQYELHIPIAINKCPKNIFKINTEVRVLLKHDDNPYVI  
GIDRGERNLLYIVVVDGKGNIVEQYSLNEIINNFNNGIRIKTDYHSLLDKKEKERFEARQNWTSIENIKELK  
AGYISQVVHKICELVEKYDAVIALEDLNSGFKNSRVKVEKQVYQKFEKMLIDKLNMYMDKKSNPCAT  
GGALKGYQITNKFESFKSMSTQNGFIFYIPAWLTSKIDPSTGFVNLLKTKYTSIADSKKFISFDRIMYVP  
EEDLFEFALDYKNFSRTDADYIKKWKLYSYGNRIRIFRNPKKNNVFDWEEVCLTSAYKELFNKYGINY  
QQGDIRALLCEQSDKAFYSSFMALMSLMLQMANSITGRTDVDFLISPVKNSDGIFYDSRNYEAQENAIL  
PKNADANGAYNIARKVLWAIGQFKKAEDEKLDKVKIAISNKEWLEYAQTSVKH

(167) In some embodiments, the Cpf1 protein is a crippled Cpf1 protein. As used herein a “crippled Cpf1” protein is a Cpf1 protein having diminished nuclease activity as compared to a wild-type Cpf1 protein. In some embodiments, the crippled Cpf1 protein preferentially cuts the target strand more efficiently than the non-target strand. For example, the Cpf1 protein preferentially cuts the strand of a duplexed nucleic acid molecule in which a nucleotide to be edited resides. In some embodiments, the crippled Cpf1 protein preferentially cuts the non-target strand more efficiently than the target strand. For example, the Cpf1 protein preferentially cuts the strand of a duplexed nucleic acid molecule in which a nucleotide to be edited does not reside. In some embodiments, the crippled Cpf1 protein preferentially cuts the target strand at least 5% more efficiently than it cuts the non-target strand. In some embodiments, the crippled Cpf1 protein preferentially cuts the target strand at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 50%, 60%, 70%, 80%, 90%, or at least 100% more efficiently than it cuts the non-target strand.

(168) In some embodiments, a crippled Cpf1 protein is a non-naturally occurring Cpf1 protein. In some embodiments, the crippled Cpf1 protein comprises one or more mutations relative to a wild-type Cpf1 protein. In some embodiments, the crippled Cpf1 protein comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 mutations relative to a wild-type Cpf1 protein. In some embodiments, the crippled Cpf1 protein comprises an R836A mutation mutation as set forth in SEQ ID NO: 763, or in a corresponding amino acid in another Cpf1 protein. It should be appreciated that a Cpf1 comprising a homologous residue (e.g., a corresponding amino acid) to R836A of SEQ ID NO: 763 could also be mutated to achieve similar results. In some embodiments, the crippled Cpf1 protein comprises a R1138A mutation as set forth in SEQ ID NO: 763, or in a corresponding amino acid in another Cpf1 protein. In some embodiments, the crippled Cpf1 protein comprises an R912A mutation mutation as set forth in SEQ ID NO: 762, or in a corresponding amino acid in another Cpf1 protein. Without wishing to be bound by any particular theory, residue R838 of SEQ ID NO: 763 (LbCpf1) and residue R912 of SEQ ID NO: 762 (AsCpf1) are examples of corresponding (e.g., homologous) residues. For example, a portion of the alignment between SEQ ID NO: 762 and 763 shows that R912 and R838 are corresponding residues.

(169) In some embodiments, any of the Cpf1 proteins provided herein comprises one or more amino acid deletions. In some embodiments, any of the Cpf1 proteins provided herein comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acid deletions. Without wishing to be bound by any particular theory, there is a helical region in Cpf1, which includes residues 661-667 of AsCpf1 (SEQ ID NO: 762), that may obstruct the function of a deaminase (e.g., APOBEC) that is fused to the Cpf1. This region comprises the amino acid sequence KKTGDQK. Accordingly, aspects of the disclosure provide Cpf1 proteins comprising mutations (e.g., deletions) that disrupt this helical region in Cpf1. In some embodiments, the Cpf1 protein comprises one or more deletions of the following residues in SEQ ID NO: 762, or one or more corresponding deletions in another Cpf1 protein: K661, K662, T663, G664, D665, Q666, and K667. In some embodiments, the Cpf1 protein comprises a T663 and a D665 deletion in SEQ ID NO: 762, or corresponding deletions in another Cpf1 protein. In some embodiments, the Cpf1 protein comprises a K662, T663, D665, and Q666 deletion in SEQ

ID NO: 762, or corresponding deletions in another Cpf1 protein. In some embodiments, the Cpf1 protein comprises a K661, K662, T663, D665, Q666 and K667 deletion in SEQ ID NO: 762, or corresponding deletions in another Cpf1 protein.

(170) TABLE-US-00020 AsCpf1 (deleted T663 and D665) (SEQ ID NO: 1065)

TQFEGFTNLYQVSKTLRFELIPQGKTLKHIQEQGFIEEDKARNDHYKELKPIIDRIYKTYADQCLQLVQL  
DWENLSAAIDSYRKEKTEETRNLALIEEQATYRNAIHQDYFIGRTDNLTDANKRHAIEYKGLFKAELFNG  
KVLKQLGTVTTTEHENALLRSFDKFTTYFSGFYENRKNVFSAEISTAIPIHRIVQDNFPKFKENCHIFTRL  
ITAVPSLREHFENVKKAIGIFVSTSIEEVFSFPFYNQLLTQTQIDLYNQLLGGISREAGTEKIKGLNEVLNL  
AIQKNDETAHIIASLPHRFIPLFKQILSDRNTLSFILEEFKSDEEVIQSFCYKTLRNENVLETAELFNL  
NSIDLTHIFISHKKLETISSALCDHWDTLRNALYERRISELTGKITKSAKEKVQRSLKHEDINLQEIISAAG  
KELSEAFKQKTSEILSHAHAAALDQPLPTTMLKKQEEKEILKSQLDSLGLYHLLDWFVAVDESNEVDPEF  
SARLTGIKLEMEPSLSFYNKARNYATKKPYSVEKFKNLFQMPTLASGWDVNKEKNNGAILFVKNGLY  
YLGIMPKQKGRYKALSFEPTTEKTSEGFDKMYDYFPDAAKMIPKCSTQLKAVTAHFQTHHTPILLSNNF  
IEPLEITKEIYDLNNPEKEPKKFQYAYAKKGQKGYREALCKWIDFTRDFLSKYTKTTSIDLSSLRPSSQY  
KDLGEYYAELNPLLYHISFQRIAEKEIMDAVETGKLYLFQIYNKDFAKGHHGKPNLHTLYWTGLFSPEN  
LAKTSIKLNGQAELFYRPSRMKRMARHLGEKMLNKKLKDQKTPIPDTLYQELYDYVNHRLSHDLSD  
EARALLPNVITKEVSHEIHKDRRFTSDKFFHVPITLNYQAANSPSKFNQRVNAYLKEHPETPIIGIDRGER  
NLIYITVIDSTGKILEQRSLNTIQQFDYQKKLDNREKERVAAARQAWSVVGTIKDLKQGYLSQVIHEIVDL  
MIHYQAVVVLENLNFQFKSKRTGIAEKAVYQQFEKMLIDKLNCLVLKDYPAEKVGGVLNPNYQLTDQF  
TSFAKMGTQSGFLFYVPAPYTSKIDPLTGFVDPFVWKTIKNHESRKHFLLEGFDLHYDVKTGDFILHFK  
MNRNLSFQRGLPGFMPAWDIVFEKNETQFDAQGTPFIAGKRIVPVIENTHRFTGRYRDLYPANELIALLE  
EKGIVFRDGSNILPKLLENDSDHAIDTMVALIRSVLQMRNSNAATGEDYINSPVRDLNGVCFDSRFQNP  
EWPMDADANGAYHIALKGQLLLNHLKESKDLKLQNGISNQDWLAYIQELRN AsCpf1 (deleted K662,  
T663, D665, and Q666) (SEQ ID NO: 1066)

TQFEGFTNLYQVSKTLRFELIPQGKTLKHIQEQGFIEEDKARNDHYKELKPIIDRIYKTYADQCLQLVQL  
DWENLSAAIDSYRKEKTEETRNLALIEEQATYRNAIHQDYFIGRTDNLTDANKRHAIEYKGLFKAELFNG  
KVLKQLGTVTTTEHENALLRSFDKFTTYFSGFYENRKNVFSAEISTAIPIHRIVQDNFPKFKENCHIFTRL  
ITAVPSLREHFENVKKAIGIFVSTSIEEVFSFPFYNQLLTQTQIDLYNQLLGGISREAGTEKIKGLNEVLNL  
AIQKNDETAHIIASLPHRFIPLFKQILSDRNTLSFILEEFKSDEEVIQSFCYKTLRNENVLETAELFNL  
NSIDLTHIFISHKKLETISSALCDHWDTLRNALYERRISELTGKITKSAKEKVQRSLKHEDINLQEIISAAG  
KELSEAFKQKTSEILSHAHAAALDQPLPTTMLKKQEEKEILKSQLDSLGLYHLLDWFVAVDESNEVDPEF  
SARLTGIKLEMEPSLSFYNKARNYATKKPYSVEKFKNLFQMPTLASGWDVNKEKNNGAILFVKNGLY  
YLGIMPKQKGRYKALSFEPTTEKTSEGFDKMYDYFPDAAKMIPKCSTQLKAVTAHFQTHHTPILLSNNF  
IEPLEITKEIYDLNNPEKEPKKFQYAYAKKGQKGYREALCKWIDFTRDFLSKYTKTTSIDLSSLRPSSQYKDL  
GEYYAELNPLLYHISFQRIAEKEIMDAVETGKLYLFQIYNKDFAKGHHGKPNLHTLYWTGLFSPENLAK  
TSIKLNGQAELFYRPSRMKRMARHLGEKMLNKKLKDQKTPIPDTLYQELYDYVNHRLSHDLSD  
EARALLPNVITKEVSHEIHKDRRFTSDKFFHVPITLNYQAANSPSKFNQRVNAYLKEHPETPIIGIDRGER  
NLIYITVIDSTGKILEQRSLNTIQQFDYQKKLDNREKERVAAARQAWSVVGTIKDLKQGYLSQVIHEIVDLMI  
HYQAVVVLENLNFQFKSKRTGIAEKAVYQQFEKMLIDKLNCLVLKDYPAEKVGGVLNPNYQLTDQFTS  
FAKMGTQSGFLFYVPAPYTSKIDPLTGFVDPFVWKTIKNHESRKHFLLEGFDLHYDVKTGDFILHFKMN  
RNLSFQRGLPGFMPAWDIVFEKNETQFDAQGTPFIAGKRIVPVIENTHRFTGRYRDLYPANELIALLEEK  
GIVFRDGSNILPKLLENDSDHAIDTMVALIRSVLQMRNSNAATGEDYINSPVRDLNGVCFDSRFQNP  
EWPMDADANGAYHIALKGQLLLNHLKESKDLKLQNGISNQDWLAYIQELRN AsCpf1 (deleted K661, K662,  
T663, D665, Q666, and K667) (SEQ ID NO: 1067)

TQFEGFTNLYQVSKTLRFELIPQGKTLKHIQEQGFIEEDKARNDHYKELKPIIDRIYKTYADQCLQLVQL  
DWENLSAAIDSYRKEKTEETRNLALIEEQATYRNAIHQDYFIGRTDNLTDANKRHAIEYKGLFKAELFNG  
KVLKQLGTVTTTEHENALLRSFDKFTTYFSGFYENRKNVFSAEISTAIPIHRIVQDNFPKFKENCHIFTRL  
ITAVPSLREHFENVKKAIGIFVSTSIEEVFSFPFYNQLLTQTQIDLYNQLLGGISREAGTEKIKGLNEVLNL  
AIQKNDETAHIIASLPHRFIPLFKQILSDRNTLSFILEEFKSDEEVIQSFCYKTLRNENVLETAELFNL  
NSIDLTHIFISHKKLETISSALCDHWDTLRNALYERRISELTGKITKSAKEKVQRSLKHEDINLQEIISAAG  
KELSEAFKQKTSEILSHAHAAALDQPLPTTMLKKQEEKEILKSQLDSLGLYHLLDWFVAVDESNEVDPEF  
SARLTGIKLEMEPSLSFYNKARNYATKKPYSVEKFKNLFQMPTLASGWDVNKEKNNGAILFVKNGLY  
YLGIMPKQKGRYKALSFEPTTEKTSEGFDKMYDYFPDAAKMIPKCSTQLKAVTAHFQTHHTPILLSNNF  
IEPLEITKEIYDLNNPEKEPKKFQYAYAGGYREALCKWIDFTRDFLSKYTKTTSIDLSSLRPSSQYKDLGE  
YYAELNPLLYHISFQRIAEKEIMDAVETGKLYLFQIYNKDFAKGHHGKPNLHTLYWTGLFSPENLAKTS  
IKLNGQAELFYRPSRMKRMARHLGEKMLNKKLKDQKTPIPDTLYQELYDYVNHRLSHDLSD  
EARALLPNVITKEVSHEIHKDRRFTSDKFFHVPITLNYQAANSPSKFNQRVNAYLKEHPETPIIGIDRGER  
NLIYITVIDSTGKILEQRSLNTIQQFDYQKKLDNREKERVAAARQAWSVVGTIKDLKQGYLSQVIHEIVDLMIHYQ  
AVVVLENLNFQFKSKRTGIAEKAVYQQFEKMLIDKLNCLVLKDYPAEKVGGVLNPNYQLTDQFTSFAK  
MGTQSGFLFYVPAPYTSKIDPLTGFVDPFVWKTIKNHESRKHFLLEGFDLHYDVKTGDFILHFKMNRN  
LSFQRGLPGFMPAWDIVFEKNETQFDAQGTPFIAGKRIVPVIENTHRFTGRYRDLYPANELIALLEEKGIVF

RDGSLNPKLLMDSHAIDTMVALIRSVLQMRNSNAATGEDYINSPVRDLNGVCFDSRFQNPPEWPMADANGAYHIALKGQLLNHLKESKDLKLQNGISNQDWLAYIQELRN

(171) In some embodiments, the guide nucleotide sequence-programmable DNA-binding protein domain of the present disclosure has no requirements for a PAM sequence. One example of such guide nucleotide sequence-programmable DNA-binding protein may be an Argonaute protein from *Natronobacterium gregoryi* (NgAgo). NgAgo is a ssDNA-guided endonuclease. NgAgo binds 5' phosphorylated ssDNA of ~24 nucleotides (gDNA) to guide it to its target site and will make DNA double-strand breaks at gDNA site. In contrast to Cas9, the NgAgo-gDNA system does not require a protospacer-adjacent motif (PAM). Using a nuclease inactive NgAgo (dNgAgo) can greatly expand the codons that may be targeted. The characterization and use of NgAgo have been described in Gao et al., *Nat Biotechnol.* Epub 2016 May 2. PubMed PMID: 27136078; Swarts et al., *Nature.* 507(7491) (2014):258-61; and Swarts et al., *Nucleic Acids Res.* 43(10) (2015):5120-9, each of which are incorporated herein by reference. The sequence of *Natronobacterium gregoryi* Argonaute is provided in SEQ ID NO: 270.

(172) TABLE-US-00021 Wild type *Natronobacterium gregoryi* Argonaute (SEQ ID NO: 270)

MTVIDLDSTTTADELTSGHTYDISVTLTGVYDNTDEQHPRMSLAFEQDN  
GERRYITLWKNTTPKDVFTYDYATGSTYIFTNIDYEVKDG YENLTATYQ  
TTVENATAQEVGTTDEDETFAAGGEPLDHHLDDALNETPDDAETESDSGH  
VMTSFASRDQLPEWTLHTYTLTATDGAKTDTEYARRTLAYTVRQELYTD  
HDAAPVATDGLMLLTPEPLGETPLDLDCGVRVEADETRTLDYTTAKDRL  
LARELVEEGLKRSLWDDYLVRGIDEVLSKEPVLTCEFDLHERYDLSVE  
VGHSGRAYLHINFRHRFVPKLTLADIDDDNIYPGLRVKTTYRPRRGHIV  
WGLRDECATDSLNTLGNQSVVAYHRNNQTPINTDLLDAIEAADRRVVET  
RRQGHGDDAVSFPQELLAVEPNTHQIKQFASDGFHQQARSKTRLSASRC  
SEKAQFAERLDPVRLNGSTVEFSSEFFTGNNQQRLRLLYENGESVLT  
RDGARGAHPDETFSKGINPPESFEVAVVLPEQQADTCKAQWDTMADLL  
NQAGAPPTRSETVQYDAFSSPESISLNVAGAI DPSEVDAA FVVLPPDQE  
GFADLASPTETYDELKKALANMGIYSQMAYFDRFRDAKIFYTRNVALGL  
LAAAGGVAFTTEHAMPGDADMFIGIDVSRSYPEDGASGQINIAATATAV  
YKDGITILGHSSTRPQLGEKLQSTDVRDIMKNAILGYQQVTGESPTHIVI  
HRDGMNEDLDPATEFLNEQGVEYDIVEIRKQPQTRLLAVSDVQYDTPV  
KSIAAINQNEPRATVATFGAPEYLATRDGGGLPRPIQIERVAGETDIET  
LTRQVYLLSQSHIQVHNSTARLPITTAYADQASTHATKGYLVQTGAFES NVGFL

(173) Also provided herein are Cas9 variants that have relaxed PAM requirements (PAMless Cas9). PAMless Cas9 exhibits an increased activity on a target sequence that does not include a canonical PAM (e.g., NGG) at its 3'-end as compared to *Streptococcus pyogenes* Cas9 as provided by SEQ ID NO: 1, e.g., increased activity by at least 5-fold, at least 10-fold, at least 50-fold, at least 100-fold, at least 500-fold, at least 1,000-fold, at least 5,000-fold, at least 10,000-fold, at least 50,000-fold, at least 100,000-fold, at least 500,000-fold, or at least 1,000,000-fold. Such Cas9 variants that have relaxed PAM requirements are described in U.S. Provisional applications 62/245,828, 62/279,346, 62/311,763, 62/322,178, and 62/357,332, each of which is incorporated herein by reference. In some embodiments, the dCas9 or Cas9 nickase useful in the present disclosure may further comprise mutations that relax the PAM requirements, e.g., mutations that correspond to A262T, K294R, S409I, E480K, E543D, M694I, or E1219V in SEQ ID NO: 1.

(174) Other on-limiting, exemplary Cas9 variants (including dCas9, Cas9 nickase, and Cas9 variants with alternative PAM requirements) suitable for use in the nucleobase editors described herein and their respective sequence are provided below.

(175) TABLE-US-00022 VRER-nCas9 (D10A/D1135V/G1218R/R1335E/T1337R) *S. pyogenes* Cas9 Nickase (SEQ ID NO: 821)

MDKKYSIGLAIGTNSVGWAVITDEYKVPSSKKFKVLGNTDRHSIKKNLIGALLFDSGETAEATRLKRTAR  
RRYTRRKNRICYLQEIFSNEMAKVDDSFHRL EESFLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLR  
KKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDA  
KAILSARLSKSRLENLIAQLPGEKKNGLFGNLIALSLGLTPNFKSNFDLAEDAKLQLSKDTYDDDLDN  
LLAQIGDQYADFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLKALVRQQLPEK  
YKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDNGSIPHQIHL  
GELHAILRRQEDFYPLKDNREKIEKILTFRIPIYYVGPLARGNSRFAWMTRKSEETITPWNFEEVVDKGA  
SAQSFIERMNTNFDKNLPNEKVLPKHSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFK  
TNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEENEDILEDIVLTLT  
LFEDREMIEERLKTYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRN  
FMQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIKKGILQTVKVVDLKVVMGRHKPENIVI  
EMARENQTTQKGQKNSRERMKRIE EGIKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQEL  
DINRLSDYDVDHIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKMKMKNYWRQLLNAKLITQRK  
FDNLTKAERGGSELDKAGFIKRQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSD  
FRKDFQFYKVBREINNYHHAHDAYLNAVVG TALIKKYPKLESEFVYGDYKVYDVVRKMIKSEQEIGKA  
TAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVL SMPQVNIVKKTEVQ

TGGFSKESILPKRNSDKLIARKKDWDPKKYGGFVSPTVAYSVLVAKVEKGSKKLKSVKELLGITIME  
RSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELENGRKRMLASARELQKGNELALPSKYVNFLYLA  
SHYEKLKGSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVLADANLDKVLSAYNKHRRDKPIREQAENI  
IHLFTLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDATLIHQSIITGLYETRIDLSQLGGD (single underline: HNH  
domain; double underline: RuvC domain) VQR-nCas9 (D10A/D1135V/R1335Q/T1337R) *S. pyogenes*  
Cas9 Nickase (SEQ ID NO: 822)

MDKKYISGLAIGTNSVGWAVITDEYKVPSKKFKVLGNTDRHSIKKNLIGALLEDSGETAEATRLKRTAR  
RRYTRRKNRICYLQEIFSNEMAKVDDSFHRLSEESFLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLR  
KKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDA  
KAILSARLSKSRRLLENLIAQLPGEKKNGLFGNLIALSLGLTPNFKSNFDLAEDAKLQLSKDTYDDDLDN  
LLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLKALVRQQLPEK  
YKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDNGSIPHQIHL  
GELHAILRRQEDFYPLKDNREKIEKILTRIPYYVGPLARGNSRFAWMTRKSEETITPWNFEEVVDKGA  
SAQSFIERMTNFDKNLPNEKVLPKHSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFK  
TNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEENEDILEDIVLTLT  
LFEDREMIEERLKTIAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRN  
FMQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIKKGILQTVKVVDDELVKVMGRHKPENIVI  
EMARENQTTQKGQKNSRERMKRIEIEGKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQEL  
DINRLSDYDVDHIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKMKMKNYWRQLLNAKLITQRK  
FDNLTKAERGGLSELDKAGFIKRQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSD  
FRKDFQFYKVBREINNYHHAHDAYLNAVVGTAIIKKYPKLESEFVYGDYKVYDVRKMIAKSEQEIGKA  
TAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVLSPQVNVKKTEVQ  
TGGFSKESILPKRNSDKLIARKKDWDPKKYGGFVSPTVAYSVLVAKVEKGSKKLKSVKELLGITIME  
RSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLA  
SHYEKLKGSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVLADANLDKVLSAYNKHRRDKPIREQAENI  
IHLFTLTNLGAPAAFKYFDTTIDRKQYRSTKEVLDATLIHQSIITGLYETRIDLSQLGGD (single underline:  
HNH domain; double underline: RuvC domain) EQR-nCas9 (D10A/D1135E/R1335Q/T1337R) *S.*  
*pyogenes* Cas9 Nickase (SEQ ID NO: 823)

MDKKYISGLAIGTNSVGWAVITDEYKVPSKKFKVLGNTDRHSIKKNLIGALLEDSGETAEATRLKRTAR  
RRYTRRKNRICYLQEIFSNEMAKVDDSFHRLSEESFLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLR  
KKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDA  
KAILSARLSKSRRLLENLIAQLPGEKKNGLFGNLIALSLGLTPNFKSNFDLAEDAKLQLSKDTYDDDLDN  
LLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLKALVRQQLPEK  
YKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDNGSIPHQIHL  
GELHAILRRQEDFYPLKDNREKIEKILTRIPYYVGPLARGNSRFAWMTRKSEETITPWNFEEVVDKGA  
SAQSFIERMTNFDKNLPNEKVLPKHSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFK  
TNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEENEDILEDIVLTLT  
LFEDREMIEERLKTIAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRN  
FMQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIKKGILQTVKVVDDELVKVMGRHKPENIVI  
EMARENQTTQKGQKNSRERMKRIEIEGKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQEL  
DINRLSDYDVDHIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKMKMKNYWRQLLNAKLITQRK  
FDNLTKAERGGLSELDKAGFIKRQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSD  
FRKDFQFYKVBREINNYHHAHDAYLNAVVGTAIIKKYPKLESEFVYGDYKVYDVRKMIAKSEQEIGKA  
TAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVLSPQVNVKKTEVQ  
TGGFSKESILPKRNSDKLIARKKDWDPKKYGGFVSPTVAYSVLVAKVEKGSKKLKSVKELLGITIME  
RSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLA  
SHYEKLKGSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVLADANLDKVLSAYNKHRRDKPIREQAENI  
IHLFTLTNLGAPAAFKYFDTTIDRKQYRSTKEVLDATLIHQSIITGLYETRIDLSQLGGD (single underline:  
HNH domain; double underline: RuvC domain) KKH-nCas9 (D10A/E782K/N968K/R1015H) *S. aureus*  
Cas9 Nickase (SEQ ID NO: 268)

MKRNYILGLAIGITSVGYGIIDYETRDVIDAGVRLFKEANVENNEGRRSKRGARRLKRRRRHRIQRVKK  
LLFDYNLLTDHSELGINPYEARVKGLSQKLSEEFSAALLHLAKRRGVHNVNEVEEDTGNELSTKEQI  
SRNSKALEEKYVAELQLERLKKDGEVRGSINRFKTSDYVKEAKQLLKVQKAYHQLDQSFIDTYIDLL  
TRRTYYEGPGEGSPFGWKDIKEWYEMLMGHCTYFPEELRSVKYAYNADLYNALNDLNNLVITRDENE  
KLEYEYKFQIENVFKQKKKPTLKQIAKEILVNEEDIKGYRVTSTGKPEFTNLKVYHDIKDITARKEIEN  
AELLDQIAKILTIYQSSEDIQEELTNLNSLTQEEIEQISNLKGYTGTHNLSLKAINLILDELWHTNDNQIA  
IFNRLKLVPKKVDLSQQKEIPTTLVDDFILSPVVKRSFIQSIKVINAIKKYGLPNDIIELAREKNSKDAQK  
MINEMQKRNRQTNERIEEIIRTTGKENAKYLIEKIKLHDMQEGKCLYSLEAIPLEDLLNPNFNYEVDHIIP  
RSVSFDNSFNKNVLVKQEENSCKGNRTPFQYLSSSDSKISYETFKKHILNLAAGKGRISKTKKEYLLEER  
DINRFSVQKDFINRLVDTRYATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFCKERNKGY  
KHHAEDALIIANADFIFKEWKLDKAKKVMENQMFEKQAESMPEIETEQEYKEIFITPHQIKHIKDFK



acid sequence set forth in any of SEQ ID NOs: 271-292, 303, and 1072-1083, or an amino acid sequence homologous to a fragment of the amino acid sequence set forth in any of SEQ ID NOs: 271-292, 303, and 1072-1083. In some embodiments, proteins comprising a deaminase, a fragments of a deaminase, or homologs of a deaminase or a deaminase are referred to as “deaminase variants.” A deaminase variant shares homology to a deaminase, or a fragment thereof. For example a deaminase variant is at least about 70% identical, at least about 80% identical, at least about 90% identical, at least about 95% identical, at least about 96% identical, at least about 97% identical, at least about 98% identical, at least about 99% identical, at least about 99.5% identical, or at least about 99.9% to a wild type deaminase or a deaminase as set forth in any of SEQ ID NOs: 271-292, 303, and 1072-1083. In some embodiments, the deaminase variant comprises a fragment of the deaminase, such that the fragment is at least about 70% identical, at least about 80% identical, at least about 90% identical, at least about 95% identical, at least about 96% identical, at least about 97% identical, at least about 98% identical, at least about 99% identical, at least about 99.5% identical, or at least about 99.9% to the corresponding fragment of wild type deaminase or a deaminase as set forth in any of SEQ ID NOs: 271-292, 303, and 1072-1083. In some embodiments, the cytosine deaminase is at least at least about 70% identical, at least about 80% identical, at least about 90% identical, at least about 95% identical, at least about 96% identical, at least about 97% identical, at least about 98% identical, at least about 99% identical, at least about 99.5% identical, or at least about 99.9% identical to an APOBEC3G variant as set forth in SEQ ID NO: 291 or SEQ ID NO: 292, and comprises mutations corresponding to the D316E/D317R mutations in SEQ ID NO: 290.

(178) In some embodiments, the cytosine deaminase domain is fused to the N-terminus of the guide nucleotide sequence-programmable DNA-binding protein domain. For example, the fusion protein may have an architecture of NH.sub.2-[cytosine deaminase]-[guide nucleotide sequence-programmable DNA-binding protein domain]-COOH. The “[ ]-” used in the general architecture above indicates the presence of an optional linker sequence. The term “linker,” as used herein, refers to a chemical group or a molecule linking two molecules or moieties, e.g., two domains of a fusion protein, such as, for example, a dCas9 domain and a cytosine deaminase domain. Typically, the linker is positioned between, or flanked by, two groups, molecules, or other moieties and connected to each one via a covalent bond, thus connecting the two. In some embodiments, the linker is an amino acid or a plurality of amino acids (e.g., a peptide or protein). In some embodiments, the linker is an organic molecule, group, polymer, or chemical moiety. In some embodiments, the linker is 5-100 amino acids in length, for example, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 30-35, 35-40, 40-45, 45-50, 50-60, 60-70, 70-80, 80-90, 90-100, 100-150, or 150-200 amino acids in length. Longer or shorter linkers are also contemplated.

(179) In some embodiments, the cytosine deaminase domain and the Cas9 domain are fused to each other via a linker. Various linker lengths and flexibilities between the deaminase domain (e.g., APOBEC1) and the Cas9 domain can be employed (e.g., ranging from very flexible linkers of the form (GGGS).sub.n (SEQ ID NO: 337), (GGGGS).sub.n (SEQ ID NO: 308), (GGS).sub.n (SEQ ID NO: 784), and (G).sub.n (SEQ ID NO: 783) to more rigid linkers of the form (EAAAK).sub.n (SEQ ID NO: 309), SGSETPGTSESATPES (SEQ ID NO: 310) (see, e.g., Guilinger et al., *Nat. Biotechnol.* 2014; 32(6): 577-82; the entire contents are incorporated herein by reference), (SGGS).sub.n SGSETPGTSESATPES (SGGS).sub.n (SEQ ID NO: 1068), (XP).sub.1 (SEQ ID NO: 785), or a combination of any of these, wherein X is any amino acid and n is independently an integer between 1 and 30, in order to achieve the optimal length for deaminase activity for the specific application. In some embodiments, n is independently 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30, or, if more than one linker or more than one linker motif is present, any combination thereof. In some embodiments, the linker comprises a (GGS).sub.n (SEQ ID NO: 784) motif, wherein n is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15. In some embodiments, the linker comprises a (GGS).sub.n (SEQ ID NO: 784) motif, wherein n is 1, 3, or 7. In some embodiments, the linker comprises the amino acid sequence SGSETPGTSESATPES (SEQ ID NO: 310), also referred to as the XTEN linker. In some embodiments, the linker comprises an amino acid sequence chosen from the group including, but not limited to, AGVF (SEQ ID NO: 825), GFLG (SEQ ID NO: 826), FK, AL, ALAL (SEQ ID NO: 827), or ALALA (SEQ ID NO: 828). In some embodiments, suitable linker motifs and configurations include those described in Chen et al., *Fusion protein linkers: property, design and functionality.* *Adv Drug Deliv Rev.* 2013; 65(10):1357-69, which is incorporated herein by reference. In some embodiments, the linker may comprise any of the following amino acid sequences: VPFLLEPDNINGKTC (SEQ ID NO: 311), GSAGSAAGSGEF (SEQ ID NO: 312), SIVAQLSRPDPA (SEQ ID NO: 313), MKIIEQLPSA (SEQ ID NO: 314), VRHKLKRVGS (SEQ ID NO: 315), GHGTGSTGSGSS (SEQ ID NO: 316), MSRPDPA (SEQ ID NO: 317), GSAGSAAGSGEF (SEQ ID NO: 312), SGSETPGTSESA (SEQ ID NO: 318), SGSETPGTSESATPEGGSGGS (SEQ ID NO: 319), or GGSM (SEQ ID NO: 320). Additional suitable linker sequences will be apparent to those of skill in the art based on the instant disclosure.

(180) In some embodiments, the nucleobase editor comprises a guide nucleotide sequence-programmable DNA-binding protein domain and an apolipoprotein B mRNA-editing complex 1 (APOBEC1) deaminase domain, where the deaminase domain is fused to the N-terminus of the napDNAbp domain via a linker comprising the amino acid sequence SGSETPGTSESATPES (SEQ ID NO: 310). In some embodiments, the a guide nucleotide sequence-programmable DNA-binding protein domain comprises the amino acid sequence of any of the a guide nucleotide sequence-programmable DNA-binding protein domains provided herein. In some embodiments, the deaminase is rat APOBEC1 (SEQ ID NO: 288). In some embodiments, the deaminase is human APOBEC1 (SEQ ID NO: 286). In some embodiments, the deaminase is pmCDA1 (SEQ ID NO: 289). In some embodiments, the deaminase is human APOBEC3G (SEQ ID NO:



279). In some embodiments, the deaminase domain of any one of (SEQ ID NOs: 290-292). In some embodiments, the fusion protein comprises a guide nucleotide sequence-programmable DNA-binding protein domain and an apolipoprotein B mRNA-editing complex 1 catalytic polypeptide-like 3G (APOBEC3G) deaminase domain, wherein the deaminase domain is fused to the N-terminus of the a guide nucleotide sequence-programmable DNA-binding protein domain via a linker of any length or composition (e.g., an amino acid sequence, a peptide, a polymer, or a bond). In some embodiments, the linker comprises the amino acid sequence SGSETPGTSESATPES (SEQ ID NO: 310). In some embodiments, the linker comprises the amino acid sequence (SGGS).sub.2SGSETPGTSESATPES(SGGS).sub.2 (SEQ ID NO: 1069).

(181) In some embodiments, the fusion protein comprises a guide nucleotide sequence-programmable DNA-binding protein domain and a cytidine deaminase 1 (CDA1) deaminase domain, wherein the deaminase domain is fused to the N-terminus of the guide nucleotide sequence-programmable DNA-binding protein domain via a linker comprising the amino acid sequence SGSETPGTSESATPES (SEQ ID NO: 310). In some embodiments, the linker comprises the amino acid sequence (SGGS).sub.2SGSETPGTSESATPES(SGGS).sub.2 (SEQ ID NO: 1069). In some embodiments, the guide nucleotide sequence-programmable DNA-binding protein domain comprises the amino acid sequence of any of the guide nucleotide sequence-programmable DNA-binding protein domains provided herein.

(182) In some embodiments, the fusion protein comprises a guide nucleotide sequence-programmable DNA-binding protein and an activation-induced cytidine deaminase (AID) deaminase domain, where the deaminase domain is fused to the N-terminus of the guide nucleotide sequence-programmable DNA-binding protein domain via a linker comprising the amino acid sequence SGSETPGTSESATPES (SEQ ID NO: 310). In some embodiments, the linker comprises the amino acid sequence (SGGS).sub.2SGSETPGTSESATPES(SGGS).sub.2 (SEQ ID NO: 1069). In some embodiments, the guide nucleotide sequence-programmable DNA-binding protein comprises the amino acid sequence of any of the guide nucleotide sequence-programmable DNA-binding protein domains provided herein.

(183) Some aspects of the disclosure are based on the recognition that certain configurations of a guide nucleotide sequence-programmable DNA-binding protein, and a cytidine deaminase domain fused by a linker are useful for efficiently deaminating target cytidine residues. Other aspects of this disclosure relate to the recognition that a nucleobase editing fusion protein with an apolipoprotein B mRNA-editing complex 1 (APOBEC1) deaminase domain fused to the N-terminus of a guide nucleotide sequence-programmable DNA-binding protein via a linker comprising the amino acid sequence SGSETPGTSESATPES (SEQ ID NO: 310) was capable of efficiently deaminating target nucleic acids in a double stranded DNA target molecule. In some embodiments, the fusion protein comprises a guide nucleotide sequence-programmable DNA-binding protein domain and an apolipoprotein B mRNA-editing complex 1 (APOBEC1) deaminase domain, where the deaminase domain is fused to the N-terminus of the napDNAbp via a linker comprising the amino acid sequence SGSETPGTSESATPES (SEQ ID NO: 310). In some embodiments, the fusion protein comprises a guide nucleotide sequence-programmable DNA-binding protein domain and an apolipoprotein B mRNA-editing complex 1 (APOBEC1) deaminase domain, where the deaminase domain is fused to the N-terminus of the napDNAbp via a linker comprising the amino acid sequence (SGGS).sub.2SGSETPGTSESATPES(SGGS).sub.2 (SEQ ID NO: 1069).

(184) To successfully edit the desired target C base, the linker between Cas9 and APOBEC may be optimized, as described in Komor et al., *Nature*, 533, 420-424 (2016), which is incorporated herein by reference. The numbering scheme for base editing is based on the predicted location of the target C within the single stranded stretch of DNA (R-loop) displaced by a programmable guide RNA sequence occurring when a DNA-binding domain (e.g. Cas9, nCas9, dCas9) binds a genomic site (see FIG. 6). Conveniently, the sequence immediately surrounding the target C also matches the sequence of the guide RNA. The numbering scheme for base editing is based on a standard 20-mer programmable sequence, and defines position "21" as the first DNA base of the PAM sequence, resulting in position "1" assigned to the first DNA base matching the 5'-end of the 20-mer programmable guide RNA sequence. Therefore, for all Cas9 variants, position "21" is defined as the first base of the PAM sequence (e.g. NGG, NGAN, NGNG, NGAG, NGCG, NNGRRT, NGRN, NNNRRT, NNGATT, NNAGAA, NAAAC). When a longer programmable guide RNA sequence is used (e.g. 21-mer) the 5'-end bases are assigned a decreasing negative number starting at "-1". For other DNA-binding domains that differ in the position of the PAM sequence, or that require no PAM sequence, the programmable guide RNA sequence is used as a reference for numbering. A 3-aa linker gives a 2-5 base editing window (e.g., positions 2, 3, 4, or 5 relative to the PAM sequence at position 21). A 9-aa linker gives a 3-6 base editing window (e.g., positions 3, 4, 5, or 6 relative to the PAM sequence at position 21). A 16-aa linker (e.g., the SGSETPGTSESATPES (SEQ ID NO: 310) linker) gives a 4-7 base editing window (e.g., positions 4, 5, 6, or 7 relative to the PAM sequence at position 21). A 21-aa linker gives a 5-8 base editing window (e.g., positions 5, 6, 7, 8 relative to the PAM sequence at position 21). Each of these windows can be useful for editing different targeted C bases. For example, the targeted C bases may be at different distances from the adjacent PAM sequence, and by varying the linker length, the precise editing of the desired C base is ensured. One skilled in the art, based on the teachings of CRISPR/Cas9 technology, in particular the teachings of U.S. Provisional applications 62/245,828, 62/279,346, 62/311,763, 62/322,178, 62/357352, 62/370,700, and 62/398,490, and in Komor et al., *Nature*, Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage, 533, 420-424 (2016), each of which is incorporated herein by reference, will be able to determine the window of editing for his/her purpose, and properly design the linker of the cytosine deaminase-dCas9 protein for the precise targeting of the desired C base.

(185) To successfully edit the desired target C base, appropriate Cas9 domain may be selected to attached to the deaminase domain (e.g., APOBEC1), since different Cas9 domains may lead to different editing windows, as described in

in U.S. Pat. Nos. 9,068,179, US Patent Application Publications US 2015/0166980, US 2015/0166981, US 2015/0166982, US20150166984, and US20150165054, and US Provisional Applications, U.S. Ser. No. 62/245,828, filed Oct. 23, 2015; 62/279,346, filed Jan. 15, 2016; 62/311,763, filed Mar. 22, 2016; 62/322,178, filed Apr. 13, 2016; 62/357,352, filed Jun. 30, 2016, U.S. Pat. No. 62,370,700, filed Aug. 3, 2016; 62/398,490, filed Sep. 22, 2016; 62/408,686, filed Oct. 14, 2016; PCT Application PCT/US2016/058344, filed Oct. 22, 2016, U.S. patent application Ser. No. 15/311,852, filed Oct. 22, 2016; and in Komor et al., *Nature*, Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage, 533, 420-424 (2016), the entire contents of each of which are incorporated herein by reference. For example, APOBEC1-XTEN-SaCas9n-UGI gives a 1-12 base editing window (e.g., positions 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 relative to the NNNRRT PAM sequence in positions 20-26). One skilled in the art, based on the teachings of CRISPR/Cas9 technology, will be able to determine the editing window for his/her purpose, and properly determine the required Cas9 homolog and linker attached to the cytosine deaminase for the precise targeting of the desired C base.

(186) In some embodiments, the fusion protein useful in the present disclosure further comprises a uracil glycosylase inhibitor (UGI) domain. A “uracil glycosylase inhibitor” refers to a protein that inhibits the activity of uracil-DNA glycosylase. The C to T base change induced by deamination results in a U:G heteroduplex, which triggers cellular DNA-repair response. Uracil DNA glycosylase (UDG) catalyzes removal of U from DNA in cells and initiates base excision repair, with reversion of the U:G pair to a C:G pair as the most common outcome. Thus, such cellular DNA-repair response may be responsible for the decrease in nucleobase editing efficiency in cells. Uracil DNA Glycosylase Inhibitor (UGI) is known in the art to potently blocks human UDG activity. As described in Komor et al., *Nature* (2016), fusing a UGI domain to the cytidine deaminase-dCas9 fusion protein reduced the activity of UDG and significantly enhanced editing efficiency.

(187) Suitable UGI protein and nucleotide sequences are provided herein and additional suitable UGI sequences are known to those in the art, and include, for example, those published in Wang et al., Uracil-DNA glycosylase inhibitor gene of bacteriophage PBS2 encodes a binding protein specific for uracil-DNA glycosylase. *J. Biol. Chem.* 264:1163-1171(1989); Lundquist et al., Site-directed mutagenesis and characterization of uracil-DNA glycosylase inhibitor protein. Role of specific carboxylic amino acids in complex formation with *Escherichia coli* uracil-DNA glycosylase. *J. Biol. Chem.* 272:21408-21419(1997); Ravishankar et al., X-ray analysis of a complex of *Escherichia coli* uracil DNA glycosylase (EcUDG) with a proteinaceous inhibitor. The structure elucidation of a prokaryotic UDG. *Nucleic Acids Res.* 26:4880-4887(1998); and Putnam et al., Protein mimicry of DNA from crystal structures of the uracil-DNA glycosylase inhibitor protein and its complex with *Escherichia coli* uracil-DNA glycosylase. *J. Mol. Biol.* 287:331-346(1999), each of which is incorporated herein by reference. In some embodiments, the UGI domain comprises the amino acid sequence of SEQ ID NO: 304 without the N-terminal methionine (M). In some embodiments, the UGI comprises the following amino acid sequence: *Bacillus* phage PBS2 (Bacteriophage PBS2)Uracil-DNA glycosylase inhibitor  
MTNLSDIIEKETGKQLVIQESILMLPEEVVEEVIGNKPESDILVHTAYDESTDENVMMLLTSDAPEYKPWAL  
VIQDSNGENKIKML (SEQ ID NO: 304)

(188) In some embodiments, the UGI protein comprises a wild type UGI or a UGI as set forth in SEQ ID NO: 304. In some embodiments, the UGI proteins useful in the present disclosure include fragments of UGI and proteins homologous to a UGI or a UGI fragment. For example, in some embodiments, a UGI comprises a fragment of the amino acid sequence set forth in SEQ ID NO: 304. In some embodiments, a UGI comprises an amino acid sequence homologous to the amino acid sequence set forth in SEQ ID NO: 304 or an amino acid sequence homologous to a fragment of the amino acid sequence set forth in SEQ ID NO: 304. In some embodiments, proteins comprising UGI or fragments of UGI or homologs of UGI or UGI fragments are referred to as “UGI variants.” A UGI variant shares homology to UGI, or a fragment thereof. For example a UGI variant is at least about 70% identical, at least about 80% identical, at least about 90% identical, at least about 95% identical, at least about 96% identical, at least about 97% identical, at least about 98% identical, at least about 99% identical, at least about 99.5% identical, or at least about 99.9% to a wild type UGI or a UGI as set forth in SEQ ID NO: 304. In some embodiments, the UGI variant comprises a fragment of UGI, such that the fragment is at least about 70% identical, at least about 80% identical, at least about 90% identical, at least about 95% identical, at least about 96% identical, at least about 97% identical, at least about 98% identical, at least about 99% identical, at least about 99.5% identical, or at least about 99.9% to the corresponding fragment of wild type UGI or a UGI as set forth in SEQ ID NO: 304.

(189) It should be appreciated that additional proteins may be uracil glycosylase inhibitors. For example, other proteins that are capable of inhibiting (e.g., sterically blocking) a uracil-DNA glycosylase base-excision repair enzyme are within the scope of this disclosure. In some embodiments, a uracil glycosylase inhibitor is a protein that binds DNA. In some embodiments, a uracil glycosylase inhibitor is a protein that binds single-stranded DNA. For example, a uracil glycosylase inhibitor may be a *Erwinia tasmaniensis* single-stranded binding protein. In some embodiments, the single-stranded binding protein comprises the amino acid sequence (SEQ ID NO: 305). In some embodiments, a uracil glycosylase inhibitor is a protein that binds uracil. In some embodiments, a uracil glycosylase inhibitor is a protein that binds uracil in DNA. In some embodiments, a uracil glycosylase inhibitor is a catalytically inactive uracil DNA-glycosylase protein. In some embodiments, a uracil glycosylase inhibitor is a catalytically inactive uracil DNA-glycosylase protein that does not excise uracil from the DNA. For example, a uracil glycosylase inhibitor is a UdgX. In some embodiments, the UdgX comprises the amino acid sequence (SEQ ID NO: 306). As another example, a uracil glycosylase inhibitor is a catalytically inactive UDG. In some embodiments, a catalytically inactive UDG comprises the amino acid sequence (SEQ

ID NO: 307). It should be appreciated that other uracil glycosylase inhibitors would be apparent to the skilled artisan and are within the scope of this disclosure.

(190) TABLE-US-00023 *Erwinia tasmaniensis* SSB (thermostable single- stranded DNA binding protein) (SEQ ID NO: 305) MASRGVNVKLVGLGQDPEVRYMPNNGAVANITLATSESWRDKQTGET  
KEKTEWHRVVLFGKLAEVAGEYLRKGSQVYIEGALQTRKWTDQAGVEKY  
TTEVVVNVGGMQMLGGRSQGGGASAGGQNGGSNNGWGQPQQPQGGNQF  
SGGAQQQARPQQPQQNNAPANNEPIDEDDDIP UdgX (binds to Uracil in DNA but does not excise) (SEQ ID NO: 306) MAGAQDFVPHTADLAELAAAAGECRGCGLYRDATQAVFGAGGRSARIMM  
IGEQQGDKEDLAGLPFVGPAGRLLDRALEAADIDRDALYVTNAVKHFKF  
TRAAGGKRRIHKTPSRTEVVACRPWLIAEMTSVEPDVVLLGATAAKAL  
LGNDFRVTQHRGEVLHVDDVPGDPALVATVHPSSLLRGPKEERESAFAG LVDDLRVAADVVRP UDG  
(catalytically inactive human UDG, binds to Uracil in DNA but does not excise) (SEQ ID NO: 307) MIGQKTLYSFFSPSPARKRHAPSPEPAVQGTGVAGVPEESGDAAAIPAK  
KAPAGQEEPPTPSSPLSAEQLDRIQRNKAALLRLAARNVPVGFGESEW  
KKHLSGEFGKPYFIKLMGFVAERKHVTVYPPPHQVFTWTQMCDIKDVK  
VVILGQEPYHGPNQAHGLCFVQRPVPPPPSLENIYKELSTDIEDFVHP  
GHGDLGWAQGVLLNVAULTVRAHQANSHKERGWEQFTDAVVSWLNNQ  
SNGLVFLLWGSYAQKKGSAIDRKRHHVLQTAHPSPLSVYRGFFGCRHFS KTNELLQKSGKKPIDWKEL

(191) In some embodiments, the UGI domain is fused to the C-terminus of the dCas9 domain in the fusion protein. Thus, the fusion protein would have an architecture of NH.sub.2-[cytosine deaminase]-[guide nucleotide sequence-programmable DNA-binding protein domain]-[UGI]-COOH. In some embodiments, the UGI domain is fused to the N-terminus of the cytosine deaminase domain. As such, the fusion protein would have an architecture of NH.sub.2-[UGI]-[cytosine deaminase]-[guide nucleotide sequence-programmable DNA-binding protein domain]-COOH. In some embodiments, the UGI domain is fused between the guide nucleotide sequence-programmable DNA-binding protein domain and the cytosine deaminase domain. As such, the fusion protein would have an architecture of NH.sub.2-[cytosine deaminase]-[UGI]-[guide nucleotide sequence-programmable DNA-binding protein domain]-COOH. The linker sequences described herein may also be used for the fusion of the UGI domain to the cytosine deaminase-dCas9 fusion proteins.

(192) In some embodiments, the fusion protein comprises the structure: [cytosine deaminase]-[optional linker sequence]-[guide nucleotide sequence-programmable DNA binding protein]-[optional linker sequence]-[UGI]; [cytosine deaminase]-[optional linker sequence]-[UGI]-[optional linker sequence]-[guide nucleotide sequence-programmable DNA binding protein]; [UGI]-[optional linker sequence]-[cytosine deaminase]-[optional linker sequence]-[guide nucleotide sequence-programmable DNA binding protein]; [UGI]-[optional linker sequence]-[guide nucleotide sequence-programmable DNA binding protein]-[optional linker sequence]-[cytosine deaminase]; [guide nucleotide sequence-programmable DNA binding protein]-[optional linker sequence]-[cytosine deaminase]-[optional linker sequence]-[UGI]; or [guide nucleotide sequence-programmable DNA binding protein]-[optional linker sequence]-[UGI]-[optional linker sequence]-[cytosine deaminase].

(193) In some embodiments, the fusion protein comprises the structure: [cytosine deaminase]-[optional linker sequence]-[Cas9 nickase]-[optional linker sequence]-[UGI]; [cytosine deaminase]-[optional linker sequence]-[UGI]-[optional linker sequence]-[Cas9 nickase]; [UGI]-[optional linker sequence]-[cytosine deaminase]-[optional linker sequence]-[Cas9 nickase]; [UGI]-[optional linker sequence]-[Cas9 nickase]-[optional linker sequence]-[cytosine deaminase]; [Cas9 nickase]-[optional linker sequence]-[cytosine deaminase]-[optional linker sequence]-[UGI]; or [Cas9 nickase]-[optional linker sequence]-[UGI]-[optional linker sequence]-[cytosine deaminase].

(194) In some embodiments, fusion proteins provided herein further comprise a nuclear localization sequence (NLS). In some embodiments, the NLS is fused to the N-terminus of the fusion protein. In some embodiments, the NLS is fused to the C-terminus of the fusion protein. In some embodiments, the NLS is fused to the N-terminus of the UGI protein. In some embodiments, the NLS is fused to the C-terminus of the UGI protein. In some embodiments, the NLS is fused to the N-terminus of the guide nucleotide sequence-programmable DNA-binding protein domain. In some embodiments, the NLS is fused to the C-terminus of the guide nucleotide sequence-programmable DNA-binding protein domain. In some embodiments, the NLS is fused to the N-terminus of the cytosine deaminase. In some embodiments, the NLS is fused to the C-terminus of the deaminase. In some embodiments, the NLS is fused to the fusion protein via one or more linkers. In some embodiments, the NLS is fused to the fusion protein without a linker. Non-limiting, exemplary NLS sequences may be PKKKRKV (SEQ ID NO: 829) or MDSLLMNRKFLYQFKNVRWAKGRRETYLC (SEQ ID NO: 830).

(195) In some embodiments, any of the fusion proteins provided herein comprise a second UGI domain. Fusion proteins comprising two UGI domains are described in U.S. Provisional Applications, U.S. Ser. No. 62/475,830, filed Mar. 23, 2017; 62/490,587; 62/511,934, filed May 26, 2017; 62/551,951, filed Aug. 30, 2017; and Komor et al. (2017) Improved Base Excision Repair Inhibition and Bacteriophage Mu Gam Protein Yields C:G-to-T:A base editors with higher efficiency and product purity. *Sci Adv*, 3: eaao4774; the entire contents of which is incorporated herein by reference. In some embodiments, the second UGI domain comprises a wild-type UGI or a UGI as set forth in SEQ ID NO: 304. In some embodiments, the UGI proteins provided herein include fragments of UGI and proteins homologous to a UGI or a UGI fragment. For example, in some embodiments, the second UGI domain comprises a fragment of the amino acid sequence

set forth in SEQ ID NO: 304. In some embodiments, a UGI fragment comprises an amino acid sequence that comprises at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% of the amino acid sequence as set forth in SEQ ID NO: 304. In some embodiments, the second UGI domain comprises an amino acid sequence homologous to the amino acid sequence set forth in SEQ ID NO: 304 or an amino acid sequence homologous to a fragment of the amino acid sequence set forth in SEQ ID NO: 304. In some embodiments, proteins comprising UGI or fragments of UGI or homologs of UGI or UGI fragments are referred to as “UGI variants.” A UGI variant shares homology to UGI, or a fragment thereof. For example a UGI variant is at least 70% identical, at least 75% identical, at least 80% identical, at least 85% identical, at least 90% identical, at least 95% identical, at least 96% identical, at least 97% identical, at least 98% identical, at least 99% identical, at least 99.5% identical, or at least 99.9% identical to a wild type UGI or a UGI as set forth in SEQ ID NO: 304. In some embodiments, the UGI variant comprises a fragment of UGI, such that the fragment is at least 70% identical, at least 80% identical, at least 90% identical, at least 95% identical, at least 96% identical, at least 97% identical, at least 98% identical, at least 99% identical, at least 99.5% identical, or at least 99.9% to the corresponding fragment of wild-type UGI or a UGI as set forth in SEQ ID NO: 304.

(196) In some embodiments, the fusion protein comprises the structure: [deaminase]-[optional linker sequence]-[dCas9]-[optional linker sequence]-[first UGI]-[optional linker sequence]-[second UGI]; [deaminase]-[optional linker sequence]-[Cas9 nickase]-[optional linker sequence]-[first UGI]-[optional linker sequence]-[second UGI]; or [deaminase]-[optional linker sequence]-[Cas9]-[optional linker sequence]-[first UGI]-[optional linker sequence]-[second UGI].

(197) In some embodiments, the nucleobase editor comprises a guide nucleotide sequence-programmable DNA-binding protein domain and an apolipoprotein B mRNA-editing complex 1 (APOBEC1) deaminase domain, wherein the deaminase domain is fused to the N-terminus of the guide nucleotide sequence-programmable DNA-binding protein domain via a linker comprising the amino acid sequence (SGGS).sub.2SGSETPGTSESATPES(SGGS).sub.2 (SEQ ID NO: 1069). In some embodiments, the a guide nucleotide sequence-programmable DNA-binding protein domain comprises the amino acid sequence of any of the a guide nucleotide sequence-programmable DNA-binding protein domains provided herein. In some embodiments, the deaminase is rat APOBEC1 (SEQ ID NO: 288). In some embodiments, the deaminase is human APOBEC1 (SEQ ID NO: 286). In some embodiments, the deaminase is a human APOBEC3G variant of any one of (SEQ ID NOs: 290-292). In some embodiments, the nucleobase editor comprises a first UGI domain fused to the C-terminus of a guide nucleotide sequence-programmable DNA-binding protein domain via a linker comprising the amino acid sequence (GGS).sub.n(SEQ ID NO: 784), wherein n is 3. In some embodiments, the nucleobase editor comprises a second UGI domain fused to the C-terminus of a first UGI domain via a linker comprising the amino acid sequence (GGS).sub.n (SEQ ID NO: 784), wherein n is 3.

(198) In some embodiments, the fusion protein comprises the amino acid sequence of SEQ ID NO: 1084. In some embodiments, the fusion protein comprises an amino acid sequence that is at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to the amino acid sequence as set forth in SEQ ID NO: 1084.

(199) In some embodiments, any of the fusion proteins provided herein may further comprise a Gam protein. The term “Gam protein,” as used herein, refers generally to proteins capable of binding to one or more ends of a double strand break of a double stranded nucleic acid (e.g., double stranded DNA). In some embodiments, the Gam protein prevents or inhibits degradation of one or more strands of a nucleic acid at the site of the double strand break. In some embodiments, a Gam protein is a naturally-occurring Gam protein from bacteriophage Mu, or a non-naturally occurring variant thereof. Fusion proteins comprising Gam proteins are described in Komor et al. (2017) Improved Base Excision Repair Inhibition and Bacteriophage Mu Gam Protein Yields C:G-to-T:A base editors with higher efficiency and product purity. *Sci Adv*, 3: eaao4774; the entire contents of which is incorporated by reference herein. In some embodiments, the Gam protein comprises an amino acid sequence that is at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 99.5% identical to the amino acid sequence provided by SEQ ID NO: 3027. In some embodiments, the Gam protein comprises the amino acid sequence of SEQ ID NO: 3027. In some embodiments, the fusion protein (e.g., BE4-Gam of SEQ ID NO: 3028) comprises a Gam protein, wherein the Cas9 domain of BE4 is replaced with any of the Cas9 domains provided herein.

(200) TABLE-US-00024 Gam from bacteriophage Mu: (SEQ ID NO: 1070)

AKPAKRIKSAAAAYVPQNRDAVITDIKRIGDLQREASRLETEMNDAIAEITEKFAARIAPIKTDIETLSKGVQGW  
CEANRDELTNNGGKVKTNLVTGDVSWVRPPSVSIRGMDAVMETLERLGLQRFIRTKQEINKEAILLEPKAVAGV  
AGITVKSGIEDFSIIPFEQEAGI BE4-Gam: (SEQ ID NO: 1071)

**MAKPAKRIKSAAAAYVPQNRDAVITDIKRIGDLQREASRLETEMNDAIAEITEKFAARIAPIKTDIETLSKGVQGW**  
**CEANRDELTNNGGKVKTNLVTGDVSWVRPPSVSIRGMDAVMETLERLGLQRFIRTKQEINKEAILLEPKAVAGV**  
**VAGITVKSGIEDFSIIPFEQEAGISGSETPGTSESATPESSETGPVAVDPTLRRRIEPHEFEVFFDPRELKRET**  
CLLYEINWGGRRHSIWRHTSQNTNKHVEVNFIEKFTTERRYFCPNTRCSITWFLSWSPCGECSRAITEFLSRYPHVT  
LFIYIARLYHHADPRNRQGLRDLISSGVTIQIMTEQESGYCWRNRFVNYSPSNEAHWPYPHPLWVRLYVLELYCII  
LGLPPCLNILRRKQPQLTFFTIALQSCHYQRLPPHILWATGLKSGGSSGGSSGSETPGTSESATPESGGSSGGSS  
DKKYSIGLAIGTNSVGWAVITDEYKVPSKKFKVLGNTDRHSIKKNLIGALLFDSGETAEATRLKRTARRRYTRRK  
NRICYLQEIFSNEMAKVDDSFHRLLEESFLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLRKKLVDSTDKADL  
RLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDAKAILSARLSKSRLENLI

AQLPGKEALNGLFNKLSGLTNPKNFKSFDNLAQIGDYADLFLAAKNLSDAIL  
 LSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLKALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFI  
 KPILEKMDGTEELLVKLNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFYPLKDNREKIEKILTRIPYY  
 VGPLARGNSRFAWMTRKSEETITPWNFEFVVDKGASAQSFIERMTNFDKNLPNEKVLPKHSLLYEYFTVYNELTK  
 VKYVTEGMRKPAFLSGEQKKAIVDLLFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRENASLGTYHDLKII  
 KDKDFLDNEENEDILEDIVLTLTLFEDREMIEERLKYAHLEDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSG  
 KTILDFLKSDGFANRNFQMQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIIKGILQTVKVVDDELVKVM  
 GRHKPENIVIEMARENQTTQKGQKNSRERMKRIIEGKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQ  
 ELDINRLSDYDVDHIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKKMKNYWRQLLNAKLITQRKFDNLT  
 KAERGGSELDDKAGFIKRLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFRKDFQFYKVR  
 EINNYHHAHDAYLNAVVGTAIIKKYPKLESEFVYGDYKVYDVRKMIKSEQEIGKATAKYFFYSNIMNFFKTEIT  
 LANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVLSPQVNVKKTETVQTGGFSKESILPKRNSDKLIARKKDW  
 DPKKYGGFDSPTVAYSVLVAKVEKGKSKKLKSVKELLGITIMERSSEFKNPIDFLEAKGYKEVKKDLIILPKY  
 SLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLASHYEKLKGGSPEDNEQKQLFVEQHKHYLDEIEQISEF  
 SKRVILADANLDKVL SAYNKHRRDKPIREQAENIIHFLTLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDTLHQS  
 ITGLYETRIDLSQLGGDSGGSGGGSGGSTNLSIIKETGKQLVIQESILMLPEEVEEVIGNKPESDILVHTAYDE  
 STDENVMLLTSDAPEYKPWALVIQDSNGENKIKMLSGSGSGSGGSTNLSIIKETGKQLVIQESILMLPEEVEE  
 VIGNKPESDILVHTAYDESTDENVMLLTSDAPEYKPWALVIQDSNGENKIKMLSGSGSPKKKRK

(201) Some aspects of the present disclosure provide nucleobase editors associated with a guide nucleotide sequence (e.g., a guide RNA or gRNA). gRNAs can exist as a complex of two or more RNAs, or as a single RNA molecule. gRNAs that exist as a single RNA molecule may be referred to as single-guide RNAs (sgRNAs), though “gRNA” is used interchangeably to refer to guide RNAs that exist as either single molecules or as a complex of two or more molecules. Typically, gRNAs that exist as a single RNA species comprise two domains: (1) a domain that shares homology to a target nucleic acid (e.g., and directs binding of the Cas9 complex to the target); and (2) a domain that binds the Cas9 protein. In some embodiments, domain (2) corresponds to a sequence known as a tracrRNA, and comprises a stem-loop structure. For example, in some embodiments, domain (2) is identical or homologous to a tracrRNA as provided in Jinek et al., *Science* 337:816-821(2012), which is incorporated herein by reference. Other examples of gRNAs (e.g., those including domain 2) can be found in U.S. Provisional Patent Application, U.S. Ser. No. 61/874,682, filed Sep. 6, 2013, entitled “Switchable Cas9 Nucleases And Uses Thereof,” and U.S. Provisional Patent Application, U.S. Ser. No. 61/874,746, filed Sep. 6, 2013, entitled “Delivery System For Functional Nucleases,” each of which is incorporated herein by reference in their entirety. The gRNA comprises a nucleotide sequence that complements a target site, which mediates binding of the nuclease/RNA complex to said target site, providing the sequence specificity of the nuclease:RNA complex. These proteins are able to be targeted, in principle, to any sequence specified by the guide RNA. Methods of using RNA-programmable nucleases, such as Cas9, for site-specific cleavage (e.g., to modify a genome) are known in the art (see e.g., Cong, L. et al. *Science* 339, 819-823 (2013); Mali, P. et al. *Science* 339, 823-826 (2013); Hwang, W. Y. et al. *Nature biotechnology* 31, 227-229 (2013); Jinek, M. et al. *eLife* 2, e00471 (2013); Dicarolo, J. E. et al. *Nucleic acids research* (2013); Jiang, W. et al. *Nature Biotechnology* 31, 233-239 (2013); each of which are incorporated herein by reference). In particular, examples of guide nucleotide sequences (e.g., sgRNAs) that may be used to target the fusion protein of the present disclosure to its target sequence to deaminate the targeted C bases are described in Komor et al., *Nature*, 533, 420-424 (2016), which is incorporated herein by reference.

(202) The specific structure of the guide nucleotide sequences (e.g., sgRNAs) depends on its target sequence and the relative distance of a PAM sequence downstream of the target sequence. One skilled in the art will understand, that no unifying structure of guide nucleotide sequence is given, for that the target sequences are different for each and every C targeted to be deaminated.

(203) However, the present disclosure provides guidance in how to design the guide nucleotide sequence, e.g., an sgRNA, so that one skilled in the art may use such teaching to a target sequence of interest. An gRNA typically comprises a tracrRNA framework allowing for Cas9 binding, and a guide sequence, which confers sequence specificity to fusion proteins disclosed herein. In some embodiments, the guide RNA comprises a structure 5'-[guide sequence]-tracrRNA-3'. Non-limiting, exemplary tracrRNA sequences are shown in Table 11.

(204) TABLE-US-00025 TABLE 11 TracrRNA orthologues and sequences SEQ Organism tracrRNA sequence  
 ID NO S. *pyogenes* GUUUAAGAGCUAUGCUGGAAAGCCACGGUGAAAAA 322  
 GUUCAACUAUUGCCUGAUCGGAUAAAUUGAACG AUACGACAGUCGGUGCUUUUUU S. *pyogenes*  
 GUUUAAGAGCUAGAAAUAGCAAGUUUAAUAAGGC 323  
 UAGUCCGUUAUCAACUUGAAAAAGUGGCACCGAGU CGGUGCUUUUUU S. *thermophilus* CRISPR1  
 GUUUUUGUACUCUCAAGAUUCAAUAAUCUUGCAGA 324  
 AGCUACAAAGAUAAAGGCUUCAUGCCGAAAUCAACA CCCUGUCAUUUUAUGGCAGGGUGUUUU S.  
*thermophilus* CRISPR3 GUUUUAGAGCUGUGUUGUUUGUUAACAACACAG 325  
 CGAGUAAAAUAAGGCUUAGUCCGUACUCAACUUG AAAAGGUGGCACCGAUUCGGUGUUUUU C. *jejuni*  
 AAGAAAUUUAAAAAGGGACUAAAAUAAAGAGUUUG 326  
 CGGGACUCUGCGGGGUUACAAUCCCCUAAAACCGCU UUU F. *novicida*  
 AUCUAAAAUUAAUAUGUACCAAUAUAUAAUGCU 327

CUGUAACAUUUUAAAAGUAUUUUGAACGGACCCUCU GUUUGACACGUCUGAAUAACUAAAA S.  
*thermophilus*2 UGUAAGGGACGCCUACACAGUUAUUAAAUUCUUG 328  
 CAGAAGCUACAAAGAUAAAGGCUUCAUGCCGAAAUC  
 AACACCCUGUCAUUUUUAUGGCAGGGUGUUUUCGUU AUUU M. *mobile*  
 UGUUUUUCGAAAUAACAGAUACAGUUAAGAAUAC 329  
 AUAAGAAUGAUACAUCACUAAAAAAGGCUUUAUG  
 CCGUAACUACUACUUAUUUUCAAAAUAAGUAGUUU UUUUU L. *innocua*  
 AUUGUUAGUAUUCAAAAUAACAUAGCAAGUUAAAA 330  
 UAAGGCUUUGUCCGUUAUCAACUUUUAAUUAAGUA GCGCUGUUUCGGCGCUUUUUUU S. *pyogenes*  
 GUUGGAACCAUUCAAAACAGCAUAGCAAGUUAAAA 331  
 UAAGGCUAGUCCGUUAUCAACUUGAAAAAGUGGCA CCGAGUCGGUGCUUUUUUU S. *mutans*  
 GUUGGAAUCAUUCGAAACAACACAGCAAGUUAAAA 332  
 UAAGGCAGUGAUUUUUUAUCCAGUCCGUACACAAC  
 UUGAAAAAGUGCGCACCGAUUCGGUGCUUUUUUAU UU S. *thermophilus*  
 UUGUGGUUUGAAACCAUUCGAAACAACACAGCGAG 333  
 UUAAAAUAAGGCUUAGUCCGUACUCAACUUGAAAA GGUGGCACCGAUUCGGUGUUUUUUUU N.  
*meningitidis* ACAUAUUGUCGCACUGCGAAAUGAGAACCGUUGCU 334  
 ACAUAUAGGCCGUCUGAAAAGAUGUGCCGCAACGC UCUGCCCCUAAAAGCUUCUGCUUUAAGGGGCA  
 P. *multocida* GCAUAUUGUUGCACUGCGAAAUGAGAGACGUUGCU 335  
 ACAUAUAGGCUUCUGAAAAGAAUGACCGUAACGCU  
 CUGCCCCUUGUGAUUCUUAUUGCAAGGGGCAUCG UUUUU

(205) The guide sequence of the gRNA comprises a sequence that is complementary to the target sequence. The guide sequence is typically about 20 nucleotides long. For example, the guide sequence may be 15-25 nucleotides long. In some embodiments, the guide sequence is 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotides long. In some embodiments, the guide sequence is more than 25 nucleotides long. Such suitable guide RNA sequences typically comprise guide sequences that are complementary to a nucleic sequence within 50 nucleotides upstream or downstream of the target nucleotide to be edited.

(206) In some embodiments, the guide RNA is about 15-100 nucleotides long and comprises a sequence of at least 10 contiguous nucleotides that is complementary to a target sequence. In some embodiments, the guide RNA is 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides long. In some embodiments, the guide RNA comprises a sequence of 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 contiguous nucleotides that is complementary to a target sequence.

(207) Compositions

(208) Some aspects of the present disclosure relate to compositions that may be used for generating cancer vaccines in vivo or ex vivo. In some embodiments, the composition comprises: (i) a nucleobase editor or a nucleic acid molecule encoding the nucleobase editor described herein; and (ii) a guide nucleotide sequence targeting the nucleobase editor to a tumor specific antigen-encoding polynucleotide. The guide nucleotide sequence that may be used to generate heteroclitic or cryptic epitopes may be selected from SEQ ID NOs: X-X. Guide nucleotide sequences for generating specific heteroclitic or cryptic epitopes may be found in Tables 5 and 6.

(209) In some embodiments, the composition described herein further comprises a pharmaceutically acceptable carrier.

(210) As used here, the term “pharmaceutically-acceptable carrier” means a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, manufacturing aid (e.g., lubricant, talc magnesium, calcium or zinc stearate, or steric acid), or solvent encapsulating material, involved in carrying or transporting the compound from one site (e.g., the delivery site) of the body, to another site (e.g., organ, tissue or portion of the body). A pharmaceutically acceptable carrier is “acceptable” in the sense of being compatible with the other ingredients of the formulation and not injurious to the tissue of the subject (e.g., physiologically compatible, sterile, physiologic pH, etc.). Some examples of materials which can serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, methylcellulose, ethyl cellulose, microcrystalline cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) lubricating agents, such as magnesium stearate, sodium lauryl sulfate and talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol (PEG); (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) pH buffered solutions; (21) polyesters, polycarbonates and/or polyanhydrides; (22) bulking agents, such as polypeptides and amino acids (23) serum component, such as serum albumin, HDL and LDL; (22) C2-C12 alcohols, such as ethanol; and (23) other non-toxic compatible substances employed in pharmaceutical formulations. Wetting agents, coloring agents, release agents, coating agents, sweetening agents, flavoring agents, perfuming agents, preservative and antioxidants can also be present in the formulation. The terms such as “excipient”, “carrier”, “pharmaceutically acceptable carrier” or the like are used

interchangeably herein.

(211) In some embodiments, the nucleobase editors and the guide nucleotides of the present disclosure in a composition is administered by injection, by means of a catheter, by means of a suppository, or by means of an implant, the implant being of a porous, non-porous, or gelatinous material, including a membrane, such as a sialastic membrane, or a fiber. In some embodiments, the injection is directed to the liver.

(212) In other embodiments, the nucleobase editors and the guide nucleotides are delivered in a controlled release system. In one embodiment, a pump may be used (see, e.g., Langer, 1990, *Science* 249:1527-1533; Sefton, 1989, *CRC Crit. Ref Biomed. Eng.* 14:201; Buchwald et al., 1980, *Surgery* 88:507; Saudek et al., 1989, *N. Engl. J. Med.* 321:574). In another embodiment, polymeric materials can be used. (See, e.g., Medical Applications of Controlled Release (Langer and Wise eds., CRC Press, Boca Raton, Fla., 1974); Controlled Drug Bioavailability, Drug Product Design and Performance (Smolen and Ball eds., Wiley, New York, 1984); Ranger and Peppas, 1983, *Macromol. Sci. Rev. Macromol. Chem.* 23:61. See also Levy et al., 1985, *Science* 228:190; During et al., 1989, *Ann. Neurol.* 25:351; Howard et al., 1989, *J. Neurosurg.* 71:105.) Other controlled release systems are discussed, for example, in Langer, supra.

(213) In typical embodiments, the pharmaceutical composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous or subcutaneous administration to a subject, e.g., a human. Typically, compositions for administration by injection are solutions in sterile isotonic aqueous buffer. Where necessary, the pharmaceutical can also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the pharmaceutical is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the pharmaceutical is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients can be mixed prior to administration.

(214) A pharmaceutical composition for systemic administration may be a liquid, e.g., sterile saline, lactated Ringer's or Hank's solution. In addition, the pharmaceutical composition can be in solid forms and re-dissolved or suspended immediately prior to use. Lyophilized forms are also contemplated.

(215) The pharmaceutical composition can be contained within a lipid particle or vesicle, such as a liposome or microcrystal, which is also suitable for parenteral administration. The particles can be of any suitable structure, such as unilamellar or plurilamellar, so long as compositions are contained therein. Compounds can be entrapped in 'stabilized plasmid-lipid particles' (SPLP) containing the fusogenic lipid dioleoylphosphatidylethanolamine (DOPE), low levels (5-10 mol %) of cationic lipid, and stabilized by a polyethyleneglycol (PEG) coating (Zhang Y. P. et al., *Gene Ther.* 1999, 6:1438-47). Positively charged lipids such as N-[1-(2,3-dioleoyloxy)propyl]-N,N,N-trimethyl-ammoniummethylsulfate, or "DOTAP," are particularly preferred for such particles and vesicles. The preparation of such lipid particles is well known. See, e.g., U.S. Pat. Nos. 4,880,635; 4,906,477; 4,911,928; 4,917,951; 4,920,016; and 4,921,757.

(216) The pharmaceutical compositions of this disclosure may be administered or packaged as a unit dose, for example. The term "unit dose" when used in reference to a pharmaceutical composition of the present disclosure refers to physically discrete units suitable as unitary dosage for the subject, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required diluent; i.e., carrier, or vehicle.

(217) In some embodiments, the nucleobase editors or the guide nucleotides described herein may be conjugated to a therapeutic moiety, e.g., an anti-inflammatory agent. Techniques for conjugating such therapeutic moieties to polypeptides, including e.g., Fc domains, are well known; see, e.g., Amon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in Monoclonal Antibodies And Cancer Therapy, Reisfeld et al. (eds.), 1985, pp. 243-56, Alan R. Liss, Inc.); Hellstrom et al., "Antibodies For Drug Delivery", in Controlled Drug Delivery (2nd Ed.), Robinson et al. (eds.), 1987, pp. 623-53, Marcel Dekker, Inc.); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in Monoclonal Antibodies '84: Biological And Clinical Applications, Pinchera et al. (eds.), 1985, pp. 475-506); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in Monoclonal Antibodies For Cancer Detection And Therapy, Baldwin et al. (eds.), 1985, pp. 303-16, Academic Press; and Thorpe et al. (1982) "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates," *Immunol. Rev.*, 62:119-158.

(218) Further, the compositions of the present disclosure may be assembled into kits. In some embodiments, the kit comprises nucleic acid vectors for the expression of the nucleobase editors described herein. In some embodiments, the kit further comprises appropriate guide nucleotide sequences (e.g., gRNAs) or nucleic acid vectors for the expression of such guide nucleotide sequences, to target the nucleobase editors to the desired target sequences.

(219) The kit described herein may include one or more containers housing components for performing the methods described herein and optionally instructions of uses. Any of the kit described herein may further comprise components needed for performing the assay methods. Each component of the kits, where applicable, may be provided in liquid form (e.g., in solution), or in solid form, (e.g., a dry powder). In certain cases, some of the components may be reconstitutable or otherwise processible (e.g., to an active form), for example, by the addition of a suitable solvent or other species (for example, water or certain organic solvents), which may or may not be provided with the kit.

(220) In some embodiments, the kits may optionally include instructions and/or promotion for use of the components

As used herein, “instructions” can define a component of instruction and/or promotion, and typically involve written instructions on or associated with packaging of the disclosure. Instructions also can include any oral or electronic instructions provided in any manner such that a user will clearly recognize that the instructions are to be associated with the kit, for example, audiovisual (e.g., videotape, DVD, etc.), Internet, and/or web-based communications, etc. The written instructions may be in a form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals or biological products, which can also reflect approval by the agency of manufacture, use or sale for animal administration. As used herein, “promoted” includes all methods of doing business including methods of education, hospital and other clinical instruction, scientific inquiry, drug discovery or development, academic research, pharmaceutical industry activity including pharmaceutical sales, and any advertising or other promotional activity including written, oral and electronic communication of any form, associated with the disclosure. Additionally, the kits may include other components depending on the specific application, as described herein.

(221) The kits may contain any one or more of the components described herein in one or more containers. The components may be prepared sterilely, packaged in a syringe and shipped refrigerated. Alternatively it may be housed in a vial or other container for storage. A second container may have other components prepared sterilely. Alternatively the kits may include the active agents premixed and shipped in a vial, tube, or other container.

(222) The kits may have a variety of forms, such as a blister pouch, a shrink wrapped pouch, a vacuum sealable pouch, a sealable thermoformed tray, or a similar pouch or tray form, with the accessories loosely packed within the pouch, one or more tubes, containers, a box or a bag. The kits may be sterilized after the accessories are added, thereby allowing the individual accessories in the container to be otherwise unwrapped. The kits can be sterilized using any appropriate sterilization techniques, such as radiation sterilization, heat sterilization, or other sterilization methods known in the art. The kits may also include other components, depending on the specific application, for example, containers, cell media, salts, buffers, reagents, syringes, needles, a fabric, such as gauze, for applying or removing a disinfecting agent, disposable gloves, a support for the agents prior to administration, etc.

(223) Therapeutics

(224) The compositions or cancer vaccines (e.g., a whole-cell vaccine comprising a modified tumor cell) described herein may be administered to a subject in need thereof, in a therapeutically effective amount, to treat cancer. The compositions and cancer vaccines described herein induce tumor-specific adaptive responses. It is known that cancer cells exploit immune checkpoints to evade immune surveillance. Thus, in some embodiments, in addition to the compositions or cancer vaccines described herein, agents that modulate the activities of immune checkpoints are also administered to boost the tumor-specific immune response elicited by the cancer vaccines.

(225) In some embodiments, the agents that modulate the activities of immune checkpoints are immune checkpoint inhibitors. “Immune checkpoints” are proteins in the immune system that either enhance an immune response signal (co-stimulatory molecules) or reduce an immune response signal. Many cancers protect themselves from the immune system by exploiting the inhibitory immune checkpoint proteins to inhibit the T cell signal. Such inhibitory checkpoint proteins include, without limitation, Cytotoxic T-Lymphocyte-Associated protein 4 (CTLA-4), Programmed Death 1 receptor (PD-1), T-cell Immunoglobulin domain and Mucin domain 3 (TIM3), Lymphocyte Activation Gene-3 (LAG3), V-set domain-containing T-cell activation inhibitor 1 (VTVN1 or B7-H4), Cluster of Differentiation 276 (CD276 or B7-H3), B and T Lymphocyte Attenuator (BTLA), Galectin-9 (GAL9), Checkpoint kinase 1 (Chk1), Adenosine A2A receptor (A2aR), Indoleamine 2,3-dioxygenase (IDO), Killer-cell Immunoglobulin-like Receptor (KIR), Lymphocyte Activation Gene-3 (LAG3), and V-domain Ig suppressor of T cell activation (VISTA).

(226) Some of these immune checkpoint proteins need their cognate binding partners, or ligands, for their immune inhibitory activity. For example, A2AR is the receptor of adenosine A2A and binding of A2A to A2AR activates a negative immune feedback loop. As another example, PD-1 associates with its two ligands, PD-L1 and PD-L2, to down regulate the immune system by preventing the activation of T-cells. PD-1 promotes the programmed cell death of antigen specific T-cells in lymph nodes and simultaneously reduces programmed cell death of suppressor T cells, thus achieving its immune inhibitory function. As yet another example, CTLA4 is present on the surface of T cells, and when bound to its binding partner CD80 or CD86 on the surface of antigen-present cells (APCs), it transmits an inhibitory signal to T cells, thereby reducing the immune response.

(227) Cancer cells are known to exploit the immune checkpoint proteins to escape being attacked by the immune system. Therefore, the use of immune checkpoint inhibitors to enhance an immune response against cancer, and thus treating cancer, have been described. The immunotherapeutic agents in the compositions of the present disclosure may also be immune checkpoint inhibitors. In some embodiments, the immune checkpoint inhibits any one or more of Cytotoxic T-Lymphocyte-Associated protein 4 (CTLA-4), Programmed Death 1 receptor (PD-1), T-cell Immunoglobulin domain and Mucin domain 3 (TIM3), Lymphocyte Activation Gene-3 (LAG3), V-set domain-containing T-cell activation inhibitor 1 (VTVN1 or B7-H4), Cluster of Differentiation 276 (CD276 or B7-H3), B and T Lymphocyte Attenuator (BTLA), Galectin-9 (GAL9), Checkpoint kinase 1 (Chk1), Adenosine A2A receptor (A2aR), Indoleamine 2,3-dioxygenase (IDO), Killer-cell Immunoglobulin-like Receptor (KIR), Lymphocyte Activation Gene-3 (LAG3) and V-domain Ig suppressor of T cell activation (VISTA).

(228) An “immune checkpoint inhibitor” is a molecule that prevents or weakens the activity of an immune checkpoint inhibitor. For example, an immune checkpoint inhibitor may inhibit the binding of the immune checkpoint protein to its cognate binding partner, e.g., PD-1, CTLA-4, or A2aR. In some embodiments, the immune checkpoint inhibitor is a small



molecule. In some embodiments, the immune checkpoint inhibitors is a nucleic acid aptamer (e.g., a siRNA targeting any one of the immune checkpoint proteins). In some embodiments, the immune checkpoint inhibitor is a recombinant protein. In some embodiments, the immune checkpoint inhibitor is an antibody. In some embodiments, the antibody comprises an anti-CTLA-4, anti-PD-1, anti-PD-L1, anti-TIM3, anti-LAG3, anti-B7-H3, anti-B7-H4, anti-BTLA, anti-GAL9, anti-Chk, anti-A2aR, anti-IDO, anti-KIR, anti-LAG3, anti-VISTA antibody, or a combination of any two or more of the foregoing antibodies. In some embodiments, the immune checkpoint inhibitor is a monoclonal antibody. In some embodiments, the immune checkpoint inhibitor comprises anti-PD1, anti-PD-L1, anti-CTLA-4, or a combination of any two or more of the foregoing antibodies. For example, the anti-PD-1 antibody is pembrolizumab (Keytruda®) or nivolumab (Opdivo®) and the anti-CTLA-4 antibody is ipilimumab (Yervoy®). Thus, in some embodiments, the immune checkpoint inhibitor comprises pembrolizumab, nivolumab, ipilimumab, or any combination of two or more of the foregoing antibodies. The examples described herein are not meant to be limiting and that any immune checkpoint inhibitors known in the art and any combinations thereof may be used in accordance with the present disclosure.

(229) In some embodiments, the immune checkpoint may be inhibited by disrupting any one of the immune checkpoint genes (e.g., CTLA-4, PD-1, PD-L1, TIM3, LAG3, B7-H3, B7-H4, BTLA, GAL9, Chk1, or A2aR) using any of the gene editing methods known in the art (e.g., CRISPR/Cas9 mediated cleavage of any of the genes).

(230) In some embodiments, an adjuvant is further administered to the subject. Adjuvants are substances which enhance the immune response when administered together with an immunogen or antigen. Adjuvants are thought to function in several ways, including by increasing the surface area of antigen, prolonging the retention of the antigen in the body thus allowing time for the lymphoid system to have access to the antigen, slowing the release of antigen, targeting antigen to macrophages, activating macrophages, or otherwise eliciting non-specific activation of the cells of the immune system see, e.g., H. S. Warren et al, *Annu. Rev. immunol.*, 4:369 (1986). Currently, an essential role of adjuvants in vaccines is to direct CD4+ T cell subset differentiation, although how adjuvants perform this function is poorly understood.

(231) The ability of a adjuvant to induce and increase a specific type of immune response and the identification of that ability is thus a key factor in the selection of particular adjuvants for vaccine use against a particular pathogen. Typical adjuvants include water and oil emulsions, e.g., Freund's adjuvant, and chemical compounds such as aluminum hydroxide or alum. At present, alum is the only adjuvant approved in the United States for human vaccines; it has been determined that alum induces the production of TH 2 cells.

(232) Many of the most effective adjuvants include bacteria or their products, e.g., microorganisms such as the attenuated strain of *Mycobacterium bovis*, *bacillus Calmette-Guerin* (BCG); microorganism components, e.g., alum-precipitated diphtheria toxoid, bacterial lipopolysaccharide and endotoxins. However, the role that the bacteria play is ill-defined. Recently, it has been noted that many bacteria or their products, lipopolysaccharide, *Staphylococcus aureus*, *Mycobacterium tuberculosis*, and *C. parvum*, stimulate IL-12 production by macrophages A. D'Andrea et al, *J. Exp. Med.*, 176:1387 (1992).

(233) Cancers or tumors include but are not limited to neoplasms, malignant tumors, metastases, or any disease or disorder characterized by uncontrolled cell growth such that it would be considered cancerous. The cancer may be a primary or metastatic cancer. Cancers include, but are not limited to, biliary tract cancer; bladder cancer; brain cancer including glioblastomas and medulloblastomas; breast cancer; cervical cancer; choriocarcinoma; colon cancer; endometrial cancer; esophageal cancer; gastric cancer; hematological neoplasms including acute lymphocytic and myelogenous leukemia; multiple myeloma; AIDS-associated leukemias and adult T-cell leukemia lymphoma; intraepithelial neoplasms including Bowen's disease and Paget's disease; liver cancer; lung cancer; lymphomas including Hodgkin's disease and lymphocytic lymphomas; neuroblastomas; oral cancer including squamous cell carcinoma; ovarian cancer including those arising from epithelial cells, stromal cells, germ cells and mesenchymal cells; pancreatic cancer; prostate cancer; rectal cancer; sarcomas including leiomyosarcoma, rhabdomyosarcoma, liposarcoma, fibrosarcoma, and osteosarcoma; skin cancer including melanoma, Kaposi's sarcoma, basocellular cancer, and squamous cell cancer; testicular cancer including germinal tumors such as seminoma, non-seminoma, teratomas, choriocarcinomas; stromal tumors and germ cell tumors; thyroid cancer including thyroid adenocarcinoma and medullar carcinoma; and renal cancer including adenocarcinoma and Wilms' tumor. Commonly encountered cancers include breast, prostate, lung, ovarian, colorectal, and brain cancer.

(234) "A therapeutically effective amount" as used herein refers to the amount of each base-editing agent of the present disclosure required to confer therapeutic effect on the subject, either alone or in combination with one or more other therapeutic agents. Effective amounts vary, as recognized by those skilled in the art, depending on the particular condition being treated, the severity of the condition, the individual subject parameters including age, physical condition, size, gender and weight, the duration of the treatment, the nature of concurrent therapy (if any), the specific route of administration and like factors within the knowledge and expertise of the health practitioner. These factors are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is generally preferred that a maximum dose of the individual components or combinations thereof be used, that is, the highest safe dose according to sound medical judgment. It will be understood by those of ordinary skill in the art, however, that a subject may insist upon a lower dose or tolerable dose for medical reasons, psychological reasons or for virtually any other reasons. Empirical considerations, such as the half-life, generally will contribute to the determination of the dosage. For example, therapeutic agents that are compatible with the human immune system, such as polypeptides comprising regions from humanized antibodies or fully human antibodies, may be used to prolong the half-life of the polypeptide.

(235) Frequency of administration may be determined and adjusted over the course of therapy, and is generally, but not necessarily, based on treatment and/or suppression and/or amelioration and/or delay of a disease. Alternatively, sustained continuous release formulations of a polypeptide or a polynucleotide may be appropriate. Various formulations and devices for achieving sustained release are known in the art. In some embodiments, dosage is daily, every other day, every three days, every four days, every five days, or every six days. In some embodiments, dosing frequency is once every week, every 2 weeks, every 4 weeks, every 5 weeks, every 6 weeks, every 7 weeks, every 8 weeks, every 9 weeks, or every 10 weeks; or once every month, every 2 months, or every 3 months, or longer. The progress of this therapy is easily monitored by conventional techniques and assays.

(236) The dosing regimen (including the polypeptide used) can vary over time. In some embodiments, for an adult subject of normal weight, doses ranging from about 0.01 to 1000 mg/kg may be administered. In some embodiments, the dose is between 1 to 200 mg. The particular dosage regimen, i.e., dose, timing and repetition, will depend on the particular subject and that subject's medical history, as well as the properties of the polypeptide or the polynucleotide (such as the half-life of the polypeptide or the polynucleotide, and other considerations well known in the art).

(237) For the purpose of the present disclosure, the appropriate dosage of a therapeutic agent as described herein will depend on the specific agent (or compositions thereof) employed, the formulation and route of administration, the type and severity of the disease, whether the polypeptide or the polynucleotide is administered for preventive or therapeutic purposes, previous therapy, the subject's clinical history and response to the antagonist, and the discretion of the attending physician. Typically the clinician will administer a polypeptide until a dosage is reached that achieves the desired result.

(238) Administration of one or more agents can be continuous or intermittent, depending, for example, upon the recipient's physiological condition, whether the purpose of the administration is therapeutic or prophylactic, and other factors known to skilled practitioners. The administration of an agent (e.g., cancer vaccine) may be essentially continuous over a preselected period of time or may be in a series of spaced dose, e.g., either before, during, or after developing a disease. "Treat," as used herein, means to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the disease, the symptom of the disease, or the predisposition toward the disease. "Treating," as used herein refers to the application or administration of a composition or a cancer vaccine described herein to a subject in need thereof.

(239) "A subject in need thereof", refers to an individual who has a disease, a symptom of the disease, or a predisposition toward the disease. In some embodiments, the subject is a mammal. In some embodiments, the subject is a non-human primate. In some embodiments, the subject is human. Alleviating a disease includes delaying the development or progression of the disease, or reducing disease severity. Alleviating the disease does not necessarily require curative results.

(240) As used therein, "delaying" the development of a disease means to defer, hinder, slow, retard, stabilize, and/or postpone progression of the disease. This delay can be of varying lengths of time, depending on the history of the disease and/or individuals being treated. A method that "delays" or alleviates the development of a disease, or delays the onset of the disease, is a method that reduces probability of developing one or more symptoms of the disease in a given time frame and/or reduces extent of the symptoms in a given time frame, when compared to not using the method. Such comparisons are typically based on clinical studies, using a number of subjects sufficient to give a statistically significant result.

(241) "Development" or "progression" of a disease means initial manifestations and/or ensuing progression of the disease. Development of the disease can be detectable and assessed using standard clinical techniques as well known in the art. However, development also refers to progression that may be undetectable. For purpose of this disclosure, development or progression refers to the biological course of the symptoms. "Development" includes occurrence, recurrence, and onset.

(242) As used herein "onset" or "occurrence" of a disease includes initial onset and/or recurrence. Conventional methods, known to those of ordinary skill in the art of medicine, can be used to administer the isolated polypeptide or pharmaceutical composition to the subject, depending upon the type of disease to be treated or the site of the disease. This composition can also be administered via other conventional routes, e.g., administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir.

(243) The term "parenteral" as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, intraarterial, intrasynovial, intrasternal, intrathecal, intralesional, and intracranial injection or infusion techniques. In addition, it can be administered to the subject via injectable depot routes of administration such as using 1-, 3-, or 6-month depot injectable or biodegradable materials and methods.

(244) Without further elaboration, it is believed that one skilled in the art can, based on the above description, utilize the present disclosure to its fullest extent. The following specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever. All publications cited herein are incorporated by reference for the purposes or subject matter referenced herein.

## EXAMPLES

(245) In order that the disclosure described herein may be more fully understood, the following examples are set forth. The synthetic examples described in this application are offered to illustrate the compounds and methods provided herein and are not to be construed in any way as limiting their scope.

Example 1: Guide Nucleotide Sequence-Programmable DNA-Binding Protein Domains, Deaminases, and Base Editors

(246) Non-limiting examples of suitable guide nucleotide sequence-programmable DNA-binding protein domains are provided. The disclosure provides Cas9 variants, for example, Cas9 proteins from one or more organisms, which may comprise one or more mutations (e.g., to generate dCas9 or Cas9 nickase). In some embodiments, one or more of the

amino acid residues, identified below by an asterisk, of a Cas9 protein may be mutated. In some embodiments, the D10 and/or H840 residues of the amino acid sequence provided in SEQ ID NO: 1, or a corresponding mutation in any of the amino acid sequences provided in SEQ ID NOs: 11-260, are mutated. In some embodiments, the D10 residue of the amino acid sequence provided in SEQ ID NO: 1, or a corresponding mutation in any of the amino acid sequences provided in SEQ ID NOs: 11-260, is mutated to any amino acid residue, except for D. In some embodiments, the D10 residue of the amino acid sequence provided in SEQ ID NO: 1, or a corresponding mutation in any of the amino acid sequences provided in SEQ ID NOs: 11-260, is mutated to an A. In some embodiments, the H840 residue of the amino acid sequence provided in SEQ ID NO: 1, or a corresponding residue in any of the amino acid sequences provided in SEQ ID NOs: 11-260, is an H. In some embodiments, the H840 residue of the amino acid sequence provided in SEQ ID NO: 1, or a corresponding mutation in any of the amino acid sequences provided in SEQ ID NOs: 11-260, is mutated to any amino acid residue, except for H. In some embodiments, the H840 residue of the amino acid sequence provided in SEQ ID NO: 1, or a corresponding mutation in any of the amino acid sequences provided in SEQ ID NOs: 11-260, is mutated to an A. In some embodiments, the D10 residue of the amino acid sequence provided in SEQ ID NO: 1, or a corresponding residue in any of the amino acid sequences provided in SEQ ID NOs: 11-260, is a D.

(247) A number of Cas9 sequences from various species were aligned to determine whether corresponding homologous amino acid residues of D10 and H840 of SEQ ID NO: 1 or SEQ ID NO: 11 can be identified in other Cas9 proteins, allowing the generation of Cas9 variants with corresponding mutations of the homologous amino acid residues. The alignment was carried out using the NCBI Constraint-based Multiple Alignment Tool (COBALT (accessible at [st-va.ncbi.nlm.nih.gov/tools/cobalt](http://st-va.ncbi.nlm.nih.gov/tools/cobalt)), with the following parameters. Alignment parameters: Gap penalties -11, -1; End-Gap penalties -5, -1. CDD Parameters: Use RPS BLAST on; Blast E-value 0.003; Find Conserved columns and Recompute on. Query Clustering Parameters: Use query clusters on; Word Size 4; Max cluster distance 0.8; Alphabet Regular.

(248) An exemplary alignment of four Cas9 sequences is provided below. The Cas9 sequences in the alignment are: Sequence 1 (S1): SEQ ID NO: 11 WP\_0109222511 gi 4992247111 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*]; Sequence 2 (S2): SEQ ID NO: 12|WP\_039695303|gi 746743737|type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus gallolyticus*]; Sequence 3 (S3): SEQ ID NO: 13|WP\_045635197|gi 782887988|type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus mitis*]; Sequence 4 (S4): SEQ ID NO: 14|5AXW\_A|gi 924443546| *Staphylococcus Aureus* Cas9. The HNH domain (bold and underlined) and the RuvC domain (boxed) are identified for each of the four sequences. Amino acid residues 10 and 840 in S1 and the homologous amino acids in the aligned sequences are identified with an asterisk following the respective amino acid residue.

(249) TABLE-US-00026 S1 1 --MDKK-YSIGLD\*IGTNSVGWAVITDEYKVPSSKKFKVLGNTDRHSIKKNLI--  
GALLFDSG--ETAETRLKRTARRRYT 73 S2 1 --  
MTKKNYSIGLD\*IGTNSVGWAVITDDYKVPKKMKVLGNTDKKYIKKNLL--GALLFDSG--  
ETAETRLKRTARRRYT 74 S3 1 --M-KKGYSIGLD\*IGTNSVGFAVITDDYKVPSSKKMKVLGNTDKRFIKKNLI--  
GALLFDEG--TTAEARLKRRTARRRYT 73 S4 1 GSHMKRNYILGLD\*IGITSVGYGII--DYET-----  
RDVIDAGVRLFKEANVENNEGRRSKRGARRLKR 61 S1 74  
RRKNRICYLQEIFSNEMAKVDDSSFFHRLSEESFLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLRKKLVDSTDKADLRL  
153 S2 75  
RRKNRLRYLQEIFANEIAKVDESFFQRLDESFLTDDDKTFDSHPIFGNKAEEDAYHQQFPTIYHLRKHLADSSEKADLRL  
154 S3 74  
RRKNRLRYLQEIFSEEMSKVDSSFFHRLDDSLIPEDKRESKYPIFATLTEEKEYHKQFPTIYHLRKLQADSKEKTDLRL  
153 S4 62 RRRHRIQRVKKLL-----FDYNLLTD-----HSELGINPYEARVKGLSQKLSEEE 107 S1  
154  
IYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDAKAILSARLSKSRRLLENLIAQLPGEK  
233 S2 155  
VYLALAHMIKFRGHFLIEGELNAENTDVQKIFADFGVYNRTFDDSHLSEITVDVASILTEKISKSRRLLENLIKYYPTK  
234 S3 154  
IYLALAHMIKYRGHFLYEEAFDIKNNDIQKIFNEFISIYDNTFEGSSLSGQNAQVEAIFTDKISKSAKRERVLKLPDEK  
233 S4 108 FSAALLHLAKRRG-----VHNVNEVEEDT----- 131 S1 234  
KNGLFGNLIASLGLTPNFKSNFDLAEDAKLQLSKDITYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEIT  
313 S2 235  
KNTLFGNLIALLGLQPNFKTNFKLSEDAKLQFSKDTYEEDLEELLGKIGDDYADLFTSAKNLYDAILLSGILTVDDNST  
314 S3 234  
STGLFSEFLKLIVGNQADFKKHFDLEDKAPLQFSKDTYDEDLENLLGQIGDDFTDLFVSAKKLYDAILLSGILTVTDPST  
313 S4 132 ----GNELS-----TKEQISR----- 144 S1 314  
KAPLSASMIKRYDEHHQDLTLLKALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKM--  
DGTEELLV 391 S2 315  
KAPLSASMIKRYVEHHEDLEKLKEFIKANKSELYHDIFKDKNKNKYAGYIENGVKQDEFYKYLKNILSKIKIDGSDYFLD  
394 S3 314  
KAPLSASMIERYENHQNDLAALKQFIKNNLPEKYDEVFSDQSKDGYAGYIDGKTTQETFYKYIKNLLSKF--  
EGTDYFLD 391 S4 145 ----SKALEEKYVAELQ-----LERLKKDG----- 165 S1 392  
KLNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFYFPFLKDNREKIEKILTFRIPYVVGPLARGNSRFAWMTRKSEE

471 S2 395  
KIEREDFLRKQRTFDNGSIPHQIHLQEMHAILRRQG DYYPFLKEKQDRIEKILTRIPYYVGPLVRKDSRFAWAEYRSDE  
474 S3 392  
KIEREDFLRKQRTFDNGSIPHQIHLQEMNAILRRQGEYYPFLKDNKEKIEKILTRIPYYVGPLARGNRDFAWLTRNSDE  
471 S4 166 --EVRGSINRFKTS-----YVKEAKQLLKVQKAYHQLDQSFIDTYIDLLETRRTYYEGP--GEGSPFGW-  
----K 227 S1 472  
TITPWNFEVVDKGASAQSFIERMTNFDKNLPNEKVLPHKSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVD  
551 S2 475 KITPWNFDKVIDKEKSAEKFITRMTLNDLYLPEEKVLPKHS HVYETYAVYNELTKIKYVNEQGKE-  
SFFDSNMKQEIFDH 553 S3 472  
AIRPWNFEIIVDKASSAEDFINKMTNYDLYLPEEKVLPKHSLLYETFAVYNELTKVKFIAEGLRDYQFLDSGQKKQIVNQ  
551 S4 228 DIKEW-----YEMLMGHCTYFPEELRSVKYAYNADLYNALNDLNNLVITRDENEK---  
LEYEYEFQIEN 289 S1 552 LFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDR---  
FNASLGTYHDLKKIHKDKDFLDNEENEDILEDIVLTTLTFED 628 S2 554  
VFKENRKVTKEKLLNYLNKEFPEYRIKDLIGLDKENKSFNASLGTYHDLKKIL-  
DKAFLDDKVNEEVIEDIHKTLTLTFED 632 S3 552 LFKENRKVTEKDIIHYLHN-VDGYDGIELKGIEKQ---  
FNASLSTYHDLKKIHKDKDFMDDAKNEAILENIVHTLTIFED 627 S4 290  
VFKQKKKPTLKQIAKEILVNEEDIKGYRVTSTGKPEF---TNLKVYHDIKDITARKEII---ENAELLDQIAKILTIYQS  
363 S1 629 REMIEERLKYAHLFDDKVMKQLKR-  
RRTYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRNFMLIHDDSLTFKED 707 S2 633  
KDMIHERLQKYS DIFTANQLKKLER-  
RHYTGWGRLSYKLINGIRNKENKNTILDY LIDDGSANRNFMLINDDTL PFKQI 711 S3 628  
REMIKQRLAQYDSL FDEKVIKALTR-  
RHYTGWGLSAKLINGICDKQTGNTILDY LIDDGKINRNFMLINDDGLSFKEI 706 S4 364  
SEDIQEELTNLSEL TQEEIEQISNLKGYTGTHNLSLKAINLILDE-----LWHTNDNQIAIFNRLKLVP----- 428  
S1 708 IQKAQVSGQGcustom character RENQTT-----QKGQKNSRERM 781 S2 712 IQKSQVVGDV  
custom character RENQTT-----NRGRSQQQRL 784 S3 707 IQKAQVIGKTcustom character RENQTT-----  
ARGKKNSQQRY 779 S4 429 -KKVDLSQQKcustom character REKNSKDAQKMINEMQKRN RQTN 505 S1  
782 KRIEEGIKELGSQIL-----KEHPVENTQLQNEKLYLYLQNGRDMYVDQELDINRLSD---  
YDVDH\*IVPQSFLKDD 850 S2 785  
KKLQNSLKELGSNILNEEKPSYIEDKVENSHLQNDQLFLYYIQNGKDMYTGDDEL DIDHLSD---  
YDIDH\*IIPQAFIKDD 860 S3 780 KRIEDSLKILASGL---  
DSNILKENPTDNNQLQNDRLFYYLQNGKDMYTGEALDINQLSS---YDIDH\*IIPQAFIKDD 852 S4 506  
ERIEEIIRTTGK-----ENAKYLIEKIKLHDMQEGKCLYSLEAIPLEDLLNPNFNYEVDH\*IIPRSVSFDN  
570 S1 851 SIDNKVLTRSDKNRGKSDNVPSEE VVKMKNYWRQLLNAKLITQRKFEDN-LTKAERG  
custom character 922 S2 861 SIDNRVLTSSAKNRGKSDDVPSLDIVRARKAEWVRLYKSGLISKRKFDN-  
LTKAERG custom character 932 S3 853  
SLDNRVLTSSKDNRGKSDNVP SIEVVQKRKA FWQQLLSKLISERKENN-LTKAERG custom character 924  
S4 571 SFNNKVLVKQEEASKKGNRTPFQYLSSSDSKISYETFEKKHILNLA KGKGRISKTKKE  
custom character 650 S1 923 custom character 1002 S2 933 custom character 1012 S3 925 custom character  
1004 S4 651 custom character 712 S1 1003 custom character 1077 S2 1013 custom character 1083 S3 1005  
custom character 1081 S4 713 custom character 764 S1 1078 custom character  
GGFSKESILPKRNSDKLIARKD---WDPKKYGGFDSPTVAYSVLV VAKV 1149 S2 1084 custom character  
GGFSKESILPKGDSDKLIPRKTKKVYWDTKKYGGFDSPTVAYSVFV VADV 1158 S3 1082 custom character  
GGFSKESILPKGNSDKLIPRKTKDILLDTTKYGGFDSPIAYSILLIADI 1156 S4 765 custom character  
RKDDKGNTLIVNNLNGLYDKDNDKL---KKLIN-KSP---EKLLMYHH 835 S1 1078  
EKGKSKKLKSVKELLGITIMERS SFEKNPI-DFLEAKG-----  
YKEVKKD LIKLPKYS LFELENGRRKRMLASAGELQKG 1223 S2 1084  
EKGKAKKLKTVKELVGISIMERS SFEENPV-EFLENKG-----  
YHNIREDKLIKLPKYS LFEFEGGRRRL LASASELQKG 1232 S3 1157  
EKGKAKKLKTVKTLVGITIMEKA AFEENPI-TFLENKG-----  
YHNVRKENILCLPKYS LFELENGRRRL LASAKELQKG 1230 S4 836 DPQTYQKLK-----  
LIMEQYGDEKNPLYKY YEETGNYLTKYSKKDNGPVIKKIKYYGNKLN AHL DITDDYPNSRNKV 907 S1 1224  
NELALPSKYVNFLYLASHYEKLKGS PEDNEQKQLFVEQHKHYLDEIIEQISEFSKRVLADANLDKVL SAYNKH---  
--- 1297 S2 1233 NEMVLPGYLVELLYHAHRADNF-----  
NSTEYLN YVSEHKKEFEKVLSCVEDFANLYVDVEKNLSKIRAVADSM----- 1301 S3 1231  
NEIVLPVYLTLLYH SKNVHKL-----DEPGHLEYIQKHRNEFKDLLNLVSEFSQKYVLADANLEKIKSLYADN-----  
1299 S4 908 VKLSLKPYRFD-VYLDNGVYKFV----TVKNLDVIK--  
KENYYEVNSKAYEEAKLKKISNQA EFIASFYNNDLIKING 979 S1 1298  
RDKPIREQAENIIHFLTTLNLGAPAAFKYFDTTIDRKRYTSTKEVL DATLIHQ SIT--GLYETRI ---- DLSQL 1365  
S2 1302 DNFSIEEISNSFINLLT LTALGAPADNFLGEKIPRKRYTSTKECLNATLIHQ SIT-GLYETRIDL SKL  
1369 S3 1300 EQADIEILANSFINLLT FTALGAPAAFKFFGKDIDRKRYTTVSEILNATLIHQ SIT-GLYETWI 1369

S4 980 ELYR91GVNNDIRVNNDITYR-EYLENMDNKRPPRIIKTIASKT ---  
QSIKKYSTDILGNLYEVKSKKHPQIIKK 1055 S1 1366 GGD 1368 S2 1370 GEE 1372 S3 1368 GED 1370 S4 4056 G-  
- 1056

(250) The alignment demonstrates that amino acid sequences and amino acid residues that are homologous to a reference Cas9 amino acid sequence or amino acid residue can be identified across Cas9 sequence variants, including, but not limited to Cas9 sequences from different species, by identifying the amino acid sequence or residue that aligns with the reference sequence or the reference residue using alignment programs and algorithms known in the art. This disclosure provides Cas9 variants in which one or more of the amino acid residues identified by an asterisk in SEQ ID NOs: 11-14 (e.g., S1, S2, S3, and S4, respectively) are mutated as described herein. The residues D10 and H840 in Cas9 of SEQ ID NO: 1 that correspond to the residues identified in SEQ ID NOs: 11-14 by an asterisk are referred to herein as “homologous” or “corresponding” residues. Such homologous residues can be identified by sequence alignment, e.g., as described above, and by identifying the sequence or residue that aligns with the reference sequence or residue. Similarly, mutations in Cas9 sequences that correspond to mutations identified in SEQ ID NO: 1 herein, e.g., mutations of residues 10, and 840 in SEQ ID NO: 1, are referred to herein as “homologous” or “corresponding” mutations. For example, the mutations corresponding to the D10A mutation in SEQ ID NO: 1 or S1 (SEQ ID NO: 11) for the four aligned sequences above are D11A for S2, D10A for S3, and D13A for S4; the corresponding mutations for H840A in SEQ ID NO: 1 or S1 (SEQ ID NO: 11) are H850A for S2, H842A for S3, and H560A for S4.

(251) A total of 250 Cas9 sequences (SEQ ID NOs: 11-260) from different species are provided. Amino acid residues homologous to residues 10, and 840 of SEQ ID NO: 1 may be identified in the same manner as outlined above. All of these Cas9 sequences may be used in accordance with the present disclosure.

(252) TABLE-US-00027 WP\_039695303.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus gallolyticus*] SEQ ID NO: 12 WP\_045635197.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus mitis*] SEQ ID NO: 13 5AXW A Cas9, Chain A, Crystal Structure [*Staphylococcus Aureus*] SEQ ID NO: 14 WP\_009880683.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 15 WP\_010922251.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 16 WP\_011054416.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 17 WP\_011284745.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 18 WP\_011285506.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 19 WP\_011527619.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 20 WP\_012560673.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 21 WP\_014407541.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 22 WP\_020905136.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 23 WP\_023080005.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 24 WP\_023610282.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 25 WP\_030125963.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 26 WP\_030126706.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 27 WP\_031488318.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 28 WP\_032460140.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 29 WP\_032461047.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 30 WP\_032462016.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 31 WP\_032462936.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 32 WP\_032464890.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 33 WP\_033888930.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 34 WP\_038431314.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 35 WP\_038432938.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 36 WP\_038434062.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus pyogenes*] SEQ ID NO: 37 BAQ51233.1 CRISPR-associated protein, CsnI family [*Streptococcus pyogenes*] SEQ ID NO: 38 KGE60162.1 hypothetical protein MGAS2111\_0903 [*Streptococcus pyogenes* MGAS2111] SEQ ID NO: 39 KGE60856.1 CRISPR-associated endonuclease protein [*Streptococcus pyogenes* SS1447] SEQ ID NO: 40 WP\_002989955.1 MULTISPECIES: type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus*] SEQ ID NO: 41 WP\_003030002.1 MULTISPECIES: type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus*] SEQ ID NO: 42 WP\_003065552.1 MULTISPECIES: type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus*] SEQ ID NO: 43 WP\_001040076.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 44 WP\_001040078.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 45 WP\_001040080.1 type

II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 46  
 WP\_001040081.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*]  
 SEQ ID NO: 47 WP\_001040083.1 type II CRISPR RNA-guided endonuclease Cas9  
 [*Streptococcus agalactiae*] SEQ ID NO: 48 WP\_001040085.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 49 WP\_001040087.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 50  
 WP\_001040088.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*]  
 SEQ ID NO: 51 WP\_001040089.1 type II CRISPR RNA-guided endonuclease Cas9  
 [*Streptococcus agalactiae*] SEQ ID NO: 52 WP\_001040090.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 53 WP\_001040091.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 54  
 WP\_001040092.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*]  
 SEQ ID NO: 55 WP\_001040094.1 type II CRISPR RNA-guided endonuclease Cas9  
 [*Streptococcus agalactiae*] SEQ ID NO: 56 WP\_001040095.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 57 WP\_001040096.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 58  
 WP\_001040097.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*]  
 SEQ ID NO: 59 WP\_001040098.1 type II CRISPR RNA-guided endonuclease Cas9  
 [*Streptococcus agalactiae*] SEQ ID NO: 60 WP\_001040099.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 61 WP\_001040100.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 62  
 WP\_001040104.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*]  
 SEQ ID NO: 63 WP\_001040105.1 type II CRISPR RNA-guided endonuclease Cas9  
 [*Streptococcus agalactiae*] SEQ ID NO: 64 WP\_001040106.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 65 WP\_001040107.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 66  
 WP\_001040108.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*]  
 SEQ ID NO: 67 WP\_001040109.1 type II CRISPR RNA-guided endonuclease Cas9  
 [*Streptococcus agalactiae*] SEQ ID NO: 68 WP\_001040110.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 69 WP\_015058523.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 70  
 WP\_017643650.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*]  
 SEQ ID NO: 71 WP\_017647151.1 type II CRISPR RNA-guided endonuclease Cas9  
 [*Streptococcus agalactiae*] SEQ ID NO: 72 WP\_017648376.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 73 WP\_017649527.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 74  
 WP\_017771611.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*]  
 SEQ ID NO: 75 WP\_017771984.1 type II CRISPR RNA-guided endonuclease Cas9  
 [*Streptococcus agalactiae*] SEQ ID NO: 76 CFQ25032.1 CRISPR-associated protein  
 [*Streptococcus agalactiae*] SEQ ID NO: 77 CFV16040.1 CRISPR-associated protein  
 [*Streptococcus agalactiae*] SEQ ID NO: 78 KLJ37842.1 CRISPR-associated protein Csn1  
 [*Streptococcus agalactiae*] SEQ ID NO: 79 KLJ72361.1 CRISPR-associated protein Csn1  
 [*Streptococcus agalactiae*] SEQ ID NO: 80 KLL20707.1 CRISPR-associated protein Csn1  
 [*Streptococcus agalactiae*] SEQ ID NO: 81 KLL42645.1 CRISPR-associated protein Csn1  
 [*Streptococcus agalactiae*] SEQ ID NO: 82 WP\_047207273.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 83 WP\_047209694.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 84  
 WP\_050198062.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*]  
 SEQ ID NO: 85 WP\_050201642.1 type II CRISPR RNA-guided endonuclease Cas9  
 [*Streptococcus agalactiae*] SEQ ID NO: 86 WP\_050204027.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 87 WP\_050881965.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*] SEQ ID NO: 88  
 WP\_050886065.1 type II CRISPR RNA-guided endonuclease Cas9 [*Streptococcus agalactiae*]  
 SEQ ID NO: 89 AHN30376.1 CRISPR-associated protein Csn1 [*Streptococcus agalactiae* 138P]  
 SEQ ID NO: 90 EAO78426.1 reticulocyte binding protein [*Streptococcus agalactiae* H36B] SEQ  
 ID NO: 91 CCW42055.1 CRISPR-associated protein, SAG0894 family [*Streptococcus agalactiae*  
 ILRI112] SEQ ID NO: 92 WP\_003041502.1 type II CRISPR RNA-guided endonuclease Cas9  
 [*Streptococcus anginosus*] SEQ ID NO: 93 WP\_037593752.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [*Streptococcus anginosus*] SEQ ID NO: 94 WP\_049516684.1 CRISPR-  
 associated protein Csn1 [*Streptococcus anginosus*] SEQ ID NO: 95 GAD46167.1 hypothetical  
 protein ANG6\_0662 [*Streptococcus anginosus* T5] SEQ ID NO: 96 WP\_018363470.1 type II



CRISPR RNA-guided endonuclease Cas9 [Streptococcus caballi] SEQ ID NO: 97  
 WP\_003043819.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus canis] SEQ  
 ID NO: 98 WP\_006269658.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 constellatus] SEQ ID NO: 99 WP\_048800889.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus constellatus] SEQ ID NO: 100 WP\_012767106.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus dysgalactiae] SEQ ID NO: 101 WP\_014612333.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus dysgalactiae] SEQ ID NO: 102  
 WP\_015017095.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus dysgalactiae]  
 SEQ ID NO: 103 WP\_015057649.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus dysgalactiae] SEQ ID NO: 104 WP\_048327215.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus dysgalactiae] SEQ ID NO: 105 WP\_049519324.1 CRISPR-  
 associated protein Csn1 [Streptococcus dysgalactiae] SEQ ID NO: 106 WP\_012515931.1 type  
 II CRISPR RNA-guided endonuclease Cas9 [Streptococcus equi] SEQ ID NO: 107  
 WP\_021320964.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus equi] SEQ ID  
 NO: 108 WP\_037581760.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 equi] SEQ ID NO: 109 WP\_004232481.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus equinus] SEQ ID NO: 110 WP\_009854540.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus gallolyticus] SEQ ID NO: 111 WP\_012962174.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus gallolyticus] SEQ ID NO: 112  
 WP\_039695303.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus gallolyticus]  
 SEQ ID NO: 113 WP\_014334983.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus infantarius] SEQ ID NO: 114 WP\_003099269.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus iniae] SEQ ID NO: 115 AHY15608.1 CRISPR-associated  
 protein Csn1 [Streptococcus iniae] SEQ ID NO: 116 AHY17476.1 CRISPR-associated protein  
 Csn1 [Streptococcus iniae] SEQ ID NO: 117 ESR09100.1 hypothetical protein IUSA1\_08595  
 [Streptococcus iniae IUSA1] SEQ ID NO: 118 AGM98575.1 CRISPR-associated protein  
 Cas9/Csn1, subtype II/NMEMI [Streptococcus iniae SF1] SEQ ID NO: 119 ALF27331.1  
 CRISPR-associated protein Csn1 [Streptococcus intermedius] SEQ ID NO: 120 WP\_018372492.1  
 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus massiliensis] SEQ ID NO: 121  
 WP\_045618028.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus mitis] SEQ ID  
 NO: 122 WP\_045635197.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 mitis] SEQ ID NO: 123 WP\_002263549.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus mutans] SEQ ID NO: 124 WP\_002263887.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 125 WP\_002264920.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 126  
 WP\_002269043.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ  
 ID NO: 127 WP\_002269448.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 mutans] SEQ ID NO: 128 WP\_002271977.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus mutans] SEQ ID NO: 129 WP\_002272766.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 130 WP\_002273241.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 131  
 WP\_002275430.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ  
 ID NO: 132 WP\_002276448.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 mutans] SEQ ID NO: 133 WP\_002277050.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus mutans] SEQ ID NO: 134 WP\_002277364.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 135 WP\_002279025.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 136  
 WP\_002279859.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ  
 ID NO: 137 WP\_002280230.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 mutans] SEQ ID NO: 138 WP\_002281696.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus mutans] SEQ ID NO: 139 WP\_002282247.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 140 WP\_002282906.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 141  
 WP\_002283846.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ  
 ID NO: 142 WP\_002287255.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 mutans] SEQ ID NO: 143 WP\_002288990.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus mutans] SEQ ID NO: 144 WP\_002289641.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 145 WP\_002290427.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 146  
 WP\_002295753.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ  
 ID NO: 147 WP\_002296423.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus

SEQ ID NO: 148 WP\_002304487.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus mutans] SEQ ID NO: 149 WP\_002305844.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 150 WP\_002307203.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 151  
 WP\_002310390.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ  
 ID NO: 152 WP\_002352408.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 mutans] SEQ ID NO: 153 WP\_012997688.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus mutans] SEQ ID NO: 154 WP\_014677909.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 155 WP\_019312892.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 156  
 WP\_019313659.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ  
 ID NO: 157 WP\_019314093.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 mutans] SEQ ID NO: 158 WP\_019315370.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus mutans] SEQ ID NO: 159 WP\_019803776.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 160 WP\_019805234.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 161  
 WP\_024783594.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ  
 ID NO: 162 WP\_024784288.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 mutans] SEQ ID NO: 163 WP\_024784666.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus mutans] SEQ ID NO: 164 WP\_024784894.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 165 WP\_024786433.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus mutans] SEQ ID NO: 166  
 WP\_049473442.1 CRISPR-associated protein Csn1 [Streptococcus mutans] SEQ ID NO: 167  
 WP\_049474547.1 CRISPR-associated protein Csn1 [Streptococcus mutans] SEQ ID NO: 168  
 EMC03581.1 hypothetical protein SMU69\_09359 [Streptococcus mutans NLML4] SEQ ID NO: 169  
 WP\_000428612.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus oralis] SEQ  
 ID NO: 170 WP\_000428613.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 oralis] SEQ ID NO: 171 WP\_049523028.1 CRISPR-associated protein Csn1 [Streptococcus  
 parasanguinis] SEQ ID NO: 172 WP\_003107102.1 type II CRISPR RNA-guided endonuclease  
 Cas9 [Streptococcus parauberis] SEQ ID NO: 173 WP\_054279288.1 type II CRISPR RNA-  
 guided endonuclease Cas9 [Streptococcus phocae] SEQ ID NO: 174 WP\_049531101.1 CRISPR-  
 associated protein Csn1 [Streptococcus pseudopneumoniae] SEQ ID NO: 175 WP\_049538452.1  
 CRISPR-associated protein Csn1 [Streptococcus pseudopneumoniae] SEQ ID NO: 176  
 WP\_049549711.1 CRISPR-associated protein Csn1 [Streptococcus pseudopneumoniae] SEQ ID  
 NO: 177 WP\_007896501.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 pseudoporcinus] SEQ ID NO: 178 EFR44625.1 CRISPR-associated protein, Csn1 family  
 [Streptococcus pseudoporcinus SPIN 20026] SEQ ID NO: 179 WP\_002897477.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus sanguinis] SEQ ID NO: 180  
 WP\_002906454.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus sanguinis] SEQ  
 ID NO: 181 WP\_009729476.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus  
 sp. F0441] SEQ ID NO: 182 CQR24647.1 CRISPR-associated protein [Streptococcus sp.  
 FF10] SEQ ID NO: 183 WP\_000066813.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus sp. M334] SEQ ID NO: 184 WP\_009754323.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Streptococcus sp. taxon 056] SEQ ID NO: 185 WP\_044674937.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Streptococcus suis] SEQ ID NO: 186  
 WP\_044676715.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus suis] SEQ ID  
 NO: 187 WP\_044680361.1 type II CRISPR RNA-guided endonuclease Cas9 [Streptococcus suis]  
 SEQ ID NO: 188 WP\_044681799.1 type II CRISPR RNA-guided endonuclease Cas9  
 [Streptococcus suis] SEQ ID NO: 189 WP\_049533112.1 CRISPR-associated protein Csn1  
 [Streptococcus suis] SEQ ID NO: 190 WP\_029090905.1 type II CRISPR RNA-guided  
 endonuclease Cas9 [Brochothrix thermosphacta] SEQ ID NO: 191 WP\_006506696.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Catenibacterium mitsuokai] SEQ ID NO: 192  
 AIT42264.1 Cas9hc:NLS:A [Cloning vector pYB196] SEQ ID NO: 193 WP\_034440723.1 type II  
 CRISPR endonuclease Cas9 [Clostridiales bacterium S5-A11] SEQ ID NO: 194 AKQ21048.1  
 Cas9 [CRISPR-mediated gene targeting vector p (bhsp68-Cas9)] SEQ ID NO: 195  
 WP\_004636532.1 type II CRISPR RNA-guided endonuclease Cas9 [Dolosigranulum pigrum] SEQ  
 ID NO: 196 WP\_002364836.1 MULTISPECIES: type II CRISPR RNA-guided endonuclease Cas9  
 [Enterococcus] SEQ ID NO: 197 WP\_016631044.1 MULTISPECIES: type II CRISPR RNA-  
 guided endonuclease Cas9 [Enterococcus] SEQ ID NO: 198 EMS75795.1 hypothetical protein  
 H318\_06676 [Enterococcus durans IPLA 655] SEQ ID NO: 199 WP\_002373311.1 type II  
 CRISPR RNA-guided endonuclease Cas9 [Enterococcus faecalis] SEQ ID NO: 200



WP\_002378009.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecalis*] SEQ ID NO: 201 WP\_002407324.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecalis*] SEQ ID NO: 202 WP\_002413717.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecalis*] SEQ ID NO: 203 WP\_010775580.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecalis*] SEQ ID NO: 204 WP\_010818269.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecalis*] SEQ ID NO: 205 WP\_010824395.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecalis*] SEQ ID NO: 206 WP\_016622645.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecalis*] SEQ ID NO: 207 WP\_033624816.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecalis*] SEQ ID NO: 208 WP\_033625576.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecalis*] SEQ ID NO: 209 WP\_033789179.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecalis*] SEQ ID NO: 210 WP\_002310644.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecium*] SEQ ID NO: 211 WP\_002312694.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecium*] SEQ ID NO: 212 WP\_002314015.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecium*] SEQ ID NO: 213 WP\_002320716.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecium*] SEQ ID NO: 214 WP\_002330729.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecium*] SEQ ID NO: 215 WP\_002335161.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecium*] SEQ ID NO: 216 WP\_002345439.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecium*] SEQ ID NO: 217 WP\_034867970.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecium*] SEQ ID NO: 218 WP\_047937432.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus faecium*] SEQ ID NO: 219 WP\_010720994.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus hirae*] SEQ ID NO: 220 WP\_010737004.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus hirae*] SEQ ID NO: 221 WP\_034700478.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus hirae*] SEQ ID NO: 222 WP\_007209003.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus italicus*] SEQ ID NO: 223 WP\_023519017.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus mundtii*] SEQ ID NO: 224 WP\_010770040.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus phoeniculicola*] SEQ ID NO: 225 WP\_048604708.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus* sp. AM1] SEQ ID NO: 226 WP\_010750235.1 type II CRISPR RNA-guided endonuclease Cas9 [*Enterococcus villorum*] SEQ ID NO: 227 AII16583.1 Cas9 endonuclease [Expression vector pCas9] SEQ ID NO: 228 WP\_029073316.1 type II CRISPR RNA-guided endonuclease Cas9 [*Kandleria vitulina*] SEQ ID NO: 229 WP\_031589969.1 type II CRISPR RNA-guided endonuclease Cas9 [*Kandleria vitulina*] SEQ ID NO: 230 KDA45870.1 CRISPR-associated protein Cas9/Csn1, subtype II/NMEMI [*Lactobacillus animalis*] SEQ ID NO: 231 WP\_039099354.1 type II CRISPR RNA-guided endonuclease Cas9 [*Lactobacillus curvatus*] SEQ ID NO: 232 AKP02966.1 hypothetical protein ABB45\_04605 [*Lactobacillus farciminis*] SEQ ID NO: 233 WP\_010991369.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria innocua*] SEQ ID NO: 234 WP\_033838504.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria innocua*] SEQ ID NO: 235 EHN60060.1 CRISPR-associated protein, Csn1 family [*Listeria innocua* ATCC 33091] SEQ ID NO: 236 EFR89594.1 crispr-associated protein, Csn1 family [*Listeria innocua* FSL S4-378] SEQ ID NO: 237 WP\_038409211.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria ivanovii*] SEQ ID NO: 238 EFR95520.1 crispr-associated protein Csn1 [*Listeria ivanovii* FSL F6-596] SEQ ID NO: 239 WP\_003723650.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria monocytogenes*] SEQ ID NO: 240 WP\_003727705.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria monocytogenes*] SEQ ID NO: 241 WP\_003730785.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria monocytogenes*] SEQ ID NO: 242 WP\_003733029.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria monocytogenes*] SEQ ID NO: 243 WP\_003739838.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria monocytogenes*] SEQ ID NO: 244 WP\_014601172.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria monocytogenes*] SEQ ID NO: 245 WP\_023548323.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria monocytogenes*] SEQ ID NO: 246 WP\_031665337.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria monocytogenes*] SEQ ID NO: 247 WP\_031669209.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria monocytogenes*] SEQ ID NO: 248 WP\_033920898.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria monocytogenes*] SEQ ID NO: 249 AKI42028.1 CRISPR-associated protein [*Listeria monocytogenes*] SEQ ID NO: 250 AKI50529.1 CRISPR-associated protein [*Listeria monocytogenes*] SEQ ID NO: 251 EFR83390.1 crispr-associated protein Csn1 [*Listeria monocytogenes* FSL F2-208] SEQ ID NO: 252 WP\_046323366.1 type II CRISPR RNA-guided endonuclease Cas9 [*Listeria seeligeri*] SEQ ID NO: 253 AKE81011.1 Cas9 [Plant multiplex genome editing vector

pYLCRISPR/Cas9Pubi-H] SEQ ID NO: 254 CUO82355.1 Uncharacterized protein conserved in bacteria [*Roseburia hominis*] SEQ ID NO: 255 WP\_033162887.1 type II CRISPR RNA-guided endonuclease Cas9 [*Sharpea azabuensis*] SEQ ID NO: 256 AGZ01981.1 Cas9 endonuclease [synthetic construct] SEQ ID NO: 257 AKA60242.1 nuclease deficient Cas9 [synthetic construct] SEQ ID NO: 258 AKS40380.1 Cas9 [Synthetic plasmid pFC330] SEQ ID NO: 259 4UN5\_B Cas9, Chain B, Crystal Structure SEQ ID NO: 260  
(253) TABLE-US-00028 Non-limiting examples of suitable deaminase domains are provided. Human AID (SEQ ID NO: 303)

QGRPFQPDWGLDEHSQALSGRLRAI (italic: nucleic acid editing domain; underline: cytoplasmic localization signal) Human APOBEC-3G (SEQ ID NO: 279)  
*MKPHFRNTVERMYRDTESYNFYNRPILSRRNTVWLCYEVKTKGSPRPPLDAKIFRGQVYSELKYHPEMRFFHWFSKWRKLHRDQEYEVTWYISWSPCTKCTRDMAFLAEDPKVTLTIFVARLYYFWDPDYQEALRSLCQKRDGPRATMKIMNYDEFQHCWSKFVYSQRELFEWNNLPKYIILLHIMLGEILRHSMDDPPTFTFNFNNEPWVRGRHETYLCEYEVERMHNDTWVLLNQRRGFLCNQAPHKHGFLEGRHAELCFLDVIPFWKLDLDQDYRVTCFTSWSPCFSCAQEMAKFISKNHVSLCIFTARIYDDQGRCEGLRTLAEAGAKISIMTYSEFKHCWDTFVDHQGCPFQPWDGLDEHSQDLSGRLRAILQNQEN* (italic: nucleic acid editing domain; underline: cytoplasmic localization signal) Human APOBEC-3F (SEQ ID NO: 280)  
*MKPHFRNTVERMYRDTFSYNFYNRPILSRRNTVWLCYEVKTKGSPRPRLDAKIFRGQVYSQPEHHAEMCFLSWFCGNQLPAYKCFQITWVSWTPCPDCVAKLAEFLAHPNVTLTISAARLYYYWERDYRRALCRLSQA GARVTIMDYEEFAYCWENFVYSEGQPFMPWYKFDDNYAFLHRTLKEILRNPMEAMYPHIFYFHFKNLRKAYGRNESWLCFTMEVVKHHSPVSWKRGVFRNQVDPETHCHAERCFLSWFCDDILSPNTNYEVTWYTSWSPCECAGEVAEFLARHSNVNLTIFTARLYYFWDTDYQEGLRSLSQEGASVEIMGYKDFKYCWENFVYNDDEPFK PWKGLKYNFLFLDSKLQEILE* (italic: nucleic acid editing domain) Human APOBEC-3B (SEQ ID NO: 281)  
*MNPQIRNPMERMYRDTFYDNFENEPILYGRSYTWLCYEVKIKRGRSNLLWDTGVFRGQVYFKPQYHAEMCFLSWFCGNQLPAYKCFQITWVSWTPCPDCVAKLAEFLSEHPNVTLTISAARLYYYWERDYRRALCRLSQA GARVTIMDYEEFAYCWENFVYNEGQQFMPWYKFDDNYAFLHRTLKEILRYLMDPDTFTFNFNNDPL VLRR RQTYLCEYEVERLDNGTWVLMQHMGLFCNEAKNLLCGFYGRHAELRFLDLVPSLQLDPAQIYRVTFISWS PCFSWGCAGEVRAFLQENTHVRLRIFAARIYDYDPLYKEALQMLRDAGAQVSIMTYDEFEYCWDTFVYRQ GCPFQPWDGLEEHSQALSGRLRAILQNQGN* (italic: nucleic acid editing domain) Rat APOBEC-3B: (SEQ ID NO: 1073)  
*MQPQGLGPNAGMGPVCLGCSHRRPYSPIRNPLKKLYQQTFYFHFKNVRYAWGRKNNFLCYEVNGMDCAL PVPLRQGVFRKQGHIAELCFIYWFDKVLRLVLSMEEFKVTWYMSWSPCSKCAEQVARFLAAHRNLSLA IFSSRLYYYLRPNPNYQQKLCRLIQEGVHVAAMDLPFCKCWNKFVDNDGQPFPRPWLRLINFSFYDCKLQ EIFSRMNLREDVFYLQFNNSHRVKPVQNRYYRRKSYLCYQLERANGQEPLKGYLLYKKGEQHVEILFLE KMRSMELSQVRITCYLTWSPCPNCARQLAFAFKKDHDPDLILRIYTSRLYFYWRKKFQKGLCTLWRSIGHVD VMDLPQFADCWTNFVNPQRPFPRPWNELEKNSWRIQRRLRRIKESWGL* Bovine APOBEC-3B: (SEQ ID NO: 1074)  
*DGWEVAFRSGTVLKAGVLGVSMTEGWAGSGHPGQACVWTPGTRNTMNLREVLFKQQFGNQPRVPAP YYRRKTYLQYQLKQRNDLTLDRGCFRNKKQRHAEIRFIDKINSLDLNPSQSYKIICYITWSPCPNCANELVN FITRNNHLKLEIFASRLYFHWIKSFKMGLQDLQNAISVAVMTHTEFEDCWEQFVDNQSRPFQPWDKLEQY SASIRRLQRIITAPI* Chimpanzee APOBEC-3B: (SEQ ID NO: 1075)  
*MNPQIRNPMEMYQRTFYYNFENEPILYGRSYTWLCYEVKIRRGHSNLLWDTGVFRGQMYSQPEHHAEM CFLSWFCGNQLSAYKCFQITWVSWTPCPDCVAKLAKFLAHPNVTLTISAARLYYYWERDYRRALCRLS QAGARVKIMDDEEFAYCWENFVYNEGQPFMPWYKFDDNYAFLHRTLKEIIRHLMDDPDTFTFNFNNDPLVL RRHQTYLCEYEVERLDNGTWVLMQHMGLFCNEAKNLLCGFYGRHAELRFLDLVPSLQLDPAQIYRVTF ISWSPCFSWGCAGQVRAFLQENTHVRLRIFAARIYDYDPLYKEALQMLRDAGAQVSIMTYDEFEYCWDTF VYRQGC PFQPWDGLEEHSQALSGRLRAILQVRASSLCMVPHRPPPPQSPGPCLPLCSEPPLGSLPTGRPAP SLPFLLTASFSPPPASLPPLPSLSLSPGHLVPVPSFHSLSLSCSIQPPCSSRIRETGWASVSKEGRDLG* Human APOBEC-3C: (SEQ ID NO: 282)  
*MNPQIRNPMKAMYPGTFFYFQFKNLWEANDRNETWLCFTVEGIKRRSVVSWKTGVFRNQVDSETHCHAER CFLSWFCDDILSPNTKYQVTWYTSWSPCPDCAGEVAEFLARHSNVNLTIFTARLYYFQYPCYQEGLRSLSQEG VAVEIMDYEDFKYCWENFVYNDNEPFKPKWKGLKTNFRLLKRRLRESLQ* (italic: nucleic acid editing domain) Gorilla APOBEC-3C: (SEQ ID NO: 1076)  
*MNPQIRNPMKAMYPGTFFYFQFKNLWEANDRNETWLCFTVEGIKRRSVVSWKTGVFRNQVDSETHCHAER CFLSWFCDDILSPNTNYQVTWYTSWSPCECAGEVAEFLARHSNVNLTIFTARLYYFQDQDYQEGLRSLSQ EGVAVKIMDYKDFKYCWENFVYNDDEPFKPKWKGLKYNFRFLKRRLQEILE* Human APOBEC-3A: (SEQ ID NO: 283)  
*MEASPASGPRHLMDPHIFTSNFNNGIGRHKTYLCEYEVERLDNGTSVKMDQHRGFLHNQAKNLLCGFYGRH AELRFLDLVPSLQLDPAQIYRVTFISWSPCFSWGCAGEVRAFLQENTHVRLRIFAARIYDYDPLYKEALQML RDAGAQVSIMTYDEFKHCWDTFVDHQGCPFQPWDGLDEHSQALSGRLRAILQNQGN* (italic: nucleic acid editing domain) Rhesus macaque APOBEC-3A: (SEQ ID NO: 1077)  
*MDGSPASRPRHLMDDPNTFTFNFNNDLSVRGRHQTYLCEYEVERLDNGTWVPMDERRGFLCNKAKNVPCGD YGCHVELRFLCEVPSWQLDPAQTYRVTFISWSPCFRRGCAGQVRVFLQENKHVRLRIFAARIYDYDPLY QEALRTL RDAGAQVSIMTYEEFKHCWDTFVDRQGRPFQPWDGLDEHSQALSGRLRAILQNQGN* Bovine APOBEC-3A: (SEQ ID NO: 1078)  
*MDEYTFNFNQGWPSKTYLCEYEMERLDGDATIPLEDEYKGFVRNKGLDQPEKPCHAELYFLGKIHSWNL DRNQHYRLTCFISWSPCYDCAQKLTTFLKENHHISLHILASRIYTHNRFQCHQSGLCCELQAAGARITIMTFED FKHCWETFVDHKGKPFQPWEGNLVKSQALCTELQAILKTQQN* Human APOBEC-3H: (SEQ ID NO:

(284) MALLTAETFRRLAAPPYPRKALLCYQLTPQNGSTPTRGYFENKKKCHAEICFINKSMGLDETQ  
CYQVTCYLTWSPCSSCAWELVDFIKAHDHNLGIFASRLYYHWCKPQQKGLRLLCGSQVPVEVMGFPKFAD  
CWENFVDHEKPLSFNPYKMLEELDKNSRAIKRRLERIKIPGVRAQGRYMDILCDAEV (italic: nucleic acid  
editing domain) Rhesus macaque APOBEC-3H: (SEQ ID NO: 1079)  
MALLTAKTFSLQFNNKRRVKNKPYPRKALLCYQLTPQNGSTPTRGHLKNKKKDHAEIRFINKIKSMGLDET  
QCYQVTCYLTWSPCPCAGELVDFIKAHRHLNLRIFASRLYYHWRPNYQEGLLLLCGSQVPVEVMGLPEFT  
DCWENFVDHKEPPSFNPSEKLEELDKNSQAIKRRLERIKSRSDVLENGLRSLQLGPVTPSSSIRNSR Human  
APOBEC-3D (SEQ ID NO: 285)  
MNPQIRNPMERMYRDTFYDNFENEPILYGRSYTWLCYEYVKIKRGRSNLLWDTGVFRGPVLPKRQSNHRQE  
VYFRFENHAEMCFLSWFCGNRLPANRRFQITWFWVSWNPCLPCVVKVTKFLAEHPNVTLTISAARLYYYRDRD  
WRWVLLRLHKAGARVKIMDYEDFAYCWENFVCNEGQPFMPWYKFDDNYASLHRTLKEILRNPMMEAMY  
HIFYFHFKNLLKACGRNESWLCFTMEVTKHHSVFRKRGRGVFRNQVDPETHCHAERCFLSWFCDDILSPNTN  
YEVTWYTSWSPCECAGEVAEFLARHSNVNLTIFTARLCYFWDTDYQEGLCSLSQEGASVKIMGYKDFVSC  
WKNFVYSDDEPFKPKWGLQTNFRLLKRRRLREILQ (italic: nucleic acid editing domain) Human  
APOBEC-1 (SEQ ID NO: 286)  
MTSEKGPSTGDPTLRRRIEPWEFDVFYDPRELRKEACLLYEIKWGMRSRKIWRSSGKNTTNHVEVNFIEKFTS  
ERDFHPSMSCSITWFLSWSPCWECSQAIREFLSRHPGVTLVIYVARLFWHMDQQNRQGLRDLVNSGVTIQI  
MRASEYYHCWRNFVNYPGDEAHWPQYPPPLWMMLYALELHCHILSLPPCLKISRRWQNHLTFFRLHLQNC  
HYQTIPPHILLATGLIHPSVAWR Mouse APOBEC-1 (SEQ ID NO: 287)  
MSSETGPVAVDPTLRRRIEPHEFEVFFDPRELRKETCLLYEINWGGRHSVWRHTSQNTSNHVEVNFIEKFTT  
ERYFRPNTRCSITWFLSWSPCGECSRAITEFLSRHPYVTLFIYIARLYHHTDQRNRQGLRDLISSGVTIQIMTE  
QEYCYCWRNFVNYPSPNEAYWPRYPHLWVKLYVLELYCIILGLPPCLKILRRKQPQLTFFTTITLQTCHYQRI  
PPHLLWATGLK Rat APOBEC-1 (SEQ ID NO: 288)  
MSSETGPVAVDPTLRRRIEPHEFEVFFDPRELRKETCLLYEINWGGRHHSIWRHTSQNTNKHVEVNFIEKFTTE  
RYFCPNTRCSITWFLSWSPCGECSRAITEFLSRYPHVTLFIYIARLYHHADPRNRQGLRDLISSGVTIQIMTEQ  
ESGYCWRNFVNYPSPNEAHWPYPHLWVRLYVLELYCIILGLPPCLNILRRKQPQLTFFTTIALQSCHYQRLP  
PHILWATGLK Human APOBEC-2: (SEQ ID NO: 1080)  
MAQKEEA AVATEAASQNGEDLENLDDPEKLKELIELPPFEIVTGERLPANFFKFQFRNVEYSSGRNKTFCLCY  
VVEAQGGQVQASRGYLEDEHAAAHAEAAFFNTILPAFDPALRYNVTWYVSSSPCAACADRIIKTSLKTK  
NLRLLLVGRLFMWEEPEIQAALKKLKEAGCKLRIMKPQDFEYVWQNFVEQEEGESKAFQPWEDIQENFLY  
YEEKLADILK Mouse APOBEC-2: (SEQ ID NO: 1081)  
MAQKEEA AEAAPASQNGDDLENLEDPEKLKELIDLPPEIVTGVRLPVNFFKFQFRNVEYSSGRNKTFCLC  
YVVEVQSKGGQAQATQGYLEDEHAGAHAEAAFFNTILPAFDPALKYNVTWYVSSSPCAACADRILKTLKSK  
TKNLRLLLVSRLFMWEEPEVQAALKKLKEAGCKLRIMKPQDFEYIWQNFVEQEEGESKAFEPWEDIQENF  
LYYEEKLADILK Rat APOBEC-2: (SEQ ID NO: 1082)  
MAQKEEA AEAAPASQNGDDLENLEDPEKLKELIDLPPEIVTGVRLPVNFFKFQFRNVEYSSGRNKTFCLC  
YVVEAQSKGGQVQATQGYLEDEHAGAHAEAAFFNTILPAFDPALKYNVTWYVSSSPCAACADRILKTLKSK  
TKNLRLLLVSRLFMWEEPEVQAALKKLKEAGCKLRIMKPQDFEYIWQNFVEQEEGESKAFEPWEDIQENF  
LYYEEKLADILK Bovine APOBEC-2: (SEQ ID NO: 1083)  
MAQKEEA AAAAAEPASQNGEEVENLEDPEKLKELIELPPFEIVTGERLPAHYFKFQFRNVEYSSGRNKTFCLCY  
VVEAQSKGGQVQASRGYLEDEHATNHAEAAFFNSIMPTFDPALRYMVTWYVSSSPCAACADRIVKTLNKT  
KNLRLLLVGRLFMWEEPEIQAALRKLKEAGCRLRIMKPQDFEYIWQNFVEQEEGESKAFEPWEDIQENFL  
YYEEKLADILK Petromyzon marinus CDA1 (pmCDA1) (SEQ ID NO: 289)  
MTDAEYVRIHEKLDIYTFKKQFFNNKKS VSHRCYVLFELKRRGERRACFWGYAVNKPQSGTERGIHAEIFSI  
RKVEEYLRDNPQGFTINWYSSWSPCADCAEKILEWYNQELRGNGHTLKIWACKLYYEKNARNQIGLWNL  
RDNGVGLNVMVSEHYQCCRKIFIQSSHNQLNENRWLEKTLKRAEKRRSELSIMI QVKILHTTKSPAV Human  
APOBEC3G D316R\_D317R (SEQ ID NO: 290)  
MKPHFRNTVERMYRDTFSYNFYNRPILSRRNTVWLCYEYVKTGKPSRPPLDAKIFRGQVYSELKYHPEMRFF  
HWFSKWRKLHRDQEYEV TWYISWSPCTKCTRDMATFLAEDPKVTLTIFVARLYYFWDPDYQEALRSLCQ  
KRDGPRATMKIMNYDEFQHCWSKFVYSQREL FEPWNNLPKYIYILLHIMLGEILRHSM DPPTFTFNFNNEPW  
VRGRHETYL CYEVERMHNDTWVLLNQRRGFLCNQAPHKHG FLEGRHAELCFLDVIPFWKLDLDQDYRV  
TCFTSWSPCFSCAQEMAKFISK NKHVSLCIFTARIYRRQGRQCQEGLRTLAEAGAKISIMTYSEFKHCWDTFVD  
HQQGCPFPWDGLDEHSQDLSGRLRAILQNQEN Human APOBEC3G chain A (SEQ ID NO: 291)  
MDPPTFTFNFNNEPWVRGRHETYL CYEVERMHNDTWVLLNQRRGFLCNQAPHKHG FLEGRHAELCFLDV  
IPFWKLDLDQDYRVTCFTSWSPCFSCAQEMAKFISK NKHVSLCIFTARIYDDQGRQCQEGLRTLAEAGAKISI  
MTYSEFKHCWDTFVDHQQGCPFPWDGLDEHSQDLSGRLRAILQ Human APOBEC3G chain A  
D120R\_D121R (SEQ ID NO: 292)  
MDPPTFTFNFNNEPWVRGRHETYL CYEVERMHNDTWVLLNQRRGFLCNQAPHKHG FLEGRHAELCFLDV  
IPFWKLDLDQDYRVTCFTSWSPCFSCAQEMAKFISK NKHVSLCIFTARIYRRQGRQCQEGLRTLAEAGAKISI  
MTYSEFKHCWDTFVDHQQGCPFPWDGLDEHSQDLSGRLRAILQ Non-limiting examples of fusion  
proteins/nucleobase editors are provided. His6-rAPOBEC1-XTEN-dCas9 for Escherichia coli expression

(SEQ ID NO: 293)

MGSSHHHHHHMSSETGPVAVDPTLRRRIEPHEFEVFFDPRELRKETCLLYEINWGGRHSIWRHTSQNTNKH  
VEVNFIEKFTTERRYFCPNTRCSITWFLSWSPCGECSRAITEFLSRYPHVTLFIYIARLYHHADPRNRQGLRDLI  
SSGVTIQIMTEQESGYCWRNFVNYSNEAHWPYPHPLWVRLYVLELYCIILGLPPCLNILRRKQPQLTFFTI  
ALQSCHYQRLPPHILWATGLKSGSETPGTSESATPESDKKYSIGLAIGTNSVGWAVITDEYKVPSSKKFKVLG  
NTDRHSIKKNLIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIFSNEMAKVDDSSFFHRLEESFLVE  
EDKKHERHPIFGNIVDEVAYHEKYPTIYHLRKKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSD  
VDKLFQILVQTYNQLFEENPINASGVDAKAILSARLSKSRLENLIAQLPGEKKNGLFGNLIASLGLTPNFK  
SNFDLAEDAQLQSKDQYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRY  
DEHHQDLTLLKALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNRE  
DLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFYFPLKDNREKIEKILTRIPYYVGPLARGNSRFAWMTRKS  
EETITPWNFEVVDKGASAQSFIERMTNFDKNLPNEKVLPKHSLLYEYFTVYNELTKVKYVTEGMRKPAFL  
SGEQKKAIVDLLFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEEN  
EDILEDIVLTTLTFEDREMIEERLKYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLK  
SDGFANRNFMLIHDDSLTFKEDIQKAQVSGQDLSHEHIANLAGSPAIIKKGILQTVKVVDELVKVMGRH  
KPENIVIAMARENQTTQKGQKNSRERMKRIEIEGKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVD  
QELDINRLSDYDVDAIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKMKMKNYWRQLLNAKLITQR  
KFDNLTKAERGGLSELDKAGFIKRQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFR  
KDFQFYKVVREINNYHHAHDAYLNAVVG TALIKKYPKLESEFVYGDYKVYDVRKMIKSEQEIGKATAKYF  
FYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVL SMPQVNIVKKTEVQTGGFSKESI  
LPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVAKVEKGKSKKLKSVKELLGITIMERSSFEKNPIDF  
LEAKGYKEVKKDLIILPKYSLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLASHYEKLKGS PEDN  
EQKQLFVEQHKHYLDEIIEQISEFSKRVLADANLDKVL SAYNKH RDKPIREQAENIIHLFTLTNLGAPAAFK  
YFDTTIDRKRYTSTKEVLDATLIHQ SITGLYETRIDLSQLGGDSGGSPKKKRKV rAPOBEC1-XTEN-dCas9-NLS

for Mammalian expression (SEQ ID NO: 294)

MSSETGPVAVDPTLRRRIEPHEFEVFFDPRELRKETCLLYEINWGGRHSIWRHTSQNTNKHVEVNFIEKFTT  
ERYFCPNTRCSITWFLSWSPCGECSRAITEFLSRYPHVTLFIYIARLYHHADPRNRQGLRDLISSGVTIQIMTE  
QESGYCWRNFVNYSNEAHWPYPHPLWVRLYVLELYCIILGLPPCLNILRRKQPQLTFFTIALQSCHYQRL  
PPHILWATGLKSGSETPGTSESATPESDKKYSIGLAIGTNSVGWAVITDEYKVPSSKKFKVLGNTDRHSIKKN  
LIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIFSNEMAKVDDSSFFHRLEESFLVEEDKKHERHPIF  
GNIVDEVAYHEKYPTIYHLRKKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFQILVQT  
YNQLFEENPINASGVDAKAILSARLSKSRLENLIAQLPGEKKNGLFGNLIASLGLTPNFKSNFDLAEDA  
QLSKDQYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLK  
ALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDN  
GSIPHQIHLGELHAILRRQEDFYFPLKDNREKIEKILTRIPYYVGPLARGNSRFAWMTRKSEETITPWNFEV  
VDKGASAQSFIERMTNFDKNLPNEKVLPHSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDL  
LFTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEENEDILEDIVLT  
LFEDREMIEERLKYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRNF  
QLIHDDSLTFKEDIQKAQVSGQDLSHEHIANLAGSPAIIKKGILQTVKVVDELVKVMGRHKPENIVIAMARE  
NQTQKGQKNSRERMKRIEIEGKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQELDINRLSDY  
DVDAIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKMKMKNYWRQLLNAKLITQRKFDNLT  
KAERGLSELDKAGFIKRQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFRKDFQFYK  
VREINNYHHAHDAYLNAVVG TALIKKYPKLESEFVYGDYKVYDVRKMIKSEQEIGKATAKYFFYSNIMNFFKT  
EITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVL SMPQVNIVKKTEVQTGGFSKESILPKRNSDK  
LIARKKDWDPKKYGGFDSPTVAYSVLVAKVEKGKSKKLKSVKELLGITIMERSSFEKNPIDFLEAKGYKEV  
KDLIILPKYSLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLASHYEKLKGS PEDNEQKQLFVEQ  
HKHYLDEIIEQISEFSKRVLADANLDKVL SAYNKH RDKPIREQAENIIHLFTLTNLGAPAAFKYFDTTID  
RKRYTSTKEVLDATLIHQ SITGLYETRIDLSQLGGDSGGSPKKKRKV hAPOBEC1-XTEN-dCas9-NLS

for Mammalian expression (SEQ ID NO: 295)

MTSEKGPSTGDPTLRRRIEPWEFDVFDYDPRELKKEACLLYEIKWGM SRKIWRSSGKNNTNHVEVNFIIKFTSE  
RDFHPSMSCSITWFLSWSPCWECSQAIREFLSRHPGVTLVYVARLFWHMDQQNRQGLRDLVNSGVTIQI  
MRASEYYHCWRNFVNYPGDEAHWPQYPLWMMLYALELHCIIISLPPCLKISRRWQNHLTFFRLHLQNC  
HYQTIPPHILLATGLIHPSVAWRSGSETPGTSESATPESDKKYSIGLAIGTNSVGWAVITDEYKVPSSKKFKVLGN  
TDRHSIKKNLIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIFSNEMAKVDDSSFFHRLEESFLVEEDKK  
HERHPIFGNIVDEVAYHEKYPTIYHLRKKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFQI  
LVQTYNQLFEENPINASGVDAKAILSARLSKSRLENLIAQLPGEKKNGLFGNLIASLGLTPNFKSNFDLAEDA  
KLQLSKDQYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLK  
LVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDNGSIPH  
QIHLGELHAILRRQEDFYFPLKDNREKIEKILTRIPYYVGPLARGNSRFAWMTRKSEETITPWNFEVVDKGA  
SAQSFIERMTNFDKNLPNEKVLPHSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFKTNRKVT  
VKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEENEDILEDIVLTTLTFEDREMIEERL

TYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRNFMQLIHDDSLTFKEDIQK  
AQVSGQGDSLHEHIANLAGSPAIIKGILQTVKVVDDELVKVMGRHKPENIVIAMARENQTTQKGQKNSRERM  
KRIIEGIKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQELDINRLSDYDVDAIVPQSFLKDDSIDNK  
VLTRSDKNRGKSDNVPSEEVVKKMKNYWRQLLNAKLITQRKFDNLTKAERGGLSELDKAGFIKRQLVETRQIT  
KHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFRKDFQFYKVREINNYHHAHDAYLNAVVG TALIKKYP  
KLESEFVYGDYKVYDVRKMIAKSEQEIGKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKG  
RDFATVRKVL SMPQVNIVKKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKVE  
KGKSKKLKSVKELLGITIMERSSEFEKNPIDFLEAKGYKEVKKD LIIKLPKYSLFELENGRKRMLASAGELQKGNEL  
ALPSKYVNFLYLASHYEKLKGSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVLADANLDKVL SAYNKHRDKPI  
REQAENIIHLFTLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQ SITGLYETRIDLSQLGGDSGGSPKKKRK V  
rAPOBEC1-XTEN-dCas9-UGI-NLS (SEQ ID NO: 296)

MSSETGPVAVDPTLRRRIEPHEFEVFFDPREL RKETCLLYEINWGGRHSIWRHTSQNTNKHVEVNFIEKFTT  
ERYFCPNTRCSITWFLSWSPCGECSRAITEFLSRYPHVT LFIYIARLYHHADPRNRQGLRDLISSGVTIQIMTE  
QESGYCWRNFVNYSNPSNEAHWP RYPHLWVRLYVLELYCIILGLPPCLNILRRKQPQLTFFTIALQSCHYQRL  
PPHILWATGLKSGSETPGTSESATPESDKKYSIGLAIGTNSVGWAVITDEYKVPSKKFKVLGNTDRHSIKKN  
LIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIFS NEMAKVDDSFHRL EESFLVEEDKKHERHPIF  
GNIVDEVAYHEKYPTIYHLRKKLVDSTDKADLR LIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQT  
YNQLFEENPINASGVDAKAILSARLSKSRLENLIAQLPGEKKNGLFGNLIALSLGLTPNFKSNFDLAEDA KL  
QLSKD TYDDDLNLLAQIGDQYADLFLAAKNLS DAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLK  
ALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDN  
GSIPHQIHLGELHAILRRQEDFY PFLKDNREKIEKILTFRIPYYVGPLARGNSRFAWMTRKSEETITPWNFEEV  
VDKGASAQSFIERMTNFDKNLPNEKVL PKHSLLEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDL  
LFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEENEDILEDIVLT LT  
LFEDREMIEERLKYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRNFM  
QLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIIKGILQTVKVVDDELVKVMGRHKPENIVIAMARE  
NQTTQKGQKNSRERMKRIIEGIKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQELDINRLSDY  
DVDAIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKKMKNYWRQLLNAKLITQRKFDNLTKAERG  
GLSELDKAGFIKRQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFRKDFQFYKVREI  
NNYHHAHDAYLNAVVG TALIKKYPKLESEFVYGDYKVYDVRKMIAKSEQEIGKATAKYFFYSNIMNFFKT  
EITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVL SMPQVNIVKKTEVQTGGFSKESILPKRNSDKLIA  
RKKDWDPKKYGGFDSPTVAYSVLVVAKVEKGKSKKLKSVKELLGITIMERSSEFEKNPIDFLEAKGYKEVK  
KD LIIKLPKYSLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLASHYEKLKGSPEDNEQKQLFVEQH  
KHYLDEIIIEQISEFSKRVLADANLDKVL SAYNKHRDKPIREQAENIIHLFTLTNLGAPAAFKYFDTTIDRKRY  
TSTKEVLDATLIHQ SITGLYETRIDLSQLGGDSGGSTNLS DIIKETGKQLVIQESILMLPEEVVEEVIGNKPESD  
ILVHTAYDESTDENVMLLTSDAPEYKPWALVIQDSNGENKIKMLSGGSPKKKRKV rAPOBEC1-XTEN-Cas9  
nickase-UGI-NLS )(BE3, SEQ ID NO: 297

MSSETGPVAVDPTLRRRIEPHEFEVFFDPREL RKETCLLYEINWGGRHSIWRHTSQNTNKHVEVNFIEKFTT  
ERYFCPNTRCSITWFLSWSPCGECSRAITEFLSRYPHVT LFIYIARLYHHADPRNRQGLRDLISSGVTIQIMTE  
QESGYCWRNFVNYSNPSNEAHWP RYPHLWVRLYVLELYCIILGLPPCLNILRRKQPQLTFFTIALQSCHYQRL  
PPHILWATGLKSGSETPGTSESATPESDKKYSIGLAIGTNSVGWAVITDEYKVPSKKFKVLGNTDRHSIKKN  
LIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIFS NEMAKVDDSFHRL EESFLVEEDKKHERHPIF  
GNIVDEVAYHEKYPTIYHLRKKLVDSTDKADLR LIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQT  
YNQLFEENPINASGVDAKAILSARLSKSRLENLIAQLPGEKKNGLFGNLIALSLGLTPNFKSNFDLAEDA KL  
QLSKD TYDDDLNLLAQIGDQYADLFLAAKNLS DAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLK  
ALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDN  
GSIPHQIHLGELHAILRRQEDFY PFLKDNREKIEKILTFRIPYYVGPLARGNSRFAWMTRKSEETITPWNFEEV  
VDKGASAQSFIERMTNFDKNLPNEKVL PKHSLLEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDL  
LFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEENEDILEDIVLTITL  
FEDREMIEERLKYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRNFMQ  
LIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIIKGILQTVKVVDDELVKVMGRHKPENIVIAMAREN  
QTTQKGQKNSRERMKRIIEGIKELGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQELDINRLSDYD  
VDHIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVPSEEVVKKMKNYWRQLLNAKLITQRKFDNLTKAERG  
LSELDKAGFIKRQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFRKDFQFYKVREIN  
NYHHAHDAYLNAVVG TALIKKYPKLESEFVYGDYKVYDVRKMIAKSEQEIGKATAKYFFYSNIMNFFKTE  
ITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVL SMPQVNIVKKTEVQTGGFSKESILPKRNSDKLIAR  
KKDWDPKKYGGFDSPTVAYSVLVVAKVEKGKSKKLKSVKELLGITIMERSSEFEKNPIDFLEAKGYKEVK  
D LIIKLPKYSLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLASHYEKLKGSPEDNEQKQLFVEQH  
HYLDEIIIEQISEFSKRVLADANLDKVL SAYNKHRDKPIREQAENIIHLFTLTNLGAPAAFKYFDTTIDRKRYT  
STKEVLDATLIHQ SITGLYETRIDLSQLGGDSGGSTNLS DIIKETGKQLVIQESILMLPEEVVEEVIGNKPESDI  
LVHTAYDESTDENVMLLTSDAPEYKPWALVIQDSNGENKIKMLSGGSPKKKRKV pmCDA1-XTEN-dCas9-UGI  
(bacteria) (SEQ ID NO: 298)

MTDAEYVRIHEKLDIYTFKKQFFNNKKSVSHRCYVLFELKRRGERRACFWGYAVNKPQ  
SGTERGIAHEIFSIRKVEEYLRDNPQGFTINWYSSWSPCADCAEKILEWYNQELRGNGHT  
LKIWACKLYYEKNARNQIGLWNLRDNGVGLNVMVSEHYQCCRKIFIQSSHNQLNENR  
WLEKTLKRAEKRRSELSIMIQVKILHTTKSPAVSGSETPGTSESATPESDKKYSIGLAIGT  
NSVGWAVITDEYKVPSKKFKVLGNTDRHSIKKNLIGALLFDSGETAEATRLKRTARRRY  
TRRKNRICYLQEIFSNEMAKVDDSFHRLSESLVEEDKKHERHPIFGNIVDEVAYHEKY  
PTIYHLRKKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTY  
NQLFEENPINASGVDAKAILSARLSKSRLENLIAQLPGEKKNGLFGNLIASLSGLTPNFK  
SNFDLAEDAKLQLSKDTYDDDDLNDLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEIT  
KAPLSASMIKRYDEHHQDLTLLKALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEF  
YKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFYPL  
KDNREKIEKILTFRIPIYYVGPLARGNSRFAWMTRKSEETITPWNFEEVVDKGASAQSFIE  
RMTNFDKNLPNEKVLPHKSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDL  
LFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRNFNASLGTYHDLLKIIKDKDFLDNEE  
NEDILEDIVLTLTLEFEDREMIEERLKTYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIR  
DKQSGKTILDFLKSDGFANRNFQMQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSP  
AIKKGILQTVKVVDELVKVMGRHKPENIVIAMARENQTTQKGQKNSRERMKRIEEGIKE  
LGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQELDINRLSDYDVAIVPQSFLKD  
DSIDNKVLTRSDKNRGKSDNVPSEEVVKMKMKNYWRQLLNAKLITQRKFDNLTKAERGG  
LSELDKAGFIKRLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFR  
KDFQFYKVINNYHHAHDAYLNAVVGTAIIKKYPKLESEFVYGDYKVYDVRKMIAS  
EQEIGKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRK  
VLSMPQVNIVKKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVL  
VVAKVEKGKSKKLKSVKELLGITIMERSSEFKNPIDFLEAKGYKEVKKDLIIKLPKYSLFE  
LENGRKRMLASAGELQKGNELALPSKYVNFLYLASHYEKLKGSPEDEQKQLFVEQHK  
HYLDEIIEQISEFSKRVILADANLDKVL SAYNKH RD KPIREQAENIIHLFTLTNLGAPAAFK  
YFDTTIDRKRYTSTKEVL DATLIHQ SITGLYETRIDLSQLGGDSGGSMTNLSDIIEKETGK  
QLVIQESILMLPEEVEEVIGNKPESDILVHTAYDESTDENVMLLTSDAPEYKPWALVIQD SNGENKIKML

pmCDA1-XTEN-nCas9-UGI-NLS (mammalian construct) (SEQ ID NO: 299);

MTDAEYVRIHEKLDIYTFKKQFFNNKKSVSHRCYVLFELKRRGERRACFWGYAVNKPQ  
SGTERGIAHEIFSIRKVEEYLRDNPQGFTINWYSSWSPCADCAEKILEWYNQELRGNGHT  
LKIWACKLYYEKNARNQIGLWNLRDNGVGLNVMVSEHYQCCRKIFIQSSHNQLNENR  
WLEKTLKRAEKRRSELSIMIQVKILHTTKSPAVSGSETPGTSESATPESDKKYSIGLAIGT  
NSVGWAVITDEYKVPSKKFKVLGNTDRHSIKKNLIGALLFDSGETAEATRLKRTARRRY  
TRRKNRICYLQEIFSNEMAKVDDSFHRLSESLVEEDKKHERHPIFGNIVDEVAYHEKY  
PTIYHLRKKLVDSTDKADLRLIYLALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTY  
NQLFEENPINASGVDAKAILSARLSKSRLENLIAQLPGEKKNGLFGNLIASLSGLTPNFK  
SNFDLAEDAKLQLSKDTYDDDDLNDLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEIT  
KAPLSASMIKRYDEHHQDLTLLKALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEF  
YKFIKPILEKMDGTEELLVKLNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFYPL  
KDNREKIEKILTFRIPIYYVGPLARGNSRFAWMTRKSEETITPWNFEEVVDKGASAQSFIE  
RMTNFDKNLPNEKVLPHKSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDL  
LFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRNFNASLGTYHDLLKIIKDKDFLDNEE  
NEDILEDIVLTLTLEFEDREMIEERLKTYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIR  
DKQSGKTILDFLKSDGFANRNFQMQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSP  
AIKKGILQTVKVVDELVKVMGRHKPENIVIAMARENQTTQKGQKNSRERMKRIEEGIKE  
LGSQILKEHPVENTQLQNEKLYLYYLQNGRDMYVDQELDINRLSDYDVHIVPQSFLKD  
DSIDNKVLTRSDKNRGKSDNVPSEEVVKMKMKNYWRQLLNAKLITQRKFDNLTKAERGG  
LSELDKAGFIKRLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFR  
KDFQFYKVINNYHHAHDAYLNAVVGTAIIKKYPKLESEFVYGDYKVYDVRKMIAS  
EQEIGKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRK  
VLSMPQVNIVKKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVL  
VVAKVEKGKSKKLKSVKELLGITIMERSSEFKNPIDFLEAKGYKEVKKDLIIKLPKYSLFE  
LENGRKRMLASAGELQKGNELALPSKYVNFLYLASHYEKLKGSPEDEQKQLFVEQHK  
HYLDEIIEQISEFSKRVILADANLDKVL SAYNKH RD KPIREQAENIIHLFTLTNLGAPAAFK  
YFDTTIDRKRYTSTKEVL DATLIHQ SITGLYETRIDLSQLGGDSGGSTNLSDIIEKETGKQL  
VIQESILMLPEEVEEVIGNKPESDILVHTAYDESTDENVMLLTSDAPEYKPWALVIQDSN  
GENKIKMLSGGSPKKKRKV huAPOBEC3G-XTEN-dCas9-UGI (bacteria) (SEQ ID NO: 300)  
MDPPTFTFNFNNEPWVRGRHETLYLCYEVERMHNDTWVLLNQRRGFLCNQAPHKHGFL  
EGRHAELCFLDVIPFWKLDLDQDYRVTCFTSWSPCFSCAQEMAKFISK NKHVSLCIFTAR  
IYDDQGRQCQGLR TLAEAGAKISIMTYSEFKHCWDTFVDHQGCPFPWDGLDEHSQDL



SGRLRAILQSGSETPGTSESATPESDKKYSIGLAIGTNSVGWAVITDEYKVPSSKKFKVLG  
NTDRHSIKKNLIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIFSNEMAKVDDSD  
FFHRLEESFLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLRKKLVDSTDKADLRLIYL  
ALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDAKAILSARL  
SKSRLENLIAQLPGEKKNGLFGNLIASLGLTPNFKSNDLAEDAQLQSKDQYDDDL  
NLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLK  
ALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNRE  
DLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFYFPLKDNREKIEKILTRIPYYVGPLAR  
GNSRFAWMTRKSEETITPWNFEEVVDKGASAQSFIERMTNFDKNLPNEKVLPKHSLLYE  
YFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFKTNRKVTVKQLKEDYFKKIECF  
DSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEENEDILEDIVLTLTLFEDREMIEERLK  
TYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRNFMQ  
LIHDDSLTFKEDIQKAQVSGQGDLSHEHIANLAGSPAIKKGILQTVKVVDELVKVMGRH  
KPENIVIAMARENQTTQKGQKNSRERMKRIIEGKELGSQILKEHPVENTQLQNEKLYLY  
YLQNGRDMYVDQELDINRLSDYDVAIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVP  
EEVVKMKMKNYWRQLLNAKLITQRKFDNLTKAERGGLSELDKAGFIKRQLVETRQITKH  
VAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFRKDFQFYKPREINNYHHAHDAYL  
NAVVGITALIKKYPKLESEFVYGDYKVYDVRKMIAKSEQEIGKATAKYFFYSNIMNFFKT  
EITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVLSMPQVNIVKKTEVQTGGFSKE  
SILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVAKVEKGKSKKLKSVKELLGIT  
IMERSSFEKNPIDFLEAKGYKEVKDLIILPKYSLFELENKRMLASAGELQKGNELA  
LPSKYVNFLYLASHYEKLKGSPEDEQKQLFVEQHKHYLDEIIEQISEFSKRVILADANL  
DKVLSAYNKHDKPIREQAENIIHLFTLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDTLI  
HQSITGLYETRIDLSQLGGDSGGSMTNLSDIIEKETGKQLVIQESILMLPEEVVEEVIGNKPE  
SDILVHTAYDESTDENVMMLTSDAPEYKPWALVIQDSNGENKIKML huAPOBEC3G-XTEN-nCas9-UGI-NLS  
(mammalian construct) (SEQ ID NO: 301)

MDPPTFTFNFNNEPWVRGRHETYLCEYVERMHNDTWVLLNQRRGFLCNQAPHKHGFL  
EGRHAELCFLDVIPFWKLDLDQDYRVTCFTSWSPCFSCAQEMAKFISKNKHVSLCIFTAR  
IYDDQGRQCQGLRDLAEAGAKISIMTYSEFKHCWDTFVDHQGCPFQPWDGLDEHSQDL  
SGRLRAILQSGSETPGTSESATPESDKKYSIGLAIGTNSVGWAVITDEYKVPSSKKFKVLG  
NTDRHSIKKNLIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIFSNEMAKVDDSD  
FFHRLEESFLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLRKKLVDSTDKADLRLIYL  
ALAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDAKAILSARL  
SKSRLENLIAQLPGEKKNGLFGNLIASLGLTPNFKSNDLAEDAQLQSKDQYDDDL  
NLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLK  
ALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNRE  
DLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFYFPLKDNREKIEKILTRIPYYVGPLAR  
GNSRFAWMTRKSEETITPWNFEEVVDKGASAQSFIERMTNFDKNLPNEKVLPKHSLLYE  
YFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFKTNRKVTVKQLKEDYFKKIECF  
DSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEENEDILEDIVLTLTLFEDREMIEERLK  
TYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRNFMQ  
LIHDDSLTFKEDIQKAQVSGQGDLSHEHIANLAGSPAIKKGILQTVKVVDELVKVMGRH  
KPENIVIAMARENQTTQKGQKNSRERMKRIIEGKELGSQILKEHPVENTQLQNEKLYLY  
YLQNGRDMYVDQELDINRLSDYDVAIVPQSFLKDDSIDNKVLTRSDKNRGKSDNVP  
EEVVKMKMKNYWRQLLNAKLITQRKFDNLTKAERGGLSELDKAGFIKRQLVETRQITKH  
VAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFRKDFQFYKPREINNYHHAHDAYL  
NAVVGITALIKKYPKLESEFVYGDYKVYDVRKMIAKSEQEIGKATAKYFFYSNIMNFFKT  
EITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVLSMPQVNIVKKTEVQTGGFSKE  
SILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVAKVEKGKSKKLKSVKELLGIT  
IMERSSFEKNPIDFLEAKGYKEVKDLIILPKYSLFELENKRMLASAGELQKGNELA  
LPSKYVNFLYLASHYEKLKGSPEDEQKQLFVEQHKHYLDEIIEQISEFSKRVILADANL  
DKVLSAYNKHDKPIREQAENIIHLFTLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDTLI  
HQSITGLYETRIDLSQLGGDSGGSTNLSDIIEKETGKQLVIQESILMLPEEVVEEVIGNKPE  
DILVHTAYDESTDENVMMLTSDAPEYKPWALVIQDSNGENKIKMLSGGSPKKKRKV huAPOBEC3G  
(D316R\_D317R)-XTEN-nCas9-UGI-NLS (mammalian construct) (SEQ ID NO: 302)  
MDPPTFTFNFNNEPWVRGRHETYLCEYVERMHNDTWVLLNQRRGFLCNQAPHKHGFL  
EGRHAELCFLDVIPFWKLDLDQDYRVTCFTSWSPCFSCAQEMAKFISKNKHVSLCIFTAR  
IYRRQGRQCQGLRDLAEAGAKISIMTYSEFKHCWDTFVDHQGCPFQPWDGLDEHSQDL  
GRLRAILQSGSETPGTSESATPESDKKYSIGLAIGTNSVGWAVITDEYKVPSSKKFKVLGN  
TDRHSIKKNLIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIFSNEMAKVDDSD  
FHRLEESFLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLRKKLVDSTDKADLRLIYLA



LAHMIKFRGHFLIEGDLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDAKAILSARLS  
KSRRLLENLIAQLPGEKKNGLFGNLIALSLGLTPNFKSNFDLAEDAKLQLSKDTYDDDL  
NLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLTLLK  
ALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLVKLNRE  
DLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFYFPFLKDNREKIEKILTRIPYYVGPLAR  
GNSRFAWMTRKSEETITPWNFEEVVDKGASQSFIERMTNFDKNLPNEKVLPKHSLLYE  
YFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFKTNRKVTVKQLKEDYFKKIECF  
DSVEISGVEDRFNASLGTYHDLLKIIKDKDFLDNEENEDILEDIVLTLTLFEDREMIEERLK  
TYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGKTILDFLKSDGFANRNFQM  
LIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIIKKGILQTVKVVDELVKVMGRH  
KPENIVIAMARENQTTQKGQKNSRERMKRIIEGKELGSQILKEHPVENTQLQNEKLYLY  
YLQNGRDMYVDQELDINRLSDYDVDHIVPQSFLKDDSIDNKVLTRSDKNRGKSDNPVS  
EEVVKMKMKNYWRQLLNAKLITQRKFDNLTKAERGGLSELDKAGFIKRQLVETRQITKH  
VAQILDSRMNTKYDENDKLIREVKVITLKSCLVSDFRKDFQFYKVVREINNYHHAHDAYL  
NAVVG TALIKKYPKLESEFVYGDYKVYDVRKMIAKSEQEIGKATAKYFFYSNIMNFFKT  
EITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVL SMPQVNIVKKTEVQTGGFSKE  
SILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKVEKGKSKKLKSVKELLGIT  
IMERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELENGRKRMLASAGELQKGNELA  
LPSKYVNFLYLASHYEKLKGSPEDEQKQLFVEQHKHYLDEIIEQISEFSKRVILADANL  
DKVLSAYNKHHRDKPIREQAENIIHLFTLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLI  
HQSITGLYETRIDLSQLGGDSGGSTNLSDIIEKETGKQLVQIESILMLPEEVEEVIGNKPES DILVHTA  
YDESTDENVMMLTSDAPEYKPWALVIQDSNGENKIKMLSGGSPKKRKV Base Editor 4 (BE4; APOBEC1-  
linker(32 aa)-Cas9n(D10A)-linker(9 aa)- UGI-linker(9 aa)-UGI) (SEQ ID NO: 1084)  
MSSETGPVAVDPTLRRRIEPHEFEVFFDPRELRKETCLLYEINWGGRHISWRHTSQNTNKHVEVNFIEKFTT  
ERYFCPNTRCSITWFLSWSPCGECSRAITEFLSRYPHVTLFIYIARLYHHADPRNRQGLRDLISSGVTIQIMTE  
QESGYCWRNFVNYSNPSNEAHWPYPHLLWVRLYVLELYCIILGLPPCLNLRKQPQLTFFTIALQSCHYQRL  
PPHILWATGLKSGGSSGGSSGSETPGTSESATPESSGGSSGGSDKKYSIGLAIGTNSVGWAVITDEYKVPSKK  
FKVLGNTDRHSIKKNLIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIFSNEMAKVDDSSFFHRL  
SFLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLRKKLVDSTDKADRLIYLALAHMIKFRGHFLIEGDLN  
PDNSDVKLFIQLVQTYNQLFEENPINASGVDAKAILSARLSKSRRLLENLIAQLPGEKKNGLFGNLIALSLGL  
TPNFKSNFDLAEDAKLQLSKDTYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSAS  
MIKRYDEHHQDLTLLKALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILEKMDGTEELLV  
KLNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFYFPFLKDNREKIEKILTRIPYYVGPLARGNSRFAW  
MTRKSEETITPWNFEEVVDKGASQSFIERMTNFDKNLPNEKVLPKHSLLYEYFTVYNELTKVKYVTEGM  
RKPAFLSGEQKKAIVDLLFKTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRFNASLGTYHDLLKIIKDKDF  
LDNEENEDILEDIVLTLTLFEDREMIEERLKTYAHLFDDKVMKQLKRRRYTGWGRLSRKLINGIRDKQSGK  
TILDFLKSDGFANRNFQMQLIHDDSLTFKEDIQKAQVSGQGDSLHEHIANLAGSPAIIKKGILQTVKVVDELVK  
VMGRHKPENIVIAMARENQTTQKGQKNSRERMKRIIEGKELGSQILKEHPVENTQLQNEKLYLYLONGR  
DMYVDQELDINRLSDYDVDHIVPQSFLKDDSIDNKVLTRSDKNRGKSDNPSEEVVKMKMKNYWRQLLNA  
KLITQRKFDNLTKAERGGLSELDKAGFIKRQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVITLKS  
LVSDFRKDFQFYKVVREINNYHHAHDAYLNAVVG TALIKKYPKLESEFVYGDYKVYDVRKMIAKSEQEIGK  
ATAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATVRKVL SMPQVNIVKKTEVQTG  
GFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKVEKGKSKKLKSVKELLGITIMERSSFE  
KNPIDFLEAKGYKEVKKDLIIKLPKYSLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLASHYEKLK  
GSPEDNEQKQLFVEQHKHYLDEIIEQISEFSKRVILADANL DKVLSAYNKHHRDKPIREQAENIIHLFTLTNLG  
APAAFKYFDTTIDRKRYTSTKEVLDATLIHQ SITGLYETRIDLSQLGGDSGGSGGSGGSTNLSDIIEKETGKQ  
LVQIESILMLPEEVEEVIGNKPESDILVHTAYDESTDENVMMLTSDAPEYKPWALVIQDSNGENKIKMLSGG  
SGGSGGSTNLSDIIEKETGKQLVQIESILMLPEEVEEVIGNKPESDILVHTAYDESTDENVMMLTSDAPEYKP  
WALVIQDSNGENKIKMLSGGSPKKRK

Example 2: Anti-Cancer Vaccination Using CRISPR-Cas9 Genome/Base-Editing Technologies

(254) Described herein are new methods to stimulate the immune system to treat tumors and prevent metastatic lesions. By turning the genome and proteome of the malignant cells into a personalized endogenous anti-cancer vaccine in vivo.

(255) Provided herein is a new immuno-oncology methodology to raise robust T-cell and B-cell mediated immune responses against tumor-specific proteins, which are otherwise tolerated as “self” by the immune system of a cancer patient.<sup>sup.1-3</sup> (FIG. 1, Tables 1-3). This methodology is uniquely suited for programmable CRISPR-Cas9 genome- and base-editing tools,<sup>sup.4-10</sup> exploited to alter the translated sequences of tumor specific genes.<sup>sup.11</sup> to produce highly immunogenic heteroclitic and cryptic peptide epitopes in situ (Tables 5 and 7, FIG. 1). Heteroclitic epitopes are altered versions of endogenous peptide sequences engineered to elicit potent immune reactions through the MHC-I and MHC-II antigen presentation pathways,<sup>sup.12</sup> which also produce cross-reactive responses towards the parent wild-type peptide sequences (FIG. 2A).<sup>sup.2,13,14</sup> For example, the peptide epitope EAAGIGILTV (SEQ ID NO: 388) from the melanocyte differentiation and melanoma marker MART-1.<sup>sup.26-36</sup> is weakly immunogenic, whereas vaccination with a

similar peptide engineered with a hydrophobic residue.<sup>sup.15,16</sup> on the MHC-anchor position MART-1 (27L) ELAGIGILTV (SEQ ID NO: 1085) promotes robust T-cell immune responses against melanoma (Table 5).<sup>sup.17,18</sup> Cryptic epitopes arise from non-translated genomic sequences through processes that are elevated in cancer cells, such as aberrant mRNA splicing, alternative open-reading frames (ORFs), and deglycosylation of proteins.<sup>sup.19</sup> For example, LAGE-1 immunogenic antigens are expressed from ORF-2.<sup>sup.20,21</sup> of the gene NY-ESO-122-24 (Table 7). Introduction of these immunogenic protein sequences using genome/base editing is designed to break “self”-tolerance to cancer-specific antigens,<sup>sup.1-3</sup> which is known drive the infiltration of immune cells into the tumor promoting the recognition of malignant cells as foreign through 3,827,234 multiple mechanisms (FIG. 3).<sup>sup.25,26</sup> Anti-cancer vaccination using genome/base-editing is rendered cancer-specific by targeting genes that are preferentially or exclusively expressed by tumor cells to prevent autoimmunity side effects.<sup>sup.2,27</sup> Anti-cancer vaccination strategies could be particularly useful for the treatment of melanoma (Table 1),<sup>sup.28-30</sup> as well as colorectal tumors, stomach cancer, and other highly mutagenized cancers that accumulate non-synonymous hitchhiker mutations with high frequencies (FIG. 4).<sup>sup.31</sup> In such cases, spreading of the adaptive immune response towards translated “neo-epitopes”<sup>sup.32,33</sup> that are unique to the cell lineage facilitate remission.<sup>sub.34</sup> and prevent metastatic lesions (abscopal effect) 35-39 (FIG. 3). Importantly, the recently FDA-approved checkpoint inhibitors (anti-PD1 and anti-CTLA4 antibodies), which lower the threshold for T-cell stimulation,<sup>sup.27,40</sup> have shown promise for co-administration with anti-cancer vaccines,<sup>sup.41-46</sup> and could enhance the clinical effectiveness of immunization against a broader assortment of cancer types.<sup>sup.47,48</sup>

(256) The heteroclitic and cryptic epitopes programmed by genome/base-editing may be personalized to match each patient's malignancy and immune system, or alternatively a guide-RNA cocktail can be developed to engage the most frequent HLA allele supertypes that broadly cover the human population (FIG. 2B). Recent advances for the delivery of genome editing tools are enabling for anti-cancer vaccination. These methods include intracellular delivery using electroporation,<sup>sup.49-50</sup> viral vectors,<sup>sup.51,52</sup> cell penetrating peptides,<sup>sup.53,54</sup> liposomes,<sup>sup.55,56</sup> polymers,<sup>sup.57</sup> membrane deformation,<sup>sup.58</sup> and nanoparticles.<sup>sup.59</sup>; and the types of cargo include RNA transcripts, DNA expression vectors, or Cas9 protein-guide RNA complexes purified,<sup>sup.60</sup> or within cationic lipid vesicles.<sup>sup.61,62</sup> Therefore, the vaccination treatments could be potentially performed in vivo directly on tumor cells,<sup>sup.63</sup> the tissues that originated the tumor,<sup>sup.64</sup> or alternatively ex vivo for re-injection of irradiated whole-cell vaccines.<sup>sup.65-67</sup>

(257) Also provided herein are numerous specific examples of genomic target sites in tumor-associated genes (Tables 1-3), and the guide-RNAs designed to program the alteration of these translated sequences (Tables 5 and 7), in order to replicate or closely mimic known epitopes that have literature or pre-clinical precedent. The genome editing reactions were designed for one of the CRISPR/Cas9 tools: (i) “base editors” that catalyze chemical reactions on nucleobases (e.g. cytidine deaminase-Cas9 fusion<sup>4</sup>); or (ii) engineered nucleases with DNA cutting activity (e.g. WT Cas9,<sup>sup.5-7</sup> Cas9 nickases.<sup>sup.8</sup> or FokI-nuclease-dCas9 fusions.<sup>sup.9,10</sup>). Examples of other potentially useful genome-editing reactions to alter cancer-specific genes to produce heteroclitic/cryptic epitopes are shown in Tables 5 and 7. By extension, Cas9 tools and Homology-Directed Repair (HDR) pathways may also be exploited to introduce heteroclitic epitopes through DNA templates by lowering the rate of indels using several techniques.<sup>sup.68-70</sup> finally, to expand the repertoire of heteroclitic and cryptic epitopes in an unbiased high-throughput manner, the aforementioned tools could be used to screen libraries of guide-RNAs targeting all PAM sites across a tumor-associated genes of interest, which can be replicated using the genome/base-editing reactions shown in Tables 5 and 7.<sup>sup.71,72</sup>

## REFERENCES

- (258) 1. Melero, I. et al. Therapeutic vaccines for cancer: an overview of clinical trials. *Nature reviews. Clinical oncology* 11, 509-524, (2014). 2. Pardoll, D. M. Inducing autoimmune disease to treat cancer. *Proceedings of the National Academy of Sciences of the United States of America* 96, 5340-5342, (1999). 3. Buonaguro, L., Petrizzo, A., Tornesello, M. L. & Buonaguro, F. M. Translating tumor antigens into cancer vaccines. *Clinical and vaccine immunology: CVI* 18, 23-34, (2011). 4. Komor, A. C., Kim, Y. B., Packer, M. S., Zuris, J. A. & Liu, D. R. Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage. *Nature* advance online publication, (2016). 5. Cong, L. et al. Multiplex genome engineering using CRISPR/Cas systems. *Science* 339, 819-823, (2013). 6. Jinek, M. et al. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337, 816-821, (2012). 7. Mali, P. et al. RNA-guided human genome engineering via Cas9. *Science* 339, 823-826, (2013). 8. Ran, F. A. et al. Double nicking by RNA-guided CRISPR Cas9 for enhanced genome editing specificity. *Cell* 154, 1380-1389, (2013). 9. Guilinger, J. P., Thompson, D. B. & Liu, D. R. Fusion of catalytically inactive Cas9 to FokI nuclease improves the specificity of genome modification. *Nature biotechnology* 32, 577-582, (2014). 10. Tsai, S. Q. et al. Dimeric CRISPR RNA-guided FokI nucleases for highly specific genome editing. *Nature biotechnology* 32, 569-576, (2014). 11. Vigneron, N., Stroobant, V., Van den Eynde, B. J. & van der Bruggen, P. Database of T cell-defined human tumor antigens: the 2013 update. *Cancer immunity* 13, 15, (2013). 12. Borbulevych, O. Y., Baxter, T. K., Yu, Z., Restifo, N. P. & Baker, B. M. Increased immunogenicity of an anchor-modified tumor-associated antigen is due to the enhanced stability of the peptide/MHC complex: implications for vaccine design. *J Immunol* 174, 4812-4820, (2005). 13. Bakker, A. B. et al. Analogues of CTL epitopes with improved MHC class-I binding capacity elicit anti-melanoma CTL recognizing the wild-type epitope. *International journal of cancer* 70, 302-309, (1997). 14. Purcell, A. W., McCluskey, J. & Rossjohn, J. More than one reason to rethink the use of peptides in vaccine design. *Nat Rev Drug Discov* 6, 404-414, (2007). 15. Chowell, D. et al. TCR contact residue hydrophobicity is a hallmark of immunogenic CD8+ T cell epitopes. *Proceedings of the*

*National Academy of Sciences of the United States of America* 112, E1754-1762, (2015). 16. Ruppert, J. et al. Prominent role of secondary anchor residues in peptide binding to HLA-A2.1 molecules. *Cell* 74, 929-937, (1993). 17. Madura, F. et al. Structural basis for ineffective T-cell responses to MHC anchor residue-improved "heteroclitic" peptides. *European journal of immunology* 45, 584-591, (2015). 18. Rivoltini, L. et al. A superagonist variant of peptide MART1/Melan A27-35 elicits anti-melanoma CD8+ T cells with enhanced functional characteristics: implication for more effective immunotherapy. *Cancer research* 59, 301-306, (1999). 19. Andersen, R. S. et al. High frequency of T cells specific for cryptic epitopes in melanoma patients. *Oncoimmunology* 2, e25374, (2013). 20. Rimoldi, D. et al. Efficient simultaneous presentation of NY-ESO-1/LAGE-1 primary and nonprimary open reading frame-derived CTL epitopes in melanoma. *J Immunol* 165, 7253-7261, (2000). 21. Mandic, M. et al. The alternative open reading frame of LAGE-1 gives rise to multiple promiscuous HLA-DR-restricted epitopes recognized by T-helper 1-type tumor-reactive CD4+ T cells. *Cancer research* 63, 6506-6515, (2003). 22. Campos-Perez, J. et al. DNA fusion vaccine designs to induce tumor-lytic CD8+ T-cell attack via the immunodominant cysteine-containing epitope of NY-ESO 1. *International journal of cancer* 133, 1400-1407, (2013). 23. Webb, A. I. et al. Functional and structural characteristics of NY-ESO-1-related HLA A2-restricted epitopes and the design of a novel immunogenic analogue. *The Journal of biological chemistry* 279, 23438-23446, (2004). 24. Chen, J. L. et al. Identification of NY-ESO-1 peptide analogues capable of improved stimulation of tumor-reactive CTL. *J Immunol* 165, 948-955, (2000). 25. Lally, K. M. et al. Unmasking cryptic epitopes after loss of immunodominant tumor antigen expression through epitope spreading. *International journal of cancer* 93, 841-847, (2001). 26. Fridman, W. H. et al. The ultimate goal of curative anti-cancer therapies: inducing an adaptive anti-tumor immune response. *Frontiers in immunology* 2, 66, (2011). 27. Attia, P. et al. Autoimmunity correlates with tumor regression in patients with metastatic melanoma treated with anti-cytotoxic T-lymphocyte antigen-4. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology* 23, 6043-6053, (2005). 28. Rosenberg, S. A. et al. Immunologic and therapeutic evaluation of a synthetic peptide vaccine for the treatment of patients with metastatic melanoma. *Nature medicine* 4, 321-327, (1998). 29. Slingluff, C. L., Jr. et al. Clinical and immunologic results of a randomized phase II trial of vaccination using four melanoma peptides either administered in granulocyte-macrophage colony-stimulating factor in adjuvant or pulsed on dendritic cells. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology* 21, 4016-4026, (2003). 30. Slingluff, C. L., Jr. et al. Immunologic and clinical outcomes of vaccination with a multiepitope melanoma peptide vaccine plus low-dose interleukin-2 administered either concurrently or on a delayed schedule. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology* 22, 4474-4485, (2004). 31. Lawrence, M. S. et al. Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* 499, 214-218, (2013). 32. Duan, F. et al. Genomic and bioinformatic profiling of mutational neoepitopes reveals new rules to predict anticancer immunogenicity. *J Exp Med* 211, 2231-2248, (2014). 33. Schumacher, T. N. & Schreiber, R. D. Neoantigens in cancer immunotherapy. *Science* 348, 69-74, (2015). 34. Gubin, M. M. et al. Checkpoint blockade cancer immunotherapy targets tumour-specific mutant antigens. *Nature* 515, 577-581, (2014). 35. Kreiter, S. et al. Mutant MHC class II epitopes drive therapeutic immune responses to cancer. *Nature* 520, 692-696, (2015). 36. Chandra, R. A. et al. A systematic evaluation of abscopal responses following radiotherapy in patients with metastatic melanoma treated with ipilimumab. *Oncoimmunology* 4, e1046028, (2015). 37. Ma, Y. et al. Chemotherapy and radiotherapy: cryptic anticancer vaccines. *Seminars in immunology* 22, 113-124, (2010). 38. Karbach, J. et al. Long-term complete remission following radiosurgery and immunotherapy in a melanoma patient with brain metastasis: immunologic correlates. *Cancer immunology research* 2, 404-409, (2014). 39. Demaria, S. et al. Ionizing radiation inhibition of distant untreated tumors (abscopal effect) is immune mediated. *International journal of radiation oncology, biology, physics* 58, 862-870, (2004). 40. Wong, R. M. et al. Programmed death-1 blockade enhances expansion and functional capacity of human melanoma antigen-specific CTLs. *International immunology* 19, 1223-1234, (2007). 41. Fu, J. et al. Preclinical evidence that PD1 blockade cooperates with cancer vaccine TEGVAX to elicit regression of established tumors. *Cancer research* 74, 4042-4052, (2014). 42. Gibney, G. T. et al. Safety, correlative markers, and clinical results of adjuvant nivolumab in combination with vaccine in resected high-risk metastatic melanoma. *Clinical cancer research: an official journal of the American Association for Cancer Research* 21, 712-720, (2015). 43. Soares, K. C. et al. PD-1/PD-L1 blockade together with vaccine therapy facilitates effector T-cell infiltration into pancreatic tumors. *J Immunother* 38, 1-11, (2015). 44. Le, D. T. et al. Evaluation of ipilimumab in combination with allogeneic pancreatic tumor cells transfected with a GM-CSF gene in previously treated pancreatic cancer. *J Immunother* 36, 382-389, (2013). 45. Sierro, S. R. et al. Combination of lentivector immunization and low-dose chemotherapy or PD-1/PD-L1 blocking primes self-reactive T cells and induces anti-tumor immunity. *European journal of immunology* 41, 2217-2228, (2011). 46. Hodi, F. S. et al. Biologic activity of cytotoxic T lymphocyte-associated antigen 4 antibody blockade in previously vaccinated metastatic melanoma and ovarian carcinoma patients. *Proceedings of the National Academy of Sciences of the United States of America* 100, 4712-4717, (2003). 47. Morse, M. A. & Lyerly, H. K. Checkpoint blockade in combination with cancer vaccines. *Vaccine* 33, 7377-7385, (2015). 48. Vanneman, M. & Dranoff, G. Combining immunotherapy and targeted therapies in cancer treatment. *Nature reviews. Cancer* 12, 237-251, (2012). 49. Bakondi, B. et al. In Vivo CRISPR/Cas9 Gene Editing Corrects Retinal Dystrophy in the S334ter-3 Rat Model of Autosomal Dominant Retinitis Pigmentosa. *Molecular therapy: the journal of the American Society of Gene Therapy* 24, 556-563, (2016). 50. Chen, S., Lee, B., Lee, A. Y., Modzelewski, A. J. & He, L. Highly Efficient Mouse Genome Editing by CRISPR Ribonucleoprotein Electroporation of Zygotes. *The Journal of biological chemistry*, (2016). 51. Maggio, I. et al. Adenoviral vector delivery of RNA-guided CRISPR/Cas9 nuclease complexes induces targeted mutagenesis in a diverse array of human cells.

Scientific reports 4, 5105, (2014). 52. Wang, W. et al. CCR5 gene disruption via lentiviral vectors expressing Cas9 and single guided RNA renders cells resistant to HIV-1 infection. *PloS one* 9, e115987, (2014). 53. Ramakrishna, S. et al. Gene disruption by cell-penetrating peptide-mediated delivery of Cas9 protein and guide RNA. *Genome research* 24, 1020-1027, (2014). 54. Liu, J., Gaj, T., Patterson, J. T., Sirk, S. J. & Barbas, C. F., 3rd. Cell-penetrating peptide-mediated delivery of TALEN proteins via bioconjugation for genome engineering. *PloS one* 9, e85755, (2014). 55. Ye, L. et al. Seamless modification of wild-type induced pluripotent stem cells to the natural CCR5Delta32 mutation confers resistance to HIV infection. *Proceedings of the National Academy of Sciences of the United States of America* 111, 9591-9596, (2014). 56. Allen, T. M. & Cullis, P. R. Liposomal drug delivery systems: from concept to clinical applications. *Advanced drug delivery reviews* 65, 36-48, (2013). 57. *Efficient Delivery of Sigma CRISPRs via a Non-Liposomal Polymeric Transfection Reagent, TransIT®-CRISPR*, <[www.sigmaaldrich.com/technical-documents/articles/biology/transit-crispr-transfection-reagent.html](http://www.sigmaaldrich.com/technical-documents/articles/biology/transit-crispr-transfection-reagent.html)> (2016). 58. Han, X. et al. CRISPR-Cas9 delivery to hard-to-transfect cells via membrane deformation. *Science advances* 1, e1500454, (2015). 59. Cox, D. B., Platt, R. J. & Zhang, F. Therapeutic genome editing: prospects and challenges. *Nature medicine* 21, 121-131, (2015). 60. Kim, S., Kim, D., Cho, S. W., Kim, J. & Kim, J. S. Highly efficient RNA-guided genome editing in human cells via delivery of purified Cas9 ribonucleoproteins. *Genome research* 24, 1012-1019, (2014). 61. Zuris, J. A. et al. Cationic lipid-mediated delivery of proteins enables efficient protein-based genome editing in vitro and in vivo. *Nature biotechnology* 33, 73-80, (2015). 62. Wang, M. et al. Efficient delivery of genome-editing proteins using bio-reducible lipid nanoparticles. *Proceedings of the National Academy of Sciences of the United States of America*, (2016). 63. Villarreal, D. O. et al. Alarmin IL-33 acts as an immunoadjuvant to enhance antigen-specific tumor immunity. *Cancer research* 74, 1789-1800, (2014). 64. Ginsberg, B. A. et al. Immunologic response to xenogeneic gp100 DNA in melanoma patients: comparison of particle-mediated epidermal delivery with intramuscular injection. *Clinical cancer research: an official journal of the American Association for Cancer Research* 16, 4057-4065, (2010). 65. Geary, S. M., Lemke, C. D., Lubaroff, D. M. & Salem, A. K. Proposed mechanisms of action for prostate cancer vaccines. *Nature reviews. Urology* 10, 149-160, (2013). 66. Dranoff, G. et al. Vaccination with irradiated tumor cells engineered to secrete murine granulocyte-macrophage colony-stimulating factor stimulates potent, specific, and long-lasting anti-tumor immunity. *Proceedings of the National Academy of Sciences of the United States of America* 90, 3539-3543, (1993). 67. Chiang, C. L., Benencia, F. & Coukos, G. Whole tumor antigen vaccines. *Seminars in immunology* 22, 132-143, (2010). 68. Chu, V. T. et al. Increasing the efficiency of homology-directed repair for CRISPR-Cas9-induced precise gene editing in mammalian cells. *Nature biotechnology* 33, 543-548, (2015). 69. Yu, C. et al. Small molecules enhance CRISPR genome editing in pluripotent stem cells. *Cell stem cell* 16, 142-147, (2015). 70. Paquet, D. et al. Efficient introduction of specific homozygous and heterozygous mutations using CRISPR/Cas9. *Nature* 533, 125-129, (2016). 71. Chen, S. et al. Genome-wide CRISPR screen in a mouse model of tumor growth and metastasis. *Cell* 160, 1246-1260, (2015). 72. Shalem, O., Sanjana, N. E. & Zhang, F. High-throughput functional genomics using CRISPR-Cas9. *Nature reviews. Genetics* 16, 299-311, (2015). 73. Overwijk, W. W. & Restifo, N. P. B16 as a mouse model for human melanoma. *Current protocols in immunology*/edited by John E. Coligan . . . [et al.] Chapter 20, Unit 20 21, (2001). 74. Saenger, Y. M. et al. Improved tumor immunity using anti-tyrosinase related protein-1 monoclonal antibody combined with DNA vaccines in murine melanoma. *Cancer research* 68, 9884-9891, (2008). 75. Yu, Z. et al. Poor immunogenicity of a self/tumor antigen derives from peptide-MHC-I instability and is independent of tolerance. *The Journal of clinical investigation* 114, 551-559, (2004). 76. Robbins, P. F. et al. The intronic region of an incompletely spliced gp100 gene transcript encodes an epitope recognized by melanoma-reactive tumor-infiltrating lymphocytes. *J Immunol* 159, 303-308, (1997). 77. Pinilla, C. et al. Combinatorial peptide libraries as an alternative approach to the identification of ligands for tumor-reactive cytolytic T lymphocytes. *Cancer research* 61, 5153-5160, (2001). 78. Wang, R. F., Parkhurst, M. R., Kawakami, Y., Robbins, P. F. & Rosenberg, S. A. Utilization of an alternative open reading frame of a normal gene in generating a novel human cancer antigen. *J Exp Med* 183, 1131-1140, (1996). 79. Lupetti, R. et al. Translation of a retained intron in tyrosinase-related protein (TRP) 2 mRNA generates a new cytotoxic T lymphocyte (CTL)-defined and shared human melanoma antigen not expressed in normal cells of the melanocytic lineage. *J Exp Med* 188, 1005-1016, (1998). 80. Visseren, M. J. et al. Affinity, specificity and T-cell-receptor diversity of melanoma-specific CTL generated in vitro against a single tyrosinase epitope. *International journal of cancer* 72, 1122-1128, (1997). 81. Valmori, D. et al. Analysis of the cytolytic T lymphocyte response of melanoma patients to the naturally HLA-A\*0201-associated tyrosinase peptide 368-376. *Cancer research* 59, 4050-4055, (1999). 82. Rubio-Godoy, V. et al. Toward synthetic combinatorial peptide libraries in positional scanning format (PS-SCL)-based identification of CD8+ Tumor-reactive T-Cell Ligands: a comparative analysis of PS-SCL recognition by a single tumor-reactive CD8+ cytolytic T-lymphocyte clone. *Cancer research* 62, 2058-2063, (2002). 83. Skipper, J. C. et al. An HLA-A2-restricted tyrosinase antigen on melanoma cells results from posttranslational modification and suggests a novel pathway for processing of membrane proteins. *J Exp Med* 183, 527-534, (1996). 84. Bernatchez, C. et al. Altered decamer and nonamer from an HLA-A0201-restricted epitope of Survivin differentially stimulate T-cell responses in different individuals. *Vaccine* 29, 3021-3030, (2011). 85. Hirohashi, Y. et al. An HLA-A24-restricted cytotoxic T lymphocyte epitope of a tumor-associated protein, survivin. *Clinical cancer research: an official journal of the American Association for Cancer Research* 8, 1731-1739, (2002). 86. Gross, D. A. et al. High vaccination efficiency of low-affinity epitopes in antitumor immunotherapy. *The Journal of clinical investigation* 113, 425-433, (2004). 87. Scardino, A. et al. HER-2/neu and hTERT cryptic epitopes as novel targets for broad spectrum tumor immunotherapy. *J Immunol* 168, 5900-5906, (2002). 88. Aurisicchio, L. et al. A novel minigene scaffold for therapeutic cancer vaccines. *Oncoimmunology* 3, e27529, (2014). 89. Bae, J., Martinson, J. A.

Klingemann, H. G. Identification of novel CD33 antigen-specific peptides for the generation of cytotoxic T lymphocytes against acute myeloid leukemia. *Cellular immunology* 227, 38-50, (2004). 90. Bae, J., Martinson, J. A. & Klingemann, H. G. Heteroclitic CD33 peptide with enhanced anti-acute myeloid leukemic immunogenicity. *Clinical cancer research: an official journal of the American Association for Cancer Research* 10, 7043-7052, (2004). 91. Smith, H. A., Rekoske, B. T. & McNeel, D. G. DNA vaccines encoding altered peptide ligands for SSX2 enhance epitope-specific CD8<sup>+</sup> T-cell immune responses. *Vaccine* 32, 1707-1715, (2014). 92. Oka, Y., Tsuboi, A., Oji, Y., Kawase, I. & Sugiyama, H. WT1 peptide vaccine for the treatment of cancer. *Current opinion in immunology* 20, 211-220, (2008). 93. Pinilla-Ibarz, J. et al. Improved human T-cell responses against synthetic HLA-0201 analog peptides derived from the WT1 oncoprotein. *Leukemia* 20, 2025-2033, (2006). 94. Di Stasi, A., Jimenez, A. M., Minagawa, K., Al-Obaidi, M. & Rezvani, K. Review of the Results of WT1 Peptide Vaccination Strategies for Myelodysplastic Syndromes and Acute Myeloid Leukemia from Nine Different Studies. *Frontiers in immunology* 6, 36, (2015). 95. Tsuboi, A. et al. Enhanced induction of human WT1-specific cytotoxic T lymphocytes with a 9-mer WT1 peptide modified at HLA-A\*2402-binding residues. *Cancer immunology, immunotherapy: CII* 51, 614-620, (2002). 96. Trojan, A. et al. Generation of cytotoxic T lymphocytes against native and altered peptides of human leukocyte antigen-A\*0201 restricted epitopes from the human epithelial cell adhesion molecule. *Cancer research* 61, 4761-4765, (2001). 97. Volpe, G. et al. Alternative BCR/ABL splice variants in Philadelphia chromosome-positive leukemias result in novel tumor-specific fusion proteins that may represent potential targets for immunotherapy approaches. *Cancer research* 67, 5300-5307, (2007). 98. Casnici, C. et al. Out of frame peptides from BCR/ABL alternative splicing are immunogenic in HLA A2.1 transgenic mice. *Cancer letters* 276, 61-67, (2009). 99. Casnici, C. et al. Immunologic evaluation of peptides derived from BCR/ABL-out-of-frame fusion protein in HLA A2.1 transgenic mice. *J Immunother* 35, 321-328, (2012). 100. Tangri, S. et al. Structural features of peptide analogs of human histocompatibility leukocyte antigen class I epitopes that are more potent and immunogenic than wild-type peptide. *J Exp Med* 194, 833-846, (2001). 101. Barve, M. et al. Induction of immune responses and clinical efficacy in a phase II trial of IDM-210I, a 10-epitope cytotoxic T-lymphocyte vaccine, in metastatic non-small-cell lung cancer. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology* 26, 4418-4425, (2008). 102. Graff-Dubois, S. et al. Generation of CTL recognizing an HLA-A\*0201-restricted epitope shared by MAGE-A1, -A2, -A3, -A4, -A6, -A10, and -A12 tumor antigens: implication in a broad-spectrum tumor immunotherapy. *J Immunol* 169, 575-580, (2002). 103. Tourdot, S. et al. A general strategy to enhance immunogenicity of low-affinity HLA-A2. 1-associated peptides: implication in the identification of cryptic tumor epitopes. *European journal of immunology* 30, 3411-3421, (2000). 104. Tsang, K. Y., Palena, C., Gulley, J., Arlen, P. & Schlom, J. A human cytotoxic T-lymphocyte epitope and its agonist epitope from the nonvariable number of tandem repeat sequence of MUC-1. *Clinical cancer research: an official journal of the American Association for Cancer Research* 10, 2139-2149, (2004). 105. Geynisman, D. M. et al. A randomized pilot phase I study of modified carcinoembryonic antigen (CEA) peptide (CAP1-6D)/montanide/GM-CSF-vaccine in patients with pancreatic adenocarcinoma. *Journal for immunotherapy of cancer* 1, 8, (2013). 106. Kleinstiver, B. P. et al. Engineered CRISPR-Cas9 nucleases with altered PAM specificities. *Nature* 523, 481-485, (2015). 107. Guevara-Patino, J. A. et al. Optimization of a self antigen for presentation of multiple epitopes in cancer immunity. *The Journal of clinical investigation* 116, 1382-1390, (2006). 108. Duan, F. et al. Immune rejection of mouse tumors expressing mutated self. *Cancer research* 69, 3545-3553, (2009). 109. Myers, C. E. et al. Variation in cytotoxic T-lymphocyte responses to peptides derived from tyrosinase-related protein-2. *Human immunology* 69, 24-31, (2008). 110. Castle, J. C. et al. Exploiting the mutanome for tumor vaccination. *Cancer research* 72, 1081-1091, (2012). 111. Cho, H. I. & Celis, E. Optimized peptide vaccines eliciting extensive CD8 T-cell responses with therapeutic antitumor effects. *Cancer research* 69, 9012-9019, (2009).

#### EQUIVALENTS AND SCOPE

(259) In the claims articles such as “a,” “an,” and “the” may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include “or” between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The disclosure includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The disclosure includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

(260) Furthermore, the disclosure encompasses all variations, combinations, and permutations in which one or more limitations, elements, clauses, and descriptive terms from one or more of the listed claims is introduced into another claim. For example, any claim that is dependent on another claim can be modified to include one or more limitations found in any other claim that is dependent on the same base claim. Where elements are presented as lists, e.g., in Markush group format, each subgroup of the elements is also disclosed, and any element(s) can be removed from the group. It should be understood that, in general, where the disclosure, or aspects of the disclosure, is/are referred to as comprising particular elements and/or features, certain embodiments of the disclosure or aspects of the disclosure consist, or consist essentially of, such elements and/or features. For purposes of simplicity, those embodiments have not been specifically set forth in haec verba herein.

(261) It is also noted that the terms “comprising” and “containing” are intended to be open and permits the inclusion of additional elements or steps. Where ranges are given, endpoints are included. Furthermore, unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as

ranges can assume any specific value or sub-range within the stated ranges in different embodiments of the disclosure, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.

(262) This application refers to various issued patents, published patent applications, journal articles, and other publications, all of which are incorporated herein by reference. If there is a conflict between any of the incorporated references and the instant specification, the specification shall control. In addition, any particular embodiment of the present disclosure that falls within the prior art may be explicitly excluded from any one or more of the claims. Because such embodiments are deemed to be known to one of ordinary skill in the art, they may be excluded even if the exclusion is not set forth explicitly herein. Any particular embodiment of the disclosure can be excluded from any claim, for any reason, whether or not related to the existence of prior art.

(263) Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation many equivalents to the specific embodiments described herein. The scope of the present embodiments described herein is not intended to be limited to the above Description, but rather is as set forth in the appended claims. Those of ordinary skill in the art will appreciate that various changes and modifications to this description may be made without departing from the spirit or scope of the present disclosure, as defined in the following claims.

## Claims

1. A method for producing a heteroclitic epitope in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a composition comprising: (i) a fusion protein comprising (a) a guide nucleotide sequence-programmable DNA-binding protein domain, and (b) a cytosine deaminase domain; and (ii) a guide nucleotide sequence, wherein the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 724-751, 870-877, 888-905, and 907-985; wherein the guide nucleotide sequence of (ii) targets the fusion protein of (i) to a polynucleotide encoding a tumor-specific antigen in a tumor cell; and wherein the fusion protein changes a target cytosine (C) base to a thymine (T) base via deamination.
2. The method of claim 1, wherein the polynucleotide encoding the tumor-specific antigen is located in the genome of the tumor cell.
3. The method of claim 1, wherein the guide nucleotide sequence-programmable DNA-binding protein domain is selected from the group consisting of nuclease inactive Cas9 (dCas9) domains, Cas9 nickase (nCas9) domains, nuclease inactive Cpf1 domains, and nuclease inactive Argonaute domains.
4. The method of claim 3, wherein the guide nucleotide sequence-programmable DNA-binding protein domain is a nuclease inactive Cas9 (dCas9) domain or a Cas9 nickase (nCas9) domain.
5. The method of claim 1, wherein the cytosine deaminase domain is selected from the group consisting of APOBEC1, APOBEC2, APOBEC3A, APOBEC3B, APOBEC3C, APOBEC3D, APOBEC3F, APOBEC3G, APOBEC3H, APOBEC4, and activation-induced deaminase (AID).
6. The method of claim 1, wherein the fusion protein of (i) further comprises a uracil glycosylase inhibitor (UGI) domain.
7. The method of claim 6, wherein the fusion protein comprises the structure: NH.sub.2-[cytosine deaminase domain]-[optional linker sequence]-[guide nucleotide sequence-programmable DNA-binding protein domain]-[optional linker sequence]-[UGI domain]-COOH; NH.sub.2-[UGI domain]-[optional linker sequence]-[cytosine deaminase domain]-[optional linker sequence]-[guide nucleotide sequence-programmable DNA-binding protein domain]-COOH; or NH.sub.2-[cytosine deaminase domain]-[optional linker sequence]-[guide nucleotide sequence-programmable DNA-binding protein domain]-COOH.
8. The method of claim 7, wherein the optional linker sequence comprises (GGGS).sub.n, (SEQ ID NO: 337), (GGGGGS).sub.n (SEQ ID NO: 308), (G).sub.n (SEQ ID NO: 783), (EAAAK).sub.n (SEQ ID NO: 309), (GGGS).sub.n (SEQ ID NO: 784), SGSETPGTSESATPES (SEQ ID NO: 310), (XP).sub.n (SEQ ID NO: 785), or a combination of any of these, wherein n is independently an integer between 1 and 30, and wherein X is any amino acid.
9. The method of claim 1, wherein the fusion protein comprises the amino acid sequence of any one of SEQ ID NOs: 293-302, 1071, and 1084.
10. The method of claim 1, wherein the tumor-specific antigen is selected from the group consisting of: gp100; MART-1; hTERT; TyRP1; HER2; CEA-CAM; tyrosinase (TYR); CD33; MAGE-A3; MAGE-A4; NY-ESO-1; SSX-2; survivin; EpCAM; and MUC1.
11. The method of claim 1, wherein the target C base is in a target codon in a coding region of the polynucleotide encoding the tumor-specific antigen, and wherein the target codon is any one of the following target codons: CTT (Leu/L), CTC (Leu/L), ATG (Met/M), GTT (Val/V), GTA (Val/V), GTC (Val/V), GTG (Val/V), TCT (Ser/S), TCC (Ser/S), TCA (Ser/S), TCG (Ser/S), AGT (Ser/S), AGC (Ser/S), CCT (Pro/P), CCC (Pro/P), CCA (Pro/P), CCG (Pro/P), ACT (Thr/T), ACC (Thr/T), ACA (Thr/T), ACG (Thr/T), GCT (Ala/A), GCC (Ala/A), GCA (Ala/A), GCG (Ala/A), CAT (His/H), CAC (His/H), GAT (Asp/D), GAC (Asp/D), GAA (Glu/E), GAG (Glu/E), TGT (Cys/C), TGC (Cys/C), CGT (Arg/R), CGC (Arg/R), AGA (Arg/R), AGG (Arg/R), CGG (Arg/R), GGT (Gly/G), GGC (Gly/G), GGA (Gly/G), GGG (Gly/G), CAG (Gln/Q), TGG (Trp/W), CGA (Arg/R), CAA (Gln/Q), TGG (Trp/W), and CGA (Arg/R).
12. The method of claim 11, wherein the target codon is converted to a modified codon selected from any one of the following modified codons: ATA (Ile/I), ATT (Ile/I), ATC (Ile/I), ATG (Met/M), TTT (Phe/F), TTC (Phe/F), TTA (Leu/L), TTG (Leu/L), AAT (Asp/N), AAC (Asp/N), TCT (Ser/S), TCC (Ser/S), TCA (Ser/S), TCG (Ser/S), CTT (Leu/L), CTC

(Leu/L), CTA (Leu/L), CTG (Leu/L), GTT (Val/V), GTC (Val/V), GTA (Val/V), GTG (Val/V), ACT (Thr/T), ACC (Thr/T), ACA (Thr/T), ACG (Thr/T), TAT (Tyr/Y), TAC (Tyr/Y), AAA (Lys/K), AAG (Lys/K), TGT (Cys/C), TGC (Cys/C), CAG (Gln/Q), TGG (Trp/W), GAT (Asp/D), GAC (Asp/D), GAA (Glu/E), GAG (Glu/E), AGT (Ser/S), AGC (Ser/S), AGA (Arg/R), AGG (Arg/R), TAG (amber), TGA (opal), and TAA (ochre).

13. The method of claim 1, wherein the target C base is located in a non-coding region of the polynucleotide encoding the tumor-specific antigen.
  14. The method of claim 1, wherein the heteroclitic epitope is at least 5-fold more immunogenic than a native epitope from the tumor-specific antigen.
  15. The method of claim 1, wherein the heteroclitic epitope is displayed on the surface of the tumor cell via the MHC class I antigen presentation pathway.
  16. The method of claim 1, wherein the method is carried out in vivo.
  17. The method of claim 1, wherein the tumor-specific antigen is gp100.
  18. The method of claim 17, wherein the deamination of the target C base: (a) in codon T210 of gp100 results in a T210I mutation (SEQ ID NO: 786); (b) in codon A288 of gp100 results in a A288V mutation (SEQ ID NO: 818); or (c) in codon T155 of gp100 results in a T155I mutation (SEQ ID NO: 787).
  19. The method of claim 18, wherein; (a) a heteroclitic epitope comprising the amino acid sequence of IIDQVPFSV (SEQ ID NO: 786) is generated, and wherein the I at position 2 corresponds to the T210I mutation; (b) a heteroclitic epitope comprising the amino acid sequence of YLEPGPVTV (SEQ ID NO: 818) is generated, and wherein the V at position 7 corresponds to the A288V mutation; or (c) a heteroclitic epitope comprising the amino acid sequence of KIWGQYWQV (SEQ ID NO: 787) is generated, and wherein the I at position 2 corresponds to the T155I mutation.
  20. The method of claim 19, wherein the guide nucleotide sequence comprises a nucleotide sequence selected from the group consisting of SEQ ID NOs: 724, 725, 870-877, 888, and 889.
  21. The method of claim 1, the method further comprising administering to the subject a therapeutically effective amount of an immune checkpoint inhibitor.
  22. The method of claim 1, wherein the target C base is in a target codon in a coding region of the polynucleotide encoding the tumor-specific antigen.
  23. The method of claim 1, wherein the heteroclitic epitope is displayed on the surface of an antigen presenting cell (APC) via the MHC class II antigen presentation pathway.
-