



US012391978B2

(12) **United States Patent**
Oldham-Haltom et al.(10) **Patent No.:** **US 12,391,978 B2**(45) **Date of Patent:** ***Aug. 19, 2025**

- (54) **REAL TIME CLEAVAGE ASSAY**
- (71) Applicant: **Exact Sciences Corporation**, Madison, WI (US)
- (72) Inventors: **Rebecca Oldham-Haltom**, Marshall, WI (US); **Hongzhi Zou**, Middleton, WI (US); **Graham P. Lidgard**, Madison, WI (US); **Michael J. Domanico**, Middleton, WI (US); **Hatim Allawi**, Middleton, WI (US)
- (73) Assignee: **Exact Sciences Corporation**, Madison, WI (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 269 days.

This patent is subject to a terminal disclaimer.

4,735,214 A 4/1988 Berman
4,965,188 A 10/1990 Mullis et al.
5,011,769 A 4/1991 Duck et al.
5,124,246 A 6/1992 Urdea et al.
5,288,609 A 2/1994 Engelhardt
5,338,671 A 8/1994 Scalice et al.
5,403,711 A 4/1995 Walder et al.
5,409,818 A 4/1995 Davey et al.
5,475,610 A 12/1995 Atwood et al.
5,494,810 A 2/1996 Barany et al.
5,508,169 A 4/1996 Deugau et al.
5,538,871 A 7/1996 Nuovo et al.
5,541,311 A 7/1996 Dahlberg et al.
5,602,756 A 2/1997 Atwood et al.
5,612,473 A 3/1997 Wu et al.
5,614,402 A 3/1997 Dahlberg et al.
5,624,802 A 4/1997 Urdea et al.
5,639,611 A 6/1997 Wallace et al.
5,660,988 A 8/1997 Duck et al.
5,710,264 A 1/1998 Urdea et al.
5,719,028 A 2/1998 Dahlberg et al.
5,773,258 A 6/1998 Birch et al.
5,786,146 A 7/1998 Herman et al.
5,792,614 A 8/1998 Western et al.
5,795,763 A 8/1998 Dahlberg et al.

(Continued)

(21) Appl. No.: **18/045,927**(22) Filed: **Oct. 12, 2022**

(65) **Prior Publication Data**
US 2023/0183782 A1 Jun. 15, 2023

FOREIGN PATENT DOCUMENTS

CN 104781421 A 7/2015
EP 1674585 6/2006

(Continued)

Related U.S. Application Data

- (63) Continuation of application No. 16/789,276, filed on Feb. 12, 2020, now Pat. No. 11,499,179, which is a continuation of application No. 15/068,364, filed on Mar. 11, 2016, now Pat. No. 10,604,793, which is a continuation of application No. 15/019,758, filed on Feb. 9, 2016, now abandoned, which is a continuation of application No. 13/720,757, filed on Dec. 19, 2012, now Pat. No. 9,290,797, which is a continuation of application No. 12/946,737, filed on Nov. 15, 2010, now Pat. No. 8,361,720.

- (51) **Int. Cl.**
C12Q 1/68 (2018.01)
C12Q 1/6818 (2018.01)
C12Q 1/6827 (2018.01)
C12Q 1/686 (2018.01)
C12Q 1/6886 (2018.01)

- (52) **U.S. Cl.**
CPC **C12Q 1/6827** (2013.01); **C12Q 1/6818** (2013.01); **C12Q 1/686** (2013.01); **C12Q 1/6886** (2013.01); **C12Q 2600/156** (2013.01); **C12Q 2600/158** (2013.01)

- (58) **Field of Classification Search**
CPC C12Q 1/686
See application file for complete search history.

- (56) **References Cited**

U.S. PATENT DOCUMENTS

4,376,110 A 3/1983 David et al.
4,486,530 A 12/1984 David et al.
4,683,195 A 7/1987 Mullis et al.
4,683,202 A 7/1987 Mullis

20 Claims, 3 Drawing Sheets**Specification includes a Sequence Listing.****OTHER PUBLICATIONS**

Allawi, et al., "Invader plus method detects herpes simplex virus in cerebrospinal fluid and simultaneously differentiates types 1 and 2", J Clin Microbiol., 2006, 44:3443-7.

Applied Biosystems, "Methylation Analysis by Bisulfite Sequencing: Chemistry, Products and Protocols from Applied Biosystems", 2007, 52pgs.

Eads, et al., "MethyLight: a high-throughput assay to measure DNA methylation", Nucleic Acids Res., 2000, 28:E32, 8pgs.

Gomez, et al. "Werkhauser RP, Abath FG. Development of a real time polymerase chain reaction for quantitation of Schistosoma mansoni DNA", Mem Inst Oswaldo Cruz. Sep. 2006;101 Suppl 1:133-6.

(Continued)

Primary Examiner — Gary Benzion*Assistant Examiner* — Olatinka A Oyeyemi

(74) *Attorney, Agent, or Firm* — Bozicevic, Field & Francis LLP; Shweta Chandra

- (57) **ABSTRACT**

A cleavage-based real-time PCR assay method is provided. In general terms, the assay method includes subjecting a reaction mixture comprising a) PCR reagents for amplifying a nucleic acid target, and b) flap cleavage reagents for performing a flap cleavage assay on the amplified nucleic acid target to two sets of thermocycling conditions. No additional reagents are added to the reaction between said first and second sets of cycles and, in each cycle of the second set of cycles, cleavage of a flap probe is measured.

(56)

References Cited

U.S. PATENT DOCUMENTS

5,837,450	A	11/1998	Dahlberg et al.	7,384,746	B2	6/2008	Lyamichev et al.
5,843,654	A	12/1998	Heisler et al.	7,407,782	B2	8/2008	Prudent et al.
5,843,669	A	12/1998	Kaiser et al.	7,429,455	B2	9/2008	Dong et al.
5,846,717	A	12/1998	Brow et al.	7,432,048	B2	10/2008	Neri et al.
5,849,481	A	12/1998	Urdea et al.	7,435,390	B2	10/2008	Cracauer et al.
5,851,770	A	12/1998	Babon et al.	7,462,451	B2	12/2008	Skrzypczynski et al.
5,874,283	A	2/1999	Harrington et al.	7,473,773	B2	1/2009	Elagin et al.
5,882,867	A	3/1999	Ullman et al.	7,482,118	B2	1/2009	Allawi et al.
5,888,780	A	3/1999	Dahlberg et al.	7,482,127	B2	1/2009	Agarwal et al.
5,914,230	A	6/1999	Liu et al.	7,514,220	B2	4/2009	Hall et al.
5,958,692	A	9/1999	Cotton et al.	7,527,928	B2	5/2009	Neri et al.
5,965,408	A	10/1999	Short	7,527,948	B2	5/2009	Hudson et al.
5,985,557	A	11/1999	Prudent et al.	7,541,145	B2	6/2009	Prudent et al.
5,994,069	A	11/1999	Hall et al.	7,582,436	B2	9/2009	Hall et al.
6,001,567	A	12/1999	Brow et al.	7,588,891	B2	9/2009	Prudent et al.
6,013,170	A	1/2000	Meade	7,601,496	B2	10/2009	Dahlberg et al.
6,063,573	A	5/2000	Kayyem	7,662,594	B2	2/2010	Kong et al.
6,090,543	A	7/2000	Prudent et al.	7,674,924	B2	3/2010	Skrzypczynski et al.
6,090,606	A	7/2000	Kaiser et al.	7,678,542	B2	3/2010	Lyamichev et al.
6,110,677	A	8/2000	Western et al.	7,691,573	B2	4/2010	Dahlberg et al.
6,110,684	A	8/2000	Kemper et al.	7,700,750	B2	4/2010	Mast et al.
6,121,001	A	9/2000	Western et al.	7,790,393	B2	9/2010	Lyamichev et al.
6,150,097	A	11/2000	Tyagi et al.	8,304,214	B2	11/2012	Gerdes et al.
6,183,960	B1	2/2001	Lizardi	8,361,720	B2	1/2013	Oldham-Haltom et al.
6,194,149	B1	2/2001	Neri et al.	RE44,596	E	11/2013	Stroun et al.
6,210,880	B1	4/2001	Lyamichev et al.	8,715,937	B2	5/2014	Zou et al.
6,210,884	B1	4/2001	Lizardi	8,808,990	B2	8/2014	Lidgard et al.
6,214,545	B1	4/2001	Dong et al.	8,916,344	B2	12/2014	Zou et al.
6,221,583	B1	4/2001	Kayyem et al.	8,962,250	B2	2/2015	Stanley
6,235,502	B1	5/2001	Weissman et al.	9,000,146	B2	4/2015	Bruinsma et al.
6,248,229	B1	6/2001	Meade	9,096,893	B2	8/2015	Allawi et al.
6,251,594	B1	6/2001	Gonzalzo et al.	9,127,318	B2	9/2015	Oldham-haltom et al.
6,329,178	B1	12/2001	Patel et al.	9,163,278	B2	10/2015	Bruinsma et al.
6,335,167	B1	1/2002	Pinkel et al.	9,169,511	B2	10/2015	Lidgard et al.
6,348,314	B1	2/2002	Prudent et al.	9,212,392	B2	12/2015	Allawi et al.
6,355,437	B1	3/2002	Neri et al.	9,290,797	B2	3/2016	Oldham-Haltom et al.
6,358,691	B1	3/2002	Neri et al.	9,315,853	B2	4/2016	Domanico et al.
6,372,424	B1	4/2002	Brow et al.	9,422,592	B2	8/2016	Morris et al.
6,395,524	B2	5/2002	Loeb et al.	9,428,746	B2	8/2016	Holmberg et al.
6,458,535	B1	10/2002	Hall et al.	9,546,403	B1	1/2017	Warren et al.
6,555,357	B1	4/2003	Kaiser et al.	9,637,792	B2	5/2017	Ahlquist et al.
6,562,611	B1	5/2003	Kaiser et al.	9,657,511	B2	5/2017	Lidgard et al.
6,602,695	B2	8/2003	Patel et al.	9,726,670	B2	8/2017	Ataman-onal et al.
6,605,451	B1	8/2003	Marmaro et al.	10,011,878	B2	7/2018	Ahlquist et al.
6,635,463	B2	10/2003	Ma et al.	10,030,272	B2	7/2018	Ahlquist et al.
6,673,616	B1	1/2004	Dahlberg et al.	10,292,687	B2	5/2019	Maguire et al.
6,692,917	B2	2/2004	Neri et al.	10,327,742	B2	6/2019	Fitzgerald et al.
6,706,471	B1	3/2004	Brow et al.	10,370,726	B2	8/2019	Ahlquist et al.
6,709,815	B1	3/2004	Dong et al.	10,385,406	B2	8/2019	Allawi et al.
6,709,819	B2	3/2004	Lyamichev et al.	10,435,755	B2	10/2019	Ahlquist et al.
6,759,226	B1	7/2004	Ma et al.	10,465,248	B2	11/2019	Allawi et al.
6,780,585	B1	8/2004	Dong et al.	10,519,510	B2	12/2019	Ahlquist et al.
6,780,982	B2	8/2004	Lyamichev et al.	10,604,793	B2	3/2020	Oldham-Haltom et al.
6,872,816	B1	3/2005	Hall et al.	10,648,025	B2	5/2020	Allawi et al.
6,875,572	B2	4/2005	Prudent et al.	10,648,035	B2	5/2020	Agarwal et al.
6,913,881	B1	7/2005	Aizenstein et al.	10,704,081	B2	7/2020	Lidgard et al.
6,932,943	B1	8/2005	Cracauer et al.	10,822,638	B2	11/2020	Allawi et al.
7,011,944	B2	3/2006	Prudent et al.	11,118,228	B2	9/2021	Allawi et al.
7,037,650	B2	5/2006	Gonzalzo et al.	2002/0128465	A1	9/2002	Lyamichev et al.
7,045,289	B2	5/2006	Allawi et al.	2002/0142454	A1	10/2002	Cracauer et al.
7,060,436	B2	6/2006	Lyamichev et al.	2002/0156255	A1	10/2002	Cracauer et al.
7,067,643	B2	6/2006	Dahlberg et al.	2002/0198693	A1	12/2002	Marusich et al.
7,087,381	B2	8/2006	Dahlberg et al.	2003/0072689	A1	4/2003	Cracauer et al.
7,087,414	B2	8/2006	Gerdes et al.	2003/0082544	A1	5/2003	Fors et al.
7,101,672	B2	9/2006	Dong et al.	2003/0092039	A1	5/2003	Olson-Munoz et al.
7,122,364	B1	10/2006	Lyamichev et al.	2003/0104378	A1	6/2003	Allawi et al.
7,150,982	B2	12/2006	Allawi et al.	2003/0104470	A1	6/2003	Fors et al.
7,195,871	B2	3/2007	Lyamichev et al.	2003/0113236	A1	6/2003	Cracauer et al.
7,256,020	B2	8/2007	Lyamichev et al.	2003/0113237	A1	6/2003	Cracauer et al.
7,273,696	B2	9/2007	Dahlberg et al.	2003/0124526	A1	7/2003	Cracauer et al.
7,297,780	B2	11/2007	Skrzypczynski et al.	2003/0134349	A1	7/2003	Ma et al.
7,306,917	B2	12/2007	Prudent et al.	2003/0143535	A1	7/2003	Lyamichev et al.
7,312,033	B2	12/2007	Accola et al.	2003/0165954	A1	9/2003	Katagiri et al.
7,354,708	B2	4/2008	Hall et al.	2003/0186238	A1	10/2003	Allawi et al.
7,381,530	B2	6/2008	Hall et al.	2003/0219784	A1	11/2003	Ip et al.
				2003/0224040	A1	12/2003	Baylin et al.
				2003/0224437	A1	12/2003	Gerdes et al.
				2004/0014067	A1	1/2004	Lyamichev et al.
				2004/0018489	A1	1/2004	Ma et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2004/0072182 A1	4/2004	Lyamichev et al.	2009/0203018 A1	8/2009	Agarwal et al.
2004/0096874 A1	5/2004	Neville et al.	2009/0215043 A1	8/2009	Kwitek et al.
2004/0175733 A1	9/2004	Andersen et al.	2009/0253142 A1	10/2009	Allawi et al.
2004/0203035 A1	10/2004	Mast et al.	2009/0299641 A1	12/2009	Allawi et al.
2004/0219576 A1	11/2004	Skrzypczynski et al.	2009/0305283 A1	12/2009	Prudent et al.
2004/0234960 A1	11/2004	Olek et al.	2010/0075334 A1	3/2010	Kim et al.
2004/0235024 A1	11/2004	Lyamichev et al.	2010/0152431 A1	6/2010	Skrzypczynski et al.
2005/0048527 A1	3/2005	Allawi et al.	2010/0304444 A1	12/2010	Morley
2005/0074788 A1	4/2005	Dahlberg et al.	2011/0009277 A1	1/2011	Devos et al.
2005/0106596 A1	5/2005	Skrzypczynski et al.	2011/0123990 A1	5/2011	Baker et al.
2005/0130179 A1	6/2005	Lyamichev et al.	2011/0160446 A1	6/2011	Ritt et al.
2005/0158716 A1	7/2005	Dahlberg et al.	2011/0287424 A1	11/2011	Chen
2005/0164177 A1	7/2005	Neri et al.	2011/0318738 A1	12/2011	Jones et al.
2005/0181435 A1	8/2005	Prudent et al.	2012/0122088 A1	5/2012	Zou et al.
2005/0186588 A1	8/2005	Lyamichev et al.	2012/0122105 A1	5/2012	Oldham-Haltom et al.
2005/0196750 A1	9/2005	Elagin et al.	2012/0122106 A1	5/2012	Zou et al.
2005/0214926 A1	9/2005	Zielenski et al.	2012/0288868 A1	11/2012	Bruinsma et al.
2005/0239101 A1	10/2005	Sukumar et al.	2013/0231256 A1	9/2013	Oldham-Haltom et al.
2005/0277138 A1	12/2005	Skrzypczynski et al.	2013/0296738 A1	11/2013	Swain et al.
2006/0134663 A1	6/2006	Harkin et al.	2014/0087382 A1	3/2014	Allawi et al.
2006/0147938 A1	7/2006	Accola et al.	2015/0292029 A1	10/2015	Agarwal et al.
2006/0147955 A1	7/2006	Allawi et al.	2016/0010081 A1	1/2016	Allawi et al.
2006/0160074 A1	7/2006	Dorn et al.	2016/0081671 A1	3/2016	Lubinski et al.
2006/0171952 A1	8/2006	Mather et al.	2016/0090634 A1	3/2016	Kisiel et al.
2006/0183207 A1	8/2006	Lyamichev et al.	2016/0168643 A1	6/2016	Ahlquist et al.
2006/0198709 A1	9/2006	Marusich et al.	2016/0194721 A1	7/2016	Allawi et al.
2006/0199202 A1	9/2006	Lyamichev et al.	2016/0237480 A1	8/2016	Oldham-Haltom et al.
2006/0234252 A1	10/2006	Andersen	2016/0312299 A1	10/2016	Tyler et al.
2006/0240452 A1	10/2006	Skrzypczynski et al.	2016/0333424 A1	11/2016	Morris et al.
2006/0246475 A1	11/2006	Peterson et al.	2016/0340740 A1	11/2016	Zhang
2006/0252032 A1	11/2006	Aslanukov et al.	2017/0121704 A1	5/2017	Allawi et al.
2007/0048748 A1	3/2007	Williams et al.	2017/0121757 A1	5/2017	Lidgard et al.
2007/0049745 A1	3/2007	Skrzypczynski et al.	2017/0321286 A1	11/2017	Allawi et al.
2007/0087345 A1	4/2007	Olson-Munoz et al.	2017/0335401 A1	11/2017	Allawi et al.
2007/0111200 A1	5/2007	Hudson et al.	2018/0143198 A1	5/2018	Wen et al.
2007/0134249 A1	6/2007	Denney et al.	2018/0245157 A1	8/2018	Allawi et al.
2007/0161062 A1	7/2007	Tacke et al.	2019/0085406 A1	3/2019	Mortimer et al.
2007/0190540 A1	8/2007	Stanley	2019/0177769 A1	6/2019	Allawi et al.
2007/0202517 A1	8/2007	Agarwal et al.	2019/0218601 A1	7/2019	Allawi et al.
2007/0202525 A1	8/2007	Quake et al.	2019/0330702 A1	10/2019	Allawi et al.
2007/0207455 A1	9/2007	Law et al.	2020/0248233 A1	8/2020	Allawi et al.
2007/0292856 A1	12/2007	Lyamichev et al.	2020/0291458 A1	9/2020	Lidgard et al.
2008/0003571 A1	1/2008	Mckernan et al.	2022/0071605 A1	3/2022	Eisele et al.
2008/0014124 A1	1/2008	Skrzypczynski et al.	2022/0349009 A1	11/2022	Taylor et al.
2008/0015349 A1	1/2008	Skrzypczynski et al.	2023/0046033 A1	2/2023	Zubin et al.
2008/0032305 A1	2/2008	Dorn et al.			
2008/0071074 A1	3/2008	Skrzypczynski et al.			
2008/0131870 A1	6/2008	Allawi et al.			
2008/0131875 A1	6/2008	Hall et al.			
2008/0131890 A1	6/2008	Allawi et al.			
2008/0160524 A1	7/2008	Ma et al.			
2008/0176215 A1	7/2008	Hudson et al.			
2008/0181823 A1	7/2008	Iszczyszyn et al.			
2008/0182254 A1	7/2008	Hall et al.			
2008/0182980 A1	7/2008	Skrzypczynski et al.			
2008/0187919 A1	8/2008	King et al.			
2008/0187926 A1	8/2008	Dahlberg et al.			
2008/0188375 A1	8/2008	Neri et al.			
2008/0199936 A1	8/2008	Lyamichev et al.			
2008/0213767 A1	9/2008	Western et al.			
2008/0220425 A1	9/2008	Ma et al.			
2008/0226660 A1	9/2008	Bryan et al.			
2008/0261220 A1	10/2008	Cracauer et al.			
2008/0268455 A1	10/2008	Hall et al.			
2008/0293046 A1	11/2008	Allawi et al.			
2009/0029869 A1	1/2009	Skrzypczynski et al.			
2009/0041634 A1	2/2009	Cracauer et al.			
2009/0068664 A1	3/2009	Lyamichev et al.			
2009/0075256 A1	3/2009	Lyamichev et al.			
2009/0078574 A1	3/2009	Lyamichev et al.			
2009/0111092 A1	4/2009	Elagin et al.			
2009/0117576 A1	5/2009	Dong et al.			
2009/0142752 A1	6/2009	Hall et al.			
2009/0142754 A1	6/2009	Allawi et al.			
2009/0203011 A1	8/2009	Liebenberg et al.			

FOREIGN PATENT DOCUMENTS

JP	2008502890	A	1/2008	
JP	2010533853	A	10/2010	
KR	1020160128136	A	11/2016	
WO	WO 1992002258	A1	2/1992	
WO	WO 1993010820	A1	6/1993	
WO	WO 1994022892	A1	10/1994	
WO	WO 1994024144	A2	10/1994	
WO	WO 1995000669	A1	1/1995	
WO	WO 1995015373	A2	6/1995	
WO	WO 1997046705	A1	12/1997	
WO	WO 1999028498	A2	6/1999	
WO	WO 2001094634	A2	12/2001	
WO	WO 2002070755	A2	9/2002	
WO	WO 2005023091	A2	3/2005	
WO	WO 2005038041	A2	4/2005	
WO	WO 2005038051	A2	4/2005	
WO	WO 2005098050	A2	10/2005	
WO	WO-2005123913	A2	12/2005 C12N 9/1241
WO	WO 2005124356	A2	12/2005	
WO	WO 2006084132	A2	8/2006	
WO	WO 2006113770	A1	10/2006	
WO	WO 2007121489	A2	10/2007	
WO	WO 2008084219	A1	7/2008	
WO	WO 2008100913	A2	8/2008	
WO	WO 2008103763	A2	8/2008	
WO	WO 2009035447	A1	3/2009	
WO	WO 2009102788	A2	8/2009	
WO	WO 2009155271		12/2009	
WO	WO 2010074924	A1	7/2010	
WO	WO 2010086389	A1	8/2010	
WO	WO 2010089538	A2	8/2010	

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	WO 2011058316	A1	5/2011
WO	WO 2011119934	A2	9/2011
WO	WO 2012067831		5/2012
WO	WO 2012155072	A2	11/2012
WO	WO 2013058868	A2	4/2013
WO	WO 2013070950	A1	5/2013
WO	WO 2013116375	A1	8/2013
WO	WO 2013142545	A1	9/2013
WO	WO 2014039556	A1	3/2014
WO	WO 2014062218	A1	4/2014
WO	WO 2014159650	A2	10/2014
WO	WO 2014159652	A2	10/2014
WO	WO 2014160117	A1	10/2014
WO	WO 2015066695	A1	5/2015
WO	WO 2015153283	A1	10/2015
WO	WO 2015153284	A1	10/2015
WO	WO 2016094813	A1	6/2016
WO	WO 2016094839	A2	6/2016
WO	WO 2017040627	A1	3/2017
WO	WO 2017075061	A1	5/2017
WO	WO 2017129716	A1	8/2017
WO	WO 2017180886	A1	10/2017
WO	WO 2017192221	A1	11/2017
WO	WO 2017223216	A1	12/2017
WO	WO 2018017740	A1	1/2018
WO	WO 2018045322	A1	3/2018
WO	WO 2018160576	A1	9/2018
WO	WO 2019108626	A1	6/2019
WO	WO 2020112869	A1	6/2020
WO	WO 2020154665	A1	7/2020
WO	WO 2020206256	A1	10/2020
WO	WO 2021041726	A1	3/2021
WO	WO 2021055508	A1	3/2021
WO	WO 2021076969	A1	4/2021
WO	WO 2021087275	A1	5/2021
WO	WO 2021212031	A1	10/2021
WO	WO 2021226071	A2	11/2021
WO	WO 2021226074	A2	11/2021
WO	WO 2022039904	A2	2/2022
WO	WO 2022040306	A1	2/2022
WO	WO 2022165247	A1	8/2022
WO	WO 2022187695	A1	9/2022
WO	WO 2023081796	A1	5/2023

OTHER PUBLICATIONS

Hecker, et al., "High and low annealing temperatures increase both specificity and yield in touchdown and stepdown PCR", *Biotechniques*, 20(3):478-85, 2006.

Herman, et al., "Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands", *Proc Natl Acad Sci.*, 1996, 93:9821-6.

Hosono et al., "Multiplex PCR-Based Real-Time Invader Assay (mPCR-RETINA): A Novel SNP-Based Method for Detecting Allelic Asymmetries Within Copy Number Variation Regions", *Human Mutation*, 2007, 0:1-8.

Itzkowitz, et al., "A simplified, noninvasive stool DNA test for colorectal cancer detection", *Am J Gastroenterol.*, 2008, 103:2862-70.

Itzkowitz, et al., "Improved fecal DNA test for colorectal cancer screening", *Clin Gastroenterol Hepatol.*, 2007, 5:111-7.

Kearns et al., (1999) "Rapid detection of methicillin-resistant staphylococci by multiplex PCR" *J Hosp Infect*, pp. 33-37.

Layton et al., "Development of Bacteroides 16S rRNA gene TaqMan-based real-time PCR assays for estimation of total, human, and bovine fecal pollution in water" *Appl Environ Microbiology*, 72:6 pp. 4214-4224.

Li, et al., "COLD-PCR: a new platform for highly improved mutation detection in cancer and genetic testing", *Biochem Soc Trans.*, 37(Pt 2):427-32, 2009.

Lindh, et al. (2006) "Real-time Taqman PCR targeting 14 human papilloma virus types", *J Clin Virol*. Dec. 2007;40(4):321-4. Epub Nov. 5, 2007.

Melo, et al. "Development of molecular approaches for the identification of transmission sites of schistosomiasis", *Trans R Soc Trop Med Hyg*. Nov. 2006; 1 00(11): 1 049-55. Epub Apr. 18, 2006.

Minarovicova, et al. "A single-tube nested real-time polymerase chain reaction for sensitive contained detection of *Cryptosporidium parvum*", *Lett Appl Microbiol*. Nov. 2009; 49(5):568-72. Epub Jul. 31, 2009.

Nitsche et al., (1999) "Different real-time PCR formats compared for the quantitative detection of human cytomegalovirus DNA" *Clin Chem*, 45:11 pp. 1932-1937.

Parsons, et al., "Genotypic selection methods for the direct analysis of point mutations", *Mutat Res.*, 387(2):97-121, 1997.

Penders et al., (2005) "Quantification of *Bifidobacterium* spp., *Escherichia coli* and *Clostridium difficile* in faecal samples of breast-fed and formula-fed infants by real-time PCR" *FEMS Microbiol Lett*. 243:1 pp. 141-147.

Qiagen, "EpiTect® MethyLight PCR Handbook", MethyLight PCR Kit, MethyLight PCR + ROX Vial Kit, 2008, 36pgs.

Roux, et al., "One-step optimization using touchdown and stepdown PCR", *Methods Mol Biol*, 67:39-45, 1997.

Supplemental European Search Report for European Patent Application No. EP11841038, Document dated Apr. 4, 2014, 8 pages.

Tabone et al. (2009) "Temperature Switch PCR (TSP): Robust assay design for reliable amplification and genotyping of SNPs", *BMC Genomics* 10:580.

Tadokoro, et al., "Quantitation of viral load by real-time PCR-monitoring Invader reaction", *J Virol Methods.*, 2009, 155:182-6.

Yamada, et al., "Fluorometric identification of 5-methylcytosine modification in DNA: combination of photosensitized oxidation and invasive cleavage", *Bioconjug Chem.*, 2008, 19:20-3.

Zymo Research Corp., "EZ DNA Methylation-Gold™ Kit", Flyer, Catalog Nos. D5005 & D5006, Ver. 2.1.0, downloaded Feb. 23, 2011, 2pgs.

Zymo Research Corp., "EZ DNA Methylation-Gold™ Kit", Instructions, Catalog Nos. D5005 & D5006, Ver. 2.1.0, downloaded Feb. 23, 2011, 10pgs.

Zymo Research Corp., "EZ DNA Methylation™ Kit", Instruction Manual, Catalog Nos. D5001 & D5002, Ver. 1.2.2, downloaded Feb. 23, 2011, 10pgs.

Zymo Research Corp., "Material Safety Data Sheet", MSDS: CT Conversion Reagent, Creation Date: Apr. 28, 2003, Revision Date: May 4, 2009, 1-4.

International Preliminary Report on Patentability for PCT/US2020/048270, mailed Mar. 10, 2022, 11 pages.

International Search Report and Written Opinion for PCT/US2016/058875, mailed Apr. 21, 2017, 15 pages.

International Search Report and Written Opinion for PCT/US2017/024468, mailed Sep. 1, 2017, 17 pages.

International Search Report and Written Opinion for PCT/US2018/015535, mailed Jun. 25, 2018, 20 pages.

International Search Report and Written Opinion for PCT/US2019/063401, mailed Feb. 20, 2020, 12 pages.

International Search Report and Written Opinion for PCT/US2020/048270, mailed Dec. 7, 2020, 13 pages.

European Supplemental Search Report for EP17792973.4, mailed Jan. 3, 2020, 15 pages.

European Supplemental Search Report for EP18744801.4, mailed Dec. 14, 2020, 23 pages.

Extended European Search Report for EP 19890483.1, mailed Sep. 29, 2022, 10 pages.

Extended European Search Report for EP21195952.3, mailed Apr. 12, 2022, 8 pages.

Certified copy of U.S. Appl. No. 60/900,713, filed Feb. 12, 2007, in the name of Baylin et al., 188 pages.

Notice of opposition and statement filed in EP Patent 3434791, filed on Mar. 5, 2021, 16 pages.

Ahlquist et al., "Colorectal cancer screening by detection of altered human DNA in stool: feasibility of a multitarget assay panel", *Gastroenterology*, 2000, 119: 1219-1227.

Ahlquist et al., "Next-generation stool DNA test accurately detects colorectal cancer and large adenomas", *Gastroenterology*, 2012, 142: 248-256.

(56)

References Cited**OTHER PUBLICATIONS**

- Ahlquist et al., "Novel use of hypermethylated DNA markers in stool for detection of colorectal cancer: a feasibility study", *Gastroenterology*, 2002, 122(suppl a40): 2 pages.
- Ahlquist et al., "Stool DNA and occult blood testing for screen detection of colorectal neoplasia", *Annals of Internal Medicine*, 2008, 149(7): 441-450.
- Allawi et al., "Abstract 712: Detection of lung cancer by assay of novel methylated DNA markers in plasma", *Cancer Research*, Proceedings: AACR Annual Meeting Apr. 1-5, 2017, Washington, DC, 3 pages.
- Anderson et al., "Methylated DNA markers for detection of sporadic colorectal neoplasia: comparison between age groups younger than and older than 50", *Abstract Su2013, Gastroenterology*, Apr. 1, 2016, 150(4): S-611.
- Andersson et al., "Properties of targeted preamplification in DNA and cDNA quantification", *Expert Review of Molecular Diagnostics*, 2015, 15(8): 1085-1100.
- Antequera et al., "High levels of de novo methylation and altered chromatin structure at CpG islands in cell lines", *Cell*, Aug. 10, 1990, 62(3): 503-514.
- Arneson et al., "Genomeplex whole-genome amplification", *Cold Spring Harbor Protocols*, Jan. 1, 2008, 2008(2): 7 pages.
- Aronchick et al., "A novel tableted purgative for colonoscopic preparation: efficacy and safety comparisons with colyte and fleet phospho-soda", *Gastrointestinal Endoscopy*, 2000, 52(3): 346-352.
- Ballabio et al., "Screening for steroid sulfatase (STS) gene deletions by multiplex DNA amplification", *Human Genetics*, 1990, 84(6): 571-573.
- Ballester et al., "Novel methylated DNA markers for the detection of colorectal neoplasia in lynch syndrome", *Abstract 307, Gastroenterology*, 2016, 150(4): S-70.
- Barany, "Genetic disease detection and DNA amplification using cloned thermostable ligase", *Proceedings of the National Academy of Sciences of the USA*, Jan. 1991, 88: 189-193.
- Bardan E et al., "Colonoscopic resection of large colonic polyps—a prospective study", *Israel Journal of Medical Sciences*, 1997, 33(12): 777-780.
- Belinsky et al., "Promoter hypermethylation of multiple genes in sputum precedes lung cancer incidence in a high-risk cohort", *Cancer Research*, Mar. 15, 2006, 66(6): 3338-3344.
- Bentley et al., "Accurate whole human genome sequencing using reversible terminator chemistry", *Nature*, Nov. 6, 2008, 456(7218): 53-59.
- Berezikov et al., "Approaches to microRNA discovery", *Nature Genetics*, Jun. 2006, 38: S2-S7.
- Berger et al., "Stool DNA screening for colorectal neoplasia: biological and technical basis for high detection rates", *Pathology*, Feb. 2012, 44(2): 80-88.
- Bibikova, "Goldengate® assay for methylation of beadarraytm technology", *Illumina, Technical Notes: Epigenetic Analysis*, Jan. 1, 2009, 6 pages.
- Boynton et al., "DNA integrity as a potential marker for stool-based detection of colorectal cancer", *Clinical Chemistry*, 2003, 49(7): 1058-1065.
- Budd et al., "Circulating tumor cells versus imaging-predicting overall survival in metastatic breast cancer", *Clinical Cancer Research*, Nov. 1, 2006, 12(21): 6403-6409.
- Bustin, "Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays", *Journal of Molecular Endocrinology*, 2000, 25: 169-193.
- Carvalho et al., "Genome-wide DNA methylation profiling of non-small cell lung carcinomas", *Epigenetics Chromatin*, Jun. 22, 2012, 5(9): 18 pages.
- Ceska et al., "Structure-specific DNA cleavage by 5' nucleases", *Trends in Biochemical Sciences*, Sep. 1998, 23(9): 331-336.
- Chamberlain et al., "Deletion screening of the duchenne muscular dystrophy locus via multiplex DNA amplification", *Nucleic Acids Research*, 1988, 16(23): 11141-11156.
- Chapman et al., "Autoantibodies in lung cancer: possibilities for early detection and subsequent cure", *Thorax*, Mar. 2008, 63(3): 228-233.
- Chen et al., "Detection in fecal DNA of colon cancer-specific methylation of the nonexpressed vimentin", *Journal of the National Cancer Institute*, 2005, 97(15): 1124-1132.
- Chen et al., "DNA methylation identifies loci distinguishing hereditary nonpolyposis colorectal cancer without germ-line MLH1/MSH2 mutation from sporadic colorectal cancer", *Clinical and Translational Gastroenterology*, Dec. 15, 2016, 7(12): e208.
- Chen et al., "HOPX is methylated and exerts tumour-suppressive function through Ras-induced senescence in human lung cancer", *The Journal of Pathology*, Feb. 2015, 235(3): 397-407.
- Cohen et al., "Relationship of circulating tumor cells to tumor response, progression-free survival, and overall survival in patients with metastatic colorectal cancer", *Journal of Clinical Oncology*, Jul. 1, 2008, 26(19): 3213-3221.
- Cristofanilli et al., "Circulating tumor cells, disease progression, and survival in metastatic breast cancer", *The New England Journal of Medicine*, Aug. 19, 2004, 351(8): 781-791.
- Dammann et al., "The CpG island of the novel tumor suppressor gene RASSF1A is intensely methylated in primary small cell lung carcinomas", *Oncogene*, Jun. 14, 2001, 20(27): 3563-3567.
- Dassonville et al., "Expression of epidermal growth factor receptor and survival in upper aerodigestive tract cancer", *Journal of Clinical Oncology*, Oct. 1993, 11(10): 1873-1878.
- Devos et al., "Circulating methylated SEPT9 DNA in plasma is a biomarker for colorectal cancer", *Clinical Chemistry*, Jul. 2009, 55(7): 1337-1346.
- Don et al., "'Touchdown' PCR to circumvent spurious priming during gene amplification", *Nucleic Acids Research*, 1991, 19(14): 4008.
- Dowdy et al., "Statistics for Research", *John Wiley & Sons*, New York, 1983, TOC only, 6 pages.
- Eads et al., "CpG island hypermethylation in human colorectal tumors is not associated with DNA methyltransferase overexpression", *Cancer Research*, May 15, 1999, 59(10): 2302-2306.
- Ebert et al., "Aristaless-like homeobox-4 gene methylation is a potential marker for colorectal adenocarcinoma", *Gastroenterology*, 2006, 131: 1418-1430.
- Egeblad et al., "New functions for the matrix metalloproteinases in cancer progression", *Nature Reviews Cancer*, Mar. 2002, 2(3): 161-174.
- Elbashir et al., "RNA interference is mediated by 21- and 22-nucleotide RNAs", *Genes & Development*, Jan. 15, 2001, 15(2): 188-200.
- Fackler et al., "Quantitative multiplex methylation-specific PCR assay for the detection of promoter hypermethylation in multiple genes in breast cancer", *Cancer Research*, Jul. 1, 2004, 64(13): 4442-4452.
- Fasman, "Practical handbook of biochemistry and molecular biology", 1989, CRC Press, Boca Raton, FL, pp. 385-394.
- Fedurco et al., "BTA, a novel reagent for DNA attachment on glass and efficient generation of solid-phase amplified DNA colonies", *Nucleic Acids Research*, Feb. 9, 2006, 34(3)e22: 13 pages.
- Feil et al., "Methylation analysis on individual chromosomes: improved protocol for bisulphite genomic sequencing", *Nucleic Acids Research*, Feb. 25, 1994, 22(4): 695-696.
- Feng et al., "Genome-wide analysis of DNA methylation and their associations with long noncoding RNA/mRNA expression in non-small-cell lung cancer", *Epigenomics*, Jan. 2017, 9(2): 137-153.
- Finger et al., "The wonders of flap endonucleases: structure, function, mechanism and regulation", *Subcellular Biochemistry*, 2012, 62: 301-326.
- Frommer et al., "A genomic sequencing protocol that yields a positive display of 5-methylcytosine residues in individual DNA strands", *Proceedings of the National Academy of Sciences of the USA*, Mar. 1, 1992, 89(5): 1827-1831.
- Gardiner-Garden et al., "CpG Islands in vertebrate genomes", *Journal of Molecular Biology*, 1987, 196: 261-282.
- Gatlin et al., "Automated identification of amino acid sequence variations in proteins by HPLC/microspray tandem mass spectrometry", *Analytical Chemistry*, Feb. 15, 2000, 72(4): 757-763.

(56)

References Cited

OTHER PUBLICATIONS

- Gemperle et al., "Regulation of the formyl peptide receptor 1 (FPR1) gene in primary human macrophages", PLoS One, Nov. 2012, 7(11):e50195: 6 pages.
- Genebank, "*Homo sapiens* formyl peptide receptor 1 (FPR1), transcript variant 1, mRNA", National Library of Medicine, GenBank Accession No. NM_001193306.1, Apr. 9, 2019, 5 pages.
- Genebank, "*Homo sapiens* S100 calcium binding protein A12 (S100A12), Mrna", National Library of Medicine, GenBank Accession No. NM005621, Apr. 23, 2024, 3 pages.
- Genpept, "protein S100-A12 [*Homo sapiens*]", GenPeptAccession No. NP05612, Sep. 11, 2022, 3 pages.
- Gevaert et al., "Pancancer analysis of DNA methylation-driven genes using MethylMix", Genome Biology, Jan. 29, 2015, 16(17): 13 pages.
- Gonzalzo et al., "Identification and characterization of differentially methylated regions of genomic DNA by methylation-sensitive arbitrarily primed PCR", Cancer Research, Feb. 15, 1997, 57(4): 594-599.
- Gonzalzo et al., "Rapid quantitation of methylation differences at specific sites using methylation-sensitive single nucleotide primer extension (Ms-SNuPE)", Nucleic Acids Research, Jun. 15, 1997, 25(12): 2529-2531.
- Grady et al., "Detection of aberrantly methylated hMLHI promoter dna in the serum of patients with microsatellite unstable colon cancer", Cancer Research, 2001, 61: 900-902.
- Grafstrom et al., "The characteristics of DNA methylation in an in vitro DNA synthesizing system from mouse fibroblasts", Nucleic Acids Research, 1985, 13(8): 2827-2842.
- Grandis et al., "TGF-alpha and EGFR in head and neck cancer", Journal of Cellular Biochemistry, 1993, Supplement 17f: 188-191.
- Grigg et al., "Sequencing 5-methylcytosine residues in genomic DNA", BioEssays, Jun. 1994, 16(6): 431-436.
- Grigg, "Sequencing 5-methylcytosine residues by the bisulphite method", DNA sequencing, 1996, 6(4): 189-198.
- Grunau et al., "Bisulfite genomic sequencing: systematic investigation of critical experimental parameters", Nucleic Acids Research, Jul. 1, 2001, 29(13): E65: 7 pages.
- Gu et al., "Genome-scale DNA methylation mapping of clinical samples at single-nucleotide resolution", Nature Methods, Feb. 2010, 7(2): 133-136.
- Guilfoyle et al., "Ligation-mediated PCR amplification of specific fragments from a class-II restriction endonuclease total digest", Nucleic Acids Research, 1997, 25: 1854-1858.
- Hall et al., "Sensitive detection of DNA polymorphisms by the serial invasive signal amplification reaction", Proceedings of the National Academy of Sciences of the USA, 2000, 97(15): 8272-8277.
- Hardcastle et al., "Randomised controlled trial of faecal-occult-blood screening for colorectal cancer", The Lancet, 1996, 348: 1472-1477.
- Harris et al., "Single-molecule DNA sequencing of a viral genome", Science, Apr. 4, 2008, 320(5872): 106-109.
- Hayden et al., "Multiplex-Ready PCR: A new method for multiplexed SSR and SNP genotyping", BMC Genomics, 2008, 9(80): 12 pages.
- Hayes et al., "Circulating tumor cells at each follow-up time point during therapy of metastatic breast cancer patients predict progression-free and overall survival", Clinical Cancer Research, Jul. 15, 2006, 12(14): 4218-4224.
- He et al., "Development of a multiplex Methylight assay for the detection of multigene methylation in human colorectal cancer", Cancer Genetics and Cytogenetics, 2010, 202(1): 1-10.
- Heid et al., "Real time quantitative PCR", Genome Research, Oct. 1996, 6(10): 986-994.
- Heitman et al., "Colorectal cancer screening for average-risk north americans: an economic evaluation", PLoS Medicine, Nov. 2010, 7(11):e1000370: 13 pages.
- Heller et al., "Lung cancer: from single-gene methylation to methylome profiling", Cancer and Metastasis Reviews, Mar. 2010, 29(1): 95-107.
- Henegariu et al., "Multiplex PCR: critical parameters and step-by-step protocol", Biotechniques, Sep. 1997, 23(3): 504-511.
- Heresbach et al., "Review in depth and meta-analysis of controlled trials on colorectal cancer screening by faecal occult blood test", European Journal of Gastroenterology & Hepatology, 2006, 18(4): 427-433.
- Higuchi et al., "A general method of in vitro preparation and specific mutagenesis of DNA fragments: study of protein and DNA interactions", Nucleic Acids Research, 1988, 16(15): 7351-7367.
- Higuchi et al., "Kinetic PCR analysis: real-time monitoring of DNA amplification reactions", Biotechnology, 1993, 11: 1026-1030.
- Higuchi et al., "Simultaneous amplification and detection of specific DNA sequences", Biotechnology, 1992, 10: 413-417.
- Hoque et al., "Genome-wide promoter analysis uncovers portions of the cancer methylome", Cancer Research, Apr. 15, 2008, 68(8): 2661-2670.
- Hua et al., "Quantitative methylation analysis of multiple genes using methylation-sensitive restriction enzyme-based quantitative PCR for the detection of hepatocellular carcinoma", Experimental and Molecular Pathology, Aug. 2011, 91(1): 455-60.
- Huang et al., "Transactivation of the epidermal growth factor receptor by formylpeptide receptor exacerbates the malignant behavior of human glioblastoma cells", Cancer Research, Jun. 15, 2007, 67(12): 5906-5913.
- Imperiale et al., "Fecal DNA versus fecal occult blood for colorectal-cancer screening in an average-risk population", The New England Journal of Medicine, 2004, 351: 2704-2714.
- Iyer et al., "Accurate nonendoscopic detection of barrett's esophagus by methylated DNA markers: a multisite case control study", The American Journal of Gastroenterology, 2020, 115: 1201-1209.
- Iyer et al., "Accurate non-endoscopic detection of barrett's esophagus in a multicenter prospective validation cohort: the sos 2 trial", AGA Abstracts, Gastroenterology, 2018, 878:S-175-S-176.
- Iyer et al., "Independent validation of an accurate methylated DNA marker panel for the non-endoscopic detection of Barrett's esophagus: a multisite case control study", AGA Abstracts, 2020, 1084: S-211.
- Jiang et al., "Lengthening and shortening of plasma DNA in hepatocellular carcinoma patients", The Proceedings of the National Academy of Sciences, Feb. 2, 2015, 112(11): E1317-E1325.
- Jongeneel et al., "An atlas of human gene expression from massively parallel signature sequencing (MPSS)", Genome Research, Jul. 2005, 15(7): 1007-1014.
- Kaiser et al., "A comparison of eubacterial and archaeal structure-specific 5'-exonucleases", Journal of Biological Chemistry, Jul. 23, 1999, 274(30): 21387-21394.
- Kalinina et al., "Nanoliter scale PCR with TaqMan detection", Nucleic Acids Research, 1997, 25: 1999-2004.
- Kann et al., "Improved marker combination for detection of de novo genetic variation and aberrant DNA in colorectal neoplasia", Clinical Chemistry, 2006, 52: 2299-2302.
- Karl et al., "Improved diagnosis of colorectal cancer using a combination of fecal occult blood and novel fecal protein markers", Clinical Gastroenterology and Hepatology, 2008, 6(10): 1122-1128.
- Kawai et al., "Comparison of DNA methylation patterns among mouse cell lines by restriction landmark genomic scanning", Molecular and Cellular Biology, Nov. 1994, 14(11): 7421-7427.
- Kisiel et al., "New DNA methylation markers for pancreatic cancer: discovery, tissue validation, and pilot testing in pancreatic juice", Clinical Cancer Research, Oct. 1, 2015, 21(19): 4473-4481.
- Kisiel et al., "Novel stool DNA markers for inflammatory bowel disease associated colorectal cancer high grade dysplasia: high specificity across three independent international populations", Gastroenterology, Abstract 185, 2016, 150(4): S-48.
- Kling, "Ultrafast DNA sequencing", Nature Biotechnology, Dec. 2003, 21(12): 1425-1427.
- Kneip et al., "SHOX2 DNA methylation is a biomarker for the diagnosis of lung cancer in plasma", Journal of Thoracic Oncology, Oct. 2011, 6(10): 1632-1638.
- Kober et al., "Methyl-CpG binding column-based identification of nine genes hypermethylated in colorectal cancer", Molecular Carcinogenesis, Nov. 2011, 50(11): 846-856.

(56)

References Cited**OTHER PUBLICATIONS**

- Korbie et al., "Multiplex bisulfite PCR resequencing of clinical FFPE DNA", *Clinical Epigenetics*, Mar. 17, 2015, 7(28): 11 pages.
- Kronborg et al., "Randomized study of biennial screening with a faecal occult blood test: results after nine screening rounds", *Scandinavian Journal of Gastroenterology*, 2004, 39: 846-851.
- Kunte et al., "MicroRNAs as novel targets for NSAID chemoprevention of colon carcinogenesis", *Gastroenterology*, May 2011, 140(5): S-41.
- Kuppuswamy et al., "Single nucleotide primer extension to detect genetic diseases: experimental application to hemophilia B (factor IX) and cystic fibrosis genes", *The Proceedings of the National Academy of Sciences*, Feb. 15, 1991, 88(4): 1143-1147.
- Kwiatkowski et al., "Clinical, genetic, and pharmacogenetic applications of the Invader assay, *The Journal of Molecular Diagnostics*", Dec. 1999, 4(4): 353-364.
- Lange et al., "Genome-scale discovery of DNA-methylation biomarkers for blood-based detection of colorectal cancer", *PLoS One*, 2012;7(11):e50266.
- Leontiou et al., "Bisulfite conversion of DNA: performance comparison of different kits and methylation quantitation of epigenetic biomarkers that have the potential to be used in non-invasive prenatal testing", *PLoS One*, Aug. 6, 2015, 10(8):e0135058: 22 pages.
- Leung et al., "Detection of epigenetic changes in fecal DNA as a molecular screening test for colorectal cancer: a feasibility study", *Clinical Chemistry*, 2004, 50(11): 2179-2182.
- Levin et al., "Genetic biomarker prevalence is similar in fecal Immunochemical test positive and negative colorectal cancer tissue", *Digestive Diseases and Sciences*, Mar. 2017, 62(3): 678-688.
- Levin et al., "Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, 2008: a joint guideline from the american cancer society, the US multi-society task force on colorectal cancer, and the american college of radiology", *Gastroenterology*, 2008, 134(5): 1570-1595.
- Liu et al., "Flap endonuclease 1: a central component of DNA metabolism", *Annual Review of Biochemistry*, 2004, 73: 589-615.
- Lokk et al., "Methylation markers of early-stage non-small cell lung cancer", *PLoS One*, Jun. 2012, 7(6):e39813: 9 pages.
- Louwagie et al., "Feasibility of a DNA methylation assay for noninvasive CRC screening", *Clinical Cancer Research*, Oct. 2007, 13(19):B16: 4 pages.
- Lyamichev et al., "Polymorphism identification and quantitative detection of genomic DNA by invasive cleavage of oligonucleotide probes", *Nature Biotechnology*, 1999, 17: 292-296.
- Mandal et al., "Lipopolysaccharide induces formyl peptide receptor 1 gene expression in macrophages and neutrophils via transcriptional and posttranscriptional mechanisms", *Journal of Immunology*, Nov. 1, 2005, 175(9): 6085-6091.
- Mandal et al., "Signaling in lipopolysaccharide-induced stabilization of formyl peptide receptor 1 mRNA in mouse peritoneal macrophages", *Journal of Immunology*, Feb. 15, 2007, 178(4): 2542-2548.
- Mandel et al., "Reducing mortality from colorectal cancer by screening for fecal occult blood", *The New England Journal of Medicine* 1993, 328: 1365-1371.
- Marabella et al., "Serum ribonuclease in patients with lung carcinoma", *Journal of Surgical Oncology*, 1976, 8(6): 501-505.
- Margulies et al., "Genome sequencing in microfabricated high-density picolitre reactors", *Nature*, Sep. 15, 2005, 437(7057): 376-380.
- Martin et al., "Genomic sequencing indicates a correlation between DNA hypomethylation in the 5' region of the pS2 gene and its expression in human breast cancer cell lines", *Gene*, May 19, 1995, 157(1-2): 261-264.
- Medina-Aguilar et al., "Methylation landscape of human breast cancer cells in response to dietary compound resveratrol", *PLoS One*, Jun. 29, 2016, 11(6):e0157866: 12 pages.
- Meissner et al., "Patterns of colorectal cancer screening uptake among both men and women in the united states", *Cancer Epidemiology, Biomarkers & Prevention*, 2006, 15: 389-394.
- Meissner et al., "Reduced representation bisulfite sequencing for comparative high-resolution DNA methylation analysis", *Nucleic Acids Research*, Oct. 13, 2005, 33(18): 5868-5877.
- Melnikov et al., "MSRE-PCR for analysis of gene-specific DNA methylation", *Nucleic Acids Research*, Jun. 8, 2005, 33(10):e93: 7 pages.
- Mitchell et al., "A panel of genes methylated with high frequency in colorectal cancer", *BMC Cancer*, Jan. 31, 2014, 14(54): 15 pages.
- Mitchell et al., "Evaluation of methylation biomarkers for detection of circulating tumor DNA and application to colorectal cancer", *Genes*, Dec. 15, 2016, 7(12):125: 11 pages.
- Monte et al., "Cloning, chromosome mapping and functional characterization of a human homologue of murine gtse-1 (B99) gene", *Gene*, Aug. 22, 2000, 254(1-2): 229-236.
- Monte et al., "hGTSE-1 expression stimulates cytoplasmic localization of p53", *Journal of Biological Chemistry*, Mar. 19, 2004, 279(12): 11744-11752.
- Moon et al., "Identification of novel hypermethylated genes and demethylating effect of vincristine in colorectal cancer", *Journal of Experimental & Clinical Cancer Research*, 2014, 33(4): 10 pages.
- Moreno et al., "Circulating tumor cells predict survival in patients with metastatic prostate cancer", *Urology*, Apr. 2005, 65(4): 713-718.
- Morris et al., "Whole blood FPR1 mRNA expression predicts both non-small cell and small cell lung cancer", *International Journal of Cancer*, Jun. 1, 2018, 142(11): 2355-2362.
- Muller et al., "Methylation changes in faecal DNA: a marker for colorectal cancer screening?", *Lancet*, 2004, 363: 1283-1285.
- Munson et al., "Recovery of bisulfite-converted genomic sequences in the methylation-sensitive QPCR", *Nucleic Acids Research*, 2007, 35(9): 2893-2903.
- Neuwelt et al., "Possible sites of origin of human plasma ribonucleases as evidenced by isolation and partial characterization of ribonucleases from several human tissues", *Cancer Research*, Jan. 1978, 38(1): 88-93.
- Nilsson et al., "Altered DNA methylation and differential expression of genes influencing metabolism and inflammation in adipose tissue from subjects with type 2 diabetes", *Diabetes*, Sep. 2014, 63: 2962-2976.
- Noutsias et al., "Pre-amplification techniques for real-time RT-PCR analyses of endomyocardial biopsies", *BMC Molecular Biology*, Jan. 14, 2008, 9(3): 20 pages.
- Nyce et al., "Variable effects of DNA-synthesis inhibitors upon DNA methylation in mammalian cells", *Nucleic Acids Research*, May 27, 1986, 14(10): 4353-4367.
- O'Driscoll et al., "Feasibility and relevance of global expression profiling of gene transcripts in serum from breast cancer patients using whole genome microarrays and quantitative RT-PCR", *Cancer Genomics Proteomics*, Mar.-Apr. 2008, 5(2): 94-104.
- Olek et al., "A modified and improved method for bisulphite based cytosine methylation analysis", *Nucleic Acids Research*, Dec. 15, 1996, 24(24): 5064-5066.
- Olek et al., "The pre-implantation ontogeny of the H19 methylation imprint", *Nature Genetics*, Nov. 1997, 17(3): 275-276.
- Olivier, "The Invader assay for SNP genotyping", *Mutation Research*, Jun. 3, 2005, 573(1-2): 103-110.
- Olkhov-Mitsel et al., "Novel multiplex methylight protocol for detection of DNA methylation in patient tissues and bodily fluids", *Scientific Reports*, Mar. 21, 2014, 4(4432): 8 pages.
- Ooki et al., "Potential utility of HOP homeobox gene promoter methylation as a marker of tumor aggressiveness in gastric cancer", *Oncogene*, Jun. 3, 2010, 29(22): 3263-3275.
- Orpana, "Fluorescence resonance energy transfer (FRET) using ssDNA binding fluorescent dye", *Biomolecular Engineering*, Apr. 2004, 21(2): 45-50.
- Osborn et al., "Stool screening for colorectal cancer: molecular approaches", *Gastroenterology*, 2005, 128(1): 192-206.

(56)

References Cited**OTHER PUBLICATIONS**

- Osman et al., "Expression of matrix metalloproteinases and tissue inhibitors of metalloproteinases define the migratory characteristics of human monocyte-derived dendritic cells", *Immunology*, Jan. 2002, 105(1): 73-82.
- Pantel et al., "Detection, clinical relevance and specific biological properties of disseminating tumour cells", *Nature Reviews Cancer*, May 2008, 8(5): 329-340.
- Parekh et al., "As tests evolve and costs of cancer care rise: reappraising stool-based screening for colorectal neoplasia", *Alimentary Pharmacology & Therapeutics*, 2008, 27: 697-712.
- Petko et al., "Aberrantly methylated CDKN2A, MGMT, and MLH1 in colon polyps and in fecal DNA from patients with colorectal polyps", *Clinical Cancer Research*, 2005, 11: 1203-1209.
- Ponomaryova et al., "Potentialities of aberrantly methylated circulating DNA for diagnostics and post-treatment follow-up of lung cancer patients", *Lung Cancer*, Sep. 2013, 81(3): 397-403.
- Promega, "Maxwell® RSC ccfDNA plasma kit", Technical Manual, Instructions for Use of Product ASI480, Promega Corporation, Jan. 2021, 12 pages.
- Ramsahoye et al., "Non-CpG methylation is prevalent in embryonic stem cells and may be mediated by DNA methyltransferase 3a", *Proceedings of the National Academy of Sciences*, 2000, 97(10): 5237-5242.
- Reddi et al., "Elevated serum ribonuclease in patients with pancreatic cancer", *The Proceedings of the National Academy of Sciences*, Jul. 1976, 73(7): 2308-2310.
- Rein et al., "Identifying 5-methylcytosine and related modifications in DNA genomes", *Nucleic Acids Research*, May 15, 1998, 26(10): 2255-2264.
- Reinartz et al., "Massively parallel signature sequencing (MPSS) as a tool for in-depth quantitative gene expression profiling in all organisms", *Briefings in Functional Genomics*, Feb. 2002, 1(1): 95-104.
- Rex et al., "American college of gastroenterology guidelines for colorectal cancer screening 2008", *The American Journal of Gastroenterology*, 2009, 104: 739-750.
- Ronaghi et al., "A sequencing method based on real-time pyrophosphate", *Science*, Jul. 17, 1998, 281(5375): 363-365.
- Ronaghi et al., "Real-time DNA sequencing using detection of pyrophosphate release", *Analytical Biochemistry*, Nov. 1, 1996, 242(0432): 84-89.
- Rothberg et al., "An integrated semiconductor device enabling non-optical genome sequencing", *Nature*, Jul. 20, 2011, 475(7356): 348-352.
- Roux, "Using mismatched primer-template pairs in touchdown PCR", *Biotechniques*, 1994, 16(5): 812-814.
- Ruano et al., "Biphasic amplification of very dilute DNA samples via 'booster' PCR", *Nucleic Acids Research*, Jul. 11, 1989, 17(13): 5407.
- Sadri et al., "Rapid analysis of DNA methylation using new restriction enzyme sites created by bisulfite modification", *Nucleic Acids Research*, Dec. 15, 1996, 24(24): 5058-5059.
- Salomon et al., "Methylation of mouse DNA in vivo: di- and tripyrimidine sequences containing 5-methylcytosine", *Biochimica et Biophysica Acta*, Apr. 15, 1970, 204(2): 340-351.
- Santani et al., "Characterization, quantification, and potential clinical value of the epidermal growth factor receptor in head and neck squamous cell carcinomas", *Head & Neck*, 1991, 13(2): 132-139.
- Schmidt et al., "SHOX2 DNA methylation is a biomarker for the diagnosis of lung cancer based on bronchial aspirates", *BMC Cancer*, Nov. 3, 2010, 10(600): 9 pages.
- Schouten et al., "Relative quantification of 40 nucleic acid sequences by multiplex ligation-dependent probe amplification", *Nucleic Acids Research*, 2002, 30(12):e57: 13 pages.
- Schuebel et al., "Comparing the DNA hypermethylome with gene mutations in human colorectal cancer", *PLoS Genetics*, Sep. 2007, 3(9): 1709-1723.
- Schuuring et al., "Characterization of the EMS1 gene and its product, human Contactin", *Cell Adhesion & Communication*, 1998, 6(2-3): 185-209.
- Schuuring et al., "Identification and cloning of two overexpressed genes, U21B31/PRAD1 and EMS1, within the amplified chromosome 11q13 region in human carcinomas", *Oncogene*, Feb. 1992, 7(2): 355-361.
- Selvin, "Fluorescence resonance energy transfer", *Methods in Enzymology*, 1995, 246: 300-334.
- Shao et al., "Formyl peptide receptor ligands promote wound closure in lung epithelial cells", *American Journal of Respiratory Cell and Molecular Biology*, Mar. 2011, 44(3): 264-269.
- Sharaf et al., "Comparative effectiveness and cost-effectiveness of screening colonoscopy vs. Sigmoidoscopy and alternative strategies", *The American Journal of Gastroenterology*, 2013, 108: 120-132.
- Shen et al., "Multiple but dissectible functions of FEN-1 nucleases in nucleic acid processing, genome stability and diseases" *BioEssays*, Jul. 2005, 27(7): 717-729.
- Shendure et al., "Accurate multiplex polony sequencing of an evolved bacterial genome", *Science*, Sep. 9, 2005, 309(5741): 1728-1732.
- Shendure et al., "Next-generation DNA sequencing", *Nature Biotechnology*, Oct. 2008, 26(10): 1135-1145.
- Siegel et al., "Cancer Statistics, 2013", *CA: A Cancer Journal for Clinicians*, 2013, 63: 11-30.
- Singer-Sam et al., "A quantitative HpaII-PCR assay to measure methylation of DNA from a small number of cells", *Nucleic Acids Research*, Feb. 11, 1990, 18(3): 687.
- Singer-Sam et al., "A sensitive, quantitative assay for measurement of allele-specific transcripts differing by a single nucleotide", *PCR Methods and Applications*, Feb. 1992, 1(3): 160-163.
- Singh et al., "Risk of developing colorectal cancer following a negative colonoscopy examination evidence for a 10-year interval between colonoscopies", *Journal of the American Medical Association*, 2006, 295: 2366-2373.
- Straub et al., "Bases, a versatile, highly integrated high-throughput methylation profiling for methylation specific PCR based marker identification applied to colorectal cancer", *Clinical Cancer Research*, Oct. 2007, 13(19 Suppl):A61: 4 pages.
- Stryer, "Fluorescence energy transfer as a spectroscopic ruler", *Annual Review of Biochemistry*, 1978, 47: 819-846.
- Swift-Scanlan et al., "Two-color quantitative multiplex methylation-specific PCR", *Biotechniques*, Feb. 2006, 40(2): 210-219.
- Szabo et al., "Allele-specific expression and total expression levels of imprinted genes during early mouse development: implications for imprinting mechanisms", *Genes & Development*, Dec. 15, 1995, 9(24): 3097-3108.
- Tadokoro et al., "Classification of hepatitis B virus genotypes by the PCR-Invader method with genotype-specific probes", *Journal of Virological Methods*, Dec. 2006, 138(1-2): 30-39.
- Taylor et al., "Discovery of novel DNA methylation markers for the detection of colorectal neoplasia: selection by methylome-wide analysis", *Gastroenterology*, Abstract 109, May 1, 2014, 146(5): 5-30.
- Toth et al., "Detection of methylated SEPT9 in plasma is a reliable screening method for both left- and right-sided colon cancers", *PLoS One*, 2012, 7(9):e46000, 7 pages.
- Toyota et al., "Identification of differentially methylated sequences in colorectal cancer by methylated CpG island amplification", *Cancer Research*, May 15, 1999, 59(10): 2307-2312.
- Triglia et al., "A procedure for in vitro amplification of DNA segments that lie outside the boundaries of known sequences", *Nucleic Acids Research*, 1988, 16(16): 8186.
- Turcatti et al., "A new class of cleavable fluorescent nucleotides: synthesis and optimization as reversible terminators for DNA sequencing by synthesis", *Nucleic Acids Research*, Mar. 2008, 36(4):e25: 13 pages.
- Tureci et al., "Humoral immune responses of lung cancer patients against tumor antigen NY-ESO-1", *Cancer Letters*, May 8, 2006, 236(1): 64-71.

(56)

References Cited**OTHER PUBLICATIONS**

Turner et al., "Role of matrix metalloproteinase 9 in pituitary tumor behavior", *The Journal of Clinical Endocrinology and Metabolism*, Aug. 2000, 85(8): 2931-2935.

Umu et al., "A comprehensive profile of circulating RNAs in human serum", *RNA Biology*, Feb. 1, 2018, 15(2): 242-250.

Vancompernelle et al., "Expression and function of formyl peptide receptors on human fibroblast cells", *Journal of Immunology*, Aug. 15, 2003, 171(4): 2050-2056.

Vogelstein et al., "Cancer genome landscapes", *Science*, 2013, 339: 1546-1558.

Vogelstein et al., "Digital PCR", *Proceedings of the National Academy of Sciences of the USA*, 1999, 96: 9236-9241.

Wang et al., "Crosstalk to stromal fibroblasts induces resistance of lung cancer to epidermal growth factor receptor tyrosine kinase inhibitors", *Clinical Cancer Research*, Nov. 1, 2009, 15(21): 6630-6638.

Wang et al., "DNA methylation study of fetus genome through a genome-wide analysis", *BMC Medical Genomics*, Apr. 15, 2014, 7(18): 8 pages.

Weisenberger et al., "Comprehensive DNA methylation analysis on the Illumina® Infinium® Assay Platform", *Illumina, Application Note: Epigenetic Analysis*, 2008, 4 pages.

Williams et al., "Amplification of complex gene libraries by emulsion PCR", *Nature Methods*, Jul. 2006, 3(7): 545-550.

Winawer et al., "Screening for colorectal cancer with fecal occult blood testing and sigmoidoscopy", *Journal of the National Cancer Institute*, 1993, 85(16): 1311-1318.

Woodcock et al., "The majority of methylated deoxycytidines in human DNA are not in the CpG dinucleotide", *Biochemical and Biophysical Research Communications*, 1987, 145: 888-894.

Wrangle et al., "Functional identification of cancer-specific methylation of CDO1, HOXA9, and TAC1 for the diagnosis of lung cancer", *Clinical Cancer Research*, Apr. 1, 2014, 20(7): 1856-1864.

Wu et al., "Detection of colorectal cancer using a simplified SEPT9 gene methylation assay is a reliable method for opportunistic screening", *The Journal of Molecular Diagnostics*, Jul. 2016, 18(4): 535-545.

Xiong et al., "COBRA: a sensitive and quantitative DNA methylation assay", *Nucleic Acids Research*, Jun. 15, 1997, 25(12): 2532-2534.

Yoo et al., "Epigenetic therapy of cancer: past, present and future", *Nature Reviews Drug Discovery*, Jan. 2006, 5(1): 37-50.

Yu et al., "Significance of combined detection of LunX mRNA and tumor markers in diagnosis of lung carcinoma", *Chinese Journal of Cancer Research*, Feb. 2014, 26(1): 89-94.

Zeschnigk et al., "Imprinted segments in the human genome: different DNA methylation patterns in the prader willi/angelman syndrome region as determined by the genomic sequencing method", *Human Molecular Genetics*, Mar. 1997, 6(3): 387-95.

Zheng et al., "Detection of single nucleotide polymorphism genotyping by real-time polymerase chain reaction coupled with high specific invader assay in single tube", *Chinese Journal of Analytical Chemistry*, 2015, 43(7): 1001-1008.

Zhou et al., "Massively parallel signature sequencing", *Methods in Molecular Biology*, 2006, 331: 285-311.

Zou et al., "A sensitive method to quantify human long DNA in stool: relevance to colorectal cancer screening", *Cancer Epidemiology, Biomarkers & Prevention*, 2006, 15(6): 1115-1119.

Zou et al., "Quantification of methylated markers with a multiplex methylation-specific technology", *Clinical Chemistry*, Feb. 2012, 58(2): 375-383.

Zou et al., "Sensitive quantification of methylated markers with a novel methylation specific technology", *Clinical Chemistry, Abstract D-144*, 2010, 56(6 Suppl)A199: 3 pages.

Zou et al., "Detection of aberrant p16 methylation in the serum of colorectal cancer patients", *Clinical Cancer Research*, 2002, 8(1): 188-191.

* cited by examiner





FIG. 1

Amplification Curves

— C1: mutant 1e5	-- C2: 1m:10WT	----- C3: 1m:100WT	— C4: 1m:1000WT
----- C5: 1m:10000WT	--- C6: WT 1e5	----- C7: HGD 1e5	— C8: water
— D1: mutant 1e5	-- D2: 1m:10WT	----- D3: 1m:100WT	— D4: 1m:1000WT
----- D5: 1m:10000WT	--- D6: WT 1e5	----- D7: HGD 1e5	— D8: water

Sample name	Mutant copy number	Cp value of 1 stage approach
WT 1e5	0	44.56
WT 1e5	0	43.87
1M:10000WT*	10	43.99
1M:10000WT	10	43.04
1M:1000WT	100	43.39
1M:1000WT	100	43.66
1M:100WT	1000	39.7
1M:100WT	1000	39.62
1M:10WT	10000	33.68
1M:10WT	10000	34.28
Mutant 1e5	100000	27.72
Mutant 1e5	100000	28.04




*Note: M and WT represent mutant copy and wild-type copy, respectively

FIG. 2

Amplification Curves

— C1 mutant 1e5	— C2: 1m:10WT	----- C3: 1m:100WT	----- C4: 1m:1000WT
— C5: 1m:10000WT	— C6: WT 1e5	----- C7 HGD 1e5	— C8: water
— D1 mutant 1e5	— D2: 1m:10WT	----- D3: 1m:100WT	----- D4: 1m:1000WT
— D5: 1m:10000WT	— D6: WT 1e5	----- D7 HGD 1e5	— D8: water

Sample name	Mutant copy number	Cp value of 2 stage approach
WT 1e5	0	40.33
WT 1e5	0	40.27
1M:10000WT*	10	38.89
1M:10000WT	10	38.09
1M:1000WT	100	36.59
1M:1000WT	100	36.31
1M:100WT	1000	31.41
1M:100WT	1000	31.82
1M:10WT	10000	26.71
1M:10WT	10000	26.73
Mutant 1e5	100000	20.62
Mutant 1e5	100000	20.45



*Note: M and WT represent mutant copy and wild-type copy, respectively

FIG. 3

REAL TIME CLEAVAGE ASSAY

CROSS REFERENCE TO RELATED APPLICATIONS

The present application is a continuation of U.S. patent application Ser. No. 16/789,276, filed on Feb. 12, 2020, issued as U.S. Pat. No. 11,499,179 on Nov. 15, 2022, which is a continuation of Ser. No. 15/068,364, filed on Mar. 11, 2016, issued as U.S. Pat. No. 10,604,793 on Mar. 31, 2020 which is a continuation of Ser. No. 15/019,758, filed on Feb. 9, 2016, which is a continuation of Ser. No. 13/720,757, filed on Dec. 19, 2012, issued as U.S. Pat. No. 9,290,797 on Mar. 22, 2016, which is a continuation of Ser. No. 12/946,737, filed on Nov. 15, 2010, issued as U.S. Pat. No. 8,361,720 on Jan. 29, 2013, all of which are incorporated by reference in their entireties.

BACKGROUND

Several point mutations in the human genome have a direct association with a disease. For example, several germline KRAS mutations have been found to be associated with Noonan syndrome (Schubbert et al. Nat. Genet. 2006 38: 331-6) and cardio-facio-cutaneous syndrome (Niihori et al. Nat. Genet. 2006 38: 294-6). Likewise, somatic KRAS mutations are found at high rates in leukemias, colorectal cancer (Burmer et al. Proc. Natl. Acad. Sci. 1989 86: 2403-7), pancreatic cancer (Almoguera et al. Cell 1988 53: 549-54) and lung cancer (Tam et al. Clin. Cancer Res. 2006 12: 1647-53). Many point mutations in the human genome have no apparent causative association with a disease.

Methods for the detection of point mutations may be used, for example, to provide a diagnostic for diseases that are associated with the point mutations.

SUMMARY

A cleavage-based real-time PCR assay method is provided. In certain embodiments, the assay method includes subjecting a reaction mixture comprising a) PCR reagents for amplifying a nucleic acid target, and b) flap cleavage reagents for performing a flap cleavage assay on the amplified nucleic acid target to two sets of thermocycling conditions. The first set of thermocycling conditions includes a set of 5-15 cycles of: i. a first temperature of at least 90° C.; ii. a second temperature in the range of 60° C. to 75° C.; iii. a third temperature in the range of 65° C. to 75° C. The second and third temperatures may be the same. The second set of thermocycling conditions includes a set of 20-50 cycles of: i. a fourth temperature of at least 90° C.; ii. a fifth temperature that is at least 10° C. lower than the second temperature; iii. a sixth temperature in the range of 65° C. to 75° C. In certain cases, no additional reagents are added to the reaction between the first and second sets of cycles and, in each cycle of the second set of cycles, cleavage of a flap probe is measured.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 schematically illustrates some of the general principles of a flap assay.

FIG. 2 shows results of an assay done using single stage thermocycling. Detection and quantitation of the KRAS G35T mutation in the presence of the wild type sequence at levels, as indicated. Kinetic curves show all ratios of mutant to wild type other than 1:10 and 1:100 are indistinguishable.

FIG. 3 shows results of an assay done using two stage thermocycling. Detection and quantitation of the KRAS G35T mutation in the presence of the wild type sequence at levels, as indicated. Kinetic curves show resolution of ratios from 1:10 to 1:10,000.

DEFINITIONS

The term "sample" as used herein relates to a material or mixture of materials, typically, although not necessarily, in liquid form, containing one or more analytes of interest.

The term "nucleotide" is intended to include those moieties that contain not only the known purine and pyrimidine bases, but also other heterocyclic bases that have been modified. Such modifications include methylated purines or pyrimidines, acylated purines or pyrimidines, alkylated riboses or other heterocycles. In addition, the term "nucleotide" includes those moieties that contain hapten or fluorescent labels and may contain not only conventional ribose and deoxyribose sugars, but other sugars as well. Modified nucleosides or nucleotides also include modifications on the sugar moiety, e.g., wherein one or more of the hydroxyl groups are replaced with halogen atoms or aliphatic groups, are functionalized as ethers, amines, or the likes.

The term "nucleic acid" and "polynucleotide" are used interchangeably herein to describe a polymer of any length, e.g., greater than about 2 bases, greater than about 10 bases, greater than about 100 bases, greater than about 500 bases, greater than 1000 bases, up to about 10,000 or more bases composed of nucleotides, e.g., deoxyribonucleotides or ribonucleotides, and may be produced enzymatically or synthetically (e.g., PNA as described in U.S. Pat. No. 5,948,902 and the references cited therein) which can hybridize with naturally occurring nucleic acids in a sequence specific manner analogous to that of two naturally occurring nucleic acids, e.g., can participate in Watson-Crick base pairing interactions. Naturally-occurring nucleotides include guanine, cytosine, adenine and thymine (G, C, A and T, respectively).

The term "nucleic acid sample," as used herein denotes a sample containing nucleic acid.

The term "target polynucleotide," as used herein, refers to a polynucleotide of interest under study. In certain embodiments, a target polynucleotide contains one or more target sites that are of interest under study.

The term "oligonucleotide" as used herein denotes a single stranded multimer of nucleotides of about 2 to 200 nucleotides. Oligonucleotides may be synthetic or may be made enzymatically, and, in some embodiments, are 10 to 50 nucleotides in length. Oligonucleotides may contain ribonucleotide monomers (i.e., may be oligoribonucleotides) or deoxyribonucleotide monomers. An oligonucleotide may be 10 to 20, 11 to 30, 31 to 40, 41 to 50, 51 to 60, 61 to 70, 71 to 80, 80 to 100, 100 to 150 or 150 to 200 nucleotides in length, for example.

The term "duplex," or "duplexed," as used herein, describes two complementary polynucleotides that are base-paired, i.e., hybridized together.

The term "primer" as used herein refers to an oligonucleotide that has a nucleotide sequence that is complementary to a region of a target polynucleotide. A primer binds to the complementary region and is extended, using the target nucleic acid as the template, under primer extension conditions. A primer may be in the range of about 15 to about 50 nucleotides although primers outside of this length may be used. A primer can be extended from its 3' end by the action

of a polymerase. An oligonucleotide that cannot be extended from its 3' end by the action of a polymerase is not a primer.

The term "extending" as used herein refers to any addition of one or more nucleotides to the end of a nucleic acid, e.g., by ligation of an oligonucleotide or by using a polymerase.

The term "amplifying" as used herein refers to generating one or more copies of a target nucleic acid, using the target nucleic acid as a template.

The term "denaturing," as used herein, refers to the separation of a nucleic acid duplex into two single strands.

The terms "determining," "measuring," "evaluating," "assessing," "assaying," "detecting," and "analyzing" are used interchangeably herein to refer to any form of measurement, and include determining if an element is present or not. These terms include both quantitative and/or qualitative determinations. Assessing may be relative or absolute. "Assessing the presence of" includes determining the amount of something present, as well as determining whether it is present or absent.

The term "using" has its conventional meaning, and, as such, means employing, e.g., putting into service, a method or composition to attain an end.

As used herein, the term " T_m " refers to the melting temperature of an oligonucleotide duplex at which half of the duplexes remain hybridized and half of the duplexes dissociate into single strands. The T_m of an oligonucleotide duplex may be experimentally determined or predicted using the following formula $T_m = 81.5 + 16.6(\log_{10}[\text{Na}^+]) + 0.41(\text{fraction G+C}) - (60/N)$, where N is the chain length and $[\text{Na}^+]$ is less than 1 M. See Sambrook and Russell (2001; Molecular Cloning: A Laboratory Manual, 3rd ed., Cold Spring Harbor Press, Cold Spring Harbor N.Y., ch. 10). Other formulas for predicting T_m of oligonucleotide duplexes exist and one formula may be more or less appropriate for a given condition or set of conditions.

As used herein, the term " T_m -matched" refers to a plurality of nucleic acid duplexes having T_m that are within a defined range, e.g., within 5° C. or 10° C. of each other.

As used herein, the term "reaction mixture" refers to a mixture of reagents that are capable of reacting together to produce a product in appropriate external conditions over a period of time. A reaction mixture may contain PCR reagents and flap cleavage reagents, for example, the recipes for which are independently known in the art.

The term "mixture", as used herein, refers to a combination of elements, that are interspersed and not in any particular order. A mixture is heterogeneous and not spatially separable into its different constituents. Examples of mixtures of elements include a number of different elements that are dissolved in the same aqueous solution, or a number of different elements attached to a solid support at random or in no particular order in which the different elements are not spatially distinct. A mixture is not addressable. To illustrate by example, an array of spatially separated surface-bound polynucleotides, as is commonly known in the art, is not a mixture of surface-bound polynucleotides because the species of surface-bound polynucleotides are spatially distinct and the array is addressable.

As used herein, the term "PCR reagents" refers to all reagents that are required for performing a polymerase chain reaction (PCR) on a template. As is known in the art, PCR reagents essentially include a first primer, a second primer, a thermostable polymerase, and nucleotides. Depending on the polymerase used, ions (e.g., Mg^{2+}) may also be present. PCR reagents may optionally contain a template from which a target sequence can be amplified.

As used herein, the term "flap assay" refers to an assay in which a flap oligonucleotide is cleaved in an overlap-dependent manner by a flap endonuclease to release a flap that is then detected. The principles of flap assays are well known and described in, e.g., Lyamichev et al. (Nat. Biotechnol. 1999 17:292-296), Ryan et al (Mol. Diagn. 1999 4:135-44) and Allawi et al (J Clin Microbiol. 2006 44: 3443-3447). For the sake of clarity, certain reagents that are employed in a flap assay are described below. The principles of a flap assay are illustrated in FIG. 1. In the flap assay shown in FIG. 1, an invasive oligonucleotide 2 and flap oligonucleotide 4 are hybridized to target 6 to produce a first complex 8 that contains a nucleotide overlap at position 10. First complex 8 is a substrate for flap endonuclease. Flap endonuclease 12 cleaves flap oligonucleotide 4 to release a flap 14 that hybridizes with FRET cassette 16 that contains a quencher "Q" and a nearby quenched fluorophore "R" that is quenched by the quencher Q. Hybridization of flap 14 to FRET cassette 16 results in a second complex 18 that contains a nucleotide overlap at position 20. The second complex is also a substrate for flap endonuclease. Cleavage of FRET cassette 16 by flap endonuclease 12 results in release of the fluorophore 22, which produces a fluorescent signal. These components are described in greater detail below.

As used herein, the term "invasive oligonucleotide" refers to an oligonucleotide that is complementary to a region in a target nucleic acid. The 3' terminal nucleotide of the invasive oligonucleotide may or may not base pair a nucleotide in the target (e.g., which may be the site of a SNP or a mutation, for example).

As used herein, the term "flap oligonucleotide" refers to an oligonucleotide that contains a flap region and a region that is complementary to a region in the target nucleic acid. The target complementary regions on the invasive oligonucleotide and the flap oligonucleotide overlap by a single nucleotide. As is known, if the 3' terminal nucleotide of the invasive nucleotide and the nucleotide that overlaps that nucleotide in the flap oligonucleotide both base pair with a nucleotide in the target nucleic acid, then a particular structure is formed. This structure is a substrate for an enzyme, defined below as a flap endonuclease, that cleaves the flap from the target complementary region of the flap oligonucleotide. If the 3' terminal nucleotide of the invasive oligonucleotide does not base pair with a nucleotide in the target nucleic acid, or if the overlap nucleotide in the flap oligonucleotide does not base pair with a nucleotide in the target nucleic acid, the complex is not a substrate for the enzyme and there is little or no cleavage.

The term "flap endonuclease" or "FEN" for short, as used herein, refers to a class of nucleolytic enzymes that act as structure specific endonucleases on DNA structures with a duplex containing a single stranded 5' overhang, or flap, on one of the strands that is displaced by another strand of nucleic acid, i.e., such that there are overlapping nucleotides at the junction between the single and double-stranded DNA. FENs catalyze hydrolytic cleavage of the phosphodiester bond at the junction of single and double stranded DNA, releasing the overhang, or the flap. Flap endonucleases are reviewed by Ceska and Savers (*Trends Biochem. Sci.* 1998 23:331-336) and Liu et al (*Annu. Rev. Biochem.* 2004 73: 589-615). FENs may be individual enzymes, multi-subunit enzymes, or may exist as an activity of another enzyme or protein complex, e.g., a DNA polymerase. A flap endonuclease may be thermostable.

As used herein, the term “cleaved flap” refers to a single-stranded oligonucleotide that is a cleavage product of a flap assay.

As used herein, the term “FRET cassette” refers to a hairpin oligonucleotide that contains a fluorophore moiety and a nearby quencher moiety that quenches the fluorophore. Hybridization of a cleaved flap with a FRET cassette produces a secondary substrate for the flap endonuclease. Once this substrate is formed, the 5' fluorophore-containing base is cleaved from the cassette, thereby generating a fluorescence signal.

As used herein, the term “flap assay reagents” refers to all reagents that are required for performing a flap assay on a substrate. As is known in the art, flap assays include an invasive oligonucleotide, a flap oligonucleotide, a flap endonuclease and a FRET cassette, as described above. Flap assay reagents may optionally contain a target to which the invasive oligonucleotide and flap oligonucleotide bind.

DESCRIPTION OF EXEMPLARY EMBODIMENTS

Before the present invention is described in greater detail, it is to be understood that this invention is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range, is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges and are also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described.

All publications and patents cited in this specification are herein incorporated by reference as if each individual publication or patent were specifically and individually indicated to be incorporated by reference and are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited. The citation of any publication is for its disclosure prior to the filing date and should not be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided may be different from the actual publication dates which may need to be independently confirmed.

It must be noted that as used herein and in the appended claims, the singular forms “a”, “an”, and “the” include plural referents unless the context clearly dictates otherwise. It is further noted that the claims may be drafted to exclude any optional element. As such, this statement is intended to serve

as antecedent basis for use of such exclusive terminology as “solely,” “only” and the like in connection with the recitation of claim elements, or use of a “negative” limitation.

As will be apparent to those of skill in the art upon reading this disclosure, each of the individual embodiments described and illustrated herein has discrete components and features which may be readily separated from or combined with the features of any of the other several embodiments without departing from the scope or spirit of the present invention. Any recited method can be carried out in the order of events recited or in any other order which is logically possible.

Described herein is a cleavage-based real-time PCR assay method. In general terms, the assay method includes subjecting a reaction mixture comprising a) PCR reagents for amplifying a nucleic acid target, and b) flap cleavage reagents for performing a flap cleavage assay on the amplified nucleic acid target to two sets of thermocycling conditions. In certain cases, no additional reagents are added to the reaction between the first and second sets of cycles and, in each cycle of the second set of cycles, cleavage of a flap probe is measured. In further describing the method, the reagent mixture used in the method will be described first, followed by a description of the thermocycling conditions used in the method.

In the following description, the skilled artisan will understand that any of a number of polymerases and flap endonucleases could be used in the methods, including without limitation, those isolated from thermostable or hyperthermostable prokaryotic, eukaryotic, or archaeal organisms. The skilled artisan will also understand that the enzymes that are used in the method, e.g., polymerase and flap endonuclease, include not only naturally occurring enzymes, but also recombinant enzymes that include enzymatically active fragments, cleavage products, mutants, and variants of wild type enzymes.

Reaction Mixture

As noted above, the reaction mixture used in the method contains at least PCR reagents for amplifying a nucleic acid target and flap cleavage reagents for performing a flap cleavage assay on the amplified nucleic acid. The reaction mixture employed in the method may therefore contain a pair of primers as well a reaction buffer (which can be pH buffered and may include salt, e.g., $MgCl_2$ and other components necessary for PCR), nucleotides, e.g., dGTP, dATP, dTTP and dCTP and a thermostable DNA polymerase, as well as a flap oligonucleotide, a flap endonuclease and a FRET cassette, as defined above. Depending on how the assay is performed (i.e., depending on whether one of the PCR primers is used as an invasive oligonucleotide in the flap assay) the reaction mix may additionally contain an invasive oligonucleotide that is distinct from the PCR primers. The reaction mixture may further contain a nucleic acid target.

The exact identities and concentrations of the reagents present in the reaction mixture may be similar to or the same as those independently employed in PCR and flap cleavage assays, with the exception that the reaction mixture contains Mg^{2+} at a concentration that is higher than employed in conventional PCR reaction mixtures (which contain Mg^{2+} at a concentration of between about 1.8 mM and 3 mM). In certain embodiments, the reaction mixture described herein contains Mg^{2+} at a concentration in the range of 4 mM to 10 mM, e.g., 6 mM to 9 mM. Exemplary reaction buffers and DNA polymerases that may be employed in the subject reaction mixture include those described in various publications (e.g., Ausubel, et al., Short Protocols in Molecular

Biology, 3rd ed., Wiley & Sons 1995 and Sambrook et al., Molecular Cloning: A Laboratory Manual, Third Edition, 2001 Cold Spring Harbor, N.Y.). Reaction buffers and DNA polymerases suitable for PCR may be purchased from a variety of suppliers, e.g., Invitrogen (Carlsbad, CA), Qiagen (Valencia, CA) and Stratagene (La Jolla, CA). Exemplary polymerases include Taq, Pfu, Pwo, UItma and Vent, although many other polymerases may be employed in certain embodiments. Guidance for the reaction components suitable for use with a polymerase as well as suitable conditions for their use, is found in the literature supplied with the polymerase. Primer design is described in a variety of publications, e.g., Diffenbach and Dveksler (PCR Primer, A Laboratory Manual, Cold Spring Harbor Press 1995); R. Rapley, (The Nucleic Acid Protocols Handbook (2000), Humana Press, Totowa, N.J.); Schena and Kwok et al., Nucl. Acid Res. 1990 18:999-1005). Primer and probe design software programs are also commercially available, including without limitation, Primer Detective (ClonTech, Palo Alto, Calif.), Lasergene, (DNASTAR, Inc., Madison, Wis.); and Oligo software (National Biosciences, Inc., Plymouth, Minn.) and iOligo (Caesar Software, Portsmouth, N.H.).

Exemplary flap cleavage assay reagents are found in Lyamichiev et al. (Nat. Biotechnol. 1999 17:292-296), Ryan et al (Mol. Diagn. 1999 4:135-44) and Allawi et al (J Clin Microbiol. 2006 44: 3443-3447). Appropriate conditions for flap endonuclease reactions are either known or can be readily determined using methods known in the art (see, e.g., Kaiser et al., J. Biol. Chem. 274:21387-94, 1999). Exemplary flap endonucleases that may be used the method include, without limitation, *Thermus aquaticus* DNA polymerase I, *Thermus thermophilus* DNA polymerase I, mammalian FEN-1, *Archaeoglobus fulgidus* FEN-1, *Methanococcus jannaschii* FEN-1, *Pyrococcus furiosus* FEN-1, *Methanobacterium thermoautotrophicum* FEN-1, *Thermus thermophilus* FEN-1, CLEAVASE™ (Third Wave, Inc., Madison, Wis.), *S. cerevisiae* RTH1, *S. cerevisiae* RAD27, *Schizosaccharomyces pombe* rad2, bacteriophage T5 5'-3' exonuclease, *Pyrococcus horikoshii* FEN-1, human exonuclease 1, calf thymus 5'-3' exonuclease, including homologs thereof in eubacteria, eukaryotes, and archaea, such as members of the class II family of structure-specific enzymes, as well as enzymatically active mutants or variants thereof. Descriptions of cleaving enzymes can be found in, among other places, Lyamichiev et al., Science 260:778-83, 1993; Eis et al., Nat. Biotechnol. 19:673-76, 2001; Shen et al., Trends in Bio. Sci. 23:171-73, 1998; Kaiser et al. J. Biol. Chem. 274:21387-94, 1999; Ma et al., J. Biol. Chem. 275:24693-700, 2000; Allawi et al., J. Mol. Biol. 328:537-54, 2003; Sharma et al., J. Biol. Chem. 278:23487-96, 2003; and Feng et al., Nat. Struct. Mol. Biol. 11:450-56, 2004.

In particular embodiments, the reaction mix may contain reagents for assaying multiple (e.g., at least 2, 3, 4 or more) different targets sequences in parallel. In these cases, the reaction mix may contain multiple pairs of PCR primers, multiple different flap oligonucleotides having different flaps, and multiple different FRET cassettes for detecting the different flaps, once they are cleaved. In one embodiment, oligonucleotides in a mixture may have common flaps but different binding sequences to allow for, for example, a set of mutations to cleave a common FRET cassette and report a signal where a single fluorophore is indicative of the presence of a mutation. In this embodiment, which mutation is present in the sample may be determined after the presence of a mutation has identified. Optionally, the reaction may contain multiple invasive oligonucleotides if one of the PCR primers is not used as an invasive oligonucleotide.

Upon cleavage of the FRET cassettes, multiple distinguishable fluorescent signals may be observed. The fluorophore may be selected from, e.g., 6-carboxyfluorescein (FAM), which has excitation and emission wavelengths of 485 nm and 520 nm respectively, Redmond Red, which has excitation and emission wavelengths of 578 nm and 650 nm respectively and Yakima Yellow, which has excitation and emission wavelengths of 532 nm and 569 nm respectively, and Quasor670 which has excitation and emission wavelengths of 644 nm and 670 nm respectively, although many others could be employed. In certain cases, at least one of the PCR primer pairs, flap oligonucleotides and FRET cassettes may be for the detection of an internal control.

As would be apparent, the various oligonucleotides used in the method are designed so as to not interfere with each other. For example, in particular embodiments, the flap oligonucleotide may be capped at its 3' end, thereby preventing its extension. Likewise, in certain embodiments, the invasive oligonucleotide may also be capped at its 3' end if it is not used as one of the PCR primers. In particular embodiment, if the invasive oligonucleotide is not used as one of the PCR primers, then the invasive oligonucleotide may be present at a concentration that is in the range of 5% to 50%, e.g., 10% to 40% of the concentration of the PCR primers. Further, in certain cases, the T_m s of the flap portion and the target complementary regions of the flap oligonucleotide may independently be at least 10° C. lower (e.g., 10-20° C. lower) than the T_m s of the PCR primers, which results in a) less hybridization of the flap oligonucleotide to the target nucleic acid at higher temperatures (60° C. to 75° C.) and b) less hybridization of any cleaved flap to the FRET cassette at higher temperatures (60° C. to 75° C.). The lower fifth temperature is favorable for hybridization of the oligonucleotides used in the flap assay, and for the activity of the flap endonuclease.

In a multiplex reaction, the primers may be designed to have similar thermodynamic properties, e.g., similar T_m s, G/C content, hairpin stability, and in certain embodiments may all be of a similar length, e.g., from 18 to 30 nt, e.g., 20 to 25 nt in length. The other reagents used in the reaction mixture may also be T_m matched.

The assay mixture may be present in a vessel, including without limitation, a tube; a multi-well plate, such as a 96-well, a 384-well, a 1536-well plate; and a microfluidic device. In certain embodiments, multiple multiplex reactions are performed in the same reaction vessel. Depending on how the reaction is performed, the reaction mixture may be of a volume of 5 μ l to 200 e.g., 10 μ l to 100 μ l although volumes outside of this range are envisioned.

In certain embodiments, a subject reaction mix may further contain a nucleic acid sample. In particular embodiments, the sample may contain genomic DNA or an amplified version thereof (e.g., genomic DNA amplified using the methods of Lage et al, Genome Res. 2003 13: 294-307 or published patent application US20040241658, for example). In exemplary embodiments, the genomic sample may contain genomic DNA from a mammalian cell such as a human, mouse, rat or monkey cell. The sample may be made from cultured cells or cells of a clinical sample, e.g., a tissue biopsy, scrape or lavage or cells of a forensic sample (i.e., cells of a sample collected at a crime scene). In particular embodiments, the genomic sample may be from a formalin fixed paraffin embedded (FFPE) sample.

In particular embodiments, the nucleic acid sample may be obtained from a biological sample such as cells, tissues, bodily fluids, and stool. Bodily fluids of interest include but are not limited to, blood, serum, plasma, saliva, mucous,

phlegm, cerebral spinal fluid, pleural fluid, tears, lactal duct fluid, lymph, sputum, cerebrospinal fluid, synovial fluid, urine, amniotic fluid, and semen. In particular embodiments, a sample may be obtained from a subject, e.g., a human, and it may be processed prior to use in the subject assay. For example, the nucleic acid may be extracted from the sample prior to use, methods for which are known.

For example, DNA can be extracted from stool from any number of different methods, including those described in, e.g. Coll et al (J. of Clinical Microbiology 1989 27: 2245-2248), Sidransky et al (Science 1992 256: 102-105), Villa (Gastroenterology 1996 110: 1346-1353) and Nollau (Bio-Techniques 1996 20: 784-788), and U.S. Pat. Nos. 5,463, 782, 7,005,266, 6,303,304 and 5,741,650. Commercial DNA extraction kits for the extraction of DNA from stool include the QIAamp stool mini kit (QIAGEN, Hilden, Germany), Instagene Matrix (Bio-Rad, Hercules, Calif.), and RapidPrep Micro Genomic DNA isolation kit (Pharmacia Biotech Inc., Piscataway, N.J.), among others.

Method for Sample Analysis

In performing the subject method, the reaction mixture is generally subjected to the following thermocycling conditions: a first set of 5 to 15 (e.g., 8 to 12) cycles of: i. a first temperature of at least 90° C.; ii. a second temperature in the range of 60° C. to 75° C. (e.g., 65° C. to 75° C.); iii. a third temperature in the range of 65° C. to 75° C.; followed by: a second set of 20-50 cycles of: i. a fourth temperature of at least 90° C.; ii. a fifth temperature that is at least 10° C. lower than the second temperature (e.g., in the range of 50° C. to 55° C.; and iii. a sixth temperature in the range of 65° C. to 75° C. No additional reagents need to be added to the reaction mixture during the thermocycling, e.g., between the first and second sets of cycles. In particular embodiments, the thermostable polymerase is not inactivated between the first and second sets of conditions, thereby allowing the target to be amplified during each cycle of the second set of cycles. In particular embodiments, the second and third temperatures are the same temperature such that "two step" thermocycling conditions are performed. Each of the cycles may be independently of a duration in the range of 10 seconds to 3 minutes, although durations outside of this range are readily employed.

In each cycle of the second set of cycles (e.g., while the reaction is in the fifth temperature), a signal generated by cleavage of the flap probe may be measured to provide a real-time measurement of the amount of target nucleic acid in the sample (where the term "real-time" is intended to refer to a measurement that is taken as the reaction progresses and products accumulate). The measurement may be expressed as an absolute number of copies or a relative amount when normalized to a control nucleic acid in the sample.

Without being bound to any specific theory, it is believed that the higher reaction temperatures in the first set of cycles may allow the target nucleic acid to be efficiently amplified by the pair of PCR primers without significant interference by any of the flap assay reagents or their reaction products. The lower reaction temperature used in the second set of cycles (i.e., the fifth temperature) is not optimum for the polymerase used for PCR, but allows the flap oligonucleotide to efficiently hybridize to the target nucleic acid and is closer to the optimum temperature of the flap endonuclease. The lower reaction temperature used in the second set of cycles also facilitates subsequent hybridization of the cleaved flap to the FRET cassette. Thus, at a lower temperature, the target nucleic acid may be detected without significant interference from the PCR reagents.

In certain cases, fluorescence indicating the amount of cleaved flap can be detected by an automated fluorometer designed to perform real-time PCR having the following features: a light source for exciting the fluorophore of the FRET cassette, a system for heating and cooling reaction mixtures and a fluorometer for measuring fluorescence by the FRET cassette. This combination of features, allows real-time measurement of the cleaved flap, thereby allowing the amount of target nucleic acid in the sample to be quantified. Automated fluorometers for performing real-time PCR reactions are known in the art and can be adapted for use in this specific assay, for example, the ICYCLER™ from Bio-Rad Laboratories (Hercules, Calif.), the Mx3000P™, the MX3005P™ and the MX4000™ from Stratagene (La Jolla, Calif.), the ABI PRISM™ 7300, 7500, 7700, and 7900 Taq Man (Applied Biosystems, Foster City, Calif.), the SMARTCYCLER™, ROTORGENE 2000™ (Corbett Research, Sydney, Australia) and the GENE XPERT™ System (Cepheid, Sunnyvale, Calif.) and the LIGHTCYCLER™ (Roche Diagnostics Corp., Indianapolis, Ind.). The speed of ramping between the different reaction temperatures is not critical and, in certain embodiments, the default ramping speeds that are preset on thermocyclers may be employed.

In certain cases, the method may further involve graphing the amount of cleavage that occurs at each of the second set of cycles, thereby providing an estimate of the abundance of the nucleic acid target. The estimate may be calculated by determining the threshold cycle (i.e., the cycle at which this fluorescence increases above a predetermined threshold; the "Ct" value or "Cp" value). This estimate can be compared to a control (which control may be assayed in the same reaction mix as the genomic locus of interest) to provide a normalized estimate. The thermocycler may also contain a software application for determining the threshold cycle for each of the samples. An exemplary method for determining the threshold cycle is set forth in, e.g., Luu-The et al (Biotechniques 2005 38: 287-293).

A device for performing sample analysis is also provided. In certain embodiments, the device comprises: a) a thermocycler programmed to perform the above-described and b) a vessel comprising: PCR reagents for amplifying a nucleic acid target, and flap cleavage reagents for performing a flap cleavage assay on the nucleic acid target.

Utility

The method described finds use in a variety of applications, where such applications generally include sample analysis applications in which the presence of a target nucleic acid sequence in a given sample is detected.

In particular, the above-described methods may be employed to diagnose, to predict a response to treatment, or to investigate a cancerous condition or another mammalian disease, including but not limited to, leukemia, breast carcinoma, prostate cancer, Alzheimer's disease, Parkinson's disease, epilepsy, amyotrophic lateral sclerosis, multiple sclerosis, stroke, autism, mental retardation, and developmental disorders. Many nucleotide polymorphisms are associated with and are thought to be a factor in producing these disorders. Knowing the type and the location of the nucleotide polymorphism may greatly aid the diagnosis, prognosis, and understanding of various mammalian diseases. In addition, the assay conditions described herein can be employed in other nucleic acid detection applications including, for example, for the detection of infectious diseases, viral load monitoring, viral genotyping, environmental testing, food testing, forensics, epidemiology, and other areas where specific nucleic acid sequence detection is of use.

11

In some embodiments, a biological sample may be obtained from a patient, and the sample may be analyzed using the method. In particular embodiments, the method may be employed to identify and/or estimate the amount of mutant copies of a genomic locus that are in a biological sample that contains both wild type copies of a genomic locus and mutant copies of the genomic locus that have a point mutation relative to the wild type copies of the genomic locus. In this example, the sample may contain at least 100 times (e.g., at least 1,000 times, at least 5,000 times, at least 10,000 times, at least 50,000 times or at least 100,000 times) more wild type copies of the genomic locus than mutant copies said genomic locus.

In these embodiments, the method may be employed to detect an oncogenic mutation (which may be a somatic mutation) in, e.g., PIK3CA, NRAS, KRAS, JAK2, HRAS, FGFR3, FGFR1, EGFR, CDK4, BRAF, RET, PGDFRA, KIT or ERBB2, which mutation may be associated with breast cancer, melanoma, renal cancer, endometrial cancer, ovarian cancer, pancreatic cancer, leukemia, colorectal cancer, prostate cancer, mesothelioma, glioma, meulloblastoma, polythemia, lymphoma, sarcoma or multiple myeloma (see, e.g., Chial 2008 Proto-oncogenes to oncogenes to cancer. Nature Education 1:1).

In these embodiments, the reaction mixture may contain a first primer and a second primer wherein the first primer comprises a 3' terminal nucleotide that base pairs with the point mutation. The first primer may be employed as the invasive oligonucleotide in the second set of cycles or, in certain cases, there may be a separate invasive oligonucleotide present in the reaction mixture that also has a 3' terminal nucleotide that base pairs with the point mutation. Since the point mutation in the genomic locus may have a direct association with cancer, e.g., colorectal cancer, the subject method may be employed to diagnose patients with cancer, alone, or in combination with other clinical techniques (e.g., a physical examination such as a colonoscopy or immunohistochemical analysis) or molecular techniques. For example, results obtained from the subject assay may be combined with other information, e.g., information regarding the methylation status of other loci, information regarding in the same locus or at a different locus, cytogenetic information, information regarding rearrangements, gene expression information or information about the length of telomers, to provide an overall diagnosis of cancer or other diseases.

In one embodiment, a sample may be collected from a patient at a first location, e.g., in a clinical setting such as in a hospital or at a doctor's office, and the sample may be forwarded to a second location, e.g., a laboratory where it is processed and the above-described method is performed to generate a report. A "report" as described herein, is an electronic or tangible document which includes report elements that provide test results that may include a Ct or Cp value or the like that indicates the presence of mutant copies of the genomic locus in the sample. Once generated, the report may be forwarded to another location (which may the same location as the first location), where it may be interpreted by a health professional (e.g., a clinician, a laboratory technician, or a physician such as an oncologist, surgeon, pathologist), as part of a clinical diagnosis.

Kits

Also provided are kits for practicing the subject method, as described above. The components of the kit may be present in separate containers, or multiple components may be present in a single container.

12

In addition to above-mentioned components, the kit may further include instructions for using the components of the kit to practice the subject methods. The instructions for practicing the subject methods are generally recorded on a suitable recording medium. For example, the instructions may be printed on a substrate, such as paper or plastic, etc. As such, the instructions may be present in the kits as a package insert, in the labeling of the container of the kit or components thereof (i.e., associated with the packaging or subpackaging) etc. In other embodiments, the instructions are present as an electronic storage data file present on a suitable computer readable storage medium, e.g. CD-ROM, diskette, etc. In yet other embodiments, the actual instructions are not present in the kit, but means for obtaining the instructions from a remote source, e.g. via the internet, are provided. An example of this embodiment is a kit that includes a web address where the instructions can be viewed and/or from which the instructions can be downloaded. As with the instructions, this means for obtaining the instructions is recorded on a suitable substrate.

In addition to the instructions, the kits may also include one or more control samples, e.g., positive or negative controls analytes for use in testing the kit.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. The citation of any publication is for its disclosure prior to the filing date and should not be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it is readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

Example 1

KRAS G35T Assay

The assay described below is designed to detect nucleic acid sequences containing the KRAS G35T mutation in a background of wild type sequences. For reference, partial nucleotide sequences for the wild type and G35T mutant alleles of KRAS are shown below.

Partial sequence of amplification region for KRAS, wild type (position 35 underlined):

(SEQ ID NO: 1)
 ATGACTGAATATAAACTTGTGGTAGTTGGAGCTGTTGGCGTAGGCA
 AGAGTGCCTTGACGATACAGCTAATTCAGAATCATTTTGTGGACGA
 ATATGATCCAACAATAGAGGTAAATCTGTTTAAATATGCATATTA
 CTGG

Partial sequence of amplification region for KRAS, mutant G35T (position 35 underlined):

(SEQ ID NO: 2)
 ATGACTGAATATAAACTTGTGGTAGTTGGAGCTGTTGGCGTAGGCA
 AGAGTGCCTTGACGATACAGCTAATTCAGAATCATTTTGTGGACGA

13

-continued

ATATGATCCAACAATAGAGGTAAATCTTGTTTTAATATGCATATTA

CTGG

The ability to detect the KRAS mutation T at position 35 in a background of wild type G at position 35 was tested using two different thermocycling protocols, one of which uses single stage cycling and the other uses two stage cycling (see Table 1). In both protocols, at all dilutions, approximately 100,000 copies (i.e., 100,000 double stranded plasmids) of the wild type sequence were present. To the 100,000 copies of wild type, approximately 10,000, 1000, 100, and 10 copies of the mutant target gene were added. A sample containing 100,000 copies of the mutant sequence was used as a control.

Table 1 summarizes the cycling conditions for the cleavage-based assay for single stage thermocycling and two stage thermocycling. Fluorescent signal acquisition occurs at the 53° C. temperature, conducive to the cleavage reaction of the flap probe from the target and the cleavage of the fluorophore from the FRET cassette as mediated by the released flap.

TABLE 1

Single Stage Cycling Compared to 2-Stage (Headstart) Protocol					
Stage	Temperature	Time	Single Stage: Number of Cycles	2-Stage (Headstart): Number of Cycles	Fluorescent Signal Acquisition
Pre-incubation (enzyme activation)	95° C.	2 min.	1	1	None
Amplification (Pre-Amp, Headstart)	95° C.	20 sec.	NONE	10	None
	67° C.	30 sec.			None
	70° C.	30 sec.			None
Amplification	95° C.	20 sec.	50	45	None
	53° C.	1 min.			Single
	70° C.	30 sec.			None
Cooling (Hold)	40° C.	30 sec.	1	1	None

Primers for the PCR amplification of the KRAS G35T mutation were 5'-CTATTGT TGGATCATATTCGTC-3' (SEQ ID NO:3) as the reverse primer and 5'-ACTTGTGGTAGTTGGAGCTCT-3' (SEQ ID NO:4) as the forward primer. Note that in the forward primer, the 3'T base (underlined) corresponds to the mutant position 35. The penultimate C at position 34 is also a mismatch to both the mutant and wild type sequence, and is designed to increase the discrimination of the 3' base against the wild type target.

The homogeneous detection of the KRAS G35T mutation was accomplished by the use of an endonuclease cleavable flap probe, a cleavable FRET cassette, and a heat stable flap endonuclease. For the detection of the G35T mutation, the flap probe sequence was 5'-GACGCG-GAGTTGGCGTAGGCA-3'/3C6 (SEQ ID NO:5), where the mutant base is shown underlined and the 3'-end is blocked with a hexanediol group in order to inhibit extension. The cleaved flap portion, which subsequently binds the FRET cassette, and in turn releases the fluorophore away from its quencher, includes all of the bases from the 5'-end to the mutation-specific T_m. Primers and flap probes were supplied as non-catalog items by Integrated DNA Technologies (IDT, Coralville, Iowa).

The FRET cassette used was 5'-FAM/TCT/Quencher/AGCCGGTTTTCCGGCTGAGACTCCGCGTCCGT-3'/

14

3C6 (SEQ ID NO:6), where FAM is fluorescein, the quencher is the Eclipse® Dark Quencher, and the 3'-end is blocked with a hexanediol group in order to inhibit primer extension. The FRET cassette was supplied by Hologic (Madison, Wisconsin).

The PCR reactions were done in LightCycler® 480 Multiwell 96 Plates (Roche, Indianapolis) in 10 mM MOPS pH 7.5, with 7.5 mM MgCl₂, and 250 μM dNTPs (Promega, Madison, Wisconsin). Taq polymerase was the iTaq enzyme (BioRad, Hercules, California) and the cleavage enzyme was Cleavase 2.0 (Hologic, Madison, Wisconsin). Forward primer concentration was 50 nM, reverse primer concentration was 500 nM, flap probe was at 500 nM, and the FRET cassette was used at a final concentration of 200 nM. All amplification and detection was performed in the LightCycler 480 optical thermocycler (Roche, Indianapolis, Indiana).

Raw data and kinetic curves, as generated by the LightCycler, for the two different cycling conditions, as summarized in Table 1, are shown in FIGS. 2 and 3. The results, showing the improved linear quantitative response of the 2-stage cycling protocol are delineated in Table 2.

Table 2 shows the detection and quantitation of the KRAS G35T mutation in the presence of the wild type sequence at levels, as indicated, comparing two different cycling protocols. The point at which the fluorescence of a sample rises above the background fluorescence is called the “crossing point (Cp)” of the sample (Roche LightCycler 480 Manual, Indianapolis, IN), and in these assays is calculated as being the point at which fluorescence rose to 18% of the maximum fluorescence. Cp levels above 40 cycles show no detectable dose response.

TABLE 2

detection and quantitation of the KRAS G35T mutation					
Mutant 35T copies	Wild Type 35G copies	Ratio of Mutant: Wild Type	1-Stage Crossing Point (Cp)	2-Stage (Headstart) Crossing Point (Cp)	
0	100000	N/A	44.56	40.33	
0	100000	N/A	43.87	40.27	
10	99990	1:10000	43.99	38.89	
10	99990	1:10000	43.04	38.09	
100	99900	1:1000	43.39	36.59	
100	99900	1:1000	43.66	36.31	
1000	99000	1:100	43.66	31.41	

15

TABLE 2-continued

detection and quantitation of the KRAS G35T mutation				
Mutant 35T copies	Wild Type 35G copies	Ratio of Mutant: Wild Type	1-Stage Crossing Point (Cp)	2-Stage (Headstart) Crossing Point (Cp)
1000	99000	1:100	39.70	31.82
10000	90000	1:10	39.62	26.71
10000	90000	1:10	33.68	26.73

5

16

TABLE 2-continued

detection and quantitation of the KRAS G35T mutation				
Mutant 35T copies	Wild Type 35G copies	Ratio of Mutant: Wild Type	1-Stage Crossing Point (Cp)	2-Stage (Headstart) Crossing Point (Cp)
100000	0	N/A	27.72	20.62
100000	0	N/A	28.04	20.45

SEQUENCE LISTING

Sequence total quantity: 6		
SEQ ID NO: 1	moltype = DNA length = 142	
FEATURE	Location/Qualifiers	
misc_feature	1..142	
	note = amplification region for KRAS	
source	1..142	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 1		
atgactgaat ataaacttgt	ggtagtgtgga gctggtggcg taggcaagag tgccttgacg	60
atacagctaa ttcagaatca	ttttgtggac gaatatgac caacaataga ggtaaatctt	120
gttttaatat gcataattact	gg	142
SEQ ID NO: 2	moltype = DNA length = 142	
FEATURE	Location/Qualifiers	
misc_feature	1..142	
	note = amplification region for KRAS	
source	1..142	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 2		
atgactgaat ataaacttgt	ggtagtgtgga gctggtggcg taggcaagag tgccttgacg	60
atacagctaa ttcagaatca	ttttgtggac gaatatgac caacaataga ggtaaatctt	120
gttttaatat gcataattact	gg	142
SEQ ID NO: 3	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
misc_feature	1..22	
	note = Synthetic reverse primer	
source	1..22	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 3		
ctattgttgg atcatattcg	tc	22
SEQ ID NO: 4	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
misc_feature	1..21	
	note = Synthetic forward primer	
source	1..21	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 4		
acttgttgga gttggagctc	t	21
SEQ ID NO: 5	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
misc_feature	1..21	
	note = Synthetic probe	
misc_difference	21	
	note = 3'-C6 hexanediol group	
source	1..21	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 5		
gacgcggagt tggcgtaggc	a	21
SEQ ID NO: 6	moltype = DNA length = 35	
FEATURE	Location/Qualifiers	
misc_feature	1..35	
	note = Synthetic cleavable FRET cassette	
misc_difference	35	
	note = 3'-C6 hexanediol group	

-continued

```

misc_difference      1
                     note = 5'-FAM modified where FAM is fluorescein
misc_difference      3^4
                     note = nucleotide modified with quencher
source               1..35
                     mol_type = other DNA
                     organism = synthetic construct
SEQUENCE: 6
tctagccggt ttccggtg agactccgcg tccgt

```

35

What is claimed is:

1. A method for detecting a plurality of target nucleic acids in a sample, the method comprising:

(a) subjecting a PCR reaction mixture comprising a flap endonuclease activity to thermocycling conditions that comprise:

i) a first set of 5 to 15 cycles each comprising a denaturation step followed by one or more steps in which all temperatures are in the range of 60° C. to 75° C.; followed by:

ii) a second set of 20 to 50 cycles each comprising a denaturation step followed by one or more steps comprising a temperature that is in the range of 50° C. to 57° C.;

wherein the PCR reaction mixture comprises at least: a first set of PCR primers, wherein a first target nucleic acid of the plurality of target nucleic acids is amplified by the first set of PCR primers and not by any other primers present in the PCR reaction mixture;

a second set of PCR primers, wherein a second target nucleic acid of the plurality of target nucleic acids is amplified by the second set of PCR primers and not by any other primers present in the PCR reaction mixture; and

(b) detecting a first fluorescent signal in at least some cycles of ii), wherein the first fluorescent signal is generated by cleavage of a first probe by the flap endonuclease activity and indicates the presence of the first target nucleic acid; and

detecting a second fluorescent signal in at least some cycles of ii), wherein the second fluorescent signal is generated by cleavage of a second probe by the flap endonuclease activity and indicates the presence of the second target nucleic acid,

wherein no additional reagents are added to the PCR reaction mixture between said first and second sets of cycles.

2. The method of claim 1, wherein the cycles of the second set of cycles of ii) each comprises a temperature that is in the range of 50° C. to 55° C.

3. The method of claim 1, wherein the detecting step of (b) comprises measuring cleavage of a first fluorophore from an oligonucleotide in each of the second set of cycles of ii).

4. The method of claim 1, wherein step (a) comprises an initial pre-incubation step that activates a thermostable polymerase.

5. The method of claim 1, wherein the method further comprises graphing the amount of first and second fluorescent signals that are detected in each of the cycles of ii), thereby providing an estimate of the abundance of the first and second target nucleic acids in the reaction mix.

6. The method of claim 1, wherein the first fluorescent signal of (b) indicates a mutation and the second fluorescent signal of (b) indicates a control.

7. The method of claim 6, wherein the cycles of the second set of cycles of ii) each comprises a temperature that is in the range of 50° C. to 55° C.

8. The method of claim 6, wherein the detecting step of (b) comprises measuring cleavage of a first fluorophore from a first oligonucleotide in a plurality of the second set of cycles of ii) and measuring cleavage of a second fluorophore from a second oligonucleotide in a plurality of the second set of cycles of ii).

9. The method of claim 6, wherein step (a) comprises an initial pre-incubation step that activates a thermostable polymerase.

10. The method of claim 6, wherein the method further comprises graphing the amount of first and second fluorescent signals that are detected in a plurality of the cycles of ii), thereby providing an estimate of the abundance of the first and second target nucleic acids in the reaction mix.

11. A method for detecting a target nucleic acid comprising:

(a) subjecting a PCR reaction mixture comprising a flap endonuclease activity to thermocycling conditions that comprise:

i) a first set of 5 to 15 cycles each comprising a denaturation step followed by one or more steps in which all temperatures are in the range of 65° C. to 75° C.; followed by

ii) a second set of 20 to 50 cycles each comprising a denaturation step followed by one or more steps comprising a temperature that is in the range of 50° C. to 65° C.,

wherein the PCR reaction mixture comprises:

a first set of PCR primers, wherein a first target nucleic acid of the plurality of target nucleic acids is amplified by the first set of PCR primers and not by any other primers present in the PCR reaction mixture;

a second set of PCR primers, wherein a second target nucleic acid of the plurality of target nucleic acids is amplified by the second set of PCR primers and not by any other primers present in the PCR reaction mixture; and

(b) detecting a first fluorescent signal in at least some cycles of ii), wherein the first fluorescent signal is generated by cleavage of a first probe by the flap endonuclease activity and indicates the presence of the first target nucleic acid; and

detecting a second fluorescent signal in at least some cycles of ii), wherein the second fluorescent signal is generated by cleavage of a second probe by the flap endonuclease activity and indicates the presence of the second target nucleic acid,

wherein no additional reagents are added to the PCR reaction mixture between said first and second sets of cycles.

19

12. The method of claim 11, wherein the cycles of the second set of cycles of ii) each comprises a temperature that is in the range of 50° C. to 55° C.

13. The method of claim 11, wherein the detecting step of (b) comprises measuring cleavage of a first fluorophore from an oligonucleotide in each of the second set of cycles of ii).

14. The method of claim 11, wherein step (a) comprises an initial pre-incubation step that activates a thermostable polymerase.

15. The method of claim 11, wherein the method further comprises graphing the amount of first and second fluorescent signals that are detected in a plurality of the cycles of ii), thereby providing an estimate of the abundance of the first and second target nucleic acids in the reaction mix.

16. The method of claim 11, wherein the first fluorescent signal of (b) indicates a mutation and the second fluorescent signal of (b) indicates a control.

20

17. The method of claim 16, wherein the cycles of the second set of cycles of ii) each comprises a temperature that is in the range of 50° C. to 55° C.

18. The method of claim 16, wherein the detecting step of (b) comprises measuring cleavage of a first fluorophore from an oligonucleotide in a plurality of the second set of cycles of ii) and measuring cleavage of a second fluorophore from a second oligonucleotide in a plurality of the second set of cycles of ii).

19. The method of claim 16, wherein step (a) comprises an initial pre-incubation step that activates a thermostable polymerase.

20. The method of claim 16, wherein the method further comprises graphing the amount of first and second fluorescent signals that are detected in each of the cycles of ii), thereby providing an estimate of the abundance of the first and second target nucleic acids in the reaction mix.

* * * * *