



US012391928B2

(12) **United States Patent**
Sahenk

(10) **Patent No.:** **US 12,391,928 B2**

(45) **Date of Patent:** **Aug. 19, 2025**

(54) **RECOMBINANT ADENO-ASSOCIATED VIRUS PRODUCTS AND METHODS FOR TREATING LIMB GIRDLE MUSCULAR DYSTROPHY 2A**

(71) Applicant: **RESEARCH INSTITUTE AT NATIONWIDE CHILDREN'S HOSPITAL**, Columbus, OH (US)

(72) Inventor: **Zarife Sahenk**, Columbus, OH (US)

(73) Assignee: **Research Institute at Nationwide Children's Hospital**, Columbus, OH (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1224 days.

(21) Appl. No.: **17/255,488**

(22) PCT Filed: **Jun. 28, 2019**

(86) PCT No.: **PCT/US2019/039893**

§ 371 (c)(1),

(2) Date: **Dec. 23, 2020**

(87) PCT Pub. No.: **WO2020/006458**

PCT Pub. Date: **Jan. 2, 2020**

(65) **Prior Publication Data**

US 2021/0277362 A1 Sep. 9, 2021

Related U.S. Application Data

(60) Provisional application No. 62/865,081, filed on Jun. 21, 2019, provisional application No. 62/691,934, filed on Jun. 29, 2018.

(51) **Int. Cl.**

C12N 7/00 (2006.01)

A61K 48/00 (2006.01)

A61P 21/00 (2006.01)

C12N 9/64 (2006.01)

C12N 15/86 (2006.01)

(52) **U.S. Cl.**

CPC **C12N 7/00** (2013.01); **A61K 48/0058** (2013.01); **A61P 21/00** (2018.01); **C12N 9/6472** (2013.01); **C12N 15/86** (2013.01); **C12N 2750/14131** (2013.01); **C12N 2750/14143** (2013.01)

(58) **Field of Classification Search**

CPC ... **A61K 48/0058**; **A61P 21/00**; **C12N 9/6472**; **C12N 15/86**; **C12N 2750/14143**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,173,414 A 12/1992 Lebkowski et al.
5,449,616 A 9/1995 Campbell et al.

5,658,776 A 8/1997 Flotte et al.
5,672,694 A 9/1997 Campbell et al.
5,786,211 A 7/1998 Johnson
5,871,982 A 2/1999 Wilson et al.
6,204,059 B1 3/2001 Samulski et al.
6,258,595 B1 7/2001 Gao et al.
6,262,035 B1 7/2001 Campbell et al.
6,566,118 B1 5/2003 Atkinson et al.
6,632,800 B1 10/2003 Russell et al.
7,282,199 B2 10/2007 Gao et al.
7,759,314 B2 7/2010 Fallon et al.
7,790,449 B2 9/2010 Gao et al.
7,883,858 B2 2/2011 Hood et al.
9,061,059 B2 6/2015 Chakraborty et al.
9,434,928 B2 9/2016 Mendell et al.
10,105,453 B2 10/2018 Mendell et al.
11,358,993 B2 6/2022 Rodino-Klapac et al.
2001/0029040 A1 10/2001 Toyo-Oka
2003/0225260 A1 12/2003 Snyder
2006/0154250 A1 7/2006 Morris et al.
2007/0099251 A1 5/2007 Zhang et al.
2008/0249052 A1 10/2008 Duan et al.
2009/0054823 A1 2/2009 Bridges et al.
2009/0275107 A1 11/2009 Lock et al.
2009/0280103 A1 11/2009 Flueck
2010/0003218 A1 1/2010 Duan et al.
2010/0008979 A1 1/2010 Tomatsu et al.
2010/0026655 A1 2/2010 Harley
2010/0075866 A1 3/2010 Hood et al.
2010/0112694 A1 5/2010 Marban
2010/0120627 A1 5/2010 Belouchi et al.
2010/0247495 A1 9/2010 Ichim et al.
2010/0266551 A1 10/2010 Richard et al.
2011/0023139 A1 1/2011 Weinstein et al.
2011/0053221 A1 3/2011 Chen et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CO 20210000227 1/2021
EP 0 127 839 A2 12/1984

(Continued)

OTHER PUBLICATIONS

Wang, B., Li, J., Fu, F. et al. Construction and analysis of compact muscle-specific promoters for AAV vectors. *Gene Ther* 15, 1489-1499 (2008). (Year: 2008).*

(Continued)

Primary Examiner — Maria G Leavitt

Assistant Examiner — Michael Angelo Riga

(74) *Attorney, Agent, or Firm* — Foley & Lardner LLP

(57) **ABSTRACT**

Products and methods for treating limb girdle muscular dystrophy 2A are provided. In the methods, recombinant adeno-associated viruses deliver DNA encoding a protein with calpain 3 activity.

4 Claims, 11 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited**U.S. PATENT DOCUMENTS**

2011/0070210	A1	3/2011	Andrijauskas	
2011/0076744	A1	3/2011	Wright et al.	
2011/0082192	A1	4/2011	Milne et al.	
2011/0104120	A1	5/2011	Xiao et al.	
2011/0266551	A1	11/2011	Thompson et al.	
2011/0294193	A1	12/2011	Amalfitano et al.	
2011/0301226	A1	12/2011	Mendell et al.	
2012/0087862	A1	4/2012	Hood et al.	
2013/0171172	A1	7/2013	Richard et al.	
2014/0010861	A1	1/2014	Bancel et al.	
2014/0147432	A1	5/2014	Chakraborty et al.	
2014/0179770	A1	6/2014	Zhang et al.	
2014/0234255	A1	8/2014	Lai et al.	
2014/0249208	A1	9/2014	Chakraborty et al.	
2014/0256802	A1	9/2014	Boye et al.	
2014/0273231	A1	9/2014	Zhang et al.	
2014/0323956	A1	10/2014	Mendell et al.	
2015/0111955	A1	4/2015	High et al.	
2015/0125429	A1	5/2015	Perlingeiro et al.	
2015/0232883	A1	8/2015	Dahlman et al.	
2015/0238627	A1	8/2015	Leger et al.	
2016/0058890	A1 *	3/2016	Buj Bello	C12N 9/16 435/320.1
2018/0256752	A1	9/2018	Buj Bello et al.	
2019/0000998	A1	1/2019	Mendell et al.	
2019/0202880	A1	7/2019	Rodino-Klapac et al.	
2019/0343966	A1	11/2019	Wang et al.	
2020/0339960	A1	10/2020	Sahenk	
2021/0128749	A1	5/2021	Rodino-Klapac et al.	
2021/0393801	A1	12/2021	Rodino-Klapac et al.	
2023/0390417	A1 *	12/2023	Sahenk	C12N 15/86

FOREIGN PATENT DOCUMENTS

EP	0 155 476	A	9/1985	
EP	2 859 896	A1	4/2015	
JP	2006-121961	A	5/2006	
WO	WO-95/03392	A1	2/1995	
WO	1995/13365	A1	5/1995	
WO	1995/13392	A1	5/1995	
WO	1996/17947	A1	6/1996	
WO	1997/06243	A1	2/1997	
WO	1997/08298	A1	3/1997	
WO	1997/09441	A2	3/1997	
WO	1997/21825	A1	6/1997	
WO	1998/09657	A2	3/1998	
WO	WO-99/01176	A1	1/1999	
WO	1999/11764	A2	3/1999	
WO	WO-99/43360	A1	9/1999	
WO	2001/83692	A2	11/2001	
WO	2002/53703	A2	7/2002	
WO	WO-03/074714	A1	9/2003	
WO	WO-2004/058146	A2	7/2004	
WO	WO-2007/057781	A2	5/2007	
WO	WO-2009/019505	A2	2/2009	
WO	WO-2009/054725	A2	4/2009	
WO	WO-2013/016352	A1	1/2013	
WO	WO-2013/078316	A1	5/2013	
WO	WO-2013/123503	A1	8/2013	
WO	WO-2013/151665	A2	10/2013	
WO	WO-2013/176772	A1	11/2013	
WO	WO-2014/037526	A1	3/2014	
WO	WO-2014/039916	A1	3/2014	
WO	WO-2014/093622	A2	6/2014	
WO	WO-2014/093712	A1	6/2014	
WO	WO-2014/204725	A1	12/2014	
WO	WO-2015/018503	A1	2/2015	
WO	WO-2015/021457	A2	2/2015	
WO	WO-2015/110449	A1	7/2015	
WO	WO-2015/158749	A2	10/2015	
WO	WO-2015/197232	A1	12/2015	
WO	WO-2016/115543	A2	7/2016	
WO	WO-2017087395	A1 *	5/2017 A61K 38/39
WO	WO-2017/165859	A1	9/2017	

WO	2017/180976	A1	10/2017
WO	WO-2017/180857	A1	10/2017
WO	WO-2017/181014	A1	10/2017
WO	WO-2017/181015	A1	10/2017
WO	WO-2017/221145	A1	12/2017
WO	WO-2018/170408	A1	9/2018
WO	WO-2019/012336	A1	1/2019
WO	WO-2019/078916	A1	4/2019
WO	WO-2019/118806	A1	6/2019
WO	WO-2019/152474	A1	8/2019
WO	WO-2019/195362	A1	10/2019
WO	WO-2019/209777	A1	10/2019
WO	WO-2019/245973	A1	12/2019
WO	WO-2020/123645	A1	6/2020
WO	WO-2020/176614	A1	9/2020
WO	WO-2021/035120	A1	2/2021
WO	WO-2021/257655	A1	12/2021

OTHER PUBLICATIONS

Roudaut, Carinne, et al. "Restriction of calpain3 expression to the skeletal muscle prevents cardiac toxicity and corrects pathology in a murine model of limb-girdle muscular dystrophy." *Circulation* 128.10 (2013): 1094-1104. (Year: 2013).*

NCBI BLAST Tool: Pairwise Similarity 1, Instant App ('488) SEQ ID No. 1 [1-3977]: U.S. Pat. No. 9,981,049B2 SEQ ID No. 8 (CAPN3) (Year: 2024).*

NCBI BLAST Tool: Pairwise Similarity 2, Instant App ('488) SEQ ID No. 1 [1107-3572]: U.S. Pat. No. 9,981,049B2 SEQ ID No. 8 (CAPN3) (Year: 2024).*

Wu, Zhijian, Aravind Asokan, and R. Jude Samulski. "Adeno-associated virus serotypes: vector toolkit for human gene therapy." *Molecular therapy* 14.3 (2006): 316-327. (Year: 2006).*

Yalvac et al., Impaired regeneration in calpain-3 null muscle is associated with perturbations in mTORC1 signaling and defective mitochondrial biogenesis, *Skelet. Muscle*, 7:27, (2017).

Bartoli et al., Safety and efficacy of AAV-mediated calpain 3 gene transfer in a mouse model of limb-girdle muscular dystrophy type 2A, *Mol. Ther.*, 13(2):250-259 (2006).

Carinne et al: Restriction of Calpain3 Expression to the Skeletal Muscle Prevents Cardiac Toxicity and Corrects Pathology in a Murine Model of Limb-Girdle Muscular Dystrophy, *Musc. Skel. Gen. Cell. Ther.*, 128(10):1094-1104 (2013).

Carter, Adeno-associated virus vectors, *Current Opinions in Biotechnology*, 3(5):533-539 (1992).

Clark et al., A stable cell line carrying adenovirus-inducible rep and cap genes allows for infectivity titration of adeno-associated virus vectors, *Gene Therapy*, 3(12):1124-1132 (1996).

Clark et al., Highly purified recombinant adeno-associated virus vectors are biologically active and free of detectable helper and wild-type viruses, *Hum. Gene. Ther.*, 10(6):1031-1039 (1999).

Cserjesi et al., Myogenin induces the myocyte-specific enhancer binding factor MEF-2 independently of other muscle-specific gene products, *Mol. Cell. Biol.*, 11(10):4854-4862 (1991).

De et al., High levels of persistent expression of alpha1-antitrypsin mediated by the nonhuman primate serotype rh.10 adeno-associated virus despite preexisting immunity to common human adeno-associated viruses, *Mol. Ther.*, 13(1):67-76 (2006).

Flotte et al., Gene expression from adeno-associated virus vectors in airway epithelial cells, *Am. J. Respir. Cell Mol. Biol.*, 7:349-356 (1992).

Gao et al., Clades of adeno-associated viruses are widely disseminated in human tissues, *J. Virol.*, 78:6381-6388 (2004).

Hermionat et al., Use of adeno-associated virus as a mammalian DNA cloning vector: transduction of neomycin resistance into mammalian tissue culture cells, *Proc. Natl. Acad. Sci. U.S.A.*, 81(20):6466-6470 (1984).

International Application No. PCT/US19/39893, International Preliminary Report on Patentability, mailed Jan. 7, 2021.

International Application No. PCT/US19/39893, International Search Report and Written Opinion, mailed Sep. 25, 2019.

(56)

References Cited**OTHER PUBLICATIONS**

- Johnson et al., Muscle creatine kinase sequence elements regulating skeletal and cardiac muscle expression in transgenic mice, *Mol. Cell. Biol.*, 9(8):3393-3399 (1989).
- Klapac et al., Micro-dystrophin and follistatin co-delivery restores muscle function in aged DMD model, *Hum. Mol. Genet.*, 22(24):4929-4937 (2013).
- Kramerova et al., Null mutation of calpain 3 (p94) in mice causes abnormal sarcomere formation in vivo and in vitro, *Hum. Mol. Genet.*, 13(13):1373-1388 (2004).
- Laughlin et al., Cloning of infectious adeno-associated virus genomes in bacterial plasmids, *Gene*, 23(1):65-73 (1983).
- Lebkowski et al., Adeno-associated virus: a vector system for efficient introduction and integration of DNA into a variety of mammalian cell types, *Mol. Cell. Biol.*, 7:3988-96 (1988).
- Mader et al., A steroid-inducible promoter for the controlled overexpression of cloned genes in eukaryotic cells, *Proc. Natl. Acad. Sci. U.S.A.*, 90(12):5603-5607 (1993).
- Marsic et al., Vector design tour de force: integrating combinatorial and rational approaches to derive novel adeno-associated virus variants, *Mol. Ther.*, 22(11):1900-1909 (2014).
- McCarty, Self-complementary AAV vectors; advances and applications, *Mol. Ther.*, 16(10):1648-1656 (2008).
- McLaughlin et al., Adeno-associated virus general transduction vectors: analysis of proviral structures, *J. Virol.*, 62(6):1963-73 (1988).
- Mori et al., Two novel adeno-associated viruses from cynomolgus monkey: pseudotyping characterization of capsid protein, *Virology*, 330(2):375-383 (2004).
- Muscat et al., Multiple 5'-flanking regions of the human alpha-skeletal actin gene synergistically modulate muscle-specific expression, *Mol. Cell. Biol.*, 7(11):4089-4099 (1987).
- Muzyczka, Use of adeno-associated virus as a general transduction vector for mammalian cells, current topics in microbiology and immunology, 158:97-129 (1992).
- Paul et al., Increased viral titer through concentration of viral harvests from retroviral packaging lines, *Human Gene Therapy*, 4(5):609-615 (1993).
- Perrin et al., An experimental rabies vaccine produced with a new BHK-21 suspension cell culture process: use of serum-free medium and perfusion-reactor system, *Vaccine*, 13(13):1244-1250 (1995).
- Rabinowitz et al., Cross-packaging of a single adeno-associated virus (AAV) type 2 vector genome into multiple AAV serotypes enables transduction with broad specificity, *J. Virol.*, 76(2):791-801 (2002).
- Richard et al., Mutations in the Proteolytic Enzyme Calpain 3 Cause Limb-Girdle Muscular Dystrophy Type 2A, *Cell*, 81(1):27-40 (1995).
- Rodino-Klapac et al., A translational approach for limb vascular delivery of the micro-dystrophin gene without high volume or high pressure for treatment of Duchenne muscular dystrophy, *J. Transl. Med.*, 5:45-55 (2007).
- Sambrook et al., *Molecular cloning: A laboratory manual*, 2nd Ed., Cold spring harbor laboratory, (1989).
- Samulski et al., Cloning of adeno-associated virus into pBR322: rescue of intact virus from the recombinant plasmid in human cells, *Proc. Natl. Acad. Sci. U.S.A.*, 79(6):2077-2081 (1982).
- Samulski et al., Helper-free stocks of recombinant adeno-associated viruses: normal integration does not require viral gene expression, *J. Virol.*, 63(9):3822-3828 (1989).
- Schenpp et al., Highly purified recombinant adeno-associated virus vectors. Preparation and quantitation, *Methods Mol. Med.*, 69:427-443 (2002).
- Semenza et al., Hypoxia-inducible nuclear factors bind to an enhancer element located 3' to the human erythropoietin gene, *Proc. Natl. Acad. Sci. U.S.A.*, 88(13):5680-5684 (1991).
- Senapathy et al., Molecular cloning of adeno-associated virus variant genomes and generation of infectious virus by recombination in mammalian cells, *J. Biol. Chem.*, 259:4661-4666 (1984).
- Srivastava et al., Nucleotide sequence and organization of the adeno-associated virus 2 genome, *J. Virol.*, 45:555-564 (1983).
- Tratschin et al., A human parvovirus, adeno-associated virus, as a eucaryotic vector: transient expression and encapsidation of the procaryotic gene for chloramphenicol acetyltransferase, *Mol. Cell. Biol.*, 4(10):2072-2081 (1984).
- Tratschin et al., Adeno-associated virus vector for high-frequency integration, expression, and rescue of genes in mammalian cells, *Mol. Cell. Biol.*, 5(11):3251-3260 (1985).
- Wang et al., Construction and analysis of compact muscle-specific promoters for AAV vectors, *Gene Therapy*, 15(22):1489-1499 (2008).
- Wang et al., The potential of adeno-associated viral vectors for gene delivery to muscle tissue, *Exp. Opin. on Drug. Del.*, 11(3):345-364 (2014).
- Wein et al., Translation from a DMD exon 5 IRES results in a functional dystrophin isoform that attenuates dystrophinopathy in humans and mice, *Nature Medicine*, 20(9):992-1000 (2014).
- Weintraub et al., The myoD gene family: nodal point during specification of the muscle cell lineage, *Science*, 251:761-766 (1991).
- Roudaut et al., Restriction of Calpain3 Expression to the Skeletal Muscle Prevents Cardiac Toxicity and Corrects Pathology in a Murine Model of Limb-Girdle Muscular Dystrophy, *Circulation*, 128(10): 1094-1104, (Sep. 2013).
- Rodino-Klapac, et al., Micro-dystrophin and follistatin co-delivery restores muscle function in aged DMD model, *Hum. Mol. Genet.*, 22(24): 4929-4937, (Dec. 2013).
- Yalvac et al., Impaired regeneration in calpain-3 null muscle is associated with perturbations in mTORC1 signaling and defective mitochondrial biogenesis, *Skelet. Muscle*, 7:27, 18 pages (2017).
- Abadi et al., Supplementation with alpha-lipoic acid, CoQ10, and vitamin E augments running performance and mitochondrial function in female mice, *PLoS One*, 8(4):e60722 (2013).
- ABSS (Sequence Alignment; WO2020006458, SEQ ID #1; accessed Mar. 12, 2024) (Year: 2024).
- ABSS2 (Sequence Alignment; U.S. Appl. No. 17/255,488, SEQ ID #1; accessed Mar. 12, 2024) (Year: 2024).
- Allamand et al., Early adenovirus-mediated gene transfer effectively prevents muscular dystrophy in alpha-sarcoglycan-deficient mice, *Gene Ther.*, 7(16):1385-91 (2000).
- Anderson et al., "Nucleic acid hybridisation: A practical approach," Ch. 4, IRL Press Limited, Oxford, England (1 page) 1985.
- Anderson et al., "Quantitative Filter Hybridisation—Chapter 4", *Nucleic acid hybridisation a practical approach*, 1985, pp. 73-111.
- Angelini et al., The clinical spectrum of sarcoglycanopathies, *Neurology*, 52:176-179 (1999).
- Araishi et al., Loss of the sarcoglycan complex and sarcospan leads to muscular dystrophy in beta-sarcoglycan-deficient mice, *Hum. Mol. Genet.* 8: 1589-1598 (1999).
- Arnold et al., Electrophysiological Biomarkers in Spinal Muscular Atrophy: Preclinical Proof of Concept, *Ann. Clin. Transl. Neural.*, 1 (1):34-44 (Jan. 2014).
- Asokan et al., The AAV Vector Toolkit: Poised at the Clinical Crossroads; *Molecular Therapy*, 20(4):699-708 (2012).
- Au et al., "Gene therapy advances: a meta-analysis of AAV Usage in Clinical Settings," *Frontiers in Medicine*, Feb. 9, 2022, vol. 8 (pp. 1-14).
- Bang et al., The complete gene sequence of titin, expression of an unusual approximately 700- kDa titin isoform, and its interaction with obscurin identify a novel Z-line to I-band linking system, *Gire. Res.* 89:1065-72 (2001).
- Barresi et al., Disruption of heart sarcoglycan complex and severe cardiomyopathy caused by beta sarcoglycan mutations, *J. Med. Genet.* 37: 102-107 (2000).
- Bearzi et al., Human cardiac stem cells, *Proc. Natl. Acad. Sci. USA.* 104:14068-73 (2007).
- Beastrom et al., mdx(5cv) mice manifest more severe muscle dysfunction and diaphragm force deficits than do mdx Mice, *Am. J. Pathol.*, 179(5):2464-74 (2011).
- Behlke, Chemical modification of siRNAs for in vivo use, *Oligo-nucleotides*. 18:305-319 (2008).

(56)

References Cited**OTHER PUBLICATIONS**

- Belfort et al., Homing endonucleases: from genetic anomalies to programmable genomic clippers *Methods Mol. Biol.* 1123:1-26 (2014).
- Boch et al., Breaking the code of DNA binding specificity of TAL-type III effectors, *Science*, 326:1509-12 (2009).
- Boissel et al., "megaTALs a rare-cleaving nuclease architecture for therapeutic genome engineering," *Nucleic Acids Research*, 2014, vol. 42, No. 4 (pp. 2591-2601).
- Boissel et al., Assembly and characterization of megaTALs for hyperspecific genome engineering applications, *Methods Mol. Biol.* 1239:171-96 (2015).
- Bolduc et al., "Recessive Mutations in the Putative Calcium-Activated Chloride Channel Anoctamin 5 Cause Proximal LGMD2L and Distal MMD3 Muscular Dystrophies", *The American Journal of Human Genetics*, 86, Feb. 12, 2010, (pp. 213-221).
- Bonnemann et al., Betasarcoglycan (A3b) mutations cause autosomal recessive muscular dystrophy with loss of the sarcoglycan complex, *Nat. Genet.*, 11(3):266-273 (1995).
- Bonnemann et al., Genomic screening for beta-sarcoglycan gene mutations: missense mutations may cause severe limb-girdle muscular dystrophy type 2E (LGMD 2E), *Hum. Mol. Genet.* 5:1953-1961 (1996).
- Bouquet et al., Miyoshi-like distal myopathy with mutations in anoctamin 5 gene, *Rev. Neural. (Paris)*, 168(2):135-41 (Feb. 2012).
- Bramsen et al., Development of therapeutic-grade small interfering RNAs by chemical engineering, *Front. Genet.* 20:154 (2012).
- Ceccadi et al., Homologous recombination-deficient tumors are hyper-dependent on POLQ mediated repair, *Nature*, 518:258-262 (2015).
- Cekaite et al., Gene expression analysis in blood cells in response to unmodified and 2'-modified siRNAs reveals TLR-dependent and independent effects, *J. Mol. Biol.* 365:90-108 (2007).
- Centner et al., Identification of muscle specific ring finger proteins as potential regulators of the titin kinase domain, *J. Mol. Biol.* 306:717-26 (2001).
- Cermak et al., "Efficient design and assembly of custom TALEN and other TAL effector-based constructs for DNA targeting," *Nucleic Acids Research*, 2011, (pp. 1-11).
- Cermak et al., Efficient design and assembly of custom TALENs using the Golden Gate platform, *Methods Mol. Biol.* 1239:133-59 (2015).
- Ceyhan-Birsoy et al., Recessive truncating titin gene, TTN, mutations presenting as centronuclear myopathy, *Neurology*, 81:1205-14 (2013).
- Chandrasekharan et al., Genetic defects in muscular dystrophy, *Methods Enzymol.* 479:291-322 (2010).
- Chao et al., "Several log increase in therapeutic transgene delivery by distinct adeno-associated viral serotype vectors," *Molecular therapy: the journal of the American Society of Gene Therapy*, 2000, vol. 2, Issue 6, pp. 619-623.
- Chao et al., "Sustained and complete phenotype correction of hemophilia B mice following intramuscular injection of AAV1 serotype vectors," *Molecular therapy: the journal of the American Society of Gene Therapy*, 2001, vol. 4, Issue 3, pp. 217-222.
- Chauveau et al., A rising titan: TTN review and mutation update, *Human Mutation*, 35:1046-59 (2014).
- Chernolovskaya et al., Chemical modification of siRNA, *Curr. Opin. Mol. Ther.* 12:158-67 (2010).
- Chicoine et al., "Plasmapheresis eliminates the negative impact of AAV antibodies on microdystrophin gene expression following vascular delivery," *Molecular therapy: the journal of the American Society of Gene Therapy*, 2014, vol. 22, Issue 2, pp. 338-347.
- Chicoine et al., "Vascular delivery of rAAVrh74.MCK.GALGT2 to the gastrocnemius muscle of the rhesus macaque stimulates the expression of dystrophin and laminin $\alpha 2$ surrogates", *Mol. Ther.*, 22:713-24 (2014).
- Chiorini et al., Cloning and characterization of adeno-associated virus type 5, *J. Viral.*, 73(2):1309-19 (Feb. 1999).
- Chiorini et al., Cloning of adeno-associated virus type 4 (AAV4) and generation of recombinant AAV4 particles, *J. Viral.*, 71 (9):6823-33 (Sep. 1997).
- Cho et al., DNA repair: Familiar ends with alternative endings, *Nature*, 518:174-6 (2015).
- Chu et al., "SV40 DNA transfection of cells in suspension: analysis of the efficiency of transcription and translation of T-antigen", *Gene*, 13, (1981) 197-202.
- Clark et al., "Recombinant adeno-associated viral vectors mediate long-term transgene expression in muscle," *Human gene therapy*, 1997, vol. 8, Issue 6, pp. 659-669.
- Cordier et al., "Muscle-Specific Promoters May be Necessary for Adeno-Associated Virus-Mediated Gene Transfer in the Treatment of Muscular Dystrophies," *Human Gene Therapy*, Jan. 20, 2001, vol. 12, pp. 205-215.
- Cordier et al., "Rescue of Skeletal Muscles of gamma-Sarcoglycan-Deficient Mice with Adeno-Associated Virus-Mediated Gene Transfer," *Molecular Therapy*, Feb. 2000, vol. 1, No. 2 pp. 119-129.
- Cox et al., "Therapeutic genome editing: prospects and challenges," *Nature Medicine*, Feb. 21, 2015, vol. 2 (pp. 121-131).
- D'Amario et al., Functionally competent cardiac stem cells can be isolated from endomyocardial biopsies of patients with advanced cardiomyopathies, *Gire. Res.* 108:857-61 (2011).
- Database Genbank [online], Accession No. AJ277892.2, Nov. 14, 2006 issue.
- Daya et al., "Gene Therapy Using Adeno-Associated Virus Vectors," *Clinical Microbiology Reviews*, Oct. 2008, vol. 21, No. 4 (pp. 583-593).
- Deleavey et al., Chemical modification of siRNA, *Curr. Protoc. Nucleic Acid Chem.* Chapter 16:Unit 16.3 (2009).
- Doench et al., "Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9", *Nature Biotechnology*, Feb. 2016, vol. 34, No. 2 (pp. 184-191).
- Draviam et al., The beta-II-core of sarcoglycan is essential for deposition at the plasma membrane, *Muscle and Nerve*, 34:691-701 (2006).
- Dreier et al., "Development of Zinc Finger Domains for Recognition of the 5'-ANN-3' Family of DNA Sequences and Their Use in the Construction of Artificial Transcription Factors," *The Journal of Biological Chemistry*, Aug. 3, 2001, vol. 276, No. 31 (pp. 29466-29478).
- Dreier et al., Insights into the molecular recognition of the 5'-GNN-3' family of DNA sequences by zinc finger domains, *J. Mol. Biol.* 303:489-502 (2000).
- Dreier, B. et al., "Development of zinc finger domains for recognition of the 5'-CNN-3' family DNA sequence and their use in the construction of artificial transcription factors", *The Journal of Biological Chemistry*, vol. 280, No. 42, Oct. 21, 2005, pp. 35588-3597.
- Dressman et al., Delivery of alpha- and beta-sarcoglycan by recombinant adeno-associated virus: efficient rescue of muscle, but differential toxicity, *Hum. Gene Ther.*, 13(13):1631-1646 (2002).
- Dressman, AAV-Mediated gene transfer to models of muscular dystrophy: Insights into assembly of multi-subunit membrane proteins, University of Pittsburgh (1997).
- Durbecq et al., Disruption of the beta-sarcoglycan gene reveals pathogenetic complexity of limb-girdle muscular dystrophy type 2E, *Mol. Cell*, 5:141-151 (2000).
- Fanin et al., Gender difference in limb-girdle muscular dystrophy: a muscle fiber morphometric study in 101 patients, *Clin. Neuropathology*, 33:179-801 (2014).
- Fanin et al., LGMD2E patients risk developing dilated cardiomyopathy, *Neuromuscl. Disord.*, 13(4):303-309 (2003).
- Fonfara et al., "Phylogeny of Cas9 determines functional exchangeability of dual-RNA and Cas9 among orthologous type II CRISPR-Cas systems", *Nucleic Acids Research*, vol. 42, No. 4, Nov. 22, 2013, pp. 2377-2590 (14 pages).
- Forbes et al., "Skeletal muscles of ambulant children with Duchenne muscular dystrophy: validation of multicenter study of evaluation with MR imaging and MR spectroscopy", *Radiology*, 269:198-207 (2013).

(56)

References Cited**OTHER PUBLICATIONS**

- Fowler, et al., Improved knockdown from artificial microRNAs in an enhanced miR-155 Backbone: a designer's guide to potent multi-target RNAi, *Nucleic Acids Research*, 44(5): e48, (Nov. 2015).
- Foye, Whole Genome Sequencing Solved Our Family's Genetic Mystery: Titin, *Narrat. Inq. Bioeth* 5:206-8 (2015).
- Francois, et al., Accurate Titration of Infectious AAV Particles Requires Measurement of Biologically Active Vector Genomes and Suitable Controls. *Molecular Therapy—Methods & Clinical Development*, Sep. 21, 2018, vol. 10, pp. 223-236.
- Fucini et al., Adenosine modification may be preferred for reducing siRNA immune stimulation, *Nucleic Acid Ther.* 22:205-210 (2012).
- Gaglione et al., Recent progress in chemically modified siRNAs, *Mini. Rev. Med. Chem.* 10:578-9t (2010).
- Gao et al., A novel and efficient model of coronary artery ligation and myocardial infarction in the mouse, *Gire. Res.* 107:1445-53 (2010).
- Gao et al., A novel and efficient model of coronary artery ligation in the mouse, *Methods Mol. Bic* 1037:299-311 (2013).
- Gao et al., Adeno-associated viruses undergo substantial evolution in primates during natural infections, *Proc. Natl. Acad. Sci. U.S.A.*, 2003, vol. 100, pp. 6081-6086.
- Gautel et al., The central Z-disk region of titin is assembled from a novel repeat in variable copy numbers, *Journal of Cell Science.* 109:2747-2754 (1996).
- Gebeyehu, et al., "Novel biotinylated nucleotide—analogs for labeling and colorimetric detection of DNA," *Nucleic Acids Research*, vol. 15, No. 11, (Jun. 11, 1987), p. 4513-4534.
- GenBank Accession No. AF028704.1, Adena-associated virus 6, complete genome, Jan. 12, 1998.
- GenBank Accession No. AF028705.1, Adeno-associated virus 3B, complete genome, Jan. 12, 1998.
- GenBank Accession No. AF085716.1, Adeno-associated virus 5 DNA binding trs helicase (Rep22) and capsid protein (VP1) genes, complete eds, Feb. 9, 1999.
- Genbank Accession No. AX753246, Sequence 1 from Patent EP1310571, Jun. 23, 2003.
- GenBank Accession No. AX753249, Sequence 4 from Patent EP1310571, Jun. 23, 2003.
- GenBank Accession No. AX753250.1, Sequence 5 from Patent EP1310571, Jun. 23, 2003.
- GenBank Accession No. AY631965.1, Adena-associated virus 10 nonstructural protein and caps protein genes, complete eds, Nov. 30, 2004.
- GenBank Accession No. AY631966.1, Adena-associated virus 11 nonstructural protein and caps protein genes, complete eds, Nov. 30, 2004.
- GenBank Accession No. DO813647.1, Adena-associated virus 12 Rep78 and VP1 genes, complete eds, Feb. 20, 2008.
- GenBank Accession No. EU285562.1, Adena-associated virus 13 nonstructural protein and capsid protein genes, complete eds, Sep. 23, 2008.
- Genbank Accession No. NC_001401.0, Adeno-associated virus-2, complete genome, Aug. 13, 2018.
- GenBank Accession No. NC_001401.2, Adeno-associated virus—2, complete genome, Aug. 13, 2018.
- Genbank Accession No. NC_001729.1, Adeno-associated virus- 3, complete genome, Aug. 13, 2018.
- GenBank Accession No. NC_001829.1, Adeno-associated virus-4, complete genome, Aug. 13, 2018.
- GenBank Accession No. NC_001862, Adeno-associated virus 6, complete genome, Jan. 12, 2004, located at <https://www.ncbi.nlm.nih.gov/nuccore/NC_001862.1?report=genbank>.
- GenBank Accession No. NC_002077.1, Adeno-associated virus—1, complete genome, Aug. 13, 2018, located at <https://www.ncbi.nlm.nih.gov/nuccore/NC_002077>.
- GenBank Accession No. NC_006152.1, Adeno-associated virus 5, complete genome, Aug. 13, 2018.
- GenBank Accession No. NC_006260.1, Adeno-associated virus—7, complete genome, Aug. 13, 2018.
- GenBank Accession No. NC_006261.1, Adeno-associated virus—8, complete genome, Aug. 13, 2018.
- Genbank Accession No. NM_002324.4, *Homo sapiens* sarcoglycan beta {SGCB}, Mma, Feb. 20, 2019.
- Genbank Accession No. NP_000233.1, Beta Sarcoglycan {43kD dystrophin-associated glycoprotein} *Homo sapiens*, Mar. 19, 1999.
- GenBank Accession No. J01901, Adeno-associated virus 2, complete genome, Apr. 27, 1993.
- GenBank Accession No. U89790.1, Adeno-associated virus 4, complete genome, Aug. 21, 2017.
- GenBank Registered No. NG_011618, *Homo sapiens* titin (TTN), RefSeqGene (LRG_391) on chromosome 2, Apr. 5, 2020.
- Genbank Synthetic construct *Homo sapiens* titin:100069183, MGC:199194 anoctamin 5 (ANO5) mRNA, encodes complete protein GenBank: BC172489.1, Mar. 16, 2009.
- GenBank: Accession No. NP_000223.1: beta-sarcoglycan sequence, dated Mar. 3, 1999.
- Georganopoulou et al., "A Journey with LGMD: From Protein Abnormalities to Patient Impact", *The Protein Journal*, Kluwer Academic/Plenum Publishers, Dordrecht, NL, vol. 40, No. 4, Jun. 10, 2021, pp. 466-488.
- Gerull et al., Identification of a novel frameshift mutation in the giant muscle filament titin in a large Australian family with dilated cardiomyopathy, *J. Mal. Med. (Berl)*. 84:478-83 (2006).
- Gerull et al., Mutations of TTN, encoding the giant muscle filament titin, cause familial dilated cardiomyopathy, *Nat. Genet.* 30:201-4 (2002).
- Gibertini et al., Fibrosis and inflammation are greater in muscles of beta-sarcoglycan-null mouse than mdx mouse, *Cell Tissue Res.* 356:427-443 (2014).
- Goeddel, "Gene Expression Technology: Methods in Enzymology," Academic Press, vol. 185, Jun. 11, 1990, pp. 3-7.
- Gombash et al., Adeno-Associated Viral Vector Delivery to the Enteric Nervous System: A Review, *Postdoc J.*, 2015, vol. 3, Issue 8, pp. 1-12.
- Govoni et al., "Ongoing therapeutics trials and outcome measures for Duchenne muscular dystrophy", *Cell Mol. Life Sci.*, 70:4585-602 (2013).
- Graham et al., A new technique for the assay of infectivity of human adenovirus 5 DNA, *Virology*, 1973, vol. 52, Issue 2, pp. 456-467.
- Gramlich et al., "Antisense-mediated exon skipping: a therapeutic strategy for titin-based dilated cardiomyopathy," *EMBO Molecular Medicine*, 7(5): 562-76 (2015).
- Gramlich et al., "Stress-induced dilated cardiomyopathy in a knock-in mouse model mimicking human titin-based disease", *J. Mal. Cell Cadiol.* 47:352-8 (2009).
- Granzier et al., "Deleting titin's I-band/A-band junction reveals critical roles for titin in biomechanical sensing and cardiac function", *Proc. Natl. Acad. Sci. USA.* 111:14589-94 (2014).
- Greig et al., "Impact of intravenous infusion time on AAV8 vector pharmacokinetics, safety, and liver transduction in cynomolgus macaques," *Molecular Therapy—Methods & Clinical Develop.* 3:16079, 7 pages (2016).
- Grieger et al., "Production and characterization of adeno-associated viral vectors", *Nat. Protoc.* 1 :1412-1428 (2006).
- Griffin et al. Preclinical systemic delivery of adeno-associated [alpha]-sarcoglycan gene transfer for limb-girdle muscular dystrophy, *Human Gene Therapy*, 32(7-8): 390-404, (Apr. 2021).
- Griffin et al., Defective Membrane Fusion and Repair in Anoctamin5-Deficient Muscular Dystrophy, *Human Molecular Genetics*, vol. 25, No. 10, pp. 1900-1911 (Feb. 23, 2016).
- Griffin et al., "Dose-Escalation of Systemically Delivered Adeno-Associated Virus-Mediated alpha-Sarcoglycan in a Mouse Model With Limb-Girdle Muscular Dystrophy Type 2D," Presented at the 2019 Muscular Dystrophy Association Clinical and Scientific Conference, Apr. 13-17, 2019. (Retrieved from: investorrelations.sarepta.com/staticfiles/8b00e773-3b86-4769-83dc-4d2bf22ffb0c).
- Griffin et al., "Systemic Dose Escalation Study of Alpha-Sarcoglycan Provides Functional Improvement in SGCA (I-) Mouse Model of LGMD2D," *Molecular Therapy*, vol. 26, No. 5S1, May 2018, p. 166.

(56)

References Cited**OTHER PUBLICATIONS**

- Grose et al., "Homologous Recombination Mediates Functional Recovery of Dysferlin Deficiency following AAV5 Gene Transfer", *PLoS One*, Jun. 2012, vol. 7, Issue 6, e39233.
- Guilinger et al., "Fusion of catalytically inactive Cas9 to FokI nuclease improves the specificity of genome modification," *Nature Biotechnology*, vol. 32, No. 6, Jun. 2014 (pp. 577-582).
- Gutschner et al., "Genome engineering—Matching supply with demand," *Cell Cycle*, 15(11): 1395-96 2016.
- Hafez et al., "Homing endonucleases: DNA scissors on a mission", *Genome*, 55:553-69 (2012).
- Hagan, "When are mice considered old?" The Jackson Laboratory, <https://www.jax.org/news-and-insights/jax-blog/2017/november/when-are-mice-considered-old#> Nov. 7, 2017 (8 pages).
- Hakim et al., Monitoring murine skeletal muscle function for muscle gene therapy, *Methods Mol. Biol.*, 2011, vol. 709, pp. 75-89.
- Hakim et al., The passive mechanical properties of the extensor digitorum longus muscle are compromised in 2- to 20-mo-old mdx mice, *J. Appl. Physiol.* 110: 1656-1663 (2011).
- Handschin et al., Peroxisome proliferator-activated receptor gamma coactivator 1 coactivators, energy homeostasis, and metabolism, *Endocrine reviews*, 27:728-735 (2002).
- Herman et al., "Truncations of titin causing dilated cardiomyopathy", *N. Engl. J. Med.* 366:619-28, 2012.
- Herson et al., A phase I trial of adeno-associated virus serotype 1-gamma-sarcoglycan gene therapy for limb girdle muscular type 2C, *Brain*, 2012, vol. 135, Pt 2, pp. 483-492.
- Herzog et al., Stable gene transfer and expression of human blood coagulation factor IX after intramuscular injection of recombinant adeno-associated virus, *Proc. Natl. Acad. Sci. USA*, 1997, vol. 94 (pp. 5804-5809).
- Hicks et al., A founder mutation in Anoctamin 5 is a major cause of limb-girdle muscular dystrophy, *Brain*, 134 (Pt. 1):171-82 (Jan. 2011).
- Horii et al., Validation of microinjection methods for generating knockout mice by CRISPR/Cas-mediated genome engineering, *Sci Rep.* 4:4513 (2014).
- Inouye et al., "Codon optimization of genes for efficient protein expression in mammalian cells by selection of only preferred human codons," *Protein Expression and Purification*, 2015, vol. 109, pp. 47-54.
- International Application No. PCT/US20/47339, International Preliminary Report on Patentability, mailed Mar. 3, 2022, 8 pages.
- International Application No. PCT/US2016/061703, International Preliminary Report on Patentability, mailed May 15, 2018, 10 pages.
- International Application No. PCT/US2020/019892, International Preliminary Report on Patentability, mailed Sep. 10, 2021 (8 pages).
- International Preliminary Report on Patentability for Appl. Ser. No. PCT/US2016/062052 dated May 22, 2018, 9 pages.
- International Preliminary Report on Patentability for Appl. Ser. No. PCT/US2017/027583 dated Oct. 25, 2018, 5 pages.
- International Preliminary Report on Patentability on PCT Appl. No. PCT/US2012/066265 dated May 27, 2014 (9 pages).
- International Search Report and Written Opinion for Appl. Ser. No. PCT/US2017/027583 dated Jul. 14, 2017 (8 pages).
- International Search Report and Written Opinion for Appl. Ser. No. PCT/US2016/061703 dated Feb. 2, 2017 (13 pages).
- International Search Report and Written Opinion on PCT Appl. No. PCT/US2012/066265 dated Mar. 28, 2013 (7 pages).
- International Search Report and Written Opinion on PCT Appl. No. PCT/US2020/047339 dated Dec. 10, 2020 (12 pages).
- International Search Report for Appl. Ser. No. PCT/US2016/062052 dated Feb. 7, 2017 (5 pages).
- International Search Report issued in connection with PCT/US2020/019892 dated May 14, 2020 (4 pages).
- Itoh-Satoh et al., Titan mutations are the molecular basis for dilated cardiomyopathy, *Biochem. Biophys. Res. Commun.* 291:385-93 (2002).
- Jaber et al., Titin isoforms, extracellular matrix, and global chamber remodeling in experimental dilated cardiomyopathy: functional implications and mechanistic insight, *Circ. Heart Fail.* 1:192-9 (2008).
- Jinek et al., "A Programmable Dual-RNA-Guided DNA Endonuclease in Adaptive Bacterial Immunity," *Science*, Aug. 17, 2012, 337(6096):816-821.
- John Hopkins Medicine, "Types of Muscular Dystrophy and Neuromuscular diseases," 2023, 6 pages.
- Judge et al., "Design of noninflammatory synthetic siRNA mediating potent gene silencing in vivo", *Mol. Ther.* 13:494-505 (2006).
- Justison et al., Percutaneous assisted venous return isolated limb perfusion, *J. Extra Corpor. Technol.*, 2009, vol. 41, Issue 4, pp. 231-234.
- Kajigaya et al., Self-assembled B19 parvovirus caps ids, produced in a baculovirus system, are antigenically and immunogenically similar to native virions, *Proc. Natl. Acad. Sci. USA*, 88(11):4646-50 (Jun. 1991).
- Kariko et al., "Suppression of RNA recognition by Toll-like receptors: the impact of nucleoside modification and the evolutionary origin of RNA," *Immunity*, Aug. 2005, vol. 23 (pp. 165-175).
- Kennell, "Principles and Practices of Nucleic Acid Hybridization," *Progress in Nucleic Acid Research and Molecular Biology*, Academic Press, vol. 11, 1971, (pp. 259-301).
- Kent et al., "Mechanism of microhomology-mediated end-joining promoted by human DNA polymerase theta", *Nat. Struct. Mol. Biol.* 22:230-237 (2015).
- Kessler et al., "Gene delivery to skeletal muscle results in sustained expression and systemic delivery of a therapeutic protein," *PNAS*, 1996, vol. 93, pp. 14082-14087.
- Kirnbauer et al., Virus-like particles of bovine papillomavirus type 4 in prophylactic and therapeutic immunization, *Virology*, 219(1):37-44 (May 1996).
- Kleinstiver et al., The I-TevI nuclease and linker domains contribute to the specificity of monomeric TALENs, G3 (Bethesda). 4:1155-65 (2014).
- Kobayashi et al., Sarcolemma-localized nNOS is required to maintain activity after mild exercise, *Nature*. 456:511-5 (2008).
- Kole et al., "RNA therapeutics: beyond RNA interference and antisense oligonucleotides", *Nat Rev Drug Discov.* Jan. 20, 2012;11(2):125-40. doi: 10.1038/nrd3625.
- Kolmerer et al., "Genomic organization of M line titin and its tissue-specific expression in two distinct isoforms", *J. Mol. Biol.* 256:556-63 (1996).
- Kormann et al., "Expression of therapeutic proteins after delivery of chemically modified mRNA in mice," *Nature Biotechnology*, Feb. 2011, vol. 29, No. 2 (pp. 154-157).
- Kornberg et al., "The early history of DNA polymerase: a commentary by Arthur Kornberg", *Biochimica et Biophysica Acta*. 1000:53-56 (1989).
- Kotin et al., "Manufacturing Clinical Grade Recombinant Adeno-Associated Virus Using Invertebrate Cell Lines," *Human Gene Therapy*, 28(4):Abstract Only, (Apr. 1, 2017).
- Kotin et al., Manufacturing Clinical Grade Recombinant Adeno-Associated Virus Using Invertebrate Cell Lines, *Hum. Gene Ther.*, 28(4):350-360 (2017).
- Kramerova et al., Failure to up-regulate transcription of genes necessary for muscle adaptation underlies limb girdle muscular dystrophy 2A calpainopathy, *Hum. Mol. Genet.*, 25(11):2194-2207 (2016).
- Labeit et al., "Titins: giant proteins in charge of muscle ultrastructure and elasticity", *Science*. 270:293-6 (1995).
- Laws et al., Progression of kyphosis in mdx mice, *J. Appl. Physiol.* 97:1970-7 (2004).
- Lewinter et al., Cardiac titin and heart disease, *J. Cardiovasc. Pharmacol.* 63:207-12 (2014).
- Lewinter, "Titin isoforms in heart failure: are there benefits to supersizing?", *Circulation*. 110:109-11 2004.
- Lewis et al., "Generation of neutralizing activity against human immunodeficiency virus type 1 in serum by antibody gene transfer," *Journal of virology*, 2002, vol. 76, Issue 17, pp. 8769-8775.

(56)

References Cited

OTHER PUBLICATIONS

- Li et al., "Modularly assembled designer TAL effector nucleases for targeted gene knockout and gene replacement in eukaryotes," *Nucleic Acids Research*, 2011, vol. 39, No. 14 (pp. 6315-6325).
- Li et al., Electrical impedance myography for the in vivo and ex vivo assessment of muscular dystrophy (mdx) mouse muscle, *Muscle Nerve*, 49(6):829-35 (Jun. 2014).
- Li et al., Electrophysiologic biomarkers for assessing disease progression and the effect of riluzole in SOD1 G93A ALS mice, *PLoS One*, 8(6):e65976 (Jun. 2013).
- Li et al., Intracoronary administration of cardiac stem cells in mice: a new, improved technique for cell therapy in murine models, *Basic Res. Cardiol.* 106:849-64 (2011).
- Lin et al., Transcriptional co-activator PGC-1 alpha drives the formation of slow-twitch muscle fibres, *Nature*, 418:797-801 (2002).
- Liu et al., "Adeno-associated virus-mediated microdystrophin expression protects young mdx muscle from contraction-induced injury," *Molecular therapy: the journal of the American Society of Gene Therapy*, 2005, vol. 11, Issue 2, pp. 245-256.
- Liu et al., "Validated Zinc Finger Protein Designs for All 16 GNN DNA Triplet Targets," *The Journal of Biological Chemistry*, Feb. 8, 2002, vol. 277, No. 6 (pp. 3850-3856).
- Louis et al., "EM_EST:BE676391", Jan. 27, 2011 (Jan. 27, 2011), XP055708767, Retrieved from the Internet: URL:http://ibis.internal.epo.org/exam/dbfetch.jsp?id=EM_EST:BE676391 [retrieved on Jun. 25, 2020].
- Ma et al., Pol III Promoters to express small RNAs: Delineation of transcription initiation, *Mol. Ther. Nucleic Acids*. 3:e161 (2014).
- Mahmood et al., "Limb-girdle muscular dystrophies: Where next after six decades from the first proposal (review)," *Molecular Medicine reports*, 2014, vol. 9 (pp. 1515-1532).
- Mak et al., "The crystal structure of TAL effector PthXo1 bound to its DNA target," *Science*, Feb. 10, 2012, vol. 335, No. 6069 (pp. 716-719).
- Makarenko et al., Passive stiffness changes caused by upregulation of compliant titin isoforms in human dilated cardiomyopathy hearts, *Gire. Res.* 95:708-16 (2004).
- Martin et al., Overexpression of Galgt2 in skeletal muscle prevents injury resulting from eccentric contractions in both mdx and wild-type mice, *Am. J. Physiol. Cell Physiol.*, vol. 296, pp. 476-488, Dec. 24, 2008.
- Mashiko et al., Generation of mutant mice by pronuclear injection of circular plasmid expressing Cas9 and single guided RNA, *Sci. Rep.* 3:3355 (2013).
- Mateos-Gomez et al., Mammalian Polymerase theta promotes alternative-NHEJ and suppresses recombination, *Nature*. 518:254-257 (2015).
- Matsuda et al., Visualization of dystrophic muscle fibers in mdx mouse by vital staining with Evans blue: evidence of apoptosis in dystrophin-deficient muscle, *J. Biochem.*, 118(5):959-964 (1995).
- McCarty et al., "Adeno-associated virus terminal repeat (TR) mutant generates self-complementary vectors to overcome the rate-limiting step to transduction in vivo", *Gene Therapy*, vol. 10, May 30, 2003, pp. 2112-2118.
- McCarty et al., "Self-complementary recombinant adeno-associated virus (scAAV) vectors promote efficient transduction independently of DNA synthesis", *Gene Therapy*, May 22, 2001, vol. 8, pp. 1248-1254.
- McNally et al., "Mild and Severe Muscular Dystrophy Caused by a Single gamma-Sarcoglycan Mutation", *American Journal of Human Genetics*, Nov. 1996, vol. 59, No. 5, pp. 1040-1047.
- McNally et al., The genetic landscape of cardiomyopathy and its role in heart failure, *Cell. Metab.* 21:174-182 (2015).
- Meadows et al., Micro-RNA-29 Overexpression by adeno-associated virus suppresses fibrosis in mdx: utm+/- Mice (S61.003), *Neurology*, 82:S61.003 (Abstract) (2014).
- Meadows et al., Reducing Skeletal Muscle Fibrosis with AAV-Delivered miR-29, 2012, *Neurology*, vol. 78, Issue 1, Supplement PO4.089.
- Melacini et al., Heart involvement in muscular dystrophies due to sarcoglycan gene mutations, *Muscle Nerve*. 22:473-479 (1999).
- Mendell et al., "A phase 1/2a follistatin gene therapy trial for becker muscular dystrophy," *Molecular therapy: the journal of the American Society of Gene Therapy*, 2015, vol. 23, Issue 1, pp. 192-201.
- Mendell et al., "Gene Therapy for Muscular Dystrophy: Lessons Learned and Path Forward", *Neuroscience Letters*, vol. 527, No. 2, Oct. 2012, 21 pages.
- Mendell et al., "Limb-girdle muscular dystrophy type 2D gene therapy restores alpha-sarcoglycan and associated proteins," *Ann. Neural.*, 2009, vol. 66 Issue 3, pp. 290-297.
- Mendell et al., "Sustained alpha-sarcoglycan gene expression after gene transfer in limb-girdle muscular dystrophy, type 2D," *Annals of neurology*, 2010, vol. 68, Issue 5, pp. 629-638.
- Mendell et al., Gene Delivery for Limb-Girdle Muscular Dystrophy Type 2D by Isolated Limb Infusion, *Human Gene Therapy*, 2019, vol. 30, Issue 7, pp. 794-801.
- Mendell et al., Gene Therapy for Spinal Muscular Atrophy Type 1 Shows Potential to Improve Survival and Motor Functional Outcomes, *Mol. Ther.* 24:S190 (2016).
- Mendell et al., Single-Dose Gene-Replacement Therapy for Spinal Muscular Atrophy, *N. Engl. J. Med.*, 377:1713-1722 (2017).
- Merten, O.W., AAV vector production: state of the art developments and remaining challenges. *Cell and Gene Therapy Insights*, Dec. 1, 2016, vol. 2, No. 5, pp. 521-551.
- Mingozzi et al., "Therapeutic in vivo gene transfer for genetic disease using AAV: progress and challenges", *Nature Reviews Genetics*, May 2011, vol. 12 (pp. 341-355).
- Monjaret et al., "The Phenotype of Dysferlin-Deficient Mice is not Rescued by Adeno-Associated Virus-Mediated Transfer of Anoctamin 5," *Human Gene Therapy Clinical Development*, 24(2):65-76 (Jun. 1, 2013).
- Moore et al., Limb-girdle muscular dystrophy in the United States, *J. Neuropathol. Exp. Neural.*, 65(10):995-1003 (2006).
- Moorwood et al., Isometric and eccentric force generation assessment of skeletal muscles isolated from murine models of muscular dystrophies, *Journal of Visualized Experiments*. 71 :e50036 (2013).
- Moscou et al., "A Simple Cipher Governs DNA Recognition by TAL Effectors", *Science*, Dec. 11, 2009, vol. 326 (p. 1501).
- Murphy et al., "Long-term correction of obesity and diabetes in genetically obese mice by a single intramuscular injection of recombinant adeno-associated virus encoding mouse leptin," *Proceedings of the National Academy of Sciences of the United States of America*, 1997, vol. 94, Issue 25, pp. 13921-13926.
- Narayanawami et al., Evidence-based guideline summary: diagnosis and treatment of limb-girdle and distal dystrophies: report of the guideline development subcommittee of the American Academy of Neurology and the practice issues review panel of the American Association of Neuromuscular & Electrodiagnostic Medicine, *Neurology*, 83:1453-1463 (2014).
- NCBI Accession No. NG_051363.1, *Homo sapiens* TTN antisense RNA 1 (TTN-AS1), RefSeqGene on chromosome 2, dated Feb. 17, 2020.
- NCBI Accession No. XM_012650762.1, Predicted:Propithecus coquereli titin (TTN), mRNA, dated Jun. 1, 2015.
- NCBI Accession No. XM_024453100.1, Predicted:*Homo sapiens* titin (TTN), transcript variant X12, mRNA, dated Mar. 1, 2020.
- NCBI Reference Sequence: "anoctamin-5 isoform a [*Homo sapiens*]", GenPept, Mar. 15, 2015, NP_998764.1.
- NCBI, GenBank accession No. U34976.1 (Nov. 8, 1995), 2 pages.
- Noguchi S, "Human gamma-sarcoglycan mRNA, complete cds.", *NCBI*, (Nov. 8, 1995), Database accession No. U34976, 2 pages.
- Obermann et al., Molecular structure of the sarcomeric M band: mapping of titin and myosin binding domains in myomesin and the identification of a potential regulatory phosphorylation site in myomesin, *EMBO J.* 16:211-20 (1997).
- Pacak et al., Long-term Skeletal Muscle Protection After Gene Transfer in a Mouse Model of LGMD-2D, *Molecular Therapy*, 2007, vol. 15, Issue 10, pp. 1775-1781.
- Pavlovicova et al., Structure and composition of tubular aggregates of skeletal muscle fibres, *Gen. Physiol. Biophys.*, 22(4):425-40 (Dec. 2003).

(56)

References Cited

OTHER PUBLICATIONS

- Payne et al., Nutritional therapy improves function and complements corticosteroid intervention in mdx mice. *Muscle Nerve*. Jan. 2006; 33(1):66-77.
- Peer et al., Special delivery: targeted therapy with small RNAs. *Gene Ther.* 18:1127-33 (2011).
- Peled et al., Titin mutation in familial restrictive cardiomyopathy. *Int. J. Cardiol.* 171:24-30 (2014).
- Penttilä et al., Eight new mutations and the expanding phenotype variability in muscular dystrophy caused by ANOS. *Neurology*, 78(12):897-903 (Mar. 2012).
- Powers et al., Exercise-induced oxidative stress in humans: cause and consequences. *Free Radic. Biol. Med.*, 51 (5):942-50 (Sep. 2011).
- Pozsgai et al., "Beta-Sarcoglycan gene transfer decreases fibrosis and restores force in LGMD2E mice," *Gene Therapy*, 2016, vol. 23 (pp. 57-66).
- Pozsgai et al., "Beta-Sarcoglycan Gene Transfer Leads to Functional Improvement in a Model of LGMD2E," *Molecular Therapy* vol. 22, Supplement 1, May 2014 (p. S199).
- Pozsgai et al., "Pre-Clinical Efficacy Study of Beta-Sarcoglycan Gene Transfer," *Molecular Therapy*, May 1, 2013, vol. 21, No. 1 (p. S68).
- Pozsgai et al., "Systemic AAV-Mediated [beta]-Sarcoglycan Delivery Targeting Cardiac and Skeletal Muscle Ameliorates Histological and Functional Deficits in LGMD2E Mice," *Molecular Therapy, The Journal of the American Society of Gene Therapy*, Apr. 2017, vol. 25, No. 4 (pp. 855-869).
- Rafael-Fortney et al., Early treatment with lisinopril and spironolactone preserves cardiac and skeletal muscle in duchenne muscular dystrophy mice. *Circulation*. 124:582-8 (2011).
- Raj et al., "Self-complementary adeno-associated viral vectors for gene therapy of hemophilia B: progress and challenges" *Expert Review of Hematology*, Oct. 2011, vol. 4, No. 5 (pp. 539-549).
- Ran et al., "In vivo genome editing using *Staphylococcus aureus* Cas9," *Nature*, Apr. 9, 2015, vol. 520, (18 pages).
- Roberts et al., Integrated allelic, transcriptional, and phenomic dissection of the cardiac effects of titin truncations in health and disease. *Sci. Transl. Med.* 7:270ra6 (2015).
- Rodino-Klapac et al., "Persistent expression of FLAG-tagged micro dystrophin in nonhuman primates following intramuscular and vascular delivery," *Molecular therapy: the journal of the American Society of Gene Therapy*, 2010, vol. 18, Issue 1 (pp. 109-117).
- Rodino-Klapac et al., Demonstration of SGCA Expression and Related Outcomes in Phase I/IIa Safety Isolated Limb Perfusion Trial in LGMD2D Subjects, *Molecular Therapy*, 2018, vol. 26, Issue 5, Supplemental 1, p. 1, Abstract No. 250.
- Rodino-Klapac et al., Lack of toxicity of alpha-sarcoglycan overexpression supports clinical gene transfer trial in LGMD2D, *Neurology*, 2008, vol. 71, Issue 4, pp. 240-247.
- Rose, comprehensive Virology 3:1-61 (1974).
- Ruffing et al., "Mutations in the carboxy terminus of adeno-associated virus 2 capsid proteins affect viral infectivity: lack of an RGD integrin-binding motif," *Journal of General Virology*, 1994, vol. 75, pp. 3385-3392.
- Rutledge et al., Infectious clones and vectors derived from adeno-associated virus (AAV) serotypes other than AAV type 2, *J. Viral.*, 72(1):309-19 (Jan. 1998).
- Sahenk et al., Systemic delivery of AAVrh74.tMCK.hCAPN3 rescues the phenotype in a mouse model for LGMD2A/R1, *Mol. Ther. Methods Clin. Dev.*, 22:401-414 (2021).
- Salva et al., "Design of Tissue-specific Regulatory Cassettes for High-level rAAV-mediated Expression in Skeletal and Cardiac Muscle," *Mol. Ther.*, 2007, vol. 15, Issue 2, pp. 320-329.
- Sambrook et al., Cold spring harbor laboratory press, cold Spring Harbor, N.Y., (2001).
- Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2 edition (1989).
- Sander et al., CRISPR-Cas systems for editing, regulating and targeting genomes, *Nat. Biotechnol.* 32:347-55 (2014).
- Sandona et al., Sarcoglycanopathies: molecular pathogenesis and therapeutic prospects, *Exp Rev. Mol. Med.* 11:e28 (2009).
- Sanganalmath et al., Cell therapy for heart failure: a comprehensive overview of experimental and clinical studies, current challenges, and future directions, *Gire. Res.* 113:810-34 (2013).
- Sarepta Therapeutics: "Sarepta Therapeutics' Investigational Gene Therapy SRP-9003 for the Treatment of Limb-Girdle Muscular Dystrophy Type 2E Shows Sustained Expression and Functional Improvements 2 Years After Administration", Mar. 18, 2021, pp. 1-3, Retrieved from the Internet: URL: <https://investorrelations.sarepta.com/news--releases/news-release-details/sarepta-therapeutics-investigational-gene-therapy-srp-9003-0> [retrieved on Jun. 23, 2023].
- Schreiber et al., The transcriptional coactivator PGC-1 regulates the expression and activity of the orphan nuclear receptor estrogen-related receptor alpha (ERRalpha), *J. Biol. Chem.*, 278: 9013-9018 (2003).
- Segal et al., "Toward controlling gene expression at will: Selection and design of zinc finger domains recognizing each of the 5'-GNN-3' DNA target sequences," *Proceedings of the National Academy of Sciences, USA*, Mar. 1999, vol. 96 (pp. 2758-2763).
- Semplicini et al., Clinical and genetic spectrum in limb-girdle muscular dystrophy type 2E, *Neurology*, 84:1772-81 (2015).
- Shield et al., E-box sites and a proximal regulatory region of the muscle creatine kinase gene differentially regulate expression in diverse skeletal muscles and cardiac muscle of transgenic mice, *Mal. Cell. Biol.*, 16(9):5058-5068 (1996).
- Shih et al., Finding the Achilles' heel of Muscle Giant-TALEN-mediated Gene-editing in Zebrafish Titin, *Circulation Research*, Oct. 21, 2015, vol. 117, No.(suppl_1), pp. A344. DOI: https://doi.org/10.1161/res.117.suppl_1.344.
- Siu et al., Familial dilated cardiomyopathy locus maps to chromosome 2q31, *Circulation*. 99:1022-6 (1999).
- Smith et al., Modification and secretion of human interleukin 2 produced in insect cells by a baculovirus expression vector, *Proc. Natl. Acad. Sci. USA*, 82(24):8404-8 (1985).
- Sondergaard et al., "AAV.Dysferlin Overlap Vectors Restore Function in Dysferlinopathy Animal Models," *Annals of Clinical and Translational Neurology*, 2015, vol. 2, Issue 3, pp. 256-270.
- Sonntag et al., A viral assembly factor promotes AAV2 capsid formation in the nucleolus, *PNAS*, 2010, vol. 107, Issue 22, pp. 10220-10225.
- Sorimachi et al., Tissue-specific expression and alpha-actinin binding properties of the Z-disc titin: implications for the nature of vertebrate Z-discs, *J. Mol. Biol.* 270:688-95 (1997).
- Soutschek et al., "Therapeutic silencing of an endogenous gene by systemic administration of modified siRNAs," *Nature*, Nov. 2004, pp. 173-178, vol. 432.
- Steenfolt et al., Precision genome editing: a small revolution for glycobiology, *Glycobiology*. 24:663-80 (2014).
- Straub et al., Animal models for muscular dystrophy show different patterns of sarcolemmal disruption, *J. Cell Biol.* 139:375-385 (1997).
- Strobel, et al. "Antioxidant Supplementation Reduces Skeletal Muscle Mitochondrial Biogenesis", *Official Journal of the American College of Sports Medicine*, 2011, pp. 1017-1024.
- Sun et al., Correction of Multiple Striated Muscles in Murine Pompe Disease Through Adeno-Associated Virus-mediated Gene Therapy, *Mal. Ther.*, 16(8):1366-71 (2008).
- Sveen et al., Cardiac involvement in patients with limb-girdle muscular dystrophy type 2 and Becker muscular dystrophy, *Arch. Neurol.*, 65(9):1196-1201 (2008).
- Thiruvengadam et al., "Anoctamin 5 Knockout Mouse Model Recapitulates LGMD2L Muscle Pathology and Offers Insight Into in vivo Functional Deficits," *Journal of Neuromuscular Diseases*, 2021, vol. 8 (S243-S255).
- Torella, et al., "Cardiovascular development: towards biomedical applicability; Resident cardiac stem cells", *CMLS Cellular and Molecular Life Sciences* 64(6): 661-673 (2007).
- Tsai et al., "GUIDE-seq enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases," *Nature Biotechnology*, Feb. 2015, vol. 33, No. 2 (pp. 187-197).
- Tsai et al., Dimeric CRISPR RNA-guided FokI nucleases for highly specific genome editing, *Nat. Biotechnol.* 32:569-76 (2014).

(56)

References Cited

OTHER PUBLICATIONS

- Van Akkooi et al., Isolated limb perfusion for an irresectable melanoma recurrence in a Jehovah's witness, *Eur. J. Cardiothorac. Surg.*, 2006, vol. 30, Issue 2, pp. 408-410.
- Voikar et al., Long-term individual housing in C57BU6J and DBA/2 mice: assessment of behavioral consequences, *Genes Brain Behav.*, 4(4):240-52 (2005).
- Volkov et al., Selective protection of nuclease-sensitive sites in siRNA prolongs silencing effect, *Oligonucleotides*, 19:191-202 (2009).
- Wang et al., Loss of miR-29 in myoblasts contributes to dystrophic muscle pathogenesis, *Mol. Ther.*, 20(6):1222-33 (2012).
- Wang et al., Rapid and efficient assembly of transcription activator-like effector genes by USER cloning, *J. Genet. Genomics*, 41:339-47 (2014).
- Wang et al., Recombinant AAV serotype 1 transduction efficiency and tropism in the murine brain, *Gene Ther.*, 2003, vol. 10, Issue 17, pp. 1528-1534.
- Watson et al., "Recombinant DNA," *Scientific American*, Second Edition, 2001 (pp. 153-154).
- Weber et al., "A Modular Cloning System for Standardized Assembly of Multigene Constructs," Feb. 2011, vol. 6, No. 2, e16765 (11 pages).
- Whitehead et al., Silencing or stimulation? siRNA delivery and the immune system, *Annual Review of Chemical and Biomolecular Engineering*, 2:77-96 (2011).
- Wikipedia, "Adeno-associated virus," downloaded Dec. 29, 2017 (pp. 1-18).
- Wikipedia, "Limb-girdle muscular dystrophy," 11 pages, Retrieved Oct. 26, 2023, from https://en.wikipedia.org/wiki/Limb-girdle_muscular_dystrophy (11 pages).
- Winkler, Oligonucleotide conjugates for therapeutic applications, *Ther. Deliv.*, 4:791-809 (2013).
- Witting et al., "Anoctamin 5 muscular dystrophy in Denmark: prevalence, genotypes, phenotypes, cardiac findings, and muscle protein expression", *Case Reports*, May 14, 2013, PMID: 23670307 DOI: 10.1007/s00415-013-6934-y.
- Wolfs et al., MegaTevs: single-chain dual nucleases for efficient gene disruption, *Nucleic Acids Res.* 42:8816-29 (2014).
- Wong-Kisiel et al., Two siblings with limb-girdle muscular dystrophy type 2E responsive to deflazacort, *Neuromusc. Disord.* 20:122-124 (2010).
- Wu et al., Mutational analysis of the adeno-associated virus type 2 (AAV2) capsid gene and construction of AAV2 vectors with altered tropism, *J. Viral.*, 74(18):8635-47 (Sep. 2000).
- Xiao et al., "Production of High-Titer Recombinant Adeno-Associated Virus Vectors in the Absence of Helper Adenovirus," *Journal of Virology*, Mar. 1998, vol. 72 No. 3 (pp. 2224-2232).
- Xiao et al., "Efficient long-term gene transfer into muscle tissue of immunocompetent mice by adeno-associated virus vector," *Journal of virology*, 1996, vol. 70, Issue 11, pp. 8098-8108.
- Xu et al., "An Isolated Limb Infusion Method Allows for Broad Distribution of rAAVrh74.MCK.GALGT2 to Leg Skeletal Muscles in the Rhesus Macaque," *Molecular Therapy—Methods & Clinical Develop.* 10:89-104 (Sep. 2018).
- Xu et al., "Genetic disruption of Anos1 in mice does not recapitulate human ANOS1-deficient muscular dystrophy," *Skeletal Muscle*, 2015, vol. 5, No. 43 (pp. 1-14).
- Xu et al., Postnatal overexpression of the CT GalNAc transferase inhibits muscular dystrophy in mdx mice without altering muscle growth or neuromuscular development: evidence for a utrophin-independent mechanism, *Neuromuscul. Disord.*, 2007, vol. 17, Issue 3, pp. 209-220.
- Yan et al., Inverted terminal repeat sequences are important for intermolecular recombination and circularization of adeno-associated virus genomes, *J. Viral.*, 79(1):364-79 (Jan. 2005).
- Yuasa et al., "Gene therapy of muscular dystrophy: Systemic gene delivery to skeletal muscles" Jan. 2007, *Drug Delivery System* 22(2):140-147, doi.org/10.2745/dd.22.140 (English Abstract).
- Zanotti et al., Opposing roles of miR-21 and miR-29 in the progression of fibrosis in Duchenne muscular dystrophy, *Biochem. Biophys. Acta.*, 1852:1451-4 (2015).
- Zetsche et al., "Cpf1 Is a Single RNA-Guided Endonuclease of a Class 2 CRISPR-Cas System," *Cell*, Oct. 22, 2015, vol. 163, No. 3 (pp. 759-771).
- Zhang et al., Dual AAV therapy ameliorates exercise-induced muscle injury and functional ischemia in murine models of Duchenne muscular dystrophy, *Hum. Mol. Genet.* 22:3720-9 (2013).
- Zhao et al., BPV1 E2 protein enhances packaging of full-length plasmid DNA in BPV1 pseudovirions, *Virology*, 272(2):382-93 (Jul. 2000).
- Zhou et al., Pressure Overload by Transverse Aortic Constriction Induces Maladaptive Hypertrophy in a Titin-Truncated Mouse Model, *Biomed. Res. Int.* 2015:163564 (2015).
- Zou et al., "An internal promoter underlies the difference in disease severity between N- and C-terminal truncation mutations of Titin in zebrafish", *eLife*, Oct. 16, 2015, vol. 4, pp. e09406. DOI:<https://doi.org/10.7554/eLife.09406>.
- Dorange et al., "Analytical approaches to characterize AAV vector production & purification: Advances and challenges," *Cell & Gene Therapy Insights*, 4(2):119-129 (2018).
- Hou et al., "Serious Overestimation in Quantitative PCR by Circular (Supercoiled) Plasmid Standard: Microalgal pcna as the Model Gene," *PLoS One* 5(3):e9545, 8 pages (Mar. 5, 2010) doi:10.1371/journal.pone.0009545.
- Martinez-Fernandez de la Camara et al., "The accurate quantification of AAV genomic titre depends on the conformation of the plasmid reference," *ARVO Annual Meeting Abstract*, 3 pages, Jul. 2018.
- Thomas et al., "B4GALNT2 (GALGT2) Gene Therapy Reduces Skeletal Muscle Pathology in the FKRP P448L Mouse Model of Limb Girdle Muscular Dystrophy 21", *Am. J. Pathol.*, 186(9):2429-2448 (2016).
- Werling et al., "Systematic comparison and validation of quantitative real-time PCR methods for the quantitation of adeno-associated viral products," *Human Gene Therapy Methods*, 26.3:82-92 (Jun. 2015).
- Pozsgai et al., "506. [beta]—Sarcoglycan Gene Transfer Prevents Muscle Fibrosis and Inflammation in an Aged LGMD2E Mouse Model," *Molecular Therapy*, vol. 23 Supplement 1, May 2015, 2 pages.
- Chu et al., "The limb-girdle muscular dystrophies: is treatment on the horizon?" *Neurotherapeutics*, 15(4):849-862 (Oct. 2018).
- Monies et al., "A first-line diagnostic assay for limb-girdle muscular dystrophy and other myopathies", *Human Genomics*, 10(1):32, pp. 1-7 (Sep. 27, 2016).
- Theadom et al., "Prevalence of muscular dystrophies: a systematic literature review," *Neuroepidemiology* 43(3-4):259-68 (2014).
- Wagner et al., "A novel method for the quantification of adeno-associated virus vectors for RNA interference applications using quantitative polymerase chain reaction and purified genomic adeno-associated virus DNA as a standard," *Human Gene Therapy Methods*, 24(6):355-63 (Dec. 2013).
- Walter et al., "Recent developments in Duchenne muscular dystrophy: facts and numbers," *Journal of Cachexia, Sarcopenia and Muscle*, 8(5):681-685 (Oct. 2017).
- Hartigan-O'Connor et al., "Developments in gene therapy for muscular dystrophy," *Microscopy Research and Technique* 48:223-238 (2000).
- Pozsgai, E.R., Adeno-Associated Virus Mediated β -Sarcoglycan Gene Replacement Therapy for the Treatment of Limb Girdle Muscular Dystrophy Type 2E [Doctoral dissertation, Ohio State University]. OhioLink Electronic Theses and Dissertations Center. (2016) http://rave.ohiolink.edu/etdc/view?acc_num=osu147697211337827.
- Agbandje-McKenna et al., "AAV Capsid Structure and Cell Interactions", *Adeno-Associated Virus: Methods and Protocols*, in *Methods in Molecular Biology*, Ch. 3, 807:47-92 (2011).

* cited by examiner

Figure 1




Figure 2

Created with SnapGene®





Figure 3

**B**

CALPAIN 3 mRNA Expression




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8A




Figure 8B




Figure 9





Figure 10

Assessment of Cardiac Calpain 3 protein expression
Following systemic AAVrh.74 tMCK.CAPN3 Gene Delivery



Full-length Calpain 3 protein is below the limit of detection.
6 x 10¹² vg AAVrh74.tMCK.CAPN3 systemically dosed CPN3KO mice show no detectable Calpain 3 protein expression in cardiac muscle.
Positive control (hCPN3) human quadriceps muscle lysate.

1

RECOMBINANT ADENO-ASSOCIATED VIRUS PRODUCTS AND METHODS FOR TREATING LIMB GIRDLE MUSCULAR DYSTROPHY 2A

This application is a U.S. 371 National Stage Application of PCT International Application No. PCT/US19/39893, filed Jun. 28, 2019, which claims priority to U.S. Provisional Patent Application No. 62/691,934, filed Jun. 29, 2018 and U.S. Provisional Patent Application No. 62/865,081, filed Jun. 21, 2019, the contents of each of which are incorporated herein by reference in their entirety.

Provided herein are products and methods for treating limb girdle muscular dystrophy 2A. In the methods, recombinant adeno-associated viruses deliver DNA encoding a protein with calpain3 (CAPN3) activity.

INCORPORATION BY REFERENCE OF THE SEQUENCE LISTING

This application contains, as a separate part of disclosure, a Sequence Listing in computer-readable form (filename: 52684P2_SeqListing.txt; 23,755 bytes—ASCII text file created Jun. 26, 2019) which is incorporated by reference herein in its entirety.

BACKGROUND

Muscular dystrophies (MDs) are a group of genetic diseases. The group is characterized by progressive weakness and degeneration of the skeletal muscles that control movement. Some forms of MD develop in infancy or childhood, while others may not appear until middle age or later. The disorders differ in terms of the distribution and extent of muscle weakness (some forms of MD also affect cardiac muscle), the age of onset, the rate of progression, and the pattern of inheritance.

One group of MDs is the limb girdle group (LGMD) of MDs. LGMDs are rare conditions and they present differently in different people with respect to age of onset, areas of muscle weakness, heart and respiratory involvement, rate of progression and severity. LGMDs can begin in childhood, adolescence, young adulthood or even later. Both genders are affected equally. LGMDs cause weakness in the shoulder and pelvic girdle, with nearby muscles in the upper legs and arms sometimes also weakening with time. Weakness of the legs often appears before that of the arms. Facial muscles are usually unaffected. As the condition progresses, people can have problems with walking and may need to use a wheelchair over time. The involvement of shoulder and arm muscles can lead to difficulty in raising arms over head and in lifting objects. In some types of LGMD, the heart and breathing muscles may be involved.

There are at least nineteen forms of LGMD, and the forms are classified by their associated genetic defects.

Type	Pattern of Inheritance	Gene or Chromosome
LGMD1A	Autosomal dominant	Myotilin gene
LGMD1B	Autosomal dominant	Lamin A/C gene
LGMD1C	Autosomal dominant	Caveolin gene
LGMD1D	Autosomal dominant	Chromosome 7
LGMD1E	Autosomal dominant	Desmin gene
LGMD1F	Autosomal dominant	Chromosome 7
LGMD1G	Autosomal dominant	Chromosome 4
LGMD2A	Autosomal recessive	Calpain-3 gene
LGMD2B	Autosomal recessive	Dysferlin gene

2

-continued

Type	Pattern of Inheritance	Gene or Chromosome
5 LGMD2C	Autosomal recessive	Gamma-sarcoglycan gene
LGMD2D	Autosomal recessive	Alpha-sarcoglycan gene
LGMD2E	Autosomal recessive	Beta-sarcoglycan gene
LGMD2F	Autosomal recessive	Delta-sarcoglycan gene
LGMD2G	Autosomal recessive	Telethonin gene
LGMD2H	Autosomal recessive	TRIM32
LGMD2I	Autosomal recessive	FKRP gene
10 LGMD2J	Autosomal recessive	Titin gene
LGMD2K	Autosomal recessive	POMT1 gene
LGMD2L	Autosomal recessive	Fukutin gene

Specialized tests for LGMD are now available through a national scheme for diagnosis, the National Commissioning Group (NCG).

Mutations in calpain3 gene (CAPN3) lead to one of the most common limb-girdle muscular dystrophies worldwide, LGMD2A. At present, there is no treatment for this inherited disease. Previous studies have demonstrated the potential for CAPN3 gene transfer to correct the pathological signs in CAPN3-deficient mice. However expression of CAPN3 driven by desmin promoter resulted in cardiotoxicity [Bartoli et al., *Mol. Ther.*, 13: 250-259 (2006)]. In follow up studies, skeletal muscle expression of the gene was studied [Roudaut et al., *Circulation*, 128: 1094-1104 (2013)].

Adeno-associated virus (AAV) is a replication-deficient parvovirus, the single-stranded DNA genome of which is about 4.7 kb in length including two 145 nucleotide inverted terminal repeat (ITRs). There are multiple serotypes of AAV. The nucleotide sequences of the genomes of AAV serotypes are known. For example, the complete genome of AAV-1 is provided in GenBank Accession No. NC_002077; the complete genome of AAV-2 is provided in GenBank Accession No. NC_001401 and Srivastava et al., *J. Virol.*, 45: 555-564 (1983); the complete genome of AAV-3 is provided in GenBank Accession No. NC_1829; the complete genome of AAV-4 is provided in GenBank Accession No. NC_001829; the AAV-5 genome is provided in GenBank Accession No. AF085716; the complete genome of AAV-6 is provided in GenBank Accession No. NC_001862; at least portions of AAV-7 and AAV-8 genomes are provided in GenBank Accession Nos. AX753246 and AX753249, respectively; the AAV-9 genome is provided in Gao et al., *J. Virol.*, 78: 6381-6388 (2004); the AAV-10 genome is provided in *Mol. Ther.*, 13(1): 67-76 (2006); and the AAV-11 genome is provided in *Virology*, 330(2): 375-383 (2004). The sequence of the AAV rh.74 genome is provided in see U.S. Pat. No. 9,434,928, incorporated herein by reference. Cis-acting sequences directing viral DNA replication (rep), encapsidation/packaging and host cell chromosome integration are contained within the AAV ITRs. Three AAV promoters (named p5, p19, and p40 for their relative map locations) drive the expression of the two AAV internal open reading frames encoding rep and cap genes. The two rep promoters (p5 and p19), coupled with the differential splicing of the single AAV intron (at nucleotides 2107 and 2227), result in the production of four rep proteins (rep 78, rep 68, rep 52, and rep 40) from the rep gene. Rep proteins possess multiple enzymatic properties that are ultimately responsible for replicating the viral genome. The cap gene is expressed from the p40 promoter and it encodes the three capsid proteins VP1, VP2, and VP3. Alternative splicing and non-consensus translational start sites are responsible for the production of the three related capsid proteins. A single consensus polyadenylation site is located at map position 95 of the AAV

genome. The life cycle and genetics of AAV are reviewed in Muzyczka, *Current Topics in Microbiology and Immunology*, 158: 97-129 (1992).

AAV possesses unique features that make it attractive as a vector for delivering foreign DNA to cells, for example, in gene therapy. AAV infection of cells in culture is noncytopathic, and natural infection of humans and other animals is silent and asymptomatic. Moreover, AAV infects many mammalian cells allowing the possibility of targeting many different tissues in vivo. Moreover, AAV transduces slowly dividing and non-dividing cells, and can persist essentially for the lifetime of those cells as a transcriptionally active nuclear episome (extrachromosomal element). The AAV proviral genome is inserted as cloned DNA in plasmids, which makes construction of recombinant genomes feasible. Furthermore, because the signals directing AAV replication and genome encapsidation are contained within the ITRs of the AAV genome, some or all of the internal approximately 4.3 kb of the genome (encoding replication and structural capsid proteins, rep-cap) may be replaced with foreign DNA. To generate AAV vectors, the rep and cap proteins may be provided in trans. Another significant feature of AAV is that it is an extremely stable and hearty virus. It easily withstands the conditions used to inactivate adenovirus (56° to 65° C. for several hours), making cold preservation of AAV less critical. AAV may even be lyophilized. Finally, AAV-infected cells are not resistant to superinfection.

There remains a need in the art for treatments for LGMD2A.

SUMMARY

Methods and products for delivering DNA encoding a protein with calpain3 (CAPN3) activity are provided herein. Such methods and products can be used to treat various diseases, for example, LGMD2A.

Recombinant adeno-associated viruses (rAAVs) are provided encoding a protein with calpain 3 (CAPN3) activity. The recombinant adeno-associated viruses comprise a polynucleotide that comprises a nucleotide sequence encoding the protein with CAPN3 activity. The nucleotide sequence encoding the protein with CAPN3 activity, for example, is at least 90% identical to SEQ ID NO: 2 or comprises the sequence of SEQ ID NO: 2.

For example, the provided rAAV comprise a polynucleotide which comprises a first AAV inverted terminal repeat (ITR), a promoter, a nucleotide sequence encoding a protein with calpain 3 (CAPN3) activity and a second AAV ITR. The nucleotide sequence encoding the protein with CAPN3 activity, for example, is at least 90% identical to SEQ ID NO: 2, or at least 91% identical to SEQ ID NO: 2, at least 92% identical to SEQ ID NO: 2, at least 93% identical to SEQ ID NO: 2, at least 94% identical to SEQ ID NO: 2, at least 95% identical to SEQ ID NO: 2, at least 96% identical to SEQ ID NO: 2, at least 97% identical to SEQ ID NO: 2, at least 98% identical to SEQ ID NO: 2, or at least 99% identical to SEQ ID NO: 2. The rAAV comprises a nucleotide sequence encoding a protein with CAPN3 activity comprises the sequence of SEQ ID NO: 2.

In addition, the provided rAAV comprises a nucleotide sequence encoding a protein with CAPN3 activity that comprises an amino acid sequence that is at least 90% identical to SEQ ID NO:7, at least 91% identical to SEQ ID NO:7, at least 92% identical to SEQ ID NO:7, at least 93% identical to SEQ ID NO:7, at least 94% identical to SEQ ID NO:7, at least 95% identical to SEQ ID NO:7, at least 96% identical to SEQ ID NO:7, at least 97% identical to SEQ ID

NO:7, at least 98% identical to SEQ ID NO:7, or at least 99% identical to SEQ ID NO: 7. The rAAV comprises a nucleotide sequence encoding a protein with CAPN3 activity comprising the amino acid sequence of SEQ ID NO: 7.

The provided rAAV comprise a polynucleotide sequence which is at least 90% identical to SEQ ID NO: 1, at least 91% identical to SEQ ID NO: 1, at least 92% identical to SEQ ID NO: 1, at least 93% identical to SEQ ID NO: 1, at least 94% identical to SEQ ID NO: 1, at least 95% identical to SEQ ID NO: 1, at least 96% identical to SEQ ID NO: 1, at least 97% identical to SEQ ID NO: 1, at least 98% identical to SEQ ID NO: 1, or at least 99% identical to SEQ ID NO: 1. The rAAV comprises a polynucleotide sequence of SEQ ID NO: 1.

The nucleotide sequence, in one embodiment, is under the transcription control of a muscle-specific promoter. For example, the muscle-specific promoter comprises one or more of a human skeletal actin gene element, a cardiac actin gene element, a desmin promoter, a skeletal alpha-actin (ASKA) promoter, a troponin I (TNNI2) promoter, a myocyte-specific enhancer binding factor mef binding element, a muscle creatine kinase (MCK) promoter, a truncated MCK (tMCK) promoter, a myosin heavy chain (MHC) promoter, a hybrid a-myosin heavy chain enhancer-MCK enhancer-promoter (MHCK7) promoter, a C5-12 promoter, a murine creatine kinase enhancer element, a skeletal fast-twitch troponin c gene element, a slow-twitch cardiac troponin c gene element, a slow-twitch troponin i gene element, hypoxia-inducible nuclear factor (HIF)-response element (HRE), a steroid-inducible element, and a glucocorticoid response element (gre). In one embodiment, the muscle-specific promoter is a tMCK promoter, which comprises a sequence of SEQ ID NO: 3.

For example, the rAAV comprises a polynucleotide which comprises, in one embodiment, a first AAV inverted terminal repeat (ITR), a tMCK promoter, the nucleotide sequence encoding the protein with calpain 3 activity, and a second AAV inverted terminal repeat (ITR). The AAV ITR (e.g., the first and/or second AAV ITRs) is, for example, an AAV2 inverted terminal repeat. The capsid proteins of the rAAV comprise, for example, an AAV rh.74 capsid protein or an AAV9 capsid protein.

The provided rAAV comprises one or more of AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-6, AAV-7, AAV-8, AAV-9, AAV-10, AAV-11, AAV-12, AAV-13, AAV rh.74 and AAV rh.10 capsid proteins.

In another embodiment, compositions comprising any of the disclosed rAAV are provided. For example, the compositions are formulated for intramuscular injection or intravenous injection.

Methods of treating limb girdle muscular dystrophy 2A in a subject comprising administering to the subject a therapeutically effective amount of any of the disclosed rAAV or any composition comprising a disclosed rAAV are also provided. In any of the provided methods, the rAAV are administered by intramuscular injection or intravenous injection.

For example, in these methods treatment results in one or more of: (a) an increased muscle fiber diameter, (b) a decreased number of small lobulated muscle fibers, (c) a decreased number of fibers with internal nuclei, (d) a decreased endomysial connective tissue content, (e) correction of muscle atrophy, and (f) a increased muscle force generation. The muscle fiber affected by the treatment comprise one or more of slow twitch oxidative (STO) muscle fiber, fast twitch oxidative (FTO) muscle fiber, and fast twitch glycolytic (FTG) fiber.

In addition, in any of the provided methods, the treatment results in one or more of: (a) at least a 5%, 10%, 15%, 20%, 25%, 30%, or 35%, or 40% decrease of total muscle fiber number per mm² by 4 weeks after administration; (b) at least a 5%, 10%, 15%, 20%, or 25% increase of muscle fiber diameter by 4 weeks after administration; (c) at least a 5%, 10%, 15%, 20%, 25%, 30%, 35%, or 42% decrease of STO muscle fiber number per mm² by 4 weeks after administration; (d) at least a 5%, 10%, 15%, 20%, or 25% increase of STO muscle fiber diameter by 4 weeks after administration; (e) at least a 5%, 10%, 15%, or 20% decrease of FTO muscle fiber number per mm² by 4 weeks after administration; (f) at least a 5%, 10%, 15%, or 20% increase of FTO muscle fiber diameter by 4 weeks after administration; (g) at least a 5%, 10%, 15%, 20%, 25%, 30%, or 35% decrease of FTG muscle fiber number per mm² by 4 weeks after administration; and (h) at least a 5%, 10%, 15%, 20%, or 25% increase of FTG muscle fiber diameter by 4 weeks after administration.

In any of the provided methods, the heart muscle of the subject shows minimum or low calpain 3 protein expressed from any of the provided rAAV, or a composition comprising any of the provided rAAV. The muscle fiber affected by the treatment with the composition comprise one or more of slow twitch oxidative (STO) muscle fiber, fast twitch oxidative (FTO) muscle fiber, and fast twitch glycolytic (FTG) fiber.

Compositions for treating limb girdle muscular dystrophy 2A comprising a therapeutically effective amount of any of the disclosed rAAV or a composition comprising any of the disclosed rAAV are provided. These composition for treating treating limb girdle muscular dystrophy 2A are formulated for administration by intramuscular injection or intravenous injection. In addition, treatment with any of the disclosed compositions limb girdle muscular dystrophy 2A results in one or more of: (a) an increased muscle fiber diameter, (b) a decreased number of small lobulated muscle fibers, (c) a decreased number of fibers with internal nuclei, (d) a decreased endomysial connective tissue content, (e) correction of muscle atrophy, and (f) a increased muscle force generation. The muscle fiber affected by the treatment with the composition comprise one or more of slow twitch oxidative (STO) muscle fiber, fast twitch oxidative (FTO) muscle fiber, and fast twitch glycolytic (FTG) fiber.

In addition, the treatment with any of the disclosed compositions for treating limb girdle muscular dystrophy 2A results in one or more of: (a) at least a 5%, 10%, 15%, 20%, 25%, 30%, or 35%, or 40% decrease of total muscle fiber number per mm² by 4 weeks after administration; (b) at least a 5%, 10%, 15%, 20%, or 25% increase of muscle fiber diameter by 4 weeks after administration; (c) at least a 5%, 10%, 15%, 20%, 25%, 30%, 35%, or 42% decrease of STO muscle fiber number per mm² by 4 weeks after administration; (d) at least a 5%, 10%, 15%, 20%, or 25% increase of STO muscle fiber diameter by 4 weeks after administration; (e) at least a 5%, 10%, 15%, or 20% decrease of FTO muscle fiber number per mm² by 4 weeks after administration; (f) at least a 5%, 10%, 15%, or 20% increase of FTO muscle fiber diameter by 4 weeks after administration; (g) at least a 5%, 10%, 15%, 20%, 25%, 30%, or 35% decrease of FTG muscle fiber number per mm² by 4 weeks after administration; and (h) at least a 5%, 10%, 15%, 20%, or 25% increase of FTG muscle fiber diameter by 4 weeks after administration.

Further, treatment with any of the provided compositions for treatment of limb girdle muscular dystrophy 2A results in the heart muscle of the subject showing minimum or low

calpain 3 protein expressed from any of the provided rAAV, or a composition comprising any of the provided rAAV. The heart muscle, after administration with the rAAV, shows no or little toxic effect, e.g., inflammation, necrosis and/or regeneration.

The disclosure also provides for use of a therapeutically effective amount of any of the disclosed rAAV or a composition comprising any of the disclosed rAAV for the preparation of a medicament for the treatment of limb girdle muscular dystrophy 2A. For example, the medicament is formulated for administration by intramuscular injection or intravenous injection.

In any of the uses, treatment with the medicament results in one or more of: (a) an increased muscle fiber diameter, (b) a decreased number of small lobulated muscle fibers, (c) a decreased number of fibers with internal nuclei, (d) a decreased endomysial connective tissue content, (e) correction of muscle atrophy, and (f) a increased muscle force generation. The muscle fiber affected by treatment with the medicament is one or more of slow twitch oxidative (STO) muscle fiber, fast twitch oxidative (FTO) muscle fiber, and fast twitch glycolytic (FTG) fiber.

In addition, in any of the uses of a therapeutically effective amount of any of the disclosed rAAV or a composition provided, treatment with the medicament results in one or more of: (a) at least a 5%, 10%, 15%, 20%, 25%, 30%, or 35%, or 40% decrease of total muscle fiber number per mm² by 4 weeks after administration; (b) at least a 5%, 10%, 15%, 20%, or 25% increase of muscle fiber diameter by 4 weeks after administration; (c) at least a 5%, 10%, 15%, 20%, 25%, 30%, 35%, or 42% decrease of STO muscle fiber number per mm² by 4 weeks after administration; (d) at least a 5%, 10%, 15%, 20%, or 25% increase of STO muscle fiber diameter by 4 weeks after administration; (e) at least a 5%, 10%, 15%, or 20% decrease of FTO muscle fiber number per mm² by 4 weeks after administration; (f) at least a 5%, 10%, 15%, or 20% increase of FTO muscle fiber diameter by 4 weeks after administration; (g) at least a 5%, 10%, 15%, 20%, 25%, 30%, or 35% decrease of FTG muscle fiber number per mm² by 4 weeks after administration; and (h) at least a 5%, 10%, 15%, 20%, or 25% increase of FTG muscle fiber diameter by 4 weeks after administration.

The any of the uses of a therapeutically effective amount of any of the disclosed rAAV or a composition provided, after treatment with the medicament, the heart muscle of the subject shows no, minimum or low calpain 3 protein expressed from the disclosed or disclosed composition.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1F show that gene therapy restored impaired regeneration in CAPN3-KO muscle. Schematic diagram of single-stranded AAV9.CAPN3 rAAV is shown in FIG. 1A. In between the 5' and 3' single strand ITRs (inverted terminal repeats), the muscle creatine kinase (MCK) promoter (563 bp) drives the expression of CAPN3 open reading frame (2466 bp). Also labeled is polyadenylation site (Poly A, 53 bp). Tibialis anterior (TA) muscles from CAPN3-KO mice were first injected with CTX, and 2 weeks later with 1×10^{11} vg of AAV.CAPN3 to left TA (FIG. 1B) or PBS to right TA (FIG. 1C). Four weeks after rAAV injection, the muscle diameter increased and the lobulated fibers were less common compared to the untreated CAPN3-KO muscle. In FIG. 1D, lobulated fibers with a pattern of subsarcolemmal organelle, mitochondria distribution (arrows) suggest partial myotube fusion in the untreated CAPN3-KO muscle at higher magnification. Scale bar=20 μ m for B-D. In FIG. 1E, the

muscle fiber size distribution histograms (mean \pm SEM/mm² area; derived from 3 mice in each group) of the treated and untreated TA muscle from CAPN3-KO mice show a shift to larger diameter fibers with the treatment and an increase in the small diameter subpopulation present in the untreated group. In FIG. 1F, the Slow twiTch Oxidative (STO) fiber size distribution histograms show a larger number of small fibers (e.g., fiber diameters equal to or less than 30 μ m) in the untreated CAPN3-KO muscle as compared to treated CAPN3-KO muscle.

FIG. 2 shows a schematic diagram of the rAAV of this disclosure, named as "AAVrh.74.tMCK.CAPN3."

FIGS. 3A-3B provide Western Blot (panel A) and RT-PCR (panel B) data after AAVrh.74.tMCK.CAPN3 administration via intramuscular injection (1E11 vg) and systemic injection (3E12 vg and 6E12 vg). This data was compared with normal human muscle lysate (Gel load of 60% total protein as compared to mouse lysates) and untreated CAPN3-KO mice.

FIG. 4 provides representative images of SDH-stained tissue sections of CAPN3 KO (AAV.hCAPN3 gene injected and untreated) and wild type (WT) TA muscles. Mean fiber size of slow twitch oxidative (STO, dark), fast twitch oxidative (FTO, intermediate) and fast twitch glycolytic (FTG, light) fibers appeared normalized towards WT values in the TA muscle of mice treated with AAVrh.74.tMCK.CAPN3. Fiber type sizes with and without treatment are illustrated in the Table 4.

FIG. 5 provides relative CAPN3 protein expression levels in WT (Z18-14) and TA muscles from the low dose cohort (3E12 vg, Z18-13, Z18-15, Z18-16, Z18-17, Z18-18), and gastrocnemius (gastroc), heart, quadriceps, tibialis anterior (TA) and triceps from the high dose cohort (6E12 vg, Z18-20, Z18-21, Z18-23, Z18-24, Z18-22) are shown (UT: untreated).

FIG. 6 provides AAVrh74.tMCK.hCAPN3 vector copies/ μ g genomic DNA in 6E12 vector genome systemic high dose cohort in the following muscles: quadriceps (quad), heart, tibialis anterior (TA), gastrocnemius (gastroc) triceps, and the liver.

FIG. 7 provides the mean fiber diameters of slow twitch oxidative (STO, dark), fast twitch oxidative (FTO, intermediate) and fast twitch glycolytic (FTG, light) fibers from left TA muscle following systemic administration of AAVrh.74.tMCK.CAPN3 at 3E12 and 6E12 vg. Data from untreated CAPN3KO and WT mice was included.

FIG. 8 provides the data for the run-to-exhaustion test. FIG. 8A provides data for the low dose cohort, which received 3E12 vg of AAVrh.74.tMCK.CAPN3, and the high dose cohort, which received 6E12 vg of AAVrh.74.tMCK.CAPN3 4 weeks after systemic administration. Treated CAPN3 KO mice performed better on Run-to-Exhaustion test compared to untreated counterparts. FIG. 8B provides data for the high dose cohort, in which the mice were tested 20-24 weeks after systemic administration of 6E12 vg of AAVrh.74.tMCK.CAPN3 (n=5) and untreated counterparts (n=16)

FIG. 9 provides hematoxylin & eosin(H&E) stained fresh frozen sections of the left ventricles from representative heart samples of CAPN3 KO mice at 4 weeks post-systemic injection of the AAVrh7.4.tMCK.hCAPN3 vector at 3E12 vg and 6E12 vg doses with matching untreated controls.

FIG. 10 provides Western blot analysis of the cardiac tissues from the high dose cohort (which received 6E12 vg of AAVrh7.4.tMCK.hCAPN3. This analysis showed no or minimum detectable calpain 3 protein in the heart of the

treated animal. Animal identification numbers Z18-19 and 22 represent the lysates from the untreated CAPN3 KO mice.

DETAILED DESCRIPTION

Recombinant AAVs (rAAVs) provided herein comprise a polynucleotide that comprises a first AAV inverted terminal repeat (ITR), a promoter, a nucleotide sequence encoding a protein with calpain 3 (CAPN3) activity and a second AAV ITR. In one embodiment, the nucleotide encodes CAPN3. Embodiments include, but are not limited to, an rAAV comprising a nucleotide sequence encoding CAPN3 or a protein with CAPN3 activity, wherein the nucleotide sequence is at least 65%, at least 70%, at least 75%, at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, or 89% identical to the nucleotide sequence of SEQ ID NO: 2. Additional embodiments include, but are also not limited to, rAAV comprising a nucleotide sequence at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identical to the nucleotide sequence set forth in SEQ ID NO: 2 and encodes a polypeptide with a CAPN3 proteolytic activity. The CAPN3 proteolytic activity is understood in the art as the activity of proteolyzing potential substrates such as fodrin and HSP60, and/or to the activity of autolytically self-cleaving. Thus, as used herein, the term "a protein with calpain 3 (CAPN3) activity" refers to a protein with CAPN3 proteolytic activity, which includes but is not limited to the activity of proteolyzing substrates such as fodrin and HSP60, and/or to the activity of autolytically self-cleaving. The protein with CAPN3 activity can have the full or partial activity of a full length calpain 3 protein. In one embodiment, the protein with CAPN3 activity has at least 60%, 70%, 80%, 90%, 95%, or 99% of activity of a full length CAPN3 protein. In another embodiment, the protein with CAPN3 activity comprises an amino acid sequence with at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to SEQ ID NO: 7.

In some embodiments, the nucleotide sequence encoding the protein with CAPN3 activity comprises a sequence of SEQ ID NO: 2. In another embodiment, the protein with CAPN3 activity comprises an amino acid sequence with at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to SEQ ID NO: 7. In another embodiment, the protein with CAPN3 activity comprises the amino acid sequence of SEQ ID NO: 7. In another embodiment, the polynucleotide of the rAAV comprises a sequence which is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to SEQ ID NO: 1. In another embodiment, the polynucleotide comprises a sequence at least 95% identical to SEQ ID NO: 1. In one embodiment, the polynucleotide comprises the sequence of SEQ ID NO: 1.

In another aspect, described herein is a recombinant AAV comprising a nucleotide sequence that encodes a protein with CAPN3 activity and/or that comprises a nucleotide sequence that hybridizes under stringent conditions to the nucleic acid sequence of SEQ ID NO: 2, or the complement thereof. The term "stringent" is used to refer to conditions that are commonly understood in the art as stringent. Hybridization stringency is principally determined by temperature, ionic strength, and the concentration of denaturing agents such as formamide. Examples of stringent conditions for hybridization and washing are 0.015 M sodium chloride, 0.0015 M sodium citrate at 65-68° C. or 0.015 M sodium chloride, 0.0015M sodium citrate, and 50% formamide at

42° C. See Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, (Cold Spring Harbor, N.Y. 1989).

In recombinant genomes described herein, the CAPN3 polynucleotide is operatively linked to transcriptional control elements (including, but not limited to, promoters, enhancers and/or introns), specifically transcriptional control elements functional in target cells of interest. For example, various embodiment provide methods of transducing muscle cells using muscle-specific transcriptional control elements, including, but not limited to, those derived from the actin and myosin gene families, such as from the myoD gene family [See Weintraub et al., *Science*, 251: 761-766 (1991)], the myocyte-specific enhancer binding factor MEF-2 [Cserjesi and Olson, *Mol Cell Biol*, 11: 4854-4862 (1991)], control elements derived from the human skeletal actin gene [Muscat et al., *Mol Cell Biol*, 7: 4089-4099 (1987)], muscle creatine kinase sequence elements [See Johnson et al., *Mol Cell Biol*, 9:3393-3399 (1989)] and the murine creatine kinase enhancer (mCK) element, control elements derived from the skeletal fast-twitch troponin C gene, the slow-twitch cardiac troponin C gene and the slow-twitch troponin I gene: hypoxia-inducible nuclear factors [Semenza et al., *Proc Natl Acad Sci USA*, 88: 5680-5684 (1991)], steroid-inducible elements and promoters including the glucocorticoid response element (GRE) [See Mader and White, *Proc. Natl. Acad. Sci. USA*, 90: 5603-5607 (1993)], the tMCK promoter [see Wang et al., *Gene Therapy*, 15: 1489-1499 (2008)], the CK6 promoter [see Wang et al., *supra*] and other control elements. In one embodiment, the nucleotide sequence encoding a protein with calpain 3 (CAPN3) activity is operably linked to a muscle-specific promoter. In one embodiment, the muscle-specific promoter comprises one or more of a human skeletal actin gene element, a cardiac actin gene element, a desmin promoter, a skeletal alpha-actin (ASKA) promoter, a troponin I (TNNI2) promoter, a myocyte-specific enhancer binding factor mef binding element, a muscle creatine kinase (MCK) promoter, a truncated MCK (tMCK) promoter, a myosin heavy chain (MHC) promoter, a hybrid a-myosin heavy chain enhancer-/MCK enhancer-promoter (MHCK7) promoter, a C5-12 promoter, a murine creatine kinase enhancer element, a skeletal fast-twitch troponin c gene element, a slow-twitch cardiac troponin c gene element, a slow-twitch troponin i gene element, hypoxia-inducible nuclear factor (HIF)-response element (HRE), a steroid-inducible element, a glucocorticoid response element (gre). In another embodiment, the muscle-specific promoter is an MCK promoter, a tMCK promoter, or an MHCK7 promoter. In some embodiments, the muscle-specific promoter is tMCK that comprises a nucleotide sequence of SEQ ID NO: 3.

Previous studies showed that expression of CAPN3 driven by desmin promoter resulted in cardiotoxicity. In follow up studies, selective skeletal muscle expression of the gene eliminated the cardiac defects. The AAV genomes disclosed herein comprise a muscle specific promoter, tMCK to restrict CAPN3 expression to the skeletal muscle and showed no cardiac toxicity following systemic delivery of the virus at 6E12 vg (twice the proposed initial high dose) 4 weeks after gene injection.

The rAAV genomes described herein lack AAV rep and cap DNA. rAAV genomes provided comprise a CAPN3 polynucleotide as described above and one or more AAV ITRs flanking the polynucleotide. AAV DNA in the rAAV genomes may be from any AAV serotype for which a recombinant virus can be derived including, but not limited

to, AAV serotypes AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-6, AAV-7, AAV-8, AAV-9, AAV-10, AAV-11, AAV-12, AAV-13, AAV rh.74 and AAV rh.10. Other types of rAAV variants, for example rAAV with capsid mutations, are also contemplated. See, for example, Marsic et al., *Molecular Therapy*, 22(11): 1900-1909 (2014). As noted in the Background section above, the nucleotide sequences of the genomes of various AAV serotypes are known in the art. To promote skeletal muscle specific expression, AAV1, AAV5, AAV6, AAV8 or AAV9 may be used.

DNA plasmids provided comprise rAAV genomes. The DNA plasmids are transferred to cells permissible for infection with a helper virus of AAV (including, but not limited to, adenovirus, E1-deleted adenovirus or herpesvirus) for assembly of the rAAV genome into infectious viral particles. Techniques to produce rAAV particles, in which an AAV genome to be packaged, rep and cap genes, and helper virus functions are provided to a cell are standard in the art. Production of rAAV requires that the following components are present within a single cell (denoted herein as a packaging cell): a rAAV genome, AAV rep and cap genes separate from (i.e., not in) the rAAV genome, and helper virus functions. The AAV ITRs and rep and cap genes may be from any AAV serotype for which recombinant virus can be derived and may be from a different AAV serotype than the rAAV genome ITRs, including, but not limited to, AAV serotypes AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-6, AAV-7, AAV-8, AAV-9, AAV-10, AAV-11, AAV-12, AAV-13, AAV rh. 10 and AAV rh.74. Production of pseudotyped rAAV is disclosed in, for example, WO 01/83692 which is incorporated by reference herein in its entirety. Thus, in one embodiment, the rAAV comprises one or more of AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-6, AAV-7, AAV-8, AAV-9, AAV-10, AAV-11, AAV-12, AAV-13, AAV rh. 10 and AAV rh.74 capsid proteins. In another embodiment, the rAAV comprises an AAV rh.74 capsid protein or an AAV9 capsid protein.

A method of generating a packaging cell is to create a cell line that stably expresses all the necessary components for AAV particle production. For example, a plasmid (or multiple plasmids) comprising a rAAV genome lacking AAV rep and cap genes, AAV rep and cap genes separate from the rAAV genome, and a selectable marker, such as a neomycin resistance gene, are integrated into the genome of a cell. AAV genomes have been introduced into bacterial plasmids by procedures such as GC tailing [Samulski et al., *Proc. Natl. Acad. Sci. USA*, 79:2077-2081 (1982)], addition of synthetic linkers containing restriction endonuclease cleavage sites [Laughlin et al., *Gene*, 23:65-73 (1983)] or by direct, blunt-end ligation [Senapathy & Carter, *J. Biol. Chem.*, 259:4661-4666 (1984)]. The packaging cell line is then infected with a helper virus such as adenovirus. The advantages of this method are that the cells are selectable and are suitable for large-scale production of rAAV. Other examples of suitable methods employ adenovirus or baculovirus rather than plasmids to introduce rAAV genomes and/or rep and cap genes into packaging cells.

General principles of rAAV production are reviewed in, for example, Carter, *Current Opinions in Biotechnology*, 1533-1539 (1992); and Muzyczka, *Curr. Topics in Microbial. and Immunol.*, 158:97-129 (1992). Various approaches are described in Ratschin et al., *Mol. Cell. Biol.*, 4:2072 (1984); Hermonat et al., *Proc. Natl. Acad. Sci. USA*, 81:6466 (1984); Tratschin et al., *Mol. Cell. Biol.*, 5:3251 (1985); McLaughlin et al., *J. Virol.*, 62:1963 (1988); Lebkowski et al., *Mol. Cell. Biol.*, 7:349 (1988); Samulski et al., *J. Virol.*, 63:3822-3828 (1989); U.S. Pat. No. 5,173,414; WO

95/13365 and corresponding U.S. Pat. No. 5,658,776; WO 95/13392; WO 96/17947; PCT/US98/18600; WO 97/09441 (PCT/US96/14423); WO 97/08298 (PCT/US96/13872); WO 97/21825 (PCT/US96/20777); WO 97/06243 (PCT/FR96/01064); WO 99/11764; Perrin et al., *Vaccine*, 13:1244-1250 (1995); Paul et al., *Human Gene Therapy*, 4:609-615 (1993); Clark et al., *Gene Therapy* 3:1124-1132 (1996); U.S. Pat. Nos. 5,786,211; 5,871,982; 6,258,595; and McCarty, *Mol. Ther.*, 16(10): 1648-1656 (2008). The foregoing documents are hereby incorporated by reference in their entirety herein, with particular emphasis on those sections of the documents relating to rAAV production.

Thus packaging cells are provided that produce infectious rAAV. In one embodiment, packaging cells may be stably transformed cancer cells such as HeLa cells, and PerC.6 cells (a cognate 293 line). In another embodiment, packaging cells are cells that are not transformed cancer cells, such as low passage 293 cells (human fetal kidney cells transformed with E1 of adenovirus), MRC-5 cells (human fetal fibroblasts), WI-38 cells (human fetal fibroblasts), Vero cells (monkey kidney cells) and FRhL-2 cells (rhesus fetal lung cells).

Recombinant AAV provided herein are thus replication-deficient, infectious, encapsidated viral particles which comprise a recombinant genome. Examples include, but are not limited to, a rAAV including a genome comprising the sequence set out in SEQ ID NO: 1 encoding CAPN3, a rAAV including a genome consisting essentially of the sequence set out in SEQ ID NO: 1 encoding CAPN3, and a rAAV (named "AAVrh.74.tMCK.CAPN3") including a genome consisting of the sequence set out in SEQ ID NO: 1 encoding CAPN3. The genomes of the rAAV lack AAV rep and cap DNA, that is, there is no AAV rep or cap DNA between the ITRs of the rAAV genome.

The sequence of the AAVrh.74.tMCK.CAPN3 sequence is set out in SEQ ID NO: 1, in which an AAV2 ITR spans nucleotides 1-128, the tMCK promoter spans nucleotides 165-884, a chimeric intron spans nucleotides 937-1069, a Kozak Sequence spans nucleotides 1101-1106, the CAPN3 polynucleotide spans nucleotides 1107-3572, a poly A signal spans nucleotides 3581-3780, and a second AAV2 ITR spans nucleotides 3850-3977.

The rAAV may be purified by methods known in the art such as by column chromatography or cesium chloride gradients. Methods for purifying rAAV vectors from helper virus are known in the art and include methods disclosed in, for example, Clark et al., *Hum. Gene Ther.*, 10(6): 1031-1039 (1999); Schenpp and Clark, *Methods Mol. Med.*, 69: 427-443 (2002); U.S. Pat. No. 6,566,118; and WO 98/09657.

In another embodiment, compositions comprising rAAV described herein are provided. Compositions provided comprise rAAV in a pharmaceutically acceptable carrier. The compositions may also comprise other ingredients such as diluents and adjuvants. Acceptable carriers, diluents and adjuvants are nontoxic to recipients and are preferably inert at the dosages and concentrations employed, and include buffers such as phosphate, citrate, or other organic acids; antioxidants such as ascorbic acid; low molecular weight polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such

as sodium; and/or nonionic surfactants such as Tween, pluronics or polyethylene glycol (PEG).

Titers of rAAV to be administered in methods described herein can vary depending, for example, on the particular rAAV, the mode of administration, the treatment goal, the individual, and the cell type(s) being targeted, and may be determined by methods standard in the art. Titers of rAAV may range from about 1×10^{10} , about 1×10^{11} , about 1×10^{12} , about 1×10^{13} , to about 1×10^{14} , or more DNase resistant particles (DRPs) per ml. Dosages may also be expressed in units of viral genomes (vg). Exemplary disclosed doses include 1E11 vg, 3E12 vg and 6E12 vg.

Methods of transducing a target cell such as a muscle cell with rAAV, in vivo or in vitro, are contemplated herein. The in vivo methods comprise the step of administering an effective dose, or effective multiple doses, of a composition comprising a rAAV provided herein to subject (e.g., an animal including but not limited to a human patient) in need thereof. If the dose is administered prior to development of a disorder/disease, the administration is prophylactic. If the dose is administered after the development of a disorder/disease, the administration is therapeutic. An effective dose is a dose that alleviates (eliminates or reduces) at least one symptom associated with the disorder/disease state being treated, that slows or prevents progression to a disorder/disease state, that slows or prevents progression of a disorder/disease state, that diminishes the extent of disease, that results in remission (partial or total) of disease, and/or that prolongs survival. In comparison to the subject before treatment, methods herein result in one or more of: an increased muscle fiber diameter, a decreased number of small lobulated muscle fibers, a decreased number of fibers with internal nuclei, a decreased endomysial connective tissue content, correction of muscle atrophy, and an increased muscle force generation. In one embodiment, the muscle fiber comprises one or more of slow twitch oxidative (STO) muscle fiber, fast twitch oxidative (FTO) muscle fiber, and fast twitch glycolytic (FTG) fiber. In one embodiment, the treatment results in one or more of (a) at least a 5%, 10%, 15%, 20%, 25%, 30%, or 35%, or 40% decrease of total muscle fiber number per mm^2 by 4 weeks after administration; (b) at least a 5%, 10%, 15%, 20%, or 25% increase of muscle fiber diameter by 4 weeks after administration; (c) at least a 5%, 10%, 15%, 20%, 25%, 30%, 35%, or 42% decrease of STO muscle fiber number per mm^2 by 4 weeks after administration; (d) at least a 5%, 10%, 15%, 20%, or 25% increase of STO muscle fiber diameter by 4 weeks after administration; (e) at least a 5%, 10%, 15%, or 20% decrease of FTO muscle fiber number per mm^2 by 4 weeks after administration; (f) at least a 5%, 10%, 15%, or 20% increase of FTO muscle fiber diameter by 4 weeks after administration; (g) at least a 5%, 10%, 15%, 20%, 25%, 30%, or 35% decrease of FTG muscle fiber number per mm^2 by 4 weeks after administration; and (h) at least a 5%, 10%, 15%, 20%, or 25% increase of FTG muscle fiber diameter by 4 weeks after administration. The method of this disclosure, in one embodiment, leads to no, minimum or low calpain 3 protein expressed from the rAAV in the heart muscle of the subject administered with the rAAV.

Assays to examine these results are understood in the art and/or are described in the examples herein. Use of the methods described herein to prevent or treat disorders/diseases (e.g., muscular dystrophies) caused by defects in CAPN3 activity or defects in expression of CAPN3 is contemplated. LGMD2A is an example of a disease contemplated for prevention or treatment according to the methods.

Combination therapies are also contemplated. Combination as used herein includes both simultaneous treatment or sequential treatments. Combinations of methods described herein with standard medical treatments (e.g., corticosteroids) are specifically contemplated, as are combinations with novel therapies.

Administration of an effective dose of the compositions may be by routes standard in the art including, but not limited to, intramuscular, parenteral, intravenous, intrathecal, oral, buccal, nasal, pulmonary, intracranial, intraosseous, intraocular, rectal, or vaginal. Route(s) of administration and serotype(s) of AAV components of the rAAV (in particular, the AAV ITRs and capsid protein) may be chosen and/or matched by those skilled in the art taking into account the infection and/or disease state being treated and the target cells/tissue(s) that are to express the CAPN3. In one embodiment, the rAAV is administered by intramuscular injection, intravenous injection, intraperitoneal injection, subcutaneous injection, epicutaneous administration, intravaginal injection, intradermal administration, or nasal administration. In another embodiment, the rAAV is administered by intramuscular injection or intravenous injection.

In particular, actual administration of rAAV described herein may be accomplished by using any physical method that will transport the rAAV recombinant vector into the target tissue of an animal. Administration includes, but is not limited to, injection into muscle, the bloodstream, and/or directly into the liver. Simply resuspending a rAAV in phosphate buffered saline has been demonstrated to be sufficient to provide a vehicle useful for muscle tissue expression, and there are no known restrictions on the carriers or other components that can be co-administered with the rAAV. Capsid proteins of a rAAV may be modified so that the rAAV is targeted to a particular target tissue of interest such as muscle. See, for example, WO 02/053703, the disclosure of which is incorporated by reference herein. Pharmaceutical compositions can be prepared as injectable formulations or as topical formulations to be delivered to the muscles by transdermal transport. Numerous formulations for both intramuscular injection and transdermal transport have been previously developed and can be used in the practice of the methods. The rAAV can be used with any pharmaceutically acceptable carrier for ease of administration and handling.

For purposes of intramuscular injection, solutions in an adjuvant such as sesame or peanut oil or in aqueous propylene glycol can be employed, as well as sterile aqueous solutions. Such aqueous solutions can be buffered, if desired, and the liquid diluent first rendered isotonic with saline or glucose. Solutions of rAAV as a free acid (DNA contains acidic phosphate groups) or a pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. A dispersion of rAAV can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms. In this connection, the sterile aqueous media employed are all readily obtainable by standard techniques well-known to those skilled in the art.

The pharmaceutical forms suitable for systemic (e.g., intravenous) injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be

preserved against the contaminating actions of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of a dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like. In many cases it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by use of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating rAAV in the required amount in the appropriate solvent with various other ingredients enumerated above, as required, followed by filter sterilization. Generally, dispersions are prepared by incorporating the sterilized active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the methods of preparation, in some embodiments, comprises vacuum drying and/or the freeze drying technique, each of which can yield a powder of the active ingredient plus any additional desired ingredient from the previously sterile-filtered solution thereof.

Transduction with rAAV may also be carried out in vitro. In one embodiment, desired target muscle cells are removed from the subject, transduced with rAAV and reintroduced into the subject. Alternatively, syngeneic or xenogeneic muscle cells can be used where those cells will not generate an inappropriate immune response in the subject.

Suitable methods for the transduction and reintroduction of transduced cells into a subject are known in the art. In one embodiment, cells can be transduced in vitro by combining rAAV with muscle cells, e.g., in appropriate media, and screening for those cells harboring the DNA of interest using conventional techniques such as Southern blots and/or PCR, or by using selectable markers. Transduced cells can then be formulated into pharmaceutical compositions, and the composition introduced into the subject by various techniques, such as by intramuscular, intravenous, subcutaneous and intraperitoneal injection, or by injection into smooth and cardiac muscle, using e.g., a catheter.

Transduction of cells with rAAV by methods described herein results in sustained expression of CAPN3 or a protein with CAPN3 activity. Methods are thus provided for administering rAAV which expresses CAPN3 or a protein with CAPN3 activity to a subject, preferably a human being. The subject of this disclosure includes but is not limited to human, a dog, a cat, a horse, a cow, a pig, a sheep, a goat, a chicken, a rodent (e.g., rats and mice), and a primate. These methods include transducing tissues (including, but not limited to, tissues such as muscle, organs such as liver and brain, and glands such as salivary glands) with one or more rAAV described herein.

Muscle tissue is an attractive target for in vivo DNA delivery, because it is not a vital organ and is easy to access. The methods herein provide sustained expression of CAPN3 from transduced muscle cells.

By "muscle cell," "muscle fiber," or "muscle tissue" is meant a cell or group of cells derived from muscle of any

15

kind [for example, skeletal muscle and smooth muscle (e.g., from the digestive tract, urinary bladder, blood vessels or cardiac tissue)]. Such muscle cells may be differentiated or undifferentiated, such as myoblasts, myocytes, myotubes, cardiomyocytes and cardiomyoblasts.

The term "transduction" is used to refer to the administration/delivery of CAPN3 to a recipient cell either in vivo or in vitro, via a rAAV described resulting in expression of CAPN3 by the recipient cell.

Thus, methods are provided of administering an effective dose (or doses, administered essentially simultaneously or doses given at intervals) of rAAV that encode CAPN3 to a subject in need thereof.

As noted above, the methods described herein result in the subject, in comparison to the subject before treatment, one or more of: increased muscle fiber diameter, decreased number of small lobulated slow twitch oxidative (STO) muscle fibers, decreased number of fibers with internal nuclei, decreased endomysial connective tissue content, correction of muscle atrophy, and increased muscle force generation.

EXAMPLES

Aspects and embodiments are illustrated by the following examples. Example 1 describes production of AAV9.MCK.CAPN3. Example 2 describes intramuscular administration of AAV9.MCK.CAPN3. Example 3 describes production of AAVrh.74.tMCK.CAPN3. Example 4 describes intramuscular administration of AAVrh.74.tMCK.CAPN3. Example 5 describes intravenous administration of AAVrh.74.tMCK.CAPN3. Example 6 describes end point studies. Example 7 describes toxicology and biodistribution studies. Example 8 describes in vivo biopotency testing following intramuscular injection. Example 9 describes in vivo biopotency testing following systemic injection. Example 10 describes assessment of systemic AAVrh.74.tMCK.CAPN3 gene delivery. Example 11 describes assessment of cardiac toxicity following systemic injection of AAVrh.74.tMCK.CAPN3 vector. Example 12 describes in vivo physiological analysis.

Example 1

Production of AAV9.MCK.CAPN3

An AAV vector (named AAV.CAPN3) carrying the CAPN3 gene under the muscle specific MCK promoter (FIG. 1A) was produced. A DNA including the open reading frame of mouse CAPN3 (NM_007601.3) between two NotI restriction sites was synthesized by Eurofin Genomics, USA, and then subcloned into a single strand AAV.MCK (muscle creatine kinase) vector previously described in Rodino-Klapac et al., *Journal of Translational Medicine*, 5:45-55 (2007)]. rAAV vectors were produced by a modified cross-packaging approach whereby the AAV type 2 vector genome can be packaged into multiple AAV capsid serotypes. [Rabinowitz et al., *J Virol*. 76 (2):791-801 (2002)]. Production was accomplished using a standard three-plasmid DNA/CaPO₄ precipitation method using HEK293 cells. 293 cells were maintained in DMEM supplemented with 10% fetal bovine serum (FBS) and penicillin and streptomycin. The production plasmids were: (i) pAAV.MCK.microdys, (ii) rep2-capX modified AAV helper plasmids encoding cap serotypes 1, 6, or an 8-like isolate, and (iii) an adenovirus type 5 helper plasmid (pAdhelper) expressing adenovirus E2A, E4 ORF6, and VA I/II RNA genes. To allow compari-

16

sons between serotypes, a quantitative PCR-based titration method was used to determine an encapsidated vector genome (vg) titer utilizing a Prism 7500 Taqman detector system (PE Applied Biosystems). [Clark et al., *Hum Gene Ther*. 10 (6): 1031-1039 (1999)] The primer and fluorescent probe targeted the MCK promoter and were as follows: MCK forward primer, 5-CCCGAGATGCCTGGT-TATAATT-3 (SEQ ID NO: 4); MCK reverse primer, 5-GCTCAGGCAGCAGGTGTTG-3 (SEQ ID NO: 5); and MCK probe, 5-FAM-CCAGACATGTGGCTGCTCCCC-TAMRA-3 (SEQ ID NO: 6). The final titer (vg ml⁻¹) was determined by quantitative reverse transcriptase PCR using the specific primers and probes for MCK promoter utilizing a Prism 7500 Real-time detector system (PE Applied Biosystems, Grand Island, NY, USA). Aliquoted viruses were kept at -80° C. until use.

Example 2

Intramuscular Administration of AAV9.MCK.CAPN3

To demonstrate if WT CAPN3 can restore the impaired regeneration process in CAPN3 knockout (CAPN3-KO) mice, TA muscles from CAPN3-KO mice (n=4) [Kramerova et al., *Hum Mol Genet* 13(13):1373-1388 (2004)] under anesthesia were first injected with 30 µl CTX, and 2 weeks later were transduced to express wild type CAPN3 using AAV9.MCK.CAPN3 at 1×10¹¹ vg in 20 µl volume via intramuscular injection. TA muscles from another cohort of CAPN3-KO (n=4), served as controls received the same volume of PBS 2 weeks post-CTX injection.

Mice were killed at 6 weeks post-CTX injection, and TA muscles were removed and processed for cryostat sectioning. Twelve µm thick cross sections were first stained with H&E for routine histopathological evaluation; muscle fiber type specific diameter measurements were obtained from SDH stained cross sections of the TA from 3 mice in each group. Three random images of the TA (per section per animal) was photographed at X20 magnification and the fiber diameter measurements and fiber type specific histograms were generated.

Succinic dehydrogenase (SDH) enzyme histochemistry was used to assess metabolic fiber type differentiation [slow twitch oxidative (STO), fast twitch oxidative (FTO) and fast twitch glycolytic (FTG)]. Muscle fiber type specific diameter measurements were obtained using 12 µm thick-SDH stained cross sections at 4 and 12 weeks after final cardiotoxin injection. Three images, each representing three distinct zones of the gastrocnemius muscle (a deep zone predominantly composed of STO, intermediate zone showing a checkerboard appearance of STO and FTO or FTG and the superficial zone predominantly composed of FTG fibers) along the midline axis (per section per animal) was photographed at X20 magnification using an Olympus BX41 microscope and SPOT camera (Olympus BX61, Japan). This approach was chosen to capture the alterations in the oxidative state of fibers in each zone in response to metabolic changes during regeneration. Diameters of dark (STO), intermediate (FTO) and light (FTG) fibers were determined by measuring the shortest distance across the muscle fiber using Zeiss Axiovision LE4 software (v.4.8). The fiber diameter histograms were generated separately for STO; FTG and FTO were combined to represent the total fast twitch fiber population (FTG/O), derived from 3 animals and expressed as number per mm² of endomysial area (mean±SEM). The mean fiber diameter was derived from

17

combining all 3 fiber types. An average of 900-1700 fibers were measured per group. TA muscles were used for assessment of fibrosis (see below)

Four weeks after AAV9.MCK.CAPN3 injection, a significant increase in muscle diameter with an apparent decrease of internal nuclei and far less number of small fibers with lobulated pattern was observed (FIG. 1B). The untreated CAPN3-KO muscle had 31.6% more fibers per mm² area, mostly composed of small and lobulated STO fibers indicating that the treatment improved myotube fusion, therefore decreased individual small fiber number per unit area (FIGS. 1, C and D; Table 1).

TABLE 1

Tibialis anterior muscle fiber size				
	Untreated Number per mm ²	Untreated Diameter	AAV.CAPN3- treated Number per mm ²	AAV.CAPN3- treated Diameter
STO	355	32.72 ± 0.4	233	39.81 ± 0.6*
FTG/O	116	44.26 ± 0.9	99	50.40 ± 1.2*
All fibers	471	35.55 ± 0.4	322	43.08 ± 0.6*

*p < 0.0001 compared to same wild type parameter

18

Eurofin Genomics, USA, and then inserted in an AAV production plasmid. A map of the plasmid is shown in FIG. 2.

rAAV vectors were then produced by the approach described in Example 1.

Example 4

Intravenous Administration of
AAVrh.74.tMCK.CAPN3

CAPN3-KO mice, 6 months of age, received AAVrh.74.tMCK.CAPN3 at low (3×10^{12} vg) and high doses (6×10^{12} vg) via injection into tail vein. The mice were killed at 20 weeks post gene injection for endpoint studies. Age matched vehicle treated CAPN3-KO mice served as controls.

TABLE 2

Treatment cohorts						
Cohorts	Treatment	Total # of mice	Age at start of treatment	Treatment Dose (AAVrh.74.tMCK.CAPN3)	Treatment Duration	Age at End Point
CAPN3-KO		40				
Low dose	AAV.CAPN3	8	24 wks	3e12 vg in 300 µl saline, i.v.	20 wks	44 wks
	Saline treatment	8	24 wks		20 wks	44 wks
High Dose	AAV.CAPN3	8	24 wks	6e12 vg in 300 µl saline, i.v.	20 wks	44 wks
	Saline treatment	8	24 wks		20 wks	44 wks
Wild type Controls	Saline treatment	8	24 wks		20 wks	44 wks

The fiber size distribution histograms of the treated TA muscle showed a shift to larger diameter fibers with treatment and the excessive number of small fibers in the untreated CAPN3-KO control muscle are of STO histochemical fiber type (FIGS. 1E and 1F). Collectively, these findings show that CAPN3 replacement via gene therapy in the CAPN3-KO muscle rescued defective regeneration, evidenced by toward normalization of fiber size and a decrease in the number of STO fiber population.

Example 3

Production of AAVrh.74.tMCK.CAPN3

An AAV vector (named AAVrh.74.tMCK.CAPN3) carrying the CAPN3 gene under a truncated muscle specific MCK promoter (tMCK promoter) was produced. A DNA including the open reading frame of mouse CAPN3 (NM_007601.3) between two NotI restriction sites was synthesized by

End point studies performed as described in Example 7 below include muscle physiology (TA force generation or in vivo muscle contractility assay, and protection from eccentric contractions), muscle histopathology, hCAPN3 detection using qPCR, and Western blot analysis.

Example 5

Intramuscular Administration of
AAVrh.74.tMCK.CAPN3

Regenerative responses are measured in old and young CAPN3-KO muscle to cardiotoxin (CTX)-induced synchronized necrosis following the introduction of CAPN3 into regenerating muscle via rAAV treatment.

In cohorts of young (at 2 months of age) and old mice (at 6 months of age), CTX is injected into both TA muscles to induce synchronized necrosis 2 weeks prior to rAAV injection to the left TA muscle. AAVrh.74.tMCK.CAPN3 at

1×10^{11} vg in 20 μ l volume is administered via intramuscular injection. Endpoint studies are performed at 8 weeks post gene transfer (at 1×10^{11} vg dose with efficacy established in our previous studies) to assess the correction of regeneration defect by comparing quantitative histology and physiological outcomes from the left TA to untreated right TA.

TABLE 3

Treatment cohorts							
Cohorts	Treatment	Total # of mice	CTX inj-bilateral TA muscle; age/delivery route/dose	Age at start of gene therapy	Treatment Dose (AAVrh.74.tMCK.CAPN3) Left TA	Treatment Duration	Age at End Point
CAPN3-KO	CTX + AAV.CAPN3	16					
Young		8	6 wks/i.m./30 μ l	8 wks	1×10^{11} vg in 30 μ l PBS, i.m.	8 wks	16 wks
Old		8	22 wks/i.m./30 μ l	24 wks	1×10^{11} vg in 30 μ l PBS, i.m.	8 wks	32 wks
Wild type	CTX only	18					
Young		8	6 wks/i.m./30 μ l	8 wks	30 μ l PBS	8 wks	16 wks
Old		8	22 wks/i.m./30 μ l	24 wks	30 μ l PBS	8 wks	32 wks

Eight weeks post-rAAV injection, end point studies carried out as described in Example 6 below include muscle physiology (TA force generation and protection from eccentric contractions), quantitative muscle histopathology, hCAPN3 detection using qPCR and western blot analysis.

Example 6

End Point Studies

TA Force Generation and Protection from Eccentric Contractions

A protocol to assess functional outcomes in the TA muscle is performed on muscles extracted from mice [Wein et al., *Nature Medicine*, 20(9):992-1000 (2014)]. Mice are anesthetized using ketamine/xylazine mixture. Using a dissecting scope, the hind limb skin is removed to expose the TA muscle and the patella. The distal TA tendon is dissected out and a double square knot is tied around the tendon with 4-0 suture as close to the muscle as possible, and the tendon is cut. The exposed muscle is constantly dampened with saline. Mice are then transferred to a thermal-controlled platform and maintained at 37 degrees. The knee is secured to the platform with a needle through the patella tendon, the distal TA tendon suture to the level arm of the force transducer (Aurora Scientific, Aurora, ON, Canada), and the foot is secured with tape. The TA muscle contractions are elicited by stimulating the sciatic nerve via bipolar platinum electrodes. Once the muscle is stabilized, the optimal length was determined by incrementally stretching the muscle until the maximum twitch force was achieved. After a 3 min rest period, the TA is stimulated at 50,100,150 and 200 Hz, allowing a 1 min rest period between each stimulus to determine maximum tetanic force. Muscle length is measured. Following a 5 min rest, the susceptibility of the TA muscle to contraction induced damage is assessed. After 500 ms of stimulation, the muscle is lengthened by 10% of the optimal length. This includes stimulating the muscle at 150 Hz for 700 ms. After the stimulation, the muscle is returned to the optimal length. The cycle is repeated every minute for a total of 10 cycles. Specific force is calculated by dividing the maximum tetanic force by the TA muscle cross-sectional area. After the eccentric contractions, the mice are then

euthanized, and the TA muscle is dissected out, weighed and frozen for analysis. Analysis of the data is performed blindly but not randomly.

In Vivo Muscle Contractibility Assay

This assay measures the aggregate torque produced by either the plantar or dorsiflexor muscles of the lower limb

and is carried out using muscle physiology apparatus (Aurora Scientific, ON, Canada). The animal is anesthetized with isoflurane. Once the animal is anesthetized, the hair from the back and the hind limb will be removed as needed with clippers. If hair removal with clippers is insufficient, a thin layer of hair-removal cream (Nair) is applied, and the site thoroughly cleaned with warm water to prevent discomfort. The hindlimb to be measured is attached to the foot plate with adhesive tape. The limb is held rigid in a blunt clamp. Either the tibial or peroneal component of the sciatic nerve will be stimulated with two sterile, disposable 28 gauge monopolar electrodes inserted through the skin, subcutaneously near the nerve. Mouse temperature will be maintained by conductive thermoregulated heating pad (set at 37° C.) or radiant heat source and monitored by temperature probe.

Histopathology

For histological analysis all muscles and organs are embedded in 7% gum tragacanth and flash frozen in liquid nitrogen cooled isopentane. Frozen sections (12 μ m) are collected for immunohistochemistry and western blot analysis.

Western Blot Analysis for Detection of Human CAPN3

CAPN3 protein quantification in mouse muscle tissues is assessed using a Western blotting method. The CAPN3 enzyme is resolved by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and migrates as a 94 kDa band with an autolytic product at approximately 60 kDa using Novocastra's clinical-grade antibody recognizing the N-terminus, NCL-CALP-12A2. Additionally, NCL-CALP-2C4 antibody recognizes this same CAPN3 molecular weight (94 kD), and an additional fragment (30 kD) in skeletal muscle; both antibodies are suitable for protein detection. A semi-quantitative measure of CAPN3 protein expression levels within the calpain-knockout mouse samples following delivery of the therapeutic rAAV vector is performed and compared with untreated controls.

Quantitative Muscle Histology

Cross sections of TA and quad muscles from treated with AAVrh.74.tMCK.CAPN3 versus control uninjected, are stained with hematoxylin and eosin, and photographed using Zeiss Axiovision L4 software (4 random 20 \times images per section per animal). Fiber size diameters are compared between treated and controls.

21

Statistical Analysis

Student's t-test or one-way ANOVA multiple comparison tests are performed where applicable.

Example 7

Toxicology/Biodistribution Studies

Toxicology/biodistribution studies are carried out using the established efficacious dose and one log higher dose. Toxicology studies are done by systemic (tail vein) delivery of rAAV to 6-8 week old CAPN3-KO mice including comparison to normal C57B16 normal mice. Cohorts of 6-10 mice are included and full necropsies are done using GLP-like methods.

22

Mouse GAPDH was used as a reference gene and WT C57BL/6 was used to calibrate the RT-qPCR data.

In addition, quantitative histopathological analysis was carried out after intramuscular administration. As shown in, the diameter of the TA muscle fiber of the treated CAPN3 KO mice was compared to that of the untreated control (ringer lactate injected TA) muscle. Mean fiber size of slow twitch oxidative (STO, dark), fast twitch oxidative (FTO, intermediate) and fast twitch glycolytic (FTG, light) fibers appeared normalized towards WT values in the AAV.h-CAPN3 injected TA muscle. The quantification of the fiber type size is provided in Table 4 and illustrates an increase with treatment.

TABLE 4

	WT (z18-14)		Treated (z18-11)		Untreated (z18-22 L)	
	number	diameter (μm)	number	diameter (μm)	number	diameter (μm)
STO	246	28.06 ± 0.27	142	28.89 ± 0.32	240	25.57 ± 0.27
FTO	63	36.65 ± 0.53	86	36.71 ± 0.58	110	32.19 ± 0.48
FTG	82	42.55 ± 0.53	86	43.68 ± 0.66	128	35.49 ± 0.50
All fiber	391	32.45 ± 0.38	314	35.08 ± 0.45	478	29.75 ± 0.30

Serum collected from blood samples is used for Clinical Chemistries: Alanine aminotransferase, Alkaline Phosphatase, Aspartate aminotransferase, Bilirubin (Total and Direct), Blood Urea nitrogen, Creatinine, Creatine Kinase, Glucose, and Total Protein.

A full necropsy is performed with a thorough and systematic examination and dissection of the animal viscera and carcass. The tissues/organs are collected include gonads, brain, spleen, kidneys, jejunum, colon, pancreas, heart, lung, stomach, liver, inguinal lymph nodes, spinal cord gastrocnemius and quadriceps. Tissues/organs for histopathology studies are collected and fixed in 10% neutral buffered formalin (10% NBF), with the exception of all skeletal muscle specimens which are mounted on blocks with OCT, and flash frozen in liquid nitrogen-cooled methyl-butane for cryosections.

Example 8

In Vivo Biopotency Testing Following Intramuscular Injection

In vivo biopotency testing was carried out following intramuscular (IM) injection of AAVrh.74.tMCK.CAPN3 (1E11 vg) into the tibialis anterior (TA) muscle in CAPN3 KO mice (n=3) as described above in Example 5.

At 4 weeks post-administration, gene delivery was analyzed by reverse transcription quantitative PCR (RT-qPCR) and western blot analyses. For the Western blot analysis, samples corresponding to 50 μg of whole muscle protein extracts were separated on a 3-8% acrylamide, Tris-Acetate SDS gel and transferred to a PVDF membrane. Immunodetection was performed with a monoclonal antibody raised against a synthetic peptide containing AAs 1-19 of the human Calpain 3 sequence (Leica), and muscle specific actin antibody (Leica) as a loading control. FIG. 3A demonstrates that the presence of the 94 kD calpain 3 protein in the TA muscle after intramuscular injection. The RT-qPCR analysis demonstrated relative expression levels of human Calpain 3 gene 4 weeks post-gene transfer return to normalized levels as compared to WT mice (see FIG. 3B).

In summary, the in vivo biopotency testing following IM injection of the vector (1E11 vg) into tibialis anterior (TA) muscle in CAPN3 KO mice (n=2) demonstrated that 4 weeks post-gene delivery 1) RT-qPCR and western blot analyses showed expressions of CAPN3 transcripts and 94 kDa full-length calpain 3 protein and 2) histological analysis showed an increase in the muscle fiber diameter of TA compared to the control (Ringer's lactate injected TA) muscle.

Example 9

In Vivo Biopotency Testing Following Systemic Injection

In vivo biopotency testing was carried out following systemic injection of AAVrh.74.tMCK.CAPN3 (3E12 vg or 6E12 vg) via the tail vein of CAPN3-KO mice. The low dose CAPN3KO cohort (n=5; mice were denoted as Z18-13, Z18-15, Z18-16, Z18-17, Z18-18) received 3E12 vg in 300 μl Ringer's lactate. At 4 weeks post-gene injection, mice were evaluated for running fatigue by the run-to-exhaustion treadmill test and then euthanized for tissue collection. Muscles from upper and lower limbs (TA, gastrocnemius (GAS), quadriceps, triceps), heart, liver spleen, lung, ovaries and testicles were removed, and tissue samples were frozen in isopentane, and cooled in liquid nitrogen.

RT-qPCR CAPN3 expression was evaluated in TA muscles. For the 3E12 vg low dose, CAPN3 mRNA expression levels were low as observed by high CT values, >27. Western blot analysis showed undetectable corresponding protein bands. Even though low expression data was observed in this tissue for the low dose, both functional and histological benefits were demonstrated with the systemic administration of 3E12 vg.

Subsequently, a higher dose (6E12 vg) was systemically administered to investigate whether protein expression could be detected at a higher dose of vector delivery. The high dose cohort (mice denoted as Z18-20, Z18-21, Z18-23 and Z18-24) CAPN3-KO mice received 6E12 vg AAVrh7.4.tMCK.hCAPN3 vector (twice the dose used in

23

the low dose cohort via systemic injection to the tail vein), and were euthanized 4 weeks post-injection. RT-qPCR showed variable levels of CAPN3 expression in the quad, triceps, GAS, TA and cardiac muscle.

To determine relative expression of the CAPN3 mRNA, muscle tissue samples were collected from CAPN3 KO mice treated with tMCK.hCAPN3 vector at the dose of 3E12 vg (low dose cohort 1) and 6E12 vg (high dose cohort 2). Total RNA was isolated from both cohorts and qPCR of CAPN3 vs. mouse GAPDH were assayed along with the previous samples from the cohort that received the vector via IM injection (1E11 vg; see above in Example 8). The relative expression of CAPN3 was determined by the method below:

$$CT = CT_{CAPN3} - CT_{mGAPDH}$$

$$\Delta\Delta CT = \Delta CT - \Delta CT_{Calibrator}^*$$

$$\text{Relative Expression of CAPN3} = 2^{-\Delta\Delta CT}$$

The relative expression of CAPN3 in each tissue and the original CT value were shown in the Table 5 below and in FIG. 5. Table 5 provides data for IM delivery (mice nos. Z18-11 and Z18-12) and for systemic delivery

24

specific variability and lower relative expression as compared to the IM delivery at 1E11 vg (<1% of IM delivery); this was especially true for the 3E12 low dose cohort. Accordingly, the full-length 94 kDa protein was below the limit of detection by Western blot. However, robust gene expression and prominent amounts of full-length Calpain 3 protein were exhibited following systemic injection of 6E12 vg systemic dosage in the high dose cohort.

Example 10

Assessment of Systemic AAVrh74.tMCK.hCAPN3 Gene Delivery

Gene transfer efficiency was assessed by qPCR, calculating vector genome copies within CAPN3 KO mouse tissue samples following systemic delivery of AAVrh74.tMCK.hCAPN3 at 6E12 vg. The vector genome load of the lower and upper extremity skeletal muscles (quad, TA, gastroc, triceps), heart and liver was determined. Genomic DNA was isolated from frozen tissue samples. The qPCR assay was performed on an ABI 7500 (Applied Biosystems) using the following primer set: "5'-CG-

TABLE 5

CAPN3 RT-PCR:									
Mice No.	Tissue	Genotype	Dose of Treatment (DRAPs per Mice)	CT Value					
				CAPN3	mGAPDH	ΔCT	ΔΔCT	2 ^(-ΔΔCT)	
Z18-14*	TA	WT	0	22.437	22.456	15.234	15.274	7.193	0.000
Z18-19	TA	CAPN3	0	35.259	32.705	15.159	15.176	18.814	11.621
Z18-11	TA	KO	1E11	24.338	24.217	15.800	15.835	8.460	1.267
Z18-12	TA			21.030	21.104	14.906	15.058	6.085	-1.108
Z18-13	TA		3E12	32.376	32.430	15.236	15.203	17.183	9.990
Z18-15	TA			27.407	27.443	14.510	14.520	12.910	5.717
Z18-16	TA			28.609	28.333	15.229	15.259	13.227	6.034
Z18-17	TA			28.675	28.670	14.997	15.005	13.671	6.478
Z18-18	TA			27.869	28.128	14.522	14.544	13.466	6.273
Z18-20	Gastroc		6E12	22.271	22.439	15.939	15.974	6.398	-0.795
	Heart			21.996	22.051	15.267	15.315	6.732	-0.461
	Quadriceps			21.008	21.202	15.203	15.407	5.800	-1.393
	TA			23.806	24.173	16.169	16.385	7.713	0.520
	Triceps			24.083	24.361	15.978	16.097	8.185	0.992
Z18-21	Gastroc			25.330	25.221	15.461	15.462	9.814	2.621
	Heart			25.024	24.819	15.032	15.097	9.857	2.664
	Quadriceps			26.278	26.108	15.285	15.370	10.866	3.673
	TA			26.649	26.697	15.017	15.010	11.659	4.466
	Triceps			27.040	27.134	15.321	15.343	11.755	4.562
Z18-23	Gastroc			24.150	24.144	16.225	16.117	7.976	0.783
	Heart			22.799	22.495	14.593	14.502	8.099	0.906
	Quadriceps			24.248	24.076	16.511	16.504	7.655	0.462
	TA			25.554	25.338	16.124	16.054	9.357	2.164
	Triceps			24.396	24.383	15.363	15.277	9.070	1.877
Z18-24	Gastroc			24.444	24.165	18.083	18.036	6.245	-0.948
	Heart			22.769	22.425	15.100	15.077	7.508	0.315
	Quadriceps			22.754	22.521	15.637	15.672	6.983	-0.210
	TA			23.491	23.555	15.979	15.974	7.547	0.354
	Triceps			24.554	24.403	16.370	16.329	8.128	0.935
Z18-22	Gastroc		0	31.878	31.962	15.433	15.527	16.440	9.247
	Heart			31.407	32.964	16.006	15.972	16.197	9.004
	Quadriceps			32.332	33.464	16.528	16.468	16.400	9.207
	TA			35.584	33.917	16.451	16.372	18.339	11.146
	Triceps			33.615	32.786	15.742	15.628	17.516	10.323
Human Tissue	Muscle			23.547	23.539	37.743	38.262		
pAAV.tMCK.hCAPN3	1 pg/uL			16.328	16.413	UD	38.302		

*Calibrator

Overall, the CAPN3 mRNA expression in the CAPN3 KO muscle following systemic delivery had animal- and tissue-

GAGAGCAACTGCATAAG-3' (Forward; SEQ ID NO: 8); "5'-GGCTGATGATGGCTGAATAG-3' (Reverse; SEQ ID

NO: 9). The primer pair exclusively amplifies a product from the 5' region of the hCAPN3 ORF, and region downstream unique to the expression vector, including portions of an intronic element. The final results are reported as mean copy number of AAVrh74 vector per microgram of genomic DNA.

As shown in FIG. 6, the highest vector genome copy number was present in the liver following systemic vector delivery. Vector genome distribution was variable between the muscle groups. Overall the values were higher in the quadriceps and heart tissue compared to other muscles. Experimental variability was also noted; as the case with Mouse no. Z18-21 which showed relatively lower copy numbers in all muscle groups compared to other 3 mice.

Improvement in both functional and histological features were observed in the 3E12 vg systemically treated CAPN3 KO mice, however, only low levels of muscle Calpain 3 expression were detected in total RNA isolates by RT-qPCR and the full-length 94 kDa protein was undetectable by Western blot for the particular muscle tissue (See FIG. 3A). However, robust gene expression and prominent amounts of full-length Calpain 3 protein were exhibited following the 6E12 vg systemic dosage (see FIG. 3B). The data demonstrates that Calpain 3 gene expression returned to normalized levels as compared to WT mice after 4 weeks post-gene transfer of the AAVrh74.tMCK.hCAPN3 particles. Mouse GAPDH was used as a reference gene and WT C57BL/6 to calibrate the RT-qPCR data.

Histopathology

As discussed above, an efficacy trend at 4 weeks post-injection was observed. A significant increase in fiber size was observed in the TA muscle from CAPN3 KO mice following systemic delivery of AAVrh.74.tMCK.hCAPN3 at 4 week-post injection in both cohorts (3E12 and 6E12). As shown in FIG. 7, total fiber diameter was significantly increased in both of the treated cohorts compared to untreated KO counterparts ($p < 0.00001$). Treatment resulted in normalization of fiber size and there was no dose-related difference between the treatment cohorts ($p = 0.78058$). Table 6 provides the muscle fiber sizes in wild type and CAPN3 KO mice following systemic AAV.hCAPN3 gene therapy at 3E12 and 6E12 vg.

TABLE 6

	WT (n = 3)		6 E12 CAPN3 (n = 3)		3E12 CAPN3 (n = 4)		KO (n = 4)	
	number	diameter (μm)	number	diameter (μm)	number	diameter (μm)	number	diameter (μm)
STO	532	30.0 ± 0.6	441	31.7 ± 0.8	464	30.7 ± 1.2	858	27.2 ± 0.8
FTO	278	40.7 ± 1.1	345	38.8 ± 1.1	447	40.8 ± 1.4	364	35.4 ± 1.1
FTG	275	46.1 ± 1.3	226	45.8 ± 1.4	403	44.1 ± 1.6	455	39.7 ± 1.2
All fiber	1085	38.9 ± 1.8	1012	38.8 ± 2.0	1314	38.5 ± 2.4	1677	34.1 ± 1.8

There was no histopathological evidence of cardiac toxicity following systemic injection of AAVrh7.4.tMCK.hCAPN3 vector at 4 weeks in either cohort. There were variable amounts of virus found in the heart tissue, however no protein bands were detected in the heart tissue by Western blot in either cohort.

Functionality Study: Run-to-Exhaustion Test

Mice were accustomed to the treadmill (Columbus Instruments) with a 15-minute run once per day at 10 m/min for 3 days prior to data acquisition for Run-to-Exhaustion test. The protocol used required having mice on a treadmill that is at a 15-degree incline. The treadmill was turned on at a speed at 1m/min and the speed was increased by 1m every minute until the mouse got exhausted. Exhaustion was

determined when the mouse sits on the rest pad for at least 15 seconds. The time, speed and distance to exhaustion were recorded.

FIG. 8A provides data for the run-to-exhaustion test for the low dose cohort, which received 3E12 vg of AAVrh7.4.tMCK.hCAPN3, and the high dose cohort 2, which received 6E12 vg of AAVrh7.4.tMCK.hCAPN3 as assessed 4 weeks after systemic administration. Treated CAPN3 KO mice in both cohorts performed better on the Run-to-Exhaustion test compared to untreated counterparts. There was no apparent dose-related difference in the Run-to-Exhaustion test performance or statistical difference in muscle fiber diameter between the low and high dose cohorts.

Mice from the high dose cohort 2 ($n=16$) were further analyzed 20-24 weeks after administration of 6E12 vg of AAVrh7.4.tMCK.hCAPN3. As shown in FIG. 8B, the treated CAPN3 KO mice continued to perform better on the Run-to-Exhaustion test compared to untreated counterparts ($p < 0.00001$).

Example 11

Assessment of Cardiac Toxicity Following Systemic Injection of AAVrh7.4.tMCK.hCAPN3 Vector

After the mice of the cohorts were euthanized at 4 weeks post injection, serum and organ samples were collected. The low dose cohort 1 CAPN3KO cohort ($n=5$) received 3E12 vg in 300 μl Ringer's lactate of AAVrh.74.tMCK.hCAPN3 vector via tail vein injection. The high dose cohort 2CAPN3-KO mice received 6E12 vg AAVrh7.4.tMCK.hCAPN3 vector via tail vein, and both cohorts were euthanized 4 weeks post-injection. Two sections through the apex of the heart, superficial and deep regions of ventricles were examined. No inflammation, necrosis or regeneration was found in the tissue sections indicating no toxic effects were observed on the heart muscle from the systemic delivery of AAVrh7.4.tMCK.hCAPN3 vector at two different doses at 4 weeks post-injection. Mice nos. Z18-19 and Z18-22 (Ringer's lactate-injected/untreated) served as control KO animal.

FIG. 9 provides H&E stained fresh frozen sections from the heart. Muscle fiber necrosis, regeneration or inflammation was not seen. Even though there were variable amounts of viruses present in the heart tissue, no protein bands were detected by Western blot in either cohort. FIG. 10 provides the Western blot analysis which shows the full-length Calpain 3 protein is below the limit of detection in the heart tissues after the transduction.

Example 12

In Vivo Physiological Analysis

Physiological assessment is carried out after IM or systemic administration of the AAVrh7.4.tMCK.hCAPN3 vec-

tor. During the in vivo physiological assessments, the mice are anesthetized with inhaled isoflurane. Once the animal is anesthetized, the hair from the back and the hind limb is removed as needed with clippers. If hair removal with clippers is insufficient, a thin layer of hair-removal cream is applied. During in vivo physiological force measurements, torque from the hind limb is measured with a non-invasive force foot plate connected to force detecting motor (Aurora Scientific, Canada) following supramaximal stimulations of the sciatic nerve. The hind limb to be measured is attached to the foot plate with adhesive tape. The limb is held rigid in a blunt clamp. Either the tibial or peroneal component of

the sciatic nerve is stimulated with two sterile disposable 28 gauge monopolar electrodes inserted subcutaneously near the nerve. Mouse temperature is maintained by conductive thermoregulated heating plate (set at 37° C.) or radiant heat source and monitored by infrared temperature probe.

While the present disclosure provides specific embodiments, it is understood that variations and modifications will occur to those skilled in the art. Accordingly, only such limitations as appear in the claims should be placed on the invention.

All documents referred to in this application are hereby incorporated by reference in their entirety.

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 9

<210> SEQ ID NO 1
 <211> LENGTH: 7629
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polynucleotide
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <223> OTHER INFORMATION: AAVrh.74.tMCK.CAPN3 recombinant genome
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(128)
 <223> OTHER INFORMATION: AAV2 ITR
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (165)..(884)
 <223> OTHER INFORMATION: tMCK promoter
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (937)..(1069)
 <223> OTHER INFORMATION: chimeric intron
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1101)..(1106)
 <223> OTHER INFORMATION: Kozak Sequence
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1107)..(3572)
 <223> OTHER INFORMATION: CAPN3 polynucleotide
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (3581)..(3780)
 <223> OTHER INFORMATION: poly A signal
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (3850)..(3977)
 <223> OTHER INFORMATION: second AAV2 ITR

<400> SEQUENCE: 1

g'gcgcgctcgc tcgctcactg aggccgcccc ggcaaagccc gggcgctcggg cgacctttgg	60
tcgccccggcc tcagtgcgagc agcgagcgcg cagagaggga gtggccaact ccatcactag	120
gggttcctta atacgactca ctatagggcc gcaccggtaa gcttccacta cgggtctagg	180
ctgcccattgt aaggaggcaa ggccctgggga cacccgagat gcctggttat aattaacccc	240
aacacctgct gccccccccc cccaacacc tgctgcctga gcctgagcgg ttacccccacc	300
cgggtgcctg ggtcttaggc tetgtacacc atggaggaga agctcgctct aaaaaataacc	360
ctgtccctgg tggatccact acgggtctag gctgcccatg taaggaggca aggcctgggg	420
acacccgaga tgctgggta taattaaccc caacacctgc tgccccccc cccccaacac	480
ctgctgcctg agcctgagcg gttacccccc cccggtgcct gggtcttagg ctctgtacac	540
catggaggag aagctcgctc taaaaataac cctgtccctg gtggatccac tacgggtcta	600

-continued

ggctgccc	cat gtaaggag	gc aaggcctggg	gacacccgag	atgcctgggt	ataattaacc	660
ccaacacctg	ctgccccccc	ccccccaaca	cctgctgcct	gagcctgagc	ggttacccca	720
ccccggtgcc	tgggtcttag	gctctgtaca	ccatggagga	gaagctcgct	ctaaaaataa	780
ccctgtccct	ggtggatcct	ccctggggac	agccctcctc	ggctagtcac	accctgtagg	840
ctcctctata	taacccaggg	gcacaggggc	tgcccccg	tcacggtgga	tatccctgca	900
ggtaccacgc	gctgtaattg	aactgggagt	ggacacctgt	ggagagaaag	gcaaagtgga	960
tgtcagtaag	accaataggt	gcctatcaga	aacgcaagag	tcttctctgt	ctcgacaagc	1020
ccagtttcta	ttggtctcct	taaacctgtc	ttgtaacctt	gatacttacg	gagagcaact	1080
gcataagggc	tagcctcgag	gccaccatgc	cgaccgtcat	tagcgcatct	gtggctccaa	1140
ggacagcggc	tgagccccgg	tccccagggc	cagttcctca	cccggcccag	agcaaggcca	1200
ctgaggctgg	gggtggaaac	ccaagtggca	tctattcagc	catcatcagc	cgcaattttc	1260
ctattatcgg	agtgaagag	aagacattcg	agcaacttca	caagaaatgt	ctagaaaaga	1320
aagttcttta	tgtggacctc	gagttcccac	cgatgagac	ctctctcttt	tatagccaga	1380
agttcccat	ccagttcgtc	tggaagagac	ctccggaaat	ttgcgagaat	ccccgattta	1440
tcattgatgg	agccaacaga	actgacatct	gtcaaggaga	gctaggggac	tgctggtttc	1500
tcgcagccat	tgctgctcg	accctgaacc	agcaccttct	tttccgagtc	ataccccatg	1560
atcaaagttt	catcgaaaac	tacgcaggga	tcttccactt	ccagttcttg	cgctatggag	1620
agtggtgga	cgtggttata	gatgactgcc	tgccaacgta	caacaatcaa	ctggttttca	1680
ccaagtccaa	ccaccgcaat	gagttctgga	gtgctctgct	ggagaaggct	tatgctaagc	1740
tccatggttc	ctacgaagct	ctgaaagggt	ggaacaccac	agaggccatg	gaggacttca	1800
caggaggggt	ggcagagttt	tttgagatca	gggatgctcc	tagtgacatg	tacaagatca	1860
tgaagaaagc	catcgagaga	ggctccctca	tgggctgctc	cattgatgat	ggcacgaaca	1920
tgacctatgg	aacctctcct	tctggtctga	acatggggga	gttgattgca	cggatggtaa	1980
ggaatatgga	taactcactg	ctccaggact	cagacctcga	ccccagaggc	tcagatgaaa	2040
gaccgacccg	gacaatcatt	ccggttcagt	atgagacaag	aatggcctgc	gggctgggtca	2100
gaggtcacgc	ctactctgtc	acggggctgg	atgagggtccc	gttcaaagg	gagaaagtga	2160
agctgggtcg	gctcggaat	ccgtggggcc	agggtggagt	gaacggttct	tggagtgata	2220
gatggaagga	ctggagcttt	gtggacaaag	atgagaaggc	ccgtctgcag	caccaggcca	2280
ctgaggatgg	agagttcttg	atgtcctatg	aggatttcat	ctaccatttc	acaaagttag	2340
agatctgcaa	cctcacggcc	gatgctctgc	agtctgacaa	gcttcagacc	tggacagtgt	2400
ctgtgaacga	gggcccgtgg	gtacgggggt	gctctgccgg	aggctgccgc	aacttcccag	2460
atactttctg	gaccaaccct	cagtaccgtc	tgaagctcct	ggaggaggac	gatgacctg	2520
atgactcgga	ggtgatattg	agcttctctg	tggccctgat	gcagaagaac	cggcggaagg	2580
accggaagct	agggggcagt	ctcttcacca	ttggcttcgc	catctacgag	gttcccaaag	2640
agatgcacgg	gaacaagcag	cacctgcaga	aggacttctt	cctgtacaac	gcctccaagg	2700
ccaggagcaa	aacctacatc	aacatgcggg	agggtgtcca	gcgcttcgcg	ctgcctccca	2760
gcgagtacgt	catcgtgccc	tccacctacg	agccccacca	ggagggggaa	ttcatcctcc	2820
gggtcttctc	tgaagagg	aacctctctg	aggaagtga	aaataccatc	tccgtggatc	2880
ggccagtga	aaagaaaaaa	accaagccca	tcattctctg	ttcggacaga	gcaaacagca	2940
acaaggagct	gggtgtggac	caggagtcat	aggagggcaa	aggcaaaaca	agccctgata	3000

-continued

agcaaaagca gtcocacag ccacagcctg gcagctctga tcaggaaagt gaggaacagc	3060
aacaattccg gaacattttc aagcagatag caggagatga catggagatc tgtgcagatg	3120
agctcaagaa ggtccttaac acagtcgtga acaaacacaa ggacctgaag acacacgggt	3180
tcacactgga gtectgcctg agcatgattg cgctcatgga tacagatggc tctggaaagc	3240
tcaacctgca ggagttccac cacctctgga acaagattaa ggctggcag aaaattttca	3300
aacactatga cacagaccag tccggcacca tcaacagcta cgagatgcga aatgcagtca	3360
acgacgcagg attccacctc aacaaccagc tctatgacat cattaccatg cggtagcgag	3420
acaaacacat gaacatcgac ttgacagtt tcatctgctg cttcgttagg ctggagggca	3480
tggtcagagc tttcatgca ttgacaagg atggagatgg tatcatcaag ctcaacgttc	3540
tggagtggct gcagctcacc atgtatgcct gagcgccgcg ggggatccag acatgataag	3600
atacattgat gagtttgac aaaccacaac tagaatgcag tgaaaaaat gctttatttg	3660
taaaatttgt gatgctattg ctttatttgt aaccattata agctgcaata aacaagttaa	3720
caacaacaat tgcattcatt ttatgtttca gggtcagggg gaggtgtggg aggttttttc	3780
ggatcctcta gagtcgacca gagcatggct acgtagataa gtagcatggc gggttaatca	3840
ttaactacaa ggaaccccta gtgatggagt tggccactcc ctctctgcgc gctcgcctgc	3900
tcactgaggc cggcgaccca aaggctcgcc gacgcccggg ctttgccggg gcggcctcag	3960
tgagcgagcg agcgcgcagc tgcattaatg aatcggccaa cgcgcgggga gaggcggttt	4020
gcgtattggg cgctcttcctg cttcctcgct cactgactcg ctgcgctcgg tcgttcggct	4080
gcgcgagcgc gtatcagctc actcaaaggc ggtaatacgg ttatccacag aatcagggga	4140
taacgcagga aagaacatgt gagcaaaagg ccagcaaaag gccaggaacc gtaaaaggc	4200
cgcgttgctg cgcgtttttc ataggctccg ccccccgcag gagcatcaca aaaatcgacg	4260
ctcaagtcag aggtggcgaa acccgacagg actataaaga taccaggcgt tccccctgg	4320
aagctccctc gtgcgctctc ctgttccgac cctgcgcgtt accggatacc tgtccgcctt	4380
tctcccttcg ggaagcgtgg cgctttctca tagctcacgc tgtaggatc tcagttcgggt	4440
gtaggctggt cgctccaagc tgggctgtgt gcacgaaccc cccgttcagc ccgaccgctg	4500
cgccttatcc ggtaactatc gtcttgagtc caaccggta agacacgact tatcgccact	4560
ggcagcagcc actggtaaca ggattagcag agcgaggat gtaggcgggt ctacagagtt	4620
cttgaagtgg tggcctaact acggctacac tagaaggaca gtatttggtg tctgcgctct	4680
gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca acaaaaccac	4740
cgtcggtagc ggtgggtttt ttgtttgcaa gcagcagatt acgcgcagaa aaaaaggatc	4800
tcaagaagat cctttgatct tttctacggg gtctgacgct cagtggaaacg aaaactcacg	4860
ttaagggtatt ttggtcatga gattatcaaa aaggatcttc acctagatcc ttttaaatta	4920
aaaatgaagt tttaaatcaa tctaaagtat atatgagtaa aaatattccg gaattgccag	4980
ctggggcgcc ctctggtaag gttgggaagc cctgcaaaagt aaactggatg gctttcttgc	5040
cgccaaagat ctgatggcgc aggggatcaa gatctgatca agagacagga tgaggatcgt	5100
ttcgcagatg tgaacaagat ggattgcagc caggttctcc ggcgcgttgg gtggagaggc	5160
tattcggcta tgactgggca caacagacaa tcggctgctc tgatgcgcgc gtgttcgggc	5220
tgtagcgcga ggggcgcgcg gttctttttg tcaagaccga cctgtccggt gcctgaatg	5280
aactgcagga cgaggcagcg cggctatcgt ggctggccac gacgggcgtt ccttgcgcag	5340

-continued

ctgtgctcga cgttgctact gaagcgggaa gggactggct gctattgggc gaagtgccgg	5400
ggcaggatct cctgtcatcc caccttgctc ctgccgagaa agtatccatc atggctgatg	5460
caatgcggcg gctgcatacg cttgatccgg ctacctgccc attcgaccac caagcgaaac	5520
atcgcatcga gcgagcacgt actcggatgg aagccggctc tgtcgatcag gatgatctgg	5580
acgaagagca tcaggggctc gcgccagccg aactgttcgc caggctcaag gcgcgcatgc	5640
ccgacggcga ggtatctcgtc gtgacccatg gcgatgcctg cttgccgaat atcatggtgg	5700
aaaaatggccg cttttctgga ttcctcgact gtggccggct ggggtgtggcg gaccgctatc	5760
aggacatagc gttggctacc cgtgatattg ctgaagagct tggcggcgaa tgggctgacc	5820
gcttcctcgt gctttacggt atcgccgctc ccgattcgca gcgcacgccc ttctatcgcc	5880
ttcttgacga gttcttctga accggtaata ttattgaagc atttatcagg gttattgtct	5940
catgagcggg tacatatctt aatgtattta gaaaaataaa caaatagggg ttccgcgcac	6000
atttccccga aaagtgccac ctgacgtcta agaaaccatt attatcatga cattaaccta	6060
taaaaaatagg cgtatcacga ggccctttcg tctcgcgcgt ttcggtgatg acggtgaaaa	6120
cctctgacac atgcagctcc cggagacggg cacagcttgt ctgtaagcgg atgccgggag	6180
cagacaagcc cgtcagggcg cgtcagcggg tgttgggcggg tgcgggggct ggcttaacta	6240
tgcggcatca gagcagattg tactgagagt gcaccatatg cgggtgtgaaa taccgcacag	6300
atgcgtaagg agaaaatacc gcatacaggaa cttccaacat ccaataaatc atacaggcaa	6360
ggcaaagaat tagcaaaatt aagcaataaa gcctcagagc ataaagctaa atcggttgta	6420
ccaaaaacat tatgaccctg taatactttt gcgggagaag cctttatttc aacgcaagga	6480
taaaaatttt tagaaccttc atatatctta aatgcaatgc ctgagtaatg tgtaggtaaa	6540
gattcaaacg ggtgagaaag gccggagaca gtcaaatcac catcaatatg atattcaacc	6600
gttctagctg ataaattcat gccggagagg gtagctattt ttgagaggct tctacaaagg	6660
ctatcaggtc attgcctgag agtctggagc aaacaagaga atcgatgaac ggtaatcgta	6720
aaactagcat gtcaatcata tgtaccccg ttgataatca gaaaagcccc aaaaacaggga	6780
agattgtata agcaaatatt taaattgtaa acgttaatat tttgttaaaa ttcgcgttaa	6840
atttttgtta aatcagctca ttttttaacc aataggccga aatcggcaaa atcccttata	6900
aatcaaaaaga atagaccgag atagggttga gtgttggtcc agtttggaac aagagtcac	6960
tattaaagaa cgtggactcc aacgtcaaag gcgaaaaaac cgtctatcag ggcgatggcc	7020
cactacgtga accatcccc taatcaagtt ttttggggct gaggtgccgt aaatcactaa	7080
atcggaaccc taaaggagc ccccgattta gagcttgacg gggaaagccg gcgaacgtgg	7140
cgagaaagga agggaagaaa gcgaaaggag cgggcgctag ggcgctggca agtgtagcgg	7200
tcacgctcgc cgtaaccacc acaccgcgcg cgcttaatgc gccgctacag ggcgcgtact	7260
atggttgctt tgacgagcac gtataacgtg ctttctcgt tagaatcaga gcgggagcta	7320
aacaggaggc cgattaaagg gatcttagac aggaacggta cgccagaatc ctgagaagtg	7380
tttttataat cagtgaggcc accgagtaaa agagtctgtc catcacgcaa attaacgctt	7440
gtcgcaatac ttctttgatt agtaataaca tcactgcct gagtagaaga actcaaaacta	7500
tcggccttgc tggtaatatc cagaacaata ttaccgccag ccattgcaac ggaatcgcca	7560
ttcgccattc aggctgcgca actggtggga agggcgatcg gtgcgggctt cttcgctatt	7620
acgccagct	7629

-continued

```

<210> SEQ ID NO 2
<211> LENGTH: 2466
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: CAPN3 polynucleotide

<400> SEQUENCE: 2

atgccgaccg tcattagcgc atctgtggct ccaaggacag cggtgagcc ccggtcccca    60
gggccagttc ctcacccggc ccagagcaag gccactgagg ctgggggtgg aaaccaagt    120
ggcatctatt cagccatcat cagccgcaat ttctctatta tcggagtga agagaagaca    180
ttcgagcaac ttcacaagaa atgtctagaa aagaaagttc tttatgtgga ccctgagttc    240
ccaccgatg agacctctct cttttatagc cagaagttcc ccatccagtt cgtctggaag    300
agacctccgg aaattttgga gaatccccga tttatcattg atggagccaa cagaactgac    360
atctgtcaag gagagctagg ggactgctgg tttctcgag ccattgcctg cctgacctg    420
aaccagcacc ttcttttccg agtcataccc catgatcaaa gtttcacga aaactacgca    480
gggatcttcc acttcagtt ctggcgctat ggagagtggg tggacgtggt tatagatgac    540
tgctgccaa cgtacaacaa tcaactggtt ttcaccaagt ccaaccaccg caatgagttc    600
tggagtgtct tgctggagaa ggcttatgct aagctccatg gttcctacga agctctgaaa    660
ggtgggaaca ccacagaggc catggaggac ttcacaggag ggggtggcaga gttttttgag    720
atcagggatg ctctagtga catgtacaag atcatgaaga aagccatcga gagaggtccc    780
ctcatgggct gctccattga tgatggcacg aacatgacct atggaacctc tccttctggt    840
ctgaacatgg gggagtgtat tgcacggatg gtaaggaata tggataactc actgctccag    900
gactcagacc tcgaccccag aggtcagat gaaagaccga cccggacaat cattccggtt    960
cagtatgaga caagaatggc ctgcgggctg gtcagaggtc acgcctactc tgtcacgggg    1020
ctggatgagg tcccgttcaa aggtgagaaa gtgaagctgg tgcggctgcg gaatccgtgg    1080
ggccaggtgg agtggaaagg ttcttggagt gatagatgga aggactggag ctttgtggac    1140
aaagatgaga agggccgtct gcagcaccag gtcactgagg atggagagtt ctggatgtcc    1200
tatgaggatt tcctctacca ttccacaaag ttggagatct gcaacctcac ggccgatgct    1260
ctgcagtctg acaagcttca gacctggaca gtgtctgtga acgagggccg ctgggtacgg    1320
ggttgctctg ccggaggctg ccgcaacttc ccagatactt tctggacca cctcagtac    1380
cgtctgaagc tcctggagga ggaagatgac cctgatgact cggaggtgat ttgcagcttc    1440
ctggtggccc tgatgcagaa gaaccggcgg aaggaccgga agctaggggc cagtctcttc    1500
accattggct tcgccatcta cgaggttccc aaagagatgc acgggaacaa gcagcacctg    1560
cagaaggact tcttctgta caacgcctcc aaggccagga gcaaaccta catcaacatg    1620
cgggaggtgt cccagcgctt ccgctgcct cccagcgagt acgtcactgt gccctccacc    1680
tacgagcccc accaggaggg ggaattcctc ctccgggtct tctctgaaaa gaggaacctc    1740
tctgaggaag ttgaaaatac catctccgtg gatcggccag tgaaaaagaa aaaaaccaag    1800
cccatcatct tcgtttcgga cagagcaaac agcaacaagg agctgggtgt ggaccaggag    1860
tcagaggagg gcaaaggcaa aacaagccct gataagcaaa agcagtcacc acagccacag    1920
cctggcagct ctgatcagga aagtgaggaa cagcaacaat tccggaacat tttcaagcag    1980
atagcaggag atgacatgga gatctgtgca gatgagctca agaaggtcct taacacagtc    2040

```

-continued

```

gtgaacaaac acaaggacct gaagacacac ggggtcacac tggagtcctg ccgtagcatg 2100
attgcgctca tggatacaga tggtcttga aagctcaacc tgcaggagtt ccaccacctc 2160
tggacaaga ttaaggcctg gcagaaaatt ttcaaacact atgacacaga ccagtcgggc 2220
accatcaaca gctacgagat gcgaaatgca gtcaacgacg caggattcca cctcaacaac 2280
cagctctatg acatcattac catgcggtac gcagacaaac acatgaacat cgactttgac 2340
agtttcatct gctgcttcgt taggttgag ggcatgttca gagcttttca tgcatttgac 2400
aaggatggag atggtatcat caagctcaac gttctggagt ggctgcagct caccatgtat 2460
gcctga 2466

```

```

<210> SEQ ID NO 3
<211> LENGTH: 720
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: TMCK promoter

```

```

<400> SEQUENCE: 3

```

```

ccactacggg tctaggctgc ccatgtaagg aggcaaggcc tggggacacc cgagatgcct 60
ggttataatt aaccccaaca cctgctgccc ccccccccc aacacctgct gcctgagcct 120
gagcggttac cccaccocgg tgcttgggtc ttaggtcttg tacaccatgg aggagaagct 180
cgctctaaaa ataacctgt ccttggtgga tccactacgg gtctaggctg cccatgtaag 240
gaggcaaggc ctggggacac ccgagatgcc tggttataat taacccaac acctgctgcc 300
cccccccc caacacctgc tgctgagcc tgagcggtta cccaccocg gtgctgggt 360
cttaggtct gtacaccatg gaggagaagc tcgctctaaa aataacctg tccctggtgg 420
atccactacg ggtctaggct gcccatgtaa ggaggcaagg cctggggaca cccgagatgc 480
ctggttataa ttaaccccaa cacctgctgc ccccccccc ccaacacctg ctgctgagc 540
ctgagcggtt accccacccc ggtgctggg tcttaggtc tgtacaccat ggaggagaag 600
ctcgctctaa aaataacct gtccctgggt gatecctcct ggggacagcc cctcctggct 660
agtcacaccc tgtaggtctc tctatataac ccaggggcac aggggctgcc ccgggtcac 720

```

```

<210> SEQ ID NO 4
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: MCK forward primer

```

```

<400> SEQUENCE: 4

```

```

cccgagatgc ctggttataa tt 22

```

```

<210> SEQ ID NO 5
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polynucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: MCK reverse primer

```

-continued

<400> SEQUENCE: 5

gctcaggcag caggtgttg

19

<210> SEQ ID NO 6

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polynucleotide

<220> FEATURE:

<221> NAME/KEY: misc_feature

<223> OTHER INFORMATION: MCK probe

<400> SEQUENCE: 6

ccagacatgt ggctgctccc cc

22

<210> SEQ ID NO 7

<211> LENGTH: 821

<212> TYPE: PRT

<213> ORGANISM: Homo Sapiens

<400> SEQUENCE: 7

Met Pro Thr Val Ile Ser Ala Ser Val Ala Pro Arg Thr Ala Ala Glu
1 5 10 15Pro Arg Ser Pro Gly Pro Val Pro His Pro Ala Gln Ser Lys Ala Thr
20 25 30Glu Ala Gly Gly Gly Asn Pro Ser Gly Ile Tyr Ser Ala Ile Ile Ser
35 40 45Arg Asn Phe Pro Ile Ile Gly Val Lys Glu Lys Thr Phe Glu Gln Leu
50 55 60His Lys Lys Cys Leu Glu Lys Lys Val Leu Tyr Val Asp Pro Glu Phe
65 70 75 80Pro Pro Asp Glu Thr Ser Leu Phe Tyr Ser Gln Lys Phe Pro Ile Gln
85 90 95Phe Val Trp Lys Arg Pro Pro Glu Ile Cys Glu Asn Pro Arg Phe Ile
100 105 110Ile Asp Gly Ala Asn Arg Thr Asp Ile Cys Gln Gly Glu Leu Gly Asp
115 120 125Cys Trp Phe Leu Ala Ala Ile Ala Cys Leu Thr Leu Asn Gln His Leu
130 135 140Leu Phe Arg Val Ile Pro His Asp Gln Ser Phe Ile Glu Asn Tyr Ala
145 150 155 160Gly Ile Phe His Phe Gln Phe Trp Arg Tyr Gly Glu Trp Val Asp Val
165 170 175Val Ile Asp Asp Cys Leu Pro Thr Tyr Asn Asn Gln Leu Val Phe Thr
180 185 190Lys Ser Asn His Arg Asn Glu Phe Trp Ser Ala Leu Leu Glu Lys Ala
195 200 205Tyr Ala Lys Leu His Gly Ser Tyr Glu Ala Leu Lys Gly Gly Asn Thr
210 215 220Thr Glu Ala Met Glu Asp Phe Thr Gly Gly Val Ala Glu Phe Phe Glu
225 230 235 240Ile Arg Asp Ala Pro Ser Asp Met Tyr Lys Ile Met Lys Lys Ala Ile
245 250 255Glu Arg Gly Ser Leu Met Gly Cys Ser Ile Asp Asp Gly Thr Asn Met
260 265 270

Thr	Tyr	Gly	Thr	Ser	Pro	Ser	Gly	Leu	Asn	Met	Gly	Glu	Leu	Ile	Ala
275						280						285			
Arg	Met	Val	Arg	Asn	Met	Asp	Asn	Ser	Leu	Leu	Gln	Asp	Ser	Asp	Leu
290						295				300					
Asp	Pro	Arg	Gly	Ser	Asp	Glu	Arg	Pro	Thr	Arg	Thr	Ile	Ile	Pro	Val
305				310						315				320	
Gln	Tyr	Glu	Thr	Arg	Met	Ala	Cys	Gly	Leu	Val	Arg	Gly	His	Ala	Tyr
				325				330						335	
Ser	Val	Thr	Gly	Leu	Asp	Glu	Val	Pro	Phe	Lys	Gly	Glu	Lys	Val	Lys
		340						345				350			
Leu	Val	Arg	Leu	Arg	Asn	Pro	Trp	Gly	Gln	Val	Glu	Trp	Asn	Gly	Ser
		355				360						365			
Trp	Ser	Asp	Arg	Trp	Lys	Asp	Trp	Ser	Phe	Val	Asp	Lys	Asp	Glu	Lys
370						375				380					
Ala	Arg	Leu	Gln	His	Gln	Val	Thr	Glu	Asp	Gly	Glu	Phe	Trp	Met	Ser
385				390						395				400	
Tyr	Glu	Asp	Phe	Ile	Tyr	His	Phe	Thr	Lys	Leu	Glu	Ile	Cys	Asn	Leu
				405				410						415	
Thr	Ala	Asp	Ala	Leu	Gln	Ser	Asp	Lys	Leu	Gln	Thr	Trp	Thr	Val	Ser
		420						425				430			
Val	Asn	Glu	Gly	Arg	Trp	Val	Arg	Gly	Cys	Ser	Ala	Gly	Gly	Cys	Arg
		435				440						445			
Asn	Phe	Pro	Asp	Thr	Phe	Trp	Thr	Asn	Pro	Gln	Tyr	Arg	Leu	Lys	Leu
450						455				460					
Leu	Glu	Glu	Asp	Asp	Asp	Pro	Asp	Asp	Ser	Glu	Val	Ile	Cys	Ser	Phe
465				470						475				480	
Leu	Val	Ala	Leu	Met	Gln	Lys	Asn	Arg	Arg	Lys	Asp	Arg	Lys	Leu	Gly
				485				490						495	
Ala	Ser	Leu	Phe	Thr	Ile	Gly	Phe	Ala	Ile	Tyr	Glu	Val	Pro	Lys	Glu
		500						505				510			
Met	His	Gly	Asn	Lys	Gln	His	Leu	Gln	Lys	Asp	Phe	Phe	Leu	Tyr	Asn
515						520						525			
Ala	Ser	Lys	Ala	Arg	Ser	Lys	Thr	Tyr	Ile	Asn	Met	Arg	Glu	Val	Ser
530						535				540					
Gln	Arg	Phe	Arg	Leu	Pro	Pro	Ser	Glu	Tyr	Val	Ile	Val	Pro	Ser	Thr
545				550						555				560	
Tyr	Glu	Pro	His	Gln	Glu	Gly	Glu	Phe	Ile	Leu	Arg	Val	Phe	Ser	Glu
				565				570						575	
Lys	Arg	Asn	Leu	Ser	Glu	Glu	Val	Glu	Asn	Thr	Ile	Ser	Val	Asp	Arg
		580						585				590			
Pro	Val	Lys	Lys	Lys	Lys	Thr	Lys	Pro	Ile	Ile	Phe	Val	Ser	Asp	Arg
595						600				605					
Ala	Asn	Ser	Asn	Lys	Glu	Leu	Gly	Val	Asp	Gln	Glu	Ser	Glu	Glu	Gly
610						615				620					
Lys	Gly	Lys	Thr	Ser	Pro	Asp	Lys	Gln	Lys	Gln	Ser	Pro	Gln	Pro	Gln
625				630						635				640	
Pro	Gly	Ser	Ser												

-continued

690	695	700
Asp Thr Asp Gly Ser Gly Lys Leu Asn Leu Gln Glu Phe His His Leu		
705	710	715 720
Trp Asn Lys Ile Lys Ala Trp Gln Lys Ile Phe Lys His Tyr Asp Thr		
	725	730 735
Asp Gln Ser Gly Thr Ile Asn Ser Tyr Glu Met Arg Asn Ala Val Asn		
	740	745 750
Asp Ala Gly Phe His Leu Asn Asn Gln Leu Tyr Asp Ile Ile Thr Met		
	755	760 765
Arg Tyr Ala Asp Lys His Met Asn Ile Asp Phe Asp Ser Phe Ile Cys		
	770	775 780
Cys Phe Val Arg Leu Glu Gly Met Phe Arg Ala Phe His Ala Phe Asp		
	785	790 795 800
Lys Asp Gly Asp Gly Ile Ile Lys Leu Asn Val Leu Glu Trp Leu Gln		
	805	810 815
Leu Thr Met Tyr Ala		
	820	
<210> SEQ ID NO 8		
<211> LENGTH: 19		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic Primer		
<400> SEQUENCE: 8		
cggagagcaa ctgcataag		19
<210> SEQ ID NO 9		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic Primer		
<400> SEQUENCE: 9		
ggctgatgat ggctgaatag		20

I claim:

1. A recombinant adeno-associated virus (rAAV) comprising a polynucleotide which comprises nucleotides 1 to 3977 of SEQ ID NO: 1, wherein the rAAV comprises an AAV-8, an AAV-9, or an AAV rh.74 capsid protein, or a variant of each thereof.

2. The rAAV of claim 1, wherein the rAAV comprises an rh.74 capsid protein.

3. A composition comprising the rAAV of claim 1.

4. A composition comprising the rAAV of claim 2.

* * * * *