



US012391940B2

(12) **United States Patent**  
Song et al.

(10) **Patent No.:** US 12,391,940 B2  
(45) **Date of Patent:** Aug. 19, 2025

(54) **SINGLE CELL ASSAY FOR TRANSPOSEASE-ACCESSIBLE CHROMATIN**(71) Applicant: **Becton, Dickinson and Company**, Franklin Lakes, NJ (US)(72) Inventors: **Hye-Won Song**, San Jose, CA (US); **Margaret Nakamoto**, San Jose, CA (US)(73) Assignee: **Becton, Dickinson and Company**, Franklin Lakes, NJ (US)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 127 days.

(21) Appl. No.: **17/390,640**(22) Filed: **Jul. 30, 2021**(65) **Prior Publication Data**

US 2022/0033810 A1 Feb. 3, 2022

**Related U.S. Application Data**

(60) Provisional application No. 63/059,715, filed on Jul. 31, 2020.

(51) **Int. Cl.****C12N 15/10** (2006.01)**C12N 15/11** (2006.01)**C12Q 1/6806** (2018.01)(52) **U.S. Cl.**CPC ..... **C12N 15/1093** (2013.01); **C12N 15/11** (2013.01); **C12Q 1/6806** (2013.01)(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

## U.S. PATENT DOCUMENTS

4,510,244 A	4/1985	Parks et al.
4,725,536 A	2/1988	Fritsch et al.
5,124,246 A	6/1992	Urdea et al.
5,137,809 A	8/1992	Loken et al.
5,149,625 A	9/1992	Church et al.
5,200,314 A	4/1993	Urdea
5,308,990 A	5/1994	Takahashi et al.
5,424,186 A	6/1995	Fodor et al.
5,424,413 A	6/1995	Hogan et al.
5,445,934 A	8/1995	Fodor et al.
5,470,570 A	11/1995	Taylor et al.
5,604,097 A	2/1997	Brenner
5,635,352 A	6/1997	Urdea et al.
5,635,400 A	6/1997	Brenner
5,648,245 A	7/1997	Fire et al.
5,654,413 A	8/1997	Brenner
5,656,731 A	8/1997	Urdea
5,658,737 A	8/1997	Nelson et al.
5,714,330 A	2/1998	Brenner et al.
5,744,305 A	4/1998	Fodor et al.
5,759,778 A	6/1998	Li et al.
5,763,175 A	6/1998	Brenner
5,800,992 A	9/1998	Fodor et al.
5,830,712 A	11/1998	Rampersad et al.

5,846,719 A	12/1998	Brenner et al.
5,854,033 A	12/1998	Lizardi
5,871,928 A	2/1999	Fodor et al.
5,925,525 A	7/1999	Fodor et al.
5,935,793 A	8/1999	Wong
5,962,271 A	10/1999	Chenchik et al.
5,962,272 A	10/1999	Chenchik et al.
5,968,740 A	10/1999	Fodor et al.
5,981,176 A	11/1999	Wallace
5,981,179 A	11/1999	Lorinez et al.
6,013,445 A	1/2000	Albrecht et al.
6,040,138 A	3/2000	Lockhart et al.
6,046,005 A	4/2000	Ju et al.
6,060,596 A	5/2000	Lerner et al.
6,064,755 A	5/2000	Some
6,114,149 A	9/2000	Fry et al.

(Continued)

## FOREIGN PATENT DOCUMENTS

CA	2474509 A1	2/2003
CA	2961210 A1	3/2016

(Continued)

## OTHER PUBLICATIONS

10X Genomics, Inc., 2019, User Guide: Visium Spatial Gene Expression Reagent Kits, 10xGenomics.com, 76 pp.

2018 Top 10 Innovations, The Scientist Magazine® (2018). Available at: <https://thescientist.com/features/2018-top-10-innovations-65140>, 16 pp.

Achim et al., "High-throughput spatial mapping of single-cell RNA-seq data to tissue of origin," Nature Biotechnology 2015, 33(5), 503-511.

Adey et al., "Rapid, low-input, low-bias construction of shotgun fragment libraries by high-density in vitro transposition," Genome Biology 2010, 11(R19), in 17 pages.

Advisory Action dated Nov. 29, 2019 in U.S. Appl. No. 15/084,307.

Advisory Action dated Dec. 2, 2019 in U.S. Appl. No. 15/055,407.

Advisory Action dated Aug. 25, 2020 in U.S. Appl. No. 15/084,307.

(Continued)

Primary Examiner — Nancy J Leith

Assistant Examiner — Jessica D Parisi

(74) Attorney, Agent, or Firm — Dickinson Wright PLLC

## ABSTRACT

Disclosed herein include systems, methods, compositions, and kits for labeling DNA (e.g., open chromatin-associated gDNA). The method can comprise contacting double-stranded DNA (dsDNA), such as gDNA, with a transposome to generate a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang. The transposome can comprise a transposase, a first adaptor having a first 5' overhang, and a second adaptor having a second 5' overhang. The first 5' overhang can comprise a complement of a target-binding region of a bead oligonucleotide. The first 5' overhang can comprise a coupling sequence. The second 5' overhang can comprise a universal sequence. There are provided, in some embodiments, coupling oligonucleotides comprising a 5' complement of the coupling sequence and a 3' complement of a target-binding region of a bead oligonucleotide.

# US 12,391,940 B2

Page 2

---

(56)

## References Cited

### U.S. PATENT DOCUMENTS

6,117,631 A	9/2000	Nilsen	8,114,681 B2	2/2012	Martin et al.
6,124,092 A	9/2000	O'neill et al.	8,148,068 B2	4/2012	Brenner
6,138,077 A	10/2000	Brenner	8,168,385 B2	5/2012	Brenner
6,140,489 A	10/2000	Brenner	8,206,913 B1	6/2012	Kamberov et al.
6,172,214 B1	1/2001	Brenner	8,241,850 B2	8/2012	Brenner
6,197,506 B1	3/2001	Fodor et al.	8,298,767 B2	10/2012	Brenner et al.
6,197,554 B1	3/2001	Lin et al.	8,318,433 B2	11/2012	Brenner
6,214,558 B1	4/2001	Shuber et al.	8,367,051 B2	2/2013	Matyjaszewski et al.
6,235,475 B1	5/2001	Brenner et al.	8,420,324 B2	4/2013	Rashtchian et al.
6,235,483 B1	5/2001	Wolber et al.	8,445,205 B2	5/2013	Brenner
6,265,163 B1	7/2001	Albrecht et al.	8,470,996 B2	6/2013	Brenner
6,268,152 B1	7/2001	Fodor et al.	8,476,018 B2	7/2013	Brenner
6,284,460 B1	9/2001	Fodor et al.	8,481,292 B2	7/2013	Casbon et al.
6,284,485 B1	9/2001	Boyle et al.	8,535,889 B2	9/2013	Larson et al.
6,309,822 B1	10/2001	Fodor et al.	8,563,274 B2	10/2013	Brenner et al.
6,309,823 B1	10/2001	Cronin et al.	8,603,749 B2	12/2013	Gillevet et al.
6,326,148 B1	12/2001	Pauletti et al.	8,679,756 B1	3/2014	Brenner et al.
6,355,431 B1	3/2002	Chee et al.	8,685,678 B2	4/2014	Casbon et al.
6,355,432 B1	3/2002	Fodor et al.	8,685,753 B2	4/2014	Martin et al.
6,372,813 B1	4/2002	Johnson et al.	8,715,967 B2	5/2014	Casbon et al.
6,395,491 B1	5/2002	Fodor et al.	8,722,368 B2	5/2014	Casbon et al.
6,406,848 B1	6/2002	Bridgman et al.	8,728,766 B2	5/2014	Casbon et al.
6,436,675 B1	8/2002	Welch et al.	8,741,606 B2	6/2014	Casbon et al.
6,440,667 B1	8/2002	Fodor et al.	8,835,110 B2	9/2014	Wang et al.
6,440,706 B1	8/2002	Vogelstein et al.	8,841,071 B2	9/2014	Link
6,451,536 B1	9/2002	Fodor et al.	8,856,410 B2	10/2014	Park
6,458,530 B1	10/2002	Morris et al.	8,865,470 B2	10/2014	Yan et al.
6,468,744 B1	10/2002	Cronin et al.	9,150,852 B2	10/2015	Samuels et al.
6,480,791 B1	11/2002	Strathmann	9,181,582 B2	11/2015	Kurn
6,489,114 B2	12/2002	Laayoun et al.	9,188,586 B2	11/2015	Fan et al.
6,489,116 B2	12/2002	Wagner	9,228,229 B2	1/2016	Olson et al.
6,492,121 B2	12/2002	Kurane et al.	9,262,376 B2	2/2016	Tsuto
6,500,620 B2	12/2002	Yu et al.	9,297,047 B2	3/2016	Furchak et al.
6,512,105 B1	1/2003	Hogan et al.	9,371,598 B2	6/2016	Chee
6,514,699 B1	2/2003	O'neill et al.	9,582,877 B2	2/2017	Fu et al.
6,544,739 B1	4/2003	Fodor et al.	9,593,365 B2	3/2017	Frisen et al.
6,551,784 B2	4/2003	Fodor et al.	9,644,204 B2	5/2017	Hindson et al.
6,576,424 B2	6/2003	Fodor et al.	9,677,131 B2	6/2017	Fredriksson et al.
6,600,996 B2	7/2003	Webster et al.	9,689,024 B2	6/2017	Hindson et al.
6,629,040 B1	9/2003	Goodlett et al.	9,695,468 B2	7/2017	Hindson et al.
6,653,077 B1	11/2003	Brenner	9,787,810 B1	10/2017	Chiang
6,753,147 B2	6/2004	Vögelstein et al.	9,850,515 B2	12/2017	McCoy et al.
6,787,308 B2	9/2004	Balasubramanian et al.	9,856,530 B2	1/2018	Hindson et al.
6,808,906 B2	10/2004	Shen et al.	9,868,979 B2	1/2018	Chee et al.
6,849,404 B2	2/2005	Park et al.	9,879,313 B2	1/2018	Chee et al.
6,852,488 B2	2/2005	Fodor et al.	9,905,005 B2	2/2018	Fu et al.
6,858,412 B2	2/2005	Willis et al.	9,938,523 B2	4/2018	LaBaer
6,890,741 B2	5/2005	Fan et al.	9,951,386 B2	4/2018	Hindson et al.
6,946,251 B2	9/2005	Kurn	10,002,316 B2	6/2018	Fodor et al.
6,974,669 B2	12/2005	Mirkin et al.	10,011,872 B1	7/2018	Belgrader et al.
7,022,479 B2	4/2006	Wagner	10,017,761 B2	7/2018	Weissman et al.
7,034,145 B2	4/2006	Shen et al.	10,023,910 B2	7/2018	Drmanac et al.
7,155,050 B1	12/2006	Slope	10,030,267 B2	7/2018	Hindson et al.
7,294,466 B2	11/2007	McMillan	10,041,116 B2	8/2018	Hindson et al.
7,323,309 B2	1/2008	Mirkin et al.	10,131,958 B1	11/2018	Fan et al.
7,393,665 B2	7/2008	Brenner	10,138,518 B2	11/2018	Chun
7,407,757 B2	8/2008	Brenner	10,151,003 B2	12/2018	Fan et al.
7,424,368 B2	9/2008	Huang et al.	10,208,343 B2	2/2019	Hindson et al.
7,432,055 B2	10/2008	Pemov et al.	10,227,648 B2	3/2019	Hindson et al.
7,470,515 B2	12/2008	Rashtchian et al.	10,246,703 B2	4/2019	Church et al.
7,473,767 B2	1/2009	Dimitrov	10,253,364 B2	4/2019	Hindson et al.
7,476,786 B2	1/2009	Chan et al.	10,266,874 B2	4/2019	Weissleder et al.
7,537,897 B2	5/2009	Brenner et al.	10,273,541 B2	4/2019	Hindson et al.
7,544,473 B2	6/2009	Brenner	10,288,608 B2	5/2019	Kozlov et al.
7,635,566 B2	12/2009	Brenner	10,294,511 B2	5/2019	Sanches-Kuiper et al.
7,638,612 B2	12/2009	Rashtchian et al.	10,308,982 B2	6/2019	Chee
7,718,403 B2	5/2010	Kamberov et al.	10,323,278 B2	6/2019	Belgrader et al.
7,771,946 B2	8/2010	Kurn	10,337,061 B2	7/2019	Hindson et al.
7,822,555 B2	10/2010	Huang et al.	10,344,329 B2	7/2019	Hindson et al.
7,824,856 B2	11/2010	Monforte	10,450,607 B2	10/2019	Hindson et al.
7,824,889 B2	11/2010	Vogelstein et al.	10,550,429 B2	2/2020	Harada et al.
7,915,015 B2	3/2011	Vogelstein et al.	RE47,983 E	5/2020	Gao et al.
7,985,546 B2	7/2011	Church et al.	11,092,607 B2	8/2021	Gaublomme et al.
8,071,311 B2	12/2011	Kurn	11,460,468 B2	10/2022	Fan et al.
8,110,351 B2	2/2012	Bosnes	11,467,157 B2	10/2022	Fan et al.
			11,535,882 B2	12/2022	Fu et al.
			2001/0024784 A1	9/2001	Wagner
			2001/0036632 A1	11/2001	Yu et al.

(56)	References Cited				
U.S. PATENT DOCUMENTS					
2002/0019005 A1	2/2002	Kamb	2008/0070303 A1	3/2008	West et al.
2002/0051986 A1	5/2002	Baez et al.	2008/0119736 A1	5/2008	Dentinger
2002/0065609 A1	5/2002	Ashby	2008/0194414 A1	8/2008	Albert et al.
2002/0072058 A1	6/2002	Voelker et al.	2008/0261204 A1	10/2008	Lexow
2002/0094116 A1	7/2002	Forst et al.	2008/0268508 A1	10/2008	Sowlay
2002/0106666 A1	8/2002	Hayashizaki	2008/0269068 A1	10/2008	Church et al.
2002/0132241 A1	9/2002	Fan et al.	2008/0274458 A1	11/2008	Latham et al.
2002/0168665 A1	11/2002	Okawa	2008/0299609 A1	12/2008	Kwon et al.
2002/0172953 A1	11/2002	Mirkin et al.	2008/0318802 A1	12/2008	Brenner
2002/0187480 A1	12/2002	Brandon	2009/0053669 A1	2/2009	Liu et al.
2002/0192687 A1	12/2002	Mirkin et al.	2009/0061513 A1	3/2009	Andersson Svahn et al.
2003/0003490 A1	1/2003	Fan et al.	2009/0105959 A1	4/2009	Braverman et al.
2003/0013091 A1	1/2003	Dimitrov	2009/0131269 A1	5/2009	Marfin et al.
2003/0032049 A1	2/2003	Wagner	2009/0137407 A1	5/2009	Church et al.
2003/0049616 A1	3/2003	Brenner et al.	2009/0208936 A1	8/2009	Tan et al.
2003/0077611 A1	4/2003	Slepnev	2009/0220385 A1	9/2009	Brown et al.
2003/0082818 A1	5/2003	Bahnsen et al.	2009/0226891 A2	9/2009	Nova et al.
2003/0087242 A1	5/2003	Mirkin et al.	2009/0252414 A1	10/2009	Suzuki
2003/0104436 A1	6/2003	Morris et al.	2009/0253586 A1	10/2009	Nelson et al.
2003/0165935 A1	9/2003	Vann et al.	2009/0283766 A1	11/2009	Skoglund
2003/0175908 A1	9/2003	Linnarsson	2009/0290151 A1	11/2009	Agrawal et al.
2003/0186251 A1	10/2003	Dunn et al.	2009/0298709 A1	12/2009	Ma
2003/0207296 A1	11/2003	Park et al.	2009/0311694 A1	12/2009	Gallagher et al.
2003/0207300 A1	11/2003	Matray et al.	2010/0069250 A1	3/2010	White, III
2004/0047769 A1	3/2004	Tanaami	2010/0105049 A1	4/2010	Ehrich et al.
2004/0091864 A1	5/2004	French et al.	2010/0105112 A1	4/2010	Holtze et al.
2004/0096368 A1	5/2004	Davis et al.	2010/0105886 A1	4/2010	Woudenberg et al.
2004/0096892 A1	5/2004	Wang et al.	2010/0120098 A1	5/2010	Grunenwald et al.
2004/0121342 A1	6/2004	McKeown	2010/0120630 A1	5/2010	Huang et al.
2004/0146901 A1	7/2004	Morris et al.	2010/0136544 A1	6/2010	Agresti et al.
2004/0147435 A1	7/2004	Hawiger et al.	2010/0159533 A1	6/2010	Lipson et al.
2004/0157243 A1	8/2004	Huang et al.	2010/0167354 A1	7/2010	Kurn
2004/0209298 A1	10/2004	Kamberov et al.	2010/0184076 A1	7/2010	Lawton
2004/0224325 A1	11/2004	Knapp et al.	2010/0255471 A1	10/2010	Clarke
2004/0259118 A1	12/2004	Macevitz	2010/0267028 A1	10/2010	Pasche
2005/0019776 A1	1/2005	Callow et al.	2010/0291666 A1	11/2010	Collier et al.
2005/0032110 A1	2/2005	Shen et al.	2010/0300895 A1	12/2010	Nobile et al.
2005/0048500 A1	3/2005	Lawton	2010/0323348 A1	12/2010	Hamady et al.
2005/0053952 A1	3/2005	Hong et al.	2010/0330574 A1	12/2010	Whitman
2005/0105077 A1	5/2005	Padmanabhan et al.	2011/0038507 A1	2/2011	Hager
2005/0170373 A1	8/2005	Monforte	2011/0059436 A1	3/2011	Hardin et al.
2005/0175993 A1	8/2005	Wei	2011/0059556 A1	3/2011	Strey
2005/0196760 A1	9/2005	Pemov et al.	2011/0070584 A1	3/2011	Wohlgemuth et al.
2005/0214825 A1	9/2005	Stuelpnagel	2011/0072889 A1	3/2011	Albitar et al.
2005/0250146 A1	11/2005	McMillan	2011/0160078 A1	6/2011	Fodor et al.
2005/0250147 A1	11/2005	Macevitz	2011/0201507 A1	8/2011	Rava et al.
2006/0002824 A1	1/2006	Chang et al.	2011/0230358 A1	9/2011	Rava
2006/0035258 A1	2/2006	Tadakamalla et al.	2011/0244455 A1	10/2011	Larson et al.
2006/0040297 A1	2/2006	Leamon et al.	2011/0245111 A1	10/2011	Chee
2006/0041385 A1	2/2006	Bauer	2011/0263457 A1	10/2011	Krutzik et al.
2006/0057634 A1	3/2006	Rye	2011/0294689 A1	12/2011	Namsaraev
2006/0073506 A1	4/2006	Christians et al.	2011/0312511 A1	12/2011	Winquist et al.
2006/0211030 A1	9/2006	Brenner	2012/0004132 A1	1/2012	Zhang et al.
2006/0257902 A1	11/2006	Mendoza et al.	2012/0010091 A1	1/2012	Linnarson
2006/0263709 A1	11/2006	Matsumura et al.	2012/0014977 A1	1/2012	Furihata et al.
2006/0263789 A1	11/2006	Kincaid	2012/0034607 A1	2/2012	Rothberg et al.
2006/0280352 A1	12/2006	Muschler et al.	2012/0040843 A1	2/2012	Ducree et al.
2006/0281092 A1	12/2006	Wille et al.	2012/0045844 A1	2/2012	Rothberg et al.
2006/0286570 A1	12/2006	Rowlen et al.	2012/0058520 A1	3/2012	Hayashida
2007/0020640 A1	1/2007	Mccloskey et al.	2012/0058902 A1	3/2012	Livingston et al.
2007/0031829 A1	2/2007	Yasuno et al.	2012/0065081 A1	3/2012	Chee
2007/0042400 A1	2/2007	Choi et al.	2012/0071331 A1	3/2012	Casbon
2007/0042419 A1	2/2007	Barany et al.	2012/0087862 A1	4/2012	Hood et al.
2007/0065823 A1	3/2007	Dressman et al.	2012/0142018 A1	6/2012	Jiang
2007/0065844 A1	3/2007	Golub et al.	2012/0149603 A1	6/2012	Cooney et al.
2007/0105090 A1	5/2007	Cassidy et al.	2012/0156675 A1	6/2012	Lueerssen et al.
2007/0117121 A1	5/2007	Hutchison et al.	2012/0163681 A1	6/2012	Lohse
2007/0117134 A1	5/2007	Kou	2012/0165219 A1	6/2012	Van Der Zaag et al.
2007/0117177 A1	5/2007	Luo et al.	2012/0173159 A1	7/2012	Davey et al.
2007/0133856 A1	6/2007	Dutta-Choudhury	2012/0190020 A1	7/2012	Oliphant et al.
2007/0172873 A1	7/2007	Brenner et al.	2012/0202293 A1	8/2012	Martin et al.
2007/0178478 A1	8/2007	Dhallan et al.	2012/0220022 A1	8/2012	Ehrlich et al.
2007/0202523 A1	8/2007	Becker et al.	2012/0220494 A1	8/2012	Samuels et al.
2007/0281317 A1	12/2007	Becker et al.	2012/0231972 A1	9/2012	Golyshin et al.
2008/0038727 A1	2/2008	Spier	2012/0252012 A1	10/2012	Armougom et al.
			2012/0253689 A1	10/2012	Rogan
			2012/0316074 A1	12/2012	Saxonov
			2012/0322681 A1	12/2012	Kung et al.
			2013/0005585 A1	1/2013	Anderson et al.

(56)	References Cited					
U.S. PATENT DOCUMENTS						
2013/0022977 A1	1/2013 Lapidus et al.	2015/0307874 A1	10/2015 Jaitin			
2013/0045994 A1	2/2013 Shinozuka et al.	2015/0329852 A1	11/2015 Nolan			
2013/0116130 A1	5/2013 Fu et al.	2015/0360193 A1	12/2015 Fan et al.			
2013/0190206 A1	7/2013 Leonard	2015/0376609 A1	12/2015 Hindson et al.			
2013/0203047 A1	8/2013 Casbon et al.	2016/0017320 A1	1/2016 Wang et al.			
2013/0210643 A1	8/2013 Casbon et al.	2016/0026758 A1	1/2016 Jabara et al.			
2013/0210659 A1	8/2013 Watson et al.	2016/0053253 A1	2/2016 Salathia et al.			
2013/0224743 A1	8/2013 Casbon et al.	2016/0055632 A1	2/2016 Fu et al.			
2013/0225418 A1	8/2013 Watson	2016/0060621 A1	3/2016 Agresti et al.			
2013/0225623 A1	8/2013 Buxbaum et al.	2016/0060682 A1	3/2016 Pregibon et al.			
2013/0237458 A1	9/2013 Casbon et al.	2016/0068889 A1	3/2016 Gole et al.			
2013/0267424 A1	10/2013 Casbon et al.	2016/0122751 A1	5/2016 LaBaer			
2013/0274117 A1	10/2013 Church	2016/0122753 A1	5/2016 Mikkelsen et al.			
2013/0232372 A1	12/2013 Anderson et al.	2016/0138091 A1	5/2016 Chee et al.			
2014/0004569 A1	1/2014 Lambowitz et al.	2016/0145677 A1	5/2016 Chee et al.			
2014/0057799 A1	2/2014 Johnson et al.	2016/0145683 A1	5/2016 Fan et al.			
2014/0066318 A1	3/2014 Frisen et al.	2016/0153973 A1	6/2016 Smith			
2014/0147860 A1	5/2014 Kaduchak et al.	2016/0201125 A1	7/2016 Samuels et al.			
2014/0155274 A1	6/2014 Xie et al.	2016/0208322 A1	7/2016 Anderson et al.			
2014/0155295 A1	6/2014 Hindson et al.	2016/0222378 A1	8/2016 Fodor et al.			
2014/0178438 A1	6/2014 Sahin et al.	2016/0232291 A1	8/2016 Kyriazopoulou-Panagiotopoulou et al.			
2014/0194324 A1	7/2014 Gormley et al.					
2014/0206079 A1	7/2014 Malinoski et al.					
2014/0206547 A1	7/2014 Wang					
2014/0216128 A1	8/2014 Trotter et al.					
2014/0227684 A1	8/2014 Hindson et al.					
2014/0227705 A1	8/2014 Vogelstein et al.					
2014/0228239 A1	8/2014 McCoy et al.					
2014/0228255 A1	8/2014 Hindson et al.					
2014/0235506 A1	8/2014 Hindson et al.					
2014/0243242 A1	8/2014 Nicol et al.					
2014/0244742 A1	8/2014 Yu et al.					
2014/0272952 A1	9/2014 May et al.					
2014/0274811 A1	9/2014 Arnold					
2014/0287963 A1	9/2014 Hindson et al.					
2014/0303005 A1	10/2014 Samuels et al.					
2014/0309945 A1	10/2014 Park et al.					
2014/0315211 A1	10/2014 Sugino et al.					
2014/0322716 A1	10/2014 Robins					
2014/0357500 A1	12/2014 Vigneault et al.					
2014/0378322 A1	12/2014 Hindson et al.					
2014/0378345 A1	12/2014 Hindson et al.					
2014/0378349 A1	12/2014 Hindson et al.					
2014/0378350 A1	12/2014 Hindson et al.					
2015/0005199 A1	1/2015 Hindson et al.					
2015/0005200 A1	1/2015 Hindson et al.					
2015/0011396 A1	1/2015 Schroeder et al.					
2015/0017654 A1	1/2015 Gorfinkel et al.					
2015/0066385 A1	3/2015 Schnall-Levin et al.					
2015/0072867 A1	3/2015 Soldatov et al.					
2015/0072873 A1	3/2015 Heinz et al.					
2015/0080266 A1	3/2015 Volkmut et al.					
2015/0099661 A1	4/2015 Fodor et al.					
2015/0099673 A1	4/2015 Fodor et al.					
2015/0111256 A1	4/2015 Church et al.					
2015/0118680 A1	4/2015 Fodor et al.					
2015/0119255 A1	4/2015 Fodor et al.					
2015/0119256 A1	4/2015 Fodor et al.					
2015/0119257 A1	4/2015 Fodor et al.					
2015/0119258 A1	4/2015 Fodor et al.					
2015/0119290 A1	4/2015 Fodor et al.					
2015/0133319 A1	5/2015 Fu et al.					
2015/0141292 A1	5/2015 Fodor et al.					
2015/0152409 A1	6/2015 Seitz et al.					
2015/0203897 A1	7/2015 Robons et al.					
2015/0211050 A1	7/2015 Iafrate et al.					
2015/0218620 A1	8/2015 Behlke et al.					
2015/0225777 A1	8/2015 Hindson et al.					
2015/0225778 A1	8/2015 Hindson et al.					
2015/0247182 A1	9/2015 Faham et al.					
2015/0253237 A1	9/2015 Castellarnau et al.					
2015/0259734 A1	9/2015 Asbury et al.					
2015/0275295 A1	10/2015 Wang et al.					
2015/0298091 A1	10/2015 Weitz					
2015/0299784 A1	10/2015 Fan et al.					

# US 12,391,940 B2

Page 5

---

(56)	References Cited						
U.S. PATENT DOCUMENTS							
2019/0095578 A1	3/2019	Shum et al.	EP	2970958 B1	12/2017		
2019/0119726 A1	4/2019	Shum et al.	EP	3263715 A1	1/2018		
2019/0136316 A1	5/2019	Hindson et al.	EP	2670863 B1	6/2018		
2019/0136317 A1	5/2019	Hindson et al.	EP	3136103 B1	8/2018		
2019/0136319 A1	5/2019	Hindson et al.	EP	2954102 B1	12/2018		
2019/0161743 A1	5/2019	Church et al.	EP	3428290 A1	1/2019		
2019/0177788 A1	6/2019	Hindson et al.	EP	2970957 B1	4/2019		
2019/0177800 A1	6/2019	Boutet et al.	GB	3058092 B1	5/2019		
2019/0203270 A1	7/2019	Amit et al.	JP	3256606 B1	5/2019		
2019/0203291 A1	7/2019	Hindson et al.	JP	3327123 B1	8/2019		
2019/0211395 A1	7/2019	Tsao et al.	EP	3587589 A1 *	1/2020	..... C12Q 1/6806	
2019/0218276 A1	7/2019	Regev et al.	GB	2293238 A	3/1996		
2019/0218607 A1	7/2019	Love et al.	JP	H04108385	4/1992		
2019/0221287 A1	7/2019	Tsujimoto	JP	2001078768 A	3/2001		
2019/0221292 A1	7/2019	Tsujimoto	JP	2005233974 A	9/2005		
2019/0256888 A1	8/2019	Weissleder et al.	WO	2007504831 A	3/2007		
2019/0256907 A1	8/2019	Ryan et al.	WO	2008256428 A	10/2008		
2019/0292592 A1	9/2019	Shum et al.	WO	2013039275 A	2/2013		
2019/0338353 A1	11/2019	Belgrader et al.	WO	WO1989001050	2/1989		
2019/0338357 A1 *	11/2019	Fan .....	C12Q 1/6876	WO	WO1996024061	8/1996	
2019/0390253 A1	12/2019	Kennedy et al.	WO	WO1997010365	3/1997		
2020/0102598 A1	4/2020	Xie et al.	WO	WO1999015702	4/1999		
2020/0109437 A1	4/2020	Chang et al.	WO	WO1999028505	6/1999		
2020/0115753 A1	4/2020	Shalek et al.	WO	WO2000058516	10/2000		
2020/0149037 A1	5/2020	Shum	WO	WO2001020035	3/2001		
2020/0157600 A1 *	5/2020	Fu .....	C12N 15/1006	WO	WO2001048242	7/2001	
2021/0039582 A1	2/2021	Patton et al.	WO	WO2002072772	9/2002		
2021/0123044 A1	4/2021	Zhang et al.	WO	WO2002079490	10/2002		
2021/0132078 A1	5/2021	Peikon et al.	WO	WO2002083922	10/2002		
2021/0198754 A1	7/2021	Fan et al.	WO	WO2002101358	12/2002		
2021/0213413 A1	7/2021	Saligrama et al.	WO	WO2003031591	4/2003		
2021/0214770 A1	7/2021	Prosen et al.	WO	WO2003035829	5/2003		
2021/0214784 A1	7/2021	Prosen et al.	WO	WO2004017374	2/2004		
2021/0222163 A1	7/2021	Wu et al.	WO	WO2004021986	3/2004		
2021/0222244 A1	7/2021	Martin et al.	WO	WO2004043669	4/2004		
2021/0230582 A1	7/2021	Fu et al.	WO	WO2004066185	8/2004		
2021/0230583 A1	7/2021	Lam et al.	WO	WO2004081225	9/2004		
2021/0230666 A1	7/2021	Wu et al.	WO	WO2005017206	2/2005		
2021/0246492 A1	8/2021	Song et al.	WO	WO2005021731	3/2005		
2021/0263019 A1	8/2021	Martin et al.	WO	WO2005042759	5/2005		
2021/0371914 A1	12/2021	Stoeckius et al.	WO	WO2005071110	8/2005		
2022/0154288 A1	5/2022	Mortimer	WO	WO2005080604	9/2005		
2022/0162695 A1	5/2022	Sakofsky et al.	WO	WO2005111242	11/2005		
2022/0162773 A1	5/2022	Sakofsky et al.	WO	WO2006026828	3/2006		
2022/0178909 A1	6/2022	Huang et al.	WO	WO2006071776	7/2006		
2022/0214356 A1	7/2022	Henikoff et al.	WO	WO2006102264	9/2006		
2022/0219170 A1	7/2022	Khurana et al.	WO	WO2006137932	12/2006		
2022/0220549 A1	7/2022	Shum et al.	WO	WO2007087310	8/2007		
2022/0267759 A1	8/2022	Sanjana et al.	WO	WO2007087312	8/2007		
2022/0333185 A1	10/2022	Fu et al.	WO	WO2007147079	12/2007		
2022/0348904 A1	11/2022	Shum et al.	WO	WO2008047428	4/2008		
2023/0083422 A1	3/2023	Fu et al.	WO	WO2008051928	5/2008		
2023/0109336 A1	4/2023	Shum et al.	WO	WO2008057163	5/2008		
2023/0125113 A1	4/2023	Fan et al.	WO	WO2008096318	8/2008		
2023/0193372 A1	6/2023	Shum et al.	WO	WO2008104380	9/2008		
FOREIGN PATENT DOCUMENTS							
CN	106460033 A	2/2017	WO	WO2008147428	12/2008		
CN	107208158 A	9/2017	WO	WO2008150432	12/2008		
DE	102008025656	12/2009	WO	WO2009048530	4/2009		
EP	1473080 A2	11/2004	WO	WO2009148560	12/2009		
EP	1647600 A2	4/2006	WO	WO2009152928	12/2009		
EP	1845160 A1	10/2007	WO	WO2010048605	4/2010		
EP	2036989 A1	3/2009	WO	WO2010059820	5/2010		
EP	1379693 B1	5/2009	WO	WO2010117620	10/2010		
EP	2204456 A1	7/2010	WO	WO2011091393	7/2011		
EP	2431465 A1	3/2012	WO	WO2011106738	9/2011		
EP	2203749 B1	8/2012	WO	WO2011123246	10/2011		
EP	2511708 A1	10/2012	WO	WO2011127099	10/2011		
EP	2538220 A1	12/2012	WO	WO2011143659	11/2011		
EP	2623613 A1	8/2013	WO	WO2011155833	12/2011		
EP	1745155 B1	10/2014	WO	WO2012038839	3/2012		
EP	2805769 A1	11/2014	WO	WO2012041802	4/2012		
EP	2556171 B1	9/2015	WO	WO2012042374	4/2012		

(56)	References Cited		WO	WO2016118915	7/2016	
	FOREIGN PATENT DOCUMENTS		WO	WO2016126871	8/2016	
			WO	WO2016130578	8/2016	
			WO	WO2016160965	8/2016	
WO	WO2012047297	4/2012	WO	WO2016138496	9/2016	
WO	WO2012048341	4/2012	WO	WO2016138500	9/2016	
WO	WO2012083225	6/2012	WO	WO2016145409	9/2016	
WO	WO2012099896	7/2012	WO	WO2016149418	9/2016	
WO	WO2012103154	8/2012	WO	WO2016160844	10/2016	
WO	WO2012106546	8/2012	WO	WO2016168825	10/2016	
WO	WO2012108864	8/2012	WO	WO2016172373	10/2016	
WO	WO2012112804	8/2012	WO	WO2016176091	11/2016	
WO	WO2012129363	9/2012	WO	WO2016190795	12/2016	
WO	WO2012140224	10/2012	WO	WO2016191272	12/2016	
WO	WO2012142213	10/2012	WO	WO2017032808	3/2017	
WO	WO2012148477	11/2012	WO	WO2017040306	3/2017	
WO	WO2012148497	11/2012	WO	WO2017044574	3/2017	
WO	WO2012149042	11/2012	WO	WO2017053905	3/2017	
WO	WO2012156744	11/2012	WO	WO2017079593	5/2017	
WO	WO2012162267	11/2012	WO	WO2017087873	5/2017	
WO	WO2012177639	12/2012	WO	WO2017096239	6/2017	
WO	WO2013019075	2/2013	WO	WO2017097939	6/2017	
WO	WO2013070990	5/2013	WO	WO2017117358	7/2017	
WO	WO2013096802	6/2013	WO	WO2017125508	7/2017	
WO	WO2013117595	8/2013	WO	WO2017139690	8/2017	
WO	WO2013130674	9/2013	WO	WO2017164936	9/2017	
WO	WO2013137737	9/2013	WO	WO2017173328	10/2017	
WO	WO2013148525	10/2013	WO	WO2017205691	11/2017	
WO	WO2013173394	11/2013	WO	WO2018017949	1/2018	
WO	WO2013176767	11/2013	WO	WO2018018008	1/2018	
WO	WO2013177206	11/2013	WO	WO2018020489	2/2018	
WO	WO2013188831	12/2013	WO	WO2018031631	2/2018	
WO	WO2013188872	12/2013	WO	WO2018058073	3/2018	
WO	WO2013191775	12/2013	WO	WO2018064640	4/2018	
WO	WO2014015084	1/2014	WO	WO2018075693	4/2018	
WO	WO2014015098	1/2014	WO	WO2018111872	6/2018	
WO	WO2014018093	1/2014	WO	WO2018115852	6/2018	
WO	WO2014018460	1/2014	WO	WO2018119447	6/2018	
WO	WO2014028537	2/2014	WO	WO2018132635	7/2018	
WO	WO2014031997	2/2014	WO	WO2018140966	8/2018	
WO	WO2014065756	5/2014	WO	WO2018144240	8/2018	
WO	WO2014093676	6/2014	WO	WO2018144813	8/2018	
WO	WO2014108850	7/2014	WO	WO2018165366	9/2018	
WO	WO2014124046	8/2014	WO	WO2018165366 A1 *	9/2018 .....	
WO	WO2014124336	8/2014	WO	WO2018174827	9/2018	
WO	WO2014124338	8/2014	WO	WO2018218226 A1	11/2018	
WO	WO2014126937	8/2014	WO	WO2018217862	11/2018	
WO	WO2014144495	9/2014	WO	WO2018218222	11/2018	
WO	WO2014145458	9/2014	WO	WO2018222548	12/2018	
WO	WO2014176575	10/2014	WO	WO2018226293	12/2018	
WO	WO2014200767	12/2014	WO	WO2019055852	3/2019	
WO	WO2014201273	12/2014	WO	WO2019076768	4/2019	
WO	WO2014204939	12/2014	WO	WO2019099906 A1	5/2019	
WO	WO2014210223	12/2014	WO	WO2019084046	5/2019	
WO	WO2014210225	12/2014	WO	WO2019099906	5/2019	
WO	WO2014210353	12/2014	WO	WO2019113457	6/2019	
WO	WO2015002908	1/2015	WO	WO2019113499	6/2019	
WO	WO2018015365	1/2015	WO	WO2019113506	6/2019	
WO	WO2015017586	2/2015	WO	WO2019113533	6/2019	
WO	WO2015031691	3/2015	WO	WO2019118355	6/2019	
WO	WO2015035087	3/2015	WO	WO2019126789	6/2019	
WO	WO2015044428	4/2015	WO	WO2019157529	8/2019	
WO	WO2015047186	4/2015	WO	WO2019178164	9/2019	
WO	WO2015057985	4/2015	WO	WO-2019210049 A1 *	10/2019 .....	
WO	WO2014071361	5/2015	WO	WO2019/213294 A1	11/2019	
WO	WO2015061844	5/2015	WO	WO2019213237	11/2019	
WO	WO2015103339	7/2015	WO	WO2019213294	11/2019	
WO	WO-2015104302 A1 *	7/2015 .....	C07H 21/04	WO	WO2019218101	11/2019
WO	WO2015117163	8/2015	WO	WO2020028266	2/2020	
WO	WO2015134787	9/2015	WO	WO2020033164	2/2020	
WO	WO2015160439	10/2015	WO	WO2020037065	2/2020	
WO	WO2015168161	11/2015	WO	WO2020046833	3/2020	
WO	WO2015179339	11/2015	WO	WO2020072380	4/2020	
WO	WO2015200869	12/2015	WO	WO2020097315	5/2020	
WO	WO2015200893	12/2015	WO	WO2020123384	6/2020	
WO	WO2016044227	3/2016	WO	WO2020131699	6/2020	
WO	WO2016049418	3/2016	WO	WO2020154247	7/2020	
WO	WO2016061517	4/2016	WO	WO2020159757	8/2020	
WO	WO2016100976	6/2016	WO	WO2020167920	8/2020	

(56)

**References Cited**

## FOREIGN PATENT DOCUMENTS

WO	WO2020214642	10/2020
WO	WO2020219721	10/2020
WO	WO2020242377	12/2020
WO	WO2021092386	5/2021
WO	WO2021142233	7/2021
WO	WO2021146207	7/2021
WO	WO2021146219	7/2021
WO	WO2021146636	7/2021
WO	WO2021155057	8/2021
WO	WO2021155284	8/2021
WO	WO2021163374	8/2021
WO	WO2021168015	8/2021
WO	WO2021168261	8/2021
WO	WO20210178199	9/2021
WO	WO2021247593	12/2021
WO	WO2022015667	1/2022
WO	WO2022026909	2/2022
WO	WO2022040453	2/2022
WO	WO2022143221	7/2022
WO	WO2022256324	12/2022
WO	WO2023/034739	3/2023
WO	WO2023/034789	3/2023
WO	WO2023/034790	3/2023
WO	WO2023/034794	3/2023
WO	WO2023/034872	3/2023
WO	WO2023/039433	3/2023

## OTHER PUBLICATIONS

- Agasti et al., "Photocleavable DNA barcode-antibody conjugates allow sensitive and multiplexed protein analysis in single cell," *J Am Chem Soc.* 2012, 134(45), 18499-18502.
- Ahern, "Biochemical, Reagent Kits Offer Scientists Good Return on Investment," *The Scientist* 1995, 9(15), in 5 pages.
- Alexandra M. Ewing of Richards, Layton and Finger, P.A., Entry of Appearance dated Jan. 18, 2019 in the USDC District of Delaware, C.A. No. 18-1800-RGA, 1 pp.
- Alkan et al., "Personalized copy No. and segmental duplication maps using next-generation sequencing," *Nat Genet.* 2009, 41(10):1061-1067.
- Anderson, "Study Describes RNA Sequencing Applications for Molecular Indexing Methods," *GenomeWeb* 2014, 5 pp.
- Ansorge, "Next-generation DNA sequencing techniques," *New Biotechnology* 2009, 25(4), 195-203.
- Applied Biosystems, Apr. 2008, SOLiD™ System Barcoding, Application Note, 4 pp.
- Argrawal et al., "Counting Single Native Biomolecules and Intact Viruses with Color-Coded Nanoparticles," *Analytical Chemistry* 2006, 78, 1061-1070.
- Arslan et al., "An efficient algorithm for the stochastic simulation of the hybridization of DNA to microarrays," *BMC Bioinformatics* 2009, 10(411), 1-17.
- Atanur et al., "The genome sequence of the spontaneously hypertensive rat: Analysis and functional significance." *Genome Res.* 2010, 20(6), 791-803.
- Audic et al., "The Significance of Digital Gene Expression Profiles," *Genome Res.* 1997, 7, 986-995.
- Baek et al., "Development of Hydrogel TentaGel Shell-Core Beads for Ultra-high Throughput Solution Phase Screening of Encoded OBOC Combinatorial Small Molecule Libraries," *J. Comb Chem.* 2009, 11(1), 91-102.
- BD Life Sciences, 2018, BD AbSeq antibody-oligo conjugates, [www.bd.com/genomics](http://www.bd.com/genomics), 2 pp.
- BD Life Sciences, 2018, BD AbSeq on the BD Rhapsody system: Exploration of single-cell gene regulation by simultaneous digital mRNA and protein quantification, [www.bd.com/genomics](http://www.bd.com/genomics), 7 pp.
- Bendall et al., "Single-Cell Mass Cytometry of Differential Immune and Drug Responses Across a Human Hematopoietic Continuum," *Science* 2011, 332(6030), 687-696.
- BioNumbers, Aug. 21, 2010, "Useful fundamental numbers in molecular biology," 2001-2004. <http://biornumbers.hms.harvard.edu/KeyNumbers.aspx>, 1-4.
- Biosciences Product Catalogue, Dynal® Catalog 1999, Oslo, Norway, 49-51.
- Bioscribe "Massively parallel sequencing technology for single-cell gene expression published" (press release), PhysOrg 2015, 1-2.
- Blainey, "The future is now: single-cell genomics of bacteria and archaea," *FEMS Microbiol Rev.* 2013, 37(3), 407-427.
- Bogdanova et al., "Normalization of full-length enriched cDNA," *Molecular Biosystems* 2008, 4(3), 205-212.
- Bonaldo et al., "Normalization and Subtraction: Two Approaches to facilitate Gene Discovery," *Genome Res.* 1996, 6, 791-806.
- Bontoux et al., "Integrating whole transcriptome assays on a lab-on-a-chip for single cell gene profiling", *Lab on a Chip* 2008, 8(3), 443-450.
- Bose et al., "Scalable microfluidics for single-cell RNA printing and sequencing," *Genome Biology* 2015, 16(120), 1-16.
- Brady et al., "Construction of cDNA libraries from single cells", *Methods in Enzymology* 1993, (225), 611-623.
- Braha et al., "Simultaneous stochastic sensing of divalent metal ions," *Nature Biotechnology* 2000, 18, 1005-1007.
- Bratke et al., "Differential expression of human granzymes A, B, and K in natural killer cells and during CD8+ T cell differentiation in peripheral blood," *Eur J Immunol.* 2005, 35, 2608-2616.
- Brenner et al., "Gene expression analysis by massively parallel signature sequencing (MPSS) on microbead arrays," *Nature Biotechnology* 2000, 18, 630-634.
- Brenner et al., "In vitro cloning of complex mixtures of DNA on microbeads: Physical separation of differentially expressed cDNAs," *PNAS* 2000, 97(4), 1665-1670.
- Brinza et al., "Detection of somatic mutations at 0.1% frequency from cfDNA in peripheral blood with a multiplex next-generation sequencing assay," Conference Poster, AACR 107th Annual Meeting, Apr. 16-20, 2016, 1 p.
- Brisco et al., "Quantification of RNA integrity and its use for measurement of transcript number," *Nucleic Acids Research* 2012, 40(18), e144, 1-9.
- Brodin et al., "Challenges with Using Primer IDs to Improve Accuracy of Next Generation Sequencing," *PLoS One* 2015, 19(3), 1-12.
- Brouillette et al., "A Simple and Novel Method for RNA-seq Library Preparation of Single Cell cDNA Analysis by Hyperactive Tn5 Transposase," *Developmental Dynamics* 2012, 241, 1584-1590.
- Buggenom et al., "A covalent and cleavable antibody DNA conjugation strategy for sensitive protein detection via immunoPCR," *Scientific Reports* 2016, 6(22675), 1-12.
- Buschmann et al., Enhancing the detection of barcoded reads in high throughput DNA sequencing DNA by controlling the false discovery rate, *BMC Bioinformatics*, 15(1), 264, 1-16. 2014.
- Bustin, "Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays," *Journal of Molecular Endocrinology* 2000, 25, 169-193.
- Butkus, "Cellular research set to launch first gene expression platform using 'molecular indexing' technology," *GenomeWeb* 2014, 1-5.
- Cai, "Turning single cells in microarrays by super-resolution barcoding," *Briefings in Functional Genomics* 2012, 12(2), 75-80.
- Cao et al., "Comprehensive single-cell transcriptional profiling of a multicellular organism," *Science* 2017, 357, 661-667.
- Carr et al., "Inferring relative proportions of DNA variants from sequencing electropherograms," *Bioinformatics* 2009, 25(24), 3244-3250.
- Caruccio et al., "Nextera (TM) Technology for NGS DNA library Preparation: Simultaneous Fragmentation and Tagging by in Vitro Transposition," *EpiBio* 2009, 16(3), 4-6.
- Casbon et al., "A method for counting PCR template molecules with application to next-generation sequencing," *Nucleic Acids Res.* 2011, 39(12), e81, 1-8.
- Castellarnau et al., "Stochastic particle barcoding for single-cell tracking and multiparametric analysis," *Small* 2015, 11(4), 489-498.

(56)

**References Cited****OTHER PUBLICATIONS**

- Castle et al., "DNA copy No., including telomeres and mitochondria, assayed using next-generation sequencing," *BMC Genomics* 2010, 11(244), 1-11.
- Chamberlain et al., "Deletion screening of the Duchenne muscular dystrophy locus via multiplex DNA amplification," *Nucleic Acids Res.* 1988, 16(23), 11141-11156.
- Chang et al., "Detection of Allelic Imbalance in Ascitic Supernatant by Digital Single Nucleotide Polymorphism Analysis," *Clinical Cancer Research* 2002, 8, 2580-2585.
- Chapin et al., "Rapid microRNA Profiling on Encoded Gel Microparticles," *Angew Chem Int Ed Engl.* 2011, 50(10), 2289-2293.
- Chee et al., "Accessing genetic information with high-density DNA arrays," *Science* 1996, 274, 610-614.
- Chee, "Enzymatic multiplex DNA sequencing," *Nucleic Acids Research* 1991, 19(12), 3301-3305.
- Chen et al., "Spatially resolved, highly multiplexed RNA profiling in single cells," *Science Express* 2015, 348(6233), aaa6090, 1-36.
- Church et al., "Multiplex DNA sequencing," *Science* 1988, 240(4849), 185-188.
- Civil Cover Sheet filed Nov. 15, 2018 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 1 pp.
- Clontech Laboratories, Inc., "Smart<sup>TM</sup> PCR cDNA Synthesis Kit User Manual," Clontech 2007, 1-39.
- Cloonan et al., "Stem cell transcriptome profiling via massive-scale mRNA sequencing", *Nature Methods* 2008, 5(7), 613-619.
- Combined Search and Examination Report dated Aug. 6, 2014 in UK Patent Application No. 1408829.8.
- Combined Search and Examination Report dated Feb. 21, 2017 in UK Patent Application No. 1609740.4.
- Complaint filed in *Becton, Dickinson and Company and Cellular Research Inc. v. 10X Genomics, Inc.* dated Nov. 15, 2018 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 141 pp.
- Costa et al., "Single-Tube Nested Real-Time PCR as a New Highly Sensitive Approach to Trace Hazelnut," *Journal of Agricultural and Food Chemistry* 2012, 60, 8103-8110.
- Costello et al., "Discovery and characterization of artefactual mutations in deep coverage targeted capture sequencing data due to oxidative DNA damage during sample preparation," *Nucleic Acids Res* 2013, 41(6), e67, 1-12.
- Cotten et al., "Selection of proteins with desired properties from natural proteome libraries using mRNA display," *Nature Protocols* 2011, 6, 1163-1182.
- Cox, "Bar coding objects with DNA," *Analyst* 2001, 126, 545-547.
- Craig et al., "Identification of genetic variants using bar-coded multiplexed sequencing," *Nat Methods* 2008, 5(10), 887-893.
- Cusanovich et al., "Multiplex single-cell profiling of chromatin accessibility by combinatorial cellular indexing," *Science* 2015, 348(6237), 910-914.
- Custom Antibody Services, Precision Antibody, accessed Apr. 16, 2014, 2 pp.
- Daines et al., "High-throughput multiplex sequencing to discover copy No. variants in *Drosophila*," *Genetics* 2009, 182(4), 182, 935-941.
- Dalerba et al., "Single-cell dissection of transcriptional heterogeneity in human colon tumors," *Nat Biotechnol.* 2011, 29(12), 1120-1127.
- D'Antoni et al., "Rapid quantitative analysis using a single molecule counting approach," *Anal Biochem.* 2006, 352, 97-109.
- Daser et al., "Interrogation of genomes by molecular copy-number counting (MCC)," *Nature Methods* 2006, 3(6), 447-453.
- Day et al., "Immobilization of polynucleotides on magnetic particles," *Biochem. J.* 1991, 278, 735-740.
- Decision of Refusal dated Aug. 21, 2017 in Japanese Patent Application No. 2014-558975.
- Defendant 10X Genomics, Inc.'s Letter to Judge Andrews in Response to Plaintiff's Letter of Supplemental Authority, dated Jul. 11, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 2 pp.
- Defendant 10X Genomics Motion for Admission Pro Hac Vice of Paul Ehrlich, Azra Hadzimehmmedovic and Aaron Nathan, Pursuant to Local Rule 83.5, dated May 1, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 5 pp.
- Defendant 10X Genomics, Inc.'s Motion for Admission Pro Hac Vice Pursuant to Local Rule 83.5, dated Jan. 18, 2019 in the USDC District of Delaware, C.A. No. 18-1800-RGA, 5 pp.
- Defendant 10X Genomics, Inc.'s Motion to Dismiss Pursuant to Federal Rule of Civil Procedure 12(b)(6), dated Jan. 18, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 1 pp.
- Defendant 10X Genomics, Inc.'s Motion to Dismiss the First Amended Complaint Pursuant to Federal Rule of Civil Procedure 12(b)(6), dated Mar. 1, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 1 pp.
- Defendant 10X Genomics Notice of Service for Initial Disclosures served to Opposing Counsel, dated Jun. 7, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 2 pp.
- Defendant 10X Genomic Inc.'s Notice of Service for Initial Requests for Production and Interrogatories Served to Becton, Dickinson, and Company and Cellular Research, Inc., dated May 31, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 2 pp.
- Defendant 10X Genomics, Inc.'s Notice of Service of Technical Documents, dated Jul. 8, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 2 pp.
- Defendant 10X Genomics, Inc.'s Opening Brief in Support of Its Motion to Dismiss Pursuant to Federal Rule of Civil Procedure 12(b)(6), dated Jan. 18, 2019 in the USDC District of Delaware, C.A. No. 1:18-cv-01800-RGA, 25 pp.
- Defendant 10X Genomics, Inc.'s Opening Brief in Support of Its Motion to Dismiss Pursuant to Federal Rule of Civil Procedure 12(b)(6), dated Mar. 1, 2019 in the USDC District of Delaware, C.A. No. 18-1800 RGA, 26 pp.
- Defendant 10X Genomics, Inc.'s [Proposed] Order for Partial Dismissal Pursuant to Federal Rules of Civil Procedure 12(b)(6), dated Jan. 18, 2019 in the USDC District of Delaware, C.A. No. 18-1800-RGA, 1 pp.
- Defendant 10X Genomics, Inc.'s Proposed Order for Dismissal pursuant to Federal Rules of Civil Procedure 12(b)(6), filed Mar. 1, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 1 pp.
- Defendant 10X Genomics Reply Brief in Support of its Motion to Dismiss Pursuant to Federal Rule of Civil Procedure 12(b)(6), dated Apr. 12, 2019 in the USDC for the District of Delaware, C.A. No. 18-1800-RGA, 15 pp.
- Defendant 10X Genomics Request for Oral Argument Under D. Del. LR 7.1.4, dated Apr. 18, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA 2 pp.
- Defendant 10X Genomics Response Letter to Judge Richard G. Andrews re Request for a Rule 16, dated Apr. 16, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 2 pp.
- Defendant 10X Genomics, Inc.'s Rule 7.1 Disclosure Statement, dated Jan. 18, 2019 in the USDC District of Delaware, C.A. No. 18-1800-RGA, 1 pp. 1.
- Delley et al., "Combined aptamer and transcriptome sequencing of single cells," *bioRxiv* 2017, 1-10.
- De Saizieu et al., "Bacterial transcript imaging by hybridization of total RNA to oligonucleotide arrays," *Nature Biotechnology* 1988, 16, 45-48.
- Di Carlo et al., "Dynamic single-cell analysis for quantitative biology," *Analytical Chemistry* 2006, 78(23), 7918-7925.
- Dirks et al., Triggered amplification by hybridization chain reaction., *Proc Natl Acad Sci* 2014, 101(43), 15275-15278.
- Dube et al., "Mathematical Analysis of Copy Number Variation in a DNA Sample Using Digital PCR on a Nanofluidic Device," *PLoS One* 2008, 3(8) e2876.
- Eberwine et al., "Analysis of gene expression in single live neurons," *Proc. Natl. Acad. Sci.* 1992, 89, 3010-3014.
- Evanko et al., "Hybridization chain reaction," *Nature Methods* 2004, 1(3), 186-187.
- Examination Report dated Oct. 24, 2017 in Australian Patent Application No. 2013226081.
- Examination Report dated Jul. 20, 2018 in Australian Patent Application No. 2014312208.

(56)

**References Cited****OTHER PUBLICATIONS**

- Examination Report dated May 12, 2020 in Australian Patent Application No. 2018220004.
- Examination Report dated Jul. 12, 2016 in European Patent Application No. 13755319.4.
- Examination Report dated Apr. 10, 2017 in European Patent Application No. 14761937.3.
- Examination Report dated Oct. 10, 2017 in European Patent Application No. 14761937.3.
- Examination Report dated Mar. 16, 2018 in European Patent Application No. 13754428.4.
- Examination Report dated Sep. 5, 2018 in European Patent Application No. 16710357.1.
- Examination Report dated Sep. 26, 2018 in European Patent Application No. 16714081.3.
- Examination Report dated Dec. 12, 2018 in European Patent Application No. 16719706.0.
- Examination Report dated Jan. 2, 2019 in European Patent Application No. 16757986.1.
- Examination Report dated Feb. 6, 2019 in European Patent Application No. 13754428.4.
- Examination Report dated Apr. 26, 2019 in European Patent Application No. 16710357.1.
- Examination Report dated Jun. 18, 2019 in European Patent Application No. 16710551.9.
- Examination Report dated Jul. 24, 2019 in European Patent Application No. 16714081.3.
- Examination Report dated Aug. 2, 2019 in European Patent Application No. 17202409.3.
- Examination Report dated Oct. 11, 2019 in European Patent Application No. 16757986.1.
- Examination Report dated Dec. 4, 2019 in European Patent Application No. 16719706.0.
- Examination Report dated Feb. 19, 2020 in European Patent Application No. 16710551.9.
- Examination Report dated Mar. 18, 2020 in European Patent Application No. 17202409.3.
- Examination Report dated Jul. 6, 2020 in European Patent Application No. 17781265.8.
- Examination Report dated Sep. 21, 2020 in European Patent Application No. 18703156.2.
- Examination Report dated Nov. 12, 2020 in European Patent Application No. 18716877.8.
- Examination Report dated Dec. 3, 2020 in European Patent Application No. 16719706.0.
- Examination Report dated Mar. 25, 2021 in European Patent Application No. 17781265.8.
- Examination Report dated Oct. 8, 2021 in European Patent Application No. 18716877.8.
- Examination Report dated Mar. 18, 2019 in Singapore Patent Application No. 11201405274W.
- Examination Report dated Jan. 27, 2016 in United Kingdom Patent Application No. 1408829.8.
- Examination Report dated Feb. 19, 2016 in United Kingdom Patent Application No. GB1511591.8.
- Examination Report dated Jun. 8, 2016 in United Kingdom Patent Application No. 1408829.8.
- Examination Report dated Jun. 15, 2016 in United Kingdom Patent Application No. GB1511591.8.
- Examination Report dated Jan. 3, 2018 in United Kingdom Patent Application No. 1609740.4.
- Exhibit A filed Jul. 10, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 25 pp.
- Exhibits 12-32 filed Feb. 8, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 795 pp.
- Exhibits A-D filed Jan. 18, 2019 in the USDC District of Delaware, C.A. No. 1:18-cv-01800-RGA, 47 pp.
- Exhibits A-E filed Mar. 1, 2019 in the USDC District of Delaware, C.A. No. 18-1800 RGA, 75 pp.
- Extended European Search Report dated Jul. 17, 2015 in European Patent Application No. 13755319.4.
- Extended European Search Report dated Dec. 14, 2015 in European Patent Application No. 13754428.4.
- Extended European Search Report dated Feb. 8, 2018 in European Patent Application No. 17202409.3.
- Extended European Search Report dated Jun. 11, 2018 in European Patent Application No. 16740872.3.
- Extended European Search Report dated Mar. 22, 2019 in European Patent Application No. 18195513.9.
- Extended European Search Report dated May 6, 2021 in European Patent Application No. 20207621.2.
- Extended European Search Report dated May 28, 2021 in European Patent Application No. 20209777.0.
- Fan et al., "Parallel Genotyping of Human SNPs Using Generic High-density Oligonucleotide Tag Arrays," *Genome Research* 2000, 10, 853-860.
- Fan et al., "Microfluidic digital PCR enables rapid prenatal diagnosis of fetal aneuploidy," *Am Obstet Gynecol.* 2009, 200, 543e1-543e7.
- Fan, "Molecular counting: from noninvasive prenatal diagnostics to whole-genome haplotyping," Doctoral Dissertation, Stanford University 2010, 1-185.
- Fan et al., "Non-invasive Prenatal Measurement of the Fetal Genome," *Nature* 2012, 487(7407), 320- 324.
- Fan et al., "Combinatorial labeling of single cells for gene expression cytometry," *Science* 2015, 347(6222), 1258366-1258369.
- Feldhaus et al., "Oligonucleotide-conjugated beads for transdominant genetic experiments," *Nucleic Acids Res.* 2000, 28(2), 534-543.
- Final Office Action dated Sep. 1, 2015 for U.S. Appl. No. 14/540,029.
- Final Office Action dated Sep. 24, 2015 for U.S. Appl. No. 14/540,007.
- Final Office Action dated Oct. 6, 2015 in U.S. Appl. No. 14/540,018.
- Final Office Action dated Apr. 11, 2016 for U.S. Appl. No. 14/800,526.
- Final Office Action dated Jul. 20, 2016 for U.S. Appl. No. 14/281,706.
- Final Office Action dated Aug. 12, 2016 in U.S. Appl. No. 14/381,488.
- Final Office Action dated Feb. 13, 2017 in U.S. Appl. No. 14/381,488.
- Final Office Action dated May 8, 2017 in U.S. Appl. No. 15/224,460.
- Final Office Action dated Oct. 16, 2017 in U.S. Appl. No. 15/409,355.
- Final Office Action dated Nov. 16, 2017 in U.S. Appl. No. 14/381,488.
- Final Office Action dated Jan. 25, 2018 in U.S. Appl. No. 14/381,526.
- Final Office Action dated May 3, 2018 in U.S. Appl. No. 15/046,225.
- Final Office Action dated May 10, 2018 in U.S. Appl. No. 14/381,488.
- Final Office Action dated Jul. 5, 2018 in U.S. Appl. No. 15/004,618.
- Final Office Action dated Jul. 20, 2018 in U.S. Appl. No. 15/217,886.
- Final Office Action dated Nov. 16, 2018 in U.S. Appl. No. 15/134,967.
- Final Office Action dated Feb. 19, 2019 in U.S. Appl. No. 14/381,526.
- Final Office Action dated Mar. 1, 2019 in U.S. Appl. No. 16/012,584.
- Final Office Action dated Apr. 22, 2019 in U.S. Appl. No. 15/987,851.
- Final Office Action dated May 2, 2019 in U.S. Appl. No. 16/012,635.
- Final Office Action dated May 3, 2019 in U.S. Appl. No. 15/937,713.
- Final Office Action dated Sep. 18, 2019 in U.S. Appl. No. 15/055,407.
- Final Office Action dated Oct. 2, 2019 in U.S. Appl. No. 15/084,307.
- Final Office Action dated Dec. 4, 2019 in U.S. Appl. No. 15/596,364.
- Final Office Action dated Jan. 8, 2020 in U.S. Appl. No. 15/459,977.
- Final Office Action dated Jan. 16, 2020 in U.S. Appl. No. 16/012,584.
- Final Office Action dated Jan. 29, 2020 in U.S. Appl. No. 14/381,488.
- Final Office Action dated Feb. 4, 2020 in U.S. Appl. No. 15/715,028.
- Final Office Action dated Mar. 9, 2020 in U.S. Appl. No. 15/987,851.
- Final Office Action dated Apr. 28, 2020 in U.S. Appl. No. 15/134,967.
- Final Office Action dated Jun. 5, 2020 in U.S. Appl. No. 15/084,307.
- Final Office Action dated Aug. 19, 2020 in U.S. Appl. No. 15/875,816.
- Final Office Action dated Sep. 14, 2020 in U.S. Appl. No. 16/789,358.
- Final Office Action dated Sep. 22, 2020 in U.S. Appl. No. 16/789,311.
- Final Office Action dated Sep. 25, 2020 in U.S. Appl. No. 15/055,407.
- Final Office Action dated Dec. 7, 2020 in U.S. Appl. No. 16/012,584.
- Final Office Action dated Feb. 11, 2021 in U.S. Appl. No. 15/134,967.
- Final Office Action dated Mar. 16, 2021 in U.S. Appl. No. 15/715,028.
- Final Office Action dated Mar. 25, 2021 in U.S. Appl. No. 16/374,626.
- Final Office Action dated Jun. 15, 2021 in U.S. Appl. No. 15/084,307.
- Final Office Action dated Jul. 15, 2021 in U.S. Appl. No. 16/836,750.
- Final Office Action dated Aug. 10, 2021 in U.S. Appl. No. 16/012,584.
- Final Office Action dated Aug. 27, 2021 in U.S. Appl. No. 15/055,407.
- Final Office Action dated Sep. 24, 2021 in U.S. Appl. No. 16/788,743.

(56)

**References Cited**

## OTHER PUBLICATIONS

- Final Office Action dated Oct. 1, 2021 in U.S. Appl. No. 16/677,012.
- First Action Interview Pilot Program Pre-Interview Communication dated Oct. 15, 2018 in U.S. Appl. No. 15/987,851.
- First Action Interview Office Action Summary dated Jan. 25, 2019 in U.S. Appl. No. 15/987,851.
- Fitzgerald and Grivel, "A Universal Nanoparticle Cell Secretion Capture Assay," *Cytometry Part A* 2012, 83A(2), 205-211.
- Flanigon et al., "Multiplex protein detection with DNA readout via mass spectrometry," *N Biotechnol.* 2013, 30(2), 153-158.
- Forster et al., "A human gut bacterial genome and culture collection for improved metagenomic analyses," *Nature Biotechnology* 2019, 37, 186-192.
- Fox-Walsh et al., "A multiplex RNA-seq strategy to profile poly(A+) RNA: application to analysis of transcription response and 3' end formation," *Genomics* 2011, 98, 266-721.
- Fu et al., "Counting individual DNA molecules by the stochastic attachment of diverse labels," *Proc Natl Acad Sci* 2011, 108(22), 9026-9031.
- Fu et al., Digital Encoding of Cellular mRNAs Enabling Precise and Absolute Gene Expression Measurement by Single-Molecule Counting. *Anal Chem.* 2014, 86, 2867-2870.
- Fu et al., "Molecular indexing enables quantitative targeted RNA sequencing and reveals poor efficiencies in standard library preparation," *PNAS* 2014, 111(5), 1891-1896.
- GenBank Accession No. NM\_000518.5 for *Homo sapiens* hemoglobin subunit beta (HBB), mRNA. Mar. 22, 2021 [online], [retrieved on Apr. 27, 2021], retrieved from the Internet: <URL: www.ncbi.nlm.nih.gov/nucleotide/NM\_000518.5?report=Genbank (Year: 2021).
- Gerry et al., "Universal DNA Microarray Method for Multiplex Detection of Low Abundance Point Mutations," *Journal of Molecular Biology* 1999, 292, 251-262.
- Gertz et al., "Transposase mediated construction of RNA-seq libraries," *Genome Research* 2012, 22, 134-141.
- Gillespie, "Exact Stochastic Simulation of Coupled Chemical Reactions," *Journal of Physical Chemistry* 1977, 81(25), 2340-2361.
- Gong et al., "Massively parallel detection of gene expression in single cells using subnanolitre wells," *Lab Chip* 2010, 10, 2334-2337.
- Gong et al., "Simple Method Prepare Oligonucleotide-Conjugated Antibodies and Its Application in Multiplex Protein Detection in Single Cells," *Bioconjugate Chem.* 2016, 27, 217-225.
- Grant et al., "SNP genotyping on a genome-wide amplified DOP-PCR template," *Nucleic Acids Res* 2002, 30(22), e25, 1-6.
- Gratton et al., "Cell-permeable peptides improve cellular uptake and therapeutic gene delivery of replication-deficient viruses in cells and in vivo," *Nature Medicine* 2003, 9(3), 357-362.
- Grounds for Opposition dated Jul. 21, 2016 and filed in European Patent 2414548B1.
- Gu et al., "Complete workflow for detection of low frequency somatic mutations from cell-free DNA using Ion Torrent™ platforms," Conference Poster, AACR 107th Annual Meeting, Apr. 16-20, 2016, 1 p.
- Gu et al., "Depletion of abundant sequences by hybridization (DSH): using Cas9 to remove unwanted high-abundance species in sequencing libraries and molecular counting applications," *Genome Biology* 2016, 17(41) 1-13.
- Gunderson et al., "Decoding Randomly Ordered DNA Arrays," *Genome Research* 2004, 14, 870-877.
- Gundry et al., "Direct, genome-wide assessment of DNA mutations in single cells," *Nucleic Acids Research* 2011, 40(5), 2032-2040.
- Gundry et al., "Direct mutation analysis by high-throughput sequencing: from germline to low-abundant, somatic variants," *Mutat Res.* 2012, 729(1-2), 1-15.
- Hacia et al., "Determination of ancestral alleles for human single-nucleotide polymorphisms using high-density oligonucleotide arrays," *Nature Genetics* 1999, 22, 164-167.
- Haff, "Improved Quantitative PCR Using Nested Primers," *PCR Methods and Applications* 1994, 3, 332-337.
- Hamady et al., "Error-correcting barcoded primers for pyrosequencing hundreds of samples in multiplex," *Nat Methods* 2008, 5(3), 235-237.
- Han et al., "An approach to multiplexing an immunosorbent assay with antibody-oligonucleotide conjugates," *Bioconjug Chem.* 2010, 21(12), 2190-2196.
- Harbers, "The current status of cDNA cloning," *Genomics* 2008, 91, 232-242.
- Harrington et al., Cross-sectional characterization of HIV-1 env compartmentalization in cerebrospinal fluid over the full disease course, *AIDS* 2009, 23(8), 907-915.
- Hartmann, "Gene expression profiling of single cells on large-scale oligonucleotide arrays," *Nucleic Acids Research*, (Oct. 2006) vol. 34, No. 21, p. e143, 1-12.
- Hashimshony et al., "CEL-Seq: Single-Cell RNA-Seq by Multiplexed Linear Amplification," *Cell Rep.* 2012, 2(3), 666-673.
- Hensel et al., "Simultaneous identification of bacterial virulence genes by negative selection," *Science* 1995, 269(5222), 400-403.
- Hiatt et al., "Parallel, tag-directed assembly of locally derived short sequence reads," *Nat Methods* 2010, 7(2), 119-122.
- Hiatt et al., "Single molecule molecular inversion probes for targeted, high-accuracy detection of low-frequency variation," *Genome Res.* 2013, 23(5), 843-854.
- Holcomb et al., "Abstract 1853: Single-cell multiplexed profiling of protein-level changes induced by EGFR inhibitor gefitinib," *Cancer Res* 2016, 76(14 Suppl), Abstract 1853.
- Hollas et al., "A stochastic approach to count RNA molecules using DNA sequencing methods," *Algorithms in Bioinformatics. WABI 2003, Lecture Notes in Computer Science*, 2812, 55-62.
- How many species of bacteria are there? Wisegeek.org, accessed Jan. 21, 2014, 2 pp.
- Hu et al., "Dissecting Cell-Type Composition and Activity-Dependent Transcriptional State in Mammalian Brains by Massively Parallel Single-Nucleus RNA-Seq," *Molecular Cell* 2017, 68, 1006-1015.
- Hu et al., "Single Cell Multi-Omics Technology: Methodology and Application," *Frontiers in Cell and Developmental Biology* 2018, 6(28), 1-13.
- Hug et al., Measure of the Number of Molecular of a Single mRNA Species in a Complex mRNA Preparation, *Journal of Theoretical Biology* 2003, 221, 615-624.
- Ingolia et al., Genome-Wide Analysis in Vivo of Translation with Nucleotide Resolution Using Ribosome Profiling, *Science* 2009, 324(5924), 218-223.
- International Search Report and Written Opinion dated May 7, 2012 for PCT Application No. PCT/IB2011/003160.
- International Search Report and Written Opinion dated Jun. 6, 2012 in PCT Application No. PCT/US2011/065291.
- International Search Report and Written Opinion dated Jun. 14, 2013 in PCT Application No. PCT/US2013/028103.
- International Search Report and Written Opinion dated Aug. 16, 2013 for PCT Application No. PCT/US2013/027891.
- International Search Report and Written Opinion dated Dec. 19, 2014 in PCT Application No. PCT/US2014/059542.
- International Search Report and Written Opinion dated Feb. 3, 2015 in PCT Application No. PCT/US2014/053301.
- International Search Report and Written Opinion dated May 3, 2016 in PCT Application No. PCT/US2016/018354.
- International Search Report and Written Opinion dated Jun. 9, 2016 in PCT Application No. PCT/US2016/022712.
- International Search Report and Written Opinion dated Jun. 17, 2016 in PCT Application No. PCT/US2016/019962.
- International Search Report and Written Opinion dated Jun. 20, 2016 in PCT Application No. PCT/US2016/014612.
- International Search Report and Written Opinion dated Aug. 9, 2016 in PCT Application No. PCT/US2016/019971.
- International Search Report and Written Opinion dated Sep. 27, 2016 in PCT Application No. PCT/US2016/034473.
- International Search Report and Written Opinion dated Sep. 28, 2016 in PCT Application No. PCT/US2016/028694.
- International Search Report and Written Opinion dated Dec. 5, 2016 in PCT Application No. PCT/US2016/024783.

(56)

**References Cited****OTHER PUBLICATIONS**

- International Search Report and Written Opinion dated Jan. 31, 2017 in PCT Application No. PCT/US2016/050694.
- International Search Report and Written Opinion dated Aug. 7, 2017 in PCT Application No. PCT/US2017/034576.
- International Search Report and Written Opinion dated Sep. 8, 2017 in PCT Application No. PCT/US2017/030097.
- International Search Report and Written Opinion dated Mar. 20, 2018 in PCT Application No. PCT/US2017/053331.
- International Search Report and Written Opinion dated Mar. 28, 2018 in PCT Application No. PCT/US2018/014385.
- International Search Report and Written Opinion dated Jul. 16, 2018 in PCT Application No. PCT/US2018/024602.
- International Search Report and Written Opinion dated Jun. 24, 2019 in PCT Application No. PCT/US2019/030175.
- International Search Report and Written Opinion dated Oct. 8, 2019 in PCT Application No. PCT/US2019/043949.
- International Search Report and Written Opinion dated Oct. 16, 2019 in PCT Application No. PCT/US2019/030245.
- International Search Report and Written Opinion dated Nov. 27, 2019 in PCT Application No. PCT/US2019/046549.
- International Search Report and Written Opinion dated Dec. 4, 2019 in PCT Application No. PCT/US2019/053868.
- International Search Report and Written Opinion dated Jan. 27, 2020 in PCT Application No. PCT/US2019/048179.
- International Search Report and Written Opinion dated Mar. 30, 2020 in PCT Application No. PCT/US2019/060243.
- International Search Report and Written Opinion dated Mar. 30, 2020 in PCT Application No. PCT/US2019/065237.
- International Search Report and Written Opinion dated May 18, 2020 in PCT Application No. PCT/US2020/014339.
- International Search Report and Written Opinion dated Jun. 30, 2020 in PCT Application No. PCT/US2020/017890.
- International Search Report and Written Opinion dated Nov. 12, 2020 in PCT Application No. PCT/US2020/042880.
- International Search Report and Written Opinion dated Jan. 19, 2021 in PCT Application No. PCT/US2020/059419.
- International Search Report and Written Opinion dated Apr. 9, 2021 in PCT Application No. PCT/US2021/013137.
- International Search Report and Written Opinion dated Apr. 21, 2021 in PCT Application No. PCT/US2021/015571.
- International Search Report and Written Opinion dated May 4, 2021 in PCT Application No. PCT/US2021/013109.
- International Search Report and Written Opinion dated May 11, 2021 in PCT Application No. PCT/US2021/013748.
- International Search Report and Written Opinion dated Jul. 15, 2021 in PCT Application No. PCT/US2021/019475.
- International Search Report and Written Opinion dated Jul. 20, 2021 in PCT Application No. PCT/US2021/015898.
- International Search Report and Written Opinion dated Aug. 31, 2021 in PCT Application No. PCT/US2021/035270.
- International Search Report and Written Opinion dated Sep. 22, 2021, in PCT Application No. PCT/US2021/013747.
- International Search Report and Written Opinion dated Sep. 27, 2021, in PCT Application No. PCT/US2021/013747.
- International Search Report and Written Opinion dated Oct. 12, 2021, in PCT Application No. PCT/US2021/041327.
- Invitation to Pay Fees dated May 16, 2018 in PCT Application No. PCT/US2018/024602.
- Invitation to Pay Fees dated Nov. 26, 2019 in PCT Application No. PCT/US2019/048179.
- Invitation to Pay Fees dated May 25, 2021 in PCT Application No. PCT/US2021/01598.
- Invitation to Pay Additional Search Fees dated May 7, 2020 in PCT Application No. PCT/US2020/017890.
- Invitation to Pay Additional Search Fees dated Sep. 8, 2021 in PCT Application No. PCT/US2021/032319.
- Invitation to Provide Informal Clarification dated Jun. 9, 2021 in PCT Application No. PCT/US2021/019475.
- Invitation to Respond to Written Opinion dated May 26, 2017 in Singapore Patent Application No. 11201405274W.
- Islam et al., "Characterization of the single-cell transcriptional landscape by highly multiplex RNA-seq," *Genome Research* 2011, 21, 1160-1167.
- Islam et al., "Highly multiplexed and strand specific single-cell RNA 5' end sequencing," *Nature Protocols* 2012, 7(5), 813-828.
- Islam et al., "Quantitative single-cell RNA-seq with unique molecular identifiers," *Nature Methods* 2014, 11(2), 163-168.
- Jabara, "Capturing the cloud: High throughput sequencing of multiple individual genomes from retroviral population," *Biology Lunch Bunch Series, Training Initiatives in Biomedical & Biological Sciences* of the University of North Carolina at Chapel Hill 2010.
- Jabara et al., "Accurate sampling and deep sequencing of the HIV-1 protease gene using a Primer ID," *PNAS* 2011, 108(50), 20166-20171.
- Jason J. Rawnsley of Richards, Layton and Finger, P.A., Entry of Appearance dated Jan. 18, 2019 in the USDC District of Delaware, C.A. No. 18-1800-RGA, 1 pp.
- Janeway et al., "Structural variation in immunoglobulin constant regions," *Immunology: The Immune System in Health and Disease* 1999, 101-103.
- Jiang et al., "Synthetic spike-in standards for RNA-seq experiments," *Genome Res.* 2011, 21, 1543-1551.
- Joint Stipulation and Order to Extend Time to Respond to Plaintiff's First Amended Complaint, dated Feb. 21, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 2 pp.
- Joint Stipulation and Order to Request Extended Time to File Opposition to Defendant's Motion to Dismiss dated, Mar. 8, 2019 in the USDC District of Delaware, C.A. No. 18-1800 RGA, 2 pp.
- Joint Stipulation and Order to Request Extended Time to Submit a proposed Protective Order, dated Jun. 7, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 1 pp.
- Joint Stipulation and Order to Extended Time to Submit Agreed Document Production Protocol, filed Jun. 28, 2019 in the USDC for the District of Delaware, C.A. 18-1800 (RGA), 1 pp.
- Joint Stipulation and Order to Request Extended Time to Submit Agreed Document Production Protocol, dated Jul. 11, 2019 in the USDC for the District of Delaware, C.A. 18-1800 (RGA), 1 pp.
- Junker et al., "Single-Cell Transcriptomics Enters the Age of Mass Production," *Molecular Cell* 2015, 58, 563-564.
- Kanagawa, "Bias and artifacts in multi-template polymerase chain reactions (PCR)," *Journal of Bioscience and Bioengineering* 2003, 96(4), 317-323.
- Kang et al., "Targeted sequencing with enrichment PCR: a novel diagnostic method for the detection of EGFR mutations," *Oncotarget* 2015, 6(15), 13742-13749.
- Kang et al., "Application of multi-omics in single cells," *Ann Biotechnol.* 2018, 2(1007), 1-8.
- Karrer et al., "In situ isolation of mRNA from individual plant cells: creation of cell-specific cDNA libraries," *Proc. Natl. Acad. Sci. USA* 1995, 92, 3814-3818.
- Kausch et al., "Organelle Isolation by Magnetic Immunoabsorption," *BioTechniques* 1999, 26(2), 336-343.
- Kebischull et al., "Sources of PCR-induced distortions in high-throughput sequencing data sets," *Nucleic Acids Research* 2015, 1-15.
- Keys et al., Primer ID Informs Next-Generation Sequencing Platforms and Reveals Preexisting DrugResistance Mutations in the HIV-1 Reverse Transcriptase Coding Domain, *AIDS Research and Human Retroviruses* 2015, 31(6), 658-668.
- Kim et al., Polony Multiplex Analysis of Gene Expression (PMAGE) in Mouse Hypertrophic Cardiomyopathy, *Science* 2007, 316(5830), 1481-1484.
- Kinde et al., "Detection and quantification of rare mutations with massively parallel sequencing," *Proc. Natl. Acad. Sci.* 2011, 108(23), 9530-0535.
- Kirsebom et al., "Stimuli-Responsive Polymers in the 21st Century: Elaborated Architecture to Achieve High Sensitivity, Fast Response, and Robust Behavior," *Journal of Polymer Science: Part B: Polymer Physics* 2011, 49, 173-178.

(56)

**References Cited****OTHER PUBLICATIONS**

- Kivioja et al., "Counting absolute numbers of molecules using unique molecular identifiers," *Nature Proceedings* 2011, 1-18.
- Klein et al., Droplet Barcoding for Single-Cell Transcriptomics Applied to Embryonic Stem Cells, *Cell* 2015, 161, 1187-1201.
- Ko et al., "RNA-conjugated template-switching RT-PCR method for generating an *Escherichia coli* cDNA library for small RNAs," *Journal of Microbiological Methods* 2006, 64, 297-304.
- Koboldt et al., "VarScan: variant detection in massively parallel sequencing of individual and pooled samples," *Bioinformatics* 2009, 25(17), 2283-2285.
- Kolodziejczyk et al., "The Technology and Biology of Single-Cell RNA Sequencing," *Molecular Cell* 2015, 58, 610-620.
- Konig et al., "iCLIP reveals the function of hnRNP particles in splicing at individual nucleotide resolution," *Nature Structural & Molecular Biology* 2010, 17(7), 909-916.
- Kooiker & Xue, "cDNA Library Preparation," *Cereal Genomics* 2013, 1099, 29-40.
- Kotake et al., "A simple nested RT-PCR method for quantitation of the relative amounts of multiple cytokine mRNAs in small tissue samples," *Journal of Immunological Methods* 1996, 199, 193-203.
- Kozarewa & Turner, "96-Plex Molecular Barcoding for the Illumina Genome Analyzer," *High-Throughput Next Generation Sequencing: Methods in Molecular Biology (Methods and Applications)* 2011, 733, 24 pp. DOI: 10.1007/978-1-61779-089-8\_20.
- Kozlov et al., "A high-complexity, multiplexed solution-phase assay for profiling protease activity on microarrays," *Comb Chem High Throughput Screen* 2008, 11(1), 24-35.
- Kurimoto et al., "An improved single-cell cDNA amplification method for efficient high-density oligonucleotide microarray analysis," *Nucleic Acids Res.* 2006, 34(5), e42, 1-17.
- Kurimoto et al., "Global single-cell cDNA amplification to provide a template for representative high-density oligonucleotide microarray analysis," *Nature Protocols* 2007, 2(3), 739-752.
- Lamble et al., "Improved workflows for high throughput library preparation using the transposome-based nextera system," *BMC Biotechnology* 2013, 13, 104, 1-10.
- Lan et al., "Droplet barcoding for massively parallel single-molecule deep sequencing," *Nature Communications* 2016, 7(11784), in 10 pages.
- Larson et al., "A single molecule view of gene expression," *Trends Cell Biol.* 2009, 19(11), 630-637.
- Lass-Napiorkowska et al., "Detection methodology based on target molecule-induced sequence-specific binding to a single-stranded oligonucleotide," *Anal Chem.* 2012, 84(7), 3382-3389.
- Leamon et al., A massively parallel PicoTiterPlate based platform for discrete picoliter-scale polymerase chain reactions, *Electrophoresis* 2003, 24, 3769-3777.
- Lee et al., "Large-scale arrays of picolitre chambers for single-cell analysis of large cell populations," *Lab Chip* 2010, 10, 2952-2958.
- Lee et al., "Highly Multiplexed Subcellular RNA Sequencing in Situ," *Science* 2014, 343, 1360-1363.
- Lee et al., "Universal process-inert encoding architecture for polymer microparticles," *Nature Materials* 2014, 13(5), 524-529.
- Letter regarding the opposition procedure dated Jul. 22, 2015 for European Patent Application No. 11810645.9.
- Letter to Judge Richard G. Andrews Requesting a Rule 16 Conference, dated Apr. 15, 2019 in the USDC for the District of Delaware, C.A. 18-1800 (RGA), 1 pp.
- Letter to Judge Andrews regarding Agreement on Proposed Scheduling Order, dated May 7, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 1 pp.
- Letter to Judge Andrews regarding Notice of Supplemental Authority, dated Jul. 10, 2019 in the USDC for the District of Delaware, C.A. 18-1800(RGA), 2pp.
- Lin et al., "Self-Assembled Combinatorial Encoding Nanoarrays for Multiplexed Biosensin," *Nano Lett.* 2007, 7 (2), 507-512.
- Liu et al., "Single-cell transcriptome sequencing: recent advances and remaining challenges," *F1000Research* 2016, 5(F1000 Faculty Rev)(182), 1-9.
- Livingstone, "rRNA depletion, poly(A) enrichment, or exonuclease treatment?" *Tebu-Bio Blog* 2015, in 1 page.
- Lizardi et al., "Mutation detection and single-molecule counting using isothermal rolling-circle amplification," *Nat Genet.* 1998, 19, 225-232.
- Lockhart et al., "Expression monitoring by hybridization to high-density oligonucleotide arrays," *Nature Biotechnology* 1996, 14, 1675-1680.
- Lovatt et al., "Transcriptome *in vivo* analysis (TIVA) of spatially defined single cells in live tissue," *Nat Methods* 2014, 11(2), 190-196.
- Loy et al., "A rapid library preparation method with custom assay designs for detection of variants at 0.1% allelic frequency in liquid biopsy samples," *ThermoFisher Scientific*, Oct. 2, 2018, 1 p.
- Lucito et al., "Representational Oligonucleotide Microarray Analysis: A High-Resolution Method to Detect Genome Copy Number Variation," *Genome Research* 2003, 13, 2291-2305.
- Lundberg et al., "Practical innovations for high-throughput amplicon sequencing," *Nature Methods* 2013, 10(10), 999-1007.
- Lundberg et al., "Supplementary Information for: Practical innovations for high-throughput amplicon sequencing," *Nature Methods* 2013, 1-24.
- Maamar et al., "Noise in Gene Expression Determines Cell Fate in *Bacillus subtilis*," *Science* 2007, 317, 526-529.
- Macaulay et al., "Single Cell Genomics: Advances and Future Perspectives," *PLoS Genetics* 2014, 10(1), 1-9.
- Macaulay et al., "G&T-seq: parallel sequencing of single-cell genomes and transcriptomes," *Nature Methods* 2015, 1-7.
- Macosko et al., "Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets," *Cell* 2015, 161, 1202-1214.
- Maeda et al., "Development of a DNA barcode tagging method for monitoring dynamic changes in gene expression by using an ultra high-throughput sequencer," *BioTechniques* 2008, 45(1), 95-97.
- Mair et al., "A Targeted Multi-omic Analysis Approach Measures Protein Expression and Low-Abundance Transcripts on the Single-Cell Level," *Cell Reports* 2020, 31(1), 107499, in 20 pages.
- Makrigiorgos et al., "A PCR-Based amplification method retaining quantities difference between two complex genomes," *Nature Biotech* 2002, 20(9), 936-939.
- Marcus et al., "Microfluidic single-cell mRNA isolation and analysis," *Anal Chem.* 2006, 78, 3084-3089.
- Mardis, "Next-generation DNA sequencing methods", *Annu. Rev. Genomics Hum. Genet.* 2008, 9, 387-402.
- Marguerat et al., "Next-generation sequencing: applications beyond genomes," *Biochem. Soc. Trans.* 2008, 36(5), 1091-1096.
- Marguiles et al., "Genome sequencing in microfabricated high-density picolitre reactors," *Nature* 2005, 437, 376-380.
- Martinez et al., "A microfluidic approach to encapsulate living cells in uniform alginate hydrogel microparticles," *Macromol. Biosci* 2012, 12, 946-951.
- Massachusetts General Hospital, Overview of Illumina Chemistry, <http://nextgen.mgh.harvard.edu/IlluminaChemistry.html>, downloaded Jan. 28, 2020, 2 pp.
- McCloskey et al., "Encoding PCR products with batch-stamps and barcodes," *Biochem Genet.* 2007, 45(11-12), 761-767.
- Medvedev et al., "Detecting copy No. variation with mated short reads," *Genome Res.* 2010, 20, 1613-1622.
- Mei et al., "Identification of recurrent regions of Copy-No. Variants across multiple individuals," *BMC Bioinformatics* 2010, 11, 147, 1-14.
- Merriam-Webster, definition of associate: <http://merriam-webster.com/dictionary/associate>, accessed Apr. 5, 2016.
- Meyer et al., "Parallel tagged sequencing on the 454 platform," *Nature Protocols* 2008, 3(2), 267-278.
- Miller et al., "Directed evolution by *in vitro* compartmentalization," *Nature Methods* 2006, 3(7), 561-570.
- Miner et al., "Molecular barcodes detect redundancy and contamination in hairpin-bisulfite PCR," *Nucleic Acids Research* 2004, 32(17), e135, 1-4.
- Mortazavi et al., "Mapping and quantifying mammalian transcriptomes by RNA-Seq," *Nat. Methods* 2008, 5(7), 621-628.

(56)

**References Cited****OTHER PUBLICATIONS**

- Motion and Order for Admission Pro Hac Vice Pursuant to Local Rule 83.5, dated Jan. 24, 2019 in the USDC District of Delaware, C.A. No. 18-1800-RGA, 7 pp.
- Nadai et al., Protocol for nearly full-length sequencing of HIV-1 RNA from plasma, *PLoS One* 2008, 3(1), e1420, 1-6.
- Nagai et al., "Development of a microchamber array for picoletter PCR," *Anal. Chem.* 2001, 73, 1043-1047.
- Navin et al., "The first five years of single-cell cancer genomics and beyond," *Genome Research* 2015, 25, 1499-1507.
- New COVID-19 Variants, Centers for Disease Control and Prevention 2021, accessed Jan. 21, 2021, 3 pp.
- Newell et al., Cytometry by time-of-flight shows combinatorial cytokine expression and virus-specific cell niches within a continuum of CD8+ T cell phenotypes. *Immunity* 2012, 36(1), 142-152.
- Non-Final Office Action dated Oct. 3, 2013 in U.S. Appl. No. 12/969,581.
- Non-Final Office Action dated Feb. 18, 2015 for U.S. Appl. No. 14/540,007.
- Non-Final Office Action dated Feb. 26, 2015 for U.S. Appl. No. 14/540,029.
- Non-Final Office Action dated Mar. 19, 2015 in U.S. Appl. No. 14/540,018.
- Non-Final Office Action dated May 7, 2015 for U.S. Appl. No. 13/327,526.
- Non-Final Office Action dated Dec. 3, 2015 for U.S. Appl. No. 14/281,706.
- Non-Final Office Action dated Dec. 31, 2015 for U.S. Appl. No. 14/800,526.
- Non-Final Office Action dated Apr. 11, 2016 in U.S. Appl. No. 14/472,363.
- Non-Final Office Action dated May 10, 2016 in U.S. Appl. No. 14/381,488.
- Non-Final Office Action dated May 13, 2016 in U.S. Appl. No. 14/508,911.
- Non-Final Office Action dated Aug. 17, 2016 for U.S. Appl. No. 14/800,526.
- Non-Final Office Action dated Sep. 26, 2016 in U.S. Appl. No. 15/167,807.
- Non-Final Office Action dated Oct. 11, 2016 in U.S. Appl. No. 15/224,460.
- Non-Final Office Action dated Jan. 19, 2017 in U.S. Appl. No. 15/055,445.
- Non-Final Office Action dated Mar. 24, 2017 in U.S. Appl. No. 15/409,355.
- Non-Final Office Action dated Jun. 2, 2017 in U.S. Appl. No. 14/381,526.
- Non-Final Office Action dated Jun. 7, 2017 in U.S. Appl. No. 14/381,488.
- Non-Final Office Action dated Jul. 28, 2017 in U.S. Appl. No. 14/975,441.
- Non-Final Office Action dated Sep. 8, 2017 in U.S. Appl. No. 15/046,225.
- Non-Final Office Action dated Sep. 8, 2017 in U.S. Appl. No. 15/134,967.
- Non-Final Office Action dated Nov. 1, 2017 in U.S. Appl. No. 15/667,125.
- Non-Final Office Action dated Nov. 9, 2017 in U.S. Appl. No. 15/004,618.
- Non-Final Office Action dated Jan. 9, 2018 in U.S. Appl. No. 15/217,896.
- Non-Final Office Action dated Jan. 12, 2018 in U.S. Appl. No. 15/217,886.
- Non-Final Office Action dated Mar. 8, 2018 in U.S. Appl. No. 15/608,780.
- Non-Final Office Action dated Apr. 6, 2018 in U.S. Appl. No. 15/603,239.
- Non-Final Office Action dated Jul. 25, 2018 in U.S. Appl. No. 15/108,268.
- Non-Final Office Action dated Oct. 4, 2018 in U.S. Appl. No. 15/260,106.
- Non-Final Office Action dated Oct. 25, 2018 in U.S. Appl. No. 16/012,584.
- Non-Final Office Action dated Nov. 5, 2018 in U.S. Appl. No. 16/038,790.
- Non-Final Office Action dated Nov. 26, 2018 in U.S. Appl. No. 15/937,713.
- Non-Final Office Action dated Jan. 7, 2019 in U.S. Appl. No. 15/055,407.
- Non-Final Office Action dated Jan. 14, 2019 in U.S. Appl. No. 16/219,553.
- Non-Final Office Action dated Mar. 19, 2019 in U.S. Appl. No. 15/046,225.
- Non-Final Office Action dated May 15, 2019 in U.S. Appl. No. 15/084,307.
- Non-Final Office Action dated May 23, 2019 in U.S. Appl. No. 15/459,977.
- Non-Final Office Action dated Jun. 17, 2019 in U.S. Appl. No. 14/381,488.
- Non-Final Office Action dated Jul. 9, 2019 in U.S. Appl. No. 15/596,364.
- Non-Final Office Action dated Aug. 20, 2019 for U.S. Appl. No. 15/715,028.
- Non-Final Office Action dated Sep. 18, 2019 in U.S. Appl. No. 16/194,819.
- Non-Final Office Action dated Nov. 29, 2019 in U.S. Appl. No. 15/937,713.
- Non-Final Office Action dated Jan. 17, 2020 in U.S. Appl. No. 15/084,307.
- Non-Final Office Action dated Feb. 5, 2020 in U.S. Appl. No. 15/875,816.
- Non-Final Office Action dated Mar. 12, 2020 in U.S. Appl. No. 16/789,358.
- Non-Final Office Action dated Mar. 17, 2020 in U.S. Appl. No. 15/055,407.
- Non-Final Office Action dated Mar. 26, 2020 in U.S. Appl. No. 16/012,635.
- Non-Final Office Action dated Mar. 26, 2020 in U.S. Appl. No. 16/789,311.
- Non-Final Office Action dated Jun. 8, 2020 in U.S. Appl. No. 15/715,028.
- Non-Final Office Action dated Aug. 4, 2020 in U.S. Appl. No. 15/459,977.
- Non-Final Office Action dated Aug. 19, 2020 in U.S. Appl. No. 16/374,626.
- Non-Final Office Action dated Aug. 25, 2020 in U.S. Appl. No. 14/381,488.
- Non-Final Office Action dated Dec. 4, 2020 in U.S. Appl. No. 16/677,012.
- Non-Final Office Action dated Dec. 9, 2020 in U.S. Appl. No. 16/788,743.
- Non-Final Office Action dated Jan. 19, 2021 in U.S. Appl. No. 16/836,750.
- Non-Final Office Action dated Feb. 2, 2021 in U.S. Appl. No. 16/535,080.
- Non-Final Office Action dated Feb. 25, 2021 in U.S. Appl. No. 15/055,407.
- Non-Final Office Action dated Feb. 25, 2021 in U.S. Appl. No. 15/084,307.
- Non-Final Office Action dated Mar. 29, 2021 in U.S. Appl. No. 16/789,358.
- Non-Final Office Action dated Apr. 14, 2021 in U.S. Appl. No. 16/789,311.
- Non-Final Office Action dated Apr. 20, 2021 in U.S. Appl. No. 15/875,816.
- Non-Final Office Action dated May 18, 2021 in U.S. Appl. No. 16/535,080.
- Non-Final Office Action dated Jun. 9, 2021 in U.S. Appl. No. 16/588,405.
- Non-Final Office Action dated Aug. 17, 2021 in U.S. Appl. No. 16/551,620.

(56)

**References Cited**

## OTHER PUBLICATIONS

- Non-Final Office Action dated Aug. 19, 2021 in U.S. Appl. No. 16/781,814.
- Non-Final Office Action dated Aug. 31, 2021 in U.S. Appl. No. 15/715,028.
- Non-Final Office Action dated Sep. 1, 2021 in U.S. Appl. No. 16/789,358.
- Non-Final Office Action dated Sep. 14, 2021 in U.S. Appl. No. 16/707,780.
- Non-Final Office Action dated Sep. 28, 2021 in U.S. Appl. No. 16/400,885.
- Non-Final Office Action dated Sep. 30, 2021 in U.S. Appl. No. 16/374,626.
- Non-Final Office Action dated Oct. 1, 2021 in U.S. Appl. No. 16/677,012.
- Non-Final Office Action dated Oct. 8, 2021 in U.S. Appl. No. 16/400,866.
- Notice of Allowability dated Jun. 19, 2014 for U.S. Appl. No. 12/969,581.
- Notice of Allowance dated Mar. 21, 2014 for U.S. Appl. No. 12/969,581.
- Notice of Allowance dated Aug. 22, 2014 for U.S. Appl. No. 12/969,581.
- Notice of Allowance dated Dec. 15, 2015 for U.S. Appl. No. 14/540,007.
- Notice of Allowance dated Dec. 21, 2015 in U.S. Appl. No. 14/540,018.
- Notice of Allowance dated Jan. 21, 2016 for U.S. Appl. No. 13/327,526.
- Notice of Allowance dated Jan. 9, 2019 in U.S. Appl. No. 15/603,239.
- Notice of Allowance dated Mar. 20, 2019 in U.S. Appl. No. 16/219,553.
- Notice of Allowance dated Mar. 21, 2019 in U.S. Appl. No. 15/993,468.
- Notice of Allowance dated May 28, 2019 in U.S. Appl. No. 16/219,553.
- Notice of Allowance dated Sep. 24, 2019 in U.S. Appl. No. 15/217,886.
- Notice of Allowance dated Nov. 11, 2019 in Japanese Patent Application No. 2017-245295.
- Notice of Allowance dated Nov. 29, 2019 in U.S. Appl. No. 16/012,635.
- Notice of Allowance dated Dec. 27, 2019 in U.S. Appl. No. 15/260,106.
- Notice of Allowance dated Mar. 5, 2020 in U.S. Appl. No. 15/217,886.
- Notice of Allowance dated Mar. 27, 2020 in U.S. Appl. No. 15/596,364.
- Notice of Allowance dated Mar. 30, 2020 in U.S. Appl. No. 15/937,713.
- Notice of Allowance dated Apr. 15, 2020 in U.S. Appl. No. 16/012,635.
- Notice of Allowance dated Sep. 23, 2020 in Korean Patent Application No. 10-2016-7008144.
- Notice of Allowance dated Oct. 29, 2020 in U.S. Appl. No. 15/987,851.
- Notice of Allowance dated Jan. 13, 2021 in U.S. Appl. No. 14/381,488.
- Notice of Allowance dated Jan. 13, 2021 in U.S. Appl. No. 15/459,977.
- Notice of Allowance dated Apr. 26, 2021 in Japanese Patent Application No. 2019-014564.
- Notice of Allowance dated Jun. 10, 2021 in Chinese Patent Application No. 2018800377201.
- Notice of Allowance dated Aug. 16, 2021 in Japanese Patent Application No. 2018-512152.
- Notice of Allowance dated Sep. 10, 2021 in U.S. Appl. No. 16/535,080.
- Notice, Consent, and Reference of a Civil Action to a Magistrate Judge (Rule 73.1), filed Nov. 15, 2018 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 3 pp.
- Notice of Opposition dated Jul. 9, 2015 for European Patent Application No. 11810645.9.
- Notice of Opposition dated Jul. 27, 2016 for European Patent Application No. 10762102.1.
- Notice of Reasons for Rejection dated Dec. 28, 2016 in Japanese Patent Application No. 2014-558975.
- Notice of Reasons for Rejection dated Apr. 2, 2018 in Japanese Patent Application No. 2014-558975.
- Notice of Reasons for Rejection dated Jul. 30, 2018 in Japanese Patent Application No. 2016-537867.
- Notice of Reasons for Rejection dated Aug. 31, 2018 in Japanese Patent Application No. 2016-520632.
- Notice of Reasons for Rejection dated Dec. 5, 2018 in Japanese Patent Application No. 2017-245295.
- Notice of Reason for Rejection dated Nov. 21, 2019 in Korean Patent Application No. 10-2016-7008144.
- Notice of Reasons for Rejection dated Feb. 25, 2020 in Japanese Patent Application No. 2019-014564.
- Notice of Reasons for Rejection dated May 11, 2020 in Japanese Patent Application No. 2017-549390.
- Notice of Service of Disclosures to Opposing Counsel, dated Jun. 10, 2019 in the USDC for the District of Delaware, C.A. 18-1800 (RGA), 3 pp.
- Notice of Service of Interrogatories and First Request of Documents and Things to Defendant 10X Genomics, Inc., dated Jul. 5, 2019 in the USDC for the District of Delaware, C.A. 18-1800 (RGA), 3 pp.
- Notification Prior to Examination dated Nov. 27, 2019 in Israeli Patent Application No. 265478.
- Novak et al., "Single-Cell Multiplex Gene Detection and Sequencing with Microfluidically Generated Agarose Emulsions," *Angew. Chem. Int. Ed.* 2011, 50, 390-395.
- Novus Biologicals, "Fixation and Permeability in ICC IF," Novus Biologicals 2021, 1-3.
- Office Action dated Jun. 6, 2016 in Chinese Patent Application No. 201380022187.9.
- Office Action dated Dec. 27, 2016 in Chinese Patent Application No. 201380022187.9.
- Office Action dated Feb. 17, 2017 in Canadian Patent Application No. 2,865,575.
- Office Action dated Jul. 14, 2017 in Chinese Patent Application No. 201380022187.9.
- Office Action dated Dec. 19, 2017 in Chinese Patent Application No. 201480061859.1.
- Office Action dated Feb. 15, 2018 in Canadian Patent Application No. 2,865,575.
- Office Action dated Sep. 7, 2018 in Chinese Patent Application No. 201480061859.1.
- Office Action dated Dec. 13, 2018 in Canadian Patent Application No. 2,865,575.
- Office Action dated Jan. 2, 2019 in Chinese Patent Application No. 201480059505.3.
- Office Action dated Mar. 4, 2020 in Canadian Patent Application No. 2,865,575.
- Office Action dated Jun. 22, 2020 in Chinese Patent Application No. 201680007351.2.
- Office Action dated Jun. 22, 2020 in Chinese Patent Application No. 201680007652.5.
- Office Action dated Jun. 23, 2020 in Chinese Patent Application No. 2016800157452.
- Office Action dated Jul. 20, 2020 in Japanese Patent Application No. 2018-512152.
- Office Action dated Oct. 29, 2020 in Chinese Patent Application No. 2018800377201.
- Office Action dated Jan. 4, 2021 in Japanese Patent Application No. 2017-549390.
- Office Action dated Jan. 6, 2021 in Chinese Patent Application No. 201680052330.2.
- Office Action dated Jan. 14, 2021 in Japanese Patent Application No. 2019-014564.
- Office Action dated Jan. 15, 2021 in Korean Patent Application No. 10-2020-7033213.
- Office Action dated Jan. 26, 2021 in Chinese Patent Application No. 201680007351.2.
- Office Action dated Feb. 4, 2021 in Canadian Patent Application No. 2,865,575.

(56)

**References Cited****OTHER PUBLICATIONS**

- Office Action dated Feb. 20, 2021 in Chinese Patent Application No. 201680022865.5.
- Office Action dated Mar. 1, 2021 in Chinese Patent Application No. 201680007652.5.
- Office Action dated Mar. 2, 2021 in Chinese Patent Application No. 2016800157452.
- Office Action dated Mar. 8, 2021 in Japanese Patent Application No. 2018-512152.
- Office Action dated Mar. 16, 2021 in Chinese Patent Application No. 2018800377201.
- Office Action dated May 10, 2021 in Japanese Patent Application No. 2019-566787.
- Office Action dated May 21, 2021 in Chinese Patent Application No. 201680007351.2.
- Office Action dated Jul. 26, 2021 in Korean Patent Application No. 10-2019-7011635.
- Office Action dated Jul. 28, 2021 in Korean Patent Application No. 10-2020-7033213.
- Office Action dated Aug. 13, 2021 in Chinese Patent Application No. 2017800587991.
- Office Action dated Aug. 27, 2021 in Chinese Patent Application No. 2016800076525.
- Office Action dated Aug. 30, 2021 in Japanese Patent Application No. 2019-540515.
- Office Action dated Aug. 31, 2021, in Korean Patent Application No. 10-2019-7038794.
- Office Action dated Sep. 14, 2021, in Chinese Patent Application No. 2016800523302.
- Ogino et al., "Quantification of PCR bias caused by a single nucleotide polymorphism in SMN gene dosage analysis," *J Mol Diagn.* 2002, 4(4), 185-190.
- Opposition to Defendant's Motion to Dismiss Pursuant to Federal Rule of Civil Procedure 12(b)(6) dated Feb. 15, 2019, in the USDC for the District of Delaware, C.A. 18-800-RGA, 3 pp.
- Oral Order by Judge Andrews Canceling Scheduling Conference set for May 8, 2019.
- Order Setting Rule 16(b) Conference as Ordered by Judge Andrews Pursuant to Fed. R. Civ. P. 16(b), ruling dated Apr. 17, 2019 in the USDC District of Delaware, C.A. 18-1800-RGA, 1 pp.
- Order Scheduling ADR Mediation Teleconference, filed May 13, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 4pp.
- O'Shea et al., "Analysis of T Cell Receptor Beta Chain CDR3 Size Using RNA Extracted from Formalin Fixed Paraffin Wax Embedded Tissue," *Journal of Clinical Pathology* 1997, 50(10), 811-814.
- Ozkumur et al., "Inertial Focusing for Tumor Antigen—Dependent and—Independent Sorting of Rare Circulating Tumor Cells," *Science Translational Medicine* 2013, 5(179), 1-20.
- Parameswaran et al., "A pyrosequencing-tailored nucleotide barcode design unveils opportunities for large-scale sample multiplexing," *Nucleic Acids Res.* 2007, 35(19), e130, 1-9.
- Park et al., "Discovery of common Asian copy number variants using integrated high-resolution array CGH and massively parallel DNA sequencing," *Nat Genet.* 2010, 42(5), 400-405.
- Patanjali et al., "Construction of a uniform-abundance (normalized) CNDA library," *Proceedings of the National Academy of Sciences* 1991, 88(5), 1943-1947.
- Peng et al., "Reducing amplification artifacts in high multiplex amplicon sequencing by using molecular barcodes," *BMC Genomics* 2015, 16(589), 1-12.
- Pérez-Rentero et al., "Synthesis of Oligonucleotides Carrying Thiol Groups Using a Simple Reagent Derived from Threoninol," *Molecules* 2012, 17, 10026-10045.
- Peterson et al., "Multiplexed quantification of proteins and transcripts in single cells," *Nature Biotechnology* 2017, 35, 936-939.
- Pfaffl et al., "Determination of stable housekeeping genes, differentially regulated target genes andsample integrity: BestKeeper—Excel-based tool using pair-wise correlations," *Biotechnology Letters* 2004, 26(6), 505-515.
- Picelli et al., "Tn5 transposase and tagmentation procedures for massively scaled sequencing projects," *Genome Research* 2014, 24(12), 2033-2040.
- Picelli et al., "Single-cell RNA-sequencing: The future of genome biology is now," *RNA Biology* 2017, 14(5), 637-650.
- Pihlak et al., "Rapid genome sequencing with short universal tiling probes," *Nature Biotechnology* 2008, 26, 1-9.
- Pinkel et al., "Comparative Genomic Hybridization," *Annual Review of Genomics and Human Genetics* 2005, 6, 331-354.
- Plaintiff's Brief in Opposition to Defendant's Motion to Dismiss Pursuant to Fed. R. Civ. P. 12(b)(6), filed Mar. 29, 2019 in the USDC District of Delaware, C.A. No. 18-1800 (RGA), 27 pp.
- Plaintiff's First Amended Complaint filed on Feb. 8, 2019, in the USDC for the District of Delaware, C.A. 18-1800-RGA, 178 pp.
- Pleasance et al., "A small-cell lung cancer genome with complex signatures of tobacco exposure," *Nature* 2010, 463(7278), 184-190.
- Plessy et al., "Population transcriptomics with single-cell resolution: a new field made possible by microfluidics: a technology for high throughput transcript counting and data-driven definition of cell types," *Bioessays* 2012, 35, 131-140.
- Pre-interview communication dated Nov. 27, 2018 in U.S. Appl. No. 16/012,635.
- Preissl et al., "Single-nucleus analysis of accessible chromatin in developing mouse forebrain reveals cell-type-specific transcriptional regulation," *Nature Neuroscience* 2018, 21(3), 432-439.
- Prevette et al., "Polycation-Induced Cell Membrane Permeability Does Not Enhance Cellular Uptake or Expression Efficiency of Delivered DNA," *Molecular Pharmaceutics* 2010, 7(3), 870-883.
- Pringle et al., "In Situ Hybridization Demonstration of Poly-Adenylated RNA Sequences in Formalin-Fixed Parafin Sections Using a Biotinylated Oligonucleotide Poly d(T) Probe," *Journal of Pathology* 1989, 158, 279-286.
- Proposed Stipulated Protective Order Pursuant to Rule 26(c) of the Federal Rules of Civil Procedure, filed Jun. 20, 2019 In the USDC for the District of Delaware, C.A. 18-1800 (RGA), 26 pp.
- Qiu et al., "DNA Sequence-Based "Bar Codes" for Tracking the Origins of Expressed Sequence Tagsfrom a Maize cDNA Library Constructed Using Multiple mRNA Sources," *Plant Physiol.* 2003, 133, 475-481.
- Quail et al., "SASI-Seq: sample assurance Spike-Ins, and highly differentiating 384 barcoding for Illumina sequencing," *BMC Genomics* 2014, 15(110), in 13 pages.
- Raj et al., "Stochastic mRNA synthesis in mammalian cells," *PLoS Biol.* 2006, 4(10) 1707-1719.
- Raj et al., "Imaging individual mRNA molecules using multiple singly labeled probes," *Nature Methods* 2008, 5(10), 877-879.
- Raj et al., "Single-Molecule Approaches to Stochastic Gene Expression," *Annu Rev Biophys* 2009, 38, 255-270.
- Rajeevan et al., "Global amplification of sense RNA: a novel method to replicate and archive mRNA for gene expression analysis," *Genomics* 2003, 82, 491-497.
- Report on the Filing or Determination of an Action Regarding a Patent or Trademark filed Nov. 15, 2018 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 2 pp.
- Restriction Requirement dated Mar. 15, 2016 in U.S. Appl. No. 14/381,488.
- Restriction Requirement dated Mar. 17, 2016 in U.S. Appl. No. 14/472,363.
- Restriction Requirement dated Mar. 29, 2019 in U.S. Appl. No. 15/715,028.
- Restriction Requirement dated May 5, 2021 in U.S. Appl. No. 16/400,886.
- Restriction Requirement dated May 28, 2021 in U.S. Appl. No. 16/781,814.
- Restriction Requirement dated Jun. 19, 2019 in U.S. Appl. No. 15/596,364.
- Restriction Requirement dated Jun. 4, 2021 in U.S. Appl. No. 16/551,620.
- Restriction Requirement dated Sep. 20, 2019 in U.S. Appl. No. 15/875,816.
- Restriction Requirement dated Sep. 20, 2021 in U.S. Appl. No. 16/525,054.

(56)

**References Cited****OTHER PUBLICATIONS**

- Restriction Requirement dated Oct. 1, 2021 in U.S. Appl. No. 16/525,054.
- Rhee et al., "Simultaneous detection of mRNA and protein stem cell markers in live cells," *BMC Biotechnology* 2009, 9(30), 1-10.
- Roche Diagnostics GmbH, "Genome Sequencer 20 System: First to the Finish," 2006, 1-40.
- Rule 7.1 Disclosure Statement dated Nov. 15, 2018 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 1 pp.
- Sah et al., "Complete Genome Sequence of a 2019 Novel Coronavirus (SARS-CoV-2) Strain Isolated in Nepal," *Microbiol Resour Announc.* 2020, 9(11), e00169-20, 3 pp.
- Sano et al., "Immuno-PCR: Very Sensitive Antigen Detection by Means of Specific Antibody—DNA Conjugates," *Science* 1992, 258, 120-122.
- Sasagawa et al., "Quartz-Seq: a highly reproducible and sensitive single-cell RNA sequencing method, reveals non-genetic gene-expression heterogeneity," *Genome Biology* 2013, 14, R31.
- Sasuga et al., Single-cell chemical lysis method for analyses of intracellular molecules using an array of picoliter-scale microwells, *Anal Chem* 2008, 80(23), 9141-9149.
- Satija et al., Spatial reconstruction of single-cell gene expression data, *Nature Biotechnology* 2015, 33(5), 495-508.
- Scheduling Order pursuant to Local Rule 16.1(b), filed May 7, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 10 pp.
- Scheduling Order Signed by Judge Andrews, dated May 8, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 10 pp.
- Schmitt et al., "Detection of ultra-rare mutations by next-generation sequencing," *Proc Natl Acad Sci* 2012, 109(36), 1-6.
- Schouten et al., "Relative quantification of 40 nucleic acid sequences by multiplex ligation-dependent probe amplification," *Nucleic Acids Research* 2002, 30(12), e57.
- Search and Examination Report dated Aug. 26, 2015 in United Kingdom Patent Application No. 1511591.8.
- Search Report and Written Opinion dated Jan. 26, 2016 in Singapore Patent Application No. 1120140527W.
- Search Report and Written Opinion dated Aug. 26, 2020 in Singapore Patent Application No. 10201806890V.
- Sebat et al., "Large-Scale Copy Number Polymorphism in the Human Genome," *Science* 2004, 305, 525-528.
- Shahi et al., "Abseq: ultrahigh-throughput single cell protein profiling with droplet microfluidic barcoding," *Scientific Reports* 2017, 7(44447), 1-10.
- Shalek et al., "Single-cell transcriptomics reveals bimodality in expression and splicing in immune cells," *Nature* 2013, 498(7453), 236-240.
- Shapiro et al., "Single-cell sequencing-based technologies will revolutionize whole-organism science," *Nature Reviews Genetics* 2013, 14, 618-629.
- Shendure et al., "Next-generation DNA sequencing," *Nature Biotechnology* 2008, 26(10), 1135-1145.
- Shiroguchi et al., "Digital RNA sequencing minimizes sequence-dependent bias and amplification noise with optimized single-molecule barcodes," *Proc Natl Acad Sci* 2012, 109(4):1347-1352.
- S.H.Ko, "An 'equalized cDNA library' by the reassociation of short double-stranded cDNAs," *Nucleic Acids Res.* 1990, 18(19), 5705-5711.
- Shoemaker et al., "Quantitative phenotypic analysis of yeast deletion mutants using a highly parallel molecular bar-coding strategy," *Nature Genetics* 1996, 14, 450-456.
- Shortreed et al., "A thermodynamic approach to designing structure-free combinatorial DNA word sets," *Nucleic Acids Res.* 2005, 33(15), 4965-4977.
- Shum et al., "Quantitation of mRNA Transcripts and Proteins Using the BD Rhapsody™ Single-Cell Analysis System," *Adv Exp Med Biol.* 2019, 1129, 63-79.
- Simpson et al., "Copy number variant detection in inbred strains from short read sequence data," *Bioinformatics* 2010, 26(4), 565-567.
- Smith et al., "Highly-multiplexed barcode sequencing: an efficient method for parallel analysis of pooled samples," *Nucleic Acids Research* 2010, 38(13), e142, 1-7.
- Soares et al., "Construction and characterization of a normalized cDNA library," *Proc. Natl. Acad. Sci.* 1994, 91, 9228-9232.
- Sogin et al., "Microbial diversity in the deep sea and the underexplored 'rare biosphere,'" *PNAS* 2008, 103(32), 12115-12120.
- Sommer et al., "Minimal homology requirements for PCR primers," *Nucleic Acids Research* 1989, 17(16), 6749.
- Song et al., "Design rules for size-based cell sorting and sheathless cell focusing by hydrophoresis," *Journal of Chromatography A* 2013, 1302, 191-196.
- Song et al., DNase-seq: a high-resolution technique for mapping active gene regulatory elements across the genome from mammalian cells, *Cold Spring Harb Protoc* 2010, 2, in 13 pages.
- Sos et al., "Characterization of chromatin accessibility with a transposome hypersensitive sites sequencing (THS-seq) assay," *Genome Biology* 2016, 17(20), in 15 pages.
- Soumilon et al., "Characterization of directed differentiation by high-throughput single-cell RNA-Seq," *bioRxiv* 2014, 1-13.
- Speicher et al., "The new cytogenetics: blurring the boundaries with molecular biology," *Nature Reviews Genetics* 2005, 6(10), 782-792.
- Statement of Opposition of Strawman Limited filed against European Patent No. EP2414548B1 on Jul. 19, 2016.
- Statement of Opposition dated Jul. 21, 2016 filed against European Patent No. EP2414548B1.
- Statement of Opposition filed against European Patent No. EP2414548B1 on Jul. 26, 2016.
- Statement regarding Third-Party Submission filed on Jun. 6, 2018 for U.S. Appl. No. 15/847,752.
- Stipulated Protective Order Pursuant to Rule 26(c) of the Federal Rules of Civil Procedure, dated Jun. 21, 2019 in the USDC for the District of Delaware, C.A. 18-1800 (RGA), 26 pp.
- Stipulation and Order to Extend Time to File Opposition to Motion to Dismiss, and Reply in Support of the Motion, dated Jan. 28, 2019 in the USDC for the District of Delaware, C.A. 18-1800-RGA, 2 pp.
- Stoeckius et al., "Large-scale simultaneous measurement of epitopes and transcriptomes in single cells," *Nature Methods* 2017, 14(9), 865-868.
- Stoeckius et al., "Cell Hashing with barcoded antibodies enables multiplexing and doublet detection for single cell genomics," *Genome Biology* 2018, 19(224), 1-12.
- Stratagene 1988 Catalog, Gene Characterization Kits, 39.
- Subkhankulova et al., "Comparative evaluation of linear and exponential amplification techniques for expression profiling at the single cell level," *Genome Biology* 2006, 7(3), 1-16.
- Submission dated Jan. 15, 2018 in preparation for upcoming oral proceedings in opposition against European Patent No. EP2414548B1.
- Summons in a Civil Action to Defendant 10X Genomics, Inc. filed Nov. 16, 2018 in the USDC for the District of Delaware, Civil Action No. 18-1800, 2 pp.
- Summons to Attend Oral Proceedings dated Nov. 16, 2020 in European Patent Application No. 17202409.3.
- Sun et al., "Ultra-deep profiling of alternatively spliced Drosophila Dscam isoforms by circularization-assisted multi-segment sequencing," *EMBO J.* 2013, 32(14), 2029-2038.
- Takahashi et al., "Novel technique of quantitative nested real-time PCR assay for mycobacterium tuberculosis DNA," *Journal of Clinical Microbiology* 2006, 44, 1029-1039.
- Takara Bio, "SMARTer Human BCR IgG IgM H/K/L Profiling Kit User Manual," Takara Bio USA Inc. 2019, 1-22.
- Tan et al., "Genome-wide comparison of DNA hydroxymethylation in mouse embryonic stem cells and neural progenitor cells by a new comparative hMeDIP-seq method," *Nucleic Acids Res.* 2013, 41(7), e84, 1-12.
- Tang et al., "RNA-Seq analysis to capture the transcriptome landscape of a single cell," *Nature Protocols* 2010, 5(3), 516-535.
- Taudien et al., "Haplotyping and copy No. estimation of the highly polymorphic human beta-defensin locus on 8p23 by 454 amplicon sequencing," *BMC Genomics* 2010, 11, 252, 1-14.
- The Tibbs Times, UNC bioscience newsletter, Apr. 2010, 1-17.
- Third-Party Submission filed on May 21, 2018 for U.S. Appl. No. 15/847,752.

(56)

**References Cited****OTHER PUBLICATIONS**

- Tomaz et al., "Differential methylation as a cause of allele dropout at the imprinted GNAS locus," *Genet Test Mol Biomarkers* 2010, 14(4), 455-460.
- TotalSeq<sup>TMM</sup>-A0251 anti-human Hashtag 1 Antibody, BioLegend<sup>®</sup>, Jul. 2018, 1-10.
- Treutlein et al., Reconstructing lineage hierarchies of the distal lung epithelium using single-cell RNA-seq, *Nature* 2014, 509, 371-375.
- Trzupek et al., "Discovery of CD80 and CD86 as recent activation markers on regulatory T cells by protein-RNA single-cell analysis", *Genome Medicine* 2020, 12(1), in 22 pages.
- Ullal et al., "Cancer cell profiling by barcoding allows multiplexed protein analysis in fine needle aspirates," *Sci Transl Med*. 2014, 6(219), 22 pp.
- Unopposed Motion to Extend Time for Defendant's Response, dated Dec. 4, 2018 in the USDC for the District of Delaware, C.A. 18-1800-(RGA), 2 pp.
- Vandesompele et al., "Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes," *Genome Biology* 2002, 3(7), 1-12.
- Velculescu et al., "Serial Analysis of Gene Expression," *Science* 1995, 270(5235), 484-487.
- Velculescu et al., "Characterization of the Yeast Transcriptome," *Cell* 1997, 88, 243-251.
- Vestheim et al., "Application of Blocking Oligonucleotides to Improve Signal-to-Noise Ratio in a PCR," *Methods in Molecular Biology* 2011, 687, 265-274.
- Vogelstein et al., "Digital PCR," *Proc. Natl. Acad. Sci.* 1999, 96, 9236-9241.
- Vollbrecht et al., "Validation and comparison of two NGS assays for the detection of EGFR T790M resistance mutation in liquid biopsies of NSCLC patients," *Oncotarget* 2018, 9(26), 18529-18539.
- Walker et al., "Isothermal in vitro amplification of DNA by a restriction enzyme/DNA polymerase system," *Proc Natl Acad Sci* 1992, 89, 392-396.
- Walsh et al., "Detection of inherited mutations for breast and ovarian cancer using genomic capture and massively parallel sequencing," *Proc Natl Acad Sci* 2010, 107(28), 12629-12633.
- Wang et al., "Combining Gold Nanoparticles with Real-Time Immuno-PCR for Analysis of HIV p24 Antigens," *Proceedings of ICBBE* 2007, 1198-1201.
- Wang et al., "RNA-Seq: a revolutionary tool for transcriptomics," *Nature Reviews Genetics* 2009, 10(1), 57-63.
- Wang et al., "iCLIP predicts the dual splicing effects of TIA-RNA interactions," *PLoS Biol* 2010, 8(10), e1000530, 1-16.
- Wang et al., "Advances and applications of single-cell sequencing technologies," *Molecular Cell* 2015, 58, 598-609.
- Wang et al., "Fragmentation-based whole-genome bisulfite sequencing," *Nature Protocols* 2013, 8(10), 2022-2032.
- Warren et al., "Transcription factor profiling in individual hematopoietic progenitors by digital RT-PCR," *PNAS* 2006, 103(47), 17807-17812.
- Weber et al., "A real-time polymerase chain reaction assay for quantification of allele ratios and correction of amplification bias," *Anal Biochem*. 2003, 320, 252-258.
- Weibrech et al., "Proximity ligation assays: a recent addition to the proteomics toolbox," *Expert Rev. Proteomics* 2010, 7(3), 401-409.
- Weiner et al., "Kits and their unique role in molecular biology: a brief retrospective," *BioTechniques* 2008, 44(5), 701-704.
- White et al., "High-throughput microfluidic single-cell RT-qPCR," *PNAS* 2011, 108(34), 13999-14004.
- Wittes et al., "Searching for Evidence of Altered Gene Expression: a Comment on Statistical Analysis of Microarray Data," *Journal of the National Cancer Institute* 1999, 91(5), 400-401.
- Wodicka et al., "Genome-wide expression monitoring in *Saccharomyces cerevisiae*," *Nature Biotechnology* 1997, 15, 1359-1367.
- Wojdacz et al., "Primer design versus PCR bias in methylation independent PCR amplifications," *Epigenetics* 2009, 4(4), 231-234.
- Wood et al., "Using next-generation sequencing for high resolution multiplex analysis of copy number variation from nanogram quantities of DNA from formalin-fixed paraffin-embedded specimens," *Nucleic Acids Res*. 2010, 38(14), 1-14.
- Written Submission of Publications dated Jun. 14, 2018 in Japanese Patent Application No. 2016-537867.
- Wu et al., "Quantitative assessment of single-cell RNA-sequencing methods," *Nat Methods* 2014, 11(1), 41-46.
- Yandell et al., "A probabilistic disease-gene finder for personal genomes," *Genome Res*. 2011, 21(9), 1529-1542.
- Ye et al., "Fluorescent microsphere-based readout technology for multiplexed human single nucleotide polymorphism analysis and bacterial identification," *Human Mutation* 2001, 17(4), 305-316.
- Yoon et al., "Sensitive and accurate detection of copy number variants using read depth of coverage," *Genome Res*. 2009, 19, 1586-1592.
- Zeberg et al., "The major genetic risk factor for severe COVID-19 is inherited from Neanderthals," *Nature* 2020, 587(7835), 1-13.
- Zagordi et al., "Error correction of next-generation sequencing data and reliable estimation of HIV quasispecies," *Nucleic Acids Research* 2010, 38(21), 7400-7409.
- Zhang et al., "The impact of next-generation sequencing on genomics," *J Genet Genomics* 2011, 38(3), 95-109.
- Zhang et al., "DNA-based hybridization chain reaction for amplified bioelectronic signal and ultrasensitive detection of proteins," *Anal Chem*. 2012, 84, 5392-5399.
- Zhao et al., "Homozygous Deletions and Chromosome Amplifications in Human Lung Carcinomas Revealed by Single Nucleotide Polymorphism Array Analysis," *Cancer Research* 2005, 65(13), 5561-5570.
- Zhao et al., "Methylated DNA Immunoprecipitation and High-Throughput Sequencing (MeDIP-seq) Using Low Amounts of Genomic DNA," *Cellular Reprogramming* 2014, 16(3), in 20 pages.
- Zheng et al., "Haplotyping germline and cancer genomes with high-throughput linked-read sequencing," *Nature Biotechnology* 2016, 34(3), 303-311.
- Zhou et al., "Counting alleles reveals a connection between chromosome 18q loss and vascular invasion," *Nature Biotechnology* 2001, 19, 78-81.
- Zhou et al., "Photocleavable Peptide-Oligonucleotide Conjugates for Protein Kinase Assays by MALDI-TOF MS," *Mol. BioSyst*. 2012, 8, 2395-2404.
- Zhu et al., "Reverse Transcriptase Template Switching: A SMART Approach for Full-Length cDNA Library Construction," *BioTechniques* 2001, 30(4), 892-897.
- Bolivar et al., "Targeted next-generation sequencing of endometrial cancer and matched circulating tumor DNA: identification of plasma-based, tumor-associated mutations in early stage patients," *Modern Pathology* 2019, 32(3), 405-414.
- Buenrostro et al., "Transposition of native chromatin for multimodal regulatory analysis and personal epigenomics," *Nat Methods* 2013, 10(12), 1213-1218.
- Buenrostro et al., "ATAC-seq: A Method for Assaying Chromatin Accessibility Genome-Wide," *Curr Protoc Mol Biol* 2016, 109, 1-21.
- Chang et al., "Single-cell protein and gene expression profiling of stem memory T cells by BD Ab-seq," Annual Joint Meeting of the American Society for Cell Biology and the European Molecular Biology Organization 2017, 28(26), P1896.
- Chen et al., "High-throughput sequencing of the transcriptome and chromatin accessibility in the same cell," *Nature Biotechnology* 2019, 37, 1452-1457.
- De Simone et al., "Single Cell T Cell Receptor Sequencing: Techniques and Future Challenges," *Frontiers in Immunology* 2018, 9(1638), 1-7.
- Dengl et al., "Engineered hapten-binding antibody derivatives for modulation of pharmacokinetic properties of small molecules and targeted payload delivery," *Immunol Rev*. 2016, 270, 165-177.
- Dovgan et al., "Antibody- Oligonucleotide Conjugates as Therapeutic, Imaging, and Detection Agents," *Bioconjugate Chem*. 2019, 30, 2483-2501.

(56)

**References Cited****OTHER PUBLICATIONS**

- Erickson et al., "AbSeq Protocol Using the Nano-Well Cartridge-Based Rhapsody Platform to Generate Protein and Transcript Expression Data on the Single-Cell Level," *STAR Protocols* 2020, in 31 pages.
- Examination Report dated Nov. 18, 2021 in European Patent Application No. 19724003.9.
- Examination Report dated Nov. 24, 2021 in European Patent Application No. 19762517.1.
- Examination Report dated Dec. 6, 2021 in European Patent Application No. 18703156.2.
- Examination Report dated Dec. 9, 2021 in European Patent Application No. 19723988.2.
- Examination Report dated Apr. 7, 2022 in Singapore Patent Application No. 10201806890V.
- Examination Report dated Apr. 8, 2022 in Australian Patent Application No. 2018281745.
- Final Office Action dated Nov. 2, 2021 in U.S. Appl. No. 16/789,311.
- Final Office Action dated Jan. 18, 2022 in U.S. Appl. No. 16/588,405.
- Final Office Action dated Feb. 23, 2022 in U.S. Appl. No. 16/707,780.
- Final Office Action dated Mar. 15, 2022 in U.S. Appl. No. 16/374,626.
- Final Office Action dated Mar. 25, 2022 in U.S. Appl. No. 16/551,620.
- Final Office Action dated Apr. 12, 2022 in U.S. Appl. No. 15/084,307.
- Final Office Action dated May 26, 2022 in U.S. Appl. No. 16/747,737.
- Final Office Action dated Jun. 14, 2022 in U.S. Appl. No. 15/055,407.
- Final Office Action dated Aug. 23, 2022 in U.S. Appl. No. 16/012,584.
- Final Office Action dated Nov. 15, 2022 in U.S. Appl. No. 16/525,054.
- Final Office Action dated Nov. 16, 2022 in U.S. Appl. No. 16/588,405.
- Final Office Action dated Jan. 25, 2023 in U.S. Appl. No. 16/789,311.
- Final Office Action dated Jan. 26, 2023 in U.S. Appl. No. 16/459,444.
- Final Office Action dated Feb. 21, 2023 in U.S. Appl. No. 16/551,620.
- Final Office Action dated Apr. 25, 2023 in U.S. Appl. No. 16/525,054.
- Goodridge et al., "Synthesis of Albumin and Malic Enzyme in Wheat-Germ Lysates and *Xenopus laevis* Oocytes Programmed with Chicken-Liver Messenger RNA," *Eur. J. Biochem.* 1979, 96, 1-8.
- International Search Report and Written Opinion dated Oct. 29, 2021, in PCT Application No. PCT/US2021/032319.
- International Search Report and Written Opinion dated Dec. 6, 2021, in PCT Application No. PCT/US2021/046750.
- International Search Report and Written Opinion dated Nov. 12, 2021, in PCT Application No. PCT/US2021/044036.
- International Search Report and Written Opinion dated Mar. 10, 2022, in PCT Application No. PCT/US2021/060206.
- International Search Report and Written Opinion dated Apr. 12, 2022, in PCT Application No. PCT/US2021/059573.
- International Search Report and Written Opinion dated Mar. 11, 2022, in PCT Application No. PCT/US2021/060197.
- International Search Report and Written Opinion dated Apr. 5, 2022, in PCT Application No. PCT/US2021/062473.
- International Search Report and Written Opinion dated Jun. 8, 2022, in PCT Application No. PCT/US2022/021015.
- International Search Report and Written Opinion dated Jul. 29, 2022, in PCT Application No. PCT/US2022/029023.
- International Search Report and Written Opinion dated Jul. 29, 2022, in PCT Application No. PCT/US2022/029057.
- International Search Report and Written Opinion dated Dec. 5, 2022, in PCT Application No. PCT/US2022/075774.
- International Search Report and Written Opinion dated Dec. 15, 2022, in PCT Application No. PCT/US2022/075655.
- International Search Report and Written Opinion dated Dec. 20, 2022, in PCT Application No. PCT/US2022/075661.
- International Search Report and Written Opinion dated Dec. 22, 2022, in PCT Application No. PCT/US2022/075577.
- International Search Report and Written Opinion dated Jan. 9, 2023, in PCT Application No. PCT/US2022/076366.
- International Search Report and Written Opinion dated Jan. 17, 2023, in PCT Application No. PCT/US2022/076056.
- International Search Report and Written Opinion dated Feb. 13, 2023, in PCT Application No. PCT/US2022/075656.
- Jacobsen et al., "33rd Annual Meeting & Pre-Conference Programs of the Society for Immunotherapy of Cancer," *Journal for Immunotherapy of Cancer* 2018, 6(S1), 7-11.
- Lake et al., "Integrative single-cell analysis of transcriptional and epigenetic states in the human adult brain," *Nature Biotechnology* 2018, 36(1), 70-80.
- Lee et al., "Comparison of Surface Markers between Human and Rabbit Mesenchymal Stem Cells," *Plos One* 2014, 9(11), in 10 pages.
- Lutz et al., "Isolation and analysis of high quality nuclear DNA with reduced organellar DNA for plant genome sequencing and resequencing," *BMC Biotechnology* 2011, 11(54), in 9 pages.
- Mayer et al., "Obtaining deeper insights into microbiome diversity using a simple method to block host and nontargets in amplicon sequencing," *Molecular Ecology Resources* 2021, 21(6), 1952-1965.
- Minnoye et al., "Chromatin accessibility profiling methods," *Nature Reviews Method Primers* 2021, 1- 24.
- Monneron, "One-step Isolation and Characterization of Nuclear Membranes, 1974 Electron Microscopy and Composition of Biological Membranes and Envelopes," *The Royal Publishing Society* 1974, 268, 101-108.
- Non-Final Office Action dated Dec. 15, 2021 in U.S. Appl. No. 15/875,816.
- Non-Final Office Action dated Dec. 21, 2021 in U.S. Appl. No. 15/055,407.
- Non-Final Office Action dated Jan. 6, 2022 in U.S. Appl. No. 15/084,307.
- Non-Final Office Action dated Feb. 2, 2022 in U.S. Appl. No. 16/747,737.
- Non-Final Office Action dated Feb. 9, 2022 in U.S. Appl. No. 16/525,054.
- Non-Final Office Action dated Apr. 5, 2022 in U.S. Appl. No. 16/400,885.
- Non-Final Office Action dated Apr. 8, 2022 in U.S. Appl. No. 16/232,287.
- Non-Final Office Action dated May 3, 2022 in U.S. Appl. No. 16/012,584.
- Non-Final Office Action dated May 11, 2022 in U.S. Appl. No. 16/588,405.
- Non-Final Office Action dated May 19, 2022 in U.S. Appl. No. 16/459,444.
- Non-Final Office Action dated Jul. 7, 2022 in U.S. Appl. No. 16/788,743.
- Non-Final Office Action dated Jul. 7, 2022 in U.S. Appl. No. 16/677,012.
- Non-Final Office Action dated Jul. 18, 2022 in U.S. Appl. No. 16/551,620.
- Non-Final Office Action dated Jul. 27, 2022 in U.S. Appl. No. 16/747,737.
- Non-Final Office Action dated Oct. 13, 2022 in U.S. Appl. No. 17/147,272.
- Non-Final Office Action dated Nov. 17, 2022 in U.S. Appl. No. 16/551,638.
- Non-Final Office Action dated Dec. 8, 2022 in U.S. Appl. No. 16/934,530.
- Non-Final Office Action dated Dec. 21, 2022 in U.S. Appl. No. 15/055,407.
- Non-Final Office Action dated Jan. 10, 2023 in U.S. Appl. No. 17/163,177.
- Non-Final Office Action dated Jan. 19, 2023 in U.S. Appl. No. 17/091,639.
- Non-Final Office Action dated Jan. 23, 2023 in U.S. Appl. No. 17/183,840.
- Non-Final Office Action dated Jan. 24, 2023 in U.S. Appl. No. 17/157,872.
- Non-Final Office Action dated Feb. 10, 2023 in U.S. Appl. No. 17/390,640.
- Non-Final Office Action dated Feb. 23, 2023 in U.S. Appl. No. 17/408,374.
- Non-Final Office Action dated Mar. 13, 2023 in U.S. Appl. No. 17/151,050.

(56)

**References Cited**

## OTHER PUBLICATIONS

- Non-Final Office Action dated Apr. 26, 2023 in U.S. Appl. No. 16/540,971.
- Non-Final Office Action dated Apr. 26, 2023 in U.S. Appl. No. 16/374,626.
- Notice of Allowance dated Nov. 16, 2021 in U.S. Appl. No. 16/836,750.
- Notice of Allowance dated Jan. 24, 2022 in Korean Patent Application No. 16/836,750.
- Notice of Allowance dated Feb. 9, 2022 in U.S. Appl. No. 16/781,814.
- Notice of Allowance dated Feb. 11, 2022 in Chinese Patent Application No. 201680007351.2.
- Notice of Allowance dated Feb. 16, 2022 in U.S. Appl. No. 15/875,816.
- Notice of Allowance dated Feb. 21, 2022 in Korean Patent Application No. 10-2020-7033213.
- Notice of Allowance dated Apr. 11, 2022 in U.S. Appl. No. 15/134,967.
- Notice of Allowance dated Apr. 25, 2022 in Korean Patent Application No. 10-2018-7008560.
- Notice of Allowance dated Apr. 26, 2022 in Chinese Patent Application No. 201780058799.1.
- Notice of Allowance dated Apr. 27, 2022 in U.S. Appl. No. 16/400,886.
- Notice of Allowance dated May 9, 2022 in Australian Patent Application No. 2018281745.
- Notice of Allowance dated May 15, 2022 in Japanese Patent Application No. 2019-540515.
- Notice of Allowance dated May 23, 2022 in U.S. Appl. No. 15/715,028.
- Notice of Allowance dated May 26, 2022 in Korean Patent Application No. 10-2019-7038794.
- Notice of Allowance dated Jun. 6, 2022 in U.S. Appl. No. 16/789,358.
- Notice of Allowance dated Jul. 20, 2022 in U.S. Appl. No. 16/707,780.
- Notice of Allowance dated Aug. 9, 2022 in U.S. Appl. No. 16/232,287.
- Notice of Allowance dated Sep. 26, 2022, 2022 in U.S. Appl. No. 16/232,287.
- Notice of Allowance dated Oct. 17, 2022, 2022 in U.S. Appl. No. 16/400,885.
- Notice of Allowance dated Oct. 20, 2022 in Australian Patent Application No. 2019204928.
- Notice of Allowance dated Oct. 21, 2022 in European Patent Application No. 19762517.1.
- Notice of Allowance dated Oct. 24, 2022 in European Patent Application No. 20708266.0.
- Notice of Allowance dated Oct. 25, 2022 in European Patent Application No. 19724003.9.
- Notice of Allowance dated Nov. 7, 2022 in U.S. Appl. No. 16/012,584.
- Notice of Allowance dated Jan. 10, 2023 in U.S. Appl. No. 16/588,405.
- Notice of Allowance dated Jan. 19, 2023 in Korean Patent Application No. 10-2022-7004715.
- Notice of Allowance dated Jan. 31, 2023 in U.S. Appl. No. 16/747,737.
- Notice of Allowance dated Feb. 1, 2023 in U.S. Appl. No. 17/147,272.
- Notice of Allowance dated Feb. 21, 2023 in Korean Patent Application No. 10-2022-7017261.
- Notice of Allowance dated Mar. 1, 2023 in U.S. Appl. No. 17/192,814.
- Notice of Allowance dated Mar. 10, 2023 in European Patent Application No. 19762517.1.
- Notice of Allowance dated Mar. 10, 2023 in European Patent Application No. 20708266.0.
- Notice of Allowance dated Mar. 10, 2023 in European Patent Application No. 19724003.9.
- Notice of Allowance dated Mar. 13, 2023 in European Patent Application No. 17781265.8.
- Notice of Allowance dated Apr. 4, 2023 in Australian Patent Application No. 2017331459.
- Nowak et al., "Does the KIR2DS5 gene protect from some human diseases?," PLoS One 2010, 5(8), in 6 pages.
- Office Action dated Oct. 21, 2021, in Chinese Patent Application No. 2016800073512.
- Office Action dated Nov. 2, 2021, in Japanese Patent Application No. 2017-549390.
- Office Action dated Dec. 23, 2021, in Japanese Patent Application No. 2019-566787.
- Office Action dated Dec. 17, 2021 in Korean Patent Application No. 10-2018-7008560.
- Office Action dated Jan. 13, 2022 in Chinese Patent Application No. 2017800587991.
- Office Action dated Feb. 9, 2022 in Japanese Patent Application No. 2019-540515.
- Office Action dated Mar. 7, 2022 in Korean Patent Application No. 10-2022-7004715.
- Office Action dated May 5, 2022 in European Patent Application No. 19787547.9.
- Office Action dated May 17, 2022 in Australian Patent Application No. 2019204928.
- Office Action dated May 24, 2022 in European Patent Application No. 20708266.0.
- Office Action dated Jun. 28, 2022 in European Patent Application No. 16719706.0.
- Office Action dated Aug. 2, 2022 in European Patent Application No. 19765601.0.
- Office Action dated Aug. 1, 2022 in Korean Patent Application No. 10-2022-7017261.
- Office Action dated Sep. 21, 2022 in Israel Patent Application No. 265478.
- Office Action dated Jan. 30, 2023 in European Patent Application No. 19752792.2.
- Office Action dated Feb. 8, 2023 in Australian Patent Application No. 2017331459.
- Office Action dated Feb. 20, 2023 in European Patent Application No. 19723988.2.
- Office Action dated Feb. 23, 2023 in European Patent Application No. 20816802.1.
- Office Action dated Feb. 28, 2023 in Chinese Patent Application No. 2019111653930.
- Office Action dated Mar. 15, 2023 in European Patent Application No. 19787547.9.
- Office Action dated Mar. 29, 2023 in Chinese Patent Application No. 2020800144092.
- Office Action dated Apr. 10, 2023 in Japanese Patent Application No. 2022-030956.
- Office Action dated Apr. 14, 2023 in Chinese Patent Application No. 201980082680.7.
- Office Action dated Apr. 24, 2023 in Japanese Patent Application No. 2020-561800.
- Restriction Requirement dated Aug. 8, 2022 in U.S. Appl. No. 17/163,177.
- Restriction Requirement dated Aug. 11, 2022 in U.S. Appl. No. 17/091,639.
- Restriction Requirement dated Aug. 19, 2022 in U.S. Appl. No. 17/147,283.
- Restriction Requirement dated Sep. 16, 2022 in U.S. Appl. No. 17/151,050.
- Restriction Requirement dated Sep. 19, 2022 in U.S. Appl. No. 16/934,530.
- Restriction Requirement dated Oct. 21, 2022 in U.S. Appl. No. 17/320,052.
- Restriction Requirement dated Nov. 8, 2022 in U.S. Appl. No. 17/157,872.
- Restriction Requirement dated Dec. 23, 2022 in U.S. Appl. No. 17/531,618.
- Restriction Requirement dated Jan. 20, 2023 in U.S. Appl. No. 17/373,519.
- Restriction Requirement dated Feb. 27, 2023 in U.S. Appl. No. 17/151,058.
- Restriction Requirement dated Apr. 4, 2023 in U.S. Appl. No. 17/161,558.
- Uellendahl-Werth et al., "A benchmark of hemoglobin blocking during library preparation for mRNA Sequencing of human blood samples," Scientific Reports 2020, 10(1), 5630.

(56)

**References Cited****OTHER PUBLICATIONS**

- Wangsanuwat et al., "Efficient and cost-effective bacterial mRNA sequencing from low input samples through ribosomal RNA depletion," *BMC Genomics* 2020, 21(1), 1-12.
- Yang & Zhao, "Quantitative Analysis of Nonoxynol-9 in Blood," *Contraception* 1991, 43(2), 161-166.
- Zhang et al., "Immunoaffinity Purification of Plasma Membrane with Secondary Antibody Superparamagnetic Beads," *Journal of Proteome* 2006, 6, 34-43.
- 10X Genomics, Inc., 2022, "Chromium Fixed RNA Profiling Reagent Kits," 10xGenomics.com, User Guide, in 95 pages.
- Advisory Action dated May 31, 2023 in U.S. Appl. No. 16/789,311.
- Arguel et al., "A cost effective 5' selective single cell transcriptome profiling approach with improved UMI design." *Nucleic Acids Research* 2017, 45(7), e48, in 11 pages.
- Armbrecht, et al. "Single-cell protein profiling in microchambers with barcoded beads", *Microsystems & Nanoengineering*, 2019, 5:55.
- CG000209 Chromium\_NextGEM\_SingleCell\_ATAC\_ReagentKits\_v1.1\_UserGuide\_RevG.
- CG000496\_Chromium\_NextGEM\_SingleCell\_ATAC\_ReagentKits\_v2\_UserGuide\_RevB.
- CG000505\_Chromium\_Nuclei\_Isolation\_Kit\_UG\_RevA.
- Decision of Grant dated Aug. 21, 2023 in Japanese Patent Application 2020-561800.
- Delebecque et al. "Designing and using RNA scaffolds to assemble proteins *in vivo*". *Nature protocols*, 2012, 7(10), 1797-1807.
- Dickey and Giangrande. "Oligonucleotide Aptamers: A Next-Generation Technology for the Capture and Detection of Circulating Tumor Cells." *Methods*, 2016 97:94-103.
- Dua, et al. "Patents on SELEX and therapeutic aptamers. Recent patents on DNA & gene sequences," 2008, 2(3), 172-186.
- Eulberg, et al. "Development of an automated *in vitro* selection protocol to obtain RNA-based aptamers: identification of a biostable substance P antagonist," *Nucleic acids research*, 2005, 33(4), e45. <https://doi.org/10.1093/nar/gni044>.
- Extended European Search Report Dated Oct. 4, 2023 in European Patent Application No. 23166682.9.
- Pathi. P. Design and Characterization of SSDNA Aptamer Candidates to Bind *Bacteroides fragilis* Toxin Subtypes BFT-1 and BFT-2 (Doctoral dissertation, Johns Hopkins University).2017.
- Final Office Action dated May 15, 2023 in U.S. Appl. No. 16/551,638.
- Final Office Action dated May 19, 2023 in U.S. Appl. No. 17/163,177.
- Final Office Action dated May 31, 2023 in U.S. Appl. No. 16/934,530.
- Final Office Action dated Jun. 8, 2023 in U.S. Appl. No. 17/147,283.
- Final Office Action dated Oct. 5, 2023 in U.S. Appl. No. 17/151,050.
- Hoinka and Przytycka. "AptaPLEX-A Dedicated, Multithreaded Demultiplexer for Ht-Se Lex Data." *Methods*, 2016, 106:82-85.
- International Search Report and Written Opinion dated Jun. 5, 2023, in PCT Application No. PCT/US2023/061980.
- International Search Report and Written Opinion dated Jun. 23, 2023 in PCT Application No. PCT/US2023/062070.
- Ku, et al. "Nucleic Acid Aptamers: An Emerging Tool for Biotechnology and Biomedical Sensing." *Sensors* 2015, 15, 16281-16313.
- Mairal et al. "Aptamers: Molecular Tools for Analytical Applications." *Analytical and bioanalytical chemistry* 2008,390: 989-1007.
- Non-Final Office Action dated Jun. 14, 2023 In U.S. Appl. No. 17/174,249.
- Non-Final Office Action dated Jun. 30, 2023 In U.S. Appl. No. 17/684,289.
- Non-Final Office Action dated Sep. 21, 2023 in Canadian Patent Application No. 3,034,924.
- Non-Final Office Action dated Sep. 28, 2023 in U.S. Appl. No. 16/789,311.
- Non-Final Office Action Dated Sep. 28, 2023 in U.S. Appl. No. 17/184,405.
- Non-Final Office Action Dated Oct. 5, 2023 in U.S. Appl. No. 16/848,241.
- Notice of Allowance dated Jun. 8, 2023 in U.S. Appl. No. 16/459,444.
- Notice of Allowance dated Jul. 25, 2023 in European Patent Application No. 20 816 802.1.
- Notice of Allowance dated Aug. 23, 2023 in Canadian Patent Application No. 2,865,575.
- Notice of Allowance dated Aug. 25, 2023 in European Patent Application No. 22 200 785.8.
- Notice of Allowance dated Aug. 28, 2023 in U.S. Appl. No. 16/374,626.
- Notice of Allowance dated Sep. 14, 2023 in Canada Application No. 2982467.
- Notice of Allowance dated Sep. 29, 2023 in European Application No. 22165594.7.
- Office Action dated Feb. 23, 2022 in Chinese Patent Application No. 2016800523302.
- Office Action dated Nov. 24, 2022 in Chinese Patent Application No. 2018800147939.
- Office Action dated Apr. 24, 2023 in European Patent Application No. 21714995.4.
- Office Action dated Apr. 26, 2023 in European Patent Application No. 18703156.2.
- Office Action dated May 16, 2023 in European Patent Application No. 21707112.5.
- Office Action dated May 26, 2023 in Chinese Patent Application No. 2019800373421.
- Office Action dated May 27, 2023 in Chinese Patent Application No. 2019800656859.
- Office Action dated May 30, 2023 in Chinese Patent Application No. 2019800653102.
- Office Action dated Jun. 1, 2023 in Japanese Patent Application No. 2020-561807.
- Office Action dated Jun. 16, 2023 in Chinese Patent Application No. 2019800708938.
- Office Action dated Jun. 22, 2023 in Japanese Patent Application No. 2022-071002.
- Office Action dated Jun. 28, 2023 in European Patent Application 19836239.4.
- Office Action dated Jul. 12, 2023 in Chinese Patent Application No. 2020800212600.
- Office Action dated Jul. 12, 2023 in Canadian Patent Application No. 3,059,559.
- Office Action Dated Jul. 13, 2023 in Chinese Patent Application No. 202080077712.7.
- Office Action dated Jul. 28, 2023 in Chinese Patent Application No. 201880014793.9.
- Office Action dated Jul. 29, 2023 in Chinese Patent Application No. 201980073850.5.
- Office Action dated Jul. 31, 2023 in Chinese Patent Application No. 201980068704.3.
- Office Action dated Jul. 31, 2023 in Chinese Patent Application No. 201980037175.0.
- Office Action dated Aug. 11, 2023 in European Patent Application 19752792.2.
- Office Action dated Aug. 21, 2023 in Japanese Patent Application No. 2021-507836.
- Office Action dated Aug. 30, 2023 in Chinese Patent Application 2019111653930.
- Office Action dated Aug. 31, 2023 in Chinese Patent Application 2020800483617.
- Office Action dated Sep. 21, 2023 in Japanese Patent Application 2022-030956.
- Ogawa, T et al., "The Efficacy and further functional advantages of random-base molecular barcodes for absolute and digital quantification of nucleic acid molecules". *Sci Rep* 7, 2017 12576.
- Restriction Requirement dated Jun. 28, 2023 in U.S. Appl. No. 17/336,055.
- Restriction Requirement dated Oct. 5, 2023 in U.S. Appl. No. 17/373,653.
- Restriction Requirement dated Oct. 11, 2023 in U.S. Appl. No. 17/531,555.
- Summons to Attend Oral Proceedings Dated Aug. 8, 2023 in European Patent Application No. 14749671.5.

(56)

**References Cited**

OTHER PUBLICATIONS

- Wang et al., "Development of Multicolor Flow Cytometry Calibration Standards: Assignment of Equivalent Reference Fluorophores (ERF) Unit" J. Res. Natl. Inst. Stand. Technol. 2011 116, 671-683.
- Wu & Lambowitz, "Facile single-stranded DNA sequencing of human plasma DNA via thermostable group II intron reverse transcriptase template switching," Scientific Reports 2017, 7(8421), 1-14.
- Zheng, et al. "Aptamer-Functionalized Barcode Particles for the Capture and Detection of Multiple Types of Circulating Tumor Cells." Advanced materials (Weinheim), 2014, 26, 7333-7338.
- Zhou and Rossi. "Aptamers as Targeted Therapeutics: Current Potential and Challenges." Nature reviews. Drug discovery, 2017, 16:181-202.
- Zhulidov et al., "Simple cDNA normalization using kamchatka crab duplex-specific nuclease," Nucleic Acids Research. 2004, 32(3)e37.
- Office Action dated Apr. 27, 2025 in Chinese Patent Application No. 202180065616.5.
- Guan et al., "Methods and Applications of Plant Gene Cloning," Science and Technology Press of China, 2009, 21-24.

\* cited by examiner




FIG. 1





FIG. 2

**FIG. 3**




FIG. 4A





FIG. 4B

**FIG. 5A**





FIG. 5B








FIG. 5D

**FIG. 6A**

**FIG. 6B**





FIG. 6C

**FIG. 6D**

**1**
**SINGLE CELL ASSAY FOR  
TRANSPOSOSE-ACCESSIBLE CHROMATIN**
**RELATED APPLICATIONS**

This application claims the benefit under 35 U.S.C. § 119(e) of U.S. Provisional Patent Application Ser. No. 63/059,715, filed Jul. 31, 2020, the content of this related application is incorporated herein by reference in its entirety for all purposes.

**BACKGROUND**
**Field**

The present disclosure relates generally to the field of molecular biology, for example determining open chromatin regions using molecular barcoding.

**Description of the Related Art**

Current technology allows measurement of gene expression of single cells in a massively parallel manner (e.g., >10000 cells) by attaching cell specific oligonucleotide barcodes to poly(A) mRNA molecules from individual cells as each of the cells is co-localized with a barcoded reagent bead in a compartment. There is a need for systems and methods for labeling and identifying open chromatin regions of single cell genomes.

**SUMMARY**

Disclosed herein include methods for labeling DNA. In some embodiments, the method comprises: contacting double-stranded deoxyribonucleic acid (dsDNA) with a transposome to generate a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang, wherein each dsDNA fragment of the plurality of dsDNA fragments comprises a first strand comprising the first 5' overhang and a second strand comprising the second 5' overhang, wherein the transposome comprises a transposase, a first adaptor having the first 5' overhang, and a second adaptor having the second 5' overhang; and barcoding the plurality of dsDNA fragments, or products thereof, using a first plurality of oligonucleotide barcodes to generate a plurality of barcoded DNA fragments, wherein the 3' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes is associated with a solid support, and wherein the 5' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes comprises a first target-binding region capable of hybridizing to the first 5' overhang of at least one of the plurality of dsDNA fragments. In some embodiments, barcoding the plurality of dsDNA fragments, or products thereof, using a first plurality of oligonucleotide barcodes comprises: hybridizing the first 5' overhang of the first strand of a dsDNA fragment with the first target-binding region of an oligonucleotide barcode of the first plurality of oligonucleotide barcodes; and ligating the second strand of said dsDNA fragment to said hybridized oligonucleotide barcode. The method can comprise: before ligating the second strand to the oligonucleotide barcode, filling a gap between the second strand and said hybridized oligonucleotide barcode with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. In some embodiments, ligating the second strand to the oligonucleotide barcode is performed

**2**

using a DNA ligase. In some embodiments, the first 5' overhang comprises a complement of a first target-binding region.

Disclosed herein include methods for labeling DNA. In some embodiments, the method comprises: contacting double-stranded deoxyribonucleic acid (dsDNA) with a transposome to generate a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang, wherein each dsDNA fragment of the plurality of dsDNA fragments comprises a first strand comprising the first 5' overhang and a second strand comprising the second 5' overhang, wherein the transposome comprises a transposase, a first adaptor having the first 5' overhang, and a second adaptor having the second 5' overhang, and wherein the first 5' overhang comprises a coupling sequence; providing a coupling oligonucleotide comprising a 5' complement of the coupling sequence and a 3' complement of a second target-binding region; and barcoding the plurality of dsDNA fragments, or products thereof, using a second plurality of oligonucleotide barcodes to generate a plurality of barcoded DNA fragments, wherein each oligonucleotide barcode of the second plurality of oligonucleotide barcodes comprises the second target-binding region. In some embodiments, barcoding the plurality of dsDNA fragments, or products thereof, using a second plurality of oligonucleotide barcodes comprises: hybridizing the coupling sequence of the first strand of a dsDNA fragment with the 5' complement of the coupling sequence of the coupling oligonucleotide; hybridizing the 3' complement of the second target-binding region of the coupling oligonucleotide with the second target-binding region of an oligonucleotide barcode of the second plurality of oligonucleotide barcodes; and ligating the first strand of said dsDNA fragment to said hybridized oligonucleotide barcode and extending the first strand; and/or ligating the second strand of said dsDNA fragment to said coupling oligonucleotide and extending the second strand. The method can comprise: before ligating the first strand to the oligonucleotide barcode, filling a gap between the first strand and the hybridized oligonucleotide barcode with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. The method can comprise: before ligating the second strand to the coupling oligonucleotide, filling a gap between the second strand and the coupling oligonucleotide with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. In some embodiments, ligating the first strand of said dsDNA fragment to said hybridized oligonucleotide barcode and/or ligating the second strand of said dsDNA fragment to said coupling oligonucleotide is performed with a DNA ligase.

In some embodiments, the coupling oligonucleotide is a single-stranded oligonucleotide. In some embodiments, the coupling oligonucleotide comprises at least 6 nucleotides. In some embodiments, the coupling sequence comprises at least 4 nucleotides. In some embodiments, the plurality of dsDNA fragments comprise one or more gaps. The method can comprise: filling the one or more gaps with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. In some embodiments, the first 5' overhang and/or the second 5' overhang comprises at least 4 nucleotides. In some embodiments, the second 5' overhang comprises a second universal sequence. In some embodiments, the first adaptor and/or the second adaptor comprises a DNA end sequence of the transposon.

In some embodiments, contacting dsDNA with a transposome comprises permeabilizing a cell and/or isolating the nucleus of a cell, wherein the cell comprises the dsDNA. In

some embodiments, the cell comprises a single cell. In some embodiments, the dsDNA comprises genomic DNA (gDNA). In some embodiments, the plurality of dsDNA fragments comprise open chromatin-associated gDNA. In some embodiments, the permeabilizing comprises chemical or physical permeabilization. In some embodiments, the permeabilizing comprises contacting the cell with a detergent and/or a surfactant. In some embodiments, the permeabilizing comprises permeabilizing the cell by sonification. The method can comprise: permeabilizing a nucleus in the cell to generate a permeabilized nucleus. The method can comprise: fixating the cell comprising the nucleus prior to permeabilizing the nucleus. In some embodiments, the transposase comprises a Tn5 transposase.

The method can comprise: obtaining sequence data of the plurality of barcoded DNA fragments, or products thereof. The method can comprise: determining information relating to the gDNA based on the sequences of the plurality of barcoded DNA fragments, or products thereof, in the sequencing data obtained. In some embodiments, determining the information relating to the gDNA comprises determining chromatin accessibility of the gDNA based on the sequences of the plurality of barcoded DNA fragments, or products thereof, in the sequencing data obtained. In some embodiments, determining the chromatin accessibility of the gDNA comprises: aligning the sequences of the plurality of barcoded DNA fragments to a reference sequence of the gDNA; and identifying regions of the gDNA corresponding the ends of the barcoded DNA fragments of the plurality of barcoded DNA fragments to have an accessibility above a threshold. In some embodiments, determining the chromatin accessibility of the gDNA comprises: aligning the sequences of the plurality of barcoded DNA fragments to a reference sequence of the gDNA; and determining the accessibility of regions of the gDNA corresponding the ends of barcoded DNA fragments of the plurality of barcoded DNA fragments based on the numbers of the barcoded DNA fragments of the plurality of barcoded DNA fragments in the sequencing data. In some embodiments, determining the information relating to the gDNA comprises determining methylome information of the gDNA based on the sequences of the plurality of barcoded DNA fragments in the sequencing data obtained. The method can comprise: digesting nucleosomes associated with the double-stranded gDNA. The method can comprise: performing chemical conversion and/or enzymatic conversion of cytosine bases of the plurality of dsDNA fragments, or products thereof, to generate a plurality of converted dsDNA fragments with uracil bases, wherein chemical conversion comprises bisulfate treatment, and wherein enzymatic conversion comprises APOBEC-mediated conversion. In some embodiments, barcoding the plurality of dsDNA fragments, or products thereof, comprises barcoding the plurality of converted dsDNA fragments, or products thereof. In some embodiments, determining the methylome information comprises: determining a position of the plurality of barcoded DNA fragments in the sequencing data has a thymine base and the corresponding position in a reference sequence of the gDNA has a cytosine base to determine the corresponding position in the gDNA has a 5-methylcytosine (5mC) base and/or 5-hydroxymethylcytosine (5hmC) base.

In some embodiments, the cell comprises copies of a nucleic acid target. The method can comprise: contacting a third plurality of oligonucleotide barcodes with the copies of the nucleic acid target for hybridization, wherein each oligonucleotide barcode of the third plurality of oligonucleotide barcodes comprises a first universal sequence, a third

target-binding region capable of hybridizing to the copies of the nucleic acid target, and a molecular label; extending the third plurality of oligonucleotide barcodes hybridized to the copies of a nucleic acid target to generate a plurality of barcoded nucleic acid molecules each comprising a sequence complementary to at least a portion of the nucleic acid target; and obtaining sequence information of the plurality of barcoded nucleic acid molecules, or products thereof, to determine the copy number of the nucleic acid target in the cell. In some embodiments, determining the copy number of the nucleic acid target in the cell comprises determining the copy number of the nucleic acid target in the cell based on the number of molecular labels with distinct sequences associated with the plurality of barcoded nucleic acid molecules, or products thereof.

The method can comprise: contacting random primers with the plurality of barcoded nucleic acid molecules, wherein each of the random primers comprises a third universal sequence, or a complement thereof; and extending the random primers hybridized to the plurality of barcoded nucleic acid molecules to generate a plurality of extension products. The method can comprise: amplifying the plurality of extension products using primers capable of hybridizing to the first universal sequence or complements thereof, and primers capable of hybridizing the third universal sequence or complements thereof, thereby generating a first plurality of barcoded amplicons. In some embodiments, amplifying the plurality of extension products comprises adding sequences of binding sites of sequencing primers and/or sequencing adaptors, complementary sequences thereof, and/or portions thereof, to the plurality of extension products. The method can comprise: determining the copy number of the nucleic acid target in the cell based on the number of molecular labels with distinct sequences associated with the first plurality of barcoded amplicons, or products thereof.

The method can comprise: amplifying the first plurality of barcoded amplicons using primers capable of hybridizing to the first universal sequence or complements thereof, and primers capable of hybridizing the third universal sequence or complements thereof, thereby generating a second plurality of barcoded amplicons. In some embodiments, amplifying the first plurality of barcoded amplicons comprises adding sequences of binding sites of sequencing primers and/or sequencing adaptors, complementary sequences thereof, and/or portions thereof, to the first plurality of barcoded amplicons. The method can comprise: determining the copy number of the nucleic acid target in the cell based on the number of molecular labels with distinct sequences associated with the second plurality of barcoded amplicons, or products thereof. In some embodiments, the first plurality of barcoded amplicons and/or the second plurality of barcoded amplicons comprise whole transcriptome amplification (WTA) products.

The method can comprise: synthesizing a third plurality of barcoded amplicons using the plurality of barcoded nucleic acid molecules as templates to generate a third plurality of barcoded amplicons. In some embodiments, synthesizing a third plurality of barcoded amplicons comprises performing polymerase chain reaction (PCR) amplification of the plurality of the barcoded nucleic acid molecules. In some embodiments, synthesizing a third plurality of barcoded amplicons comprises PCR amplification using primers capable of hybridizing to the first universal sequence, or a complement thereof, and a target-specific primer. The method can comprise: obtaining sequence information of the third plurality of barcoded amplicons, or products thereof. In some embodiments, obtaining the

sequence information comprises attaching sequencing adaptors to the third plurality of barcoded amplicons, or products thereof. The method can comprise: determining the copy number of the nucleic acid target in the cell based on the number of molecular labels with distinct sequences associated with the third plurality of barcoded amplicons, or products thereof.

In some embodiments, the nucleic acid target comprises a nucleic acid molecule. In some embodiments, the nucleic acid molecule comprises ribonucleic acid (RNA), messenger RNA (mRNA), microRNA, small interfering RNA (siRNA), RNA degradation product, RNA comprising a poly(A) tail, a sample indexing oligonucleotide, a cellular component-binding reagent specific oligonucleotide, or any combination thereof. In some embodiments, extending the first, second, and/or third pluralities of oligonucleotide barcodes comprising extending the plurality of oligonucleotide barcodes using a reverse transcriptase and/or a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. In some embodiments, the DNA polymerase comprises a Klenow Fragment. In some embodiments, the reverse transcriptase comprises a viral reverse transcriptase, (e.g., a murine leukemia virus (MLV) reverse transcriptase or a Moloney murine leukemia virus (MMLV) reverse transcriptase). In some embodiments, the first target-binding region, second target-binding region, and/or the third target-binding region comprises a poly(dA) region, a poly(dT) region, a random sequence, a gene-specific sequence, or any combination thereof.

In some embodiments, obtaining sequence data of the plurality of barcoded DNA fragments, or products thereof, comprises attaching sequencing adaptors and/or sequencing primers complementary sequences thereof, and/or portions thereof, to the plurality of barcoded DNA fragments, or products thereof. The method can comprise: amplifying the plurality of barcoded DNA fragments using primers capable of hybridizing to the first universal sequence or complements thereof, and primers capable of hybridizing the second universal sequence or complements thereof, thereby generating a fourth plurality of barcoded DNA fragments. In some embodiments, amplifying the plurality of barcoded DNA fragments comprises adding sequences of binding sites of sequencing primers and/or sequencing adaptors, complementary sequences thereof, and/or portions thereof, to the plurality of barcoded DNA fragments. In some embodiments, obtaining sequence data of the plurality of barcoded DNA fragments, or products thereof, comprises obtaining sequence information of the fourth plurality of barcoded amplicons, or products thereof.

In some embodiments, each oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes comprises a first universal sequence. In some embodiments, the first universal sequence, the second universal sequence, and/or the third universal sequence are the same. In some embodiments, the first universal sequence, the second universal sequence, and/or the third universal sequence are different. In some embodiments, the first universal sequence, the second universal sequence, and/or the third universal sequence comprise the binding sites of sequencing primers and/or sequencing adaptors, complementary sequences thereof, and/or portions thereof. In some embodiments, the sequencing adaptors comprise a P5 sequence, a P7 sequence, complementary sequences thereof, and/or portions thereof. In some embodiments, the sequencing primers comprise a Read 1 sequencing primer, a Read 2 sequencing primer, complementary sequences thereof, and/or portions thereof.

In some embodiments, the first, second, and/or third pluralities of oligonucleotide barcodes each comprise a molecular label. In some embodiments, at least 10 of the first, second, and/or third pluralities of oligonucleotide barcodes comprise different molecular label sequences. In some embodiments, each molecular label of the first, second, and/or third pluralities of oligonucleotide barcodes comprises at least 6 nucleotides.

In some embodiments, the first, second, and/or third pluralities of oligonucleotide barcodes are associated with a solid support. In some embodiments, the first, second, and/or third pluralities of oligonucleotide barcodes associated with the same solid support each comprise an identical sample label. In some embodiments, each sample label of the first, second, and/or third pluralities of oligonucleotide barcodes comprises at least 6 nucleotides. In some embodiments, the first, second, and/or third pluralities of oligonucleotide barcodes each comprise a cell label. In some embodiments, each cell label of the first, second, and/or third pluralities of oligonucleotide barcodes comprises at least 6 nucleotides. In some embodiments, oligonucleotide barcodes of the first, second, and/or third pluralities of oligonucleotide barcodes associated with the same solid support comprise the same cell label. In some embodiments, oligonucleotide barcodes of the first, second, and/or third pluralities of oligonucleotide barcodes associated with different solid supports comprise different cell labels.

In some embodiments, the solid support comprises a synthetic particle, a planar surface, or a combination thereof. The method can comprise: associating a synthetic particle comprising the first, second, and/or third pluralities of oligonucleotide barcodes with the cell. The method can comprise: lysing the cell after associating the synthetic particle with the cell. In some embodiments, lysing the cell comprises heating the cell, contacting the cell with a detergent, changing the pH of the cell, or any combination thereof. In some embodiments, the synthetic particle and the single cell are in the same partition. In some embodiments, the partition is a well or a droplet.

In some embodiments, at least one oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes is immobilized or partially immobilized on the synthetic particle, or at least one oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes is enclosed or partially enclosed in the synthetic particle. In some embodiments, the synthetic particle is disruptable (e.g., a disruptable hydrogel particle). In some embodiments, the synthetic particle comprises a bead. In some embodiments, the bead comprises a sepharose bead, a streptavidin bead, an agarose bead, a magnetic bead, a conjugated bead, a protein A conjugated bead, a protein G conjugated bead, a protein A/G conjugated bead, a protein L conjugated bead, an oligo(dT) conjugated bead, a silica bead, a silica-like bead, an anti-biotin microbead, an anti-fluorochrome microbead, or any combination thereof. In some embodiments, the synthetic particle comprises a material selected from the group consisting of polydimethylsiloxane (PDMS), polystyrene, glass, polypropylene, agarose, gelatin, hydrogel, paramagnetic, ceramic, plastic, glass, methylstyrene, acrylic polymer, titanium, latex, sepharose, cellulose, nylon, silicone, and any combination thereof. In some embodiments, each oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes comprises a linker functional group. In some embodiments, the synthetic particle comprises a solid support functional group. In some embodiments, the support functional group and the linker functional group are associated

with each other. In some embodiments, the linker functional group and the support functional group are individually selected from the group consisting of C6, biotin, streptavidin, primary amine(s), aldehyde(s), ketone(s), and any combination thereof.

Disclosed herein include kits. In some embodiments, the kit comprises: a transposome comprising a transposase, a first adaptor having a first 5' overhang, and a second adaptor having a second 5' overhang; and a first plurality of oligonucleotide barcodes, wherein the 3' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes is associated with a solid support, wherein the 5' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes comprises a first target-binding region capable of hybridizing to the first 5' overhang.

Disclosed herein include kits. In some embodiments, the kit comprises: a transposome comprising a transposase, a first adaptor having a first 5' overhang, and a second adaptor having a second 5' overhang, wherein the first 5' overhang comprises a coupling sequence; and a coupling oligonucleotide comprising a 5' complement of the coupling sequence and a 3' complement of a second target-binding region.

The kit can comprise: a second plurality of oligonucleotide barcodes, wherein each oligonucleotide barcode of the second plurality of oligonucleotide barcodes comprises the second target-binding region. In some embodiments, the coupling oligonucleotide is a single-stranded oligonucleotide. In some embodiments, the coupling oligonucleotide comprises at least 6 nucleotides. In some embodiments, the coupling sequence comprises at least 4 nucleotides. In some embodiments, the first 5' overhang and/or the second 5' overhang comprises at least 4 nucleotides. In some embodiments, the second 5' overhang comprises a second universal sequence. In some embodiments, the first adaptor and/or the second adaptor comprises a DNA end sequence of the transposon. The kit can comprise: a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. In some embodiments, wherein the DNA polymerase comprises a Klenow Fragment. The kit can comprise: a reverse transcriptase. In some embodiments, the reverse transcriptase comprises a viral reverse transcriptase (e.g., a murine leukemia virus (MLV) reverse transcriptase or a Moloney murine leukemia virus (MMLV) reverse transcriptase). The kit can comprise: a ligase. The kit can comprise: a detergent and/or a surfactant. The kit can comprise: a buffer, a cartridge, or both. The kit can comprise: one or more reagents for a reverse transcription reaction and/or an amplification reaction. In some embodiments, the transposase comprises a Tn5 transposase.

The kit can comprise: a third plurality of oligonucleotide barcodes, wherein each oligonucleotide barcode of the third plurality of oligonucleotide barcodes comprises a third target-binding region. In some embodiments, the first target-binding region, second target-binding region, and/or the third target-binding region comprises a poly(dA) region, a poly(dt) region, a random sequence, a gene-specific sequence, or any combination thereof. In some embodiments, the first, second, and/or third pluralities of oligonucleotide barcodes each comprise a molecular label. In some embodiments, the molecular label comprises at least 6 nucleotides. In some embodiments, at least 10 of the first, second, and/or third pluralities of oligonucleotide barcodes comprise different molecular label sequences. In some embodiments, the first, second, and/or third pluralities of oligonucleotide barcodes are associated with a solid support. In some embodiments, the first, second, and/or third pluralities of oligonucleotide barcodes each comprise a cell

label. In some embodiments, oligonucleotide barcodes of the first, second, and/or third pluralities of oligonucleotide barcodes associated with the same solid support comprise the same cell label. In some embodiments, oligonucleotide barcodes of the first, second, and/or third pluralities of oligonucleotide barcodes associated with different solid supports comprise different cell labels.

In some embodiments, the solid support comprises a synthetic particle, a planar surface, or a combination thereof. 10 In some embodiments, at least one oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes is immobilized or partially immobilized on the synthetic particle, or at least one oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes is enclosed or partially enclosed in the synthetic particle. 15 In some embodiments, the synthetic particle is disruptable (e.g., a disruptable hydrogel particle). In some embodiments, the synthetic particle comprises a bead. In some embodiments, the bead comprises a sepharose bead, a streptavidin bead, an agarose bead, a magnetic bead, a conjugated bead, a protein A conjugated bead, a protein G conjugated bead, a protein A/G conjugated bead, a protein L conjugated bead, an oligo(dT) conjugated bead, a silica bead, a silica-like bead, an anti-biotin microbead, an anti-fluorochrome microbead, or any combination thereof. 20 In some embodiments, the synthetic particle comprises a material selected from the group consisting of polydimethylsiloxane (PDMS), polystyrene, glass, polypropylene, agarose, gelatin, hydrogel, paramagnetic, ceramic, plastic, glass, methylstyrene, acrylic polymer, titanium, latex, Sepharose, cellulose, nylon, silicone, and any combination thereof. 25 In some embodiments, each oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes comprises a linker functional group. In some embodiments, the synthetic particle comprises a solid support functional group. In some embodiments, the support functional group and the linker functional group are associated with each other. In some embodiments, the linker functional group and the support functional group are individually selected from the group consisting of C6, biotin, streptavidin, primary amine(s), aldehyde(s), ketone(s), and any combination thereof.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 illustrates a non-limiting exemplary barcode.

FIG. 2 shows a non-limiting exemplary workflow of barcoding and digital counting.

FIG. 3 is a schematic illustration showing a non-limiting exemplary process for generating an indexed library of targets barcoded at the 3'-ends from a plurality of targets.

FIGS. 4A-B are non-limiting schematic illustrations of compositions and methods disclosed herein for labeling DNA (e.g., gDNA) in a single cell.

FIGS. 5A-5D show a schematic illustration of a non-limiting exemplary workflow for labeling DNA (e.g., gDNA) in a single cell.

FIGS. 6A-6D show a schematic illustration of a non-limiting exemplary workflow for labeling DNA (e.g., gDNA) in a single cell.

#### DETAILED DESCRIPTION

In the following detailed description, reference is made to the accompanying drawings, which form a part hereof. In the drawings, similar symbols typically identify similar components, unless context dictates otherwise. The illustra-

tive embodiments described in the detailed description, drawings, and claims are not meant to be limiting. Other embodiments may be utilized, and other changes may be made, without departing from the spirit or scope of the subject matter presented herein. It will be readily understood that the aspects of the present disclosure, as generally described herein, and illustrated in the Figures, can be arranged, substituted, combined, separated, and designed in a wide variety of different configurations, all of which are explicitly contemplated herein and made part of the disclosure herein.

All patents, published patent applications, other publications, and sequences from GenBank, and other databases referred to herein are incorporated by reference in their entirety with respect to the related technology.

Quantifying small numbers of nucleic acids, for example messenger ribonucleotide acid (mRNA) molecules, is clinically important for determining, for example, the genes that are expressed in a cell at different stages of development or under different environmental conditions. However, it can also be very challenging to determine the absolute number of nucleic acid molecules (e.g., mRNA molecules), especially when the number of molecules is very small. One method to determine the absolute number of molecules in a sample is digital polymerase chain reaction (PCR). Ideally, PCR produces an identical copy of a molecule at each cycle. However, PCR can have disadvantages such that each molecule replicates with a stochastic probability, and this probability varies by PCR cycle and gene sequence, resulting in amplification bias and inaccurate gene expression measurements. Stochastic barcodes with unique molecular labels (also referred to as molecular indexes (MIs)) can be used to count the number of molecules and correct for amplification bias. Stochastic barcoding, such as the Precise™ assay (Cellular Research, Inc. (Palo Alto, CA)) and Rhapsody™ assay (Becton, Dickinson and Company (Franklin Lakes, NJ)), can correct for bias induced by PCR and library preparation steps by using molecular labels (MLs) to label mRNAs during reverse transcription (RT).

The Precise™ assay can utilize a non-depleting pool of stochastic barcodes with large number, for example 6561 to 65536, unique molecular label sequences on poly(T) oligonucleotides to hybridize to all poly(A)-mRNAs in a sample during the RT step. A stochastic barcode can comprise a universal PCR priming site. During RT, target gene molecules react randomly with stochastic barcodes. Each target molecule can hybridize to a stochastic barcode resulting to generate stochastically barcoded complementary ribonucleotide acid (cDNA) molecules. After labeling, stochastically barcoded cDNA molecules from microwells of a microwell plate can be pooled into a single tube for PCR amplification and sequencing. Raw sequencing data can be analyzed to produce the number of reads, the number of stochastic barcodes with unique molecular label sequences, and the numbers of mRNA molecules.

Disclosed herein include methods for labeling DNA. In some embodiments, the method comprises: contacting double-stranded deoxyribonucleic acid (dsDNA) with a transposome to generate a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang, wherein each dsDNA fragment of the plurality of dsDNA fragments comprises a first strand comprising the first 5' overhang and a second strand comprising the second 5' overhang, wherein the transposome comprises a transposase, a first adaptor having the first 5' overhang, and a second adaptor having the second 5' overhang; and barcoding the plurality of dsDNA fragments, or products thereof, using a

first plurality of oligonucleotide barcodes to generate a plurality of barcoded DNA fragments, wherein the 3' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes is associated with a solid support, and wherein the 5' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes comprises a first target-binding region capable of hybridizing to the first 5' overhang of at least one of the plurality of dsDNA fragments. In some embodiments, barcoding the plurality of dsDNA fragments, or products thereof, using a first plurality of oligonucleotide barcodes comprises: hybridizing the first 5' overhang of the first strand of a dsDNA fragment with the first target-binding region of an oligonucleotide barcode of the first plurality of oligonucleotide barcodes; and ligating the second strand of said dsDNA fragment to said hybridized oligonucleotide barcode. The method can comprise: before ligating the second strand to the oligonucleotide barcode, filling a gap between the second strand and said hybridized oligonucleotide barcode with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. In some embodiments, ligating the second strand to the oligonucleotide barcode is performed using a DNA ligase. In some embodiments, the first 5' overhang comprises a complement of a first target-binding region.

Disclosed herein include methods for labeling DNA. In some embodiments, the method comprises: contacting double-stranded deoxyribonucleic acid (dsDNA) with a transposome to generate a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang, wherein each dsDNA fragment of the plurality of dsDNA fragments comprises a first strand comprising the first 5' overhang and a second strand comprising the second 5' overhang, wherein the transposome comprises a transposase, a first adaptor having the first 5' overhang, and a second adaptor having the second 5' overhang, and wherein the first 5' overhang comprises a coupling sequence; providing a coupling oligonucleotide comprising a 5' complement of the coupling sequence and a 3' complement of a second target-binding region; and barcoding the plurality of dsDNA fragments, or products thereof, using a second plurality of oligonucleotide barcodes to generate a plurality of barcoded DNA fragments, wherein each oligonucleotide barcode of the second plurality of oligonucleotide barcodes comprises the second target-binding region. In some embodiments, barcoding the plurality of dsDNA fragments, or products thereof, using a second plurality of oligonucleotide barcodes comprises: hybridizing the coupling sequence of the first strand of a dsDNA fragment with the 5' complement of the coupling sequence of the coupling oligonucleotide; hybridizing the 3' complement of the second target-binding region of the coupling oligonucleotide with the second target-binding region of an oligonucleotide barcode of the second plurality of oligonucleotide barcodes; and ligating the first strand of said dsDNA fragment to said hybridized oligonucleotide barcode and extending the first strand; and/or ligating the second strand of said dsDNA fragment to said coupling oligonucleotide and extending the second strand. The method can comprise: before ligating the first strand to the oligonucleotide barcode, filling a gap between the first strand and the hybridized oligonucleotide barcode with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. The method can comprise: before ligating the second strand to the coupling oligonucleotide, filling a gap between the second strand and the coupling oligonucleotide with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5'

**11**

exonuclease activity. In some embodiments, ligating the first strand of said dsDNA fragment to said hybridized oligonucleotide barcode and/or ligating the second strand of said dsDNA fragment to said coupling oligonucleotide is performed with a DNA ligase.

Disclosed herein include kits. In some embodiments, the kit comprises: a transposome comprising a transposase, a first adaptor having a first 5' overhang, and a second adaptor having a second 5' overhang; and a first plurality of oligonucleotide barcodes, wherein the 3' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes is associated with a solid support, wherein the 5' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes comprises a first target-binding region capable of hybridizing to the first 5' overhang.

Disclosed herein include kits. In some embodiments, the kit comprises: a transposome comprising a transposase, a first adaptor having a first 5' overhang, and a second adaptor having a second 5' overhang, wherein the first 5' overhang comprises a coupling sequence; and a coupling oligonucleotide comprising a 5' complement of the coupling sequence and a 3' complement of a second target-binding region.

**Definitions**

Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the present disclosure belongs. See, e.g., Singleton et al., Dictionary of Microbiology and Molecular Biology 2nd ed., J. Wiley & Sons (New York, NY 1994); Sambrook et al., Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press (Cold Spring Harbor, NY 1989). For purposes of the present disclosure, the following terms are defined below.

As used herein, the term "adaptor" refers to a sequence that can facilitate amplification or sequencing of associated nucleic acids. The associated nucleic acids can comprise target nucleic acids. The associated nucleic acids can comprise one or more of spatial labels, target labels, sample labels, indexing label, or barcode sequences (e.g., molecular labels). The adaptors can be linear. The adaptors can be preadenylated adaptors. The adaptors can be double- or single-stranded. One or more adaptor can be located on the 5' or 3' end of a nucleic acid. When the adaptors comprise known sequences on the 5' and 3' ends, the known sequences can be the same or different sequences. An adaptor located on the 5' and/or 3' ends of a polynucleotide can be capable of hybridizing to one or more oligonucleotides immobilized on a surface. An adaptor can, in some embodiments, comprise a universal sequence. A universal sequence can be a region of nucleotide sequence that is common to two or more nucleic acid molecules. The two or more nucleic acid molecules can also have regions of different sequence. Thus, for example, the 5' adaptors can comprise identical and/or universal nucleic acid sequences and the 3' adaptors can comprise identical and/or universal sequences. A universal sequence that may be present in different members of a plurality of nucleic acid molecules can allow the replication or amplification of multiple different sequences using a single universal primer that is complementary to the universal sequence. Similarly, at least one, two (e.g., a pair) or more universal sequences that may be present in different members of a collection of nucleic acid molecules can allow the replication or amplification of multiple different sequences using at least one, two (e.g., a pair) or more single universal primers that are complementary to the universal sequences. Thus, a universal primer includes a sequence that can hybridize to such a universal sequence. The target nucleic acid sequence-bearing molecules may be modified to

**12**

attach universal adaptors (e.g., non-target nucleic acid sequences) to one or both ends of the different target nucleic acid sequences. The one or more universal primers attached to the target nucleic acid can provide sites for hybridization of universal primers. The one or more universal primers attached to the target nucleic acid can be the same or different from each other.

As used herein, the term "associated" or "associated with" mean that two or more species are identifiable as being co-located at a point in time. An association can mean that two or more species are or were within a similar container. An association can be an informatics association. For example, digital information regarding two or more species can be stored and can be used to determine that one or more of the species were co-located at a point in time. An association can also be a physical association. In some embodiments, two or more associated species are "tethered", "attached", or "immobilized" to one another or to a common solid or semisolid surface. An association may refer to covalent or non-covalent means for attaching labels to solid or semi-solid supports such as beads. An association may be a covalent bond between a target and a label. An association can comprise hybridization between two molecules (such as a target molecule and a label).

As used herein, the term "complementary" refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a given position of a nucleic acid is capable of hydrogen bonding with a nucleotide of another nucleic acid, then the two nucleic acids are considered to be complementary to one another at that position. Complementarity between two single-stranded nucleic acid molecules may be "partial," in which only some of the nucleotides bind, or it may be complete when total complementarity exists between the single-stranded molecules. A first nucleotide sequence can be said to be the "complement" of a second sequence if the first nucleotide sequence is complementary to the second nucleotide sequence. A first nucleotide sequence can be said to be the "reverse complement" of a second sequence, if the first nucleotide sequence is complementary to a sequence that is the reverse (i.e., the order of the nucleotides is reversed) of the second sequence. As used herein, a "complementary" sequence can refer to a "complement" or a "reverse complement" of a sequence. It is understood from the disclosure that if a molecule can hybridize to another molecule it may be complementary, or partially complementary, to the molecule that is hybridizing.

As used herein, the term "digital counting" refers to a method for estimating a number of target molecules in a sample. Digital counting can include the step of determining a number of unique labels that have been associated with targets in a sample. This methodology, which can be stochastic in nature, transforms the problem of counting molecules from one of locating and identifying identical molecules to a series of yes/no digital questions regarding detection of a set of predefined labels.

As used herein, the term "label" or "labels" refers to nucleic acid codes associated with a target within a sample. A label can be, for example, a nucleic acid label. A label can be an entirely or partially amplifiable label. A label can be entirely or partially sequencable label. A label can be a portion of a native nucleic acid that is identifiable as distinct. A label can be a known sequence. A label can comprise a junction of nucleic acid sequences, for example a junction of a native and non-native sequence. As used herein, the term "label" can be used interchangeably with the terms, "index", "tag," or "label-tag." Labels can convey information. For example, in various embodiments, labels can be used to

determine an identity of a sample, a source of a sample, an identity of a cell, and/or a target.

As used herein, the term “non-depleting reservoirs” refers to a pool of barcodes (e.g., stochastic barcodes) made up of many different labels. A non-depleting reservoir can comprise large numbers of different barcodes such that when the non-depleting reservoir is associated with a pool of targets each target is likely to be associated with a unique barcode. The uniqueness of each labeled target molecule can be determined by the statistics of random choice, and depends on the number of copies of identical target molecules in the collection compared to the diversity of labels. The size of the resulting set of labeled target molecules can be determined by the stochastic nature of the barcoding process, and analysis of the number of barcodes detected then allows calculation of the number of target molecules present in the original collection or sample. When the ratio of the number of copies of a target molecule present to the number of unique barcodes is low, the labeled target molecules are highly unique (i.e., there is a very low probability that more than one target molecule will have been labeled with a given label).

As used herein, the term “nucleic acid” refers to a polynucleotide sequence, or fragment thereof. A nucleic acid can comprise nucleotides. A nucleic acid can be exogenous or endogenous to a cell. A nucleic acid can exist in a cell-free environment. A nucleic acid can be a gene or fragment thereof. A nucleic acid can be DNA. A nucleic acid can be RNA. A nucleic acid can comprise one or more analogs (e.g., altered backbone, sugar, or nucleobase). Some non-limiting examples of analogs include: 5-bromouracil, peptide nucleic acid, xeno nucleic acid, morpholinos, locked nucleic acids, glycol nucleic acids, threose nucleic acids, dideoxynucleotides, cordycepin, 7-deaza-GTP, fluorophores (e.g., rhodamine or fluorescein linked to the sugar), thiol containing nucleotides, biotin linked nucleotides, fluorescent base analogs, CpG islands, methyl-7-guanosine, methylated nucleotides, inosine, thiouridine, pseudouridine, dihydrouridine, queuosine, and wyosine. “Nucleic acid”, “polynucleotide”, “target polynucleotide”, and “target nucleic acid” can be used interchangeably.

A nucleic acid can comprise one or more modifications (e.g., a base modification, a backbone modification), to provide the nucleic acid with a new or enhanced feature (e.g., improved stability). A nucleic acid can comprise a nucleic acid affinity tag. A nucleoside can be a base-sugar combination. The base portion of the nucleoside can be a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides can be nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to the 2', the 3', or the 5' hydroxyl moiety of the sugar. In forming nucleic acids, the phosphate groups can covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn, the respective ends of this linear polymeric compound can be further joined to form a circular compound; however, linear compounds are generally suitable. In addition, linear compounds may have internal nucleotide base complementarity and may therefore fold in a manner as to produce a fully or partially double-stranded compound. Within nucleic acids, the phosphate groups can commonly be referred to as forming the internucleoside backbone of the nucleic acid. The linkage or backbone can be a 3' to 5' phosphodiester linkage.

A nucleic acid can comprise a modified backbone and/or modified internucleoside linkages. Modified backbones can include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. Suitable modified nucleic acid backbones containing a phosphorus atom therein can include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkyl phosphotriesters, methyl and other alkyl phosphonate such as 3'-alkylene phosphonates, 5'-alkylene phosphonates, chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkyl phosphoramidates, phosphorodiamidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, selenophosphates, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', a 5' to 5' or a 2' to 2' linkage.

A nucleic acid can comprise polynucleotide backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These can include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts.

A nucleic acid can comprise a nucleic acid mimetic. The term “mimetic” can be intended to include polynucleotides wherein only the furanose ring or both the furanose ring and the internucleotide linkage are replaced with non-furanose groups, replacement of only the furanose ring can also be referred as being a sugar surrogate. The heterocyclic base moiety or a modified heterocyclic base moiety can be maintained for hybridization with an appropriate target nucleic acid. One such nucleic acid can be a peptide nucleic acid (PNA). In a PNA, the sugar-backbone of a polynucleotide can be replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleotides can be retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. The backbone in PNA compounds can comprise two or more linked aminoethylglycine units which gives PNA an amide containing backbone. The heterocyclic base moieties can be bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone.

A nucleic acid can comprise a morpholino backbone structure. For example, a nucleic acid can comprise a 6-membered morpholino ring in place of a ribose ring. In some of these embodiments, a phosphorodiamidate or other non-phosphodiester internucleoside linkage can replace a phosphodiester linkage.

A nucleic acid can comprise linked morpholino units (e.g., morpholino nucleic acid) having heterocyclic bases attached to the morpholino ring. Linking groups can link the morpholino monomeric units in a morpholino nucleic acid. Non-ionic morpholino-based oligomeric compounds can have less undesired interactions with cellular proteins. Morpholino-based polynucleotides can be nonionic mimics of nucleic acids. A variety of compounds within the morpholino class can be joined using different linking groups. A further class of polynucleotide mimetic can be referred to as

cyclohexenyl nucleic acids (CeNA). The furanose ring normally present in a nucleic acid molecule can be replaced with a cyclohexenyl ring. CeNA DMT protected phosphoramidite monomers can be prepared and used for oligomeric compound synthesis using phosphoramidite chemistry. The incorporation of CeNA monomers into a nucleic acid chain can increase the stability of a DNA/RNA hybrid. CeNA oligoadenylylates can form complexes with nucleic acid complements with similar stability to the native complexes. A further modification can include Locked Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 4' carbon atom of the sugar ring thereby forming a 2'-C, 4'-C-oxymethylene linkage thereby forming a bicyclic sugar moiety. The linkage can be a methylene ( $-\text{CH}_2-$ ) group bridging the 2' oxygen atom and the 4' carbon atom wherein n is 1 or 2. LNA and LNA analogs can display very high duplex thermal stabilities with complementary nucleic acid ( $T_m=+3$  to  $+10^\circ \text{ C}.$ ), stability towards 3'-exonucleaseolytic degradation and good solubility properties.

A nucleic acid may also include nucleobase (often referred to simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases can include the purine bases, (e.g., adenine (A) and guanine (G)), and the pyrimidine bases, (e.g., thymine (T), cytosine (C) and uracil (U)). Modified nucleobases can include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl ( $-\text{C}\equiv\text{C}-\text{CH}_3$ ) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-aminoadenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Modified nucleobases can include tricyclic pyrimidines such as phenoxazine cytidine(1H-pyrimido(5,4-b)(1,4)benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido(5,4-b)(1,4)benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g., 9-(2-aminoethoxy)-H-pyrimido(5,4-b)(1,4)benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido(5,4-b)(1,4)benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g., 9-(2-aminoethoxy)-H-pyrimido(5,4-b)(1,4)benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido(4,5-b)indol-2-one), pyridoindole cytidine (H-pyrido(3',2':4,5)pyrrolo[2,3-d]pyrimidin-2-one).

As used herein, the term "sample" refers to a composition comprising targets. Suitable samples for analysis by the disclosed methods, devices, and systems include cells, tissues, organs, or organisms.

As used herein, the term "sampling device" or "device" refers to a device which may take a section of a sample and/or place the section on a substrate. A sample device can refer to, for example, a fluorescence activated cell sorting (FACS) machine, a cell sorter machine, a biopsy needle, a biopsy device, a tissue sectioning device, a microfluidic device, a blade grid, and/or a microtome.

As used herein, the term "solid support" refers to discrete solid or semi-solid surfaces to which a plurality of barcodes (e.g., stochastic barcodes) may be attached. A solid support may encompass any type of solid, porous, or hollow sphere,

ball, bearing, cylinder, or other similar configuration composed of plastic, ceramic, metal, or polymeric material (e.g., hydrogel) onto which a nucleic acid may be immobilized (e.g., covalently or non-covalently). A solid support may comprise a discrete particle that may be spherical (e.g., microspheres) or have a non-spherical or irregular shape, such as cubic, cuboid, pyramidal, cylindrical, conical, oblong, or disc-shaped, and the like. A bead can be non-spherical in shape. A plurality of solid supports spaced in an array may not comprise a substrate. A solid support may be used interchangeably with the term "bead."

As used herein, the term "stochastic barcode" refers to a polynucleotide sequence comprising labels of the present disclosure. A stochastic barcode can be a polynucleotide sequence that can be used for stochastic barcoding. Stochastic barcodes can be used to quantify targets within a sample. Stochastic barcodes can be used to control for errors which may occur after a label is associated with a target. For example, a stochastic barcode can be used to assess amplification or sequencing errors. A stochastic barcode associated with a target can be called a stochastic barcode-target or stochastic barcode-tag-target.

As used herein, the term "gene-specific stochastic barcode" refers to a polynucleotide sequence comprising labels and a target-binding region that is gene-specific. A stochastic barcode can be a polynucleotide sequence that can be used for stochastic barcoding. Stochastic barcodes can be used to quantify targets within a sample. Stochastic barcodes can be used to control for errors which may occur after a label is associated with a target. For example, a stochastic barcode can be used to assess amplification or sequencing errors. A stochastic barcode associated with a target can be called a stochastic barcode-target or stochastic barcode-tag-target.

As used herein, the term "stochastic barcoding" refers to the random labeling (e.g., barcoding) of nucleic acids. Stochastic barcoding can utilize a recursive Poisson strategy to associate and quantify labels associated with targets. As used herein, the term "stochastic barcoding" can be used interchangeably with "stochastic labeling."

As used here, the term "target" refers to a composition which can be associated with a barcode (e.g., a stochastic barcode). Exemplary suitable targets for analysis by the disclosed methods, devices, and systems include oligonucleotides, DNA, RNA, mRNA, microRNA, tRNA, and the like. Targets can be single or double stranded. In some embodiments, targets can be proteins, peptides, or polypeptides. In some embodiments, targets are lipids. As used herein, "target" can be used interchangeably with "species."

As used herein, the term "reverse transcriptases" refers to a group of enzymes having reverse transcriptase activity (i.e., that catalyze synthesis of DNA from an RNA template). In general, such enzymes include, but are not limited to, retroviral reverse transcriptase, retrotransposon reverse transcriptase, retroplasmid reverse transcriptases, retrorv reverse transcriptases, bacterial reverse transcriptases, group II intron-derived reverse transcriptase, and mutants, variants or derivatives thereof. Non-retroviral reverse transcriptases include non-LTR retrotransposon reverse transcriptases, retroplasmid reverse transcriptases, retrorv reverse transcriptases, and group II intron reverse transcriptases. Examples of group II intron reverse transcriptases include the *Lactococcus lactis* LI.LtrB intron reverse transcriptase, the *Thermosynechococcus elongatus* Tel4c intron reverse transcriptase, or the *Geobacillus stearothermophilus* Gs-IIC intron reverse transcriptase. Other classes of reverse transcriptases can include many classes of non-retroviral

reverse transcriptases (i.e., retrons, group II introns, and diversity-generating retroelements among others).

The terms “universal adaptor primer,” “universal primer adaptor” or “universal adaptor sequence” are used interchangeably herein to refer to a nucleotide sequence that can be used to hybridize to barcodes (e.g., stochastic barcodes) to generate gene-specific barcodes. A universal adaptor sequence can, for example, be a known sequence that is universal across all barcodes used in methods of the disclosure. For example, when multiple targets are being labeled using the methods disclosed herein, each of the target-specific sequences may be linked to the same universal adaptor sequence. In some embodiments, more than one universal adaptor sequences may be used in the methods disclosed herein. For example, when multiple targets are being labeled using the methods disclosed herein, at least two of the target-specific sequences are linked to different universal adaptor sequences. A universal adaptor primer and its complement may be included in two oligonucleotides, one of which comprises a target-specific sequence and the other comprises a barcode. For example, a universal adaptor sequence may be part of an oligonucleotide comprising a target-specific sequence to generate a nucleotide sequence that is complementary to a target nucleic acid. A second oligonucleotide comprising a barcode and a complementary sequence of the universal adaptor sequence may hybridize with the nucleotide sequence and generate a target-specific barcode (e.g., a target-specific stochastic barcode). In some embodiments, a universal adaptor primer has a sequence that is different from a universal PCR primer used in the methods of this disclosure.

#### Barcodes

Barcodeing, such as stochastic barcoding, has been described in, for example, Fu et al., *Proc Natl Acad Sci U.S.A.*, 2011 May 31, 108(22):9026-31; U.S. Patent Application Publication No. US2011/0160078; Fan et al., *Science*, 2015 Feb. 6, 347(6222):1258367; US Patent Application Publication No. US2015/0299784; and PCT Application Publication No. WO2015/031691; the content of each of these, including any supporting or supplemental information or material, is incorporated herein by reference in its entirety. In some embodiments, the barcode disclosed herein can be a stochastic barcode which can be a polynucleotide sequence that may be used to stochastically label (e.g., barcode, tag) a target. Barcodes can be referred to stochastic barcodes if the ratio of the number of different barcode sequences of the stochastic barcodes and the number of occurrence of any of the targets to be labeled can be, or be about, 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 11:1, 12:1, 13:1, 14:1, 15:1, 16:1, 17:1, 18:1, 19:1, 20:1, 30:1, 40:1, 50:1, 60:1, 70:1, 80:1, 90:1, 100:1, or a number or a range between any two of these values. A target can be an mRNA species comprising mRNA molecules with identical or nearly identical sequences. Barcodes can be referred to as stochastic barcodes if the ratio of the number of different barcode sequences of the stochastic barcodes and the number of occurrence of any of the targets to be labeled is at least, or is at most, 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 11:1, 12:1, 13:1, 14:1, 15:1, 16:1, 17:1, 18:1, 19:1, 20:1, 30:1, 40:1, 50:1, 60:1, 70:1, 80:1, 90:1, or 100:1. Barcode sequences of stochastic barcodes can be referred to as molecular labels.

A barcode, for example a stochastic barcode, can comprise one or more labels. Exemplary labels can include a universal label, a cell label, a barcode sequence (e.g., a molecular label), a sample label, a plate label, a spatial label, and/or a pre-spatial label. FIG. 1 illustrates an exemplary

barcode 104 with a spatial label. The barcode 104 can comprise a 5' amine that may link the barcode to a solid support 105. The barcode can comprise a universal label, a dimension label, a spatial label, a cell label, and/or a molecular label. The order of different labels (including but not limited to the universal label, the dimension label, the spatial label, the cell label, and the molecule label) in the barcode can vary. For example, as shown in FIG. 1, the universal label may be the 5'-most label, and the molecular label may be the 3'-most label. The spatial label, dimension label, and the cell label may be in any order. In some embodiments, the universal label, the spatial label, the dimension label, the cell label, and the molecular label are in any order. The barcode can comprise a target-binding region. The target-binding region can interact with a target (e.g., target nucleic acid, RNA, mRNA, DNA) in a sample. For example, a target-binding region can comprise an oligo (dT) sequence which can interact with poly(A) tails of mRNAs. In some instances, the labels of the barcode (e.g., universal label, dimension label, spatial label, cell label, and barcode sequence) may be separated by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 or more nucleotides.

A label, for example the cell label, can comprise a unique set of nucleic acid sub-sequences of defined length, e.g., seven nucleotides each (equivalent to the number of bits used in some Hamming error correction codes), which can be designed to provide error correction capability. The set of error correction sub-sequences comprise seven nucleotide sequences can be designed such that any pairwise combination of sequences in the set exhibits a defined “genetic distance” (or number of mismatched bases), for example, a set of error correction sub-sequences can be designed to exhibit a genetic distance of three nucleotides. In this case, review of the error correction sequences in the set of sequence data for labeled target nucleic acid molecules (described more fully below) can allow one to detect or correct amplification or sequencing errors. In some embodiments, the length of the nucleic acid sub-sequences used for creating error correction codes can vary, for example, they can be, or be about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 31, 40, 50, or a number or a range between any two of these values, nucleotides in length. In some embodiments, nucleic acid sub-sequences of other lengths can be used for creating error correction codes.

The barcode can comprise a target-binding region. The target-binding region can interact with a target in a sample. The target can be, or comprise, ribonucleic acids (RNAs), messenger RNAs (mRNAs), microRNAs, small interfering RNAs (siRNAs), RNA degradation products, RNAs each comprising a poly(A) tail, or any combination thereof. In some embodiments, the plurality of targets can include deoxyribonucleic acids (DNAs).

In some embodiments, a target-binding region can comprise an oligo(dT) sequence which can interact with poly(A) tails of mRNAs. One or more of the labels of the barcode (e.g., the universal label, the dimension label, the spatial label, the cell label, and the barcode sequences (e.g., molecular label)) can be separated by a spacer from another one or two of the remaining labels of the barcode. The spacer can be, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20, or more nucleotides. In some embodiments, none of the labels of the barcode is separated by spacer.

#### Universal Labels

A barcode can comprise one or more universal labels. In some embodiments, the one or more universal labels can be

the same for all barcodes in the set of barcodes attached to a given solid support. In some embodiments, the one or more universal labels can be the same for all barcodes attached to a plurality of beads. In some embodiments, a universal label can comprise a nucleic acid sequence that is capable of hybridizing to a sequencing primer. Sequencing primers can be used for sequencing barcodes comprising a universal label. Sequencing primers (e.g., universal sequencing primers) can comprise sequencing primers associated with high-throughput sequencing platforms. In some embodiments, a universal label can comprise a nucleic acid sequence that is capable of hybridizing to a PCR primer. In some embodiments, the universal label can comprise a nucleic acid sequence that is capable of hybridizing to a sequencing primer and a PCR primer. The nucleic acid sequence of the universal label that is capable of hybridizing to a sequencing or PCR primer can be referred to as a primer binding site. A universal label can comprise a sequence that can be used to initiate transcription of the barcode. A universal label can comprise a sequence that can be used for extension of the barcode or a region within the barcode. A universal label can be, or be about, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or a number or a range between any two of these values, nucleotides in length. For example, a universal label can comprise at least about 10 nucleotides. A universal label can be at least, or be at most, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 100, 200, or 300 nucleotides in length. In some embodiments, a cleavable linker or modified nucleotide can be part of the universal label sequence to enable the barcode to be cleaved off from the support.

#### Dimension Labels

A barcode can comprise one or more dimension labels. In some embodiments, a dimension label can comprise a nucleic acid sequence that provides information about a dimension in which the labeling (e.g., stochastic labeling) occurred. For example, a dimension label can provide information about the time at which a target was barcoded. A dimension label can be associated with a time of barcoding (e.g., stochastic barcoding) in a sample. A dimension label can be activated at the time of labeling. Different dimension labels can be activated at different times. The dimension label provides information about the order in which targets, groups of targets, and/or samples were barcoded. For example, a population of cells can be barcoded at the G0 phase of the cell cycle. The cells can be pulsed again with barcodes (e.g., stochastic barcodes) at the G1 phase of the cell cycle. The cells can be pulsed again with barcodes at the S phase of the cell cycle, and so on. Barcodes at each pulse (e.g., each phase of the cell cycle), can comprise different dimension labels. In this way, the dimension label provides information about which targets were labelled at which phase of the cell cycle. Dimension labels can interrogate many different biological times. Exemplary biological times can include, but are not limited to, the cell cycle, transcription (e.g., transcription initiation), and transcript degradation. In another example, a sample (e.g., a cell, a population of cells) can be labeled before and/or after treatment with a drug and/or therapy. The changes in the number of copies of distinct targets can be indicative of the sample's response to the drug and/or therapy.

A dimension label can be activatable. An activatable dimension label can be activated at a specific time point. The activatable label can be, for example, constitutively activated (e.g., not turned off). The activatable dimension label can be, for example, reversibly activated (e.g., the activatable dimension label can be turned on and turned off). The dimension label can be, for example, reversibly activatable

at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more times. The dimension label can be reversibly activatable, for example, at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more times. In some embodiments, the dimension label can be activated with fluorescence, light, a chemical event (e.g., cleavage, ligation of another molecule, addition of modifications (e.g., pegylated, sumoylated, acetylated, methylated, deacetylated, demethylated), a photochemical event (e.g., photocaging), and introduction of a non-natural nucleotide.

The dimension label can, in some embodiments, be identical for all barcodes (e.g., stochastic barcodes) attached to a given solid support (e.g., a bead), but different for different solid supports (e.g., beads). In some embodiments, at least 60%, 70%, 80%, 85%, 90%, 95%, 97%, 99% or 100% of barcodes on the same solid support can comprise the same dimension label. In some embodiments, at least 60% of barcodes on the same solid support can comprise the same dimension label. In some embodiments, at least 95% of barcodes on the same solid support can comprise the same dimension label.

There can be as many as  $10^6$  or more unique dimension label sequences represented in a plurality of solid supports (e.g., beads). A dimension label can be, or be about 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or a number or a range between any two of these values, nucleotides in length. A dimension label can be at least, or be at most, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 100, 200, or 300 nucleotides in length. A dimension label can comprise between about 5 to about 200 nucleotides. A dimension label can comprise between about 10 to about 150 nucleotides. A dimension label can comprise between about 20 to about 125 nucleotides in length.

#### Spatial Labels

A barcode can comprise one or more spatial labels. In some embodiments, a spatial label can comprise a nucleic acid sequence that provides information about the spatial orientation of a target molecule which is associated with the barcode. A spatial label can be associated with a coordinate in a sample. The coordinate can be a fixed coordinate. For example, a coordinate can be fixed in reference to a substrate. A spatial label can be in reference to a two or three-dimensional grid. A coordinate can be fixed in reference to a landmark. The landmark can be identifiable in space. A landmark can be a structure which can be imaged. A landmark can be a biological structure, for example an anatomical landmark. A landmark can be a cellular landmark, for instance an organelle. A landmark can be a non-natural landmark such as a structure with an identifiable identifier such as a color code, bar code, magnetic property, fluorescents, radioactivity, or a unique size or shape. A spatial label can be associated with a physical partition (e.g., A well, a container, or a droplet). In some embodiments, multiple spatial labels are used together to encode one or more positions in space.

The spatial label can be identical for all barcodes attached to a given solid support (e.g., a bead), but different for different solid supports (e.g., beads). In some embodiments, the percentage of barcodes on the same solid support comprising the same spatial label can be, or be about, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 99%, 100%, or a number or a range between any two of these values. In some embodiments, the percentage of barcodes on the same solid support comprising the same spatial label can be at least, or be at most, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 99%, or 100%. In some embodiments, at least 60% of barcodes on the same solid support can comprise the same spatial label.

21

In some embodiments, at least 95% of barcodes on the same solid support can comprise the same spatial label.

There can be as many as  $10^6$  or more unique spatial label sequences represented in a plurality of solid supports (e.g., beads). A spatial label can be, or be about, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or a number or a range between any two of these values, nucleotides in length. A spatial label can be at least or at most 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 100, 200, or 300 nucleotides in length. A spatial label can comprise between about 5 to about 200 nucleotides. A spatial label can comprise between about 10 to about 150 nucleotides. A spatial label can comprise between about 20 to about 125 nucleotides in length.

#### Cell Labels

A barcode (e.g., a stochastic barcode) can comprise one or more cell labels. In some embodiments, a cell label can comprise a nucleic acid sequence that provides information for determining which target nucleic acid originated from which cell. In some embodiments, the cell label is identical for all barcodes attached to a given solid support (e.g., a bead), but different for different solid supports (e.g., beads). In some embodiments, the percentage of barcodes on the same solid support comprising the same cell label can be, or be about 60%, 70%, 80%, 85%, 90%, 95%, 97%, 99%, 100%, or a number or a range between any two of these values. In some embodiments, the percentage of barcodes on the same solid support comprising the same cell label can be, or be about 60%, 70%, 80%, 85%, 90%, 95%, 97%, 99%, or 100%. For example, at least 60% of barcodes on the same solid support can comprise the same cell label. As another example, at least 95% of barcodes on the same solid support can comprise the same cell label.

There can be as many as  $10^6$  or more unique cell label sequences represented in a plurality of solid supports (e.g., beads). A cell label can be, or be about, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or a number or a range between any two of these values, nucleotides in length. A cell label can be at least, or be at most, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 100, 200, or 300 nucleotides in length. For example, a cell label can comprise between about 5 to about 200 nucleotides. As another example, a cell label can comprise between about 10 to about 150 nucleotides. As yet another example, a cell label can comprise between about 20 to about 125 nucleotides in length.

#### Barcode Sequences

A barcode can comprise one or more barcode sequences. In some embodiments, a barcode sequence can comprise a nucleic acid sequence that provides identifying information for the specific type of target nucleic acid species hybridized to the barcode. A barcode sequence can comprise a nucleic acid sequence that provides a counter (e.g., that provides a rough approximation) for the specific occurrence of the target nucleic acid species hybridized to the barcode (e.g., target-binding region).

In some embodiments, a diverse set of barcode sequences are attached to a given solid support (e.g., a bead). In some embodiments, there can be, or be about,  $10^2$ ,  $10^3$ ,  $10^4$ ,  $10^5$ ,  $10^6$ ,  $10^7$ ,  $10^8$ ,  $10^9$ , or a number or a range between any two of these values, unique molecular label sequences. For example, a plurality of barcodes can comprise about 6561 barcode sequences with distinct sequences. As another example, a plurality of barcodes can comprise about 65536 barcode sequences with distinct sequences. In some embodiments, there can be at least, or be at most,  $10^2$ ,  $10^3$ ,  $10^4$ ,  $10^5$ ,  $10^6$ ,  $10^7$ ,  $10^8$ , or  $10^9$ , unique barcode sequences. The unique molecular label sequences can be attached to a given solid support (e.g., a bead). In some embodiments, the unique

22

molecular label sequence is partially or entirely encompassed by a particle (e.g., a hydrogel bead).

The length of a barcode can be different in different implementations. For example, a barcode can be, or be about, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or a number or a range between any two of these values, nucleotides in length. As another example, a barcode can be at least, or be at most, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 100, 200, or 300 nucleotides in length.

#### 10 Molecular Labels

A barcode (e.g., a stochastic barcode) can comprise one or more molecular labels. Molecular labels can include barcode sequences. In some embodiments, a molecular label can comprise a nucleic acid sequence that provides identifying information for the specific type of target nucleic acid species hybridized to the barcode. A molecular label can comprise a nucleic acid sequence that provides a counter for the specific occurrence of the target nucleic acid species hybridized to the barcode (e.g., target-binding region).

In some embodiments, a diverse set of molecular labels are attached to a given solid support (e.g., a bead). In some embodiments, there can be, or be about,  $10^2$ ,  $10^3$ ,  $10^4$ ,  $10^5$ ,  $10^6$ ,  $10^7$ ,  $10^8$ ,  $10^9$ , or a number or a range between any two of these values, of unique molecular label sequences. For example, a plurality of barcodes can comprise about 6561 molecular labels with distinct sequences. As another example, a plurality of barcodes can comprise about 65536 molecular labels with distinct sequences. In some embodiments, there can be at least, or be at most,  $10^2$ ,  $10^3$ ,  $10^4$ ,  $10^5$ ,  $10^6$ ,  $10^7$ ,  $10^8$ , or  $10^9$ , unique molecular label sequences. Barcodes with unique molecular label sequences can be attached to a given solid support (e.g., a bead).

For barcoding (e.g., stochastic barcoding) using a plurality of stochastic barcodes, the ratio of the number of different molecular label sequences and the number of occurrence of any of the targets can be, or be about, 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 11:1, 12:1, 13:1, 14:1, 15:1, 16:1, 17:1, 18:1, 19:1, 20:1, 30:1, 40:1, 50:1, 60:1, 70:1, 80:1, 90:1, 100:1, or a number or a range between any two of these values. A target can be an mRNA species comprising mRNA molecules with identical or nearly identical sequences. In some embodiments, the ratio of the number of different molecular label sequences and the number of occurrence of any of the targets is at least, or is at most, 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 11:1, 12:1, 13:1, 14:1, 15:1, 16:1, 17:1, 18:1, 19:1, 20:1, 30:1, 40:1, 50:1, 60:1, 70:1, 80:1, 90:1, or 100:1.

A molecular label can be, or be about, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or a number or a range between any two of these values, nucleotides in length. A molecular label can be at least, or be at most, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 100, 200, or 300 nucleotides in length.

#### Target-Binding Region

A barcode can comprise one or more target binding regions, such as capture probes. In some embodiments, a target-binding region can hybridize with a target of interest. In some embodiments, the target binding regions can comprise a nucleic acid sequence that hybridizes specifically to a target (e.g., target nucleic acid, target molecule, e.g., a cellular nucleic acid to be analyzed), for example to a specific gene sequence. In some embodiments, a target binding region can comprise a nucleic acid sequence that can attach (e.g., hybridize) to a specific location of a specific target nucleic acid. In some embodiments, the target binding region can comprise a nucleic acid sequence that is capable of specific hybridization to a restriction enzyme site overhang (e.g., an EcoRI sticky-end overhang). The barcode can

then ligate to any nucleic acid molecule comprising a sequence complementary to the restriction site overhang.

In some embodiments, a target binding region can comprise a non-specific target nucleic acid sequence. A non-specific target nucleic acid sequence can refer to a sequence that can bind to multiple target nucleic acids, independent of the specific sequence of the target nucleic acid. For example, target binding region can comprise a random multimer sequence, a poly(dA) sequence, a poly(dT) sequence, a poly(dG) sequence, a poly(dC) sequence, or a combination thereof. For example, the target binding region can be an oligo(dT) sequence that hybridizes to the poly(A) tail on mRNA molecules. A random multimer sequence can be, for example, a random dimer, trimer, quatramer, pentamer, hexamer, septamer, octamer, nonamer, decamer, or higher multimer sequence of any length. In some embodiments, the target binding region is the same for all barcodes attached to a given bead. In some embodiments, the target binding regions for the plurality of barcodes attached to a given bead can comprise two or more different target binding sequences. A target binding region can be, or be about, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or a number or a range between any two of these values, nucleotides in length. A target binding region can be at most about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50 or more nucleotides in length. For example, an mRNA molecule can be reverse transcribed using a reverse transcriptase, such as Moloney murine leukemia virus (MMLV) reverse transcriptase, to generate a cDNA molecule with a poly(dC) tail. A barcode can include a target binding region with a poly(dG) tail. Upon base pairing between the poly(dG) tail of the barcode and the poly(dC) tail of the cDNA molecule, the reverse transcriptase switches template strands, from cellular RNA molecule to the barcode, and continues replication to the 5' end of the barcode. By doing so, the resulting cDNA molecule contains the sequence of the barcode (such as the molecular label) on the 3' end of the cDNA molecule.

In some embodiments, a target-binding region can comprise an oligo(dT) which can hybridize with mRNAs comprising polyadenylated ends. A target-binding region can be gene-specific. For example, a target-binding region can be configured to hybridize to a specific region of a target. A target-binding region can be, or be about, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26 27, 28, 29, 30, or a number or a range between any two of these values, nucleotides in length. A target-binding region can be at least, or be at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26 27, 28, 29, or 30, nucleotides in length. A target-binding region can be about 5-30 nucleotides in length. When a barcode comprises a gene-specific target-binding region, the barcode can be referred to herein as a gene-specific barcode.

#### Orientation Property

A stochastic barcode (e.g., a stochastic barcode) can comprise one or more orientation properties which can be used to orient (e.g., align) the barcodes. A barcode can comprise a moiety for isoelectric focusing. Different barcodes can comprise different isoelectric focusing points. When these barcodes are introduced to a sample, the sample can undergo isoelectric focusing in order to orient the barcodes into a known way. In this way, the orientation property can be used to develop a known map of barcodes in a sample. Exemplary orientation properties can include, electrophoretic mobility (e.g., based on size of the barcode), isoelectric point, spin, conductivity, and/or self-assembly. For example, barcodes with an orientation property of

self-assembly, can self-assemble into a specific orientation (e.g., nucleic acid nanostructure) upon activation.

#### Affinity Property

A barcode (e.g., a stochastic barcode) can comprise one or 5 more affinity properties. For example, a spatial label can comprise an affinity property. An affinity property can include a chemical and/or biological moiety that can facilitate binding of the barcode to another entity (e.g., cell receptor). For example, an affinity property can comprise an 10 antibody, for example, an antibody specific for a specific moiety (e.g., receptor) on a sample. In some embodiments, the antibody can guide the barcode to a specific cell type or molecule. Targets at and/or near the specific cell type or molecule can be labeled (e.g., stochastically labeled). The 15 affinity property can, in some embodiments, provide spatial information in addition to the nucleotide sequence of the spatial label because the antibody can guide the barcode to a specific location. The antibody can be a therapeutic antibody, for example a monoclonal antibody or a polyclonal 20 antibody. The antibody can be humanized or chimeric. The antibody can be a naked antibody or a fusion antibody.

The antibody can be a full-length (i.e., naturally occurring or formed by normal immunoglobulin gene fragment recombinatorial processes) immunoglobulin molecule (e.g., an 25 IgG antibody) or an immunologically active (i.e., specifically binding) portion of an immunoglobulin molecule, like an antibody fragment.

The antibody fragment can be, for example, a portion of 30 an antibody such as F(ab')2, Fab', Fab, Fv, sFv and the like. In some embodiments, the antibody fragment can bind with the same antigen that is recognized by the full-length antibody. The antibody fragment can include isolated fragments consisting of the variable regions of antibodies, such as the "Fv" fragments consisting of the variable regions of 35 the heavy and light chains and recombinant single chain polypeptide molecules in which light and heavy variable regions are connected by a peptide linker ("scFv proteins"). Exemplary antibodies can include, but are not limited to, antibodies for cancer cells, antibodies for viruses, antibodies 40 that bind to cell surface receptors (CD8, CD34, CD45), and therapeutic antibodies.

#### Universal Adaptor Primer

A barcode can comprise one or more universal adaptor 45 primers. For example, a gene-specific barcode, such as a gene-specific stochastic barcode, can comprise a universal adaptor primer. A universal adaptor primer can refer to a nucleotide sequence that is universal across all barcodes. A universal adaptor primer can be used for building gene-specific barcodes. A universal adaptor primer can be, or be 50 about, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26 27, 28, 29, 30, or a number or a range between any two of these nucleotides in length. A universal adaptor primer can be at least, or be at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26 27, 28, 29, or 30 nucleotides in length. A universal adaptor primer can be from 5-30 nucleotides in length.

#### Linker

When a barcode comprises more than one of a type of 60 label (e.g., more than one cell label or more than one barcode sequence, such as one molecular label), the labels may be interspersed with a linker label sequence. A linker label sequence can be at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50 or more nucleotides in length. A linker label sequence 65 can be at most about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50 or more nucleotides in length. In some instances, a linker label sequence is 12 nucleotides in length. A linker label sequence

can be used to facilitate the synthesis of the barcode. The linker label can comprise an error-correcting (e.g., Hamming) code.

#### Solid Supports

Barcodes, such as stochastic barcodes, disclosed herein can, in some embodiments, be associated with a solid support. The solid support can be, for example, a synthetic particle. In some embodiments, some or all of the barcode sequences, such as molecular labels for stochastic barcodes (e.g., the first barcode sequences) of a plurality of barcodes (e.g., the first plurality of barcodes) on a solid support differ by at least one nucleotide. The cell labels of the barcodes on the same solid support can be the same. The cell labels of the barcodes on different solid supports can differ by at least one nucleotide. For example, first cell labels of a first plurality of barcodes on a first solid support can have the same sequence, and second cell labels of a second plurality of barcodes on a second solid support can have the same sequence. The first cell labels of the first plurality of barcodes on the first solid support and the second cell labels of the second plurality of barcodes on the second solid support can differ by at least one nucleotide. A cell label can be, for example, about 5-20 nucleotides long. A barcode sequence can be, for example, about 5-20 nucleotides long. The synthetic particle can be, for example, a bead.

The bead can be, for example, a silica gel bead, a controlled pore glass bead, a magnetic bead, a Dynabead, a Sephadex/Sepharose bead, a cellulose bead, a polystyrene bead, or any combination thereof. The bead can comprise a material such as polydimethylsiloxane (PDMS), polystyrene, glass, polypropylene, agarose, gelatin, hydrogel, paramagnetic, ceramic, plastic, glass, methylstyrene, acrylic polymer, titanium, latex, Sepharose, cellulose, nylon, silicone, or any combination thereof.

In some embodiments, the bead can be a polymeric bead, for example a deformable bead or a gel bead, functionalized with barcodes or stochastic barcodes (such as gel beads from 10x Genomics (San Francisco, CA). In some implementation, a gel bead can comprise a polymer based gels. Gel beads can be generated, for example, by encapsulating one or more polymeric precursors into droplets. Upon exposure of the polymeric precursors to an accelerator (e.g., tetramethylethylenediamine (TEMED)), a gel bead may be generated.

In some embodiments, the particle can be disruptable (e.g., dissolvable, degradable). For example, the polymeric bead can dissolve, melt, or degrade, for example, under a desired condition. The desired condition can include an environmental condition. The desired condition may result in the polymeric bead dissolving, melting, or degrading in a controlled manner. A gel bead may dissolve, melt, or degrade due to a chemical stimulus, a physical stimulus, a biological stimulus, a thermal stimulus, a magnetic stimulus, an electric stimulus, a light stimulus, or any combination thereof.

Analytes and/or reagents, such as oligonucleotide barcodes, for example, may be coupled/immobilized to the interior surface of a gel bead (e.g., the interior accessible via diffusion of an oligonucleotide barcode and/or materials used to generate an oligonucleotide barcode) and/or the outer surface of a gel bead or any other microcapsule described herein. Coupling/immobilization may be via any form of chemical bonding (e.g., covalent bond, ionic bond) or physical phenomena (e.g., Van der Waals forces, dipole-dipole interactions, etc.). In some embodiments, coupling/immobilization of a reagent to a gel bead or any other microcapsule described herein may be reversible, such as,

for example, via a labile moiety (e.g., via a chemical cross-linker, including chemical cross-linkers described herein). Upon application of a stimulus, the labile moiety may be cleaved and the immobilized reagent set free. In some embodiments, the labile moiety is a disulfide bond. For example, in the case where an oligonucleotide barcode is immobilized to a gel bead via a disulfide bond, exposure of the disulfide bond to a reducing agent can cleave the disulfide bond and free the oligonucleotide barcode from the bead. The labile moiety may be included as part of a gel bead or microcapsule, as part of a chemical linker that links a reagent or analyte to a gel bead or microcapsule, and/or as part of a reagent or analyte. In some embodiments, at least one barcode of the plurality of barcodes can be immobilized on the particle, partially immobilized on the particle, enclosed in the particle, partially enclosed in the particle, or any combination thereof.

In some embodiments, a gel bead can comprise a wide range of different polymers including but not limited to: polymers, heat sensitive polymers, photosensitive polymers, magnetic polymers, pH sensitive polymers, salt-sensitive polymers, chemically sensitive polymers, polyelectrolytes, polysaccharides, peptides, proteins, and/or plastics. Polymers may include but are not limited to materials such as poly(N-isopropylacrylamide) (PNIPAAm), poly(styrene sulfonate) (PSS), poly(allyl amine) (PAAM), poly(acrylic acid) (PAA), poly(ethylene imine) (PEI), poly(diallyldimethyl-ammonium chloride) (PDADMAC), poly(pyrorrrole) (PPy), poly(vinylpyrrolidone) (PVPON), poly(vinyl pyridine) (PVP), poly(methacrylic acid) (PMAA), poly(methyl methacrylate) (PMMA), polystyrene (PS), poly(tetrahydrofuran) (PTHF), poly(phthalaldehyde) (PTHF), poly(hexyl viologen) (PHV), poly(L-lysine) (PLL), poly(L-arginine) (PARG), poly(lactic-co-glycolic acid) (PLGA).

Numerous chemical stimuli can be used to trigger the disruption, dissolution, or degradation of the beads. Examples of these chemical changes may include, but are not limited to pH-mediated changes to the bead wall, disintegration of the bead wall via chemical cleavage of cross-link bonds, triggered depolymerization of the bead wall, and bead wall switching reactions. Bulk changes may also be used to trigger disruption of the beads.

Bulk or physical changes to the microcapsule through various stimuli also offer many advantages in designing capsules to release reagents. Bulk or physical changes occur on a macroscopic scale, in which bead rupture is the result of mechano-physical forces induced by a stimulus. These processes may include, but are not limited to pressure induced rupture, bead wall melting, or changes in the porosity of the bead wall.

Biological stimuli may also be used to trigger disruption, dissolution, or degradation of beads. Generally, biological triggers resemble chemical triggers, but many examples use biomolecules, or molecules commonly found in living systems such as enzymes, peptides, saccharides, fatty acids, nucleic acids and the like. For example, beads may comprise polymers with peptide cross-links that are sensitive to cleavage by specific proteases. More specifically, one example may comprise a microcapsule comprising GFLGK peptide cross links. Upon addition of a biological trigger such as the protease Cathepsin B, the peptide cross links of the shell well are cleaved and the contents of the beads are released. In other cases, the proteases may be heat-activated. In another example, beads comprise a shell wall comprising cellulose. Addition of the hydrolytic enzyme chitosan serves

as biologic trigger for cleavage of cellulosic bonds, depolymerization of the shell wall, and release of its inner contents.

The beads may also be induced to release their contents upon the application of a thermal stimulus. A change in temperature can cause a variety changes to the beads. A change in heat may cause melting of a bead such that the bead wall disintegrates. In other cases, the heat may increase the internal pressure of the inner components of the bead such that the bead ruptures or explodes. In still other cases, the heat may transform the bead into a shrunken dehydrated state. The heat may also act upon heat-sensitive polymers within the wall of a bead to cause disruption of the bead.

Inclusion of magnetic nanoparticles to the bead wall of microcapsules may allow triggered rupture of the beads as well as guide the beads in an array. A device of this disclosure may comprise magnetic beads for either purpose. In one example, incorporation of  $\text{Fe}_3\text{O}_4$  nanoparticles into polyelectrolyte containing beads triggers rupture in the presence of an oscillating magnetic field stimulus.

A bead may also be disrupted, dissolved, or degraded as the result of electrical stimulation. Similar to magnetic particles described in the previous section, electrically sensitive beads can allow for both triggered rupture of the beads as well as other functions such as alignment in an electric field, electrical conductivity or redox reactions. In one example, beads containing electrically sensitive material are aligned in an electric field such that release of inner reagents can be controlled. In other examples, electrical fields may induce redox reactions within the bead wall itself that may increase porosity.

A light stimulus may also be used to disrupt the beads. Numerous light triggers are possible and may include systems that use various molecules such as nanoparticles and chromophores capable of absorbing photons of specific ranges of wavelengths. For example, metal oxide coatings can be used as capsule triggers. UV irradiation of polyelectrolyte capsules coated with  $\text{SiO}_2$  may result in disintegration of the bead wall. In yet another example, photo switchable materials such as azobenzene groups may be incorporated in the bead wall. Upon the application of UV or visible light, chemicals such as these undergo a reversible cis-to-trans isomerization upon absorption of photons. In this aspect, incorporation of photon switches result in a bead wall that may disintegrate or become more porous upon the application of a light trigger.

For example, in a non-limiting example of barcoding (e.g., stochastic barcoding) illustrated in FIG. 2, after introducing cells such as single cells onto a plurality of microwells of a microwell array at block 208, beads can be introduced onto the plurality of microwells of the microwell array at block 212. Each microwell can comprise one bead. The beads can comprise a plurality of barcodes. A barcode can comprise a 5' amine region attached to a bead. The barcode can comprise a universal label, a barcode sequence (e.g., a molecular label), a target-binding region, or any combination thereof.

The barcodes disclosed herein can be associated with (e.g., attached to) a solid support (e.g., a bead). The barcodes associated with a solid support can each comprise a barcode sequence selected from a group comprising at least 100 or 1000 barcode sequences with unique sequences. In some embodiments, different barcodes associated with a solid support can comprise barcode with different sequences. In some embodiments, a percentage of barcodes associated with a solid support comprises the same cell label. For example, the percentage can be, or be about 60%, 70%,

80%, 85%, 90%, 95%, 97%, 99%, 100%, or a number or a range between any two of these values. As another example, the percentage can be at least, or be at most 60%, 70%, 80%, 85%, 90%, 95%, 97%, 99%, or 100%. In some embodiments, barcodes associated with a solid support can have the same cell label. The barcodes associated with different solid supports can have different cell labels selected from a group comprising at least 100 or 1000 cell labels with unique sequences.

10 The barcodes disclosed herein can be associated to (e.g., attached to) a solid support (e.g., a bead). In some embodiments, barcoding the plurality of targets in the sample can be performed with a solid support including a plurality of synthetic particles associated with the plurality of barcodes. 15 In some embodiments, the solid support can include a plurality of synthetic particles associated with the plurality of barcodes. The spatial labels of the plurality of barcodes on different solid supports can differ by at least one nucleotide. The solid support can, for example, include the plurality of barcodes in two dimensions or three dimensions. The synthetic particles can be beads. The beads can be silica gel beads, controlled pore glass beads, magnetic beads, Dynabeads, Sephadex/Sepharose beads, cellulose beads, polystyrene beads, or any combination thereof. The solid support 20 can include a polymer, a matrix, a hydrogel, a needle array device, an antibody, or any combination thereof. In some embodiments, the solid supports can be free floating. In some embodiments, the solid supports can be embedded in a semi-solid or solid array. The barcodes may not be associated with solid supports. The barcodes can be individual nucleotides. The barcodes can be associated with a substrate.

As used herein, the terms "tethered," "attached," and "immobilized," are used interchangeably, and can refer to covalent or non-covalent means for attaching barcodes to a solid support. Any of a variety of different solid supports can be used as solid supports for attaching pre-synthesized barcodes or for in situ solid-phase synthesis of barcode.

In some embodiments, the solid support is a bead. The 40 bead can comprise one or more types of solid, porous, or hollow sphere, ball, bearing, cylinder, or other similar configuration which a nucleic acid can be immobilized (e.g., covalently or non-covalently). The bead can be, for example, composed of plastic, ceramic, metal, polymeric material, or any combination thereof. A bead can be, or comprise, a discrete particle that is spherical (e.g., microspheres) or have a non-spherical or irregular shape, such as cubic, cuboid, pyramidal, cylindrical, conical, oblong, or disc-shaped, and the like. In some embodiments, a bead can be non-spherical 45 in shape.

Beads can comprise a variety of materials including, but not limited to, paramagnetic materials (e.g., magnesium, molybdenum, lithium, and tantalum), superparamagnetic materials (e.g., ferrite ( $\text{Fe}_3\text{O}_4$ ; magnetite) nanoparticles), 55 ferromagnetic materials (e.g., iron, nickel, cobalt, some alloys thereof, and some rare earth metal compounds), ceramic, plastic, glass, polystyrene, silica, methylstyrene, acrylic polymers, titanium, latex, Sepharose, agarose, hydrogel, polymer, cellulose, nylon, or any combination thereof.

60 In some embodiments, the bead (e.g., the bead to which the labels are attached) is a hydrogel bead. In some embodiments, the bead comprises hydrogel.

Some embodiments disclosed herein include one or more particles (for example, beads). Each of the particles can 65 comprise a plurality of oligonucleotides (e.g., barcodes). Each of the plurality of oligonucleotides can comprise a barcode sequence (e.g., a molecular label sequence), a cell

label, and a target-binding region (e.g., an oligo(dT) sequence, a gene-specific sequence, a random multimer, or a combination thereof). The cell label sequence of each of the plurality of oligonucleotides can be the same. The cell label sequences of oligonucleotides on different particles can be different such that the oligonucleotides on different particles can be identified. The number of different cell label sequences can be different in different implementations. In some embodiments, the number of cell label sequences can be, or be about 10, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, 10000, 20000, 30000, 40000, 50000, 60000, 70000, 80000, 90000, 100000,  $10^6$ ,  $10^7$ ,  $10^8$ ,  $10^9$ , a number or a range between any two of these values, or more. In some embodiments, the number of cell label sequences can be at least, or be at most 10, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, 10000, 20000, 30000, 40000, 50000, 60000, 70000, 80000, 90000, 100000,  $10^6$ ,  $10^7$ ,  $10^8$ , or  $10^9$ . In some embodiments, no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, or more of the plurality of the particles include oligonucleotides with the same cell sequence. In some embodiment, the plurality of particles that include oligonucleotides with the same cell sequence can be at most 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, or more. In some embodiments, none of the plurality of the particles has the same cell label sequence.

The plurality of oligonucleotides on each particle can comprise different barcode sequences (e.g., molecular labels). In some embodiments, the number of barcode sequences can be, or be about 10, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, 10000, 20000, 30000, 40000, 50000, 60000, 70000, 80000, 90000, 100000,  $10^6$ ,  $10^7$ ,  $10^8$ ,  $10^9$ , or a number or a range between any two of these values. In some embodiments, the number of barcode sequences can be at least, or be at most 10, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000, 3000, 4000, 5000, 6000, 7000, 8000, 9000, 10000, 20000, 30000, 40000, 50000, 60000, 70000, 80000, 90000, 100000,  $10^6$ ,  $10^7$ ,  $10^8$ , or  $10^9$ . For example, at least 100 of the plurality of oligonucleotides comprise different barcode sequences. As another example, in a single particle, at least 100, 500, 1000, 5000, 10000, 15000, 20000, 50000, a number or a range between any two of these values, or more of the plurality of oligonucleotides comprise different barcode sequences. Some embodiments provide a plurality of the particles comprising barcodes. In some embodiments, the ratio of an occurrence (or a copy or a number) of a target to be labeled and the different barcode sequences can be at least 1:1, 1:2, 1:3, 1:4, 1:5, 1:6, 1:7, 1:8, 1:9, 1:10, 1:11, 1:12, 1:13, 1:14, 1:15, 1:16, 1:17, 1:18, 1:19, 1:20, 1:30, 1:40, 1:50, 1:60, 1:70, 1:80, 1:90, or more. In some embodiments, each of the plurality of oligonucleotides further comprises a sample label, a universal label, or both. The particle can be, for example, a nanoparticle or microparticle.

The size of the beads can vary. For example, the diameter of the bead can range from 0.1 micrometer to 50 micrometer. In some embodiments, the diameter of the bead can be, or be about, 0.1, 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50 micrometer, or a number or a range between any two of these values.

The diameter of the bead can be related to the diameter of the wells of the substrate. In some embodiments, the diameter of the bead can be, or be about, 10%, 20%, 30%, 40%,

50%, 60%, 70%, 80%, 90%, 100%, or a number or a range between any two of these values, longer or shorter than the diameter of the well. The diameter of the beads can be related to the diameter of a cell (e.g., a single cell entrapped by a well of the substrate). In some embodiments, the diameter of the bead can be at least, or be at most, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% longer or shorter than the diameter of the well. The diameter of the beads can be related to the diameter of a cell (e.g., a single cell entrapped by a well of the substrate). In some embodiments, the diameter of the bead can be, or be about, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 150%, 200%, 250%, 300%, or a number or a range between any two of these values, longer or shorter than the diameter of the cell. In some embodiments, the diameter of the beads can be at least, or be at most, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 150%, 200%, 250%, or 300% longer or shorter than the diameter of the cell.

A bead can be attached to and/or embedded in a substrate. 20 A bead can be attached to and/or embedded in a gel, hydrogel, polymer and/or matrix. The spatial position of a bead within a substrate (e.g., gel, matrix, scaffold, or polymer) can be identified using the spatial label present on the barcode on the bead which can serve as a location address.

Examples of beads can include, but are not limited to, streptavidin beads, agarose beads, magnetic beads, Dynabeads®, MACS® microbeads, antibody conjugated beads (e.g., anti-immunoglobulin microbeads), protein A conjugated beads, protein G conjugated beads, protein A/G conjugated beads, protein L conjugated beads, oligo(dT) conjugated beads, silica beads, silica-like beads, anti-biotin microbeads, anti-fluorochrome microbeads, and BcMag™ Carboxyl-Terminated Magnetic Beads.

A bead can be associated with (e.g., impregnated with) 35 quantum dots or fluorescent dyes to make it fluorescent in one fluorescence optical channel or multiple optical channels. A bead can be associated with iron oxide or chromium oxide to make it paramagnetic or ferromagnetic. Beads can be identifiable. For example, a bead can be imaged using a camera. A bead can have a detectable code associated with the bead. For example, a bead can comprise a barcode. A bead can change size, for example, due to swelling in an organic or inorganic solution. A bead can be hydrophobic. A bead can be hydrophilic. A bead can be biocompatible.

A solid support (e.g., a bead) can be visualized. The solid support can comprise a visualizing tag (e.g., fluorescent dye). A solid support (e.g., a bead) can be etched with an identifier (e.g., a number). The identifier can be visualized through imaging the beads.

A solid support can comprise an insoluble, semi-soluble, or insoluble material. A solid support can be referred to as “functionalized” when it includes a linker, a scaffold, a building block, or other reactive moiety attached thereto, whereas a solid support may be “nonfunctionalized” when it lacks such a reactive moiety attached thereto. The solid support can be employed free in solution, such as in a microtiter well format; in a flow-through format, such as in a column; or in a dipstick.

The solid support can comprise a membrane, paper, plastic, coated surface, flat surface, glass, slide, chip, or any combination thereof. A solid support can take the form of resins, gels, microspheres, or other geometric configurations. A solid support can comprise silica chips, microparticles, nanoparticles, plates, arrays, capillaries, flat supports such as glass fiber filters, glass surfaces, metal surfaces (steel, gold silver, aluminum, silicon and copper), glass supports, plastic supports, silicon supports, chips, filters,

31

membranes, microwell plates, slides, plastic materials including multiwell plates or membranes (e.g., formed of polyethylene, polypropylene, polyamide, polyvinylidene-fluoride), and/or wafers, combs, pins or needles (e.g., arrays of pins suitable for combinatorial synthesis or analysis) or beads in an array of pits or nanoliter wells of flat surfaces such as wafers (e.g., silicon wafers), wafers with pits with or without filter bottoms.

The solid support can comprise a polymer matrix (e.g., gel, hydrogel). The polymer matrix may be able to permeate intracellular space (e.g., around organelles). The polymer matrix may be able to be pumped throughout the circulatory system.

#### Substrates and Microwell Array

As used herein, a substrate can refer to a type of solid support. A substrate can refer to a solid support that can comprise barcodes or stochastic barcodes of the disclosure. A substrate can, for example, comprise a plurality of microwells. For example, a substrate can be a well array comprising two or more microwells. In some embodiments, a microwell can comprise a small reaction chamber of defined volume. In some embodiments, a microwell can entrap one or more cells. In some embodiments, a microwell can entrap only one cell. In some embodiments, a microwell can entrap one or more solid supports. In some embodiments, a microwell can entrap only one solid support. In some embodiments, a microwell entraps a single cell and a single solid support (e.g., a bead). A microwell can comprise barcode reagents of the disclosure.

#### Methods of Barcoding

The disclosure provides for methods for estimating the number of distinct targets at distinct locations in a physical sample (e.g., tissue, organ, tumor, cell). The methods can comprise placing barcodes (e.g., stochastic barcodes) in close proximity with the sample, lysing the sample, associating distinct targets with the barcodes, amplifying the targets and/or digitally counting the targets. The method can further comprise analyzing and/or visualizing the information obtained from the spatial labels on the barcodes. In some embodiments, a method comprises visualizing the plurality of targets in the sample. Mapping the plurality of targets onto the map of the sample can include generating a two dimensional map or a three dimensional map of the sample. The two dimensional map and the three dimensional map can be generated prior to or after barcoding (e.g., stochastically barcoding) the plurality of targets in the sample. Visualizing the plurality of targets in the sample can include mapping the plurality of targets onto a map of the sample. Mapping the plurality of targets onto the map of the sample can include generating a two dimensional map or a three dimensional map of the sample. The two dimensional map and the three dimensional map can be generated prior to or after barcoding the plurality of targets in the sample. In some embodiments, the two dimensional map and the three dimensional map can be generated before or after lysing the sample. Lysing the sample before or after generating the two dimensional map or the three dimensional map can include heating the sample, contacting the sample with a detergent, changing the pH of the sample, or any combination thereof.

In some embodiments, barcoding the plurality of targets comprises hybridizing a plurality of barcodes with a plurality of targets to create barcoded targets (e.g., stochastically barcoded targets). Barcoding the plurality of targets can comprise generating an indexed library of the barcoded targets. Generating an indexed library of the barcoded targets can be performed with a solid support comprising the plurality of barcodes (e.g., stochastic barcodes).

32

#### Contacting a Sample and a Barcode

The disclosure provides for methods for contacting a sample (e.g., cells) to a substrate of the disclosure. A sample comprising, for example, a cell, organ, or tissue thin section, can be contacted to barcodes (e.g., stochastic barcodes). The cells can be contacted, for example, by gravity flow wherein the cells can settle and create a monolayer. The sample can be a tissue thin section. The thin section can be placed on the substrate. The sample can be one-dimensional (e.g., forms a planar surface). The sample (e.g., cells) can be spread across the substrate, for example, by growing/culturing the cells on the substrate.

When barcodes are in close proximity to targets, the targets can hybridize to the barcode. The barcodes can be contacted at a non-depletable ratio such that each distinct target can associate with a distinct barcode of the disclosure. To ensure efficient association between the target and the barcode, the targets can be cross-linked to barcode.

#### Cell Lysis

Following the distribution of cells and barcodes, the cells can be lysed to liberate the target molecules. Cell lysis can be accomplished by any of a variety of means, for example, by chemical or biochemical means, by osmotic shock, or by means of thermal lysis, mechanical lysis, or optical lysis. Cells can be lysed by addition of a cell lysis buffer comprising a detergent (e.g., SDS, Li dodecyl sulfate, Triton X-100, Tween-20, or NP-40), an organic solvent (e.g., methanol or acetone), or digestive enzymes (e.g., proteinase K, pepsin, or trypsin), or any combination thereof. To increase the association of a target and a barcode, the rate of the diffusion of the target molecules can be altered by for example, reducing the temperature and/or increasing the viscosity of the lysate.

In some embodiments, the sample can be lysed using a filter paper. The filter paper can be soaked with a lysis buffer on top of the filter paper. The filter paper can be applied to the sample with pressure which can facilitate lysis of the sample and hybridization of the targets of the sample to the substrate.

In some embodiments, lysis can be performed by mechanical lysis, heat lysis, optical lysis, and/or chemical lysis. Chemical lysis can include the use of digestive enzymes such as proteinase K, pepsin, and trypsin. Lysis can be performed by the addition of a lysis buffer to the substrate. A lysis buffer can comprise Tris HCl. A lysis buffer can comprise at least about 0.01, 0.05, 0.1, 0.5, or 1 M or more Tris HCl. A lysis buffer can comprise at most about 0.01, 0.05, 0.1, 0.5, or 1 M or more Tris HCl. A lysis buffer can comprise about 0.1 M Tris HCl. The pH of the lysis buffer can be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more. The pH of the lysis buffer can be at most about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more. In some embodiments, the pH of the lysis buffer is about 7.5. The lysis buffer can comprise a salt (e.g., LiCl). The concentration of salt in the lysis buffer can be at least about 0.1, 0.5, or 1 M or more. The concentration of salt in the lysis buffer can be at most about 0.1, 0.5, or 1 M or more. In some embodiments, the concentration of salt in the lysis buffer is about 0.5M. The lysis buffer can comprise a detergent (e.g., SDS, Li dodecyl sulfate, triton X, tween, and NP-40). The concentration of the detergent in the lysis buffer can be at least about 0.0001%, 0.0005%, 0.001%, 0.005%, 0.01%, 0.05%, 0.1%, 0.5%, 1%, 2%, 3%, 4%, 5%, 6%, or 7%, or more. The concentration of the detergent in the lysis buffer can be at most about 0.0001%, 0.0005%, 0.001%, 0.005%, 0.01%, 0.05%, 0.1%, 0.5%, 1%, 2%, 3%, 4%, 5%, 6%, or 7%, or more. In some embodiments, the concentration of the deter-

gent in the lysis buffer is about 1% Li dodecyl sulfate. The time used in the method for lysis can be dependent on the amount of detergent used. In some embodiments, the more detergent used, the less time needed for lysis. The lysis buffer can comprise a chelating agent (e.g., EDTA and EGTA). The concentration of a chelating agent in the lysis buffer can be at least about 1, 5, 10, 15, 20, 25, or 30 mM or more. The concentration of a chelating agent in the lysis buffer can be at most about 1, 5, 10, 15, 20, 25, or 30 mM or more. In some embodiments, the concentration of chelating agent in the lysis buffer is about 10 mM. The lysis buffer can comprise a reducing reagent (e.g., beta-mercaptoethanol, and DTT). The concentration of the reducing reagent in the lysis buffer can be at least about 1, 5, 10, 15, or 20 mM or more. The concentration of the reducing reagent in the lysis buffer can be at most about 1, 5, 10, 15, or 20 mM or more. In some embodiments, the concentration of reducing reagent in the lysis buffer is about 5 mM. In some embodiments, a lysis buffer can comprise about 0.1M TrisHCl, about pH 7.5, about 0.5M LiCl, about 1% lithium dodecyl sulfate, about 10 mM EDTA, and about 5 mM DTT.

Lysis can be performed at a temperature of about 4, 10, 15, 20, 25, or 30° C. Lysis can be performed for about 1, 5, 10, 15, or 20 or more minutes. A lysed cell can comprise at least about 100000, 200000, 300000, 400000, 500000, 600000, or 700000 or more target nucleic acid molecules. A lysed cell can comprise at most about 100000, 200000, 300000, 400000, 500000, 600000, or 700000 or more target nucleic acid molecules.

#### Attachment of Barcodes to Target Nucleic Acid Molecules

Following lysis of the cells and release of nucleic acid molecules therefrom, the nucleic acid molecules can randomly associate with the barcodes of the co-localized solid support. Association can comprise hybridization of a barcode's target recognition region to a complementary portion of the target nucleic acid molecule (e.g., oligo(dT) of the barcode can interact with a poly(A) tail of a target). The assay conditions used for hybridization (e.g., buffer pH, ionic strength, temperature, etc.) can be chosen to promote formation of specific, stable hybrids. In some embodiments, the nucleic acid molecules released from the lysed cells can associate with the plurality of probes on the substrate (e.g., hybridize with the probes on the substrate). When the probes comprise oligo(dT), mRNA molecules can hybridize to the probes and be reverse transcribed. The oligo(dT) portion of the oligonucleotide can act as a primer for first strand synthesis of the cDNA molecule. For example, in a non-limiting example of barcoding illustrated in FIG. 2, at block 216, mRNA molecules can hybridize to barcodes on beads. For example, single-stranded nucleotide fragments can 50 hybridize to the target-binding regions of barcodes.

Attachment can further comprise ligation of a barcode's target recognition region and a portion of the target nucleic acid molecule. For example, the target binding region can comprise a nucleic acid sequence that can be capable of specific hybridization to a restriction site overhang (e.g., an EcoRI sticky-end overhang). The assay procedure can further comprise treating the target nucleic acids with a restriction enzyme (e.g., EcoRI) to create a restriction site overhang. The barcode can then be ligated to any nucleic acid molecule comprising a sequence complementary to the restriction site overhang. A ligase (e.g., T4 DNA ligase) can be used to join the two fragments.

For example, in a non-limiting example of barcoding illustrated in FIG. 2, at block 220, the labeled targets from a plurality of cells (or a plurality of samples) (e.g., target-barcode molecules) can be subsequently pooled, for

example, into a tube. The labeled targets can be pooled by, for example, retrieving the barcodes and/or the beads to which the target-barcode molecules are attached.

The retrieval of solid support-based collections of attached target-barcode molecules can be implemented by use of magnetic beads and an externally-applied magnetic field. Once the target-barcode molecules have been pooled, all further processing can proceed in a single reaction vessel. Further processing can include, for example, reverse transcription reactions, amplification reactions, cleavage reactions, dissociation reactions, and/or nucleic acid extension reactions. Further processing reactions can be performed within the microwells, that is, without first pooling the labeled target nucleic acid molecules from a plurality of cells.

#### Reverse Transcription or Nucleic Acid Extension

The disclosure provides for a method to create a target-barcode conjugate using reverse transcription (e.g., at block 224 of FIG. 2) or nucleic acid extension. The target-barcode conjugate can comprise the barcode and a complementary sequence of all or a portion of the target nucleic acid (i.e., a barcoded cDNA molecule, such as a stochastically barcoded cDNA molecule). Reverse transcription of the associated RNA molecule can occur by the addition of a reverse transcription primer along with the reverse transcriptase. The reverse transcription primer can be an oligo(dT) primer, a random hexanucleotide primer, or a target-specific oligonucleotide primer. Oligo(dT) primers can be, or can be about, 12-18 nucleotides in length and bind to the endogenous poly(A) tail at the 3' end of mammalian mRNA. Random hexanucleotide primers can bind to mRNA at a variety of complementary sites. Target-specific oligonucleotide primers typically selectively prime the mRNA of interest.

In some embodiments, reverse transcription of an mRNA molecule to a labeled-RNA molecule can occur by the addition of a reverse transcription primer. In some embodiments, the reverse transcription primer is an oligo(dT) primer, random hexanucleotide primer, or a target-specific oligonucleotide primer. Generally, oligo(dT) primers are 12-18 nucleotides in length and bind to the endogenous poly(A) tail at the 3' end of mammalian mRNA. Random hexanucleotide primers can bind to mRNA at a variety of complementary sites. Target-specific oligonucleotide primers typically selectively prime the mRNA of interest.

In some embodiments, a target is a cDNA molecule. For example, an mRNA molecule can be reverse transcribed using a reverse transcriptase, such as Moloney murine leukemia virus (MMLV) reverse transcriptase, to generate a cDNA molecule with a poly(dC) tail. A barcode can include a target binding region with a poly(dG) tail. Upon base pairing between the poly(dG) tail of the barcode and the poly(dC) tail of the cDNA molecule, the reverse transcriptase switches template strands, from cellular RNA molecule to the barcode, and continues replication to the 5' end of the barcode. By doing so, the resulting cDNA molecule contains the sequence of the barcode (such as the molecular label) on the 3' end of the cDNA molecule.

Reverse transcription can occur repeatedly to produce multiple labeled-cDNA molecules. The methods disclosed herein can comprise conducting at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 reverse transcription reactions. The method can comprise conducting at least about 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 reverse transcription reactions.

**Amplification**

One or more nucleic acid amplification reactions (e.g., at block 228 of FIG. 2) can be performed to create multiple copies of the labeled target nucleic acid molecules. Amplification can be performed in a multiplexed manner, wherein multiple target nucleic acid sequences are amplified simultaneously. The amplification reaction can be used to add sequencing adaptors to the nucleic acid molecules. The amplification reactions can comprise amplifying at least a portion of a sample label, if present. The amplification reactions can comprise amplifying at least a portion of the cellular label and/or barcode sequence (e.g., a molecular label). The amplification reactions can comprise amplifying at least a portion of a sample tag, a cell label, a spatial label, a barcode sequence (e.g., a molecular label), a target nucleic acid, or a combination thereof. The amplification reactions can comprise amplifying 0.5%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 100%, or a range or a number between any two of these values, of the plurality of nucleic acids. The method can further comprise conducting one or more cDNA synthesis reactions to produce one or more cDNA copies of target-barcode molecules comprising a sample label, a cell label, a spatial label, and/or a barcode sequence (e.g., a molecular label).

In some embodiments, amplification can be performed using a polymerase chain reaction (PCR). As used herein, PCR can refer to a reaction for the in vitro amplification of specific DNA sequences by the simultaneous primer extension of complementary strands of DNA. As used herein, PCR can encompass derivative forms of the reaction, including but not limited to, RT-PCR, real-time PCR, nested PCR, quantitative PCR, multiplexed PCR, digital PCR, and assembly PCR.

Amplification of the labeled nucleic acids can comprise non-PCR based methods. Examples of non-PCR based methods include, but are not limited to, multiple displacement amplification (MDA), transcription-mediated amplification (TMA), nucleic acid sequence-based amplification (NASBA), strand displacement amplification (SDA), real-time SDA, rolling circle amplification, or circle-to-circle amplification. Other non-PCR-based amplification methods include multiple cycles of DNA-dependent RNA polymerase-driven RNA transcription amplification or RNA-directed DNA synthesis and transcription to amplify DNA or RNA targets, a ligase chain reaction (LCR), and a Q $\beta$  replicase (%) method, use of palindromic probes, strand displacement amplification, oligonucleotide-driven amplification using a restriction endonuclease, an amplification method in which a primer is hybridized to a nucleic acid sequence and the resulting duplex is cleaved prior to the extension reaction and amplification, strand displacement amplification using a nucleic acid polymerase lacking 5' exonuclease activity, rolling circle amplification, and ramification extension amplification (RAM). In some embodiments, the amplification does not produce circularized transcripts.

In some embodiments, the methods disclosed herein further comprise conducting a polymerase chain reaction on the labeled nucleic acid (e.g., labeled-RNA, labeled-DNA, labeled-cDNA) to produce a labeled amplicon (e.g., a stochastically labeled amplicon). The labeled amplicon can be double-stranded molecule. The double-stranded molecule can comprise a double-stranded RNA molecule, a double-stranded DNA molecule, or a RNA molecule hybridized to a DNA molecule. One or both of the strands of the double-

stranded molecule can comprise a sample label, a spatial label, a cell label, and/or a barcode sequence (e.g., a molecular label). The labeled amplicon can be a single-stranded molecule. The single-stranded molecule can comprise DNA, RNA, or a combination thereof. The nucleic acids of the disclosure can comprise synthetic or altered nucleic acids.

Amplification can comprise use of one or more non-natural nucleotides. Non-natural nucleotides can comprise photolabile or triggerable nucleotides. Examples of non-natural nucleotides can include, but are not limited to, peptide nucleic acid (PNA), morpholino and locked nucleic acid (LNA), as well as glycol nucleic acid (GNA) and threose nucleic acid (TNA). Non-natural nucleotides can be added to one or more cycles of an amplification reaction. The addition of the non-natural nucleotides can be used to identify products as specific cycles or time points in the amplification reaction.

Conducting the one or more amplification reactions can comprise the use of one or more primers. The one or more primers can comprise, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 or more nucleotides. The one or more primers can comprise at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 or more nucleotides. The one or more primers can comprise less than 12-15 nucleotides. The one or more primers can anneal to at least a portion of the plurality of labeled targets (e.g., stochastically labeled targets). The one or more primers can anneal to the 3' end or 5' end of the plurality of labeled targets. The one or more primers can anneal to an internal region of the plurality of labeled targets. The internal region can be at least about 50, 100, 150, 200, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 650, 700, 750, 800, 850, 900 or 1000 nucleotides from the 3' ends the plurality of labeled targets. The one or more primers can comprise a fixed panel of primers. The one or more primers can comprise at least one or more custom primers. The one or more primers can comprise at least one or more control primers. The one or more primers can comprise at least one or more gene-specific primers.

The one or more primers can comprise a universal primer. The universal primer can anneal to a universal primer binding site. The one or more custom primers can anneal to a first sample label, a second sample label, a spatial label, a cell label, a barcode sequence (e.g., a molecular label), a target, or any combination thereof. The one or more primers can comprise a universal primer and a custom primer. The custom primer can be designed to amplify one or more targets. The targets can comprise a subset of the total nucleic acids in one or more samples. The targets can comprise a subset of the total labeled targets in one or more samples. The one or more primers can comprise at least 96 or more custom primers. The one or more primers can comprise at least 960 or more custom primers. The one or more primers can comprise at least 9600 or more custom primers. The one or more custom primers can anneal to two or more different labeled nucleic acids. The two or more different labeled nucleic acids can correspond to one or more genes.

Any amplification scheme can be used in the methods of the present disclosure. For example, in one scheme, the first round PCR can amplify molecules attached to the bead using a gene specific primer and a primer against the universal Illumina sequencing primer 1 sequence. The second round of PCR can amplify the first PCR products using a nested gene specific primer flanked by Illumina sequencing primer 2 sequence, and a primer against the universal Illumina

sequencing primer 1 sequence. The third round of PCR adds P5 and P7 and sample index to turn PCR products into an Illumina sequencing library. Sequencing using 150 bpx2 sequencing can reveal the cell label and barcode sequence (e.g., molecular label) on read 1, the gene on read 2, and the sample index on index 1 read.

In some embodiments, nucleic acids can be removed from the substrate using chemical cleavage. For example, a chemical group or a modified base present in a nucleic acid can be used to facilitate its removal from a solid support. For example, an enzyme can be used to remove a nucleic acid from a substrate. For example, a nucleic acid can be removed from a substrate through a restriction endonuclease digestion. For example, treatment of a nucleic acid containing a dUTP or ddUTP with uracil-d-glycosylase (UDG) can be used to remove a nucleic acid from a substrate. For example, a nucleic acid can be removed from a substrate using an enzyme that performs nucleotide excision, such as a base excision repair enzyme, such as an apurinic/aprimidinic (AP) endonuclease. In some embodiments, a nucleic acid can be removed from a substrate using a photocleavable group and light. In some embodiments, a cleavable linker can be used to remove a nucleic acid from the substrate. For example, the cleavable linker can comprise at least one of biotin/avidin, biotin/streptavidin, biotin/neutravidin, Ig-protein A, a photo-labile linker, acid or base labile linker group, or an aptamer.

When the probes are gene-specific, the molecules can hybridize to the probes and be reverse transcribed and/or amplified. In some embodiments, after the nucleic acid has been synthesized (e.g., reverse transcribed), it can be amplified. Amplification can be performed in a multiplex manner, wherein multiple target nucleic acid sequences are amplified simultaneously. Amplification can add sequencing adaptors to the nucleic acid.

In some embodiments, amplification can be performed on the substrate, for example, with bridge amplification. cDNAs can be homopolymer tailed in order to generate a compatible end for bridge amplification using oligo(dT) probes on the substrate. In bridge amplification, the primer that is complementary to the 3' end of the template nucleic acid can be the first primer of each pair that is covalently attached to the solid particle. When a sample containing the template nucleic acid is contacted with the particle and a single thermal cycle is performed, the template molecule can be annealed to the first primer and the first primer is elongated in the forward direction by addition of nucleotides to form a duplex molecule consisting of the template molecule and a newly formed DNA strand that is complementary to the template. In the heating step of the next cycle, the duplex molecule can be denatured, releasing the template molecule from the particle and leaving the complementary DNA strand attached to the particle through the first primer. In the annealing stage of the annealing and elongation step that follows, the complementary strand can hybridize to the second primer, which is complementary to a segment of the complementary strand at a location removed from the first primer. This hybridization can cause the complementary strand to form a bridge between the first and second primers secured to the first primer by a covalent bond and to the second primer by hybridization. In the elongation stage, the second primer can be elongated in the reverse direction by the addition of nucleotides in the same reaction mixture, thereby converting the bridge to a double-stranded bridge. The next cycle then begins, and the double-stranded bridge can be denatured to yield two single-stranded nucleic acid molecules, each having one end attached to the particle

surface via the first and second primers, respectively, with the other end of each unattached. In the annealing and elongation step of this second cycle, each strand can hybridize to a further complementary primer, previously unused, on the same particle, to form new single-strand bridges. The two previously unused primers that are now hybridized elongate to convert the two new bridges to double-strand bridges.

The amplification reactions can comprise amplifying at least 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 100% of the plurality of nucleic acids.

Amplification of the labeled nucleic acids can comprise PCR-based methods or non-PCR based methods. Amplification of the labeled nucleic acids can comprise exponential amplification of the labeled nucleic acids. Amplification of the labeled nucleic acids can comprise linear amplification of the labeled nucleic acids. Amplification can be performed by polymerase chain reaction (PCR). PCR can refer to a reaction for the in vitro amplification of specific DNA sequences by the simultaneous primer extension of complementary strands of DNA. PCR can encompass derivative forms of the reaction, including but not limited to, RT-PCR, real-time PCR, nested PCR, quantitative PCR, multiplexed PCR, digital PCR, suppression PCR, semi-suppressive PCR and assembly PCR.

In some embodiments, amplification of the labeled nucleic acids comprises non-PCR based methods. Examples of non-PCR based methods include, but are not limited to, multiple displacement amplification (MDA), transcription-mediated amplification (TMA), nucleic acid sequence-based amplification (NASBA), strand displacement amplification (SDA), real-time SDA, rolling circle amplification, or circle-to-circle amplification. Other non-PCR-based amplification methods include multiple cycles of DNA-dependent RNA polymerase-driven RNA transcription amplification or RNA-directed DNA synthesis and transcription to amplify DNA or RNA targets, a ligase chain reaction (LCR), a Q $\beta$  replicase (%), use of palindromic probes, strand displacement amplification, oligonucleotide-driven amplification using a restriction endonuclease, an amplification method in which a primer is hybridized to a nucleic acid sequence and the resulting duplex is cleaved prior to the extension reaction and amplification, strand displacement amplification using a nucleic acid polymerase lacking 5' exonuclease activity, rolling circle amplification, and/or ramification extension amplification (RAM).

In some embodiments, the methods disclosed herein further comprise conducting a nested polymerase chain reaction on the amplified amplicon (e.g., target). The amplicon can be double-stranded molecule. The double-stranded molecule can comprise a double-stranded RNA molecule, a double-stranded DNA molecule, or a RNA molecule hybridized to a DNA molecule. One or both of the strands of the double-stranded molecule can comprise a sample tag or molecular identifier label. Alternatively, the amplicon can be a single-stranded molecule. The single-stranded molecule can comprise DNA, RNA, or a combination thereof. The nucleic acids of the present invention can comprise synthetic or altered nucleic acids.

In some embodiments, the method comprises repeatedly amplifying the labeled nucleic acid to produce multiple amplicons. The methods disclosed herein can comprise conducting at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amplification reactions. Alternatively, the method comprises conducting at least

39

about 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amplification reactions.

Amplification can further comprise adding one or more control nucleic acids to one or more samples comprising a plurality of nucleic acids. Amplification can further comprise adding one or more control nucleic acids to a plurality of nucleic acids. The control nucleic acids can comprise a control label.

Amplification can comprise use of one or more non-natural nucleotides. Non-natural nucleotides can comprise photolabile and/or triggerable nucleotides. Examples of non-natural nucleotides include, but are not limited to, peptide nucleic acid (PNA), morpholino and locked nucleic acid (LNA), as well as glycol nucleic acid (GNA) and threose nucleic acid (TNA). Non-natural nucleotides can be added to one or more cycles of an amplification reaction. The addition of the non-natural nucleotides can be used to identify products as specific cycles or time points in the amplification reaction.

Conducting the one or more amplification reactions can comprise the use of one or more primers. The one or more primers can comprise one or more oligonucleotides. The one or more oligonucleotides can comprise at least about 7-9 nucleotides. The one or more oligonucleotides can comprise less than 12-15 nucleotides. The one or more primers can anneal to at least a portion of the plurality of labeled nucleic acids. The one or more primers can anneal to the 3' end and/or 5' end of the plurality of labeled nucleic acids. The one or more primers can anneal to an internal region of the plurality of labeled nucleic acids. The internal region can be at least about 50, 100, 150, 200, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 650, 700, 750, 800, 850, 900 or 1000 nucleotides from the 3' ends the plurality of labeled nucleic acids. The one or more primers can comprise a fixed panel of primers. The one or more primers can comprise at least one or more custom primers. The one or more primers can comprise at least one or more control primers. The one or more primers can comprise at least one or more housekeeping gene primers. The one or more primers can comprise a universal primer. The universal primer can anneal to a universal primer binding site. The one or more custom primers can anneal to the first sample tag, the second sample tag, the molecular identifier label, the nucleic acid or a product thereof. The one or more primers can comprise a universal primer and a custom primer. The custom primer can be designed to amplify one or more target nucleic acids. The target nucleic acids can comprise a subset of the total nucleic acids in one or more samples. In some embodiments, the primers are the probes attached to the array of the disclosure.

In some embodiments, barcoding (e.g., stochastically barcoding) the plurality of targets in the sample further comprises generating an indexed library of the barcoded targets (e.g., stochastically barcoded targets) or barcoded fragments of the targets. The barcode sequences of different barcodes (e.g., the molecular labels of different stochastic barcodes) can be different from one another. Generating an indexed library of the barcoded targets includes generating a plurality of indexed polynucleotides from the plurality of targets in the sample. For example, for an indexed library of the barcoded targets comprising a first indexed target and a second indexed target, the label region of the first indexed polynucleotide can differ from the label region of the second indexed polynucleotide by, by about, by at least, or by at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, or a number

40

or a range between any two of these values, nucleotides. In some embodiments, generating an indexed library of the barcoded targets includes contacting a plurality of targets, for example mRNA molecules, with a plurality of oligonucleotides including a poly(T) region and a label region; and conducting a first strand synthesis using a reverse transcriptase to produce single-strand labeled cDNA molecules each comprising a cDNA region and a label region, wherein the plurality of targets includes at least two mRNA molecules of different sequences and the plurality of oligonucleotides includes at least two oligonucleotides of different sequences. Generating an indexed library of the barcoded targets can further comprise amplifying the single-strand labeled cDNA molecules to produce double-strand labeled cDNA molecules; and conducting nested PCR on the double-strand labeled cDNA molecules to produce labeled amplicons. In some embodiments, the method can include generating an adaptor-labeled amplicon.

Barcode (e.g., stochastic barcoding) can include using nucleic acid barcodes or tags to label individual nucleic acid (e.g., DNA or RNA) molecules. In some embodiments, it involves adding DNA barcodes or tags to cDNA molecules as they are generated from mRNA. Nested PCR can be performed to minimize PCR amplification bias. Adaptors can be added for sequencing using, for example, next generation sequencing (NGS). The sequencing results can be used to determine cell labels, molecular labels, and sequences of nucleotide fragments of the one or more copies of the targets, for example at block 232 of FIG. 2.

FIG. 3 is a schematic illustration showing a non-limiting exemplary process of generating an indexed library of the barcoded targets (e.g., stochastically barcoded targets), such as barcoded mRNAs or fragments thereof. As shown in step 1, the reverse transcription process can encode each mRNA molecule with a unique molecular label sequence, a cell label sequence, and a universal PCR site. In particular, RNA molecules 302 can be reverse transcribed to produce labeled cDNA molecules 304, including a cDNA region 306, by hybridization (e.g., stochastic hybridization) of a set of barcodes (e.g., stochastic barcodes) 310 to the poly(A) tail region 308 of the RNA molecules 302. Each of the barcodes 310 can comprise a target-binding region, for example a poly(dT) region 312, a label region 314 (e.g., a barcode sequence or a molecule), and a universal PCR region 316.

In some embodiments, the cell label sequence can include 3 to 20 nucleotides. In some embodiments, the molecular label sequence can include 3 to 20 nucleotides. In some embodiments, each of the plurality of stochastic barcodes further comprises one or more of a universal label and a cell label, wherein universal labels are the same for the plurality of stochastic barcodes on the solid support and cell labels are the same for the plurality of stochastic barcodes on the solid support. In some embodiments, the universal label can include 3 to 20 nucleotides. In some embodiments, the cell label comprises 3 to 20 nucleotides.

In some embodiments, the label region 314 can include a barcode sequence or a molecular label 318 and a cell label 320. In some embodiments, the label region 314 can include one or more of a universal label, a dimension label, and a cell label. The barcode sequence or molecular label 318 can be, can be about, can be at least, or can be at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, or a number or a range between any of these values, of nucleotides in length. The cell label 320 can be, can be about, can be at least, or can be at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, or a number or a range between any of these values, of nucleotides in length. The

universal label can be, can be about, can be at least, or can be at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, or a number or a range between any of these values, of nucleotides in length. Universal labels can be the same for the plurality of stochastic barcodes on the solid support and cell labels are the same for the plurality of stochastic barcodes on the solid support. The dimension label can be, can be about, can be at least, or can be at most 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, or a number or a range between any of these values, of nucleotides in length.

In some embodiments, the label region **314** can comprise, comprise about, comprise at least, or comprise at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, or a number or a range between any of these values, different labels, such as a barcode sequence or a molecular label **318** and a cell label **320**. Each label can be, can be about, can be at least, or can be at most 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, or a number or a range between any of these values, of nucleotides in length. A set of barcodes or stochastic barcodes **310** can contain, contain about, contain at least, or can be at most, 10, 20, 40, 50, 70, 80, 90,  $10^2$ ,  $10^3$ ,  $10^4$ ,  $10^5$ ,  $10^6$ ,  $10^7$ ,  $10^8$ ,  $10^9$ ,  $10^{10}$ ,  $10^{11}$ ,  $10^{12}$ ,  $10^{13}$ ,  $10^{14}$ ,  $10^{15}$ ,  $10^{20}$ , or a number or a range between any of these values, barcodes or stochastic barcodes **310**. And the set of barcodes or stochastic barcodes **310** can, for example, each contain a unique label region **314**. The labeled cDNA molecules **304** can be purified to remove excess barcodes or stochastic barcodes **310**. Purification can comprise Ampure bead purification.

As shown in step **2**, products from the reverse transcription process in step **1** can be pooled into 1 tube and PCR amplified with a 1<sup>st</sup> PCR primer pool and a 1<sup>st</sup> universal PCR primer. Pooling is possible because of the unique label region **314**. In particular, the labeled cDNA molecules **304** can be amplified to produce nested PCR labeled amplicons **322**. Amplification can comprise multiplex PCR amplification. Amplification can comprise a multiplex PCR amplification with 96 multiplex primers in a single reaction volume. In some embodiments, multiplex PCR amplification can utilize, utilize about, utilize at least, or utilize at most, 10, 20, 40, 50, 70, 80, 90,  $10^2$ ,  $10^3$ ,  $10^4$ ,  $10^5$ ,  $10^6$ ,  $10^7$ ,  $10^8$ ,  $10^9$ ,  $10^{10}$ ,  $10^{11}$ ,  $10^{12}$ ,  $10^{13}$ ,  $10^{14}$ ,  $10^{15}$ ,  $10^{20}$ , or a number or a range between any of these values, multiplex primers in a single reaction volume. Amplification can comprise using a 1<sup>st</sup> PCR primer pool **324** comprising custom primers **326A-C** targeting specific genes and a universal primer **328**. The custom primers **326** can hybridize to a region within the cDNA portion **306'** of the labeled cDNA molecule **304**. The universal primer **328** can hybridize to the universal PCR region **316** of the labeled cDNA molecule **304**.

As shown in step **3** of FIG. 3, products from PCR amplification in step **2** can be amplified with a nested PCR primers pool and a 2<sup>nd</sup> universal PCR primer. Nested PCR can minimize PCR amplification bias. In particular, the nested PCR labeled amplicons **322** can be further amplified by nested PCR. The nested PCR can comprise multiplex PCR with nested PCR primers pool **330** of nested PCR primers **332a-c** and a 2<sup>nd</sup> universal PCR primer **328'** in a single reaction volume. The nested PCR primer pool **328** can contain, contain about, contain at least, or contain at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, or a number or a range between any of these values, different nested PCR primers **330**. The nested PCR primers **332** can contain an adaptor **334** and hybridize to a region within the cDNA

portion **306''** of the labeled amplicon **322**. The universal primer **328'** can contain an adaptor **336** and hybridize to the universal PCR region **316** of the labeled amplicon **322**. Thus, step **3** produces adaptor-labeled amplicon **338**. In some embodiments, nested PCR primers **332** and the 2<sup>nd</sup> universal PCR primer **328'** may not contain the adaptors **334** and **336**. The adaptors **334** and **336** can instead be ligated to the products of nested PCR to produce adaptor-labeled amplicon **338**.

As shown in step **4**, PCR products from step **3** can be PCR amplified for sequencing using library amplification primers. In particular, the adaptors **334** and **336** can be used to conduct one or more additional assays on the adaptor-labeled amplicon **338**. The adaptors **334** and **336** can be hybridized to primers **340** and **342**. The one or more primers **340** and **342** can be PCR amplification primers. The one or more primers **340** and **342** can be sequencing primers. The one or more adaptors **334** and **336** can be used for further amplification of the adaptor-labeled amplicons **338**. The one or more adaptors **334** and **336** can be used for sequencing the adaptor-labeled amplicon **338**. The primer **342** can contain a plate index **344** so that amplicons generated using the same set of barcodes or stochastic barcodes **310** can be sequenced in one sequencing reaction using next generation sequencing (NGS).

#### Methods and Compositions for Labeling DNA

Disclosed herein include methods and compositions to identify accessible DNA regions in single cells using ATAC-seq (Assay for Transposase-Accessible Chromatin)-seq. These methods and compositions are amendable to a variety of single cell analysis platforms, such as, for example, the BD Rhapsody System. ATAC-seq is a widely used epigenomics tool to identify the open chromatin regions of genome. But single cell ATAC-Seq (scATAC-Seq) is not part of the currently available single cell RNAseq platforms such as the BD Rhapsody System. In some embodiments of the methods and compositions provided herein, Tn5 transposase is employed for fragmentation (e.g., cut the DNA on open chromatin region) and the fragmented DNA can be captured by (i) solid supports (e.g., Rhapsody beads) through oligonucleotides specific for the ATACseq DNA fragment (e.g., as shown in FIGS. 5A-5D) and/or (ii) through a coupling oligonucleotide (e.g., SPLINT oligo) to the solid supports (e.g., Rhapsody beads) comprising oligonucleotides with a capture sequence either the same or different than the one used to capture mRNA (e.g., as shown in FIGS. 6A-6D).

The compositions and methods provided herein achieve scATAC-Seq at a much higher throughput than currently available methods. For example, the compositions and methods disclosed herein, when employed on the BD Rhapsody platform, can achieve up to about 20,000 cells or nuclei per cartridge. This level of throughput is far greater than, for example, the only 800 cells per run that can be done with the C1 Fluidigm platform, or the at most 10,000 nuclei with 10 $\times$  Chromium. Moreover, the 10 $\times$  chromium ATAC-Seq analyzes DNA sequence only, while the methods and compositions provided herein can enable the analysis of DNA and RNA (e.g., mRNA) simultaneously. There are provided, in some embodiments, solid supports (e.g., Rhapsody capture bead) that can capture both DNA and RNA information, such as, for example, beads loaded with oligonucleotides with dT sequence to capture mRNAs and a specific sequence to capture the DNA fragments generated by ATAC-Seq.

Methods and compositions for labeling gDNA and determining information relating to the gDNA have been described in U.S. Patent Application Publication No. 2019/0338357, and in U.S. patent application Ser. No. 16/934,

530, filed Jul. 21, 2020, entitled "SINGLE CELL CHROMATIN IMMUNOPRECIPITATION SEQUENCING ASSAY"; the content of each of these applications is incorporated herein by reference in its entirety.

The methods and systems described herein can be used with methods and systems using antibodies associated with (e.g., attached to or conjugated with) oligonucleotides (also referred to herein as AbOs or AbOligos). Embodiments of using AbOs to determine protein expression profiles in single cells and tracking sample origins have been described in U.S. patent application Ser. No. 15/715,028, published as U.S. Patent Application Publication No. 2018/0088112, and U.S. patent application Ser. No. 15/937,713, published as U.S. Patent Application Publication No. 2018/0346970; the content of each is incorporated by reference herein in its entirety.

There are provided, in some embodiments, single cell assays for transposase-accessible chromatin (scATAC)-Seq. The methods and compositions provided herein can enable the labeling and identification of single cell DNA (e.g., open chromatin-associated gDNA). FIGS. 4A-B are non-limiting schematic illustrations of compositions and methods provided herein for labeling DNA (e.g., gDNA) in a single cell. A single cell **402** can undergo a permeabilizing step. The single cell can comprise a plasma membrane **404** and a nuclear membrane **406**. The plasma membrane **404** and/or the nuclear membrane **406** can be permeabilized. Cell permeabilization can be performed using digitonin. In some embodiments, the workflow comprises nuclear isolation. The single cell can comprise gDNA. The gDNA can be associated with a histone **408** to form closed chromatin **410**. The gDNA can comprise open chromatin **412** (e.g., not associated with nucleosomes). In some embodiments, a transposome **414** is provided. The transposome **414** can comprise a homodimer or a heterodimer. The transposome can comprise a transposase. The transposase can comprise Tn5 transposase. Tn5 transposase can comprise a first and second transposase monomers **416**. The transposome **414** can comprise a first adaptor **418a** having a first 5' overhang and a second adaptor **418b** having a second 5' overhang. The workflow can comprise introducing **400a** the transposome into the permeabilized cell and/or nucleus (which in, some embodiments, are isolated) thereby generating a cell and/or nucleus **420** wherein the transposome is associated with open chromatin and subsequently generates a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang. The workflow can comprise bulk tagging, or can comprise tagging of single cells and/or single nuclei in partitions. The plurality of dsDNA fragments generated can be barcoded according to the methods disclosed herein. The tagging step can comprise an about 30 min incubation at a temperature of about 37° C. The workflow can comprise partitioning a plurality of solid supports and a plurality of single cells and/or single nuclei into a plurality of partitions (e.g., a microwell array comprising a plurality of microwells) as shown in FIG. 4B. For example, a first partition **422a** (e.g., a microwell) of the plurality of partitions (e.g., a microwell array) can comprise a first solid support **424a** and a first single cell **420a**. The first single cell **420a** can comprise a plurality of dsDNA fragments generated by the transposome. The first solid support **424a** can comprise a first, second, and/or third plurality of oligonucleotide barcodes **426a**. A second partition **422b** of the plurality of partitions can comprise a second solid support **424b** and a second single cell **420b**. The second single cell **420b** can comprise a plurality of dsDNA fragments generated by the transposome. The second solid

support **406b** can comprise a first, second, and/or third plurality of oligonucleotide barcodes **426b**. Each of the oligonucleotide barcodes can comprise a cell label sequence. The cell labels of the plurality of oligonucleotide barcodes situated in the same partition can comprise the same cell label sequence. The cellular label sequences of the pluralities of oligonucleotide barcodes situated in different partitions can different from each other. The workflow can comprise cell lysis and release of the plurality of dsDNA fragments. Fragmented DNA from single cell and/or single nuclei can captured by of oligonucleotide barcodes **426**. Cell lysis can comprise contacting cells with SDS and/or Proteinase K at 65° C.

FIGS. 5A-5D is a schematic illustration of a non-limiting exemplary workflow for labeling DNA (e.g., gDNA) in a single cell. In some embodiments, the transposome is provided comprising a transposase, a first adaptor having a first 5' overhang, and a second adaptor having a second 5' overhang. The first 5' overhang can comprise a complement of a first target-binding region. The second 5' overhang can comprise a universal sequence (e.g., a second universal sequence). In some embodiments, the workflow comprises contacting double-stranded deoxyribonucleic acid (dsDNA) with a transposome to generate a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang. Each dsDNA fragment of the plurality of dsDNA fragments can comprise a first strand comprising the first 5' overhang and a second strand comprising the second 5' overhang. In some embodiments, the plurality of dsDNA fragments can comprise three species which vary with regards to their overhangs due to being generated by a transposome comprising: (i) two copies of a first adaptor; (ii) one copy of a first adaptor and one copy of a second adaptor, or (iii) two copies of a second adaptor. For example, the plurality of dsDNA fragments can comprise: (i) dsDNA fragment **524a** comprising a gDNA sequence **526** flanked by Tn5 transposase recognition sequences (e.g., Mosaic End (ME **528**)), wherein dsDNA fragment **524a** comprises a first strand comprising a first 5' overhang comprising a complement of a first target-binding region **516rc** and a second strand comprising a second 5' overhang comprising a complement of a first target-binding region **516rc**; (ii) dsDNA fragment **524b** comprising a gDNA sequence **526** flanked by Tn5 transposase recognition sequences (e.g., Mosaic End (ME **528**)), wherein dsDNA fragment **524b** comprises a first strand comprising a first 5' overhang comprising a complement of a first target-binding region **516rc** and a second strand comprising a second 5' overhang comprising a second universal sequence **530**; and (iii) dsDNA fragment **524c** comprising a gDNA sequence **526** flanked by Tn5 transposase recognition sequences (e.g., Mosaic End (ME **528**)), wherein dsDNA fragment **524c** comprises a first strand comprising a first 5' overhang comprising a second universal sequence **530** and a second strand comprising a second 5' overhang comprising a second universal sequence **530**. In some embodiments, dsDNA fragment **524a** will not be amplify due to suppressive PCR. In some embodiments, dsDNA fragment **524b** is included in the ATAC-Seq library. In some embodiments, dsDNA fragment **524c** will not be captured by oligonucleotide barcodes. In some embodiments, there can be issues of secondary hairpin structure formation of ATAC-Seq library if the same seq tag was added in both ends. In some embodiments, the overhangs disclosed herein and/or PCR primer design can eliminate this issue. The first universal sequence **506** and second universal sequence **530** can be different (e.g., comprise different Illumina primer sequences, complements

thereof, and/or portions thereof). The ME **528** can be about 19 nucleotides in length. The plurality of dsDNA fragments can comprise a gap **525**. The gap **525** can be about 9 nucleotides in length. The 5' overhangs can be about 14 nucleotides in length.

The workflow can, in some embodiments, comprise simultaneous ATAC-Seq and RNAseq. The workflow can comprise cell lysis to release Tn5 fragments (e.g., under conditions optimized to preserve mRNAs, such as, for example, 72° C. incubation for 5 minutes) and mRNA and capturing with beads (e.g., Rhapsody beads) through dT oligos. The workflow can comprise one or more RNase inhibitors to protect mRNAs. A barcode (e.g., a stochastic barcode, an oligonucleotide barcode of the third plurality of oligonucleotide barcodes, an oligonucleotide barcode **504**) can comprise a third target-binding region **512** (e.g., a poly(dT)) that can bind to nucleic acid targets (e.g., polyadenylated RNA transcripts **518** or other nucleic acid targets, such as for example, binding reagent oligonucleotides, whether associated with a binding reagent or dissociated from the binding reagent) via a poly(dA) tail **522**, or other nucleic acid targets, for labeling or barcoding (e.g., unique labeling). The third target-binding region **512** can comprise a gene-specific sequence, an oligo(dT) sequence, a random multimer, or any combination thereof. The oligonucleotide barcode **504** can also comprise a number of labels. The oligonucleotide barcode **504** can include molecular label (ML) **510** and a sample label (e.g., partition label, cell label (CL) **508**) for labeling the transcripts and/or tracking sample origins of the RNA transcripts (or nucleic acid targets, such as for example, antibody oligonucleotides, whether associated with antibodies or have dissociated from antibodies), respectively, along with one or more additional sequences flanking the molecular label **510**/cell label **508** region of each oligonucleotide barcode **504** for subsequent reactions, such as, for example, a first universal sequence **506** (e.g., Read 1 sequence). In some embodiments, the oligonucleotide barcode **504** is associated with a solid support (e.g., a particle **502**). The oligonucleotide barcode **504** can be associated with particle **502** via its 5' end. A plurality of barcodes **504** can be associated with particle **502**. A barcode (e.g., a stochastic barcode, an oligonucleotide barcode of the first plurality of oligonucleotide barcodes, an oligonucleotide barcode **514**) can comprise a first target-binding region **516** (e.g., a poly(dT)) that can bind to first 5' overhang **516rc** of the first strand of a dsDNA fragment. The first target-binding region **516** can comprise a gene-specific sequence, an oligo(dT) sequence, a random multimer, any predetermined sequence, or any combination thereof. The oligonucleotide barcode **514** can also comprise a number of labels. The oligonucleotide barcode **514** can include molecular label (ML) **510** and a sample label (e.g., partition label, cell label (CL) **508**) for labeling the transcripts and/or tracking sample origins dsDNA fragments, respectively, along with one or more additional sequences flanking the molecular label **510**/cell label **508** region of each oligonucleotide barcode **514** for subsequent reactions, such as, for example, a first universal sequence **506** (e.g., Read 1 sequence). In some embodiments, the oligonucleotide barcode **514** is associated with a solid support (e.g., a particle **502**). The oligonucleotide barcode **514** can be associated with particle **502** via its 3' end. A plurality of barcodes **514** can be associated with particle **502**.

In some embodiments, the particle is a bead. The bead can be a polymeric bead, for example a deformable bead or a gel bead, functionalized with barcodes or stochastic barcodes (such as hydrogel beads from 10x Genomics, San Francisco,

CA). In some implementation, a gel bead can comprise a polymer-based gels. Gel beads can be generated, for example, by encapsulating one or more polymeric precursors into droplets. Upon exposure of the polymeric precursors to an accelerator (e.g., tetramethylethylenediamine (TEMED)), a hydrogel bead may be generated.

Methods for immobilization of oligonucleotides to solid substrates are well established. A variety of chemical modifications can be used for covalent coupling (directly or via a linker) of nucleic acids to solid supports (e.g., beads) via their 5' and/or 3' ends. To allow binding of nucleic acid molecules, the solid support can be functionalized to expose nucleophilic groups, which can react with reactive groups on the nucleic acid. Alternatively, reactive groups can be introduced into the solid support to react with nucleophiles present in the nucleic acid. Suitable groups or moieties include hydroxyl, thiol, amino and activated carboxylic acid groups, while groups capable of reacting with these include dichlorotriazinyl, alkyl epoxy, maleimide, bromoacetyl and others. Strategies for coupling nucleic acids via their 5' and/or 3' ends to different types of solid supports (e.g., functionalized solid supports) are described, for example, in Gosh and Musso (Nuc. acid Res., 15(13), 5353-5372, 1987); "Strategies for Attaching Oligonucleotides to Solid Supports", Technical Bulletin, Integrated DNA Technologies, 2014(v6); Hermanson, Bioconjugate Techniques, Second Edition (Academic Press, 2008); Chrisey, et al. Covalent attachment of synthetic DNA to self-assembled monolayer films (1996) Nucleic Acids Research, vol. 24, No. 15 3031-3039; Guo et al., Nucleic Acids Res. 22:5456-5465 (1994); Pease et al., Proc. Natl. Acad. Sci. USA 91(11): 5022-5026 (1994); Khrapko et al., Mol Biol (Mosk) (USSR) 25:718-730 (1991); Stimpson et al., Proc. Natl. Acad. Sci. USA 92:6379-6383 (1995); U.S. Pat. No. 5,871,928 to Fodor et al.; U.S. Pat. No. 5,654,413 to Brenner; U.S. Pat. Nos. 5,429,807; and 5,599,695 to Pease et al; the content of each is incorporated by reference in its entirety. Depending on the presence of reactive or nucleophilic groups on the solid support and nucleic acid molecule, the coupling may be performed directly or with bifunctional reagents or cross-linking agents. Bifunctional and coupling reagents are well known in the art and are available from commercial sources such as Thermo Fisher Scientific, Sigma-Aldrich, ProteoChem, and the like.

The workflow can comprise hybridization **500a** of the dsDNA fragment **524b** and oligonucleotide barcode **514**. The workflow can comprise hybridization **500a** of the polyadenylated RNA transcript **518** and oligonucleotide barcode **504**. In some embodiments, the extension reaction **500b** can comprise extending the oligonucleotide barcode **504** hybridized to the polyadenylated RNA transcript **518** to generate a barcoded nucleic acid molecule **532** comprising cDNA **520c** (the reverse complementary sequence of RNA sequence **520r**).

In some embodiments, the workflow can comprise, before ligating the second strand of the dsDNA fragment **524b** to oligonucleotide barcode **514**, filling a gap **527** between the second strand of the dsDNA fragment **524b** and hybridized oligonucleotide barcode **514b** with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity (e.g., by Klenow extension **500b**). In some embodiments, the workflow can comprise filling the one or more gaps **525** with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity (e.g., by Klenow extension **500b**). The workflow can comprise barcoding dsDNA fragment **524b** using a oligonucleotide barcode **514** to generate a barcoded

DNA fragment **536** (e.g., the workflow can comprise ligating **500b** the second strand of dsDNA fragment **524b** to hybridized oligonucleotide barcode **514**).

The workflow can comprise, in some embodiments, barcoding dsDNA fragment **524a** using an oligonucleotide barcode **514** to generate a barcoded DNA fragment **534** (e.g., the workflow can comprise ligating **500b** the second strand of dsDNA fragment **524a** to hybridized oligonucleotide barcode **514**). In some embodiments, barcoded DNA fragment **534** is subjected to one or more downstream extension, amplification, and/or sequencing reactions (e.g., amplification using primers hybridizing to **506** and **516rc**, or complements thereof).

The workflow can comprise denaturation **500c** (e.g., with use of heating and/or chemicals). The workflow can comprise downstream **500d** primer extension, amplification and/or sequencing of barcoded DNA fragment **536** and/or barcoded nucleic acid molecule **532**. Barcoded DNA fragment **536** and/or barcoded nucleic acid molecule **532** can serve as a template for one or more extension reactions (e.g., random priming and extension) and/or amplification reactions (e.g., PCR). For example, barcoded nucleic acid molecule **532** can be contacted with random primers **540** and a primer **538** that can anneal to first universal sequence **506** (or a complement thereof). For example, barcoded DNA fragment **536** can undergo a first round of amplification ("PCR1") employing amplification primers **538** and **542** that can anneal to first universal sequence and second universal sequence (or complements thereof), respectively, thereby generating a fourth plurality of barcoded DNA fragments. In some embodiments, amplifying the plurality of barcoded DNA fragments comprises adding sequences of binding sites of sequencing primers and/or sequencing adaptors, complementary sequences thereof, and/or portions thereof, to the plurality of barcoded DNA fragments. In some embodiments, obtaining sequence data of the plurality of barcoded DNA fragments, or products thereof, comprises obtaining sequence information of the fourth plurality of barcoded amplicons, or products thereof.

FIGS. 6A-6D is a schematic illustration of a non-limiting exemplary workflow for labeling DNA (e.g., gDNA) in a single cell. In some embodiments, the transposome is provided comprising a transposase, a first adaptor having a first 5' overhang, and a second adaptor having a second 5' overhang. The first 5' overhang can comprise a complement of a second target-binding region. The second 5' overhang can comprise a universal sequence (e.g., a second universal sequence). In some embodiments, the workflow comprises contacting double-stranded deoxyribonucleic acid (dsDNA) with a transposome to generate a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang. Each dsDNA fragment of the plurality of dsDNA fragments can comprise a first strand comprising the first 5' overhang and a second strand comprising the second 5' overhang. There is provided, in some embodiments, a coupling oligonucleotide **634** comprising a 5' complement of the coupling sequence **630rc** and a 3' complement of a second target-binding region **616rc**. In some embodiments, the plurality of dsDNA fragments can comprise three species which vary with regards to their overhangs due to being generated by a transposome comprising: (i) two copies of a first adaptor, (ii) one copy of a first adaptor and one copy of a second adaptor, or (iii) two copies of a second adaptor. For example, the plurality of dsDNA fragments can comprise: (i) dsDNA fragment **624a** comprising a gDNA sequence **626** flanked by Tn5 transposase recognition sequences (e.g., Mosaic End (ME **628**)), wherein dsDNA fragment **624a**

comprises a first strand comprising a first 5' overhang comprising a coupling sequence **630** and a second strand comprising a second 5' overhang comprising a coupling sequence **630**; (ii) dsDNA fragment **624b** comprising a gDNA sequence **626** flanked by Tn5 transposase recognition sequences (e.g., Mosaic End (ME **628**)), wherein dsDNA fragment **624b** comprises a first strand comprising a first 5' overhang comprising a coupling sequence **630** and a second strand comprising a second 5' overhang comprising a second universal sequence **632**; and (iii) dsDNA fragment **624c** comprising a gDNA sequence **626** flanked by Tn5 transposase recognition sequences (e.g., Mosaic End (ME **628**)), wherein dsDNA fragment **624c** comprises a first strand comprising a first 5' overhang comprising a second universal sequence **632** and a second strand comprising a second 5' overhang comprising a second universal sequence **632**. In some embodiments, dsDNA fragment **624a** will not be amplified due to suppressive PCR. In some embodiments, dsDNA fragment **624b** is included in the ATAC-Seq library. In some embodiments, dsDNA fragment **624c** will not be captured by oligonucleotide barcodes. In some embodiments, there can be issues of secondary hairpin structure formation of ATAC-Seq library if the same seq tag was added in both ends. In some embodiments, the overhangs disclosed herein and/or PCR primer design can eliminate this issue. The first universal sequence **606** and second universal sequence **632** can be different (e.g., comprise different Illumina primer sequences, complements thereof, and/or portions thereof). The ME **628** can be about 19 nucleotides in length. The plurality of dsDNA fragments can comprise a gap **625**. The gap **625** can be about 9 nucleotides in length. The 5' overhangs can be about 14 nucleotides in length.

The workflow can comprise simultaneous ATAC-Seq and RNAseq. The workflow can comprise cell lysis to release Tn5 fragments (e.g., under conditions optimized to preserve mRNAs, such as, for example, 72° C. incubation for 5 minutes) and mRNA and capturing with beads (e.g., Rhapsody beads) through dT oligos. The workflow can comprise one or more RNase inhibitors to protect mRNAs. A barcode (e.g., a stochastic barcode, an oligonucleotide barcode of the third plurality of oligonucleotide barcodes, an oligonucleotide barcode **604**) can comprise a third target-binding region **612** (e.g., a poly(dT)) that can bind to nucleic acid targets (e.g., poly-adenylated RNA transcripts **618** or other nucleic acid targets, such as for example, binding reagent oligonucleotides, whether associated with a binding reagent or dissociated from the binding reagent) via a poly(dA) tail **622**, or other nucleic acid targets, for labeling or barcoding (e.g., unique labeling). The third target-binding region **612** can comprise a gene-specific sequence, an oligo(dT) sequence, a random multimer, or any combination thereof. The oligonucleotide barcode **604** can also comprise a number of labels. The oligonucleotide barcode **604** can include molecular label (ML) **610** and a sample label (e.g., partition label, cell label (CL) **608**) for labeling the transcripts and/or tracking sample origins of the RNA transcripts (or nucleic acid targets, such as for example, antibody oligonucleotides, whether associated with antibodies or have dissociated from antibodies), respectively, along with one or more additional sequences flanking the molecular label **610**/cell label **608** region of each oligonucleotide barcode **604** for subsequent reactions, such as, for example, a first universal sequence **606** (e.g., Read 1 sequence). In some embodiments, the oligonucleotide barcode **604** is associated with a solid support (e.g., a particle **602**). The oligonucleotide barcode **604** can associate with particle **602** via its 5' end. A

plurality of barcodes **604** can be associated with particle **602**. A barcode (e.g., a stochastic barcode, an oligonucleotide barcode of the second plurality of oligonucleotide barcodes, an oligonucleotide barcode **614**) can comprise a second target-binding region **616** (e.g., a poly(dT)) that can bind to the 3' complement of a second target-binding region **616rc** of coupling oligonucleotide **634**. The second target-binding region **616** can comprise a gene-specific sequence, an oligo(dT) sequence, a random multimer, any predetermined sequence, or any combination thereof. The oligonucleotide barcode **614** can also comprise a number of labels. The oligonucleotide barcode **614** can include molecular label (ML) **610** and a sample label (e.g., partition label, cell label (CL) **608**) for labeling the transcripts and/or tracking sample origins dsDNA fragments, respectively, along with one or more additional sequences flanking the molecular label **610**/cell label **608** region of each oligonucleotide barcode **614** for subsequent reactions, such as, for example, a first universal sequence **606** (e.g., Read 1 sequence). In some embodiments, the oligonucleotide barcode **614** is associated with a solid support (e.g., a particle **602**). The oligonucleotide barcode **614** can associate with particle **602** via its 5' end. A plurality of barcodes **614** can be associated with particle **602**. In some embodiments, the particle is a bead. The bead can be a polymeric bead, for example a deformable bead or a gel bead, functionalized with barcodes or stochastic barcodes (such as hydrogel beads from 10x Genomics, San Francisco, CA). In some implementation, a gel bead can comprise a polymer-based gels. Gel beads can be generated, for example, by encapsulating one or more polymeric precursors into droplets. Upon exposure of the polymeric precursors to an accelerator (e.g., tetramethylethylenediamine (TEMED)), a hydrogel bead may be generated.

The workflow can comprise hybridization **600a** of the coupling sequence of the first strand of a dsDNA fragment **624b** with the a 5' complement of the coupling sequence **630rc** of the coupling oligonucleotide **634**. The workflow can comprise hybridization **600a** of 3' complement of the second target-binding region **616rc** of the coupling oligonucleotide **634** with the second target-binding region **616** of oligonucleotide barcode **614**. The workflow can comprise hybridization **600a** of the poly-adenylated RNA transcript **618** and oligonucleotide barcode **604**.

In some embodiments, the extension reaction **600b** can comprise extending the oligonucleotide barcode **604** hybridized to the poly-adenylated RNA transcript **618** to generate a barcoded nucleic acid molecule **640** comprising cDNA **620c** (the reverse complementary sequence of RNA sequence **620r**). In some embodiments, the workflow can comprise, before ligating the first strand of the dsDNA fragment **624b** to oligonucleotide barcode **614**, filling a gap **627** between the first strand of the dsDNA fragment **624b** and the hybridized oligonucleotide barcode **614** with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity (e.g., by Klenow extension **600b**).

The workflow can comprise barcoding dsDNA fragment **624b** using a oligonucleotide barcode **614** to generate barcoded DNA fragment **638a** and/or barcoded DNA fragment **638b**. In some embodiments, the workflow can comprise, before ligating the second strand of the dsDNA fragment **624b** to coupling oligonucleotide **634**, filling a gap **629** between the second strand of the dsDNA fragment **624b** and the coupling oligonucleotide **634** with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity (e.g., by Klenow extension **600b**).

The workflow can comprise ligating **600b** the first strand of the dsDNA fragment **624b** to the hybridized oligonucleotide barcode **614** and/or ligating **600b** second strand of the dsDNA fragment **624b** to coupling oligonucleotide **634** (e.g., with a DNA ligase). In some embodiments, the workflow can comprise filling the one or more gaps **625** with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity (e.g., by Klenow extension **600b**). The workflow can comprise extension **600b** of the first and/or second strands of dsDNA fragment **624b**.

The workflow can comprise barcoding dsDNA fragment **624a** using a oligonucleotide barcode **614** to generate barcoded DNA fragment **636a** and/or barcoded DNA fragment **636b** as shown in FIGS. 6A-6D. In some embodiments, barcoded DNA fragment **636a** and/or barcoded DNA fragment **636b** is subjected to one or more downstream extension, amplification, and/or sequencing reactions (e.g., amplification using primers hybridizing to **606** and **630rc**, or complements thereof).

The workflow can comprise denaturation **600c** (e.g., with use of heating and/or chemicals). The workflow can comprise downstream **600d** primer extension, amplification and/or sequencing of barcoded nucleic acid molecule **640**, barcoded DNA fragment **638a** and/or barcoded DNA fragment **638b**. Barcoded nucleic acid molecule **640**, barcoded DNA fragment **638a** and/or barcoded DNA fragment **638b** can serve as a template for one or more extension reactions (e.g., random priming and extension) and/or amplification reactions (e.g., PCR). For example, barcoded nucleic acid molecule **640** can be contacted with random primers **644** and a primer **642** that can anneal to first universal sequence **606** (or a complement thereof). For example, barcoded DNA fragment **638a** and/or barcoded DNA fragment **638b** an undergo a first round of amplification ("PCR1") employing amplification primers **642** and **646** that can anneal to first universal sequence and second universal sequence (or complements thereof), respectively, thereby generating a fourth plurality of barcoded DNA fragments. In some embodiments, amplifying the plurality of barcoded DNA fragments comprises adding sequences of binding sites of sequencing primers and/or sequencing adaptors, complementary sequences thereof, and/or portions thereof, to the plurality of barcoded DNA fragments. In some embodiments, obtaining sequence data of the plurality of barcoded DNA fragments, or products thereof, comprises obtaining sequence information of the fourth plurality of barcoded amplicons, or products thereof.

There are provided, in some embodiments, compositions and methods for labeling DNA (e.g., open chromatin-associated gDNA). In some embodiments, the method comprises: contacting double-stranded deoxyribonucleic acid (dsDNA) with a transposome to generate a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang, wherein each dsDNA fragment of the plurality of dsDNA fragments comprises a first strand comprising the first 5' overhang and a second strand comprising the second 5' overhang, wherein the transposome comprises a transposase, a first adaptor having the first 5' overhang, and a second adaptor having the second 5' overhang; and barcoding the plurality of dsDNA fragments, or products thereof, using a first plurality of oligonucleotide barcodes to generate a plurality of barcoded DNA fragments, wherein the 3' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes is associated with a solid support, and wherein the 5' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes comprises a

first target-binding region capable of hybridizing to the first 5' overhang of at least one of the plurality of dsDNA fragments. In some embodiments, barcoding the plurality of dsDNA fragments, or products thereof, using a first plurality of oligonucleotide barcodes comprises: hybridizing the first 5' overhang of the first strand of a dsDNA fragment with the first target-binding region of an oligonucleotide barcode of the first plurality of oligonucleotide barcodes; and ligating the second strand of said dsDNA fragment to said hybridized oligonucleotide barcode. The method can comprise: before ligating the second strand to the oligonucleotide barcode, filling a gap between the second strand and said hybridized oligonucleotide barcode with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. Ligating the second strand to the oligonucleotide barcode can be performed using a DNA ligase. The first 5' overhang can comprise a complement of a first target-binding region.

There are provided, in some embodiments, compositions and methods for labeling DNA (e.g., open chromatin-associated gDNA). In some embodiments, the method comprises: contacting double-stranded deoxyribonucleic acid (dsDNA) with a transposome to generate a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang, wherein each dsDNA fragment of the plurality of dsDNA fragments comprises a first strand comprising the first 5' overhang and a second strand comprising the second 5' overhang, wherein the transposome comprises a transposase, a first adaptor having the first 5' overhang, and a second adaptor having the second 5' overhang, and wherein the first 5' overhang comprises a coupling sequence; providing a coupling oligonucleotide comprising a 5' complement of the coupling sequence and a 3' complement of a second target-binding region; and barcoding the plurality of dsDNA fragments, or products thereof, using a second plurality of oligonucleotide barcodes to generate a plurality of barcoded DNA fragments, wherein each oligonucleotide barcode of the second plurality of oligonucleotide barcodes comprises the second target-binding region. In some embodiments, barcoding the plurality of dsDNA fragments, or products thereof, using a second plurality of oligonucleotide barcodes comprises: hybridizing the coupling sequence of the first strand of a dsDNA fragment with the 5' complement of the coupling sequence of the coupling oligonucleotide; hybridizing the 3' complement of the second target-binding region of the coupling oligonucleotide with the second target-binding region of an oligonucleotide barcode of the second plurality of oligonucleotide barcodes; and ligating the first strand of said dsDNA fragment to said hybridized oligonucleotide barcode and extending the first strand; and/or ligating the second strand of said dsDNA fragment to said coupling oligonucleotide and extending the second strand. The method can comprise: before ligating the first strand to the oligonucleotide barcode, filling a gap between the first strand and the hybridized oligonucleotide barcode with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. The method can comprise: before ligating the second strand to the coupling oligonucleotide, filling a gap between the second strand and the coupling oligonucleotide with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. Ligating the first strand of said dsDNA fragment to said hybridized oligonucleotide barcode and/or ligating the second strand of said dsDNA fragment to said coupling oligonucleotide can be performed with a DNA ligase.

The coupling oligonucleotide can be a single-stranded oligonucleotide or a double-stranded oligonucleotide. The coupling oligonucleotide can be, can be about, can be at least, or can be at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, or a number or a range between any of these values, of nucleotides in length. The coupling sequence can be, can be about, can be at least, or can be at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, or a number or a range between any of these values, of nucleotides in length. The plurality of dsDNA fragments can comprise one or more gaps. The one or more gaps can be, can be about, can be at least, or can be at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, or a number or a range between any of these values, of nucleotides in length. The one or more gaps can be about 9 nucleotides in length. The method can comprise: filling the one or more gaps with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. The first 5' overhang and/or the second 5' overhang can be, can be about, can be at least, or can be at most, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, or a number or a range between any of these values, of nucleotides in length. The first 5' overhang and/or the second 5' overhang can comprise about 14 nucleotides. The second 5' overhang can comprise a second universal sequence. The first adaptor and/or the second adaptor can comprise a DNA end sequence of the transposon (e.g., ME).

A cell can comprise the dsDNA. Contacting dsDNA with a transposome can comprise permeabilizing a cell and/or isolating the nucleus of a cell comprising the dsDNA. The cell can comprise a single cell. The dsDNA can comprise genomic DNA (gDNA). The plurality of dsDNA fragments can comprise open chromatin-associated gDNA. The permeabilizing can comprise chemical or physical permeabilization. The permeabilizing can comprise contacting the cell with a detergent and/or a surfactant. The permeabilizing can comprise permeabilizing the cell by sonification. The method can comprise: permeabilizing a nucleus in the cell to generate a permeabilized nucleus. The method can comprise: fixating the cell comprising the nucleus prior to permeabilizing the nucleus.

The transposase and/or transposome can vary. For example, the transposase can comprise a Tn5 transposase. The transposase can be, for example, a Tn transposase (e.g., Tn3, Tn5, Tn7, Tn10, Tn552, Tn903), a MuA transposase, a Vibhar transposase (e.g., from *Vibrio harveyi*), Ac-Ds, Ascot-1, Bsl, Cin4, Copia, En/Spm, F element, hobo, Hsmar1, Hsmar2, IN (HIV), IS1, IS2, IS3, IS4, IS5, IS6, IS10, IS21, IS30, IS50, IS51, IS150, IS256, IS407, IS427, IS630, IS903, IS911, IS982, IS1031, ISL2, L1, Mariner, P element, Tam3, Tc1, Tc3, Tel, THE-1, Tn/O, TnA, Tn3, Tn5, Tn7, Tn10, Tn552, Tn903, Tol1, Tol2, Tn10, Ty1, any prokaryotic transposase, or any transposase related to and/or derived from those listed above. In some embodiments, a transposase related to and/or derived from a parent transposase can comprise a peptide fragment with at least about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% amino acid sequence homology to a corresponding peptide fragment of the parent transposase. The peptide fragment can be at least about 10, about 15, about 20, about 25, about 30, about 35, about 40, about 45, about 50, about 60, about 70, about 80, about 90, about 100, about 150, about 200, about 250, about 300, about 400, or about 500 amino acids in length. For example, a transposase derived from Tn5 can comprise a peptide

53

fragment that is 50 amino acids in length and about 80% homologous to a corresponding fragment in a parent Tn5 transposase. In some cases, the insertion can be facilitated and/or triggered by addition of one or more cations. The cations can be divalent cations such as, for example, Ca<sup>2+</sup>, Mg<sup>2+</sup> and Mn<sup>2+</sup>.

The method can comprise: obtaining sequence data of the plurality of barcoded DNA fragments, or products thereof. The method can comprise: determining information relating to the gDNA based on the sequences of the plurality of barcoded DNA fragments, or products thereof, in the sequencing data obtained. Determining the information relating to the gDNA can comprise determining chromatin accessibility of the gDNA based on the sequences of the plurality of barcoded DNA fragments, or products thereof, in the sequencing data obtained. In some embodiments, determining the chromatin accessibility of the gDNA comprises: aligning the sequences of the plurality of barcoded DNA fragments to a reference sequence of the gDNA; and identifying regions of the gDNA corresponding the ends of the barcoded DNA fragments of the plurality of barcoded DNA fragments to have an accessibility above a threshold. In some embodiments, determining the chromatin accessibility of the gDNA comprises: aligning the sequences of the plurality of barcoded DNA fragments to a reference sequence of the gDNA; and determining the accessibility of regions of the gDNA corresponding the ends of barcoded DNA fragments of the plurality of barcoded DNA fragments based on the numbers of the barcoded DNA fragments of the plurality of barcoded DNA fragments in the sequencing data.

Determining the information relating to the gDNA can comprise determining methylome information of the gDNA based on the sequences of the plurality of barcoded DNA fragments in the sequencing data obtained. The method can comprise: digesting nucleosomes associated with the double-stranded gDNA. The method can comprise: performing chemical conversion and/or enzymatic conversion of cytosine bases of the plurality of dsDNA fragments, or products thereof, to generate a plurality of converted dsDNA fragments with uracil bases, wherein chemical conversion comprises bisulfate treatment, and wherein enzymatic conversion comprises APOBEC-mediated conversion. Barcoding the plurality of dsDNA fragments, or products thereof, can comprise barcoding the plurality of converted dsDNA fragments, or products thereof. In some embodiments, determining the methylome information comprises: determining a position of the plurality of barcoded DNA fragments in the sequencing data has a thymine base and the corresponding position in a reference sequence of the gDNA has a cytosine base to determine the corresponding position in the gDNA has a 5-methylcytosine (5mC) base and/or 5-hydroxymethylcytosine (5hmC) base.

The cell can comprise copies of a nucleic acid target. In some embodiments, the method comprises contacting a third plurality of oligonucleotide barcodes with the copies of the nucleic acid target for hybridization, wherein each oligonucleotide barcode of the third plurality of oligonucleotide barcodes comprises a first universal sequence, a third target-binding region capable of hybridizing to the copies of the nucleic acid target, and a molecular label; extending the third plurality of oligonucleotide barcodes hybridized to the copies of a nucleic acid target to generate a plurality of barcoded nucleic acid molecules each comprising a sequence complementary to at least a portion of the nucleic acid target; and obtaining sequence information of the plurality of barcoded nucleic acid molecules, or products

54

thereof, to determine the copy number of the nucleic acid target in the cell. Determining the copy number of the nucleic acid target in the cell can comprise determining the copy number of the nucleic acid target in the cell based on the number of molecular labels with distinct sequences associated with the plurality of barcoded nucleic acid molecules, or products thereof.

In some embodiments, the method comprises contacting random primers with the plurality of barcoded nucleic acid molecules, wherein each of the random primers comprises a third universal sequence, or a complement thereof; and extending the random primers hybridized to the plurality of barcoded nucleic acid molecules to generate a plurality of extension products. In some embodiments, the method comprises amplifying the plurality of extension products using primers capable of hybridizing to the first universal sequence or complements thereof, and primers capable of hybridizing the third universal sequence or complements thereof, thereby generating a first plurality of barcoded amplicons. Amplifying the plurality of extension products can comprise adding sequences of binding sites of sequencing primers and/or sequencing adaptors, complementary sequences thereof, and/or portions thereof, to the plurality of extension products.

In some embodiments, the method comprises determining the copy number of the nucleic acid target in the cell based on the number of molecular labels with distinct sequences associated with the first plurality of barcoded amplicons, or products thereof. In some embodiments, the method comprises amplifying the first plurality of barcoded amplicons using primers capable of hybridizing to the first universal sequence or complements thereof, and primers capable of hybridizing the third universal sequence or complements thereof, thereby generating a second plurality of barcoded amplicons. Amplifying the first plurality of barcoded amplicons can comprise adding sequences of binding sites of sequencing primers and/or sequencing adaptors, complementary sequences thereof, and/or portions thereof, to the first plurality of barcoded amplicons. In some embodiments, the method comprises determining the copy number of the nucleic acid target in the cell based on the number of molecular labels with distinct sequences associated with the second plurality of barcoded amplicons, or products thereof. The first plurality of barcoded amplicons and/or the second plurality of barcoded amplicons can comprise whole transcriptome amplification (WTA) products.

In some embodiments, the method comprises synthesizing a third plurality of barcoded amplicons using the plurality of barcoded nucleic acid molecules as templates to generate a third plurality of barcoded amplicons. Synthesizing a third plurality of barcoded amplicons can comprise performing polymerase chain reaction (PCR) amplification of the plurality of the barcoded nucleic acid molecules. Synthesizing a third plurality of barcoded amplicons can comprise PCR amplification using primers capable of hybridizing to the first universal sequence, or a complement thereof, and a target-specific primer. The method can comprise: obtaining sequence information of the third plurality of barcoded amplicons, or products thereof. Obtaining the sequence information can comprise attaching sequencing adaptors to the third plurality of barcoded amplicons, or products thereof. The methods provided herein can comprise determining the copy number of the nucleic acid target in the cell based on the number of molecular labels with distinct sequences associated with the third plurality of barcoded amplicons, or products thereof. The nucleic acid target can comprise a nucleic acid molecule. The nucleic acid molecule

55

can comprise ribonucleic acid (RNA), messenger RNA (mRNA), microRNA, small interfering RNA (siRNA), RNA degradation product, RNA comprising a poly(A) tail, a sample indexing oligonucleotide, a cellular component-binding reagent specific oligonucleotide, or any combination thereof.

In some embodiments, extending the first, second, and/or third pluralities of oligonucleotide barcodes comprising extending the plurality of oligonucleotide barcodes using a reverse transcriptase and/or a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. The DNA polymerase can comprise a Klenow Fragment. The reverse transcriptase can comprise a viral reverse transcriptase, (e.g., a murine leukemia virus (MLV) reverse transcriptase or a Moloney murine leukemia virus (MMLV) reverse transcriptase). The first target-binding region, second target-binding region, and/or the third target-binding region can comprise a poly(dA) region, a poly(dT) region, a random sequence, a gene-specific sequence, a predetermined sequence of any length and composition, or any combination thereof. Obtaining sequence data of the plurality of barcoded DNA fragments, or products thereof, can comprise attaching sequencing adaptors and/or sequencing primers complementary sequences thereof, and/or portions thereof, to the plurality of barcoded DNA fragments, or products thereof.

The method can comprise: amplifying the plurality of barcoded DNA fragments using primers capable of hybridizing to the first universal sequence or complements thereof, and primers capable of hybridizing the second universal sequence or complements thereof, thereby generating a fourth plurality of barcoded DNA fragments. Amplifying the plurality of barcoded DNA fragments can comprise adding sequences of binding sites of sequencing primers and/or sequencing adaptors, complementary sequences thereof, and/or portions thereof, to the plurality of barcoded DNA fragments. Obtaining sequence data of the plurality of barcoded DNA fragments, or products thereof, can comprise obtaining sequence information of the fourth plurality of barcoded amplicons, or products thereof.

Determining the information relating to the gDNA can comprise determining chromatin accessibility of the gDNA based on the sequences of the fourth plurality of barcoded amplicons, or products thereof, in the sequencing data obtained. In some embodiments, determining the chromatin accessibility of the gDNA comprises: aligning the sequences of the fourth plurality of barcoded amplicons (or products thereof) to a reference sequence of the gDNA; and identifying regions of the gDNA corresponding the ends of the barcoded amplicons (or products thereof) of the fourth plurality of barcoded amplicons (or products thereof) to have an accessibility above a threshold. In some embodiments, determining the chromatin accessibility of the gDNA comprises: aligning the sequences of the fourth plurality of barcoded amplicons (or products thereof) to a reference sequence of the gDNA; and determining the accessibility of regions of the gDNA corresponding the ends of barcoded amplicons (or products thereof) of the fourth plurality of barcoded amplicons (or products thereof) based on the numbers of the barcoded amplicons (or products thereof) of the fourth plurality of barcoded amplicons (or products thereof) in the sequencing data.

Determining the information relating to the gDNA can comprise determining methylome information of the gDNA based on the sequences of the fourth plurality of barcoded amplicons (or products thereof) in the sequencing data obtained. The method can comprise: digesting nucleosomes

56

associated with the double-stranded gDNA. The method can comprise: performing chemical conversion and/or enzymatic conversion of cytosine bases of the plurality of dsDNA fragments, or products thereof, to generate a plurality of converted dsDNA fragments with uracil bases, wherein chemical conversion comprises bisulfite treatment, and wherein enzymatic conversion comprises APOBEC-mediated conversion. Barcoding the plurality of dsDNA fragments, or products thereof, can comprise barcoding the plurality of converted dsDNA fragments, or products thereof. In some embodiments, determining the methylome information comprises: determining a position of the plurality of barcoded DNA fragments in the sequencing data has a thymine base and the corresponding position in a reference sequence of the gDNA has a cytosine base to determine the corresponding position in the gDNA has a 5-methylcytosine (5mC) base and/or 5-hydroxymethylcytosine (5hmC) base.

Each oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes can comprise a first universal sequence. The first universal sequence, the second universal sequence, and/or the third universal sequence can be the same. The first universal sequence, the second universal sequence, and/or the third universal sequence can be different. The first universal sequence, the second universal sequence, and/or the third universal sequence can comprise the binding sites of sequencing primers and/or sequencing adaptors, complementary sequences thereof, and/or portions thereof. The sequencing adaptors can comprise a P5 sequence, a P7 sequence, complementary sequences thereof, and/or portions thereof. The sequencing primers can comprise a Read 1 sequencing primer, a Read 2 sequencing primer, complementary sequences thereof, and/or portions thereof. The first, second, and/or third pluralities of oligonucleotide barcodes each can comprise a molecular label. At least 10 of the first, second, and/or third pluralities of oligonucleotide barcodes can comprise different molecular label sequences. Each molecular label of the first, second, and/or third pluralities of oligonucleotide barcodes can comprise at least 6 nucleotides. The first, second, and/or third pluralities of oligonucleotide barcodes can be associated with a solid support. The first, second, and/or third pluralities of oligonucleotide barcodes associated with the same solid support each can comprise an identical sample label. Each sample label of the first, second, and/or third pluralities of oligonucleotide barcodes can comprise at least 6 nucleotides. The first, second, and/or third pluralities of oligonucleotide barcodes each can comprise a cell label. Each cell label of the first, second, and/or third pluralities of oligonucleotide barcodes can comprise at least 6 nucleotides. Oligonucleotide barcodes of the first, second, and/or third pluralities of oligonucleotide barcodes associated with the same solid support can comprise the same cell label. Oligonucleotide barcodes of the first, second, and/or third pluralities of oligonucleotide barcodes associated with different solid supports can comprise different cell labels.

The solid support can comprise a synthetic particle, a planar surface, or a combination thereof. The method can comprise: associating a synthetic particle comprising the first, second, and/or third pluralities of oligonucleotide barcodes with the cell. The method can comprise: lysing the cell after associating the synthetic particle with the cell. Lysing the cell can comprise heating the cell, contacting the cell with a detergent, changing the pH of the cell, or any

combination thereof. The synthetic particle and the single cell can be in the same partition. The partition can be a well or a droplet.

At least one oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes can be immobilized or partially immobilized on the synthetic particle, or at least one oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes can be enclosed or partially enclosed in the synthetic particle. The synthetic particle can be disruptable (e.g., a disruptable hydrogel particle). The synthetic particle can comprise a bead. The bead can comprise a Sepharose bead, a streptavidin bead, an agarose bead, a magnetic bead, a conjugated bead, a protein A conjugated bead, a protein G conjugated bead, a protein A/G conjugated bead, a protein L conjugated bead, an oligo(dT) conjugated bead, a silica bead, a silica-like bead, an anti-biotin microbead, an anti-fluorochrome microbead, or any combination thereof. The synthetic particle can comprise a material selected from the group consisting of polydimethylsiloxane (PDMS), polystyrene, glass, polypropylene, agarose, gelatin, hydrogel, paramagnetic, ceramic, plastic, glass, methylstyrene, acrylic polymer, titanium, latex, Sepharose, cellulose, nylon, silicone, and any combination thereof. Each oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes can comprise a linker functional group. The synthetic particle can comprise a solid support functional group. The support functional group and the linker functional group can be associated with each other. The linker functional group and the support functional group can be individually selected from the group consisting of C6, biotin, streptavidin, primary amine(s), aldehyde(s), ketone(s), and any combination thereof.

#### Kits

Disclosed herein include kits. In some embodiments, the kit comprises: a transposome comprising a transposase, a first adaptor having a first 5' overhang, and a second adaptor having a second 5' overhang; and a first plurality of oligonucleotide barcodes, wherein the 3' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes is associated with a solid support, wherein the 5' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes comprises a first target-binding region capable of hybridizing to the first 5' overhang.

Disclosed herein include kits. In some embodiments, the kit comprises: a transposome comprising a transposase, a first adaptor having a first 5' overhang, and a second adaptor having a second 5' overhang, wherein the first 5' overhang comprises a coupling sequence; and a coupling oligonucleotide comprising a 5' complement of the coupling sequence and a 3' complement of a second target-binding region.

The kit can comprise: a second plurality of oligonucleotide barcodes, wherein each oligonucleotide barcode of the second plurality of oligonucleotide barcodes comprises the second target-binding region. The coupling oligonucleotide can be a single-stranded oligonucleotide. The coupling oligonucleotide can comprise at least 6 nucleotides. The coupling sequence can comprise at least 4 nucleotides. The first 5' overhang and/or the second 5' overhang can comprise at least 4 nucleotides. The second 5' overhang can comprise a second universal sequence. The first adaptor and/or the second adaptor can comprise a DNA end sequence of the transposon. The kit can comprise: a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity. The DNA polymerase can comprise a Klenow Fragment. The kit can comprise: a reverse transcriptase. The reverse transcriptase can comprise a viral

reverse transcriptase (e.g., a murine leukemia virus (MLV) reverse transcriptase or a Moloney murine leukemia virus (MMLV) reverse transcriptase). The kit can comprise: a ligase. The kit can comprise: a detergent and/or a surfactant. The kit can comprise: a buffer, a cartridge, or both. The kit can comprise: one or more reagents for a reverse transcription reaction and/or an amplification reaction. The transposase can comprise a Tn5 transposase.

The kit can comprise: a third plurality of oligonucleotide barcodes, wherein each oligonucleotide barcode of the third plurality of oligonucleotide barcodes can comprise a third target-binding region. The first target-binding region, second target-binding region, and/or the third target-binding region can comprise a poly(dA) region, a poly(dT) region, a random sequence, a gene-specific sequence, or any combination thereof. The first, second, and/or third pluralities of oligonucleotide barcodes each can comprise a molecular label. The molecular label can comprise at least 6 nucleotides. At least 10 of the first, second, and/or third pluralities of oligonucleotide barcodes can comprise different molecular label sequences.

The first, second, and/or third pluralities of oligonucleotide barcodes can be associated with a solid support. The first, second, and/or third pluralities of oligonucleotide barcodes each can comprise a cell label. Oligonucleotide barcodes of the first, second, and/or third pluralities of oligonucleotide barcodes associated with the same solid support can comprise the same cell label. Oligonucleotide barcodes of the first, second, and/or third pluralities of oligonucleotide barcodes associated with different solid supports can comprise different cell labels. The solid support can comprise a synthetic particle, a planar surface, or a combination thereof.

In some embodiments, at least one oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes is immobilized or partially immobilized on the synthetic particle, or at least one oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes is enclosed or partially enclosed in the synthetic particle. The synthetic particle can be disruptible (e.g., a disruptable hydrogel particle). The synthetic particle can comprise a bead. The bead can comprise a Sepharose bead, a streptavidin bead, an agarose bead, a magnetic bead, a conjugated bead, a protein A conjugated bead, a protein G conjugated bead, a protein A/G conjugated bead, a protein L conjugated bead, an oligo(dT) conjugated bead, a silica bead, a silica-like bead, an anti-biotin microbead, an anti-fluorochrome microbead, or any combination thereof. The synthetic particle can comprise a material selected from the group consisting of polydimethylsiloxane (PDMS), polystyrene, glass, polypropylene, agarose, gelatin, hydrogel, paramagnetic, ceramic, plastic, glass, methylstyrene, acrylic polymer, titanium, latex, Sepharose, cellulose, nylon, silicone, and any combination thereof. Each oligonucleotide barcode of the first, second, and/or third pluralities of oligonucleotide barcodes can comprise a linker functional group. The synthetic particle can comprise a solid support functional group. The support functional group and the linker functional group can be associated with each other. The linker functional group and the support functional group can be individually selected from the group consisting of C6, biotin, streptavidin, primary amine(s), aldehyde(s), ketone(s), and any combination thereof.

#### Terminology

In at least some of the previously described embodiments, one or more elements used in an embodiment can interchangeably be used in another embodiment unless such a replacement is not technically feasible. It will be appreciated

by those skilled in the art that various other omissions, additions and modifications may be made to the methods and structures described above without departing from the scope of the claimed subject matter. All such modifications and changes are intended to fall within the scope of the subject matter, as defined by the appended claims.

One skilled in the art will appreciate that, for this and other processes and methods disclosed herein, the functions performed in the processes and methods can be implemented in differing order. Furthermore, the outlined steps and operations are only provided as examples, and some of the steps and operations can be optional, combined into fewer steps and operations, or expanded into additional steps and operations without detracting from the essence of the disclosed embodiments.

With respect to the use of substantially any plural and/or singular terms herein, those having skill in the art can translate from the plural to the singular and/or from the singular to the plural as is appropriate to the context and/or application. The various singular/plural permutations may be expressly set forth herein for sake of clarity. As used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural references unless the context clearly dictates otherwise. Any reference to "or" herein is intended to encompass "and/or" unless otherwise stated.

It will be understood by those within the art that, in general, terms used herein, and especially in the appended claims (e.g., bodies of the appended claims) are generally intended as "open" terms (e.g., the term "including" should be interpreted as "including but not limited to," the term "having" should be interpreted as "having at least," the term "includes" should be interpreted as "includes but is not limited to," etc.). It will be further understood by those within the art that if a specific number of an introduced claim recitation is intended, such an intent will be explicitly recited in the claim, and in the absence of such recitation no such intent is present. For example, as an aid to understanding, the following appended claims may contain usage of the introductory phrases "at least one" and "one or more" to introduce claim recitations. However, the use of such phrases should not be construed to imply that the introduction of a claim recitation by the indefinite articles "a" or "an" limits any particular claim containing such introduced claim recitation to embodiments containing only one such recitation, even when the same claim includes the introductory phrases "one or more" or "at least one" and indefinite articles such as "a" or "an" (e.g., "a" and/or "an" should be interpreted to mean "at least one" or "one or more"); the same holds true for the use of definite articles used to introduce claim recitations. In addition, even if a specific number of an introduced claim recitation is explicitly recited, those skilled in the art will recognize that such recitation should be interpreted to mean at least the recited number (e.g., the bare recitation of "two recitations," without other modifiers, means at least two recitations, or two or more recitations). Furthermore, in those instances where a convention analogous to "at least one of A, B, and C, etc." is used, in general such a construction is intended in the sense one having skill in the art would understand the convention (e.g., "a system having at least one of A, B, and C" would include but not be limited to systems that have A alone, B alone, C alone, A and B together, A and C together, B and C together, and/or A, B, and C together, etc.). In those instances where a convention analogous to "at least one of A, B, or C, etc." is used, in general such a construction is intended in the sense one having skill in the art would

understand the convention (e.g., "a system having at least one of A, B, or C" would include but not be limited to systems that have A alone, B alone, C alone, A and B together, A and C together, B and C together, and/or A, B, and C together, etc.). It will be further understood by those within the art that virtually any disjunctive word and/or phrase presenting two or more alternative terms, whether in the description, claims, or drawings, should be understood to contemplate the possibilities of including one of the terms, either of the terms, or both terms. For example, the phrase "A or B" will be understood to include the possibilities of "A" or "B" or "A and B."

In addition, where features or aspects of the disclosure are described in terms of Markush groups, those skilled in the art will recognize that the disclosure is also thereby described in terms of any individual member or subgroup of members of the Markush group.

As will be understood by one skilled in the art, for any and all purposes, such as in terms of providing a written description, all ranges disclosed herein also encompass any and all possible sub-ranges and combinations of sub-ranges thereof. Any listed range can be easily recognized as sufficiently describing and enabling the same range being broken down into at least equal halves, thirds, quarters, fifths, tenths, etc. As a non-limiting example, each range discussed herein can be readily broken down into a lower third, middle third and upper third, etc. As will also be understood by one skilled in the art all language such as "up to," "at least," "greater than," "less than," and the like include the number recited and refer to ranges which can be subsequently broken down into sub-ranges as discussed above. Finally, as will be understood by one skilled in the art, a range includes each individual member. Thus, for example, a group having 1-3 articles refers to groups having 1, 2, or 3 articles. Similarly, a group having 1-5 articles refers to groups having 1, 2, 3, 4, or 5 articles, and so forth.

From the foregoing, it will be appreciated that various embodiments of the present disclosure have been described herein for purposes of illustration, and that various modifications may be made without departing from the scope and spirit of the present disclosure. Accordingly, the various embodiments disclosed herein are not intended to be limiting, with the true scope and spirit being indicated by the following claims.

What is claimed is:

1. A method of labeling DNA, the method comprising: contacting double-stranded deoxyribonucleic acid (dsDNA) with a transposome to generate a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang, wherein each dsDNA fragment of the plurality of dsDNA fragments comprises a first strand comprising the first 5' overhang and a second strand comprising the second 5' overhang, wherein the transposome comprises a transposase, a first adaptor having the first 5' overhang, and a second adaptor having the second 5' overhang; barcoding the plurality of dsDNA fragments, or products thereof, using a first plurality of oligonucleotide barcodes to generate a plurality of barcoded DNA fragments, wherein the 3' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes is associated with a solid support, and wherein the 5' end of each oligonucleotide barcode of the first plurality of oligonucleotide barcodes comprises a first target-binding region capable of hybridizing to the first 5' overhang of at least one of the plurality of dsDNA fragments.

**61**

2. The method of claim 1, wherein barcoding the plurality of dsDNA fragments, or products thereof, using a first plurality of oligonucleotide barcodes comprises:

hybridizing the first 5' overhang of the first strand of a dsDNA fragment with the first target-binding region of an oligonucleotide barcode of the first plurality of oligonucleotide barcodes; and  
ligating the second strand of said dsDNA fragment to said hybridized oligonucleotide barcode using a DNA ligase.

3. The method of claim 2, comprising, before ligating the second strand to the oligonucleotide barcode, filling a gap between the second strand and said hybridized oligonucleotide barcode with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity.

4. The method of claim 1, wherein the first 5' overhang comprises a complement of a first target-binding region.

5. The method of claim 1, wherein the plurality of dsDNA fragments comprise one or more gaps, the method comprising filling the one or more gaps with a DNA polymerase lacking at least one of 5' to 3' exonuclease activity and 3' to 5' exonuclease activity.

6. The method of claim 1, wherein: (i) the first 5' overhang and/or the second 5' overhang comprises at least 4 nucleotides; (ii) the second 5' overhang comprises a second universal sequence; and/or (iii) the first adaptor and/or the second adaptor comprises a DNA end sequence of the transposon.

7. The method of claim 1, wherein contacting dsDNA with a transposome comprises permeabilizing a cell and/or isolating the nucleus of a cell, wherein the cell comprises the dsDNA, and wherein the cell comprises a single cell.

8. The method of claim 7, comprising:

permeabilizing a nucleus in the cell to generate a permeabilized nucleus; and/or  
fixating the cell comprising the nucleus prior to permeabilizing the nucleus.

9. The method of claim 1, wherein the dsDNA comprises genomic DNA (gDNA), and wherein the plurality of dsDNA fragments comprise open chromatin-associated gDNA.

10. The method of claim 1, comprising:

obtaining sequence data of the plurality of barcoded DNA fragments, or products thereof; and  
determining information relating to the gDNA based on the sequences of the plurality of barcoded DNA fragments, or products thereof, in the sequencing data obtained.

11. The method of claim 10, wherein determining the information relating to the gDNA comprises determining chromatin accessibility of the gDNA based on the sequences of the plurality of barcoded DNA fragments, or products thereof, in the sequencing data obtained.

12. The method of claim 11, wherein determining the chromatin accessibility of the gDNA comprises:

aligning the sequences of the plurality of barcoded DNA fragments to a reference sequence of the gDNA; and identifying regions of the gDNA corresponding the ends of the barcoded DNA fragments of the plurality of barcoded DNA fragments to have an accessibility above a user-defined threshold.

13. The method of claim 11, wherein determining the chromatin accessibility of the gDNA comprises:

aligning the sequences of the plurality of barcoded DNA fragments to a reference sequence of the gDNA; and determining the accessibility of regions of the gDNA corresponding the ends of barcoded DNA fragments of

**62**

the plurality of barcoded DNA fragments based on the numbers of the barcoded DNA fragments of the plurality of barcoded DNA fragments in the sequencing data.

14. The method of claim 10, wherein determining the information relating to the gDNA comprises determining methylome information of the gDNA based on the sequences of the plurality of barcoded DNA fragments in the sequencing data obtained, further comprising:

digesting nucleosomes associated with the double-stranded gDNA.

15. The method of claim 14, wherein determining the methylome information comprises:

determining a position of the plurality of barcoded DNA fragments in the sequencing data has a thymine base and the corresponding position in a reference sequence of the gDNA has a cytosine base to determine the corresponding position in the gDNA has a 5-methylcytosine (5mC) base and/or 5-hydroxymethylcytosine (5hmC) base.

16. The method of claim 14, comprising performing chemical conversion and/or enzymatic conversion of cytosine bases of the plurality of dsDNA fragments, or products thereof, to generate a plurality of converted dsDNA fragments with uracil bases, wherein chemical conversion comprises bisulfite treatment, wherein enzymatic conversion comprises APOBEC-mediated conversion, and wherein barcoding the plurality of dsDNA fragments, or products thereof, comprises barcoding the plurality of converted dsDNA fragments, or products thereof.

17. A method of labeling DNA, the method comprising: contacting double-stranded deoxyribonucleic acid (dsDNA) with a transposome to generate a plurality of dsDNA fragments each comprising a first 5' overhang and a second 5' overhang, wherein each dsDNA fragment of the plurality of dsDNA fragments comprises a first strand comprising the first 5' overhang and a second strand comprising the second 5' overhang, wherein the transposome comprises a transposase, a first adaptor having the first 5' overhang, and a second adaptor having the second 5' overhang, and wherein the first 5' overhang comprises a coupling sequence; providing a coupling oligonucleotide comprising a 5' complement of the coupling sequence and a 3' complement of a second target-binding region; and barcoding the plurality of dsDNA fragments, or products thereof, using a second plurality of oligonucleotide barcodes to generate a plurality of barcoded DNA fragments, wherein each oligonucleotide barcode of the second plurality of oligonucleotide barcodes comprises the second target-binding region.

18. The method of claim 17, wherein barcoding the plurality of dsDNA fragments, or products thereof, using a second plurality of oligonucleotide barcodes comprises:

hybridizing the coupling sequence of the first strand of a dsDNA fragment with the 5' complement of the coupling sequence of the coupling oligonucleotide;

hybridizing the 3' complement of the second target-binding region of the coupling oligonucleotide with the second target-binding region of an oligonucleotide barcode of the second plurality of oligonucleotide barcodes; and

ligating the first strand of said dsDNA fragment to said hybridized oligonucleotide barcode and extending the first strand; and/or

**63**

ligating the second strand of said dsDNA fragment to  
said coupling oligonucleotide and extending the sec-  
ond strand.

- 19.** The method of claim **18**, comprising,  
before ligating the first strand to the oligonucleotide <sup>5</sup>  
barcode, filling a gap between the first strand and the  
hybridized oligonucleotide barcode with a DNA poly-  
merase lacking at least one of 5' to 3' exonuclease  
activity and 3' to 5' exonuclease activity; and/or  
before ligating the second strand to the coupling oligo-  
nucleotide, filling a gap between the second strand and  
the coupling oligonucleotide with a DNA polymerase  
lacking at least one of 5' to 3' exonuclease activity and  
3' to 5' exonuclease activity. <sup>10</sup>
- 20.** The method of claim **17**, wherein the coupling oligo-  
nucleotide is a single-stranded oligonucleotide, wherein the  
coupling oligonucleotide comprises at least 6 nucleotides,  
and wherein the coupling sequence comprises at least 4  
nucleotides. <sup>15</sup>

**64**

- 21.** A kit comprising:  
(i) a transposome comprising a transposase, a first adaptor  
having a first 5' overhang, and a second adaptor having  
a second 5' overhang; and  
a first plurality of oligonucleotide barcodes, wherein  
the 3' end of each oligonucleotide barcode of the first  
plurality of oligonucleotide barcodes is associated  
with a solid support, wherein the 5' end of each  
oligonucleotide barcode of the first plurality of oli-  
gonucleotide barcodes comprises a first target-bind-  
ing region capable of hybridizing to the first 5'  
overhang; or  
(ii) a transposome comprising a transposase, a first adap-  
tor having a first 5' overhang, and a second adaptor  
having a second 5' overhang, wherein the first 5'  
overhang comprises a coupling sequence; and  
a coupling oligonucleotide comprising a 5' complement  
of the coupling sequence and a 3' complement of a  
second target-binding region.

\* \* \* \* \*