



US 20250255824A1

(19) United States

(12) Patent Application Publication

RIESEL et al.

(10) Pub. No.: US 2025/0255824 A1

(43) Pub. Date: Aug. 14, 2025

## (54) PHARMACEUTICAL COMPOSITIONS

(71) Applicant: CALLIDITAS THERAPEUTICS AB,  
Stockholm (SE)(72) Inventors: Eva Kristina RIESEL, Mölndal (SE);  
Lena Margareta PERESWETOFF-MORATH, Spånga  
(SE); Kari SANDVOLD, Uppsala (SE);  
Christian Olle Andreas PEDERSEN,  
Uppsala (SE)

(21) Appl. No.: 19/195,082

(22) Filed: Apr. 30, 2025

## Related U.S. Application Data

- (63) Continuation of application No. 18/934,978, filed on Nov. 1, 2024, now Pat. No. 12,311,057, which is a continuation of application No. 18/392,602, filed on Dec. 21, 2023, now Pat. No. 12,171,882, which is a continuation of application No. 18/100,396, filed on Jan. 23, 2023, now Pat. No. 11,896,719.  
(60) Provisional application No. 63/302,226, filed on Jan. 24, 2022, provisional application No. 63/302,216, filed on Jan. 24, 2022.

## (30) Foreign Application Priority Data

Nov. 16, 2022 (GB) ..... 2217146.6  
Nov. 16, 2022 (GB) ..... 2217150.8

## Publication Classification

## (51) Int. Cl.

A6IK 9/48 (2006.01)  
A6IK 9/28 (2006.01)

A6IK 9/50 (2006.01)

A6IK 31/58 (2006.01)


A6IP 13/12 (2006.01)

(52) U.S. Cl.  
CPC ..... A6IK 9/4891 (2013.01); A6IK 9/2846  
(2013.01); A6IK 9/5047 (2013.01); A6IK  
31/58 (2013.01); A6IP 13/12 (2018.01)

## (57) ABSTRACT

The present invention provides for a method of treatment of IgA nephropathy, which method comprises:

- (i) identifying a pharmaceutically acceptable composition intended to treat IgA nephropathy comprising budesonide and one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, which composition fulfils the following requirements in a standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test using a dissolution apparatus according to Apparatus 2 (Paddle Apparatus) of said test;
- (a) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 120 minutes, when the dissolution medium is aqueous and has a pH of about 1.2;
- (b) the composition fulfils the requirement that no more than about 10% of the budesonide is released into a pharmaceutically-relevant dissolution medium within about 30 minutes; and
- (c) the composition fulfils the requirement that at least about 70% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 120 minutes;
- (ii) wherein the method comprises the step of administering said composition to a patient with IgA nephropathy in need of said treatment.






Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6






Figure 7



**Figure 8**



**Figure 9**




Figure 10




Figure 11




Figure 12





Figure 13



**Figure 14**




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21






Figure 22



**Figure 23A**



**Figure 23B**



Figure 24

## PHARMACEUTICAL COMPOSITIONS

[0001] This application is a continuation of U.S. patent application Ser. No. 18/934,978, filed Nov. 1, 2024, which is a continuation of U.S. patent application Ser. No. 18/392,602, filed Dec. 21, 2023, now U.S. Pat. No. 12,171,882 issued Dec. 24, 2024, which is a continuation of U.S. patent application Ser. No. 18/100,396, filed Jan. 23, 2023, now U.S. Pat. No. 11,896,719 issued Feb. 13, 2024, which claims the priority benefit of U.S. Provisional Patent Application Ser. Nos. 63/302,226 and 63/302,216, both filed on Jan. 24, 2022, which are hereby incorporated by reference in their entirety. The application also claims the priority benefit of GB Application Nos. 2217150.8 and 2217146.6, both filed on Nov. 16, 2022.

### FIELD OF THE INVENTION

[0002] The invention relates to a method of treating IgA nephropathy and a method of determining whether a pharmaceutical composition is capable of safely and efficaciously treating IgA nephropathy. The invention also relates to compositions for use in treating IgA nephropathy and methods for producing those compositions.

### PRIOR ART AND BACKGROUND

[0003] The listing or discussion of an apparently prior-published document in this specification should not necessarily be taken as an acknowledgement that the document is part of the state of the art or common general knowledge.

[0004] IgA nephropathy (IgAN), sometimes referred to as Berger's disease, is a serious progressive autoimmune disease of the kidney in which up to 50% of patients end up at risk of developing end-stage renal disease (ESRD) within ten to twenty years.

[0005] IgAN is an orphan disease and it is estimated that approximately 130,000 to 150,000 people are affected by the disease in the United States and approximately 200,000 people are affected in Europe. A significantly higher prevalence has been observed in Asia, including in Greater China, where IgAN has historically been a leading cause of ESRD. It is estimated that IgAN affects approximately two million people in Greater China.

[0006] Although IgAN manifests in the kidney, most scientific studies have found that the pathogenesis of IgAN begins in the ileum, which is the final part of the small intestine before the large intestine. Masses of lymphatic tissue, known as Peyer's patches, are predominantly found in the ileum where they produce secretory IgA antibodies. IgA antibodies play a key role in the immune system by protecting the body from foreign substances, such as food-derived factors, bacteria and viruses.

[0007] Patients with IgAN have elevated levels of a subclass of IgA antibodies produced in the gut that lack units of galactose at their hinge region. The hinge region is a flexible amino acid stretch in the central part of the heavy chains of the IgA antibody. In IgAN patients, a combination of genetic predisposition and environmental, bacterial or dietary factors are presumed to lead to an increased production of these galactose-deficient IgA antibodies, potentially in combination with increased intestinal permeability, leading to these antibodies appearing in the blood. The galactose-deficient IgA antibodies (also referred to herein as poorly O-galactosylated IgA1) are immunogenic when found in the circulation, which triggers autoantibodies, or antibodies created by

the body in response to a constituent of its own tissue. This in turn leads to the formation of pathogenic immune complexes, or clusters of antibodies, which deposit in the membranes of the glomeruli, the kidney's filtration apparatus. These trapped immune complexes initiate an inflammatory cascade that damages the membranes, resulting in protein and blood leaking into the urine. Ultimately the glomeruli are destroyed, reducing the kidney's ability to remove waste products from the blood. As the disease progresses, waste products that are normally removed from the blood accumulate, resulting in potentially life-threatening complications that in many patients will lead to the need for dialysis or kidney transplant.

[0008] The standard of care for ESRD is dialysis or kidney transplant, which represents a significant health economic burden as well as a material impact on patients' quality of life.

[0009] Despite a need for new therapies, there have been few new drugs developed for chronic kidney diseases during the last decade and, until recently, there has been no approved therapy for the direct treatment of IgAN per se. Patients with IgAN are typically initially given antihypertensive medications. This treatment regimen initially attempts to manage the symptoms of IgAN by decreasing blood pressure and reducing proteinuria but has not been proven to address the underlying cause of IgAN. Over time, physicians attempt to control disease progression with a variety of off-label treatments, such as statins, omega-3-acids and diuretics, but a significant proportion of patients experience continued deterioration of kidney function, and until recently no approved treatment options have been available.

[0010] For IgAN patients whose disease has progressed, clinicians may treat patients with systemic immunosuppressive agents, primarily consisting of high doses of systemic corticosteroids, such as prednisone, prednisolone and methylprednisolone. While some published reports indicate that these agents may reduce proteinuria, this high dosing of systemic corticosteroids is also associated with a wide range of adverse events, including high blood pressure, weight gain, diabetes, serious infections and osteoporosis. Also, any potential impact on the underlying disease in terms of kidney function, as measured by estimated Glomerular Filtration Rate (eGFR), has yet to be proven.

[0011] Thus, in attempting to meet the present clinical need for an effective treatment of IgAN, there is a clear need for new and/or improved treatments for IgAN in which an effective local treatment with an immunosuppressive agent without such undesired side effects is obtained.

[0012] Peyer's patches (aggregated lymphoid nodules) are small masses of lymphatic tissue found throughout the ileum region of the small intestine. They are an important part of the immune system as they monitor intestinal bacteria populations and prevent the growth of pathogenic bacteria in the intestine.

[0013] As Peyer's patches are responsible for the synthesis of the bulk of IgA in the body, a targeted dose of locally-acting immunosuppressive agent to the ileum (and particularly the terminal/distal ileum), where Peyer's patches are predominantly found, may serve to reduce the formation of the IgA molecules that ultimately drive immune complex formation in IgAN, by reducing the formation of secretory galactose-deficient IgA antibodies and their appearance in the blood. Such targeted release will also likely limit sys-

temic exposure of locally-acting immunosuppressants, such as certain corticosteroids, in order to avoid undesirable side-effects.

[0014] Peyer's Patches are sites of intense activation of B cells in the human body. Therefore, it stands to reason that monitoring survival factors related to B-cell activation would provide an indication as to the efficacy of treatments with locally-acting immunosuppressive agents.

[0015] The tumour necrosis factor (TNF) family members, B cell activating factor (BAFF) and its homolog a proliferation-inducing ligand (APRIL) are crucial survival factors for peripheral B-cells, and are expressed by cells including monocytes, dendritic cells, neutrophils, basophils, stromal cells, activated T-cells, cells of bowel mucosa, activated and malignant B-cells, and epithelial cells (Mackay and Schneider, 2009. *Nat. Rev. Immunol.*, 9:491-502; Schneider et al, 1999. *J. Exp. Med.*, 189:1747-1756; Yu et al, 2000. *Nat. Immunol.*, 1:252-256).

[0016] BAFF is a ligand for the receptors Transmembrane activator and CAML interactor (TACI), (also known as tumour necrosis factor receptor superfamily member 13B (TNFRSF13B)); B-cell maturation antigen (BCMA) (also known as tumour necrosis factor receptor superfamily member 17 (TNFRSF17)); and B-cell activating factor receptor (BAFF-R) (also known as tumour necrosis factor receptor superfamily member 13C (TNFRSF13C)). BAFF-R is specific to BAFF, whereas TACI and BCMA also bind to APRIL (Mackay and Schneider, 2009. *Nat. Rev. Immunol.*, 9:491-502).

[0017] BAFF is a potent B cell activator and is crucial for B-cell homeostasis and for the regulation of B-cell selection. An excess of BAFF has been shown to be associated with the development of autoimmune disorders, such as IgAN, in animal models, and high levels of BAFF have been detected in the serum of patients with various autoimmune conditions. Increased levels of BAFF have been associated with an upregulation of humoral immunity through increased levels of B-cells and immunoglobulins (Steri et al, 2017. *N. Engl. J. Med.*, 376:1615-1626).

[0018] Increased serum levels of BAFF and APRIL are found in patients suffering from IgAN and this has led to the development of drugs that seek to inhibit those molecules. The focus of those drugs has been blocking the interaction between BAFF and/or APRIL and their receptors. For example, but not limited to, the BAFF inhibitor Blisibimod (Anthera Pharmaceuticals, discontinued) is a fusion protein consisting of four BAFF binding domains fused to the N-terminus of the Fc region of a human antibody, which binds to BAFF and inhibits interaction with BAFF receptors. Similarly, the combined BAFF/APRIL antagonist Atacicept (Merck Serono, licensed by Vera Therapeutics) is also a recombinant fusion protein that combines BAFF and APRIL binding domains with an antibody Fc region, and blocks interaction with TACI. The APRIL antagonist VIS649 (Visterra, a subsidiary of Otsuka) and the BAFF inhibitor Belimumab (GlaxoSmithKline) are monoclonal antibodies that bind directly to APRIL and BAFF respectively and block interaction with their receptors. Thus, drugs targeting BAFF and APRIL seek to block the activity of endogenous BAFF/APRIL molecules in order to reduce activation and proliferation of B-cells, and the associated immunological effects.

[0019] The current understanding of the pathogenesis and current treatments of IgAN are summarised in J. Barratt et

al., Treatment of IgA Nephropathy: Evolution Over Half a Century, *Seminars in Nephrology*, 2018, 38 (5), 531-540, see also Boyd et al., Kidney International, 2012, 81, 833-843. The current understanding of the pathogenesis of IgAN is also outlined in Seikrit et al., The Immune Landscape of IgA Induction in the Gut, *Seminars in Immunopathology*, 2021, 43, 627-637.

[0020] Surprisingly, we found that oral administration of a budesonide formulation with a distinct in vitro release profile leads to a pronounced decrease in the serum level of BAFF in those subjects relative to the level observed prior to administration of budesonide. Furthermore, the observed decrease in serum BAFF level may occur concomitantly with decreases in the levels of biomarkers associated with B-cell activation and proliferation. Accordingly, the in vitro release profile is indicative of successful targeted release in the intestine (i.e., successful targeted release to the distal ileum) of subjects.

[0021] In addition, what is particularly interesting is that it has been shown that treatment of IgAN with systemic glucocorticoids lowers both total serum IgA and poorly O-galactosylated IgA1 (Kosztyu P et al.: Glucocorticoids Reduce Aberrant O-Glycosylation of IgA1 in IgA Nephropathy Patients. *Kidney Blood Press Res* 2018; 43:350-359. However, with the present treatment of oral administration of a budesonide formulation as defined herein there were no differences observed in levels of total level of functional IgA antibodies, including IgA1 and IgG with budesonide capsule treatment, but serum levels of galactose-deficient IgA (poorly O-galactosylated IgA) did drop. This finding led to the conclusion that the effect of local ileal treatment with budesonide capsules was selective for the pathogenic antibodies but not effective on the general pool of IgA, IgA1 and IgG.

[0022] These results show that treatment with the budesonide formulation as defined herein is supportive of a direct effect on the underlying pathogenic pathways in IgAN and that the budesonide payload has a predominantly topical effect rather than a systemic effect, leading to reduced side effects for patients when treated with Nefcon budesonide.

[0023] To support this, in silico modelling of the budesonide formulation with the distinct in vitro release profile shows that the payload is released predominantly to the ileum, in particular the distal ileum. With budesonide having a high first pass rate, primarily through gut wall metabolism in the small intestine, (Seidegård J et. al., *Presystemic elimination of budesonide in man when administered locally at different levels in the gut, with and without local inhibition by ketoconazole*. Eur J Pharm Sci. 2008 Nov. 15; 35 (4): 264-70; Raje et al. *Evaluation of separate role of intestine and liver in first pass metabolism of budesonide in rat Xenobiotica*. 2018 December; 48 (12): 1206-1214)), these results are further indicative of the budesonide formulation having a topical rather than systemic effect.

[0024] Taken together, the results described herein show that budesonide formulations exhibiting the distinct in vitro release profile as defined herein are an effective treatment of IgAN. Due to their targeted local release and effect of the topical corticosteroid, a lower level of undesired side effects is obtained.

[0025] Formulations of the corticosteroid budesonide have previously been described in international patent application WO 2009/138716 A1.

## BRIEF DESCRIPTION OF THE FIGURES

- [0026] FIG. 1 is a flow diagram detailing the preparation of cured polymer coated beads.
- [0027] FIG. 2 shows the in vitro dissolution profiles of budesonide modified release capsules in the presence of added surfactant in the pH 6.8 buffer medium at a paddle rotation speed of 100 rpm; data shown for different capsule batches prepared according to the invention.
- [0028] FIG. 3 is a repeat of the in vitro dissolution profile shown in FIG. 2 focusing only on the dissolution at pH 6.8.
- [0029] FIG. 4: relatively shows the in vitro dissolution profile for seven separate batches of budesonide modified release capsules according to the present invention with the dissolution in the buffered pH 6.8 medium being in the absence of surfactant and with a paddle rotation speed of 50 rpm.
- [0030] FIG. 5: shows the in vitro dissolution profiles of budesonide modified release capsules according to the present invention along with three other marketed budesonide-containing formulations in the pH 6.8 buffer medium in the presence of added surfactant and at a paddle rotation speed of 100 rpm.
- [0031] FIG. 6: shows the in vitro dissolution profiles of budesonide modified release capsules according to the present invention and three other marketed budesonide-containing formulations in the pH 6.8 buffer medium in the absence of added surfactant and at a paddle rotation speed of 100 rpm.
- [0032] FIG. 7: Percentage change in BAFF level relative to baseline level following treatment. Percentage changes in BAFF relative to baseline levels were measured in patients following 9 months of treatment with: (a) placebo; (b) Nefecon-budesonide (8 mg/day); and (c) Nefecon-budesonide (16 mg/day). The dotted line indicates no percentage change following intervention with placebo or Nefecon-budesonide.
- [0033] FIG. 8: Percentage change in BAFF level relative to end of treatment level after the follow-up phase. Percentage changes in BAFF were measured in patients given Nefecon-budesonide (16 mg/day) for 9 months at: (a) month 9 at the end of the treatment phase compared to baseline; and (b) at month 12 after the follow-up phase compared to end of treatment. The dotted line indicates no percentage change following intervention with placebo or Nefecon-budesonide.
- [0034] FIG. 9: Percentage change in APRIL level relative to baseline level following treatment. Percentage changes in APRIL levels relative to baseline levels were measured in patients following 9 months of treatment with: (a) placebo; (b) Nefecon-budesonide (8 mg/day); and (c) Nefecon-budesonide (16 mg/day). The dotted line indicates no percentage change following intervention with placebo or Nefecon-budesonide.
- [0035] FIG. 10: Percentage change in TACI level relative to baseline level following treatment. Percentage changes in TACI levels relative to baseline levels were measured in patients following 9 months of treatment with: (a) placebo; (b) Nefecon-budesonide (8 mg/day); and (c) Nefecon-budesonide (16 mg/day). The dotted line indicates no percentage change following intervention with placebo or Nefecon-budesonide.
- [0036] FIG. 11: Percentage change in BCMA level relative to baseline level following treatment. Percentage changes in BCMA levels relative to baseline levels were measured in patients following 9 months of treatment with: (a) placebo; (b) Nefecon-budesonide (8 mg/day); and (c) Nefecon-budesonide (16 mg/day). The dotted line indicates no percentage change following intervention with placebo or Nefecon-budesonide.
- [0037] FIG. 12: Percentage change in CD27 level relative to baseline level following treatment. Percentage changes in CD27 levels relative to baseline levels were measured in patients following 9 months of treatment with: (a) placebo; (b) Nefecon-budesonide (8 mg/day); and (c) Nefecon-budesonide (16 mg/day). The dotted line indicates no percentage change following intervention with placebo or Nefecon-budesonide.
- [0038] FIG. 13: Percentage change in CD27 level relative to end of treatment level after the follow-up phase. Percentage changes in CD27 were measured in patients given Nefecon-budesonide (16 mg/day) for 9 months at: (a) month 9 at the end of the treatment phase compared to baseline; and (b) at month 12 after the follow-up phase compared to end of treatment. The dotted line indicates no % change following intervention with placebo or Nefecon-budesonide.
- [0039] FIG. 14: Percentage change in CD30 level relative to baseline level following treatment. Percentage changes in CD30 levels relative to baseline levels were measured in patients following 9 months of treatment with: (a) placebo; (b) Nefecon-budesonide (8 mg/day); and (c) Nefecon-budesonide (16 mg/day). The dotted line indicates no percentage change following intervention with placebo or Nefecon-budesonide.
- [0040] FIG. 15: shows that there were significant ( $p<0.05$ ) budesonide capsule-dependent decreases in the serum levels of secretory IgA. Whereas serum levels of IgA did not change.
- [0041] FIG. 16: shows that there was a significant ( $p<0.05$ ) budesonide capsule dose-dependent decrease in the serum levels of IgA-IgG immune complexes.
- [0042] FIG. 17: shows that there was a significant ( $p<0.05$ ) budesonide capsule dose-dependent decrease in the levels of poorly O-galactosylated IgA1.
- [0043] FIG. 18: shows that there were no differences observed in levels of total IgA, IgA1 and IgG with budesonide capsule treatment.
- [0044] FIG. 19: shows the in vitro dissolution profile of budesonide modified release capsules in the presence of added surfactant in the Level 1 fasted state simulated intestinal fluid at a pH of about 6.5.
- [0045] FIG. 20: shows the in vitro dissolution profile of budesonide modified release capsules in FaSSIF in the presence of added surfactant in the Level 1 fasted state simulated intestinal fluid at a pH of about 6.5 as compared with three other marketed budesonide-containing formulations.
- [0046] FIG. 21: shows the in vitro dissolution profiles of core-shell bead formulations comprising budesonide in the absence of an enteric coated capsule.
- [0047] FIG. 22: shows the in vitro dissolution profile of budesonide modified release capsules in FaSSIF in the absence of added surfactant in the Level 1 fasted state simulated intestinal fluid at a pH of about 6.5 as compared with three other marketed budesonide-containing formulations.
- [0048] FIGS. 23A-B: 23A shows the results of a PBPK model based on the dissolution of the modified release capsules; and 23B shows the results of a PBPK model based on the dissolution of comparative product Entocort®.

[0049] FIG. 24: Coronal MRI images of T2\*/T1 weighted TRUFI with detection of iron oxide loaded capsules at 15 and 90 minutes, as well as dispersion of iron oxide in the ileum at 270 minutes.

#### DISCLOSURE OF THE INVENTION

[0050] According to a first aspect of the invention, there is provided a method of treatment of IgA nephropathy, which method comprises:

[0051] (i) identifying a pharmaceutically acceptable composition intended to treat IgA nephropathy comprising budesonide and one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, which composition fulfils the following requirements in a standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test using a dissolution apparatus according to Apparatus 2 (Paddle Apparatus) of said test (as described hereinafter);

[0052] (a) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 120 minutes, when the dissolution medium is aqueous and has a pH of about 1.2;

[0053] (b) the composition fulfils the requirement that no more than about 10% of the budesonide is released into a pharmaceutically-relevant dissolution medium within about 30 minutes; and

[0054] (c) the composition fulfils the requirement that at least about 70% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 120 minutes, then

[0055] (ii) administering said composition to a patient with IgA nephropathy in need of said treatment,

[0056] which method is referred to hereinafter as “the method of the invention”.

[0057] By the term “pharmaceutically-relevant dissolution medium” we include media which is suitable for use in an in vitro dissolution assay the results of which are indicative of in vivo release at the relevant part of the intestinal tract. For example, the pharmaceutically-relevant dissolution medium may, in the alternative, be termed “enterically pharmaceutically-relevant dissolution medium” or “pharmaceutically-relevant enteric dissolution medium” may be any such medium that simulates dissolution and release in the small intestine or a relevant part thereof.

[0058] The pharmaceutically-relevant dissolution medium is preferably aqueous.

[0059] The pharmaceutically-relevant dissolution medium may have a pH of from about 6.2 to about 7.5, such as from about 6.5 to about 6.8.

[0060] The pharmaceutically-relevant dissolution medium may be a phosphate buffer medium at a pH of about 6.2, a Level 1 Fasted State Simulated Intestinal Fluid (FaSSIF) at a pH of about 6.5 (for example a FaSSIF buffer as defined below under the heading “Release in Level 1 Fasted State Simulated Intestinal Fluid at a pH of about 6.5”), a phosphate buffer medium at a pH of about 6.8 (for example the phosphate buffer defined below under the heading “Release in medium at pH 6.8”), or a phosphate buffer medium at a pH of about 7.2 or about 7.5.

[0061] The method of the invention may comprise (I) combining budesonide with one or more pharmaceutically-acceptable excipients that provide for a modified release of

said budesonide after administration to the gastrointestinal tract to make a pharmaceutically-acceptable composition intended to treat IgA nephropathy, and then (II) testing the composition in the standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test as set out above and, if the composition fulfils the requirements (a) to (c) as set out above, administering said composition to a patient with IgA nephropathy in need of said treatment.

[0062] As an alternative embodiment of the invention, there is provided a composition comprising a combination of budesonide with one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, wherein said composition fulfils the dissolution profile of step (i) outlined above for use in the treatment of IgA nephropathy.


[0063] As a further alternative embodiment of the invention, there is provided the use of a composition comprising a combination of budesonide with one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, wherein said composition fulfils the dissolution profile of step (i) outlined above for the manufacture of a medicament for the treatment of IgA nephropathy.

[0064] As referred to herein, the term “treatment” of IgA nephropathy, we further include the prophylaxis, or the diagnosis of the relevant condition in addition to therapeutic, symptomatic and/or palliative treatment.

[0065] For the avoidance of doubt, when referencing USP<711> we are referring to the test as published on 1 May 2016 and when referencing Ph. Eur. 2.9.3 we are referring to chapter 2.9.3 of the European Pharmacopoeia 10.0.

[0066] For the avoidance of doubt, step (ii) of administering the composition to a patient will only take place if the average (mean) of the tested compositions fulfils each and all of criteria (a), (b) and (c) of step (i).

[0067] As used herein, the term “budesonide” refers to a compound according to formula I:



[0068] Budesonide is also commonly referred to under its IUPAC name (16a, 17-[1(RS)-butylidenebis(oxy)]-11β,21-dihydroxypregna-1,4-diene-3,20-dione).

[0069] Although the composition of the invention comprises budesonide, it is understood that the composition may alternatively comprise a different corticosteroid that is capable of having topical action in a similar fashion to budesonide. Such suitable alternative corticosteroids include, but are not limited to, aclometasone, beclomethasone, betamethasone, clobetasol, hydrocortisone, dexamethasone, flunisolide, methylprednisolone, mometasone,

prednisolone, triamcinolone, fluticasone, ciclesonide, fludrocortisone and mixtures thereof, including mixtures comprising budesonide.

[0070] The Paddle Apparatus of Apparatus 2 may be operated at about 50 revolutions per minute (rpm), about 75 rpm or about 100 rpm. Preferably, the Paddle Apparatus of Apparatus 2 is operated at about 100 rpm, or at about 50 rpm.

[0071] The pharmaceutically-relevant dissolution medium of criterion b) and criterion c) may comprise a surfactant in an amount of about 0.5 mg/ml (0.05% w/v). The surfactant may be a polysorbate, preferably wherein the surfactant is polysorbate 80 (e.g., Tween 80).

[0072] In criterion a) of step (i) of the method the amount of budesonide released may be no more than about 5%, such as no more than about 2.5%, within about 120 minutes.

[0073] In criterion a) of step (i) of the method, the amount of budesonide released may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5%, within about 120 minutes.

[0074] In criterion b) of step (i) of the method the amount of budesonide released may be no more than about 5%, such as no more than about 2.5%, within about 30 minutes.

[0075] In criterion b) of step (i) of the method, the amount of budesonide released may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5%, within about 30 minutes.

[0076] In criterion c) of step (i) of the method the amount of budesonide released may be at least about 75%, for example about 80%, such as about 84% or about 85%, within about 120 minutes.

[0077] In criterion c) of step (i) of the method, the amount of budesonide released may be from about 70% to about 100%, such as from 75% to about 100%, for example from about 84% to about 100%, such as from about 85% to 100%, within about 120 minutes.

[0078] In criterion b) of step (i) of the method, the composition may further fulfil the requirement that no more than about 10% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 37.5 minutes, such as no more than about 5%, for example no more than about 2.5% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 37.5 minutes. For example, the amount of budesonide released into the pharmaceutically-relevant dissolution medium may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5%, within about 37.5 minutes. Optionally, the release within about 37.5 minutes is in the absence of surfactant in the dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50 rpm.

[0079] In criterion b) of step (i) of the method, the composition may further fulfil the requirement that at least about 20% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 75 minutes, such as at least about 21%, for example at least about 22% or 23% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 75 minutes. For example, the amount of budesonide released into the pharmaceutically-relevant dissolution medium may be from about 23% to about 74% within about 75 minutes. Optionally, the release within about 75 minutes is in the absence of surfactant in the dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50 rpm.

[0080] In criterion c) of the method, the composition may further fulfil the requirement that at least about 75% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 150 minutes, such as at least about 76%, for example at least about 77% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 150 minutes. For example, the amount of budesonide released into the pharmaceutically-relevant dissolution medium may be from about 77% to about 100% within about 150 minutes. Optionally, the release within about 150 minutes is in the absence of surfactant in the dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50 rpm.

[0081] In criterion b) of the method, the composition may further fulfil the requirement that no more than about 10% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 45 minutes, such as no more than about 5%, such as no more than about 2.5% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 45 minutes. For example, the amount of budesonide released into the pharmaceutically-relevant dissolution medium may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5% within about 45 minutes. Optionally, the release within about 45 minutes is in the absence of surfactant in the pharmaceutically-relevant dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50, 75 or 100 rpm.

[0082] In criterion b) of the method, the composition may further fulfil the requirement that no more than about 10% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 60 minutes, such as no more than about 5%, such as no more than about 2.5% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 60 minutes. For example, the amount of budesonide released into the pharmaceutically-relevant dissolution medium may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5% within about 60 minutes. Optionally, the release within about 60 minutes is in the absence of surfactant in the pharmaceutically-relevant dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50, 75 or 100 rpm.

[0083] In criterion b) of the method, the composition may further fulfil the requirement that from 50 to 90% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 90 minutes. Optionally, the release within about 90 minutes is in the absence of surfactant in the pharmaceutically-relevant dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50, 75 or 100 rpm.

[0084] In criterion c) of the method, the composition may further fulfil the requirement that at least about 80% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 180 minutes, such as at least about 85%, for example about 80% to about 100%, or about 85% to about 100% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 180 minutes. Optionally, the release within about 180 minutes is in the absence of surfactant in the pharmaceutically-relevant dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50, 75 or 100 rpm.

[0085] In criterion c) of the method, the composition may further fulfil the requirement that at least about 85% of the

budesonide is released into the pharmaceutically-relevant dissolution medium within about 240 minutes, such at least about 90%, for example about 90% to about 100% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 240 minutes. Optionally, the release within about 240 minutes is in the absence of surfactant in the pharmaceutically-relevant dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50, 75 or 100 rpm.

[0086] In criterion c) of the method, the composition may further fulfil the requirement that at least about 90% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 360 minutes, such at least about 95%, for example about 95% to about 100% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 360 minutes. Optionally, the release within about 360 minutes is in the absence of surfactant in the pharmaceutically-relevant dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50, 75 or 100 rpm.

[0087] In criterion c) of the method, the composition may further fulfil the requirement that at least about 90% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 480 minutes, such at least about 95%, for example about 95% to about 100% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 480 minutes. Optionally, the release within about 480 minutes is in the absence of surfactant in the pharmaceutically-relevant dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50, 75 or 100 rpm.

[0088] In criterion c) of the method, the composition may further fulfil the requirement that at least about 90% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 600 minutes, such at least about 95%, for example about 95% to about 100% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 600 minutes. Optionally, the release within about 600 minutes is in the absence of surfactant in the pharmaceutically-relevant dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50, 75 or 100 rpm.

[0089] In an embodiment, in criterion a) of step (i) of the method the dissolution in the acid resistance medium may be assessed according to the acceptance criteria in Acceptance Table 2 and/or Acceptance Table 3 of USP<711>/Table 2.9.3-2 and/or Table 2.9.3. -3 of Ph.Eur. 2.9.3.

[0090] In an embodiment, in criterion b) of step (i) of the method the dissolution in the pharmaceutically-relevant dissolution medium may be assessed according to the acceptance criteria in Acceptance Table 2 and/or Acceptance Table 3 of USP<711>/Table 2.9.3-2 and/or Table 2.9.3. -3 of Ph.Eur. 2.9.3.

[0091] In an embodiment, in criterion c) of step (i) of the method the dissolution in the pharmaceutically-relevant dissolution medium may be assessed according to the acceptance criteria in Acceptance Table 2 and/or Acceptance Table 4 of USP<711>/Table 2.9.3-2 and/or Table 2.9.3. -4 of Ph.Eur. 2.9.3.

[0092] For the avoidance of doubt, the amount of budesonide released in criterion b) at 30 minutes and criterion c) at 120 minutes is achieved in the presence and in the absence of added surfactant, such as added polysorbate 80 (e.g. Tween 80) at a concentration of about 0.5 mg/mL in the

pharmaceutically-relevant dissolution medium. In addition, the amount of budesonide released at 37.5 minutes, 60 minutes, 75 minutes, 90 minutes and 150 minutes is achieved in the presence and in the absence of added surfactant, such as added polysorbate 80 (e.g. Tween 80) at a concentration of about 0.5 mg/mL in the pharmaceutically-relevant dissolution medium.

[0093] In an embodiment the method comprises:

[0094] (i) identifying a pharmaceutically acceptable composition intended to treat IgA nephropathy comprising budesonide and one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, which composition fulfils the following requirements in a standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test using a dissolution apparatus according to Apparatus 2 (Paddle Apparatus) of said test (as described hereinafter) operated at 50 rpm;

[0095] (a) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 120 minutes, when the dissolution medium is aqueous and has a pH of about 1.2;

[0096] (b) the composition fulfils the requirement that no more than about 10% of the budesonide is released into a pharmaceutically-relevant dissolution medium within about 30 minutes, when the pharmaceutically-relevant dissolution medium is without surfactant;

[0097] (c) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 37.5 minutes, when the pharmaceutically-relevant dissolution medium is without surfactant;

[0098] (d) the composition fulfils the requirement that from about 23% to about 74% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 75 minutes, when the pharmaceutically-relevant dissolution medium is without surfactant;

[0099] (e) the composition fulfils the requirement that at least about 77% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 150 minutes, when the pharmaceutically-relevant dissolution medium is without surfactant; optionally

[0100] (f) wherein the composition fulfils the requirement that at least about 70% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 120 minutes, when the pharmaceutically-relevant dissolution medium is without surfactant.

[0101] In another embodiment the method comprises:

[0102] (i) identifying a pharmaceutically acceptable composition intended to treat IgA nephropathy comprising budesonide and one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, which composition fulfils the following requirements in a standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test using a dissolution apparatus according to Apparatus 2 (Paddle Apparatus) of said test (as described hereinafter) operated at 100 rpm;

[0103] (a) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 120 minutes, when the dissolution medium is aqueous and has a pH of about 1.2;

[0104] (b) the composition fulfils the requirement that no more than about 10% of the budesonide is released into a pharmaceutically-relevant dissolution medium within about 30 minutes, when the pharmaceutically-relevant dissolution medium is without surfactant;

[0105] (c) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 60 minutes, when the pharmaceutically-relevant dissolution medium is without surfactant;

[0106] (d) the composition fulfils the requirement that from about 50% to about 90% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 90 minutes, when the pharmaceutically-relevant dissolution medium is without surfactant; and

[0107] (e) the composition fulfils the requirement that at least about 70%, for example at least 75%, of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 120 minutes, when the pharmaceutically-relevant dissolution medium is without surfactant.

[0108] It is known from pharmacokinetic studies of drug absorption in the fasted state, that ingesting 200 to 250 ml of water along with a dosage form, a maximum total volume of about 300 to 500 mL will be available in the proximal small intestine (see Klein, *AAPS J.*, 12, 397, (2010)). Therefore, the dissolution tests that are employed in the method of the invention should employ volumes of dissolution media that is at least about 500 mL (e.g. about 900 mL). The initial volume of dissolution media used in criteria a), b) and c) may be about 900 mL.

[0109] The procedure used for testing the composition may be essentially in accordance with delayed-release solid dosage forms Method B of USP<711>/Ph.Eur. 2.9.3.

[0110] The temperature of the dissolution media in criteria a), b) and c) may be held at about 37° C.±0.5° C.

[0111] The number of compositions tested may be 6, or greater than 6, such as 12 or 24.

[0112] At each time point in criteria (a), (b) and (c) the amount of volume withdrawn from the dissolution medium may be 10 mL or 15 mL, optionally the volume withdrawn is not replaced. The withdrawal of dissolution medium does not affect the overall dissolution profile of the composition. That is to say, preferably the dissolution test is operated under sink conditions and that the amount of solvent outweighs that of the solute meaning that the withdrawal of a small amount for analysis purposes does not affect the dissolution.

Release in medium at pH 6.8

[0113] According to an alternative aspect of the invention, there is provided a method of treatment of IgA nephropathy, which method comprises:

[0114] (i) identifying a pharmaceutically acceptable composition intended to treat IgA nephropathy comprising budesonide and one or more pharmaceutically-acceptable excipients that provide for a modified

release of said budesonide after administration to the gastrointestinal tract, which composition fulfils the following requirements in a standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test using a dissolution apparatus according to Apparatus 2 (Paddle Apparatus) of said test (as described hereinafter);

[0115] (a) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 120 minutes, when the dissolution medium is aqueous and has a pH of about 1.2;

[0116] (b) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 30 minutes, when the dissolution medium is aqueous and has a pH of about 6.8; and

[0117] (c) the composition fulfils the requirement that at least about 70% of the budesonide is released into the dissolution medium within about 120 minutes, when the dissolution medium is aqueous and has a pH of about 6.8, then

[0118] (ii) administering said composition to a patient with IgA nephropathy in need of said treatment, and which method is referred to hereinafter as "the method of the invention".

[0119] The method of the invention may comprise (I) combining budesonide with one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract to make a pharmaceutically-acceptable composition intended to treat IgA nephropathy, and then (II) testing the composition in the standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test as set out above and, if the composition fulfils the requirements (a) to (c) as set out above (i.e. in relation to release in medium at pH 6.8), administering said composition to a patient with IgA nephropathy in need of said treatment.

[0120] As an alternative embodiment, there is provided a composition comprising a combination of budesonide with one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, wherein said composition fulfils the dissolution profile of step (i) outlined above (i.e. in relation to release in the above pharmaceutically-relevant medium at pH 6.8) for use in the treatment of IgA nephropathy.

[0121] As a further alternative embodiment, there is provided the use of a composition comprising a combination of budesonide with one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, wherein said composition fulfils the dissolution profile of step (i) outlined above (i.e. in relation to release in the pharmaceutically-relevant medium at pH 6.8) for the manufacture of a medicament for the treatment of IgA nephropathy.

[0122] For the avoidance of doubt, step (ii) of administering the composition to a patient will only take place if the average (mean) of the tested compositions fulfils all of criteria (a), (b) and (c) of step (i).

[0123] The Paddle Apparatus of Apparatus 2 may be operated at about 50 revolutions per minute (rpm), about 75

rpm or about 100 rpm. Preferably, the Paddle Apparatus of Apparatus 2 is operated at about 100 rpm, or at about 50 rpm.

[0124] The aqueous dissolution medium of criterion b) and criterion c) may comprise a surfactant in an amount of about 0.5 mg/ml (0.05% w/v). The surfactant may be a polysorbate, preferably wherein the surfactant is polysorbate 80 (e.g., Tween 80).

[0125] The aqueous dissolution medium of criterion b) and criterion c) may be a phosphate buffer medium, such as a sodium phosphate buffer solution at a concentration of about 50 mM.

[0126] The phosphate buffer medium at a pH of about 6.8 may be prepared by first preparing a 0.2 M sodium phosphate tribasic solution, then adding one part of the 0.2 M sodium phosphate tribasic solution to three parts of a 0.1 N hydrochloric acid solution. After mixing the two solutions together, the pH may be checked and adjusted if necessary to a pH of about 6.8 by adding either hydrochloric acid or sodium hydroxide.

[0127] In criterion a) of step (i) of the method the amount of budesonide released may be no more than about 5%, such as no more than about 2.5%, within about 120 minutes.

[0128] In criterion a) of step (i) of the method, the amount of budesonide released may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5%, within about 120 minutes.

[0129] In criterion b) of step (i) of the method the amount of budesonide released may be no more than about 5%, such as no more than about 2.5%, within about 30 minutes.

[0130] In criterion b) of step (i) of the method, the amount of budesonide released may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5%, within about 30 minutes.

[0131] In criterion c) of step (i) of the method the amount of budesonide released may be at least about 75%, for example about 80%, such as about 84% or about 85%, within about 120 minutes.

[0132] In criterion c) of step (i) of the method, the amount of budesonide released may from about 70% to about 100%, such as from about 75% to about 100%, for example from about 84% to about 100%, such as from about 85% to 100%, within about 120 minutes.

[0133] In criterion b) of step (i) of the method, the composition may further fulfil the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 37.5 minutes when the dissolution medium is aqueous and has a pH of about 6.8, such as no more than about 5%, for example no more than about 2.5% of the budesonide is released within about 37.5 minutes when the dissolution medium is aqueous and has a pH of about 6.8. For example, the amount of budesonide released may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5%, within about 37.5 minutes when the dissolution medium is aqueous and has a pH of about 6.8. Optionally, the release within about 37.5 minutes is in the absence of surfactant in the dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50 rpm.

[0134] In criterion b) of step (i) of the method, the composition may further fulfil the requirement that at least about 20% of the budesonide is released into the dissolution medium within about 75 minutes when the dissolution medium is aqueous and has a pH of about 6.8, such at least

about 21%, for example at least about 22% or 23% of the budesonide is released within about 75 minutes when the dissolution medium is aqueous and has a pH of about 6.8. For example, the amount of budesonide released may be from about 23% to about 74% within about 75 minutes when the dissolution medium is aqueous and has a pH of about 6.8. Optionally, the release within about 75 minutes is in the absence of surfactant in the dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50 rpm.

[0135] In criterion c) of the method, the composition may further fulfil the requirement that at least about 75% of the budesonide is released into the dissolution medium within about 150 minutes when the dissolution medium is aqueous and has a pH of about 6.8, such at least about 76%, for example at least about 77% of the budesonide is released within about 150 minutes when the dissolution medium is aqueous and has a pH of about 6.8. For example, the amount of budesonide released may be from about 77% to about 100% within about 150 minutes when the dissolution medium is aqueous and has a pH of about 6.8. Optionally, the release within about 150 minutes is in the absence of surfactant in the dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 50 rpm.

[0136] In criterion b) of the method, the composition may further fulfil the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 60 minutes when the dissolution medium is aqueous and has a pH of about 6.8, such as no more than about 5%, such as no more than about 2.5% of the budesonide is released within about 60 minutes when the dissolution medium is aqueous and has a pH of about 6.8. For example, the amount of budesonide released may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5% within about 60 minutes when the dissolution medium is aqueous and has a pH of about 6.8. Optionally, the release within about 60 minutes is in the absence of surfactant in the dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 100 rpm.

[0137] In criterion b) of the method, the composition may further fulfil the requirement that from 50 to 90% of the budesonide is released within about 90 minutes when the dissolution medium is aqueous and has a pH of about 6.8. Optionally, the release within about 90 minutes is in the absence of surfactant in the dissolution medium and at a paddle rotation speed of the Paddle Apparatus 2 of 100 rpm.

[0138] In an embodiment, in criterion a) of step (i) of the method the dissolution in the acid resistance medium may be assessed according to the acceptance criteria in Acceptance Table 2 and/or Acceptance Table 3 of USP<711>/Table 2.9.3-2 and/or Table 2.9.3. -3 of Ph.Eur. 2.9.3.

[0139] In an embodiment, in criterion b) of step (i) of the method the dissolution in the buffer stage medium may be assessed according to the acceptance criteria in Acceptance Table 2 and/or Acceptance Table 3 of USP<711>/Table 2.9.3-2 and/or Table 2.9.3. -3 of Ph.Eur. 2.9.3.

[0140] In an embodiment, in criterion c) of step (i) of the method the dissolution in the buffer stage medium may be assessed according to the acceptance criteria in Acceptance Table 2 and/or Acceptance Table 4 of USP<711>/Table 2.9.3-2 and/or Table 2.9.3. -4 of Ph.Eur. 2.9.3.

[0141] For the avoidance of doubt, the amount of budesonide released in criterion b) at 30 minutes and criterion c) at 120 minutes is achieved in the presence and in the absence

of added surfactant, such as added polysorbate 80 (e.g. Tween 80) at a concentration of about 0.5 mg/mL in the dissolution medium. In addition, the amount of budesonide released at 37.5 minutes, 60 minutes, 75 minutes, 90 minutes and 150 minutes is achieved in the presence and in the absence of added surfactant, such as added polysorbate 80 (e.g. Tween 80) at a concentration of about 0.5 mg/mL in the dissolution medium.

[0142] In an embodiment the method comprises:

[0143] (i) identifying a pharmaceutically acceptable composition intended to treat IgA nephropathy comprising budesonide and one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, which composition fulfils the following requirements in a standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test using a dissolution apparatus according to Apparatus 2 (Paddle Apparatus) of said test (as described hereinafter) operated at 50 rpm;

[0144] (a) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 120 minutes, when the dissolution medium is aqueous and has a pH of about 1.2;

[0145] (b) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 30 minutes, when the dissolution medium is aqueous, without surfactant and has a pH of about 6.8;

[0146] (c) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 37.5 minutes, when the dissolution medium is aqueous, without surfactant and has a pH of about 6.8

[0147] (d) the composition fulfils the requirement that from about 23% to about 74% of the budesonide is released into the dissolution medium within about 75 minutes, when the dissolution medium is aqueous, without surfactant and has a pH of about 6.8;

[0148] (e) the composition fulfils the requirement that at least about 77% of the budesonide is released into the dissolution medium within about 150 minutes, when the dissolution medium is aqueous, without surfactant and has a pH of about 6.8; optionally

[0149] (f) wherein the composition fulfils the requirement that at least about 70% of the budesonide is released into the dissolution medium within about 120 minutes, when the dissolution medium is aqueous, without surfactant and has a pH of about 6.8.

[0150] In another embodiment the method comprises:

[0151] (i) identifying a pharmaceutically acceptable composition intended to treat IgA nephropathy comprising budesonide and one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, which composition fulfils the following requirements in a standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test using a dissolution apparatus according to Apparatus 2 (Paddle Apparatus) of said test (as described hereinafter) operated at 100 rpm;

[0152] (a) the composition fulfils the requirement that no more than about 10% of the budesonide is

released into the dissolution medium within about 120 minutes, when the dissolution medium is aqueous and has a pH of about 1.2;

[0153] (b) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 30 minutes, when the dissolution medium is aqueous, without surfactant and has a pH of about 6.8;

[0154] (c) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 60 minutes, when the dissolution medium is aqueous, without surfactant and has a pH of about 6.8;

[0155] (d) the composition fulfils the requirement that from about 50% to about 90% of the budesonide is released into the dissolution medium within about 90 minutes, when the dissolution medium is aqueous, without surfactant and has a pH of about 6.8; and

[0156] (e) the composition fulfils the requirement that at least about 70%, for example at least 75%, of the budesonide is released into the dissolution medium within about 120 minutes, when the dissolution medium is aqueous, without surfactant and has a pH of about 6.8.

[0157] It is known from pharmacokinetic studies of drug absorption in the fasted state, that ingesting 200 to 250 ml of water along with a dosage form, a maximum total volume of about 300 to 500 mL will be available in the proximal small intestine (see Klein, *AAPS J.*, 12, 397, (2010)). Therefore, the dissolution tests that are employed in the method of the invention should employ volumes of dissolution media that is at least about 500 mL (e.g. about 900 mL). The initial volume of dissolution media used in criteria a), b) and c) may be about 900 mL.

[0158] The procedure used for testing the composition may be essentially in accordance with prolonged-release and/or delayed-release solid dosage forms Method B of USP<711>/Ph.Eur. 2.9.3.

[0159] The temperature of the dissolution media in criteria a), b) and c) may be held at about 37° C. $\pm$ 0.5° C.

[0160] The number of compositions tested may be 6, or greater than 6, such as 12 or 24.

[0161] At each time point in criteria (a), (b) and (c) the amount of volume withdrawn from the dissolution medium may be 10 mL or 15 mL, optionally the volume withdrawn is not replaced. The withdrawal of dissolution medium does not affect the overall dissolution profile of the composition. That is to say, preferably the dissolution test is operated under sink conditions and that the amount of solvent outweighs that of the solute meaning that the withdrawal of a small amount for analysis purposes does not affect the dissolution.

Release in Level 1 Fasted State Simulated Intestinal Fluid at a pH of about 6.5

[0162] According to a further alternative aspect of the invention, there is provided a method of treatment of IgA nephropathy, which method comprises:

[0163] (i) identifying a pharmaceutically acceptable composition intended to treat IgA nephropathy comprising budesonide and one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, which composition fulfils the

following requirements in a standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test using a dissolution apparatus according to Apparatus 2 (Paddle Apparatus) of said test;

[0164] (a) the composition fulfils the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 120 minutes, when the dissolution medium is aqueous and has a pH of about 1.2; and

[0165] (b) the composition fulfils the requirement that no more than about 10% of the budesonide is released into a dissolution medium comprising Level 1 fasted state simulated intestinal fluid at a pH of about 6.5 within about 30 minutes; and

[0166] (c) the composition fulfils the requirement that at least about 70% of the budesonide is released into a dissolution medium comprising Level 1 fasted state simulated intestinal fluid at a pH of about 6.5 within about 120 minutes; then

[0167] (ii) administering said composition to a patient with IgA nephropathy in need of said treatment, and which method is referred to hereinafter as "the method of the invention".

[0168] The method of the invention may comprise (I) combining budesonide with one or more pharmaceutically acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract to make a pharmaceutically acceptable composition intended to treat IgA nephropathy, and then (II) testing the composition in the standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test as set out above and, if the composition fulfils the requirements (a) to (c) as set out above (i.e. in relation to release in Level 1 fasted state simulated intestinal fluid at a pH of about 6.5), administering said composition to a patient with IgA nephropathy in need of said treatment.

[0169] As an alternative embodiment, there is provided a composition comprising a combination of budesonide with one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, wherein said composition fulfils the dissolution profile of step (i) outlined above (i.e. in relation to release in Level 1 fasted state simulated intestinal fluid at a pH of about 6.5) for use in the treatment of IgA nephropathy.

[0170] As a further alternative embodiment, there is provided the use of a composition comprising a combination of budesonide with one or more pharmaceutically-acceptable excipients that provide for a modified release of said budesonide after administration to the gastrointestinal tract, wherein said composition fulfils the dissolution profile of step (i) outlined above (i.e. in relation to release in Level 1 fasted state simulated intestinal fluid at a pH of about 6.5) for the manufacture of a medicament for the treatment of IgA nephropathy.

[0171] As referred to herein, the term "treatment" of IgA nephropathy, we further include the prophylaxis, or the diagnosis of the relevant condition in addition to therapeutic, symptomatic and/or palliative treatment.

[0172] The term 'Level 1 fasted state simulated intestinal fluid' (Level 1 FaSSIF-V1) will be understood by those skilled in the art to include a biorelevant dissolution medium that has a lower pH and buffering capacity than standard simulated intestinal fluid (the medium that is typically employed in standard USP/Ph.Eur.tests; pH 6.8), and which

has been developed specifically to simulate fasting conditions in the proximal small intestine (see, for example, Markopoulous et al, In-vitro simulation of luminal conditions for evaluation of performance of oral drug products: Choosing the appropriate test media, European Journal of Pharmaceutics and Biopharmaceutics, 93, 2015, 173-182).

[0173] Level 1 FaSSIF-V1 comprises a phosphate buffer system, such as one comprising NaH<sub>2</sub>PO<sub>4</sub> (at a concentration of about 28.5 mM), NaOH (at a concentration of about 13.8 mM), HCl (qs) and deionized water (qs), which produces a medium with an osmolality of about 270 mOsmol/kg and a buffer capacity of about 12 mEq/pH/L.

[0174] Surfactant may be added to the FaSSIF medium in the present method of the invention. The surfactant may be a polysorbate, such as polysorbate 80 (e.g. Tween 80). The surfactant may be present at a concentration of about 0.05% w/v (0.5 mg/mL) to facilitate the analysis.

[0175] Thus, there is further provided:

[0176] the method of the invention as hereinbefore defined;

[0177] a composition that fulfils the dissolution profile of step (i) outlined above for use in the treatment of IgA nephropathy; and

[0178] the use of a composition that fulfils the dissolution profile of step (i) outlined above for the manufacture of a medicament for the treatment of IgA nephropathy,

provided that, in each case, the FaSSIF medium employed in step (i) comprises a surfactant, such as a polysorbate, including polysorbate 80 (e.g. Tween 80), optionally present at a concentration of about 0.05% w/v (0.5 mg/mL) to facilitate the analysis.

[0179] For the avoidance of doubt, step (ii) of administering the composition to a patient will only take place if the average (mean) of the tested compositions fulfils all of criteria (a), (b) and (c) of step (i).

[0180] In criterion a) of step (i) of the method, the amount of budesonide released may be no more than about 5%, such as no more than about 2.5%, within about 120 minutes.

[0181] In criterion a) of step (i) of the method, the amount of budesonide released may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5%, within about 120 minutes.

[0182] In criterion b) of step (i) of the method, the amount of budesonide released may be no more than about 5%, such as no more than about 2.5%, within about 30 minutes.

[0183] In criterion b) of step (i) of the method, the amount of budesonide released may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5%, within about 30 minutes.

[0184] In criterion c) of step (i) of the method, the amount of budesonide released may be at least about 75%, for example about 80%, such as about 84% or about 85%, within about 120 minutes.

[0185] In criterion c) of step (i) of the method, the amount of budesonide released may be from about 70% to about 99%, such as from about 70% to about 90%, within about 120 minutes.

[0186] For the avoidance of doubt, the amount of budesonide released in criterion b) at 30 minutes and criterion c) at 120 minutes is achieved in the presence and in the absence of added polysorbate 80 (e.g. Tween 80) at a concentration of about 0.05% w/v (about 0.5 mg/mL) in the FaSSIF medium.

[0187] In criterion b) of step (i) of the method, the composition may further fulfil the requirement that no more than about 10% of the budesonide is released into the dissolution medium within about 60 minutes, such as no more than about 5%, for example no more than about 2.5% of the budesonide is released within about 60 minutes. For example, the amount of budesonide released may be from about 0% to about 10%, such as from about 0% to about 5%, for example from about 0% to about 2.5%, within about 60 minutes. The amount of budesonide released within about 60 minutes is achieved in the presence and absence of added polysorbate 80 (e.g. Tween 80) at a concentration of about 0.05% w/v (about 0.5 mg/mL) in the FaSSIF medium.

[0188] In criterion c) of step (i) of the method, in the presence of added polysorbate 80 (e.g. Tween 80) at a concentration of about 0.05% w/v (about 0.5 mg/mL) in the FaSSIF medium, the composition may further fulfil the requirement that at least about 20%, for example 25%, or 30%, such as 35% of the budesonide is released into the dissolution medium, for example at least about 40% of the budesonide is released, such as from about 30 to about 65% of the budesonide is released, such as about 35% to about 65% of the budesonide is released, including from about 40% to about 60% of the budesonide is released, for example from about 45% to about 55% of the budesonide is released into the dissolution medium, within about 90 minutes.

[0189] In criterion c) of step (i) of the method, in the absence of added polysorbate 80 (e.g. Tween 80) at a concentration of about 0.05% (about 0.5 mg/mL) w/v in the FaSSIF medium, the composition may further fulfil the requirement that at least about 10%, for example at least about 15% of the budesonide is released, such as from about 10 to about 50% of the budesonide is released, such as about 10% to about 40% of the budesonide is released, including from about 10% to about 30% of the budesonide is released, for example from about 15% to about 30% of the budesonide is released into the dissolution medium, within about 90 minutes.

[0190] In criterion c) of step (i) of the method, the composition may further fulfil the requirement that at least about 90% of the budesonide is released into the dissolution medium within about 180 minutes, for example at least about 95% of the budesonide is released within about 180 minutes. The amount of budesonide released within about 180 minutes is achieved in the presence and in the absence of added polysorbate 80 (e.g. Tween 80) at a concentration of about 0.05% w/v (about 0.5 mg/mL) in the FaSSIF medium.

[0191] The Paddle Apparatus of Apparatus 2 may be operated at about 50 rpm, about 75 rpm or about 100 rpm. Preferably, the Paddle Apparatus of Apparatus 2 is operated at about 100 rpm.

[0192] It is known from pharmacokinetic studies of drug absorption in the fasted state, that ingesting 200 to 250 ml of water along with a dosage form, a maximum total volume of about 300 to 500 mL will be available in the proximal small intestine (see Klein, *AAPS J.*, 12, 397, (2010)). Therefore, the dissolution tests that are employed in the method of the invention should employ volumes of dissolution media (including FaSSIF) that is at least about 500 mL (e.g. about 900 mL). Preferably, the initial volume of dissolution media used in criteria a), b) and c) is about 900 mL.

[0193] The procedure used for testing the composition may be essentially in accordance with delayed-release solid dosage forms Method B of USP<711>/Ph.Eur. 2.9.3.

[0194] The temperature of the dissolution media in criteria a), b) and c) may be held at about 37° C. $\pm$ 0.5° C.

[0195] The number of compositions tested may be at least 3, such as 6, or greater than 6, such as 12 or 24.

[0196] At each time point in criteria (a), (b) and (c) the amount of volume withdrawn from the dissolution medium may be 10 mL or 15 mL, optionally the volume withdrawn is not replaced. The withdrawal of dissolution medium does not affect the overall dissolution profile of the composition. That is to say, preferably the dissolution test is operated under sink conditions and that the amount of solvent outweighs that of the solute meaning that the withdrawal of a small amount for analysis purposes does not affect the dissolution.

[0197] Particularly, but not exclusively, when the composition of the invention is a core-shell composition as defined below, the method of the invention as defined herein may comprise the additional steps of:

[0198] (1) providing budesonide with the same extended release excipients, but in the absence of the delayed release excipients, as described herein; and

[0199] (2) identifying that the composition in a standard in vitro USP<711>/Ph.Eur. 2.9.3 dissolution test using a dissolution apparatus according to Apparatus 2 (Paddle Apparatus) of said test, fulfils the requirement that from about 20 to about 60%, such as from about 25 to about 50%, of the budesonide is released into a dissolution medium comprising Level 1 fasted state simulated intestinal fluid at a pH of about 6.5 within about 15 minutes.

[0200] In an embodiment, from about 70 to about 90% of the budesonide is released into the Level 1 FaSSIF-V1 dissolution medium defined herein within about 30 minutes in the absence of the delayed release coating (e.g., without an enteric coating) and, more preferably, from about 75 and about 85% of the budesonide is released into that dissolution medium within about 45 minutes.

[0201] In the absence of the delayed release (e.g., enteric) coating, from about 80% to about 90% of the budesonide may be released into the Level 1 FaSSIF-V1 dissolution medium as defined herein within about 60 minutes, more especially, from about 90% (such as about 95% (including about 97% and about 100%)) of the budesonide may be released into that dissolution medium within about 90 minutes, such as within about 120 minutes, including within about 180 minutes.

[0202] The budesonide composition in the absence of the delayed release excipients having the dissolution profile as described above is further confirmation that the bulk of the budesonide will be released in vivo to the ileum.

#### Effect on Biomarkers

[0203] The method of the invention, in all aspects outlined above, may lead to a statistically significant reduction in the level of serum B-cell activating factor (BAFF) (also known as tumour necrosis factor ligand superfamily member 13B (TNFSF13B)) in a subject relative to the baseline level of serum BAFF in the subject prior to treatment.

[0204] By "statistically significant reduction" we include the meaning of a reduction that is statistically significant to a p value $<$ 0.05 following use of a one-way Analysis of

Variance (ANOVA) when comparing the change seen in the treated patient group with the change seen in patients who received a placebo.

[0205] By "relative to the baseline level" we include the meaning that the measured level of a molecule (e.g. BAFF) is lower than the level measured at the beginning of the study (i.e. prior to administration of a drug). The baseline level is the level immediately prior to the start of the treatment and is used as a comparator for subsequently measured levels (e.g. immediately following a course of treatment, or at a timepoint following conclusion of a course of treatment). Thus, such a reduction is specific for the subject or group of subjects in question and is not an absolute value.

[0206] The reduction in the serum level of BAFF in the subject may be at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% relative to the baseline serum level of BAFF in the subject prior to treatment. For example, the reduction in the serum level of BAFF in the subject may be at least about 5%. In particular the reduction in the level of serum BAFF in the subject may be at least about 10%, such as the reduction in the serum level of BAFF in the subject may be at least about 14%.

[0207] The reduction in the serum level of BAFF in the subject may be from about 1% to about 70%. For example, the reduction in the serum level of BAFF in the subject may be from about 5% to about 50%. In particular, the reduction in the serum level of BAFF in the subject may be from about 5% to about 25%, such as the reduction in the serum level of BAFF in the subject may be from about 10% to about 25%. For example, the reduction in the serum level of BAFF in the subject may be from about 14% to about 23%.

[0208] The statistically significant reduction in the level of serum BAFF observed following the method of the invention may be associated with a statistically significant reduction in the serum level of one or more biomarkers associated with B-cell activation and/or proliferation, relative to the serum baseline level of the one or more biomarkers in the subject prior to treatment. Reduction in the serum levels of one or more biomarkers associated with B-cell activation and/or proliferation would indicate a beneficial effect in the treatment of diseases in which overactivity, overabundance, and/or over-proliferation of B-cells is linked to pathogenicity. For example, a reduction in the serum level of a biomarker produced by an active and/or proliferating B-cell may indicate that the activity/proliferation of B-cells is reduced and may be an indicator of a beneficial effect in the treatment of diseases, such as IgAN, in which overactivity, overabundance, and/or over-proliferation of B-cells is linked to pathogenicity.

[0209] The one or more biomarkers may include: transmembrane activator and CAML interactor (TACI) (also known as tumour necrosis factor receptor superfamily member 13B (TNFRSF13B)); B-cell maturation antigen (BCMA) (also known as tumour necrosis factor receptor superfamily member 17 (TNFRSF17)); BAFF-R (also known as tumour necrosis factor receptor superfamily member 13C (TNFRSF13C)); CD27; CD30; C—X—C motif chemokine 12 (CXCL12); C—X—C motif chemokine 13 (CXCL13); Chemokine (C-C motif) ligand 19 (CCL19); Interleukin 2 (IL-2); Interleukin 6 (IL-6); Chemokine (C-C

motif) ligand 3 (CCL3); Chemokine (C-C motif) ligand 4 (CCL4); soluble CD23 (sCD23); secretory IgA; IgA-IgG immune complexes; poorly O-galactosylated IgA1; or combinations thereof.

[0210] By "biomarker" (also known as "biological marker") we include the meaning of a measurable indicator of some biological state or condition. Often biomarkers are naturally occurring biological molecules, e.g. proteins, amino acids, antibodies, nucleic acids (e.g. RNA or DNA), nucleotides, lipids, carbohydrates/sugar, primary metabolites, or secondary metabolites. Such biomarkers can be associated with a particular pathological or physiological process, a disease, a pharmacological response to a drug, and may be used to predict disease incidence and prevalence, or to predict outcomes of disease and therapeutic intervention, etc.

[0211] The reduction in the serum level of the biomarker in the subject may be at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% relative to the baseline serum level of the biomarker in the subject prior to treatment. For example, the reduction in the serum level of the biomarker may be from about 1% to about 90%, or from about 5 to about 70%, or from about 10 to about 50%.

[0212] The reduction in the serum level of TACI in the subject may be at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 65%, 70%, or 75%, relative to the baseline serum level of TACI in the subject prior to treatment. For example, the reduction in the serum level of TACI in the subject may be at least about 5%. In particular, the reduction in the serum level of TACI in the subject may be at least about 11%.

[0213] The reduction in the serum level of TACI in the subject may be from about 1% to about 70%. For example, the reduction in the serum level of TACI in the subject may be from about 5% to about 50%. In particular, the reduction in the serum level of TACI in the subject may be from about 5% to about 20%, such as the reduction in the serum level of TACI in the subject may be from about 10% to about 20%. For example, the reduction in the serum level of TACI in the subject may be from about 11% to about 17%.

[0214] The reduction in the serum level of BCMA in the subject may be at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, or 60%, relative to the baseline serum level of BCMA in the subject prior to treatment. For example, the reduction in the serum level of BCMA in the subject may be at least about 1%. In particular, the reduction in the serum level of BCMA in the subject may be at least about 6%.

[0215] Furthermore, the reduction in the serum level of BCMA in the subject may be between about 1% and about 60%. For example, the reduction in the serum level of BCMA in the subject may be from about 1% to about 20%. In particular, the reduction in the serum level of BCMA in the subject may be from about 1% to about 10%, such as the reduction in the serum level of BCMA in the subject may be from about 5% to about 10%. For example, the reduction in the serum level of BCMA in the subject may be from about 6% to about 7%.

[0216] The reduction in the serum level of BAFF-R in the subject may be at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, relative to the baseline serum level of BAFF-R in the subject prior to treatment.

[0217] The reduction in the serum level of CD27 in the subject may be at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, or 65%, relative to the baseline serum level of CD27 in the subject prior to treatment. For example, the reduction in the serum level of CD27 in the subject may be at least about 5%. In particular, the reduction in the serum level of CD27 in the subject may be at least about 10%, such as the reduction in the serum level of CD27 in the subject may be at least about 15%.

[0218] The reduction in the serum level of CD27 in the subject may be from about 1% to about 60%. For example, the reduction in the serum level of CD27 in the subject may be from about 1% to about 25%. In particular, the reduction in the serum level of CD27 in the subject may be from about 5% to about 25%, such as the reduction in the serum level of CD27 in the subject may be from about 10% to about 20%. For example, the reduction in the serum level of CD27 in the subject may be from about 15% to about 19%.

[0219] The reduction in the serum level of CD30 in the subject may be at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, or 75%, relative to the baseline serum level of CD30 in the subject prior to treatment. For example, the reduction in the level of serum CD30 in the subject may be at least about 1%. In particular the reduction in the serum level of CD30 in the subject may be at least about 3%, such as the reduction in the serum level of CD30 in the subject may be at least about 5%.

[0220] The reduction in the serum level of CD30 in the subject may be from about 1% to about 75%. For example, the reduction in the serum level of CD30 in the subject may be from about 1% to about 25%. In particular, the reduction in the level of serum CD30 in the subject may be from about 1% to about 10%, such as the reduction in the serum level of CD30 in the subject may be from about 5% to about 10%. For example, the reduction in the serum level of CD30 in the subject may be from about 5% to about 8%.

[0221] The reduction in the serum level of secretory IgA in the subject may be at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, or 75%, relative to the baseline serum level of secretory IgA in the subject prior to treatment. For example, the reduction in the serum level of secretory IgA in the subject may be at least about 1%. In particular the reduction in the serum level of secretory IgA in the subject may be at least about 2%, such as the reduction in the serum level of secretory IgA in the subject may be at least about 3%.

[0222] The reduction in the serum level of secretory IgA in the subject may be from about 1% to about 75%. For example, the reduction in the serum level of secretory IgA in the subject may be from about 1% to about 25%. In particular, the reduction in the level of serum secretory IgA

in the subject may be from about 1% to about 10%, such as the reduction in the serum level of secretory IgA in the subject may be from about 1% to about 5%. For example, the reduction in the serum level of secretory IgA in the subject may be from about 1% to about 3%.

[0223] The reduction in the serum level of IgA-IgG immune complexes in the subject may be at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, or 75%, relative to the baseline serum level of IgA-IgG immune complexes in the subject prior to treatment. For example, the reduction in the serum level of IgA-IgG immune complexes in the subject may be at least about 1%. In particular the reduction in the serum level of IgA-IgG immune complexes in the subject may be at least about 5%, such as the reduction in the serum level of IgA-IgG immune complexes in the subject may be at least about 8%.

[0224] The reduction in the serum level of IgA-IgG immune complexes in the subject may be from about 1% to about 75%. For example, the reduction in the serum level of IgA-IgG immune complexes in the subject may be from about 1% to about 25%. In particular, the reduction in the level of serum IgA-IgG immune complexes in the subject may be from about 1% to about 20%, such as the reduction in the serum level of IgA-IgG immune complexes in the subject may be from about 2% to about 20%. For example, the reduction in the serum level of IgA-IgG immune complexes in the subject may be from about 2% to about 15%.

[0225] The reduction in the serum level of poorly O-galactosylated IgA1 in the subject may be at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, or 75%, relative to the baseline serum level of poorly O-galactosylated IgA1 in the subject prior to treatment. For example, the reduction in the serum level of poorly O-galactosylated IgA1 in the subject may be at least about 1%. In particular the reduction in the serum level of poorly O-galactosylated IgA1 in the subject may be at least about 5%, such as the reduction in the serum level of poorly O-galactosylated IgA1 in the subject may be at least about 8%.

[0226] The reduction in the serum level poorly O-galactosylated IgA1 in the subject may be from about 1% to about 75%. For example, the reduction in the serum level of poorly O-galactosylated IgA1 in the subject may be from about 1% to about 25%. In particular, the reduction in the level of serum poorly O-galactosylated IgA1 in the subject may be from about 1% to about 20%, such as the reduction in the serum level of poorly O-galactosylated IgA1 in the subject may be from about 2% to about 20%. For example, the reduction in the serum level of poorly O-galactosylated IgA1 in the subject may be from about 2% to about 15%.

[0227] The above-mentioned reduction in biomarkers is associated with the method of the invention, which requires that the budesonide-containing composition both exhibits a specified in vitro release profile and is intended for the treatment of, and/or capable of treating, IgAN.

[0228] As we have found that compositions exhibiting that in vitro release profile also exhibit appropriate reductions in the relevant biomarkers, this indicates that:

[0229] budesonide is being released to the region of the gastrointestinal tract within which the Peyer's Patches are predominantly located (e.g. the ileum); and

[0230] such compositions are accordingly capable of treating IgAN safely and efficaciously at an appropriate dose of budesonide.

#### Budesonide Containing Compositions

[0231] Compositions that may be employed in the method of the invention may comprise any combination of budesonide and one or more excipients that gives rise to the desired in vitro release profile in all aspects as described herein. This may be a combination of extended and/or delayed release coatings, which may be applied by a variety of formulative principles as described hereinafter.

[0232] In any event we prefer that compositions comprise at least one delayed release coating that is preferably located at the exterior of the composition to ensure that the active ingredient is not released within the stomach and/or until the small intestine is reached.

[0233] Such a delayed release coating may thus comprise a so-called "enteric coating", which refers to a material having gastroresistant properties, that is to say the material prevents dissolution or disintegration in a gastric environment, thus allowing the composition to pass through the stomach towards the ileum region of the small intestine.

[0234] The enteric coating may comprise azopolymers, disulphide polymers, cellulose acetate, cellulose acetate succinate, cellulose acetate phthalate, cellulose acetate tetrahydrophthalate, polyvinyl acetate phthalate, hydroxyethyl ethyl cellulose phthalate, methacrylic acid copolymers, polymethacrylic acid/acrylic acid copolymers, styrol maleic acid copolymers, hydroxypropyl methyl cellulose phthalate, acrylic resins, cellulose acetate trimellitate, hydroxypropyl methylcellulose trimellitate, shellac, hydroxyethyl ethyl cellulose phthalate, carboxymethylcellulose and hydroxypropyl methyl cellulose acetate succinate.

[0235] Particular enteric coating substances include cellulose acetate, cellulose acetate succinate, cellulose acetate phthalate, cellulose acetate tetrahydrophthalate, polyvinyl acetate phthalate, hydroxyethyl ethyl cellulose phthalate, methacrylic acid copolymers, polymethacrylic acid/acrylic acid copolymers, styrol maleic acid copolymers, hydroxypropyl methyl cellulose phthalate, acrylic resins, cellulose acetate trimellitate, hydroxypropyl methylcellulose trimellitate, shellac, hydroxyethyl ethyl cellulose phthalate, carboxymethylcellulose and hydroxypropyl methyl cellulose acetate succinate.

[0236] Preferred enteric coating substances include polyvinyl acetate phthalate and, particularly, methacrylic acid copolymers.

[0237] The skilled person will understand that the enteric coating can comprise other commonly used materials, such as talc (as plasticiser), dibutyl sebacate (as plasticiser), and a blend of HMPC and PEG as a sub-coating agent.

[0238] The composition may comprise one or more cores comprising budesonide that are encapsulated by a combination of delayed and extended release excipients, in order to substantially prevent release of the contents of said composition until the distal region of the small intestine (e.g., the ileum, such as the distal ileum) is reached. Such compositions are referred to hereinafter as "core-shell compositions of the invention", which encompasses "beads" and "encapsulated cores".

[0239] The cores comprising budesonide may be loaded into a capsule. When using a capsule, the delayed release coating (e.g., the enteric coating) may be on the capsule and not directly on the cores.

[0240] When the enteric coating is on the capsule, such as a size 1 capsule, the enteric coating may be present in an amount of from about 34 to about 46 mg per capsule, such as about 34 to about 42 mg per capsule, for example about 36 to about 40 mg per capsule.

[0241] In order to ensure in such core-shell compositions of the invention that the bulk of the budesonide is substantially released to the distal region of the small intestine (e.g., the ileum, such as the distal ileum), they may be (or may further be) individually coated with an extended release polymer coating.

[0242] For the avoidance of doubt, the extended release polymer coating is different to the delayed release coating.

[0243] Such an extended release coating may ensure that the bulk of the budesonide is released extensively throughout the ileum, and in combination with a delayed release coating (e.g. the enteric coating) may further ensure that such release is obtained substantially, and/or primarily, to the ileum region of the small intestine.

[0244] By "substantially released to the ileum region", we include that at least about 51%, such as at least about 60%, including at least about 70% or at least about 75%, such as at least about 80%, including at least about 90% of the initial content of active ingredient within a composition is released to that region.

[0245] The skilled person will understand that any pharmaceutical composition should be taken according to the prescription guidance and that if taken in such a manner that differs from the prescribing information then the desired effect may not be achieved. In the context of the present invention, it is preferable that the composition is taken orally at least one hour before a meal, and more preferably that the composition is taken orally in the morning at least one hour before the first meal of the day.

[0246] The extended release coating may comprise a pharmaceutically-acceptable polymeric blend comprising a water-insoluble polymer and a pore-forming polymer that is applied directly on the budesonide-containing cores. The resultant cores, or a composition comprising a plurality of cores, may then be encapsulated within a delayed release coating, which combination of excipients substantially prevents release of the contents of said composition until the ileum region of the small intestine is reached.

[0247] As used herein, the term "water-insoluble polymer" refers to a polymer having a solubility in aqueous solvents, such as water at about 25° C. of less than about 0.1 mg/mL. The presence of the water-insoluble polymer allows for the rate of release of budesonide in the composition to be controlled.

[0248] The water-insoluble polymer may be an alkyl cellulose or a derivative thereof, for example the water-insoluble polymer may be ethylcellulose (or a derivative thereof).

[0249] By the term, "alkyl cellulose or derivative thereof" we refer to chemical compounds derived from cellulose wherein a proton on at least some of the cellulose hydroxy groups has been replaced with an alkyl group.

[0250] As used herein, the term "pore-forming polymer" refers to polymers which have a higher water solubility than the water-insoluble polymer and are, therefore, capable of

dissolving first leaving pores in the coating, thus allowing a certain amount of water to penetrate towards the core.

[0251] Therefore, the pore-forming polymer may be defined as being “water-soluble”. That is to say, the pore-forming polymer has a solubility in aqueous solvents, such as water at 25° C. of at least about 10 mgmL<sup>-1</sup>.

[0252] The pore-forming polymer may have a nominal viscosity of from about 1 to about 300 mPa\*s, for example from about 1 to about 50 mPa\*s, such as from about 1 to about 30 mPa\*s, for example from about 1 to about 20 mPa\*s, such as from about 2 to about 9 mPa\*s, for example from about 2 to about 7 mPa\*s, preferably from about 2 to about 6 mPa\*s. The nominal viscosity of the pore-forming polymer may be measured at 20° C. as a 2 wt. % solution of the polymer in water by the standard Ph.Eur. 2.2.9 capillary viscometer method.

[0253] Furthermore, the pore-forming polymer may have a gelling temperature of from about 35 to about 65° C., for example from about 55 to about 65° C., such as from about 58 to about 64° C.

[0254] The pore-forming polymer may comprise a polymer selected from the list consisting of polyethylene glycol (PEG), hydroxypropylmethyl cellulose (HPMC) and hydroxypropyl cellulose (HPC). Preferably, the pore-forming polymer is hydroxypropylmethyl cellulose.

[0255] The degree of substitution of the HPMC with methoxy groups may be from about 15 to about 35 wt. %, for example from about 25 to about 35 wt. %, or about 27 to about 31 wt. %, such as from about 27 to about 30 wt. %. Furthermore, the degree of substitution of the HPMC with hydroxypropyl groups may be from about 4 to about 32 wt. %, for example from about 4 to about 20 wt. %, or from about 5 to about 15 wt. %, such as from about 7 to about 12 wt. %.

[0256] By the term “degree of substitution of the HPMC” this refers to the average level of substitution of the hydroxy groups on the cellulose chain and is expressed herein in percentage terms, i.e. the percentage of hydroxy groups that have been substituted with the moiety in question.

[0257] The water-insoluble polymer may present in an amount of from about 45 wt. % to about 90 wt. % of the total extended release coating and the pore-forming polymer may present in an amount of from about 35 wt. % to about 5 wt. % of the total extended release coating. Such as, the water-insoluble polymer may present in an amount of from about 45 wt. % to about 65 wt. % of the total extended release coating and the pore-forming polymer may present in an amount of from about 35 wt. % to about 15 wt. % of the total extended release coating. For example, the water-insoluble polymer may present in an amount of from about 47 wt. % to about 56 wt. % of the total extended release coating and the pore-forming polymer may present in an amount of from about 32 wt. % to about 22 wt. % of the total extended release coating.

[0258] The pharmaceutically-acceptable polymeric blend of the extended release coating may comprise a fatty acid in an amount of from about 2 wt. % to about 8 wt. %, such as from about 3 wt. % to about 7 wt. % of the total extended release coating.

[0259] The fatty acid may be an unsaturated fatty acid, for example a C<sub>4</sub> to C<sub>28</sub> unsaturated fatty acid, such as a C<sub>13</sub> to C<sub>22</sub> unsaturated fatty acid. For example, the unsaturated fatty acid may be selected from the group consisting of myristoleic acid, palmitoleic acid, sapienic acid, oleic acid,

elaidic acid, vaccenic acid, and erucic acid. Preferably, the unsaturated fatty acid is oleic acid.

[0260] The pharmaceutically-acceptable polymeric blend of the extended release coating may comprise a medium-chain triglyceride in an amount of from about 3 wt. % to about 12 wt. %, such as from about 5 wt. % to about 12 wt. %, for example from about 5 to 8 about wt. % of the total extended release coating.

[0261] By the term “medium-chain triglyceride” this refers to triglycerides having an aliphatic tail of from 6 to 12 carbons. For example, the medium-chain triglyceride may be selected from the list consisting of caproic acid, caprylic acid, capric acid and lauric acid.

[0262] The pharmaceutically-acceptable polymeric blend of the extended release coating may comprise a further water-soluble polymer in an amount of from about 1 wt. % to about 5 wt. %, such as from about 2 wt. % to about 3 wt. % of the total extended release coating. For the avoidance of doubt, the further water-soluble polymer is different to the pore-forming polymer. Preferably, the further water-soluble polymer is poly (ethylene glycol) having a molecular weight in the range of from about 200 to about 1000 g/mol.

[0263] The polymeric blend that coats the one or more cores may be coalescable. By the term “coalescable” when referring to the extended release polymeric blend, this refers to the polymers of the extended release blend being capable of blending to form a single polymeric phase coating.

[0264] Therefore, the extended release polymeric blend that coats the one or more cores may comprise one or more coalescable polymers. The one or more coalescable polymers may comprise the water-insoluble polymer.

[0265] The extended release polymer blend may be present in an amount of from about 5 to about 18 wt. % of the total bead/core, for example from about 6 to about 16 wt. % of the total bead/core, for example from about 6 to about 13 wt. %, such as about 6 to about 12 wt. % of the total bead/core.

[0266] It has been found that in the polymer blend of the extended release coating that the pore-forming polymer dissolves first, before the water-insoluble polymer, in an aqueous solution leaving pores in the coating, thus allowing a certain amount of water to penetrate towards the core in a controlled fashion.

[0267] The cores may have an average size in the range of from about 0.5 to about 3 mm, for example about 0.5 to about 2 mm, such as about 0.8 to about 1.5 mm.

[0268] The core-shell compositions of the invention may be prepared by a process which comprises:

[0269] (a) providing one or more cores comprising budesonide;

[0270] (b) which one or more cores are individually coated with by an extended release pharmaceutically-acceptable polymeric blend comprising a water-insoluble polymer in an amount of from about 45 wt. % to about 90 wt. % and a pore-forming polymer in an amount of from about 35 wt. % to about 5 wt. %; and wherein the composition is encapsulated within a delayed release coating, such as to substantially prevent release of the contents of said composition until the ileum region of the small intestine is reached.

[0271] The core-shell compositions of the above-described process may comprise any of the features as outlined above with respect to the method of the invention.

[0272] The cores may be prepared by providing inert (e.g., sugar) beads and coating them with an aqueous budesonide suspension. The inert beads may have a size of from about 1 to about 2 mm, such as about 1 to about 1.5 mm, for example about 1 to about 1.2 mm.

[0273] As referred to herein, the term "inert bead" includes a single pharmaceutically inert bead that provides the starting material for the preparation of the core-shell compositions of the invention.

[0274] The inert bead is preferably a commercially-available sugar sphere (often termed non-pareil). Sugar spheres predominantly comprise sucrose with smaller amounts of other materials added, such as starch. Suppliers of sugar spheres include Paulaur Corporation (USA), Chr. Hansen (Denmark), NP Pharm (France), Emilio Castelli (Italy) and JRS Pharma (Germany).

[0275] Prior to adding the extended release polymeric blend, the cores may be coated with a seal coating comprising a stabilised and a water-soluble polymer. The stabiliser may be an acid, wherein the acid is preferably citric acid and the soluble polymer is the same polymer as that used as the pore-forming polymer in the polymeric blend. Alternative stabilisers for inclusion in the seal coating include poly(vinyl pyrrolidone) (PVP) with the soluble polymer being the same polymer as that used as the pore-forming polymer in the polymeric blend.

[0276] The extended release polymeric blend may be applied to the cores as an aqueous polymer suspension and spraying the suspension onto the cores. The aqueous polymeric blend may be sprayed onto the cores at a temperature of from about 30° C. to about 65° C., such as from about 30° C. to about 50° C.

[0277] Spraying at a temperature towards the upper end of this region, such as about 50 to about 65° C. may avoid the requirement of a separate curing/coalescing step as outlined below.

[0278] The core-shell compositions of the invention may be obtainable by the process of coating the cores in a fluidized bed apparatus as defined above. That is to say, the coating with the extended release polymer blend may be carried out in a fluidized bed apparatus. Suitably fluidized bed apparatus are readily available from suppliers such as Glatt GmbH.

[0279] Following coating with the polymeric blend, the polymers may be coalesced, wherein the coalescing may be carried out by curing.

[0280] Curing may be carried out at a temperature of from about 55° C. to about 75° C., such as from about 60° C. to about 70° C., for example from about 63° C. to about 66° C. Furthermore, curing may be carried out for from about 1 hour to about 10 hours, for example from about 1 to about 5 hours, such as from about 2 to about 4 hours.

[0281] It was found that the extended release polymer coating blend allowed for the budesonide cores to be prepared in an economical fashion using a fluidized bed apparatus while arrive at compositions with the desired release profiles.

[0282] Therefore, in addition, the curing of the core-shell compositions of the invention may be carried out in a fluidized bed apparatus.

[0283] According to a further aspect of the invention there is provided a composition that gives rise to the desired in vitro release profile in all aspects as described herein, which comprises a plurality of beads comprising:

[0284] (a) budesonide-containing cores, in which cores budesonide is presented as a coating upon one or more inert core substrates (e.g., sugar beads) as described herein;

[0285] (b) an extended release coating presented upon said budesonide-containing cores, in an amount of between about 6 to about 12 wt. % of the total bead weight and which coating comprises a coalesced blend of at least two polymers (i) and (ii):

[0286] (i) any one of the water-insoluble polymers described herein (e.g., ethylcellulose), and

[0287] (ii) any one of the pore-forming polymers that described herein (e.g., hydroxypropylmethyl cellulose, with a degree of substitution with methoxy groups from about 27 to about 30 wt. % and/or a degree of substitution of with hydroxypropyl groups from about 7 to about 12 wt. %),

[0288] which blend of polymers (i) and (ii) is in any one of the proportions mentioned hereinbefore (e.g., the water-insoluble polymer (i) in an amount of from about 47 wt. % to about 56 wt. %, and the pore-forming polymer (ii) in an amount of from about 32 wt. % to about 22 wt. %, each of the total extended release coating),

which beads are then loaded into a capsule that is coated with a delayed release coating (e.g., any one of the enteric coatings described herein, such as polyvinyl acetate phthalate or, particularly, methacrylic acid copolymers).

[0289] The above composition may comprise one of more of the other preferred features disclosed in respect of core-shell compositions and/or beads as described herein, such as:

[0290] any one of the fatty acids described herein, e.g., in an amount of from about 3 wt. % to about 7 wt. % of the total extended release coating;

[0291] any one of the medium-chain triglycerides described herein, e.g., in an amount of from about 5 to about 8 wt. % of the total extended release coating;

[0292] one or more further water-soluble polymers, such as poly(ethylene glycol) having a molecular weight in the range of from about 200 to about 1000 g/mol, in an amount of from 2 wt. % to about 3 wt. % of the total extended release coating.

[0293] The above composition may also comprise a seal coating solution of an appropriate acid (e.g., a citric acid), located between the budesonide-containing cores and the extended release coating.

[0294] Each capsule may comprise about 4 mg of budesonide and when administered to the patient a total of about 16 mg may be delivered orally by way of the patient taking four capsules.

[0295] The coalesced blend of at least two polymers (i) and (ii), which is obtained prior to loading the coated beads into the capsule is preferably obtained by coating the budesonide-containing cores in a fluidized bed apparatus by way of the application of an aqueous dispersion comprising polymers (i) and (ii) as defined above at a temperature that is between about 30° C. to about 65° C., such as from about 30° C. to about 50° C., particularly about 50 to about 65° C. and then, if necessary, further coalescing the so coated polymeric extended release coating within the fluidized bed apparatus by application of a temperature of from about 60°

C. to about 70° C. (e.g., from about 63° C. to about 66° C.) for an appropriate time (e.g., from about 2 to about 4 hours) in order to cure it.

[0296] Alternatively, a polymeric blend may be sprayed as an organic solution. In this embodiment it is not necessary to carry out a further curing step.

[0297] As alluded to above, the core(s) that is/are coated by the aforementioned extended release polymeric blend may be loaded into a capsule, and the capsule may be coated with said delayed release coating.

[0298] However they are made, compositions that may be employed in the methods of the invention may be useful in the treatment of IgAN because they meet the in vitro release characteristics defined above.

[0299] Accordingly, there is further provided:

[0300] any one of the compositions as defined herein for use in the treatment of IgAN;

[0301] the use of any one of the compositions as defined herein for manufacture of a medicament for the treatment of IgAN; and

[0302] a method of treatment of IgAN, which method comprises the administration of any one of the compositions as defined herein to a patient in need of such treatment.

[0303] According to a further aspect of the invention, there is provided a core-shell composition of the invention, as described hereinbefore.

[0304] As an alternative to the core-shell composition, the composition may comprise a tablet comprising budesonide, which tablet is encapsulated within one or more excipients that substantially prevent release of the contents of said composition until the ileum region of the small intestine is reached. Such compositions are referred to hereinafter as "encapsulated tablet compositions of the invention"

[0305] The one or more excipients that encapsulate the tablet to substantially prevent release of the contents of said composition until the ileum region of the small intestine is reached may be an enteric coating as defined above.

[0306] The enteric coating in the tablet composition may be present in an amount of from about 5 wt. % to about 15 wt. %, such as from about 7 wt. % to about 13 wt. %, for example from about 8 wt. % to about 12 wt. % of the total tablet.

[0307] Tablet compositions of the invention may comprise a wet granulate of budesonide along with a filler. Optionally, the filler comprises dicalcium phosphate, microcrystalline cellulose, mannitol, or a mixture thereof.

[0308] The filler may be present in an amount of from about 50 wt. % to about 80 wt. %, such as about 60 wt. % to about 75 wt. %, for example about 65 wt. % to about 75 wt. % of the total tablet.

[0309] Tablet compositions of the invention may be a compressed tablet that further comprises a lubricant, optionally wherein the lubricant is magnesium stearate, aluminium stearate, calcium stearate, sodium stearate, zinc stearate, stearic acid, decanoic acid, dodecanoic acid, sodium stearyl fumarate or a mixture thereof.

[0310] The lubricant may be present in an amount of from about 0.1 wt. % to about 2 wt. %, such as from about 0.1 wt. % to about 1 wt. % of the total tablet.

[0311] Tablet compositions of the invention may further comprise a disintegrant, optionally wherein the disintegrant

is selected from crospovidone, croscarmellose sodium or sodium starch glycolate. Alternatively, the tablet may not comprise a disintegrant.

[0312] The disintegrant may be present in an amount of from about 0.5 wt. % to about 5 wt. %, such as from about 0.5 wt. % to about 4 wt. %, for example from about 0.8 wt. % to about 3.5 wt. % of the total tablet.

[0313] Tablet compositions of the invention may further comprise a binder, optionally wherein the binder is selected from hydroxyethyl cellulose, hydroxypropyl cellulose, copovidone, or mixtures thereof.

[0314] The binder may be present in an amount of from about 5 wt. % to about 10 wt. %, such as from about 6 wt. % to about 9 wt. %, for example from about 7 wt. % to about 9 wt. % of the total tablet.

[0315] Such tablet compositions of the invention may be gelling matrix tablets that further comprise a gelling matrix material, such as a low molecular weight HPMC (e.g., Hypromellose).

[0316] The gelling matrix material may be present in an amount of about 10 to about 25 wt. %, such as about 10 to about 20 wt. %, for example about 15 to about 20 wt. % of the total tablet.

[0317] The gelling matrix material may be diluted with a water-soluble filler, which water-soluble filler may comprise lactose, dextrose, mannitol, and combinations thereof.

[0318] The tablet may comprise an enteric coating comprising any of the materials as outlined above. The enteric coating may be present in an amount of from about 5 to about 15 wt. % of the total tablet, such as about 8 to about 12 wt. % of the total tablet.

[0319] Each tablet may comprise from about 2 to about 20 mg of budesonide, such as from about 4 to about 16 mg of budesonide. Preferably, each tablet comprises about 4 mg of budesonide.

[0320] Compositions when characterised by the method of the invention (in all aspects) are useful because they are more effective treatments of IgAN. Such compositions, pharmaceutical formulations, uses and methods described herein may also have the advantage that, in the treatment of IgAN, they may be more convenient for the physician and/or patient than, be more efficacious than, be less toxic than, have a broader range of activity than, be more potent than, produce fewer side effects than, have a lower inter-patient variability, or may have other useful pharmacological properties over, similar formulations or methods (treatments) known in the prior art, whether for use in the treatment of IgAN or otherwise.

[0321] Wherever the word 'about' is employed herein in the context of amounts, for example absolute amounts, such as weights, volumes, sizes, diameters, etc., or relative amounts (e.g. percentages) of individual constituents in a composition or a component of a composition (including concentrations and ratios), timeframes, and parameters such as temperatures, etc., it will be appreciated that such variables are approximate and as such may vary by  $\pm 10\%$ , for example  $\pm 5\%$  and preferably  $\pm 2\%$  (e.g.  $\pm 1\%$ ) from the actual numbers specified herein. This is the case even if such numbers are presented as percentages in the first place (for example 'about 10%' may mean  $\pm 10\%$  about the number 10, which is anything between 9% and 11%).

## EXAMPLES

## Example 1: Manufacturing of Core-Shell Composition

[0322] Opadry OY-7240 as referred to below is a dry powder polymer blend having the following components:

% w/w	Ingredients/Compendial Reference	Grade/Dye Strength	E Number	CFR Reference
90.910	HPMC 2910/Hypromellose (USP, Ph. Eur., JP, FCC)	5 mPas	E464	172.874
9.090	Macrogol/PEG (NF, FCC, Ph. Eur., JECFA, JP)	MW 400	E1521	172.820

[0323] Surelease is a polymer dispersion having the following components:

Test	Minimum	maximum
Oleic Acid, %	1.6	2.2
Ethylcellulose, %	17.0	20.0
Oleic Acid to Ethylcellulose Ratio	0.00	0.14
Medium Chain Triglycerides, %	0.80	4.00
MCT to Ethylcellulose Ratio	0.00	0.24
Glycerin, %	0.0	0.6
Solids, %	23.0	26.0
Ph	9.5	11.5
Viscosity Brookfield, cps	400.00	1500.00

[0324] For the avoidance of doubt, the minimum and maximum values in the table above refer to the minimum and maximum amounts of these components in different batches of Surelease.

[0325] A flow diagram detailing the preparation of budesonide beads according to the core-shell composition of the invention is provided in FIG. 1 and explained in further detail below.

[0326] A budesonide coating suspension was prepared by dissolving the Opadry OY-7240 Clear (2.29 kg) in purified water (26.5 kg) and then adding micronized budesonide (0.640 kg) to the solution while mixing continuously.

[0327] Sugar spheres (40.3 kg) having a mesh size of 16 to 18 were loaded into a pre-heated product bowl of a fluidized bed. When the product temperature reached the target of 45° C. the active coating suspension was sprayed onto the sugar spheres/inert core. The operation was monitored and controlled by a process computer. After the completion of spraying the required amount of active coating suspension, the active coated beads were dried and cooled.

[0328] A seal coating solution was prepared by dissolving citric acid monohydrate (0.093 kg) and Opadry OY-7240 Clear (2.26 kg) in purified water (21.8 kg) while mixing continuously.

[0329] The seal coating solution was applied onto the pre-warmed active coated beads. The operation was moni-

tored and controlled by a process computer. The coating was stopped when the required amount of solution has been sprayed, and the seal coated beads were dried and cooled. The fluid bed was emptied, and the beads were screened using 1.4 mm (14 Mesh) and 0.5 mm (35 Mesh) screens to remove any oversize and undersize particles. The beads were weighed, and the yield of the accepted fraction was calculated.

[0330] An extended release coating solution was prepared by adding Opadry OY-7240 (1.37 kg) to purified water (16.3 kg) while mixing continuously. An ethylcellulose suspension dispersion Type B (Surelease, 12.8 kg) was added to the Opadry solution during continued mixing.

[0331] The accepted fraction of the seal coated beads was loaded into the pre-heated product bowl of the fluidized bed with Wurster columns. When the product temperature reached the target temperature the extended release polymer coating suspension was applied onto the beads. The operation was monitored and controlled by a process computer. The amount sprayed was calculated from the amount of the accepted fraction of seal coated beads from the previous step.

[0332] The resulting polymer blend of the extended release coating on the beads comprised about 27.3 wt. % HPMC of the total blend and about 51.8 wt. % ethylcellulose of the total blend. The ethylcellulose is a water-insoluble polymer as defined above and the HPMC acts as the pore forming polymer.

[0333] After completion of the spraying, the beads were dried and cooled. The fluid bed was emptied, and the beads were screened using 1.4 mm (14 Mesh) and 0.5 mm (35 Mesh) screens to remove any oversize and undersize particles. The beads were weighed, and the yield of the accepted fraction was calculated.

[0334] The accepted fraction of the polymer coated beads was loaded into a pre-heated drying bowl of the fluid bed equipment. The beads were cured at target temperature of 65° C. for 3 hours. The operation was monitored and controlled by a process computer. The fluid bed was emptied, and a sample of the cured beads was taken for assay and dissolution testing. The cured beads were weighed, and the yield was calculated. The beads were filled into a stainless steel in-bin hopper.

[0335] The cured beads were then filled into capsules of size 1 using an automated encapsulator and the capsules were then coated with an enteric coating. The enteric coating of the capsules used was a blend of methacrylic acid and methyl methacrylate copolymers 1:1, and 1:2. The amount of enteric coating applied to each capsule was in the range of from about 34 to about 42 mg per capsule. The total amount of budesonide in each capsule was about 4 mg.

Example 2: General Process for Standard In Vitro Dissolution Test According to USP<711>/pH.Eur.  
2.9.3 in the Presence of Surfactant in the Buffer Stage and at a Paddle Rotation Speed of 100 Rpm

[0336] The in vitro dissolution of the encapsulated budesonide core-shell beads of Example 1 were analysed as described in Ph.Eur. 2.9.3 Dissolution test for solid dosage forms (using Apparatus 2) and as described in USP <711> Dissolution (using Apparatus 2). The measurement was carried out as described below.

## Dissolution Apparatus Setup

Dissolution Apparatus Setup	
Apparatus:	USP <711>/Ph. Eur. 2.9.3 Apparatus 2
Vessel Size/Type:	1000 mL/clear glass, round-bottom
Rotation Speed:	100 rpm
Media Volume:	900 mL-Acid Resistance
	900 mL-Buffer Dissolution
Test Temperature:	37.0 ± 0.5° C.
Pull Volume:	15 mL
Replacement:	No
Volume Discard:	5 mL
Sampling Time Point:	Acid Resistance Stage: 2 hours Buffer Stage: Specified time points 0.5 and 2 hours

[0337] Budesonide release was measured using Ultra Performance Liquid Chromatography (UPLC).

## Reagents and Standards

## Standards and Reference Materials:

[0338] Budesonide, Ph. Eur. CSR or suitable secondary standard.

## Other Reagents:

[0339] Tween 80, (Polyoxyethylene (20), Polysorbate (80), Fisher Scientific or equivalent.

## Dissolution Media and Diluents

## Acid Resistance Media

[0340] 0.1 N HCl Solution. To prepare 6 L of acid resistance media, 50 mL of concentrated HCl was combined with 6000 mL of water and the resulting solution was mixed well.

## 0.2 M Sodium Phosphate Tribasic Buffer Solution

[0341] To prepare 1 L of 0.2 M sodium phosphate tribasic buffer solution, about 76.02 g of sodium phosphate tribasic was added and dissolved into 1000 ml of water, followed by mixing.

Buffer Dissolution Media: 50 mM Sodium Phosphate Buffer, pH 6.8, with Tween 80.

[0342] To make 6 L of buffer dissolution media, 4500 mL Acid Resistance Media was combined with 1500 mL 0.2 M Sodium Phosphate Tribasic Buffer Solution and 3 g of Tween 80, followed by mixing. The pH was checked and, if necessary, adjusted pH to  $6.8 \pm 0.05$  using either Hydrochloric Acid or Sodium Hydroxide.

## Analytical Procedure

## Acid Resistance Procedure

[0343] Note: Care should be taken not to scrape or damage the capsules when placing into sinkers.

[0344] 900 mL of preheated degassed Acid Resistance media was placed into each of 6 dissolution vessels. The media was held at a temperature of  $37^\circ \text{C} \pm 0.5^\circ \text{C}$ .

[0345] The apparatus was operated as per the USP<711>/Pharmacopeia test no. 2.9.3 Rotating Paddle Apparatus method at 100 rpm.

[0346] 6 capsules were then placed each into separate coil sinkers and then into individual vessels.

[0347] At 2 hours, a 15-mL aliquot of the Acid Resistance Solution was withdrawn using a syringe.

[0348] The test solution was filtered with a Whatman GF/F with GMF filter, with the first 5 mL being discarded and the remaining solution being collected in a test tube.

[0349] The two steps below were completed after the Buffer Stage Dissolution has been initiated.

[0350] 5.0 mL of the filtered Acid Resistance Sample solution was pipetted into a 10 mL volumetric flask and diluted to volume with Acetonitrile.

[0351] The solution was mixed well and an aliquot was transferred to a HPLC vial and analysed.

[0352] The budesonide release in the acid resistance stage was assessed based on the acceptance criteria in USP<711>/Ph.Eur. 2.9.3.

## Buffer Stage Dissolution Procedure

[0353] After Acid Resistance samples were pulled, forceps were used to transfer each coil沉器 containing a budesonide capsule to a different set of dissolution vessels containing 900 mL of buffer dissolution media at temperature  $37^\circ \text{C} \pm 0.5^\circ \text{C}$ .

[0354] The apparatus was operated as per the USP<711>/Pharmacopeia test no. 2.9.3 Rotating Paddle Apparatus method at 100 rpm.

[0355] Specified timepoint sampling: At 0.5 and 2 hours, a 15-mL aliquot of the dissolution solution was withdrawn using a syringe. The fluid withdrawn was not replaced.

[0356] The test solution was filtered with a Whatman GF/F with GMF filter, with the first 5 mL being discarded and the remaining solution being collected in a test tube.

[0357] 5.0 mL of the filtered Dissolution Sample Solution was pipetted into a 10 mL volumetric flask and diluted to volume with acetonitrile.

[0358] The solution was mixed well and an aliquot was transferred to a HPLC vial.

[0359] Budesonide release in the buffer stage was assessed based on the acceptance criteria in USP<711>/Ph.Eur. 2.9.3.

Example 3: Dissolution Profile Analysis According  
to In Vitro USP<711>/Pharmacopeia Test No. 2.9.3  
in the Presence of Surfactant in the Buffer Stage  
and at a Paddle Rotation Speed of 100 Rpm

[0360] The capsules prepared in Example 1 ("budesonide capsules" or "nefcon budesonide") were tested under the dissolution conditions outlined in Example 2.

[0361] The overall dissolution profile for three separate batches can be seen in FIG. 2, with the time from 0 to 2 hours being at acid pH (pH 1.2) and the time from 2 to 4 hours being at buffered pH 6.8. FIG. 3 is a repeat of FIG. 2, showing only the dissolution at buffered pH 6.8 with time 0 to 120 minutes in FIG. 3 corresponding to time 2 to 4 hours in FIG. 2. Twelve capsules were tested per batch.

[0362] The quantitative results of the dissolution of budesonide in the various media at time points 2 hours at pH 1.2, 0.5 hours at pH 6.8 and 2 hours at pH 6.8 are provided in Table 1 below.

TABLE 1

		Batch 1	Batch 2	Batch 3
Dissolution (%)	pH 1.2, 2.0 h	Average	0	0
		SD	0	0
	pH 6.8, 0.5 h	Average	0	0
		SD	0	0
	pH 6.8, 2.0 h	Average	88	87
		SD	2.2	2.6

\*SD = standard deviation

[0363] The budesonide release in the acid resistance stage and the buffer stage was assessed based on the acceptance criteria in <711>/Ph.Eur. 2.9.3.

Example 4: General Process for Standard In Vitro Dissolution Test According to USP<711>/pH.Eur. 2.9.3 in the Absence of Surfactant in the Buffer Stage and at a Paddle Rotation Speed of 50 Rpm

[0364] The in vitro dissolution of the encapsulated budesonide core-shell beads of Example 1 were analysed as described in Ph.Eur. 2.9.3 Dissolution test for solid dosage forms (using Apparatus 2) and as described in USP <711> Dissolution (using Apparatus 2). The measurement was carried out as described below.

#### Dissolution Apparatus Setup

Dissolution Apparatus Setup	
Apparatus:	USP <711>/Ph. Eur. 2.9.3 Apparatus 2
Vessel Size/Type:	1000 mL/clear glass, round-bottom
Rotation Speed:	50 rpm
Media Volume:	900 mL-Acid Resistance 900 mL-Buffer Dissolution
Test Temperature:	37.0 ± 0.5° C.
Pull Volume:	15 mL
Replacement:	No
Volume Discard:	5 mL
Sampling Time Point:	Acid Resistance Stage: 2 hours Buffer Stage: Specified time points 0.625, 1.25 and 2.5 hours.

[0365] Budesonide release was measured using Ultra Performance Liquid Chromatography (UPLC).

#### Reagents and Standards

##### Standards and Reference Materials:

[0366] Budesonide, Ph. Eur. CSR or suitable secondary standard.

#### Dissolution Media and Diluents

##### Acid Resistance Media

[0367] 0.1 N HCl Solution. To prepare 6 L of acid resistance media, 50 mL of concentrated HCl was combined with 6000 ml of water and the resulting solution was mixed well.

##### 0.2 M Sodium Phosphate Tribasic Buffer Solution

[0368] To prepare 1 L of 0.2 M sodium phosphate tribasic buffer solution, about 76.02 g of sodium phosphate tribasic was added and dissolved into 1000 ml of water, followed by mixing.

Buffer Dissolution Media: 50 mM Sodium Phosphate Buffer, pH 6.8, with Tween 80.

[0369] To make 6 L of buffer dissolution media, 4500 mL Acid Resistance Media was combined with 1500 mL 0.2 M Sodium Phosphate Tribasic Buffer Solution. The pH was checked and, if necessary, adjusted pH to 6.8+0.05 using either Hydrochloric Acid or Sodium Hydroxide.

#### Analytical Procedure

##### Acid Resistance Procedure

[0370] Note: Care should be taken not to scrape or damage the capsules when placing into sinkers.

[0371] 900 mL of preheated degassed Acid Resistance media was placed into each of 6 dissolution vessels. The media was held at a temperature of 37° C.±0.5° C.

[0372] The apparatus was operated as per the USP<711>/ Pharmacopeia test no. 2.9.3 Rotating Paddle Apparatus method at 50 rpm.

[0373] 6 capsules were then placed each into separate coil sinkers and then into individual vessels.

[0374] At 2 hours, a 15-mL aliquot of the Acid Resistance Solution was withdrawn using a syringe.

[0375] The test solution was filtered with a Whatman GF/F with GMF filter, with the first 5 mL being discarded and the remaining solution being collected in a test tube.

[0376] The two steps below were completed after the Buffer Stage Dissolution has been initiated.

[0377] 5.0 mL of the filtered Acid Resistance Sample solution was pipetted into a 10 mL volumetric flask and diluted to volume with Acetonitrile.

[0378] The solution was mixed well and an aliquot was transferred to a HPLC vial and analysed.

[0379] The budesonide release in the acid resistance stage was assessed based on the acceptance criteria in USP<711>/ Ph.Eur. 2.9.3.

[0380] The capsules prepared in Example 1 (“budesonide capsules” or “nefecon budesonide”) were tested under the dissolution conditions outlined in this example.

[0381] The overall dissolution profile for seven separate batches can be seen in FIG. 4, with the time from 0 to 2 hours being at acid pH (pH 1.2) and the time from 2 to 4.5 hours being at buffered pH 6.8.

[0382] For all samples no more than 10% of the budesonide was released at time point 0.625 hours.

[0383] The quantitative results of the dissolution of budesonide at time point 1.25 hours (75 minutes) at pH 6.8 are provided in the table below.

Capsule batch no.	Average (%)	Min (%)	Max (%)	SD
3160733R	71	66	74	3.2
3166190R	48	43	53	4.1
3181645R	34	23	39	5.6
3197634R	61	57	71	5.1
3197635R	56	46	63	5.7
3197636R	55	49	60	3.9
3197637R	51	48	53	1.7

\*SD = standard deviation

[0384] At 1.25 h (75 minutes) from 23 to 74% of the budesonide is released.

**[0385]** The quantitative results of the dissolution of budesonide at time point 2.5 hours (150 minutes) at pH 6.8 are provided in the table below.

Capsule batch no.	Average (%)	Min (%)	Max (%)	SD
3160733R	101	99	103	1.5
3166190R	90	87	93	2.2
3181645R	80	77	83	2.1
3197634R	93	90	95	2.1
3197635R	93	88	98	3.9
3197636R	92	88	95	2.4
3197637R	89	87	92	1.7

**[0386]** The lowest amount of release observed at 2.5 hours was 77%.

**Example 5: Comparative Test According to USP<711>/Ph.Eur. 2.9.3 in the Presence of Surfactant in the Buffer Stage and at a Paddle Rotation Speed of 100 Rpm**

**[0387]** Outlined below is a variation of the in vitro test of Example 2. In this test the budesonide modified release capsules according to the present invention were analysed along with three other marketed budesonide containing formulations. The three formulations being Entocort® (Tillotts Pharma), Budenofalk® (Dr Falk Pharma GmbH) and Cortiment® (Ferring Pharmaceuticals, CH).

Dissolution Apparatus	USP <711>/Ph. Eur. 2.9.3
Apparatus:	Apparatus 2
Vessel Size/Type:	1000 mL/clear glass, round-bottom
Rotation Speed:	100 rpm
Media Volume:	900 mL-Acid Resistance
Test Temperature:	900 mL-Buffer Dissolution Profile 37.0 ± 0.5° C.
Pull Volume:	10 mL
Replacement:	No
Sampling Time Points	Acid Resistance Stage: 2 hours Buffer Stage: 15, 30, 45, 60, 90, 120, 180 minutes Check the pH of the dissolution medium after preparation, and again in each vessel after each test and record the result (Buffer Dissolution Medium only).
pH check	

#### Standards and Reference Materials

**[0388]** Budesonide, Ph. Eur. CSR.

#### Other Reagents:

**[0389]** Tween 80, (Polysorbate (80)), Fisher Scientific or equivalent.

#### Dissolution Media, Mobile Phase and Diluents

#### Acid Resistance Media

**[0390]** 0.1N HCl Solution. For example, to prepare 10 L, combine 82 mL of concentrated HCl and 10000 mL of water, mix well.

#### Buffer Dissolution Media

**[0391]** Sodium Phosphate pH 6.80 dissolution media was prepared by diluting one bottle (961.5 mL) of Reagecon DBC09-960 concentrate to a total volume of 25 L. See Sodium Phosphate pH 6.80 Dissolution Media 6x961.5 mL (reagecon.com) for more details.

**[0392]** The pH of the buffer solution was checked after preparation.

**[0393]** After Acid Resistance samples were pulled, the capsules were taken out of the solution with pincers and placed aside while the vessels were emptied, cleaned and filled with preheated buffer medium. 0.05% Tween 80 (or equivalent) was added to each dissolution vessel; for example 450 mg of Tween 80 was added to the dissolution vessels after they had been filled with 900 mL of pre-heated buffer, to arrive at a surfactant concentration of 0.05% w/v.

**[0394]** After all vessels reached the target temperature, the experiment was started by adding a capsule to each vessel.

#### Modification of Procedure for Budenofalk

**[0395]** A different procedure was used when the capsule broke during the acid stage. Most of the acid phase was carefully decanted and then the remaining acid was carefully removed with a pipette in order to remove as few pellets from the vessel as possible. The buffer stage was started by adding 900 mL of pre-heated buffer medium, followed by addition of Tween.

#### Sampling for Both Stages

**[0396]** 10 mL aspirated, 8 mL discarded (through Whatman filter (0.7 µm)), 1 mL sampled into HPLC vial.

**[0397]** The variations in this test compared to Example 2 do not have an effect on the overall dissolution profiles of the products tested and the comparative dissolution profiles of the three other marketed budesonide containing formulations can be seen in FIG. 5.

**Example 6: Comparative Test According to USP<711>/Ph.Eur. 2.9.3 in the Absence of Surfactant in the Buffer Stage and at a Paddle Rotation Speed of 100 Rpm**

**[0398]** Outlined below is a variation of the in vitro test of Example 2 and Example 5. In this test the enteric coated capsules filled with the core-shell beads of Example 1 were analysed along with the three other marketed budesonide containing formulations in a buffer stage solution at pH 6.8 and a paddle rotation speed of 100 rpm. The three formulations being Entocort® (Tillotts Pharma), Budenofalk® (Dr Falk Pharma GmbH) and Cortiment® (Ferring Pharmaceuticals, CH).

Dissolution Apparatus	USP <711>/Ph. Eur. 2.9.3
Apparatus:	Apparatus 2
Vessel Size/Type:	1000 mL/clear glass, round-bottom
Rotation Speed:	100 rpm
Media Volume:	900 mL-Acid Resistance
Test Temperature:	900 mL-Buffer Dissolution Profile 37.0 ± 0.5° C.
Pull Volume:	10 mL
Replacement:	No

-continued

Sampling Time Points	Acid Resistance Stage: 2 hours Buffer Stage: 15, 30, 45, 60, 90, 120, 180 minutes
pH check	Check the pH of the dissolution medium after preparation, and again in each vessel after each test and record the result (Buffer Dissolution Medium only).

#### Standards and Reference Materials

[0399] Budesonide, Ph. Eur. CSR.

#### Dissolution Media, Mobile Phase and Diluents

##### Acid Resistance Media

[0400] N HCl Solution. For example, to prepare 10 L, combine 82 mL of concentrated HCl and 10000 ml of water, mix well.

##### Buffer Dissolution Media

[0401] Sodium Phosphate pH 6.80 dissolution media was prepared by diluting one bottle (961.5 mL) of Reagecon DBC09-960 concentrate to a total volume of 25 L.

[0402] See Sodium Phosphate pH 6.80 Dissolution Media 6×961.5 ml (reagecon.com) for more details.

[0403] The pH of the buffer solution was checked after preparation.

[0404] After Acid Resistance samples were pulled, the capsules were taken out of the solution with pincers and placed aside while the vessels were emptied, cleaned and filled with 900 mL preheated buffer medium.

[0405] After all vessels reached the target temperature, the experiment was started by adding a capsule to each vessel.

#### Modification of Procedure for Budenofalk

[0406] A different procedure was used when the capsule broke during the acid stage. Most of the acid phase was carefully decanted and then the remaining acid carefully removed with a pipette in order to remove as few pellets from the vessel as possible. The buffer stage was started by adding 900 mL of pre-heated buffer medium.

#### Sampling for Both Stages

[0407] 10 mL aspirated, 8 mL discarded (through Whatman filter (0.7 µm)), 1 mL sampled into HPLC vial.

[0408] The dissolution profiles of the products tested and the comparative dissolution profiles of the budesonide core-shell beads of Example 1 and the three other marketed budesonide containing formulations can be seen in FIG. 6.

[0409] The table below shows the f2 values for comparison of the budesonide capsules according to the invention tested under this method with each of the other commercially available products. An f2 value of 50 or greater is required to demonstrate similarity of the profiles (FDA SUPAC Guidances 1995, 1997).

Nefecon	Entocort EC	Budenofalk	Cortiment
F2 value (USP)	18.1	16.0	15.8

[0410] It is clear that the release profiles of budesonide differ widely among the four commercial products. None of the f2 comparisons between Nefecon and the other products demonstrated similarity, which would require an f2 value of 50 or greater. In fact, based on the f2 evaluation as well as visual inspection of the graphical profiles, their release profiles must be considered strongly dissimilar.

#### Example 7: Administration of Bead Filled Capsules and Measurement of Biomarkers

[0411] For the study detailed in Example 7 to 16, enteric coated capsules filled with cured beads containing budesonide were used. The capsules are referred to below as "budesonide capsules".

#### Study Design

[0412] A randomised, double-blind, placebo-controlled experiment was carried out in which subjects with biopsy-confirmed primary IgA nephropathy and overt proteinuria were administered the budesonide capsules and the levels of various biomarkers were measured in blood taken from those patients: before the commencement of treatment; at the completion of treatment; and, at a timepoint following the completion of treatment.

#### Patients

[0413] Men or women aged at least 18 years with biopsy-confirmed primary IgA nephropathy and overt proteinuria were recruited for the run-in phase. All patients provided written informed consent before enrolment. Inclusion criteria for randomisation to treatment included estimated GFR (eGFR) of at least 45 mL/min per 1.73 m<sup>2</sup> and a urine protein creatinine ratio (UPCR) of more than 0.5 g/g or urinary total protein of at least 0.75 g/day—levels that would be considered to increase the risk of progression to end-stage renal disease. Either 24-h protein excretion or UPCR on the 24-h collection of urine were used to determine eligibility to overcome possible collection errors and deviations from normal creatinine excretion (e.g. physically active and muscular men), thus minimising the risk of unintentionally excluding patients.

#### Procedures

[0414] Medication was an oral capsule formulation (the budesonide capsule) or placebo, designed to provide sustained release of active compound that was delayed until the capsule reached the ileum, particularly the distal ileum, targeting the site with a high density of Peyer's Patches.

[0415] After screening, eligible patients were enrolled into a 6-month run-in phase, a 9-month treatment phase, and a 3-month follow-up phase; patient eligibility was assessed before run-in and treatment phases. During run-in, RAS blockade was optimised by up-titrating ACE inhibitors (ACEIs) and angiotensin II receptor blockers (ARBs) to a maximum recommended dose or maximum tolerated dose (in keeping with established clinical practice), to a target blood pressure of less than 130/80 mm Hg, UPCR of less than 0.5 g/g, and urine protein of less than 0.75 g/day. At the end of run-in, patients with persistent proteinuria (UPCR ≥0.5 g/g or proteinuria ≥0.75 g/day) despite optimised RAS blockade, eGFR (estimated by the Chronic Kidney Disease Epidemiology Collaboration [CKD-EPI] serum creatinine equation ≥45 mL/min) or measured GFR ≥45 mL/min per 1.73 m<sup>2</sup>, and blood pressure 160/100 mm Hg or less were eligible for randomisation to treatment.

**[0416]** An independent Data and Safety Monitoring Board (DSMB) monitored all safety issues and reviewed data at the interim analysis.

#### Randomisation and Masking

**[0417]** Patients were stratified according to their baseline UPCR ( $\leq 0.9$  g/g and  $> 0.9$  g/g) at month 0 (baseline). Patients were randomly allocated to treatment groups using a computer algorithm method of permuted blocks. Within each block, patients were allocated in a 1:1:1 ratio to 16 mg/day budesonide capsules, 8 mg/day budesonide capsules, or placebo. All patients continued optimised RAS blockade treatment throughout the treatment phase.

**[0418]** In total, 50 patients were given placebo, 51 patients were given 8 mg/day budesonide capsules, and 48 patients were given 16 mg/day budesonide capsules. Randomisation was done by Pharma Consulting Group AB (Uppsala, Sweden). Demographics and baseline characteristics of the recruited patients recruited are shown in Table 2.

**[0419]** The experiment was double-blind. Therefore, throughout the experiment and the analyses, allocation to treatment groups was unknown to each patient, all experimental staff (including the investigators and other staff who performed the randomisation and analyses), the sponsor, and the DSMB (the DSMB reviewed masked safety data and unmasked data were available should there be concerns).

**[0420]** To ensure masking, placebo capsules provided by the sponsor had the same appearance and route of administration as the active capsules. Patients self-administered masked capsules, once daily, 1 h before breakfast during the treatment phase. During follow-up (months 9-12), patients who received 16 mg/day budesonide capsules during months 0-9 were tapered to 8 mg/day for 2 weeks while all other patients (i.e. those who received budesonide capsules 8 mg/day or placebo during months 0-9) received placebo to maintain masking. No further experiment medication was administered after tapering.

TABLE 2

Patient demographics and baseline characteristics.				
Parameter	Placebo (n = 50)	Budesonide Capsules 8 mg/day (n = 51)	Budesonide Capsules 16 mg/day (n = 48)	Total (n = 149)
Age	38.9 (12.0)	40.6 (13.0)	37.5 (11.9)	39.0 (12.3)
Sex				
Male	35 (70%)	37 (73%)	33 (69%)	105 (71%)
Female	15 (30%)	14 (27%)	15 (31%)	44 (29%)
BMI (kg/m <sup>2</sup> )	27.5 (5.4)	26.5 (4.4)	27.8 (5.2)	27.3 (5.0)
Weight	85.2 (18.9)	80.9 (14.5)	86.7 (16.9)	84.2 (16.9)
Race				
Asian	1 (2%)	0 (0%)	1 (2%)	2 (1%)
Caucasian	48 (96%)	49 (96%)	47 (98%)	144 (97%)
Other	1 (2%)	2 (4%)	0 (0%)	3 (2%)

TABLE 2-continued

Patient demographics and baseline characteristics.				
Parameter	Placebo (n = 50)	Budesonide Capsules 8 mg/day (n = 51)	Budesonide Capsules 16 mg/day (n = 48)	Total (n = 149)
<b>Ethnicity</b>				
Hispanic/Latino	3 (6%)	11 (22%)	7 (15%)	21 (14%)
No-hispanic/Non-Latino	47 (94%)	40 (78.4%)	41 (85.4%)	128 (85.9%)
<b>Blood Pressure (mm Hg)</b>				
Systolic	128.1 (11.9)	127.7 (13.6)	126.7 (11.6)	127.5 (12.3)
Diastolic	80.2 (10.1)	80.3 (9.7)	78.1 (9.6)	79.6 (9.8)
UPCR (g/g)	0.8 (0.5-1.6)	0.8 (0.5-1.2)	0.8 (0.5-1.3)	0.8 (0.5-1.3)
24 h protein excretion (g)	1.2 (1.0-3.2)	1.1 (0.9-1.8)	1.3 (0.9-2.1)	1.2 (0.9-2.01)
UACR (g/g)	0.7 (0.4-1.3)	0.7 (0.5-1.0)	0.7 (0.4-1.2)	0.7 (0.4-1.1)
24 h albumin excretion (g)	1.1 (0.8-2.2)	1.0 (0.7-1.6)	1.1 (0.8-1.8)	1.0 (0.8-1.8)
eGFR CKD-EPI (creatinine formula; mL/min per 1.73 m <sup>2</sup> )	76.5 (23.2)	74.1 (25.8)	83.8 (25.9)	78.3 (25.1)
Time from diagnosis to start of treatment (days)	1101 (294-2870)	1972 (623-4188)	1219 (498-2573)	1499 (496-3162)

Data are n (%), mean (SD), or median (IQR).

Acronyms:

BMI = body-mass index;

CKD-EPI = Chronic Kidney Disease Epidemiology Collaboration Equation;

eGFR = estimated glomerular filtration rate;

UACR = urine albumin creatinine ratio;

UPCR = urine protein creatinine ratio.

**[0421]** Blood samples were taken from patients at the start of the treatment phase (month 0, before any treatment given), at the conclusion of the treatment phase (month 9), and at the conclusion of the follow-up phase (month 12). The samples obtained were tested for the levels of a number of biomarkers including: BAFF; APRIL; TACI; BCMA; CD27, CD30; secretory IgA; IgA-IgG Immune Complex; and poorly O-galactosylated IgA1.

**[0422]** Treatment code envelopes were provided for each randomised patient. In case of emergency, the code envelope could be opened. Any unmasked patient had to be withdrawn from the experiment.

#### Example 8: Treatment of Patients with the Budesonide Capsules Leads to a Reduction in Serum Levels of BAFF

#### Materials and Methods

**[0423]** Biomarkers were measured using a custom-designed bead-based multiplex Luminex® Assay (R & D Systems) as per the manufacturer's instructions.

**[0424]** The concept of the Luminex® Assay is based on fluorescent tagged microspheres selectively binding to the molecules of interest, thus allowing detection and quantification of multiple biomarkers simultaneously in a very small volume of serum.

**[0425]** In this study, biomarkers were selected and divided into panels (Panel 1: BAFF; APRIL; Panel 2: TACI; BCMA; CD27; CD30) according to the assay dynamic range.

**[0426]** A panel 1-specific tagged microparticle cocktail, or a panel 2-specific tagged microparticle cocktail was diluted 1:10 with the reagent diluent (provided with the microparticles by R & D Systems), serum samples were diluted 1:2 in reagent diluent and serial dilutions of a standard solution (provided with the microparticles by R & D Systems) were used to create a standard curve.

**[0427]** 50 µl of microparticle cocktail was applied to the Luminex® Assay plate followed by 50 µl of standard or sample. The plate was incubated on a microplate shaker at 800 rpm at room temperature for 2 hours. Proteins not bound to the microbeads were removed by washing with washing buffer (provided with the microparticles by R & D Systems).

**[0428]** A biotinylated antibody cocktail (provided with the microparticles by R & D Systems) was added to each well and incubated at room temperature for 1 hour. The plate was washed with washing buffer (provided with the microparticles by R & D Systems) and then incubated with 50 µl per well of streptavidin-Phycoerythrin (provided with the microparticles by R & D Systems) for 30-min, followed by a final wash step.

**[0429]** The microparticles were resuspended in 100 µl of wash buffer and the fluorescence in each well was read within 90 mins on the MAGPIX® Luminex machine.

**[0430]** Comparisons of differences in biomarker levels were conducted using a one-way Analysis of variance (ANOVA) statistical test with a p value<0.05.

## Results

**[0431]** As can be seen in FIG. 7 and Table 3, treatment of patients with the budesonide capsules at 8 mg/day and 16 mg/day led to statistically significant decreases in the serum level of BAFF compared with placebo treated patients in samples taken at the end of the 9-month treatment phase.

TABLE 3

Percentage change in serum levels of BAFF from start to end of treatment.				
Biomarker measured	Placebo	Budesonide Capsules (8 mg/day)	Budesonide Capsules (16 mg/day)	Combined budesonide results
BAFF	+0% (+48% to -36%)	-14% (+57% to -69%)	-23% (+38% to -66%)	-18% (+57% to -69%)

Mean values (Placebo, n = 50; budesonide capsules 8 mg/day n = 51; budesonide capsules 16 mg/day, n = 48; budesonide capsules 8 mg/day and 16 mg/day, n = 99) and range of values shown.

**[0432]** Furthermore, FIG. 8 and Table 4 show that in serum samples taken after completion of the follow-on phase (i.e. 3 months after the end of the treatment phase), the serum level of BAFF in patients previously treated with 16 mg/day budesonide capsules increases again indicating that the observed reduction was dependant on exposure to budesonide capsules.

TABLE 4

Percentage change in serum levels of BAFF from end of treatment to end of the follow-on phase.				
Biomarker measured	Placebo	Budesonide Capsules (8 mg/day)	Budesonide Capsules (16 mg/day)	Combined budesonide results
BAFF	+70% (-4% to -129%)	+29% (-9% to +82%)	+152% (-4% to +442%)	+87% (-9% to +442%)

Mean values (Placebo, n = 50; budesonide capsules 8 mg/day n = 51; budesonide capsules 16 mg/day, n = 48; budesonide capsules 8 mg/day and 16 mg/day, n = 99) and range of values shown.

**[0433]** The reduction in serum levels of BAFF following treatment with budesonide capsules is consistent with a disease-modifying action in IgAN.

Example 9: Treatment of Patients with Budesonide Capsules does not Lead to a Reduction in Serum Levels of APRIL

## Materials and Methods

**[0434]** The same materials and methods were used as described in Example 8 above.

**[0435]** Comparisons of differences in APRIL levels were conducted using a one-way Analysis of variance (ANOVA) statistical test with a p value<0.05.

## Results

**[0436]** As can be seen in FIG. 9 and Table 5, treatment of patients with budesonide capsules at 8 mg/day and 16 mg/day did not lead to an observable change in the serum level of APRIL in samples taken at the end of the 9-month treatment phase, suggesting that the effect produced by treatment with the budesonide capsules is specific to BAFF. Similarly, no change was observed in the placebo group.

TABLE 5

Percentage change in serum levels of APRIL from start to end of treatment.				
Biomarker measured	Placebo	Budesonide capsules (8 mg/day)	Budesonide capsules (16 mg/day)	Combined budesonide results
APRIL	-2% (+79% to -43%)	+2% (+148% to -72%)	+3% (+41% to -66%)	+3% (+148% to -72%)

Mean values (Placebo, n = 50; budesonide capsules 8 mg/day n = 51; budesonide capsules 16 mg/day, n = 48; budesonide capsules 8 mg/day and 16 mg/day, n = 99) and range of values shown.

Example 10: Decrease in Serum Levels of BAFF Following Budesonide Capsule Treatment is Associated with a Reduction in Serum Levels of TACI

## Materials and Methods

**[0437]** The same materials and methods were used as described in Example 8 above.

**[0438]** Comparisons of differences in TACI levels were conducted using a one-way Analysis of variance (ANOVA) statistical test with a p value<0.05.

## Results

**[0439]** As can be seen in FIG. 10 and Table 6, treatment of patients with budesonide capsules at 8 mg/day and 16 mg/day led to statistically significant decreases in the serum level of TACI compared with placebo treated patients in samples taken at the end of the 9-month treatment phase. In the placebo group, the level of TACI was actually seen to increase slightly, relative to the baseline level.

TABLE 6

Percentage change in serum levels of TACI from start to end of treatment.				
Biomarker measured	Placebo	Budesonide capsules (8 mg/day)	Budesonide capsules (16 mg/day)	Combined budesonide results
TACI	+16% (+143% to -61%)	-17% (+26% to -54%)	-11% (+60% to -65%)	-14% (+60% to -65%)

Mean values (Placebo, n = 50; budesonide capsules 8 mg/day n = 51; budesonide capsules 16 mg/day, n = 48; budesonide capsules 8 mg/day and 16 mg/day, n = 99) and range of values shown.

Example 11: Decrease in Serum Levels of BAFF Following the Budesonide Capsule Treatment is Associated with a Reduction in Serum Levels of BCMA

## Materials and Methods

**[0440]** The same materials and methods were used as described in Example 6 above.

**[0441]** Comparisons of differences in BCMA levels were conducted using a one-way Analysis of variance (ANOVA) statistical test with a p value<0.05.

## Results

**[0442]** As can be seen in FIG. 11 and Table 7, treatment of patients with budesonide capsules at 8 mg/day and 16 mg/day led to a statistically significant decrease in the serum level of BCMA compared with placebo treated patients in samples taken at the end of the 9-month treatment phase. In the placebo group, the level of BCMA was actually seen to increase very slightly relative to the baseline level.

TABLE 7

Percentage change in serum levels of BCMA from start to end of treatment.				
Biomarker measured	Placebo	Budesonide capsules (8 mg/day)	Budesonide capsules (16 mg/day)	Combined budesonide results
BCMA	+3% (+68% to -27%)	-7% (+8% to -53%)	-6% (+16% to -39%)	-7% (+16% to -53%)

Mean values (Placebo, n = 50; budesonide capsules 8 mg/day n = 51; budesonide capsules 16 mg/day, n = 48; budesonide capsules 8 mg/day and 16 mg/day, n = 99) and range of values shown.

Example 12: Decrease in Serum Levels of BAFF Following Budesonide Capsule Treatment is Associated with a Reduction in Serum Levels of CD27

**[0443]** As the study was restricted to evaluating changes in the circulating, rather than tissue, levels of each biomarker, and were therefore uncertain of the actual site where modulation occurred, pathway analyses were undertaken to determine whether biomarkers that were significantly modulated by budesonide capsules, including those surrogate biomarkers of immune cell activation (including sCD27, and sCD30 (see Example 13)), were associated with any particular biologic processes and pathways.

## Materials and Methods

**[0444]** The same materials and methods were used as described in Example 8 above.

**[0445]** Comparisons of differences in CD27 levels were conducted using a one-way Analysis of variance (ANOVA) statistical test with a p value<0.05.

## Results

**[0446]** As can be seen in FIG. 12 and Table 8, treatment of patients with budesonide capsules at 8 mg/day and 16 mg/day led to a statistically significant decrease in the serum level of CD27 compared with placebo treated patients in samples taken at the end of the 9-month treatment phase. In the placebo group, the level of CD27 was actually seen to increase slightly relative to the baseline level.

TABLE 8

Percentage change in serum levels of CD27 from start to end of treatment.				
Biomarker measured	Placebo	Budesonide capsules (8 mg/day)	Budesonide capsules (16 mg/day)	Combined budesonide results
CD27	+6% (+84% to -30%)	-15% (+10% to -56%)	-19% (+4% to -44%)	-17% (+10% to -56%)

Mean values (Placebo, n = 50; budesonide capsules 8 mg/day n = 51; budesonide capsules 16 mg/day, n = 48; budesonide capsules 8 mg/day and 16 mg/day, n = 99) and range of values shown.

**[0447]** Furthermore, FIG. 13 and Table 9 show that in samples taken after completion of the follow-on phase (i.e. 3 months after the end of the treatment phase), the serum level of CD27 in patients previously treated with 16 mg/day budesonide capsules increases again indicating that the observed reduction was dependant on exposure to the budesonide capsules.

TABLE 9

Percentage change in serum levels of CD27 from end of treatment to end of the follow-on phase.				
Biomarker measured	Placebo	Budesonide capsules (8 mg/day)	Budesonide capsules (16 mg/day)	Combined budesonide results
CD27	+16% (-14% to +65%)	+20% (-2% to +183%)	+36% (0% to +132%)	+28% (-2% to +183%)

Mean values (Placebo, n = 50; budesonide capsules 8 mg/day n = 51; budesonide capsules 16 mg/day, n = 48; budesonide capsules 8 mg/day and 16 mg/day, n = 99) and range of values shown.

**Example 13: Decrease in Serum Levels of BAFF Following Budesonide Capsule Treatment is Associated with a Reduction in Serum Levels of CD30**

#### Materials and Methods

[0448] The same materials and methods were used as described in Example 8 above.

[0449] Comparisons of differences in CD30 levels were conducted using a one-way Analysis of variance (ANOVA) statistical test with a p value<0.05.

#### Results

[0450] As can be seen in FIG. 14 and Table 10, treatment of patients with budesonide capsules at 8 mg/day and 16 mg/day led to a small but statistically significant decrease in the serum level of CD30 compared with placebo treated patients in samples taken at the end of the 9-month treatment phase. In the placebo group, the level of CD30 was actually seen to increase slightly relative to the baseline level.

BSA for 1 hour at room temperature. Serum samples and standards (high, medium, and low) were diluted 1:10 in PBS and applied to the plate, following washing, and incubated at 4° C. overnight. Following this, the plate was washed and polyclonal rabbit anti-human IgA HRP (Sigma, 1:2000) was added to each well, and the plate was incubated at room temperature for 90 minutes. The plate was then washed again, and levels of secretory IgA were visualized with o-phenylenediamine dihydrochloride substrate. OD492 of the high, medium, and low standards on each plate was used to normalize the values of the plates to a standard plate.

#### Results

[0453] There were significant (p<0.05) budesonide capsule-dependent decreases in the serum levels of secretory IgA. Serum levels of secretory IgA did not change (FIG. 15). This is indicative of local release and local gut action of the targeted release of the budesonide capsules, rather than systemic exposure of budesonide

TABLE 10

Percentage change in serum levels of CD30 from start to end of treatment.				
Biomarker measured	Placebo	Budesonide capsules (8 mg/day)	Budesonide capsules (16 mg/day)	Combined budesonide results
CD30	+11% (+88% to -46%)	-8% (+80% to -72%)	-5% (+92% to -67%)	-6% (+92% to -72%)

Mean values (Placebo, n = 50; budesonide capsules 8 mg/day n = 51; budesonide capsules 16 mg/day, n = 48; budesonide capsules 8 mg/day and 16 mg/day, n = 99) and range of values shown.

[0451] Consistent with the large meta-analysis of genome-wide association (GWA) studies (Gesualdo L, Di Leo V, Coppo R. The mucosal immune system and IgA nephropathy. Semin Immunopathol. 2021; 43:657-668; Coppo R. The gut-renal connection in IgA nephropathy. Semin Nephrol. 2018; 38:504-512.), this study identified the intestinal immune network for IgA production as one of the most enriched Kyoto Encyclopaedia of Genes and Genomes (KEGG) pathways, indicating that the mechanism of action of budesonide capsules is, at least in part, driven by an effect within the ileal gut-associated lymphoid tissue (GALT).

**Example 14:-Analysis of Secretory IgA Levels Following Budesonide Capsule Treatment**

#### Materials and Methods

[0452] Monoclonal mouse anti-human secretory component (Sigma) diluted 1:10,000 in coating buffer was applied to the wells of an immunoplate and incubated at 4° C. overnight. The plate was then washed and blocked with 2%

**Example 15: Analysis of IgA-IgG Immune Complex and Poorly O-Galactosylated IgA1 Levels Following Budesonide Capsule Treatment**

#### Materials and Methods

[0454] IgA-IgG immune complexes: Wells from a 96-well immunoplate were coated with AffiniPure F(ab')2 fragment goat anti-human serum IgA (a chain specific) (Jackson Immunology) diluted to 5 µg/mL in coating buffer. Following overnight incubation at 4° C., the plate was washed and non-specific protein binding was blocked with 2% BSA in PBS for 1 hour at room temperature. The test serum samples and standards (high, medium, and low) were diluted 1:500 in PBS, added to duplicate wells, and incubated at 4° C. overnight. The plate was then washed and incubated for 90 minutes with polyclonal rabbit anti-human IgG-HRP (Dako) diluted 1:2000 in PBS. The plate was washed again for 4 cycles, and levels of IgA/IgG ICs in the serum samples were visualized using o-phenylenediamine dihydrochloride sub-

strate. OD492 of the high, medium, and low standards on each plate was used to normalize the values of the plates to a standard plate.

[0455] Undergalactosylated IgA: Levels of poorly O-galactosylated IgA1 were measured using a commercially available KM55 ELISA (cat no 27600, Immuno-Biological Laboratories, Inc. Minneapolis, MN 55432, USA)

### Results

[0456] At the end of the treatment there was a significant ( $p<0.05$ ) budesonide capsule dose-dependent decrease in the serum levels of IgA-IgG immune complexes. IgA-IgG immune complex levels returned to baseline levels over the 3 months following cessation of budesonide capsule treatment (FIG. 16). There was a similar, although less pronounced, change in the levels of poorly O-galactosylated IgA1 (FIG. 17).

[0457] What is particularly interesting is that it has been shown that treatment of IgAN with systemic glucocorticoids lowers both total serum IgA and O-galactosylated IgA1 (Kosztyu P et al.: Glucocorticoids Reduce Aberrant O-Glycosylation of IgA1 in IgA Nephropathy Patients. Kidney Blood Press Res 2018; 43:350-359. However, with the present treatment there were no differences observed in levels of total IgA, IgA1 and IgG with budesonide capsule treatment (FIG. 18), which led us to conclude that the effect of local ileal treatment with budesonide capsules was selective for the pathogenic antibodies but not effective on the general pool of IgA, IgA1 and IgG.

[0458] These results show that treatment with budesonide capsules is supportive of a direct effect of budesonide capsules on the underlying pathogenic pathways in IgAN and that the budesonide payload has a predominantly topical effect rather than a systemic effect, leading to reduced side effects for patients when treated with budesonide capsules.

#### Example 16: Comparison of Lag Times for Onset of Plasma Profiles Between the Budesonide Capsules and Reference Commercial Product

[0459] A test product of the budesonide capsules and a reference commercial product (Entocort® EC; AstraZeneca) containing the same active ingredient were administered to 24 subjects in the fasted state in a randomized, crossover clinical study.

[0460] The lag times in hours for the onset of blood levels for each subject after administration of the test formulation on two separate occasions, and for the administration of the Reference (REF) product (with suitable wash-out period of 7 to 14 days), are shown in Table 11 below.

TABLE 11

Subject No.	Lag times for onset of plasma profiles			
	Test Formulation Administration 1 (F1)	Test Formulation Administration 2 (F2)	REF product	
14	8	4	0.67	
38	3.5	3	0.67	
40	4	4.5	1	
41	4.5	4	0.67	
19	1	2.5	0.67	
28	2.5	6	0.67	
29	4	4	0.67	
36	3	3.5	1.5	
01	4.5	6	1.5	

TABLE 11-continued

Subject No.	Lag times for onset of plasma profiles		
	Test Formulation Administration 1 (F1)	Test Formulation Administration 2 (F2)	REF product
03	4	4.5	1
15	3.5	3.5	1
20	4	4	0.67
04	4.5	3.5	1
08	2.5	3	0.67
10	4	3.5	1
12	4.5	5	1
06	5.5	/	/
17	5	4	1
22	4	3.5	0.33
34	3.5	4	1
09	3	4	2
11	2.5	3.5	0.67
16	4	4.5	0.67
42	3	6	1.5

### Statistical Analysis

[0461] Data were analysed with SigmaPlot for Windows, version 11.0.

### Descriptive Statistics

[0462] Median values were calculated for the lag times to onset of plasma level for each arm of the study and are presented in Table 12 below. (Median values are reported rather than mean values because these are discrete data i.e., lag times can only correspond to the sampling times in the study).

TABLE 12

Group	N	Median lag time values		
		Median	25 <sup>th</sup> percentile	75 <sup>th</sup> percentile
F1	24	4.000	3.000	4.500
F2	24	4.000	3.500	4.500
REF	24	1.000	0.670	1.000

[0463] It was concluded that the Median value (50<sup>th</sup> percentile) of the lag time to onset of plasma levels was 4 hours for both administrations of test formulation and 1 hour for the reference commercial formulation.

[0464] To determine whether the lag times observed for the test formulation were statistically different to those for the Reference product, two Repeated Measures Analysis of Variance (ANOVA) tests were applied to the data-one assuming that the data are normally distributed, and one with no assumption about how the data are distributed.

[0465] The Repeated Measures Analysis of Variance assuming a normally distributed data set resulted in an F value of 51.815 and a P value of  $P<0.001$ , which is highly statistically significant. A post-hoc comparison using a Tukey test to determine which arms of the study were different to each other gave the results presented in Table 13 below:

TABLE 13

Post-hoc comparison between arms of study				
Comparison	Difference of Means	q-statistic	P Value	P < 0.050
F2 vs. REF	3.151	18.005	<0.001	Yes
F2 vs. F1	0.261	1.517	0.820	No
F1 vs. REF	2.890	16.770	<0.001	Yes

[0466] This showed that, whereas there is no difference in the lag time to onset of plasma levels whenever the test formulation was administered on separate occasions, the lag time to onset of plasma levels following administration of test formulation is highly statistically different ( $P<0.001$ ) compared to when the commercial reference product is administered.

[0467] A Repeated Measures Analysis of Variance was also applied to rankings of the individual lag time values using the Friedman test (see e.g. Stanton, A Glanz, Primer of Biostatistics, 5th Edition, McGraw Hill 2002, ISBN 0-07-137946-0, pages 370-380).

[0468] Using this test, the Chi-square value was 52.950, resulting in a P value of <0.001, which is highly statistically significant. A post-hoc comparison using a Tukey test to determine which arms of the study were different to each other gave the results presented in Table 14 below:

TABLE 14

Post-hoc comparison between arms of study			
Comparison	Difference of Ranks	q-statistic	P < 0.05
F2 vs REF	68.500	9.237	Yes
F2 vs F1	9.500	1.281	No
F1 vs REF	59.000	7.956	Yes

[0469] This test shows (again) that, whereas there is no difference in the lag time to onset of plasma levels when test formulation is administered on separate occasions, the lag time to onset of plasma levels when test formulation is administered is statistically different ( $P<0.05$ ) to when the commercial reference formulation is administered.

[0470] In conclusion, the test formulation has a significantly different lag time to onset of plasma levels compared to the commercial reference formulation, Entocort® EC, irrespective of the type of statistical analysis that is applied. The median lag time to onset of plasma levels was 4 hours in each of the arms of the study in which F1, F2 was administered, while the median lag time for Entocort® EC was 1 hour.

[0471] This analysis clearly demonstrates that, unlike the Reference product, the test formulation as described and claimed herein does not release the bulk of its active ingredient until the distal part of the small intestine (e.g., the ileum, such as the distal ileum) is reached.

Example 17: General Process for Standard In Vitro Dissolution Test According to USP<711>/Ph.Eur. 2.9.3 in the Presence of Added Surfactant Tween 80

[0472] The in vitro dissolution of the encapsulated budesonide core-shell beads of Example 1 were analysed as described in Ph.Eur. 2.9.3 Dissolution test for solid dosage forms (using Apparatus 2) and as described in USP <711>

Dissolution (using Apparatus 2). The measurement was carried out as described below.

[0473] Three marketed budesonide-containing formulations were also analysed in this test. The three formulations being Entocort® (Tillotts Pharma), Budenofalk® (Dr Falk Pharma GmbH) and Cortiment® (Ferring Pharmaceuticals, CH).

#### Dissolution Apparatus Setup

Dissolution Apparatus:	USP <711>/Ph. Eur. 2.9.3 Apparatus 2
Vessel Size/Type:	1000 mL/clear glass, round-bottom
Rotation Speed:	100 rpm
Media Volume:	900 mL-Acid Resistance 900 mL-Buffer Dissolution Profile
Test Temperature:	37.0 ± 0.5° C.
Pull Volume:	15 mL
Replacement:	No
Sampling Time Point:	Acid Resistance Stage: 2 hours Buffer Stage: 15, 30, 45, 60, 90, 120, 180 minutes Check the pH of the dissolution medium before and in each vessel after each test and record the result (Buffer Dissolution Medium only)
pH check	

[0474] Budesonide release was measured using Ultra Performance Liquid Chromatography (UPLC).

#### Reagents and Standards

##### Standards and Reference Materials:

[0475] Budesonide, Ph. Eur. CSR or suitable secondary standard.

##### Other Reagents:

[0476] Tween 80, (Polysorbate (80)), Fisher Scientific or equivalent.

##### Dissolution Media and Diluents

##### Acid Resistance Media

[0477] 0.1 N HCl Solution. To prepare 6 L of acid resistance media, 50 mL of concentrated HCl was combined with 6000 mL of water and the resulting solution was mixed well.

##### Buffer Dissolution Media

[0478] FaSSIF Buffer Concentrate (from Biorelevant.com, product code FASBUF01) was used to prepare the Buffer solution.

[0479] A 0.05 w/v % (0.5 mg/mL) Tween 80 to the Buffer Solution: for example, To make 6 L of the Buffer Dissolution Medium, 3 g of Tween 80 was added to 6L of the Buffer solution, to arrive at a Tween 80 concentration of 0.05 w/v %.

[0480] The resulting solution was mixed well and the pH was checked. If necessary, the pH was adjusted to 6.5±0.05 using either Hydrochloric Acid or Sodium Hydroxide.

[0481] Budesonide release was assessed based on the acceptance criteria in USP<711>/Ph.Eur. 2.9.3.

**Example 18: Dissolution Profile Analysis of Budesonide Capsules According to In Vitro USP<711>/Pharmacopeia Test No. 2.9.3 in the Presence of Added Surfactant Tween 80**

[0482] The enteric coated capsules filled with cured beads (“budesonide capsules”) prepared as described in Example 1 above were tested under the dissolution conditions outlined in Example 17.

[0483] The overall average dissolution profile for three samples in the buffer stage can be seen in FIG. 19. No budesonide release was observed in the acid resistance stage at the 2 hour sampling time point.

[0484] The quantitative results of the dissolution of budesonide in the various media at time points 2 hours at pH 1.2, and 15, 30, 45, 60, 90, 120, and 180 minutes in the buffer stage at pH 6.5 are provided in the table below.

Time (min)	120	Acid Budesonide Release (%)						
		15	30	45	60	90	120	180
Vessel 1	0	0	0	0	0	56.9	85.0	97.6
Vessel 2	0	0	0	0	0	50.0	83.7	97.5
Vessel 3	0	0	0	0	0	45.8	85.0	101.4
Mean	0	0	0	0	0	50.9	84.6	98.8
Range	0-0	0-0	0-0	0-0	0-0	46-57	84-85	97-101
SD	0	0	0	0	0	5.6	0.8	2.3

\*SD = standard deviation

[0485] The budesonide release was assessed based on the acceptance criteria in USP<711>/Ph.Eur. 2.9.3.

[0486] For comparison, the dissolution profiles of budesonide capsules and three other budesonide-containing formulations were obtained according to the protocol outlined below. The three other budesonide-containing formulations being Entocort® (Tillots Pharma), Budenofalk® (Dr Falk Pharma GmbH) and Cortiment® (Ferring Pharmaceuticals, CH).

#### Method for Capsules

Dissolution Apparatus	USP <711>/Ph. Eur. 2.9.3
Apparatus:	Apparatus 2
Vessel Size/Type:	1000 mL/clear glass, round-bottom
Rotation Speed:	100 rpm
Media Volume:	900 mL-Acid Resistance 900 mL-Buffer Dissolution Profile
Test Temperature:	37.0 ± 0.5° C.
Pull Volume:	10 mL
Replacement:	No
Sampling Time Points	Acid Resistance Stage: 2 hours Buffer Stage: 15, 30, 45, 60, 90, 120, 180 minutes Check the pH of the dissolution medium after preparation, and again in each vessel after each test and record the result (Buffer Dissolution Medium only).
pH check	

#### Standards and Reference Materials

[0487] Budesonide, Ph. Eur. CSR.

#### Other Reagents:

[0488] Tween 80, (Polysorbate (80)), Fisher Scientific or equivalent.

#### Dissolution Media, Mobile Phase and Diluents

#### Acid Resistance Media

[0489] 0.1N HCl Solution. For example, to prepare 10 L, 82 mL of concentrated HCl was combined with 10000 ml of water, and mixed well.

#### Buffer Dissolution Media

[0490] FaSSIF Buffer Concentrate from Biorelevant.com was used to prepare the Buffer solution according to the instructions provided. The resulting solution was mixed well. The pH of the buffer solution was checked after preparation. If necessary, the pH was adjusted to 6.5±0.05 using either Hydrochloric Acid or Sodium Hydroxide.

[0491] After Acid Resistance samples were pulled, the capsules were taken out of the solution with pincers and placed aside while the vessels are emptied, cleaned and filled with preheated buffer medium. 0.05 w/v % Tween 80 (or equivalent) was added to each dissolution vessel; for example 450 mg of Tween 80 was added to the dissolution vessels after they have been filled with 900 mL of pre-heated buffer, to arrive at a surfactant concentration of 0.05%.

[0492] After all vessels reached the target temperature, the experiment was started by adding a capsule to each vessel.

#### Modification of Procedure for Budenofalk

[0493] A different procedure was used when the capsule broke during the acid stage. Most of the acid phase was carefully decanted and then the remaining acid carefully removed with a pipette in order to remove as few pellets from the vessel as possible. The buffer stage was started by adding 900 mL of pre-heated buffer medium, followed by addition of Tween.

#### Sampling for Both Stages

[0494] 10 mL aspirated, 8 mL discarded (through Whatman filter), 1 mL sampled into HPLC vial.

[0495] For the avoidance of doubt, the variations in this test compared to Example 2 do not have an effect on the overall dissolution profiles of the products tested.

[0496] FIG. 20 shows the dissolution profile of the budesonide capsules as compared to the three other budesonide-containing formulations. From this figure it can be clearly seen that within the FaSSIF medium that mimics the environment in the small intestine, the budesonide capsules have a release profile that is distinguished from all other marketed budesonide-containing formulations.

#### Example 19: Dissolution Profile Analysis of Core-Shell Beads in the Absence of Enteric Coated Capsules According to In Vitro USP<711>/Pharmacopeia Test No. 2.9.3

[0497] The cured core-shell beads as prepared in Example 1 in the absence of an enteric coated capsule were also tested under the buffer stage dissolution conditions (only) as outlined in Example 17.

**[0498]** The overall average dissolution profile for three samples in the buffer stage can be seen in FIG. 21.

**[0499]** The quantitative results of the dissolution of budesonide from the core-shell beads in the buffer stage at pH 6.5 at time points 15, 30, 45, 60, 90, 120, and 180 minutes are provided in the table below.

Time (min)	Buffer Stage Budesonide Release (%)						
	15	30	45	60	90	120	180
Vessel 1	43.7	72.0	81.5	88.9	96.6	99.9	101.6
Vessel 2	29.3	64.9	78.0	84.9	93.1	96.8	99.1
Vessel 3	47.6	68.2	79.6	86.4	93.9	97.7	99.9
Mean	40.2	68.3	79.7	86.7	94.5	98.1	100.2
Range	29-48	65-72	78-82	85-89	93-97	97-100	99-102
SD	9.7	3.6	1.8	2.0	1.8	1.6	1.3

\*SD = standard deviation

**[0500]** The release profile of the core-shell beads in the absence of enteric coated capsules in the FaSSIF Buffer Concentrate further confirms that the bulk of the budesonide will be released in vivo to the ileum. That is to say, the majority of the release from the beads occurs over a 90 minute window and combined with the delay in release from the enteric coating to arrive at the desired dissolution profile of the entire formulation, the bulk of the budesonide will be released in vivo to the ileum as confirmed by the biomarker data obtained and the modelling results provided below in Example 25.

#### Example 20: General Process for Standard In Vitro Dissolution Test According to USP<711>/Ph.Eur. 2.9.3 in the Absence of Added Surfactant Tween 80

**[0501]** The in vitro dissolution of the encapsulated budesonide core-shell beads of Example 1 were analysed as described in Ph.Eur. 2.9.3 Dissolution test for solid dosage forms (using Apparatus 2) and as described in USP <711> Dissolution (using Apparatus 2). The measurement was carried out as described below.

**[0502]** The three marketed budesonide-containing formulations were also analysed in this test. The three formulations being Entocort® (Tillotts Pharma), Budenofalk® (Dr Falk Pharma GmbH) and Cortiment® (Ferring Pharmaceuticals, CH).

#### Dissolution Apparatus Setup

Dissolution Apparatus:	USP <711>/Ph. Eur. 2.9.3 Apparatus 2
Vessel Size/Type:	1000 mL/clear glass, round-bottom
Rotation Speed:	100 rpm
Media Volume:	900 mL-Acid Resistance 900 mL-Buffer Dissolution Profile
Test Temperature:	37.0 ± 0.5° C.
Pull Volume:	10 mL
Replacement:	No
Sampling Time Point:	Acid Resistance Stage: 2 hours Buffer Stage: 15, 30, 45, 60, 90, 120, 180 minutes Check the pH of the dissolution medium after preparation, and again in each vessel after each test and record the result (Buffer Dissolution Medium only)
pH check	

**[0503]** Budesonide release was measured using Ultra Performance Liquid Chromatography (UPLC).

#### Reagents and Standards

##### Standards and Reference Materials:

**[0504]** Budesonide, Ph. Eur. CSR or suitable secondary standard.

#### Dissolution Media and Diluents

##### Acid Resistance Media

**[0505]** 0.1 N HCl Solution. To prepare 10 L of acid resistance media, 82 mL of concentrated HCl was combined with 10000 mL of water and the resulting solution was mixed well.

##### Buffer Dissolution Media

**[0506]** FaSSIF Buffer Concentrate from Biorelevant.com was used to prepare the Buffer solution according to the instructions provided. The resulting solution was mixed well. The pH of the buffer solution was checked after preparation. If necessary, the pH was adjusted to 6.5±0.05 using either Hydrochloric Acid or Sodium Hydroxide.

**[0507]** After Acid Resistance samples were pulled, the capsules were taken out of the solution with pincers and placed aside while the vessels were emptied, cleaned and filled with 900 mL of pre-heated buffer medium.

**[0508]** After all vessels reached the target temperature, the experiment was started by adding a capsule to each vessel.

#### Modification of Procedure for Budenofalk

**[0509]** A different procedure was used when the capsule broke during the acid stage. Most of the acid phase was carefully decanted and then the remaining acid carefully removed with a pipette in order to remove as few pellets from the vessel as possible. The buffer stage was started by adding 900 mL of pre-heated buffer medium.

#### Sampling for Both Stages

**[0510]** 10 mL aspirated, 8 mL discarded (through Whatman filter), 1 mL sampled into HPLC vial.

**[0511]** Budesonide release was assessed based on the acceptance criteria in USP<711>/Ph.Eur. 2.9.3.

#### Example 21: Dissolution Profile Analysis of Budesonide Capsules According to In Vitro USP<711>/Pharmacopeia Test No. 2.9.3 in the Absence of Added Surfactant Tween 80

**[0512]** The enteric coated capsules filled with cured beads (“budesonide capsules”) prepared as described in Example 1 above were tested under the dissolution conditions outlined in Example 20.

**[0513]** The overall average dissolution profile for the budesonide capsules can be seen in FIG. 22. FIG. 22 also contains the dissolution profiles of three other budesonide-containing formulations under the same test as a comparative. From this figure it can be clearly seen that within the FaSSIF medium that mimics the environment in the small intestine, the budesonide capsules have a release profile that is distinguished from all other marketed budesonide-containing formulations.

**[0514]** For the budesonide capsules and Cortiment, no budesonide release was observed in the acid resistance stage at the 2 hour sampling time point. For Entocort budesonide release amounting to 0.8% was observed, and for Budeno-

small budesonide release amounting to 0.6% was observed in the acid resistance stage at the 2 hour sampling time point.

**[0515]** The quantitative results of the dissolution of budesonide in the various media at time points 2 hours at pH 1.2, and 15, 30, 45, 60, 90, 120, and 180 minutes in the buffer stage at pH 6.5 are provided in the table below for both the budesonide capsules and the three comparative formulations.

Time (min)	120	Buffer Stage Budesonide Release (%)						
		15	30	45	60	90	120	180
Budesonide Capsules								
Mean	0.0	0.0	0.0	0.0	0.0	19.4	77.7	100.1
Range	0-0	0-0	0-0	0-0	0-0	1-44	72-87	98-103
SD	0.0	0.0	0.0	0.0	0.0	14.9	5.5	1.8
Entocort								
Mean	0.8	34.2	55.4	69.4	78.4	89.3	95.5	102.8
Range	1-1	32-37	53-57	66-71	75-82	85-92	91-98	96-109
SD	0.3	1.5	1.7	2.0	2.1	2.5	2.7	4.1
Budenofalk								
Mean	0.6	0.6	0.6	0.6	0.6	0.8	0.8	1.9
Range	0-1	0-1	0-1	0-1	0-1	1-1	1-1	2-3
SD	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.4
Cortiment								
Mean	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5
Range	0-0	0-0	0-0	0-0	0-0	0-0	0-0	0-1
SD	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2

\*SD = standard deviation

**[0516]** The budesonide release was assessed based on the acceptance criteria in USP<711>/Ph.Eur. 2.9.3.

**[0517]** The table below shows the f2 values for comparison of the budesonide capsules according to the invention tested under this method with each of the other commercially available products. An f2 value of 50 or greater is required to demonstrate similarity of the profiles (FDA SUPAC Guidance 1995, 1997).

Nefcon	Entocort EC	Budenofalk	Cortiment
F2 value (biorelevant)	11.7	16.1	15.8

**[0518]** It is clear that the release profiles of budesonide differ widely among the four commercial products. None of the f2 comparisons between Nefcon and the other products demonstrated similarity, which would require an f2 value of 50 or greater. In fact, based on the f2 evaluation as well as visual inspection of the graphical profiles, their release profiles must be considered strongly dissimilar.

#### Example 22: First Tablet Formulation

**[0519]** Tablets are manufactured via the following process steps:

**[0520]** 1. Wet granulation. Budesonide, mannitol, hydroxyethyl cellulose, hydroxypropyl cellulose and sodium starch glycolate are blended. An ethanol-water mixture is thereafter sprayed on the powder during blending. The resulting granules are thereafter dried.

**[0521]** 2. Final blending. The dried granules are blended with sodium stearyl fumarate.

**[0522]** 3. Tableting. Tablets are compressed using a tableting machine.

**[0523]** 4. Enteric coating. Methacrylic acid and methyl methacrylate copolymers, talc and dibutyl sebacate are dispersed in an isopropyl alcohol-water mixture, during mixing. The coating dispersion is thereafter sprayed onto the tablets using a fluid bed apparatus.

**[0524]** An example of a tablet composition according to the present invention is shown in the table below.

#### Tablet Composition.

Component	Amount (mg)	Function
Budesonide	4.00	API (intragrangular)
Mannitol	81.50	Filler (intragrangular)
Hydroxyethyl cellulose	5.00	Binder (intragrangular)
Hydroxypropyl cellulose	5.00	Binder (intragrangular)
Sodium starch glycolate	4.00	Disintegrant (intragrangular)
Purified water	*	Wet granulation solvent
Ethanol	*	Wet granulation solvent
Sodium stearyl fumarate	0.50	Lubricant (extragrangular)
Total (uncoated tablet)	100.00 mg	
Methacrylic acid and methyl methacrylate copolymer (1:1)	9.85	Enteric coating polymer
Methacrylic acid and methyl methacrylate copolymer (1:2)	3.31	Enteric coating polymer
Talc	3.31	Glidant, coating uniformity
Dibutyl sebacate	2.55	Plasticizer
Isopropyl alcohol	*	Coating solvent
Purified water	*	Coating solvent
Total (coated tablet)	119.02 mg	

\*Removed during process.

#### Example 23: Second Tablet Formulation

**[0525]** Tablets are manufactured via the following process steps:

**[0526]** 1. Wet granulation. Budesonide is initially blended with colloidal silica. Microcrystalline cellulose and dibasic calcium phosphate are thereafter added followed by additional blending. A solution of hydroxypropyl cellulose in an ethanol-water mixture is thereafter sprayed onto the powder during blending. The resulting granules are thereafter dried.

**[0527]** 2. Blending. The dried granules are blended with microcrystalline cellulose, sodium starch glycolate and Copovidone. As a final blending step, magnesium stearate is added to the blend, followed by final blending.

**[0528]** 3. Tableting. Tablets are compressed using a tableting machine.

**[0529]** 4. Enteric coating. Methacrylic acid and methyl methacrylate copolymers, talc and dibutyl sebacate are dispersed in an isopropyl alcohol-water mixture, during mixing. The coating dispersion is thereafter sprayed onto the tablets using a pan coater.

[0530] An example of a tablet composition according to the present invention is shown in the table below.

#### Tablet Composition.

Component	Amount (mg)	Function
Budesonide, micronized	4.00	API (intragranular)
Colloidal silica	0.50	Glidant (intragranular)
Dibasic calcium phosphate	16.00	Filler (intragranular)
Microcrystalline cellulose	15.50	Filler (intragranular)
Hydroxypropyl cellulose	4.00	Binder (intragranular)
Ethanol	*	Wet granulation solvent
Purified water	*	Wet granulation solvent
Microcrystalline cellulose	53.50	Filler (extragranular)
Copovidone	5.00	Binder (extragranular)
Sodium starch glycolate	1.00	Disintegrant (extragranular)
Magnesium stearate	0.50	Lubricant (extragranular)
Total (uncoated tablet)	100.00 mg	
Methacrylic acid and methyl methacrylate copolymer (1:1)	9.85	Enteric coating polymer
Methacrylic acid and methyl methacrylate copolymer (1:2)	3.31	Enteric coating polymer
Talc	3.31	Glidant, coating uniformity
Dibutyl sebacate	2.55	Plasticizer
Isopropyl alcohol	*	Coating solvent
Purified water	*	Coating solvent
Total (coated tablet)	119.02 mg	

\*Removed during process.

#### Example 24: Third Tablet Formulation

#### Tablet Components:

Component	Amount (mg)	Function
Budesonide, micronized	4	Active Pharmaceutical Ingredient (API)
Microcrystalline cellulose	61.5	Filler
Calcium hydrogen phosphate dihydrate	10	Filler
Hypromellose, 100 mPas	10	Filler/gelling agent
Hypromellose, 4000 mPas	10	Filler/gelling agent
Crospovidone	4	Disintegrant
Purified water	*	Wet granulation solvent
Sodium stearyl fumarate	0.5	Lubricant
Total, uncoated tablet	100	
Methacrylic acid-methyl methacrylate copolymer (1:1)	8.7	Enteric coating polymer
Methacrylic acid-methyl methacrylate copolymer (1:2)	2.9	Enteric coating polymer
Talc	2.9	Glidant, coating uniformity
Dibutyl sebacate	2.2	Plasticizer
Isopropyl alcohol	*	Coating solvent
Purified water	*	Coating solvent
Total, coated tablet	117	

\*Removed during manufacturing.

#### Manufacturing of Uncoated Tablets (200 g Batch Size)

[0531] 1. Budesonide and all excipients except Sodium stearyl fumarate were blended with a Turbula mixer (blending at 46 rpm for 75 min).

[0532] 2. The powder blend was granulated by spraying water on the powder during blending. The amount of water sprayed was 15% of the weight of the dry powder.

[0533] 3. The granules were dried over night at 50° C.

[0534] 4. The dried granules were mixed with sodium stearyl fumarate (lubricant) for 10 min at 46 rpm, using a Turbula mixer.

[0535] 5. Tablets were compressed, to an uncoated tablet weight of 100 mg.

#### Preparation of Coating Dispersion

[0536] A coating dispersion was prepared with the composition shown in the table below. The dispersion was prepared according to the following steps:

[0537] 1. A diluent mixture was prepared by blending isopropyl alcohol, water and dibutyl sebacate in a vessel (Vessel A).

[0538] 2. A Eudragit suspension was prepared by transferring approximately half of the diluent mixture (in Vessel A) to another vessel (Vessel B). Eudragit L100 and Eudragit S100 were thereafter added slowly to Vessel B during mixing. The Eudragit suspension in Vessel B was thereafter mixed for another 30-60 min.

[0539] 3. Talc was added slowly to the remaining diluent mixture in Vessel A, during mixing with a high shear mixer. The talc suspension was thereafter mixed for an additional 10 min with high shear mixer.

[0540] 4. The talc suspension in Vessel A was thereafter slowly poured into the Eudragit suspension (Vessel B) during mixing.

[0541] 5. The mixture in Vessel B was thereafter stirred for 24 h at room temperature.

[0542] 6. The mixture was passed through a 0.5 mm sieve.

[0543] 7. The final coating dispersion was thereafter stored at room temperature under continuous stirring until and during coating.

Commodity name	Amount (g)	Supplier
Eudragit L100 (Methacrylic acid copolymer type A)	39.4	Evonik
Eudragit S100 (Methacrylic acid copolymer type B)	13.22	Evonik
Talc, Pharma M grade	13.22	Imerys
Dibutyl sebacate USP/NF	10.18	Merck
Isopropyl alcohol	594.36	—
Water purified	20.44	—
Total	690.82	

#### Coating of Tablets (50 g Batch Size)

[0544] Tablets were coated using a stainless-steel pan coater with 12 cm diameter, and a standard spray nozzle. Coating weight gain was determined by weighing tablet samples after different coating durations. The following spraying parameters were used:

[0545] Rotation speed of coating pan: 20-30 rpm

[0546] Product temperature: 24-27° C.

[0547] Pump flow: 250-260 µl/min

[0548] Spray nozzle air pressure: 0.25 bar

## In Vitro Dissolution of Tablets

**[0549]** The in vitro dissolution profiles of the tablets were analysed according to the protocol outlined in Example 6 above (Comparative test according to USP<711>/Ph.Eur. 2.9.3 in the absence of surfactant in the phosphate buffer stage and at a paddle rotation speed of 100 rpm) and the protocol outlined in Example 20 above (General Process For Standard in vitro dissolution test according to USP<711>/Ph.Eur. 2.9.3 in FaSSIF buffer in the absence of added surfactant Tween 80 at 100 rpm). The dissolution values at certain time points can be seen below.

		Dissolution (%)				
		HCl 2 h	buffer 15 min	buffer 30 min	buffer 60 min	buffer 120 min
Phosphate buffer, pH 6.8	Tablet 1	0	0	0	14.3	75.8
	Tablet 2	0	0	0	44.9	87.0
	Average	0	0	0	29.6	81.4
FaSSIF	Tablet 1	0	0	0	36.1	75.7
	Tablet 2	0	0	0	30.3	72.2
	Average	0	0	0	33.2	73.9

**[0550]** The budesonide release from these tablets matches the dissolution profile required to achieve the majority of release to the ileum. What is particularly surprising that the combination of a gelling agent (hypromellose) with a disintegrant (crospovidone) allowed the achievement of the correct in vitro dissolution profile. Although it is expected that other tablet formulations are possible to achieve the desired release profile.

**Example 25:** In Silico Modelling of Site of Release of Budesonide Capsules

**[0551]** A physiologically based pharmacokinetic (PBPK) model of the capsules prepared in Example 1 (“budesonide capsules” or “nefcon budesonide”) was run using GastroPlus® software (Simulations Plus, CA; version 9.8.3002).

**[0552]** The in vitro release of the budesonide capsules with an acid stage (first 2 hours) followed by a buffer stage according to the protocol outlined in Example 2 was uploaded in the PBPK software. The IVIVC (in vitro in vivo correlation) was obtained to correlate the PBPK model prediction with measured pharmacokinetic results. The predicted Cmax was faster and higher than observed. However, Budesonide is known to undergo gut wall metabolism in the small intestine (Seidegård J et. al., Presystemic elimination of budesonide in man when administered locally at different levels in the gut, with and without local inhibition by ketoconazole. Eur J Pharm Sci. 2008 Nov. 15; 35 (4): 264-70; Raje et al. Evaluation of separate role of intestine and liver in first pass metabolism of budesonide in rat Xenobiotica. 2018 December; 48 (12): 1206-1214). After introducing gut wall metabolism to the PBPK model a good fit of the data was achieved.

**[0553]** As can be seen in FIG. 23(a), the IVIVC PBPK model confirmed that the budesonide capsules began releasing budesonide only once the ileum was reached and at least about 90% of the budesonide was released throughout the entire ileum, with the small remainder being released in the section of the intestines following the ileum (i.e. the caecum). Therefore, this model confirms that a composition having an in vitro release profile as defined by the invention achieves release of the budesonide payload to the site of

highest concentration of Peyer’s patches in the intestinal tract, being the ileum and, therefore, any composition fitting this release profile will be effective in treating IgA nephropathy.

**[0554]** Release of budesonide from Entocort® (Tillotts Pharma) was also modelled using the GastroPlus® software and the results showed that all budesonide was released prior to the formulation entering the ileum section of the small intestines (see FIG. 23(b)).

**Example 26:** Study on In Vivo Release of Capsule Contents

**[0555]** A study was conducted to evaluate where and when capsules coated with the same enteric coating as the capsules described in Example 1 release their contents in the gastrointestinal tract.

**[0556]** In this study, VCaps plus size 1 HPMC capsules were filled with 75 mg of caffeine, 10 mg black iron oxide, and 87.5 mg manganese gluconate dehydrate. 140.9 mg of sugar beads (also called pellets) were added to replicate the total weight of the cores in the capsules described in Example 1. The capsules were then coated with the same enteric coating, using the same coating process and the same equipment and facilities, as the capsules described in Example 1.

**[0557]** The caffeine was placed inside the capsules to use as a marker to determine when the contents of the capsule were released, by measuring appearance of caffeine in the saliva at various time points. Caffeine is rapidly absorbed after release in the intestinal tract, and as such, the appearance of caffeine in saliva provides a sensitive marker of capsule opening (Sager et al. Low dose caffeine as a salivary tracer for the determination of gastric water emptying in fed and fasted state: A MRI validation study. Eur J Pharm Biopharm 127:443-452 (2018)). The iron oxide was placed inside the capsule so that magnetic resonance imaging (MRI) could be used to locate the position of the capsule as it moves through the gastrointestinal tract by visualizing the iron oxide.

**[0558]** The study was performed at the University Clinic in Greifswald, Germany, as an open-label, single-center study in 12 healthy young human subjects. The subjects stopped ingesting caffeine-containing foods and drinks for three days prior to their participation in the study and fasted overnight prior to the study for at least 10 hours.

**[0559]** An MRI scan was performed, and a saliva sample obtained, prior to each participant’s ingestion of the enteric coated capsule with a glass of water. Subsequently, MRI scans were performed every 15 minutes for the first four hours and thereafter every 30 minutes until the study was completed. Saliva samples were obtained one minute after each MRI scan. MRI imaging was conducted with a Siemens MAGNETOM Avanto MR-scanner (Siemens Healthcare, Erlangen, Germany) with a field strength of 1.5 Tesla and imaging data were analyzed using Horos 2.2.0 (The Horos Project). All measurements were performed with the subject in the supine position (lying on their back, head up). Saliva samples were analyzed using a LCMS 8060 system (Shimadzu Corporation, Kyoto, Japan) and appropriately prepared for this purpose.

**[0560]** FIG. 24 shows MRI images for different locations within the gastrointestinal tract. At 15 and 90 minutes the capsule is intact and located in the stomach and jejunum, respectively, while at 270 minutes the capsule has released its contents and the iron oxide has dispersed in the ileum.

**[0561]** The average gastric emptying time of the capsules (time at which the capsule moved out of the stomach) was

$58 \pm 30$  min (highest observed gastric emptying time was 112.5 min). These values are consistent with the usual gastric emptying times for large, non-disintegrating dosage forms (Wilson et al. Chapter 3. Gastrointestinal Transit and Drug Absorption (pages 41 to 65) In "Oral Drug Absorption: Prediction and Assessment" Edited by J. Dressman and C. Reppas—2nd. Edition Drug and the Pharmaceutical Sciences Vol. 193 Marcel Dekker, NY, NY ISBN-13: 978-1-4200-7733-9 (2010)). None of the capsules showed disintegration in the stomach.

[0562] The first time point with a measured concentration  $\geq 10$  ng/ml of caffeine in undiluted saliva was considered as the "salivary caffeine appearance". As shown in the table below, the average time to the first appearance of caffeine in saliva was 238 minutes after capsule ingestion, with a standard deviation of 47 minutes and a range of 158-345 minutes. After subtracting the individual's gastric emptying time from the time at which caffeine first appeared in their saliva, the time for release of caffeine after entering the small intestine was calculated as averaging 181+31 minutes (range 120-233 minutes). These values indicate that opening of the enteric coated capsule and release of caffeine fell well within the usual range of small intestinal transit times (3.5 to 4.5 hours (210 to 270 minutes)) (Wilson et al. Chapter 3. Gastrointestinal Transit and Drug Absorption (pages 41 to 65) In "Oral Drug Absorption: Prediction and Assessment" Edited by J. Dressman and C. Reppas—2nd. Edition Drug and the Pharmaceutical Sciences Vol. 193 Marcel Dekker, NY, NY ISBN-13: 978-1-4200-7733-9 (2010)).

[0563] The table below shows the individual and average results of gastric emptying time (as determined by MRI), time of salivary caffeine appearance and time of salivary caffeine appearance after gastric emptying.

Subject	Gastric Emptying Time (min)	Time to Salivary Caffeine Appearance after ingestion (min)	Time to Salivary Caffeine Appearance after gastric emptying (min)
001	53	203	150
002	83	255	173
003	68	255	188
004	23	218	195
005	38	218	180
006	113	345	233
007	8	218	210
008	53	218	165
009	68	285	218
010	53	233	180
011	98	255	158
012	38	158	120
mean	58	238	181
SD	30	47	31

[0564] Based on the first time of appearance of caffeine in the saliva and the location of the iron oxide in the corresponding MRI image, the enteric coated capsules opened and released their contents in the ileum in 10 of the 12 subjects. The study thus confirms that the enteric coating and capsule described in Example 1 consistently results in release of the capsule's contents to the ileum. The beads contained in Example 1 would start releasing budesonide after the capsule opens, with the majority of release occurring within one hour. Comparing the average small intestinal transit time for beads of 3.5 to 4.5 hours (Wilson et al. Chapter 3. Gastrointestinal Transit and Drug Absorption

(pages 41 to 65) In "Oral Drug Absorption: Prediction and Assessment" Edited by J. Dressman and C. Reppas—2nd. Edition Drug and the Pharmaceutical Sciences Vol. 193 Marcel Dekker, NY, NY ISBN-13: 978-1-4200-7733-9 (2010)) with the average time for the capsules to open once they reach the small intestine (181 minutes in this study), plus the time required for release of the majority of the budesonide from the beads (about one hour), it can be concluded that the majority of the budesonide would be released from the beads to the distal ileum and thus be targeted to the Peyer's Patches located there.

1. (canceled)
2. A pharmaceutical composition comprising:  
a plurality of cores comprising budesonide encapsulated within a capsule, wherein the plurality of cores are each coated with an extended release pharmaceutically-acceptable polymeric blend comprising a water-insoluble polymer having a solubility in water (at 25° C.) of less than about 0.1 mg/mL and a water-soluble polymer having a solubility in water (at 25° C.) of at least about 10 mg/mL;  
wherein the extended release pharmaceutically-acceptable polymeric blend is cured and is present in an amount of from 5 wt. % to about 18 wt. % of the total coated core weight;  
wherein the pharmaceutical composition further comprises an enteric coating.
3. The pharmaceutical composition according to claim 2, wherein the pharmaceutical composition meets the following release profile in a standard in vitro USP<711> dissolution test using a dissolution apparatus according to Apparatus 2 (Paddle Apparatus) at a paddle rotation speed of 100 rpm:
  - a) no more than about 10% of the budesonide is released into an aqueous dissolution medium with a pH of about 1.2 within about 120 minutes;
  - b) no more than about 10% of the budesonide is released into a pharmaceutically-relevant dissolution medium within about 30 minutes, wherein the pharmaceutically-relevant dissolution medium is a Level 1 Fasted State Simulated Intestinal Fluid at a pH of about 6.5, or a phosphate buffer medium at a pH of about 6.8; and
  - c) at least about 70% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 120 minutes.
4. The pharmaceutical composition according to claim 2, wherein the combination of the enteric coating and the pharmaceutically-acceptable polymeric blend in the extended-release component result in the pharmaceutical composition releasing budesonide substantially to the ileum region of the small intestine to reduce the formation by mucosal B cells in Peyer's patches in the ileum of secretory galactose-deficient IgA antibodies that drive immune complex formation in IgA nephropathy, to treat said IgA nephropathy, and limiting systemic exposure of said budesonide.

5. The pharmaceutical composition according to claim 2, wherein the water-insoluble polymer is present in an amount of from about 45 wt. % to about 90 wt. % of the extended release pharmaceutically-acceptable polymeric blend and the water-soluble polymer is present in an amount of from about 35 wt. % to about 5 wt. % of the extended release pharmaceutically-acceptable polymeric blend.

6. The pharmaceutical composition according to claim 2, wherein the water-insoluble polymer is present in an amount of from about 45 wt. % to about 65 wt. % of the extended release pharmaceutically-acceptable polymeric blend and the water-soluble polymer is present in an amount of from

about 35 wt. % to about 15 wt. % of the extended release pharmaceutically-acceptable polymeric blend.

**7.** The pharmaceutical composition according to claim 2, wherein the water-insoluble polymer is present in an amount of from about 47 wt. % to about 56 wt. % of the extended release pharmaceutically-acceptable polymeric blend and the water-soluble polymer is present in an amount of from about 32 wt. % to about 22 wt. % of the extended release pharmaceutically-acceptable polymeric blend.

**8.** The pharmaceutical composition according to claim 2, wherein the extended release pharmaceutically-acceptable polymeric blend is present in an amount of from about 6 wt. % to about 16 wt. % of the total coated core weight.

**9.** The pharmaceutical composition according to claim 2, wherein the extended release pharmaceutically-acceptable polymeric blend is present in an amount of from about 6 wt. % to about 12 wt. % of the total coated core weight.

**10.** The pharmaceutical composition according to claim 2, wherein the water-insoluble polymer is an alkyl cellulose.

**11.** The pharmaceutical composition according to claim 10, wherein the alkyl cellulose is an ethyl cellulose.

**12.** The pharmaceutical composition according to claim 2, wherein the water-soluble polymer is selected from polyethylene glycol (PEG), hydroxypropylmethyl cellulose (HPMC), and hydroxypropyl cellulose (HPC).

**13.** The pharmaceutical composition according to claim 2, wherein the water-insoluble polymer is an alkyl cellulose and the water-soluble polymer is selected from polyethylene glycol (PEG), hydroxypropylmethyl cellulose (HPMC), and hydroxypropyl cellulose (HPC).

**14.** The pharmaceutical composition according to claim 2, wherein the enteric coating is on the capsule.

**15.** The pharmaceutical composition according to claim 14, where the enteric coating is present in an amount of from about 34 mg to about 46 mg per capsule.

**16.** The pharmaceutical composition according to claim 14, where the enteric coating is present in an amount of from about 34 mg to about 42 mg per capsule.

**17.** The pharmaceutical composition according to claim 14, where the enteric coating is present in an amount of from about 36 mg to about 40 mg per capsule.

**18.** The pharmaceutical composition according to claim 2, wherein the capsule is a size 1 capsule.

**19.** The pharmaceutical composition according to claim 2, wherein the capsule comprises about 4 mg of budesonide.

**20.** The pharmaceutical composition according to claim 2, wherein the pharmaceutical composition meets the following release profile in a standard in vitro USP<711> dissolution test using a dissolution apparatus according to Apparatus 2 (Paddle Apparatus) at a paddle rotation speed of 100 rpm:

- a) no more than about 10% of the budesonide is released into an aqueous dissolution medium with a pH of about 1.2 within about 120 minutes;
- b) no more than about 10% of the budesonide is released into a pharmaceutically-relevant dissolution medium within about 30 minutes, wherein the pharmaceutically-relevant dissolution medium is a Level 1 Fasted State Simulated Intestinal Fluid at a pH of about 6.5, or a phosphate buffer medium at a pH of about 6.8; and
- c) at least about 70% of the budesonide is released into the pharmaceutically-relevant dissolution medium within about 120 minutes.

**21.** The pharmaceutical composition according to claim 20, wherein in criterion a) the release into the aqueous dissolution medium with a pH of about 1.2 is assessed according to the acceptance criteria in Acceptance Table 2 and/or Acceptance Table 3 of USP<711>.

**22.** The pharmaceutical composition according to claim 20, wherein in criterion b) the release into the pharmaceutically-relevant dissolution medium is assessed according to the acceptance criteria in Acceptance Table 2 and/or Acceptance Table 3 of USP<711>.

**23.** The pharmaceutical composition according to claim 20, wherein in criterion c) the release into the pharmaceutically-relevant dissolution medium is assessed according to the acceptance criteria in Acceptance Table 2 and/or Acceptance Table 4 of USP<711>.

**24.** The pharmaceutical composition according to claim 20, wherein the temperature of the dissolution media in criteria a), b) and c) is held at a temperature of about 37° C. $\pm$ 0.5° C.

**25.** The pharmaceutical composition according to claim 20, wherein the number of pharmaceutical compositions tested is 6.

**26.** The pharmaceutical composition according to claim 20, wherein at each time point in criteria (a), (b) and (c) the amount of volume withdrawn from the dissolution medium is 10 mL or 15 mL.

\* \* \* \* \*