

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12404334
Kind Code	B2
Date of Patent	September 02, 2025
Inventor(s)	Harwin; Peter Evan et al.

Methods of treating gastrointestinal inflammatory disease

Abstract

Provided herein are variant $\alpha 4\beta 7$ integrin antibodies and methods of use.

Inventors: Harwin; Peter Evan (Dorado, PR), Kiselak; Tomas (Dorado, PR), Shaheen; Hussam Hisham (Auburn, NH), Zhu; Eric Franklin (Cambridge, MA)

Applicant: Paragon Therapeutics, Inc. (Waltham, MA)

Family ID: 91664105

Assignee: Paragon Therapeutics, Inc. (Waltham, MA)

Appl. No.: 19/182382

Filed: April 17, 2025

Prior Publication Data

Document Identifier	Publication Date
US 20250243280 A1	Jul. 31, 2025

Related U.S. Application Data

continuation parent-doc US 18774123 20240716 US 12304956 child-doc US 19182382
continuation parent-doc WO PCT/US2024/031569 20240530 PENDING child-doc US 18774123
us-provisional-application US 63559081 20240228
us-provisional-application US 63554886 20240216
us-provisional-application US 63599922 20231116
us-provisional-application US 63505962 20230602
us-provisional-application US 63504966 20230530

Publication Classification

Int. Cl.: C07K16/28 (20060101); **A61K9/00** (20060101); **A61K47/18** (20170101); **A61K47/26** (20060101); **A61P1/00** (20060101); A61K39/00 (20060101)

U.S. Cl.:

CPC C07K16/2839 (20130101); **A61K9/0019** (20130101); **A61K47/183** (20130101); **A61K47/26** (20130101); **A61P1/00** (20180101); A61K2039/505 (20130101); A61K2039/545 (20130101)

Field of Classification Search

USPC: None

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
5811523	12/1997	Trinchieri et al.	N/A	N/A
5840299	12/1997	Bendig et al.	N/A	N/A
7083784	12/2005	Dall'Acqua et al.	N/A	N/A
7090845	12/2005	Fong et al.	N/A	N/A
7147851	12/2005	Ponath et al.	N/A	N/A
7211252	12/2006	Mundy et al.	N/A	N/A
7402410	12/2007	Ponath et al.	N/A	N/A
7435802	12/2007	Bendig et al.	N/A	N/A
7528236	12/2008	Fong et al.	N/A	N/A
7682613	12/2009	Fabene et al.	N/A	N/A
7687605	12/2009	Fong et al.	N/A	N/A
7803904	12/2009	Briskin et al.	N/A	N/A
8124082	12/2011	Fong et al.	N/A	N/A
8246958	12/2011	Bendig et al.	N/A	N/A
8444981	12/2012	Hsu et al.	N/A	N/A
8449489	12/2012	Thorn et al.	N/A	N/A
8454961	12/2012	Hsu et al.	N/A	N/A
8454962	12/2012	Hsu et al.	N/A	N/A
8722019	12/2013	Jefferies et al.	N/A	N/A
8871490	12/2013	Hsu et al.	N/A	N/A
8962804	12/2014	Williams et al.	N/A	N/A
9028815	12/2014	Stavnhagen et al.	N/A	N/A
9040295	12/2014	Kon et al.	N/A	N/A
9193790	12/2014	Arthos et al.	N/A	N/A
9364567	12/2015	Vitalis et al.	N/A	N/A
9441041	12/2015	Arthos et al.	N/A	N/A
9499620	12/2015	Hsu et al.	N/A	N/A
9662490	12/2016	Tracey et al.	N/A	N/A
9663579	12/2016	Fox et al.	N/A	N/A
9678071	12/2016	Escalante et al.	N/A	N/A
9717453	12/2016	Cadavid et al.	N/A	N/A
9764033	12/2016	Diluzio et al.	N/A	N/A

9821049	12/2016	Richardson et al.	N/A	N/A
9873742	12/2017	Keir et al.	N/A	N/A
9896509	12/2017	Arthos et al.	N/A	N/A
9914779	12/2017	Borie et al.	N/A	N/A
9976166	12/2017	Schellenberger et al.	N/A	N/A
10004808	12/2017	Fox et al.	N/A	N/A
10040855	12/2017	Diluzio	N/A	A61K 47/183
10143752	12/2017	Fox et al.	N/A	N/A
10221251	12/2018	Humphreys et al.	N/A	N/A
10273542	12/2018	Hackney et al.	N/A	N/A
10324088	12/2018	Singh et al.	N/A	N/A
10519228	12/2018	Srinivasan et al.	N/A	N/A
10532051	12/2019	Ebsworth et al.	N/A	N/A
10654931	12/2019	Hosen et al.	N/A	N/A
10669587	12/2019	Hackney et al.	N/A	N/A
10844416	12/2019	Tescione et al.	N/A	N/A
10918716	12/2020	Lissoos et al.	N/A	N/A
10947254	12/2020	Hudson et al.	N/A	N/A
10981965	12/2020	Mumm	N/A	N/A
10981966	12/2020	Mumm	N/A	N/A
10988540	12/2020	Hosen et al.	N/A	N/A
11045469	12/2020	Campbell et al.	N/A	N/A
11053303	12/2020	Lebert et al.	N/A	N/A
11059911	12/2020	Humphreys et al.	N/A	N/A
11091551	12/2020	Keir et al.	N/A	N/A
11261493	12/2021	Hackney et al.	N/A	N/A
11268119	12/2021	Mitchelson et al.	N/A	N/A
11286308	12/2021	Waldman et al.	N/A	N/A
11377499	12/2021	Wu	N/A	N/A
11389533	12/2021	Rosario et al.	N/A	N/A
11560434	12/2022	Diluzio et al.	N/A	N/A
11596688	12/2022	Rosario et al.	N/A	N/A
11639390	12/2022	Min et al.	N/A	N/A
11753668	12/2022	Mitchelson et al.	N/A	N/A
11760803	12/2022	Brake et al.	N/A	N/A
11780852	12/2022	Hudson et al.	N/A	N/A
11806507	12/2022	Herzlinger	N/A	N/A
11884731	12/2023	Lasch	N/A	N/A
11911409	12/2023	Yoneyama	N/A	N/A
11987620	12/2023	Johnson et al.	N/A	N/A
11999791	12/2023	Kao et al.	N/A	N/A
12024561	12/2023	Jayaraman et al.	N/A	N/A
12030948	12/2023	Jayaraman et al.	N/A	N/A
12053526	12/2023	Scholz et al.	N/A	N/A
12103975	12/2023	Gawlitsek et al.	N/A	N/A
12133966	12/2023	Herzlinger	N/A	N/A
12304956	12/2024	Harwin	424/144.1	C07K 16/00
2001/0046496	12/2000	Brettman et al.	N/A	N/A

2002/0197233	12/2001	Relton et al.	N/A	N/A
2004/0132642	12/2003	Hwang	N/A	N/A
2005/0053598	12/2004	Burke et al.	N/A	N/A
2005/0074451	12/2004	Yednock et al.	N/A	N/A
2005/0095238	12/2004	Brettman et al.	N/A	N/A
2005/0233387	12/2004	Artis et al.	N/A	N/A
2005/0255118	12/2004	Wehner	N/A	N/A
2005/0260193	12/2004	Lieberburg	N/A	N/A
2005/0276803	12/2004	Chan et al.	N/A	N/A
2006/0029600	12/2005	Rubin et al.	N/A	N/A
2006/0128607	12/2005	Bosserhoff et al.	N/A	N/A
2006/0234306	12/2005	Artis et al.	N/A	N/A
2007/0048219	12/2006	Hsei et al.	N/A	N/A
2007/0122403	12/2006	Dall'Acqua et al.	N/A	N/A
2007/0122404	12/2006	O'Keefe	N/A	N/A
2008/0075719	12/2007	Chan et al.	N/A	N/A
2008/0153756	12/2007	Krissansen et al.	N/A	N/A
2009/0011464	12/2008	Ponath et al.	N/A	N/A
2009/0208492	12/2008	O'Connor et al.	N/A	N/A
2009/0214527	12/2008	Burgess et al.	N/A	N/A
2009/0263421	12/2008	Spetz-Holmgren et al.	N/A	N/A
2009/0298195	12/2008	Ruker et al.	N/A	N/A
2010/0119517	12/2009	Burgess et al.	N/A	N/A
2010/0129353	12/2009	DeLuca	N/A	N/A
2010/0247431	12/2009	Armour et al.	N/A	N/A
2010/0255508	12/2009	Gelzleichter et al.	N/A	N/A
2010/0266587	12/2009	McLachlan	N/A	N/A
2010/0266636	12/2009	Richardson et al.	N/A	N/A
2010/0267934	12/2009	VandeWinkel et al.	N/A	N/A
2010/0297699	12/2009	Li et al.	N/A	N/A
2011/0002851	12/2010	Haas et al.	N/A	N/A
2011/0158982	12/2010	Segal et al.	N/A	N/A
2012/0034243	12/2011	Brettman et al.	N/A	N/A
2012/0039907	12/2011	Armour et al.	N/A	N/A
2012/0121580	12/2011	Bhambhani et al.	N/A	N/A
2012/0282249	12/2011	Fox et al.	N/A	N/A
2013/0052642	12/2012	Alvarez et al.	N/A	N/A
2013/0059337	12/2012	Bendig et al.	N/A	N/A
2013/0108617	12/2012	Kappos et al.	N/A	N/A
2013/0109032	12/2012	Gelzleichter et al.	N/A	N/A
2013/0337470	12/2012	Chackerian	N/A	N/A
2014/0004130	12/2013	Bordignon	N/A	N/A
2014/0080762	12/2013	Hazlehurst et al.	N/A	N/A
2014/0120084	12/2013	Anand et al.	N/A	N/A
2014/0135483	12/2013	Seito et al.	N/A	N/A
2014/0154252	12/2013	Thompson et al.	N/A	N/A
2014/0170157	12/2013	Agarwal et al.	N/A	N/A
2014/0186345	12/2013	Brettman et al.	N/A	N/A
2014/0212425	12/2013	Chang et al.	N/A	N/A

2014/0377253	12/2013	Harding et al.	N/A	N/A
2015/0010563	12/2014	Hamrah et al.	N/A	N/A
2015/0056205	12/2014	Lieberburg	N/A	N/A
2015/0064177	12/2014	Bendig et al.	N/A	N/A
2015/0093399	12/2014	Jefferies	N/A	N/A
2015/0197560	12/2014	Fong et al.	N/A	N/A
2015/0238602	12/2014	Cadavid et al.	N/A	N/A
2016/0129112	12/2015	Neelon	N/A	N/A
2016/0209426	12/2015	Diehl et al.	N/A	N/A
2016/0340435	12/2015	Chang et al.	N/A	N/A
2016/0340443	12/2015	Rossi et al.	N/A	N/A
2016/0375133	12/2015	Bhambhani et al.	N/A	N/A
2017/0002077	12/2016	Tam et al.	N/A	N/A
2017/0020867	12/2016	Moussy et al.	N/A	N/A
2017/0102393	12/2016	Gelzleichter et al.	N/A	N/A
2017/0210807	12/2016	Fong et al.	N/A	N/A
2017/0258869	12/2016	Li	N/A	N/A
2017/0275365	12/2016	Hsu et al.	N/A	N/A
2017/0315133	12/2016	Alexander et al.	N/A	N/A
2017/0327584	12/2016	Lasch	N/A	N/A
2017/0360926	12/2016	Rosario et al.	N/A	N/A
2018/0051086	12/2017	Abhyanker	N/A	N/A
2018/0086833	12/2017	Hassanali et al.	N/A	N/A
2018/0142017	12/2017	Hamrah et al.	N/A	N/A
2018/0148514	12/2017	Williams	N/A	N/A
2018/0207279	12/2017	Fox et al.	N/A	N/A
2018/0289811	12/2017	Fox et al.	N/A	N/A
2018/0291104	12/2017	Fong et al.	N/A	N/A
2018/0327497	12/2017	Diluzio et al.	N/A	N/A
2018/0346578	12/2017	Diluzio et al.	N/A	N/A
2019/0010242	12/2018	Eckelman et al.	N/A	N/A
2019/0038744	12/2018	Cadavid et al.	N/A	N/A
2019/0040140	12/2018	Brake et al.	N/A	N/A
2019/0076532	12/2018	Diluzio et al.	N/A	N/A
2019/0077868	12/2018	Sachs et al.	N/A	N/A
2019/0192683	12/2018	Jefferies	N/A	N/A
2019/0231878	12/2018	Brown et al.	N/A	N/A
2019/0310266	12/2018	Gelzleichter et al.	N/A	N/A
2019/0330318	12/2018	Bennett et al.	N/A	N/A
2019/0330366	12/2018	Eckelman et al.	N/A	N/A
2019/0345240	12/2018	Srinivasan et al.	N/A	N/A
2020/0002422	12/2019	Sachs et al.	N/A	N/A
2020/0002423	12/2019	Brettman et al.	N/A	N/A
2020/0018751	12/2019	Singh et al.	N/A	N/A
2020/0025775	12/2019	Diehl et al.	N/A	N/A
2020/0031937	12/2019	Lasch	N/A	N/A
2020/0048354	12/2019	Fong et al.	N/A	N/A
2020/0087401	12/2019	Rosario et al.	N/A	N/A
2020/0148773	12/2019	Taylor et al.	N/A	N/A
2020/0155673	12/2019	Rosario et al.	N/A	N/A

2020/0165677	12/2019	Stappenbeck et al.	N/A	N/A
2020/0179486	12/2019	Rosario et al.	N/A	N/A
2020/0206353	12/2019	Fox et al.	N/A	N/A
2020/0241006	12/2019	Naik et al.	N/A	N/A
2020/0253506	12/2019	Jones et al.	N/A	N/A
2020/0265940	12/2019	Dulai	N/A	N/A
2020/0276303	12/2019	Cadavid et al.	N/A	N/A
2020/0283533	12/2019	Hosen et al.	N/A	N/A
2020/0323772	12/2019	Jones et al.	N/A	N/A
2020/0376092	12/2019	Vitalis et al.	N/A	N/A
2020/0392231	12/2019	Anania et al.	N/A	N/A
2020/0392232	12/2019	Hassanali et al.	N/A	N/A
2021/0024636	12/2020	Mansour-Awad	N/A	N/A
2021/0052733	12/2020	Diluzio et al.	N/A	N/A
2021/0147555	12/2020	Jayaraman et al.	N/A	N/A
2021/0238288	12/2020	Anand et al.	N/A	N/A
2021/0252141	12/2020	Lissoos et al.	N/A	N/A
2021/0253714	12/2020	Jayaraman et al.	N/A	N/A
2021/0260185	12/2020	Cadavid et al.	N/A	N/A
2021/0262032	12/2020	Chowers et al.	N/A	N/A
2021/0278416	12/2020	Ananthakrishnan et al.	N/A	N/A
2021/0290604	12/2020	Williams	N/A	N/A
2021/0301002	12/2020	Bennett et al.	N/A	N/A
2021/0322410	12/2020	Campbell et al.	N/A	N/A
2021/0338836	12/2020	Yu et al.	N/A	N/A
2021/0340261	12/2020	Diluzio et al.	N/A	N/A
2021/0340584	12/2020	Tescione et al.	N/A	N/A
2021/0388092	12/2020	Abhyankar	N/A	N/A
2021/0396753	12/2020	Rubin et al.	N/A	N/A
2021/0401982	12/2020	Jayaraman et al.	N/A	N/A
2022/0010019	12/2021	Swirski et al.	N/A	N/A
2022/0041735	12/2021	Hosen et al.	N/A	N/A
2022/0048992	12/2021	Gabathuler et al.	N/A	N/A
2022/0062372	12/2021	Srinivasan et al.	N/A	N/A
2022/0098329	12/2021	Santich et al.	N/A	N/A
2022/0162297	12/2021	Basi	N/A	N/A
2022/0170028	12/2021	Li et al.	N/A	N/A
2022/0186314	12/2021	Hackney et al.	N/A	N/A
2022/0193235	12/2021	Herzlinger	N/A	N/A
2022/0206012	12/2021	Chackerian	N/A	N/A
2022/0220556	12/2021	Khatri et al.	N/A	N/A
2022/0259291	12/2021	Oppenheim et al.	N/A	N/A
2022/0267448	12/2021	Gowtham et al.	N/A	N/A
2022/0267449	12/2021	Ameli et al.	N/A	N/A
2022/0289855	12/2021	Ginsberg et al.	N/A	N/A
2022/0290241	12/2021	McGovern et al.	N/A	N/A
2022/0298232	12/2021	Srinivasan et al.	N/A	N/A
2022/0306735	12/2021	Dekosky et al.	N/A	N/A
2022/0349903	12/2021	Gelzleichter et al.	N/A	N/A

2022/0357343	12/2021	Diehl et al.	N/A	N/A
2022/0370617	12/2021	Diluzio et al.	N/A	N/A
2022/0378739	12/2021	Rezvani et al.	N/A	N/A
2022/0403033	12/2021	Anand et al.	N/A	N/A
2022/0403034	12/2021	Hassanali et al.	N/A	N/A
2023/0033021	12/2022	Jones et al.	N/A	N/A
2023/0043949	12/2022	Rosario et al.	N/A	N/A
2023/0048046	12/2022	Li et al.	N/A	N/A
2023/0093155	12/2022	Li et al.	N/A	N/A
2023/0096620	12/2022	Mora et al.	N/A	N/A
2023/0142437	12/2022	Ling et al.	N/A	N/A
2023/0287075	12/2022	Mumm	N/A	N/A
2023/0295336	12/2022	Eckelman et al.	N/A	N/A
2023/0310712	12/2022	Ling et al.	N/A	N/A
2023/0312727	12/2022	Diluzio et al.	N/A	N/A
2023/0313305	12/2022	Stappenbeck et al.	N/A	N/A
2023/0340131	12/2022	Jayaraman et al.	N/A	N/A
2023/0391887	12/2022	Williams	N/A	N/A
2023/0399415	12/2022	Preyer et al.	N/A	N/A
2024/0084016	12/2023	Jayaraman et al.	N/A	N/A
2024/0100158	12/2023	Jayaraman et al.	N/A	N/A
2024/0101679	12/2023	Jayaraman et al.	N/A	N/A
2024/0103008	12/2023	Baribaud et al.	N/A	N/A
2024/0173402	12/2023	Rosario et al.	N/A	N/A
2024/0182581	12/2023	Rosario et al.	N/A	N/A
2024/0199729	12/2023	Johnson et al.	N/A	N/A
2024/0239900	12/2023	Jayaraman et al.	N/A	N/A
2024/0242807	12/2023	Minar et al.	N/A	N/A
2024/0247067	12/2023	Brake et al.	N/A	N/A
2024/0254237	12/2023	Jaremicz et al.	N/A	N/A
2024/0279271	12/2023	Kumar et al.	N/A	N/A
2024/0316105	12/2023	Sun et al.	N/A	N/A
2024/0355416	12/2023	Ghiassian et al.	N/A	N/A
2024/0360222	12/2023	Hosen et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
102020016890	12/2021	BR	N/A
1173201	12/2004	EP	N/A
1133315	12/2005	EP	N/A
1214091	12/2005	EP	N/A
0808367	12/2006	EP	N/A
1113810	12/2007	EP	N/A
1078006	12/2008	EP	N/A
1699826	12/2008	EP	N/A
1835922	12/2008	EP	N/A
1123086	12/2009	EP	N/A
1485127	12/2010	EP	N/A
1784426	12/2010	EP	N/A

2458990	12/2011	EP	N/A
1278543	12/2011	EP	N/A
1896504	12/2011	EP	N/A
2140881	12/2012	EP	N/A
2279004	12/2014	EP	N/A
0918797	12/2014	EP	N/A
2021025	12/2015	EP	N/A
2298348	12/2015	EP	N/A
2109480	12/2016	EP	N/A
2704742	12/2016	EP	N/A
2704798	12/2016	EP	N/A
2739649	12/2016	EP	N/A
2632492	12/2016	EP	N/A
2854845	12/2017	EP	N/A
3224619	12/2018	EP	N/A
3131559	12/2018	EP	N/A
2903691	12/2018	EP	N/A
3268742	12/2018	EP	N/A
2408816	12/2018	EP	N/A
2782599	12/2018	EP	N/A
3254697	12/2018	EP	N/A
3321281	12/2018	EP	N/A
3326645	12/2019	EP	N/A
3167902	12/2019	EP	N/A
2970372	12/2019	EP	N/A
1587540	12/2020	EP	N/A
3530673	12/2021	EP	N/A
3411120	12/2022	EP	N/A
3554539	12/2022	EP	N/A
3615001	12/2022	EP	N/A
3204424	12/2022	EP	N/A
3237004	12/2023	EP	N/A
4456917	12/2023	EP	N/A
WO 1990007321	12/1989	WO	N/A
WO 2000030681	12/1999	WO	N/A
WO 2001080833	12/2000	WO	N/A
WO 2002077610	12/2001	WO	N/A
WO 2005046731	12/2004	WO	N/A
WO 2007007151	12/2006	WO	N/A
WO 2007007152	12/2006	WO	N/A
WO 2007007159	12/2006	WO	N/A
WO 2007007160	12/2006	WO	N/A
WO 2007061679	12/2006	WO	N/A
WO 2013142857	12/2012	WO	N/A
WO 2014049044	12/2013	WO	N/A
WO 2014182532	12/2013	WO	N/A
WO 2014193625	12/2013	WO	N/A
WO 2015174978	12/2014	WO	N/A
WO 2017019673	12/2016	WO	N/A
WO 2017087735	12/2016	WO	N/A

WO 2017158393	12/2016	WO	N/A
WO 2017165742	12/2016	WO	N/A
WO 2017218434	12/2016	WO	N/A
WO 2018104893	12/2017	WO	N/A
WO 2019180163	12/2018	WO	N/A
WO 2019196522	12/2018	WO	N/A
WO 2020260315	12/2019	WO	N/A
WO 2021003553	12/2020	WO	N/A
WO 2021060425	12/2020	WO	N/A
WO 2021070203	12/2020	WO	N/A
WO 2021188814	12/2020	WO	N/A
WO 2021188827	12/2020	WO	N/A
WO 2022026699	12/2021	WO	N/A
WO 2022036422	12/2021	WO	N/A
WO 2022123603	12/2021	WO	N/A
WO 2022157806	12/2021	WO	N/A
WO 2022192225	12/2021	WO	N/A
WO 2022234594	12/2021	WO	N/A
WO 2022253994	12/2021	WO	N/A
WO 2022272033	12/2021	WO	N/A
WO 2023288203	12/2022	WO	N/A
WO 2023007516	12/2022	WO	N/A
WO 2023012828	12/2022	WO	N/A
WO 2023021532	12/2022	WO	N/A
WO 2023044402	12/2022	WO	N/A
WO 2023058051	12/2022	WO	N/A
WO 2023064278	12/2022	WO	N/A
WO 2023064945	12/2022	WO	N/A
WO 2023073469	12/2022	WO	N/A
WO 2023081809	12/2022	WO	N/A
WO 2023086910	12/2022	WO	N/A
WO 2023097219	12/2022	WO	N/A
WO 2023116099	12/2022	WO	N/A
WO 2023126411	12/2022	WO	N/A
WO 2023166179	12/2022	WO	N/A
2023150731	12/2022	WO	N/A
WO 2023194837	12/2022	WO	N/A
WO 2023196866	12/2022	WO	N/A
WO 2023215882	12/2022	WO	N/A
WO 2023217072	12/2022	WO	N/A
2023248207	12/2022	WO	N/A
2024009205	12/2023	WO	N/A
2024081269	12/2023	WO	N/A
2024112618	12/2023	WO	N/A
2024123793	12/2023	WO	N/A
2024129960	12/2023	WO	N/A
2024151852	12/2023	WO	N/A
2024182425	12/2023	WO	N/A
2024188443	12/2023	WO	N/A
2024205498	12/2023	WO	N/A

2024206308	12/2023	WO	N/A
2024222859	12/2023	WO	N/A
2024229094	12/2023	WO	N/A
WO 2025038708	12/2024	WO	N/A

OTHER PUBLICATIONS

Aliprantis et al., “A Phase 1 Randomized, Double-Blind, Placebo-Controlled Trial to Assess the Safety, Tolerability, and Pharmacokinetics of a Respiratory Syncytial Virus Neutralizing Monoclonal Antibody MK-1654 in Healthy Adults”, *Clinical Pharm in Drug Development*, vol. 10, No. 5, pp. 556-566, 2021. cited by applicant

Arijs et al., “Effect of Vedolizumab (anti- $\alpha 4\beta 7$ -integrin) Therapy on Histological Healing and Mucosal Gene Expression in Patients with UC”, *Gut* Published Online First: Oct. 7, 2016 doi: 10.1136/gutjnl-2016-312293. cited by applicant

Ayalew et al., “C-Terminal Lysine Processing of IgG in Human Suction Blister Fluid: Implications for Subcutaneous Administration”, *Molecular Pharmaceutics*, vol. 19, pp. 4043-4054, 2022. cited by applicant

Boden et al., “Identification of Candidate Biomarkers Associated with Response to Vedolizumab in Inflammatory Bowel Disease”, *Digestive Diseases and Sciences*, vol. 63, pp. 2419-2429, 2018. cited by applicant

Boden et al., “Vedolizumab Efficacy Is Associated With Decreased Intracolonic Dendritic Cells, Not Memory T Cells”, *Inflammatory Bowel Diseases*, vol. 30, Issue 5, pp. 704-717, May 2024. cited by applicant

Bokemeyer et al., “Real-World Effectiveness of Vedolizumab vs Anti-TNF in Biologic-naïve Crohn's Disease Patients: A 2-year Propensity-score-adjusted Analysis from the VEDO IBD-Study”, *Inflammatory Bowel Diseases*, vol. 30, No. 5, pp. 746-756, May 2024. cited by applicant

Bressler et al., “Vedolizumab and Anti-Tumour Necrosis Factor α Real-World Outcomes in Biologic-Naïve Inflammatory Bowel Disease Patients: Results from the EVOLVE Study”, *Journal of Crohn's and Colitis*, vol. 15, No. 10, ppl 1694-1706, Oct. 2021. cited by applicant

Buer et al., “Combining Anti-TNF- α and Vedolizumab in the Treatment of Inflammatory Bowel Disease: A Case Series”, *Inflamm Bowel Dis.*, vol. 24, No. 5, pp. 997-1004, May 2018. cited by applicant

Casanova et al., “Persistence, Effectiveness and Safety of Ustekinumab and Vedolizumab Therapy for Complex Perianal Fistula in Crohn's Disease: The HEAL Study from GETECCU”, *Digestive and Liver Disease*, DOI: 10.1016/j.dld.2024.05.009, Jun. 2024. cited by applicant

Colombel et al., “Vedolizumab, Adalimumab, and Methotrexate Combination Therapy in Crohn's Disease (EXPLORER)”, *Clinical Gastroenterology and Hepatology*, vol. 22, No. 7, pp. 1487-1496, Jul. 2024. cited by applicant

D'Haens et al., Exposure-Efficacy Relationship of Vedolizumab Subcutaneous and Intravenous Formulations in Crohn's Disease and Ulcerative Colitis, *Expert Review of Clinical Pharmacology*, vol. 17, No. 4, pp. 403-412, Apr. 2024. cited by applicant

Dosing, “A Randomized Trial of Vedolizumab Dose Optimization in Patients With Moderate to Severe Ulcerative Colitis Who Have Early Nonresponse and High Drug Clearance: The ENTERPRET Trial”, *Gastroenterol. Hepatol.*, vol. 18, pp. 7-8, Jul. 2022. cited by applicant

Dreesen et al., “Evidence to Support Monitoring of Vedolizumab Trough Concentrations in Patients With Inflammatory Bowel Diseases”, *Clinical Gastroenterology and Hepatology*, vol. 16, No. 12, pp. 1937-1946, 2018. cited by applicant

Engel et al., “Vedolizumab in IBD-Lessons From Real-World Experience; A Systematic Review and Pooled Analysis”, *Journal of Crohn's and Colitis*, vol. 12, No. 2, pp. 245-257, Jan. 24, 2018. cited by applicant

Feagan et al., “Vedolizumab as Induction and Maintenance Therapy for Ulcerative Colitis”, *The*

New England Journal of Medicine, vol. 369, No. 8, pp. 699-710, Aug. 22, 2013. cited by applicant

Gabriels et al., "Using Fluorescently Labeled Vedolizumab to Visualize Local Drug Distribution During Colonoscopy and Identify Mucosal Target Cells in Patients with Inflammatory Bowel Disease". cited by applicant

Gonzalez et al. "Crohn's Patient Serum Proteomics Reveals Response Signature for Infliximab but not Vedolizumab". *Inflamm Bowel Dis.* Feb. 1, 2024;7:izae016. doi: 10.1093/ibd/izae016. Epub ahead of print. PMID: 38367209. cited by applicant

Hanzel et al., "Pharmacokinetic-Pharmacodynamic Model of Vedolizumab for Targeting Endoscopic Remission in Patients With Crohn Disease: Posthoc Analysis of the LOVE-CD Study", *Inflammatory Bowel Diseases*, vol. 28, pp. 689-699, 2022. cited by applicant

Haraya et al., "Translational Approach for Predicting Human Pharmacokinetics of Engineered Therapeutic Monoclonal Antibodies with Increased FcRn-Binding Mutations", *BioDrugs*, vol. 37, pp. 99-108, 2023. cited by applicant

Honap et al., "An Update on the Safety of Long-Term Vedolizumab Use in Inflammatory Bowel Disease", *Expert Opinion on Drug Safety*, vol. 22, No. 9, pp. 767-776, 2023. cited by applicant

Hu et al., "Evaluation of Heavy Chain C-Terminal Deletions on Productivity and Product Quality of Monoclonal Antibodies in Chinese Hamster Ovary (CHO) Cells", *Biotech Progress*, vol. 33, No. 3, pp. 786-794, Mar. 24, 2017. cited by applicant

Hu et al., "Combination Therapy Does Not Improve Rate of Clinical or Endoscopic Remission in Patients with Inflammatory Bowel Diseases Treated with Vedolizumab or Ustekinumab", *Clinical Gastroenterology and Hepatology*, vol. 19, pp. 1366-1376, 2021. cited by applicant

Hui et al., "Vedolizumab for Induction and Maintenance of Remission in Crohn's Disease", *Cochrane Database of Systematic Reviews*, Issue 7, Art. No. CD013611, DOI: 10.1002/14651858.CD013611.pub2, pp. 1-66, 2023. cited by applicant

Jiang et al., "Evaluation of Heavy-Chain C-Terminal Deletion on Product Quality and Pharmacokinetics of Monoclonal Antibodies", *Journal of Pharmaceutical Sciences*, vol. 105, No. 7, pp. 2066-2072, Jul. 1, 2016. cited by applicant

Laursen et al., "Pain Perception After Subcutaneous Injections of Media Containing Different Buffers", *Basic and Clinical Pharmacology and Toxicology*, vol. 98, pp. 218-221 (Year: 2006). cited by applicant

Liu et al., "Ustekinumab and Vedolizumab Dual Biologic Therapy in the Treatment of Crohn's Disease", *Case Reports in Medicine*, vol. 2017, Article ID 5264216, 2 pages, Nov. 2017. cited by applicant

Loftus et al., "Long-Term Safety of Vedolizumab for Inflammatory Bowel Disease", *Alimentary Pharmacology & Therapeutics*, vol. 52, pp. 1353-1365, 2020. cited by applicant

Mao et al., "Safety of Dual Biological Therapy in Crohn's Disease: A Case Series of Vedolizumab in Combination with Other Biologics", *BMJ Open Gastro*, vol. 5, No. 1, e000243. doi:10.1136/bmjgast-2018-000243. cited by applicant

Nowak et al., "A Phase 1 Randomized Dose-Escalation Study of a Human Monoclonal Antibody to IL-6 in CKD", *Kidney360*, vol. 2, pp. 224-235, Feb. 2021. cited by applicant

Orito et al., "A Phase I Study to Evaluate Safety, Pharmacokinetics, and Pharmacodynamics of Respiratory Syncytial Virus Neutralizing Monoclonal Antibody MK-1654 in Healthy Japanese Adults" *Clinical and Translational Science*, vol. 15, No. 7, pp. 1753-1763, Jul. 2022. cited by applicant

Osterman et al., "Vedolizumab Exposure Levels and Clinical Outcomes in Ulcerative Colitis: Determining the Potential for Dose Optimisation", *Aliment Pharmacol Ther.*, vol. 49, pp. 408-418, 2019. cited by applicant

Parikh et al., "Vedolizumab for the Treatment of Active Ulcerative Colitis: A Randomized Controlled Phase 2 Dose-Ranging Study", *Inflamm Bowel Dis.*, vol. 18, No. 8, pp. 1470-1479, Aug. 2012. cited by applicant

Parrot et al., "Systematic Review with Meta-Analysis: the Effectiveness of Either Ustekinumab or Vedolizumab in Patients with Crohn's Disease Refractory to Anti-Tumour Necrosis Factor", *Aliment Pharmacol Therap.*, vol. 55, pp. 380-388, 2022. cited by applicant

Peyrin-Biroulet et al., "Loss of Response to Vedolizumab and Ability of Dose Intensification to Restore Response in Patients with Crohn's Disease or Ulcerative Colitis: A Systematic Review and Meta-Analysis", *Clinical Gastroenterology and Hepatology*, vol. 17, No. 5, pp. 838-846, Apr. 2019. cited by applicant

Poole et al., "Vedolizumab: First Global Approval", *Drugs*, vol. 74, pp. 1293-1303, 2014. cited by applicant

Ramdani et al., "Monoclonal Antibody Engineering and Design to Modulate FcRn Activities: A Comprehensive Review", *International Journal of Molecular Sciences*, vol. 23, No., 17, pp. 9604 (1-12), Aug. 2022. cited by applicant

Ribaldone et al., "Dual biological therapy with Anti-TNF, Vedolizumab or Ustekinumab in Inflammatory Bowel Disease: A Systematic Review with Pool Analysis", *Scandinavian Journal of Gastroenterology*, vol. 54, No. 4, pp. 407-413, 2019. cited by applicant

Rosario et al., "Population Pharmacokinetics-Pharmacodynamics of Vedolizumab in Patients with Ulcerative Colitis and Crohn's Disease", *Alimentary Pharmacology & Therapeutics*, vol. 42, Issue 2, pp. 188-202, Jul. 2015. cited by applicant

Rosario et al., "Vedolizumab Pharmacokinetics, Pharmacodynamics, Safety, and Tolerability Following Administration of a Single, Ascending, Intravenous Dose to Healthy Volunteers", *Clin. Drug Investig.*, vol. 36, pp. 913-923, 2016. cited by applicant

Sandborn et al., "Vedolizumab as Induction and Maintenance Therapy for Crohn's Disease", *The New England Journal of Medicine*, vol. 369, No. 8, pp. 711-721, Aug. 22, 2013. cited by applicant

Sands et al., "Effects of Vedolizumab Induction Therapy for Patients With Crohn's Disease in Whom Tumor Necrosis Factor Antagonist Treatment Failed", *Gastroenterology*, vol. 147, pp. 618-627, 2014. cited by applicant

Sands et al., "Vedolizumab versus Adalimumab for Moderate-to-Severe Ulcerative Colitis", *N.E. Journal of Medicine*, vol. 381, No. 13, pp. 1215-1226, 2019. cited by applicant

Schmidt et al., "Predictors and Management of Loss of Response to Vedolizumab in Inflammatory Bowel Disease", *Inflamm Bowel Dis.*, vol. 24, No. 11, pp. 2461-2467, Nov. 2018. cited by applicant

Sengupta et al., "Higher Vedolizumab Serum Levels do not Increase the Risk of Adverse Events in Patients with Inflammatory Bowel Disease", *Scand. Journal of Gastroenterology*, vol. 55, No. 7, pp. 800-805, 2020. cited by applicant

Singh et al., "A Phase 1 Study of the Long-Acting Anti-IL-5 Monoclonal Antibody GSK3511294 in Patients with Asthma", *Br. J. Clin. Pharmacol.*, vol. 88, pp. 702-712, 2022. cited by applicant

Ungar et al., "Association of Vedolizumab Level, Anti-Drug Antibodies, and a4b7 Occupancy With Response in Patients With Inflammatory Bowel Diseases", *Clinical Gastroenterology and Hepatology*, vol. 16, pp. 697-705, 2018. cited by applicant

Ungar et al., "Dose Optimisation for Loss of Response to Vedolizumab—Pharmacokinetics and Immune Mechanisms", *Journal of Crohn's and Colitis*, vol. 15, No. 10, pp. 1707-1719, Oct. 2021. cited by applicant

Vande Casteele et al., "Real-world multicentre observational study including population pharmacokinetic modelling to evaluate the exposure-response relationship of vedolizumab in inflammatory bowel disease: ERELATE Study", *Alimentary Pharm & Therapeutics*, vol. 56, No. 3, pp. 463-476, Aug. 2022. cited by applicant

Veisman et al., "Association of Infliximab and Vedolizumab Trough Levels with Reported Rates of Adverse Events: A Cross-Sectional Study", *Journal of Clinical Medicine*, vol. 10, No. 18, pp. 4265, Sep. 2021. cited by applicant

Vermeire et al., "Efficacy and Safety of Subcutaneous Vedolizumab in Patients With Moderately to

Severely Active Crohn's Disease: Results From the VISIBLE 2 Randomised Trial", Journal of Crohn's and Colitis, vol. 16, No. 1, pp. 27-38, Jan. 2022. cited by applicant

Verstockt et al., "Expression Levels of 4 Genes in Colon Tissue Might Be Used to Predict Which Patients Will Enter Endoscopic Remission After Vedolizumab Therapy for Inflammatory Bowel Diseases", Clinical Gastroenterology and Hepatology, vol. 18, pp. 1142-1151, 2020. cited by applicant

Wyant et al., "Development and Validation of Receptor Occupancy Pharmacodynamic Assays Used in the Clinical Development of the Monoclonal Antibody Vedolizumab", Cytometry Part B (Clinical Cytometry), vol. 90B, pp. 168-176, 2016. cited by applicant

Wyant et al., "Vedolizumab Affects Antibody Responses to Immunization Selectively in the Gastrointestinal Tract: Randomised Controlled Trial Results", Gut, vol. 64, No. 1, pp. 77-83, 2015. cited by applicant

Xu et al., "PCSK9 Inhibitor Recaticimab for Hypercholesterolemia on Stable Statin Dose: a Randomized, Double-Blind, Placebo-Controlled Phase 1b/2 Study", BMC Medicine, vol. 20, No. 13, pp. 1-13, 2022. cited by applicant

Yarur et al., "Vedolizumab Concentrations Are Associated with Long-Term Endoscopic Remission in Patients with Inflammatory Bowel Diseases", Digestive Diseases and Sciences, vol. 64, pp. 1651-1659, 2019. cited by applicant

Zeissig et al., "Vedolizumab is Associated with Changes in Innate Rather than Adaptive Immunity in Patients with Inflammatory Bowel Disease", Gut, vol. 68, No. 1, pp. 25-39, Jan. 2019. cited by applicant

Zhu et al., "A Novel Monoclonal Antibody Drug Candidate SPY001 Targeting Integrin $\alpha 4\beta 7$ for the Treatment of IBD Demonstrates Prolonged Half-Life in Non-Human Primates", Abstract citation ID: jjad212.0895 P765. cited by applicant

Zhu et al., "Development and Characterization of a Novel Extended Half-Life Monoclonal Antibody Drug Candidate Targeting Integrin $\alpha 4\beta 7$ for the Treatment of IBD", Abstract citation ID: jjad212.0717 P587. cited by applicant

International Search Report and Written Opinion in PCT International Application No. PCT/US2024/031569 mailed Sep. 23, 2024. (14 pages). cited by applicant

Sandborn et al., "Efficacy and safety of vedolizumab subcutaneous formulation in a randomized trial of patients with ulcerative colitis", Gastroenterology, vol. 158, No. 3, pp. 562-572, Feb. 1, 2020. cited by applicant

Primary Examiner: Duffy; Brad

Attorney, Agent or Firm: Greenberg Traurig, LLP

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS (1) This application is a continuation of U.S. application Ser. No. 18/774,123, filed Jul. 16, 2024, which is a continuation of International Application No. PCT/US2024/031569, filed May 30, 2024, which claims the benefit of and priority to U.S. Provisional Application No. 63/504,966, filed on May 30, 2023; U.S. Provisional Application No. 63/505,962, filed on Jun. 2, 2023; U.S. Provisional Application No. 63/599,922, filed on Nov. 16, 2023; U.S. Provisional Application No. 63/554,886, filed on Feb. 16, 2024; and U.S. Provisional Application No. 63/559,081, filed on Feb. 28, 2024, the entire contents of each of which are incorporated herein by reference.

SEQUENCE LISTING

(1) This application contains a Sequence Listing which has been submitted electronically in XML format and is hereby incorporated by reference in its entirety. The XML copy, created on Apr. 14, 2025, is titled 220703-010211.xml and is 23.8 kilobytes in size.

BACKGROUND

(2) Integrins are cell-adhesion transmembrane receptors that function as extracellular matrix (ECM)-cytoskeletal linkers and transducers of biochemical and mechanical signals between cells and their environment. Due to their exposure on the cell surface and sensitivity to molecular inhibition, integrins such as $\alpha 4\beta 7$ integrins have been investigated as pharmacological targets for treating various diseases including cancer and inflammatory diseases (e.g., inflammatory bowel disease). However, current integrin therapies have been associated with serious side effects given the role of integrins in important biological processes and/or require multiple and frequent doses to maintain therapeutic efficacy. As such, improved $\alpha 4\beta 7$ therapies are needed.

SUMMARY

(3) The disclosure provides, in one aspect, a multidose regimen for use in treating a disease in a subject in need thereof comprising: (a) a first injectable liquid formulation comprising a total dosage amount of at least 500 mg of an $\alpha 4\beta 7$ binding antibody; and (b) a second injectable liquid formulation comprising a total dosage amount of at least 120 mg of the $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody consists of: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(4) In certain embodiments, the first injectable liquid formulation comprises a total dosage amount of from about 500 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody and the second injectable liquid formulation comprises a total dosage amount of from about 120 mg to about 450 mg of the $\alpha 4\beta 7$ binding antibody.

(5) In some embodiments, the first injectable liquid formulation comprises a total dosage amount of about 600 mg of the $\alpha 4\beta 7$ binding antibody and the second injectable liquid formulation comprises a total dosage amount of about 300 mg of the $\alpha 4\beta 7$ binding antibody.

(6) In some embodiments, the first injectable liquid formulation is for intravenous administration.

(7) In certain embodiments, the first injectable liquid formulation is for subcutaneous administration. In some embodiments, the first injectable liquid formulation is suitable for administration as a single injection or multiple injections.

(8) In certain embodiments, the second injectable liquid formulation is for subcutaneous administration. In some embodiments, the second injectable liquid formulation is suitable for administration as a single injection or multiple injections.

(9) In some embodiments, the disease is an inflammatory bowel disease. In some embodiments, the inflammatory bowel disease is Crohn's disease. In some embodiments, the inflammatory bowel disease is ulcerative colitis.

(10) In one aspect, the disclosure provides a dosing regimen for use in treating a disease in a subject in need thereof, the dosing regimen comprising: (a) a first formulation comprising a total dosage amount of at least about 500 mg of an $\alpha 4\beta 7$ binding antibody for administration to the subject; (b) a second formulation comprising a total dosage amount of at least about 120 mg of the $\alpha 4\beta 7$ binding antibody for subcutaneous administration to the subject after the first formulation, and thereafter as a maintenance dose at least eight weeks after administration of the second formulation, wherein the $\alpha 4\beta 7$ binding antibody consists of: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(11) In some embodiments, the first formulation is for subcutaneous administration.

(12) In some embodiments, the first formulation is for intravenous administration.

- (13) In certain embodiments, the first formulation comprises a total dosage amount of from about 500 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody. For example, in some embodiments, the first formulation comprises a total dosage amount of about 600 mg of the $\alpha 4\beta 7$ binding antibody.
- (14) In certain embodiments, the second formulation comprises a total dosage amount of from about 120 mg to about 450 mg of the $\alpha 4\beta 7$ binding antibody. For example, in some embodiments, the second formulation comprises a total dosage amount of about 300 mg of the $\alpha 4\beta 7$ binding antibody.
- (15) In certain embodiments, the second formulation is administered at least two weeks after the first formulation, and thereafter as a maintenance dose at least eight weeks after administration of the second formulation.
- (16) In some embodiments, the first formulation is suitable for administration as a single injection or multiple injections. In some embodiments, the second formulation is suitable for administration as a single injection or multiple injections.
- (17) In certain embodiments, the first formulation and the second formulation do not contain citrate.
- (18) In certain embodiments, the disease is an inflammatory bowel disease. In some embodiments, the inflammatory bowel disease is Crohn's disease. In some embodiments, the inflammatory bowel disease is ulcerative colitis.
- (19) In another aspect, the disclosure provides an injectable dosage form of an $\alpha 4\beta 7$ binding antibody consisting of (a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and (b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3, wherein the $\alpha 4\beta 7$ binding antibody has an average serum half-life greater than about 10 days in cynomolgus monkeys.
- (20) In some embodiments, the average serum half-life is about 17 days or greater in cynomolgus monkeys.
- (21) In certain embodiments, the injectable dosage form is an injectable liquid formulation.
- (22) In certain embodiments, the $\alpha 4\beta 7$ binding antibody is present at a concentration of at least 180 mg/mL.
- (23) In some embodiments, the injectable liquid formulation does not comprise citrate.
- (24) In one aspect, the disclosure provides a method of treating a disease in a patient in need thereof, the method comprising: administering to a subject in need thereof (a) an effective amount of an induction dose of an $\alpha 4\beta 7$ binding antibody, and (b) an effective amount of one or more maintenance doses of the $\alpha 4\beta 7$ binding antibody, wherein the one or more maintenance doses are administered subcutaneously at least eight weeks apart, wherein the $\alpha 4\beta 7$ binding antibody consists of: (a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and (b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.
- (25) In certain embodiments, the disease is an inflammatory bowel disease. In some embodiments, the inflammatory bowel disease is Crohn's disease. In some embodiments, the inflammatory bowel disease is ulcerative colitis.
- (26) In certain embodiments, the effective amount of the induction dose of the $\alpha 4\beta 7$ binding antibody is at least 600 mg, and the effective amount of each of the one or more maintenance doses of the $\alpha 4\beta 7$ binding antibody is at least 300 mg.
- (27) In certain embodiments, the effective amount of the induction dose of the $\alpha 4\beta 7$ binding antibody is about 600 mg and the effective amount of each of the one or more maintenance doses of the $\alpha 4\beta 7$ binding antibody is about 300 mg.
- (28) In some embodiments, the effective amount of the induction dose of the $\alpha 4\beta 7$ binding antibody is administered subcutaneously.
- (29) In some embodiments, the effective amount of the induction dose of the $\alpha 4\beta 7$ binding antibody is administered intravenously.
- (30) In some embodiments, the one or more maintenance doses are administered subcutaneously

about 12 weeks apart.

(31) In some embodiments, the one or more maintenance doses are administered subcutaneously about 26 weeks apart.

(32) In some embodiments, the induction dose comprises a dosage amount of the $\alpha 4\beta 7$ binding antibody that is about 2 or more times higher than a dosage amount of each of the one or more maintenance doses.

(33) In one aspect, the disclosure provides a method of achieving C.sub.trough of 35 $\mu\text{g/mL}$ or greater for a $\alpha 4\beta 7$ binding antibody in a subject in need thereof six weeks following administration of the $\alpha 4\beta 7$ binding antibody to the subject, the method comprising administering to the subject in need thereof at least about 600 mg of the $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody consists of: (a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and (b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(34) In certain embodiments, from about 600 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody is administered intravenously.

(35) In certain embodiments, from about 600 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody is administered subcutaneously.

(36) In some embodiments, about 600 mg of the $\alpha 4\beta 7$ binding antibody is administered intravenously.

(37) In some embodiments, about 600 mg of the $\alpha 4\beta 7$ binding antibody is administered subcutaneously.

(38) In some embodiments, the method further comprises administering to the subject in need thereof at least about 300 mg of the $\alpha 4\beta 7$ binding antibody, wherein the administration of the at least about 300 mg of the $\alpha 4\beta 7$ binding antibody results in a steady state C.sub.trough of at least 6 $\mu\text{g/mL}$. In certain embodiments, the administration results in a serum C.sub.avgW0-W12 of at least 60 g/mL.

(39) In some embodiments, the administration of the at least about 600 mg of the $\alpha 4\beta 7$ binding antibody results in a serum C.sub.avgW30-W40 of at least 30 g/mL.

(40) In some embodiments, the method further comprises administering to the subject in need thereof at least about 300 mg of the $\alpha 4\beta 7$ binding antibody and wherein administration results in an average serum concentration of the $\alpha 4\beta 7$ binding antibody above 35 g/mL for ten weeks following the administration of the at least about 300 mg of the $\alpha 4\beta 7$ binding antibody.

(41) In some embodiments, the method further comprises administering subcutaneously to the subject in need thereof at least about 300 mg of the $\alpha 4\beta 7$ binding antibody four weeks after administration of last dosage amount of the $\alpha 4\beta 7$ binding antibody.

(42) In some embodiments, the method further comprises administering subcutaneously to the subject a maintenance dose of at least about 600 mg of the $\alpha 4\beta 7$ binding antibody every twenty-six weeks after administration of the last dose of the $\alpha 4\beta 7$ binding antibody.

(43) In one aspect, the disclosure provides an $\alpha 4\beta 7$ binding antibody consisting of: (a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(44) In some embodiments, the disclosure provides an isolated nucleic acid encoding the heavy chain and/or light chain of the $\alpha 4\beta 7$ binding antibody, which antibody consists of (a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the isolated nucleic acid encodes SEQ ID NO: 1. In some embodiments, the isolated nucleic acid encodes SEQ ID NO: 3.

(45) In certain embodiments, the disclosure provides a recombinant host cell comprising an isolated nucleic acid encoding the heavy chain and/or light chain of the $\alpha 4\beta 7$ binding antibody, which antibody consists of (a) a heavy chain consisting of an amino acid sequence according to SEQ ID

NO: 1, and (b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the host cell comprises an isolated nucleic acid encodes SEQ ID NO: 1. In some embodiments, the host cell comprises an isolated nucleic acid encodes SEQ ID NO: 3.

(46) In certain embodiments, the disclosure provides an expression vector comprising an isolated nucleic acid encoding the heavy chain and/or light chain of the $\alpha 4\beta 7$ binding antibody, which antibody consists of (a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the expression vector comprises an isolated nucleic acid encodes SEQ ID NO: 1. In some embodiments, the expression vector comprises an isolated nucleic acid encodes SEQ ID NO: 3.

(47) In some embodiments, the disclosure provides a multidose regimen, a dosing regimen, an injectable dosage form, and/or a method, each as provided herein, wherein the disease is a gastrointestinal inflammatory disease.

(48) In certain embodiments, the disclosure provides a multidose regimen, a dosing regimen, an injectable dosage form, a method, and/or an antibody, each as provided herein, wherein the $\alpha 4\beta 7$ binding antibody has an average serum half-life greater than about 10 days in cynomolgus monkeys. In some embodiments, the average serum half-life is about 17 days or greater in cynomolgus monkeys.

(49) In certain embodiments, the disclosure provides a multidose regimen, a dosing regimen, an injectable dosage form, a method, and/or an antibody, each as provided herein, wherein the $\alpha 4\beta 7$ binding antibody has one or more of the following characteristics: (a) a melting temperature T_m Onset greater than about 55° C., (b) a melting temperature T_m Onset greater than 56.2° C., (c) a proportion of GOF from about 55.1% to about 60.4%, (d) a proportion of G1F from about 16.6% to about 18.8%, (e) a proportion of Man5 from about 7.3% to about 8.8%, (f) does not induce T cell activation markers CD25 or CD69, (g) does not result in release of cytokines, which cytokines are one or more of IL-6, IL-8, IL-10, IFN γ , IL-4, IL-17, IL-2, IL-23p70 and TNF, (h) does not induce complement-dependent cytotoxicity (CDC) in primary human PBMCs and in $\alpha 4\beta 7$ -expressing human β -lymphoid cell, (i) does not induce antibody dependent cellular cytotoxicity (ADCC) in human NK cells, and (j) does not impact suppressive activity of regulatory T cells expressing $\alpha 4\beta 7$ integrin, wherein the suppressive activity is measured by presence of elevated CD71, CD25, Ki67, granzyme B, or OX40.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

(1) FIG. 1A depicts cell binding of an $\alpha 4\beta 7$ binding antibody. Cell binding was determined (Relative MFI) for increasing concentrations of $\alpha 4\beta 7$ binding antibody Ab001 (dashed line, open shapes) or a comparator antibody (solid line, filled shapes) to PBMCs from three donors (Donor 1=circles; Donor 2=squares; Donor 3=triangles). EC₅₀ values were calculated for each donor combination and indicated in Table 2.

(2) FIG. 1B and FIG. 1C depict binding of the $\alpha 4\beta 7$ binding antibody Ab001 and a comparator antibody to $\alpha 4\beta 7$ -expressing peripheral blood mononuclear cells. FIG. 1B: Labeled (AlexaFluor-647) Ab001 binds to $\alpha 4\beta 7$ -expressing cells isolated from PBMCs. FIG. 1C: Modification competing off AF647-Ab001 with unlabeled Ab001 demonstrates receptor occupancy. N=3 donors.

(3) FIGS. 2A and 2B depict blocking of MAdCAM-1 and VCAM-1 adhesion by an $\alpha 4\beta 7$ binding antibody Ab001 (i.e., $\alpha 4\beta 7$ binding antibody comprising a heavy chain consisting of a sequence according to SEQ ID NO: 1 and a light chain consisting of a sequence according to SEQ ID NO: 3). FIG. 2A depicts the percent inhibition of total adhesion of integrin-mediated adhesion by MAdCAM-1. The percent inhibition was calculated after treatment with increasing concentrations

of either an $\alpha 4\beta 7$ binding antibody or comparator antibody. The IC_{sub.50} of the $\alpha 4\beta 7$ binding antibody was 86 μ M and the IC_{sub.50} of the comparator antibody was 83 μ M. FIG. 2B depicts the percent inhibition of total adhesion of integrin-mediated adhesion by VCAM-1, a potential off-target interaction. The percent inhibition was calculated after treatment with increasing concentrations of either an $\alpha 4\beta 7$ binding antibody, comparator antibody or a positive control antibody. The positive control antibody was able to inhibit adhesion by VCAM-1, but neither the $\alpha 4\beta 7$ binding antibody nor the comparator antibody were able to.

(4) FIG. 3 depicts the half-life of an $\alpha 4\beta 7$ binding antibody Ab001 in non-human primates (NHPs). NHPs were intravenously injected with either an $\alpha 4\beta 7$ binding antibody or comparator antibody and monitored out to 21 days post-injection. Normalized concentration of $\alpha 4\beta 7$ binding antibody or comparator antibody was calculated at each timepoint and used to determine the β -elimination half-life. Also plotted is data from an external study of the comparator antibody (dashed line) illustrating comparable results.

(5) FIG. 4 depicts simulated maintenance of trough concentrations at 6 weeks post-induction by IV associated with higher rates of remission in ulcerative colitis. The percentage of patients with C_{sub.trough} values ≤ 17 , 17-25, 25-35, and ≥ 35 μ g/mL were plotted in a stacked bar chart for patients treated with a comparator antibody during a clinical trial at 6 weeks post-induction. Also shown are the percentage of patients with C_{sub.trough} values from a simulated trial of 1000 patients treated with the comparator antibody, and an $\alpha 4\beta 7$ binding antibody described herein (e.g., Ab001) at doses of 300 mg, 350 mg, and 400 mg.

(6) FIG. 5 depicts quarterly maintenance dosing of $\alpha 4\beta 7$ binding antibody to maintain clinically relevant AUC levels. Relative AUC_{sub.ss,1 year} values were calculated using simulated trial of 1000 patients for an $\alpha 4\beta 7$ binding antibody described herein (e.g., Ab001) with a half-life of 35 or 50 days and compared to clinical trial data for intravenously (IV) and subcutaneously (SC) dosed comparator antibody. Relative AUC_{sub.ss,1 year} values were calculated for a quarterly dosing frequency ranging from 2 weeks (Q2W) to 16 weeks (Q16W).

(7) FIG. 6 depicts data from patients maintaining clinically relevant C_{sub.trough} values during maintenance dosing. In a simulated trial of 1000 patients, the percent of patients with a C_{sub.trough}-6 μ g/mL was calculated for patients treated with comparator antibody dosed intravenously (IV; 300 mg dose) and subcutaneously (SC; 108 mg or 320 mg dose) and patients subcutaneously dosed with 300 mg of an $\alpha 4\beta 7$ binding antibody described herein (e.g., Ab001) with a half-life of 43, 53 or 56 days. The percentage of patients with C_{sub.trough} ≥ 6 μ g/mL were plotted for a dosing frequency ranging from Q2W to Q20W.

(8) FIGS. 7A and 7B depict simulation of serum concentrations of the $\alpha 4\beta 7$ binding antibody described herein (e.g., Ab001) and a comparator antibody. FIG. 7A depicts an induction and maintenance dosing regimen by subcutaneous (SC) route of administration based on dosing at W0 and W2 during induction. FIG. 7B depicts serum concentrations based on Q8W IV dosing of comparator antibody and Q12W SC dosing of the $\alpha 4\beta 7$ binding antibody described herein (e.g., Ab001) based on the half-life of the antibodies.

(9) FIG. 7C (left) is a graph showing week 6 clinical remission rates in ulcerative colitis (%) (left); FIG. 7C (right) depicts induction simulation of serum concentrations of the $\alpha 4\beta 7$ binding antibody described herein (e.g., Ab001) and a comparator antibody.

(10) FIG. 7D is a graph showing human half-life predictions by NHP allometric scaling.

(11) FIG. 7E depicts maintenance simulation of serum concentrations of the $\alpha 4\beta 7$ binding antibody described herein (e.g., Ab001, 300 mg Q12W) and a comparator antibody (108 mg Q2W) when administered subcutaneously (300 mg Q12W for Ab001, 108 mg Q2W for comparator antibody).

(12) FIG. 7F show depicts simulation of serum concentrations of the $\alpha 4\beta 7$ binding antibody described herein (e.g., Ab001, SC dosing, 4-6 injections per year for maintenance) and a comparator antibody (IV/SC dosing, 26 injections per year for maintenance).

(13) FIGS. 8A-8C are graphs showing human PK simulation (Simulated Using 75 mg/kg NHP—

SC Dose (n=6, M/F)). FIG. 8A is a graph showing the $\alpha 4\beta 7$ binding antibody described herein (e.g., Ab001) simulated human profiles at 600 mg. FIG. 8B is a graph showing $\alpha 4\beta 7$ binding antibody described herein (e.g., Ab001) simulated human profiles 600 mg Q26W. FIG. 8C is a graph showing $\alpha 4\beta 7$ binding antibody described herein (e.g., Ab001) simulated human profiles 400 mg, 200 mg, Q12W.

(14) FIG. 9 is a graph showing serum concentration time profiles following subcutaneous and intravenous single-dose administration of $\alpha 4\beta 7$ binding antibody Ab001 in male and female monkeys. The male and female monkeys were administered a single subcutaneous dose of either 30 mg/kg, 75 mg/kg or 150 mg/kg of Ab001, or a single intravenous dose of 150 mg/kg of Ab001.

(15) FIGS. 10A-10C are graphs showing human PK induction simulation studies for $\alpha 4\beta 7$ binding antibody Ab001 relative to comparator antibody based on NHP t.sub.1/2 data. FIG. 10A is a graph showing induction simulation of serum concentrations of Ab001 in human at 600 mg of subcutaneous dose at week 0 and 300 mg of subcutaneous dose at week 2. FIG. 10B is a graph showing induction simulation of serum concentrations of Ab001 in human at 600 mg of subcutaneous dose at week 0, and 300 mg of subcutaneous doses at week 2 and week 6. FIG. 10C is a graph showing induction simulation of serum concentrations of the comparator antibody in human at 300 mg of intravenous dose at week 0, week 2 and week 6.

(16) FIGS. 11A-11D are graphs showing human PK maintenance simulation studies for $\alpha 4\beta 7$ binding antibody Ab001 relative to the comparator antibody based on NHP t.sub.1/2 data. FIG. 11A is a graph showing maintenance simulation of serum concentrations of Ab001 in human at 600 mg of subcutaneous dose every 26 weeks. FIG. 11B is a graph showing maintenance simulation of serum concentrations of Ab001 in human at 300 mg of subcutaneous dose every 12 weeks. FIG. 11C is a graph showing maintenance simulation of serum concentrations of the comparator antibody in human at 300 mg of intravenous dose every 8 weeks. FIG. 11D is a graph showing maintenance simulation of serum concentrations of the comparator antibody in human at 108 mg of subcutaneous dose every 2 weeks.

(17) FIG. 12 is a graph showing human PK simulation profile for $\alpha 4\beta 7$ binding antibody described herein such as Ab001 at 600 mg and 300 mg in a single ascending dose study.

(18) FIG. 13 is a graph showing human PK simulation profile for $\alpha 4\beta 7$ binding antibody described herein such as Ab001 at 600 mg and 300 mg in a multiple ascending dose study.

DETAILED DESCRIPTION

(19) To facilitate an understanding of the present disclosure, a number of terms and phrases are defined below.

(20) As used herein, unless otherwise indicated, the term “antibody” is understood to mean an intact antibody (e.g., an intact monoclonal antibody), or a fragment thereof, such as a Fc fragment of an antibody (e.g., an Fc fragment of a monoclonal antibody), or an antigen-binding fragment of an antibody (e.g., an antigen-binding fragment of a monoclonal antibody), including an intact antibody, antigen-binding fragment, or Fc fragment that has been modified, engineered, or chemically conjugated. In general, antibodies are multimeric proteins that contain four polypeptide chains. Two of the polypeptide chains are called immunoglobulin heavy chains (H chains), and two of the polypeptide chains are called immunoglobulin light chains (L chains). The immunoglobulin heavy and light chains are connected by an interchain disulfide bond. The immunoglobulin heavy chains are connected by interchain disulfide bonds. A light chain consists of one variable region (VL) and one constant region (CL). The heavy chain consists of one variable region (VH) and at least three constant regions (CH1, CH2 and CH3). The variable regions determine the binding specificity of the antibody. Each variable region contains three hypervariable regions known as complementarity determining regions (CDRs) flanked by four relatively conserved regions known as framework regions (FRs). The extent of the FRs and CDRs has been defined (Kabat, E. A., et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; and Chothia, C. et al. (1987) J. Mol. Biol.

196:901-917). The three CDRs, referred to as CDR1, CDR2, and CDR3, contribute to the antibody binding specificity. Naturally occurring antibodies have been used as starting material for engineered antibodies, such as chimeric antibodies and humanized antibodies. Examples of antibody-based antigen-binding fragments include Fab, Fab', (Fab')₂, Fv, single chain antibodies (e.g., scFv), minibodies, and diabodies. Examples of antibodies that have been modified or engineered include chimeric antibodies, humanized antibodies, and multispecific antibodies (e.g., bispecific antibodies). An example of a chemically conjugated antibody is an antibody conjugated to a toxin moiety.

(21) The terms “variable domain” and “variable region” are used interchangeably and refer to the portions of the antibody or immunoglobulin domains that exhibit variability in their sequence and that are involved in determining the specificity and binding affinity of a particular antibody. Variability is not evenly distributed throughout the variable domains of antibodies; it is concentrated in sub-domains of each of the heavy and light chain variable regions. These sub-domains are called “hypervariable regions” or “complementarity determining regions” (CDRs). The more conserved (i.e., non-hypervariable) portions of the variable domains are called the “framework” regions (FRM or FR) and provide a scaffold for the six CDRs in three-dimensional space to form an antigen-binding surface.

(22) An “Fc polypeptide” of a dimeric Fc as used herein refers to one of the two polypeptides forming the dimeric Fc domain, i.e. a polypeptide comprising C-terminal constant regions of an immunoglobulin heavy chain, capable of stable self-association. For example, an Fc polypeptide of a dimeric IgG Fc comprises an IgG CH₂ and an IgG CH₃ constant domain sequence. An Fc can be of the class IgA, IgD, IgE, IgG, and IgM. These classes are also designated α , δ , ϵ , γ , and μ , respectively. Several of these may be further divided into subclasses (isotypes), e.g., IgG.sub.1, IgG.sub.2, IgG.sub.3, IgG.sub.4, IgA.sub.1, and IgA.sub.2.

(23) The terms “Fc receptor” and “FcR” are used to describe a receptor that binds to the Fc region of an antibody. For example, an FcR can be a native sequence human FcR. Generally, an FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the Fc γ RI, Fc γ RII, and Fc γ RII subclasses, including allelic variants and alternatively spliced forms of these receptors. Fc γ RII receptors include Fc γ RIIA (an “activating receptor”) and Fc γ RIIB (an “inhibiting receptor”), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Immunoglobulins of other isotypes can also be bound by certain FcRs (see, e.g., Janeway et al., *Immuno Biology: the immune system in health and disease*, (Elsevier Science Ltd., NY) (4th ed., 1999)). Activating receptor Fc γ RIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor Fc γ RIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain (reviewed in Daeron, *Annu. Rev. Immunol.* 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-92 (1991); Capel et al., *Immunomethods* 4:25-34 (1994); and de Haas et al., *J. Lab. Clin. Med.* 126:330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term “FcR” herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer et al., *J. Immunol.* 117:587 (1976); and Kim et al., *J. Immunol.* 24:249 (1994)).

(24) The terms “recipient”, “individual”, “subject”, “host”, and “patient”, are used interchangeably herein and in some embodiments, refer to any mammalian subject for whom diagnosis, treatment, or therapy is desired, particularly humans. “Mammal” for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and laboratory, zoo, sports, or pet animals, such as dogs, horses, cats, cows, sheep, goats, pigs, mice, rats, rabbits, guinea pigs, monkeys etc. In some embodiments, the mammal is human. None of these terms require the supervision of medical personnel.

(25) As used herein, the term “effective amount” refers to the amount of a compound (e.g., a compound of the present disclosure) sufficient to effect beneficial or desired results. An effective

amount can be administered in one or more administrations, applications or dosages and is not intended to be limited to a particular formulation or administration route. As used herein, the term “treating” includes any effect, e.g., lessening, reducing, modulating, ameliorating or eliminating, that results in the improvement of the condition, disease, disorder, and the like, or ameliorating a symptom thereof.

(26) As used herein, the term “pharmaceutical composition” refers to the combination of an active agent with a carrier, inert or active, making the composition especially suitable for diagnostic or therapeutic use in vivo or ex vivo.

(27) As used herein, the term “pharmaceutically acceptable carrier” refers to any of the standard pharmaceutical carriers, such as a phosphate buffered saline solution, water, emulsions (e.g., such as an oil/water or water/oil emulsions), and various types of wetting agents. The compositions also can include stabilizers and preservatives. For examples of carriers, stabilizers and adjuvants, see e.g., Martin, Remington's Pharmaceutical Sciences, 15th Ed., Mack Publ. Co., Easton, PA (1975).

(28) The terms “a” and “an” as used herein mean “one or more” and include the plural unless the context is inappropriate.

(29) As used herein, all numerical values or numerical ranges include whole integers within or encompassing such ranges and fractions of the values or the integers within or encompassing ranges unless the context clearly indicates otherwise. Thus, for example, reference to a range of 90-100%, includes 91%, 92%, 93%, 94%, 95%, 95%, 96%, 97%, etc., as well as 91.1%, 91.2%, 91.3%, 91.4%, 91.5%, etc., 92.1%, 92.2%, 92.3%, 92.4%, 92.5%, etc., and so forth. In another example, reference to a range of 1-5,000-fold includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, fold, etc., as well as 1.1, 1.2, 1.3, 1.4, 1.5, fold, etc., 2.1, 2.2, 2.3, 2.4, 2.5, fold, etc., and so forth.

(30) “About” a number, as used herein, refers to range including the number and ranging from 10% below that number to 10% above that number. “About” a range refers to 10% below the lower limit of the range, spanning to 10% above the upper limit of the range.

(31) “Percent (%) identity” refers to the extent to which two sequences (nucleotide or amino acid) have the same residue at the same positions in an alignment. For example, “an amino acid sequence is X % identical to SEQ ID NO: Y” refers to % identity of the amino acid sequence to SEQ ID NO: Y and is elaborated as X % of residues in the amino acid sequence are identical to the residues of sequence disclosed in SEQ ID NO: Y. Generally, computer programs are employed for such calculations. Exemplary programs that compare and align pairs of sequences, include ALIGN (Myers and Miller, 1988), FASTA (Pearson and Lipman, 1988; Pearson, 1990) and gapped BLAST (Altschul et al., 1997), BLASTP, BLASTN, or GCG (Devereux et al., 1984).

(32) Throughout the description, where compositions are described as having, including, or comprising specific components, or where processes and methods are described as having, including, or comprising specific steps, it is contemplated that, additionally, there are compositions of the present disclosure that consist essentially of, or consist of, the recited components, and that there are processes and methods according to the present disclosure that consist essentially of, or consist of, the recited processing steps.

(33) As a general matter, compositions specifying a percentage are by weight unless otherwise specified. Further, if a variable is not accompanied by a definition, then the previous definition of the variable controls.

(34) $\alpha 4\beta 7$ Integrin Binding Antibodies

(35) Provided herein are $\alpha 4\beta 7$ binding antibodies (e.g., $\alpha 4\beta 7$ binding proteins, long acting $\alpha 4\beta 7$ binding antibodies, long acting $\alpha 4\beta 7$ binding molecules or long acting $\alpha 4\beta 7$ targeting molecules. The $\alpha 4\beta 7$ binding antibodies provided herein have a binding specificity for target antigen human $\alpha 4\beta 7$ integrin). $\alpha 4\beta 7$ binding antibodies described herein may have improved life and/or specificity. $\alpha 4\beta 7$ binding antibodies described herein may have specificity for $\alpha 4\beta 7$ and not related integrins including $\alpha 4\beta 1$ and $\alpha E\beta 7$.

antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain comprising an amino acid sequence having at least 95% sequence identity with an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain comprising an amino acid sequence having at least 96% sequence identity with an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain comprising an amino acid sequence having at least 97% sequence identity with an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain comprising an amino acid sequence having at least 98% sequence identity with an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain comprising an amino acid sequence having at least 99% sequence identity with an amino acid sequence according to SEQ ID NO: 3.

(41) In some embodiments, the antigen-binding site comprises the HCDR1, HCDR2, HCDR3, LCDR1, LCDR2, and LCDR3, determined under Kabat (see Kabat et al., (1991) Sequences of Proteins of Immunological Interest, NIH Publication No. 91-3242, Bethesda), Chothia (see, e.g., Chothia C & Lesk A M, (1987), *J Mol Biol* 196: 901-917), MacCallum (see MacCallum R M et al., (1996) *J Mol Biol* 262: 732-745), IMGT (see Lefranc, (1999) *The Immunologist*, 7, 132-136), or any other CDR determination method known in the art, of the heavy chain and light chain sequences of an antibody disclosed in Table 1A.

(42) TABLE-US-00001 TABLE 1A Sequences of $\alpha 4\beta 7$ binding antibodies SEQ ID NO Name Sequence 1 YTE Heavy QVQLVQSGAEVKKPGASVKVSCKGSGYTFTSYWM Chain HWVRQAPGQRLEWIGEIDPSESNTNYNQKFKGRVT LTVDISASTAYMELSSLRSED TAVYYCARGGYD GW DY AIDYWGQGTLVTVSSASTKGPSVFPLAPSSKSTS GGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFP AVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSN TKVDKKVEPKSCDKTHTCPPCPAELAGAPSVFLFPP KPKDTLYITREPEVTCVVVDVSHEDPEVKFNWYVD GVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYT LPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTTPVLDS DGSFFLYSKLTVDKSRWQQGN VFSCSV MHEALHNHYTQKSLSLSPG 2 LS Heavy QVQLVQSGAEVKKPGASVKVSCKGSGYTFTSYWM Chain HWVRQAPGQRLEWIGEIDPSESNTNYNQKFKGRVT LTVDISASTAYMELSSLRSED TAVYYCARGGYD GW DY AIDYWGQGTLVTVSSASTKGPSVFPLAPSSKSTS GGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFP AVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSN TKVDKKVEPKSCDKTHTCPPCPAELAGAPSVFLFPP KPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVD GVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYT LPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTTPVLDS DGSFFLYSKLTVDKSRWQQGN VFSCSV LHEALHSHYTQKSLSLSPG 3 Light Chain DVVMTQSPLSLPVTPGEPASISCRSSQSLAKSYGNTY LSWY LQKPGQSPQLLIYGISNRFS GVPDRFSGSGSGT DFTLKISRVEAEDVGVYYCLQGTHQPYTFGQGTKVE

IKRTVASFIFPPSDEQLKSGTASVVCLLNNFYPR
EAKVQWKVDNALQSGNSQESVTEQDSKDYSLSS
TLTLISKADYEKHKVYACEVTHQGLSSPVTKSFNRGE C

(43) In some embodiments, the antigen-binding site comprises the HCDR1, HCDR2, HCDR3, LCDR1, LCDR2, and LCDR3 disclosed in Table 1B.

(44) TABLE-US-00002 TABLE 1B SEQ ID NO Name Sequence
6 HCDR1 SYWMH
7 HCDR2 EIDPSESNTNYNQKFKG
8 HCDR3 GGYDGWDYAIDY
9 LCDR1 RSSQSLAKSYGNTYLS
10 LCDR2 GISNRFS
11 LCDR3 LQGTHQPYT

(45) In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 80% sequence identity with an amino acid sequence according to SEQ ID NO: 4. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 85% sequence identity with an amino acid sequence according to SEQ ID NO: 4. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 90% sequence identity with an amino acid sequence according to SEQ ID NO: 4. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 95% sequence identity with an amino acid sequence according to SEQ ID NO: 4. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 96% sequence identity with an amino acid sequence according to SEQ ID NO: 4. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 97% sequence identity with an amino acid sequence according to SEQ ID NO: 4. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 98% sequence identity with an amino acid sequence according to SEQ ID NO: 4. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 99% sequence identity with an amino acid sequence according to SEQ ID NO: 4. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having 100% sequence identity with an amino acid sequence according to SEQ ID NO: 4.

(46) In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a light chain comprising an amino acid sequence having at least 80% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a light chain comprising an amino acid sequence having at least 85% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a light chain comprising an amino acid sequence having at least 90% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a light chain comprising an amino acid sequence having at least 95% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a light chain comprising an amino acid sequence having at least 96% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a light chain comprising an amino acid sequence having at least 97% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a light chain comprising an amino acid sequence having at least 98% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a light chain comprising an amino acid sequence having at least 99% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a light chain comprising an amino acid sequence having 100% sequence identity with an amino acid sequence according to SEQ ID NO: 5.

(47) In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 80% sequence identity with an amino acid sequence according

to SEQ ID NO: 4; and a light chain comprising an amino acid sequence having at least 80% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 85% sequence identity with an amino acid sequence according to SEQ ID NO: 4; and a light chain comprising an amino acid sequence having at least 85% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 90% sequence identity with an amino acid sequence according to SEQ ID NO: 4; and a light chain comprising an amino acid sequence having at least 90% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 95% sequence identity with an amino acid sequence according to SEQ ID NO: 4; and a light chain comprising an amino acid sequence having at least 95% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 96% sequence identity with an amino acid sequence according to SEQ ID NO: 4; and a light chain comprising an amino acid sequence having at least 96% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 97% sequence identity with an amino acid sequence according to SEQ ID NO: 4; and a light chain comprising an amino acid sequence having at least 97% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 98% sequence identity with an amino acid sequence according to SEQ ID NO: 4; and a light chain comprising an amino acid sequence having at least 98% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence having at least 99% sequence identity with an amino acid sequence according to SEQ ID NO: 4; and a light chain comprising an amino acid sequence having at least 99% sequence identity with an amino acid sequence according to SEQ ID NO: 5. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence according to SEQ ID NO: 4; and a light chain comprising an amino acid sequence according to SEQ ID NO: 5.

(48) In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain comprising an amino acid sequence consisting of SEQ ID NO: 3.

(49) In some embodiments, the $\alpha 4\beta 7$ binding antibody Ab001 comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence consisting of SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody Ab001 consists of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence consisting of SEQ ID NO: 3.

(50) TABLE-US-00003 The $\alpha 4\beta 7$ binding antibody of the disclosure can have the following properties: Property Range Exemplary Value Solubility (30° C.) 100 mg/ml-300 mg/ml 200 mg/ml Solubility (5° C.) 90 mg/ml-270 mg/ml 206.5 mg/ml Viscosity at 150 6-16 mPa 11.1 mPa mg/ml, 25° C. (161 mg/mL, 25° C.) pI 7.5-7.9 7.7 Expression titer 1-10 4.7 mg/mL

(51) The $\alpha 4\beta 7$ binding antibody of the disclosure can have the following N-glycan profiling (determined using enzymatically released N-glycans, labeled and monitored by hydrophilic interaction liquid chromatography (HILIC) coupled with a fluorescence detector (FLD)):

(52) TABLE-US-00004 Glycan Range Exemplary Value G0F 40-75% 60.4% (sample 1), 55.1% (sample 2) G1F 10-35% 16.6% (sample 1) 18.8% (sample 2) Man5 1-15% 7.3% (sample 1) 8.8% (sample 2)

(53) In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises one or more of the following characteristics: (a) an average serum half-life greater than about 10 days in cynomolgus monkeys,

(b) an average serum half-life of about 17 days or greater in cynomolgus monkeys, (c) a melting temperature T_{mOnset} greater than about 55° C., (d) a melting temperature T_{mOnset} greater than 56.2° C., (e) does not induce T cell activation markers CD25 or CD69, (f) does not result in release of cytokines at least 6 hours after the administration, wherein the cytokines comprise one or more of IL-6, IL-8, IL-10, IFN γ , IL-4, IL-17, IL-2, IL-23p70 and TNF, (g) does not induce complement-dependent cytotoxicity (CDC) in primary human PBMCs and in $\alpha 4\beta 7$ -expressing human β -lymphoid cell, (h) does not induce antibody dependent cellular cytotoxicity (ADCC) in human NK cells, and (i) does not impact suppressive activity of regulatory T cells expressing $\alpha 4\beta 7$ integrin, wherein the suppressive activity is measured by presence of elevated CD71, CD25, Ki67, granzyme B, or OX40.

Fc Modifications

(54) Provided herein are $\alpha 4\beta 7$ binding antibodies comprising modified Fc regions. Unless otherwise specified herein, numbering of amino acid residues in the Fc region or constant region is according to the EU numbering system, also called the EU index, as described in Kabat et al, Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD, 1991.

(55) In some embodiments, the $\alpha 4\beta 7$ binding antibodies comprise a modified Fc comprising one or more modifications. In some embodiments, the one or more modifications promote selective binding of Fc-gamma receptors.

(56) In some embodiments, one or more modifications in the modified Fc is selected from the group consisting of: S298A, E333A, K334A, K326A, F243L, R292P, Y300L, V3051, P396L, F243L, R292P, Y300L, L235V, P396L, F243L, S239D, I332E, A330L, S267E, L328F, D265S, S239E, K326A, A327H, G237F, K326E, G236A, D270L, H268D, S324T, L234F, N325L, V266L, and S267D. In some embodiments, one or more modifications in the modified Fc is selected from the group consisting of S228P, M252Y, S254T, T256E, T256D, T250Q, H285D, T307A, T307Q, T307R, T307W, L309D, Q411H, Q311V, A378V, E380A, M428L, N434A, N434S, N297A, D265A, L234A, L235A, and N434W.

(57) In some embodiments, the modified Fc comprises a specific combination of amino acid substitutions selected from the group consisting of: L234A/L235A; V234A/G237A; L235A/G237A/E318A; S228P/L236E; H268Q/V309L/A330S/A331S; C220S/C226S/C229S/P238S; C226S/C229S/E3233P/L235V/L235A; L234F/L235E/P331S; C226S/P230S; L234A/G237A; L234A/L235A/G237A; L234A/L235A/P329G.

(58) In some embodiments, the modified Fc comprises a specific combination of amino acid substitutions selected from the group consisting of M428L/N434S (LS); M252Y/S254T/T256E (YTE); T250Q/M428L; T307A/E380A/N434A; T256D/T307Q (DQ); T256D/T307W (DW); M252Y/T256D (YD); T307Q/Q311V/A378V (QVV); T256D/H285D/T307R/Q311V/A378V (DDRVV); L309D/Q311H/N434S (DHS); S228P/L235E (SPLE); L234A/L235A (LA); M428L/N434A; L234A/G237A (LAGA); L234A/L235A/G237A; L234A/L235A/P329G; D265A/YTE; LALA/YTE; LAGA/YTE; LALAGA/YTE; LALAPG/YTE; N297A/LS; D265A/LS; LALA/LS; LALAGA/LS; LALAPG/LS; N297A/DHS; D265A/DHS; LALA/DHS; LAGA/DHS; LALAGA/DHS; LALAPG/DHS; SP/YTE; SPLE/YTE; SP/LS; SPLE/LS; SP/DHS; SPLE/DHS; N297A/LA; D265A/LA; LALA/LA; LAGA/LA; LALAGA/LA; LALAPG/LA; N297A/N434A; D265A/N434A; LALA/N434A; LAGA/N434A; LALAGA/N434A; LALAPG/N434A; N297A/N434W; D265A/N434W; LALA/N434W; LAGA/N434W; LALAGA/N434W; LALAPG/N434W; N297A/DQ; D265A/DQ; LALA/DQ; LAGA/DQ; LALAGA/DQ; LALAPG/DQ; N297A/DW; D265A/DW; LALA/DW; LAGA/DW; LALAGA/DW; LALAPG/DW; N297A/YD; D265A/YD; LALA/YD; LAGA/YD; LALAGA/YD; LALAPG/YD; N297A/QVV; D265A/QVV; LALA/QVV; LAGA/QVV, LALAGA/QVV; LALAPG/QVV; DDRVV; N297A/DDRVV; D265A/DDRVV; LALA/DDRVV; LAGA/DDRVV; LALAGA/DDRVV; and LALAPG/DDRVV. In some embodiments, the modified Fc comprises a

specific combination of amino acid substitutions selected from the group consisting of M428L/N434S (LS) and M252Y/S254T/T256E (YTE). In some embodiments, the modified Fc comprises M428L/N434S (LS) modifications. In some embodiments, the modified Fc comprises M252Y/S254T/T256E (YTE) modifications.

(59) In some embodiments, the Fc of the $\alpha 4\beta 7$ binding antibodies described herein lacks a C-terminal lysine residue (e.g., Ab001), the lack of which can be expected to decrease pI. In some embodiments, composition comprising the $\alpha 4\beta 7$ binding antibodies wherein the Fc lacks C-terminal lysine residue (e.g., Ab001) has a homogeneous charge profile as compared to a composition comprising an $\alpha 4\beta 7$ binding antibody, wherein the Fc comprises a C-terminal lysine residue. While the lack of C-terminal lysine residue can be expected to decrease solubility or decrease recombinant production yield, it has been discovered that truncation of C-terminal lysine in the $\alpha 4\beta 7$ binding antibody yields a highly soluble protein and allows a relatively high production yield of $\alpha 4\beta 7$ binding antibodies. In particular, Ab001 achieved surprising high yields (greater than 4.5 g/L) and high solubility (greater than 150 g/L).

(60) In some embodiments, the methods provided herein wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain comprising an amino acid sequence according to SEQ ID NO: 3 (e.g., Ab001) may increase FcRn binding, while exhibiting satisfactory bioavailability when the $\alpha 4\beta 7$ binding antibody is administered subcutaneously.

(61) In some embodiments, the $\alpha 4\beta 7$ binding antibodies described herein include modifications to improve its ability to mediate effector function. Such modifications are known in the art and include afucosylation, or engineering of the affinity of the Fc towards an activating receptor, mainly FCGR3a for antibody-dependent cellular cytotoxicity (ADCC), and towards C1q for complement-dependent cytotoxicity (CDC).

(62) In some aspects, an antibody provided herein comprises an IgG1 domain with reduced fucose content at position Asn 297 (EU numbering) compared to a naturally occurring IgG1 domain. Such Fc domains are known to have improved ADCC. In some aspects, such antibodies do not comprise any fucose at position Asn 297.

(63) In some embodiments, the $\alpha 4\beta 7$ binding antibody provided herein comprises glycan moieties. In some embodiments, the $\alpha 4\beta 7$ binding antibody provided herein comprises an IgG1 domain with high total fucose. The major N-glycan types are GOF and GIF. In some embodiments, the $\alpha 4\beta 7$ binding antibody of the disclosure has one or more of the following N-glycan profiling: the proportion of GOF can be greater than about 40%. For example, the proportion of GOF can range from about 40% to about 75%; the proportion of G1F can range from about 10% to about 35%; the proportion of Man5 can be less than about 20%.

(64) In some embodiments, the proportion of Man5 can range from about 1% to about 15%. In some embodiments, the proportion of GOF in the $\alpha 4\beta 7$ binding antibody is from about 40% to about 45%, from about 45% to about 50%, from about 50% to about 55%, from about 55% to about 60%, from about 60% to about 65%, from about 65% to about 70%, from about 70% to about 75%.

(65) In some embodiments, the proportion of G1F in the $\alpha 4\beta 7$ binding antibody is from about 10% to about 15%, from about 15% to about 20%, from about 20% to about 25%, from about 25% to about 30%, from about 30% to about 35%.

(66) In some embodiments, the proportion of Man5 in the $\alpha 4\beta 7$ binding antibody is from about 1% to about 5%, from about 5% to about 10%, from about 10% to about 15%.

(67) In some embodiments, the $\alpha 4\beta 7$ binding antibodies described herein comprise an Fc region with one or more amino acid substitutions which improve ADCC, such as a substitution at one or more of positions 298, 333, and 334 of the Fc region. In some embodiments, an antibody provided herein comprises an Fc region with one or more amino acid substitutions at positions 239, 332, and 330.

(68) In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise an Fc region with at least one galactose residue in the oligosaccharide attached to the Fc region. Such antibody variants may have improved CDC function.

(69) In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise one or more alterations that improves or diminishes C1q binding and/or CDC.

(70) In certain embodiments, the Fc region comprises one or more amino acid substitutions, wherein the one or more substitutions result in an increase in one or more of antibody half-life, ADCC activity, ADCP activity, or CDC activity compared with the Fc without the one or more substitutions. In certain embodiments, the one or more amino acid substitutions results in increased antibody half-life at pH 6.0 compared to an antibody comprising a wild-type Fc region. In certain embodiments, the antibody has an increased half-life that is about 10,000-fold, 1,000-fold, 500-fold, 100-fold, 50-fold, 20-fold, 10-fold, 9-fold, 8-fold, 7-fold, 6-fold, 5-fold, 4.5-fold, 4-fold, 3.5-fold, 3-fold, 2.5-fold, 2-fold, 1.95-fold, 1.9-fold, 1.85-fold, 1.8-fold, 1.75-fold, 1.7-fold, 1.65-fold, 1.6-fold, 1.55-fold, 1.50-fold, 1.45-fold, 1.4-fold, 1.35-fold, 1.3-fold, 1.25-fold, 1.2-fold, 1.15-fold, 1.1-fold, or 1.05-fold longer compared to an antibody comprising a wild-type Fc region.

(71) In certain embodiments, the Fc region comprises one or more amino acid substitutions, wherein the one or more substitutions result in a decrease in one or more of ADCC activity, ADCP activity, or CDC activity compared with the Fc without the one or more substitutions.

(72) In certain embodiments, the Fc region binds an Fc γ Receptor selected from the group consisting of: Fc γ RI, Fc γ RIIa, Fc γ RIIb, Fc γ RIIc, Fc γ RIIIa, and Fc γ RIIIb. In certain embodiments, the Fc region binds an Fc γ Receptor with higher affinity at pH 6.0 compared to an antibody comprising a wild-type Fc region.

(73) In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise an extended half-life (i.e., serum half-life). In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life of at least about 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 94, 96, or more than 96 days in humans. In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life in a range of about 28 days to about 96 days, about 28 days to about 84 days, about 28 days to about 70 days, about 28 days to about 56 days, about 28 days to about 42 days, of about 35 days to about 96 days, about 35 days to about 84 days, about 35 days to about 70 days, about 35 days to about 56 days, about 35 days to about 42 days, of about 42 days to about 96 days, about 42 days to about 84 days, about 42 days to about 70 days, or about 42 days to about 56 days in humans. In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life in a range of about 42 days to about 56 days in humans. In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life of at least about 50 days in humans. In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life of about 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56 or more than 56 days in humans.

(74) In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life of from about 10 days to about 25 days, from about 17 days to 23 days, from about 20 days to 22 days in non-human primates (NHPs). In some embodiments, the NHP is cynomolgus monkey. In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life of from about 11 days to 23 days in cynomolgus monkeys. In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life of from about 17 days to 23 days in cynomolgus monkeys. In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life of from about 19 days to 23 days in cynomolgus monkeys. In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life of from about 20 days to 23 days in cynomolgus monkeys. In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life of about 17 days in cynomolgus monkeys. In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life of about 20 days in cynomolgus monkeys. In some embodiments, the $\alpha\beta 7$ binding antibodies described herein comprise a half-life of about 22 days in

cynomolgus monkeys. In some embodiments, the $\alpha 4\beta 7$ binding antibodies described herein comprise a half-life of from about 5 days to about 15 days, or from about 10 to 12 days in transgenic mice expressing human neonatal Fc receptor (hFcRn). In some embodiments, the transgenic mice expressing hFcRn are Tg276 mice. In some embodiments, the $\alpha 4\beta 7$ binding antibodies described herein comprise a half-life of about 12 days in Tg276 mice.

(75) Methods of measuring half-life are known in the art. In some embodiments, the half-life is measured in a rodent model, such as Tg276 mice. In some embodiments, the half-life is measured in a non-human primate, such as cynomolgus monkeys. Other animals can also be used to measure PK of $\alpha 4\beta 7$ binding antibodies. In some embodiments, the half-life is measured in human. In some embodiments, the half-life is following intravenous administration. In some embodiments, the half-life is following subcutaneous administration.

(76) In some embodiments, the $\alpha 4\beta 7$ binding antibodies described herein have a half-life that is at least 20% longer than a comparator antibody. In some embodiments, the comparator antibody comprises the same complementarity determining regions and variable regions but different Fc regions. In some embodiments, the half-life of the $\alpha 4\beta 7$ binding antibodies described herein protein is at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% longer, at least 95% longer, at least 100% longer, at least 150% longer, at least 200% longer, at least 250% longer, at least 300% longer or at least 350% longer than the half-life of the comparator antibody. In some embodiments, the half-life of the $\alpha 4\beta 7$ binding antibodies described herein is longer than the half-life of the comparator antibody by at least 2 fold, at least 3 fold, at least 4 fold, at least 5 fold, at least 6 fold, at least 7 fold, at least 8 fold, at least 9 fold, or at least 10 fold.

(77) In some embodiments, the $\alpha 4\beta 7$ binding antibodies described herein bind $\alpha 4\beta 7$ with a $K_{sub.D}$ lower than or equal to 20 nM, 15 nM, 10 nM, 9 nM, 8 nM, 7 nM, 6 nM, 5 nM, 4 nM, 3 nM, 2 nM, 1 nM, 0.9 nM, 0.8 nM, 0.7 nM, 0.6 nM, 0.5 nM, 0.4 nM, 0.3 nM, 0.2 nM, 0.1 nM, 90 pM, 80 pM, 70 pM, 60 pM, 50 pM, 40 pM, 30 pM, 20 pM, or 10 pM. For example, in certain embodiments, the first antigen-binding site binds $\alpha 4\beta 7$ with a $K_{sub.D}$ within the range of about 10 pM-about 1 nM, about 10 pM-about 0.9 nM, about 10 pM-about 0.8 nM, about 10 pM-about 0.7 nM, about 10 pM-about 0.6 nM, about 10 pM-about 0.5 nM, about 10 pM-about 0.4 nM, about 10 pM-about 0.3 nM, about 10 pM-about 0.2 nM, about 10 pM-about 0.1 nM, about 10 pM-about 50 pM, 0.1 nM-about 10 nM, about 0.1 nM-about 9 nM, about 0.1 nM-about 8 nM, about 0.1 nM-about 7 nM, about 0.1 nM-about 6 nM, about 0.1 nM-about 5 nM, about 0.1 nM-about 4 nM, about 0.1 nM-about 3 nM, about 0.1 nM-about 2 nM, about 0.1 nM-about 1 nM, about 0.1 nM-about 0.5 nM, about 0.5 nM-about 10 nM, about 0.5 nM-about 9 nM, about 0.5 nM-about 8 nM, about 0.5 nM-about 7 nM, about 0.5 nM-about 6 nM, about 0.5 nM-about 5 nM, about 0.5 nM-about 4 nM, about 0.5 nM-about 3 nM, about 0.5 nM-about 2 nM, about 0.5 nM-about 1 nM, about 1 nM-about 10 nM, about 1 nM-about 9 nM, about 1 nM-about 8 nM, about 1 nM-about 7 nM, about 1 nM-about 6 nM, about 1 nM-about 5 nM, about 1 nM-about 4 nM, about 1 nM-about 3 nM, about 1 nM-about 2 nM, about 2 nM-about 10 nM, about 3 nM-about 10 nM, about 4 nM-about 10 nM, about 5 nM-about 10 nM, about 6 nM-about 10 nM, about 7 nM-about 10 nM, about 8 nM-about 10 nM, or about 9 nM-about 10 nM. In some embodiments, the affinity is measured by SPR. In some embodiments, the affinity is measured by BLI. In some embodiments, the affinity is measured by KinExA.

(78) Methods of Treatment and Dosage Regimens

(79) Described herein, in certain embodiments, are methods of treating a disease or disorder in a patient in need thereof, the method comprising subcutaneously or intravenously administering to the patient an effective amount of an $\alpha 4\beta 7$ binding antibody disclosed herein.

(80) Provided herein are methods of treating a disease or disorder in a patient in need thereof comprising subcutaneously administering to the patient one or more initial doses comprising an effective amount of the $\alpha 4\beta 7$ binding antibody (also referred to herein as “induction dose”, “induction regimen” or “induction therapy”) and one or more subsequent doses comprising an

effective amount of the $\alpha 4\beta 7$ binding antibody (also referred to herein as “maintenance dose”). (81) Also provided herein are methods of treating a disease or disorder in a patient in need thereof comprising intravenously administering to the patient one or more initial doses comprising an effective amount of the $\alpha 4\beta 7$ binding antibody (also referred to herein as “induction dose”, “induction regimen” or “induction therapy”) and one or more subsequent doses comprising an effective amount of the $\alpha 4\beta 7$ binding antibody (also referred to herein as “maintenance dose”). As used herein, “maintenance dose”, “maintenance regimen” and “maintenance therapy” are used interchangeably and is administered after induction dose(s) to continue the response achieved by induction dose(s) of $\alpha 4\beta 7$ binding antibody.

(82) In some embodiments, a single or more than one induction dose is administered, followed by maintenance doses. The disclosure contemplates one, two, three, or more induction doses. An induction dose can be administered intravenously or subcutaneously.

(83) The induction dosing can involve administration of a higher dose than a maintenance dose, more frequent administration than a maintenance dose, or both, of the $\alpha 4\beta 7$ binding antibody, e.g., in case potentially by a factor of 2. Alternatively, the induction dosing amount is the same as the maintenance dosing amount.

(84) In some embodiments, one or more induction dose(s) is greater than the maintenance dose. In some embodiments, the method comprises administering a single induction dose that is greater than the maintenance dose. For example, the induction dose can be 1.5, 2, 3, 4, 5 times higher than the maintenance dose. Alternatively, the one or more induction dose amount(s) is the same amount as the maintenance dose. In some embodiments, one or more induction dose(s) is administered subcutaneously to the patient in need thereof. Yet, in other embodiments, the one or more induction dose(s) is administered intravenously. In some embodiments, an induction dose may require one or more than one subcutaneous injection or infusion on the same day.

(85) In some embodiments, the method comprises administering one or more induction doses, wherein the induction doses are the same or greater than the maintenance dose and/or are administered at a frequency that is greater than the administration of the maintenance dose. For example, the method comprises administering one, two, three or four induction doses of the $\alpha 4\beta 7$ binding antibody. In some embodiments, a first dose is administered at day 0 and a second dose at a subsequent day, for example, day 1 to week 6, e.g. day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, day 14, days 21, day 28, day 35, day 42. In some embodiments, the one or more induction dose(s) may require one or more than one subcutaneous injection, or one or more than one infusion on the same day.

(86) In some embodiments, a total induction dosage amount corresponding to the sum of the one or more administrations is from about 400 mg to about 1000 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, a total induction dosage amount corresponding to the sum of the one or more administrations is from about 500 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the total dosage amount is about 400 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the total dosage amount is about 500 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the total dosage amount about 600 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the total dosage amount is about 700 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the dosage amount is about 800 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the total dosage amount is about 900 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the total dosage amount is about 1000 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the total dosage amount is about 1100 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the total dosage amount is about 1200 mg of the $\alpha 4\beta 7$ binding antibody.

(87) In some embodiments, the one or more induction dose(s) is administered subcutaneously to the patient in need thereof. Yet, in other embodiments, the one or more induction dose(s) is administered intravenously.

(88) In some embodiments, the maintenance regimen comprises lower doses than the induction

[illegible]

subcutaneously at day zero. In some embodiments, one or more induction dose(s) of about 850 mg of the $\alpha 4\beta 7$ binding antibody is administered intravenously or subcutaneously at day zero. In some embodiments, one or more induction dose(s) of about 900 mg of the $\alpha 4\beta 7$ binding antibody is administered intravenously or subcutaneously at day zero. In some embodiments, one or more induction dose(s) of about 950 mg of the $\alpha 4\beta 7$ binding antibody is administered intravenously or subcutaneously at day zero. In some embodiments, one or more induction dose(s) of about 1000 mg of the $\alpha 4\beta 7$ binding antibody is administered intravenously or subcutaneously at day zero. In some embodiments, one or more induction dose(s) of about 1050 mg of the $\alpha 4\beta 7$ binding antibody is administered intravenously or subcutaneously at day zero. In some embodiments, one or more induction dose(s) of about 1100 mg of the $\alpha 4\beta 7$ binding antibody is administered intravenously or subcutaneously at day zero. In some embodiments, one or more induction dose(s) of about 1150 mg of the $\alpha 4\beta 7$ binding antibody is administered intravenously or subcutaneously at day zero. In some embodiments, one or more induction dose(s) of about 1200 mg of the $\alpha 4\beta 7$ binding antibody is administered intravenously or subcutaneously at day zero. In some embodiments, the one or more induction dose(s) is administered subcutaneously.

(90) In some embodiments, the method comprises administering two or more induction doses, wherein the first induction dose is greater than the second induction dose, and wherein the two or more induction doses are administered at a frequency that is greater than the administration of the maintenance dose. In some embodiments, the first induction dose is greater than second induction dose and greater than the maintenance dose, and the second induction is the same or greater than the maintenance dose. In some embodiments, one or more induction dose(s) of from about 150 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody is administered intravenously or subcutaneously at day zero. In some embodiments, the method comprises administering intravenously or subcutaneously a first induction dose is that greater than the second induction dose of the $\alpha 4\beta 7$ binding antibody (e.g. a first induction dose of about 600 mg) at day zero, an administering a second induction of the $\alpha 4\beta 7$ binding antibody (e.g. a second induction dose of about 150 mg) at day 14, and administering a maintenance dose that is the same or lower than the second inductions dose (e.g. about 150 mg) every two weeks thereafter. In one example, the induction doses are about 600 mg/mL at week 0 and about 300 mg/mL at week 2, with maintenance doses of about 300 mg/mL every 8 to 12 weeks after the second induction dose. In one example, the induction doses are about 600 mg/mL at week 0 and about 300 mg/mL at week 2, and about 300 mg/mL at week 6. In some embodiments, administration of the induction doses can result in plasma or serum C.sub.avgW0-W6 of at least about 65 $\mu\text{g/mL}$, e.g. from about 65 $\mu\text{g/mL}$ to about 75 $\mu\text{g/mL}$ or greater, about 65 $\mu\text{g/mL}$, about 70 $\mu\text{g/mL}$, about 75 $\mu\text{g/mL}$ or more. In some embodiments, administration of the induction doses can result in plasma or serum C.sub.avgW0-W6 of about 70 $\mu\text{g/mL}$. In some embodiments, administration of the induction doses can result in plasma or serum C.sub.avgW0-W12 of at least about 60 $\mu\text{g/mL}$, e.g. from about 60 $\mu\text{g/mL}$ to about 70 $\mu\text{g/mL}$ or greater, about 60 $\mu\text{g/mL}$, about 65 $\mu\text{g/mL}$, about 70 $\mu\text{g/mL}$ or more. In some embodiments, administration of the induction doses can result in plasma or serum C.sub.avgW0-W12 of about 65 $\mu\text{g/mL}$. Maintenance doses can be about 300 mg/mL at least every 8 weeks, for example every 12 weeks or every 24 weeks after the final induction dose. In some embodiments, administration of the induction doses can result in a plasma or serum C.sub.avgW30-W40 of at least about 30 $\mu\text{g/mL}$, e.g. about 30 $\mu\text{g/mL}$, about 40 $\mu\text{g/mL}$, about 50 $\mu\text{g/mL}$, about 60 $\mu\text{g/mL}$, about 70 $\mu\text{g/mL}$ or more.

(91) In some embodiments, the plasma or serum concentration of at least about 35 $\mu\text{g/mL}$ (e.g. from 35 to 40 $\mu\text{g/mL}$, from 40 to 45 $\mu\text{g/mL}$, from 45 to 50 $\mu\text{g/mL}$, from 50 to 55 $\mu\text{g/mL}$, from 55 to 60 $\mu\text{g/mL}$, e.g. 36 $\mu\text{g/mL}$, 41 $\mu\text{g/mL}$, 46 $\mu\text{g/mL}$, 51 $\mu\text{g/mL}$, 56 $\mu\text{g/mL}$, 60 $\mu\text{g/mL}$ or more) is achieved in the patient for at least 4, at least 8, at least 12, at least 16, at least 20, at least 24 weeks or more (e.g. from 4 weeks to 6 weeks, from 6 weeks to 8 weeks, from 8 weeks to 10 weeks, from 10 weeks to 12 weeks, from 12 weeks to 14 weeks, from 14 weeks to 16 weeks, from 16 weeks to

18 weeks, from 18 weeks to 20 weeks, from 20 weeks to 22 weeks, from 22 weeks to 24 weeks or more) upon administration of an effective amount of the $\alpha 4\beta 7$ binding antibody during the induction phase.

(92) In some embodiments, the average plasma or serum concentration of at least about 35 $\mu\text{g/mL}$ (e.g. from 35 to 40 $\mu\text{g/mL}$, from 40 to 45 $\mu\text{g/mL}$, from 45 to 50 $\mu\text{g/mL}$, from 50 to 55 $\mu\text{g/mL}$, from 55 to 60 $\mu\text{g/mL}$, e.g. 36 $\mu\text{g/mL}$, 41 $\mu\text{g/mL}$, 46 $\mu\text{g/mL}$, 51 $\mu\text{g/mL}$, 56 $\mu\text{g/mL}$, 60 $\mu\text{g/mL}$ or more) is achieved in the patient over the course of the induction period. In some embodiments, the average plasma or serum concentration of at least about 45 $\mu\text{g/mL}$ is achieved in the patient over the course of the induction period. In some embodiments, the average plasma or serum concentration of at least about 60 $\mu\text{g/mL}$ is achieved in the patient over the course of the induction period.

(93) In some embodiments, a maintenance dose of the $\alpha 4\beta 7$ binding antibody is administered subcutaneously at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24 weeks or more (e.g. from 4 weeks to 6 weeks, from 6 weeks to 8 weeks, from 8 weeks to 10 weeks, from 10 weeks to 12 weeks, from 12 weeks to 14 weeks, from 14 weeks to 16 weeks, from 16 weeks to 18 weeks, from 18 weeks to 20 weeks, from 20 weeks to 22 weeks, from 22 weeks to 24 weeks or more) after the administration of the final induction dose. In some embodiments, the maintenance dose is lower than the induction dose (e.g. twice, 3 times or more lower than the induction dose). In some embodiments, the maintenance dose is the same as the induction dose. In some embodiments, a maintenance dose of the $\alpha 4\beta 7$ binding antibody is administered subcutaneously every 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26 or more weeks. In some embodiments, a maintenance dose of the $\alpha 4\beta 7$ binding antibody is administered subcutaneously every 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26 or more weeks. In some embodiments, the maintenance dose comprises an effective amount of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the maintenance dose comprises from about 150 mg to about 300 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, upon administration of the maintenance dose results, an effective minimum serum or plasma concentration of the $\alpha 4\beta 7$ binding antibody is achieved. In some embodiments, the plasma or serum concentration of at least about 6 $\mu\text{g/mL}$ (e.g. from about 6 $\mu\text{g/mL}$ to about 10 $\mu\text{g/mL}$, from about 6 $\mu\text{g/mL}$ to about 20 $\mu\text{g/mL}$, from about 6 $\mu\text{g/mL}$ to about 30 $\mu\text{g/mL}$ e.g. 6 $\mu\text{g/mL}$, 10 $\mu\text{g/mL}$, 15 $\mu\text{g/mL}$, 20 $\mu\text{g/mL}$, 30 $\mu\text{g/mL}$ or more) is achieved in the patient for at least 4, at least 8, at least 12, at least 16, at least 20, at least 24 weeks or more (e.g. from 4 weeks to 6 weeks, from 6 weeks to 8 weeks, from 8 weeks to 10 weeks, from 10 weeks to 12 weeks, from 12 weeks to 14 weeks, from 14 weeks to 16 weeks, from 16 weeks to 18 weeks, from 18 weeks to 20 weeks, from 20 weeks to 22 weeks, from 22 weeks to 24 weeks or more) upon the administration upon administration of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the plasma or serum concentration does not fall below 6 $\mu\text{g/mL}$ between maintenance dosing intervals. In some embodiments, the plasma or serum concentration does not fall below 10 $\mu\text{g/mL}$ between maintenance dosing intervals. In some embodiments, the plasma or serum concentration does not fall below 20 $\mu\text{g/mL}$ between maintenance dosing intervals. In some embodiments, the plasma or serum concentration does not fall below 30 $\mu\text{g/mL}$ between maintenance dosing intervals.

(94) In some embodiments, the method comprises administering subcutaneously to the subject in need thereof one or more induction dose(s) of an $\alpha 4\beta 7$ binding antibody, and administering subcutaneously to the subject in need thereof one or more maintenance dose(s) of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the induction dose comprises an amount to the $\alpha 4\beta 7$ binding antibody about 2 or more times higher than the maintenance dose. In some embodiments, the induction dose comprises from about 150 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody. For example, the induction dose comprises from about 400 mg to about 600 mg of the $\alpha 4\beta 7$ binding antibody and the maintenance dose of from about 150 mg to about 300 mg of the $\alpha 4\beta 7$ binding antibody. In another example, the induction dose comprises about 400 mg of the $\alpha 4\beta 7$ binding

antibody and wherein the maintenance dose of about 200 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the induction dose comprises from about 500 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody. For example, in some embodiments, the induction dose comprises about 600 mg of the $\alpha 4\beta 7$ binding antibody and the maintenance dose of about 300 mg of the $\alpha 4\beta 7$ binding antibody. In another example, the induction dose comprises about 1000 mg of the $\alpha 4\beta 7$ binding antibody and the maintenance dose of about 300 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the one or more induction dose comprises a total dosage amount of from about 150 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody. For example, the induction dose comprises one or more induction dose comprises a total dosage amount of from about 400 mg to about 600 mg of the $\alpha 4\beta 7$ binding antibody and the maintenance dose of from about 150 mg to about 300 mg of the $\alpha 4\beta 7$ binding antibody. In another example, the one or more induction dose comprises a total dosage amount of about 400 mg of the $\alpha 4\beta 7$ binding antibody and the maintenance dose of about 200 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the one or more induction dose comprises a total dosage amount of from about 500 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody. In another example the one or more induction dose comprises a total dosage amount of about 600 mg of the $\alpha 4\beta 7$ binding antibody and the maintenance dose of about 300 mg of the $\alpha 4\beta 7$ binding antibody. In another example, the one or more induction dose comprises a total dosage amount of about 1000 mg of the $\alpha 4\beta 7$ binding antibody and the maintenance dose of about 300 mg of the $\alpha 4\beta 7$ binding antibody.

(95) In some embodiments, upon administration of the one or more induction dose(s), an effective minimum average serum or plasma concentration of the $\alpha 4\beta 7$ binding antibody is achieved over the course of the induction period. In some embodiments, the average serum or plasma concentration of the $\alpha 4\beta 7$ binding antibody of at least about 35 $\mu\text{g/mL}$ is achieved in the patient over the course of the induction period, for example 36 $\mu\text{g/mL}$, is achieved in the patient over the course of the induction period. In some embodiments, the serum or plasma concentration of the $\alpha 4\beta 7$ binding antibody of at least about 35 $\mu\text{g/mL}$ is achieved in the patient for at least 12 weeks or more upon the administration of the induction dose. In some embodiments, the serum or plasma concentration of the $\alpha 4\beta 7$ binding antibody of at least about 6 $\mu\text{g/mL}$ (e.g. from about 6 $\mu\text{g/mL}$ to about 10 $\mu\text{g/mL}$ from about 6 $\mu\text{g/mL}$ to about 20 $\mu\text{g/mL}$, from about 6 $\mu\text{g/mL}$ to about 30 $\mu\text{g/mL}$ e.g. 6 $\mu\text{g/mL}$, 10 $\mu\text{g/mL}$, 15 $\mu\text{g/mL}$, 20 $\mu\text{g/mL}$, 30 $\mu\text{g/mL}$ or more) is achieved in the patient for at least 12 weeks upon the administration of the maintenance dose. In some embodiments, the serum or plasma concentration of the $\alpha 4\beta 7$ binding antibody of at least about 10 $\mu\text{g/mL}$ is achieved in the patient for at least 12 weeks upon the administration of the maintenance dose.

(96) Further described herein, in certain embodiments, are methods of treating a disease or disorder in a patient in need thereof comprising subcutaneously administering to the patient about 108 mg or about 300 mg of an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2 and a light chain comprising an amino acid sequence according to SEQ ID NO: 3; and wherein a plasma or serum concentration of the $\alpha 4\beta 7$ binding antibody of at least about 35 $\mu\text{g/mL}$ is achieved in the patient for at least 2 weeks or more upon the administration.

(97) In some embodiments, the method of treating a disease or disorder in a patient in need thereof comprises subcutaneously administering to the patient one or more induction doses and one or more maintenance doses of an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain comprising an amino acid sequence according to SEQ ID NO: 3; and wherein an average serum or plasma concentration of the $\alpha 4\beta 7$ binding antibody of at least about 60 $\mu\text{g/mL}$ is achieved in the patient during the induction period, and wherein a serum or plasma concentration of the $\alpha 4\beta 7$ binding antibody of at least about 10 $\mu\text{g/mL}$ is achieved in the patient for at least 12 weeks upon the administration of the maintenance dose.

(98) In some embodiments, the average plasma or serum concentration of at least about 35 $\mu\text{g/mL}$

is achieved in the patient during the induction period, for example from about 35 to about 60 $\mu\text{g/mL}$.

(99) In some embodiments, the plasma or serum concentration of at least about 35 $\mu\text{g/mL}$ is achieved in the patient for at least 2 weeks or more upon the administration of the one or more induction doses(s). In some embodiments, the plasma or serum concentration of at least about 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 48, 59, 60, or more than 60 $\mu\text{g/mL}$ is achieved in the patient for at least 2 weeks or more upon the administration of the one or more induction doses(s).

(100) In some embodiments, the plasma or serum concentration of at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, or more than 60 $\mu\text{g/mL}$ is achieved in the patient for at least 2 weeks or more upon the administration of the one or more induction dose(s) and/or one or more of the maintenance dose(s) of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the plasma or serum concentration of at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 100, 150, or more than 150 $\mu\text{g/mL}$ is achieved in the patient for at least 4 weeks or more upon administration of the one or more induction dose(s) and/or one or more of the maintenance dose(s) of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the plasma or serum concentration of at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 100, 150, or more than 150 $\mu\text{g/mL}$ is achieved in the patient for at least 6 weeks or more upon the administration of the one or more induction dose(s) and/or one or more of the maintenance dose(s) of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the plasma or serum concentration of at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 100, 150, or more than 150 $\mu\text{g/mL}$ is achieved in the patient for at least 8 weeks or more upon the administration of the one or more induction dose(s) and/or one or more of the maintenance dose(s) of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the plasma or serum concentration of at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 100, 150, or more than 150 $\mu\text{g/mL}$ is achieved in the patient for at least 10 weeks or more upon the administration of the one or more induction dose(s) and/or one or more of the maintenance dose(s) of the $\alpha 4\beta 7$ binding antibody.

(101) In some embodiments, the plasma or serum concentration of at least about 35 $\mu\text{g/mL}$ is achieved in at least about 80% of the patients for at least 2 weeks or more upon the administration of the one or more induction dose(s) of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the plasma or serum concentration of at least about 35 $\mu\text{g/mL}$ is achieved in at least about 85% of the patients for at least 2 weeks or more upon the administration of the one or more induction dose(s) of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the plasma or serum concentration of at least about 35 $\mu\text{g/mL}$ is achieved in at least about 90% of the patients for at least 2 weeks or more upon the administration of the one or more induction dose(s) of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the plasma or serum concentration of at least about 35 $\mu\text{g/mL}$ is achieved in at least about 95% of the patients for at least 2 weeks or more upon the administration of the one or more induction dose(s) of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the plasma or serum concentration of at least about 35 $\mu\text{g/mL}$ is achieved in at least about 99% of the patients for at least 2 weeks or more upon the administration of the one or more induction dose(s) of the $\alpha 4\beta 7$ binding antibody.

(102) In some embodiments, the plasma or serum concentration of at least about 60 $\mu\text{g/mL}$ is achieved in at least about 80% of the patients for at least 2 weeks or more upon the administration of the one or more induction doses(s). In some embodiments, the plasma or serum concentration of at least about 60 $\mu\text{g/mL}$ is achieved in at least about 85% of the patients for at least 2 weeks or more upon the administration of the one or more induction doses(s). In some embodiments, the plasma or serum concentration of at least about 60 $\mu\text{g/mL}$ is achieved in at least about 90% of the patients for at least 2 weeks or more upon the administration of the one or more induction doses(s). In some embodiments, the plasma or serum concentration of at least about 60 $\mu\text{g/mL}$ is achieved in at least about 95% of the patients for at least 2 weeks or more upon the administration of the one or more induction doses(s). In some embodiments, the plasma or serum concentration of at least about

60 µg/mL is achieved in at least about 99% of the patients for at least 2 weeks or more upon the administration of the one or more induction doses(s).

(103) In some embodiments, the average plasma or serum concentration of at least about 35 µg/mL is achieved in at least about 80% of the patients during the induction period. In some embodiments, the average plasma or serum concentration of at least about 35 µg/mL is achieved in at least about 85% of the patients during the induction period. In some embodiments, the average plasma or serum concentration of at least about 35 µg/mL is achieved in at least about 90% of the patients during the induction period. In some embodiments, the average plasma or serum concentration of at least about 35 µg/mL is achieved in at least about 95% of the patients during the induction period. In some embodiments, the average plasma or serum concentration of at least about 35 µg/mL is achieved in at least about 99% of the patients during the induction period.

(104) In some embodiments, the average plasma or serum concentration of at least about 60 µg/mL is achieved in at least about 80% of the patients during the induction period. In some embodiments, the average plasma or serum concentration of at least about 60 µg/mL is achieved in at least about 85% of the patients during the induction period. In some embodiments, the average plasma or serum concentration of at least about 60 µg/mL is achieved in at least about 90% of the patients during the induction period. In some embodiments, the average plasma or serum concentration of at least about 60 µg/mL is achieved in at least about 95% of the patients during the induction period. In some embodiments, the average plasma or serum concentration of at least about 60 µg/mL is achieved in at least about 99% of the patients during the induction period.

(105) Further described herein, in certain embodiments, are methods of treating a disease or disorder in a patient in need thereof comprising subcutaneously administering to the patient from about 75 mg to about 300 mg of an α4β7 binding antibody, wherein the α4β7 binding antibody comprises a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2 and a light chain comprising an amino acid sequence according to SEQ ID NO: 3; and wherein a maintenance C.sub.trough of the α4β7 binding antibody of at least about 6 µg/mL is achieved in the patient for at least 2 weeks or more upon the administration. In some embodiments, the α4β7 antibody is a long acting engineered α4β7 binding antibody consisting of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In a further embodiment, the dose and administration frequency provide for a maintenance C.sub.trough of the α4β7 binding antibody of at least about 6 µg/mL in the patient perpetually upon the administration, and the dosing is no more frequent than every 4 weeks, or alternatively every 8 weeks, or alternatively every 12 weeks, or alternatively every 16 weeks, or alternatively every 26 weeks. In an embodiment, the dose is sufficient for annual (Q52) dosing. In a further embodiment, the dose and administration frequency provide for a maintenance C.sub.trough of the α4β7 binding antibody of at least about 10 µg/mL in the patient perpetually upon the administration, and the dosing is no more frequent than every 4 weeks, or alternatively every 8 weeks, or alternatively every 12 weeks, or alternatively every 16 weeks, or alternatively every 26 weeks. The methods may be by intravenous (IV) or subcutaneous (SC) administration, or both (an IV induction dose or doses and SC maintenance dose or doses); a particular advantage of the disclosure is that it provides for subcutaneous administration of both the induction and maintenance doses.

(106) In some embodiments, the maintenance C.sub.trough of at least about 6 µg/mL (for example 10 µg/mL) is achieved in the patient upon administration of the one or more maintenance dose(s). In some embodiments, the maintenance C.sub.trough of at least about 2, 4, 6, 8, 10, 12, 14, 16, or more than 16 µg/mL is achieved in the patient for at least 2 weeks or more upon the administration. In some embodiments, the maintenance C.sub.trough of at least about 2, 4, 6, 8, 10, 12, 14, 16, or more than 16 µg/mL is achieved in the patient for at least 4 weeks or more upon administration. In some embodiments, the maintenance C.sub.trough of at least about 2, 4, 6, 8, 10, 12, 14, 16, or more than 16 µg/mL is achieved in the patient for at least 6 weeks or more upon the administration.

In some embodiments, the maintenance of at least about 2, 4, 6, 8, 10, 12, 14, 16, or more than 16 µg/mL is achieved in the patient for at least 8 weeks or more upon the administration. In some embodiments, the maintenance C.sub.trough of at least about 2, 4, 6, 8, 10, 12, 14, 16, or more than 16 µg/mL is achieved in the patient for at least 10 weeks or more upon the administration.

(107) In some embodiments, the maintenance C.sub.trough of at least about 6 µg/mL is achieved in at least about 80% of the patients for at least 2 weeks or more upon the administration. In some embodiments, the maintenance C.sub.trough of at least about 6 µg/mL is achieved in at least about 85% of the patients for at least 2 weeks or more upon the administration. In some embodiments, the maintenance C.sub.trough of at least about 6 µg/mL is achieved in at least about 90% of the patients for at least 2 weeks or more upon the administration. In some embodiments, the maintenance C.sub.trough of at least about 6 µg/mL is achieved in at least about 95% of the patients for at least 2 weeks or more upon the administration. In some embodiments, the maintenance C.sub.trough of at least about 6 µg/mL is achieved in at least about 99% of the patients for at least 2 weeks or more upon the administration.

(108) In some embodiments, the C.sub.6-wks, induction of at least about 30 µg/mL is achieved in the patient. In some embodiments, the C.sub.6-wks, induction of at least about 35 µg/mL is achieved in the patient. In some embodiments, the C.sub.6-wks, induction of at least about 20, 25, 30, 35, 40, 45, 50, 55, 60 or more than 60 µg/mL is achieved in the patient. In some embodiments, the C.sub.7-wks, induction of at least about 20, 25, 30, 35, 40, 45, 50, 55, 60 or more than 60 µg/mL is achieved in the patient. In some embodiments, the C.sub.8-wks, induction of at least about 20, 25, 30, 35, 40, 45, 50, 55, 60 or more than 60 µg/mL is achieved in the patient. In some embodiments, the C.sub.9-wks, induction of at least 20, 25, 30, 35, 40, 45, 50, 55, 60 or more than 60 µg/mL is achieved in the patient. In some embodiments, the C.sub.10-wks, induction of at least about 20, 25, 30, 35, 40, 45, 50, 55, 60 or more than 60 45 µg/mL is achieved in the patient. In some embodiments, the C.sub.11-wks, induction of at least about 20, 25, 30, 35, 40, 45, 50, 55, 60 or more than 60 µg/mL is achieved in the patient. In some embodiments, the C.sub.12-wks, induction of at least about 20, 25, 30, 35, 40, 45, 50, 55, 60 or more than 60 µg/mL is achieved in the patient.

(109) In some embodiments, the C.sub.6-wks, induction of at least about 35 µg/mL is achieved in at least about 80% of the patients. In some embodiments, the C.sub.6-wks, induction of at least about 35 µg/mL is achieved in at least about 85% of the patients. In some embodiments, the C.sub.6-wks, induction of at least about 35 µg/mL is achieved in at least about 90% of the patients. In some embodiments, the C.sub.6-wks, induction of at least about 35 µg/mL is achieved in at least about 95% of the patients. In some embodiments, the C.sub.6-wks, induction of at least about 35 µg/mL is achieved in at least about 99% of the patients.

(110) In some embodiments, the C.sub.6-wks, induction of at least about 60 µg/mL is achieved in at least about 80% of the patients. In some embodiments, the C.sub.6-wks, induction of at least about 60 µg/mL is achieved in at least about 85% of the patients. In some embodiments, the C.sub.6-wks, induction of at least about 60 µg/mL is achieved in at least about 90% of the patients. In some embodiments, the C.sub.6-wks, induction of at least about 60 µg/mL is achieved in at least about 95% of the patients. In some embodiments, the C.sub.6-wks, induction of at least about 60 µg/mL is achieved in at least about 99% of the patients. Provided herein is a method of achieving C.sub.avgW0-W12 of from 60 µg/mL to 70 µg/mL or greater for an α4β7 binding antibody in a subject in need thereof, the method comprising administering to the subject in need thereof a first dosage amount of at least 600 mg of the α4β7 binding antibody, wherein the α4β7 binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the α4β7 binding antibody consists of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3

(111) In some embodiments, administration results in a C.sub.avgW0-W12 of at least 60 µg/mL. In some embodiments, administration results in a C.sub.avgW0-W12 of about 65 µg/mL.

(112) In some embodiments, administration comprises administering to the subject in need thereof the first dosage amount of at least 600 mg of the α4β7 binding antibody and a second dosage amount of at least 300 mg of the α4β7 binding antibody at least two weeks after administration of the first dosage amount of the α4β7 binding antibody.

(113) In some embodiments, the method comprises administering the first dosage amount of at least 600 mg intravenously and the second dosage amount of at least 300 mg of the α4β7 binding antibody subcutaneously.

(114) In some embodiments, the method comprises the first dosage amount of at least 600 mg subcutaneously and the second dosage amount of at least 300 mg of the α4β7 binding antibody subcutaneously.

(115) In some embodiments, the method further comprises administering to the subject in need thereof one or more dosage amount of at least 300 mg of the α4β7 binding antibody at least four weeks after administration of the second dosage amount of the α4β7 binding antibody. In some embodiments, the one or more dosage amount of the α4β7 binding antibody is administered subcutaneously.

(116) In some embodiments, the method comprises administering to the subject in need thereof at least 600 mg of the α4β7 binding antibody every twenty-six weeks after administration of the second dosage amount of the α4β7 binding antibody.

(117) Provided herein is a method of achieving C.sub.avgW0-W6 of from 65 µg/mL to 75 µg/mL or greater for an α4β7 binding antibody in a subject in need thereof, the method comprising administering to the subject in need thereof a first dosage amount of at least 600 mg of the α4β7 binding antibody, wherein the α4β7 binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the α4β7 binding antibody consists of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(118) In some embodiments, administration results in a C.sub.avgW0-W6 of about 70 µg/mL. In some embodiments, administration results in a C.sub.avgW0-W12 of about 65 µg/mL.

(119) In some embodiments, the administering comprises administering to the subject in need thereof a first dosage amount of at least 600 mg of the α4β7 binding antibody and a second dosage amount of at least 300 mg of the α4β7 binding antibody at least two weeks after administration of the first dosage amount of at least 600 mg of the α4β7 binding antibody.

(120) In some embodiments, the method comprises administering the first dosage amount of the α4β7 binding antibody intravenously and the second dosage amount of the α4β7 binding antibody subcutaneously.

(121) In some embodiments, the method comprises administering the first dosage amount and the second dosage amount of the α4β7 binding antibody subcutaneously.

(122) In some embodiments, the method further comprises administering to the subject in need thereof one or more additional dosage amount of at least 300 mg of the α4β7 binding antibody at least four weeks after administration of the second dosage amount of the α4β7 binding antibody.

(123) In some embodiments, the method comprises administering the one or more additional dosage amount of the α4β7 binding antibody subcutaneously.

(124) In some embodiments, the method further comprises administering to the subject in need thereof at least 600 mg of the α4β7 binding antibody every twenty-six weeks after administration of second dosage amount of the α4β7 binding antibody.

(125) In some embodiments, the administration of the one or more dosage amount of the α4β7 binding antibody results in C.sub.avg of from 40 µg/mL to 50 µg/mL or greater. In some embodiments, the administration of the one or more dose of the α4β7 binding antibody results in

C.sub.avg of about 45 µg/mL.

(126) Provided herein is a method of achieving C.sub.trough of 35 µg/mL or greater for an α4β7 binding antibody in a subject in need thereof six weeks following administration of the α4β7 binding antibody to the subject, the method comprising administering to the subject in need thereof at least 600 mg of the α4β7 binding antibody, wherein the α4β7 binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the α4β7 binding antibody consists of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(127) In some embodiments, the administration of the at least 600 mg of the α4β7 binding antibody results in a serum C.sub.avgW0-W12 of at least 60 µg/mL. In some embodiments, the administration of the at least 600 mg of the α4β7 binding antibody results in a serum C.sub.avgW30-W40 of at least 30 µg/mL.

(128) In some embodiments, the at least 600 mg of the α4β7 binding antibody is administered intravenously. In other embodiments, the at least 600 mg of the α4β7 binding antibody is administered subcutaneously.

(129) In some embodiments, the method further comprises administering to the subject in need thereof at from about 120 to about 450 mg of the α4β7 binding antibody, wherein the administration of the from about 120 to about 450 mg of the α4β7 binding antibody results in a steady state C.sub.trough of at least 6 µg/mL. In some embodiments, the method further comprises administering to the subject in need thereof at least 120 mg of the α4β7 binding antibody, wherein the administration of the at least 120 mg of the α4β7 binding antibody results in a steady state C.sub.trough of at least 6 µg/mL. In some embodiments, the method further comprises administering to the subject in need thereof at least 300 mg of the α4β7 binding antibody, wherein the administration of the at least 300 mg of the α4β7 binding antibody results in a steady state C.sub.trough of at least 6 µg/mL.

(130) In some embodiments, the method further comprises administering to the subject in need thereof at least 300 mg of the α4β7 binding antibody resulting in an average serum concentration of the α4β7 binding antibody above 35 µg/mL for ten weeks following the administration of the at least 300 mg of the α4β7 binding antibody. In some embodiments, the method results in a serum C.sub.avgW0-W12 of at least 60 g/mL. In some embodiments, the administration of the at least 300 mg of the α4β7 binding antibody results in a serum C.sub.avgW30-W40 of at least 30 g/mL.

(131) In some embodiments, the method further comprises administering to the subject in need thereof at least 300 mg of the α4β7 binding antibody and results in an average serum concentration of the α4β7 binding antibody above 35 g/mL for ten weeks following the administration of the at least 300 mg of the α4β7 binding antibody.

(132) In some embodiments, the method further comprises administering subcutaneously to the subject in need thereof at least 300 mg of the α4β7 binding antibody four weeks after administration of last dosage amount of the α4β7 binding antibody.

(133) In some embodiments, the method further comprises administering subcutaneously to the subject a maintenance dose of at least 600 mg of the α4β7 binding antibody every twenty-six weeks after administration of the last dose of the α4β7 binding antibody.

(134) Provided herein are multidose regimen for use in treating a disease in a subject in need thereof comprising a first injectable liquid formulation comprising a total dosage amount of at least 500 mg of an α4β7 binding antibody; and a second injectable liquid formulation comprising a total dosage amount of at least 120 mg of the α4β7 binding antibody, wherein the α4β7 binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the α4β7 binding antibody consists of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence

according to SEQ ID NO: 3. In some embodiments, the second injectable liquid formulation comprises a total dosage amount of about 300 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the first injectable liquid formulation is for intravenous administration. In some embodiments, the first injectable liquid formulation is a single dose or a multidose formulation. In some embodiments, the second injectable liquid formulation is for subcutaneous administration. In some embodiments, the second injectable liquid formulation is a single dose or a multidose formulation.

(135) Provided herein are multidose regimen for use in treating a disease in a subject in need thereof comprising (a) a first injectable liquid formulation comprising a total dosage amount of at least 500 mg of a $\alpha 4\beta 7$ binding antibody; and (b) a second injectable liquid formulation comprising a total dosage amount of at least 300 mg of the $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody consists of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(136) In some embodiments, the first injectable liquid formulation comprises a total dosage amount of about 600 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the first injectable liquid formulation comprises a total dosage amount of about 1000 mg of the $\alpha 4\beta 7$ binding antibody.

(137) In some embodiments, the second injectable liquid formulation comprises a total dosage amount of about 300 mg of the $\alpha 4\beta 7$ binding antibody.

(138) In some embodiments, the first injectable liquid formulation is for intravenous administration. In some embodiments, the first injectable liquid formulation is a single dose or a multidose formulation. In some embodiments, the second injectable liquid formulation is for subcutaneous administration. In some embodiments, the second injectable liquid formulation is a single dose or a multidose formulation.

(139) Provided herein is a dosing regimen for use in treating a disease in a subject in need thereof, the dosing regimen comprising a first formulation comprising a total dosage amount of at least about 5500 mg of an $\alpha 4\beta 7$ binding antibody for administration to the subject; a second formulation comprising a total dosage amount of at least about 120 mg of the $\alpha 4\beta 7$ binding antibody for subcutaneous administration to the subject after the first formulation, and thereafter as a maintenance dose at least eight weeks after administration of the second formulation, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody consists of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the first injectable liquid formulation comprises a total dosage amount of from about 500 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the first injectable liquid formulation comprises a total dosage amount of about 600 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the first injectable liquid formulation comprises a total dosage amount of about 1000 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the second injectable liquid formulation comprises a total dosage amount of about 120 mg to about 450 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the second injectable liquid formulation comprises a total dosage amount of about 300 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the second formulation is administered at least two weeks after the first formulation, and thereafter as a maintenance dose at least eight weeks after administration of the second formulation.

(140) Provided herein is a dosing regimen for use in treating a disease in a subject in need thereof, the dosing regimen comprising: (a) a first formulation comprising a total dosage amount of at least about 600 mg of an $\alpha 4\beta 7$ binding antibody for administration to the subject; (b) a second

formulation comprising a total dosage amount of at least about 300 mg of the $\alpha 4\beta 7$ binding antibody for subcutaneous administration to the subject at least two weeks after the first formulation, and thereafter as a maintenance dose at least eight weeks after administration of the second formulation, wherein the $\alpha 4\beta 7$ binding antibody comprises: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody consists of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(141) In some embodiments, the first formulation is for subcutaneous administration. On other embodiments, the first formulation is for intravenous administration.

(142) In some embodiments, the first formulation comprises a total dosage amount of about 600 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the second formulation comprises a total dosage amount of about 300 mg of the $\alpha 4\beta 7$ binding antibody.

(143) In some embodiments, the first formulation dosing regimen comprises a single or multiple injections. In some embodiments, the second formulation dosing regimen comprises a single or multiple injections.

(144) In some embodiments, the first formulation and the second formulation do not contain citrate. In some embodiments, the first formulation or second formulation or both formulations comprise one or more of: histidine; arginine or a salt thereof; ethylenediaminetetraacetic acid (EDTA); and polysorbate 80.

(145) Provided herein is an injectable dosage form of an $\alpha 4\beta 7$ binding antibody comprising: a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3, wherein the $\alpha 4\beta 7$ binding antibody has an average serum half-life greater than 6 days in cynomolgus monkeys. In some embodiments, the $\alpha 4\beta 7$ binding antibody has an average serum half-life greater than 10 days in cynomolgus monkeys. In some embodiments, the $\alpha 4\beta 7$ binding antibody consists of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(146) In some embodiments, the average serum half-life is about 17 days or greater in cynomolgus monkeys. In some embodiments, the average serum half-life is 22 days in cynomolgus monkeys.

(147) In some embodiments, the injectable dosage form comprises at least about 100 mg/ml, at least about 150 mg/ml, at least about 180 mg/ml, or at least about 200 mg/ml of the $\alpha 4\beta 7$ binding antibody. For example, the injectable dosage form comprises from at least about 100 mg/ml to at least 110 mg/ml, from at least about 110 mg/ml to at least 120 mg/ml, from at least about 120 mg/ml to at least 130 mg/ml, from at least about 130 mg/ml to at least 140 mg/ml, from at least about 140 mg/ml to at least 150 mg/ml, from at least about 150 mg/ml to at least 160 mg/ml, from at least about 160 mg/ml to at least 170 mg/ml, from at least about 170 mg/ml to at least 180 mg/ml, from at least about 180 mg/ml to at least 190 mg/ml, from at least about 190 mg/ml to at least 200 mg/ml, from at least about 200 mg/ml to at least 210 mg/ml or more of the $\alpha 4\beta 7$ binding antibody.

(148) In some embodiments, the injectable dosage form comprises about 600 mg of the $\alpha 4\beta 7$ binding antibody. In other embodiments, the injectable dosage form comprises about 1000 mg of the $\alpha 4\beta 7$ binding antibody.

(149) In some embodiments, the injectable dosage form is an injectable liquid formulation. In some embodiments, the injectable liquid formulation does not comprise citrate. In some embodiments, the injectable liquid formulation comprises one or more of: histidine; arginine or a salt thereof; ethylenediaminetetraacetic acid (EDTA); and polysorbate 80.

(150) Provided herein are methods of treating a disease in a subject in need thereof, the method comprising administering to the subject in need thereof (a) an effective amount of an induction dose of an $\alpha 4\beta 7$ binding antibody, and (b) an effective amount of one or more maintenance doses

of the $\alpha 4\beta 7$ binding antibody, wherein the one or more maintenance doses are administered subcutaneously at least eight weeks apart. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody consists of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(151) In some embodiments, the effective amount of each of the one or more maintenance doses of the $\alpha 4\beta 7$ binding antibody is at least about 120 mg. In some embodiments, the effective amount of each of the one or more maintenance doses of the $\alpha 4\beta 7$ binding antibody is at least about 300 mg. In some embodiments, the effective amount of the induction dose of the $\alpha 4\beta 7$ binding antibody is about 600 mg. In some embodiments, the effective amount of the induction dose of the $\alpha 4\beta 7$ binding antibody is about 1000 mg. In some embodiments, the effective amount of the induction dose of the $\alpha 4\beta 7$ binding antibody is administered subcutaneously. In some embodiments, the effective amount of the induction dose of the $\alpha 4\beta 7$ binding antibody is administered intravenously. In some embodiments, the one or more maintenance doses are administered subcutaneously about 12 weeks apart. In some embodiments, the one or more maintenance doses are administered subcutaneously about 26 weeks apart. In some embodiments, the induction dose comprises a dosage amount of the $\alpha 4\beta 7$ binding antibody that is about 2 or more times higher than a dosage amount of each of the one or more maintenance doses.

(152) In some embodiments, the administration of the $\alpha 4\beta 7$ binding antibody follows a biphasic decline in serum concentration. In some embodiments, repeated dosing of $\alpha 4\beta 7$ binding antibody does not have a significant influence on clearance (CL) and steady-state volume of distribution (Vss). In some embodiments, pharmacokinetic is not impacted by disease state of the patient.

(153) In some embodiments, the $\alpha 4\beta 7$ binding antibody provided herein and the $\alpha 4\beta 7$ binding antibody in the regimen, injectable dosage form, method provided herein has one or more of the following characteristics: (a) an average serum half-life greater than about 6 days in cynomolgus monkeys, (b) an average serum half-life of 22 days in cynomolgus monkeys, (c) a melting temperature T_{mOnset} greater than about 55° C., (d) a melting temperature T_{mOnset} greater than 56.2° C., (e) a proportion of GOF from about 55.1% to about 60.4%, (f) a proportion of G1F from about 16.6% to about 18.8%, (g) a proportion of Man5 from about 7.3% to about 8.8%, (h) does not induce T cell activation markers CD25 or CD69, (i) does not result in release of cytokines at least 6 hours after the administration, wherein the cytokines comprise one or more of IL-6, IL-8, IL-10, IFN γ , IL-4, IL-17, IL-2, IL-23p70 and TNF, (j) does not induce complement-dependent cytotoxicity (CDC) in primary human PBMCs and in $\alpha 4\beta 7$ -expressing human β -lymphoid cell, (k) does not induce antibody dependent cellular cytotoxicity (ADCC) in human NK cells, and (l) does not impact suppressive activity of regulatory T cells expressing $\alpha 4\beta 7$ integrin, wherein the suppressive activity is measured by presence of elevated CD71, CD25, Ki67, granzyme B, or OX40.

(154) Described herein, in certain embodiments, are methods of treating a gastrointestinal inflammatory disease in a patient in need thereof. As used herein, the term “gastrointestinal inflammatory disease” refers to a disease of the gastrointestinal tract that involves inflammatory pathways. For example, the gastrointestinal inflammatory disease includes, but is not limited to, inflammatory bowel disease, ulcerative colitis (with or without exposure to anti-tumor necrosis factor (anti-TNF), Crohn's disease (including fistulizing Crohn's Disease), chronic pouchitis, collagenous gastritis, microscopic or collagenous colitis, colitis (including immune mediated colitis), sclerosing cholangitis (including in subjects with underlying inflammatory bowel disease, celiac enteritis, ileitis. In other aspects of the disclosure, provided herein are methods of treating Intestinal Acute Graft Versus Host Disease (aGVHD) (e.g. in subjects undergoing allogeneic hematopoietic stem cell transplantation (Allo-HSCT)), steroid-refractory acute intestinal graft-

versus-host disease (GVHD) (e.g. in subjects who have undergone Allo-HSCT), Type 1 diabetes (T1D) (e.g. with or without anti-TNF pre-treatment), immune checkpoint inhibitor-related colitis in subjects with genitourinary cancer or melanoma.

(155) In some embodiments, the disease is a gastrointestinal inflammatory disease. In some embodiments, the disease is an inflammatory bowel disease. In some embodiments, the inflammatory bowel disease is Crohn's disease. In some embodiments, the inflammatory bowel disease is ulcerative colitis.

(156) In some embodiments, the subject in need thereof has moderately to severely active ulcerative colitis. The Mayo score is a commonly used disease activity index in UC. The complete Mayo score is composed of four parts: rectal bleeding, stool frequency, physician assessment, and endoscopy appearance, each part being rated from 0 to 3, giving a total score of 0 to 12. In some embodiments, the subject in need thereof has a Mayo score 6 to 12 with endoscopic sub score ≥ 2 . In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody results in clinical response defined as a reduction in complete Mayo score of 3 or more points and 30% from baseline, (or a partial Mayo score reduction of 2 or more points and 25% or greater from baseline, if the complete Mayo score was not performed at the visit) with an accompanying decrease in rectal bleeding subscore of 1 or 0. In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody results in clinical remission, for example results in a complete Mayo score of 2 or less points and no individual subscore greater than 1 point. Alternatively, a modified or adapted Mayo score can be used for inclusion and for the clinical remission endpoint for UC. Clinical remission by the modified or adapted Mayo score is: stool frequency subscore 0 or 1 and not greater than baseline, rectal bleeding subscore of 0, and endoscopic subscore 0 or 1 without friability.

(157) In some embodiments, the subject in need thereof has moderately to severely active Crohn's disease. The Crohn's disease activity index (CDAI) is a commonly used disease activity index in Crohn's disease (CD) and is calculated from eight independent variables. Crohn's disease "clinical remission" refers to a CDAI score of 150 points or less. The American College of Gastroenterology and the European Crohn's and Colitis Organization define CDAI <150, 250-220, 220-450, and >450 as reflecting remission, mild disease activity, moderate disease activity, and severe disease activity, respectively. In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody results in clinical response defined as a ≥ 100 -point decrease in the CDAI score from baseline. In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody results in clinical remission defined as a CDAI score of 150 points or less.

(158) Described herein, in certain embodiments, are methods of treating an inflammatory bowel disease in a patient in need thereof, the method comprising subcutaneously or intravenously administering to the patient an effective amount of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3.

(159) In some embodiments, the inflammatory bowel disease is Crohn's disease or ulcerative colitis. In some embodiments, the inflammatory bowel disease is ulcerative colitis.

(160) Described herein, in certain embodiments, are methods of treating an inflammatory bowel disease in a patient in need thereof, the method comprising subcutaneously or intravenously administering to the patient an effective amount of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody is a long acting engineered $\alpha 4\beta 7$ binding antibody, consisting of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(161) In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 75 mg to about 150 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 150 mg to about 300 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a

dose of about 250 mg to about 750 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg to about 700 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 700 mg to about 900 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg to about 600 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg to about 500 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg to about 400 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 400 mg to about 700 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 400 mg to about 600 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg to about 500 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 500 mg to about 700 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 700 mg to about 900 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 500 mg to about 600 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 600 mg to about 700 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 75 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, about 250 mg, about 275 mg, or about 300 mg, about 350 mg, about 400 mg, about 450 mg, about 500 mg, about 550 mg, about 600 mg, about 650 mg, about 700 mg, about 750 mg, about 800 mg, about 850 mg, about 900 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered 30 at a dose of about 300 mg.

(162) In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody is intravenous, intratumoral, intramuscular, subcutaneous, intralesional, intrainestinal, intracolonic, intrarectal, intrapouch, or intraperitoneal. In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody is through a parenteral route such as intravenous, intramuscular, subcutaneous, intraarterial, or intraperitoneal administration. In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody is intravenous or subcutaneous. In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody is intravenous. In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody is subcutaneous.

(163) Administration of the $\alpha 4\beta 7$ binding antibody can occur at various intervals. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered to the patient at least once at an interval more than 8 weeks. In some embodiments, the interval is about 8 to about 12 weeks. In some embodiments, the interval is about 8 to about 16 weeks. In some embodiments, the interval is about 8 to about 20 weeks. In some embodiments, the interval is about 8 to about 24 weeks. In some embodiments, the interval is about 12 to about 26 weeks. In some embodiments, the interval is about 12 to about 22 weeks. In some embodiments, the interval is about 12 to about 18 weeks. In some embodiments, the interval is about 12 to about 14 weeks. In some embodiments, the interval is about 16 to about 26 weeks. In some embodiments, the interval is about 16 to about 22 weeks. In some embodiments, the interval is about 16 to about 18 weeks. In some embodiments, the interval is about 20 to about 26 weeks. In some embodiments, the interval is about 20 to about 22 weeks. In some embodiments, the interval is about 22 to about 26 weeks. In some embodiments, the interval is about 24 to about 26 weeks. In some embodiments, the interval is about 12 weeks. In some embodiments, the interval is about 14 weeks. In some embodiments, the interval is about 13 weeks. In some embodiments, the interval is about 16 weeks. In some embodiments, the interval is about 17 weeks. In some embodiments, the interval is about 18 weeks. In some embodiments, the interval is about 19 weeks. In some embodiments, the interval is about 20 weeks. In some embodiments, the interval is about 21 weeks. In some embodiments, the interval is about 22 weeks. In some embodiments, the interval is about 23 weeks. In some embodiments, the interval is about 24 weeks. In some embodiments, the interval is about 25 weeks. In some embodiments, the interval is about 26 weeks. In an embodiment, the dose is sufficient for annual (Q52) dosing.

(164) Described herein, in certain embodiments, are methods of treating a disease or disorder in a patient in need thereof comprising subcutaneously administering to the patient about 108 mg or

about 300 mg of an $\alpha 4\beta 7$ binding antibody every four or six weeks, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence according to SEQ ID NO: 4 and a light chain comprising an amino acid sequence according to SEQ ID NO: 5; and wherein no intervening subcutaneous administration of the $\alpha 4\beta 7$ binding antibody occurs during the four or six weeks. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, about 250 mg, about 275 mg, or about 300 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered every one week, two weeks, three weeks, four weeks, five weeks, six week, 7 weeks, 8 weeks, 10 weeks, 12 weeks, 14 weeks, 16 weeks, 20 weeks, 22 weeks, 24 weeks, 26 weeks, or more than 26 weeks.

(165) In some embodiments, the induction dose comprises from about 400 mg to about 1000 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of from about 150 to about 300 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the induction dose comprises about 600 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of about 300 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the induction dose comprises about 1000 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of about 300 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the induction dose comprises about 400 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of about 200 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the induction dose comprises about 600 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of about 600 mg of the $\alpha 4\beta 7$ binding antibody.

(166) In some embodiments, the method comprises administering subcutaneously to the subject an induction dose of about 600 mg of the $\alpha 4\beta 7$ binding antibody at week 0 and administering subcutaneously to the subject a maintenance dose of about 600 mg of the $\alpha 4\beta 7$ binding antibody at week 26.

(167) In some embodiments, upon administration of the one or more induction dose(s) an average serum or plasma concentration at least about 35 $\mu\text{g/mL}$ of the $\alpha 4\beta 7$ binding antibody is achieved in the patient during an induction period. In some embodiments, upon administration of the one or more maintenance dose(s) a serum or plasma concentration at least about 6 $\mu\text{g/mL}$, at least about 10 $\mu\text{g/mL}$, of the $\alpha 4\beta 7$ binding antibody is achieved in the patient in the patient for at least 2 weeks or more.

(168) Pharmaceutical Compositions

(169) The present disclosure also features pharmaceutical compositions that contain a therapeutically effective amount of the $\alpha 4\beta 7$ binding antibodies described herein. The composition can be formulated for use in a variety of drug delivery systems. One or more physiologically acceptable excipients or carriers can also be included in the composition for proper formulation. Suitable formulations for use in the present disclosure are found in Remington's Pharmaceutical Sciences, Mack Publishing Company, Philadelphia, Pa., 17th ed., 1985. For a brief review of methods for drug delivery, see, e.g., Langer (Science 249:1527-1533, 1990).

(170) In some embodiments, the pharmaceutical composition is a liquid composition. In some embodiments, the pharmaceutical composition is formulated for subcutaneous administration. In some embodiments, the pharmaceutical composition is formulated for intravenous administration. In some embodiments, the liquid composition is a colorless to brown; in some embodiments, the liquid composition is clear to opalescent. In some embodiments, the liquid composition contains at least 150 mg/ml, or at least 180 mg/ml, or at least 200 mg/ml of an $\alpha 4\beta 7$ binding antibody. In some embodiments, the liquid composition contains from about 150 mg/ml to about 160 mg/ml, from about 160 mg/ml to about 170 mg/ml, from about 170 mg/ml to about 10 mg/ml, from about 180 mg/ml to about 190 mg/ml, from about 190 mg/ml to about 200 mg/ml or more of a $\alpha 4\beta 7$ binding antibody.

(171) In some embodiments, a pharmaceutical composition may contain formulation materials for modifying, maintaining or preserving, for example, the pH, osmolarity, viscosity, clarity, color,

isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption or penetration of the composition. In such embodiments, suitable formulation materials include, but are not limited to, amino acids or salt thereof (such as glycine, glutamine, asparagine, arginine (e.g., arginine-HCl), histidine, or lysine); antimicrobials; antioxidants (such as ascorbic acid, sodium sulfite or sodium hydrogen-sulfite); buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates or other organic acids); bulking agents (such as mannitol or glycine); chelating agents (such as ethylenediamine tetraacetic acid (EDTA), pentetic acid (DTPA)); complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin); fillers; monosaccharides; disaccharides; and other carbohydrates (such as glucose, mannose or dextrans); proteins (such as serum albumin, gelatin or immunoglobulins); coloring, flavoring and diluting agents; emulsifying agents; hydrophilic polymers (such as polyvinylpyrrolidone); low molecular weight polypeptides; salt-forming counterions (such as sodium); preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid or hydrogen peroxide); solvents (such as glycerin, propylene glycol or polyethylene glycol); sugar alcohols (such as mannitol or sorbitol); suspending agents; surfactants or wetting agents (such as pluronics, PEG, sorbitan esters, polysorbates such as polysorbate 20, polysorbate 80, triton, tromethamine, lecithin, cholesterol, tyloxapal); stability enhancing agents (such as sucrose or sorbitol); tonicity enhancing agents (such as alkali metal halides, preferably sodium or potassium chloride, mannitol sorbitol); delivery vehicles; diluents; excipients and/or pharmaceutical adjuvants (see, Remington's Pharmaceutical Sciences, 18th ed. (Mack Publishing Company, 1990).

(172) In some embodiments, a pharmaceutical composition is citrate-free. In some embodiments, a pharmaceutical composition may contain nanoparticles, e.g., polymeric nanoparticles, liposomes, or micelles.

(173) In some embodiments, a pharmaceutical composition may contain a sustained- or controlled-delivery formulation. Techniques for formulating sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. Sustained-release preparations may include, e.g., porous polymeric microparticles or semipermeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained release matrices may include polyesters, hydrogels, polylactides, copolymers of L-glutamic acid and gamma ethyl-L-glutamate, poly (2-hydroxyethyl-methacrylate), ethylene vinyl acetate, or poly-D(-)-3-hydroxybutyric acid. Sustained release compositions may also include liposomes that can be prepared by any of several methods known in the art.

(174) Pharmaceutical compositions containing an $\alpha 4\beta 7$ binding antibody disclosed herein can be presented in a dosage unit form and can be prepared by any suitable method. A pharmaceutical composition should be formulated to be compatible with its intended route of administration. Examples of routes of administration are intravenous (IV), subcutaneous (SC), intradermal, inhalation, transdermal, topical, transmucosal, intrathecal and rectal administration. In some embodiments, the $\alpha 4\beta 7$ binding antibody disclosed herein is administered by intravenously. In some embodiments, the $\alpha 4\beta 7$ binding antibody disclosed herein is administered intravenously or subcutaneously. In specific embodiments, the $\alpha 4\beta 7$ binding antibody disclosed herein is administered subcutaneously.

(175) Aspects of the disclosure relate to injectable dosage forms of $\alpha 4\beta 7$ binding antibody. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody consists of a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the injectable dosage form is a liquid formulation. In some embodiments, the liquid formulation comprises one or more of histidine, arginine or a salt thereof, EDTA and polysorbate 80. In some

embodiments, the injectable liquid formulation does not comprise citrate.

(176) Described herein, in certain embodiments, are subcutaneous dosage forms comprising at least about 150 mg of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the subcutaneous dosage forms comprising at least about 300 mg of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3.

(177) In some embodiments, provided herein are subcutaneous dosage forms comprising at least about 150 mg of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. In some embodiments, provided herein are subcutaneous dosage forms comprising at least about 150 mg of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain consisting an amino acid sequence according to SEQ ID NO: 1; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3.

(178) In some embodiments, provided herein are subcutaneous dosage forms comprising at least about 300 mg of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. In some embodiments, provided herein are subcutaneous dosage forms comprising at least about 300 mg of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain consisting an amino acid sequence according to SEQ ID NO: 1; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3.

(179) Useful formulations can be prepared by methods known in the pharmaceutical art. For example, see *Remington's Pharmaceutical Sciences*, 18th ed. (Mack Publishing Company, 1990). Formulation components suitable for parenteral administration include a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerin, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as EDTA; buffers such as acetates, citrates or phosphates; and agents for the adjustment of tonicity such as sodium chloride or dextrose. In some embodiments, the formulation for parenteral administration is citrate-free.

(180) For intravenous or subcutaneous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor ELTM (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). The carrier should be stable under the conditions of manufacture and storage, and should be preserved against microorganisms. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol), and suitable mixtures thereof.

(181) An intravenous or subcutaneous drug delivery formulation may be contained in a syringe, pen, or bag. In some embodiments, the bag is connected to a channel comprising a tube and/or a needle. In some embodiments, the formulation is a lyophilized formulation or a liquid formulation. In some embodiments, the formulation is a liquid formulation. Various devices can be used to deliver liquid formulations by subcutaneous route of administration, including on-body infusion devices, autoinjector devices, prefilled syringes, and syringes. Generally, administration time depends on volume and device, and can range from seconds to minutes.

(182) These compositions may be sterilized by conventional sterilization techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as-is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration.

(183) A polyol, which acts as a tonicifier and may stabilize the $\alpha 4\beta 7$ binding antibody, may also be included in the formulation. The polyol is added to the formulation in an amount which may vary with respect to the desired isotonicity of the formulation. In some embodiments, the aqueous

formulation is isotonic. The amount of polyol added may also be altered with respect to the molecular weight of the polyol. For example, a lower amount of a monosaccharide (e.g., mannitol) is added, compared to a disaccharide (such as trehalose). In some embodiments, the polyol which is used in the formulation as a tonicity agent is mannitol.

(184) A detergent or surfactant may also be added to the formulation. Exemplary detergents include nonionic detergents such as polysorbates (e.g., polysorbates 20, 80 etc.) or poloxamers (e.g., poloxamer 188). The amount of detergent added is such that it reduces aggregation of the formulated antibody and/or minimizes the formation of particulates in the formulation and/or reduces adsorption. In some embodiments, the formulation may include a surfactant which is a polysorbate. In some embodiments, the formulation may contain the detergent polysorbate 80 or Tween 80. Tween 80 is a term used to describe polyoxyethylene (20) sorbitanmonooleate (see Fiedler, Lexikon der Hilfsstoffe, Editio Cantor Verlag Aulendorf, 4th ed., 1996).

(185) In embodiments, the protein product of the present disclosure is formulated as a liquid formulation. In some embodiments, the liquid formulation is prepared in combination with a sugar at stabilizing levels. In some embodiments, the liquid formulation is prepared in an aqueous carrier. In some embodiments, a stabilizer is added in an amount no greater than that which may result in a viscosity undesirable or unsuitable for intravenous administration. In some embodiments, the sugar is disaccharides, e.g., sucrose. In some embodiments, the liquid formulation may also include one or more of a buffering agent, a surfactant, and a preservative.

(186) In some embodiments, the pH of the liquid formulation is set by addition of a pharmaceutically acceptable acid and/or base. In some embodiments, the pharmaceutically acceptable acid is hydrochloric acid. In some embodiments, the base is sodium hydroxide. In some embodiments, the pH of the liquid formulation is from about 3.5 to 9, 5.5 to 6.5, for example 6.0. In some embodiments, the pH of the liquid formulation is about 6.

(187) The aqueous carrier of interest herein is one which is pharmaceutically acceptable (safe and non-toxic for administration to a human) and is useful for the preparation of a liquid formulation. Illustrative carriers include sterile water for injection (SWFI), bacteriostatic water for injection (BWFI), a pH buffered solution (e.g., phosphate-buffered saline), sterile saline solution, Ringer's solution or dextrose solution.

(188) A preservative may be optionally added to the formulations herein to reduce bacterial action. The addition of a preservative may, for example, facilitate the production of a multi-use (multiple-dose) formulation.

(189) The $\alpha 4\beta 7$ binding antibody may be lyophilized to produce a lyophilized formulation including the proteins and a lyoprotectant. The lyoprotectant may be sugar, e.g., disaccharides. In some embodiments, the lyoprotectant is sucrose or maltose. The lyophilized formulation may also include one or more of a buffering agent, a surfactant, a bulking agent, and/or a preservative.

(190) The amount of sucrose or maltose useful for stabilization of the lyophilized drug product may be in a weight ratio of at least 1:2 protein to sucrose or maltose. In some embodiments, the protein to sucrose or maltose weight ratio is of from 1:2 to 1:5. In some embodiments, the pH of the formulation, prior to lyophilization, is set by addition of a pharmaceutically acceptable acid and/or base. In some embodiments, the pharmaceutically acceptable acid is hydrochloric acid. In some embodiments, the pharmaceutically acceptable base is sodium hydroxide.

(191) In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 75 mg to about 150 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 250 mg to about 750 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg to about 700 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg to about 600 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 700 mg to about 900 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 600 mg to about 1200 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg to about 500

mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg to about 400 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 400 mg to about 700 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 400 mg to about 600 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg to about 500 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 500 mg to about 700 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 500 mg to about 600 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 600 mg to about 700 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 75 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, about 250 mg, about 275 mg, or about 300 mg, about 350 mg, about 400 mg, about 450 mg, about 500 mg, about 550 mg, about 600 mg, about 650 mg, about 700 mg, about 750 mg, about 800 mg, about 850 mg, about 900 mg or more. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg. Actual dosage levels of the active ingredients in the pharmaceutical compositions of this disclosure may be about 250 mg, 350 mg, 400 mg, 450 mg, 500 mg, 550 mg, 600 mg, 650 mg, 700 mg, 750 mg, 800 mg, 850 mg, 900 mg, 950 mg, 1000 mg, 1050 mg, 1100 mg, 1150 mg, or 1200 mg so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

(192) The specific dose can be a uniform dose for each patient of about 150 mg, of about 200 mg, of about 250 mg, about 300 mg, about 350 mg, about 400 mg, about 450 mg, about 500 mg, about 550 mg, about 600 mg, about 650 mg, about 700 mg, about 750 mg, about 800 mg, about 850 mg, about 900 mg, about 1000 mg, about 1100 mg, about 1200 mg or more of $\alpha 4\beta 7$ binding antibody. Alternatively, a patient's dose can be tailored to the approximate body weight or surface area of the patient. Other factors in determining the appropriate dosage can include the disease or condition to be treated or prevented, the severity of the disease, the route of administration, and the age, sex and medical condition of the patient. Further refinement of the calculations necessary to determine the appropriate dosage for treatment is routinely made by those skilled in the art, especially in light of the dosage information and assays disclosed herein. The dosage can also be determined through the use of known assays for determining dosages used in conjunction with appropriate dose-response data. An individual patient's dosage can be adjusted as the progress of the disease is monitored. Blood levels of the targetable construct or complex in a patient can be measured to see if the dosage needs to be adjusted to reach or maintain an effective concentration. Pharmacogenomics may be used to determine which targetable constructs and/or complexes, and dosages thereof, are most likely to be effective for a given individual (Schmitz et al., *Clinica Chimica Acta* 308: 43-53, 2001; Steimer et al., *Clinica Chimica Acta* 308: 33-41, 2001).

(193) Kits

(194) Provided herein is a kit for treating a disease in a subject in need thereof comprising: (a) an injectable liquid formulation comprising an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3, and (b) instructions to administer from about 500 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody, instructions to administer from about 120 mg to about 450 mg of the $\alpha 4\beta 7$ binding antibody at least two weeks after the administration of the about 500 mg to about 1200 mg of the $\alpha 4\beta 7$ binding antibody, and instructions to administer from about 120 mg to about 450 mg of the $\alpha 4\beta 7$ binding antibody at least four weeks thereafter. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(195) Provided herein is a kit for treating a disease in a subject in need thereof comprising: (a) an injectable liquid formulation comprising an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding

antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3, and (b) instructions to administer at least 600 mg of the $\alpha 4\beta 7$ binding antibody, at least 300 mg of the $\alpha 4\beta 7$ binding antibody at least two weeks after the administration of the 600 mg of the $\alpha 4\beta 7$ binding antibody, and at least 300 mg of the $\alpha 4\beta 7$ binding antibody at least four weeks thereafter. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(196) Provided herein is a kit for treating a disease in a subject in need thereof comprising: (a) an injectable subcutaneous liquid formulation comprising an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3, and (b) instructions to administer subcutaneously at least 600 mg of $\alpha 4\beta 7$ binding antibody, at least 300 mg of the $\alpha 4\beta 7$ binding antibody at least two weeks after the administration of the 600 mg of $\alpha 4\beta 7$ binding antibody, and at least 300 mg of $\alpha 4\beta 7$ binding antibody at least four weeks thereafter. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(197) Methods of Preparation

(198) The $\alpha 4\beta 7$ binding antibodies described above can be made using recombinant DNA technology well known to a skilled person in the art. For example, one or more isolated polynucleotides encoding the $\alpha 4\beta 7$ binding antibody can be ligated to other appropriate nucleotide sequences, including, for example, constant region coding sequences, and expression control sequences, to produce conventional gene expression constructs (i.e., expression vectors) encoding the desired $\alpha 4\beta 7$ binding antibodies. Production of defined gene constructs is within routine skill in the art.

(199) Nucleic acids encoding desired $\alpha 4\beta 7$ binding antibodies can be incorporated (ligated) into expression vectors, which can be introduced into host cells through conventional transfection or transformation techniques. Exemplary host cells are *E. coli* cells, Chinese hamster ovary (CHO) cells, human embryonic kidney 293 (HEK 293) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), and myeloma cells that do not otherwise produce IgG protein. Transformed host cells can be grown under conditions that permit the host cells to express the genes that encode $\alpha 4\beta 7$ binding antibodies. In some embodiments, the nucleic acid comprises a sequence as set forth in SEQ ID NOs: 12-17.

(200) Specific expression and purification conditions will vary depending upon the expression system employed. For example, if a gene is to be expressed in *E. coli*, it is first cloned into an expression vector by positioning the engineered gene downstream from a suitable bacterial promoter, e.g., Trp or Tac, and a prokaryotic signal sequence. The expressed protein may be secreted. The expressed protein may accumulate in refractile or inclusion bodies, which can be harvested after disruption of the cells by French press or sonication. The refractile bodies then are solubilized, and the protein may be refolded and/or cleaved by methods known in the art.

(201) If the engineered gene is to be expressed in eukaryotic host cells, e.g., CHO cells, it is first inserted into an expression vector containing a suitable eukaryotic promoter, a secretion signal, a poly A sequence, and a stop codon. Optionally, the vector or gene construct may contain enhancers and introns. In embodiments involving fusion proteins comprising an $\alpha 4\beta 7$ binding antibody or portion thereof, the expression vector optionally contains sequences encoding all or part of a constant region, enabling an entire, or a part of, a heavy or light chain to be expressed. The gene construct can be introduced into eukaryotic host cells using conventional techniques.

(202) In some embodiments, in order to express an $\alpha 4\beta 7$ binding antibody, an N-terminal signal

sequence is included in the protein construct. Exemplary N-terminal signal sequences include signal sequences from interleukin-2, CD-5, IgG kappa light chain, trypsinogen, serum albumin, and prolactin.

(203) After transfection, single clones can be isolated for cell bank generation using methods known in the art, such as limited dilution, ELISA, FACS, microscopy, or Clonepix. Clones can be cultured under conditions suitable for bio-reactor scale-up and maintained expression of the $\alpha 4\beta 7$ binding antibodies.

(204) The $\alpha 4\beta 7$ binding antibodies can be isolated and purified using methods known in the art including centrifugation, depth filtration, cell lysis, homogenization, freeze-thawing, affinity purification, gel filtration, ion exchange chromatography, hydrophobic interaction exchange chromatography, and mixed-mode chromatography.

(205) Additional Embodiments

(206) Described herein, in certain embodiments, are subcutaneous dosage forms comprising an effective amount of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the effective amount achieves a serum or plasma average concentration of about 35 $\mu\text{g/mL}$ and a trough serum or plasma concentration of at least about 6 $\mu\text{g/mL}$, e.g., about 10 $\mu\text{g/mL}$. In some embodiments, the heavy chain comprises an amino acid sequence according to SEQ ID NO: 1 and the light chain comprises an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the heavy chain comprises an amino acid sequence according to SEQ ID NO: 2 and the light chain comprises an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the subcutaneous dosage forms comprise from about 75 mg to about 900 mg or more of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the subcutaneous dosage forms comprise about 75 mg, 100 mg, 150 mg, 200 mg, 350 mg, 400 mg, 450 mg, 500 mg, 550 mg, 600 mg, 650 mg, 700 mg, 750 mg, 800 mg, 850 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg or more of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the subcutaneous dosage forms comprise about 300 mg to about 1200 mg for an induction dose and about 150 mg to 350 mg for a maintenance dose of the $\alpha 4\beta 7$ binding antibody.

(207) Described herein, in certain embodiments, are methods of administering one or more of the dosage described herein. More particularly, the methods of administering one or more of the dosage forms described herein provide for dosing no more frequently than every eight weeks (Q8 dosing), or no more frequently than every twelve weeks (Q12 dosing), or no more frequently than every sixteen weeks (Q16 dosing), and in some embodiments no more frequently than twice per year (Q26 dosing) for maintenance therapy after one to three doses for induction therapy. In some embodiments, the dose is sufficient for annual (Q52) dosing. The methods of administering may further include an induction dose containing 2 or more times the dose of an $\alpha 4\beta 7$ binding antibody of the disclosure. The methods may be by intravenous (IV) or subcutaneous (SC) administration, or both (an IV induction dose or doses and SC maintenance dose); a particular advantage of the disclosure is that it provides for subcutaneous administration of both the induction and maintenance doses.

(208) Described herein, in certain embodiments, are methods of treating an inflammatory bowel disease in a patient in need thereof, the method comprising subcutaneously or intravenously administering to the patient an effective amount of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the inflammatory bowel disease is Crohn's disease or ulcerative colitis. In some embodiments, the inflammatory bowel disease is ulcerative colitis. In some embodiments, the heavy chain comprises an amino acid sequence according to SEQ ID NO: 1 and the light chain comprises an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the heavy chain comprises an amino acid sequence according to SEQ ID NO: 2 and the light chain comprises

an amino acid sequence according to SEQ ID NO: 3. In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody is subcutaneous. In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody is intravenous. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of from about 100 mg to about 900 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of from about 100 mg to about 600 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of from about 100 mg to about 300 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 350 mg, 400 mg, 450 mg, 500 mg, 550 mg, 600 mg, 650 mg, 700 mg, 750 mg, 800 mg, 850 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg or more.

(209) Described herein, in certain embodiments, are methods of treating a disease or disorder in a patient in need thereof, the method comprising administering to the patient at least once at an interval more than 8 weeks, an effective amount of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the interval is about 12 to about 26 weeks. In some embodiments, the interval is about 12 weeks. In some embodiments, the interval is about 16 weeks. In some embodiments, the interval is about 26 weeks. In some embodiments, the disease or disorder is an inflammatory bowel disease. In some embodiments, the inflammatory bowel disease is Crohn's disease or ulcerative colitis. In some embodiments, the inflammatory bowel disease is ulcerative colitis. In some embodiments, the heavy chain comprises an amino acid sequence according to SEQ ID NO: 1 and the light chain comprises an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the heavy chain comprises an amino acid sequence according to SEQ ID NO: 2 and the light chain comprises an amino acid sequence according to SEQ ID NO: 3. In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody is subcutaneous. In some embodiments, administration of the $\alpha 4\beta 7$ binding antibody is intravenous. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg. In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 350 mg, 400 mg, 450 mg, 500 mg, 550 mg, 600 mg, 650 mg, 700 mg, 750 mg, 800 mg, 850 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg or more. In some embodiments, the disease or disorder is an inflammatory bowel disease and administration of the $\alpha 4\beta 7$ binding antibody is at an induction dose of about 600 mg, either by intravenous or subcutaneous administration, and a maintenance dose of about 300 mg and is subcutaneous.

(210) Described herein, in certain embodiments, are methods of treating a disease or disorder in a patient in need thereof comprising administering to the patient about 108 mg or about 300 mg of an $\alpha 4\beta 7$ binding antibody every four or six weeks, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence according to SEQ ID NO: 4 and a light chain comprising an amino acid sequence according to SEQ ID NO: 5; and wherein no intervening subcutaneous administration of the $\alpha 4\beta 7$ binding antibody occurs during the four or six weeks.

(211) Described herein, in certain embodiments, are methods of treating a disease or disorder in a patient in need thereof comprising administering to the patient about 108 mg or about 300 mg of an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2 and a light chain comprising an amino acid sequence according to SEQ ID NO: 3; and wherein a serum or plasma concentration of the $\alpha 4\beta 7$ binding antibody of at least about 35 $\mu\text{g/mL}$ is achieved in the patient for at least 2 weeks or more upon the administration.

(212) Described herein, in certain embodiments, are methods of treating a disease or disorder in a patient in need thereof comprising subcutaneously administering to the patient an induction dose of about 150 mg to about 1200 mg, for example about 300 mg to about 1000 mg, of an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2 and a light chain comprising an amino acid

sequence according to SEQ ID NO: 3; and wherein a serum or plasma concentration of the $\alpha 4\beta 7$ binding antibody of at least about 35 $\mu\text{g/mL}$ is achieved in the patient for at least 2 weeks or more upon the administration and a trough serum or plasma concentration is at least about 6 $\mu\text{g/mL}$, for example 10 $\mu\text{g/mL}$.

(213) Described herein, in certain embodiments, are methods of treating a disease or disorder in a patient in need thereof comprising subcutaneously administering to the patient about 108 mg or about 300 mg of an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2 and a light chain comprising an amino acid sequence according to SEQ ID NO: 3; and wherein a maintenance C.sub.trough of the $\alpha 4\beta 7$ binding antibody of at least about 6 $\mu\text{g/mL}$, e.g., about 10 $\mu\text{g/mL}$, is achieved in the patient upon the administration. In a further embodiment, the dose and administration frequency provide for a C.sub.trough of the $\alpha 4\beta 7$ binding antibody of at least about 6 $\mu\text{g/mL}$, e.g., about 10 $\mu\text{g/mL}$, in the patient perpetually upon the administration, and the dosing is no more frequent than every 4 weeks, or alternatively every 8 weeks, or alternatively every 12 weeks, or alternatively every 16 weeks, or alternatively every 26 weeks.

(214) In one embodiment, a single or more than one induction dose is administered, followed by maintenance doses. An induction dose can be administered intravenously or subcutaneously. The induction dose may be higher than a maintenance dose or more frequent than a maintenance dose, e.g., in case potentially by a factor of 2. Aspects of the disclosure contemplate one, two, three, or more induction doses. In one example, the induction doses are 600 mg/mL at week 0 and 300 mg/mL at week 2, with maintenance doses of 300 mg/mL every 12 weeks after the second induction dose.

(215) Aspects of the disclosure relate to isolated nucleic acids, expression vector comprising the nucleic acid sequences or recombinant host cell comprising the nucleic acid sequences encoding the heavy chain and/or light chain of the $\alpha 4\beta 7$ binding antibodies described herein. In some embodiments, the nucleic acid sequence encodes SEQ ID NO: 1. In some embodiments, the nucleic acid sequence encodes SEQ ID NO: 3.

(216) Aspects of the disclosure relate to an $\alpha 4\beta 7$ binding antibody comprising a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain comprising an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the $\alpha 4\beta 7$ binding antibody comprises a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

(217) Aspects of the disclosure relate to an $\alpha 4\beta 7$ binding antibody comprising a heavy chain and a light chain, wherein the heavy chain consists of an amino acid sequence according to SEQ ID NO: 1. In some embodiments, the light chain consists of an amino acid sequence according to SEQ ID NO: 3.

(218) According to aspects of the disclosure, the $\alpha 4\beta 7$ binding antibody has an increased half-life as compared to an $\alpha 4\beta 7$ binding antibody comprising a heavy chain sequence consisting of SEQ ID NO: 4. In some embodiments, the $\alpha 4\beta 7$ binding antibody has a half-life of from 42 days to 56 days or more in humans.

(219) Aspects of the disclosure relate to a pharmaceutical composition comprising an $\alpha 4\beta 7$ binding antibody and a pharmaceutically acceptable carrier, the $\alpha 4\beta 7$ binding antibody comprising a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain comprising an amino acid sequence according to SEQ ID NO: 3. In some embodiments, pharmaceutical composition comprises an $\alpha 4\beta 7$ binding antibody and a pharmaceutically acceptable carrier, the $\alpha 4\beta 7$ binding antibody comprising a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the pharmaceutical composition is citrate free.

(220) Aspects of the disclosure relate to a subcutaneous dosage form comprising at least about 150 mg of an $\alpha 4\beta 7$ binding antibody comprising a heavy chain consisting of an amino acid sequence

according to SEQ ID NO: 1; and a light chain comprising an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the subcutaneous dosage form comprises at least about 150 mg of an $\alpha 4\beta 7$ binding antibody comprising a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. In some embodiments, the dosage form is citrate free. In some embodiments, the subcutaneous dosage comprises from about 150 mg to about 900 mg of the $\alpha 4\beta 7$ binding antibody. (221) Methods for treating inflammatory bowel disease in a subject in need thereof are provided. In some embodiments, the method comprises administering subcutaneously to the subject one or more of the dosage forms of aspects the disclosure. In some embodiments, the inflammatory bowel disease is Crohn's disease or ulcerative colitis. In some embodiments, the inflammatory bowel disease is ulcerative colitis.

(222) In some embodiments, a method of treating an inflammatory bowel disease in a subject in need thereof comprises administering subcutaneously to the subject an effective amount of an $\alpha 4\beta 7$ binding antibody comprising a heavy chain and a light chain, wherein the heavy chain consists of an amino acid sequence according to SEQ ID NO: 1; and the light chain comprises the variable region of SEQ ID NO: 3.

(223) In some embodiments, a method for treating inflammatory bowel disease in a subject in need thereof, the method comprising administering subcutaneously to the subject an effective amount of an $\alpha 4\beta 7$ binding antibody comprising a heavy chain and a light chain, wherein the heavy chain consists of an amino acid sequence according to SEQ ID NO: 1; and the light chain consists of the variable region of SEQ ID NO: 3.

(224) In some embodiments, the $\alpha 4\beta 7$ binding antibody is administered at a dose of from about 150 mg to about 900 mg. In some embodiments, the method comprises administering to the subject the effective amount of an $\alpha 4\beta 7$ binding antibody at an interval of more than 8 weeks, for example about 12 to about 26 weeks, about 12 weeks, about 16 weeks, about 26 weeks.

(225) In some embodiments, the inflammatory bowel disease is Crohn's disease or ulcerative colitis. In some embodiments, the inflammatory bowel disease is ulcerative colitis.

(226) In some embodiments, an average serum or plasma or serum concentration of the $\alpha 4\beta 7$ binding antibody of at least about 35 $\mu\text{g/mL}$ is achieved in the subject during an induction period.

(227) In some embodiments, a maintenance C.sub.trough of the $\alpha 4\beta 7$ binding antibody of at least about 6 $\mu\text{g/mL}$ is achieved in the subject.

(228) Aspects of the disclosure relate to a method for treating inflammatory bowel disease in a subject in need thereof, the method comprising (a) administering subcutaneously to the subject one or more induction dose comprising an effective amount of an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising a heavy chain and a light chain, wherein the heavy chain consists of an amino acid sequence according to SEQ ID NO: 1; and the light chain consists of the variable region of SEQ ID NO: 3; (b) administering subcutaneously to the subject one or more maintenance dose comprising an effective amount of the $\alpha 4\beta 7$ binding antibody.

(229) In some embodiments, the one or more maintenance dose of the $\alpha 4\beta 7$ binding antibody is administered at an interval of more than 8 weeks.

(230) In some embodiments, the induction dose comprises an amount to the $\alpha 4\beta 7$ binding antibody about 2 or more times higher than the maintenance dose.

(231) In some embodiments, the induction dose comprises from about 400 mg to about 600 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of from about 150 to about 300 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the induction dose comprises about 600 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of about 300 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the induction dose comprises about 400 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of about 200 mg of the $\alpha 4\beta 7$ binding antibody. In some embodiments, the induction dose comprises about 600 mg of the $\alpha 4\beta 7$ binding antibody and

wherein the maintenance dose of about 600 mg of the $\alpha 4\beta 7$ binding antibody.

(232) In some embodiments, the method comprises administering subcutaneously to the subject an induction dose of about 600 mg of the $\alpha 4\beta 7$ binding antibody at week 0 and administering subcutaneously to the subject a maintenance dose of about 600 mg of the $\alpha 4\beta 7$ binding antibody at week 26.

(233) In some embodiments, upon administration of the one or more induction dose an average serum or plasma concentration at least about 35 $\mu\text{g/mL}$ of the $\alpha 4\beta 7$ binding antibody is achieved in the patient during an induction period. In some embodiments, upon administration of the one or more maintenance dose a serum or plasma concentration at least about 6 $\mu\text{g/mL}$ of the $\alpha 4\beta 7$ binding antibody is achieved in the patient in the patient for at least 2 weeks or more.

(234) While the present disclosure provides for subcutaneous administration of the $\alpha 4\beta 7$ binding antibody, it is also possible to administer the $\alpha 4\beta 7$ binding antibody by the intravenous route, i.e., by infusion. It is further contemplated that induction or loading dose or doses may be administered by the subcutaneous (SC) or intravenous (IV) route of administration.

(235) Specific Embodiments

(236) Non-limiting specific embodiments are described below, each of which is considered to be within the present disclosure. Embodiment 1. A multidose regimen comprising liquid formulations for use in treating a disease in a subject in need thereof comprising: (a) a first injectable liquid formulation comprising a total dosage amount of at least 500 mg of a long acting $\alpha 4\beta 7$ targeting molecule; and (b) a second injectable liquid formulation comprising a total dosage amount of at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule, wherein the long acting $\alpha 4\beta 7$ targeting molecule comprises: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

Embodiment 2. The multidose regimen of Embodiment 1, wherein the first injectable liquid formulation comprises a total dosage amount of from about 500 mg to about 1200 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 3. The multidose regimen of Embodiment 1, wherein the first injectable liquid formulation comprises a total dosage amount of about 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 4. The multidose regimen of Embodiment 1, wherein the first injectable liquid formulation comprises a total dosage amount of about 1000 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 5. The multidose regimen of anyone of Embodiments 1-4, wherein the second injectable liquid formulation comprises a total dosage amount of from about 300 mg to about 450 mg of the long acting $\alpha 4\beta 7$ targeting molecule.

Embodiment 6. The multidose regimen of anyone of Embodiments 1-4, wherein the second injectable liquid formulation comprises a total dosage amount of about 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 7. The multidose regimen of anyone of Embodiments 1-6, wherein the first injectable liquid formulation is for intravenous administration. Embodiment 8.

The multidose regimen of anyone of Embodiments 1-6, wherein the first injectable liquid formulation is for subcutaneous administration. Embodiment 9. The multidose regimen of Embodiment 8, wherein the first injectable liquid formulation is a single dose or a multidose formulation. Embodiment 10. The multidose regimen of anyone of Embodiments 1-9, wherein the second injectable liquid formulation is for subcutaneous administration. Embodiment 11. The multidose regimen of Embodiment 10, wherein the second injectable liquid formulation is suitable for administration as a single injection or multiple injections. Embodiment 12. The multidose regimen of any one of Embodiments 1-11, wherein the disease is a gastrointestinal inflammatory disease. Embodiment 13. The multidose regimen of any one of Embodiments 1-11, wherein the disease is an inflammatory bowel disease. Embodiment 14. The multidose regimen of Embodiment 13, wherein the inflammatory bowel disease is Crohn's disease. Embodiment 15. The multidose regimen of Embodiment 13, wherein the inflammatory bowel disease is ulcerative colitis.

Embodiment 16. A dosing regimen for use in treating a disease in a subject in need thereof, the dosing regimen comprising: (a) a first formulation comprising a total dosage amount of at least

about 600 mg of a long acting $\alpha 4\beta 7$ targeting molecule for administration to the subject; (b) a second formulation comprising a total dosage amount of at least about 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule for subcutaneous administration to the subject at least two weeks after the first formulation, and thereafter as a maintenance dose at least eight weeks after administration of the second formulation, wherein the long acting $\alpha 4\beta 7$ targeting molecule comprises: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. Embodiment 17. The dosing regimen of Embodiment 16, wherein the first formulation is for subcutaneous administration. Embodiment 18. The dosing regimen of Embodiment 16, wherein the first formulation is for intravenous administration. Embodiment 19. The dosing regimen of any one of Embodiments 16-18, wherein the first formulation comprises a total dosage amount of from about 600 mg to about 1200 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 20. The dosing regimen of any one of Embodiments 16-18, wherein the first formulation comprises a total dosage amount of about 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 21. The dosing regimen of any one of Embodiments 16-20, wherein the second formulation comprises a total dosage amount of from about 300 mg to about 450 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 22. The dosing regimen of any one of Embodiments 16-20, wherein the second formulation comprises a total dosage amount of about 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 23. The dosing regimen of Embodiment 19 or Embodiment 20, wherein the first formulation is suitable for administration as a single injection or multiple injections. Embodiment 24. The dosing regimen of any one of Embodiments 16-23, wherein the second formulation is a single dose or multidose injectable formulation. Embodiment 25. The dosing regimen of any one of Embodiments 16-24, wherein the first formulation and the second formulation do not contain citrate. Embodiment 26. The dosing regimen of Embodiment 25, wherein the first formulation or second formulation or both formulations comprise one or more of: (a) histidine; (b) arginine or a salt thereof; (c) ethylenediaminetetraacetic acid (EDTA); and (d) polysorbate 80. Embodiment 27. The dosing regimen of any one of Embodiments 16-26, wherein the disease is a gastrointestinal inflammatory disease. Embodiment 28. The dosing regimen of any one of Embodiments 16-26, wherein the disease is an inflammatory bowel disease. Embodiment 29. The dosing regimen of Embodiment 28, wherein the inflammatory bowel disease is Crohn's disease. Embodiment 30. The dosing regimen of Embodiment 28, wherein the inflammatory bowel disease is ulcerative colitis. Embodiment 31. An injectable dosage form of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and a) a light chain consisting of an amino acid sequence according to SEQ ID NO, wherein the $\alpha 4\beta 7$ binding antibody has an average serum half-life greater than 6 days in cynomolgus monkeys. Embodiment 32. The injectable dosage form of Embodiment 31, wherein the average serum half-life is about 17 days or greater in cynomolgus monkeys. Embodiment 33. The injectable dosage form of Embodiments 31 or 32 comprising at least 100 mg/ml of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 34. The injectable dosage form of Embodiments 31 or 32 comprising at least 180 mg/ml of the $\alpha 4\beta 7$ binding antibody. Embodiment 35. The injectable dosage form of Embodiments 31 or 32 comprising at least 150 mg/ml of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 36. The injectable dosage form of Embodiment 31 or Embodiment 32 comprising at least 200 mg/ml of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 37. The injectable dosage form of any one of Embodiments 31-32, comprising from about 600 mg to about 1200 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 38. The injectable dosage form of any one of Embodiments 31-32, comprising about 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 39. The injectable dosage form of any one of Embodiments 31-32, comprising about 1000 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 40. The injectable dosage form of any one of Embodiments 31-39, wherein the injectable dosage form is an injectable liquid formulation. Embodiment 41. The injectable dosage form of Embodiment 40, wherein the injectable liquid formulation does not comprise

citrate. Embodiment 42. The injectable dosage form of Embodiment 41, wherein the injectable liquid formulation comprises one or more of: (a) histidine; (b) arginine or a salt thereof; (c) ethylenediaminetetraacetic acid (EDTA); and (d) polysorbate 80. Embodiment 43. The injectable liquid formulation of any one of Embodiments 1-30 or the Embodiments 40-42, wherein the injectable liquid formulation is a pharmaceutical composition. Embodiment 44. A method of treating a disease in a patient in need thereof, the method comprising: administering to a subject in need thereof (a) an effective amount of an induction dose of a long acting $\alpha 4\beta 7$ targeting molecule, and (b) an effective amount of one or more maintenance doses of the long acting $\alpha 4\beta 7$ targeting molecule, wherein the one or more maintenance doses are administered subcutaneously at least eight weeks apart, wherein the long acting $\alpha 4\beta 7$ targeting molecule comprises: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. Embodiment 45. The method of Embodiment 44, wherein the disease is a gastrointestinal inflammatory disease. Embodiment 46. The method of Embodiment 44, wherein the disease is an inflammatory bowel disease. Embodiment 47. The method of Embodiment 45, wherein the inflammatory bowel disease is Crohn's disease. Embodiment 48. The method of Embodiment 45, wherein the inflammatory bowel disease is ulcerative colitis. Embodiment 49. The method of any one of Embodiments 44-48, wherein the effective amount of each of the one or more maintenance doses of the long acting $\alpha 4\beta 7$ targeting molecule is at least about 300 mg. Embodiment 50. The method of any one of Embodiments 44-48, wherein the effective amount of each of the one or more maintenance doses of the long acting $\alpha 4\beta 7$ targeting molecule is from about 300 mg to about 450 mg. Embodiment 51. The method of any one of Embodiments 44-48, wherein the effective amount of the induction dose of the long acting $\alpha 4\beta 7$ targeting molecule is about 600 mg. Embodiment 52. The method of any one of Embodiments 44-48, wherein the effective amount of the induction dose of the long acting $\alpha 4\beta 7$ targeting molecule is about 1000 mg. Embodiment 53. The method of any one of Embodiments 44-48, wherein the effective amount of the induction dose of the long acting $\alpha 4\beta 7$ targeting molecule is from 600 mg to about 1200 mg. Embodiment 54. The method of any one of Embodiments 51-53 comprising administering the effective amount of the induction dose in a single or multiple injections. Embodiment 55. The method of any one of Embodiments 44-54, wherein the effective amount of the induction dose of the long acting $\alpha 4\beta 7$ targeting molecule is administered subcutaneously. Embodiment 56. The method of any one of Embodiments 44-54, wherein the effective amount of the induction dose of the long acting $\alpha 4\beta 7$ targeting molecule is administered intravenously. Embodiment 57. The method of any one of Embodiments 44-56, wherein the one or more maintenance doses are administered subcutaneously about 12 weeks apart. Embodiment 58. The method of any one of Embodiments 44-56, wherein the one or more maintenance doses are administered subcutaneously about 26 weeks apart. Embodiment 59. The method of any one of Embodiments 44-58, wherein the induction dose comprises a dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule that is about 2 or more times higher than a dosage amount of each of the one or more maintenance doses. Embodiment 60. The method of any one of Embodiments 44-59, wherein: (a) the long acting $\alpha 4\beta 7$ targeting molecule follows a biphasic decline in serum concentration; (b) repeated dosing of the long acting $\alpha 4\beta 7$ targeting molecule does not have a significant influence on clearance (CL) and steady-state volume of distribution (Vss); (c) pharmacokinetic is not impacted by disease state of the patient; or (d) a combination thereof. Embodiment 61. A method of achieving C.sub.avgW0-W12 of from 60 μ g/ml to 70 μ g/ml or greater for a long acting $\alpha 4\beta 7$ targeting molecule in a subject in need thereof, the method comprising administering to the subject in need thereof a first dosage amount of at least 600 mg of the $\alpha 4\beta 7$ binding antibody, wherein the long acting $\alpha 4\beta 7$ targeting molecule comprises: (a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and (b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. Embodiment 62. The method of Embodiment 61, wherein the administering results in a C.sub.avgW0-W12 of about 65 μ g/ml.

Embodiment 63. The method of Embodiment 62, wherein the administering comprises administering to the subject in need thereof the first dosage amount of at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule and a second dosage amount of at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule at least two weeks after administration of the first dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 64. The method of Embodiment 63, comprising administering the first dosage amount of at least 600 mg intravenously and the second dosage amount of at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule subcutaneously. Embodiment 65. The method of Embodiment 63, comprising administering the first dosage amount of at least 600 mg subcutaneously and the second dosage amount of at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule subcutaneously. Embodiment 66. The method of any one of Embodiments 61-65 further comprising administering to the subject in need thereof one or more dosage amount of at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule at least four weeks after administration of the second dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 67. The method of Embodiment 66 comprising administering the one or more dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule subcutaneously. Embodiment 68. The method of any one of Embodiments 61-67 comprising administering to the subject in need thereof at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule every twenty-six weeks after administration of the second dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 69. A method of achieving C.sub.avgW0-W6 of from 65 $\mu\text{g/ml}$ to 75 $\mu\text{g/ml}$ or greater for a long acting $\alpha 4\beta 7$ targeting molecule in a subject in need thereof, the method comprising administering to the subject in need thereof a first dosage amount of at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule, wherein the long acting $\alpha 4\beta 7$ targeting molecule comprises: (a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and (b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. Embodiment 70. The method of Embodiment 69, wherein the administering results in a C.sub.avgW0-W6 of about 70 $\mu\text{g/ml}$. Embodiment 71. The method of Embodiment 69, wherein the administering results in a C.sub.avgW0-W6 of about 65 $\mu\text{g/ml}$. Embodiment 72. The method of any one of Embodiments 69-71, wherein the administering comprises administering to the subject in need thereof a first dosage amount of at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule and a second dosage amount of at least 300 mg of long acting $\alpha 4\beta 7$ targeting molecule at least two weeks after administration of the first dosage amount of at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 73. The method of Embodiment 72, comprising administering the first dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule intravenously and the second dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule subcutaneously. Embodiment 74. The method of Embodiment 72, comprising administering the first dosage amount and the second dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule subcutaneously. Embodiment 75. The method of any one of Embodiments 72-74 further comprising administering to the subject in need thereof one or more additional dosage amount of at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule at least four weeks after administration of the second dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 76. The method of Embodiment 75 comprising administering the one or more additional dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule subcutaneously. Embodiment 77. The method of Embodiment 75 or Embodiment 68 further comprising administering to the subject in need thereof at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule every twenty-six weeks after administration of second dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule. Embodiment 78. The method of any one of Embodiments 66-68 or 75-77, wherein the administration of the one or more dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule results in C.sub.avg of from 40 $\mu\text{g/ml}$ to 50 $\mu\text{g/ml}$ or greater. Embodiment 79. The method of any one of Embodiments 66-68 or 75-77, wherein the administration of the one or more dose of the long acting $\alpha 4\beta 7$ targeting molecule results in C.sub.avg of about 45 $\mu\text{g/ml}$. Embodiment 80. A method of achieving C.sub.trough of 35 $\mu\text{g/ml}$ or greater for a long acting $\alpha 4\beta 7$ targeting molecule in a

subject in need thereof six weeks following administration of the long acting $\alpha 4\beta 7$ targeting molecule to the patient, the method comprising administering to the patient in need thereof at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule, wherein the long acting $\alpha 4\beta 7$ targeting molecule comprises: (a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and (b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

Embodiment 81. The method of Embodiment 80, wherein the administration of the at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule results in a serum C.sub.avgW0-W12 of at least 60 $\mu\text{g/ml}$.

Embodiment 82. The method of Embodiment 80, wherein the administration of the at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule results in a serum C.sub.avgW30-W40 of at least 30 $\mu\text{g/ml}$.

Embodiment 83. The method of Embodiment 80, wherein the at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule is administered intravenously.

Embodiment 84. The method of Embodiment 80, wherein the at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule is administered subcutaneously.

Embodiment 85. The method of any one of Embodiments 80-84 further comprising administering to the subject in need thereof at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule, wherein the administration of the at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule results in a steady state C.sub.trough of at least 6 $\mu\text{g/ml}$.

Embodiment 86. The method of any one of Embodiments 80-84, further comprising administering to the subject in need thereof at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule results in an average serum concentration of the $\alpha 4\beta 7$ binding antibody above 35 $\mu\text{g/ml}$ for ten weeks following the administration of the at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule.

Embodiment 87. The method of any one of Embodiments 85-86 further comprising administering subcutaneously to the subject in need thereof at least 300 mg of long acting $\alpha 4\beta 7$ targeting molecule four weeks after administration of last dosage amount of the long acting $\alpha 4\beta 7$ targeting molecule.

Embodiment 88. The method of any one of Embodiments 85-86 further comprising administering subcutaneously to the patient a maintenance dose of at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule every twenty-six weeks after administration of the last dose of the $\alpha 4\beta 7$ binding antibody.

Embodiment 89. A long acting $\alpha 4\beta 7$ targeting molecule comprising: (a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

Embodiment 90. An isolated nucleic acid encoding the heavy chain and/or light chain of the long acting $\alpha 4\beta 7$ targeting molecule of Embodiment 89.

Embodiment 91. The isolated nucleic acid of Embodiment 90 encoding SEQ ID NO: 1.

Embodiment 92. The isolated nucleic acid of Embodiment 90 encoding SEQ ID NO: 3.

Embodiment 93. A recombinant host cell comprising the isolated nucleic acid of any one of Embodiments 90-92, or an expression vector comprising the isolated nucleic acid of any one of Embodiments 90-92.

Embodiment 95. A kit for treating a disease in a subject in need thereof comprising: (a) an injectable liquid formulation comprising a long acting $\alpha 4\beta 7$ targeting molecule, wherein the long acting $\alpha 4\beta 7$ targeting molecule comprises: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3, and (b) instructions to administer at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule, at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule at least two weeks after the administration of the 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule, and at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule at least four weeks thereafter.

Embodiment 96. A kit for treating a disease in a subject in need thereof comprising: (a) an injectable subcutaneous liquid formulation comprising a long acting $\alpha 4\beta 7$ targeting molecule, wherein the long acting $\alpha 4\beta 7$ targeting molecule comprises: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3, and (b) instructions to administer subcutaneously at least 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule, at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule at least two weeks after the administration of the 600 mg of the long acting $\alpha 4\beta 7$ targeting molecule, and at least 300 mg of the long acting $\alpha 4\beta 7$ targeting molecule at least four weeks

thereafter. Embodiment 97. The kit of Embodiment 95 or Embodiment 96, wherein the disease is a gastrointestinal inflammatory disease. Embodiment 98. The kit of Embodiment 95 or Embodiment 96, wherein the disease is an inflammatory bowel disease, optionally Crohn's disease or ulcerative colitis. Embodiment 99. The long acting $\alpha 4\beta 7$ targeting molecule of Embodiment 89, the long acting $\alpha 4\beta 7$ targeting molecule of the regimen of any one of Embodiments 1-30, long acting $\alpha 4\beta 7$ targeting molecule of the injectable dosage form of any one of Embodiments 31-42, the long acting $\alpha 4\beta 7$ targeting molecule of the method of any one of Embodiments 44-88, the long acting $\alpha 4\beta 7$ targeting molecule of the kit of any one of Embodiments 95-98, wherein the long acting $\alpha 4\beta 7$ targeting molecule comprises one or more of the following characteristics: i. an average serum half-life greater than about 6 days in cynomolgus monkeys, ii. an average serum half-life of about 17 days or greater in cynomolgus monkeys, iii. a melting temperature T_{mOnset} greater than about 55°C ., iv. a melting temperature T_{mOnset} greater than 56.2°C ., v. does not induce T cell activation markers CD25 or CD69, vi. does not result in release of cytokines at least 6 hours after the administration, wherein the cytokines comprise one or more of IL-6, IL-8, IL-10, IFN γ , IL-4, IL-17, IL-2, IL-23p70 and TNF, vii. does not induce complement-dependent cytotoxicity (CDC) in primary human PBMCs and in $\alpha 4\beta 7$ -expressing human β -lymphoid cell, viii. does not induce antibody dependent cellular cytotoxicity (ADCC) in human NK cells, ix. does not impact suppressive activity of regulatory T cells expressing $\alpha 4\beta 7$ integrin, wherein the suppressive activity is measured by presence of elevated CD71, CD25, Ki67, granzyme B, or OX40, x. a proportion of GOF from about 55.10% to about 60.4%, xi. a proportion of G1F from about 16.6% to about 18.8%, xii. a proportion of Man5 from about 7.3% to about 8.8%, Embodiment 100. A subcutaneous dosage form comprising at least about 300 mg of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. Embodiment 101. The dosage form of Embodiment 100, wherein the heavy chain comprises an amino acid sequence according to SEQ ID NO: 1 and the light chain comprises an amino acid sequence according to SEQ ID NO: 3. Embodiment 102. The dosage form of Embodiment 100, wherein the heavy chain comprises an amino acid sequence according to SEQ ID NO: 2 and the light chain comprises an amino acid sequence according to SEQ ID NO: 3. Embodiment 103. The dosage form of any one of Embodiments 100-102, comprising about 350 mg, 400 mg, 450 mg, 500 mg, 550 mg, 600 mg, 650 mg, or 700 mg of the $\alpha 4\beta 7$ binding antibody. Embodiment 104. A method of administering one or more of the dosage forms of any one of Embodiments 100-103. Embodiment 105. A method of treating an inflammatory bowel disease in a patient in need thereof, the method comprising subcutaneously or intravenously administering to the patient an effective amount of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. Embodiment 106. The method of Embodiment 105, wherein the inflammatory bowel disease is Crohn's disease or ulcerative colitis. Embodiment 107. The method of Embodiment 105, wherein the inflammatory bowel disease is ulcerative colitis. Embodiment 108. The method of any one of Embodiments 105-107, wherein the heavy chain comprises an amino acid sequence according to SEQ ID NO: 1 and the light chain comprises an amino acid sequence according to SEQ ID NO: 3. Embodiment 109. The method of any one of Embodiments 105-108, wherein the heavy chain comprises an amino acid sequence according to SEQ ID NO: 2 and the light chain comprises an amino acid sequence according to SEQ ID NO: 3. Embodiment 110. The method of any one of claims Embodiments 105-109, wherein administration of the $\alpha 4\beta 7$ binding antibody is subcutaneous. Embodiment 111. The method of any one of Embodiments 105-110, wherein administration of the $\alpha 4\beta 7$ binding antibody is intravenous. Embodiment 112. The method of any one of Embodiments 105-111, wherein the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg. Embodiment 113. The method of any one of Embodiments 105-112, wherein the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 350

mg, 400 mg, 450 mg, 500 mg, 550 mg, 600 mg, 650 mg, or 700 mg. Embodiment 114. A method of treating a disease or disorder in a patient in need thereof, the method comprising administering to the patient at least once at an interval more than 8 weeks, an effective amount of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. Embodiment 115. The method of Embodiment 114, wherein the interval is about 12 to about 26 weeks. Embodiment 116. The method of Embodiment 115, wherein the interval is about 12 weeks. Embodiment 117. The method of Embodiment 115, wherein the interval is about 16 weeks. Embodiment 118. The method of Embodiment 115, wherein the interval is about 26 weeks. Embodiment 119. The method of Embodiment 115, wherein the disease or disorder is an inflammatory bowel disease. Embodiment 120. The method of Embodiment 119, wherein the inflammatory bowel disease is Crohn's disease or ulcerative colitis. Embodiment 121. The method of Embodiment 119, wherein the inflammatory bowel disease is ulcerative colitis. Embodiment 122. The method of any one of Embodiments 115-121, wherein the heavy chain comprises an amino acid sequence according to SEQ ID NO: 1 and the light chain comprises an amino acid sequence according to SEQ ID NO: 3. Embodiment 123. The method of any one of Embodiments 115-121, wherein the heavy chain comprises an amino acid sequence according to SEQ ID NO: 2 and the light chain comprises an amino acid sequence according to SEQ ID NO: 3. Embodiment 124. The method of any one of Embodiments 115-123, wherein administration of the $\alpha 4\beta 7$ binding antibody is subcutaneous. Embodiment 125. The method of any one of Embodiments 115-123, wherein administration of the $\alpha 4\beta 7$ binding antibody is intravenous. Embodiment 126. The method of any one of Embodiments 115-125, wherein the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 300 mg. Embodiment 127. The method of any one of Embodiments 115-125, wherein the $\alpha 4\beta 7$ binding antibody is administered at a dose of about 350 mg, 400 mg, 450 mg, 500 mg, 550 mg, 600 mg, 650 mg, or 700 mg. Embodiment 128. The method of any one of Embodiments 115-127, wherein the disease or disorder is an inflammatory bowel disease and administration of the $\alpha 4\beta 7$ binding antibody is at a dose of about 300 mg and is subcutaneous. Embodiment 129. A method of treating a disease or disorder in a patient in need thereof comprising subcutaneously administering to the patient about 108 mg or about 300 mg of an $\alpha 4\beta 7$ binding antibody every four or six weeks, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence according to SEQ ID NO: 4 and a light chain comprising an amino acid sequence according to SEQ ID NO: 5; and wherein no intervening subcutaneous administration of the $\alpha 4\beta 7$ binding antibody occurs during the four or six weeks. Embodiment 130. A method of treating a disease or disorder in a patient in need thereof comprising subcutaneously administering to the patient about 108 mg or about 300 mg of an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2 and a light chain comprising an amino acid sequence according to SEQ ID NO: 3; and wherein a plasma concentration of the $\alpha 4\beta 7$ binding antibody of at least about 35 $\mu\text{g/mL}$ is achieved in the patient for at least 2 weeks or more upon the administration. Embodiment 131. A method of treating a disease or disorder in a patient in need thereof comprising subcutaneously administering to the patient about 108 mg or about 300 mg of an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising an amino acid sequence according to SEQ ID NO: 1 or SEQ ID NO: 2 and a light chain comprising an amino acid sequence according to SEQ ID NO: 3; and wherein a C.sub.trough of the $\alpha 4\beta 7$ binding antibody of at least about 6 $\mu\text{g/mL}$ is achieved in the patient for at least 2 weeks or more upon the administration. Embodiment 132. The method of any one of Embodiments 130-131, wherein the disorder is inflammatory bowel disease. Embodiment 133. The method of any one of Embodiments 130-131, wherein the disorder is Crohn's disease or ulcerative colitis. Embodiment 134. An isolated nucleic acid, wherein the nucleic acid encodes the heavy chain and/or light chain of the $\alpha 4\beta 7$ binding antibody of any one of Embodiments 100-134. Embodiment 135. The isolated nucleic acid of Embodiment 134, wherein

the nucleic acid sequence encodes SEQ ID NO: 1. Embodiment 136. The isolated nucleic acid of Embodiment 134, wherein the nucleic acid encodes SEQ ID NO: 3. Embodiment 137. A recombinant host cell comprising the nucleic acid sequences of any one of Embodiments 135-136 or an expression vector comprising the nucleic acid sequences of any one of Embodiments 135-136. Embodiment 138. An $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. Embodiment 139. An $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. Embodiment 140. An $\alpha 4\beta 7$ binding antibody comprising a heavy chain and a light chain, wherein the heavy chain consists of an amino acid sequence according to SEQ ID NO: 1. Embodiment 141. The $\alpha 4\beta 7$ binding antibody of Embodiment 140, wherein the light chain consists of an amino acid sequence according to SEQ ID NO: 3. Embodiment 142. The $\alpha 4\beta 7$ binding antibody of any one of Embodiments 138-141, wherein the $\alpha 4\beta 7$ binding antibody has an increased half-life as compared to an $\alpha 4\beta 7$ binding antibody comprising a heavy chain sequence consisting of SEQ ID NO: 4. Embodiment 143. The $\alpha 4\beta 7$ binding antibody of any one of Embodiments 138-142, wherein the $\alpha 4\beta 7$ binding antibody has a half-life of from 42 days to 56 days or more in humans. Embodiment 144. A pharmaceutical composition comprising an $\alpha 4\beta 7$ binding antibody and a pharmaceutically acceptable carrier, the $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. Embodiment 145. A pharmaceutical composition comprising an $\alpha 4\beta 7$ binding antibody and a pharmaceutically acceptable carrier, the $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. Embodiment 146. The pharmaceutical composition of any one of Embodiment 144-145, wherein the composition is citrate free. Embodiment 147. A subcutaneous dosage form comprising at least about 150 mg of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and b) a light chain comprising an amino acid sequence according to SEQ ID NO: 3. Embodiment 148. A subcutaneous dosage form comprising at least about 150 mg of an $\alpha 4\beta 7$ binding antibody comprising: a) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and b) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3. Embodiment 149. The subcutaneous dosage form of Embodiment 147 or Embodiment 148, wherein the dosage form is citrate free. Embodiment 150. The subcutaneous dosage form of Embodiment 147, Embodiment 148, or Embodiment 149, comprising from about 150 mg to about 900 mg of the $\alpha 4\beta 7$ binding antibody. Embodiment 151. A method for treating inflammatory bowel disease in a subject in need thereof, the method comprising administering subcutaneously to the subject one or more of the dosage forms of any one of Embodiments 147-150. Embodiment 152. The method of Embodiment 151, wherein the inflammatory bowel disease is Crohn's disease or ulcerative colitis. Embodiment 153. The method of Embodiment 151, wherein the inflammatory bowel disease is ulcerative colitis. Embodiment 154. A method of treating an inflammatory bowel disease in a subject in need thereof, the method comprising administering subcutaneously to the subject an effective amount of an $\alpha 4\beta 7$ binding antibody comprising a heavy chain and a light chain, wherein the heavy chain consists of an amino acid sequence according to SEQ ID NO: 1; and the light chain comprises the variable region of SEQ ID NO: 3. Embodiment 155. A method for treating inflammatory bowel disease in a subject in need thereof, the method comprising administering subcutaneously to the subject an effective amount of an $\alpha 4\beta 7$ binding antibody comprising a heavy chain and a light chain, wherein the heavy chain consists of an amino acid sequence according to SEQ ID NO: 1; and the light chain consists of the variable region of SEQ ID NO: 3. Embodiment 156. The method of Embodiment 154 or Embodiment 155, wherein the $\alpha 4\beta 7$ binding antibody is administered at a dose of from about 150 mg to about 900 mg. Embodiment

157. The method of Embodiment 154 or Embodiment 155, comprising administering to the subject the effective amount of an $\alpha 4\beta 7$ binding antibody at an interval of more than 8 weeks. Embodiment 158. The method of Embodiment 157, wherein the interval is about 12 to about 26 weeks. Embodiment 159. The method of Embodiment 157, wherein the interval is about 12 weeks. Embodiment 160. The method of Embodiment 157, wherein the interval is about 16 weeks. Embodiment 161. The method of Embodiment 157, wherein the interval is about 26 weeks. Embodiment 162. The method of Embodiment 157, wherein the inflammatory bowel disease is Crohn's disease or ulcerative colitis. Embodiment 163. The method of Embodiment 157, wherein the inflammatory bowel disease is ulcerative colitis. Embodiment 164. The method of Embodiment 155 or Embodiment 156, wherein an average plasma or serum concentration of the $\alpha 4\beta 7$ binding antibody of at least about 35 $\mu\text{g/mL}$ is achieved in the subject during an induction period. Embodiment 165. The method of Embodiment 154 or Embodiment 155, wherein a maintenance C.sub.trough of the $\alpha 4\beta 7$ binding antibody of at least about 6 $\mu\text{g/mL}$ is achieved in the subject upon the administration. Embodiment 166. A method for treating inflammatory bowel disease in a subject in need thereof, the method comprising (a) administering subcutaneously to the subject one or more induction dose comprising an effective amount of an $\alpha 4\beta 7$ binding antibody, wherein the $\alpha 4\beta 7$ binding antibody comprises a heavy chain comprising a heavy chain and a light chain, wherein the heavy chain consists of an amino acid sequence according to SEQ ID NO: 1; and the light chain consists of the variable region of SEQ ID NO: 3; (b) administering subcutaneously to the subject one or more maintenance dose comprising an effective amount of the $\alpha 4\beta 7$ binding antibody. Embodiment 167. The method of Embodiment 166, wherein the induction dose comprises an amount to the $\alpha 4\beta 7$ binding antibody about 2 or more times higher than the maintenance dose. Embodiment 168. The method of Embodiment 166, wherein the induction dose comprises from about 400 mg to about 600 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of from about 150 to about 300 mg of the $\alpha 4\beta 7$ binding antibody. Embodiment 169. The method of Embodiment 166, wherein the induction dose comprises about 600 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of about 300 mg of the $\alpha 4\beta 7$ binding antibody. Embodiment 170. The method of Embodiment 166, wherein the induction dose comprises about 400 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of about 200 mg of the $\alpha 4\beta 7$ binding antibody. Embodiment 171. The method of Embodiment 166, wherein the induction dose comprises about 600 mg of the $\alpha 4\beta 7$ binding antibody and wherein the maintenance dose of about 600 mg of the $\alpha 4\beta 7$ binding antibody. Embodiment 172. The method of any one of Embodiments 166-171, wherein upon administration of the one or more induction dose an average serum or plasma concentration at least about 35 $\mu\text{g/mL}$ of the $\alpha 4\beta 7$ binding antibody is achieved in the patient during an induction period. Embodiment 173. The method of any one of Embodiments 166-171, wherein upon administration of the one or more maintenance dose a serum or plasma concentration at least about 6 $\mu\text{g/mL}$ of the $\alpha 4\beta 7$ binding antibody is achieved in the patient in the patient for at least 2 weeks or more. Embodiment 174. The method of any one of Embodiments 166-171, wherein the one or more maintenance dose of the $\alpha 4\beta 7$ binding antibody is administered at an interval of more than 8 weeks.

EXAMPLES

(237) The disclosure now being generally described, will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present disclosure, and is not intended to limit the disclosure.

Example 1: $\alpha 4\beta 7$ Binding Antibodies Efficacy

(238) This Example describes the efficacy of the $\alpha 4\beta 7$ binding antibodies described herein. PBMCs contain cell types that express $\alpha 4\beta 7$ integrin and CD4⁺ helper T cells are associated with inflammatory bowel diseases. The ability of $\alpha 4\beta 7$ binding antibody described herein (i.e., $\alpha 4\beta 7$ binding antibody Ab001 comprising a heavy chain comprising a sequence according to SEQ ID NO: 1 and a light chain comprising a sequence according to SEQ ID NO: 3) to bind PBMCs was

analyzed. Cell binding to PBMCs isolated from three human donors was determined by staining with either comparator antibody (solid line; filled shapes) or Ab001 (dashed line; open shapes) at various increasing concentrations (FIG. 1A and FIG. 1B). FIG. 1C shows the receptor occupancy on $\alpha 4\beta 7$ -expressing peripheral blood mononuclear cells. The comparator antibody described in this Example and the remaining Examples comprises a heavy chain comprising a sequence according to SEQ ID NO: 4 and a light chain comprising a sequence according to SEQ ID NO: 5. A basic immunophenotyping T-cell panel indicated that both the comparator antibody and $\alpha 4\beta 7$ binding antibody exhibit binding on the memory T-helper CD3+CD4+CD45RO+ population. EC.sub.50 values for the comparator antibody and $\alpha 4\beta 7$ binding antibody is seen in (Table 2A) and EC.sub.50 values for both antibodies correspond well with what has previously been reported for binding to the same population: EC.sub.50 0.3 nM.

(239) TABLE-US-00005 TABLE 2A EC50 data for binding to PBMCs Comparator $\alpha 4\beta 7$ binding antibody Antibody (Ab001) EC.sub.50 (nM) EC.sub.50 (nM) Donor 1 0.22 0.21 Donor 2 0.27 0.30 Donor 3 0.24 0.22

(240) The IC.sub.50 values for competitive inhibition were determined by titrating unlabeled Ab001 into samples equilibrated with EC50 concentrations of fluorescently labeled Ab001. Ab001 and Comparator Antibody Receptor Occupancy (IC.sub.50) on CD4+, CD20+, and CD45RO+ Cells from Human and Monkey Donors are shown in Table 2B.

(241) TABLE-US-00006 TABLE 2B Receptor Occupancy by Cell Type Human whole blood Human PBMCs Monkey PBMCs Comparator Comparator Comparator Cell Type Attribute Ab001 Antibody Ab001 Antibody Ab001 Antibody CD4 + IC 50 (nM, 0.41-0.64 0.40-0.46 0.37-0.39 0.35-0.40 0.49-0.54 0.48-0.60 T.sub.h Cells range, n = 3 donors) Maximum 85-97% 86-97% 68-84% 88-91% 86-88% 88-93% inhibition (% , range, n = 3 donors) CD20 + IC.sub.50 (nM, 0.44-0.60 0.40-0.49 0.35-0.38 0.36-0.38 0.46-0.50 0.42-0.66 B Cells range, n = 3 donors) Maximum 85-98% 83-98% 68-83% 90-91% 76-83% 80-85% inhibition (% , range, n = 3 donors) CD45RO + IC.sub.50 (nM, 0.41-0.64 0.40-0.47 0.37-0.40 0.31-0.44 0.53 0.58 Memory T range, n = 1 Cells donor) Maximum 86-97% 87-97% 67-84% 84-89% 89% 93% inhibition (% , range, n = 1 donor) All cell IC.sub.50 (nM, 0.53 $\pm$ 0.08 0.44 $\pm$ 0.03 0.38 $\pm$ 0.02 0.38 $\pm$ 0.03 0.50 $\pm$ 0.03 0.56 $\pm$ 0.08 types mean, n = 7) All cell Maximum 90 $\pm$ 5% 90 $\pm$ 5% 81 $\pm$ 3% 90 $\pm$ 1% 84% $\pm$ 4% 87% $\pm$ 5% types inhibition (% , mean, n = 7)

(242) $\alpha 4\beta 7$ integrin is a ligand for mucosal addressin cell adhesion molecule 1 (MAdCAM1) and the interaction of these two molecules leads to gut homing of lymphocytes. Blocking this interaction is a therapeutic target for inflammatory bowel diseases including ulcerative colitis and Crohn's disease. $\alpha 4\beta 7$ -mediated adhesion of MAdCAM-1 was tracked in cells treated with various increasing concentrations of $\alpha 4\beta 7$ binding antibody Ab001 or comparator antibody to determine percent inhibition of total adhesion of MAdCAM-1 (FIG. 2A). $\alpha 4\beta 7$ binding antibody Ab001 preformed as effectively as the comparator antibody. $\alpha 4\beta 1$ integrin is a related integrin that is a ligand for VCAM-1 that mediates homing of lymphocytes to vascular endothelium. Importantly, in a similar assay tracking VCAM-1 adhesion the $\alpha 4\beta 7$ binding antibody Ab001 did not block VCAM-1 adhesion. As a positive control an antibody that can target $\alpha 4\beta 1$ integrin did inhibit VCAM-1 adhesion (FIG. 2B). $\alpha 4\beta 7$ binding antibody Ab001 was evaluated in human whole blood assays designed to detect effects on cytokine release and T cell activation. In whole blood assays from three human donors, $\alpha 4\beta 7$ binding antibody Ab001 did not induce T cell activation markers CD25 or CD69 as measured by FACS. Furthermore, no release of cytokines (Luminex assays of IL-6, IL-8, IL-1, IFN γ , IL-4, IL-17, IL-2, IL-23p70 and TNF) was observed at time points taken 6 hours or 24 hours after addition of $\alpha 4\beta 7$ binding antibody Ab001 (400 μ g/mL) to the samples. In contrast, the above markers were induced when human whole blood was incubated with lipopolysaccharide (LPS) or phytohemagglutinin (PHA), known mitogens that target T and B cells and activate cytokine release.

(243) The ability of $\alpha 4\beta 7$ binding antibody Ab001 to induce CDC was evaluated in primary human

PBMCs and in the $\alpha 4\beta 7$ -expressing human B-lymphoid cell line RPMI-8866. PBMCs were incubated (4 hr) with (i) $\alpha 4\beta 7$ binding antibody Ab001, (ii) comparator antibody, (iii) OKT3 (known to induce CDC in T cells), and (iv) an IgG1 isotype negative control antibody at a series of concentrations up to 10 $\mu\text{g/mL}$. RPMI-8866 cells were incubated (4 hr) with (i) $\alpha 4\beta 7$ binding antibody Ab001, (ii) comparator antibody, (iii) rituximab (known to induce CDC in B cells), and (iv) the same IgG1 isotype negative control antibody as in the PBMC cell assay. In both the PBMC and RPMI-8866 assays, cell viability was measured and complement-induced specific death (%) was reported versus test antibody concentration for all experimental conditions. Ab001, like the comparator antibody, did not induce CDC in either PBMCs or RPMI-8866 cells.

(244) $\alpha 4\beta 7$ binding antibody Ab001 was evaluated for its ability to induce ADCC in human NK cells. Primary human NK cells were isolated and used as effector cells along with an $\alpha 4\beta 7$ expressing human B-lymphoid cell line (RPMI-8866) used as the target cell population. Effector NK cells and RPMI-8866 cells together were incubated (3 hr) with (i) $\alpha 4\beta 7$ binding antibody Ab001, (ii) comparator antibody, (iii) rituximab (known to induce ADCC), and (iv) an IgG1 isotype negative control antibody at a series of concentrations from 300 $\mu\text{g/mL}$ to 10 g/mL . Cell viability was measured and ADCC-induced specific death (%) was reported versus test antibody concentration for all experimental conditions. Neither $\alpha 4\beta 7$ binding antibody Ab001 nor the comparator antibody induced NK cell-mediated ADCC in RPMI-8866 cells. $\alpha 4\beta 7$ binding antibody Ab001 showed similar cell death ranges as comparator antibody.

(245) The ability of $\alpha 4\beta 7$ binding antibody Ab001 to impact the normal activity of regulatory T cells (Treg) was tested using human PBMCs. Regulatory T cells expressing $\alpha 4\beta 7$ integrin were isolated by FACS and co-cultured with CD8⁺T responder cells in the presence or absence of test antibodies and a T cell stimulation reagent. Markers of T cell activation (CD71, CD25, Ki67, granzyme B, or OX40) were measured. $\alpha 4\beta 7$ binding antibody Ab001 did not impact Treg suppressive activity at concentrations more than 10-fold above its known EC50. As Treg cell numbers increased from a ratio of 0:16 (None) to 8:16 Treg: CD8⁺, stimulated CD8⁺ cells were suppressed based on one marker of T cell activation (CD25⁺). Similar results were observed when treated with $\alpha 4\beta 7$ binding antibody Ab001, media alone, the IgG1 isotype negative control, and the comparator antibody. Similar results were also seen for Treg suppressive activity when measured by the markers CD71, Ki67, granzyme B, and OX40.

(246) The ability of $\alpha 4\beta 7$ binding antibody Ab001 to bind and induce internalization of $\alpha 4\beta 7$ integrin was measured using CD4⁺ memory T Cells. Human memory CD4⁺ T cells were exposed to fluorescently labeled test antibodies, including $\alpha 4\beta 7$ binding antibody Ab001-A647 and the comparator antibody, at concentrations ranging from 0 nM to 100 nM, then incubated (24 hr) at temperatures that either allow (37° C.) or restrict (4° C.) receptor internalization. Cells were then washed with either buffer (FACS), or a gentle acid (0.5 M NaCl+0.2 M acetic acid), the latter to remove membrane-bound antibodies and associated fluorescence signal. Cells incubated with A647-labeled Ab001 under internalization conditions (37° C.) retained greater fluorescence after acid wash than cells stained and incubated under conditions when internalization was restricted (4° C.). This indicated that $\alpha 4\beta 7$ integrin was internalized after binding to fluorescently comparator antibody.

(247) The ability for functional $\alpha 4\beta 7$ to be rapidly expressed after antibody-induced internalization was determined by incubating $\alpha 4\beta 7$ binding antibody Ab001 or the comparator antibody (24 hr at 37° C.) at concentrations of 0 nM (control), 0.11 nM, 0.33 nM, or 1.0 nM with human PBMCs. After incubation, to allow internalization, cells were washed twice with PBS to remove unbound, unlabeled test antibodies. The cells were then resuspended in suitable media (RPMI 1640+10% fetal bovine serum), incubated for 96 hr with fluorescently-labeled $\alpha 4\beta 7$ binding antibody Ab001 or comparator antibody and monitored for fluorescence at 0 hr, 24 hr, and 96 hr. The results were similar for both $\alpha 4\beta 7$ binding antibody Ab001 and comparator antibody, demonstrating that functional $\alpha 4\beta 7$ could be re-expressed on lymphocytes within 24 to 96 hours of antibody-induced

internalization.

(248) Using non-human primates (NHPs) the half-life of Ab001 was studied. NHPs were intravenously injected with comparator antibody or Ab001 and normalized concentration of each protein was calculated at multiple timepoints out to 21 days post injection (FIG. 3). Data from an external study using the comparator antibody was also plotted ($t_{sub.1/2} \approx 10$ days). The β -elimination half-life of $\alpha 4\beta 7$ binding antibody Ab001 ($t_{sub.1/2} \approx 17$ days) was 70% longer than observed for the comparator antibody.YTE modification used for the $\alpha 4\beta 7$ binding antibody Ab001 described herein typically increases mAb half-life by 2-4 $\times$ versus wild-type in both NHP and human. Human mAb half-life is at least $\sim 2 \times$ NHP half-life. Therefore the human half-life based on initial allometric scaling based on these data is projected to be 43-56 days.

(249) In additional experiments, half-life extension was evaluated via pharmacokinetic analysis in Tg276 transgenic mice (hemizygous for human FcRn) following a single intravenous bolus dose of the $\alpha 4\beta 7$ binding antibody Ab001 (10 mg/kg, 5 mg/kg, 1 mg/kg) and in cynomolgus monkeys following a single bolus dose of the $\alpha 4\beta 7$ binding antibody Ab001 (150 mg/kg), given by either intravenous (IV) and subcutaneous (SC) administration. PK simulations in humans were based on allometric scaling of the $\alpha 4\beta 7$ binding antibody Ab001 clearance observed in cynomolgus monkey and conducted in MATLAB.

(250) In Tg276 mice, the half-life of the $\alpha 4\beta 7$ binding antibody Ab001 was 10-11 days across all dose cohorts compared to approximately 3 days for the comparator antibody. In cynomolgus monkeys, the half-life of the $\alpha 4\beta 7$ binding antibody Ab001 was 19-23 days following a single IV or SC infusion, a significant increase over the reported half-life of 10 days for the comparator antibody in cynomolgus monkeys. Based on allometric scaling of the clearance of the $\alpha 4\beta 7$ binding antibody Ab001 observed in this study, predictive simulations of the $\alpha 4\beta 7$ binding antibody Ab001 PK in humans suggested that Q8W-Q12W SC dosing at 300 mg would be able to achieve a $C_{sub.6-wks}$, $induction \geq 35 \mu g/mL$ and $C_{sub.trough}$, $ss \geq 6 \mu g/mL$ for simulated patients.

Example 2: Subcutaneous Dose Regimens for $\alpha 4\beta 7$ Binding Antibodies

(251) Pharmacokinetic (PK)-pharmacodynamic (PD) modeling and simulation was conducted to determine dose and regimens of subcutaneous doses that maintain certain serum concentrations above a specific level at a given time during the regimen or at steady state. Achieving these thresholds has been demonstrated to be an important metric for patients to achieve clinical remission in ulcerative colitis. Specifically, maintaining a trough concentration $C_{sub.trough} \geq 35 \mu g/mL$ after 6 weeks post-induction is associated with significantly higher rates of remission in ulcerative colitis.

(252) A two compartment linear model with parameters for central volume of distribution (V_c), peripheral volume of distribution (V_p), clearance from the central volume (Cl), first-order absorption rate constant from a subcutaneous depot into the central volume (k_a), and subcutaneous bioavailability relative to the intravenous administration was used to identify dosing regimens. Given the expectation of the $\alpha 4\beta 7$ binding antibodies disclosed herein to have a longer serum half-life than normal IgG, the effective human half-life used in these models was 53 days.

(253) In order to achieve higher clinical remission overall, dosing regimens where $>95\%$ of patients achieved a minimum concentration of $\geq 35 \mu g/mL$ at 6 weeks (Table 3) or $C_{sub.trough} \geq 6 \mu g/mL$ at steady state (Table 4) were identified. The model was further refined by adding the effect of covariates including body weight and serum albumin.

(254) To simulate the serum concentrations of the $\alpha 4\beta 7$ binding antibodies over time with specified dosing regimens in a large population of diverse patients, 1000 patients were uniquely assigned a body weight and serum albumin concentration. These two covariates were assigned independently and randomly chosen using a cumulative distribution function (CDF) of body weights and a separate CDF for serum albumin concentration. To derive each CDF, the percentiles for body weight and serum albumin concentration reported in a clinical trial of a comparator antibody were used to fit a generalized extreme value distribution to allow for a variety of probability distribution

behaviors that could be expected in a real-world clinical trial. Body weights values were log-transformed while serum albumin concentration values were used directly.

(255) TABLE-US-00007 TABLE 3 Minimum induction regimens allowing for >95% of patients to achieve a serum concentration of ≥ 35 $\mu\text{g/mL}$ at 6 weeks Minimum Minimum x Dose y Dose Strategy IV (Dose/Time) (mg) SC (Dose/Time) (mg) Strategy 1 x W0 550 None N/A Strategy 2 300 mg W0, x W2 200 None N/A Strategy 3 600 mg W0, x W2 0 None N/A Strategy 4 None N/A y W0 620 Strategy 5 None N/A 300 mg W0, y W2 280 Strategy 6 None NA 600 mg W0, y W2 100 Strategy 7 None N/A 300 mg W0, y W1, 150 W2, W3

(256) TABLE-US-00008 TABLE 4 Minimum maintenance regimens allowing for >90% of patients to achieve a serum C.sub.trough of ≥ 6 $\mu\text{g/mL}$ at steady-state Frequency Route Dose (mg) Every 4 weeks IV 40 SQ 45 Every 8 weeks IV 110 SQ 120 Every 12 weeks IV 220 SQ 275 Every 26 weeks IV 1800 SQ 2000

Example 3: Dose Responses of $\alpha 4\beta 7$ Binding Antibodies

(257) This Example describes the dose responses of the $\alpha 4\beta 7$ binding antibodies described herein.

(258) A simulated trial of 1000 patients was simulated using induction scenarios with an $\alpha 4\beta 7$ binding antibody with YTE modifications such as $\alpha 4\beta 7$ binding antibody Ab001 at intravenous doses of 300 mg, 350 mg, and 400 mg to determine the C.sub.trough values at 6 weeks post-induction (FIG. 4). Data generated in a clinical trial of a comparator antibody dosed intravenously at 300 mg is also shown. The comparator antibody was also run through the same simulation. A human half-life of 25.5 days was used for the comparator antibody and a human half-life of 50 days was used for Ab001 in these simulations. At most, 25% of patients dosed with the comparator antibody met the C.sub.trough ≥ 35 $\mu\text{g/mL}$ threshold at 6 weeks post-induction. In comparison, the vast majority of simulated patients dosed with all concentrations of Ab001 tested maintained a C.sub.trough ≥ 35 $\mu\text{g/mL}$.

(259) The best in class efficacy maintenance dosing of the comparator antibody is based on area under the curve at steady state (AUC.sub.ss) data of Q8W IV maintenance dosing at 300 mg (FIG. 5). When dosed subcutaneously the comparator antibody can only be dosed at 108 mg Q2W without dropping below the clinically relevant AUC. In contrast, Ab001 is predicted to maintain clinically relevant AUC with quarterly maintenance subcutaneous dosing. These data were generated using a human half-life of 35 and 50 days for Ab001 and plotted with the comparator antibody data (FIG. 5). Depending on human half-life, this could be achieved even as infrequently as Q16W.

(260) A post-hoc analysis of the comparator antibody trial data suggests a conservative target of C.sub.trough ≥ 6 $\mu\text{g/mL}$ during treatment maintains long-term efficacy. Best-in-class 300 mg IV Q8W dosing of the comparator antibody showed 75% of patients had C.sub.trough ≥ 6 $\mu\text{g/mL}$. The comparator antibody dosed subcutaneously at 108 mg with a half-life of 25.5 days, dosed IV at 300 mg with a half-life of 25.5 days, and Ab001 dosed subcutaneously at 300 mg with a half-life of 43, 53, or 56 days was used to simulate a trial of 1000 patients (FIG. 6). Similar to clinical results, 65% of simulated patients with Q8W IV achieve C.sub.trough ≥ 6 $\mu\text{g/mL}$ (65% is marked with a dashed line). Ab001 with a minimum 43-day half-life enables Q12W SC dosing. This matches the simulated comparator antibody IV Q8W dosing. Ab001 with a 50-day half-life at Q12W SC dosing results in more than 90% of patient with C.sub.trough ≥ 6 $\mu\text{g/mL}$. Dosing frequencies that maintained a clinically relevant AUC.sub.ss in FIG. 5 are indicated with stars.

(261) Taken together, these data indicate Ab001 with a half-life of at least 35 days can be dosed subcutaneously with a Q12W frequency and maintain both a clinically relevant C.sub.trough and AUC.sub.ss.

Example 4: Binding Affinity of $\alpha 4\beta 7$ Binding Antibodies

(262) This Example describes the binding affinity of the $\alpha 4\beta 7$ binding $\alpha 4\beta 7$ binding antibody Ab001.

(263) Briefly, binding affinity of the $\alpha 4\beta 7$ binding $\alpha 4\beta 7$ binding antibody Ab001 to $\alpha 4\beta 7$ was

determined by Kinetic Exclusion Assay (KinExA) and flow cytometry, and functional activity was determined by inhibition of MAdCAM-mediated cellular adhesion.

(264) The $\alpha 4\beta 7$ binding $\alpha 4\beta 7$ binding antibody Ab001 binds specifically to $\alpha 4\beta 7$ and not to the related integrins $\alpha 4\beta 1$ and $\alpha E\beta 7$. In cellular assays, Ab001 binds $\alpha 4\beta 7$ on the surface of RPMI-8866 cells with an affinity matching that of a comparator antibody, but it does not bind to Ramos cells expressing $\alpha 4\beta 1$. The $\alpha 4\beta 7$ binding $\alpha 4\beta 7$ binding antibody Ab001 demonstrates high affinity for $\alpha 4\beta 7$, with a K_{sub}.D of 836 μ M determined by KinExA. Functionally, the $\alpha 4\beta 7$ binding $\alpha 4\beta 7$ binding antibody Ab001 potently inhibits MAdCAM-1-mediated cellular adhesion with an IC_{sub}.50 comparable to that of a comparator antibody but has no inhibitory activity for VCAM-1-mediated cell adhesion.

Example 5: Simulation of Serum Concentrations of the $\alpha 4\beta 7$ Binding Antibody

(265) A simulation was performed using induction scenarios as follows: (1) with an $\alpha 4\beta 7$ binding $\alpha 4\beta 7$ binding antibody described herein such as Ab001 at subcutaneous doses of 600 mg at week 0 (W0) and 300 mg at week 2 (W2) to determine the average serum concentration during the induction period of 12 weeks; (2) with the comparator antibody at intravenous dose of 300 mg at week 0, week 2 and week 6; and (3) with the comparator antibody at intravenous dose of 300 mg at week 0 and week 2, and a subcutaneous dose of 108 mg Q2W starting at week 6. FIG. 7A shows that C_{sub}.avgW0-12 of Ab001 is greater than C_{sub}.avgW0-12 of comparator antibody. FIG. 7C shows that C_{sub}.avgW0-12 of Ab001 is greater than C_{sub}.avgW0-12 of comparator antibody. Simulated patients with Q12W SC achieve a C_{sub}.trough $\geq$ 36 μ g/mL which was higher than C_{sub}.trough of Simulated patients with IV administration of the comparator antibody and which was higher than C_{sub}.trough of simulated patients with IV/SC administration of the comparator antibody. In addition, simulated patients with Q12W SC achieve maintain the C_{sub}.trough $\geq$ 36 μ g/mL during the induction period.

(266) FIG. 7B depicts serum concentrations based on 300 mg Q8W IV dosing of comparator antibody and 300 mg Q12W SC dosing of the $\alpha 4\beta 7$ binding antibody described herein such as Ab001 based on the half-life of the antibodies. FIG. 7B shows that Q12W SC dosing of the $\alpha 4\beta 7$ binding antibody described herein such as Ab001 results in a maintenance target C_{sub}.trough $\geq$ 6 μ g/mL.

(267) FIG. 7D and FIG. 7E show that simulated patients with Q12W SC achieve an effective minimum average serum or plasma concentration of the $\alpha 4\beta 7$ binding antibody described herein such as Ab001.

(268) FIG. 7F depicts serum concentrations based on dosing of comparator antibody at intravenous dose of 300 mg at week 0 and week 2, and a subcutaneous dose of 108 mg Q2W starting at week 6 (26 injections per year for maintenance), and based on dosing of the $\alpha 4\beta 7$ binding antibody described herein such as Ab001 at subcutaneous doses of 300 mg Q12W (4-6 injections per year for maintenance), based on the half-life of the antibodies. FIG. 7F shows that the $\alpha 4\beta 7$ binding antibody described herein such as Ab001 has a superior steady state profile for patients.

Example 6: PK Simulation of Serum Concentrations of the $\alpha 4\beta 7$ Binding Antibody

(269) FIGS. 8A-8C are graphs showing human PK simulation (Simulated Using 75 mg/kg NHP—SC Dose (n=6, M/F)). FIG. 8A is a graph showing the $\alpha 4\beta 7$ binding antibody described herein such as Ab001 simulated human profiles at 600 mg. FIG. 8B is a graph showing $\alpha 4\beta 7$ binding antibody described herein such as Ab001 simulated human profiles 600 mg Q26W. FIG. 8C is a graph showing $\alpha 4\beta 7$ binding antibody described herein such as Ab001 simulated human profiles 400 mg, 200 mg, Q12W.

(270) FIG. 8A shows that a single 600 mg dose remains above the induction threshold concentration for 12 weeks and drops below the maintenance concentration at 32 weeks.

(271) FIG. 8B shows that simulation dosing of 600 mg $\alpha 4\beta 7$ binding antibody described herein such as Ab001 Q26W (Q6M) maintains induction and maintenance concentration.

(272) FIG. 8C shows that simulation dosing of 400 mg, 200 mg $\alpha 4\beta 7$ binding antibody described

herein such as Ab001 Q12W maintain induction and maintenance concentration.

Example 7: Summary of Key Binding Characteristics, Preclinical In Vitro and In Vivo Studies for the $\alpha 4\beta 7$ Binding Antibody Ab001

(273) TABLE-US-00009 TABLE 5 Study Summary Study Description Study Name and Methods Key Results and Conclusions Functional Test system: $\alpha 4\beta 7$ - the $\alpha 4\beta 7$ binding antibody Ab001 potency, $\alpha 4\beta 7$: Characterization positive RPMI-8866 EC.sub.50 = 0.15 nM of the $\alpha 4\beta 7$ cells, $\alpha 4\beta 1$ - positive the $\alpha 4\beta 7$ binding antibody Ab001 potency, binding Ramos cells $\alpha 4\beta 1$: EC.sub.50 = NB antibody $\alpha 4\beta 7$ and $\alpha 4\beta 1$ double the $\alpha 4\beta 7$ binding antibody Ab001 potency, $\alpha 4\beta 7$, Ab001 on $\alpha 4\beta 7$ positive HuT-78 cells; $\alpha 4\beta 1$: EC.sub.50 = 0.25 nM & $\alpha 4\beta 1$ fluorescence-activated Comparator antibody potency, $\alpha 4\beta 7$: cell sorting (FACS) EC.sub.50 = 0.14 nM Concentration range: 5 Comparator antibody potency, $\alpha 4\beta 1$: EC.sub.50 = NB pM-100 nM Comparator antibody potency, $\alpha 4\beta 1$, 3 cell lines; 6 $\alpha 4\beta 7$: EC.sub.50 = 0.24 nM concentrations; three The $\alpha 4\beta 7$ binding antibody Ab001 exhibits replicates functional, high-potency binding to $\alpha 4\beta 7$ but not to $\alpha 4\beta 1$. Natalizumab was included in this study as a positive control for functional $\alpha 4\beta 1$ binding. Binding Potencies Test system: Receptor the $\alpha 4\beta 7$ binding antibody Ab001; Human whole and Receptor Occupancy (IC.sub.50) on blood: IC.sub.50 for all cell types = 0.53 nM Occupancy of the CD4+, CD20+, and the $\alpha 4\beta 7$ binding antibody Ab001; Human PBMCs: $\alpha 4\beta 7$ binding CD45RO+ Cells from IC.sub.50 for all cell types = 0.38 nM antibody Ab001 Human Donors PBMCs the $\alpha 4\beta 7$ binding antibody Ab001; Monkey and Monkey Donors PBMCs: IC.sub.50 for all cell types = 0.50 nM PBMCs; titration of Comparator antibody; Human whole blood: IC.sub.50 for unlabeled Ab001 into all cell types = 0.44 nM samples equilibrated with Comparator antibody; Human PBMCs: IC.sub.50 for all EC50 concentrations of cell types = 0.38 nM fluorescently labeled Comparator antibody; Monkey PBMCs: IC.sub.50 for all Ab001 cell types = 0.56 nM Determination of Test system: KinExA The $\alpha 4\beta 7$ binding antibody Ab001 K.sub.D v. K.sub.D against $\alpha 4\beta 7$, and surface plasmon $\alpha 4\beta 7$ = 836 pM $\alpha 4\beta 1$, and $\alpha E\beta 7$ resonance The $\alpha 4\beta 7$ binding antibody Ab001 K.sub.D v. by SPR or Concentration range: $\alpha 4\beta 1$ = NB KinExA Varied by target The $\alpha 4\beta 7$ binding antibody Ab001 K.sub.D v. n = 5 concentrations $\alpha E\beta 7$ = NB Comparator antibody K.sub.D v. $\alpha 4\beta 7$ = 562 pM Comparator antibody K.sub.D v. $\alpha 4\beta 1$ = NB Comparator antibody K.sub.D v. $\alpha E\beta 7$ = NB Natalizumab was included in this study as a positive control for functional $\alpha 4\beta 1$ binding. Species Cross- Test system: SPR with the $\alpha 4\beta 7$ binding antibody Ab001 Cyno- $\alpha 4\beta 7$ K.sub.D = reactivity and multiple cycle kinetics 2.72 nM Binding Affinity the $\alpha 4\beta 7$ binding antibody Ab001 Dog- $\alpha 4\beta 7$ K.sub.D = Determination 4.81 nM the $\alpha 4\beta 7$ binding antibody Ab001 Rabbit- $\alpha 4\beta 7$ K.sub.D = 6.26 nM the $\alpha 4\beta 7$ binding antibody Ab001 Rat- $\alpha 4\beta 7$ K.sub.D = NB the $\alpha 4\beta 7$ binding antibody Ab001 Mouse- $\alpha 4\beta 7$ K.sub.D = NB the $\alpha 4\beta 7$ binding antibody Ab001 Human- $\alpha 4\beta 7$ K.sub.D = 4.65 nM Comparator antibody Cyno- $\alpha 4\beta 7$ K.sub.D = 1.2 nM Comparator antibody Dog- $\alpha 4\beta 7$ K.sub.D = 1.77 nM Comparator antibody Rabbit- $\alpha 4\beta 7$ K.sub.D = 5.34 nM Comparator antibody Rat- $\alpha 4\beta 7$ K.sub.D = NB Comparator antibody Mouse- $\alpha 4\beta 7$ K.sub.D = NB Comparator antibody Human- $\alpha 4\beta 7$ K.sub.D = 4.16 nM Binding of Ab001 Test system: binding to In human PBMCs, a dose-dependent increase in to Cells Isolated CD4.sup.+ T cells, Ab001 binding was observed in CD4.sup.+ T cells, from Human, CD4.sup.+CD45RO+ (human CD4.sup.+ CD45RO.sup.+ + memory T cells, and CD20.sup.+ B Rabbit, Mouse, only), and either CD20.sup.+ cells. In rabbits, a dose-dependent increase in and Rat PBMCs B cells or IgG.sup.+ cells from Ab001 binding was observed in a subset of PBMCs. cryopreserved PBMCs No binding above background was observed in cells from human, mouse, rat, isolated from either mice or rats. The results were and rabbit, using flow comparable to those of comparator antibody. cytometry Binding of Ab001 Test system: binding to Ab001 preferentially bound to $\alpha 4\beta 7$ -expressing B to Human primary human cells, NK cells, and T cells (naïve, TEMRA, Monocytes, B monocytes, B cells, T effector memory, and central memory) similar to Cells, NK Cells, cells (naïve, TEMRA, comparator antibody and T cell effector memory cells Subtypes (EM), central memory cells (CM) and a human B-lymphoid cell line expressing $\alpha 4\beta 7$ and $\alpha 4\beta 1$ (RPMI-8226) was evaluated using fluorescence-activated cell

sorting (FACS) and fluorescently labeled antibody Functional Test system: $\alpha 4\beta 1$ and MAdCAM-1 Inhibition IC₅₀: Characterization of $\alpha 4\beta 7$ expressing HuT-78 the $\alpha 4\beta 7$ binding antibody Ab001 = 0.086 nM the $\alpha 4\beta 7$ binding cells with coating of Comparator antibody = 0.083 nM antibody Ab001 recombinant VCAM-1 Inhibition IC₅₀: on MAdCAM-1 MAdCAM-1 the $\alpha 4\beta 7$ binding antibody Ab001 = None and VCAM-1 or VCAM-1 Comparator antibody = None Mediated Concentration range: 5 Natalizumab = 0.23 nM Adhesion pM-100 nM Conclusion: the $\alpha 4\beta 7$ binding antibody Ab001 and n = 6 replicates a comparator antibody exhibit functional activities that are distinct from those of natalizumab. Pharmacokinetic Doses: 10 mg/kg T.sub.1/2 = 17 days (experiment 1) study of the $\alpha 4\beta 7$ Route of Administration: T.sub.1/2 = 22 days (experiment 2) binding antibody IV Conclusion: The half-life of the $\alpha 4\beta 7$ binding Ab001 in N = 6 per group antibody Ab001 in male cynomolgus monkeys is cynomolgus 70% longer than that of a comparator antibody. monkeys Mouse Tg276 Dose: 1, 5, and 10 mg/kg Conclusion: the $\alpha 4\beta 7$ binding antibody Ab001 pharmacokinetic Route of administration: exhibited extended pharmacokinetic half-life in study IV Tg276 mice (>200% longer than a comparator N = 5 males per group antibody). K.sub.D: equilibrium dissociation constant between the antibody and its antigen IC.sub.50: half-maximal inhibitory concentration EC.sub.50: half maximal effective concentration MAdCAM-1: Mucosal vascular addressin cell adhesion molecule 1 VCAM-1: Vascular cell adhesion protein 1

Example 8: Formulations

(274) $\alpha 4\beta 7$ binding antibody Ab001 drug substance has a deglycosylated molecular weight of 146,698.3 daltons. It is soluble to greater than 200 mg/mL as a colorless to brown, clear to opalescent liquid.

(275) $\alpha 4\beta 7$ binding antibody Ab001 drug product is a sterile liquid for injection supplied in 2 mL glass vials. Each single-use drug product vial contains a nominal volume of 2 mL of 150 mg/mL Ab001 in aqueous solution with histidine, arginine, ethylenedinitrotetraacetic acid disodium salt dihydrate (EDTA), and polysorbate 80.

Example 9: Toxicology and Toxicokinetics of the $\alpha 4\beta 7$ Binding Antibody Ab001

(276) Toxicokinetic assessment after single dose IV and SC dosing of $\alpha 4\beta 7$ binding antibody Ab001 at 30, 75, and 150 mg/kg was conducted in male and female cynomolgus monkeys over 27 days. No PK outliers were observed, thus all animals were considered in the statistical analysis. $\alpha 4\beta 7$ binding antibody Ab001 serum concentration-time profiles in male monkeys dosed intravenously (IV) displayed a biphasic decline with a mean terminal phase t.sub.1/2 of 6.19±1.72 days, mean AUC.sub.last of 27200 4880 day.Math.µg/mL and a mean Vss of 53.0±4.30 mL/kg. A sharp increase in CL was noted after 14 days, consistent with the development of ADAs. When male and female monkeys were dosed SC, T.sub.max values ranged between 1 and 7 days. Ab001 exhibited linear dose proportionality when considering C.sub.max and AUC.sub.last. Mean terminal t.sub.1/2 values ranged from 10.2 to 24.2 days. There were no statistical differences (t test, p=0.05) in exposure at the 75 mg/kg dose between male and female monkeys when considering C.sub.max and AUC.

(277) $\alpha 4\beta 7$ binding antibody Ab001 serum concentration-time profiles in male monkeys dosed IV (150 mg/kg) displayed a biphasic decline with a mean (+SD) terminal phase t.sub.1/2 of 6.19±1.72 days, mean (+SD) AUC.sub.last of 27200±3200 day.Math.µg/mL and a mean (+SD) Vss of 53.0 4.30 mL/kg. A sharp increase in CL was noted after 14 days in this cohort, indicating the probable formation of ADAs.

(278) T.sub.max values for male and female monkeys dosed SC ranged between 1 and 7 days. $\alpha 4\beta 7$ binding antibody Ab001 exhibited approximately dose-linear pharmacokinetics when considering C.sub.max and AUC.sub.last. Mean terminal t.sub.1/2 values ranged from 10.2 to 24.2 days. There were no statistical differences (t-test, p=0.05) in exposure at 75 mg/kg between male and female monkeys when considering C.sub.max and AUC.sub.last.

(279) Overall, serum exposures of $\alpha 4\beta 7$ binding antibody Ab001 after SC dosing were proportional to dose when considering C.sub.max (maximum mean concentration) and AUC.sub.last (area under

the concentration/time curve from 0 to 648 hr) (Table 15). Mean ($\pm$ SD) C.sub.max values ranged from 297 ± 46.9 to 1780 ± 315 $\mu\text{g/mL}$, and AUC.sub.last ranged from 4590 ± 600 to 30500 ± 6460 day.Math. $\mu\text{g/mL}$. Exposures (AUC.sub.last and C.sub.max) were similar in male and female animals after a 75 mg/kg SC dose. Ab001 IV dosing at 150 mg/kg resulted in a mean AUC.sub.last exposure of 27200 ± 3200 day.Math. $\mu\text{g/mL}$ and a mean C.sub.max of 3670 ± 433 $\mu\text{g/mL}$.

(280) $\alpha 4\beta 7$ binding antibody Ab001 was then administered to male and female cynomolgus monkeys ($n=3/\text{sex}/\text{group}$) every two weeks (Q2W) (doses on Day 1, Day 15, and Day 29) for 1 month at 0 (SC and IV; vehicle control), 25 (SC), 80 (SC), 160 (SC), and 160 mg/kg (IV). All animals survived until their scheduled necropsy on Study Day 30. Bioanalytical samples were collected, processed, and analyzed using a validated GyroLab® bioanalytical method, and toxicokinetic parameters were determined using noncompartmental analysis. AUC.sub.0-28 d and C.sub.max increased in a slightly more than dose-proportional manner after Q2W SC doses of 25, 80, and 160 mg/kg. No sex differences in exposure were observed. Following a single SC administration, median T.sub.max values were observed between 3.00 to 7.00 days post dose across dose levels, while T.sub.max values after the Day 15 dose were between 1.00 and 7.00 days. Mean ($\pm$ SD) AUC.sub.0-28 d exposures (combined sex) after biweekly SC dosing were 6180 ± 2950 , 28200 ± 2260 , and 71500 ± 17000 day.Math. $\mu\text{g/mL}$, respectively. The C.sub.max ($\pm$ SD) after the SC doses on Day 15 was 430 ± 291 , 1380 ± 151 , and 4040 ± 1380 $\mu\text{g/mL}$, respectively. The mean AUC.sub.0-28 d ($\pm$ SD) exposure (combined sex) after biweekly IV dosing at 160 mg/kg was 73200 ± 12500 day.Math. $\mu\text{g/mL}$, while the C.sub.max ($\pm$ SD) was 4980 ± 905 $\mu\text{g/mL}$. The mean accumulation ratio after a second biweekly dose was <1.4 (AUC) for all doses.

(281) Anti- $\alpha 4\beta 7$ binding antibody Ab001 binding antibodies were associated with decreased exposure in two females and one male animal at the 25 mg/kg SC dose, although these animals were still included in the TK calculations for this dose. The formation of ADAs had no impact on exposure at either the 80 mg/kg or the 160 mg/kg dose (NOAEL). Throughout the evaluation of the study, serum $\alpha 4\beta 7$ binding antibody Ab001 concentrations (at C.sub.min for 25 mg/kg dose) were a minimum of $3300\times$ above the EC50 of Ab001 for saturable binding. Exposures (C.sub.max and AUC.sub.0-28 d) were slightly more than dose-proportional, and no sex-related differences were observed.

(282) Mean ($\pm$ SD) AUC.sub.0-28 d exposures (combined sex) after biweekly SC dosing were 6180 ± 2950 , 28200 ± 2260 , and 71500 ± 17000 day.Math. $\mu\text{g/mL}$ at 25, 80, and 160 mg/kg, respectively. Mean ($\pm$ SD) C.sub.max after SC doses on Day 15 were 430 ± 291 , 1380 ± 151 , and 4040 ± 1380 $\mu\text{g/mL}$ at 25, 80, and 160 mg/kg, respectively. The mean ($\pm$ SD) AUC.sub.0-28 d exposure (combined sex) after biweekly IV dosing at 160 mg/kg was 73200 ± 12500 day.Math. $\mu\text{g/mL}$, while the mean ($\pm$ SD) C.sub.max was 4980 ± 905 $\mu\text{g/mL}$. The mean accumulation ratio after a second biweekly dose was ≤ 1.40 (AUC) for all doses.

(283) It was not clear whether $\alpha 4\beta 7$ binding antibody Ab001 would demonstrate toxicity because of its half-life extension. The results showed that, despite this prior uncertainty, there were no macroscopic (e.g., organs) or microscopic (e.g., cells/tissues) findings at any dose.

Example 10: Human PK

(284) Serum $\alpha 4\beta 7$ binding antibody Ab001 concentration-time profiles in humans were predicted for single and multiple SC and IV injections of $\alpha 4\beta 7$ binding antibody Ab001 over the dose range of 100 to 1000 mg. Steady-state $\alpha 4\beta 7$ binding antibody Ab001 serum exposures were predicted using the following two resources: (1) $\alpha 4\beta 7$ binding antibody Ab001 PK parameters in cynomolgus monkeys with interspecies extrapolation of CL parameters using published allometric exponents for half-life engineered monoclonal antibodies; and (2) literature values of PK parameters reported for vedolizumab, an approved therapeutic human IgG1 monoclonal antibody targeting $\alpha 4\beta 7$ (Rosario et al. 2016, Clinical Drug Investigation 36 (11): 913-23).

(285) $\alpha 4\beta 7$ binding antibody assumptions were made: (i) Ab001 follows a biphasic decline after administration; (ii) pharmacokinetic parameters including CL and Vss do not change upon repeated

dosing and do not change over the dose range of 100 to 1000 mg in a 70 kg human (as observed with vedolizumab); (iii) disease state does not have a significant influence on PK; (iv) target-mediated drug disposition (TMDD) has a small influence on PK; (v), there will be no anti- $\alpha 4\beta 7$ binding antibody Ab001 antibody development or, if present, anti $\alpha 4\beta 7$ binding antibody Ab001 antibodies do not have an impact on $\alpha 4\beta 7$ binding antibody Ab001 PK.

(286) The predicted terminal phase $t_{sub.1/2}$ value of $\alpha 4\beta 7$ binding antibody Ab001 in humans based on the above assumptions is approximately 64 days and is consistent with the expected extended half-life resulting from incorporation of the YTE mutation into the Fc domain.

Vedolizumab has a reported human half-life of 25.5 days (Rosario et al. 2015, *Alimentary Pharmacology & Therapeutics* 42 (2): 188-202; Rosario et al. 2016). The predicted steady-state $C_{sub.max}$ and AUC values for $\alpha 4\beta 7$ binding antibody Ab001 in humans for single and multiple IV and SC dosing are shown in Table 6.

(287) TABLE-US-00010 TABLE 6 Predicted Exposure in Humans After Single and Multiple SC and IV Administration Clinical Single Dose Two Doses Given Q2W Dose $C_{sub.max}$ AUC_{sub.0-28d} AUC_{sub.0-inf} $C_{sub.max}$ AUC_{sub.0-28d} AUC_{sub.0-inf} (mg/70 kg) Route $\mu g/mL$ day • $\mu g/mL$ day • $\mu g/mL$ day • $\mu g/mL$ day • $\mu g/mL$ 100 SC 12.0 281 568 22.3 436 1450 300 37.2 900 2530 69.6 1360 5830 600 75.0 1830 5880 141 2760 12800 900 113 2760 9370 212 4150 19800 1000 125 3070 10500 236 4620 22100 100 IV 33.3 432 874 47.6 695 2150 300 100 1360 3640 145 2140 8170 600 200 2740 8230 291 4320 17500 900 300 4130 12900 437 6490 26800 1000 333 4590 14500 486 7220 29900

Example 11: PK Induction Simulation of Serum Concentrations of the $\alpha 4\beta 7$ Binding Antibody

(288) A simulation was performed using the following induction scenarios (1) with an $\alpha 4\beta 7$ binding antibody described herein such as Ab001 at subcutaneous doses of 600 mg at week 0 (W0) and 300 mg at week 2 (W2) to determine the average serum concentration during the induction period of 12 weeks, (2) with an $\alpha 4\beta 7$ binding antibody described herein such as Ab001 at subcutaneous doses of 600 mg at week 0 (W0), 300 mg at week 2 (W2) and 300 mg at week 6 (W6) to determine the average serum concentration during the induction period of 12 weeks, (3) with the comparator antibody at intravenous dose of 300 mg at week 0 (W0), week 2 (W2) and week 6 (W6) to determine the average serum concentration during the induction period of 12 weeks.

(289) An analysis of FIG. 10A indicates that $C_{sub.maxW0-12}$, $C_{sub.minW0-12}$, and $C_{sub.avgW0-12}$ of Ab001 at the subcutaneous induction dose of 600 mg at week 0 (W0) and 300 mg at week 2 (W2) was calculated to be 89.7 $\mu g/mL$, 37.4 $\mu g/mL$, and 60.7 $\mu g/mL$, respectively. An analysis of FIG. 10B indicates that $C_{sub.maxW0-12}$, $C_{sub.minW0-12}$, and $C_{sub.avgW0-12}$ of Ab001 at the subcutaneous induction dose of 600 mg at week 0 (W0), 300 mg at week 2 (W2) and 300 mg at week 6 (W6) was calculated to be 94.7 $\mu g/mL$, 57.6 $\mu g/mL$, and 73.9 $\mu g/mL$, respectively. An analysis of FIG. 10C indicates that $C_{sub.maxW0-12}$, $C_{sub.minW0-12}$, and $C_{sub.avgW0-12}$ of the comparator antibody at the intravenous dose of 300 mg at week 0 (W0), week 2 (W2) and week 6 (W6) was calculated to be 164 $\mu g/mL$, 22.9 $\mu g/mL$, and 65.1 $\mu g/mL$, respectively.

Example 12: PK Maintenance Simulation of Serum Concentrations of the $\alpha 4\beta 7$ Binding Antibody

(290) A simulation was performed using maintenance scenarios (1) with an $\alpha 4\beta 7$ binding antibody described herein such as Ab001 at subcutaneous doses of 600 mg every 26 weeks to determine the average serum concentration during the maintenance period from week 30 through week 60, (2) with an $\alpha 4\beta 7$ binding antibody described herein such as Ab001 at subcutaneous doses of 300 mg every 12 weeks to determine the average serum concentration during the maintenance period from week 30 through week 60, (3) with the comparator antibody at intravenous dose of 300 mg every 8 weeks to determine the average serum concentration during the maintenance period from week 30 through week 60, and (4) with the comparator antibody at subcutaneous dose of 108 mg every 2 weeks to determine the average serum concentration during the maintenance period from week 30

through week 60.

(291) An analysis of FIG. 11A indicates that C.sub.maxW30-60, C.sub.minW30-60, and C.sub.avgW30-60 of Ab001 at the subcutaneous maintenance dose of 600 mg every 26 weeks was calculated to be 74.8 µg/mL, 7.13 µg/mL, and 31.2 µg/mL, respectively. An analysis of FIG. 11B indicates that C.sub.maxW30-60, C.sub.minW30-60, and C.sub.avgW30-60 of Ab001 at the subcutaneous maintenance dose of 300 mg every 12 weeks was calculated to be 50.6 µg/mL, 17.8 µg/mL, and 31.3 µg/mL, respectively. An analysis of FIG. 11C indicates that C.sub.maxW30-60, C.sub.minW30-60, and C.sub.avgW30-60 of the comparator antibody at the intravenous dose of 300 mg every 8 weeks was calculated to be 136 µg/mL, 10.1 µg/mL, and 37.6 µg/mL, respectively. An analysis of FIG. 11D indicates that C.sub.maxW30-60, C.sub.minW30-60, and C.sub.avgW30-60 of the comparator antibody at the subcutaneous dose of 108 mg every 2 weeks was calculated to be 33.7 µg/mL, 27.2 µg/mL, and 31.1 µg/mL, respectively.

Example 13: Single Ascending Dose Simulation of α4β7 Binding Antibody

(292) A single ascending dose simulation was performed for two scenarios: (1) with an α4β7 binding antibody described herein such as Ab001 at 600 mg to determine serum concentration of Ab001 during the period of 84 days from the administration, and (2) with an α4β7 binding antibody described herein such as Ab001 at 300 mg to determine serum concentration of Ab001 during the period of 84 days from the administration.

(293) FIG. 12 shows a comparison of simulated serum concentration of Ab001 at 600 mg and 300 mg dose of Ab001. An analysis of FIG. 12 indicates that serum concentration of Ab001 in both scenarios remain within concentration range of 10 µg/mL and 40 µg/mL after 84 days of administration.

Example 14: Multiple Ascending Dose Simulation of α4β7 Binding Antibody

(294) A multiple ascending dose simulation was performed for two scenarios: (1) with an α4β7 binding antibody described herein such as Ab001 at subcutaneous doses of 600 mg on Day 0 (D0) followed by 600 mg subcutaneous dose on Day 14 (D14) to determine serum concentration of Ab001 during the period of 56 days from the administration, and (2) with an α4β7 binding antibody described herein such as Ab001 at subcutaneous doses of 600 mg on Day 0 (D0) followed by 600 mg subcutaneous dose on Day 14 (D14) to determine serum concentration of Ab001 during the period of 56 days from the administration.

(295) FIG. 13 shows a comparison of simulated serum concentration of Ab001 at 600 mg and 300 mg dose of Ab001. An analysis of FIG. 13 indicates that serum concentration of Ab001 in both scenarios remain above 40 µg/mL after 56 days of administration.

Example 15: Clinical Efficacy of α4β7 Binding Antibody

(296) Vedolizumab has demonstrated some efficacy in treatment of ulcerative colitis (UC) and Crohn's disease (CD) (ENTYVIO Product Monograph). This antibody has a serum half-life of 26 days in human subjects (25 days according to the product insert). Various factors increase clearance of vedolizumab, including albumin, body weight, fecal calprotectin, prior treatment with TNF antagonist drugs, and presence of anti-vedolizumab antibody. In patients administered 300 mg vedolizumab as a 30-minute intravenous infusion on Weeks 0 and 2, median serum trough concentration at Week 6 was 25.6 mcg/mL (range 0.9 to 140.0) in ulcerative colitis and 24.5 mcg/mL (range 1.1 to 177.0) in Crohn's Disease. Median steady state serum trough concentrations were 9.8 mcg/mL (range 2.4 to 42.8) and 11.2 mcg/mL (0.4 to 54.5), respectively, in patients with ulcerative colitis and Crohn's disease, when 300 mg vedolizumab was administered intravenously every eight weeks starting at week 6.

(297) Clinical trials evaluated efficacy and safety of intravenous (for induction) and intravenous or subcutaneous vedolizumab for the maintenance treatment of adult patients with moderately to severely active ulcerative colitis (Mayo score 6 to 12 with endoscopic sub score ≥2). The primary endpoint of the induction phase was proportion of patients with clinical response at Week 6. The primary endpoint for the subcutaneous study was the proportion of patients in clinical remission

(complete Mayo score of ≤ 2 points and no individual subscore >1 point) at Week 52. Secondary endpoints were mucosal healing (Mayo endoscopic subscore of ≤ 1 point) at Week 52; durable clinical response (clinical response at Weeks 6 and 52); durable clinical remission (clinical remission at Weeks 6 and 52); and corticosteroid-free clinical remission (patients using oral corticosteroids at baseline who had discontinued corticosteroids and were in clinical remission) at Week 52.

(298) The efficacy and safety of intravenous (for induction) and subcutaneous (for maintenance) vedolizumab for the treatment of adult patients with moderately to severely active Crohn's disease (CDAI score of 220 to 450) was evaluated in a randomized, double-blind, placebo-controlled study evaluating efficacy endpoints at Week 6 (induction) and Week 52 (maintenance). The primary endpoint was the proportion of patients with clinical remission (CDAI score ≤ 150). The secondary endpoints were enhanced clinical response; corticosteroid free remission; and clinical remission in TNF α antagonist naïve patients, at Week 52.

(299) $\alpha 4\beta 7$ binding antibody Ab001 is evaluated in clinical trials in patients with ulcerative colitis and Crohn's disease with various primary and secondary endpoints, such as clinical remission in both ulcerative colitis and Crohn's disease, endoscopic improvement (mucosal healing) in ulcerative colitis, durable clinical response in ulcerative colitis, durable clinical remission in ulcerative colitis, enhanced clinical response in Crohn's disease, clinical remission in advanced therapy (such as a TNF α antagonist) naïve patients in Crohn's disease, and corticosteroid-free clinical remission in both ulcerative colitis and Crohn's disease. Biomarker tests are conducted at Week 6, including C-reactive protein (CRP), fecal calprotectin, and symptoms.

(300) In a clinical study, the characteristics of the $\alpha 4\beta 7$ binding antibody Ab001 provide for greater proportion of patients achieving clinical remission in both ulcerative colitis and Crohn's disease than previously observed with vedolizumab.

(301) In a clinical study, the characteristics of the $\alpha 4\beta 7$ binding antibody Ab001 provide for greater endoscopic improvement in ulcerative colitis than previously observed for vedolizumab.

(302) In a clinical study, the characteristics of the $\alpha 4\beta 7$ binding antibody Ab001 provide for greater mucosal healing in ulcerative colitis than previously observed for vedolizumab.

(303) In a clinical study, the characteristics of the $\alpha 4\beta 7$ binding antibody Ab001 provide for greater durable clinical response in ulcerative colitis than previously observed for vedolizumab.

(304) In a clinical study, the characteristics of the $\alpha 4\beta 7$ binding antibody Ab001 provide for greater durable clinical remission in ulcerative colitis than previously observed for vedolizumab.

(305) In a clinical study, the characteristics of the $\alpha 4\beta 7$ binding antibody Ab001 provide for greater enhanced clinical response in Crohn's disease than previously observed for vedolizumab.

(306) In a clinical study, the characteristics of the $\alpha 4\beta 7$ binding antibody Ab001 provide for greater clinical remission in TNF α antagonist naïve patients in Crohn's disease than previously observed for vedolizumab.

(307) In a clinical study, the characteristics of the $\alpha 4\beta 7$ binding antibody Ab001 provide for greater corticosteroid-free clinical remission in both ulcerative colitis and Crohn's disease than previously observed for vedolizumab.

(308) In a clinical study, the characteristics of the $\alpha 4\beta 7$ binding antibody Ab001 provide for improvements in biomarkers, including CRP, fecal calprotectin, and symptoms at Week 6 than previously observed for vedolizumab.

Example 16: Two-Compartment PK Model

(309) To inform first-in-human dosing of Ab001, a two-compartment pharmacokinetic (PK) model was built and calibrated to data from cynomolgus monkeys (N=34). The model included mechanisms for intercompartmental transport and elimination of drug, and it addressed target-mediated drug disposition (TMDD) through an empirical Michaelis-Menten elimination function. The model parameters were allometrically scaled to humans using scaling exponents specific toYTE-containing mAbs. Single and biweekly administration of drug via IV and SC were simulated.

PK metrics, including the maximum drug concentration (C.sub.max), area under the PK curve (AUC) and trough concentration (C.sub.min), were calculated for each dosing scenario and are reported here.

(310) The two-compartment PK model was calibrated to preclinical PK datasets for cynomolgus monkeys for both drug molecules, Ab001 and the comparator antibody. The parameters $T_{1/2}$, V_{max} , and K_m were fitted to the mean measurements of the IV-administered data set for Ab001. For the SC administered dataset, the parameters $T_{sub.1/2}$, $V_{sub.max}$, and $K_{sub.m}$ were taken from IV fits, and optimized bioavailability (F) came out to be 100%, and absorption half-life ($T_{sub.1/2}$, SC) to be 1.08 days. The fitted parameters were used to simulate cynomolgus monkey preclinical data for both IV and SC routes.

(311) For comparator antibody preclinical fits, 10 and 50 mg/kg datasets were available for both IV and SC-administered routes. The data in the 10 mg/kg IV dose group ≥ 20 days were not included in calibration, as these points showed an anomalous increase in serum drug levels. The parameters $T_{sub.1/2}$, $V_{sub.max}$, $K_{sub.m}$, $P_{sub.dist12}$, and $V_{sub.c}$ were fitted to capture the IV-administered dataset. A lower value of optimized central volume was found for the comparator antibody, which was required to capture the C.sub.max for the IV dataset. For the SC administered dataset, the parameters $T_{sub.1/2}$, $V_{sub.max}$, $K_{sub.m}$, $P_{sub.dist12}$, and $V_{sub.c}$ were taken from the IV calibrations and the model simulations were overlaid to SC preclinical data. The bioavailability (F) and absorption half-life ($T_{sub.1/2}$, SC) were set to those from Ab001 SC dose calibrated values.

(312) The Ab001 half-life in cynomolgus monkeys was estimated to be 17.1 days, and for the comparator antibody it was estimated to be 9.74 days. Based on the respective value for each antibody and applying the allometric scaling, the half-life in humans was estimated in this study to be 63.7 days, and the critical concentration above which TMDD has a negligible effect is estimated to be 8 $\mu\text{g/mL}$. In a prior analysis the estimated half-life of Ab001 in humans was 54 days using the FcRn scaling factors. In this study the human serum half-life of the comparator antibody was estimated to be 38.5 days.

INCORPORATION BY REFERENCE

(313) All publications and patents cited throughout the text of this specification (including all patents, patent applications, scientific publications, manufacturer's specifications, instructions, etc.), whether supra or infra, are hereby incorporated by reference in their entirety for all purposes. To the extent the material incorporated by reference contradicts or is inconsistent with this specification, the specification will supersede any such material.

EQUIVALENTS

(314) The disclosure may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The foregoing embodiments are therefore to be considered in all respects illustrative rather than limiting the disclosure described herein. Scope of the disclosure is thus indicated by the appended claims rather than by the foregoing description, and all changes that come within the meaning and range of equivalency of the claims are intended to be embraced therein.

Claims

1. A method of treating a gastrointestinal inflammatory disease in a subject in need thereof, the method comprising: administering to the subject in need thereof two or more doses of a citrate-free pharmaceutical composition, wherein the citrate-free pharmaceutical composition comprises from about 150 mg/ml to about 200 mg/ml of a long acting $\alpha 4\beta 7$ targeting molecule, wherein the two or more doses are administered at least eight weeks apart, and wherein the long acting $\alpha 4\beta 7$ targeting molecule comprises: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1; and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.
2. The method of claim 1, wherein the two or more doses comprise from about 300 mg to about

450 mg of a long acting $\alpha 4\beta 7$ targeting molecule.

3. The method of claim 1, wherein the two or more doses are administered at least 12 weeks apart.

4. The method of claim 1, wherein the two or more doses are administered subcutaneously.

5. The method of claim 1, further comprising administering one or more additional doses at least two weeks before administration of the two or more doses of the citrate-free pharmaceutical composition.

6. The method of claim 5, wherein the one or more additional doses comprise a citrate-free pharmaceutical composition comprising from about 150 mg/ml to about 200 mg/ml of a long acting $\alpha 4\beta 7$ targeting molecule.

7. The method of claim 5, wherein the one or more additional doses comprise from about 500 mg to about 1200 mg of the long acting $\alpha 4\beta 7$ targeting molecule.

8. The method of claim 5, wherein the one or more additional doses are administered intravenously.

9. The method of claim 5, wherein the one or more additional doses are administered subcutaneously.

10. The method of claim 1, wherein the gastrointestinal inflammatory disease is inflammatory bowel disease.

11. The method of claim 10, wherein the inflammatory bowel disease is Crohn's disease.

12. The method of claim 10, wherein the inflammatory bowel disease is ulcerative colitis.

13. A method of treating a gastrointestinal inflammatory disease in a subject in need thereof, the method comprising: administering subcutaneously to the subject in need thereof a citrate-free pharmaceutical composition comprising at least about 180 mg/ml of a long acting $\alpha 4\beta 7$ targeting molecule that comprises: (i) a heavy chain consisting of an amino acid sequence according to SEQ ID NO: 1, and (ii) a light chain consisting of an amino acid sequence according to SEQ ID NO: 3.

14. The method of claim 13, wherein the citrate-free formulation is administered as two or more doses at least twelve weeks apart.

15. The method of claim 14, wherein the two or more doses comprise from about 300 mg to about 450 mg of the long acting $\alpha 4\beta 7$ targeting molecule.

16. The method of claim 14, further comprising administering one or more additional doses at least two weeks before subcutaneous administration of the two or more doses of the citrate-free pharmaceutical composition.

17. The method of claim 16, wherein the one or more additional doses comprise from about 500 mg to about 1200 mg of the long acting $\alpha 4\beta 7$ targeting molecule.

18. The method of claim 13, wherein the gastrointestinal inflammatory disease is inflammatory bowel disease.

19. The method of claim 18, wherein the inflammatory bowel disease is Crohn's disease.

20. The method of claim 18, wherein the inflammatory bowel disease is ulcerative colitis.
