

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12398430
Kind Code	B2
Date of Patent	August 26, 2025
Inventor(s)	Kondou; Satoshi et al.

Breast cancer detection kit or device, and detection method

Abstract

It is intended to provide a kit or a device for the detection of breast cancer and a method for detecting breast cancer. The present invention provides a kit or a device for the detection of breast cancer, comprising nucleic acid(s) capable of specifically binding to a miRNA in a sample of a subject, and a method for detecting breast cancer, comprising measuring the miRNA in vitro.

Inventors: Kondou; Satoshi (Kamakura, JP), Nobumasa; Hitoshi (Kamakura, JP), Kozono; Satoko (Kamakura, JP), Sudo; Hiroko (Kamakura, JP), Kawauchi; Junpei (Kamakura, JP), Ochiya; Takahiro (Tokyo, JP), Kosaka; Nobuyoshi (Tokyo, JP), Ono; Makiko (Tokyo, JP), Tamura; Kenji (Tokyo, JP)

Applicant: TORAY INDUSTRIES, INC. (Tokyo, JP); NATIONAL CANCER CENTER (Tokyo, JP)

Family ID: 1000008779167

Assignee: TORAY INDUSTRIES, INC. (Tokyo, JP); NATIONAL CANCER CENTER (Tokyo, JP)

Appl. No.: 18/514603

Filed: November 20, 2023

Prior Publication Data

Document Identifier	Publication Date
US 20240093311 A1	Mar. 21, 2024

Foreign Application Priority Data

JP	2014-122672	Jun. 13, 2014
JP	2015-069321	Mar. 30, 2015

Related U.S. Application Data

division parent-doc US 17944779 20220914 US 11859255 child-doc US 18514603
division parent-doc US 16797625 20200221 US 11479822 20221025 child-doc US 17944779
division parent-doc US 15318328 US 10597726 20200324 WO PCT/JP2015/066986 20150612
child-doc US 16797625

Publication Classification

Int. Cl.: **C12Q1/68** (20180101); **C12M1/00** (20060101); **C12P19/34** (20060101); **C12Q1/6886** (20180101); C12N15/09 (20060101)

U.S. Cl.:

CPC **C12Q1/6886** (20130101); **C12M1/00** (20130101); **C12Q1/68** (20130101); C12N15/09 (20130101); C12Q2600/158 (20130101); C12Q2600/16 (20130101); C12Q2600/178 (20130101)

Field of Classification Search

CPC: C12Q (2600/158); C12Q (1/6886); C12Q (2600/178)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
8603744	12/2012	Croce et al.	N/A	N/A
2001/0053519	12/2000	Fodor et al.	N/A	N/A
2007/0299030	12/2006	Dmitrovsky et al.	N/A	N/A
2008/0076674	12/2007	Litman et al.	N/A	N/A
2011/0028332	12/2010	Kuroda et al.	N/A	N/A
2012/0115139	12/2011	Kuroda et al.	N/A	N/A
2013/0035251	12/2012	Keller et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
101921760	12/2009	CN	N/A
101988061	12/2010	CN	N/A
101988064	12/2010	CN	N/A
2 336 353	12/2010	EP	N/A
2 354 246	12/2010	EP	N/A
2008-500837	12/2007	JP	N/A
WO 2012-507300	12/2011	JP	N/A
WO 2005/118806	12/2004	WO	N/A
WO 2009/082744	12/2008	WO	N/A
WO 2009/119809	12/2008	WO	N/A
WO 2010/062706	12/2009	WO	N/A
WO 2010/100328	12/2009	WO	N/A
WO 2010/123043	12/2009	WO	N/A

WO 2012/048236	12/2011	WO	N/A
WO 2012/070037	12/2011	WO	N/A
WO 2013/057567	12/2012	WO	N/A
WO 2013/190091	12/2012	WO	N/A
WO 2014/048441	12/2013	WO	N/A
WO 2014/057279	12/2013	WO	N/A
WO 2014/081507	12/2013	WO	N/A

OTHER PUBLICATIONS

Wu, Qian, et al. "Next-Generation Sequencing of MicroRNAs for Breast Cancer Detection", BioMed Research International, 2011, 597145, pp. 1-7, 2011. (Year: 2011). cited by examiner

Chinese Office Action and Search Report for Chinese Application No. 202110418361.8, dated Jun. 27, 2024. cited by applicant

American Cancer Society, "Breast Cancer", pp. 6-9, 13, 27-28, 41-46, 52-54, 63-64, and 106 (2013). cited by applicant

Berillo et al., "Binding of intronic miRNAs to the mRNAs of host genes encoding intronic miRNAs and proteins that participate in tumourigenesis," Computers in Biology and Medicine (2013), vol. 43, pp. 1374-1381. cited by applicant

Buffa et al. "microRNA-Associated Progression Pathways and Potential Therapeutic Targets identified by integrated mRNA and microRNA Expression Profiling in Breast Cancer," Cancer Res. Sep. 1, 2011; 71(17): 5635-45. (Year:2011). cited by applicant

Carson et al. "Breast Imaging in Coronal Planes with Simultaneous Pulse Echo and Transmission Ultrasound," Science 214.4525 (1981): 1141-1143. (Year: 1981). cited by applicant

Cheung, V. G. et al., "Natural variation in human gene expression assessed in lymphoblastoid cells," Nature Genetics (2003), vol. 33, pp. 422-425. cited by applicant

Cobb et al., "Sepsis gene expression profiling: Murine splenic compared with hepatic responses determined by using complementary DNA microarrays," Crit. Care Med. (2002), vol. 30, No. 12, pp. 2711-2721. cited by applicant

Communication Pursuant to Rule 164(1) EPC issued Jan. 2, 2018, in European Patent Application No. 15805922.0. cited by applicant

Cookson et al., "Circulating microRNA profiles reflect the presence of breast tumours but not the profiles of microRNAs within the tumours", Cellular Oncology, vol. 35, No. 4, 2012, pp. 301-308. cited by applicant

Cuk et al., "Circulating microRNAs in plasma as early detection markers for breast cancer.", Int. J. Cancer, vol. 132, No. 7, 2013, pp. 1602-1612. cited by applicant

Cuk et al., "Plasma microRNA panel for minimally invasive detection of breast cancer.", Plos One, vol. 8, No. 10, e76729, 2013, 10 pages. cited by applicant

Eto et al. "Prospect of microRNA toward laboratory medicine Gastrointestinal Cancer and microRNA," Clinical Chemistry, vol. 43, 2014, 99-105. cited by applicant

Fu et al., "miRNA Biomarkers in Breast Cancer Detection and Management," Journal of Cancer (2011), vol. 2, pp. 116-122. cited by applicant

Godfrey et al., "Serum microRNA expression as an early marker for breast cancer risk in prospectively collected samples from the Sister Study cohort," Breast Cancer Research, vol. 15, No. 3, 2013, 10 pages. cited by applicant

Guadagni et al., "A Re-Evaluation of Carcinoembryonic Antigen (CEA) as a Serum Marker for Breast Cancer: A Prospective Longitudinal Study," Clinical Cancer Research, vol. 7, 2001, p. 2357-2362. cited by applicant

Hoshikawa et al., "Hypoxia induces different genes in the lungs of rats compared with mice," Physiol. Genomics (2003), vol. 12, pp. 209-219. cited by applicant

Inazawa, "Gan ni Okeru Morateki miRNA Kaiseki to Shindan Chiryo eno Oyo," Journal of

Clinical and Experimental Medicine, vol. 249, No. 10, 2014, pp. 1119-1124. cited by applicant

International Search Report for PCT/JP2015/066986 (PCT/ISA/210) mailed on Sep. 1, 2015. cited by applicant

Japanese Office Action for Japanese Application No. 2022-057348, dated Apr. 25, 2023. cited by applicant

Kojima et al., MicroRNA markers for the diagnosis of pancreatic and biliary-tract cancers, PLOS ONE (2015), vol. 10, No. 2, e0118220 (pp. 1-22). cited by applicant

Leidner et al., "Dampening Enthusiasm for Circulating MicroRNA in Breast Cancer," PloS One, vol. 8, No. 3, e57841, 2013, 11 pages. cited by applicant

MiScript™ miRNA PCR Array (384-well, 384HC) Human miRBase Profiler HC Plate 4 from Qiagen, from <https://b2b.qiagen.com/-/media/genetable/mi/hs/34/mihs-3404z> (Year: 2012). cited by applicant

Mitchell et al., "Circulating microRNAs as stable blood-based markers for cancer detection.", Proc. Natl. Acad. Sci. USA, vol. 105, No. 30, 2008, pp. 10513-10518. cited by applicant

Ng et al., "Circulating microRNAs as specific biomarkers for breast cancer detection.", Plos One, vol. 8, No. 1, e53141, 2013, 10 pages. cited by applicant

Office Action issued Jun. 29, 2021, in Japanese Patent Application No. 2020-079623. cited by applicant

Office Action issued Sep. 24, 2021, in Republic of Korea Patent Application No. 10-2017-7000876. cited by applicant

Ono et al., E-2022 "Circulating microRNA markers for detection of breast cancer," Digital Abstract for The 73rd Annual Meeting of the Japanese Cancer Association, Published (online) Sep. 19, 2014. cited by applicant

Ono et al., E-2022 "Circulating microRNA markers for detection of breast cancer," English Oral Session for The 73rd Annual Meeting of the Japanese Cancer Association, Published (online) Sep. 26, 2014. cited by applicant

Perez-Sanchez et al., "Clinical utility of microRNAs in Exhaled Breath Condensate as Biomarkers for Lung Cancer," J. Pers. Med. (2021), vol. 11, 111, pp. 1-14. cited by applicant

Persson et al., "Identification of new microRNAs In paired normal and tumor breast tissue suggests a dual role for the ERBB2/Her2 gene," Cancer Res. (2011), vol. 71, No. 1, pp. 78-86. cited by applicant

Qiagen Product description "mi Script™ miRNA PCR Array (384-well, 384HC) Human miRBase Profiler HC Plate 5", document 1073798, Aug. 2012, from <https://b2b.qiagen.com/-/media/genetable/mi/hs/34/mihs-3405z> (Year: 2012). cited by applicant

Sato, "microRNA chip ni yoru Biomarker Kaihatsu," Molecular Targeted Therapy for Cancer (2015), vol. 12, No. 4, pp. 456-465. cited by applicant

Schrauder et al., "Circulating Micro-RNAs as Potential Blood-Based Markers for Early Stage Breast Cancer Detection," PloS One, vol. 7, No. 1, e29770, 2012, 9 pages. cited by applicant

Sobin et al., "TNM Classification of Malignant Tumours," 7th edition, 2010, pp. 171-181. cited by applicant

Song et al., "Bioinformatic Prediction of SNPs within miRNA Binding Sites of Inflammatory Genes Associated with Gastric Cancer," Asian Pac. J. Cancer Prev. (2014), vol. 15, pp. 937-943. cited by applicant

Tamaki et al., "The Challenge to Reduce Breast Cancer Mortality in Okinawa: Consensus of the first Okinawa Breast Oncology Meeting," Japanese Journal of Clinical Oncology, vol. 43, No. 2, 2013, pp. 208-213. cited by applicant

Wang et al., "Plasma miR-601 and miR-760 are novel biomarkers for the early detection of colorectal cancer.", Plos One, vol. 7, No. 9, e44398, 2012, 8 pages. cited by applicant

Written Opinion of the International Searching Authority for PCT/JP2015/066986 (PCT/ISA/237) mailed on Sep. 1, 2015. cited by applicant

U.S. Appl. No. 17/944,779, filed Sep. 14, 2022. cited by applicant
U.S. Appl. No. 16/797,625, filed Feb. 21, 2020. cited by applicant
U.S. Appl. No. 15/318,328, filed Dec. 12, 2016. cited by applicant
Extended European Search Report for European Application No. 24215251.0, dated Mar. 17, 2025.
cited by applicant
Gao et al., “miR-615-5p is epigenetically inactivated and functions as a tumor suppressor in
pancreatic ductal adenocarcinoma,” *Oncogene*, vol. 34, No. 13, XP037748183, Apr. 28, 2014, pp.
1629-1640. cited by applicant
Tayebi et al., “miR-615-5p is restrictedly expressed in cirrhotic and cancerous liver tissues and its
overexpression alleviates the tumorigenic effects in hepatocellular carcinoma,” *FEBS Letters*, vol.
586, No. 19, XP071253316, Jul. 20, 2012, pp. 3309-3316. cited by applicant

Primary Examiner: Kapushoc; Stephen T

Attorney, Agent or Firm: Birch, Stewart, Kolasch & Birch, LLP

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS (1) This application is a Divisional of U.S. application Ser. No. 17/944,779 filed Sep. 14, 2022, which is a Divisional of U.S. application Ser. No. 16/797,625 filed Feb. 21, 2020 (now U.S. Pat. No. 11,479,822), which is a Divisional of U.S. application Ser. No. 15/318,328, filed on Dec. 12, 2016 (now U.S. Pat. No. 10,597,726), which is the National Phase under 35 U.S.C. § 371 of International Application No. PCT/JP2015/066986 filed on Jun. 12, 2015, which claims the benefit under 35 U.S.C. § 119(a) to Japanese Patent Application Nos. 2014-122672 filed Jun. 13, 2014, and 2015-069321 filed Mar. 30, 2015, all of which are hereby expressly incorporated by reference into the present application.

REFERENCE TO ELECTRONIC SEQUENCE LISTING

(1) The application contains a Sequence Listing which has been submitted electronically in .XML format and is hereby incorporated by reference in its entirety. Said .XML copy, created on Sep. 12, 2022, is named “PH-6234-PCT-US-DIV1-DIV1 Sequence Listing ST26” and is 784,115 bytes in size. The sequence listing contained in this .XML file is part of the specification and is hereby incorporated by reference herein in its entirety.

TECHNICAL FIELD

(2) The present invention relates to a kit or a device for the detection of breast cancer, comprising nucleic acid(s) capable of specifically binding to a particular miRNA, which is used for examining the presence or absence of breast cancer in a subject, and a method for detecting breast cancer, comprising measuring an expression level of the miRNA using the nucleic acid.

BACKGROUND ART

(3) The breast is constituted by a mammary gland which produces mother milk, lobules which arise from the mammary gland, mammary ducts which arise from the lobules and deliver milk, and fat which supports these constituents, etc. Approximately 90% of breast cancer cases originate in the mammary ducts, while approximately 5 to 10% of the breast cancer cases originate in the lobules (Non-Patent Literature 1). According to the 2011 statistics of cancer type-specific mortality in Japan disclosed by the Center for Cancer Control and Information Services, National Cancer Center, the number of breast cancer deaths climbed to 12,731 people. It is estimated that one out of 14 Japanese females will experience breast cancer. The number of incidences of this cancer in females takes the 1st place by cancer type. It is estimated that one out of 8 American females will experience breast cancer. The estimated number of American individuals affected by breast cancer

climbed to 232,670 people in 2014, among which approximately 40,000 people reportedly died (Non-Patent Literature 1).

(4) The stages of breast cancer progression are defined in Non-Patent Literature 2 and classified into stages 0, IA, IB, IIA, IIB, IIIA, IIIB, IIIC, and IV according to tumor size, infiltration, lymph node metastasis, distant metastasis, etc. The 5-year relative survival rate of breast cancer largely depends on the stages of cancer progression and is reportedly 100% for stage 0 and stage I, 93% for stage II, 72% for stage III, and 22% for stage IV (Non-Patent Literature 1). Thus, the early detection of breast cancer leads to improvement in the survival rate. Therefore, an approach that permits the early detection is strongly desired.

(5) The treatment of breast cancer is basically surgical treatment, which is used in combination with drug therapy or radiotherapy depending on the progressed stage, metastasis, general health conditions, and breast cancer classification. Particularly, for early breast cancer of stage I or II, breast conservation therapy may be selected with a combined use with radiotherapy (Non-Patent Literature 1).

(6) According to Non-Patent Literature 1, initial diagnostic tests of breast cancer include inspection and palpation as well as imaging tests such as mammography, which is breast-dedicated X-ray examination, and ultrasonography (echo examination). When there are findings on suspected breast cancer by the initial test, pathological examination which involves inserting a needle into a lesion and collecting cells or tissues to be examined under a microscope, is carried out as a secondary test. If necessary, imaging tests such as CT, MRI, abdominal ultrasonography, bone scintigraphy, and PET are also carried out in order to examine the state or spread of the lesion.

(7) For example, CEA, CA-15-3, and CA27-29 are known as tumor markers for the detection of breast cancer. These tumor markers in blood have been reported to elevate when breast cancer has metastasized to other organs such as the bone or the liver. However, these tumor markers do not elevate in some patients and may thus be limited by their usefulness (Non-Patent Literature 1).

(8) As shown in Patent Literatures 1 to 4, there are reports, albeit at a research stage, on the detection of breast cancer using the expression levels of microRNAs (miRNAs) or combinations of the expression levels of miRNAs and the expression levels of additional protein markers in biological samples including blood.

(9) Specifically, Patent Literature 1 discloses a method for detecting prostate cancer or other cancers including breast cancer by combining hsa-miR-602 or hsa-miR-135a-3p with known protein markers in blood.

(10) Patent Literature 2 discloses a method for detecting various cancers including breast cancer by combining hsa-miR-23b-3p or hsa-miR-135a-3p with 5 or more other miRNAs in blood or tissues.

(11) Patent Literature 3 discloses a method for detecting breast cancer using hsa-miR-92a-3p, hsa-miR-92a-2-5p, hsa-miR-92b-5p, and the like in blood cells.

(12) Patent Literature 4 discloses a method for detecting breast cancer using hsa-miR-451a, hsa-miR-296-5p, hsa-miR-16-5p, and the like in tissues.

(13) Non-Patent Literature 3 discloses that hsa-miR-760 and the like in blood are significantly expressed in breast cancer patients.

(14) Non-Patent Literature 4 discloses that hsa-miR-423-5p, hsa-miR-486-5p, and the like in blood are decreased after surgery of breast cancer.

(15) Non-Patent Literature 5 discloses that hsa-miR-4257, hsa-miR-1915-3p, hsa-miR-718, and the like in blood are significantly expressed in breast cancer patients.

(16) Non-Patent Literature 6 discloses that hsa-miR-940 and the like in blood are significantly expressed in breast cancer patients.

CITATION LIST

Patent Literature

(17) Patent Literature 1: JP Patent Publication (Kohyo) No. 2012-507300 A (2012) Patent

Literature 2: JP Patent Publication (Kohyo) No. 2008-500837 A (2008) Patent Literature 3:

Non-Patent Literature

(18) Non-Patent Literature 1: American Cancer Society, "Breast Cancer", 2013, p. 6-9, 13, 27-28, 41-46, 52-54, 63-64, and 106 Non-Patent Literature 2: Sobin, L. et al., "TNM Classification of Malignant Tumours, the 7th edition", 2010, p. 171 to 181 Non-Patent Literature 3: Godfrey, A C. et al., 2013, Breast Cancer Research, Vol. 15 (3), p. R42 Non-Patent Literature 4: Cookson, V J. et al., 2012, Cellular Oncology, Vol. 35 (4), p. 301-8 Non-Patent Literature 5: Schrauder, M G. et al., 2012, PLoS One, Vol. 7 (1), p. e29770 Non-Patent Literature 6: Leidner, R S. et al., 2013, PLoS One, Vol. 8 (3), p. e57841 Non-Patent Literature 7: Tamaki, K. et al., 2013, Japanese Journal of Clinical Oncology, Vol. 43 (2), p. 208-213 Non-Patent Literature 8: Guadagni, F. et al., 2001, Clinical Cancer Research, Vol. 7, p. 2357 to 2362

SUMMARY OF INVENTION

Problem to be Solved by Invention

(19) An object of the present invention is to find novel tumor markers for breast cancer and to provide a method that can effectively detect breast cancer using nucleic acid(s) capable of specifically binding to the markers. As described in Non-Patent Literature 1, initial diagnostic tests of breast cancer include inspection and palpation as well as imaging tests such as mammography, which is breast-dedicated X-ray examination, and ultrasonography. The mammography is reportedly effective as breast cancer examination targeting women aged 40 or older, and the American Cancer Society recommends that women in this age range take mammography every year (Non-Patent Literature 1). The mammography, however, has been reported to have limitations in the visualization of breast cancer present in the dense breast before menopause or a very small tumor of early breast cancer (Non-Patent Literature 1). In Japan, the mammography rate was only 24.3% in 2010, and a challenge to improvement in breast cancer survival rate will be to increase this mammography rate (Non-Patent Literature 7).

(20) For example, CEA, CA-15-3, and CA27-29 mentioned above are known as tumor markers for the detection of breast cancer. These tumor markers, however, are helpful in confirming therapeutic effects on recurrent breast cancer, but rarely elevate in early breast cancer. Therefore, the tumor markers may not be useful for the purpose of breast cancer examination (Non-Patent Literature 1). According to Non-Patent Literature 8, the specific sensitivity of CEA and CA15-3 is uselessly 6.4% and 12.2%, respectively, for stage 1 and is only 25.0% and 62.5%, respectively, even for stage 4. Thus, the tumor marker measurement is less significant as a preoperative test. Since these blood tumor markers may elevate for reasons other than breast cancer, the presence or absence of breast cancer is difficult to determine. The false diagnosis of other cancers as breast cancer wastes appropriate therapeutic opportunity or places unnecessary economical and physical burdens on patients due to the application of wrong medicine.

(21) As described below, there are reports, albeit at a research stage, on the determination of breast cancer using the expression levels of microRNAs (miRNAs) in biological samples including blood, none of which, however, have yet been brought into practical use.

(22) Patent Literature 1 discloses a method for detecting prostate cancer or other cancers including breast cancer by combining hsa-miR-602 or hsa-miR-135a-3p with known protein markers in blood. The measurement of both miRNA and protein markers, however, brings about increase in examination costs and a complicated process and is therefore not favorable. This detection method does not describe specific detection performance such as accuracy, sensitivity, or specificity for determining breast cancer and is thus industrially less practical.

(23) Patent Literature 2 discloses a method for detecting various cancers including breast cancer by combining hsa-miR-23b-3p or hsa-miR-135a-3p with 5 or more other miRNAs in blood or tissues. This detection method does not describe specific detection performance such as accuracy, sensitivity, or specificity for determining breast cancer and is thus industrially less practical.

(24) Patent Literature 3 describes a method for detecting breast cancer using hsa-miR-92a-3p, hsa-miR-92a-2-5p, hsa-miR-92b-5p, and the like. This detection method does not describe specific detection performance such as accuracy, sensitivity, or specificity for determining breast cancer and is thus industrially less practical. In addition, these miRNA markers were not validated in an independent sample group and are therefore less reliable.

(25) Patent Literature 4 discloses a method for detecting breast cancer using hsa-miR-451a, hsa-miR-296-5p, hsa-miR-16-5p, and the like in tissues. For this detection method, however, tissue resection by surgical operation is essential for obtaining samples, and this step places a heavy physical burden on patients. Therefore, this method is not favorable as an examination method. In addition, this detection method does not describe specific detection performance such as accuracy, sensitivity, or specificity for determining breast cancer and is thus industrially less practical.

(26) Non-Patent Literature 3 discloses that hsa-miR-760 and the like in blood are significantly expressed in breast cancer patients. This literature, however, neither describes detection performance such as accuracy, sensitivity, or specificity for determining breast cancer nor describes a specific method for detecting breast cancer. Therefore, this approach is industrially less practical.

(27) Non-Patent Literature 4 discloses that hsa-miR-423-5p, hsa-miR-486-5p, and the like in blood are decreased after surgery of breast cancer. This literature, however, neither describes detection performance such as accuracy, sensitivity, or specificity for determining breast cancer nor describes a specific method for detecting breast cancer. Therefore, this approach is industrially less practical.

(28) Non-Patent Literature 5 discloses that hsa-miR-4257, hsa-miR-1915-3p, hsa-miR-718, and the like in blood are significantly expressed in breast cancer patients. This approach, however, employed as many as 240 miRNAs for detecting breast cancer and might cause increase in examination cost and complicated discriminant algorithms. Thus, this approach is not industrially practical.

(29) Non-Patent Literature 6 discloses that hsa-miR-940 and the like in blood are significantly expressed in breast cancer patients. The authors, however, concluded that this marker is less reproducible, and finally abandoned the marker in the study. In addition, this literature neither describes detection performance such as accuracy, sensitivity, or specificity for determining breast cancer nor describes a specific method for detecting breast cancer. Therefore, this approach is industrially less practical.

(30) As mentioned above, the existing tumor markers exhibit low performance in the detection of breast cancer, or neither performance nor detection methods are specifically shown as to the markers at a research stage. Therefore, use of these markers might impose implementation of needless extra examination due to the false detection of healthy subjects as being breast cancer patients, or might waste therapeutic opportunity because of overlooking breast cancer patients. In addition, the measurement of dozens of miRNAs increases examination cost and is therefore difficult to use in large-scale screening such as medical checkup. Furthermore, the collection of breast tissues for measuring the tumor markers is highly invasive to patients and is not favorable. Hence, there is a demand for a highly accurate breast cancer marker that is detectable from blood, which can be collected with limitedly invasiveness, and is capable of correctly determining a breast cancer patient as a breast cancer patient and a healthy subject as a healthy subject. Particularly, the early detection and treatment of breast cancer can drastically reduce the risk of recurrence and also permit breast conservation therapy. Therefore, a highly sensitive breast cancer marker capable of detecting breast cancer even at a low progressed stage is desired. Moreover, the mammography rate is presumably increased by providing a more convenient initial screening of breast cancer.

Solution to Problem

(31) The present inventors have conducted diligent studies to attain the object and consequently completed the present invention by finding several genes usable as markers for the detection of breast cancer from blood, which can be collected with limitedly invasiveness, and finding that breast cancer can be significantly detected by using nucleic acids capable of specifically binding to

thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (g) a polynucleotide comprising a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251, (h) a polynucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (i) a polynucleotide comprising a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, and (j) a polynucleotide hybridizing under stringent conditions to any of the polynucleotides (f) to (i).

(39) (7) The kit according to any of (1) to (6), wherein the kit further comprises nucleic acid(s) capable of specifically binding to at least one polynucleotide selected from the group consisting of other breast cancer markers miR-658, miR-6842-5p, miR-6124, miR-6765-3p, miR-7106-5p, miR-4534, miR-92b-3p, miR-3135b, miR-4687-3p, miR-762, miR-3619-3p, miR-4467, miR-557, miR-1237-5p, miR-1908-5p, miR-4286, miR-6885-5p and miR-6763-5p.

(40) (8) The kit according to (7), wherein miR-658 is hsa-miR-658, miR-6842-5p is hsa-miR-6842-5p, miR-6124 is hsa-miR-6124, miR-6765-3p is hsa-miR-6765-3p, miR-7106-5p is hsa-miR-7106-5p, miR-4534 is hsa-miR-4534, miR-92b-3p is hsa-miR-92b-3p, miR-3135b is hsa-miR-3135b, miR-4687-3p is hsa-miR-4687-3p, miR-762 is hsa-miR-762, miR-3619-3p is hsa-miR-3619-3p, miR-4467 is hsa-miR-4467, miR-557 is hsa-miR-557, miR-1237-5p is hsa-miR-1237-5p, miR-1908-5p is hsa-miR-1908-5p, miR-4286 is hsa-miR-4286, miR-6885-5p is hsa-miR-6885-5p, and miR-6763-5p is hsa-miR-6763-5p.

(41) (9) The kit according to (7) or (8), wherein the nucleic acid is a polynucleotide selected from the group consisting of the following polynucleotides (k) to (o): (k) a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (l) a polynucleotide comprising a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269, (m) a polynucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (n) a polynucleotide comprising a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, and (o) a polynucleotide hybridizing under stringent conditions to any of the polynucleotides (k) to (n).

(42) (10) The kit according to any of (1) to (9), wherein the kit comprises at least two nucleic acids capable of specifically binding to at least two polynucleotides, respectively, selected from the group consisting of all of the breast cancer markers according to (1) or (2).

(43) (11) A device for the detection of breast cancer, comprising nucleic acid(s) capable of specifically binding to at least one polynucleotide selected from the group consisting of the following breast cancer markers: miR-4783-3p, miR-4730, miR-1307-3p, miR-4634, miR-663a, miR-4532, miR-7704, miR-3178, miR-6729-5p, miR-6090, miR-4732-5p, miR-3184-5p, miR-6727-5p, miR-6088, miR-4674, miR-8073, miR-4787-5p, miR-1469, miR-125a-3p, miR-1233-5p, miR-885-3p, miR-6802-5p, miR-328-5p, miR-6787-5p, miR-8069, miR-6875-5p, miR-1246, miR-4734, miR-6757-5p, miR-6756-5p, miR-3665, miR-6836-3p, miR-6821-5p, miR-6805-5p, miR-4728-5p, miR-6726-5p, miR-197-5p, miR-149-3p, miR-6850-5p, miR-4476, miR-6858-5p, miR-564, miR-4763-3p, miR-575, miR-6771-5p, miR-1231, miR-1908-3p, miR-150-3p, miR-3937, miR-887-3p, miR-3940-5p, miR-4741, miR-6808-5p, miR-6869-5p, miR-5090, miR-615-5p, miR-8072, miR-128-1-5p, miR-1238-5p, miR-365a-5p, miR-204-3p, miR-4492, miR-6785-5p, miR-

miR-4270 is hsa-miR-4270, miR-6721-5p is hsa-miR-6721-5p, miR-7111-5p is hsa-miR-7111-5p, miR-6791-5p is hsa-miR-6791-5p, miR-7109-5p is hsa-miR-7109-5p, miR-4258 is hsa-miR-4258, miR-6515-3p is hsa-miR-6515-3p, miR-6851-5p is hsa-miR-6851-5p, miR-6125 is hsa-miR-6125, miR-4749-5p is hsa-miR-4749-5p, miR-4726-5p is hsa-miR-4726-5p, miR-4513 is hsa-miR-4513, miR-6089 is hsa-miR-6089, miR-6816-5p is hsa-miR-6816-5p, miR-4466 is hsa-miR-4466, miR-4488 is hsa-miR-4488, miR-6752-5p is hsa-miR-6752-5p, and miR-4739 is hsa-miR-4739.

(45) (13) The device according to (11) or (12), wherein the nucleic acid is a polynucleotide selected from the group consisting of the following polynucleotides (a) to (e): (a) a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235 and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (b) a polynucleotide comprising a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235 and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, (c) a polynucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235 and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (d) a polynucleotide comprising a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235 and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, and (e) a polynucleotide hybridizing under stringent conditions to any of the polynucleotides (a) to (d).

(46) (14) The device according to any of (11) to (13), wherein the device further comprises nucleic acid(s) capable of specifically binding to at least one polynucleotide selected from the group consisting of other breast cancer markers miR-760, miR-602, miR-423-5p, miR-92a-2-5p, miR-16-5p, miR-451a, miR-135a-3p, miR-486-5p, miR-4257, miR-92b-5p, miR-1915-3p, miR-718, miR-940, miR-296-5p, miR-23b-3p and miR-92a-3p.

(47) (15) The device according to (14), wherein miR-760 is hsa-miR-760, miR-602 is hsa-miR-602, miR-423-5p is hsa-miR-423-5p, miR-92a-2-5p is hsa-miR-92a-2-5p, miR-16-5p is hsa-miR-16-5p, miR-451a is hsa-miR-451a, miR-135a-3p is hsa-miR-135a-3p, miR-486-5p is hsa-miR-486-5p, miR-4257 is hsa-miR-4257, miR-92b-5p is hsa-miR-92b-5p, miR-1915-3p is hsa-miR-1915-3p, miR-718 is hsa-miR-718, miR-940 is hsa-miR-940, miR-296-5p is hsa-miR-296-5p, miR-23b-3p is hsa-miR-23b-3p, and miR-92a-3p is hsa-miR-92a-3p.

(48) (16) The device according to (14) or (15), wherein the nucleic acid is a polynucleotide selected from the group consisting of the following polynucleotides (f) to (j): (f) a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (g) a polynucleotide comprising a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251, (h) a polynucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (i) a polynucleotide comprising a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, and (j) a polynucleotide hybridizing under stringent conditions to any of the polynucleotides (f) to (i).

(49) (17) The device according to any of (11) to (16), wherein the device further comprises nucleic acid(s) capable of specifically binding to at least one polynucleotide selected from the group consisting of other breast cancer markers miR-658, miR-6842-5p, miR-6124, miR-6765-3p, miR-7106-5p, miR-4534, miR-92b-3p, miR-3135b, miR-4687-3p, miR-762, miR-3619-3p, miR-4467,

miR-557, miR-1237-5p, miR-1908-5p, miR-4286, miR-6885-5p and miR-6763-5p.
(50) (18) The device according to (17), wherein miR-658 is hsa-miR-658, miR-6842-5p is hsa-miR-6842-5p, miR-6124 is hsa-miR-6124, miR-6765-3p is hsa-miR-6765-3p, miR-7106-5p is hsa-miR-7106-5p, miR-4534 is hsa-miR-4534, miR-92b-3p is hsa-miR-92b-3p, miR-3135b is hsa-miR-3135b, miR-4687-3p is hsa-miR-4687-3p, miR-762 is hsa-miR-762, miR-3619-3p is hsa-miR-3619-3p, miR-4467 is hsa-miR-4467, miR-557 is hsa-miR-557, miR-1237-5p is hsa-miR-1237-5p, miR-1908-5p is hsa-miR-1908-5p, miR-4286 is hsa-miR-4286, miR-6885-5p is hsa-miR-6885-5p, and miR-6763-5p is hsa-miR-6763-5p.

(51) (19) The device according to (17) or (18), wherein the nucleic acid is a polynucleotide selected from the group consisting of the following polynucleotides (k) to (o): (k) a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (l) a polynucleotide comprising a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269, (m) a polynucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (n) a polynucleotide comprising a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, and (o) a polynucleotide hybridizing under stringent conditions to any of the polynucleotides (k) to (n).

(52) (20) The device according to any one of (11) to (19), wherein the device is for measurement based on a hybridization technique.

(53) (21) The device according to (20), wherein the hybridization technique is a nucleic acid array technique.

(54) (22) The device according to any one of (11) to (21), wherein the device comprises at least two nucleic acids capable of specifically binding to at least two polynucleotides, respectively, selected from all of the breast cancer markers according to (11) or (12).

(55) (23) A method for detecting breast cancer, comprising measuring an expression level(s) of a target nucleic acid(s) in a sample from a subject using a kit according to any one of (1) to (10) or a device according to any one of (11) to (22), and evaluating in vitro whether or not the subject has breast cancer using both of the measured expression level(s) and a control expression level(s) in a sample from a healthy subject measured in the same way.

(56) (24) The method according to (23), wherein the subject is a human.

(57) (25) The method according to (23) or (24), wherein the sample is blood, serum, or plasma.

DEFINITION OF TERMS

(58) The terms used herein are defined as described below.

(59) Abbreviations or terms such as nucleotide, polynucleotide, DNA, and RNA used herein abide by “Guidelines for the preparation of specification which contain nucleotide and/or amino acid sequences” (edited by Japan Patent Office) and common use in the art.

(60) The term “polynucleotide” used herein refers to a nucleic acid including any of RNA, DNA, and RNA/DNA (chimera). The DNA includes any of cDNA, genomic DNA, and synthetic DNA. The RNA includes any of total RNA, mRNA, rRNA, miRNA, siRNA, snoRNA, snRNA, non-coding RNA and synthetic RNA. The “synthetic DNA” and the “synthetic RNA” used herein refer to DNA and RNA artificially prepared using, for example, an automatic nucleic acid synthesizer, on the basis of predetermined nucleotide sequences (which may be any of natural and non-natural sequences). The “non-natural sequence” used herein is intended to be used in a broad sense and includes, for example, a sequence comprising substitution, deletion, insertion, and/or addition of one or more nucleotide(s) (i.e., a variant sequence) and a sequence comprising one or more

modified nucleotide(s) (i.e., a modified sequence), which are different from the natural sequence. The term “polynucleotide” used herein is used interchangeably with the term “nucleic acid”. (61) The term “fragment” used herein refers to a polynucleotide (including oligonucleotides) having a nucleotide sequence that consists of a consecutive portion of a polynucleotide and desirably has a length of 15 or more nucleotides, preferably 17 or more nucleotides, more preferably 19 or more nucleotides.

(62) The term “gene” used herein is intended to include not only RNA and double-stranded DNA but also each single-stranded DNA such as a plus strand (or a sense strand) or a complementary strand (or an antisense strand) constituting the duplex. The gene is not particularly limited by its length. Thus, The “gene” used herein includes any of double-stranded DNA including human genomic DNA, single-stranded DNA (plus strand), single-stranded DNA having a sequence complementary to the plus strand (complementary strand) including cDNA, microRNA (miRNA), and their fragments, and their transcripts, unless otherwise specified. The “gene” includes not only a “gene” represented by a particular nucleotide sequence (or SEQ ID NO) but also “nucleic acids” encoding RNAs having biological functions equivalent to an RNA encoded by the gene, for example, a congener (i.e., a homolog or an ortholog), a variant (e.g., a genetic polymorph), and a derivative. Specific examples of such a “nucleic acid” encoding a congener, a variant, or a derivative can include a “nucleic acid” having a nucleotide sequence hybridizing under stringent conditions described later to a complementary sequence of a nucleotide sequence represented by any of SEQ ID NOs: 1 to 871 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t. Regardless whether or not there is a difference in functional region, the “gene(s)” can comprise, for example, expression regulatory region(s), coding region(s), exon(s), or intron(s). The “gene” may be contained in a cell or may exist alone after being released into the outside of a cell. Alternatively, the “gene” may be in a state enclosed in a vesicle called exosome.

(63) The term “exosome” used herein refers to a vesicle that is encapsulated by a lipid bilayer and secreted from a cell. The exosome is derived from a multivesicular endosome and may incorporate biomaterials such as “gene(s)” (e.g., RNA or DNA) or protein(s) when released into an extracellular environment. The exosome is known to be contained in a body fluid such as blood, serum, plasma, or lymph.

(64) The term “transcript” used herein refers to RNA synthesized with the DNA sequence of a gene as a template. RNA polymerase binds to a site called a promoter located upstream of the gene and adds ribonucleotides complementary to the nucleotide sequence of the DNA to the 3' end to synthesize an RNA. This RNA contains not only the gene itself but also the whole sequence from a transcription initiation site to the end of a polyA sequence, including expression regulatory region(s), coding region(s), exon(s), or intron(s).

(65) Unless otherwise specified, the term “microRNA (miRNA)” used herein is intended to mean a 15- to 25-nucleotide non-coding RNA that is transcribed as an RNA precursor which has a hairpin-like structure, cleaved by a dsRNA-cleaving enzyme which has RNase III cleavage activity, and integrated into a protein complex called RISC, and is involved in the suppression of translation of mRNA. The term “miRNA” used herein includes not only a “miRNA” represented by a particular nucleotide sequence (or SEQ ID NO) but also a precursor of the “miRNA” (pre-miRNA or pri-miRNA), and miRNAs that have biological functions equivalent thereto, for example, a congener (i.e., a homolog or an ortholog), a variant (e.g., a genetic polymorph), and a derivative. Such a precursor, a congener, a variant, or a derivative can be specifically identified using miRBase Release 20 (<http://www.mirbase.org/>), and examples thereof can include a “miRNA” having a nucleotide sequence hybridizing under stringent conditions described later to a complementary sequence of any particular nucleotide sequence represented by any of SEQ ID NOs: 1 to 871. The term “miRNA” used herein may be a gene product of a miR gene. Such a gene product includes a mature miRNA (e.g., a 15- to 25-nucleotide or 19- to 25-nucleotide non-coding RNA involved in the suppression of translation of mRNA as described above) or a miRNA precursor (e.g., pre-

miRNA or pri-miRNA as described above).

(66) The term “probe” used herein includes a polynucleotide that is used for specifically detecting an RNA resulting from the expression of a gene or a polynucleotide derived from the RNA, and/or a polynucleotide complementary thereto.

(67) The term “primer” used herein includes a polynucleotide that specifically recognizes and amplifies RNA resulting from the expression of a gene or a polynucleotide derived from the RNA, and/or a polynucleotide complementary thereto. In this context, the “complementary polynucleotide (complementary strand or reverse strand)” means a polynucleotide in a complementary relationship of A:T (U) and G:C base pairs with the full-length sequence of a polynucleotide consisting of a nucleotide sequence defined by any of SEQ ID NOs: 1 to 871 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, or a partial sequence thereof (here, this full-length or partial sequence is referred to as a plus strand for the sake of convenience). However, such a complementary strand is not limited to a sequence completely complementary to the nucleotide sequence of the target plus strand and may have a complementary relationship to an extent that permits hybridization under stringent conditions to the target plus strand.

(68) The term “stringent conditions” used herein refers to conditions under which a nucleic acid probe hybridizes to its target sequence to a larger extent (e.g., a measurement value equal to or larger than “(a mean of background measurement values)+(a standard deviation of the background measurement values) $\times$ 2”) than that for other sequences. The stringent conditions are dependent on a sequence and differ depending on an environment where hybridization is performed. A target sequence complementary 100% to the nucleic acid probe can be identified by controlling the stringency of hybridization and/or washing conditions. Specific examples of the “stringent conditions” will be mentioned later.

(69) The term “T_m value” used herein means a temperature at which the double-stranded moiety of a polynucleotide is denatured into single strands so that the double strands and the single strands exist at a ratio of 1:1.

(70) The term “variant” used herein means, in the case of a nucleic acid, a natural variant attributed to polymorphism, mutation, or the like; a variant containing the deletion, substitution, addition, or insertion of 1 or 2 or more nucleotides in a nucleotide sequence represented by any of SEQ ID NOs: 1 to 871 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, or a partial sequence thereof; a variant that exhibits percent (%) identity of approximately 90% or higher, approximately 95% or higher, approximately 97% or higher, approximately 98% or higher, approximately 99% or higher to each of these nucleotide sequences or the partial sequence thereof; or a nucleic acid hybridizing under the stringent conditions defined above to a polynucleotide or an oligonucleotide comprising each of these nucleotide sequences or the partial sequence thereof.

(71) The term “multiple” used herein means an integer of approximately 10, 9, 8, 7, 6, 5, 4, 3, or 2.

(72) The “variant” used herein can be prepared by use of a well-known technique such as site-directed mutagenesis or PCR-based mutagenesis.

(73) The term “percent (%) identity” used herein can be determined with or without an introduced gap, using a protein or gene search system based on BLAST or FASTA described above (Zheng Zhang et al., 2000, J. Comput. Biol., Vol. 7, p. 203-214; Altschul, S. F. et al., 1990, Journal of Molecular Biology, Vol. 215, p. 403-410; and Pearson, W. R. et al., 1988, Proc. Natl. Acad. Sci. U.S.A., Vol. 85, p. 2444-2448).

(74) The term “derivative” used herein is meant to include a modified nucleic acid, for example, a derivative labeled with a fluorophore or the like, a derivative containing a modified nucleotide (e.g., a nucleotide containing a group such as halogen, alkyl such as methyl, alkoxy such as methoxy, thio, or carboxymethyl, and a nucleotide that has undergone base rearrangement, double bond saturation, deamination, replacement of an oxygen molecule with a sulfur atom, etc.), PNA

(peptide nucleic acid; Nielsen, P. E. et al., 1991, Science, Vol. 254, p. 1497-500), and LNA (locked nucleic acid; Obika, S. et al., 1998, Tetrahedron Lett., Vol. 39, p. 5401-5404) without any limitation.

(75) The “nucleic acid” used herein capable of specifically binding to a polynucleotide selected from the breast cancer marker miRNAs described above is a synthesized or prepared nucleic acid and specifically includes a “nucleic acid probe” or a “primer”. The “nucleic acid” is utilized directly or indirectly for detecting the presence or absence of breast cancer in a subject, for diagnosing the presence or absence of breast cancer, or the severity of breast cancer, the presence or absence of amelioration of breast cancer, or the degree of amelioration of breast cancer, or the therapeutic sensitivity of breast cancer, or for screening for a candidate substance useful in the prevention, amelioration, or treatment of breast cancer. The “nucleic acid” includes a nucleotide, an oligonucleotide, and a polynucleotide capable of specifically recognizing and binding to a transcript represented by any of SEQ ID NOs:1 to 871 or a synthetic cDNA nucleic acid thereof in vivo, particularly, in a sample such as a body fluid (e.g., blood or urine), in relation to the development of breast cancer. The nucleotide, the oligonucleotide, and the polynucleotide can be effectively used as probes for detecting the aforementioned gene expressed in vivo, in tissues, in cells, or the like on the basis of the properties described above, or as primers for amplifying the aforementioned gene expressed in vivo.

(76) The term “detection” used herein is interchangeable with the term “examination”, “measurement”, or “detection or decision support”. The term “evaluation” used herein is meant to include diagnosing or evaluation-supporting on the basis of examination results or measurement results.

(77) The term “subject” used herein means a mammal such as a primate including a human and a chimpanzee, a pet animal including a dog and a cat, a livestock animal including cattle, a horse, sheep, and a goat, and a rodent including a mouse and a rat. The term “healthy subject” also means such a mammal without the cancer to be detected.

(78) The term “P” or “P value” used herein refers to a probability at which a more extreme statistic than that is actually calculated from data under a null hypothesis is observed in a statistical test. Thus, smaller “P” or “P value” is regarded as being a more significant difference between subjects to be compared.

(79) The term “sensitivity” used herein means a value of $(\text{the number of true positives})/(\text{the number of true positives} + \text{the number of false negatives})$. High sensitivity allows breast cancer to be detected early, leading to the complete resection of cancer sites and reduction in the rate of recurrence.

(80) The term “specificity” used herein means a value of $(\text{the number of true negatives})/(\text{the number of true negatives} + \text{the number of false positives})$. High specificity prevents needless extra examination for healthy subjects who might have been misjudged as being breast cancer patients, leading to reduction in burden on patients and reduction in medical expense.

(81) The term “accuracy” used herein means a value of $(\text{the number of true positives} + \text{the number of true negatives})/(\text{the total number of cases})$. The accuracy indicates the ratio of samples that are identified correctly to all samples, and serves as a primary index for evaluating detection performance.

(82) The “sample” used herein, that is subjected to determination, detection, or diagnosis refers to a tissue and a biological material in which the expression of the gene of the present invention varies as breast cancer develops, as breast cancer progresses, or as therapeutic effects on breast cancer are exerted. Specifically, the “sample” refers to a breast tissue, a perimammary vascular channel, lymph node, and organ, an organ suspected of having metastasis, the skin, a body fluid such as blood, urine, saliva, sweat, or tissue exudates, serum or plasma prepared from blood, feces, hair, and the like. The “sample” further refers to a biological sample extracted therefrom, specifically, a gene such as RNA or miRNA.

- (83) The term “hsa-miR-4783-3p gene” or “hsa-miR-4783-3p” used herein includes the hsa-miR-4783-3p gene (miRBase Accession No. MIMAT0019947) described in SEQ ID NO: 1, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4783-3p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4783” (miRBase Accession No. MI0017428, SEQ ID NO: 270) having a hairpin-like structure is known as a precursor of “hsa-miR-4783-3p”.
- (84) The term “hsa-miR-4730 gene” or “hsa-miR-4730” used herein includes the hsa-miR-4730 gene (miRBase Accession No. MIMAT0019852) described in SEQ ID NO: 2, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4730 gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4730” (miRBase Accession No. MI0017367, SEQ ID NO: 271) having a hairpin-like structure is known as a precursor of “hsa-miR-4730”.
- (85) The term “hsa-miR-1307-3p gene” or “hsa-miR-1307-3p” used herein includes the hsa-miR-1307-3p gene (miRBase Accession No. MIMAT0005951) described in SEQ ID NO: 3, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1307-3p gene can be obtained by a method described in Morin R D et al., 2008, Genome Res, Vol. 18, p. 610-621. Also, “hsa-mir-1307” (miRBase Accession No. MI0006444, SEQ ID NO: 272) having a hairpin-like structure is known as a precursor of “hsa-miR-1307-3p”.
- (86) The term “hsa-miR-4634 gene” or “hsa-miR-4634” used herein includes the hsa-miR-4634 gene (miRBase Accession No. MIMAT0019691) described in SEQ ID NO: 4, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4634 gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4634” (miRBase Accession No. MI0017261, SEQ ID NO: 273) having a hairpin-like structure is known as a precursor of “hsa-miR-4634”.
- (87) The term “hsa-miR-663a gene” or “hsa-miR-663a” used herein includes the hsa-miR-663a gene (miRBase Accession No. MIMAT0003326) described in SEQ ID NO: 5, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-663a gene can be obtained by a method described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-663a” (miRBase Accession No. MI0003672, SEQ ID NO: 274) having a hairpin-like structure is known as a precursor of “hsa-miR-663a”.
- (88) The term “hsa-miR-4532 gene” or “hsa-miR-4532” used herein includes the hsa-miR-4532 gene (miRBase Accession No. MIMAT0019071) described in SEQ ID NO: 6, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4532 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4532” (miRBase Accession No. MI0016899, SEQ ID NO: 275) having a hairpin-like structure is known as a precursor of “hsa-miR-4532”.
- (89) The term “hsa-miR-7704 gene” or “hsa-miR-7704” used herein includes the hsa-miR-7704 gene (miRBase Accession No. MIMAT0030019) described in SEQ ID NO: 7, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7704 gene can be obtained by a method described in Swaminathan S et al., 2013, Biochem Biophys Res Commun, Vol. 434, p. 228-234. Also, “hsa-mir-7704” (miRBase Accession No. MI0025240, SEQ ID NO: 276) having a hairpin-like structure is known as a precursor of “hsa-miR-7704”.
- (90) The term “hsa-miR-3178 gene” or “hsa-miR-3178” used herein includes the hsa-miR-3178 gene (miRBase Accession No. MIMAT0015055) described in SEQ ID NO: 8, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3178 gene can be obtained by a method described in Stark M S et al., 2010, PLoS One, Vol. 5, e9685. Also, “hsa-mir-3178” (miRBase Accession No. MI0014212, SEQ ID NO: 277) having a hairpin-like structure is known as a precursor of “hsa-miR-3178”.
- (91) The term “hsa-miR-6729-5p gene” or “hsa-miR-6729-5p” used herein includes the hsa-miR-6729-5p gene (miRBase Accession No. MIMAT0027359) described in SEQ ID NO: 9, a homolog

or an ortholog of a different organism species, and the like. The hsa-miR-6729-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6729” (miRBase Accession No. MI0022574, SEQ ID NO: 278) having a hairpin-like structure is known as a precursor of “hsa-miR-6729-5p”.

(92) The term “hsa-miR-6090 gene” or “hsa-miR-6090” used in herein includes the hsa-miR-6090 gene (miRBase Accession No. MIMAT0023715) described in SEQ ID NO: 10, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6090 gene can be obtained by a method described in Yoo J K et al., 2012, *Stem Cells Dev*, Vol. 21, p. 2049-2057. Also, “hsa-mir-6090” (miRBase Accession No. MI0020367, SEQ ID NO: 279) having a hairpin-like structure is known as a precursor of “hsa-miR-6090”.

(93) The term “hsa-miR-4732-5p gene” or “hsa-miR-4732-5p” used in herein includes the hsa-miR-4732-5p gene (miRBase Accession No. MIMAT0019855) described in SEQ ID NO: 11, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4732-5p gene can be obtained by a method described in Persson H et al., 2011, *Cancer Res*, Vol. 71, p. 78-86. Also, “hsa-mir-4732” (miRBase Accession No. MI0017369, SEQ ID NO: 280) having a hairpin-like structure is known as a precursor of “hsa-miR-4732-5p”.

(94) The term “hsa-miR-3184-5p gene” or “hsa-miR-3184-5p” used herein includes the hsa-miR-3184-5p gene (miRBase Accession No. MIMAT0015064) described in SEQ ID NO: 12, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3184-5p gene can be obtained by a method described in Stark M S et al., 2010, *PLoS One*, Vol. 5, e9685. Also, “hsa-mir-3184” (miRBase Accession No. MI0014226, SEQ ID NO: 281) having a hairpin-like structure is known as a precursor of “hsa-miR-3184-5p”.

(95) The term “hsa-miR-6727-5p gene” or “hsa-miR-6727-5p” used herein includes the hsa-miR-6727-5p gene (miRBase Accession No. MIMAT0027355) described in SEQ ID NO: 13, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6727-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6727” (miRBase Accession No. MI0022572, SEQ ID NO: 282) having a hairpin-like structure is known as a precursor of “hsa-miR-6727-5p”.

(96) The term “hsa-miR-6088 gene” or “hsa-miR-6088” used herein includes the hsa-miR-6088 gene (miRBase Accession No. MIMAT0023713) described in SEQ ID NO: 14, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6088 gene can be obtained by a method described in Yoo J K et al., 2012, *Stem Cells Dev*, Vol. 21, p. 2049-2057. Also, “hsa-mir-6088” (miRBase Accession No. MI0020365, SEQ ID NO: 283) having a hairpin-like structure is known as a precursor of “hsa-miR-6088”.

(97) The term “hsa-miR-4674 gene” or “hsa-miR-4674” used herein includes the hsa-miR-4674 gene (miRBase Accession No. MIMAT0019756) described in SEQ ID NO: 15, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4674 gene can be obtained by a method described in Persson H et al., 2011, *Cancer Res*, Vol. 71, p. 78-86. Also, “hsa-mir-4674” (miRBase Accession No. MI0017305, SEQ ID NO: 284) having a hairpin-like structure is known as a precursor of “hsa-miR-4674”.

(98) The term “hsa-miR-8073 gene” or “hsa-miR-8073” used herein includes the hsa-miR-8073 gene (miRBase Accession No. MIMAT0031000) described in SEQ ID NO: 16, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-8073 gene can be obtained by a method described in Wang H J et al., 2013, *Shock*, Vol. 39, p. 480-487. Also, “hsa-mir-8073” (miRBase Accession No. MI0025909, SEQ ID NO: 285) having a hairpin-like structure is known as a precursor of “hsa-miR-8073”.

(99) The term “hsa-miR-4787-5p gene” or “hsa-miR-4787-5p” used herein includes the hsa-miR-4787-5p gene (miRBase Accession No. MIMAT0019956) described in SEQ ID NO: 17, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4787-5p gene can be obtained by a method described in Persson H et al., 2011, *Cancer Res*, Vol. 71, p. 78-86. Also,

“hsa-mir-4787” (miRBase Accession No. MI0017434, SEQ ID NO: 286) having a hairpin-like structure is known as a precursor of “hsa-miR-4787-5p”.

(100) The term “hsa-miR-1469 gene” or “hsa-miR-1469” used herein includes the hsa-miR-1469 gene (miRBase Accession No. MIMAT0007347) described in SEQ ID NO: 18, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1469 gene can be obtained by a method described in Kawaji H et al., 2008, BMC Genomics, Vol. 9, p. 157. Also, “hsa-mir-1469” (miRBase Accession No. MI0007074, SEQ ID NO: 287) having a hairpin-like structure is known as a precursor of “hsa-miR-1469”.

(101) The term “hsa-miR-125a-3p gene” or “hsa-miR-125a-3p” used herein includes the hsa-miR-125a-3p gene (miRBase Accession No. MIMAT0004602) described in SEQ ID NO: 19, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-125a-3p gene can be obtained by a method described in Lagos-Quintana M et al., 2002, Curr Biol, Vol. 12, p. 735-739. Also, “hsa-mir-125a” (miRBase Accession No. MI0000469, SEQ ID NO: 288) having a hairpin-like structure is known as a precursor of “hsa-miR-125a-3p”.

(102) The term “hsa-miR-1233-5p gene” or “hsa-miR-1233-5p” used herein includes the hsa-miR-1233-5p gene (miRBase Accession No. MIMAT0022943) described in SEQ ID NO: 20, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1233-5p gene can be obtained by a method described in Berezikov E et al., 2007, Mol Cell, Vol. 28, p. 328-336. Also, “hsa-mir-1233-1 and hsa-mir-1233-2” (miRBase Accession Nos. MI0006323 and MI0015973, SEQ ID NOs: 289 and 290) having a hairpin-like structure are known as precursors of “hsa-miR-1233-5p”.

(103) The term “hsa-miR-885-3p gene” or “hsa-miR-885-3p” used herein includes the hsa-miR-885-3p gene (miRBase Accession No. MIMAT0004948) described in SEQ ID NO: 21, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-885-3p gene can be obtained by a method described in Berezikov E et al., 2006, Genome Res, Vol. 16, p. 1289-1298. Also, “hsa-mir-885” (miRBase Accession No. MI0005560, SEQ ID NO: 291) having a hairpin-like structure is known as a precursor of “hsa-miR-885-3p”.

(104) The term “hsa-miR-6802-5p gene” or “hsa-miR-6802-5p” used herein includes the hsa-miR-6802-5p gene (miRBase Accession No. MIMAT0027504) described in SEQ ID NO: 22, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6802-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6802” (miRBase Accession No. MI0022647, SEQ ID NO: 292) having a hairpin-like structure is known as a precursor of “hsa-miR-6802-5p”.

(105) The term “hsa-miR-328-5p gene” or “hsa-miR-328-5p” used herein includes the hsa-miR-328-5p gene (miRBase Accession No. MIMAT0026486) described in SEQ ID NO: 23, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-328-5p gene can be obtained by a method described in Kim J et al., 2004, Proc Natl Acad Sci USA, Vol. 101, p. 360-365. Also, “hsa-mir-328” (miRBase Accession No. MI0000804, SEQ ID NO: 293) having a hairpin-like structure is known as a precursor of “hsa-miR-328-5p”.

(106) The term “hsa-miR-6787-5p gene” or “hsa-miR-6787-5p” used herein includes the hsa-miR-6787-5p gene (miRBase Accession No. MIMAT0027474) described in SEQ ID NO: 24, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6787-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6787” (miRBase Accession No. MI0022632, SEQ ID NO: 294) having a hairpin-like structure is known as a precursor of “hsa-miR-6787-5p”.

(107) The term “hsa-miR-8069 gene” or “hsa-miR-8069” used herein includes the hsa-miR-8069 gene (miRBase Accession No. MIMAT0030996) described in SEQ ID NO: 25, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-8069 gene can be obtained by a method described in Wang H J et al., 2013, Shock, Vol. 39, p. 480-487. Also, “hsa-mir-8069” (miRBase Accession No. MI0025905, SEQ ID NO: 295) having a hairpin-like structure is known

as a precursor of “hsa-miR-8069”.

(108) The term “hsa-miR-6875-5p gene” or “hsa-miR-6875-5p” used herein includes the hsa-miR-6875-5p gene (miRBase Accession No. MIMAT0027650) described in SEQ ID NO: 26, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6875-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6875” (miRBase Accession No. MI0022722, SEQ ID NO: 296) having a hairpin-like structure is known as a precursor of “hsa-miR-6875-5p”.

(109) The term “hsa-miR-1246 gene” or “hsa-miR-1246” used herein includes the hsa-miR-1246 gene (miRBase Accession No. MIMAT0005898) described in SEQ ID NO: 27, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1246 gene can be obtained by a method described in Morin R D et al., 2008, *Genome Res*, Vol. 18, p. 610-621. Also, “hsa-mir-1246” (miRBase Accession No. MI0006381, SEQ ID NO: 297) having a hairpin-like structure is known as a precursor of “hsa-miR-1246”.

(110) The term “hsa-miR-4734 gene” or “hsa-miR-4734” used herein includes the hsa-miR-4734 gene (miRBase Accession No. MIMAT0019859) described in SEQ ID NO: 28, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4734 gene can be obtained by a method described in Persson H et al., 2011, *Cancer Res*, Vol. 71, p. 78-86. Also, “hsa-mir-4734” (miRBase Accession No. MI0017371, SEQ ID NO: 298) having a hairpin-like structure is known as a precursor of “hsa-miR-4734”.

(111) The term “hsa-miR-6757-5p gene” or “hsa-miR-6757-5p” used herein includes the hsa-miR-6757-5p gene (miRBase Accession No. MIMAT0027414) described in SEQ ID NO: 29, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6757-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6757” (miRBase Accession No. MI0022602, SEQ ID NO: 299) having a hairpin-like structure is known as a precursor of “hsa-miR-6757-5p”.

(112) The term “hsa-miR-6756-5p gene” or “hsa-miR-6756-5p” used herein includes the hsa-miR-6756-5p gene (miRBase Accession No. MIMAT0027412) described in SEQ ID NO: 30, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6756-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6756” (miRBase Accession No. MI0022601, SEQ ID NO: 300) having a hairpin-like structure is known as a precursor of “hsa-miR-6756-5p”.

(113) The term “hsa-miR-3665 gene” or “hsa-miR-3665” used herein includes the hsa-miR-3665 gene (miRBase Accession No. MIMAT0018087) described in SEQ ID NO: 31, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3665 gene can be obtained by a method described in Xie X et al., 2005, *Nature*, Vol. 434, p. 338-345. Also, “hsa-mir-3665” (miRBase Accession No. MI0016066, SEQ ID NO: 301) having a hairpin-like structure is known as a precursor of “hsa-miR-3665”.

(114) The term “hsa-miR-6836-3p gene” or “hsa-miR-6836-3p” used herein includes the hsa-miR-6836-3p gene (miRBase Accession No. MIMAT0027575) described in SEQ ID NO: 32, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6836-3p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6836” (miRBase Accession No. MI0022682, SEQ ID NO: 302) having a hairpin-like structure is known as a precursor of “hsa-miR-6836-3p”.

(115) The term “hsa-miR-6821-5p gene” or “hsa-miR-6821-5p” used herein includes the hsa-miR-6821-5p gene (miRBase Accession No. MIMAT0027542) described in SEQ ID NO: 33, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6821-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6821” (miRBase Accession No. MI0022666, SEQ ID NO: 303) having a hairpin-like structure is known as a precursor of “hsa-miR-6821-5p”.

(116) The term “hsa-miR-6805-5p gene” or “hsa-miR-6805-5p” used herein includes the hsa-miR-

6805-5p gene (miRBase Accession No. MIMAT0027510) described in SEQ ID NO: 34, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6805-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6805” (miRBase Accession No. MI0022650, SEQ ID NO: 304) having a hairpin-like structure is known as a precursor of “hsa-miR-6805-5p”.

(117) The term “hsa-miR-4728-5p gene” or “hsa-miR-4728-5p” used herein includes the hsa-miR-4728-5p gene (miRBase Accession No. MIMAT0019849) described in SEQ ID NO: 35, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4728-5p gene can be obtained by a method described in Persson H et al., 2011, *Cancer Res*, Vol. 71, p. 78-86. Also, “hsa-mir-4728” (miRBase Accession No. MI0017365, SEQ ID NO: 305) having a hairpin-like structure is known as a precursor of “hsa-miR-4728-5p”.

(118) The term “hsa-miR-6726-5p gene” or “hsa-miR-6726-5p” used herein includes the hsa-miR-6726-5p gene (miRBase Accession No. MIMAT0027353) described in SEQ ID NO: 36, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6726-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6726” (miRBase Accession No. MI0022571, SEQ ID NO: 306) having a hairpin-like structure is known as a precursor of “hsa-miR-6726-5p”.

(119) The term “hsa-miR-197-5p gene” or “hsa-miR-197-5p” used herein includes the hsa-miR-197-5p gene (miRBase Accession No. MIMAT0022691) described in SEQ ID NO: 37, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-197-5p gene can be obtained by a method described in Lagos-Quintana M et al., 2003, *RNA*, Vol. 9, p. 175-179. Also, “hsa-mir-197” (miRBase Accession No. MI0000239, SEQ ID NO: 307) having a hairpin-like structure is known as a precursor of “hsa-miR-197-5p”.

(120) The term “hsa-miR-149-3p gene” or “hsa-miR-149-3p” used herein includes the hsa-miR-149-3p gene (miRBase Accession No. MIMAT0004609) described in SEQ ID NO: 38, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-149-3p gene can be obtained by a method described in Lagos-Quintana M et al., 2002, *Curr Biol*, Vol. 12, p. 735-739. Also, “hsa-mir-149” (miRBase Accession No. MI0000478, SEQ ID NO: 308) having a hairpin-like structure is known as a precursor of “hsa-miR-149-3p”.

(121) The term “hsa-miR-6850-5p gene” or “hsa-miR-6850-5p” used herein includes the hsa-miR-6850-5p gene (miRBase Accession No. MIMAT0027600) described in SEQ ID NO: 39, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6850-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6850” (miRBase Accession No. MI0022696, SEQ ID NO: 309) having a hairpin-like structure is known as a precursor of “hsa-miR-6850-5p”.

(122) The term “hsa-miR-4476 gene” or “hsa-miR-4476” used herein includes the hsa-miR-4476 gene (miRBase Accession No. MIMAT0019003) described in SEQ ID NO: 40, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4476 gene can be obtained by a method described in Lima D D et al., 2010, *Blood*, Vol. 116, e118-e127. Also, “hsa-mir-4476” (miRBase Accession No. MI0016828, SEQ ID NO: 310) having a hairpin-like structure is known as a precursor of “hsa-miR-4476”.

(123) The term “hsa-miR-6858-5p gene” or “hsa-miR-6858-5p” used herein includes the hsa-miR-6858-5p gene (miRBase Accession No. MIMAT0027616) described in SEQ ID NO: 41, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6858-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6858” (miRBase Accession No. MI0022704, SEQ ID NO: 311) having a hairpin-like structure is known as a precursor of “hsa-miR-6858-5p”.

(124) The term “hsa-miR-564 gene” or “hsa-miR-564” used herein includes the hsa-miR-564 gene (miRBase Accession No. MIMAT0003228) described in SEQ ID NO: 42, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-564 gene can be obtained by a method

described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-564” (miRBase Accession No. MI0003570, SEQ ID NO: 312) having a hairpin-like structure is known as a precursor of “hsa-miR-564”.

(125) The term “hsa-miR-4763-3p gene” or “hsa-miR-4763-3p” used herein includes the hsa-miR-4763-3p gene (miRBase Accession No. MIMAT0019913) described in SEQ ID NO: 43, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4763-3p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4763” (miRBase Accession No. MI0017404, SEQ ID NO: 313) having a hairpin-like structure is known as a precursor of “hsa-miR-4763-3p”.

(126) The term “hsa-miR-575 gene” or “hsa-miR-575” used herein includes the hsa-miR-575 gene (miRBase Accession No. MIMAT0003240) described in SEQ ID NO: 44, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-575 gene can be obtained by a method described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-575” (miRBase Accession No. MI0003582, SEQ ID NO: 314) having a hairpin-like structure is known as a precursor of “hsa-miR-575”.

(127) The term “hsa-miR-6771-5p gene” or “hsa-miR-6771-5p” used herein includes the hsa-miR-6771-5p gene (miRBase Accession No. MIMAT0027442) described in SEQ ID NO: 45, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6771-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6771” (miRBase Accession No. MI0022616, SEQ ID NO: 315) having a hairpin-like structure is known as a precursor of “hsa-miR-6771-5p”.

(128) The term “hsa-miR-1231 gene” or “hsa-miR-1231” used herein includes the hsa-miR-1231 gene (miRBase Accession No. MIMAT0005586) described in SEQ ID NO: 46, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1231 gene can be obtained by a method described in Berezikov E et al., 2007, Mol Cell, Vol. 28, p. 328-336. Also, “hsa-mir-1231” (miRBase Accession No. MI0006321, SEQ ID NO: 316) having a hairpin-like structure is known as a precursor of “hsa-miR-1231”.

(129) The term “hsa-miR-1908-3p gene” or “hsa-miR-1908-3p” used herein includes the hsa-miR-1908-3p gene (miRBase Accession No. MIMAT0026916) described in SEQ ID NO: 47, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1908-3p gene can be obtained by a method described in Bar M et al., 2008, Stem Cells, Vol. 26, p. 2496-2505. Also, “hsa-mir-1908” (miRBase Accession No. MI0008329, SEQ ID NO: 317) having a hairpin-like structure is known as a precursor of “hsa-miR-1908-3p”.

(130) The term “hsa-miR-150-3p gene” or “hsa-miR-150-3p” used herein includes the hsa-miR-150-3p gene (miRBase Accession No. MIMAT0004610) described in SEQ ID NO: 48, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-150-3p gene can be obtained by a method described in Lagos-Quintana M et al., 2002, Curr Biol, Vol. 12, p. 735-739. Also, “hsa-mir-150” (miRBase Accession No. MI0000479, SEQ ID NO: 318) having a hairpin-like structure is known as a precursor of “hsa-miR-150-3p”.

(131) The term “hsa-miR-3937 gene” or “hsa-miR-3937” used herein includes the hsa-miR-3937 gene (miRBase Accession No. MIMAT0018352) described in SEQ ID NO: 49, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3937 gene can be obtained by a method described in Liao J Y et al., 2010, PLoS One, Vol. 5, e10563. Also, “hsa-mir-3937” (miRBase Accession No. MI0016593, SEQ ID NO: 319) having a hairpin-like structure is known as a precursor of “hsa-miR-3937”.

(132) The term “hsa-miR-887-3p gene” or “hsa-miR-887-3p” used herein includes the hsa-miR-887-3p gene (miRBase Accession No. MIMAT0004951) described in SEQ ID NO: 50, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-887-3p gene can be obtained by a method described in Berezikov E et al., 2006, Genome Res, Vol. 16, p. 1289-1298. Also, “hsa-mir-887” (miRBase Accession No. MI0005562, SEQ ID NO: 320) having a hairpin-like

structure is known as a precursor of “hsa-miR-887-3p”.

(133) The term “hsa-miR-3940-5p gene” or “hsa-miR-3940-5p” used herein includes the hsa-miR-3940-5p gene (miRBase Accession No. MIMAT0019229) described in SEQ ID NO: 51, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3940-5p gene can be obtained by a method described in Liao J Y et al., 2010, PLoS One, Vol. 5, e10563. Also, “hsa-mir-3940” (miRBase Accession No. MI0016597, SEQ ID NO: 321) having a hairpin-like structure is known as a precursor of “hsa-miR-3940-5p”.

(134) The term “hsa-miR-4741 gene” or “hsa-miR-4741” used herein includes the hsa-miR-4741 gene (miRBase Accession No. MIMAT0019871) described in SEQ ID NO: 52, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4741 gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4741” (miRBase Accession No. MI0017379, SEQ ID NO: 322) having a hairpin-like structure is known as a precursor of “hsa-miR-4741”.

(135) The term “hsa-miR-6808-5p gene” or “hsa-miR-6808-5p” used herein includes the hsa-miR-6808-5p gene (miRBase Accession No. MIMAT0027516) described in SEQ ID NO: 53, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6808-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6808” (miRBase Accession No. MI0022653, SEQ ID NO: 323) having a hairpin-like structure is known as a precursor of “hsa-miR-6808-5p”.

(136) The term “hsa-miR-6869-5p gene” or “hsa-miR-6869-5p” used herein includes the hsa-miR-6869-5p gene (miRBase Accession No. MIMAT0027638) described in SEQ ID NO: 54, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6869-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6869” (miRBase Accession No. MI0022716, SEQ ID NO: 324) having a hairpin-like structure is known as a precursor of “hsa-miR-6869-5p”.

(137) The term “hsa-miR-5090 gene” or “hsa-miR-5090” used herein includes the hsa-miR-5090 gene (miRBase Accession No. MIMAT0021082) described in SEQ ID NO: 55, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-5090 gene can be obtained by a method described in Ding N et al., 2011, J Radiat Res, Vol. 52, p. 425-432. Also, “hsa-mir-5090” (miRBase Accession No. MI0017979, SEQ ID NO: 325) having a hairpin-like structure is known as a precursor of “hsa-miR-5090”.

(138) The term “hsa-miR-615-5p gene” or “hsa-miR-615-5p” used herein includes the hsa-miR-615-5p gene (miRBase Accession No. MIMAT0004804) described in SEQ ID NO: 56, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-615-5p gene can be obtained by a method described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-615” (miRBase Accession No. MI0003628, SEQ ID NO: 326) having a hairpin-like structure is known as a precursor of “hsa-miR-615-5p”.

(139) The term “hsa-miR-8072 gene” or “hsa-miR-8072” used herein includes the hsa-miR-8072 gene (miRBase Accession No. MIMAT0030999) described in SEQ ID NO: 57, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-8072 gene can be obtained by a method described in Wang H J et al., 2013, Shock, Vol. 39, p. 480-487. Also, “hsa-mir-8072” (miRBase Accession No. MI0025908, SEQ ID NO: 327) having a hairpin-like structure is known as a precursor of “hsa-miR-8072”.

(140) The term “hsa-miR-128-1-5p gene” or “hsa-miR-128-1-5p” used herein includes the hsa-miR-128-1-5p gene (miRBase Accession No. MIMAT0026477) described in SEQ ID NO: 58, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-128-1-5p gene can be obtained by a method described in Lagos-Quintana M et al., 2002, Curr Biol, Vol. 12, p. 735-739. Also, “hsa-mir-128-1” (miRBase Accession No. MI0000447, SEQ ID NO: 328) having a hairpin-like structure is known as a precursor of “hsa-miR-128-1-5p”.

(141) The term “hsa-miR-1238-5p gene” or “hsa-miR-1238-5p” used herein includes the hsa-miR-

1238-5p gene (miRBase Accession No. MIMAT0022947) described in SEQ ID NO: 59, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1238-5p gene can be obtained by a method described in Berezikov E et al., 2007, Mol Cell, Vol. 28, p. 328-336. Also, “hsa-mir-1238” (miRBase Accession No. MI0006328, SEQ ID NO: 329) having a hairpin-like structure is known as a precursor of “hsa-miR-1238-5p”.

(142) The term “hsa-miR-365a-5p gene” or “hsa-miR-365a-5p” used herein includes the hsa-miR-365a-5p gene (miRBase Accession No. MIMAT0009199) described in SEQ ID NO: 60, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-365a-5p gene can be obtained by a method described in Xie X et al., 2005, Nature, Vol. 434, p. 338-345. Also, “hsa-mir-365a” (miRBase Accession No. MI0000767, SEQ ID NO: 330) having a hairpin-like structure is known as a precursor of “hsa-miR-365a-5p”.

(143) The term “hsa-miR-204-3p gene” or “hsa-miR-204-3p” used herein includes the hsa-miR-204-3p gene (miRBase Accession No. MIMAT0022693) described in SEQ ID NO: 61, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-204-3p gene can be obtained by a method described in Lim L P et al., 2003, Science, Vol. 299, p. 1540. Also, “hsa-mir-204” (miRBase Accession No. MI0000284, SEQ ID NO: 331) having a hairpin-like structure is known as a precursor of “hsa-miR-204-3p”.

(144) The term “hsa-miR-4492 gene” or “hsa-miR-4492” used herein includes the hsa-miR-4492 gene (miRBase Accession No. MIMAT0019027) described in SEQ ID NO: 62, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4492 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4492” (miRBase Accession No. MI0016854, SEQ ID NO: 332) having a hairpin-like structure is known as a precursor of “hsa-miR-4492”.

(145) The term “hsa-miR-6785-5p gene” or “hsa-miR-6785-5p” used herein includes the hsa-miR-6785-5p gene (miRBase Accession No. MIMAT0027470) described in SEQ ID NO: 63, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6785-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6785” (miRBase Accession No. MI0022630, SEQ ID NO: 333) having a hairpin-like structure is known as a precursor of “hsa-miR-6785-5p”.

(146) The term “hsa-miR-6511a-5p gene” or “hsa-miR-6511a-5p” used herein includes the hsa-miR-6511a-5p gene (miRBase Accession No. MIMAT0025478) described in SEQ ID NO: 64, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6511a-5p gene can be obtained by a method described in Joyce C E et al., 2011, Hum Mol Genet, Vol. 20, p. 4025-4040. Also, “hsa-mir-6511a-1, hsa-mir-6511a-2, hsa-mir-6511a-3, and hsa-mir-6511a-4” (miRBase Accession Nos. MI0022223, MI0023564, MI0023565, and MI0023566, SEQ ID NOs: 334, 335, 336, and 337) having a hairpin-like structure are known as precursors of “hsa-miR-6511a-5p”.

(147) The term “hsa-miR-4525 gene” or “hsa-miR-4525” used herein includes the hsa-miR-4525 gene (miRBase Accession No. MIMAT0019064) described in SEQ ID NO: 65, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4525 gene can be obtained by a method described in Lima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4525” (miRBase Accession No. MI0016892, SEQ ID NO: 338) having a hairpin-like structure is known as a precursor of “hsa-miR-4525”.

(148) The term “hsa-miR-1915-5p gene” or “hsa-miR-1915-5p” used herein includes the hsa-miR-1915-5p gene (miRBase Accession No. MIMAT0007891) described in SEQ ID NO: 66, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1915-5p gene can be obtained by a method described in Bar M et al., 2008, Stem Cells, Vol. 26, p. 2496-2505. Also, “hsa-mir-1915” (miRBase Accession No. MI0008336, SEQ ID NO: 339) having a hairpin-like structure is known as a precursor of “hsa-miR-1915-5p”.

(149) The term “hsa-miR-3180 gene” or “hsa-miR-3180” used herein includes the hsa-miR-3180 gene (miRBase Accession No. MIMAT0018178) described in SEQ ID NO: 67, a homolog or an

ortholog of a different organism species, and the like. The hsa-miR-3180 gene can be obtained by a method described in Creighton Cl et al., 2010, PLoS One, Vol. 5, e9637. Also, “hsa-mir-3180-4 and hsa-mir-3180-5” (miRBase Accession Nos. MI0016408 and MI0016409, SEQ ID NOs: 340 and 341) having a hairpin-like structure are known as precursors of “hsa-miR-3180”.

(150) The term “hsa-miR-6879-5p gene” or “hsa-miR-6879-5p” used herein includes the hsa-miR-6879-5p gene (miRBase Accession No. MIMAT0027658) described in SEQ ID NO: 68, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6879-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6879” (miRBase Accession No. MI0022726, SEQ ID NO: 342) having a hairpin-like structure is known as a precursor of “hsa-miR-6879-5p”.

(151) The term “hsa-miR-1199-5p gene” or “hsa-miR-1199-5p” used herein includes the hsa-miR-1199-5p gene (miRBase Accession No. MIMAT0031119) described in SEQ ID NO: 69, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1199-5p gene can be obtained by a method described in Salvi A et al., 2013, Int J Oncol, Vol. 42, p. 391-402. Also, “hsa-mir-1199” (miRBase Accession No. MI0020340, SEQ ID NO: 343) having a hairpin-like structure is known as a precursor of “hsa-miR-1199-5p”.

(152) The term “hsa-miR-6746-5p gene” or “hsa-miR-6746-5p” used herein includes the hsa-miR-6746-5p gene (miRBase Accession No. MIMAT0027392) described in SEQ ID NO: 70, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6746-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6746” (miRBase Accession No. MI0022591, SEQ ID NO: 344) having a hairpin-like structure is known as a precursor of “hsa-miR-6746-5p”.

(153) The term “hsa-miR-711 gene” or “hsa-miR-711” used herein includes the hsa-miR-711 gene (miRBase Accession No. MIMAT0012734) described in SEQ ID NO: 71, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-711 gene can be obtained by a method described in Artzi S et al., 2008, BMC Bioinformatics, Vol. 9, p. 39. Also, “hsa-mir-711” (miRBase Accession No. MI0012488, SEQ ID NO: 345) having a hairpin-like structure is known as a precursor of “hsa-miR-711”.

(154) The term “hsa-miR-663b gene” or “hsa-miR-663b” used herein includes the hsa-miR-663b gene (miRBase Accession No. MIMAT0005867) described in SEQ ID NO: 72, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-663b gene can be obtained by a method described in Takada S et al., 2008, Leukemia, Vol. 22, p. 1274-1278. Also, “hsa-mir-663b” (miRBase Accession No. MI0006336, SEQ ID NO: 346) having a hairpin-like structure is known as a precursor of “hsa-miR-663b”.

(155) The term “hsa-miR-4707-3p gene” or “hsa-miR-4707-3p” used herein includes the hsa-miR-4707-3p gene (miRBase Accession No. MIMAT0019808) described in SEQ ID NO: 73, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4707-3p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4707” (miRBase Accession No. MI0017340, SEQ ID NO: 347) having a hairpin-like structure is known as a precursor of “hsa-miR-4707-3p”.

(156) The term “hsa-miR-6893-5p gene” or “hsa-miR-6893-5p” used herein includes the hsa-miR-6893-5p gene (miRBase Accession No. MIMAT0027686) described in SEQ ID NO: 74, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6893-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6893” (miRBase Accession No. MI0022740, SEQ ID NO: 348) having a hairpin-like structure is known as a precursor of “hsa-miR-6893-5p”.

(157) The term “hsa-miR-4675 gene” or “hsa-miR-4675” used herein includes the hsa-miR-4675 gene (miRBase Accession No. MIMAT0019757) described in SEQ ID NO: 75, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4675 gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4675”

(miRBase Accession No. MI0017306, SEQ ID NO: 349) having a hairpin-like structure is known as a precursor of “hsa-miR-4675”.

(158) The term “hsa-miR-4638-5p gene” or “hsa-miR-4638-5p” used herein includes the hsa-miR-4638-5p gene (miRBase Accession No. MIMAT0019695) described in SEQ ID NO: 76, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4638-5p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4638” (miRBase Accession No. MI0017265, SEQ ID NO: 350) having a hairpin-like structure is known as a precursor of “hsa-miR-4638-5p”.

(159) The term “hsa-miR-4651 gene” or “hsa-miR-4651” used herein includes the hsa-miR-4651 gene (miRBase Accession No. MIMAT0019715) described in SEQ ID NO: 77, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4651 gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4651” (miRBase Accession No. MI0017279, SEQ ID NO: 351) having a hairpin-like structure is known as a precursor of “hsa-miR-4651”.

(160) The term “hsa-miR-6087 gene” or “hsa-miR-6087” used herein includes the hsa-miR-6087 gene (miRBase Accession No. MIMAT0023712) described in SEQ ID NO: 78, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6087 gene can be obtained by a method described in Yoo J K et al., 2012, Stem Cells Dev, Vol. 21, p. 2049-2057. Also, “hsa-mir-6087” (miRBase Accession No. MI0020364, SEQ ID NO: 352) having a hairpin-like structure is known as a precursor of “hsa-miR-6087”.

(161) The term “hsa-miR-4665-5p gene” or “hsa-miR-4665-5p” used herein includes the hsa-miR-4665-5p gene (miRBase Accession No. MIMAT0019739) described in SEQ ID NO: 79, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4665-5p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4665” (miRBase Accession No. MI0017295, SEQ ID NO: 353) having a hairpin-like structure is known as a precursor of “hsa-miR-4665-5p”.

(162) The term “hsa-miR-4758-5p gene” or “hsa-miR-4758-5p” used herein includes the hsa-miR-4758-5p gene (miRBase Accession No. MIMAT0019903) described in SEQ ID NO: 80, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4758-5p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4758” (miRBase Accession No. MI0017399, SEQ ID NO: 354) having a hairpin-like structure is known as a precursor of “hsa-miR-4758-5p”.

(163) The term “hsa-miR-6887-5p gene” or “hsa-miR-6887-5p” used herein includes the hsa-miR-6887-5p gene (miRBase Accession No. MIMAT0027674) described in SEQ ID NO: 81, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6887-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6887” (miRBase Accession No. MI0022734, SEQ ID NO: 355) having a hairpin-like structure is known as a precursor of “hsa-miR-6887-5p”.

(164) The term “hsa-miR-3620-5p gene” or “hsa-miR-3620-5p” used herein includes the hsa-miR-3620-5p gene (miRBase Accession No. MIMAT0022967) described in SEQ ID NO: 82, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3620-5p gene can be obtained by a method described in Witten D et al., 2010, BMC Biol, Vol. 8, p. 58. Also, “hsa-mir-3620” (miRBase Accession No. MI0016011, SEQ ID NO: 356) having a hairpin-like structure is known as a precursor of “hsa-miR-3620-5p”.

(165) The term “hsa-miR-1909-3p gene” or “hsa-miR-1909-3p” used herein includes the hsa-miR-1909-3p gene (miRBase Accession No. MIMAT0007883) described in SEQ ID NO: 83, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1909-3p gene can be obtained by a method described in Bar M et al., 2008, Stem Cells, Vol. 26, p. 2496-2505. Also, “hsa-mir-1909” (miRBase Accession No. MI0008330, SEQ ID NO: 357) having a hairpin-like structure is known as a precursor of “hsa-miR-1909-3p”.

(166) The term “hsa-miR-7641 gene” or “hsa-miR-7641” used herein includes the hsa-miR-7641 gene (miRBase Accession No. MIMAT0029782) described in SEQ ID NO: 84, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7641 gene can be obtained by a method described in Yoo J K et al., 2013, Arch Pharm Res, Vol. 36, p. 353-358. Also, “hsa-mir-7641-1 and hsa-mir-7641-2” (miRBase Accession Nos. MI0024975 and MI0024976, SEQ ID NOs: 358 and 359) having a hairpin-like structure are known as precursors of “hsa-miR-7641”.

(167) The term “hsa-miR-6724-5p gene” or “hsa-miR-6724-5p” used herein includes the hsa-miR-6724-5p gene (miRBase Accession No. MIMAT0025856) described in SEQ ID NO: 85, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6724-5p gene can be obtained by a method described in Li Y et al., 2012, Gene, Vol. 497, p. 330-335. Also, “hsa-mir-6724” (miRBase Accession No. MI0022559, SEQ ID NO: 360) having a hairpin-like structure is known as a precursor of “hsa-miR-6724-5p”.

(168) The term “hsa-miR-1343-3p gene” or “hsa-miR-1343-3p” used herein includes the hsa-miR-1343-3p gene (miRBase Accession No. MIMAT0019776) described in SEQ ID NO: 86, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1343-3p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-1343” (miRBase Accession No. MI0017320, SEQ ID NO: 361) having a hairpin-like structure is known as a precursor of “hsa-miR-1343-3p”.

(169) The term “hsa-miR-6780b-5p gene” or “hsa-miR-6780b-5p” used herein includes the hsa-miR-6780b-5p gene (miRBase Accession No. MIMAT0027572) described in SEQ ID NO: 87, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6780b-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6780b” (miRBase Accession No. MI0022681, SEQ ID NO: 362) having a hairpin-like structure is known as a precursor of “hsa-miR-6780b-5p”.

(170) The term “hsa-miR-4484 gene” or “hsa-miR-4484” used herein includes the hsa-miR-4484 gene (miRBase Accession No. MIMAT0019018) described in SEQ ID NO: 88, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4484 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4484” (miRBase Accession No. MI0016845, SEQ ID NO: 363) having a hairpin-like structure is known as a precursor of “hsa-miR-4484”.

(171) The term “hsa-miR-4690-5p gene” or “hsa-miR-4690-5p” used herein includes the hsa-miR-4690-5p gene (miRBase Accession No. MIMAT0019779) described in SEQ ID NO: 89, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4690-5p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4690” (miRBase Accession No. MI0017323, SEQ ID NO: 364) having a hairpin-like structure is known as a precursor of “hsa-miR-4690-5p”.

(172) The term “hsa-miR4429 gene” or “hsa-miR-4429” used herein includes the hsa-miR-4429 gene (miRBase Accession No. MIMAT0018944) described in SEQ ID NO: 90, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4429 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4429” (miRBase Accession No. MI0016768, SEQ ID NO: 365) having a hairpin-like structure is known as a precursor of “hsa-miR4429”.

(173) The term “hsa-miR-1227-5p gene” or “hsa-miR-1227-5p” used herein includes the hsa-miR-1227-5p gene (miRBase Accession No. MIMAT0022941) described in SEQ ID NO: 91, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1227-5p gene can be obtained by a method described in Berezikov E et al., 2007, Mol Cell, Vol. 28, p. 328-336. Also, “hsa-mir-1227” (miRBase Accession No. MI0006316, SEQ ID NO: 366) having a hairpin-like structure is known as a precursor of “hsa-miR-1227-5p”.

(174) The term “hsa-miR-4725-3p gene” or “hsa-miR-4725-3p” used herein includes the hsa-miR-4725-3p gene (miRBase Accession No. MIMAT0019844) described in SEQ ID NO: 92, a homolog

or an ortholog of a different organism species, and the like. The hsa-miR-4725-3p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4725” (miRBase Accession No. MI0017362, SEQ ID NO: 367) having a hairpin-like structure is known as a precursor of “hsa-miR-4725-3p”.

(175) The term “hsa-miR-6861-5p gene” or “hsa-miR-6861-5p” used herein includes the hsa-miR-6861-5p gene (miRBase Accession No. MIMAT0027623) described in SEQ ID NO: 93, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6861-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6861” (miRBase Accession No. MI0022708, SEQ ID NO: 368) having a hairpin-like structure is known as a precursor of “hsa-miR-6861-5p”.

(176) The term “hsa-miR-6812-5p gene” or “hsa-miR-6812-5p” used herein includes the hsa-miR-6812-5p gene (miRBase Accession No. MIMAT0027524) described in SEQ ID NO: 94, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6812-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6812” (miRBase Accession No. MI0022657, SEQ ID NO: 369) having a hairpin-like structure is known as a precursor of “hsa-miR-6812-5p”.

(177) The term “hsa-miR-3197 gene” or “hsa-miR-3197” used herein includes the hsa-miR-3197 gene (miRBase Accession No. MIMAT0015082) described in SEQ ID NO: 95, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3197 gene can be obtained by a method described in Stark M S et al., 2010, PLoS One, Vol. 5, e9685. Also, “hsa-mir-3197” (miRBase Accession No. MI0014245, SEQ ID NO: 370) having a hairpin-like structure is known as a precursor of “hsa-miR-3197”.

(178) The term “hsa-miR-8059 gene” or “hsa-miR-8059” used herein includes the hsa-miR-8059 gene (miRBase Accession No. MIMAT0030986) described in SEQ ID NO: 96, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-8059 gene can be obtained by a method described in Wang H J et al., 2013, Shock, Vol. 39, p. 480-487. Also, “hsa-mir-8059” (miRBase Accession No. MI0025895, SEQ ID NO: 371) having a hairpin-like structure is known as a precursor of “hsa-miR-8059”.

(179) The term “hsa-miR-3185 gene” or “hsa-miR-3185” used herein includes the hsa-miR-3185 gene (miRBase Accession No. MIMAT0015065) described in SEQ ID NO: 97, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3185 gene can be obtained by a method described in Stark M S et al., 2010, PLoS One, Vol. 5, e9685. Also, “hsa-mir-3185” (miRBase Accession No. MI0014227, SEQ ID NO: 372) having a hairpin-like structure is known as a precursor of “hsa-miR-3185”.

(180) The term “hsa-miR-4706 gene” or “hsa-miR-4706” used herein includes the hsa-miR-4706 gene (miRBase Accession No. MIMAT0019806) described in SEQ ID NO: 98, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4706 gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4706” (miRBase Accession No. MI0017339, SEQ ID NO: 373) having a hairpin-like structure is known as a precursor of “hsa-miR-4706”.

(181) The term “hsa-miR-4497 gene” or “hsa-miR-4497” used herein includes the hsa-miR-4497 gene (miRBase Accession No. MIMAT0019032) described in SEQ ID NO: 99, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4497 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4497” (miRBase Accession No. MI0016859, SEQ ID NO: 374) having a hairpin-like structure is known as a precursor of “hsa-miR-4497”.

(182) The term “hsa-miR-3131 gene” or “hsa-miR-3131” used herein includes the hsa-miR-3131 gene (miRBase Accession No. MIMAT0014996) described in SEQ ID NO: 100, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3131 gene can be obtained by a method described in Stark M S et al., 2010, PLoS One, Vol. 5, e9685. Also, “hsa-mir-3131”

(miRBase Accession No. MI0014151, SEQ ID NO: 375) having a hairpin-like structure is known as a precursor of “hsa-miR-3131”.

(183) The term “hsa-miR-6806-5p gene” or “hsa-miR-6806-5p” used herein includes the hsa-miR-6806-5p gene (miRBase Accession No. MIMAT0027512) described in SEQ ID NO: 101, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6806-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6806” (miRBase Accession No. MI0022651, SEQ ID NO: 376) having a hairpin-like structure is known as a precursor of “hsa-miR-6806-5p”.

(184) The term “hsa-miR-187-5p gene” or “hsa-miR-187-5p” used herein includes the hsa-miR-187-5p gene (miRBase Accession No. MIMAT0004561) described in SEQ ID NO: 102, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-187-5p gene can be obtained by a method described in Lim L P et al., 2003, Science, Vol. 299, p. 1540. Also, “hsa-mir-187” (miRBase Accession No. MI0000274, SEQ ID NO: 377) having a hairpin-like structure is known as a precursor of “hsa-miR-187-5p”.

(185) The term “hsa-miR-3180-3p gene” or “hsa-miR-3180-3p” used herein includes the hsa-miR-3180-3p gene (miRBase Accession No. MIMAT0015058) described in SEQ ID NO: 103, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3180-3p gene can be obtained by a method described in Creighton C J et al., 2010, PLoS One, Vol. 5, e9637. Also, “hsa-mir-3180-1, hsa-mir-3180-2, and hsa-mir-3180-3” (miRBase Accession Nos. MI0014214, MI0014215, and MI0014217, SEQ ID NOs: 378, 379, and 380) having a hairpin-like structure are known as precursors of “hsa-miR-3180-3p”.

(186) The term “hsa-miR-6848-5p gene” or “hsa-miR-6848-5p” used herein includes the hsa-miR-6848-5p gene (miRBase Accession No. MIMAT0027596) described in SEQ ID NO: 104, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6848-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6848” (miRBase Accession No. MI0022694, SEQ ID NO: 381) having a hairpin-like structure is known as a precursor of “hsa-miR-6848-5p”.

(187) The term “hsa-miR-6820-5p gene” or “hsa-miR-6820-5p” used herein includes the hsa-miR-6820-5p gene (miRBase Accession No. MIMAT0027540) described in SEQ ID NO: 105, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6820-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6820” (miRBase Accession No. MI0022665, SEQ ID NO: 382) having a hairpin-like structure is known as a precursor of “hsa-miR-6820-5p”.

(188) The term “hsa-miR-6800-5p gene” or “hsa-miR-6800-5p” used herein includes the hsa-miR-6800-5p gene (miRBase Accession No. MIMAT0027500) described in SEQ ID NO: 106, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6800-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6800” (miRBase Accession No. MI0022645, SEQ ID NO: 383) having a hairpin-like structure is known as a precursor of “hsa-miR-6800-5p”.

(189) The term “hsa-miR-6717-5p gene” or “hsa-miR-6717-5p” used herein includes the hsa-miR-6717-5p gene (miRBase Accession No. MIMAT0025846) described in SEQ ID NO: 107, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6717-5p gene can be obtained by a method described in Li Y et al., 2012, Gene, Vol. 497, p. 330-335. Also, “hsa-mir-6717” (miRBase Accession No. MI0022551, SEQ ID NO: 384) having a hairpin-like structure is known as a precursor of “hsa-miR-6717-5p”.

(190) The term “hsa-miR-6795-5p gene” or “hsa-miR-6795-5p” used herein includes the hsa-miR-6795-5p gene (miRBase Accession No. MIMAT0027490) described in SEQ ID NO: 108, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6795-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6795” (miRBase Accession No. MI0022640, SEQ ID NO: 385) having a

hairpin-like structure is known as a precursor of “hsa-miR-6795-5p”.

(191) The term “hsa-miR-4632-5p gene” or “hsa-miR-4632-5p” used herein includes the hsa-miR-4632-5p gene (miRBase Accession No. MIMAT0022977) described in SEQ ID NO: 109, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4632-5p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4632” (miRBase Accession No. MI0017259, SEQ ID NO: 386) having a hairpin-like structure is known as a precursor of “hsa-miR-4632-5p”.

(192) The term “hsa-miR-665 gene” or “hsa-miR-665” used herein includes the hsa-miR-665 gene (miRBase Accession No. MIMAT0004952) described in SEQ ID NO: 110, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-665 gene can be obtained by a method described in Berezikov E et al., 2006, Genome Res, Vol. 16, p. 1289-1298. Also, “hsa-mir-665” (miRBase Accession No. MI0005563, SEQ ID NO: 387) having a hairpin-like structure is known as a precursor of “hsa-miR-665”.

(193) The term “hsa-miR-6778-5p gene” or “hsa-miR-6778-5p” used herein includes the hsa-miR-6778-5p gene (miRBase Accession No. MIMAT0027456) described in SEQ ID NO: 111, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6778-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6778” (miRBase Accession No. MI0022623, SEQ ID NO: 388) having a hairpin-like structure is known as a precursor of “hsa-miR-6778-5p”.

(194) The term “hsa-miR-3663-3p gene” or “hsa-miR-3663-3p” used herein includes the hsa-miR-3663-3p gene (miRBase Accession No. MIMAT0018085) described in SEQ ID NO: 112, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3663-3p gene can be obtained by a method described in Liao J Y et al., 2010, PLoS One, Vol. 5, e10563. Also, “hsa-mir-3663” (miRBase Accession No. MI0016064, SEQ ID NO: 389) having a hairpin-like structure is known as a precursor of “hsa-miR-3663-3p”.

(195) The term “hsa-miR-4689 gene” or “hsa-miR-4689” used herein includes the hsa-miR-4689 gene (miRBase Accession No. MIMAT0019778) described in SEQ ID NO: 113, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4689 gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4689” (miRBase Accession No. MI0017322, SEQ ID NO: 390) having a hairpin-like structure is known as a precursor of “hsa-miR-4689”.

(196) The term “hsa-miR-211-3p gene” or “hsa-miR-211-3p” used herein includes the hsa-miR-211-3p gene (miRBase Accession No. MIMAT0022694) described in SEQ ID NO: 114, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-211-3p gene can be obtained by a method described in Lim L P et al., 2003, Science, Vol. 299, p. 1540. Also, “hsa-mir-211” (miRBase Accession No. MI0000287, SEQ ID NO: 391) having a hairpin-like structure is known as a precursor of “hsa-miR-211-3p”.

(197) The term “hsa-miR-6511b-5p gene” or “hsa-miR-6511b-5p” used herein includes the hsa-miR-6511b-5p gene (miRBase Accession No. MIMAT0025847) described in SEQ ID NO: 115, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6511b-5p gene can be obtained by a method described in Li Y et al., 2012, Gene, Vol. 497, p. 330-335. Also, “hsa-mir-6511b-1 and hsa-mir-6511b-2” (miRBase Accession Nos. MI0022552 and MI0023431, SEQ ID NOs: 392 and 393) having a hairpin-like structure are known as precursors of “hsa-miR-6511b-5p”.

(198) The term “hsa-miR-4750-5p gene” or “hsa-miR-4750-5p” used herein includes the hsa-miR-4750-5p gene (miRBase Accession No. MIMAT0019887) described in SEQ ID NO: 116, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4750-5p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4750” (miRBase Accession No. MI0017389, SEQ ID NO: 394) having a hairpin-like structure is known as a precursor of “hsa-miR-4750-5p”.

(199) The term “hsa-miR-6126 gene” or “hsa-miR-6126” used herein includes the hsa-miR-6126 gene (miRBase Accession No. MIMAT0024599) described in SEQ ID NO: 117, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6126 gene can be obtained by a method described in Smith J L et al., 2012, J Virol. Vol. 86, p. 5278-5287. Also, “hsa-mir-6126” (miRBase Accession No. MI0021260, SEQ ID NO: 395) having a hairpin-like structure is known as a precursor of “hsa-miR-6126”.

(200) The term “hsa-miR-614 gene” or “hsa-miR-614” used herein includes the hsa-miR-614 gene (miRBase Accession No. MIMAT0003282) described in SEQ ID NO: 118, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-614 gene can be obtained by a method described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-614” (miRBase Accession No. MI0003627, SEQ ID NO: 396) having a hairpin-like structure is known as a precursor of “hsa-miR-614”.

(201) The term “hsa-miR-7110-5p gene” or “hsa-miR-7110-5p” used herein includes the hsa-miR-7110-5p gene (miRBase Accession No. MIMAT0028117) described in SEQ ID NO: 119, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7110-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-7110” (miRBase Accession No. MI0022961, SEQ ID NO: 397) having a hairpin-like structure is known as a precursor of “hsa-miR-7110-5p”.

(202) The term “hsa-miR-744-5p gene” or “hsa-miR-744-5p” used herein includes the hsa-miR-744-5p gene (miRBase Accession No. MIMAT0004945) described in SEQ ID NO: 120, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-744-5p gene can be obtained by a method described in Berezikov E et al., 2006, Genome Res, Vol. 16, p. 1289-1298. Also, “hsa-mir-744” (miRBase Accession No. MI0005559, SEQ ID NO: 398) having a hairpin-like structure is known as a precursor of “hsa-miR-744-5p”.

(203) The term “hsa-miR-6769a-5p gene” or “hsa-miR-6769a-5p” used herein includes the hsa-miR-6769a-5p gene (miRBase Accession No. MIMAT0027438) described in SEQ ID NO: 121, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6769a-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6769a” (miRBase Accession No. MI0022614, SEQ ID NO: 399) having a hairpin-like structure is known as a precursor of “hsa-miR-6769a-5p”.

(204) The term “hsa-miR-4792 gene” or “hsa-miR-4792” used herein includes the hsa-miR-4792 gene (miRBase Accession No. MIMAT0019964) described in SEQ ID NO: 122, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4792 gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4792” (miRBase Accession No. MI0017439, SEQ ID NO: 400) having a hairpin-like structure is known as a precursor of “hsa-miR-4792”.

(205) The term “hsa-miR-5787 gene” or “hsa-miR-5787” used herein includes the hsa-miR-5787 gene (miRBase Accession No. MIMAT0023252) described in SEQ ID NO: 123, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-5787 gene can be obtained by a method described in Yoo H et al., 2011, Biochem Biophys Res Commun, Vol. 415, p. 567-572. Also, “hsa-mir-5787” (miRBase Accession No. MI0019797, SEQ ID NO: 401) having a hairpin-like structure is known as a precursor of “hsa-miR-5787”.

(206) The term “hsa-miR-6798-5p gene” or “hsa-miR-6798-5p” used herein includes the hsa-miR-6798-5p gene (miRBase Accession No. MIMAT0027496) described in SEQ ID NO: 124, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6798-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6798” (miRBase Accession No. MI0022643, SEQ ID NO: 402) having a hairpin-like structure is known as a precursor of “hsa-miR-6798-5p”.

(207) The term “hsa-miR-6781-5p gene” or “hsa-miR-6781-5p” used herein includes the hsa-miR-6781-5p gene (miRBase Accession No. MIMAT0027462) described in SEQ ID NO: 125, a

homolog or an ortholog of a different organism species, and the like. The hsa-miR-6781-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6781” (miRBase Accession No. MI0022626, SEQ ID NO: 403) having a hairpin-like structure is known as a precursor of “hsa-miR-6781-5p”.

(208) The term “hsa-miR-4419b gene” or “hsa-miR-4419b” used herein includes the hsa-miR-4419b gene (miRBase Accession No. MIMAT0019034) described in SEQ ID NO: 126, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4419b gene can be obtained by a method described in Jima D D et al., 2010, *Blood*, Vol. 116, e118-e127. Also, “hsa-mir-4419b” (miRBase Accession No. MI0016861, SEQ ID NO: 404) having a hairpin-like structure is known as a precursor of “hsa-miR-4419b”.

(209) The term “hsa-miR-4446-3p gene” or “hsa-miR-4446-3p” used herein includes the hsa-miR-4446-3p gene (miRBase Accession No. MIMAT0018965) described in SEQ ID NO: 127, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4446-3p gene can be obtained by a method described in Jima D D et al., 2010, *Blood*, Vol. 116, e118-e127. Also, “hsa-mir-4446” (miRBase Accession No. MI0016789, SEQ ID NO: 405) having a hairpin-like structure is known as a precursor of “hsa-miR-4446-3p”.

(210) The term “hsa-miR-4259 gene” or “hsa-miR-4259” used herein includes the hsa-miR-4259 gene (miRBase Accession No. MIMAT0016880) described in SEQ ID NO: 128, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4259 gene can be obtained by a method described in Goff L A et al., 2009, *PLoS One*, Vol. 4, e7192. Also, “hsa-mir-4259” (miRBase Accession No. MI0015858, SEQ ID NO: 406) having a hairpin-like structure is known as a precursor of “hsa-miR-4259”.

(211) The term “hsa-miR-5572 gene” or “hsa-miR-5572” used herein includes the hsa-miR-5572 gene (miRBase Accession No. MIMAT0022260) described in SEQ ID NO: 129, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-5572 gene can be obtained by a method described in Tandon M et al., 2012, *Oral Dis*, Vol. 18, p. 127-131. Also, “hsa-mir-5572” (miRBase Accession No. MI0019117, SEQ ID NO: 407) having a hairpin-like structure is known as a precursor of “hsa-miR-5572”.

(212) The term “hsa-miR-6075 gene” or “hsa-miR-6075” used herein includes the hsa-miR-6075 gene (miRBase Accession No. MIMAT0023700) described in SEQ ID NO: 130, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6075 gene can be obtained by a method described in Voellenkle C et al., 2012, *RNA*, Vol. 18, p. 472-484. Also, “hsa-mir-6075” (miRBase Accession No. MI0020352, SEQ ID NO: 408) having a hairpin-like structure is known as a precursor of “hsa-miR-6075”.

(213) The term “hsa-miR-296-3p gene” or “hsa-miR-296-3p” used herein includes the hsa-miR-296-3p gene (miRBase Accession No. MIMAT0004679) described in SEQ ID NO: 131, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-296-3p gene can be obtained by a method described in Houbaviy H B et al., 2003, *Dev Cell*, Vol. 5, p. 351-358. Also, “hsa-mir-296” (miRBase Accession No. MI0000747, SEQ ID NO: 409) having a hairpin-like structure is known as a precursor of “hsa-miR-296-3p”.

(214) The term “hsa-miR-6891-5p gene” or “hsa-miR-6891-5p” used herein includes the hsa-miR-6891-5p gene (miRBase Accession No. MIMAT0027682) described in SEQ ID NO: 132, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6891-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6891” (miRBase Accession No. MI0022738, SEQ ID NO: 410) having a hairpin-like structure is known as a precursor of “hsa-miR-6891-5p”.

(215) The term “hsa-miR-4745-5p gene” or “hsa-miR-4745-5p” used herein includes the hsa-miR-4745-5p gene (miRBase Accession No. MIMAT0019878) described in SEQ ID NO: 133, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4745-5p gene can be obtained by a method described in Persson H et al., 2011, *Cancer Res*, Vol. 71, p. 78-86.

Also, “hsa-mir-4745” (miRBase Accession No. MI0017384, SEQ ID NO: 411) having a hairpin-like structure is known as a precursor of “hsa-miR-4745-5p”.

(216) The term “hsa-miR-6775-5p gene” or “hsa-miR-6775-5p” used herein includes the hsa-miR-6775-5p gene (miRBase Accession No. MIMAT0027450) described in SEQ ID NO: 134, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6775-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6775” (miRBase Accession No. MI0022620, SEQ ID NO: 412) having a hairpin-like structure is known as a precursor of “hsa-miR-6775-5p”.

(217) The term “hsa-miR-6870-5p gene” or “hsa-miR-6870-5p” used herein includes the hsa-miR-6870-5p gene (miRBase Accession No. MIMAT0027640) described in SEQ ID NO: 135, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6870-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6870” (miRBase Accession No. MI0022717, SEQ ID NO: 413) having a hairpin-like structure is known as a precursor of “hsa-miR-6870-5p”.

(218) The term “hsa-miR-920 gene” or “hsa-miR-920” used herein includes the hsa-miR-920 gene (miRBase Accession No. MIMAT0004970) described in SEQ ID NO: 136, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-920 gene can be obtained by a method described in Novotny G W et al., 2007, *Int J Androl*, Vol. 30, p. 316-326. Also, “hsa-mir-920” (miRBase Accession No. MI0005712, SEQ ID NO: 414) having a hairpin-like structure is known as a precursor of “hsa-miR-920”.

(219) The term “hsa-miR-4530 gene” or “hsa-miR-4530” used herein includes the hsa-miR-4530 gene (miRBase Accession No. MIMAT0019069) described in SEQ ID NO: 137, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4530 gene can be obtained by a method described in Jima D D et al., 2010, *Blood*, Vol. 116, e118-e127. Also, “hsa-mir-4530” (miRBase Accession No. MI0016897, SEQ ID NO: 415) having a hairpin-like structure is known as a precursor of “hsa-miR-4530”.

(220) The term “hsa-miR-6819-5p gene” or “hsa-miR-6819-5p” used herein includes the hsa-miR-6819-5p gene (miRBase Accession No. MIMAT0027538) described in SEQ ID NO: 138, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6819-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6819” (miRBase Accession No. MI0022664, SEQ ID NO: 416) having a hairpin-like structure is known as a precursor of “hsa-miR-6819-5p”.

(221) The term “hsa-miR-6825-5p gene” or “hsa-miR-6825-5p” used herein includes the hsa-miR-6825-5p gene (miRBase Accession No. MIMAT0027550) described in SEQ ID NO: 139, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6825-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6825” (miRBase Accession No. MI0022670, SEQ ID NO: 417) having a hairpin-like structure is known as a precursor of “hsa-miR-6825-5p”.

(222) The term “hsa-miR-7847-3p gene” or “hsa-miR-7847-3p” used herein includes the hsa-miR-7847-3p gene (miRBase Accession No. MIMAT0030422) described in SEQ ID NO: 140, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7847-3p gene can be obtained by a method described in Pie H et al., 2012, *PLoS One*, Vol. 7, e50746. Also, “hsa-mir-7847” (miRBase Accession No. MI0025517, SEQ ID NO: 418) having a hairpin-like structure is known as a precursor of “hsa-miR-7847-3p”.

(223) The term “hsa-miR-6131 gene” or “hsa-miR-6131” used herein includes the hsa-miR-6131 gene (miRBase Accession No. MIMAT0024615) described in SEQ ID NO: 141, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6131 gene can be obtained by a method described in Dannemann M et al., 2012, *Genome Biol Evol*, Vol. 4, p. 552-564. Also, “hsa-mir-6131” (miRBase Accession No. MI0021276, SEQ ID NO: 419) having a hairpin-like structure is known as a precursor of “hsa-miR-6131”.

(224) The term “hsa-miR-4433-3p gene” or “hsa-miR-4433-3p” used herein includes the hsa-miR-4433-3p gene (miRBase Accession No. MIMAT0018949) described in SEQ ID NO: 142, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4433-3p gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4433” (miRBase Accession No. MI0016773, SEQ ID NO: 420) having a hairpin-like structure is known as a precursor of “hsa-miR-4433-3p”.

(225) The term “hsa-miR-1228-5p gene” or “hsa-miR-1228-5p” used herein includes the hsa-miR-1228-5p gene (miRBase Accession No. MIMAT0005582) described in SEQ ID NO: 143, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1228-5p gene can be obtained by a method described in Berezikov E et al., 2007, Mol Cell, Vol. 28, p. 328-336. Also, “hsa-mir-1228” (miRBase Accession No. MI0006318, SEQ ID NO: 421) having a hairpin-like structure is known as a precursor of “hsa-miR-1228-5p”.

(226) The term “hsa-miR-6743-5p gene” or “hsa-miR-6743-5p” used herein includes the hsa-miR-6743-5p gene (miRBase Accession No. MIMAT0027387) described in SEQ ID NO: 144, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6743-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6743” (miRBase Accession No. MI0022588, SEQ ID NO: 422) having a hairpin-like structure is known as a precursor of “hsa-miR-6743-5p”.

(227) The term “hsa-miR-1268a gene” or “hsa-miR-1268a” used herein includes the hsa-miR-1268a gene (miRBase Accession No. MIMAT0005922) described in SEQ ID NO: 145, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1268a gene can be obtained by a method described in Morin R D et al., 2008, Genome Res, Vol. 18, p. 610-621. Also, “hsa-mir-1268a” (miRBase Accession No. MI0006405, SEQ ID NO: 423) having a hairpin-like structure is known as a precursor of “hsa-miR-1268a”.

(228) The term “hsa-miR-3917 gene” or “hsa-miR-3917” used herein includes the hsa-miR-3917 gene (miRBase Accession No. MIMAT0018191) described in SEQ ID NO: 146, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3917 gene can be obtained by a method described in Creighton C) et al., 2010, PLoS One, Vol. 5, e9637. Also, “hsa-mir-3917” (miRBase Accession No. MI0016423, SEQ ID NO: 424) having a hairpin-like structure is known as a precursor of “hsa-miR-3917”.

(229) The term “hsa-miR-6786-5p gene” or “hsa-miR-6786-5p” used herein includes the hsa-miR-6786-5p gene (miRBase Accession No. MIMAT0027472) described in SEQ ID NO: 147, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6786-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6786” (miRBase Accession No. MI0022631, SEQ ID NO: 425) having a hairpin-like structure is known as a precursor of “hsa-miR-6786-5p”.

(230) The term “hsa-miR-3154 gene” or “hsa-miR-3154” used herein includes the hsa-miR-3154 gene (miRBase Accession No. MIMAT0015028) described in SEQ ID NO: 148, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3154 gene can be obtained by a method described in Berezikov E et al., 2006, Genome Res, Vol. 16, p. 1289-1298. Also, “hsa-mir-3154” (miRBase Accession No. MI0014182, SEQ ID NO: 426) having a hairpin-like structure is known as a precursor of “hsa-miR-3154”.

(231) The term “hsa-miR-638 gene” or “hsa-miR-638” used herein includes the hsa-miR-638 gene (miRBase Accession No. MIMAT0003308) described in SEQ ID NO: 149, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-638 gene can be obtained by a method described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-638” (miRBase Accession No. MI0003653, SEQ ID NO: 427) having a hairpin-like structure is known as a precursor of “hsa-miR-638”.

(232) The term “hsa-miR-6741-5p gene” or “hsa-miR-6741-5p” used herein includes the hsa-miR-6741-5p gene (miRBase Accession No. MIMAT0027383) described in SEQ ID NO: 150, a

homolog or an ortholog of a different organism species, and the like. The hsa-miR-6741-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6741” (miRBase Accession No. MI0022586, SEQ ID NO: 428) having a hairpin-like structure is known as a precursor of “hsa-miR-6741-5p”.

(233) The term “hsa-miR-6889-5p gene” or “hsa-miR-6889-5p” used herein includes the hsa-miR-6889-5p gene (miRBase Accession No. MIMAT0027678) described in SEQ ID NO: 151, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6889-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6889” (miRBase Accession No. MI0022736, SEQ ID NO: 429) having a hairpin-like structure is known as a precursor of “hsa-miR-6889-5p”.

(234) The term “hsa-miR-6840-3p gene” or “hsa-miR-6840-3p” used herein includes the hsa-miR-6840-3p gene (miRBase Accession No. MIMAT0027583) described in SEQ ID NO: 152, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6840-3p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6840” (miRBase Accession No. MI0022686, SEQ ID NO: 430) having a hairpin-like structure is known as a precursor of “hsa-miR-6840-3p”.

(235) The term “hsa-miR-6510-5p gene” or “hsa-miR-6510-5p” used herein includes the hsa-miR-6510-5p gene (miRBase Accession No. MIMAT0025476) described in SEQ ID NO: 153, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6510-5p gene can be obtained by a method described in Joyce C E et al., 2011, Hum Mol Genet, Vol. 20, p. 4025-4040. Also, “hsa-mir-6510” (miRBase Accession No. MI0022222, SEQ ID NO: 431) having a hairpin-like structure is known as a precursor of “hsa-miR-6510-5p”.

(236) The term “hsa-miR-3188 gene” or “hsa-miR-3188” used herein includes the hsa-miR-3188 gene (miRBase Accession No. MIMAT0015070) described in SEQ ID NO: 154, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3188 gene can be obtained by a method described in Stark M S et al., 2010, PLoS One, Vol. 5, e9685. Also, “hsa-mir-3188” (miRBase Accession No. MI0014232, SEQ ID NO: 432) having a hairpin-like structure is known as a precursor of “hsa-miR-3188”.

(237) The term “hsa-miR-551b-5p gene” or “hsa-miR-551b-5p” used herein includes the hsa-miR-551b-5p gene (miRBase Accession No. MIMAT0004794) described in SEQ ID NO: 155, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-551b-5p gene can be obtained by a method described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-551b” (miRBase Accession No. MI0003575, SEQ ID NO: 433) having a hairpin-like structure is known as a precursor of “hsa-miR-551b-5p”.

(238) The term “hsa-miR-5001-5p gene” or “hsa-miR-5001-5p” used herein includes the hsa-miR-5001-5p gene (miRBase Accession No. MIMAT0021021) described in SEQ ID NO: 156, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-5001-5p gene can be obtained by a method described in Hansen T B et al., 2011, RNA Biol, Vol. 8, p. 378-383. Also, “hsa-mir-5001” (miRBase Accession No. MI0017867, SEQ ID NO: 434) having a hairpin-like structure is known as a precursor of “hsa-miR-5001-5p”.

(239) The term “hsa-miR-1268b gene” or “hsa-miR-1268b” used herein includes the hsa-miR-1268b gene (miRBase Accession No. MIMAT0018925) described in SEQ ID NO: 157, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1268b gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-1268b” (miRBase Accession No. MI0016748, SEQ ID NO: 435) having a hairpin-like structure is known as a precursor of “hsa-miR-1268b”.

(240) The term “hsa-miR-7107-5p gene” or “hsa-miR-7107-5p” used herein includes the hsa-miR-7107-5p gene (miRBase Accession No. MIMAT0028111) described in SEQ ID NO: 158, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7107-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-

1645. Also, “hsa-mir-7107” (miRBase Accession No. MI0022958, SEQ ID NO: 436) having a hairpin-like structure is known as a precursor of “hsa-miR-7107-5p”.

(241) The term “hsa-miR-6824-5p gene” or “hsa-miR-6824-5p” used herein includes the hsa-miR-6824-5p gene (miRBase Accession No. MIMAT0027548) described in SEQ ID NO: 159, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6824-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6824” (miRBase Accession No. MI0022669, SEQ ID NO: 437) having a hairpin-like structure is known as a precursor of “hsa-miR-6824-5p”.

(242) The term “hsa-miR-6732-5p gene” or “hsa-miR-6732-5p” used herein includes the hsa-miR-6732-5p gene (miRBase Accession No. MIMAT0027365) described in SEQ ID NO: 160, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6732-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6732” (miRBase Accession No. MI0022577, SEQ ID NO: 438) having a hairpin-like structure is known as a precursor of “hsa-miR-6732-5p”.

(243) The term “hsa-miR-371a-5p gene” or “hsa-miR-371a-5p” used herein includes the hsa-miR-371a-5p gene (miRBase Accession No. MIMAT0004687) described in SEQ ID NO: 161, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-371a-5p gene can be obtained by a method described in Suh M R et al., 2004, Dev Biol, Vol. 270, p. 488-498. Also, “hsa-mir-371a” (miRBase Accession No. MI0000779, SEQ ID NO: 439) having a hairpin-like structure is known as a precursor of “hsa-miR-371a-5p”.

(244) The term “hsa-miR-6794-5p gene” or “hsa-miR-6794-5p” used herein includes the hsa-miR-6794-5p gene (miRBase Accession No. MIMAT0027488) described in SEQ ID NO: 162, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6794-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6794” (miRBase Accession No. MI0022639, SEQ ID NO: 440) having a hairpin-like structure is known as a precursor of “hsa-miR-6794-5p”.

(245) The term “hsa-miR-6779-5p gene” or “hsa-miR-6779-5p” used herein includes the hsa-miR-6779-5p gene (miRBase Accession No. MIMAT0027458) described in SEQ ID NO: 163, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6779-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6779” (miRBase Accession No. MI0022624, SEQ ID NO: 441) having a hairpin-like structure is known as a precursor of “hsa-miR-6779-5p”.

(246) The term “hsa-miR-4271 gene” or “hsa-miR-4271” used herein includes the hsa-miR-4271 gene (miRBase Accession No. MIMAT0016901) described in SEQ ID NO: 164, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4271 gene can be obtained by a method described in Goff L A et al., 2009, PLoS One, Vol. 4, e7192. Also, “hsa-mir-4271” (miRBase Accession No. MI0015879, SEQ ID NO: 442) having a hairpin-like structure is known as a precursor of “hsa-miR-4271”.

(247) The term “hsa-miR-5195-3p gene” or “hsa-miR-5195-3p” used herein includes the hsa-miR-5195-3p gene (miRBase Accession No. MIMAT0021127) described in SEQ ID NO: 165, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-5195-3p gene can be obtained by a method described in Schotte D et al., 2011, Leukemia, Vol. 25, p. 1389-1399. Also, “hsa-mir-5195” (miRBase Accession No. MI0018174, SEQ ID NO: 443) having a hairpin-like structure is known as a precursor of “hsa-miR-5195-3p”.

(248) The term “hsa-miR-6762-5p gene” or “hsa-miR-6762-5p” used herein includes the hsa-miR-6762-5p gene (miRBase Accession No. MIMAT0027424) described in SEQ ID NO: 166, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6762-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6762” (miRBase Accession No. MI0022607, SEQ ID NO: 444) having a hairpin-like structure is known as a precursor of “hsa-miR-6762-5p”.

(249) The term “hsa-miR-939-5p gene” or “hsa-miR-939-5p” used herein includes the hsa-miR-939-5p gene (miRBase Accession No. MIMAT0004982) described in SEQ ID NO: 167, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-939-5p gene can be obtained by a method described in Lui W O et al., 2007, Cancer Res, Vol. 67, p. 6031-6043. Also, “hsa-mir-939” (miRBase Accession No. MI0005761, SEQ ID NO: 445) having a hairpin-like structure is known as a precursor of “hsa-miR-939-5p”.

(250) The term “hsa-miR-1247-3p gene” or “hsa-miR-1247-3p” used herein includes the hsa-miR-1247-3p gene (miRBase Accession No. MIMAT0022721) described in SEQ ID NO: 168, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1247-3p gene can be obtained by a method described in Morin R D et al., 2008, Genome Res, Vol. 18, p. 610-621. Also, “hsa-mir-1247” (miRBase Accession No. MI0006382, SEQ ID NO: 446) having a hairpin-like structure is known as a precursor of “hsa-miR-1247-3p”.

(251) The term “hsa-miR-6777-5p gene” or “hsa-miR-6777-5p” used herein includes the hsa-miR-6777-5p gene (miRBase Accession No. MIMAT0027454) described in SEQ ID NO: 169, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6777-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6777” (miRBase Accession No. MI0022622, SEQ ID NO: 447) having a hairpin-like structure is known as a precursor of “hsa-miR-6777-5p”.

(252) The term “hsa-miR-6722-3p gene” or “hsa-miR-6722-3p” used herein includes the hsa-miR-6722-3p gene (miRBase Accession No. MIMAT0025854) described in SEQ ID NO: 170, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6722-3p gene can be obtained by a method described in Li Y et al., 2012, Gene, Vol. 497, p. 330-335. Also, “hsa-mir-6722” (miRBase Accession No. MI0022557, SEQ ID NO: 448) having a hairpin-like structure is known as a precursor of “hsa-miR-6722-3p”.

(253) The term “hsa-miR-3656 gene” or “hsa-miR-3656” used herein includes the hsa-miR-3656 gene (miRBase Accession No. MIMAT0018076) described in SEQ ID NO: 171, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3656 gene can be obtained by a method described in Meiri E et al., 2010, Nucleic Acids Res, Vol. 38, p. 6234-6246. Also, “hsa-mir-3656” (miRBase Accession No. MI0016056, SEQ ID NO: 449) having a hairpin-like structure is known as a precursor of “hsa-miR-3656”.

(254) The term “hsa-miR-4688 gene” or “hsa-miR-4688” used herein includes the hsa-miR-4688 gene (miRBase Accession No. MIMAT0019777) described in SEQ ID NO: 172, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4688 gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4688” (miRBase Accession No. MI0017321, SEQ ID NO: 450) having a hairpin-like structure is known as a precursor of “hsa-miR-4688”.

(255) The term “hsa-miR-3195 gene” or “hsa-miR-3195” used herein includes the hsa-miR-3195 gene (miRBase Accession No. MIMAT0015079) described in SEQ ID NO: 173, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3195 gene can be obtained by a method described in Stark M S et al., 2010, PLoS One, Vol. 5, e9685. Also, “hsa-mir-3195” (miRBase Accession No. MI0014240, SEQ ID NO: 451) having a hairpin-like structure is known as a precursor of “hsa-miR-3195”.

(256) The term “hsa-miR-6766-5p gene” or “hsa-miR-6766-5p” used herein includes the hsa-miR-6766-5p gene (miRBase Accession No. MIMAT0027432) described in SEQ ID NO: 174, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6766-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6766” (miRBase Accession No. MI0022611, SEQ ID NO: 452) having a hairpin-like structure is known as a precursor of “hsa-miR-6766-5p”.

(257) The term “hsa-miR-4447 gene” or “hsa-miR-4447” used herein includes the hsa-miR-4447 gene (miRBase Accession No. MIMAT0018966) described in SEQ ID NO: 175, a homolog or an

ortholog of a different organism species, and the like. The hsa-miR-4447 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4447” (miRBase Accession No. MI0016790, SEQ ID NO: 453) having a hairpin-like structure is known as a precursor of “hsa-miR-4447”.

(258) The term “hsa-miR-4656 gene” or “hsa-miR-4656” used herein includes the hsa-miR-4656 gene (miRBase Accession No. MIMAT0019723) described in SEQ ID NO: 176, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4656 gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4656” (miRBase Accession No. MI0017284, SEQ ID NO: 454) having a hairpin-like structure is known as a precursor of “hsa-miR-4656”.

(259) The term “hsa-miR-7108-5p gene” or “hsa-miR-7108-5p” used herein includes the hsa-miR-7108-5p gene (miRBase Accession No. MIMAT0028113) described in SEQ ID NO: 177, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7108-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-7108” (miRBase Accession No. MI0022959, SEQ ID NO: 455) having a hairpin-like structure is known as a precursor of “hsa-miR-7108-5p”.

(260) The term “hsa-miR-3191-3p gene” or “hsa-miR-3191-3p” used herein includes the hsa-miR-3191-3p gene (miRBase Accession No. MIMAT0015075) described in SEQ ID NO: 178, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3191-3p gene can be obtained by a method described in Stark M S et al., 2010, PLoS One, Vol. 5, e9685. Also, “hsa-mir-3191” (miRBase Accession No. MI0014236, SEQ ID NO: 456) having a hairpin-like structure is known as a precursor of “hsa-miR-3191-3p”.

(261) The term “hsa-miR-1273g-3p gene” or “hsa-miR-1273g-3p” used herein includes the hsa-miR-1273g-3p gene (miRBase Accession No. MIMAT0022742) described in SEQ ID NO: 179, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1273g-3p gene can be obtained by a method described in Reshmi G et al., 2011, Genomics, Vol. 97, p. 333-340. Also, “hsa-mir-1273g” (miRBase Accession No. MI0018003, SEQ ID NO: 457) having a hairpin-like structure is known as a precursor of “hsa-miR-1273g-3p”.

(262) The term “hsa-miR-4463 gene” or “hsa-miR-4463” used herein includes the hsa-miR-4463 gene (miRBase Accession No. MIMAT0018987) described in SEQ ID NO: 180, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4463 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4463” (miRBase Accession No. MI0016811, SEQ ID NO: 458) having a hairpin-like structure is known as a precursor of “hsa-miR-4463”.

(263) The term “hsa-miR-2861 gene” or “hsa-miR-2861” used herein includes the hsa-miR-2861 gene (miRBase Accession No. MIMAT0013802) described in SEQ ID NO: 181, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-2861 gene can be obtained by a method described in Li H et al., 2009, Clin Invest, Vol. 119, p. 3666-3677. Also, “hsa-mir-2861” (miRBase Accession No. MI0013006, SEQ ID NO: 459) having a hairpin-like structure is known as a precursor of “hsa-miR-2861”.

(264) The term “hsa-miR-3196 gene” or “hsa-miR-3196” used herein includes the hsa-miR-3196 gene (miRBase Accession No. MIMAT0015080) described in SEQ ID NO: 182, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3196 gene can be obtained by a method described in Stark M S et al., 2010, PLoS One, Vol. 5, e9685. Also, “hsa-mir-3196” (miRBase Accession No. MI0014241, SEQ ID NO: 460) having a hairpin-like structure is known as a precursor of “hsa-miR-3196”.

(265) The term “hsa-miR-6877-5p gene” or “hsa-miR-6877-5p” used herein includes the hsa-miR-6877-5p gene (miRBase Accession No. MIMAT0027654) described in SEQ ID NO: 183, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6877-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-

1645. Also, “hsa-mir-6877” (miRBase Accession No. MI0022724, SEQ ID NO: 461) having a hairpin-like structure is known as a precursor of “hsa-miR-6877-5p”.

(266) The term “hsa-miR-3679-5p gene” or “hsa-miR-3679-5p” used herein includes the hsa-miR-3679-5p gene (miRBase Accession No. MIMAT0018104) described in SEQ ID NO: 184, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3679-5p gene can be obtained by a method described in Creighton C J et al., 2010, PLoS One, Vol. 5, e9637. Also, “hsa-mir-3679” (miRBase Accession No. MI0016080, SEQ ID NO: 462) having a hairpin-like structure is known as a precursor of “hsa-miR-3679-5p”.

(267) The term “hsa-miR-4442 gene” or “hsa-miR-4442” used herein includes the hsa-miR-4442 gene (miRBase Accession No. MIMAT0018960) described in SEQ ID NO: 185, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4442 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4442” (miRBase Accession No. MI0016785, SEQ ID NO: 463) having a hairpin-like structure is known as a precursor of “hsa-miR-4442”.

(268) The term “hsa-miR-6789-5p gene” or “hsa-miR-6789-5p” used herein includes the hsa-miR-6789-5p gene (miRBase Accession No. MIMAT0027478) described in SEQ ID NO: 186, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6789-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6789” (miRBase Accession No. MI0022634, SEQ ID NO: 464) having a hairpin-like structure is known as a precursor of “hsa-miR-6789-5p”.

(269) The term “hsa-miR-6782-5p gene” or “hsa-miR-6782-5p” used herein includes the hsa-miR-6782-5p gene (miRBase Accession No. MIMAT0027464) described in SEQ ID NO: 187, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6782-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6782” (miRBase Accession No. MI0022627, SEQ ID NO: 465) having a hairpin-like structure is known as a precursor of “hsa-miR-6782-5p”.

(270) The term “hsa-miR-486-3p gene” or “hsa-miR-486-3p” used herein includes the hsa-miR-486-3p gene (miRBase Accession No. MIMAT0004762) described in SEQ ID NO: 188, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-486-3p gene can be obtained by a method described in Fu H et al., 2005, FEBS Lett, Vol. 579, p. 3849-3854. Also, “hsa-mir-486 and hsa-mir-486-2” (miRBase Accession Nos. MI0002470 and MI0023622, SEQ ID NOs: 466 and 467) having a hairpin-like structure are known as precursors of “hsa-miR-486-3p”.

(271) The term “hsa-miR-6085 gene” or “hsa-miR-6085” used herein includes the hsa-miR-6085 gene (miRBase Accession No. MIMAT0023710) described in SEQ ID NO: 189, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6085 gene can be obtained by a method described in Voellenkle C et al., 2012, RNA, Vol. 18, p. 472-484. Also, “hsa-mir-6085” (miRBase Accession No. MI0020362, SEQ ID NO: 468) having a hairpin-like structure is known as a precursor of “hsa-miR-6085”.

(272) The term “hsa-miR-4746-3p gene” or “hsa-miR-4746-3p” used herein includes the hsa-miR-4746-3p gene (miRBase Accession No. MIMAT0019881) described in SEQ ID NO: 190, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4746-3p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4746” (miRBase Accession No. MI0017385, SEQ ID NO: 469) having a hairpin-like structure is known as a precursor of “hsa-miR-4746-3p”.

(273) The term “hsa-miR-619-5p gene” or “hsa-miR-619-5p” used herein includes the hsa-miR-619-5p gene (miRBase Accession No. MIMAT0026622) described in SEQ ID NO: 191, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-619-5p gene can be obtained by a method described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-619” (miRBase Accession No. MI0003633, SEQ ID NO: 470) having a hairpin-like structure is known as a precursor of “hsa-miR-619-5p”.

(274) The term “hsa-miR-937-5p gene” or “hsa-miR-937-5p” used herein includes the hsa-miR-937-5p gene (miRBase Accession No. MIMAT0022938) described in SEQ ID NO: 192, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-937-5p gene can be obtained by a method described in Lui W O et al., 2007, Cancer Res, Vol. 67, p. 6031-6043. Also, “hsa-mir-937” (miRBase Accession No. MI0005759, SEQ ID NO: 471) having a hairpin-like structure is known as a precursor of “hsa-miR-937-5p”.

(275) The term “hsa-miR-6803-5p gene” or “hsa-miR-6803-5p” used herein includes the hsa-miR-6803-5p gene (miRBase Accession No. MIMAT0027506) described in SEQ ID NO: 193, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6803-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6803” (miRBase Accession No. MI0022648, SEQ ID NO: 472) having a hairpin-like structure is known as a precursor of “hsa-miR-6803-5p”.

(276) The term “hsa-miR-4298 gene” or “hsa-miR-4298” used herein includes the hsa-miR-4298 gene (miRBase Accession No. MIMAT0016852) described in SEQ ID NO: 194, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4298 gene can be obtained by a method described in Goff L A et al., 2009, PLoS One, Vol. 4, e7192. Also, “hsa-mir-4298” (miRBase Accession No. MI0015830, SEQ ID NO: 473) having a hairpin-like structure is known as a precursor of “hsa-miR-4298”.

(277) The term “hsa-miR-4454 gene” or “hsa-miR-4454” used herein includes the hsa-miR-4454 gene (miRBase Accession No. MIMAT0018976) described in SEQ ID NO: 195, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4454 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4454” (miRBase Accession No. MI0016800, SEQ ID NO: 474) having a hairpin-like structure is known as a precursor of “hsa-miR-4454”.

(278) The term “hsa-miR-4459 gene” or “hsa-miR-4459” used herein includes the hsa-miR-4459 gene (miRBase Accession No. MIMAT0018981) described in SEQ ID NO: 196, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4459 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4459” (miRBase Accession No. MI0016805, SEQ ID NO: 475) having a hairpin-like structure is known as a precursor of “hsa-miR-4459”.

(279) The term “hsa-miR-7150 gene” or “hsa-miR-7150” used herein includes the hsa-miR-7150 gene (miRBase Accession No. MIMAT0028211) described in SEQ ID NO: 197, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7150 gene can be obtained by a method described in Oulas A et al., 2009, Nucleic Acids Res, Vol. 37, p. 3276-3287. Also, “hsa-mir-7150” (miRBase Accession No. MI0023610, SEQ ID NO: 476) having a hairpin-like structure is known as a precursor of “hsa-miR-7150”.

(280) The term “hsa-miR-6880-5p gene” or “hsa-miR-6880-5p” used herein includes the hsa-miR-6880-5p gene (miRBase Accession No. MIMAT0027660) described in SEQ ID NO: 198, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6880-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6880” (miRBase Accession No. MI0022727, SEQ ID NO: 477) having a hairpin-like structure is known as a precursor of “hsa-miR-6880-5p”.

(281) The term “hsa-miR-4449 gene” or “hsa-miR-4449” used herein includes the hsa-miR-4449 gene (miRBase Accession No. MIMAT0018968) described in SEQ ID NO: 199, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4449 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4449” (miRBase Accession No. MI0016792, SEQ ID NO: 478) having a hairpin-like structure is known as a precursor of “hsa-miR-4449”.

(282) The term “hsa-miR-8063 gene” or “hsa-miR-8063” used herein includes the hsa-miR-8063 gene (miRBase Accession No. MIMAT0030990) described in SEQ ID NO: 200, a homolog or an

ortholog of a different organism species, and the like. The hsa-miR-8063 gene can be obtained by a method described in Wang H J et al., 2013, Shock, Vol. 39, p. 480-487. Also, “hsa-mir-8063” (miRBase Accession No. MI0025899, SEQ ID NO: 479) having a hairpin-like structure is known as a precursor of “hsa-miR-8063”.

(283) The term “hsa-miR-4695-5p gene” or “hsa-miR-4695-5p” used herein includes the hsa-miR-4695-5p gene (miRBase Accession No. MIMAT0019788) described in SEQ ID NO: 201, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4695-5p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4695” (miRBase Accession No. MI0017328, SEQ ID NO: 480) having a hairpin-like structure is known as a precursor of “hsa-miR-4695-5p”.

(284) The term “hsa-miR-6132 gene” or “hsa-miR-6132” used herein includes the hsa-miR-6132 gene (miRBase Accession No. MIMAT0024616) described in SEQ ID NO: 202, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6132 gene can be obtained by a method described in Dannemann M et al., 2012, Genome Biol Evol, Vol. 4, p. 552-564. Also, “hsa-mir-6132” (miRBase Accession No. MI0021277, SEQ ID NO: 481) having a hairpin-like structure is known as a precursor of “hsa-miR-6132”.

(285) The term “hsa-miR-6829-5p gene” or “hsa-miR-6829-5p” used herein includes the hsa-miR-6829-5p gene (miRBase Accession No. MIMAT0027558) described in SEQ ID NO: 203, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6829-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6829” (miRBase Accession No. MI0022674, SEQ ID NO: 482) having a hairpin-like structure is known as a precursor of “hsa-miR-6829-5p”.

(286) The term “hsa-miR-4486 gene” or “hsa-miR-4486” used herein includes the hsa-miR-4486 gene (miRBase Accession No. MIMAT0019020) described in SEQ ID NO: 204, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4486 gene can be obtained by a method described in Lima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4486” (miRBase Accession No. MI0016847, SEQ ID NO: 483) having a hairpin-like structure is known as a precursor of “hsa-miR-4486”.

(287) The term “hsa-miR-6805-3p gene” or “hsa-miR-6805-3p” used herein includes the hsa-miR-6805-3p gene (miRBase Accession No. MIMAT0027511) described in SEQ ID NO: 205, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6805-3p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6805” (miRBase Accession No. MI0022650, SEQ ID NO: 304) having a hairpin-like structure is known as a precursor of “hsa-miR-6805-3p”.

(288) The term “hsa-miR-6826-5p gene” or “hsa-miR-6826-5p” used herein includes the hsa-miR-6826-5p gene (miRBase Accession No. MIMAT0027552) described in SEQ ID NO: 206, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6826-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6826” (miRBase Accession No. MI0022671, SEQ ID NO: 484) having a hairpin-like structure is known as a precursor of “hsa-miR-6826-5p”.

(289) The term “hsa-miR-4508 gene” or “hsa-miR-4508” used herein includes the hsa-miR-4508 gene (miRBase Accession No. MIMAT0019045) described in SEQ ID NO: 207, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4508 gene can be obtained by a method described in Lima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4508” (miRBase Accession No. MI0016872, SEQ ID NO: 485) having a hairpin-like structure is known as a precursor of “hsa-miR-4508”.

(290) The term “hsa-miR-1343-5p gene” or “hsa-miR-1343-5p” used herein includes the hsa-miR-1343-5p gene (miRBase Accession No. MIMAT0027038) described in SEQ ID NO: 208, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1343-5p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86.

Also, “hsa-mir-1343” (miRBase Accession No. MI0017320, SEQ ID NO: 361) having a hairpin-like structure is known as a precursor of “hsa-miR-1343-5p”.

(291) The term “hsa-miR-7114-5p gene” or “hsa-miR-7114-5p” used herein includes the hsa-miR-7114-5p gene (miRBase Accession No. MIMAT0028125) described in SEQ ID NO: 209, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7114-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-7114” (miRBase Accession No. MI0022965, SEQ ID NO: 486) having a hairpin-like structure is known as a precursor of “hsa-miR-7114-5p”.

(292) The term “hsa-miR-3622a-5p gene” or “hsa-miR-3622a-5p” used herein includes the hsa-miR-3622a-5p gene (miRBase Accession No. MIMAT0018003) described in SEQ ID NO: 210, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3622a-5p gene can be obtained by a method described in Witten D et al., 2010, BMC Biol, Vol. 8, p. 58. Also, “hsa-mir-3622a” (miRBase Accession No. MI0016013, SEQ ID NO: 487) having a hairpin-like structure is known as a precursor of “hsa-miR-3622a-5p”.

(293) The term “hsa-miR-6765-5p gene” or “hsa-miR-6765-5p” used herein includes the hsa-miR-6765-5p gene (miRBase Accession No. MIMAT0027430) described in SEQ ID NO: 211, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6765-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6765” (miRBase Accession No. MI0022610, SEQ ID NO: 488) having a hairpin-like structure is known as a precursor of “hsa-miR-6765-5p”.

(294) The term “hsa-miR-7845-5p gene” or “hsa-miR-7845-5p” used herein includes the hsa-miR-7845-5p gene (miRBase Accession No. MIMAT0030420) described in SEQ ID NO: 212, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7845-5p gene can be obtained by a method described in Pie H et al., 2012, PLoS One, Vol. 7, e50746. Also, “hsa-mir-7845” (miRBase Accession No. MI0025515, SEQ ID NO: 489) having a hairpin-like structure is known as a precursor of “hsa-miR-7845-5p”.

(295) The term “hsa-miR-3960 gene” or “hsa-miR-3960” used herein includes the hsa-miR-3960 gene (miRBase Accession No. MIMAT0019337) described in SEQ ID NO: 213, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3960 gene can be obtained by a method described in Hu R et al., 2011, J Biol Chem, Vol. 286, p. 12328-12339. Also, “hsa-mir-3960” (miRBase Accession No. MI0016964, SEQ ID NO: 490) having a hairpin-like structure is known as a precursor of “hsa-miR-3960”.

(296) The term “hsa-miR-6749-5p gene” or “hsa-miR-6749-5p” used herein includes the hsa-miR-6749-5p gene (miRBase Accession No. MIMAT0027398) described in SEQ ID NO: 214, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6749-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6749” (miRBase Accession No. MI0022594, SEQ ID NO: 491) having a hairpin-like structure is known as a precursor of “hsa-miR-6749-5p”.

(297) The term “hsa-miR-1260b gene” or “hsa-miR-1260b” used herein includes the hsa-miR-1260b gene (miRBase Accession No. MIMAT0015041) described in SEQ ID NO: 215, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1260b gene can be obtained by a method described in Stark M S et al., 2010, PLoS One, Vol. 5, e9685. Also, “hsa-mir-1260b” (miRBase Accession No. MI0014197, SEQ ID NO: 492) having a hairpin-like structure is known as a precursor of “hsa-miR-1260b”.

(298) The term “hsa-miR-6799-5p gene” or “hsa-miR-6799-5p” used herein includes the hsa-miR-6799-5p gene (miRBase Accession No. MIMAT0027498) described in SEQ ID NO: 216, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6799-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6799” (miRBase Accession No. MI0022644, SEQ ID NO: 493) having a hairpin-like structure is known as a precursor of “hsa-miR-6799-5p”.

(299) The term “hsa-miR-4723-5p gene” or “hsa-miR-4723-5p” used herein includes the hsa-miR-4723-5p gene (miRBase Accession No. MIMAT0019838) described in SEQ ID NO: 217, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4723-5p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4723” (miRBase Accession No. MI0017359, SEQ ID NO: 494) having a hairpin-like structure is known as a precursor of “hsa-miR-4723-5p”.

(300) The term “hsa-miR-6784-5p gene” or “hsa-miR-6784-5p” used herein includes the hsa-miR-6784-5p gene (miRBase Accession No. MIMAT0027468) described in SEQ ID NO: 218, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6784-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6784” (miRBase Accession No. MI0022629, SEQ ID NO: 495) having a hairpin-like structure is known as a precursor of “hsa-miR-6784-5p”.

(301) The term “hsa-miR-5100 gene” or “hsa-miR-5100” used herein includes the hsa-miR-5100 gene (miRBase Accession No. MIMAT0022259) described in SEQ ID NO: 219, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-5100 gene can be obtained by a method described in Tandon M et al., 2012, Oral Dis, Vol. 18, p. 127-131. Also, “hsa-mir-5100” (miRBase Accession No. MI0019116, SEQ ID NO: 496) having a hairpin-like structure is known as a precursor of “hsa-miR-5100”.

(302) The term “hsa-miR-6769b-5p gene” or “hsa-miR-6769b-5p” used herein includes the hsa-miR-6769b-5p gene (miRBase Accession No. MIMAT0027620) described in SEQ ID NO: 220, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6769b-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6769b” (miRBase Accession No. MI0022706, SEQ ID NO: 497) having a hairpin-like structure is known as a precursor of “hsa-miR-6769b-5p”.

(303) The term “hsa-miR-1207-5p gene” or “hsa-miR-1207-5p” used herein includes the hsa-miR-1207-5p gene (miRBase Accession No. MIMAT0005871) described in SEQ ID NO: 221, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1207-5p gene can be obtained by a method described in Huppi K et al., 2008, Mol Cancer Res, Vol. 6, p. 212-221. Also, “hsa-mir-1207” (miRBase Accession No. MI0006340, SEQ ID NO: 498) having a hairpin-like structure is known as a precursor of “hsa-miR-1207-5p”.

(304) The term “hsa-miR-642a-3p gene” or “hsa-miR-642a-3p” used herein includes the hsa-miR-642a-3p gene (miRBase Accession No. MIMAT0020924) described in SEQ ID NO: 222, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-642a-3p gene can be obtained by a method described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-642a” (miRBase Accession No. MI0003657, SEQ ID NO: 499) having a hairpin-like structure is known as a precursor of “hsa-miR-642a-3p”.

(305) The term “hsa-miR-4505 gene” or “hsa-miR-4505” used herein includes the hsa-miR-4505 gene (miRBase Accession No. MIMAT0019041) described in SEQ ID NO: 223, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4505 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4505” (miRBase Accession No. MI0016868, SEQ ID NO: 500) having a hairpin-like structure is known as a precursor of “hsa-miR-4505”.

(306) The term “hsa-miR-4270 gene” or “hsa-miR-4270” used herein includes the hsa-miR-4270 gene (miRBase Accession No. MIMAT0016900) described in SEQ ID NO: 224, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4270 gene can be obtained by a method described in Goff L A et al., 2009, PLoS One, Vol. 4, e7192. Also, “hsa-mir-4270” (miRBase Accession No. MI0015878, SEQ ID NO: 501) having a hairpin-like structure is known as a precursor of “hsa-miR-4270”.

(307) The term “hsa-miR-6721-5p gene” or “hsa-miR-6721-5p” used herein includes the hsa-miR-6721-5p gene (miRBase Accession No. MIMAT0025852) described in SEQ ID NO: 225, a

homolog or an ortholog of a different organism species, and the like. The hsa-miR-6721-5p gene can be obtained by a method described in Li Y et al., 2012, *Gene*, Vol. 497, p. 330-335. Also, “hsa-mir-6721” (miRBase Accession No. MI0022556, SEQ ID NO: 502) having a hairpin-like structure is known as a precursor of “hsa-miR-6721-5p”.

(308) The term “hsa-miR-7111-5p gene” or “hsa-miR-7111-5p” used herein includes the hsa-miR-7111-5p gene (miRBase Accession No. MIMAT0028119) described in SEQ ID NO: 226, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7111-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-7111” (miRBase Accession No. MI0022962, SEQ ID NO: 503) having a hairpin-like structure is known as a precursor of “hsa-miR-7111-5p”.

(309) The term “hsa-miR-6791-5p gene” or “hsa-miR-6791-5p” used herein includes the hsa-miR-6791-5p gene (miRBase Accession No. MIMAT0027482) described in SEQ ID NO: 227, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6791-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6791” (miRBase Accession No. MI0022636, SEQ ID NO: 504) having a hairpin-like structure is known as a precursor of “hsa-miR-6791-5p”.

(310) The term “hsa-miR-7109-5p gene” or “hsa-miR-7109-5p” used herein includes the hsa-miR-7109-5p gene (miRBase Accession No. MIMAT0028115) described in SEQ ID NO: 228, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7109-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-7109” (miRBase Accession No. MI0022960, SEQ ID NO: 505) having a hairpin-like structure is known as a precursor of “hsa-miR-7109-5p”.

(311) The term “hsa-miR-4258 gene” or “hsa-miR-4258” used herein includes the hsa-miR-4258 gene (miRBase Accession No. MIMAT0016879) described in SEQ ID NO: 229, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4258 gene can be obtained by a method described in Goff L A et al., 2009, *PLoS One*, Vol. 4, e7192. Also, “hsa-mir-4258” (miRBase Accession No. MI0015857, SEQ ID NO: 506) having a hairpin-like structure is known as a precursor of “hsa-miR-4258”.

(312) The term “hsa-miR-6515-3p gene” or “hsa-miR-6515-3p” used herein includes the hsa-miR-6515-3p gene (miRBase Accession No. MIMAT0025487) described in SEQ ID NO: 230, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6515-3p gene can be obtained by a method described in Joyce C E et al., 2011, *Hum Mol Genet*, Vol. 20, p. 4025-4040. Also, “hsa-mir-6515” (miRBase Accession No. MI0022227, SEQ ID NO: 507) having a hairpin-like structure is known as a precursor of “hsa-miR-6515-3p”.

(313) The term “hsa-miR-6851-5p gene” or “hsa-miR-6851-5p” used herein includes the hsa-miR-6851-5p gene (miRBase Accession No. MIMAT0027602) described in SEQ ID NO: 231, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6851-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6851” (miRBase Accession No. MI0022697, SEQ ID NO: 508) having a hairpin-like structure is known as a precursor of “hsa-miR-6851-5p”.

(314) The term “hsa-miR-6125 gene” or “hsa-miR-6125” used herein includes the hsa-miR-6125 gene (miRBase Accession No. MIMAT0024598) described in SEQ ID NO: 232, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6125 gene can be obtained by a method described in Smith J L et al., 2012, *J Virol*, Vol. 86, p. 5278-5287. Also, “hsa-mir-6125” (miRBase Accession No. MI0021259, SEQ ID NO: 509) having a hairpin-like structure is known as a precursor of “hsa-miR-6125”.

(315) The term “hsa-miR-4749-5p gene” or “hsa-miR-4749-5p” used herein includes the hsa-miR-4749-5p gene (miRBase Accession No. MIMAT0019885) described in SEQ ID NO: 233, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4749-5p gene can be obtained by a method described in Persson H et al., 2011, *Cancer Res*, Vol. 71, p. 78-86.

Also, “hsa-mir-4749” (miRBase Accession No. MI0017388, SEQ ID NO: 510) having a hairpin-like structure is known as a precursor of “hsa-miR-4749-5p”.

(316) The term “hsa-miR-4726-5p gene” or “hsa-miR-4726-5p” used herein includes the hsa-miR-4726-5p gene (miRBase Accession No. MIMAT0019845) described in SEQ ID NO: 234, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4726-5p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4726” (miRBase Accession No. MI0017363, SEQ ID NO: 511) having a hairpin-like structure is known as a precursor of “hsa-miR-4726-5p”.

(317) The term “hsa-miR-4513 gene” or “hsa-miR-4513” used herein includes the hsa-miR-4513 gene (miRBase Accession No. MIMAT0019050) described in SEQ ID NO: 235, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4513 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4513” (miRBase Accession No. MI0016879, SEQ ID NO: 512) having a hairpin-like structure is known as a precursor of “hsa-miR-4513”.

(318) The term “hsa-miR-760 gene” or “hsa-miR-760” used herein includes the hsa-miR-760 gene (miRBase Accession No. MIMAT0004957) described in SEQ ID NO: 236, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-760 gene can be obtained by a method described in Berezikov E et al., 2006, Genome Res, Vol. 16, p. 1289-1298. Also, “hsa-mir-760” (miRBase Accession No. MI0005567, SEQ ID NO: 513) having a hairpin-like structure is known as a precursor of “hsa-miR-760”.

(319) The term “hsa-miR-602 gene” or “hsa-miR-602” used herein includes the hsa-miR-602 gene (miRBase Accession No. MIMAT0003270) described in SEQ ID NO: 237, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-602 gene can be obtained by a method described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-602” (miRBase Accession No. MI0003615, SEQ ID NO: 514) having a hairpin-like structure is known as a precursor of “hsa-miR-602”.

(320) The term “hsa-miR-423-5p gene” or “hsa-miR-423-5p” used herein includes the hsa-miR-423-5p gene (miRBase Accession No. MIMAT0004748) described in SEQ ID NO: 238, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-423-5p gene can be obtained by a method described in Kasashima K et al., 2004, Biochem Biophys Res Commun, Vol. 322, p. 403-410. Also, “hsa-mir-423” (miRBase Accession No. MI0001445, SEQ ID NO: 515) having a hairpin-like structure is known as a precursor of “hsa-miR-423-5p”.

(321) The term “hsa-miR-92a-2-5p gene” or “hsa-miR-92a-2-5p” used herein includes the hsa-miR-92a-2-5p gene (miRBase Accession No. MIMAT0004508) described in SEQ ID NO: 239, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-92a-2-5p gene can be obtained by a method described in Mourelatos Z et al., 2002, Genes Dev, Vol. 16, p. 720-728. Also, “hsa-mir-92a-2” (miRBase Accession No. MI0000094, SEQ ID NO: 516) having a hairpin-like structure is known as a precursor of “hsa-miR-92a-2-5p”.

(322) The term “hsa-miR-16-5p gene” or “hsa-miR-16-5p” used herein includes the hsa-miR-16-5p gene (miRBase Accession No. MIMAT0000069) described in SEQ ID NO: 240, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-16-5p gene can be obtained by a method described in Lagos-Quintana M et al., 2001, Science, Vol. 294, p. 853-858. Also, “hsa-mir-16-1 and hsa-mir-16-2” (miRBase Accession Nos. MI0000070 and MI0000115, SEQ ID NOs: 517 and 518) having a hairpin-like structure are known as precursors of “hsa-miR-16-5p”.

(323) The term “hsa-miR-451a gene” or “hsa-miR-451a” used herein includes the hsa-miR-451a gene (miRBase Accession No. MIMAT0001631) described in SEQ ID NO: 241, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-451a gene can be obtained by a method described in Altuvia Y et al., 2005, Nucleic Acids Res, Vol. 33, p. 2697-2706. Also, “hsa-mir-451a” (miRBase Accession No. MI0001729, SEQ ID NO: 519) having a hairpin-like structure is known as a precursor of “hsa-miR-451a”.

(324) The term “hsa-miR-135a-3p gene” or “hsa-miR-135a-3p” used herein includes the hsa-miR-135a-3p gene (miRBase Accession No. MIMAT0004595) described in SEQ ID NO: 242, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-135a-3p gene can be obtained by a method described in Lagos-Quintana M et al., 2002, *Curr Biol*, Vol. 12, p. 735-739. Also, “hsa-mir-135a” (miRBase Accession No. MI0000452, SEQ ID NO: 520) having a hairpin-like structure is known as a precursor of “hsa-miR-135a-3p”.

(325) The term “hsa-miR-486-5p gene” or “hsa-miR-486-5p” used herein includes the hsa-miR-486-5p gene (miRBase Accession No. MIMAT0002177) described in SEQ ID NO: 243, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-486-5p gene can be obtained by a method described in Fu H et al., 2005, *FEBS Lett*, Vol. 579, p. 3849-3854. Also, “hsa-mir-486 and hsa-mir-486-2” (miRBase Accession Nos. MI0002470 and MI0023622, SEQ ID NOs: 466 and 467) having a hairpin-like structure are known as precursors of “hsa-miR-486-5p”.

(326) The term “hsa-miR-4257 gene” or “hsa-miR-4257” used herein includes the hsa-miR-4257 gene (miRBase Accession No. MIMAT0016878) described in SEQ ID NO: 244, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4257 gene can be obtained by a method described in Goff L A et al., 2009, *PLoS One*, Vol. 4, e7192. Also, “hsa-mir-4257” (miRBase Accession No. MI0015856, SEQ ID NO: 521) having a hairpin-like structure is known as a precursor of “hsa-miR-4257”.

(327) The term “hsa-miR-92b-5p gene” or “hsa-miR-92b-5p” used herein includes the hsa-miR-92b-5p gene (miRBase Accession No. MIMAT0004792) described in SEQ ID NO: 245, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-92b-5p gene can be obtained by a method described in Cummins J M et al., 2006, *Proc Natl Acad Sci USA*, Vol. 103, p. 3687-3692. Also, “hsa-mir-92b” (miRBase Accession No. MI0003560, SEQ ID NO: 522) having a hairpin-like structure is known as a precursor of “hsa-miR-92b-5p”.

(328) The term “hsa-miR-1915-3p gene” or “hsa-miR-1915-3p” used herein includes the hsa-miR-1915-3p gene (miRBase Accession No. MIMAT0007892) described in SEQ ID NO: 246, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1915-3p gene can be obtained by a method described in Bar M et al., 2008, *Stem Cells*, Vol. 26, p. 2496-2505. Also, “hsa-mir-1915” (miRBase Accession No. MI0008336, SEQ ID NO: 339) having a hairpin-like structure is known as a precursor of “hsa-miR-1915-3p”.

(329) The term “hsa-miR-718 gene” or “hsa-miR-718” used herein includes the hsa-miR-718 gene (miRBase Accession No. MIMAT0012735) described in SEQ ID NO: 247, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-718 gene can be obtained by a method described in Artzi S et al., 2008, *BMC Bioinformatics*, Vol. 9, p. 39. Also, “hsa-mir-718” (miRBase Accession No. MI0012489, SEQ ID NO: 523) having a hairpin-like structure is known as a precursor of “hsa-miR-718”.

(330) The term “hsa-miR-940 gene” or “hsa-miR-940” used herein includes the hsa-miR-940 gene (miRBase Accession No. MIMAT0004983) described in SEQ ID NO: 248, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-940 gene can be obtained by a method described in Lui W O et al., 2007, *Cancer Res*, Vol. 67, p. 6031-6043. Also, “hsa-mir-940” (miRBase Accession No. MI0005762, SEQ ID NO: 524) having a hairpin-like structure is known as a precursor of “hsa-miR-940”.

(331) The term “hsa-miR-296-5p gene” or “hsa-miR-296-5p” used herein includes the hsa-miR-296-5p gene (miRBase Accession No. MIMAT0000690) described in SEQ ID NO: 249, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-296-5p gene can be obtained by a method described in Houbaviy H B et al., 2003, *Dev Cell*, Vol. 5, p. 351-358. Also, “hsa-mir-296” (miRBase Accession No. MI0000747, SEQ ID NO: 409) having a hairpin-like structure is known as a precursor of “hsa-miR-296-5p”.

(332) The term “hsa-miR-23b-3p gene” or “hsa-miR-23b-3p” used herein includes the hsa-miR-23b-3p gene (miRBase Accession No. MIMAT0000418) described in SEQ ID NO: 250, a homolog

or an ortholog of a different organism species, and the like. The hsa-miR-23b-3p gene can be obtained by a method described in Lagos-Quintana M et al., 2002, *Curr Biol*, Vol. 12, p. 735-739. Also, “hsa-mir-23b” (miRBase Accession No. MI0000439, SEQ ID NO: 525) having a hairpin-like structure is known as a precursor of “hsa-miR-23b-3p”.

(333) The term “hsa-miR-92a-3p gene” or “hsa-miR-92a-3p” used herein includes the hsa-miR-92a-3p gene (miRBase Accession No. MIMAT0000092) described in SEQ ID NO: 251, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-92a-3p gene can be obtained by a method described in Mourelatos Z et al., 2002, *Genes Dev*, Vol. 16, p. 720-728. Also, “hsa-mir-92a-1 and hsa-mir-92a-2” (miRBase Accession Nos. MI0000093 and MI0000094, SEQ ID NOs: 526 and 527) having a hairpin-like structure are known as precursors of “hsa-miR-92a-3p”.

(334) The term “hsa-miR-658 gene” or “hsa-miR-658” used herein includes the hsa-miR-658 gene (miRBase Accession No. MIMAT0003336) described in SEQ ID NO: 252, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-658 gene can be obtained by a method described in Cummins J M et al., 2006, *Proc Natl Acad Sci USA*, Vol. 103, p. 3687-3692. Also, “hsa-mir-658” (miRBase Accession No. MI0003682, SEQ ID NO: 528) having a hairpin-like structure is known as a precursor of “hsa-miR-658”.

(335) The term “hsa-miR-6842-5p gene” or “hsa-miR-6842-5p” used herein includes the hsa-miR-6842-5p gene (miRBase Accession No. MIMAT0027586) described in SEQ ID NO: 253, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6842-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6842” (miRBase Accession No. MI0022688, SEQ ID NO: 529) having a hairpin-like structure is known as a precursor of “hsa-miR-6842-5p”.

(336) The term “hsa-miR-6124 gene” or “hsa-miR-6124” used herein includes the hsa-miR-6124 gene (miRBase Accession No. MIMAT0024597) described in SEQ ID NO: 254, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6124 gene can be obtained by a method described in Smith J L et al., 2012, *J Virol*, Vol. 86, p. 5278-5287. Also, “hsa-mir-6124” (miRBase Accession No. MI0021258, SEQ ID NO: 530) having a hairpin-like structure is known as a precursor of “hsa-miR-6124”.

(337) The term “hsa-miR-6765-3p gene” or “hsa-miR-6765-3p” used herein includes the hsa-miR-6765-3p gene (miRBase Accession No. MIMAT0027431) described in SEQ ID NO: 255, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6765-3p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-6765” (miRBase Accession No. MI0022610, SEQ ID NO: 531) having a hairpin-like structure is known as a precursor of “hsa-miR-6765-3p”.

(338) The term “hsa-miR-7106-5p gene” or “hsa-miR-7106-5p” used herein includes the hsa-miR-7106-5p gene (miRBase Accession No. MIMAT0028109) described in SEQ ID NO: 256, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-7106-5p gene can be obtained by a method described in Ladewig E et al., 2012, *Genome Res*, Vol. 22, p. 1634-1645. Also, “hsa-mir-7106” (miRBase Accession No. MI0022957, SEQ ID NO: 532) having a hairpin-like structure is known as a precursor of “hsa-miR-7106-5p”.

(339) The term “hsa-miR-4534 gene” or “hsa-miR-4534” used herein includes the hsa-miR-4534 gene (miRBase Accession No. MIMAT0019073) described in SEQ ID NO: 257, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4534 gene can be obtained by a method described in Jima D D et al., 2010, *Blood*, Vol. 116, e118-e127. Also, “hsa-mir-4534” (miRBase Accession No. MI0016901, SEQ ID NO: 533) having a hairpin-like structure is known as a precursor of “hsa-miR-4534”.

(340) The term “hsa-miR-92b-3p gene” or “hsa-miR-92b-3p” used herein includes the hsa-miR-92b-3p gene (miRBase Accession No. MIMAT0003218) described in SEQ ID NO: 258, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-92b-3p gene can be

obtained by a method described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-92b” (miRBase Accession No. MI0003560, SEQ ID NO: 522) having a hairpin-like structure is known as a precursor of “hsa-miR-92b-3p”.

(341) The term “hsa-miR-3135b gene” or “hsa-miR-3135b” used herein includes the hsa-miR-3135b gene (miRBase Accession No. MIMAT0018985) described in SEQ ID NO: 259, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3135b gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-3135b” (miRBase Accession No. MI0016809, SEQ ID NO: 534) having a hairpin-like structure is known as a precursor of “hsa-miR-3135b”.

(342) The term “hsa-miR-4687-3p gene” or “hsa-miR-4687-3p” used herein includes the hsa-miR-4687-3p gene (miRBase Accession No. MIMAT0019775) described in SEQ ID NO: 260, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4687-3p gene can be obtained by a method described in Persson H et al., 2011, Cancer Res, Vol. 71, p. 78-86. Also, “hsa-mir-4687” (miRBase Accession No. MI0017319, SEQ ID NO: 535) having a hairpin-like structure is known as a precursor of “hsa-miR-4687-3p”.

(343) The term “hsa-miR-762 gene” or “hsa-miR-762” used herein includes the hsa-miR-762 gene (miRBase Accession No. MIMAT0010313) described in SEQ ID NO: 261, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-762 gene can be obtained by a method described in Berezikov E et al., 2006, Genome Res, Vol. 16, p. 1289-1298. Also, “hsa-mir-762” (miRBase Accession No. MI0003892, SEQ ID NO: 536) having a hairpin-like structure is known as a precursor of “hsa-miR-762”.

(344) The term “hsa-miR-3619-3p gene” or “hsa-miR-3619-3p” used herein includes the hsa-miR-3619-3p gene (miRBase Accession No. MIMAT0019219) described in SEQ ID NO: 262, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-3619-3p gene can be obtained by a method described in Witten D et al., 2010, BMC Biol, Vol. 8, p. 58. Also, “hsa-mir-3619” (miRBase Accession No. MI0016009, SEQ ID NO: 537) having a hairpin-like structure is known as a precursor of “hsa-miR-3619-3p”.

(345) The term “hsa-miR-4467 gene” or “hsa-miR-4467” used herein includes the hsa-miR-4467 gene (miRBase Accession No. MIMAT0018994) described in SEQ ID NO: 263, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4467 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4467” (miRBase Accession No. MI0016818, SEQ ID NO: 538) having a hairpin-like structure is known as a precursor of “hsa-miR-4467”.

(346) The term “hsa-miR-557 gene” or “hsa-miR-557” used herein includes the hsa-miR-557 gene (miRBase Accession No. MIMAT0003221) described in SEQ ID NO: 264, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-557 gene can be obtained by a method described in Cummins J M et al., 2006, Proc Natl Acad Sci USA, Vol. 103, p. 3687-3692. Also, “hsa-mir-557” (miRBase Accession No. MI0003563, SEQ ID NO: 539) having a hairpin-like structure is known as a precursor of “hsa-miR-557”.

(347) The term “hsa-miR-1237-5p gene” or “hsa-miR-1237-5p” used herein includes the hsa-miR-1237-5p gene (miRBase Accession No. MIMAT0022946) described in SEQ ID NO: 265, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1237-5p gene can be obtained by a method described in Berezikov E et al., 2007, Mol Cell, Vol. 28, p. 328-336. Also, “hsa-mir-1237” (miRBase Accession No. MI0006327, SEQ ID NO: 540) having a hairpin-like structure is known as a precursor of “hsa-miR-1237-5p”.

(348) The term “hsa-miR-1908-5p gene” or “hsa-miR-1908-5p” used herein includes the hsa-miR-1908-5p gene (miRBase Accession No. MIMAT0007881) described in SEQ ID NO: 266, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-1908-5p gene can be obtained by a method described in Bar M et al., 2008, Stem Cells, Vol. 26, p. 2496-2505. Also, “hsa-mir-1908” (miRBase Accession No. MI0008329, SEQ ID NO: 541) having a hairpin-

like structure is known as a precursor of “hsa-miR-1908-5p”.

(349) The term “hsa-miR-4286 gene” or “hsa-miR-4286” used herein includes the hsa-miR-4286 gene (miRBase Accession No. MIMAT0016916) described in SEQ ID NO: 267, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4286 gene can be obtained by a method described in Goff L A et al., 2009, PLoS One, Vol. 4, e7192. Also, “hsa-mir-4286” (miRBase Accession No. MI0015894, SEQ ID NO: 542) having a hairpin-like structure is known as a precursor of “hsa-miR-4286”.

(350) The term “hsa-miR-6885-5p gene” or “hsa-miR-6885-5p” used herein includes the hsa-miR-6885-5p gene (miRBase Accession No. MIMAT0027670) described in SEQ ID NO: 268, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6885-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6885” (miRBase Accession No. MI0022732, SEQ ID NO: 543) having a hairpin-like structure is known as a precursor of “hsa-miR-6885-5p”.

(351) The term “hsa-miR-6763-5p gene” or “hsa-miR-6763-5p” used herein includes the hsa-miR-6763-5p gene (miRBase Accession No. MIMAT0027426) described in SEQ ID NO: 269, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6763-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6763” (miRBase Accession No. MI0022608, SEQ ID NO: 544) having a hairpin-like structure is known as a precursor of “hsa-miR-6763-5p”.

(352) The term “hsa-miR-6089 gene” or “hsa-miR-6089” used herein includes the hsa-miR-6089 gene (miRBase Accession No. MIMAT0023714) described in SEQ ID NO: 851, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6089 gene can be obtained by a method described in Yoo J K et al., 2012, Stem Cells Dev, Vol. 21, p. 2049-2057. Also, “hsa-mir-6089-1 and hsa-mir-6089-2” (miRBase Accession Nos. MI0020366 and MI0023563, SEQ ID NOs: 857 and 858) having a hairpin-like structure are known as precursors of “hsa-miR-6089”.

(353) The term “hsa-miR-6816-5p gene” or “hsa-miR-6816-5p” used herein includes the hsa-miR-6816-5p gene (miRBase Accession No. MIMAT0027532) described in SEQ ID NO: 852, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6816-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res., Vol. 22, p. 1634-1645. Also, “hsa-mir-6816” (miRBase Accession No. MI0022661, SEQ ID NO: 859) having a hairpin-like structure is known as a precursor of “hsa-miR-6816-5p”.

(354) The term “hsa-miR-4466 gene” or “hsa-miR-4466” used herein includes the hsa-miR-4466 gene (miRBase Accession No. MIMAT0018993) described in SEQ ID NO: 853, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4466 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4466” (miRBase Accession No. MI0016817, SEQ ID NO: 860) having a hairpin-like structure is known as a precursor of “hsa-miR-4466”.

(355) The term “hsa-miR-4488 gene” or “hsa-miR-4488” used herein includes the hsa-miR-4488 gene (miRBase Accession No. MIMAT0019022) described in SEQ ID NO: 854, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-4488 gene can be obtained by a method described in Jima D D et al., 2010, Blood, Vol. 116, e118-e127. Also, “hsa-mir-4488” (miRBase Accession No. MI0016849, SEQ ID NO: 861) having a hairpin-like structure is known as a precursor of “hsa-miR-4488”.

(356) The term “hsa-miR-6752-5p gene” or “hsa-miR-6752-5p” used herein includes the hsa-miR-6752-5p gene (miRBase Accession No. MIMAT0027404) described in SEQ ID NO: 855, a homolog or an ortholog of a different organism species, and the like. The hsa-miR-6752-5p gene can be obtained by a method described in Ladewig E et al., 2012, Genome Res, Vol. 22, p. 1634-1645. Also, “hsa-mir-6752” (miRBase Accession No. MI0022597, SEQ ID NO: 862) having a hairpin-like structure is known as a precursor of “hsa-miR-6752-5p”.

(357) The term “hsa-miR-4739 gene” or “hsa-miR-4739” used herein includes the hsa-miR-4739

MIMAT0022259 220 hsa-miR-6769b-5p MIMAT0027620 221 hsa-miR-1207-5p MIMAT0005871
222 hsa-miR-642a-3p MIMAT0020924 223 hsa-miR-4505 MIMAT0019041 224 hsa-miR-4270
MIMAT0016900 225 hsa-miR-6721-5p MIMAT0025852 226 hsa-miR-7111-5p MIMAT0028119
227 hsa-miR-6791-5p MIMAT0027482 228 hsa-miR-7109-5p MIMAT0028115 229 hsa-miR-4258
MIMAT0016879 230 hsa-miR-6515-3p MIMAT0025487 231 hsa-miR-6851-5p MIMAT0027602
232 hsa-miR-6125 MIMAT0024598 233 hsa-miR-4749-5p MIMAT0019885 234 hsa-miR-4726-5p
MIMAT0019845 235 hsa-miR-4513 MIMAT0019050 236 hsa-miR-760 MIMAT0004957 237 hsa-
miR-602 MIMAT0003270 238 hsa-miR-423-5p MIMAT0004748 239 hsa-miR-92a-2-5p
MIMAT0004508 240 hsa-miR-16-5p MIMAT0000069 241 hsa-miR-451a MIMAT0001631 242
hsa-miR-135a-3p MIMAT0004595 243 hsa-miR-486-5p MIMAT0002177 244 hsa-miR-4257
MIMAT0016878 245 hsa-miR-92b-5p MIMAT0004792 246 hsa-miR-1915-3p MIMAT0007892
247 hsa-miR-718 MIMAT0012735 248 hsa-miR-940 MIMAT0004983 249 hsa-miR-296-5p
MIMAT0000690 250 hsa-miR-23b-3p MIMAT0000418 251 hsa-miR-92a-3p MIMAT0000092 252
hsa-miR-658 MIMAT0003336 253 hsa-miR-6842-5p MIMAT0027586 254 hsa-miR-6124
MIMAT0024597 255 hsa-miR-6765-3p MIMAT0027431 256 hsa-miR-7106-5p MIMAT0028109
257 hsa-miR-4534 MIMAT0019073 258 hsa-miR-92b-3p MIMAT0003218 259 hsa-miR-3135b
MIMAT0018985 260 hsa-miR-4687-3p MIMAT0019775 261 hsa-miR-762 MIMAT0010313 262
hsa-miR-3619-3p MIMAT0019219 263 hsa-miR-4467 MIMAT0018994 264 hsa-miR-557
MIMAT0003221 265 hsa-miR-1237-5p MIMAT0022946 266 hsa-miR-1908-5p MIMAT0007881
267 hsa-miR-4286 MIMAT0016916 268 hsa-miR-6885-5p MIMAT0027670 269 hsa-miR-6763-5p
MIMAT0027426 270 hsa-mir-4783 MI0017428 271 hsa-mir-4730 MI0017367 272 hsa-mir-1307
MI0006444 273 hsa-mir-4634 MI0017261 274 hsa-mir-663a MI0003672 275 hsa-mir-4532
MI0016899 276 hsa-mir-7704 MI0025240 277 hsa-mir-3178 MI0014212 278 hsa-mir-6729
MI0022574 279 hsa-mir-6090 MI0020367 280 hsa-mir-4732 MI0017369 281 hsa-mir-3184
MI0014226 282 hsa-mir-6727 MI0022572 283 hsa-mir-6088 MI0020365 284 hsa-mir-4674
MI0017305 285 hsa-mir-8073 MI0025909 286 hsa-mir-4787 MI0017434 287 hsa-mir-1469
MI0007074 288 hsa-mir-125a MI0000469 289 hsa-mir-1233-1 MI0006323 290 hsa-mir-1233-2
MI0015973 291 hsa-mir-885 MI0005560 292 hsa-mir-6802 MI0022647 293 hsa-mir-328
MI0000804 294 hsa-mir-6787 MI0022632 295 hsa-mir-8069 MI0025905 296 hsa-mir-6875
MI0022722 297 hsa-mir-1246 MI0006381 298 hsa-mir-4734 MI0017371 299 hsa-mir-6757
MI0022602 300 hsa-mir-6756 MI0022601 301 hsa-mir-3665 MI0016066 302 hsa-mir-6836
MI0022682 303 hsa-mir-6821 MI0022666 304 hsa-mir-6805 MI0022650 305 hsa-mir-4728
MI0017365 306 hsa-mir-6726 MI0022571 307 hsa-mir-197 MI0000239 308 hsa-mir-149
MI0000478 309 hsa-mir-6850 MI0022696 310 hsa-mir-4476 MI0016828 311 hsa-mir-6858
MI0022704 312 hsa-mir-564 MI0003570 313 hsa-mir-4763 MI0017404 314 hsa-mir-575
MI0003582 315 hsa-mir-6771 MI0022616 316 hsa-mir-1231 MI0006321 317 hsa-mir-1908
MI0008329 318 hsa-mir-150 MI0000479 319 hsa-mir-3937 MI0016593 320 hsa-mir-887
MI0005562 321 hsa-mir-3940 MI0016597 322 hsa-mir-4741 MI0017379 323 hsa-mir-6808
MI0022653 324 hsa-mir-6869 MI0022716 325 hsa-mir-5090 MI0017979 326 hsa-mir-615
MI0003628 327 hsa-mir-8072 MI0025908 328 hsa-mir-128-1 MI0000447 329 hsa-mir-1238
MI0006328 330 hsa-mir-365a MI0000767 331 hsa-mir-204 MI0000284 332 hsa-mir-4492
MI0016854 333 hsa-mir-6785 MI0022630 334 hsa-mir-6511a-1 MI0022223 335 hsa-mir-6511a-2
MI0023564 336 hsa-mir-6511a-3 MI0023565 337 hsa-mir-6511a-4 MI0023566 338 hsa-mir-4525
MI0016892 339 hsa-mir-1915 MI0008336 340 hsa-mir-3180-4 MI0016408 341 hsa-mir-3180-5
MI0016409 342 hsa-mir-6879 MI0022726 343 hsa-mir-1199 MI0020340 344 hsa-mir-6746
MI0022591 345 hsa-mir-711 MI0012488 346 hsa-mir-663b MI0006336 347 hsa-mir-4707
MI0017340 348 hsa-mir-6893 MI0022740 349 hsa-mir-4675 MI0017306 350 hsa-mir-4638
MI0017265 351 hsa-mir-4651 MI0017279 352 hsa-mir-6087 MI0020364 353 hsa-mir-4665
MI0017295 354 hsa-mir-4758 MI0017399 355 hsa-mir-6887 MI0022734 356 hsa-mir-3620
MI0016011 357 hsa-mir-1909 MI0008330 358 hsa-mir-7641-1 MI0024975 359 hsa-mir-7641-2

MI0021259 510 hsa-mir-4748 MI0017388 511 hsa-mir-4726 MI0017363 512 hsa-mir-4513
MI0016879 513 hsa-mir-760 MI0005567 514 hsa-mir-602 MI0003615 515 hsa-mir-423
MI0001445 516 hsa-mir-92a-2 MI0000094 517 hsa-mir-16-1 MI0000070 518 hsa-mir-16-2
MI0000115 519 hsa-mir-451a MI0001729 520 hsa-mir-135a MI0000452 521 hsa-mir-4257
MI0015856 522 hsa-mir-92b MI0003560 523 hsa-mir-718 MI0012489 524 hsa-mir-940
MI0005762 525 hsa-mir-23b MI0000439 526 hsa-mir-92a-1 MI0000093 527 hsa-mir-92a-2
MI0000094 528 hsa-mir-658 MI0003682 529 hsa-mir-6842 MI0022688 530 hsa-mir-6124
MI0021258 531 hsa-mir-6765 MI0022610 532 hsa-mir-7106 MI0022957 533 hsa-mir-4534
MI0016901 534 hsa-mir-3135b MI0016809 535 hsa-mir-4687 MI0017319 536 hsa-mir-762
MI0003892 537 hsa-mir-3619 MI0016009 538 hsa-mir-4467 MI0016818 539 hsa-mir-557
MI0003563 540 hsa-mir-1237 MI0006327 541 hsa-mir-1908 MI0008329 542 hsa-mir-4286
MI0015894 543 hsa-mir-6885 MI0022732 544 hsa-mir-6763 MI0022608 545 isomiR example 1 of
SEQ ID NO: 1 — 546 isomiR example 2 of SEQ ID NO: 1 — 547 isomiR example 1 of SEQ ID
NO: 2 — 548 isomiR example 2 of SEQ ID NO: 2 — 549 isomiR example 1 of SEQ ID NO: 3 —
550 isomiR example 2 of SEQ ID NO: 3 — 551 isomiR example 1 of SEQ ID NO: 4 — 552
isomiR example 2 of SEQ ID NO: 4 — 553 isomiR example 1 of SEQ ID NO: 5 — 554 isomiR
example 2 of SEQ ID NO: 5 — 555 isomiR example 1 of SEQ ID NO: 6 — 556 isomiR example 2
of SEQ ID NO: 6 — 557 isomiR example 1 of SEQ ID NO: 8 — 558 isomiR example 2 of SEQ ID
NO: 8 — 559 isomiR example 1 of SEQ ID NO: 11 — 560 isomiR example 2 of SEQ ID NO: 11
— 561 isomiR example 1 of SEQ ID NO: 14 — 562 isomiR example 2 of SEQ ID NO: 14 — 563
isomiR example 1 of SEQ ID NO: 15 — 564 isomiR example 2 of SEQ ID NO: 15 — 565 isomiR
example 1 of SEQ ID NO: 19 — 566 isomiR example 2 of SEQ ID NO: 19 — 567 isomiR
example 1 of SEQ ID NO: 20 — 568 isomiR example 2 of SEQ ID NO: 20 — 569 isomiR
example 1 of SEQ ID NO: 21 — 570 isomiR example 2 of SEQ ID NO: 21 — 571 isomiR
example 1 of SEQ ID NO: 23 — 572 isomiR example 2 of SEQ ID NO: 23 — 573 isomiR
example 1 of SEQ ID NO: 27 — 574 isomiR example 2 of SEQ ID NO: 27 — 575 isomiR
example 1 of SEQ ID NO: 28 — 576 isomiR example 2 of SEQ ID NO: 28 — 577 isomiR
example 1 of SEQ ID NO: 31 — 578 isomiR example 2 of SEQ ID NO: 31 — 579 isomiR
example 1 of SEQ ID NO: 35 — 580 isomiR example 2 of SEQ ID NO: 35 — 581 isomiR
example 1 of SEQ ID NO: 37 — 582 isomiR example 2 of SEQ ID NO: 37 — 583 isomiR
example 1 of SEQ ID NO: 38 — 584 isomiR example 2 of SEQ ID NO: 38 — 585 isomiR
example 1 of SEQ ID NO: 40 — 586 isomiR example 2 of SEQ ID NO: 40 — 587 isomiR
example 1 of SEQ ID NO: 42 — 588 isomiR example 2 of SEQ ID NO: 42 — 589 isomiR
example 1 of SEQ ID NO: 43 — 590 isomiR example 2 of SEQ ID NO: 43 — 591 isomiR
example 1 of SEQ ID NO: 47 — 592 isomiR example 2 of SEQ ID NO: 47 — 593 isomiR
example 1 of SEQ ID NO: 48 — 594 isomiR example 2 of SEQ ID NO: 48 — 595 isomiR
example 1 of SEQ ID NO: 50 — 596 isomiR example 2 of SEQ ID NO: 50 — 597 isomiR
example 1 of SEQ ID NO: 51 — 598 isomiR example 2 of SEQ ID NO: 51 — 599 isomiR
example 1 of SEQ ID NO: 52 — 600 isomiR example 2 of SEQ ID NO: 52 — 601 isomiR
example 1 of SEQ ID NO: 55 — 602 isomiR example 2 of SEQ ID NO: 55 — 603 isomiR
example 1 of SEQ ID NO: 56 — 604 isomiR example 2 of SEQ ID NO: 56 — 605 isomiR
example 1 of SEQ ID NO: 58 — 606 isomiR example 2 of SEQ ID NO: 58 — 607 isomiR
example 1 of SEQ ID NO: 60 — 608 isomiR example 2 of SEQ ID NO: 60 — 609 isomiR
example 1 of SEQ ID NO: 61 — 610 isomiR example 2 of SEQ ID NO: 61 — 611 isomiR example
1 of SEQ ID NO: 62 — 612 isomiR example 2 of SEQ ID NO: 62 — 613 isomiR example 1 of
SEQ ID NO: 64 — 614 isomiR example 2 of SEQ ID NO: 64 — 615 isomiR example 1 of SEQ ID
NO: 65 — 616 isomiR example 2 of SEQ ID NO: 65 — 617 isomiR example 1 of SEQ ID NO: 66
— 618 isomiR example 2 of SEQ ID NO: 66 — 619 isomiR example 1 of SEQ ID NO: 67 — 620
isomiR example 2 of SEQ ID NO: 67 — 621 isomiR example 1 of SEQ ID NO: 71 — 622 isomiR
example 2 of SEQ ID NO: 71 — 623 isomiR example 1 of SEQ ID NO: 72 — 624 isomiR

example 2 of SEQ ID NO: 247 — 825 isomiR example 1 of SEQ ID NO: 248 — 826 isomiR example 2 of SEQ ID NO: 248 — 827 isomiR example 1 of SEQ ID NO: 249 — 828 isomiR example 2 of SEQ ID NO: 249 — 829 isomiR example 1 of SEQ ID NO: 250 — 830 isomiR example 2 of SEQ ID NO: 250 — 831 isomiR example 1 of SEQ ID NO: 251 — 832 isomiR example 2 of SEQ ID NO: 251 — 833 isomiR example 1 of SEQ ID NO: 252 — 834 isomiR example 2 of SEQ ID NO: 252 — 835 isomiR example 1 of SEQ ID NO: 254 — 836 isomiR example 2 of SEQ ID NO: 254 — 837 isomiR example 1 of SEQ ID NO: 258 — 838 isomiR example 2 of SEQ ID NO: 258 — 839 isomiR example 1 of SEQ ID NO: 259 — 840 isomiR example 2 of SEQ ID NO: 259 — 841 isomiR example 1 of SEQ ID NO: 260 — 842 isomiR example 2 of SEQ ID NO: 260 — 843 isomiR example 1 of SEQ ID NO: 263 — 844 isomiR example 2 of SEQ ID NO: 263 — 845 isomiR example 1 of SEQ ID NO: 265 — 846 isomiR example 2 of SEQ ID NO: 265 — 847 isomiR example 1 of SEQ ID NO: 266 — 848 isomiR example 2 of SEQ ID NO: 266 — 849 isomiR example 1 of SEQ ID NO: 267 — 850 isomiR example 2 of SEQ ID NO: 267 — 851 hsa-miR-6089 MIMAT0023714 852 hsa-miR-6816-5p MIMAT0027532 853 hsa-miR-4466 MIMAT0018993 854 hsa-miR-4488 MIMAT0019022 855 hsa-miR-6752-5p MIMAT0027404 856 hsa-miR-4739 MIMAT0019868 857 hsa-mir-6089-1 MI0020366 858 hsa-mir-6089-2 MI0023563 859 hsa-mir-6816 MI0022661 860 hsa-mir-4466 MI0016817 861 hsa-mir-4488 MI0016849 862 hsa-mir-6752 MI0022597 863 hsa-mir-4739 MI0017377 864 isomiR example 1 of SEQ ID NO: 851 — 865 isomiR example 2 of SEQ ID NO: 851 — 866 isomiR example 1 of SEQ ID NO: 853 — 867 isomiR example 2 of SEQ ID NO: 853 — 868 isomiR example 1 of SEQ ID NO: 854 — 869 isomiR example 2 of SEQ ID NO: 854 — 870 isomiR example 1 of SEQ ID NO: 856 — 871 isomiR example 2 of SEQ ID NO: 856 — (362) The present specification encompasses the contents described in the specifications and/or drawings of Japanese Patent Application No. 2014-122672 and No. 2015-069321 from which the present application claims priorities.

Advantageous Effect of Invention

(363) According to the present invention, breast cancer can be detected easily and in high accuracy. For example, the presence or absence of breast cancer in patients can be easily detected by using, as indicators, the measurement values of several miRNAs in bloods, sera, and/or plasmas of the patients, which can be collected with limitedly invasiveness.

Description

BRIEF DESCRIPTION OF DRAWINGS

- (1) FIG. 1 This figure shows the relationship between the nucleotide sequences of hsa-miR-6805-5p represented by SEQ ID NO: 34 and hsa-miR-6805-3p represented by SEQ ID NO: 205, which are produced from a precursor hsa-mir-6805 represented by SEQ ID NO: 304.
- (2) FIG. 2 Left diagram: the expression level measurement values of hsa-miR-4783-3p (SEQ ID NO: 1) in healthy subjects (100 persons) and breast cancer patients (62 persons) selected as a training cohort were each plotted on the ordinate. The horizontal line in the diagram depicts a threshold (6.63) that was optimized by Fisher's discriminant analysis and discriminated between the two groups. Right diagram: the expression level measurement values of hsa-miR-4783-3p (SEQ ID NO: 1) in healthy subjects (50 persons) and breast cancer patients (31 persons) selected as a validation cohort were each plotted on the ordinate. The horizontal line in the diagram depicts the threshold (6.63) that was set in the training cohort and discriminated between the two groups.
- (3) FIG. 3 Left diagram: the expression level measurement values of hsa-miR-4783-3p (SEQ ID NO: 1) in healthy subjects (100 persons, circles) and breast cancer patients (62 persons, triangles) selected as a training cohort were each plotted on the abscissa against their expression level measurement values of hsa-miR-4730 (SEQ ID NO: 2) on the ordinate. The line in the diagram

depicts a discriminant function ($0=1.41x+y+0.77$) that was optimized by Fisher's discriminant analysis and discriminated between the two groups. Right diagram: the expression level measurement values of hsa-miR-4783-3p (SEQ ID NO: 1) in healthy subjects (50 persons, circles) and breast cancer patients (31 persons, triangles) selected as a validation cohort were each plotted on the abscissa against their expression level measurement values of hsa-miR-4730 (SEQ ID NO: 2) on the ordinate. The line in the diagram depicts the threshold ($0=1.41x+y+0.77$) that was set for the training cohort and discriminated between the two groups.

(4) FIG. 4 Upper diagram: a discriminant ($1.87 \times \text{hsa-miR-4730} + 0.42 \times \text{hsa-miR-602} - 18.58$) was prepared by use of Fisher's discriminant analysis from the expression level measurement values of hsa-miR-602 (SEQ ID NO: 237) and hsa-miR-4730 (SEQ ID NO: 2) in 62 breast cancer patients, 102 healthy subjects, and 33 prostate cancer patients selected as a training cohort, and discriminant scores obtained from the discriminant were plotted on the ordinate against the sample groups on the abscissa. The dotted line in the diagram depicts a discriminant boundary that offered a discriminant score of 0 and discriminated between the groups. Lower diagram: discriminant scores obtained from the discriminant prepared for the training cohort as to the expression level measurement values of hsa-miR-602 (SEQ ID NO: 237) and hsa-miR-4730 (SEQ ID NO: 2) in 31 breast cancer patients, 48 healthy subjects, and 19 prostate cancer patients selected as a validation cohort were plotted on the ordinate against the sample groups on the abscissa. The dotted line in the diagram depicts the discriminant boundary that offered a discriminant score of 0 and discriminated between the two groups.

DESCRIPTION OF EMBODIMENTS

(5) Hereinafter, the present invention will be further described in detail.

(6) 1. Target Nucleic Acid for Breast Cancer

(7) Primary target nucleic acids as breast cancer markers for detecting the presence and/or absence of breast cancer or breast cancer cells using the nucleic acid probes or the primers for the detection of breast cancer defined above according to the present invention, at least one miRNA selected from the group consisting of the following miRNAs: hsa-miR-4783-3p, hsa-miR-4730, hsa-miR-1307-3p, hsa-miR-4634, hsa-miR-663a, hsa-miR-4532, hsa-miR-7704, hsa-miR-3178, hsa-miR-6729-5p, hsa-miR-6090, hsa-miR-4732-5p, hsa-miR-3184-5p, hsa-miR-6727-5p, hsa-miR-6088, hsa-miR-4674, hsa-miR-8073, hsa-miR-4787-5p, hsa-miR-1469, hsa-miR-125a-3p, hsa-miR-1233-5p, hsa-miR-885-3p, hsa-miR-6802-5p, hsa-miR-328-5p, hsa-miR-6787-5p, hsa-miR-8069, hsa-miR-6875-5p, hsa-miR-1246, hsa-miR-4734, hsa-miR-6757-5p, hsa-miR-6756-5p, hsa-miR-3665, hsa-miR-6836-3p, hsa-miR-6821-5p, hsa-miR-6805-5p, hsa-miR-4728-5p, hsa-miR-6726-5p, hsa-miR-197-5p, hsa-miR-149-3p, hsa-miR-6850-5p, hsa-miR-4476, hsa-miR-6858-5p, hsa-miR-564, hsa-miR-4763-3p, hsa-miR-575, hsa-miR-6771-5p, hsa-miR-1231, hsa-miR-1908-3p, hsa-miR-150-3p, hsa-miR-3937, hsa-miR-887-3p, hsa-miR-3940-5p, hsa-miR-4741, hsa-miR-6808-5p, hsa-miR-6869-5p, hsa-miR-5090, hsa-miR-615-5p, hsa-miR-8072, hsa-miR-128-1-5p, hsa-miR-1238-5p, hsa-miR-365a-5p, hsa-miR-204-3p, hsa-miR-4492, hsa-miR-6785-5p, hsa-miR-6511a-5p, hsa-miR-4525, hsa-miR-1915-5p, hsa-miR-3180, hsa-miR-6879-5p, hsa-miR-1199-5p, hsa-miR-6746-5p, hsa-miR-711, hsa-miR-663b, hsa-miR-4707-3p, hsa-miR-6893-5p, hsa-miR-4675, hsa-miR-4638-5p, hsa-miR-4651, hsa-miR-6087, hsa-miR-4665-5p, hsa-miR-4758-5p, hsa-miR-6887-5p, hsa-miR-3620-5p, hsa-miR-1909-3p, hsa-miR-7641, hsa-miR-6724-5p, hsa-miR-1343-3p, hsa-miR-6780b-5p, hsa-miR-4484, hsa-miR-4690-5p, hsa-miR-4429, hsa-miR-1227-5p, hsa-miR-4725-3p, hsa-miR-6861-5p, hsa-miR-6812-5p, hsa-miR-3197, hsa-miR-8059, hsa-miR-3185, hsa-miR-4706, hsa-miR-4497, hsa-miR-3131, hsa-miR-6806-5p, hsa-miR-187-5p, hsa-miR-3180-3p, hsa-miR-6848-5p, hsa-miR-6820-5p, hsa-miR-6800-5p, hsa-miR-6717-5p, hsa-miR-6795-5p, hsa-miR-4632-5p, hsa-miR-665, hsa-miR-6778-5p, hsa-miR-3663-3p, hsa-miR-4689, hsa-miR-211-3p, hsa-miR-6511b-5p, hsa-miR-4750-5p, hsa-miR-6126, hsa-miR-614, hsa-miR-7110-5p, hsa-miR-744-5p, hsa-miR-6769a-5p, hsa-miR-4792, hsa-miR-5787, hsa-miR-6798-5p, hsa-miR-6781-5p, hsa-miR-4419b, hsa-miR-4446-3p, hsa-miR-4259, hsa-miR-5572, hsa-miR-6075, hsa-miR-296-3p,

described above).

(247) The 238th target gene is the hsa-miR-423-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Non-Patent Literature 4 described above).

(248) The 239th target gene is the hsa-miR-92a-2-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Patent Literature 3 described above).

(249) The 240th target gene is the hsa-miR-16-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Patent Literature 4 described above).

(250) The 241st target gene is the hsa-miR-451a gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Patent Literature 4 described above).

(251) The 242nd target gene is the hsa-miR-135a-3p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Patent Literatures 1 and 2 described above).

(252) The 243rd target gene is the hsa-miR-486-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Non-Patent Literature 4 described above).

(253) The 244th target gene is the hsa-miR-4257 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Non-Patent Literature 5 described above).

(254) The 245th target gene is the hsa-miR-92b-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Patent Literature 3 described above).

(255) The 246th target gene is the hsa-miR-1915-3p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Non-Patent Literature 5 described above).

(256) The 247th target gene is the hsa-miR-718 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Non-Patent Literature 5 described above).

(257) The 248th target gene is the hsa-miR-940 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Non-Patent Literature 6 described above).

(258) The 249th target gene is the hsa-miR-296-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Patent Literature 4 described above).

(259) The 250th target gene is the hsa-miR-23b-3p gene, a congener thereof, a transcript thereof, or

a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Patent Literature 2 described above).

(260) The 251st target gene is the hsa-miR-92a-3p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. The previously known report shows that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer (Patent Literature 3 described above).

(261) The 252nd target gene is the hsa-miR-658 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(262) The 253rd target gene is the hsa-miR-6842-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(263) The 254th target gene is the hsa-miR-6124 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(264) The 255th target gene is the hsa-miR-6765-3p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(265) The 256th target gene is the hsa-miR-7106-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(266) The 257th target gene is the hsa-miR-4534 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(267) The 258th target gene is the hsa-miR-92b-3p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(268) The 259th target gene is the hsa-miR-3135b gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(269) The 260th target gene is the hsa-miR-4687-3p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(270) The 261st target gene is the hsa-miR-762 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(271) The 262nd target gene is the hsa-miR-3619-3p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(272) The 263rd target gene is the hsa-miR-4467 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(273) The 264th target gene is the hsa-miR-557 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(274) The 265th target gene is the hsa-miR-1237-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(275) The 266th target gene is the hsa-miR-1908-5p gene, a congener thereof, a transcript thereof,

or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(276) The 267th target gene is the hsa-miR-4286 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(277) The 268th target gene is the hsa-miR-6885-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(278) The 269th target gene is the hsa-miR-6763-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(279) The 270th target gene is the hsa-miR-6089 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(280) The 271st target gene is the hsa-miR-6816-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(281) The 272nd target gene is the hsa-miR-4466 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(282) The 273rd target gene is the hsa-miR-4488 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(283) The 274th target gene is the hsa-miR-6752-5p gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(284) The 275th target gene is the hsa-miR-4739 gene, a congener thereof, a transcript thereof, or a variant or a derivative thereof. None of the previously known reports show that change in the expression of the gene or the transcript thereof can serve as a marker for breast cancer.

(285) 2. Nucleic Acid Probe or Primer for Detection of Breast Cancer

(286) In the present invention, nucleic acid(s) capable of specifically binding to any of the target nucleic acids as the breast cancer markers described above can be used as a nucleic acid, for example, a nucleic acid probe or a primer, for the detection or diagnosis of breast cancer.

(287) In the present invention, the nucleic acid probes or the primers that can be used for detecting breast cancer or for diagnosing breast cancer enables qualitative and/or quantitative measurement of the presence, expression level, or existing amount (abundance) of any of the target nucleic acids as the breast cancer markers described above, for example, human-derived hsa-miR-4783-3p, hsa-miR-4730, hsa-miR-1307-3p, hsa-miR-4634, hsa-miR-663a, hsa-miR-4532, hsa-miR-7704, hsa-miR-3178, hsa-miR-6729-5p, hsa-miR-6090, hsa-miR-4732-5p, hsa-miR-3184-5p, hsa-miR-6727-5p, hsa-miR-6088, hsa-miR-4674, hsa-miR-8073, hsa-miR-4787-5p, hsa-miR-1469, hsa-miR-125a-3p, hsa-miR-1233-5p, hsa-miR-885-3p, hsa-miR-6802-5p, hsa-miR-328-5p, hsa-miR-6787-5p, hsa-miR-8069, hsa-miR-6875-5p, hsa-miR-1246, hsa-miR-4734, hsa-miR-6757-5p, hsa-miR-6756-5p, hsa-miR-3665, hsa-miR-6836-3p, hsa-miR-6821-5p, hsa-miR-6805-5p, hsa-miR-4728-5p, hsa-miR-6726-5p, hsa-miR-197-5p, hsa-miR-149-3p, hsa-miR-6850-5p, hsa-miR-4476, hsa-miR-6858-5p, hsa-miR-564, hsa-miR-4763-3p, hsa-miR-575, hsa-miR-6771-5p, hsa-miR-1231, hsa-miR-1908-3p, hsa-miR-150-3p, hsa-miR-3937, hsa-miR-887-3p, hsa-miR-3940-5p, hsa-miR-4741, hsa-miR-6808-5p, hsa-miR-6869-5p, hsa-miR-5090, hsa-miR-615-5p, hsa-miR-8072, hsa-miR-128-1-5p, hsa-miR-1238-5p, hsa-miR-365a-5p, hsa-miR-204-3p, hsa-miR-4492, hsa-miR-6785-5p, hsa-miR-6511a-5p, hsa-miR-4525, hsa-miR-1915-5p, hsa-miR-3180, hsa-miR-6879-5p, hsa-miR-1199-5p, hsa-miR-6746-5p, hsa-miR-711, hsa-miR-663b, hsa-miR-4707-3p, hsa-miR-

6893-5p, hsa-miR-4675, hsa-miR-4638-5p, hsa-miR-4651, hsa-miR-6087, hsa-miR-4665-5p, hsa-miR-4758-5p, hsa-miR-6887-5p, hsa-miR-3620-5p, hsa-miR-1909-3p, hsa-miR-7641, hsa-miR-6724-5p, hsa-miR-1343-3p, hsa-miR-6780b-5p, hsa-miR-4484, hsa-miR-4690-5p, hsa-miR-4429, hsa-miR-1227-5p, hsa-miR-4725-3p, hsa-miR-6861-5p, hsa-miR-6812-5p, hsa-miR-3197, hsa-miR-8059, hsa-miR-3185, hsa-miR-4706, hsa-miR-4497, hsa-miR-3131, hsa-miR-6806-5p, hsa-miR-187-5p, hsa-miR-3180-3p, hsa-miR-6848-5p, hsa-miR-6820-5p, hsa-miR-6800-5p, hsa-miR-6717-5p, hsa-miR-6795-5p, hsa-miR-4632-5p, hsa-miR-665, hsa-miR-6778-5p, hsa-miR-3663-3p, hsa-miR-4689, hsa-miR-211-3p, hsa-miR-6511b-5p, hsa-miR-4750-5p, hsa-miR-6126, hsa-miR-614, hsa-miR-7110-5p, hsa-miR-744-5p, hsa-miR-6769a-5p, hsa-miR-4792, hsa-miR-5787, hsa-miR-6798-5p, hsa-miR-6781-5p, hsa-miR-4419b, hsa-miR-4446-3p, hsa-miR-4259, hsa-miR-5572, hsa-miR-6075, hsa-miR-296-3p, hsa-miR-6891-5p, hsa-miR-4745-5p, hsa-miR-6775-5p, hsa-miR-6870-5p, hsa-miR-920, hsa-miR-4530, hsa-miR-6819-5p, hsa-miR-6825-5p, hsa-miR-7847-3p, hsa-miR-6131, hsa-miR-4433-3p, hsa-miR-1228-5p, hsa-miR-6743-5p, hsa-miR-1268a, hsa-miR-3917, hsa-miR-6786-5p, hsa-miR-3154, hsa-miR-638, hsa-miR-6741-5p, hsa-miR-6889-5p, hsa-miR-6840-3p, hsa-miR-6510-5p, hsa-miR-3188, hsa-miR-551b-5p, hsa-miR-5001-5p, hsa-miR-1268b, hsa-miR-7107-5p, hsa-miR-6824-5p, hsa-miR-6732-5p, hsa-miR-371a-5p, hsa-miR-6794-5p, hsa-miR-6779-5p, hsa-miR-4271, hsa-miR-5195-3p, hsa-miR-6762-5p, hsa-miR-939-5p, hsa-miR-1247-3p, hsa-miR-6777-5p, hsa-miR-6722-3p, hsa-miR-3656, hsa-miR-4688, hsa-miR-3195, hsa-miR-6766-5p, hsa-miR-4447, hsa-miR-4656, hsa-miR-7108-5p, hsa-miR-3191-3p, hsa-miR-1273g-3p, hsa-miR-4463, hsa-miR-2861, hsa-miR-3196, hsa-miR-6877-5p, hsa-miR-3679-5p, hsa-miR-4442, hsa-miR-6789-5p, hsa-miR-6782-5p, hsa-miR-486-3p, hsa-miR-6085, hsa-miR-4746-3p, hsa-miR-619-5p, hsa-miR-937-5p, hsa-miR-6803-5p, hsa-miR-4298, hsa-miR-4454, hsa-miR-4459, hsa-miR-7150, hsa-miR-6880-5p, hsa-miR-4449, hsa-miR-8063, hsa-miR-4695-5p, hsa-miR-6132, hsa-miR-6829-5p, hsa-miR-4486, hsa-miR-6805-3p, hsa-miR-6826-5p, hsa-miR-4508, hsa-miR-1343-5p, hsa-miR-7114-5p, hsa-miR-3622a-5p, hsa-miR-6765-5p, hsa-miR-7845-5p, hsa-miR-3960, hsa-miR-6749-5p, hsa-miR-1260b, hsa-miR-6799-5p, hsa-miR-4723-5p, hsa-miR-6784-5p, hsa-miR-5100, hsa-miR-6769b-5p, hsa-miR-1207-5p, hsa-miR-642a-3p, hsa-miR-4505, hsa-miR-4270, hsa-miR-6721-5p, hsa-miR-7111-5p, hsa-miR-6791-5p, hsa-miR-7109-5p, hsa-miR-4258, hsa-miR-6515-3p, hsa-miR-6851-5p, hsa-miR-6125, hsa-miR-4749-5p, hsa-miR-4726-5p, hsa-miR-4513, hsa-miR-6089, hsa-miR-6816-5p, hsa-miR-4466, hsa-miR-4488, hsa-miR-6752-5p and hsa-miR-4739 or a combination thereof, congeners thereof, transcripts thereof, or variants or derivatives thereof; and, optionally in combination therewith, hsa-miR-760, hsa-miR-602, hsa-miR-423-5p, hsa-miR-92a-2-5p, hsa-miR-16-5p, hsa-miR-451a, hsa-miR-135a-3p, hsa-miR-486-5p, hsa-miR-4257, hsa-miR-92b-5p, hsa-miR-1915-3p, hsa-miR-718, hsa-miR-940, hsa-miR-296-5p, hsa-miR-23b-3p and hsa-miR-92a-3p or a combination thereof, transcripts thereof, or variants or derivatives thereof; and, optionally in combination therewith, hsa-miR-658, hsa-miR-6842-5p, hsa-miR-6124, hsa-miR-6765-3p, hsa-miR-7106-5p, hsa-miR-4534, hsa-miR-92b-3p, hsa-miR-3135b, hsa-miR-4687-3p, hsa-miR-762, hsa-miR-3619-3p, hsa-miR-4467, hsa-miR-557, hsa-miR-1237-5p, hsa-miR-1908-5p, hsa-miR-4286, hsa-miR-6885-5p and hsa-miR-6763-5p or a combination thereof; congeners thereof; transcripts thereof; or variants or derivatives thereof.

(288) The expression level of the target nucleic acids described above are increased or decreased (hereinafter, referred to as “increased/decreased”) depending on the types of the target nucleic acids in subjects who have breast cancer, as compared with healthy subjects. Hence, the nucleic acid of the present invention can be effectively used for measuring expression levels of the target nucleic acids in body fluids derived from subjects (e.g., humans) suspected of having breast cancer and body fluids derived from healthy subjects and thereby detecting breast cancer through the comparison thereof.

(289) The nucleic acid probe or the primer that can be used in the present invention is a nucleic acid probe capable of specifically binding to a polynucleotide consisting of a nucleotide sequence represented by at least one of SEQ ID NOs: 1 to 235 and 851 to 856, or a primer for amplifying a

polynucleotide consisting of a nucleotide sequence represented by at least one of SEQ ID NOs: 1 to 235 and 851 to 856.

(290) The nucleic acid probe or the primer that can be used in the present invention may further comprise a nucleic acid probe capable of specifically binding to a polynucleotide consisting of a nucleotide sequence represented by at least one of SEQ ID NOs: 236 to 251, or a primer for amplifying a polynucleotide consisting of a nucleotide sequence represented by at least one of SEQ ID NOs: 236 to 251.

(291) The nucleic acid probe or the primer that can be used in the present invention may further comprise a nucleic acid probe capable of specifically binding to a polynucleotide that consists of a nucleotide sequence represented by at least one of SEQ ID NOs: 252 to 269, or a primer for amplifying a polynucleotide that consists of a nucleotide sequence represented by at least one of SEQ ID NOs: 252 to 269.

(292) Specifically, these nucleic acid probes or primers comprise a combination of one or more polynucleotides selected from: a group of polynucleotides that comprise nucleotide sequences represented by any of SEQ ID NOs: 1 to 871 or nucleotide sequences derived from the nucleotide sequences by the replacement of u with t, and a group of complementary polynucleotides thereof, a group of polynucleotides that respectively hybridize under stringent conditions (mentioned later) to DNAs consisting of nucleotide sequences complementary to these nucleotide sequences; and a group of complementary polynucleotides thereof, and a group of polynucleotide comprising 15 or more, preferably 17 or more consecutive nucleotides in the nucleotide sequences of these polynucleotides. These polynucleotides can be used as nucleic acid probes and primers for detecting the breast cancer markers as target nucleic acids.

(293) More specifically, examples of the nucleic acid probes or the primers that can be used in the present invention include one or more polynucleotides selected from the group consisting of the following polynucleotides (a) to (e): (a) a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235, and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (b) a polynucleotide comprising a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235, and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, (c) a polynucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235, and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (d) a polynucleotide comprising a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235, and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, and (e) a polynucleotide hybridizing under stringent conditions to any of the polynucleotides (a) to (d).

(294) In addition to at least one polynucleotide selected from the group consisting of the polynucleotides (a) to (e), the nucleic acid probes or the primers that can be used in the present invention may further comprise a polynucleotide selected from the group consisting of the following polynucleotides (f) to (j): (f) a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (g) a polynucleotide comprising a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251, (h) a polynucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (i) a polynucleotide comprising a nucleotide sequence

complementary to a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, and (j) a polynucleotide hybridizing under stringent conditions to any of the polynucleotides (f) to (i). (295) In addition to at least one polynucleotide selected from the group consisting of (a) to (j) described above, the nucleic acid probes or the primers that can be used in the present invention may further comprise a polynucleotide selected from the group consisting of the following polynucleotides (k) to (o): (k) a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (l) a polynucleotide comprising a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269, (m) a polynucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (n) a polynucleotide comprising a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, and (o) a polynucleotide hybridizing under stringent conditions to any of the polynucleotides (k) to (n).

(296) For these polynucleotides, the “fragment thereof comprising 15 or more consecutive nucleotides” can comprise, but is not limited to, the number of nucleotides in the range from, for example, 15 consecutive nucleotides to less than the total number of nucleotides of the sequence, from 17 consecutive nucleotides to less than the total number of nucleotides of the sequence, or from 19 consecutive nucleotides to less than the total number of nucleotides of the sequence, in the nucleotide sequence of each polynucleotide.

(297) These polynucleotides or fragments thereof used in the present invention may each be DNA or may each be RNA.

(298) The polynucleotides that can be used in the present invention can each be prepared by use of a general technique such as a DNA recombination technique, a PCR method, or a method using an automatic DNA/RNA synthesizer.

(299) The DNA recombination technique and the PCR method may employ techniques described in, for example, Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, US (1993); and Sambrook et al., *Molecular Cloning—A Laboratory Manual*, Cold Spring Harbor Laboratory Press, US (1989).

(300) The human-derived hsa-miR-4783-3p, hsa-miR-4730, hsa-miR-1307-3p, hsa-miR-4634, hsa-miR-663a, hsa-miR-4532, hsa-miR-7704, hsa-miR-3178, hsa-miR-6729-5p, hsa-miR-6090, hsa-miR-4732-5p, hsa-miR-3184-5p, hsa-miR-6727-5p, hsa-miR-6088, hsa-miR-4674, hsa-miR-8073, hsa-miR-4787-5p, hsa-miR-1469, hsa-miR-125a-3p, hsa-miR-1233-5p, hsa-miR-885-3p, hsa-miR-6802-5p, hsa-miR-328-5p, hsa-miR-6787-5p, hsa-miR-8069, hsa-miR-6875-5p, hsa-miR-1246, hsa-miR-4734, hsa-miR-6757-5p, hsa-miR-6756-5p, hsa-miR-3665, hsa-miR-6836-3p, hsa-miR-6821-5p, hsa-miR-6805-5p, hsa-miR-4728-5p, hsa-miR-6726-5p, hsa-miR-197-5p, hsa-miR-149-3p, hsa-miR-6850-5p, hsa-miR-4476, hsa-miR-6858-5p, hsa-miR-564, hsa-miR-4763-3p, hsa-miR-575, hsa-miR-6771-5p, hsa-miR-1231, hsa-miR-1908-3p, hsa-miR-150-3p, hsa-miR-3937, hsa-miR-887-3p, hsa-miR-3940-5p, hsa-miR-4741, hsa-miR-6808-5p, hsa-miR-6869-5p, hsa-miR-5090, hsa-miR-615-5p, hsa-miR-8072, hsa-miR-128-1-5p, hsa-miR-1238-5p, hsa-miR-365a-5p, hsa-miR-204-3p, hsa-miR-4492, hsa-miR-6785-5p, hsa-miR-6511a-5p, hsa-miR-4525, hsa-miR-1915-5p, hsa-miR-3180, hsa-miR-6879-5p, hsa-miR-1199-5p, hsa-miR-6746-5p, hsa-miR-711, hsa-miR-663b, hsa-miR-4707-3p, hsa-miR-6893-5p, hsa-miR-4675, hsa-miR-4638-5p, hsa-miR-4651, hsa-miR-6087, hsa-miR-4665-5p, hsa-miR-4758-5p, hsa-miR-6887-5p, hsa-miR-3620-5p, hsa-miR-1909-3p, hsa-miR-7641, hsa-miR-6724-5p, hsa-miR-1343-3p, hsa-miR-6780b-5p, hsa-miR-4484, hsa-miR-4690-5p, hsa-miR-4429, hsa-miR-1227-5p, hsa-miR-4725-3p, hsa-miR-6861-

5p, hsa-miR-6812-5p, hsa-miR-3197, hsa-miR-8059, hsa-miR-3185, hsa-miR-4706, hsa-miR-4497, hsa-miR-3131, hsa-miR-6806-5p, hsa-miR-187-5p, hsa-miR-3180-3p, hsa-miR-6848-5p, hsa-miR-6820-5p, hsa-miR-6800-5p, hsa-miR-6717-5p, hsa-miR-6795-5p, hsa-miR-4632-5p, hsa-miR-665, hsa-miR-6778-5p, hsa-miR-3663-3p, hsa-miR-4689, hsa-miR-211-3p, hsa-miR-6511b-5p, hsa-miR-4750-5p, hsa-miR-6126, hsa-miR-614, hsa-miR-7110-5p, hsa-miR-744-5p, hsa-miR-6769a-5p, hsa-miR-4792, hsa-miR-5787, hsa-miR-6798-5p, hsa-miR-6781-5p, hsa-miR-4419b, hsa-miR-4446-3p, hsa-miR-4259, hsa-miR-5572, hsa-miR-6075, hsa-miR-296-3p, hsa-miR-6891-5p, hsa-miR-4745-5p, hsa-miR-6775-5p, hsa-miR-6870-5p, hsa-miR-920, hsa-miR-4530, hsa-miR-6819-5p, hsa-miR-6825-5p, hsa-miR-7847-3p, hsa-miR-6131, hsa-miR-4433-3p, hsa-miR-1228-5p, hsa-miR-6743-5p, hsa-miR-1268a, hsa-miR-3917, hsa-miR-6786-5p, hsa-miR-3154, hsa-miR-638, hsa-miR-6741-5p, hsa-miR-6889-5p, hsa-miR-6840-3p, hsa-miR-6510-5p, hsa-miR-3188, hsa-miR-551b-5p, hsa-miR-5001-5p, hsa-miR-1268b, hsa-miR-7107-5p, hsa-miR-6824-5p, hsa-miR-6732-5p, hsa-miR-371a-5p, hsa-miR-6794-5p, hsa-miR-6779-5p, hsa-miR-4271, hsa-miR-5195-3p, hsa-miR-6762-5p, hsa-miR-939-5p, hsa-miR-1247-3p, hsa-miR-6777-5p, hsa-miR-6722-3p, hsa-miR-3656, hsa-miR-4688, hsa-miR-3195, hsa-miR-6766-5p, hsa-miR-4447, hsa-miR-4656, hsa-miR-7108-5p, hsa-miR-3191-3p, hsa-miR-1273g-3p, hsa-miR-4463, hsa-miR-2861, hsa-miR-3196, hsa-miR-6877-5p, hsa-miR-3679-5p, hsa-miR-4442, hsa-miR-6789-5p, hsa-miR-6782-5p, hsa-miR-486-3p, hsa-miR-6085, hsa-miR-4746-3p, hsa-miR-619-5p, hsa-miR-937-5p, hsa-miR-6803-5p, hsa-miR-4298, hsa-miR-4454, hsa-miR-4459, hsa-miR-7150, hsa-miR-6880-5p, hsa-miR-4449, hsa-miR-8063, hsa-miR-4695-5p, hsa-miR-6132, hsa-miR-6829-5p, hsa-miR-4486, hsa-miR-6805-3p, hsa-miR-6826-5p, hsa-miR-4508, hsa-miR-1343-5p, hsa-miR-7114-5p, hsa-miR-3622a-5p, hsa-miR-6765-5p, hsa-miR-7845-5p, hsa-miR-3960, hsa-miR-6749-5p, hsa-miR-1260b, hsa-miR-6799-5p, hsa-miR-4723-5p, hsa-miR-6784-5p, hsa-miR-5100, hsa-miR-6769b-5p, hsa-miR-1207-5p, hsa-miR-642a-3p, hsa-miR-4505, hsa-miR-4270, hsa-miR-6721-5p, hsa-miR-7111-5p, hsa-miR-6791-5p, hsa-miR-7109-5p, hsa-miR-4258, hsa-miR-6515-3p, hsa-miR-6851-5p, hsa-miR-6125, hsa-miR-4749-5p, hsa-miR-4726-5p, hsa-miR-4513, hsa-miR-6089, hsa-miR-6816-5p, hsa-miR-4466, hsa-miR-4488, hsa-miR-6752-5p, hsa-miR-4739, hsa-miR-760, hsa-miR-602, hsa-miR-423-5p, hsa-miR-92a-2-5p, hsa-miR-16-5p, hsa-miR-451a, hsa-miR-135a-3p, hsa-miR-486-5p, hsa-miR-4257, hsa-miR-92b-5p, hsa-miR-1915-3p, hsa-miR-718, hsa-miR-940, hsa-miR-296-5p, hsa-miR-23b-3p, hsa-miR-92a-3p, hsa-miR-658, hsa-miR-6842-5p, hsa-miR-6124, hsa-miR-6765-3p, hsa-miR-7106-5p, hsa-miR-4534, hsa-miR-92b-3p, hsa-miR-3135b, hsa-miR-4687-3p, hsa-miR-762, hsa-miR-3619-3p, hsa-miR-4467, hsa-miR-557, hsa-miR-1237-5p, hsa-miR-1908-5p, hsa-miR-4286, hsa-miR-6885-5p and hsa-miR-6763-5p represented by SEQ ID NOs: 1 to 269, and 851 to 856 are known in the art, and their acquisition methods are also known as mentioned above. Therefore, each polynucleotide that can be used as a nucleic acid probe or a primer in the present invention can be prepared by cloning the gene.

(301) Such nucleic acid probes or primers can be chemically synthesized using an automatic DNA synthesizer. In general, the phosphoramidite method is used in this synthesis, and single-stranded DNA up to approximately 100 nucleotides can be automatically synthesized by this method. The automatic DNA synthesizer is commercially available from, for example, Polygen GmbH, ABI, or Applied Biosystems, Inc.

(302) Alternatively, the polynucleotides of the present invention can also be prepared by a cDNA cloning methods. The cDNA cloning technique may employ, for example, microRNA Cloning Kit Wako.

(303) In this context, the sequences of the nucleic acid probes and the primers for detecting the polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 1 to 269, and 851 to 856 do not exist as miRNAs or precursors thereof in the living body or in vivo. For example, the nucleotide sequences represented by SEQ ID NO: 34 and SEQ ID NO: 205 are produced from the precursor represented by SEQ ID NO: 304. This precursor has a hairpin-like structure as shown in FIG. 1, and the nucleotide sequences represented by SEQ ID NO: 34 and

SEQ ID NO: 205 have mismatch sequences with each other. As such, a nucleotide sequence completely complementary to the nucleotide sequence represented by SEQ ID NO: 34 or SEQ ID NO: 205 is not naturally produced in vivo. Likewise, the nucleic acid probe and the primer for detecting the nucleotide sequence represented by any of SEQ ID NOs: 1 to 269, and 851 to 856 have artificial nucleotide sequences that do not exist in the living body or in vivo.

(304) 3. Kit or Device for Detection of Breast Cancer

(305) The present invention also provides a kit or a device for the detection of breast cancer, comprising one or more polynucleotides (which may include a variant, a fragment, or a derivative thereof; hereinafter, also referred to as polynucleotide for detection) that can be used as nucleic acid probes or primers in the present invention for measuring target nucleic acids as breast cancer markers.

(306) The target nucleic acid as a breast cancer marker according to the present invention is selected from the following group 1: miR-4783-3p, miR-4730, miR-1307-3p, miR-4634, miR-663a, miR-4532, miR-7704, miR-3178, miR-6729-5p, miR-6090, miR-4732-5p, miR-3184-5p, miR-6727-5p, miR-6088, miR-4674, miR-8073, miR-4787-5p, miR-1469, miR-125a-3p, miR-1233-5p, miR-885-3p, miR-6802-5p, miR-328-5p, miR-6787-5p, miR-8069, miR-6875-5p, miR-1246, miR-4734, miR-6757-5p, miR-6756-5p, miR-3665, miR-6836-3p, miR-6821-5p, miR-6805-5p, miR-4728-5p, miR-6726-5p, miR-197-5p, miR-149-3p, miR-6850-5p, miR-4476, miR-6858-5p, miR-564, miR-4763-3p, miR-575, miR-6771-5p, miR-1231, miR-1908-3p, miR-150-3p, miR-3937, miR-887-3p, miR-3940-5p, miR-4741, miR-6808-5p, miR-6869-5p, miR-5090, miR-615-5p, miR-8072, miR-128-1-5p, miR-1238-5p, miR-365a-5p, miR-204-3p, miR-4492, miR-6785-5p, miR-6511a-5p, miR-4525, miR-1915-5p, miR-3180, miR-6879-5p, miR-1199-5p, miR-6746-5p, miR-711, miR-663b, miR-4707-3p, miR-6893-5p, miR-4675, miR-4638-5p, miR-4651, miR-6087, miR-4665-5p, miR-4758-5p, miR-6887-5p, miR-3620-5p, miR-1909-3p, miR-7641, miR-6724-5p, miR-1343-3p, miR-6780b-5p, miR-4484, miR-4690-5p, miR-4429, miR-1227-5p, miR-4725-3p, miR-6861-5p, miR-6812-5p, miR-3197, miR-8059, miR-3185, miR-4706, miR-4497, miR-3131, miR-6806-5p, miR-187-5p, miR-3180-3p, miR-6848-5p, miR-6820-5p, miR-6800-5p, miR-6717-5p, miR-6795-5p, miR-4632-5p, miR-665, miR-6778-5p, miR-3663-3p, miR-4689, miR-211-3p, miR-6511b-5p, miR-4750-5p, miR-6126, miR-614, miR-7110-5p, miR-744-5p, miR-6769a-5p, miR-4792, miR-5787, miR-6798-5p, miR-6781-5p, miR-4419b, miR-4446-3p, miR-4259, miR-5572, miR-6075, miR-296-3p, miR-6891-5p, miR-4745-5p, miR-6775-5p, miR-6870-5p, miR-920, miR-4530, miR-6819-5p, miR-6825-5p, miR-7847-3p, miR-6131, miR-4433-3p, miR-1228-5p, miR-6743-5p, miR-1268a, miR-3917, miR-6786-5p, miR-3154, miR-638, miR-6741-5p, miR-6889-5p, miR-6840-3p, miR-6510-5p, miR-3188, miR-551b-5p, miR-5001-5p, miR-1268b, miR-7107-5p, miR-6824-5p, miR-6732-5p, miR-371a-5p, miR-6794-5p, miR-6779-5p, miR-4271, miR-5195-3p, miR-6762-5p, miR-939-5p, miR-1247-3p, miR-6777-5p, miR-6722-3p, miR-3656, miR-4688, miR-3195, miR-6766-5p, miR-4447, miR-4656, miR-7108-5p, miR-3191-3p, miR-1273g-3p, miR-4463, miR-2861, miR-3196, miR-6877-5p, miR-3679-5p, miR-4442, miR-6789-5p, miR-6782-5p, miR-486-3p, miR-6085, miR-4746-3p, miR-619-5p, miR-937-5p, miR-6803-5p, miR-4298, miR-4454, miR-4459, miR-7150, miR-6880-5p, miR-4449, miR-8063, miR-4695-5p, miR-6132, miR-6829-5p, miR-4486, miR-6805-3p, miR-6826-5p, miR-4508, miR-1343-5p, miR-7114-5p, miR-3622a-5p, miR-6765-5p, miR-7845-5p, miR-3960, miR-6749-5p, miR-1260b, miR-6799-5p, miR-4723-5p, miR-6784-5p, miR-5100, miR-6769b-5p, miR-1207-5p, miR-642a-3p, miR-4505, miR-4270, miR-6721-5p, miR-7111-5p, miR-6791-5p, miR-7109-5p, miR-4258, miR-6515-3p, miR-6851-5p, miR-6125, miR-4749-5p, miR-4726-5p, miR-4513, miR-6089, miR-6816-5p, miR-4466, miR-4488, miR-6752-5p and miR-4739.

(307) An additional target nucleic acid(s) that may be optionally used in the measurement is selected from the following group 2: miR-760, miR-602, miR-423-5p, miR-92a-2-5p, miR-16-5p, miR-451a, miR-135a-3p, miR-486-5p, miR-4257, miR-92b-5p, miR-1915-3p, miR-718, miR-940, miR-296-5p, miR-23b-3p and miR-92a-3p.

(308) An additional target nucleic acid(s) that may be optionally used in the measurement is further selected from the following group 3: miR-658, miR-6842-5p, miR-6124, miR-6765-3p, miR-7106-5p, miR-4534, miR-92b-3p, miR-3135b, miR-4687-3p, miR-762, miR-3619-3p, miR-4467, miR-557, miR-1237-5p, miR-1908-5p, miR-4286, miR-6885-5p and miR-6763-5p.

(309) The kit or the device of the present invention comprises nucleic acid(s) capable of specifically binding to any of the target nucleic acids as the breast cancer markers described above, preferably one or more polynucleotides selected from the nucleic acid probes or the primers described in the preceding Section 2, specifically, the polynucleotides described in the preceding Section 2, or variants thereof, etc.

(310) Specifically, the kit or the device of the present invention can comprise at least one polynucleotides comprising (or consisting of) a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235, and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, polynucleotide(s) comprising (or consisting of) a complementary sequence thereof, polynucleotide(s) hybridizing under stringent conditions to any of these polynucleotides, or variant(s) or fragment(s) comprising 15 or more consecutive nucleotides of any of these polynucleotide sequences.

(311) The kit or the device of the present invention can further comprise one or more polynucleotides comprising (or consisting of) a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, polynucleotide(s) comprising (or consisting of) a complementary sequence thereof, polynucleotide(s) hybridizing under stringent conditions to any of these polynucleotides, variant(s) or fragment(s) comprising 15 or more consecutive nucleotides of any of these polynucleotide sequences.

(312) The kit or the device of the present invention can further comprise one or more polynucleotide(s) comprising (or consisting of) a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, polynucleotide(s) comprising (or consisting of) a complementary sequence thereof, polynucleotide(s) hybridizing under stringent conditions to any of these polynucleotides, variant(s) or fragment(s) comprising 15 or more consecutive nucleotides of any of these polynucleotide sequences.

(313) The fragment(s) that can be comprised in the kit or the device of the present invention is/are, for example, one or more polynucleotides, preferably two or more polynucleotides selected from the group consisting of the following polynucleotides (1) to (3): (1) a polynucleotide comprising 15 or more consecutive nucleotides in a nucleotide sequence that is derived from a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235, and 851 to 856 by the replacement of u with t, or a complementary sequence thereof; (2) a polynucleotide comprising 15 or more consecutive nucleotides in a nucleotide sequence that is derived from a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 by the replacement of u with t, or a complementary sequence thereof; and (3) a polynucleotide comprising 15 or more consecutive nucleotides in a nucleotide sequence that is derived from a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 by the replacement of u with t, or a complementary sequence thereof.

(314) In a preferred embodiment, the polynucleotide is a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235, and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a polynucleotide consisting of a complementary sequence thereof, a polynucleotide hybridizing under stringent conditions to any of these polynucleotides, or a variant thereof comprising 15 or more, preferably 17 or more, more preferably 19 or more consecutive nucleotides.

(315) In a preferred embodiment, the polynucleotide is a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a polynucleotide consisting of a

complementary sequence thereof, a polynucleotide hybridizing under stringent conditions to any of these polynucleotides, or a variant thereof comprising 15 or more, preferably 17 or more, more preferably 19 or more consecutive nucleotides.

(316) In a preferred embodiment, the polynucleotide is a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a polynucleotide consisting of a complementary sequence thereof, a polynucleotide hybridizing under stringent conditions to any of these polynucleotides, or a variant thereof comprising 15 or more, preferably 17 or more, more preferably 19 or more consecutive nucleotides.

(317) In a preferred embodiment, the fragment can be a polynucleotide comprising 15 or more, preferably 17 or more, more preferably 19 or more consecutive nucleotides.

(318) In the present invention, the size of the polynucleotide fragment is the number of nucleotides in the range from, for example, 15 consecutive nucleotides to less than the total number of nucleotides of the sequence, from 17 consecutive nucleotides to less than the total number of nucleotides of the sequence, or from 19 consecutive nucleotides to less than the total number of nucleotides of the sequence, in the nucleotide sequence of each polynucleotide.

(319) Specific examples of the aforementioned polynucleotide combination that constitutes the kit or the device of the present invention can include combinations of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more of the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 269 and 851 to 856 shown in Table 1 described above. However, these are given merely for illustrative purposes, and all of various other possible combinations are included in the present invention.

(320) The aforementioned combination constituting the kit or the device for discriminating a breast cancer patient from a healthy subject according to the present invention is desirably, for example, a combination of two or more polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs shown in Table 1. Usually, a combination of two of these polynucleotides can produce adequate performance. Specifically, any two of the aforementioned polynucleotides that consist of the aforementioned nucleotide sequences represented by SEQ ID NOs: 1 to 269 and 851 to 856 may be combined. For such a combination, it is preferred to select at least one of the newly found polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 235 and 851 to 856. More specifically, the combination is more preferably a combination comprising at least one of polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 20, 24, 26, 27, 30, 33, 182, 194, 206, and 208, among the combinations of the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 269 and 851 to 856.

(321) The combination of polynucleotides with cancer type specificity capable of discriminating a breast cancer patient not only from a healthy subject but also from other cancer patients is preferably, for example, a combination of multiple polynucleotides comprising at least one polynucleotide selected from the group consisting of polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 16, 17, 20, 21, 22, 23, 24, 25, 26, 27, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 43, 45, 46, 47, 49, 50, 51, 52, 53, 54, 55, 58, 59, 60, 62, 63, 64, 65, 67, 68, 69, 70, 71, 72, 73, 75, 77, 79, 80, 81, 82, 83, 86, 88, 89, 90, 92, 93, 94, 96, 98, 99, 100, 103, 104, 106, 107, 108, 110, 111, 113, 114, 115, 116, 118, 119, 121, 122, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 135, 136, 139, 140, 143, 145, 146, 147, 149, 150, 155, 157, 160, 161, 165, 167, 171, 173, 174, 175, 177, 178, 181, 182, 186, 190, 193, 194, 199, 204, 205, 206, 208, 211, 218, 225, 232, 236, 237, 238, 239, 242, 243, 244, 246, 247, 252, 260, 265, 266, 851, 852, 853, 854, 855, 856 (hereinafter, this group is referred to as “cancer type-specific polynucleotide group 1”), with any of the polynucleotides consisting of nucleotide sequences represented by the other SEQ ID NOs.

(322) The combination of polynucleotides with cancer type specificity capable of discriminating a

breast cancer patient not only from a healthy subject but also from other cancer patients is more preferably a combination of multiple polynucleotides, any of which are selected from the cancer type-specific polynucleotide group 1 described above.

(323) The combination of polynucleotides with cancer type specificity capable of discriminating a breast cancer patient not only from a healthy subject but also from other cancer patients is more preferably a combination comprising at least one polynucleotide selected from, particularly, the group consisting of polynucleotides of SEQ ID NOs: 1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 16, 17, 20, 21, 22, 23, 24, 25, 26, 27, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 43, 45, 46, 47, 49, 50, 51, 52, 54, 55, 58, 59, 60, 62, 63, 64, 65, 67, 68, 69, 71, 72, 73, 75, 77, 79, 80, 82, 83, 86, 88, 92, 93, 96, 99, 103, 104, 106, 110, 111, 114, 116, 118, 119, 122, 124, 125, 127, 130, 132, 133, 135, 139, 143, 145, 147, 149, 157, 160, 173, 177, 181, 182, 186, 211, 218, 232, 236, 237, 238, 239, 242, 243, 246, 247, 260, 266, 851, 852, 853, 854 (hereinafter, this group is referred to as “cancer type-specific polynucleotide group 2”), among the combinations of multiple polynucleotides selected from the cancer type-specific polynucleotide group 1 described above.

(324) The number of the polynucleotides with cancer type specificity in the combination described above can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more and is preferably 2 or more for the combination.

(325) Non-limiting examples of the combination of two polynucleotides that consist of one polynucleotide that consists of a nucleotide sequence selected from the cancer type-specific polynucleotide group 2 or a complementary sequence thereof and one polynucleotide that consists of a nucleotide sequence selected from the cancer type-specific polynucleotide group 1 or a complementary sequence thereof are listed below.

(326) (1-1) a combination of SEQ ID NOs: 2 and 1 (markers: hsa-miR-4730 and hsa-miR-4783-3p); (1-2) a combination of SEQ ID NOs: 2 and 237 (markers: hsa-miR-4730 and hsa-miR-602); (1-3) a combination of SEQ ID NOs: 2 and 4 (markers: hsa-miR-4730 and hsa-miR-4634); (1-4) a combination of SEQ ID NOs: 2 and 3 (markers: hsa-miR-4730 and hsa-miR-1307-3p); (1-5) a combination of SEQ ID NOs: 2 and 51 (markers: hsa-miR-4730 and hsa-miR-3940-5p); (2-1) a combination of SEQ ID NOs: 1 and 237 (markers: hsa-miR-4783-3p and hsa-miR-602); (2-2) a combination of SEQ ID NOs: 1 and 4 (markers: hsa-miR-4783-3p and hsa-miR-4634); (2-3) a combination of SEQ ID NOs: 1 and 3 (markers: hsa-miR-4783-3p and hsa-miR-1307-3p); (2-4) a combination of SEQ ID NOs: 1 and 51 (markers: hsa-miR-4783-3p and hsa-miR-3940-5p); (2-5) a combination of SEQ ID NOs: 1 and 6 (markers: hsa-miR-4783-3p and hsa-miR-4532); (3-1) a combination of SEQ ID NOs: 4 and 237 (markers: hsa-miR-4634 and hsa-miR-602); (3-2) a combination of SEQ ID NOs: 3 and 237 (markers: hsa-miR-1307-3p and hsa-miR-602); (3-3) a combination of SEQ ID NOs: 51 and 237 (markers: hsa-miR-3940-5p and hsa-miR-602); (3-4) a combination of SEQ ID NOs: 237 and 6 (markers: hsa-miR-602 and hsa-miR-4532); (3-5) a combination of SEQ ID NOs: 237 and 12 (markers: hsa-miR-602 and hsa-miR-3184-5p); (4-1) a combination of SEQ ID NOs: 3 and 4 (markers: hsa-miR-1307-3p and hsa-miR-4634); (4-2) a combination of SEQ ID NOs: 4 and 51 (markers: hsa-miR-4634 and hsa-miR-3940-5p); (4-3) a combination of SEQ ID NOs: 4 and 6 (markers: hsa-miR-4634 and hsa-miR-4532); (4-4) a combination of SEQ ID NOs: 4 and 12 (markers: hsa-miR-4634 and hsa-miR-3184-5p); (4-5) a combination of SEQ ID NOs: 4 and 15 (markers: hsa-miR-4634 and hsa-miR-4674); (5-1) a combination of SEQ ID NOs: 3 and 51 (markers: hsa-miR-1307-3p and hsa-miR-3940-5p); (5-2) a combination of SEQ ID NOs: 3 and 6 (markers: hsa-miR-1307-3p and hsa-miR-4532); (5-3) a combination of SEQ ID NOs: 3 and 12 (markers: hsa-miR-1307-3p and hsa-miR-3184-5p); (5-4) a combination of SEQ ID NOs: 3 and 15 (markers: hsa-miR-1307-3p and hsa-miR-4674); (5-5) a combination of SEQ ID NOs: 3 and 8 (markers: hsa-miR-1307-3p and hsa-miR-3178); (6-1) a combination of SEQ ID NOs: 51 and 6 (markers: hsa-miR-3940-5p and hsa-miR-4532); (6-2) a combination of SEQ ID NOs: 51 and 12 (markers: hsa-miR-3940-5p and hsa-miR-3184-5p); (6-3) a combination of SEQ ID NOs: 51 and 15 (markers: hsa-miR-3940-5p and hsa-miR-4674); (6-4) a combination of SEQ ID NOs: 51 and 8 (markers: hsa-miR-3940-5p and hsa-miR-3178); (6-5) a

combination of SEQ ID NOs: 51 and 34 (markers: hsa-miR-3940-5p and hsa-miR-6805-5p); (7-1) a combination of SEQ ID NOs: 2 and 6 (markers: hsa-miR-4730 and hsa-miR-4532); (7-2) a combination of SEQ ID NOs: 12 and 6 (markers: hsa-miR-3184-5p and hsa-miR-4532); (7-3) a combination of SEQ ID NOs: 15 and 6 (markers: hsa-miR-4674 and hsa-miR-4532); (7-4) a combination of SEQ ID NOs: 8 and 6 (markers: hsa-miR-3178 and hsa-miR-4532); (7-5) a combination of SEQ ID NOs: 6 and 34 (markers: hsa-miR-4532 and hsa-miR-6805-5p); (8-1) a combination of SEQ ID NOs: 2 and 12 (markers: hsa-miR-4730 and hsa-miR-3184-5p); (8-2) a combination of SEQ ID NOs: 1 and 12 (markers: hsa-miR-4783-3p and hsa-miR-3184-5p); (8-3) a combination of SEQ ID NOs: 12 and 15 (markers: hsa-miR-3184-5p and hsa-miR-4674); (8-4) a combination of SEQ ID NOs: 8 and 12 (markers: hsa-miR-3178 and hsa-miR-3184-5p); (8-5) a combination of SEQ ID NOs: 12 and 34 (markers: hsa-miR-3184-5p and hsa-miR-6805-5p); (9-1) a combination of SEQ ID NOs: 2 and 15 (markers: hsa-miR-4730 and hsa-miR-4674); (9-2) a combination of SEQ ID NOs: 1 and 15 (markers: hsa-miR-4783-3p and hsa-miR-4674); (9-3) a combination of SEQ ID NOs: 237 and 15 (markers: hsa-miR-602 and hsa-miR-4674); (9-4) a combination of SEQ ID NOs: 8 and 15 (markers: hsa-miR-3178 and hsa-miR-4674); (9-5) a combination of SEQ ID NOs: 15 and 34 (markers: hsa-miR-4674 and hsa-miR-6805-5p); (10-1) a combination of SEQ ID NOs: 2 and 8 (markers: hsa-miR-4730 and hsa-miR-3178); (10-2) a combination of SEQ ID NOs: 1 and 8 (markers: hsa-miR-4783-3p and hsa-miR-3178); (10-3) a combination of SEQ ID NOs: 237 and 8 (markers: hsa-miR-602 and hsa-miR-3178); (10-4) a combination of SEQ ID NOs: 4 and 8 (markers: hsa-miR-4634 and hsa-miR-3178); (10-5) a combination of SEQ ID NOs: 8 and 34 (markers: hsa-miR-3178 and hsa-miR-6805-5p); (11-1) a combination of SEQ ID NOs: 2 and 34 (markers: hsa-miR-4730 and hsa-miR-6805-5p); (11-2) a combination of SEQ ID NOs: 1 and 34 (markers: hsa-miR-4783-3p and hsa-miR-6805-5p); (11-3) a combination of SEQ ID NOs: 237 and 34 (markers: hsa-miR-602 and hsa-miR-6805-5p); (11-4) a combination of SEQ ID NOs: 4 and 34 (markers: hsa-miR-4634 and hsa-miR-6805-5p); (11-5) a combination of SEQ ID NOs: 3 and 34 (markers: hsa-miR-1307-3p and hsa-miR-6805-5p); (12-1) a combination of SEQ ID NOs: 2 and 9 (markers: hsa-miR-4730 and hsa-miR-6729-5p); (12-2) a combination of SEQ ID NOs: 1 and 9 (markers: hsa-miR-4783-3p and hsa-miR-6729-5p); (12-3) a combination of SEQ ID NOs: 9 and 237 (markers: hsa-miR-6729-5p and hsa-miR-602); (12-4) a combination of SEQ ID NOs: 4 and 9 (markers: hsa-miR-4634 and hsa-miR-6729-5p); (12-5) a combination of SEQ ID NOs: 3 and 9 (markers: hsa-miR-1307-3p and hsa-miR-6729-5p); (13-1) a combination of SEQ ID NOs: 2 and 143 (markers: hsa-miR-4730 and hsa-miR-1228-5p); (13-2) a combination of SEQ ID NOs: 1 and 143 (markers: hsa-miR-4783-3p and hsa-miR-1228-5p); (13-3) a combination of SEQ ID NOs: 237 and 143 (markers: hsa-miR-602 and hsa-miR-1228-5p); (13-4) a combination of SEQ ID NOs: 4 and 143 (markers: hsa-miR-4634 and hsa-miR-1228-5p); (13-5) a combination of SEQ ID NOs: 3 and 143 (markers: hsa-miR-1307-3p and hsa-miR-1228-5p); (14-1) a combination of SEQ ID NOs: 2 and 13 (markers: hsa-miR-4730 and hsa-miR-6727-5p); (14-2) a combination of SEQ ID NOs: 1 and 13 (markers: hsa-miR-4783-3p and hsa-miR-6727-5p); (14-3) a combination of SEQ ID NOs: 237 and 13 (markers: hsa-miR-602 and hsa-miR-6727-5p); (14-4) a combination of SEQ ID NOs: 4 and 13 (markers: hsa-miR-4634 and hsa-miR-6727-5p); (14-5) a combination of SEQ ID NOs: 3 and 13 (markers: hsa-miR-1307-3p and hsa-miR-6727-5p); (15-1) a combination of SEQ ID NOs: 2 and 125 (markers: hsa-miR-4730 and hsa-miR-6781-5p); (15-2) a combination of SEQ ID NOs: 1 and 125 (markers: hsa-miR-4783-3p and hsa-miR-6781-5p); (15-3) a combination of SEQ ID NOs: 237 and 125 (markers: hsa-miR-602 and hsa-miR-6781-5p); (15-4) a combination of SEQ ID NOs: 4 and 125 (markers: hsa-miR-4634 and hsa-miR-6781-5p); (15-5) a combination of SEQ ID NOs: 3 and 125 (markers: hsa-miR-1307-3p and hsa-miR-6781-5p); (16-1) a combination of SEQ ID NOs: 2 and 236 (markers: hsa-miR-4730 and hsa-miR-760); (16-2) a combination of SEQ ID NOs: 1 and 236 (markers: hsa-miR-4783-3p and hsa-miR-760); (16-3) a combination of SEQ ID NOs: 237 and 236 (markers: hsa-miR-602 and hsa-miR-760); (16-4) a combination of SEQ ID NOs: 4 and 236 (markers: hsa-miR-4634 and hsa-miR-760); (16-5) a

combination of SEQ ID NOs: 3 and 236 (markers: hsa-miR-1307-3p and hsa-miR-760); (17-1) a combination of SEQ ID NOs: 2 and 46 (markers: hsa-miR-4730 and hsa-miR-1231); (17-2) a combination of SEQ ID NOs: 1 and 46 (markers: hsa-miR-4783-3p and hsa-miR-1231); (17-3) a combination of SEQ ID NOs: 237 and 46 (markers: hsa-miR-602 and hsa-miR-1231); (17-4) a combination of SEQ ID NOs: 4 and 46 (markers: hsa-miR-4634 and hsa-miR-1231); (17-5) a combination of SEQ ID NOs: 3 and 46 (markers: hsa-miR-1307-3p and hsa-miR-1231); (18-1) a combination of SEQ ID NOs: 2 and 32 (markers: hsa-miR-4730 and hsa-miR-6836-3p); (18-2) a combination of SEQ ID NOs: 1 and 32 (markers: hsa-miR-4783-3p and hsa-miR-6836-3p); (18-3) a combination of SEQ ID NOs: 237 and 32 (markers: hsa-miR-602 and hsa-miR-6836-3p); (18-4) a combination of SEQ ID NOs: 4 and 32 (markers: hsa-miR-4634 and hsa-miR-6836-3p); (18-5) a combination of SEQ ID NOs: 3 and 32 (markers: hsa-miR-1307-3p and hsa-miR-6836-3p); (19-1) a combination of SEQ ID NOs: 2 and 62 (markers: hsa-miR-4730 and hsa-miR-4492); (19-2) a combination of SEQ ID NOs: 1 and 62 (markers: hsa-miR-4783-3p and hsa-miR-4492); (19-3) a combination of SEQ ID NOs: 237 and 62 (markers: hsa-miR-602 and hsa-miR-4492); (19-4) a combination of SEQ ID NOs: 4 and 62 (markers: hsa-miR-4634 and hsa-miR-4492); (19-5) a combination of SEQ ID NOs: 3 and 62 (markers: hsa-miR-1307-3p and hsa-miR-4492); (20-1) a combination of SEQ ID NOs: 2 and 88 (markers: hsa-miR-4730 and hsa-miR-4484); (20-2) a combination of SEQ ID NOs: 1 and 88 (markers: hsa-miR-4783-3p and hsa-miR-4484); (20-3) a combination of SEQ ID NOs: 237 and 88 (markers: hsa-miR-602 and hsa-miR-4484); (20-4) a combination of SEQ ID NOs: 4 and 88 (markers: hsa-miR-4634 and hsa-miR-4484); (20-5) a combination of SEQ ID NOs: 3 and 88 (markers: hsa-miR-1307-3p and hsa-miR-4484); (21-1) a combination of SEQ ID NOs: 2 and 52 (markers: hsa-miR-4730 and hsa-miR-4741); (21-2) a combination of SEQ ID NOs: 1 and 52 (markers: hsa-miR-4783-3p and hsa-miR-4741); (21-3) a combination of SEQ ID NOs: 237 and 52 (markers: hsa-miR-602 and hsa-miR-4741); (21-4) a combination of SEQ ID NOs: 4 and 52 (markers: hsa-miR-4634 and hsa-miR-4741); (21-5) a combination of SEQ ID NOs: 3 and 52 (markers: hsa-miR-1307-3p and hsa-miR-4741); (22-1) a combination of SEQ ID NOs: 2 and 7 (markers: hsa-miR-4730 and hsa-miR-7704); (22-2) a combination of SEQ ID NOs: 1 and 7 (markers: hsa-miR-4783-3p and hsa-miR-7704); (22-3) a combination of SEQ ID NOs: 237 and 7 (markers: hsa-miR-602 and hsa-miR-7704); (22-4) a combination of SEQ ID NOs: 4 and 7 (markers: hsa-miR-4634 and hsa-miR-7704); (22-5) a combination of SEQ ID NOs: 3 and 7 (markers: hsa-miR-1307-3p and hsa-miR-7704); (23-1) a combination of SEQ ID NOs: 2 and 26 (markers: hsa-miR-4730 and hsa-miR-6875-5p); (23-2) a combination of SEQ ID NOs: 1 and 26 (markers: hsa-miR-4783-3p and hsa-miR-6875-5p); (23-3) a combination of SEQ ID NOs: 237 and 26 (markers: hsa-miR-602 and hsa-miR-6875-5p); (23-4) a combination of SEQ ID NOs: 4 and 26 (markers: hsa-miR-4634 and hsa-miR-6875-5p); (23-5) a combination of SEQ ID NOs: 3 and 26 (markers: hsa-miR-1307-3p and hsa-miR-6875-5p); (24-1) a combination of SEQ ID NOs: 2 and 25 (markers: hsa-miR-4730 and hsa-miR-8069); (24-2) a combination of SEQ ID NOs: 1 and 25 (markers: hsa-miR-4783-3p and hsa-miR-8069); (24-3) a combination of SEQ ID NOs: 237 and 25 (markers: hsa-miR-602 and hsa-miR-8069); (24-4) a combination of SEQ ID NOs: 4 and 25 (markers: hsa-miR-4634 and hsa-miR-8069); (24-5) a combination of SEQ ID NOs: 3 and 25 (markers: hsa-miR-1307-3p and hsa-miR-8069); (25-1) a combination of SEQ ID NOs: 2 and 54 (markers: hsa-miR-4730 and hsa-miR-6869-5p); (25-2) a combination of SEQ ID NOs: 1 and 54 (markers: hsa-miR-4783-3p and hsa-miR-6869-5p); (25-3) a combination of SEQ ID NOs: 237 and 54 (markers: hsa-miR-602 and hsa-miR-6869-5p); (25-4) a combination of SEQ ID NOs: 4 and 54 (markers: hsa-miR-4634 and hsa-miR-6869-5p); (25-5) a combination of SEQ ID NOs: 3 and 54 (markers: hsa-miR-1307-3p and hsa-miR-6869-5p); (26-1) a combination of SEQ ID NOs: 2 and 92 (markers: hsa-miR-4730 and hsa-miR-4725-3p); (26-2) a combination of SEQ ID NOs: 1 and 92 (markers: hsa-miR-4783-3p and hsa-miR-4725-3p); (26-3) a combination of SEQ ID NOs: 237 and 92 (markers: hsa-miR-602 and hsa-miR-4725-3p); (26-4) a combination of SEQ ID NOs: 4 and 92 (markers: hsa-miR-4634 and hsa-miR-4725-3p); (26-5) a

combination of SEQ ID NOs: 3 and 92 (markers: hsa-miR-1307-3p and hsa-miR-4730-3p); (27-1) a combination of SEQ ID NOs: 2 and 14 (markers: hsa-miR-4730 and hsa-miR-6088); (27-2) a combination of SEQ ID NOs: 1 and 14 (markers: hsa-miR-4783-3p and hsa-miR-6088); (27-3) a combination of SEQ ID NOs: 237 and 14 (markers: hsa-miR-602 and hsa-miR-6088); (27-4) a combination of SEQ ID NOs: 4 and 14 (markers: hsa-miR-4634 and hsa-miR-6088); (27-5) a combination of SEQ ID NOs: 3 and 14 (markers: hsa-miR-1307-3p and hsa-miR-6088); (28-1) a combination of SEQ ID NOs: 2 and 242 (markers: hsa-miR-4730 and hsa-miR-135a-3p); (28-2) a combination of SEQ ID NOs: 1 and 242 (markers: hsa-miR-4783-3p and hsa-miR-135a-3p); (28-3) a combination of SEQ ID NOs: 237 and 242 (markers: hsa-miR-602 and hsa-miR-135a-3p); (28-4) a combination of SEQ ID NOs: 4 and 242 (markers: hsa-miR-4634 and hsa-miR-135a-3p); (28-5) a combination of SEQ ID NOs: 3 and 242 (markers: hsa-miR-1307-3p and hsa-miR-135a-3p); (29-1) a combination of SEQ ID NOs: 2 and 47 (markers: hsa-miR-4730 and hsa-miR-1908-3p); (29-2) a combination of SEQ ID NOs: 1 and 47 (markers: hsa-miR-4783-3p and hsa-miR-1908-3p); (29-3) a combination of SEQ ID NOs: 237 and 47 (markers: hsa-miR-602 and hsa-miR-1908-3p); (29-4) a combination of SEQ ID NOs: 4 and 47 (markers: hsa-miR-4634 and hsa-miR-1908-3p); (29-5) a combination of SEQ ID NOs: 3 and 47 (markers: hsa-miR-1307-3p and hsa-miR-1908-3p); (30-1) a combination of SEQ ID NOs: 2 and 45 (markers: hsa-miR-4730 and hsa-miR-6771-5p); (30-2) a combination of SEQ ID NOs: 1 and 45 (markers: hsa-miR-4783-3p and hsa-miR-6771-5p); (30-3) a combination of SEQ ID NOs: 237 and 45 (markers: hsa-miR-602 and hsa-miR-6771-5p); (30-4) a combination of SEQ ID NOs: 4 and 45 (markers: hsa-miR-4634 and hsa-miR-6771-5p); (30-5) a combination of SEQ ID NOs: 3 and 45 (markers: hsa-miR-1307-3p and hsa-miR-6771-5p); (31-1) a combination of SEQ ID NOs: 2 and 39 (markers: hsa-miR-4730 and hsa-miR-6850-5p); (31-2) a combination of SEQ ID NOs: 1 and 39 (markers: hsa-miR-4783-3p and hsa-miR-6850-5p); (31-3) a combination of SEQ ID NOs: 237 and 39 (markers: hsa-miR-602 and hsa-miR-6850-5p); (31-4) a combination of SEQ ID NOs: 4 and 39 (markers: hsa-miR-4634 and hsa-miR-6850-5p); (31-5) a combination of SEQ ID NOs: 3 and 39 (markers: hsa-miR-1307-3p and hsa-miR-6850-5p); (32-1) a combination of SEQ ID NOs: 2 and 21 (markers: hsa-miR-4730 and hsa-miR-885-3p); (32-2) a combination of SEQ ID NOs: 1 and 21 (markers: hsa-miR-4783-3p and hsa-miR-885-3p); (32-3) a combination of SEQ ID NOs: 237 and 21 (markers: hsa-miR-602 and hsa-miR-885-3p); (32-4) a combination of SEQ ID NOs: 4 and 21 (markers: hsa-miR-4634 and hsa-miR-885-3p); (32-5) a combination of SEQ ID NOs: 3 and 21 (markers: hsa-miR-1307-3p and hsa-miR-885-3p); (33-1) a combination of SEQ ID NOs: 2 and 17 (markers: hsa-miR-4730 and hsa-miR-4787-5p); (33-2) a combination of SEQ ID NOs: 1 and 17 (markers: hsa-miR-4783-3p and hsa-miR-4787-5p); (33-3) a combination of SEQ ID NOs: 237 and 17 (markers: hsa-miR-602 and hsa-miR-4787-5p); (33-4) a combination of SEQ ID NOs: 4 and 17 (markers: hsa-miR-4634 and hsa-miR-4787-5p); (33-5) a combination of SEQ ID NOs: 3 and 17 (markers: hsa-miR-1307-3p and hsa-miR-4787-5p); (34-1) a combination of SEQ ID NOs: 2 and 83 (markers: hsa-miR-4730 and hsa-miR-1909-3p); (34-2) a combination of SEQ ID NOs: 1 and 83 (markers: hsa-miR-4783-3p and hsa-miR-1909-3p); (34-3) a combination of SEQ ID NOs: 237 and 83 (markers: hsa-miR-602 and hsa-miR-1909-3p); (34-4) a combination of SEQ ID NOs: 4 and 83 (markers: hsa-miR-4634 and hsa-miR-1909-3p); (34-5) a combination of SEQ ID NOs: 3 and 83 (markers: hsa-miR-1307-3p and hsa-miR-1909-3p); (35-1) a combination of SEQ ID NOs: 2 and 149 (markers: hsa-miR-4730 and hsa-miR-638); (35-2) a combination of SEQ ID NOs: 1 and 149 (markers: hsa-miR-4783-3p and hsa-miR-638); (35-3) a combination of SEQ ID NOs: 237 and 149 (markers: hsa-miR-602 and hsa-miR-638); (35-4) a combination of SEQ ID NOs: 4 and 149 (markers: hsa-miR-4634 and hsa-miR-638); (35-5) a combination of SEQ ID NOs: 3 and 149 (markers: hsa-miR-1307-3p and hsa-miR-638); (36-1) a combination of SEQ ID NOs: 2 and 246 (markers: hsa-miR-4730 and hsa-miR-1915-3p); (36-2) a combination of SEQ ID NOs: 1 and 246 (markers: hsa-miR-4783-3p and hsa-miR-1915-3p); (36-3) a combination of SEQ ID NOs: 237 and 246 (markers: hsa-miR-602 and hsa-miR-1915-3p); (36-4) a combination of SEQ ID NOs: 4 and 246 (markers: hsa-miR-4634 and hsa-miR-1915-3p); (36-5) a

combination of SEQ ID NOs: 3 and 246 (markers: hsa-miR-1307-3p and hsa-miR-1915-3p); (37-1) a combination of SEQ ID NOs: 2 and 22 (markers: hsa-miR-4730 and hsa-miR-6802-5p); (37-2) a combination of SEQ ID NOs: 1 and 22 (markers: hsa-miR-4783-3p and hsa-miR-6802-5p); (37-3) a combination of SEQ ID NOs: 237 and 22 (markers: hsa-miR-602 and hsa-miR-6802-5p); (37-4) a combination of SEQ ID NOs: 4 and 22 (markers: hsa-miR-4634 and hsa-miR-6802-5p); (37-5) a combination of SEQ ID NOs: 3 and 22 (markers: hsa-miR-1307-3p and hsa-miR-6802-5p); (38-1) a combination of SEQ ID NOs: 2 and 55 (markers: hsa-miR-4730 and hsa-miR-5090); (38-2) a combination of SEQ ID NOs: 1 and 55 (markers: hsa-miR-4783-3p and hsa-miR-5090); (38-3) a combination of SEQ ID NOs: 237 and 55 (markers: hsa-miR-602 and hsa-miR-5090); (38-4) a combination of SEQ ID NOs: 4 and 55 (markers: hsa-miR-4634 and hsa-miR-5090); (38-5) a combination of SEQ ID NOs: 3 and 55 (markers: hsa-miR-1307-3p and hsa-miR-5090); (39-1) a combination of SEQ ID NOs: 2 and 182 (markers: hsa-miR-4730 and hsa-miR-3196); (39-2) a combination of SEQ ID NOs: 1 and 182 (markers: hsa-miR-4783-3p and hsa-miR-3196); (39-3) a combination of SEQ ID NOs: 237 and 182 (markers: hsa-miR-602 and hsa-miR-3196); (39-4) a combination of SEQ ID NOs: 4 and 182 (markers: hsa-miR-4634 and hsa-miR-3196); (39-5) a combination of SEQ ID NOs: 3 and 182 (markers: hsa-miR-1307-3p and hsa-miR-3196); (40-1) a combination of SEQ ID NOs: 2 and 73 (markers: hsa-miR-4730 and hsa-miR-4707-3p); (40-2) a combination of SEQ ID NOs: 1 and 73 (markers: hsa-miR-4783-3p and hsa-miR-4707-3p); (40-3) a combination of SEQ ID NOs: 237 and 73 (markers: hsa-miR-602 and hsa-miR-4707-3p); (40-4) a combination of SEQ ID NOs: 4 and 73 (markers: hsa-miR-4634 and hsa-miR-4707-3p); (40-5) a combination of SEQ ID NOs: 3 and 73 (markers: hsa-miR-1307-3p and hsa-miR-4707-3p); (41-1) a combination of SEQ ID NOs: 2 and 77 (markers: hsa-miR-4730 and hsa-miR-4651); (41-2) a combination of SEQ ID NOs: 1 and 77 (markers: hsa-miR-4783-3p and hsa-miR-4651); (41-3) a combination of SEQ ID NOs: 237 and 77 (markers: hsa-miR-602 and hsa-miR-4651); (41-4) a combination of SEQ ID NOs: 4 and 77 (markers: hsa-miR-4634 and hsa-miR-4651); (41-5) a combination of SEQ ID NOs: 3 and 77 (markers: hsa-miR-1307-3p and hsa-miR-4651); (42-1) a combination of SEQ ID NOs: 2 and 24 (markers: hsa-miR-4730 and hsa-miR-6787-5p); (42-2) a combination of SEQ ID NOs: 1 and 24 (markers: hsa-miR-4783-3p and hsa-miR-6787-5p); (42-3) a combination of SEQ ID NOs: 237 and 24 (markers: hsa-miR-602 and hsa-miR-6787-5p); (42-4) a combination of SEQ ID NOs: 4 and 24 (markers: hsa-miR-4634 and hsa-miR-6787-5p); (42-5) a combination of SEQ ID NOs: 3 and 24 (markers: hsa-miR-1307-3p and hsa-miR-6787-5p); (43-1) a combination of SEQ ID NOs: 2 and 103 (markers: hsa-miR-4730 and hsa-miR-3180-3p); (43-2) a combination of SEQ ID NOs: 1 and 103 (markers: hsa-miR-4783-3p and hsa-miR-3180-3p); (43-3) a combination of SEQ ID NOs: 237 and 103 (markers: hsa-miR-602 and hsa-miR-3180-3p); (43-4) a combination of SEQ ID NOs: 4 and 103 (markers: hsa-miR-4634 and hsa-miR-3180-3p); (43-5) a combination of SEQ ID NOs: 3 and 103 (markers: hsa-miR-1307-3p and hsa-miR-3180-3p); (44-1) a combination of SEQ ID NOs: 2 and 49 (markers: hsa-miR-4730 and hsa-miR-3937); (44-2) a combination of SEQ ID NOs: 1 and 49 (markers: hsa-miR-4783-3p and hsa-miR-3937); (44-3) a combination of SEQ ID NOs: 237 and 49 (markers: hsa-miR-602 and hsa-miR-3937); (44-4) a combination of SEQ ID NOs: 4 and 49 (markers: hsa-miR-4634 and hsa-miR-3937); (44-5) a combination of SEQ ID NOs: 3 and 49 (markers: hsa-miR-1307-3p and hsa-miR-3937); (45-1) a combination of SEQ ID NOs: 2 and 239 (markers: hsa-miR-4730 and hsa-miR-92a-2-5p); (45-2) a combination of SEQ ID NOs: 1 and 239 (markers: hsa-miR-4783-3p and hsa-miR-92a-2-5p); (45-3) a combination of SEQ ID NOs: 237 and 239 (markers: hsa-miR-602 and hsa-miR-92a-2-5p); (45-4) a combination of SEQ ID NOs: 4 and 239 (markers: hsa-miR-4634 and hsa-miR-92a-2-5p); (45-5) a combination of SEQ ID NOs: 3 and 239 (markers: hsa-miR-1307-3p and hsa-miR-92a-2-5p); (46-1) a combination of SEQ ID NOs: 2 and 23 (markers: hsa-miR-4730 and hsa-miR-328-5p); (46-2) a combination of SEQ ID NOs: 1 and 23 (markers: hsa-miR-4783-3p and hsa-miR-328-5p); (46-3) a combination of SEQ ID NOs: 237 and 23 (markers: hsa-miR-602 and hsa-miR-328-5p); (46-4) a combination of SEQ ID NOs: 4 and 23 (markers: hsa-miR-4634 and hsa-miR-328-

5p); (46-5) a combination of SEQ ID NOs: 3 and 23 (markers: hsa-miR-1307-3p and hsa-miR-328-5p); (47-1) a combination of SEQ ID NOs: 2 and 58 (markers: hsa-miR-4730 and hsa-miR-128-1-5p); (47-2) a combination of SEQ ID NOs: 1 and 58 (markers: hsa-miR-4783-3p and hsa-miR-128-1-5p); (47-3) a combination of SEQ ID NOs: 237 and 58 (markers: hsa-miR-602 and hsa-miR-128-1-5p); (47-4) a combination of SEQ ID NOs: 4 and 58 (markers: hsa-miR-4634 and hsa-miR-128-1-5p); (47-5) a combination of SEQ ID NOs: 3 and 58 (markers: hsa-miR-1307-3p and hsa-miR-128-1-5p); (48-1) a combination of SEQ ID NOs: 2 and 211 (markers: hsa-miR-4730 and hsa-miR-6765-5p); (48-2) a combination of SEQ ID NOs: 1 and 211 (markers: hsa-miR-4783-3p and hsa-miR-6765-5p); (48-3) a combination of SEQ ID NOs: 237 and 211 (markers: hsa-miR-602 and hsa-miR-6765-5p); (48-4) a combination of SEQ ID NOs: 4 and 211 (markers: hsa-miR-4634 and hsa-miR-6765-5p); (48-5) a combination of SEQ ID NOs: 3 and 211 (markers: hsa-miR-1307-3p and hsa-miR-6765-5p); (49-1) a combination of SEQ ID NOs: 2 and 147 (markers: hsa-miR-4730 and hsa-miR-6786-5p); (49-2) a combination of SEQ ID NOs: 1 and 147 (markers: hsa-miR-4783-3p and hsa-miR-6786-5p); (49-3) a combination of SEQ ID NOs: 237 and 147 (markers: hsa-miR-602 and hsa-miR-6786-5p); (49-4) a combination of SEQ ID NOs: 4 and 147 (markers: hsa-miR-4634 and hsa-miR-6786-5p); (49-5) a combination of SEQ ID NOs: 3 and 147 (markers: hsa-miR-1307-3p and hsa-miR-6786-5p); (50-1) a combination of SEQ ID NOs: 2 and 65 (markers: hsa-miR-4730 and hsa-miR-4525); (50-2) a combination of SEQ ID NOs: 1 and 65 (markers: hsa-miR-4783-3p and hsa-miR-4525); (50-3) a combination of SEQ ID NOs: 237 and 65 (markers: hsa-miR-602 and hsa-miR-4525); (50-4) a combination of SEQ ID NOs: 4 and 65 (markers: hsa-miR-4634 and hsa-miR-4525); (50-5) a combination of SEQ ID NOs: 3 and 65 (markers: hsa-miR-1307-3p and hsa-miR-4525); (51-1) a combination of SEQ ID NOs: 2 and 31 (markers: hsa-miR-4730 and hsa-miR-3665); (51-2) a combination of SEQ ID NOs: 1 and 31 (markers: hsa-miR-4783-3p and hsa-miR-3665); (51-3) a combination of SEQ ID NOs: 237 and 31 (markers: hsa-miR-602 and hsa-miR-3665); (51-4) a combination of SEQ ID NOs: 4 and 31 (markers: hsa-miR-4634 and hsa-miR-3665); (51-5) a combination of SEQ ID NOs: 3 and 31 (markers: hsa-miR-1307-3p and hsa-miR-3665); (52-1) a combination of SEQ ID NOs: 2 and 72 (markers: hsa-miR-4730 and hsa-miR-663b); (52-2) a combination of SEQ ID NOs: 1 and 72 (markers: hsa-miR-4783-3p and hsa-miR-663b); (52-3) a combination of SEQ ID NOs: 237 and 72 (markers: hsa-miR-602 and hsa-miR-663b); (52-4) a combination of SEQ ID NOs: 4 and 72 (markers: hsa-miR-4634 and hsa-miR-663b); (52-5) a combination of SEQ ID NOs: 3 and 72 (markers: hsa-miR-1307-3p and hsa-miR-663b); (53-1) a combination of SEQ ID NOs: 2 and 63 (markers: hsa-miR-4730 and hsa-miR-6785-5p); (53-2) a combination of SEQ ID NOs: 1 and 63 (markers: hsa-miR-4783-3p and hsa-miR-6785-5p); (53-3) a combination of SEQ ID NOs: 237 and 63 (markers: hsa-miR-602 and hsa-miR-6785-5p); (53-4) a combination of SEQ ID NOs: 4 and 63 (markers: hsa-miR-4634 and hsa-miR-6785-5p); (53-5) a combination of SEQ ID NOs: 3 and 63 (markers: hsa-miR-1307-3p and hsa-miR-6785-5p); (54-1) a combination of SEQ ID NOs: 2 and 80 (markers: hsa-miR-4730 and hsa-miR-4758-5p); (54-2) a combination of SEQ ID NOs: 1 and 80 (markers: hsa-miR-4783-3p and hsa-miR-4758-5p); (54-3) a combination of SEQ ID NOs: 237 and 80 (markers: hsa-miR-602 and hsa-miR-4758-5p); (54-4) a combination of SEQ ID NOs: 4 and 80 (markers: hsa-miR-4634 and hsa-miR-4758-5p); (54-5) a combination of SEQ ID NOs: 3 and 80 (markers: hsa-miR-1307-3p and hsa-miR-4758-5p); (55-1) a combination of SEQ ID NOs: 2 and 37 (markers: hsa-miR-4730 and hsa-miR-197-5p); (55-2) a combination of SEQ ID NOs: 1 and 37 (markers: hsa-miR-4783-3p and hsa-miR-197-5p); (55-3) a combination of SEQ ID NOs: 237 and 37 (markers: hsa-miR-602 and hsa-miR-197-5p); (55-4) a combination of SEQ ID NOs: 4 and 37 (markers: hsa-miR-4634 and hsa-miR-197-5p); (55-5) a combination of SEQ ID NOs: 3 and 37 (markers: hsa-miR-1307-3p and hsa-miR-197-5p); (56-1) a combination of SEQ ID NOs: 2 and 67 (markers: hsa-miR-4730 and hsa-miR-3180); (56-2) a combination of SEQ ID NOs: 1 and 67 (markers: hsa-miR-4783-3p and hsa-miR-3180); (56-3) a combination of SEQ ID NOs: 237 and 67 (markers: hsa-miR-602 and hsa-miR-3180); (56-4) a combination of SEQ ID NOs: 4 and 67

(markers: hsa-miR-4634 and hsa-miR-3180); (56-5) a combination of SEQ ID NOs: 3 and 67 (markers: hsa-miR-1307-3p and hsa-miR-3180); (57-1) a combination of SEQ ID NOs: 2 and 232 (markers: hsa-miR-4730 and hsa-miR-6125); (57-2) a combination of SEQ ID NOs: 1 and 232 (markers: hsa-miR-4783-3p and hsa-miR-6125); (57-3) a combination of SEQ ID NOs: 237 and 232 (markers: hsa-miR-602 and hsa-miR-6125); (57-4) a combination of SEQ ID NOs: 4 and 232 (markers: hsa-miR-4634 and hsa-miR-6125); (57-5) a combination of SEQ ID NOs: 3 and 232 (markers: hsa-miR-1307-3p and hsa-miR-6125); (58-1) a combination of SEQ ID NOs: 2 and 127 (markers: hsa-miR-4730 and hsa-miR-4446-3p); (58-2) a combination of SEQ ID NOs: 1 and 127 (markers: hsa-miR-4783-3p and hsa-miR-4446-3p); (58-3) a combination of SEQ ID NOs: 237 and 127 (markers: hsa-miR-602 and hsa-miR-4446-3p); (58-4) a combination of SEQ ID NOs: 4 and 127 (markers: hsa-miR-4634 and hsa-miR-4446-3p); (58-5) a combination of SEQ ID NOs: 3 and 127 (markers: hsa-miR-1307-3p and hsa-miR-4446-3p); (59-1) a combination of SEQ ID NOs: 2 and 145 (markers: hsa-miR-4730 and hsa-miR-1268a); (59-2) a combination of SEQ ID NOs: 1 and 145 (markers: hsa-miR-4783-3p and hsa-miR-1268a); (59-3) a combination of SEQ ID NOs: 237 and 145 (markers: hsa-miR-602 and hsa-miR-1268a); (59-4) a combination of SEQ ID NOs: 4 and 145 (markers: hsa-miR-4634 and hsa-miR-1268a); (59-5) a combination of SEQ ID NOs: 3 and 145 (markers: hsa-miR-1307-3p and hsa-miR-1268a); (60-1) a combination of SEQ ID NOs: 2 and 16 (markers: hsa-miR-4730 and hsa-miR-8073); (60-2) a combination of SEQ ID NOs: 1 and 16 (markers: hsa-miR-4783-3p and hsa-miR-8073); (60-3) a combination of SEQ ID NOs: 237 and 16 (markers: hsa-miR-602 and hsa-miR-8073); (60-4) a combination of SEQ ID NOs: 4 and 16 (markers: hsa-miR-4634 and hsa-miR-8073); (60-5) a combination of SEQ ID NOs: 3 and 16 (markers: hsa-miR-1307-3p and hsa-miR-8073); (61-1) a combination of SEQ ID NOs: 2 and 11 (markers: hsa-miR-4730 and hsa-miR-4732-5p); (61-2) a combination of SEQ ID NOs: 1 and 11 (markers: hsa-miR-4783-3p and hsa-miR-4732-5p); (61-3) a combination of SEQ ID NOs: 237 and 11 (markers: hsa-miR-602 and hsa-miR-4732-5p); (61-4) a combination of SEQ ID NOs: 4 and 11 (markers: hsa-miR-4634 and hsa-miR-4732-5p); (61-5) a combination of SEQ ID NOs: 3 and 11 (markers: hsa-miR-1307-3p and hsa-miR-4732-5p); (62-1) a combination of SEQ ID NOs: 2 and 186 (markers: hsa-miR-4730 and hsa-miR-6789-5p); (62-2) a combination of SEQ ID NOs: 1 and 186 (markers: hsa-miR-4783-3p and hsa-miR-6789-5p); (62-3) a combination of SEQ ID NOs: 237 and 186 (markers: hsa-miR-602 and hsa-miR-6789-5p); (62-4) a combination of SEQ ID NOs: 4 and 186 (markers: hsa-miR-4634 and hsa-miR-6789-5p); (62-5) a combination of SEQ ID NOs: 3 and 186 (markers: hsa-miR-1307-3p and hsa-miR-6789-5p); (63-1) a combination of SEQ ID NOs: 2 and 50 (markers: hsa-miR-4730 and hsa-miR-887-3p); (63-2) a combination of SEQ ID NOs: 1 and 50 (markers: hsa-miR-4783-3p and hsa-miR-887-3p); (63-3) a combination of SEQ ID NOs: 237 and 50 (markers: hsa-miR-602 and hsa-miR-887-3p); (63-4) a combination of SEQ ID NOs: 4 and 50 (markers: hsa-miR-4634 and hsa-miR-887-3p); (63-5) a combination of SEQ ID NOs: 3 and 50 (markers: hsa-miR-1307-3p and hsa-miR-887-3p); (64-1) a combination of SEQ ID NOs: 2 and 69 (markers: hsa-miR-4730 and hsa-miR-1199-5p); (64-2) a combination of SEQ ID NOs: 1 and 69 (markers: hsa-miR-4783-3p and hsa-miR-1199-5p); (64-3) a combination of SEQ ID NOs: 237 and 69 (markers: hsa-miR-602 and hsa-miR-1199-5p); (64-4) a combination of SEQ ID NOs: 4 and 69 (markers: hsa-miR-4634 and hsa-miR-1199-5p); (64-5) a combination of SEQ ID NOs: 3 and 69 (markers: hsa-miR-1307-3p and hsa-miR-1199-5p); (65-1) a combination of SEQ ID NOs: 2 and 33 (markers: hsa-miR-4730 and hsa-miR-6821-5p); (65-2) a combination of SEQ ID NOs: 1 and 33 (markers: hsa-miR-4783-3p and hsa-miR-6821-5p); (65-3) a combination of SEQ ID NOs: 237 and 33 (markers: hsa-miR-602 and hsa-miR-6821-5p); (65-4) a combination of SEQ ID NOs: 4 and 33 (markers: hsa-miR-4634 and hsa-miR-6821-5p); (65-5) a combination of SEQ ID NOs: 3 and 33 (markers: hsa-miR-1307-3p and hsa-miR-6821-5p); (66-1) a combination of SEQ ID NOs: 2 and 247 (markers: hsa-miR-4730 and hsa-miR-718); (66-2) a combination of SEQ ID NOs: 1 and 247 (markers: hsa-miR-4783-3p and hsa-miR-718); (66-3) a combination of SEQ ID NOs: 237 and 247 (markers: hsa-miR-602 and hsa-miR-718); (66-4) a combination of SEQ ID NOs: 4 and 247

(markers: hsa-miR-4634 and hsa-miR-718); (66-5) a combination of SEQ ID NOs: 3 and 247 (markers: hsa-miR-1307-3p and hsa-miR-718); (67-1) a combination of SEQ ID NOs: 2 and 36 (markers: hsa-miR-4730 and hsa-miR-6726-5p); (67-2) a combination of SEQ ID NOs: 1 and 36 (markers: hsa-miR-4783-3p and hsa-miR-6726-5p); (67-3) a combination of SEQ ID NOs: 237 and 36 (markers: hsa-miR-602 and hsa-miR-6726-5p); (67-4) a combination of SEQ ID NOs: 4 and 36 (markers: hsa-miR-4634 and hsa-miR-6726-5p); (67-5) a combination of SEQ ID NOs: 3 and 36 (markers: hsa-miR-1307-3p and hsa-miR-6726-5p); (68-1) a combination of SEQ ID NOs: 2 and 218 (markers: hsa-miR-4730 and hsa-miR-6784-5p); (68-2) a combination of SEQ ID NOs: 1 and 218 (markers: hsa-miR-4783-3p and hsa-miR-6784-5p); (68-3) a combination of SEQ ID NOs: 237 and 218 (markers: hsa-miR-602 and hsa-miR-6784-5p); (68-4) a combination of SEQ ID NOs: 4 and 218 (markers: hsa-miR-4634 and hsa-miR-6784-5p); (68-5) a combination of SEQ ID NOs: 3 and 218 (markers: hsa-miR-1307-3p and hsa-miR-6784-5p); (69-1) a combination of SEQ ID NOs: 2 and 43 (markers: hsa-miR-4730 and hsa-miR-4763-3p); (69-2) a combination of SEQ ID NOs: 1 and 43 (markers: hsa-miR-4783-3p and hsa-miR-4763-3p); (69-3) a combination of SEQ ID NOs: 237 and 43 (markers: hsa-miR-602 and hsa-miR-4763-3p); (69-4) a combination of SEQ ID NOs: 4 and 43 (markers: hsa-miR-4634 and hsa-miR-4763-3p); (69-5) a combination of SEQ ID NOs: 3 and 43 (markers: hsa-miR-1307-3p and hsa-miR-4763-3p); (70-1) a combination of SEQ ID NOs: 2 and 29 (markers: hsa-miR-4730 and hsa-miR-6757-5p); (70-2) a combination of SEQ ID NOs: 1 and 29 (markers: hsa-miR-4783-3p and hsa-miR-6757-5p); (70-3) a combination of SEQ ID NOs: 237 and 29 (markers: hsa-miR-602 and hsa-miR-6757-5p); (70-4) a combination of SEQ ID NOs: 4 and 29 (markers: hsa-miR-4634 and hsa-miR-6757-5p); (70-5) a combination of SEQ ID NOs: 3 and 29 (markers: hsa-miR-1307-3p and hsa-miR-6757-5p); (71-1) a combination of SEQ ID NOs: 2 and 110 (markers: hsa-miR-4730 and hsa-miR-665); (71-2) a combination of SEQ ID NOs: 1 and 110 (markers: hsa-miR-4783-3p and hsa-miR-665); (71-3) a combination of SEQ ID NOs: 237 and 110 (markers: hsa-miR-602 and hsa-miR-665); (71-4) a combination of SEQ ID NOs: 4 and 110 (markers: hsa-miR-4634 and hsa-miR-665); (71-5) a combination of SEQ ID NOs: 3 and 110 (markers: hsa-miR-1307-3p and hsa-miR-665); (72-1) a combination of SEQ ID NOs: 2 and 20 (markers: hsa-miR-4730 and hsa-miR-1233-5p); (72-2) a combination of SEQ ID NOs: 1 and 20 (markers: hsa-miR-4783-3p and hsa-miR-1233-5p); (72-3) a combination of SEQ ID NOs: 237 and 20 (markers: hsa-miR-602 and hsa-miR-1233-5p); (72-4) a combination of SEQ ID NOs: 4 and 20 (markers: hsa-miR-4634 and hsa-miR-1233-5p); (72-5) a combination of SEQ ID NOs: 3 and 20 (markers: hsa-miR-1307-3p and hsa-miR-1233-5p); (73-1) a combination of SEQ ID NOs: 2 and 157 (markers: hsa-miR-4730 and hsa-miR-1268b); (73-2) a combination of SEQ ID NOs: 1 and 157 (markers: hsa-miR-4783-3p and hsa-miR-1268b); (73-3) a combination of SEQ ID NOs: 237 and 157 (markers: hsa-miR-602 and hsa-miR-1268b); (73-4) a combination of SEQ ID NOs: 4 and 157 (markers: hsa-miR-4634 and hsa-miR-1268b); (73-5) a combination of SEQ ID NOs: 3 and 157 (markers: hsa-miR-1307-3p and hsa-miR-1268b); (74-1) a combination of SEQ ID NOs: 2 and 75 (markers: hsa-miR-4730 and hsa-miR-4675); (74-2) a combination of SEQ ID NOs: 1 and 75 (markers: hsa-miR-4783-3p and hsa-miR-4675); (74-3) a combination of SEQ ID NOs: 237 and 75 (markers: hsa-miR-602 and hsa-miR-4675); (74-4) a combination of SEQ ID NOs: 4 and 75 (markers: hsa-miR-4634 and hsa-miR-4675); (74-5) a combination of SEQ ID NOs: 3 and 75 (markers: hsa-miR-1307-3p and hsa-miR-4675); (75-1) a combination of SEQ ID NOs: 2 and 82 (markers: hsa-miR-4730 and hsa-miR-3620-5p); (75-2) a combination of SEQ ID NOs: 1 and 82 (markers: hsa-miR-4783-3p and hsa-miR-3620-5p); (75-3) a combination of SEQ ID NOs: 237 and 82 (markers: hsa-miR-602 and hsa-miR-3620-5p); (75-4) a combination of SEQ ID NOs: 4 and 82 (markers: hsa-miR-4634 and hsa-miR-3620-5p); (75-5) a combination of SEQ ID NOs: 3 and 82 (markers: hsa-miR-1307-3p and hsa-miR-3620-5p); (76-1) a combination of SEQ ID NOs: 2 and 106 (markers: hsa-miR-4730 and hsa-miR-6800-5p); (76-2) a combination of SEQ ID NOs: 1 and 106 (markers: hsa-miR-4783-3p and hsa-miR-6800-5p); (76-3) a combination of SEQ ID NOs: 237 and 106 (markers: hsa-miR-602 and hsa-miR-6800-5p); (76-4) a combination of SEQ ID NOs: 4

and 106 (markers: hsa-miR-4634 and hsa-miR-6800-5p); (76-5) a combination of SEQ ID NOs: 3 and 106 (markers: hsa-miR-1307-3p and hsa-miR-6800-5p); (77-1) a combination of SEQ ID NOs: 2 and 111 (markers: hsa-miR-4730 and hsa-miR-6778-5p); (77-2) a combination of SEQ ID NOs: 1 and 111 (markers: hsa-miR-4783-3p and hsa-miR-6778-5p); (77-3) a combination of SEQ ID NOs: 237 and 111 (markers: hsa-miR-602 and hsa-miR-6778-5p); (77-4) a combination of SEQ ID NOs: 4 and 111 (markers: hsa-miR-4634 and hsa-miR-6778-5p); (77-5) a combination of SEQ ID NOs: 3 and 111 (markers: hsa-miR-1307-3p and hsa-miR-6778-5p); (78-1) a combination of SEQ ID NOs: 2 and 96 (markers: hsa-miR-4730 and hsa-miR-8059); (78-2) a combination of SEQ ID NOs: 1 and 96 (markers: hsa-miR-4783-3p and hsa-miR-8059); (78-3) a combination of SEQ ID NOs: 237 and 96 (markers: hsa-miR-602 and hsa-miR-8059); (78-4) a combination of SEQ ID NOs: 4 and 96 (markers: hsa-miR-4634 and hsa-miR-8059); (78-5) a combination of SEQ ID NOs: 3 and 96 (markers: hsa-miR-1307-3p and hsa-miR-8059); (79-1) a combination of SEQ ID NOs: 2 and 266 (markers: hsa-miR-4730 and hsa-miR-1908-5p); (79-2) a combination of SEQ ID NOs: 1 and 266 (markers: hsa-miR-4783-3p and hsa-miR-1908-5p); (79-3) a combination of SEQ ID NOs: 237 and 266 (markers: hsa-miR-602 and hsa-miR-1908-5p); (79-4) a combination of SEQ ID NOs: 4 and 266 (markers: hsa-miR-4634 and hsa-miR-1908-5p); (79-5) a combination of SEQ ID NOs: 3 and 266 (markers: hsa-miR-1307-3p and hsa-miR-1908-5p); (80-1) a combination of SEQ ID NOs: 2 and 124 (markers: hsa-miR-4730 and hsa-miR-6798-5p); (80-2) a combination of SEQ ID NOs: 1 and 124 (markers: hsa-miR-4783-3p and hsa-miR-6798-5p); (80-3) a combination of SEQ ID NOs: 237 and 124 (markers: hsa-miR-602 and hsa-miR-6798-5p); (80-4) a combination of SEQ ID NOs: 4 and 124 (markers: hsa-miR-4634 and hsa-miR-6798-5p); (80-5) a combination of SEQ ID NOs: 3 and 124 (markers: hsa-miR-1307-3p and hsa-miR-6798-5p); (81-1) a combination of SEQ ID NOs: 2 and 68 (markers: hsa-miR-4730 and hsa-miR-6879-5p); (81-2) a combination of SEQ ID NOs: 1 and 68 (markers: hsa-miR-4783-3p and hsa-miR-6879-5p); (81-3) a combination of SEQ ID NOs: 237 and 68 (markers: hsa-miR-602 and hsa-miR-6879-5p); (81-4) a combination of SEQ ID NOs: 4 and 68 (markers: hsa-miR-4634 and hsa-miR-6879-5p); (81-5) a combination of SEQ ID NOs: 3 and 68 (markers: hsa-miR-1307-3p and hsa-miR-6879-5p); (82-1) a combination of SEQ ID NOs: 2 and 71 (markers: hsa-miR-4730 and hsa-miR-711); (82-2) a combination of SEQ ID NOs: 1 and 71 (markers: hsa-miR-4783-3p and hsa-miR-711); (82-3) a combination of SEQ ID NOs: 237 and 71 (markers: hsa-miR-602 and hsa-miR-711); (82-4) a combination of SEQ ID NOs: 4 and 71 (markers: hsa-miR-4634 and hsa-miR-711); (82-5) a combination of SEQ ID NOs: 3 and 71 (markers: hsa-miR-1307-3p and hsa-miR-711); (83-1) a combination of SEQ ID NOs: 2 and 35 (markers: hsa-miR-4730 and hsa-miR-4728-5p); (83-2) a combination of SEQ ID NOs: 1 and 35 (markers: hsa-miR-4783-3p and hsa-miR-4728-5p); (83-3) a combination of SEQ ID NOs: 237 and 35 (markers: hsa-miR-602 and hsa-miR-4728-5p); (83-4) a combination of SEQ ID NOs: 4 and 35 (markers: hsa-miR-4634 and hsa-miR-4728-5p); (83-5) a combination of SEQ ID NOs: 3 and 35 (markers: hsa-miR-1307-3p and hsa-miR-4728-5p); (84-1) a combination of SEQ ID NOs: 2 and 173 (markers: hsa-miR-4730 and hsa-miR-3195); (84-2) a combination of SEQ ID NOs: 1 and 173 (markers: hsa-miR-4783-3p and hsa-miR-3195); (84-3) a combination of SEQ ID NOs: 237 and 173 (markers: hsa-miR-602 and hsa-miR-3195); (84-4) a combination of SEQ ID NOs: 4 and 173 (markers: hsa-miR-4634 and hsa-miR-3195); (84-5) a combination of SEQ ID NOs: 3 and 173 (markers: hsa-miR-1307-3p and hsa-miR-3195); (85-1) a combination of SEQ ID NOs: 2 and 5 (markers: hsa-miR-4730 and hsa-miR-663a); (85-2) a combination of SEQ ID NOs: 1 and 5 (markers: hsa-miR-4783-3p and hsa-miR-663a); (85-3) a combination of SEQ ID NOs: 237 and 5 (markers: hsa-miR-602 and hsa-miR-663a); (85-4) a combination of SEQ ID NOs: 4 and 5 (markers: hsa-miR-4634 and hsa-miR-663a); (85-5) a combination of SEQ ID NOs: 3 and 5 (markers: hsa-miR-1307-3p and hsa-miR-663a); (86-1) a combination of SEQ ID NOs: 2 and 851 (markers: hsa-miR-4730 and hsa-miR-6089); (86-2) a combination of SEQ ID NOs: 1 and 851 (markers: hsa-miR-4783-3p and hsa-miR-6089); (86-3) a combination of SEQ ID NOs: 237 and 851 (markers: hsa-miR-602 and hsa-miR-6089); (86-4) a combination of SEQ ID NOs: 4 and 851

(markers: hsa-miR-4634 and hsa-miR-6089); (86-5) a combination of SEQ ID NOs: 3 and 851 (markers: hsa-miR-1307-3p and hsa-miR-6089); (87-1) a combination of SEQ ID NOs: 2 and 852 (markers: hsa-miR-4730 and hsa-miR-6816-5p); (87-2) a combination of SEQ ID NOs: 1 and 852 (markers: hsa-miR-4783-3p and hsa-miR-6816-5p); (87-3) a combination of SEQ ID NOs: 237 and 852 (markers: hsa-miR-602 and hsa-miR-6816-5p); (87-4) a combination of SEQ ID NOs: 4 and 852 (markers: hsa-miR-4634 and hsa-miR-6816-5p); (87-5) a combination of SEQ ID NOs: 3 and 852 (markers: hsa-miR-1307-3p and hsa-miR-6816-5p); (88-1) a combination of SEQ ID NOs: 2 and 30 (markers: hsa-miR-4730 and hsa-miR-6756-5p); (88-2) a combination of SEQ ID NOs: 1 and 30 (markers: hsa-miR-4783-3p and hsa-miR-6756-5p); (88-3) a combination of SEQ ID NOs: 237 and 30 (markers: hsa-miR-602 and hsa-miR-6756-5p); (88-4) a combination of SEQ ID NOs: 4 and 30 (markers: hsa-miR-4634 and hsa-miR-6756-5p); (88-5) a combination of SEQ ID NOs: 3 and 30 (markers: hsa-miR-1307-3p and hsa-miR-6756-5p); (89-1) a combination of SEQ ID NOs: 2 and 93 (markers: hsa-miR-4730 and hsa-miR-6861-5p); (89-2) a combination of SEQ ID NOs: 1 and 93 (markers: hsa-miR-4783-3p and hsa-miR-6861-5p); (89-3) a combination of SEQ ID NOs: 237 and 93 (markers: hsa-miR-602 and hsa-miR-6861-5p); (89-4) a combination of SEQ ID NOs: 4 and 93 (markers: hsa-miR-4634 and hsa-miR-6861-5p); (89-5) a combination of SEQ ID NOs: 3 and 93 (markers: hsa-miR-1307-3p and hsa-miR-6861-5p); (90-1) a combination of SEQ ID NOs: 2 and 27 (markers: hsa-miR-4730 and hsa-miR-1246); (90-2) a combination of SEQ ID NOs: 1 and 27 (markers: hsa-miR-4783-3p and hsa-miR-1246); (90-3) a combination of SEQ ID NOs: 237 and 27 (markers: hsa-miR-602 and hsa-miR-1246); (90-4) a combination of SEQ ID NOs: 4 and 27 (markers: hsa-miR-4634 and hsa-miR-1246); (90-5) a combination of SEQ ID NOs: 27 and 208 (markers: hsa-miR-1246 and hsa-miR-1343-5p); (91-1) a combination of SEQ ID NOs: 2 and 853 (markers: hsa-miR-4730 and hsa-miR-4466); (91-2) a combination of SEQ ID NOs: 1 and 853 (markers: hsa-miR-4783-3p and hsa-miR-4466); (91-3) a combination of SEQ ID NOs: 237 and 853 (markers: hsa-miR-602 and hsa-miR-4466); (91-4) a combination of SEQ ID NOs: 4 and 853 (markers: hsa-miR-4634 and hsa-miR-4466); (91-5) a combination of SEQ ID NOs: 3 and 853 (markers: hsa-miR-1307-3p and hsa-miR-4466); (92-1) a combination of SEQ ID NOs: 2 and 238 (markers: hsa-miR-4730 and hsa-miR-423-5p); (92-2) a combination of SEQ ID NOs: 1 and 238 (markers: hsa-miR-4783-3p and hsa-miR-423-5p); (92-3) a combination of SEQ ID NOs: 237 and 238 (markers: hsa-miR-602 and hsa-miR-423-5p); (92-4) a combination of SEQ ID NOs: 4 and 238 (markers: hsa-miR-4634 and hsa-miR-423-5p); (92-5) a combination of SEQ ID NOs: 3 and 238 (markers: hsa-miR-1307-3p and hsa-miR-423-5p); (93-1) a combination of SEQ ID NOs: 2 and 130 (markers: hsa-miR-4730 and hsa-miR-6075); (93-2) a combination of SEQ ID NOs: 1 and 130 (markers: hsa-miR-4783-3p and hsa-miR-6075); (93-3) a combination of SEQ ID NOs: 237 and 130 (markers: hsa-miR-602 and hsa-miR-6075); (93-4) a combination of SEQ ID NOs: 4 and 130 (markers: hsa-miR-4634 and hsa-miR-6075); (93-5) a combination of SEQ ID NOs: 3 and 130 (markers: hsa-miR-1307-3p and hsa-miR-6075); (94-1) a combination of SEQ ID NOs: 2 and 177 (markers: hsa-miR-4730 and hsa-miR-7108-5p); (94-2) a combination of SEQ ID NOs: 1 and 177 (markers: hsa-miR-4783-3p and hsa-miR-7108-5p); (94-3) a combination of SEQ ID NOs: 237 and 177 (markers: hsa-miR-602 and hsa-miR-7108-5p); (94-4) a combination of SEQ ID NOs: 4 and 177 (markers: hsa-miR-4634 and hsa-miR-7108-5p); (94-5) a combination of SEQ ID NOs: 3 and 177 (markers: hsa-miR-1307-3p and hsa-miR-7108-5p); (95-1) a combination of SEQ ID NOs: 2 and 64 (markers: hsa-miR-4730 and hsa-miR-6511a-5p); (95-2) a combination of SEQ ID NOs: 1 and 64 (markers: hsa-miR-4783-3p and hsa-miR-6511a-5p); (95-3) a combination of SEQ ID NOs: 237 and 64 (markers: hsa-miR-602 and hsa-miR-6511a-5p); (95-4) a combination of SEQ ID NOs: 4 and 64 (markers: hsa-miR-4634 and hsa-miR-6511a-5p); (95-5) a combination of SEQ ID NOs: 3 and 64 (markers: hsa-miR-1307-3p and hsa-miR-6511a-5p); (96-1) a combination of SEQ ID NOs: 2 and 114 (markers: hsa-miR-4730 and hsa-miR-211-3p); (96-2) a combination of SEQ ID NOs: 1 and 114 (markers: hsa-miR-4783-3p and hsa-miR-211-3p); (96-3) a combination of SEQ ID NOs: 237 and 114 (markers: hsa-miR-602 and hsa-miR-211-3p); (96-4) a combination of SEQ ID NOs: 4

and 114 (markers: hsa-miR-4634 and hsa-miR-211-3p); (96-5) a combination of SEQ ID NOs: 3 and 114 (markers: hsa-miR-1307-3p and hsa-miR-211-3p); (97-1) a combination of SEQ ID NOs: 2 and 119 (markers: hsa-miR-4730 and hsa-miR-7110-5p); (97-2) a combination of SEQ ID NOs: 1 and 119 (markers: hsa-miR-4783-3p and hsa-miR-7110-5p); (97-3) a combination of SEQ ID NOs: 237 and 119 (markers: hsa-miR-602 and hsa-miR-7110-5p); (97-4) a combination of SEQ ID NOs: 4 and 119 (markers: hsa-miR-4634 and hsa-miR-7110-5p); (97-5) a combination of SEQ ID NOs: 3 and 119 (markers: hsa-miR-1307-3p and hsa-miR-7110-5p); (98-1) a combination of SEQ ID NOs: 2 and 135 (markers: hsa-miR-4730 and hsa-miR-6870-5p); (98-2) a combination of SEQ ID NOs: 1 and 135 (markers: hsa-miR-4783-3p and hsa-miR-6870-5p); (98-3) a combination of SEQ ID NOs: 237 and 135 (markers: hsa-miR-602 and hsa-miR-6870-5p); (98-4) a combination of SEQ ID NOs: 4 and 135 (markers: hsa-miR-4634 and hsa-miR-6870-5p); (98-5) a combination of SEQ ID NOs: 3 and 135 (markers: hsa-miR-1307-3p and hsa-miR-6870-5p); (99-1) a combination of SEQ ID NOs: 2 and 243 (markers: hsa-miR-4730 and hsa-miR-486-5p); (99-2) a combination of SEQ ID NOs: 1 and 243 (markers: hsa-miR-4783-3p and hsa-miR-486-5p); (99-3) a combination of SEQ ID NOs: 237 and 243 (markers: hsa-miR-602 and hsa-miR-486-5p); (99-4) a combination of SEQ ID NOs: 4 and 243 (markers: hsa-miR-4634 and hsa-miR-486-5p); (99-5) a combination of SEQ ID NOs: 3 and 243 (markers: hsa-miR-1307-3p and hsa-miR-486-5p); (100-1) a combination of SEQ ID NOs: 2 and 122 (markers: hsa-miR-4730 and hsa-miR-4792); (100-2) a combination of SEQ ID NOs: 1 and 122 (markers: hsa-miR-4783-3p and hsa-miR-4792); (100-3) a combination of SEQ ID NOs: 237 and 122 (markers: hsa-miR-602 and hsa-miR-4792); (100-4) a combination of SEQ ID NOs: 4 and 122 (markers: hsa-miR-4634 and hsa-miR-4792); (100-5) a combination of SEQ ID NOs: 3 and 122 (markers: hsa-miR-1307-3p and hsa-miR-4792); (101-1) a combination of SEQ ID NOs: 2 and 260 (markers: hsa-miR-4730 and hsa-miR-4687-3p); (101-2) a combination of SEQ ID NOs: 1 and 260 (markers: hsa-miR-4783-3p and hsa-miR-4687-3p); (101-3) a combination of SEQ ID NOs: 237 and 260 (markers: hsa-miR-602 and hsa-miR-4687-3p); (101-4) a combination of SEQ ID NOs: 4 and 260 (markers: hsa-miR-4634 and hsa-miR-4687-3p); (101-5) a combination of SEQ ID NOs: 3 and 260 (markers: hsa-miR-1307-3p and hsa-miR-4687-3p); (102-1) a combination of SEQ ID NOs: 2 and 59 (markers: hsa-miR-4730 and hsa-miR-1238-5p); (102-2) a combination of SEQ ID NOs: 1 and 59 (markers: hsa-miR-4783-3p and hsa-miR-1238-5p); (102-3) a combination of SEQ ID NOs: 237 and 59 (markers: hsa-miR-602 and hsa-miR-1238-5p); (102-4) a combination of SEQ ID NOs: 4 and 59 (markers: hsa-miR-4634 and hsa-miR-1238-5p); (102-5) a combination of SEQ ID NOs: 3 and 59 (markers: hsa-miR-1307-3p and hsa-miR-1238-5p); (103-1) a combination of SEQ ID NOs: 2 and 854 (markers: hsa-miR-4730 and hsa-miR-4488); (103-2) a combination of SEQ ID NOs: 1 and 854 (markers: hsa-miR-4783-3p and hsa-miR-4488); (103-3) a combination of SEQ ID NOs: 237 and 854 (markers: hsa-miR-602 and hsa-miR-4488); (103-4) a combination of SEQ ID NOs: 4 and 854 (markers: hsa-miR-4634 and hsa-miR-4488); (103-5) a combination of SEQ ID NOs: 3 and 854 (markers: hsa-miR-1307-3p and hsa-miR-4488); (104-1) a combination of SEQ ID NOs: 2 and 132 (markers: hsa-miR-4730 and hsa-miR-6891-5p); (104-2) a combination of SEQ ID NOs: 1 and 132 (markers: hsa-miR-4783-3p and hsa-miR-6891-5p); (104-3) a combination of SEQ ID NOs: 237 and 132 (markers: hsa-miR-602 and hsa-miR-6891-5p); (104-4) a combination of SEQ ID NOs: 4 and 132 (markers: hsa-miR-4634 and hsa-miR-6891-5p); (104-5) a combination of SEQ ID NOs: 3 and 132 (markers: hsa-miR-1307-3p and hsa-miR-6891-5p); (105-1) a combination of SEQ ID NOs: 2 and 181 (markers: hsa-miR-4730 and hsa-miR-2861); (105-2) a combination of SEQ ID NOs: 1 and 181 (markers: hsa-miR-4783-3p and hsa-miR-2861); (105-3) a combination of SEQ ID NOs: 237 and 181 (markers: hsa-miR-602 and hsa-miR-2861); (105-4) a combination of SEQ ID NOs: 4 and 181 (markers: hsa-miR-4634 and hsa-miR-2861); (105-5) a combination of SEQ ID NOs: 3 and 181 (markers: hsa-miR-1307-3p and hsa-miR-2861); (106-1) a combination of SEQ ID NOs: 2 and 79 (markers: hsa-miR-4730 and hsa-miR-4665-5p); (106-2) a combination of SEQ ID NOs: 1 and 79 (markers: hsa-miR-4783-3p and hsa-miR-4665-5p); (106-3) a combination of SEQ ID NOs: 237

and 79 (markers: hsa-miR-602 and hsa-miR-4665-5p); (106-4) a combination of SEQ ID NOs: 4 and 79 (markers: hsa-miR-4634 and hsa-miR-4665-5p); (106-5) a combination of SEQ ID NOs: 3 and 79 (markers: hsa-miR-1307-3p and hsa-miR-4665-5p); (107-1) a combination of SEQ ID NOs: 2 and 133 (markers: hsa-miR-4730 and hsa-miR-4745-5p); (107-2) a combination of SEQ ID NOs: 1 and 133 (markers: hsa-miR-4783-3p and hsa-miR-4745-5p); (107-3) a combination of SEQ ID NOs: 237 and 133 (markers: hsa-miR-602 and hsa-miR-4745-5p); (107-4) a combination of SEQ ID NOs: 4 and 133 (markers: hsa-miR-4634 and hsa-miR-4745-5p); (107-5) a combination of SEQ ID NOs: 3 and 133 (markers: hsa-miR-1307-3p and hsa-miR-4745-5p); (108-1) a combination of SEQ ID NOs: 2 and 41 (markers: hsa-miR-4730 and hsa-miR-6858-5p); (108-2) a combination of SEQ ID NOs: 1 and 41 (markers: hsa-miR-4783-3p and hsa-miR-6858-5p); (108-3) a combination of SEQ ID NOs: 237 and 41 (markers: hsa-miR-602 and hsa-miR-6858-5p); (108-4) a combination of SEQ ID NOs: 4 and 41 (markers: hsa-miR-4634 and hsa-miR-6858-5p); (108-5) a combination of SEQ ID NOs: 3 and 41 (markers: hsa-miR-1307-3p and hsa-miR-6858-5p); (109-1) a combination of SEQ ID NOs: 2 and 139 (markers: hsa-miR-4730 and hsa-miR-6825-5p); (109-2) a combination of SEQ ID NOs: 1 and 139 (markers: hsa-miR-4783-3p and hsa-miR-6825-5p); (109-3) a combination of SEQ ID NOs: 237 and 139 (markers: hsa-miR-602 and hsa-miR-6825-5p); (109-4) a combination of SEQ ID NOs: 4 and 139 (markers: hsa-miR-4634 and hsa-miR-6825-5p); (109-5) a combination of SEQ ID NOs: 3 and 139 (markers: hsa-miR-1307-3p and hsa-miR-6825-5p); (110-1) a combination of SEQ ID NOs: 2 and 118 (markers: hsa-miR-4730 and hsa-miR-614); (110-2) a combination of SEQ ID NOs: 1 and 118 (markers: hsa-miR-4783-3p and hsa-miR-614); (110-3) a combination of SEQ ID NOs: 237 and 118 (markers: hsa-miR-602 and hsa-miR-614); (110-4) a combination of SEQ ID NOs: 4 and 118 (markers: hsa-miR-4634 and hsa-miR-614); (110-5) a combination of SEQ ID NOs: 3 and 118 (markers: hsa-miR-1307-3p and hsa-miR-614); (111-1) a combination of SEQ ID NOs: 2 and 86 (markers: hsa-miR-4730 and hsa-miR-1343-3p); (111-2) a combination of SEQ ID NOs: 1 and 86 (markers: hsa-miR-4783-3p and hsa-miR-1343-3p); (111-3) a combination of SEQ ID NOs: 237 and 86 (markers: hsa-miR-602 and hsa-miR-1343-3p); (111-4) a combination of SEQ ID NOs: 4 and 86 (markers: hsa-miR-4634 and hsa-miR-1343-3p); (111-5) a combination of SEQ ID NOs: 3 and 86 (markers: hsa-miR-1307-3p and hsa-miR-1343-3p); (112-1) a combination of SEQ ID NOs: 2 and 60 (markers: hsa-miR-4730 and hsa-miR-365a-5p); (112-2) a combination of SEQ ID NOs: 1 and 60 (markers: hsa-miR-4783-3p and hsa-miR-365a-5p); (112-3) a combination of SEQ ID NOs: 237 and 60 (markers: hsa-miR-602 and hsa-miR-365a-5p); (112-4) a combination of SEQ ID NOs: 4 and 60 (markers: hsa-miR-4634 and hsa-miR-365a-5p); (112-5) a combination of SEQ ID NOs: 3 and 60 (markers: hsa-miR-1307-3p and hsa-miR-365a-5p); (113-1) a combination of SEQ ID NOs: 2 and 116 (markers: hsa-miR-4730 and hsa-miR-4750-5p); (113-2) a combination of SEQ ID NOs: 1 and 116 (markers: hsa-miR-4783-3p and hsa-miR-4750-5p); (113-3) a combination of SEQ ID NOs: 237 and 116 (markers: hsa-miR-602 and hsa-miR-4750-5p); (113-4) a combination of SEQ ID NOs: 4 and 116 (markers: hsa-miR-4634 and hsa-miR-4750-5p); (113-5) a combination of SEQ ID NOs: 3 and 116 (markers: hsa-miR-1307-3p and hsa-miR-4750-5p); (114-1) a combination of SEQ ID NOs: 2 and 160 (markers: hsa-miR-4730 and hsa-miR-6732-5p); (114-2) a combination of SEQ ID NOs: 1 and 160 (markers: hsa-miR-4783-3p and hsa-miR-6732-5p); (114-3) a combination of SEQ ID NOs: 237 and 160 (markers: hsa-miR-602 and hsa-miR-6732-5p); (114-4) a combination of SEQ ID NOs: 4 and 160 (markers: hsa-miR-4634 and hsa-miR-6732-5p); (114-5) a combination of SEQ ID NOs: 3 and 160 (markers: hsa-miR-1307-3p and hsa-miR-6732-5p); (115-1) a combination of SEQ ID NOs: 2 and 38 (markers: hsa-miR-4730 and hsa-miR-149-3p); (115-2) a combination of SEQ ID NOs: 1 and 38 (markers: hsa-miR-4783-3p and hsa-miR-149-3p); (115-3) a combination of SEQ ID NOs: 237 and 38 (markers: hsa-miR-602 and hsa-miR-149-3p); (115-4) a combination of SEQ ID NOs: 4 and 38 (markers: hsa-miR-4634 and hsa-miR-149-3p); (115-5) a combination of SEQ ID NOs: 3 and 38 (markers: hsa-miR-1307-3p and hsa-miR-149-3p); (116-1) a combination of SEQ ID NOs: 2 and 99 (markers: hsa-miR-4730 and hsa-miR-4497); (116-2) a combination of SEQ ID NOs: 1 and 99

(markers: hsa-miR-4783-3p and hsa-miR-4497); (116-3) a combination of SEQ ID NOs: 237 and 99 (markers: hsa-miR-602 and hsa-miR-4497); (116-4) a combination of SEQ ID NOs: 4 and 99 (markers: hsa-miR-4634 and hsa-miR-4497); (116-5) a combination of SEQ ID NOs: 3 and 99 (markers: hsa-miR-1307-3p and hsa-miR-4497); (117-1) a combination of SEQ ID NOs: 2 and 104 (markers: hsa-miR-4730 and hsa-miR-6848-5p); (117-2) a combination of SEQ ID NOs: 1 and 104 (markers: hsa-miR-4783-3p and hsa-miR-6848-5p); (117-3) a combination of SEQ ID NOs: 237 and 104 (markers: hsa-miR-602 and hsa-miR-6848-5p); (117-4) a combination of SEQ ID NOs: 4 and 104 (markers: hsa-miR-4634 and hsa-miR-6848-5p); and (117-5) a combination of SEQ ID NOs: 3 and 104 (markers: hsa-miR-1307-3p and hsa-miR-6848-5p).

(327) The kit or the device of the present invention may also comprise a polynucleotide(s) that is/are already known or that will be found in the future, to enable detection of breast cancer, in addition to the polynucleotide(s) (which may include variant(s), fragment(s), or derivative(s)) according to the present invention

(328) The kit of the present invention may also comprise an antibody for measuring marker(s) for breast cancer examination known in the art, such as CEA, CA-15-3, and CA27-29, in addition to the polynucleotide(s) as described above.

(329) The polynucleotides described above contained in the kit of the present invention may be packaged in different containers either individually or in any combination.

(330) The kit of the present invention may comprise a kit for extracting nucleic acids (e.g., total RNA) from body fluids, cells, or tissues, a fluorescent material for labeling, an enzyme and a medium for nucleic acid amplification, an instruction manual, etc.

(331) The device of the present invention is a device for cancer marker measurement in which nucleic acids such as the polynucleotides according to the present invention described above, variants thereof, derivatives thereof, or fragments thereof are bound or attached to, for example, a solid phase. Examples of the material for the solid phase include plastics, paper, glass, and silicon. The material for the solid phase is preferably a plastic from the viewpoint of easy processability. The solid phase has any shape and is, for example, square, round, reed-shaped, or film-shaped. The device of the present invention includes, for example, a device for measurement by a hybridization technique. Specific examples thereof include blotting devices and nucleic acid arrays (e.g., microarrays, DNA chips, and RNA chips).

(332) The nucleic acid array technique is a technique which involves binding or attaching the nucleic acids one by one by use of a method [e.g., a method of spotting the nucleic acids using a high-density dispenser called spotter or arrayer onto the surface of the solid phase surface-treated, if necessary, by coating with L-lysine or the introduction of a functional group such as an amino group or a carboxyl group; a method of spraying the nucleic acids onto the solid phase using an inkjet which injects very small liquid droplets by a piezoelectric element or the like from a nozzle; or a method of sequentially synthesizing nucleotides on the solid phase] to prepare an array such as a chip and measuring a target nucleic acid through the use of hybridization using this array.

(333) The kit or the device of the present invention comprises nucleic acids capable of specifically binding to the polynucleotides of at least one, preferably at least two, more preferably at least three, most preferably at least five to all of the breast cancer marker miRNAs, respectively, of the group 1 described above. The kit or the device of the present invention may optionally further comprise nucleic acids capable of specifically binding to the polynucleotides of at least one, preferably at least two, more preferably at least three, most preferably at least five to all of the breast cancer marker miRNAs, respectively, of the group 2 described above. The kit or the device of the present invention may optionally further comprise nucleic acids capable of specifically binding to the polynucleotides of at least one, preferably at least two, more preferably at least three, most preferably at least five to all of the breast cancer marker miRNAs, respectively, of the group 3 described above. The kit or the device of the present invention can be used for detecting breast cancer as described in Section 4 below.

(334) 4. Method for Detecting Breast Cancer

(335) The present invention further provides a method for detecting breast cancer, comprising using the kit or the device of the present invention (comprising the aforementioned nucleic acid(s) that can be used in the present invention) as described in Section “3. Kit or device for detection of breast cancer” to measure an expression level of one or more breast cancer-derived genes represented by: expression level(s) of breast cancer-derived genes selected from the following group of miRNAs, i.e., miR-4783-3p, miR-4730, miR-1307-3p, miR-4634, miR-663a, miR-4532, miR-7704, miR-3178, miR-6729-5p, miR-6090, miR-4732-5p, miR-3184-5p, miR-6727-5p, miR-6088, miR-4674, miR-8073, miR-4787-5p, miR-1469, miR-125a-3p, miR-1233-5p, miR-885-3p, miR-6802-5p, miR-328-5p, miR-6787-5p, miR-8069, miR-6875-5p, miR-1246, miR-4734, miR-6757-5p, miR-6756-5p, miR-3665, miR-6836-3p, miR-6821-5p, miR-6805-5p, miR-4728-5p, miR-6726-5p, miR-197-5p, miR-149-3p, miR-6850-5p, miR-4476, miR-6858-5p, miR-564, miR-4763-3p, miR-575, miR-6771-5p, miR-1231, miR-1908-3p, miR-150-3p, miR-3937, miR-887-3p, miR-3940-5p, miR-4741, miR-6808-5p, miR-6869-5p, miR-5090, miR-615-5p, miR-8072, miR-128-1-5p, miR-1238-5p, miR-365a-5p, miR-204-3p, miR-4492, miR-6785-5p, miR-6511a-5p, miR-4525, miR-1915-5p, miR-3180, miR-6879-5p, miR-1199-5p, miR-6746-5p, miR-711, miR-663b, miR-4707-3p, miR-6893-5p, miR-4675, miR-4638-5p, miR-4651, miR-6087, miR-4665-5p, miR-4758-5p, miR-6887-5p, miR-3620-5p, miR-1909-3p, miR-7641, miR-6724-5p, miR-1343-3p, miR-6780b-5p, miR-4484, miR-4690-5p, miR-4429, miR-1227-5p, miR-4725-3p, miR-6861-5p, miR-6812-5p, miR-3197, miR-8059, miR-3185, miR-4706, miR-4497, miR-3131, miR-6806-5p, miR-187-5p, miR-3180-3p, miR-6848-5p, miR-6820-5p, miR-6800-5p, miR-6717-5p, miR-6795-5p, miR-4632-5p, miR-665, miR-6778-5p, miR-3663-3p, miR-4689, miR-211-3p, miR-6511b-5p, miR-4750-5p, miR-6126, miR-614, miR-7110-5p, miR-744-5p, miR-6769a-5p, miR-4792, miR-5787, miR-6798-5p, miR-6781-5p, miR-4419b, miR-4446-3p, miR-4259, miR-5572, miR-6075, miR-296-3p, miR-6891-5p, miR-4745-5p, miR-6775-5p, miR-6870-5p, miR-920, miR-4530, miR-6819-5p, miR-6825-5p, miR-7847-3p, miR-6131, miR-4433-3p, miR-1228-5p, miR-6743-5p, miR-1268a, miR-3917, miR-6786-5p, miR-3154, miR-638, miR-6741-5p, miR-6889-5p, miR-6840-3p, miR-6510-5p, miR-3188, miR-551b-5p, miR-5001-5p, miR-1268b, miR-7107-5p, miR-6824-5p, miR-6732-5p, miR-371a-5p, miR-6794-5p, miR-6779-5p, miR-4271, miR-5195-3p, miR-6762-5p, miR-939-5p, miR-1247-3p, miR-6777-5p, miR-6722-3p, miR-3656, miR-4688, miR-3195, miR-6766-5p, miR-4447, miR-4656, miR-7108-5p, miR-3191-3p, miR-1273g-3p, miR-4463, miR-2861, miR-3196, miR-6877-5p, miR-3679-5p, miR-4442, miR-6789-5p, miR-6782-5p, miR-486-3p, miR-6085, miR-4746-3p, miR-619-5p, miR-937-5p, miR-6803-5p, miR-4298, miR-4454, miR-4459, miR-7150, miR-6880-5p, miR-4449, miR-8063, miR-4695-5p, miR-6132, miR-6829-5p, miR-4486, miR-6805-3p, miR-6826-5p, miR-4508, miR-1343-5p, miR-7114-5p, miR-3622a-5p, miR-6765-5p, miR-7845-5p, miR-3960, miR-6749-5p, miR-1260b, miR-6799-5p, miR-4723-5p, miR-6784-5p, miR-5100, miR-6769b-5p, miR-1207-5p, miR-642a-3p, miR-4505, miR-4270, miR-6721-5p, miR-7111-5p, miR-6791-5p, miR-7109-5p, miR-4258, miR-6515-3p, miR-6851-5p, miR-6125, miR-4749-5p, miR-4726-5p, miR-4513, miR-6089, miR-6816-5p, miR-4466, miR-4488, miR-6752-5p and miR-4739; and optionally expression level(s) of breast cancer-derived gene(s) selected from the following group of miRNA: i.e., miR-760, miR-602, miR-423-5p, miR-92a-2-5p, miR-16-5p, miR-451a, miR-135a-3p, miR-486-5p, miR-4257, miR-92b-5p, miR-1915-3p, miR-718, miR-940, miR-296-5p, miR-23b-3p and miR-92a-3p, and optionally expression level(s) of breast cancer-derived gene(s) selected from the following group of miRNA: i.e., miR-658, miR-6842-5p, miR-6124, miR-6765-3p, miR-7106-5p, miR-4534, miR-92b-3p, miR-3135b, miR-4687-3p, miR-762, miR-3619-3p, miR-4467, miR-557, miR-1237-5p, miR-1908-5p, miR-4286, miR-6885-5p, and miR-6763-5p in a sample in vitro, further comparing, for example, the expression level(s) of the aforementioned gene(s) in the sample (e.g., blood, serum, or plasma) collected from a subject suspected of having breast cancer with a control expression level in the sample collected from a healthy subject (including a non-breast cancer patient(s)), and evaluating the subject as

having breast cancer when the expression level(s) of the target nucleic acid(s) is different between the samples.

(336) This method of the present invention enables limitedly-invasive early diagnosis of the breast cancer with high sensitivity and specificity and thereby brings about early treatment and improved prognosis. In addition, exacerbation of the disease or the effectiveness of surgical, radiotherapeutic, and chemotherapeutic treatments can be monitored.

(337) The method for extracting the breast cancer-derived gene from the sample such as blood, serum, or plasma according to the present invention is particularly preferably a method in which the breast cancer-derived gene is prepared by the addition of a reagent for RNA extraction in 3D-Gener™ RNA extraction reagent from liquid sample kit (Toray Industries, Inc.). A general acidic phenol method (acid guanidinium-phenol-chloroform (AGPC)) may be used, or Trizol™ (Life Technologies Corp.) may be used. The breast cancer-derived gene(s) may be prepared by the addition of a reagent for RNA extraction containing acidic phenol, such as Trizol (Life Technologies Corp.) or Isogen (Nippon Gene Co., Ltd., Japan). Alternatively, a kit such as miRNeasy® Mini Kit (Qiagen N.V.) can be used, though the method is not limited thereto.

(338) The present invention also provides use of the kit or the device of the present invention for detecting in vitro an expression product(s) of a breast cancer-derived miRNA gene in a sample derived from a subject.

(339) In the method of the present invention, the used kit or the device comprises a single polynucleotide or any possible combination of polynucleotides that can be used in the present invention as described above.

(340) In the detection or (genetic) diagnosis of breast cancer according to the present invention, each polynucleotide contained in the kit or the device of the present invention can be used as a probe or a primer. In the case of using the polynucleotide as a primer, TaqMan™ MicroRNA Assays from Life Technologies Corp., miScript PCR System from Qiagen N.V., or the like can be used, though the method is not limited thereto.

(341) The polynucleotide contained in the kit or the device of the present invention can be used as a primer or a probe according to a routine method in a method known in the art for specifically detecting the particular gene, for example, a hybridization technique such as Northern blot, Southern blot, in situ hybridization, Northern hybridization, or Southern hybridization, or a quantitative amplification technique such as quantitative RT-PCR. A body fluid such as blood, serum, plasma, or urine from a subject is collected as a sample to be assayed according to the type of the detection method used. Alternatively, total RNA prepared from such a body fluid by the method described above may be used, and various polynucleotides including cDNA prepared on the basis of the RNA may be used.

(342) The kit or the device of the present invention is useful for the diagnosis of breast or the detection of the presence or absence of breast cancer. Specifically, the detection of breast cancer using the kit or the device can be performed by detecting in vitro an expression level of a gene using the nucleic acid probe or the primer contained in the kit or the device in a sample such as blood, serum, plasma, or urine from a subject suspected of having breast cancer. The subject suspected of having breast cancer can be evaluated as having breast cancer when the expression level(s) of a target miRNA marker(s) measured using a polynucleotide(s) (including variant(s), fragment(s), or derivative(s) thereof) consisting of a nucleotide sequence(s) represented by at least one of SEQ ID NOs: 1 to 235, and 851 to 856 or a complementary sequence(s) thereof, and optionally a nucleotide sequence(s) represented by one or more of SEQ ID NOs: 236 to 251 or a complementary sequence(s) thereof, and optionally a nucleotide sequence(s) represented by one or more of SEQ ID NOs: 252 to 269 or a complementary sequence thereof in the sample such as blood, serum, plasma, or urine of the subject has a statistically significantly higher or lower than the expression level(s) thereof in the sample such as blood, serum, or plasma, or urine of a healthy subject.

(343) The method of the present invention can be combined with a diagnostic imaging method such as mammography, ultrasonography (echo examination), CT, MRI, abdominal ultrasonography, bone scintigraphy, or PET, or pathological examination which involves analyzing a lesion tissue under a microscope. The method of the present invention is capable of specifically detecting breast cancer and can substantially discriminate breast cancer from the other cancers.

(344) The method for detecting the absence of an expression product(s) of a breast cancer-derived gene(s) or the presence of the expression product(s) of a breast cancer-derived gene in a sample using the kit or the device of the present invention comprises; collecting a body fluid such as blood, serum, plasma, or urine of a subject; measuring the expression level(s) of the target gene(s) contained therein using one or more polynucleotides (including variant(s), fragment(s), or derivative(s)) selected from the polynucleotide group of the present invention; and evaluating the presence or absence of breast cancer or to detect breast cancer. The method for detecting breast cancer according to the present invention can also evaluate or diagnose, for example, the presence or absence of amelioration of the disease or the degree of amelioration thereof in a breast cancer patient in the case that a therapeutic drug is administered to the patient for amelioration of the disease.

(345) The method of the present invention can comprise, for example, the following steps (a), (b), and (c): (a) a step of contacting in vitro a sample from a subject with a polynucleotide(s) contained in the kit or the device of the present invention in vitro; (b) a step of measuring an expression level(s) of the target nucleic acid(s) in the sample using the polynucleotide(s) as a nucleic acid probe(s) or a primer(s); and (c) a step of evaluating the presence or absence of breast cancer (cells) in the subject on the basis of the measurement results in the step (b).

(346) Specifically, the present invention provides a method for detecting breast cancer, comprising measuring an expression level(s) of a target nucleic acid(s) in a sample of a subject using nucleic acid(s) capable of specifically binding to at least one (preferably at least two) polynucleotides selected from the group consisting of miR-4783-3p, miR-4730, miR-1307-3p, miR-4634, miR-663a, miR-4532, miR-7704, miR-3178, miR-6729-5p, miR-6090, miR-4732-5p, miR-3184-5p, miR-6727-5p, miR-6088, miR-4674, miR-8073, miR-4787-5p, miR-1469, miR-125a-3p, miR-1233-5p, miR-885-3p, miR-6802-5p, miR-328-5p, miR-6787-5p, miR-8069, miR-6875-5p, miR-1246, miR-4734, miR-6757-5p, miR-6756-5p, miR-3665, miR-6836-3p, miR-6821-5p, miR-6805-5p, miR-4728-5p, miR-6726-5p, miR-197-5p, miR-149-3p, miR-6850-5p, miR-4476, miR-6858-5p, miR-564, miR-4763-3p, miR-575, miR-6771-5p, miR-1231, miR-1908-3p, miR-150-3p, miR-3937, miR-887-3p, miR-3940-5p, miR-4741, miR-6808-5p, miR-6869-5p, miR-5090, miR-615-5p, miR-8072, miR-128-1-5p, miR-1238-5p, miR-365a-5p, miR-204-3p, miR-4492, miR-6785-5p, miR-6511a-5p, miR-4525, miR-1915-5p, miR-3180, miR-6879-5p, miR-1199-5p, miR-6746-5p, miR-711, miR-663b, miR-4707-3p, miR-6893-5p, miR-4675, miR-4638-5p, miR-4651, miR-6087, miR-4665-5p, miR-4758-5p, miR-6887-5p, miR-3620-5p, miR-1909-3p, miR-7641, miR-6724-5p, miR-1343-3p, miR-6780b-5p, miR-4484, miR-4690-5p, miR-4429, miR-1227-5p, miR-4725-3p, miR-6861-5p, miR-6812-5p, miR-3197, miR-8059, miR-3185, miR-4706, miR-4497, miR-3131, miR-6806-5p, miR-187-5p, miR-3180-3p, miR-6848-5p, miR-6820-5p, miR-6800-5p, miR-6717-5p, miR-6795-5p, miR-4632-5p, miR-665, miR-6778-5p, miR-3663-3p, miR-4689, miR-211-3p, miR-6511b-5p, miR-4750-5p, miR-6126, miR-614, miR-7110-5p, miR-744-5p, miR-6769a-5p, miR-4792, miR-5787, miR-6798-5p, miR-6781-5p, miR-4419b, miR-4446-3p, miR-4259, miR-5572, miR-6075, miR-296-3p, miR-6891-5p, miR-4745-5p, miR-6775-5p, miR-6870-5p, miR-920, miR-4530, miR-6819-5p, miR-6825-5p, miR-7847-3p, miR-6131, miR-4433-3p, miR-1228-5p, miR-6743-5p, miR-1268a, miR-3917, miR-6786-5p, miR-3154, miR-638, miR-6741-5p, miR-6889-5p, miR-6840-3p, miR-6510-5p, miR-3188, miR-551b-5p, miR-5001-5p, miR-1268b, miR-7107-5p, miR-6824-5p, miR-6732-5p, miR-371a-5p, miR-6794-5p, miR-6779-5p, miR-4271, miR-5195-3p, miR-6762-5p, miR-939-5p, miR-1247-3p, miR-6777-5p, miR-6722-3p, miR-3656, miR-4688, miR-3195, miR-6766-5p, miR-4447, miR-4656, miR-7108-5p, miR-3191-3p, miR-1273g-

thereof comprising 15 or more consecutive nucleotides, (b) a polynucleotide comprising a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235 and 851 to 856, (c) a polynucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235 and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (d) a polynucleotide comprising a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235 and 851 to 856 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, and (e) a polynucleotide hybridizing under stringent conditions to any of the polynucleotides (a) to (d).

(350) The method of the present invention can further employ nucleic acid(s) capable of specifically binding to at least one polynucleotide selected from the group consisting of miR-760, miR-602, miR-423-5p, miR-92a-2-5p, miR-16-5p, miR-451a, miR-135a-3p, miR-486-5p, miR-4257, miR-92b-5p, miR-1915-3p, miR-718, miR-940, miR-296-5p, miR-23b-3p and miR-92a-3p.

(351) As for such a nucleic acid, specifically, miR-760 is hsa-miR-760, miR-602 is hsa-miR-602, miR-423-5p is hsa-miR-423-5p, miR-92a-2-5p is hsa-miR-92a-2-5p, miR-16-5p is hsa-miR-16-5p, miR-451a is hsa-miR-451a, miR-135a-3p is hsa-miR-135a-3p, miR-486-5p is hsa-miR-486-5p, miR-4257 is hsa-miR-4257, miR-92b-5p is hsa-miR-92b-5p, miR-1915-3p is hsa-miR-1915-3p, miR-718 is hsa-miR-718, miR-940 is hsa-miR-940, miR-296-5p is hsa-miR-296-5p, miR-23b-3p is hsa-miR-23b-3p, and miR-92a-3p is hsa-miR-92a-3p.

(352) In a preferred embodiment, such a nucleic acid is specifically selected from the group consisting of the following polynucleotides (f) to (j): (f) a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (g) a polynucleotide comprising a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251, (h) a polynucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (i) a polynucleotide comprising a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, and (j) a polynucleotide hybridizing under stringent conditions to any of the polynucleotides (f) to (i).

(353) The nucleic acid(s) further used in the method of the present invention can comprise nucleic acid(s) capable of specifically binding to at least one polynucleotide selected from the group consisting of miR-658, miR-6842-5p, miR-6124, miR-6765-3p, miR-7106-5p, miR-4534, miR-92b-3p, miR-3135b, miR-4687-3p, miR-762, miR-3619-3p, miR-4467, miR-557, miR-1237-5p, miR-1908-5p, miR-4286, miR-6885-5p and miR-6763-5p.

(354) Specifically, miR-658 is hsa-miR-658, miR-6842-5p is hsa-miR-6842-5p, miR-6124 is hsa-miR-6124, miR-6765-3p is hsa-miR-6765-3p, miR-7106-5p is hsa-miR-7106-5p, miR-4534 is hsa-miR-4534, miR-92b-3p is hsa-miR-92b-3p, miR-3135b is hsa-miR-3135b, miR-4687-3p is hsa-miR-4687-3p, miR-762 is hsa-miR-762, miR-3619-3p is hsa-miR-3619-3p, miR-4467 is hsa-miR-4467, miR-557 is hsa-miR-557, miR-1237-5p is hsa-miR-1237-5p, miR-1908-5p is hsa-miR-1908-5p, miR-4286 is hsa-miR-4286, miR-6885-5p is hsa-miR-6885-5p, and miR-6763-5p is hsa-miR-6763-5p.

(355) Further, in a preferred embodiment, such nucleic acid(s) is/are specifically a polynucleotide selected from the group consisting of the following polynucleotides (k) to (o): (k) a polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (l) a

polynucleotide comprising a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269, (m) a polynucleotide consisting of a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, a variant thereof, a derivative thereof, or a fragment thereof comprising 15 or more consecutive nucleotides, (n) a polynucleotide comprising a nucleotide sequence complementary to a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a nucleotide sequence derived from the nucleotide sequence by the replacement of u with t, and (o) a polynucleotide hybridizing under stringent conditions to any of the polynucleotides (k) to (n).

(356) Examples of the sample used in the method of the present invention can include samples prepared from living tissues (preferably breast tissues) or body fluids such as blood, serum, plasma, and urine from the subjects. Specifically, for example, an RNA-containing sample prepared from the tissue, a polynucleotide-containing sample further prepared therefrom, a body fluid such as blood, serum, plasma, or urine, a portion or the whole of a living tissue collected from the subject by biopsy or the like, or a living tissue excised by surgery can be used, and the sample for measurement can be prepared therefrom.

(357) The subject used herein refers to a mammal, for example, a human, a monkey, a mouse or a rat, and is preferably a human.

(358) The steps of the method of the present invention can be changed according to the type of the sample to be assayed.

(359) In the case of using RNA as an analyte, the detection of breast cancer (cells) can comprise, for example, the following steps (a), (b), and (c): (a) a step of binding RNA prepared from a sample from a subject or complementary polynucleotides (cDNAs) transcribed from the RNA to a polynucleotide(s) in the kit or the device of the present invention; (b) a step of measuring the sample-derived RNA or the cDNAs synthesized from the RNA, which is/are bound to the polynucleotide(s) by hybridization using the polynucleotide(s) as nucleic acid probe(s) or by quantitative RT-PCR using the polynucleotide(s) as primer(s); and (c) a step of evaluating the presence or absence of breast cancer (or breast cancer-derived gene expression) on the basis of the measurement results of the step (b).

(360) For example, various hybridization methods can be used for detecting, examining, evaluating, or diagnosing breast cancer (or breast cancer-derived gene expression) in vitro according to the present invention. For example, Northern blot, Southern blot, RT-PCR, DNA chip analysis, in situ hybridization, Northern hybridization, or Southern hybridization can be used as such a hybridization method.

(361) In the case of using the Northern blot, the presence or absence of expression of each gene or the expression level thereof in the RNA can be detected or measured by use of the nucleic acid probe(s) that can be used in the present invention. Specific examples thereof can include a method which comprises labeling the nucleic acid probe (or a complementary strand) with a radioisotope (.sup.32P, .sup.33P, .sup.35S, etc.), a fluorescent material, or the like, that hybridizes the labeled product with the tissue-derived RNA from a subject, which is transferred to a nylon membrane or the like according to a routine method, and then detecting and measuring a signal derived from the label (radioisotope or fluorescent material) on the formed DNA/RNA duplex using a radiation detector (examples thereof can include BAS-1800 II (Fujifilm Corp.)) or a fluorescence detector (examples thereof can include STORM 865 (GE Healthcare Japan Corp.)).

(362) In the case of using the quantitative RT-PCR, the presence or absence of expression of each gene or the expression level thereof in the RNA can be detected or measured by use of the primer that can be used in the present invention. Specific examples thereof can include a method which comprises; preparing cDNAs from the tissue-derived RNA of a subject according to a routine procedure; hybridizing a pair of primers (consisting of a plus strand and a reverse strand that bind to the cDNA) of the present invention with the cDNA such that the region of each target gene can

be amplified with the cDNA as a template; and performing PCR according to a routine method to detect the obtained double-stranded DNA. The method for detecting the double-stranded DNA can include; a method of performing the PCR using the primers labeled in advance with a radioisotope or a fluorescent material; a method of electrophoresing the PCR product on an agarose gel and staining the double-stranded DNA with ethidium bromide or the like for detection; and a method of transferring the produced double-stranded DNA to a nylon membrane or the like according to a routine method and hybridizing the double-stranded DNA to a labeled nucleic acid probe for detection.

(363) In the case of using the nucleic acid array analysis, an RNA chip or a DNA chip in which the nucleic acid probes (single-stranded or double-stranded) are attached to a substrate (solid phase) is used. Regions having the attached nucleic acid probes are referred to as probe spots, and regions having no attached nucleic acid probe are referred to as blank spots. A gene group immobilized on a solid-phase substrate is generally called a nucleic acid chip, a nucleic acid array, a microarray, or the like. The DNA or RNA array includes a DNA or RNA macroarray and a DNA or RNA microarray. The term “chip” used herein includes all of these arrays. 3D-Gene™ Human miRNA Oligo chip (Toray Industries, Inc.) can be used as the DNA chip, though the DNA chip is not limited thereto.

(364) Examples of the measurement using the DNA chip can include, but are not limited to, a method of detecting and measuring a signal derived from the label on the nucleic acid probe using an image detector (examples thereof can include Typhoon 9410 (GE Healthcare) and 3D-Gene™ scanner (Toray Industries, Inc.)).

(365) The “stringent conditions” used herein are, as mentioned above, conditions under which a nucleic acid probe hybridizes to its target sequence to a larger extent (e.g., a measurement value equal to or larger than “(a mean of background measurement values)+(a standard deviation of the background measurement values) $\times 2$ ”) than that for other sequences.

(366) The stringent conditions are defined by hybridization and subsequent washing conditions. Examples of the hybridization conditions include, but not limited to 30° C. to 60° C. for 1 to 24 hours in a solution containing SSC, a surfactant, formamide, dextran sulfate, a blocking agent(s), etc. In this context, 1 $\times$ SSC is an aqueous solution (pH 7.0) containing 150 mM sodium chloride and 15 mM sodium citrate. The surfactant includes, for example, SDS (sodium dodecyl sulfate), Triton, or Tween. The hybridization conditions more preferably comprise 3-10 $\times$ SSC and 0.1-1% SDS. Examples of the conditions for the washing, following the hybridization, which is another condition to define the stringent conditions, can include conditions comprising continuous washing at 30° C. in a solution containing 0.5 $\times$ SSC and 0.1% SDS, at 30° C. in a solution containing 0.2 $\times$ SSC and 0.1% SDS, and at 30° C. in a 0.05 $\times$ SSC solution. It is desirable that the complementary strand should maintain its hybridized state with a target plus strand even by the washing under such conditions. Specifically, examples of such a complementary strand can include a strand consisting of a nucleotide sequence in a completely complementary relationship with the nucleotide sequence of the target plus (+) strand, and a strand consisting of a nucleotide sequence having at least 80%, preferably at least 85%, more preferably at least 90% or at least 95%, for example, at least 98% or at least 99% identity to the strand.

(367) Other examples of the “stringent conditions” for the hybridization are described in, for example, Sambrook, J. & Russel, D., Molecular Cloning, A LABORATORY MANUAL, Cold Spring Harbor Laboratory Press, published on Jan. 15, 2001, Vol. 1, 7.42 to 7.45 and Vol. 2, 8.9 to 8.17, and can be used in the present invention.

(368) Examples of the conditions for carrying out PCR using polynucleotide fragments contained in the kit of the present invention as primers include treatment for approximately 15 seconds to 1 minute at 5 to 10° C. plus a T_m value calculated from the sequences of the primers, using a PCR buffer having composition such as 10 mM Tris-HCL (pH 8.3), 50 mM KCL, and 1 to 2 mM MgCl₂. Examples of the method for calculating such a T_m value include T_m value=2 $\times$ (the

number of adenine residues+the number of thymine residues)+4×(the number of guanine residues+the number of cytosine residues).

(369) In the case of using the quantitative RT-PCR, a commercially available kit for measurement specially designed for quantitatively measuring miRNA, such as TaqMan™ MicroRNA Assays (Life Technologies Corp.); LNA™-based MicroRNA PCR (Exiqon); or Ncode™ miRNA qRT-PCT kit (Invitrogen Corp.) may be used.

(370) For the calculation of gene expression levels, statistical analysis described in, for example, Statistical analysis of gene expression microarray data (Speed T., Chapman and Hall/CRC), and A beginner's guide Microarray gene expression data analysis (Causton H. C. et al., Blackwell publishing) can be used in the present invention, though the calculation method is not limited thereto. For example, twice, preferably 3 times, more preferably 6 times the standard deviation of the measurement values of the blank spots are added to the average measurement value of the blank spots on the DNA chip, and probe spots having a signal value equal to or larger than the resulting value can be regarded as detection spots. Alternatively, the average measurement value of the blank spots is regarded as a background and can be subtracted from the measurement values of the probe spots to determine gene expression levels. A missing value for a gene expression level can be excluded from the analyte, preferably replaced with the smallest value of the gene expression level in each DNA chip, or more preferably replaced with a value obtained by subtracting 0.1 from a logarithmic value of the smallest value of the gene expression level. In order to eliminate low-signal genes, only a gene that shows a gene expression level of 2.sup.6, preferably 2.sup.8, more preferably 2.sup.10, or larger, in 20% or more, preferably 50% or more, more preferably 80% or more of the number of measurement samples can be selected as the analyte. Examples of the normalization of the gene expression level include, but are not limited to, global normalization and quantile normalization (Bolstad, B. M. et al., 2003, Bioinformatics, Vol. 19, p. 185-193).

(371) The present invention also provides a method comprising measuring target genes or gene expression levels in a sample from a subject using the polynucleotide, the kit, or the device (e.g., chip) for detection of the present invention, or a combination thereof, preparing a discriminant (discriminant function) with gene expression levels in a sample from a breast cancer patient and a sample from a healthy subject as supervising samples, and determining or evaluating the presence and/or absence of the breast cancer-derived genes in the sample.

(372) Specifically, the present invention further provides the method comprising: a first step of measuring in vitro expression levels of target genes in multiple samples known to determine or evaluate the presence and/or absence of the breast cancer-derived genes in the samples, using the polynucleotides, the kit, or the device (e.g., chip) for detection of the present invention, or a combination thereof; a second step of preparing a discriminant with the measurement values of the expression levels of the target genes (target nucleic acid) obtained in the first step as supervising samples; a third step of measuring in vitro expression levels of the target genes in a sample derived from a subject in the same way as in the first step; and a fourth step of assigning the measurement values of the expression levels of the target genes obtained in the third step to the discriminant obtained in the second step, and determining or evaluating the presence and/or absence of the breast cancer-derived gene in the sample on the basis of the results obtained from the discriminant, wherein the target genes can be detected using the polynucleotides or using polynucleotides for detection contained in the kit or the device (e.g., chip). In this context, the discriminant can be prepared by use of Fisher's discriminant analysis, nonlinear discriminant analysis based on Mahalanobis' distance, neural network, Support Vector Machine (SVM), or the like, though the method is not limited thereto.

(373) When a clustering boundary is a straight line or a hyperplane, the linear discriminant analysis is a method for determining the association of a cluster using Formula 1 as a discriminant. In this formula, x represents an explanatory variable, w represents a coefficient of the explanatory variable, and wo represents a constant term.

$$(374) f(x) = w_0 + \sum_{i=1}^n w_i x_i \quad \text{Formula1}$$

(375) Values obtained from the discriminant are referred to as discriminant scores. The measurement values of a newly offered data set can be assigned as explanatory variables to the discriminant to determine clusters by the signs of the discriminant scores.

(376) The Fisher's discriminant analysis, one type of linear discriminant analysis, is a dimension reduction method for selecting a dimension suitable for classification, and constructs a highly discriminating synthetic variable by focusing on the variance of synthetic variables and minimizing the variance of data having the same label (Venables, W. N. et al., Modern Applied Statistics with S. Fourth edition. Springer., 2002). In the Fisher's discriminant analysis, direction w of projection is determined so as to maximize Formula 2. In this formula, μ represents an average input, n_g represents the number of data associated to class g , and μ_g represents an average input of the data associated to class g . The numerator and the denominator are the interclass variance and the intraclass variance, respectively, when each data is projected in the direction of the vector w . Discriminant coefficient w_i is determined by maximizing this ratio (Takafumi Kanamori et al., "Pattern Recognition", Kyoritsu Shuppan Co., Ltd. (2009); and Richard O. et al., Pattern Classification Second Edition., Wiley-Interscience, 2000).

$$(377) J(w) = \frac{\sum_{g=1}^G n_g (w^T \mu_g - w^T \mu)^T (w^T \mu_g - w^T \mu)}{\sum_{g=1}^G \sum_{i: y_i = g} (w^T x_i - w^T \mu_g)^T (w^T x_i - w^T \mu_g)} \quad \text{Formula2}$$

$$\text{subject to} \quad \sum_{i=1}^n \frac{x_i}{n}, \quad \mu_g = \sum_{i: u_i = g}^n \frac{x_i}{n_g}$$

(378) The Mahalanobis' distance is calculated according to Formula 3 in consideration of data correlation and can be used as nonlinear discriminant analysis for determining a cluster to which a data point is associated, based on a short Mahalanobis' distance from the data point to that cluster. In this formula, μ represents a central vector of each cluster, and S^{-1} represents an inverse matrix of the variance-covariance matrix of the cluster. The central vector is calculated from explanatory variable x , and an average vector, a median value vector, or the like can be used.

$$D(x, \mu) = \{(x - \mu).sup.t S.sup.-1(x - \mu)\}.sup.1/2 \quad \text{Formula 3}$$

(379) SVM is a discriminant analysis method devised by V. Vapnik (The Nature of Statistical Learning Theory, Springer, 1995). Particular data points of a data set having known classes are defined as explanatory variables, and classes are defined as objective variables. A boundary plane called hyperplane for correctly classifying the data set into the known classes is determined, and a discriminant for data classification is determined using the boundary plane. Then, the measurement values of a newly offered data set can be assigned as explanatory variables to the discriminant to determine classes. In this respect, the result of the discriminant analysis may be the associated classe, may be a probability of being classified into correct classes, or may be the distance from the hyperplane. In SVM, a method of nonlinearly converting a feature vector to a high dimension and performing linear discriminant analysis in the space is known as a method for tackling nonlinear problems. An expression in which an inner product of two factors in a nonlinearly mapped space is expressed only by inputs in their original spaces is called kernel. Examples of the kernel can include a linear kernel, a RBF (Radial Basis Function) kernel, and a Gaussian kernel. While highly dimensional mapping is performed according to the kernel, the optimum discriminant, i.e., a discriminant, can be actually constructed by mere calculation according to the kernel, which avoids calculating features in the mapped space (e.g., Hideki Aso et al., Frontier of Statistical Science 6 "Statistics of pattern recognition and learning—New concepts and approaches", Iwanami Shoten, Publishers (2004); Nello Cristianini et al., Introduction to SVM, Kyoritsu Shuppan Co., Ltd. (2008)).

(380) C-support vector classification (C-SVC), one type of SVM, comprises preparing a hyperplane by supervising a data set with the explanatory variables of two groups and classifying

an unknown data set into either of the groups (C. Cortes et al., 1995, Machine Learning, Vol. 20, p. 273-297).

(381) Exemplary calculation of the C-SVC discriminant that can be used in the method of the present invention will be given below. First, all subjects are divided into two groups, i.e., a breast cancer patient group and a healthy subject group. For example, breast tissue examination can be used for a reference under which each subject is confirmed either as a breast cancer patient or as a healthy subject.

(382) Next, a data set consisting of comprehensive gene expression levels of serum-derived samples of the two divided groups (hereinafter, this data set is referred to as a training cohort) is prepared, and a C-SVC discriminant is determined by explanatory variables that are genes found to differ clearly in their gene expression levels between the two groups, and objective variables (e.g., -1 and +1) that are the grouping. An optimizing objective function is represented by Formula 4 wherein e represents all input vectors, y represents an objective variable, a represents a Lagrange's undetermined multiplier vector, Q represents a positive definite matrix, and C represents a parameter for adjusting constrained conditions.

$$(383) \min_a \frac{1}{2} a^T Q a - e^T a \quad \text{Formula 4} \quad \text{subject to } y^T a = 0, 0 \leq a_i \leq C, i = 1, \dots, l,$$

(384) Formula 5 is a finally obtained discriminant, and a group to which the data point is associated can be determined on the basis of the sign of a value obtained according to the discriminant. In this formula, x represents a support vector, y represents a label indicating the association of a group, a represents the corresponding coefficient, b represents a constant term, and K represents a kernel function.

$$(385) f(x) = \text{sgn} \left(\sum_{i=1}^l y_i a_i K(x_i, x) + b \right) \quad \text{Formula 5}$$

(386) For example, a RBF kernel defined by Formula 6 can be used as the kernel function. In this formula, x represents a support vector, and y represents a kernel parameter for adjusting the complexity of the hyperplane.

$$K(x_{\text{sub}.i}, x_{\text{sub}.j}) = \exp(-r \|x_{\text{sub}.i} - x_{\text{sub}.j}\|_{\text{sup}.2}), r < 0 \quad \text{Formula 6}$$

(387) In addition, an approach such as neural network, k-nearest neighbor algorithms, decision trees, or logistic regression analysis can be selected as a method for determining or evaluating the presence and/or absence of expression of a breast cancer-derived target gene in a sample derived from a subject, or for evaluating the expression level thereof by comparison with a control derived from a healthy subject.

(388) The method of the present invention can comprise, for example, the following steps (a), (b), and (c): (a) a step of measuring expression level(s) of target gene(s) in tissues containing breast cancer-derived genes derived from breast cancer patients and/or samples already known to be tissues containing no breast cancer-derived gene(s) derived from healthy subjects, using the polynucleotide(s), the kit, or the device (e.g., DNA chip) for detection according to the present invention; (b) a step of preparing the discriminants of Formulas 1 to 3, 5, and 6 described above from the measurement values of the expression level measured in the step (a); and (c) a step of measuring expression level(s) of the target gene(s) in a sample derived from a subject using the polynucleotide(s), the kit, or the device (e.g., DNA chip) for diagnosis (detection) according to the present invention, substituting the obtained measurement value c into the discriminants prepared in the step (b), and determining or evaluating the presence and/or absence of the breast cancer-derived target gene in the sample, or evaluating the expression levels thereof by comparison with a healthy subject-derived control, on the basis of the obtained results. In this context, in the discriminants of Formulas 1 to 3, 5, and 6, x represents an explanatory variable and includes a value obtained by measuring polynucleotide(s) selected from the polynucleotides described in the Section 2 above, or any fragment thereof. Specifically, the explanatory variable for discriminating a breast cancer patient from a healthy subject according to the present invention is gene expression level(s)

selected from, for example, the following expression levels (1) to (3): (1) gene expression level(s) in the serum of a breast cancer patient or a healthy subject measured by any DNA comprising 15 or more consecutive nucleotides in a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235 and 851 to 856 or a complementary sequence thereof, (2) gene expression level(s) in the serum of a breast cancer patient or a healthy subject measured by any DNA comprising 15 or more consecutive nucleotides in a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251 or a complementary sequence thereof, (3) gene expression level(s) in the serum of a breast cancer patient or a healthy subject measured by any DNA comprising 15 or more consecutive nucleotides in a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269 or a complementary sequence thereof.

(389) As described above, for the method for determining or evaluating the presence and/or absence of breast cancer-derived gene(s) in a sample derived from a subject, the preparation of a discriminant requires a discriminant prepared from a training cohort. For enhancing the accuracy of the discriminant, it is necessary for the discriminant to use genes that show clear difference between two groups in the training cohort.

(390) Each gene that is used for an explanatory variable in a discriminant is preferably determined as follows. First, comprehensive gene expression levels of a breast cancer patient group and comprehensive gene expression levels of a healthy subject group, both of which are in a training cohort, are used as a data set, the degree of difference in the expression level of each gene between the two groups is determined through the use of, for example, the P value of t test, which is parametric analysis, or the P value of Mann-Whitney's U test or Wilcoxon test, which is nonparametric analysis.

(391) The gene can be regarded as being statistically significant when the critical rate (significance level) of the P value obtained by the test is smaller than, for example, 5%, 1%, or 0.01%.

(392) In order to correct an increased probability of type I error attributed to the repetition of a test, a method known in the art, for example, Bonferroni or Holm method, can be used for the correction (e.g., Yasushi Nagata et al., "Basics of statistical multiple comparison methods", Scientist Press Co., Ltd. (2007)). As an example of the Bonferroni correction, for example, the P value obtained by a test is multiplied by the number of repetitions of the test, i.e., the number of genes used in the analysis, and the obtained value can be compared with a desired significance level to suppress a probability of causing type I error in the whole test.

(393) Instead of the statistical test, the absolute value (fold change) of an expression ratio of a median value of each gene expression level between gene expression levels of a breast cancer patient group and gene expression levels of a healthy subject group may be calculated to select a gene that is used for an explanatory variable in a discriminant. Alternatively, ROC curves may be prepared using gene expression levels of a breast cancer patient group and a healthy subject group, and a gene that is used for an explanatory variable in a discriminant can be selected on the basis of an AUROC value.

(394) Next, a discriminant that can be calculated by various methods described above is prepared using any number of genes that show large difference in their gene expression levels determined here. Examples of the method for constructing a discriminant that produces the largest discriminant accuracy include a method of constructing a discriminant in every combination of genes that satisfy the significance level of P value, and a method of repetitively evaluating the genes for use in the construction of a discriminant while increasing the number of genes one by one in a descending order of the difference in gene expression level (Furey T S. et al., 2000, Bioinformatics., Vol. 16, p. 906-14). A gene expression level of another independent breast cancer patient or healthy subject is assigned as an explanatory variable to this discriminant to calculate discriminant results of the group to which this independent breast cancer patient or healthy subject is associated. Specifically, the found gene set for diagnosis and the discriminant constructed using the gene set for diagnosis can be evaluated in an independent sample cohort to find a more universal gene set for diagnosis

capable of detecting breast cancer and a more universal method for discriminating breast cancer. (395) Split-sample method is preferably used for evaluating the performance (generality) of the discriminant. Specifically, a data set is divided into a training cohort and a validation cohort, and gene selection by a statistical test and discriminant construction are performed using the training cohort. Accuracy, sensitivity, and specificity are calculated using results of discriminant analysis in the validation cohort according to the discriminant constructed and a true group to which the validation cohort is associated, to evaluate the discriminant performance of the discriminant. On the other hand, instead of dividing a data set, gene selection by a statistical test and discriminant construction may be performed using all of samples, and accuracy, sensitivity, and specificity can be calculated by the discriminant analysis using a newly prepared cohorts for evaluation of the performance of the discriminant.

(396) The present invention provides polynucleotides for detection and disease diagnosis useful in the diagnosis and treatment of breast cancer, a method for detecting breast cancer using the polynucleotide(s), and a kit and a device for the detection of breast cancer, comprising the polynucleotide(s). Particularly, in order to select gene(s) for diagnosis and prepare a discriminant so as to exhibit accuracy beyond the breast cancer diagnosis method using an existing tumor marker CEA, a gene set for diagnosis and a discriminant for the method of the present invention can be constructed, which exhibit accuracy beyond CEA, for example, by comparing expressed genes in serum from a patient confirmed to be negative using CEA but finally found to have breast cancer by detailed examination such as computed tomography using a contrast medium, with genes expressed in serum from a patient having no breast cancer.

(397) For example, the gene set for diagnosis is set to any combination selected from one or two or more of the polynucleotides based on a nucleotide sequence represented by any of SEQ ID NOs: 1 to 235, and 851 to 856, or a complementary sequence thereof as described above; and optionally one or two or more of the polynucleotides based on a nucleotide sequence represented by any of SEQ ID NOs: 236 to 251, or a complementary sequence thereof; and optionally one or two or more of the polynucleotides based on a nucleotide sequence represented by any of SEQ ID NOs: 252 to 269, or a complementary sequence thereof. Further, a discriminant is constructed using expression levels of the gene set for diagnosis in samples from class I breast cancer patients as a result of tissue diagnosis and samples from class II healthy subjects as a result of tissue diagnosis. As a result, the presence or absence of breast cancer-derived genes in an unknown sample can be determined with 100% accuracy at the maximum by measuring expression levels of the gene set for diagnosis in an unknown sample.

EXAMPLES

(398) Hereinafter, the present invention will be described further specifically with reference to Examples below. However, the scope of the present invention is not intended to be limited by these Examples.

Reference Example 1

(399) <Collection of Samples from Breast Cancer Patients and Healthy Subjects>

(400) Sera were collected using VENOJECT II vacuum blood collecting tube VP-AS109K60 (Terumo Corp.) from 100 healthy subjects (92 males and 8 females) and 62 breast cancer patients (20 cases with stage I, 24 cases with stage IIA, 7 cases with stage IIB, 2 cases with stage IIIA, 3 cases with stage IIIB, 1 case with stage IIIC, and 5 cases with stage IV) who were confirmed to have no primary cancer other than breast cancer after acquisition of informed consent, and used as a training cohort. Likewise, Sera were collected using VENOJECT II vacuum blood collecting tube VP-AS109K60 (Terumo Corp.) from 50 healthy subjects (44 males and 6 females) and 31 breast cancer patients (9 cases with stage I, 13 cases with stage IIA, 5 cases with stage IIB, 1 case with stage IIIA, 1 case with stage IIIB, 1 case with stage IIIC, and 1 case with stage IV) who were confirmed to have no primary cancer other than breast cancer after acquisition of informed consent, and used as a validation cohort.

(401) <Extraction of Total RNA>

(402) Total RNA was obtained from 300 μ L of the serum sample obtained from each of 243 persons in total of 150 healthy subjects and 93 breast cancer patients included in the training cohort and the validation cohort, using a reagent for RNA extraction in 3D-Gene™ RNA extraction reagent from liquid sample kit (Toray Industries, Inc.) according to the protocol provided by the manufacturer.

(403) <Measurement of Gene Expression Level>

(404) miRNAs in the total RNA obtained from the serum samples of each of 243 persons in total of 150 healthy subjects and 93 breast cancer patients included in the training cohort and the validation cohort were fluorescently labeled using 3D-Gene™ miRNA Labeling kit (Toray Industries, Inc.) according to the protocol (ver 2.20) provided by the manufacturer. The oligo DNA chip used was 3D-Gene™ Human miRNA Oligo chip (Toray Industries, Inc.) with attached probes having sequences complementary to 2,555 miRNAs among the miRNAs registered in miRBase Release 20. Hybridization between the miRNAs in the total RNA and the probes on the DNA chip under stringent conditions and washing following the hybridization were performed according to the protocol provided by the manufacturer. The DNA chip was scanned using 3D-Gene™ scanner (Toray Industries, Inc.) to obtain images. Fluorescence intensity was digitized using 3D-Gene™ Extraction (Toray Industries, Inc.). The digitized fluorescence intensity was converted to a logarithmic value with a base of 2 and used as a gene expression level, from which a blank value was subtracted. A missing value was replaced with a value obtained by subtracting 0.1 from a logarithmic value of the smallest value of the gene expression level in each DNA chip. As a result, the comprehensive gene expression levels of the miRNAs in the sera were obtained for the 93 breast cancer patients and the 150 healthy subjects. Calculation and statistical analysis using the digitized gene expression levels of the miRNAs were carried out using R language 3.0.2 (R Development Core Team (2013). R: A language and environment for statistical computing. R Foundation for Statistical Computing, URL <http://www.R-project.org/>) and MASS package 7.3-30 (Venables, W. N. & Ripley, B. D. (2002) Modern Applied Statistics with S. Fourth Edition. Springer, New York. ISBN 0-387-95457-0).

Reference Example 2

(405) <Collection of Samples from Patients with Cancer Other than Breast Cancer>

(406) Sera were collected using VENOJECT II vacuum blood collecting tube VP-AS109K60 (Terumo Corp.) from each of 33 prostate cancer patients confirmed to have no cancer in other organs after acquisition of informed consent, and used as a training cohort together with the samples of 62 breast cancer patients and 102 healthy subjects of Reference Example 1. Likewise, sera were collected using VENOJECT 11 vacuum blood collecting tube VP-AS109K60 (Terumo Corp.) from each of 19 prostate cancer patients confirmed to have no cancer in other organs after acquisition of informed consent, and used as a validation cohort together with the samples of 31 breast cancer patients confirmed to have no cancer in organs other than the breast and 48 healthy subjects of Reference Example 1. Subsequent operations were conducted in the same way as in Reference Example 1.

Example 1

(407) <Selection of Gene Marker Using the Training Cohort, and Method for Evaluating Breast Cancer Discriminant Performance of the Single Gene Marker Using the Validation Cohort>

(408) In this Example, a gene marker for discriminating a breast cancer patient from a healthy subject was selected from the training cohort and studied in the validation cohort independent of the training cohort, for a method for evaluating the breast cancer discriminant performance of each selected gene marker alone.

(409) Specifically, first, the miRNA expression levels of the training cohort and the validation cohort obtained in the preceding Reference Examples were combined and normalized by quantile normalization.

(410) Next, genes for diagnosis were selected using the training cohort. Here, in order to acquire diagnostic markers with higher reliability, only genes that show the gene expression level of 2.sup.6 or higher in 50% or more of the samples in either of the breast cancer patient group in the training cohort or the healthy subject group in the training cohort were selected. In order to further acquire statistically significant genes for discriminating a breast cancer patient group from a healthy subject group, the P value obtained by two-tailed t-test assuming equal variance as to each gene expression level was corrected by the Bonferroni method, and genes that satisfied $p < 0.01$ were acquired as gene markers for use in explanatory variables of a discriminant. The obtained genes are described in Table 2.

(411) In this way, hsa-miR-4783-3p, hsa-miR-4730, hsa-miR-1307-3p, hsa-miR-4634, hsa-miR-663a, hsa-miR-4532, hsa-miR-7704, hsa-miR-3178, hsa-miR-6729-5p, hsa-miR-6090, hsa-miR-4732-5p, hsa-miR-3184-5p, hsa-miR-6727-5p, hsa-miR-6088, hsa-miR-4674, hsa-miR-8073, hsa-miR-4787-5p, hsa-miR-1469, hsa-miR-125a-3p, hsa-miR-1233-5p, hsa-miR-885-3p, hsa-miR-6802-5p, hsa-miR-328-5p, hsa-miR-6787-5p, hsa-miR-8069, hsa-miR-6875-5p, hsa-miR-1246, hsa-miR-4734, hsa-miR-6757-5p, hsa-miR-6756-5p, hsa-miR-3665, hsa-miR-6836-3p, hsa-miR-6821-5p, hsa-miR-6805-5p, hsa-miR-4728-5p, hsa-miR-6726-5p, hsa-miR-197-5p, hsa-miR-149-3p, hsa-miR-6850-5p, hsa-miR-4476, hsa-miR-6858-5p, hsa-miR-564, hsa-miR-4763-3p, hsa-miR-575, hsa-miR-6771-5p, hsa-miR-1231, hsa-miR-1908-3p, hsa-miR-150-3p, hsa-miR-3937, hsa-miR-887-3p, hsa-miR-3940-5p, hsa-miR-4741, hsa-miR-6808-5p, hsa-miR-6869-5p, hsa-miR-5090, hsa-miR-615-5p, hsa-miR-8072, hsa-miR-128-1-5p, hsa-miR-1238-5p, hsa-miR-365a-5p, hsa-miR-204-3p, hsa-miR-4492, hsa-miR-6785-5p, hsa-miR-6511a-5p, hsa-miR-4525, hsa-miR-1915-5p, hsa-miR-3180, hsa-miR-6879-5p, hsa-miR-1199-5p, hsa-miR-6746-5p, hsa-miR-711, hsa-miR-663b, hsa-miR-4707-3p, hsa-miR-6893-5p, hsa-miR-4675, hsa-miR-4638-5p, hsa-miR-4651, hsa-miR-6087, hsa-miR-4665-5p, hsa-miR-4758-5p, hsa-miR-6887-5p, hsa-miR-3620-5p, hsa-miR-1909-3p, hsa-miR-7641, hsa-miR-6724-5p, hsa-miR-1343-3p, hsa-miR-6780b-5p, hsa-miR-4484, hsa-miR-4690-5p, hsa-miR-4429, hsa-miR-1227-5p, hsa-miR-4725-3p, hsa-miR-6861-5p, hsa-miR-6812-5p, hsa-miR-3197, hsa-miR-8059, hsa-miR-3185, hsa-miR-4706, hsa-miR-4497, hsa-miR-3131, hsa-miR-6806-5p, hsa-miR-187-5p, hsa-miR-3180-3p, hsa-miR-6848-5p, hsa-miR-6820-5p, hsa-miR-6800-5p, hsa-miR-6717-5p, hsa-miR-6795-5p, hsa-miR-4632-5p, hsa-miR-665, hsa-miR-6778-5p, hsa-miR-3663-3p, hsa-miR-4689, hsa-miR-211-3p, hsa-miR-6511b-5p, hsa-miR-4750-5p, hsa-miR-6126, hsa-miR-614, hsa-miR-7110-5p, hsa-miR-744-5p, hsa-miR-6769a-5p, hsa-miR-4792, hsa-miR-5787, hsa-miR-6798-5p, hsa-miR-6781-5p, hsa-miR-4419b, hsa-miR-4446-3p, hsa-miR-4259, hsa-miR-5572, hsa-miR-6075, hsa-miR-296-3p, hsa-miR-6891-5p, hsa-miR-4745-5p, hsa-miR-6775-5p, hsa-miR-6870-5p, hsa-miR-920, hsa-miR-4530, hsa-miR-6819-5p, hsa-miR-6825-5p, hsa-miR-7847-3p, hsa-miR-6131, hsa-miR-4433-3p, hsa-miR-1228-5p, hsa-miR-6743-5p, hsa-miR-1268a, hsa-miR-3917, hsa-miR-6786-5p, hsa-miR-3154, hsa-miR-638, hsa-miR-6741-5p, hsa-miR-6889-5p, hsa-miR-6840-3p, hsa-miR-6510-5p, hsa-miR-3188, hsa-miR-551b-5p, hsa-miR-5001-5p, hsa-miR-1268b, hsa-miR-7107-5p, hsa-miR-6824-5p, hsa-miR-6732-5p, hsa-miR-371a-5p, hsa-miR-6794-5p, hsa-miR-6779-5p, hsa-miR-4271, hsa-miR-5195-3p, hsa-miR-6762-5p, hsa-miR-939-5p, hsa-miR-1247-3p, hsa-miR-6777-5p, hsa-miR-6722-3p, hsa-miR-3656, hsa-miR-4688, hsa-miR-3195, hsa-miR-6766-5p, hsa-miR-4447, hsa-miR-4656, hsa-miR-7108-5p, hsa-miR-3191-3p, hsa-miR-1273g-3p, hsa-miR-4463, hsa-miR-2861, hsa-miR-3196, hsa-miR-6877-5p, hsa-miR-3679-5p, hsa-miR-4442, hsa-miR-6789-5p, hsa-miR-6782-5p, hsa-miR-486-3p, hsa-miR-6085, hsa-miR-4746-3p, hsa-miR-619-5p, hsa-miR-937-5p, hsa-miR-6803-5p, hsa-miR-4298, hsa-miR-4454, hsa-miR-4459, hsa-miR-7150, hsa-miR-6880-5p, hsa-miR-4449, hsa-miR-8063, hsa-miR-4695-5p, hsa-miR-6132, hsa-miR-6829-5p, hsa-miR-4486, hsa-miR-6805-3p, hsa-miR-6826-5p, hsa-miR-4508, hsa-miR-1343-5p, hsa-miR-7114-5p, hsa-miR-3622a-5p, hsa-miR-6765-5p, hsa-miR-7845-5p, hsa-miR-3960, hsa-miR-6749-5p, hsa-miR-1260b, hsa-miR-6799-5p, hsa-miR-4723-5p, hsa-miR-6784-5p, hsa-miR-5100, hsa-miR-6769b-5p, hsa-miR-1207-5p, hsa-miR-642a-3p, hsa-miR-4505, hsa-miR-4270, hsa-miR-6721-5p, hsa-miR-

7111-5p, hsa-miR-6791-5p, hsa-miR-7109-5p, hsa-miR-4258, hsa-miR-6515-3p, hsa-miR-6851-5p, hsa-miR-6125, hsa-miR-4749-5p, hsa-miR-4726-5p, hsa-miR-4513, hsa-miR-760, hsa-miR-602, hsa-miR-423-5p, hsa-miR-92a-2-5p, hsa-miR-16-5p, hsa-miR-451a, hsa-miR-135a-3p, hsa-miR-486-5p, hsa-miR-4257, hsa-miR-92b-5p, hsa-miR-1915-3p, hsa-miR-718, hsa-miR-940, hsa-miR-296-5p, hsa-miR-23b-3p and hsa-miR-92a-3p genes, and polynucleotides consisting of the nucleotide sequences of SEQ ID NOs: 1 to 251 related thereto were found.

(412) Among them, genes newly found as markers for examining the presence or absence of breast cancer are polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 235.

(413) A discriminant for determining the presence or absence of breast cancer was further prepared by Fisher's discriminant analysis with the expression levels of these genes as indicators.

Specifically, any newly found polynucleotide consisting of a nucleotide sequence represented by any of SEQ ID NOs: 1 to 251 in the training cohort was applied for Formula 2 above to prepare a discriminant. Calculated accuracy, sensitivity, and specificity are shown in Table 3. In this respect, a discriminant coefficient and a constant term are shown in Table 4. Here, all of the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 251 were selected as markers capable of determining not only invasive ductal breast cancer (56 cases), which is a main type of breast cancer, but also invasive lobular cancer (3 cases) and unusual metastatic carcinoma with poor prognosis (1 case).

(414) Accuracy, sensitivity, and specificity in the validation cohort were calculated using the discriminant thus prepared, and the discriminant performance of the selected polynucleotides was validated using the independent samples (Table 3). For example, the expression level measurement value of the gene that consists of the nucleotide sequence represented by SEQ ID NO: 1 was compared between the healthy subjects (100 persons) and the breast cancer patients (62 persons) in the training cohort. As a result, the expression level measurement values were found to be significantly lower in the breast cancer patient group than in the healthy subject group (see the left diagram of FIG. 2). These results were also reproducible for the healthy subjects (50 persons) and the breast cancer patients (31 persons) in the validation cohort (see the right diagram of FIG. 2).

Likewise, the results obtained about the other polynucleotides shown in SEQ ID NOs: 2 to 251 showed that the expression level measurement values were significantly lower (−) or higher (+) in the breast cancer patient group than in the healthy subject group (Table 2). These results were able to be validated in the validation cohort. For example, as for this nucleotide sequence represented by SEQ ID NO: 1, the number of correctly or incorrectly identified samples in the detection of breast cancer was calculated using the threshold (6.63) that was set in the training cohort and discriminated between the two groups. As a result, 31 true positives, 50 true negatives, 0 false positives, and 0 false negatives were obtained. From these values, 100% accuracy, 100% sensitivity, and 100% specificity were obtained as the detection performance. In this way, the detection performance was calculated as to all of the polynucleotides shown in SEQ ID NOs: 1 to 251, and described in Table 3. Likewise, the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 2 to 251 shown in Table 2 exhibited sensitivity of 100%, 100%, 90.3%, 96.8%, 100%, 100%, 96.8%, 96.8%, 100%, 100%, 100%, 100%, 96.8%, 96.8%, 100%, 100%, 96.8%, 100%, 93.5%, 96.8%, 96.8%, 100%, 96.8%, 96.8%, 93.5%, 96.8%, 93.5%, 87.1%, 83.9%, 96.8%, 96.8%, 96.8%, 100%, 80.6%, 87.1%, 87.1%, 100%, 90.3%, 90.3%, 90.3%, 100%, 96.8%, 100%, 87.1%, 87.1%, 96.8%, 96.8%, 90.3%, 96.8%, 83.9%, 77.4%, 90%, 83.9%, 96.8%, 93.5%, 80.6%, 96.8%, 90.3%, 93.5%, 90.3%, 87.1%, 87.1%, 96.8%, 83.9%, 87.1%, 77.4%, 90.3%, 77.4%, 90.3%, 83.9%, 74.2%, 93.5%, 87.1%, 93.5%, 93.5%, 77.4%, 90.3%, 87.1%, 87.1%, 83.9%, 87.1%, 93.5%, 77.4%, 93.5%, 74.2%, 83.9%, 100%, 90.3%, 74.2%, 83.9%, 80.6%, 87.1%, 77.4%, 83.9%, 71%, 96.8%, 77.4%, 87.1%, 77.4%, 71%, 90.3%, 80.6%, 67.7%, 77.4%, 87.1%, 74.2%, 83.9%, 77.4%, 71%, 87.1%, 74.2%, 90.3%, 80.6%, 74.2%, 83.9%, 83.9%, 71%, 87.1%, 61.3%, 61.3%, 83.9%, 61.3%, 90.3%, 80.6%, 61.3%, 64.5%, 80.6%, 74.2%, 80.6%, 71%, 71%, 77.4%,

64.5%, 71%, 71%, 83.9%, 74.2%, 83.9%, 63.3%, 64.5%, 71%, 67.7%, 71%, 71%, 74.2%, 71%, 64.5%, 83.9%, 71%, 83.9%, 61.3%, 61.3%, 67.7%, 64.5%, 64.5%, 54.8%, 64.5%, 74.2%, 58.1%, 58.1%, 58.1%, 61.3%, 67.7%, 61.3%, 67.7%, 58.1%, 58.1%, 54.8%, 67.7%, 58.1%, 64.5%, 61.3%, 67.7%, 58.1%, 58.1%, 48.4%, 61.3%, 54.8%, 38.7%, 35.5%, 64.5%, 54.8%, 64.5%, 54.8%, 61.3%, 35.5%, 48.4%, 61.3%, 61.3%, 54.8%, 71%, 61.3%, 45.2%, 48.4%, 29%, 54.8%, 41.9%, 67.7%, 29%, 29%, 53.3%, 51.6%, 45.2%, 35.5%, 41.9%, 41.9%, 48.4%, 41.9%, 35.5%, 41.9%, 35.5%, 48.4%, 32.3%, 41.9%, 41.9%, 41.9%, 35.5%, 35.5%, 41.9%, 61.3%, 32.3%, 45.2%, 38.7%, 51.6%, 29%, 35.5%, 38.7%, 54.8%, 58.1%, 51.6%, 29%, 41.9%, 38.7%, 96.8%, 96.8%, 96.8%, 100%, 96.8%, 87.1%, 80.6%, 100%, 87.1%, 93.5%, 67.7%, 67.7%, 61.3%, 67.7%, 38.7% and 54.8%, respectively, in the validation cohort (Table 3). As seen from Comparative Example mentioned later, the existing marker CEA had sensitivity of 19.4% in the validation cohort (Table 5-2), demonstrating that all of the polynucleotides consisting of the nucleotide sequences represented by SEQ ID Nos: 1 to 251 can discriminate, each alone, breast cancer in the validation cohort with sensitivity beyond CEA.

(415) For example, the 83 polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 31, 32, 33, 34, 35, 36, 38, 40, 42, 44, 45, 47, 48, 49, 50, 55, 59, 64, 65, 66, 68, 70, 73, 75, 78, 79, 80, 81, 84, 88, 92, 93, 95, 97, 98, 99, 102, 109, 113, 119, 122, 124, 128, 130, 133, 145, 149, 169, 236, 237, 238, 239, 240, 241, 242, 243, 244 were able to correctly determine breast cancer in the 9 breast cancer samples of stage 1 contained in the validation cohort. Thus, these polynucleotides can detect even early breast cancer and contributes to the early diagnosis of breast cancer.

Example 2

(416) <Method a for Evaluating Breast Cancer Discriminant Performance by Combination of Multiple Gene Markers Using the Samples in the Validation Cohort>

(417) In this Example, a method for evaluating breast cancer discriminant performance by a combination of the gene markers selected in Example 1 was studied. Specifically, Fisher's discriminant analysis was conducted as to 31,255 polynucleotide combinations comprising at least one of the newly found polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 235 among the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 251 selected in Example 1, to construct a discriminant for determining the presence or absence of breast cancer. Next, accuracy, sensitivity, and specificity in the validation cohort were calculated using the discriminant thus prepared, and the discriminant performance of the selected polynucleotides was validated using the independent samples.

(418) For example, the gene expression level measurement values of the nucleotide sequences represented by SEQ ID NO: 1 and SEQ ID NO: 2 were compared between the healthy subjects and the breast cancer patients. As a result, a scatter diagram that significantly separated the expression level measurement values of the breast cancer patient group from those of the healthy subject group was obtained (see the left diagram of FIG. 3). These results were also reproducible for the healthy subjects (50 persons) and the breast cancer patients (31 persons) in the validation cohort (see the right diagram of FIG. 3). Likewise, a scatter diagram that significantly separated the expression level measurement values of the breast cancer patient group from those of the healthy subject group was also obtained as to the other polynucleotide combinations comprising at least one of the newly found polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 235 among the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 251. These results were able to be validated in the validation cohort. For example, as for these nucleotide sequences represented by SEQ ID NO: 1 and SEQ ID NO: 2, the number of samples that were correctly or incorrectly identified breast cancer was calculated using the discriminant function ($0=1.41x+y+0.77$) that was set in the training cohort and discriminated between the two groups. As a result, 31 true positives, 50 true negatives, 0 false positives, and 0

false negatives were obtained. From these values, 100% accuracy, 100% sensitivity, and 100% specificity were obtained as the detection performance. In this way, the detection performance was calculated as to all of the polynucleotide combinations comprising at least one of the newly found polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 235 among the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 251. Among them, 250 combinations comprising the expression level measurement value of the polynucleotide consisting of the nucleotide sequence represented by SEQ ID NO: 1 and the detection performance thereof were described in Table 6 as an example. For example, all of the combinations of the expression level measurement values of the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 and 2, SEQ ID NOs: 1 and 3, SEQ ID NOs: 1 and 4, and SEQ ID NOs: 1 and 5 exhibited sensitivity of 100% in the validation cohort. Likewise, all of the combinations of two polynucleotides consisting of the nucleotide sequence represented by SEQ ID NO: 1 and a nucleotide sequence represented by any of SEQ ID NOs: 6 to 251 also exhibited sensitivity of 100%. In addition, the combinations of two polynucleotides consisting of the nucleotide sequences except for SEQ ID NO: 1 were described in Table 7 as an example. As the specific combinations of two polynucleotides, for example, the combinations represented by SEQ ID NOs: 3 and 20, SEQ ID NOs: 6 and 20, SEQ ID NOs: 7 and 20, SEQ ID NOs: 10 and 20, SEQ ID NOs: 20 and 22, SEQ ID NOs: 20 and 238, SEQ ID NOs: 20 and 239, SEQ ID NOs: 12 and 24, SEQ ID NOs: 20 and 24, SEQ ID NOs: 24 and 27, SEQ ID NOs: 24 and 33, SEQ ID NOs: 24 and 236, SEQ ID NOs: 24 and 240, SEQ ID NOs: 3 and 26, SEQ ID NOs: 12 and 26, SEQ ID NOs: 13 and 26, SEQ ID NOs: 17 and 26, SEQ ID NOs: 19 and 26, SEQ ID NOs: 3 and 27, SEQ ID NOs: 5 and 27, SEQ ID NOs: 13 and 27, SEQ ID NOs: 20 and 27, SEQ ID NOs: 26 and 27, SEQ ID NOs: 27 and 120, SEQ ID NOs: 27 and 206, SEQ ID NOs: 27 and 237, SEQ ID NOs: 3 and 30, SEQ ID NOs: 17 and 30, SEQ ID NOs: 27 and 30, SEQ ID NOs: 27 and 33, SEQ ID NOs: 30 and 39, SEQ ID NOs: 30 and 117, SEQ ID NOs: 3 and 33, SEQ ID NOs: 7 and 33, SEQ ID NOs: 10 and 33, SEQ ID NOs: 11 and 33, SEQ ID NOs: 13 and 33, SEQ ID NOs: 25 and 33, SEQ ID NOs: 33 and 244, SEQ ID NOs: 3 and 182, SEQ ID NOs: 6 and 182, SEQ ID NOs: 7 and 182, SEQ ID NOs: 12 and 182, SEQ ID NOs: 27 and 182, SEQ ID NOs: 182 and 236, SEQ ID NOs: 2 and 194, SEQ ID NOs: 7 and 194, SEQ ID NOs: 27 and 194, SEQ ID NOs: 194 and 236, SEQ ID NOs: 2 and 206, SEQ ID NOs: 7 and 206, SEQ ID NOs: 206 and 236, SEQ ID NOs: 2 and 208, SEQ ID NOs: 7 and 208, SEQ ID NOs: 13 and 208, SEQ ID NOs: 20 and 208, and SEQ ID NOs: 27 and 208 exhibited accuracy of 96% or higher for discriminating the breast cancer patients from the healthy subjects in both of the training cohort and the validation cohort. Thus, the combinations of two of the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 251 also produced excellent detection sensitivity for breast cancer. Markers for the detection of breast cancer with better sensitivity are obtained by further combining 3, 4, 5, 6, 7, 8, 9, 10 or more of the measurement values of the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 251. For example, the newly found polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 235 among the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 251 selected in Example 1, were measured to obtain their expression levels between the healthy subject group and the breast cancer group in the validation cohort. All of the polynucleotides were ranked in the descending order of their P values based on the Student's t-test which indicates statistical significance of a difference between groups (i.e., one having the lowest P value was ranked in the first place), and breast cancer detection sensitivity was evaluated using combinations of one or more polynucleotides to which the polynucleotides were added one by one from the top to bottom according to the rank. In short, the order in which the polynucleotides were combined in this evaluation is in reverse in terms of SEQ ID NOs, such as SEQ ID NO: 235 to SEQ ID NOs: 234, 233, . . . shown in Table 2 in order. As a result, the sensitivity in the validation cohort was 38.7% for 1 polynucleotide (SEQ ID NO: 235), 48.4% for 2 polynucleotides (SEQ ID NOs: 234 and 235), 74.2% for 4 polynucleotides (SEQ ID

NOs: 232 to 235), 87.1% for 6 polynucleotides (SEQ ID NOs: 230 to 235), 90.3% for 10 polynucleotides (SEQ ID NOs: 226 to 235), 93.5% for 13 polynucleotides (SEQ ID NOs: 223 to 235), 96.8% for 16 polynucleotides (SEQ ID NOs: 220 to 235), 100% for 20 polynucleotides (SEQ ID NOs: 216 to 235), 100% for 30 polynucleotides (SEQ ID NOs: 206 to 235), 100% for 50 polynucleotides (SEQ ID NOs: 186 to 235), 100% for 100 polynucleotides (SEQ ID NOs: 136 to 235), 100% for 200 polynucleotides (SEQ ID NOs: 36 to 235), and 100% for 235 polynucleotides (SEQ ID NOs: 1 to 235).

(419) These results demonstrated that a combination of multiple polynucleotides can produce higher breast cancer discriminant performance than that of each polynucleotide alone or a combination of a fewer number of polynucleotides. In this context, the combinations of multiple polynucleotides are not limited to the combinations of the polynucleotides added in the order of statistically significant difference as described above, and any combination of multiple polynucleotides can be used in the detection of breast cancer.

(420) From these results, it can be concluded that all of the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 251 serve as excellent markers for breast cancer detection.

(421) TABLE-US-00002 TABLE 2 Expression level in P value after breast cancer Bonferroni patient with respect SEQ ID NO: Gene name correction to healthy subject 1 hsa-miR-4783-3p 1.88 .Math. E-98 + 2 hsa-miR-4730 1.39 .Math. E-95 - 3 hsa-miR-1307-3p 9.90 .Math. E-81 + 4 hsa-miR-4634 1.73 .Math. E-79 - 5 hsa-miR-663a 2.71 .Math. E-76 + 6 hsa-miR-4532 2.24 .Math. E-75 + 7 hsa-miR-7704 4.18 .Math. E-74 - 8 hsa-miR-3178 2.19 .Math. E-71 - 9 hsa-miR-6729-5p 1.88 .Math. E-69 - 10 hsa-miR-6090 2.55 .Math. E-68 + 11 hsa-miR-4732-5p 1.36 .Math. E-67 + 12 hsa-miR-3184-5p 8.11 .Math. E-67 - 13 hsa-miR-6727-5p 3.02 .Math. E-64 - 14 hsa-miR-6088 3.54 .Math. E-64 + 15 hsa-miR-4674 9.34 .Math. E-64 - 16 hsa-miR-8073 1.47 .Math. E-63 + 17 hsa-miR-4787-5p 5.85 .Math. E-62 - 18 hsa-miR-1469 2.97 .Math. E-61 + 19 hsa-miR-125a-3p 9.95 .Math. E-61 - 20 hsa-miR-1233-5p 2.94 .Math. E-60 + 21 hsa-miR-885-3p 6.54 .Math. E-60 + 22 hsa-miR-6802-5p 4.56 .Math. E-59 + 23 hsa-miR-328-5p 1.83 .Math. E-58 + 24 hsa-miR-6787-5p 5.55 .Math. E-58 + 25 hsa-miR-8069 2.26 .Math. E-57 - 26 hsa-miR-6875-5p 2.72 .Math. E-53 - 27 hsa-miR-1246 9.57 .Math. E-53 + 28 hsa-miR-4734 2.06 .Math. E-52 + 29 hsa-miR-6757-5p 4.71 .Math. E-51 + 30 hsa-miR-6756-5p 5.35 .Math. E-51 + 31 hsa-miR-3665 1.70 .Math. E-50 - 32 hsa-miR-6836-3p 8.38 .Math. E-50 - 33 hsa-miR-6821-5p 1.19 .Math. E-49 + 34 hsa-miR-6805-5p 3.17 .Math. E-49 - 35 hsa-miR-4728-5p 3.69 .Math. E-49 + 36 hsa-miR-6726-5p 5.62 .Math. E-49 - 37 hsa-miR-197-5p 1.43 .Math. E-46 + 38 hsa-miR-149-3p 3.23 .Math. E-46 + 39 hsa-miR-6850-5p 4.58 .Math. E-46 - 40 hsa-miR-4476 5.54 .Math. E-46 - 41 hsa-miR-6858-5p 9.61 .Math. E-46 + 42 hsa-miR-564 1.62 .Math. E-44 - 43 hsa-miR-4763-3p 2.01 .Math. E-44 + 44 hsa-miR-575 3.61 .Math. E-44 - 45 hsa-miR-6771-5p 2.62 .Math. E-43 - 46 hsa-miR-1231 7.46 .Math. E-43 - 47 hsa-miR-1908-3p 9.60 .Math. E-43 - 48 hsa-miR-150-3p 2.19 .Math. E-42 - 49 hsa-miR-3937 9.92 .Math. E-42 - 50 hsa-miR-887-3p 2.38 .Math. E-41 - 51 hsa-miR-3940-5p 3.44 .Math. E-41 - 52 hsa-miR-4741 4.16 .Math. E-41 - 53 hsa-miR-6808-5p 5.42 .Math. E-41 + 54 hsa-miR-6869-5p 1.03 .Math. E-40 - 55 hsa-miR-5090 5.03 .Math. E-40 - 56 hsa-miR-615-5p 1.94 .Math. E-39 - 57 hsa-miR-8072 2.71 .Math. E-39 + 58 hsa-miR-128-l-5p 2.72 .Math. E-39 - 59 hsa-miR-1238-5p 7.46 .Math. E-39 + 60 hsa-miR-365a-5p 9.50 .Math. E-39 + 61 hsa-miR-204-3p 1.32 .Math. E-38 - 62 hsa-miR-4492 3.33 .Math. E-37 - 63 hsa-miR-6785-5p 4.21 .Math. E-37 - 64 hsa-miR-6511a-5p 7.68 .Math. E-37 + 65 hsa-miR-4525 1.16 .Math. E-36 - 66 hsa-miR-1915-5p 1.34 .Math. E-36 - 67 hsa-miR-3180 1.07 .Math. E-35 - 68 hsa-miR-6879-5p 1.56 .Math. E-35 + 69 hsa-miR-1199-5p 1.15 .Math. E-34 - 70 hsa-miR-6746-5p 5.65 .Math. E-34 + 71 hsa-miR-711 5.82 .Math. E-34 - 72 hsa-miR-663b 1.42 .Math. E-33 - 73 hsa-miR-4707-3p 2.19 .Math. E-33 - 74 hsa-miR-6893-5p 6.31 .Math. E-33 - 75 hsa-miR-4675 6.39 .Math. E-33 + 76 hsa-miR-4638-5p 6.40 .Math. E-33 - 77 hsa-miR-4651 8.91 .Math. E-33 - 78 hsa-miR-6087 1.91 .Math. E-32 + 79 hsa-miR-4665-5p 3.57

.Math. E-32 - 80 hsa-miR-4758-5p 4.55 .Math. E-32 + 81 hsa-miR-6887-5p 4.45 .Math. E-31 + 82 hsa-miR-3620-5p 4.64 .Math. E-31 - 83 hsa-miR-1909-3p 5.74 .Math. E-31 - 84 hsa-miR-7641 8.30 .Math. E-31 - 85 hsa-miR-6724-5p 1.02 .Math. E-30 + 86 hsa-miR-1343-3p 1.19 .Math. E-30 + 87 hsa-miR-6780b-5p 1.22 .Math. E-30 + 88 hsa-miR-4484 2.77 .Math. E-30 - 89 hsa-miR-4690-5p 3.50 .Math. E-30 + 90 hsa-miR-4429 2.05 .Math. E-29 + 91 hsa-miR-1227-5p 3.84 .Math. E-29 + 92 hsa-miR-4725-3p 5.39 .Math. E-29 - 93 hsa-miR-6861-5p 5.43 .Math. E-29 + 94 hsa-miR-6812-5p 7.48 .Math. E-29 + 95 hsa-miR-3197 8.20 .Math. E-29 + 96 hsa-miR-8059 9.29 .Math. E-29 + 97 hsa-miR-3185 9.34 .Math. E-29 + 98 hsa-miR-4706 1.69 .Math. E-28 + 99 hsa-miR-4497 2.22 .Math. E-28 - 100 hsa-miR-3131 3.64 .Math. E-28 + 101 hsa-miR-6806-5p 9.04 .Math. E-28 - 102 hsa-miR-187-5p 4.89 .Math. E-27 - 103 hsa-miR-3180-3p 7.10 .Math. E-27 - 104 hsa-miR-6848-5p 7.26 .Math. E-27 - 105 hsa-miR-6820-5p 7.77 .Math. E-27 - 106 hsa-miR-6800-5p 1.80 .Math. E-26 - 107 hsa-miR-6717-5p 1.97 .Math. E-26 + 108 hsa-miR-6795-5p 5.16 .Math. E-26 + 109 hsa-miR-4632-5p 8.43 .Math. E-26 + 110 hsa-miR-665 2.56 .Math. E-25 - 111 hsa-miR-6778-5p 8.69 .Math. E-25 - 112 hsa-miR-3663-3p 1.09 .Math. E-24 + 113 hsa-miR-4689 3.36 .Math. E-24 + 114 hsa-miR-211-3p 8.65 .Math. E-24 + 115 hsa-miR-6511b-5p 9.67 .Math. E-24 + 116 hsa-miR-4750-5p 1.07 .Math. E-23 + 117 hsa-miR-6126 2.06 .Math. E-23 + 118 hsa-miR-614 2.46 .Math. E-22 + 119 hsa-miR-7110-5p 3.56 .Math. E-22 - 120 hsa-miR-744-5p 5.83 .Math. E-22 + 121 hsa-miR-6769a-5p 1.44 .Math. E-21 + 122 hsa-miR-4792 2.04 .Math. E-21 - 123 hsa-miR-5787 3.93 .Math. E-21 + 124 hsa-miR-6798-5p 5.13 .Math. E-21 - 125 hsa-miR-6781-5p 2.43 .Math. E-20 - 126 hsa-miR-4419b 2.59 .Math. E-20 + 127 hsa-miR-4446-3p 7.52 .Math. E-20 - 128 hsa-miR-4259 8.07 .Math. E-20 + 129 hsa-miR-5572 1.06 .Math. E-19 - 130 hsa-miR-6075 1.78 .Math. E-19 - 131 hsa-miR-296-3p 4.75 .Math. E-19 + 132 hsa-miR-689l-5p 8.62 .Math. E-19 - 133 hsa-miR-4745-5p 1.02 .Math. E-18 - 134 hsa-miR-6775-5p 1.17 .Math. E-18 + 135 hsa-miR-6870-5p 1.24 .Math. E-18 - 136 hsa-miR-920 1.78 .Math. E-18 + 137 hsa-miR-4530 3.26 .Math. E-18 - 138 hsa-miR-6819-5p 3.67 .Math. E-18 + 139 hsa-miR-6825-5p 5.28 .Math. E-18 - 140 hsa-miR-7847-3p 7.32 .Math. E-18 + 141 hsa-miR-6131 1.09 .Math. E-17 + 142 hsa-miR-4433-3p 1.35 .Math. E-17 + 143 hsa-miR-1228-5p 9.82 .Math. E-17 - 144 hsa-miR-6743-5p 1.40 .Math. E-16 + 145 hsa-miR-1268a 2.20 .Math. E-16 - 146 hsa-miR-3917 1.73 .Math. E-15 + 147 hsa-miR-6786-5p 1.89 .Math. E-15 - 148 hsa-miR-3154 2.12 .Math. E-15 + 149 hsa-miR-638 2.19 .Math. E-15 - 150 hsa-miR-6741-5p 5.44 .Math. E-15 + 151 hsa-miR-6889-5p 9.29 .Math. E-15 - 152 hsa-miR-6840-3p 1.61 .Math. E-14 - 153 hsa-miR-6510-5p 3.23 .Math. E-14 - 154 hsa-miR-3188 4.44 .Math. E-14 + 155 hsa-miR-551b-5p 1.71 .Math. E-13 + 156 hsa-miR-5001-5p 2.07 .Math. E-13 + 157 hsa-miR-1268b 2.24 .Math. E-13 - 158 hsa-miR-7107-5p 2.31 .Math. E-13 - 159 hsa-miR-6824-5p 3.30 .Math. E-13 + 160 hsa-miR-6732-5p 3.62 .Math. E-13 - 161 hsa-miR-371a-5p 9.05 .Math. E-13 + 162 hsa-miR-6794-5p 9.74 .Math. E-13 + 163 hsa-miR-6779-5p 1.37 .Math. E-12 - 164 hsa-miR-4271 1.69 .Math. E-12 - 165 hsa-miR-5195-3p 1.79 .Math. E-12 + 166 hsa-miR-6762-5p 3.61 .Math. E-12 + 167 hsa-miR-939-5p 4.78 .Math. E-12 - 168 hsa-miR-1247-3p 7.37 .Math. E-12 + 169 hsa-miR-6777-5p 9.79 .Math. E-12 + 170 hsa-miR-6722-3p 1.21 .Math. E-11 + 171 hsa-miR-3656 1.27 .Math. E-11 + 172 hsa-miR-4688 1.86 .Math. E-11 + 173 hsa-miR-3195 2.02 .Math. E-11 - 174 hsa-miR-6766-5p 6.97 .Math. E-11 + 175 hsa-miR-4447 1.08 .Math. E-10 + 176 hsa-miR-4656 1.12 .Math. E-10 - 177 hsa-miR-7108-5p 1.51 .Math. E-10 - 178 hsa-miR-3191-3p 2.67 .Math. E-10 + 179 hsa-miR-1273g-3p 2.89 .Math. E-10 - 180 hsa-miR-4463 4.62 .Math. E-10 + 181 hsa-miR-2861 4.97 .Math. E-10 + 182 hsa-miR-3196 5.22 .Math. E-10 - 183 hsa-miR-6877-5p 6.47 .Math. E-10 - 184 hsa-miR-3679-5p 1.33 .Math. E-09 + 185 hsa-miR-4442 1.56 .Math. E-09 - 186 hsa-miR-6789-5p 1.93 .Math. E-09 - 187 hsa-miR-6782-5p 1.97 .Math. E-09 + 188 hsa-miR-486-3p 2.12 .Math. E-09 - 189 hsa-miR-6085 4.04 .Math. E-09 + 190 hsa-miR-4746-3p 8.57 .Math. E-09 - 191 hsa-miR-619-5p 1.13 .Math. E-08 - 192 hsa-miR-937-5p 1.65 .Math. E-08 + 193 hsa-miR-6803-5p 2.32 .Math. E-08 + 194 hsa-miR-4298 2.33 .Math. E-08 + 195 hsa-miR-4454 2.63 .Math. E-08 + 196 hsa-miR-4459 1.83 .Math. E-07 + 197 hsa-miR-7150 2.60

98.4 100 100 100 100 100 1236 100 100 100 100 100 1237 100 100 100 100 100 100 1238 100
100 100 100 100 100 1239 100 100 100 100 100 100 1240 100 100 100 100 100 100 1241 100 100
100 100 100 100 1242 100 100 100 100 100 100 1243 100 100 100 100 100 100 1244 100 100 100
100 100 100 1245 99.4 98.4 100 100 100 100 1246 99.4 98.4 100 100 100 100 1247 99.4 98.4 100
100 100 100 1248 99.4 98.4 100 100 100 100 1249 99.4 98.4 100 100 100 100 1250 99.4 98.4 100
100 100 100 1251 99.4 98.4 100 100 100 100

(427) TABLE-US-00008 TABLE 7 Training cohort Validation cohort Accuracy Sensitivity
Specificity Accuracy Sensitivity Specificity SEQ ID NO: (%) (%) (%) (%) (%) (%) 3_20 100 100
100 100 100 100 6_20 100 100 100 100 100 100 7_20 100 100 100 100 100 100 10_20 100 100
100 100 100 100 20_22 98.8 96.7 100 98.8 96.8 100 20_238 98.8 96.7 100 98.8 96.8 100 20_239
100 100 100 100 100 100 12_24 100 100 100 98.8 96.8 100 20_24 100 100 100 98.8 96.8 100
24_27 98.8 96.8 100 98.8 96.8 100 24_33 100 100 100 100 100 100 24_236 100 100 100 98.8 96.8
100 24_240 100 100 100 100 100 100 3_26 100 100 100 100 100 100 12_26 100 100 100 98.8
96.8 100 13_26 100 100 100 98.8 96.8 100 17_26 100 100 100 98.8 100 98 19_26 100 100 100
98.8 100 98 3_27 100 100 100 100 100 100 5_27 100 100 100 100 100 100 13_27 99.4 98.4 100
100 100 100 20_27 98.1 96.7 99 100 100 100 26_27 99.4 98.4 100 96.3 93.5 98 27_120 98.8 98.4
99 100 100 100 27_206 98.1 95.2 100 97.5 96.8 98 27_237 99.4 98.4 100 100 100 100 3_30 100
100 100 100 100 100 17_30 99.4 98.4 100 100 100 100 27_30 100 100 100 98.8 96.8 100 27_33
98.8 98.4 99 100 100 100 30_39 99.4 98.4 100 100 100 100 30_117 99.4 98.4 100 98.8 96.8 100
3_33 99.4 98.4 100 100 100 100 7_33 100 100 100 100 100 100 10_33 100 100 100 100 100 100
11_33 100 100 100 98.8 100 98 13_33 100 100 100 100 100 100 25_33 100 100 100 98.8 100 98
33_244 100 100 100 100 100 100 3_182 99.4 98.4 100 100 100 100 6_182 99.4 98.4 100 100 100
100 7_182 100 100 100 100 100 100 12_182 100 100 100 100 100 100 27_182 98.1 96.8 99 97.5
96.8 98 182_236 100 100 100 98.8 96.8 100 2_194 99.4 98.4 100 100 100 100 7_194 100 100 100
100 100 100 27_194 98.1 95.2 100 97.5 96.8 98 194_236 100 100 100 98.8 96.8 100 2_206 99.4
98.4 100 100 100 100 7_206 100 100 100 100 100 100 206_236 100 100 100 98.8 96.8 100 2_208
99.4 98.4 100 100 100 100 7_208 100 100 100 100 100 100 13_208 99.4 98.4 100 100 100 100
20_208 98.1 96.7 99 97.5 93.5 100 27_208 98.1 95.2 100 98.8 96.8 100

Example 3

(428) <Selection of Gene Markers Using all Samples and Method for Evaluating Breast Cancer Discriminant Performance of Acquired Gene Markers>

(429) In this Example, the samples of the training cohort and the validation cohort used in Examples 1 and 2 were integrated, and selection of a gene marker and evaluation of its breast cancer discriminant performance were conducted using all of the samples.

(430) Specifically, the miRNA expression levels in the sera of the 93 breast cancer patients and the 150 healthy subjects obtained in the preceding Reference Examples were normalized by quantile normalization. In order to acquire diagnostic markers with higher reliability, only genes having a gene expression level of 2.sup.6 or higher in 50% or more of the samples in either of the breast cancer patient group or the healthy subject group, were selected in the gene marker selection. In order to further acquire statistical significance for discriminating a breast cancer patient group from a healthy subject group, the P value obtained by two-tailed t-test assuming equal variance as to each gene expression level was corrected by the Bonferroni method, and genes that satisfied $p < 0.01$ were selected as gene markers for use in explanatory variables of a discriminant. The obtained genes are described in Table 7. In this way, hsa-miR-658, hsa-miR-6842-5p, hsa-miR-6124, hsa-miR-6765-3p, hsa-miR-7106-5p, hsa-miR-4534, hsa-miR-92b-3p, hsa-miR-3135b, hsa-miR-4687-3p, hsa-miR-762, hsa-miR-3619-3p, hsa-miR-4467, hsa-miR-557, hsa-miR-1237-5p, hsa-miR-1908-5p, hsa-miR-4286, hsa-miR-6885-5p and hsa-miR-6763-5p genes, and the nucleotide sequences of SEQ ID NOs: 252 to 269 related thereto were found in addition to the genes described in Table 2. As with the nucleotide sequences of SEQ ID NOs: 1 to 251, the results obtained about the polynucleotides represented by the nucleotide sequences of SEQ ID NOs: 252 to

269 also showed that the measurement values were significantly lower (–) or higher (+) in the breast cancer patient group than in the healthy subject group (Table 8). These results were able to be validated in the validation cohort. Thus, the presence or absence of breast cancer in the newly obtained samples can be determined by the methods described in Examples 1 and 2 by using the gene expression level measurement values described in Table 8 either alone or in combination with the gene expression level measurement values described in Table 2

(431) TABLE-US-00009 TABLE 8 Expression level in P value after breast cancer Bonferroni patient with respect SEQ ID NO: Gene name correction to healthy subject 1 hsa-miR-4783-3p 1.81 .Math. E-151 + 2 hsa-miR-4730 1.00 .Math. E-148 – 3 hsa-miR-1307-3p 5.85 .Math. E-126 + 4 hsa-miR-4634 9.92 .Math. E-114 – 5 hsa-miR-663a 2.55 .Math. E-112 – 6 hsa-miR-4532 8.96 .Math. E-113 + 7 hsa-miR-7704 4.82 .Math. E-114 + 8 hsa-miR-3178 5.69 .Math. E-112 – 9 hsa-miR-6729-5p 3.52 .Math. E-112 – 10 hsa-miR-6090 1.64 .Math. E-100 – 11 hsa-miR-4732-5p 7.04 .Math. E-96 – 12 hsa-miR-3184-5p 2.08 .Math. E-103 + 13 hsa-miR-6727-5p 3.11 .Math. E-99 + 14 hsa-miR-6088 4.52 .Math. E-100 – 15 hsa-miR-4674 2.03 .Math. E-94 + 16 hsa-miR-8073 5.49 .Math. E-98 – 17 hsa-miR-4787-5p 1.53 .Math. E-96 + 18 hsa-miR-1469 3.95 .Math. E-87 + 19 hsa-miR-125a-3p 3.12 .Math. E-91 – 20 hsa-miR-1233-5p 2.43 .Math. E-93 + 21 hsa-miR-885-3p 4.92 .Math. E-87 – 22 hsa-miR-6802-5p 5.06 .Math. E-87 – 23 hsa-miR-328-5p 6.06 .Math. E-96 + 24 hsa-miR-6787-5p 2.26 .Math. E-85 + 25 hsa-miR-8069 5.26 .Math. E-78 + 26 hsa-miR-6875-5p 2.38 .Math. E-74 + 27 hsa-miR-1246 1.05 .Math. E-75 + 28 hsa-miR-4734 6.82 .Math. E-87 – 29 hsa-miR-6757-5p 9.57 .Math. E-75 + 30 hsa-miR-6756-5p 4.28 .Math. E-70 – 31 hsa-miR-3665 5.78 .Math. E-78 – 32 hsa-miR-6836-3p 7.87 .Math. E-76 – 33 hsa-miR-6821-5p 1.18 .Math. E-80 – 34 hsa-miR-6805-5p 3.26 .Math. E-79 – 35 hsa-miR-4728-5p 9.76 .Math. E-65 + 36 hsa-miR-6726-5p 4.07 .Math. E-72 + 37 hsa-miR-197-5p 5.63 .Math. E-71 – 38 hsa-miR-149-3p 3.94 .Math. E-70 + 39 hsa-miR-6850-5p 2.45 .Math. E-70 – 40 hsa-miR-4476 8.32 .Math. E-64 + 41 hsa-miR-6858-5p 4.09 .Math. E-67 – 42 hsa-miR-564 5.81 .Math. E-69 + 43 hsa-miR-4763-3p 2.83 .Math. E-72 + 44 hsa-miR-575 1.96 .Math. E-69 – 45 hsa-miR-6771-5p 1.90 .Math. E-65 – 46 hsa-miR-1231 8.50 .Math. E-61 – 47 hsa-miR-1908-3p 3.70 .Math. E-64 + 48 hsa-miR-150-3p 2.41 .Math. E-58 – 49 hsa-miR-3937 5.29 .Math. E-66 – 50 hsa-miR-887-3p 1.78 .Math. E-64 – 51 hsa-miR-3940-5p 2.92 .Math. E-65 – 52 hsa-miR-4741 1.21 .Math. E-57 + 53 hsa-miR-6808-5p 5.95 .Math. E-62 – 54 hsa-miR-6869-5p 9.36 .Math. E-66 – 55 hsa-miR-5090 2.20 .Math. E-62 – 56 hsa-miR-615-5p 1.59 .Math. E-59 – 57 hsa-miR-8072 1.93 .Math. E-53 + 58 hsa-miR-128-l-5p 2.45 .Math. E-57 + 59 hsa-miR-1238-5p 3.00 .Math. E-60 – 60 hsa-miR-365a-5p 5.03 .Math. E-59 + 61 hsa-miR-204-3p 1.93 .Math. E-49 – 62 hsa-miR-4492 6.71 .Math. E-60 – 63 hsa-miR-6785-5p 3.81 .Math. E-56 + 64 hsa-miR-6511a-5p 6.21 .Math. E-62 + 65 hsa-miR-4525 2.38 .Math. E-55 – 66 hsa-miR-1915-5p 8.81 .Math. E-57 – 67 hsa-miR-3180 1.19 .Math. E-50 – 68 hsa-miR-6879-5p 2.38 .Math. E-58 – 69 hsa-miR-1199-5p 7.83 .Math. E-51 – 70 hsa-miR-6746-5p 9.76 .Math. E-50 – 71 hsa-miR-711 1.44 .Math. E-49 + 72 hsa-miR-663b 4.49 .Math. E-49 – 73 hsa-miR-4707-3p 2.18 .Math. E-52 + 74 hsa-miR-6893-5p 2.49 .Math. E-43 – 75 hsa-miR-4675 5.39 .Math. E-56 + 76 hsa-miR-4638-5p 1.64 .Math. E-51 + 77 hsa-miR-4651 6.25 .Math. E-50 – 78 hsa-miR-6087 6.15 .Math. E-53 – 79 hsa-miR-4665-5p 3.35 .Math. E-48 – 80 hsa-miR-4758-5p 1.52 .Math. E-54 – 81 hsa-miR-6887-5p 8.54 .Math. E-46 – 82 hsa-miR-3620-5p 4.32 .Math. E-42 – 83 hsa-miR-1909-3p 6.17 .Math. E-53 + 84 hsa-miR-7641 2.63 .Math. E-43 – 85 hsa-miR-6724-5p 3.58 .Math. E-49 + 86 hsa-miR-1343-3p 1.04 .Math. E-45 – 87 hsa-miR-6780b-5p 5.24 .Math. E-50 – 88 hsa-miR-4484 5.63 .Math. E-49 + 89 hsa-miR-4690-5p 2.68 .Math. E-43 – 90 hsa-miR-4429 7.93 .Math. E-44 – 91 hsa-miR-1227-5p 1.26 .Math. E-46 – 92 hsa-miR-4725-3p 3.25 .Math. E-44 + 93 hsa-miR-6861-5p 2.28 .Math. E-46 + 94 hsa-miR-6812-5p 3.34 .Math. E-37 + 95 hsa-miR-3197 8.32 .Math. E-45 + 96 hsa-miR-8059 6.38 .Math. E-41 + 97 hsa-miR-3185 7.20 .Math. E-46 + 98 hsa-miR-4706 4.07 .Math. E-39 + 99 hsa-miR-4497 7.65 .Math. E-39 + 100 hsa-miR-3131 4.60 .Math. E-42 – 101 hsa-miR-6806-5p 8.93 .Math. E-34 + 102 hsa-miR-187-5p 5.69 .Math. E-42 +

103 hsa-miR-3180-3p 1.37 .Math. E-41 - 104 hsa-miR-6848-5p 9.99 .Math. E-33 - 105 hsa-miR-6820-5p 2.48 .Math. E-37 + 106 hsa-miR-6800-5p 2.58 .Math. E-42 + 107 hsa-miR-6717-5p 2.75 .Math. E-37 - 108 hsa-miR-6795-5p 2.83 .Math. E-36 - 109 hsa-miR-4632-5p 1.50 .Math. E-40 + 110 hsa-miR-665 3.04 .Math. E-37 - 111 hsa-miR-6778-5p 3.57 .Math. E-36 - 112 hsa-miR-3663-3p 2.48 .Math. E-39 + 113 hsa-miR-4689 1.46 .Math. E-43 + 114 hsa-miR-211-3p 1.36 .Math. E-39 + 115 hsa-miR-6511b-5p 4.55 .Math. E-39 + 116 hsa-miR-4750-5p 1.70 .Math. E-32 + 117 hsa-miR-6126 6.52 .Math. E-41 + 118 hsa-miR-614 1.36 .Math. E-31 + 119 hsa-miR-7110-5p 2.57 .Math. E-36 - 120 hsa-miR-744-5p 1.36 .Math. E-32 - 121 hsa-miR-6769a-5p 2.68 .Math. E-28 + 122 hsa-miR-4792 3.78 .Math. E-29 - 123 hsa-miR-5787 5.62 .Math. E-30 + 124 hsa-miR-6798-5p 8.69 .Math. E-34 - 125 hsa-miR-6781-5p 1.34 .Math. E-35 + 126 hsa-miR-4419b 3.49 .Math. E-28 - 127 hsa-miR-4446-3p 2.83 .Math. E-32 - 128 hsa-miR-4259 9.25 .Math. E-30 - 129 hsa-miR-5572 7.92 .Math. E-32 - 130 hsa-miR-6075 1.06 .Math. E-34 - 131 hsa-miR-296-3p 7.28 .Math. E-27 - 132 hsa-miR-6891-5p 1.19 .Math. E-28 + 133 hsa-miR-4745-5p 1.47 .Math. E-30 + 134 hsa-miR-6775-5p 1.24 .Math. E-28 - 135 hsa-miR-6870-5p 1.50 .Math. E-29 - 136 hsa-miR-920 2.07 .Math. E-27 + 137 hsa-miR-4530 4.79 .Math. E-26 + 138 hsa-miR-6819-5p 4.33 .Math. E-27 - 139 hsa-miR-6825-5p 3.99 .Math. E-29 + 140 hsa-miR-7847-3p 1.49 .Math. E-26 + 141 hsa-miR-6131 3.83 .Math. E-28 - 142 hsa-miR-4433-3p 8.75 .Math. E-27 - 143 hsa-miR-1228-5p 4.12 .Math. E-20 - 144 hsa-miR-6743-5p 4.05 .Math. E-31 - 145 hsa-miR-1268a 1.93 .Math. E-25 + 146 hsa-miR-3917 3.71 .Math. E-24 - 147 hsa-miR-6786-5p 3.71 .Math. E-27 + 148 hsa-miR-3154 1.49 .Math. E-21 + 149 hsa-miR-638 2.61 .Math. E-28 + 150 hsa-miR-6741-5p 2.54 .Math. E-24 - 151 hsa-miR-6889-5p 3.14 .Math. E-26 + 152 hsa-miR-6840-3p 4.16 .Math. E-20 - 153 hsa-miR-6510-5p 2.15 .Math. E-19 + 154 hsa-miR-3188 3.01 .Math. E-21 + 156 hsa-miR-5001-5p 1.47 .Math. E-22 - 157 hsa-miR-1268b 1.93 .Math. E-20 + 158 hsa-miR-7107-5p 6.37 .Math. E-21 - 159 hsa-miR-6824-5p 9.39 .Math. E-21 - 160 hsa-miR-6732-5p 3.49 .Math. E-17 - 161 hsa-miR-371a-5p 1.21 .Math. E-19 - 162 hsa-miR-6794-5p 1.02 .Math. E-15 + 163 hsa-miR-6779-5p 6.91 .Math. E-17 + 164 hsa-miR-4271 2.14 .Math. E-22 + 165 hsa-miR-5195-3p 1.17 .Math. E-19 - 166 hsa-miR-6762-5p 5.65 .Math. E-18 + 167 hsa-miR-939-5p 4.70 .Math. E-21 + 168 hsa-miR-1247-3p 4.05 .Math. E-14 + 169 hsa-miR-6777-5p 4.37 .Math. E-16 - 170 hsa-miR-6722-3p 9.50 .Math. E-18 - 171 hsa-miR-3656 1.77 .Math. E-21 + 172 hsa-miR-4688 4.75 .Math. E-17 - 173 hsa-miR-3195 3.33 .Math. E-18 - 174 hsa-miR-6766-5p 1.29 .Math. E-18 - 175 hsa-miR-4447 1.50 .Math. E-14 + 176 hsa-miR-4656 7.31 .Math. E-14 + 177 hsa-miR-7108-5p 1.06 .Math. E-16 - 179 hsa-miR-1273g-3p 3.79 .Math. E-17 + 180 hsa-miR-4463 4.63 .Math. E-18 - 181 hsa-miR-2861 2.60 .Math. E-15 + 182 hsa-miR-3196 6.55 .Math. E-12 + 183 hsa-miR-6877-5p 2.44 .Math. E-17 + 184 hsa-miR-3679-5p 6.45 .Math. E-18 + 185 hsa-miR-4442 3.11 .Math. E-15 - 186 hsa-miR-6789-5p 1.50 .Math. E-14 - 187 hsa-miR-6782-5p 1.14 .Math. E-13 - 188 hsa-miR-486-3p 8.80 .Math. E-15 + 189 hsa-miR-6085 3.88 .Math. E-15 + 190 hsa-miR-4746-3p 2.87 .Math. E-13 - 191 hsa-miR-619-5p 2.89 .Math. E-12 - 192 hsa-miR-937-5p 2.36 .Math. E-13 + 193 hsa-miR-6803-5p 2.14 .Math. E-12 + 194 hsa-miR-4298 1.25 .Math. E-14 + 195 hsa-miR-4454 9.63 .Math. E-11 + 196 hsa-miR-4459 3.66 .Math. E-13 - 197 hsa-miR-7150 1.76 .Math. E-10 + 198 hsa-miR-6880-5p 3.57 .Math. E-11 - 199 hsa-miR-4449 4.62 .Math. E-11 + 200 hsa-miR-8063 4.89 .Math. E-13 + 201 hsa-miR-4695-5p 5.30 .Math. E-09 - 202 hsa-miR-6132 2.15 .Math. E-10 + 203 hsa-miR-6829-5p 3.96 .Math. E-09 - 204 hsa-miR-4486 2.13 .Math. E-07 + 205 hsa-miR-6805-3p 8.96 .Math. E-10 + 206 hsa-miR-6826-5p 7.62 .Math. E-08 - 207 hsa-miR-4508 6.79 .Math. E-08 + 208 hsa-miR-1343-5p 8.96 .Math. E-07 + 209 hsa-miR-7114-5p 1.55 .Math. E-09 + 210 hsa-miR-3622a-5p 2.52 .Math. E-08 - 211 hsa-miR-6765-5p 6.46 .Math. E-07 + 212 hsa-miR-7845-5p 1.78 .Math. E-04 - 213 hsa-miR-3960 2.10 .Math. E-10 - 214 hsa-miR-6749-5p 1.32 .Math. E-07 - 215 hsa-miR-1260b 1.44 .Math. E-04 + 216 hsa-miR-6799-5p 1.14 .Math. E-07 + 217 hsa-miR-4723-5p 6.30 .Math. E-07 - 218 hsa-miR-6784-5p 4.08 .Math. E-08 + 219 hsa-miR-5100 1.78 .Math. E-06 - 220 hsa-miR-6769b-5p 6.44 .Math. E-06 + 221 hsa-miR-1207-5p 8.21

.Math. E-07 - 222 hsa-miR-642a-3p 2.06 .Math. E-09 - 223 hsa-miR-4505 3.06 .Math. E-06 + 224 hsa-miR-4270 9.73 .Math. E-07 + 225 hsa-miR-6721-5p 1.84 .Math. E-08 - 226 hsa-miR-7111-5p 3.24 .Math. E-09 + 227 hsa-miR-6791-5p 2.88 .Math. E-06 - 228 hsa-miR-7109-5p 7.96 .Math. E-05 + 229 hsa-miR-4258 1.36 .Math. E-05 + 230 hsa-miR-6515-3p 3.81 .Math. E-03 + 231 hsa-miR-6851-5p 1.63 .Math. E-06 - 232 hsa-miR-6125 1.36 .Math. E-05 - 233 hsa-miR-4749-5p 7.65 .Math. E-05 - 234 hsa-miR-4726-5p 5.88 .Math. E-04 + 235 hsa-miR-4513 4.79 .Math. E-06 - 236 hsa-miR-760 1.45 .Math. E-111 + 237 hsa-miR-602 1.06 .Math. E-88 - 238 hsa-miR-423-5p 9.24 .Math. E-79 - 239 hsa-miR-92a-2-5p 2.54 .Math. E-85 - 240 hsa-miR-16-5p 1.34 .Math. E-69 + 241 hsa-miR-451a 5.22 .Math. E-54 + 242 hsa-miR-135a-3p 8.02 .Math. E-58 + 243 hsa-miR-486-5p 8.22 .Math. E-52 + 244 hsa-miR-4257 4.02 .Math. E-46 + 245 hsa-miR-92b-5p 1.86 .Math. E-45 - 246 hsa-miR-1915-3p 4.13 .Math. E-28 - 247 hsa-miR-718 1.29 .Math. E-25 + 248 hsa-miR-940 5.23 .Math. E-17 - 249 hsa-miR-296-5p 3.77 .Math. E-16 - 250 hsa-miR-23b-3p 1.91 .Math. E-04 + 251 hsa-miR-92a-3p 3.74 .Math. E-07 + 252 hsa-miR-658 4.76 .Math. E-43 + 253 hsa-miR-6842-5p 3.63 .Math. E-12 - 254 hsa-miR-6124 8.99 .Math. E-06 + 255 hsa-miR-6765-3p 2.30 .Math. E-05 - 256 hsa-miR-7106-5p 4.48 .Math. E-05 - 257 hsa-miR-4534 2.10 .Math. E-04 - 258 hsa-miR-92b-3p 2.53 .Math. E-04 - 259 hsa-miR-3135b 3.35 .Math. E-04 + 260 hsa-miR-4687-3p 3.90 .Math. E-04 + 261 hsa-miR-762 4.64 .Math. E-04 - 262 hsa-miR-3619-3p 1.13 .Math. E-03 + 263 hsa-miR-4467 1.30 .Math. E-03 - 264 hsa-miR-557 1.49 .Math. E-03 + 265 hsa-miR-1237-5p 1.57 .Math. E-03 + 266 hsa-miR-1908-5p 4.45 .Math. E-03 - 267 hsa-miR-4286 5.37 .Math. E-03 - 268 hsa-miR-6885-5p 7.57 .Math. E-03 + 269 hsa-miR-6763-5p 9.02 .Math. E-03 +

Example 4

(432) <Method for Evaluating Breast Cancer-Specific Discriminant Performance by Combination of Multiple Gene Markers Using Samples in the Validation Cohort>

(433) In this Example, gene expression levels of miRNAs in sera were compared between breast cancer patients and a control group consisting of healthy subjects and prostate cancer patients in the same way as the method described in Example 1 using the gene markers selected in Example 1 with respect to the training cohort described in Reference Example 2, to select a gene marker for diagnosis. The polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 851 to 856 thus newly selected were further combined with the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 269 to study a method for evaluating breast cancer-specific discriminant performance.

(434) Specifically, first, the miRNA expression levels in the training cohort and the validation cohort obtained in Reference Example 2 mentioned above were combined and normalized by quantile normalization. Next, Fisher's discriminant analysis was conducted as to combinations of 1 to 2 measurement values that comprise at least one of the measurement values of the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 269, and 851 to 856 to construct a discriminant for determining the presence or absence of breast cancer. Next, accuracy, sensitivity, and specificity in the validation cohort were calculated using the discriminant thus prepared, with a positive sample group that is the breast cancer patient group and a negative sample group that is a combination of the healthy subject group and the prostate cancer patient group. The discriminant performance of the selected polynucleotides was validated using independent samples.

(435) Most of polynucleotides consisting of the nucleotide sequences represented by these SEQ ID NOs (SEQ ID NOs: 1 to 269, and 851 to 856 corresponding to the miRNA markers of Table 1) or complementary sequences thereof mentioned above were able to provide relatively high accuracy, sensitivity, and specificity in the determination of the presence or absence of breast cancer, and furthermore, were able to specifically discriminate breast cancer from the other cancers.

(436) At least one polynucleotide selected from the group consisting of polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15,

16, 17, 20, 21, 22, 23, 24, 25, 26, 27, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 43, 45, 46, 47, 49, 50, 51, 52, 53, 54, 55, 58, 59, 60, 62, 63, 64, 65, 67, 68, 69, 70, 71, 72, 73, 75, 77, 79, 80, 81, 82, 83, 86, 88, 89, 90, 92, 93, 94, 96, 98, 99, 100, 103, 104, 106, 107, 108, 110, 111, 113, 114, 115, 116, 118, 119, 121, 122, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 135, 136, 139, 140, 143, 145, 146, 147, 149, 150, 155, 157, 160, 161, 165, 167, 171, 173, 174, 175, 177, 178, 181, 182, 186, 190, 193, 194, 199, 204, 205, 206, 208, 211, 218, 225, 232, 236, 237, 238, 239, 242, 243, 244, 246, 247, 252, 260, 265, 266, 851, 852, 853, 854, 855 and 856, or complementary sequences thereof (the cancer type-specific polynucleotide group 1) was capable of specifically binding to the target marker. Among the combinations of multiple polynucleotides selected from the cancer type-specific polynucleotide group 1, particularly, combinations comprising at least one polynucleotide selected from the group consisting of polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 16, 17, 20, 21, 22, 23, 24, 25, 26, 27, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 43, 45, 46, 47, 49, 50, 51, 52, 54, 55, 58, 59, 60, 62, 63, 64, 65, 67, 68, 69, 71, 72, 73, 75, 77, 79, 80, 82, 83, 86, 88, 92, 93, 96, 99, 103, 104, 106, 110, 111, 114, 116, 118, 119, 122, 124, 125, 127, 130, 132, 133, 135, 139, 143, 145, 147, 149, 157, 160, 173, 177, 181, 182, 186, 211, 218, 232, 236, 237, 238, 239, 242, 243, 246, 247, 260, 266, 851, 852, 853 and 854, or complementary sequences thereof (the cancer type-specific polynucleotide group 2) were able to specifically discriminate breast cancer from the other cancer with high accuracy.

(437) The number of the polynucleotides with cancer type specificity in the combination mentioned above can be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more for the combination. The combinations of 2 or more of these polynucleotides were able to exhibit discriminant accuracy of 95% or higher.

(438) Specifically, the discriminant accuracy of the measurement using combinations of one or two polynucleotide(s) that consists of a nucleotide sequence represented by any of SEQ ID NOs: 1 to 269 and 851 to 856 or a complementary sequence thereof is shown in Table 9.

(439) For example, the combinations of SEQ ID NOs: 2 and 1, SEQ ID NOs: 2 and 237, SEQ ID NOs: 2 and 4, SEQ ID NOs: 2 and 3, SEQ ID NOs: 2 and 51, SEQ ID NOs: 1 and 237, SEQ ID NOs: 1 and 4, SEQ ID NOs: 1 and 3, SEQ ID NOs: 1 and 51, SEQ ID NOs: 1 and 6, SEQ ID NOs: 4 and 237, SEQ ID NOs: 3 and 237, SEQ ID NOs: 51 and 237, SEQ ID NOs: 237 and 6, SEQ ID NOs: 237 and 12, SEQ ID NOs: 3 and 4, SEQ ID NOs: 4 and 51, SEQ ID NOs: 4 and 6, SEQ ID NOs: 4 and 12, SEQ ID NOs: 4 and 15, SEQ ID NOs: 3 and 51, SEQ ID NOs: 3 and 6, SEQ ID NOs: 3 and 12, SEQ ID NOs: 3 and 15, SEQ ID NOs: 3 and 8, SEQ ID NOs: 51 and 6, SEQ ID NOs: 51 and 12, SEQ ID NOs: 51 and 15, SEQ ID NOs: 51 and 8, SEQ ID NOs: 51 and 34, SEQ ID NOs: 2 and 6, SEQ ID NOs: 12 and 6, SEQ ID NOs: 15 and 6, SEQ ID NOs: 8 and 6, SEQ ID NOs: 6 and 34, SEQ ID NOs: 2 and 12, SEQ ID NOs: 1 and 12, SEQ ID NOs: 12 and 15, SEQ ID NOs: 8 and 12, SEQ ID NOs: 12 and 34, SEQ ID NOs: 2 and 15, SEQ ID NOs: 1 and 15, SEQ ID NOs: 237 and 15, SEQ ID NOs: 8 and 15, SEQ ID NOs: 15 and 34, SEQ ID NOs: 2 and 8, SEQ ID NOs: 1 and 8, SEQ ID NOs: 237 and 8, SEQ ID NOs: 4 and 8, SEQ ID NOs: 8 and 34, SEQ ID NOs: 2 and 34, SEQ ID NOs: 1 and 34, SEQ ID NOs: 237 and 34, SEQ ID NOs: 4 and 34, SEQ ID NOs: 3 and 34, SEQ ID NOs: 2 and 9, SEQ ID NOs: 1 and 9, SEQ ID NOs: 9 and 237, SEQ ID NOs: 4 and 9, SEQ ID NOs: 3 and 9, SEQ ID NOs: 2 and 143, SEQ ID NOs: 1 and 143, SEQ ID NOs: 237 and 143, SEQ ID NOs: 4 and 143, SEQ ID NOs: 3 and 143, SEQ ID NOs: 2 and 13, SEQ ID NOs: 1 and 13, SEQ ID NOs: 237 and 13, SEQ ID NOs: 4 and 13, SEQ ID NOs: 3 and 13, SEQ ID NOs: 2 and 125, SEQ ID NOs: 1 and 125, SEQ ID NOs: 237 and 125, SEQ ID NOs: 4 and 125, SEQ ID NOs: 3 and 125, SEQ ID NOs: 2 and 236, SEQ ID NOs: 1 and 236, SEQ ID NOs: 237 and 236, SEQ ID NOs: 4 and 236, SEQ ID NOs: 3 and 236, SEQ ID NOs: 2 and 46, SEQ ID NOs: 1 and 46, SEQ ID NOs: 237 and 46, SEQ ID NOs: 4 and 46, SEQ ID NOs: 3 and 46, SEQ ID NOs: 2 and 32, SEQ ID NOs: 1 and 32, SEQ ID NOs: 237 and 32, SEQ ID NOs: 4 and 32, SEQ ID NOs: 3 and 32, SEQ ID NOs: 2 and 62, SEQ ID NOs: 1 and 62, SEQ ID NOs: 237 and 62, SEQ ID NOs: 4 and 62, SEQ ID NOs: 3 and 62, SEQ ID NOs: 2 and 88, SEQ ID NOs: 1 and 88, SEQ ID

NOs: 237 and 88, SEQ ID NOs: 4 and 88, SEQ ID NOs: 3 and 88, SEQ ID NOs: 2 and 52, SEQ ID NOs: 1 and 52, SEQ ID NOs: 237 and 52, SEQ ID NOs: 4 and 52, SEQ ID NOs: 3 and 52, SEQ ID NOs: 2 and 7, SEQ ID NOs: 1 and 7, SEQ ID NOs: 237 and 7, SEQ ID NOs: 4 and 7, SEQ ID NOs: 3 and 7, SEQ ID NOs: 2 and 26, SEQ ID NOs: 1 and 26, SEQ ID NOs: 237 and 26, SEQ ID NOs: 4 and 26, SEQ ID NOs: 3 and 26, SEQ ID NOs: 2 and 25, SEQ ID NOs: 1 and 25, SEQ ID NOs: 237 and 25, SEQ ID NOs: 4 and 25, SEQ ID NOs: 3 and 25, SEQ ID NOs: 2 and 54, SEQ ID NOs: 1 and 54, SEQ ID NOs: 237 and 54, SEQ ID NOs: 4 and 54, SEQ ID NOs: 3 and 54, SEQ ID NOs: 2 and 92, SEQ ID NOs: 1 and 92, SEQ ID NOs: 237 and 92, SEQ ID NOs: 4 and 92, SEQ ID NOs: 3 and 92, SEQ ID NOs: 2 and 14, SEQ ID NOs: 1 and 14, SEQ ID NOs: 237 and 14, SEQ ID NOs: 4 and 14, SEQ ID NOs: 3 and 14, SEQ ID NOs: 2 and 242, SEQ ID NOs: 1 and 242, SEQ ID NOs: 237 and 242, SEQ ID NOs: 4 and 242, SEQ ID NOs: 3 and 242, SEQ ID NOs: 2 and 47, SEQ ID NOs: 1 and 47, SEQ ID NOs: 237 and 47, SEQ ID NOs: 4 and 47, SEQ ID NOs: 3 and 47, SEQ ID NOs: 2 and 45, SEQ ID NOs: 1 and 45, SEQ ID NOs: 237 and 45, SEQ ID NOs: 4 and 45, SEQ ID NOs: 3 and 45, SEQ ID NOs: 2 and 39, SEQ ID NOs: 1 and 39, SEQ ID NOs: 237 and 39, SEQ ID NOs: 4 and 39, SEQ ID NOs: 3 and 39, SEQ ID NOs: 2 and 21, SEQ ID NOs: 1 and 21, SEQ ID NOs: 237 and 21, SEQ ID NOs: 4 and 21, SEQ ID NOs: 3 and 21, SEQ ID NOs: 2 and 17, SEQ ID NOs: 1 and 17, SEQ ID NOs: 237 and 17, SEQ ID NOs: 4 and 17, SEQ ID NOs: 3 and 17, SEQ ID NOs: 2 and 83, SEQ ID NOs: 1 and 83, SEQ ID NOs: 237 and 83, SEQ ID NOs: 4 and 83, SEQ ID NOs: 3 and 83, SEQ ID NOs: 2 and 149, SEQ ID NOs: 1 and 149, SEQ ID NOs: 237 and 149, SEQ ID NOs: 4 and 149, SEQ ID NOs: 3 and 149, SEQ ID NOs: 2 and 246, SEQ ID NOs: 1 and 246, SEQ ID NOs: 237 and 246, SEQ ID NOs: 4 and 246, SEQ ID NOs: 3 and 246, SEQ ID NOs: 2 and 22, SEQ ID NOs: 1 and 22, SEQ ID NOs: 237 and 22, SEQ ID NOs: 4 and 22, SEQ ID NOs: 3 and 22, SEQ ID NOs: 2 and 55, SEQ ID NOs: 1 and 55, SEQ ID NOs: 237 and 55, SEQ ID NOs: 4 and 55, SEQ ID NOs: 3 and 55, SEQ ID NOs: 2 and 182, SEQ ID NOs: 1 and 182, SEQ ID NOs: 237 and 182, SEQ ID NOs: 4 and 182, SEQ ID NOs: 3 and 182, SEQ ID NOs: 2 and 73, SEQ ID NOs: 1 and 73, SEQ ID NOs: 237 and 73, SEQ ID NOs: 4 and 73, SEQ ID NOs: 3 and 73, SEQ ID NOs: 2 and 77, SEQ ID NOs: 1 and 77, SEQ ID NOs: 237 and 77, SEQ ID NOs: 4 and 77, SEQ ID NOs: 3 and 77, SEQ ID NOs: 2 and 24, SEQ ID NOs: 1 and 24, SEQ ID NOs: 237 and 24, SEQ ID NOs: 4 and 24, SEQ ID NOs: 3 and 24, SEQ ID NOs: 2 and 103, SEQ ID NOs: 1 and 103, SEQ ID NOs: 237 and 103, SEQ ID NOs: 4 and 103, SEQ ID NOs: 3 and 103, SEQ ID NOs: 2 and 49, SEQ ID NOs: 1 and 49, SEQ ID NOs: 237 and 49, SEQ ID NOs: 4 and 49, SEQ ID NOs: 3 and 49, SEQ ID NOs: 2 and 239, SEQ ID NOs: 1 and 239, SEQ ID NOs: 237 and 239, SEQ ID NOs: 4 and 239, SEQ ID NOs: 3 and 239, SEQ ID NOs: 2 and 23, SEQ ID NOs: 1 and 23, SEQ ID NOs: 237 and 23, SEQ ID NOs: 4 and 23, SEQ ID NOs: 3 and 23, SEQ ID NOs: 2 and 58, SEQ ID NOs: 1 and 58, SEQ ID NOs: 237 and 58, SEQ ID NOs: 4 and 58, SEQ ID NOs: 3 and 58, SEQ ID NOs: 2 and 211, SEQ ID NOs: 1 and 211, SEQ ID NOs: 237 and 211, SEQ ID NOs: 4 and 211, SEQ ID NOs: 3 and 211, SEQ ID NOs: 2 and 147, SEQ ID NOs: 1 and 147, SEQ ID NOs: 237 and 147, SEQ ID NOs: 4 and 147, SEQ ID NOs: 3 and 147, SEQ ID NOs: 2 and 65, SEQ ID NOs: 1 and 65, SEQ ID NOs: 237 and 65, SEQ ID NOs: 4 and 65, SEQ ID NOs: 3 and 65, SEQ ID NOs: 2 and 31, SEQ ID NOs: 1 and 31, SEQ ID NOs: 237 and 31, SEQ ID NOs: 4 and 31, SEQ ID NOs: 3 and 31, SEQ ID NOs: 2 and 72, SEQ ID NOs: 1 and 72, SEQ ID NOs: 237 and 72, SEQ ID NOs: 4 and 72, SEQ ID NOs: 3 and 72, SEQ ID NOs: 2 and 63, SEQ ID NOs: 1 and 63, SEQ ID NOs: 237 and 63, SEQ ID NOs: 4 and 63, SEQ ID NOs: 3 and 63, SEQ ID NOs: 2 and 80, SEQ ID NOs: 1 and 80, SEQ ID NOs: 237 and 80, SEQ ID NOs: 4 and 80, SEQ ID NOs: 3 and 80, SEQ ID NOs: 2 and 37, SEQ ID NOs: 1 and 37, SEQ ID NOs: 237 and 37, SEQ ID NOs: 4 and 37, SEQ ID NOs: 3 and 37, SEQ ID NOs: 2 and 67, SEQ ID NOs: 1 and 67, SEQ ID NOs: 237 and 67, SEQ ID NOs: 4 and 67, SEQ ID NOs: 3 and 67, SEQ ID NOs: 2 and 232, SEQ ID NOs: 1 and 232, SEQ ID NOs: 237 and 232, SEQ ID NOs: 4 and 232, SEQ ID NOs: 3 and 232, SEQ ID NOs: 2 and 127, SEQ ID NOs: 1 and 127, SEQ ID NOs: 237 and 127, SEQ ID NOs: 4 and 127, SEQ ID NOs: 3 and 127, SEQ ID NOs: 2 and 145, SEQ ID NOs: 1 and 145, SEQ ID NOs: 237 and 145, SEQ ID NOs: 4 and 145,

SEQ ID NOs: 3 and 145, SEQ ID NOs: 2 and 16, SEQ ID NOs: 1 and 16, SEQ ID NOs: 237 and 16, SEQ ID NOs: 4 and 16, SEQ ID NOs: 3 and 16, SEQ ID NOs: 2 and 11, SEQ ID NOs: 1 and 11, SEQ ID NOs: 237 and 11, SEQ ID NOs: 4 and 11, SEQ ID NOs: 3 and 11, SEQ ID NOs: 2 and 186, SEQ ID NOs: 1 and 186, SEQ ID NOs: 237 and 186, SEQ ID NOs: 4 and 186, SEQ ID NOs: 3 and 186, SEQ ID NOs: 2 and 50, SEQ ID NOs: 1 and 50, SEQ ID NOs: 237 and 50, SEQ ID NOs: 4 and 50, SEQ ID NOs: 3 and 50, SEQ ID NOs: 2 and 69, SEQ ID NOs: 1 and 69, SEQ ID NOs: 237 and 69, SEQ ID NOs: 4 and 69, SEQ ID NOs: 3 and 69, SEQ ID NOs: 2 and 33, SEQ ID NOs: 1 and 33, SEQ ID NOs: 237 and 33, SEQ ID NOs: 4 and 33, SEQ ID NOs: 3 and 33, SEQ ID NOs: 2 and 247, SEQ ID NOs: 1 and 247, SEQ ID NOs: 237 and 247, SEQ ID NOs: 4 and 247, SEQ ID NOs: 3 and 247, SEQ ID NOs: 2 and 36, SEQ ID NOs: 1 and 36, SEQ ID NOs: 237 and 36, SEQ ID NOs: 4 and 36, SEQ ID NOs: 3 and 36, SEQ ID NOs: 2 and 218, SEQ ID NOs: 1 and 218, SEQ ID NOs: 237 and 218, SEQ ID NOs: 4 and 218, SEQ ID NOs: 3 and 218, SEQ ID NOs: 2 and 43, SEQ ID NOs: 1 and 43, SEQ ID NOs: 237 and 43, SEQ ID NOs: 4 and 43, SEQ ID NOs: 3 and 43, SEQ ID NOs: 2 and 29, SEQ ID NOs: 1 and 29, SEQ ID NOs: 237 and 29, SEQ ID NOs: 4 and 29, SEQ ID NOs: 3 and 29, SEQ ID NOs: 2 and 110, SEQ ID NOs: 1 and 110, SEQ ID NOs: 237 and 110, SEQ ID NOs: 4 and 110, SEQ ID NOs: 3 and 110, SEQ ID NOs: 2 and 20, SEQ ID NOs: 1 and 20, SEQ ID NOs: 237 and 20, SEQ ID NOs: 4 and 20, SEQ ID NOs: 3 and 20, SEQ ID NOs: 2 and 157, SEQ ID NOs: 1 and 157, SEQ ID NOs: 237 and 157, SEQ ID NOs: 4 and 157, SEQ ID NOs: 3 and 157, SEQ ID NOs: 2 and 75, SEQ ID NOs: 1 and 75, SEQ ID NOs: 237 and 75, SEQ ID NOs: 4 and 75, SEQ ID NOs: 3 and 75, SEQ ID NOs: 2 and 82, SEQ ID NOs: 1 and 82, SEQ ID NOs: 237 and 82, SEQ ID NOs: 4 and 82, SEQ ID NOs: 3 and 82, SEQ ID NOs: 2 and 106, SEQ ID NOs: 1 and 106, SEQ ID NOs: 237 and 106, SEQ ID NOs: 4 and 106, SEQ ID NOs: 3 and 106, SEQ ID NOs: 2 and 111, SEQ ID NOs: 1 and 111, SEQ ID NOs: 237 and 111, SEQ ID NOs: 4 and 111, SEQ ID NOs: 3 and 111, SEQ ID NOs: 2 and 96, SEQ ID NOs: 1 and 96, SEQ ID NOs: 237 and 96, SEQ ID NOs: 4 and 96, SEQ ID NOs: 3 and 96, SEQ ID NOs: 2 and 266, SEQ ID NOs: 1 and 266, SEQ ID NOs: 237 and 266, SEQ ID NOs: 4 and 266, SEQ ID NOs: 3 and 266, SEQ ID NOs: 2 and 124, SEQ ID NOs: 1 and 124, SEQ ID NOs: 237 and 124, SEQ ID NOs: 4 and 124, SEQ ID NOs: 3 and 124, SEQ ID NOs: 2 and 68, SEQ ID NOs: 1 and 68, SEQ ID NOs: 237 and 68, SEQ ID NOs: 4 and 68, SEQ ID NOs: 3 and 68, SEQ ID NOs: 2 and 71, SEQ ID NOs: 1 and 71, SEQ ID NOs: 237 and 71, SEQ ID NOs: 4 and 71, SEQ ID NOs: 3 and 71, SEQ ID NOs: 2 and 35, SEQ ID NOs: 1 and 35, SEQ ID NOs: 237 and 35, SEQ ID NOs: 4 and 35, SEQ ID NOs: 3 and 35, SEQ ID NOs: 2 and 173, SEQ ID NOs: 1 and 173, SEQ ID NOs: 237 and 173, SEQ ID NOs: 4 and 173, SEQ ID NOs: 3 and 173, SEQ ID NOs: 2 and 5, SEQ ID NOs: 1 and 5, SEQ ID NOs: 237 and 5, SEQ ID NOs: 4 and 5, SEQ ID NOs: 3 and 5, SEQ ID NOs: 2 and 851, SEQ ID NOs: 1 and 851, SEQ ID NOs: 237 and 851, SEQ ID NOs: 4 and 851, SEQ ID NOs: 3 and 851, SEQ ID NOs: 2 and 852, SEQ ID NOs: 1 and 852, SEQ ID NOs: 237 and 852, SEQ ID NOs: 4 and 852, SEQ ID NOs: 3 and 852, SEQ ID NOs: 2 and 30, SEQ ID NOs: 1 and 30, SEQ ID NOs: 237 and 30, SEQ ID NOs: 4 and 30, SEQ ID NOs: 3 and 30, SEQ ID NOs: 2 and 93, SEQ ID NOs: 1 and 93, SEQ ID NOs: 237 and 93, SEQ ID NOs: 4 and 93, SEQ ID NOs: 3 and 93, SEQ ID NOs: 2 and 27, SEQ ID NOs: 1 and 27, SEQ ID NOs: 237 and 27, SEQ ID NOs: 4 and 27, SEQ ID NOs: 3 and 27, SEQ ID NOs: 2 and 853, SEQ ID NOs: 1 and 853, SEQ ID NOs: 237 and 853, SEQ ID NOs: 4 and 853, SEQ ID NOs: 3 and 853, SEQ ID NOs: 2 and 238, SEQ ID NOs: 1 and 238, SEQ ID NOs: 237 and 238, SEQ ID NOs: 4 and 238, SEQ ID NOs: 3 and 238, SEQ ID NOs: 2 and 130, SEQ ID NOs: 1 and 130, SEQ ID NOs: 237 and 130, SEQ ID NOs: 4 and 130, SEQ ID NOs: 3 and 130, SEQ ID NOs: 2 and 177, SEQ ID NOs: 1 and 177, SEQ ID NOs: 237 and 177, SEQ ID NOs: 4 and 177, SEQ ID NOs: 3 and 177, SEQ ID NOs: 2 and 64, SEQ ID NOs: 1 and 64, SEQ ID NOs: 237 and 64, SEQ ID NOs: 4 and 64, SEQ ID NOs: 3 and 64, SEQ ID NOs: 2 and 114, SEQ ID NOs: 1 and 114, SEQ ID NOs: 237 and 114, SEQ ID NOs: 4 and 114, SEQ ID NOs: 3 and 114, SEQ ID NOs: 2 and 119, SEQ ID NOs: 1 and 119, SEQ ID NOs: 237 and 119, SEQ ID NOs: 4 and 119, SEQ ID NOs: 3 and 119, SEQ ID NOs: 2 and 135, SEQ ID NOs: 1 and 135, SEQ ID NOs:

100 100 1_133 99.5 98.4 100 100 100 100 237_133 98.5 95.2 100 99 96.8 100 4_133 99 96.8 100
99 96.8 100 3_133 99.5 98.4 100 100 100 100 41 91.4 91.4 91.4 91.4 91.4 2_41 99.5 98.4 100
100 100 100 1_41 99.5 98.4 100 100 100 100 237_41 99 96.8 100 99 96.8 100 4_41 99.5 98.4 100
99 96.8 100 3_41 99.5 98.4 100 100 100 100 139 88.3 88.3 88.3 88.3 88.3 2_139 100 100 100
100 100 100 1_139 99.5 98.4 100 100 100 100 237_139 100 100 100 98 93.5 100 4_139 100 100
100 99 96.8 100 3_139 99.5 98.4 100 100 100 100 118 87.8 87.8 87.8 87.8 87.8 2_118 99.5
98.4 100 100 100 100 1_118 99.5 98.4 100 100 100 100 237_118 99 96.8 100 100 100 100 4_118
99.5 98.4 100 100 100 100 3_118 99.5 98.4 100 100 100 100 86 85.3 85.3 85.3 85.3 85.3
2_86 99.5 98.4 100 100 100 100 1_86 99.5 98.4 100 100 100 100 237_86 99 96.8 100 100 100 100
4_86 99 96.8 100 100 100 100 3_86 99.5 98.4 100 99 96.8 100 60 84.8 84.8 84.8 84.8 84.8
2_60 99.5 98.4 100 100 100 100 1_60 99.5 98.4 100 100 100 100 237_60 99 96.8 100 99 96.8 100
4_60 99.5 98.4 100 99 96.8 100 3_60 99.5 98.4 100 100 100 100 116 80.2 80.2 80.2 80.2 80.2
2_116 100 100 100 100 100 1_116 99.5 98.4 100 100 100 100 237_116 99.5 98.4 100 100 100
100 4_116 99.5 98.4 100 99 96.8 100 3_116 99.5 98.4 100 100 100 100 160 80.2 80.2 80.2 80.2
80.2 80.2 2_160 99.5 98.4 100 100 100 100 1_160 99.5 98.4 100 100 100 100 237_160 99.5 98.4
100 98 93.5 100 4_160 99 96.8 100 99 96.8 100 3_160 99.5 98.4 100 100 100 100 38 79.2 79.2
79.2 79.2 79.2 2_38 99.5 98.4 100 100 100 100 1_38 99.5 98.4 100 100 100 100 237_38 98.5
95.2 100 100 100 100 4_38 99 96.8 100 99 96.8 100 3_38 99.5 98.4 100 99 96.8 100 99 86.2 86.2
86.2 86.2 86.2 2_99 99.5 98.4 100 100 100 100 1_99 99.5 98.4 100 100 100 100 237_99 99
96.7 100 99 96.8 100 4_99 99 96.7 100 98 93.5 100 3_99 99.5 98.4 100 100 100 100 104 86.3 86.3
86.3 86.3 86.3 2_104 99.5 98.4 100 100 100 100 1_104 100 100 100 100 100 100 237_104
98.5 95.2 100 98 93.5 100 4_104 99 96.8 100 99 96.8 100 3_104 100 100 100 100 100 100

Comparative Example 1

(442) <Breast Cancer Discriminant Performance of an Existing Tumor Marker in Blood>

(443) The concentrations of the existing tumor marker CEA in blood were measured in the training cohort and the validation cohort obtained in the preceding Reference Examples. When the concentrations of these tumor markers in blood are higher than the reference values described in Non-Patent Literature 3 above (CEA: 5 ng/mL), subjects are usually suspected of having cancer. Thus, whether or not the concentration of CEA in blood exceeded its reference value was confirmed for each sample, and the results were assessed for the ability of the tumor marker to detect cancer in breast cancer patients. The sensitivity of the existing marker in the training cohort and the validation cohort was calculated. The results are shown in Table 5. The sensitivity of CEA was as low as 11.3% in the training cohort and 19.4% in the validation cohort, demonstrating that the marker is not useful in the detection of breast cancer (Table 5).

(444) On the other hand, as shown above in Tables 3 and 6 of Examples 1 and 2, it can be concluded that in all of the polynucleotides consisting of the nucleotide sequences represented by SEQ ID NOs: 1 to 251, combinations of 1 or 2 polynucleotides exhibiting sensitivity beyond the existing breast cancer markers are present, and thus such polynucleotides serve as excellent diagnosis markers.

(445) As shown in these Examples and Comparative Example, the kit, etc., and the method of the present invention can detect breast cancer with higher sensitivity than the existing tumor marker and therefore permit early detection and treatment of breast cancer. As a result, improvement in the survival rate because of the reduced risk of recurrence, and breast conservation therapy as a therapeutic option can also be provided.

INDUSTRIAL APPLICABILITY

(446) According to the present invention, breast cancer can be effectively detected by a simple and inexpensive method. This enables early detection, diagnosis and treatment of breast cancer. The method of the present invention can detect breast cancer with limited invasiveness using the blood of a patient and therefore allows breast cancer to be detected conveniently and rapidly. All

publications, patents, and patent applications cited herein are incorporated herein by reference in their entirety.

Claims

1. A method for detecting breast cancer in a human subject, comprising: measuring expression level of miR-6757-5p in a blood, serum or plasma sample from the subject, comparing the measured expression level of hsa-miR-6757-5p to a control expression level for a healthy subject; detecting an increased level of hsa-miR-6757-5p in the sample from the subject as compared to the control expression level from the sample from the healthy subject; wherein the increased level of hsa-miR-6757-5p indicates that the subject has breast cancer; and wherein the method further comprises treating the subject for the breast cancer or performing a diagnostic procedure on the subject with the breast cancer; wherein the treatment comprises surgery, radiotherapy, chemotherapy or a combination thereof; and wherein the diagnostic procedure comprises mammography, ultrasonography, CT, MRI, abdominal ultrasonography, bone scintigraphy, PET, pathological examination which involves analyzing a lesion tissue under a microscope, or a combination thereof.

2. The method according to claim 1, wherein the expression level of hsa-miR-6757-5p in the sample is measured by using a kit comprising a nucleic acid(s) that specifically binds to hsa-miR-6757-5p.

3. The method according to claim 2, wherein the kit further comprises at least one nucleic acid capable of specifically binding to at least one polynucleotide selected from the group consisting of other breast cancer markers: miR-1307-3p, miR-4634, miR-663a, miR-4532, miR-7704, miR-3178, miR-6729-5p, miR-6090, miR-4732-5p, miR-3184-5p, miR-6727-5p, miR-6088, miR-4674, miR-8073, miR-4787-5p, miR-1469, miR-125a-3p, miR-885-3p, miR-6802-5p, miR-328-5p, miR-6787-5p, miR-8069, miR-6875-5p, miR-1246, miR-4734, miR-6756-5p, miR-3665, miR-6836-3p, miR-6821-5p, miR-6805-5p, miR-4728-5p, miR-6726-5p, miR-197-5p, miR-149-3p, miR-6850-5p, miR-4476, miR-6858-5p, miR-564, miR-4763-3p, miR-575, miR-6771-5p, miR-1231, miR-1908-3p, miR-150-3p, miR-3937, miR-887-3p, miR-3940-5p, miR-4741, miR-6808-5p, miR-6869-5p, miR-5090, miR-615-5p, miR-8072, miR-128-1-5p, miR-1238-5p, miR-365a-5p, miR-204-3p, miR-4492, miR-6785-5p, miR-6511a-5p, miR-4525, miR-1915-5p, miR-3180, miR-6879-5p, miR-1199-5p, miR-6746-5p, miR-711, miR-663b, miR-4707-3p, miR-6893-5p, miR-4675, miR-4638-5p, miR-4651, miR-6087, miR-4665-5p, miR-4758-5p, miR-6887-5p, miR-3620-5p, miR-1909-3p, miR-7641, miR-6724-5p, miR-1343-3p, miR-6780b-5p, miR-4484, miR-4690-5p, miR-4429, miR-1227-5p, miR-4725-3p, miR-6861-5p, miR-6812-5p, miR-3197, miR-8059, miR-3185, miR-4706, miR-4497, miR-3131, miR-6806-5p, miR-187-5p, miR-3180-3p, miR-6848-5p, miR-6820-5p, miR-6800-5p, miR-6717-5p, miR-6795-5p, miR-4632-5p, miR-665, miR-6778-5p, miR-3663-3p, miR-4689, miR-211-3p, miR-6511b-5p, miR-4750-5p, miR-6126, miR-614, miR-7110-5p, miR-744-5p, miR-6769a-5p, miR-4792, miR-5787, miR-6798-5p, miR-6781-5p, miR-4419b, miR-4446-3p, miR-4259, miR-5572, miR-6075, miR-296-3p, miR-6891-5p, miR-4745-5p, miR-6775-5p, miR-6870-5p, miR-920, miR-4530, miR-6819-5p, miR-6825-5p, miR-7847-3p, miR-6131, miR-4433-3p, miR-1228-5p, miR-6743-5p, miR-1268a, miR-3917, miR-6786-5p, miR-3154, miR-638, miR-6741-5p, miR-6889-5p, miR-6840-3p, miR-6510-5p, miR-3188, miR-551b-5p, miR-5001-5p, miR-1268b, miR-7107-5p, miR-6824-5p, miR-6732-5p, miR-371a-5p, miR-6794-5p, miR-6779-5p, miR-4271, miR-5195-3p, miR-6762-5p, miR-939-5p, miR-1247-3p, miR-6777-5p, miR-6722-3p, miR-3656, miR-4688, miR-3195, miR-6766-5p, miR-4447, miR-4656, miR-7108-5p, miR-3191-3p, miR-1273g-3p, miR-4463, miR-2861, miR-3196, miR-6877-5p, miR-3679-5p, miR-4442, miR-6789-5p, miR-6782-5p, miR-486-3p, miR-6085, miR-4746-3p, miR-619-5p, miR-937-5p, miR-6803-5p, miR-4298, miR-4454, miR-4459, miR-7150, miR-6880-5p, miR-4449, miR-8063, miR-4695-5p, miR-6132, miR-6829-5p, miR-4486, miR-6805-3p, miR-6826-5p, miR-

4508, miR-1343-5p, miR-7114-5p, miR-3622a-5p, miR-6765-5p, miR-7845-5p, miR-3960, miR-6749-5p, miR-1260b, miR-6799-5p, miR-4723-5p, miR-6784-5p, miR-5100, miR-6769b-5p, miR-1207-5p, miR-642a-3p, miR-4505, miR-4270, miR-6721-5p, miR-7111-5p, miR-6791-5p, miR-7109-5p, miR-4258, miR-6515-3p, miR-6851-5p, miR-6125, miR-4749-5p, miR-4726-5p, miR-4513, miR-6089, miR-6816-5p, miR-4466, miR-4488, miR-6752-5p and miR-4739, and/or miR-760, miR-602, miR-423-5p, miR-92a-2-5p, miR-16-5p, miR-451a, miR-135a-3p, miR-486-5p, miR-4257, miR-92b-5p, miR-1915-3p, miR-718, miR-940, miR-296-5p, miR-23b-3p, miR-92a-3p, miR-658, miR-6842-5p, miR-6124, miR-6765-3p, miR-7106-5p, miR-4534, miR-92b-3p, miR-3135b, miR-4687-3p, miR-762, miR-3619-3p, miR-4467, miR-557, miR-1237-5p, miR-1908-5p, miR-4286, miR-6885-5p and miR-6763-5p.

4. The method according to claim 1, comprising performing the diagnostic procedure on the subject.

5. The method according to claim 1, wherein the expression level of hsa-miR-6757-5p in the sample is measured by using a device comprising a nucleic acid(s) that specifically binds to hsa-miR-6757-5p.

6. The method according to claim 5, wherein the device further comprises at least one nucleic acid capable of specifically binding to at least one polynucleotide selected from the group consisting of other breast cancer markers: miR-1307-3p, miR-4634, miR-663a, miR-4532, miR-7704, miR-3178, miR-6729-5p, miR-6090, miR-4732-5p, miR-3184-5p, miR-6727-5p, miR-6088, miR-4674, miR-8073, miR-4787-5p, miR-1469, miR-125a-3p, miR-885-3p, miR-6802-5p, miR-328-5p, miR-6787-5p, miR-8069, miR-6875-5p, miR-1246, miR-4734, miR-6756-5p, miR-3665, miR-6836-3p, miR-6821-5p, miR-6805-5p, miR-4728-5p, miR-6726-5p, miR-197-5p, miR-149-3p, miR-6850-5p, miR-4476, miR-6858-5p, miR-564, miR-4763-3p, miR-575, miR-6771-5p, miR-1231, miR-1908-3p, miR-150-3p, miR-3937, miR-887-3p, miR-3940-5p, miR-4741, miR-6808-5p, miR-6869-5p, miR-5090, miR-615-5p, miR-8072, miR-128-1-5p, miR-1238-5p, miR-365a-5p, miR-204-3p, miR-4492, miR-6785-5p, miR-6511a-5p, miR-4525, miR-1915-5p, miR-3180, miR-6879-5p, miR-1199-5p, miR-6746-5p, miR-711, miR-663b, miR-4707-3p, miR-6893-5p, miR-4675, miR-4638-5p, miR-4651, miR-6087, miR-4665-5p, miR-4758-5p, miR-6887-5p, miR-3620-5p, miR-1909-3p, miR-7641, miR-6724-5p, miR-1343-3p, miR-6780b-5p, miR-4484, miR-4690-5p, miR-4429, miR-1227-5p, miR-4725-3p, miR-6861-5p, miR-6812-5p, miR-3197, miR-8059, miR-3185, miR-4706, miR-4497, miR-3131, miR-6806-5p, miR-187-5p, miR-3180-3p, miR-6848-5p, miR-6820-5p, miR-6800-5p, miR-6717-5p, miR-6795-5p, miR-4632-5p, miR-665, miR-6778-5p, miR-3663-3p, miR-4689, miR-211-3p, miR-6511b-5p, miR-4750-5p, miR-6126, miR-614, miR-7110-5p, miR-744-5p, miR-6769a-5p, miR-4792, miR-5787, miR-6798-5p, miR-6781-5p, miR-4419b, miR-4446-3p, miR-4259, miR-5572, miR-6075, miR-296-3p, miR-6891-5p, miR-4745-5p, miR-6775-5p, miR-6870-5p, miR-920, miR-4530, miR-6819-5p, miR-6825-5p, miR-7847-3p, miR-6131, miR-4433-3p, miR-1228-5p, miR-6743-5p, miR-1268a, miR-3917, miR-6786-5p, miR-3154, miR-638, miR-6741-5p, miR-6889-5p, miR-6840-3p, miR-6510-5p, miR-3188, miR-551b-5p, miR-5001-5p, miR-1268b, miR-7107-5p, miR-6824-5p, miR-6732-5p, miR-371a-5p, miR-6794-5p, miR-6779-5p, miR-4271, miR-5195-3p, miR-6762-5p, miR-939-5p, miR-1247-3p, miR-6777-5p, miR-6722-3p, miR-3656, miR-4688, miR-3195, miR-6766-5p, miR-4447, miR-4656, miR-7108-5p, miR-3191-3p, miR-1273g-3p, miR-4463, miR-2861, miR-3196, miR-6877-5p, miR-3679-5p, miR-4442, miR-6789-5p, miR-6782-5p, miR-486-3p, miR-6085, miR-4746-3p, miR-619-5p, miR-937-5p, miR-6803-5p, miR-4298, miR-4454, miR-4459, miR-7150, miR-6880-5p, miR-4449, miR-8063, miR-4695-5p, miR-6132, miR-6829-5p, miR-4486, miR-6805-3p, miR-6826-5p, miR-4508, miR-1343-5p, miR-7114-5p, miR-3622a-5p, miR-6765-5p, miR-7845-5p, miR-3960, miR-6749-5p, miR-1260b, miR-6799-5p, miR-4723-5p, miR-6784-5p, miR-5100, miR-6769b-5p, miR-1207-5p, miR-642a-3p, miR-4505, miR-4270, miR-6721-5p, miR-7111-5p, miR-6791-5p, miR-7109-5p, miR-4258, miR-6515-3p, miR-6851-5p, miR-6125, miR-4749-5p, miR-4726-5p, miR-4513, miR-6089, miR-6816-5p, miR-4466, miR-4488, miR-6752-5p and miR-4739, and/or miR-

760, miR-602, miR-423-5p, miR-92a-2-5p, miR-16-5p, miR-451a, miR-135a-3p, miR-486-5p, miR-4257, miR-92b-5p, miR-1915-3p, miR-718, miR-940, miR-296-5p, miR-23b-3p, miR-92a-3p, miR-658, miR-6842-5p, miR-6124, miR-6765-3p, miR-7106-5p, miR-4534, miR-92b-3p, miR-3135b, miR-4687-3p, miR-762, miR-3619-3p, miR-4467, miR-557, miR-1237-5p, miR-1908-5p, miR-4286, miR-6885-5p and miR-6763-5p.
