

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12409167
Kind Code	B2
Date of Patent	September 09, 2025
Inventor(s)	Clozel; Martine et al.

Pharmaceutical combination comprising a selective S1P.SUB.1 .receptor agonist

Abstract

The present invention relates to a pharmaceutical combination comprising a first active ingredient which is (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]propylimino)-3-o-tolyl-thiazolidin-4-one or a pharmaceutically acceptable salt thereof and a second active ingredient which is selected from the group consisting of methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, and 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate, or a pharmaceutically acceptable salt thereof.

Inventors:	Clozel; Martine (Allschwil, CH), Piali; Luca (Basel, CH)
Applicant:	Vanda Pharmaceuticals Inc. (Washington, DC)
Family ID:	54848562
Assignee:	Vanda Pharmaceuticals Inc. (Washington, DC)
Appl. No.:	18/315954
Filed:	May 11, 2023

Prior Publication Data

Document Identifier	Publication Date
US 20240139160 A1	May. 02, 2024

Foreign Application Priority Data

WO	PCT/EP2014/077469	Dec. 11, 2014
----	-------------------	---------------

Related U.S. Application Data

continuation parent-doc US 17325912 20210520 US 11672783 child-doc US 18315954
continuation parent-doc US 16277829 20190215 US 11026927 20210608 child-doc US 17325912

Publication Classification

Int. Cl.: **A61K31/426** (20060101); **A61K9/00** (20060101); **A61K31/22** (20060101); **A61K31/225** (20060101); **A61K31/27** (20060101); **A61K31/4015** (20060101)

U.S. Cl.:

CPC **A61K31/426** (20130101); **A61K9/0053** (20130101); **A61K31/22** (20130101); **A61K31/225** (20130101); **A61K31/27** (20130101); **A61K31/4015** (20130101); **A61K31/426** (20130101); **A61K2300/00** (20130101); **A61K31/22** (20130101); **A61K2300/00** (20130101); **A61K31/225** (20130101); **A61K2300/00** (20130101); **A61K31/4015** (20130101); **A61K2300/00** (20130101);

Field of Classification Search

CPC: A61K (31/426); A61K (9/0053)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
4959389	12/1989	Speiser et al.	N/A	N/A
7435828	12/2007	Binkert et al.	N/A	N/A
7626037	12/2008	Binkert et al.	N/A	N/A
7875726	12/2010	Binkert et al.	N/A	N/A
7879821	12/2010	Hauser	N/A	N/A
8263780	12/2011	Abele et al.	N/A	N/A
8273779	12/2011	Binkert et al.	N/A	N/A
RE43728	12/2011	Binkert et al.	N/A	N/A
RE43833	12/2011	Binkert et al.	N/A	N/A
8399514	12/2012	Lukashev et al.	N/A	N/A
8492441	12/2012	Legangneux	N/A	N/A
8524752	12/2012	Binkert et al.	N/A	N/A
8669281	12/2013	Zeidan et al.	N/A	N/A
8785484	12/2013	Brossard et al.	N/A	N/A
RE45174	12/2013	Binkert et al.	N/A	N/A
8912340	12/2013	Abele et al.	N/A	N/A
9000018	12/2014	Binkert et al.	N/A	N/A
9062014	12/2014	Bonham et al.	N/A	N/A
9340518	12/2015	Herse	N/A	N/A
10220023	12/2018	Dingemanse et al.	N/A	N/A
10245253	12/2018	Clozel	N/A	A61P 37/02
11026927	12/2020	Clozel	N/A	A61P 1/00
11672783	12/2022	Clozel	514/369	A61P 37/02

2009/0182047	12/2008	Joshi et al.	N/A	N/A
2010/0048651	12/2009	Gangakhedkar et al.	N/A	N/A
2010/0160259	12/2009	Schmouder	N/A	N/A
2010/0160369	12/2009	Canne Bannen	N/A	N/A
2010/0260755	12/2009	Gammans et al.	N/A	N/A
2011/0112196	12/2010	Lukashev	N/A	N/A
2011/0200557	12/2010	Grima et al.	N/A	N/A
2011/0274657	12/2010	Aguilar et al.	N/A	N/A
2011/0311485	12/2010	Giulio et al.	N/A	N/A
2012/0034303	12/2011	Nilsson et al.	N/A	N/A
2012/0238028	12/2011	Reszka et al.	N/A	N/A
2013/0259856	12/2012	Kaye et al.	N/A	N/A
2013/0314669	12/2012	Levin et al.	N/A	N/A
2013/0316003	12/2012	Nilsson et al.	N/A	N/A
2014/0056973	12/2013	Ma et al.	N/A	N/A
2014/0057917	12/2013	Cundy et al.	N/A	N/A
2014/0099364	12/2013	Nilsson	424/464	A61K 9/2027
2014/0163100	12/2013	Dawson et al.	N/A	N/A
2014/0179778	12/2013	Mao et al.	N/A	N/A
2014/0303217	12/2013	Brossard et al.	N/A	N/A
2014/0315964	12/2013	Brossard et al.	N/A	N/A
2014/0316140	12/2013	Brossard et al.	N/A	N/A
2015/0183741	12/2014	Guckian et al.	N/A	N/A
2015/0203493	12/2014	Guckian et al.	N/A	N/A
2015/0220693	12/2014	Cadavid et al.	N/A	N/A
2015/0238602	12/2014	Cadavid et al.	N/A	N/A
2015/0265580	12/2014	Brossard et al.	N/A	N/A
2019/0091190	12/2018	Terwey et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
07527/33	12/2001	AU	N/A
2013329421	12/2014	AU	N/A
2834455	12/2011	CA	N/A
102009008851	12/2009	DE	N/A
102010048788	12/2010	DE	N/A
102010026879	12/2010	DE	N/A
0312697	12/1988	EP	N/A
2137537	12/2008	EP	N/A
2177521	12/2009	EP	N/A
2202232	12/2009	EP	N/A
2210890	12/2009	EP	N/A
2305660	12/2010	EP	N/A
2316430	12/2010	EP	N/A
2343287	12/2010	EP	N/A
2366702	12/2010	EP	N/A
2390252	12/2010	EP	N/A

2455080	12/2011	EP	N/A
2692344	12/2013	EP	N/A
2005535679	12/2004	JP	N/A
2012-500285	12/2011	JP	N/A
2014-515373	12/2013	JP	N/A
102427123	12/2021	KR	N/A
00/30622	12/1999	WO	N/A
2004010987	12/2003	WO	N/A
2005005421	12/2004	WO	N/A
2005/054215	12/2004	WO	N/A
2005082841	12/2004	WO	N/A
2006058316	12/2005	WO	N/A
2008/062376	12/2007	WO	N/A
2008/097596	12/2007	WO	N/A
2009/115954	12/2008	WO	N/A
2010/022177	12/2009	WO	N/A
WO-2010046835	12/2009	WO	A61K 31/426
2010/079222	12/2009	WO	N/A
2011/035900	12/2010	WO	N/A
2011/067243	12/2010	WO	N/A
2011/069647	12/2010	WO	N/A
2011/113578	12/2010	WO	N/A
2011/144338	12/2010	WO	N/A
2012/162669	12/2011	WO	N/A
2012/164026	12/2011	WO	N/A
2013/119677	12/2012	WO	N/A
2013/148690	12/2012	WO	N/A
2014/018891	12/2013	WO	N/A
2014/020156	12/2013	WO	N/A
2014/027330	12/2013	WO	N/A
2014/028299	12/2013	WO	N/A
2014/031844	12/2013	WO	N/A
2014/031892	12/2013	WO	N/A
2014/058875	12/2013	WO	N/A
2014/071371	12/2013	WO	N/A
2014/152494	12/2013	WO	N/A
2015/066515	12/2014	WO	N/A
2016/091996	12/2015	WO	N/A
2016/092042	12/2015	WO	N/A

OTHER PUBLICATIONS

Curtin et al., Novel therapeutic options for multiple sclerosis, Exert Rev. Clin. Pharmacol., 7(1), 91-104 (2014). cited by examiner

Brennan et al. Dimethyl Fumarate and Monomethyl Fumarate are Distinguished by Non-Overlapping Pharmacodynamic Effects In Vivo (P1.206), Neurology, Apr. 8, 2014; 82 (10 Supplement). cited by examiner

“Abstracts—ACTRIMS Forum 2017”, Multiple Sclerosis Journal, vol. 23, p. 2-90, (2017). cited by applicant

Brennan et al., “Dimethyl Fumarate and Monomethyl Fumarate are Distinguished by Non-

Overlapping Pharmacodynamic Effects In Vivo (P1.206)", *Neurology*, 2014, vol. 82, Suppl. 10. cited by applicant

Brossard et al., "Pharmacokinetics and Pharmacodynamics Of Ponesimod, A Selective S1P1 Receptor Modulator, In the first-in-human study", *British Journal of Clinical Pharmacology*, 2013, vol. 76, No. 6, pp. 888-896. cited by applicant

Cohen et al., "The Future Of Multiple Sclerosis Treatment", *Journal of the Neurological Sciences*, 2009, vol. 277, No. 51, pp. S55-S61. cited by applicant

Curtin et al., "Novel Therapeutic Options For Multiple Sclerosis", *Exert Rev. Clin. Pharmacol.*, 2014, vol. 7, No. 1, pp. 91-104. cited by applicant

Ellwardt et al., "Molecular Mechanisms Linking Neuroinflammation and Neurodegeneration in MS," *Experimental Neurology*, 2014, vol. 262, pp. 8-17. cited by applicant

Fernandez et al., "Combination Therapy In Multiple Sclerosis", *Journal of the Neurological Sciences*, 2007, vol. 259, pp. 95-103. cited by applicant

Fernandez et al., "Review of the novelties presented at the 28th Congress of the European Committee for Treatment and Research in Multiple Sclerosis (ECTRIMS) (III)", *Rev. Neurol.*, 2013, vol. 57, No. 7, pp. 317-329. cited by applicant

Olsson et al., "Oral Ponesimod In Relapsing-Remitting Multiple Sclerosis: A Randomised Phase II Trial", *Journal of Neurology, Neurosurgery and Psychiatry*, 2014, vol. 85, pp. 1198-1208. cited by applicant

Pitarokoili et al., "Lewis Rat Model of Experimental Autoimmune Encephalomyelitis," *Current Protocols in Neuroscience*, 2017, vol. 81, pp. 1-22. cited by applicant

Pouzol et al., "Complete Resolution Of Clinical Signs And Synergism Of The Combination Ponesimod-Dimethyl Fumarate In Rat Models Of Multiple Sclerosis", *Actelion Pharmaceuticals Ltd—Poster DMF-ponesimod ACTRIMS*, 2017. cited by applicant

Remington, "The Science and Practice of Pharmacy", 21st Ed., Part 5, *Pharmaceutical Manufacturing*, 2005. cited by applicant

Shin et al., "Mechanisms of Experimental Autoimmune Encephalomyelitis in Lewis Rats: Recent Insights from Macrophages," *Anat Cell Biol*, 2012, vol. 45, pp. 141-148. cited by applicant

Shurin et al., "Immune-Mediated Diseases: Where Do We Stand?", Part Of The *Advances In Experimental Medicine And Biology Book Series AEMB*, 2007, vol. 601, pp. 1-12. cited by applicant

Stahl et al., "Handbook of Pharmaceutical Salts, Properties, Selection and Use", 2008, pp. 330-350. cited by applicant

Weaver et al., "An Elevated Matrix Metalloproteinase (MMP) In An Animal Model Of Multiple Sclerosis Is Protective By Affecting Th1/Th2 polarization", *The FASEB Journal*, 2005, vol. 19, No. 12, pp. 1668-1670. cited by applicant

Wouters et al., "Pharmaceutical Salts and Co-crystals", *RSC Publishing*, 2012. cited by applicant

Wustrow et al., "Comparison of the Efficacy and Tolerability of a Novel Methyl Hydrogenfumarate Prodrug with Dimethyl Fumarate in Rodent EAE and GI Irritation Models", *Neurology*, 2011, vol. 76, No. 9, Suppl. 4, pp. A387. cited by applicant

"Actelion's Orally Active Selective S1P1 Receptor Agonist to be Jointly Developed/ Promoted with Roche in Autoimmune Disorders and Transplantation; Deal Potentially Worth well over US\$630 Million to Actelion", *USCULO Eletal Report; Translating Research into Practice*, p. 1, (2009). cited by applicant

Brinkmann, "Sphingosine 1-phosphate receptors in health and disease: Mechanistic insights from gene deletion studies and reverse pharmacology", *Pharmacology & Therapeutics*, vol. 115, p. 84-105, (2007). cited by applicant

Brossard et al., "Effects on Heart Rate of Three Different Up-titration Regimens of Ponesimod, A Selective S1P1 Receptor Modulator, and of Re-Initiation of Treatment in Healthy Male and Female Subjects," *Multiple Sclerosis Journal*, vol. 18, No. S4, p. 192, Abstract P468, 2 pages, (2012). cited

by applicant

Brossard et al., “Effects of three different up-titration regimens of ponesimod, a selective S1P1 receptor agonist, on heart rate”, ECTRIMS Lyon France, p. 109, (2012). cited by applicant

Brossard et al., “Ascending multiple-dose study with ponesimod, a selective S1P1 receptor agonist: tolerability, safety, pharmacokinetics, and pharmacodynamics”, ECTRIMS Lyon France, p. 109, (2012). cited by applicant

Brossard et al., “Ascending single-dose study with ponesimod, a selective S1P1 receptor agonist: safety, pharmacokinetics, pharmacodynamics, and modeling”, ECTRIMS Lyon France, p. 111, (2012). cited by applicant

Brossard et al., “Pharmacokinetics and pharmacodynamics of ponesimod, a selective S1P1 receptor modulator, in the first-in-human study”, British Journal of Clinical Pharmacology, vol. 76(6), p. 888-896, (2013). cited by applicant

Brossard et al., “A randomised, double-blind, ascending multiple-dose study with ponesimod, a selective S1P1 receptor modulator: safety, pharmacokinetics, pharmacodynamics”, ECTRIMS Lyon France, (2012). cited by applicant

Brossard P et al, “Multiple-dose tolerability, pharmacokinetics, and pharmacodynamics of ponesimod, an S1P1 receptor modulator: favorable impact of dose up-titration.”, Journal of Clinical Pharmacology Feb. 2014, (Feb. 2014), vol. 54, No. 2, ISSN 1552-4604, pp. 179-188, XP002754016. cited by applicant

Brossard et al., “PII-109; Effects of Three Different Up-Titration Regimens of Ponesimod, a Selective S1p1 Receptor Agonist, on Heart Rate”, Nature, vol. 91, p. S92-S93, (2012). cited by applicant

Brossard et al., “PII-111; Ascending Single-Dose Study With Ponesimod, a Selective S1p1 Receptor Agonist: Safety, Pharmacokinetics, Pharmacodynamics, and Modeling”, Nature, vol. 91, p. S93, (2012). cited by applicant

Brossard et al., “PII-110; Ascending Multiple-Dose Study With Ponesimod, A Selective S 1p1 Receptor Agonist: Tolerability, Safety, Pharmacokinetics, And Pharmacodynamics”, Nature, vol. 91, p. S93, (2012). cited by applicant

Bunemann et al., “Activation of Muscarinic K⁺ Current in Guinea-Pig Atrial Myocytes by sphingosine-1-phosphate”, Journal of Physiology, vol. 489.3, p. 701-707, (1995). cited by applicant

Davidov et al., “Chronic nitric oxide synthase blockade desensitizes the heart to the negative metabolic effects of nitric oxide”, Life Sciences, vol. 79, p. 1674-1680, (2006). cited by applicant

Frolkis et al., “The Role of Invertors (Intracellular Activators) in Age-Related Changes in Cell Response to Hormones”, Experimental Gerontology, vol. 30(¾), p. 401-414, (1995). cited by applicant

Fujishiro et al., “Use of Sphingosine-1-Phosphate 1 Receptor Agonist, KRP-203, in Combination with a Subtherapeutic Dose of Cyclosporine A for Rat Renal Transplantation”, Transplantation, vol. 82(6), p. 804-812, (2006). cited by applicant

Guo et al., “Effects of Sphingosine 1-phosphate on pacemaker activity in rabbit sino-atrial node cells”, European Journal of Physiology, vol. 438, p. 642-648, (1999). cited by applicant

Hale et al., “Selecting against S1P3 enhances the acute cardiovascular Tolerability of 3-(N-benzyl) aminopropylphosphonic acid S 1P receptor agonists”, Bioorganic & Medicinal Chemistry Letters, vol. 14, p. 3501-3505, (2004). cited by applicant

Himmel et al., “Evidence for Edg-3 Receptor-Mediated Activation of IK.ACh by Sphingosine-1-Phosphate in Human Atrial Cardiomyocytes”, Molecular Pharmacology, vol. 58, p. 449-454, (2000). cited by applicant

Hoch et al., “Effect of Ponesimod, a Selective S1P1 Receptor Modulator, on the QT Interval in Healthy Individuals”, Basic & Clinical Pharmacology & Toxicology, vol. 116, p. 429-437, (2015). cited by applicant

Hoch et al., Oc015—Up-Titration Study With Ponesimod, a Selective S1p1 Receptor Modulator, to

Assess Its Maximum Tolerated Dose in Healthy Subjects, *Clinical Therapeutics*, vol. 35(8S), p. e6-e7, (2013). cited by applicant

Hoch et al., “The Proceedings of the 12th Conference of the European Association for Clinical Pharmacology and Therapeutics”, *Clinical Therapeutics*, vol. XX(xx), (2015). cited by applicant

Hoch et al., “PP162—Thorough QT study with ponesimod, a selective S1P1 receptor modulator”, *Clinical Therapeutics*, vol. 35(8S), p. e68-e69, (2013). cited by applicant

Hoch et al., “A Novel Gradual Up-Titration Regimen Mitigates The First-Dose Effects Of Ponesimod, A Selective S1P1 Receptor Modulator”, *Clinical Therapeutics*, vol. 37 (8S), e36e37, (2015). cited by applicant

Hoch et al., “Clinical pharmacology of ponesimod, a selective S1P1 receptor modulator, after up titration to supratherapeutic doses in healthy subjects”, *European Journal of Pharmaceutical Sciences*, vol. 63, p. 147-153, (2014). cited by applicant

Huwiler et al., “New players on the center stage: Sphingosine 1-phosphate and its receptors as drug targets”, *Biochemical Pharmacology*, p. 1-8, (2008). cited by applicant

International Search Report of International Application No. PCT/EP2015/079208, dated Mar. 4, 2016, 4 pages. cited by applicant

Kappos et al., “Siponimod (BAF312) for the treatment of Secondary Progressive Multiple Sclerosis: Design of the Phase 3 Expand Trial”, *Multiple Sclerosis and Related Disorders*, vol. 3(6), p. 752, (2014). (<http://dx.doi.org/10.1016/j.msard.2014.09.185>). cited by applicant

Keller et al., “Immunomodulator FTY720 Induces Myofibroblast Differentiation via the Lysophospholipid Receptor S1P3 and Smad3 Signaling”, *The American Journal of Pathology*, vol. 170, p. 281-292, (2007). cited by applicant

Kovarik et al., “A Mechanistic Study to Assess Whether Isoproterenol Can Reverse the Negative Chronotropic Effect of Fingolimod”, *Pharmacodynamics*, p. 1-8, (2008). cited by applicant

Koyrakh et al., “The Heart Rate Decrease Caused by Acute FTY720 Administration is Mediated by the G-Protein-Gated Potassium Channel IKACH”, *American Journal of Transplantation*, vol. 5, p. 529-536, (2005). cited by applicant

Krause et al., “Population modeling of the pharmacokinetics and pharmacodynamics of ponesimod, a selective S1P1 receptor agonist”, *University of Auckland, New Zealand*, p. 78, (2012). cited by applicant

Krause et al., “Population modeling of the pharmacokinetics and pharmacodynamics of ponesimod, a selective S1P1 receptor agonist”, *Nature*, vol. 91, p. S36-S37, (2012). cited by applicant

Krause et al., “Population pharmacokinetics and pharmacodynamics of ponesimod, a selective S1P1 receptor modulator”, *Journal of Pharmacokinetics Pharmacodynamics*, vol. 41, p. 261-278, (2014). cited by applicant

Legangneux et al., “BAF312 Dose-Titration Scheme Significantly Attenuates Initial Heart Rate Reducing Effects in Healthy Subjects”, *63rd American Academy of Neurology Annual Meeting, Honolulu Hawaii*, (2011). cited by applicant

Legangneux et al., “BAF312 Dose-Titration Scheme Significantly Attenuates Initial Heart Rate Reducing Effects in Healthy Subjects”, *British Journal of Clinical Pharmacology*, vol. 75(3), p. 831-841, (2012). cited by applicant

Ochi et al., “Sphingosine-1-phosphate Effects on Guinea Pig Atrial Myocytes: Alterations in Action Potentials and K⁺ Currents”, *Cardiovascular Research*, vol. 70, p. 88-96, (2006). cited by applicant

T. Olsson, Boster A., Fernandez O., Freedman M. S., Pozzilli C., Bach D., Berkani O., Mueller M. S., Sidorenko T., Radue E.-W., Melanson M., “Oral ponesimod in relapsing-remitting multiple sclerosis: a randomised phase II trial”, *Journal of Neurology Neurosurgery & Psychiatry*, GB , (Nov. 1, 2014), vol. 85, No. 11, doi: 10.1136/jnnp-2013-307282, ISSN 0022-3050, pp. 1198-1208, XP055734579. cited by applicant

Peters et al., “Sphingosine-1-phosphate signaling in the cardiovascular system”, *Current Opinion in Pharmacology*, vol. 7, p. 1-7, (2007). cited by applicant

Rey et al., “Desensitization by Progressive Up-Titration Prevents First-Dose Effects on the Heart: Guinea Pig Study with Ponesimod, a Selective S1P1 Receptor Modulator”, *PlosOne*, vol. 8(9), p. e74285 (1-7), (2013). cited by applicant

Reyes et al., “Effects of ponesimod, a selective S1P1 receptor modulator, on the pharmacokinetics of a hormonal combination contraceptive”, *European Journal of Clinical Pharmacology*, vol. 70, p. 287-293, (2014). cited by applicant

Sanna et al., “Sphingosine 1-Phosphate (S1P) Receptor Subtypes S1P1 and S1P3, Respectively, Regulate Lymphocyte Recirculation and Heart Rate”, *The Journal of Biological Chemistry*, vol. 279(4), p. 13839-13848, (2004). cited by applicant

Scherz Michael W., Brossard Patrick, D'ambrosio Daniele, Ipek Murat, Dingemanse Jasper, “Three different up-titration regimens of ponesimod, an S1P 1 receptor modulator, in healthy subjects : *Journal of Clinical Pharmacology*”, *The Journal of clinical pharmacology*, US , (Jun. 1, 2015), vol. 55, No. 6, doi:10.1002/jcph.467, ISSN 0091-2700, pp. 688-697, XP055980306. cited by applicant

Schmouder et al., “FTY720: Placebo-Controlled Study of the Effect on Cardiac Rate and Rhythm in Healthy Subjects”, *Journal of Clinical Pharmacology*, vol. 46, p. 895-904, (2006). cited by applicant

Vaclavkova et al., “Oral ponesimod in patients with chronic plaque psoriasis: a randomised, double-blind, placebo-controlled phase 2 trial”, *The Lancet*, p. 1-10, (2014). cited by applicant

Wouters et al., “Pharmaceutical Salts and Cocrystals”, *RSC Drug Discovery*, p. 1-10, (2012). cited by applicant

Juif et al., “Mitigation of Initial Cardiodynamic Effects of the S1P1 Receptor Modulator Ponesimod Using a Novel Up-Titration Regimen”, *The Journal of Clinical Pharmacology*, vol. 57(3), p. 401-410, (2017). cited by applicant

Lott et al., “Tolerance modeling: effects of the selective S1P1 receptor modulator ponesimod on heart rate”, *Universitat des Saarlandes—Population Approach Group Europe*, 26th Meeting, Budapest 2017, p. 1, (2017). cited by applicant

Lott et al., “Modeling Tolerance Development for the Effect on Heart Rate of the Selective S 1Pi Receptor Modulator Ponesimod,” *Clinical Pharmacology & Therapeutics*, 10 pages (2017); DOI: 10.1002/cpt.877. cited by applicant

Piali et al., “The Selective Sphingosine 1-Phosphate Receptor 1 Agonist Ponesimod Protects against Lymphocyte-Mediated Tissue Inflammation,” *The Journal of Pharmacology and Experimental Therapeutics*, vol. 337, No. 2, p. 547-556, (2011). cited by applicant

Basile Jan N., “Titration of beta-blockers in heart failure. How to maximize benefit while minimizing adverse events”, *Postgraduate Medicine*, (Mar. 1, 2003), vol. 113, No. 3, pp. 63-70, XP055980347. cited by applicant

De Vries et al., “Guide to Good Prescribing: A Practical Manual,” *WHO Action Programme on Essential Drugs*, 11 pages, (1994). cited by applicant

James p. Mc Cormack, G. Michael Allan Md, Adil S. Virani, “Is Bigger Better? An argument for very low starting doses”, *CMAJ*, (Jan. 11, 2011), vol. 183, No. 1, pp. 65-69, XP055913813. cited by applicant

International Search Report of International Application No. PCT/EP2015/079311, dated Feb. 25, 2016, 7 pages. cited by applicant

Brossard, vol. 54, Issue2 Feb. 2014 pp. 179-188 (Year: 2014). cited by applicant

Daniele D'Ambrosio, Ponesimod, a selective S1P1 receptor modulator: a potential treatment for multiple sclerosis and other immune-mediated diseases, *Therapeutic Advances in Chronic Disease* vol. 7, Issue 1, Jan. 2016, pp. 18-33 (Year: 2016). cited by applicant

De Vries T.P.G.M., Henning R.H., Hogerzeil H.V., Fresle D.A., “Annex 1—Essentials of pharmacology in daily practice”, *Guide to Good Prescribing—A practical manual*, (Jan. 1, 1998), pp. 98-104, XP055980341. cited by applicant

Notice of Allowance dated Sep. 30, 2024 for U.S. Appl. No. 18/046,492 (pp. 1-8). cited by

applicant

Notice of Allowance dated Oct. 9, 2024 for U.S. Appl. No. 18/046,492 (pp. 1-6). cited by applicant

U.S. Appl. No. 17/325,912, filed May 20, 2021. cited by applicant

U.S. Appl. No. 16/277,829, filed Feb. 15, 2019. cited by applicant

U.S. Appl. No. 15/534,951, filed Jun. 9, 2017. cited by applicant

Primary Examiner: Ivanova; Svetlana M

Attorney, Agent or Firm: Hoffman Warnick LLC

Background/Summary

CROSS REFERENCE TO RELATED APPLICATIONS (1) This application is a continuation of U.S. application Ser. No. 17/325,912 filed May 20, 2021, which is a continuation application of U.S. patent application Ser. No. 16/277,829, filed Feb. 15, 2019, now U.S. Pat. No. 11,026,927, which is a divisional application of U.S. application Ser. No. 15/534,951, filed on Jun. 9, 2017, now U.S. Pat. No. 10,245,253, which claims the benefit of United States Application under 35 U.S.C. 371 claiming benefit of PCT Application No. PCT/EP2015/079311, filed on Dec. 10, 2015, which claims the benefit of PCT Application No. PCT/EP2014/077469, filed on Dec. 11, 2014, the contents of each of which are incorporated herein by reference.

(1) The present invention relates to a pharmaceutical combination comprising a first active ingredient which is (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one or a pharmaceutically acceptable salt thereof and a second active ingredient which is selected from the group consisting of methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, and 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate, or a pharmaceutically acceptable salt thereof.

Description

DESCRIPTION OF THE FIGURE

(1) FIG. 1: Efficacy of ponesimod, DMF or its combination in rat EAE (n=11/group). Clinical scoring was performed daily by an independent examiner in a blinded manner. SEM=standard error of the mean.

DESCRIPTION OF THE INVENTION

(2) 1) In a first embodiment the present invention relates to a pharmaceutical combination comprising a first active ingredient which is (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one (hereinafter also referred to as "COMPOUND 1") or a pharmaceutically acceptable salt thereof and a second active ingredient which is selected from the group consisting of methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, and 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate, or a pharmaceutically acceptable salt thereof.

(3) WO 2010/046835 discloses different crystalline forms of COMPOUND 1; it is to be understood that the present invention encompasses COMPOUND 1 in any form including amorphous as well as crystalline forms of COMPOUND 1. It is further to be understood that crystalline forms of COMPOUND 1 encompasses all types of crystalline forms of COMPOUND 1 including polymorphs of the mere molecule, solvates and hydrates, molecular salts and co-crystals (when the

same molecule can be co-crystallized with different co-crystal formers) provided they are suitable for pharmaceutical administration. In a preferred embodiment, COMPOUND 1 is in crystalline form A or C as described in WO 2010/046835. In a most preferred embodiment, COMPOUND 1 is in crystalline form C.

(4) Likewise it is to be understood that the present invention encompasses methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, and 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate in any form including amorphous as well as crystalline forms as described for COMPOUND 1 in the preceding paragraph. The term “methyl fumarate” as used herein refers to (E)-4-methoxy-4-oxobut-2-enoic acid and/or pharmaceutically acceptable salts thereof.

(5) COMPOUND 1 is a selective SIP, receptor agonist and oral administration thereof results in a consistent, sustained, and dose-dependent reduction in the number of peripheral blood lymphocytes. COMPOUND 1 has been described to be useful in the treatment and/or prevention of diseases or disorders associated with an activated immune system (see e.g. WO 2005/054215 and WO 2009/115954). In particular COMPOUND 1 (INN: ponesimod) has shown clinical benefit in phase II trials in patients with moderate to severe chronic plaque psoriasis and in patients with relapsing-remitting multiple sclerosis. COMPOUND 1 may be prepared according to any procedure as disclosed in WO 2005/054215, WO 2008/062376 and WO 2014/027330.

(6) Dimethyl fumarate (also termed “DMF” or “BG-12”) has been described in WO 00/030622 to be useful for the treatment of autoimmune diseases. In particular dimethyl fumarate (Tecfidera®) has been approved for the treatment of relapsing forms of multiple sclerosis, including relapsing-remitting multiple sclerosis, which is the most common form of the disease. Dimethyl fumarate can be prepared according to procedures known in the art for example as described in EP 0312697 A2.

(7) Methyl fumarate (also termed “monomethyl fumarate” or “MMF”) has been shown to be a pharmacologically active metabolite of dimethyl fumarate. Methyl fumarate can be prepared according to procedures known in the art for example as described in EP 0312697 A2.

(8) (N,N-Diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate (also termed “XP23829”) is a prodrug that is rapidly converted to monomethyl fumarate. XP23829 is currently in clinical development for the treatment of moderate-to-severe chronic plaque-type psoriasis and for the treatment of relapsing forms of multiple sclerosis. (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate and the preparation thereof is described in WO 2010/022177. 2-(2,5-Dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate (also termed “ALKS 8700”) is a prodrug that rapidly converts to monomethyl fumarate. ALKS 8700 is currently in clinical development for the treatment of multiple sclerosis. 2-(2,5-Dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate and the preparation thereof is described in WO 2014/152494.

(9) 2) A further embodiment of the invention relates to a pharmaceutical combination according to embodiment 1), wherein the first active ingredient is (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one and the second active ingredient is methyl fumarate or a pharmaceutically acceptable salt thereof.

(10) 3) A further embodiment of the invention relates to a pharmaceutical combination according to embodiment 1), wherein the first active ingredient is (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one and the second active ingredient is dimethyl fumarate.

(11) 4) A further embodiment of the invention relates to a pharmaceutical combination according to embodiment 1), wherein the first active ingredient is (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one and the second active ingredient is (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate.

(12) 5) A further embodiment of the invention relates to a pharmaceutical combination according to embodiment 1), wherein the first active ingredient is (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one and the second active ingredient is

2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate.

(13) 6) A further embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 5), wherein the first and the second active ingredient are comprised in a single pharmaceutical composition.

(14) In a special case of embodiment 6) where e.g. the first active ingredient is administered once daily and the second ingredient is administered twice daily, only one of the two pharmaceutical compositions needed per day will contain both the first and the second active ingredient whereas the other will only contain the second active ingredient.

(15) Moreover, in case of a pharmaceutical combination according to embodiment 6) where the first and/or the second active ingredient is administered according to a dose up-titration regimen (see for example the up-titration regimen disclosed in WO 2009/115954 for COMPOUND 1) the pharmaceutical compositions needed for the dose up-titration will contain the amounts of active ingredient required for the different steps of the dose up-titration regimen.

(16) 7) A further embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 5), wherein the first and the second active ingredient are comprised in separated pharmaceutical compositions.

(17) In case the first and the second active ingredient are comprised in separated pharmaceutical compositions, they can be administered simultaneously, sequentially or separately; preferably the separated pharmaceutical compositions are administered simultaneously or sequentially, especially sequentially. In case the first active ingredient is for example administered once daily and the second active ingredient twice daily, then the separated pharmaceutical compositions are preferably administered one time per day simultaneously or sequentially, especially sequentially. If administered sequentially or separately, the separated pharmaceutical compositions may be administered in one or the other order. The number of administrations per day may be the same or different for the separated pharmaceutical compositions. For instance, one pharmaceutical composition may be administered twice daily and the other pharmaceutical composition may be administered once or twice daily. Preferably the pharmaceutical composition comprising COMPOUND 1 is administered once daily and the pharmaceutical composition comprising the second active ingredient is administered twice daily. Further, the separated pharmaceutical compositions may be administered by the same or different routes of administration, preferably by the same route of administration. Most preferably the separated pharmaceutical compositions are administered orally.

(18) 8) A further embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 7) for use as a medicament.

(19) 9) A further embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 7) for use in the prevention and/or treatment of a disease or disorder associated with an activated immune system.

(20) 10) A further embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 7) for use in the prevention and/or treatment of a disease or disorder selected from the group consisting of rejection of transplanted organs such as kidney, liver, heart, lung, pancreas, cornea, and skin; graft-versus-host disease; autoimmune syndromes including rheumatoid arthritis, multiple sclerosis, inflammatory bowel diseases such as Crohn's disease and ulcerative colitis, psoriasis, psoriatic arthritis, thyroiditis such as Hashimoto's thyroiditis, and uveo-retinitis; atopic diseases such as rhinitis, conjunctivitis, and dermatitis; asthma; type I diabetes; post-infectious autoimmune diseases including rheumatic fever and post-infectious glomerulonephritis; solid cancers; and tumor metastasis.

(21) 11) A further embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 7) for use in the prevention and/or treatment of a disease or disorder selected from the group consisting of rejection of transplanted organs selected from kidney, liver, heart and lung; graft-versus-host disease; autoimmune syndromes selected from

rheumatoid arthritis, multiple sclerosis, psoriasis, psoriatic arthritis, Crohn's disease, and Hashimoto's thyroiditis; and atopic dermatitis.

(22) 12) A preferred embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 7) for use in the prevention and/or treatment of graft-versus-host disease.

(23) 13) A most preferred embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 7) for use in the prevention and/or treatment of multiple sclerosis.

(24) 14) Another very preferred embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 7) for use in the prevention and/or treatment of relapsing multiple sclerosis.

(25) 15) Another very preferred embodiment of the invention relates to a pharmaceutical combination according to any one of embodiments 1) to 7) for use in the prevention and/or treatment of relapsing-remitting multiple sclerosis.

(26) The present invention also relates to a method for the prevention or treatment of a disease or disorder listed in any one of embodiments 9) to 15) comprising administering to a subject (preferably a human subject) in need thereof a pharmaceutically active amount of a pharmaceutical combination according to any one of embodiments 1) to 7).

(27) 16) A further embodiment of the invention relates to a pharmaceutical composition containing, as active principle, (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one and at least one therapeutically inert excipient, wherein the pharmaceutical composition is to be administered in combination with a second pharmaceutical composition containing, as active principle, methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, or 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate, or a pharmaceutically acceptable salt thereof and at least one therapeutically inert excipient.

(28) 17) A further embodiment of the invention relates to a pharmaceutical composition containing, as active principle, (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one and at least one therapeutically inert excipient, wherein the pharmaceutical composition is to be administered in combination with a second pharmaceutical composition containing, as active principle, dimethyl fumarate and at least one therapeutically inert excipient.

(29) 18) A further embodiment of the invention relates to a pharmaceutical composition according to embodiment 16) or 17) for use as a medicament.

(30) 19) A further embodiment of the invention relates to a pharmaceutical composition according to embodiment 16) or 17) for use in the prevention and/or treatment of a disease or disorder listed in any one of embodiments 9) to 15).

(31) 20) A further embodiment of the invention relates to a pharmaceutical composition containing, as active principle, methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, or 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate, or a pharmaceutically acceptable salt thereof, and at least one therapeutically inert excipient, wherein the pharmaceutical composition is to be administered in combination with a second pharmaceutical composition containing, as active principle, (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one and at least one therapeutically inert excipient.

(32) 21) A further embodiment of the invention relates to a pharmaceutical composition containing, as active principle, dimethyl fumarate and at least one therapeutically inert excipient, wherein the pharmaceutical composition is to be administered in combination with a second pharmaceutical composition containing, as active principle, (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one and at least one therapeutically

inert excipient.

(33) 22) A further embodiment of the invention relates to a pharmaceutical composition according to embodiment 20) or 21) for use as a medicament.

(34) 23) A further embodiment of the invention relates to a pharmaceutical composition according to embodiment 20) or 21) for use in the prevention and/or treatment of a disease or disorder listed in any one of embodiments 9) to 15).

(35) 24) A further embodiment of the invention relates to a kit of parts comprising a first pharmaceutical composition containing, as active principle, (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one and at least one therapeutically inert excipient; and a second pharmaceutical composition containing, as active principle, methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, or 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate, or a pharmaceutically acceptable salt thereof, and at least one therapeutically inert excipient.

(36) 25) A further embodiment of the invention relates to a kit of parts according to embodiment 24), wherein the second pharmaceutical composition contains, as active principle, dimethyl fumarate and at least one therapeutically inert excipient.

(37) 26) A further embodiment of the invention relates to a kit of parts according to embodiment 24) or 25) further comprising instructions for the simultaneous, sequential or separate administration of the pharmaceutical compositions.

(38) 27) A further embodiment of the invention relates to a kit of parts according to any one of embodiments 24) to 26) for use as a medicament.

(39) 28) A further embodiment of the invention relates to a kit of parts according to any one of embodiments 24) to 26) for use in the prevention and/or treatment of a disease or disorder listed in any one of embodiments 9) to 15).

(40) 29) A further embodiment of the invention relates to the use of (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one and a second active ingredient which is selected from the group consisting of methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, and 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for use in the prevention and/or treatment of a disease or disorder listed in any one of embodiments 9) to 15).

(41) 30) A further embodiment of the invention relates to the use according to embodiment 29), wherein the second active ingredient is dimethyl fumarate.

(42) 31) A further embodiment of the invention relates to the use of (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one for the manufacture of a medicament for use, in combination with a second medicament comprising methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, or 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate, or a pharmaceutically acceptable salt thereof, in the prevention and/or treatment of a disease or disorder listed in any one of embodiments 9) to 15).

(43) 32) A further embodiment of the invention relates to the use according to embodiment 31), wherein the second medicament comprises dimethyl fumarate.

(44) 33) A further embodiment of the invention relates to the use of methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, or 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate, or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for use, in combination with a second medicament comprising (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one, in the prevention and/or treatment of a disease or disorder listed in any one of embodiments 9) to 15).

(45) 34) A further embodiment of the invention relates to the use of dimethyl fumarate for the

manufacture of a medicament for use, in combination with a second medicament comprising (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-thiazolidin-4-one, in the prevention and/or treatment of a disease or disorder listed in any one of embodiments 9) to 15).

(46) Based on the dependencies of the different embodiments 1) to 34) as disclosed hereinabove, the following embodiments are thus possible and intended and herewith specifically disclosed in individualised form: 1, 2+1, 3+1, 4+1, 5+1, 6+1, 6+2+1, 6+3+1, 6+4+1, 6+5+1, 7+1, 7+2+1, 7+3+1, 7+4+1, 7+5+1, 8+1, 8+2+1, 8+3+1, 8+4+1, 8+5+1, 8+6+1, 8+6+2+1, 8+6+3+1, 8+6+4+1, 8+6+5+1, 8+7+1, 8+7+2+1, 8+7+3+1, 8+7+4+1, 8+7+5+1, 9+1, 9+2+1, 9+3+1, 9+4+1, 9+5+1, 9+6+1, 9+6+2+1, 9+6+3+1, 9+6+4+1, 9+6+5+1, 9+7+1, 9+7+2+1, 9+7+3+1, 9+7+4+1, 9+7+5+1, 10+1, 10+2+1, 10+3+1, 10+4+1, 10+5+1, 10+6+1, 10+6+2+1, 10+6+3+1, 10+6+4+1, 10+6+5+1, 10+7+1, 10+7+2+1, 10+7+3+1, 10+7+4+1, 10+7+5+1, 11+1, 11+2+1, 11+3+1, 11+4+1, 11+5+1, 11+6+1, 11+6+2+1, 11+6+3+1, 11+6+4+1, 11+6+5+1, 11+7+1, 11+7+2+1, 11+7+3+1, 11+7+4+1, 11+7+5+1, 12+1, 12+2+1, 12+3+1, 12+4+1, 12+5+1, 12+6+1, 12+6+2+1, 12+6+3+1, 12+6+4+1, 12+6+5+1, 12+7+1, 12+7+2+1, 12+7+3+1, 12+7+4+1, 12+7+5+1, 13+1, 13+2+1, 13+3+1, 13+4+1, 13+5+1, 13+6+1, 13+6+2+1, 13+6+3+1, 13+6+4+1, 13+6+5+1, 13+7+1, 13+7+2+1, 13+7+3+1, 13+7+4+1, 13+7+5+1, 14+1, 14+2+1, 14+3+1, 14+4+1, 14+5+1, 14+6+1, 14+6+2+1, 14+6+3+1, 14+6+4+1, 14+6+5+1, 14+7+1, 14+7+2+1, 14+7+3+1, 14+7+4+1, 14+7+5+1, 15+1, 15+2+1, 15+3+1, 15+4+1, 15+5+1, 15+6+1, 15+6+2+1, 15+6+3+1, 15+6+4+1, 15+6+5+1, 15+7+1, 15+7+2+1, 15+7+3+1, 15+7+4+1, 15+7+5+1, 16, 17, 18+16, 18+17, 19+16, 19+17, 20, 21, 22+20, 22+21, 23+20, 23+21, 24, 25+24, 26+24, 26+25+24, 27+24, 27+25+24, 27+26+24, 27+26+25+24, 28+24, 28+25+24, 28+26+24, 28+26+25+24, 29, 30+29, 31, 32+31, 33, and 34.

(47) In the list above the numbers refer to the embodiments according to their numbering provided hereinabove whereas “+” indicates the dependency from another embodiment. The different individualised embodiments are separated by commas. In other words, “6+4+1” for example refers to embodiment 6) depending on embodiment 4), depending on embodiment 1), i.e. embodiment “6+4+1” corresponds to the pharmaceutical combination of embodiment 1) further limited by the features of the embodiments 4) and 6).

(48) Definitions provided herein are intended to apply uniformly to the subject matter as defined in any one of embodiments 1) to 34), and, mutatis mutandis, throughout the description and the claims unless an otherwise expressly set out definition provides a broader or narrower definition. It is well understood that a definition or preferred definition of a term or expression defines and may replace the respective term or expression independently of (and in combination with) any definition or preferred definition of any or all other terms or expressions as defined herein.

(49) Any reference to an active ingredient as defined in any one of embodiments 1) to 34) is to be understood as referring also to the pharmaceutically acceptable salts of such active ingredient, as appropriate and expedient.

(50) The term “pharmaceutically acceptable salts” refers to salts that retain the desired biological activity of the subject compound and exhibit minimal undesired toxicological effects. Such salts include inorganic or organic acid and/or base addition salts depending on the presence of basic and/or acidic groups in the subject compound. For reference see for example ‘Handbook of Pharmaceutical Salts. Properties, Selection and Use.’, P. Heinrich Stahl, Camille G. Wermuth (Eds.), Wiley-VCH, 2008 and ‘Pharmaceutical Salts and Co-crystals’, Johan Wouters and Luc Quéré (Eds.), RSC Publishing, 2012.

(51) The term “pharmaceutical combination”, as used herein, refers to a combination of two or more, preferably two, active ingredients, wherein the active ingredients are comprised in a single pharmaceutical composition or in separated pharmaceutical compositions.

(52) The term “active ingredient”, as used herein, refers to the pharmaceutically active component of a pharmaceutical composition. Examples of active ingredients, as used herein, are in a first group (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-o-tolyl-

thiazolidin-4-one (COMPOUND 1) and in a second group methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, and 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate, or a pharmaceutically acceptable salt thereof (preferably dimethyl fumarate).

(53) The term “simultaneous” or “simultaneously”, when used in relation to the administration of active ingredients or of pharmaceutical compositions, means that the administration of a first active ingredient (or of a first pharmaceutical composition, respectively) is still ongoing when the administration of a second active ingredient (or of a second pharmaceutical composition, respectively) is started. Especially, the term “simultaneous” or “simultaneously” means that two active ingredients (or two pharmaceutical compositions, respectively) are administered at the same time, i.e. with the same starting and end time, as is for instance the case for the administration of two active ingredients comprised in a single pharmaceutical composition.

(54) The term “sequential” or “sequentially”, when used in relation to the administration of active ingredients or of pharmaceutical compositions, means that the administration of a second active ingredient (or of a second pharmaceutical composition, respectively) is started less than one hour after the administration of a first active ingredient (or of a first pharmaceutical composition, respectively) has been finalized.

(55) The term “separate” or “separately”, when used in relation to the administration of active ingredients or of pharmaceutical compositions, means that the administration of a second active ingredient (or of a second pharmaceutical composition, respectively) is started one hour or more (and up to about twelve hours, up to about 24 hours, or up to about 7 days) after the administration of a first active ingredient (or of a first pharmaceutical composition, respectively) has been finalized.

(56) The expressions “to be administered in combination” or “for use, in combination” mean simultaneous, sequential or separate, preferably sequential, administration of active ingredients or pharmaceutical compositions.

(57) The term “route of administration”, as used herein, refers to the path by which an active ingredient (e.g. in form of a pharmaceutical composition in a particular dosage form) enters the body. The active ingredients may be administered by enteral (especially oral) or parenteral administration (including topical application or inhalation). Examples of dosage forms which may be used for the administration of the active ingredients are tablets, capsules, pills, powders, solutions, suspensions, emulsions, injectable aqueous or oily solutions or suspensions, suppositories, creams, gels, ear or eye drops, nasal spray, skin patches, or aerosols. Dosage forms for oral administration, such as tablets, capsules, pills, solutions or suspensions are preferred. In case the two active ingredients are comprised in separated pharmaceutical compositions, said separated pharmaceutical compositions may be administered by the same or different routes of administration using the same or different dosage forms.

(58) For the sake of clarity, relapsing multiple sclerosis means relapsing forms of multiple sclerosis which includes forms of multiple sclerosis with relapses. Examples of relapsing multiple sclerosis are relapsing-remitting multiple sclerosis, secondary progressive multiple sclerosis with relapses, and progressive relapsing multiple sclerosis.

(59) The production of pharmaceutical compositions can be effected in a manner which will be familiar to any person skilled in the art (see for example Remington, The Science and Practice of Pharmacy, 21st Edition (2005), Part 5, “Pharmaceutical Manufacturing” [published by Lippincott Williams & Wilkins]) by bringing COMPOUND 1 and/or a fumarate derivative which is selected from the group consisting of methyl fumarate, dimethyl fumarate, (N,N-diethylcarbamoyl)methyl methyl (2E)but-2-ene-1,4-dioate, and 2-(2,5-dioxopyrrolidin-1-yl)ethyl methyl (2E)but-2-ene-1,4-dioate, or a pharmaceutically acceptable salt thereof, optionally in combination with other therapeutically valuable substances, into a galenical administration form together with suitable, non-toxic, inert, therapeutically compatible solid or liquid carrier materials and, if desired, usual

pharmaceutical adjuvants. Formulations for dimethyl fumarate are described in e.g. WO 00/030622 and WO 2010/079222.

(60) The optimal dosing regimen (i.e., the magnitude of the dose and the dosing frequency) for each of the two active ingredients of the pharmaceutical combination according to the present invention may vary depending upon the route of administration, the dosage form, the disease or disorder to be treated, and the particular second active ingredient (fumarate derivative) applied. Further, the dose and/or the dosing frequency may be different during the initial phase and the later phase of the treatment for the first and/or the second active ingredient of the pharmaceutical combination. A preferred dosing regimen for COMPOUND 1 has been disclosed in WO 2009/115954. A preferred maintenance dose for COMPOUND 1 is 10 mg or 20 mg orally once daily, such as especially 20 mg orally once daily. A preferred dosing regimen for dimethyl fumarate is disclosed in WO 2008/097596 (see also U.S. Pat. No. 8,399,514). Most preferably dimethyl fumarate is administered at a starting dose of 120 mg twice a day, orally, for 7 days and a maintenance dose after 7 days of 240 mg twice a day, orally, especially in case of an immediate release formulation such as the one used for Tecfidera®. A preferred dosing regimen for a controlled release pharmaceutical composition of monomethyl fumarate or dimethyl fumarate is disclosed in EP 2316430. A preferred dose for XP23829 is 800 mg orally once daily or 400 mg orally twice daily. A preferred dose for ALKS 8700 is 420 mg orally twice daily.

(61) The present invention also includes isotopically labelled, especially .sup.2H (deuterium) labelled active ingredients, which active ingredients are identical to the active ingredients as defined in embodiment 1) except that one or more atoms have each been replaced by an atom having the same atomic number but an atomic mass different from the atomic mass usually found in nature. Isotopically labelled, especially .sup.2H (deuterium) labelled active ingredients and pharmaceutically acceptable salts thereof are within the scope of the present invention. Substitution of hydrogen with the heavier isotope .sup.2H (deuterium) may lead to greater metabolic stability, resulting e.g. in increased in-vivo half-life or reduced dosage requirements, or may lead to reduced inhibition of cytochrome P450 enzymes, resulting e.g. in an improved safety profile. In one embodiment only one of the two active ingredients of the pharmaceutical combination is isotopically labelled. In a preferred embodiment of the invention, the active ingredients are not isotopically labelled, or one active ingredient is not isotopically labelled and the other active ingredient is labelled only with one or more deuterium atoms, or both active ingredients are each labelled only with one or more deuterium atoms. In a most preferred embodiment, the active ingredients are not isotopically labelled at all. Isotopically labelled active ingredients may be prepared in analogy to the methods described for the not isotopically labelled active ingredients, but using the appropriate isotopic variation of suitable reagents or starting materials.

(62) The term “about” placed before a numerical value “X” refers in the current application to an interval extending from X minus 10% of X to X plus 10% of X.

(63) Biological Assay:

(64) The efficacy of dimethyl fumarate (DMF) in acute monophasic models of experimental autoimmune encephalomyelitis (EAE) can be determined in a pilot experiment.

(65) Female Lewis rats are immunized with an emulsion of myelin basic protein (MBP) from Guinea pig in complete Freund's adjuvant. A total of 200 micrograms of MBP is injected per rat at two sites subcutaneously (right paw and base of tail). Within 10 days the rats develop signs of paralysis which will be graded on a scale from 0 to 15 assessing the tail and each limb individually according to the method originally described by Weaver A. et al., *FASEB J.*, 2005, 19(12): 1668-1670. The disease takes an acute monophasic course and is self-remitting. Usually by day 21 clinical scores reach values below 3.0.

(66) Groups of 10 to 14 rats are dosed starting at the day of disease induction (day 0) with different doses of DMF in the range between 40 and 160 mg/kg once daily (q.d.). Clinical scores are assessed on a daily basis and disease development is compared between vehicle-treated rats and

rats receiving DMF (vehicle: 0.5% Methylcellulose/0.5% Tween® 80). In parallel, body weights of the rats are monitored on a daily basis to follow general health. The experiment is terminated at day 21 whereupon hematology can be measured.

(67) Plasma samples can be taken for compound concentration determination and for the measurement of protein markers. Different organs may be isolated and fixed for histopathological examination, particularly the spinal cord and the brain. Spinal cord tissue preparations can be used to assess the degree of demyelination, neuronal loss and inflammatory cell infiltration by different staining methods (H&E, Luxol Fast Blue for myelin, MBP, NeuN, CD3, Nrf2 and other markers by immunohistochemistry for demyelination, neuronal loss, T cell infiltration, antioxidant pathway activation and other pathways, respectively). Additional tissue samples may be preserved in RNAlater® solution for analysis of gene expression.

(68) In parallel to the measurement of efficacy of DMF in the EAE model, a second pilot experiment is required. Healthy female Lewis rats will receive ponesimod at different doses in the range from 0.3 to 100 mg/kg. The goal is to assess the dose-effect relationship of ponesimod on lymphocyte count in peripheral blood in this strain of rat.

(69) Based on the first pilot experiment described above one dose of DMF is selected for the combination efficacy experiment. The dose selected exerts partial efficacy between none and strongest efficacy on EAE scores (dose 120 mg/kg).

(70) Based on the second pilot experiment using ponesimod, the dose selected shows full efficacy on lymphocyte count reduction (dose 100 mg/kg).

(71) The combination efficacy experiment is performed in the same rat EAE model as described for the first pilot experiment. It consists of four treatment groups: 1. Vehicle (0.5% Methylcellulose/0.5% Tween® 80) q.d., from day 0 2. DMF (120 mg/kg) q.d., from day 0 3. Ponesimod (100 mg/kg) q.d., from day 0 4. DMF (120 mg/kg) q.d., from day 0+ponesimod (100 mg/kg) q.d., from day 0

(72) In life and termination endpoints are the same as described for the first pilot experiment.

(73) The goal of this experiment is to show that the addition of a dose of ponesimod, that is fully efficacious on lymphocyte count, shows added benefit to a partially efficacious dose of DMF in a model of EAE.

(74) The results from the combination efficacy experiment are shown in FIG. 1. As can be seen from FIG. 1, both compounds, ponesimod and DMF exhibit efficacy on the clinical course of EAE in the rat monophasic EAE model. Ponesimod showed a marked and statistically significant effect while DMF alone showed a moderate effect during the ascending phase of the disease. When combined, the two compounds showed synergistic efficacy from day 16 onwards, where DMF alone had no effect on clinical score anymore.

Claims

1. A pharmaceutical combination comprising a first active ingredient which is about 10 mg of (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-O-tolyl-thiazolidin-4 one or a pharmaceutically acceptable salt thereof and a second active ingredient which is dimethyl fumarate or a pharmaceutically acceptable salt thereof.

2. A pharmaceutical combination according to claim 1, wherein the first and the second active ingredient are comprised in a single pharmaceutical composition.

3. A pharmaceutical combination according to claim 1, wherein the first and the second active ingredient are comprised in separated pharmaceutical compositions.

4. A medicament comprising the pharmaceutical combination according to claim 1.

5. A pharmaceutical combination comprising a first active ingredient which is about 20 mg of (R)-5-[3-chloro-4-(2,3-dihydroxy-propoxy)-benz[Z]ylidene]-2-([Z]-propylimino)-3-O-tolyl-thiazolidin-4 one or a pharmaceutically acceptable salt thereof and a second active ingredient

which is dimethyl fumarate or a pharmaceutically acceptable salt thereof.

6. A pharmaceutical combination according to claim 5, wherein the first and the second active ingredient are comprised in a single pharmaceutical composition.

7. A pharmaceutical combination according to claim 5, wherein the first and the second active ingredient are comprised in separated pharmaceutical compositions.

8. A medicament comprising the pharmaceutical combination according to claim 5.
