

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12409211
Kind Code	B2
Date of Patent	September 09, 2025
Inventor(s)	Abiad; Maurice et al.

Non-protein Clostridial toxin compositions

Abstract

Pharmaceutical compositions that stabilize a Clostridial toxin active ingredient are described. The compositions can be liquid or solid compositions, and comprise a surfactant and an antioxidant. In some embodiments, the compositions comprise a surfactant selected from a poloxamer and a polysorbate; an antioxidant selected from methionine, N-acetyl cysteine, ethylenediaminetetraacetic acid and combinations thereof, and, optionally, a tonicity agent and/or a lyoprotector selected from, for example, trehalose, sucrose.

Inventors: Abiad; Maurice (Anaheim, CA), Dani; Bhas (San Diego, CA), ShalaeV; Evgenyi (Dana Point, CA)

Applicant: Allergan, Inc. (Irvine, CA)

Family ID: 59974865

Assignee: Allergan, Inc. (Irvine, CA)

Appl. No.: 18/629715

Filed: April 08, 2024

Prior Publication Data

Document Identifier	Publication Date
US 20240252604 A1	Aug. 01, 2024

Related U.S. Application Data

continuation parent-doc US 16531800 20190805 US 10973890 20210413 child-doc US 17185312
division parent-doc US 18136048 20230418 US 12144847 child-doc US 18629715
division parent-doc US 17185312 20210225 ABANDONED child-doc US 18136048
division parent-doc US 15703527 20170913 US 10369190 20190806 child-doc US 16531800
us-provisional-application US 62394009 20160913

Publication Classification

Int. Cl.: **A61K38/48** (20060101); **A61K9/00** (20060101); **A61K9/19** (20060101); **A61K38/16** (20060101); **A61K47/10** (20170101); **A61K47/18** (20170101); **A61K47/26** (20060101); **A61P9/06** (20060101); **A61P13/00** (20060101); **A61P21/00** (20060101); **A61P25/24** (20060101); **A61P29/00** (20060101)

U.S. Cl.:

CPC **A61K38/4893** (20130101); **A61K9/0019** (20130101); **A61K9/19** (20130101); **A61K38/164** (20130101); **A61K38/48** (20130101); **A61K47/10** (20130101); **A61K47/183** (20130101); **A61K47/26** (20130101); **A61P9/06** (20180101); **A61P13/00** (20180101); **A61P21/00** (20180101); **A61P25/24** (20180101); **A61P29/00** (20180101);

Field of Classification Search

CPC: A61K (38/4893)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
4891319	12/1989	Roser	N/A	N/A
5472706	12/1994	Friedman et al.	N/A	N/A
5512547	12/1995	Johnson et al.	N/A	N/A
5556771	12/1995	Shen et al.	N/A	N/A
5616554	12/1996	Beardsley	N/A	N/A
5696077	12/1996	Johnson et al.	N/A	N/A
5714468	12/1997	Binder	N/A	N/A
5756468	12/1997	Johnson et al.	N/A	N/A
5945098	12/1998	Sarno et al.	N/A	N/A
5962637	12/1998	Shone et al.	N/A	N/A
5989545	12/1998	Foster et al.	N/A	N/A
6043042	12/1999	Shone et al.	N/A	N/A
6087327	12/1999	Pearce et al.	N/A	N/A
6203794	12/2000	Dolly et al.	N/A	N/A
6306423	12/2000	Donovan et al.	N/A	N/A
6337386	12/2001	Shone et al.	N/A	N/A
6461617	12/2001	Shone et al.	N/A	N/A
6506399	12/2002	Donovan	N/A	N/A
6555558	12/2002	Chen et al.	N/A	N/A
6599504	12/2002	Wadstrom et al.	N/A	N/A
6632440	12/2002	Quinn et al.	N/A	N/A
6653062	12/2002	DePablo et al.	N/A	N/A
6777196	12/2003	Klein et al.	N/A	N/A
6843998	12/2004	Steward et al.	N/A	N/A
6977080	12/2004	Donovan	N/A	N/A
7001602	12/2005	Schmidt	N/A	N/A
7132259	12/2005	Dolly et al.	N/A	N/A

7160699	12/2006	Wang et al.	N/A	N/A
7183066	12/2006	Fernandez-Salas et al.	N/A	N/A
7205018	12/2006	Sherwood et al.	N/A	N/A
7208166	12/2006	Marchini et al.	N/A	N/A
7208285	12/2006	Steward et al.	N/A	N/A
7211261	12/2006	Moyer et al.	N/A	N/A
7223577	12/2006	Steward et al.	N/A	N/A
7255865	12/2006	Walker	N/A	N/A
7273722	12/2006	Lin et al.	N/A	N/A
7309602	12/2006	David	N/A	N/A
7332567	12/2007	Steward et al.	N/A	N/A
7341843	12/2007	Atassi	N/A	N/A
7354740	12/2007	Xiang et al.	N/A	N/A
7375079	12/2007	Gordon et al.	N/A	N/A
7399607	12/2007	Williams et al.	N/A	N/A
7419676	12/2007	Dolly et al.	N/A	N/A
7442686	12/2007	Lasko et al.	N/A	N/A
7473559	12/2008	Lee	N/A	N/A
7491403	12/2008	Borodic	N/A	N/A
7531193	12/2008	Demarne et al.	N/A	N/A
7560100	12/2008	Pinchasi et al.	N/A	N/A
7579010	12/2008	Hunt	N/A	N/A
7579171	12/2008	Porro et al.	N/A	N/A
7591806	12/2008	Xu	N/A	N/A
7598027	12/2008	Fernandez-Salas et al.	N/A	N/A
7601740	12/2008	Weiner et al.	N/A	N/A
7604802	12/2008	O'Hagan et al.	N/A	N/A
7642079	12/2009	Cayouette et al.	N/A	N/A
7645570	12/2009	Fernandez-Salas et al.	N/A	N/A
7674454	12/2009	Huizinga et al.	N/A	N/A
7727537	12/2009	Modi	N/A	N/A
7758872	12/2009	Finzi	N/A	N/A
7776844	12/2009	Yu et al.	N/A	N/A
7780967	12/2009	Hunt	N/A	N/A
7829525	12/2009	Frevert	N/A	N/A
7838011	12/2009	Modi	N/A	N/A
7846722	12/2009	Williams et al.	N/A	N/A
7867480	12/2010	Cevc et al.	N/A	N/A
7875436	12/2010	Fournie-Zaluski et al.	N/A	N/A
7879341	12/2010	Taylor	N/A	N/A
7906136	12/2010	Wong et al.	N/A	N/A
7910554	12/2010	McKerracher	N/A	N/A
7927622	12/2010	Cevc et al.	N/A	N/A
7998489	12/2010	Steward et al.	N/A	N/A
8057807	12/2010	Schmidt	N/A	N/A
8057808	12/2010	Ackerman	N/A	N/A
8067200	12/2010	Foster et al.	N/A	N/A
8124357	12/2011	Fernandez-Salas et al.	N/A	N/A
8137677	12/2011	Hunt	N/A	N/A
8143212	12/2011	Stogniew et al.	N/A	N/A

8158360	12/2011	Heise et al.	N/A	N/A
8168206	12/2011	Hunt	N/A	N/A
8173138	12/2011	Moyer et al.	N/A	N/A
8192979	12/2011	Borodic et al.	N/A	N/A
8241640	12/2011	Borodic	N/A	N/A
8257699	12/2011	Bookbinder et al.	N/A	N/A
8277822	12/2011	Dott et al.	N/A	N/A
8283133	12/2011	Franciskovich et al.	N/A	N/A
8309077	12/2011	Murthy et al.	N/A	N/A
8323666	12/2011	Hunt	N/A	N/A
8343508	12/2012	Pomato et al.	N/A	N/A
8357541	12/2012	Ton et al.	N/A	N/A
8372645	12/2012	Taylor	N/A	N/A
8398997	12/2012	Dake et al.	N/A	N/A
8398998	12/2012	Bigalke et al.	N/A	N/A
8399006	12/2012	de Juan, Jr. et al.	N/A	N/A
8404249	12/2012	Dake et al.	N/A	N/A
8420081	12/2012	Fraunhofer et al.	N/A	N/A
8444991	12/2012	Randolph et al.	N/A	N/A
8470330	12/2012	Maddon et al.	N/A	N/A
8476024	12/2012	Mahrhold et al.	N/A	N/A
8486467	12/2012	Prescott	N/A	N/A
8501195	12/2012	Turkel et al.	N/A	N/A
8512679	12/2012	Hyde et al.	N/A	N/A
8512979	12/2012	Dunson, Jr. et al.	N/A	N/A
8529939	12/2012	Masters et al.	N/A	N/A
8580250	12/2012	Hunt	N/A	N/A
8580252	12/2012	Bookbinder et al.	N/A	N/A
8586020	12/2012	Song et al.	N/A	N/A
8591884	12/2012	Soparkar	N/A	N/A
8609112	12/2012	Blumenfeld et al.	N/A	N/A
8617568	12/2012	Jung et al.	N/A	N/A
8618261	12/2012	Ester et al.	N/A	N/A
8623419	12/2013	Malakhov et al.	N/A	N/A
8632785	12/2013	Hunt	N/A	N/A
8642047	12/2013	Hunt	N/A	N/A
8647638	12/2013	Clarke et al.	N/A	N/A
8669091	12/2013	Gentshev et al.	N/A	N/A
8685381	12/2013	Schlessinger et al.	N/A	N/A
8685684	12/2013	Lee et al.	N/A	N/A
8697155	12/2013	Levy et al.	N/A	N/A
8697644	12/2013	Prestrelski et al.	N/A	N/A
8703755	12/2013	Adams et al.	N/A	N/A
8715706	12/2013	Barak	N/A	N/A
8716240	12/2013	DeFrees et al.	N/A	N/A
8729241	12/2013	Liu et al.	N/A	N/A
8734826	12/2013	Barak	N/A	N/A
8796217	12/2013	Amari et al.	N/A	N/A
8808710	12/2013	Randolph et al.	N/A	N/A
8821879	12/2013	Babuka et al.	N/A	N/A

8828432	12/2013	van Lengerich	N/A	N/A
8840905	12/2013	Schmidt	N/A	N/A
8841110	12/2013	Xiang et al.	N/A	N/A
8883146	12/2013	Fraunhofer et al.	N/A	N/A
8889151	12/2013	Turkel et al.	N/A	N/A
8889826	12/2013	Leese et al.	N/A	N/A
8920795	12/2013	Jung et al.	N/A	N/A
8936790	12/2014	Turkel et al.	N/A	N/A
8940308	12/2014	Turkel et al.	N/A	N/A
8968747	12/2014	Turkel et al.	N/A	N/A
8993268	12/2014	Jung et al.	N/A	N/A
9006388	12/2014	Paul et al.	N/A	N/A
9012177	12/2014	Glass et al.	N/A	N/A
9017726	12/2014	Song et al.	N/A	N/A
9040074	12/2014	Holzer et al.	N/A	N/A
9044477	12/2014	Blanda et al.	N/A	N/A
9050336	12/2014	Blanda et al.	N/A	N/A
9050367	12/2014	Taylor	N/A	N/A
9061025	12/2014	Burstein et al.	N/A	N/A
9066943	12/2014	Schmidt	N/A	N/A
9072688	12/2014	Boyden et al.	N/A	N/A
9078892	12/2014	Turkel et al.	N/A	N/A
9078893	12/2014	Turkel et al.	N/A	N/A
9085619	12/2014	Fraunhofer et al.	N/A	N/A
9107815	12/2014	Hunt	N/A	N/A
9109244	12/2014	Pompejus et al.	N/A	N/A
9125804	12/2014	Webb et al.	N/A	N/A
9157875	12/2014	Warner et al.	N/A	N/A
9161970	12/2014	Tezel et al.	N/A	N/A
9173944	12/2014	Taylor et al.	N/A	N/A
9180081	12/2014	Dake et al.	N/A	N/A
9198856	12/2014	Burger et al.	N/A	N/A
9198945	12/2014	Kim et al.	N/A	N/A
9198958	12/2014	Jung et al.	N/A	N/A
9211248	12/2014	Dake et al.	N/A	N/A
9211315	12/2014	Bookbinder et al.	N/A	N/A
9220780	12/2014	Jung et al.	N/A	N/A
9220783	12/2014	Taylor	N/A	N/A
9226959	12/2015	Kramps et al.	N/A	N/A
9233080	12/2015	Golubovic-Liakopoulos et al.	N/A	N/A
9233238	12/2015	Buysman et al.	N/A	N/A
9248168	12/2015	Blumenfeld	N/A	N/A
9265722	12/2015	Donovan	N/A	N/A
9278140	12/2015	Hunt	N/A	N/A
9283217	12/2015	Dibas et al.	N/A	N/A
9284566	12/2015	Liao et al.	N/A	N/A
9284579	12/2015	Green et al.	N/A	N/A
9302008	12/2015	Hunt	N/A	N/A
9327105	12/2015	Ramdas et al.	N/A	N/A

9340587	12/2015	Thompson et al.	N/A	N/A
9389162	12/2015	Chen et al.	N/A	N/A
9394234	12/2015	Chen et al.	N/A	N/A
9415006	12/2015	Baker, Jr. et al.	N/A	N/A
9447401	12/2015	Wei et al.	N/A	N/A
9452157	12/2015	Viscomi et al.	N/A	N/A
9458536	12/2015	Felts et al.	N/A	N/A
9474688	12/2015	Weeks et al.	N/A	N/A
9480731	12/2015	Jung et al.	N/A	N/A
9486408	12/2015	Edelson et al.	N/A	N/A
9512495	12/2015	Eyal et al.	N/A	N/A
9517202	12/2015	Chen et al.	N/A	N/A
9517255	12/2015	Turzi	N/A	N/A
9539233	12/2016	Ohtake et al.	N/A	N/A
9554968	12/2016	Weikart et al.	N/A	N/A
9567345	12/2016	Eickhoff et al.	N/A	N/A
9603799	12/2016	Sorayya et al.	N/A	N/A
9629904	12/2016	Hunt	N/A	N/A
9650608	12/2016	Khan et al.	N/A	N/A
9657279	12/2016	Voigt et al.	N/A	N/A
9662450	12/2016	Jones et al.	N/A	N/A
9669107	12/2016	Kim et al.	N/A	N/A
9687541	12/2016	Middaugh et al.	N/A	N/A
9695248	12/2016	Maddon et al.	N/A	N/A
9713643	12/2016	Friedman et al.	N/A	N/A
9724397	12/2016	Cherif-Cheikh et al.	N/A	N/A
9725705	12/2016	Xiang	N/A	N/A
9744237	12/2016	Moazed	N/A	N/A
9750797	12/2016	Brandon et al.	N/A	N/A
9757329	12/2016	Webb et al.	N/A	N/A
9763898	12/2016	Schiffrin et al.	N/A	N/A
9764031	12/2016	Ilyinskii et al.	N/A	N/A
9790511	12/2016	Desfougeres et al.	N/A	N/A
9795614	12/2016	Grosskreutz et al.	N/A	N/A
9815886	12/2016	Grossman et al.	N/A	N/A
9878036	12/2017	Coulter et al.	N/A	N/A
9890400	12/2017	Walther et al.	N/A	N/A
9901627	12/2017	Borodic	N/A	N/A
9937099	12/2017	Weikart et al.	N/A	N/A
9950042	12/2017	Borodic	N/A	N/A
9956194	12/2017	Ohlstein et al.	N/A	N/A
9956435	12/2017	Ruegg et al.	N/A	N/A
9981022	12/2017	Hunt	N/A	N/A
10016338	12/2017	Weikart et al.	N/A	N/A
10022384	12/2017	Giliyar et al.	N/A	N/A
10030238	12/2017	Cossins et al.	N/A	N/A
10047384	12/2017	Medoff	N/A	N/A
10071103	12/2017	Sengupta et al.	N/A	N/A
10080786	12/2017	Dake et al.	N/A	N/A
10081654	12/2017	Verdine et al.	N/A	N/A

10086013	12/2017	Dong et al.	N/A	N/A
10087432	12/2017	Rummel	N/A	N/A
10098840	12/2017	Mosqueira et al.	N/A	N/A
10105421	12/2017	Taylor	N/A	N/A
10111939	12/2017	Thompson et al.	N/A	N/A
10117941	12/2017	Manoharan et al.	N/A	N/A
10118965	12/2017	Kim et al.	N/A	N/A
10143728	12/2017	Jung et al.	N/A	N/A
10201594	12/2018	Ruegg et al.	N/A	N/A
10245317	12/2018	Biemans et al.	N/A	N/A
10258673	12/2018	Pokushalov et al.	N/A	N/A
10260111	12/2018	Etchebarne	N/A	N/A
10286044	12/2018	Bookbinder et al.	N/A	N/A
10293034	12/2018	Jung et al.	N/A	N/A
10307468	12/2018	Palan et al.	N/A	N/A
10325685	12/2018	Apte et al.	N/A	N/A
10327423	12/2018	Ala'Aldeen et al.	N/A	N/A
10329526	12/2018	Salmons et al.	N/A	N/A
10335370	12/2018	Xu et al.	N/A	N/A
10342812	12/2018	Wang	N/A	N/A
10366789	12/2018	Apte et al.	N/A	N/A
10366793	12/2018	Apte et al.	N/A	N/A
10369190	12/2018	Abiad et al.	N/A	N/A
10377785	12/2018	Ala'Aldeen et al.	N/A	N/A
10381112	12/2018	Apte et al.	N/A	N/A
10383519	12/2018	Apte et al.	N/A	N/A
10406139	12/2018	Austad et al.	N/A	N/A
10410749	12/2018	Apte et al.	N/A	N/A
10456473	12/2018	Manoharan et al.	N/A	N/A
10465197	12/2018	McClain et al.	N/A	N/A
10471150	12/2018	Konorty et al.	N/A	N/A
10532019	12/2019	Edelson et al.	N/A	N/A
10973890	12/2020	Abiad et al.	N/A	N/A
12144847	12/2023	Abiad et al.	N/A	N/A
2001/0038858	12/2000	Roser et al.	N/A	N/A
2002/0037833	12/2001	Donovan	N/A	N/A
2002/0044968	12/2001	van Lengerich	N/A	N/A
2002/0064536	12/2001	Hunt	N/A	N/A
2002/0107199	12/2001	Walker	N/A	N/A
2003/0118598	12/2002	Hunt	N/A	N/A
2003/0138437	12/2002	Hunt	N/A	N/A
2003/0138460	12/2002	Hunt	N/A	N/A
2003/0180289	12/2002	Foster et al.	N/A	N/A
2004/0009180	12/2003	Donovan	N/A	N/A
2004/0043374	12/2003	DePablo et al.	N/A	N/A
2004/0086532	12/2003	Donovan	N/A	N/A
2004/0143213	12/2003	Hunter et al.	N/A	N/A
2004/0161776	12/2003	Maddon et al.	N/A	N/A
2004/0204471	12/2003	Seibert	N/A	N/A
2004/0213813	12/2003	Ackerman	N/A	N/A

2004/0214753	12/2003	Britten et al.	N/A	N/A
2004/0220386	12/2003	Steward et al.	N/A	N/A
2005/0025778	12/2004	Cormier et al.	N/A	N/A
2005/0069562	12/2004	Donovan	N/A	N/A
2005/0112135	12/2004	Cormier et al.	N/A	N/A
2005/0123565	12/2004	Subramony et al.	N/A	N/A
2005/0147690	12/2004	Masters et al.	N/A	N/A
2005/0153873	12/2004	Chan et al.	N/A	N/A
2005/0214325	12/2004	David	N/A	N/A
2005/0220854	12/2004	Maa et al.	N/A	N/A
2005/0238663	12/2004	Hunt	N/A	N/A
2005/0238668	12/2004	Wang et al.	N/A	N/A
2005/0244358	12/2004	Hermida Ochoa	N/A	N/A
2005/0260186	12/2004	Bookbinder et al.	N/A	N/A
2005/0266011	12/2004	Maa et al.	N/A	N/A
2005/0271684	12/2004	Trautman et al.	N/A	N/A
2006/0002862	12/2005	Truong-Le et al.	N/A	N/A
2006/0018931	12/2005	Taylor	N/A	N/A
2006/0057165	12/2005	Dimitrakoudis et al.	N/A	N/A
2006/0073207	12/2005	Masters et al.	N/A	N/A
2006/0073333	12/2005	Anderson	N/A	N/A
2006/0182767	12/2005	Borodic	N/A	N/A
2006/0211619	12/2005	Steward et al.	N/A	N/A
2006/0269575	12/2005	Hunt	N/A	N/A
2006/0292224	12/2005	Moore et al.	N/A	N/A
2007/0020295	12/2006	Donovan	N/A	N/A
2007/0026019	12/2006	Hunt	N/A	N/A
2007/0134199	12/2006	Frevert	N/A	N/A
2007/0166332	12/2006	Steward et al.	N/A	N/A
2008/0020000	12/2007	McKerracher	N/A	N/A
2008/0044390	12/2007	Jin et al.	N/A	N/A
2008/0057575	12/2007	Fernandez-Salas et al.	N/A	N/A
2008/0069841	12/2007	Panjwani et al.	N/A	N/A
2008/0096248	12/2007	Steward et al.	N/A	N/A
2008/0138893	12/2007	Steward et al.	N/A	N/A
2008/0187960	12/2007	Foster et al.	N/A	N/A
2008/0220021	12/2007	Modi	N/A	N/A
2008/0241881	12/2007	Steward et al.	N/A	N/A
2008/0274194	12/2007	Miller et al.	N/A	N/A
2008/0279896	12/2007	Heinen et al.	N/A	N/A
2009/0155314	12/2008	Tezel et al.	N/A	N/A
2009/0181007	12/2008	Gennero et al.	N/A	N/A
2009/0181083	12/2008	Holm et al.	N/A	N/A
2009/0324584	12/2008	Hoerr et al.	N/A	N/A
2009/0324647	12/2008	Borodic	N/A	N/A
2010/0150994	12/2009	Kotyla	N/A	N/A
2010/0158951	12/2009	Randolph et al.	N/A	N/A
2010/0184689	12/2009	Panjwani et al.	N/A	N/A
2010/0203559	12/2009	Ester et al.	N/A	N/A
2010/0233741	12/2009	Wang et al.	N/A	N/A

2010/0233802	12/2009	Zhu et al.	N/A	N/A
2010/0260796	12/2009	Belin-Poput et al.	N/A	N/A
2010/0266638	12/2009	Turkel et al.	N/A	N/A
2010/0279953	12/2009	Hunt	N/A	N/A
2010/0291136	12/2009	Jung et al.	N/A	N/A
2010/0330123	12/2009	Thompson et al.	N/A	N/A
2011/0091503	12/2010	Taylor	N/A	N/A
2011/0091921	12/2010	Aigner et al.	N/A	N/A
2011/0200657	12/2010	Baker	N/A	N/A
2012/0039862	12/2011	Borodic	N/A	N/A
2012/0093866	12/2011	Burger et al.	N/A	N/A
2012/0107361	12/2011	Thompson et al.	N/A	N/A
2012/0114625	12/2011	Wiessel et al.	N/A	N/A
2012/0122128	12/2011	Fernandez-Salas et al.	N/A	N/A
2012/0122802	12/2011	Hunt	N/A	N/A
2012/0148562	12/2011	Ho et al.	N/A	N/A
2012/0258126	12/2011	Scholler et al.	N/A	N/A
2013/0017587	12/2012	Nierlich et al.	N/A	N/A
2013/0046275	12/2012	Holzer et al.	N/A	N/A
2013/0121987	12/2012	Taylor	N/A	N/A
2013/0171282	12/2012	Kim et al.	N/A	N/A
2013/0203148	12/2012	Steward et al.	N/A	N/A
2013/0224248	12/2012	Taylor et al.	N/A	N/A
2013/0230856	12/2012	Schneider et al.	N/A	N/A
2013/0330321	12/2012	Turkel et al.	N/A	N/A
2013/0344167	12/2012	Chery et al.	N/A	N/A
2014/0030241	12/2013	Greenberg	N/A	N/A
2014/0030248	12/2013	Turkel et al.	N/A	N/A
2014/0037603	12/2013	Bolster et al.	N/A	N/A
2014/0044695	12/2013	Hunt	N/A	N/A
2014/0056863	12/2013	Greenberg et al.	N/A	N/A
2014/0086900	12/2013	Jung et al.	N/A	N/A
2014/0112908	12/2013	Hunt	N/A	N/A
2014/0120077	12/2013	Ruegg et al.	N/A	N/A
2014/0154237	12/2013	Waugh et al.	N/A	N/A
2014/0161783	12/2013	Jung et al.	N/A	N/A
2014/0199288	12/2013	Hunt	N/A	N/A
2014/0228782	12/2013	Barak	N/A	N/A
2014/0294788	12/2013	Bailey et al.	N/A	N/A
2014/0294803	12/2013	Turkel et al.	N/A	N/A
2014/0301455	12/2013	Lee et al.	N/A	N/A
2014/0356340	12/2013	Burbidge et al.	N/A	N/A
2015/0004149	12/2014	Burbidge et al.	N/A	N/A
2015/0064166	12/2014	Jung et al.	N/A	N/A
2015/0086532	12/2014	Borodic	N/A	N/A
2015/0165003	12/2014	Jung et al.	N/A	N/A
2015/0175707	12/2014	De Jong et al.	N/A	N/A
2015/0238527	12/2014	Chang et al.	N/A	N/A
2015/0297684	12/2014	Borodic	N/A	N/A
2015/0297800	12/2014	Weikart et al.	N/A	N/A

2015/0308988	12/2014	Babuka et al.	N/A	N/A
2015/0313973	12/2014	Forssen et al.	N/A	N/A
2015/0330972	12/2014	Boyle et al.	N/A	N/A
2015/0335576	12/2014	Chan et al.	N/A	N/A
2015/0352226	12/2014	Hunt	N/A	N/A
2016/0000737	12/2015	Zemel et al.	N/A	N/A
2016/0022764	12/2015	Sharma et al.	N/A	N/A
2016/0073650	12/2015	Wiessel et al.	N/A	N/A
2016/0073659	12/2015	Zemel et al.	N/A	N/A
2016/0089442	12/2015	Jung et al.	N/A	N/A
2016/0131669	12/2015	Li et al.	N/A	N/A
2016/0151482	12/2015	Carnes et al.	N/A	N/A
2016/0175408	12/2015	Chang et al.	N/A	N/A
2016/0184413	12/2015	Taylor et al.	N/A	N/A
2016/0194669	12/2015	Argyros et al.	N/A	N/A
2016/0206710	12/2015	Jung et al.	N/A	N/A
2016/0213757	12/2015	Edelson et al.	N/A	N/A
2016/0228409	12/2015	Cross, III	N/A	N/A
2016/0250302	12/2015	Ruegg et al.	N/A	N/A
2016/0354418	12/2015	Quintens et al.	N/A	N/A
2017/0007675	12/2016	Prestrelski et al.	N/A	N/A
2017/0037438	12/2016	Helman et al.	N/A	N/A
2017/0042983	12/2016	Jung et al.	N/A	N/A
2017/0136105	12/2016	Ho et al.	N/A	N/A
2017/0181952	12/2016	Edelson et al.	N/A	N/A
2017/0202954	12/2016	Jezek	N/A	N/A
2017/0204394	12/2016	Slovic et al.	N/A	N/A
2017/0209389	12/2016	Toth et al.	N/A	N/A
2017/0209553	12/2016	Kaspar et al.	N/A	N/A
2017/0216414	12/2016	Tezel et al.	N/A	N/A
2017/0224786	12/2016	Hunt	N/A	N/A
2017/0275279	12/2016	Buckner et al.	N/A	N/A
2017/0290778	12/2016	Waugh	N/A	N/A
2017/0304601	12/2016	Gardner et al.	N/A	N/A
2017/0326242	12/2016	Da Silveira Moreira et al.	N/A	N/A
2017/0342395	12/2016	Xiang	N/A	N/A
2017/0347664	12/2016	Thompson et al.	N/A	N/A
2017/0348226	12/2016	Webb et al.	N/A	N/A
2017/0348263	12/2016	Ohlstein et al.	N/A	N/A
2017/0356002	12/2016	Thompson et al.	N/A	N/A
2018/0027833	12/2017	Wiessel et al.	N/A	N/A
2018/0055779	12/2017	Park et al.	N/A	N/A
2018/0065966	12/2017	Bhattacharjee et al.	N/A	N/A
2018/0071361	12/2017	Abiad et al.	N/A	N/A
2018/0078589	12/2017	Kyle et al.	N/A	N/A
2018/0092851	12/2017	Hassett et al.	N/A	N/A
2018/0147357	12/2017	Marashi et al.	N/A	N/A
2018/0161406	12/2017	Singh	N/A	N/A
2018/0164298	12/2017	Ali et al.	N/A	N/A

2018/0177831	12/2017	Borody	N/A	N/A
2018/0214557	12/2017	Tan et al.	N/A	N/A
2018/0214717	12/2017	Ruegg et al.	N/A	N/A
2018/0237847	12/2017	Culler et al.	N/A	N/A
2018/0243543	12/2017	Baek et al.	N/A	N/A
2018/0258159	12/2017	Grossman et al.	N/A	N/A
2018/0271959	12/2017	Hunt	N/A	N/A
2018/0282309	12/2017	Fieldhouse et al.	N/A	N/A
2018/0296582	12/2017	von Maltzahn et al.	N/A	N/A
2018/0303945	12/2017	Adams et al.	N/A	N/A
2018/0333475	12/2017	Dorsey et al.	N/A	N/A
2018/0344820	12/2017	Jacky et al.	N/A	N/A
2019/0002507	12/2018	Giacalone et al.	N/A	N/A
2019/0008795	12/2018	Waugh	N/A	N/A
2019/0008845	12/2018	Canedo	N/A	N/A
2019/0032105	12/2018	Quiring et al.	N/A	N/A
2019/0038726	12/2018	Turkel et al.	N/A	N/A
2019/0050534	12/2018	Apte et al.	N/A	N/A
2019/0076518	12/2018	Singh	N/A	N/A
2019/0083434	12/2018	Ohlstein et al.	N/A	N/A
2019/0099474	12/2018	Jung et al.	N/A	N/A
2019/0111119	12/2018	Moon et al.	N/A	N/A
2019/0142936	12/2018	Cui et al.	N/A	N/A
2019/0142940	12/2018	Moazed	N/A	N/A
2019/0151465	12/2018	Kim et al.	N/A	N/A
2019/0183785	12/2018	Edelson et al.	N/A	N/A
2019/0183986	12/2018	Sanders	N/A	N/A
2019/0183988	12/2018	Jarstad et al.	N/A	N/A
2019/0194630	12/2018	Bradshaw et al.	N/A	N/A
2019/0201505	12/2018	Palan et al.	N/A	N/A
2019/0201506	12/2018	Lee et al.	N/A	N/A
2019/0201630	12/2018	Vogt	N/A	N/A
2019/0216862	12/2018	Goodman et al.	N/A	N/A
2019/0224253	12/2018	Sandy et al.	N/A	N/A
2019/0231750	12/2018	Schanbacher	N/A	N/A
2019/0231844	12/2018	Chen et al.	N/A	N/A
2019/0233167	12/2018	Felts et al.	N/A	N/A
2019/0240263	12/2018	Goodman et al.	N/A	N/A
2019/0247293	12/2018	Dake et al.	N/A	N/A
2019/0247476	12/2018	Abiad et al.	N/A	N/A
2019/0262266	12/2018	Moen et al.	N/A	N/A
2019/0262438	12/2018	Borodic	N/A	N/A
2019/0274746	12/2018	Toth et al.	N/A	N/A
2019/0276859	12/2018	Bestel-Corre et al.	N/A	N/A
2019/0290740	12/2018	Thompson et al.	N/A	N/A
2019/0298813	12/2018	Jung et al.	N/A	N/A
2019/0321355	12/2018	Anavi-Goffer	N/A	N/A
2019/0321427	12/2018	Ganz et al.	N/A	N/A
2019/0343139	12/2018	Snow et al.	N/A	N/A
2019/0345436	12/2018	Tracy et al.	N/A	N/A

2019/0350987	12/2018	Goodman et al.	N/A	N/A
2019/0351034	12/2018	Abiad et al.	N/A	N/A
2019/0381160	12/2018	Petit et al.	N/A	N/A
2019/0381185	12/2018	Kim et al.	N/A	N/A
2019/0388541	12/2018	Thyagarajan et al.	N/A	N/A
2020/0009473	12/2019	Jung et al.	N/A	N/A
2020/0108129	12/2019	Wu et al.	N/A	N/A
2021/0290739	12/2020	Abiad et al.	N/A	N/A
2021/0369821	12/2020	Abiad et al.	N/A	N/A
2023/0173041	12/2022	Abiad et al.	N/A	N/A
2023/0355723	12/2022	Abiad et al.	N/A	N/A
2023/0355724	12/2022	Abiad et al.	N/A	N/A
2024/0299509	12/2023	Abiad et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
2009339292	12/2010	AU	N/A
1215084	12/1998	CN	N/A
103705913	12/2013	CN	N/A
105833254	12/2015	CN	N/A
1112082	12/2000	EP	N/A
1253932	12/2001	EP	N/A
1390053	12/2003	EP	N/A
1747672	12/2006	EP	N/A
2248518	12/2009	EP	N/A
2373294	12/2010	EP	N/A
2511844	12/2011	EP	N/A
2679217	12/2013	EP	N/A
2692350	12/2013	EP	N/A
2709657	12/2013	EP	N/A
2720712	12/2013	EP	N/A
2943077	12/2014	EP	N/A
3380124	12/2017	EP	N/A
3380125	12/2017	EP	N/A
3405585	12/2017	EP	N/A
3436054	12/2018	EP	N/A
3448399	12/2018	EP	N/A
3538080	12/2018	EP	N/A
2419527	12/2005	GB	N/A
H10203997	12/1997	JP	N/A
H11507072	12/1998	JP	N/A
2005-527470	12/2004	JP	N/A
2012-512162	12/2011	JP	N/A
2019-526609	12/2018	JP	N/A
2011-0106346	12/2010	KR	N/A
2012-0112248	12/2011	KR	N/A
2014-0041460	12/2013	KR	N/A
2535115	12/2013	RU	N/A
WO-1995/017904	12/1994	WO	N/A
WO-1995/33850	12/1994	WO	N/A

WO-96/11699	12/1995	WO	N/A
WO-1996/039166	12/1995	WO	N/A
WO-1997/035604	12/1996	WO	N/A
WO-01/37656	12/2000	WO	N/A
WO-01/58472	12/2000	WO	N/A
WO-01/93827	12/2000	WO	N/A
WO-2004/060384	12/2003	WO	N/A
WO-2005/007185	12/2004	WO	N/A
WO-2005/084410	12/2004	WO	N/A
WO-2005/084705	12/2004	WO	N/A
WO-2005/121354	12/2004	WO	N/A
WO-2006/005910	12/2005	WO	N/A
WO-2006/013357	12/2005	WO	N/A
WO-2006/020208	12/2005	WO	N/A
WO-2006/042149	12/2005	WO	N/A
WO-2006/059093	12/2005	WO	N/A
WO-2006/059105	12/2005	WO	N/A
WO-2006/106823	12/2005	WO	N/A
WO-2006/122123	12/2005	WO	N/A
WO-2007/041664	12/2006	WO	N/A
WO-2007/044809	12/2006	WO	N/A
WO-2007/048638	12/2006	WO	N/A
WO-2008/049631	12/2007	WO	N/A
WO-2008/118691	12/2007	WO	N/A
WO-2008/133884	12/2007	WO	N/A
WO-2009/008595	12/2008	WO	N/A
WO-2009/051837	12/2008	WO	N/A
WO-2009/128644	12/2008	WO	N/A
WO-2010/011597	12/2009	WO	N/A
WO-2010/078242	12/2009	WO	N/A
WO-2010/090677	12/2009	WO	N/A
WO-2010/094463	12/2009	WO	N/A
WO-2010/096134	12/2009	WO	N/A
WO-2010/118888	12/2009	WO	N/A
WO-2010/146591	12/2009	WO	N/A
WO-2010/151840	12/2009	WO	N/A
WO-2011/103465	12/2010	WO	N/A
WO-2011/110948	12/2010	WO	N/A
WO-2012/015912	12/2011	WO	N/A
WO-2012/048854	12/2011	WO	N/A
WO-2012/094163	12/2011	WO	N/A
WO-2012/116209	12/2011	WO	N/A
WO-2012/134240	12/2011	WO	N/A
WO-2013/028781	12/2012	WO	N/A
WO-2013/071138	12/2012	WO	N/A
WO-2013/170052	12/2012	WO	N/A
WO-2014/005039	12/2013	WO	N/A
WO-2014/006217	12/2013	WO	N/A
WO-2014/008138	12/2013	WO	N/A
WO-2014/026161	12/2013	WO	N/A

WO-2014/068317	12/2013	WO	N/A
WO-2014/124096	12/2013	WO	N/A
WO-2014/144842	12/2013	WO	N/A
WO-2014/165617	12/2013	WO	N/A
WO-2014/184746	12/2013	WO	N/A
WO-2014/190944	12/2013	WO	N/A
WO-2015/020982	12/2014	WO	N/A
WO-2015/089452	12/2014	WO	N/A
WO-2015/120231	12/2014	WO	N/A
WO-2015/157595	12/2014	WO	N/A
WO-2015/166242	12/2014	WO	N/A
WO-2016/014750	12/2015	WO	N/A
WO-2016/048689	12/2015	WO	N/A
WO-2016/083549	12/2015	WO	N/A
WO-2016/154112	12/2015	WO	N/A
WO-2016/191545	12/2015	WO	N/A
WO-2017/066121	12/2016	WO	N/A
WO-2017/089894	12/2016	WO	N/A
WO-2017/179775	12/2016	WO	N/A
WO-2017/203038	12/2016	WO	N/A
WO-2017/220553	12/2016	WO	N/A
WO-2018/038301	12/2017	WO	N/A
WO-2018/038585	12/2017	WO	N/A
WO-2018/039318	12/2017	WO	N/A
WO-2018/053004	12/2017	WO	N/A
WO-2018/053021	12/2017	WO	N/A
WO-2018/065972	12/2017	WO	N/A
WO-2018/083692	12/2017	WO	N/A
WO-2018/102343	12/2017	WO	N/A
WO-2018/112138	12/2017	WO	N/A
WO-2018/112419	12/2017	WO	N/A
WO-2018/130099	12/2017	WO	N/A
WO-2018/135722	12/2017	WO	N/A
WO-2018/136617	12/2017	WO	N/A
WO-2018/175899	12/2017	WO	N/A
WO-2018/195097	12/2017	WO	N/A
WO-2018/206772	12/2017	WO	N/A
WO-2018/216974	12/2017	WO	N/A
WO-2018/222652	12/2017	WO	N/A
WO-2018/234645	12/2017	WO	N/A
WO-2019/005773	12/2018	WO	N/A
WO-2019/046311	12/2018	WO	N/A
WO-2019/046529	12/2018	WO	N/A
WO-2019/051204	12/2018	WO	N/A
WO-2019/051380	12/2018	WO	N/A
WO-2019/070850	12/2018	WO	N/A
WO-2019/075452	12/2018	WO	N/A
WO-2019/092504	12/2018	WO	N/A
WO-2019/104135	12/2018	WO	N/A
WO-2019/104136	12/2018	WO	N/A

WO-2019/118393	12/2018	WO	N/A
WO-2019/127719	12/2018	WO	N/A
WO-2019/135244	12/2018	WO	N/A
WO-2019/147799	12/2018	WO	N/A
WO-2019/152667	12/2018	WO	N/A
WO-2019/155391	12/2018	WO	N/A
WO-2019/164134	12/2018	WO	N/A
WO-2019/166514	12/2018	WO	N/A
WO-2019/174560	12/2018	WO	N/A
WO-2019/178055	12/2018	WO	N/A
WO-2019/178359	12/2018	WO	N/A
WO-2019/178487	12/2018	WO	N/A
WO-2019/178490	12/2018	WO	N/A
WO-2019/178494	12/2018	WO	N/A
WO-2019/211382	12/2018	WO	N/A
WO-2019/226599	12/2018	WO	N/A

OTHER PUBLICATIONS

Akers, “Excipient-drug interactions in parental formulations,” *Journal of Pharmaceutical Sciences*, 91(11): 2283-2300 (2002). cited by applicant

Alam et al., “Pain Associated With Injection of Botulinum A Exotoxin Reconstituted Using Isotonic Sodium Chloride With and Without Preservative: A Double-Blind, Randomized Controlled Trial,” *Arch. Dermatol.*, 138(4): 510-514 (2002). cited by applicant

Allergan Inc. “Stability of lyophilized botulinum toxin compositions: Compositions comprising Sucrose”, European Patent No. 13178507.3, Jun. 2015. cited by applicant

Allergan Inc. “Stability of lyophilized botulinum toxin compositions: Compositions of Sucrose”, European Patent No. 13178507.3, Nov. 2019. cited by applicant

Amersdorfer et al., “Molecular Characterization of Murine Humoral Immune Response to Botulinum Neurotoxin Type A Binding Domain as Assessed by Using Phage Antibody Libraries”, *Infect. Immun.*, 65(9): 3743-3752 (1997). cited by applicant

Ansel, “Pharmaceutical Dosage Forms and Drug Delivery Systems” 1999, Lippincott Williams & Wilkins 7th edition, 2 Pages. cited by applicant

Atmaca, “Antioxidant Effects of Sulfur-Containing Amino Acids”, *Yonsei Medical Journal*, 45(5): 776-788 (2004). cited by applicant

Borodic et al., “Effects of Repeated Botulinum Toxin Injections on Orbicularis Oculi Muscle”, *J. Clin. Neuro-ophthalmology*, 12(2): 121-127 (1992). cited by applicant

Borodic et al., “Therapy with Botulinum Toxin”, *Neurological Disease and Therapy*, Ed., Jankovic, pp. 119-157 (1994). cited by applicant

Boyd et al., “The Effect of Botulinum Neurotoxins on the Release of Insulin from the Insulinoma Cell Lines HIT-15 and RINm5F”, *J. Biol. Chem.*, 270(31): 18216-18218 (1995). cited by applicant

Bradshaw et al., “Regulation of Neurotoxin Complex Expression in Clostridium Botulinum Strains 62A, Hall A—hyper, and NCTC 2916”, *Anaerobe*, 10(6): 321-333 (2004). cited by applicant

Brandt et al., “Efficacy and Safety Evaluation of a Novel Botulinum Toxin Topical Gel for the Treatment of Moderate to Severe Lateral Canthal Lines”, *Dermatol. Surg.*, 36(Suppl. 4): 2111-2118 (2010). cited by applicant

Brashear et al., “A multicenter, double-blind, randomized, placebo-controlled, parallel study of the safety and efficacy of Botox (botulinum toxin type A) purified neurotoxin in the treatment of focal upper limb spasticity poststroke”, *Neurology*, 56(Suppl. 3): A78 (2001) Abstract only. cited by applicant

Carpenter et al., “Interactions of stabilizers with proteins during freezing and drying”, *Formulation*

and Delivery of Protein and Peptides, ACS Symposiums Series, vol. 567, Eds. Cleland, J.L. and Langer, R., Washington D.C., American Chemical Society, pp. 134-147 (1994). cited by applicant

Carpenter et al., "Interactions of Stabilizing Additives with Proteins During Freeze-Thawing and Freeze-Drying," International Symposium on Biological Product Freeze-Drying and Formulation, Oct. 24-26, 1990; Karger (1992), 225-239. cited by applicant

Carpenter et al., "Rational Design of Stable Lyophilized Protein Formulations: Theory and Practice," Pharm. Biotechnol., 13: 109-133 (2002). cited by applicant

Carruthers et al., "Botox: Beyond Wrinkles," Clinics in Dermatology, 22(1): 89-93 (2004). cited by applicant

Carruthers et al., "Improvement of Tension-Type Headache When Treating Wrinkles With Botulinum Toxin A Injections," Headache, 39(9): 662-665 (1999). cited by applicant

Chen et al., "Applications of emulsifiers in pharmaceutical formulations" Daily Chemical Industry, 44(10): 590-593 w/ English translation (2014). cited by applicant

Cheng et al., "Unlabeled Uses of Botulinum Toxins: A Review, Part 1," Am. J. Health Syst. Pharm., 63(2): 145-152 (2006). cited by applicant

Choi et al., "Effective Botulinum Toxin Injection Guide for Treatment of Temporal Headache", Toxins (Basel), 8(9): Art. 265, 10 pages (2016). cited by applicant

Decision of the Opposition Division regarding the validity of EP patent No. 2679217, Dated Mar. 12, 2019. cited by applicant

Definition of "Saline" by Oxford Dictionary on Lexico.com, Also meaning of Saline, Online article obtained from <http://www.lexico.com/saline> (2020). cited by applicant

Depiereux et al., "Match-Box: A Fundamentally New Algorithm for the Simultaneous Alignment of Several Protein Sequences," Comput. Appl. Biosci., 8(5): 501-509 (1992). cited by applicant

Doft et al., "Treatment of Axillary Hyperhidrosis With Botulinum Toxin: A Single Surgeon's Experience With 53 Consecutive Patients," Aesthetic Plast. Surg., 35(6): 1079-1086 (2011). cited by applicant

Dressler, "Five-year Experience With incobotulinumtoxinA (Xeomin®): The First Botulinum Toxin Drug Free of Complexing Proteins," European Journal of Neurology, 19(3): 385-389 (2012). cited by applicant

Edgar, "Muscle: Multiple Sequence Alignment with High Accuracy and High Throughput," Nucleic Acids Research, 32(5): 1792-1797 (2004). cited by applicant

European Commission Guidelines: Medicinal products for human use safety, environment and information; Excipients in the label and package leaflet of medicinal products for human use, vol. 3B; Jul. 2003. cited by applicant

European Patent Office, Interlocutory decision dated Mar. 12, 2019, in opposition proceedings of EP 2679217, 29 pages. cited by applicant

Experimental Report D10, Filed by the Patentee Allergan During Opposition Proceedings Corresponding to EP application No. 13178507.3, Allergan ref. No. 166929 u7/011, 1 page, Document filed Nov. 2018. cited by applicant

Experimental Report Filed by the Patentee Allergan During Opposition Proceedings Corresponding to EP application No. 09768315.5, Allergan ref. No. 149691 011 /u7, 2 pages, Document filed May 2018. cited by applicant

Extended European Search Report for EP Application No. EP 19190978.7 dated Nov. 13, 2020. cited by applicant

Finzi et al., "Treatment of Depression with Botulinum Toxin A: A Case Series," Dermatologic Surgery, 32: 645-650 (2006). cited by applicant

Foran et al., "Botulinum Neurotoxin C1 Cleaves Both Syntaxin and SNAP-25 in Intact and Permeabilized Chromaffin Cells: Correlation With Its Blockade of Catecholamine Release," Biochemistry, 35: 2630-2636 (1996). cited by applicant

Foster et al., "Botulinum Toxin Can Smooth Out the Wrinkles," Ophthalmology Times, pp. 13-15

(1997). cited by applicant

Garcia-Rodriguez et al., "Molecular Evolution of Antibody Cross-Reactivity for Two Subtypes of Type A Botulinum Neurotoxin," *Nature Biotech.*, 25(1): 107-116 (2007). cited by applicant

Gennaro, Alfonso R., *Pharmaceutical Sciences*, Remington's 17th Edition, 1985, Mack Publishing Company, Easton, PA. cited by applicant

Gennaro, Alfonso, *Remington: The Science and Practice*, 2000, Lippincott Williams & Wilkins, 2 Pages, 20th edition. cited by applicant

Glogau et al., "Results of a Randomized, Double-Blind, Placebo-Controlled Study to Evaluate the Efficacy and Safety of a Botulinum Toxin Type A Topical Gel for the Treatment of Moderate-To-Severe Lateral Canthal Lines," *J. Drugs Dermatol.*, 11(1): 38-45 (2012). cited by applicant

Goodnough et al., "Recovery of type-A botulinum toxin following lyophilization," *ACS Symposium Series*, 567: 193-203 (1994). cited by applicant

Gotoh, "Significant Improvement in Accuracy of Multiple Protein Sequence Alignments by Iterative Refinement as Assessed by Reference to Structural Alignments", *J. Mol. Biol.*, 264(4): 823-838 (1996). cited by applicant

Grosse et al., "Success of Repeat Detrusor Injections of Botulinum a Toxin in Patients With Severe Neurogenic Detrusor Overactivity and Incontinence," *European Urology*, 47(5): 653-659 (2005). cited by applicant

Hallis et al., "Development of Novel Assays for Botulinum Type A and B Neurotoxins Based on Their Endopeptidase Activities," *J. Clin. Microbiol.*, 34(8): 1934-1938 (1996). cited by applicant

Hardman, Joel G., *Goodman & Gilman's the Pharmacological Basis of Therapeutics*, McGraw-Hill, 2001, 10th Edition, 2 pages. cited by applicant

Humeau et al., "How Botulinum and Tetanus Neurotoxins Block Neurotransmitter Release," *Biochimie*, 82(5): 427-446 (2000). cited by applicant

Innotox, Botulinum toxin products, retrieved from https://www.asushit.com/index.php?route=product/product&product_id=26092, on Jan. 10, 2021, 6 pages. cited by applicant

Jabbari et al., "Treatment of Refractory Pain With Botulinum Toxins—An Evidence-Based Review," *Pain Med.*, 12(11): 1594-1606 (2011). cited by applicant

Johnson et al., "Clostridial botulinum and its Neurotoxins: A Metabolic and Cellular Perspective," *Toxicon*, 39(11):1703-1722 (2001). cited by applicant

Johnson, "Preparation of Peptide and Protein Powders for Inhalation," *Adv. Drug Del. Rev.*, 26(1): 3-15 (1997). cited by applicant

Jones et al., "Development of Improved SNAP-25 Endopeptidase Immunoassays for Botulinum Type A and E Toxins," *J. Immunol. Methods*, 329(1-2): 92-101 (2008). cited by applicant

Kim et al., "The Efficacy and Safety of Liquid-Type Botulinum Toxin Type A for the Management of Moderate to Severe Glabellar Frown Lines", *Plast. Reconstr. Surg.*, 135(3): 732-741 (2015). cited by applicant

Kohl et al., "Comparison of the effect of botulinum toxin A (Botox®) with the highly-purified neurotoxin (NT 201) in the extensor digitorum brevis muscle test," *Mov. Disord.*, 15(Suppl 3): Abst. 165 (2000). cited by applicant

Lalli et al., "The Journey of Tetanus and Botulinum Neurotoxins in Neurons," *Trends in Microbiol.*, 11(9): 431-437 (2003). cited by applicant

Lang "History and Uses of Botox (Botulinum Toxin Type A)," *Lippincott's Case Management*, 9(2): 109-112 (2004). cited by applicant

Lawrence et al., "Detecting Subtle Sequence Signals: A Gibbs Sampling Strategy for Multiple Alignment," *Science*, 262(5131): 208-214 (1993). cited by applicant

Lipham, "Cosmetic and Clinical Applications of Botulinum Toxin," *Slack Incorporated*, 1st Ed., 2 pages, (2004). cited by applicant

Malhotra et al., "Botulinum Toxin and Human Serum Albumin," *Arch. Ophthalmol.*, 121(11): 1661-1662 (2003). cited by applicant

Marconi et al., "A protein-chip Membrane-Capture Assay for Botulinum Neurotoxin Activity," *Toxicol. App. Pharmacol.*, 233(3): 439-446 (2008). cited by applicant

Marini et al., "SiMa, a New Neuroblastoma Cell Line Combining Poor Prognostic Cytogenetic Markers with High Adrenergic Differentiation," *Cancer Genet. Cytogenet.*, 112(2): 161-164 (1999). cited by applicant

Masumoto et al., "Involvement of SNAP-25 in TRH-induced Exocytosis in Pituitary GH4C1 Cells," *Journal of Endocrinology*, 153(1): R5-R10 (1997). cited by applicant

McLeod et al., "Loss of factor VIII activity during storage in PVC containers due to adsorption," *Hemophilia*, 6(2): 89-92 (2000). cited by applicant

Medytox, History of Medytox, retrieved from www.medytox.com/page/history_en?site_id=en, on Jan. 10, 2021, 4; pages. cited by applicant

Mocellin et al., "DNA Array-Based Gene Profiling From Surgical Specimen to the Molecular Portrait of Cancer," *Ann Surg*, 241(1): 16-26 (2005). cited by applicant

Morgenstern et al., "Multiple DNA and Protein Sequence Alignment Based on Segment-to-Segment Comparison," *Proc. Natl. Acad. Sci.*, 93(22): 12098-12103 (1996). cited by applicant

Nabokina et al., "Intracellular Location of SNAP-25 in Human Neutrophils," *Biochem Biophys. Res. Comm.*, 239: 592-597 (1997). cited by applicant

Naumann et al., "Botulinum toxin type A in the treatment of focal, axillary and palmar hyperhidrosis and other hyperhidrotic conditions," *European J. Neurology*, 6(Supp 4): s111-s115 (1999). cited by applicant

Nizai, "Handbook of Pharmaceutical Manufacturing Formulations: Liquid Products," Informa Healthcare, USA, Inc., 52 Vanderbilt Avenue, New York, NY, 10017, vol. 3, p. 58, (2004). cited by applicant

Notredame et al., "T-Coffee: A Novel Method for Fast and Accurate Multiple Sequence Alignment", *J Mol. Biol.*, 302(1): 205-217 (2000). cited by applicant

Ortisi et al., "Blepharospasm and Hemifacial Spasm: A Protocol for Titration of Botulinum Toxin Dose to the Individual Patient and for the Management of Refractory Cases," *Eye (Lond)*, 20(8): 916-922 (2006). cited by applicant

Osterberg et al., "Development of Freeze-Dried Albumin-Free Formulation of Recombinant Factor VIII SQ," *Pharm. Res.*, 14(7): 892-898 (1997). cited by applicant

Panicker et al., "Botulinum Toxins: Pharmacology and Its Current Therapeutic Evidence for Use," *Neurol. India*, 51(4): 455-460 (2003). cited by applicant

Parish, "Commercial Preparations and Handling of Botulinum Toxin Type A and Type B," *Clin. Dermat.*, 21(6): 481-484 (2003). cited by applicant

Parkins et al., "The Formulation of Biopharmaceutical Products," *Pharm. Sci. Technol. Today*, 3(4): 129-137 (2000). cited by applicant

Pellett et al., "A Neuronal Cell-Based Botulinum Neurotoxin Assay for Highly Sensitive and Specific Detection of Neutralizing Serum Antibodies," *FEBS Lett.*, 581(25): 4803-4808 (2007). cited by applicant

Pickar et al., Excerpt, "Example 3", Dosage Calculation, 7th Edition, Thomson, Delmar Learning, 5 Maxwell Drive, Clifton Park, NY, 3 pages (2004). cited by applicant

Pokushalov et al., Long-Term Suppression of Atrial Fibrillation by Botulinum Toxin Injection Into Epicardial Fat Pads in Patients Undergoing Cardiac Surgery One-Year Follow-Up of a Randomized Pilot Study, *Circ Arrhythm Electrophysiol*, 1334-1341 (2015). cited by applicant

Poliziani et al., "Striving for More Good Days: Patient Perspectives on Botulinum Toxin for the Treatment of Cervical Dystonia," *Patient Prefer. Adherence*, 10(10): 1601-1608 (2016). cited by applicant

Raffestin et al., "Organization and Regulation of the Neurotoxin Genes in *Clostridium Botulinum* and *Clostridium Tetani*," *Anaerobe*, 10(2): 93-100 (2004). cited by applicant

Ragona et al., "Management of Parotid Sialoceles with Botulinum Toxin", *Laryngoscope*, vol. 109,

No. 8, pp. 1344-1346 (1999). cited by applicant

Ramirez-Castaneda et al., "Long-term Efficacy and Safety of Botulinum Toxin Injections in Dystonia," *Toxins* (Basel), 5(2): 249-266 (2013). cited by applicant

Rasooly et al., "Development of an In Vitro Assay as an Alternative to the Mouse Bioassay for Clostridium botulinum Neurotoxin Type A," *App. Environ. Microbial.*, 74(14): 4309-4313 (2008). cited by applicant

Rodriguez, "Botox Fights Depression," *Scientific American*, 2012, 1-6, retrieved from <https://www.scientificamerican.com/article/botox-fights-depression/>, on Jan. 12, 2021. cited by applicant

Rowe, Raymond et al., *Aliphatic Polyesters, Handbook of Pharmaceutical Excipients*, Nov. 2003, 4th Edition, pp. 19-21. cited by applicant

Saito., "Pluronic Surfactants," *Journal of Japan Oil Chemists' Society*, 49(10): 1071-1080 pages w/ English abstract (2000). cited by applicant

Schantz et al., "Preparation and Characterization of Botulinum Toxin Type A for Human Treatment," *Neurological Disease and Therapy*, 25: 41-49 (1994). cited by applicant

Schantz et al., "Properties and Use of Botulinum Toxin and Other Microbial Neurotoxins in Medicine," *Microbial. Rev.*, 56(1): 80-99 (1992). cited by applicant

Schulte-Baukloh et al., "Persistence of the Synaptosomal-Associated Protein-25 Cleavage Product After Intrathecal Botulinum Toxin A Injections in Patients with Myelomeningocele Showing an Inadequate Response to Treatment," *BJU Int.*, 100(5):1075-1080 (2007). cited by applicant

Sellers et al., "Principles of Biopharmaceutical Protein Formulation: An Overview," *Meth. Mol. Biol.*, 308: 243-263 (2005). cited by applicant

Sesardic et al., "Role for Standards in Assays of Botulinum Toxins: International Collaborative Study of Three Preparations of Botulinum Type A Toxin," *Biologicals*, 31(4): 256-276 (2003). cited by applicant

Shih et al., "Expression Profiling by Microarrays in Colorectal Cancer (Review)," *Oncol. Rep.*, 13(3): 517-524 (2005). cited by applicant

Shimazaki et al., "Phosphorylation of 25-kDa Synaptosome-Associated Protein. Possible Involvement in Protein Kinase C-mediated Regulation of Neurotransmitter Release," *J. Biol. Chem.*, 271(24):14548-14533 (1996). cited by applicant

Silberstein et al., "Botulinum Toxin Type A in the Prophylactic Treatment of Chronic Tension-Type Headache: A Multicentre, Double-Blind, Randomized, Placebo-Controlled, Parallel-Group Study," *Cephalgia*, 26(7): 790-800 (2006). cited by applicant

Singh "Critical Aspects of Bacterial Protein Toxins," *Adv. Exp. Med. Biol.*, 391: 63-84 (1996). cited by applicant

Subramanian et al., "DIALign-T: An Improved Algorithm for Segment-Based Multiple Sequence Alignment," *BMC Bioinformatics*, 6(Art. 66): 13 Pages (2005). cited by applicant

Thompson et al., "Clustal W: Improving the Sensitivity of Progressive Multiple Sequence Alignment Through Sequence Weighting, Position-Specific Gap Penalties and Weight Matrix Choice", *Nucleic Acids Res.*, 22(22): 4673-4680 (1994). cited by applicant

Trompetto et al., "Do Flexible Inter-Injection Intervals Improve the Effects of Botulinum Toxin A Treatment in Reducing Impairment and Disability in Patients With Spasticity?" *Medical Hypotheses*, 102: 28-32 (2017). cited by applicant

Troost, "Botulinum Toxin Type A (Botox) in the Treatment of Migraine and Other Headaches," *Expert. Rev. Neurother.*, 4(1): 27-31 (2004). cited by applicant

Tumber et al., "Botulinum Toxin Type a Therapy and Human Serum Albumin", *Anesthesiology*, 104(5): 1108-1109 (2006). cited by applicant

Turton et al., "Botulinum and Tetanus Neurotoxins: Structure, Function and Therapeutic Utility," *Trends Biochem. Sci.*, 27(11): 552-558 (2002). cited by applicant

Van Ingen, "Sistema: Sucrose esters, not a 'sweet' solution", *White Paper, Flavours of Asia*, pp. 1-11, downloaded 2013 from flavours.asia/uploads/7/9/8/9/7/7989988/white_paper_-

_sucrose_esters_not_a_sweet_solution. cited by applicant

Veeratterapillay et al., "Discontinuation Rates and Inter-Injection Interval for Repeated Intravesical Botulinum Toxin Type A Injections for Detrusor Overactivity," Int. J. Urol., 21(2): 175-178 (2014). cited by applicant

Walle et al., "Align-m A New Algorithm for Multiple Alignment of Highly Divergent Sequences," Bioinformatics, 20(9): 1428-1435 (2004). cited by applicant

Wieder, J., et al., Understanding Botulinum Toxin, Derma. Surg., 1998, 1172-1174, 24. cited by applicant

Williamson et al., "Clostridial Neurotoxins and Substrate Proteolysis in Intact Neurons," J. Biol. Chem., 271(13) : 7694-7699 (1996). cited by applicant

Yowler et al., "Botulin um Neurotoxin A Activity Is Dependent Upon the Presence of Specific Gangliosides in Neuroblastoma Cells Expressing Synaptotagmin I," J. Biol. Chem., 277(36): 32815-32819 (2002). cited by applicant

Zhou et al., "Biotherapeutic formulation factors affecting metal leachables from stainless steel studied by design experiments," AAPS PharmSciTech, 13(1): 284-294 (2012). cited by applicant

Whelan et al., "The complete amino acid sequence of the Clostridium botulinum type-E neurotoxin, derived by nucleotide-sequence analysis of the encoding gene," European Journal of Biochemistry 204 (1992): 657-667. cited by applicant

Primary Examiner: Navarro; Albert M

Attorney, Agent or Firm: Foley Hoag LLP

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS (1) This application is a divisional of U.S. patent application Ser. No. 18/136,048, filed Apr. 18, 2023, which is a divisional of U.S. patent application Ser. No. 17/185,312, filed Feb. 25, 2021, which is a continuation of U.S. patent application Ser. No. 16/531,800, filed Aug. 5, 2019, which is a divisional of U.S. patent application Ser. No. 15/703,527, filed Sep. 13, 2017, which claims the benefit of U.S. Provisional Application No. 62/394,009, filed Sep. 13, 2016, the entire content of each application is incorporated herein by reference.

TECHNICAL FIELD

(1) The present disclosure relates to solid and liquid pharmaceutical compositions comprising a Clostridial toxin active ingredient and one or more non-protein excipient.

BACKGROUND

(2) A pharmaceutical composition is a formulation which contains at least one active ingredient (such as a Clostridial toxin) as well as, for example, one or more excipients, buffers, carriers, stabilizers, preservatives and/or bulking agents, and is suitable for administration to a patient to achieve a desired diagnostic result or therapeutic effect. The pharmaceutical compositions disclosed herein have diagnostic, therapeutic and/or research utility.

(3) For storage stability and convenience of handling, a pharmaceutical composition can be formulated as a lyophilized (i.e. freeze dried) or vacuum dried powder which can be reconstituted with a suitable fluid, such as saline or water, prior to administration to a patient. Alternately, the pharmaceutical composition can be formulated as an aqueous solution or suspension. A pharmaceutical composition can contain a proteinaceous active ingredient. Unfortunately, a protein active ingredient can be very difficult to stabilize (i.e. maintained in a state where loss of biological activity is minimized), thereby resulting in a loss of protein and/or loss of protein activity during

the formulation, reconstitution (if required) and storage of the pharmaceutical composition prior to use. Stability problems can arise due to surface adsorption of a protein active ingredient, physical instability, such as, e.g., denaturation or aggregation, or chemical instability, such as, e.g., cross-linking, deamidation, isomerization, oxidation, formation of acidic or basic species, Maillard reaction, and fragmentation. To prevent such instability, various protein-based excipients, such as albumin and gelatin, have been used to stabilize a protein active ingredient present in a pharmaceutical composition.

(4) Unfortunately, despite their known stabilizing effects, significant drawbacks exist to the use of protein excipients, such as albumin or gelatin, in a pharmaceutical composition. For example, albumin and gelatin are expensive and increasingly difficult to obtain. Furthermore, blood products or animal derived products such as albumin and gelatin, when administered to a patient can subject the patient to a potential risk of receiving blood borne pathogens or infectious agents. Thus, it is known that the possibility exists that the presence of an animal-derived protein excipient in a pharmaceutical composition can result in inadvertent incorporation of infectious elements into the pharmaceutical composition. For example, it has been reported that use of human serum albumin may transmit prions into a pharmaceutical composition. Thus, it is desirable to find suitable non-protein excipients, such as, e.g., stabilizers, cryo-protectants and lyoprotectants, which can be used to stabilize the protein active ingredient present in a pharmaceutical composition.

(5) The unique characteristics of Clostridial toxins further constrain and hinder the selection of suitable non-protein excipients for a pharmaceutical composition comprising a Clostridial toxin. For example, Clostridial toxins are large proteins having an average molecular weight of approximately 150 kDa, and are further complexed with non-toxin associated proteins that increase the size to approximately 300-900-kDa. The size of a Clostridial toxin complex makes it much more fragile and labile than smaller, less complex proteins, thereby compounding the formulation and handling difficulties if Clostridial toxin stability is to be maintained. Hence, the use of non-protein excipients, such as, e.g., stabilizers, cryo-protectants and lyoprotectants must be able to interact with the Clostridial toxin in a manner which does not denature, fragment or otherwise inactivate the toxin or cause disassociation of the non-toxin associated proteins present in the toxin complex.

(6) Another problem associated with a Clostridial toxin, is the exceptional safety, precision, and accuracy that is necessary for at all steps of the formulation process. Thus, a non-protein excipient should not itself be toxic or difficult to handle so as to not exacerbate the already extremely stringent requirements.

(7) Still another difficulty linked with a Clostridial toxin, is the small amount of Clostridial toxin that is used in a pharmaceutical composition. As with enzymes generally, the biological activities of the Clostridial toxins are dependent, at least in part, upon their three-dimensional conformation. Thus, a Clostridial toxin is detoxified by heat, various chemicals, surface stretching, and surface drying. Additionally, it is known that dilution of a Clostridial toxin complex obtained by the known culturing, fermentation and purification methods to the much lower concentration used in a pharmaceutical composition results in rapid inactivation of the toxin. The low amount of a Clostridial toxin that is used in a pharmaceutical composition, makes this active ingredient very susceptible to adsorption to, e.g., the surfaces of laboratory glassware, vessels, to the vial in which the pharmaceutical composition is reconstituted and to the inside surface of a syringe used to inject the pharmaceutical composition. Such adsorption of a Clostridial toxin to surfaces can lead to a loss of active ingredient and to denaturation of the remaining Clostridial toxin, both of which reduce the total activity of the active ingredient present in the pharmaceutical composition. Hence, the use of non-protein excipients, such as, e.g., stabilizers, cryo-protectants and lyoprotectants must be able to act as surface blockers to prevent the adsorption of a Clostridial toxin to a surface.

(8) Yet another problem connected to a Clostridial toxin, is the pH-sensitivity associated with complex formation. For example, the 900-kDa BoNT/A complex is known to be soluble in dilute

aqueous solutions at pH 3.5-6.8. However, at a pH above about 7 the non-toxic associated proteins dissociate from the 150-kDa neurotoxin, resulting in a loss of toxicity, particularly as the pH rises above pH 8.0. See Edward J. Schantz et al., pp. 44-45, *Preparation and characterization of botulinum toxin type A for human treatment*, in Jankovic, J., et al., *Therapy with Botulinum Toxin* (Marcel Dekker, Inc., 1994). As the non-toxic associated proteins are believed to preserve or help stabilize the secondary and tertiary structures upon which toxicity depends, the dissociation of these proteins results in a more unstable Clostridial toxin. Thus, non-protein excipients useful to formulate a pharmaceutical composition comprising a Clostridial toxin must be able to operate within the confines of a pH level necessary to maintain the activity a Clostridial toxin.

(9) What is needed therefore is a Clostridial toxin pharmaceutical composition wherein a Clostridial toxin active ingredient (such as a botulinum toxin) is stabilized by a non-protein excipient. The present compositions relate to solid and liquid Clostridial toxin active ingredient pharmaceutical compositions with one or more non-protein excipients which functions to stabilize the Clostridial toxin active ingredient present in the solid or liquid pharmaceutical composition.

BRIEF SUMMARY

(10) In one aspect, a pharmaceutical composition comprising a Clostridial toxin active ingredient, a tonicity agent, a surfactant and an antioxidant is provided.

(11) In another aspect, a pharmaceutical composition comprising a Clostridial toxin active ingredient, a surfactant and an antioxidant is provided.

(12) In another aspect, a pharmaceutical composition comprising a Clostridial toxin active ingredient, a lyoprotector, a surfactant and an antioxidant is provided.

(13) In some embodiments, the pharmaceutical compositions comprise a botulinum toxin. In some embodiments, the pharmaceutical composition comprises trehalose. In some embodiments, the pharmaceutical composition comprises sodium chloride. In some embodiments, the composition comprises a poloxamer and/or a polysorbate. In some embodiments, the composition comprises poloxamer 188 and/or polysorbate 20. In some embodiments, the antioxidant is selected from the group consisting of L-methionine, N-acetyl-cysteine (NAC), butylated hydroxytoluene (BHT), ethylene diamine tetraacetic acid sodium salt (EDTA), an EDTA analog, ethylene glycol-bis(2-aminoethylether)-N,N,N',N'-tetraacetic acid (EGTA), an EGTA analog, diethylenetriaminepentaacetic acid (DTPA), a DTPA analog, ascorbic acid, and combinations thereof. In some embodiments, the composition further comprises a buffering agent. In one embodiment, the buffering agent includes histidine buffer. In some embodiments, the composition has a pH of from 5 to 7. In some embodiments, the composition is a liquid formulation. In some embodiments, the composition is a solid lyophilized formulation.

(14) In another aspect, a liquid pharmaceutical composition comprising a Clostridial toxin active ingredient, trehalose, a poloxamer or a polysorbate, and L-methionine or NAC is provided. In some embodiments, the liquid pharmaceutical composition comprises a botulinum toxin. In some embodiments, the liquid pharmaceutical composition further comprises EDTA, EGTA, DTPA or analogs thereof. In some embodiments, the liquid pharmaceutical composition comprises a histidine buffer. In some embodiments, the pH of the liquid pharmaceutical composition ranges from 5 to 7. In some embodiments, the relative weight amount of L-methionine ranges from about 0.1% to about 0.3%. In some embodiments, the relative weight amount of NAC ranges from about 0.1% to about 0.5%. In some embodiments, the relative weight amount of EDTA ranges from about 0.01% to about 0.05%. In some embodiments, the relative weight amount of trehalose ranges from about 1.0 to about 10%. In some embodiments, the relative weight amount of poloxamer 188 ranges from about 0.5% to about 5%. In some embodiments, the relative weight amount of polysorbate ranges from about 0.02% to about 0.06%.

(15) In another embodiment, a liquid pharmaceutical composition is provided. The composition comprises a Clostridial toxin active ingredient; a tonicity agent selected from trehalose, sucrose and combinations thereof; a surfactant selected from a poloxamer, a polysorbate and combinations

thereof; and an antioxidant selected from methionine, NAC, ascorbic acid, butylated hydroxytoluene, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof. In one embodiment, the composition excludes albumin, and in another embodiment, when the antioxidant is methionine the composition excludes a polysorbate. In one embodiment, the composition excludes animal protein stabilizers.

(16) In another embodiment, a liquid pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; and an antioxidant selected from methionine, NAC, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof. In one embodiment, the composition excludes albumin.

(17) In another embodiment, the liquid composition is an animal protein free composition that comprises a botulinum toxin, a poloxamer, and methionine, and optionally includes a disaccharide. In one embodiment, the liquid composition excludes a disaccharide.

(18) In a further embodiment, the liquid composition is an animal protein free composition that comprises a botulinum toxin, a surfactant selected from a poloxamer, a polysorbate and combinations thereof; and an antioxidant selected from the group consisting of a chelating agent, a sacrificial antioxidant, a chain terminator, and combinations thereof. In one embodiment, the antioxidant includes a combination of a chelating agent and a chain terminator.

(19) In yet another embodiment, the liquid composition is an animal-protein free composition comprising a botulinum toxin, a poloxamer surfactant, and methionine, and optionally a disaccharide. In one embodiment, the liquid composition excludes a disaccharide.

(20) In another embodiment, a liquid pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; trehalose or sucrose; a poloxamer; and methionine. In one embodiment, the composition excludes albumin.

(21) In another embodiment, a liquid pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient, trehalose in an amount between 1-15 wt %; a poloxamer in an amount between 0.5-8 wt %; and methionine in an amount between 0.05-5 wt %. In one embodiment, the composition excludes albumin.

(22) In another embodiment, a liquid pharmaceutical composition is contemplated. The composition comprises a botulinum toxin; a disaccharide; a poloxamer; and an antioxidant selected from methionine, NAC, ascorbic acid, butylated hydroxytoluene, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof. In one embodiment, the composition excludes albumin.

(23) In another embodiment, the liquid composition is an animal protein free composition that comprises a botulinum toxin; a disaccharide; a poloxamer; and an antioxidant selected from the group consisting of a chelating agent, a sacrificial antioxidant, a chain terminator, and combinations thereof.

(24) In another embodiment, a liquid composition comprised of a Clostridial toxin active ingredient, a poloxamer, a chelating agent and a chain terminator is provided.

(25) In another embodiment, a liquid composition comprised of a Clostridial toxin active ingredient; a poloxamer; a chelating agent selected from EDTA, EGTA, DTPA and analogs thereof; and NAC is provided.

(26) In another embodiment, a liquid composition comprised of a Clostridial toxin active ingredient, a poloxamer, and methionine is provided.

(27) In another embodiment, a liquid composition comprised of a Clostridial toxin active ingredient, a tonicity agent selected from trehalose, sucrose and combinations thereof, a surfactant selected from a poloxamer, a polysorbate and combinations thereof, a chelating agent and a chain terminator is provided.

(28) In another embodiment, a liquid composition comprised of a Clostridial toxin active ingredient; a tonicity agent selected from trehalose, sucrose and combinations thereof; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; a chelating agent selected from

EDTA, EGTA, DTPA and analogs thereof; and NAC, is provided.

(29) In any of the foregoing embodiments, it is contemplated that the composition is not, in some embodiments, an emulsion and/or excludes nanoparticles comprising an amphiphilic entity.

(30) In another aspect, the present disclosure provides a solid pharmaceutical composition comprising a botulinum toxin; trehalose; a poloxamer or a polysorbate; NAC; and a chelating agent selected from EDTA, EGTA, DTPA and analogs thereof. In an alternative embodiment, the solid pharmaceutical composition comprises a botulinum toxin, trehalose, a poloxamer and L-methionine. In some embodiments, the solid pharmaceutical composition further comprises histidine buffer. In some embodiments, the relative weight amount of L-methionine ranges from about 0.1% to about 0.3%. In some embodiments, the relative weight amount of NAC ranges from about 0.01% to about 0.5%. In some embodiments, the relative weight amount of EDTA ranges from about 0.01% to about 0.05%. In some embodiments, the relative weight amount of trehalose ranges from about 1.0 to about 10%. In some embodiments, the relative weight amount of poloxamer ranges from about 0.5% to about 5%. In some embodiments, the relative weight amount of polysorbate ranges from about 0.02% to about 0.06%.

(31) In one embodiment, a solid or lyophilized pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; a disaccharide selected from trehalose, sucrose and combinations thereof; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; and an antioxidant selected from methionine, N-acetyl cysteine, BHT, EDTA, EGTA, DTPA, ascorbic acid, analogs thereof, and combinations thereof. In one embodiment, the composition excludes albumin, a hydroxyalkyl starch, glutamic acid, glutamine, aspartic acid, asparagine, a polyalcohol, glycine, and/or polyvinylpyrrolidone.

(32) In another embodiment, the solid or lyophilized composition is an animal protein free composition that comprises a botulinum toxin; a disaccharide selected from trehalose, sucrose and combinations thereof; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; and an antioxidant selected from the group consisting of a chelating agent, a sacrificial antioxidant, a chain terminator, and combinations thereof.

(33) In another embodiment, a solid or lyophilized pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; trehalose or sucrose; a poloxamer; and methionine. In one embodiment, the composition excludes albumin, a hydroxyalkyl starch, glutamic acid, glutamine, aspartic acid, asparagine, a polyalcohol, glycine, and/or polyvinylpyrrolidone.

(34) In another embodiment, a solid or lyophilized pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; trehalose in an amount between 1-15 wt %; a poloxamer in an amount between 0.5-8 wt %; and methionine in an amount between 0.05-5 wt %. In one embodiment, the composition excludes albumin, a hydroxyalkyl starch, glutamic acid, glutamine, aspartic acid, asparagine, a polyalcohol, glycine, and/or polyvinylpyrrolidone.

(35) In another embodiment, a solid or lyophilized pharmaceutical composition is contemplated. The composition comprises a botulinum toxin; a disaccharide; a poloxamer; and an antioxidant selected from methionine, N-acetyl cysteine, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof. In one embodiment, the composition excludes albumin, a hydroxyalkyl starch, glutamic acid, glutamine, aspartic acid, asparagine, a polyalcohol, glycine, and/or polyvinylpyrrolidone.

(36) In another embodiment, the solid or lyophilized composition is an animal protein free composition that comprises a botulinum toxin, a disaccharide, a poloxamer; and an antioxidant selected from the group consisting of a chelating agent, a sacrificial antioxidant, a chain terminator, and combinations thereof.

(37) In another embodiment, a lyophilized composition comprised of a Clostridial toxin active ingredient; a lyoprotector selected from sucrose, trehalose, mannitol, sorbitol, glucose, and

combinations thereof; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; a chelating agent; and a chain terminator; is provided.

(38) In another embodiment, a lyophilized composition comprised of a Clostridial toxin active ingredient; a lyoprotector selected from sucrose, trehalose, mannitol, sorbitol, glucose, and combinations thereof; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; a chelating agent selected from EDTA, EGTA, DTPA, and analogs thereof; and NAC; is provided.

(39) In another embodiment, a lyophilized composition comprised of a Clostridial toxin active ingredient; a lyoprotector selected from sucrose, trehalose, mannitol, sorbitol, glucose, and combinations thereof; a poloxamer; and a chain terminator is provided. In one embodiment, the lyophilized composition excludes a chelating agent. In one embodiment, the chain terminator is NAC.

(40) In another embodiment, a lyophilized composition comprised of a Clostridial toxin active ingredient; a lyoprotector selected from sucrose, trehalose, mannitol, sorbitol, glucose, and combinations thereof; a poloxamer; and methionine is provided.

(41) In another embodiment, a lyophilized composition comprised of a Clostridial toxin active ingredient; a lyoprotector selected from sucrose, trehalose, mannitol, sorbitol, glucose, and combinations thereof; a poloxamer; a chelating agent; and a chain terminator is provided.

(42) In another embodiment, a lyophilized composition comprised of a Clostridial toxin active ingredient; a lyoprotector selected from sucrose, trehalose, mannitol, sorbitol, glucose, and combinations thereof; a poloxamer; a chelating agent selected from EDTA, EGTA, DTPA and analogs thereof; and NAC is provided.

(43) In another embodiment, the lyophilized composition is an animal protein free composition that comprises a Clostridial toxin active ingredient; a lyoprotector selected from sucrose, trehalose, mannitol, sorbitol, glucose, and combinations thereof; a poloxamer; and NAC; and optionally includes EDTA, EGTA, DTPA or analogs thereof. In one embodiment, the lyophilized composition excludes EDTA, EGTA, DTPA and analogs thereof.

(44) In another embodiment, a lyophilized composition comprised of a Clostridial toxin active ingredient; a lyoprotector selected from sucrose, trehalose, mannitol, sorbitol, glucose, and combinations thereof; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; and NAC is provided.

(45) In certain embodiments, the lyophilized composition is reconstituted with a tonicity agent selected from trehalose, sucrose, sodium chloride, mannitol, sorbitol, glucose, and combinations thereof. In at least one embodiment, the lyophilized composition is reconstituted with a reconstitution vehicle comprising NaCl prior to administration to a patient.

(46) In any of the foregoing embodiments of solid or liquid compositions, it is contemplated that one or more, in any combination, of these ingredients are excluded polyvinylpyrrolidone, diblock copolymers of polypropylene glycol and polyethylene glycol, and/or a polyalcohol such as inositol, lactitol, isomalt, xylitol, erythritol. In any of the foregoing embodiments of solid or liquid compositions, it is contemplated that the composition is free of animal proteins.

(47) In another aspect, a pharmaceutical composition comprising a Clostridial toxin active ingredient; a tonicity agent and/or a lyoprotector selected from trehalose, sucrose and combinations thereof; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; and an antioxidant selected from methionine, N-acetyl cysteine, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof, is contemplated. In one embodiment, the composition excludes albumin, and in embodiments when the composition is a liquid and the antioxidant is methionine, the surfactant excludes a polysorbate. The composition can be liquid or solid.

(48) In another aspect, a pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; trehalose or sucrose; a poloxamer; and methionine. In one embodiment, the composition excludes albumin. The composition can be liquid or solid.

- (49) In another aspect, a pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient, trehalose in an amount between 1-15 wt %, a poloxamer in an amount between 0.5-8 wt %, and methionine in an amount between 0.05-5 wt %. In one embodiment, the composition excludes albumin. The composition can be liquid or solid.
- (50) In another aspect, a pharmaceutical composition is contemplated. The composition comprises a botulinum toxin; a disaccharide; a poloxamer; and an antioxidant selected from methionine, NAC, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof. In one embodiment, the composition excludes albumin. The composition can be liquid or solid.
- (51) In another aspect, a pharmaceutical composition comprising a Clostridial toxin active ingredient; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; and an antioxidant selected from methionine, N-acetyl cysteine, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof, is provided. In one embodiment, the composition excludes albumin, and in embodiments when the composition is a liquid and the antioxidant is methionine, the surfactant excludes a polysorbate. The composition can be liquid or solid. In one embodiment, the composition further comprises a tonicity agent and/or a lyoprotector. In some embodiments, the tonicity agent is selected from trehalose, sucrose, sodium chloride, mannitol, sorbitol, glucose, and combinations thereof. In alternative embodiments, the lyoprotector is selected from trehalose, sucrose, mannitol, sorbitol, glucose, and combinations thereof. In one embodiment, the tonicity agent and/or lyoprotector is a disaccharide. In one embodiment, the disaccharide is selected from trehalose and sucrose.
- (52) In another aspect, a method for treating depression is contemplated. The method comprises providing for administration, instructing to administer, or administering a composition according to any of the embodiments and aspects described herein.
- (53) In another aspect, a method for treating cardiac arrhythmia is contemplated. The method comprises providing for administration, instructing to administer, or administering a composition according to any of the embodiments and aspects described herein.
-

Description

DETAILED DESCRIPTION

- (1) Compositions described herein, in embodiments, are directed to stable liquid and/or stable solid pharmaceutical compositions of a Clostridial toxin active ingredient, where the composition comprises, in some embodiments, a surfactant and an antioxidant, and optionally a tonicity agent and/or a lyoprotector. In certain liquid compositions, the disaccharide is optional.
- (2) Also as will be described below, the compositions are useful in methods for the treatment of various diseases, disorders, and conditions, including, for example, depression (e.g. major depressive disorder), headache (e.g. migraine, tension headache, and the like), pain, atrial fibrillation, hyperhidrosis, muscle spasticity, cervical dystonia, blepharospasm, overactive bladder (e.g. neurogenic detrusor over-activity, and idiopathic overactive bladder), bladder pain (e.g. interstitial cystitis, and bladder pain syndrome), skin conditions (e.g. wrinkles, fine wrinkles, excess sebum production, acne, and rosacea), irregularities, and the like using the compositions provided herein. Embodiments can include various administration techniques, including, for example, injection, such as intramuscular, intracutaneous, subcutaneous, or the like, instillation, intravenous, transdermal, and topical.

Definitions

- (3) As used herein, the words or terms set forth below have the following definitions:
- (4) The articles “a” and “an” are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

- (5) “About” or “approximately” as used herein means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, (i.e., the limitations of the measurement system). For example, “about” can mean within 1 or more than 1 standard deviations, per practice in the art. Where particular values are described in the application and claims, unless otherwise stated, the term “about” means within an acceptable error range for the particular value. The term “about” when qualifying a value of a stated item, number, percentage, or term refers to a range of plus or minus ten percent of the value of the stated item, percentage, parameter, or term.
- (6) “Administration”, or “to administer” means the step of giving (i.e. administering) a pharmaceutical composition to a subject, or alternatively a subject receiving a pharmaceutical composition. The pharmaceutical compositions disclosed herein can be locally administered by various methods. For example, intramuscular, intradermal, subcutaneous administration, intrathecal administration, intraperitoneal administration, topical (transdermal), instillation, and implantation (for example, of a slow-release device such as polymeric implant or miniosmotic pump) can all be appropriate routes of administration.
- (7) “Alleviating” means a reduction in the occurrence of a pain, of a headache, or of any symptom or cause of a condition or disorder. Thus, alleviating includes some reduction, significant reduction, near total reduction, and total reduction.
- (8) “Animal protein free” means the absence of blood derived, blood pooled and other animal derived products or compounds. “Animal” means a mammal (such as a human), bird, reptile, fish, insect, spider or other animal species. “Animal” excludes microorganisms, such as bacteria. Thus, an animal protein free pharmaceutical composition can include a botulinum neurotoxin. For example, an “animal protein free” pharmaceutical composition means a pharmaceutical composition which is either substantially free or essentially free or entirely free of a serum derived albumin, gelatin and other animal derived proteins, such as immunoglobulins. An example of an animal protein free pharmaceutical composition is a pharmaceutical composition which comprises or which consists of a botulinum toxin (as the active ingredient) and a suitable polysaccharide as a stabilizer or excipient.
- (9) “Antioxidant” refers to any compound which protects an active ingredient from reaction with oxygen. Antioxidants can be broadly divided into three categories: (i) sacrificial antioxidants, which react with oxygen more readily than a particular active ingredient and therefore can scavenge oxygen, e.g., ascorbic acid and sulfites; (ii) chain terminators, which are molecules that form stable radicals due to weak bonds to hydrogen atoms that are attacked in a propagation of radical chains by consumption of oxygen, e.g., methionine, NAC, glutathione, lipoic acid, butylated hydroxytoluene (BHT), and cysteine, (iii) chelating agents, which reduce catalytic activity of transition metals by forming complexes with the metals, e.g., EDTA, EGTA and DTPA and analogs thereof.
- (10) “Biological activity” describes the beneficial or adverse effects of a drug on living matter. When a drug is a complex chemical mixture, this activity is exerted by the substance's active ingredient but can be modified by the other constituents. Biological activity can be assessed as potency or as toxicity by an in vivo LD₅₀ or ED₅₀ assay, or through an in vitro assay such as, for example, cell-based potency assays as described in U.S. 2010/0203559 and U.S. 2010/0233802.
- (11) “Botulinum toxin” means a neurotoxin produced by *Clostridium botulinum*, as well as a botulinum toxin (or the light chain or the heavy chain thereof) made recombinantly by a non-Clostridial species. The phrase “botulinum toxin”, as used herein, encompasses the botulinum toxin serotypes A, B, C, D, E, F and G, and their subtypes and any other types of subtypes thereof, or any re-engineered proteins, analogs, derivatives, homologs, parts, sub-parts, variants, or versions, in each case, of any of the foregoing. “Botulinum toxin”, as used herein, also encompasses a “modified botulinum toxin”. Further “botulinum toxin” as used herein also encompasses a

botulinum toxin complex, (for example, the 300, 600 and 900 kDa complexes), as well as the neurotoxic component of the botulinum toxin (150 kDa) that is unassociated with the complex proteins.

(12) “Clostridial toxin” refers to any toxin produced by a Clostridial toxin strain that can execute the overall cellular mechanism whereby a Clostridial toxin intoxicates a cell and encompasses the binding of a Clostridial toxin to a low or high affinity Clostridial toxin receptor, the internalization of the toxin/receptor complex, the translocation of the Clostridial toxin light chain into the cytoplasm and the enzymatic modification of a Clostridial toxin substrate. Non-limiting examples of Clostridial toxins include Botulinum toxins, such as a BoNT/A, a BoNT/B, a BoNT/C.sub.1, a BoNT/D, a BoNT/E, a BoNT/F, a BoNT/G, a Tetanus toxin (TeNT), a *Baratii* toxin (BaNT), and a *Butyricum* toxin (BuNT). The BoNT/C.sub.2 cytotoxin and BoNT/C.sub.3 cytotoxin, not being neurotoxins, are excluded from the term “Clostridial toxin.” The term Clostridial toxin also includes the approximately 150-kDa Clostridial toxin alone (i.e. without the NAPs). A Clostridial toxin includes naturally occurring Clostridial toxin variants, such as, e.g., Clostridial toxin isoforms and Clostridial toxin subtypes; non-naturally occurring Clostridial toxin variants, such as, e.g., conservative Clostridial toxin variants, non-conservative Clostridial toxin variants, Clostridial toxin chimeric variants and active Clostridial toxin fragments thereof, or any combination thereof. A Clostridial toxin also includes Clostridial toxin complexes, which refers to a complex comprising a Clostridial toxin and non-toxin associated proteins (NAPs), such as, e.g., a Botulinum toxin complex, a Tetanus toxin complex, a *Baratii* toxin complex, and a *Butyricum* toxin complex. Non-limiting examples of Clostridial toxin complexes include those produced by a *Clostridium botulinum*, such as, e.g., a 900-kDa BoNT/A complex, a 500-kDa BoNT/A complex, a 300-kDa BoNT/A complex, a 500-kDa BoNT/B complex, a 500-kDa BoNT/C.sub.1 complex, a 500-kDa BoNT/D complex, a 300-kDa BoNT/D complex, a 300-kDa BoNT/E complex, and a 300-kDa BoNT/F complex.

(13) “Clostridial toxin active ingredient” refers to a molecule which contains any part of a Clostridial toxin that exerts an effect upon or after administration to a subject or patient. As used herein, the term “Clostridial toxin” encompasses (i) a Clostridial toxin complex comprising the approximately 150-kDa Clostridial toxin and other proteins collectively called non-toxin associated proteins (NAPs), (ii) the approximately 150-kDa Clostridial toxin alone (i.e. without the NAPs), or (iii) a modified Clostridial toxin, such as, e.g., a re-targeted Clostridial toxins.

(14) “Deformity” means a cosmetic, physical or functional irregularity, defect, abnormality, imperfection, malformation, depression, or distortion.

(15) “Effective amount” as applied to the biologically active ingredient means that amount of the ingredient which is generally sufficient to effect a desired change in the subject. For example, where the desired effect is a reduction in an autoimmune disorder symptom, an effective amount of the ingredient is that amount which causes at least a substantial reduction of the autoimmune disorder symptom, and without resulting in significant toxicity.

(16) “Effective amount” when used in reference to the amount of an excipient or specific combination of excipients added to a Clostridial toxin composition, refers to the amount of each excipient that is necessary to achieve the desired initial recovered potency of a Clostridial toxin active ingredient. In aspects of this embodiment, an effective amount of an excipient or combination of excipients results in an initial recovered potency of, e.g., at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90% or at least 100%. In other aspects of this embodiment, a therapeutically effective concentration of a Clostridial toxin active ingredient reduces a symptom associated with the ailment being treated by, e.g., at most 10%, at most 20%, at most 30%, at most 40%, at most 50%, at most 60%, at most 70%, at most 80%, at most 90% or at most 100%.

(17) “Heavy chain” means the heavy chain of a botulinum neurotoxin. It has a molecular weight of about 100 kDa and can be referred to as the H chain, or as H.

- (18) H.sub.C means a fragment (about 50 kDa) derived from the H chain of a botulinum neurotoxin which is approximately equivalent to the carboxyl end segment of the H chain, or the portion corresponding to that fragment in the intact H chain. It is believed to be immunogenic and to contain the portion of the natural or wild type botulinum neurotoxin involved in high affinity, presynaptic binding to motor neurons.
- (19) H.sub.N means a fragment (about 50 kDa) derived from the H chain of a botulinum neurotoxin which is approximately equivalent to the amino end segment of the H chain, or the portion corresponding to that fragment in the intact in the H chain. It is believed to contain the portion of the natural or wild type botulinum neurotoxin involved in the translocation of the L chain across an intracellular endosomal membrane.
- (20) "Light chain" means the light chain of a Clostridial neurotoxin. It has a molecular weight of about 50 kDa, and can be referred to as the L chain, L, or as the proteolytic domain (amino acid sequence) of a botulinum neurotoxin.
- (21) LH.sub.N or L-H.sub.N means a fragment derived from a Clostridial neurotoxin that contains the L chain, or a functional fragment thereof coupled to the H.sub.N domain. It can be obtained from the intact Clostridial neurotoxin by proteolysis, so as to remove or to modify the H.sub.C domain.
- (22) "Implant" means a controlled release (e.g., pulsatile or continuous) composition or drug delivery system. The implant can be, for example, injected, inserted or implanted into a human body.
- (23) "Liquid composition", "liquid pharmaceutical composition", or "liquid formulation" refers to a pharmaceutically active preparation of drug or biological which is capable of being stored in a liquid pharmaceutical excipient, such as a buffering agent, for an extended period of time, such that it can be ready-to-use as needed by a clinician. The liquid pharmaceutical composition is manufactured without a lyophilization process.
- (24) "Local administration" means direct administration of a pharmaceutical at or to the vicinity of a site on or within an animal body, at which site a biological effect of the pharmaceutical is desired, such as via, for example, intramuscular or intra- or subdermal injection or topical administration. Local administration excludes systemic routes of administration, such as intravenous or oral administration. Topical administration is a type of local administration in which a pharmaceutical agent is applied to a patient's skin.
- (25) "Lyoprotector" or "lyoprotectant" means a substance that is included in a lyophilized formulation to protect a Clostridial toxin active ingredient during the freeze-drying process. Lyoprotectors include for example polyhydroxy compounds such as sugars (mono-, di-, and polysaccharides), polyalcohols, and their derivatives. Exemplary lyoprotectors which can be used with the lyophilized formulations disclosed herein include sucrose, trehalose, mannitol, sorbitol, glucose, raffinose, maltose, glycerol, lactose, fructose, galactose, and combinations thereof.
- (26) "Lyophilized composition", "lyophilized pharmaceutical composition", "lyophilized formulation", or "solid composition" refers to a formulation containing a Clostridial toxin active ingredient which has been subjected to a lyophilization, freeze-drying or vacuum-drying process; and can be reconstituted with a reconstitution vehicle, such as for example saline or water, prior to administration to a patient. The lyophilized composition can be a freeze-dried composition or a vacuum-dried composition.
- (27) "Modified botulinum toxin" means a botulinum toxin that has had at least one of its amino acids deleted, modified, or replaced, as compared to a native botulinum toxin. Additionally, the modified botulinum toxin can be a recombinantly produced neurotoxin, or a derivative or fragment of a recombinantly made neurotoxin. A modified botulinum toxin retains at least one biological activity of the native botulinum toxin, such as, the ability to bind to a botulinum toxin receptor, or the ability to inhibit neurotransmitter release from a neuron. One example of a modified botulinum toxin is a botulinum toxin that has a light chain from one botulinum toxin serotype (such as

serotype A), and a heavy chain from a different botulinum toxin serotype (such as serotype B). Another example of a modified botulinum toxin is a botulinum toxin coupled to a neurotransmitter, such as substance P.

(28) "Mutation" means a structural modification of a naturally occurring protein or nucleic acid sequence. For example, in the case of nucleic acid mutations, a mutation can be a deletion, addition or substitution of one or more nucleotides in the DNA sequence. In the case of a protein sequence mutation, the mutation can be a deletion, addition or substitution of one or more amino acids in a protein sequence. For example, a specific amino acid comprising a protein sequence can be substituted for another amino acid, for example, an amino acid selected from a group which includes the amino acids alanine, asparagine, cysteine, aspartic acid, glutamic acid, phenylalanine, glycine, histidine, isoleucine, lysine, leucine, methionine, proline, glutamine, arginine, serine, threonine, valine, tryptophan, tyrosine or any other natural or non-naturally occurring amino acid or chemically modified amino acids. Mutations to a protein sequence can be the result of mutations to DNA sequences that when transcribed, and the resulting mRNA translated, produce the mutated protein sequence. Mutations to a protein sequence can also be created by fusing a peptide sequence containing the desired mutation to a desired protein sequence.

(29) "Patient" means a human or non-human subject receiving medical or veterinary care. Accordingly, the compositions as disclosed herein can be used in treating any animal, such as, for example, mammals, or the like.

(30) "Peripherally administering" or "peripheral administration" means subdermal, intradermal, transdermal, or subcutaneous administration, but excludes intramuscular administration.

"Peripheral" means in a subdermal location, and excludes visceral sites.

(31) "Pharmaceutical composition" means a composition comprising an active pharmaceutical ingredient, such as, for example, a Clostridial toxin active ingredient such as a botulinum toxin, and at least one additional ingredient, such as, for example, a stabilizer or excipient or the like. A pharmaceutical composition is therefore a formulation which is suitable for diagnostic or therapeutic administration to a subject, such as a human patient. The pharmaceutical composition can be, for example, in a lyophilized or vacuum dried condition, a solution formed after reconstitution of the lyophilized or vacuum dried pharmaceutical composition, or as a solution or solid which does not require reconstitution.

(32) "Pharmacologically acceptable excipient" is synonymous with "pharmacological excipient" or "excipient" and refers to any excipient that has substantially no long term or permanent detrimental effect when administered to mammal and encompasses compounds such as, e.g., stabilizing agent, a bulking agent, a cryo-protectant, a lyoprotectant, an additive, a vehicle, a carrier, a diluent, or an auxiliary. An excipient generally is mixed with an active ingredient, or permitted to dilute or enclose the active ingredient and can be a solid, semi-solid, or liquid agent. It is also envisioned that a pharmaceutical composition comprising a Clostridial toxin active ingredient can include one or more pharmaceutically acceptable excipients that facilitate processing of an active ingredient into pharmaceutically acceptable compositions. Insofar as any pharmacologically acceptable excipient is not incompatible with the Clostridial toxin active ingredient, its use in pharmaceutically acceptable compositions is contemplated. Non-limiting examples of pharmacologically acceptable excipients can be found in, e.g., *Pharmaceutical Dosage Forms and Drug Delivery Systems* (Howard C. Ansel et al., eds., Lippincott Williams & Wilkins Publishers, 7^{sup}.th ed. 1999); *Remington: The Science and Practice of Pharmacy* (Alfonso R. Gennaro ed., Lippincott, Williams & Wilkins, 20^{sup}.th ed. 2000); *Goodman & Gilman's The Pharmacological Basis of Therapeutics* (Joel G. Hardman et al., eds., McGraw-Hill Professional, 10^{sup}.th ed. 2001); and *Handbook of Pharmaceutical Excipients* (Raymond C. Rowe et al., APhA Publications, 4^{sup}.th edition 2003), each of which is hereby incorporated by reference in its entirety.

(33) The constituent ingredients of a pharmaceutical composition can be included in a single composition (that is, all the constituent ingredients, except for any required reconstitution fluid, are

present at the time of initial compounding of the pharmaceutical composition) or as a two-component system, for example a vacuum-dried composition reconstituted with a reconstitution vehicle which can, for example, contain an ingredient not present in the initial compounding of the pharmaceutical composition. A two-component system can provide several benefits, including that of allowing incorporation of ingredients which are not sufficiently compatible for long-term shelf storage with the first component of the two-component system. For example, the reconstitution vehicle may include a preservative which provides sufficient protection against microbial growth for the use period, for example one-week of refrigerated storage, but is not present during the two-year freezer storage period during which time it might degrade the toxin. Other ingredients, which may not be compatible with a botulinum toxin or other ingredients for long periods of time, can be incorporated in this manner; that is, added in a second vehicle (e.g. in the reconstitution vehicle) at the approximate time of use. A pharmaceutical composition can also include preservative agents such as benzyl alcohol, benzoic acid, phenol, parabens and sorbic acid. Pharmaceutical compositions can include, for example, excipients, such as surface active agents; dispersing agents; inert diluents; granulating and disintegrating agents; binding agents; lubricating agents; preservatives; physiologically degradable compositions such as gelatin; aqueous vehicles and solvents; oily vehicles and solvents; suspending agents; dispersing or wetting agents; emulsifying agents, demulcents; buffers; salts; thickening agents; fillers; antioxidants; stabilizing agents; and pharmaceutically acceptable polymeric or hydrophobic materials and other ingredients known in the art and described, for example in Genaro, ed., 1985, Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pa., which is incorporated herein by reference.

(34) "Polysaccharide" means a polymer of more than two saccharide molecule monomers. The monomers can be identical or different.

(35) "Stabilizing agent", "stabilization agent" or "stabilizer" means a substance that acts to stabilize a Clostridial toxin active ingredient such that the potency of the pharmaceutical composition is increased relative to an unstabilized composition.

(36) "Stabilizers" can include excipients, and can include protein and non-protein molecules.

(37) "Surfactant" refers to a natural or synthetic amphiphilic compound. A surfactant can be non-ionic, zwitterionic, or ionic. Non-limiting examples of surfactants include a poloxamer, a polysorbate, and combinations thereof.

(38) "Therapeutic formulation" means a formulation can be used to treat and thereby alleviate a disorder or a disease, such as, for example, a disorder or a disease characterized by hyperactivity (i.e. spasticity) of a peripheral muscle.

(39) "Therapeutically effective concentration", "therapeutically effective amount," "effective amount," "effective dose," and "therapeutically effective dose" refer to the minimum dose of a Clostridial toxin active ingredient necessary to achieve the desired therapeutic effect and includes a dose sufficient to reduce a symptom associated with ailment being treated.

(40) "TEM" as used herein, is synonymous with "Targeted Exocytosis Modulator" or "retargeted endopeptidase." Generally, a TEM comprises an enzymatic domain from a Clostridial toxin light chain, a translocation domain from a Clostridial toxin heavy chain, and a targeting domain. The targeting domain of a TEM provides an altered cell targeting capability that targets the molecule to a receptor other than the native Clostridial toxin receptor utilized by a naturally-occurring Clostridial toxin. This re-targeted capability is achieved by replacing the naturally-occurring binding domain of a Clostridial toxin with a targeting domain having a binding activity for a non-Clostridial toxin receptor. Although binding to a non-Clostridial toxin receptor, a TEM undergoes all the other steps of the intoxication process including internalization of the TEM/receptor complex into the cytoplasm, formation of the pore in the vesicle membrane and di-chain molecule, translocation of the enzymatic domain into the cytoplasm, and exerting a proteolytic effect on a component of the SNARE complex of the target cell.

(41) "Tonicity agent" means a low molecular weight excipient which is included in a formulation

to provide isotonicity. Non-limiting examples of a tonicity agent include a disaccharide such as trehalose or sucrose; a polyalcohol such as sorbitol or mannitol; a monosaccharide such as glucose; and a salt such as sodium chloride.

(42) “Topical administration” excludes systemic administration of the neurotoxin. In other words, and unlike conventional therapeutic transdermal methods, topical administration of botulinum toxin does not result in significant amounts, such as the majority of, the neurotoxin passing into the circulatory system of the patient.

(43) “Treating” means to alleviate (or to eliminate) at least one symptom of a condition or disorder, such as, for example, wrinkles, spasticity, depression, pain (such as, for example, headache pain), bladder overactivity, or the like, either temporarily or permanently.

(44) As used herein, the term “unit” or “U” refers to the LD_{sub}50 dose or the dose determined by a cell based potency assay (CBPA). The LD_{sub}50 dose is defined as the amount of a Clostridial toxin active ingredient, Clostridial toxin complex or modified Clostridial toxin that killed 50% of the mice injected with the Clostridial toxin, Clostridial toxin complex or modified Clostridial toxin. The CBPA dose is determined as described in U.S. Pat. No. 8,618,261, the assay details of which are incorporated by reference herein.

(45) “Variant” means a Clostridial neurotoxin, such as wild-type botulinum toxin serotype A, B, C, D, E, F or G, that has been modified by the replacement, modification, addition or deletion of at least one amino acid relative to wild-type botulinum toxin, which is recognized by a target cell, internalized by the target cell, and catalytically cleaves a SNARE (SNAP (Soluble NSF Attachment Protein) Receptor) protein in the target cell.

(46) An example of a variant neurotoxin component can comprise a variant light chain of a botulinum toxin having one or more amino acids substituted, modified, deleted and/or added. This variant light chain may have the same or better ability to prevent exocytosis, for example, the release of neurotransmitter vesicles. Additionally, the biological effect of a variant may be decreased compared to the parent chemical entity. For example, a variant light chain of a botulinum toxin type A having an amino acid sequence removed may have a shorter biological persistence than that of the parent (or native) botulinum toxin type A light chain.

(47) Pharmaceutical Compositions

(48) In a first aspect, a pharmaceutical composition comprising (or consisting of, or consisting essentially of) a Clostridial toxin active ingredient, a disaccharide, a surfactant and an antioxidant is described. The composition can, in one embodiment, be a solid composition, such as a lyophilized powder that is reconstituted prior to use. In another embodiment, the composition is a liquid composition; that is, the composition is manufactured and stored in liquid form. Studies were conducted demonstrating that the compositions stabilize the Clostridial toxin active ingredient to retain its potency, as will now be described with respect to Examples 1-11.

(49) In one study, described in Example 1, compositions were prepared that comprised botulinum toxin as a model Clostridial toxin active ingredient, a disaccharide, a surfactant, and an antioxidant. The formulations were lyophilized and stored under various temperatures for selected periods of time, ranging from 1 month to 7.5 months. Potency of the lyophilized formulations after storage and reconstitution with saline was tested by a cell based potency assay. The results were normalized to target potency. The potencies of the solid compositions are shown in Tables 1.1, 1.2, 2 and 3.

(50) TABLE-US-00001 TABLE 1.1 Lyophilized formulations Excipient, % w/w Normalized potency, TWEEN® storage at 25° C. Formulation Treh 20 P 188 NaCl Met NAC Buffer T0 3 mo 6 mo Comparator 1 3 0.04 0.9 0.2 — water 80.6% 86.4% 80.8% Comparator 2 2 — 4 — — 20 mM 81.5% 68.6% 68.0% Histidine pH 5.5 Formulation 1 2 — 4 — — 0.03 20 mM 98.6% 91.9% 86.9% Histidine pH 5.5 Treh = trehalose; P 188 = poloxamer P 188; Met = L-methionine; NAC = N-acetyl-L-cysteine.

(51) TABLE-US-00002 TABLE 1.2 Lyophilized formulations Excipient, % w/w Normalized

potency, TWEEN® storage at 40° C. Formulation Treh 20 P 188 NaCl Met NAC Buffer 1 mo 3 mo 6 mo Comparator 1 3 0.04 0.9 0.2 Water 74.8% 70.3% 24.0% Comparator 2 2 4 20 mM 64.8% 57.8% 46.8% Histidine pH 5.5 Formulation 1 2 4 0.03 20 mM 86.5% 77.9% 62.3% Histidine pH 5.5 Treh = trehalose; P 188 = poloxamer P 188; Met = L-methionine; NAC = N-acetyl-L-cysteine.

(52) The data in Table 1.1 demonstrates that a lyophilized botulinum toxin composition comprised of trehalose, poloxamer, and NAC retained more than 85% potency after storage for six months at 25° C. The formulation referred to as Comparator 2, which was identical to Formulation 1 except that it lacked an antioxidant (i.e., no methionine, no NAC), had only 68% retained potency after storage for six months at 25° C. Formulation 1 had about 7.5% higher potency than Comparator 1 after storage for six months at 25° C. (calculated as a percent difference, $((86.9\%-80.8\%)/80.8\%)$). Formulation 1 had 27.8% higher potency than Comparator 2 after storage for six months at 25° C. (calculated as a percent difference, $((86.9\%-68\%)/68\%)$). The data in Table 1.2 shows potency data for the same three formulations as Table 1.1 (Formulation 1 and Comparator 1 and Comparator 2), demonstrating that a lyophilized botulinum toxin composition comprised of trehalose, poloxamer, and NAC retained more than 60% potency after storage for six months at 40° C. The formulation referred to as Comparator 2, which was identical to Formulation 1 except that it lacked an antioxidant (e.g., no methionine, no NAC), had only 46.8% retained potency after storage for six months at 40° C. Formulation 1 had about 160% higher potency than Comparator 1 after storage for six months at 40° C. (calculated as a percent difference). Formulation 1 had about 33% higher potency than Comparator 2 after storage for six months at 40° C. (calculated as a percent difference). Accordingly, in one embodiment, a lyophilized composition is contemplated, where the composition is comprised of a Clostridial toxin active ingredient, such as a botulinum toxin; a disaccharide; a surfactant; and an antioxidant. In one embodiment, the composition retains potency after storage for six months at 25° C. that is at least about 15%, 20%, or 25% greater than a formulation lacking the antioxidant. In another embodiment, a lyophilized composition is contemplated, where the composition is comprised of a Clostridial toxin active ingredient, such as a botulinum toxin; a disaccharide; a surfactant; and an antioxidant. In one embodiment, the composition retains potency after storage for six months at 40° C. that is at least about 20%, 25%, or 30% greater than a formulation lacking the antioxidant. In another embodiment, a lyophilized composition is contemplated, where the composition is comprised of a Clostridial toxin active ingredient, such as a botulinum toxin; a disaccharide; a surfactant; and an antioxidant. In one embodiment, the composition retains at least about 55% or 60% potency after storage for six months at 40° C., where potency is measured in a cell-based potency assay.

(53) It will be appreciated that potency of the Clostridial toxin can be ascertained in various ways, such as the in vitro cell-based potency assay used in the studies discussed herein or in an in vivo assay. An exemplary in vivo assay is the LD.sub.50 potency assay, where the potency of the toxin is expressed as a multiple of the LD.sub.50 value for a mouse, one unit (U) of toxin being defined as being the equivalent amount of toxin that kills 50% of a group of 18 to 20 female Swiss-Webster mice, weighing about 20 grams each. Any known method for assessing potency can be used to determine whether a composition or a particular composition components stabilize a toxin. The in vitro cell-based potency assay used herein is not intended to be limiting to the possible approaches to ascertain potency.

(54) In another study, with continuing reference to Example 1, lyophilized botulinum toxin compositions were prepared with trehalose; a polysorbate or a poloxamer; and methionine or NAC. Comparator formulations were prepared with sucrose (Comparator 3) and with no antioxidant (Comparator 2). The lyophilized formulations and lyophilized comparator formulations were placed in storage for 1 month at -20° C. or at 40° C. Potency of the botulinum toxin was measured using a cell-based potency assay and the results are shown in Table 2.

(55) TABLE-US-00003 TABLE 2 Lyophilized formulations Normalized potency Excipient, % w/w (% of target) TWEEN® 1 mo -20° 1 mo 40° Formulation Treh Sucr 20 P 188 NaCl Met NAC

Buffer T0 C. C. Comparator 2 2 4 20 mM 87.15% 88.95% 76.32% Histidine pH 5.5 Comparator 3 3 0.04 0.9 0.2 Water 84.06% 85.18% 72.86% Formulation 2 2 4 0.2 20 mM 98.60% 120.85% 91.17% Histidine pH 6.0 Formulation 3 8 0.04 0.03 20 mM 105.2% 110.05% 96.89% Histidine pH 6.0 Treh = trehalose; Sucr = sucrose; P 188 = poloxamer P 188; Met = L-methionine; NAC = N-acetyl-L-cysteine.

(56) Inspection of the data in Table 2 for compositions identified as Formulation 2 and Comparator 2 demonstrates the stabilizing effect on the botulinum toxin by the combination of a disaccharide, a surfactant, and an antioxidant. Formulation 2 and Comparator 2 were identical in all respects other than Comparator 2 lacked an antioxidant. The potency data for these two compositions shows that the antioxidant acts to stabilize the toxin, as Formulation 2 retained more than 90% potency after storage for one month at 40° C. whereas the comparative formulation (Comparator 2) lacking the antioxidant had only 76% potency after storage for one month at 40° C. Formulation 3, comprised of trehalose, polysorbate surfactant (TWEEN®-20), and an antioxidant NAC, retained more than 95% potency after storage for one month at 40° C., considerably better than either comparative formulation. Accordingly, in one embodiment, a lyophilized composition is contemplated, where the composition is comprised of a Clostridial toxin active ingredient, such as a botulinum toxin; a disaccharide; a surfactant; and an antioxidant. In one embodiment, the composition retains at least about 80%, 85% or 90% potency after storage for one month at 40° C., where potency is measured in a cell-based potency assay. In another embodiment, a lyophilized composition is contemplated, where the composition is comprised of a Clostridial toxin active ingredient, such as a botulinum toxin; a disaccharide; a surfactant selected from a poloxamer and a polysorbate; and an antioxidant selected from methionine, NAC, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof. In one embodiment, the composition retains at least about 80%, 85% or 90% potency after storage for one month at 40° C., where potency is measured in a cell-based potency assay. Such compositions, in another embodiment, retain potency after storage for one month at 40° C. that is at least about 18%, 20% or 24% greater than a formulation lacking the antioxidant.

(57) With continued reference to Example 1, compositions comprised of a Clostridial toxin active ingredient (botulinum toxin), trehalose, a poloxamer surfactant (e.g. KOLLIPHOR® P-188) and methionine as a stabilizing antioxidant were prepared. A comparative formulation lacking the methionine was prepared (Comparator 2). The compositions were lyophilized and placed in storage for 7.5 months at 25° C. Potency of the botulinum toxin was measured after 3 months and after 7.5 months of storage using a cell-based potency assay and normalized to the target potency. The results are shown in Table 3.

(58) TABLE-US-00004 TABLE 3 Lyophilized formulations Normalized potency, Excipient, % w/w storage at 25° C. Formulation Trehalose P-188 Met Buffer T0 3 mo 7.5 mo Comparator 2 2 4 20 mM 87.1% 78.0% 78.0% Histidine, pH 5.5 Formulation 2 2 4 0.2 20 mM 98.6% 97.0% 98.0% Histidine, pH 6.0 Formulation 4 8 0.6 0.2 20 mM 86.5% 83.0% 84.0% Histidine, pH 6.0 P-188 = Poloxamer P-188; Met = L-methionine.

(59) Inspection of the data in Table 3 for compositions identified as Formulation 2 and Comparator 2 demonstrates the stabilizing effect on the botulinum toxin by the combination of a tonicity agent, a surfactant, and an antioxidant. Formulation 2 and Comparator 2 were identical in all respects other than Comparator 2 lacked the antioxidant and had a slightly lower pH. The potency data for these two compositions shows that the antioxidant acts to stabilize the toxin, as Formulation 2 retained more than 95% potency after storage for 3 months and after storage for 7.5 months at 25° C. whereas the comparative formulation lacking the antioxidant had only 78% potency after storage for 3 months and after storage for 7.5 months at 25° C. Formulation 4 was comprised of the same components as Formulation 2, but with a higher weight percent of trehalose and a lower weight percent of poloxamer surfactant. Formulation 4 also stabilized the botulinum toxin better than the Comparator 2 formulation that lacked methionine. Accordingly, in one embodiment, a lyophilized composition is contemplated, where the composition is comprised of a Clostridial toxin active

ingredient, such as a botulinum toxin; trehalose; a poloxamer surfactant; and methionine. In one embodiment, the composition retains at least about 80%, 85% or 90% potency after storage for three months at 25° C., where potency is measured in a cell-based potency assay. In another embodiment, a lyophilized composition is contemplated, where the composition is comprised of a Clostridial toxin active ingredient, such as a botulinum toxin; trehalose; a poloxamer surfactant; and methionine. In one embodiment, the composition retains potency after storage for three months at 25° C. that is at least about 10%, 15%, or 20% greater than a formulation lacking the antioxidant. (60) Example 2 describes another study where liquid compositions comprised of a Clostridial toxin active ingredient, a tonicity agent, a surfactant, and an antioxidant were prepared. Liquid solutions of a Clostridial toxin active ingredient, using botulinum toxin as a model, were prepared with a disaccharide tonicity agent, a poloxamer surfactant, and an antioxidant were prepared. In this particular study, the disaccharide tonicity agent was trehalose and the poloxamer surfactant was poloxamer P188. Three formulations were prepared, each with the same amount of botulinum toxin, 8 w/w % trehalose, and 4 w/w % poloxamer P188 in histidine buffer. The target potency was 100 Units/ml. Formulation 10 had no antioxidant; Formulation 11 contained NAC, and Formulation 12 contained L-methionine. The liquid compositions were placed in glass vials and stored at -70° C., 5° C., 25° C. and 40° C. for one month. Potency of the botulinum toxin was measured prior to storage and after one month of storage using a cell-based potency assay. Results are shown in Table 4.

(61) TABLE-US-00005 TABLE 4 Liquid formulations Potency, U/mL 1 mo. -70° 1 mo. 5° 1 mo. 25° 1 mo. 40° Formulation No.* Antioxidant T0 C. C. C. C. Formulation 10 None 128 135 135 106 0.225 Formulation 11 N-acetyl-L- 128 133 129 61 0.2 cysteine, 0.2% w/w Formulation 12 L-methionine, 133 146 146 145 138 0.2% w/w *Each formulation contained the same amount of botulinum toxin, 8 w/w % trehalose, and 4 w/w % poloxamer P188 in histidine buffer.

(62) The liquid composition comprising methionine (Formulation 12) retained its potency after one month storage at all four temperatures, including 40° C., whereas the liquid composition with no antioxidant (Formulation 10) lost approximately 17% potency after storage for 1 month at 25° C. and lost essentially all activity (i.e., complete inactivation of the toxin) after storage for 1 month at 40° C. The liquid composition comprising NAC lost over 50% potency after storage for 1 month at 25° C. and lost essentially all activity (i.e., complete inactivation of the toxin) after storage for 1 month at 40° C., suggesting that N-acetyl-L-cysteine can act as a pro-oxidant in this formulation. In contrast, as shown in Tables 1.1 and 1.2, the lyophilized compositions comprising NAC lost 7% ((98.6-91.9)/98.6) potency after storage for 3 months at 25° C. and lost 12% ((98.6-86.5)/98.6) potency after storage for 1 month at 40° C. This demonstrates that NAC by itself can function as a stabilizer in the lyophilized compositions.

(63) Further studies were conducted to investigate the impact of antioxidant compounds on the liquid compositions. In Example 3, liquid compositions were prepared with 100 U/mL botulinum toxin, 8 w/w % trehalose, and 4 w/w % poloxamer P188 in histidine buffer at pH 6.0. Each formulation had a different antioxidant or a combination of antioxidants, as set forth in Table 5 below and in Example 3. The antioxidants tested included NAC, L-methionine, L-tryptophan, L-glutathione, sodium sulfite, propyl gallate, and EDTA sodium salt. Potency of the formulations was tested by a cell based potency assay after filling (time zero, t0) and after storage at 40° C. for 2 weeks and 1 month. Potency test results are shown in Table 5.

(64) TABLE-US-00006 TABLE 5 Liquid formulations Antioxidant.sup.2 Potency U/mL NAC Met TRP GSH NaSul PrpGal EDTA 2 wks 40° 1 mo. 40° Formulation No..sup.1 % % % % % % % T0 C. C. Formulation 20 0.2 126 129 130 Formulation 21 0.2 127 13.71 .sup. NT.sup.3 Formulation 22 0.2 123 3.76 NT Formulation 23 0.2 23.7 0.161 NT Formulation 24 0.2 0.164 0.150 NT Formulation 25 0.2 0.2 133 0.253 NT Formulation 26 0.2 0.03 129 127 127 Formulation 27 0.2 0.2 0.03 129 125 122 Formulation 28 0.2 0.2 126 2.45 NT .sup.1Each formulation contained 100 U/mL botulinum toxin, 8 w/w % trehalose, and 4 w/w % poloxamer P188 in 20 mM histidine buffer, pH

6.0 and the specified antioxidant. .sup.2NAC = N-acetyl-L-cysteine; Met = L-methionine; TRP = L-tryptophan; GSH = L-glutathione; NaSul = sodium sulfite; PrpGal = propyl gallate; EDTA = ethylene diamine tetraacetic acid, sodium salt. .sup.3NT = not tested

(65) The stabilizing effect of the antioxidant, or lack of stabilizing effect for certain antioxidants, is apparent from this study. Compositions comprising methionine (Formulation 20) retained full potency after storage at 40° C. for 2 weeks and 1 month, in contrast to compositions comprising tryptophan (Formulation 21) or glutathione (Formulation 22), indicating that not all amino acids function to stabilize a Clostridial toxin active ingredient in a liquid composition. Compositions comprising sodium sulfite (Formulation 23) or propyl gallate (Formulation 24) lost essentially all potency after 2 weeks storage at 40° C. Liquid compositions comprising NAC and EDTA sodium salt (Formulation 26) retained full potency after storage at 40° C. for 2 weeks and 1 month, as did a composition comprising NAC, EDTA sodium salt and tryptophan (Formulation 27). In contrast, as shown in Table 4, liquid compositions comprising NAC but not EDTA (Formulation 25) lost essentially all potency after 2 weeks storage at 40° C. This demonstrates that the combination of antioxidants—a chelating agent (e.g., EDTA, EGTA, DTPA or analogs thereof) and/or a chain terminator antioxidant (e.g., methionine, cysteine, NAC or BHT) provides a stabilizing effect on the botulinum toxin. Accordingly, compositions in liquid or solution form comprising a Clostridial toxin active ingredient; a disaccharide; a surfactant; and an antioxidant selected from methionine, NAC, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof, are contemplated. In one embodiment, the composition retains potency of the toxin for a period of at least about 2 weeks when stored at about 40° C., where the potency is measured in a cell-based potency assay. In another embodiment, the composition retains potency of the toxin for a period of at least about 1 month when stored at about 40° C., where the potency is measured in a cell-based potency assay. In a further embodiment, the composition is in liquid or solution form comprising a Clostridial toxin active ingredient; a disaccharide; a surfactant; and an antioxidant selected from (i) methionine, and (ii) NAC and a chelating agent selected from EDTA, EGTA, DTPA, and analogs thereof. In one embodiment, the composition retains potency of the toxin for a period of at least about 1 month when stored at about 40° C., where the potency is measured in a cell-based potency assay.

(66) In another embodiment, a liquid composition comprises a Clostridial toxin active ingredient; a tonicity agent; a surfactant; and an antioxidant selected from the group consisting of a sacrificial antioxidant, a chelating agent antioxidant, a chain terminator antioxidant, and combinations thereof.

(67) Example 4 details another study designed to investigate the effect of a tonicity agent on stability of the toxin in the composition. In this study liquid compositions were prepared with 100 U/mL botulinum toxin, 4 w/w % poloxamer P188, 0.2 w/w % methionine, and either 8 w/w % trehalose or sucrose, in histidine buffer at pH 6.0. The two compositions were stored at 25° C. and potency was tested by a cell based potency assay prior to storage (time zero, t0) and after storage. Potency test results are shown in Table 6.

(68) TABLE-US-00007 TABLE 6 Liquid formulations Potency, U/mL at T0 and after indicated time (in months) at 25° C. Formulation No.* Disaccharide T0 1 mo 5.5 mo 6 mo 8 mo 10.5 mo
Formulation 30 trehalose (8 w/w %) 133 145 — 147 159 43
Formulation 31 sucrose (8 w/w %) 117 — 10 — — —
*Each formulation contained 100 U/mL botulinum toxin, 4 w/w % poloxamer P188 and 0.2 w/w % methionine in histidine buffer.

(69) Comparing the potency of the toxin in Formulation 30 after 6 months of storage at 25° C. with that in Formulation 31 after 5.5 months of storage at 25° C. it is apparent that trehalose affords a stabilizing effect in a composition with poloxamer and methionine, as Formulation 30 with trehalose retained 110% of its potency (147 U/mL/133 U/mL) after 6 months storage at 25° C., whereas Formulation 31 with sucrose had only 8.5% of the target toxin potency (10 U/mL/117 U/mL) after 5.5 months storage at 25° C. Accordingly, in one embodiment, a liquid composition comprised of a Clostridial toxin active ingredient, such as a botulinum toxin; trehalose; a

poloxamer surfactant; and methionine is contemplated. In one embodiment, the composition retains at least about 80%, 85%, 90% or 95% potency after storage for 1, 2, 3, 4, 5, or 6 months at 25° C., where potency is measured in a cell-based potency assay. In an alternative embodiment, the composition retains at least about 80%, 85%, 90% or 95% potency after storage for 1, 2, 3, 4, 5, or 6 months at 25° C., where potency is measured using an in vivo LD.sub.50 potency assay.

(70) Further studies were conducted on liquid compositions as described in Example 5. In this study, compositions comprising a Clostridial toxin active ingredient, poloxamer P188 (4 w/w %) or polysorbate (TWEEN® 20, 0.04 w/w %), trehalose (8% w/w), and methionine (0.2 w/w %), in 20 mM histidine buffer at pH 6.0, were prepared. The composition with poloxamer P188 was identified as Formulation No. 30 and the composition with polysorbate was identified as Formulation No. 32. Potency was tested by a cell based potency assay prior to storage (time zero, t0) and after 1 month of storage 40° C. for 1 month. Potency test results are shown in Table 7.

(71) TABLE-US-00008 TABLE 7 Liquid formulations Potency, U/mL at time zero Formulation and after 1 month storage at 40° C. No.* Surfactant T0 1 mo. Formulation 30 poloxamer P188 133 138 (4 w/w %) Formulation 32 polysorbate 114 9 (TWEEN® 20, 0.04 w/w %) *Each formulation contained 100 U/mL botulinum toxin, 8 w/w % trehalose, and 0.2 w/w % methionine in histidine buffer.

(72) Comparing the potency of the toxin in Formulation 30 after 1 month of storage at 40° C. with that in Formulation 32 it is apparent that a poloxamer surfactant affords a stabilizing effect in a composition with trehalose and methionine, as Formulation 30 with poloxamer P188 retained about 104% of its initial potency (138 U/mL/133 U/mL) after one month of storage at 40° C., whereas Formulation 32 with polysorbate had only about 8% of the initial toxin potency (9 U/mL/114 U/mL) after one month of storage at 40° C. Accordingly, in one embodiment, a liquid composition comprised of a Clostridial toxin active ingredient, such as a botulinum toxin; trehalose; a poloxamer surfactant; and methionine is provided. In one embodiment, the composition retains at least about 80%, 85%, or 90% potency after storage for 1 month at 40° C., where potency is measured in a cell-based potency assay.

(73) In another study, five liquid compositions were prepared with botulinum toxin as a model Clostridial toxin active ingredient. The compositions were prepared with or without poloxamer surfactant, with or without trehalose, and with or without methionine. Details of the compositions are given in Example 6. Potency of Formulations 33-37 was tested by a cell based potency assay prior to storage (time zero, t0) and after 1 month of storage at -70° C. and 40° C. for 1 month. Potency test results are shown in Table 8.

(74) TABLE-US-00009 TABLE 8 Liquid formulations Composition Components* Poloxamer Potency U/mL Toxin Trehalose P188 Methionine 1 mo. -70° 1 mo. 40° Formulation No. U/mL % % % T0 C. C. Formulation 33 100 8 4 0 128 135 0.225 Formulation 34 150 0 0 0.2 52 13 0 Formulation 35 150 8 0 0.2 131 145 30 Formulation 36 100 0 4 0.2 128 142 103 Formulation 37 100 8 4 0.2 126 — 130 *Each formulation was in 20 mM histidine buffer at pH 6.0.

(75) The potency of the botulinum toxin in Formulation 36 and in Formulation 37 after 1 month of storage at -70° C. or at 40° C. reveals that liquid compositions with poloxamer and methionine, with or without a disaccharide such as trehalose, act to stabilize the toxin. This demonstrates that poloxamer and methionine can stabilize the botulinum toxin in liquid compositions without a disaccharide. The results from Formulation 33 suggest that methionine is a stabilizer, as Formulation 33 had little toxin potency after storage in contrast to Formulation 37 and Formulation 36. The results from Formulations 34 and 35 suggest that a surfactant also has a stabilizing effect on the toxin. Accordingly, in one embodiment, a liquid composition comprised of a Clostridial toxin active ingredient, such as a botulinum toxin; a poloxamer surfactant; and an antioxidant such as methionine is contemplated. In one embodiment, the composition retains at least about 70%, 75%, or 85%, potency after storage for 1 month at 40° C., where potency is measured in a cell-based potency assay. In another embodiment, a disaccharide is also included. In one embodiment,

the composition retains at least about 80%, 85% or 90% potency after storage for 1 month at 40° C., where potency is measured in a cell-based potency assay. In an alternative embodiment, the composition retains at least about 80%, 85% or 90% potency after storage for 1 month at 40° C., where potency is measured using an in vivo LD.sub.50 potency assay.

(76) In another study, described in Example 7, liquid compositions were prepared with botulinum toxin as a model Clostridial toxin active ingredient. The compositions were prepared with a poloxamer surfactant or with a polysorbate surfactant, with or without a disaccharide. All formulations comprised methionine. Details of the compositions are given in Example 7. Potency of the compositions, identified as Formulations 31, 38, 39 and 40, was tested by a cell based potency assay prior to storage (time zero, t0) and after storage at 5° C. for 5.5 months, 7.5 months and 12 months. Potency test results are shown in Table 9.

(77) TABLE-US-00010 TABLE 9 Liquid formulations Potency U/mL after storage for Composition Components.sup.1 indicated time TWEEN® Buffer.sup.2 (in months) at 5° C. Treh Suc NaCl P188 20 Met or 5.5 7.5 12 Formulation No. % % % % % % water T0 mo mo mo
 Formulation 31 8 4 0.2 buffer 117 88 NT 94 Formulation 38 8 4 0.2 water 114 .sup. NT.sup.3 108
 91 Formulation 39 0.9 4 0.2 buffer 94 NT 80 62 Formulation 40 8 0.04 0.2 buffer 114 NT 117 91
 .sup.1All formulations contained 100 U/mL botulinum toxin; Treh = trehalose; Suc = sucrose; P-188 = poloxamer P188; TWEEN® 20 = polysorbate; Met = L-methionine; .sup.2Buffer = 20 mM His, pH 6.0; .sup.3NT = not tested

(78) The potency of the botulinum toxin in Formulation 31 and in Formulation 38 after 12 months of storage at 5° C. reveals that the histidine buffer is not acting to stabilize the composition, as Formulation 38 was prepared with water and was stabilized to the same extent by the surfactant, disaccharide and methionine as Formulation 31 which was prepared with histidine buffer. Formulation 40 prepared with trehalose, polysorbate and methionine successfully stabilized the botulinum toxin as evident from the 91 U/mL potency 12 months of storage at 5° C. Accordingly, in one embodiment, a liquid composition comprised of a Clostridial toxin active ingredient, such as a botulinum toxin; a poloxamer or polysorbate surfactant; trehalose or sucrose; and an antioxidant such as methionine is contemplated. In one embodiment, the composition retains at least about 75%, 80%, or 85%, potency after storage for 1, 3, 6, 9, or 12 months at 5° C., where potency is measured in a cell-based potency assay. In an alternative embodiment, the composition retains at least about 75%, 80%, or 85%, potency after storage for 1, 3, 6, 9, or 12 months at 5° C., where potency is measured using an in vivo LD.sub.50 potency assay.

(79) In another study, described in Example 8, liquid compositions with no tonicity agent were prepared. Botulinum toxin was used as a model Clostridial toxin active ingredient. The compositions were prepared with either a poloxamer surfactant or with a polysorbate surfactant, and with methionine as the antioxidant. Details of the compositions are given in Example 8. Potency of the compositions, identified as Formulations 41 and 42, was tested by a cell based potency assay prior to storage (time zero, t0) and after storage for two weeks at -70° C. and at 40° C. Potency test results are shown in Table 10.

(80) TABLE-US-00011 TABLE 10 Liquid formulations Composition Components* Potency U/mL Polysorbate after storage Toxin P188 (TWEEN®20) methionine for 2 weeks Formulation No. U/mL % % % -70° C. 40° C. Formulation 41 100 0 0.04 0.2 25 1 Formulation 42 100 4 0 0.2 105 72 *Both formulations were in 20 mM histidine buffer at pH 6.0. P-188 = poloxamer P188; Met = L-methionine

(81) The potency of the botulinum toxin in Formulation 41 and in Formulation 42 after 2 weeks of storage at -70° C. and at 40° C. reveals higher potency recoveries were achieved in liquid compositions comprising poloxamer as the surfactant, as compared to liquid compositions comprising polysorbate as the surfactant.

(82) In another study, liquid compositions were prepared comprising a Clostridial toxin active ingredient; poloxamer P188 (4% w/w or 0.6% w/w); trehalose (2% w/w or 8% w/w); and an

antioxidant—(i) EDTA and NAC (0.03% w/w and 0.2% w/w, respectively), or (ii) methionine (0.2% w/w); in 20 mM histidine buffer at pH 6.0. Each formulation had between 30-200U botulinum toxin per vial. A summary of the compositions is set forth in Table 11.

(83) TABLE-US-00012 TABLE 11 Liquid formulations

	Tonicity Agent	Trehalose	Surfactant	or Poloxamer	Antioxidant	Toxin	Sucrose	NaCl	P188	Methionine	EDTA	NAC	U/mL	wt %	wt %	wt %
wt %	wt %	wt %	wt %	wt %	wt %	wt %	wt %	wt %	wt %	wt %	wt %	wt %	wt %	wt %	wt %	wt %
30-200	8	4	0.2	20 mM His, pH 6.0	30-200	8	0.6	0.2	20 mM His, pH 6.0	30-200	2	0.6	4	0.2	20 mM His, pH 6.0	30-200
8	4	0.03	0.2	20 mM His, pH 6.0	30-200	8	4	0.2	0.03	0.2	20 mM His, pH 6.0	30-200	8	4	0.2	0.03

(84) Accordingly, in one embodiment, a liquid pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; a tonicity agent selected from trehalose, sucrose, sodium chloride, mannitol, sorbitol, glucose and combinations thereof; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; and an antioxidant selected from methionine, NAC, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof. In one embodiment, the composition excludes albumin, and in another embodiment, when the antioxidant is methionine the composition excludes a polysorbate.

(85) In another embodiment, a liquid pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; and an antioxidant selected from methionine, NAC, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof. In one embodiment, the composition excludes albumin. In one embodiment, the composition excludes a tonicity agent.

(86) In another embodiment, a liquid pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; a tonicity agent selected from trehalose, sucrose, sodium chloride, mannitol, sorbitol, glucose, and combinations thereof; a poloxamer; and methionine. In one embodiment, the composition excludes albumin.

(87) In another embodiment, a liquid pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; a disaccharide tonicity agent in an amount between 1-15 wt %; a poloxamer in an amount between 0.5-8 wt %; and an antioxidant in an amount between 0.05-5 wt %. In another embodiment, a composition comprises a Clostridial toxin active ingredient; trehalose in an amount between 1-15 wt %; a poloxamer in an amount between 0.5-8 wt %; and methionine in an amount between 0.05-5 wt %. In one embodiment, the composition excludes albumin. In another embodiment, a liquid pharmaceutical composition comprises a Clostridial toxin active ingredient; trehalose in an amount between 2-15 wt % or 1-10 wt %; a poloxamer in an amount between 0.5-8 wt %; and methionine in an amount between 0.05-5 wt %. In one embodiment, the composition excludes albumin.

(88) In another embodiment, a liquid pharmaceutical composition is contemplated. The composition comprises a botulinum toxin; a tonicity agent selected from trehalose, sucrose, sodium chloride, mannitol, sorbitol, glucose, and combinations thereof; a poloxamer; and an antioxidant selected from methionine, NAC, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof. In one embodiment, the composition excludes albumin.

(89) In any of the foregoing embodiments, it is contemplated that the composition is not, in some embodiments, an emulsion and/or excludes nanoparticles comprising an amphiphilic entity.

(90) Lyophilized compositions were prepared for a study described in Example 9. The compositions that were prepared comprised botulinum toxin as a model Clostridial toxin; a disaccharide; a surfactant; and methionine as an antioxidant. The formulations were lyophilized and stored for two weeks at -20° C. or at 40° C. Potency of the lyophilized formulations after storage and reconstitution with saline was tested by a cell based potency assay. The potencies of the solid compositions are shown in Table 12.

(91) TABLE-US-00013 TABLE 12 Lyophilized formulations

Composition	Components*	TWEEN®	Poloxamer	Potency	U/mL	Trehalose	Sucrose	20 P188	Methionine	2 weeks -20°	2 weeks 40°						
Formulation No.	%	%	%	%	%	C.	C.	Formulation	43	8	0	0	4	0.2	166	162	
0.2	147	137	Formulation	45	0	8	0	4	0.2	149	134	Formulation	46	0	8	0.04	0
0.2	147	137	Formulation	45	0	8	0	4	0.2	149	134	Formulation	46	0	8	0.04	0

*Each

formulation comprised 200 U/vial BoNT/A and was in 20 mM histidine buffer at pH 6.0.

(92) A comparison of the potency data for lyophilized Formulation 43 and Formulation 45 permits analysis of the impact of trehalose versus sucrose as the lyoprotector on stability of botulinum toxin in lyophilized compositions, with a poloxamer as the surfactant, and methionine as an antioxidant. Formulations 43 and 45 differ only in that the former has trehalose and the latter has sucrose; both formulations comprised a poloxamer and methionine. The potency of the botulinum toxin in Formulation 43 and in Formulation 45 after 2 weeks of storage at -20°C . and at 40°C . reveals a higher potency recovery was achieved in lyophilized compositions comprising trehalose, as compared to lyophilized compositions comprising sucrose. Accordingly, in one embodiment, a lyophilized composition is provided, where the composition comprises a Clostridial toxin active ingredient, a poloxamer, methionine, and trehalose. In one embodiment, the toxin has a potency recovery of at least about 75% or at least about 80% after storage at two weeks at -20°C . and/or at 40°C .

(93) A comparison of the potency data for Formulation 43 and Formulation 44 permits analysis of polysorbate and poloxamer as surfactants on stability of botulinum toxin in lyophilized compositions, with trehalose as a lyoprotector, and methionine as an antioxidant. Formulations 43 and 44 differ only in that the former has a poloxamer surfactant and the latter has polysorbate as the surfactant. The potency of the botulinum toxin in Formulation 43 and in Formulation 44 after 2 weeks of storage at -20°C . and at 40°C . reveals a higher potency recovery was achieved in lyophilized compositions comprising a poloxamer surfactant, as compared to lyophilized compositions comprising a polysorbate surfactant, when trehalose and methionine are present. Accordingly, in one embodiment, a lyophilized composition is provided, where the composition comprises a Clostridial toxin active ingredient, a poloxamer, methionine, and trehalose. In one embodiment, the toxin has a potency recovery of at least about 75% or at least about 80% after storage at two weeks at -20°C . and/or at 40°C .

(94) Accordingly, in one embodiment, a solid or lyophilized pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; a tonicity agent and/or lyoprotector selected from trehalose, sucrose, mannitol, sorbitol, glucose, and combinations thereof; a surfactant selected from a poloxamer, a polysorbate and combinations thereof; and an antioxidant selected from methionine, NAC, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof. In some embodiments, the solid composition comprises a lyoprotector. In some embodiments, the lyoprotector includes sucrose, trehalose, mannitol, sorbitol, glucose, or combinations thereof. In certain embodiments, the lyophilized composition is reconstituted with a tonicity agent selected from trehalose, sucrose, sodium chloride, mannitol, sorbitol, glucose, and combinations thereof. In at least one embodiment, the lyophilized composition is reconstituted with a reconstitution vehicle comprising NaCl prior to administration to a patient. In at least one embodiment, NaCl is present in an amount of 0.9% (w/w) in the reconstitution vehicle. In one embodiment, the composition excludes albumin, a hydroxyalkyl starch, glutamic acid, glutamine, aspartic acid, asparagine, a polyalcohol, glycine, and/or polyvinylpyrrolidone.

(95) In another embodiment, a solid or lyophilized pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; trehalose or sucrose; a poloxamer; and methionine. In one embodiment, the composition excludes albumin, a hydroxyalkyl starch, glutamic acid, glutamine, aspartic acid, asparagine, a polyalcohol, glycine, and/or polyvinylpyrrolidone. In one embodiment, the lyophilized pharmaceutical composition comprises botulinum toxin as the Clostridial toxin active ingredient, trehalose, a poloxamer, and methionine.

(96) In another embodiment, a solid or lyophilized pharmaceutical composition is contemplated. The composition comprises a Clostridial toxin active ingredient; trehalose in an amount between 1-15 wt %; a poloxamer in an amount between 0.5-8 wt %; and methionine in an amount between 0.05-5 wt %. In another embodiment, the composition comprises a Clostridial toxin active ingredient, such as botulinum toxin; 8 wt % trehalose; 4 wt % poloxamer; and 0.2 wt %

methionine. In one embodiment, the botulinum toxin is present in an amount of about 200 units. In another embodiment, the botulinum toxin is present in an amount of about 50 units. In one embodiment, the composition excludes albumin, a hydroxyalkyl starch, glutamic acid, glutamine, aspartic acid, asparagine, a polyalcohol, glycine, and/or polyvinylpyrrolidone.

(97) In another embodiment, a solid or lyophilized pharmaceutical composition is contemplated. The composition comprises a botulinum toxin; a disaccharide; a poloxamer; and an antioxidant selected from methionine, NAC, EDTA, EGTA, DTPA, analogs thereof, and combinations thereof. In one embodiment, the composition excludes albumin, a hydroxyalkyl starch, glutamic acid, glutamine, aspartic acid, asparagine, a polyalcohol, glycine, and/or polyvinylpyrrolidone.

(98) In another embodiment, a lyophilized composition comprises a Clostridial toxin active ingredient; a tonicity agent and/or a lyoprotector; a surfactant; and an antioxidant selected from the group consisting of a sacrificial antioxidant, a chelating agent antioxidant, a chain terminator antioxidant, and combinations thereof. In some embodiments, the lyophilized Clostridial pharmaceutical composition comprises a lyoprotector. In some embodiments, the lyoprotector includes sucrose, trehalose, mannitol, sorbitol, glucose, or combinations thereof.

(99) In any of the foregoing embodiments of solid or liquid compositions, it is contemplated that one or more, in any combination, of these ingredients are excluded: polyvinylpyrrolidone, diblock copolymers of polypropylene glycol and polyethylene glycol, and/or a polyalcohol such as inositol, lactitol, isomalt, xylitol, or erythritol.

(100) Pharmaceutical Composition Components

(101) The present pharmaceutical compositions include a Clostridial toxin or a Clostridial toxin active ingredient. A skilled artisan will appreciate that the description herein refers to a Clostridial toxin active ingredient, however, a Clostridial toxin may also be used in the compositions described herein. Accordingly, the term Clostridial toxin active ingredient will be used; however it should be understood that a Clostridial toxin is equally contemplated. In one embodiment, a therapeutically effective concentration of a Clostridial toxin active ingredient is present in the composition. In one embodiment, the Clostridial toxin active ingredient reduces a symptom associated with the aliment being treated by, e.g., at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90% or at least 100%. In other aspects of this embodiment, a therapeutically effective concentration of a Clostridial toxin active ingredient reduces a symptom associated with the aliment being treated by, e.g., at most 10%, at most 20%, at most 30%, at most 40%, at most 50%, at most 60%, at most 70%, at most 80%, at most 90% or at most 100%.

(102) It is envisioned that any amount of Clostridial toxin active ingredient can be added in formulating a Clostridial toxin active ingredient pharmaceutical compositions disclosed herein, with the proviso that a therapeutically effective amount of Clostridial toxin active ingredient is recoverable. In aspects of this embodiment, the amount of Clostridial toxin active ingredient added to the formulation is at least 0.1 U/ml, at least 1.0 U/ml, at least 10 U/ml, at least 50 U/ml, at least 100 U/ml, at least 200 U/ml, or at least 1000 U/ml. In other aspects of this embodiment, the amount of Clostridial toxin active ingredient added to the formulation is at most 0.1 U/ml, at most 1.0 U/ml, at most 10 U/ml, at most 50 U/ml, at most 100 U/ml, at most 200 U/ml, or at most 1000 U/ml. In yet other aspects of this embodiment, the amount of Clostridial toxin active ingredient added to the formulation is from about 0.1 U/ml to about 1000 U/ml, or about 1.0 U/ml to about 1000 U/ml. In still other aspects of this embodiment, the amount of Clostridial toxin active ingredient added to the formulation is from about 0.001 U/ml to about 100 U/ml, about 0.01 U/ml to about 100 U/ml, about 0.1 U/ml to about 100 U/ml, or about 1.0 U/ml to about 100 U/ml.

(103) In other aspects of this embodiment, the amount of Clostridial toxin active ingredient added to the formulation is at least 1.0 pg, at least 10 pg, at least 100 pg, at least 1.0 ng, at least 10 ng, at least 100 ng, at least 1.0 μ g, at least 10 μ g, at least 100 μ g, or at least 1.0 mg. In still other aspects of this embodiment, the amount of Clostridial toxin active ingredient added to the formulation is at most 1.0 pg, at most 10 pg, at most 100 pg, at most 1.0 ng, at most 10 ng, at most 100 ng, at most

1.0 µg, at most 10 µg, at most 100 µg, or at most 1.0 mg. In still other aspects of this embodiment, the amount of Clostridial toxin active ingredient added to the formulation is about 1.0 pg to about 10 µg, about 10 pg to about 10 µg, about 100 pg to about 10 µg, about 1.0 ng to about 10 µg, about 10 ng to about 10 µg, or about 100 ng to about 10 µg. In still other aspects of this embodiment, the amount of Clostridial toxin active ingredient added to the formulation is about 1.0 pg to about 1.0 µg, about 10 pg to about 1.0 µg, about 100 pg to about 1.0 µg, about 1.0 ng to about 1.0 µg, about 10 ng to about 1.0 µg, or about 100 ng to about 1.0 µg. In further aspects of this embodiment, the amount of Clostridial toxin active ingredient added to the formulation is about 1.0 pg to about 5.0 µg, about 10 pg to about 5.0 µg, about 100 pg to about 5.0 µg, about 1.0 ng to about 5.0 µg, about 10 ng to about 5.0 µg, or about 100 ng to about 5.0 µg. In further aspects of this embodiment, the amount of Clostridial toxin active ingredient added to the formulation is about 1.0 pg to about 10 µg, about 10 pg to about 10 µg, about 100 pg to about 10 µg, about 1.0 ng to about 10 µg, about 10 ng to about 10 µg, or about 100 ng to about 10 µg.

(104) In aspects of this embodiment, a Clostridial toxin pharmaceutical composition comprises a BoNT/A, a BoNT/B, a BoNT/C.sub.1, a BoNT/D, a BoNT/E, a BoNT/F, a BoNT/G, a mosaic BONT such as for example BoNT/DC, a TeNT, a BaNT, or a BuNT. In another embodiment, a Clostridial toxin pharmaceutical composition comprises a Clostridial toxin variant as the Clostridial toxin. In aspects of this embodiment, a Clostridial toxin pharmaceutical composition comprises naturally-occurring Clostridial toxin active ingredient variant or a non-naturally-occurring Clostridial toxin variant. In other aspects of this embodiment, a Clostridial toxin pharmaceutical composition comprises a BoNT/A variant, a BoNT/B variant, a BoNT/C.sub.1 variant, a BoNT/D variant, a BoNT/E variant, a BoNT/F variant, a BoNT/G variant, a TeNT variant, a BaNT variant, or a BuNT variant, where the variant is either a naturally-occurring variant or a non-naturally-occurring variant.

(105) Aspects of the present pharmaceutical compositions provide, in part, a Clostridial toxin complex as a Clostridial toxin active ingredient. As used herein, the term “Clostridial toxin complex” refers to a complex comprising a Clostridial toxin and associated NAPs, such as, e.g., a Botulinum toxin complex, a Tetanus toxin complex, a *Baratii* toxin complex, and a *Butyricum* toxin complex. Non-limiting examples of Clostridial toxin complexes include those produced by a *Clostridium botulinum*, such as, e.g., a 900-kDa BoNT/A complex, a 500-kDa BoNT/A complex, a 300-kDa BoNT/A complex, a 500-kDa BoNT/B complex, a 500-kDa BoNT/C.sub.1 complex, a 500-kDa BoNT/D complex, a 300-kDa BoNT/D complex, a 300-kDa BoNT/E complex, and a 300-kDa BoNT/F complex. Clostridial toxin complexes can be purified using the methods described in Schantz, supra, (1992); Hui Xiang et al., *Animal Product Free System and Process for Purifying a Botulinum Toxin*, U.S. Pat. No. 7,354,740, each of which is hereby incorporated by reference in its entirety. Clostridial toxin complexes can be obtained from, e.g., List Biological Laboratories, Inc. (Campbell, California), the Centre for Applied Microbiology and Research (Porton Down, U.K), Wako (Osaka, Japan), and Sigma Chemicals (St Louis, Missouri).

(106) Aspects of the present pharmaceutical compositions provide, in part, a non-protein excipient. As used herein, the term “non-protein excipient” refers to any excipient that is not a polypeptide comprising at least fifteen amino acids. It is envisioned that any non-protein excipient is useful in formulating a Clostridial toxin active ingredient pharmaceutical compositions disclosed in the present specification, with the proviso that a therapeutically effective amount of the Clostridial toxin active ingredient is recovered using this non-protein excipient.

(107) Aspects of the present pharmaceutical compositions provide, in part, a sugar. As used herein, the term “sugar” refers to a compound comprising one to 10 monosaccharide units, e.g., a monosaccharide, a disaccharide, a trisaccharide, and an oligosaccharide comprising four to ten monosaccharide units. It is envisioned that any sugar is useful in formulating a Clostridial toxin active ingredient pharmaceutical compositions disclosed in the present specification, with the proviso that a therapeutically effective amount of the Clostridial toxin active ingredient is

recovered using this sugar. In some embodiments, for example in a lyophilized composition, the sugar can function as a lyoprotector. In some other embodiments, for example in a lyophilized formulation or in a liquid formulation, the sugar can function as a tonicity agent. Monosaccharides are polyhydroxy aldehydes or polyhydroxy ketones with three or more carbon atoms, including aldoses, dialdoses, aldoketoses, ketoses and diketoses, as well as cyclic forms, deoxy sugars and amino sugars, and their derivatives, provided that the parent monosaccharide has a (potential) carbonyl group. Monosaccharides include trioses, like glyceraldehyde and dihydroxyacetone; tetroses, like erythrose, erythrulose and threose; pentoses, like arabinose, lyxose, ribose, ribulose, xylose, xylulose; hexoses, like allose, altrose, fructose, fucose, galactose, glucose, gulose, idose, mannose, psicose, rhamnose, sorbose, tagatose, talose and trehalose; heptoses, like sedoheptulose and mannoheptulose; octoses, like octulose and 2-keto-3-deoxy-manno-octonate; nonoses like sialose; and decose. Oligosaccharides are compounds in which at least two monosaccharide units are joined by glycosidic linkages. According to the number of units, they are called disaccharides, trisaccharides, tetrasaccharides, pentasaccharides, hexasaccharides, heptosaccharides, octosaccharides, monosaccharides, decosaccharides, etc. An oligosaccharide can be unbranched, branched or cyclic. Common disaccharides include, without limitation, sucrose, lactose, maltose, trehalose, cellobiose, gentiobiose, kojibiose, laminaribiose, mannobiose, melibiose, nigerose, rutinose, and xylobiose. Common trisaccharides include, without limitation, raffinose, acarbose, maltotriose, and melezitose. Other non-limiting examples of specific uses of sugar excipients can be found in, e.g., Ansel, supra, (1999); Gennaro, supra, (2000); Hardman, supra, (2001); and Rowe, supra, (2003), each of which is hereby incorporated by reference in its entirety

(108) In an embodiment, a Clostridial toxin active ingredient pharmaceutical composition comprises a sugar. In aspects of this embodiment, a Clostridial toxin active ingredient pharmaceutical composition comprises a monosaccharide. In other aspects of this embodiment, a Clostridial toxin active ingredient pharmaceutical composition comprises a disaccharide, a trisaccharide, a tetrasaccharide, a pentasaccharide, a hexasaccharide, a heptosaccharide, an octosaccharide, a monosaccharide, or a decosaccharide. In yet other aspects of this embodiment, a Clostridial toxin active ingredient pharmaceutical composition comprises an oligosaccharide comprising two to ten monosaccharide units.

(109) It is envisioned that any amount of sugar is useful in formulating a Clostridial toxin active ingredient pharmaceutical compositions disclosed in the present specification, with the proviso that a therapeutically effective amount of the Clostridial toxin active ingredient is recovered using this sugar amount. In aspects of this embodiment, the amount of sugar added to the formulation is about 0.1% (w/w), about 0.5% (w/w), about 1.0% (w/w), about 1.5% (w/w), about 2.0% (w/w), about 2.5% (w/w), about 3.0% (w/w), about 3.5% (w/w), about 4.0% (w/w), about 4.5% (w/w), about 5.0% (w/w), about 5.5% (w/w), about 6.0% (w/w), about 6.5% (w/w), about 7.0% (w/w), about 7.5% (w/w), about 8.0% (w/w), about 8.5% (w/w), about 9.0% (w/w), about 9.5% (w/w), about 10% (w/w), about 15% (w/w), about 20% (w/w), about 25% (w/w), about 30% (w/w), or about 35% (w/w). In other aspects of this embodiment, the amount of sugar added to the formulation is at least 0.1% (w/w), at least 0.5% (w/w), at least 1.0% (w/w), at least 1.5% (w/w), at least 2.0% (w/w), at least 2.5% (w/w), at least 3.0% (w/w), at least 3.5% (w/w), at least 4.0% (w/w), at least 4.5% (w/w), at least 5.0% (w/w), at least 5.5% (w/w), at least 6.0% (w/w), at least 6.5% (w/w), at least 7.0% (w/w), at least 7.5% (w/w), at least 8.0% (w/w), at least 8.5% (w/w), at least 9.0% (w/w), at least 9.5% (w/w), at least 10% (w/w), at least 15% (w/w), at least 20% (w/w), at least 25% (w/w), at least 30% (w/w), or at least 35% (w/w). In yet other aspects of this embodiment, the amount of sugar added to the formulation is at most 0.1% (w/w), at most 0.5% (w/w), at most 1.0% (w/w), at most 1.5% (w/w), at most 2.0% (w/w), at most 2.5% (w/w), at most 3.0% (w/w), at most 3.5% (w/w), at most 4.0% (w/w), at most 4.5% (w/w), at most 5.0% (w/w), at most 5.5% (w/w), at most 6.0% (w/w), at most 6.5% (w/w), at most 7.0% (w/w), at most 7.5% (w/w), at most 8.0% (w/w), at most 8.5% (w/w), at most 9.0% (w/w), at most 9.5% (w/w), at most 10% (w/w), at most

15% (w/w), at most 20% (w/w), at most 25% (w/w), at most 30% (w/w), or at most 35% (w/w). (110) In an embodiment, the present Clostridial toxin active ingredient pharmaceutical composition comprises a disaccharide. Common disaccharides include, without limitation, sucrose, lactose, maltose, trehalose, cellobiose, gentiobiose, kojibiose, laminaribiose, mannobiose, melibiose, nigerose, rutinose, and xylobiose. In aspects of this embodiment, the Clostridial toxin active ingredient pharmaceutical composition comprises sucrose. In one specific embodiment, the Clostridial toxin active ingredient pharmaceutical composition comprises trehalose. In aspects of this embodiment, the amount of disaccharide added to the formulation added to the formulation is about 0.1% (w/w), about 0.5% (w/w), about 1.0% (w/w), about 1.5% (w/w), about 2.0% (w/w), about 2.5% (w/w), about 3.0% (w/w), about 3.5% (w/w), about 4.0% (w/w), about 4.5% (w/w), about 5.0% (w/w), about 5.5% (w/w), about 6.0% (w/w), about 6.5% (w/w), about 7.0% (w/w), about 7.5% (w/w), about 8.0% (w/w), about 8.5% (w/w), about 9.0% (w/w), about 9.5% (w/w), about 10% (w/w), about 15% (w/w), about 20% (w/w), about 25% (w/w), about 30% (w/w), or about 35% (w/w).

(111) Aspects of the present pharmaceutical compositions provide, in part, a surfactant. It is envisioned that any surfactant is useful in formulating a Clostridial toxin active ingredient pharmaceutical compositions disclosed in the present specification, with the proviso that a therapeutically effective amount of the Clostridial toxin active ingredient is recovered using this surfactant amount. Non-limiting examples of surfactants include polysorbates like polysorbate 20 (TWEEN® 20), polysorbate 40 (TWEEN® 40), polysorbate 60 (TWEEN® 60), polysorbate 61 (TWEEN® 61), polysorbate 65 (TWEEN® 65), polysorbate 80 (TWEEN® 80), and polysorbate 81 (TWEEN® 81); poloxamers (polyethylene-polypropylene copolymers), like Poloxamer 124 (PLURONIC® L44), Poloxamer 181 (PLURONIC® L61), Poloxamer 182 (PLURONIC® L62), Poloxamer 184 (PLURONIC® L64), Poloxamer 188 (PLURONIC® F68), Poloxamer 237 (PLURONIC® F87), Poloxamer 338 (PLURONIC® L108), Poloxamer 407 (PLURONIC® F127), polyoxyethyleneglycol dodecyl ethers, like BRIJ® 30, and BRIJ® 35; 2-dodecoxyethanol (LUBROL®-PX); polyoxyethylene octyl phenyl ether (TRITON® X-100); sodium dodecyl sulfate (SDS); solutol HS15; 3-[(3-Cholamidopropyl)dimethylammonio]-1-propanesulfonate (CHAPS); 3-[(3-Cholamidopropyl)dimethylammonio]-2-hydroxy-1-propanesulfonate (CHAPSO); sucrose monolaurate; and sodium cholate. Other non-limiting examples of surfactant excipients can be found in, e.g., Ansel, supra, (1999); Gennaro, supra, (2000); Hardman, supra, (2001); and Rowe, supra, (2003), each of which is hereby incorporated by reference in its entirety.

(112) Thus in an embodiment, a Clostridial toxin active ingredient pharmaceutical composition comprises a surfactant. In aspects of this embodiment, a Clostridial toxin active ingredient pharmaceutical 1 comprises a polysorbate, a poloxamer, a polyoxyethyleneglycol dodecyl ether, 2-dodecoxyethanol, polyoxyethylene octyl phenyl ether, sodium dodecyl sulfate, 3-[(3-Cholamidopropyl)dimethylammonio]-1-propanesulfonate, 3-[(3-Cholamidopropyl)dimethylammonio]-2-hydroxy-1-propanesulfonate, sucrose monolaurate; or sodium cholate.

(113) It is envisioned that any amount of surfactant is useful in formulating a Clostridial toxin active ingredient pharmaceutical compositions disclosed in the present specification, with the proviso that a therapeutically effective amount of the Clostridial toxin active ingredient is recovered using this surfactant amount. In aspects of this embodiment, the amount of surfactant added to the formulation is about 0.01% (w/w), about 0.02% (w/w), about 0.03% (w/w), about 0.04% (w/w), about 0.05% (w/w), about 0.06% (w/w), about 0.07% (w/w), about 0.08% (w/w), about 0.09% (w/w), about 0.1% (w/w), about 0.5% (w/w), about 1.0% (w/w), about 1.5% (w/w), about 2.0% (w/w), about 2.5% (w/w), about 3.0% (w/w), about 3.5% (w/w), about 4.0% (w/w), about 4.5% (w/w), about 5.0% (w/w), about 5.5% (w/w), about 6.0% (w/w), about 6.5% (w/w), about 7.0% (w/w), about 7.5% (w/w), about 8.0% (w/w), about 8.5% (w/w), about 9.0% (w/w), about 9.5% (w/w), about 10% (w/w), about 15% (w/w), about 20% (w/w), about 25% (w/w), about 30% (w/w), or about 35% (w/w). In other aspects of this embodiment, the amount of surfactant

added to the formulation is at least 0.01% (w/w), at least 0.02% (w/w), at least 0.03% (w/w), at least 0.04% (w/w), at least 0.05% (w/w), at least 0.06% (w/w), at least 0.07% (w/w), at least 0.08% (w/w), at least 0.09% (w/w), at least 0.1% (w/w), at least 0.5% (w/w), at least 1.0% (w/w), at least 1.5% (w/w), at least 2.0% (w/w), at least 2.5% (w/w), at least 3.0% (w/w), at least 3.5% (w/w), at least 4.0% (w/w), at least 4.5% (w/w), at least 5.0% (w/w), at least 5.5% (w/w), at least 6.0% (w/w), at least 6.5% (w/w), at least 7.0% (w/w), at least 7.5% (w/w), at least 8.0% (w/w), at least 8.5% (w/w), at least 9.0% (w/w), at least 9.5% (w/w), at least 10% (w/w), at least 15% (w/w), at least 20% (w/w), at least 25% (w/w), at least 30% (w/w), or at least 35% (w/w). In yet other aspects of this embodiment, the amount of surfactant added to the formulation is at most 0.01% (w/w), at most 0.02% (w/w), at most 0.03% (w/w), at most 0.04% (w/w), at most 0.05% (w/w), at most 0.06% (w/w), at most 0.07% (w/w), at most 0.08% (w/w), at most 0.09% (w/w), at most 0.1% (w/w), at most 0.5% (w/w), at most 1.0% (w/w), at most 1.5% (w/w), at most 2.0% (w/w), at most 2.5% (w/w), at most 3.0% (w/w), at most 3.5% (w/w), at most 4.0% (w/w), at most 4.5% (w/w), at most 5.0% (w/w), at most 5.5% (w/w), at most 6.0% (w/w), at most 6.5% (w/w), at most 7.0% (w/w), at most 7.5% (w/w), at most 8.0% (w/w), at most 8.5% (w/w), at most 9.0% (w/w), at most 9.5% (w/w), at most 10% (w/w), at most 15% (w/w), at most 20% (w/w), at most 25% (w/w), at most 30% (w/w), or at most 35% (w/w).

(114) In some embodiments, the Clostridial toxin active ingredient pharmaceutical composition comprises a poloxamer. Poloxamers which can be used with the present pharmaceutical composition include Poloxamer 124 (PLURONIC® L44), Poloxamer 181 (PLURONIC® L61), Poloxamer 182 (PLURONIC® L62), Poloxamer 184 (PLURONIC® L64), Poloxamer 188 (e.g., PLURONIC® F68, KOLLIPHOR® P 188), Poloxamer 237 (PLURONIC® F87), Poloxamer 338 (PLURONIC® L108), Poloxamer 407 (PLURONIC® F127). In some embodiments, poloxamer 188 may be more advantageous.

(115) In some embodiments, the Clostridial toxin active ingredient pharmaceutical composition comprises a polysorbate. Polysorbates which can be used with the present pharmaceutical composition includes polysorbate 20 (TWEEN® 20), polysorbate 40 (TWEEN® 40), polysorbate 60 (TWEEN® 60), polysorbate 61 (TWEEN® 61), polysorbate 65 (TWEEN® 65), polysorbate 80 (TWEEN® 80), and polysorbate 81 (TWEEN® 81). In some embodiments, polysorbate 20 may be more advantageous than some other polysorbates.

(116) Aspects of the present pharmaceutical compositions provide, in part, at least an antioxidant. Non-limiting examples of antioxidant include, without limitation, methionine, cysteine, NAC, glutathione, lipoic acid, sodium metabisulfite, sodium thiosulfate, ascorbic acid, butylated hydroxyanisole, butylated hydroxytoluene, vitamin E and analogs including Trolox C; chelators such as EDTA (ethylene diamine tetraacetic acid sodium salt), EGTA (ethylene glycol-bis(2-aminoethylether)-N,N,N',N'-tetraacetic acid), DTPA (diethylenetriaminepentaacetic acid), analogs or derivatives thereof; and combinations thereof. In aspects of this embodiment, the amount of antioxidant added to the formulation ranges from about 0.01% (w/w) to about 0.10% (w/w).

(117) It is further envisioned that a Clostridial toxin active ingredient pharmaceutical composition disclosed in the present specification can optionally include, without limitation, other pharmaceutically acceptable components (or pharmaceutical components), including, without limitation, buffers, preservatives, tonicity adjusters, salts, antioxidants, osmolality adjusting agents, emulsifying agents, sweetening or flavoring agents, and the like. Various buffers and means for adjusting pH can be used to prepare a pharmaceutical composition disclosed in the present specification, provided that the resulting preparation is pharmaceutically acceptable. Such buffers include, without limitation, acetate buffers, borate buffers, citrate buffers, phosphate buffers, neutral buffered saline, and phosphate buffered saline. It is understood that acids or bases can be used to adjust the pH of a pharmaceutical composition as needed. It is envisioned that any buffered pH level can be useful in formulating a Clostridial toxin active ingredient pharmaceutical composition, with the proviso that a therapeutically effective amount of the Clostridial toxin active

ingredient is recovered using this effective pH level. In an aspect of this embodiment, an effective pH level is at least about pH 5.0, at least about pH 5.5, at least about pH 6.0, at least about pH 6.5, at least about pH 7.0 or at about pH 7.5. In another aspect of this embodiment, an effective pH level is at most about pH 5.0, at most about pH 5.5, at most about pH 6.0, at most about pH 6.5, at most about pH 7.0 or at most about pH 7.5. In yet another aspect of this embodiment, an effective pH level is about pH 5.0 to about pH 8.0, an effective pH level is about pH 5.0 to about pH 7.0, an effective pH level is about pH 5.0 to about pH 6.0, is about pH 5.5 to about pH 8.0, an effective pH level is about pH 5.5 to about pH 7.0, an effective pH level is about pH 5.5 to about pH 5.0, is about pH 5.5 to about pH 7.5, an effective pH level is about pH 5.5 to about pH 6.5.

(118) The pharmaceutical compositions disclosed herein can have a pH of between about 5 and 8 when reconstituted or upon injection. In certain embodiments, the composition will have a pH below 8, such as, for example, 7.9, or 7.8, or 7.7, or 7.6, or 7.5, or 7.4, or 7.3, or 7.2, or 7.1, or 7.0, or 6.9, or 6.8, or 6.7, or 6.6, or 6.5, or 6.4, or 6.3, or 6.2, or 6.1, or 6.0, or 5.9, or 5.8, or 5.7, or 5.6, or 5.5, or 5.4, or 5.3, or 5.2, or 5.1, or the like. In some embodiments, the pH ranges from 5 to 7.

(119) It is envisioned that any concentration of a buffer can be useful in formulating a Clostridial toxin active ingredient pharmaceutical composition, with the proviso that a therapeutically effective amount of the Clostridial toxin active ingredient is recovered using this effective concentration of buffer. In aspects of this embodiment, an effective concentration of buffer is at least 0.1 mM, at least 0.2 mM, at least 0.3 mM, at least 0.4 mM, at least 0.5 mM, at least 0.6 mM, at least 0.7 mM, at least 0.8 mM, or at least 0.9 mM. In other aspects of this embodiment, an effective concentration of buffer is at least 1.0 mM, at least 2.0 mM, at least 3.0 mM, at least 4.0 mM, at least 5.0 mM, at least 6.0 mM, at least 7.0 mM, at least 8.0 mM, or at least 9.0 mM. In yet other aspects of this embodiment, an effective concentration of buffer is at least 10 mM, at least 20 mM, at least 30 mM, at least 40 mM, at least 50 mM, at least 60 mM, at least 70 mM, at least 80 mM, or at least 90 mM. In still other aspects of this embodiment, an effective concentration of buffer is at least 100 mM, at least 200 mM, at least 300 mM, at least 400 mM, at least 500 mM, at least 600 mM, at least 700 mM, at least 800 mM, or at least 900 mM. In further aspects of this embodiment, an effective concentration of buffer is at most 0.1 mM, at most 0.2 mM, at most 0.3 mM, at most 0.4 mM, at most 0.5 mM, at most 0.6 mM, at most 0.7 mM, at most 0.8 mM, or at most 0.9 mM. In still other aspects of this embodiment, an effective concentration of buffer is at most 1.0 mM, at most 2.0 mM, at most 3.0 mM, at most 4.0 mM, at most 5.0 mM, at most 6.0 mM, at most 7.0 mM, at most 8.0 mM, or at most 9.0 mM. In yet other aspects of this embodiment, an effective concentration of buffer is at most 10 mM, at most 20 mM, at most 30 mM, at most 40 mM, at most 50 mM, at most 60 mM, at most 70 mM, at most 80 mM, or at most 90 mM. In still other aspects of this embodiment, an effective concentration of buffer is at most 100 mM, at most 200 mM, at most 300 mM, at most 400 mM, at most 500 mM, at most 600 mM, at most 700 mM, at most 800 mM, or at most 900 mM. In still further aspects of this embodiment, an effective concentration of buffer is about 0.1 mM to about 900 mM, 0.1 mM to about 500 mM, 0.1 mM to about 100 mM, 0.1 mM to about 90 mM, 0.1 mM to about 50 mM, 1.0 mM to about 900 mM, 1.0 mM to about 500 mM, 1.0 mM to about 100 mM, 1.0 mM to about 90 mM, or 1.0 mM to about 50 mM.

(120) Embodiments described herein can be practiced with a composition that comprises a plurality of botulinum toxin serotypes, such as botulinum toxin serotypes selected from the group consisting of botulinum toxin serotypes A, B, C.sub.1 D, E, F and G. In certain embodiments, purified botulinum toxins, can be used. In other embodiments, modified botulinum toxins may be used.

(121) In some embodiments, the composition may optionally also include NaCl. NaCl may particularly preferably be included in compositions comprising botulinum toxin, trehalose or sucrose, poloxamer 188, and methionine, and is particularly preferably included in liquid compositions comprising botulinum toxin, trehalose or sucrose, poloxamer 188, and methionine. In some lyophilized formulations, NaCl may function as a tonicity agent in a reconstitution vehicle. In one embodiment, NaCl may be present in an amount of 0.9% (w/w) in the reconstitution vehicle.

(122) In some embodiments, the Clostridial toxin active ingredient pharmaceutical composition can be formulated as a lyophilized (i.e. freeze dried) or vacuum dried powder which can be reconstituted with a suitable fluid, such as saline or water, prior to administration to a patient. In alternative embodiments, the pharmaceutical composition can be formulated as an aqueous solution or suspension.

(123) In some embodiments, the solid Clostridial toxin active ingredient pharmaceutical composition comprises a botulinum toxin, a tonicity agent and/or a lyoprotector, a poloxamer and/or a polysorbate and an antioxidant. In some embodiments, the Clostridial toxin active ingredient pharmaceutical composition comprises a botulinum toxin. In some embodiments, the Clostridial toxin active ingredient pharmaceutical composition comprises trehalose. In some embodiments, the Clostridial toxin active ingredient pharmaceutical composition comprises poloxamer 188 or polysorbate 20. In some embodiments, the composition comprises EDTA EGTA, DTPA, or analogs thereof. In alternative embodiments, the composition comprises methionine and/or NAC. In aspects of these alternative embodiments, the composition further comprises EDTA, EGTA, DTPA, or analogs thereof. In some embodiments, the composition further comprises a buffering agent. In one embodiment, the composition comprises histidine buffer. In some embodiments, the relative weight amounts of disaccharide, poloxamer and antioxidant are within the following ranges: trehalose: 1 to 15%, 1 to 10%, or 2-15% or 2-10%; poloxamer: 0.5-8% or 0.5 to 5%; methionine: 0.01 to 5%, 0.02 to 3%, 0.05 to 1%, 0.05 to 0.5%. In some embodiments, the relative weight amounts of trehalose, poloxamer and methionine are within the following ranges: trehalose: 1 to 15%, 1 to 10%, or 2-15% or 2-10%; poloxamer: 0.5-8% or 0.5 to 5%; methionine: 0.01 to 5%, 0.02 to 3%, 0.05 to 1%, 0.05 to 0.5%. In some embodiments, the relative weight amounts of trehalose, poloxamer and methionine are within the following ranges respectively: 1 to 10%; 0.5 to 5% and 0.1 to 0.3%. In other embodiments, the relative weight amounts of trehalose, polysorbate and methionine are within the following ranges respectively: 1 to 15%; 0.02% to 0.06%; and 0.1 to 0.3%. In other embodiments, the relative weight amounts of trehalose, polysorbate and methionine are within the following ranges respectively: 1 to 10%; 0.02% to 0.06%; and 0.1 to 0.3%. In some embodiments, the relative weight amount of EDTA or an EDTA analog is from about 0.01 to 0.10%. In some embodiments, the relative weight amount of NAC ranges from 0.01 to 0.5%.

(124) In aspects of these embodiments, the Clostridial toxin active ingredient pharmaceutical composition is formulated as a solid (i.e., lyophilized or vacuum dried) composition. In some embodiments, the solid Clostridial pharmaceutical composition comprises a lyoprotector. In some embodiments, the preferred lyoprotector includes sucrose, trehalose, mannitol, sorbitol, glucose, or combinations thereof. In some embodiments, the solid pharmaceutical composition comprises NAC in a relative weight amount of 0.01 to 0.5%. In some embodiments, the pharmaceutical composition further comprises EDTA, EGTA, DTPA, or analogs thereof. In alternative embodiments, the solid pharmaceutical composition comprises methionine and EDTA, EGTA, DTPA, or analogs thereof.

(125) In one embodiment, the composition is a solid composition consisting of botulinum toxin type A, 8% (w/w) trehalose, 4% (w/w) poloxamer 188, 0.2% (w/w) methionine, and a Histidine buffer. In one embodiment, the solid composition is reconstituted with a reconstitution vehicle comprising NaCl prior to administration to a patient. In one embodiment, NaCl may be present in an amount of 0.9% (w/w) in the reconstitution vehicle.

(126) In an alternative aspect of these embodiments, the Clostridial toxin active ingredient pharmaceutical composition is formulated as a liquid. In some embodiments, the liquid pharmaceutical composition comprises a tonicity agent selected from trehalose, sucrose, sodium chloride, mannitol, sorbitol, glucose, and combinations thereof. In some embodiments, the liquid pharmaceutical composition comprises NAC in a relative weight amount of 0.1 to 0.5%. In some embodiments, the liquid pharmaceutical composition comprises NAC and a chelating agent

selected from EDTA, EGTA, DTPA, and analogs thereof. In some embodiments, the liquid pharmaceutical composition comprises histidine buffer. In some embodiments, the liquid pharmaceutical composition has a pH from 5 to 7.

(127) Methods of Treatment

(128) In embodiments, methods of treating diseases, disorders, conditions, and the like, are described which comprise administering a pharmaceutical composition as described herein to a subject in need thereof in an amount sufficient to produce improved patient function. In certain embodiments, the diseases are of a neuromuscular nature, such as, for example, those diseases that affect muscles and nerve control thereof, such as, for example, overactive bladder, and the like. Certain embodiments relate to the treatment of pain, such as, for example, treatment of headache pain, or back pain, or muscle pain, or the like. In certain embodiments, methods encompass the treatment of psychological disorders, including, for example, depression, anxiety, and the like.

(129) Compositions and methods described herein can be useful for the treatment, reduction of symptoms, and/or prevention of, for example, achalasia, anal fissure, anismus, blepharospasm, cerebral palsy, cervical dystonia, cervicogenic headache, hemifacial spasm, dyshidrotic eczema, dysphagia, dysphonia, esophageal dysmotility, esophageal muscular ring, esotropia (infantile), eyelift, facial myokymia, gait disturbances (idiopathic toe-walking), generalized dystonia, hemifacial spasm, hyperfunctional facial lines (glabellar, forehead, crows' feet, down-turned angles of the mouth), hyperhidrosis, incontinence (idiopathic or neurogenic), medication overuse headache, migraine headache, myoclonus, muscle mass or activity reduction, involving, for example, the masseter or the like, myofascial pain syndrome, obstructive urinary symptoms, pancreas divisum pancreatitis, Parkinson's disease, puborectalis syndrome, reduction of surgical scar tension, salivary hypersecretion, sialoceles, sixth nerve palsy, spasticity, speech/voice disorders, strabismus, surgery adjunct (ophthalmic), tardive dyskinesia, temporomandibular joint disorders, tension headache, thoracic outlet syndrome, torsion dystonia, torticollis, Tourette's syndrome, tremor, whiplash-associated neck pain, pain, itching, inflammation, allergy, cancer and benign tumors, fever, obesity, infectious diseases, viral and bacterial, hypertension, cardiac arrhythmias, vasospasm, atherosclerosis, endothelial hyperplasia, venous thrombosis, varicose veins, aphthous stomatitis, hypersalivation, temporomandibular joint syndrome, hyperhidrosis, bromhidrosis, acne, rosacea, hyperpigmentation, hypertrophic scars, keloids, calluses and corns, skin wrinkling, excessive sebum production, psoriasis, dermatitis, allergic rhinitis, nasal congestion, post nasal drip, sneezing, ear wax, serous and suppurative otitis media, tonsil and adenoid hypertrophy, tinnitus, dizziness, vertigo, hoarseness, cough, sleep apnea, snoring, glaucoma, conjunctivitis, uveitis, strabismus, Grave's disease, excessive hair growth, hair loss, asthma, bronchitis, emphysema, mucus production, pleuritis, coagulation disorders, myeloproliferative disorders, disorders involving eosinophils, neutrophils, macrophages and lymphocytes, immune tolerance and transplantation, autoimmune disorders, dysphagia, acid reflux, hiatal hernia, gastritis and hyperacidity, diarrhea and constipation, hemorrhoids, urinary incontinence, prostatic hypertrophy, erectile dysfunction, priapism and Peyronie's disease, epididymitis, contraception, menstrual cramps, preventing premature delivery, endometriosis and fibroids, arthritis, osteoarthritis, rheumatoid, bursitis, tendonitis, tenosynovitis, fibromyalgia, seizure disorders, spasticity, headache, and neuralgias.

(130) In certain embodiments, patients are limited to a maximum of 360 U of botulinum toxin administered over any 90-day period.

(131) Treatment of Nerve/Muscle Conditions

(132) In an embodiment, the neuromuscular disease is hyperhidrosis. A subject suffering from hyperhidrosis, for example, receives about 59 U per axilla, or about 58 U per axilla, or about 57 U per axilla, or about 56 U per axilla, or about 55 U per axilla, or about 54 U per axilla, or about 53 U per axilla, or about 52 U per axilla, or about 51 U per axilla, or about 50 U per axilla, or about 49 U per axilla, or about 48 U per axilla, or about 47 U per axilla, or about 46 U per axilla, or about 45 U

per axilla, or about 44 U per axilla, or about 43 U per axilla, or about 42 U per axilla, or about 41 U per axilla, or about 40 U per axilla, or about 39 U per axilla, or about 38 U per axilla, or about 37 U per axilla, or about 36 U per axilla, or less, per treatment of a pharmaceutical composition described herein. In an embodiment, 50 U total are injected intradermally into 10-15 sites spaced approximately 1-2 cm apart.

(133) In an embodiment, the neuromuscular disease is hemifacial spasm. A subject suffering from hemifacial spasm, for example receives between about 1.5 to 15 U per treatment of a pharmaceutical composition described herein. In a further example, the subject receives between about 1.5 to 3 U, 1.5 to 5 U, 1.5 to 7 U, 1.5 to 10 U, 1.5 to 12 U, 1.5 to 15 U, 5 to 10 U, 5 to 15 U, or 10 to 15 U per treatment are administered to a patient with hemifacial spasm. In a still further example, the subject receives about 1.5 U, about 2 U, about 2.5 U, about 3 U, about 3.5 U, about 4 U, about 4.5 U about 5 U, about 5.5 U, about 6 U, about 6.5 U, about 7 U, about 7.5 U, about 8 U, about 8.5 U, about 9 U, about 9.5 U, about 10 U, about 10.5 U, about 11 U, about 11.5 U, about 12 U, about 12.5 U, about 13 U, about 13.5 U, about 14 U, about 14.5 U, or about 15 U per treatment are administered to a patient with hemifacial spasm. Dosages greater than 15 U per treatment may also be administered to patients with hemifacial spasm to achieve a therapeutic response. A treatment session can comprise multiple treatments.

(134) In an embodiment, the neuromuscular disease is cervical dystonia. A subject suffering from cervical dystonia, for example, receives between about 15 to 300 U per treatment of a pharmaceutical composition described herein. In a further example, the subject receives between about 35 to 250 U, 65 to 200 U, 85 to 175 U, 105 to 160 U, or 125 to 145 U are administered to a patient with cervical dystonia. In an embodiment, dosages to the sternocleidomastoideole is limited to 100 U or less. Dosages greater than 300 U per treatment may also be administered to patients with cervical dystonia to achieve a therapeutic response. A treatment session can comprise multiple treatments.

(135) In an embodiment, the neuromuscular disease is blepharospasm. A subject suffering from blepharospasm, for example, receives between about 1.25 to 2.5 U of a pharmaceutical composition described herein injected into the medial and lateral pretarsal orbicularis oculi of the upper lid and into the lateral pretarsal orbicularis oculi of the lower lid. In a further example, the subject receives about 1.5 U, about 1.6 U, about 1.7 U, about 1.8 U, about 1.9 U, about 2.0 U, about 2.1 U, about 2.2 U, about 2.3 U, about 2.4 U, about 2.5 U, or more, per injection site. A treatment session can comprise multiple treatments.

(136) In an embodiment, the neuromuscular disease is strabismus. A subject suffering from strabismus, for example, receives between about 1.25 to 2.5 U per injection site of a pharmaceutical composition described herein. In a further example, the subject receives about 1.5 U, about 1.6 U, about 1.7 U, about 1.8 U, about 1.9 U, about 2.0 U, about 2.1 U, about 2.2 U, about 2.3 U, about 2.4 U, about 2.5 U, or more, per injection site to achieve a therapeutic response. In embodiments, lower doses are used for treatment of small deviations. In embodiments, vertical muscles and horizontal strabismus of less than 20 prism diameters can be treated with 1.25 to 2.5 U per injection site. A treatment session can comprise multiple treatments.

(137) In an embodiment, the neuromuscular disease is muscle spasticity. A subject suffering from muscle spasticity, for example, receives between about 20 to 200 U per treatment of a pharmaceutical composition described herein. In a further example, the subject receives between about 20 to 30 U, 20 to 40 U, 20 to 60 U, 20 to 80 U, 20 to 100 U, 20 to 125 U, 20 to 150 U, or 20 to 175 U per treatment are administered to a patient with muscle spasticity. In a still further example, the subject receives about 20 U, about 25 U, about 30 U, about 35 U, about 40 U, about 45 U, about 50 U, about 55 U, about 60 U, about 65 U, about 70 U, about 75 U, about 80 U, about 85 U, about 90 U, about 95 U, about 100 U, about 105 U, about 110 U, about 115 U, about 120 U, about 125 U, about 130 U, about 135 U, about 140 U, about 145 U, about 150 U, about 155 U, about 160 U, about 165 U, about 170 U, about 175 U, about 180 U, about 185 U, about 190 U,

about 195 U, or about 200 U per treatment are administered to a patient with muscle spasticity. In an embodiment, the biceps brachii can be injected with between 100 U and 200 U divided into 4 injection sites. In an embodiment, the flexor carpi radialis can be injected with between 12.5 U and 50 U in 1 injection site. In an embodiment, the flexor carpi ulnaris can be injected with between 12.5 U and 50 U in 1 injection site. In an embodiment, the flexor digitorum *profundus* can be injected with between 30 U and 50 U in one injection site. In an embodiment, the flexor digitorum *sublimis* can be injected with between 30 U and 50 in a single injection site. Dosages greater than 200 U per treatment may also be administered to patients with muscle spasticity to achieve a therapeutic response. A treatment session can comprise multiple treatments.

(138) Treatment of Pain

(139) In another embodiment, methods for treating pain comprising the step of administering a pharmaceutical composition described herein to a subject in need thereof in an amount sufficient to reduce pain. In another embodiment, the patient suffers from myofascial pain, migraine headache pain, tension headache pain, neuropathic pain, facial pain, lower-back pain, sinus-headache pain, pain associated with temporomandibular joint disease, pain associated with spasticity or cervical dystonia, post-surgical wound pain, or neuralgia. A treatment session can comprise multiple treatments.

(140) In an embodiment, the patient suffers from facial pain. A subject suffering from facial pain, for example, receives between about 4 to 40 U per treatment of a pharmaceutical composition described herein. In a further example, the subject receives between about 4 to 10 U, 4 to 15 U, 4 to 20 U, 4 to 25 U, 4 to 30 U, 4 to 35 U, 7 to 15 U, 7 to 20 U, 7 to 25 U, 7 to 30 U, 7 to 35 U, or 7 to 40 U per treatment are administered to a patient suffering from facial pain. In a still further example, the subject receives about 4 U, about 5 U, about 7.5 U, about 10 U, about 12.5 U, about 15 U, about 17.5 U, about 20.0 U, about 22.5 U, about 25.0 U, about 27.5 U, about 30.0 U, about 32.5 U, about 35 U, about 37.5 U, or about 40 U per treatment are administered to a patient with facial pain. Dosages greater than 40 U per treatment may also be administered to patients with facial pain to achieve a therapeutic response. A treatment session can comprise multiple treatments.

(141) In an embodiment, the patient suffers from myofascial pain. A subject suffering from myofascial pain, for example, receives between about 5 to 100 U per treatment of a pharmaceutical composition described herein. In a further example, the subject receives between about 5 to 10 U, 5 to 20 U, 5 to 30 U, 5 to 40 Units, 5 to 50 Units, 5 to 60 Units, 5 to 70 Units, 5 to 80 Units, 5 to 90 U, 10 to 20 U, 10 to 30 U, 10 to 50 U, or 10 to 60 U, or 10 to 70 U, or 10 to 80 U, 10 to 90 U, or 10 to 100 U per treatment are administered to a patient suffering from myofascial pain. In a further example, the subject receives about 5 U, about 10 U, about 15 U, about 20 U, about 25 U, about 30 U, about 35 U, about 40 U, about 45 U, about 50 U, about 55 U, about 60 U, about 65 U, about 70 U, about 75 U, about 80 U, about 85 U, about 90 U, about 95 U, or about 100 U per treatment. Dosages greater than 100 U per treatment may also be administered to patients with myofascial pain to achieve a therapeutic response. A treatment session can comprise multiple treatments.

(142) In an embodiment, the subject suffers from lower-back pain. A subject suffering from lower-back pain, for example, receives between about 15 to 150 U per treatment of a pharmaceutical composition described herein. In a further example, the subject receives between about 15 to 30 U, 15 to 50 U, 15 to 75 U, 15 to 100 U, 15 to 125 U, 15 to 150 U, 20 to 100 U, 20 to 150 U, or 100 to 150 U per treatment. In a still further example, the subject receives about 15 U, about 20 U, about 25 U, about 30 U, about 35 U, about 40 U, about 45 U, about 50 U, about 55 U, about 60 U, about 65 U, about 70 U, about 75 U, about 80 U, about 85 U, about 90 U, about 95 U, about 100 U, about 105 U, about 110 U, about 115 U, about 120 U, about 125 U, about 130 U, about 135 U, about 140 U, about 145 U, or about 150 U per treatment to alleviate lower-back pain. Dosages greater than 150 U per treatment may also be administered to patients with lower-back pain to achieve a therapeutic response. A treatment session can comprise multiple treatments.

(143) In an embodiment, the patient suffers from migraine headache pain, including wherein the

patient suffers from migraine headaches of 4 hours or more 15 or more days per month. A subject suffering from migraine-headache pain, for example, receives between about 0.5 to 200 U per treatment of a pharmaceutical composition described herein. In a further example, the subject receives between about 5 to 190 U, 15 to 180 U, 25 to 170 U, 35 to 160 U, 45 to 150 U, 55 to 140 U, 65 to 130 U, 75 to 120 U, 85 to 110 U, or 95 to 105 U per treatment to alleviate migraine headache pain. A treatment session can comprise multiple treatments.

(144) For example, about 0.5 U, about 1.0 U, about 1.5 U, about 2.0 U, about 2.5 U, about 3.0 U, about 3.5 U, about 4.0 U, about 4.5 U, about 5.0 U, about 5.5 U, about 6.0 U, about 6.5 U, about 7.0 U, about 7.5 U, about 8.0 U, about 8.5 U, about 9.0 U, about 9.5 U, about 10.0 U, about 12 U, about 15 U, about 17 U, about 20 U, about 22 U, about 25 U, about 27 U, about 30 U, about 32 U, about 35 U, about 37 U, about 40 U, about 42 U, about 45 U, about 47 U, or about 50 U per treatment site are administered to a patient with migraine-headache pain. A patient can be treated at multiple sites, such as, for example, 2 sites, 3 sites, 4 sites, 5 sites, 6 sites, 7 sites, 8 sites, 9 sites, 10 sites, 11 sites, 12 sites, 13 sites, 14 sites, 15 sites, 16 sites, 17 sites, 18 sites, 19 sites, 20 sites, 21 sites, 22 sites, 23 sites, 24 sites, 25 sites, 26 sites, 27 sites, 28 sites, 29 sites, 30 sites, 31 sites, 32 sites, or more, or the like. In an embodiment, a patient suffering from migraine is injected 31 times with 5 U per 0.1 mL injection, across the corrugator (2 injections of 5 U each), procerus (1 injection of 5 U), *frontalis* (4 injections of 5 U each), temporalis (8 injections of 5 U each), occipitalis (6 injections of 5 U each), cervical paraspinal (4 injections of 5 U each), and trapezius (6 injections of 5 U each) muscles. With the exception of the procerus muscle which can be injected at the midline, all muscles can, in certain embodiments, be injected bilaterally with half of the injection sites to the left and half to the right side of the head and neck. Dosages greater than 200 U per treatment may also be administered to patients with migraine-headache pain to achieve a therapeutic response. A treatment session can comprise multiple treatments.

(145) In an embodiment, the patient suffers from sinus-headache pain. A subject suffering from sinus-headache pain, for example, receives between about 4 to 40 U per treatment of a pharmaceutical composition described herein. In a further example, the subject receives between about 4 to 10 U, 4 to 15 U, 4 to 20 U, 4 to 25 U, 4 to 30 U, 4 to 35 U, 7 to 15 U, 7 to 20 U, 7 to 25 U, 7 to 30 U, 7 to 35 U, or 7 to 40 U per treatment to alleviate sinus-headache pain. In a still further example, the subject receives about 4 U, about 5 U, about 7.5 U, about 10 U, about 12.5 U, about 15 U, about 17.5 U, about 20.0 U, about 22.5 U, about 25.0 U, about 27.5 U, about 30.0 U, about 32.5 U, about 35 U, about 37.5 U, or about 40 U per treatment. Dosages greater than 40 U per treatment may also be administered to patients with sinus headache-pain to achieve a therapeutic response. A treatment session can comprise multiple treatments.

(146) In an embodiment, the patient suffers from tension-headache pain. A subject suffering from tension-headache pain, for example, receives between about 5 to 50 U per treatment of a pharmaceutical composition described herein. In a further example, between about 5 to 10 U, 5 to 15 U, 5 to 20 U, 5 to 25 U, 5 to 30 U, 5 to 35 U, 5 to 40 U, 5 to 45 U, 10 to 20 U, 10 to 25 U, 10 to 30 U, 10 to 35 U, 10 to 40 U, or 10 to 45 U per treatment are administered to a patient with tension-headache pain. In a still further example, the subject receives about 5 U, about 10 U, about 20 U, about 25 U, about 30 U, about 35 U, about 40 U, about 45 U, or about 50 U per treatment administered to alleviate tension-headache pain. In an embodiment, a patient suffering from tension headache is injected 31 times with 5 U per 0.1 mL injection, across the corrugator (2 injections of 5 U each), procerus (1 injection of 5 U), *frontalis* (4 injections of 5 U each), temporalis (8 injections of 5 U each), occipitalis (6 injections of 5 U each), cervical paraspinal (4 injections of 5 U each), and trapezius (6 injections of 5 U each) muscles. With the exception of the procerus muscle which can be injected at the midline, all muscles can, in certain embodiments, be injected bilaterally with half of the injection sites to the left and half to the right side of the head and neck. Dosages greater than 200 U per treatment may also be administered to patients with tension headache pain to achieve a therapeutic response. A treatment session can comprise multiple treatments.

(147) In an embodiment, the patient suffers from sinus headache pain or facial pain associated with acute or recurrent chronic sinusitis. For example, a pharmaceutical composition as described herein can be administered to the nasal mucosa or to the subcutaneous structures overlying the sinuses, wherein the administration of the formulation reduces the headache and/or facial pain associated with acute recurrent or chronic sinusitis. In further embodiments, any of the pharmaceutical formulations described herein can be administered to the nasal mucosa or to the subcutaneous structures overlying the sinuses, such as over one or more of the sinuses selected from the group consisting of: ethmoid; maxillary; mastoid; frontal; and sphenoid. In another embodiment, subcutaneous structures overlying the sinuses lie within one or more of the areas selected from the group consisting of: forehead; malar; temporal; post auricular; and lip. In embodiments, multiple injections of 5 U each are administered to treat the sinus headache pain or facial pain associated with acute or recurrent chronic sinusitis.

(148) In another embodiment, a patient suffering from sinus headache pain or facial pain associated with acute or recurrent chronic sinusitis is treated by administering any of the pharmaceutical formulations described herein to an afflicted area of the patient. In a further embodiment, the pharmaceutical formulations disclosed herein are administered to the projections of a trigeminal nerve innervating a sinus.

(149) Patients suffering from sinus headache pain or facial pain associated with acute or recurrent chronic sinusitis often exhibit symptoms including rhinitis, sinus hypersecretion and/or purulent nasal discharge. In one embodiment, patients treated with the pharmaceutical compositions described herein exhibit, prior to treatment, symptoms of sinus hypersecretion and purulent nasal discharge.

(150) Embodiments contemplated herein provide methods for treating a patient suffering from sinus headache pain or facial pain associated with acute or recurrent chronic sinusitis, wherein the subject suffers from neuralgia. In certain embodiments, the neuralgia is trigeminal neuralgia. In another embodiment, the neuralgia is: associated with compressive forces on a sensory nerve; associated with intrinsic nerve damage, demyelinating disease, or a genetic disorder; associated with a metabolic disorder; associated with central neurologic vascular disease; or associated with trauma. In another embodiment, the pain is associated with dental extraction or reconstruction.

(151) Treatment of Urological Disorders

(152) In an embodiment, methods for treating a patient suffering from overactive bladder (OAB), such as, for example, that due to a neurologic condition (NOAB), or idiopathic OAB (IOAB) are provided. For example, pharmaceutical formulations described herein can be administered to the bladder or its vicinity, e.g. the detrusor, wherein the administration of the formulation reduces the urge incontinence associated with overactive bladder. In certain embodiments, the dosage can be, for example, 200 U, or more, or less, or the like. For example, the dosage can be about 15 U, about 20 U, about 25 U, about 30 U, about 35 U, about 40 U, about 45 U, about 50 U, about 55 U, about 60 U, about 65 U, about 70 U, about 75 U, about 80 U, about 85 U, about 90 U, about 95 U, about 100 U, about 105 U, about 110 U, about 115 U, about 120 U, about 125 U, about 130 U, about 135 U, about 140 U, about 145 U, about 150 U, about 160 U, about 170 U, about 180 U, about 190 U, about 200 U, about 210 U, about 220, about 230 U, about 240 U, or more, or the like, per treatment. A patient can be injected at multiple sites, such as, for example, 2 sites, 3 sites, 4 sites, 5 sites, 6 sites, 7 sites, 8 sites, 9 sites, 10 sites, 11 sites, 12 sites, 13 sites, 14 sites, 15 sites, 16 sites, 17 sites, 18 sites, 19 sites, 20 sites, 21 sites, 22 sites, 23 sites, 24 sites, 25 sites, 26 sites, 27 sites, 28 sites, 29 sites, 30 sites, 31 sites, 32 sites, 33 sites, 34 sites, 35 sites, 36 sites, 37 sites, 38 sites, or more, or the like. In an embodiment, patients suffering from OAB are treated with 30 1 mL injections of approximately 6.7 U per injection into the detrusor muscle.

(153) In an embodiment, methods for treating a patient suffering from neurogenic detrusor overactivity (NDO), such as that due to a neurologic condition, are provided. For example, pharmaceutical formulations can be administered to the bladder or its vicinity, e.g. the detrusor,

wherein the administration of the formulation reduces the urge incontinence associated with overactive bladder. In certain embodiments, the dosage can be, for example, 200 U, or more, or less, or the like. For example, the dosage can be about 15 U, about 20 U, about 25 U, about 30 U, about 35 U, about 40 U, about 45 U, about 50 U, about 55 U, about 60 U, about 65 U, about 70 U, about 75 U, about 80 U, about 85 U, about 90 U, about 95 U, about 100 U, about 105 U, about 110 U, about 115 U, about 120 U, about 125 U, about 130 U, about 135 U, about 140 U, about 145 U, about 150 U, about 160 U, about 170 U, about 180 U, about 190 U, about 200 U, about 210 U, about 220, about 230 U, about 240 U, or more, or the like, per treatment. A patient can be injected at multiple sites, such as, for example, 2 sites, 3 sites, 4 sites, 5 sites, 6 sites, 7 sites, 8 sites, 9 sites, 10 sites, 11 sites, 12 sites, 13 sites, 14 sites, 15 sites, 16 sites, 17 sites, 18 sites, 19 sites, 20 sites, 21 sites, 22 sites, 23 sites, 24 sites, 25 sites, 26 sites, 27 sites, 28 sites, 29 sites, 30 sites, 31 sites, 32 sites, or more, or the like. In an embodiment, patients suffering from NDO are treated with 30 1 mL injections of approximately 6.7 U per injection into the detrusor muscle.

(154) Treatment of Cosmetic Features

(155) In another embodiment, methods for cosmetically modifying soft-tissue features comprising the step of administering at least one pharmaceutical composition as described herein to a subject in need thereof in an amount sufficient to modify said features are provided. In a further embodiment, the pharmaceutical composition is administered via transcutaneous or transmucosal injection either at a single focus or multiple foci.

(156) In embodiments, pharmaceutical formulations are administered to the face or neck of the subject. In a further embodiment, the pharmaceutical formulations are administered to the subject in an amount sufficient to reduce rhytides. For example, the formulation can be administered between eyebrows of the subject in an amount sufficient to reduce vertical lines between the eyebrows and on a bridge of a nose. The pharmaceutical formulations can also be administered near either one or both eyes of the subject in an amount sufficient to reduce lines at corners of the eyes. In an embodiment, compositions can be injected locally to smooth skin. In another embodiment, the pharmaceutical formulations can also be administered to a forehead of the subject in an amount sufficient to reduce horizontal lines on said forehead. In yet another embodiment, the pharmaceutical formulation is administered to the neck of the subject in an amount sufficient to reduce muscle bands in the neck. In an embodiment, a pharmaceutical composition is applied to the masseter muscle to relax the muscle and/or decrease masseter mass.

(157) In a further embodiment, the patient suffers from facial wrinkles. A subject suffering from facial wrinkles, for example, can receive between about 1 to 100 U per treatment of a pharmaceutical formulation. In a further example, the subject receives between about 1 to 10 U, 1 to 20 U, 1 to 30 U, 1 to 40 U, 1 to 50 U, 1 to 60 U, 1 to 70 U, 1 to 80 U, 1 to 90 U, 5 to 20 U, 5 to 30 U, 5 to 40 U, 5 to 50 U, 5 to 60 U, 5 to 70 U, 5 to 80 U, 5 to 90 U, or 5 to 100 U per treatment. In a still further example, the subject receives about 1 U, about 10 U, about 20 U, about 30 U, about 40 U, about 50 U, about 60 U, about 70 U, about 80 U, about 90 U, or about 100 U per treatment. Dosages greater than 100 U per treatment may also be administered to patients suffering from inflammation or an inflammatory disorder to achieve a therapeutic response.

(158) Treatment of Inflammation

(159) In another embodiment, methods for treating inflammation comprising the step of administering a pharmaceutical composition as described herein to a subject in need thereof in an amount sufficient to reduce inflammation. In certain embodiments, pharmaceutical formulations are administered to a patient without producing muscle weakness. In an embodiment, the pharmaceutical formulations are administered to patients with an inflammatory condition. In certain embodiments, the inflammatory condition is neurogenic inflammation. In another embodiment, the subject suffers from rheumatoid arthritis or a gastro-intestinal inflammatory disease.

(160) In a further embodiment, the patient suffers from an inflammatory disorder. A subject

suffering from an inflammatory disorder, for example, receives between about 1 to 100 U per treatment of a pharmaceutical composition as described herein. In a further example, the subject receives between about 1 to 10 U, 1 to 20 U, 1 to 30 U, 1 to 40 U, 1 to 50 U, 1 to 60 U, 1 to 70 U, 1 to 80 U, 1 to 90 U, 5 to 20 U, 5 to 30 U, 5 to 40 U, 5 to 50 U, 5 to 60 U, 5 to 70 U, 5 to 80 U, 5 to 90 U, or 5 to 100 U per treatment. In a still further example, the subject receives about 1 U, about 10 U, about 20 U, about 30 U, about 40 U, about 50 U, about 60 U, about 70 U, about 80 U, about 90 U, or about 100 U per treatment. Dosages greater than 100 U per treatment may also be administered to patients suffering from inflammation or an inflammatory disorder to achieve a therapeutic response.

(161) Treatment of Skin Conditions

(162) A method for treating a skin disorder can have the step of local administration of a botulinum neurotoxin to a location of a skin disorder of a patient, such as to a face, hand or foot of a patient. The neurotoxin can be locally administered in an amount of between about 10-3 units/kg of patient weight and about 35 units/kg of patient weight. For example, the neurotoxin is locally administered in an amount of between about 10-2U/kg and about 25U/kg of patient weight. In a further example, the neurotoxin is administered in an amount of between about 10-1U/kg and about 15U/kg. In one method, the neurotoxin is locally administered in an amount of between about 1U/kg and about 10U/kg in a composition as described herein. In a clinical setting, it can be advantageous to administer from 1 U to 3000 U of a neurotoxin, such as botulinum toxin type A or B, to a skin disorder location by topical application or by subdermal administration, to effectively treat the skin disorder.

(163) Administration of botulinum toxin can be carried out at multiple sites in the skin, wherein the sites of adjacent injections are separated by about 0.1 to 10 cm, or about 0.5 to about 5 cm, for example, by about 1.5 to about 3 cm. The toxins may be any of the botulinum toxins A, B, C, D, E, F, G or a mosaic toxin. The amounts administered may vary between 0.1 and 1000 U, or about 1 to about 40, or from about 5 to about 10 U, depending on the manufactures specifications, the class of the toxin and the mode of administration. The repeat time range for these administrations for maintenance of the desired change varies substantially according to the location of the injection, the condition to be adjusted and the condition of the patient. Thus, the repeat time may vary from about 1 week to about 50 weeks, however, a common range is about 4 to about 25 weeks, or even about 12 weeks to about 16 weeks.

(164) The distances between administration sites, such as, for example, injection sites, can vary from about 1 mm to about 10 cm, suitably from about 5 mm to about 5 cm, and more usually from about 1 cm to about 3 cm. Thus, for example botulinum A may be suitably administered by intradermal injection between about 0.1 to about 10 U at a separation of from about 0.5 to about 10 cm.

(165) In another embodiment, methods for treating cutaneous disorders comprising the step of administering a pharmaceutical composition as described herein to a subject in need thereof in an amount sufficient to reduce a sebaceous or mucous secretion is provided. In further embodiments, the pharmaceutical compositions as described herein are administered to a patient without producing muscle weakness. In certain embodiments, the pharmaceutical composition as described herein are injected into one or more sites of an eyelid or conjunctiva. In another embodiment, the formulations are administered to a body surface.

(166) In another embodiment, the pharmaceutical formulations are administered in an amount sufficient to reduce cutaneous bacterial or fungal growth, including but not limited to *Staphylococcus*; *Streptococcus* and *Moraxella*. For example, the pharmaceutical compositions as described herein are administered to an area selected from the group consisting of: eyelid; scalp; feet; groin; and armpit to reduce cutaneous infection.

(167) Treatment of Depression

(168) In another embodiment, a method for treating depression is provided. Depression is a general

term for recognized forms of depression that are defined and are separately diagnosed according to criteria given in handbooks for psychiatry, for example in the *Diagnostic and Statistical Manual of Mental Disorders* 4th edition (DSM-IV) published by the American Psychiatric Association, Washington, D.C. (1994). In the DSM-IV, depressive disorders are classified under mood disorders and are divided into three types: major depressive disorder, dysthymic disorder and depressive disorder not otherwise specified (or “atypical”). In general, regardless of whether or not the depressive syndrome is melancholic, atypical, or some admixture of the two, a diagnosis of major depression is given when depressed mood is present, or loss of interest or pleasure in all activities is present, for at least two weeks.

(169) Depression is often associated with psychomotor abnormalities, such as increased or retarded motor activity. Many depressed persons can also be recognized by their “depressed facies” in which the muscles of facial expression assume a distressed or sad appearance. For example, the brow may be furrowed, the inner ends of the eyebrows raised, and the angles of the mouth lowered such that the facial appearance is recognizably sad and/or anxious. Four major muscle groups are involved in frowning: the frontal, procerus, corrugator supercilii and orbicularis oculi (Weider et al. *Derm Surg.* 24:1172-1174, 1998. The corrugator supercilii is also known as the “scowl” muscle.

(170) A subject diagnosed with depression or experiencing a depressive episode is treated by administering any of the pharmaceutical compositions described herein. In a further embodiment, the pharmaceutical formulations disclosed herein are administered to the patient via injection subcutaneously. Example 10 describes treatment of a person with a form of depression by administering a therapeutically effective amount of botulinum toxin in a composition comprising botulinum toxin, trehalose, a poloxamer surfactant, and methionine. The composition is administered to a facial muscle involved in frowning or scowling. The neurotoxin affects the ability of the subject to frown and/or scowl, thereby treating depression. More generally, a therapeutically effective amount of Botulinum toxin A can be injected into one or more of the *frontalis*, procerus, the corrugator supercilii, orbicularis oculi, or the depressor anguli oris (*triangularis* muscle).

(171) In another embodiment, the method comprises administering a therapeutically effective amount of a Clostridial toxin active ingredient in a composition as described herein to a facial muscle involved in frowning, scowling, or a sad appearance. The Clostridial toxin active ingredient causes partial or complete paralysis of the facial muscle, thereby affecting the ability of the subject to frown and/or scowl, or appear sad, and thereby treat depression. For example, a therapeutically effective amount of a composition comprising the Clostridial toxin active ingredient botulinum toxin, along with a surfactant, an antioxidant, and optionally a tonicity modifier, can be injected into one or more of the orbicularis oculi, *frontalis*, procerus, the corrugator supercilii, or the depressor anguli oris (*triangularis* muscle). In a specific example, the composition comprising Botulinum toxin A is injected into the procerus muscle over the *glabella*. Other administration points and paradigms are disclosed, for example, in U.S. Pat. No. 7,758,872, which is incorporated by reference herein.

(172) In other embodiments of the method, adult subjects with moderate to severe major depressive disorder (MDD), either single episode or recurrent are contemplated for treatment, where the MDD diagnosis is based upon the DSM-IV-TR criteria. In one embodiment, a single treatment is contemplated, and in other embodiments, a single, repeated treatment is contemplated, with the treatment is repeated at intervals of 2-6, 2-4, or 3-6 months. The amount of Clostridial toxin active ingredient dosed is, for example, 30 U or 50 U, where, in some embodiments, the dose is divided into a plurality of injections. In one embodiment, the plurality of injections is 6 and in another embodiment, is 8. In one embodiment, 30 U is divided into 6 injections to the glabellar region of the forehead (procerus and corrugator muscles). In one embodiment, 50 U is divided into 8 injections to the glabellar region of the forehead (procerus and corrugator muscles).

(173) Effective treatment is indicated by, for example, a primary efficacy measure known in the art, such as the clinical assessment known as Montgomery-Asberg Depression Rating scale. Additional

efficacy measures: clinic CGI-S score (Clinical Global Impression of Change scores), clinic HAM-D17 total score (Hamilton Rating Scale for Depression).

(174) Treatment of Cardiac Arrhythmia

(175) In another embodiment, a method for treating cardiac arrhythmia is provided. Arrhythmias are caused by a disruption of the normal functioning of the electrical conduction system of the heart. Normally, the chambers of the heart (atria and ventricles) contract in a coordinated manner. The signal to contract is an electrical impulse that begins in the sinoatrial nod, and the impulse is conducted through the atria and stimulates them to contract. The impulse passes through the atrioventricular node, then travels through the ventricles and stimulates them to contract. Problems can occur anywhere along the conduction system, causing various arrhythmias. Problems can also occur in the heart muscle itself, causing it to respond differently to the signal to contract, also causing arrhythmias, or causing the ventricles to contract independently of the normal conduction system.

(176) Arrhythmias include tachycardias, bradycardias and true arrhythmias of disturbed rhythm. Arrhythmias are classified as lethal if they cause a severe decrease in the pumping function of the heart. When the pumping function is severely decreased for more than a few seconds, blood circulation is essentially stopped, and organ damage (such as brain damage) can occur within a few minutes. Lethal arrhythmias include ventricular fibrillation, also ventricular tachycardia that is rapid and sustained, or pulseless, and may include sustained episodes of other arrhythmias. Additional types of arrhythmias include atrial fibrillation or flutter, multifocal atrial tachycardia, paroxysmal supraventricular tachycardia, Wolff-Parkinson-White syndrome, sinus tachycardia, sinus bradycardia, bradycardia associated with heart block, sick sinus syndrome, and ectopic heartbeat.

(177) Accordingly, a method for treating cardiac arrhythmia is provided, the method comprising the step of administering a composition as described herein that comprises a therapeutically effective amount of Clostridial toxin active ingredient, the composition administered locally to the heart of a patient with a cardiac arrhythmia or at risk of a cardiac arrhythmia. Particular arrhythmias treatable include bradycardia and tachycardia. In one embodiment, the composition is locally administered, by which is meant administration directly to, in, or to the vicinity of, the cardiac muscle to be treated. Local administration includes intrapericardial, intracardiac cardiac catheterization and direct cardiac muscle injection routes of administration for the composition.

(178) Example 11 describes treatment of a person undergoing cardiac surgery by administering a therapeutically effective amount of botulinum toxin in a composition comprising botulinum toxin, trehalose, a poloxamer surfactant, and NAC. In one embodiment, the composition is administered via injection into one or more epicardial fat pads of the heart. The dose administered, in one exemplary embodiment, 25 U per epicardial fat pad, to a total dose of 125U. In another exemplary embodiment, 50 U per epicardial fat pad, to a total dose of 250 U, is administered.

(179) Effective treatment is indicated by, for example, a primary efficacy endpoint of, for example, incidence of atrial fibrillation (AF) as measured by ECG for 4 weeks or at 4 weeks post treatment. Additional efficacy endpoints include length of hospital stay, length of stay in ICU, rehospitalization rate, anticoagulant medication use, need for interventional procedures for post operative atrial fibrillation, such as ablation, pacemaker implantation, electrical or pharmacologic cardioversion.

EXAMPLES

(180) The following examples illustrate embodiments and aspects of the present compositions and methods and are not intended to limit the scope thereof.

Example 1

(181) Activities and Stabilities of Exemplary Solid Clostridial Pharmaceutical Compositions

(182) Bulk solutions of botulinum toxin were prepared by mixing an appropriate aliquot of a botulinum toxin type A with a solution having the ingredients set forth in Tables 1-3. Four

formulations were prepared for testing, referred to as Formulation 1, Formulation 2, Formulation 3 and Formulation 4, and three comparative formulations were prepared, referred to as Comparator 1, Comparator 2, and Comparator 3. Formulations 1, 2, 3 and 4 were composed of a disaccharide, a surfactant, an antioxidant and a histidine buffer. Each formulation was filled into a glass vial and lyophilized using conventional freeze-drying conditions. Portions of the lyophilized formulations were stored under several conditions; for example, at 25° C. and at 40° C. for six months (Tables 1.1 and 1.2), or at -20° C. and 40° C. for 1 month (Table 2), or at 25° C. for 7.5 months (Table 3). Potency of the lyophilized formulations after storage and reconstitution with saline was tested by a cell based potency assay, as described in U.S. Pat. No. 8,618,261, the assay details of which are incorporated by reference herein. The results were normalized to target. The potencies of the solid compositions are shown in Tables 1-3.

Example 2

(183) Activities of Liquid Clostridial Pharmaceutical Compositions in the Presence or Absence of Antioxidants

(184) Bulk drug product solutions were prepared by mixing an appropriate aliquot of a botulinum toxin type A with three different vehicle solutions as shown in Table 4. All three formulations contained 8% w/w trehalose, 4% w/w poloxamer P188 and 20 mM histidine buffer at pH 6.0. Formulation 10 contained no antioxidant. Formulations 11 and 12 contained NAC and methionine, respectively. The bulk solutions were filled into 2 mL glass vials (1.25 mL fill), and sealed with rubber stopper and aluminum shell. Potency of the formulations was tested by a cell based potency assay after filling (time zero, t0) and after storage for one month at four temperatures (-70° C., 5° C., 25° C. and 40° C.). The target potency was 100 Units/mL. Potency test results are given in Table 4.

Example 3

(185) Impact of Exemplary Antioxidants on the Stability of Exemplary Liquid Formulations

(186) Bulk drug product solutions were prepared by mixing an appropriate aliquot of the BoNT/A drug substance with different antioxidants as shown in Table 3.1. All compositions contained 8% w/w trehalose, 4% w/w poloxamer P188, 20 mM histidine buffer at pH 6.0 and 0.2 wt % of one or more of the following antioxidants: NAC, L-methionine, L-tryptophan, L-glutathione, sodium sulfite, and/or propyl gallate, and/or EDTA. The formulations are specified in Table 3.1. Target potency was 100 U/mL. The bulk solutions were filled into 2 mL glass vials (1.25 mL fill), and sealed with rubber stopper and aluminum shell. Potency of the formulations was tested by a cell based potency assay after filling (time zero, t0) and after storage at 40° C. for 2 weeks and 1 month. Potency test results are given in Table 5, above.

(187) TABLE-US-00014 TABLE 3.1 Toxin Tre P188 NAC Met TRP GSH NaSul PrpGal EDTA
 Formulation No. U/mL % % Buffer % % % % % % % % Formulation 20 100 8 4 20 mM 0.2 His, pH 6.0
 Formulation 21 100 8 4 20 mM 0.2 His, pH 6.0 Formulation 22 100 8 4 20 mM 0.2 His, pH 6.0
 Formulation 23 100 8 4 20 mM 0.2 His, pH 6.0 Formulation 24 100 8 4 20 mM 0.2 His, pH 6.0
 Formulation 25 100 8 4 20 mM 0.2 0.2 His, pH 6.0 Formulation 26 100 8 4 20 mM 0.2 0.03 His, pH 6.0
 Formulation 27 100 8 4 20 mM 0.2 0.2 0.03 His, pH 6.0 Formulation 28 100 8 4 20 mM 0.2 0.2 His, pH 6.0
 Tre = trehalose; P-188 = poloxamer P188; His = L-histidine; NAC = N-acetyl-L-cysteine; Met = L-methionine; TRP = L-tryptophan; GSH = L-glutathione; NaSul = sodium sulfite; PrpGal = propyl gallate; EDTA = ethylene diamine tetraacetic acid, sodium salt.

Example 4

(188) Impact of Exemplary Disaccharides on the Stability of Exemplary Liquid Compositions

(189) Liquid compositions were prepared by mixing an appropriate aliquot of BoNT/A (100 U/mL) with solutions of one of 8% w/w trehalose or 8% w/w sucrose, and of 4% w/w poloxamer P188 and 0.2 w/w % L-methionine, in 20 mM histidine buffer at pH 6.0. The composition with trehalose was assigned Formulation No. 30 and the composition with sucrose was assigned Formulation No. 31. The compositions were stored at 25° ° C. and potency was tested by cell based potency assay prior

to storage (time zero, t0) and at selected time points after storage. Potency test results are shown in Table 6.

Example 5

(190) Impact of Exemplary Surfactants on the Stability of Exemplary Liquid Compositions

(191) Liquid compositions were prepared by mixing an appropriate aliquot of BoNT/A (100 U/mL) with solutions comprising poloxamer P188 (4 w/w %) or polysorbate (TWEEN® 20, 0.04 w/w %) and 8% w/w trehalose, 0.2 w/w % L-methionine, in 20 mM histidine buffer at pH 6.0. The composition with poloxamer P188 was identified as Formulation No. 30 and the composition with polysorbate was identified as Formulation No. 32. Potency was tested by cell based potency assay prior to storage (time zero, t0) and after 1 month of storage at 40° ° C. Potency test results are shown in Table 7.

Example 6

(192) Stability of Toxin in Compositions with No Tonicity Agent

(193) Five liquid compositions were prepared by mixing an appropriate aliquot of BoNT/A (100 U/mL or 150 U/mL) with solutions comprising or lacking poloxamer P188 (4 w/w %), 8% w/w trehalose, and 0.2 w/w % L-methionine, in 20 mM histidine buffer at pH 6.0. Details of the compositions are given in Table 6.1:

(194) TABLE-US-00015 TABLE 6.1 Formulation Toxin Trehalose Poloxamer Methionine No. U/mL % P188% % Buffer Formulation 100 8 4 0 20 mM His, 33 pH 6.0 Formulation 150 0 0 0.2 20 mM His, 34 pH 6.0 Formulation 150 8 0 0.2 20 mM His, 35 pH 6.0 Formulation 100 0 4 0.2 20 mM His, 36 pH 6.0 Formulation 100 8 4 0.2 20 mM His, 37 pH 6.0

(195) Potency of Formulations 33-37 was tested by cell based potency assay prior to storage (time zero, t0) and after 1 month of storage at -70° C. and 40° ° C. for 1 month. Potency test results are shown in Table 8.

Example 7

(196) Stability of Toxin in Liquid Compositions

(197) Liquid compositions were prepared by mixing an appropriate aliquot of BoNT/A (100 U/mL) with solutions comprising poloxamer P188 (4 w/w %) or polysorbate (TWEEN®-20, 0.04 w/w %), sucrose or trehalose (8% w/w) or NaCl (0.9 w/w %), and 0.2 w/w % L-methionine, in water or 20 mM histidine buffer at pH 6.0. Details of the compositions are given in Table 7.1:

(198) TABLE-US-00016 TABLE 7.1 Polysorbate Buffer* Toxin Treh Suc NaCl P188 (TWEEN®20) Met or Formulation No. U/mL % % % % % water Formulation 31 100 8 4 0.2 buffer Formulation 38 100 8 4 0.2 water Formulation 39 100 0.9 4 0.2 buffer Formulation 40 100 8 0.04 0.2 buffer Tre = trehalose; Suc = sucrose; P-188 = poloxamer P188; Met = L-methionine *Buffer = 20 mM His, pH 6.0

(199) Potency of Formulations 31 and 38-40 was tested by a cell based potency assay prior to storage (time zero, t0) and after 5.5 months, 7.5 months and 12 months of storage at 5° C. Potency test results are shown in Table 9.

Example 8

(200) Stability of Toxin in Liquid Compositions with No Tonicity Agent

(201) Liquid compositions were prepared by mixing an appropriate aliquot of BoNT/A (200 U/mL) with solutions comprising poloxamer P188 (4 w/w %) or polysorbate (TWEEN®-20, 0.04 w/w %) and 0.2 w/w % L-methionine, in 20 mM histidine buffer at pH 6.0. Details of the compositions are given in Table 8.1.

(202) TABLE-US-00017 TABLE 8.1 Toxin Polysorbate Formulation No. U/mL P188% (TWEEN® 20) % methionine % Buffer Formulation 41 200 0 0.04 0.2 20 mM His, pH 6.0 Formulation 42 200 4 0 0.2 20 mM His, pH 6.0 P-188 = poloxamer P188; Met = L-methionine

(203) Potency of Formulations 41 and 42 was tested by a cell based potency assay prior to storage (time zero, t0) and 2 weeks of storage at -70° C. and 40° ° C. Potency test results are shown in Table 10.

Example 9

(204) Impact of Exemplary Lyoprotectors and Surfactants on the Stability of Exemplary Lyophilized Formulations

(205) Lyophilized compositions were prepared as follows. Bulk solutions of botulinum toxin were prepared by mixing an appropriate aliquot of a botulinum toxin type A (BoNT/A, 200 U/mL) with solutions comprising poloxamer P188 (4 w/w %) or polysorbate (TWEEN®20, 0.04 w/w %), with trehalose or sucrose (8 w/w %), and 0.2 w/w % L-methionine, in 20 mM histidine buffer at pH 6.0. Details of the compositions are given in Table 9.1. The solutions were filled into glass vials and lyophilized using conventional freeze-drying conditions.

(206) TABLE-US-00018 TABLE 9.1 Toxin Target TWEEN ® Poloxamer Potency Trehalose Sucrose 20 P188 Methionine Formulation No. U/vial % % % % Buffer Formulation 43 200 8 0 4 0.2 20 mM His, pH 6.0 Formulation 44 200 8 0.04 0 0.2 20 mM His, pH 6.0 Formulation 45 200 0 8 0 4 0.2 20 mM His, pH 6.0 Formulation 46 200 0 8 0.04 0 0.2 20 mM His, pH 6.0

(207) Potency of the lyophilized compositions, designated as Formulations 43-46, was tested by a cell based potency assay after reconstitution of the lyophiles subsequent to storage for 2 weeks at -20° C. and 40° C. Potency test results are shown in Table 12.

Example 10

(208) Method for Treating Depression

(209) A lyophilized composition comprised of botulinum toxin type A (BoNT/A, 200U), 8 wt % trehalose, 4 wt % poloxamer P188, 0.2 wt % methionine, in 20 mM histidine buffer, pH 6.0, is prepared as set forth in Example 9. The lyophilized composition is reconstituted in saline prior to administration.

(210) A 50-year-old male reports a history of depression for two years. He is presently taking antidepressant medications (PAXIL® 20 mg orally per day and WELLBUTRIN 225 mg orally per day). The male subject reports continued depression. He is treated by injecting into the frown/glabellar lines 30 Unit equivalents of the reconstituted composition. Four months later he reports a much-improved mood with no feelings of depression. His frown/glabellar lines are re-treated with 30 Unit equivalents of the reconstituted composition.

Example 11

(211) Method for Treating Cardia Arrhythmia

(212) A formulation similar to Formulation 1 as described in Example 1 is prepared. The lyophilized composition is comprised of botulinum toxin type A (BoNT/A, 50U), 2 wt % trehalose, 4 wt % poloxamer P188, 0.3 wt % NAC, in 20 mM histidine buffer, pH 5.5. The lyophilized composition is reconstituted in saline prior to administration.

(213) A Caucasian male is undergoing cardiac surgery. As part of the surgical procedure, an amount of the reconstituted composition to provide a total dose of 75 U botulinum toxin is evenly divided for administration via injection into three epicardial fat pads of his heart. The medical staff reports no arrhythmia during or post-surgery.

(214) Many alterations and modifications may be made by those having ordinary skill in the art, without departing from the spirit and scope of the disclosure. Therefore, it must be understood that the described embodiments have been set forth only for the purposes of examples, and that the embodiments should not be taken as limiting the scope of the following claims. The following claims are, therefore, to be read to include not only the combination of elements which are literally set forth, but all equivalent elements for performing substantially the same function in substantially the same way to obtain substantially the same result. The claims are thus to be understood to include those that have been described above, those that are conceptually equivalent, and those that incorporate the ideas of the disclosure.

Claims

1. A method of making a stable solid pharmaceutical composition which is free of animal protein excipients, the method comprising: compounding a Clostridial botulinum toxin active ingredient with two or more non-protein excipients to form a compounded mixture; wherein the two or more non-protein excipients comprise: (i) a poloxamer surfactant; and (ii) about 0.01 to 5 wt % methionine; and lyophilizing or vacuum drying the compounded mixture to produce the solid pharmaceutical composition; wherein the composition is suitable for subcutaneous or intramuscular injection after reconstitution with an aqueous carrier.
2. The method of claim 1, wherein the Clostridial botulinum toxin active ingredient is botulinum toxin serotype E (BoNT/E).
3. The method of claim 2, wherein the composition comprises about 1 unit to about 1000 mouse LD.sub.50 units of BoNT/E.
4. The method of claim 2, wherein the composition comprises about 1 ng to about 1 µg of BoNT/E.
5. The method of claim 2, wherein the composition further comprises NaCl.
6. The method of claim 2, wherein the poloxamer surfactant comprises poloxamer 188.
7. The method of claim 2, wherein the composition further comprises a disaccharide.
8. The method of claim 7, wherein the disaccharide is selected from sucrose or trehalose.
9. The method of claim 7, wherein the disaccharide comprises trehalose.
10. The method of claim 2, wherein the composition comprises a buffering agent.
11. The method of claim 10, wherein the buffering agent comprises histidine buffer.
12. The method of claim 2, wherein the composition comprises: (i) botulinum toxin serotype E (BoNT/E); (ii) about 0.5 to 8 wt % of a poloxamer surfactant; and (iii) about 0.01 to 5 wt % methionine.
13. The method of claim 2, wherein the composition comprises: (i) botulinum toxin serotype E (BoNT/E); (ii) histidine buffer; (iii) about 0.5 to 8 wt % of a poloxamer surfactant; and (iv) about 0.01 to 5 wt % methionine.
14. The method of claim 2, wherein the composition comprises: (i) botulinum toxin serotype A (BoNT/A); (ii) histidine buffer; (iii) about 0.5 to 8 wt % Poloxamer 188; and (iv) about 0.01 to 5 wt % methionine.
15. The method of claim 2, wherein the composition comprises: (i) botulinum toxin serotype E (BoNT/E); (ii) histidine buffer; (iii) about 0.5 to 8 wt % Poloxamer 188; (iv) about 0.01 to 5 wt % methionine; and (v) trehalose.
16. The method of claim 2, wherein the composition comprises: (i) botulinum toxin serotype E (BoNT/E); (ii) histidine buffer; (iii) about 4 wt % Poloxamer 188; (iv) about 0.2 wt % methionine; (v) about 8 wt % trehalose.
17. The method of claim 16, wherein the composition comprises: (i) botulinum toxin serotype E (BoNT/E); (ii) histidine buffer; (iii) about 4 wt % Poloxamer 188; (iv) about 0.2 wt % methionine; (v) about 8 wt % trehalose, and, wherein the composition comprises about 1 ng to about 1 µg of BoNT/E, and wherein the composition is lyophilized.
18. The method of claim 2, wherein the method comprises lyophilizing the compounded mixture.
19. The method of claim 2, wherein the method comprises vacuum drying the compounded mixture.
20. A method of making a stable solid pharmaceutical composition which is free of animal protein excipients, the method comprising: compounding a Clostridial botulinum toxin active ingredient with three or more non-protein excipients to form a compounded mixture; wherein the three or more non-protein excipients comprise: (i) a poloxamer surfactant, (ii) methionine, and (iii) trehalose, and lyophilizing or vacuum drying the compounded mixture to produce the solid pharmaceutical composition; wherein the toxin has a potency recovery of at least about 75% or at least about 80% after storage at two weeks at —20° C. and/or at 40° C., and wherein the composition is suitable for subcutaneous or intramuscular injection after reconstitution with an

aqueous carrier.

21. The method of claim 20, wherein the Clostridial botulinum toxin active ingredient is botulinum toxin serotype E (BoNT/E).
