



US012410249B2

(12) **United States Patent**
Cicic

(10) **Patent No.:** **US 12,410,249 B2**

(45) **Date of Patent:** ***Sep. 9, 2025**

(54) **LOW DOSE ANTIBODY-BASED METHODS
FOR TREATING HEMATOLOGIC
MALIGNANCIES**

(71) Applicant: **Actinium Pharmaceuticals, Inc.**, New
York, NY (US)

(72) Inventor: **Dragan Cicic**, Brooklyn, NY (US)

(73) Assignee: **Actinium Pharmaceuticals, Inc.**, New
York, NY (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 870 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **17/679,213**

(22) Filed: **Feb. 24, 2022**

(65) **Prior Publication Data**

US 2022/0251196 A1 Aug. 11, 2022

Related U.S. Application Data

(63) Continuation of application No. 16/300,672, filed as
application No. PCT/US2017/034406 on May 25,
2017, now Pat. No. 11,292,835.

(60) Provisional application No. 62/342,568, filed on May
27, 2016.

(51) **Int. Cl.**

C07K 16/28 (2006.01)

A61K 39/00 (2006.01)

A61K 51/10 (2006.01)

A61P 35/02 (2006.01)

(52) **U.S. Cl.**

CPC **C07K 16/2803** (2013.01); **A61K 51/1069**
(2013.01); **A61K 51/1093** (2013.01); **A61K**
51/1096 (2013.01); **A61P 35/02** (2018.01);
A61K 2039/505 (2013.01)

(58) **Field of Classification Search**

CPC **C07K 16/2896**; **A61K 51/1093**; **A61K**
51/1096; **A61K 51/1069**; **A61P 35/02**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,923,985 A 5/1990 Gansow et al.
6,683,162 B2 1/2004 Scheinberg et al.
11,292,835 B2 4/2022 Cicic

OTHER PUBLICATIONS

The abstract of Jurcic et al (Journal of Clinical Oncology, May 20,
2015, vol. 33, No. 15, Suppl 1, abstract No. 7050) (Year: 2015).
Thol et al (Blood, 2015, vol. 126, pp. 319-327). (Year: 2015).
The abstract of McDevitt et al (Blood, Nov. 18, 2011), vol. 118, No.
21, abstract No. 768) (Year: 2011).*

Burnett, et al. "A comparison of low-dose cytarabine and hydroxyurea
with or without all-trans retinoic acid for acute myeloid leukemia
and high-risk myelodysplastic syndrome in patients not considered
fit for intensive treatment." Cancer 109.6 (2007): 1114-1124.

Caron, et al., "A Phase IB Trial of Humanized Monoclonal Antibody
M195 (Anti-CD33) in Myeloid Leukemia: Specific Targeting With-
out Immunogenicity", Blood, 1994, vol. 83, pp. 1760-1768.

Co, et al., "Chimeric and humanized antibodies with specificity for
the CD33 antigen.", J. Immunol. 148: 1149-1154, 1992.

De Kruijff, et al., "A Critical Review of Alpha Radionuclide
Therapy—How to Deal with Recoiling Daughters?", Pharmaceuti-
cals, Jun. 2015, ISSN 1424-8247, 321-336.

Deng, et al., "Monoclonal antibodies: what are the pharmacokinetic
and pharmacodynamic considerations for drug development?", Expert
Opinion on Drug Metabolism and Toxicology, 2012, vol. 8, pp.
141-160.

Dutta, et al., "The Expression Pattern of CD33 Antigen Can
Differentiate Leukemic from Normal Progenitor Cells in Acute
Myeloid Leukemia", Indian Journal of Hematology and Blood
Transfusion, 2014, vol. 30, pp. 130-134.

Harousseau, et al., "A randomized phase 3 study of tipifarnib
compared with best supportive care, including hydroxyurea, in the
treatment of newly diagnosed acute myeloid leukemia in patients 70
years or older", Blood, Aug. 6, 2009, vol. 114, No. 6, 1166-1173.
Jurcic, et al., "Phase I Trial of Targeted Alpha-Particle Therapy with
Actinium-225 (225Ac)-Lintuzumab and Low-Dose Cytarabine (LDAC)
in Patients Age 60 or Older with Untreated Acute Myeloid Leuke-
mia (AML)", The American Society of Hematology, Abstract, vol.
128, Issue 22, Dec. 2, 2016, 7 pgs.

Jurcic, et al. "Targeted alpha-particle nano-generator actinium-225
(225Ac)-lintuzumab (anti-CD33) in acute myeloid leukemia (AML)." Clinical Lymphoma, Myeloma and Leukemia 13 (2013): S379-
S380.

Jurcic, et al. "Phase I trial of the targeted alpha-particle nano-
generator actinium-225 (225Ac)-lintuzumab (anti-CD33) in com-
bination with low-dose cytarabine (LDAC) for older patients with
untreated acute myeloid leukemia (AML)." Blood (2013) 122(21):
1460-1460.

Krause, et al., "Differential regulation of myeloid leukemias by the
bone marrow microenvironment", Nature Medicine, 2013, vol. 19,
pp. 1513-1517.

(Continued)

Primary Examiner — Karen A. Canella

(74) *Attorney, Agent, or Firm* — Alan J. Morrison

(57)

ABSTRACT

This invention provides a method for treating a subject
afflicted with a hematologic malignancy comprising admin-
istering to the subject an agent targeting a hematologic
malignancy-associated antigen, wherein the subject has a
low peripheral cancerous cell burden. This invention also
provides a method for treating a subject afflicted with a
hematologic malignancy and having a high peripheral can-
cerous cell burden, comprising (i) medically lowering the
subject's peripheral cancerous cell burden, and (ii) while the
subject's peripheral cancerous cell burden is still low,
administering to the subject an agent targeting a hematologic
malignancy-associated antigen. Particularly envisioned are
the subject methods for treating acute myeloid leukemia
using an anti-CD33 antibody labeled with an alpha-emitting
isotope, such as ²²⁵Ac-HuM195.

13 Claims, 7 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

Kumar, "Genetic Abnormalities and Challenges in the Treatment of Acute Myeloid Leukemia", *Genes & Cancer*, vol. 2, No. 2, 95-107, 2011.

Maguire, et al., "Efficient 1-Step Radiolabeling of Monoclonal Antibodies to High Specific Activity with ^{225}Ac for α -Particle Radioimmunotherapy of Cancer", *Journal of Nuclear Medicine*, 2014, vol. 55, pp. 1492-1498.

McDevitt, et al., "Design and synthesis of ^{225}Ac radioimmunopharmaceuticals", *Applied Radiation and Isotopes*, 57 (2002) 841-847.

Mulford et al., "The Promise of Targeted α -Particle Therapy", *The Journal of Nuclear Medicine*, vol. 46, No. 1 (Suppl), Jan. 2005, 199S-204S.

Pollard, et al., "Correlation of CD33 expression level with disease characteristics and response to gemtuzumab ozogamicin containing chemotherapy in childhood AML", *Blood*, Apr. 1, 2012, 119(16):3705-3711.

Rosenblat, et al., "Sequential cytarabine and α -particle immunotherapy with bismuth-213-lintuzumab (H uM 195) for acute myeloid leukemia", *Clinical Cancer Research* 16.21 (2010): 5303-5311.

SWOG Oncology Research Professional (ORP) Manual, vol. I, Chapter 11A, Leukemia (2014).

Topp, et al., "Targeted therapy with the T-cell—engaging antibody blinatumomab of chemotherapy-refractory minimal residual disease in B-lineage acute lymphoblastic leukemia patients results in high response rate and prolonged leukemia-free survival", *Journal of Clinical Oncology*, vol. 29, No. 18, Jun. 20, 2011, pp. 2493-2498.

V.H.J. van der Velden, et al., "High CD33-antigen loads in peripheral blood limit the efficacy of gemtuzumab ozogamicin (Mylotarg) treatment in acute myeloid leukemia patients", *Leukemia*, 2004, May 18(5):983-988.

Walter, et al., "Acute myeloid leukemia stem cells and CD33-targeted immunotherapy", *Blood*, 2012, vol. 119, pp. 6198-6208.

Zhu, et al., "Blinatumomab, a bispecific T-cell engager (BiTE®) for CD-19 targeted cancer immunotherapy: clinical pharmacology and its implications", *Clinical pharmacokinetics* 55.10 (2016): 1271-1288.

* cited by examiner




Figure 1

624 TCTAGACCACCATGGAGAAAGACACACTCCTGCTATGGGTCTACTTCTCTGGGTTCAGGTTCCACAGGTGACATTTCAG
M E K D T L L L W V L L L W V P G S T G D I Q
704 ATGACCCAGTCTCCGAGCTCTCTGTCCGCATCAGTAGGAGACAGGGTCACCATCACATGCAGAGCCAGCGAAAGTGTCGA
M T Q S P S S L S A S V G D R V T I T C R A S E S V D
784 CAATTATGGCATAGCTTTATGAAGTGGTTCCAACAGAAACCCGGGAAGGCTCCTAAGCTTCTGATTTACGCTGCATCCA
N Y G I S F M N W F Q Q K P G K A P K L L I Y A A S
864 ACCAAGGCTCCGGGGTACCTCTCGCTTCTCAGGCAGTGGATCTGGGACAGACTTCACTCTCACCATTTCATCTCTGCAG
N Q G S G V P S R F S G S G S G T D F T L T I S S L Q
944 CCTGATGACTTCGCAACCTATTACTGTCAGCAAAGTAAGGAGGTTCCGTGGACGTTCCGTCAAGGGACCAAGGTGGAGAT
P D D F A T Y Y C Q Q S K E V P W T F G Q G T K V E I
1024 CAAACGTAAGTAGAATCCAAAGTCTAGAAATCTAAACTCTGAGGGGTCGGATGACGTGGCCATTCTTTGCCTAAAGCA
K R
1104 TTGAGTTTACTGCAAGGTCAGAAAAGCATGCAAAGCCCTCAGAATGGCTGCAAAGAGCTCCAACAAAACAATTTAGAACT
1184 TTATTAAGGAATAGGGGAAGCTAGGAAGAAACTCAAACATCAAGATTTTAAATACGCTTCTTGGTCTCCTTGCTATAA
1264 TTATCTGGGATAAGCATGCTGTTTTCTGTCTGTCCCTAACATGCTCTGTGATTATCCGCAAACAACACACCCAAGGGCAG
1344 AACTTTGTTACTTAAACACCATCCTGTTTGCTTCTTTCTCAGGAACTGTGGCTGCACCATCTGTCTTCATCTTCCCGCC
T V A A P S V F I F P P
1424 ATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTAC
S D E Q L K S G T A S V V C L L N N F Y P R E A K V
1504 AGTGAAGGTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTCACAGAGCAGGACAGCAAGGACAGCACCTAC
Q W K V D N A L Q S G N S Q E S V T E Q D S K D S T Y
1584 AGCCTCAGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCAGGG
S L S S T L T L S K A D Y E K H K V Y A C E V T H Q G
1664 CCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTTAGAGGGAGAAGTGCCCCACCTGCTCCTCAGTTCCA
L S S P V T K S F N R G E C .
1744 GCCTGACCCCTCCCATCCTTTGGCCTCTGACCCTTTTCCACAGGGGACCTACCCCTATTGCGGTCTCCAGCTCATCT
1824 TTCACCTCACCCCCCTCCTCCTCCTTGGCTTTAATTATGCTAATGTTGGAGGAGAATGAATAAATAAAGTGAATCTTTGC
1904 ACCTGTGGTTTCTCTCTTTCCTCATTTAATAATTATTATCTGTTGTTTTACCAACTACTCAATTTCTCTTATAAGGGACT
1984 AAATATGTAGTCATCCTAAGGCGCATAACCATTATATAAAATCATCCTTCATTCTATTTTACCCTATCATCCTCTGCAAG
2064 ACAGTCTCCCTCAAACCCACAAGCCTTCTGTCTCAGCTCCCCTGGGCCATGGTAGGAGAGACTTGCTTCCTTGTTTT
2144 CCCCTCTCAGCAAGCCCTCATAGTCCTTTTAAAGGTGACAGGTCTTACAGTCATATATCCTTTGATTCAATTCCCTGA
2224 GAATCAACCAAAGCAAATTTTTCAAAGAAGAAACCTGCTATAAAGAGAATCATTCATTGCAACATGATATAAAATAACA
2304 ACACAATAAAAGCAATTAAATAAACAACAATAGGGAAATGTTTAAAGTTCATCATGGTACTTAGACTTAATGGAATGTCA
2384 TGCCTTATTTACATTTTTAAACAGGTACTGAGGGACTCCTGTCTGCCAAGGGCCGTATTGAGTACTTCCACAACCTAAT
2464 TTAATCCACACTATACTGTGAGATTAAAAACATTCATTAAAAATGTTGCAAAGGTTCTATAAAGCTGAGAGACAAATATAT
2544 TCTATAACTCAGCAATCCCACTTCTAGGATCC

Figure 2

624 TCTAGACCACCATGGGATGGAGCTGGATCTTTCTCTTCTCCTCTGTGAGAACTGCTGGCGTCCACTCTCAGGTTCACTG
M G W S W I F L F L L S G T A G V H S Q V Q L
704 GTGCAGTCTGGAGCTGAGGTGAAGAAGCCTGGGAGCTCAGTGAAGGTTTCTGCAAAGCTTCTGGCTACACCTTCACTGA
V Q S G A E V K K P G S S V K V S C K A S G Y T F T D
784 CTACAAATGCACTGGGTGAGGCAGGCTCCTGGCCAAGGCTGGAATGCATTGGATATATTTATCCTTACAATGGTGTA
Y N M H W V R Q A P G Q G L E W I G Y I Y P Y N G G
864 CCGGCTACAACCAGAAGTCAAGAGCAAGGCCACAATTACAGCAGACGAGAGTACTAACACAGCCTACATGGAACCTCTCC
T G Y N Q K F K S K A T I T A D E S T N T A Y M E L S
944 AGCCTGAGGTCTGAGGACACTGCAGTCTATTACTGCGCAAGAGGGCGCCCCGCTATGGACTACTGGGGCCAAGGGACTCT
S L R S E D T A V Y Y C A R G R P A M D Y W G Q G T L
1024 GGTCACTGTCTCTTTCAGGTAAGAATGGCCTCTAGACCACCATGGGATGGAGCTTCTGGGGCAGGCCAGGCCTGACCTTG
V T V S S
1104 GCTTTGGGGCAGGGAGGGGGCTAAGGTGAGGCAGGTGGCGCCAGCCAGGTGCACACCCAATGCCCATGAGCCCAGACACT
1184 GGACGCTGAACCTCGCGGACAGTTAAGAACCAGGGGCCCTCTGCGCCCTGGGGCCAGCTCTGTCCCACACCGCGGTCA
1264 TGGCACCACCTCTCTTGCAGCCTCCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCTCCTCCAAGAGCACCTCTGGGG
A S T K G P S V F P L A P S S K S T S G
1344 GCACAGCGGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGCTGTCTGGAACCTCAGGCGCCCTGACC
G T A A L G C L V K D Y F P E P V T V S W N S G A L T
1424 AGCGGCGTGCACACCTTCCCGGCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAG
S G V H T F P A V L Q S S G L Y S L S S V V T V P S S
1504 CAGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACAAGCCAGCAACACCAAGGTGGACAAGAAAGTTGGTGAGA
S L G T Q T Y I C N V N H K P S N T K V D K K V
1584 GGGCAGCACAGGGAGGGAGGGTGTCTGCTGGAAGCCAGGCTCAGCGCTCCTGCCCTGGACGCATCCCGGCTATGCAGCCCC
1664 AGTCCAGGGCAGCAAGGCAGGCCCCGTCTGCCTCTTACCCGGAGGCCTCTGCCCCCCCCACTCATGCTCAGGGAGAGGG
1744 TCTTCTGGCTTTTTCAGGCTCTGGGCAGGCACAGGCTAGGTGCCCTAACCAGGCCCTGCACACAAAGGGGCAGGT
1824 GCTGGGCTCAGACCTGCCAAGAGCCATATCCGGGAGGACCCTGCCCCTGACCTAAGCCCACCCCAAGGCCAAACTCTCC
1904 ACTCCCTCAGCTCGGACACCTTCTCTCTCTCCAGATTCCAGTAACCTCCAATCTTCTCTCTGCAGAGCCCAATCTTG
E P K S C
1984 ACAAACCTCACACATGCCCACCGTGCCAGGTAAGCCAGCCAGGCCTCGCCCTCAGCTCAAGGCGGGACAGGTGCCCT
D K T H T C P P C P
2064 AGAGTAGCCTGCATCCAGGACAGGCCCCAGCCGGGTGCTGACACGTCCACCTCCATCTTCTCCTCAGCACCTGAACTCC
A P E L
2144 TGGGGGACCGTCAGTCTTCTCTTCCCCCAAAACCAAGGACACCCCTCATGATCTCCCGGACCCCTGAGGTCACATGC
L G G P S V F L F P P K P K D T L M I S R T P E V T C
2224 GTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAA
V V V D V S H E D P E V K F N W Y V D G V E V H N A K
2304 GACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCCTACCGTCTGCACCAGGACTGGCTGA
T K P R E E Q Y N S T Y R V V S V L T V L H Q D W L
2384 ATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAGCCCTCCAGCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGT
N G K E Y K C K V S N K A L P A P I E K T I S K A K
2464 GGGACCCGTGGGGTGGCAGGGCCACATGGACAGAGGGCCGCTCGGCCACCCCTGCCCCTGAGAGTGACCGCTGTACCAA

Figure 3

2544 CCTCTGTCCCTACAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAG
G Q P R E P Q V Y T L P P S R D E L T K N Q
2624 GTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAA
V S L T C L V K G F Y P S D I A V E W E S N G Q P E N
2704 CAACTACAAGACCACGCCTCCCGTGTGGACTCCGACGGCTCCTTCTCCTCTACAGCAAGCTCACCGTGGACAAGAGCA
N Y K T T P P V L D S D G S F F L Y S K L T V D K S
2784 GGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCC
R W Q Q G N V F S C S V M H E A L H N H Y T Q K S L S
2864 CTGTCTCCGGGTAAATGAGTGCGACGGCCGGCAAGCCCCGCTCCCCGGGCTCTCGCGGTGCGACGAGGATGCTTGGCAC
L S P G K *
2944 GTACCCCTGTACATACTTCCCGGGCGCCAGCATGGAATAAAGCAACCAGCGCTGCCCTGGGCCCCGTGCGAGACTGTG
3024 ATGGTTCTTTCCACGGGTCAAGGCCGAGTCTGAGGCCTGAGTGGCATGAGGGAGGCAGAGCGGGTCCCCTGTCCCCACAC
3104 TGGCCCAAGCTGTGCAGGTGTGCCTGGGCCGCTAGGGTGGGGCTCAGCCAGGGGCTGCCCTCGGCAGGGTGGGGGATTT
3184 GCCAGCGTGGCCCTCCCTCCAGCAGCACCTGCCCTGGGCTGGGCCACGGGAAGCCCTAGGAGCCCCTGGGGACAGACACA
3264 CAGCCCCCTGCCTCTGTAGGAGACTGTCTGTTCTGTGAGCGCCCTGTCTCCGACCTCCATGCCCACTCGGGGGCATGCC
3344 TAGTCCATGTGCGTAGGGACAGGCCCTCCCTCACCCTATCTACCCCCACGGCACTAACCCTGGCTGCCCTGCCAGCCTC
3424 GCACCCGATGGGGACACAACCGACTCCGGGGACATGCACTCTCGGGCCCTGTGGAGGGACTGGTGCAGATGCCACACA
3504 CAACTCAGCCCAGACCCGTTCAACAAACCCGCACTGAGGTTGGCCGGCCACACGGCCACCACACACACAGTGCACGC
3584 CTCACACACGGAGCCTCAGCCGGGGAAGTGCACAGCAGCCAGACCAGAGCAAGGTCCTCGCACACGTGAACACTCCTCG
3664 GACACAGCCCCCAGAGCCCCACGGGCACCTCAAGGCCACGAGCCTCTCGGCAGCTTCTCCACATGCTGACCTGCTC
3744 AGACAAACCCAGCCCTCCTCTCACAAGGGTGGCCCTGCAGCCGCCACACACACAGGGGATCACACACCACGTACGTC
3824 CCTGGCCCTGGCCCACTTCCCAGTGCCGCCCTTCCCTGCAGGATCC

Figure 3
Continued




Figure 4




Figure 5

Protocol Schema




Figure 6

1

LOW DOSE ANTIBODY-BASED METHODS FOR TREATING HEMATOLOGIC MALIGNANCIES

This application is a continuation of U.S. Ser. No. 16/300, 672, filed Nov. 12, 2018, which is a § 371 national stage entry of PCT Application No. PCT/US2017/034406, filed May 25, 2017, which claims priority of U.S. Provisional Application No. 62/342,568, filed May 27, 2016, the contents of all of which are incorporated herein by reference in their entirety.

Throughout this application, various publications are cited. The disclosure of these publications is hereby incorporated by reference into this application to describe more fully the state of the art to which this invention pertains.

FIELD OF THE INVENTION

The present invention relates to treating a subject afflicted with a hematologic malignancy using a low dose of a potent agent (such as a toxin-conjugated antibody) targeting a hematologic malignancy-associated antigen, wherein the subject has a low peripheral cancerous cell burden.

BACKGROUND OF THE INVENTION

Cytotoxic agent-conjugated antibodies have recently become a promising tool for treating hematologic malignancies such as acute myeloid leukemia ("AML"). Of particular interest are therapeutic conjugates of cancer cell-specific antibodies with potent cytotoxic agents like alpha-emitting isotopes such as 225-actinium (²²⁵Ac).

However, doses of monoclonal antibodies labeled with potent cytotoxic agents cannot safely be escalated above maximum tolerated doses established in clinical trials. This safety-based dosing constraint prevents the escalation of antibody conjugate doses to saturation levels.

Therefore, for these conjugates, there is an unmet need for treating AML and other hematologic cancers by administering them in doses low enough to avoid toxicity while high enough to be therapeutically effective. There is also an unmet need to achieve this balance in a manner independent of patient age, co-morbidities and disease severity.

SUMMARY OF THE INVENTION

This invention provides a method for treating a subject afflicted with a hematologic malignancy comprising administering to the subject an agent targeting a hematologic malignancy-associated antigen, wherein the subject has a low peripheral cancerous cell burden.

This invention also provides a method for treating a subject afflicted with a hematologic malignancy and having a high peripheral cancerous cell burden, comprising (i) medically lowering the subject's peripheral cancerous cell burden, and (ii) while the subject's peripheral cancerous cell burden is still low, administering to the subject an agent targeting a hematologic malignancy-associated antigen.

This invention further provides a method for treating a human subject afflicted with acute myeloid leukemia comprising administering to the subject an anti-CD33 antibody labeled with an alpha-emitting isotope, wherein (i) the subject has a low peripheral blast burden, and (ii) the antibody is administered in sub-saturating dose.

This invention still further provides a method for treating a human subject afflicted with acute myeloid leukemia and having a high peripheral blast burden, comprising (i) medi-

2

cally lowering the subject's peripheral blast burden, and (ii) while the subject's peripheral blast burden is still low, administering to the subject an anti-CD33 antibody labeled with an alpha-emitting isotope, wherein the antibody is administered in a sub-saturating dose.

This invention still further provides a method for treating a human subject afflicted with acute myeloid leukemia comprising intravenously administering ²²⁵Ac-labeled HuM195 to the subject, wherein (i) the subject has a low peripheral blast burden, and (ii) the ²²⁵Ac-labeled HuM195 is administered in a sub-saturating dose.

Finally, this invention provides a method for treating a human subject afflicted with acute myeloid leukemia and having a high peripheral blast burden, comprising (i) medically lowering the subject's peripheral blast burden, and (ii) while the subject's peripheral blast burden is still low, intravenously administering ²²⁵Ac-labeled HuM195 to the subject, wherein the ²²⁵Ac-labeled HuM195 is administered in a sub-saturating dose.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1

This figure shows a schematic diagram of the expression plasmids for HuM195. The humanized VL and VH exons of HuM195 are flanked by XbaI sites. The VL exon was inserted into mammalian expression vector pVk, and the VH exon into pVg1 (Co, et al., J. Immunol. 148:1149-1154, 1992).

FIG. 2

This figure shows the complete sequence of the HuM195 light chain gene cloned in pVk between the XbaI and BamHI sites. The nucleotide number indicates its position in the plasmid pVk-HuM195. The VL and CK exons are translated in single letter code; the dot indicates the translation termination codon. The mature light chain begins at the double-underlined aspartic acid (D). The intron sequence is in italics. The polyA signal is underlined.

FIG. 3

This figure shows the complete sequence of the HuM195 heavy chain gene cloned in pVg1 between the XbaI and BamHI sites. The nucleotide number indicates its position in the plasmid pVg1-HuM195. The VH, CH1, H, CH2 and CH3 exons are translated in single letter code; the dot indicates the translation termination codon. The mature heavy chain begins at the double-underlined glutamine (Q). The intron sequences are in italics. The polyA signal is underlined.

FIG. 4

This figure shows the structure of ²²⁵Ac-Lintuzumab (²²⁵Ac-HuM195).

FIG. 5

This figure shows a flowchart for the production of ²²⁵Ac-HuM195.

FIG. 6

This figure shows a dosing protocol for ²²⁵Ac-Lintuzumab (²²⁵Ac-HuM195) treatment of AML.

DETAILED DESCRIPTION OF THE INVENTION

This invention provides surprisingly effective methods for treating a subject afflicted with a hematologic malignancy, such as AML. These methods use a potent agent, such as an antibody conjugated with an alpha-emitting isotope, targeting a hematologic malignancy-associated antigen, such as CD33. The subject who is being treated has a low peripheral

cancerous cell burden and, ideally, only a sub-saturating dose of agent is needed to treat the subject.

Definitions

In this application, certain terms are used which shall have the meanings set forth as follows.

As used herein, “administer”, with respect to an agent, means to deliver the agent to a subject’s body via any known method. Specific modes of administration include, without limitation, intravenous, oral, sublingual, transdermal, subcutaneous, intraperitoneal, intrathecal and intra-tumoral administration.

In addition, in this invention, the various antibodies and other antigen-targeting agents used can be formulated using one or more routinely used pharmaceutically acceptable carriers. Such carriers are well known to those skilled in the art. For example, injectable drug delivery systems include solutions, suspensions, gels, microspheres and polymeric injectables, and can comprise excipients such as solubility-altering agents (e.g., ethanol, propylene glycol and sucrose) and polymers (e.g., polycaprylactones and PLGA’s). Implantable systems include rods and discs, and can contain excipients such as PLGA and polycaprylactone.

As used herein, the “agent” targeting a hematologic malignancy-associated antigen can be any type of compound or composition useful for such purpose. Types of agents include, without limitation, antibodies, other protein-based drugs, peptides, nucleic acids, carbohydrates and small molecules drugs.

As used herein, the term “antibody” includes, without limitation, (a) an immunoglobulin molecule comprising two heavy chains and two light chains and which recognizes an antigen; (b) polyclonal and monoclonal immunoglobulin molecules; (c) monovalent and divalent fragments thereof, and (d) bi-specific forms thereof. Immunoglobulin molecules may derive from any of the commonly known classes, including but not limited to IgA, secretory IgA, IgG and IgM. IgG subclasses are also well known to those in the art and include, but are not limited to, human IgG1, IgG2, IgG3 and IgG4. Antibodies can be both naturally occurring and non-naturally occurring. Furthermore, antibodies include chimeric antibodies, wholly synthetic antibodies, single chain antibodies, and fragments thereof. Antibodies may be human, humanized or nonhuman.

As used herein, an “anti-CD33 antibody” is an antibody that binds to any available epitope of CD33. In one embodiment, the anti-CD33 antibody binds to the epitope recognized by the antibody HuM195.

As used herein, the term “burden”, when used in connection with a cancerous cell, means quantity. So, a cancerous cell “burden” means the quantity of cancerous cells. Cancerous cells have a burden with respect to their tissue of origin (i.e., the primary site of disease), such as the “bone marrow blast burden” in the case of AML. Cancerous cells also have a burden with respect to one or more tissues other than those of origin, such as the blast burden in blood, liver and spleen in the case of AML. The term “peripheral burden” relates to such cells. The peripheral burden of cancerous cells, such as blasts in the case of AML, can be measured in different ways with different outcomes. For example, in the case of AML, the “peripheral blast burden” can be measured as the total blast population outside of the bone marrow, or the total blast population of the blood, spleen and liver combined, or simply the blast population of the blood as measured in cells per unit volume. As used herein in connection with AML and other cancers originat-

ing in the bone marrow, and unless stated otherwise, the term “peripheral cancerous cell burden” (e.g., peripheral blast burden) refers to the cancerous cell population of the blood as measured in cells per unit volume (e.g., cells/ μ l). This blood-based measurement is a useful proxy for the more cumbersome measurements of spleen and liver burdens, for example.

Herein, a peripheral cancerous cell burden in a subject is “high” if, when the subject is administered an agent (e.g., an antibody) targeting a hematologic malignancy-associated antigen at the maximum safe dose, the agent does not reach the primary site of disease in a sufficient amount to bind to more than 90% of its target antigens at that site. Conversely, a peripheral cancerous cell burden in a subject is “low” if, when the subject is administered an agent (e.g., an antibody) targeting a hematologic malignancy-associated antigen at the maximum safe dose, the agent reaches the primary site of disease in a sufficient amount to bind to more than 90% of its target antigens at that site. In the case of AML, examples of low peripheral blast burden are those yielding blood blast burdens at or below 1,000 blast cells/ μ l, at or below 500 blast cells/ μ l, at or below 400 blast cells/ μ l, at or below 300 blast cells/ μ l, at or below 200 blast cells/ μ l, at or below 100 blast cells/ μ l, and at or below 50 blast cells/ μ l.

As used herein, the term “cytotoxic agent” includes, without limitation, a radionuclide, a protein-based toxin, a non-protein based toxin, and a chemotherapeutic agent. Radionuclides include, for example, alpha-emitting isotopes (i.e., alpha-emitting isotopes, such as ^{225}Ac , ^{213}Bi and ^{213}Po), beta-emitting isotopes (e.g., ^{90}Y), and gamma-emitting isotopes (e.g., ^{137}Cs (which emits gamma rays via its decay product ^{137}Ba)). Protein-based toxins include, without limitation, ricin and toxic portions thereof, and botulinum toxin and toxic portions thereof. Methods for affixing a cytotoxic agent to an antibody (i.e., “labeling” an antibody with a cytotoxic agent) are well known.

A “hematologic malignancy”, also known as a blood cancer, is a cancer that originates in blood-forming tissue, such as the bone marrow or other cells of the immune system. Hematologic malignancies include, without limitation, leukemias (such as AML, acute promyelocytic leukemia, acute lymphoblastic leukemia, acute mixed lineage leukemia, chronic myeloid leukemia, chronic lymphocytic leukemia, hairy cell leukemia, large granular lymphocytic leukemia), myelodysplastic syndrome (MDS), myeloproliferative disorders (polycythemia vera, essential thrombocythosis, primary myelofibrosis and chronic myeloid leukemia), lymphomas, multiple myeloma, and MGUS and similar disorders.

As used herein, a “hematologic malignancy-associated antigen” can be, for example, a protein and/or carbohydrate marker found exclusively or predominantly on the surface of a cancer cell associated with that particular malignancy. Examples of hematologic malignancy-associated antigens include, without limitation, CD20, CD33, CD38, CD45, CD52, CD123 and CD319.

The antibody “HuM195” (also known as lintuzumab) is known, as are methods of making it. Likewise, methods of labeling HuM195 with ^{225}Ac are known. These methods are exemplified, for example, in Scheinberg, et al., U.S. Pat. No. 6,683,162. This information is also exemplified in the examples and figures below.

A low peripheral cancerous cell burden can be “medically induced” using any known method for doing so. These methods include, for example, pharmaceutical induction (such as by oral administration of hydroxyurea at 10-70 mg/kg/day), and spleen removal.

As used herein, the term “subject” includes, without limitation, a mammal such as a human, a non-human primate, a dog, a cat, a horse, a sheep, a goat, a cow, a rabbit, a pig, a rat and a mouse. Where the subject is human, the subject can be of any age. For example, the subject can be 60 years or older, 65 or older, 70 or older, 75 or older, 80 or older, 85 or older, or 90 or older. For a human subject afflicted with AML, the subject’s bone marrow blast burden can be, for example, over 5%, over 10%, over 20%, over 30%, over 40%, over 50%, over 60%, over 70%, over 80%, or over 90%. As an additional example, for a human subject afflicted with AML, the subject can be newly diagnosed, or relapsed and/or refractory, or in remission.

As used herein, a “sub-saturating dose” of an agent targeting a hematologic malignancy-associated antigen is one that introduces into the subject’s body fewer target antigen-binding sites (e.g., Fab’s) than there are target antigens. By way of example, for an anti-CD33 antibody, a sub-saturating dose is one that introduces into the subject’s body fewer CD33-binding sites than there are CD33 molecules. In one embodiment, a sub-saturating dose of an agent targeting a hematologic malignancy-associated antigen is one where the ratio of target antigen-binding sites to target antigens is less than or equal to 9:10. In another embodiment, the ratio of target antigen-binding sites to target antigens is less than or equal to 1:2, less than or equal to 1:5, less than or equal to 1:10, less than or equal to 1:20, or less than or equal to 1:100.

For an agent such as an antibody labeled with an alpha-emitting isotope or other cytotoxic agent, the majority of the drug administered to a subject typically consists of non-labeled antibody, with the minority being the labeled antibody. Thus, in one embodiment, a sub-saturating dose of an agent targeting a hematologic malignancy-associated antigen is one where the ratio of total (i.e., labeled and unlabeled) target antigen-binding sites to target antigens is less than or equal to 9:10 (and can be less than or equal to 1:2, less than or equal to 1:5, less than or equal to 1:10, less than or equal to 1:20, or less than or equal to 1:100). In another embodiment, a sub-saturating dose of an agent targeting a hematologic malignancy-associated antigen is one where the ratio of labeled target antigen-binding sites to target antigens is less than or equal to 9:10 (and can be less than or equal to 1:2, less than or equal to 1:5, less than or equal to 1:10, less than or equal to 1:20, or less than or equal to 1:100).

Sub-saturating doses used in connection with this invention include, for example, a single administration, and two or more administrations (i.e., fractions). The amount administered in each dose can be measured, for example, by radiation (e.g., $\mu\text{Ci/kg}$) or weight (e.g., mg/kg or mg/m^2). In the case of ^{225}Ac -HuM195 (also known as “Actimab-A”) for treating AML, dosing regimens include the following, without limitation: (i) $2 \times 0.5 \mu\text{Ci/kg}$, $2 \times 1.0 \mu\text{Ci/kg}$, $2 \times 1.5 \mu\text{Ci/kg}$, or $2 \times 2.0 \mu\text{Ci/kg}$, where the fractions are administered one week apart; (ii) $1 \times 0.5 \mu\text{Ci/kg}$, $1 \times 1.0 \mu\text{Ci/kg}$, $1 \times 2.0 \mu\text{Ci/kg}$, $1 \times 3.0 \mu\text{Ci/kg}$, or $1 \times 4.0 \mu\text{Ci/kg}$; (iii) $1 \times 15\text{--}20 \mu\text{g/kg}$ ($0.03\text{--}0.06 \mu\text{g/kg}$ labeled); and (iv) less than or equal to approximately 2 mg per subject (approximately 0.04 mg labeled antibody per subject).

As used herein, “treating” a subject afflicted with a disorder shall include, without limitation, (i) slowing, stopping or reversing the disorder’s progression, (ii) slowing, stopping or reversing the progression of the disorder’s symptoms, (iii) reducing the likelihood of the disorder’s recurrence, and/or (iv) reducing the likelihood that the disorder’s symptoms will recur. In the preferred embodiment, treating a subject afflicted with a disorder means (i)

reversing the disorder’s progression, ideally to the point of eliminating the disorder, and/or (ii) reversing the progression of the disorder’s symptoms, ideally to the point of eliminating the symptoms and/or (iii) reducing or eliminating the likelihood of relapse (i.e. consolidation, which is a common goal of post remission therapy for AML and, ideally, results in the destruction of any remaining leukemia cells).

The treatment of hematologic malignancy, such as the treatment of AML, can be measured according to a number of clinical endpoints. These include, without limitation, survival time (such as weeks, months or years of improved survival time, e.g., one, two or more months of additional survival time), and response status (such as complete remission (CR), near complete remission (nCR), very good partial remission (VGPR) and partial remission (PR)).

In one embodiment, treatment of hematologic malignancy, such as the treatment of AML, can be measured in terms of remission. Included here are the following non-limiting examples. (1) Morphologic complete remission (“CR”): ANC $\geq 1,000/\text{mcl}$, platelet count $100,000/\text{mcl}$, $<5\%$ bone marrow blasts, no Auer rods, no evidence of extramedullary disease. (No requirements for marrow cellularity, hemoglobin concentration). (2) Morphologic complete remission with incomplete blood count recovery (“CRi”): Same as CR but ANC may be $<1,000/\text{mcl}$ and/or platelet count $<100,000/\text{mcl}$. (3) Partial remission (PR): ANC $1,000/\text{mcl}$, platelet count $>100,000/\text{mcl}$, and at least a 50% decrease in the percentage of marrow aspirate blasts to 5–25%, or marrow blasts $<5\%$ with persistent Auer rods. These criteria and others are known, and are described, for example, in SWOG Oncology Research Professional (ORP) Manual Volume I, Chapter 11A, Leukemia (2014).

Embodiments of the Invention

This invention provides a first method for treating a subject afflicted with a hematologic malignancy comprising administering to the subject an agent targeting a hematologic malignancy-associated antigen, wherein the subject has a low peripheral cancerous cell burden.

In one embodiment of the first method, the subject’s low peripheral cancerous cell burden is medically induced, such as by administering a pharmaceutical agent like hydroxyurea. Alternatively, the subject’s low peripheral cancerous cell burden can be naturally occurring.

This invention also provides a second method for treating a subject afflicted with a hematologic malignancy and having a high peripheral cancerous cell burden, comprising (i) medically lowering the subject’s peripheral cancerous cell burden, and (ii) while the subject’s peripheral cancerous cell burden is still low, administering to the subject an agent targeting a hematologic malignancy-associated antigen.

Preferably, in these methods, the subject is human. In one embodiment of these methods, the hematologic malignancy is acute myeloid leukemia, and the cancerous cells are leukemic blasts.

The antigen-targeting agent is preferably administered in a sub-saturating dose, and can be administered intravenously or via another route. This agent can be, for example, an anti-CD33 antibody labeled with a cytotoxic agent (e.g., an alpha-emitting isotope like ^{225}Ac).

This invention further provides a third method for treating a human subject afflicted with acute myeloid leukemia comprising administering to the subject an anti-CD33 antibody labeled with an alpha-emitting isotope, wherein (i) the

subject has a low peripheral blast burden, and (ii) the antibody is administered in a sub-saturating dose.

In one embodiment, the subject's low peripheral blast burden is medically induced, and is preferably pharmaceutically induced (e.g., via administering hydroxyurea). Alternatively, the subject's low peripheral blast burden can be naturally occurring.

This invention still further provides a fourth method for treating a human subject afflicted with acute myeloid leukemia and having a high peripheral blast burden, comprising (i) medically lowering the subject's peripheral blast burden, and (ii) while the subject's peripheral blast burden is still low, administering to the subject an anti-CD33 antibody labeled with an alpha-emitting isotope, wherein the antibody is administered in a sub-saturating dose.

Preferably, medically lowering the subject's peripheral blast burden comprises pharmaceutically lowering the subject's peripheral blast burden (e.g., via administering hydroxyurea). The antibody can be administered intravenously or via another route.

This invention still further provides a method for treating a human subject afflicted with acute myeloid leukemia comprising intravenously administering ^{225}Ac -labeled HuM195 to the subject, wherein (i) the subject has a low peripheral blast burden, and (ii) the ^{225}Ac -labeled HuM195 is administered in a sub-saturating dose.

In one embodiment, the subject's low peripheral blast burden is medically induced, and is preferably pharmaceutically induced (e.g., via administering hydroxyurea). Alternatively, the subject's low peripheral cancerous cell burden can be naturally occurring. The ^{225}Ac -labeled HuM195 is preferably administered intravenously.

Finally, this invention provides a method for treating a human subject afflicted with acute myeloid leukemia and having a high peripheral blast burden, comprising (i) medically lowering the subject's peripheral blast burden, and (ii) while the subject's peripheral blast burden is still low, intravenously administering ^{225}Ac -labeled HuM195 to the subject, wherein the ^{225}Ac -labeled HuM195 is administered in a sub-saturating dose.

Preferably, medically lowering the subject's peripheral blast burden comprises pharmaceutically lowering the subject's peripheral blast burden (e.g., via administering hydroxyurea). Also, the ^{225}Ac -labeled HuM195 is preferably administered intravenously.

This invention will be better understood by reference to the examples which follow, but those skilled in the art will readily appreciate that the specific examples detailed are only illustrative of the invention as described more fully in the claims which follow thereafter.

EXAMPLES

Example 1—Structure of ^{225}Ac -Lintuzumab (^{225}Ac -HuM195)

^{225}Ac -Lintuzumab includes three key components; humanized monoclonal antibody HuM195 (generic name, lintuzumab), the alpha-emitting radioisotope ^{225}Ac , and the bi-functional chelate 2-(p-isothiocyanatobenzyl)-1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (p-SCN-Bn-DOTA). As depicted in FIG. 4, HuM195 is radiolabeled using the bi-functional chelate p-SCN-Bn-DOTA that binds to ^{225}Ac and that is covalently attached to the IgG via a lysine residue on the antibody.

Example 2—p-SCN-Bn-DOTA

DOTA, 2-(4-Isothiocyanatobenzyl)-1,4,7,10-tetraazacyclododecane tetraacetic acid (Macrocyclics item code B205-

GMP) is synthesized by a multi-step organic synthesis that is fully described in U.S. Pat. No. 4,923,985.

Example 3—Preparation of ^{225}Ac -Lintuzumab (^{225}Ac -HuM195)

The procedure for preparing ^{225}Ac -Lintuzumab is based on the method described by Michael R. McDevitt, "Design and synthesis of ^{225}Ac radioimmunopharmaceuticals", Applied Radiation and Isotope, 57 (2002), 841-847. The procedure involves radiolabeling the bi-functional chelate, p-SCN-Bn-DOTA, with the radioisotope ^{225}Ac , followed by binding of the radiolabeled p-SCN-Bn-DOTA to the antibody (HuM195). The construct, ^{225}Ac -p-SCN-Bn-DOTA-HuM195, is purified using 10 DG size exclusion chromatography and eluted with 1% human serum albumin (HSA). The resulting drug product, Ac225-Lintuzumab, is then passed through a 0.2 μm sterilizing filter.

Example 4—Process Flow for Preparation of ^{225}Ac -Lintuzumab (^{225}Ac -HuM195)

The procedure, shown in FIG. 5, begins with confirming the identity of all components and the subsequent QC release of the components to production. The ^{225}Ac is assayed to confirm the level of activity and is reconstituted to the desired activity concentration with hydrochloric acid. A vial of lyophilized p-SCNBn-DOTA is reconstituted with metal-free water to a concentration of 10 mg/mL. To the actinium reaction vial, 0.02 ml of ascorbic acid solution (150 mg/mL) and 0.05 ml of reconstituted p-SCN-Bn-DOTA are added and the pH adjusted to between 5 and 5.5 with 2M tetramethylammonium acetate (TMAA). The mixture is then heated at $55\pm 4^\circ\text{C}$. for 30 minutes.

To determine the labeling efficiency of the ^{225}Ac -p-SCN-Bn-DOTA, an aliquot of the reaction mixture is removed and applied to a 1 ml column of Sephadex C25 cation exchange resin. The product is eluted in 2-4 ml fractions with a 0.9% saline solution. The fraction of ^{225}Ac activity that elutes is ^{225}Ac -p-SCN-Bn-DOTA and the fraction that is retained on the column is un-chelated, unreactive ^{225}Ac . Typically, the labeling efficiency is greater than 95%.

To the reaction mixture, 0.22 ml of previously prepared HuM195 in DTPA (1 mg HuM195) and 0.02 ml of ascorbic acid are added. The DTPA is added to bind any trace amounts of metals that may compete with the labeling of the antibody. The ascorbic acid is added as a radio-protectant. The pH is adjusted with carbonate buffer to pH 8.5-9. The mixture is heated at $37\pm 3^\circ\text{C}$. for 30 minutes. The final product is purified by size exclusion chromatography using 10DG resin and eluted with 2 ml of 1% HSA. Typical reaction yields are 10%.

Example 5—Actimab-A Treatment in a Phase 1 Clinical Trial for Relapsed and Refractory AML Patients

Relapsed and refractory adult AML patients of all ages were enrolled in this single treatment group trial. Treatment consisted of a single infusion of Actimab-A administered at escalating doses as shown in Table 1 below.

TABLE 1

Dose Level	^{225}Ac Activity	HuM195 Dose
1	0.5 $\mu\text{Ci/kg}$	15 $\mu\text{g/kg}$
2	1 $\mu\text{Ci/kg}$	15 $\mu\text{g/kg}$

9

TABLE 1-continued

Dose Level	²²⁵ Ac Activity	HuM195 Dose
3	2 μ Ci/kg	20 μ g/kg
4	4 μ Ci/kg	25 μ g/kg
5	3 μ Ci/kg	20 μ g/kg

Each cohort was planned for three patients. Dose escalation proceeded if none of the three patients experienced dose-limiting toxicities (DLTs). If one of the three patients experienced DLTs, the cohort was expanded with additional three patients. If none experienced DLTs, the dose escalation continued. If two out of six patients experienced DLTs, dose level was decreased. The total of 20 patients were treated and their baseline characteristics were collected, including circulating blasts counts. Circulating blasts counts were available for 18 patients. After the treatment, patients were followed up for safety and efficacy, including anti-leukemic effects, which primarily comprises blasts as a percentage of all cells in patients' red bone marrow.

Example 6—Actimab-A Treatment in a Phase 1/2 Trial in Newly Diagnosed AML Patients

In the Phase 1 portion of the Actimab-A Phase 1/2 multicenter trial, newly diagnosed older AML patients ineligible for treatment with intensive chemotherapy regimens were treated with a course of low dose cytarabine followed by escalating fractionated doses (two fractions) of Actimab-A per Table 2 below. If the patient's disease did not progress after treatment with Actimab-A, and if the patient remained otherwise eligible, additional up to 11 low dose cytarabine cycles of treatment were to be administered.

TABLE 2

Dose Level	Ac-225 Activity per fractionated dose (μ Ci/Kg)	Total Dose (μ Ci/Kg)	HuM195 per fractionated dose (μ g/Kg)	Total HuM195 dose (μ g/Kg)
1	0.5	1.0	7.5	15
2	1	2	10	20
3	1.5	3	10	20
4	2	4	12.5	25

Each cohort was planned to enroll 3 patients. Dose escalation proceeded if none of the three patients experienced dose-limiting toxicities ("DLTs"). If one of the three patients experienced DLTs, the cohort was expanded with an additional three patients. If none experienced DLTs, the dose escalation continued. If two out of six patients experienced DLTs, the dose level was decreased. A total of 18 patients were treated and their baseline characteristics were collected, including circulating blast counts. Circulating blast counts were available for all 18 patients. After the treatment, patients were followed up for safety and efficacy, including anti-leukemic effects primarily comprising blasts as a percentage of all cells in patients' red bone marrow.

Example 7—Efficacy Outcomes in an Actimab-A Phase 1 Trial for Relapsed and Refractory AML Patients and a Phase 1/2 Trial for Older Newly Diagnosed AML Patients

Initial anti-leukemic efficacy was analyzed as a function of achieving a composite complete response ("CRc"), where

10

CRc is defined as achieving a post-treatment patient bone marrow blast percentage of <5%. Response rates per dose level in both referenced trials are shown in Tables 3 and 4 below. Table 3 shows efficacy outcomes for the Phase 1 single dose trial in relapsed and refractory AML patients.

TABLE 3

Patients in the Trial			
	Patients Treated	Achieved CRc	% CRc
Dose level 1 (0.5 μ Ci/kg)	3	0	0%
Dose level 2 (1.0 μ Ci/kg)	3	1	33%
Dose level 3 (2.0 μ Ci/kg)	5	0	0%
Dose level 4 (3.0 μ Ci/kg)	5	1	20%
Dose level 5 (4.0 μ Ci/kg)	2	1	50%

Table 4 shows efficacy outcomes for the Phase 1 fractionated dose trial in older newly diagnosed AML patients.

TABLE 4

Patients in the Trial			
	Patients Treated	Achieved CRc	% CRc
Dose level 1 (0.5 μ Ci/kg $\times$ 2)	3	0	0%
Dose level 2 (1.0 μ Ci/kg $\times$ 2)	6	1	17%
Dose level 3 (1.5 μ Ci/kg $\times$ 2)	3	2	67%
Dose level 4 (2.0 μ Ci/kg $\times$ 2)	6	2	33%
Total	18	5	28%

Example 8—Efficacy Analysis by Commonly Used Risk Factors of an Actimab-A Phase 1/2 Trial for Newly Diagnosed AML Patients

There are several commonly used metrics used as prognostic factors to determine the likelihood of response in newly diagnosed AML. The most commonly used prognostic factors are age, cytogenetics and the presence of antecedent hematologic disease (most often myelodysplastic syndrome (MDS, in which case AML is referred to as secondary AML ("sAML")). Those factors were applied to analyze efficacy outcomes in the Phase 1 older newly diagnosed AML trial. Results of those analyses are presented in Table 5 below.

TABLE 5

	Responders	Non-responders
Age (Median)	76	77
Genetic risk	High	Intermediate
Response %	29%	27%
De novo vs sAML	De novo	sAML
Response %	50%	17%

It was concluded that none of the above commonly used prognostic factors correlates with statistically significant differences in response outcomes.

Example 9—Peripheral Blast Burden Correlation with Efficacy Outcomes in Actimab-A Phase 1 and Phase 1/2 Clinical Trials for AML Patients

Additional analyses were performed, and it was surprisingly discovered that the number of circulating blasts has a strong and statistically significant correlation with the clini-

11

cal outcome. This discovery was observed in both trials, i.e., in both relapsed/refractory and newly diagnosed patients. A circulating blast count of fewer than 200 per microliter corresponded with responses. However, when the circulating blast count was greater than or equal to 200 per microliter, no patients responded. These data are presented in Table 6 below.

TABLE 6

Circulating blasts per microliter	≥200	<200
Phase 1 single dose trial	0%	43%
Phase 1/2 fractionated dosing trial	0%	42%

Example 10—Responses in Newly Diagnosed and Relapsed/Refractory AML Patients According to Peripheral Blast Status Prior to Actimab-A Treatment

Response outcomes in AML patients were compared by dose level in the two referenced trials by selecting only patients with <200 circulating blasts per microliter of peripheral blood. Tables 7-9 below summarize these findings.

Table 7 shows efficacy outcomes per peripheral blast counts by dose level in the Phase 1 trial for relapsed and refractory AML patients.

TABLE 7

	Pts pb ≥ 200	Pts with CRc	% CRc	Pts pb < 200	Pts with CRc	% CRc
Dose level 1 (0.5 μCi/kg)	2	0	0%	1	0	0%
Dose level 2 (1 μCi/kg)	0	0	0%	3	1	33%
Dose level 3 (2 μCi/kg)	5	0	0%	0	0	NA
Dose level 4 (3 μCi/kg)	3	0	0%	2	1	50%
Dose level 5 (4 μCi/kg)	1	0	0%	1	1	100%
Total	11	0	0%	7	3	43%

Table 8 shows efficacy outcomes per peripheral blast counts by dose level in the Phase 1/2 trial for newly diagnosed older AML patients.

TABLE 8

	Pts pb ≥ 200	Pts with CRc	% CRc	Pts pb < 200	Pts with CRc	% CRc
Dose level 1 (0.5 μCi/kg × 2)	1	0	0%	2	0	0%
Dose level 2 (1 μCi/kg × 2)	3	0	0%	3	1	33%
Dose level 3 (1.5 μCi/kg × 2)	0	0	0%	3	2	67%
Dose level 4 (2 μCi/kg × 2)	2	0	0%	4	2	50%
Total	6	0	0%	12	5	42%

Table 9 shows a statistical analysis of the responses in both AML trials.

TABLE 9

Patients with:	CRc rates	CRc %
<200 pb	8/19	42%
≥200 pb	0/17	0%

* p value: 0.002416 (highly significant)

12

Example 11—Anti-Leukemic Effect of Actimab-A Based on Significant Leukemic Blast Cell Killing

In early stage clinical trials, anti-leukemic effect is often measure as the ability of the studied therapy to reduce leukemic blast burden in the bone marrow by ≥50%. The ability of Actimab-A to have this effect was analyzed by comparing patients with peripheral blast burdens of ≥200 and patients with peripheral blast burdens of <200 per microliter of peripheral blood. Results of these analyses are presented in Table 10 below. These results represent analyses of 38 patients wherein data were available for 31 of them.

TABLE 10

	≥50% reduced patients (%)	<50% reduced patients (%)
pb < 200	12 (80%)	3 (20%)
pb ≥ 200	4 (25%)	12 (75%)

* p value: 0.002197 (highly significant)

REFERENCES

1. Burnett, et al., "A Comparison of Low-Dose Cytarabine and Hydroxyurea With or Without All-trans Retinoic Acid for Acute Myeloid Leukemia and High-Risk Myelodys-

plastic Syndrome in Patients Not Considered Fit for Intensive Treatment", Cancer, Mar. 15, 2007, Vol. 109, No 6.

2. Co, et al., J. Immunol. 148:1149-1154, 1992.
3. Domino and Baldor, "The 5-Minute Clinical Consult 2014."
4. Gansow, et al., U.S. Pat. No. 4,923,985.
5. Harousseau, et al., "A randomized phase 3 study of tipifarnib compared with best supportive care, including hydroxyurea, in the treatment of newly diagnosed acute myeloid leukemia in patients 70 years or older", Blood, 6 Aug. 2009, Vol. 114, No. 6.
6. Jurcic, et al., "Phase I Trial of Targeted Alpha-Particle Therapy with Actinium-225 (²²⁵Ac)-Lintuzumab and

13

- Low-Dose Cytarabine (LDAC) in Patients Age 60 or Older with Untreated Acute Myeloid Leukemia (AML)", ASH 2016 Abstract.
7. Kumar, "Genetic Abnormalities and Challenges in the Treatment of Acute Myeloid Leukemia", Genes & Cancer 5 2(2):95-107, 2011.
8. McDevitt, "Design and synthesis of ²²⁵Ac radioimmuno-pharmaceuticals", Applied Radiation and Isotope, 57 (2002), 841-847.
9. Mulford et al., "The Promise of Targeted a-Particle 10 Therapy", The Journal of Nuclear Medicine, Vol. 46, No. 1 (Suppl), January 2005.

14

10. Mylotarg Wyeth Product Monograph (2005).
11. Pollard, et al., "Correlation of CD33 expression level with disease characteristics and response to gemtuzumab ozogamicin containing chemotherapy in childhood AML", Blood, 2012, April 19, 119(16):3705-3711.
12. V. H. J. van der Velden, et al., "High CD33-antigen loads in peripheral blood limit the efficacy of gemtuzumab ozogamicin (Mylotarg) treatment in acute myeloid leukemia patients", Leukemia, 2004, May, 18(5):983-8.
13. Scheinberg, et al., U.S. Pat. No. 6,683,162.
14. SWOG Oncology Research Professional (ORP) Manual, Volume I, Chapter 11A, Leukemia (2014).

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 4

<210> SEQ ID NO 1

<211> LENGTH: 1952

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (12)..(407)

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (766)..(1083)

<400> SEQUENCE: 1

```

tctagaccac c atg gag aaa gac aca ctc ctg cta tgg gtc cta ctt ctc      50
      Met Glu Lys Asp Thr Leu Leu Leu Trp Val Leu Leu Leu
          1              5              10

tgg gtt cca ggt tcc aca ggt gac att cag atg acc cag tct ccg agc      98
Trp Val Pro Gly Ser Thr Gly Asp Ile Gln Met Thr Gln Ser Pro Ser
      15              20              25

tct ctg tcc gca tca gta gga gac agg gtc acc atc aca tgc aga gcc      146
Ser Leu Ser Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala
      30              35              40              45

agc gaa agt gtc gac aat tat ggc att agc ttt atg aac tgg ttc caa      194
Ser Glu Ser Val Asp Asn Tyr Gly Ile Ser Phe Met Asn Trp Phe Gln
          50              55              60

cag aaa ccc ggg aag gct cct aag ctt ctg att tac gct gca tcc aac      242
Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ala Ala Ser Asn
          65              70              75

caa ggc tcc ggg gta ccc tct cgc ttc tca ggc agt gga tct ggg aca      290
Gln Gly Ser Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr
          80              85              90

gac ttc act ctc acc att tca tct ctg cag cct gat gac ttc gca acc      338
Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp Asp Phe Ala Thr
          95              100              105

tat tac tgt cag caa agt aag gag gtt ccg tgg acg ttc ggt caa ggg      386
Tyr Tyr Cys Gln Gln Ser Lys Glu Val Pro Trp Thr Phe Gly Gln Gly
      110              115              120              125

acc aag gtg gag atc aaa cgt aagtagaato caaagtctag aaattctaaa      437
Thr Lys Val Glu Ile Lys Arg
          130

ctctgagggg gtcggatgac gtggccattc ttgcctaaa gcattgagtt tactgcaagg      497

tcagaaaagc atgcaaagcc ctccagaatgg ctgcaaagag ctccaacaaa acaatttaga      557

actttattaa ggaatagggg gaagctagga agaaactcaa aacatcaaga ttttaaatac      617

gcttcttggt ctcccttgcta taattatctg ggataagcat gctgttttct gtctgtccct      677

```


-continued

```

aacatgctct gtgattatcc gcaaacaaca cacccaaggg cagaactttg ttacttaaac 737
accatectgt ttgcttcttt cctcagga act gtg gct gca cca tct gtc ttc 789
                               Thr Val Ala Ala Pro Ser Val Phe
                               135                               140
atc ttc ccg cca tct gat gag cag ttg aaa tct gga act gcc tct gtt 837
Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val
                               145                               150                               155
gtg tgc ctg ctg aat aac ttc tat ccc aga gag gcc aaa gta cag tgg 885
Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp
                               160                               165                               170
aag gtg gat aac gcc ctc caa tcg ggt aac tcc cag gag agt gtc aca 933
Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr
                               175                               180                               185
gag cag gac agc aag gac agc acc tac agc ctc agc agc acc ctg acg 981
Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr
                               190                               195                               200
ctg agc aaa gca gac tac gag aaa cac aaa gtc tac gcc tgc gaa gtc 1029
Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val
205                               210                               215                               220
acc cat cag ggc ctg agc tcg ccc gtc aca aag agc ttc aac agg gga 1077
Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly
                               225                               230                               235
gag tgt tagagggaga agtgcccca cctgctctc agttccagcc tgacccctc 1133
Glu Cys
ccatcctttg gcctctgacc ctttttcac aggggacctt cccctattgc ggtcctcag 1193
ctcatctttc acctcacccc cctcctctc cttggcttta attatgctaa tgttgaggga 1253
gaatgaataa ataaagttaa tctttgcacc tgtggtttct ctctttcctc atttaataat 1313
tattatctgt tgttttacca actactcaat ttctcttata agggactaaa tatgtagtca 1373
tcctaaggcg cataaccatt tataaaaaatc atccttcatt ctattttacc ctatcctcct 1433
ctgcaagaca gtctctcctc aaaccacaaa gccttctgtc ctcacagtcc cctggggccat 1493
ggtaggagag acttgcttcc ttgttttccc ctctcagca agccctcata gtctttttta 1553
aggggtgacag gtcttacagt catatacct ttgattcaat tccctgagaa tcaaccaaag 1613
caaatTTTTc aaaagaagaa acctgctata aagagaatca ttcattgcaa catgatataa 1673
aataacaaca caataaaagc aattaaataa acaaacaata gggaaatgtt taagttcatc 1733
atggtactta gacttaatgg aatgtcatgc cttatttaca tttttaaaca ggtactgagg 1793
gactcctgtc tgccaagggc cgtattgagt actttccaca acctaattta atccacacta 1853
tactgtgaga ttaaaaacat tcattaaaat gttgcaaagg ttctataaag ctgagagaca 1913
aatatattct ataactcagc aatcccactt ctaggatcc 1952

```

<210> SEQ ID NO 2

<211> LENGTH: 238

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 2

```

Met Glu Lys Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
1           5           10           15

```

```

Gly Ser Thr Gly Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
20           25           30

```

```

Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Glu Ser

```

-continued

35	40	45
Val Asp Asn Tyr Gly Ile Ser Phe Met Asn Trp Phe Gln Gln Lys Pro		
50	55	60
Gly Lys Ala Pro Lys Leu Leu Ile Tyr Ala Ala Ser Asn Gln Gly Ser		
65	70	75 80
Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr		
	85	90 95
Leu Thr Ile Ser Ser Leu Gln Pro Asp Asp Phe Ala Thr Tyr Tyr Cys		
	100	105 110
Gln Gln Ser Lys Glu Val Pro Trp Thr Phe Gly Gln Gly Thr Lys Val		
	115	120 125
Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro		
	130	135 140
Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu		
145	150	155 160
Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn		
	165	170 175
Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser		
	180	185 190
Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala		
	195	200 205
Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly		
	210	215 220
Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys		
225	230	235
<210> SEQ ID NO 3		
<211> LENGTH: 3246		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic		
polynucleotide		
<220> FEATURE:		
<221> NAME/KEY: CDS		
<222> LOCATION: (12)..(416)		
<220> FEATURE:		
<221> NAME/KEY: CDS		
<222> LOCATION: (660)..(953)		
<220> FEATURE:		
<221> NAME/KEY: CDS		
<222> LOCATION: (1345)..(1389)		
<220> FEATURE:		
<221> NAME/KEY: CDS		
<222> LOCATION: (1508)..(1837)		
<220> FEATURE:		
<221> NAME/KEY: CDS		
<222> LOCATION: (1935)..(2255)		
<400> SEQUENCE: 3		
tctagaccac c atg gga tgg agc tgg atc ttt ctc ttc ctc ctg tca gga		50
Met Gly Trp Ser Trp Ile Phe Leu Phe Leu Leu Ser Gly		
1 5 10		
act gct ggc gtc cac tct cag gtt cag ctg gtg cag tct gga gct gag		98
Thr Ala Gly Val His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu		
15 20 25		
gtg aag aag cct ggg agc tca gtg aag gtt tcc tgc aaa gct tct ggc		146
Val Lys Lys Pro Gly Ser Ser Val Lys Val Ser Cys Lys Ala Ser Gly		
30 35 40 45		
tac acc ttc act gac tac aac atg cac tgg gtg agg cag gct cct ggc		194
Tyr Thr Phe Thr Asp Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly		
50 55 60		

-continued

caa ggc ctg gaa tgg att gga tat att tat cct tac aat ggt ggt acc Gln Gly Leu Glu Trp Ile Gly Tyr Ile Tyr Pro Tyr Asn Gly Gly Thr 65 70 75	242
ggc tac aac cag aag ttc aag agc aag gcc aca att aca gca gac gag Gly Tyr Asn Gln Lys Phe Lys Ser Lys Ala Thr Ile Thr Ala Asp Glu 80 85 90	290
agt act aac aca gcc tac atg gaa ctc tcc agc ctg agg tct gag gac Ser Thr Asn Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp 95 100 105	338
act gca gtc tat tac tgc gca aga ggg cgc ccc gct atg gac tac tgg Thr Ala Val Tyr Tyr Cys Ala Arg Gly Arg Pro Ala Met Asp Tyr Trp 110 115 120 125	386
ggc caa ggg act ctg gtc act gtc tct tca ggtaagaatg gcctctagac Gly Gln Gly Thr Leu Val Thr Val Ser Ser 130 135	436
caccatggga tggagctttc tggggcaggc caggcctgac cttggctttg gggcagggag	496
ggggctaagg tgaggcaggt ggcgccagcc aggtgcacac ccaatgccca tgagcccaga	556
cactggacgc tgaacctgcg ggacagttaa gaacccaggg gcctctgcgc cctgggccc	616
gctctgtccc acaccgcggt cacatggcac cacctctctt gca gcc tcc acc aag Ala Ser Thr Lys	671
ggc cca tcg gtc ttc ccc ctg gca ccc tcc tcc aag agc acc tct ggg Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly 140 145 150 155	719
ggc aca gcg gcc ctg ggc tgc ctg gtc aag gac tac ttc ccc gaa ccg Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro 160 165 170	767
gtg acg gtg tcg tgg aac tca ggc gcc ctg acc agc ggc gtg cac acc Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr 175 180 185	815
ttc ccg gct gtc cta cag tcc tca gga ctc tac tcc ctc agc agc gtg Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val 190 195 200	863
gtg acc gtg ccc tcc agc agc ttg ggc acc cag acc tac atc tgc aac Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn 205 210 215	911
gtg aat cac aag ccc agc aac acc aag gtg gac aag aaa gtt Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val 220 225 230	953
ggtgagaggc cagcacaggg agggagggtg tctgctggaa gccaggctca gcgtcctgc	1013
ctggacgcat cccggctatg cagccccagt ccagggcagc aaggcaggcc ccgtctgcct	1073
cttcaccccg aggcctctgc ccgccccact catgctcagg gagagggtct tctggctttt	1133
tccccaggct ctgggcaggc acaggctagg tgcccctaac ccaggccctg cacacaaagg	1193
ggcaggtgct gggctcagac ctgccaaagag ccatatccgg gaggaccctg cccctgacct	1253
aagccccccc caaaggccaa actctccact ccctcagctc ggacaccttc tctctcccca	1313
gattccagta actcccaatc ttctctctgc a gag ccc aaa tct tgt gac aaa Glu Pro Lys Ser Cys Asp Lys 235 240	1365
act cac aca tgc cca ccg tgc cca ggtaagccag ccaggccctc gccctccagc Thr His Thr Cys Pro Pro Cys Pro 245	1419
tcaaggcggg acagggtgcc tagagtagcc tgcattccagg gacaggcccc agccgggtgc	1479
tgacacgtcc acctccatct ctctctca gca cct gaa ctc ctg ggg gga ccg Ala Pro Glu Leu Leu Gly Gly Pro 250 255	1531

-continued

tca gtc ttc ctc ttc ccc cca aaa ccc aag gac acc ctc atg atc tcc	1579
Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser	
260 265 270	
cgg acc cct gag gtc aca tgc gtg gtg gtg gac gtg agc cac gaa gac	1627
Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp	
275 280 285	
cct gag gtc aag ttc aac tgg tac gtg gac ggc gtg gag gtg cat aat	1675
Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn	
290 295 300	
gcc aag aca aag ccg cgg gag gag cag tac aac agc acg tac cgt gtg	1723
Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val	
305 310 315 320	
gtc agc gtc ctc acc gtc ctg cac cag gac tgg ctg aat ggc aag gag	1771
Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu	
325 330 335	
tac aag tgc aag gtc tcc aac aaa gcc ctc cca gcc ccc atc gag aaa	1819
Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys	
340 345 350	
acc atc tcc aaa gcc aaa ggtgggaccc gtgggggtgag agggccacat	1867
Thr Ile Ser Lys Ala Lys	
355	
ggacagaggc cggtctcgcc caccctctgc cctgagagtg accgctgtac caacctctgt	1927
ccctaca ggg cag ccc cga gaa cca cag gtg tac acc ctg ccc cca tcc	1976
Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser	
360 365 370	
cgg gat gag ctg acc aag aac cag gtc agc ctg acc tgc ctg gtc aaa	2024
Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys	
375 380 385	
ggc ttc tat ccc agc gac atc gcc gtg gag tgg gag agc aat ggg cag	2072
Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln	
390 395 400	
ccg gag aac aac tac aag acc acg cct ccc gtg ctg gac tcc gac ggc	2120
Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly	
405 410 415 420	
tcc ttc ttc ctc tac agc aag ctc acc gtg gac aag agc agg tgg cag	2168
Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln	
425 430 435	
cag ggg aac gtc ttc tca tgc tcc gtg atg cat gag gct ctg cac aac	2216
Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn	
440 445 450	
cac tac acg cag aag agc ctc tcc ctg tct ccg ggt aaa tgagtgcgac	2265
His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys	
455 460 465	
ggcccggaag ccccgctcc ccgggctctc gcggctgcac gaggatgctt ggcacgtacc	2325
ccctgtacat acttcccggg cgcccagcat ggaaataaag caccagcgc tgcctggggc	2385
ccctgcgaga ctgtgatggt tctttccacg ggtcaggccg agtctgaggc ctgagtggca	2445
tgagggaggc agagcgggtc ccaactgtccc cacactggcc caggctgtgc aggtgtgcct	2505
ggggcgccta ggggtgggct cagccagggg ctgccctcgg cagggtgggg gatttgccag	2565
cgtggccctc cctccagcag cacctgccct gggctgggccc acgggaagcc ctaggagccc	2625
ctggggacag acacacagcc cctgcctctg taggagactg tctgttctg tgagcgcct	2685
gtcctccgac ctccatgccc actcgggggc atgcctagtc catgtgcgta gggacaggcc	2745
ctccctcacc catctacccc cagggcacta acccctggct gccctgcccc gcctcgcacc	2805
cgcattgggga cacaaccgac tccggggaca tgcaactctcg ggccctgtgg agggactggt	2865

-continued

```
gcagatgccc acacacacac tcagcccaga cccgttcaac aaaccccgca ctgaggttgg 2925
cgggccacac ggccaccaca cacacacgtg cagcctcac acacggagcc tcaccggggc 2985
gaactgcaca gcacccagac cagagcaagg tctctgcaca cgtgaacact cctcggacac 3045
aggcccccac gagccccacg cggcacctca aggcccacga gcctctcggc agcttctcca 3105
catgctgacc tgctcagaca aacccagccc tctctcaca aggggtgcccc tgcagccgcc 3165
acacacacac aggggatcac acaccacgtc acgtccctgg ccctggccca cttcccagtg 3225
ccgcccttcc ctgcaggatc c 3246
```

```
<210> SEQ ID NO 4
<211> LENGTH: 465
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide
```

```
<400> SEQUENCE: 4
```

```
Met Gly Trp Ser Trp Ile Phe Leu Phe Leu Leu Ser Gly Thr Ala Gly
 1             5             10            15
Val His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
      20             25            30
Pro Gly Ser Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
      35             40            45
Thr Asp Tyr Asn Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu
      50            55            60
Glu Trp Ile Gly Tyr Ile Tyr Pro Tyr Asn Gly Gly Thr Gly Tyr Asn
      65            70            75            80
Gln Lys Phe Lys Ser Lys Ala Thr Ile Thr Ala Asp Glu Ser Thr Asn
      85            90            95
Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
      100           105           110
Tyr Tyr Cys Ala Arg Gly Arg Pro Ala Met Asp Tyr Trp Gly Gln Gly
      115           120           125
Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
      130           135           140
Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
      145           150           155           160
Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
      165           170           175
Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
      180           185           190
Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
      195           200           205
Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro
      210           215           220
Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys
      225           230           235           240
Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro
      245           250           255
Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
      260           265           270
Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp
      275           280           285
```

-continued

Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn
290						295					300				
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val
305					310					315					320
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu
			325						330					335	
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys
			340					345					350		
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr
		355					360					365			
Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr
	370					375					380				
Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu
385					390					395					400
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu
			405						410					415	
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys
			420					425					430		
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu
		435				440						445			
Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly
	450					455					460				

Lys
465

What is claimed is:

1. A method for treating relapsed and/or refractory acute myeloid leukemia in a human subject, comprising the step of:

intravenously administering a therapeutically effective dose of ²²⁵Ac-labeled HuM195 to a human subject afflicted with relapsed and/or refractory acute myeloid leukemia, wherein the human subject has a peripheral blast burden below 200 blast cells/μl.

2. The method of claim 1, further comprising the step of before administering the ²²⁵Ac-labeled HuM195 to the human subject, determining that the human subject has a peripheral blast burden below 200 blast cells/μl.

3. The method of claim 1, wherein the human subject is ineligible to receive high intensity chemotherapy.

4. The method of claim 2, wherein the human subject is ineligible to receive high intensity chemotherapy.

5. The method of claim 1, further comprising the step of: before administering the ²²⁵Ac-labeled HuM195 to the human subject, administering low dose cytarabine to the human subject.

6. The method of claim 2, further comprising the step of: before administering the ²²⁵Ac-labeled HuM195 to the human subject, administering low dose cytarabine to the human subject.

7. The method of claim 3, further comprising the step of: before administering the ²²⁵Ac-labeled HuM195 to the human subject, administering low dose cytarabine to the human subject.

8. The method of claim 4, further comprising the step of: before administering the ²²⁵Ac-labeled HuM195 to the human subject, administering low dose cytarabine to the human subject.

9. The method of claim 1, wherein the human subject's peripheral blast burden is 50 blast cells/μl or lower.

10. The method of claim 1, further comprising the step of: before administering the ²²⁵Ac-labeled HuM195 to the human subject, medically lowering the human subject's peripheral blast burden from at or above 200 blast cells/μl to below 200 blast cells/μl,

wherein the administering step is performed while the human subject's peripheral blast burden is still below 200 blast cells/μl.

11. The method of claim 10, wherein the human subject's peripheral blast burden is medically lowered to 50 blast cells/μl or lower.

12. The method of claim 5, further comprising the step of: before administering the ²²⁵Ac-labeled HuM195 to the human subject, medically lowering the human subject's peripheral blast burden from at or above 200 blast cells/μl to below 200 blast cells/μl,

wherein the administering step is performed while the human subject's peripheral blast burden is still below 200 blast cells/μl.

13. The method of claim 12, wherein the human subject's peripheral blast burden is medically lowered to 50 blast cells/μl or lower.

* * * * *