

US Patent & Trademark Office

Patent Public Search | Text View

United States Plant Patent

PP36958

Kind Code

P2

Date of Patent

September 16, 2025

Inventor(s)

Reynbery; Jared et al.

***Cannabis* plant named ‘PTC8510V’**

Abstract

A new *Cannabis* variety distinguished by its maturity time, apical dominance, internode length, flower trichome density, aroma, very high THCV:THC ratio, and total cannabinoid content.

Latin Name: Cannabis sativa PTC8510V

Inventors: Reynbery; Jared (Portland, OR), Bakker; Erica (Portland, OR)

Applicant: Phylos Bioscience, Inc. (Portland, OR)

Assignee: Phylos Bioscience, Inc. (Portland, OR)

Appl. No.: 18/902526

Filed: September 30, 2024

Publication Classification

Int. Cl.: A01H5/02 (20180101); A01H6/28 (20180101)

U.S. Cl.:

USPC PLT/258

Field of Classification Search

CPC: A01H (5/02); A01H (5/12); A01H (5/00); A01H (6/28)

USPC: PLT/258; PLT/263.1

Background/Summary

(1) Latin name of the genus and species of the plant claimed: *Cannabis sativa*.

(2) Variety denomination: ‘PTC8510V’.

BACKGROUND

(3) *Cannabis* is an annual, dioecious, flowering herb. *Cannabis* plants produce a variety of secondary metabolites, including cannabinoids, terpenoids, and other compounds, which are often secreted by glandular trichomes that occur most abundantly on the floral calyxes and bracts of female plants. Cannabinoids are the most studied group of secondary metabolites in *Cannabis*. Most exist in two forms, as acids and in neutral (decarboxylated) forms. The acid form is designated by an “A” at the end of its acronym (i.e. THCA). The biologically active forms for human consumption are the neutral forms. Cannabinoids are typically synthesized in the plant as acid forms. Although some decarboxylation can occur in the plant, it can be significantly increased post-harvest by thorough drying of plant material followed by heating it, often by either combustion, vaporization, or heating in an oven. For most cannabinoid producing purposes, only female plants are desired. The presence of male flowers is considered undesirable as pollination is known to reduce cannabinoid yield. For this reason, many *Cannabis* varieties are grown “sinsemilla” through vegetative (i.e., asexual) propagation.

SUMMARY

(4) The present application relates to a new and distinct variety of *Cannabis* plant named ‘PTC8510V’. This new *Cannabis* variety was selected when growing in a cultivated area in Hillsboro, Oregon. ‘PTC8510V’ is commercially known as “Powerhouse,” and is also known by its breeder name 22TP1C-141-105.

(5) ‘PTC8510V’ is a selection resulting from pollen receiver plant (‘21TP1B-3-7’, unpatented), which has a THCV content of 14.17% and a THC content of 4.41%, crossed with pollen donor plant (‘22TX1-2-59’, unpatented), which has a THCV content of 10.55% and THC content of 15.65%. 112 seeds of this cross were planted Nov. 7, 2022. ‘PTC8510V’ was discovered and selected as a single plant from F1 seed and was subsequently clonally propagated.

(6) Parent cultivar ‘21TP1B-3-7’ is an early maturing (7-9 weeks) cultivar with low apical dominance, short internode lengths, flowers with moderate trichome density, a THCV content of 10-15%, and a THC content of 4-6%, resulting in a THCV:THC ratio of roughly 3:1. Parent cultivar ‘22TX1-2-59’ is a medium maturing variety (8-9 weeks) with high apical dominance, long internode lengths, flowers with low trichome density, a THCV content of 10-13%, and a THC content of 13-16%, resulting in a THCV:THC ratio of roughly 1:1. For comparison, ‘PTC8510V’ is a medium maturing variety (8-10 weeks) with moderately low apical dominance, long internode length, flowers with moderate trichome density, a THCV content of 14-19%, and a THC content of 2-4%, resulting in a THCV:THC ratio of roughly 5:1.

(7) ‘PTC8505V’ is a cultivar that has been commercially available and is of similar pedigree to ‘PTC8510V’, but is not a direct parent or sibling. ‘PTC8510V’ is distinguished from ‘PTC8505V’ at least on the basis that ‘PTC8510V’ has a very high THCV:THC ratio of 5:1, a total THCV of 14.29-16.54%, and a spicy-lime like aroma. In contrast, ‘PTC8505V’ has at most a 3:1 ratio of THCV:THC, a total THCV of about 13.44%, and a piney, grassy aroma.

(8) ‘PTC8510V’ can be distinguished from other known *Cannabis* varieties at least by the following unique combination of characteristics: spicy lime-like aroma with hints of cedar, very high THCV:THC ratio (5:1), total THCV of 14.29-16.54%, total cannabinoid content of about 20%, medium maturity time, moderately low apical dominance, long internode length, and moderate flower trichome density.

(9) ‘PTC8510V’ was asexually reproduced for at least three cycles by apical and sub-apical semi-herbaceous stem cuttings in Hillsboro, Oregon. Asexual reproduction of ‘PTC8510V’ shows that the characteristics as herein described come true to form, are firmly fixed, and are established and transmitted through succeeding propagations.

Description

BRIEF DESCRIPTION OF THE DRAWING(S)

(1) FIG. 1 shows an 86-day old greenhouse grown plant of 'PTC8510V'. The photograph was taken under ambient greenhouse light conditions in September 2024.

(2) FIG. 2 shows a close-up of the flowers of 'PTC8510V'. The photograph was taken under ambient greenhouse light conditions in September 2024.

(3) The color photographs show typical specimens of the new variety and depict the color as nearly true as is reasonably possible to make the same in a color illustration of this character. It should be noted that colors may vary, for example due to lighting conditions at the time the photograph is taken. Therefore, color characteristics of this new variety should be determined with reference to the observations described herein, rather than from the photograph alone.

DETAILED DESCRIPTION

(4) The following detailed description of the 'PTC8510V' variety is based on observations of asexually reproduced progeny. Measurements were taken 35 days after flower initiation, unless otherwise indicated. The following detailed description concerns plants growing in a greenhouse in Hillsboro, Oregon. All color codes are hexadecimal color codes. Colors were detected using the Color Detector Android® smart phone application by Sadens Studio™ under ambient greenhouse light conditions in Hillsboro, Oregon.

(5) Certain characteristics of this variety may change with changing environmental conditions (such as photoperiod, temperature, moisture, soil conditions, nutrient availability, or other factors) without change to the genotype. Color descriptions and other terminology are used in accordance with their ordinary dictionary descriptions, unless the context clearly indicates otherwise. Scientific name: *Cannabis sativa*. Parentage: *Pollen receiver plant*.—'21TP1B-3-7'. *Pollen plant*.—'22TX1-2-59'. Plant: *Growth habit*.—An upright, tap-rooted annual plant, forming fibrous roots when asexually propagated. *Shape*.—Simple Erect. *Plant height*.—1.53-1.74 m. *Plant width*.—1-1.1 m. *Flowering plant height*.—23-31 cm. *Harvest plant height*.—85-101 cm. *Plant stretch*.—54-76 cm. *Plant % stretch*.—174-309. *Plant wet biomass*.—0.51-0.72 kg. *Plant wet bucked weight*.—0.38-0.61 kg. *Proportion female plants*.—100% Pistillate. *Proportion male plants*.—No staminate flowers produced naturally; however, can be induced with stress or ethylene-blocking compounds. *Proportion hermaphrodite plants*.—Less than 5%. *Time to harvest*.—10-12 weeks. *Hardiness*.—Hardy to tolerant of warm and dry conditions without showing stress. Not tolerant to soil oversaturated with water for long periods. This plant has not been tested in outdoor conditions and therefore the USDA hardiness zone is unknown. Stem: *Stem shape*.—Hollow, ribbed, texture. *Stem length*.—121-130 cm. *Stem strength*.—Strong and upright. *Stem diameter at base*.—2-2.1 cm. *Stem color*.—C9E782. *Stem texture*.—Smooth with shallow grooves. *Depth of main stem ribs/grooves*.—Shallow. *Internode length*.—3.2-8.4 cm. Leaf: *Shape*.—Palmately compound. *Structure*.—Serrated margins, lightly acicular to lanceolate leaflets, tapering to an acuminate apex. *Length at maturity (including petiole)*.—17-21 cm. *Average width*.—10.5 cm. *Arrangement*.—Alternate. *Margins*.—Jaggedly serrate with each tooth apex angled toward leaflet apex. *Leaf color (adaxial)*.—578C67. *Leaf color (abaxial)*.—70926A. *Glossiness*.—Slightly glossy on the upper surface. *Leaf hairs*.—Present on both upper and lower surfaces, predominant leaf hairs are glandular capitate-stalked. *Petiole color*.—d9f577. *Petiole anthocyanin color and intensity*.—E5D4BA; medium intensity during vegetative stage; strong during flowering stage. *Petiole length*.—5.2-6.9 cm. *Petiole diameter*.—1-2 mm. *Petiole trichome type*.—Predominantly non-glandular. *Stipule length at maturity*.—6-7 mm. *Stipule width at maturity*.—<1 mm. *Stipule shape*.—Linear with slight curvaceous growth. *Stipule color*.—F0F3B5. *Stipule trichome type*.—Predominantly glandular capitate-stalked. *Number of leaflets*.—3-7. *Central leaflet length*.—13-14 cm. *Central leaflet width*.—2.6-3.2 cm. *Central leaflet length/width ratio*.—4-4.8. *Central leaflet number of teeth*.—22-24.

Vein/midrib shape.—Obliquely continuous throughout leaflet. *Vein/midrib color*.—749855. *Typical venation pattern*.—Pinnately veined leaflets. *Sugar leaf*.—Palmately compound, 39 mm long and 32 mm wide. Predominant trichomes are glandular capitate-stalked. Upper-side sugar leaf color is 578C67 and lower-side leaf color is 70926A. *Aroma*.—Spicy lime-like aroma with hints of cedar. Female/pistillate flower: *Inflorescence type*.—Elongated compound cymes. *Inflorescence position*.—Terminal and axillary. *Number of inflorescences per plant*.—110-170. *Length of inflorescences*.—2-23 cm. *Flower arrangement*.—Cymose. *Flower shape*.—Urceolate. *Flower (individual pistillate) length*.—4-6 mm. *Flower (individual pistillate) depth*.—2-4 mm. *Flower (compound cyme) diameter*.—5-5.5 cm. *Petal description*.—Apetalous. *Bract shape*.—Urceolate. *Bract color*.—Upper surfaces and lower surfaces are 8FBF63. *Typical bract number*.—3-200 bracts per inflorescence. *Bract length*.—4-7 mm. *Bract width*.—2-3 mm. *Bract apex and base*.—Bracts have a rounded base and caudate apex. *Bract texture*.—Smooth and covered with sticky trichomes. *Bracteoles*.—Not present. *Calyx shape*.—No defined calyx. *Corolla*.—No defined corolla. *Pistils per flower*.—One. *Pistil length*.—8-10 mm. *Stigma shape*.—Acute. *Number of stigmas per flower*.—Two. *Stigma length*.—3-8 mm. *Stigma width*.—<1 mm. *Stigma color*.—E8EDB0. *Style*.—Average length is 5 mm, EB6734 in color at maturity. *Trichome shape*.—Capitate-Stalked Glandular. *Trichome color*.—A4B897. *Terminal inflorescence shape*.—Oblong. *Terminal inflorescence color*.—2D5416 & EBEEB4. *Terminal inflorescence length*.—18-22 cm. *Terminal inflorescence width*.—4-5.5 cm. *Pedicel*.—Absent. *Peduncle length*.—5 mm. *Peduncle diameter*.—2 mm. *Peduncle texture*.—Lightly ribbed. *Peduncle color*.—559E57. *Peduncle trichome*.—Predominantly non-glandular. *Staminate shape*.—No staminate flowers produced naturally; however, male flower (staminate) can be induced with stress or ethylene-blocking compounds. *Numbers of flowers per plant*.—Hundreds to thousands flowers per plant, multitudinous, congested with high concentrations on bushier wide inflorescence. *Natural blooming time*.—Autumn. *Fresh flower fragrance*.—Citrus/Pine. Seed: *Seed shape*.—Striped, smooth and globular. *Seed size*.—1.6-2.3 mm.

(6) TABLE-US-00001 TABLE 4 Cannabinoid Content (% dry weight of flower) as determined by High Performance Liquid Chromatography (HPLC). Metabolite Minimum Maximum total CBD 0 0 total CBDV 0.17 0.19 total THC 2.61 3.28 total THCV 14.29 16.54 total cannabinoids 17.07 19.58

(7) TABLE-US-00002 TABLE 5 Terpene Content (% dry weight of flower) as determined by Gas Chromatography. Metabolite Minimum Maximum alpha-pinene 0.063 0.138 camphene 0.009 0.011 beta-pinene 0.073 0.124 delta-3-carene <LOQ <LOQ beta-myrcene 0.572 1.739 alpha-terpinene <LOQ <LOQ D-limonene 0.119 0.247 beta-ocimene 0.094 0.342 p-cymene <LOQ <LOQ gamma-terpinene <LOQ <LOQ terpinolene <LOQ <LOQ linalool 0.044 0.076 isopulegol <LOQ <LOQ geraniol <LOQ <LOQ beta-caryophyllene 0.232 0.309 alpha-humulene 0.061 0.081 nerolidol 1 <LOQ <LOQ guaial 0.01 0.01 caryophyllene oxide <LOQ <LOQ alpha-bisabolol 0.034 0.043 eucalyptol <LOQ <LOQ nerolidol 2 0.104 0.167 total terpenes 1.4 3.19 Top terpenes: Beta-myrcene, beta-ocimene, beta-caryophyllene, and D-limonene. Disease/pest resistance: Unknown. Molecular markers: A unique combination of 29 single nucleotide polymorphism (SNP) markers was identified for variety 'PTC8510V'. The markers are provided in Table 6. The combination of markers disclosed for 'PTC8510V' are unique to this variety, thus these markers can be used, for example, to genotype *Cannabis* plants to identify variety 'PTC8510V'.

(8) TABLE-US-00003 TABLE 6 Single Nucleotide Polymorphism (SNP) Markers of 'PTC8510V'. The following table refers to chromosome positions of the publicly available Abacus Cannabis reference genome (Csat_AbacusV2, NCBI assembly accession GCA_025232715.1; also referred to as "CsaAba2"). Chromosome Position SNP ID REF ALT 'PTC8510V' CsaAba2_1 37927202 141916_45118 T C C/C CsaAba2_1 42508653 262_823098 G T G/G CsaAba2_1 45592057 121785_197 G A A/A CsaAba2_2 4020302 167_3219271 A C A/A CsaAba2_2 6539925 167_940812 T C C/C CsaAba2_2 78649906 347_110280 T A A/A CsaAba2_2 91155632

158_767308 G A G/G CsaAba2_2 92450873 366_239268 C T C/C CsaAba2_3 14960943
120687_9128 A G A/A CsaAba2_3 53074525 141717_593569 T A T/T CsaAba2_3 76672303
142214_2388819 C G G/G CsaAba2_4 14480629 142704_8363492 T C T/T CsaAba2_4 33805380
142250_6863690 G A G/G CsaAba2_4 68786255 142078_499253 C A C/C CsaAba2_4 72692194
142078_3894722 C T T/T CsaAba2_4 73898859 142078_4996873 T C T/T CsaAba2_5 4965109
142050_513504 C T C/C CsaAba2_5 75432249 422_2180389 A C A/A CsaAba2_6 74959100
164_2634629 A G G/G CsaAba2_8 9471523 171_8847435 C A C/C CsaAba2_8 52088446
141527_667545 A G G/G CsaAba2_9 9551328 424_4709922 G C G/G CsaAba2_9 10568626
424_5631426 A G A/A CsaAba2_X 23172429 142055_4365541 C T T/T CsaAba2_X 51310004
142293_5209347 A G G/G CsaAba2_X 51844299 142293_5615610 C T T/T CsaAba2_X
58645424 132592_6808 G T T/T CsaAba2_X 60242150 102945_11717 T C C/C CsaAba2_X
71949822 421_5214874 C T T/T

Claims

1. A new and distinct variety of *Cannabis* plant named ‘PTC8510V’, substantially as illustrated and described herein.
